

## SPAIN

### Submission of information on forest management reference levels by Spain

as requested by the Cancún decisions, i.e. „Consideration of further commitments for Annex I Parties under the Kyoto Protocol, Draft conclusions proposed by the Chair”, contained in FCCC/KP/AWG/2010/L.8, and its Addendum: Draft decision [-/CMP.6], Land use, land-use change and forestry, contained in FCCC/KP/AWG/2010/L.8/Add.2

#### **1 Forest management reference level value**

The values reflected in table 1 for forest management reference levels were calculated as averages of the projected business as usual emissions and removals in forest area under management (or forest land remaining forest land) for the period 2013-2020, taking account of policies implemented before April 2009. The Reference level A includes HWP using the first order decay functions whereas Reference level B assumes instant oxidation for HWP.

**Table 1. Value of proposed reference levels (Gg CO<sub>2</sub>eq).**

Reference level*	
(A)	(B)
-23.889	-21.442

\* The reported values are averages of the projected FM data series for the period 2013-2020, taking account of policies implemented before April 2009. (A) with emissions/removals from HWP using the first order decay functions; (B) assuming instant oxidation (provided for transparency reasons only)

#### **2 General description**

Projections for Spain are provided by the Joint Research Centre of the European Commission (JRC), based on elaboration of the results of independent EU modeling groups, coordinated by the International Institute for Applied Systems Analysis (IIASA), assisted by the JRC and funded by the European Commission Directorate General of Climate Action (DG CLIM). Spain is in the process of improving its national projections for the LULUCF sector, and these improvements are not going to be available on time for the submission on the 28th of February, this is the reason for deciding using the JRC projections.

When constructing the RL, all elements mentioned in footnote 1 of paragraph 4 of the decision -/CMP.6 on LULUCF were taken into account:

- Removals or emissions from forest management as shown in greenhouse gas inventories and relevant historical data: taken into account by adjusting results of the modeling exercise through an “ex-post processing of models results” (see section 5 “Description of construction of reference levels”). This ex-post processing also took into account the need for consistency with the inclusion of carbon pools.
- Age-class structure: models used the latest available country specific age-class structure data (see section 5 “Description of construction of reference levels”).

- c) Forest management activities already undertaken: indirectly taken into account through the use of the latest available forest time series data (from national forest inventory or other country statistics), and the estimation of the evolution of harvest demand by 2020 based on macroeconomic drivers and the application of policies implemented in the Member States by April 2009 and legislative provisions adopted by April 2009 (see section 6, “Policies included”). In the case of Spain, the National Forest Plan has been taken into account; this NFP has a validity of 30 years (from 2002 to 2032) and it is available in Spanish in [http://www.mma.es/portal/secciones/biodiversidad/montes\\_politica\\_forestal/estrategia\\_monte/pfe.htm](http://www.mma.es/portal/secciones/biodiversidad/montes_politica_forestal/estrategia_monte/pfe.htm). Spain plans to review this Plan during 2011, depending on available resources. The revision should then be ready for 2012.
- d) Projected forest management activities under a business as usual scenario: taken into account through the estimation of the evolution of harvest demand by 2020 based on macroeconomic drivers and the application of policies implemented in the Member States by April 2009 and legislative provisions adopted by April 2009 (see section 6 “Policies included”). As mentioned before, the framework policy for forest actions in Spain is the National Forest Plan,
- e) Continuity with the treatment of forest management in the first commitment period; not relevant.
- f) The need to exclude removals from accounting in accordance with decision 16/CMP.1, paragraph 1. The projections included in this submission follow the general principles that govern the treatment of land use, land-use change and forestry activities.

Reference levels including and excluding “force majeure” should be provided. Force majeure has not been defined yet, therefore, it is difficult to exclude the emissions from extreme events (forest fires, in the case of Spain) from the reference level. Depending on the definition adopted, and in the case Spain decides to use force majeure clause, the consistency between RL and the accounting in the period 2013-2020 will have to be ensured. At this moment, all the past emissions from forest fires are included in the reference level as part of historical emissions/removals.

### **3 Pools and gases**

The projected reference level includes the emissions and removals of the following pools: aboveground biomass and belowground biomass. The rest of the pools (carbon organic soil, dead wood and litter) haven’t been considered, and, following the provision included in Decision 16/CMP.1, it has been demonstrated in the National Inventory Report that these pools are not GHG sources.

The reason for omitting these pools is that the emissions/removals associated to those pools are not included in the historical data, therefore, their influence can not be projected in the future.

As mentioned before for force majeure, the consistency between reference level and the accounting in the period 2013-2020 regarding pools will be ensured.

**Table 2. Carbon pools and GHG sources included in the reference level.**

Change in C pool included in the reference level						GHG sources included in the reference level					
Above-ground biomass	Below-ground biomass	Litter	Dead wood	Soil		Fertilization	Drainage of soils	Liming	Biomass burning		
				mineral	organic				CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
yes	yes	no	no	no	no	no	no	no	yes	yes	yes

Yes/No indicate if the pool or gas is included or not in the projections used to set the reference level. The information on the coverage of pools and gases is taken by the JRC from the national GHG inventory report under the KP / UNFCCC (FL remaining FL). In the latter case, if "living biomass" is reported, it is assumed that it contains both aboveground and belowground biomass.

In the case of Spain, in the National Inventory Report is demonstrated that litter, dead wood and soil are pools that are not sources, therefore, they are not included in the estimation of emissions/removals from LULUCF activities. It is worth mentioning that there is no significant area of organic soils in Spain.

Fertilization, drainage and liming are not forest management practices used in Spanish forests; therefore, associated emissions are not included either in the historical data or in the projected reference level.

#### **4 Approaches, methods and models used**

The models used to project emissions and removals from FM are G4M (from IIASA) and EFISCEN (from the European Forest Institute, EFI). Table 3 and figure 1 below provide the essential features of the main models involved and an overview of the modeling architecture.

The reference level builds on macro projections of GDP and population which are exogenous to the models used. They reflect the recent economic downturn, followed by sustained economic growth resuming after 2010. This data is entering GLOBIOM model that uses these projections to translate them into demand for timber (see main assumptions for the BASELINE scenario on pp.13-16 in Capros et al. (2010)<sup>1</sup> for more information). Bioenergy demand was projected by the PRIMES biomass model (see [http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/THE\\_NEW\\_PRIMES\\_BIOMASS\\_MODEL.pdf](http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/THE_NEW_PRIMES_BIOMASS_MODEL.pdf)). The biomass system model is incorporated in the baseline scenario of the PRIMES large scale energy model for Europe (see [http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The PRIMES\\_MODEL\\_2008.pdf](http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The_PRIMES_MODEL_2008.pdf)). It is an economic supply model that computes the optimal use of resources and investment in secondary and final transformation, so as to meet a given demand of final biomass energy products, driven by the rest of sectors as in PRIMES model. The primary supply of biomass and waste has been linked with resource origin, availability and concurrent use (land, forestry, municipal or industrial waste etc). The total primary production levels for each primary commodity are restricted by the technical potential of the appropriate primary resource.

Data on potential yields and GHG emissions and removals for diverse forest management alternatives are derived from the more detailed forestry models (G4M and EFISCEN). For baseline scenario (BAU), the economic land use models project domestic production and consumption, net exports and prices of wood products and

<sup>1</sup> P. Capros, L. Mantzos, N. Tasios, A. De Vita, N. Kouvaritakis (2010), EU energy trends to 2030 — UPDATE 2009, European Commission, Directorate-General for Energy in collaboration with Climate Action DG and Mobility and Transport DG. Luxembourg: Publications Office of the European Union, 2010. ISBN 978-92-79-16191-9. Available online: [http://ec.europa.eu/energy/observatory/trends\\_2030/doc/trends\\_to\\_2030\\_update\\_2009.pdf](http://ec.europa.eu/energy/observatory/trends_2030/doc/trends_to_2030_update_2009.pdf)


changes in land use for EU member states and other world regions. The sector specific information from the economic models is used by the forest models to project GHG emissions and removals.

A more detailed description of modeling steps is provided in following sections. More detailed descriptions of each model are provided in the Annexes.

**Table 3. Essential features of the main models involved in projection of FM emissions and removals.**

<b>G4M</b>	The Global Forest Model (G4M) provides spatially explicit estimates of annual above- and belowground wood increment, development of above- and belowground forest biomass and costs of forestry options such as forest management, afforestation and deforestation by comparing the income of alternative land uses.
<b>EFISCEN</b>	The European Forest Information Scenario Model (EFISCEN) is a large-scale model that assesses the supply of wood and biomass from forests and projects forest resource development on regional to European scale, based on forest inventory data. EFISCEN provides projections on basic forest inventory data (stemwood volume, increment, age-structure), as well as carbon in forest biomass and soil.
<b>GLOBIOM</b>	GLOBIOM is a global static partial equilibrium model integrating the agricultural, livestock, bioenergy and forestry sectors with the aim to give policy advice on global issues concerning land use competition between the major land-based production sectors.

**FIGURE 1: synthetic flowchart of information exchange between models.**



The modelling approach essentially included the following steps:

1) Selection of relevant input data

- Forest area used by the models is taken from national forest inventories and scaled to match the area reported in GHG inventories (EFISCEN) or from recent literature (G4M), see Table 4.
- Main forest and forest management parameters (age structure, increment, historical harvest) are taken from national forest inventories and other country statistics (see Figure 2, and Tables 9 and 11). Other forest parameters and management characteristics taken from relevant sources (see Table 10).
- Future harvest demand under a business as usual (BAU) scenario (see Table 11) was derived from key macroeconomic drivers (GDP, population), based only on policies and measures enacted by Member States up to April 2009 (the EU 2020 renewable target and the 20% GHG reduction targets are not included in this baseline). In particular, the bio-energy demand was estimated by the Primes model and the timber demand was estimated by the Globiom model. See section 6 “Policies included” and the Annex for more information.

2) Elaboration of input data: the input data (area, age structure, increment, management characteristics, rotation length, future harvest demand,...) were elaborated by the two forest models (G4M and EFISCEN) to produce estimates of emissions and removals from FM till 2020 (for the above and below ground biomass carbon pools). The two models differ in the way they allocate harvest demand to thinnings and final fellings (including rotation lengths) with implications on emissions and removals from the forest. In general, both models follow the rules of sustainable forest management, securing sustainable yields. Further they follow different growth concepts (EFISCEN forest growth is based in inventory data, whereas G4M estimates growth from productivity maps, i.e. NPP maps) representing alternative approaches of forest growth estimation and projection. Given the unavoidable uncertainties which characterize any projections of emissions and removals from the forest sector, we think that taking the average of two different models makes the future trend illustrated below (see table 8) more robust. Elaborations also included a simulation of the impact of +/-20% harvest as compared as BAU harvest (see sensitivity analysis in table 8). See Annex I for more details on the models.

3) Ex-post processing of models' results: In order to ensure consistency between models' results and historical data reported by the country, the emissions and removals estimated by the models for the entire time series (up to 2020) were “calibrated” (i.e. adjusted) using historical data from the country for the period 2000-2008 (for which we had both data from the GHG inventories and data projected by the models). To this aim, an “offset” was calculated for two components:

- biomass: offset calculated as difference between [average of country's emissions and removals from biomass for the period 2000-2008 (table 5)] and [average of models' estimated emissions and removals from biomass for the period 2000-2008 (table 8)]
- non-biomass pools and GHG sources: offset calculated as the sum of non-biomass pools and GHG sources as reported by the country for the period 2000-2008 (table 5), and not estimated by models.

The calibrated average of models, which is used for the setting of reference level, is obtained by adding the total offset (biomass offset + non-biomass pools and GHG sources offset) to the models' average. In other words, models' results were adjusted to match the average historical data provided by each country for the period 2000-2008. This ensures consistency between country data and models' data in terms of: (i) absolute level of emissions and removals from biomass, i.e. the calibration „reconciles” differences in estimates which may be due to a large variety of factors, including different input data, different parameters, different estimation methods (e.g., some country uses a „stock-change approach”, while the models use a „gain-loss approach”); (ii) coverage of non-biomass pools and GHG sources. The calibration procedure automatically incorporates into the projections the average rate (for the period 2000-2008) of the GHG impact of past disturbances, not estimated by the model (e.g. emissions from fires,...).

The future trend of emissions and removals up to 2020 as predicted by the model is not affected by this calibration procedure, but only by the current forest characteristics (e.g., age structure,...) and the future harvest demand.

It is important to note that, to maintain consistency in the future, technical corrections (as referred in para 15 quarter and 15 quinquies of the document FCCC/KP/AWG/2010/CRP.4/Rev.4 ) will be needed in the following cases: (i) if recalculations of emissions and removals from FM (or forest land remaining forest land) for the period 2000-2008 will be carried out in any future submission of annual GHG inventories; (ii) if any future threshold selected for "force majeure" indicates that an event in the 2000-2008 period can be considered "force majeure", the impact of event (in terms of GHG) should be removed from historical FM emissions/removals (according to provisions of any future force majeure decision) , thus affecting the calibration procedure described above. For transparency reasons, the section "disturbances in the context of force majeure" reports the emissions from forest fires from 1990-2008 (expressed in Gg CO<sub>2</sub>-eq. and as % of 1990 total GHG emissions excluding LULUCF).

Spain wants to mention here that, due to time constraints, it was not possible to re-run models with the latest data available, and to correct some assumptions made by modellers. Spain

## 5 Description of construction of reference levels

### I. Description of how each of the following elements were considered or treated in the construction of the forest management reference level, taking into account the principles in decision 16/CMP.1

#### (a) Area under forest management

All the forest area in Spain is considered as managed forest. Therefore, the historical data for forest land remaining forest land reported under the Convention can be considered the historical data for area under forest management under the Kyoto Protocol.

Spain uses the "20 years transition rule" as considered good practice guidance in IPCC GPG for LULUCF (2003), this means that an afforested or reforested land is considered forest in transition, and only 20 years after the afforestation or reforestation this land becomes forest land remaining forest land, or land under forest management.

As can be appreciated, the area used by G4M is significantly bigger than the real area of forest to be considered under the Kyoto Protocol. This is due to the different definition of forest used by G4M (different FCC) and different source of data. This problem is something that Spain, together with JRC, will try to solve in the near future, and clarification and new data would be provided during the review process.

**Table 4. Area for FM as used by models (kha).**

	2000	2005	2008	2010	2015	2020	Source of historical data (up to 2008)	Projected data (2010-2020)
<b>G4M</b>	15.678	15.675	15.675	15.675	15.673	15.670	(1)	(4)
<b>EFISCEN</b>	12.581	12.579	12.577	12.576	12.573	12.570	(3)	

(1) G4M model: Gallaun, H., G. Zanchi, G. J. Nabuurs, G. Hengeveld, M. Schardt and P. J. Verkerk (2010). "EU-wide maps of growing stock and above-ground biomass in forests based on remote sensing and field measurements." Forest Ecology and Management 260(3): 252-261 (Based on CORINE and TBFRA). G4M is a spatially explicit forestry model and relies on the information from forest maps for its initialisation. This map served as a basis that was adjusted to the degree possible to data reported by countries (see points 2 and 3 below)

(3) Taken from FM area reported in latest available KP submission for the yr 2008. 2000 and 2005 were estimated based on deforestation area reported to KP.

(4) Data of 2008 minus the area of Deforestation projected by G4M.

## (b) Emissions and removals from forest management

### 1) Historical emissions and removals from forest management

The data below (table 5) are the data as estimated in the National GHG Inventory for 2011 (1990-2009). There is a big difference with the estimations for forest land remaining forest land and forest management data submitted in the 2010 GHG National inventory, where a mistake was detected only after the submission of that inventory to UNFCCC. New data are in line with previous submissions of Spanish National GHG Inventories. Due to a high level political decision in Spain, the FCC selected to define forest under the Kyoto Protocol (see Initial Report for the Establishment of the Assigned Amount) was not the same that the definition used to report land remaining forest land under the Convention. As a consequence of this decision, there was a variation in the annual increment of biomass. Trying to unify the definitions before the supplementary information for LULUCF under the Kyoto Protocol was mandatory (GHG inventory submitted in 2010), and taking into account that the forest definition for the Kyoto Protocol can not be changed once the Initial Report has been submitted, Spain modified the FCC in the Convention reporting. In the process of changing the data (areas and increments) from FCC of 10% to a FCC of 20%, a mathematical error took place that led to this big difference in figures.

**Table 5. Spain's historical emissions and removals from FM (all pools and GHGs, Gg CO2eq)**

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	av. 2000- 2008
Biomass (1)	-18665	-18664	-18662	-18660	-18658	-18656	-18654	-18652	-18650	-18648	-18646	-18644	-18643	-18641	-18639	-18637	-18635	-18633	-18631	-18639
Non-biomass pools	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
GHG sources (2)	190	296	104	85	571	159	46	139	136	85	177	72	113	156	125	260	563	38	24	170
TOTAL	-18475	-18368	-18558	-18574	-18087	-18497	-18608	-18513	-18514	-18563	-18470	-18573	-18530	-18485	-18514	-18377	-18072	-18595	-18607	-18469

(1) Above and below-ground

(2) as listed in table 2.

Note that CO2 emissions from forest fires and harvesting are already included in the annual increment of biomass (calculated as the difference in stocks between two different national forest inventories), therefore, GHG sources include only CH4 and N2O.

**2) The relationship between forest management and forest land remaining forest land as shown in GHG inventories and relevant historical data, including information provided under Article 3.3., and, if applicable, Article 3.4 forest management of the Kyoto Protocol and under forest land remaining forest land under the Convention**

All the forest area in Spain is considered as managed forest. Therefore, the historical data for forest land remaining forest land reported under the Convention can be considered the historical data for area under forest management emissions and removals under the Kyoto Protocol.

**Table 6. Spain's historical emissions and removals from FL remaining FL (Gg CO<sub>2</sub>eq), based on 2011 GHG inventory.**

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	av. 2000- 2008
Biomass (1)	-18665	-18664	-18662	-18660	-18658	-18656	-18654	-18652	-18650	-18648	-18646	-18644	-18643	-18641	-18639	-18637	-18635	-18633	-18631	-18639
Non-biomass pools	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
GHG sources (2)	190	296	104	85	571	159	46	139	136	85	177	72	113	156	125	260	563	38	24	170
TOTAL	-18475	-18368	-18558	-18574	-18087	-18497	-18608	-18513	-18514	-18563	-18470	-18573	-18530	-18485	-18514	-18377	-18072	-18595	-18607	-18469

(1) Above and below-ground

(2) as listed in table 2.

Due to the application of the „20 years transition rule” mentioned in the IPCC Good Practice Guidance for LULUCF (2003), all the afforested or reforested areas are considered forests in transition until they reach 20 years. Therefore, area of forest under forest management in Spain will remain constant, except the reduction of this area due to deforestation, until 2011, when the area considered „forest in transition” will become area under forest land remaining forest land.

**Table 7. Emissions and removals (Gg CO<sub>2</sub>eq) from AR, D and FM (if elected), based on 2011 submission of GHG National Inventory.**

	A. Article 3.3 activities			B.1 Forest management
	A.1 Aff/Reforestation		A.2. Deforestation	
	A.1.1 Lands not harvested	A.1.2 Lands harvested		
2008	-6.407,6	NA,NO	106,17	-18607
2009	-6.545,24	NA,NO	107,10	-18.562

**3) Modeled emissions and removals from forest management**

In the following table, the results of the modeling process, including calibration, is reflected.



**Table 8. Emissions and removals from FM as estimated by models (above and below-ground biomass, Gg CO<sub>2</sub>eq), calibration of models' results, and sensitivity analysis.**

		av. 2000-2008	2000	2005	2010	2015	2020	av. 2013-2020
Step 1: models' results (only biomass)	EFISCEN (1)	-20503	-20873	-20012	-21544	-21294	-21179	-21270
	G4M	-24860	-22833	-25372	-27884	-29695	-30981	-30041
	Average of models	-22682	-21853	-22692	-24714	-25494	-26080	-25655
Step 2: ex-post processing	Offset (2)	4043						
		170						
		<b>4213</b>						
	Calibrated average of models (3)	-18469	-17640	-18479	-20501	-21281	-21867	<b>-21443</b>
Sensitivity analysis (4)		+10% harvest			-17468	-18416	-19071	-18578
		-10% harvest			-23512	-23759	-24104	-23870

- (1) Efiscen does not estimate data for all countries for 2000 and 2005. When data were missing, backward extrapolation was applied as follow: sink in 2005 = sink in 2010 x ratio of harvest 2010/2005; this approach assumes that in the short term harvest is the main factor determining the sink. Estimates were extrapolated for the following countries: Bulgaria, Czech Republic, Estonia, Hungary, Italy, Latvia, Lithuania, and Netherlands.
- (2) The "offset" is distinguished between:
- biomass: calculated as difference between [average of country's emissions and removals from biomass for the period 2000-2008 (table 5)] and [average of models' estimated emissions and removals from biomass for the period 2000-2008 (table 8)]
  - non-biomass pools and GHG sources: calculated as the sum of non-biomass pools and GHG sources as reported by the country for the period 2000-2008 (table 5).
- (3) The calibrated average of models, used for the setting of reference level (see grey cell), is obtained by adding the offset to the average of models. See "ex-post processing of model's results" for details.
- (4) Preliminary simulation of the impact of +/-10% harvest as compared as BAU harvest on the emissions and removals from FM. Data are calibrated averages of models' results.

The yearly data for the calibrated average of models (assuming instantaneous oxidation for HWP) are:

**Table 8-bis. Emissions and removals from FM as estimated by models (above and below-ground biomass, Gg CO<sub>2</sub>eq), calibration of models' results, and sensitivity analysis (yearly data)**


2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
-20.097	-20.501	-20.657	-20.813	-20.969	-21.126	-21.282	-21.399	-21.516	-21.633	-21.750	-21.867

Yearly data on HWP can be found in table 15.

(c) Forest characteristics and related management

1) age class structure

Figure 2. Evolution of the forest age class structure (in yrs) as modelled by EFISCEN.



This is an approximation to the real situation. One of the main characteristics of Spanish forests is that most of them are unevenaged forests, that can't be clasified in age class structures. But for the estimation of the projected reference level we assume EFFISCEN clasification of ages of Spanish trees, although Spain will try to have more accurate information on age class structure to be considered during the review process.

2) increment

Table 9. Increments as estimated by models (m3 ha-1 yr-1)

	2000	2005	2010	2015	2020
G4M	2,9	3,2	3,4	3,5	3,7
EFISCEN	2,3	2,6	2,7	2,7	2,7

Models calculate the increments using national data. Sources for these increments can be found in table 10.

This increments are higher than the increments used for 2011 National GHG Inventory (probably due to the mistake made in calculations last year). Due to time constraints, it was not possible to re-run models with the correct data. Spain is planning to provide new increments reflecting the corrections made in 2011 GHG Inventory to be considered during the review process.

### 3) rotation length

See information in table 10.

Spain will try to provide more information on rotation lengths during the review.

### 4) information on forest management activities under “business as usual”

In the case of Spain, the National Forest Plan has been taken into account; this NFP has a validity of 30 years (from 2002 to 2032) and it is available in Spanish in [http://www.mma.es/portal/secciones/biodiversidad/montes\\_politica\\_forestal/estrategia\\_monte/pfe.htm](http://www.mma.es/portal/secciones/biodiversidad/montes_politica_forestal/estrategia_monte/pfe.htm). A description of the activities of forest management under business as usual can be found in the Spanish National Inventory Report.

### 5) other relevant information

**Table 10. Source of the main forest parameters and characteristics as used by the models.**

Model/country	Forest parameters and characteristics				
	Area (ha) by species group and age class	Growing stock (m <sup>3</sup> ) by species group and age class	Increment (m <sup>3</sup> ha <sup>-1</sup> y <sup>-1</sup> ) by species group and age class	BEF, root/shoot ratio, <b>wood density</b> by species and age-class	
				BEF and R/S ratio (dimensionless)	Wood density (t dry matter/ m <sup>3</sup> fresh volume)
<b>EFISCEN</b>	Recent inventory data were provided by national correspondents and agencies for Belgium, Czech Republic, Hungary, Ireland, Italy, Latvia. For other countries we used data collected by (13) based on data provided by national correspondents during the European Forest Sector Outlook Study in 2001		Increment functions are generally based on national forest inventory data. In case increment data was not available, yield tables have been used.	Species-specific and age-dependent BEFs have been developed for selected number of countries for EFISCEN by Vilén et al. 2005 (5) and national reports (22) and are applied to neighbouring countries	Basic wood densities are based on IPCC defaults (1)
<b>G4M</b>	Input data for all countries: for area GLC 2000 (61) and for forest area (62, scaled to JRC data to the degree possible); for the increment NPP (63, scaled to MCPFE 2005); BEF and root/shoot ratio are assumed to be constant; carbon in biomass, soil, litter and dead trees are from Kindermann et al., based on FAO and GLC 2000 (64); the age structure is desumed from NFI.				
	Management regimes have been derived from a country-wise compilation of guidelines, handbooks and personal communication (6).				

GLOBIOM	Same input data of G4M				Input data from G4M
SPAIN	23, 48, (60), 62, 63, 64	All species	3 (roots), 4 (total aboveground biomass, coarse roots)	1 (IPCC default values)	Pelkonen, P., et al. 1999. Forestry in Changing Societies in Europe- Country Reports. Part II. Silva network

- 1 IPCC, 2003. Good practice guidance for land use, land-use change and forestry. IPCC national greenhouse gas inventories programme. In: Penman, J., Gytarsky, M., Hiraishi, T., Krug, T., Kruger, D., Pipatti, R., Buendia, L., Miwa, K., Ngara, T., Tanabe, K., Wagner, F. (Eds.). Institute for Global Environmental strategies for the IPCC, Hayama, Kanagawa.
- 3 Mokany K, Raison RJ, Prokushkin AS (2006) Critical analysis of root:shoot ratios in terrestrial biomes. *Global Change Biology*, 12, 84–96.
- 4 Sabate S, Gracia CA, Sanchez A (2002) Likely effects of climate change on growth of *Quercus ilex*, *Pinus halepensis*, *Pinus pinaster*, *Pinus sylvestris* and *Fagus sylvatica* forests in the Mediterranean region. *Forest Ecology and Management*, 162, 23–37.
- 5 Vilén, T., Meyer, J., Thürig, E., Lindner, M., Green, T., 2005. Improved regional and national level estimates of the carbon stock and stock change of tree biomass for six European countries, (Deliverable 6.1). Improved national estimates of the carbon stock and stock change of the forest soils for six European countries (Deliverable 6.2). CarboInvent Project: [http://www.joanneum.at/carboinvent/D\\_6\\_1\\_6\\_2.pdf](http://www.joanneum.at/carboinvent/D_6_1_6_2.pdf) European Forest Institute, Joensuu, p. 31.
- 6 Nabuurs, G., Pussinen, A., van Brusselen, J., Schelhaas, M., 2007. Future harvesting pressure on European forests. *European Journal of Forest Research* 126, 391-400.
- 13 Schelhaas, M.J., Brusselen, J.V., Pussinen, A., Pesonen, E. Schuck, A., Nabuurs, G.J., Sasse, V. (2006). Outlook for the development of European forest resources. UN-ECE
- 22 Romano D., Arcarese, C., Bernetti, A. Caputo, A., Córdor, R.D., Contaldi, M., De Lauretis, R., Di Cristofaro, E., Federici, S., Gagna, A., Gonella, B., Liburdi, R., Taurino, E., Vitullo, M., 2009. Italian Greenhouse Gas Inventory 1990-2007. National Inventory Report 2009. ISPRA - Institute for Environmental Protection and Research, Rome.
- 23 Schelhaas, M.J., Varis, S., Schuck, A. and Nabuurs, G.J. 1999. EFISCEN's European Forest Resource Database. European Forest Institute, Joensuu, Finland.
- 48 Segundo inventario forestal nacional (IFN2) 1986-1996. No reference
- 60 Global Forest Model (the output may be used as input data for area for EU FASOM and GLOBIOM)
- 61 Gallaun, H., G. Zanchi, G. J. Nabuurs, G. Hengeveld, M. Schardt and P. J. Verkerk (2010). "EU-wide maps of growing stock and above-ground biomass in forests based on remote sensing and field measurements." *Forest Ecology and Management* 260(3): 252-261. (Based on CORINE and TBFRA)
- 62 Global Land Cover 2000
- 63 Cramer . W, D. W. Kicklighter, A, Bondeau, B., Moore, III, G. Churkina, B., Nermy, A. Ruimy, A., L., Schloss, and the Participants of the Potsdam NPP Model Intercomparison (1999). Comparing global models of terrestrial net primary productivity (NPP) : overview and key results. *Global Change Biology*, Volume 5 Issue 51, pp 1-15
- 64 Kindermann G., McCallum I., Fritz S., Obersteiner M., 2008. A global forest growing stock, biomass and carbon map based on FAO statistics. *Silva Fennica*, Vol 42(3), pp 387-396.

#### **(d) Harvesting rates**

##### **1) Historical harvesting rates, and assumed future harvesting rates**

Data for Spanish historical harvest rates are extracted from FAO database for harvested wood products.

**Table 11. Historical harvest rate and projected BAU harvest demand used by models (roundwood overbark 1000 m3)**

2000	2005	2010	2015	2020	ratio (av. 2013-2020)/2005	Source of historical data (until 2007)
17.023	17.755	18.246	18.738	19.229	1,06	FAO June 2010

Notes: values in the table express 5-yrs average (e.g. 2000 is the average 1998-2002, 2005 is the average 2003-2007). Until 2007, data are from national statistics or other country data. Data for 2020 were estimated by the models Primes (wood for bioenergy) and Globiom (timber). Data between 2008 and 2020 are interpolated. The harvest rate used by each model may slightly deviate from harvest demand (e.g. if the model did not “find” all the wood in the forests).

A general assumption has been done that all the harvest predicted till 2020 is allocated to FM, i.e. it was assumed that the harvest up to 2020 on areas afforested/reforested or deforested after 1990 is negligible as compared to the harvest of forest areas which qualify as FM.

#### **(e) Harvested wood products**

The contribution of HWP to the reference level of Spain would amount to -2,447 Mt CO<sub>2</sub>.

It was calculated using the C-HWP-Model, which estimates delayed emissions on the basis of the annual stock change of semi-finished wood products as outlined in the 2006 GL (Rüter, 2011). The estimation uses the product categories, half lives and methodologies as suggested in para 27, page 31 of FCCC/KP/AWG/2010/CRP.4/Rev.4.

The activity data (production and trade of sawnwood, wood based panels and paper and paperboard) is derived from the TIMBER database (UNECE 2011) (time series 1964-2009).

In order to achieve accurate results, the HWP numbers have been calculated applying the sub-categories of sawnwood, wood based panels and paper and paperboard as specified in Table 1. Sawnwood includes the Items 1632 and 1633, wood based panels comprising of Items 1634, 1640, 1646, 1647, 1648, 1649 and 1650, and paper and paperboard corresponds to Item 1876.

Following conversion factors have been used:

**Table 12: Conversion factors of considered commodities\***

Classification		Description of commodity	Air dry density	C conv. factor	Source
FAO	UNECE		[g/cm³]	[Gg C/1000m³]	
1866	1.2.C	Industrial roundwood, coniferous	0,450	2,250E-01	Kollmann (1982), (oak, beech)
1867	1.2.NC	Industrial roundwood, non-coniferous	0,670	3,350E-01	Kollmann (1982), (oak, beech)
1632	5.C	Sawnwood, coniferous	0,450	2,250E-01	Kollmann (1982), (oak, beech)
1633	5.NC	Sawnwood, non-coniferous	0,670	3,350E-01	Kollmann (1982), (oak, beech)
1634	6.1	Veneer sheets	0,590	2,950E-01	IPCC (2003)
1640	6.2	Plywood	0,480	2,402E-01	IPCC (2003)
1646	6.3	Particle board	0,630	2,898E-01	Hasch (2002), Barbu (2011)
1647	6.4.1	Hardboard	0,850	4,165E-01	Kollmann (1982), Barbu (2011)
1648	6.4.2	Medium density fibreboard	0,725	3,190E-01	Hasch (2002), Barbu (2011)
1649	6.4.x	Fibreboard, compressed	0,788	3,504E-01	(50 % hardboard / 50 % medium density fibreboard)
1650	6.4.3	Other board (Insulating board)	0,270	1,148E-01	Kollmann (1982), Barbu (2011)
1876	10	Paper and paperboard	0,900**	4,500E-01**	IPCC (2006)

\* Items 1866 and 1867 are needed for methodological reasons only (see following section), \*\* in [g/g] and [Gg C/1000t]

In order to only estimate emissions from HWP removed from forests which are accounted for by Spain under Article 3, in a first step, the annual share of carbon in HWP coming from domestic forests has been calculated.

Following equations were used as industrial roundwood is assumed to serve as raw material for the production of HWP.

$$(1) \quad \text{ratio}_{\text{INDRW consumption from dom harvest}} = \frac{(\text{Production}_{\text{INDRW}} - \text{Export}_{\text{INDRW}})}{(\text{Production}_{\text{INDRW}} + \text{Import}_{\text{INDRW}} - \text{Export}_{\text{INDRW}})}$$

$$(2) \quad \text{Production}_{\text{HWP from dom harvest}} = \text{Production}_{\text{HWP}} \cdot \text{ratio}_{\text{INDRW consumption from domestic harvest}}$$

The ratio (Equation 1) was calculated both for coniferous and non-coniferous industrial roundwood (*INDRW*, Items 1866 and 1867). For coniferous sawnwood and paper and paperboard, the ratio for coniferous industrial roundwood was applied. For non-coniferous sawnwood the ratio for non-coniferous industrial roundwood was applied. For the other HWP, the ratio of the annual mass weighted average of coniferous and non-coniferous industrial roundwood was applied.

As a result, this share of HWP produced from domestically harvested timber is presented as a percentage in Table 2.

The presented approach follows the initial assumption that all forests in Spain are managed, and in order to simplify matters, it is presumed that all harvest is allocated to forest management. This assumption is to be verified and corrected where necessary. The final allocation of carbon in HWP to forests which are accounted for under Article 3 shall be part of a technical correction as suggested in para 15 quater, page 27 of FCCC/KP/AWG/2010/CRP.4/Rev.4.

**Table 13: Historic time series of amounts and share of accountable carbon Inflow to the HWP pool [in 1000t C and %]**

1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980
613	836	889	1046	1137	1157	1302	1499	1740	1841	1869	1798	2021	2080	2138	2370	2216
97,9%	96,2%	95,7%	96,5%	96,3%	95,3%	97,1%	98,5%	98,9%	96,4%	93,7%	99,0%	98,9%	99,0%	96,9%	93,1%	94,0%

1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
2407	2516	2404	2484	2539	2744	2787	2782	2988	3172	3211	3000	2985	3176	3152	3222	3179
99,4%	99,4%	99,4%	99,4%	99,3%	99,3%	99,3%	98,9%	98,9%	99,0%	98,8%	99,0%	95,9%	96,6%	95,5%	97,8%	96,8%

1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
3079	3607	3918	4167	4224	4282	4346	4341	4754	4464	4198	4165
90,3%	91,7%	87,5%	86,6%	88,5%	89,7%	88,9%	87,5%	85,4%	82,3%	88,2%	87,0%

The annual carbon Inflow (= carbon in produced HWP) to the HWP pool prior to the year 1964 (first year for which activity data from TIMBER database (UNECE 2011) is available for Spain) has been calculated from the 5 years average from 1964 to 1968 and was assumed to be the constant carbon pool Inflow for the time period 1900-1963.

In order to provide a projection for the development of the HWP pool consistent with the assumptions on the future harvest, the rates of change of the Projected harvest (Model GLOBIOM) as compared to the last 5 years average of historic harvest, for which up-to-date data is available, was calculated (cf Table 3).

These projected growth rates as cp. to the average of the years 2003-2007 for Spain were applied to the same 5 years average of historic carbon Inflow to the HWP pool in order to receive the future Inflow to the HWP pool.

**Table 14: Projection of carbon Inflow to the HWP pool**

Average of historic harvest (2003-2007) [in 1000m3]	17.755										
Average HWP pool Inflow* (2003-2007) [in 1000t C]	4437										

years	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Projected harvest rate [in 1000m3]	18246,3	18344,6	18442,8	18541,1	18639,4	18737,7	18836	18934	19032	19131	19229
Change as cp to historic harvest (2003-2007) [in %]	2,77%	3,32%	3,87%	4,43%	4,98%	5,53%	6,09%	6,64%	7,20%	7,75%	8,30%
Projected carbon Inflow to HWP pool [in 1000t C]	4560,21	4584,77	4609,33	4633,89	4658,46	4683,02	4707,58	4732,14	4756,7	4781,26	4805,82

\*a similar approach was chosen by Kangas and Baudin (2003): ECE/TIM/DP/30

For calculating the pool of HWP in use, three half-lives for application in the first order decay function have been used as suggested by para 7, page 31 of FCCC/KP/AWG/2010/CRP.4/Rev.4.

- Sawnwood: 35 years

- Wood based panels: 25 years
- Paper and paperboard: 2 years

The projected net-emissions are calculated from the annual stock change estimates following the calculation method provided in IPCC 2006, Vol.4, Ch. 12 (Equation 12.1).

**Table 15: Historic (up to 2009) and projected net-emissions from HWP pool [in 1000t CO<sub>2</sub>]**

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
-3571	-3513	-2586	-2488	-3132	-2905	-2974	-2667	-2303	-4088	-4882	-5429	-5202	-4974	-4829	-4496	-5528

2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
-4017	-2720	-4599	-3402	-3097	-2877	-2715	-2596	-2507	-2439	-2386	-2344	-2310	-2281

#### (f) Disturbances in the context of force majeure

The calibration procedure described above automatically incorporates the average rate of past disturbances (for the period 2000-2008) into the projections. See further comments in section „Ex-post processing of models’ results” on the need of future consistency. For transparency reasons, the tables below report the emissions from forest fires from 1990-2008 (expressed in Gg CO<sub>2</sub>-eq. and as % of 1990 total GHG emissions excluding LULUCF). Depending on the definition adopted for force majeure, some or none of the events that provoked the emissions shown below should be excluded in order to maintain consistency with the accounting of the commitment period, in the case that the provision of force majeure is available and Spain decides voluntarily to use it.

**Table 16. Emissions from forest fires (Gg CO<sub>2</sub>eq and % of 1990 total GHG without LULUCF)**

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	av. 2000-2008
GgCO <sub>2</sub> eq (1)	2077	3226	1134	930	6228	1729	504	1519	1487	926	1928	781	1231	1699	1366	2840	6140	417	260	1851
% 1990 GHG	0,7	1,1	0,4	0,3	2,2	0,6	0,2	0,5	0,5	0,3	0,7	0,3	0,4	0,6	0,5	1,0	2,2	0,1	0,1	0,6

(1) Data are taken from the last available CRF tables, rows "wildfires" of Tables LULUCF 5(V) (Biomass burning). Some countries reported CH<sub>4</sub> and N<sub>2</sub>O in this table but did not explicitly included CO<sub>2</sub> emissions (i.e. CO<sub>2</sub> is implicitly included in tab 5A); in these cases, the JRC indirectly derived CO<sub>2</sub> emissions from CH<sub>4</sub> and N<sub>2</sub>O reported emissions, using default factors from IPCC Good Practice Guidance for LULUCF (IPCC GPG 2003, table 3A.1.16). In the case of Spain, data for CO<sub>2</sub> emissions associated to forest fires have been provided by the country.

#### (g) Factoring out in accordance with paragraph 1(h) (i) and 1(h) (ii) of decision 16/CMP.1

Not relevant.



**II. Description of any other relevant elements considered or treated in the construction of the forest management reference level, including any additional information related to footnote 1 in paragraph 4 of decision [-/CMP.6]**

**6 Policies included**

**I. Pre-2010 domestic policies included**

Policy assumptions are made in the baseline scenario of the PRIMES model which underpins the projections for the construction of the Reference Level. For the purpose of this submission, policies and measures included are those implemented by April 2009 and legislative provisions adopted by April 2009 that are defined in such a way that there is almost no uncertainty how they should be implemented in the future. An inventory of legal measures and EU financial support included in the PRIMES model is reproduced from Capros et al. (2010) in Annex II to this submission. However more details are provided on pp.17-21 ("BASELINE") of the publication *EU energy trends to 2030 - UPDATE 2009*.<sup>2</sup>

As mentioned before, the policy framework for forest action in Spain is the National Forest Plan, with validity from 2002 to 2032. The first review of NFP will take place in 2012.

More detail of policies and measures included in the reference level can be found in Annex II.

**II. Confirmation of factoring out policies after 2009**

No national forest related policies have been adopted since the National Forest Plan was endorsed by the Council of Ministries of Spain in July 2002. EU policies have been considered by modelers only up to April 2009.

---

<sup>2</sup> P. Capros, L. Mantzos, N. Tasios, A. De Vita, N. Kouvaritakis (2010), *EU energy trends to 2030 — UPDATE 2009*, European Commission, Directorate-General for Energy in collaboration with Climate Action DG and Mobility and Transport DG. Luxembourg: Publications Office of the European Union, 2010. ISBN 978-92-79-16191-9. Available online: [http://ec.europa.eu/energy/observatory/trends\\_2030/doc/trends\\_to\\_2030\\_update\\_2009.pdf](http://ec.europa.eu/energy/observatory/trends_2030/doc/trends_to_2030_update_2009.pdf)

## **ANNEX I – Description of models**

### **GLOBIOM**

GLOBIOM is a global static partial equilibrium model integrating the agricultural, bioenergy and forestry sectors with the aim to give policy advice on global issues concerning land use competition between the major land-based production sectors. Concept and structure of GLOBIOM are similar to the US Agricultural Sector and Mitigation of Greenhouse Gas (ASMGHG) model (Schneider, McCarl and Schmid 2007). The global agricultural and forest market equilibrium is computed by choosing land use and processing activities to maximize the sum of producer and consumer surplus subject to resource, technological, and political restrictions, as described by McCarl and Spreen (1980). Prices and international trade flows are endogenously computed for 28 world regions.

The market is represented through implicit product supply functions based on detailed, geographically explicit, Leontief production functions, explicit resource supply functions (land and water), and product demand functions.

Land and its characteristics are the key elements of our modeling approach. In order to enable global bio-physical process modeling of agricultural and forest production, a comprehensive database has been built (Skalsky et al., 2008), which contains geo-spatial data on soil, climate/weather, topography, land cover/use, and crop management (e.g. fertilization, irrigation). The data are available from various research institutes (NASA, JRC, FAO, USDA, IFRPI, etc.) and significantly vary with respect to spatial, temporal, and attribute resolutions, thematic relevance, accuracy, and reliability. Therefore, data were harmonized into several common spatial resolution layers including 5 and 30 arcmin as well as country layers. Consequently, Homogeneous Response Units (HRU) have been delineated by including only those parameters of landscape, which are almost constant over time. At the global scale, we have included five altitude classes, seven slope classes, and six soil classes. In a second step, the HRU layer is merged with other relevant information such as global climate map, land category/use map, irrigation map, etc. to delineate Simulation Units, which are actually input into the Environmental Policy Integrated Climate model (EPIC, Williams 1995, Izaurralde et al. 2006). This HRU concept assures consistent aggregation of geo-spatially explicit bio-physical impacts that are simulated with EPIC (e.g. crop yields, nitrogen leaching, soil carbon sequestration).

Currently, two major land cover types are represented in the model: cropland and forest. Crop production accounts for about 20 globally most important crops. The data are taken from FAOSTAT, where national averages over the years 2001-2005 are used to define base levels for yields, harvested areas, prices, production, consumption, trade, and supply utilization. Irrigated crop yields, crop specific irrigation water requirements, and costs for five irrigation systems are derived from a variety of sources as described in Sauer et al. (2008). For selected crops (corn, sugarcane and wheat), management and land quality specific yields have been estimated with EPIC. Four management systems are currently represented which correspond to the IFRPI crop distribution data classification (irrigated, high input - rainfed, low input - rainfed and subsistence management systems). The number of crops, systems, and parameters (especially environmental parameters like soil carbon, erosion, and nutrient leakage) estimated with EPIC is being expanded.

Crop supply can enter one of three processing/demand channels: consumption, livestock production or biofuel production. Consumption is modeled by constant elasticity demand functions parameterized using FAOSTAT data. Only a preliminary regional livestock production representation is applied in the present version of the model where a bundle of livestock products is assimilated to a generic commodity - “animal calories”. Feed requirements have been calculated from the Supply Utilisation Accounts, FAOSTAT. Demand for livestock products is represented through upward sloping demand curves. Biofuel options from crops include first generation technologies for a) ethanol from sugarcane or corn, and b) biodiesel from soya or rapeseed. The processing data are based on Hermann and Patel (2007) for ethanol and Haas et al. (2006) for biodiesel. Market demand for ethanol and biodiesel is represented through vertical demand functions.

Primary forest production is characterized also on the basis of HRUs and the resulting Simulation Units. The most important parameters for the model are mean annual increment, maximum share of sawlogs in the mean annual increment, and harvesting cost. These parameters are shared with the G4M Model – a successor of the

model described by Kindermann et al (2006). More specifically, mean annual increment for the management, is obtained by downscaling the biomass stock data from the Global Forest Resources Assessment (FAO, 2005) from the country level to the grid using the method described in Kindermann et al. (2008). This downscaled biomass stock data is subsequently used to parameterize the increment curves Kindermann (2008). Finally, sawnwood share is estimated by the tree size which in turn depends on yield and rotation time. Harvesting costs is adjusted for slope and tree size as well.

Five primary forest products are defined: sawlogs, pulplogs, other industrial logs, firewood, and energy biomass. Sawlogs, pulplogs and energy biomass are further processed. Sawnwood and woodpulp production, and demand parameters rely on the 4DSM model described in Rametsteiner et al. (2007). FAO data and other secondary sources have been used for quantities and prices of sawnwood and woodpulp. For production cost estimates of these products, for example, mill costs, an internal IIASA database and purchased data were used. The energy biomass can be converted into methanol and heat or electricity and heat, where processing costs and conversion coefficients are obtained from Leduc et al. (2008), Hamelinck and Faaij (2001), Sørensen (2005), and Biomass Technology Group (2005). Demand for woody bioenergy production is implemented through minimum quantity restrictions, similarly as demand for other industrial logs and for firewood.

The final model calibration, supposed to correct data imperfections and get the baseline solution close to the observed values, is done by adjusting the cost parameters of selected activities so that for the baseline activity levels, their marginal cost equals to their marginal revenue, as assumed by the microeconomic theory. The controlled activities are crop areas, primary forest products supply and animal calories supply.

#### Input

- Baseline prices and quantities of considered products
- Supply and demand elasticities
- Ressource requirements (land, water,...)
- Production cost
- Transformation cost
- Transport cost
- Conversion coefficients from primary to final products
- Initial land use

#### Output

- supply and demand quantities
- equilibrium prices
- volumes traded between the regions
- land use change
- water consumption

## References

- Biomass Technology Group (2005). Handbook Biomass Gasification. H.A.M. Knoef. ISBN: 90-810068-1-9.
- FAO. 2005. "Global Forest Resources Assessment 2005, Progress towards sustainable forest management." Vol. 147 of FAO Forestry Paper. FAO, Rome.
- FAO. 2007. FAOSTAT. Available online at [faostat.fao.org](http://faostat.fao.org).
- Haas, M.J., McAloon, A.J., Yee, W.C. and T.A. Foglia. 2006. "A process model to estimate biodiesel production costs." *Bioresource Technology* 97: 671-678.
- Hamelinck, C.N. and A.P.C. Faaij (2001). "Future Prospects for Production of Methanol and Hydrogen from Biomass." Utrecht University, Copernicus Institute, Science Technology and Society, Utrecht, Netherlands 2001.
- Hermann, B.G. and M. Patel. 2007. "Today's and tomorrow's bio-based bulk chemicals from white biotechnology - A techno-economic analysis." *Applied Biochemistry and Biotechnology* 136: 361-388.
- Izaurrealde, R.C., J.R. Williams, W.B. McGill, N.J. Rosenberg, and M.C. Quiroga Jakas. 2006. Simulating soil C dynamics with EPIC: model description and testing against long-term data. *Ecological Modelling*, 192:362-384.
- Kindermann, G.E. 2008. "Getting forest growth functions out of yield tables."
- Kindermann, G.E., McCallum, I. and S. Fritz. 2008. "A global forest growing stock, biomass and carbon map based on FAO statistics." Forthcoming.
- Kindermann, G.E., Obersteiner, M., Rametsteiner, E. and I. McCallum. 2006. "Predicting the deforestation-trend under different carbon-prices." *Carbon Balance and Management* 1: 15.
- Leduc, S., D. Schwab, E. Dotzauer, E. Schmid, M. Obersteiner (2008). "Optimal Location of Wood Gasification Plants for Methanol Production with Heat Recovery." *International Journal of Energy Research*, IGEC-III special issue.
- McCarl, B.A. and T.H. Spreen. 1980. "Price Endogenous Mathematical Programming as a Tool for Sector Analysis." *American Journal of Agricultural Economics* 62: 87-102.
- Rametsteiner E, Nilsson S, Boettcher H, Havlik P, Kraxner F, Leduc S, Obersteiner M, Rydzak F, Schneider U, Schwab D, Willmore L. 2007. Study of the Effects of Globalization on the Economic Viability of EU Forestry. Final Report of the AGRI Tender Project: AGRI-G4-2006-06 [2007].
- Sauer, T., Havlík, P., Kindermann, G., and Schneider, U.A. 2008. Agriculture, Population, Land and Water Scarcity in a changing World - the Role of Irrigation. Paper prepared for the 2008 Congress of the European Association of Agricultural Economists in Gent, Belgium.
- Schneider, U.A., B.A. McCarl, and E. Schmid. 2007. "Agricultural sector analysis on greenhouse gas mitigation in US agriculture and forestry." *Agricultural Systems*. 94:128-140.
- Skalsky, R., Z. Tarasovic(ová, J. Balkovic(, E. Schmid, M. Fuchs, E. Moltchanova, G. Kindermann, and P. Scholtz. 2008. GEO-BENE global database for bio-physical modeling v. 1.0 – concepts, methodologies and data. The GEO-BENE database report. International Institute for Applied Systems Analysis (IIASA), Austria, pp. 58.
- Sørensen, Å.L (2005) "Economies of Scale in Biomass Gasification Systems." IIASA, Interim Report 2005, IR-05-030.

Williams, J.R. 1995. The EPIC Model. In Computer Models of Watershed Hydrology (Ed.: V.P. Singh). Water Resources Publications, Highlands Ranch, Colorado, 1995, pp 909-1000.

### **EFISCEN**

The European Forest Information Scenario (EFISCEN) model (Sallnäs 1990; Schelhaas et al. 2007) is a large-scale model that assesses the supply of wood and biomass from forests and projects forest resource development on regional to European scale (Eggers et al. 2008; Ľupek et al. 2010).. The core of the model was developed in the late 1980s, as a forest resource projection model for Sweden.

EFISCEN uses forest inventory data as an input, including:

- area (ha);
- average standing volume of growing stock (m<sup>3</sup>/ha);
- net annual increment (m<sup>3</sup>/ha/y).

Based on this data, the state of the forest is described as an area distribution over age- and volume-classes in matrices. During simulations, forest area moves between matrix cells, describing different natural processes (e.g. growth and mortality) and human actions (e.g. forest management). Growth dynamics are simulated by shifting area proportions between matrix cells. In each 5-year time step, the area in each matrix cell moves up one age-class to simulate ageing. Part of the area of a cell also moves to a higher volume-class, thereby simulating volume increment. Growth dynamics are estimated by the model's growth functions whose coefficients are based on inventory data.

Management scenarios are specified at two levels in the model. First, a basic management regime defines the period during which thinnings can take place and a minimum age for final fellings. These regimes can be regarded as constraints on the total harvest level. Thinnings are implemented by moving area to a lower volume class and final fellings by moving area outside the matrix to a bare-forest-land class, from where it can re-enter the matrix. The applied management regimes are based on a country level compilation of management guidelines (Nabuurs et al. 2007). Second, the demand for wood is specified for thinnings and for final felling separately and EFISCEN may fell the demanded wood volume if available. If wood demand is high, management is intensive and rotation lengths are close to the lower limit defined in the management regimes. If wood demand is low, rotation lengths are longer, because less fellings are needed to fulfill the demand.

EFISCEN projects (i) stemwood volume, (ii) increment, (iii) age-classes and (iv) wood removals for five year time-steps. To assess biomass carbon stocks, stemwood volume is converted into carbon in stems, branches, foliage, coarse and fine roots, using basic wood densities, a generic carbon content, and age-dependent biomass distribution factors. Felling residues and litter production of trees, due to turnover and natural mortality, are used as input data for the dynamic soil model YASSO (Liski et al., 2005) and incorporated as independent module.

The soil model **YASSO** is used to estimate changes in the soil C pool by EFISCEN model. YASSO consists of three litter compartments and five decomposition compartments. For the soil carbon module, the litter is grouped as non-woody litter (foliage and fine roots), fine woody litter (branches and coarse roots) and coarse woody litter (stems and stumps). Each of the litter compartments has a fractionation rate determining the proportion of its contents released to the decomposition compartments in a time step. For the compartment of non-woody litter, this rate is equal to 1 which means that all of its contents is released in one time step, whereas for the woody litter compartments this rate is smaller than 1. Litter is distributed over the decomposition compartments of extractives, celluloses and lignin-like

compounds according to its chemical composition. Each decomposition compartment has a specific decomposition rate, determining the proportional loss of its contents in a time step. Fractions of the losses from the decomposition compartments are transferred into the subsequent decomposition compartments having slower decomposition rates while the rest is removed from the system. The fractionation rates of woody litter and the decomposition rates are controlled by temperature and water availability and are based on litterbag data across Europe (Liski et al., 2003).

The model is especially suited for simulating managed, even-aged forests at large scales. The model has been validated for Finland (Nabuurs et al. 2001) and Switzerland (Thürig and Schelhaas 2006) by running EFISCEN on historic data. Other validations have been performed by comparing its growth functions against growth functions of other models and by comparing projections against projections of other models (e.g. Ľupek et al. 2010).

#### Some reference

-[http://www.efi.int/portal/virtual\\_library/databases/efiscen/](http://www.efi.int/portal/virtual_library/databases/efiscen/)

- <http://www.environment.fi/default.asp?contentid=250208&lan=EN>

- Eggers J, Lindner M., Zudin S, Zaehle S, Liski J. (2008). Impact of changing wood demand, climate and land use on European forest resources and carbon stocks during the 21st century. *Global Change Biology*, 14: 2288-2303.

Liski, J., Nissinen, A., Erhard, M., Taskinen, O., 2003. Climatic effects on litter decomposition from arctic tundra to tropical rainforest. *Global Change Biology* 9, 575-584.

- Liski J., Palosuo T., Peltoniemi M., Sievänen R. (2005). Carbon and decomposition model Yasso for forest soils. *Ecological Modelling*, 189 (1-2): 168-182

-Nabuurs, G.J., Schelhaas, M.J., Pussinen, A., 2000. Validation of the European Forest Information Scenario Model (EFISCEN) and a projection of Finnish forests. *Silva Fennica* 34, 167-179.

- Nabuurs G.J., Pussinen A., van Brusselen J., Schelhaas M. J. (2007). Future harvesting pressure on European forests. *Eur. J Forest Res*, 126: 391-400.

-Sallnäs, O., 1990. A matrix model of the Swedish forest. *Studia Forestalia Suecica* 183, 23.

-Schelhaas, M.J., Eggers, J., Lindner, M., Nabuurs, G.J., Päivinen, R., Schuck, A., Verkerk, P.J., Werf, D.C.v.d., Zudin, S., 2007. Model documentation for the European Forest Information Scenario model (EFISCEN 3.1.3). Alterra-rapport 1559/EFI Technical Report 26, Alterra, Wageningen, 118 pp.

- Thürig E., Schelhaas M. J. (2006). Evaluation of a large scale forest scenario model in heterogeneous forests: a case study for Switzerland. *Canadian Journal of Forest Research*, 36 (3): 671-683.

-Ľupek, B., Zanchi, G., Verkerk, P.J., Churkina, G., Viovy, N., Hughes, J.K., Lindner, M., 2010. A comparison of alternative modelling approaches to evaluate the European forest carbon fluxes. *Forest Ecology and Management* 260, 241-251.

#### **GLOBAL FORESTRY MODEL -G4M**

##### *General description*

The Global Forest Model (G4M) is applied and developed by IIASA and estimates the annual above ground wood increment and harvesting costs. By comparing the income of managed forest (difference of wood price and harvesting costs, income by storing carbon in forests) with income by alternative land use on the same place,

the decision of afforestation or deforestation is made. As G4M is spatially explicit (currently on a 30"x30" resolution) the different deforestation pressure at the forest frontier can also be handled. The model can use external information (like wood prices, prescribed land-use change) from other models or data bases, which guarantee food security and land for urban development or account for disturbances. As outputs, G4M produces estimates of land-use change, carbon sequestration/emissions in forests, impacts of carbon incentives (e.g., avoided deforestation), and supply of biomass for bio-energy and timber.

The model handles age classes with one year width. Afforestation and disasters cause an uneven age-class distribution over a forest landscape. The model performs final cuts in a manner, that all age classes have the same area after one rotation period. During this age class harmonization time the standing biomass, increment and amount of harvest is fluctuating due to changes in age-class distribution and afterwards stabilizing.

The main forest management options considered by G4M are variation of thinning and choice of rotation length. G4M does not model species explicitly but a change of species can be emulated by adapting NPP, wood price and harvesting costs. The rotation length can be individually chosen but the model can estimate optimal rotation lengths to maximize increment, maximize stocking biomass or maximize harvestable biomass.

### *Adjustments and harmonisation*

An EU-wide forest/ non-forest map was generated, consistent with the Temperate and Boreal Forest Resource Assessment –TBFRA 2000 (UNECE-FAO, 2000) at the national level. For areas where CORINE land cover data are available, the CORINE dataset was aggregated from the original 100 meters to 500 meters spatial resolution. Firstly, the number of forest pixels within each 5 by 5 pixel aggregation unit was calculated. Secondly, a threshold with the minimum number of forested pixels within the aggregation units was determined for each country. This threshold was selected accordingly, to generate a forest map in agreement with the total forest area given by TBFRA 2000 at the national level. For areas not covered by CORINE data, a similar approach was applied with Vegetation Continuous Fields (VCF) data (Hansen et al. 2003). The area covered with woody vegetation in the VCF data is given in percent. A percentage threshold of the minimum area covered by woody vegetation was defined for each country to match total forest area from TBFRA 2000. Based on FAO data the map distinguishes between managed and unmanaged forest. Criteria of wilderness and remoteness were used to locate the unmanaged forest areas on the map. The initial growing stock per grid cell was taken from the European forest biomass map from Gallaun et al. (in press). For countries outside Europe the forest biomass map compiled by Kindermann et al. (2008) was used.

Increment is determined by a potential NPP map (Cramer et al. 1999) and translated into mean annual increment (MAI). At present this increment map is static but can be changed to a dynamic growth model which reacts to changes of temperature, precipitation or CO<sub>2</sub> concentration. For the purpose of this study the increment map was scaled at country level to match either MCPFE or reported country data. Age structure and stocking degree are used as additional information for adjusting MAI. If stocking degree of forest modelled with a given age structure (country average) in a cell is greater than 1.05 age structure of the modelled forest is shifted iteratively by a few age classes towards older forest. If stocking degree of forest modelled in a cell is smaller than 0.5 age structure of the modelled forest is shifted iteratively by a few age classes towards younger forest. It is required that the shifts are symmetrical to keep country average age structure close to statistical value. If the age structure shift distribution within a country is skewed towards older forest, the country's average MAI is increased iteratively. If the age structure shift distribution within a country is skewed towards younger forest country MAI is decreased iteratively.

The model uses external projections of wood demand per country to calculate total harvest iteratively. The potential harvest amount per country under a scenario of rotation lengths that maintain current biomass stocks is estimated. If total harvest is smaller than wood demand the model changes grid per grid (starting from the most productive forest) management to a rotation length that optimizes forest increment and thus allows for more harvest. This mimics the typical observation that managed forests in Europe are currently not managed optimally with respect to yield. The rotation length is changed at maximum by 5 years per time step. If harvest still too small and unmanaged forest is available the status of the unmanaged forest will change to managed. If total harvest greater than demand the model changes management to maximum biomass rotation length, i.e. manages forests for carbon sequestration. If wood demand is still lower than potential harvest managed forest

can be transferred into unmanaged forest. Thinning is applied to all managed forests. The stands are thinned to maintain a stocking degree specified (between 0.5 and 1.05), i.e. thinning mimics natural mortality along the self-thinning line. The model can consider the use of harvest residues e.g. for bioenergy purposes.

Despite the harmonization efforts to reproduce observed data on increment, area and harvest, the forest carbon balance as described in the model might still deviate from the observed forest carbon sink or source. This might be due to differences in forest management or forest disturbances. The model cannot account for such effects. To compensate for processes affecting the carbon balance that cannot be modelled, an adjustment algorithm has been introduced. Rotation length of unmanaged forest is set to the value that yields constant biomass (equal to observed biomass in 2000). If modelled carbon sink/source from forest management (averaged over 1990-1995) is smaller/larger than reported by a country, the rotation length of unmanaged forest is changed to maximizing biomass. The procedure is applied cell by cell within the country's unmanaged forest until the reported stock change is matched.

### **Some references**

- Böttcher H., Aoki K., De Cara S., Gusti M., Havlik P., Kindermann G., Schneider U., Obersteiner M. (2008). GAINS GHG mitigation potentials costs from land-use, land-use change and forestry (LULUCF) in Annex 1 countries. Methodology. International Institute for Applied Systems Analysis, Laxenburg, Austria, 39 pp.
- Gusti M., Havlik P., Obersteiner M. (2008). Technical description of the IIASA model cluster. IIASA. 12 p.
- Kindermann G., McCallum I., Fritz S., Obersteiner M. (2008). A global forest growing stock, biomass and carbon map based on FAO statistics. *Silva Fennica*. Vol.42(3), pp.387-396.
- Kindermann G., Obersteiner M., Rametsteiner E. and McCallum I. (2006). Predicting the Deforestation-Trend under Different Carbon-Prices. *Carbon Balance and Management*, 1:15; doi:10.1186/1750-0680-1-15.

### **The PRIMES Energy Systems Model**

#### *General Description*

A summary description of the energy systems model for is provided on [http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The PRIMES MODEL 2008.pdf](http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The_PRIMES_MODEL_2008.pdf) and of the biomass system model, which is incorporated in the large scale model, on [http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The PRIMES MODEL 2008.pdf](http://www.e3mlab.ntua.gr/e3mlab/PRIMES%20Manual/The_PRIMES_MODEL_2008.pdf).



## **ANNEX II – Description policies and measures included in the Reference Level**

This table has been extracted from pp.17-19 in P. Capros, L. Mantzos, N. Tasios, A. De Vita, N. Kouvaritakis (2010), EU energy trends to 2030 — UPDATE 2009, European Commission, Directorate-General for Energy in collaboration with Climate Action DG and Mobility and Transport DG. Luxembourg: Publications Office of the European Union, 2010. ISBN 978-92-79-16191-9. Available online: [http://ec.europa.eu/energy/observatory/trends\\_2030/doc/trends\\_to\\_2030\\_update\\_2009.pdf](http://ec.europa.eu/energy/observatory/trends_2030/doc/trends_to_2030_update_2009.pdf).

**TABLE: INVENTORY OF LEGAL MEASURES AND COMMUNITY FINANCIAL SUPPORT INCLUDED IN PRIMES**

Measure	How the measure is reflected in PRIMES
Regulatory measures	
Energy efficiency	
Eco-design implementing measures	
Eco-design Framework Directive 2005/32/EC	Adaptation of modelling parameters for different product groups. As requirements concern only new products, the effect will be gradual (marginal in 2010; rather small in 2015 and up to full effect by 2030). The potential envisaged in the Eco-design supporting studies and the relationship between cost and efficiency improvements in the model's database were cross-checked.
Stand-by regulation 2008/1275/EC	
Simple Set-to boxes regulation 2009/107/EC	
Office/street lighting regulation 2009/245/EC	
Household lighting regulation 2009/244/EC	
External power supplies regulation 2009/278/EC	
Other energy efficiency	
Labelling Directive 2003/66/EC	Enhancing the price mechanism mirrored in the model
Cogeneration Directive 2004/8/EC	National measures supporting cogeneration are reflected
Directive 2006/32/EC on end-use energy efficiency and energy services	National implementation measures are reflected
Buildings Directive 2002/91/EC	National measures e.g. on strengthening of building codes and integration of RES are reflected
Energy Star Program (voluntary labelling program)	Enhancing the price mechanism mirrored in the model
Energy markets and power generation	
Completion of the internal energy market (including provisions of the 3rd package)	The model reflects the full implementation of the Second Internal market Package by 2010 and Third Internal Market Package by 2015. It simulates liberalised market regime for electricity and gas (decrease of mark-ups of power generation operators; third party access; regulated tariffs for infrastructure use; producers and suppliers are considered as separate companies) with optimal use of interconnectors
EU ETS directive 2003/87/EC as amended by Directive 2008/101/EC and Directive 2009/29/EC	The ETS carbon price is modelled so that the cumulative cap set for GHGs covered by the ETS is respected <sup>3</sup> . The permissible total CDM amount over 2008-2020 is conservatively estimated at 1600 Mt. Banking of allowances is reflected. The model endogenously calculates carbon prices clearing the ETS market that allow to match cumulative emissions over the period 2008-2030 with cumulative allowances assuming the maximum permissible use of CDMs. Resulting carbon prices in the baseline 2009 are: 25 €/08/t CO2eq in 2020 and 39 €/08/t CO2eqin 2030.
Energy Taxation Directive 2003/96/EC	Tax rates (EU minimal rates or higher national ones) are kept constant in real term. The modelling reflects the practice of MS to

<sup>3</sup> For the allocation regime for allowances in 2010, the current system based on National Allocation Plans and essentially cost-free allowances is assumed, with price effects stemming from different investment and dispatch patterns triggered by need to submit allowances. For the further time periods, in the power sector there will be a gradual introduction of full auctioning, which will be fully applicable from 2020 onwards, in line with the specifications of the amended ETS directive. For the other sectors (aviation and industry), the baseline follows a conservative approach which reflects the specifications in the directive on the evolution of auctioning shares and the provisions for free allocation for energy intensive sectors based on benchmarking.

Measure	How the measure is reflected in PRIMES
<b>Regulatory measures</b>	
	increase tax rates above the minimum rate due to i.e. inflation.
Large Combustion Plant directive 2001/80/EC	Emission limit values laid down in part A of Annexes III to VII in respect of sulphur dioxide, nitrogen oxides and dust are respected. Some existing power plants had a derogation which provided them with 2 options to comply with the Directive: either to operate only a limited number of hours or to be upgraded. The model selected between the two options on a case by case basis. The upgrading is reflected through higher capital costs.
IPPC Directive 2008/1/EC	Costs of filters and other devices necessary for compliance are reflected in the parameters of the model
Directive on the geological storage of CO2 2009/31/EC	Enabling measure allowing economic modelling to determine CCS penetration
Directive on national emissions' ceilings for certain pollutants 2001/81/EC	PRIMES model takes into account results of RAINS/GAINS modelling regarding classical pollutants (SO2, NOx). Emission limitations are taken into account bearing in mind that full compliance can also be achieved via additional technical measures in individual MS.
Water Framework Directive 2000/60/EC	Hydro power plants in PRIMES respect the European framework for the protection of all water bodies as defined by the Directive
Landfill Directive 99/31/EC	Provisions on waste treatment and energy recovery are reflected
<b>Transport</b>	
Regulation on CO2 from cars 2009/443/EC	Limits on emissions from new cars: 135 gCO2/km in 2015, 115 in 2020, 95 in 2025 – in test cycle. The 2015 target should be achieved gradually with a compliance of 65% of the fleet in 2012, 75% in 2013, 80% in 2014 and finally 100% in 2015. Penalties for non-compliance are dependent on the number of grams until 2018; starting in 2019 the maximum penalty is charged from the first gram.
Regulation EURO 5 and 6 2007/715/EC	Emission limits introduced for new cars and light commercial vehicles
Fuel Quality Directive 2009/30/EC	Modelling parameters reflect the Directive, taking into account the uncertainty related to the scope of the Directive addressing also parts of the energy chain outside the area of PRIMES modelling (e.g. oil production outside EU).
Biofuels directive 2003/30/EC	Support to biofuels such as tax exemptions and obligation to blend fuels is reflected in the model The requirement of 5.75% of all transportation fuels to be replaced with biofuels by 2010 has not been imposed as the target is indicative. Support to biofuels is assumed to continue. The biofuel blend is assumed to be available on the supply side.
Implementation of MARPOL Convention ANNEX VI - 2008 amendments - revised Annex VI	Amendment of Annex VI of the MARPOL Convention reduce sulphur content in marine fuels which is reflected in the model by a change in refineries output
<b>Financial support</b>	
TEN-E guidelines (Decision 1364/2006)	The model takes into account all TEN-E realised infrastructure projects
European Energy programme for Re-covery (Regulation 2009/663/EC)	Financial support to CCS demonstration plants; off-shore wind and gas and electricity interconnections is reflected in the model. For modelling purposes the following amounts for CCS power plants were assumed, following as-sumptions of summer 2009: Germany: 950 MW (450MW coal post-combustion, 200MW lignite post-combustion and 300MW lignite oxy-fuel), Italy 660 MW (coal post-combustion), Netherlands 1460 MW (800MW coal post-combustion, 660MW coal integrated gasification pre-combustion), Spain 500 MW (coal oxy-fuel), UK 3400 MW (1600MW coal post-combustion, 1800MW coal integrated gasification pre-combustion), Poland 896 MW (306MW coal post-combustion, 590MW lignite post-combustion).
RTD support (7th framework pro-gramme- theme 6)	Financial support to R&D for innovative technologies such as CCS, RES, nuclear and energy efficiency is reflected by technology learning and economies of scale leading to cost reductions of these technologies
State aid Guidelines for Environmental Protection	Financial support to R&D for innovative technologies such as CCS, RES, nuclear and energy efficiency is reflected by technology

Measure	How the measure is reflected in PRIMES
<b>Regulatory measures</b>	
and 2008 Block Exemption Regulation	learning and economies of scale leading to cost reductions of these technologies
Cohesion Policy – ERDF, ESF and Cohesion Fund	Financial support to national policies on energy efficiency and renewables is reflected by facilitating and speeding up the uptake of energy efficiency and renewables technologies.
<b>National measures</b>	
National Forest Plan	This is the framework for forest policy in Spain. It was adopted in 2002, and has validity to 2032. The national forest plan can be found in <a href="http://www.mma.es/portal/secciones/biodiversidad/montes_politica_forestal/estrategia_monte/pfe.htm">http://www.mma.es/portal/secciones/biodiversidad/montes_politica_forestal/estrategia_monte/pfe.htm</a> and in and a summary of the policies included in this plan can be found in the 5th National Communication <a href="http://unfccc.int/national_reports/annex_i_natcom/submitted_natcom/items/4903.php">http://unfccc.int/national_reports/annex_i_natcom/submitted_natcom/items/4903.php</a>
Strong national RES policies	National policies on e.g. feed-in tariffs, quota systems, green certificates, subsidies and other cost incentives are reflected
Nuclear	<p>Nuclear, including the replacement of plants due for retirement, is modelled on its economic merit and in competition with other energy sources for power generation except for MS with legislative provisions on nuclear phase out. Several constraints are put on the model such as decisions of Member States not to use nuclear at all (Austria, Cyprus, Denmark, Estonia, Greece, Ireland, Latvia, Luxembourg, Malta and Portugal) and closure of existing plants in some new Member States according to agreed schedules (Bulgaria 1760 MW, Lithuania 2600 MW and Slovakia 940 MW).</p> <p>The nuclear phase-out in Belgium and Germany is respected while lifetime of nuclear power plants was extended to 60 years in Sweden.</p> <p>Nuclear investments are possible in Bulgaria, the Czech Republic, France, Finland, Hungary, Lithuania, Romania, Slovakia, Slovenia and Spain. For modelling the following plans on new nuclear plants were taken into account: Bulgaria (1000 MW by 2020 and 1000 MW by 2025), Finland (1600 MW by 2015), France (1600 MW by 2015 and 1600 MW by 2020), Lithuania (800 MW by 2020 and 800 MW by 2025), Romania (706 MW by 2010, 776 MW by 2020 and 776 MW by 2025), Slovakia (880 MW by 2015).</p> <p>Member States experts were invited to provide information on new nuclear investments/programmes in spring 2009 and commented on the PRIMES baselines results in summer 2009, which had a significant impact on the modelling results for nuclear capacity.</p>