



Environmental Protection Agency
An Ghníomhaireacht um Chaomhnú Comhshaoil

Ireland's National Inventory Report 2015

Greenhouse Gas Emissions 1990-2013



ENVIRONMENTAL PROTECTION AGENCY

The Environmental Protection Agency (EPA) is responsible for protecting and improving the environment as a valuable asset for the people of Ireland. We are committed to protecting people and the environment from the harmful effects of radiation and pollution.

The work of the EPA can be divided into three main areas:

Regulation: *We implement effective regulation and environmental compliance systems to deliver good environmental outcomes and target those who don't comply.*

Knowledge: *We provide high quality, targeted and timely environmental data, information and assessment to inform decision making at all levels.*

Advocacy: *We work with others to advocate for a clean, productive and well protected environment and for sustainable environmental behaviour.*

Our Responsibilities

Licensing

We regulate the following activities so that they do not endanger human health or harm the environment:

- waste facilities (e.g. landfills, incinerators, waste transfer stations);
- large scale industrial activities (e.g. pharmaceutical, cement manufacturing, power plants);
- intensive agriculture (e.g. pigs, poultry);
- the contained use and controlled release of Genetically Modified Organisms (GMOs);
- sources of ionising radiation (e.g. x-ray and radiotherapy equipment, industrial sources);
- large petrol storage facilities;
- waste water discharges;
- dumping at sea activities.

National Environmental Enforcement

- Conducting an annual programme of audits and inspections of EPA licensed facilities.
- Overseeing local authorities' environmental protection responsibilities.
- Supervising the supply of drinking water by public water suppliers.
- Working with local authorities and other agencies to tackle environmental crime by co-ordinating a national enforcement network, targeting offenders and overseeing remediation.
- Enforcing Regulations such as Waste Electrical and Electronic Equipment (WEEE), Restriction of Hazardous Substances (RoHS) and substances that deplete the ozone layer.
- Prosecuting those who flout environmental law and damage the environment.

Water Management

- Monitoring and reporting on the quality of rivers, lakes, transitional and coastal waters of Ireland and groundwaters; measuring water levels and river flows.
- National coordination and oversight of the Water Framework Directive.
- Monitoring and reporting on Bathing Water Quality.

Monitoring, Analysing and Reporting on the Environment

- Monitoring air quality and implementing the EU Clean Air for Europe (CAFE) Directive.
- Independent reporting to inform decision making by national and local government (e.g. *periodic reporting on the State of Ireland's Environment and Indicator Reports*).

Regulating Ireland's Greenhouse Gas Emissions

- Preparing Ireland's greenhouse gas inventories and projections.
- Implementing the Emissions Trading Directive, for over 100 of the largest producers of carbon dioxide in Ireland.

Environmental Research and Development

- Funding environmental research to identify pressures, inform policy and provide solutions in the areas of climate, water and sustainability.

Strategic Environmental Assessment

- Assessing the impact of proposed plans and programmes on the Irish environment (e.g. *major development plans*).

Radiological Protection

- Monitoring radiation levels, assessing exposure of people in Ireland to ionising radiation.
- Assisting in developing national plans for emergencies arising from nuclear accidents.
- Monitoring developments abroad relating to nuclear installations and radiological safety.
- Providing, or overseeing the provision of, specialist radiation protection services.

Guidance, Accessible Information and Education

- Providing advice and guidance to industry and the public on environmental and radiological protection topics.
- Providing timely and easily accessible environmental information to encourage public participation in environmental decision-making (e.g. *My Local Environment, Radon Maps*).
- Advising Government on matters relating to radiological safety and emergency response.
- Developing a National Hazardous Waste Management Plan to prevent and manage hazardous waste.

Awareness Raising and Behavioural Change

- Generating greater environmental awareness and influencing positive behavioural change by supporting businesses, communities and householders to become more resource efficient.
- Promoting radon testing in homes and workplaces and encouraging remediation where necessary.

Management and structure of the EPA

The EPA is managed by a full time Board, consisting of a Director General and five Directors. The work is carried out across five Offices:

- Office of Climate, Licensing and Resource Use
- Office of Environmental Enforcement
- Office of Environmental Assessment
- Office of Radiological Protection
- Office of Communications and Corporate Services

The EPA is assisted by an Advisory Committee of twelve members who meet regularly to discuss issues of concern and provide advice to the Board.



IRELAND

NATIONAL INVENTORY REPORT 2015

GREENHOUSE GAS EMISSIONS 1990 - 2013 REPORTED TO THE UNITED NATIONS FRAMEWORK CONVENTION ON CLIMATE CHANGE

P. Duffy, E. Hanley, K. Black, P. O'Brien, B. Hyde, J. Ponzi and S. Alam.

Environmental Protection Agency
An Ghníomhaireacht um Chaomhnú Comhshaoil
PO Box 3000, Johnstown Castle, Co. Wexford, Ireland

Telephone: +353 53 9160600 Fax: +353 53 9160699
Email: info@epa.ie Website: www.epa.ie
LoCall 1890 33 55 99

Contents

List of Tables.....	xv
List of Figures.....	xviii
Acknowledgements.....	xxi
EXECUTIVE SUMMARY	1
ES.1 Background.....	1
ES.2 Summary of National Emission and Removal-related Trends.....	2
ES.3 Overview of Source and Sink Category Emission Estimates and Trends	4
ES.4 Indirect Greenhouse Gases	4
Chapter 1 Introduction.....	7
1.1 Background and Context	7
1.2 Introduction and Reporting Requirements under the UNFCCC	7
1.2.1 Scope of Greenhouse Gas Inventories.....	8
1.2.1.1 Greenhouse Gases and Global Warming Potential.....	8
1.2.1.2 Common Reporting Format	9
1.2.1.3 Supplementary Information.....	10
1.3 National Inventory Arrangements.....	10
1.3.1 Institutional, Legal and Procedural Arrangements	10
1.3.2 Overview of Inventory Planning, Preparation and Management.....	13
1.3.3 Quality Assurance, Quality Control and Verification Plan.....	13
1.3.4 Changes in the National Inventory Arrangements since Previous Annual GHG Inventory Submission	15
1.4 Inventory Preparation, and Data Collection, Processing and Storage	15
1.4.1 GHG Inventory and KP-LULUCF Inventory.....	15
1.4.2 Data Collection, Processing and Storage	15
1.5 Methodologies and Emission Factors.....	17
1.5.1 Carbon dioxide (CO ₂)	17
1.5.2 Methane (CH ₄).....	18
1.5.3 Nitrous oxide (N ₂ O).....	18
1.6 Overview of Key Categories.....	18
1.6.1 Key Categories at IPCC Level 2	19
1.6.2 Disaggregated Key Categories.....	23
1.6.3 Use of Key Category Analysis	25
1.7 Uncertainty Evaluation	25
1.8 Completeness and Time-Series Consistency	28
Chapter 2 Trends in greenhouse gas emissions	45
2.1 Description and interpretation of emission trends for aggregated GHG emissions	45
2.2 Trends by Gas	48
2.3 Description and interpretation of emission trends by sector	50
2.3.1 Trends in Energy (IPCC Sector 1).....	51
2.3.2 Trends in Industrial Processes and Product Use (IPCC Sector 2)	54
2.3.3 Trends in Agriculture (IPCC Sector 3)	56
2.3.4 Trends in Land Use, Land Use Change and Forestry (IPCC Sector 4).....	57
2.3.5 Trends in Waste (IPCC Sector 5)	58
2.4 Emissions of Indirect Greenhouse Gases.....	59
Chapter 3 Energy.....	63
3.1 Overview of Energy Sector	63
3.1.1 Emissions Overview	63
3.1.2 Methodology Overview.....	64

3.1.3	Quality Assurance and Quality Control.....	68
3.2	Emissions from Fuel Combustion (1.A).....	69
3.2.1	Comparison of the Sectoral Approach with the Reference Approach.....	69
3.2.2	International Bunker Fuels	70
3.2.3	Feedstocks and Non-energy Use of Fuels.....	71
3.2.4	Energy Industries (1.A.1).....	72
3.2.4.1	Public electricity and heat production (1.A.1.a).....	72_Toc435439315
3.2.4.1.1	Category Description	72
3.2.4.1.2	Methodological Issues	72
3.2.4.1.3	Uncertainties and Time-series Consistency	73
3.2.4.1.4	Category-specific QA/QC and verification	73
3.2.4.1.5	Category-specific Recalculations	74
3.2.4.1.6	Category-specific Planned Improvements	74
3.2.4.2	Petroleum Refining (1.A.1.b).....	74
3.2.4.2.1	Category Description	74
3.2.4.2.2	Methodological Issues	74
3.2.4.2.3	Uncertainties and Time-series Consistency	75
3.2.4.2.4	Category-specific QA/QC and verification	75
3.2.4.2.5	Category-specific Recalculations	75
3.2.4.2.6	Category-specific Planned Improvements	75
3.2.4.3	Manufacture of Solid Fuels and Other Energy Industries (1.A.1.c).....	75
3.2.4.3.1	Category Description	75
3.2.4.3.2	Methodological Issues	76
3.2.4.3.3	Uncertainties and Time-series Consistency	76
3.2.4.3.4	Category-specific QA/QC and verification	76
3.2.4.3.5	Category-specific Recalculations	76
3.2.4.3.6	Category-specific Planned Improvements	76
3.2.5	Manufacturing Industries and Construction (1.A.2)	76
3.2.5.1	Category Description.....	76
3.2.5.2	Methodological Issues.....	77
3.2.5.3	Uncertainties and Time-series Consistency	78
3.2.5.4	Category-specific QA/QC and verification.....	78
3.2.5.5	Category-specific Recalculations.....	78
3.2.5.6	Category-specific Planned Improvements	78
3.2.6	Transport (1.A.3)	79
3.2.6.1	Civil Aviation (1.A.3.a)	79
3.2.6.1.1	Category Description	79
3.2.6.1.2	Methodological Issues	79
3.2.6.1.3	Uncertainties and Time-series Consistency	81
3.2.6.1.4	Category-specific QA/QC and verification	81
3.2.6.1.5	Category-specific Recalculations	83
3.2.6.1.6	Category-specific Planned Improvements	83
3.2.6.2	Road Transportation (1.A.3.b).....	83
3.2.6.2.1	Category Description	83
3.2.6.2.2	Methodological Issues	84
3.2.6.2.3	Uncertainties and Time-series Consistency	96
3.2.6.2.4	Category-specific QA/QC and verification	97
3.2.6.2.5	Category-specific Recalculations	97
3.2.6.2.6	Category-specific Planned Improvements	98
3.2.6.3	Railways (1.A.3.c), Navigation (1.A.3.d) and Other Transportation (1.A.3.e)	98
3.2.6.3.1	Category Description	98

3.2.6.3.2 Methodological Issues	98
3.2.6.3.3 Uncertainties and Time-series Consistency	98
3.2.6.3.4 Category-specific QA/QC and verification	98
3.2.6.3.5 Category-specific Recalculations	99
3.2.6.3.6 Category-specific Planned Improvements	99
3.2.7 Other Sectors (1.A.4)	99
3.2.7.1 Category Description	99
3.2.7.2 Methodological Issues	100
3.2.7.3 Uncertainties and Time-series Consistency	100
3.2.7.4 Category-specific QA/QC and verification	101
3.2.7.5 Category-specific Recalculations	101
3.2.7.6 Category-specific Planned Improvements	101
3.3 Fugitive Emissions (1.B)	101
3.3.1 Coal Mining and Handling (1.B.1.a)	101
3.3.1.1 Category Description	101
3.3.1.2 Methodological Issues	102
3.3.1.3 Uncertainties and Time-series Consistency	103
3.3.1.4 Category-specific QA/QC and verification	103
3.3.1.5 Category-specific Recalculations	103
3.3.1.6 Category-specific Planned Improvements	103
3.3.2 Oil and Natural Gas (NFR 1.B.2)	103
3.3.2.1 Category Description	103
3.3.2.2 Methodological Issues	103
3.3.2.3 Uncertainties and Time-series Consistency	104
3.3.2.4 Category-specific QA/QC and verification	104
3.3.2.5 Category-specific Recalculations	104
3.3.2.6 Category-specific Planned Improvements	104
3.4 CO ₂ Transport and Storage (1.C)	104
Chapter 4 Industrial Processes and Product Use	105
4.1 Overview of the Industrial Processes and Product Use Sector	105
4.1.1 Emissions Overview	105
4.1.2 Methodology Overview	107
4.2 Emissions from Mineral Industry (2.A)	110
4.2.1 Cement Production (2.A.1)	111
4.2.1.1 Category Description	111
4.2.1.2 Methodological Issues	111
4.2.1.3 Uncertainties and Time-series Consistency	112
4.2.1.4 Category-specific QA/QC and verification	112
4.2.1.5 Category-specific Recalculations	113
4.2.1.6 Category-specific Planned Improvements	113
4.2.2 Lime Production (2.A.2)	113
4.2.2.1 Category Description	113
4.2.2.2 Methodological Issues	113
4.2.2.3 Uncertainties and Time-series Consistency	113
4.2.2.4 Category-specific QA/QC and verification	114
4.2.2.5 Category-specific Recalculations	114
4.2.2.6 Category-specific Planned Improvements	114
4.2.3 Glass Production (2.A.3)	114
4.2.3.1 Category Description	114
4.2.3.2 Methodological Issues	115
4.2.3.3 Uncertainties and Time-series Consistency	115

4.2.3.4 Category-specific QA/QC and verification	115
4.2.3.5 Category-specific Recalculations.....	115
4.2.3.6 Category-specific Planned Improvements	115
4.2.4 Other Process Uses of Carbonates (2.A.4)	116
4.2.4.1 Category Description.....	116
4.2.4.2 Methodological Issues	116
4.2.4.3 Uncertainties and Time-series Consistency	117
4.2.4.4 Category-specific QA/QC and verification	117
4.2.4.5 Category-specific Recalculations.....	117
4.2.4.6 Category-specific Planned Improvements	117
4.3 Emissions from Chemical Industry (2.B)	117
4.3.1 Ammonia Production (2.B.1)	118
4.3.1.1 Category Description.....	118
4.3.1.2 Methodological Issues	118
4.3.1.3 Uncertainties and Time-series Consistency	118
4.3.1.4 Category-specific QA/QC and verification	119
4.3.1.5 Category-specific Recalculations.....	119
4.3.1.6 Category-specific Planned Improvements	119
4.3.2 Nitric Acid Production (2.B.2).....	119
4.3.2.1 Category Description.....	119
4.3.2.2 Methodological Issues	119
4.3.2.3 Uncertainties and Time-series Consistency	119
4.3.2.4 Category-specific QA/QC and verification	119
4.3.2.5 Category-specific Recalculations.....	119
4.3.2.6 Category-specific Planned Improvements	120
4.3.3 Adipic Acid Production (2.B.3)	120
4.3.4 Caprolactam, Glyoxal and Glyoxylic Acid Production (2.B.4)	120
4.3.5 Carbide Production (2.B.5)	120
4.3.6 Titanium Dioxide Production (2.B.6).....	120
4.3.7 Soda Ash Production (2.B.7).....	120
4.3.8 Petrochemical and Carbon Black Production (2.B.8)	120
4.3.9 Fluorochemical Production (2.B.9)	120
4.3.10 Other Chemical Industry (2.B.10)	120
4.4 Emissions from Metal Industry (2.C).....	120
4.4.1 Iron and Steel Production (2.C.1)	120
4.4.2 Ferroalloys Production (2.C.2)	121
4.4.3 Aluminium Production (2.C.3).....	121
4.4.4 Magnesium Production (2.C.4)	121
4.4.5 Lead Production (2.C.5).....	121
4.4.6 Zinc Production (2.C.6).....	121
4.4.7 Other Metal Industry (2.C.7)	121
4.5 Emissions from Non-energy Products from Fuels and Solvent Use (2.D).....	121
4.5.1 Lubricant Use (2.D.1)	121
4.5.1.1 Category Description.....	121
4.5.1.2 Methodological Issues	121
4.5.1.3 Uncertainties and Time-series Consistency	122
4.5.1.4 Category-specific QA/QC and verification	122
4.5.1.5 Category-specific Recalculations.....	122
4.5.1.6 Category-specific Planned Improvements	122
4.5.2 Paraffin Wax Use (2.D.2)	122
4.5.2.1 Category Description.....	122

4.5.2.2 Methodological Issues	122
4.5.2.3 Uncertainties and Time-series Consistency	123
4.5.2.4 Category-specific QA/QC and verification	123
4.5.2.5 Category-specific Recalculations.....	123
4.5.2.6 Category-specific Planned Improvements	123
4.5.3 Other Non-energy Products from Fuels and Solvent Use (2.D.3)	123
4.5.3.1 Category Description.....	123
4.5.3.2 Methodological Issues	123
4.5.3.3 Uncertainties and Time-series Consistency	123
4.5.3.4 Category-specific QA/QC and verification	123
4.5.3.5 Category-specific Recalculations.....	124
4.5.3.6 Category-specific Planned Improvements	124
4.6 Emissions from Electronics Industry (2.E)	124
4.6.1 Integrated Circuit or Semiconductor (2.E.1)	124
4.6.1.1 Category Description.....	124
4.6.1.2 Methodological Issues	124
4.6.1.3 Uncertainties and Time-series Consistency	124
4.6.1.4 Category-specific QA/QC and verification	124
4.6.1.5 Category-specific Recalculations.....	125
4.6.1.6 Category-specific Planned Improvements	125
4.6.2 TFT Flat Panel Display Industry (2.E.2)	125
4.6.3 Photovoltaics Industry (2.E.3)	125
4.6.4 Heat Transfer Fluid Use (2.E.4)	125
4.6.5 Other Electronics Industry (2.E.5).....	125
4.7 Emissions from Product Uses as Substitutes for ODS (2.F)	125
4.7.1 Refrigeration and air conditioning (2.F.1)	127
4.7.1.1 Category Description.....	127
4.7.1.2 Methodological Issues	128
4.7.1.3 Uncertainties and Time-series Consistency	128
4.7.1.4 Category-specific QA/QC and verification	128
4.7.1.5 Category-specific Recalculations.....	129
4.7.1.6 Category-specific Planned Improvements	129
4.7.2 Foam Blowing Agents (2.F.2)	129
4.7.2.1 Category Description.....	129
4.7.2.2 Methodological Issues	129
4.7.3 Fire Protection (2.F.3).....	129
4.7.3.1 Category Description.....	129
4.7.3.2 Methodological Issues	130
4.7.3.3 Uncertainties and Time-series Consistency	130
4.7.3.4 Category-specific QA/QC and verification	130
4.7.3.5 Category-specific Recalculations.....	130
4.7.3.6 Category-specific Planned Improvements	130
4.7.4 Aerosols (2.F.4).....	130
4.7.4.1 Category Description.....	130
4.7.4.2 Methodological Issues	131
4.7.4.3 Uncertainties and Time-series Consistency	132
4.7.4.4 Category-specific QA/QC and verification	132
4.7.4.5 Category-specific Recalculations.....	132
4.7.4.6 Category-specific Planned Improvements	132
4.7.5 Solvents (2.F.5)	132
4.7.6 Other Product Uses as Substitutes for ODS (2.F.6).....	132

4.8 Emissions from Other Product Manufacture and Use (2.G)	132
4.8.1 Electrical Equipment (2.G.1)	133
4.8.1.1 Category Description	133
4.8.1.2 Methodological Issues	133
4.8.1.3 Uncertainties and Time-series Consistency	133
4.8.1.4 Category-specific QA/QC and verification	134
4.8.1.5 Category-specific Recalculations	134
4.8.1.6 Category-specific Planned Improvements	134
4.8.2 SF ₆ and PFCs from Other Product Uses (2.G.2)	134
4.8.2.1 Category Description	134
4.8.2.2 Methodological Issues	134
4.8.2.3 Uncertainties and Time-series Consistency	135
4.8.2.4 Category-specific QA/QC and verification	135
4.8.2.5 Category-specific Recalculations	135
4.8.2.6 Category-specific Planned Improvements	135
4.8.3 N ₂ O from Product Use (2.G.3)	135
4.8.3.1 Category Description	135
4.8.3.2 Methodological Issues	136
4.8.3.3 Uncertainties and Time-series Consistency	136
4.8.3.4 Category-specific QA/QC and verification	136
4.8.3.5 Category-specific Recalculations	136
4.8.3.6 Category-specific Planned Improvements	136
4.8.4 Other – Other Product Manufacture and Use (2.G.4)	136
4.9 Other – (2.H)	136
Chapter 5 Agriculture	137
5.1 Overview of Agriculture Sector	137
5.1.1 Emissions Overview	137
5.1.2 Methodology Overview	138
Emissions from livestock (3.1)	141
5.2 Emissions from Enteric Fermentation (3.A)	142
5.2.1 Enteric Fermentation, Cattle (3.A.1)	142
5.2.1.1 Category Description	142
5.2.1.2 Methodological Issues	142
5.2.1.3 Uncertainties and Time-series Consistency	146
5.2.1.4 Category-specific QA/QC and verification	146
5.2.1.5 Category-specific Recalculations	146
5.2.1.6 Category-specific Planned Improvements	146
5.2.2 Enteric Fermentation, All Other Livestock (3.A.2-3.A.4)	146
5.2.2.1 Category Description	146
5.2.2.2 Methodological Issues	146
5.2.2.3 Uncertainties and Time-series Consistency	147
5.2.2.4 Category-specific QA/QC and verification	147
5.2.2.5 Category-specific Recalculations	147
5.2.2.6 Category-specific Planned Improvements	147
5.3 Emissions from Manure Management (3.B)	147
5.3.1 Manure Management, Cattle (3.B.1)	147
5.3.1.1 Category Description	147
5.3.1.2 Methodological Issues	148
5.3.1.3 Uncertainties and Time-series Consistency	150
5.3.1.4 Category-specific QA/QC and verification	150
5.3.1.5 Category-specific Recalculations	150

5.3.1.6 Category-specific Planned Improvements	150
5.3.2 Manure Management, All Other Livestock (3.B.2-3.B.4)	150
5.3.2.1 Category Description.....	150
5.3.2.2 Methodological Issues	150
5.3.2.3 Uncertainties and Time-series Consistency	150
5.3.2.4 Category-specific QA/QC and verification	151
5.3.2.5 Category-specific Recalculations.....	151
5.3.2.6 Category-specific Planned Improvements	151
5.4 Emissions from Rice Cultivation (3.C)	151
5.5 Emissions from Agricultural Soils (3.D)	151
5.5.1 Direct N ₂ O Emissions From Managed Soils (3.D.1)	151
5.5.1.1 Category Description.....	151
5.5.1.2 Methodological Issues	152
5.5.1.3 Uncertainties and Time-series Consistency	153
5.5.1.4 Category-specific QA/QC and verification	153
5.5.1.5 Category-specific Recalculations.....	153
5.5.1.6 Category-specific Planned Improvements	153
5.5.2 Indirect N ₂ O Emissions from Managed Soils (3.D.2).....	154
5.5.2.1 Category Description.....	154
5.5.2.2 Methodological Issues	154
5.5.2.3 Uncertainties and Time-series Consistency	155
5.5.2.4 Category-specific QA/QC and verification	155
5.5.2.5 Category-specific Recalculations.....	155
5.5.2.6 Category-specific Planned Improvements	155
5.6 Emissions from Prescribed Burning of Savannas (3.E)	156
5.7 Emissions from Field Burning of Agricultural Residues (3.F)	156
5.8 Emissions from Liming (3.G)	156
5.8.1 Category Description	156
5.8.2 Methodological Issues.....	156
5.8.3 Uncertainties and Time-series Consistency	156
5.8.4 Category-specific QA/QC and verification.....	156
5.8.5 Category-specific Recalculations	156
5.8.6 Category-specific Planned Improvements	156
5.9 Emissions from Urea Application (3.H)	157
5.9.1 Category Description	157
5.9.2 Methodological Issues.....	157
5.9.3 Uncertainties and Time-series Consistency	157
5.9.4 Category-specific QA/QC and verification.....	157
5.9.5 Category-specific Recalculations	157
5.9.6 Category-specific Planned Improvements	157
5.10 Emissions from Other Carbon-Containing Fertilisers (3.I)	157
5.11 Emissions from Other Agricultural Sources (3.J).....	157
Chapter 6 Land-Use, Land-Use Change and Forestry	159
6.1 Introduction	159
6.2 Overview of LULUCF Sector	159
6.2.1 Sector Coverage	159
6.2.2 Land Use Definitions and Land Use Change Matrices	161
6.2.2.1 Land use classification hierarchy	162
6.2.3 Land use change trends	163
6.3 Forest Land (Category 4.A).....	169
6.3.1 Overall approach and activity data.....	169

6.3.2 Detailed description of activity data	171
6.3.2.1 FIPS95	171
6.3.2.2 IFORIS	171
6.3.2.3 The National Forest Inventory	172
6.3.2.4 Harvests and Deforestation	174
6.3.2.5 Activity Data for Afforestation Areas	178
6.3.2.6 Definition of carbon pools	180
6.3.3 Description of models used	180
6.3.3.1 CARBWARE	180
6.3.3.1.1 Pre-processing, Growth Simulator and Stand Modifier Modules	181
6.3.3.1.2 The Carbon Flow Sub-model	181
6.3.3.1.3 Scaling and Aggregation of PSP into Different Reporting Categories	187
6.3.3.1.4 Datasets Used to Develop the CARBWARE Models	188
6.3.3.2 FORCARB	188
6.3.4 Forest land remaining forest land (CRF 4.A.1)	190
6.3.4.1 Time Series Adjustment of Living Biomass and DOM Pools	192
6.3.4.2 Mineral soils	194
6.3.4.3 Organic Soils	194
6.3.4.4 Emissions from Biomass Burning	196
6.3.4.5 Direct and indirect emissions of N ₂ O from organic and synthetic fertilisers	196
6.3.4.6 Emissions of N ₂ O and CH ₄ from drainage and rewetted organic soils	197
6.3.4.7 N ₂ O emissions from mineral soils as a result of land use change of management (F _{SOM})	199
6.3.4.8 Uncertainty Analysis for Category 4.A.1	200
6.3.5 Forest land remaining forest land (CRF 4.A.2)	205
6.3.5.1 Time Series Adjustment of Living Biomass and DOM Pools	205
6.3.5.2 Mineral Soils	207
6.3.5.3 Organic Soils	207
6.3.5.4 Emissions from Biomass Burning	208
6.3.5.5 Emissions of N ₂ O from Fertilization	208
6.3.5.6 Emissions of N ₂ O and CH ₄ from drainage and rewetted organic soils	209
6.3.5.7 N ₂ O emissions from mineral soils as a result of land use change of management (F _{SOM})	210
6.3.5.8 CO ₂ emissions from urea application to soils	210
6.3.5.9 Uncertainty Analysis for Category 4.A.2	210
6.3.6 Deforestation Areas (CRF 4.B.2 to 4.F.2)	212
6.3.6.1 Deforestation Losses	213
6.3.6.1.1 Mineral soils	215
6.3.6.1.2 Organic soils	216
6.3.6.1.3 Uncertainty for deforestation estimates	217
6.3.7.1 Harvested wood products (4.G)	219
6.3.7.2 HWP uncertainties	221
6.4 Cropland (4.B)	224
6.4.1 Description	224
6.4.2 Soil Type and Soil Organic Carbon	228
6.4.3 Cropland Areas	229
6.4.4 Soil	233
6.4.5 Carbon Stock Change in Biomass	235
6.4.6 Cropland Dead Organic Matter/Litter	238
6.4.7 Carbon Stock Change in Soils	238
6.4.8 Estimation of Emissions from Soils	238

6.4.9 Cropland emissions due to Biomass Burning.....	240
6.4.10 Uncertainties and time-series consistency in Cropland.....	244
6.4.11 Category-specific QA/QC and verification.....	245
6.4.12 Cropland Category-specific recalculations and impact on emission trend	245
6.4.13 Cropland recalculations and impact on emission trend.....	246
6.4.14 Cropland Category-specific planned improvements	247
6.5 Grassland (4.C)	247
6.5.1 Grassland Areas.....	247
6.5.2 Methodological issues.....	251
6.5.2.1 Carbon Stock Changes in Grassland.....	251
6.5.2.2 Carbon Stock Changes in Living Biomass	251
6.5.2.3 Dead Organic Matter/Litter	252
6.5.2.4 Carbon Stock Changes in Soils	252
6.5.2.5 CH ₄ emissions due to drainage of grasslands on organic soils	257
6.5.3 Land converted to Grassland.....	259
6.5.3.1 Forest Land converted to Grassland	259
6.5.3.2 Wetland converted to Grassland	259
6.5.4 Grassland emissions due to Biomass Burning	260
6.5.5 Uncertainties and time-series consistency	263
6.5.6 Category-specific QA/QC and verification.....	264
6.5.7 Grassland -specific recalculations and impact on emission trend.....	264
6.5.8 Grassland Category-specific planned improvements	266
6.6 Wetlands (4.D)	266
6.6.1 Wetland Areas.....	266
6.6.2 Unmanaged Wetland Areas.....	267
6.6.3 Exploited Peatland Areas	267
6.6.4 Carbon Stock Changes in Wetland	268
6.6.4.1 Biomass	268
6.6.4.2 Soils	268
6.6.5 Emissions of Non-CO ₂ Gases.....	269
6.6.5.1 N ₂ O emissions due to drainage of peatland for peat extraction	269
6.6.5.2 CH ₄ emissions due to drainage of peatland for peat extraction.....	270
6.6.6 Emissions due to Extraction and Use of Horticultural Peat.....	271
6.6.6.1 On-site emissions.....	272
6.6.6.2 Off-site emissions.....	272
6.6.7 Wetland emissions due to Biomass Burning.....	273
6.6.8 Wetland recalculations and impact on emission trend.....	276
6.6.9 Wetland planned improvements.....	277
6.7 Settlements (Category 4.E)	277
6.7.1 Areas of Settlements	277
6.7.2 Carbon Stock Changes in Settlements	278
6.7.2.1 Biomass	278
6.7.2.2 Soils	279
6.7.2.3 Biomass Burning on Settlement.....	279
6.7.3 Settlements recalculations and impact on emission trend	279
6.8 Other Land (Category 4.F).....	280
6.8.1 Areas of Other Land.....	280
6.8.2 Carbon Stock Changes in Other Land.....	280
6.8.3 Biomass Burning on Other Land	280
6.8.4 Other Land recalculations and impact on emission trend.....	280
6.9 Uncertainty in non-Forest LULUCF categories.....	281

6.9.1 Uncertainty in Cropland	282
6.9.2 Uncertainty in Grassland	283
6.9.3 Uncertainty in Wetlands	284
6.9.4 Uncertainty in Settlements	284
6.9.5 Uncertainty in Other Land	285
6.10 Quality Assurance and Quality Control	285
6.10.1 Category specific QA/QC for Forest Lands	285
6.10.2 QC plan for Activity Data	285
6.10.3 Emission Factors, Models and Calculations	286
6.10.4 Completeness and error checks in compilation of the CRF tables	286
6.10.5 Validation and QA/QC Links to Uncertainty Analysis	286
6.10.6 Validation of Reported Estimates (Category 4.A.1)	287
6.10.7 Validation of reported estimates (Category 4.A.2)	288
6.10.8 Independent External Reviews	290
6.11 Recalculations in LULUCF	290
6.11.1 Forest lands (4.A)	290
6.11.2 Cropland (4.B)	291
6.11.3 Grassland (4.C)	291
6.11.4 Wetlands (4.D)	292
6.12 Improvements in LULUCF	292
6.12.1 Planned Improvements for Forest Lands	293
6.12.2 Planned Improvements for Other Categories	294
Chapter 7 Waste	295
7.1 Overview of the Waste Sector	295
7.1.1 Emissions Overview	295
7.1.2 Methodology Overview	296
7.2 Emissions from Solid Waste Disposal (5.A)	299
7.2.1 Managed Waste Disposal Sites (5.A.1)	299
7.2.1.1 Category Description	299
7.2.1.2 Methodological Issues	299
7.2.1.3 Uncertainties and Time-series Consistency	303
7.2.1.4 Category-specific QA/QC and verification	303
7.2.1.5 Category-specific Recalculations	303
7.2.1.6 Category-specific Planned Improvements	303
7.2.2 Unmanaged Waste Disposal Sites (5.A.2)	303
7.2.2.1 Category Description	303
7.2.2.2 Methodological Issues	303
7.2.2.3 Uncertainties and Time-series Consistency	304
7.2.2.4 Category-specific QA/QC and verification	304
7.2.2.5 Category-specific Recalculations	304
7.2.2.6 Category-specific Planned Improvements	304
7.3 Emissions from Biological Treatment of Solid Waste (5.B)	304
7.3.1 Composting (5.B.1)	304
7.3.1.1 Category Description	304
7.3.1.2 Methodological Issues	304
7.3.1.3 Uncertainties and Time-series Consistency	305
7.3.1.4 Category-specific QA/QC and verification	305
7.3.1.5 Category-specific Recalculations	305
7.3.1.6 Category-specific Planned Improvements	305
7.3.2 Anaerobic Digestion at Biogas Facilities (5.B.2)	305
7.4 Emissions from Incineration and Open Burning of Waste (5.C)	306

7.4.1 Waste Incineration (5.C.1)	306
7.4.1.1 Category Description.....	306
7.4.1.2 Methodological Issues	306
7.4.1.3 Uncertainties and Time-series Consistency	308
7.4.1.4 Category-specific QA/QC and verification	308
7.4.1.5 Category-specific Recalculations.....	308
7.4.1.6 Category-specific Planned Improvements	308
7.4.2 Open Burning of Waste (5.C.2)	308
7.4.2.1 Category Description.....	308
7.4.2.2 Methodological Issues	308
7.4.2.3 Uncertainties and Time-series Consistency	309
7.4.2.4 Category-specific QA/QC and verification	309
7.4.2.5 Category-specific Recalculations.....	309
7.4.2.6 Category-specific Planned Improvements	309
7.5 Emissions from Wastewater Treatment and Discharge (5.D).....	309
7.5.1 Domestic Wastewater (5.D.1)	309
7.5.1.1 Category Description.....	309
7.5.1.2 Methodological Issues	309
7.5.1.3 Uncertainties and Time-series Consistency	313
7.5.1.4 Category-specific QA/QC and verification	313
7.5.1.5 Category-specific Recalculations.....	313
7.5.1.6 Category-specific Planned Improvements	313
7.5.2 Industrial Wastewater (5.D.2)	314
7.6 Emissions from Other Waste Sources (5.E)	314
Chapter 8 Other Sources	315
Chapter 9 Indirect CO₂ and N₂O Emissions	317
9.1 Description of Sources of Indirect Emissions in GHG Inventory.....	317
9.2 Methodological Issues	317
9.3 Uncertainties and Time-series Consistency.....	318
9.4 Category-specific QA/QC and Verification.....	318
9.5 Category-specific Recalculations	318
9.6 Category-specific Planned Improvements.....	318
Chapter 10 Recalculations and Improvements	319
10.1 Introduction	319
10.2 Explanation and Justification for Recalculations	319
10.2.1 Recalculations in Energy	320
10.2.2 Recalculations in Industrial Processes and Product Use.....	320
10.2.3 Recalculations in Agriculture.....	321
10.2.4 Recalculations in LULUCF.....	322
10.2.4.1 Forest lands (4.A)	322
10.2.4.2 Cropland (4.B).....	322
10.2.4.3 Grassland (4.C)	323
10.2.4.4 Wetlands (4.D).....	324
10.2.4.5 Settlement (4.E)	324
10.2.4.6 Other Land (4.F)	324
10.2.5 Recalculations in Waste.....	325
10.3 Effects on Emission Levels, Trends and Time-Series Consistency.....	326
10.4 Response to the Review Process and Planned Improvements.....	332

PART II SUPPLEMENTARY INFORMATION REQUIRED UNDER ARTICLE 7.1 OF THE KYOTO PROTOCOL	333
Chapter 11 Emissions and Removals from LULUCF Activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol	335
11.1 General Information	335
11.1.1 Introduction	335
11.1.2 Institutional Arrangements	335
11.1.3 Definitions and Application	340
11.1.4 The geographical location of the boundaries of the areas that encompass activities under Article 3, paragraphs 3 and 4	342
11.1.5 Classification hierarchy and continuity of accounting reported activities over time	342
11.1.6.1 The information on identifiable units of land under mandatory activities and spatial assessment units.	343
11.1.6.2 The information on identifiable units of land under elected activities and spatial assessment units.	344
11.1.7 Information that demonstrated elected activities are directly human induced and have occurred since 1990	345
11.2 Methodologies and description of data	347
11.2.1 Afforestation	347
11.2.2 Forest management	348
11.2.3 Deforestation	351
11.2.3.1 Information on how harvesting or forest disturbance that is followed by the re- establishment of a forest is distinguished from deforestation	351
11.2.3.2 Deforestation Information	351
11.2.4 Cropland management	352
11.2.5 Grazing land management	352
11.2.6 Direct and indirect emissions from N fertilisation	353
11.2.7 N ₂ O and CH ₄ from drained and rewetted organic soils	353
11.2.8 N ₂ O losses from mineralization of soils due C loss associated with land use change	353
11.2.9 Biomass burning from Forest ARD and FM Fires	353
11.2.10 Biomass burning from CM Fires	354
11.2.11 Biomass burning from GM Fires	354
11.3 Justification for Omitting a Carbon Pool	354
11.3.1 Afforestation: Mineral Soils for grassland, cropland and forest land transitions	354
11.3.2 Forest management: Mineral Soils	359
11.3.3 Cropland management Matter Carbon Pools	360
11.3.4 Grazing land management Carbon Pools	361
11.4 Natural disturbances	362
11.4.1 Calculation of background and margin	362
11.4.2 Trigger test for implementation of the natural disturbance provision	364
11.4.3 Exclusion of emissions from salvage logging	364
11.5 FMRL and technical corrections	365
11.6 Harvested wood products	367
11.6.1 Information on activity data for the harvested wood products categories used for estimating the harvested wood products pool removed from domestic forests, for domestic consumption and for export, (para 2g(i and vii) of annex II of 2/CMP.8)	368
11.6.2 Information on half-lives used in estimating the emissions and removals for these categories in accordance with decision 2/CMP.7, annex, paragraph 29 or 30	368
11.6.3 Information on whether emissions from harvested wood products originating from forests prior to the start of the second commitment period have been included in the accounting (see para 2g(iii) of annex II to decision 2.CMP.8).	368

11.6.4 Information on how emissions from the harvested wood products pool that have been accounted for during the first commitment period on the basis of instantaneous oxidation have been excluded from the accounting for the second commitment period; (see para 2g(iv) of annex II to decision 2.CMP.8).	368
11.6.5 Information showing that harvested wood products resulting from deforestation have been accounted on the basis of instantaneous oxidation; (see para 2g(v) of annex II to decision 2.CMP.8).....	369
11.6.6 Information showing that carbon dioxide emissions from harvested wood products in solid waste disposal sites, where these emissions are separately accounted for, and from wood harvested for energy purposes have been accounted on the basis of instantaneous oxidation;	370
11.7 Uncertainty Analysis	370
11.8 Quality Assurance and Quality Control (QA\QC)	371
11.9 Recalculations in KP LULUCF	371
Chapter 12 Information on Accounting of Kyoto Units.....	373
12.1 Background Information	373
12.2 Summary of Information Reported in the SEF Tables.....	373
12.3 Discrepancies and notifications	374
12.4 Publicly Accessible Information	375
12.5 Calculation of the Commitment Period Reserve.....	378
12.6 Accounting for Activities under Article 3.3	378
Chapter 13 Changes in National System	379
13.1 Changes in National System since previous submission	379
Chapter 14 Changes in National Registry.....	381
14.1 Introduction	381
14.2 Information on Changes in National Registry	381
Chapter 15 Minimisation of Adverse Impacts under Article 3, paragraph14	387
15.1 Introduction	387
15.2 Context.....	387
15.3 Specific Elements	387
Glossary.....	391
References.....	393
 Annex 1	 405
Key Category Analyses	405
Annex 2	423
Assessment of Uncertainty	423
Annex 3.1.A	433
Energy - Combustion (IPCC Sector 1.A).....	433
Annex 3.1.B	447
Energy - Transport (IPCC Sector 1.A.3)	447
Annex 3.2	457
Industrial Processes (IPCC Sector 2).....	457
Annex 3.3	463
Agriculture (IPCC Sector 3).....	463
Annex 3.4	473
Activity Data for LULUCF (IPCC Sector 4.A) Forest Land	473
Annex 3.5	527
Waste (IPCC Sector 5)	527
Annex 4	533
Ireland's Energy Balance 1990-2013.....	533

Annex 5.1	547
Ireland's Response to the Recommendations in the UNFCCC Annual Review Reports	547
Annex 5.3	569
Standard Electronic Format (SEF) 2014	569
Annex 5.4	577
Greenhouse Gases GWP and IPCC Reporting Format	577
Appendix 1	581
Standard Independent Assessment Report	581
(Electronic Appendix)	581

List of Tables

Table 1.1 Key Data Providers and Information covered by MOU	16
Table 1.2 Key Categories at IPCC Level 2 in 1990	20
Table 1.3 Key Categories at IPCC Level 2 in 2013	20
Table 1.4 Summary of Methods	21
Table 1.5 Summary of Emission Factors	22
Table 1.6 Key Category Analysis Level Assessment 1990 (excluding LULUCF)	29
Table 1.7 Key Category Analysis Level Assessment 2013 (excluding LULUCF)	30
Table 1.8 Key Category Analysis Level Assessment 1990 (including LULUCF)	31
Table 1.9 Key Category Analysis Level Assessment 2013 (including LULUCF)	32
Table 1.10 Key Category Analysis Trend Assessment 1990-2013 (excluding LULUCF)	33
Table 1.11 Key Category Analysis Trend Assessment 2013 (including LULUCF)	34
Table 1.12 Tier 1 Uncertainty Estimates 2013 excluding LULUCF (continued on following pages).....	35
Table 1.13 Tier 1 Uncertainty Estimates 2013 including LULUCF (continued on following pages)	39
Table 1.14 Summary of Completeness	43
Table 2.1. Greenhouse Gas Emissions 1990, 1995, 2000-2013 (kt CO ₂ equivalent).....	47
Table 2.2 Emissions of NO _x , SO ₂ , NMVOC and CO 1990-2013 (Tonnes).....	60
Table 3.1 Level 3 Source Methodology for Energy	66
Table 3.2 Emissions from Energy 1990-2013.....	67
Table 3.3 Allocated CO ₂ emissions from fuel used for non-energy purpose	71
Table 3.4 EURO class vehicle commencement years.....	84
Table 3.5 Emission factors for Rail and Navigation	98
Table 3.6 Emission factors for Agriculture.....	100
Table 3.7 Emission factors for Charcoal use in Residential	100
Table 3.8 Emission factors for underground mining and post-mining activities	102
Table 3.9 Emission factors for Abandoned underground mines (1.B.1.a.1(ii)).....	103
Table 4.1 Level 3 Source Methodology for IPPU	108
Table 4.2 Emissions from Industrial Processes and Product Use 1990-2013	109
Table 4.3. Emissions of HFC, PFC, SF ₆ and NF ₃ from IPPU 1990-2013 (kt CO ₂ eq	126
Table 5.1 Level 3 Source Methodology for Agriculture	138
Table 5.2 Emissions from Agriculture 1990-2013	140
Table 5.3 Animal Classifications for Cattle Population	143
Table 5.4 Tier 2 CH ₄ Enteric Fermentation Emission Factors for 1990, 2007- 2013	146
Table 5.5 Tier 2 CH ₄ Manure Management Emission Factors for 1990, 2007- 2013.....	149
Table 5.6 Information related to Direct N ₂ O Emissions from Managed Soils (3.D.1).....	153
Table 5.7 Information related to Indirect N ₂ O Emissions from Managed Soils (3.D.2)	155
Table 6.1 Level 3 Source Category Coverage for Land Use, Land-Use Change and Forestry.....	160
Table 6.2 Emissions ^a and Removals ^a from Land Use Land-Use Change and Forestry 1990-2013 (kt CO ₂ eq)....	164
Table 6.3 Description of Land Use Categories	165
Table 6.4 Summary Land Use Matrices 1990-2013 (units ha)	166
Table 6.5.1 Definition of carbon pools used in LULUCF and KP-LULUCF reporting	180
Table 6.5.2 Breakdown of species used in the pre-1990 and post-1990 forest categories.....	189
Table 6.5.3 Yield class, silviculture and rotation criteria selected for periods 1990-1999 and 2000-2012	189
Table 6.5.4 Time series for the forest category 4.A.1.....	191
Table 6.5.5 Area (in kHa) and emissions from organic soils over the times series for forest land remaining forests.....	195
Table 6.5.6 Area statistics and emission profiles over the time series 1990 to 2013 for wild fires in categories 4.A.1 and 4.A.2 and reported in table 4(V).....	198
Table 6.5.7 The area activity data and N ₂ O and CH ₄ emissions from drainage of forest land remaining forest land	199
Table 6.5.8 Uncertainty estimates for individual activity and area data sets for forest land remaining forest land	201
Table 6.5.9 Combined uncertainty estimates for forest land remaining forest land pools.....	201
Table 6.5.10 NFI external validation of CARBWARE models	202

Table 6.5.11 Uncertainty analysis for forest land remaining forest land since 1990 category	203
Table 6.5.12 Time series for forest category 4.A.2.....	206
Table 6.5.13 Area (in kha) and emissions from different organic soils types over the times series for land converted to forests	208
Table 6.5.14 The area activity data and N ₂ O and CH ₄ emissions from drainage of land converted to forest land	209
Table 6.5.15 Uncertainty analysis of lands converted to forest land as reported in CRF 4.A.2.....	210
Table 6.5.16 The new deforestation, land use change and soil type matrix showing annual deforestation areas (kha/ year) associated with different land uses and soils categories	213
Table 6.5.17 Deforestation carbon stock changes and harvest over the time series.....	215
Table 6.5.18 Uncertainty estimates for individual activity and area data sets for deforested lands	217
Table 6.5.19 Combined uncertainty estimates for deforested land.....	217
Table 6.5.20 Uncertainty estimates for deforested land.....	218
Table 6.5.21 Annual inflows of sawnwood (SW), wood-based panels (WBP) , paper and paper board (PPB) from domestic harvest.	220
Table 6.5.22 Detailed inflows and CSC for different HWPs in the land converted to forest land (i.e. AR sub-category or 4A2). Note: Harvest in the AR category starts from 2007 onwards.	222
Table 6.5.23 Detailed inflows and CSC for different HWPs in the forestland remaining forest land remaining (i.e .FM sub-category or 4A1).	222
Table 6.5.24 Detailed inflows and CSC for different HWPs from category 4A	223
Table 6.5.25 Uncertainty of HWP estimates for categories 4A1 and 4A2	223
Table 6.6.1 Soil Class Coverage and Soil Organic Carbon	229
Table 6.6.2 Examples of binary coding of cropland parcel history	233
Table 6.6.3 Summary of Cropland parcel histories.....	233
Table 6.6.4 Adjustment Factors for SOC.....	239
Table 6.6.5 Carbon content of cropland soils as a function of period under grass	240
Table 6.6.6 Land Cover/Use associated with NASA FIRMS fires	242
Table 6.6.7 Tier 1 Default parameters for use in Eq 6.5.4	262
Table 6.7 Summary of Uncertainty analysis	281
Table 6.8.1 Comparison of 2013 inventory year IEFs reported for other countries and those reported by Ireland forest land remaining forest land (4A1)	287
Table 6.8.2 Comparisons of 2010 inventory year IEFs reported for other countries and those reported by Ireland land converted forest land	288
Table 6.8.3 Validation of net biome productivity	289
Table 7.1 Level 3 Source Methodology for Waste	296
Table 7.2 Emissions from Waste 1990-2013	297
Table 7.3. Methane Emissions from Solid Waste Disposal 1990-2013.....	302
Table 7.4 Information related to Managed Waste Disposal (5.A.1)	302
Table 7.5 Information related to Unmanaged Waste Disposal Sites (5.A.2)	304
Table 7.6 Information related to Composting (5.B.1).....	305
Table 7.7 Information related to Waste Incineration (5.C.1)	307
Table 7.8 Information related to Open Burning of Waste (5.C.2)	308
Table 7.9 Information related to Domestic Wastewater (5.D.1).....	311
Table 7.10 Estimates of N ₂ O emissions from human sewage 1990-2013	313
Table 10.1. Changes in Methodological Descriptions compared to 2014 NIR.....	327
Table 10.2. Recalculations by Gas 1990-2012	328
Table 10.3 Recalculations by IPCC Sector 1990-2012	330
Table 11.1 Reported Activities and Pools (CRF Table NIR 1)	337
Table 11.2 Land Transition Matrix (CRF Table NIR 2) for inventory year 2013.....	338
Table 11.3 Key Categories for Article 3.3 and A.4 Activities (CRF Table NIR 3)	339
Table 11.4 Information Table on Reporting of emission/removals for Activities under Article 3, paragraphs 3 and 4 of the Kyoto Protocol.....	339
Table 11.5 Forest category codes used in CRF Tables 5 (KP1)	344
Table 11.6 Detailed areas and CSC for level 3 forest categories under AR activities in 2013	349
Table 11.7 Detailed areas and CSC for level 3 forest categories under FM activities in 2013	350
Table 11.8 CSC for forestland converted to other land categories	352
Table 11.9 Record sample plots taken from mineral soils sites	356

Table 11.10 Results from the hierarchical analysis of variance on soils C at a depth of 0-30cm	356
Table 11.11a Total emissions from wild fires and area specific emissions from disturbances for the calibration period for FM.....	363
Table 11.11b Total and area specific emissions from disturbances for the calibration period for AR	363
Table 11.12 Calculation steps used to derive the background and harvest for FM and AR for the second commitment period.....	364
Table 11.13 A comparison of previous activity data for 1990-2009 used for the FMRL submission and the new data used in the 2013 inventory.....	366
Table 11.14 Summary of the FMRL recalculated (FMRLcorr) values for different pools in 2013, compared to the FMRL and reported FM (reported FM) emission/removal for 2013	367
Table 11.15 Adjusted HWP stock and stock changes for AR activities since 2007	368
Table 11.16 The estimated fractions of HWP feedstock originating from different forest activities under article 3.3 and 3.4	369
Table 11.17 Uncertainty estimates of forest activity estimates provided in CRF tables.....	370
Table 12.1 Information on the SEF tables.....	374
Table 12.2 Discrepancies and notifications	374
Table 12.3 Publicly Accessible Information	375

List of Figures

Figure ES.1 National total Greenhouse Gas emissions (excluding LULUCF) 1990-2013	3
Figure 1.1 National Inventory System Overview	12
Figure 2.1 National total Greenhouse Gas emissions (excluding LULUCF) 1990-2013.....	45
Figure 2.2 Inter annual change in national total emissions 1990-2013.....	46
Figure 2.3 Greenhouse Gas emissions-by Gas 1990-2013.....	48
Figure 2.4 Total Primary Energy Requirement (TPER) 1990-2013	52
Figure 2.5 Trend in Emissions from Energy 1990-2013	52
Figure 2.6 Fuel use in Road Transport 1990-2013	53
Figure 2.7 Vehicle numbers and Census of Population 1990-2013	54
Figure 2.8 Trend in Emissions from Industrial Processes and Product Use 1990-2013	55
Figure 2.9 Trend in Emissions from Agriculture 1990-2013	57
Figure 2.10 Trend in Emissions and Removals from Land Use Land-Use Change and Forestry 1990-2013	58
Figure 2.11 Trend in Emissions from Waste 1990-2013	59
Figure 2.12 Trend in Indirect Greenhouse Gases 1990-2013	61
Figure 3.1 Total Emissions from Energy by Category, 1990-2013	68
Figure 3.2 Total Emissions from Energy by Gas, 1990-2013.....	68
Figure 3.3 Emissions from 1.A.1.a Public Electricity and Heat Production 1990-2013.....	72
Figure 3.4 Emissions from 1.A.2 Manufacturing Industries and Construction 1990-2013	77
Figure 3.5 Emissions from 1.A.3 Transport 1990-2013.....	79
Figure 3.6 Number of LTOs from Irish airports 1990-2013	80
Figure 3.7 National LTO data and Eurocontrol LTO data for 2004-2010	81
Figure 3.8 National LTO fuel data and Eurocontrol LTO fuel data for 2004-2010	82
Figure 3.9 National Cruise fuel data and Eurocontrol Cruise fuel data for 2004-2010.....	82
Figure 3.10 Emissions from 1.A.3.b Road Transport 1990-2013	83
Figure 3.10.1(a) Historic passenger car fleet in Irish transport sector, Petrol	85
Figure 3.10.1(b) Historic passenger car fleet in Irish transport sector, Diesel	85
Figure 3.10.2 (a) Historic LDV fleet in Irish transport sector	86
Figure 3.10.2 (b) Historic HDV fleet in Irish transport sector	86
Figure 3.10.3(a) Historic Buses and coaches fleet in Irish transport sector	87
Figure 3.10.3 (b) Historic Mopeds and Motorcycle fleet in Irish transport sector	87
Figure 3.10.4 Average Vehicle mileage for Petrol PC (2000-2013).....	88
Figure 3.10.5 Average Vehicle mileage for Diesel LDV (2008-2013)	89
Figure 3.10.6 Average Vehicle mileage for HD Vehicle categories (2008-2013)	89
Figure 3.10.7 Average vehicle mileage by vehicle type (1990-2013)	90
Figure 3.10.8 Average vehicle mileage for LD and HD by vehicle type (1990-2013)	91
Figure 3.10.9 Average Vehicle mileage for bus and coach industry (1990-2013)	92
Figure 3.10.10 Average mileage for motorcycle (1990-2013)	92
Figure 3.10.11 Vehicle speeds by vehicle type and road type.....	93
Figure 3.11 Emissions from 1.A.4 Other Sectors 1990-2013	99
Figure 3.12 Fugitive emissions from Underground Coal Mines 1990-Methodological Issues	102
Figure 4.1 Total Emissions from IPPU by Category, 1990-2013.....	110
Figure 4.2 Total Emissions from IPPU by Gas, 1990-2013	110
Figure 4.3 Emissions of HFC, PFC, SF ₆ and NF ₃	127
Figure 5.1 Total Emissions from Agriculture by Sector, 1990-2013	141
Figure 5.2 Total Emissions from Agriculture by Gas, 1990-2013.....	141
Figure 6.1 Methodologies and hierarchy of determining land use areas and transitions	162
Figure 6.2 Overview of land use change between 1990 and 2013	163
Figure 6.3.1 Activity data and models used to derive CSC for forest lands	170
Figure 6.3.2 The process involved in deriving the total forest area and afforestation areas since 1990 using the IFORIS database	172
Figure 6.3.3 Overview of the NFI classification system (taken from NFI, 2007a)	172
Figure 6.3.4 The concentric plot design and mapping of individual trees NFI, 2007a	173
Figure 6.3.5 Methodology used to derive harvest information for post-1990 State Forests.....	175

Figure 6.3.6 Procedure used to derive harvest activity data for private forested areas	176
Figure 6.3.7 Procedure to derive activity data for Afforestation Areas after 2006	179
Figure 6.3.8 Schematic Overview of CARBWARE Functionality	181
Figure 6.3.9 Validation of optimised age-class distributions	192
Figure 6.3.10 Adjusted time series for forest category 4.A.1.	194
Figure 6.3.11 The adjusted time series for forest category 4.A.2.....	207
Figure 6.4.1 Long term time series of areas under crops in Ireland since 1847	225
Figure 6.4.2 Spatial pattern of long term consolidation of tillage activities in well define regions of Ireland ...	227
Figure 6.4.3 GIS layers showing attribute data associated with Land Parcel Information System.....	227
Figure 6.4.4 Analysis indicating low rates of conversion of Cropland to permanent Grassland	228
Figure 6.4.5 Spatial distribution of all Cropland land parcels.	231
Figure 6.4.6 Time Series of Cropland from 1990-2013	234
Figure 6.4.7 Estimated area under Apple Orchards 1989-2013	237
Figure 6.4.8 Carbon Stock change in Biomass of Perennial Crops.....	238
Figure 6.4.9 Validation of use of Forest Fires as a scalable proxy for fire events on other Land Uses.....	241
Figure 6.4.10 Validation of the use of Forest Service Statistics as a scalable proxy for all fires in a given year.	241
Figure 6.4.11 Estimate of Cropland area burnt by wildfire	243
Figure 6.4.12 Estimated CH ₄ emissions from wildfires on Croplands	244
Figure 6.4.13 Estimated N ₂ O emissions from wildfires on cropland	244
Figure 6.4.14 Comparison between 2014 and 2015 submissions of estimated total emissions associated with Croplands.....	246
Figure 6.5.1 Plot of original CSO data for Pasture areas and adjusted data for 2010, 2011 and 2012 based on extrapolation of long term trend from 1991 to 2009	249
Figure 6.5.2 Trends in Grassland use 1990-2013.....	251
Figure 6.5.3 Time Series of carbon losses from drained organic soils under grassland	253
Figure 6.5.4 Area of Rewetted Organic Soils under Grassland	254
Figure 6.5.5 Carbon uptake to rewetted drained organic soils under grassland.....	255
Figure 6.5.6 Carbon losses due to DOC emissions associated with rewetting of drained organic soils under grassland.....	256
Figure 6.5.7 Time series of methane emissions associated with rewetting of drained organic soils under grassland.....	257
Figure 6.5.8 shows the estimate of CH ₄ emissions from this analysis.	258
Figure 6.5.9 Estimate of methane emissions due to drainage of organic soils under grasslands	258
Figure 6.5.10 Time Series of conversion of Peat Extraction Sites to Grassland.....	259
Figure 6.5.11 Change in carbon polls due to the conversion of peat extraction sites to Grassland on drained organic soils. Note negative values indicate a loss of carbon to the atmosphere.	260
Figure 6.5.12 The estimated area of grassland subjected to controlled burning from 1990-2013.....	262
Figure 6.5.13 Estimated CH ₄ emissions from wildfires on Grassland	263
Figure 6.5.14 Estimated N ₂ O emissions from wildfires on Grassland.....	263
Figure 6.5.15 Grassland recalculations between the 2014 and 2015 submissions for the period 1990-2012...	265
Figure 6.6.1 N ₂ O emissions associated with the drainage for Peat Extraction.....	269
Figure 6.6.2 Estimated area of peatland drained for Peat Extraction	270
Figure 6.6.3 Estimate of methane emissions due to drainage of organic soils under grasslands	271
Figure 6.6.4 Time Series of Exports of Horticultural Peat products by weight	272
Figure 6.6.5 Off-Site emissions due to use of Horticultural Peat.	273
Figure 6.6.6 Estimate of area of peatland subject to wildfire	274
Figure 6.6.7 Estimated CO ₂ emission due to peatland fires	275
Figure 6.6.8 Estimated CH ₄ emission due to peatland fires.....	276
Figure 6.6.9 Estimated N ₂ O emission due to peatland fires.....	276
Figure 6.6.10 Comparison between 2014 and 2015 submissions of estimated total emissions associated with wetlands	277
Figure 6.7.1 Estimated Area of Settlements 1990 to 2013.....	278
Figure 6.7.2 Comparison between 2014 and 2015 submissions of estimated total emissions associated with Settlements	279
Figure 6.8.1 Recalculations between 2014 and 2015 submissions of estimated total emissions associated with Other Land	281
Figure 7.1 Total Emissions from Waste by Sector, 1990-2013	298

Figure 7.2 Total Emissions from Waste by Gas, 1990-2013	298
Figure 7.3 Methane Emissions from Solid Waste Disposal 1990-2013	301
Figure 10.1 Impact of Recalculations in Energy between annual Submissions 1990-2012	320
Figure 10.2 Impact of Recalculations in IPPU between annual Submissions 1990-2012	321
Figure 10.3 Impact of Recalculations in Agriculture between annual Submissions 1990-2012	321
Figure 10.4 Impact of Recalculations in LULUCF between annual Submissions 1990-2012.....	325
Figure 10.5 Impact of Recalculations in Waste between annual Submissions 1990-2012.....	326
Figure 10.6 Total Impact of Recalculations between annual Submissions 1990-2012.....	331
Figure 11.2.1 Time series of grasslands classified according to GM defintion and Other grassland (unmanaged semi natural).....	353
Figure 11.1 Variation in mineral soil carbon stocks and estimation of ΔC_{so} using the nationally derived data (n = 30).....	355
Figure 11.2 Mean soil C reference values for forest and grasslands at steady state across different mineral soil types	357
Figure 11.3 Comparison of mean SOC values (histograms) and standard deviations (error bars) across different land use categories. Mean values with different alphabetical characters indicate a significant difference (p<0.05)	358
Figure 11.4 Regression analysis showing trends in SOC stocks for different soils types as a function of forest age	359
Figure 11.5 Hedgerow established under a series of rural protection schemes	361

Acknowledgements

The authors wish to express their appreciation to the various Government Departments, Sustainable Energy Authority of Ireland, the Central Statistics Office, Teagasc, Aether (UK), the Electricity Supply Board and the many other data suppliers and individuals who have contributed to the subject matter of this national inventory report.

EXECUTIVE SUMMARY

ES.1 Background

The present report constitutes Ireland's National Inventory Report for 2015 and refers to the greenhouse gas inventory time-series for the years 1990-2013.

This is the first submission of the inventory under the Revision of the UNFCCC Inventory Reporting Guidelines on annual inventories for Parties included in Annex I to the Convention adopted by COP at Warsaw (Decision 24/CP.19). The estimates presented here were estimated in accordance with the guidelines in Annex I of the decision using methodologies provided in the 2006 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories (IPCC 2006) and GWPs listed in table 2.14 of the errata to the contribution of Working Group 1 to the Fourth Assessment Report of the IPCC as contained in Annex III of the decision. The Common Reporting Format (CRF) tables reported in this submission were generated by the CRF Reporter software and submitted via the UNFCCC submission portal and are in accordance with Annex II of the decision. The UNFCCC guidelines require that Parties prepare a National Inventory Report (NIR) as one of the key components of their annual submissions to the UNFCCC secretariat. The purpose of the NIR is to describe the input data, methodologies, emission factors, quality assurance and quality control procedures and other information underlying the inventory compilation for greenhouse gases and to give details of any recalculations of inventories previously submitted. It is needed to assess the transparency, completeness and overall quality of the inventories as part of the rigorous on-going technical review of submissions from Annex I Parties. The structure of this report is consistent with the Appendix in Annex I of Decision 24/CP.19.

Ireland's submission under the UNFCCC in 2015 is **not** its official submission under the Kyoto Protocol, with 2013 being the first year of the second commitment period 2013-2020. According to Decision 13/CP.20 of the Conference of the Parties to the UNFCCC, the CRF Reporter version 5.0.0 was not functioning in order to enable Annex I Parties to submit their CRF tables for the year 2015. CRF reporter version 5.10 still contains issues in the common reporting format tables and XML format in relation to Kyoto Protocol requirements (activities under Article 3, paragraphs 3 and 4), and it is therefore not yet functioning to allow submission of all the information required under Kyoto Protocol. Recalling the Conference of Parties invitation to submit as soon as practically possible, and considering that CRF reporter 5.10 allows sufficiently accurate reporting under the UNFCCC, the present report is the official submission for the year 2015 under the UNFCCC.

The NIR is prepared according to the Appendix in Annex I to Decision 24/CP.19. Part I includes sections describing the national system for inventory preparation and management, emission trends, key emission categories, recalculations and on-going improvements. In addition, detailed documentation of methods, activity data and emission factors used for each of the five source categories, as defined by the Intergovernmental Panel on Climate Change (IPCC), are included. Part II contains the supplementary information required under Article 7, paragraph 1 of the Kyoto Protocol, which refers mainly to the reporting and accounting of emissions and removals for activities under Article 3, paragraph 3 (Afforestation, Reforestation and Deforestation) and Article 3, paragraph 4 (Forest management, Cropland management and Grazing land management). The report contains several annexes, which include calculation sheets, activity data, emission factors and other

appropriate reference material to support the descriptions of inventory estimation methods given in both Part I and Part II and to provide adequate transparency for review purposes, as required by the UNFCCC reporting guidelines.

The Environmental Protection Agency has overall responsibility for the national greenhouse gas inventory in Ireland's national system, which was established in 2007 under Article 5 of the Kyoto Protocol. The EPA Office of Climate, Licensing, Research and Resource Use (OCLRR) performs the role of inventory agency in Ireland and undertakes all aspects of inventory preparation and management as well as the reporting of Ireland's submissions annually in accordance with the requirements Regulation [No. 525/2013](#) of the European Parliament and of the Council and the UNFCCC. In addition to complying with the UNFCCC reporting guidelines, the 2015 NIR is intended to inform Irish Government departments and institutions involved in the national system, as well as other relevant stakeholders in Ireland, of the level of emissions and the state-of-the-art of Irish greenhouse gas inventories. The in-depth analysis of key categories and the up-to-date data on emissions trends provides essential information for the implementation of the National Climate Change Strategy and the development of emissions projections. The detailed NIR, together with activities provided for in the national system, allows data providers to become fully aware of the importance of their contributions to the inventory process and it serves to identify areas where improvements in input data can be achieved.

Ireland's commitment on greenhouse gases under the Kyoto Protocol pursuant to its Article 3, paragraph 9, the Doha Amendment ([1/CMP.8](#)) is set out in Annex B of the protocol. Ireland's quantified emission limitation reduction commitment (QELRCs) for the period 2013 to 2020 is 80 percent of its base year emissions. The QELRCs for the European Union and its Member States for the second commitment period under the Kyoto Protocol are based on the understanding that these will be fulfilled jointly with the European Union and its member States and Iceland, in accordance with Article 4 of the Kyoto Protocol. The legislative agreements setting out joint fulfilment under Article 4 of the Kyoto Protocol between the European Union and its Member States ([Council Decision EU 2015/1339](#)), and the European Union and its Member States and Iceland ([Council Decision EU 2015/1340](#)) were finalised in July 2015.

The European Union's Effort Sharing Decision ([No. 406/2009/EC](#)) established binding annual targets for Member States for the period 2013–2020. These targets cover emissions from most sectors not included in the EU Emissions Trading System (EU ETS), such as transport (except aviation and international maritime shipping), buildings, agriculture and waste. Ireland's binding target is set out in Annex II of the decision and limits emissions to -20 per cent compared to 2005 greenhouse gas levels. Ireland's actual annual emissions allocations (AEAs) for each year of the period 2013 to 2020 are set out in Annex II to Decision [2013/162/EU](#) as adjusted by the amounts in Annex II to Decision [2013/634/EU](#).

ES.2 Summary of National Emission and Removal-related Trends

In 2013, total emissions of greenhouse gases including indirect emissions from solvent use (excluding the *LULUCF* sector) in Ireland were 58,820.88 kt CO₂ equivalent, which is 3.6 per cent higher than emissions in 1990 as presented in Figure ES.1. The total for 2013 is 18.1 per cent lower than the peak of 71,802.64 kt CO₂ equivalent in 2001 when emissions reached a maximum following a period of unprecedented economic growth. The *Energy* sector accounted for 60.8 per cent of total emissions in 2013, *Agriculture* contributed 32.2 per cent while a further 4.7 per cent emanated from

Industrial Processes and Product Use and 2.3 per cent was due to *Waste*. Emissions of CO₂ accounted for 63.1 per cent of the national total in 2013, with CH₄ and N₂O contributing 22.5 per cent and 12.1 per cent, respectively. The combined emissions of HFC, PFC, SF₆ and NF₃ accounted for 2.3 per cent of total emissions in 2013.

Figure ES.1 National total Greenhouse Gas emissions (excluding LULUCF) 1990-2013

An approach 1 level assessment of emission source categories (ranking on the basis of their contribution to total emissions) identified 27 key categories in 2013 (excluding the *LULUCF* sector). There were 17 key categories of CO₂, accounting for 61.4 per cent of total emissions. There were six key categories of CH₄, three key categories of N₂O and 1 key category of HFC in level assessment, which accounted for 21.2 per cent, 10.8 per cent and 1.9 per cent of total emissions, respectively. The results of the Tier 1 key category analysis clearly show the impact of CO₂ emissions from energy consumption on total emissions in Ireland. These combustion sources of CO₂ emissions accounted for 15 out of 27 key categories identified by level assessment in 2013 and for approximately two-thirds (58.6 per cent) of total emissions. The top ten key categories contributed 73.3 per cent of total emissions in 2013 with emissions of CO₂ from the combustion of liquid fuels (petrol and diesel) by road traffic being the single largest source, accounting for 17.8 per cent of the total national emissions.

The application of uncertainty analysis for Irish greenhouse gas inventories using the IPCC approach indicates an overall level uncertainty of 10.2 per cent in the 2013 inventory (excluding the *LULUCF* sector) and a trend uncertainty of 3.2 per cent for the period 1990 to 2013. These values are determined largely by the low uncertainty in the estimates of CO₂ emissions from the energy sector, which is the major source category in Ireland and for which the input data and methodologies are most reliable. The 63.1 per cent of emissions contributed by CO₂ in 2013 are estimated to have an uncertainty of 1.3 per cent. Emissions of CH₄ from 3.B Manure Management and N₂O from 3.D.1 Direct N₂O Emissions from Managed Soils sectors combined account for majority of the level

uncertainty (contributing 3.1 per cent and 9.6 per cent, respectively to each gas uncertainty) in the 2013 inventory. The impact of HFC, PFC, SF₆ and NF₃ on inventory uncertainty in the year 2013 was negligible (0.5 per cent) because they account for only 2.3 per cent of total emissions.

ES.3 Overview of Source and Sink Category Emission Estimates and Trends

Chapter 2 of the NIR describes the trends in Ireland's time-series of greenhouse gas inventories for the years 1990 through 2013. The emissions time-series is available as a complete set of Common Reporting Format (CRF) files, generated by the online CRF Reporter GHG inventory software web application, to be used for annual data submissions to the European Union and the UNFCCC secretariat. The annual inventories are complete with respect to both the coverage of the seven direct greenhouse gases for which information is required and the coverage of the five IPCC source categories. Some recalculations have again been undertaken for the purposes of the 2015 submission and the latest inventories for the years 1990-2013 indicate revisions and improvements in some areas due to these recalculations.

Fuel combustion in the Energy sector is the principal source of emissions in Ireland and major increases in fuel use have driven the increase in emissions in the 1990-2013 time-series. The largest increase took place in transport with an increase of 115.5 per cent on 1990 levels, while there were increases of 6.8 per cent and 1.3 per cent in the emissions from the industrial sectors and energy industries, respectively. The emissions from Agriculture sector, the other main source category, increased during the 1990s but have decreased to 8.5 per cent below 1990 levels in 2013. As the emissions from energy increased, the contribution of agriculture to total national emissions decreased from 36.5 per cent in 1990 to 32.2 per cent in 2013. This is primarily as a result of falling livestock numbers since 1998 due to reform of the Common Agricultural Policy (CAP).

ES.4 Indirect Greenhouse Gases

The inventory reporting process requires the inclusion of a number of gases whose indirect effects are also relevant to the assessment of human-induced impacts on climate. They include sulphur dioxide (SO₂), nitrogen oxides (NO_x), carbon monoxide (CO) and volatile organic compounds (VOC). Emissions of SO₂ contribute to the formation of aerosols, which may offset the effects of greenhouse gases, while CO, NO_x and VOC are precursors of ozone, another naturally occurring greenhouse gas. This NIR does not describe the methods used to estimate emissions of SO₂, NO_x, CO and NMVOC but the annual emissions estimates over the period 1990-2013 are included in the submission.

Indirect CO₂ emissions from NMVOCs from solvent use (category 2.D.3 in the IPPU sector) are included in Ireland's national total for greenhouse gas emissions to be consistent with reporting under the Kyoto Protocol for the first commitment period (previous CRF sector 3, solvent and other product use) and are reported in CRF Table 6.

The emissions of most of the indirect gases have decreased substantially in the period 1990-2013 under various forms of control legislation emanating from the European Commission and the Convention on Long Range Transboundary Air Pollution. The reductions achieved between 1990 and 2013 in Ireland are of the order of 86.2 per cent in the case of SO₂, 64.8 per cent for CO and 41.0 per cent for NO_x and 33.5 per cent for NMVOC.

PART I

ANNUAL INVENTORY SUBMISSION 2015

Chapter 1 Introduction

1.1 Background and Context

This report constitutes Ireland's National Inventory Report (NIR), for the years 1990-2013, as required under the United Nations Framework Convention on Climate Change. Ireland's submission under the UNFCCC in 2015 is **not** to be considered its official submission under the Kyoto Protocol, with 2013 being the first year of the Kyoto commitment period 2013-2020 due to the reporting issues relating to KP LULUCF CRF tables.

The objective of the NIR is to describe the methodologies, input data, background information and the entire process of inventory compilation for greenhouse gases and to give explanations for any improvements and recalculations of the inventories reported in previous submissions. The report is a key component of the UN review process which assesses the transparency, completeness and overall quality of the inventories from Annex I Parties.

1.2 Introduction and Reporting Requirements under the UNFCCC

The United Nations Framework Convention on Climate Change (UNFCCC) (Articles 4 and 12), hereafter referred to as the Convention, requires Annex I Parties to develop, publish and make available to the Conference of the Parties (COP), the Convention's implementation body, their national inventories of emissions and removals of all greenhouse gases not controlled by the Montreal Protocol. The revision of the UNFCCC Reporting Guidelines on annual inventories for Parties included in Annex I to the Convention ([Decision 24/CP.19](#)), hereafter referred to as the UNFCCC reporting guidelines, describe the scope and reporting of the emissions inventories. They specify the methodologies and procedures to be followed for submitting consistent and comparable data on an annual basis in a timely, efficient and transparent manner to meet the needs of the Convention. Under the UNFCCC reporting guidelines, Parties are required to compile a National Inventory Report (NIR) and up-to-date annual inventories in an electronic Common Reporting Format (CRF) as the key components of their annual submissions.

The NIR is compiled according to the structure adopted by the Appendix to Annex I of Decision 24/CP.19.

- Part I includes sections describing the national system for inventory preparation and management, emission trends, key emission categories, recalculations and on-going improvements. In addition, detailed documentation of methods, activity data and emission factors used for each of the five source categories as defined by the Intergovernmental Panel on Climate Change (IPCC) is provided.
- Part II contains the supplementary information required under Article 7, paragraph 1 of the Kyoto Protocol, which refers mainly to the reporting and accounting of emissions and removals for activities under Article 3, paragraph 3 (Afforestation, Reforestation and Deforestation) and Article 3, paragraph 4 (Forest management, Cropland management and Grazing land management), i.e. emissions by sources and removals by sinks of GHGs resulting from LULUCF activities.

The NIR addresses the full range of reporting requirements related to annual inventories set down in the UNFCCC reporting guidelines and responds to issues identified in the UNFCCC annual review process. Furthermore, the report captures the cyclical nature of the reporting process and clarifies the chronology of changes and revisions that are part of normal inventory development, including those that are implemented in response to the UNFCCC review process. In this way, the report continues to improve the basis for technical assessment and expert review of Irish greenhouse gas inventories. An attempt has been made to provide all the primary inventory information, including calculation sheets as appropriate, to facilitate replication of the emission estimates for the most recent year of the inventory time-series so that the annual submission is fully transparent.

In addition to complying with the UNFCCC reporting guidelines, the report is intended to inform Government Departments, national institutions and other stakeholders of the state of the art of Irish greenhouse gas inventories as they address the challenges to comply with commitments under the Kyoto Protocol. In this context, it provides some additional background on relevant emission sources in Ireland, the standard reporting format and other issues for the benefit of those not entirely familiar with the agreed content of the NIR or the general reporting requirements under the Convention and the Kyoto Protocol. The report is also aimed at all the key data providers, with a view to making them fully aware of the importance of their contributions to the inventory process and to provide a means of identifying areas where improvements in input data may be possible. The in-depth analysis of key categories and the up-to-date data on emissions trends provides essential information for the implementation of the National Climate Change Strategy and the development of emissions projections.

The NIR is updated annually in accordance with the UNFCCC guidelines and is published on the web site of the EPA [<http://erc.epa.ie/ghg/nirdownloads.jsp>]. Such updating is necessary to keep the UNFCCC secretariat and other interested parties informed of the status of Irish greenhouse gas inventories and to document on-going improvements, recalculations and other developments affecting the estimates of emissions. The structure of the report is designed to facilitate year-on-year revision in a manner that allows for systematic and efficient assessment of progress towards the achievement of greenhouse gas emission inventories that meet the guiding principles of transparency, consistency, comparability, completeness and accuracy.

1.2.1 Scope of Greenhouse Gas Inventories

1.2.1.1 Greenhouse Gases and Global Warming Potential

The full range of greenhouse gases for which emissions data are required under the Convention is given in Table 5.4.1 of Annex 5.4. It includes carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O), the most widely known and most ubiquitous of the anthropogenic greenhouse gases, along with 19 hydrofluorocarbons (HFC), 9 perfluorocarbons (PFC), sulphur hexafluoride (SF₆) and nitrogen trifluoride (NF₃). The global warming potentials (GWPs) of the various greenhouse gases vary greatly, and are as listed in table 2.14 of the errata to the contribution of Working Group 1 to the Fourth Assessment Report of the IPCC as contained in Annex III of the decision 24/CP.19. The GWP of a gas is a measure of the cumulative warming over a specified time period, e.g. 100 years, resulting from a unit mass of the gas emitted at the beginning of that time period, expressed relative to an absolute GWP of 1 for the reference gas carbon dioxide (IUCC, 1998). The mass emission of any gas multiplied by its GWP gives the equivalent emission of the gas as carbon dioxide. Therefore, while CO₂, CH₄ and N₂O are important because they are normally emitted in large amounts, HFCs, PFCs, SF₆ and NF₃ are included in the inventory process mainly because of their comparatively much larger GWP values.

The inventory reporting process allows for the inclusion of a number of additional gases whose indirect effects are also relevant to the assessment of human-induced impacts on climate. These include sulphur dioxide (SO₂), nitrogen oxides (NO_x), carbon monoxide (CO) and non-methane volatile organic compounds (NMVOC). Emissions of SO₂ contribute to the formation of aerosols, which may offset the effects of greenhouse gases, while CO, NO_x and NMVOC are precursors of ozone formation, another naturally occurring greenhouse gas. This NIR does not describe the methods used to estimate emissions of SO₂, NO_x, CO and NMVOC but up-to-date estimates of total emissions are included for information purposes. These estimates are taken from Ireland's submission to the Convention on Long Range Transboundary Air Pollution (CLRTAP), which are produced annually in a manner that is fully consistent with the inventory for greenhouse gases.

1.2.1.2 Common Reporting Format

Greenhouse gas emissions are reported under the Convention in a multi-level reporting format adopted by the Intergovernmental Panel on Climate Change (IPCC). This is a standard table format that forms the basis of the new Common Reporting Format (CRF), Annex II to the UNFCCC reporting guidelines, which assigns all potential sources of emissions and removals of a Party's national total to five Level 1 broad source categories. A further category is provided for the reporting of any additional sources that may be specific to individual Parties. Table 5.4.2 of Annex 5.4 lists the Level 1 and Level 2 source/sink categories. Level 2 source/sink categories are further sub divided to a finer level of disaggregation, level 3. The Level 3 categories are detailed in the description of category coverage and inventory methods and data in the respective sectoral chapters of this NIR. The computation of emissions is usually undertaken at Level 3 or lower, using further appropriate disaggregation (for example, by using fuel type in the case of combustion sources under 1.A Energy-Fuel Combustion) while summary results are normally published at Level 2.

The reporting format also accommodates the reporting of emissions and removals under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (i.e. emissions by sources and removals by sinks of GHGs resulting from KP LULUCF activities) for the years 2013-2020 of the second commitment period (CP2). The additional tables use a hierarchical system similar to that for reporting under the Convention, with flexibility for Parties to provide as much disaggregation as is necessary to reflect the variation in the parameters underlying the estimates of emissions and removals for the Article 3, paragraphs 3 and 4, activities applicable in their territories. The Kyoto reporting tables also include the accounting quantity for each relevant activity i.e. the quantity of units to be added or subtracted from a Party's assigned amount in accordance with the provisions of Article 7, paragraph 4, of the Protocol. This submission does not include the KP LULUCF CRF tables due to the erroneous reporting by the CRF Reporter software when generating these tables.

The IPCC reporting format also includes a number of Memo Item entries. These items refer to sources of emissions whose contributions are not included in a Party's national total but which are to be reported because of their importance in relation to the overall assessment of emissions and for comparisons among Parties.

The national total of emissions that is commonly used under the Convention excludes the estimates for the Land Use Land-Use Change and Forestry (LULUCF) sector in Table 5.4.2 of Annex 5.4, this total being consistent with that for the categories included in Annex A of the Kyoto Protocol. Ireland's national total during the second commitment period also includes indirect CO₂ emissions from NMVOCs from solvent use (category 2.D.3 in the IPPU sector) to be consistent with reporting

under the Kyoto Protocol for the first commitment period (previous CRF sector 3, solvent and other product use) and are reported in CRF Table 6.

1.2.1.3 Supplementary Information

For a Party to the Kyoto Protocol, the annual inventory submission under the Convention is also its annual inventory submission under the Protocol. Supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol comprises the GHG emissions and removals under Article 3, paragraphs 3 and 4, of the Kyoto Protocol, details of all Kyoto units for the year subsequent to the inventory year as generated by the national registry and compiled in the Standard Electronic Format, changes in the national system and national registry and information on the minimization of adverse impacts of climate change and response measures on developing countries in accordance with Article 3, paragraph 14. This submission is not considered to be an official submission under the Kyoto Protocol and as such Part II of this report is provided for information purposes only.

1.3 National Inventory Arrangements

1.3.1 Institutional, Legal and Procedural Arrangements

The Environmental Protection Agency is required to establish and maintain databases of information on the environment and to disseminate such information to interested parties (Section 52 of the Environmental Protection Agency Act of 1992 (DOE, 1992)). The Act states that the Agency must provide, of its own volition or upon request, information and advice to Ministers of the Government in the performance of their duties (Section 55). This includes making available such data and materials as are necessary to comply with Ireland's reporting obligations and commitments within the framework of international agreements. These requirements are the regulatory basis on which the EPA prepares annual inventories of greenhouse gases and other important emissions to air in Ireland. It is in this context that in 1995 the Department of the Environment, Community and Local Government (DECLG) designated the EPA as the inventory agency with responsibility for the submission of emissions data to the UNFCCC Secretariat and to the Secretariat for the Convention on Long-Range Transboundary Air Pollution (CLRTAP).

The establishment of Ireland's national inventory system was completed by Government Decision in early 2007, building on the framework that had been applied for many years. The EPA's Office of Climate, Licensing, Resource and Research (OCLRR) was designated the inventory agency and the EPA was also designated as the single national entity with overall responsibility for the annual greenhouse gas inventory. Within the OCLRR, the Climate Resource and Research Programme (CRRP), compiles the national greenhouse gas emission inventories for submission on behalf of DECLG under the Framework Convention on Climate Change and Regulation (EU) 525/2013, the latter being the basis for EU Member States' reporting under the Convention and the Kyoto Protocol. All formal mechanisms together with the QA/QC procedures are fully operational since they were established in the 2007 reporting cycle.

Following establishment of the national system, institutional arrangements directed towards national inventory reporting that involve the EPA, DECLG and other stakeholders were reorganised, extended and legally consolidated across all participating institutions to strengthen inventory capacity within the EPA. This ensured that more formal and comprehensive mechanisms of data collection and processing were established and maintained for long term implementation. In particular, the system puts in place formal procedures for the planning, preparation and

management of the national atmospheric inventory and identifies the roles and responsibilities of all the organisations involved in its compilation. This was achieved through extensive discussions with all key data providers leading to the adoption of Memoranda of Understanding (MOU) between the key data providers and the inventory agency. These MOUs stipulate the scope, timing and quality of the inputs necessary for inventory compilation in accordance with the guidelines for national systems. Secondary MOUs are, in turn, used by some key data providers to formalise the receipt of data from their own particular sources. Table 1.1 lists the key data providers and indicates the range of data covered by MOU in the national system. A QA/QC plan is an integral part of the national system.

Figure 1.1 provides a schematic overview of the institutions, procedures and information flows involved in the national system. In addition to the primary data received from the key data providers, the inventory team draws on various other data streams available within the EPA, such as the National Waste Database, reports on wastewater treatment, Annual Environmental Reports from companies subject to Integrated Pollution Prevention Control and submissions prepared under the European Pollutant Release and Transfer Register and also obtains information from other diverse sources to prepare the inventories for fluorinated gases and solvent use. The inventory team also draws on national research related to greenhouse gas emissions and special studies undertaken from time to time to acquire the information needed to improve the estimates for particular categories and gases.

The Emissions Trading Unit (ETU), also within the Climate Resource and Research Programme of the OCLRR, is a key component of the national system. The ETU are responsible for administering the European Union Emissions Trading Scheme (ETS), under Directive 2003/87/EC (EP and CEU, 2003), in Ireland and, as such, provide annual verified emissions data to the inventory team.

The estimates of emissions and removals for forest lands under the Convention, as well as those in respect of Article 3, paragraph 3, activities under the Kyoto Protocol, are prepared by consultants contracted to the Department of Agriculture, Food and Marine (DAFM). These are delivered to the inventory agency under a Memorandum of Understanding between DAFM and OCLRR. A research fellow contracted directly to OCLRR is responsible for completion of the annual inventory for all other land categories in LULUCF for the annual inventory under the Convention and elected activities under Article 3, paragraph 4, of the Kyoto Protocol (Cropland management and Grazing land management). The deliverables received by OCLRR from DAFM and the research fellow include the completed CRF tables and draft NIR sections for their respective areas of responsibility.

The approval of the completed annual inventory involves sign-off by the QA/QC manager and the inventory manager before it is transmitted to the Board of the EPA via the Programme Manager of the Climate Resource and Research Programme in OCLRR. Any issues arising from the Board's examination of the estimates are communicated to the inventory experts for resolution before final adoption of the inventory. The results for the inventory year are normally released at national level in autumn of the following year. This is in advance of their official submission to the European Commission in accordance with Regulation (EU) 525/2013 in January of the reporting year and subsequently to the UNFCCC secretariat.

The national system is also exploited for the purpose of parallel inventory preparation and reporting under the LRTAP Convention ensuring efficiency and consistency in the compilation of emission inventories for a wide range of substances using common datasets and inputs.




Figure 1.1 National Inventory System Overview

1.3.2 Overview of Inventory Planning, Preparation and Management

The inventory agency plans for preparation of the annual inventory as soon as possible after completion of the annual reporting cycle in April following submission to the UNFCCC secretariat. Planning largely involves the identification of improvements to be undertaken by way of revised methodologies and updated activity data or emission factors as well as addressing the issues and recommendations in the review of the previous inventory submission.

Planning also considers the further development of inventory reporting for the LULUCF sector and for Kyoto Protocol activities under Article 3, paragraphs 3 and 4, as new data becomes available through national research and development of the forest inventory.

In addition, any changes required by the outcome of review activities conducted among the Member States of the European Union, or by the need to report in a manner consistent with other Member States for the purposes of Regulation (EU) 525/2013, are taken into account in inventory planning.

The first version of the latest annual inventory, produced in autumn of the following year, and a short National Inventory Report are used to comply with the subsequent 15 January deadline prescribed by Regulation (EU) 525/2013, which governs the reporting of greenhouse gases and implementation of the Kyoto Protocol by the European Union and its EU Member States.

The inventory preparation and management process thereafter involves making any revisions subsequent to the receipt of updated or outstanding information nationally. In addition, any observations or amendments following initial assessment at EU level of the 15 January submission by Member States to the European Commission are incorporated into the inventory between 15 January and 15 March.

The complete and final inventory submission, including the National Inventory Report, is submitted to the European Commission by 15 March as required under Regulation (EU) 525/2013. This version of the latest inventory is fixed and retained for submission to the UNFCCC secretariat by 15 April to complete the reporting cycle. Ireland's national system is operating very successfully and the timeliness of inventory preparation has benefited from the implementation of more formal arrangements and enhanced engagement among the various institutions and contributors.

1.3.3 Quality Assurance, Quality Control and Verification Plan

In early 2005, the inventory agency in Ireland commissioned a project with UK consultants NETCEN to establish formal QA/QC procedures that would meet the needs of the UNFCCC reporting requirements. The project developed a QA/QC system including a documented QA/QC plan and procedures along with a QA/QC manual.

The manual provides a general overview of the QA/QC system. In addition, the manual provides guidance and templates for appropriate quality checking, documentation and traceability. The selection of source data, calculation methodologies, peer and expert review of inventory data and the annual requirements for continuous improvement for the inventory are also outlined in the manual.

The QA/QC plan identifies the specific data quality objectives related to the principles of transparency, consistency, completeness, comparability and accuracy required for Ireland's national inventory and provides specific guidance and documentation forms and templates for the practical implementation of QA/QC procedures. The QA/QC procedures cover such elements as data selection and acquisition, data processing and reporting.

The inventory agency initiated a new approach to QA/QC in the 2006 reporting cycle. Its application was completed and consolidated in delivering the submissions up to 2012. This involved the allocation of responsibilities linked to the national system mentioned in section 1.2.1 and the use of a template spread sheet system to record the establishment and maintenance of general inventory checking and management activities covering the overall compilation process, as well as the undertaking of specific annual activities and any necessary periodic activities in response to specific events or outcomes in inventory reporting and review. The system facilitates record keeping related to the chain of activities from data capture, through emissions calculations and checking, to archiving and the identification of improvements. The system has been carried forward for use in completing the 2015 submission.

Ireland's calculation spread sheets in all sectors are structured and organised to facilitate the QA/QC process and more efficient time-series analysis and also to ensure ease of transfer of the outputs to the CRF Reporter Tool. This facilitates rapid year-on-year extension of the time-series, rapid inter-annual comparisons and efficient updating and recalculation, where appropriate, in the annual reporting cycle. Internal aggregation to various levels corresponding to the CRF tables provides immediate and complete checks on the results.

External reviews of the agriculture sector and of the entire ETS results for 2005 were conducted as important new components of quality assurance at the beginning of 2007. The review for the agriculture sector was performed by a Technical Inspector in the Department of Agriculture, Food and Marine. This review used the new calculation files to assess the consistency of the time series which had been subject to considerable improvement and recalculation in the 2006 reporting cycle. These improvements and recalculations were part of a move to higher tier methods for enteric fermentation in cattle as well as advice from the Department on various aspects of input data and calculation parameters. A detailed bilateral review with UK agricultural experts took place in the offices of the EPA in July 2014 to review, in particular, the changes to the agriculture inventory with respect to the use of the 2006 IPCC guidelines in this reporting submission. The inventory agency also continues to work closely with the Department of Agriculture, Food and Marine and seeks advice and guidance from experts in Teagasc, an Agency of the Department of Agriculture, Food and Marine.

The inventory team contracted an external service provider, Aether, to assist in aspects of inventory compilation in 2013 and subsequent submissions. The transparency, robustness and accessibility of the inventory data within the electronic filing structures were assessed by Aether, who concluded that the system is very well organised.

The ETS returns to the ETU provide for the complete coverage of CO₂ estimates in a number of sub-categories under 1.A.1 Energy Industries and 2.A. Mineral Products. When the allocation to these categories from the ETS raw data is completed, the output is returned to the ETS administrator in OCLRR for final checking against the source data. This ensures the efficient and consistent transfer of the verified ETS emissions estimates into the national inventory. Inventory development continues to benefit from the internal review procedures that are on-going with regard to the EU and its Member States. In 2014, experts from the inventory team attended 2 workshops, in March and June, organised by UBA Germany and the European Commission to facilitate the implementation of the 2006 IPCC guidelines for inventory reporting for this first submission for the second commitment period.

1.3.4 Changes in the National Inventory Arrangements since Previous Annual GHG Inventory Submission

There has been no change in the national inventory arrangements since the previous annual inventory submission in April 2014.

1.4 Inventory Preparation, and Data Collection, Processing and Storage

1.4.1 GHG Inventory and KP-LULUCF Inventory

An emissions inventory database normally contains information on measured emission quantities, activity statistics (populations, fuel consumption, vehicle/kilometres of travel, industrial production and land areas), emission factors and the associated emission estimates for a specified list of source categories. In practice, very few measured data are available for greenhouse gases and, consequently, the emissions from most activities are estimated by applying emission factors for each source/gas combination to appropriate activity data for the activity concerned. Virtually all emissions and removals estimates may be ultimately derived on the basis of such simple product of activity data and emission factor. However, a certain amount of data analysis and preparatory calculations are generally needed in order to make available suitable combinations of activity data and emission factors at the level of disaggregation that gives the best estimates of emissions and removals. In the case of some source/gas combinations, such as methane emissions from solid waste landfills and CO₂ sequestration by forest biomass, it may be necessary to apply sophisticated models to generate the activity data, the emission factors or the emissions. The methods recommended by 2006 IPCC Guidelines for national greenhouse gas inventories use a tier system to take account of these issues and other factors, such as data availability, technical expertise, inventory capacity and other circumstances, which may vary considerably across countries.

1.4.2 Data Collection, Processing and Storage

Preparation for the annual GHG inventory takes place in an Excel spread sheet system where activity data stored in Source Data files are linked to calculation sheets in Data Processing files that produce the emissions estimates at the lowest possible level of disaggregation. These are combined and allocated according to IPCC requirements for direct transmission into the CRF Reporter online application for the generation of the CRF tables and Party submissions. These results are stored in Outputs files while supporting QA/QC sheets, extracted from Data Processing files, are held in summary QA/QC record files. The Data Processing files hold the emission factors and they are structured on a time-series basis, which facilitates efficient recalculation and output to the CRF Reporter. This procedure applies to all IPCC sectors of the GHG inventory for which the calculations are made by the inventory team and the full set of files applicable to each year under the four headings is stored using appropriate version control on the OCLRR servers.

Table 1.1 lists the principal data suppliers and the information that they are required to deliver to the inventory agency annually under MOU for the preparation of the GHG inventory. In some cases, e.g. the national energy balance, the input file received from the data supplier may be linked directly to the Data Processing files, but generally some degree of preparation and pre-processing is needed before the activity data are used in inventory preparation. In collating and compiling the activity data, the inventory team collects data from the various data streams e.g. Annual Emissions Reports (AERs) under the European Pollutant Release and Transfer Register.

Table 1.1 Key Data Providers and Information covered by MOU

Key Data Provider	Data Supplied	Deadline	Sector in which data are used
Sustainable Energy Authority of Ireland	National Energy Balance; Detailed national energy consumption disaggregated by economic sector and fuel	30 September	Energy, Waste
Department of Agriculture, Food and Marine	Table 1.1-1.4 Statistical data for cattle compiled under the Animal Identification and Movement (AIM) scheme Fertiliser and lime statistics Poultry statistics Sheep statistics	30 September	Agriculture
Department of Agriculture, Food and Marine (Forest Sector Development Division)	Table 2.1 GHG emission/removal estimates from all pools for forest lands under the Convention Statistical data on Afforestation, Reforestation, Deforestation and harvesting for forest land lands under Article 3.3 of KP GHG emission/removal estimates from all biomass pools for KP Article 3.3 and elected activities under Article 3, paragraph 4, of the Kyoto Protocol (Cropland management and Grazing land management).	30 September	LULUCF and Article 3.3 and 3.4 of the Kyoto Protocol
Central Statistics Office	Annual population, livestock populations, crop statistics, housing survey data	30 September	Agriculture, IPPU, Waste
Bord Gais	Analysis results for indigenous and imported natural gas	30 September	Energy
Marine Institute	Annual Report on Discharges, Spills and Emissions from Offshore Gas Production Installations	30 October	Energy
Emissions Trading Unit	Verified CO ₂ estimates and related fuel and production data for installations covered by the EU ETS ¹	30 April	Energy, IPPU
*Department of Communications, Energy and Natural Resources	National Oil Balance (as a component of the Energy Balance)	30 September	Energy
*Road Safety Authority	Road transport statistics from the National Car Test (NCT)	30 April	Energy
**Forest Service	(i) GIS data base on premiums and grants afforestation areas (iFORIS) with associated attributes (ii) NFI database	30 September 2007, 2012	LULUCF and Article 3.3 & 3.4 activities
**Coillte	GIS data base of intersected of NFI permanent sample plot points (Coillte-NFI plots) with sub-compartment and management unit data.	30 September	LULUCF and Article 3.3 & 3.4 activities

¹ETS – Emissions Trading Scheme

*These bodies have MOUs with SEAI rather than with OCLRR

**These bodies have MOUs with the Department of Agriculture, Food and Marine rather than with OCLRR

A national model called CARBWARE is used to derive the estimates of emissions and removals for forest lands, which are incorporated in the overall scheme for LULUCF reporting under the Convention following the procedure outlined above. A variety of databases related to land cover, soil type and forest areas are applied for the LULUCF inventory under the Convention. These include the National Forest Inventory (NFI), the Forest Inventory and Planning System (FIPS), the Land Parcels

Information System (LPIS), CORINE Land Cover Maps and the General Soil Map of Ireland. These are supported by statistical information from Bord na Móna and the National Roads Authority.

The static national model, CARBWARE has been extensively developed to a dynamic version to provide the necessary estimates for Article 3, paragraph 3 and paragraph 4, selected activities under the Kyoto Protocol. This work was undertaken by FERs Ltd, the consultants working to Department of Agriculture, Food and Marine, who supply the estimates from these activities to OCLRR under an agreed MOU (Table 1.1). Secondary MOUs between Department of Agriculture, Food and Marine and its data suppliers formalise annual data collection for this area of the inventory. The model contains a multitude of component modules needed to produce estimates of the carbon stock changes for the various carbon pools under afforestation and deforestation areas and for reporting any relevant emissions of CH₄ and N₂O. The model processes detailed spatially explicit data on forest species and soil type obtained from the NFI, FIPS, soils maps, supported by the Grants and Premiums Administration System (GPAS), and felling license records. The model uses complex pre-processing functions, growth models, allometric equations and pool allocation and transfers to produce the results required for Article 3, paragraph 3 and paragraph 4, selected activities.

The annual ETS compilation serves as an important source of activity-specific and company-specific data on CO₂ emissions, fuel use and emission factors for major combustion sources and industrial processes. The emissions trading scheme covers approximately 100 installations in Ireland with combined CO₂ emissions of 15,685.38 kt in 2013, accounting for 26.7 per cent of total greenhouse gas emissions (58,820.88 kt CO₂ equivalent). Guidance provided under the associated Decision 2004/156/EC (EP and CEU, 2004) on methodologies for estimating and reporting greenhouse gas emissions to support Directive 2003/87/EC, together with monitoring and verification mechanisms administered by the ETU, consolidates and improves the information in relation to a substantial proportion of CO₂ emissions for the purposes of reporting national GHG inventories under the Convention and the Protocol.

All of the data used in the compilation of the national GHG inventory submission is stored on an EPA data server located in the Monaghan Regional Inspectorate of the EPA where key staff involved in inventory compilation is located. All background data for recent years are available in electronic format, with a transparent file structure. All data (emission estimates, AD, inventory submissions, references, QA/QC) on the data server are backed up daily.

1.5 Methodologies and Emission Factors

Table 1.4 and Table 1.5 present summaries of the methodologies and emission factors used by Ireland to estimate GHG emissions reported for the years 1990-2013. More than 80 per cent of the total emissions (excluding LULUCF) are covered by Tier 2 methods or higher in Ireland's GHG inventory under the Convention and a Tier 3 model is applied for carbon stock changes for Article 3, paragraph 3 and paragraph 4, activities under the Kyoto Protocol.

1.5.1 Carbon dioxide (CO₂)

Tier 2 or Tier 3 methods are used for the majority of CO₂ combustion source categories and country-specific emission factors are used for all fuels. Even for those combustion categories where data limitations dictate the use of Tier 1 methods, such as 1.A.2 Manufacturing Industries and Construction and 1.A.4 Other Sectors, the CO₂ emissions obtained using the energy balance fuel data and country-specific emission factors are reliable. Tier 2 methods also apply to important process

sources of CO₂ emissions, such as cement and lime production, where country-specific circumstances are again taken fully into account.

The national model used to estimate carbon stock change in the various carbon pools for forest lands in respect of both Convention reporting and Article 3, paragraph 3 and paragraph 4, activities is a Tier 3 methodology. The methods for CO₂ in other LULUCF categories and for relevant CH₄ and N₂O emissions in this sector are invariably Tier 1.

1.5.2 Methane (CH₄)

Ireland's national circumstances are well captured in the Tier 2 methods applied for the major sources of CH₄ in the inventory, which are enteric fermentation and manure management associated with cattle and the CH₄ emissions from solid waste disposal sites.

Tier 2 and Tier 3 methods are used for CH₄ emissions from 1.A.1 Energy Industries and 1.A.3.b Road Transport, respectively, while Tier 1 methods and IPCC default emission factors are used for other CH₄ emissions.

1.5.3 Nitrous oxide (N₂O)

Ireland relies on the simplified IPCC Tier 1 methodologies and default emission factors to estimate all N₂O emissions in agriculture, which is the main source of N₂O in the inventory.

Tier 2 and Tier 3 methods are used for N₂O emissions from 1.A.1 Energy Industries and 1.A.3.b Road Transport, respectively, while Tier 1 methods and IPCC default emission factors are used for other N₂O emissions.

1.6 Overview of Key Categories

The 2006 IPCC guidelines defines a key category as one that is prioritised within the national inventory system because its estimate has a significant influence on the Party's total inventory of greenhouse gases in terms of the absolute level of emissions and removals, the trend in emissions and removals or uncertainty in emissions or removals. Information about key categories is considered to be crucial to the choice of methodology for individual sources and to the management and reduction of overall inventory uncertainty. The identification of such categories is recommended in order that inventory agencies can give them priority in the preparation of annual inventories, especially in cases where resources may be limited. Information on key categories is clearly also vital for the development of policies and measures for emissions reduction. The 2006 IPCC guidelines provide two approaches for undertaking the analysis of key categories that can be applied at any appropriate level of source aggregation, depending on the information available. The simplest approach, approach 1, is again used for 2013 data to further highlight which sources of emissions are the most important in Ireland. This approach identifies key categories using a pre-determined cumulative emissions threshold. Key categories are those that, when summed together in descending order of magnitude, add up to 95 percent of the total level.

The 2006 IPCC guidelines encourage inventory agencies to use approach 2 for its key category analysis, and this has also been suggested in previous annual inventory review reports. In response to this, initial work on using approach 2 was carried out, which highlighted differences between the level of disaggregation found in the approach 1 key category analysis compared to the approach 1 uncertainty assessment. Some sub-categories are reported at a more detailed level in the key category analysis compared to the Uncertainty Analysis (such as transport). Due to resource

constraints, it was not possible to complete this work for this year's submission so the finalisation of the approach 2 key category analysis and the further disaggregation of the approach 1 uncertainty assessment are planned improvements for the 2016 submission.

1.6.1 Key Categories at IPCC Level 2

As inventories of CO₂, CH₄ and N₂O were developed in Ireland during the 1990s, it was quickly established that CO₂ emissions from fuel combustion was by far the largest contributor to the combined national total for these three primary greenhouse gases. It was also evident that CH₄ emissions produced by large cattle herds and the N₂O emissions from agricultural soils, associated with intensive farming practices and large inputs of nitrogen to agricultural soils, were also major sources, even if the estimates were more uncertain than those for CO₂. A preliminary estimate of key categories is therefore provided by considering the emissions aggregated at the IPCC Level 2 source category classification, which clearly indicates the importance of CO₂ emissions from fuel combustion and CH₄ and N₂O emissions from agriculture.

The results at the IPCC Level 2 source category classification may be readily drawn from the CRF table Summary 2. Those for 1990 and 2013 are shown in Table 1.2 and Table 1.3, respectively. It can be seen that there are seven highly significant key categories of emissions in Ireland in the 1990-2013 trend including; CO₂ combustion sources in 1.A.1 Energy Industries, 1.A.2 Manufacturing Industries and Construction 1.A.3 Transport and 1.A.4 Other Sectors, along with the CH₄ emissions from categories 3.A Enteric Fermentation and 3.B Manure Management and N₂O emissions from 3.D Agricultural Soils. These seven categories accounted for 88.1 per cent and 90.2 per cent of total emissions in 1990 and 2013, respectively. In the case of 2013 emissions, three additional Level 2 source categories are needed to reach the cumulative 95 per cent threshold that defines key categories: 2.F.1 Refrigeration and air-conditioning with HFC emissions, 2.A.1 Cement Production with CO₂ emissions and 5.A Solid Waste Disposal with CH₄ emissions. Category 2.F.1 is key in 2013 level analysis and not in 1990, whereas categories 2.B.1 and 2.B.2 are key in 1990 level analysis and not in 2013. The increase in the contribution of CO₂ emissions from category 1.A.3 Transport from 8.8 per cent in 1990 to 18.6 per cent in 2013 is notable, along with the corresponding reductions in the contributions from the three categories: two (3.A and 3.D) in Agriculture and (1.A.4) in Energy. This simple analysis of key categories continues to prove useful to the formulation of mitigation strategies and for prioritising work on inventories in Ireland.

When LULUCF is accounted for in the Level 2 analysis, CO₂ emissions in four LULUCF categories (4.A Forest land, 4.B Cropland, 4.C. Grassland, 4.D Wetlands) become key categories in 1990, and the same three categories and associated gas (less CO₂ from 4.B Cropland), plus CO₂ emissions in LULUCF category 4.G Harvested wood products, become key categories in 2013.

Table 1.2 Key Categories at IPCC Level 2 in 1990

IPCC Level 2 Source Category		GHG	Emissions in 1990 (kt CO ₂ eq)	1990 Level Assessment (%)	Cumulative Total of Level (%)
3.A	Enteric Fermentation	CH ₄	11,356.97	20.01	20.01
1.A.1	Energy Industries	CO ₂	11,145.01	19.64	39.65
1.A.4	Other Sectors(Comm/Resid/Agric)	CO ₂	10,030.94	17.68	57.33
3.D.	Agricultural Soils	N ₂ O	7,140.60	12.58	69.91
1.A.3	Transport	CO ₂	5,021.69	8.85	78.76
1.A.2	Manufacturing Industries and Construction	CO ₂	3,942.64	6.95	85.70
5.A	Solid Waste Disposal	CH ₄	1,396.49	2.46	88.16
3.B	Manure Management	CH ₄	1,342.29	2.37	90.53
2.B.2	Nitric Acid Production*	N ₂ O	995.32	1.75	92.28
2.B.1	Ammonia Production*	CO ₂	990.23	1.74	94.03
2.A.1	Cement Production	CO ₂	884.00	1.56	95.59

* Nitric acid and Ammonia plants ceased operation in 2002 and 2001, respectively

Table 1.3 Key Categories at IPCC Level 2 in 2013

IPCC Level 2 Source Category		GHG	Emissions in 2013 (kt CO ₂ eq)	2013 Level Assessment (%)	Cumulative Total of Level (%)
1.A.1	Energy Industries	CO ₂	11,239.25	19.11	19.11
1.A.3	Transport	CO ₂	10,935.10	18.59	37.70
3.A	Enteric Fermentation	CH ₄	10,553.71	17.94	55.64
1.A.4	Other Sectors(Comm/Resid/Agric)	CO ₂	8,755.82	14.89	70.53
3.D.	Agricultural Soils	N ₂ O	6,113.27	10.39	80.92
1.A.2	Manufacturing Industries and Construction	CO ₂	4,209.13	7.16	88.07
3.B	Manure Management	CH ₄	1,242.33	2.11	90.19
2.F.1	Refrigeration and air-con (incl. MAC)	HFC	1,124.16	1.91	92.10
2.A.1	Cement Production	CO ₂	1,111.75	1.89	93.99
5.A	Solid Waste Disposal	CH ₄	1,106.03	1.88	95.87

Table 1.4 Summary of Methods

IPCC SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	NF ₃
1. Energy	T1,T2,T3	T1,T2,T3	T1,T2,T3				
A. Fuel Combustion (Sectoral Approach)	T1,T2,T3	T1,T2,T3	T1,T2,T3				
1. Energy Industries	T1, T3	T1, T2	T1, T2				
2. Manufacturing Industries and Construction	T1, T2, T3	T1	T1				
3. Transport	T2, T3	T1, T2, T3	T1, T2, T3				
4. Other Sectors	T1, T2	T1	T1				
5. Other							
B. Fugitive Emissions from Fuels	NA	T1, T2	NA				
1. Solid Fuels	NA	T1	NA				
2. Oil and Natural Gas	NA	T1, T2	NA				
C. Carbon Dioxide Transport and Storage	NA						
2. Industrial Processes and Product Use	T1,T3	NA	D	T1, T2, T3	T2	T2	T2
A. Mineral Industry	T3	NA	NA	NA	NA	NA	NA
B. Chemical Industry	NA	NA	NA				
C. Metal Production	NA	NA	NA				
D. Non-Energy Products from Fuels and Solvent Use	T1	NA	NA				
E. Electronic Industry	NA	NA	NA	T2	T2	T2	T2
F. Product Uses as Substitutes for ODS	NA	NA	NA	T1, T2, T3	NA	NA	NA
G. Other Product Manufacture and Use			T1			T1	
H. Other							
3. Agriculture	T1	T1,T2	T1,T2				
A. Enteric Fermentation		T1, T2	NA				
B. Manure Management		T1, T2	T2				
C. Rice Cultivation		NA	NA				
D. Agricultural Soils		NA	T1				
E. Prescribed Burning of Savannas		NA	NA				
F. Field Burning of Agricultural Residues		NA	NA				
G. Liming	T1						
H. Urea Application	T1						
I. Other	NA						
4. Land-Use Land-Use Change and Forestry	T1,T2,T3	T1	T1				
A. Forest Land	T1,T2,T3	T1	T1				
B. Cropland							
C. Grassland	T1, T3		T1				
D. Wetlands	T1	T1	T1				
E. Settlements	T1, T3		T1				
F. Other Land	T1, T3						
G. Harvested wood products	T2						
H. Other							
5. Waste	T1	T1,T2	T1				
A. Solid Waste Disposal	NA	T2	NA				
B. Biological treatment of solid waste	NA	T1	T1				
C. Incineration and open burning of waste	T1	T1	T1				
D. Wastewater treatment and discharge	NA	T1,T2	T1				
E. Other							
6. Other							
Article 3.3 Afforestation and Deforestation	T3	T1	T1				
International Bunkers							
Aviation	T1	T1	T1				
Navigation	T1	T1	T1				
Multilateral Operations	NA	NA	NA				
CO₂ Emissions from Biomass	T1	T1	T1				
CO₂ captured	NA	NA	NA				
Long-term storage of C in waste disposal sites	NA	NA	NA				
Indirect N₂O	NA	NA	NA				
Indirect CO₂	T1	NA	NA				

T1: IPCC Tier 1 or equivalent

T2: IPCC Tier 2 or equivalent

T3: IPCC Tier 3 or equivalent

Table 1.5 Summary of Emission Factors

IPCC SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	NF ₃
1. Energy	CS,D,M,PS	CS,D,M	D,M				
A. Fuel Combustion (Sectoral Approach)	CS,D,M,PS	D,M	D,M				
1. Energy Industries	CS,D,PS	D	D				
2. Manufacturing Industries and Construction	CS,D,PS	D	D				
3. Transport	CS,M	D,M	D,M				
4. Other Sectors	CS,D	D	D				
5. Other							
B. Fugitive Emissions from Fuels	NA	CS,D	NA				
1. Solid Fuels	NA	D	NA				
2. Oil and Natural Gas	NA	CS,D	NA				
C. Carbon Dioxide Transport and Storage	NA	NA	NA				
2. Industrial Processes and Product Use	D,PS	NA	D	CS	NA	NA	NA
A. Mineral Industry	PS						
B. Chemical Industry	NA	NA	NA				
C. Metal Production	NA	NA					
D. Non-Energy Products from Fuels and Solvent Use	D	NA	NA	NA	NA	NA	NA
E. Electronic Industry							
F. Product Uses as Substitutes for ODS				CS	NA	NA	NA
G. Other Product Manufacture and Use			D	NA		NA	
H. Other							
3. Agriculture	D	CS,D	CS,D				
A. Enteric Fermentation		CS,D	NA				
B. Manure Management		CS,D	CS,D				
C. Rice Cultivation		NA	NA				
D. Agricultural Soils		NA	CS,D				
E. Prescribed Burning of Savannas							
F. Field Burning of Agricultural Residues		NA	NA				
G. Liming	D						
H. Urea Application	D						
I. Other	NA						
4. Land-Use Land-Use Change and Forestry	CS,D,OTH	D	D				
A. Forest Land	CS	D	D				
B. Cropland							
C. Grassland	CS,D		D				
D. Wetlands	CS,D	D	D				
E. Settlements	CS,D, OTH		D				
F. Other Land	CS						
G. Harvested wood products	D						
H. Other							
5. Waste	D	CS,D	D				
A. Solid Waste Disposal	NA	CS,D	NA				
B. Biological treatment of solid waste	NA	D	D				
C. Incineration and open burning of waste	D	D	D				
D. Wastewater treatment and discharge	NA	CS,D	D				
E. Other							
6. Other							
Article 3.3 Afforestation and Deforestation	CS	D	D				
International Bunkers							
Aviation	CS	CR	CR				
Marine	CS	D	D				
Multilateral Operations	NA	NA	NA				
CO₂ Emissions from Biomass	CS, D	D, M, CR	D, M, CR				
CO₂ captured	NA	NA	NA				
Long-term storage of C in waste disposal sites	NA	NA	NA				
Indirect N₂O	NA	NA	NA				
Indirect CO₂	CS, CR, D	NA	NA				

PS: Plant specific
D: Default
CS: Country specific
M: Model
CR: CORINAIR

1.6.2 Disaggregated Key Categories

Ireland uses the approach 1 from the 2006 IPCC guidelines to extend the analysis above to identify key categories that may be treated separately at a more disaggregated level, level 3. This gives more information about the individual sources or combination of sources and gases that are of most importance within a Level 2 category. The disaggregation corresponds generally to that at which the emissions are calculated and to that used for estimating uncertainty. The results of the analysis for the approach 1 level 3 assessment in relation to emissions excluding LULUCF in both 1990 and 2013 are presented in Table 1.6 and Table 1.7, respectively. Tables 1.8 and 1.9 present the approach 1 level 3 assessment including LULUCF. Ranking in this way identifies those categories that should be prioritised in the inventory process itself and also the individual components of emissions that could be targeted by specific abatement measures. Results for approach 1 trend assessment for 1990-2013 excluding LULUCF are shown in Table 1.10 and the trend assessment including LULUCF is presented in Table 1.11. The complete tables of ranked sources for 2013 key category analysis are provided in Tables 1.A-D in Annex 1.

The results of the level and trend assessments for 2013 **excluding LULUCF** categories may be summarised as follows:

- (i) The **level assessment** identifies 27 key categories, all of which but five (CO₂ emissions in 1.A.2 Manufacturing Industries & Construction – solid fuels; CH₄ emissions in Manure Management: 3.B.1. Non-Dairy Cattle and 3.B.1 Dairy Cattle; N₂O emissions in 3.D.2 Agricultural Soils (indirect emissions) and 3.B.2.5 Manure Management - Indirect N₂O emissions) are also key categories by trend assessment;
- (ii) There are 17 key categories of CO₂ in level assessment, accounting for 61.4 per cent of total emissions;
- (iii) There are 6 key categories of CH₄, three key categories of N₂O and one category of HFC in level assessment, which account for 21.2 per cent, 10.8 per cent and 1.9 per cent, respectively, of total emissions;
- (iv) Energy accounts for 15 key categories, Agriculture for 9, while Industrial Processes and Product Use contributes two and Waste contributes one;
- (v) The **trend assessment** identifies 27 key categories, all of which but five (CO₂ emissions in 1.A.2. Manufacturing Industries & Construction - Non-renewable waste fuels; CH₄ emissions in: 1.A.4.b. Residential – peat fuel, 1.A.4.b. Residential – solid fuels and 1.B.2.b Fugitive emissions – natural gas; HFC emissions from 2.F.4 Product Uses - Aerosols & MDIs) are key categories for 2013 level assessment;
- (vi) There are 17 key categories of CO₂ in trend assessment, accounting for 79.0 per cent of the total trend;
- (vii) There are 7 key categories of CH₄, one key category of N₂O and two key categories of HFC in trend assessment, which account for 7.2 per cent, 4.3 per cent and 4.5 per cent, respectively, of the total trend.

The results of the level and trend assessment for 2013 **including LULUCF** categories may be summarised as follows:

- (i) The **level assessment** identifies 34 key categories, 22 of these are sources of CO₂ emissions, accounting for 66.9 per cent of total emissions;

- (ii) There are seven additional categories that are not present in the assessment excluding LULUCF, of which six are LULUCF and one is N₂O emissions from Manure Management: 3.B.1. Non-Dairy Cattle;
- (iii) The six additional LULUCF categories are: CO₂ emissions from five categories: 4.C.1 Grassland Remaining Grassland, 4.A.2 Land converted to Forest Land, 4.D.1 Wetlands Remaining Wetlands, 4.G Harvested Wood Products and 4.B.1 Cropland Remaining Cropland and CH₄ emissions from one category: 4.C.1 Grassland Remaining Grassland;
- (iv) There are seven key categories from sources of CH₄, four key categories from N₂O and one category of HFC, which account for 17.8 per cent, 9.1 per cent and 1.6 per cent, respectively, of total emissions;
- (v) Energy accounts for 15 key categories, Agriculture for ten, LULUCF for six, while Industrial Processes contributes 2 and Waste contributes 1;
- (vi) The **trend assessment** identifies 33 key categories, seven of which were not present in the assessment excluding LULUCF: CO₂ emissions from LULUCF categories: 4.A.2 Land converted to Forest Land, 4.A.1 Forest land Remaining Forest Land, 4.C.1 Grassland Remaining Grassland, 4.B.1 Cropland Remaining Cropland, 4.D.1 Wetlands Remaining Wetlands and 4.G Harvested Wood Products and N₂O emissions from: 4.A.2 Land converted to Forest Land. There was also one category present in the assessment excluding LULUCF but not present in assessment including LULUCF: HFC emissions from 2.F.4 Product Uses - Aerosols & MDIs;
- (vii) There are 24 key categories of CO₂ in the trend assessment, accounting for 83.8 per cent of the total trend;
- (viii) There are six key categories of CH₄, one key category of N₂O and one key category of HFC in the trend assessment, which account for 4.9 per cent, 3.4 per cent and 3.0 per cent, respectively, of the total trend.

The list of key categories given by level 3 assessment in 2013 is very similar to that for 1990. However, the higher ranking of the main CO₂ sources in Energy, at the expense of CH₄ and N₂O sources in Agriculture, is notable in 2013. The top ten key categories (excluding LULUCF) were in a different order but identical apart from three (3 key sectors in 2013: CO₂ emissions from 1.A.1 Energy Industries – gaseous fuels, 1.A.4.b. Residential - liquid fuels and 1.A.2. Manufacturing Industries & Construction - gaseous fuels replaced 3 key sectors in 1990: CO₂ emissions from 1.A.4.b. Residential - peat fuel, 1.A.4.b. Residential - solid fuels and 1.A.4.a Commercial/Institutional - liquid fuels). Those seven key categories contributed 55.6 and 57.7 per cent, of total emissions in 1990 and 2013, respectively. The emissions of CO₂ from the use of petrol and diesel by road traffic (1.A.3.b) and CH₄ emissions from 3.A.1. Enteric Fermentation - Non-Dairy Cattle were the largest source categories of greenhouse gas emissions in Ireland in 2013, accounting for 17.8 and 11.3 per cent of the total, respectively.

The CO₂ removals in five categories (4.C.1 Grassland Remaining Grassland, 4.A.2 Land converted to Forest Land, 4.D.1 Wetlands Remaining Wetlands, 4.G Harvested Wood Products and 4.B.1 Cropland Remaining Cropland) and CH₄ emissions from one category (4.C.1 Grassland Remaining Grassland) are key categories in level assessment when the LULUCF sector is included in the detailed analysis. CO₂ removals in category KP A.1 Afforestation (which is determined largely by 5.A.1 Forest Land Remaining Forest Land as well as 5.A.2 Land converted to Forest Land under LULUCF) is a key category in 2013 when Article 3, paragraph 3 and paragraph 4, activities are included in the analysis.

1.6.3 Use of Key Category Analysis

The approach 1 used to determine key categories is based on the principle that the cumulative uncertainty in their emissions represents 90 per cent of the total inventory uncertainty and that 95 per cent of total emissions account for this cumulative fraction of uncertainty. This quantitative approach may therefore result in a much larger number of key categories than might be expected using simpler qualitative criteria. In effect, an inventory with only a small number of major emission sources will require the inclusion of many source categories in order to reach the 95 per cent emissions threshold.

This is well shown by the results of key category determination for Ireland, based on approach 1 level assessment, in Table 1.9. The results including LULUCF indicate that 20 of the 31 key categories in 2013 each accounted for less than 3 per cent of the total emissions and that only seven key categories contributed more than 5 per cent each to the total. The approach 1 analysis adequately identifies the specific sources of emissions that are significant in terms of the overall uncertainty of the inventory but it provides little direction on where to focus priority when the number is large. In these circumstances, information on the uncertainty in the individual source categories and other factors must be taken into account in making decisions regarding the most cost-effective use of inventory capacity related to key categories.

The results of the approach 1 key category analysis in Table 1.7 and 1.9 (excluding LULUCF) clearly show the impact of CO₂ emissions from energy consumption on total emissions in Ireland. These emissions account for 16 of the key categories listed in Table 1.10 (trend, excluding LULUCF) and for 78.6 per cent of total emissions in 2013. While key categories determined by CO₂ emissions from energy consumption have a major bearing on total emissions in Ireland, the remaining potential for significant reduction in the uncertainties associated with these sources is rather limited. The activity data and CO₂ emission factors for Energy source categories in general are among the most reliable items of input data in the inventory and there is consequently little scope for improving the accuracy of the emission estimates. The application of a robust Tier 2 methodology for emissions of CH₄ from enteric fermentation in cattle (non-dairy and dairy) and the use of verified estimates for CO₂ emissions from cement production means that the contributions from these three additional key categories (ranked 18, 20 and 21 in Table 1.10, respectively), making up a further 3.3 per cent of the total, are also known with probably the highest certainty now achievable. The HFC emissions from 2.F.1 Refrigeration and air-conditioning, N₂O emissions from 3.D.1 Agricultural Soils – direct soil emissions and CH₄ emissions from both 3.A.2 Enteric fermentation in sheep and 5.A Solid Waste Disposal account for most of the remaining important key categories in Table 1.10. The uncertainties in the estimates for these complex sources (section 1.6) will remain high due to the large number of factors that influence their emissions and the relatively simple methods that continue to be used.

1.7 Uncertainty Evaluation

The approach 1, propagation of error, method provided by the 2006 IPCC guidelines has been used to make an assessment of uncertainty in the emissions inventory data for 2013 in the same way as for previous years. This method estimates uncertainties for the entire inventory in a particular year and the uncertainty in the trend over time by combining the uncertainties in activity data and emission factors for each source category. The analysis for 2013 data is presented in Table 1.12 (excluding LULUCF) and Table 1.13 (including LULUCF), using emissions on a GWP basis and a level of

source category disaggregation that corresponds closely to the level used for emissions calculation and for key category analysis. This disaggregation level limits the likely dependency and correlation between source categories.

The input values of uncertainty for activity data and emission factors in the GHG inventory have been assigned largely on the basis of general information related to the methodological descriptions in the 2006 IPCC guidelines, supported by opinions elicited from the principal data suppliers, such as statistical offices, energy agencies, Government Departments and individual experts who contributed to certain parts of the inventory.

Where higher tier methods are used for combustion sources, such as those covered by ETS and road transport, the activity data uncertainty estimates are those indicated for the tier concerned. Accordingly, low estimates of uncertainty apply to the activity data for categories such as 1.A.1 Energy Industries and 1.A.3 Transport, as shown on Table 1.12. Slightly higher uncertainty levels are used for energy activity data in sub-categories under 1.A.2 Manufacturing Industries and Construction and 1.A.4 Other Sectors, where the end use of fuels is not as well quantified in the top-down methods used. Low activity data uncertainties are justified in respect of CO₂ emissions sources in Industrial Processes, for which bottom-up data are applied in most cases and the major sources of emissions are covered by ETS. Country-specific CO₂ emission factors are used for all combustion sources, which gives a basis for assigning the uncertainties for emission factors while again taking into account the applicable tiers. Uncertainties in the emission factors for CH₄ and N₂O released from combustion sources are high and not well established quantitatively. For CH₄ and N₂O emission factors for combustion categories, the most up-to-date IPCC publications are used and an indicative uncertainty of 50 per cent is used for both gases.

The Agriculture sector is the second most important sector in Ireland's GHG inventory and has a major influence on overall uncertainty due to its large contribution in terms of CH₄ and N₂O emissions. Ireland has long-established and robust statistical data collection procedures in place for agriculture in general, which guides the selection of 1 per cent as the activity data uncertainty for all agriculture sub-categories. The 2006 IPCC guidelines indicate that the emission factor estimates for the Tier 2 method to determine CH₄ emissions from enteric fermentation in cattle are likely to have an uncertainty of 20 per cent. Following the opinion of national agriculture experts, a value of 15 per cent has been adopted for these emissions to take into account Ireland's detailed Tier 2 method and use of reliable data. In some of the other important emissions sources in Agriculture (such as manure management and agricultural soils) the activity data or emission factors ultimately used are determined by several specific component inputs, which are individually subject to varying degrees of uncertainty. The uncertainty estimates used for emission factors for these sources have been derived by assigning uncertainties to the key component parameters and combining them at the level of activity data or emission factors, as appropriate, using equations 3.1 and 3.2 in chapter 3 of the 2006 IPCC guidelines Volume 1 for each activity to obtain the input to the Tier 1 uncertainty assessment. The footnotes to Table 1.12 show how some of these uncertainty inputs are obtained.

Category 5.A Solid Waste is the principal source of CH₄ emissions outside Agriculture. Under the revised methodology used for category 5.A in this submission, the component uncertainties for both activity data and emission factor for CH₄ generation are derived using equations 3.1 and 3.2 in chapter 3 of the 2006 IPCC guidelines Volume 1 as shown in the footnotes to Table 1.12. These are combined with uncertainties of 30 per cent and 10 per cent for flaring and utilisation respectively to obtain the overall uncertainty using equation 3.2.

Equations 3.1 and 3.2 are both applied as appropriate in a hierarchical approach to derive uncertainty for LULUCF under the Convention and for activities under Article 3, paragraph 3 and paragraph 4, of the Kyoto Protocol. This is achieved by developing uncertainties for carbon pools, which are combined to give the values for the individual land-use categories, which are then combined with uncertainties for other reported activities to give the totals for LULUCF and Article 3, paragraph 3 and paragraph 4, separately. Additional information on uncertainties for LULUCF is provided in chapters 6 and 11.

The F-gas inventory has been substantially revised following work by consultants in 2013, and new data sources were established. The uncertainties associated with the F-gas emission estimates were reviewed, and still considered to be appropriate.

The approach 1 uncertainty analysis (excluding LULUCF) for Ireland's 2013 inventory under the Convention gives an overall uncertainty of 10.25 per cent in total emissions and a trend uncertainty of 3.17 per cent for the period 1990 to 2013. It is an overall increase on level and trend as compared to the values reported in 2014 of 6.79 and 2.71 per cent, respectively.

The reason for the overall increase from 2012 to 2013 is mainly due to the application of revised methodology in two Agricultural sectors: Direct and Indirect N₂O emissions from managed soils as well as new category emissions in Agriculture: 3.D.1.6 Cultivation of Organic Soils (N₂O). These sectors contribute 88 per cent of the overall level uncertainty in the inventory in 2013.

CH₄ emissions from cattle in two agricultural sectors: 3.A Enteric Fermentation and 3.B Manure Management (higher emission factor uncertainty in both sectors) also contributed to the higher overall uncertainty.

The reason for the trend increase from 2012 to 2013 is mainly due to revised (higher) N₂O uncertainty estimates for liquid fuels in four transport sectors (1.A.3. a-d) with road transport the most significant category.

Relatively low estimates are determined largely by the low uncertainties in the estimate of CO₂ emissions, which account for 63.1 per cent of total national emissions in 2013 and which are estimated to have a level uncertainty of 1.32 per cent (excluding LULUCF). When CH₄ is included, bringing the proportion of total emissions up to 85.6 per cent, the total uncertainty estimate is 3.48 per cent (excluding LULUCF), even though there are large uncertainties assigned to the CH₄ emission factors in some source categories. However, it is the influence of N₂O that leads to a higher uncertainty in total emissions bringing it to 10.24 per cent. This influence is not as large in the case of the trend, due to the modest change in emissions of N₂O from 1990 to 2013 and the relatively small share of this gas in total emissions. The impact of HFCs, PFCs, SF₆ and NF₃ on inventory uncertainty remains negligible because these gases account for only 2.3 per cent of total emissions in Ireland.

The approach 1 uncertainty analysis (including LULUCF) for Ireland's 2013 inventory under the Convention (Table 1.13) gives an overall level uncertainty of 9.89 per cent in total emissions and a trend uncertainty of 10.88 per cent for the period 1990 to 2013.

The overall level uncertainty (including LULUCF) of the 2013 inventory is a decrease on the last submission. The corresponding value in 2014 submission (2012 data) was 11.34 per cent. The reason for the decrease from 2012 to 2013 is due to revised uncertainty estimates for CO₂ in LULUCF sector changes: (4.B.2 Land converted to Cropland and 4.E.1 Settlement remaining Settlement). The

decrease is due to the removal of two sectors for CO₂ estimates was slightly offset by an increase due to the new sector being estimated: 4.G Harvested Wood Products. The overall uncertainty increase for N₂O was due to the same reasons as described above in the uncertainty analysis excluding LULUCF. The overall trend uncertainty in 2013 was higher as compared to the 2012 values reported in 2014. The reason for the increase from 2012 to 2013 is due to revised uncertainty estimates for CO₂ in LULUCF sectors in the whole trend.

1.8 Completeness and Time-Series Consistency

Table 1.14 gives an overview of the level of completeness of the 2015 GHG inventory submission with respect to the greenhouse gases covered by the revised UNFCCC reporting guidelines, the IPCC Level 2 source-category split in operation since 2005 for reporting under the Convention and Article 3, paragraph 3, activities under the Kyoto Protocol. Further detail on source/gas coverage at IPCC Level 3 is provided in the individual chapters describing the inventory methods and data for each Level 1 source-category.

The availability of new, more detailed, input data and further methodological guidance in the 2006 IPCC guidelines has allowed some emission calculations to be undertaken at a more detailed level. This has improved the accuracy of the emission estimates, and in some cases the completeness of the inventory has been improved – although not at the sectoral level:

- Fugitive emissions of CH₄ from Solid fuels (coal mines) in Energy sector 1.B.1a have been estimated for the full time series for the first time in this submission.
- In IPPU sector the new categories and gases include: 2.D.1 Lubricant use (CO₂), 2.D.2 Paraffin Wax use (CO₂), 2.G.3 Medical application – anaesthesia (N₂O) and new gas NF₃ in sector 2.E.1 Semiconductor manufacture. Also solvents emissions are now merged into IPPU sector and associated indirect CO₂ emissions from NMVOC from solvents are reported under category 2.D.3 are included in the national total.
- Three new category emissions in Agriculture: 3.D.1.6 Cultivation of Organic Soils (N₂O), 3.G.1 Liming (CO₂) and 3.H Urea Application (CO₂).
- In Waste sector two new categories: 5.B.1 Treatment of solid waste - composting and 5.C.2 Open burning of waste.
- Revised GWPs for CH₄, N₂O and all fluorinated gases.

The work done for the current reporting cycle serves to maintain a complete and consistent emissions time-series by improving the inventories for the years 1990-2012 to bring them fully into line with that for 2013.

Table 1.6 Key Category Analysis Level Assessment 1990 (excluding LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions exclud. LULUCF (kt CO ₂ eq)	1990 Level assessment exclud. LULUCF (%)	Cumulative Total (%)
1	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6,702.59	11.81	11.81
2	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6,581.38	11.60	23.41
3	1.A.1	Energy Industries - Solid Fuels	CO2	4,844.66	8.54	31.94
4	1.A.3.b	Road Transport - Liquid Fuels	CO2	4,690.42	8.26	40.21
5	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3,398.80	5.99	46.20
6	1.A.1	Energy Industries - Peat Fuel	CO2	3,164.78	5.58	51.77
7	1.A.4.b	Residential - Peat Fuel	CO2	3,123.37	5.50	57.28
8	1.A.4.b	Residential - Solid Fuels	CO2	2,483.42	4.38	61.65
9	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2,198.38	3.87	65.53
10	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1,957.00	3.45	68.98
11	1.A.1	Energy Industries - Gaseous Fuels	CO2	1,880.66	3.31	72.29
12	5.A	Solid Waste Disposal	CH4	1,396.49	2.46	74.75
13	1.A.1	Energy Industries - Liquid Fuels	CO2	1,254.90	2.21	76.96
14	3.A.2	Enteric Fermentation - Sheep	CH4	1,176.34	2.07	79.03
15	1.A.4.b	Residential - Liquid Fuels	CO2	1,175.35	2.07	81.11
16	2.B.2	Chemical Industry - Nitric Acid Production	N2O	995.32	1.75	82.86
17	2.B.1	Chemical Industry - Ammonia Production	CO2	990.23	1.74	84.60
18	2.A.1	Cement Production	CO2	884.00	1.56	86.16
19	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	873.02	1.54	87.70
20	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	1.54	89.24
21	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	684.58	1.21	90.44
22	1.A.4.c	Agriculture/Fishing - Liquid Fuels	CO2	660.30	1.16	91.61
23	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	559.22	0.99	92.59
24	3.G.1	Liming - Limestone CaCO3	CO2	355.04	0.63	93.22
25	3.B.1.1	Manure Management - Dairy Cattle	CH4	354.22	0.62	93.84
26	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	0.48	94.32
27	1.A.4.b	Residential - Peat Fuel	CH4	227.65	0.40	94.72
28	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	0.39	95.11

Table 1.7 Key Category Analysis Level Assessment 2013 (excluding LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	2013 Emissions excluding LULUCF (kt CO ₂ eq)	2013 Level assessment excluding LULUCF (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	10,483.36	17.82	17.82
2	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6,626.21	11.27	29.09
3	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	5,633.17	9.58	38.66
4	1.A.1	Energy Industries - Gaseous Fuels	CO2	4,373.09	7.43	46.10
5	1.A.1	Energy Industries - Solid Fuels	CO2	3,797.89	6.46	52.56
6	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3,131.86	5.32	57.88
7	1.A.4.b	Residential - Liquid Fuels	CO2	2,753.85	4.68	62.56
8	1.A.1	Energy Industries - Peat Fuel	CO2	2,579.98	4.39	66.95
9	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	2,039.08	3.47	70.41
10	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	1,709.71	2.91	73.32
11	1.A.4.b	Residential - Gaseous Fuels	CO2	1,420.84	2.42	75.74
12	2.F.1	Refrigeration and air-con (incl. MAC)	HFC	1,124.16	1.91	77.65
13	2.A.1	Cement Production	CO2	1,111.75	1.89	79.54
14	5.A	Solid Waste Disposal	CH4	1,106.03	1.88	81.42
15	1.A.4.b	Residential - Solid Fuels	CO2	1,098.84	1.87	83.29
16	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1,015.60	1.73	85.01
17	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	995.39	1.69	86.71
18	1.A.4.b	Residential - Peat Fuel	CO2	929.40	1.58	88.29
19	3.A.2	Enteric Fermentation - Sheep	CH4	696.09	1.18	89.47
20	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	632.26	1.07	90.54
21	1.A.4.c	Agriculture/Fishing - Liquid Fuels	CO2	541.91	0.92	91.46
22	3.G.1	Liming - Limestone CaCO ₃	CO2	515.69	0.88	92.34
23	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	480.10	0.82	93.16
24	1.A.1	Energy Industries - Liquid Fuels	CO2	401.78	0.68	93.84
25	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	325.65	0.55	94.39
26	3.B.1.1	Manure Management - Dairy Cattle	CH4	286.76	0.49	94.88
27	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	225.44	0.38	95.27

Table 1.8 Key Category Analysis Level Assessment 1990 (including LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions excluding LULUCF (kt CO ₂ eq)	1990 Emissions for LULUCF (kt CO ₂ eq)	Absolute Values (kt CO ₂ eq)	1990 Level assessment including LULUCF (%)	Cumulative Total (%)
1	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6,702.59	0.00	6,702.59	9.49	9.49
2	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6,581.38	0.00	6,581.38	9.32	18.82
3	4.C.1	LULUCF - Grassland Remaining Grassland	CO2	0.00	6,067.33	6,067.33	8.59	27.41
4	1.A.1	Energy Industries - Solid Fuels	CO2	4,844.66	0.00	4,844.66	6.86	34.27
5	1.A.3.b	Road Transport - Liquid Fuels	CO2	4,690.42	0.00	4,690.42	6.64	40.92
6	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3,398.80	0.00	3,398.80	4.81	45.73
7	1.A.1	Energy Industries - Peat Fuel	CO2	3,164.78	0.00	3,164.78	4.48	50.21
8	1.A.4.b	Residential - Peat Fuel	CO2	3,123.37	0.00	3,123.37	4.42	54.64
9	4.A.1	LULUCF - Forest land Remaining Forest Land	CO2	0.00	-2,719.66	2,719.66	3.85	58.49
10	1.A.4.b	Residential - Solid Fuels	CO2	2,483.42	0.00	2,483.42	3.52	62.01
11	4.D.1	LULUCF - Wetlands Remaining wetlands	CO2	0.00	2,291.19	2,291.19	3.25	65.25
12	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2,198.38	0.00	2,198.38	3.11	68.37
13	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1,957.00	0.00	1,957.00	2.77	71.14
14	1.A.1	Energy Industries - Gaseous Fuels	CO2	1,880.66	0.00	1,880.66	2.66	73.80
15	4.B.1	LULUCF - Cropland Remaining Cropland	CO2	0.00	-1,524.52	1,524.52	2.16	75.96
16	5.A	Solid Waste Disposal	CH4	1,396.49	0.00	1,396.49	1.98	77.94
17	1.A.1	Energy Industries - Liquid Fuels	CO2	1,254.90	0.00	1,254.90	1.78	79.72
18	3.A.2	Enteric Fermentation - Sheep	CH4	1,176.34	0.00	1,176.34	1.67	81.38
19	1.A.4.b	Residential - Liquid Fuels	CO2	1,175.35	0.00	1,175.35	1.66	83.05
20	2.B.2	Chemical Industry - Nitric Acid Production	N2O	995.32	0.00	995.32	1.41	84.46
21	2.B.1	Chemical Industry - Ammonia Production	CO2	990.23	0.00	990.23	1.40	85.86
22	2.A.1	Cement Production	CO2	884.00	0.00	884.00	1.25	87.11
23	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	873.02	0.00	873.02	1.24	88.35
24	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	0.00	871.24	1.23	89.58
25	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	684.58	0.00	684.58	0.97	90.55
26	1.A.4.c	Agriculture/Fishing - Liquid Fuels	CO2	660.30	0.00	660.30	0.94	91.49
27	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	559.22	0.00	559.22	0.79	92.28
28	4.G	LULUCF - Harvested wood products	CO2	0.00	-410.59	410.59	0.58	92.86
29	3.G.1	Liming - Limestone CaCO3	CO2	355.04	0.00	355.04	0.50	93.37
30	3.B.1.1	Manure Management - Dairy Cattle	CH4	354.22	0.00	354.22	0.50	93.87
31	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	0.00	269.73	0.38	94.25
32	4.C.1	LULUCF - Grassland Remaining Grassland	CH4	0.00	263.55	263.55	0.37	94.62
33	1.A.4.b	Residential - Peat Fuel	CH4	227.65	0.00	227.65	0.32	94.95
34	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	0.00	223.49	0.32	95.26

Table 1.9 Key Category Analysis Level Assessment 2013 (including LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	2013 Emissions excluding LULUCF (kt CO ₂ eq)	2013 Emissions for LULUCF (kt CO ₂ eq)	Absolute Values (kt CO ₂ eq)	2013 Level assessment including LULUCF (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	10,483.36	0.00	10,483.36	14.50	14.50
2	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6,626.21	0.00	6,626.21	9.16	23.66
3	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	5,633.17	0.00	5,633.17	7.79	31.45
4	4.C.1	LULUCF - Grassland Remaining Grassland	CO2	0.00	4,763.15	4,763.15	6.59	38.04
5	1.A.1	Energy Industries - Gaseous Fuels	CO2	4,373.09	0.00	4,373.09	6.05	44.09
6	1.A.1	Energy Industries - Solid Fuels	CO2	3,797.89	0.00	3,797.89	5.25	49.34
7	4.A.2	LULUCF - Land Converted to Forest Land	CO2	0.00	-3,746.98	3,746.98	5.18	54.52
8	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3,131.86	0.00	3,131.86	4.33	58.85
9	1.A.4.b	Residential - Liquid Fuels	CO2	2,753.85	0.00	2,753.85	3.81	62.66
10	4.D.1	LULUCF - Wetlands Remaining wetlands	CO2	0.00	2,684.47	2,684.47	3.71	66.37
11	1.A.1	Energy Industries - Peat Fuel	CO2	2,579.98	0.00	2,579.98	3.57	69.94
12	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	2,039.08	0.00	2,039.08	2.82	72.76
13	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	1,709.71	0.00	1,709.71	2.36	75.12
14	1.A.4.b	Residential - Gaseous Fuels	CO2	1,420.84	0.00	1,420.84	1.96	77.09
15	2.F.1	Refrigeration and air-con (incl. MAC)	HFC	1,124.16	0.00	1,124.16	1.55	78.64
16	2.A.1	Cement Production	CO2	1,111.75	0.00	1,111.75	1.54	80.18
17	5.A	Solid Waste Disposal	CH4	1,106.03	0.00	1,106.03	1.53	81.71
18	1.A.4.b	Residential - Solid Fuels	CO2	1,098.84	0.00	1,098.84	1.52	83.23
19	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	995.39	0.00	995.39	1.38	84.61
20	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	941.07	0.00	941.07	1.30	85.91
21	1.A.4.b	Residential - Peat Fuel	CO2	929.40	0.00	929.40	1.29	87.19
22	3.A.2	Enteric Fermentation - Sheep	CH4	696.09	0.00	696.09	0.96	88.16
23	4.G	LULUCF - Harvested wood products	CO2	0.00	-691.95	691.95	0.96	89.11
24	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	632.26	0.00	632.26	0.87	89.99
25	1.A.4.c	Agriculture/Fishing - Liquid Fuels	CO2	616.44	0.00	616.44	0.85	90.84
26	3.G.1	Liming - Limestone CaCO3	CO2	515.69	0.00	515.69	0.71	91.55
27	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	480.10	0.00	480.10	0.66	92.22
28	1.A.1	Energy Industries - Liquid Fuels	CO2	401.78	0.00	401.78	0.56	92.77
29	4.C.1	LULUCF - Grassland Remaining Grassland	CH4	0.00	374.97	374.97	0.52	93.29
30	4.B.1	LULUCF - Cropland Remaining Cropland	CO2	0.00	-364.38	364.38	0.50	93.79
31	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	325.65	0.00	325.65	0.45	94.24
32	3.B.1.1	Manure Management - Dairy Cattle	CH4	286.76	0.00	286.76	0.40	94.64
33	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	225.44	0.00	225.44	0.31	94.95
34	3.B.2.1	Manure Management - Non-Dairy Cattle	N2O	217.03	0.00	217.03	0.30	95.25

Table 1.10 Key Category Analysis Trend Assessment 1990-2013 (excluding LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions excluding LULUCF (kt CO ₂ eq)	2013 Emissions excluding LULUCF (kt CO ₂ eq)	2013 Level assessment excluding LULUCF (%)	2013 Trend assessment excluding LULUCF (%)	Contribution to Trend (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	4690.42	10483.36	17.82	9.22	20.26	20.26
2	1.A.1	Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	7.43	3.98	8.73	28.99
3	1.A.4.b	Residential - Peat Fuel	CO2	3123.37	929.40	1.58	3.79	8.32	37.31
4	1.A.4.b	Residential - Liquid Fuels	CO2	1175.35	2753.85	4.68	2.52	5.53	42.84
5	1.A.4.b	Residential - Solid Fuels	CO2	2483.42	1098.84	1.87	2.42	5.32	48.16
6	1.A.1	Energy Industries - Solid Fuels	CO2	4844.66	3797.89	6.46	2.01	4.41	52.57
7	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6581.38	5633.17	9.58	1.95	4.28	56.85
8	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	1420.84	2.42	1.87	4.11	60.96
9	1.A.2	Refrigeration and air-con (incl. MAC)	CO2	873.02	2039.08	3.47	1.86	4.09	65.05
10	2.F.1	Product Uses as Substitutes for ODS -Refrigeration and air-con (incl. MAC)	HFC	0.00	1124.16	1.91	1.84	4.05	69.10
11	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1957.00	941.07	1.60	1.78	3.92	73.02
12	1.A.1	Energy Industries - Liquid Fuels	CO2	1254.90	401.78	0.68	1.47	3.24	76.26
13	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	995.39	1.69	1.25	2.75	79.01
14	1.A.1	Energy Industries - Peat Fuel	CO2	3164.78	2579.98	4.39	1.15	2.52	81.53
15	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	325.65	0.55	0.95	2.08	83.61
16	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2198.38	1709.71	2.91	0.93	2.05	85.66
17	3.A.2	Enteric Fermentation - Sheep	CH4	1176.34	696.09	1.18	0.86	1.89	87.55
18	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3398.80	3131.86	5.32	0.64	1.41	88.96
19	5.A	Solid Waste Disposal	CH4	1396.49	1106.03	1.88	0.56	1.23	90.19
20	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6702.59	6626.21	11.27	0.53	1.16	91.34
21	2.A.1	Cement Production	CO2	884.00	1111.75	1.89	0.32	0.70	92.05
22	1.A.4.b	Residential - Peat Fuel	CH4	227.65	68.42	0.12	0.27	0.60	92.65
23	3.G.1	Liming - Limestone CaCO3	CO2	355.04	515.69	0.88	0.24	0.53	93.18
24	1.B.2.b	Fugitive emissions - Natural gas	CH4	156.05	23.26	0.04	0.23	0.50	93.68
25	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	0.00	132.53	0.23	0.22	0.48	94.16
26	1.A.4.b	Residential - Solid Fuels	CH4	196.51	85.73	0.15	0.19	0.42	94.59
27	2.F.4	Product Uses as Substitutes for ODS -Aerosols (incl. MDIs)	HFC	0.00	114.66	0.19	0.19	0.41	95.00

Table 1.11 Key Category Analysis Trend Assessment 2013 (including LULUCF)

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions including LULUCF (kt CO2 eq)	2013 Emissions including LULUCF (kt CO2 eq)	2013 Level assessment including LULUCF (%)	2013 Trend assessment including LULUCF (%)	Contribution to Trend (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	4690.42	10483.36	14.50	7.67	15.10	15.10
2	4.A.2	LULUCF - Land Converted to Forest Land	CO2	27.26	3746.98	5.18	5.02	9.89	24.98
3	4.A.1	LULUCF - Forest land Remaining Forest Land	CO2	2719.66	74.00	0.10	3.66	7.21	32.19
4	1.A.1	Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	6.05	3.30	6.50	38.70
5	1.A.4.b	Residential - Peat Fuel	CO2	3123.37	929.40	1.29	3.06	6.03	44.73
6	1.A.4.b	Residential - Liquid Fuels	CO2	1175.35	2753.85	3.81	2.09	4.12	48.85
7	4.C.1	LULUCF - Grassland Remaining Grassland	CO2	6067.33	4763.15	6.59	1.96	3.86	52.71
8	1.A.4.b	Residential - Solid Fuels	CO2	2483.42	1098.84	1.52	1.95	3.84	56.55
9	4.B.1	LULUCF - Cropland Remaining Cropland	CO2	1524.52	364.38	0.50	1.62	3.18	59.73
10	1.A.1	Energy Industries - Solid Fuels	CO2	4844.66	3797.89	5.25	1.57	3.10	62.83
11	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	873.02	2039.08	2.82	1.55	3.04	65.87
12	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	1420.84	1.96	1.55	3.04	68.91
13	2.F.1	Refrigeration and air-con (incl. MAC)	HFC	0.00	1124.16	1.55	1.52	2.99	71.90
14	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6581.38	5633.17	7.79	1.50	2.95	74.84
15	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1957.00	941.07	1.30	1.44	2.83	77.67
16	1.A.1	Energy Industries - Liquid Fuels	CO2	1254.90	401.78	0.56	1.19	2.35	80.02
17	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	995.39	1.38	1.03	2.04	82.06
18	1.A.1	Energy Industries - Peat Fuel	CO2	3164.78	2579.98	3.57	0.89	1.76	83.82
19	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	325.65	0.45	0.77	1.51	85.32
20	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2198.38	1709.71	2.36	0.73	1.44	86.76
21	3.A.2	Enteric Fermentation - Sheep	CH4	1176.34	696.09	0.96	0.69	1.35	88.12
22	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3398.80	3131.86	4.33	0.47	0.93	89.05
23	4.D.1	LULUCF - Wetlands Remaining wetlands	CO2	2291.19	2684.47	3.71	0.46	0.90	89.94
24	5.A	Solid Waste Disposal	CH4	1396.49	1106.03	1.53	0.44	0.86	90.80
25	4.G	LULUCF - Harvested wood products	CO2	410.59	691.95	0.96	0.37	0.72	91.53
26	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6702.59	6626.21	9.16	0.32	0.64	92.16
27	2.A.1	Cement Production	CO2	884.00	1111.75	1.54	0.28	0.55	92.71
28	4.A.2	LULUCF - Land converted to Forest Land	N2O	0.00	171.66	0.24	0.23	0.46	93.17
29	1.A.4.b	Residential - Peat Fuel	CH4	227.65	68.42	0.09	0.22	0.44	93.60
30	3.G.1	Liming - Limestone CaCO3	CO2	355.04	515.69	0.71	0.21	0.40	94.01
31	1.B.2.b	Fugitive emissions - Natural gas	CH4	156.05	23.26	0.03	0.18	0.36	94.37
32	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	0.00	132.53	0.18	0.18	0.35	94.72
33	1.A.4.b	Residential - Solid Fuels	CH4	196.51	85.73	0.12	0.16	0.31	95.03

Table 1.12 Tier 1 Uncertainty Estimates 2013 excluding LULUCF (continued on following pages)

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO ₂	1880.66	4373.09	1.00	2.50	2.69	0.04	0.00	0.11	0.11	0.15	0.02
2	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO ₂	1254.90	401.78	1.00	2.50	2.69	0.00	0.00	0.01	-0.04	0.04	0.00
3	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO ₂	0.00	86.50	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Peat	CO ₂	3164.78	2579.98	1.00	5.00	5.10	0.04	0.00	0.06	-0.06	0.09	0.01
5	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO ₂	4844.66	3797.89	1.00	5.00	5.10	0.10	0.01	0.09	-0.11	0.14	0.02
6	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO ₂	873.02	2039.08	7.00	3.00	7.62	0.06	0.00	0.36	0.06	0.36	0.13
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO ₂	2198.38	1709.71	10.00	2.50	10.31	0.08	0.01	0.43	-0.03	0.43	0.18
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO ₂	0.00	132.53	1.00	5.00	5.10	0.00	0.00	0.00	0.01	0.01	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CO ₂	0.00	2.16	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO ₂	871.24	325.65	2.00	5.00	5.39	0.00	0.00	0.02	-0.05	0.05	0.00
11	1.A.3.a Domestic Aviation	CO ₂	51.13	10.01	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.3.b Road Transportation	CO ₂	4690.42	10483.36	1.25	3.00	3.25	0.30	0.09	0.33	0.30	0.44	0.19
13	1.A.3.c Railways	CO ₂	133.19	117.65	1.00	1.00	1.41	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.d Domestic Navigation - Liquid Fuels	CO ₂	84.90	175.55	1.00	2.00	2.24	0.00	0.00	0.00	0.00	0.01	0.00
15	1.A.3.e Other Transportation	CO ₂	62.04	148.53	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.01	0.00
16	1.A.4 Other Sectors - Gaseous Fuels	CO ₂	493.22	2416.22	2.50	2.50	3.54	0.02	0.00	0.15	0.08	0.17	0.03
17	1.A.4 Other Sectors - Liquid Fuels	CO ₂	3792.65	4311.36	10.00	5.00	11.18	0.59	0.35	1.07	0.03	1.07	1.16
18	1.A.4 Other Sectors - Peat	CO ₂	3259.11	929.40	10.00	20.00	22.36	0.11	0.01	0.23	-0.86	0.89	0.80
19	1.A.4 Other Sectors - Solid Fuels	CO ₂	2485.97	1098.84	5.00	10.00	11.18	0.04	0.00	0.14	-0.26	0.29	0.09
20	2.A.1 Cement Production	CO ₂	884.00	1111.75	1.50	1.50	2.12	0.00	0.00	0.04	0.01	0.04	0.00
21	2.A.2 Lime Production	CO ₂	214.08	189.64	5.00	5.00	7.07	0.00	0.00	0.02	0.00	0.02	0.00
22	2.A.3 Glass Production	CO ₂	13.33	0.00	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
23	2.A.4 Other Process Uses of Carbonates	CO ₂	5.32	0.31	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
24	2.B.1 Ammonia Production	CO ₂	990.23	0.00	1.00	5.00	5.10	0.00	0.00	0.00	-0.09	0.09	0.01
25	2.D Non-energy Products from Fuels and Solvent Use	CO ₂	84.42	102.06	30.00	5.00	30.41	0.00	0.00	0.08	0.00	0.08	0.01
26	3.G Liming	CO ₂	355.04	515.69	5.00	50.00	50.25	0.17	0.03	0.06	0.13	0.15	0.02
27	3.H Urea Application	CO ₂	44.47	21.66	5.00	50.00	50.25	0.00	0.00	0.00	-0.02	0.02	0.00
28	4.A.1 Forest Land Remaining Forest Land	CO ₂	-2719.66	74.00	51.00	114.00	124.89	0.15	0.02	0.09	5.32	5.32	28.26
29	4.B.1 Cropland Remaining Cropland	CO ₂	-1524.52	-364.38	20.59	69.15	72.15	-0.42	0.18	-0.17	1.35	1.36	1.85
30	4.C.1 Grassland Remaining Grassland	CO ₂	6067.33	4763.15	12.22	90.00	90.83	6.90	47.60	1.34	-2.12	2.51	6.29
31	4.C.2 Land Converted to Grassland	CO ₂	97.76	138.31	43.95	101.11	110.24	0.24	0.06	0.14	0.06	0.15	0.02
32	4.D.1 Wetlands Remaining Wetland	CO ₂	2291.19	2684.47	21.49	101.45	103.70	4.44	19.71	1.33	0.56	1.45	2.09
33	4.D.2 Land Converted to Wetlands	CO ₂	0.00	7.12	2.50	50.00	50.06	0.01	0.00	0.00	0.01	0.01	0.00
34	4.E.2 Land Converted to Settlements	CO ₂	73.88	54.96	39.97	81.83	91.07	0.08	0.01	0.05	-0.03	0.06	0.00
35	4.F.2 Land Converted to Other Land	CO ₂	0.55	11.06	51.93	75.00	91.23	0.02	0.00	0.01	0.01	0.02	0.00
36	4.G Harvested Wood Products	CO ₂	-410.59	-691.95	25.00	26.92	36.74	-0.41	0.16	-0.40	-0.12	0.42	0.17
37	5.C Incineration and Open Burning of Waste	CO ₂	90.61	43.05	10.00	5.00	11.18	0.00	0.00	0.01	0.00	0.01	0.00
Total CO₂			36724.97	40053.22				5.09					110.11
						Level uncertainty, CO₂			2.26	Trend uncertainty, CO₂			10.49

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	CH4	0.00	1.40	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CH4	3.43	2.11	1.00	70.00	70.01	0.00	0.00	0.00	0.00	0.00	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CH4	0.39	0.11	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CH4	0.00	0.72	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	CH4	1.86	1.56	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CH4	0.91	0.71	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	CH4	1.91	4.71	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH4	0.40	0.91	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH4	2.06	1.47	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH4	0.00	0.11	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CH4	0.00	0.00	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH4	2.30	0.86	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	CH4	0.02	0.01	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	CH4	47.47	16.47	1.25	71.00	71.01	0.00	0.00	0.00	-0.04	0.04	0.00
15	1.A.3.c Railways	CH4	0.19	0.17	1.00	60.00	60.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	CH4	0.20	0.42	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	CH4	0.14	0.33	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	CH4	14.08	16.96	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	CH4	1.12	5.40	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	CH4	11.36	13.62	10.00	66.00	66.75	0.00	0.00	0.00	0.00	0.00	0.00
21	1.A.4 Other Sectors - Peat	CH4	227.98	68.42	10.00	50.00	50.99	0.00	0.00	0.02	-0.15	0.15	0.02
22	1.A.4 Other Sectors - Solid Fuels	CH4	196.51	85.73	5.00	50.00	50.25	0.00	0.00	0.01	-0.10	0.10	0.01
23	1.B.1 Fugitive emissions from Solid Fuels	CH4	55.56	20.15	10.00	50.00	50.99	0.00	0.00	0.01	-0.03	0.03	0.00
24	1.B.2.a Fugitive Emissions from Fuels - Oil and Natural Gas - Oil	CH4	0.21	0.33	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
25	1.B.2.b Fugitive Emissions from Fuels - Oil and Natural Gas - Natural Gas	CH4	156.05	23.26	10.00	50.00	50.99	0.00	0.00	0.01	-0.12	0.12	0.01
26	3.A Enteric Fermentation	CH4	11356.97	10553.71	1.00	17.00	17.03	8.21	67.49	0.26	-0.36	0.45	0.20
27	3.B Manure Management	CH4	1342.29	1242.33	1.00	19.00	19.03	0.14	0.02	0.03	-0.05	0.06	0.00
28	4.A LULUCF - Forest Land	CH4	58.58	77.27	30.00	100.00	104.40	0.13	0.02	0.05	0.03	0.06	0.00
29	4.B.1 LULUCF - Cropland remaining Cropland	CH4	0.07	0.08	100.00	39.10	107.37	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	CH4	263.55	374.97	96.40	91.20	132.70	0.79	0.63	0.83	0.16	0.85	0.72
31	4.D.1 LULUCF - Wetlands remaining Wetlands	CH4	152.31	147.37	86.00	66.50	108.71	0.26	0.07	0.29	-0.01	0.29	0.09
32	5.A Solid Waste Disposal	CH4	1396.49	1119.29	34.64	34.64	48.99	0.76	0.58	0.97	-0.20	0.99	0.97
33	5.C Incineration and Open Burning of Waste	CH4	0.83	0.07	10.00	30.00	31.62	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	CH4	61.10	50.86	10.00	30.00	31.62	0.00	0.00	0.01	-0.01	0.01	0.00
Total CH4			15356.35	13831.90				10.31				2.04	
						Level uncertainty, CH4			3.21	Trend uncertainty, CH4			1.43
Combined CO2 and CH4			52081.32	53885.12				15.40				112.15	
						Level uncertainty, CO2 and CH4			3.92	Trend uncertainty, CO2 & CH4			10.59

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	N2O	0.00	5.45	1.00	63.00	63.01	0.00	0.00	0.00	0.01	0.01	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	N2O	10.21	67.46	1.00	50.00	50.01	0.00	0.00	0.00	0.05	0.05	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	N2O	1.47	0.24	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	N2O	0.00	1.14	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	N2O	52.07	43.93	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.01	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	N2O	7.74	6.05	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	N2O	3.04	7.49	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N2O	0.47	1.09	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N2O	4.83	3.36	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N2O	0.00	0.26	1.00	20.00	20.02	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	N2O	0.00	0.01	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N2O	4.12	1.54	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	N2O	0.55	0.16	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	N2O	48.09	98.33	1.25	68.00	68.01	0.01	0.00	0.00	0.06	0.06	0.00
15	1.A.3.c Railways	N2O	15.49	13.68	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	N2O	0.67	1.43	1.00	90.00	90.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	N2O	0.67	1.58	1.00	25.00	25.02	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	N2O	2.23	2.68	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	N2O	0.27	1.29	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	N2O	76.42	66.32	10.00	50.00	50.99	0.00	0.00	0.02	-0.01	0.02	0.00
21	1.A.4 Other Sectors - Peat	N2O	13.22	3.81	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
22	1.A.4 Other Sectors - Solid Fuels	N2O	11.72	5.11	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
23	2.B.2 Nitric Acid Production	N2O	995.32	0.00	1.00	10.00	10.05	0.00	0.00	0.00	-0.18	0.18	0.03
24	2.G Other Product Manufacture and Use	N2O	31.34	41.06	5.00	5.00	7.07	0.00	0.00	0.01	0.00	0.01	0.00
25	3.B Manure Management	N2O	496.01	517.94	11.22	50.00	51.24	0.18	0.03	0.14	0.00	0.14	0.02
26	3.D.1 Direct N2O Emissions From Managed Soils	N2O	6581.38	5633.17	7.80	100.00	100.30	81.20	6592.83	1.09	-2.09	2.36	5.57
27	3.D.2 Indirect N2O Emissions From Managed Soils	N2O	559.22	480.10	11.18	50.00	51.23	0.15	0.02	0.13	-0.09	0.16	0.03
28	4.A LULUCF - Forest Land	N2O	92.86	171.66	30.00	100.00	104.40	0.29	0.08	0.12	0.13	0.17	0.03
29	4.B.1 LULUCF - Cropland remaining Cropland	N2O	0.02	0.02	100.00	100.00	141.42	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	N2O	0.90	6.66	91.02	100.00	135.22	0.01	0.00	0.01	0.01	0.02	0.00
31	4.C.2 LULUCF - Land converted to Grassland	N2O	0.08	3.74	90.80	100.00	135.07	0.01	0.00	0.01	0.01	0.01	0.00
32	4.D.1 LULUCF - Wetlands remaining Wetlands	N2O	47.16	43.71	86.00	100.00	131.89	0.09	0.01	0.09	-0.01	0.09	0.01
33	5.C Incineration and Open Burning of Waste	N2O	1.04	12.30	10.00	10.00	14.14	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	N2O	95.64	118.70	10.00	10.00	14.14	0.00	0.00	0.03	0.00	0.03	0.00
Total N2O			9173.01	7490.36				82.13					5.71
						Level uncertainty, N2O			9.06	Trend uncertainty, N2O			2.39
Combined CO2, CH4 and N2O			61254.32	61375.48				97.52					117.86
						Level uncertainty, CO2, CH4 & N2O			9.88	Trend uncertainty, CO2, CH4 & N2O			10.86

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	2.E Electronics Industry	Aggregate F-gases	1.17	1307.97	20.00	10.00	22.36	0.22	0.05	0.65	0.23	0.69	0.48
2	2.G Other Product Manufacture and Use	Aggregate F-gases	33.42	21.52	10.00	0.00	10.00	0.00	0.00	0.01	0.00	0.01	0.00
Total F-gases			34.59	1329.50				0.22					0.48
						Level uncertainty, F-gases			0.47	Trend uncertainty, F-gases			0.69
TOTAL for all gases			61288.92	62704.98				97.74					118.34
						Total level uncertainty for all GHGs			9.89	Total trend uncertainty for all GHGs			10.88

Equation 3.1 (chapter 3 of the 2006 IPCC guidelines Volume 1):

$$U_{total} = \sqrt{U_1^2 + U_2^2 + \dots + U_n^2}$$

Where:

U_{total} = the percentage uncertainty in the product of the quantities (half the 95 per cent confidence interval divided by the total and expressed as a percentage);

U_n = the percentage uncertainties associated with each of the quantities.

Equation 3.2 (chapter 3 of the 2006 IPCC guidelines Volume 1):

$$U_{total} = \frac{\sqrt{(U_1 \cdot x_1)^2 + (U_2 \cdot x_2)^2 + \dots + (U_n \cdot x_n)^2}}{|x_1 + x_2 + \dots + x_n|}$$

Where:

U_{total} = the percentage uncertainty in the sum of the quantities (half the 95 per cent confidence interval divided by the total (i.e., mean) and expressed as a percentage). This term 'uncertainty' is thus based upon the 95 per cent confidence interval;

x_n and U_n = the uncertain quantities and the percentage uncertainties associated with them, respectively.

Table 1.13 Tier 1 Uncertainty Estimates 2013 including LULUCF (continued on following pages)

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	1.00	2.50	2.69	0.04	0.00	0.11	0.11	0.15	0.02
2	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO2	1254.90	401.78	1.00	2.50	2.69	0.00	0.00	0.01	-0.04	0.04	0.00
3	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO2	0.00	86.50	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Peat	CO2	3164.78	2579.98	1.00	5.00	5.10	0.04	0.00	0.06	-0.06	0.09	0.01
5	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO2	4844.66	3797.89	1.00	5.00	5.10	0.10	0.01	0.09	-0.11	0.14	0.02
6	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO2	873.02	2039.08	7.00	3.00	7.62	0.06	0.00	0.36	0.06	0.36	0.13
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO2	2198.38	1709.71	10.00	2.50	10.31	0.08	0.01	0.43	-0.03	0.43	0.18
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO2	0.00	132.53	1.00	5.00	5.10	0.00	0.00	0.00	0.01	0.01	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CO2	0.00	2.16	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO2	871.24	325.65	2.00	5.00	5.39	0.00	0.00	0.02	-0.05	0.05	0.00
11	1.A.3.a Domestic Aviation	CO2	51.13	10.01	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.3.b Road Transportation	CO2	4690.42	10483.36	1.25	3.00	3.25	0.30	0.09	0.33	0.30	0.44	0.19
13	1.A.3.c Railways	CO2	133.19	117.65	1.00	1.00	1.41	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.d Domestic Navigation - Liquid Fuels	CO2	84.90	175.55	1.00	2.00	2.24	0.00	0.00	0.00	0.00	0.01	0.00
15	1.A.3.e Other Transportation	CO2	62.04	148.53	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.01	0.00
16	1.A.4 Other Sectors - Gaseous Fuels	CO2	493.22	2416.22	2.50	2.50	3.54	0.02	0.00	0.15	0.08	0.17	0.03
17	1.A.4 Other Sectors - Liquid Fuels	CO2	3792.65	4311.36	10.00	5.00	11.18	0.59	0.35	1.07	0.03	1.07	1.16
18	1.A.4 Other Sectors - Peat	CO2	3259.11	929.40	10.00	20.00	22.36	0.11	0.01	0.23	-0.86	0.89	0.80
19	1.A.4 Other Sectors - Solid Fuels	CO2	2485.97	1098.84	5.00	10.00	11.18	0.04	0.00	0.14	-0.26	0.29	0.09
20	2.A.1 Cement Production	CO2	884.00	1111.75	1.50	1.50	2.12	0.00	0.00	0.04	0.01	0.04	0.00
21	2.A.2 Lime Production	CO2	214.08	189.64	5.00	5.00	7.07	0.00	0.00	0.02	0.00	0.02	0.00
22	2.A.3 Glass Production	CO2	13.33	0.00	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
23	2.A.4 Other Process Uses of Carbonates	CO2	5.32	0.31	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
24	2.B.1 Ammonia Production	CO2	990.23	0.00	1.00	5.00	5.10	0.00	0.00	0.00	-0.09	0.09	0.01
25	2.D Non-energy Products from Fuels and Solvent Use	CO2	84.42	102.06	30.00	5.00	30.41	0.00	0.00	0.08	0.00	0.08	0.01
26	3.G Liming	CO2	355.04	515.69	5.00	50.00	50.25	0.17	0.03	0.06	0.13	0.15	0.02
27	3.H Urea Application	CO2	44.47	21.66	5.00	50.00	50.25	0.00	0.00	0.00	-0.02	0.02	0.00
28	4.A.1 Forest Land Remaining Forest Land	CO2	-2719.66	74.00	51.00	114.00	124.89	0.15	0.02	0.09	5.32	5.32	28.26
29	4.B.1 Cropland Remaining Cropland	CO2	-1524.52	-364.38	20.59	69.15	72.15	-0.42	0.18	-0.17	1.35	1.36	1.85
30	4.C.1 Grassland Remaining Grassland	CO2	6067.33	4763.15	12.22	90.00	90.83	6.90	47.60	1.34	-2.12	2.51	6.29
31	4.C.2 Land Converted to Grassland	CO2	97.76	138.31	43.95	101.11	110.24	0.24	0.06	0.14	0.06	0.15	0.02
32	4.D.1 Wetlands Remaining Wetland	CO2	2291.19	2684.47	21.49	101.45	103.70	4.44	19.71	1.33	0.56	1.45	2.09
33	4.D.2 Land Converted to Wetlands	CO2	0.00	7.12	2.50	50.00	50.06	0.01	0.00	0.00	0.01	0.01	0.00
34	4.E.2 Land Converted to Settlements	CO2	73.88	54.96	39.97	81.83	91.07	0.08	0.01	0.05	-0.03	0.06	0.00
35	4.F.2 Land Converted to Other Land	CO2	0.55	11.06	51.93	75.00	91.23	0.02	0.00	0.01	0.01	0.02	0.00
36	4.G Harvested Wood Products	CO2	-410.59	-691.95	25.00	26.92	36.74	-0.41	0.16	-0.40	-0.12	0.42	0.17
37	5.C Incineration and Open Burning of Waste	CO2	90.61	43.05	10.00	5.00	11.18	0.00	0.00	0.01	0.00	0.01	0.00
Total CO2			36724.97	40053.22				5.09				110.11	
						Level uncertainty, CO2			2.26	Trend uncertainty, CO2			10.49

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	CH4	0.00	1.40	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CH4	3.43	2.11	1.00	70.00	70.01	0.00	0.00	0.00	0.00	0.00	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CH4	0.39	0.11	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CH4	0.00	0.72	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	CH4	1.86	1.56	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CH4	0.91	0.71	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	CH4	1.91	4.71	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH4	0.40	0.91	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH4	2.06	1.47	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH4	0.00	0.11	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CH4	0.00	0.00	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH4	2.30	0.86	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	CH4	0.02	0.01	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	CH4	47.47	16.47	1.25	71.00	71.01	0.00	0.00	0.00	-0.04	0.04	0.00
15	1.A.3.c Railways	CH4	0.19	0.17	1.00	60.00	60.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	CH4	0.20	0.42	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	CH4	0.14	0.33	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	CH4	14.08	16.96	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	CH4	1.12	5.40	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	CH4	11.36	13.62	10.00	66.00	66.75	0.00	0.00	0.00	0.00	0.00	0.00
21	1.A.4 Other Sectors - Peat	CH4	227.98	68.42	10.00	50.00	50.99	0.00	0.00	0.02	-0.15	0.15	0.02
22	1.A.4 Other Sectors - Solid Fuels	CH4	196.51	85.73	5.00	50.00	50.25	0.00	0.00	0.01	-0.10	0.10	0.01
23	1.B.1 Fugitive emissions from Solid Fuels	CH4	55.56	20.15	10.00	50.00	50.99	0.00	0.00	0.01	-0.03	0.03	0.00
24	1.B.2.a Fugitive Emissions from Fuels - Oil and Natural Gas - Oil	CH4	0.21	0.33	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
25	1.B.2.b Fugitive Emissions from Fuels - Oil and Natural Gas - Natural Gas	CH4	156.05	23.26	10.00	50.00	50.99	0.00	0.00	0.01	-0.12	0.12	0.01
26	3.A Enteric Fermentation	CH4	11356.97	10553.71	1.00	17.00	17.03	8.21	67.49	0.26	-0.36	0.45	0.20
27	3.B Manure Management	CH4	1342.29	1242.33	1.00	19.00	19.03	0.14	0.02	0.03	-0.05	0.06	0.00
28	4.A LULUCF - Forest Land	CH4	58.58	77.27	30.00	100.00	104.40	0.13	0.02	0.05	0.03	0.06	0.00
29	4.B.1 LULUCF - Cropland remaining Cropland	CH4	0.07	0.08	100.00	39.10	107.37	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	CH4	263.55	374.97	96.40	91.20	132.70	0.79	0.63	0.83	0.16	0.85	0.72
31	4.D.1 LULUCF - Wetlands remaining Wetlands	CH4	152.31	147.37	86.00	66.50	108.71	0.26	0.07	0.29	-0.01	0.29	0.09
32	5.A Solid Waste Disposal	CH4	1396.49	1119.29	34.64	34.64	48.99	0.76	0.58	0.97	-0.20	0.99	0.97
33	5.C Incineration and Open Burning of Waste	CH4	0.83	0.07	10.00	30.00	31.62	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	CH4	61.10	50.86	10.00	30.00	31.62	0.00	0.00	0.01	-0.01	0.01	0.00
Total CH4			15356.35	13831.90	Level uncertainty, CH4			10.31	Trend uncertainty, CH4			2.04	1.43
Combined CO2 and CH4			52081.32	53885.12	Level uncertainty, CO2 and CH4			15.40	Trend uncertainty, CO2 & CH4			112.15	10.59

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	N2O	0.00	5.45	1.00	63.00	63.01	0.00	0.00	0.00	0.01	0.01	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	N2O	10.21	67.46	1.00	50.00	50.01	0.00	0.00	0.00	0.05	0.05	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	N2O	1.47	0.24	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	N2O	0.00	1.14	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	N2O	52.07	43.93	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.01	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	N2O	7.74	6.05	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	N2O	3.04	7.49	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N2O	0.47	1.09	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N2O	4.83	3.36	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N2O	0.00	0.26	1.00	20.00	20.02	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	N2O	0.00	0.01	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N2O	4.12	1.54	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	N2O	0.55	0.16	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	N2O	48.09	98.33	1.25	68.00	68.01	0.01	0.00	0.00	0.06	0.06	0.00
15	1.A.3.c Railways	N2O	15.49	13.68	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	N2O	0.67	1.43	1.00	90.00	90.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	N2O	0.67	1.58	1.00	25.00	25.02	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	N2O	2.23	2.68	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	N2O	0.27	1.29	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	N2O	76.42	66.32	10.00	50.00	50.99	0.00	0.00	0.02	-0.01	0.02	0.00
21	1.A.4 Other Sectors - Peat	N2O	13.22	3.81	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
22	1.A.4 Other Sectors - Solid Fuels	N2O	11.72	5.11	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
23	2.B.2 Nitric Acid Production	N2O	995.32	0.00	1.00	10.00	10.05	0.00	0.00	0.00	-0.18	0.18	0.03
24	2.G Other Product Manufacture and Use	N2O	31.34	41.06	5.00	5.00	7.07	0.00	0.00	0.01	0.00	0.01	0.00
25	3.B Manure Management	N2O	496.01	517.94	11.22	50.00	51.24	0.18	0.03	0.14	0.00	0.14	0.02
26	3.D.1 Direct N2O Emissions From Managed Soils	N2O	6581.38	5633.17	7.80	100.00	100.30	81.20	6592.83	1.09	-2.09	2.36	5.57
27	3.D.2 Indirect N2O Emissions From Managed Soils	N2O	559.22	480.10	11.18	50.00	51.23	0.15	0.02	0.13	-0.09	0.16	0.03
28	4.A LULUCF - Forest Land	N2O	92.86	171.66	30.00	100.00	104.40	0.29	0.08	0.12	0.13	0.17	0.03
29	4.B.1 LULUCF - Cropland remaining Cropland	N2O	0.02	0.02	100.00	100.00	141.42	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	N2O	0.90	6.66	91.02	100.00	135.22	0.01	0.00	0.01	0.01	0.02	0.00
31	4.C.2 LULUCF - Land converted to Grassland	N2O	0.08	3.74	90.80	100.00	135.07	0.01	0.00	0.01	0.01	0.01	0.00
32	4.D.1 LULUCF - Wetlands remaining Wetlands	N2O	47.16	43.71	86.00	100.00	131.89	0.09	0.01	0.09	-0.01	0.09	0.01
33	5.C Incineration and Open Burning of Waste	N2O	1.04	12.30	10.00	10.00	14.14	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	N2O	95.64	118.70	10.00	10.00	14.14	0.00	0.00	0.03	0.00	0.03	0.00
Total N2O			9173.01	7490.36				82.13					5.71
						Level uncertainty, N2O			9.06	Trend uncertainty, N2O			2.39
Combined CO2, CH4 and N2O			61254.32	61375.48				97.52					117.86
						Level uncertainty, CO2, CH4 & N2O			9.88	Trend uncertainty, CO2, CH4 & N2O			10.86

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	2.E Electronics Industry	Aggregate F-gases	1.17	1307.97	20.00	10.00	22.36	0.22	0.05	0.65	0.23	0.69	0.48
2	2.G Other Product Manufacture and Use	Aggregate F-gases	33.42	21.52	10.00	0.00	10.00	0.00	0.00	0.01	0.00	0.01	0.00
Total F-gases			34.59	1329.50				0.22					0.48
Level uncertainty, F-gases								0.47	Trend uncertainty, F-gases				0.69
TOTAL for all gases			61288.92	62704.98				97.74					118.34
Total level uncertainty for all GHGs								9.89	Total trend uncertainty for all GHGs				10.88

Equation 3.1 (chapter 3 of the 2006 IPCC guidelines Volume 1):

$$U_{total} = \sqrt{U_1^2 + U_2^2 + \dots + U_n^2}$$

Where:

U_{total} = the percentage uncertainty in the product of the quantities (half the 95 per cent confidence interval divided by the total and expressed as a percentage);

U_n = the percentage uncertainties associated with each of the quantities.

Equation 3.2 (chapter 3 of the 2006 IPCC guidelines Volume 1):

$$U_{total} = \frac{\sqrt{(U_1 \cdot x_1)^2 + (U_2 \cdot x_2)^2 + \dots + (U_n \cdot x_n)^2}}{|x_1 + x_2 + \dots + x_n|}$$

Where:

U_{total} = the percentage uncertainty in the sum of the quantities (half the 95 per cent confidence interval divided by the total (i.e., mean) and expressed as a percentage).

This term ‘uncertainty’ is thus based upon the 95 per cent confidence interval;

x_n and U_n = the uncertain quantities and the percentage uncertainties associated with them, respectively.

Table 1.14 Summary of Completeness

IPCC SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	NF ₃
1. Energy							
A. Fuel Combustion (Sectoral Approach)	All	All	All	NA	NA	NA	NA
1. Energy Industries	All	All	All	NA	NA	NA	NA
2. Manufacturing Industries and Construction	All	All	All	NA	NA	NA	NA
3. Transport	All	All	All	NA	NA	NA	NA
4. Other Sectors	All	All	All	NA	NA	NA	NA
5. Other	NO	NO	NO	NA	NA	NA	NA
B. Fugitive Emissions from Fuels							
1. Solid Fuels	NO	All	NO	NA	NA	NA	NA
2. Oil and Natural Gas	All	All	Part	NA	NA	NA	NA
C. Carbon Dioxide Transport and Storage	NO	NO	NO	NA	NA	NA	NA
2. Industrial Processes and Product Use							
A. Mineral Industry	All	Part	Part	NA	NA	NA	NA
B. Chemical Industry	Part	NO	Part	NO	NO	NO	NO
C. Metal Production	NO	NO	NO	NO	NO	NO	NO
D. Non-Energy Products from Fuels and Solvent Use	Part	NA	NA	NA	NA	NA	NA
E. Electronic Industry	NA	NA	NA	All	All	All	All
F. Product Uses as Substitutes for ODS	NA	NA	NA	All	NO	NO	NO
G. Other Product Manufacture and Use	NO	NO	Part	NO	NO	All	NO
H. Other	NO	NO	NO	NA	NA	NA	NA
3. Agriculture							
A. Enteric Fermentation	NA	All	NA	NA	NA	NA	NA
B. Manure Management	NA	All	All	NA	NA	NA	NA
C. Rice Cultivation	NA	NO	NA	NA	NA	NA	NA
D. Agricultural Soils	NA	NE	All	NA	NA	NA	NA
E. Prescribed Burning of Savannas	NO	NO	NO	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NO	NO	NO	NA	NA	NA	NA
G. Liming	NO	NO	NO	NA	NA	NA	NA
H. Urea Application	All	NO	NO	NA	NA	NA	NA
I. Other	All	NO	NO	NA	NA	NA	NA
4. Land-Use Land-Use Change and Forestry							
A. Forest Land	All	Part	Part	NA	NA	NA	NA
B. Cropland	All	NO	All	NA	NA	NA	NA
C. Grassland	All	NO	IE	NA	NA	NA	NA
D. Wetlands	All	NE	All	NA	NA	NA	NA
E. Settlements	Part	NO	NO	NA	NA	NA	NA
F. Other Land	All	NE	NE	NA	NA	NA	NA
G. Harvested Wood Products	All	NO	NO	NA	NA	NA	NA
H. Other	NO	NO	NO	NA	NA	NA	NA
5. Waste							
A. Solid Waste Disposal	NO	All	NA	NA	NA	NA	NA
B. Biological Treatment of Solid Waste	NA	All	All	NA	NA	NA	NA
C. Waste Incineration and Open Burning of Waste	All	All	All	NA	NA	NA	NA
D. Wastewater Treatment and Discharge	NO	All	All	NA	NA	NA	NA
E. Other	NO	NO	NO	NA	NA	NA	NA
6. Other	NO	NO	NO	NA	NA	NA	NA
Article 3.3 Afforestation and Deforestation	All	All	All	NA	NA	NA	NA
Memo Items:							
International Bunkers							
Aviation	All	All	All	NA	NA	NA	NA
Navigation	All	All	All	NA	NA	NA	NA
Multilateral Operations	NO	NO	NO	NA	NA	NA	NA
CO₂ Emissions from Biomass	All	NA	NA	NA	NA	NA	NA
CO₂ captured	NO	NO	NO	NA	NA	NA	NA
Long-term storage of C in waste disposal sites	NE	NO	NO	NA	NA	NA	NA
Indirect N₂O	NO	NO	NE	NA	NA	NA	NA
Indirect CO₂	Part	NO	NO	NA	NA	NA	NA

All : Emissions of the gas are covered for all sources under the source category/memo item

NA : Emissions of the gas not applicable to the source category/memo item

NO : Emissions of the gas does not occur in Ireland for the source category/memo item

NE : Emissions on the gas not estimated for the source category/memo item

Part : Emissions of the gas estimated for some activities in the source category

Chapter 2 Trends in greenhouse gas emissions

2.1 Description and interpretation of emission trends for aggregated GHG emissions

The trends in emissions of the greenhouse gases in Ireland over the period 1990-2013 are shown in Figure 2.1 and Table 2.1. The estimates reported here show some changes on those reported in the 2014 submission, which reflect recalculations that are fully described in subsequent chapters. The trends in the principal emission components, shown as CO₂ equivalents, within the five IPCC sectors are shown on Figure 2.4 through Figure 2.11. Total emissions of the seven greenhouse gases in Ireland (excluding net emissions from Land Use Land Use Change and Forestry) increased steadily from 56,751.48 kt CO₂ eq in 1990 to 71,802.64 kt CO₂ eq in 2001 and then decreased slightly to 69,347.40 kt CO₂ eq in 2004. Total emissions increased again in 2005 to 71,427.64 kt CO₂ eq and then decreased for six consecutive years to 58,827.88 in 2011, then increased in 2012 to 59,599.34 kt CO₂ eq and decreased to 58,820.88 kt CO₂ eq in 2013. The largest annual change occurred from 2008 to 2009 when emissions decreased by 5,403.31 kt CO₂ eq from 68,879.27 to 63,475.96 kt CO₂ equivalent, a reduction of 7.8 per cent. Total emissions in 2013 were 3.6 per cent higher than in 1990 and 18.1 per cent lower than the peak level in 2001. The estimated total for 2013 is 58,820.88 kt CO₂ eq, equating to 778.46 kt CO₂ eq or 1.3 per cent lower than that for 2012. Inter annual changes to national total emission estimates are shown in Figure 2.2.




Figure 2.1 National total Greenhouse Gas emissions (excluding LULUCF) 1990-2013




Figure 2.2 Inter annual change in national total emissions 1990-2013

In 2013, the total Energy sector accounted for 60.8 per cent of total emissions, Agriculture contributed 32.2 per cent while a further 4.7 per cent emanated from Industrial Processes and Product Use and 2.3 per cent was due to Waste. The Energy accounted for the bulk of the CO₂ emissions in 2013 (94.7 per cent), IPPU and Agriculture sectors contributed further 3.8 per cent and 1.4 per cent, respectively and Waste contributed the remainder 0.1 per cent. CH₄ emissions are produced mainly in the Agriculture (89.1 per cent) and Waste (8.8 per cent), Energy contributed the remainder 2.0 per cent. Most of the N₂O emissions are generated in Agriculture (92.9 per cent) and Energy (4.7 per cent) with Waste and IPPU contributing further 1.8 per cent and 0.6 per cent, respectively. IPPU sector is responsible for 100 per cent of F-gas emissions.

The large increase in emissions during the period 1990-2001 was clearly driven by the growth in CO₂ emissions from energy use. CO₂ from energy use increased its share of national total emissions from 53.1 per cent in 1990 to 61.0 per cent share in 2001. The bulk of this increase occurred in the years between 1994 and 2001, during which Ireland experienced a period of unprecedented economic growth with energy CO₂ emissions increasing by an average of 4.4 per cent annually. The rate of economic growth slowed down from 2002 to 2004, which together with the closure of ammonia and nitric acid production plants and the continued decline in cattle populations and fertiliser use resulted in a reduction in the emission levels in the period 2002 to 2004. The increase in 2005 was largely due to increased emissions from road transport and from electricity generation from two new peat-fired stations. The declining trend between 2005 and 2008 is largely attributable to decreases in the agriculture and waste sectors and in 2008 to reduced emissions from mineral products in the industrial processes sector. In addition, the sustained increase in transport emissions, the major contributor to the trend, came to an end in 2008 and together with the recent economic downturn caused a major decrease in emissions in 2009 to 2011, before rising in 2012 and decreasing in 2013, with 2009 emissions reaching the level pre-1998.

Table 2.1. Greenhouse Gas Emissions 1990, 1995, 2000-2013 (kt CO₂ equivalent)

(a) Emissions by Gas

GREENHOUSE GAS EMISSIONS	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Change from base to latest reported year
CO ₂ emissions without net CO ₂ from LULUCF	32,742.45	35,693.95	45,094.54	47,470.72	45,952.73	45,554.32	46,042.40	47,952.20	47,403.08	47,489.84	47,170.87	41,981.59	41,553.77	37,898.34	38,066.63	37,057.31	13.18%
CO ₂ emissions with net CO ₂ from LULUCF	36,645.64	41,831.18	50,214.62	55,193.49	52,931.65	52,775.22	49,730.36	50,629.22	51,826.12	51,911.76	50,737.49	42,902.87	44,512.18	43,465.18	42,870.83	39,987.08	9.12%
CH ₄ emissions without CH ₄ from LULUCF	14,881.87	15,129.90	14,532.19	14,585.95	14,525.92	15,178.67	14,248.66	13,980.69	13,991.96	13,362.99	13,206.40	12,870.52	12,632.35	12,593.36	12,890.28	13,232.26	-11.08%
CH ₄ emissions with CH ₄ from LULUCF	15,356.39	15,696.78	15,027.68	15,263.33	14,971.43	15,879.21	14,870.44	14,449.93	14,577.24	13,989.07	13,802.80	13,365.41	13,577.72	13,542.40	13,369.30	13,831.95	-9.93%
N ₂ O emissions without N ₂ O from LULUCF	9,013.24	8,597.90	8,585.93	8,843.86	8,171.43	8,221.26	7,926.38	8,134.84	7,806.17	6,931.49	7,103.19	7,318.93	7,476.14	7,061.28	7,395.64	7,135.66	-20.83%
N ₂ O emissions with N ₂ O from LULUCF	9,173.01	8,805.06	8,814.32	9,116.53	8,419.20	8,522.16	8,221.16	8,416.61	8,123.38	7,264.18	7,461.99	7,661.32	7,900.08	7,478.82	7,731.37	7,490.42	-18.34%
HFCs	0.59	41.12	303.60	353.64	434.07	589.38	732.79	939.29	975.68	984.05	1,128.30	1,108.08	1,127.64	1,147.98	1,135.47	1,276.74	35.93%
PFCs	0.12	97.61	397.76	379.51	267.89	285.95	234.81	216.39	190.96	168.10	136.14	83.63	46.58	15.88	9.56	8.32	-96.15%
SF ₆	33.88	79.11	51.76	64.63	64.48	109.95	65.34	96.78	60.21	62.94	54.69	39.18	33.08	45.45	37.39	43.53	-55.02%
NF ₃	NO	4.37	49.17	21.78	46.58	46.63	18.08	28.38	28.21	37.67	NO	NO	NO	NO	0.78	0.90	-96.83%
Total (without LULUCF)	56,672.15	59,643.97	69,014.95	71,720.09	69,463.11	69,986.16	69,268.46	71,348.57	70,456.26	69,037.07	68,799.59	63,401.94	62,869.55	58,762.29	59,535.75	58,754.73	3.67%
Total (with LULUCF)	61,209.63	66,555.25	74,858.91	80,392.92	77,135.32	78,208.50	73,872.98	74,776.59	75,781.79	74,417.77	73,321.41	65,160.49	67,197.29	65,695.70	65,154.69	62,638.94	2.34%
Total (without LULUCF, with indirect)	56,751.48	59,728.00	69,091.27	71,802.64	69,543.75	70,065.88	69,347.40	71,427.64	70,540.55	69,128.51	68,879.27	63,475.96	62,936.70	58,827.88	59,599.34	58,820.88	3.65%
Total (with LULUCF, with indirect)	61,288.96	66,639.28	74,935.22	80,475.47	77,215.96	78,288.22	73,951.92	74,855.66	75,866.08	74,509.21	73,401.09	65,234.52	67,264.44	65,761.28	65,218.28	62,705.09	2.31%

(b) Emissions by IPCC Source Category

GREENHOUSE GAS SOURCE AND CATEGORIES	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Change from base to latest reported year
1. Energy	31,117.76	33,892.07	42,525.09	44,626.25	43,402.57	44,034.48	43,826.37	45,648.99	45,152.07	45,115.66	45,210.74	40,731.28	40,359.13	36,879.60	36,950.03	35,737.69	14.85%
2. Industrial Processes	3,173.30	3,112.34	4,491.39	4,553.48	4,033.34	3,442.48	3,630.74	3,954.32	3,869.92	3,919.34	3,697.21	2,795.47	2,577.97	2,452.10	2,650.44	2,708.17	-14.66%
3. Agriculture	20,735.38	20,663.57	20,246.85	20,635.52	19,973.97	20,349.23	19,891.73	19,959.03	19,551.32	18,565.73	18,576.15	18,705.05	18,758.03	18,165.33	18,750.34	18,964.61	-8.54%
4. LULUCF	4,537.48	6,911.28	5,843.96	8,672.83	7,672.21	8,222.34	4,604.52	3,428.03	5,325.53	5,380.70	4,521.82	1,758.56	4,327.74	6,933.40	5,618.94	3,884.21	-14.40%
5. Waste	1,645.71	1,975.99	1,751.62	1,904.84	2,053.23	2,159.96	1,919.62	1,786.23	1,882.96	1,436.34	1,315.50	1,170.14	1,174.42	1,265.26	1,184.94	1,344.27	-18.32%
6. Other	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NA
Total (including LULUCF)	61,209.63	66,555.25	74,858.91	80,392.92	77,135.32	78,208.50	73,872.98	74,776.59	75,781.79	74,417.77	73,321.41	65,160.49	67,197.29	65,695.70	65,154.69	62,638.94	2.34%

2.2 Trends by Gas

Emissions of CO₂ accounted for 63.1 per cent of the total (excluding LULUCF) of 58,820.88 kt CO₂ equivalent in 2013, with CH₄ and N₂O contributing 22.5 per cent and 12.1 per cent, respectively. The combined emissions of HFC, PFC, SF₆ and NF₃ accounted for 2.3 per cent of total emissions in 2013. In 1990 emissions of CO₂, CH₄, N₂O and the combined emissions of HFCs, PFCs, SF₆ and NF₃ accounted for 57.8, 26.2, 15.9 and less than 0.1 per cent, respectively of total emissions of 56,751.48 kt CO₂ equivalent as presented in Figure 2.3.




Figure 2.3 Greenhouse Gas emissions-by Gas 1990-2013

CO₂ is the most significant contributor to the greenhouse gas emissions with 1.A.1 Energy Industries and 1.A.3 Transport sectors responsible for 30.3 per cent and 29.5 per cent of total CO₂ emissions in 2013, respectively. 1.A.4 Other Sectors represents a share of 23.6 per cent, 1.A.2 Manufacturing Industries and Construction has an 11.3 per cent share and the remainder of CO₂ emissions (5.3 per cent share) fall into other sectors. Emissions of CO₂ increased from 32,821.78 kt in 1990 to 37,123.45 kt in 2013, which equates to an increase of 13.1 per cent. The main driver behind this increase in emissions is primarily fuel combustion in Transport followed by Energy Industries. Over the period 1990-2013, emissions of CO₂ from transport, predominantly road traffic in Ireland, increased by 117.8 per cent. This trend is exaggerated somewhat in later years by so-called fuel-tourism. In 2013 it is estimated that approximately 7.2 per cent of automotive fuel sold in Ireland is used in vehicles in the UK and other countries. Over the time-series, emissions of CO₂ from 1.A.1 Energy Industries increased in the first decade by 54.7 per cent until they peaked in 2001 and decreased by 34.8 to 2013, showing an overall increase of 0.8 per cent CO₂ over the 1990-2013 period. In addition, even though Ireland has only a small number of energy intensive industries, CO₂ emissions from combustion in the industrial sector 1.A.2 Manufacturing Industries and Construction increased by 6.8 per cent between 1990 and 2013.

Methane is the second most significant contributor to greenhouse gas emissions in Ireland which is due to the large population of cattle. In 2013 emissions of CH₄ were 13,232.26 CO₂ equivalent, indicating a decrease of 11.1 per cent on the 1990 level of 14,881.87 kt CO₂ equivalent. Emissions of CH₄ increased progressively from 1990, reaching a peak in 1998 of 15,593.92 kt CO₂ equivalent, which reflects an increase in livestock numbers and therefore increased emissions from source categories 3.A Enteric Fermentation and 3.B Manure Management. Between 1998 and 2011 CH₄ emissions decreased as a result of falling livestock numbers due to reform of the Common Agricultural Policy (CAP). However, total CH₄ emissions in the period 2001-2013 fluctuated to some extent on a yearly basis. This trend is a direct result of fluctuating CH₄ emissions from 1.A.4 Other Sectors and 1.B Fugitive Emissions from Fuels. The main contributor to the CH₄ trend has been Agriculture and in 2013 the sector accounted for 89.1 per cent of the total methane emissions (compared to 85.3 per cent share in 1990 when emissions from Waste had a larger share in the methane trend). Nevertheless, the sectoral methane emissions from Agriculture decreased by 7.1 per cent between 1990 (12,699.26 kt CO₂ equivalent) and 2013 (11,796.04 kt CO₂ equivalent). Another significant source of methane emissions is Waste sector, especially from landfill gas in category 5.A Solid Waste Disposal on Land. CH₄ emissions from Waste decreased from 9.8 per cent share of total methane emissions (1,458.42 kt CO₂ equivalent) in 1990 to 8.8 per cent share (1,170.22 kt CO₂ equivalent) in 2013. This decrease is a result of improved management of landfill facilities, including increased recovery of landfill gas utilised for electricity generation and flaring.

Emissions of N₂O decreased by 20.8 per cent from their 1990 level of 9,013.24 kt CO₂ equivalent in 1990 to 7,135.66 kt CO₂ equivalent in 2013. Similar to CH₄, emissions of N₂O increased during the 1990s to reach peak level of 9,308.11 kt CO₂ equivalent in 1998 reflecting increased use of synthetic fertilisers and increased amounts of animal manures associated with increasing animal numbers over that period. Emissions of N₂O subsequently show a clear downward trend following reductions in synthetic fertiliser use and organic nitrogen applications on land as a result of the effect of CAP reform on animal numbers as well the closure of Ireland's only nitric acid plant in 2002. The largest contributor to the trend is Agriculture sector with 92.9 per cent share of the total N₂O emissions (6,631.21 kt CO₂ equivalent) in 2013. This reflects an increase from 84.7 per cent share (7,636.62 kt CO₂ equivalent) in 1990 despite being a lower absolute number. Emissions from IPPU in chemical industry used to be the second largest contributor to the trend contributing 11.4 per cent to total N₂O emissions in 1990 and an average of 9.6 per cent share to the trend between 1990 and 2000, before falling to 3.9 per cent share in 2002 – the year of nitric acid plant closure. Energy and Waste sectors contribute 4.7 per cent and 1.8 per cent respectively to the rest of the N₂O trend.

Emissions of the F-gases (HFCs, PFCs, SF₆ and NF₃) were 1,329.50 kt CO₂ equivalent in 2013 compared to 34.59 kt CO₂ equivalent in 1990, a 38 fold increase over the time series. However, F-gas emissions only account for 2.3 per cent of the national total. F-gases include a wide range of substances that are used in a diverse range of products and manufacturing processes. Therefore it can be difficult to identify the factors contributing to actual trends in emissions over time. However, it is possible to establish the main contributory sub-categories underlying these trends.

The main causative factor of the increase in F-gas emissions has been the growth in HFC emissions from 2.F.1 Refrigeration and Air Conditioning through their use as replacement refrigerants across virtually all refrigeration sub-categories since 1996. Increased use of HFCs in 2.F.4 categories: Metered Dose Inhalers (MDIs) and Aerosols is also an important component of the trend. On the

other hand, following a 2013 study on F-gases, emissions from 2.F.2 Foams were proven to be not occurring in manufacturing process and consequently were removed from the whole time series. Similar was the finding in 2.F.3 Fire extinguishers between 1990-1996 (incl.) and significant emission reductions for the following years in the trend have been applied. Sector 2.E.1 Semiconductor Manufacture was the only source in 1990 until 2.F.4 Aerosols entered the market in 1992, followed by 2.F.1 MAC in 1993, 2.F.1 Refrigeration and Air Conditioning in 1995 and both 2.F.3 Fire extinguishers and 2.F.4 MDIs in 1996. Emissions from HFCs increased steadily from 0.59 kt CO₂ equivalent in 1990 CO₂ to 1,276.74 kt CO₂ in 2013.

Emissions of PFCs increased from 0.12 kt CO₂ equivalent in 1990 up to 130.82 kt CO₂ equivalent in 1997 through their use in the semiconductor manufacturing process in 2.E.1 Semiconductor Manufacture. Emissions subsequently decreased, only to significantly increase to reach their peak at 397.76 kt CO₂ equivalent in 2000. Semiconductor manufacturers continue to investigate various reduction initiatives through gas substitution and new process technologies which is reflected in the downward trend in PFC emissions between 2000 and 2013 (8.32 kt CO₂ equivalent in 2013).

SF₆ is used in a diverse number of products and processes and is therefore included in a number of IPCC source sub-categories including 2.E.1 Semiconductor Manufacture, 2.G.1 Electrical Equipment and four subcategories under 2.G.2 Other. Emissions of SF₆ were 33.88 kt CO₂ equivalent and 43.53 kt CO₂ equivalent in 1990 and 2013, respectively. However, total emissions of SF₆ across the time series vary considerably, primarily because the two largest sources (Semiconductor Manufacture and Electrical Equipment) vary considerably from year to year. Emissions of SF₆ grew steadily from 1990, peaking at 126.12 kt CO₂ equivalent in 1997. The increase over the period 1990-1997 was largely due to increased use of SF₆ in Semiconductor Manufacture. Emissions from both Semiconductor Manufacture and Electrical Equipment then show a steady decline across the time series (although there are peaks in 2003 and 2005 due to elevated emissions from Semiconductor Manufacture). Emissions in 2013 (43.53 kt CO₂ equivalent) were lower than those in 1992.

Similar to PFCs, semiconductor manufacturers have undertaken to reduce the use of SF₆ through gas substitution and new process technologies. In 2.E.1 Electrical Equipment, where SF₆ is used for electrical insulation, arc quenching and current interruption, a leak reduction programme has been in place since 1997, when peak emissions are observed. SF₆ use and methodology, similar like HFC gases has undergone a revision resulting in some changes mainly in sector 2.G.2 Other - window soundproofing with reduction of emissions in the first decade followed by increased emissions between 2001-2013.

NF₃ are solely released from 2.E.1 Semiconductor Manufacture. Emissions of NF₃ were reported since 1995 (4.37 kt CO₂ eq.) when use of this gas commenced in the industry and peaked in 2000 (49.17 kt CO₂ eq.), followed by fluctuations until 2008 when NF₃ was phased out from Semiconductor Manufacture for four consecutive years. In 2012 and 2013 small amounts of NF₃ were used again in Semiconductor Manufacture resulting in low emission levels (0.78 kt CO₂ eq. and 0.90 kt CO₂ eq, respectively).

2.3 Description and interpretation of emission trends by sector

Greenhouse gas emissions broken down by IPCC sector are presented in Table 2.1 (b). It can be seen that the largest contribution is from the Energy sector, which in 2013 contributes 60.8 per cent of

total greenhouse gas emissions (excluding LULUCF). The second largest sector is Agriculture, which accounted for 32.2 per cent of total greenhouse gas emissions in 2013. Emissions from Industrial Processes and Product Use accounted for 4.7 per cent and Waste accounted for 2.3 per cent of total emissions in 2013. The following sub-sections discuss the main contributors to trends within each IPCC source sector including LULUCF sector. Emissions of indirect gases are discussed in section 2.4.

2.3.1 Trends in Energy (IPCC Sector 1)

Emissions from the Energy sector increased by 14.8 per cent from 31,117.76 kt CO₂ equivalent in 1990 to 35,737.69 kt CO₂ equivalent in 2013. The most significant increases occurred between 1994 and 2001, driven by major increases in emissions from 1.A.1 Energy Industries and 1.A.3 Transport. Emissions were comparatively stable between 2001 and 2008, reaching a peak in 2005 with 45,648.99 kt CO₂ equivalent. A major decrease occurred between 2008 and 2009 when the sectoral emissions fell by 9.9 per cent. A further reduction of 12.3 per cent has occurred to give 35,737.69 kt CO₂ equivalent in 2013.

Energy Industries (1.A.1) accounted for 19.8 per cent and 19.3 per cent of total national greenhouse gas emissions in 1990 and 2013, respectively. Total greenhouse gas emissions from this sub-sector increased by 54.5 per cent from 11,223.13 CO₂ equivalent in 1990 to 17,334.22 CO₂ equivalent in 2001. Some reductions were achieved in 2002, 2003 and 2004 from improvements in energy efficiency and fuel switching as new electricity producers entered the market with the result that emissions decreased to 15,335.43 kt CO₂ equivalent in 2004. Emissions subsequently increased in 2005 to 15,766.71 kt CO₂ equivalent as levels of peat use returned to former levels with the entry into service of two new peat fired power plants. Emissions in 2006 decreased to 15,024.40 kt CO₂ equivalent due to a reduction in the use of Moneypoint coal-fired station during the installation of pollutant control measures, while further reductions in 2007 (14,530.69 kt CO₂ equivalent) are largely a result of the displacement of oil by natural gas. In 2008, emissions increased by 0.8 per cent or 116.03 kt CO₂ equivalent to 14,646.72 kt CO₂ equivalent, then decreased in 2009 by 10.8 per cent to 13,071.55 kt CO₂ equivalent reflecting the impact of the economic recession in Ireland. There was a slight increase in emissions (2.0 per cent) in 2010 to reach 13,326.90 kt CO₂ equivalent which reflects a reduction in the share of renewables in gross electricity consumption from 14.3 per cent in 2009 to 12.9 per cent in 2010. Wind and hydro resources were less in 2010 which resulted in more electricity generation from coal and gas-fired power stations. By 2012, wind and hydro energy generation had grown substantially, resulting in a renewables contribution to gross electricity consumption of 18.9 per cent. However, these changes combined with increased consumption of coal and reduction of natural gas resulted in an increase in emissions from the Energy industries sector of 7.0 per cent between 2011 and 2012, from 11,935.22 kt CO₂ equivalent to 12,774.63 kt CO₂ equivalent, respectively. In 2013 emissions from this sector decreased by 11.0 per cent on 2012 levels to reach 11,370.18 kt CO₂ equivalent, which reflects further increase in the share of renewables in gross electricity consumption with 20.1 per cent contribution in 2013. Overall drivers and trends in emissions from the Energy sector are presented in Figure 2.4 and Figure 2.5.




Figure 2.4 Total Primary Energy Requirement (TPER) 1990-2013




Figure 2.5 Trend in Emissions from Energy 1990-2013

There are only a small number of energy intensive industries in Ireland under sub-category 1.A.2 Manufacturing Industries and Construction. This sub-category accounted for 7.0 per cent (3,961.76 kt CO₂ equivalent) and 7.2 per cent (4,230.93 kt CO₂ equivalent) of total national greenhouse gas emissions in 1990 and 2013, respectively. However, the trend shows an increase of 6.8 per cent over

the same period as a result of large increases in use of petroleum coke in 1.A.2.f Non-metallic minerals and natural gas in 1.A.2.b Non-ferrous metals, 1.A.2.e Food Processing, Beverages and Tobacco and 1.A.2.g Other Industries. Emissions from the sector were increasing in the trend and remained at their highest between years 2000 and 2008 with their peak at 5,870.82 kt CO₂ equivalent in 2005. Following an economic downturn emissions sharply declined by 20.5 per cent between 2008 and 2009, from 5,654.19 kt CO₂ equivalent to 4,493.97 kt CO₂ equivalent, respectively and continued to decline until 2011 (4,166.68 kt CO₂ equivalent), followed by a small increases in 2012 and 2013, by 0.2 per cent and 1.5 per cent as compared to 2011 levels when manufacturing industry started to recover from recession.

Fuel combustion emissions in 1.A.3 Transport accounted for 9.0 per cent and 18.8 per cent of total national greenhouse gas emissions in 1990 and 2013, respectively. The overall sector's emissions increased by 115.5 per cent from 5,135.18 kt CO₂ equivalent in 1990 to 11,067.68 kt CO₂ equivalent in 2013. This is largely accounted for by a 121.4 per cent increase in road transport emissions over the same period, due to sustained growth in the use of passenger cars and goods vehicles. The trend is however, somewhat exaggerated by so-called fuel tourism whereby a proportion of the automotive fuel sold in the Republic of Ireland is used in vehicles in the UK and other countries. Fuel tourism is estimated to account for 7.2 per cent of automotive fuels in 2013. It is worth noting that in the years 1990-1995 inclusive there was cross border movement of automotive fuels into the Republic of Ireland. The principal drivers in road transport emission trends are shown in Figures 2.6 and 2.7. Transport emissions were 3.32 million tonnes CO₂ equivalent lower in 2013 than in 2007. This represents a decrease of 23.1 per cent, following sustained increases in this sector since 1990. The decrease primarily reflects the impact of the economic downturn plus the changes in vehicle registration tax and road tax introduced in mid-2008 and the Biofuels Obligation Scheme. Emissions from domestic aviation decreased by 80.3 per cent between 1990 (51.73 kt CO₂ equivalent) and 2013 (10.18 kt CO₂ equivalent), having peaked in 2006 at 77.29 kt CO₂ equivalent. However, their overall effect on transport emission trends is negligible.




Figure 2.6 Fuel use in Road Transport 1990-2013




Figure 2.7 Vehicle numbers and Census of Population 1990-2013

Emissions from category 1.A.4 Other Sectors decreased by 14.7 per cent from 10,585.87 kt CO₂ equivalent in 1990 to 9,025.16 kt CO₂ equivalent in 2013. Emissions from the Commercial (1.A.4.a), Residential (1.A.4.b) and Agriculture/Fishing (1.A.4.c) sub-categories decreased by 13.0, 15.0 and 17.9 per cent, respectively. Although residential fossil fuel consumption increased by 8.5 per cent from 1990 to 2013 there has been a decline in the use of carbon-intensive fuels, such as peat and coal, and greater use of oil and natural gas. The emissions of CO₂ from coal and peat use in the residential sector decreased by 63.7 per cent between 1990 and 2013 while those from oil and natural gas increased by 200.8 per cent over this period.

2.3.2 Trends in Industrial Processes and Product Use (IPCC Sector 2)

The contribution from Industrial Processes and Product Use (IPPU) is relatively small, accounting for 5.7 per cent of total greenhouse gases in 1990 and 4.7 per cent in 2013. Total emissions from the sector were 3,252.63 kt CO₂ equivalent in 1990 and 2,774.31 kt CO₂ equivalent in 2013. This is a decrease of 14.7 per cent in emissions over the time series. Overall trends in emissions from IPPU are presented in Figure 2.8.

In the early 1990's (1990 to 1994) the contribution of 2.B Chemical Industry to overall sectoral emissions was on average 60.0 per cent. By the late 1990's (1995 to 1999) this proportion had fallen to 49.8 per cent on average of total emissions from the sector. In 1990 emissions from 2.B. Chemical Industry were 1,985.55 kt CO₂ equivalent, however by 2000 they had reduced by 16.2 per cent to 1,663.30 kt CO₂ equivalent and by further 34.4 per cent in 2002 that was the last year of the chemical plant being operational for a full year before being closed in 2003. Over the same period Ireland was experiencing increased levels of economic growth, the knock-on effect of which was an increase in construction and therefore an increased need for building products such as cement. In the period 1990-2000 emissions from cement production (2.A.1), which are reported under 2.A

Mineral Products, increased by 92.4 per cent; from 884.00 kt CO₂ to 1,700.90 kt CO₂. Economic growth was sustained into the early years of the new millennium with associated increases in emissions from the sector, during which two new cement production plants were commissioned, with one opening in 2000 and the other in 2003. This resulted in further growth in emissions from the cement sector to reach peak of 2,374.06 kt CO₂ in 2007 (an increase of 168.6 per cent from 1990). Due to the economic recession, emissions from sector 2.A.1 decreased by 59.3 per cent between 2007 and 2011 to reach 966.27 kt CO₂. Emissions then rose to 1,111.75 kt CO₂ in 2013 (and increase of 15.1 per cent between 2013 and 2011), reflecting economic recovery.




Figure 2.8 Trend in Emissions from Industrial Processes and Product Use 1990-2013

The closure of Ireland's ammonia and nitric acid plants in 2003 and 2002 significantly changed the level of process emissions in Ireland. As a result CO₂ emissions from cement production (2.A.1) became the single major component of sector emissions and these emissions increased steadily during the period of economic growth up to 2007, the year when they reached a peak of 2,374.06 kt CO₂ equivalent (and 59.2 per cent share of the IPPU sector). Emissions from cement manufacture then decreased in line with the economic downturn, accounting for 55.8 per cent of total emissions from IPPU sector in 2008, falling to a 38.4 per cent contribution in 2011. However emissions in 2012 increased, reflecting economic recovery and were followed by a small decrease in 2013. The contribution from cement manufacture to emissions from IPPU sector in 2013 is now 40.1 per cent. Other sources of emissions within 2.A Mineral Products in Ireland are 2.A.2 Lime Production, 2.A.3 Glass Production (ceased in 2009) and 2.A.7 Other Mineral Products (Bricks, Ceramics, Soda Ash and Limestone use), which collectively accounted for 6.8 per cent of total IPPU sector emissions in 2013. The emissions from these sub-categories are small and their effect on overall trends is negligible.

The Non-Energy Products from Fuels and Solvent Use sector 2.D includes emissions from 2.D.1 Lubricant use, 2.D.2 Paraffin Wax use and indirect CO₂ emissions from 2.D.3 Solvent use. In 2013 sector 2.D accounted for 3.7 per cent of IPPU sector, having increased by 20.9 per cent from 84.42 CO₂ equivalent in 1990 to 102.06 CO₂ equivalent in 2013. However, Non-Energy Products from Fuels and Solvent Use sector with 0.2 per cent share of total national greenhouse gas emissions in 2013 does not affect the overall trend in greenhouse gases in Ireland. The largest contributor to overall emissions in this sector is indirect CO₂ emissions from 2.D.3 Solvent use, largely represented by domestic use of solvents, paint application, degreasing, dry cleaning, printing and chemical products manufacture and processing.

Emissions from 2.F Product Uses as Substitutes for Ozone Depleting Substances were estimated to be 1,329.50 kt CO₂ equivalent in 2013, compared to 34.59 kt CO₂ equivalent in 1990. This represents a 38 fold increase over the time series with the result that the contribution of this category to the sectoral total for IPPU increased from 1.1 per cent in 1990 to 47.9 per cent in 2013.

2.3.3 Trends in Agriculture (IPCC Sector 3)

The trend in emissions from the Agriculture sector is presented in Figure 2.9. Emissions of greenhouse gases from the Agriculture sector amounted to 20,735.38 kt CO₂ equivalent in 1990 and 18,964.61 kt CO₂ equivalent in 2013, a reduction of 8.5 per cent. Between 1990 and 1998, the total emissions from the Agriculture sector increased by 5.0 per cent, reflecting an increase in animal numbers and increased synthetic nitrogen use on farms. Following this peak in emission levels of 21,769.63 kt CO₂ equivalent in 1998, emissions from the sector decreased by 16.6 per cent to 18,165.33 kt CO₂ equivalent in 2011, to reach a level of emissions lower than those in 1990. The decrease post-1998 was a result of reductions in animal numbers and synthetic nitrogen fertiliser use due to reforms of the Common Agricultural Policy. Emissions in 2013 were 18,964.61 kt CO₂ equivalent, representing a 4.4 per cent increase on the total emissions in 2011. This was primarily driven by an increase in cattle number of 6.3% between 2011 and 2013.

Methane emissions from Agriculture emanate from two sectors 3.A Enteric Fermentation and 3.B Manure Management and are dependent on the type and number of livestock present on farms and in Ireland's case, the amounts are largely determined by a large cattle population. Agriculture accounted for 89.1 per cent of total methane emissions in 2013. The combined total of emissions of CH₄ from enteric fermentation and manure management 12,699.26 kt CO₂ equivalent was 12,699.26 in 1990. This increased by 7.1 per cent to reach 13,604.77 kt CO₂ equivalent in 1998 and subsequently decreased by 13.3 per cent to 11,796.04 kt CO₂ equivalent in 2013. Cattle account for 90.5 per cent of CH₄ emissions in Irish agriculture in 2013.

The emissions of N₂O from the Agriculture sector follow similar trends to those of CH₄ because cattle also largely determine the amount of nitrogen inputs to agricultural soils from synthetic fertiliser (sector 3.D) and animal manures (sector 3.B), which combined produce the bulk of N₂O emissions (92.9 per cent of total N₂O emissions in 2013). Nitrous oxide emissions in Agriculture increased from 7,636.62 kt CO₂ equivalent in 1990 by 2.3 per cent in the period 1990-1998 with emissions in 1998 totalling 7,815.41 kt CO₂ equivalent. Nitrous oxide emissions totalling 6,631.21 CO₂ equivalent in 2013 represent a reduction of 15.2 per cent on the 1998 level and 13.2 per cent on the 1990 level. Crops contribute very little to N₂O emissions in Ireland and the amount fluctuates annually in response to varying production of the relevant crops.

Carbon dioxide emissions were 537.35 kt CO₂ equivalent in 2013 compared to 399.51 kt CO₂ equivalent in 1990, a 34.5 per cent increase over the time series. 3.G Liming and 3.H Urea Application are the two subsectors responsible for CO₂ emissions from Agriculture sector accounting for 1.4 per cent share of total CO₂ emissions in 2013.




Figure 2.9 Trend in Emissions from Agriculture 1990-2013

2.3.4 Trends in Land Use, Land Use Change and Forestry (IPCC Sector 4)

The full assessment of emissions and removals in the LULUCF sector has given a new understanding of the relative contributions of sub-categories in this sector. In addition, this assessment has identified a number of land-use categories that are important in terms of either emissions or removals of CO₂. This sector is a net source of carbon in all years (Table 2.1 and Figure 2.10). This result is determined largely by the CO₂ emissions from 5.A Forest Land, which is a major carbon sink, and 5.C Grasslands and 5.D Wetlands which are major sources of emissions due to drainage of organic soils, Harvested Wood Products are a sink of carbon for all years. The complex dynamics of land-use changes between categories and the relative contributions from biomass and soils lead to highly fluctuating estimates of sectoral emissions and removals over the period 1990- 2013.

The increase in carbon stocks in living biomass in the category 5.A.1 Forest Land remaining Forest Land is the dominant removal that offsets CO₂ emissions. The Settlements and Other Land categories are comparatively less important in terms of emissions or removals but Cropland contribute significant inter annual variability due to sectoral response to external drivers such as potential economic returns for produce.




Figure 2.10 Trend in Emissions and Removals from Land Use Land-Use Change and Forestry 1990-2013

2.3.5 Trends in Waste (IPCC Sector 5)

The Waste sector remains an important source of CH₄ emissions (Figure 2.11) due to the continued dominance of landfills as a means of solid waste disposal in Ireland. Emissions from the waste sector increased by 31.2 per cent from 1,645.71 kt CO₂ equivalent in 1990 to 2,159.96 CO₂ equivalent in 2003 (peak) and then decreased by 37.8 per cent to 1,344.27 kt CO₂ equivalent in 2013. Overall, emissions in the Waste sector have decreased by 18.3 per cent from 1990 to 2013. The main contributor to trends in the Waste sector is the CH₄ emissions from municipal solid wastes (MSW) disposed of in solid waste landfills (5.A Solid Waste Disposal on Land) responsible for 82.3 per cent share of Waste emissions. The decrease in emission levels reflects increasing recovery of landfill gas for energy production and particularly through flaring at landfill sites, without which emissions in this sector would be considerably larger. Biological Treatment of Solid Waste – Composting, however small (1.9 per cent share of Waste emissions) is a growing source of emissions in Ireland since it commenced in 2001 with emission levels of 4.21 kt CO₂ equivalent, increasing to 25.10 kt CO₂ equivalent in 2013. The contribution of this sub-category to overall sectoral trends is negligible.

Since 1990 the population of Ireland increased by 31.0 per cent by 2013, giving an associated increase in the quantity of MSW produced and sustaining the amount of MSW disposed to landfills at close to 2 million tonnes per annum. However the quantities of MSW disposed of at landfills have decreased since 2007 due to lower personal consumption and increased recycling rates. Indeed total MSW disposed to landfill decreased by 63.5 per cent (and by 1.34 million tonnes) between 2007 and 2013. The proportion of organic materials in MSW has decreased from 40 per cent in 1990 to 18.3 per cent in 2013. The proportions of paper and textiles changed from 30 per cent and 10 per cent, respectively in 1990 to 10.9 per cent and 6.3 per cent, respectively in 2013, reflecting a significant diversion of paper products from landfills. This reduces CH₄ potential, as paper products are the

main source of degradable organic carbon in landfills. A major increase in the use of flares as a means of odour control in landfills in recent years offsets a large proportion of the CH₄ generated. This offset from flares and utilisation was 61.2 per cent in 2013, hence there was 10.8 fold increase in flaring and utilisation since 1996 (9.1 per cent first year of methane recovery).

Emissions from 5.C Incineration and Open Burning of Waste combined accounted for 92.48 kt CO₂ equivalent in 1990 and 43.57 kt CO₂ equivalent in 2013 a decrease of 52.9 per cent which equates to 5.6 and 3.2 per cent of total emissions from the waste sector, respectively. Emissions are reported for clinical waste incineration for all years from 1990-1997, when all hospital waste incinerators were closed. Emissions are also reported for industrial waste incineration, solvent destruction by thermal oxidisers, open burning of waste for all years from 1990-2013. The contribution of this sub-category to the overall sectoral trend is negligible.

Emissions of CH₄ and N₂O from 5.D Wastewater Treatment and Discharge accounted for 156.74 kt CO₂ equivalent in 1990 and 169.57 kt CO₂ equivalent in 2013 (8.2 per cent increase on 1990), which equates to 9.5 and 12.6 per cent of total emissions from the waste sector, respectively. The contribution of this sub-category to overall sectoral trends is negligible.




Figure 2.11 Trend in Emissions from Waste 1990-2013

2.4 Emissions of Indirect Greenhouse Gases

The total emissions of SO₂, NO_x, NMVOC and CO for the years 1990 to 2013 are summarised in Table 2.2 and Figure 2.12. As in the case of CO₂, the emissions of SO₂, NO_x and CO in Ireland are dominated by those emanating from fuel combustion activities, while the bulk of NMVOC emissions are generated by agriculture, solvent use and transport. Since 1990, substantial decreases have occurred in the emissions of SO₂ (86.2 per cent) and CO (64.8 per cent). Significant reductions have also taken

place in NO_x emissions (41.0 per cent) and emissions of NMVOC in 2013 were 33.5 per cent lower than those in 1990.

Table 2.2 Emissions of NO_x, SO₂, NMVOC and CO 1990-2013 (Tonnes)

	NO _x	SO ₂	NMVOC	CO
1990	133,903.31	183,748.93	135,272.89	350,150.61
1991	135,516.91	183,155.41	135,870.93	342,813.48
1992	143,783.87	171,090.01	132,293.82	328,398.85
1993	134,999.29	162,092.36	130,507.00	311,403.08
1994	133,516.80	176,863.37	129,243.96	295,523.70
1995	130,024.68	162,666.76	128,233.13	291,003.42
1996	132,718.23	150,770.69	130,002.17	287,868.28
1997	128,648.59	168,464.96	127,488.23	268,119.28
1998	133,606.69	179,666.13	128,251.06	272,635.99
1999	134,343.73	160,738.24	117,837.00	256,779.75
2000	137,391.35	141,985.95	112,282.26	246,818.25
2001	137,712.25	136,698.27	113,123.32	243,392.81
2002	131,036.85	103,594.48	109,770.66	232,362.51
2003	131,287.24	81,496.35	107,159.80	223,400.12
2004	134,078.22	74,120.75	105,685.23	219,234.17
2005	135,877.97	73,975.91	105,876.12	217,291.95
2006	131,912.57	62,955.09	105,501.25	200,858.86
2007	129,099.41	56,935.51	105,572.18	189,057.92
2008	116,336.99	47,499.35	100,489.58	178,501.29
2009	92,700.96	34,355.58	96,707.95	159,306.89
2010	85,014.47	28,256.89	91,401.62	146,460.76
2011	76,361.91	26,680.10	88,724.38	134,921.08
2012	78,498.25	25,191.43	88,203.53	129,123.90
2013	79,064.24	25,393.35	90,000.71	123,123.16

Total SO₂ emissions decreased from 168,465 tonnes in 1990 to 25,393 tonnes in 2013. This decrease in emissions reflects the economic downturn in recent years, reductions in the sulphur content of fuels, fuel switching and use of abatement technologies. Power stations (1.A.1a) remain the largest source of SO₂ emissions for the whole time series. However, from 2010 onwards, the commercial/residential (1.A.4.a, 1.A.4.b) sectors contributed a comparable amount to the total emissions (34.9 per cent compared to 34.2 per cent from power stations in 2013). Combustion sources in the industrial (1.A.2) sector largely account for the remainder of emissions, with a contribution of 19.1 per cent in 2013. In 1990, coal combustion accounted for 51.2 per cent of SO₂ emissions and fuel oil contributed 30.7 per cent. By 2013, the share of SO₂ emissions from coal had increased to 54.2 per cent and that from fuel oil had decreased to 11.5 per cent.

Road transport (1.A.3.b) is the principal source of NO_x emissions, contributing 45.0 per cent of the total in 2013. The reductions in NO_x emissions delivered by catalytic converters in cars and heavy-duty vehicles have been offset by large increases in vehicle numbers and fuel use in the past 10 years. This effect is exaggerated in latter years by so-called fuel-tourism, whereby a significant proportion of the automotive fuel sold in Ireland is used by vehicles in the UK and possibly to some extent in other countries. Combustion in the industrial (1.A.2) sector is another source of NO_x emissions, in 2013 accounting for 11.1 per cent of emissions, followed by power generation with

10.8 per cent share and combined commercial/residential sectors' 10.2 per cent share in the same year.

The emissions of NMVOC are determined mainly by agriculture sectors (3.B Manure management and 3.D Inorganic fertilisers) contributing 47.3 per cent share of total in 2013. Solvent use (2.D) was responsible for 23.6 per cent share and combined commercial/residential sectors produced 12.9 per cent of the 2013 total NMVOC emissions in Ireland. Technological controls for NMVOCs in motor vehicles have been more successful than in the case of NO_x, and have given a significant reduction in emissions from road transport over recent years with contributions of transport to the national total of 26.0 per cent in 1990, falling to 7.6 per cent in 2013.

Emissions of CO continue to decline, driven by major reductions due to catalysts in gasoline cars, which is the principal source of CO with 54.0 per cent share of total in 2013, and a large decrease in the use of solid fuels for space heating in the residential sector. Combined commercial/residential sectors contributed 22.9 per cent to 2013 total. Further reductions in the emissions of SO₂, NO_x and NMVOC will occur in the coming years as Ireland continues to implement programmes to comply with various EU legislation aimed at air quality and emissions control.




Figure 2.12 Trend in Indirect Greenhouse Gases 1990-2013

Chapter 3 Energy

3.1 Overview of Energy Sector

The list of activities under *Energy* in the IPCC reporting format is given in Table 3.1 below. A summary of emissions from these activities are given in Table 3.2, Figure 3.1 and Figure 3.2 below.

The *Energy* source category covers all combustion sources of CO₂, CH₄ and N₂O emissions and the fugitive emissions of these gases associated with the production, transport and distribution of fossil fuels.

Estimates are included for all emission sources that occur in the country and the required level of disaggregation is achieved for detailed completion of the CRF tables.

3.1.1 Emissions Overview

A summary of emissions from this sector is given in Table 3.2, Figure 3.1 and Figure 3.2 below.

Emissions from *Energy* accounted for 54.8 per cent and 60.8 per cent of total national emissions in 1990 and 2013, respectively. This sector accounted for 94.7 of total CO₂ emissions and 6.7 per cent of combined total CH₄ and N₂O emissions in 2013. CO₂ emissions make up 98.3 per cent of the total for the sector.

There are 16 key categories in this sector (see Annex 1 for further details) all of which are encompassed in the following categories:

- **1.A.1 Energy Industries** is a significant activity in Ireland, which peaked in 2001 corresponding to a peak in the consumption of coal and has since decreased with the increased use of natural gas and renewables. There was an increase in emissions in 2012 due to the increasing the use of cheaper solid fuels.
- **1.A.2 Manufacturing Industries and Construction** emissions peaked in 2005 with a significant drop between 2008 and 2009 due to the impact of the economic downturn. Emissions have since plateaued.
- **1.A.3.b Road Transport** liquid fuel consumption increased until it peaked in 2007 after which it declined until 2012. This corresponds to the pattern of emissions and is due to the effect of the economic downturn in Ireland and increases in biofuel use.
- **1.A.4 Other Sectors** dominated by Residential fuel combustion peaked in 2008 and showed a downward trend in the following years. Economic downturn combined with a switch from coal and peat to less carbon intensive fuels (natural gas and oil) and renewables were the reasons for the decrease in emissions.
- **1.B.2.b Fugitive Emissions from Oil and Natural Gas** emissions have decreased considerably across the timeseries due to the introduction of polyethylene pipes across the pipeline network. These are considered to result in negligible losses.

Other non-key categories in this sector include:

- **1.A.3.a Domestic Aviation** emissions peaked in 2006 after which emissions have significantly declined due to the reduction in the number of domestic flights due to the completion of the national motorway network.
- **1.A.3.c Railways** is a minor source of emissions and has remained relatively stable across the timeseries with no significant changes to the rail network in Ireland over this time.
- **1.A.3.d Domestic Navigation** emissions from this minor source steadily grew across the timeseries until peaking in 2006 after which emissions dropped due to the effect of the economic downturn.
- **1.A.3.e Other Transportation** account for emissions from pipeline transportation of natural gas.
- **1.B Fugitive emissions** include emissions from coal mining and handling and emissions from the oil and natural gas industries.

The greenhouse gases relevant to *Energy* sector are as follows:

- **Carbon Dioxide** emissions which make up 98.3 per cent of total GHG emissions from this sector and originate from all activities involving the combustion of fossil fuels. There was a significant decrease in emissions from 2007-2008 due to the economic downturn.
- **Nitrous Oxide** emissions originate from all combustion sources with emissions from road transport being the most significant source.
- **Methane** emissions originate from all combustion sources with emissions from *1.A.4 Other Sectors* and *1.B Fugitive emissions* being the most significant sources.

3.1.2 Methodology Overview

The combustion of fossil fuels accounts for the bulk of CO₂ emissions in most countries. The CO₂ emissions are quantified with reasonable accuracy as the fuel amounts are detailed in the energy balance sheets and information on their carbon contents is well established. The total amount of CO₂ released on combustion can therefore be readily ascertained. Only small amounts of CH₄ and N₂O are associated with fuel combustion activities. The emissions of these gases are generally not quantified with the same reliability as the emissions of CO₂ because the rates of CH₄ and N₂O production depend on several factors, in addition to fuel type, and consequently there is considerable uncertainty in the available emission factors for these gases.

The overall approach and methodologies used to estimate emissions in the *Energy* sector for 2013 have been updated for the first time in accordance with the new format of the 2006 IPCC GHG Inventory Guidelines. As for all years since 2005, CO₂ estimates reported under the EU Emissions Trading System (ETS) for 2013 data are used to achieve complete bottom-up results in respect of some important sub-categories in this sector for the 2015 inventory submission. This is a significant advance in terms of accuracy as the EU ETS estimates are verified and they represent a large proportion of the total emissions from the *Energy* sector.

Ireland's energy data in the expanded energy balance sheets (Table 4.B of Annex 4) are well disaggregated according to fuel and sector for the purposes of calculating emissions in the IPCC Level

3 source categories in a top-down approach. Supplementary sources of information facilitate the use of bottom-up methods in some important sub-categories and they provide greater detail in the overall fuel-sector matrix, making it more compatible with the inventory reporting format required for the Sectoral Approach. The simple calculation spread sheet given in Table 3.1.1 of Annex 3.1.A shows how the emissions from combustion sources are computed for the year 2013 using the activity data and emission factors described below. The complete allocation to IPCC Level 1 source categories is readily achieved from this compilation, as shown in Table 3.1.2 of Annex 3.1.A. The correspondence between the national disaggregation of sources and IPCC combustion source categories is given in Table 3.1.3 of Annex 3.1.A.

All CO₂ emission factors for fuel combustion in the present submission, except in the case of biomass, are country-specific values, regardless of methodological tier used, which are determined directly from information on the carbon contents and net calorific values of the fuels used in stationary and mobile sources. Information on CO₂ emission factors, net calorific value is available for liquid, solid and gaseous fossil fuels in Table 4.C of Annex 4. The CO₂ emission factor for natural gas takes into account the increasing contribution of imported gas in the national total given by the energy balance. The importation of natural gas from the UK began around 1993 and imported gas accounted for 96.1 per cent of the total in 2013. The CO₂ emission factor appropriate to the split between domestic and imported natural gas, which is more carbon intensive, is now used for all years from 1993 to 2013.

The annual returns to the EPA's Climate Resource and Research Programme (CRRP) by participants in the EU Emissions Trading Scheme under Directive 2009/29/EC (EP and CEU, 2009, amending Directive 2003/87/EC so as to improve and extend the greenhouse gas emission allowance trading scheme of the Community) comprise an important source of information on CO₂ emissions and emission factors that is now fully utilised for the national inventory compilation. The fuel combustion CO₂ emission factors for solid fuels used by participants under ETS take account of the fact that a very small fraction (typically less than 1 per cent) of fuel carbon may remain un-oxidised and IPCC oxidation factors appropriate to these fuels are applied when computing the emissions under the scheme. Complete oxidation of carbon is assumed in the case of liquid and gaseous fuels. For other stationary combustion sources, where activity data are in general top-down fuel use quantities taken from the energy balance, the inventory agency adopts the approach that no specific allowance is needed for un-oxidised carbon in the calculation of CO₂ emissions. Default CO₂ emission factors from new 2006 GHG IPCC guidelines are used only for biomass, which almost invariably refers to wood and wood wastes. For stationary sources and all mobile sources except road traffic, Ireland relied largely on the default emission factors for CH₄ and N₂O available from the CORINAIR/EMEP Emission Factor Guidebook (McInnes, 1996 and Richardson, 1999) in preparing the submissions up to 2009.

A comprehensive internal review of CH₄ and N₂O emission factors was undertaken in 2009 (Annex C NIR 2011), which led to substantial revisions of these emission factors across stationary combustion sources in general so that they now conform to the latest available 2006 IPCC guidelines values.

Table 4.B of Annex 4 shows the national energy balance sheets for 2013, published by Sustainable Energy Authority of Ireland (SEAI), which form the key activity data for the *Energy* sector. The energy statistics are compiled using a combination of top-down and bottom-up methods and the 2013 example indicates the same form of expanded balance sheet as previously used for all years from 1990 to 2012.

A full description of the stakeholders and the process used to compile energy statistics in Ireland is described in Annex 4.A. The balance sheets reflect revisions made by SEAI over recent years following a programme to harmonise national energy balances in compliance with the needs of the International Energy Agency (IEA) and EUROSTAT and to facilitate their wider use nationally. The energy balances incorporate additional sectoral disaggregation specific to the needs of the greenhouse gas inventory, following close collaboration between SEAI and the inventory agency. The annual submission of up-to-date energy balances from SEAI to the inventory agency is one of the primary data inputs covered by MOU in Ireland's national system. A fully consistent set of energy balance sheets for the years 1990-2013 underlies the estimates of emissions for *Energy* in this submission.

Table 3.1 Level 3 Source Methodology for Energy

1. Energy	CO₂	CH₄	N₂O
A. Fuel Combustion			
1. Energy Industries			
a. Public Electricity and Heat Production*	T1, T3*	T1, T2	T1, T2
b. Petroleum Refining	T3	T1	T1
c. Manufacture of Solid Fuels and Other Energy Industries	T3	T1	T1
2. Manufacturing Industries and Construction			
a. Iron and Steel	T2	T1	T1
b. Non-Ferrous Metals*	T1, T2*	T1	T1
c. Chemicals*	T1, T2*	T1	T1
d. Pulp, Paper and Print	T2	T1	T1
e. Food Processing, Beverages and Tobacco*	T1, T2*	T1	T1
f. Non-metallic minerals*	T1, T2, T3*	T1	T1
g. Other*	T1, T2, T3*	T1	T1
3. Transport			
a. Domestic Aviation	T3	T2	T2
b. Road Transportation*	T2, T3*	T3	T3
c. Railways	T2	T1	T1
d. Domestic navigation	T2	T1	T1
e. Other transportation	T2	T1	T1
4. Other Sectors			
a. Commercial/Institutional*	T1, T2*	T1	T1
b. Residential*	T2*	T1*	T1
c. Agriculture/Fishing*	T2*	T1	T1
5. Other	IE	IE	IE
B. Fugitive Emissions from Fuels			
1. Solid Fuels			
a. Coal mining and handling	NO	T1	NO
b. Solid Fuel Transformation	NO	NO	NO
c. Other	NO	NO	NO
2. Oil and Natural Gas			
a. Oil	NO	T1	NA
b. Natural gas*	NO	T2*	NA
c. Venting and Flaring	NO	T1	NA
d. Other	NO	NO	NO

* Key Category

T1, T2, T3: Tier 1, Tier 2, Tier 3 as described in the 2006 IPCC Guidelines

NE: emissions not estimated;

NO: activity not occurring; NA: not applicable (emissions of the gas do not occur in the source category); E: emissions included elsewhere

Table 3.2 Emissions from Energy 1990-2013

		Gas	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
1.A.1.a	Public Electricity and Heat Production	CO2, CH4, N2O	kt CO2 eq	10953.9	11441.0	12108.5	12126.2	12448.5	13132.9	13852.0	14490.6	14813.8	15498.3	15754.4	16893.9	15934.5	15222.1	14836.2	15244.8	14527.0	14055.8	14155.1	12610.6	12895.1	11556.5	12356.3	10952.9
1.A.1.b	Petroleum Refining	CO2, CH4, N2O	kt CO2 eq	168.7	166.7	171.8	172.7	178.3	181.3	179.4	218.7	247.8	223.9	274.8	321.5	340.3	337.9	337.0	411.9	377.1	360.8	367.5	315.4	310.5	285.4	313.5	294.5
1.A.1.c	Manufacture of Solid Fuels and Other Energy Industries	CO2, CH4, N2O	kt CO2 eq	100.5	76.5	65.3	62.6	72.2	69.4	72.2	51.6	80.0	77.9	87.2	118.8	145.6	166.0	162.2	110.1	120.2	114.1	124.1	145.5	121.3	93.3	104.8	122.7
1.A.2.a	Iron and Steel	CO2, CH4, N2O	kt CO2 eq	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	16.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.2.b	Non-Ferrous Metals	CO2, CH4, N2O	kt CO2 eq	811.5	828.6	968.5	995.7	1264.0	1207.4	1160.3	1265.9	1279.3	1375.2	1437.9	1305.3	1295.9	1341.3	1433.5	1152.6	1459.2	1541.7	1544.4	1227.7	1519.0	1484.1	1487.0	1466.3
1.A.2.c	Chemicals	CO2, CH4, N2O	kt CO2 eq	411.4	345.1	338.4	351.0	366.1	357.2	334.7	343.8	346.8	397.9	485.1	490.1	460.8	444.4	461.6	450.8	366.6	324.4	330.0	283.8	276.0	256.4	245.4	233.3
1.A.2.d	Pulp, Paper and Print	CO2, CH4, N2O	kt CO2 eq	28.5	29.3	36.1	44.0	53.2	62.6	64.3	73.3	83.6	88.9	102.7	97.6	88.1	81.9	79.0	50.4	32.6	12.4	22.2	23.3	21.3	18.1	16.6	16.0
1.A.2.e	Food Processing, Beverages and Tobacco	CO2, CH4, N2O	kt CO2 eq	1021.4	1094.8	1010.9	1079.2	1140.4	1175.6	1141.4	1263.6	1300.3	1349.7	1608.6	1605.1	1361.5	1214.6	1062.3	1296.3	1203.4	1107.7	1125.8	1076.2	983.8	818.3	808.9	857.7
1.A.2.f	Non-metallic minerals	CO2, CH4, N2O	kt CO2 eq	938.5	847.1	556.1	619.6	448.6	505.3	560.3	553.6	513.8	502.6	720.8	657.0	920.4	1379.0	1784.7	1923.2	1806.1	1871.6	1677.9	1104.8	921.1	832.8	895.3	921.8
1.A.2.g	Other	CO2, CH4, N2O	kt CO2 eq	734.1	913.1	842.3	880.8	953.8	1023.1	905.3	1033.9	1049.4	1079.7	1270.9	1427.8	1179.9	1052.7	873.0	997.6	884.8	953.9	954.0	778.2	775.9	756.9	723.7	735.8
1.A.3.a	Domestic Aviation	CO2, CH4, N2O	kt CO2 eq	51.7	46.9	46.5	40.0	41.6	48.9	52.3	54.9	60.7	68.8	74.4	73.9	73.3	76.0	71.9	65.4	77.3	71.5	67.2	55.2	41.0	19.3	11.5	10.2
1.A.3.b	Road Transportation	CO2, CH4, N2O	kt CO2 eq	4786.0	4975.8	5411.0	5403.4	5657.6	5887.2	6889.0	7297.9	8659.8	9317.1	10366.0	10830.8	11034.2	11205.3	11858.1	12555.0	13184.6	13839.9	13085.6	11897.1	10984.0	10735.4	10365.6	10598.2
1.A.3.c	Railways	CO2, CH4, N2O	kt CO2 eq	148.9	144.6	129.7	142.3	134.1	124.5	145.1	139.9	144.1	138.6	137.6	150.2	131.4	145.1	153.0	136.6	136.6	147.7	156.5	137.4	136.3	136.5	132.0	131.5
1.A.3.d	Domestic navigation	CO2, CH4, N2O	kt CO2 eq	85.8	82.6	92.1	92.1	104.8	92.1	105.0	108.1	117.8	130.6	152.7	152.6	162.0	174.6	227.1	211.2	250.1	197.5	204.7	199.5	200.1	173.7	183.8	177.4
1.A.3.e	Other transportation	CO2, CH4, N2O	kt CO2 eq	62.9	64.4	65.0	43.3	36.5	118.7	130.3	101.0	65.5	93.7	57.8	104.0	105.3	108.6	120.2	153.3	153.2	132.0	147.5	152.5	166.7	155.3	143.8	150.4
1.A.4.a	Commercial /Institutional	CO2, CH4, N2O	kt CO2 eq	2331.6	2368.7	2316.7	2328.2	2517.9	2259.7	2339.2	2359.4	2316.5	2438.7	2476.9	2536.6	2462.6	2574.3	2382.4	2573.2	2422.3	2494.3	2704.1	2396.5	2393.8	2171.7	2191.4	2029.1
1.A.4.b	Residential	CO2, CH4, N2O	kt CO2 eq	7523.7	7565.9	6717.8	6667.0	6496.6	6452.1	6576.7	6235.9	6744.8	6377.9	6462.6	6732.3	6658.6	6812.6	6992.5	7272.0	7157.5	6928.5	7521.6	7467.0	7800.9	6609.8	6232.8	6396.4
1.A.4.c	Agriculture/Fishing	CO2, CH4, N2O	kt CO2 eq	730.6	758.7	769.3	772.8	878.1	1008.1	811.4	839.5	832.5	878.1	909.8	920.3	923.8	927.3	888.7	953.6	914.2	868.0	939.2	796.6	753.5	721.9	688.5	599.6
1.B.1.a	Coal mining and handling	CH4	kt CO2 eq	55.6	44.9	40.7	37.6	35.3	33.3	31.5	30.1	29.0	27.9	27.0	26.2	25.4	24.7	24.1	23.5	23.0	22.5	22.1	21.7	21.2	20.9	20.5	20.1
1.B.2.a	Oil	CO2, CH4	kt CO2 eq	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.4	0.3
1.B.2.b	Natural gas	CO2, CH4	kt CO2 eq	156.1	151.6	146.2	145.0	140.4	135.9	131.1	125.6	109.6	144.7	101.3	165.2	82.1	745.5	78.5	67.4	55.8	70.9	61.0	42.0	37.3	32.7	28.3	23.3
Total Energy				31,118	31,963	31,850	32,020	32,985	33,892	35,498	36,604	38,812	40,227	42,525	44,626	43,403	44,034	43,826	45,649	45,152	45,116	45,211	40,731	40,359	36,880	36,950	35,738




Figure 3.1 Total Emissions from Energy by Category, 1990-2013




Figure 3.2 Total Emissions from Energy by Gas, 1990-2013

3.1.3 Quality Assurance and Quality Control

Extensive QA/QC procedures have again been followed for the Energy sector during the present reporting cycle by fully implementing the plan that underpins Ireland's formal national system. The inventory agency continues to apply a system of quality control checks and documentation spread

sheets to the front of all calculation workbooks. These workbooks correspond directly to the disaggregation given by the CRF sectoral background data tables and are designed so that calculations may be made on a time-series basis, rather than by individual year. This increases efficiency in the use of the time-series energy data provided by SEAI and allows for rapid recalculation and checking across the time-series and facilitates the transfer of the output emission estimates and energy quantities to the CRF Reporter software. Additional summary sheets are used for aggregation to various levels to provide full cross-checking with completed CRF tables for any year.

The quality checks at inventory level build on the extensive upgrading and quality control of energy balances completed by SEAI in recent years. This work, together with further collaboration with inventory experts and thorough evaluation of the SEAI role in relation to the national system and QA/QC procedures, has resulted in substantial improvements that are now taken into account in the emissions for *Energy* for the years 1990 through 2013 included in the present submission.

In recognition of its role as a key data provider, SEAI is continuing to develop its own procedures to ensure that energy balances fully harmonised with Eurostat and IEA requirements are made available in a timely manner to facilitate the annual reporting of greenhouse gas emissions estimates. Arrangements have been established whereby the bottom-up energy data reported to the EPA for individual enterprises in all relevant energy-use sectors covered by the EU emissions trading scheme may be reconciled at an early stage with the corresponding top-down information collected by SEAI. This procedure aims to progressively minimise differences between the energy amounts reported by SEAI and that supplied for particular sub-categories and fuels.

The incorporation of the ETS data in the *Energy* sector for the last several submissions is again considered an important step towards improved reliability and accuracy of the estimates for categories 1.A.1 and 1.A.2. Thorough checking of this input is achieved in collaboration with colleagues in the Climate, Resource and Research Programme (CRRP) of the EPA, which acts as the competent authority for the ETS in Ireland. Following receipt of the raw ETS data from CRRP, the inventory experts allocate the CO₂ estimates and corresponding energy amounts to the appropriate sub-categories for CRF reporting and then return the compilation to the CRRP contact person for final checking and accounting of any amendments following the ETS verification process. This ensures that where ETS emissions estimates cover a category completely, such as in 1.A.1, the verified CO₂ values are transferred directly to the national inventory and consistency of results is guaranteed. In the case where the CO₂ estimates from ETS do not completely cover the category, as for 1.A.2, the benefit is realised as better information on fuels and more representative emission factors, which improves the top-down estimates of emissions obtained using the energy balance.

3.2 Emissions from Fuel Combustion (1.A)

3.2.1 Comparison of the Sectoral Approach with the Reference Approach

Following the methods decision tree of the 2006 IPCC guidelines for combustion sources, the information in Table 4.B of Annex 4 allows for the full application of the two available IPCC methods for emission sources in *Energy*, i.e. the Sectoral Approach and the Reference Approach.

The Sectoral Approach uses the detailed sectoral breakdown of fuel consumption by all end users as the basis of the calculations for CO₂, CH₄ and N₂O. The relevant activity data are represented by the disaggregated entries below TPER (Total Primary Energy Requirement) in Table 4.B of Annex 4. A

combination of top-down and bottom-up methods is used in the sectoral application of the national statistics on fuel consumption to derive the emission estimates in the various sub-categories.

The IPCC Reference Approach is a top-down methodology for CO₂ that estimates emissions by accounting for the overall production of primary fuels, the external trade in primary and secondary fuels, stock changes and for the carbon that may enter long-term storage in non-energy products and feedstocks.

It can be used to report national emissions in cases where the detailed activity data required for the Sectoral Approach are not available but it is more usually applied for verification of the results of the latter for those countries that have the information to apply both methods.

The Reference Approach is used in Ireland as a verification procedure for CO₂ emissions from fuel combustion activities. The calculation sheet for the Reference Approach (Table 1.A (b) of the 2013 CRF) is reproduced as Table 3.1.4 of Annex 3.1.A of this report. The apparent consumption of fuels, the basic activity data in this case, is determined as:

Apparent Consumption = Production + Imports - Exports - International Bunkers - Stock Changes

where production applies only to primary fuels.

The default value of 1.00 is used for the proportion of carbon stored in paraffin wax, lubricants, bitumen and white spirit as outlined in CRF table 1.A(b). Ireland's only oil refinery is a small hydroskimming refinery where there is no production of other petroleum products normally used for non-energy purposes, such as bitumen, lubricants, plastics and asphalt. The associated emissions with the non-energy use of these fuels are presented in section 3.2.3 and the IPPU sector, chapter 4 of this report.

The expanded SEAI energy balance sheets now record the import of some of these products, thereby allowing improved completeness in the Reference Approach estimation of CO₂ emissions and carbon storage.

A significant amount of natural gas feedstock was traditionally used in ammonia production in Ireland but the company closed in 2003 and there is consequently no feedstock use of natural gas since then.

The national energy consumption and CO₂ emissions estimates obtained using the Sectoral Approach usually differ to some extent from the corresponding values resulting from the Reference Approach. According to the UNFCCC reporting guidelines, discrepancies between the approaches (greater than 2 per cent) should be investigated and documented to see whether they indicate systematic underestimation or overestimation of energy consumption by one or other of the methods.

The differences in the Reference Approach for 2013, energy use (excluding non-energy use, reductants and feedstocks) and CO₂ emissions were; zero and 0.48 per cent, respectively than in the Sectoral Approach. The differences between the two approaches for liquid, solid, gaseous, peat and other fuels are presented in Table 3.1.5 of Annex 3.1.A and CRF Table 1.A(c) for 2013.

3.2.2 International Bunker Fuels

The memo items of the IPCC reporting format refer to activities for which the emissions are excluded from national totals. The use of fuels in international aviation and marine bunkers is the most important of these activities.

Some of the associated emissions, particularly CO₂ emissions from international aviation, are increasing very rapidly and it is therefore important that they are closely monitored for comparison with other sources and for the benefit of the international organisations that will have to develop control strategies for them in the future.

The national energy balance sheets include marine bunkers and international aviation as specific items and the emissions may be calculated directly. The allocation of fuels to marine bunkers in the national energy balance is achieved on the basis of particular tax and excise rates applicable to the sale of such fuels.

The allocation of jet kerosene use to international aviation (bunker fuel) is done by subtracting jet kerosene used in civil aviation estimated by the inventory agency from total jet kerosene fuel sales compiled by SEAI. In 2013, the amount of jet kerosene fuel allocated to domestic aviation was 0.8 per cent of the total recorded under air transport in the energy balance.

Emissions of CH₄ and N₂O have been estimated for all years for fuel used in marine bunkers. Emissions factors from the Revised 1996 IPCC guidelines of 7 kg/TJ and 2 kg/TJ, for CH₄ and N₂O respectively, have been used to estimate emissions.

3.2.3 Feedstocks and Non-energy Use of Fuels

This category includes fossil fuels used for non-energy purposes; without the combustion and oxidation process.

There are a number of fuel types applicable in Ireland:

- Lubricants – IPCC default oxidation value of 0.2 is used, see category 2.D.1;
- Bitumen – IPCC default value of 1.0 is used for the proportion of carbon stored;
- Paraffin wax – IPCC oxidation value of 0.9 is used for candles and 0.2 for all other paraffin wax, see category 2.D.2;
- White spirit – IPCC default value of 1.0 is used for the proportion of carbon stored;
- Natural Gas– a significant amount of natural gas feedstock was used in ammonia production from 1990-2003.

Emissions from the non-energy use of fossil fuels have been included in the Industrial Processes and Product Use sector, CRF Category 2.D (Chapter 4 of this report).

Table 3.3 Allocated CO₂ emissions from fuel used for non-energy purpose

CO ₂ emitting process	CRF Category (Sectoral Approach)	Type of fuel used for non-energy purpose such as feedstock	Emission factor (t C/TJ)	Net Calorific Value (TJ/ktonne)
Automobile engine oils	2.D.1	Lubricants	20.00	42.29
NA*	NA (RA)	Bitumen	22.00	37.70
Candle production and other	2.D.2	Paraffin wax	20.00	40.20
Ammonia production	2.B.1	Natural Gas	14.98	49.00
Indirect CO ₂ from NMVOC	NA (RA)	White spirit	20.00	44.00

*All carbon is stored

3.2.4 Energy Industries (1.A.1)

The emission categories relevant under 1.A.1 *Energy Industries* are: 1.A.1.a *Public electricity and heat production*, 1.A.1.b *Petroleum refining*, 1.A.1.c *Manufacture of solid fuels and other energy industries*.

3.2.4.1 Public electricity and heat production (1.A.1.a)

3.2.4.1.1 Category Description

The emissions data from a total of 19 electricity generating stations are the basis for compiling the results in this important category. The verified CO₂ estimates reported by the ETS participants were used directly and the corresponding fuel use, as given in the national energy balance, was used to estimate CH₄ and N₂O emissions using the appropriate IPCC emission factors mentioned in the previous section.




Figure 3.3 Emissions from 1.A.1.a Public Electricity and Heat Production 1990-2013

3.2.4.1.2 Methodological Issues

The CO₂ emissions are obtained from AEMs 2013 (Annual Emission Installation Reports (AIERs) for previous years 2008-2012) are estimated by ETS operators using tier 3 methodologies in accordance with the monitoring and verification guidelines for combustion activities set down in Decision 2004/156/EC (EP and CEU, 2004), which were developed for the implementation of Directive 2003/87/EC and amended by Directive 2009/29/EC.

Three types of biomass fuel are also used in this sub-category which are not reported under ETS; landfill gas (LFG) used in engines at managed waste disposal sites, wood biomass co-fired at peat power plants and municipal solid waste (MSW) used in a waste to energy plant which was commissioned in 2011. Detailed information on these biomass fuels and the non-biogenic MSW fraction are shown in Annex 3.1.A Tables 3.1.1-3.

The bottom-up CO₂ emission estimates received from the ETS participants, along with the emissions of CH₄ and N₂O estimated by the inventory agency, are aggregated on the basis of six main fuel types (peat, coal, oil, natural gas, biomass and other fuels (MSW)) in the calculation sheets shown in Annex

3.1.A and also by solid, liquid, gaseous, biomass and other fuels for reporting in the CRF. However, the corresponding energy use as reported in the CRF is taken from the national energy balance, rather than from the ETS returns, following Ireland's established practice to always reflect the published official national energy data in emission inventories. The resulting implied emission factors (IEFs) appearing in the CRF may have large inter-annual fluctuations, which are often identified in the UNFCCC review process. These IEF fluctuations are a consequence of the difference between energy data reported to the inventory agency through the ETS and that reported by SEAI in the national energy balance. The inventory agency is working closely with SEAI to minimise these differences so that the IEF will better represent the reported emissions and activity data in future years. The inventory agency meets with SEAI regularly to resolve any issues regarding the national energy balance pending the outcome of the latest UNFCCC review. The national energy balance data now corresponds more closely to the data supplied directly to the inventory agency from ETS returns in sub-category 1.A.1.a which can be seen by the IEF comparison for liquid and solid fuels for this sub-category in Tables 3.1.6 and 3.1.7 of Annex 3.1.A.

3.2.4.1.3 Uncertainties and Time-series Consistency

The ETS data almost fully (except WtE MSW incineration and LFG used for energy production) cover sub-category 1.A.1.a and that these estimates match those reported separately under parallel arrangements that have been in place for many years for the same plants, it is assumed that time-series consistency is not seriously affected and that there is no impact on the emission trend from using the ETS data.

Where higher tier methods are used for combustion sources, such as those covered by ETS and road transport, the activity data uncertainty estimates are those indicated for the tier concerned. Accordingly, low estimates of uncertainty apply to the activity data for category 1.A.1.a. Country-specific CO₂ emission factors are used for all combustion sources, which gives a basis for assigning the uncertainties for emission factors while again taking into account the applicable tiers. Uncertainties in the emission factors for CH₄ and N₂O released from combustion sources are high and not well established quantitatively. For CH₄ and N₂O emission factors for combustion categories, the most up-to-date IPCC publications are used and an indicative uncertainty of 50 per cent is used for both gases.

3.2.4.1.4 Category-specific QA/QC and verification

These methods involve a rigorous accounting of fuel consumption and detailed information on fuel properties based on fuel sampling protocols agreed in the greenhouse gas emission permits for each installation and the application of specific emission factors for each fuel determined by accredited laboratories. The summarised CO₂ emissions compiled in the ETS database according to fuel type for all installations that constituted sub-category 1.A.1.a in 2013 are aggregated to report the CO₂ emissions for this category.

The implementation of the ETS incorporates two layers of verification. The operator's report for the installation is verified independently in accordance with requirements specified in Directive 2009/29/EC before being submitted to the competent authority. This verification assesses whether the report contains omissions, misrepresentations or errors that lead to material misstatement of the reported information. Verification undertaken by the competent authority involves resolution of issues identified in the verified reports through consultation and installation site visits. The CO₂ emissions estimates compiled through ETS for sub-category 1.A.1.a are cross-checked with a

separate long-standing data flow to the inventory agency covering plant-specific emissions for electricity generating stations that are used to report on the Large Combustion Plant Directive and the Convention on Long-Range Transboundary Air Pollution. The aggregated CO₂ emissions reported in the latter data-flow correspond to the compilation available under the ETS for all years since the ETS data became available.

The rigour of the monitoring and verification process for CO₂ emissions under the ETS provide for estimates for sub-category 1.A.1.a that are more accurate and reliable than previously reported plant-specific estimates for the same source activities. The ETS estimates are available only since 2005 and the detailed information that underlies these data cannot reasonably be acquired by the inventory agency for historical years of the relevant UNFCCC time-series. As such, the application of the improved methodology introduces a degree of inconsistency in the time-series that is unavoidable in this instance.

3.2.4.1.5 Category-specific Recalculations

Recalculations undertaken in the Public Electricity and Heat Production 1.A.1.a sector for the years 1990-2012 summed up to 89.12 kt CO₂ eq lower emissions in 1990-2012 time-series than those reported in 2014 submission for the same period but the impact on the total was negligible. Recalculations were due to revised default emission factors to bring them in line with the 2006 IPCC guidelines values for the following gases and fuels: CO₂ for LFG and biomass; CH₄ for diesel; N₂O for diesel and LFG. Additionally the annual plant-specific CO₂ emission factors for waste incineration were revised for 2011 and 2012.

3.2.4.1.6 Category-specific Planned Improvements

CO₂ from this sector, which accounts for 98.8 per cent of this category emissions, are accurately quantified and there is therefore little scope for further improvement in the inventories as delivered in the 2015 submission.

3.2.4.2 Petroleum Refining (1.A.1.b)

3.2.4.2.1 Category Description

The Annual Emission Monitoring Reports (AEM Reports) submitted online (via ETSWAP EPA application) by ETS participants in respect of their CO₂ emissions and fuel combustion in 2013 for the first time under new Directive 2009/29/EC were used to report the inventory for this category. The emissions data from one oil refinery in 1.A.1.b is the basis for compiling the results in this category.

3.2.4.2.2 Methodological Issues

One small oil refinery accounts for the emissions reported under 1.A.1.b *Petroleum Refining*.

The reported CO₂ emissions are those available from the ETS database. These emissions are estimated using tier 3 methodologies in accordance with the monitoring and verification guidelines for combustion activities set down in Decision 2004/156/EC. The emissions are estimated from the use of high-pressure gas, low-pressure gas (refinery gas), Natural Gas, LPG and small amounts of other gases as well as gasoil and residual fuel oil using country-specific emission factors. However, those fuels are aggregated in national energy balance into fewer and hence less detailed categories than fuels reported under ETS. Since activity data is derived from the energy balance and CO₂ emissions originate from ETS the resulting carbon implied emission factors fluctuate significantly. National energy balance fuel proportions need to be harmonised with those to match more accurate ETS fuel proportions in this category.

The use of residual fuel oil had been phased out at this plant in the last three years and replaced with natural gas. The CH₄ and N₂O emissions are estimated by the inventory agency using the IPCC default emission factors. Because refinery gas (high-pressure gas and low-pressure gas) and LPG account for the bulk of the emissions in 1.A.1.b in all years and the emission factors for these fuels do not fluctuate significantly, the emissions reported using ETS data are consistent with the annual estimates for historical years.

3.2.4.2.3 Uncertainties and Time-series Consistency

The ETS results fully cover sub-category 1.A.1.b and these estimates match those reported separately under parallel arrangements that have been in place for many years for the same plant, hence it is assumed that time-series consistency is not seriously affected and that there is no impact on the emission trend from using the ETS data.

Low estimates of uncertainty apply to the activity data for category 1.A.1.a. Country-specific CO₂ emission factors are used for all combustion sources, which gives a basis for assigning the uncertainties for emission factors while again taking into account the applicable tiers. Uncertainties in the emission factors for CH₄ and N₂O released from combustion sources are high and not well established quantitatively. For CH₄ and N₂O emission factors for combustion categories, the most up-to-date IPCC publications are used and an indicative uncertainty of 50 per cent is used for both gases.

3.2.4.2.4 Category-specific QA/QC and verification

As described in section 3.2.4.1.4.

3.2.4.2.5 Category-specific Recalculations

Revised default carbon emission factor (2006 IPCC guidelines) value for refinery gas applied to the pre-ETS period between 1990 and 2004 resulted in recalculations for the same years. As a result the CO₂ emissions in Petroleum Refining 1.A.1.b category decreased by 310.04 kt (8.0 per cent on average) in the combined 15 years before 2005. The revised proportion of refinery gas to natural gas in the national energy balance in the three years, 2010-2012, impacted emissions of CH₄ and N₂O which together for the three years decreased by 8.0 per cent on average but the effect on total sector's emissions was negligible.

3.2.4.2.6 Category-specific Planned Improvements

Emissions of CO₂ from this sector, which accounts for 99.9 per cent of this category's emissions, are accurately quantified and there is therefore little scope for further improvement in the inventories as delivered in the 2015 submission.

3.2.4.3 Manufacture of Solid Fuels and Other Energy Industries (1.A.1.c)

3.2.4.3.1 Category Description

The Annual Emission Monitoring Reports (AEM Reports) submitted online (via ETSWAP EPA application) by ETS participants in respect of their CO₂ emissions and fuel combustion in 2013 for the first time under new Directive 2009/29/EC were used to report the inventory for this category. The emissions data from two peat briquetting plants are the basis for compiling the results in this category.

3.2.4.3.2 Methodological Issues

Emissions for *1.A.1.c Manufacture of Solid Fuels and Other Energy Industries* refer to the production of peat briquettes from milled peat in two plants.

The values for CO₂ are taken from ETS returns which are based on tier 3 methodologies in accordance with the monitoring and verification guidelines for combustion activities set down in Decision 2004/156/EC. The CH₄ and N₂O estimates are estimated by the inventory agency using IPCC default emission factors.

3.2.4.3.3 Uncertainties and Time-series Consistency

Milled peat is the principal fuel used in this sub-category. While the plant-specific annual carbon emission factor may fluctuate in response to peat quality and moisture content, both the emission factor and activity data are sufficiently well established to ensure that the emissions time-series for this sub-category is consistent.

3.2.4.3.4 Category-specific QA/QC and verification

The inventory experts continue to collaborate with colleagues managing annual ETS returns from all participants to fully consolidate and formalise data gathering in respect of this category using the prescribed monitoring and verification mechanisms to ensure full consistency with reporting of CO₂ estimates under ETS, under the Convention and MMR 525/2013.

3.2.4.3.5 Category-specific Recalculations

Revised peat consumption data in national energy balance in the years: 2005-2012 resulted in increased emissions of CH₄ and N₂O together by 7.6 per cent on average but the effect on total sector's emissions was negligible (0.02 per cent of CO₂ equivalent). Revised Global Warming Potentials (GWPs) for those two gases had also little impact on the CO₂ equivalent totals.

3.2.4.3.6 Category-specific Planned Improvements

Emissions of CO₂ from this sector, which account for 99.6 per cent of this category's emissions, are accurately quantified and there is therefore little scope for further improvement in the inventories as delivered in the 2015 submission.

3.2.5 Manufacturing Industries and Construction (1.A.2)

3.2.5.1 Category Description

The emission categories relevant under *1.A.2 Manufacturing Industries and Construction* are: *1.A.2.a Iron and Steel; 1.A.2.b Non-Ferrous Metals; 1.A.2.c Chemicals; 1.A.2.d Pulp, Paper and Print; 1.A.2.e Food Processing, Beverages and Tobacco; 1.A.2.f Non-metallic minerals* and *1.A.2.g Other*.

Figure 3.4 shows the trend in emissions from *1.A.2 Manufacturing Industries and Construction* over the period 1990-2013. The emissions from this category show a large decrease between 2008 and 2009 reflecting the impact of the recent economic downturn in Ireland particularly in the cement production sector. The overall sectoral trend of emissions has stabilised since 2011.




Figure 3.4 Emissions from 1.A.2 Manufacturing Industries and Construction 1990-2013

3.2.5.2 Methodological Issues

The expanded annual energy balance sheets published by SEAI incorporate a mapping of industrial fuel use in combustion into the CRF sub-categories *a-g* under *1.A.2 Manufacturing Industries and Construction*. This facilitates the complete disaggregation of emissions in this source category for completion of the CRF Table 1.A (a).s2.

The combustion CO₂ emissions in a variety of installations across the CRF sub-categories *1.A.2.a* through *1.A.2.g* are covered by the ETS Directive 2009/29/EC but the total CO₂ emissions in any sub-category cannot be reported for Ireland using ETS data alone, as in the case of the sub-categories under *1.A.1*.

The ETS data are instead used to compare fuel quantities reported under ETS with corresponding amounts given in the preliminary national energy balance and to determine improved country-specific emission factors that can be applied for particular fuels and sub-categories. The emissions of CO₂ are estimated by the inventory agency on a top-down basis using the agreed final energy balance activity data and country-specific emission factors as shown in Table 3.1.8 of Annex 3.1.A. The emissions of CH₄ and N₂O are estimated using the 2006 IPCC emission factors adopted following the review of emission factors referred to in section 3.1.2.

Information provided from the ETS on fuel data have been used to develop an annual country-specific CO₂ emission factor for petroleum coke since 2005. Petroleum coke is used in sub-categories and years: *1.A.2.b* (1991-2000), *1.A.2.e* (1991-2003), *1.A.2.f* (1990-2000, 2002-2013) and *1.A.2.g* (1991-2003, 2008, 2009). The 2006 IPCC emission factor of 97.5 t CO₂/TJ compares well with the year specific emission factors which vary from 92.87 to 95.13 CO₂/TJ. The average of the five years between 2005 and 2009 of yearly specific emission factors is applied to all years from 1990 to 2004, as ETS data is only available from 2005 onwards.

When the country-specific emission factor for petroleum coke is taken into account, the implied emission factors for liquid fuels in category 1.A.2.f fluctuate significantly depending on the proportion of petroleum coke in liquid fuels. It is mostly evident in sector 1.A.2.f as the biggest proportion of petroleum coke was used in this sector (50 per cent share of sector liquid fuels in the full time series apart from 2001 when petroleum coke was not used in 1.A.2.f). Other sectors with smaller proportion of this fuel to their liquid fuel totals were less affected by fluctuating CO₂ implied emission factor.

For sub-category 1.A.2.e, the largest quantities of petroleum coke are used in 2000 to 2002, giving rise to a peak in the liquid fuels implied emission factor of 79.83 t CO₂/TJ in 2001. However, the average implied emission factor for years 2004-2013 was 72.18 t CO₂/TJ as no petroleum coke is consumed since 2004.

In 1.A.2.f, the implied emission factor for liquid fuels decreases from 83.79 t CO₂/TJ in 1990 to 73.40 t CO₂/TJ in 2001 as no petroleum coke was consumed that year, to increase and reach maximum at 89.46 t CO₂/TJ in 2006 but then decreases to 83.59 t CO₂/TJ in 2010 reflecting the decline in petroleum coke use in cement production and rise again to 85.49 t CO₂/TJ in 2013.

For sub-category 1.A.2.g, the largest quantities of petroleum coke are used in 2001, giving rise to a peak in the liquid fuels implied emission factor of 80.36 t CO₂/TJ in 2001. However, the average implied emission factor for years 2004-2013 was 69.61 t CO₂/TJ almost no petroleum coke is consumed since 2004.

3.2.5.3 Uncertainties and Time-series Consistency

The ETS data partially cover category 1.A.2 and this data is provided to SEAI annually to help improve the disaggregation of fuel amounts within the sector. All emissions are estimated based on data provided in Ireland's national energy balances provided by SEAI.

Where higher tier methods are used for combustion sources, such as those covered by ETS and road transport, the activity data uncertainty estimates are those indicated for the tier concerned. Accordingly, low estimates of uncertainty apply to the activity data for category 1.A.2. Country-specific CO₂ emission factors are used for most combustion sources, which gives a basis for assigning the uncertainties for emission factors while again taking into account the applicable tiers. Uncertainties in the emission factors for CH₄ and N₂O from combustion sources are high and not well established quantitatively. For CH₄ and N₂O emission factors for combustion categories, the most up-to-date 2006 IPCC guidelines are used and an indicative uncertainty of 50 per cent is used for both gases.

3.2.5.4 Category-specific QA/QC and verification

3.2.5.5 Category-specific Recalculations

Revised fuel consumption in national energy balance for fuels and years: coal (2010-2012), diesel (2006-2012) and natural gas (2010-2012) combined with fuel reallocation between subsectors (i.e. cement sector split from other industry) and change of GWPs for CH₄ and N₂O resulted in subsector recalculations but had little impact on the totals. Sector 1.A.2 decreased in total by 231.55 kt CO₂ eq. and by 7.7 per cent on average in the full time series 1990-2012.

3.2.5.6 Category-specific Planned Improvements

There are no planned improvements for this category.

3.2.6 Transport (1.A.3)

Figure 3.5 shows the trend in emissions from 1.A.3 *Transport* over the time series. Road transport is the main driver in the trend. Overall Transport emissions have declined since 2007 reflecting the impact of the recent economic downturn in Ireland.




Figure 3.5 Emissions from 1.A.3 Transport 1990-2013

3.2.6.1 Civil Aviation (1.A.3.a)

3.2.6.1.1 Category Description

This source category includes emissions from all civil commercial use of airplanes, including private jets and helicopters. Operations of aircraft are divided into; Landing/Take-off (LTO) cycle and Cruise. All international aviation is reported as a Memo item.

3.2.6.1.2 Methodological Issues

The fuel consumption within Ireland associated with sub-category 1.A.3.a *Civil Aviation* is estimated using a Tier 3a approach (Table 3.6.2, 2006 IPCC guidelines) based on origin and destination data for domestic air travel provided by the Irish Aviation Authority (IAA), the fuel consumption rates given by the EMEP/EEA emission inventory guidebook appropriate to the type of aircraft concerned and the length of the flights within Ireland. This approach is used for all years from 2004 to 2013 where airport pair data are available.

The inventory agency receives annual flight data for all Irish airports from the IAA. These data are fully consistent with data received in previous years from the IAA since 2004. These data included all flights, domestic and international, on an origin and destination basis and by aircraft type for over 25 different Irish origin airports.

For the years 1990 to 2003, the number of flights for each airport was estimated based on domestic passenger and aircraft movement statistics as well as the relationship between all Irish airports and Dublin airport which is the principal destination of all civil flights.

For data handling purposes, the inventory agency aggregated approximately 15 small regional airport/aerodrome pairs to “Other” which account for approximately 2 per cent of all domestic flights along with nine Irish airports which account for the remaining 98 per cent of all domestic flights.

The tier 3a methodology estimates both LTO and cruise emissions based on origin and destination, flight distances and by aircraft type. The inventory agency estimated fuel consumption for the LTO and cruise phases of each flight based on 37 aircraft types using fuel consumption emission factors from the EMEP/EEA emission inventory guidebook. Table 3.1.11 of Annex 3.1.B outlines the emission factors used for LTO/cruise for fuel, CH₄ and N₂O by aircraft type. CH₄ and N₂O emission factors by aircraft type are from Table 3.6.9 of the 2006 IPCC guidelines. Table 3.1.12 of Annex 3.1.B presents implied emission factors (IEF) for fuel consumption used in the cruise phase of flights weighted by number of flights per airport.

Figure 3.6 and Table 3.1.9 of Annex 3.1.B shows the number of LTOs for each of these nine airports and all remaining airports together under “other”. Table 3.1.10 of Annex 3.1.B outlines the distance between the airport pairs in nautical miles (nm) used in estimating fuel used in the cruise phase.




Figure 3.6 Number of LTOs from Irish airports 1990-2013

3.2.6.1.3 Uncertainties and Time-series Consistency

The activity data uncertainty for this source category is considered to be very low as the data provided to the inventory agency accurately splits all flights based on airport pairs, both domestic and international. An emission factor uncertainty of 2.5 per cent is used as the data supplied to the agency identifies both aircraft and engine type.

3.2.6.1.4 Category-specific QA/QC and verification

The inventory agency completed a verification exercise comparing civil aviation flight and LTO fuel estimates for 2004 to 2010 using data sourced from Eurocontrol through the EU's Working Group 1 of the Climate Change Committee and national data. The verification exercise showed close agreement between the two datasets for the number of civil LTOs and fuel used for both LTO and cruise phases. The only significant difference in the two datasets was for 2004 which showed Eurocontrol data 20 per cent lower than Ireland's national data. The main findings of this verification procedure are outlined in Figures 3.7 to 3.9. It is intended to repeat this verification exercise for the 2016 inventory submission.




Figure 3.7 National LTO data and Eurocontrol LTO data for 2004-2010




Figure 3.8 National LTO fuel data and Eurocontrol LTO fuel data for 2004-2010




Figure 3.9 National Cruise fuel data and Eurocontrol Cruise fuel data for 2004-2010

3.2.6.1.5 Category-specific Recalculations

Revision of jet kerosene and aviation gasoline amounts in energy balance for years 2006-2012 resulted in slight changes in CO₂ and N₂O emissions in those seven years but impact on totals was negligible.

3.2.6.1.6 Category-specific Planned Improvements

The inventory agency intends to revise fuel consumption rates for aircraft when updates to the EMEP/EEA emission inventory guidebook are available.

3.2.6.2 Road Transportation (1.A.3.b)

3.2.6.2.1 Category Description

Emissions of CO₂ reported under *1.A.3.b Road Transportation* are computed from the amounts of petrol, diesel, LPG and biofuels given under road transport in the national energy balance and country-specific emission factors for these fuels as shown in Table 3.1.1 of Annex 3.1.A.

Following the 2006 IPCC guidelines, the activity data are based on fuel sales within Ireland, even though a significant proportion of automotive fuels purchased in Ireland are used in the UK. For CO₂ emission estimates, complete oxidation of carbon content of the fuel was considered as per 2006 IPCC guidelines; however the proportion of emissions by vehicle category type was estimated using the COPERT model. The CH₄ and N₂O emissions from road traffic were estimated directly from the COPERT 4v.11.0 model (Pastramas N. et al., 2014), developed within the CORINAIR programme for estimating a range of emissions from this important source. Figure 3.10 shows the trend in emissions from *1.A.3.b Road Transport* over the time series.




Figure 3.10 Emissions from 1.A.3.b Road Transport 1990-2013

3.2.6.2.2 Methodological Issues

The COPERT 4v.11.0 model estimates emissions of CH₄ and N₂O on the basis of distance travelled using a detailed bottom-up approach (Tier 3) that accounts for such factors as fuel type, fuel consumption, engine capacity, driving speed and a range of applicable technological emission controls that may be applied on the basis of the age of the vehicle. The model is applied annually in Ireland to derive CO₂ emission proportions between vehicle categories and CH₄ and N₂O emissions estimates. The resultant 2013 emission factors have been converted to national average values per fuel type for the purpose of Table 3.1.1 of Annex 3.1.A. The COPERT 4 methodology is part of the EMEP/EEA air pollutant emission inventory guidebook for the calculation of air pollutant emissions and is consistent with the 2006 IPCC Guidelines for the calculation of greenhouse gas emissions. An overview of the methodology has been provided below, however, a detailed methodology for emissions estimation and recalculation can be obtained from a conference publication (Alam, Duffy, et al. 2015).

Data Modelling: Fleet and Mileage

Detailed information on vehicle population by type is presented in Table 3.1.13 of Annex 3.1.B. The historic vehicle fleet and mileage was recalculated from the year 1990 to 2013 from national statistics- Vehicle Bulletin of Driver Statistics (DOE, DELG, DEHLG, DOT, DOTTS, 1990-2013). The restructuring of fleet was consistent with the vehicle category structure and subsequent emissions by each fuel in given category corresponding with the 2006 IPCC guidelines. Vehicles were subsequently derived from national statistics into disaggregated level; firstly vehicle category (e.g. passenger car), then fuel technology (e.g. petrol) and subsequently engine size (e.g. <2 litre). The final split of vehicle categories was based on Emission bands using following formula for the number of vehicles in Emission band E_i:

$$N_{E_i, q} = \sum_{x=p}^{x=q} N_x$$

Where, x represents the vehicle registration year. 'i' represents, emissions Band: Pre-Euro to Euro-V.

Each vehicle class was bounded by the technology commencement year 'p' and new technology commencement year 'q' in the Table 3.4 below. The results were presented in Figures 3.10.1, 3.10.2 and 3.10.3.

Table 3.4 EURO class vehicle commencement years

Technology	Passenger car	LDV	HDV	Buses/Coaches	Moped and Motorcycle
Pre-ECE	Up to 1969	--			
ECE 15/00-01	1970-1978				
ECE 15/02	1979-1980				
ECE 15/03	1981-1985				
ECE 15/04	1986-1991				
Conventional	--	Up to 1993	Up to 1994	Up to 1993	Up to 1999
Euro-I	1992-1996	1994-1997	1995-1997	1994-1996	2000-2003
Euro-II	1997-2001	1998-2001	1998-2001	1997-2001	2004-2006
Euro-III	2002-2005	2002-2005	2002-2005	2002-2006	2007 to date
Euro-IV	2006-2010	2006-2010	2006-2010	2007-2009	--
Euro-V	2011 -to date	2011-todate	2011-todate	2010- to date	--

Note: Euro 5 will apply to passenger cars and light duty vehicles of categories and will be mandatory for vehicles registered from the 1st January 2011 or from 1st January 2012 for some vehicles. Euro 6 will apply to new vehicle registrations from 2015 (RSA, 2009)




Figure 3.10.1(a) Historic passenger car fleet in Irish transport sector, Petrol




Figure 3.10.1(b) Historic passenger car fleet in Irish transport sector, Diesel




Figure 3.10.2 (a) Historic LDV fleet in Irish transport sector




Figure 3.10.2 (b) Historic HDV fleet in Irish transport sector




Figure 3.10.3(a) Historic Buses and coaches fleet in Irish transport sector




Figure 3.10.3 (b) Historic Mopeds and Motorcycle fleet in Irish transport sector

The average mileage for each vehicle category such as petrol powered or diesel powered passenger cars, light duty vehicles and heavy duty vehicles was classified in the following equation according to Euro class split above. Some of the results were presented in the Figure 3.10.4, 3.10.5 and 3.10.6.

It is noticeable from the mileage values that fleet average for different technology and size of vehicle is degrading with each consecutive year.

$$M_{E_{i_p}^{q,Y}} = \frac{\sum_{z=l}^{z=m} M_z * N_z}{\sum_{z=l}^{z=m} N_z}$$

Where, $M_{E_{i_p}^{q,Y}}$ represents Mileage for Emission Band i (vehicles penetrated the market between year p and year q), Y is the year of calculation for the mileage where $Y=p, p+1, p+2, \dots, p+n=q$ (q=new technology commencement year). M_z and N_z represent the mileage and corresponding number of vehicles in Emissions band 'i', respectively. Subscript 'z' corresponds to the different vehicle tested numbers assigned during national car testing in the year Y for the emissions band 'i'.




Figure 3.10.4 Average Vehicle mileage for Petrol PC (2000-2013)




Figure 3.10.5 Average Vehicle mileage for Diesel LDV (2008-2013)




Figure 3.10.6 Average Vehicle mileage for HD Vehicle categories (2008-2013)

For the purpose of extrapolation of the mileage data, vehicle mileages were regressed against 34 relevant variables which were selected from World Development Indicators (WB 2013). Mileage with variable such as GDP growth (annual %) & Long-term unemployment (% of total unemployment) was found to be highly correlated, Variable influential factor was acceptable (VIF<4) and can be included

into regression models for each category of vehicle (Figure 3.10.7). The model fitting R^2 and validation R^2 were acceptable (See legend in Figure 3.10.7) apart from Diesel >2L vehicle category.

In general, Diesel passenger car category does not show high co-relation with the economic activity data, which may be related to policy-induced general switch from petrol to diesel engines in Ireland.




Figure 3.10.7 Average vehicle mileage by vehicle type (1990-2013)

Although back extrapolation was attempted according to the least disaggregated categories (i.e. LDV and HDV), no appropriate predictors were found to be correlated with the mileage data. Therefore, a model was generated with average mileage data from all LDV and HDV against GDP (constant LCU). This explains somewhat variation around the mean ($R^2 = 0.38$, Validation $R^2 = 0.38$). It is expected that with more NCT testing carried out in the next few years, more data will be available and the model will be further refined.

Using the regression model built for average mileage values for LDV and HDV, back extrapolation was conducted until as far as 1987. A ratio between the average and different classes of LDV and HDV (from 2008-2013) were derived and applied to the extrapolated data. Figure 3.10.8 presents the average mileage data from 2008-2013 as reported in NCT data, and extrapolated data from 1987-2008. Due to the lack of sufficient time series data for goods vehicles, mileage degradation earlier than 2008 cannot be accurately determined according to their starting year, and thus mileage degradation of petrol and diesel passenger cars was applied for that data as a surrogate measure in Figure 3.10.8 while inputted into the COPERT model.




Figure 3.10.8 Average vehicle mileage for LD and HD by vehicle type (1990-2013)

The largest operators of buses and coaches were taken into consideration for mileage data for the bus and coach industry. The average mileage data were available since 2006, and total fleet mileage and passenger trips were available since 1999. Passengers trips were regressed against the fleet data and average mileage data were derived since 1999. The adjusted R^2 of the trip-fleet regression models for Dublin Bus and Bus Éireann were 0.52 and 0.64, respectively. For back extrapolation, predictors applied earlier were assessed against the bus and coach mileage data. After the assessment, the best fitted models (Bus Éireann Model: variables: GDP (current US\$) & Population (Total); Adjusted R^2 : 0.89; VIF<5; Validation R^2 :0.95, and Dublin Bus Model: variables: Road sector energy consumption (% of total) & Urban population (% of total), Adjusted R^2 : 0.95; VIF<2; Validation R^2 :0.94) were developed for Dublin Bus and Bus Éireann and extrapolated results were found as shown in Figure 3.10.9. The Dublin bus mileage could be representative for all bus industry in Ireland as found from different statistics, however, the mileage of Bus Éireann for coach industry was not. Thus, the following equation was applied and results were presented in Figure 3.10.9.

$$\text{Coach mileage} = (M_B * B_F + M_B * 0.5 * P_F) / (B_F + P_F)$$

Where, M_B = Bus Éireann mileage, B_F = Bus Éireann fleet size & P_F = Fleet size of the private coaches.




Figure 3.10.9 Average Vehicle mileage for bus and coach industry (1990-2013)

The average mileage values for motorcycles were obtained from CSO, and back extrapolated using a regression model where predictor variables were: length of the total road network (in km) and Long-term unemployment (% of total unemployment) and the result is presented in Figure 3.10.10. The Model fitting R^2 was 0.59 and validation R^2 was 0.58 (Max. VIF<8). Due to insufficient data on motorcycle size and type, the average mileage in each year was applied for all mopeds and motorcycles in the COPERT model.




Figure 3.10.10 Average mileage for motorcycle (1990-2013)

Parameter selection and boundary conditions

The reported speed data by the RSA for different categories of roads were reclassified in Figure 3.10.11 below for use in COPERT based on the understanding of the road classification definition, mode classification and survey location.




Figure 3.10.11 Vehicle speeds by vehicle type and road type

Other parameters like vehicle share, fuel tank size, canister size and percentage of fuel injected vehicles, sulphur content in fuel, etc. have been followed based on the historical input from the previous year.

In the estimation of emissions, the COPERT model considers different effects on emission estimation, such as hot emissions, cold emissions, evaporation emissions, load effects, slope effect, A/C factors and Lube oil effects.

Emissions modelling using COPERT

COPERT 4 Background

COPERT 4 (COMputer Programme to calculate Emissions from Road Transport) is a Microsoft Windows application to calculate emissions from the road transport sector. It draws its origins in a methodology developed by a working group which was set up explicitly for this purpose in 1989 (COPERT 85). This was then followed by COPERT 90 (1993), COPERT II (1997) and COPERT III (1999). The current version is a synthesis of results of several large-scale activities and dedicated projects, such as:

- Dedicated projects funded by the Joint Research Centre / Transport and Air Quality Unit;
- The annual work-programme of the European Topic Centre for Air Pollution and Climate Change Mitigation (ETC/ACM);
- The European Research Group on Mobile Emission Sources (ERMES) work programme;

- The MEET project (Methodologies to Estimate Emissions from Transport), a European Commission (DG VII) sponsored project within 4th Framework Program (1996-1998);
- The PARTICULATES project (Characterisation of Exhaust Particulate Emissions from Road Vehicles), a European Commission (DG Transport) PROJECT within the 5th Framework Program (2000-2003);
- The ARTEMIS project (Assessment and Reliability of Transport Emission Models and Inventory Systems), a European Commission (DG Transport) PROJECT within the 5th Framework Program (2000-2007);
- A joint JRC/CONCAWE/ACEA project on fuel evaporation from gasoline vehicles (2005-2007).

COPERT 4 Methodology

A detailed methodology supports the software application. Relevant to GHGs methodology is called “Methodology for the calculation of exhaust emissions” (published by European Environment Agency, Guidebook 2009) and covers two GHG pollutants (CH₄ and N₂O) in four CRF categories:

- 1.A.3.b.i Passenger cars;
- 1.A.3.b.ii Light-duty trucks (< 3.5 t);
- 1.A.3.b.iii Heavy-duty vehicles (> 3.5 t and buses);
- 1.A.3.b.iv Motorcycles (and mopeds).

Exhaust emissions from road transport arise from the combustion of fuels such as gasoline, diesel, liquefied petroleum gas (LPG) and natural gas in internal combustion engines. The air/fuel charge may be ignited by a spark (‘spark-ignition’ or ‘positive-ignition’ engines), or it may ignite spontaneously when compressed (compression-ignition ‘engines’). For more detailed emission estimation methods the above four CRF categories (1.A.3.b.i-iv) are often subdivided according to the fuel used (in Irish model there are three fuel types: gasoline, diesel and LPG), and by the engine size, weight or technology level of the vehicle, giving a total of 23 vehicle categories. For certain pollutants, the emission factors for these vehicle categories can be further sub-divided according to three types of driving: “highway”, “rural” and “urban”.

CH₄ and N₂O are in the group (Group 1) of pollutants for which a detailed methodology exists, is based on specific emission factors and is covering different traffic situations (i.e. urban, rural, highway) and engine conditions. Ireland uses detailed Tier 3 method, where country specific information is available for: vehicle km and mean travelling speed per mode and vehicle technology.

In the following Tier 3 approach, total exhaust emissions from road transport are calculated as the sum of ‘hot’ emissions (when the engine is at its normal operating temperature) and emissions during transient thermal engine operation (named ‘cold-start’ emissions). It should be noted that, in this context, the word “engine” is used as shorthand for “engine and any exhaust after treatment devices”. The distinction between emissions during the ‘hot’ stabilised phase and the transient ‘warming-up’ phase is necessary because of the substantial difference in vehicle emission performance during these two conditions. Concentrations of some pollutants during the warming-up period are many times higher than during hot operation, and a different methodological approach is required to estimate the additional emissions during this period.

To summarise, total emissions can be calculated by means of the following equation:

$$E_{\text{TOTAL}} = E_{\text{HOT}} + E_{\text{COLD}}$$

where,

E_{TOTAL} = total emissions (g) of any pollutant for the spatial and temporal resolution of the application,

E_{HOT} = emissions (g) during stabilised (hot) engine operation,

E_{COLD} = emissions (g) during transient thermal engine operation (cold start).

Hot exhaust emissions depend upon a variety of factors, including the distance that each vehicle travels, its speed (or road type), its age, its engine size and its weight. The basic formula for estimating hot emissions for a given time period, and using experimentally obtained emission factors, is:

emission [g] = EF [g/km] × number of vehicles [veh] × mileage per vehicle [km/veh]

Different emission factors, numbers of vehicles and mileages per vehicle need to be used for each vehicle category and class. The time period (month, year, etc.) depends upon the application. Therefore, the formula to be applied for the calculation of hot emissions of pollutants in Group 1 (like CH_4 and N_2O), and in the case of an annual emission estimation, yields:

$$E_{HOT; i, k, r} = N_k \times M_{k,r} \times e_{HOT; i, k, r}$$

where,

$e_{HOT; i, k, r}$ = hot exhaust emissions of the pollutant i [g], produced in the period concerned by vehicles of technology k driven on roads of type r,

N_k = number of vehicles [veh] of technology k in operation in the period concerned,

$M_{k,r}$ = mileage per vehicle [km/veh] driven on roads of type r by vehicles of technology k,

$e_{HOT; i, k, r}$ = emission factor in [g/km] for pollutant i, relevant for the vehicle technology k, operated on roads of type r.

Cold starts result in additional exhaust emissions. They take place under all three driving conditions. However, they seem to be most likely for urban and rural driving, as the number of starts in highway conditions is relatively limited (in principle starts from parking lots next to highways). In principle, they occur for all vehicle categories, but emission factors are only available, or can be reasonably estimated, for gasoline, diesel and LPG cars and - assuming that these vehicles behave like passenger cars - light-duty vehicles, so that only these categories are covered by the methodology. Moreover, they are not considered to be a function of vehicle age. Cold-start emissions are calculated as an extra emission over the emissions that would be expected if all vehicles were only operated with hot engines and warmed-up catalysts. A relevant factor, corresponding to the ratio of cold over hot emissions, is applied to the fraction of kilometres driven with a cold engine. This factor varies from country to country. Driving behaviour (varying trip lengths) and climatic conditions affect the time required to warm up the engine and/or the catalyst, and hence the fraction of a trip driven with a cold engine.

Cold-start emissions are introduced into the calculation as additional emissions per km using the following formula:

$$E_{COLD; i, j} = \beta_{i, k} \times N_k \times M_k \times e_{HOT; i, k} \times (e^{COLD} / e^{HOT} |_{i, k} - 1)$$

where,

$E_{\text{COLD}; i, k}$ = cold-start emissions of pollutant i (for the reference year), produced by vehicle technology k ,

$\beta_{i, k}$ = fraction of mileage driven with a cold engine or the catalyst operated below the light-off temperature for pollutant i and vehicle technology k ,

N_k = number of vehicles [veh] of technology k in circulation,

M_k = total mileage per vehicle [km/veh] in vehicle technology k ,

$e^{\text{COLD}} / e^{\text{HOT}}|_{i,k}$ = cold/hot emission quotient for pollutant i and vehicles of k technology.

Vehicle emissions are heavily dependent on the engine operation conditions. Different driving situations impose different engine operation conditions, and therefore a distinct emission performance. In this respect, a distinction is made between urban, rural and highway driving. Different activity data and emission factors are attributed to each driving situation. Cold-start emissions are attributed mainly to urban driving (and secondarily to rural driving), as it is expected that a limited number of trips start at highway conditions. Therefore, as far as driving conditions are concerned, total emissions can be calculated by means of the equation:

$$E_{\text{TOTAL}} = E_{\text{URBAN}} + E_{\text{RURAL}} + E_{\text{HIGHWAY}}$$

where,

E_{URBAN} , E_{RURAL} and E_{HIGHWAY} are the total emissions (g) of any pollutant for the respective driving situations.

Total emissions are calculated by combining activity data for each vehicle category with appropriate emission factors. The emission factors vary according to the input data (driving situations, climatic conditions). Also, information on fuel consumption and fuel specification is required to maintain a fuel balance between the figures provided by the user and the calculations.

COPERT4 version 11.0

The changes introduced in this version of COPERT software (from previous version 10.0) and relevant to greenhouse gases included new N_2O emission factors for emission standards Euro 5 & 6 in light duty and heavy duty vehicle categories and also N_2O emission factors update for older Euro standards (1 to 4).

More details on the methods, vehicle specifications, calculation algorithms and other parameters used for calculating relevant road traffic exhaust emissions can be found in EMEP/EEA emission inventory guidebook 2013, published August 2013 (<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>).

3.2.6.2.3 Uncertainties and Time-series Consistency

Usually variations in emissions factors for CO_2 , CH_4 and N_2O in road transport sectors are less uncertain. The CO_2 emission factor uncertainty is 2.5 per cent and is subject to fuel consumption and fuel blends as per the 2006 IPCC guidelines. Uncertainties in emission factors for CH_4 and N_2O are in the range of 2 to 3 per cent and depend on a few factors including fuel composition (e.g. fuel adulteration, sulphur content), uncertainties in fleet age distribution and technical characteristics of vehicle stock, uncertainties in combustion conditions (climate, altitude), driving practices, such as speed, proportion of running distance to cold starts, or load factors, etc. These sources of uncertainty may be classified into three broad categories: fuel related, modelling parameters related

and activity data related (i.e. stock and mileage). The fuel data has been taken from national energy balance where fuel sales data is well known. The COPERT software covered most of the parameters (e.g. temperature, load factors etc.) that reduced uncertainty in emission estimations. On the other hand, vehicle stock and mileage were recalculated in this submission at the most disaggregated level of data for most of the vehicle classes and thus uncertainty associated with misclassification of data was controlled and connection of vehicle class commencement years and fleet mileage were well maintained.

A consistent time series of fuel data was obtained from the national energy balance. In addition, the historic vehicle fleet from national statistics (Vehicle Bulletin of Driver Statistics) provided a very detailed dataset which were further disaggregated with the additional information from other published sources as well as expert judgment. The final product of this process provided a consistent time series of fleet data from 1990 to 2013.

Different forms of disaggregated mileage data were available for different time series: passenger cars since 2000, LDV and HDV since 2008, bus and coaches since 2005 and mopeds and motorcycles since 2000. These datasets were back extrapolated using appropriate regression method with macro-economic variables, e.g. GDP. A consistent time series was generated for mileage data. In short time series consistency covers following points:

- The division technique applied ensured that total number of vehicles in all sub groups in each year was now consistent in the time series;
- Consistency was ensured between reported mileage data in NCT (and other sources) data and starting of EURO class vehicles;
- A time series consistency for Speed and other parameters was also ensured in order to get a consistent time series of estimated emissions.

3.2.6.2.4 Category-specific QA/QC and verification

From the year 1990 to 2001 historic fleet data were re-calculated. The process involved converting multiple tables of data from hard copies as provided by national bulletin into soft copies. Reproduced tables were then checked and ensured that all the tables are formatted correctly. Every step of analysis included cross checks with the previous years reported data as well as checks of internal consistency of the fleet distribution throughout the reporting period.

In case of vehicle mileage estimation, NCT data provided by SEAI of Ireland were analysed carefully. As the mileage data were reported at the Euro-class level, there were many missing data observed. Software applications like SPSS, R, and MS excel were for data analysis, cross checking and regression modelling.

3.2.6.2.5 Category-specific Recalculations

Recalculations in road transport category were a result of COPERT 4 software version upgrade as well as activity data revisions in the national energy balance for gasoline (2006-2012), diesel (2006-2012) and biofuels (2011-2012). In addition to those two factors, new GWPs for CH₄ and N₂O, revised fleet distribution, mileage and speed improved the accuracy of activity data and resulted in recalculations in the full time series but their overall effect on emission totals was negligible. As a result of all revisions, road transport emissions increased by 0.2 per cent on average between 1990 and 2005 and decreased by 0.7 per cent between 2006 and 2012 on average as compared to the 2014 submission.

3.2.6.2.6 Category-specific Planned Improvements

The inventory agency intends to use the latest COPERT software when available.

3.2.6.3 Railways (1.A.3.c), Navigation (1.A.3.d) and Other Transportation (1.A.3.e)

3.2.6.3.1 Category Description

Emissions from railways (1.A.3.c) are estimated for diesel used in shunting or yard locomotives, railcars and line haul locomotives. There are no coal fired steam locomotives in regular use in Ireland. Emissions from navigation (1.A.3.d) are estimated for residual oil and diesel used in all water borne transport including recreational craft.

3.2.6.3.2 Methodological Issues

The CO₂ emissions under *1.A.3.c Railways* and *1.A.3.d Navigation* are estimated using a Tier 1 approach, equations 3.4.1 and 3.5.1 from the 2006 IPCC guidelines, from the amount of oil used by these activities as recorded in the energy balance and the country specific emission factors for oil. The emissions of CH₄ and N₂O are estimated using the 2006 IPCC default emission factors adopted following the review of emission factors in 2009, referred to in 2011 NIR. Emissions factors used in these two sub-categories are presented in Table 3.5.

Table 3.5 Emission factors for Rail and Navigation

IPCC category	Fuel	CO ₂ t/TJ	Reference	CH ₄ kg/TJ	N ₂ O kg/TJ	Reference
Railways	Gasoil	73.30	CS	4.15	28.60	2006 IPCC Guidelines Table 3.4.1 Default
Navigation	Fuel Oil	76.00	CS	7.00	2.00	2006 IPCC Guidelines Table 3.5.3
Navigation	Gasoil	73.30	CS	7.00	2.00	

The emissions reported in sub-category *1.A.3.e Other Transportation* are those due to the use of natural gas at off-shore production platforms and in pipeline compressor stations. The fuel use is estimated as the difference between the value given for natural gas under own use/losses in the national energy balance (Table 4.B of Annex 4) and the amount of gas estimated to be lost from the distribution network, as reported under fugitive emissions in sub-category *1.B.2.b Natural Gas*. The country-specific emission factor for CO₂ and the default values for CH₄ and N₂O referred to in section 3.1.2 are used.

3.2.6.3.3 Uncertainties and Time-series Consistency

The uncertainties applicable to Railways, Navigation and Other Transportation are provided in Annex 2. The emission time series for 1990–2013 is consistent. Key activity data such as fuel use statistics are available for all years and are used in a consistent manner.

3.2.6.3.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to these categories. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

3.2.6.3.5 Category-specific Recalculations

Recalculations in these three combined categories were minor and mostly related to diesel consumption revisions in energy balance between 2006 and 2012. A change of CH₄ and N₂O GWPs in conjunction with activity data revisions resulted in an average of 0.1 per cent decrease (and 10.72 kt CO₂ eq. in total) in the time series 1990-2012.

3.2.6.3.6 Category-specific Planned Improvements

There are no planned improvements for this category.

3.2.7 Other Sectors (1.A.4)

3.2.7.1 Category Description

The CRF sub-category *1.A.4 Other Sectors* covers combustion sources in the commercial/institutional (1.A.4.a), residential (1.A.4.b) and Agriculture/Fishing (1.A.4.c) sectors. The residential sub-category *1.A.4.b* remains the most important source of emissions in this category in Ireland. This is evident from Figure 3.11, which shows the trend in the principal components of emissions in *1.A.4 Other Sectors* over the time series.

While the shift from carbon-intensive fuels, such as coal and peat, to oil and natural gas in *1.A.4.b* has been sufficient to maintain sectoral emissions relatively constant up to 2007, the benefits from fuel switching have been fully realised and the emissions from oil and gas are increasing in line with higher overall fuel consumption resulting from greater housing stock and population.

Emissions in *1.A.4* sector decreased from 2012 to 2013 due to milder than normal winter months.




Figure 3.11 Emissions from 1.A.4 Other Sectors 1990-2013

3.2.7.2 Methodological Issues

Table 3.1.1 of Annex 3.1.A shows the estimation of emissions for sub-category *1.A.4 Other Sectors*, using the fuel quantities as given by the energy balance (Table 4.B of Annex 4).

The inventory agency uses country-specific emission factors for CO₂, including that for petroleum coke referred to in section 3.2.5.2, and 2006 IPCC default values for CH₄ and N₂O. The energy balance provides no indication on the specific end-use of gasoil in the agricultural sector *1.A.4.c(i-ii)*. Consequently, a split based on information from agricultural experts (10 per cent stationary sources and 90 per cent mobile sources) is used by the inventory agency to distinguish between the use of this fuel in stationary and mobile combustion sources. This split has no bearing on emissions of CO₂, but it is important in relation to CH₄ or N₂O and the indirect greenhouse gases.

Emissions factors used for stationary and mobile sources in sub-category *1.A.4.c(i-ii) agriculture*, are presented in Table 3.6. No biomass is used as fuel in sub-category *1.A.4.c(i-ii) agriculture*.

Emissions from charcoal used for cooking are reported in sub-category *1.A.4.b* for all years. The quantity of charcoal used in Ireland was provided by the CSO and emission factors used for estimating emissions from this biomass fuel are presented in Table 3.7.

Table 3.6 Emission factors for Agriculture

IPCC category	Fuel	CO ₂ t/TJ	Reference	CH ₄ kg/TJ	N ₂ O kg/TJ	Reference
Agriculture Stationary (1.A.4.c(i))	Gasoil	73.30	CS	10.00	0.60	2006 IPCC Guidelines Table 3.4.1 Default
Agriculture Mobile (1.A.4.c(ii))	Gasoil	73.30	CS	4.15	28.60	2006 IPCC Guidelines Table 3.3.1 Default
Fishing (1.A.4.c(iii))	Gasoil	73.30	CS	7.00	2.00	1996 IPCC Guidelines Table 1-48 2006 IPCC Guidelines Table 3.5.3

Table 3.7 Emission factors for Charcoal use in Residential

IPCC category	Fuel	Gas	kg/TJ	Reference
Residential	Charcoal	CO ₂	112,000	2006 IPCC Guidelines Table 2.3 Chapter 2 Solid Biomass (Wood/Wood Waste)
Residential	Charcoal	CH ₄	200	2006 IPCC Guidelines Table 2.5 Chapter 2 1996 IPCC Guidelines Table 1-7 Vol III
Residential	Charcoal	N ₂ O	1	2006 IPCC Guidelines Table 2.5 Chapter 2 1996 IPCC Guidelines Table 1-8 Vol III

3.2.7.3 Uncertainties and Time-series Consistency

The uncertainties applicable to sub-category 1.A.4 Other Sectors are provided in Annex 2. The emission time series for 1990–2013 is consistent. Key activity data such as fuel use statistics are available for all years and are used in a consistent manner.

3.2.7.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to these categories. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

3.2.7.5 Category-specific Recalculations

Anthracite and lignite carbon emission factors have been adapted from the new 2006 guidelines as well as new GWPs for CH₄ and N₂O CO₂ equivalent. Those changes resulted with recalculations in sectors: 1.A.4.b and 1.A.4.c 0.6 per cent (867.57 kt CO₂ eq. in total) and 5.2 per cent (1,032.53 kt CO₂ eq. in total) increase, respectively in time series 1990-2012. On the other hand sector 1.A.4.a decreased by 1.9 per cent on average (1,092.73 kt CO₂ eq. in total) in the same period.

3.2.7.6 Category-specific Planned Improvements

It is planned to complete the time series for diesel use in Fishing 1.A.4.c *iii*, for the years 1990-2002 which are currently reported as "included elsewhere", IE in the next submission. Fuel use data is now available in Ireland's 1990-2014 energy balance.

3.3 Fugitive Emissions (1.B)

Ireland has no coal or oil industries and therefore fugitive emissions of greenhouse gases are limited to those associated with oil refining/storage, natural gas production and distribution and coal mining until 1995 (only emissions from abandoned mines are reported after 1995).

3.3.1 Coal Mining and Handling (1.B.1.a)

3.3.1.1 Category Description

Emissions from the NFR subcategory 1.B.1.a *Coal mining and handling* are reported for the first time in this submission. The national energy balance now includes coal mined in the years 1990 to 1995 when the last commercial coal mine was closed in Ireland. Ireland had no surface coal mines hence all emissions are associated with *underground mines*, CRF category 1.B.1.a.1. The CH₄ emissions from underground mines are calculated for three sub-categories:

- 1.B.1.a.1(i) Emissions from *Underground mining activities* for years 1990-1995;
- 1.B.1.a.1(ii) Emissions from *Post-mining activities* for years 1990-1995;
- 1.B.1.a.1(iii) Emissions from *Abandoned underground mines* for full 1990-2013 time series.

Only three mines were active in 1990 when production was reported at 25 kilotonnes. The same year Arigna mine closed down and production of coal in the next five years was reported at a mere one kilotonne per year between 1991 and 1995 when the last two mines: Rossmore and Castlecomer, ceased operation.

Emissions from underground mines for three activity sub-categories are presented in Figure 3.12.



Figure 3.12 Fugitive emissions from Underground Coal Mines 1990-Methodological Issues

3.3.1.2 Methodological Issues

The emission factors used in category 1.B.1.a Coal Mining and Handling and the resulting time series of fugitive CH₄ emissions are based on the 2006 IPCC guidelines default values and are presented in Tables 3.8 and 3.9.

The first two categories, *Underground mining activities* and *Post-mining activities* were applicable during the years of operation of Irish coal mines (1990-1995).

Table 3.8 Emission factors for underground mining and post-mining activities

IPCC category	CH ₄ EF	Unit	CH ₄ Conversion Factor	Unit	Reference
Underground mining activities (1.B.1.a.1(i))	10.00	m ³ /t	$0.67 \bullet 10^{-6}$	kt CH ₄ /m ³	2006 IPCC Guidelines Equation 4.1.3
Post-mining activities (1.B.1.a.1(ii))	0.90	m ³ /t	$0.67 \bullet 10^{-6}$	kt CH ₄ /m ³	2006 IPCC Guidelines Equation 4.1.4

After mining has ceased, abandoned coal mines may also continue to emit methane, hence the third category *Abandoned underground mines* is applicable for the emission time series 1990-2013. This category is based on the number of existing abandoned mines (remaining unflooded) that were closed-down within the five time-bands:

- Years 1990 – 1925;
- Years 1926 – 1950;
- Years 1951 – 1976;
- Years 1976 – 2000;
- Years 2001 – present.

In the first time band (years 1900-1925) the default lower percentage of gassy mines is zero and the consequent emissions are not occurring. In the last time band (2001-present) there were no mines in Ireland closed down within that period hence there were no emissions resulting from this time band. Emissions are calculated for the middle three time bands.

Table 3.9 Emission factors for Abandoned underground mines (1.B.1.a.1(ii))

Time band	Number of existing abandoned mines	Fraction of gassy mines (%)	CH ₄ EF	Unit	CH ₄ Conversion Factor	Unit	Reference
1926 - 1950	9	3	0.343 - 0.279	Mm ³ /year/mine	0.67	kt CH ₄ /Mm ³	2006 IPCC Guidelines, Eq. 4.1.10 Table 4.1.5 (lower value), Table 4.1.6 & Table 4.1.10
1951 - 1975	19	5	0.478 - 0.340	Mm ³ /year/mine	0.67	kt CH ₄ /Mm ³	2006 IPCC Guidelines, Eq. 4.1.10 Table 4.1.5 (lower value), Table 4.1.6 & Table 4.1.10
1976 - 2000	20	8	1.561 - 0.469	Mm ³ /year/mine	0.67	kt CH ₄ /Mm ³	2006 IPCC Guidelines, Eq. 4.1.10 Table 4.1.5 (lower value), Table 4.1.6 & Table 4.1.10

3.3.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to Coal Mining and Handling are provided in Annex 2. The emission time series for 1990–2013 is consistent. Key activity data such as coal mined and other mine statistics are available for all applicable years and are used in a consistent manner.

3.3.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to this category. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

3.3.1.5 Category-specific Recalculations

Recalculations are not applicable for the 2015 submission.

3.3.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

3.3.2 Oil and Natural Gas (NFR 1.B.2)

3.3.2.1 Category Description

Natural gas has been produced from gas fields off the south coast of Ireland since the 1970s but this source is being rapidly depleted. Substantial reserves of natural gas have been discovered off the west coast and which will soon come into production in late 2015.

3.3.2.2 Methodological Issues

Bord Gais Eireann (BGE), Ireland's gas company has assessed gas losses in the pipeline network in the context of the needs of annual inventory reporting and a long-term programme to replace cast-iron mains with polyethylene pipe in all urban areas served by natural gas. The change to polyethylene pipe is considered to result in negligible losses.

The gas company indicated that gas loss in 1995, determined as the difference between system input and metered sales, was 1.92 million therms, which equates to 4,085 tonnes of methane, when

the amounts of indigenous and imported gas and their respective properties are taken into account. This value implied a loss of the order of 0.2 per cent of total sales.

Projections made by BGE for five-year intervals from 2000 show losses decreasing to zero by 2020 on completion of the pipe replacement programme.

The BGE data continue to be used as the best available for this particular fugitive emission source. The rate of loss implied by the 1995 value and the projections is applied to give an emission for all years of the inventory time-series referred to in this report. The gas consumption recorded in the energy balance for the industrial, commercial and residential sectors is used as activity data rather than total sales and the appropriate split between indigenous and imported gas is applied for all years. The inventory agency was informed by BGE in 2004 that natural gas losses from the distribution network were so small that they could not be measured.

Only one company is involved in natural gas production in Ireland. Emissions to the atmosphere from this company's offshore gas production platforms are reported to the Department of Communications Energy and Natural Resources (DCENR) under the OSPAR Convention. Such reports have been obtained for several years in the time series and are currently covered by MOU with the inventory agency.

The available data, which relate largely to gas extraction but which also account for a small amount of flaring in some years, indicate a close relationship between emissions and the amount of gas produced. This relationship has been applied in terms of the indicative emission rates of CO₂ and CH₄ per unit of gas extracted to estimate the emissions for those years for which no reports were received. A report on emissions was supplied to the inventory agency for 2013. Fugitive CH₄ emissions from flaring in natural gas production are reported only for 1999 when a third mobile drilling unit (Glomar Arctic 3) was operating in the Kinsale field and in 2001 when a drilling vessel (Noble Ton van Langevald) was operating offshore at Kinsale. For other years in the time series, Ireland reports these fugitive emissions as "NO".

3.3.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to Oil and Natural gas are provided in Annex 2. The emission time series for 1990–2013 is consistent. Key activity data such as gas and oil statistics are available for all applicable years and are used in a consistent manner.

3.3.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to this category. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

3.3.2.5 Category-specific Recalculations

There were no recalculations in this category (other than change of GWP for CH₄). These recalculations resulted in total emissions 458.79 kt CO₂ eq. higher in the full 1990-2012 time-series.

3.3.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

3.4 CO₂ Transport and Storage (1.C)

This activity does not occur in Ireland since 1990. Emissions are reported as Not Occurring (NO).

Chapter 4 Industrial Processes and Product Use

4.1 Overview of the Industrial Processes and Product Use Sector

The list of activities under *Industrial Processes and Product Use* in the IPCC reporting format is given in Table 4.1 below. A summary of emissions from these activities are given in Table 4.2, Figure 4.1 and Figure 4.2 below.

Some of these activities are well known sources of one particular greenhouse gas, such as cement production for CO₂ or adipic acid production in the case of N₂O, while others may be more important in terms of their indirect greenhouse gas emissions, such as the use of solvents.

Major industrial processes within the chemical sector and metal production that are common to many other developed countries have never been an important part of the Irish economy. Consequently, many of the production processes listed in Table 4.1 are not relevant to the inventories of greenhouse gases in Ireland. Historically, the four key industrial sources are cement and lime production under 2.A Mineral Products and ammonia and nitric acid production under 2.B Chemical Industry. The nitric acid and ammonia plants, both operated by Irish Fertiliser Industries, ceased production in 2002 and 2003, respectively. 2.A.3 Glass Production was a relevant activity up to 2009 when production ceased. 2.A.4 Other process uses of carbonates includes emissions from ceramics, bricks and tiles, clay pipe products, soda ash use as well as limestone used to abate SO₂ emissions in peat-fired electricity generating stations.

A number of studies have been performed to improve and update the emission estimates in this sector. These continual updates ensure that the specified categories are kept up-to-date and that there are regular reviews of the assumptions and activity data availability. Improvement studies for the use of solvents include: Barry (2014), CTC (2005), Finn *et al.* (2001). Improvement studies for emissions from fluorinated gases include: Goodwin *et al.* (2013), Adams *et al.* (2005), O'Leary *et al.* (2002).

Industrial Processes and Product Use is the only sector for which emissions of HFCs, PFCs, SF₆ and NF₃ (collectively known as fluorinated gases) are reported in air emission inventories. There is no production of fluorinated gases in Ireland, but these substances are used in activities such as Ireland's electronics industry and for refrigeration and air conditioning.

All relevant sub-categories are fully covered in Ireland's inventories as shown in Table 4.1 below.

4.1.1 Emissions Overview

A summary of emissions from this sector is given in Table 4.2, Figure 4.1 and Figure 4.2 below.

Emissions from *Industrial Processes and Product Use* accounted for 5.7 per cent and 4.7 per cent of total national emissions in 1990 and 2013, respectively. This sector accounted for 100 per cent of fluorinated gas emissions (HFCs, PFCs, SF₆ and NF₃), 3.8 per cent of CO₂ emissions and 0.6 per cent of N₂O emissions in 2013.

There are two key categories in this sector (see Annex 1 for further details), which are both trend and level key categories:

- **2.A.1 Cement Production** is a significant activity in Ireland, which peaked in 2007 prior to the economic downturn in 2008.
- **2.F.1 Refrigeration and Air-Conditioning** has become a significant source in Ireland due to the growth in HFC use as replacement refrigerants across virtually all refrigeration sub-categories since 1991.

Other categories present in this sector include limestone, dolomite and other carbonate uses in:

- **2.A.2 Lime Production** emissions originated from three companies up to 1999 and two companies thereafter.
- **2.A.3 Glass Production** ceased in Ireland in 2009 prior to which the industry included the production of crystal glass, bottle glass and glass-based insulation.
- **2.A.4 Other process uses of carbonates** includes the production of bricks and roof tiles, ceramics, vitrified clay pipes, clay products, wall and floor tiles and the use of limestone to abate SO₂ emissions in peat-fired electricity generating stations.
- **2.B Chemical Industry** was a relevant activity in Ireland accounting for approximately two-thirds of the total in 1990 from the nitric acid and ammonia plants, both operated by Irish Fertiliser Industries, which ceased production in 2002 and 2003, respectively.
- **2.D Non-energy products from fuels and solvent use** is a relevant activity in Ireland due to the use of lubricants, paraffin wax and solvents. Solvent use is a significant source of NMVOC emissions, whilst lubricants and paraffin wax are minor sources of CO₂ emissions. Indirect CO₂ emissions associated with NMVOCs are included in national totals (without LULUCF with Indirect).
- **2.E.1 Integrated Circuit or Semiconductor Industry** is responsible for all emissions of PFC, as well as some emissions of HFC, SF₆ and NF₃. Emissions continue to follow the downward trend post-2000, which is due to process optimization, use of alternative chemicals, employment of alternative manufacturing processes and improved abatement systems in the sector.
- **2.G Other product manufacture and use** includes emissions of SF₆ and N₂O. The sources of SF₆ include electrical equipment, which is the most significant activity, and double glazing, medical applications, sporting goods and gas-air tracers, which are minor sources. N₂O emissions originate from Medical Application through the use of anaesthesia.

The greenhouse gases relevant to *Industrial Processes and Product Use* are as follows.

- **Carbon dioxide** emissions originate from *2.A Mineral Production* and *2.D Non-energy products from fuels and solvent use* sectors: *2.D.1 Lubricant Use* and *2.D.2 Paraffin Wax Use*. Historically, *2.B Chemical Production* has also been a source. There was a significant decrease in emissions from 2007-2009 due to the economic downturn after which emissions have been relatively stable. Indirect CO₂ emissions (not counted in *IPPU* but in the national total) originate from NMVOC emissions from sector *2.D.3 Solvents*.
- **Nitrous Oxide** emissions originate from *2.G.3 Medical Application* through the use of anaesthesia. Historically, emissions have also come from *2.B.2 Nitric Acid Production* with the plant closing in 2002.

- **HFCs** mainly originate from *2.F Product uses as ODS substitutes* and the use of these gases in refrigeration and air-conditioning systems, as well as fire protection equipment, aerosols and metered dose inhalers. Emissions have risen significantly since 1991 due to the use of HFCs as a replacement refrigerant. There is also a minor source from *2.E Electronic Industry*.
- **PFCs** are solely released from *2.E.1 Integrated Circuit or Semiconductor Industry*.
- **SF₆** emissions originate from a number of sources with the most significant being *2.E.1 Integrated Circuit or Semiconductor Industry* and emissions from *2.G.1 Electrical Equipment*. Emissions peaked in 2003 but have steadily fallen due to efficiency improvements in these two activities. Other sources of emissions include double glazing, medical applications, sporting goods and gas-air tracers.
- **NF₃** are solely released from *2.E.1 Integrated Circuit or Semiconductor Industry*.
- **CH₄** emissions are not occurring in IPPU sector.

The emission estimates clearly indicate that the combined emissions of HFC, PFC, SF₆ and NF₃ have generally increased year on year. This overall trend largely reflects the increasing use of HFCs across a range of applications (e.g. often as replacements in applications where the use of CFC and HCFCs is no longer permitted under the Montreal Protocol) and hence the presence of larger fluid banks from which operational leakage potentially occurs.

4.1.2 Methodology Overview

A summary of the Tier methods, consistent with the 2006 IPCC Guidelines, is provided in Table 4.1 below, along with a summary of the activities applicable to Ireland.

The process CO₂ emissions for the relevant source categories under *2.A Mineral Products* are largely covered by Directive 2003/87/EC (EP and CEU, 2003) on emissions trading in the EU and full use is made of this data source for the compilation of the national inventory. In general, the annual verified CO₂ emissions in respect of the installations concerned are used directly for the years covered by the EU ETS. The category-level emission factors indicated by EU ETS data are used together with the best available production data to obtain the emissions estimates for years previous to 2005.

In the chemical industry sector, emissions from 2.B.1 Ammonia production were estimated based on natural gas feedstock data from Ireland's energy statistics. Nitrous oxide emissions from 2.B.2 Nitric acid production 2.B.2 were estimated using plant data.

Emissions from 2.D.1 Lubricant use and 2.D.2 Paraffin wax use were estimated using energy data provided in Ireland's energy statistics.

Emissions of HFCs and PFCs from the 2.E.1 Integrated circuit or semiconductor industries use installation specific emissions data methodology. This is expected to give considerably more accurate emission estimates, and therefore a more certain trend with time.

Table 4.1 Level 3 Source Methodology for IPPU

2. Industrial Processes and Product Use	CO₂	CH₄	N₂O	HFC	PFC	SF₆	NF₃
A. Mineral industry							
1. Cement production	T3*	NA	NA	NA	NA	NA	NA
2. Lime production	T3	NA	NA	NA	NA	NA	NA
3. Glass production	T3	NO	NO	NO	NO	NO	NA
4. Other process uses of carbonates	T3	NA	NA	NA	NA	NA	NA
B. Chemical industry							
1. Ammonia production	T1	NO	NA	NA	NA	NA	NA
2. Nitric acid production	NA	NA	T1	NA	NA	NA	NA
3. Adipic acid production	NO	NO	NA	NA	NA	NA	NA
4. Caprolactam, glyoxal and glyoxylic acid production	NO	NO	NA	NA	NA	NA	NA
5. Carbide production	NO	NO	NA	NA	NA	NA	NA
6. Titanium dioxide production	NO	NO	NA	NA	NA	NA	NA
7. Soda ash production	NO	NO	NA	NA	NA	NA	NA
8. Petrochemical and carbon black production	NO	NO	NA	NA	NA	NA	NA
9. Fluorochemical production	NO	NO	NA	NO	NO	NO	NA
10. Other	NO	NO	NO	NO	NO	NO	NA
C. Metal industry							
1. Iron and steel production	NO	NO	NA	NA	NA	NA	NA
2. Ferroalloys production	NO	NO	NA	NA	NA	NA	NA
3. Aluminium production	NO	NO	NA	NA	NO	NA	NA
4. Magnesium production	NO	NO	NA	NA	NA	NO	NA
5. Lead production	NO	NO	NA	NA	NA	NA	NA
6. Zinc production	NO	NO	NA	NA	NA	NA	NA
7. Other	NO	NO	NO	NO	NO	NO	NO
D. Non-energy products from fuels and solvent use							
1. Lubricant use	T1	NA	NA	NA	NA	NA	NA
2. Paraffin wax use	T2	NA	NA	NA	NA	NA	NA
3. Solvent use	T1	NO	NO	NO	NO	NO	NO
E. Electronics industry							
1. Integrated circuit or semiconductor	NO	NO	NO	T2	T2	T2	T2
2. TFT flat panel display	NO	NO	NO	NO	NO	NO	NO
3. Photovoltaics	NO	NO	NO	NO	NO	NO	NO
4. Heat transfer fluid	NO	NO	NO	NO	NO	NO	NO
5. Other	NO	NO	NO	NO	NO	NO	NO
F. Product uses as substitutes for ODS							
1. Refrigeration and air conditioning	NO	NO	NO	T2*	NO	NO	NO
2. Foam blowing agents	NO	NO	NO	NO	NO	NO	NO
3. Fire protection	NO	NO	NO	T1	NO	NO	NO
4. Aerosols	NO	NO	NO	T1	NO	NO	NO
5. Solvents	NO	NO	NO	NO	NO	NO	NO
6. Other applications	NO	NO	NO	NO	NO	NO	NO
G. Other product manufacture and use							
1. Electrical equipment	NO	NO	NO	NO	NO	T1	NO
2. SF ₆ and PFCs from other product use	NO	NO	NO	NO	NO	T1	NO
3. N ₂ O from product uses	NO	NO	T1	NO	NO	NO	NO
4. Other							
H. Other	NO	NO	NO	NO	NO	NO	NO

* Key Category

T1,2,3: Tier 1, Tier 2, Tier 3 as described in the 2006 IPCC Guidelines

NE: emissions not estimated; NO: activity not occurring; NA: not applicable (emissions of the gas do not occur in the source category);

IE: emissions included elsewhere

Table 4.2 Emissions from Industrial Processes and Product Use 1990-2013

		Gas	Unit	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
2.A.1	Cement Production	CO ₂	kt	884.0	879.0	1700.9	1851.2	1859.8	2127.0	2295.1	2357.1	2347.9	2374.1	2106.7	1326.8	1105.1	966.3	1177.0	1111.7
2.A.2	Lime Production	CO ₂	kt	214.1	187.5	190.4	189.4	190.3	206.3	201.5	183.5	180.3	199.1	189.3	157.2	193.4	200.5	215.9	189.6
2.A.3	Glass Production	CO ₂	kt	13.3	12.0	10.7	10.1	5.1	0.6	0.6	0.5	0.5	0.5	0.3	0.0	NO	NO	NO	NO
2.A.4.a	Other- Ceramics	CO ₂	kt	5.23	5.64	6.66	6.30	6.05	6.26	6.36	7.53	7.66	7.04	4.18	0.53	0.42	0.83	0.03	0.03
2.A.4.b	Other- Soda Ash Use	CO ₂	kt	0.10	0.07	0.07	0.09	0.05	0.06	0.08	0.08	0.06	0.06	0.04	0.05	0.07	0.07	0.09	0.06
2.A.4.d	Other- Limestone use	CO ₂	kt	NO	NO	NO	4.33	2.04	2.24	3.42	4.17	2.38	2.11	2.52	1.54	1.03	1.04	0.44	0.21
2.B.1	Ammonia Production	CO ₂	kt	990.2	973.4	882.3	1041.2	810.9	0.3	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2.B.2	Nitric Acid Production	N ₂ O	kt CO ₂ eq	995.3	781.0	781.0	561.7	280.9	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2.D.1	Lubricant Use	CO ₂	kt	NO	11.78	70.08	19.85	11.78	14.88	17.37	59.54	19.23	23.57	20.47	22.39	16.82	18.73	18.28	19.03
2.D.2	Paraffin Wax Use	CO ₂	kt	5.09	7.53	13.05	15.33	18.36	17.49	19.13	24.19	19.05	21.10	14.40	15.52	13.13	14.42	14.53	16.88
2.D.3	Solvent use*	Indirect CO ₂	kt	79.33	84.03	76.32	82.55	80.64	79.72	78.94	79.07	84.29	91.44	79.68	74.03	67.15	65.59	63.59	66.15
2.E.1	Integrated Semiconductor	Circuit or HFCs, PFCs, SF ₆ , NF ₃	kt CO ₂ eq	1.17	145.33	491.70	424.71	344.12	393.08	285.75	310.12	249.41	238.87	179.86	107.30	68.19	41.13	31.55	34.63
2.F.1	Refrigeration and Conditioning	Air HFCs	kt CO ₂ eq	NO	11.68	149.99	199.45	287.11	431.83	581.96	748.30	779.10	802.52	937.79	925.39	951.11	979.22	975.48	1124.16
2.F.2	Fire Protection	HFCs	kt CO ₂ eq	NO	NO	7.33	8.75	10.16	11.55	12.93	34.48	34.49	34.50	34.50	34.51	34.51	34.52	34.53	34.53
2.F.3	Aerosols	HFCs	kt CO ₂ eq	NO	27.13	131.15	141.52	134.42	142.65	136.18	153.69	158.10	142.74	151.42	144.48	137.82	131.12	122.66	114.66
2.G.1	Electrical Equipment	SF ₆	kt CO ₂ eq	20.52	25.08	7.43	30.57	21.74	36.68	20.56	22.44	26.81	28.45	10.40	13.34	12.31	20.70	16.22	18.60
2.G.2	SF ₆ and PFCs from Other Product Uses	SF ₆	kt CO ₂ eq	12.90	12.99	14.69	14.56	15.48	16.12	13.64	11.81	7.14	5.68	5.15	5.87	3.35	2.61	2.76	2.92
2.G.3.a	Medical Applications	N ₂ O	kt CO ₂ eq	31.34	32.20	33.88	34.39	35.02	35.58	36.16	36.96	37.84	39.12	40.10	40.53	40.72	40.90	40.99	41.06
Total	IPPU		kt CO₂ eq	3252.6	3196.4	4567.7	4636.0	4114.0	3522.2	3709.7	4033.4	3954.2	4010.8	3776.9	2869.5	2645.1	2517.7	2714.0	2774.3

*Indirect CO₂ emissions from Solvent Use, 2.D.3, are not included in the total IPPU sector emissions but are included in Ireland's national total emissions. See CRF table 6.




Figure 4.1 Total Emissions from IPPU by Category, 1990-2013




Figure 4.2 Total Emissions from IPPU by Gas, 1990-2013

4.2 Emissions from Mineral Industry (2.A)

The emission categories relevant under 2.A *Mineral Products* are: 2.A.1 *Cement production*, 2.A.2 *Lime production*, 2.A.3 *Glass production*, 2.A.4 *Other process uses of carbonates*.

Cement production continues to be a key category in the national inventory. The production of glass ceased in Ireland in 2009.

4.2.1 Cement Production (2.A.1)

4.2.1.1 Category Description

Cement Production is a **key category** for Ireland, both in terms of the level assessment (2013) and the trend assessment (1990-2013).

During the cement manufacturing process, CO₂ is produced during the production of clinker. Clinker is produced when limestone, mainly calcium carbonate (CaCO₃) and small amounts of magnesium carbonate (MgCO₃), undergo calcination at high temperature to produce lime (CaO and MgO) and CO₂. The activated lime that results from this process combines with silica and alumina in the kiln feed to form cement clinker. The emissions of CO₂ are usually calculated from the amount of clinker produced and the stoichiometric ratio of CO₂ to CaO and MgO. A small amount of raw material may be converted into cement kiln dust (CKD) due to incomplete calcination. If the CKD is not recycled as part of subsequent kiln input, the CO₂ emissions based on clinker production must be corrected to account for the carbonate fraction lost in CKD.

Up until the year 2000, one company operated two cement plants in Ireland. A second company opened a new cement plant in 2000 and a third cement producer entered the market in 2003, bringing the total number of plants to four.

Process emissions of CO₂ from cement production have declined since 2007. This is a reflection of the recent economic downturn. However, emissions increased in 2012, which could be a sign of economic recovery.

4.2.1.2 Methodological Issues

A **Tier 3 approach** is used to estimate emissions from this category as described in the 2006 IPCC Guidelines. This methodology is based on collecting disaggregated data on the types and quantities of carbonates used to produce clinker at each cement plant as well as the respective emission factors of the carbonates consumed. Emissions are estimated using equation 2.3 from 2006 IPCC guidelines.

This method has been used for all years from 2005 to 2013 as the EU ETS became operational, plant specific CO₂ emissions and corresponding clinker production data are also available for all cement plants for the years 2004 through 2013 and these data are used directly to report emissions for category 2.A.1 in Ireland. The annual results incorporate verification of fuel use, limestone and carbonate use, combustion and process CO₂ estimates in accordance with Decision 2004/156/EC.

Information on calcium oxide (CaO) and magnesium oxide (MgO) content of clinker, for each of the four cement plants, has been provided to the inventory agency by the plant operators for all years from 2008 to 2013 as recommended in the previous annual inventory review reports. This information is not published in this inventory report as the cement producers deem it to be confidential. The data are available to the expert review teams for annual GHG inventory reviews upon request.

Prior to the implementation of the EU ETS, in 2004, plant-specific information relating to CO₂ emissions in 2002 and 2003 was obtained by the EPA for all cement plants for the development of Ireland's First National Allocation Plan (NAP1) under Directive 2003/87/EC (EP and CEU, 2003) on emissions trading in the EU. The reported process CO₂ emissions for each plant in 2002 and 2003 were calculated according to the guidelines for the monitoring and reporting of greenhouse gas emissions in Decision 2004/156/EC that supports Directive 2003/87/EC. The method used was fully

consistent with the Tier 3 method described above and its application employs reliable data on clinker production, corrected as appropriate for CKD, and CaO/MgO content of the clinker.

For the two original cement plants that were operated by the single cement producer, the company concerned supplied estimates of process emissions for the years 1990-2001 that it had calculated internally in line with the specific information provided for the years 2002 and 2003 and used for NAP1. The associated values of annual clinker production were not provided. For the purposes of complete and consistent reporting, the inventory agency estimated annual clinker production for the years 1990-2001 based on the plant specific process emission factors available for the two plants for the years from 2002 onwards. This is appropriate, as the company has always used the same local on-site supply of limestone, and the time-series of process CO₂ emissions for cement production overall may therefore be considered consistent for the period 1990-2013.

The revised estimates for category 2.A.1 were included in the 2006 submission and no further recalculations have been made since the EU ETS data were adopted as the best available for inventory purposes.

Additional information on clinker production, emissions and IEFs is provided in Table 3.2.A of Annex 3.2.

4.2.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 1.5 per cent in line with Table 2.3 of the 2006 IPCC guidelines. Production of clinker data are available, so the uncertainty associated with these data is 1-3%, based on plant level weighing of raw materials.

The uncertainty of the emission factor is 1.5 per cent in line with Table 2.3 of the 2006 IPCC guidelines. Overall chemical analysis/composition pertaining to carbonate content/mass/type is known (Tier 3), with an uncertainty range of 1-3%.

4.2.1.4 Category-specific QA/QC and verification

As the emissions are estimated from individual plant data, which are subject to verification under Directive 2003/87/EC, their validity is fully established in the context of the companies' documented methods and data and the associated guidance on emissions estimation methods provided by Decision 2004/156/EC (CEC, 2004). Such verification allows for accurate accounting of combustion emissions and process emissions separately.

Data from each plant for the most recent year in the inventory are checked for consistency with historic data from that plant. Implied emission factors are also calculated and checked for variability or step changes across the time series.

Comparisons are also made across the different plants in the same source sector, to check for consistency. Typically implied emission factors are compared. These checking procedures help to identify any erroneous point source data, and are readily undertaken due to the limited number of plants in Ireland.

Data reported under ETS for plants in this category are also cross checked with data supplied by the same operators for other reporting requirements, such as, Integrated Pollution Prevention and Control directive (IPPC) and under the European Pollutant Release and Transfer Register (E-PRTR) for consistency.

4.2.1.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.2.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.2.2 Lime Production (2.A.2)

4.2.2.1 Category Description

Calcium oxide (CaO or quicklime) is formed by heating limestone to decompose the carbonates. This is usually done in shaft or rotary kilns at high temperatures and the process releases CO₂. Dolomite and dolomitic (high magnesium) limestone may also be processed at high temperature to obtain dolomitic lime with a loss of CO₂. Quicklime can then be further treated by the addition of water, a process called slaking, to produce slaked lime (Ca(OH)₂ and Ca(OH)₂.Mg(OH)₂), which generates large amounts of heat and steam. The finished product can then be packaged and distributed for use.

Currently, there are two companies operating 3 lime plants in Ireland and a fourth that operated until 1999. It is understood that all three utilised limestone quarries and kilns to burn the limestone raw material. The nature of the fuel used and the abatement in place varies from plant to plant.

4.2.2.2 Methodological Issues

Since 2005, emissions from lime production are based on a Tier 3 input-based carbonate approach and equation 2.7 from the 2006 IPCC guidelines. The CO₂ estimates for lime production in 2013 have been obtained from the ETS returns to the EPA.

Historically, statistical data on lime production in Ireland were obtained annually from the lime manufacturers (three companies up to 1999 and two companies thereafter) and form the basis for emissions over the period 1990-2004. As in the case of cement production, lime producers now provide their own estimates of CO₂ emissions from lime manufacture under Directive 2003/87/EC on ETS. These estimates were calculated in accordance with the methods described in the supporting Decision 2004/156/EC, equivalent to Tier 3 approach, thus providing detailed information on emission estimates and activity data for another important source of CO₂ emissions in Industrial Processes and Product Use.

The implied emission factor for aggregated lime production was 0.75 t CO₂/t lime in 2013, which is similar to that for the other years for which ETS data are available. Additional detailed information on lime production, emissions and IEFs is available in Table 3.2.B in Annex 3.2.

4.2.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 5 per cent and the uncertainty of the emission factor is 5 per cent.

4.2.2.4 Category-specific QA/QC and verification

As the emissions are estimated from individual plant data, which are subject to verification under Directive 2003/87/EC, their validity is fully established in the context of the companies' documented methods and data and the associated guidance on emissions estimation methods provided by Decision 2004/156/EC (CEC, 2004). Such verification allows for accurate accounting of combustion emissions and process emissions separately.

Data from each plant for the most recent year in the inventory are checked for consistency with historic data from that plant. Implied emission factors are also calculated and checked for variability or step changes across the time series.

Comparisons are also made across the different plant in the same source sector, to check for consistency. Typically implied emission factors are compared. These checking procedures help to identify any erroneous point source data, and are readily undertaken due to the limited number of plant in Ireland.

Data reported under ETS for plants in this category are also cross checked with data supplied by the same operators for other reporting requirements, such as, Integrated Pollution Prevention and Control directive (IPPC) and under the European Pollutant Release and Transfer Register (E-PRTR) for consistency.

4.2.2.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.2.2.6 Category-specific Planned Improvements

It is planned to revise the uncertainty associated with lime production to bring it in line with the information provided in Table 2.5 of the 2006 IPCC guidelines.

4.2.3 Glass Production (2.A.3)

4.2.3.1 Category Description

There are many kinds of glass articles and compositions in use commercially, but the glass industry can be divided into four main categories: containers, flat (window) glass, fibre glass, and specialty glass. The great bulk of commercial glass is in the first two categories, and is almost entirely soda-lime glass, consisting of silica (SiO_2), soda (Na_2O), and lime (CaO), with small amounts of alumina (Al_2O_3), and other alkalies and alkaline earths, plus some minor ingredients. Insulation grade fibre glass - a major component of the third category - is similar in composition. This source category also includes emissions from the production of glass wool, a category of mineral wool, where the production process is similar to glass making.

The production of glass ceased in Ireland in 2009. The manufacture of glass was not a predominant industry in Ireland, being limited to three sectors: crystal glass, bottle glass, and glass-based insulation.

The only bottle glass plant closed in 2002, a crystal glass plant closed in early 2006, the glass-based insulation plant closed in 2008 and the last one, a second crystal glass plant closed down in 2009.

4.2.3.2 Methodological Issues

A Tier 2 approach is used based on different glass manufacturing processes undertaken in the Ireland. Similar to other categories under 2.A, information from 2 individual crystal glass plants that were participants in the Emissions Trading Scheme were also used to compile the emissions estimates for this category for the years 2005 to 2009.

The production of **bottle glass** was the major source of emissions in this category. The CO₂ emissions are estimated from the annual production quantities obtained from the company for the development of annual inventories for heavy metals. Equation 2.11 of the 2006 IPCC guidelines and the emission factor of 0.21 kg CO₂/kg glass (Table 2.6 of 2006 IPCC guidelines) are used. Allowance is made for recycled glass, which is assumed to be 5 per cent in 1990, increasing to 30 per cent in 2002 when the plant closed.

In the case of **crystal glass**, the CO₂ emissions are based on the use of potassium carbonate and sodium carbonate use (soda ash) as reported under ETS, using the emission factors of 0.415 t CO₂/t Na₂CO₃ and 0.267 t CO₂/t K₂CO₃, provided by the ETS monitoring and reporting guidelines. The company concerned has supplied estimates for all years up to and including 2009, when the plant closed.

Emissions from the production of **glass-based insulation** materials are also based largely on soda ash use although small amounts of dolomite and limestone were also used up to 2005.

The emissions of CO₂ from glass production amounted to 13.3 kt in 1990 and reduced to 0.02 kt in 2009, due to closure of the last glass manufacturing plant. Additional detailed information on glass production, emissions and IEFs is available in Table 3.2.C in Annex 3.2.

4.2.3.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 5 per cent and the uncertainty of the emission factor is 2.5 per cent.

4.2.3.4 Category-specific QA/QC and verification

As the emissions are estimated from individual plant data, which are subject to verification under Directive 2003/87/EC, their validity is fully established in the context of the companies' documented methods and data and the associated guidance on emissions estimation methods provided by Decision 2004/156/EC (CEC, 2004). Such verification allows for accurate accounting of combustion emissions and process emissions separately.

4.2.3.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.2.3.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.2.4 Other Process Uses of Carbonates (2.A.4)

4.2.4.1 Category Description

Limestone (CaCO_3), dolomite ($\text{CaMg}(\text{CO}_3)_2$) and other carbonates (e.g., MgCO_3 and FeCO_3) are basic raw materials having commercial applications in a number of industries. In addition to those industries already discussed individually (cement production, lime production and glass production), carbonates also are consumed in metallurgy (e.g., iron and steel), agriculture, construction and environmental pollution control (e.g., flue gas desulphurisation.) Soda ash (sodium carbonate, Na_2CO_3) is a white crystalline solid that is used as a raw material in a large number of industries including glass manufacture, soap and detergents, pulp and paper production as well as food additive, drinking water treatment (softener) and treatment for a low pH reading in wastewater stream. The CO_2 emissions reported under this category refer to those emissions associated with:

- Limestone (CaCO_3) used for flue gas desulphurisation,
- Limestone used for purification in sugar manufacture,
- Limestone used in the manufacture of bricks, flues and tiles,
- Clays and shale used as a raw material in the manufacture of bricks, flues and ceramics,
- Soda ash use (non-glass manufacture).

Since 2008, when the last ceramics and tile manufacturing plants closed, the only two sources of emissions in this category are from a brick manufacturing plant and from the use of limestone for flue gas desulphurisation at peat fired power plants. The emission trend in recent years is almost entirely due to the amount of desulphurisation required at these power plants.

4.2.4.2 Methodological Issues

Emissions of CO_2 have been estimated using a Tier 3, carbonate input approach, for sources in this category. Limestone has been used as environmental pollution control to reduce the sulphur emitted from peat burning in one electricity generating station since 2001 and in a second such plant since 2007. The CO_2 emissions estimates are taken from ETS reports to the EPA. They are estimated on the basis of limestone quantity used by the companies and reported process emissions, giving an implied emission factor in the range from 0.43 to 0.44 t CO_2 /t limestone between 2001 and 2013. The stoichiometric ratio of CO_2 to CaCO_3 is 0.44.

A further minor use of limestone in Ireland is its application in the purification of sugar produced from sugar beet. However, sugar production ceased in 2006 and the only information on emissions is that obtained under ETS in respect of 2005 and 2006. Additionally limestone was used for tile manufacturing by one company in the three years of its operation (2006-2008) and for brick manufacturing by another company until its closure (1990-2008). Data was reported by both companies for relevant years of trading under the ETS and for the preceding years it was sourced by the inventory agency from the companies.

The emissions of CO_2 from the use of clays and shale as a raw material in the manufacture of bricks and ceramics are estimated using information from individual plants that are participants in the ETS.

The emissions associated with soda ash use by one company in Ireland are reported by the company under ETS for the years 2005-2013 and have been used directly in the inventory. The other uses of soda ash are already reported under 2.A.3 glass production. Activity data for years prior to the ETS

data were sourced by the inventory agency from the company. Estimates of CO₂ for all years from 1990-2004 were calculated using an emission factor of 0.41 t CO₂/t soda ash, indicated by the average 2005-2008 ETS data. This approach has allowed a full 1990-2013 time series of emissions to be included in the inventory. Additional detailed information on activity data, emissions and EFs is available in Table 3.2.E in Annex 3.2.

In the case of bricks and ceramics, the ETS data for one company provide estimates of emissions for the years 2005-2013, a further one company for the years 2005-2011 and a further two companies for the years 2005-2008 which all three have now ceased trading, along with the corresponding quantities of carbonate input materials and the relevant emission factors. The emission factors for clay bricks and flue liners are in the range 0.027 to 0.053 tonne CO₂/tonne carbonate input while the emission factor for ceramic tiles averages 0.062 tonne CO₂/tonne carbonate input. The emissions for the years prior to ETS are calculated from the companies' estimates of material use and their respective average ETS emission factors. Additional detailed information on raw material use, emissions and IEFs is available in Table 3.2.D in Annex 3.2.

4.2.4.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 5 per cent and the uncertainty of the emission factor is 2.5 per cent.

4.2.4.4 Category-specific QA/QC and verification

As the emissions are estimated from individual plant data, which are subject to verification under Directive 2003/87/EC, their validity is fully established in the context of the companies' documented methods and data and the associated guidance on emissions estimation methods provided by Decision 2004/156/EC (CEC, 2004). Such verification allows for accurate accounting of combustion emissions and process emissions separately.

4.2.4.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.2.4.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.3 Emissions from Chemical Industry (2.B)

The emission categories relevant under *2.B Chemical Industry* are: *2.B.1 Ammonia Production* and *2.B.2 Nitric Acid Production*. All other Chemical Industry activities have not occurred in Ireland since 1990 and are reported as Not Occurring (NO).

Ammonia and nitric acid production in Ireland was undertaken by two plants, both of which were operated by Irish Fertiliser Industries for the production of nitrogenous fertilisers. However, during 1999 and 2000 the major fertiliser manufacturers introduced severe rationalisation and restructuring measures, which resulted in the closure of the nitric acid and ammonia plants in 2002 and 2003, respectively.

Fertiliser manufacture in Ireland no longer takes place and all fertilisers are either imported as a finished product or only undergo further blending in Ireland.

4.3.1 Ammonia Production (2.B.1)

4.3.1.1 Category Description



Ammonia (NH₃) is a major industrial chemical and the most important nitrogenous material produced. Ammonia production requires a source of nitrogen (N) and hydrogen (H). Nitrogen is obtained from air through liquid air distillation or an oxidative process where air is burnt and the residual nitrogen is recovered. Ammonia is the basis of all nitrogen fertilisers and is normally manufactured by synthesis of nitrogen and hydrogen, with natural gas (methane CH₄) as the basic raw material. Utilising the Haber Bosch process, natural gas, air and water were reacted to produce ammonia in liquid form and CO₂ as a by-product.

Urea was one of the main end products of the plant, which was formed when the ammonia produced and the CO₂ by-product reacted together to form prills (small particles) of urea. The other main product, anhydrous ammonia was stored and transported to Irish Fertiliser Industries other plant where it underwent further processing (discussed in section 4.3.2 Nitric Acid Production below).

4.3.1.2 Methodological Issues

The ammonia industry uses predominantly natural gas, the following outline of the process and sources of greenhouse emissions (CO₂) is based on production using natural gas. Anhydrous ammonia produced by catalytic steam reforming of natural gas (mostly CH₄) involves the following reactions with carbon dioxide produced as a by-product.


Primary steam reforming:




Secondary air reforming:




Overall reaction:



Ammonia synthesis:



Secondary reformer process gas shift conversion:



Emissions of CO₂ from ammonia production are estimated using a Tier 2/3 approach based on country specific data on fuel type and carbon content of the fuel supplied to the plant. Data on the natural gas feedstocks to the plant are indicated in the national energy balance provided by SEAI. No feedstock carbon is sequestered in urea and the emission factor is 54.94 kg CO₂/TJ, the value for indigenous natural gas, which equates to 2.3 tonne CO₂/tonne natural gas. The CO₂ emissions from ammonia production were 990.23 kt in 1990 and 0.30 kt in 2003, the last year of operation.

4.3.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 1 per cent and the uncertainty of the emission factor is 5 per cent.

4.3.1.4 Category-specific QA/QC and verification

4.3.1.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.3.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.3.2 Nitric Acid Production (2.B.2)

4.3.2.1 Category Description

Nitric acid is used as a raw material mainly in the manufacture of nitrogenous-based fertiliser. Nitric acid may also be used in the production of adipic acid and explosives (e.g., dynamite), for metal etching and in the processing of ferrous metals. During the production of nitric acid (HNO_3), nitrous oxide (N_2O) is generated as an unintended by-product of the high temperature catalytic oxidation of ammonia (NH_3).

Nitric acid production in Ireland ceased in 2002 due to the closure of Irish Fertiliser Industries. Ammonia, transported from Irish Fertiliser Industries ammonia production plant (section 4.3.1) to the ammonium nitrate production plant, was oxidised over a catalyst to form nitric acid. The nitric acid was then combined with more ammonia to produce ammonium nitrate which, when solidified into granules or made into bead-like prills, is applied to land using a fertiliser spreader. Other fertiliser blends were also manufactured at the plant.

4.3.2.2 Methodological Issues

For the years 1990-1995, the inventory agency received direct correspondence from the plant operator specifying the quantities of nitric acid produced and the company's estimates of N_2O emitted during the production process.

Four units at this plant produced 338,000 tonnes of nitric acid in 1990 with associated N_2O emissions of 3,340 tonnes. The emissions were estimated from nitrogen loading and the type of catalyst used in the process.

4.3.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 1 per cent and the uncertainty of the emission factor is 10 per cent.

4.3.2.4 Category-specific QA/QC and verification

There is no country specific QA/QC for this category as the plant is closed since 2002, before the establishment of Ireland's National Atmospheric Inventory System.

4.3.2.5 Category-specific Recalculations

A change of GWP for N_2O resulted in 3.9 per cent decrease in emissions for the existing time series 1990-2002, or aggregated 388.49 kt CO_2 equivalent in total.

4.3.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.3.3 Adipic Acid Production (2.B.3)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.4 Caprolactam, Glyoxal and Glyoxylic Acid Production (2.B.4)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.5 Carbide Production (2.B.5)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.6 Titanium Dioxide Production (2.B.6)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.7 Soda Ash Production (2.B.7)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.8 Petrochemical and Carbon Black Production (2.B.8)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.9 Fluorochemical Production (2.B.9)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.3.10 Other Chemical Industry (2.B.10)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

4.4 Emissions from Metal Industry (2.C)

This section covers emissions of greenhouse gases that result from the production of metals in the following categories;

- Iron and steel production 2.C.1
- Ferroalloys production 2.C.2
- Aluminium production 2.C.3
- Magnesium production 2.C.4
- Lead production 2.C.5
- Zinc production 2.C.6

4.4.1 Iron and Steel Production (2.C.1)

Ireland had one Electric Arc Furnace (EAF) in operation in the years 1990 to 2001 producing steel from scrap and recycled metal. It is planned to report process CO₂ emissions from this category in the next submission based on the emission factor provided in table 4.5 of the 2006 IPCC guidelines, 0.08 t CO₂/t steel.

4.4.2 Ferroalloys Production (2.C.2)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.4.3 Aluminium Production (2.C.3)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.4.4 Magnesium Production (2.C.4)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.4.5 Lead Production (2.C.5)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.4.6 Zinc Production (2.C.6)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.4.7 Other Metal Industry (2.C.7)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

4.5 Emissions from Non-energy Products from Fuels and Solvent Use (2.D)

4.5.1 Lubricant Use (2.D.1)

4.5.1.1 Category Description

Lubricants are mostly used in industrial and transportation applications. Lubricants are produced either at refineries through separation from crude oil or at petrochemical facilities. They can be subdivided into (a) motor oils and industrial oils, and (b) greases, which differ in terms of physical characteristics (e.g., viscosity), commercial applications, and environmental fate. The use of lubricants in engines is primarily for their lubricating properties and associated emissions are therefore considered as non-combustion emissions and are reported here in the IPPU Sector. Most waste lubricant oil is collected in Ireland and disposed of in an environmental way or energy recovery takes place and the waste lubricant is used for energy purposes. These waste oil handling emissions, however, are to be reported in the Waste Sector (or in the Energy Sector when energy recovery takes place). A small proportion of lubricant oils oxidise during use, and CO₂ emissions from this category are reported in 2.D.1 Lubricant use.

4.5.1.2 Methodological Issues

Ireland uses a Tier 1 method to estimate emissions of CO₂ from non-energy use of lubricants based on equation 5.2 in the 2006 IPCC guidelines and an ODU (Oxidising During Use) default factor 0.2 from table 5.2 shown below. The national energy balance provides data on lubricant consumption for the full time series 1992-2013 (consumption was not occurring in 1990 and 1991). The carbon content of lubricants value is 20.0 tonne carbon/TJ. Emissions of CO₂ estimated for this category are presented in Table 4.2.

Equation 5.2 Lubricants – Tier 1 Method

$$\text{CO}_2 \text{ Emissions} = \text{LC} \bullet \text{CC}_{\text{Lubricant}} \bullet \text{ODU}_{\text{Lubricant}} \bullet 44 / 12$$

Where:

CO₂ Emissions = CO₂ emissions from lubricants, tonne CO₂

LC = total lubricant consumption, TJ

CC_{Lubricant} = carbon content of lubricants (default), tonne C/TJ

ODU_{Lubricant} = ODU factor (based on default composition of oil and grease), fraction

44/12 = mass ratio of CO₂/C

4.5.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 30 per cent and the uncertainty of the emission factor is 5 per cent.

4.5.1.4 Category-specific QA/QC and verification

See Chapter 9.

4.5.1.5 Category-specific Recalculations

Emissions from Lubricant Use were included for the first time in this submission.

4.5.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.5.2 Paraffin Wax Use (2.D.2)

4.5.2.1 Category Description

The category, as defined here, includes such products as petroleum jelly, paraffin waxes and other waxes, including ozokerite (mixtures of saturated hydrocarbons, solid at ambient temperature). Paraffin waxes are separated from crude oil during the production of light (distillate) lubricating oils. Paraffin waxes are categorised by oil content and the amount of refinement. Paraffin waxes are used in applications such as: candles, corrugated boxes, paper coating, board sizing, food production, wax polishes, surfactants (as used in detergents) and many others. Emissions from the use of waxes derive primarily when the waxes or derivatives of paraffins are combusted during use (e.g., candles), and when they are incinerated with or without heat recovery or in wastewater treatment (for surfactants).

4.5.2.2 Methodological Issues

Ireland uses a Tier 2 method to estimate emissions of CO₂ from non-energy use of paraffin wax based on equation 5.5 in the 2006 IPCC guidelines and an ODU (Oxidising During Use) factor 0.9 for paraffin wax candles and an ODU factor of 0.2 for all other uses of paraffin wax. The national energy balance provides data on paraffin wax consumption for the full time series 1990-2013. The carbon content of paraffin wax value is 20.0 tonne carbon/TJ. Emissions of CO₂ estimated for this category are presented in Table 4.2. CO₂ emissions estimated for this category are presented in Table 4.2.

Equation 5.5 Waxes – Tier 2 Method

$$\text{CO}_2 \text{ Emissions} = \sum_i (\text{PW}_i \bullet \text{CC}_i \bullet \text{ODU}_i) \bullet 44 / 12$$

Where:

CO₂ Emissions = CO₂ emissions from waxes, tonne CO₂

PW_i = consumption of was type i (candle wax and residual wax), TJ

CC_i = carbon content of wax type i, tonne C/TJ

ODU_i = ODU factor for wax type i, fraction

44/12 = mass ratio of CO_2/C

4.5.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 30 per cent, the uncertainty of the emission factor is 5 per cent.

4.5.2.4 Category-specific QA/QC and verification

See Chapter 9.

4.5.2.5 Category-specific Recalculations

Emissions from Paraffin wax Use were included for the first time in this submission.

4.5.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.5.3 Other Non-energy Products from Fuels and Solvent Use (2.D.3)

4.5.3.1 Category Description

The use of solvents manufactured using fossil fuels as feedstocks can lead to evaporative emissions of various non-methane volatile organic compounds (NMVOC), which are subsequently further oxidised in the atmosphere. White spirit is used as an extraction solvent, as a cleaning solvent, as a degreasing solvent and as a solvent in aerosols, paints, wood preservatives, lacquers, varnishes and asphalt products. In Western Europe about 60 per cent of the total white spirit consumption is used in paints, lacquers and varnishes. White spirit is the most widely used solvent in the paint industry.

Emissions of Non-methane volatile organic compounds are reported in this category. NMVOCs are indirect greenhouse gases which result from the use of solvents and various other volatile compounds. The indirect CO_2 emissions associated with these NMVOC emissions are reported in CRF table 6. The indirect CO_2 emissions are included in Ireland's national total, without LULUCF with indirect. Ireland's national total is fully consistent with the national total reported in the first commitment period in this regard.

4.5.3.2 Methodological Issues

Methodologies for estimating these NMVOC emissions can be found in the EMEP/CORINAIR Emission Inventory Guidebook (EEA, 2013). Further information on emissions of NMVOCs and indirect CO_2 emissions can be found in Chapter 9 of this report. Estimates of indirect CO_2 emissions are derived from NMVOCs by assuming that 85 per cent of the mass of NMVOCs is converted to CO_2 .

4.5.3.3 Uncertainties and Time-series Consistency

See Chapter 9.

4.5.3.4 Category-specific QA/QC and verification

See chapter 9.

4.5.3.5 Category-specific Recalculations

Major recalculations occurred in this submission due to the completion of a study designed to update the NMVOC emission estimates for 2006 to 2013. This project focused on updating activity data, emission factors and in some cases methodologies as outlined in Barry, S. and O'Regan, B. (2014). All recalculations are presented in detail in chapter 5 of Ireland's IIR 2015.

4.5.3.6 Category-specific Planned Improvements

A description of the planned improvements for NMVOC emissions from this sector can be found in Ireland's Informative Inventory Report (EPA, 2015).

4.6 Emissions from Electronics Industry (2.E)

4.6.1 Integrated Circuit or Semiconductor (2.E.1)

4.6.1.1 Category Description

The semiconductor industry uses HFCs, PFCs, SF₆ and NF₃ in manufacturing processes. Both HFCs and PFCs are used in the cleaning of chambers used for chemical vapour deposition processes, dry plasma etching, vapour phase soldering and vapour phase blanketing, leak testing of hermetically sealed components and as coolants. Cleaning and etching during semiconductor manufacture account for the majority of emissions from the category. In addition, SF₆ and NF₃ are used in the etching process.

PFC emissions peaked in 2000 after which they have gradually decreased. NF₃ emission levels were highest in the period 2000-2007 and have been negligible from 2008 onwards. This is due to the economic downturn as well as the voluntary agreement implemented by the European Semiconductor Industry Association ([ESIA, 2011](#)) for the reduction of PFC emissions. Emission estimates for Electronics Industry category 2.E.1 are presented in Table 4.3 and Figure 4.3 below.

4.6.1.2 Methodological Issues

Ireland uses a Tier 2a method to estimate emissions from this category using company specific data based on gas consumption and emission control technologies in use in the process, as outlined in the 2006 IPCC guidelines. There are a small number of large semiconductor manufacturers in Ireland. These installations provide data on the annual use and estimated emissions of HFCs, PFCs, SF₆ and NF₃ in their plants over the full time series 1990-2013.

4.6.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 20 per cent and the uncertainty of the emission factor is 10 per cent.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams et al., 2005) and reviewed in 2013 (Goodwin et al., 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.6.1.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory

approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.6.1.5 Category-specific Recalculations

NF₃ gas was estimated for the first time in this 2015 submission. NF₃ emission estimates in the time series are presented in Table 4.3 and Figure 4.3. Other changes were mostly related to revised category-specific recalculations and revised GWPs and have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.6.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.6.2 TFT Flat Panel Display Industry (2.E.2)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.6.3 Photovoltaics Industry (2.E.3)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.6.4 Heat Transfer Fluid Use (2.E.4)

This activity has not existed in Ireland since 1990. This category is reported as Not Occurring (NO).

4.6.5 Other Electronics Industry (2.E.5)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

4.7 Emissions from Product Uses as Substitutes for ODS (2.F)

The compilation of emission estimates for fluorinated gases presents major challenges for inventory agencies because they emanate from diverse sources that are entirely different to those traditionally covered by atmospheric emissions inventories. In addition, the use of many of the substances concerned is continuing to change very rapidly in the marketplace. This sector covers the following categories;

- Refrigeration and air conditioning 2.F.1,
- Foam blowing agents 2.F.2,
- Fire protection 2.F.3,
- Aerosols 2.F.4,
- Solvents 2.F.5,
- Other applications 2.F.6.

Emission estimates for category 2.F are presented in Table 4.3 and Figure 4.3 below.

Table 4.3. Emissions of HFC, PFC, SF₆ and NF₃ from IPPU 1990-2013 (kt CO₂ eq)

IPCC Source Category	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
2.F.1 Refrigeration and Air-Conditioning	NO	0.71	71.90	106.13	178.63	307.24	436.99	576.34	584.26	579.79	691.06	667.26	685.69	712.07	717.08	864.95
2.F.1 Mobile Air Conditioning	NO	10.97	78.09	93.32	108.49	124.59	144.97	171.96	194.84	222.72	246.73	258.14	265.42	267.16	258.40	259.20
2.F.2 Foams	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2.F.3 Fire-extinguishers	NO	NO	7.33	8.75	10.16	11.55	12.93	34.48	34.49	34.50	34.50	34.51	34.51	34.52	34.53	34.53
2.F.4 Aerosols	NO	27.13	94.86	102.41	92.25	99.41	91.77	108.59	107.75	95.73	101.65	95.01	87.69	80.38	72.40	65.16
2.F.4 Metered Dose Inhalers	NO	NO	36.29	39.11	42.17	43.24	44.41	45.10	50.36	47.00	49.78	49.47	50.13	50.75	50.27	49.50
2.E.1 Semiconductor manufacture	0.59	2.31	15.13	3.92	2.38	3.34	1.72	2.81	4.00	4.29	4.59	3.70	4.19	3.11	2.80	3.39
TOTAL HFC	0.59	41.12	303.60	353.64	434.07	589.38	732.79	939.29	975.68	984.05	1128.30	1108.08	1127.64	1147.98	1135.47	1276.74
2.E.1 Semiconductor manufacture	0.12	97.61	397.76	379.51	267.89	285.95	234.81	216.39	190.96	168.10	136.14	83.63	46.58	15.88	9.56	8.32
TOTAL PFC	0.12	97.61	397.76	379.51	267.89	285.95	234.81	216.39	190.96	168.10	136.14	83.63	46.58	15.88	9.56	8.32
2.E.1 Semiconductor manufacture	0.46	41.04	29.64	19.49	27.27	57.16	31.14	62.54	26.25	28.81	39.14	19.97	17.41	22.15	18.41	22.01
2.G.1 Electrical equipment	20.52	25.08	7.43	30.57	21.74	36.68	20.56	22.44	26.81	28.45	10.40	13.34	12.31	20.70	16.22	18.60
2.G.2 Other - window soundproofing	0.52	0.61	0.41	0.28	0.28	0.27	0.42	0.58	0.73	0.88	1.03	1.18	1.33	1.48	1.63	1.78
2.G.2 Other - medical applications	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76
2.G.2 Other - sporting goods	NO	NO	1.89	1.89	2.81	3.46	6.64	10.47	5.66	4.04	3.37	2.05	1.26	0.36	0.37	0.37
2.G.2 Other - gas-air tracers	11.63	11.63	11.63	11.63	11.63	11.63	5.81	NO	NO	NO	NO	1.88	NO	NO	NO	NO
TOTAL SF₆	33.88	79.11	51.76	64.63	64.48	109.95	65.34	96.78	60.21	62.94	54.69	39.18	33.08	45.45	37.39	43.53
2.E.1 Semiconductor manufacture	NO	4.37	49.17	21.78	46.58	46.63	18.08	28.38	28.21	37.67	NO	NO	NO	NO	0.78	0.90
TOTAL NF₃	NO	4.37	49.17	21.78	46.58	46.63	18.08	28.38	28.21	37.67	NO	NO	NO	NO	0.78	0.90
TOTAL HFC, PFC, SF₆ and NF₃	34.59	222.22	802.29	819.56	813.03	1031.91	1051.02	1280.83	1255.05	1252.75	1319.13	1230.89	1207.30	1209.31	1183.20	1329.50




Figure 4.3 Emissions of HFC, PFC, SF₆ and NF₃

4.7.1 Refrigeration and air conditioning (2.F.1)

4.7.1.1 Category Description

Refrigeration and air conditioning is a **key category** for Ireland, both in terms of the level assessment (2013) and the trend assessment (1990-2013). It includes the following sub-categories;

- Commercial refrigeration 2.F.1.a,
- Domestic refrigeration 2.F.1.b,
- Industrial refrigeration 2.F.1.c,
- Transport refrigeration 2.F.1.d,
- Mobile air-conditioning 2.F.1.e,
- Stationary air-conditioning 2.F.1.f.

HFCs and HFC blends have been widely used as replacement refrigerants for CFC and HCFC refrigerants across virtually all refrigeration sub-sectors (i.e. domestic refrigeration, small commercial distribution systems, industrial systems, building air conditioning systems and refrigerated transport).

The first HFC refrigerant on the market was R134a in the 1990s. The composition of the HFC refrigerants present on the Irish market has undergone some significant changes across the time series. These changes are due to the rapid phase-in of different HFC refrigerants in various applications, and the introduction of new refrigerant blends i.e. R404A, 407A, 407C, R410A, R404A, R134a and R407C, which have been the main refrigerants since 2000.

In the early part of the time series (1995 through to 2000) large quantities of HCFCs were used as refrigerants (mainly R22, which are not subject to greenhouse gas emission reporting as controlled under the Montreal Protocol).

4.7.1.2 Methodological Issues

All emissions from stationary refrigeration and stationary air conditioning are reported under *2.F.1.a Commercial refrigeration* as it has not been possible to disaggregate these emissions.

Data on the quantities of industrial gases supplied to the refrigeration sector is obtained from chemical suppliers and manufacturers of refrigeration units. Sales data is provided for a range of HFCs and blends corresponding to the individual HFC species: HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a, HFC-152a.

There is no manufacture of fluorinated gases in Ireland. Imported HFCs are calculated using the data supplied as described above. Exports are calculated on the basis of refrigeration unit manufacturers' share of exports. In Ireland there is no known destruction of HFCs. Recovered gas is used either in other equipment or exported for recycling or destruction.

A bottom-up approach is not feasible for estimating emissions from stationary refrigeration and air conditioning in Ireland due to the lack of data available on equipment types and HFC sales data in equipment sub-categories. Therefore emissions are estimated using a top-down approach based on reported sales data and information on market shares, which are applied to calculate estimates of total HFC sales in the Irish stationary refrigeration and air-conditioning sectors. As a result, emissions arising from sub-categories *2.F.1.b Domestic refrigeration*, *2.F.1.c Industrial refrigeration*, *2.F.1.d Transport refrigeration* and *2.F.1.f Stationary air-conditioning* are reported under *2.F.1.a Commercial Refrigeration*.

Emissions of HFCs from sub-category *2.F.1.e Mobile Air-Conditioning* are estimated using a Tier 3b bottom-up analysis which uses national vehicle fleet statistics (Table 3.1.13, Annex 3.1.B) and assumed rates of air-conditioning unit penetration in the national vehicle fleet (AEA, 2011). The methodology used takes account of vehicle lifetime (12 years), the percentage of vehicles having HFC in their air-conditioning systems, average charge per unit, product manufacturing emissions (AEA, 2011), effective lifetime leakage rates (incorporating emissions from normal operating losses and accidental releases arising from collision damage) and decommissioning losses (EP and CEU, 2006).

4.7.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 20 per cent and the uncertainty of the emission factor is 10 per cent.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams et al., 2005) and reviewed in 2013 (Goodwin et al., 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.7.1.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.7.1.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.7.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.7.2 Foam Blowing Agents (2.F.2)

4.7.2.1 Category Description

Increasingly, HFCs are being used as replacements for CFCs and HCFCs in foams and particularly in insulation applications. This category includes the following sub-categories;

- Closed Cells 2.F.2.a,
- Open Cells 2.F.2.b.

Open-celled foams are used for applications such as household furniture cushioning, mattresses, automotive seating and for moulded products such as car steering wheels and office furniture. Closed-cell foams, on the other hand, are primarily used for insulating applications where the gaseous thermal conductivity of the chosen blowing agent (lower than air) is used to contribute to the insulating performance of the product throughout its lifetime.

4.7.2.2 Methodological Issues

Following a new 2013 study on F-gases, emissions from *2.F.2 Foam Blowing Agents* were shown to be not occurring in Ireland and consequently were removed from the whole time series, emissions were reported in for the time series 1990-2011 in submission 2013. During this study, it was identified that no manufacturing of open-cell foams have occurred in Ireland.

The production of closed-cell foams takes place in Ireland by one company that used HCFC-141b and now uses pentane. Therefore, no manufacturing emissions occur in Ireland.

4.7.3 Fire Protection (2.F.3)

4.7.3.1 Category Description

There are two general types of fire protection (fire suppression) equipment that use HFCs and/or PFCs: portable (streaming) equipment, and fixed (flooding) equipment. HFCs, PFCs and more recently a fluoroketone are mainly used as substitutes for halons, typically halon 1301, in flooding equipment.

HFCs are most commonly used in fixed flooding systems in the protection of electronic and telecommunications equipment, in data centres, military applications, records offices, bank vaults and oil production facilities. There are a number of companies operating these systems in Ireland.

Although HFC-23 can be used in some systems, the 2013 study (Goodwin et al., 2013) identified none within Ireland so the only HFC used is HFC-227ea. The majority of emissions occur when fire protection systems are triggered either accidentally or due to the occurrence of a fire. Smaller emissions occur during maintenance and filling.

4.7.3.2 Methodological Issues

Activity data on the use of HFCs in this sector has been provided by industry. From this information the number of systems and the quantity of HFCs present in the market has been estimated for the time series.

These systems were first introduced into the Irish market in 1996 so emissions are not occurring (NO) prior to 1996. The emission calculation methodology used for this category is a Tier 2 emission model. The model estimates emissions from three situations where emissions may occur:

- The first situation is from discharge (intentional and accidental). Although a major company within this sector has not recorded any discharges, they do apply the assumption that each system will discharge once over a ten year period. This conservative assumption has been applied within the model.
- The second source covers leakage emissions and is estimated as 1 per cent of the total charge for all systems present.
- The third source is from the decommissioning of systems, but this activity has not yet taken place in Ireland.

4.7.3.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 20 per cent and the uncertainty of the emission factor is 10 per cent.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams et al., 2005) and reviewed in 2013 (Goodwin et al., 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.7.3.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.7.3.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.7.3.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.7.4 Aerosols (2.F.4)

4.7.4.1 Category Description

For the purposes of estimating emissions, Aerosols and Metered Dose Inhalers are treated separately in the emission estimations. This category includes the following sub-categories;

- Metered dose inhalers 2.F.4.a,
- Other-Aerosols 2.F.4.b.

Most aerosol packages contain hydrocarbon (HC) as propellants but, in a small fraction of the total, HFCs and PFCs may be used as propellants or solvents. Emissions from aerosols usually occur shortly after production, on average six months after sale.

4.7.4.2 Methodological Issues

Emission estimates for Metered Dose Inhalers (MDI) 2.F.4.a are made on the basis of data received from industry for manufacturing emissions, and population data coupled with emission factors for emissions from use. The HFCs used in MDI's in Ireland are HFC-134a and HFC-227ea.

Process losses are based on an analysis of gross stock minus closing stock and usage data of the gases. The MDI market in Ireland is supplied by both Irish manufactured products and imported products. Irish manufactured products only contain HFC-134a based on annual industry returns and Adams et al. (2005). Imported products on the other hand can contain HFC-134a and HFC-227ea. As a result there is no emissions from manufacture in for HFC-227ea in CRF Table2(II)B-Hs2. Total emissions are calculated based on reported manufacturing losses (for HFC-134a) in conjunction with in-life emissions.

Ireland has a high prevalence of asthma and in order to reflect this country-specific circumstance, a bottom-up approach to estimating in-life emissions is applied. Approximately 10 per cent of the Irish population are suffering from asthma (Asthma Survey by the Irish Asthma Society) and about 80 per cent of the asthma medication sold relates to MDIs (Asthma Support Team of a large pharmacy chain) with the remaining 20 per cent relating to DPIs.

A calculation based on population and these data was undertaken in order to establish an estimate for the total annual demand. This demand is catered for by imported products from a number of manufacturers as well as those manufacturing in Ireland.

Information on the amount of HFCs contained in MDIs per patient was determined empirically at approximately 0.074kg per user per annum (Schwarz et al., 2010). Furthermore, it was estimated that of the HFCs used in MDIs in Ireland, HFC-134a accounted for 90 per cent and HFC-227ea for 10 per cent. HFC-227ea is mainly used by a non-Irish, European MDI producer. These data were used for the estimation of lifetime emissions.

The category Other-Aerosols 2.F.4.b, is one which can cover a large number of products, however HFC's are generally only used as propellants where the use of HFCs is considered critical. The two HFCs of interest are HFC-134a and HFC-152a.

In submissions up to and including 2011 the assumed species ratio was 90 per cent: 10 per cent, respectively for HFC-134a and HFC-152a. That assumption was revised in 2012 submission (1990-2010 data) on the basis of updated information (AEA 2011) which suggests a species ratio of 97 per cent: 3 per cent in 1990 to 2004, 99 per cent: 1 per cent for 2005 and 96 per cent: 4 per cent for 2006 onwards, respectively for HFC-134a and HFC-152.

There is no trade association for aerosol manufacturers or importers in Ireland. Furthermore Adams *et al* (2005) found that importation of HFC containing aerosols is carried out independently by retailers. As a result little information exists in relation to the Irish market for these products.

Following consultations with the British Trade Association (BAMA), O’Leary *et al.* (2002) and Adams *et al.* (2005) recommended the use of a population based proxy to estimate Irish emissions from those for the UK, which are based on trade data for the UK, on the assumption that the market for aerosols would be similar in Ireland. Emissions of HFC-134a and HFC-152a from aerosols are therefore derived using the UK estimates for lifetime and decommissioning emissions (as used in the UK national GHG inventory) and the ratio of the Irish population (CSO) to the UK population (Office of National Statistics, UK) in each year. The estimate for potential emissions is calculated using the UK trade data and the population ratio.

4.7.4.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 20 per cent and the uncertainty of the emission factor is 10 per cent.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams *et al.*, 2005) and reviewed in 2013 (Goodwin *et al.*, 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.7.4.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.7.4.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.7.4.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.7.5 Solvents (2.F.5)

There are no known emissions from solvents in Ireland. This category is reported as Not Occurring.

4.7.6 Other Product Uses as Substitutes for ODS (2.F.6)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

4.8 Emissions from Other Product Manufacture and Use (2.G)

Emission estimates for category 2.G are presented in Table 4.3 and Figure 4.3 above. This category includes the following sub-categories;

- Electric equipment 2.G.1,
- SF₆ and PFCs from other product use 2.G.2,

1. Soundproof windows 2.G.2.c,
 2. Adiabatic properties: shoes and tyres 2.G.2.d,
 3. Other-Medical Applications and Tracer in Leak Detection 2.G.2.e,
- N₂O from Product Uses,
 1. Medical Application 2.G.3.a.

4.8.1 Electrical Equipment (2.G.1)

4.8.1.1 Category Description

SF₆ is used for electrical insulation, arc quenching, and for current interruption in equipment used in the transmission and distribution of electricity. The Electricity Supply Board (ESB) is the owner of both the high and low voltage distribution systems and the owner and operator of the medium and lower voltage distribution systems in Ireland.

Electrical equipment containing SF₆ is imported into Ireland. Quantities of SF₆ are needed for servicing and repair of existing equipment. There are, therefore, no manufacturing emissions.

Significant reduction in emissions in the years 2008 to 2010 are attributed to the network operators investment in staff training, leak detection equipment and closed cycle SF₆ handling equipment. This resulted in 3 year rolling average losses. The latest increase in 2011 is due the highest installed inventory stock levels occurring in the period 2009 to 2011, but losses remain low around 0.5 per cent.

4.8.1.2 Methodological Issues

Emissions are estimated using a Tier 1 approach based on an analysis of opening and closing stocks of SF₆. The inventory estimates assume that the usage of SF₆ in equipment maintenance for one year is equal to the leakage emissions from electrical equipment in the same year. This method was reviewed by the project team and deemed to be acceptable and in line with IPCC GPG.

The company supplies an estimate of SF₆ emissions from their equipment maintenance operations to the inventory agency on a yearly basis. Those annual SF₆ usage returns include:

- The number of cylinders that are booked out for each year;
- Cylinder size: 40kg, 63kg and 5kg;
- Assumption that for the 40kg and 63kg cylinders, 60 per cent of the contained SF₆ is used for maintenance and the remaining 40 per cent is either unused or used for new works and thus not related to leakage emissions;
- Assumptions that one third of the SF₆ in the smaller cylinders (5kg) is used for maintenance whilst the remaining quantity is used for new works.

4.8.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams et al., 2005) and reviewed in 2013 (Goodwin et al., 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.8.1.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.8.1.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.8.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.8.2 SF₆ and PFCs from Other Product Uses (2.G.2)

4.8.2.1 Category Description

Emissions of SF₆ are included in this category from the following activities:

- **Soundproof windows 2.G.2.c** – SF₆ was previously used as an insulation gas in double-glazing; however its use has been phased out in response to F-gas regulations and is assumed not to have occurred since 2000
- **Adiabatic properties 2.G.2.d**– SF₆ is used as a cushioning agent in sports shoes due to its chemically and biologically inert properties and its high molecular weight, which means that it does not diffuse across membranes; thus the gas is not released until the sports shoe is destroyed at the end of its useful life
- **Medical applications 2.G.2.e** – SF₆ is used in certain medical applications such as eye surgery where it is used to seal retinal holes internally and to hold reattached retina in place
- **Tracer in Leak Detection 2.G.2.e**–SF₆ has been used as a tracer gas for leak detection in agricultural research to determine the rates of methane emissions from enteric fermentation in cattle.

4.8.2.2 Methodological Issues

Emission estimations from **Soundproof windows 2.G.2.c** account for opening and closing stock of the gas, assembly losses for Irish manufactured products, stocks in imported windows, leakage once installed and disposal emissions. Even though the use of SF₆ was discontinued in window insulation after 2000, the bank of gas in installed units is an emission source and is therefore accounted for in emission estimates.

A life-time of 25 years was applied; therefore, emissions at disposal are calculated as 100 per cent of the remaining charge after 25 years of leakage at a rate of 1 per cent per annum. The entire quantity of SF₆ remaining inside the window at the end of life is emitted, because to-date no recovery process exists.

There is no specific information available in relation to the use of SF₆ in **Adiabatic properties 2.G.2.d** (sports goods, shoes) in Ireland, so a population-proxy is used to estimate emissions based on UK inventory data for the release of SF₆ upon disposal of sporting goods, as the market share of such

products is assumed to be similar to that in the UK. Emissions are therefore derived using the annual UK per capita sales for sporting goods and the Irish population in each year.

The use of SF₆ in sporting goods was discontinued in 2007, however, emissions from their disposal continued until 2013 based on an average lifetime of eight years.

Use of SF₆ in **Medical applications 2.G.2.e** is small with one hospital reporting the use of one 10-litre cylinder every three years. Based on this data, it is assumed that a similar quantity is used in a total of 10 hospitals, which undertake similar procedures. It is assumed that actual and potential emissions are equal on the basis that in each of the 10 hospitals once a cylinder is used (over a three year period) it is replaced.

The use of SF₆ as a **Tracer in Leak Detection 2.G.2.e** was previously a relatively large source in the period 1990-2004. However the company who used SF₆ for the purpose of leak detection has since ceased trading.

A number of research projects, conducted in 2009, were identified and included in the inventory: maize experiment – emission rate of 1.8 mg SF₆/day from 60 capsules (1/animal) for 105 days; whole-crop wheat experiment – emission rate of 3.14 mg SF₆/day from 90 capsules (1/animal) for 154 days. Calculated emissions from these two experiments were used to estimate emissions from a third research project similar to these two. No projects since have been identified so this sub-category is no longer a source of emissions of SF₆ in the Irish inventory.

4.8.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

An uncertainty analysis was performed for the aggregated emissions derived from a specific consideration of the individual sector uncertainty estimates (Adams et al., 2005) and reviewed in 2013 (Goodwin et al., 2013). An iterative Monte Carlo simulation procedure was used to estimate uncertainties in total and aggregated HFC, PFC and SF₆ emissions.

4.8.2.4 Category-specific QA/QC and verification

Quality control checks are in place to ensure that the sectoral emissions total in calculation sheets is the same as that in the final inventory dataset that is reported to the UNFCCC. The current inventory approach was reviewed at the start of the F-gas 2013 project. All calculations and cell references were checked to assess the current approach.

4.8.2.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

4.8.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.8.3 N₂O from Product Use (2.G.3)

4.8.3.1 Category Description

Evaporative/fugitive emissions of nitrous oxide (N₂O) can arise from various types of product use, including;

- Medical applications (anaesthetic use, analgesic use and veterinary use);
- Use as a propellant in aerosol products, primarily in food industry (pressure-packaged whipped cream, etc);
- Oxidising agent and etchant used in semiconductor manufacturing;
- Oxidising agent used, with acetylene, in atomic absorption spectrometry;
- Production of sodium azide, which is used to inflate airbags;
- Fuel oxidant in auto racing; and
- Oxidising agent in blowtorches used by jewellers and others.

In general, medical applications and use as a propellant in aerosol products are likely to be larger sources than others.

The use of N₂O as an anaesthetic in hospitals is a source of emissions and has been estimated in this submission. Emission estimates for Medical application of N₂O category 2.G.3.a are presented in Table 4.2.

4.8.3.2 Methodological Issues

In absence of methodologies or emission factors in the existing guidelines, population-based activity data has been developed with assumed usage of 3 grams of N₂O per capita per year and emission factor of 1 (as all used gas is emitted into the atmosphere). This assumption is the same as amongst other Annex I Parties calculating emissions from this category.

4.8.3.3 Uncertainties and Time-series Consistency

The uncertainties applicable to category N₂O from Product Use are provided in Annex 2. The emission time series for 1990–2013 is consistent. Key activity data such as Ireland's population statistics are available for all applicable years and are used in a consistent manner.

4.8.3.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to this category. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

4.8.3.5 Category-specific Recalculations

Emissions of N₂O from Anaesthesia were included for the first time in this submission.

4.8.3.6 Category-specific Planned Improvements

There are no planned improvements for this category.

4.8.4 Other – Other Product Manufacture and Use (2.G.4)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

4.9 Other – (2.H)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

Chapter 5 Agriculture

5.1 Overview of Agriculture Sector

The list of activities under *Agriculture* in the IPCC reporting format is given in Table 5.1 below. A summary of emissions from these activities are given in Table 5.2, Figure 5.1 and Figure 5.2 below.

Enteric fermentation, Manure Management and Agricultural Soils are the main activities that give rise to greenhouse gas emissions in the *Agricultural* sector (Table 5.1).

Estimates are included for all emission sources that occur in the country and the required level of disaggregation is achieved for detailed completion of the CRF tables.

5.1.1 Emissions Overview

There are seven key categories in this sector:

- **3.A.1 Enteric Fermentation, Cattle** (Trend and Level)
- **3.A.2 Enteric Fermentation, Sheep** (Trend and Level)
- **3.B.1.1 Manure Management (CH₄), Cattle** (Level)
- **3.B.2.5 Manure Management (N₂O), Indirect N₂O Emissions** (Level)
- **3.D.1 Direct Soil Emissions** (Trend and Level)
- **3.D.2 Indirect Soil Emissions** (Level)
- **3.G Liming** (Trend and Level)

Other categories present in this sector include:

- **3.A.3 Enteric Fermentation, Swine**
- **3.A.4 Enteric Fermentation, Other Livestock**
- **3.B.1.2 Manure Management (CH₄), Sheep**
- **3.B.1.3 Manure Management (CH₄), Swine**
- **3.B.1.4 Manure Management (CH₄), Other Livestock**
- **3.B.2.1 Manure Management (N₂O), Cattle**
- **3.B.2.2 Manure Management (N₂O), Sheep**
- **3.B.2.3 Manure Management (N₂O), Swine**
- **3.B.2.4 Manure Management (N₂O), Other Livestock**
- **3.H Urea Application**

The greenhouse gases relevant to *Agriculture* are as follows:

- **Carbon dioxide** emissions originate from *3.G Liming* and *3.H Urea Application*. Carbon dioxide emissions have increased by 34.5 per cent over 1990-2013, mostly due to a 45.3 per cent increase in emissions from liming.
- **Nitrous Oxide** emissions originate from *3.B Manure Management* and *3.D Agricultural Soils*.

- **Methane** emissions originate from 3.A *Enteric Fermentation* and 3.B *Manure Management*. Methane is the most significant GHG in agriculture, and contributed 62.2 per cent of agricultural emissions in 2013.

The 2015 submission shows total GHG emissions of 18,964.61 kt CO₂ equivalent in the *Agriculture* sector in 2013, of which 3.A *Enteric Fermentation* accounts for 55.6 per cent 3.D *Agricultural Soils* 32.2 per cent, 3.B *Manure Management* 9.3 per cent, 3.G *Liming* 2.7 per cent, and 3.H *Urea Application* 0.1 per cent. The latest estimates show that emissions in the *Agriculture* sector have decreased by 8.5 per cent from 1990 to 2013 mainly due to a 7.1 per cent decrease in CH₄ emissions from 3.A *Enteric Fermentation* and 14.4 per cent decrease in N₂O emissions from 3.D *Agricultural Soils*.

5.1.2 Methodology Overview

A summary of the Tier methods, consistent with the 2006 IPCC Guidelines, is provided in Table 5.1 below, along with a summary of the activities applicable to Ireland.

Table 5.1 Level 3 Source Methodology for Agriculture

3. Agriculture	CO ₂	CH ₄	N ₂ O
A. Enteric Fermentation			
1. Cattle*	NA	T2*	NA
2. Sheep*	NA	T1*	NA
3. Swine	NA	T1	NA
4. Other Livestock	NA	T1	NA
B. Manure Management			
1. Cattle*	NA	T2*	T1
2. Sheep	NA	T1	T1
3. Swine	NA	T1	T1
4. Other Livestock	NA	T1	T1
5. Indirect N ₂ O emissions*	NA	NA	T2*
C. Rice Cultivation	NO	NO	NO
D. Agricultural Soils			
1. Direct N ₂ O from Managed Soils*	NA	NA	T1*
2. Indirect N ₂ O from Managed Soils*	NA	NA	T1*
E. Prescribed Burning of Savannas	NO	NO	NO
F. Field Burning of Agricultural Residues	NO	NA	NA
G. Liming*	T1*	NO	NO
H. Urea Application	T1	NO	NO
I. Other Carbon-containing fertilisers	NA	NO	NO
J. Other	NO	NO	NO

* Key Category.

T1,2,3: Tier 1, Tier 2, Tier 3 as described in the 2006 IPCC Guidelines;

NE : emissions not estimated;

NO : activity not occurring;

NA : not applicable (no emissions of the gas occur in the source category);

IE : emissions included elsewhere.

There is extensive and up-to-date statistical data on all aspects of the agriculture sector in Ireland. The majority of this data is compiled and published by the CSO and is the official source of the basic data for inventory purposes. The exception is for statistics on synthetic fertiliser use and the poultry population which are obtained from the Department of Agriculture Food and the Marine (DAFM). The CSO and DAFM are key data providers whose annual statistical inputs to the inventory agency are covered by Memorandum of Understanding (MOU) in Ireland's national system. The time-series of key agricultural statistics, as used for the various activity data (e.g. livestock populations and fertiliser use) are given in Table 3.3.A of Annex 3.3.

There is significant collaboration between inventory experts, agriculture researchers and the Department of Agriculture, Food and Marine, which grew out of the improved inventory methodologies for both CH₄ and NH₃. These collaborations are maintained by the inventory agency and are an important part of the overall QA/QC procedures being undertaken on an annual basis.

Table 5.2 Emissions from Agriculture 1990-2013

		Gas	Unit	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
3.A.1	Cattle	CH ₄	kt CO ₂ eq	10101.4	10190.7	10049.7	10026.4	9958.9	9948.2	9931.1	9839.8	9835.6	9709.4	9723.2	9613.3	9424.1	9298.9	9589.3	9758.1
3.A.2	Sheep	CH ₄	kt CO ₂ eq	1176.3	1195.2	1114.4	1055.9	990.6	963.4	960.8	904.7	852.3	778.8	716.1	666.7	637.3	650.6	695.1	696.1
3.A.3	Swine	CH ₄	kt CO ₂ eq	41.4	52.0	56.9	57.3	58.4	56.9	56.1	55.2	55.3	51.1	50.1	48.9	51.1	51.8	51.3	49.9
3.A.4	Other livestock	CH ₄	kt CO ₂ eq	37.9	42.3	39.8	40.2	40.6	39.7	40.4	43.4	46.2	47.7	51.2	52.2	53.5	52.7	54.8	49.6
3.B.1	Cattle	CH ₄	kt CO ₂ eq	1038.8	1004.8	949.1	945.5	936.3	923.1	917.8	924.6	918.8	910.4	916.9	904.4	879.8	870.1	908.7	919.0
3.B.1	Cattle	N ₂ O	kt CO ₂ eq	241.1	252.2	254.4	257.1	257.9	256.6	255.6	261.0	256.9	256.4	257.9	256.5	245.3	239.6	252.6	257.6
3.B.2	Sheep	CH ₄	kt CO ₂ eq	81.9	84.3	78.3	73.6	67.6	65.6	66.8	62.6	59.6	54.5	49.8	46.1	43.1	44.1	47.7	48.0
3.B.2	Sheep	N ₂ O	kt CO ₂ eq	18.3	18.4	17.7	17.0	16.5	16.0	15.5	14.9	13.8	12.6	11.7	11.0	10.9	11.1	11.6	11.5
3.B.3	Swine	CH ₄	kt CO ₂ eq	160.2	198.7	219.2	223.0	225.3	218.2	215.3	212.6	208.1	195.4	189.1	184.3	192.5	195.6	193.1	190.4
3.B.3	Swine	N ₂ O	kt CO ₂ eq	10.1	12.5	13.6	13.8	13.9	13.5	13.4	13.2	13.1	12.2	11.9	11.6	12.1	12.2	12.0	11.8
3.B.4	Other livestock	CH ₄	kt CO ₂ eq	61.3	49.9	54.6	57.1	55.3	62.1	63.2	63.8	64.1	60.2	60.7	68.7	68.9	67.3	80.9	84.9
3.B.4	Other livestock	N ₂ O	kt CO ₂ eq	10.3	9.6	9.6	9.8	9.7	9.8	10.2	10.3	10.4	10.2	10.5	11.3	11.5	11.4	12.2	11.6
3.B.5	Indirect N ₂ O emissions	N ₂ O	kt CO ₂ eq	216.2	226.4	228.3	230.2	230.7	229.4	228.5	228.2	226.3	222.5	221.7	219.8	216.2	215.2	222.7	225.4
3.D.1	Direct N ₂ O Emissions From Managed Soils	N ₂ O	kt CO ₂ eq	6581.4	6252.9	6214.0	6637.4	6261.7	6562.9	6307.5	6479.7	6178.8	5383.3	5533.4	5758.5	5922.3	5573.2	5871.5	5633.2
3.D.2	Indirect N ₂ O Emissions from Managed Soils	N ₂ O	kt CO ₂ eq	559.2	539.4	538.4	567.6	539.6	561.0	538.1	550.2	527.6	460.9	478.9	503.4	516.3	478.6	496.2	480.1
3.G.1	Limestone CaCO ₃	CO ₂	kt	355.0	494.6	366.4	385.3	273.9	386.8	240.8	266.7	254.9	376.8	262.2	307.3	427.9	360.7	229.4	515.7
3.H	Urea Application	CO ₂	kt	44.5	39.7	42.2	38.5	37.2	36.1	30.8	27.9	29.6	23.4	30.8	40.9	45.2	32.3	21.3	21.7
3	Total Agriculture		kt CO₂eq	20735.4	20663.6	20246.8	20635.5	19974.0	20349.2	19891.7	19959.0	19551.3	18565.7	18576.1	18705.0	18758.0	18165.3	18750.3	18964.6




Figure 5.1 Total Emissions from Agriculture by Sector, 1990-2013




Figure 5.2 Total Emissions from Agriculture by Gas, 1990-2013

Emissions from livestock (3.1)

The two IPCC Level 2 emission source categories under 3.1 *Livestock* in 2013 are 3.A *Enteric Fermentation* and 3.B *Manure Management*. Total emissions from these activities amounted to 12,313.99 kt CO₂eq in 2013.

Two large research projects have greatly contributed to improving the estimation of emissions from enteric fermentation and manure management in Ireland:

- O'Mara (2007), major research funded by the EPA;
- Hyde et al. (2008), an extensive Farm Facilities Survey.

This research, along with other relevant work related to the development of a nitrogen-flow approach to NH₃ emissions, has facilitated the application of a large amount of country-specific information underlying the various estimates of emissions.

The livestock types relevant for Ireland are as follows;

- Dairy Cattle
- Non-Dairy Cattle
- Sheep
- Swine
- Other livestock;
 - Deer
 - Goats
 - Horses
 - Mules and Asses
 - Poultry
 - Fur-bearing Animals

5.2 Emissions from Enteric Fermentation (3.A)

The IPCC Level 3 emission source categories relevant under *3.A Enteric Fermentation* in 2013 are *3.A.1 Cattle*, *3.A.2 Sheep*, *3.A.3 Swine*, and *3.A.4 Other Livestock*. Total emissions from these activities amounted to 10,553.71 kt CO₂eq in 2013.

5.2.1 Enteric Fermentation, Cattle (3.A.1)

5.2.1.1 Category Description

Methane is produced in herbivores as a by-product of enteric fermentation, a digestive process by which carbohydrates are broken down by micro-organisms into simple molecules for absorption into the bloodstream. The amount of methane that is released depends on the type of digestive tract, age, and weight of the animal, and the quality and quantity of the feed consumed. Ruminant livestock (e.g., cattle, sheep) are major sources of methane with moderate amounts produced from non-ruminant livestock (e.g., pigs, horses).

Enteric fermentation from cattle is both a trend and level key category of CH₄ in Ireland.

5.2.1.2 Methodological Issues

The Tier 2 approach in the 2006 IPCC Guidelines is used for *Enteric Fermentation, Cattle*. The Tier 2 approach has been used for 1990 and for the years 2003 to 2013. Interpolation has been used to complete the time series.

In the Tier 2 approach, the Irish cattle herd is characterised by 11 principal animal classifications as shown in Table 5.3 for which annual census data are published by the CSO. In-depth analysis of production systems and the associated animal feed and energy requirements was conducted for all categories within the Irish cattle population to determine CH₄ production. Substantial further subdivision was incorporated for dairy and beef cattle to adequately describe the wide range of cattle rearing and finishing systems applicable in Ireland. In total, dairy cows were covered by 12 systems and 18 system types were analysed for suckler cows, while up to 30 systems were examined for both male and female beef cattle (O'Mara, 2007).

Table 5.3 Animal Classifications for Cattle Population

Cattle Type	Classification		
	Dairy cows	Suckler (Beef) cows	
Beef cattle	Male < 1 year	Male 1 – 2 years	Male > 2 years
	Female < 1 year	Female 1 – 2 years	Female > 2 years
Other cattle	Breeding bulls	Dairy in-calf heifers	Beef in-calf heifers

For both **dairy cows** and **suckler cows**, the country is divided into three regions: (1) south and east, (2) west and midlands, and (3) north-west, coinciding with the regions used for the implementation of regulations on Good Agricultural Practices for the protection of Waters:

- SI 788 of 2005 (DEHLG, 2005)
- SI 378 of 2006 (DEHLG, 2006)
- SI 101 of 2009 (DEHLG, 2009)
- SI 610 of 2010 (DEHLG, 2010)
- SI 31 of 2014 (DECLG, 2014)

This division facilitates in-depth analysis of separate regions with different lengths of winter housing and takes account of different animal feeding practices. The cattle production systems in each region are defined in terms of calving date, the dates of winter housing and spring turn-out to grass, milk yield and composition, forage and concentrate feeding level, cow live-weight and live-weight change and lactation period. The number of cows in each category, given by CSO statistics, is allocated to the three regions identified above using the Cattle Movement Monitoring System (CMMS) and Animal Identification and Movement (AIM) system reports published by the Department of Agriculture, Fisheries and Food (DAFF, 2004; 2005; 2006; 2007; 2008, 2009, 2010) and the Department of Agriculture Food and the Marine (DAFM, 2011, 2012, 2013, 2014). The CSO produces two censuses of animal numbers per year, one reflecting the number of animals nationally in June and the other referring to populations in December. For the purposes of calculating emissions from breeding cattle, an average of the number in each category of breeding animals present in the national herd in June and December is used¹.

In the approach outlined by O'Mara (2007), the daily energy requirement of cows in each region is calculated by month or part thereof based on maintenance requirements, milk yield and composition, requirements for foetal growth and gain or loss of bodyweight using the French energy system (INRA, 1989). In this system, net energy requirement is defined in terms of *unites fourragere lait* (UFL), where 1 UFL is the net energy value of 1 kg of barley at 86 per cent dry matter and is equal to 7.11 MJ net energy for lactation (NE_l). This international energy system, which is well established and used locally in Ireland, was considered more appropriate to the local conditions than the system and equations used by the IPCC guidelines and IPCC good practice guidance. The energy gains and losses refer to intra-annual changes for the animal and do not mean that average body weight for

¹ The publication of separate census data for June and December annually and the application of these statistics in order to achieve the most representative annual average population related to cattle explains differences that are often seen between national and FAO statistics for agriculture. Ireland has high quality agricultural statistics and differences with FAO are to be expected, but they are of no consequence to the emissions estimates.

animals in the dairy herd is increasing from year to year. The live-weight of 535 kg for dairy cows is an indicative weight supplied by the Department of Agriculture, Food and the Marine, as dairy cow live-weights are not in general monitored on farms. The live-weight is adopted as the reference point for the annual emission factor derivation for the herd and is chosen to be consistent with other parameters relevant to the estimation of emissions from cattle, e.g. manure production.

The important equations are:

Maintenance NE_l requirements (MJ) = $9.96 + (0.6 \times LW/100)$, where LW is live-weight. A 10 per cent activity allowance was added for the housed period and a 20 per cent allowance was added for the grazing period as outlined by INRA (1989);

NE_l (MJ) required per kg milk = $0.376 \times \text{fat content} + 0.209 \times \text{protein content} + 0.948$;

Pregnancy: mean of 12.1 MJ NE_l /day for the last 3 months of pregnancy;

Live-weight change: each kg live-weight lost contributed 24.9 MJ NE_l to energy requirements, while each kg of live-weight gained required 32 MJ NE_l .

The composition of the diet of cows in each region was described by month or part thereof and daily intake was calculated by reference to the daily energy requirement. The concentrate allowance was fixed while forage intake varied according to energy requirements. Daily methane emissions (MJ/day) were calculated from digestible energy intake using the equation of Yan et al. (2000).

$$CH_4 = DEI * [0.096 + (0.035 \times S_{DMI}/T_{DMI})] - 2.298 * (FL - 1)$$

where DEI is digestible energy intake (MJ/day), S_{DMI} and T_{DMI} are silage and total dry matter intakes (kg/day), respectively, and FL is feeding level (multiples of the maintenance energy requirement).

A constant methane conversion rate of 0.065 of gross energy intake is applied when the diet consists of grazed grass and 3 kg or less of concentrate supplement per day. This is based on a large New Zealand database of measurements for grazing animals on similar production systems to those in Ireland. A methane output of 21.6 g/kg DM is used for pasture diets with a grass GE content of 18.45 MJ/kg, which is equivalent to 6.5 per cent of GE (Harry Clark, AgResearch New Zealand Personal Communication). Daily CH_4 emissions are summed to give annual emissions for cows in each region, and a weighted national average emission factor is then calculated.

Emission factors for the **beef cattle** categories, given in Table 5.3, are determined by calculating lifetime emissions for the animal and by partitioning between the first, second and third years of the animal's life. This approach allows the published CSO animal populations for June to be used directly as the activity data most representative of the inventory year for enteric fermentation while taking into account the movement of cattle from one category to another (i.e. from 0-1 year old to 1-2 year old to over 2 years old), as enumerated by the June census, up to two times in their three-year lifetime (O'Mara 2007).

Important parameters such as housing dates (expert opinion and Hyde et al., 2008), turnout dates (expert opinion and Hyde et al., 2008) and live-weight gains (expert opinion reconciled with actual national carcass weights) during winter housing periods and grazing seasons are defined for each system (O'Mara, 2007). The most important parameter for beef cattle is live-weight gain, as it directly affects the energy requirement and thus the feed intake. There is little statistical information on the live-weight gain of the different types of cattle in the cattle herd, but the weight of carcasses of all slaughtered cattle is recorded by the Department of Agriculture, Food and the Marine. Using data for the average carcass weight of male and female cattle, appropriate live-weight gains are applied to the various life stages of each animal category, such that when all categories are

combined, that data is consistent with the national statistics for carcass weight (plus or minus 10 kg difference).

Given these data for live-weight and live-weight gain, O'Mara, (2007) estimated the energy requirements of animals during the winter housing periods and grazing seasons of the animals lifetime using the INRAtion computer programme, version 3.0. This programme was devised by the French research organisation Institute National de la Recherche Agronomique (INRA) and is based on the net energy system for cattle. In version 3 of INRAtion, some adaptation for Irish conditions was made to the equations for estimating the energy requirements of growing and finishing animals (O'Mara, 1997, Crowley, 2001 and Crowley *et al*, 2002). Net energy requirements of growing beef cattle are defined in terms of UFL, as in the case of dairy cattle, while for finishing cattle, net energy requirements are defined in terms of UFV (from the French *unite fourragere viande*) where 1 UFV is the net energy value of 1 kg of barley for meat production and is equal to 7.61 MJ NE_{mg}.

The composition of the diet in each system is described by grazing season and winter housing period and daily intake is calculated by reference to the daily energy requirement. The concentrate allowance is fixed while forage intake is varied according to energy requirements. The Irish modifications to the INRAtion programme were predominantly for animals at weanling and finishing stages (i.e. at times that concentrates were likely to be fed). No modifications were made for 'heavy' growing animals, (typically animals in their second grazing season or later that were not being finished). For animals in these stages, intakes were adjusted as appropriate by expert opinion. Daily methane emissions were calculated using the equation of Yan *et al*. (2000), however a constant of 0.065 of gross energy intake was applied when the diet was grazed grass plus 3 kg or less of concentrate supplement/day. Daily emissions are aggregated to give annual emissions per system and a weighted national average emission factor is then calculated.

Bulls for breeding and in-calf heifers account for on average 6 per cent of the national cattle herd. Separate production systems were not defined for these categories because of the lack of published data on their feed intake and the small number of animals involved (O'Mara, 2007). Bulls for breeding are mostly of continental breeds, and their emission factors are based on those for late maturing male beef cattle of suckler origin in their second year. The emission factor for animals in this category is determined by an applicable period of 310 days in their second year, which is adjusted upwards to the full period of 365 days in the case of breeding bulls.

In-calf heifers are assigned the same emission factors as female beef cattle in their second year (i.e. corresponding to the category 1–2 years old). In-calf heifers only require emissions associated with the period March – December of their second year to be accounted for, as they are subsequently enumerated as dairy or suckler cows in the CSO animal census thereafter. Female beef cattle in the category 1-2 years old are assumed to be slaughtered on 3rd February of their third year (O'Mara, 2007). Adjustment for the slightly longer period is not made in respect of in-calf heifers, as they are carrying a calf in addition to normal growth which is reflected in the calculation methodology.

Table 5.4 Tier 2 CH₄ Enteric Fermentation Emission Factors for 1990, 2007- 2013

	Enteric Fermentation (kg/head/year)							
	1990	2007	2008	2009	2010	2011	2012	2013
Dairy cows	101.38	111.29	109.99	108.54	113.06	113.22	110.84	111.57
Suckler cows	74.03	73.22	74.92	72.78	72.95	74.07	75.53	73.12
Male cattle < 1 year	30.46	29.69	29.71	29.77	29.11	29.82	30.15	30.26
Male cattle 1 - 2 years	62.22	59.19	59.07	58.57	59.96	58.01	56.63	56.20
Male cattle > 2 years	55.08	38.58	36.98	38.84	39.79	38.29	37.25	37.32
Female cattle < 1 year	27.05	27.61	27.60	27.57	27.55	27.60	27.70	27.73
Female cattle 1 - 2 years	53.54	46.60	47.00	47.71	48.62	47.93	47.99	48.08
Female cattle > 2 years	21.65	22.42	22.55	22.63	22.63	22.72	22.73	22.64
Bulls for breeding	86.38	81.55	81.55	81.55	81.55	81.55	81.55	81.55
Dairy in-calf heifers	51.82	50.16	50.16	50.16	50.16	50.16	50.16	50.16
Beef in-calf heifers	55.42	53.68	53.68	53.68	53.68	53.68	53.68	53.68

5.2.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Enteric Fermentation* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as disaggregated animal numbers are available for all years and are used in a consistent manner.

5.2.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Enteric Fermentation, Cattle*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.2.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.2.1.6 Category-specific Planned Improvements

The Department of Agriculture, Food and the Marine has funded the establishment of The Agricultural Greenhouse Gas Research initiative for Ireland (AGRI-I). This is an organisational and collaborative framework designed to: build a critical mass of scientific expertise in GHG research, co-ordinate uniform measurement protocols, and address a specific set of research issues. The AGRI-I network has a specific set of research aims, primarily focussed on the inclusion of validated GHG emissions mitigation strategies into the national inventory. This research will also include a review of feed intake parameters and assumed nitrogen content of feeds and updates as necessary. A separate but related research project is investigating the development of country specific B₀ and MCF values using a range of cattle manures and environmental conditions.

5.2.2 Enteric Fermentation, All Other Livestock (3.A.2-3.A.4)

5.2.2.1 Category Description

This grouping includes **sheep 3.A.2**, **swine 3.A.3**, and **other livestock 3.A.4**. Enteric fermentation from other livestock in Ireland consists of **deer, goats, horses, mules and asses**, and **fur-bearing animals**. Enteric Fermentation emissions of CH₄ are not occurring for poultry and fur-bearing animals. *Enteric fermentation, sheep 3.A.2* is a key category in Ireland.

5.2.2.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC guidelines is used for *Enteric Fermentation; Sheep, Swine, and Other Livestock*.

The type of information used to derive the Tier 2 emission factors for cattle is not available for other important livestock categories in Ireland, such as **sheep** and **swine**. Therefore, the inventory agency continues to use the Tier 1 approach for enteric fermentation for all livestock categories other than cattle. The emission factors used are generally those for Western Europe given in Table 10.10 of the 2006 IPCC guidelines. However, in order to fully utilise Irish national statistics and the detailed CSO breakdown in respect of sheep and swine populations, the base emission factors from IPCC are adjusted as shown in Table 3.3.B of Annex 3.3. For sheep, the emission factor for lambs is calculated on the basis that lambs have an assumed lifetime of 180 days before slaughter and a CH₄ conversion rate (Y_m) of 0.06 as per Table 10.13 of the 2006 IPCC guidelines. For swine the default emission factor of 1.5 kg CH₄ per head (Table 10.10 of 2006 IPCC guidelines) per year is adjusted for each subcategory of swine on the basis of a default swine weight of 82 kg and the known average weight of each subcategory of swine in Ireland. As a result, the implied emission factors produced by the CRF related to total populations of sheep and swine in Ireland are relative to the IPCC base default values for these animal categories.

5.2.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Enteric Fermentation* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as disaggregated animal numbers available for all years and are used in a consistent manner.

5.2.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Enteric Fermentation*, *All Other Livestock*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.2.2.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.2.2.6 Category-specific Planned Improvements

The inventory agency is in the process of investigating the applicability of developing Tier 2 estimates of CH₄ from enteric fermentation from sheep as recommended in the previous annual inventory review reports. Discussions are on-going with agricultural research institutions in relation to sourcing appropriate country specific data with respect to feed intake for sheep in Ireland.

5.3 Emissions from Manure Management (3.B)

The IPCC Level 3 emission source categories relevant under *3.B Manure Management* in 2013 are *3.B.1 Cattle*, *3.B.2 Sheep*, *3.B.3 Swine*, *3.B.4 Other Livestock*, and *3.B.5 Indirect N₂O Emissions*. Total emissions of CH₄ and N₂O from these activities amounted to 1,760.28 kt CO₂eq in 2013.

5.3.1 Manure Management, Cattle (3.B.1)

5.3.1.1 Category Description

This category describes how to estimate CH₄ produced during the storage and treatment of manure, and from manure deposited on pasture. The term 'manure' is used here collectively to include both dung and urine (both the solids and the liquids) produced by livestock. The main factors affecting CH₄ emissions are the amount of manure produced and the portion of the manure that decomposes anaerobically. The former depends on the rate of waste production per animal and the number of animals, and the latter on how the manure is managed. When manure is stored or treated as a liquid (e.g., in lagoons, ponds, tanks, or pits), it decomposes anaerobically and can produce a significant

quantity of CH₄. The temperature and the retention time of the storage unit greatly affect the amount of methane produced. When manure is handled as a solid (e.g., in stacks or piles) or when it is deposited on pastures and rangelands, it tends to decompose under more aerobic conditions and less CH₄ is produced.

The decomposition of the organic material in cattle manures is both a level and trend key category of CH₄ emissions in Ireland.

This category also includes N₂O produced, directly and indirectly, during the storage and treatment of manure before it is applied to land or otherwise used for feed, fuel, or construction purposes. The N₂O emissions generated by manure in the system 'pasture, range, and paddock' occur directly and indirectly from the soil, and are therefore reported under the category '*N₂O Emissions from Managed Soils*'. Direct N₂O emissions occur via combined nitrification and denitrification of nitrogen contained in the manure. The emission of N₂O from manure during storage and treatment depends on the nitrogen and carbon content of manure, and on the duration of the storage and type of treatment. Indirect emissions result from volatile nitrogen losses that occur primarily in the forms of ammonia and NO_x. The fraction of excreted organic nitrogen that is mineralised to ammonia nitrogen during manure collection and storage depends primarily on time, and to a lesser degree temperature.

5.3.1.2 Methodological Issues

The Farm Facilities Survey (Hyde et al., 2008) provided detailed data on manure management practices to support the adoption of a higher tier method. The Farm Facilities Survey was conducted on a representative sample of farms, the results of which are available at both national level and for each of the three designated Nitrates Directive regions. The proportioning of Manure Management Systems (MMS) within the model is undertaken on an individual subsystem basis. The partitioning of the year into pasture and housing periods is based on expert opinion in conjunction with the results of the Farm Facilities Survey (Hyde et al., 2008) for each particular subsystem. Having derived the time spent at pasture and the time spent in housing for cattle, the Farm Facilities Survey is used to determine the partitioning of liquid and solid manures to MMS within the housing period, and the estimation of the number of animals that are out-wintered (i.e. at pasture all year round).

Information obtained from a national farm facilities survey (Hyde et al., 2008) and the work on emission factors for enteric fermentation in cattle described in section 5.2.1 above is the basis of the CH₄ emission factors for manure management. The results from Hyde et al. (2008) provide a much improved representation of manure allocation among the relevant waste management systems in the country while the excretion of organic matter by cattle is fully characterised as part of the analysis of their feed and energy requirements relating to enteric fermentation (O'Mara, 2007). The main results of the farm facilities survey pertinent to inventory calculations are outlined in Tables 3.3.D.1 and 3.3.D.2 of Annex 3.3.

The analysis of the feeding regime for cattle (O'Mara, 2007) included a full evaluation of the organic matter content of the feeds applicable to the 11 classifications that characterise the national herd (Table 5.2), which facilitates the estimation of their respective levels of organic matter excretion. The emission factors for manure management are derived using the quantified organic matter excretion as volatile solids (VS), the methane production potential (B₀) of manure, the allocation to manure management systems based on the farm facilities survey (Hyde et al., 2008) and the corresponding values of MCF (methane conversion factor) given for the cool climate ($\leq 10^{\circ}\text{C}$) zone in Table 10.17 of the 2006 IPCC guidelines. Ireland uses the value of 0.24 m³ CH₄/kg VS as outlined in Table 10A-4 of

the 2006 IPCC guidelines for B₀ for all cattle based on input from agricultural experts who advise that the methane potential of dairy cattle manures and non-dairy cattle manures in Ireland is the same, given the similarity of their grass-based feeding systems. Volatile solids values for dairy cows and non-dairy cattle are estimated using the information provided in O'Mara (2007). These values differ from the default values provided in the IPCC Good Practice Guidance due to the higher digestibility of feeds in Ireland. The default digestibility of 60 per cent is very low in comparison to the digestibility of silage (70 per cent), grazed grass (80 per cent) and concentrates (80 per cent). Grazed grass and silage make up the majority of feed intake of cattle in Ireland due grass based production systems. The emission factors for cattle are given in Table 5.4.

Table 5.4 Tier 2 CH₄ Manure Management Emission Factors for 1990, 2007- 2013

	Manure Management (kg/head/year)							
	1990	2007	2008	2009	2010	2011	2012	2013
Dairy cows	10.57	10.23	10.18	10.08	10.32	10.33	10.20	10.22
Suckler cows	6.59	6.45	6.61	6.40	6.42	6.53	6.66	6.44
Male cattle < 1 year	4.05	3.81	3.81	3.81	3.80	3.81	3.81	3.81
Male cattle 1 - 2 years	7.09	5.91	5.89	5.85	6.01	5.71	5.46	5.41
Male cattle > 2 years	2.96	1.34	1.20	1.35	1.41	1.31	1.22	1.23
Female cattle < 1 year	4.05	3.81	3.81	3.81	3.80	3.81	3.81	3.81
Female cattle 1 - 2 years	6.26	4.20	4.27	4.42	4.56	4.40	4.42	4.47
Female cattle > 2 years	0.25	0.26	0.26	0.26	0.26	0.26	0.26	0.26
Bulls for breeding	10.48	8.38	8.38	8.38	8.38	8.38	8.38	8.38
Dairy in-calf heifers	4.59	3.78	3.78	3.78	3.78	3.78	3.78	3.78
Beef in-calf heifers	5.32	4.43	4.43	4.43	4.43	4.43	4.43	4.43

Nitrogen excretion rates have been adopted in Ireland for all animal categories for which annual census data are published by the CSO. In 2011, the inventory agency reviewed the applicability of the nitrogen excretion rates used in the inventory in collaboration with the Department of Agriculture, Food and the Marine, agricultural researchers and animal nutritionists. Nitrogen excretion rates are provided in Table 3.3.E of Annex 3.3. In the case of cattle, the excretion rates are consistent with the nitrogen content of cattle feed intake as analysed in conjunction with the determination of Tier 2 CH₄ emission factors for cattle. The nitrogen excretion rates are used by the inventory agency, along with the information on the allocation of animal manures to each applicable manure management system (MMS) from the Farm Facilities Survey (Hyde et al., 2008) as the basis of CRF Table 3.B (b) and data provided in Annex 3.3.

Approximately two-thirds of animal manure nitrogen is excreted at pasture annually, reflecting the relatively short period that cattle are housed in Ireland. Animal manures excreted at pasture and the associated emissions are accounted for under N₂O emissions from managed agricultural soils (Section 5.5.1). In 2013 the bulk of cattle manures in housing were managed in pit storage systems (93.8 per cent and 76.0 per cent for dairy cattle and other cattle respectively) for eventual spreading on agricultural lands. The remainder of animal manures produced in-house are in deep bedding systems. The emission factors given by the 2006 IPCC guidelines, Table 10.21, 0.002 kg N₂O-N/N excreted for pit storage and 0.01 kg N₂O-N/N excreted for deep bedding manure management systems are used for cattle manures.

5.3.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Manure Management* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as disaggregated animal numbers are available for all years and are used in a consistent manner.

5.3.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Manure Management, Cattle*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.3.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.3.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

5.3.2 Manure Management, All Other Livestock (3.B.2-3.B.4)

5.3.2.1 Category Description

This grouping includes **sheep, swine, and other livestock**. Manure management from other livestock in Ireland consists of **deer, goats, horses, mules and asses, poultry and fur animals**.

5.3.2.2 Methodological Issues

The Tier 2 approach in the 2006 IPCC Guidelines is used for *Manure Management; Sheep, Swine, and Other Livestock* using detailed livestock characterisation data and information on manure management systems.

The allocations to manure management systems are again based on the national farm facilities survey (Hyde et al., 2008) and appropriate values of B_0 and VS are taken from the 2006 IPCC guidelines while MCFs are again as given in Table 10.17 of the 2006 IPCC guidelines. The application of the manure management emission factors for sheep, goats, swine, horses, mules, poultry and fur animals means that all CH_4 emissions from livestock are included in the national inventory.

Approximately two-thirds of animal manure nitrogen is excreted at pasture annually, reflecting the relatively short period that swine are housed in Ireland. Animal manures excreted at pasture and the associated emissions are accounted for under N_2O emissions from managed agricultural soils (Section 5.5.1). In 2013, 92.3 per cent of sheep manure is on pasture with the remainder in deep bedding system. All swine manure is in pit storage systems whereas for the other livestock categories (deer, goats, horses, mules and asses, poultry and fur animals) only 24.4 per cent of manures is on pasture. The remainder of animal manures produced in-house are in different MMS as outlined with CH_4 emission factors for manure management in Annex 3.3.

The emission factors given by the 2006 IPCC guidelines, Table 10.21, 0.002 kg N_2O -N/N excreted for pit storage and 0.01 kg N_2O -N/N excreted for deep bedding, 0.005 kg N_2O -N/N excreted for liquid system, 0.005 kg N_2O -N/N excreted for solid storage and dry lot, 0.001 kg N_2O -N/N excreted for litter manure management systems are used for all other livestock categories as presented in Annex 3.3.

5.3.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Manure Management* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as disaggregated animal numbers are available for all years and are used in a consistent manner.

5.3.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Manure Management, Other Livestock*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.3.2.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.3.2.6 Category-specific Planned Improvements

The inventory agency is in the process of investigating the availability of new data for manure management system practices in Ireland.

5.4 Emissions from Rice Cultivation (3.C)

No activities have been identified in Ireland for inclusion under this category. This category is reported as Not Occurring (NO).

5.5 Emissions from Agricultural Soils (3.D)

The IPCC Level 3 emission source categories relevant under *3.D Agricultural Soils* in 2013 are *3.D.1 Direct N₂O Emissions from Managed Soils* and *3.D.2 Indirect N₂O Emissions from Managed Soils*. Total emissions from these activities amounted to 6,113.27 kt CO₂eq in 2013.

The emissions of N₂O that result from anthropogenic N inputs or N mineralisation occur through both a direct pathway (i.e., directly from the soils to which the N is added/released), and through two indirect pathways: (i) following volatilisation of NH₃ and NO_x from managed soils and the subsequent redeposition of these gases and their products NH₄⁺ and NO₃⁻ to soils and waters; and (ii) after leaching and runoff of N, mainly as NO₃⁻, from managed soils.

5.5.1 Direct N₂O Emissions From Managed Soils (3.D.1)

5.5.1.1 Category Description

Direct N₂O Emissions from Managed Soils is a key category in Ireland. This category includes emissions from **inorganic N fertilisers, organic N fertilisers, urine and dung deposited by grazing, crop residues, cultivation of organic soils and mineralisation/immobilization associated with loss/gain of soil organic matter**.

The following N sources are included in the methodology for estimating direct N₂O emissions from managed soils:

- synthetic N fertilisers (F_{SN});
- organic N applied as fertiliser (e.g., animal manure, compost, sewage sludge, rendering waste) (F_{ON});
- urine and dung N deposited on pasture, range and paddock by grazing animals (F_{PRP});
- N in crop residues (above-ground and below-ground), including from N-fixing crops and from forages during pasture renewal (F_{CR});
- N mineralisation associated with loss of soil organic matter resulting from change of land use or management of mineral soils (F_{SOM}); and
- drainage/management of organic soils (i.e., Histosols)(F_{OS}).

5.5.1.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC Guidelines is used for *Direct N₂O Emissions from Managed Soils*.

The estimates of direct N₂O emissions from agricultural soils take into account the nitrogen inputs from all of these sources. Tillage farming in Ireland is concentrated in the south and southeast of the country while the bulk of organic soils occur in the midlands and west. The overarching equation used for estimating *Direct N₂O Emissions from Managed Soils* is equation 11.1 in Volume 4, Chapter 11 of the 2006 IPCC Guidelines, customised to Ireland's circumstances:

$$N_2O_{Direct} - N = N_2O - N_{N\ inputs} + N_2O - N_{OS} + N_2O - N_{PRP}$$

Where:

$$N_2O - N_{N\ inputs} = [(F_{SN} + F_{ON} + F_{CR} + F_{SOM}) \times EF_1]$$

$$N_2O - N_{OS} = [(F_{OS,G,Temp,NP}) \times EF_2]$$

$$N_2O - N_{PRP} = [(F_{PRP,CPP} \times EF_{3PRP,CPP}) + (F_{PRP,SO} \times EF_{3PRP,SO})]$$

Where:

F_{SN} = annual amount of synthetic fertiliser N applied to soils, kg N yr⁻¹

F_{ON} = annual amount of animal manure and sewage sludge applied to soils kg N yr⁻¹

F_{CR} = annual amount of N in crop residues returned to soils, kg N yr⁻¹

F_{SOM} = annual amount of N in mineral soils that is mineralised, in association with loss of soil C from soil organic matter as a result of changes to land use or management, kg N yr⁻¹

F_{PRP} = annual amount of urine and dung N deposited by grazing animals, kg N yr⁻¹ (Note: the subscripts CPP and SO refer to Cattle, Poultry and Pigs, and Sheep and Other animals, respectively)

F_{OS} = annual area of managed/drained organic soils, ha (Note: the subscripts G, Temp, and NP refer to Grassland, Temperate and Nutrient Poor, respectively)

EF_1 = emission factor for N₂O emissions from N inputs, kg N₂O–N (kg N input)⁻¹

EF_2 = emission factor for N₂O emissions from drained/managed organic soils, kg N₂O–N ha⁻¹ yr⁻¹

EF_{3PRP} = emission factor for N₂O emissions from urine and dung N deposited on pasture, range and paddock by grazing animals, kg N₂O–N (kg N input)⁻¹; (Note: the subscripts CPP and SO refer to Cattle, Poultry and Pigs, and Sheep and Other animals, respectively)

Emissions from **inorganic fertilisers** (F_{SN}) are estimated using total synthetic fertiliser application per year and the 2006 IPCC Guidelines default emission factor of 0.01 kg N₂O–N/N kg (EF_1). The annual statistics on nitrogen fertiliser use (F_{SN}) are obtained from the Department of Agriculture, Food and the Marine.

Organic fertilisers (F_{ON}) consist of animal manure applied to soils (F_{AM}) and sewage sludge applied to soils (F_{SEW}). Through calculations made for *Indirect N₂O emissions from Managed Soils (3.D.2)* the quantity of these fertilisers which are volatilised as NH₃ and NO_x are subtracted. Published estimates of sludge production (O'Leary et al, 1997; O'Leary and Carty, 1998; O'Leary et al, 2000; Smith et al, 2003; Smith et al, 2004; Smith et al, 2007; Monaghan et al, 2009) and the proportion applied on agricultural lands are used to estimate FS on the basis of 3 per cent nitrogen content in sewage sludge with typical dry solids content of 25 per cent (Fehily Timoney, 1985). Although the amount of sludge spreading on land is increasing, it contributed less than 1 per cent of the organic nitrogen input to agricultural soils in 2013. Table 3.3.G of Annex 3.3 shows the total quantity of nitrogen applied each year to agricultural soils through sewage sludge for the time series 1990-2013.

Emissions from **urine and dung deposited by grazing** (F_{PRP}) consist of emissions from cattle, poultry and pigs, which use the 2006 IPCC Guidelines default emission factor of 0.02 kg N₂O-N/N kg ($EF_{3PRP, CPP}$), and emissions from sheep and other livestock (horses, mules, goats and deer), which use the 2006 IPCC Guidelines default emission factor of 0.01 kg N₂O-N/N kg ($EF_{3PRP, SO}$).

Emissions from **crop residues** (F_{CR}) are estimated using equation 11.6 in Volume 4, Chapter 11 of the 2006 IPCC Guidelines, and uses annual crop production statistics provided by the CSO. The crops considered in Ireland are wheat, oats, barley, beans and peas, potatoes, turnips, sugar beet, and fodder beet. The contribution from crops in Ireland is small relative to other nitrogen sources and it fluctuates significantly in response to the production level of the relevant crops. Additional information on data used to estimate N₂O emissions from crop residues returned to soils is provided in Tables 3.3.H of Annex 3.3.

Emissions from **mineralisation/immobilization associated with loss/gain of soil organic matter** (F_{SOM}) are estimated using equation 11.8 in Volume 4, Chapter 11 of the 2006 IPCC Guidelines. The default C:N ratio of the soil organic matter of 10 is used. Tier 1 is used so a single value for all land-uses is applied.

Emissions from **drainage/management of organic soils (i.e., Histosols)** (F_{OS}) are estimated using the area of drained/managed organic soils from official national statistics and EF_2 from Table 2.5 of the 2013 IPCC Wetland Supplement for nutrient poor grasslands.

Table 5.5 Information related to Direct N₂O Emissions from Managed Soils (3.D.1)

Parameter	Emission Factor	Emission Factor Reference
EF_1	0.01kg N ₂ O-N/N kg	Table 11.1, Volume 4, Chapter 11 of the 2006 IPCC Guidelines
$EF_{3PRP, CPP}$	0.02kg N ₂ O-N/N kg	Table 11.1, Volume 4, Chapter 11 of the 2006 IPCC Guidelines
$EF_{3PRP, SO}$	0.01kg N ₂ O-N/N kg	Table 11.1, Volume 4, Chapter 11 of the 2006 IPCC Guidelines
EF_2	4.3 kg N ₂ O-N/ha	Table 11.1, Volume 4, Chapter 11 of the 2006 IPCC Guidelines & Table 2.5, 2013 IPCC Wetland Supplement

5.5.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Direct N₂O Emissions from Managed Soils* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as fertiliser use statistics are available for all years and are used in a consistent manner.

5.5.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Direct N₂O Emissions from Managed Soils*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.5.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.5.1.6 Category-specific Planned Improvements

A much more in-depth model approach is needed to take account of all the factors that determine soil emissions and to capture the inter-annual variation in the national emission rate. The inventory agency continues to engage with researchers working on N₂O emissions from soils, with a view to

adopting a methodology that systematically accounts for the influences of soil type, fertiliser type and application rates, temperature and rainfall, which are not captured by the current IPCC methodology. However, the lack of reliable data in relation to the key soil properties including bulk density and organic carbon content has delayed the application of such a methodology at national level. Other countries are in similar positions, in that they are using relatively sophisticated methods for estimating emissions from enteric fermentation and manure management, but do not have the data to allow a Tier 2 approach for estimating emissions of N₂O from soils.

5.5.2 Indirect N₂O Emissions from Managed Soils (3.D.2)

5.5.2.1 Category Description

Indirect N₂O Emissions from Managed Soils is a key category in Ireland. This category includes emissions from **atmospheric deposition** and **nitrogen leaching and run-off** from two indirect pathways: (i) following volatilisation of NH₃ and NO_x from managed soils and the subsequent redeposition of these gases and their products NH₄⁺ and NO₃⁻ to soils and waters; and (ii) after leaching and runoff of N, mainly as NO₃⁻, from managed soils.

5.5.2.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC Guidelines is used for *Indirect N₂O Emissions from Managed Soils*.

The IPCC methodology for indirect emissions is based on a simple approach that allocates emissions of N₂O due to nitrogen deposition resulting from NH₃ and NO_x emissions in agriculture and from nitrogen leaching to the country that generated the source nitrogen. The contributions from NH₃ and NO_x emission sources in other sectors, such as transport and stationary combustion, are excluded and the import of nitrogen from other countries through atmospheric transport and runoff is not considered.

Emissions from **atmospheric deposition** (N₂O_(ATD)) arise due to the volatilisation of nitrogen applied to soils in synthetic fertilisers and animal manures. The proportions of these fertilisers that are volatilised are Frac_{GASF} and Frac_{GASM} respectively. The volatilisation rates for Ireland are determined in an elaborate NH₃ inventory for agriculture (Duffy et al, 2015). It is assumed that nitrogen lost as NO_x is negligible in comparison to NH₃. Frac_{GASM} is split into Frac_{GASM1} and Frac_{GASM2} with Frac_{GASM1} referring to NH₃-N losses from animal manures in housing, storage and landspreading and Frac_{GASM2} being the proportion of nitrogen from sewage sludge applied to soils that is volatilised as NH₃. These values are presented in Table 5.7. Equation 11.9 in Volume 4, Chapter 11 of the 2006 IPCC Guidelines is used to estimate the emissions:

$$N_2O_{(ATD)} - N = [(F_{SN} \times Frac_{GASF}) + ((F_{ON} + F_{PRP}) \times Frac_{GASM1}) + (F_{SS}) \times Frac_{GASM2}] \times EF_4$$

Where:

N₂O_(ATD)-N = annual amount of N₂O-N produced from atmospheric deposition of N volatilised from managed soils, kg N₂O-N yr⁻¹

Frac_{GASF} = fraction of synthetic fertiliser N that volatilises as NH₃ and NO_x, kg N volatilised (kg of N applied)⁻¹

Frac_{GASM1} = fraction of applied organic N fertiliser materials (F_{ON}) and of urine and dung N deposited by grazing animals (F_{PRP}) that volatilises as NH₃ and NO_x, kg N volatilised (kg of N applied or deposited)⁻¹

Frac_{GASM2} = fraction of applied sewage sludge N (F_{SS}) that volatilises as NH₃ and NO_x, kg N volatilised (kg of N applied or deposited)⁻¹

EF₄ = emission factor for N₂O emissions from atmospheric deposition of N on soils and water surfaces, [0.01 kg N–N₂O (kg NH₃–N + NO_x–N volatilised)⁻¹]

Conversion of N₂O_(ATD)-N emissions to N₂O emissions for reporting purposes is performed by using the following equation:

$$N_2O_{(ATD)} = N_2O_{(ATD)} - N \times 44/28$$

Emissions from **leaching and run-off** are estimated through equation 11.10 in Volume 4, Chapter 11 of the 2006 IPCC Guidelines:

$$N_2O_{(L)} - N = (F_{SN} + F_{ON} + F_{CR} + F_{SOM} + F_{PRP}) \times \text{Frac}_{LEACH-(H)} \times EF_5$$

Where:

N₂O_(L)-N = annual amount of N₂O–N produced from leaching and runoff of N additions to managed soils in regions where leaching/runoff occurs, kg N₂O–N yr⁻¹

Frac_{LEACH-(H)} = fraction of all N added to/mineralised in managed soils in regions where leaching/runoff occurs that is lost through leaching and runoff, kg N (kg of N additions)⁻¹

EF₅ = emission factor for N₂O emissions from N leaching and runoff, 0.0075 kg N₂O–N (kg N leached and runoff)⁻¹

Estimates of the nitrogen loads in Irish rivers reported under the OSPAR Convention (NEUT, 1999) suggest that approximately 10 per cent of all applied nitrogen in Irish agriculture is lost through leaching. More recent research (Ryan et al., 2006; Del Prado et al., 2006 and Richards et al., 2009) also suggest an average value of 10%. The value of 0.1 is thus considered to be a more realistic estimate of Frac_{LEACH-(H)} for Irish conditions than the default value of 0.3 and it is used in this submission.

Table 5.6 Information related to Indirect N₂O Emissions from Managed Soils (3.D.2)

Parameter	Emission Factor	Emission Factor Reference
Frac _{GASF}	0.019	Calculated
Frac _{GASM1}	0.078	Calculated
Frac _{GASM2}	0.2	Table 11.3, Volume 4, Chapter 11 of the 2006 IPCC Guidelines
Frac _{LEACH-(H)}	0.1	OSPAR Convention (NEUT, 1999); Ryan et al., 2006; Del Prado et al., 2006 and Richards et al., 2009

5.5.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Indirect N₂O Emissions from Managed Soils* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent.

5.5.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Indirect N₂O Emissions from Managed Soils*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.5.2.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.5.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

5.6 Emissions from Prescribed Burning of Savannas (3.E)

No activities have been identified in Ireland for inclusion under this category. This category is reported as Not Occurring (NO).

5.7 Emissions from Field Burning of Agricultural Residues (3.F)

The practice of field burning of agricultural residues does not occur in Ireland. This is as a result of requirements imposed on farmers who are in receipt of payments under the Common Agricultural Policy and national environmental schemes². This category is reported as Not Occurring (NO).

5.8 Emissions from Liming (3.G)

5.8.1 Category Description

Liming is used to reduce soil acidity and improve plant growth in managed systems, and is applied to **cropland** and **grassland** in Ireland. *Liming* is a key category in Ireland. In Ireland, emissions from liming only occur from **Limestone CaCO_3** , with no activities identified for **Dolomite $\text{CaMg}(\text{CO}_3)_2$** which is reported as Not Occurring (NO). Total emissions from *Liming* amounted to 515.69 kt CO₂eq in 2013.

5.8.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC Guidelines is used for *Liming*.

Annual sales of lime are used to infer the quantity applied to soils, assuming that all limed sold to farmers is applied during the same year. In Ireland, lime is applied to grassland and cropland in an approximately 82:18 ratio. The default emission factor of 0.12 is used for the proportion of carbon in lime.

5.8.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Liming* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as fertiliser use statistics are available for all years and are used in a consistent manner.

5.8.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Liming*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.8.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.8.6 Category-specific Planned Improvements

There are no planned improvements for this category.

² <http://www.agriculture.gov.ie/farmerschemespayments/crosscompliance/>

5.9 Emissions from Urea Application (3.H)

5.9.1 Category Description

Adding urea to soils during fertilisation leads to a loss of CO₂ that was fixed in the industrial production process. Total emissions from these activities amounted to 21.66 kt CO₂eq in 2013.

5.9.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC Guidelines is used for *Urea Application*. The amount of Urea based fertilisers is available from national fertiliser statistics.

The default emission factor of 0.20 is used for the proportion of carbon in the urea applied to land.

5.9.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Urea Application* are provided in Annex 2. The emission time series for agriculture 1990–2013 is consistent. Key activity data such as fertiliser use statistics are available for all years and are used in a consistent manner.

5.9.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Urea Application*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

5.9.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

5.9.6 Category-specific Planned Improvements

There are no planned improvements for this category.

5.10 Emissions from Other Carbon-Containing Fertilisers (3.I)

No activities have been identified in Ireland for inclusion under this category. This category is reported as Not Occurring (NO).

5.11 Emissions from Other Agricultural Sources (3.J)

No activities have been identified in Ireland for inclusion under this category. This category is reported as Not Occurring (NO).

Chapter 6 Land-Use, Land-Use Change and Forestry

6.1 Introduction

The source category classification for reporting on the LULUCF sector was revised by Decision 24/CP.19 to that given in Table 6.1. The six top-level categories are used to represent managed land areas and they are broadly defined to accommodate all land areas in most countries, taking into account possible differences in national classification systems. Each category is split into two sub-categories, which are, in some cases, further sub-divided to reflect national circumstances and the level of detail considered most appropriate for the estimation of relevant emissions and removals. The conversion sub-categories allow for the tracking of land to the principal fixed categories using 1990 as a base year. Hence, the two sub-categories cover lands remaining the initial land use before 1990 and lands converted to other land uses since 1990. Previous inventories for Ireland used the 20 year transition before lands converted to other land uses are reclassified into the lands remaining the land use category, unless otherwise stated. The revised approach attempts to address issues of consistency and comparability of activities reported under the convention and those reported under the Kyoto Protocol. The area-based approach is intended to make the best use of the various types of data likely to be available for the given categories of land and reduce possible overlaps and omissions in reporting for national total land areas.

The net CO₂ emissions to, or removals from, the atmosphere are to be reported with respect to overall carbon gain or loss for up to five relevant carbon pools for the defined land categories. These pools are above-ground biomass, below-ground biomass, dead organic matter (litter and dead wood) and soils. For Convention reporting above-ground biomass and below-ground biomass are reported together as living biomass and litter and deadwood are reported together as dead organic matter (DOM). The 2006 IPCC guidelines provides basic methodologies for calculating changes in carbon pools where land areas form the basic activity data and carbon stock change is determined from a number of other parameters. Various levels of land sub-division may be used to capture differences due to climate, management system, vegetation type or other factors influencing carbon exchange. As for other sectors of the inventory, the 2006 IPCC guidelines also provides higher tier methods for estimating emissions and removals, which may be used if the necessary data are available. The liming of agricultural lands, which produces CO₂ emissions, is now reported under the Agriculture sector as are direct and indirect emissions of N₂O from organic and inorganic fertiliser applications to soils. Emissions of N₂O and CH₄ in the LULUCF sector are reported for such activities as soil disturbance not reported under agriculture and drainage and rewetting of mineral and organic soils. Emissions of N₂O and CH₄ are also to be reported for biomass burning.

6.2 Overview of LULUCF Sector

6.2.1 Sector Coverage

The 2015 inventory submission includes the revisions prompted by the adoption of the 2006 IPCC guidelines and relevant sections of the 2013 Wetlands Supplement to the 2006 Guidelines. Complete coverage of the relevant gases has been achieved for the years 1990-2013 in all IPCC land categories, as indicated by Table 6.1. This chapter presents a broad description of data treatment and the methodologies used to estimate emissions and removals for the relevant land categories in the time-series 1990-2013. The estimates for 4.A Forest Land are prepared under the responsibility of

COFORD/Department of Agriculture Food and the Marine (DAFM) and submitted to the inventory agency in accordance with the memorandum of understanding (MOU) between DAFM and the Office of Climate, Licensing, Research and Resource Use (OCLRR) of the EPA (see section 1.3 of this report). All other emissions and removals estimates are prepared by a research fellow working directly to the inventory agency in OCLR. A detailed report on the work undertaken to report for the 2006 inventory submission on the LULUCF sector is available ([O'Brien, 2008](#)), with subsequent revisions to methodology reported in this report where necessary.

Table 6.1 Level 3 Source Category Coverage for Land Use, Land-Use Change and Forestry

4 Land Use Land-Use Change and Forestry	Carbon Stock Change Emissions of CO ₂			Wood products	CH ₄	N ₂ O
	Biomass	DOM	Soils			
A. Forest Land						
1. Forest Land remaining Forest Land	All	All	All, NA		All	Part, IE
2. Land converted to Forest Land	All	All	All, NA		All	Part, IE
B. Cropland						
1. Cropland remaining Cropland	All	NO	All		NA	IE
2. Land converted to Cropland	NO	NO	NO		NA	NO
C. Grassland						
1. Grassland remaining Grassland	NO	NO	All, NO*		NO	All, IE
2. Land converted to Grassland	All	All	All		All	Part, IE
D. Wetlands						
1. Wetlands remaining Wetlands	All	NO	All		NO	IE
2. Land converted to Wetlands	All	All	All		All	All
E. Settlements						
1. Settlements remaining Settlements	NO	NO	NA		NO	NA
2. Land converted to Settlements	All	All	All		All	All
F. Other Land						
1. Other Land remaining Other Land	NO	NO	NO*		NO	NO
2. Land converted to Other Land	All	All	All		All	All
G. Harvested wood products				All**		

Biomass - includes above and below ground biomass

DOM - dead organic matter (deadwood and litter)

All - all emission sources covered; NE - emissions not estimated; NO - activity not occurring; NA - not applicable (no emissions of the gas occur in the pool/source category); IE - emissions included elsewhere.

* Under the Tier 1 method, there is no carbon stock change in soil for these land categories, if there is no change in management

** HWP reported based of domestic production approach and excluding Harvest from deforested lands

The 1990-2013 inventory for LULUCF (submission 2015) follows the same general approach and methodologies as those used for the submissions from 2008 to 2013. However, in response to previous annual review reports (ARR) reports ([2010](#), [2011](#), [2013](#) and [2014](#)) there have been significant changes to all LULUCF categories to ensure transparent and consistent reporting of activities and land use transition under the Convention and under the Kyoto Protocol. In particular, in the 2013 submission, there was a major reappraisal of the transition of areas remaining in a land use category and those lands converted to other land uses. Previous CRF tables 4.A to 4.F reported land use transition based on a 20-year transition. This new approach reports all land areas converted to another land use after the 1st of January 1990, and lands not subject to land use change before the 1st January 1990. For example, forest land remaining forests (4.A) includes all forest area remaining

forest before 1990 and lands converted to forests after 1990. This now ensures consistent and transparent comparison of areas reported under Convention reporting and those elected under KP LULUCF.

In the 2015 submission, a significant revision of methodologies has been applied using the 2006 IPCC guidelines, the 2013 IPCC Wetland supplement, 2013 Revised Supplementary Methods and GPG Arising from the Kyoto protocol, particularly in relation to methane and nitrous oxide emissions from soils, harvested wood products, new CO₂ emissions factors from drained organic soils and carbon loss from extraction and use of horticultural peat.

The estimates of emissions and removals from LULUCF over the period 1990-2013 are presented in Table 6.2 for all land-use categories. The LULUCF sector is a net source of emissions in all years, with the losses of carbon dominated by the impact of drainage of organic soils in Grasslands and Wetlands, and gains in the biomass carbon increasingly evident in Forest Land.

In previous submissions, there appeared to be a balance between the removals in category 4.A *Forest Land* and the emissions from 4.C *Grassland*. However, the revised emission factors for drainage organic soils provided in the 2013 Wetland Supplement has had a significant impact on the estimation of emissions from *Grassland*. The increase in carbon stocks in living biomass in the category 4.A.1 *Forest Land remaining Forest Land* is the dominant removal that offsets CO₂ emissions.

Cropland constitutes a net sink of carbon to the atmosphere for most years, due to the management of temporary grasslands within crop rotation systems.

The Wetland land use category has increased in importance relative to previous submissions. This is due to the additional reporting of the carbon loss associated with the extraction and use of peat for horticulture consistent with the 2006 IPCC guidelines in the 2015 submission.

The 2015 submission also includes a revised assessment of emissions associated with biomass burning across all land use categories with the exception of Settlements.

Settlements and Other Land categories are comparatively less important in terms of emissions or removals. The inclusion of CH₄ and N₂O through the coverage of additional emission sources has a minor effect on total emissions from LULUCF.

6.2.2 Land Use Definitions and Land Use Change Matrices

Table 6.3 summarises the definitions and coverage of the IPCC land-use categories in the LULUCF sector as they relate to Ireland along with the data sources that are used for estimating the respective areas remaining in the categories before 1990, the areas converted to the categories since 1990 and their associated greenhouse gas emissions and removals. The IPCC *Wetlands* category has been sub divided into natural unexploited wetlands (unmanaged), and exploited peatlands, the latter being managed wetland areas that are drained for the purpose of commercial and domestic harvesting of peat for combustion or horticultural use.

Table 6.4 records the net change in land-use area among the various categories over the period 1990-2013 in the form of land-use change matrices for the individual years relative to the total national area of 7.11 million hectares, based on a combination of CORINE land-cover data, forest statistics and digital afforestation maps (see Figure 6.1 and Section 6.3.1). Annex 3.4.A.B gives a more detailed breakdown of the annual exchange of land between land use types and the cumulative change over time. The matrices of land use are intended to show the dynamism of changes in Irish land use and to identify the conversions that are most significant in terms of their

potential to contribute to either emissions or removals of greenhouse gases over the inventory time-series. As mentioned, the relationship between areas afforested since 1990 (KP Article 3.3 sub-category AR) are now fully consistent with convention area in lands converted to forest land since 1990.

6.2.2.1 Land use classification hierarchy

The flow diagram shown in Figure 6.1 illustrates how different data sources are used to derive land use categories in a hierarchal manner. Forest lands are initially derived using forest datasets and statistics. This is primarily based on Forest Information and Planning System which used 1995 as the baseline (FIPS 95), afforestation and deforestation data (see section 6.3.1 and Annex 3.4.A). The areas under forest land include open areas within forest boundaries. The submission includes biomass CSC for these areas using information obtained from the 2006 and 2012 national forest inventories and a reconstruction of historical age class distributions (see section 6.3.2) Emissions from soils are reported for all areas besides open areas within forest boundaries (e.g. forest roads, biodiversity areas not covered by trees) where no drainage occurs. Identification of land cover type converted to forest land (L-F) is based on an analysis of the EU Coordination of Information on the Environment (CORINE) land cover data set. Deforestation in identified forests areas is assessed using a combination of CORINE, National Forest Inventory data (NFI), maps and aerial photography datasets to obtain information on transitions to other land use categories (see section 6.3.1).

Other land use categories (i.e. non-forest land) are then allocated to other land uses using other data sources such as annual publication of agricultural areas from the Central Statistics Office of Ireland, the Land parcel Information System from the DAFM, or specific information from industry experts, as in the case of areas of industrial drainage of peatland for exploitation. Additional spatial databases such as the Land Parcel Information System, CORINE, and the Indicative Soils Map of Ireland (Fealy and Green, 2009), are used estimate the soil types associated with each land use. However, these data do not have sufficient resolution, spatially or temporally, to allow land use tracking for all land use categories. Table 6.3 details the data sources used to estimate land use areas and soil types typical of each land use type.




Figure 6.1 Methodologies and hierarchy of determining land use areas and transitions

See table 6.3 for a detailed outline of data sources. Other Land is derived from the land not included in the forest, cropland, wetland and settlement areas and as such is the residual land area not included on the other land categories.

6.2.3 Land use change trends

The reclassification of the transition period pre and post 1990 for all LULUCF categories has resulted in a significant change in sub-category land use change trends.

Figure 6.2 shows the presents a summary of land use change across all categories between 1990 and 2015. The expansion of the analysis of Cropland to improve estimates of the extent of temporary grasslands within crop rotation systems has significantly impacted on the estimate of conversion of Grassland to Cropland, and vice versa.

Grassland is the dominant land-use category in all years, accounting for 58.3 per cent of total area in 1990, followed by Wetland accounting for 18.3 per cent. The Forest Land covered 6.8 per cent, followed and Cropland at 9.8 per cent and Settlements at 1.4 per cent. Other Land is the residual land use at 5.5 per cent. The major land-use changes since 1990 have been the conversion of Grassland and Wetland to Forest Land. In 2013, Grassland accounted for 56.9 per cent of land area, Wetland 16.1 per cent, Forest Land 10.5 per cent, Cropland at 9.5 per cent, with Other Land 5.4 per cent and Settlement 1.7 per cent.




Figure 6.2 Overview of land use change between 1990 and 2013

Table 6.2 Emissions^a and Removals^a from Land Use Land-Use Change and Forestry 1990-2013 (kt CO₂ eq)

LULUCF	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
4A Forestland	-2,540.95	-2,186.93	-2,229.88	-1,702.02	-1,560.13	-1,432.23	-1,934.26	-3,096.93	-2,809.24	-2,750.74	-3,274.44	-4,662.22	-4,751.69	-3,844.74	-3,749.46	-3,171.42
A. Forest Land CO2	-2,692.40	-1,971.33	-1,908.47	-1,782.88	-1,644.59	-2,173.87	-3,327.71	-3,033.06	-2,978.25	-3,505.85	-4,896.65	-4,985.99	-4,104.46	-3,994.96	-3,411.92	-3,672.98
A. Forest Land CH4	2.34	2.81	2.82	3.25	2.70	3.64	3.22	2.85	2.85	2.89	2.90	2.80	3.64	3.04	2.79	3.09
A. Forest Land N2O	0.31	0.40	0.46	0.48	0.49	0.50	0.50	0.51	0.52	0.53	0.54	0.55	0.57	0.57	0.57	0.58
4B Cropland	-1,524.42	-102.96	-539.35	153.65	1,843.21	1,156.20	1,252.43	-760.76	-1,659.64	-1,605.69	-39.21	592.84	-1,234.96	-1,240.94	1,040.79	1,457.63
B. Cropland CO2	-1,524.52	160.64	153.57	1,843.00	1,156.18	1,252.22	-760.90	-1,659.67	-1,605.81	-39.35	592.73	-1,234.99	-1,241.31	1,040.41	1,457.62	-364.38
B. Cropland CH4	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.00
B. CroplandN2O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4C Grassland	6,429.63	5,876.69	5,923.84	6,128.31	6,039.98	5,865.46	5,998.61	5,831.79	5,872.98	5,645.08	5,656.86	5,568.87	5,825.00	5,599.56	5,432.55	5,220.01
C. Grassland CO2	6,165.09	5,776.20	5,831.65	5,673.98	5,575.13	5,624.06	5,469.62	5,560.63	5,280.68	5,279.68	5,196.60	5,485.07	5,098.72	4,922.45	4,877.37	4,901.47
C. Grassland CH4	10.54	11.86	11.57	14.38	11.27	14.71	14.16	12.17	14.16	14.77	14.55	13.26	19.70	19.95	13.24	15.00
C. Grassland N2O	0.00	0.01	0.03	0.02	0.03	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.03
4D Wetlands	2,490.65	2,427.58	2,012.89	2,072.14	3,070.55	2,660.19	3,647.94	3,262.47	2,642.31	2,870.09	3,332.57	2,870.38	2,215.79	4,228.75	4,731.38	2,373.37
D. Wetlands CO2	2,291.19	2,739.64	1,894.27	2,763.85	2,532.36	3,334.34	3,019.27	2,518.92	2,662.17	3,093.44	2,662.46	2,093.19	3,763.62	4,250.57	2,270.43	2,691.59
D. Wetlands CH4	6.09	8.00	5.43	9.46	3.85	9.67	7.49	3.74	6.39	7.38	6.40	3.74	14.46	14.96	3.13	5.89
D. Wetlands N2O	0.16	0.20	0.14	0.24	0.11	0.24	0.19	0.10	0.16	0.18	0.16	0.10	0.35	0.36	0.08	0.15
4E Settlements	79.64	97.90	154.04	194.43	256.06	244.65	295.91	315.36	366.80	414.71	637.00	667.18	266.85	312.58	129.90	325.64
E. Settlements CO2	73.88	88.07	177.77	229.64	213.65	255.79	268.43	310.74	361.04	574.98	589.30	188.61	231.35	62.20	258.89	54.96
E. Settlements CH4	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE	IE
E. Settlements N2O	0.02	0.03	0.06	0.09	0.10	0.13	0.16	0.19	0.18	0.21	0.26	0.26	0.27	0.23	0.22	0.23
4F Other Land	0.62	0.96	29.52	31.91	46.11	48.50	50.89	53.29	55.68	1,487.72	46.86	70.73	62.21	62.13	62.04	61.96
F. Other Land CO2	0.55	0.85	18.12	30.35	30.77	31.20	31.62	32.04	1,450.34	9.49	19.62	11.10	11.09	11.08	11.07	11.06
F. Other Land CH4	IE	IE	IE	IE	IE	IE	NE	IE	IE	IE	NE	IE	IE	IE	IE	IE
F. Other Land N2O	0.00	0.01	0.05	0.05	0.06	0.07	0.07	0.08	0.13	0.13	0.17	0.17	0.17	0.17	0.17	0.17
4G Harvested Wood Prodcuts	-410.59	-645.05	-956.23	-1,046.83	-1,035.17	-884.59	-1,102.83	-1,012.36	-1,052.58	-747.13	-990.47	-597.45	-635.71	-800.59	-724.91	-659.28
	-410.59	-656.83	-1,046.83	-1,035.17	-884.59	-1,102.83	-1,012.36	-1,052.58	-747.13	-990.47	-597.45	-635.71	-800.59	-724.91	-659.28	-691.95
Total LULUCF kt CO₂ eq	4524.58	5468.19	4394.84	5831.59	8660.60	7658.18	8208.68	4592.86	3416.31	5314.04	5369.18	4510.34	1747.49	4316.73	6922.30	5607.91

^a positive values indicate emissions and negative values indicate removals

^b the emissions from lime application to grassland and cropland are reported in CRF Table 5(IV) rather than under Grassland in CRF Tables 5.B and 5.C, respectively. These emissions are not included in the totals for 5.C Grassland and 5.B Cropland

Table 6.3 Description of Land Use Categories

Land Use Category	Definition and Coverage	Area 1990 (ha)	Area 2013 (ha)	Percentage change 1990-2013	Sources of Information	Principal Conversions	
						To	From
Forest Land**	All public and private plantation forests. Forest land is an area of land where tree crown cover is greater than 20% of the total area occupied. It has a minimum width of 20m and a minimum area of 0.1ha and includes all trees with a potential to reach 5m in height in situ. Trees grown for fruit or horticulture are excluded, as are non-tree woody species such as furze and rhododendron. The forest area includes open areas within forest boundaries, assumed to be 15% based on NFI statistics.	481,074	748,604	+55.6%	National Forest Inventory (NFI) 2006 and 2012 FIPS (Forest Inventory and Planning System) 1995 COILLTE database Forest Service Premiums database CORINE Land Cover General Soil Map Deforestation statistics	Grassland Wetland Settlement Other land	Grassland Wetland
Cropland	Spatial location of cropland and temporary grasslands are identified from the history of parcels used for crops in the period 2000-2013. The parcels are the gross boundary of the parcels; actual utilised areas are based on the aggregate figures from the CSO annual statistics.	698,397	673,602	-3.6%	Central Statistics Office (CSO), NFI Land Parcel Information System Indicative Soil Map of Ireland	Forest land Settlement	Forest land
Grassland	Areas of improved grassland (pasture and areas used for the harvesting of hay and silage) and unimproved grassland in use (rough grazing) as recorded by CSO annual statistics. Semi-natural grassland as estimated using CORINE Land Cover.	4,144,251	4,044,647	-2.4%	CSO, CORINE Land Cover, NFI LPIS (Land Parcels information System) Indicative Soil Map for Ireland	Forest land Settlement	Forest land
Unmanaged Wetlands	Natural unexploited wetlands	1,206,258	1,065,968	-11.6%	CORINE Land Cover, NFI Indicative Soil Map for Ireland	Managed Peatland, Forest land	Forest land
Managed Wetland	Wetland areas commercially exploited for public and private extraction of peat and areas used for domestic harvesting of peat. The quantity of peat extracted for horticultural use is estimated from export trade data.	90,547	77,989	-15.1%	Bord na Mona (BNM) area statistics; NFI, Expert opinion Central Statistics Office	Wetlands, Grassland	Forest, land managed wetland
Settlements	Urban areas, roads, airports and the footprint of industrial, commercial/institutional and residential buildings	98,145	118,248	+20.5%	CORINE Land Cover; National Roads Authority (NRA) road construction statistics; CSO housing stock, house completions and other construction floor area statistics; General Soil Map , NFI		Grassland, Forest land
Other Land	Residual when all other land use area have been determined	391,841	382,727	-2.3%	CORINE, (includes, water bodies, bare rock etc.), NFI	Forest land	
Total Land	National territorial area (including inland water bodies and salt marshes and intertidal zones)	7,111,785	7,111,785		CORINE Land Cover		

Table 6.4 Summary Land Use Matrices 1990-2013 (units ha)

	Forest	Grassland	Cropland	Managed Wetland	Unmanaged Wetlands	Settlements	Other land	Total	Total Wetlands
1989	465,278	4,149,237	700,007	90,547	1,216,607	97,777	392,334	7,111,785	1,307,154
Forest	465,278	- 4,737	- 1,582	- 320	- 8,696	10	- 472		
Grassland	4,737	4,149,237	-	- 61	-	310	-		
Cropland	1,582	-	700,007	-	-	28	-		
Managed Wetland	320	61	-	90,547	- 1,653	-	-		
Unmanaged Wetland	8,696	-	-	1,653	1,216,607	-	-		
Settlements	- 10	- 310	- 28	-	-	97,777	- 20		
Other Land	472	-	-	-	-	20	392,334		
Other									
1990	481,074	4,144,251	698,397	91,820	1,206,258	98,145	391,841	7,111,785	1,298,077
Forest	481,074	- 5,736	- 1,915	- 320	- 10,594	10	- 572		
Grassland	5,736	4,144,251	-	- 61	-	282	-		
Cropland	1,915	-	698,397	-	-	25	-		
Managed Wetland	320	61	-	91,820	- 1,309	-	-		
Unmanaged Wetland	10,594	-	-	1,309	1,206,258	-	-		
Settlements	- 10	- 282	- 25	-	-	98,145	- 18		
Other Land	572	-	-	-	-	18	391,841		
Other									
1991	500,201	4,138,294	696,457	92,748	1,194,355	98,480	391,250	7,111,785	1,287,103
Forest	500,201	- 5,002	- 1,670	- 320	- 9,198	10	- 499		
Grassland	5,002	4,138,294	-	- 61	-	286	-		
Cropland	1,670	-	696,457	-	-	26	-		
Managed Wetland	320	61	-	92,748	2,369	-	-		
Unmanaged Wetland	9,198	-	-	- 2,369	1,194,355	-	-		
Settlements	- 10	- 286	- 26	-	-	98,480	- 19		
Other Land	499	-	-	-	-	19	391,250		
Other									
1992	516,879	4,133,067	694,761	89,998	1,187,526	98,821	390,733	7,111,785	1,277,524
Forest	516,879	- 4,791	- 1,600	- 320	- 8,799	10	- 478		
Grassland	4,791	4,133,067	-	- 61	-	306	-		
Cropland	1,600	-	694,761	-	-	28	-		
Managed Wetland	320	61	-	89,998	- 8,924	-	-		
Unmanaged Wetland	8,799	-	-	8,924	1,187,526	-	-		
Settlements	- 10	- 306	- 28	-	-	98,821	- 20		
Other Land	478	-	-	-	-	20	390,733		
Other									
1993	532,857	4,128,030	693,134	98,541	1,169,803	99,185	390,235	7,111,785	1,268,344
Forest	532,857	- 5,830	- 1,946	- 320	- 10,772	10	- 582		
Grassland	5,830	4,128,030	-	- 61	-	380	-		
Cropland	1,946	-	693,134	-	-	34	-		
Managed Wetland	320	61	-	98,541	3,492	-	-		
Unmanaged Wetland	10,772	-	-	- 3,492	1,169,803	-	-		
Settlements	- 10	- 380	- 34	-	-	99,185	- 25		
Other Land	582	-	-	-	-	25	390,235		
Other	552,295	4,121,882	691,154	94,668	1,162,523	99,634	389,629		1,257,192
1994	552,295	4,121,882	691,154	94,668	1,162,523	99,634	389,629	7,111,785	1,257,192
Forest	552,295	- 6,846	- 2,371	- 140	- 13,375	-	- 645		
Grassland	6,846	4,121,882	-	- 30	-	392	-		
Cropland	2,371	-	691,154	-	-	35	-		
Managed Wetland	140	30	-	94,668	- 4,795	-	-		
Unmanaged Wetland	13,375	-	-	4,795	1,162,523	-	-		
Settlements	-	- 392	- 35	-	-	99,634	- 26		
Other Land	645	-	-	-	-	26	389,629		
Other									
1995	575,672	4,114,674	688,748	99,293	1,144,354	100,086	388,959	7,111,785	1,243,647
Forest	575,672	- 6,028	- 2,098	- 140	- 11,819	-	- 563		
Grassland	6,028	4,114,674	-	- 30	-	497	-		
Cropland	2,098	-	688,748	-	-	45	-		
Managed Wetland	140	30	-	99,293	2,963	-	-		
Unmanaged Wetland	11,819	-	-	- 2,963	1,144,354	-	-		
Settlements	-	- 497	- 45	-	-	100,086	- 32		
Other Land	563	-	-	-	-	32	388,959		
Other									

	Forest	Grassland	Cropland	Managed Wetland	Unmanaged Wetlands	Settlements	Other land	Total	Total Wetlands
1996	596,319	4,108,180	686,605	96,160	1,135,497	100,660	388,363	7,111,785	1,231,658
Forest	596,319	- 3,164	- 1,143	- 140	- 6,377	-	- 276		
Grassland	3,164	4,108,180	-	- 30	-	544	-		
Cropland	1,143	-	686,605	-	-	49	-		
Managed Wetland	140	30	-	96,160	- 618	-	-		
Unmanaged Wetland	6,377	-	-	618	1,135,497	-	-		
Settlements	-	- 544	- 49	-	-	100,660	- 36		
Other Land	276	-	-	-	-	36	388,363		
Other									
1997	607,420	4,104,502	685,413	96,608	1,128,502	101,289	388,052	7,111,785	1,225,110
Forest	607,420	- 3,612	- 1,293	- 140	- 7,229	-	- 321		
Grassland	3,612	4,104,502	-	- 30	-	587	-		
Cropland	1,293	-	685,413	-	-	53	-		
Managed Wetland	140	30	-	96,608	1,147	-	-		
Unmanaged Wetland	7,229	-	-	- 1,147	1,128,502	-	-		
Settlements	-	- 587	- 53	-	-	101,289	- 38		
Other Land	321	-	-	-	-	38	388,052		
Other									
1998	620,015	4,100,333	684,067	95,291	1,122,421	101,967	387,692	7,111,785	1,217,711
Forest	620,015	- 3,534	- 1,267	- 140	- 7,081	-	- 313		
Grassland	3,534	4,100,333	-	- 30	-	629	-		
Cropland	1,267	-	684,067	-	-	57	-		
Managed Wetland	140	30	-	95,291	- 342	-	-		
Unmanaged Wetland	7,081	-	-	342	1,122,421	-	-		
Settlements	-	- 629	- 57	-	-	101,967	- 41		
Other Land	313	-	-	-	-	41	387,692		
Other									
1999	632,350	4,096,200	682,744	95,462	1,114,998	102,694	387,337	7,111,785	1,210,461
Forest	632,350	- 4,308	- 1,570	- 327	- 8,449	171	- 356		
Grassland	4,308	4,096,200	-	- 75	-	517	-		
Cropland	1,570	-	682,744	-	-	46	-		
Managed Wetland	327	75	-	95,462	7,006	-	-		
Unmanaged Wetland	8,449	-	-	- 7,006	1,114,998	-	-		
Settlements	- 171	- 517	- 46	-	-	102,694	- 34		
Other Land	356	-	-	-	-	34	387,337		
Other									
2000	647,187	4,091,450	681,128	88,054	1,113,556	103,462	386,948	7,111,785	1,201,610
Forest	647,187	- 4,433	- 1,456	- 327	- 8,163	171	- 401		
Grassland	4,433	4,091,450	-	- 75	-	738	-		
Cropland	1,456	-	681,128	-	-	66	-		
Managed Wetland	327	75	-	88,054	622	-	-		
Unmanaged Wetland	8,163	-	-	- 622	1,113,556	-	-		
Settlements	- 171	- 738	- 66	-	-	103,462	- 48		
Other Land	401	-	-	-	-	48	386,948		
Other									
2001	661,795	4,086,355	679,605	87,031	1,106,015	104,486	386,499	7,111,785	1,193,046
Forest	661,795	- 4,492	- 1,330	- 327	- 7,782	171	- 437		
Grassland	4,492	4,086,355	-	- 75	-	663	-		
Cropland	1,330	-	679,605	-	-	60	-		
Managed Wetland	327	75	-	87,031	- 13,191	-	-		
Unmanaged Wetland	7,782	-	-	13,191	1,106,015	-	-		
Settlements	- 171	- 663	- 60	-	-	104,486	- 43		
Other Land	437	-	-	-	-	43	386,499		
Other									
2002	675,992	4,081,274	678,216	99,820	1,085,042	105,424	386,019	7,111,785	1,184,862
Forest	675,992	- 2,670	- 751	- 327	- 4,415	171	- 249		
Grassland	2,670	4,081,274	-	- 75	-	848	-		
Cropland	751	-	678,216	-	-	76	-		
Managed Wetland	327	75	-	99,820	2,173	-	-		
Unmanaged Wetland	4,415	-	-	- 2,173	1,085,042	-	-		
Settlements	- 171	- 848	- 76	-	-	105,424	- 55		
Other Land	249	-	-	-	-	55	386,019		
Other									
2003	684,232	4,077,830	677,389	97,246	1,082,800	106,575	385,714	7,111,785	1,180,045

	Forest	Grassland	Cropland	Managed Wetland	Unmanaged Wetlands	Settlements	Other land	Total	Total Wetlands
Forest	684,232	- 3,008	- 747	- 327	- 4,664	171	- 307		
Grassland	3,008	4,077,830	-	- 75	-	901	-		
Cropland	747	-	677,389	-	-	81	-		
Managed Wetland	327	75	-	97,246	13,869	-	-		
Unmanaged Wetland	4,664	-	-	- 13,869	1,082,800	-	-		
Settlements	- 171	- 901	- 81	-	-	106,575	- 59		
Other Land	307	-	-	-	-	59	385,714		
Other									
2004	693,114	4,073,995	676,561	82,975	1,092,004	107,788	385,348	7,111,785	1,174,979
Forest	693,114	- 3,260	- 715	- 133	- 4,946	171	- 356		
Grassland	3,260	4,073,995	-	- 88	-	1,087	-		
Cropland	715	-	676,561	-	-	98	-		
Managed Wetland	133	88	-	82,975	- 689	-	-		
Unmanaged Wetland	4,946	-	-	689	1,092,004	-	-		
Settlements	- 171	- 1,087	- 98	-	-	107,788	- 71		
Other Land	356	-	-	-	-	71	385,348		
Other									
2005	702,353	4,069,737	675,748	83,442	1,086,369	109,215	384,921	7,111,785	1,169,812
Forest	702,353	- 3,617	- 241	- 133	- 3,483	400	1,037		
Grassland	3,617	4,069,737	-	- 88	-	950	-		
Cropland	241	-	675,748	-	-	85	-		
Managed Wetland	133	88	-	83,442	1,482	-	-		
Unmanaged Wetland	3,483	-	-	- 1,482	1,086,369	-	-		
Settlements	- 400	- 950	- 85	-	-	109,215	- 62		
Other Land	- 1,037	-	-	-	-	62	384,921		
Other									
2006	708,390	4,065,259	675,422	81,739	1,084,368	110,712	385,897	7,111,785	1,166,107
Forest	708,390	- 3,229	- 215	- 133	- 2,695	1,200	- 502		
Grassland	3,229	4,065,259	-	- 88	-	642	-		
Cropland	215	-	675,422	-	-	58	-		
Managed Wetland	133	88	-	81,739	- 508	-	-		
Unmanaged Wetland	2,695	-	-	508	1,084,368	-	-		
Settlements	- 1,200	- 642	- 58	-	-	110,712	- 42		
Other Land	502	-	-	-	-	42	385,897		
Other									
2007	713,965	4,061,477	675,149	82,026	1,081,164	112,653	385,353	7,111,785	1,163,190
Forest	713,965	- 2,412	- 187	- 133	- 2,679	1,200	- 37		
Grassland	2,412	4,061,477	-	- 88	-	763	-		
Cropland	187	-	675,149	-	-	69	-		
Managed Wetland	133	88	-	82,026	100	-	-		
Unmanaged Wetland	2,679	-	-	- 100	1,081,164	-	-		
Settlements	- 1,200	- 763	- 69	-	-	112,653	- 50		
Other Land	37	-	-	-	-	50	385,353		
Other									
2008	718,214	4,058,389	674,893	81,704	1,078,586	114,735	385,265	7,111,785	1,160,289
Forest	718,214	- 2,591	- 199	-	- 2,591	-	- 466		
Grassland	2,591	4,058,389	-	- 186	-	784	-		
Cropland	199	-	674,893	-	-	71	-		
Managed Wetland	-	186	-	81,704	2,783	-	-		
Unmanaged Wetland	2,591	-	-	- 2,783	1,078,586	-	-		
Settlements	-	- 784	- 71	-	-	114,735	- 51		
Other Land	466	-	-	-	-	51	385,265		
Other									
2009	724,062	4,055,200	674,623	78,735	1,078,777	115,641	384,748	7,111,785	1,157,512
Forest	724,062	- 2,941	- 249	-	- 3,741	-	- 582		
Grassland	2,941	4,055,200	-	- 186	-	975	-		
Cropland	249	-	674,623	-	-	88	-		
Managed Wetland	-	186	-	78,735	220	-	-		
Unmanaged Wetland	3,741	-	-	- 220	1,078,777	-	-		
Settlements	-	- 975	- 88	-	-	115,641	- 64		
Other Land	582	-	-	-	-	64	384,748		
Other									
2010	731,576	4,051,469	674,286	78,329	1,075,256	116,767	384,103	7,111,785	1,153,585
Forest	731,576	- 1,794	- 200	-	- 2,594	-	- 466		
Grassland	1,794	4,051,469	-	- 186	-	221	-		
Cropland	200	-	674,286	-	-	20	-		

	Forest	Grassland	Cropland	Managed Wetland	Unmanaged Wetlands	Settlements	Other land	Total	Total Wetlands
Managed Wetland	-	186	-	78,329	- 913	-	-		
Unmanaged Wetland	2,594	-	-	913	1,075,256	-	-		
Settlements	-	- 221	- 20	-	-	116,767	- 14		
Other Land	466	-	-	-	-	14	384,103		
Other									
2011	736,629	4,049,641	674,066	79,056	1,071,749	117,022	383,623	7,111,785	1,150,806

Forest	736,629	- 2,899	- 244	-	- 3,069	800	- 440		
Grassland	2,899	4,049,641	-	- 493	-	187	-		
Cropland	244	-	674,066	-	-	17	-		
Managed Wetland	-	493	-	79,056	82	-	-		
Unmanaged Wetland	3,069	-	-	- 82	1,071,749	-	-		
Settlements	- 800	- 187	- 17	-	-	117,022	- 12		
Other Land	440	-	-	-	-	12	383,623		
Other									
2012	742,481	4,047,048	673,806	78,481	1,068,763	118,037	383,170	7,111,785	1,147,244

Forest	742,481	- 2,740	- 192	-	- 2,821	65	- 435		
Grassland	2,740	4,047,048	-	- 465	-	127	-		
Cropland	192	-	673,806	-	-	11	-		
Managed Wetland	-	465	-	78,481	27	-	-		
Unmanaged Wetland	2,821	-	-	- 27	1,068,763	-	-		
Settlements	- 65	- 127	- 11	-	-	118,037	- 8		
Other Land	435	-	-	-	-	8	383,170		
Other									
2013	748,604	4,044,647	673,602	77,989	1,065,968	118,248	382,727	7,111,785	1,143,957

6.3 Forest Land (Category 4.A)

6.3.1 Overall approach and activity data

Ireland adopts the gains and losses approach for reporting biomass carbon stock changes (CSC) using tier 3 models. The reporting of other C pools is done using a C flow modelling framework. The activity data for identification of changes in forest area is based on a combination of different approaches using the following data sources (also see section 6.2.2):

- The 1995 forest information parcel data (FIPS95);
- The grant and premiums application system (GPAS) and spatial database (iFORIS) for identification of afforested lands since 1990. Information on identification of land uses converted to forest is derived from the CORINE land cover change 1990 to 2006 data set;
- Deforestation data is derived from a combination of sources including CORINE 1990 and 2000, FIPS95, National Forest Inventory (NFI) data, felling licence information and aerial photography;
- The forest fires database;
- Stratification of forest areas into different soil strata is done using NFI information and the EPA indicative soil map (IFS map).

The activity data used to derive state variables for the modelling framework is primarily derived from the FIPS95 data, harvest statistics and the 2006 NFI. The first Irish NFI was completed in 2006 and the second NFI inventory was completed in 2012. The NFI data is the primary activity data used to provide initial state variables within different forest strata for calculation of carbon stock changes (CSC) from 2006 onwards using the CARBWARE model (section 6.3.3.1 and Annex 3.4.A.5). Estimation of CSC in the forest lands before 2006 cannot be determined using the CARBWARE model due to limited historical activity data on stand variables. Therefore, a more generalised stand model (FORCARB), based on British Forestry commission yield tables, is used to provide CSC estimates prior to 2006 (Edwards and Christy, 1981; Black et al., 2012).

Figure 6.3.1 shows a schematic overview of the activity data used by the different models and the different time series the model outputs represent. The CARBWARE model has been subject to external validation and uncertainty analysis and is considered more accurately represent CSC in Irish forests, when compared to the British forestry commission (BFC) based FORCARB model. The FORCARB model is based on static management interventions (i.e. set clearfelling at maximum mean annual increment and thinnings at a 5 year marginal thinning intensity cycle), which do not reflect management interventions in Ireland (Broad and Lynch, 2006; Black et al., 2008, 2012). In addition, it is well documented that the productivity index or yield class of the major species in Ireland, Sitka spruce, is higher than those in the UK, exceeding the highest documented BFC yield class (YC 24) table (Farrelly et al., 2011). Therefore, the CARBWARE v 5 model, which has been used for KP LULUCF reporting since 2008, is now also used for convention reporting from 2007 onward. Use of two different models for the historic and post 2006 time series does offer the potential of introducing a time series bias or inconsistency. However, this is addressed by re-scaling the historic (FORCARB) time series by interpolation against the CARBWARE model outputs as indicated in Figure 6.3.1 (also see section 6.3.4.1).




Figure 6.3.1 Activity data and models used to derive CSC for forest lands

Figure 6.3.1 shows the data sources used for different forest activities (clear boxes) are represented in relation to the time series. For example FIPS95 was collected in 1995 and is used to derive information of species and forest areas in forest land from 1990 to 2006 as indicated by the black arrows. The vertical brackets show which activity data is used by different modelling frameworks FORCARB and CARBWARE. The red open box and yellow box in figure 6.3.1 indicated interpolation and adjustment of the historic data against CARBWARE outputs to ensure a consistent representation of the entire time series.

6.3.2 Detailed description of activity data

6.3.2.1 FIPS95

A full survey of the private and state forests was completed in 1996 under the Forest Service's Forest Planning and Inventory System (FIPS 95). It provides information on areas by species as identified by remote sensing (Fogarty et al 1999). This activity data is used for the determination of forest areas, species and broad age class categories for 1995 used in the FORCARB model for forest land remaining forest category (Gallagher et al., 2004). The forest area going back to 1990 and projected forward to 2005 are derived from FIPS95 minus afforestation since 1990 (iFORIS data) and deforestation since 1990. The age class structure and yield class distribution for each year was reconstructed based on felling and replanting statistics and annual harvest data (see section 6.3). The FIPS 95 data provides no information on volume, stocking density or management of forest lands and cannot be used by the new CARBWARE model. However, it is used to provide historic CSC estimates for the period 1990 to 2006 using the FORCARB model, which are then subject to re-scaling using CARBWARE estimates (Figure 6.3.1 and section 6.3.4.1).

6.3.2.2 IFORIS

The IFORIS database is used to derive the total area of forests established before 1990 and afforestation areas of lands converted to forests since 1990. Ireland adopts combined approaches 2 and 3 as set out by the IPCC GPG (2003). Spatially explicit GIS polygons, representing all forest areas in 1995, were derived from the available FIPS 95 spatial layer. Digitised maps of afforested areas since 1990 using the Grants and Premiums Administration System (GPAS), archived in the iFORIS database (Figure 6.3.2). After attributing the species information with the unique ID from the Species Data table, the spatial and attribute data were joined in the Premiums layer, representing all afforested land since 1990. The data was quality controlled and the reasons for records not meeting the data validation criteria were recorded by the Forest Service. There were four separate stages in the data validation process, which occurred in successive iterations. The validated data were appended together and then reformatted and quality controlled. The FIPS95 afforested areas was then erased from the resulting Premiums table to produce the Forestry07 layer. These data sources are being updated for the new grant aided afforestation scheme areas. For example, the Forestry08 layer is derived from the GPAS08 data and the Forestry07 layer (Figure 6.3.2). Finally, the total forest areas and afforestation area is derived directly from the GPAS and IFORIS database after removal of areas identified as deforested (see deforestation data).




Figure 6.3.2 The process involved in deriving the total forest area and afforestation areas since 1990 using the IFORIS database

6.3.2.3 The National Forest Inventory

Ireland's first National Forest Inventory was completed in 2006 using a sampling approach, based on a randomised systematic grid sample design. The second inventory was completed in 2012. This system is also designed to track land use change trends. A pilot study in Co. Wexford showed that a grid resolution of 2 km x 2 km was required to provide the density of plots needed to achieve a national estimate of timber volume with a precision of 95 per cent at the 95 per cent confidence level. This grid resolution equates to 17,423 points nationally, each representing approximately 400 ha.

There are three stages of land-use classification undertaken in the NFI, primarily to identify forest areas according to the forest definition (see chapter 11). These stages are land-use type, land-use category and land-use class (Figure 6.3.3). They form the basis of the NFI, as the classification process dictates whether the sample points are included in the NFI or not, and also the range of attributes to be collected at the individual sample points.





Figure 6.3.3 Overview of the NFI classification system (taken from NFI, 2007a)

The 2 km x 2 km grid is overlaid on the total land base map of the Republic of Ireland to facilitate land-use type (LUT) interpretation using colour air photographs (OSI, 2005, 2011/12). The primary focus of the interpretation is to identify forest land transitions. In tandem with this, other land-use types are identified for LULUCF reporting under the Convention. The grid is permanent and this allows for the re-assessment of primary sample points at future dates to monitor forest and other land-use change (i.e. afforestation and deforestation) when the OSI produces the next range of ortho-rectified aerial photos (NFI, 2007).

Once a forest plot has been identified, field measurements are undertaken in established permanent plots. The exact location of the centre of ground survey plots is identified in the field by navigating to a six digit Irish national grid co-ordinate using both GPS and electronic compass/laser technology. The total area of the circular sample plot is 500 m² (i.e. 25.24 m in diameter). Adjustments for slope are automatically made by the laser/range-finding equipment. The concentric circle approach, comprising three concentric circles with different radii is used for tree assessment. Trees of different dimensions are mapped and described on each particular plot (Figure 6.3.4). Individual trees in the plot are mapped and treemetric data are collected and archived in a GPS format. Forest mensuration measurements are made on selected individual trees within the plot based in the position within the plot and the threshold diameter (Figure 6.3.4). This information is used to estimate plot-level parameters and to scale up to 1 ha (section 6.3.3.1.3). The permanent plot data describing single tree dimensions, deadwood and plot level information, is used to initiate the CARBWARE model.



Note: The concentric plot sampling approach used (see Fig 6.3.4) has implications for uncertainty (see validation and uncertainty sections).

Figure 6.3.4 The concentric plot design and mapping of individual trees NFI, 2007a

Soil surveys were also conducted in permanent sample plots. The soil group classification used in the NFI was a modification of the great soil groups employed in the National Soil Survey (Gardiner and Radford, 1980), with the addition of sand, making 11 great soil groups. These are brown earth, gley,

regosol, grey brown podzolic, rendzina, sand, brown podzolic, basin peat, lithosol, podzol and blanket peat. For a soil to be classified as peat, the peat depth had to be greater than 30 cm. Soil categories were aggregated into three major groups on the basis of their soil carbon characteristics, which can be used to estimate carbon stock change in soils. All mineral soils were grouped together. All organic soils with a depth greater than 30 cm were classified as peats soils. Mineral soils with an organic layer less than 30 cm were classified as mineral/peat soils.

6.3.2.4 Harvests and Deforestation

Harvest before 2006

EUROSTAT information is compiled by a contractor on behalf of the DAFM. The EUROSTAT harvest is obtained from timber mills and information from the industry (e.g. Coillte and the private sector). Harvest data from 1961 to 2013 were compiled using data national data submitted to the FAO and EUROSTAT. For the 1990-2013 time series the FAO/EUROSTAT harvested volume was used to simulate harvest in the FORECARB model. This was done by adjusting age class distributions using optimisation procedures based the prescribed rotation age, thinning intervals and total harvest volume for each species cohort (see section 6.3.3.2). The simulated harvest was validated against the official FAO/EUROSTAT data as shown in section 6.3.4 (Table 6.5.4)

Harvests between 2006 and 2012

The 2006 and 2012 NFIs were used to derive harvest data for the periods after 2005. The NFI records individual trees within PSP that are harvested and the indicative date of harvest based on:

- I. The previous DBH and height of the tree in the 2006/12 inventories;
- II. The estimate year of harvest. This is based on assessment of condition of stumps and deadwood on site;
- III. The volume at year of harvest is then estimated using the DBH and height in 2006 and 2012 and growth is interpolated between inventory years and extrapolated after the last available inventory 2012 using the CARBWARE model (see section 6.3.3.1). Models are validated when a new inventory cycle is completed;
- IV. The simulated harvest was validated against the official FAO/EUROSTAT data as shown in section 6.3.4 (Table 6.5.4).

Harvest from forest land remaining forest land (CRF 4.A.1) increased from ca. 1.5 Mm³ in 1990 to ca. 3.02 Mm³ by 2013 due to changes in the age class structure and clearfell of more crops at rotation age (Table 6.5.2, Figure 6.3.9). Harvest from lands converted to forest land (i.e. all forests established since 1990, CRF 4.A.2) only occurred from 2007 onwards due to the young age class structure of this category. All harvests occurring on afforested land since 1990 are carried out as first thinnings of more productive conifer crops. The total timber volumes harvested from the areas afforested since 1990 was 82,903 m³ in 2007 increasing to 188,070 m³ by 2013. Harvesting from the Coillte lands represented ca. 80 per cent of the total timber harvest from post-1990 forests (afforestation areas only). However, approximately 65 per cent of the afforestation area is privately owned, where thinnings are not commonly carried out because of the small fragmented nature of private forest, making it economically unviable to thin forest stands. NFI analysis suggests that 70 per cent of stands, which are suitable for thinning, are not thinned.

Harvests since 2012

To derive harvest since 2012 from Coillte (State owned) forests, the NFI sample plot co-ordinates and Coillte sub-compartment polygons were intersected to produce a layer representing NFI-Coillte plots with harvest management statistics (Figure 6.3.5). Harvested volume and basal area removed during harvest was assigned to individual NFI plots, representing 400 ha, based on Coillte Forecast plans. The total volume removed in a given year was validated against independently derived FAO/Eurostat data and Coillte sales invoice information. An 'EventsTable' table for use in the CARBWARE database was created for input into the stand modification functions within the CARBWARE model to simulate the harvesting of trees. A final validation was performed on the individual tree tables (see Figure 6.3.8) to ensure adequate timber was removed during a thinning simulation. It will be possible in the future to re-evaluate the activity data 'ground truthed data' from repeat NFI inventories of harvested plots, where adjustment can be made to the harvest volumes based on new PSP information.




Figure 6.3.5 Methodology used to derive harvest information for post-1990 State Forests

To derive harvests from private forests, a GIS layer was created by intersection of Town land boundaries and names (OSI) and the GPAS layer compartments (see Figure 6.3.6) that contain NFI plots. This layer contains attributes which identifies permanent sample plots which may be subjected to harvesting activities as supplied on felling licence application forms (Figure 6.3.6). Once this layer is updated every year the Forest Service carries out the following checks:

- i. Forest inspectors open the GIS attribute table to check if the Town land in question (as specified on felling licence application) contains a sample compartment.
- ii. If there is a sample compartment in the Town land, then an aerial photo layer is used to locate the compartment as indicated in the OS map in the hardcopy of the felling licence application.
- iii. Once the compartment is located, a shaded area within or covering the entire area should be identified once the GIS layer is switched on. The shaded area will contain a unique number which is used as a reference (name - FID number).

- iv. The inspector can then contact the contractor or owner to obtain information on area, species, volume and basal area removed due to harvest.

The scaled up total volume removed in a given year was compared against independently derived FAO/Eurostat information and adjusted if required. An 'EventsTable' table for in the CARBWARE database was created for input into the stand modification functions within the CARBWARE model to simulate the harvesting of trees. A final validation was performed on the individual tree tables to ensure adequate timber was removed during a thinning simulation. It will be possible in the future to re-evaluate the activity data 'ground truthed data' from repeat NFI inventories of harvested plots.




Figure 6.3.6 Procedure used to derive harvest activity data for private forested areas

Deforestation

Clearfelled areas, which were not restocked within 5 years or if there was clear indication of land use change, were deemed to be deforested. The following approaches are used to determine deforestation areas. (see Annex 3.4.A):

- 1) Sampling approach: NFI grid points and aerial photography (see Annex 3.4.A.2)

This is a modification of IPCC GPG approach 3, where the grids or centroids are sampled using a systematic sampling procedure adopted in the NFI. Assessment of ca.18000 NFI point intersects with aerial photographs from 2000 and 2006 provides the opportunity to report deforestation for this period. This method identified 15 NFI PSP grid samples, which were deemed to be deforested between 2000 and 2006. The current land uses of these previously deforested lands were determined from photo interpretation using the 2006 images.

Assessments of deforestation from 1995 to 2000 were based on a GIS intersection of the 18,000 NFI plots with the FIPS 95 forest parcel polygon layer. This exercise produced 105 forest parcels, which were classified as forest in the FIPS 1995 dataset, but then re-classified as non-forest land in the NFI aerial photography 2000 interpretation. These 105 polygons were cross-checked with 1995 black and white aerial photographs to verify that they were forests in 1995. However, most of the sampled forest polygons were deemed to not be deforested or were originally other land uses in 1995. This

was due to original FIPS 95 interpretation inconsistencies of photographs and mapping errors in the FIPS95 layer. Only 5 NFI sample points were identified to be deforested between 1995 and 2000. Although it is recognised that a grid based sample introduces a high level of uncertainty due to the poor resolution of detecting highly fragmented deforestation, this is the only available data set for this time series. Importantly these uncertainties should not introduce bias, because deforestation could be both over and under estimated using this approach.

The final deforestation-land use change-soils matrices for 1995-2000 and 2000-2006 were obtained by intersecting identified deforested sample points with the national soils map database (see Annex 3.4.A.2).

2) Tracking deforestation using CORINE Land cover (CLC) data sets (see Annex 3.4.A.1)

Although the reporting of LUC matrices uses CORINE, classification and resolution problems have been highlighted comparative studies across Europe (Black et al., 2009; Hazeu and de Wit 2004, Cruickshank and Tomlinson 1996). Despite the abovementioned inappropriateness of CLC for reporting areas under LULUCF in a representative and accurate manner, this is the only data currently available to track historic deforestation prior to FIPS 95 (see method 1 above).

For this exercise we extracted CLC codes 311 (conifers), 312 (broadleaves) and 313 (mixed woodlands) to represent forest land area that were present in 1990. The transitional land cover classes were re-classified into the LULUCF land use categories to identify land uses following deforestation. The resulting polygons were then intersected with a national soils map using ARCGIS to derive a land use change and soil type matrix to the periods 1990 to 1995.

3) Modification to deforestation records from 2006-2012 using the NFI

The new NFI 2012 and previous NFI 2006 are now used to derive deforestation data for the period 2006 to 2012. This method replaced the previously used activity data from felling licence applications. The national forest inventory programme will continue to monitor whether clear felled forest land is replanted. The NFI performs land use transition analysis based on a 2 x 2 km grid using aerial photography every 5 years. The first NFI was completed in 2006 with a follow up completed at the end of 2012. A unit of land is defined as deforested land if there is a clear indication of land use change, either from limited felling licences or aerial photography and a permanent sample point, which was recorded as unplanted previously clearfelled land in the previous inventory, is still unplanted at the time of the subsequent inventory.

A QA exercise conducted in 2013 highlighted that the previously used felling licence record approach underestimated the areas and C stock of deforested land. Therefore, the new NFI data is now used to derive both the area and C stock activity data, derived directly from NFI permanent sample plots before deforestation occurred using the CARBWARE model.

The land use transitions due to deforested lands from 2006-2012 are derived from the corresponding NFI data. According to the deforestation definition, a total of 1600 ha of forests, which were clearfelled before the 2006 NFI and were not replanted by the repeat inventory in 2012, were classified and deforested other land.

4) Deforestation records from 2013 onwards (see Annex 3.4.A.3)

The Forestry Act legally requires a formal application to the Forest Service to fell trees under either a limited or a general felling license. General felling licences cover forestry activities associated with silvicultural management, such as thinnings or clearfell and replanting. Limited felling licences capture areas and volumes felled and land use transitions for all forest land converted to other land

uses. All limited felling licence applications for 2013 were considered as deforestation and record provide the basis for estimating emissions for all biomass pools at the time of harvest.

Felling stands younger than 10 years old are not subject to the forestry act felling licence application. However, records were kept because these areas were previously subject to premium payments under the afforestation scheme. Owners in receipt of these payments are obliged to notify the Forest service if these areas are taken out ('lands taken out') of the premiums payment due to deforestation. A data base of these records has been compiled to capture the land use change and soil categories. The biomass, litter and DOM losses associated with deforestation are based on the NFI, PSP average of all 10 year old forest areas.

The national forest inventory programme will continue to monitor whether clear felled forest land under general licences are replanted. The NFI performs land use transition analysis based on a 2 x 2 km grid using aerial photography every 5 years.

6.3.2.5 Activity Data for Afforestation Areas

Afforestation areas were derived from IFORIS data see Figure 6.3.2. Activity data of land afforested since 2006 is derived from the NFI 2006 and 2012.

Activity Data for Afforestation Areas after 2012

Activity data of land afforested since 2012, after the completion of the second NFI, was derived by GIS analysis of the updated Premium Layer (Figure 6.3.2), a digitised map of indicative forest soils (IFS) and intersection with NFI grid co-ordinates (Figure 6.3.7). The resulting species/soil matrix was used to derive productivity classes and individual tree height values based on CARBWARE growth models. These tables were used as inputs into the CARBWARE software to generate carbon gains and losses (see Figure 6.3.8)

The soils and land cover datasets were derived from a number of map sources, remotely-sensed and ground-truthed data. A land cover map with a minimum resolution of 1 ha was derived using aerial photography and satellite imagery (Fealy et al., 2006). The land cover mapping exercise used the known occurrence of grassland types in Ireland and their relation to soils. Thematic classes include grassland, bog and heath, rocky complexes, bare rock, forest (unenclosed) and scrub, urban land, coastal complexes, and water bodies. The land cover dataset was derived primarily from remotely sensed data, including 1995 Landsat TM satellite imagery, 1995 black and white stereo aerial photography and 2001 ETM satellite imagery.

The digital soil mapping project delivered soil and subsoil/parent material maps by extending information obtained from various surveys using a soil cover model (Fealy et al., 2006). Over 40 per cent of the dataset is a direct derivative of the National Soil Survey (Gardiner and Radford 1980) and has a minimum mapping unit of 1 ha. Subsequently, the FIPS-IFS project produced a first-approximation soil classification for those areas not previously surveyed by the National Soil Survey (NSS), using a methodology based on remote sensing and GIS. A modelling approach was then adopted to produce a projected map for Ireland using a modular system based on different soil/peat forming factors, such as sub-soils, parent material, vegetation and topography (Fealy et al., 2006 and Loftus et al., 2002). These maps were then combined to create a predictive model of soil/peat occurrence, which is represented in GIS map form.




Figure 6.3.7 Procedure to derive activity data for Afforestation Areas after 2006

Previous land use 1990-2000

Initially, the lands converted to forestry were of relatively poor quality, with marginal potential for economic returns under agricultural practices. In more recent years, and especially with the increase in private afforestation, land of higher quality has been converted to forestry, reflecting improved grant-aid under the afforestation programme, the decline in economic returns for conventional farming practices and a preference for less labour-intensive land usage. For deriving the previous land use prior to afforestation between 1990 and 2000 the CORINE 1990-2000 Land Cover Map of Ireland (level 6) was overlaid on NFI sample plots. This overlay combination delineated the individual areas and underlying soil type of afforested lands. It also revealed the plantation date and gave an indication of the previous land use. The previous land use given by CORINE was used as a general guidance.

Based on this analysis of *4.A.2.Land Converted to Forest Land* a constant proportion for land use transitions were applied, where *4.A.2.3 Wetlands Converted to Forest Land* account for 57 per cent of the total area; *4.A.2.2 Grassland Converted to Forest Land* account for 30 per cent of the total area; *4.A.2.1 Cropland Converted to Forest Land* account for 10 per cent of the total area; and *4.A.2.5 Other land Converted to Forest Land* account for 3 per cent of the total area converted to forest in any given year between 1990 and 2000. Additional disaggregation into soil types under each land use transition is also applied to enable the calculation of emissions from organic soils.

Previous land use 2006-2013

The land use prior to afforestation for 2006-2012 was derived using the 2006 and 2012 NFI data (see section 6.3.2.3 and Figure 6.3.3). Based on this analysis *4.A.2.3 Wetlands Converted to Forest Land* account for 45 per cent of the total area; *4.A.2.2 Grassland Converted to Forest Land* account for 45 per cent of the total area; *4.A.2.1 Cropland Converted to Forest Land* account for 8 per cent of the total area; and *4.A.2.5 Other land Converted to Forest Land* account for 2 per cent of the total area converted to forest. Additional disaggregation into soil types based on NFI data under each land use transition is also applied to enable the calculation of emissions from organic soils.

Previous land use 2000-2006

The percentage of previous land use between 2000 and 2006 were derived from interpolation of the 1990-2000 and 2006-2012 time series. This resulted in a constant decline in wetland conversion to forest land by 1 per cent of the total annual afforestation area, and increase in annual afforestation of grasslands by 1.3 per cent of the total area, a decrease in annual afforestation of croplands by 0.6 per cent of the total area, and a decrease in conversion of other land to forests by 0.1 per cent of the total area converted to forest in any year between 2000 and 2006.

6.3.2.6 Definition of carbon pools

Table 6.5.1 Definition of carbon pools used in LULUCF and KP-LULUCF reporting

LULUCF	KP LULUCF	Definition
Living biomass	Aboveground biomass	All biomass above stump height (1 % of tree height)
	Belowground biomass	Biomass below stump height including roots up to a diameter of 2mm
Dead organic matter	Deadwood	Standing deadwood, dead stumps, roots (min 2 mm) and logs (min 7cm diameter)
	Litter ^a	Needles, leaves and branches up to a diameter of 7cm
Mineral soil	Mineral soil	SOC of less than 20% (reported to max depth of 30cm)
Organic soil	Organic soil	SOC of > 20% and depth > 30cm
Organo-mineral soil	Organo-mineral soil	Mineral soil with a top organic soil of depth < 30cm


^aNote: For LULUCF reporting in the CRF table 4A1 and 4A2, litter pools are reported as IE under deadwood. This is because the FORECARB model use to estimate CSC for the historical time series does not differentiate between litter and deadwood pools.

6.3.3 Description of models used

6.3.3.1 CARBWARE

The CARBWARE model is used to derive net emissions/removals for all pools all forest categories since 2006. The CARBWARE system is initialised using individual tree data from the NFI and other activity data (See Figures 6.3.1 and 6.3.8). The growth and mortality model was developed using permanent sample plots established by Coillte in the 1950s. Following model parameterisation and extensive validation (see sections on uncertainty and QA/QC under each reporting), software was developed to facilitate reporting of pools using model functions and input activity data (Figure 6.3.8). The software system was developed as part of a QC initiative to reduce calculation errors when input data is formatted and processed. When the software was developed extensive testing and validation of the code functionally on different operating systems was carried out by FERS Ltd and an independent validator (PTR Ltd) under the COFORD funded CARBWARE project (2007-2012). The software is made available to the EPA and DAFM with a user manual.

The reporting system includes an on-going QA/QC system, whereby model outputs are validated against repeated NFI measurements on a 5 year rolling basis. Additional, external data checks on activity data are carried out by the data suppliers. The first repeat forest inventory on one-fifth of the forest area was completed 2011, with the remaining completed by the end of 2012.



The red box indicates the operational domain. The white boxes represent input data sources and data bases and the grey boxes indicate software modules which carry out different procedures.

Figure 6.3.8 Schematic Overview of CARBWARE Functionality

6.3.3.1.1 Pre-processing, Growth Simulator and Stand Modifier Modules

The pre-processing module carries out formatting of NFI input files in the Microsoft Access environment to ensure that individual tree and stand information can be used by the growth simulator and stand modifier module. The model itself comprises of a growth simulator (DBH increment model see Annex 3.4.A.5), a modifier module (see Annex 3.4.A.5.2), which facilitates inclusion of natural mortality and harvests and a biomass allocation module which facilitated carbon flow between different pools.

6.3.3.1.2 The Carbon Flow Sub-model

The total carbon stock changes for a given forest plot is calculated as the sum of the gains and losses in the above-ground biomass (AB), below-ground biomass (BB), Litter (Li), deadwood (DW) and soil (So) carbon pools (Equation 2.3 in Chapter 2 of the 2006 IPCC guidelines):

$$\Delta C_{lu} = \Delta C_{AB} + \Delta C_{BB} + \Delta C_{Li} + \Delta C_{Dw} + \Delta C_{So} \dots\dots\dots(6.3.1)$$

Biomass estimates include biomass for trees only, non-tree vegetation is assumed to be in steady state following canopy closure. The definition of C pools is the same for Convention and KP reporting. Below ground biomass includes all roots up to a diameter of 5cm. Litter is defined as deadwood with a diameter of less than 7cm. This includes abscised needles and leaves. The dead wood pool included all lying and standing deadwood, dead roots and stumps with a diameter greater than 7cm. organic and mineral/organic soils are reported.

a) Biomass Carbon gains and losses

Biomass carbon stock changes are calculated using a tier 3 gain and loss method, corresponding to the process-based approach given by equation 2.4 in Chapter 2 of the 2006 IPCC guidelines, which gives the net carbon stock change as the sum of carbon gains and carbon losses for each NFI PSP

$$\Delta C = \Delta C_G + \Delta C_L \dots\dots\dots(6.3.2)$$

The biomass carbon gains (ΔC_G) for both above-ground biomass (AB) and below-ground biomass (BB) are calculated for each NFI PSP using

$$\Delta C_G = GTOTAL \times CF \dots\dots\dots(6.3.3)$$

where GTOTAL is the biomass gain (t dm. ha⁻¹ yr⁻¹) in a PSP and CF is the carbon fraction of biomass dry matter, which is taken as 50 per cent for all carbon pools (Black et al., 2007). GTOTAL is derived from the sum of all living individual tree components (i.e. AB or BB) within the NFI PSP, for example:

$$GTOTAL_{AB} = AB_n - AB_{n-1} \dots\dots\dots(6.3.4)$$

where n is the year of inventory. The GTOTAL value for each NFI permanent sample plot normalised to 1 ha. The AB and BB of individual trees were calculated using biomass algorithms for different species cohorts based on national research information (Annex 3.4.A.4, Table 3.4.A.4.a), where diameter at breast height (DBH) and tree height (H) are used as dependent variables. These variables are input data in the NFI 'individual tree table' for the first NFI (2005/6, See Figure 6.3.4). The increases in DBH and H of individual trees between NFI years were simulated in the single tree growth models (See Annex 3.4.A.5.2,). The stocking (number of trees in a plot) is adjusted after every growth simulation cycle using the stand modification module (Figure 6.3.4), which removes trees based on natural mortality models and harvest activity data (Annex 3.4.A.5.2).

Biomass carbon losses from the above-ground biomass pool ($\Delta C_{L(AB)}$) were calculated based on harvest (L_{timber}), harvest residue (L_{HR}), litter fall (L_{LF}), above-ground losses due to mortality ($L_{mort(AB)}$) and fire (L_{fire}):

$$\Delta C_{L(AB)} = L_{timber} + L_{HR} + L_{LF} + L_{mort(AB)} + L_{fire} \dots\dots\dots(6.3.5)$$

L_{timber} is calculated based on the above-ground biomass removed from harvest, simulated in the stand modification module (Annex 3.4.5.2). The allocation algorithms for timber based on harvested AB, H or DBH were derived from national research information (see Annex 3.4.A.4, Table 3.4.A.4.a). L_{HR} includes the harvest residue representing all stems and branches with a DBH less than 7cm and litter left on site after timber is removed:

$$L_{HR} = AG - L_{timber} \dots\dots\dots(6.3.6)$$

L_{LF} reflects the transfer of carbon from the AB pool to the litter pool. This is calculated in the allocation module (Figure 6.3.5), based on nationally derived leaf/needle biomass (LB) and the foliage turnover rates (F_t) (Tobin et al., 2006):

$$L_{LF} = LB \times F_t \dots\dots\dots(6.3.7)$$

Allometric equations and coefficients used for the calculation of LB for different species cohorts, with either AB or DBH as dependent variables, are shown in Annex 3.4.A.4, Table 3.4.A.4.a. The F_t rate was assumed to be 6.7 years (i.e. $F_t = 0.15$) for conifer crops and 1 year for broadleaf crops (Tobin et al., 2006). The mortality of trees is based on nationally derived single tree mortality models (Annex 3.4.A.5.2.1).

The above-ground biomass loss from mortality ($L_{mort(AB)}$) was calculated using DBH and H as dependent variables in biomass algorithms (Annex 3.4.A.4, Table 3.4.A.4.a). The AB carbon losses associated with fires (L_{fire}) was determined as described in section 6.3.2. These losses are estimated in respect of total biomass burned and reported under a separate forest category in CRF Table (V). The above-ground biomass gains in previously burned forest areas are assumed to be zero.

The biomass pools allocated to $L_{HR} + L_{LF} + L_{mort(AB)}$ pools are transferred to the litter and deadwood pools. Timber biomass harvested (L_{timber}) is assumed to be immediately oxidised in the year of harvest.

Biomass carbon losses from the below-ground biomass pool ($\Delta C_{L(BB)}$) were calculated as the sum of losses due to death of roots after harvest (L_{HRroot}), natural mortality of roots ($L_{mort(BB)}$) and root death following fire (L_{fire}):

$$\Delta C_{L(BB)} = L_{HRroot} + L_{mort(BB)} + L_{fire} \dots\dots\dots(6.3.8)$$

L_{HRroot} is the root biomass transferred to the deadwood pool following harvest as is $L_{mort(BB)}$ following tree death. All roots are assumed to die and decompose following harvest. The mortality of roots is assumed to follow that for trees, as estimated from nationally derived single tree mortality models (Annex K.2). The below-ground biomass loss from mortality ($L_{mort(BB)}$) was calculated using above-ground and total biomass algorithms (Annex 3.4.A.4, Table 3.4.A.4.a). The BB biomass losses associated with fires (L_{fire}) was determined in the same way as described above for AB losses due to fires and reported in Table 4(V). The below-ground biomass gains in burned forest are assumed to be zero.

Carbon stock changes associated with deforestation reported in CRF and KP tables include those for the total standing biomass of all trees, including roots, removed at clear fell (i.e. all biomass carbon is assumed to be immediately oxidised). Since activity data and methods used to derive deforestation estimates are now based on NFI measurements, AB, BB, litter and deadwood C losses are directly estimated using allometric equations and C flow models in CARBWARE (Annex 3.4.A.4, Table 3.4.A.4.a and sections below).

Where deforestation occurred after 2012 (i.e. harvested after the last NFI in 2012), there was no plot data for estimating C stock before harvest. Therefore, the carbon stock losses in the AB and BB pools for deforestation were calculated from an estimation of standing volume (V) of these NFI plots, as specified in the Coillte inventory, a basic density (D) in the range 0.35 to 0.55 (depending on tree species), a biomass expansion factor (BEF, total biomass to timber biomass see section 7) of 1.68 to 4 t/t⁻¹ (Black et al., 2004), a carbon fraction (CF) of 0.5 and a root to shoot ratio R of 0.2 (Black et al., 2009), as follows

$$TOTAL_{(AB)} = (V \times D \times BEF \times CF) \times 1/(1 - R) \dots\dots\dots(6.3.9)$$

$$TOTAL_{(BB)} = TOTAL_{(AB)} \times R \dots\dots\dots(6.3.10)$$

The equations are similar to those used in the IPCC GPG (2006) eq. 2.8 PG 2.12 Chapter 2. However, the term (1-R) is included for above ground biomass because BEF is defined as the ratio of total biomass (including roots) to timber biomass. Similarly, the term R is included in the below ground biomass calculation.

There is no activity data for deforested areas before 2006, therefore the 2006-2013 mean AB (65.9.81 t C/ha), BB (17.2 tC/Ha), litter and deadwood (16.4 tC/Ha) C stock was applied as an IEF for these deforested areas.

b) Litter Carbon Stock Change

Net litter stock change (ΔC_{Li}) was calculated based on litter inputs (gains) due to litterfall (L_{LF}), as given by equation 6.3.7, harvest residue litter input (L_{HR}) in equation 6.3.6, mortality litter inputs (M_{Li}), and losses associated with decomposition of the litter pool (L_{decomp}):

$$\Delta C_{Li} = (L_{LF} + L_{HR} + M_{Li}) - L_{decomp} \dots\dots\dots(6.3.11)$$

where M_{Li} is the input to the litter pool from natural mortality (i.e. all aboveground dead material with a diameter less than 7 cm). This is derived from the $L_{mort(AB)}$ minus the timber fraction of the new dead pool ($L_{mort(tim)}$):

$$M_{Li} = L_{mort(AB)} - L_{mort(tim)} \dots\dots\dots(6.3.12)$$

The decomposition losses of the new input litter (L_{decomp}) and existing litter pool (L_{old}) are calculated using decomposition factors of 0.14 taken from national research (Saiz et al. 2007; Black et al. 2009b):

$$L_{decomp} = 1 - \left[\sum [L_{LF}, L_{HR}, M_{Li}, L_{old}]^{-D_{Li}} \right] \dots\dots\dots (6.3.13)$$

$$L_{old} = \sum [\{ (L_{LF}, L_{HR}, M_{Li})_{n-1, n-2, n-x} \}, L_{ini}]^{-D_{Li}} \dots\dots\dots (6.3.14)$$

where, L_{ini} is the initial litter pool estimated following the completion of the first NFI in 2005 using constructed lookup stand attribute tables based on the FORECARB model. The remaining litter from the newly input litter, harvest residue and mortality pools from the previous years (n-1, n-2 etc) were accumulated following decomposition.

The accumulated litter pool was assumed to be immediately oxidised when deforestation occurs (i.e. reported as an emission in the CRF and KP tables):

$$\text{Deforested } \Delta C_{Li} = L_{old} \times -1 \dots\dots\dots(6.3.15)$$

The accumulated litter pool for these deforestation events is derived from the initial litter pool look up tables as described above.

c) Deadwood Carbon Stock Change

Net deadwood stock changes (ΔC_{DW}) were derived from carbon inputs associated with timber extraction residue (L_{tr}), timber from mortality (M_{timber}), dead roots from mortality ($L_{mort(BB)}$), roots from harvest (L_{HRroot}) and carbon loss due to decomposition of the new and previously existing deadwood pool (D_{DW}):

$$\Delta C_{DW} = (L_{tr} + M_{timber} + L_{mort(BB)} + L_{HRroot}) - D_{DW} \dots\dots\dots(6.3.16)$$

A small amount (approximately 4 per cent, Tarleton (PTR Ltd) personal communication) of harvested timber is assumed to be left on site following harvest and this is used to estimate L_{tr} :

$$L_{tr} = L_{timber} \times RF \dots\dots\dots(6.3.17)$$

The deadwood input from natural mortality (M_{timber}) is derived from allometric equations applied to the DBH and H of dead trees after mortality iterations (see Annex 3.4.A.5.2), while $L_{mort(BB)}$ and L_{HRroot} are known from the analysis for the litter pool. The decomposition losses from the new input deadwood carbon pool, existing decaying logs (DL_{old}) and decaying stumps (DS_{old}) are calculated using equation 6.3.18 based on decomposition factors of 0.095 for stumps and 0.076 for roots (Tobin et al., 2007):

$$D_{DW} = 1 - \left[\sum [L_{tr}, M_{timber}, DL_{old}]^{-D_{log} \times t} + \sum [L_{mort(BB)}, L_{HRroot}, DS_{old}]^{-D_{st}} \right] \dots\dots\dots (6.3.18)$$

The volume and decay class of logs and stumps, measured in permanent sample plots during the NFI in 2005 and 2006, are used to calculate the carbon stocks in the decaying deadwood pools DL_{old} and DS_{old} , respectively. In the case of decaying logs

$$DL_{old} = \sum_i [VL_i \times DDC_i \times CF]^{-D_{log} \times t} + \left[\sum [L_{tr}, M_{timber}]_{(n-1, n-2, \dots, n-x)}^{-D_{log}} \right] \dots\dots\dots (6.3.19)$$

where VL is the log volume of the specific decay class ($i, n=4$), DDC is the density of the specific decay class (i) and CF is the carbon fraction (0.5). The density and decay classes described by Tobin et al (2007) were used to calculate the deadwood carbon pools in the NFI permanent sample plots (NFI, 2007b). L_{tr} and M_{timber} ($n-1, n-2, \dots, x$) is the accumulated deadwood from the stand modifier functions (equation 6.3.16 and Figure 6.3.5) within the CARBWARE model for previous years (n). Similarly, decay class and volume functions were used to derive the carbon pool of decaying stumps in NFI sample plots (Tobin et al 2007, NFI, 2007b):

$$DS_{old} = \sum_j [VS_j \times DDC_j \times CF]^{-D_{st} \times t} + \left[\sum [L_{mort(BB)}, L_{HRroot}]_{(n-1, n-2, \dots, n-x)}^{-D_{st}} \right] \dots\dots\dots(6.3.20)$$

where VS is the stump volume of the specific decay class ($j, n=4$), DDC is the density of the specific decay class (i) and CF is the carbon fraction (0.5). The density and decay classes described by Tobin et al (2007) were used to calculate the deadwood carbon pools in the NFI permanent sample plots (NFI, 2007b). $L_{mort(BB)}$ and L_{HRroot} ($n-1, n-2, \dots, x$) is the accumulated deadwood from the stand modifier functions (equation 6.3.16 and Figure 6.3.5) within the CARBWARE model for previous years (n). The carbon stock of the deadwood pool in NFI plots were attributed to each permanent sample plot using a deadwood look up function in the stand attribute table of CARBWARE (Figure 6.3.4). The decomposition emissions of the old and new deadwood carbon pools was then calculated using decay constant described by Tobin et al. (2007).

The accumulated deadwood and litter pools (DS_{old} and DL_{old}) were assumed to be immediately oxidised when deforestation occurs so that

$$\text{Deforested } \Delta C_{DW} = (DL_{old} + DS_{old}) \times -1 \dots\dots\dots(6.3.21)$$

The accumulated deadwood pool for these deforestation events is derived from the mean deadwood carbon pool of the forest category and age class, based on analysis of the NFI permanent sample plots.

d) Soils

Soils are classified into three major groups; mineral, peat and peaty/mineral soils. Peat soils are organic soils with a depth greater than 30 cm and peaty/mineral soils are a continuum between the peat and mineral categories. Current research information suggests that mineral soils in Ireland do not represent a source of carbon emissions, and therefore soil carbon stock changes are reported only for peats and peaty/mineral soils (see Ch 11). On site emissions from peat soils given by equation 6.3.22 is now based on new published data (Byrne and Farrell, 2005), but information on soil classification and peat depth available from the NFI is also taken into account.

$$\Delta C_{So} = \sum_i (A_i \times EF_{soil}) \dots\dots\dots(6.3.22)$$

The area (A_i) of the 0.05 ha plots with peat soils is multiplied by 20 to scale the measurement up to 1 ha. The on-site EF_{soil} is 0.58 t C/ha⁻¹.yr⁻¹ for the first 50 years following afforestation and is zero thereafter. Emissions from peaty/mineral soils are calculated in the same way (equation 6.23), but a soils depth function (SD) is applied to the emission factor to account for the smaller organic carbon pool available. If soil depth is less than 30 cm then,

$$\Delta C_{So} = \sum_j (A_j \times EF_{soil} \times SD) \dots\dots\dots(6.3.23)$$

and

$$SD = \frac{depth(cm)}{30cm} \dots\dots\dots(6.3.24)$$

Ireland uses a country specific emission factor for organic forest soils (Byrne and Farrell, 2005). This is calculated as the mean on site organic soil EF of 0.59 t C/ha/year over the first rotation (assumed to be 50 years for peatland forests). Byrne and Farrell (2005) demonstrate that organic soils are not a source following successive rotations. These EFs are based on total soil respiration measurements, which include respiratory inputs from autotrophic respiration and litter decomposition. Therefore, these EFs are considered to overestimate since autotrophic respiration is accounted for in NPP estimates (i.e. below ground biomass growth) and litter decomposition is accounted for in the litter pool. Other studies suggest that autotrophic respiration accounts for up to 40 per cent of total soils respiration (Siaz et al., 2007). There is currently no research information on the partitioning of soil respiration between heterotrophic and autotrophic processes in peatland soils. Therefore a conservative EF will be applied until new research information becomes available. While the EF rate is lower compared to the default rate of 0.68 t C/ha/year for organic soils in cold wet temperate conditions and the region specific value used in previous submissions of 4 t C/ha/year, the transition period is much longer than the previously used default periods. The accumulated default emission of 29.5 t/ha over 50 years is now more than 2 fold higher than the previously used methods (i.e. 13.6 t C/ha to Tier 1 and 14 t C/ha for previously used tier 2, (Hargreaves et al, 2003, NIR, 2011). A country specific transition period of fifty years is therefore considered appropriate to afforested areas on organic soils (See Byrne and Farrell, 2005). This EF is applied to all first rotation forests going back to 1940 assuming that 60 per cent of afforestation occurred on peat soils before 1990 (Black et al., 2009). All forest lands planted before 1940 are assumed to be second rotation crops or are older than 50 years by 1990 and organic soils emissions from these forests are deemed to be zero (Byrne and Farrell, 2005).

An additional off-site emission factor of 0.31 tC/ha/year for dissolved organic carbon (DOC) runoff from drain organic and organo-mineral soils was applied based on new guidance in the 2013

Supplement to the 2006 Guidelines for National Greenhouse Gas Inventories: Wetlands (2013 Wetland supplement, Eq 2.4 and Table 2.2, Ch2). This EF is simply multiplied by the area of drained organic and mineral forest soils. These emissions have been applied to all forest over the entire time series regardless of forest age.

6.3.3.1.3 Scaling and Aggregation of PSP into Different Reporting Categories

Tree measurements within NFI plots were systematically sampled (see Figure 6.3.4), so all trees were not measured in a plot. The sampling method, in conjunction with an assumption of homogeneous spatial distribution of diameters within a stand, informs the calculation of a sampling weight or *expansion factor* (EF) which is used to allow for the possibility that some trees on a given plot were not sampled. The expansion factor is inversely proportional to the prior probability that a given tree is included in the sample, based on the diameter class of the tree (see Figure 6.3.3). Each tree in the sample is thus replicated a number of times equal to its expansion factor. This replication is allowed for when calculating variables derived at plot level, such as density, by incorporating the expansion factor into the equations. For example, the estimated number of trees on a plot with a single sampled tree of greater than 70 mm is $(12.62/3)^2$. Figure 6.3.4 shows that trees of three diameter classes are only recorded if they are observed within a certain distance from the plot centre. The expansion factor used by the NFI assumes a random distribution for tree diameter in the plot. Because of that assumption, the weight assigned to a tree in the i th diameter class is:

$$\frac{R_3^2}{R_i^2} \dots\dots\dots (6.3.25)$$

where R_i denotes the radius of the concentric circle associated with the i th diameter class.

In practice, the expansion factor, or weight, is used to estimate plot-level features, e.g. basal area. In such calculations, the number of trees of the i th diameter class that were not included in the sample

is estimated by $\frac{R_3^2}{R_i^2} \times n_i$, where n_i is the number of trees of the i th class that are included in the sample. The expansion factor therefore defines the relationship between each included tree and the estimated number of trees of the same class that were not included (Equation 6.3.25).

$$n_{ij} \times EF_{ij} = \hat{N}_{ij} \dots\dots\dots (6.3.26)$$

where $n_{ij} \times EF_{ij}$ is the product of the expansion factor for the j th tree in the i th class, and \hat{N}_{ij} is the corresponding estimate. In the terminology of the NFI, the RHS of Equation 6.3.26 is the representative tree number. With minor and obvious changes to the equation, we can calculate other tree-level estimates, including representative basal area, and individual-tree estimates can be aggregated for the entire plot to give plot-level estimates, including representative density. For example the aboveground biomass carbon of a plot (t C/ha) $GTOTAL_{(AB)}$ of a plot is calculated as:

$$GTOTAL_{(AB)} = \frac{\sum [AB_{ij} \times EXF_{ij}] \times 20}{1000} \dots\dots\dots (6.3.27)$$

where, 20 is the factor used to scale up to 1 ha and 1000 is used to convert kilogrammes of biomass carbon to tonnes.

For convention reporting the total gains or losses for each pool and soil category is calculated as the sum of the pool scales up using representative area of PSP within respective categories. A PSP

represents 400 ha based on a 2 x 2 km grid sample. Since the NFI only detects forest areas at a 400 ha resolution the adjustment is done using the spatial GPAS data. The same adjustment is done for all other categories and KP tables.

So for example, if the area of organic soils under forest land remaining forest land is estimated to be 4.8 kha based NFI PSP (i.e. 12 plots out of 650 (representing a total of 260 kha) plots for the afforestation categories) and the total IFORIS area is 260.47 kha, then the area is readjusted as follows:

$$\text{New sub-category area (4.809 kha)} = \left(\frac{12}{650}\right) \times 260.47$$

These calculations are carried out automatically by the CARBWARE software. The calculation steps were subject to QA/QC checks during the coding of the software.

6.3.3.1.4 Datasets Used to Develop the CARBWARE Models

a) Permanent Sample Plot

The pre-processing, growth and mortality model was calibrated on data extracted from the permanent sample plot record system of Coillte Teoranta (the Irish Forestry Board state commercial forestry company). Broad and Lynch (2006b) provide details of the dataset in the context of modelling plot volume. The database consists of records of many silvicultural and thinning trials. These longitudinal trials were established from the 1950s onwards, and were initially established as replicated and blocked experimental designs (Broad and Lynch, 2006a).

b) Pre-processing functions

Raw data in the single tree tables and stand attributes are pre-processed by the CARBWARE software to provide variables used in the growth and modification models. In some cases, not all required variables, such as tree height (H) and crown ratio (CR) are measured. These missing values are estimated using functions described in Annex 3.4.A.5.1.

c) Growth models

The availability of only one NFI cycle meant that that the CARBWARE model had to be developed and adapted to estimate carbon stock changes. This has been done by using diameter increment models for all trees with a DBH greater than 5cm and H increment models for trees with DBH less than 5cm (Annex 3.4.A.5.2). The generated DBH and H values, produced after each growth iteration, were then used to derive biomass estimates for a range of different biomass functions (Annex 3.4.A.4, Table 3.4.A.4.a).

d) Stand modification functions

The NFI permanent plots structure is modified at the end of each growth cycle to simulate the losses associated with natural mortality and harvest (see Annex 3.4.A.5.2).

6.3.3.2 FORCARB

The FORCARB model is used to calculate CSC for the historic time series from 1990 to 2006. This is then adjusted (see figure 6.3.1) to ensure a time series consistency. The FORCARB model uses a similar C flow modelling approach as described for CARBWARE, but the main difference is that the growth, harvest and mortality is derived from stand level BFC yield tables as described by Black et al., 2012. The breakdown of species distributions was derived from an intersection of NFI and Coillte sub-compartments as described by Black et al., 2012. Species were grouped into cohorts and a

representative species table was selected from the BFC yield tables to derive stand variables such as DBH, stocking etc.

Table 6.5.2 Breakdown of species used in the pre-1990 and post-1990 forest categories

Cohort	Species table	Proportion
Spruce	Sitka spruce	0.593
Pine	Lodgepole pine	0.307
Larch	Japanese Larch	0.081
FGB	Sycamore, Ash, Birch	0.004
SGB	Beech	0.016

The yield class categories, silviculture and rotation age for each species within the pre-1990 and post-1990 categories for the period 1990-1999 were derived from the FIPS 95 dataset, modified from Gallagher et al, (2004, see Table 6.5.2). The matrix was modified for the period 2000-2012 using NFI and Coillte sub-compartment information as described by Black et al, 2012 (Table 6.5.3).

Table 6.5.3 Yield class, silviculture and rotation criteria selected for periods 1990-1999 and 2000-2012

Period :1990-1999 (Source FIPS 95)				
Species cohort	Yield class	Proportion of cohort	Silviculture	Rotation
Spruce	10	0.37	No thinning	MMAI
	16	0.26	No thinning	MMAI
	20	0.20	Thin	MMAI less 20%
	24	0.17	Thin	MMAI less 20%
Pines	10	1.00	Thin	MMAI
Larch	10	1.00	Thin	MMAI
FGB	6	1.00	Thin	MMAI
SBG	6	1.00	Thin	MMAI
Period :2000-2012 (Source NFI-Coillte intersect)				
Spruce	10	0.37	No thinning	MMAI
	16	0.13	No thinning	MMAI
	20	0.20	Thin	MMAI less 20%
	24	0.17	Thin	MMAI less 20%
	16	0.13	No thinning	MMAI less 30%
	10	0.30	No thinning	MMAI
Pines	10	0.80	No thinning	30% less MMAI
Larch	10	1.00	Thin	MMAI
FGB	6	1.00	Thin	MMAI
SBG	6	1.00	Thin	MMAI

MMAI is maximum mean annual increment, which determines the age of clearfell.

The FORCARB growth model describes gains and losses in biomass pools on mean tree-level allometric functions (DBH and height, see annex 3.4.A.4) and stand attributes (stocking) for representative species, according to Forestry Commission yield models (Edwards and Christy 1981, Black et al 2012). Stand attributes, such as age, mean DBH, top height, stocking and timber harvested, for five species cohorts (spruce, larch, pine, slow growing and fast growing broadleaves), were used as inputs for the calculation of cumulative stand biomass using species-specific allometric relationships (as described for CARBWARE models above). Harvest, thinning's and stocking changes

associated with mortality are specified in the static yield class tables (Edwards and Christy 1981, Black et al 2012).

A modified expo-linear growth function (Monteith 2000) was used to more accurately simulate biomass during the early years of the rotation and interpolate growth over time, since static models provide data at 5 year intervals and do not consider growth of young forest (<10 years old).

Stand biomass (St) was expressed as:

$$St = Mt \left[\frac{1 - e^{-k_s(k_t - t)}}{1 - e^{-k_s k_t}} \right] \dots\dots\dots (6.3.28)$$

where:

$$Mt = \frac{Cm}{Rm} \ln \left[1 + \frac{Co}{Cm} e^{Rmt} \right] \dots\dots\dots (6.3.29)$$

where:

Mt is Monteith’s function, Cm is maximum growth rate, Co is initial absolute growth rate and Rm is the initial relative growth rate and t is time (years). Parameters Cm , Rm , Co , k_s and k_t were fitted using the least squares optimisation method to estimated stand biomass values.

The current annual increment in above or below ground biomass for any given year was then calculated as:

$$\Delta C_b = St_{n+1} - St_n \dots\dots\dots (6.3.30)$$

The same C allocation models described in for the CARBWARE models were applied to simulate the biomass gains and losses and the transfer of C between pools. The resulting static tables with carbon gains, losses for biomass, net litter, deadwood pools and harvest volume were used to derive estimates of CSC from areas and age class distributions for reporting in categories 4.A.1 *Forest Land Remaining Forest Land* and 4.A.2 *Land Converted to Forest Land* (see section 6.3.4).

Age class distributions were derived from afforestation records for the category 4.A.2 *Land Converted to Forest Land*. For 4.A.1 *For the Forest Land Remaining Forest Land* category, age class distributions were initially derived from afforestation data before 1990 and felled/restocked areas. The age class distributions were then adjusted using optimisation procedures using the prescribed total harvest volume for each species cohort. The age class distributions were validated against data obtained age class distributions for 1998, 2006 and 2012 (see section 6.3.4).

For the time series adjustment of derived for C pools, the FORCARB model was run until 2012 and the 1990 to 2006 time series data was re-scaled using the CARBWARE 2006 to 2011 data (see section 6.3.4.1). Emissions form soils were not rescaled because this was derived directly using eq. 6.3.23 and 6.24 once areas on mineral, peaty mineral and peat soils was determined (see section 6.3.3.1.2).

6.3.4 Forest land remaining forest land (CRF 4.A.1)

Table 6.5.4 shows the net biomass, dead organic matter, soil C and to CO₂ emissions/removals for the time series 1990-2013 for forest land remaining forest land (i.e. all forest established before 1990 reported in CRF 4.A.1). For the historical time series 1990 to 2006, the adjusted FORCARB estimates reported. For the 2007 to 2013 time series, the CARBWARE model estimates are reported (Table 6.5.4).

The only significant change to the reported historical (1990-2012) C emissions/removals for this sub-category are related to new methods used to estimate DOC emissions from drained organic soils. For example in 2012, emissions from organic soils are 75.3 kt C higher for the 2015, compared to the 2014 submission.

Table 6.5.4 Time series for the forest category 4.A.1

Year	Area (kHa)		CSC (kt C)						Net CO ₂ kt	Harvests ¹ (M m ³)	
			Living biomass			DOM	Mineral Soils	Organic soils	Total	EUROSTAT	Modelled
	Total	Organic	Gain	Loss	Net	Net	Net	Net			
1990	465.26	277.88	2628.45	-1633.17	995.28	-74.46	NO	-151.44	-2821.05	1.68	1.67
1991	465.24	277.86	2657.59	-1651.63	1005.96	-79.15	NO	-151.33	-2843.46	1.77	1.76
1992	465.22	277.85	2686.00	-1849.25	836.75	-106.40	NO	-151.21	-2123.50	2.08	2.08
1993	465.20	277.84	2703.29	-1767.39	935.91	-19.44	NO	-150.98	-2806.78	2.10	2.10
1994	465.18	277.83	2714.29	-1903.31	810.98	-42.05	NO	-150.87	-2266.24	2.29	2.28
1995	464.84	277.63	2724.43	-2016.14	708.29	-61.91	NO	-150.62	-1817.76	2.38	2.32
1996	464.51	277.43	2732.25	-2088.57	643.68	-72.85	NO	-150.35	-1541.79	2.46	2.40
1997	464.18	277.23	2743.65	-1920.54	823.11	23.82	NO	-149.81	-2556.09	2.32	2.26
1998	463.84	277.03	2765.65	-2129.50	636.15	-21.63	NO	-149.57	-1704.85	2.64	2.57
1999	463.51	276.83	2777.54	-2205.00	572.54	5.54	NO	-149.11	-1572.86	2.78	2.71
2000	462.65	276.32	2766.10	-2429.68	336.43	58.36	NO	-147.97	-905.00	3.01	2.84
2001	461.79	275.81	2826.15	-2505.04	321.11	5.82	NO	-147.62	-657.44	2.84	2.67
2002	460.94	275.30	2778.41	-2582.06	196.35	-4.50	NO	-146.96	-164.59	2.91	2.74
2003	460.08	274.78	2766.18	-2508.17	258.02	60.11	NO	-145.31	-633.66	3.00	2.83
2004	459.22	274.27	2750.41	-2351.70	398.72	135.71	NO	-144.86	-1428.40	2.85	2.68
2005	458.37	273.76	2767.79	-2526.01	241.78	103.95	NO	-143.40	-741.88	2.94	2.77
2006	456.37	272.57	2768.56	-2575.94	192.62	69.42	NO	-142.01	-440.13	2.97	2.03
2007	454.77	271.61	2789.64	-2529.67	259.98	126.47	NO	-140.63	-901.33	2.90	2.48
2008	452.77	270.42	2816.55	-2229.10	587.45	74.22	NO	-138.19	-1919.42	2.46	1.97
2009	451.97	269.94	2831.88	-2320.25	511.63	99.55	NO	-137.26	-1737.71	2.46	2.14
2010	451.17	269.46	2864.00	-2668.44	195.56	149.08	NO	-135.55	-766.67	2.74	2.61
2011	449.57	268.50	2802.58	-2682.68	119.91	130.12	NO	-133.24	-428.20	2.72	2.60
2012	449.57	268.50	2749.39	-2881.78	-132.39	195.11	NO	-131.32	251.54	2.72	2.63
2013	449.53	268.49	2563.41	-2667.36	-103.95	231.26	NO	-129.78	9.07	2.83	2.83

¹ The harvest volumes show a comparison of the EUROSTAT and modelled harvest using FORCARB and the CARBWARE model. Note: the harvest volumes from 2007 onwards are calculated as total harvest (FAO/EUROSTAT) minus post-1990 forest harvests

The FORCARB model (see 6.3.3.2) was initially run to determine net emissions/removals in pools for the entire time series. Since the initial age class distribution in 1990 and changes in age class could not be determined from the FIPS 95 data, age class was modelled using a partial least squares optimisation based on total harvest volume (EUROSTAT harvest volume). The optimisation essentially adjusts the age class distribution until the least difference between EUROSTAT and modelled FORCARB harvests is obtained (i.e. the minimum RMSE is obtained after at least 100 iterations). The optimisation procedure was initially performed on the 1990 data set, followed by repeated optimisation procedures in the following years. The priority age class distribution for 1990 (blue histograms) was based on and incomplete Coillte inventory for 1986 (Black et al., 2012, Figure 6.3.9). Figure 6.3.9 also shows the posterior age-class distribution (red histograms) following harvest optimisation for the year 1990. To ensure that the derived FORCARB age-class distributions over the entire time series were realistic, validations were made against independent age class data for 1998, 2006 and 2012 data (Black et al, 2012, see Figure 6.3.9).




Figure 6.3.9 Validation of optimised age-class distributions

It can be seen from Figure 6.3.9 that both the FORCARB and published age-class distributions (Black et al., 2012) show the same trends over the time series. There is a right shift in the age-class distribution from 1990 to 1998, which suggest a transition from a younger to an older-aged forest estate. From 1998 to 2006, this trend is reversed because of a larger occurrence of clearfelling and restocking of sites. The slight reversed trends over the period 2006 to 2012 suggests and increase in mean age, which is consistent with a higher proportion of fillings coming from thinned stands (Black et al., 2012). These trends in combination with the increase harvest trends and higher emissions from harvest residues in the DOM pools over the time series appears to be the main driver of the observed decrease in removals by the pre-1990 forest category as suggested by Black et al., 2012.

6.3.4.1 Time Series Adjustment of Living Biomass and DOM Pools

To ensure that there is no bias introduced in estimates over the time series due to the use of the different models, the 1990 to 2006 FORCARB series was adjusted (Table 6.5.4) and rescaled using tier 1 2006 IPCC Guidelines time series overlap approaches:

- Living biomass gains (LB_{gain} , kt C) from the 2007 to 2012 time series for the CARBWARE and FORCARB model outputs were compared. The ratio (2.19) of the total CARBWARE and FORCARB LB_{gain} values for 2007-2012 was used to adjust the time series:

$$LB_{gain_{adj}} = LB_{gain_{ini}} \times 2.19 \dots \dots \dots (eq\ 6.3.31)$$

where, $LBgain_{adj}$ is the adjusted living biomass gain value and $LBgain_{ini}$ is initial FORCARB estimate. This method is consistent with eq 5.1 Ch5 of Time series in the 2006 GPG.

- b) The adjusted biomass losses ($LBloss$) were scaled using the ratio of living biomass gains to living biomass losses, derived for each year in the 1990-2006 time series. For example the adjustment for 1990 is:

$$LB\ loss_{adj(1990)} = LBgain_{adj(1990)} \times \frac{LB\ loss_{ini(1990)}}{LBgain_{ini(1990)}} \dots \dots \dots (eq\ 6.3.32)$$


- c) For dead organic matter (DOM), the ratio (-1.51) of the average CARBWARE to average FORCARB values for 2007-2012 was used to adjust the time series:

$$DOM_{adj} = DOM_{ini} \times -1.51 \dots \dots \dots (eq\ 6.3.33)$$

- d) There were no adjustments to the soil EF, since the FORCARB and CARBWARE estimates were identical.

Figure 6.3.10 shows the initial FORCARB estimates (blue symbols) and the time series adjustment as (red symbols) reported in the CRF table 4.A.1 and Table 6.5.4. Both time series show the same trend but the adjusted values show a higher net removal of CO₂. This is due to fundamental differences in the model input variables and the spatial scale at which the FORCARB and CARBWARE models operate. There are also known underestimated biases in the FORCARB model introduced when BFC yield tables are used. These are introduced by:

- a) Use of prescribed thinning cycles and clearfell regimes which do not occur in practice. The CARBWARE model imposes harvest when this is indicated in the NFI or felling licence records, so gives a clear indication the land owner intends to harvest a site. Also rotation ages as prescribed in the BFC are generally higher than those imposed under current management practice (Black et al., 2007; 2012);
- b) Predefined stocking rates in the FORCARB model, which is generally under estimated, when compared to the real situation as evident from NFI data and national research (Black et al., 2007). This would result in an underestimation of LBgains when the FORCARB model is run;
- c) Differences in the current annual increment when BFC yield table (as used in FORCARB) are compared to NFI (CARBWARE) and national research information;
- d) The CARBWARE model provides a more accurate assessment of increment in younger stand than the FORCARB, BFC based model;
- e) Although the average yield class of the major species, Sitka spruce is similar for both the FORCARB and CARBWARE based estimates. The median is higher for the NFI based assessment, which would also result in a higher increment when compared to the FORCARB model.



The final adjusted time series comprises of adjusted FORCARB estimates for the period 1990-2006 and CARBWARE estimated for the period 2007-2012.

Figure 6.3.10 Adjusted time series for forest category 4.A.1.

6.3.4.2 Mineral soils

The tier 1 approach is applied, which assumes that the carbon stock change (CSC) in mineral soil organic matter for category 4.A.1 Forest Land remaining Forest Land (FL-FL) is zero. Therefore, the notation key NO is therefore used for mineral soils under this land category in CRF Table 4.A.

6.3.4.3 Organic Soils

Emissions from the drainage of organic soils are reported using eq.6.3.23 and 6.3.24, described in section 6.2. Forest soils are classified as organic soils or (peats) if the peat depth is greater than 30 cm and the organic content is greater than 20 per cent. If the organic or peat layer is less than 30cm then the soils is classified as organo-mineral (or peaty-mineral) soils. For previous submissions, it was assumed that afforestation occurs on mineral and organic soils in the proportions 60 per cent and 40 per cent, respectively. The allocation to mineral, organo-mineral and organic soils is now determined separately for each year using PSP data from the 2006 and 2012 NFI, based on soil type and forest age attributes. The area of forest soils subjected to emissions/removals is obtained from a matrix of the three general soils types and the forest areas according to FIPS 07 and NFI information. The sample provides a breakdown in percentage of soil types in the FL-FL (pre-1990 forests younger than 50 years) and L-FL (post-1990 forest) areas. The total area is scaled up using the annual area in each category. The scaled up area is adjusted (i.e. reduced) to account for open areas in forest areas (ca. 10 per cent of the total area, NFI, 2007), since these are not planted or drained and emissions are assume to be zero. Forests older than 50 years old are assumed to be in steady state regardless of the soil type (see justifications in section 6.3.3.1.2, d) soils).

Table 6.5.5 Area (in kHa) and emissions from organic soils over the times series for forest land remaining forests

					Drained productive		Sites < 50 years old (Drained productive)		On-site emissions		Off-site DOC emissions		Total
Year	Total Area ¹	Mineral	Org.	Open area ²	Org. ³	Org.-mineral ³	Org. ⁴	Org.-mineral ⁴	Org. ⁵	Org.-mineral ⁵	Org. ⁶	Org.-mineral ⁶	
	(kHa)								kt C				kt CO ₂
1990	465.3	187.4	277.9	41.4	221.5	31.8	107.6	28.8	-63.5	-9.4	-68.7	-9.9	555.3
1991	465.2	187.4	277.9	41.4	221.5	31.8	107.4	28.8	-63.4	-9.4	-68.7	-9.9	554.9
1992	465.2	187.4	277.9	41.4	221.5	31.8	107.2	28.7	-63.3	-9.4	-68.7	-9.9	554.4
1993	465.2	187.4	277.8	41.5	221.5	31.8	106.9	28.6	-63.1	-9.4	-68.7	-9.9	553.6
1994	465.2	187.3	277.8	41.5	221.5	31.8	106.7	28.6	-63.0	-9.4	-68.7	-9.9	553.2
1995	464.8	187.2	277.6	41.5	221.3	31.8	106.5	28.5	-62.8	-9.3	-68.6	-9.9	552.3
1996	464.5	187.1	277.4	41.0	221.4	31.8	106.0	28.4	-62.6	-9.3	-68.6	-9.9	551.3
1997	464.2	186.9	277.2	41.1	221.2	31.8	105.3	28.3	-62.1	-9.3	-68.6	-9.9	549.3
1998	463.8	186.8	277.0	41.1	221.0	31.8	105.1	28.2	-62.0	-9.2	-68.5	-9.8	548.4
1999	463.5	186.7	276.8	41.2	220.8	31.7	104.5	28.1	-61.7	-9.2	-68.4	-9.8	546.7
2000	462.7	186.3	276.3	41.3	220.2	31.7	103.0	27.8	-60.8	-9.1	-68.3	-9.8	542.5
2001	461.8	186.0	275.8	41.5	219.7	31.6	102.8	27.8	-60.6	-9.1	-68.1	-9.8	541.3
2002	460.9	185.6	275.3	41.6	219.2	31.5	102.1	27.6	-60.2	-9.0	-68.0	-9.8	538.8
2003	460.1	185.3	274.8	41.7	218.7	31.4	99.8	27.1	-58.9	-8.9	-67.8	-9.7	532.8
2004	459.2	185.0	274.3	41.8	218.2	31.4	99.4	27.0	-58.7	-8.8	-67.6	-9.7	531.2
2005	458.4	184.6	273.8	42.0	217.7	31.3	97.5	26.6	-57.5	-8.7	-67.5	-9.7	525.8
2006	456.4	183.8	272.6	42.3	216.5	31.1	96.2	25.9	-56.8	-8.5	-67.1	-9.6	520.7
2007	454.8	183.2	271.6	42.4	215.5	31.0	94.7	25.3	-55.9	-8.3	-66.8	-9.6	515.6
2008	452.8	182.4	270.4	42.7	214.3	30.8	91.9	24.3	-54.2	-8.0	-66.4	-9.5	506.7
2009	452.0	182.0	269.9	42.9	213.9	30.7	90.8	23.8	-53.6	-7.9	-66.3	-9.5	503.3
2010	451.2	181.7	269.5	43.0	213.4	30.7	88.5	23.2	-52.2	-7.7	-66.1	-9.5	497.0
2011	449.6	181.1	268.5	43.2	212.4	30.5	85.6	22.3	-50.5	-7.4	-65.9	-9.5	488.6
2012	449.6	181.1	268.5	43.2	212.4	30.5	82.8	21.4	-48.9	-7.1	-65.9	-9.5	481.5
2013	449.5	181.0	268.5	43.6	212.2	30.5	80.6	21.1	-47.5	-7.0	-65.8	-9.5	475.9

¹ Total area includes open areas

² Open area within forest areas (roads, extraction routes, biodiversity etc).

³ Area of drained organic (org.) and organo-mineral soils based in NFI 2006 and 2012 (excluding open areas). Organic soils include all soils with a > 20% C and an organic layer greater than 30 cm (e.g. Blanket peats, fens, cutaway peats. Organo-mineral soils are mineral soils with an organic overlay of < 30cm. These include peaty podsols and peaty gleys (Source NFI).

⁴ No emissions from drained organic soils on sites older than 50 years old, (Data source NFI)

⁵ On-site emissions are calculated using Eq 6.3.23 and 6.2.24 and areas of productive and drained organic and organo-mineral soils less than 50 years old.

⁶ Off-site emissions are calculated using and EF of -0.31 tC /ha and the area of drained productive organic and organo-mineral soils using Eq 2.4 and Table 2.2, Ch2 of the 2013 Wet land supplement.

6.3.4.4 Emissions from Biomass Burning

Estimates of emissions from forest biomass burning in Ireland relate to forest wildfires. The estimates are recalculated in this submission based on new biomass, and DOM input estimates from the CARBWARE model for forest in 2012. In order to incorporate the effect of forest fires into CARBWARE, the following assumptions were made:

- 1) All fires are assumed to occur in all forest land classes under *4.A.1 Forest Land Remaining Forest Land* and *4.A.2 Land Converted to Forest Land*. However, because no geographically explicit data on fires are available to distinguish between fires occurring in these categories, these are equally distributed between the two categories based on the proportional area of these categories from 2007 onwards (Table 6.5.5). This assumption is made because there is evidence that fires generally only occur in forest at the pre-thicket stage of growth when there is enough woody biomass to act as a source for combustion by wild fires;
- 2) Emissions from the burning of forest biomass and DOM pools are calculated using tier 2 approaches. A carbon release factor of 0.4 is used for wildfires, with emission ratios for methane and nitrous oxide of 0.012 and 0.007, respectively (GPG 2003 Table 3 A 1.15). For nitrous oxide a C:N ratio of 0.01 is assumed. The overall implied emission factor for all GHGs as reported in CRF 4(V) is 290 t CO₂ eq/ha compared to an IEF of 39 t CO₂ eq/ha when the default values applied as specified in Eq 2.27, Table 2.4, 2.5 and 2.6 in Ch 2 (vol 4) of the 2006 IPCC Guidelines;
- 3) Emissions directly resulting from fire (i.e. combustion) are included for all years from 1990 (Table 6.5.6). Data on forest areas were obtained from the Forest assessment reports, reconstitution grant data for grant aided forests and the state owned forest company (Coillte);
- 4) Biomass burned per ha includes all aboveground biomass, litter and deadwood. However, no activity data exists documenting the amount of timber or biomass burned. Therefore, for the forest land remaining forest land category, the average biomass input for combustion is based on an average aboveground biomass C stock for a YC 16 crop over a standard rotation- 74.2 tC ha⁻¹, equivalent to 149,450 kg biomass d.wt ha⁻¹. The average C stock for litter and deadwood is estimated to be 14.1 tC ha⁻¹, equivalent to 28,263 kg biomass d.wt ha⁻¹. For the land converted to forest land category, the average aboveground biomass C stock of a 18 year old YC 16 crop is 45.3 tC ha⁻¹, equivalent to 90,526 kg biomass d.wt ha⁻¹. The average C stock for litter and deadwood is estimated to be 6.5 tC ha⁻¹, equivalent to 12,959 kg biomass d.wt ha⁻¹;
- 5) Emissions from soils are assumed to negligible and do not occur (NO);
- 6) The indirect effect of fires on carbon stock changes include those associated with loss of productivity of the area after fire and re-growth following re-planting, which is assumed to occur in the following year. It is assumed that changes in the area of forest remaining forest due to fire before 1995 were already captured by the FIPS 1995 data underlying the FORCARB model. Therefore, the indirect effects of fires and replanting on carbon stock changes, excluding the direct emission due to combustion, were only applied for the years from 1995 onwards. These are included in CRF Table 4.A.1 since they represent areas replanted.

6.3.4.5 Direct and indirect emissions of N₂O from organic and synthetic fertilisers

Ireland does not report separately the emissions of N₂O due to fertiliser use for *4.A Forest Land*. The amount of synthetic fertiliser used in forests is negligible compared to that used in Agriculture. N₂O emissions from fertiliser applications are based on national fertiliser use data reported under Agriculture. The notation key IE is therefore used in CRF Table 4(I).

6.3.4.6 Emissions of N₂O and CH₄ from drainage and rewetted organic soils

a) N₂O from drained organic soils

Tier 1 estimates of N₂O emissions due to the drainage of organic soils in forest lands were first reported in 2009. Nitrous oxide emission estimates for drained forest soils are now improved for this submission to meet requirements set out by the IPCC 2006 guidelines and the 2013 wetland supplement. The NFI data was used to derive a breakdown of areas for drained rich organic and poor organic soils over the time-series, based on planting year, soil type and cultivation type. Soils were categorised into mineral (soils with no organic layer), rich N organic (peaty-gleys or organo-mineral soils) and poor N organic (blanket peats and fen peats). Soils were assumed not to be drained if there was no cultivation, no drainage or if pit planting was employed during forest establishment as specified in the NFI database. Some upland previously degraded peatland sites are not drained prior to afforestation. Some of these lands were drained in the 1970s due to the arterial drainage scheme before they were afforested. In addition cutaway peats were drained before afforestation occurred. The total area subjected to drainage excluded open areas within forest areas, where no drainage occurs. The proportion of the three tier 1 soil types subjected to drainage for the time-series are determined from this soil/drainage matrix (table 6.5.7). The productive drained areas of the 2 organic soil categories was used to estimate N₂O emissions using the equation 11.1 in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (2006 IPCC Guidelines Ch 11).

Previous submissions used the default emission factors for poor organic and rich organic soils (2003 IPCC GPG for LULUCF CH 3 Appendix 3a.2; Table 3a.2.1 pp 3.275). However, the 2013 IPCC Wetland supplement recommend only one EFs for drained temperate forest (2.8 kg N₂O-N per ha per year) for both nutrient rich and nutrient poor organic soils (Wetland supplement Table 2.5). However, in the quoted literature used to derive these emission factors (Yamulki et al., 2013), these authors suggest the EF for nutritionally poor organic forest soils in Scotland is 0.7kg N₂O per ha per year. Therefore, we have adopted to use the default EF for nutrient rich organic soils (2.8 N₂O-N per ha per year) and a country specific EF of 0.7 kg N₂O per ha per year for nutrient poor organic soils, since this is more reflective of national circumstances (Table 6.5.7). The decline in N₂O emissions from organic soils in the forest land remaining forest land category since 1990 is due to a reduction in drained areas due to deforestation activities (Table 6.5.7).

Table 6.5.6 Area statistics and emission profiles over the time series 1990 to 2013 for wild fires in categories 4.A.1 and 4.A.2 and reported in table 4(V)

	F-F land (pre-1990)							F-L (post 1990)					
	Fire area	Prop area burned	Biom&DO M	CO2	CH4	N2O	ktCO2 eq	Prop area burned	Biom&DO M	CO2	CH4	N2O	ktCO2 eq
	Ha		Mg	kt						Mg		kt	
1990	389.00	1.00	69130.75	101.39	0.44	2.58E-03	113.22	NO	NO	NO	NO	NO	NO
1991	250.00	1.00	44428.51	65.16	0.28	1.66E-03	72.76	NO	NO	NO	NO	NO	NO
1992	160.70	1.00	28558.64	41.89	0.18	1.07E-03	46.77	NO	NO	NO	NO	NO	NO
1993	324.00	1.00	57579.34	84.45	0.37	2.15E-03	94.30	NO	NO	NO	NO	NO	NO
1994	372.00	1.00	66109.62	96.96	0.42	2.47E-03	108.27	NO	NO	NO	NO	NO	NO
1995	508.00	1.00	90278.72	132.41	0.58	3.37E-03	147.86	NO	NO	NO	NO	NO	NO
1996	565.00	1.00	100408.42	147.27	0.64	3.75E-03	164.45	NO	NO	NO	NO	NO	NO
1997	309.00	1.00	54913.63	80.54	0.35	2.05E-03	89.94	NO	NO	NO	NO	NO	NO
1998	163.00	1.00	28967.39	42.49	0.19	1.08E-03	47.44	NO	NO	NO	NO	NO	NO
1999	133.00	1.00	23635.96	34.67	0.15	8.82E-04	38.71	NO	NO	NO	NO	NO	NO
2000	334.00	1.00	59356.48	87.06	0.38	2.22E-03	97.21	NO	NO	NO	NO	NO	NO
2001	666.00	1.00	118357.54	173.59	0.76	4.42E-03	193.85	NO	NO	NO	NO	NO	NO
2002	153.00	1.00	27190.25	39.88	0.17	1.02E-03	44.53	NO	NO	NO	NO	NO	NO
2003	944.00	1.00	167762.04	246.05	1.07	6.26E-03	274.76	NO	NO	NO	NO	NO	NO
2004	550.00	1.00	97742.71	143.36	0.63	3.65E-03	160.08	NO	NO	NO	NO	NO	NO
2005	200.00	1.00	35542.80	52.13	0.23	1.33E-03	58.21	NO	NO	NO	NO	NO	NO
2006	200.00	1.00	35542.80	52.13	0.23	1.33E-03	58.21	NO	NO	NO	NO	NO	NO
2007	224.83	1.00	39955.44	58.60	0.26	1.49E-03	65.44	NO	NO	NO	NO	NO	NO
2008	273.55	0.64	31056.17	45.55	0.20	1.16E-03	50.86	0.36	10223.96	15.00	0.07	3.82E-04	16.74
2009	154.48	0.63	17392.65	25.51	0.11	6.49E-04	28.49	0.37	5858.42	8.59	0.04	2.19E-04	9.59
2010	1013.09	0.63	113040.74	165.79	0.72	4.22E-03	185.14	0.37	39014.69	57.22	0.25	1.46E-03	63.90
2011	375.55	0.63	42047.05	61.67	0.27	1.57E-03	68.86	0.37	14379.80	21.09	0.09	5.37E-04	23.55
2012	95.00	0.63	10636.18	15.60	0.07	3.97E-04	17.42	0.37	3637.50	5.34	0.02	1.36E-04	5.96
2013	408.3573	0.61	44268.20	64.93	0.28	1.65E-03	72.50	0.39	16480.97	24.17	0.11	6.15E-04	26.99

b) CH₄ from drained lands and ditches

Estimation of CH₄ emissions from drained organic lands and forest drain ditches are based on the same activity data used for determination of N₂O emissions with additional information on the fraction of land covered by drain ditches using Eq. 2.6 of CH 2 of the Wetland supplement 2013.

The default emission factors for EF_{CH₄land} (2.5 kg CH₄ per ha per year, Table 2.3 of the Wetland supplement 2013) and EF_{CH₄ditch} (217 kg CH₄ per ha per year, Table 2.4 of the Wetland supplement 2013) are used. The fraction of the total areas which is occupied by ditches (Frac_{Ditch}) was derived using country specific information (Forestry Scheme manual, 2003; Mulqueen et al., 1999) which specifies drain spacing's. For poor organic soils, such a blanket peats, these typically have 0.3m drains every 12m, which equates to a Frac_{Ditch} of 0.0249. This derived country specific Frac_{Ditch} for forest bogs are within the ranges reported for forest bogs and peats reported in Table 2A.1 in Annex 2A.2 of the IPCC Wetland supplement 2013. Richer organo-mineral soils, such as peaty gleys or peaty-podzols require drains every 80m, which is equivalent to a Frac_{Ditch} of 0.00375.

The decline in CH₄ emissions from organic soils in the forest land remaining forest land category since 1990 is due to a reduction in drained areas due to deforestation activities (Table 6.5.7).

c) Rewetting of organic soils

Forest soils are managed to maintain drains so that nutrient uptake and crop productivity is maintained. Therefore, forest soils are not rewetted.

6.3.4.7 N₂O emissions from mineral soils as a result of land use change of management (F_{SOM})

Emissions of N₂O from mineral soils are based on mineralisation rates due to loss of organic C from mineral soils. Based on available scientific evidence there we show that management and afforestation of mineral soils results in no significant change in soil organic carbon (SOC, See Ch 11). Therefore, emissions due to mineralisation of forest soil do not occur (NO).

Table 6.5.7 The area activity data and N₂O and CH₄ emissions from drainage of forest land remaining forest land

Year	Area (kHa)				kt N ₂ O			kt CH ₄		
	Total Area	Open area	Organic N-poor	Organic N-rich	Organic N-poor	Organic N-rich	Total N ₂ O	Drained lands	Ditches	Total CH ₄
1990	465.3	41.4	221.5	31.8	0.16	0.14	0.30	0.62	1.22	1.84
1991	465.2	41.4	221.5	31.8	0.16	0.14	0.30	0.62	1.22	1.84
1992	465.2	41.4	221.5	31.8	0.16	0.14	0.30	0.62	1.22	1.84
1993	465.2	41.5	221.5	31.8	0.16	0.14	0.30	0.62	1.22	1.84
1994	465.2	41.5	221.5	31.8	0.16	0.14	0.30	0.62	1.22	1.84
1995	464.8	41.5	221.3	31.8	0.15	0.14	0.29	0.62	1.22	1.84
1996	464.5	41.0	221.4	31.8	0.15	0.14	0.29	0.62	1.22	1.84
1997	464.2	41.1	221.2	31.8	0.15	0.14	0.29	0.62	1.22	1.84
1998	463.8	41.1	221.0	31.8	0.15	0.14	0.29	0.62	1.22	1.84
1999	463.5	41.2	220.8	31.7	0.15	0.14	0.29	0.62	1.22	1.84
2000	462.7	41.3	220.2	31.7	0.15	0.14	0.29	0.62	1.22	1.83
2001	461.8	41.5	219.7	31.6	0.15	0.14	0.29	0.61	1.21	1.83
2002	460.9	41.6	219.2	31.5	0.15	0.14	0.29	0.61	1.21	1.82
2003	460.1	41.7	218.7	31.4	0.15	0.14	0.29	0.61	1.21	1.82
2004	459.2	41.8	218.2	31.4	0.15	0.14	0.29	0.61	1.20	1.81
2005	458.4	42.0	217.7	31.3	0.15	0.14	0.29	0.61	1.20	1.81
2006	456.4	42.3	216.5	31.1	0.15	0.14	0.29	0.61	1.19	1.80
2007	454.8	42.4	215.5	31.0	0.15	0.14	0.29	0.60	1.19	1.79
2008	452.8	42.7	214.3	30.8	0.15	0.14	0.29	0.60	1.18	1.78
2009	452.0	42.9	213.9	30.7	0.15	0.14	0.28	0.60	1.18	1.78
2010	451.2	43.0	213.4	30.7	0.15	0.13	0.28	0.60	1.18	1.77
2011	449.6	43.2	212.4	30.5	0.15	0.13	0.28	0.59	1.17	1.77
2012	449.6	43.2	212.4	30.5	0.15	0.13	0.28	0.59	1.17	1.77
2013	449.5	43.6	212.2	30.5	0.15	0.13	0.28	0.59	1.17	1.76

6.3.4.8 Uncertainty Analysis for Category 4.A.1

Characterisation of uncertainties associated with individual activity and area information was obtained directly or derived from already published studies. If no estimates were available expert judgement was applied (Table 6.5.8). Some uncertainties cannot be quantified due to a lack of validation data. These include uncertainties associated with mortality models. However, mortality factors are selected where a tree has a 95 per cent probability of being dead (Annex K2_A). Other assumptions regarding the number of locations and amount of timber removed during harvest cannot be evaluated until the repeat NFI is completed in 2012/3.

The IPCC tier 1 approach is applied to estimate uncertainties for the Convention reporting and Article 3.3 activities described in this chapter using the methods for combining uncertainties given in section 6.3 of the IPCC good practice guidance for LULUCF. However, many of the input variables are auto correlated with each other, and therefore violate the basic assumption in this approach that inputs are statistically independent. For example, biomass and litter pools are derived from DBH increment models and biomass equations. However the simple tier 1 method is adopted until the capacity to develop Monte Carlo approaches is developed and reported in future submissions.

The percentage input uncertainties in the various methodological parameters used for the analysis of carbon stock change in the relevant carbon pools and for the emissions of non-CO₂ gases are listed in Table 6.5.9. The combined uncertainties of the products of the respective parameters associated with each component pool are calculated using equation 6.3.34 (equation 3.1 of the IPCC good practice guidance):

$$U_{\text{total}} = \sqrt{U_1^2 + U_2^2 + U_3^2 + U_n^2} \dots\dots\dots(6.3.34)$$

Where:

U_{total} is the combined uncertainty of the product of the input values U₁, U₂, U₃ and U_n given table 6.5.7. The calculated percentage uncertainties for pools are given in Table 6.5.8, which also indicates the associated input parameters whose uncertainties have been combined. The uncertainties in the reported carbon stock changes reported in the CRF tables are calculated in Table 6.5.15 as the sum of the uncertainties for carbon pools using equation 6.3.35 (equation 3.2 of the 2006 IPCC Guidelines):

$$U_{\text{total}} = \frac{\sqrt{(U_1 \times x_1)^2 + (U_2 \times x_2)^2 + (U_n \times x_n)^2}}{|x_1 + x_2 + x_n|} \dots\dots\dots(6.3.35)$$

Where:

U_{total} is the combined uncertainty, U₁, U₂ and U_n are the uncertainties of pool estimates (Table 6.5.9) and x₁, x₂ and x_n are the mean values for the respective pools reported in the CRF tables.

Table 6.5.8 Uncertainty estimates for individual activity and area data sets for forest land remaining forest land

Code	Component	Sub-category	% Uncertainty	Source
A	Biomass algorithms	AB and BB	12.0	Black et al 2007
B	C fraction	All biomass pools	0.9	Black et al 2007
C	DBH and H Increment	AB and BB	11.9	Black et al 2007, Black 2008, Black et al 2009 section 6.3.2
D	Area data	GPAS data	0.6	Derived from Black et al 2009a Table 2 Comparison of NFI and GPAS data
E	Litter	Li	3.1	Tobin et al, 2006
F	Deadwood	DW	22.0	Tobin et al, 2007
G	Peat soils	So	90.0	Assume same as Tier 1 (Table 2.3, 2.3.1 CH ₂ , 2006 IPCC Guidelines)
H	Fire C stocks	fire	15.0	95 % confidence interval for biomass stocks (NFI)
I	Areas burned	fire area	50.0	Expert Judgement, guess
J	re-scaling of FORECARB	LB	3.1	% sd of ration of CABWARE/FORECARB
K	re-scaling of FORECARB	DOM	28.5	% sd of ration of CABWARE/FORECARB
L	N ₂ O	N ₂ O and CH ₄ area	12.3	Conf. interval of NFI analysis
M	N ₂ O EF	N ₂ O emissions drained	119.0	Wetland supplement Table 2.5 and Yamulki et al., 2013
O	CH ₄ LAND EF	CH ₄ emissions drainage	87.2	Wetland supplement Table 2.4
P	CH ₄ DITCH EF	CH ₄ emissions drainage	126.0	Wetland supplement Table 2.3
Q	Soils DOC	So	43.5	Wetland supplement Table 2.1

Table 6.5.9 Combined uncertainty estimates for forest land remaining forest land pools

	Component	Equation in NIR	% uncertainty	Uncertainty of combined products (code)
LB net	Biomass	Eq 6.3.2	17.2	A+B+C+D+E+J
DOM	DOM	6.3.16 and 6.3.11	36.2	D+E+F+K
SO	Soils	6.3.24	100.0	D+G+Q
Fires	Fire	Section 6.3.3.4	59.4	H+I
N ₂ O	Drainage of soils	2006 IPCC Guidelines	119.6	L+M
CH ₄	Drainage of soils		153.7	L+O+P

For deriving uncertainties for code C in Table 6.5.9, CARBWARE DBH and H growth models were validated using repeated NFI permanent sample plot data taken in 2012. These represent repeat measurement of 1150 plots taken at a 3-6 year interval. Since modelling errors include NFI measurement and sampling errors, specific consideration was focused towards identifying sampling errors associated with the methodology employed by the Forest service (NFI data providers) and assessing model error (both sources of error in assessing biomass stock changes):

a) Accuracy of repeated DBH measurements

An infield validation check was used to ensure the corresponding tree was measured in the repeat inventory based on a spatial query of mapped trees. Measurement error of diameter and height was not checked infield or validated before entry into the data base. This resulted in a significant occurrence of negative increment data (5 to 12 per cent of data) was removed prior to model validation. In addition, trees with a DBH increment > 15 cm over 5 year cycle and with increment

values higher than 2 times the plot standard deviation were removed from the database. Zero increments from harvested trees were also removed from the database to ensure Wilcoxon ranked tests could be interpreted properly (see validation section). However, no further attempts were made to clean data with erroneous measurements in the remaining data.

b) Partial sampling of trees within a plot

Data on trees within three diameter classes (<12, 12-20 and >20 cm) are recorded if they are observed within a certain distance from the plot centre in three concentric plots within the 0.5 ha plot. This represents an additional sampling area and increases the probability of a lower representative sample of smaller, compared to larger trees. Performance of model calibration was assessed using root mean squared error (RMSE), accuracy (a measure of bias), precision and theoretical excess error.

Data were further stratified to investigate reasons for the large variation in growth increment prediction residuals across different species cohorts, DBH size classes, forest types and management regimes. Comparisons of model accuracy, bias and precision across different species cohorts and size classes show poor performance of the model in some cases (Table 6.4.9). Stratified cohort groups all had lower empirical excess error (Table 6.5.10), when compared to the theoretical excess error except for the SGB cohort, suggesting that the variation in the NFI model residuals is smaller than the random theatrically expected variation in the calibration dataset.

For all DBH categories, Spruce, pines, OC and SGB shows good agreement with the model with no significant difference between observed and simulated values ($P > 0.05$). In contrast, FGB and Larch showed poor agreement with the model predictions significant differences between observed and predicted values (Table 6.5.10). Larch and FBG showed a 27 per cent lower and 128 per cent higher growth rate than the model prediction, respectively.

This analysis (Table 6.5.10) and the uncertainty of biomass equations (annex 3.4.A.4) show that the largest uncertainty is associated with broadleaf cohorts. Current national research is being conducted to improve biomass gains and loss estimated for these cohorts.

Table 6.5.10 NFI external validation of CARBWARE models

Cohort	<12 cm	12-20 cm	20-30 cm	30-40 cm	>40 cm	All classes
Spruce						
Accuracy	-0.42	0.09	0.28	0.09	-0.73	0.17 (4.8%)
Precision	1.94	1.9	1.86	1.91	2.09	2.04
P-value	<0.01	0.37	0.14	0.55	0.03	0.36
E_{imp}						9.80%
N	204	1234	1092	226	48	2804
Pines						
Accuracy	-0.3	0.13	0.14	-0.59	ND	-0.21 (-9.4 %)
Precision	1.37	1.62	1.61	3.17	ND	2.25
P-value	0.037	0.23	0.52	<0.01	ND	0.29
E_{imp}						0.40%
N	56	342	379	44	6	827
Larch						
Accuracy	ND	-1.59	0.48	ND	ND	-0.88 (-27.8 %)
Precision	ND	2.13	1.38	ND	ND	2.14
P-value	ND	<0.001	0.05	ND	ND	<0.001
E_{imp}						7.90%
N	8	54	36	4	0	102
OC						
Accuracy %	ND	-0.21	-0.53	-1.14	ND	-0.51 (-21.4 %)

Cohort	<12 cm	12-20 cm	20-30 cm	30-40 cm	>40 cm	All classes
Precision	ND	1.34	1.69	1.83	ND	1.65
P-value	ND	0.544	0.05	0.02	ND	0.06
E _{imp}						14.70%
N	5	77	66	31	19	198
FGB						
Accuracy	<0.001	1.44	3.06	4.19	ND	2.0 (128.1 %)
Precision	1.49	1.85	1.87	2.47	ND	2.28
P-value	0.2	<0.001	<0.0001	<0.0001	ND	<0.0001
E _{imp}						8.70%
N	64	194	183	35	19	495
SGB						
Accuracy	ND	-0.28	-0.23	-0.67	-1.24	-0.50 (-30.5 %)
Precision	ND	1.27	1.73	1.7	1.91	1.68
P-value	ND	0.37	0.75	0.17	<0.001	0.11
E _{imp}						55.10%

Tier 1 time series trend analysis was also conducted as specified in section 6.3 of the IPCC good practice guidance for LULUCF (Table 6.5.11). The increased uncertainty in the trends based on the 1990 base year are associated with increase harvests and age class distribution shifts over the time series, as described in text relating to Table 6.5.2 and Figure 6.3.5.

Table 6.5.11 Uncertainty analysis for forest land remaining forest land since 1990 category³

Year	Category	Year emission/ reductions (kt CO ₂ eq)	Base year emission/ reductions (kt CO ₂ eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
1990	CRF 4A.1	-2821.0	-2821.0	29.9	0.8	na
	CRF 4 (II)	134.0	134.0	78.5	0.1	na
	CRF4(V)	113.2	113.2	59.4	0.1	na
	Total	-2573.8	-2573.8	25.8	1.0	na
1991	CRF 4A.1	-2843.5	-2821.0	29.9	0.9	0.8
	CRF 4 (II)	134.0	134.0	78.5	0.1	0.0
	CRF4(V)	72.8	113.2	59.4	0.0	-35.7
	Total	-2636.7	-2573.8	25.8	1.0	2.4
1992	CRF 4A.1	-2123.5	-2821.0	36.7	0.9	-24.7
	CRF 4 (II)	134.0	134.0	78.5	0.1	0.0
	CRF4(V)	46.8	113.2	59.4	0.0	-58.7
	Total	-1942.7	-2573.8	26.9	1.0	-24.5
1993	CRF 4A.1	-2806.8	-2821.0	28.9	0.8	-0.5
	CRF 4 (II)	134.0	134.0	78.5	0.1	0.0
	CRF4(V)	94.3	113.2	59.4	0.1	-16.7
	Total	-2578.5	-2573.8	26.0	1.0	0.2
1994	CRF 4A.1	-2266.2	-2821.0	33.3	0.8	-19.7
	CRF 4 (II)	134.0	134.0	78.5	0.1	0.0
	CRF4(V)	108.3	113.2	59.4	0.1	-4.4
	Total	-2024.0	-2573.8	25.8	1.0	-21.4
1995	CRF 4A.1	-1817.8	-2821.0	39.3	0.8	-35.6
	CRF 4 (II)	133.9	134.0	78.5	0.1	-0.1
	CRF4(V)	147.9	113.2	59.4	0.1	30.6
	Total	-1536.0	-2573.8	26.0	1.0	-40.3
1996	CRF 4A.1	-1541.8	-2821.0	44.8	0.8	-45.3
	CRF 4A II	133.9	134.0	78.5	0.1	-0.1
	CRF4(V)	164.4	113.2	59.4	0.1	45.2
	Total	-1243.4	-2573.8	26.5	1.0	-51.7

³ Note that uncertainties for category 4A(II) and 4A(V) include land in the forest remaining forest land category

Year	Category	Year emission/ reductions (kt CO2 eq)	Base year emission/ reductions (kt CO2 eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
1997	CRF 4A.1	-2556.1	-2821.0	29.6	0.8	-9.4
	CRF 4 (II)	133.8	133.8	78.5	0.1	0.0
	CRF4(V)	89.9	113.2	59.4	0.1	-20.6
	Total	-2332.4	-2574.0	26.0	1.0	-9.4
1998	CRF 4A.1	-1704.8	-2821.0	39.9	0.8	-39.6
	CRF 4 (II)	133.7	134.0	78.5	0.1	-0.3
	CRF4(V)	47.4	113.2	59.4	0.0	-58.1
	Total	-1523.7	-2573.8	27.3	1.0	-40.8
1999	CRF 4A.1	-1572.9	-2821.0	41.7	0.8	-44.2
	CRF 4 (II)	133.6	134.0	78.5	0.1	-0.4
	CRF4(V)	38.7	113.2	59.4	0.0	-65.8
	Total	-1400.6	-2573.8	27.8	1.0	-45.6
2000	CRF 4A.1	-905.0	-2821.0	64.9	0.8	-67.9
	CRF 4 (II)	133.2	134.0	78.5	0.1	-0.6
	CRF4(V)	97.2	113.2	59.4	0.1	-14.1
	Total	-674.5	-2573.8	29.4	1.0	-73.8
2001	CRF 4A.1	-657.4	-2821.0	87.9	0.7	-76.7
	CRF 4 (II)	132.9	134.0	78.5	0.1	-0.8
	CRF4(V)	193.8	113.2	59.4	0.1	71.2
	Total	-330.7	-2573.8	32.5	1.0	-87.2
2002	CRF 4A.1	-164.6	-2821.0	335.8	0.8	-94.2
	CRF 4 (II)	132.6	134.0	78.5	0.2	-1.1
	CRF4(V)	44.5	113.2	59.4	0.0	-60.7
	Total	12.6	-2573.8	48.8	1.0	-100.5
2003	CRF 4A.1	-633.7	-2821.0	88.8	0.7	-77.5
	CRF 4 (II)	132.3	134.0	78.5	0.1	-1.3
	CRF4(V)	274.8	113.2	59.4	0.2	142.7
	Total	-226.6	-2573.8	34.5	1.0	-91.2
2004	CRF 4A.1	-1428.4	-2821.0	43.0	0.8	-49.4
	CRF 4 (II)	132.0	134.0	78.5	0.1	-1.5
	CRF4(V)	160.1	113.2	59.4	0.1	41.4
	Total	-1136.3	-2573.8	26.8	1.0	-55.9
2005	CRF 4A.1	-741.9	-2821.0	76.1	0.8	-73.7
	CRF 4 (II)	131.7	131.7	78.5	0.1	0.0
	CRF4(V)	-58.2	113.2	59.4	0.0	-151.4
	Total	-668.4	-2576.1	31.8	1.0	-74.1
2006	CRF 4A.1	-440.1	-2821.0	123.2	0.8	-84.4
	CRF 4A II	131.0	134.0	78.5	0.2	-2.3
	CRF4(V)	58.2	113.2	59.4	0.1	-48.6
	Total	-251.0	-2573.8	37.0	1.0	-90.2
2007	CRF 4A.1	-901.3	-2821.0	61.1	0.8	-68.0
	CRF 4 (II)	130.4	134.0	78.5	0.1	-2.7
	CRF4(V)	65.4	113.2	59.4	0.1	-42.2
	Total	-705.5	-2573.8	29.9	1.0	-72.6
2008	CRF 4A.1	-1919.4	-2821.0	32.9	0.8	-32.0
	CRF 4A II	129.7	134.0	78.5	0.1	-3.3
	CRF4(V)	50.9	113.2	59.4	0.0	-55.1
	Total	-1738.9	-2573.8	26.9	1.0	-32.4
2009	CRF 4A.1	-1737.7	-2821.0	34.7	0.7	-38.4
	CRF 4 (II)	129.4	134.0	78.5	0.2	-3.5
	CRF4(V)	-28.5	113.2	59.4	0.0	-125.2
	Total	-1636.8	-2573.8	27.7	1.0	-36.4
2010	CRF 4A.1	-766.7	-2821.0	68.6	0.7	-72.8
	CRF 4 (II)	129.1	134.0	78.5	0.1	-3.7
	CRF4(V)	185.1	113.2	59.4	0.1	63.5
	Total	-452.5	-2573.8	31.0	1.0	-82.4
2011	CRF 4A.1	-428.2	-2821.0	118.0	0.8	-84.8
	CRF 4 (II)	128.5	134.0	78.5	0.2	-4.1
	CRF4(V)	68.9	113.2	59.4	0.1	-39.2
	Total	-230.8	-2573.8	36.4	1.0	-91.0

Year	Category	Year emission/ reductions (kt CO ₂ eq)	Base year emission/ reductions (kt CO ₂ eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
2012	CRF 4A.1	251.5	-2821.0	204.3	0.8	-108.9
	CRF 4 (II)	128.5	134.0	78.5	0.2	-4.1
	CRF4(V)	17.4	113.2	59.4	0.0	-84.6
	Total	397.5	-2573.8	133.8	1.0	-115.4
2013	CRF 4A.1	9.1	-2821.0	5687.5	0.8	-100.3
	CRF 4 (II)	128.4	134.0	78.5	0.2	-4.2
	CRF4(V)	72.5	113.2	59.4	0.1	-36.0
	Total	209.9	-2573.8	55.2	1.0	-108.2

Table 6.5.11 shows that the uncertainty of estimates for forest land remaining forest land was 25 per cent in 1990, increasing to 55.24 per cent by 2013. This is because the net emission in 2013 was close to zero, which makes the percentage uncertainty higher when the absolute uncertainty is much lower, when compared to other years.

The overall uncertainty for the 1990-2012 time series for the current submission was higher when compared to previous submissions because of the inclusion of CH₄, N₂O and DOC CO₂ emissions from organic soils in the new submission. The emission factors associated with these estimates carry a very high uncertainty (see Table 6.5.9).

6.3.5 Forest land remaining forest land (CRF 4.A.2)

Table 6.5.12 shows the net biomass, dead organic matter, soil C and net CO₂ emissions/removals for the time series 1990-2013 for lands converted to forest land (i.e. all forest established after 1990 reported in category 4.A.2). For the data time series pre 2006, the adjusted FORCARB estimates are reported. For the data time series 2007 to 2013, the CARBWARE model estimates are reported (Table 6.5.12). The methods used and values reported in category 4.A.2 and now fully consistent and comparable with KP emission/removals reported for AR activities, for the years 2008-2013 (see chapter 11).

The increase on removals by the post 1990 forest is due to an increase in forests area and productivity as new established forests mature. The slight decrease in the slope of the change in removals from 2007 onward is due to thinning harvests in productive forests at age 17 years old and onwards.

6.3.5.1 Time Series Adjustment of Living Biomass and DOM Pools

To ensure that there is no bias introduced in estimates over the time series due to the use of the different models, the 1990 to 2006 FORCARB series was adjusted (Table 6.5.12) and rescaled using tier 1 IPCC GPG (2006) time series overlap approaches:

- Living biomass gains (*LB_{gain}*, kt C) from the 2007 to 2012 time series for the CARBWARE and FORCARB model outputs were compared. The ratio (1.586) of the total CARBWARE and FORCARB *LB_{gain}* values for 2007-2012 was used to adjust the time series:

$$LB\ gain_{adj} = LB\ gain_{ini} \times 1.58 \dots \dots \dots (eq\ 6.3.36)$$

where, *LB_{gain_{adj}}* is the adjusted living biomass gain value and *LB_{gain_{ini}}* is initial FORCARB estimate. This method is consistent with eq 5.1 Chapter 5 of time series in the 2006 GPG;

- The adjusted biomass losses (*LBloss*) were also determined using equation 6.3.36 but using a ratio of 1.585 scaled using the ratio of living biomass losses of the total CARBWARE and FORCARB *LBloss* values for 2007-2012. The ratio of gains to losses (as applied to forest land remaining forest land) was not used because there are no harvest losses for the FORCARB time series 1990-2007;
- For dead organic matter (DOM), the ratio (1.911) of the average CARBWARE to average FORCARB values for 2007-2012 was used to adjust the time series:

$$DOM_{adj} = DOM_{ini.} \times 1.911 \dots \dots \dots (eq\ 6.3.37);$$

- There were no adjustments to the soil EF, since the FORCARB and CARBWARE estimates were identical.

Table 6.5.12 Time series for forest category 4.A.2

Year	Area (kHa)		CSC (kt C)						Net CO ₂ kt	Harvests ¹ (M m3)	
			Living biomass			DOM	Mineral Soils	Organic soils	Total		
	Gain	Loss	Net	Net	Net	Net					
	Total	Organic								Thinnings	Modelled
1990	15.82	9.92	0.00	-0.26	-0.26	0.01	NO	-7.19	27.26	NO	
1991	34.96	20.92	19.37	-0.71	18.66	0.12	NO	-14.83	-14.49	NO	
1992	51.66	30.51	41.33	-1.29	40.03	0.32	NO	-22.28	-66.28	NO	
1993	67.66	39.70	61.16	-1.97	59.19	0.56	NO	-28.32	-115.26	NO	
1994	87.12	50.88	87.94	-4.85	83.09	5.42	NO	-36.17	-191.94	NO	
1995	110.83	64.51	123.65	-8.66	114.99	11.38	NO	-48.38	-285.97	NO	
1996	131.81	76.57	165.64	-12.72	152.92	16.78	NO	-57.35	-411.97	NO	
1997	143.25	83.14	209.23	-17.11	192.12	21.92	NO	-61.33	-559.94	NO	
1998	156.17	90.57	255.19	-22.49	232.69	27.96	NO	-66.04	-713.57	NO	
1999	168.84	97.85	305.86	-29.07	276.79	35.25	NO	-70.24	-886.60	NO	
2000	184.54	106.88	367.71	-36.13	331.58	41.85	NO	-76.02	-1090.53	NO	
2001	200.00	115.62	433.89	-42.74	391.15	45.76	NO	-82.63	-1299.03	NO	
2002	215.06	123.98	502.11	-58.30	443.81	58.81	NO	-88.11	-1519.88	NO	
2003	224.15	128.94	583.98	-66.18	517.80	62.07	NO	-92.71	-1786.27	NO	
2004	233.89	134.16	664.95	-87.96	576.99	77.16	NO	-97.06	-2042.67	NO	
2005	243.99	139.46	757.80	-95.81	661.99	78.65	NO	-101.56	-2343.31	NO	
2006	252.02	143.12	853.49	-140.81	712.68	97.46	NO	-103.71	-2590.25	NO	
2007	259.20	146.39	954.36	-238.41	715.95	116.08	NO	-105.72	-2663.12	0.08	0.08
2008	265.45	149.23	1073.96	-262.76	811.20	125.00	NO	-107.72	-3037.77	0.02	0.02
2009	272.10	152.25	1185.35	-350.33	835.02	170.73	NO	-110.55	-3282.39	0.21	0.22
2010	280.41	156.04	1285.32	-383.79	901.53	182.52	NO	-112.92	-3560.80	0.14	0.15
2011	287.06	159.07	1347.64	-438.09	909.55	201.97	NO	-116.20	-3649.53	0.18	0.18
2012	292.91	161.73	1377.42	-461.09	916.33	207.18	NO	-118.68	-3684.39	0.11	0.11
2013	299.07	164.50	1406.22	-462.92	943.30	205.02	NO	-119.82	-3771.15	0.21	0.20

The harvest volumes show a comparison of the EUROSTAT and modelled harvest using FORECARB and the CARBWARE model.

Figure 6.3.11 shows the initial FORCARB estimates (blue symbols) and the time series adjustment as (red symbols) reported in the CRF table 4.A.2 and Table 6.5.12. Both time series show the same trend but the adjusted values show a higher net removal of CO₂. This is due to fundamental differences in the model input variables and the spatial scale at which the FORCARB and CARBWARE models operate. There are also known underestimated biases in the FORCARB model introduced when BFC yield tables are used. These are introduced by the same factors as those discussed under the forest land converted to forest land section. The final adjusted time series comprises of adjusted FORCARB estimates for the period 1990-2006 and CARBWARE estimated for the period 2007-2013.




Figure 6.3.11 The adjusted time series for forest category 4.A.2

6.3.5.2 Mineral Soils

Land converted to forest land (L-FL since 1990) on mineral soils are demonstrated not to be a source (see chapter 11.4), regardless of forest type and disturbance. Mineral CSC is reported as NO under this land category in CRF Table 4.A.

6.3.5.3 Organic Soils

The same approaches as described for forest land remaining forest land were used to estimate emissions from organic soils in lands converted to forest land since 1990 (section 6.3.4.3). However, there are no forests older than 50 years-old in the categories reported in 4.A.2, so this was not considered (Table 6.5.13). The allocation of emission estimates for the sub-categories CRF 4.A.2. 2.1 to 2.5 are based on the proportion of lands converted to forests described above.

Table 6.5.13 Area (in kha) and emissions from different organic soils types over the times series for land converted to forests

					Drained productive		On-site emissions		Off-site DOC emissions		Total
Year	Total Area ¹	Mineral	Org.	Open area ²	Org. ³	Org.-mineral ³	Org. ⁴	Org.-mineral ⁴	Org. ⁵	Org.-mineral ⁵	
	(kHa)						kt C				kt CO ₂
1990	15.8	5.9	9.9	1.9	6.6	2.1	-3.9	-0.6	-2.0	-0.7	26.4
1991	35.0	14.0	20.9	4.2	12.8	5.4	-7.6	-1.6	-4.0	-1.7	54.4
1992	51.7	21.2	30.5	6.1	19.8	7.4	-11.7	-2.2	-6.1	-2.3	81.7
1993	67.7	28.0	39.7	8.1	25.0	9.6	-14.7	-2.8	-7.7	-3.0	103.8
1994	87.1	36.2	50.9	10.4	32.6	11.3	-19.2	-3.3	-10.1	-3.5	132.6
1995	110.8	46.3	64.5	13.2	43.0	16.0	-25.4	-4.7	-13.3	-5.0	177.4
1996	131.8	55.2	76.6	15.7	52.0	17.4	-30.7	-5.1	-16.1	-5.4	210.3
1997	143.2	60.1	83.1	17.0	55.8	18.3	-32.9	-5.4	-17.3	-5.7	224.9
1998	156.2	65.6	90.6	18.6	59.5	20.7	-35.1	-6.1	-18.4	-6.4	242.2
1999	168.8	71.0	97.9	20.1	62.6	22.9	-36.9	-6.8	-19.4	-7.1	257.5
2000	184.5	77.7	106.9	22.0	67.2	25.7	-39.6	-7.6	-20.8	-8.0	278.7
2001	200.0	84.4	115.6	23.8	72.4	28.9	-42.7	-8.5	-22.4	-9.0	303.0
2002	215.1	91.1	124.0	25.6	76.6	31.6	-45.2	-9.3	-23.8	-9.8	323.1
2003	224.2	95.2	128.9	26.7	80.9	32.9	-47.7	-9.7	-25.1	-10.2	339.9
2004	233.9	99.7	134.2	27.8	84.8	34.3	-50.0	-10.1	-26.3	-10.6	355.9
2005	244.0	104.5	139.5	29.0	88.6	36.0	-52.3	-10.6	-27.5	-11.2	372.4
2006	252.0	108.9	143.1	31.4	88.8	39.1	-52.4	-11.6	-27.5	-12.1	380.3
2007	259.2	112.8	146.4	32.5	90.1	41.4	-53.1	-11.8	-27.9	-12.8	387.7
2008	265.4	116.2	149.2	33.4	90.8	43.7	-53.6	-12.4	-28.2	-13.6	395.0
2009	272.1	119.8	152.3	34.4	92.2	45.7	-54.4	-13.4	-28.6	-14.2	405.3
2010	280.4	124.4	156.0	35.6	94.3	47.7	-55.6	-13.3	-29.2	-14.8	414.1
2011	287.1	128.0	159.1	36.6	96.1	49.1	-56.7	-14.5	-29.8	-15.2	426.1
2012	292.9	131.2	161.7	37.6	97.8	50.2	-57.7	-15.1	-30.3	-15.6	435.2
2013	299.1	134.6	164.5	38.6	98.3	50.5	-58.0	-15.7	-30.5	-15.7	439.3

¹ Total area includes open areas

² Open area within forest areas (roads, extraction routes, biodiversity etc).

³ Productive area of drained organic (org.) and organo-mineral soils based in NFI 2006 and 2012 (excluding open areas). Organic soils include all soils with a > 20% C and an organic layer greater than 30 cm (e.g. Blanket peats, fens, cutaway peats. Organo-mineral soils are mineral soils with an organic overlay of < 30cm. These include peaty podsols and peaty gleys (Source NFI).

⁴ On-site emissions are calculated using Eq 6.3.23 and 6.2.24 and areas of productive and drained organic and organo-mineral soils less than 50 years old.

⁵ Off-site emissions are calculated using and EF of -0.31 tC /ha and the area of drained productive organic and organo-mineral soils using Eq 2.4 and Table 2.2, Ch2 of the 2013 Wet land supplement.

6.3.5.4 Emissions from Biomass Burning

The methodology for estimating emissions from biomass burning is discussed in category 4.A.1 (see section 6.3.4.4). Fires are only reported from 2008 onwards because fires generally only occur in forest at the pre-thicket stage of growth when there is enough woody biomass to act as a source for combustion by wild fires (Table 6.5.6). Fires on land afforested since 1990 represent 37 per cent to 39 per cent of the total fire areas over the period 2008 to 2013 (Table 6.5.6).

6.3.5.5 Emissions of N₂O from Fertilization

Ireland does not report separately the emissions of N₂O due to fertiliser use for 4.A Forest Land. The amount of synthetic fertiliser used in forests is negligible compared to that used in agriculture and therefore all N₂O emissions from fertiliser applications are reported under agriculture. The notation key IE is therefore used in CRF Table 4(I).

6.3.5.6 Emissions of N₂O and CH₄ from drainage and rewetted organic soils

The methodology for estimating N₂O and CH₄ emissions from drainage of organic soils are discussed section 6.3.4.6.

The methodology for estimating N₂O and CH₄ emissions from drainage of organic soils are discussed section 6.3.4.6.

a) N₂O from drained organic soils

The increase in N₂O emissions from organic soils in the land converted to forest land category since 1990 is due to an increase in afforestation of organic soils under the grants and premiums scheme (Table 6.5.14).

b) CH₄ from drained lands and ditches

The increase in CH₄ emissions from organic soils in land converted to forest land category since 1990 is due to an increase in afforestation of organic soils under the grants and premiums scheme (Table 6.5.14).

c) Rewetting of organic soils

Forest soils are managed to maintain drains so that nutrient uptake and crop productivity is maintained. Therefore, forest soils are not rewetted.

Table 6.5.14 The area activity data and N₂O and CH₄ emissions from drainage of land converted to forest land

Year	Area (kHa)				kt N ₂ O			kt CH ₄		
	Total Area	Open area	Organic N-poor	Organic N-rich	Organic N-poor	Organic N-rich	Total N ₂ O	Drained lands	Ditches	Total CH ₄
1990	15.8	1.9	6.6	2.1	0.00	0.01	0.01	0.02	0.04	0.06
1991	35.0	4.2	12.8	5.4	0.01	0.02	0.03	0.04	0.07	0.12
1992	51.7	6.1	19.8	7.4	0.01	0.03	0.05	0.07	0.11	0.18
1993	67.7	8.1	25.0	9.6	0.02	0.04	0.06	0.08	0.14	0.23
1994	87.1	10.4	32.6	11.3	0.02	0.05	0.07	0.11	0.19	0.29
1995	110.8	13.2	43.0	16.0	0.03	0.07	0.10	0.14	0.25	0.39
1996	131.8	15.7	52.0	17.4	0.04	0.08	0.11	0.17	0.30	0.47
1997	143.2	17.0	55.8	18.3	0.04	0.08	0.12	0.18	0.32	0.50
1998	156.2	18.6	59.5	20.7	0.04	0.09	0.13	0.20	0.34	0.53
1999	168.8	20.1	62.6	22.9	0.04	0.10	0.14	0.21	0.36	0.57
2000	184.5	22.0	67.2	25.7	0.05	0.11	0.16	0.23	0.38	0.61
2001	200.0	23.8	72.4	28.9	0.05	0.13	0.18	0.25	0.41	0.66
2002	215.1	25.6	76.6	31.6	0.05	0.14	0.19	0.27	0.44	0.71
2003	224.2	26.7	80.9	32.9	0.06	0.14	0.20	0.28	0.46	0.74
2004	233.9	27.8	84.8	34.3	0.06	0.15	0.21	0.29	0.49	0.78
2005	244.0	29.0	88.6	36.0	0.06	0.16	0.22	0.31	0.51	0.81
2006	252.0	31.4	88.8	39.1	0.06	0.17	0.23	0.31	0.51	0.83
2007	259.2	32.5	90.1	41.4	0.06	0.18	0.25	0.32	0.52	0.84
2008	265.4	33.4	90.8	43.7	0.06	0.19	0.26	0.33	0.53	0.86
2009	272.1	34.4	92.2	45.7	0.06	0.20	0.27	0.34	0.54	0.87
2010	280.4	35.6	94.3	47.7	0.07	0.21	0.28	0.35	0.55	0.90
2011	287.1	36.6	96.1	49.1	0.07	0.22	0.28	0.36	0.56	0.92
2012	292.9	37.6	97.8	50.2	0.07	0.22	0.29	0.36	0.57	0.93
2013	299.1	38.6	98.3	50.5	0.07	0.22	0.29	0.37	0.57	0.94

6.3.5.7 N₂O emissions from mineral soils as a result of land use change of management (F_{SOM})

Emissions of N₂O from mineral soils are based on mineralisation rates due to loss of organic C from mineral soils. Based on available scientific evidence there we show that management and afforestation of mineral soils results in no significant change in soil organic carbon (SOC, See Ch 11). Therefore, emissions due to mineralisation of forest soil do not occur (NO).

6.3.5.8 CO₂ emissions from urea application to soils

All fertiliser application related emissions, including CO₂ emissions from urea application are reported under Agriculture because these are based on national sales data (IE).

6.3.5.9 Uncertainty Analysis for Category 4.A.2

The same uncertainty analysis was carried out for lands converted to forest land as was done for forests remaining forest land (Tables 6.3.15). The only different sources of uncertainty in this analysis (see Table 6.5.8 and 6.3.9) was the uncertainty due to re-adjustment scaling factor uncertainty for biomass (LB), litter and dead wood (DOM), i.e. Codes J and K Table 6.5.8), which were 3.7 and 9.0 per cent for lands converted to forest land.

Table 6.5.14 shows that the uncertainty of estimates for land converted to forest land was 96 per cent in 1990, decreasing to 20.2 per cent by 2013. This is because the net emission reduction in 1990 was close to zero, which makes the percentage uncertainty higher when the absolute uncertainty is much lower, when compared to other years. The same issue arises when trend uncertainty compared to a base year is analysed. Hence the use of a base year, where absolute values are zero, or where values are very small, and expressing these as a percentage does not truly reflect the absolute uncertainty.

The overall uncertainty for the 1990-2012 time series for the current submission was 4 to 7 per cent higher when compared to previous submissions because of the inclusion of CH₄, N₂O and DOC CO₂ emissions from organic soils in the new submission. The emission factors associated with these estimates carry a very high uncertainty (see Table 6.5.9).

Table 6.5.15 Uncertainty analysis of lands converted to forest land as reported in CRF 4.A.2

Year	Category	Year emission/reductions (kt CO ₂ eq)	Base year emission/reductions (kt CO ₂ eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
1990	CRF 4A.2	27.3	27.3	96.6	0.8	na
	CRF 4(II)	8.7	8.7	88.4	0.2	na
	CRF4(V)	0.0	0.0	0.0	0.0	na
	Total	35.9	35.9	96.6	1.0	na
1991	CRF 4A.2	-14.5	27.3	383.7	0.8	-153.2
	CRF 4(II)	18.3	8.7	91.8	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	3.8	35.9	74.3	1.0	-153.2
1992	CRF 4A.2	-66.3	27.3	129.0	0.8	-343.2
	CRF 4(II)	27.2	8.7	90.4	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-39.1	35.9	129.0	1.0	-343.2
1993	CRF 4A.2	-115.3	27.3	95.7	0.8	-522.9
	CRF 4(II)	34.6	8.7	90.7	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-80.6	35.9	95.7	1.0	-522.9

Year	Category	Year emission/reductions (kt CO2 eq)	Base year emission/reductions (kt CO2eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
1994	CRF 4A.2	-191.9	27.3	74.3	0.8	-804.2
	CRF 4(II)	43.9	43.9	89.4	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-148.0	71.2	74.3	1.0	-804.2
1995	CRF 4A.2	-286.0	27.3	67.1	0.8	-1149.2
	CRF 4(II)	59.0	8.7	90.2	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-227.0	35.9	67.1	1.0	-1149.2
1996	CRF 4A.2	-412.0	27.3	56.3	0.8	-1611.5
	CRF 4(II)	69.4	8.7	88.9	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-342.5	35.9	56.3	1.0	-1611.5
1997	CRF 4A.2	-559.9	27.3	45.8	0.8	-2154.4
	CRF 4(II)	74.1	74.1	88.7	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-485.8	101.4	45.8	1.0	-2154.4
1998	CRF 4A.2	-713.6	27.3	39.9	0.8	-2718.0
	CRF 4(II)	80.2	8.7	89.4	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-633.4	35.9	39.9	1.0	-2718.0
1999	CRF 4A.2	-886.6	27.3	35.3	0.8	-3352.8
	CRF 4(II)	85.6	8.7	90.1	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-801.0	35.9	35.3	1.0	-3352.8
2000	CRF 4A.2	-1090.5	27.3	32.2	0.8	-4101.0
	CRF 4(II)	92.9	8.7	90.6	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-997.6	35.9	32.2	1.0	-4101.0
2001	CRF 4A.2	-1299.0	27.3	30.3	0.8	-4866.0
	CRF 4(II)	101.3	8.7	91.1	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-1197.7	35.9	30.3	1.0	-4866.0
2002	CRF 4A.2	-1519.9	27.3	28.4	0.8	-5676.3
	CRF 4(II)	108.3	8.7	91.5	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-1411.6	35.9	28.4	1.0	-5676.3
2003	CRF 4A.2	-1786.3	27.3	26.7	0.8	-6653.6
	CRF 4(II)	113.8	8.7	91.4	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-1672.5	35.9	26.7	1.0	-6653.6
2004	CRF 4A.2	-2042.7	27.3	25.2	0.8	-7594.3
	CRF 4(II)	119.1	8.7	91.3	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-1923.6	35.9	25.2	1.0	-7594.3
2005	CRF 4A.2	-2343.3	27.3	24.2	0.8	-8697.3
	CRF 4(II)	124.7	124.7	91.3	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-2218.7	151.9	24.2	1.0	-8697.3
2006	CRF 4A.2	-2590.3	27.3	23.1	0.8	-9603.4
	CRF 4(II)	128.0	8.7	92.3	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-2462.3	35.9	23.1	1.0	-9603.4
2007	CRF 4A.2	-2663.1	27.3	22.8	0.8	-9870.7
	CRF 4(II)	131.5	8.7	92.9	0.2	
	CRF4(V)	0.0	0.0	0.0	0.0	
	Total	-2531.6	35.9	22.8	1.0	-9870.7
2008	CRF 4A.2	-3037.8	27.3	21.7	0.8	-11045.3
	CRF 4(II)	134.6	8.7	93.4	0.2	
	CRF4(V)	16.5	0.0	52.2	0.0	

Year	Category	Year emission/reductions (kt CO2 eq)	Base year emission/reductions (kt CO2eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
	Total	-2886.7	35.9	21.8	1.0	-8133.7
2009	CRF 4A.2	-3282.4	27.3	20.9	0.8	-12142.7
	CRF 4A II	137.8	8.7	93.8	0.2	
	CRF4(V)	9.4	0.0	52.2	0.0	
	Total	-3135.1	35.9	20.9	1.0	-8824.9
2010	CRF 4A.2	-3560.8	27.3	20.4	0.8	-13164.2
	CRF 4(II)	141.9	8.7	94.0	0.1	
	CRF4(V)	62.9	0.0	52.2	0.0	
	Total	-3355.9	35.9	20.7	1.0	-9439.5
2011	CRF 4A.2	-3649.5	27.3	20.3	0.8	-13489.7
	CRF 4(II)	145.2	8.7	94.1	0.2	
	CRF4(V)	23.2	0.0	52.2	0.0	
	Total	-3481.1	35.9	20.4	1.0	-9787.9
2012	CRF 4A.2	-3684.4	27.3	20.3	0.8	-13617.6
	CRF 4(II)	148.0	8.7	94.2	0.2	
	CRF4(V)	5.9	0.0	52.2	0.0	
	Total	-3530.5	35.9	20.4	1.0	-9925.3
2013	CRF 4A.2	-3771.2	27.3	20.3	0.8	-13935.9
	CRF 4(II)	148.8	8.7	94.2	0.1	
	CRF4(V)	72.5	0.0	52.2	0.0	
	Total	-3549.9	35.9	20.3	1.0	-9979.3

6.3.6 Deforestation Areas (CRF 4.B.2 to 4.F.2)

This section describes deforestation areas reported under forest converted to other lands under sub-categories 4.B.2 to 4.F.2. In previous submissions, deforestation was reported only in respect of forest land converted to settlements, derived from CORINE data. Following the ERT team recommendations in the 2010 centralised review, Ireland has developed a new system to track deforestation trends going back to 1990 (see sections 6.3.2.4). Historical deforestation trends show a marked increase in deforestation from 2000 to 2006 and a shift in the major land use transitions into grassland before 2000 and to settlements, wetlands and other land after 2000. These findings are consistent with a) an increase in building and infrastructural developments on forest land due to high economic growth in the late 1990s to mid-2000s; and b) an increase in deforestation of peatland forests following the introduction of EU life peatland restoration scheme in the 2004.

The new NFI data for 2012 allowed the recalculation of forest areas for 2006 to 2012. The development of the new methodology resulted in an increase in the reported area of deforestation from 1.38 to 12.512 kha in 2008, with a further 0.8 kha being deforested in 2009, 0.8 kha in 2010, 1.6 kha in 2011 and 0.8 kha in 2012. In 2013, the limited felling licence records and lands taken out database identified deforested areas of 0.129 kha to grasslands and settlements (see chapter 11 and Table 7.3.16). The increase in conversion of forest land to other lands after 2005 is associated with clearfelled forest land which was not replanted within a 5 year period as determined using the 2006 and 2012 NFI. The lands have not been converted any known land use but are classified as deforestation and reported as forest converted to other land.

The estimate of final land use after deforestation is based on an analysis of the CORINE land cover change from 1990 to 2000, the National Forest Inventory and the Forest Inventory and Planning Strategy, FIPS) data up to 2005. Post 2006 analysis is based on detailed information from the 2006 and 2012 NFI. The 2013 data is based on felling licence information and the lands taken out database (see 6.3.2.4).

Table 6.5.16 The new deforestation, land use change and soil type matrix showing annual deforestation areas (kha/ year) associated with different land uses and soils categories

	TOTAL Area		Grassland		Settlement		Wetland		Other	
	Total	Organic	Total	Organic	Total	Organic	Total	Organic	Total	Organic
1990	0.021	0.006	0.008	0.006	0.010	NO	NO	NO	0.002	NO
1991	0.041	0.012	0.016	0.012	0.020	NO	NO	NO	0.004	NO
1992	0.062	0.017	0.025	0.017	0.031	NO	NO	NO	0.007	NO
1993	0.082	0.023	0.033	0.023	0.041	NO	NO	NO	0.009	NO
1994	0.103	0.029	0.041	0.029	0.051	NO	NO	NO	0.011	NO
1995	0.436	0.029	0.308	0.029	0.051	NO	NO	NO	0.078	NO
1996	0.770	0.029	0.574	0.029	0.051	NO	NO	NO	0.144	NO
1997	1.103	0.029	0.841	0.029	0.051	NO	NO	NO	0.211	NO
1998	1.436	0.029	1.108	0.029	0.051	NO	NO	NO	0.277	NO
1999	1.770	0.029	1.375	0.029	0.051	NO	NO	NO	0.344	NO
2000	2.627	0.258	1.775	0.086	0.222	NO	0.171	0.114	0.459	0.057
2001	3.484	0.487	2.175	0.144	0.394	NO	0.341	0.228	0.574	0.115
2002	4.341	0.715	2.575	0.201	0.565	NO	0.512	0.342	0.688	0.172
2003	5.198	0.944	2.975	0.259	0.737	NO	0.682	0.456	0.803	0.230
2004	6.055	1.173	3.376	0.316	0.908	NO	0.853	0.570	0.918	0.287
2005	6.912	1.402	3.776	0.373	1.079	NO	1.024	0.684	1.033	0.344
2006	8.912	3.002	3.776	0.373	1.479	0.400	1.024	0.684	2.633	1.544
2007	10.512	4.202	3.776	0.373	2.679	1.200	1.424	1.084	2.633	1.544
2008	12.512	5.002	4.176	0.373	3.879	2.000	1.424	1.084	3.033	1.544
2009	13.312	5.402	4.576	0.373	3.879	2.000	1.824	1.484	3.033	1.544
2010	14.112	5.402	5.376	0.373	3.879	2.000	1.824	1.484	3.033	1.544
2011	15.712	5.802	6.576	0.373	3.879	2.000	2.224	1.884	3.033	1.544
2012	16.512	6.602	6.576	0.373	4.679	2.800	2.224	1.884	3.033	1.544
2013	16.640	6.615	6.640	0.374	4.744	2.812	2.224	1.884	3.033	1.544

* No transition from forests to croplands were detected

6.3.6.1 Deforestation Losses

Carbon stock changes associated with deforestation reported in all CRF tables include those for the total standing biomass of all trees removed at clear fell (i.e. all biomass carbon is assumed to be immediately oxidised):

$$C_{L(Total)} = C_{L(AB)} + C_{L(BB)} = TOTAL_{(Biomass)} lost \quad (6.3.38)$$

The carbon stock **losses (C_L)** in the **above ground (AB)** and **below ground (BB)** pools were calculated differently depending on the activity data available, but this was done in a hierarchical order:

- 1) Total biomass and DOM losses were directly determined from the NFI permanent sample plot tree data and allometric equations as described in section 6.3.3.1.2 above.
- 2) Where plots were clearfelled before 2006 from the standing volume (V) of the forest stand, as specified by Coillte plot queries, a basic density (D) in the range 0.35 to 0.55 (depending on tree species), a biomass expansion factor (BEF) of 1.68 to 4 t/t⁻¹ (Black at al., 2004) a carbon fraction (CF) of 0.5 and a root to shoot ratio R of 0.2, as described in Eq6.3.9 and 6.3.10 above). A list of plot data from Coillte provided information of deforestation area (including open areas), species, age, standing volume before clearfell.

- 3) There is no activity data for deforested areas before 2006, therefore the 2006-2013 mean AB (65.9.81 t C/ha), BB (17.2 tC/Ha), litter and deadwood (16.4 tC/Ha) C stock was applied as an IEF for these deforested areas. (see section 6.3.3.1.2).

It is important to note that many deforested lands are not fully stocked before clearfell and land use change, with the exception of:

- Cleafelled non-regenerated land within a 5 year period (1600 ha since 2006 with a mean biomass stock of 230 t C ha⁻¹),
- EU life bog restoration projects in 2007 (400 ha , biomass stock of 176 t C ha⁻¹),
- Wind farm conversions in 2007 (400 ha, biomass stock of 230 t C ha⁻¹),
- Conversion to settlements (ESB lines) in 2008,
- Grassland conversion in 2009.

All other deforestation events over the period 2006-2012, representing 6400 ha of the total 9600 ha had a lower biomass stock ranging from 1.1 to 112 t C ha⁻¹. These were younger aged crops, which were prematurely clearfelled for deforestation or scrub land forests converted to settlements.

The accumulated litter and DOM pool was assumed to be immediately oxidised when deforestation occurs. The approach adopted to apply an instantiations oxidation to litter and DOM (i.e. harvest residue, stumps and roots) in forests land converted to other land is based on the **conservativeness** principal. The rationale for this assumption is explained for the land use transitions for forestry indicated below:

- a) *Forest conversion to rewetted wetlands.* Most forest conversion to wetland involves EU wetland conservation measures, where drains are blocked to encourage peat vegetation regeneration. This would create anaerobic condition for remaining harvest residues (stumps, lying deadwood and litter) resulting in very low decay at rates than those used in Ch11 for 1st rotation crops.
- i. Organic soils emissions due to rewetting are now estimated using the new Wetland supplement guidelines (see section 6.3.6.1.1). Biomass gains after conversion to rewetted and regenerating wetlands are included in on-site removals (see section 3.2.1 of the wetland supplement) and are therefore reported as IE.
 - ii. The remaining forest conversions to wetlands occur for peat extraction (i.e. 400 ha in 2007). The tier 1 default of zero emissions/removals for biomass are applied to peat extraction sites (IPCC 2006 GPG and wetland supplement). The emissions from organic soils wetland soil in peat extraction are outlined in section 6.3.6.1.2 below).
- b) Recent evidence of forests conversion to grassland and settlements suggests that harvest residues are removed after harvest. The current common practice is to chip woody residues for bio-fuel or horticultural purposes (expert opinion, Forest Service). In this case, we would argue that instant oxidation should be applied since these are in essence harvested wood products and in the case of compost would decay relatively quickly. In some cases it is possible that forest residues are ploughed, piled up and left on site to decay over time. However, we have no data supporting this, so apply the conservative approach of instantiations oxidation.

Table 6.5.17 Deforestation carbon stock changes and harvest over the time series

Area (kha)		CSC (kt C)						Net CO ₂ kt	Harvests (m ³)
Year	Total	Living biomass			DOM Net	Mineral Soils Net	Organic soils Net	Total	Modelled
		Gain	Loss	Net					
1990	0.02	0.06	-1.71	-1.66	-0.50	-0.01	-0.02	8.06	4040
1991	0.04	0.06	-1.71	-1.66	-0.50	-0.03	-0.05	8.19	4040
1992	0.06	0.06	-1.71	-1.66	-0.50	-0.04	-0.07	8.33	4040
1993	0.08	0.06	-1.71	-1.66	-0.50	-0.05	-0.09	8.46	4040
1994	0.10	0.06	-1.71	-1.66	-0.43	-0.07	-0.11	8.33	4040
1995	0.44	1.81	-27.71	-25.89	-4.41	-0.14	-0.11	112.04	65364
1996	0.77	1.81	-27.71	-25.89	-5.46	-0.21	-0.11	116.17	65364
1997	1.10	1.81	-27.71	-25.89	-5.46	-0.29	-0.11	116.44	65364
1998	1.44	1.81	-27.71	-25.89	-5.46	-0.36	-0.11	116.71	65364
1999	1.77	1.81	-27.71	-25.89	-4.81	-0.43	-0.11	114.57	65364
2000	2.63	2.72	-71.24	-68.52	-10.85	-0.69	-0.44	295.15	168068
2001	3.48	2.72	-71.24	-68.52	-14.05	-0.94	-0.77	309.01	168068
2002	4.34	2.72	-71.24	-68.52	-14.05	-1.19	-1.10	311.15	168068
2003	5.20	2.72	-71.24	-68.52	-14.05	-1.44	-1.43	313.28	168068
2004	6.05	2.72	-71.24	-68.52	-14.05	-1.69	-1.76	315.41	168068
2005	6.91	2.72	-71.24	-68.52	-13.63	-1.94	-2.09	316.00	168068
2006	8.91	0.00	-392.16	-392.16	-39.79	-2.38	-3.34	1604.82	936429
2007	10.51	0.00	-177.56	-177.56	-8.12	-2.82	-4.09	706.21	417156
2008	12.51	2.72	-102.81	-100.09	-17.40	-3.70	-4.77	461.91	232740
2009	13.31	2.72	-92.44	-89.72	-6.16	-3.70	-6.02	387.21	222262
2010	14.11	5.44	-45.69	-40.25	-6.01	-3.69	-6.02	205.21	111220
2011	15.71	8.16	-56.57	-48.41	-28.58	-3.68	-6.21	318.55	114959
2012	16.51	0.00	-42.22	-42.22	-12.80	-3.66	-6.93	240.57	101250
2013	16.64	0.44	-5.88	-5.45	-0.79	-3.71	-6.94	61.89	13925

Note: Harvests from deforestation are immediately oxidised as biomass losses and therefore are not included in harvested wood product carbon stock changes

6.3.6.1.1 Mineral soils

Ireland has chosen not to account for carbon stock changes in mineral soils converted to grasslands. Verifiable information demonstrating that deforested grassland mineral soils are not a source of anthropogenic GHG is provided in chapter 11. The notation key NO is therefore used for mineral soils under this land category in CRF Table 4.C.

For deforested settlement (4E) and other land (4F) categories we use a **conservative estimate**, as used for other countries (e.g. Finland, Sweden), that 20% of SOC is emitted over a 20 year period in these soils. A mean SOC stock of 110 t C ha⁻¹ was used based on best available soil data (see Figure 11.3, Ch 11). It should be noted that this is a conservative approach since:

- All deforested land allocated to the other land use category (Table H2 appendix H) are forest lands which have been clearfelled but not replanted within a 5 year period. These lands have not undergone a land use transition but are defined as deforestation to comply with the requirements set out in the annex to decision of 2CMP/8 (see CH 11);
- Land converted to settlement contains green areas which will not reduce SOC as a result of deforestation. However it is assumed that the total deforested area emits CO₂ from mineral soils because there is no activity data to determine the percentage green area in urban areas.

Mineralisation emissions of N_2O due to the loss of SOC (F_{som}) due to deforestation to settlement and other land is estimated using Eq. 11.8 in Ch. 11 of the 2006 IPCC Guidelines, CSC for mineral soils (see above), and the default C:N ratio of 15.

6.3.6.1.2 Organic soils

Drained organic soils

Grasslands (4C): Emissions of CO_2 from deforested grasslands are assumed to occur because lands are likely to be shallowly drained because they are temperate rich organic soils. The default on-site emission factor on 3.6 t C/ha (Table 2.1 of the wetland supplement), and off-site EF_{DOC} of 0.31 t C /ha is used (Table 2.2 of the 2013 Wetland supplement).

The default emission factors and methods for temperate shallow drained nutrient rich organic soils are used for CH_4 emissions from deforested grasslands (Eq. 2.6 of IPCC wetland supplement, 2013). For CH_4 -land emissions the default EF_{CH_4-land} of 39 kg CH_4 /ha and $FRAC_{ditch}$ of 0.05 is used for shallow drained grasslands (Tables 2.3 and 2.4 of the Wetland supplement). The emission factor from shallow drains EF_{drain} of 527 kg CH_4 /ha is used (Table 2.4 of wetland supplement).

Default emission factors (1.6 kg N- N_2O /ha, Table 2.5 of the Wetland supplement) and methods (Eq. 2.7) for temperate shallow drained nutrient rich organic soils are used for N_2O emissions from deforested grasslands.

Settlements (4E) and other lands: The 2013 wetland supplement and 2006 IPCC Guidelines provide no methodology for drained organic soils under settlement. Therefore, emissions from organic soils converted to settlement and other land are assumed to continue using the on-site and DOC EFs and methods outlined in reported using eq. 6.3.23 and 6.3.24, described in section 6.3.3.1.2 (Soils). Emissions of CH_4 and N_2O in deforested settlements and other lands are not reported under Convention but are reported under the KP.

Peat extraction (4D): For the deforestation of land to peat extraction the default emission factors and methods are used (CH 2 of 2013 Wetland supplement). For CO_2 emissions, default on-site EF_{-land} 2.8 tC /ha and the EF_{DOC} of 0.31 t C/ha is used (Table 2.1 and 2.2 of the 2013 Wetland supplement).

Wetlands (4D): For CH_4 , CH_4 -land emissions the default EF_{CH_4-land} of 6.1 kg CH_4 /ha and $FRAC_{ditch}$ of 0.05 is used (Tables 2.3 and 2.4 of the 2013 Wetland supplement). The emission factor drained peat extraction sites of EF_{drain} of 542 kg CH_4 /ha is used (Table 2.4 of wetland supplement).

Default emission factors (1.6 kg N- N_2O /ha, Table 2.5 of the Wetland supplement) and methods (Eq. 2.7) for temperate shallow drained nutrient rich organic soils are used for N_2O emissions from deforested grasslands.

Rewetting of organic soils

Emissions from organic soils following forest conversion back to wetlands (4D i.e. rewetting of organic soils) include on-site emission/removals (i.e. C-composite) and off-site DOC emissions (section 3.2.1 of the 2013 wetland supplement). On site removals due to non-woody vegetation/organic soils bio-geochemical reactions are assumed to not occur (NO) because peat soils are re-saturated since drainage does not occur on regenerated wetlands (as part of EU life peatland regeneration projects). On-site emissions are estimated based on the area of rewetted soils, the default emission factor EF_{CO_2} of -0.23 (Table 3.1 and Eq 3.4 of the wetland supplement). Off-site DOC emissions are estimated using Eq. 3.5 and the default $EF_{DOC-rewetted}$ of -0.24 (Table 3.2 of the wetland

supplement). EF for CH₄ due to rewetting is 92 kg CH₄ per ha per (Eq 3.8 and table 3.3 of the 2013 wetland supplement).

6.3.6.1.3 Uncertainty for deforestation estimates

The same uncertainty analysis was carried out for lands converted to forest land as was done for forests remaining forest land (Tables 6.3.20). The only different sources of uncertainty in this analysis (see Table 6.5.18 and 6.3.19) was the uncertainty due different activity data used for deforestation areas and additional pools, particularly for category 4(III).

It is important to note that the uncertainty estimates and net emissions for deforestation are a sub-total of the total emissions presented in CRF 4(II) and 4(III) (i.e. this does not include emissions from other land uses (deforestation transitions)).

Table 6.5.18 Uncertainty estimates for individual activity and area data sets for deforested lands

Code	Component	Sub-category	% Uncertainty	Source
A	Biomass algorithms	AB and BB	12.00	Black et al 2007
B	C fraction	All biomass pools	0.87	Black et al 2007
C	Volume to biomass	Defor losses	38.50	Felling licences and BEF uncertainty, Black 2004
D	Area deforestation	NFI, OSI aerial photos	46.70	Sample strata uncertainly analysis/ new deforestation methods
E	Litter	Li	3.10	Tobin et al, 2006
F	Deadwood	DW	22.00	Tobin et al, 2007
G	Peat soils and DOC	So	90.00	Assume same as Tier 1 (Table 2.3,2.3.1 CH ₂ , 2006 IPCC Guidelines)
I	Drained area	N ₂ O and CH ₄ drained	12.30	Conf. interval of NFI analysis
J	N ₂ O emission factors drainage	N ₂ O emissions	119.00	Wetland supplement Table 2.5 and Yamulki et al., 2013
K	CH ₄ EF ditches	CH ₄ emissions	87.20	Wetland supplement Table 2.4
L	CH ₄ EF lands	CH ₄ emissions	126.00	Wetland supplement Table 2.3
M	N ₂ O EF mineralisation	N ₂ O emissions	66.00	2006 IPCC Guidelines Eq 11.8
N	DOC emissions from drained soils	So	43.50	Wetland supplement Table 2.1
O	Mineral soil EF to settlement	So	50.00	Guess, unknown
P	Peat extraction EF CO ₂	So	69.81	Wetland supplement Table 2.1 and 2.2
Q	Rewetting CO ₂	So	125.36	Wetland supplement Table 3.1and 3.2
R	Rewetting CH ₄	CH ₄ emissions	240.00	Wetland supplement Table 3.3

Table 6.5.19 Combined uncertainty estimates for deforested land

	Component	Equation in NIR	% uncertainty	Uncertainty of combined products (code)
LB net	Biomass	Eq 6.3.2 and 6.3.9	61.71	A, B, C, D
DOM	DOM (deadwood and litter)	Eq 6.3.16	64.48	B, C, D, E, F
SO	Soils	Eq 11.23 (So) and W/supplement	187.78	D, G, N, O, P, Q
N ₂ O	N ₂ O drainage	Wetland supplement	128.43	D, I, J
CH ₄	CH ₄ drain and rewetting	Wetland supplement	285.01	I, K, L, R

Table 6.5.20 Uncertainty estimates for deforested land

Year	Category	Year emission/ reductions (kt CO2 eq)	Base year emission/reductions (kt CO2eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base- year (% mean trend)
1990	CRF 4B-E	8.06	8.06	48.91	0.93	na
	CRF 4 (II)	0.01	0.01	197.87	0.01	na
	CRF4 (III)	0.43	0.43	66.00	0.07	na
	Total	8.50	8.50	47.61	1.00	na
1991	CRF 4B-E	8.19	8.06	48.41	0.87	1.65
	CRF 4 (II)	0.03	0.01	197.87	0.01	100.00
	CRF4 (III)	0.85	0.43	66.00	0.12	100.00
	Total	9.07	8.50	58.14	1.00	6.74
1992	CRF 4B-E	8.33	8.06	48.10	0.81	3.30
	CRF 4 (II)	0.04	0.01	197.87	0.02	200.00
	CRF4 (III)	1.28	0.43	66.00	0.17	200.00
	Total	9.64	8.50	50.90	1.00	13.47
1993	CRF 4B-E	8.46	8.06	47.99	0.77	4.96
	CRF 4 (II)	0.05	0.01	197.87	0.02	300.00
	CRF4 (III)	1.70	0.43	66.00	0.21	300.00
	Total	10.22	8.50	54.04	1.00	20.21
1994	CRF 4B-E	8.33	8.06	48.99	0.73	3.40
	CRF 4 (II)	0.07	0.07	197.87	0.02	0.00
	CRF4 (III)	2.13	0.43	66.00	0.25	400.00
	Total	10.53	8.55	58.14	1.00	23.12
1995	CRF 4B-E	112.04	8.06	53.14	0.95	1290.23
	CRF 4 (II)	0.07	0.01	197.87	0.00	400.00
	CRF4 (III)	4.42	0.43	66.00	0.05	937.10
	Total	116.52	8.50	47.71	1.00	1271.12
1996	CRF 4B-E	116.17	8.06	51.68	0.93	1341.60
	CRF 4 (II)	0.07	0.01	197.87	0.00	400.00
	CRF4 (III)	6.70	0.43	66.00	0.07	1474.19
	Total	122.95	8.50	47.62	1.00	1346.74
1997	CRF 4B-E	116.44	8.06	51.57	0.91	1344.93
	CRF 4 (II)	0.07	0.07	197.87	0.00	0.00
	CRF4 (III)	8.99	0.43	66.00	0.09	2011.29
	Total	125.50	8.55	47.81	1.00	1367.46
1998	CRF 4B-E	116.71	8.06	51.48	0.89	1348.26
	CRF 4 (II)	0.07	0.01	197.87	0.00	400.00
	CRF4 (III)	11.28	0.43	66.00	0.11	2548.39
	Total	128.06	8.50	48.25	1.00	1406.89
1999	CRF 4B-E	114.57	8.06	52.19	0.87	1321.66
	CRF 4 (II)	0.07	0.01	197.87	0.00	400.00
	CRF4 (III)	13.56	0.43	66.00	0.13	3085.48
	Total	128.20	8.50	49.01	1.00	1408.58
2000	CRF 4B-E	295.15	8.06	53.31	0.91	3562.46
	CRF 4 (II)	0.63	0.01	223.63	0.01	4562.82
	CRF4 (III)	21.42	0.43	66.00	0.08	4930.65
	Total	317.20	8.50	47.75	1.00	3632.61
2001	CRF 4B-E	309.01	8.06	51.45	0.88	3734.51
	CRF 4 (II)	1.19	0.01	225.12	0.01	8725.64
	CRF4 (III)	29.28	0.43	66.00	0.11	6775.81
	Total	339.49	8.50	48.27	1.00	3894.86
2002	CRF 4B-E	311.15	8.06	51.21	0.85	3760.96
	CRF 4 (II)	1.76	0.01	225.66	0.02	12888.45
	CRF4 (III)	37.14	0.43	66.00	0.13	8620.97
	Total	350.04	8.50	49.22	1.00	4019.03
2003	CRF 4B-E	313.28	8.06	51.00	0.82	3787.41
	CRF 4 (II)	2.32	0.01	225.94	0.03	17051.27
	CRF4 (III)	44.99	0.43	66.00	0.15	10466.13
	Total	360.59	8.50	50.47	1.00	4143.20
2004	CRF 4B-E	315.41	8.06	50.83	0.79	3813.86
	CRF 4 (II)	2.89	0.01	226.11	0.03	21214.09
	CRF4 (III)	52.85	0.43	66.00	0.17	12311.29
	Total	371.15	8.50	51.95	1.00	4267.38
2005	CRF 4B-E	316.00	8.06	50.87	0.77	3821.17
	CRF 4 (II)	3.45	3.45	226.22	0.04	0.00
	CRF4 (III)	60.71	0.43	66.00	0.19	14156.45
	Total	380.15	8.50	53.66	1.00	4373.40
2006	CRF 4B-E	1604.82	8.06	55.65	0.94	19814.04
	CRF 4 (II)	5.68	0.01	156.68	0.01	41848.92

Year	Category	Year emission/ reductions (kt CO2 eq)	Base year emission/reductions (kt CO2eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base- year (% mean trend)
2007	CRF4 (III)	74.44	0.43	66.00	0.05	17382.26
	Total	1,684.95	8.50	47.62	1.00	19727.29
	CRF 4B-E	706.21	8.06	57.35	0.85	8663.25
	CRF 4 (II)	8.02	0.01	159.77	0.03	59145.38
	CRF4 (III)	88.18	0.43	66.00	0.12	20608.06
2008	Total	802.41	8.50	49.68	1.00	9342.21
	CRF 4B-E	461.91	8.06	51.41	0.73	5631.79
	CRF 4 (II)	9.14	0.01	148.45	0.04	67381.38
	CRF4 (III)	115.65	0.43	66.00	0.23	27059.68
	Total	586.70	8.50	58.51	1.00	6803.88
2009	CRF 4B-E	387.21	8.06	55.33	0.72	4704.79
	CRF 4 (II)	9.52	0.01	151.56	0.04	70226.15
	CRF4 (III)	115.65	0.43	66.00	0.24	27059.68
	Total	512.38	8.50	62.65	1.00	5929.35
	CRF 4B-E	205.21	8.06	55.48	0.56	2446.44
2010	CRF 4 (II)	9.52	0.01	151.56	0.07	70226.15
	CRF4 (III)	115.23	0.43	66.00	0.37	26959.68
	Total	329.96	8.50	83.98	1.00	3782.74
	CRF 4B-E	318.55	8.06	45.70	0.61	3852.85
	CRF 4 (II)	10.75	0.01	164.05	0.07	79286.60
2011	CRF4 (III)	114.80	0.43	66.00	0.32	26859.68
	Total	444.10	8.50	68.20	1.00	5125.86
	CRF 4B-E	240.57	8.06	51.51	0.57	2885.18
	CRF 4 (II)	11.86	0.01	154.56	0.08	87522.61
	CRF4 (III)	114.38	0.43	66.00	0.35	26759.68
2012	Total	366.81	8.50	78.04	1.00	4216.34
	CRF 4B-E	61.89	8.06	120.10	0.44	667.96
	CRF 4 (II)	11.88	0.01	154.43	0.11	87664.71
	CRF4 (III)	115.74	0.43	66.00	0.45	27079.84
	Total	189.51	8.50	140.13	1.00	2130.02

6.3.7.1 Harvested wood products (4.G)

Harvested wood products (HWP) are reported for the first time in 2015 based on the domestic production approach outlined in the 2013 IPCC supplementary guidance for the Kyoto protocol. The approach adopted is broadly consistent with the 2006 GPG guidance for approach B1 but there are some differences because all harvest from deforested lands are immediately oxidised. This is done to ensure a consistent approach between Kyoto and convention reporting of HWP.

The primary activity data used for estimating HWP CSC is the EUROSTAT and FAO data from 1961 to 2013. The FAO/EUROSTAT data is used to calibrate the FORCARB and CARBWARE model harvests as described in section 6.3.2.4. The domestic harvest, imported and exported timber flows from 1961 to 2013 are shown in CRF Table 4Gs.2. The methods used to derive HWP for afforested (land converted to forests), land remaining forest domestically produced HWP is outlined in the following step below.

Derive sawnwood (SW), wood based panel (WBP), paper and paper board (PPB) HWP feed stock from FAO/EUROSTAT data using Eq 2.8.1 and 2.8.2 of the IPCC Supplementary guidance 2013. This uses the data produced in CRF Gs4 and f_{irw} and f_{pulp} ratios to derive the volume of SW, WBP and PPB (see Table 6.5.21).

- Volumes of the SW and WBP HWP from domestic harvest are converted to tC using default conversion factors. The aggregate value of 0.458 and 0.595 Mg/m³ is used for SW and WBP, respectively (Table 2.8.1 2013 IPCC supplement). A carbon fraction of 0.5 is used for SW and a C fraction of 0.454 for WBP (Table 2.8.1 of the 2013 IPCC supplement). The final inflows of different domestically produced HWP are shown in table 6.5.18.

- b) Harvest product data was further extrapolated back to 1900 using regression equations using exponential function for each wood product (WP_j):

$$WP_j = e^{0.015(year-1961) \times tC1961_j} \quad (6.3.39)$$

where *year* is the specific year before 1961 and $tC1961_j$ is the tC feedstock for the wood product *j* in 1961. Historic consumption rates from 1900-1960, using a growth rate of 1.15 per cent y^{-1} , were used to estimate emissions from products entering the system prior to 1961, as outlined in 2006 GPG for AFOLU.

- c) The estimation of the annual fraction of harvest originating from the different forest activities (i.e. forest remaining forest (FM), land converted to forest (AR) and deforested (D) harvest) are then derived using Eq. 2.8.3 in Ch 2 of the IPCC supplementary guidance. The input information for the different activities (*j*) are derived from harvest data shown in Tables 6.3.4, 6.3.12 and 6.3.17 in the sections above. All harvests for deforested land are assumed to be immediately oxidised, so CSC in the CRF under HWP are reported as IE under sub categories 4.C.2, 4.D.2, 4.E.2 or 4.F.2 as part of biomass losses.
- d) The estimation of the annual amount of HWP being produced from domestic harvest, which is related to the 3 different forest activities is then determined using Eq. 2.8.4 of the 2013 IPCC supplementary guidance.

Table 6.5.21 Annual inflows of sawnwood (SW), wood-based panels (WBP), paper and paper board (PPB) from domestic harvest.

Year	f_{IRW} Eq. 2.8.1	SW m^3	WBP m^3	SW tC	WBP tC	f_{pulp} Eq 2.8.2	PPB tC
1961	0.94	45317	20487	10378	5511	0.72	3535
1970	0.93	47635	112575	10908	30283	0.07	3591
1980	0.88	125589	62355	28760	16774	0.19	7797
1990	0.98	378570	235380	86693	63317	NO	0
1991	0.99	382438	237785	87578	63964	NO	0
1992	0.99	568477	243209	130181	65423	NO	0
1993	0.97	618230	239722	141575	64485	NO	0
1994	0.97	685479	256209	156975	68920	NO	0
1995	0.97	660025	327092	151146	87988	NO	0
1996	1.00	685958	433342	157084	116569	NO	0
1997	0.96	614850	517163	140801	139117	NO	0
1998	0.95	640854	626612	146755	168559	NO	0
1999	0.88	713252	667520	163335	179563	NO	0
2000	0.96	852697	715358	195268	192431	NO	0
2001	0.96	887805	704592	203307	189535	NO	0
2002	0.95	773286	662076	177082	178098	NO	0
2003	0.93	939125	778170	215060	209328	NO	0
2004	0.92	865597	775377	198222	208576	NO	0
2005	0.91	921476	794376	211018	213687	NO	0
2006	0.92	1005030	898464	230152	241687	NO	0
2007	0.90	984351	869180	225416	233809	NO	0
2008	0.86	596051	708934	136496	190703	NO	0
2009	0.91	705648	650035	161593	174859	NO	0
2010	0.95	731414	758484	167494	204032	NO	0
2011	0.95	726243	703852	166310	189336	NO	0
2012	0.92	715888	655557	163938	176345	NO	0
2013	0.90	741510	643795	169806	173181	NO	0

- a) The tier 2 first order decay model Forestry production and trade data from 1961-2011 from FAO, projected HWP inflows (see above) and historical growth for timber utilisation (see below) were used to estimate harvested wood product (HWP) emissions/removals in Ireland for 1900-2030 using a model based on 2006 IPCC Guidelines ; i.e. the Pingoud and Wagner 2006 model:

$$C_{i+1} = e^{-k} \times C_i + \left[\frac{(1-e^{-k})}{k} \right] \times Inflow_i \quad (6.3.40)$$

$$\Delta C_i = C_{i+1} - C_i \quad (6.3.41)$$

Where:

i = year

C_i = the carbon stock in the particular HWP category from a particular forest activity at the beginning of year i , kt C

k = decay constant of first-order decay for HWP category given in units yr^{-1} ($k = \ln(2)/\text{HL}$, where HL is half-life of the HWP pool in years (see below).

$Inflow_i$ = the inflow to the particular HWP category (HWP_j) during year i , kt C yr^{-1}

ΔC_i = carbon stock change of the HWP category during year i , kt C yr^{-1}

Default half-lives of two years for paper, 25 years for wood-based panels, and 35 years for saw[n] wood⁴ were used to estimate emissions resulting from products coming out of use.

The final HWP data for lands converted to forest land and land remaining forests is shown in Tables 6.3.22 and 6.3.23.

6.3.7.2 HWP uncertainties

Sources of uncertainties related to the FAO were considered to be 15 % because national data is based on a systematic survey (Table 6.5.24). The 2006 IPCC Guidelines provides no HWP (CH 12) category specific uncertainties for allocation into HWP categories, C conversion factors or product density conversion factors for biomass, so the same uncertainty was used for all HWP categories (Table 6.5.24).

The uncertainty associated with domestically produced HWP from category 4A1 (or FM) and 4A2 (AR, Table 6.5.25) were derived using the same approach adopted for other forest categories (see eq. 6.3.34 and 6.3.35) using sources of uncertainty shown in Table 6.5.24).

The base year removals were zero for HWP from category 4A2 (AR) because there were no activities prior to 2007 (Table 6.5.25). Therefore, the mean trend change for the category (expressed as a percentage) is undefined (divided by zero).

⁴ Product categories, half-lives and methodologies outlined in para 27, page 31 of FCCC/KP/AWG/2010/CRP.4/Rev.4

Table 6.5.22 Detailed inflows and CSC for different HWP's in the land converted to forest land (i.e. AR sub-category or 4A2). Note: Harvest in the AR category starts from 2007 onwards.

	inflow tC			inflow tC			Δstock tC			Δstock ktCO ₂	Solid wood	Kt C			Yrs
Year	SW	WBP	PPB	SW	WBP	PPB	SW	WBP	PPB	Total	Initial stock	Cgain	C loss	Net	Half-life
2007	6269.3	6502.7	NO	6207.6	6413.4	0.0	6207.6	6413.4	NO	-46.3	12.4	12.8	-0.2	12.6	29.9
2008	1325.5	1851.9	NO	7398.4	8064.5	0.0	1190.8	1651.1	NO	-10.4	15.2	3.2	-0.3	2.8	29.2
2009	14041.0	15193.7	NO	21156.2	22829.0	0.0	13757.8	14764.4	NO	-104.6	43.1	29.2	-0.7	28.5	29.8
2010	8947.7	10899.6	NO	29600.9	32954.6	0.0	8444.8	10125.6	NO	-68.1	61.3	19.8	-1.3	18.6	29.5
2011	10292.3	11717.3	NO	39211.5	43609.8	0.0	9610.6	10655.2	NO	-74.3	81.2	22.0	-1.7	20.3	29.7
2012	6400.0	6884.3	NO	44779.6	49207.0	0.0	5568.1	5597.3	NO	-40.9	92.1	13.3	-2.1	11.2	29.8
2013	10544.2	10753.8	NO	54342.1	58467.6	0.0	9562.4	9260.5	NO	-69.0	110.6	21.3	-2.5	18.8	30.0

Table 6.5.23 Detailed inflows and CSC for different HWP's in the forestland remaining forest land remaining (i.e. FM sub-category or 4A1).

	inflow tC			inflow tC			Δstock tC			Δstock ktCO ₂	Solid wood	ktC			Yrs
Year	SW	WBP	PPB	SW	WBP	PPB	SW	WBP	PPB	Total	Initial stock	Cgain	C loss	Net	Half-life
1990	86483.6	63164.7	NO	931852.3	702954.1	816.7	68707.1	44285.8	-338.3	-413.1	1603.5	149.6	-37.0	112.7	30.8
1991	87378.3	63818.2	NO	1000098.0	746673.4	577.5	68245.7	43719.3	-239.2	-409.7	1713.1	151.2	-39.5	111.7	30.8
1992	129928.9	65296.5	NO	1109137.4	790655.2	408.3	109039.4	43981.8	-169.1	-560.5	1862.9	195.2	-42.4	152.9	31.7
1993	141302.4	64361.1	NO	1227300.3	832511.8	288.7	118162.9	41856.6	-119.6	-586.3	2019.7	205.7	-45.8	159.9	31.9
1994	156697.3	68798.3	NO	1358389.6	877600.0	204.2	131089.3	45088.3	-84.6	-645.7	2192.3	225.5	-49.4	176.1	31.9
1995	146998.9	85573.7	NO	1477305.2	938000.3	144.4	118915.6	60400.3	-59.8	-657.3	2368.1	232.6	-53.3	179.3	31.3
1996	152918.9	113477.8	NO	1599750.8	1024269.7	102.1	122445.5	86269.4	-42.3	-765.1	2572.7	266.4	-57.7	208.7	30.7
1997	136837.9	135201.7	NO	1703872.5	1129605.6	72.2	104121.7	105335.9	-29.9	-767.9	2778.0	272.0	-62.6	209.4	30.0
1998	143119.2	164382.3	NO	1812171.9	1260840.8	51.0	108299.4	131235.2	-21.1	-878.2	3012.8	307.5	-68.0	239.5	29.7
1999	159490.4	175336.4	NO	1934557.8	1399291.1	36.1	122385.9	138450.3	-14.9	-956.3	3268.5	334.8	-74.0	260.8	29.8
2000	184358.9	181681.0	NO	2079167.6	1540212.9	25.5	144609.8	140921.8	-10.6	-1046.9	3548.4	366.0	-80.5	285.5	30.0
2001	191258.9	178302.9	NO	2227773.9	1673949.5	18.0	148606.3	133736.6	-7.5	-1035.2	3825.2	369.6	-87.2	282.3	30.2
2002	166857.5	167814.8	NO	2349304.7	1793685.1	12.8	121530.8	119735.5	-5.3	-884.6	4061.8	334.7	-93.4	241.3	30.0
2003	203011.5	197600.7	NO	2504250.8	1939523.2	9.0	154946.0	145838.1	-3.7	-1102.9	4356.6	400.6	-99.8	300.8	30.1
2004	186517.9	196261.3	NO	2639827.0	2080052.3	6.4	135576.2	140529.1	-2.6	-1012.4	4627.3	382.8	-106.7	276.1	29.9
2005	198963.2	201479.9	NO	2785067.6	2221885.6	4.5	145240.6	141833.3	-1.9	-1052.6	4908.8	400.4	-113.4	287.1	30.0
2006	157531.7	165426.9	NO	2886436.2	2324282.7	3.2	101368.6	102397.1	-1.3	-747.1	5108.5	323.0	-119.2	203.8	29.9
2007	187600.8	194585.8	NO	3015590.2	2452638.1	2.3	129154.1	128355.4	-0.9	-944.2	5361.0	382.2	-124.7	257.5	29.9
2008	120898.9	168912.1	NO	3076166.0	2552162.6	1.6	60575.7	99524.5	-0.7	-587.0	5518.0	289.8	-129.7	160.1	29.2
2009	133647.6	144619.4	NO	3148177.1	2625006.5	1.1	72011.1	72843.9	-0.5	-531.1	5660.0	278.3	-133.4	144.9	29.8
2010	152077.4	185252.7	NO	3237024.7	2735933.6	0.8	88847.6	110927.1	-0.3	-732.5	5855.8	337.3	-137.6	199.8	29.5
2011	149422.4	170110.7	NO	3321501.2	2828893.5	0.6	84476.5	92959.9	-0.2	-650.6	6029.8	319.5	-142.1	177.4	29.7
2012	151690.5	163170.0	NO	3406566.9	2912466.0	0.4	85065.7	83572.5	-0.2	-618.3	6195.1	314.9	-146.2	168.6	29.8
2013	158480.7	161630.9	NO	3496687.9	2992235.3	0.3	90121.1	79769.2	-0.1	-622.9	6361.7	320.1	-150.2	169.9	30.0

Table 6.5.24 Detailed inflows and CSC for different HWP categories from category 4A

Code	Component	Sub-category	% Uncertainty	Source
A	HWP categories	SW, WBP, Pulp	15.00	Pg 2.135 Section 2.8.3 Ch IPCC 2013 supplementary GPG
B	FAO data	All	15.00	Table 12.6 Ch 12 2006 IPCC Guidelines
C	C conversion factor from dry weight	All	10.00	Table 12.6 Ch 12 2006 IPCC Guidelines
D	Density	All HWP categories	25.00	Table 12.6 Ch 12 2006 IPCC Guidelines
	Combined uncertainty		34.28	

Table 6.5.25 Uncertainty of HWP estimates for categories 4A1 and 4A2

Year	Category	Year emission/reductions (kt CO ₂ eq)	Base year emission/reductions (kt CO ₂ eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
1990	CRF 4A.1	-413.07	-413.07	24.87	1.00	na
	CRF 4A.2					na
	Total	-413.07	-413.07	24.87	1.00	na
1991	CRF 4A.1	-409.66	-413.07	24.87	1.00	-0.82
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-409.66	-413.07	24.87	1.00	-0.82
1992	CRF 4A.1	-560.46	-413.07	26.37	1.00	35.68
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-560.46	-413.07	26.37	1.00	35.68
1993	CRF 4A.1	-586.30	-413.07	26.87	1.00	41.94
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-586.30	-413.07	26.87	1.00	41.94
1994	CRF 4A.1	-645.67	-413.07	26.99	1.00	56.31
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-645.67	-413.07	26.99	1.00	56.31
1995	CRF 4A.1	-657.27	-413.07	25.50	1.00	59.12
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-657.27	-413.07	25.50	1.00	59.12
1996	CRF 4A.1	-765.13	-413.07	24.60	1.00	85.23
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-765.13	-413.07	24.60	1.00	85.23
1997	CRF 4A.1	-767.90	-413.07	24.24	1.00	85.90
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-767.90	-413.07	24.24	1.00	85.90
1998	CRF 4A.1	-878.22	-413.07	24.35	1.00	112.61
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-878.22	-413.07	24.35	1.00	112.61
1999	CRF 4A.1	-956.34	-413.07	24.29	1.00	131.52
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-956.34	-413.07	24.29	1.00	131.52
2000	CRF 4A.1	-1046.91	-413.07	24.24	1.00	153.45
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-1,046.91	-413.07	24.24	1.00	153.45
2001	CRF 4A.1	-1035.23	-413.07	24.27	1.00	150.62
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-1,035.23	-413.07	24.27	1.00	150.62
2002	CRF 4A.1	-884.62	-413.07	24.24	1.00	114.16
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-884.62	-413.07	24.24	1.00	114.16
2003	CRF 4A.1	-1102.86	-413.07	24.25	1.00	166.99
	CRF 4A.2 NO		NO	NO	NO	NO
	Total	-1,102.86	-413.07	24.25	1.00	166.99
2004	CRF 4A.1	-1012.38	-413.07	24.24	1.00	145.09
	CRF 4A.2 NO		NO	NO	NO	NO

Year	Category	Year emission/reductions (kt CO2 eq)	Base year emission/reductions (kt CO2eq)	Combined uncertainty in year (±%)	Contribution to total variance in year (fraction)	Mean trend in year in relation to base-year (% mean trend)
Total		-1,012.38	-413.07	24.24	1.00	145.09
2005	CRF 4A.1	-1052.60	-413.07	24.24	1.00	154.82
	CRF 4A.2 NO	NO	NO	NO	NO	NO
Total		-1,052.60	-413.07	24.24	1.00	154.82
2006	CRF 4A.1	-747.14	-413.07	24.24	1.00	80.88
	CRF 4A.2 NO	NO	NO	NO	NO	NO
Total		-747.14	-413.07	24.24	1.00	80.88
2007	CRF 4A.1	-944.20	-413.07	24.24	0.95	128.58
	CRF 4A.2	-46.28	0.00	24.24	0.05	#DIV/0!
Total		-990.48	-413.07	23.13	1.00	139.79
2008	CRF 4A.1	-587.03	-413.07	24.95	0.98	42.12
	CRF 4A.2	-10.42	0.00	24.55	0.02	#DIV/0!
Total		-597.45	-413.07	24.51	1.00	44.64
2009	CRF 4A.1	-531.13	-413.07	24.24	0.81	28.58
	CRF 4A.2	-104.58	0.00	24.25	0.19	#DIV/0!
Total		-635.71	-413.07	20.64	1.00	53.90
2010	CRF 4A.1	-732.51	-413.07	24.39	0.92	77.33
	CRF 4A.2	-68.09	0.00	24.34	0.08	#DIV/0!
Total		-800.60	-413.07	22.41	1.00	93.82
2011	CRF 4A.1	-650.60	-413.07	24.24	0.90	57.50
	CRF 4A.2	-74.31	0.00	24.27	0.10	#DIV/0!
Total		-724.91	-413.07	21.92	1.00	75.49
2012	CRF 4A.1	-618.34	-413.07	24.24	0.94	49.69
	CRF 4A.2	-40.94	0.00	24.24	0.06	#DIV/0!
Total		-659.28	-413.07	22.78	1.00	59.61
2013	CRF 4A.1	-622.93	-413.07	24.28	0.90	50.81
	CRF 4A.2	-69.02	0.00	24.24	0.10	#DIV/0!
Total		-691.95	-413.07	21.99	1.00	67.51

6.4 Cropland (4.B)

6.4.1 Description

The Definition of Cropland includes “*all annual and perennial crops as well as temporary fallow land*”. This definition includes crops and temporary grassland managed as part of crop rotation systems. The definition also includes hedgerows associated with cropland systems.

Figure 6.4.1 shows the long term historic record in areas under crops since 1847. The historic data and more recent data are based on different methodologies, but common, underlying trends are evident.




Figure 6.4.1 Long term time series of areas under crops in Ireland since 1847

The analysis of cropland area has been revised significantly for the 2015 submission. Previously, the area of land associated with cropland was based solely on the annual total utilised agriculture area of crops reported by the CSO. Changes in cropland areas were based on the inter-annual variation in this reported area. The approach led to large inter-annual transitions between Cropland and Grassland land use categories. However, in the majority of cases, the transition was temporary and not a permanent transition to grassland.

Therefore the approach did not present a complete analysis of the role of temporary grasslands managed as part of a rotational cropland system. This was due to under recording of fallow / setaside areas, and a lack of analysis of the dynamic history of land management on individual field level. As such the previous assessment of cropland area did not fully represent those lands which would fall under the 2006 IPCC guidelines for Cropland category: *“Cropland includes all annual and perennial crops as well as temporary fallow land (i.e., land set at rest for one or several years before being cultivated again).”*⁵

The revised approach for the 2015 submission is based on detailed analysis of the Land Parcel Information System data, LPIS, collated annually by the Department of Agriculture Food and Marine. The LPIS is a description of all parcels of land covered under various agricultural and rural environmental administrative schemes, in Ireland, since 2000. This effectively covers all agricultural lands in Ireland. The system is subject to auditing, and provides robust and detail information on croplands. Although the LPIS was not designed to enable tracking of land use over time, careful post-processing and analysis of the data has demonstrated that the tracking of land use, at the resolution of individual parcels is possible with a high degree of consistency. Table 6.6.2 shows a number of examples of tracking of individual parcels of agricultural land use based on LPIS.

⁵ 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4, Chapter 5, Section 5.1

It is clear that cropland land parcels are managed in a wide variety of ways, from those which are recorded as under crops in all years, indicative of continuous cultivation, to those which have spent only short periods under crops.


Based on the analysis of LPIS, the definition of Cropland has been revised for reporting purposes as those lands which have been cultivated in the reporting year, and those which lands which are under temporary grassland, but have been recorded as having used to cultivate a crop at some time since 2000. Crops and temporary grasslands combine to an area of suitable lands which represent a stable cohort area of Cropland land use. No distinction is made between crop types, and it is assumed that the main factor influencing changes in long term carbon profile is period under grass.

The definition excludes permanent grasslands which have been managed exclusively for grazing (pasture) or grass fodder (silage and hay). However, the temporary grasslands included in Cropland continue to be considered actively managed in the reporting year, often for livestock.

The Central Statistics Office, CSO, provides annual statistics for Utilised Agricultural Area under various land uses, including a detailed breakdown on various crop types and grass management. However, the CSO data does not differentiate between permanent and temporary grasslands.

An analysis of historical areas under crops shows a significant decline in crops over a sustained period of decades. This is consistent with major changes in the agricultural economy and rural demographic in Ireland over several generations.

Figure 6.4.2 shows an analysis of changes in the spatial extent of croplands in Ireland over recent decades. It is clear that over time, cropland activities have consolidated into regions with suitable soils and benign climate.



use. Each line represents the start year in which the initial switch to grass occurred, and the proportion remaining as grass in each subsequent year. Although there is some variability in the return rates, there is clear convergence on all land parcels returning to crops within a 12 to 14 year period.




Figure 6.4.4 Analysis indicating low rates of conversion of Cropland to permanent Grassland

There is no evidence, in this analysis, of any permanent transition from cropland to grassland. This may reflect two important features of land ownership and land use in Ireland. Firstly, there is a low turnover of land sales in Ireland, with farms remaining in family ownership. Secondly, as a consequence, there is a high level of land rental and leasing on short term agreements. This means it is relatively easy for a tillage farmer to expand production area in response to projected market conditions and sentiment, without the need for major investment in land purchases. It also means that individual parcels on soils suitable for crops may remain under grass for longer periods, due to existing leasing arrangements with grass-based farmers.

Hedgerows are maintained as an integral system of cropland systems to protect crops against livestock incursion, and in many cases to define parcel boundaries. There is anecdotal evidence of hedgerow removal to facilitate access and traffic of machinery; however, recent hedgerow surveys across Ireland suggest the removal has not occurred on the same scale as other parts of Europe. Additional work is required to quantify change in hedgerow in Ireland, both in terms of extent and condition, and the EPA has funded a number of research initiatives on this topic. At present, a consistent time series of changes in hedgerow extent or condition is not available.

6.4.2 Soil Type and Soil Organic Carbon

For all non-forest land use categories, soil organic carbon (SOC) is the basic parameter in the default IPCC estimation methods for determining carbon stock changes in soils, which is a significant source of carbon emissions in land management and conversion categories in LULUCF. With the exception of forest soils, the organic carbon status of Irish soils under native vegetation is established from the soil type and the default reference soil organic carbon stocks (SOC_{ref}) for cold, temperate moist regions (Tables 2.3, 5.5 of the 2006 IPCC good practice guidance on LULUCF, and the relevant sections of the 2013 Wetlands Supplement). The Indicative Soils Map of Ireland (Fealy and Green, 2009, Teagasc, 2010) is the basic data source used in this analysis for soil type information in Ireland.

Mineral soils as identified from the soil map are allocated to the HAC (high activity clay), LAC (low activity clay), sandy and humic soil classes used by the IPCC, while drained peats are allocated to the IPCC wetlands class as shown on Table 6.6.1, based on detailed national assessment of soil carbon stocks in Ireland (Tomlinson, 2003). The values of SOC_{ref} appropriate to each soil association may then be assigned using the correspondence to IPCC classes given in Table 6.6.1. The distribution of CORINE Land Cover over IPCC soil classes was established in the same way to facilitate complete correspondence between land use, soil and SOC_{ref} .

Table 6.6.1 Soil Class Coverage and Soil Organic Carbon

General Soil Map Soil Association	Proportion of IPCC Soil Class					Proportion of Soil Association in Area of Ireland
	HAC	LAC	Peaty/ Humic	Sandy Soil	Wetlands Soil	
basin peat					0.34	0.06
brown earth		0.19				0.13
brown podzolic		0.21				0.15
Gley		0.30			0.02	0.22
grey brown podzolic		0.30				0.21
Lithosol			0.22	1.00		0.04
lowland blanket peat					0.31	0.05
Podzol			0.78			0.08
Renzinas	1.00					0.01
upland blanket peat					0.33	0.06
Proportion of IPCC Soil Class in Area of Ireland	0.01	0.71	0.10	0.01	0.17	
SOC_{ref} (t C/ha)	95	85	115	71	NA	

Choice of Methods

Ireland has adopted a Tier 1 approach to reporting greenhouse gas emissions associated with those areas defined as Cropland land use.

Activity data

The primary sources of activity data for Cropland used for the 2015 submission:

- Central Statistics Office annual statistics of Utilised Agriculture Area (1980-2013);
- Land Parcel Information System data (2000-2012) Maintained by the DAFM, the LPIS is integrated with the forestry, IFORIS data system;
- EPA/Teagasc Indicative Soil Map (2008);
- Activity within the Construction Sector from DEHLG (1990-2013);
- Fire Information for Resource Management System (FIRMS) NASA;
- National forest fire statistics see Table 6.5.6;
- Expert opinion ⁶ was sought on the use of histosols for crops in Ireland.

6.4.3 Cropland Areas

The area of cropland in a given year is the sum of the area of crops and the area of temporary grassland. The sum can be viewed as the areas of land whose current GHG emissions and removals are influenced by previous and current crop cultivation.

The area of land under crops in any given years is based on the CSO statistics.

⁶ Expert opinion was elicited from the delegates to a one day workshop:- "GHG fluxes in terrestrial ecosystems in Ireland - Grasslands, croplands, peatlands and forests " held on 20th September 2007
<http://www.ucc.ie/en/hydromet/celticflux/#Anchor1>

The CSO changed methodologies for estimation of area under crops in 1991. The methodology prior to 1991 was not consistent with the methodology used from 1991 to 2009. This results in a stepwise break point in the CSO between 1990 and 1991. A linear scaling adjustment has been applied to reconcile the 1990 estimate for crop areas with those reported under the 1991-2009 methodology.

In additional, the crop areas were revised by the inventory agency to account for inconsistencies due to the impact of changes in total farmed area reported in 1997, as described in the 2007 National Inventory Report, and reprinted in Section 6.13 Annex 3.4.

In 2010 the CSO adopted the LPIS as the primary source of data for estimation of agricultural land. Analysis of the time series indicates that this revision did not introduce a significant bias with the areas for both 2010-2013 consistent with previous trends and inter-annual variability within this sector. In the estimation of crops, and therefore there is no need to make adjustments to the Cropland time series data to take account for the 2010 methodological change.

In a given year, the area of temporary grassland is estimated as the difference between the CSO estimate of crops in that year, and the total cohort of lands used for cropland.

The LPIS has been used to provide estimates of the area of temporary grassland included under the definition of Cropland land use.

The analysis of the LPIS data from 2000 to 2012 provides robust identification of all parcels used for crops in this period. In total there are 118,957 parcels included in the Cropland cohort. Figure 6.4.5 shows the distribution of these lands. They are clearly concentrated within the specific geographic regions on a limited range of soil types and similar climatic conditions, as a result of the consolidation processes outline in Section 6.4.1.




Figure 6.4.5 Spatial distribution of all Cropland land parcels.

During the period 2000-2012, it is possible to explicitly identified parcels converted to Forest land use, as these parcels will either transfer to the IFORIS database, maintain a presence in the LPIS database as Forest, or both.

For the period 1990-1999, it is necessary to adjust the area of the cohort to accommodate the known incidences of conversion of Cropland to Forest land and Settlement. The details on the estimate of Cropland to Forest land conversion is given in Section 6.3. In order to meet this demand, the cohort of Cropland in 1990 must contain these lands in the first instance. Although the geographic location afforestation areas are known, the previous land use at each site can only be inferred. Therefore, during the period 1990-1999, the exact location of cropland to forest conversion cannot be inferred from existing analysis.

Analysis of the conversion of Cropland to Settlement has not been completed to the same level of spatial detail. The demand for settlement on croplands is currently based on estimates of activities in the construction sector which is disaggregated based on proportion of national land use in other lands uses. See section 6.7 for more detail on the attribution of previous land use for new Settlement. It is assumed; these lands will be excluded automatically from the LPIS and the CSO statistics, or assigned an appropriate attribute: e.g. farm building, dwelling, etc. Preliminary exploration of the LPIS data, postal service geodirectory data and contemporary aerial and satellite imagery largely confirm this assumption. However, additional analysis is required.

There is an important consequence of using this approach for Cropland, which includes all crop and temporary grassland land parcels identified within the 2000-2012 LPIS data, and extrapolating this area back to 1990. By definition, therefore, there has been no land converted between Cropland and Grassland land uses, and by corollary all agricultural grasslands within the Grassland land use category are defined as permanent grasslands.

There has been no evidence of deforestation to Cropland. Likewise, the analysis does not identify an instance of conversion of Wetlands, Settlement or Other Land to Cropland. Therefore, transition of land to Cropland does not occur.

All changes in emissions and removals are associated with short term transitions between crops and temporary grasslands, and reporting as occurring in the Cropland remaining Cropland land use category.

The analysis of the LPIS provides the history of each land parcel. However, it is not feasible to produce estimates of emissions and removals for each parcel. Therefore it is necessary to devise a consistent approach to summarise the spatial data. The crop types were aggregated into two broad classifications: Crop or Grass. For ease of analysis these were further codified into “0” for Crop and “1” for Grass. This allows the compression of the history of each parcel into a binary code. This allows for immediate grouping of parcels based on similar patterns of land use history. The total of 118,957 identified cropland parcels were condensed to 2199 management patterns. Table 6.6.1 shows some examples of this coding and grouping. Though quite successful in condensing the spatial data into a more manageable form, with 420 patterns accounting for 95 per cent of the total area. However, this is also considered to be too complex for routine analysis.

The next step was to consolidate the patterns further, based on the number of years of grass within each pattern, regardless of order in which this occurs. Therefore pattern B and I in Table 6.6.1 each have 8 years as grass, though in different sequences. Similarly, patterns E, F and G have 11 years as grass. This approach to consolidating the data is valid only where the default linear approach to transition from one management type to another is adopted. In this way, the cropland land parcels can be condensed to a set of 26 groups, based on the total number of years as grass, and whether the parcel is crop or temporary grassland in the given year.

Table 6.6.2 Examples of binary coding of cropland parcel history

Pattern Id	Code	Number of Parcels	Sum Area (ha)	Number of years of Grass
A	0000000000000	35897	159930.6	0
B	0000011111111	3558	11180.48	8
C	1101111111111	1511	7695.15	12
D	1100000000000	1431	5569.52	2
E	1111111100111	899	3159.8	11
F	1100111111111	840	3317.04	11
G	1110011111111	824	2880.86	11
H	0000000000100	211	636.19	1
I	1111100000111	127	416.92	8
J	0000011000000	93	295.82	2
K	0000000111100	83	300.05	4
L	1111000000001	83	299.76	5
M	0010000000000	81	341.9	1
N	0001011111111	79	373.34	8
O	0000101111111	77	474.12	7
P	0000001011111	75	210.82	6
Q	0011000000000	73	309.55	2

It is interesting to note the difference in histories of crop parcels and temporary grassland parcels, shown in Table 6.6.4. Not surprisingly, if in 2012 a parcel was identified under a crop, then it is more likely to have been under crops in previous years, and spent relatively less time under temporary grass, and vice versa. This reflects the situation that tillage farmers in Ireland will concentrate their efforts on the lands they own, and therefore these lands will spend more time under crops. While, temporary grassland will include a high proportion of lands which are rented or leased for crops on an ad hoc, demand driven, basis, and therefore are less intensively used for crops.

Table 6.6.3 Summary of Cropland parcel histories

2012	Years as Grass	Crops	Grass
13	0	61.0%	0.0%
12	1	4.1%	1.2%
11	2	3.8%	2.4%
10	3	3.1%	4.9%
9	4	2.9%	3.9%
8	5	2.4%	5.0%
7	6	2.7%	6.0%
6	7	3.2%	8.7%
5	8	4.9%	10.1%
4	9	2.4%	10.0%
3	10	2.2%	13.1%
2	11	4.4%	17.4%
1	12	3.0%	17.3%
		100.0%	100.0%

6.4.4 Soil

The LPIS and Indicative Soil Map were overlaid to provide an indication of the soil types associated with parcels within the Cropland cohort. The Indicative Soil Map was produced at a resolution of

1:250,000. As such, caution must be taken when attempting to assign parameters to the much high resolution LPIS data. Over 98 per cent of parcels associated with crops were associated with Low Activity Soils. The other 2 per cent were associated with High Activity Soils.

The large uncertainty associated with identification of crop land parcels on high activity soils prompted the decision that it is sufficient to estimate the change in carbon pools based on the assumption that all crop management occurs on low activity soils, without introducing significant bias in the estimation of emissions and removals. In previous submissions Croplands were assumed to revert to managed grassland during set-aside or fallow periods (the temporary exclusion of tillage areas from production) but stay within the category *4.B Croplands Remaining Croplands*, as a land parcel that is given over to set-aside in one year will likely be tilled in subsequent years. The CSO data included set-aside areas within what is termed “Other Crops”. However, with the inclusion of information based on the detailed analysis of the LPIS, and effective tracking of all cropland parcels, this assumption is no longer required. The analysis now provides parcel level information on the non-crop utilisation of parcels during fallow or setaside periods.




Figure 6.4.6 Time Series of Cropland from 1990-2013

Figure 6.4.6 shows the time series Cropland area from 1990 – 2013, split between area under crops in a given year, and temporary grassland. The total area decreases gradually over time, reflecting on-going conversion of Cropland to Forest Land and Settlement. Figure 6.4.1 shows analysis of the long term trend in croplands in Ireland over the last century shows a steady decline in tillage area (with exceptional temporary reversal related to WW1 and WW2), and a consolidation of activity on most the suitable soil types and local climate zones. From the graph, this long term trend appears to have achieved a steady state, and it is reasonable to assume that no lands were in transition to cropland at the beginning of the reporting period, 1990. Therefore, it is assumed, with the exception of land

conversion between Cropland and Forest land and Settlement, the Cropland cohort, identified from the analysis of the LPIS from 2000-2013, has been stable since 1990.

6.4.5 Carbon Stock Change in Biomass

Estimation of changes in above ground biomass is described below. It is assumed, by the Tier 1 methodology that below ground biomass remains constant if there is no change in long term management.

Annual Crops

Changes in above ground biomass are based on the areas transitioning between crops and temporary grassland in the given year. It is assumed there is no significant to change in below ground biomass.

The areas for crop to temporary grass and temporary grass to cropland are estimated separately, based on the actual proportion of parcels transitioning in either direction. This has been estimated on an annual basis for all years from 2000-2012 from the LPIS database.

As a first approximation, the values for the period 1990 to 1999 are estimated from the average proportions observed from the LPIS. On average 7.3 per cent of cropland in a given year was temporary grassland in the previous year. Similarly, on average 13.3 per cent of temporary grasslands were under crops in the previous year.

Analysis of the 2013 LPIS data is not complete, therefore transition for 2013 are based on analysis of the average values of the period 2000- 2012.

The areas from 1990 to 1999 are driven by net changes in the area of total crops, and an inverse correlation with temporary grasslands. In the period 2000-2012, the conversions are more variable, driven by external factors, leading to variability in the demand for crop areas and the rate at which temporary grasslands revert to crop.

The stock change relates only to above-ground biomass and its estimation is based on the difference between initial and final carbon content of biomass for the lands converted. In the conversion of land to cropland, it is assumed under the Tier 1 approach that the dominant vegetation from the initial land use is removed entirely. The carbon stock change is then quantified as the net sum of carbon lost on conversion and the carbon added by the first year's growth of crops. It is assumed that temporary grasslands are managed in the same manner as improved permanent grasslands. The dry matter content of grassland is taken as 13.6 tonnes/ha and the carbon content of dry matter is 0.5 per cent. The default value of 5 t dry matter/ha from Table 6.4 Vol 4. of the 2006 IPCC Guidelines is adopted for the carbon stock in crop biomass after one year. The carbon stock change in biomass on the area (A) converted to cropland is then calculated eq 6.4.2, derived from eq 2.15 from Ch 2 Vol 4 of the 2006 IPCC Guidelines as follows:

$$\Delta C = A * [(C_{\text{after}} - C_{\text{before}}) + DC_{\text{growth}}] \quad (6.4.2)$$

$$\Delta C = A * [(0.0 - 13.6.0 * 0.5) + 5.0]$$

Where A is the area of crops converted to temporary grassland.

Similarly, the inverse relationship is applied where the transition is from temporary grassland to annual crops.

Perennial Woody Crops

The area of some woody crops and bioenergy crops (including Christmas trees and miscanthus) are included in the CSO “Other Crops” category. However, this category is dominated in the period 1993 to 2007 by the additional reporting of fallow or setaside lands in this category.

The areas of fruit orchards is included in the CSO “Fruit” category, however, this category also includes soft, non-woody fruit plants, including the strawberry crop. The area under Fruit is in a long term decline, however, this appears to reflect the industry move to indoor, protective environments for the production of strawberry.

Therefore, the annual CSO statistics are not a suitable proxy for woody crops.

A self-consistent time series of activity within the Christmas tree sector is not available at the present time. Christmas trees are defined as a horticultural crop, and so included in the CSO annual statistics within the broader horticultural sector. A variety of sources of information have been explored, with some widely different estimates of the national plantation area. There is however, a high level of consistency in the market for Christmas trees. In 1997, O’Reilly et al produced a report for COFORD on opportunities within the Christmas tree sector which estimated a plantation area of 1,500ha to supply a market of 450,000 trees. In 2002, an All Island report from InterTrade Ireland⁷ published estimates from Bord Glas and Goodbody Economic Consultants of some 3,000 ha of plantations in 1998, falling to 2,428 in 2001. In 2006, the Teagasc Fact Sheet on Christmas Tree Production estimates between 300,000 and 500,000 trees were planted each year. Bord Bia⁸ currently estimates the market for Christmas trees from Irish producers to be between 500,00 and 700,000 plants. Typically trees are grown at a density of between 4,500 and 7500 plants per hectare. This implies an annual demand for the harvesting of between 68 and 144 ha from Christmas tree plantation. 2m trees are harvested at ages between 7 to 10 years. From this the total area of Christmas tree plantations can be estimated at between 475 ha and 1,440 ha, with an average of 960 ha. Therefore, Christmas tree plantations in Ireland may cover between 1,000 and 4,000 hectares, with a best estimate of 1,500 ha. The market is quite stable, with perhaps some evidence of growth in market since 1997. Therefore it is reasonable to assume that national Christmas tree plantation areas are in long term equilibrium with respect to total area and greenhouse gas emissions and removals.

In Ireland, the dominant commercial permanent woody fruit crop is apples. Annual statistics on area of apple orchards are not available. Census of Apple Orchards in Ireland data are available for years 1991, 1997, 2002, 2007 and 2012. There are an estimated to be about 45 specialised apple growers in Ireland. There was to have a significant decline in area under apple orchards during the 1990’s, however, the sector appears to have stabilised, with no significant trend in area under orchard since 1997. Annual areas for years between censuses data has been based on linear interpolation between data points. It is reasonable to assume the decline in orchards observed in the 1990s was initiated earlier. Therefore the areas for 1989 and 1990 area based on an extrapolation of the rate of decline between 1991 and 1997. Since there is no significant observed rate of change in area of orchards between 1997 and 2012, it is assumed there is no change area of orchards for 2013.

⁷<http://www.intertradeireland.com/media/A%20Review%20of%20the%20All-Island%20Horticulture%20Industry.pdf>

⁸<http://www.bordbia.ie/industry/manufacturers/insight/publications/MarketReviews/Documents/Export-Performance-and-Prospect-2015.pdf>




Figure 6.4.7 Estimated area under Apple Orchards 1989-2013

The estimate of biomass gains and losses associated with these transitions between perennial crops and other crop types is based on the Tier 1 approach described in Section 5.2.1.1, Vol 4 of the 2006 IPCC Guidelines.

Biomass in transition from perennial woody crop to annual crops is estimated using eq 2.15 from Ch 2 Vol 4 of the 2006 IPCC Guidelines

$$\Delta C_B = \Delta C_G + \Delta C_{conversion} - \Delta C_L \quad 6.4.3$$


Where ΔC_B = annual change in biomass stock in perennial crops

ΔC_G = annual increase in carbon stocks in biomass under perennial crops. This is assumed to be zero for well-established perennial crop areas. For lands in transition, the accumulation rate is given by the reference carbon stock (63 tC ha^{-1}) divided by the default 30 year period it is assumed it takes for woody crop to reach maturity/equilibrium.

$\Delta C_{conversion}$ = the initial decrease in biomass from perennial woody crops to annual crops. This is equal to the net change due to a loss of 63 tC ha^{-1} in the year of transition, from Table 5.1 Ch 5 Vol 4 of the 2006 IPCC Guidelines, and a gain in biomass due to subsequent growth of crops in the year of transition. This is equal 5.0 tC yr^{-1} , from Table 5.8 of Ch 5 Vol 4 of the 2006 IPCC Guidelines.

ΔC_L = is the annual loss due to harvesting, fuel wood gathering and disturbance. This is assumed to be zero for well-established perennial crops.

Figure 6.4.8 shows the time series of estimated carbon stock change in above ground biomass based on this analysis.



Negative is a source of emissions to the atmosphere

Figure 6.4.8 Carbon Stock change in Biomass of Perennial Crops

6.4.6 Cropland Dead Organic Matter/Litter

Tier 1 assumption is applied, with default estimation of zero emissions or removals associated with dead organic matter/ litter..

6.4.7 Carbon Stock Change in Soils

The spatial distribution of cropland areas over IPCC soil class is derived from GIS analysis of the LPIS 2004 dataset provided by the Department of Agriculture, superimposed on the Indicative Soils Map of Ireland (Fealy and Green, 2009). The GIS analysis shows that a very high proportion (98 per cent) of croplands are located on Low Activity Clay (LAC) soils. It is assumed that only grasslands on LAC soils are suitable for direct conversion to croplands, which is consistent with the requirement for cropland productivity. It is therefore reasonable to assume that all grassland areas converted to croplands are also on LAC soils and that no other land categories are converted to croplands. The research noted in 6.4.5 will also analyse the validity of this assumption.

6.4.8 Estimation of Emissions from Soils

Mineral Soils

The annual change in SOC in mineral soils over the transition period is based on the Tier 1 methodology, described in Section 2.3.3.1, of the 2006 IPCC Guidelines. From which the carbon emissions or removals for the various land-use conversion categories from equation 2.25 in the guidelines, as follows:

$$\Delta C = A * (SOC_0 - SOC_{0-T}) / T(6.4.1)$$

$$SOC = SOC_{ref} * F_{LU} * F_{MG} * F_I$$

where

- ΔC = annual change in carbon stocks
- A = area of land converted from a former land use
- SOC_0 = soil organic carbon stock for current land use
- SOC_{0-T} = soil organic carbon stock for former land use
- SOC_{ref} = reference soil organic carbon under native vegetation for a given soil type in area A
- T = transition period
- F_{LU} = stock change factor for land use or land-use change type
- F_{MG} = stock change factor for management regime
- F_I = stock change factor for organic matter input

The factors F_{LU} , F_{MG} and F_I account for changes in SOC due to management practices that impact on soil carbon. Table 6.6.4 shows the adjustment factors derived from the product of F_{LU} , F_{MG} and F_I taken from Table 2.3 in Ch 2 of the 2006 IPCC Guidelines for the land uses defined for Ireland (Table 6.3). Equation 6.4.1 is the basic Tier 1 methodology used for estimating emissions from mineral soils for all land-use categories as described in the following sections. The default transition period of 20 years is applied for all mineral soils. The estimation procedure is performed following a simple approach that provides estimates of emissions from soils for the defined land uses in accordance with the IPCC good practice guidance for LULUCF and the available information for the country. It involves the identification and quantification of the land areas subject to a change of use, the application of the data in Table 6.6.1 to assign SOC_{ref} for the soil types in those land areas and the calculation of carbon stock change on the basis of the factors given in Table 6.6.4.

Table 6.6.4 Adjustment Factors for SOC

Land Use	F_{LU}	F_{MG}	F_I	Adjustment factor, AF
Cropland	0.71	1.09	1.11	0.86
Improved grassland	1.0	1.0	1.14	1.14
Unimproved grassland	1.0	1.0	NA	1.0
Rough grazing	1.0	0.95	NA	0.95
Other non-agricultural land (Native grassland)	1.0	1.0	NA	1.0

Carbon stock changes in mineral soils are estimated using methodology outline in Section 6.4.3 and Equation 6.4.1. Farm management and input practices for crop and temporary grasslands are assumed to have been constant over the inventory period for lands within the cropland cohort. Therefore the SOC will change for mineral soils, only in response to variations in the period lands spend under temporary grasslands. In line with expert opinion it is assumed that no cultivation occurs on organic soils, as discussed in 6.4.2.

The approach taken is to estimate change in soil carbon stocks based on total area under crops and total area under temporary grassland.

In any given year the total carbon stock under crops is estimated from the weighted average of years spent as temporary grass as part of the national average rotation pattern.

Table 6.6.4 shows the average proportion of crop parcels (by area) spent as temporary grassland in previous years. It is interesting to note that a high proportion of crop parcels have a continuous history of management under crops. These are assumed to be in long term equilibrium with respect to soil carbon.

The default transition period for conversion of cropland to grassland is 20 years, over which it is assumed that the impact on soil carbon occurs. Because the change in carbon is assumed to be linear with time in the default model, it is consistent to assume all parcels with the same number of years as under temporary grassland will have the same averaged soil carbon stocks.

Table 6.6.2 shows the average carbon stocks for crop and temporary grass parcels based on years spent under grass. Clearly, the more years a parcel spends under a crop, the closer its carbon content is to the reference level for continuous cropping. Likewise, the more years a parcel spend under grass, the closer the carbon levels are to the reference content for permanent grassland.

Table 6.6.5 Carbon content of cropland soils as a function of period under grass

Carbon content of Soil (tC ha ⁻¹)		
Years	Crops Under Grass	Grass under Crop
0	73.0	102.8
1	73.1	102.8
2	73.3	102.5
3	73.7	102.2
4	74.2	101.6
5	74.9	101.0
6	75.7	100.2
7	76.7	99.2
8	77.8	98.1
9	79.1	96.8
10	80.5	95.4
11	82.0	93.8
12	83.8	92.1
13	85.6	90.2
14	87.6	88.2
15	89.8	86.1
16	92.1	83.8
17	94.6	81.3
18	97.2	78.7
19	99.9	75.9
20	102.8	73.0

6.4.9 Cropland emissions due to Biomass Burning

Activity data on the occurrence of fire on cropland is limited. The NASA FIRMS data set for region of Ireland was interrogated. It identified approximately 1.5 per cent instances of likely fire events coincide with cropland locations. There are significant limitations to the satellite product mainly related to the relatively low spatial resolution and high probability of cloud interference in any signal over Ireland.

The NASA FIRMS data was overlaid on spatial land cover and land use data including CORINE, and LPIS. From this a table of the probability of fire on each land use type was constructed. Fire events on Forest location account for 9.5 per cent of all fire detections. Figure 6.4.9 illustrates the strong correlation between the fires detected at forest location and those detected on other land uses. This provides confidence in use of Forest Fire statistics to estimate fires on other land uses.




Figure 6.4.9 Validation of use of Forest Fires as a scalable proxy for fire events on other Land Uses

Activity data for forest fire is described in section 6.3.4.4. These data are considered more robust for fire detection relative to the low resolution NASA FIRMS data, because the collation method is more likely to record incidences of relatively minor fires, difficult to detection in the satellite data. Figure 6.4.10 is a comparison of the time series of both satellite and Forest service data, and shows, although the remote sensing product under-records fires, there remains a strong correlation between the two data.




Figure 6.4.10 Validation of the use of Forest Service Statistics as a scalable proxy for all fires in a given year

It is assumed that the principle driver for fire across all land uses is the incidence of suitable weather conditions. Therefore, to a first approximation, the number of fires recorded on forest land is a fixed proportion on the total number of fires. Table 6.6.6 provides a summary of this analysis. Therefore, a simple scaling has been implemented to estimate the number of fires on grassland, based on the recorded number of fires on forest, and the relative proportion of fires detected in the satellite data which correspond to cropland.

Table 6.6.6 Land Cover/Use associated with NASA FIRMS fires

Land Use	Proportion of All Fires Detected
Forest	9.5%
Cropland	1.5%
Grassland	17.8%
Wetlands	71.0%
Settlement	0.3%
Other	0.0%

Meteorological conditions determine the suitable conditions for fire, however remote sensing cannot establish whether the actual fires are due to natural causes or direct human interventions. Although not prohibited by law, it is not practice to deploy controlled burning as a cropland management tool. Landowners are required to inform local authorities and fire services of their intention of initiating a controlled fire, however this information has not been collated at a national level.

Therefore, the incidence of fires on croplands is assumed to be as a result of an accidental fire outbreak. Therefore, all fires on cropland are classified as wildfire.

The emissions associated with fires are estimated based on the generic approach outlined in 2006 IPCC guidelines Vol 4, Section 2.4 and additional details provided in Vol 4 Ch 5, Section 5.2.4.2 for cropland remaining cropland.

Note, the Tier 1 approach assumes there are no long term losses of biomass carbon due to fires on cropland, and emissions are estimated for CH₄ and N₂O only.

Figure 6.4.11 shows the activity data is the estimate of cropland area burnt.




Figure 6.4.11 Estimate of Cropland area burnt by wildfire

Emissions of CH₄ and N₂O are calculated using eq 2.27 from Ch2 2 Vol 4 of the 2006 IPCC Guidelines, and shown here:

$$L_{fire} = A \cdot M_B \cdot C_f \cdot G_{ef} \cdot 10^{-3} \text{eq 6.4.4}$$

Where L_{fire} = amount of greenhouse gas emissions from fire, in tonnes of gas (CH₄, N₂O),

A = area burnt, ha,

M_B = mass of fuel available for combustion, tonnes ha⁻¹. This includes biomass, litter and dom. For Tier 1 Litter and DOM are assumed zero for croplands remaining croplands.

C_f = combustion factor, dimensionless,

The default value for $M_B \cdot C_f$ is 4.0 from Table 2.4 in the Ch 2 Vol 4 2006 IPCC Guidelines

G_{ef} = is the emission factor, g kg⁻¹ dry matter burnt. The default values for cropland are CH₄ = 2.7 g kg⁻¹ dm_{burnt}, N₂O = 0.07 g kg⁻¹ dm_{burnt}.

Figures 6.4.13 and 6.4.14 show the time series of estimated emission of CH₄ and N₂O due to biomass burning on croplands.




Figure 6.4.12 Estimated CH₄ emissions from wildfires on Croplands




Figure 6.4.13 Estimated N₂O emissions from wildfires on cropland

6.4.10 Uncertainties and time-series consistency in Cropland

Time Series Consistency

As indicated previously, there were three events identifies at which there was potential for time series inconsistencies to arise in the primary CSO utilised agricultural areas statistics. In 1991 and 2010 the CSO made documented changes to the methodology used to estimate crop areas. In 1997, there is a stepwise increase in reported crop area, which was not due to CSO methodological

changes. However, discussion with agricultural experts confirm that the introduction of the development phase of the LPIS data collection system in 1995 and 1996 would have changed the land owners perception of land parcel and utilised areas, particularly tillage farmers who were first to engage with LPIS.

Figure 6.4.1 shows the original crop area data from the CSO from 1980 to 2013, and the adjusted data produced by linear extrapolation of trends across the breakpoints.

The Land Parcel Information System is used to estimate the impact of short term land management practices and temporary transitions between crop and grassland. The LPIS data from 2000-2012 was used in the analysis presented in this submission Figure 6.4.6 shows crop and temporary grass areas for this period. There is large inter-annual variation, but no evidence of longer term trends in this period. This is consistent with the CSO data for crops, which show the period back to 1990 to have a similar trend. However, as noted in previous submission, the CSO data only captures net transitions between crop and grassland and back again.

Uncertainty in Area

The uncertainty in areas for the period 2000-2013 can be estimated from the requirements for submission of data to the DAFM under the various farm payment schemes associated with LPIS. The requirement for submitting data to the LPIS is for an accuracy of 0.1 ha for parcels. The mean parcel size is 4.3 ha. Therefore the average uncertainty for each parcel is of the order of 0.1/4.3, or 2.3 per cent.

Uncertainty in areas for the period 1990-1999 is more difficult to quantify. The uncertainty in areas is based on the uncertainty in the CSO methodology adoption during 1991-2012. This is estimated at approximately 2.0%. However, it is clear from the discussion on consistency in time series that in the gross dynamic is not captured, and consequently there is probably a systematic underestimate of the activity data underlying the estimation of N₂O emissions associated with mobilisation of carbon, i.e. the areas in transition from crop to temporary grassland. Net carbon emission and removals should not be biased.

Uncertainty in Emissions

The uncertainties associated with estimation of greenhouse gas emissions and removals due to activities under cropland land use are based on those appropriate to the adoption of the Tier 1 methodologies, land use and management factors and emission factors set out in the 2006 IPCC guidelines.

6.4.11 Category-specific QA/QC and verification

There are no QA/QC or verification procedures specific to the Cropland category

6.4.12 Cropland Category-specific recalculations and impact on emission trend

There have been significant changes made within the Cropland category. These relate to the incorporation of LPIS data into the analysis of areas of crop and temporary grassland. This has led to recalculation of emissions and removals for all years in the reporting period.

Figure 6.4.5.1 shows the difference between the 2014 and 2015 submissions for the period 1990-2012. Two assumptions which formed the basis for the trend in the previous submission have been addressed in the 2015 submission, based on the relationship between crop and temporary grassland identified in the analysis of the LPIS data.

In previous submissions it was assumed:

- A net increase in crop area report by the CSO was a conservative, permanent land use conversion, from Grassland.
- A net decrease in crop area, not immediately associated with afforestation or settlement, was a conversion to Grassland.

This led to the untenable situation that it appeared that new lands were being brought under crops on a continual basis, will the total area of Cropland was essentially constant.

Current analysis of the LPIS data indicates little or no evidence for long term conversion into or out of Cropland. However, there is short term rotation between crops and temporary grassland, which can impact on the biomass and soil carbon stocks. It is these short duration transition which are capture in the analysis in the LPIS and given rise to revised estimates of emissions and removals associated with Cropland land use category.

6.4.13 Cropland recalculations and impact on emission trend

Figure 6.4.14 shows a comparison between 2014 and 2015 submissions of estimated total emissions associated with Croplands. The difference is driven by the impact of the complete reassessment of the definition of Cropland and the inclusion of temporary grassland. Made possible by the detailed analysis of the LPIS spatial dataset, and represent a major step forward in the assessment of land use in this category. While the revised analysis is a more accurate assessment of impact of the management of croplands, it also reflects a high degree of inter-annual variability within the category. This variability requires additional analysis to ensure it does not result from an artefact in the transition to the revised methodology for assessing activity data (areas), but is reflective of actual potential emissions and removals.




Figure 6.4.14 Comparison between 2014 and 2015 submissions of estimated total emissions associated with Croplands

6.4.14 Cropland Category-specific planned improvements

A major project on Irish soils, the Irish Soils Information System, has recently published its data and produced a new, complete map of soils in Ireland. It is proposed to revise the attribution of soil type and soil carbon, and land use, based on this new, comprehensive dataset. It is hoped this will lead to the adoption of Tier 2 country specific value for reference soil carbon stock and management factors.

The 2015 submission is the first step towards incorporation of the Land Parcel Information System into the reporting methodologies. The stock change approach adopted in this first instance indicates large inter-annual variability in carbon stocks. This needs to be further assessed in the context of a relatively stable Cropland cohort area.

The current activity data for perennial woody crops is limited. Additional data discovery and analysis will be undertaken to improve these data.

The extrapolation of LPIS analysis to the period 1990-1999 presented in the 2015 submission is relatively simplistic. Further analysis will be undertaken, including exploration of pre-2000 agricultural spatial databases, to further refine understanding of land use within Cropland category during this period.

The EPA is funding research into the remote sensing technologies and analytical techniques for the quantification of non-forest wood biomass in the landscape. In the context of Cropland, this refers to primarily hedgerows. Hedgerows are an important feature of the Irish landscape. They are traditional means of establishing field and ownership boundaries and protecting crops from livestock incursion. In recent years, environmental payment schemes have included incentives for hedgerow plantation, maintenance and protection. The aim of the research is to exploration cost effective measurement and monitoring systems to quantify the impact of such policy incentives on biomass and carbon.

6.5 Grassland (4.C)

Definition: Grassland land use includes improved grasslands, unimproved grasslands and semi-natural grasslands. Improved grasslands include areas identified as lands managed for permanent identified as rough grazing for livestock. Semi-natural grasslands are those lands identified as dominated by grass habitats, but not currently managed for livestock. The hierarchy of land use identification is outlined in section 6.2.2.1. With this hierarchy, those lands identified as under grass, but with a recent history of crop management, are classified as temporary grassland within the Cropland land use category. All grasslands, including semi-natural grasslands are considered to be present as the result of direct management decisions. The definition of grasslands also includes hedgerows which are an integral part of livestock and land management practice in Ireland.

6.5.1 Grassland Areas

Grassland is the dominant land-use category in Ireland. Anthropogenic influence on grasslands is long standing and profound. There has been a long term trends towards livestock production in Ireland since the mid-1800s. The main driver was an increased demand for dairy and meat products from the industrial population centres in Britain. However, the trend also reflects a response to major changes in rural labour force, and a move to less labour intensive activities. Between 1850 and 1965 the number of cattle increased from approximately 2 million animals to 7.0 million. The reported areas of pasture and hay for this period did increased from approximately 3.5 million hectares to a maximum of 4.5 million hectares in the 1900s, and gradually dropping to 4.0 million again in the mid-

1990s. This points to a significant intensification the management and use of grassland through the 20th century leading to increased productivity.

In recent decades, changes in agriculture, have been driven by measures under the Common Agriculture Policy, where for example the “headage payment” subsidy lead to a very dramatic increase in sheep numbers from 3.5 million animals in the early 1980s to 8.9 million by 1990. This had a severe environmental impact due to over-grazing on hill sides. Reform of the scheme, in the mid-1990s, led to a sharp decline in sheep numbers, and the reported area of rough grazing. Similarly, production quotas on milk effectively led to the compression and stagnation of the dairy sector in Ireland

Choice of Methods

Ireland has adopted a Tier 1 approach to reporting greenhouse gas emissions associated with those area defined as Grassland land use.

Activity data

The primary sources of activity data for Grassland used for the 2015 submission:

- Central Statistics Office annual statistics of Utilised Agriculture Area (1980-2013);
- Land Parcel Information System data (2000-2012) Maintained by the Department of Agriculture, Food and the Marine, the LPIS is integrated with the Irish FORest Information System data (IFORIS);
- EPA/Teagasc Indicative Soil Map (Fealy and Green 2009);
- Activity within the Construction Sector from Department of Environment, Community and Local Government (1990-2013);
- Fire Information for Resource Management System (FIRMS) NASA;
- National forest fire statistics see Table 6.5.6;
- National Forest Inventory 2006 and 2012.

The estimate of the area of grasslands are based on CSO annual agriculture statistics for improved grassland (pastures and areas harvested for silage and hay) and unimproved grassland, which refers to rough grazing, and ancillary spatial data on semi-natural grasslands.

The definition of Grassland includes hedgerows and small wooded areas (non-forest), which are maintained as an integral component of livestock management and to establish field boundaries. Analysis of the National Forest Inventory for 2006 and 2012 includes estimates of hedgerows and non-forest wooded areas in the agricultural landscape. However, further research is required to complete a robust time series of hedgerow extent and condition in Ireland. Preliminary studies to this construct this time series have been funded by the EPA.

The CSO changed methodologies for estimation of area under grassland in 1991. The methodology prior to 1991 was not consistent with the methodology used from 1991 to 2009. This results in a stepwise break point in the CSO between 1990 and 1991. A linear scaling adjustment has been applied to reconcile the 1990 estimate for crop areas with those reported under the 1991-2009 methodology. The main impact of the change in methodology in 1991 was a reduction in the area reported under rough grazing, as commonage areas were excluded from the assessment.

In 2010 the CSO revised the methodology for the estimation of utilised agricultural land. The 2015 submission includes analysis of data from the CSO revised to include a hindcast to provide revised estimates for agricultural grasslands for all years from 2008. In the 2014 submission, the CSO had

only proved hindcast to 2010. At stated previously, the revised methodology did not introduce a discernible bias in the estimate on CSO estimates of crops. However, there is a significant impact on the CSO estimate of grassland areas within the improved grassland category, especially in the estimate of pasture. There is no corroborating evidence that the increase in improved grassland area reported by the CSO reflects a real change in land use. Additional analysis of the revised CSO methodology is required to determine the best means to achieve continuity between the new methodology and areas published in previous years. In the 2015 submission, it has been decided to extrapolate the long trend trends observed in the pasture area to estimated area for 2010 to 2013. Figure 6.5.1 shows the original data and the impact of the adjustment. It is worth noting, this change in methodology has increased the total area of agricultural grassland to levels similar to those recorded in the pre-1991, and the last major change in methodology. The research noted in section 6.4.2 will also analyse this issue.




Figure 6.5.1 Plot of original CSO data for Pasture areas and adjusted data for 2010, 2011 and 2012 based on extrapolation of long term trend from 1991 to 2009

It is important to note that both improved and unimproved grassland areas are estimates of grasslands *in use* for agricultural purposes. Rough grazing areas *in use* are native grasslands that are unmanaged with regard to drainage or other factors, such as fertiliser application, but which are grazed by cattle or sheep. The CSO annual statistics for rough grazing exclude other areas of grassland not reported to be in use for agricultural purposes. These grasslands are assumed to be semi-natural grasslands with limited human management interventions, in a carbon-stable state, with no associated emission or sink activity. However, they do represent a reserve of lands available for conversion to other land uses.

In previous submissions, natural grasslands were included in the Other Land land use category. In response to comments from the ERT, it is agreed that this was inconsistent with IPCC guidelines. Therefore, in the 2014 submission, natural grassland areas are included in the Grassland land use

category, with a baseline estimate for 1990 based on CORINE 1990 analysis of natural grassland classification types. When there is a demand for new agricultural grassland for use as rough grazing, it is met by a transition from natural grassland.

In the 2015 submission, a revised methodology has been applied to estimate the area of temporary grassland within the Cropland land use category. The CSO estimate of agricultural grassland does not include this sub-division of grasslands. Therefore, the area of improved grasslands reported by the CSO is adjusted to take account of these areas of temporary grasslands reportable the under Cropland category.

Overall, the area of grassland has decreased slightly, see Figure 6.5.2. The area of improved pasture has been near steady state, while the area of rough grazing, or unimproved grassland, has been decreasing. The dominant driver has been the conversion of grassland to Forest Land, and to a lesser extent, conversion to Settlement.

The area of natural grassland has increased. This is probably in response to government policy for hill farming, which in recent years has sought to decrease over grazing on vulnerable commonage and mountain areas, and other policies and market drivers which encourage consolidation of livestock activities and more intense management of grasslands. Therefore, there has been a decline in the reported agricultural land area, with conversion to Forest Land and “abandonment” to natural grassland the principle drivers of trends. The grazing of unimproved grasslands leads to degradation of the soil, with consequent emission of carbon.

From the data available, it is difficult to estimate the changes in managed area within the category *4.C.1 Grassland Remaining Grassland*. The annual CSO figures refer to the areas of land that farmers have declared to be “in use” under the specified types of use. Given the economic investment required to maintain “improved” grassland, it is probable that the declared “in use” areas are a good indicator of the actual extent of well-maintained managed grasslands. Therefore, significant changes in the improved grassland areas do represent changes in land use, with lands either being neglected, or actively managed, depending on the potential for good economic return. The neglect of improved grasslands will cause the land to revert to the nominally unmanaged or semi-natural native grassland state over time. The transition to rough grazing causes a degradation of the soil, leading to an emission of carbon. However, it is assumed that the average biomass remains constant. This is a potential underestimate of the effect of grazing, but insufficient analysis data exists to quantify the impact.




Figure 6.5.2 Trends in Grassland use 1990-2013

Hedgerows are maintained as an integral system of grassland systems to control the movement of livestock, manage grazing fodder stock, and in many cases to define parcel boundaries. There is anecdotal evidence of hedgerow removal to facilitate access, traffic of machinery and deploying alternative methods to management of grazing intensity. However, recent hedgerow surveys across Ireland suggest the removal has not occurred on the same scale as has occurred in other parts of Europe. Additional work is required to quantified change over time in hedgerows in Ireland, both in terms of extent and condition, and the EPA has funded a number of research initiatives on this topic. At present, a consistent time series of changes in hedgerow extent or condition is not available.

6.5.2 Methodological issues

6.5.2.1 Carbon Stock Changes in Grassland

The relevant carbon stock changes are for living biomass under 4.C.2 *Land Converted to Grassland* and for soils under both 4.C.1 *Grassland Remaining Grassland* and 4.C.2 *Land Converted to Grassland*.

6.5.2.2 Carbon Stock Changes in Living Biomass

The calculation steps for Tier 1 methodology are described in Section 6.2.1, it assumes that for grassland remaining grassland there is zero biomass carbon stock change. This approach is adopted here and the notation NO is entered in CRF Table 4.C. However, as Grassland is the dominant land use category for Ireland, the 2006 IPCC Guidelines suggest it is appropriate to develop national specific data to estimate the impact of management and disturbance. Ireland has funded a number of studies to achieve this, and the country specific analysis will be incorporated into the methodology as appropriate.

The category 4.C.2.5 *Other Land Converted to Grassland* is the most important conversion category in most years with small areas of land converted from forest and exhausted peat extraction sites. Carbon stock changes are estimated using the Tier 1 methodology. In the case of peatlands there is

no initial biomass at the time of conversion to grassland and therefore the carbon stock change is due only to the first year's growth at 6.0 t C/ha. The conversion of natural grassland for agricultural use is in effect the transition of semi-natural native grassland to improved or unimproved pasture, as indicated in section 6.5.1 above. There may be a change in carbon stock associated with conversion to improved grassland, as the land may be subject to ploughing and reseedling. This is accounted for through Equation 6.4.2. However, using the default values, a loss of 6.0 t C/ha for standing biomass followed by a gain of 6.0 t C/ha through growth in the first year⁹. Table 6.5.15 in section 6.3.6 above gives the area of forest land converted to grassland for the years 1990-2006. The immediate oxidation of biomass, litter and dead wood for years prior to 2006 were derived using the mean IEF for 2006 to 2009 (see section 6.3.3).

6.5.2.3 Dead Organic Matter/Litter

Tier 1 assumption is applied, with default estimation of zero emissions or removals associated with dead organic matter/ litter.

6.5.2.4 Carbon Stock Changes in Soils

The soil type distribution of grassland areas converted from other land uses over the IPCC soil classes is determined from GIS analysis of CORINE 1990 land cover data superimposed on the Indicative Soils Map (Fealy and Green, 2009). Mineral soils as identified from the general soil map were allocated to the five IPCC soil groups and their organic carbon status is established from the soil type and the default reference soil organic carbon stocks (Table 6.6.1). Table 6.6.4 shows the adjustment factors applied to the default SOC_{ref} to correct for land use and farming practice. The principal conversion affecting carbon stock change in soils is that from native grassland to rough grazing, which causes a decrease in soil carbon.

Organic Soils

Carbon emissions and removals are considered from two source activities:

- Drainage of organic soil under Grasslands;
- Rewetting of previously drained grassland.

Drainage of organic soil under Grasslands

A significant source of carbon emission is the use of organic soil types as pasture. It is assumed here that the wetlands soils under pasture are to some extent artificially drained, and so enables the emission of carbon from this organic soil type.

The Tier 1 methodology from the 2006 IPCC Guidelines, described in Section 2.3.3.1, and eq 2.26 in the guidelines (see eg 6.54.1 below) for estimating emissions from organic soils is to assign a direct annual carbon loss rate that accounts for the oxidation of organic matter due to drainage, tillage or disturbance of the land area concerned. For example, the default emission factors of 5.3 t C/ha per year for shallow drained managed grassland soils in cold temperate climatic regions given in the Ch 2 of the 2013 Wetlands Supplement to the 2006 IPCC Guidelines are adopted for Ireland. Tier 2 EF are used for forest organic based on country specific information.

$$L_{organic} = A \cdot EF \quad \text{eq 6.5.1}$$

⁹ There appears to be some inconsistency between default biomass carbon stocks given in Table 3.4.9 and those derived from Tables 3.4.2 and 3.4.3 of the IPCC good practice guidance on LULUCF. The inventory agency believes that the value of 13.6 tonnes DM/ha for the cold wet temperate climate zone should be 12 tonnes DM/ha.

Where L_{organic} is the annual carbon loss due to drainage

A is the area of grasslands on drained organic soils

EF is the emission factor for the template climate in t C yr^{-1} .

The adoption of the revised emission factor has a profound impact on the estimation of emissions associated with the agricultural use of grasslands. Figure 6.5.3 shows the time series of carbon losses from drained organic soils under agricultural grassland.




Figure 6.5.3 Time Series of carbon losses from drained organic soils under grassland

New estimates have been provided for forest conversion to grassland organic soils for the period 1990 to 2009 using the tier 1 emission factor of 0.25 t C per ha per year, (see CRF 4C and 4 KP-1 A2). Emissions from peaty mineral soils are adjusted according to peat depth as described in equation 6.3.2.

Rewetting of previously drained organic soils under grassland area

There has been a gradual decline in the total area of grasslands utilised for agriculture. It is inappropriate to classify the lands no longer identified in the agricultural statistics as “abandoned” lands, as this implies a release from ownership, and responsibility. Rather these lands are considered to revert to a semi-natural grassland status, which involves rewetting due to poor maintenance of drainage systems. The decision to allow an area of land rewet in this way, is a deliberate response to policy and market drivers, and is reversible.

The estimate of the area of rewetted grassland on organic soils is based on the assumption that a fixed proportion of the change in area of utilised grasslands is organic soil, which will revert to a wet status. Figure 6.5.4 shows the estimated areas of rewetted drained organic soils under grassland.




Figure 6.5.4 Area of Rewetted Organic Soils under Grassland

The Tier 1 methodologies for emissions and removals associated with rewetting of previously drained organic soils are presented in Ch 3 of the 2013 Wetlands Supplement to the 2006 IPCC Guidelines

There are three sources of emissions and removals associated with this rewetting:

1. CO₂ removals from the atmosphere due to uptake to soil;
2. CO₂ losses in the form of dissolved organic carbon into the water system;
3. CH₄ emissions due to rewetting.

$$CO_2 - C_{rewetted\ organic\ soil} = CO_2 - C_{composite} + CO_2 - C_{DOC} + L_{fire} - CO_2 - C$$

Where $CO_2 - C_{rewetted\ organic\ soil}$ = CO₂ as C removals and emissions from rewetted organic soils, tC yr⁻¹.

$CO_2 - C_{composite}$ = emissions and removals from soil and non-tree vegetation tC yr⁻¹

$CO_2 - C_{DOC}$ = emissions from dissolved organic carbon exported from rewetted organic soils, tC yr⁻¹

$L_{fire} - CO_2 - C$ = emissions from fires on these soils.

CO₂ removals from the atmosphere due to uptake to soil ($CO_2 - C_{composite}$)

From simplified version of Eq 3.4 in the 2013 Wetlands Supplement

$$CO_2 - C_{composite} = A \cdot EF_{CO_2}$$

Where A = area of nutrient poor rewetted organic soils, and EF_{CO₂} = emission factor for nutrient poor, soils in temperate climate zone and is equal to -0.23 tC ha⁻¹, from Table 3.1 in the 2013 Wetlands Supplement. The minus sign indicate a sink of carbon.

Figure 6.5.5 shows that time series estimate of carbon uptake to rewetted drained organic soils under grassland.




Figure 6.5.5 Carbon uptake to rewetted drained organic soils under grassland

CO₂ losses in the form of dissolved organic carbon into the water system.

A simplified version of Eq 3.5 from the 2013 Wetlands Supplements can be applied to Ireland.

$$CO_2 - C_{DOC} = A \cdot EF_{DOC \text{ rewetted}}$$

Where $CO_2 - C_{DOC}$ = off-site emissions of CO₂ from dissolved organic carbon exported from rewetted soils.

A is the area of nutrient poor rewetted organic soils, and $EF_{DOC \text{ rewetted}}$ = the emission factor for nutrient poor soils in a temperate climate zone, the default value from Table 3.2 in the Wetlands Supplement is approximately 0.24 t CO₂-C ha⁻¹ yr⁻¹.

Figure 6.5.6 shows that time series estimate of carbon loss due to DOC emissions associated with rewetting of drained organic soils under grassland.




Figure 6.5.6 Carbon losses due to DOC emissions associated with rewetting of drained organic soils under grassland

CH₄ emissions/removals due to rewetting

Rewetting or organic introducing anaerobic conditions to the soil, which leads to decomposition of organic matter with the release of CH₄. Eq 3.7 in the 2013 Wetlands Supplement provides the basis for Tier 1 approach to estimation of these emissions.

$$CH_4 - C_{rewetted\ org\ soil} = CH_4 - C_{soil} + L_{fire}$$

Where L_{fire} is the estimate of emission from fires, and is included in the Biomass Burning section 6.5.6

A simplified version of Eq 3.8 from the 2013 Wetlands Supplement can be applied to estimate emissions from the soils

$$CH_4 - C_{soil} = A_{rewetted\ org\ soil} \cdot EF_{NP,T}$$

Where A is the area of rewetted organic soils under grassland, and $EF_{NP,T}$ is the emissions factor for CH₄ emissions from nutrient poor soils in the temperate climate zone. The default value for $EF_{NP,T}$ is 92 kg CH₄-C ha⁻¹ yr⁻¹.

Figure 6.5.7 shows that time series estimate of methane emissions associated with rewetting of drained organic soils under grassland.




Figure 6.5.7 Time series of methane emissions associated with rewetting of drained organic soils under grassland

6.5.2.5 CH₄ emissions due to drainage of grasslands on organic soils

Section 2.2.2.1 of the 2013 Wetland Supplement to the 2006 IPCC Guidelines provides methodology for estimation of CH₄ emissions and removals from drained inland organic soils. The approach requires an estimate of the area impacted by drainage, and an estimate of the density of drainage ditches constructed and maintained to achieve this drainage. For Ireland, Eq 2.6 from the 2013 Wetlands Supplement can be simplified to the following

$$CH_{4_organic} = \left(A_{T, NP, O} \cdot \left((1 - Frac_{ditch}) \cdot EF_{CH_4_land} + Frac_{ditch} \cdot EF_{CH_4_ditch} \right) \right) \text{ eq 6.5.5}$$

Where CH₄_{organic} = emissions of methane due to the drainage of peatland under grassland.

A_{T, NP, O} = Area of nutrient poor, drained organic soils, in Ireland's temperate climate zone.

EF_{CH₄_land} = emission factor for methane emissions from nutrient poor soils serviced by drainage ditches in temperate zone. The default value for EF_{CH₄_land} is 1.8 kg CH₄ ha⁻¹ yr⁻¹ from table 2.3 of the 2013 Wetlands Supplement for shallow drained soils, which is typical drainage for Ireland.

EF_{CH₄_ditch} = emission factor for methane emissions from ditches in temperate zone, draining nutrient poor soils. The default value for EF_{CH₄_land} is 527 kg CH₄ ha⁻¹ yr⁻¹ from table 2.4 of the 2013 Wetlands Supplement, for shallow drained soils, which is typical drainage for Ireland.

Frac_{ditch} = Fraction of total area of drained organic soil which is occupied by ditches. The default value suggested in the 2013 Wetland supplement is Frac_{ditch} = 0.05.

Figure 6.5.8 shows the time series of estimated area of nutrient poor grassland drained for use as agricultural grassland.

Figure 6.5.9 shows the time series of estimated methane emissions due to the drainage infrastructure maintained to facilitate poor grassland use as agricultural grassland.




Figure 6.5.8 shows the estimate of CH₄ emissions from this analysis.




Figure 6.5.9 Estimate of methane emissions due to drainage of organic soils under grasslands

6.5.3 Land converted to Grassland

In the period 1990-2013, limited areas of Forest land and Wetland have been converted to Grassland.

6.5.3.1 Forest Land converted to Grassland

For details of the analysis of greenhouse gas emissions and removals associated with deforestation and conversion to grassland land use see section 6.3.6.1.2.

6.5.3.2 Wetland converted to Grassland

BnM has supplied data related the after use of sites under their management which have been exhausted for industrial peat extraction and converted to various other land uses. Although industrial peat extraction is no longer viable on these sites, there remains a significant depth of peat, and these lands remain classified as drained organic soils. In most instances, the after use of peat extraction sites involve rewetting or restoration of wetland function. However, BnM also converted some lands to agricultural grassland, see Figure 6.5.10, Emissions associated with this land use conversion are considered here.




Figure 6.5.1 Time Series of conversion of Peat Extraction Sites to Grassland

Biomass emissions and removals Wetland to Grassland conversion

Industrial extraction sites are completed denuded of vegetation, therefore the living biomass is zero before conversion to grassland, It is assumed that the grass is established to reference level of 6 tC ha⁻¹ living biomass for grass during the year of conversion. Therefore biomass increase is equal to 6 tC ha⁻¹ in the year of conversion. Figure 6.5.11 shows the estimate of biomass increase over time series.




Figure 6.5.11 Change in carbon pools due to the conversion of peat extraction sites to Grassland on drained organic soils. Note negative values indicate a loss of carbon to the atmosphere.

DOM and Litter emissions and removals Wetland to Grassland conversion

DOM and Litter are indistinguishable from the organic matter in organic soils. Therefore it is assumed to be included in the assessment of carbon emissions and removals estimated for soils.

Soil emissions and removals Carbon Wetland to Grassland conversion

In the conversion of peat extraction sites to grassland it is assumed that drainage of the site is maintained. Therefore carbon losses due to drained of organic soil continue. The conversion to grassland increases the default emission factor from 2.8 tC ha⁻¹ yr⁻¹, for temperate, peat extraction land use, to 5.3 tC ha⁻¹ yr⁻¹, for temperate, nutrient poor, grassland, see Table 2.1 in the 2013 Wetland Supplement of the 2006 IPCC Guidelines. Figure 6.5.11 shows the estimate of carbon loss to the atmosphere over time series.

6.5.4 Grassland emissions due to Biomass Burning

Activity data on the occurrence of fire on grassland is limited. The NASA FIRMS data set for region of Ireland was interrogated. It identified approximately 18 per cent instances of likely fire events coincided with grassland locations. There are significant limitations to the satellite product mainly

related to the relatively low spatial resolution and high probability of cloud interference in any signal over Ireland.

From this a table of the probability of fire on each land use type was constructed. Fire events on Forest location account for 9.5 per cent of all fire detections. See Section 6.4.9 for more detailed discussion of the analysis of satellite and Forest Service data as a proxy for estimation of fires on other land uses.

Also the very small proportion of fires detected on Settlement and Other Land use categories are aggregated into the Grassland proportion.

The NASA FIRMS data was overlaid on spatial land cover and land use data including CORINE, and LPIS. From this a table of the probability of fire on each land use type was constructed. The analysis suggests a very high proportion of fires are on peatlands. Although peatland fires are a feature of natural fire activity in Ireland, the land cover data has difficulty in distinguishing natural vegetation on peatlands and rough grazing, especially on blanket bog. Therefore, an ad hoc decision has been made to classify 50 per cent of peatland fires as actually occurring to manage rough grazing, and therefore included in the grassland fire area. Therefore the incidence of grassland fires increases to 53 per cent of all fires.

Activity data for forest fire is described in section 6.3.4.4. These data are considered more robust for fire detection relative to the low resolution FIRMS data, because the collation method is more likely to record incidences of relatively minor fires, difficult to detection in the satellite data. Comparison of the time series of both satellite and Forest service data, it can be seen than, although the remote sensing product under-records fires, there remains a strong correlation between the two data.

It is assumed that the principle driver for fire across all land uses is the incidence of suitable weather conditions. Therefore, to a first approximation, the number of fires recorded on forest land is a fixed proportion on the total number of fires. Therefore, a simple scaling has been implemented to estimate the number of fires on grassland, based on the recorded number of fires on forest, and the relative proportion of fires detected in the satellite data which correspond to cropland.

Although meteorological conditions provide suitable conditions for fire, remote sensing cannot establish whether the actual fires are due to natural causes or direct human interventions. Although not encouraged, controlled burning is deployed, in limited circumstances, as a grassland management tool, particularly in the control of low level scrub vegetation on poor or inaccessible grasslands. Landowners are required to inform local authorities and fire services of their intention of initiating a controlled fire, however this information has not been collated at a national level.

Therefore, the incidence of fires on grasslands is assumed to be as a result of land management. Therefore, all fires on grassland are classified as controlled fire. Figure 6.5.12 shows the estimated area of grassland burned.

The emissions associated with fires are estimated based on the 2006 IPCC guidelines Vol 4 Chp 6, Section 6.2.4 2.




Figure 6.5.12 The estimated area of grassland subjected to controlled burning from 1990-2013

Emissions of CH₄ and N₂O are calculated using eq 2.27 from Ch2 2 Vol 4 of the 2006 IPCC Guidelines, and shown below. The approach requires the area of grassland burnt to be stratified between mineral soils, drained organic soils and undrained organic soils, and provide appropriate default values for each of the parameters.

$$L_{fire} = A \cdot M_B \cdot C_f \cdot G_{ef} \cdot 10^{-3} \text{ eq 6.5.4}$$

Where L_{fire} = amount of greenhouse gas emissions from fire, in tonnes of gas (CH₄, N₂O),

A = area burnt, ha,

M_B = mass of fuel available for combustion, tonnes ha⁻¹. This includes biomass, litter and dom. For Tier 1 Litter and DOM are assumed zero for croplands remaining croplands.

C_f = combustion factor, dimensionless,

The default values for grassland for M_B are from Table 2.4 in the Ch 2 Vol 4 2006 IPCC Guidelines, and Table 2.7 in the 2013 Wetlands Supplement. See Table 6.6.7 C_f is assumed equal to 1.0, that is all available fuel is burned.

G_{ef} = is the emission factor, g kg⁻¹ dry matter burnt. The default values for grassland are CH₄=2.3 g kg⁻¹ dm_{burnt}, N₂O=0.21 g kg⁻¹ dm_{burnt}.

Table 6.6.7 Tier 1 Default parameters for use in Eq 6.5.4

	Mineral Soil	Drained Organic soil	Undrained Organic soil	Comment
$M_B \cdot C_f$ (t ha ⁻¹)	4.1	336	66	
$G_{ef} \text{ CO}_2$ (g kg ⁻¹ dm _{burnt})	0	362	362	It can be assumed there is no long term loss of biomass
$G_{ef} \text{ CH}_4$ (g kg ⁻¹ dm _{burnt})	2.3	9	9	
$G_{ef} \text{ N}_2\text{O}$ (g kg ⁻¹ dm _{burnt})	0.21	0	0	Assumes that N ₂ O emissions from drained organic soils is negligible

The Tier 1 approximation, there are no long term losses of biomass carbon due to fires on grassland, and emissions are estimated for CH₄ and N₂O only, see Figure 6.5.13 and 6.5.14.




Figure 6.5.13 Estimated CH₄ emissions from wildfires on Grassland




Figure 6.5.14 Estimated N₂O emissions from wildfires on Grassland

6.5.5 Uncertainties and time-series consistency

The dominant uncertainty in relation to Grassland is the history and impact of changes in land management on greenhouse gas emissions. Analysis of the archive of LPIS data for permanent grasslands is not as advanced as that for Cropland (and temporary grassland). The allocation of soil types under grasslands is based on extrapolation of the analysis of a single point in time. Therefore, where there are changes reported in the area of grassland, either through the demand from other land uses, or changes in management, it has been assumed these impact proportionately across all

soil types. For example, if there is demand for 1000ha to transition between improved and unimproved grassland, and 10 per cent of grasslands are on drained organic soils, then 100ha of the conversion is assumed to be related to these drained organic soils. It is likely that that soil type itself is a major influence on the type of changes in land use and management which can occur. It is not possible to provide a quantitative assessment of the uncertainty associated with the current assumption. This will be subject of future analysis and development of the inventory.

A separate, but related, source of uncertainty relates to the management of grasslands which are no longer reported as in use for agriculture. In the 2015 submission, it was assumed that un-utilised grasslands revert to a semi-natural state, in particular that the drainage systems for organic soils fail and rewetted conditions rapidly ensue. The validity of this assumption will be the subject of future analysis. Therefore it is reasonable to assign a large uncertainty to this component of the analysis, as this may lead to an overestimation of the impact of changes in management on these lands. Much of the net-net change in losses from grasslands, between 1990 and 2013, is due to the finding here that the area of drained organic soil under grassland has decreased in this period. The validity of this finding needs to be verified for future submissions.

As discussed above, the time series for agricultural grasslands have been adjusted in response to three discontinuities in the CSO data. The 1990/1991 and 2009/2010 discontinuities have been implemented in response to known changes in methodology. The 1996/1997 discontinuity is in response to a unrealistic stepwise change in areas based on the survey returns, but with no change in methodology. Anecdote evidence points to changes in the knowledge base of those filling in the survey, and improved on-farm information on actual field sizes and areas farmed.

The solution to the discontinuity arising from the most recent change in CSO methodology is not tenable in the long term, as it makes adjustments to the more recent data based on historic trends. However, the revised methodology is accepted to be a more complete approach to the estimation of utilised grassland. It is also worth noting that the revised methodology results in grassland areas more in line with pre-1991 estimates. This will be subject of future analysis and development of the inventory, to devise a more robust method to assure consistency across the time series.

6.5.6 Category-specific QA/QC and verification

There are no QA/QC or verification procedures specific to the Grassland category.

6.5.7 Grassland -specific recalculations and impact on emission trend

There have been significant changes made within the Grassland category. These relate to the incorporation of LPIS data into the analysis of areas of crop and temporary grassland, and the adoption of the revised emission factor associated with drained of organic soils provide in the 2013 Wetlands Supplement.

The adoption of the revised emission factors for drained organic soils under Grassland has increased the estimated loss of carbon from these soils very significantly. The emission factor has increase from 0.25 tC ha^{-1} to 5.3 tC ha^{-1} , for shallow drained grassland, i.e. by a factor of over 50 times. This elevates drainage of organic soils under grassland to key category status. The impact on the trend of carbon emissions and removals is also profound. Relative to 1990, the area of drained organic soils in this category has decreased.

This incorporation of LPIS data into the analysis has led to recalculation of emissions and removals for all years in the reporting period. The area of temporary grassland within the Cropland cohort, have been subtracted from the total agricultural Grassland areas.

Figure 6.5.15 shows the difference between the 2014 and 2015 submissions for the period 1990-2012.




Figure 6.5.15 Grassland recalculations between the 2014 and 2015 submissions for the period 1990-2012

Two assumptions which formed the basis for the trend in the previous submission have been addressed in the 2015 submission, based on the relationship between crop and temporary grassland identified in the analysis of the LPIS data.

In previous submissions it was assumed:

- A net increase in crop area report by the CSO was a conservative, permanent land use conversion, from Grassland;
- A net decrease in crop area, not immediately associated with afforestation or settlement, was a conversion to Grassland.

This led to the untenable situation that it appeared that new lands were being lost to cropland on a continual basis.

Current analysis of the LPIS data indicates little or no evidence for long term conversion between cropland and permanent grassland. However, there is short term rotation between crops and temporary grassland. These short duration transitions which are captured in the analysis in the LPIS and give rise to revised estimates of emissions and removals associated with Cropland land use category, and are no longer reported under Grassland.

A third assumption, that where organic soils are un-utilised they revert to a rewetted state, has not been refined in the 2015 assessment. This may lead to an overestimation of the impact of changes in management on these lands. Much of the net-net change in losses from grasslands, between 1990 and 2013, is due to the finding here that the area of drained organic soil under grassland has decreased in this period. The validity of this finding needs to be verified for future submissions.

6.5.8 Grassland Category-specific planned improvements

A major project on Irish soils, the Irish Soils Information System, has recently published its data and produced a new, complete map of soils in Ireland. It is proposed to revise the attribution of soil type and soil carbon, and land use, based on this new, comprehensive dataset. It is hoped this will lead to the adoption of Tier 2 country specific value for reference soil carbon stock and management factors.

The 2015 submission is the first step towards incorporation of the Land Parcel Information System into the reporting methodologies. Next steps will include analysis of history of permanent grasslands, in particular assessing changes in condition of grassland no longer reported within the LPIS.

A revision is required in the procedure adopted to adjust for methodological changes in the annual CSO agricultural statistics.

The EPA is funding research into the remote sensing technologies and analytical techniques for the quantification of non-forest wood biomass in the landscape. In the context of grassland, this refers to primarily hedgerows. Hedgerows are an important feature of the Irish landscape. They are a traditional means of establishing field and ownership boundaries and protecting crops from livestock incursion. In recent years, environmental payment schemes have included incentives for hedgerow plantation, maintenance and protection. The aim of the research is to exploration cost effective measurement and monitoring systems to quantify the impact of such policy incentives on biomass and carbon.

The emission and management factors associated with the drainage of organic and wet soils in Ireland will be assessed, based on the findings of country specific research.

6.6 Wetlands (4.D)

6.6.1 Wetland Areas

The term Wetlands as applied to Ireland refer to natural unmanaged wetlands and exploited peatlands, which are those wetland areas drained for the purpose of commercial exploitation and harvesting of peat. The national wetland area is therefore split into two types, unmanaged wetland and managed peatland (Table 6.4).

In this analysis, Wetlands in Ireland are characterised as either natural unmanaged wetlands or managed peatlands. Managed peatlands are those wetland areas drained for the purpose of commercial and domestic exploitation and harvesting of peat for use in energy or horticulture. The national wetland area is therefore split into two types, unmanaged wetland and managed peatland (Table 6. 4). In the 2013 submission, *4.D.1 Wetlands Remaining Wetlands* have been subdivided into two categories: Exploited Wetlands Remaining Wetlands and Unexploited Wetlands Remaining Wetlands.

The activities of enterprises engaged in the drainage of peat for extraction for horticulture use have been estimated for the first time. The estimate is based on analysis of export and domestic sales and extrapolation of the area of peatland required to meet this demand.

A limited area of forest land on peat have been deforested and rewetted as part of a wetland restoration project supported by the EU LIFE programme.

6.6.2 Unmanaged Wetland Areas

The initial 1990 unmanaged wetland area is based on the total area of peatland (excluding exploited areas) estimated from the CORINE 1990 land cover map. The main land use change out of unmanaged wetlands has been demand from forestry.

A small area of land is reported as converted to wetland due to the deforestation. These are included in the unmanaged wetland area, to differentiated the managed from peat extraction, and to reflect the land use intent of the action, which is to restore natural ecosystem function. It is assumed natural regeneration of biomass occurs over a period of five years to a maximum biomass of 3tC ha^{-1} , (see Table 6.5.15 section 6.6.2). It is also assumed soil carbon loss from these rewetted lands has ceased.

6.6.3 Exploited Peatland Areas

The commercial exploitation of peatlands is dominated by Bord na Mona (the Irish Peat Board) according to the land-use definition in Table 6.3. Commercial extraction proceeds in three separate stages, all of which may lead to changes in carbon stocks. Drainage is the first management activity, followed after several years by removal of the top layers of plant growth in the first season of peat extraction and then by the industrial extraction and harvesting of a layer of 10 to 15 cm of peat annually. The average working life of commercially developed Irish peatland is of the order of 30-50 years. Conversion to grasslands or forest land has been the historically favoured use of cutaway peatland. However, in recent years wetland reclamation has been investigated, and achieved with some success. The areas reported under category *5.D.1 Wetlands Remaining Wetlands* refer to all lands drained, whether the peat remains covered by vegetation or is exposed. Bord na Mona manages its peat reserves to meet present demand and is therefore progressing to extract peat from new sites only when an older field is exhausted. It is assumed that the decrease in reserves of peatland indicate new extraction areas, and therefore they are an estimate of the area from which biomass has been removed. Until recently, Bord na Mona held a small area of un-drained wetlands in reserved. However, these lands have been transferred to the National Parks and Wildlife Service for conservation.

Also, a small area of exploited wetland has been restored to ecosystem function through drainage management and rewetting. This has led to an uptake of carbon in the revitalised biomass.

Bord na Mona supplies the area estimates for the company's commercial peat harvesting activities. The data is commercially sensitive, and therefore not presented in this report. For the period from 1990 to 2012, the data for Bord na Mona commercial peat extraction areas are given as totals for consecutive five-year periods for a variety of peatland categories. Thereafter, BnM has provided annual statistics. The annual average value obtained from this total is used for each of the five years periods to obtain the full time series. Domestic harvesting of peat bogs by private landowners for their own household use is a strong tradition in many parts of Ireland, and although well documented in a social and cultural context, the amount of such peat extraction is poorly quantified. Previously estimates of the land area devoted to private harvesting of peat was estimated to be in the region of 400 ha per year based on the assumption that the area under private commercial and domestic use was of the order of one eighth that of Bord na Mona lands. The extraction of peat for use for residential heating is estimated from the national energy balance and a bulk density estimate of 0.25 t/m^3 for peat m^{-3} (McGoff et al. 2007). This approach ensures consistency between the quantities of peat combusted in *1.A.4.b Residential* and the area of private peat exploitation in LULUCF.

The 2015 submission includes an analysis of GHG emissions and removals associated with peat extraction and use for Horticultural use. BnM is also the dominantly player in this market, and most of the area of peatland exploited for extraction of horticultural peat is included in the areas supplied by BnM. However, an additional area of drained peat is included in the estimate to take account of non-BnM commercial activities. This estimate is based on market share and assumes the other commercial enterprises adopt similar management practices as BnM.

6.6.4 Carbon Stock Changes in Wetland

6.6.4.1 Biomass

Carbon stock changes in biomass are determined by the balance between carbon loss due to the removal of vegetation on preparation for peat harvesting and gain on areas of restored peatland. These changes have been estimated on the basis that the entire cover of vegetation is removed to prepare for peat harvesting and that an equivalent amount of biomass is returned on restoration of cutaway areas. In the 2006 NIR, it was assumed that the restoration of biomass occurred in the year of conversion. However, discussions with experts from Bord na Mona suggest a more appropriate biomass transition period of 5 years.

The area from which vegetation is removed is taken to be the amount of peatland reserve that is drained to come under production annually and the restoration area is taken as the annual increase in cutaway wetland given by Table 6.6.1. The vegetation is typically heather-dominated bog or heathland cover for which a biomass carbon content of 3 t C/ha is adopted (Cruickshank et al, 2000).

Table 6.5.15 in section 6.3.7 above gives the area of forest land converted to wetlands for the years 1990-2010. The immediate oxidation of biomass, litter and dead wood for years prior to 2006 were derived using the mean IEF for 2006 to 2010 (see section 6.3.3). Thereafter, it is assumed that natural vegetation cover will gradually recover over a period of five years at the rate of 0.6 t C ha⁻¹yr⁻¹ up to an equilibrium of 3 t C ha⁻¹.

DOM and Litter emissions and removals

DOM and Litter are indistinguishable from the organic matter in organic soils. Therefore it is assumed to be included in the assessment of carbon emissions and removals estimated for soils. It is also worth noting that the material removed from peat extraction sites for energy also incorporates DOM and litter, and therefore the carbon losses in these off-site activities are included in the Energy sector.

6.6.4.2 Soils

There is a loss of carbon associated with drainage and the exposure of the new peat surface annually after harvesting takes place. The annual activity data are the active production areas of Bord na Mona bog (Table 6.6.1), together with the areas of peatland in use by private commercial enterprises and by domestic users. Additional areas drained for the extraction of peat to supply the horticultural market, as outlined in Section 6.4.4, are included in the total area drained. These peatlands are nutrient-poor raised bogs or rain-fed blanket bogs for which the appropriate carbon emission factor is 2.8 t C/ha, for boreal and temperate climatic regions provided in Table 2.1, in the 2013 Wetlands Supplement to the IPCC good practice guidance. The activity land area in respect of the soils carbon pool is the value that appears in CRF Table 4.D.1. This area is significantly larger than that relevant to the estimation of carbon stock change in biomass above, as the land is drained on a continuous basis, whilst biomass change occurs in the first year of extraction.

Loss of Carbon via Dissolved Organic Carbon

The 2013 Wetlands Supplement also provides guidance on estimation of carbon loss through dissolved carbon entering the drainage system. This is based on the assumption of flows of carbon through extensive drainage systems. The Tier 1 methodology assigns an emission factor of carbon loss per hectare drained.

6.6.5 Emissions of Non-CO₂ Gases

6.6.5.1 N₂O emissions due to drainage of peatland for peat extraction

The Tier 1 methodology for estimation of N₂O emissions from drainage of organic soils for peat extraction is revised in Section 2.2.2.2 of the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. Eq 2.7 in the 2013 wetlands Supplement can be simplified to eq 6.5.3, for Ireland, and nutrient poor soils.

$$N_2O - N_{os} = F_{OS,G,temp,NP} \cdot EF_{2,G,Temp,NP} \text{eq 6.5.3}$$

Where N₂O-N = N₂O emissions, as N, for drained organic soils,

F_{os}= Area of drained organic soils,

EF₂= is the emission factor for N₂O losses from drained organic soils

G = Peat extraction land use

Temp= temperate climate zone

NP= indicates nutrient poor soils, which are typical of grassland types on organic soils in Ireland.

The default emission factor EF₂ of 0.3 kg N₂O-N ha⁻¹ yr⁻¹ for nutrient poor drained organic soils drained for peat extraction from Table 2.5 in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. Figure 6.6.1 shows the time series of the estimate of these emissions.




Figure 6.6.1 N₂O emissions associated with the drainage for Peat Extraction

6.6.5.2 CH₄ emissions due to drainage of peatland for peat extraction

Section 2.2.2.1 of the 2013 Wetland Supplement to the 2006 IPCC Guidelines provides methodology for estimation of CH₄ emissions and removals from drained inland organic soils. The approach requires an estimate of the area impacted by drainage, and an estimate of the density of drainage ditches constructed and maintained to achieve this drainage.




Figure 6.6.2 Estimated area of peatland drained for Peat Extraction

For Ireland, Eq 2.6 from the 2013 Wetlands Supplement can be simplified to the following:

$$CH_{4_organic} = \left(A_{T, NP, O} \cdot \left((1 - Frac_{ditch}) \cdot EF_{CH_4_{land}} + Frac_{ditch} \cdot EF_{CH_4_{ditch}} \right) \right) \text{ eq 6.5.5}$$

Where CH₄_{organic} = emissions of methane due to the drainage of peatland

A_{T, NP, O} = Area of nutrient poor, drained organic soils, in Ireland's temperate climate zone.

EF_{CH₄_land} = emission factor for methane emissions from nutrient poor soils serviced by drainage ditches in temperate zone. The default value for EF_{CH₄_land} is 6.1 kg CH₄ ha⁻¹ yr⁻¹ from table 2.3 of the 2013 Wetlands Supplement for shallow drained soils, which is typical drainage for Ireland.

EF_{CH₄_ditch} = emission factor for methane emissions from ditches in temperate zone, draining nutrient poor soils. The default value for EF_{CH₄_land} is 542 kg CH₄ ha⁻¹ yr⁻¹ from table 2.4 of the 2013 Wetlands Supplement, for shallow drained soils, which is typical drainage for Ireland.

Frac_{ditch} = Fraction of total area of drained organic soil which is occupied by ditches. The default value suggested in the 2013 Wetland supplement is Frac_{ditch} = 0.05.

Figure 6.6.3 shows the estimate of CH₄ emissions from this analysis.




Figure 6.6.3 Estimate of methane emissions due to drainage of organic soils under grasslands

6.6.6 Emissions due to Extraction and Use of Horticultural Peat

The carbon loss to the atmosphere due to the extraction and use of horticultural peat has been estimated for the first time in the 2015 submission. Ireland has developed a significant domestic and international market for horticultural peat products. The dominant producer in the market is the state owned Bord Na Mona, which, as stated above is also the dominant producer in the industrial extraction of peat for energy.

In private communication industry experts, it is estimated that 80% of horticultural peat products are exported. The quantity of peat exported is capture in the national imports and exports trade figures as provided by the CSO to the UN Commodity Trade Statistics Database, see Figure 6.6.4. <http://comtrade.un.org/db/default.aspx>




Figure 6.6.4 Time Series of Exports of Horticultural Peat products by weight

6.6.6.1 On-site emissions

BnM provide the details on the area of peatland which are in drained for extraction, which includes areas used for horticultural products. Therefore, the direct emissions associated with BnM lands are already included in the analysis above.

The activities of private commercial industry also supplying domestic and international markets are not included in the above analysis. It is estimated that non-BnM enterprises comprise 20 per cent of the export market, based the difference between recent BnM sales of horticultural products and total national sales.

It is necessary to estimate the area of peatland needed to be drained in order to supply this private industry market. It is assumed that the private industry adopt similar extraction techniques to those employed by BnM. That is a depth of peat of between 10-15cm is extracted each year.

6.6.6.2 Off-site emissions

The CSO activity data records exports of peat products by weight. It is assumed that all products are 100 per cent air dried peat, unless other information of product constituents is available.

The 2006 IPCC guidelines, Vol 4 Chp 7 section 7.2.1, provide a Tier 1 methodology and emission factor for carbon loss by weight of product. It is noted that this value is considered by national experts to be quite high, and may not be consistent with the Tier 1 methodology provided for carbon loss by volume of product. Ireland has commission a small scale study to analyse a representative sample of peat extracted by BnM for its products with the aim to provide country specific emissions factors for this source.

The export figures are scaled to include domestic use of products.

Product consistency:

In response to customer and regulatory demands in international markets, BnM have begun a programme of product development for the retail market, which includes the incorporation of non-

peat material into their products. BnM communicated confidential information regarding their programme of product development, including actions to reduce peat content in their retail products. This information is commercially sensitive, however, it has been included in the analysis estimated carbon loss from BnM products.

Figure 6.6.5 shows the time series for carbon loss based on this analysis. Note the impact of product development and changes to the constituents of the product relative to a baseline of no change in product.




Figure 6.6.5 Off-Site emissions due to use of Horticultural Peat.

6.6.7 Wetland emissions due to Biomass Burning

Activity data on the occurrence of fire on wetlands is limited. The NASA FIRMS data set for region of Ireland was interrogated. It identified approximately 71 per cent instances of likely fire events coincided with wetland location. There are significant limitations to the satellite product mainly related to the relatively low spatial resolution and high probability of cloud interference in any signal over Ireland.

From this Table of the probability of fire on each land use type was constructed. Fire events on Forest location account for 9.5 per cent of all fire detections. See Section 6.4.9 for more detailed discussion of the analysis of satellite and Forest Service data as a proxy for estimation of fires on other land uses.

The NASA FIRMS data was overlaid on spatial land cover and land use data including CORINE, and LPIS. Peatland fires tend to burn over larger areas than other fires. The areas in which fires occur tend to be under populated, with limited infrastructure at risk. Therefore fires can spread to larger areas, before being noticed, and therefore more readily detected with remote sensing. From this a table of the probability of fire on each land use type was constructed. The analysis suggests a very high proportion of fires are on peatlands. Although peatland fires are a feature of natural fire activity

in Ireland, the land cover data has difficulty in distinguishing natural vegetation on peatlands and rough grazing, especially on blanket bog. Therefore, an ad hoc decision has been made to classify 50 per cent of peatland fires as actually occurring to manage rough grazing, and therefore included in the grassland fire area. Therefore the incidence of fires on wetlands decreases to 36 per cent.

Activity data for forest fire is described in section 6.3.4.4. These data are considered more robust for fire detection relative to the low resolution FIRMS data, because the collation method is more likely to record incidences of relatively minor fires, difficult to detection in the satellite data. Comparison of the time series of both satellite and Forest service data, it can be seen than, although the remote sensing product under-records fires, there remains a strong correlation between the two data.

It is assumed that the principle driver for fire across all land uses is the incidence of suitable weather conditions. Therefore, to a first approximation, the number of fires recorded on forest land is a fixed proportion on the total number of fires. Therefore, a simple scaling has been implemented to estimate the number of fires on wetlands, based on the recorded number of fires on forest, and the relative proportion of fires detected in the satellite data which correspond to cropland.

Although meteorological conditions provide suitable conditions for fire, remote sensing cannot establish whether the actual fires are due to natural causes or direct human interventions. Unlike in other areas, such as Scotland, it is not common practice to deploy controlled burning as a peatland/heathland management tool to maintain game habitat. However, these areas, especially in mountain areas, are of high amenity value, and attract numerous visitors during fine weather, which can give rise to accidental or malicious fire setting.

Therefore, the incidence of fires on wetlands is assumed to be accidental, and all fires on wetland are classified as wildfire.




Figure 6.6.6 Estimate of area of peatland subject to wildfire

The emissions associated with fires are estimated based on the 2006 IPCC guidelines and relevant sections of the 2013 Wetlands Supplement to these.

Emissions of CH₄ and N₂O are calculated using eq 2.27 from Ch2 2 Vol 4 of the 2006 IPCC Guidelines, and shown below. The approach requires the area of grassland burnt to be stratified between mineral soils, drained organic soils and undrained organic soils, and provide appropriate default values for each of the parameters.

$$L_{fire} = A \cdot M_B \cdot C_f \cdot G_{ef} \cdot 10^{-3} \text{eq 6.4.4}$$

Where L_{fire} = amount of greenhouse gas emissions from fire, in tonnes of gas (CH₄, N₂O),

A = area burnt, ha,

M_B = mass of fuel available for combustion, tonnes ha⁻¹. This includes biomass, litter and dom. For Tier 1 Litter and DOM are assumed zero for croplands remaining croplands.

C_f = combustion factor, dimensionless,

The default value for $M_B \cdot C_f$ is 336 t dm ha⁻¹ from Table 2.4 in the Ch 2 Vol 4 2006 IPCC Guidelines.

G_{ef} = is the emission factor, g kg⁻¹ dry matter burnt. The default values for cropland are CO₂=362 gkg⁻¹ dm_{burnt}; CH₄= 9.0 g kg⁻¹ dm_{burnt}; N₂O =0.21 g kg⁻¹ dm_{burnt}.




Figure 6.6.7 Estimated CO₂ emission due to peatland fires




Figure 6.6.8 Estimated CH₄ emission due to peatland fires




Figure 6.6.9 Estimated N₂O emission due to peatland fires

6.6.8 Wetland recalculations and impact on emission trend

The main recalculation with the Wetland land use category is the inclusion for the first time of estimates of emissions and removals associated with the extraction and use of peat for horticultural use. This has had a significant impact on the absolute emissions of carbon to the atmosphere, and a change in the trends in carbon losses, which are now strongly influence by the export market for peat products.

Figure 6.6.10 shows a comparison between 2014 and 2015 submissions of estimated total emissions associated with wetlands




Figure 6.6.10 Comparison between 2014 and 2015 submissions of estimated total emissions associated with wetlands

Another major impact on the absolute emissions has been the adoption of emission factors from the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. In general, the revised emission factors are larger than the previous default emission factors relevant to Ireland. However, adoption of these emission factors does not impact on trend relative to 1990.

In additional, emissions and removals of greenhouse gases (CO_2 , N_2O and CH_4) due to biomass burning, drainage ditches are also included for the first time. These have limited impact on the overall emissions trends.

6.6.9 Wetland planned improvements

A number of studies are on-going which are investigating improved activity data collation with respect to area and condition of wetlands, including the impact of current drainage systems on raised and blanket bogs, continued domestic removal of peat for use in residential heat generation, and a national assessment on the condition and on-going impact of human interventions on all wetlands. The findings of these studies will be incorporated into the inventory assessment as appropriate.

A study has been funded to verify the default emission factor for carbon losses from the use of horticultural peat, or to establish a country specific emission factor if appropriate.

6.7 Settlements (Category 4.E)

6.7.1 Areas of Settlements

The area of settlements in 1990 is based on the urban categories within CORINE 1990. Land converted to settlements is the area demanded for new road building, available from the National Roads Authority, and the area covered by new residential, commercial and industrial construction

based on CSO annual statistics, which report floor area of development projects. An incomplete time series of housing types (for the years 1995-2010) was used to estimate the residential building footprint from floor area. It was assumed that approximately 50 per cent of the planning permits granted for construction were for green-field sites previously not part of the urban fabric.

With the exception of *Forest converted to Settlement*, the identification of previous land use from which settlement areas are converted is based on an analysis of the distribution of land use classes given by CORINE 1990. The extent of deforestation associated with conversion to settlement has been independently assessed, and is outlined in section 6.3.2 with the exclusion of wetland, water bodies, existing continuous urban fabric and other marginal lands unsuitable for development. The remaining change in Settlement area is assumed to have occurred in proportion to the respective categories in CORINE 1990.




Figure 6.7.1 Estimated Area of Settlements 1990 to 2013

It is assumed Settlement remaining Settlement is constant since 1990. All new settlement activity is categorised as “in transition”.

6.7.2 Carbon Stock Changes in Settlements

6.7.2.1 Biomass

The assumption is made of complete removal of biomass in the year of conversion. The biomass loss from grassland and cropland is as per guidelines using the Tier 1 approach. It is assumed that those lands converted from “Other Land” had a biomass equivalent to natural vegetation. The relative loss of biomass from forest per hectare is large. No account has been made of the potential increased carbon stock in biomass in urban areas, e.g. in parks or roadside planting. This may be a significant carbon sink, especially under the policy of actively encouraging urban tree planting along new roads and in new housing developments, but no data is available.

Table 6.5.15 in section 6.3.7 above gives the area of forest land converted to settlements for the years 1990-2009. The immediate oxidation of biomass, litter and dead wood for years prior to 2006

were derived using the mean IEF for 2006 to 2009 (see section 6.3.3). It is assumed there is no recovery of biomass in these areas deforested to Settlement.

6.7.2.2 Soils

With the exception of Forest converted to Settlement the estimate of change in soil carbon during conversion to settlement is based on a review of approaches taken by other reporting parties. The revised 2006 IPCC guidelines also provide some additional insight into this potential source of emissions. It is assumed that 50 per cent of the soil carbon present in the soil prior to conversion to Settlement is lost to the atmosphere, and this occurs in the year of conversion. A 100 per cent uncertainty is attached to this emission factor. The methodology applied to Forest converted to Settlement is outlined in section 6.3.6.

The estimate of soil types under settlement is based on the national distribution of soil types associated with the previous land use. It is assumed that Wetland is unsuitable for conversion to Settlement, and therefore conversion does not occur. The soil carbon content prior to conversion to Settlement is based on the land management factors and soil organic carbon reference values given in Tables 6.3.1 and 6.3.2.

6.7.2.3 Biomass Burning on Settlement

See section 6.4.9 for a detailed discussion of the analysis of areas of biomass burning. Only a very small proportion of burnt areas can be assigned to Settlement, and this includes a very high uncertainty. Given the assumption that Settlement would have low level of biomass available for burning, here it is assumed that the GIS exercise which assigned detected fires to Other Land is probably better assigned to Grassland. Therefore emissions associated with Settlement are aggregated into the analysis for Grassland.

6.7.3 Settlements recalculations and impact on emission trend

Figure 6.7.2 shows a comparison between 2014 and 2015 submissions of estimated total emissions associated with Settlements. The difference is driven by a change in the assessment of emissions and removals associated with deforestation to Settlement.




Figure 6.7.2 Comparison between 2014 and 2015 submissions of estimated total emissions associated with Settlements

6.8 Other Land (Category 4.F)

6.8.1 Areas of Other Land

The category *4.F Other Land* includes all lands not classified under the categories 4.A through 4.E. It represents the difference between the sum of categories 4.A through 4.E and the total land area of Ireland. A large part of *4.F Other Land* is not relevant in terms of its potential for emissions or removals. With the inclusion of unmanaged, natural grassland areas in Grassland reporting in this submission, the previous assumption that significant areas of afforestation occurred on “Other Land” has been revised. Much of this afforestation is now, more accurately, attributed to conversion from grasslands.

6.8.2 Carbon Stock Changes in Other Land

The degradation of lands reverting to rough grazing not in use, results in carbon losses from the soil. The soil classes are identified for *4.F.2.3 Grassland Converted to Other Land* in the same way as for other land-use categories. For mineral soils, SOC_{ref} is assigned according to Table 6.5 while Table 6.6 is used to apply the SOC adjustment factors and the carbon stock change is calculated using Equation 6.4.1. The default emission factor of 0.25 t C/ha is used to calculate carbon loss from organic soils.

Table 6.5.15 in show the transition of forest land to other land, which are not classified as crop, grassland, settlements or wetlands for the years 1990-2013. These forest conversions are small areas being converted to quarries or telecommunication masks. More recently, these areas also include forest conversions into windfarms, but these are only the areas for roads and turbine platforms. Areas in the turbulence zone are generally clearfelled and replanted.

The immediate oxidation of biomass, litter and dead wood for years prior to 2006 were derived using the mean IEF for 2006 to 2009 (see section 6.3.3). It is assumed that these deforested lands revert to a natural grassland state, and recover an above ground biomass of the order of 6 t C ha⁻¹ in the year of conversion.

6.8.3 Biomass Burning on Other Land

See section 6.4.9 for a detailed discussion of the analysis of areas of biomass burning. Only a very small proportion of burnt areas can be assigned to Other Land, and this includes a very high uncertainty. Given the assumption that Other Land would have low level of biomass available for burning, here it is assumed that the GIS exercise which assigned detected fires to Other Land is probably better assigned to Grassland. Therefore emissions associated with Other Land are aggregated into the analysis for Grassland.

6.8.4 Other Land recalculations and impact on emission trend

Figure 6.8 1 shows a comparison between 2014 and 2015 submissions of estimated total emissions associated with Other Land. The difference is driven by a change in the assessment of emissions and removals associated with deforestation to Other Land.




Figure 6.8.1 Recalculations between 2014 and 2015 submissions of estimated total emissions associated with Other Land

6.9 Uncertainty in non-Forest LULUCF categories

The purpose uncertainty analysis is to identify those key categories which contribute significantly to the uncertainty in the overall estimate emissions and removals. The results of the formal Tier 1 approach to uncertainty analysis are presented in Table 6.7. These are consistent with the findings of the qualitative discussion provided here.

Categories of land use can be identified as potential key categories for uncertainty in the estimate of greenhouse gas emissions within LULUCF by virtue of uncertainty in the activity data or uncertainty in the emission factor, or a combination of both.

Table 6.7 Summary of Uncertainty analysis

IPCC Source Category	Gas	Activity Data (AD) Uncert.	Emission Factor (EF) Uncert.	Reference Activity Data	Reference Emission Factor
Category/ Sub-category		%	%		
4.A Forest land	CO ₂	51.0	114.0	See Sections 6.3.4.7 and 6.3.5.7	Country Specific value cf Chapter 6.3
4.B.1 Cropland Remaining Cropland	CO ₂	7.2	69.1	Teagasc Soil and Sub-soil Map 2008, Trend analysis of LPIS and CSO UAA areas	Default value from IPCC GPG
4.B.2 Cropland In Transition	CO ₂	7.2	69.1	Teagasc Soil and Sub-soil Map 2008, Trend analysis of LPIS and CSO UAA areas	Default value from IPCC GPG
4.C.1 Grassland remaining Grassland	CO ₂	12.2	30.2	Teagasc Soil and Sub-soil Map 2008, Trend analysis of LPIS and CSO UAA areas	Default value from IPCC GPG
4.C.2 Grassland in Transition	CO ₂	666.7	401.8	Teagasc Soil and Sub-soil Map 2008, Trend analysis of LPIS and CSO UAA areas	Default value from IPCC GPG
4.D.1 Wetlands remaining wetlands	CO ₂	6.1	26.7	CORINE, BnM, SEAI, Expert opinion	Default value from IPCC GPG
4.D.2 Land Converted to Wetland	CO ₂	2.5	50.0	Deforestation data, Chapter 6.3	Country Specific value cf Chapter 6.3

	IPCC Source Category	Gas	Activity Data (AD) Uncert.	Emission Factor (EF) Uncert.	Reference Activity Data	Reference Emission Factor
4.E.1	Settlement remaining Settlement	CO ₂	40.4	75.0	Expert assessment of Dept of Environment Construction figures and National Road Authority infrastructure activity, CORINE	Default value from IPCC GPG
4.E.2	Settlement in Transition	CO ₂	40.4	92.5	Expert assessment of Dept of Environment Construction figures and National Road Authority infrastructure activity, CORINE	Default value from IPCC GPG
4.F.1	Other Land remaining Other Land	CO ₂	30.9	90.0	Uncertainty in Other Land Area based on combined uncertainty of land use change in other land use categories	Default value from IPCC GPG
4.F.2	Lands converted to Other Land	CO ₂	136.8	75.0	Uncertainty in Other Land Area based on combined uncertainty of land use change in other land use categories	Default value from IPCC GPG
4.A	Forest Land	CH ₄	30.0	100.0		
4.B	Cropland	CH ₄	100.0	39.1	Uncertainty in area of burning based level of detection by remote sensing, scaled by reported Forest fire	Default value from IPCC GPG
4.C	Grassland	CH ₄	96.4	91.2	Uncertainty in area of burning based level of detection by remote sensing, scaled by reported Forest fire	Default value from IPCC GPG
4.D	Wetland	CH ₄	86.0	66.5	Uncertainty in area of burning based level of detection by remote sensing, scaled by reported Forest fire	Default value from IPCC GPG
4.E	Settlement	CH ₄	0.0	0.0	Uncertainty in area of burning based level of detection by remote sensing, scaled by reported Forest fire	Default value from IPCC GPG
4.F	Other Land	CH ₄	0.0	0.0	Uncertainty in area of burning based level of detection by remote sensing, scaled by reported Forest fire	Default value from IPCC GPG
4.A	Forest Land	N ₂ O	30.0	100.0		Default value from IPCC GPG
4.B	Cropland	N ₂ O	75.0	100.0	N2O emissions associated with burning only, Activity data same as CH4 emissions for Burning	Default value from IPCC GPG
4.C	Grassland	N ₂ O	17.4	100.0	Combined uncertainty in carbon loss from drained organic soils under grassland. The uncertainty from the carbon estimate cascades to the Activity Data Uncertainty in this approach	Default value from IPCC GPG
4.D	Wetland	N ₂ O	56.8	92.7	Combined uncertainty in carbon loss from drained organic soils within Wetlands. The uncertainty from the carbon estimate cascades to the Activity Data Uncertainty in this approach	Default value from IPCC GPG
4.E	Settlement	N ₂ O	45.2	54.7	Combined uncertainty in carbon loss from drained organic soils within Wetlands. The uncertainty from the carbon estimate cascades to the Activity Data Uncertainty in this approach	Default value from IPCC GPG

6.9.1 Uncertainty in Cropland

The dominant contribution to the 2013 uncertainty is estimate of emissions and removals within the Cropland category is uncertainty in the emission factor for the loss of carbon due to conversion to cropland from other land uses, most notably conversion from temporary grassland, accounting for approximately 90% of the overall uncertainty in this category. However this analysis does not include a number of quantified elements which may prove to just as significant.

Tier 1 methodology for croplands remaining cropland assumes zero net emissions of carbon where the land management practices are well established. In general, cropland land use area in Ireland is decreasing, with croplands concentrated in well-defined regions. This supports the assumption that

the lands on which crops are grown are well established within this farming system, and takes place on the most suitable, productive soils and therefore the assumption of zero emissions is reasonable. It is difficult to quantify the uncertainty associated with this assumption.

The area of cropland in Ireland has shown a downward long term trend. The proportion of reported agricultural land under croplands is of the order of 10%, this also shows a long term downward trend. The main uncertainty in the total area of cropland is the area of non-permanent grassland which should be included in the cropland category in the context of typical crop rotation practices. In addition, the practice of land leasing gives cropland farmers access to additional land resources which allows them to respond to market drivers, and leads to the temporary conversion of land use between cropland and grassland. The 2015 submission incorporates the findings of a research study to examine this issue, using high spatial and temporal resolution data to attempt to identify a suite of representative crop rotation and farming practices which would characterise the dynamic land use pattern associated with cropland regions. It is difficult to quantify the uncertainty associated with this approach and further analysis is required.

6.9.2 Uncertainty in Grassland

Grassland has the potential for large uncertainty by virtue of large areal extent, and estimates for drainage of organic soils in this category. The dominant sources for uncertainty in this category are the activity data and emission factors associated with the use of drained organic soils within agriculture.

Grass based agriculture accounts for 90 per cent of agricultural area in Ireland. The area of permanent grasslands is very stable over time, with known afforestation accounting for a high proportion of the observed decrease in reported grassland area. Reported grassland areas include both utilised agricultural areas and natural grassland. Carbon stock changes are estimated using Tier 1 methodologies based on interannual changes in reported areas.

The area of drained organic soils under grassland is based on the proportion of agricultural grassland land cover overlaid on a soil map. There are the usual issues of matching mapping scale and interpretation of land use from land cover. Additional analysis is required to confirm these findings and the uncertainty is high.

Country specific emission factors for grasslands on drained organic soils are the subject of on-going research. Findings from which will be published in 2014. Preliminary findings indicate that the default IPCC Tier 1 emission factor may be as much as an order of magnitude in too low under conditions in Ireland. This introduces considerable uncertainty in the analysis of emissions, which cannot be quantified at this time. We welcome the publication of the IPCC 2013 Wetlands Supplement, to complement the country specific research, to help inform the analysis. Pending publication and review of this research, the 1996 IPCC GPG default factor is applied, with the recognition that the reported emissions from this activity are likely to be underestimated, and the associated uncertainty is very high.

Tier 1 methodology for grassland remaining grassland on mineral soils assumes zero net emissions where management practices are well established. There is emerging research which indicates that improved grasslands on mineral soils in Ireland continue to act as a sink of carbon. This appears to be a sustained impact of increased intensity of land management (fertiliser usage and manure management, grazing practice). However additional analysis required to confirm this result on a national scale and link it to activity data related to management practices. However, the analysis to

date is sufficient to demonstrate that Grassland remaining Grassland on mineral soils is “not a source”, but the quantitative uncertainty in this assessment is high. Khalil et al. (2012), Peichl, et al (2011). Kiely et al (2009). Byrne, and Kiely (2006).

6.9.3 Uncertainty in Wetlands

Drainage of organic soils within Wetland land use category is significant by virtue of uncertainty in areal extent and emission factors. Uncertainty analysis reveals these two components to contribute in equal measure to overall uncertainty. However, there is concern over the appropriateness of the 1996 IPCC GPG default emission factor for Ireland.

The area of peatland drained for peat extraction is dominated by the activities of the semi state commercial company Bord na Mona (BnM) which owns approximately 80kha of land of which 60kha is currently in production. There are a number of smaller commercial enterprises, mainly involved in peat extraction for horticulture which compete in the export market with BnM. BnM estimate these players have 12 per cent of the market. This is consistent with proxy data from sales export figures from Central Statistics Office. There is uncertainty in the conversion of volume of sales of peat to an equivalent area of drained lands to meet this product demand. It is assumed that the competitive operators employ similar extraction methods as BnM and therefore require an area of land in proportion to their market share. This is likely an overestimate of area drained as the extraction methods deployed are likely to be more vigorous than the approach taken by BnM. A similar issue arising with the use of proxy data from the energy sector to estimate the area of peatland drained to meet demand for residential heating by private, non-commercial sector.

In the analysis of carbon losses due to the use of horticultural peat, Ireland has adopted the Tier 1 approach based on an estimate of production from figures available in units of weights of product exported. It has been noted that the default emission factor for this approach is relatively high, and national export opinion suggests this should be verified by country specific analysis. Therefore at present Ireland considers the estimate of losses due to this source highly uncertain.

6.9.4 Uncertainty in Settlements

The area of settlement in the 1990 base year is based on the CORINE 1990 estimate of urban, industrial and other manmade environments. Change in settlement area since 1990 is based on construction statistics, national road infrastructure development and specific deforestation activities identified earlier.

There is a critical assumption which limits the potential for carbon stock change to only the specific footprint of the buildings, i.e. the sealed area, as captured in the planning permission declarations, with additional assumptions with respect to minimum new paving requirements and hedgerow removal required for new builds. This means there is an implicit assumption of no carbon stock change in lands adjacent to new constructions (green areas, etc.) relative to previous land use. Additional analysis is required to address this issue, however it is unlikely that this analysis would elevate land use change to Settlement to key category status. It is worth noting that these lands are reported as part of the “Other Land” category by default as they would not be captured in Agricultural, Forestry or Wetland statistics.

Reporting of potential change in soil carbon during conversion to settlement is based on a review of approaches taken by other reporting parties. The revised 2006 IPCC guidelines also provide some additional insight into this potential source of emissions. It is assumed that 50 per cent of the soil

carbon present in the soil prior to conversion to Settlement is lost to the atmosphere, and this occurs in the year of conversion. A 100 per cent uncertainty is attached to this emission factor.

6.9.5 Uncertainty in Other Land

In the absence of a “wall to wall” land use mapping system in Ireland, the Other Land area is estimated from the residual area required to maintain a reporting of constant total national land area once estimates for all other land use categories have been taken into account. As such, this category will be subject to the cascade of uncertainty in estimates of land use area from the other land use categories.

6.10 Quality Assurance and Quality Control

The entire compilation for this submission for both LULUCF (Chapter 6) and activities under Article 3.3 of the Kyoto Protocol (Chapter 11) were reviewed externally by an independent consultant, qualified as a UNFCCC expert reviewer for LULUCF/KP-LULUCF in March 2012. This provides an important element of quality assurance for this 2012 submission. Following the findings of this independent peer review, both chapter 6 and 11 of this report have been substantially improved to provide additional transparency and consistency between Convention and KP reporting for LULUCF.

6.10.1 Category specific QA/QC for Forest Lands

Category specific QA/QC plans and documentation for forest land are carried out by FERS Ltd on behalf of the DAFM and EPA using 2006 IPCC Guidelines(Chapter 6), these include.

6.10.2 QC plan for Activity Data

- Evaluation of required data from external sources (Forest service, Collite);
- Set up of memoranda of understanding between DAFM, EPA and data providers including:
 - Deadlines for data delivery;
 - Internalised QA/QC checks and procedures;
 - Metadata;
 - Notification of changes to methods used for collecting activity data;
 - Identification of contact points and responsible parties.
- Correspondence with data providers 2 months before agreed delivery dates to notify of new requirements, request notification of changes to any activity data and to remind providers of deadlines;
- QC checks of reference sources for national activity data by evaluation of documentation with regard to activity data. For example, is data collection or sampling regimes adequate and un-bias? Does the agency have any information on uncertainties?
- Comparisons of input data with independent data sets such as harvest statistics (FAO/Eurostat), land cover data such as CORINE (see Black et al., 2009a);
- Time series consistency checks of activity data;
- Collation and initial completeness checks of activity data required;
- Pre-processing activity data and compiling data bases to be used by CARBWARE.

6.10.3 Emission Factors, Models and Calculations

QC checks on the background data used to develop emission factors: assessment of the adequacy of the emission factors and the QA/QC performed during their development. (e.g. Byrne and Farrell, 2005-organic soil emissions; Tobin et al 2007-litter turnover).

QC checks on Models: Both the FORCARB and CABEWARE models were developed specifically for GHG inventory reporting. When these models were designed and developed the following was considered;

- Appropriateness of model assumptions, extrapolations, interpolations;
- Model calibration: models have been calibrated (see Annex 3.4.A.5) using historic (1950-2000) Irish forestry data (Hawkins et al., 2012);
- Calibration of the age class distributions used in the FORCARB model was checked against independently derived information (see Figure 6.3.9, Black et al., 2012);
- Model design specifically considered the activity data characteristics, and their applicability to the greenhouse gas inventory. For example, the key activity data for reporting is the NFI. The CARBWARE model was designed to specifically deal with single tree input data, and not stand-based data, because of NFI limitations (see Annex 3.4.A.5);
- If model descriptions, assumptions, rationale, and scientific evidence and references supporting the approach and parameters used for modelling have not been published, detailed descriptions are supplied in the Annex (3.4.A.5) to the NIR;
- Models are re-evaluated and updated annually using any new research information or if uncertainty analysis and validations indicate large uncertainties of bias in the assessment of any pool of forest subcategory. For example, the improvement to litter flow and turnover rates in conifer crops for this submission, as identified by the NIR 2012;
- All pools are included in the models, so are complete in relation to the IPCC source/sink categories. Where categories or pools are not reported, this is justified in chapter 11.

QA/QC of calculations is facilitated by the software and database management system designed to run the CARBWARE model (see Figure 6.3.8, section 6.3.3.1). This reduces the risk of calculation errors or manual error over the time series. Correct coding and calculation QA/QC was carried out by three independent parties using identical data during development of the software under the COFORD funded CARBWARE project (2007-2011).

6.10.4 Completeness and error checks in compilation of the CRF tables

Transcription of data to the CRF reported and compilation of data in the required format can result in error of inconsistencies. A check on the final CRF table is performed on completion of data transcription. Following recommendations from previous ARRs corrections or adjustments are made and documented in the NIR A QA/QC check list is documented every year to record problems detected and corrective actions.

6.10.5 Validation and QA/QC Links to Uncertainty Analysis

Comparisons of emission factors between countries: this is carried out for forest remaining forest land and land converted to forests see tables 6.10.1 and 6.10.2.

Uncertainty analysis or validation is used to identify where improvements should be made to pool or categories estimates and methods. For example, improvements are planned following the identified issue bias in estimating broadleaf biomass changes (Table 6.5.10).

Uncertainty analysis includes trend analysis to determine if there are any time series inconsistencies.

Time series adjustments are applied if there are fundamental differences in the activity data being used or methods applied over a time series.

6.10.6 Validation of Reported Estimates (Category 4.A.1)

In addition to the DBH growth model uncertainty and model validations shown in Annex 3.4.A.5, IEFs reported in the CRF table 4.A were compared to other countries with similar forest characteristics for the inventory year 2014 in the 2014 submission using the Locator Tool (Table 6.8.1). It is important to note that changes in methodology due to the introduction of new pools and EFs as part of the 2006 IPCC Guidelines, the 2013 Wetland Supplement and the 2013 KP Supplementary Guidance has resulted in significant differences in IEFs for drained organic soils and N₂O from drainage. These comparisons do not take these changes into account.

Table 6.8.1 Comparison of 2013 inventory year IEFs reported for other countries and those reported by Ireland forest land remaining forest land (4A1)

Pool	IEF (Mg/ha)			
	Ireland	EU28	UK	Range
LB net	-0.29	0.67	1.04	-0.29 to 6.77
DOM	0.43	-0.001	0.22	-0.23 to 0.51
Organic Soils	-0.22	-0.38	-0.4	-0.71 to 1.86
Fire CO ₂	260.64	IE,NO,NA	Mg/kg biomass	8.3 to 260.64
Fire CH ₄	1.14	IE,NO,NA	Mg/kg biomass	0.03 to 1.14
Fire N ₂ O	0.01	IE,NO,NA	Mg/kg biomass	<0.001 to 0.3008

The same methodologies for LB and DOB are applied in the current submission. Ireland has the lowest reported net LB IEF and second highest DOM IEF, when compared to other countries. This is because of the change in age class distribution and reduction in productivity, but harvests are maintained at the same rates, resulting in negative increment. The change in age class distributions are driven by an increase in clearfell harvest resulting in an increase in the IEF for net losses in category 4A1 from -3.5 to -6.4 from 1990 to 2012 and an increase allocation of harvest residues to the DOM pool.

Ireland is the second highest IEF for LB gains, higher than the UK but lower than Malta. Ireland has the higher LB losses than any other country. The high LB gains are consistent with experimental validation studies (Black et al., 2009a, See Table 6.8.2)

Reported IEF from organic soils are within the ranges reported.

The IEFs for wildfires in Ireland is the highest reported value under the convention.

It should be stressed that IEFs have been validated against other sources such as eddy covariance, NFI and research information and show good agreement (see Table 6.8.3). The research also shows that Irish forests have a higher NEP, NPP and GPP when compared to most published values in the literature (see Black et al., 2009a, Luyssaert et al 2007).

The higher biomass gains have been attributed to:

- a) The mild oceanic climate in Ireland and the large percentage of high yielding Sitka spruce plantations-planted at high stocking rates (2500 stems per ha).
- b) Yield classes experienced in Ireland are much higher than that in the UK. For example YC 28 to 30 m³/ha/yr can be obtained on some mineral soils, compared to a max of 24 in the British yield class tables.

Models used in the UK are based on the BFC yield tables, we show that the individual tree model used in CARBWARE provide a better estimate than BFC models (see a comparison of FORECARB and CARBWARE in section 6.3.4.1). It should be stressed that the CARBWARE single tree model has been validated against a partial sample of the new NFI (see section 6.4.3.7 Table 6.5.10). In addition, the growth models were developed using a historic permanent sample data base going back to 1950 (i.e the Coillte PSP, see Annex 3.4.A.5).

6.10.7 Validation of reported estimates (Category 4.A.2)

IEFs reported in the CRF table 4A2 were compared to other countries with similar forest characteristics for the inventory year 2010 in the 2012 submission (Table 6.8.2).

Table 6.8.2 Comparisons of 2010 inventory year IEFs reported for other countries and those reported by Ireland land converted forest land

Pool	IEF (Mg/ha)			
	Ireland	EU28	UK	Range
LB net	3.12	1.34	0.91	-0.187 to 8.79
DOM	0.7	0.21	0.03	-0.22 to 2.55
Organic Soils	-0.45	-0.65	2.77	-10.8 to 2.8
Fire CO ₂	151.8	IE,NO,NA	Mg/kg biomass	0.0009 to 151.8
Fire CH ₄	0.66	IE,NO,NA	Mg/kg biomass	0.29 to 0.03
Fire N ₂ O	0.003	IE,NO,NA	Mg/kg biomass	0.008 to <0.001

All of the reported IEFs for all pools are within the ranges reported for other countries. Additional validations of the CARBWARE growth and C flow models were carried out by comparisons to eddy covariance data (a micrometeorological measure of stand net carbon balance including all pools) from the COFORD funded CARBiFOR project. The eddy covariance measurements and standard inventory assessments, used as inputs in to the CARBWARE single tree growth and C flow model, were carried out for 2 chronosequences (Table 6.8.3):

- a) A range of Sitka spruce stands on a mineral surface water gley soil, including 2 thinning cycles;
- b) Two Ash sites aged 6 and 12 on brown earth soils.

Eddy covariance provides an estimate of net ecosystem exchange (NEE, positive values represent a net removal) excluding emissions related to immediate oxidation of harvested timber. For comparison to the CARBWARE estimated net biome productivity (i.e. NEE minus harvest losses) are shown in bold in Table 6.8.3 below.

Table 6.8.3 Validation of net biome productivity

(NBP, i.e. net C emissions/removals) estimates using CARBWARE against eddy covariance derived estimates across 2 chronosequences 2010 inventory year IEFs reported for other countries and those reported by Ireland land converted forest land

Species	Yield class	Silviculture	Forest age	Year	E.covariance	± Uncertainty	NEE - harvest	± Uncertainty	Carbware	± Uncertainty	Wlilcoxin p-value
					NEE		NBP		NBP		
					(t C ha ⁻¹ yr ⁻¹)						
Sitka spruce	24	un-thinned	20	2006	8.81	1.09	8.81	1.09	8.50	1.09	<0.01
Sitka spruce	24	1st Thin	21	2007	10.33	1.41	-3.09	2.67	-4.20	3.60	<0.05
Sitka spruce	24	1st Thin	22	2008	6.75	1.19	6.75	1.19	9.80	0.50	0.12
Sitka spruce	24	2ndThin	23	2009	8.14	1.94	-3.06	1.90	-3.90	0.59	<0.08
Sitka spruce	24	2ndThin	24	2010	8.18	1.47	8.18	1.47	7.80	0.16	<0.01
Sitka spruce	24	2ndThin	25	2011	8.54	1.11	8.54	1.11	9.30	0.72	<0.05
Sitka spruce	24	un-thinned	14	2009	8.52	1.46	8.52	1.46	7.15	0.36	<0.05
Sitka spruce	24	un-thinned	7	2009	2.21	0.46	2.21	0.46	3.58	2.54	<0.05
Ash	6	un-thinned	6	2010	1.38	0.29	1.38	0.29	-1.23	1.97	0.25
Ash	12	un-thinned	10	2010	4.67	0.71	4.67	0.71	2.14	1.54	0.14

This analysis shows that there is good agreement between the CARBWARE and eddy covariance based estimates across different age classes, species and silvicultural treatments, as evident from the signed rank Wilcoxin p-value (p-values <0.05 include no significant difference between the two estimates). It is evident that there are, however, differences in the following cases:

- The NBP for Ash sites are underestimated by CARBWARE, when compared to the eddy covariance approach. This is due to:
 - The previously mentioned under-estimation of DBH increment for the fast growing broadleaf cohort (FGB), where DBH is < 12cm (see Table 6.5.10);
 - Overestimation of litter and mortality losses in the FGB cohort models. Future improvements to the CARBWARE model are planned once more research from the COFORD research programme;
 - CARBWARE assumes that there is no significant change in mineral soil stock changes following afforestation, but NEE based estimates include emissions/ removals from soils, which in some cases can represent a net removal of C over time, although this is not always significant (see chapter 11, justification for not reporting soil CSC);
 - Eddy covariance based estimates include non-forest vegetation gains and losses, which are not estimated in the CARBWARE model. It is feasible that non-forest biomass in the early stages of forest establishment can represent a net removal of C, but this assumed to be zero at steady state since non-forest vegetation is decomposed after canopy closure.
- In one of the Sitka spruce sites (22 year old stand, Table 6.8.2), CARBWARE overestimated NBP, when compared to eddy covariance. Research from the CARBiFOR project shows that the lower NBP for this site is due to climatic inter-annual variability, which is not captured by CARBWARE.

The CARBWARE models are being continually validated against NFI data and updated as new research information from the COFORD funded programme becomes available.

Planned improvements include:

- Re-evaluation of CSC in mineral soils and emission factors for organic soils (ForCRep project 2013-2016);
- Development of a remote sensing system for tracking deforestation and land use change from forests to other land (ForCRep project 2013-2016);
- Re-evaluation of FGB cohort model using information from the BetterFOR project (2013-2016);
- Development of a remote sensing system to identify forest areas subjected to wild fires and improvement to currently used biomass combustion and EFs used in national reporting.

6.10.8 Independent External Reviews

An external review of the CARBWARE system was completed in 2007 as part of the design and methodology development research programme, funded by COFORD.

6.11 Recalculations in LULUCF

The recalculations for LULUCF includes a complete revision of the time series, now based on land remaining a land use category for the periods before 1990 and lands converted to other land uses since 1990. This required a major methodological change for forest lands resulting mainly from wider use of the national forest inventory data in the CARBWARE model for forest land and its development to ensure consistency between the LULUCF submissions under the Convention and the Kyoto Protocol. The following are the principal items leading to recalculations for the years 1990-2012 due to methodological improvements and in response to recommendations made previous annual inventory review reports. Most recalculations include the use of new EF and methodologies outlined in the 2006 IPCC guidelines, the 2013 wetland supplement, and 2013 supplementary guidance. It should also be noted that GWP of CH₄ and N₂O have been changed in the Revised UNFCCC Reporting guidelines, so some changes are not due to methodological changes. The main recalculations for 2013 took place in the following categories:

6.11.1 Forest lands (4.A)

- A major recalculations of organic soil estimates resulted in an increase in emissions from 56 kt C to 131 kt C (134 per cent increase) for category 4A1 in 2012, when 2014 and the latest submission is compared. Emissions from organic soils in category 4A2 increased from 72 kt C to 118 kt C in 2012, when 2014 and the latest submission is compared. These are due to:
 - New methods and EFs for estimating DOC losses from drained organic soils based on 2013 Wetland supplement (Eq 2.4 and Table 2.2, Ch2).
 - Re-evaluation of areas drained and those greater than 50 years old based on the 2012 NFI.
- Use of new N₂O EFs and methods for drainage of soils and inclusion of CH₄ emissions in CRF 4II, based on new GPG (see section 6.3.4.6).
- Reporting of HWP emissions/removals (CRF 4G.s.1) for the first time using harvest from domestic production, but excluding deforestation harvests.

Forests converted to other land uses

- An increase in deforestation C-CO₂eq emissions (i.e. in Tables 4C2.1 to 4F2.1) in 2012 from 223 kt in the previous submission to 237 kt CO₂ in this submission. This is due to inclusion of DOC emissions from organic soils, based on new GPG (see above).
- CO₂ and CH₄ from forest land converted to peat extraction area is now included as specified in the 2013 wetland supplement (see section 6.3.6.1.2).
- Inclusion of N₂O emission estimated from mineralisation associated with C loss from mineral forest soils (CRF 4III) converted to settlements and other lands in accordance with 2006 GPG (see section 6.3.6.1.1).
- Use of new N₂O EFs and methods for drainage of soils and inclusion of CH₄ emissions in CRF 4II, based on new GPG (see section 6.3.4.6).
- Inclusion of N₂O and CH₄ emissions associated with rewetting of wetlands in peatland regeneration projects on previously forested lands (see section 6.3.6.1.2).

6.11.2 Cropland (4.B)

- There has been a major re-estimate of Temporary Grassland associated with long term rotation patterns within the Cropland land use category. All years have recalculated.
- CH₄ and N₂O associated with fires on cropland have assessed for the first time. It is assumed that fires on croplands are accidental, and therefore have been classified as “wildfire”.
- N₂O emissions associated with the loss of carbon due to transition between temporary grassland and cropland are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.
- CO₂ emissions associated with the use of liming products on croplands are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.

6.11.3 Grassland (4.C)

- Semi-Natural, low managed, grasslands are included in the Grassland land use category for the first time. Previously these lands have been included as unmanaged grassland. However, this classification is ambiguous, as it does not reflect the decision to allow these lands to revert to a semi-natural state. The revision means there is increase in the total area of Grassland, and a corresponding decrease in area of Other Land. All previous transitions between Grassland and Other Land, and the GHG emissions associated with these, are now reported under Grassland Remaining Grassland.
- Due to the revise of the analysis of Cropland areas to include temporary grassland, there has been a knock on revision of the area of permanent agriculture grassland. This has caused a recalculation for all years.
- Estimates of changes in soil carbon loss due to drainage of organic soils under Grassland have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have recalculated.
- CH₄ and N₂O associated with drainage of organic soils on Grassland have assessed for the first time, based on the methodology and emission factors in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines.

- Estimates of changes in soil carbon loss due to drainage of organic soils for peat extraction have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have been recalculated.
- CH₄ and N₂O associated with fires on Grassland have been assessed for the first time. It is assumed that fires on Grasslands are associated with land management practices, and therefore have been classified as “controlled fires”.
- An estimate of soil carbon loss through Dissolved Organic Carbon flows into water ways has been made based on the Tier 1 methodology provided in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have been estimated.
- CO₂ emissions associated with the use of liming products on grassland are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.

6.11.4 Wetlands (4.D)

- Estimates of changes in soil carbon loss due to drainage of organic soils for peat extraction have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have been recalculated.
- CH₄ and N₂O associated with drainage of organic soils on Grassland have been assessed for the first time, based on the methodology and emission factors in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines.
- An estimate of soil carbon loss through Dissolved Organic Carbon flows into water ways has been made based on the Tier 1 methodology provided in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have been estimated.
- An estimate of carbon loss to the atmosphere associated with the extraction and use of peat for horticulture has been presented for the first time based on the Tier 1 methodology provided in the 2006 IPCC Guidelines for all years.
- An estimate of emissions associated with wildfires on wetlands has been presented for the first time based on the Tier 1 methodology provided in the 2006 IPCC Guidelines for all years.

6.12 Improvements in LULUCF

The coverage of sources of emissions and removals by Ireland in the LULUCF sector under the Convention is complete for the years 1990-2013. This submission also contains estimates for 2008-2013 in respect of activities under Article 3.3 and 3.4 (Forest Management, Cropland Management and Grazing land management) of the Kyoto Protocol (chapter 11), which are now fully consistent with Convention reporting for LULUCF. Even though a rather simplified approach has had to be followed for many land-use categories due to the level of information available, the assessment of emissions and removals according to the reporting requirements of Decision 24/CP.19 has identified a number of important CO₂ emission sources, in addition to the well-known carbon sink in forests. The inventory agency is continuing to collaborate with the bodies from which the key land-use and forestry datasets are obtained and has established formal arrangements for the provision of the data within the national system, in the same way as for other sectors. The inventory agency's capacity on GIS continues to be developed, which facilitates the assessment and integration of available

datasets. It is intended to apply this capacity in a more detailed treatment of soils for future submissions.

The results of the national forest inventory are now being applied more extensively in the LULUCF inventory and this submission reflects further improvements given by this data source and by supporting research projects on climate change and forestry being undertaken over the period from 2007 to 2014. The CARBWARE development project has improved forest carbon stock change reporting tools and software to make available an integrated system that meets the reporting needs of the Convention and the Kyoto Protocol with respect to forest land. It also draws on data from the now completed CARBiFOR II project and other related research projects, to continually refine estimates of carbon stock change for reporting purposes and for projecting carbon sinks into the future. A new research project ForCRep (2012-2016) has been funded by COFORD and the DAFM to specifically address reporting of emissions associated wild fires and further investigate soil stock changes in mineral following conversions to and from forestry. This new research project will also explore the development of methods to deforestation using a wall to wall approach based on new remote sensing products. This will be integrated with other EPA projects, using similar approached for tracking land use transitions in grasslands and crop lands.

There has also been extensive validation and verification of the models used for LULUCF and Kyoto reporting. This is part of an on-going QA/QC procedure. The LULUCF sector now adopts a tier 1 QA/QC system for LULUCF.

A new research project, funded by DAFM, has been initiated to track deforestation using a wall to wall approach based on new remote sensing products. This will be integrated with other EPA projects, using similar approached for tracking land use transitions in grasslands and crop lands.

On-going work on developing a single forest cover and attribute data set has been progressing in the Forest Service. The most recent data set has been compiled for 2010, apart from a subset of grant and premium data that needs to have species attributes input manually. Annual versions will include data on location, planting year, species area and open space area attributes, for all forest greater than 0.5 ha in area (with the post 1990 afforestation data for areas down to 0.1 ha). The Forest Service will have a system in place for access to and use of the data.

Research is on-going into the extent, and condition, of hedgerows in Ireland, which will be classified as settlement biomass in future submissions. Further research is required in this area. New research has been instigated to determine country specific emission factors associated with agricultural and forestry practices on drained organic soils. The land use conversion to settlements, particularly as regards new construction, remains a coarse estimate. Additional analysis is required to determine the real dynamic rate of conversion between grassland and croplands, and *vice versa*. This analysis will be undertaken in collaboration with the Department of Agriculture, Food and Marine.

In 2012, new research has been funded by the EPA to investigate the potential of the other data sources, especially the Land Parcel Information System, as a resource for more complete description of patterns of agricultural land use and land management in Ireland.

6.12.1 Planned Improvements for Forest Lands

- Re-evaluation of CSC in mineral soils and emission factors for organic soils (ForCRep project 2013-2016);

- Development of a remote sensing system for tracking deforestation and land use change from forests to other land (ForCRep project 2013-2016);
- Re-evaluation of FGB cohort model using information from the Betterfor project (2013-2015);
- Development of a remote sensing system to identify forest areas subjected to wild fires and improvement to currently used biomass combustion and EFs used in national reporting.

6.12.2 Planned Improvements for Other Categories

Specific issues which are to be addressed include:

- Previous land use and soil type analysis of areas converted to croplands, currently assumed to be LAC soils and that no other land categories are converted to croplands;
- The current activity data for perennial woody crops is limited. Additional data discovery and analysis will be undertaken to improve these data.
- The extrapolation of LPIS analysis to the period 1990-1999 presented in the 2015 submission is relatively simplistic. Further analysis will be undertaken, including exploration of pre-2000 agricultural spatial databases, to further refine understanding of land use within Cropland category during this period.
- Additional analysis of the revised CSO methodology is required to determine the best means to achieve continuity between the new methodology and areas published in previous years.
- A major project on Irish soils, the Irish Soils Information System, has recently published its data and produced a new, complete map of soils in Ireland. It is proposed to revise the attribution of soil type and soil carbon, and land use, based on this new, comprehensive dataset. It is hoped this will lead to the adoption of Tier 2 country specific value for reference soil carbon stock and management factors.
- A study has been funded to verify the default emission factor for carbon losses from the use of horticultural peat, or to establish a country specific emission factor if appropriate.

Chapter 7 Waste

7.1 Overview of the Waste Sector

The list of activities under *Waste* in the IPCC reporting format is given in Table 7.1 below. A summary of emissions from these activities are given in Table 7.2 and Figure 7.1.

Solid waste disposal in landfill sites, *wastewater treatment*, *waste incineration*, and *biological treatment of solid waste* are the main activities that give rise to greenhouse gas emissions in the *Waste* sector (Table 7.1).

The largest of these sources is usually *solid waste disposal* on land where CH₄ is the gas concerned. Landfills represent a key emission category in Ireland and the emission estimates of CH₄ are considered to be reasonably well quantified in the national inventory.

7.1.1 Emissions Overview

A summary of emissions from these activities are given in Table 7.2.

There is one key category in this sector, which is both a trend and level key category:

- **5.A Solid Waste Disposal (CH₄)** is a significant activity in Ireland, which peaked in 2003.

Other categories present in this sector include:

- **5.B.1 Composting** is a small but increasing sector, consisting of household organic waste collected at kerbside and brought to civic amenity/temporary collection sites, as well as organic material composted at households;
- **5.C.1 Waste Incineration** includes emissions from clinical waste up to 1997 when all hospital waste incinerators were closed, and industrial/hazardous waste which covers emissions from incineration of solvents or liquid/vapour destruction in thermal oxidisers at chemical or pharmaceutical plants, which peaked in 2006;
- **5.C.2 Open Burning of Waste** peaked in 2003;
- **5.D Wastewater Treatment and Discharge** includes treatment of domestic wastewater and human sewage.

The greenhouse gases relevant to *Waste* are as follows:

- **Carbon dioxide** emissions originate from 5.C.1 *Waste Incineration* and 5.C.2 *Open Burning of Waste*;
- **Nitrous Oxide** emissions originate from 5.B.1 *Composting*, 5.C.1 *Waste Incineration*, 5.C.2 *Open Burning of Waste*, and 5.D.1 *Human Sewage*;
- **Methane** emissions originate from 5.A.1 *Managed Waste Disposal Sites*, 5.A.2 *Unmanaged Waste Disposal Sites*, 5.B.1 *Composting*, 5.C.1 *Waste Incineration*, 5.C.2 *Open Burning of Waste*, and 5.D.1 *Domestic Wastewater*.

The 2015 submission shows total GHG emissions of 1,344.27 kt CO₂ equivalent in the *Waste* sector in 2013, of which 5.A *Solid waste disposal* accounts for 82.3 per cent, 5.B *Biological treatment of solid waste* 1.9 per cent, 5.C *Incineration and open burning of waste* 3.2 per cent and 5.D *Wastewater treatment and discharge* 12.6 per cent. The latest estimates show that emissions in the *Waste* sector

have decreased by 18.3 per cent from 1990 to 2013 mainly due to a 20.8 per cent decrease in CH₄ emissions from 5.A solid waste disposal.

7.1.2 Methodology Overview

A summary of the Tier methods consistent with the 2006 IPCC Guidelines is provided in Table 7.1 below, along with a summary of the activities applicable to Ireland.

Ireland's first waste to energy municipal solid waste (MSW) incinerator commenced operation in 2011 and emissions from this new plant have been reported under public electricity and heat production (1.A.1.a) in chapter 3 in accordance with the 2006 IPCC guidelines.

Table 7.1 Level 3 Source Methodology for Waste

5. Waste	CO₂	CH₄	N₂O
A. Solid Waste Disposal*			
1. Managed Waste Disposal Sites	NA	T2	NA
2. Unmanaged Waste Disposal Sites	NA	T2	NA
B. Biological Treatment of Solid Waste			
1. Composting	NA	T1	T1
2. Anaerobic Digestion at Biogas Facilities	NO	NO	NO
C. Incineration and Open Burning of Waste			
1. Waste Incineration	T1	T1	T1
2. Open Burning of Waste	T1,T2	T1,T2	T1,T2
D. Wastewater Treatment and Discharge			
1. Domestic Wastewater	NO	T2	T1
2. Industrial Wastewater	NO	IE	NO
E. Other	NO	NO	NO

*Key Category

T1,2,3: Tier 1, Tier 2, Tier 3 as described in the 2006 IPCC Guidelines;

NE : emissions not estimated;

NO : activity not occurring;

NA : not applicable (no emissions of the gas occur in the source category);

IE : emissions included elsewhere.

Table 7.2 Emissions from Waste 1990-2013

		Gas	Unit	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
5.A.1	Managed Waste Disposal Sites	CH ₄	kt CO ₂ eq	NO	NO	1132.8	1289.8	1425.5	1518.6	1315.6	1213.2	1336.1	951.8	855.3	717.3	745.1	860.2	787.6	958.5
5.A.2	Unmanaged Waste Disposal Sites	CH ₄	kt CO ₂ eq	1396.5	1725.8	375.4	354.0	333.5	310.4	285.7	263.5	243.6	225.6	209.4	194.6	181.2	169.0	157.8	147.5
5.B.1	Composting	CH ₄	kt CO ₂ eq	NO	NO	NO	2.2	3.4	4.7	5.0	8.0	8.0	7.3	9.6	12.3	12.2	13.4	13.1	13.3
5.B.1	Composting	N ₂ O	kt CO ₂ eq	NO	NO	NO	2.0	3.0	4.2	4.4	7.2	7.1	6.5	8.6	11.0	10.9	11.9	11.7	11.8
5.C.1	Waste incineration	CH ₄	kt CO ₂ eq	0.01	0.01	0.0003	0.0003	0.0003	0.0005	0.0005	0.0005	0.0005	0.0004	0.0003	0.0003	0.0003	0.0002	0.0002	0.0002
5.C.1	Waste incineration	N ₂ O	kt CO ₂ eq	0.83	0.83	0.60	0.64	0.65	0.98	1.11	1.08	1.04	0.83	0.62	0.64	0.54	0.38	0.45	0.43
5.C.1	Waste incineration	CO ₂	kt	83.0	83.0	58.7	63.0	64.0	96.3	109.8	106.3	102.8	82.0	61.3	62.7	53.5	37.0	44.4	42.4
5.C.2	Open burning of waste	CH ₄	kt CO ₂ eq	0.8	1.0	1.6	2.0	4.5	5.9	3.6	2.3	2.3	0.1	0.1	0.1	0.1	0.5	0.1	0.1
5.C.2	Open burning of waste	N ₂ O	kt CO ₂ eq	0.2	0.3	0.4	0.5	1.1	1.5	0.9	0.6	0.6	0.02	0.02	0.02	0.02	0.12	0.02	0.02
5.C.2	Open burning of waste	CO ₂	kt	7.6	9.4	14.6	18.9	41.7	54.7	33.0	21.0	21.6	0.7	0.6	0.7	0.7	4.5	0.7	0.7
5.D.1	Domestic wastewater	CH ₄	kt CO ₂ eq	61.1	62.7	62.4	63.0	65.2	50.3	48.3	49.3	44.4	44.2	50.9	51.5	50.3	50.1	50.7	50.9
5.D.1	Domestic wastewater	N ₂ O	kt CO ₂ eq	95.6	93.1	105.2	108.6	110.6	112.4	112.3	113.8	115.5	117.3	119.1	119.3	119.9	118.2	118.5	118.7
Total Waste				1645.7	1976.0	1751.6	1904.8	2053.2	2160.0	1919.6	1786.2	1883.0	1436.3	1315.5	1170.1	1174.4	1265.3	1184.9	1344.3




Figure 7.1 Total Emissions from Waste by Sector, 1990-2013




Figure 7.2 Total Emissions from Waste by Gas, 1990-2013

7.2 Emissions from Solid Waste Disposal (5.A)

The IPCC Level 3 emission source categories relevant under 5.A *Solid Waste Disposal* in 2013 are 5.A.1 *Managed Waste Disposal Sites* and 5.A.2 *Unmanaged Waste Disposal Sites*. Total CH₄ emissions from these activities amounted to 1,106.03 kt CO₂eq in 2013.

7.2.1 Managed Waste Disposal Sites (5.A.1)

7.2.1.1 Category Description

Treatment and disposal of municipal, industrial and other solid waste at solid waste disposal sites (SWDS) produces significant amounts of methane (CH₄). In addition to CH₄, SWDS also produce biogenic carbon dioxide (CO₂) and non-methane volatile organic compounds (NMVOCs) as well as smaller amounts of nitrous oxide (N₂O), nitrogen oxides (NO_x) and carbon monoxide (CO). Waste minimisation and recycling/reuse policies have been introduced to reduce the amount of waste generated, and increasingly, alternative waste management practices to solid waste disposal on land have been implemented to reduce the environmental impacts of waste management. Also, landfill gas recovery has become more common as a measure to reduce CH₄ emissions from SWDS.

Landfills are an important component of waste management in Ireland, but the development of a national waste management strategy for Ireland recognised the need to reduce the amount of municipal solid waste being placed in landfills (DELG, 1998).

7.2.1.2 Methodological Issues

The Tier 2 approach in the 2006 IPCC Guidelines is used for *Managed Waste Disposal Sites*. The model is a simple first-order decay spread sheet model that keeps a running total of the amount of degradable organic carbon (DOC) available in a landfill as the basis for calculating the amount of DOC converted to CH₄ and CO₂ annually. Analyses undertaken, as part of the improved methodology introduced in the 2010 submission, for both individual sites and groups of landfills shows annual MCF values increasing over time to reflect the change from generally shallow, poorly-managed landfills before 1998 (and therefore pre-landfill licensing) to well controlled and engineered landfills in subsequent years.

The EPA commenced the development of the National Waste Database (NWD) in the early 1990s to address a severe lack of information on waste production and waste management practices in Ireland. The database was needed to support radical reform of national policy and legislation on waste pursuant to the Waste Management Act of 1996 and subsequent Government strategies on sustainable development (DELG, 1997) and waste management (DELG, 1998). National statistics generated from this database published on a three-year cycle, and interim reports published on a yearly basis since 2001 by the EPA are the primary basis for establishing the historical time-series of municipal solid waste (MSW) placed in landfills from 1995 onwards. These reports include:

- Carey et al, 1996;
- Crowe et al, 2000;
- Meaney et al, 2003;
- Collins et al, 2004a; Collins et al, 2004b; Collins et al, 2005;
- Le Bolloch et al, 2006; Le Bolloch et al, 2007; Le Bolloch et al, 2009;
- McCooole et al, 2009; McCooole et al, 2011; McCooole et al, 2012; McCooole et al, 2013.

Identification and risk assessment of historical landfills under S.I. No. 524 of 2008 (DEHLG, 2008) serves as the main source of information on landfilling of waste prior to 1995. The results of other surveys undertaken in previous years (Boyle, 1987, ERL, 1993, MCOS, 1994 and DOE, 1994) have also been used to some extent in compiling the MSW time-series.

The NWD reports, published since 1995, provide a good starting point for assigning waste quantities to individual landfills and provide a representation of waste composition. However, assumptions on waste quantities and composition are still required to establish the basic historical information, given the extended time-frame that must be taken into account for a number of the models. The waste quantities for each of the 16 IPCC spread sheet model analyses are determined by adding up the amounts of household and commercial waste for the relevant landfills for each year where this is given by the NWD. The quantities of waste for other years, which are not available from the NWD, are estimated by using a variety of documents and published reports.

Waste paper products are the key determinant of degradable carbon in landfills. The NWD shows a significant decline in the proportion of waste paper products in waste going to landfills which reflects the increase in recycling of paper. The NWD is used to give the values for all years in the period 1995 to 2013. In the analysis for historical years, the paper content was fixed at 40 per cent for 1980 and previous years and decreases linearly from 40 per cent in 1980 to 30.1 per cent in 1995. The proportion of organics, the other principal constituent of waste, was estimated in the same way for each year.

In response to a recommendation from a previous review, organic waste is now separated into food and garden waste. Additional information on the composition of solid waste disposed at landfills is provided in Annex 3.5, tables 3.5.A and 3.5.B.

The waste constituents of MSW that contribute to DOC, food waste, waste paper, wood, textiles and disposable nappies, are identified in the available NWD breakdown for 1995, 1998, 2001 through 2013. The IPCC default proportions of DOC content are used for all these constituents (Annex 3.5). Street cleansing composition data is available from the NWD, and the DOC content is therefore calculated from its constituent components. In addition, a DOC content of 5 per cent has been assumed for sewage sludge.

The 2006 IPCC Guidelines provide narrow ranges for the value of decay rate constant appropriate to the individual waste components under different climatic zones. Ireland has chosen the highest values given for the Western Europe wet temperate conditions for all waste constituents, as the value of the ratio MAP:PET (Mean Annual Precipitation: Potential Evapotranspiration) is greater than 2 in Ireland.

A value of 0.6 is considered appropriate for the fraction of organic carbon that ultimately decomposes in solid waste landfills in general in Ireland, given that decomposition is not significantly inhibited by lignin, (which is one of the most slowly decomposing components of vegetation such as wood). A higher value of 0.75 has been applied in the models for two major landfills that are less than 10 years old (Annex 3.5) where site conditions and management are conducive to the enhanced degradation of organic carbon.

The choice of MCF in each of the model runs is made by assigning the individual landfill or group of landfills to the IPCC management category considered to reflect the applicable level of management for each year of their lifetime. The licensing of landfill sites came into effect around 1998, which ultimately resulted in the closure of approximately 250 sites. All landfills that continued in operation

under licence after 1998, together with all new sites, are assumed to come within the IPCC description of a managed site and the MCF of 1.0 applies. The larger landfills that were in existence prior to the introduction of waste licensing were subject to some level of management but not to the extent of fully managed licensed sites after 1998. These large sites are assigned to the IPCC category of unmanaged deep sites for the years up to 1998 with a MCF of 0.8 and to the managed category with a MCF of 1.0 for the remainder of their lifetime. The 250 sites that operated primarily as small open town dumps and shallow uncontrolled disposal sites with significant aerobic conditions up to the introduction of waste licensing are assigned to the IPCC category of unmanaged shallow sites up to 1998, for which the appropriate MCF is 0.4. A transition from unmanaged shallow classification in 1960 to one-third unmanaged shallow and two-thirds unmanaged deep sites in 1998 is applied to the remainder of sites, giving an increasing MCF from 0.4 to 0.67 over this period.

Information on the number of flares in use, together with data relating to flare capacity, run time and performance was used to estimate the volume of landfill gas flared at each site. The tonnage of CH₄ flared was calculated from landfill gas volume by accounting for gas temperature (assumed to be ambient air temperature) and suction pressure (provided in survey returns) and by using methane destruction efficiencies of 50 per cent for open flares and 98 per cent for closed flares.

The study and the surveys found that there were 13 methane utilisation plants at landfills in Ireland in 2013 with a total of 32 engines. The amount of methane input to landfill gas utilisation plants is calculated from their known electricity outputs as obtained by SEAI from EIRGRID (Electricity Transmission System Operator) and validated against survey returns using an overall efficiency of 34.6 per cent for the engines, which is considered typical of the engine types in general use. The overall results of CH₄ production, utilisation and flaring are presented in Table 7.3.




Figure 7.3 Methane Emissions from Solid Waste Disposal 1990-2013

Table 7.3. Methane Emissions from Solid Waste Disposal 1990-2013

	Methane Generation (Tonnes)	Methane Flaring (Tonnes)	Methane Utilisation (Tonnes)	Methane Recovery (Tonnes)	Percent Methane Recovery	Methane Emissions (Tonnes)	Methane Emissions (kt CO ₂ eq)
1990	55,859.50	-	-	-	-	55,859.50	1,396.49
1991	59,606.84	-	-	-	-	59,606.84	1,490.17
1992	62,498.51	-	-	-	-	62,498.51	1,562.46
1993	64,861.66	-	-	-	-	64,861.66	1,621.54
1994	67,141.19	-	-	-	-	67,141.19	1,678.53
1995	69,031.40	-	-	-	-	69,031.40	1,725.79
1996	71,113.99	598.85	5,877.35	6,476.19	9.11	64,637.80	1,615.94
1997	73,812.57	591.84	18,354.17	18,946.01	25.67	54,866.56	1,371.66
1998	76,362.12	770.11	17,632.04	18,402.15	24.10	57,959.97	1,449.00
1999	79,213.62	1,096.60	19,317.01	20,413.61	25.77	58,800.01	1,470.00
2000	83,999.69	3,855.11	19,818.49	23,673.60	28.18	60,326.09	1,508.15
2001	89,811.71	3,900.65	20,159.50	24,060.15	26.79	65,751.56	1,643.79
2002	95,239.62	8,773.44	16,107.54	24,880.98	26.12	70,358.64	1,758.97
2003	100,027.32	13,083.99	13,780.67	26,864.66	26.86	73,162.66	1,829.07
2004	103,982.82	23,180.45	16,749.43	39,929.88	38.40	64,052.94	1,601.32
2005	108,655.09	28,638.38	20,947.12	49,585.50	45.64	59,069.59	1,476.74
2006	113,567.46	29,033.88	21,346.10	50,379.98	44.36	63,187.49	1,579.69
2007	118,049.63	40,395.90	30,558.07	70,953.98	60.11	47,095.65	1,177.39
2008	122,136.82	46,639.68	32,908.74	79,548.42	65.13	42,588.40	1,064.71
2009	123,933.19	52,050.80	35,403.54	87,454.33	70.57	36,478.85	911.97
2010	124,029.12	49,886.65	37,090.88	86,977.53	70.13	37,051.59	926.29
2011	122,118.63	44,205.15	36,744.73	80,949.88	66.29	41,168.75	1,029.22
2012	119,040.83	45,121.03	36,102.70	81,223.72	68.23	37,817.11	945.43
2013	114,103.63	38,988.32	30,873.94	69,862.26	61.23	44,241.37	1,106.03

Table 7.3 and Figure 7.3 present the results for methane emissions from 5.A Solid Waste Disposal. These estimates of CH₄ generation obtained using the model in the IPCC 2006 Guidelines are considered more robust than estimates developed previous to the 2010 submission. The estimates show a steady increase in CH₄ production over the period 1990-2010, reflecting Ireland's strong dependence on solid waste disposal to landfills. The utilisation of CH₄ remained generally constant up to 2006 since becoming established in 1996. The quantity of CH₄ utilised has subsequently almost doubled with the installation of engines at a number of the newer larger landfills and expansion at other sites. The quantity of CH₄ flared increased sharply from 2003 onwards. This reflects the proliferation of the use of enclosed flares as a means of odour control at landfills throughout the country, all of which operate under EPA licence and stringent environmental controls. Methane recovery through flaring and utilisation peaked in 2009.

Table 7.4 Information related to Managed Waste Disposal (5.A.1)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	Emission Factor Reference
5.A.1	Managed Waste Disposal	T2	First Order Decay (FOD) model	2006 IPCC Guidelines

7.2.1.3 Uncertainties and Time-series Consistency

The methodologies used in the derivation of emissions estimates from the waste sector are consistent over the time-series. In the case of category 5.A, this consistency applies to all three components that determine the ultimate emissions, i.e. CH₄ generation, CH₄ flared and CH₄ utilised.

Despite continuous improvements in national data, the overall uncertainty associated with estimating CH₄ emissions from source category 5.A is high at 32.0 per cent. This uncertainty is primarily due to the length of the historical period that must be taken into account. Uncertainty estimates for the source category are calculated using equations 6.3 and 6.4 of the IPCC Good Practice Guidance. Uncertainties of 20 per cent are assumed in relation to the quantity of MSW, its composition and DOC contents, giving a combined uncertainty of 34.6 per cent for activity data using equation 6.4. The emission factor uncertainty is also 34.6 per cent, when 20 per cent is taken as the uncertainty for the fraction of DOC dissimilated, MCF and decay rate constant. This gives an uncertainty of 48.9 per cent for CH₄ generation again using equation 6.4, which is combined with uncertainties of 30 per cent and 10 per cent for CH₄ flaring and utilisation, respectively using equation 6.3 to give an uncertainty of 32.0 per cent for emissions. The Tier 1 uncertainty analysis is presented in Annex 2 of this report.

7.2.1.4 Category-specific QA/QC and verification

The inventory agency intends to continue its annual surveys of landfill operations to determine landfill gas flaring and utilisation statistics. All survey returns with respect to landfill gas flaring and utilisation that was undertaken as part of this submission were reviewed by a member of the inventory team and clarifications were sought directly from landfill operators. This data is collated with other units involved in reporting within the EPA such as annual environmental reports and E-PRTR and this collaboration ensures an element of consistency in environmental reporting in this area.

7.2.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.2.1.6 Category-specific Planned Improvements

The inventory agency intends to reconsider the uncertainty estimates for this category to account for information in the 2006 IPCC guidelines in the next submission.

7.2.2 Unmanaged Waste Disposal Sites (5.A.2)

7.2.2.1 Category Description

Solid waste disposal sites that are unmanaged are typically open dump sites or shallow uncontrolled disposal sites with significant aerobic conditions.

7.2.2.2 Methodological Issues

The Tier 2 approach in the 2006 IPCC Guidelines is used for *Unmanaged Waste Disposal Sites* as described in section 7.2.1.2. The 250 sites that operated primarily as small open town dumps and shallow uncontrolled disposal sites with significant aerobic conditions up to the introduction of waste licensing are assigned to the IPCC category of unmanaged shallow sites up to 1998, for which the appropriate MCF is 0.4. A transition from unmanaged shallow classification in 1960 to one-third unmanaged shallow and two-thirds unmanaged deep sites in 1998 is applied to the remainder of sites, giving an increasing MCF from 0.4 to 0.67 over this period.

Table 7.5 Information related to Unmanaged Waste Disposal Sites (5.A.2)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	Emission Factor Reference
5.A.2	Unmanaged Waste Disposal	T2	First Order Decay (FOD) model	2006 IPCC Guidelines

7.2.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Unmanaged Waste Disposal Sites* are provided in Annex 2.

7.2.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Unmanaged Waste Disposal Sites*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

7.2.2.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.2.2.6 Category-specific Planned Improvements

The inventory agency intends to reconsider the uncertainty estimates for this category to account for information in the 2006 IPCC guidelines in the next submission.

7.3 Emissions from Biological Treatment of Solid Waste (5.B)

Composting (5.B.1) is the only source of emissions in this category. *Anaerobic digestion at biogas facilities* (5.B.2) does not occur in Ireland. Total CH₄ and N₂O emissions from these activities amounted to 25.10 kt CO₂eq in 2013.

7.3.1 Composting (5.B.1)

7.3.1.1 Category Description

Composting is an aerobic process and a large fraction of the degradable organic carbon (DOC) in the waste material is converted into carbon dioxide (CO₂). CH₄ is formed in anaerobic sections of the compost, but it is oxidised to a large extent in the aerobic sections of the compost. The estimated CH₄ released into the atmosphere ranges from less than 1 percent to a few per cent of the initial carbon content in the material (Beck-Friis, 2001; Detzel et al., 2003; Arnold, 2005).

Composting can also produce emissions of N₂O. The range of the estimated emissions varies from less than 0.5 percent to 5 percent of the initial nitrogen content of the material (Petersen et al., 1998; Hellebrand 1998; Vesterinen, 1996; Beck-Friis, 2001; Detzel et al., 2003). Poorly working composts are likely to produce more of both of CH₄ and N₂O (e.g., Vesterinen, 1996).

Composting is composed of household organic waste collected at kerbside and brought to civic amenity/temporary collection sites, as well as organic material composted at households.

7.3.1.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC guidelines is used for *Composting* using equations 4.1 and 4.2.

Equation 4.1;

$$CH_4 \text{ Emissions} = \sum (M_i * E_i) * 10^3 - R$$

Where:

CH₄ Emissions = total CH₄ emissions in inventory year, kt CH₄

M_i = mass of organic waste treated by biological treatment type *i*, kt

EF = emission factor for treatment *i*, g CH₄/kg waste treated

i = composting

R = total amount of CH₄ recovered in inventory year, kt CH₄

Equation 4.2;

$$N_2O \text{ Emissions} = \sum (M_i * E_i) * 10^3$$

Where:

N₂O Emissions = total N₂O emissions in inventory year, kt N₂O

M_i = mass of organic waste treated by biological treatment type *i*, kt

EF = emission factor for treatment *i*, g N₂O/kg waste treated

i = composting

The EFs used are from 4.1 of the 2006 IPCC guidelines as presented in Table 7.6.

Table 7.6 Information related to Composting (5.B.1)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	N ₂ O Emission Factor	Emission Factor Reference
5.B.1	Composting	T1	4 g CH ₄ /kg	0.3 g N ₂ O/kg	2006 IPCC Guidelines Table 4.1

7.3.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Composting* are provided in Annex 2.

7.3.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Composting*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

7.3.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.3.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

7.3.2 Anaerobic Digestion at Biogas Facilities (5.B.2)

No activities have been identified for inclusion under this category. This category is reported as Not Occurring (NO).

7.4 Emissions from Incineration and Open Burning of Waste (5.C)

The emission categories relevant under *5.C Incineration and Open Burning of Waste* are *5.C.1 Waste Incineration* and *5.C.2 Open Burning of Waste*. Total emissions from these activities amounted to 45.60 kt CO₂eq in 2013.

7.4.1 Waste Incineration (5.C.1)

7.4.1.1 Category Description

Waste incineration is defined as the combustion of solid and liquid waste in controlled incineration facilities. Modern refuse combustors have tall stacks and specially designed combustion chambers, which provide high combustion temperatures, long residence times, and efficient waste agitation while introducing air for more complete combustion. Types of waste incinerated include municipal solid waste (MSW), industrial waste, hazardous waste, clinical waste and sewage sludge. The practice of MSW incineration is currently more common in developed countries, while it is common for both developed and developing countries to incinerate clinical waste.

Emissions from waste incineration without energy recovery are reported in the Waste Sector, while emissions from incineration with energy recovery are reported in the Energy Sector, both with a distinction between fossil and biogenic carbon dioxide (CO₂) emissions.

The category of *Waste Incineration* in Ireland encompasses emissions from **clinical waste** incineration and **hazardous waste** incineration from industry. The incineration of clinical waste was discontinued after 1997. Ireland's first waste to energy MSW incinerator commenced operation in 2011 and emissions from this new plant have been reported under public electricity and heat production (1.A.1.a) in chapter 3.

7.4.1.2 Methodological Issues

The Tier 1 approach in the 2006 IPCC Guidelines is used for *Waste Incineration*.

In the early 1990s, the majority of hospitals operated on-site incinerator units where hazardous **clinical waste** was incinerated. A number of hospitals operated the practice of incinerating both hazardous and non-hazardous waste. Due to the implementation of stricter standards on incineration and the requirement for facilities to be licensed by the EPA, all incinerators were closed by the mid- to late-1990s. Prior to the closure of these facilities, a number of applications were made to the EPA in respect of IPPC licences. National reports and Government records also contain some information on the quantity of health-care waste incinerated during the period of operation of the incinerators. From these sources, it was determined that an estimated 4,000 tonnes of health-care waste was incinerated per annum. This value was used across the time series for the period 1990–1997, after which negligible quantities of health-care waste were incinerated up until the closure of the two remaining incinerators in 2000. Since 1997, the bulk of clinical waste in Ireland is treated using non-incineration technologies (namely sterilisation and shredding), with the remaining waste disposed of through landfilling, exported for incineration or used as a fuel in cement kilns.

Emissions from clinical waste incineration (biogenic and non-biogenic) are estimated using the tier 1 method and equation 5.1 from the 2006 IPCC guidelines.

Equation 5.1;

$$CO_2Emissions = \sum_i (SW_i * dm_i * CF_i * FCF_i * OF_i) * 44/12$$

Where:

CO₂ Emissions = CO₂ emissions in inventory year, kt/yr

SW_i = total amount of solid waste of type *i* (wet weight) incinerated or open-burned, kt/yr

dmi = dry matter content in the waste (wet weight) incinerated or open-burned, (fraction)

CF_i = fraction of carbon in the dry matter (total carbon content), (fraction)

FCF_i = fraction of fossil carbon in the total carbon, (fraction)

O_F_i = oxidation factor, (fraction)

44/12 = conversion factor from C to CO₂

i = type of waste incinerated/open-burned specified as follows:

ISW: industrial solid waste, SS: sewage sludge, HW: hazardous waste, CW: clinical waste, others (that must be specified).

Values for the above parameters were taken from of the 2006 IPCC guidelines and are presented in Table 7.7. Methane (CH₄) and nitrous oxide (N₂O) emission factors were taken from Tables 5.3 and 5.4 of the 2006 IPCC guidelines.

Table 7.7 Information related to Waste Incineration (5.C.1)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	N ₂ O Emission Factor	CO ₂ Emission Factor	Emission Factor Reference
5.C.1	Clinical Waste	T1	60 kg/CH ₄ /kt waste (wet), 100 % oxidation factor	20 g N ₂ O/t waste (wet); 100 % oxidation factor	40 % Fossil Carbon (as % of Total Carbon); 60 % C Content of Waste	2006 IPCC Guidelines Table 5.2 pg.5.18, Table 5.3, pg 5.20, Table 5.4, pg 5.21
5.C.1	Solvent (liquid/vapour) waste	T1	0.56 g CH ₄ /tonne (wet); 100 % oxidation factor	100 g N ₂ O/tonne waste (wet); 100 % oxidation factor	100 % Fossil Carbon (as % of Total Carbon); 80 % C Content of Waste	2006 IPCC Guidelines Table 5.3, pg 5.20, Table 5.4, pg 5.22 (Netherlands)

There are currently only a small number of facilities based in the pharmaceutical and chemical sectors that operate incinerators or thermal oxidisers for the treatment of **hazardous waste**, mainly for solvent or liquid/vapour destruction. The facilities that operate these units report emissions to the atmosphere to the EPA as part of IPPC licensing requirements. Estimates of the quantity of hazardous waste incinerated at the relevant facilities are determined from returns to the National Waste Database (Carey et al, 1996; Crowe et al, 2000; Meaney et al, 2003; Collins et al, 2004a; Collins et al, 2004b; Collins et al, 2005; Le Bolloch et al, 2006; Le Bolloch et al, 2007; Le Bolloch et al, 2008; McCoolle et al, 2009; McCoolle et al, 2010; McCoolle et al, 2011; McCoolle et al, 2012 ;McCoolle et al, 2013 and McCoolle et al, in prep).

Emissions from solvent waste incineration are estimated using the tier 1 method and equation 5.3 from the 2006 IPCC guidelines. Additional information on emissions, EFs and parameters used is available in Table 3.5.D of Annex 3.5.

Equation 5.3;

$$CO_2Emissions = \sum_i (AL_i * CL_i * OF_i) * 44/12$$

Where:

CO₂ Emissions = CO₂ emissions in inventory year, kt/yr

AL_i = amount of incinerated fossil liquid waste type *i*, kt

CL_i = carbon content of fossil liquid waste type *i*, (fraction)

OF_i = oxidation factor for fossil liquid waste type *i*, (fraction)

44/12 = conversion factor from C to CO₂

7.4.1.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Waste Incineration* are provided in Annex 2.

7.4.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Waste Incineration*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

7.4.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.4.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

7.4.2 Open Burning of Waste (5.C.2)

7.4.2.1 Category Description

Open Burning of Waste in Ireland consists of the open burning of household waste.

7.4.2.2 Methodological Issues

A combination of Tier 1 and Tier 2 approaches in the 2006 IPCC guidelines is used for *Open Burning of Waste*. Statistics on open burning of waste are not available in Ireland and estimates are made based on data for uncollected household waste and proportions that are buried or burned in the open. The emission factors used to estimate emissions from open burning of waste are presented in Table 7.8.

Table 7.8 Information related to Open Burning of Waste (5.C.2)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	N ₂ O Emission Factor	CO ₂ Emission Factor	Emission Factor Reference
5.C.2	Open Burning of Waste	T1,T2	6.5 kg/CH ₄ /t waste (wet); 58 % oxidation factor	150 g N ₂ O/t waste (dry); 58 % oxidation factor	Plastics: 100 % Fossil Carbon (as % of Total Carbon) & 75 % C Content of Waste; Textiles: 20 % Fossil Carbon (as % of Total Carbon) & 50 % C Content of Waste	2006 IPCC Guidelines Table 5.3 pg 5.20, Table 5.6, Table 2.4, p. 2.14 of Ch.2 Waste, Vol.5

7.4.2.3 Uncertainties and Time-series Consistency

The uncertainties applicable to *Open Burning of Waste* are provided in Annex 2.

7.4.2.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Open Burning of Waste*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

7.4.2.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.4.2.6 Category-specific Planned Improvements

There are no planned improvements for this category.

7.5 Emissions from Wastewater Treatment and Discharge (5.D)

The IPCC Level 3 emission source categories relevant under *5.D Wastewater Treatment and Discharge* in 2013 are *5.D.1 Domestic Wastewater* (CH₄) and (N₂O). Total CH₄ and N₂O emissions from these activities amounted to 169.57 kt CO₂eq in 2013.

7.5.1 Domestic Wastewater (5.D.1)

7.5.1.1 Category Description

Wastewater can be a source of methane (CH₄) when treated or disposed anaerobically. It can also be a source of nitrous oxide (N₂O) emissions. Carbon dioxide (CO₂) emissions from wastewater are not considered in the 2006 IPCC guidelines because these are of biogenic origin and should not be included in national total emissions. Domestic wastewater is defined as wastewater from household water use, which includes emissions from commercial wastewater as the methodology is on a per person basis.

7.5.1.2 Methodological Issues

A combination of Tier 1 and Tier 2 approaches in the 2006 IPCC guidelines is used for *Domestic Wastewater*.

Approximately two-thirds of the population in Ireland is served by sewerage treatment plants, the other one-third of the population uses septic tanks to treat wastewater mainly for individual houses in non-urban areas (Smith et al., 2004). Since 2003, there are between 6 to 9 urban wastewater treatment plants with biogas recovery for heat only or CHP.

Sludge is produced in all of the primary, secondary and tertiary stages of wastewater treatment. The anaerobic stabilisation of sludge makes it safe for disposal and is a source of CH₄ in Ireland. The amount of wastewater sludge produced in Ireland is available from biennial reports on urban wastewater treatment. The sludge arising from the secondary treatment of over half of the population equivalent served by urban wastewater treatment plants is anaerobically digested. The CH₄ produced at these plants is used for electricity and heat generation for use on site since 2003. It is reported that approximately three per cent of this sludge is treated anaerobically (O' Leary et al. 1997, 2000; O'Leary and Carty, 1998; Smith et al. 2003; 2004, 2007; Monaghan et al. 2009). The average BOD of industrial wastewater sludge is 60 kg/t (40 per cent of the typical BOD content of treated industrial wastewater) and DOC is estimated as the product of average BOD content and tonnes of dry solids of sludge. The sludge from wastewater treatment is disposed of in landfills, used

as organic fertiliser on agricultural lands or in composting. The quantity of sludge that is disposed of in landfills contributes to CH₄ emissions from SWDS and is accounted for in 5.A *Solid Waste Disposal*. The sludge applied to agricultural land contributes to N₂O emissions from soils and is included in emission estimates for 3.D.1 *Direct Emissions to Soil*.

The total emissions of CH₄ from wastewater are estimated using equation 6.1 from the 2006 IPCC guidelines.

Equation 6.1;

$$CH_4 \text{ Emissions} = [\sum (U_i * T_{i,j} * EF_j)] (TOW - S) - R$$

Where:

CH₄ Emissions = CH₄ emissions in inventory year, kg CH₄/yr

TOW = total organics in wastewater in inventory year, kg BOD/yr

S = organic component removed as sludge in inventory year, kg BOD/yr

U_i = fraction of population in income group *i* in inventory year

T_{i,j} = degree of utilisation of treatment/discharge pathway or system, *j*, for each income group fraction *i* in inventory year

i = income group: rural, urban high income and urban low income

j = each treatment/discharge pathway or system

EF_j = emission factor, kg CH₄ / kg BOD

R = amount of CH₄ recovered in inventory year, kg CH₄/yr

The total organics in wastewater (TOW) in Ireland is estimated based on population equivalent data from urban waste discharge reports (equation 6.3) and is disaggregated by the degree of utilisation of treatment/discharge pathway or system (T_{i,j}) as follows;

1. Organically degradable material in wastewater at treatment plant with biogas facility (kg BOD y⁻¹)
2. Organically degradable material in wastewater at treatment plant without biogas facility (kg BOD y⁻¹)
3. Organically degradable material in wastewater at septic tanks (kg BOD y⁻¹)

Equation 6.3;

$$TOW = P \cdot BOD \cdot 0.001 \cdot I \cdot 365$$

Where:

TOW = total organics in wastewater in inventory year, kg BOD/yr

P = country population in inventory year, (person)

BOD = country-specific per capita BOD in inventory year, g/person/day, 60 g (Europe)

0.001 = conversion from grams BOD to kg BOD

I = correction factor for additional industrial BOD discharged into sewers (for collected the default is 1.25, for uncollected the default is 1.00.)

On-site domestic septic tanks consist of an underground tank (over 1 metre deep) and a percolation area for the treatment of the effluent. Prevailing soil temperatures at the depths where methanogenesis is assumed to occur (i.e. the bottom of the septic tank) only exceed 15°C for two months of the year in Ireland. Thus, the low prevailing temperatures in septic tanks means that the CH₄ correction factor (MCF) has been revised down from the 2006 IPCC guidelines default value, from 0.5 to 0.083. The CH₄ emission factor for septic tanks is estimated based on equation 6.2 from the 2006 IPCC guidelines and an EF of 0.05 kg CH₄/kg BOD.

Equation 6.2;

$$EF_j = B_o * MCF_j$$

Where:

EF_j = emission factor, kg CH₄/kg BOD

j = each treatment/discharge pathway or system

B_o = maximum CH₄ producing capacity, kg CH₄/kg BOD (0.6 kg CH₄/kg BOD)

MCF_j = methane correction factor (fraction), (0.5*2/12 = 0.083)

EFs are presented in Table 7.9 below.

The CH₄ emission factor for urban wastewater treatment plants without biogas recovery is also estimated using equation 6.2 from the 2006 IPCC guidelines and an EF of 0.018 kg CH₄/kg BOD; based on a MCF of 0.03 (assumed some anaerobic) and a B_o 0.6 kg CH₄/kg BOD.

Table 7.9 Information related to Domestic Wastewater (5.D.1)

IPCC category	Category Description	Method used	CH ₄ Emission Factor	N ₂ O Emission Factor	Emission Factor Reference
5.D.1	Septic tank	T1,T2	0.05 (kg CH ₄ /kg BOD)		2006 IPCC Guidelines, Table 6.2, modified for Ireland's cold climate
5.D.1	Urban wastewater treatment plant (without biogas facility)	T1,T2	0.018 (kg CH ₄ /kg BOD)		2006 IPCC Guidelines Table 6.2; Monaghan et al. 2009
5.D.1	Sewage	T1		3.2 g N ₂ O/person/year	2006 IPCC Guidelines Table 6.11

Wastewater treatment plants with heat or CHP account for on average over 45 percent of the BOD loading in Ireland since 2003. Emissions resulting from the biogas use/recovery are reported in the Energy sector.

Human consumption of food results in the production of sewage, which is processed in septic tanks or in wastewater treatment facilities. This treated waste is disposed of directly onto land, into the soil through percolation areas or discharged to a water body. Nitrous oxide can be produced during these processes through nitrification and denitrification. N₂O emissions are estimated using equation 6.7 and 6.8 of the 2006 IPCC guidelines.

Equation 6.7;

$$N_2O \text{ Emissions} = N_{EFFLUENT} * EF_{EFFLUENT} * 44/28$$

Where:

N_2O emissions = N_2O emissions in inventory year, kg N_2O /yr

$N_{EFFLUENT}$ = nitrogen in the effluent discharged to aquatic environments, kg N/yr

$EF_{EFFLUENT}$ = emission factor for N_2O emissions from discharged to wastewater, 0.005 kg N_2O -N/kg N (Table 6.11)

The factor 44/28 is the conversion of kg N_2O -N into kg N_2O .

$N_{EFFLUENT}$ is estimated from equation 6.8.

Equation 6.8;

$$N_{EFFLUENT} = P \cdot Protein \cdot F_{NPR} \cdot F_{NON-CON} \cdot F_{IND-COM} - N_{SLUDGE}$$

Where:

$N_{EFFLUENT}$ = total annual amount of nitrogen in the wastewater effluent, kg N/yr

P = human population

Protein = annual per capita protein consumption, kg/person/yr

F_{NPR} = fraction of nitrogen in protein, default = 0.16, kg N/kg protein

$F_{NON-CON}$ = factor for non-consumed protein added to the wastewater, 1.4 (Table 6.11)

$F_{IND-COM}$ = factor for industrial and commercial co-discharged protein into the sewer system, 1.25 (Table 6.11)

N_{SLUDGE} = nitrogen removed with sludge (default = zero), kg N/yr

Emission estimates of N_2O are provided in Table 7.10.

Table 7.10 Estimates of N₂O emissions from human sewage 1990-2013

Year	Population (million)	Days	Per capita protein consumption (g/day)	N fraction in protein (IPCC default)	Effluent EF kg N ₂ O-N/kg -N (IPCC default)	Non- consumed protein (IPCC default)	Industrial co- discharge (IPCC default)	N ₂ O * (kt)
	A	B	C	D	E	F	G	
1990	3.506	365	114.0	0.16	0.005	1.4	1.25	0.321
1991	3.526	365	113.0	0.16	0.005	1.4	1.25	0.320
1992	3.555	365	114.0	0.16	0.005	1.4	1.25	0.325
1993	3.574	365	113.0	0.16	0.005	1.4	1.25	0.324
1994	3.586	365	110.0	0.16	0.005	1.4	1.25	0.317
1995	3.601	365	108.0	0.16	0.005	1.4	1.25	0.312
1996	3.626	365	108.0	0.16	0.005	1.4	1.25	0.314
1997	3.664	365	109.0	0.16	0.005	1.4	1.25	0.321
1998	3.703	365	112.0	0.16	0.005	1.4	1.25	0.333
1999	3.742	365	115.0	0.16	0.005	1.4	1.25	0.346
2000	3.790	365	116.0	0.16	0.005	1.4	1.25	0.353
2001	3.847	365	118.0	0.16	0.005	1.4	1.25	0.365
2002	3.917	365	118.0	0.16	0.005	1.4	1.25	0.371
2003	3.980	365	118.0	0.16	0.005	1.4	1.25	0.377
2004	4.045	365	116.0	0.16	0.005	1.4	1.25	0.377
2005	4.134	365	115.0	0.16	0.005	1.4	1.25	0.382
2006	4.233	365	114.0	0.16	0.005	1.4	1.25	0.387
2007	4.376	365	112.0	0.16	0.005	1.4	1.25	0.394
2008	4.485	365	111.0	0.16	0.005	1.4	1.25	0.400
2009	4.533	365	110.0	0.16	0.005	1.4	1.25	0.400
2010	4.555	365	110.0	0.16	0.005	1.4	1.25	0.402
2011	4.575	365	108.0	0.16	0.005	1.4	1.25	0.397
2012	4.585	365	108.0	0.16	0.005	1.4	1.25	0.398
2013	4.593	365	108.0	0.16	0.005	1.4	1.25	0.398

*emissions calculated as A * B * C * D * E * F * G * 44 / 28000

7.5.1.3 Uncertainties and Time-series Consistency

Uncertainties in estimates of emissions from the source category 5.D arise due to the quality of source data, wastewater production estimates, its chemical parameters in terms of COD or BOD, the methane producing capacity and its treatment. Uncertainty estimates of 10 per cent and 30 per cent are assigned to the activity data and emission factor used, respectively.

The uncertainties applicable to *Domestic Wastewater* are provided in Annex 2.

7.5.1.4 Category-specific QA/QC and verification

Standard QA/QC procedures have been applied to *Domestic Wastewater*. Details of Ireland's QA/QC process can be found in Chapter 1 of this report.

7.5.1.5 Category-specific Recalculations

Recalculations are not being assessed for the 2015 submission.

7.5.1.6 Category-specific Planned Improvements

There are no planned improvements for this category.

7.5.2 Industrial Wastewater (5.D.2)

Emissions from *Industrial Wastewater (5.D.2)* are included in *Domestic Wastewater (5.D.1)*. This category is reported as Included Elsewhere (IE).

7.6 Emissions from Other Waste Sources (5.E)

No activities have been identified in Ireland for inclusion under this category. This category is reported as Not Occurring (NO).

Chapter 8 Other Sources

The sector *Other* in the IPCC source sector classification (Table A.2, Annex A) that is the basis for the CRF reporting tables provides for the inclusion of greenhouse gas emission sources that may be particular to individual Parties. There are no such sources to report in Ireland.

Chapter 9 Indirect CO₂ and N₂O Emissions

9.1 Description of Sources of Indirect Emissions in GHG Inventory

The use of solvents manufactured using fossil fuels as feedstocks can lead to evaporative emissions of various non-methane volatile organic compounds (NMVOC), which are subsequently further oxidised in the atmosphere. White spirit is used as an extraction solvent, as a cleaning solvent, as a degreasing solvent and as a solvent in aerosols, paints, wood preservatives, lacquers, varnishes and asphalt products. In Western Europe about 60 per cent of the total white spirit consumption is used in paints, lacquers and varnishes. White spirit is the most widely used solvent in the paint industry.

The IPCC source sector 2.D.3, *Solvent and Other Produce Use*, is important in relation to the emissions of NMVOC (non-methane volatile organic compounds). Non-methane volatile organic compounds are indirect greenhouse gases which result from the use of solvents and various other volatile compounds and are therefore reported as CO₂ equivalent emissions included in national totals.

The levels of solvent use and the emissions from solvents have changed substantially in response to product replacement and reformulation and emission controls being implemented under Integrated Pollution Prevention Control (IPPC) and the Solvents Directive (CEC, 1999).

Indirect CO₂ emissions from NMVOC accounted for 0.1 per cent (79.33 kt of CO₂ equivalent) and 0.2 per cent (66.15 kt of CO₂ equivalent) of total national emissions in 1990 and 2013, respectively. The national total for Ireland includes indirect CO₂ emissions from this category and is fully consistent with the national total emissions reported in the first commitment period of the Kyoto Protocol.

There are no key categories in this sector. Categories present in this sector include:

- 2.D.3.a Domestic solvent use including fungicides,
- 2.D.3.d Coating Applications,
- 2.D.3.e Degreasing and surface cleaning,
- 2.D.3.f Dry Cleaning,
- 2.D.3.g Chemical Products, Manufacture and Processing,
- 2.D.3.h Printing.

The emission estimates have negligible effect on national totals.

9.2 Methodological Issues

Methodologies for estimating these NMVOC emissions can be found in the EEA/EMEP Emission Inventory Guidebook (EEA, 2013). The UNFCCC reporting format explicitly provides for the inclusion of CO₂ emissions that result from the oxidation of the carbon in NMVOC emissions. This approach is consistent with the overall sectoral approach adopted for estimating CO₂ from the combustion of fuels (Section 3.2), where the CO₂ emissions are based on the full carbon content of the fuel even though some of the carbon is usually emitted as NMVOC or CO. CO₂ emission estimates are derived from NMVOCs by assuming that 85 per cent of the mass of NMVOCs is converted to CO₂.

The activity data used for computing estimates of CO₂ emissions in *Solvent and Other Product Use* are the mass emissions of NMVOC determined for the relevant source categories. The Irish data used for

this purpose are the NMVOC emissions compiled according to the EEA/EMEP Guidebook 2013 used for reporting to the UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP) (UNECE, 1999) and the National Emissions Ceilings Directive (EP and CEU, 2001).

Emissions from domestic solvent use have steadily increased across the time series, while those from the majority of other sub-categories have decreased. The main drivers for the increasing emissions from domestic solvent use are considered to be the increased per capita consumption of cosmetics, toiletries and household products. The drivers for the decrease in other sub-categories include improved management practices and abatement technologies, legislation such as the Deco Paints Directive (EP and CEU, 2004b; DEHLG, 2007) and the Solvents Directive (CEC 1999, and the economic downturn in recent years.

A detailed description of the methodology behind the NMVOC emissions from this sector can be found in Ireland's Informative Inventory Report 2015 (EPA, 2015).

9.3 Uncertainties and Time-series Consistency

The uncertainties applicable to this category can be found in Annex 2.

The uncertainty of the activity data is 30 per cent.

The uncertainty of the emission factor is 5 per cent.

There are a large number of NMVOC sources within this sector, and hence a wide range of methodologies and input datasets. For many of the methodologies, it is not possible to obtain a full time series of the input data. As a result, extrapolation, interpolation and surrogate data is used to complete the time series of emissions.

All calculations requiring extrapolation, interpolation and the use of surrogate data are clearly presented in the data processing sheets and are accompanied by comments and explanatory text from the inventory compilers to ensure transparency. In particular the use of colour coding to indicate where extrapolation and interpolation is used allows a high degree of transparency.

Some methodologies draw on point source data. This is always checked for consistency with historic data and for consistency across the different point sources within the same source sector.

9.4 Category-specific QA/QC and Verification

Quality control checks have been installed to ensure that the emission estimates calculated in the data processing sheets are the same as those in the inventory dataset that is used for reporting purposes.

9.5 Category-specific Recalculations

Category-specific recalculations have not been assessed in this report due to the significant impact of the changes in the reporting guidelines. A higher level assessment of the recalculations can be found in Chapter 10.

9.6 Category-specific Planned Improvements

There are no planned improvements for this category.

Chapter 10 Recalculations and Improvements

10.1 Introduction

On-going demands for more complete and more accurate estimates of greenhouse gas emissions means that the methodologies being used are subject to regular revision and refinement as inventory capacity is increased and better data become available. The general improvement in inventories over time may therefore introduce inconsistencies between the emissions estimates for recent years and those for years much earlier in the time-series. Recalculated estimates are often needed to eliminate these inconsistencies and to ensure that the inventories for all years in a time-series are directly comparable with respect to the sources and gases covered and that the methods, activity data and emission factors are applied in a transparent and consistent manner. In this way, the results can be used with greater confidence in identifying trends and in monitoring progress towards the commitments that have been defined with reference to emissions in the base year. The UNFCCC reporting guidelines provide for the reporting of recalculations as part of the annual submissions from Annex 1 Parties. Justification for the recalculations should be provided, as well as explanations of the changes that have been made and the numerical values of the original and revised estimates must be compared to show the impact of the changes.

10.2 Explanation and Justification for Recalculations

The foregoing chapters describe recalculations and improvements for the individual Level 1 source sectors of the inventory undertaken for the 2015 submission and they present the corresponding quantitative changes in emissions and removals within the individual sectors. The recalculations are either due to the national circumstances; revised activity data and country specific emissions factors change or the application of the 2006 IPCC Guidelines involving revised default emission factors, parameters, new sources and sinks, changes in reporting categories, new gases, changes in GWPs, and have an impact across the whole inventory. Table 10.1 records the major changes and where they are described in the 2015 NIR. This section summarises the recalculations and assesses their effect in relation to total national emissions to record the updates and the most recent emissions estimates as they appear in the 2015 submission CRF tables. The original and revised numerical values of the emissions estimates for the years 1990-2012, along with the changes related to methods, activity data and emission factors are detailed in the respective CRF Tables 8s1 to 8s4. The principal changes that give rise to recalculated estimates for the years 1990-2012 included in the 2015 submission are outlined below (Figures 10.1 to 10.6).

10.2.1 Recalculations in Energy

The overall effect of recalculations on Energy sector emissions was an increase by 0.1 per cent on average and 942.66 kt CO₂ eq. in total in the 1990-2012 trend. The main reasons for change between submissions were confined to application of the 2006 IPCC guidelines and were mostly evident in sector 1.B Fugitive Emissions from Fuels with new category emissions (coal mining and handling) and revised GWPs (CH₄) from oil and natural gas. The detailed results of the recalculations are given in CRF Tables 8s1 for the relevant years. The impact of the recalculations in the Energy sector between annual Submissions in the 1990-2012 time series is outlined below in Figure 10.1.




Figure 10.1 Impact of Recalculations in Energy between annual Submissions 1990-2012

10.2.2 Recalculations in Industrial Processes and Product Use

The overall impact of recalculations in the IPPU sector resulted in 5.0 per cent on average and 3,724.07 kt CO₂ eq. in total in the 1990-2012 trend. The results of the recalculations are given in CRF Tables 8s1 and 8s4 for the relevant years. The main reasons for the recalculation between the two submissions were; the application of the 2006 IPCC Guidelines, the revised GWP values in the UNFCCC reporting guidelines, new gases (including NF₃) in F-gases, the inclusion of Solvent Use in IPPU sector, N₂O from anaesthesia and CO₂ from Lubricant Use and Paraffin Wax Use. The impact of the recalculations in the IPPU sector between annual submissions in the 1990-2012 time series is outlined below in Figure 10.2.




Figure 10.2 Impact of Recalculations in IPPU between annual Submissions 1990-2012

10.2.3 Recalculations in Agriculture

The overall impact of recalculations in the Agriculture sector resulted in 3.2 per cent on average and 14,154.28 kt CO₂ eq. in total in the 1990-2012 trend. The main reasons for the change between the two submissions were; the application of the 2006 IPCC guidelines, the revised GWPs in the UNFCCC Reporting guidelines, the estimation of new sources including, Liming, Urea Application and Organic Soils. Additional information on recalculations is provided in CRF Table 8s2 for the relevant years. The impact of the recalculations in the Agriculture sector between annual Submissions in the 1990-2012 time series is outlined below in Figure 10.3.




Figure 10.3 Impact of Recalculations in Agriculture between annual Submissions 1990-2012

10.2.4 Recalculations in LULUCF

The recalculations for LULUCF includes a complete revision of the time series, now based on land remaining a land use category for the periods before 1990 and lands converted to other land uses since 1990. A major reassessment of the carbon losses and gains associated with the management of organic soils across all relevant land uses has had a profound impact on the emissions profile of the LULUCF sector, most notably the Grassland category. In addition, the revisions of assessment of carbon losses from organic soils within Forest Land, and the inclusion for the first time for carbon losses associated with the peat extraction and use for Horticulture, within the Wetland category, also contributed significantly to an increase in estimate of GHG emissions to air for this sector.

10.2.4.1 Forest lands (4.A)

- A major recalculations of organic soil estimates resulted in an increase in emissions from 56 kt C to 131 kt C (134 per cent increase) for category 4.A.1 in 2012, when 2014 and the latest submission is compared. Emissions from organic soils in category 4.A.2 increased from 72 kt C to 118 kt C in 2012, when 2014 and the latest submission is compared. These are due to:
 - New methods and EFs for estimating DOC losses from drained organic soils based on 2013 Wetland supplement (Eq 2.4 and Table 2.2, Ch2);
 - Re-evaluation of areas drained and those greater than 50 years old based on the 2012 NFI.
- Use of new N₂O EFs and methods for drainage of soils and inclusion of CH₄ emissions in CRF 4II, based on 2006 IPCC guidelines (see section 6.3.4.6).
- Reporting of HWP emissions/removals (CRF 4G.s.1) for the first time using harvest from domestic production, but excluding deforestation harvests.

Forests converted to other land uses

- An increase in deforestation C-CO₂eq emissions (i.e. in Tables 4C2.1 to 4F2.1) in 2012 from 223 kt in the previous submission to 237 kt CO₂ in this submission. This is due to inclusion of DOC emissions from organic soils, based on new GPG (see above).
- CO₂ and CH₄ from forest land converted to peat extraction area is now included as specified in the 2013 wetland supplement (see section 6.3.6.1.2).
- Inclusion of N₂O emission estimated from mineralisation associated with C loss from mineral forest soils (CRF 4III) converted to settlements and other lands in accordance with 2006 IPCC guidelines (see section 6.3.6.1.1).
- Use of new N₂O EFs and methods for drainage of soils and inclusion of CH₄ emissions in CRF 4II, based on new 2006 IPCC guidelines (see section 6.3.4.6).
- Inclusion of N₂O and CH₄ emissions associated with rewetting of wetlands in peatland regeneration projects on previously forested lands (see section 6.3.6.1.2).

10.2.4.2 Cropland (4.B)

- A major revision in the assessment of areas of lands which fall under the definition of croplands was implemented based on detailed analysis of new spatial data for arable land at a field scale. This has allowed reporting of areas under crops and temporary grasslands as Cropland, consistent with the definition of the category in the 2006 IPCC guidelines. The

revision impacts on the full time series. This revision also has an impact on the areas reported under Grassland (6.4.1).

- For the first time, an estimate has been provided for emissions and sinks associated with areas under commercial woody crops, full time series, which do not fulfil the definition of Forest Land. (6.4.5).
- For the first time, an estimate has been provided for emissions associated with incidents of fire on croplands, for the full time series (6.4.9).
- CO₂ emissions associated with the use of liming products on croplands are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.
- N₂O emissions associated with the loss of carbon due to transition between temporary grassland and cropland are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.

10.2.4.3 Grassland (4.C)

- Semi-Natural, low managed, grasslands are included in the Grassland land use category for the first time. Previously these lands have been included as unmanaged grassland. However, this classification is ambiguous, as it does not reflect the decision to allow these lands to revert to a semi-natural state. The revision means there is increase in the total area of Grassland, and a corresponding decrease in area of Other Land. All previous transitions between Grassland and Other Land, and the GHG emissions associated with these, are now reported under Grassland Remaining Grassland (6.5.1).
- Due to the revision of the analysis of Cropland areas to include temporary grassland, there has been a knock revision of the area of permanent agriculture grassland. This has caused a recalculation for all years (6.5.1).
- Estimates of changes in soil carbon loss due to drainage of organic soils under Grassland have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have recalculated. The revised emission factors relevant to Ireland are over an order of magnitude greater than previous default values. Therefore, this has had a profound impact on the estimation of carbon loss to the atmosphere from Grassland for all years in the time series.(6.5.2.4).
- CH₄ and N₂O associated with drainage of organic soils on Grassland have assessed for the first time, based on the methodology and emission factors in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines.
- Estimates of changes in soil carbon loss due to drainage of organic soils for peat extraction have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have recalculated.
- CH₄ and N₂O associated with fires on Grassland have assessed for the first time. It is assumed that fire on Grasslands is associated with land management practices, and therefore has been classified as “controlled fires” (6.5.4).
- An estimate of soil carbon loss soils through Dissolved Organic Carbon flows into water ways has been made based on the Tier 1 methodology provided in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have been estimated (6.3.2).

- CO₂ emissions associated with the use of liming products on grassland are no longer reported under LULUCF, but have been transposed to the Agriculture Sector.
- Greater clarity has been provided in the 2006 guidelines, and subsequent reporting documentation on how to report on N₂O emissions associated with carbon loss from grassland soils. These are now reported under Agriculture.

10.2.4.4 Wetlands (4.D)

- Estimates of changes in soil carbon loss due to drainage of organic soils for peat extraction have been recalculated to reflect the revision of the emission factors for this land use published in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have recalculated (6.6.3).
- CH₄ and N₂O associated with drainage of organic soils on Wetlands have assessed for the first time, based on the methodology and emission factors in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines (6.5.5).
- An estimate of soil carbon loss soils through Dissolved Organic Carbon flows into water ways has been made based on the Tier 1 methodology provided in the 2013 Wetlands Supplement to the 2006 IPCC Guidelines. All years have estimated. (6.6.4.2)
- An estimate of carbon loss to the atmosphere associated with the extraction and use of peat for horticulture has presented for the first time based on the Tier 1 methodology provided in the 2006 IPCC Guidelines. All years (6.6.6).
- An estimate of emission associated with wildfires on wetlands has presented for the first time based on the Tier 1 methodology provided in the 2006 IPCC Guidelines. All years. (6.6.7)

10.2.4.5 Settlement (4.E)

Settlements recalculations are driven by a revision in the assessment of emissions and removals associated with deforestation to Settlement.

10.2.4.6 Other Land (4.F)

Recalculations for all years are driven by a revision in the assessment of emissions and removals associated with deforestation to Other Land.




Figure 10.4 Impact of Recalculations in LULUCF between annual Submissions 1990-2012

The net effect of the recalculations and additional information on recalculations is provided in chapter 6 and CRF Table 8s2 for the relevant years.

10.2.5 Recalculations in Waste

The overall impact of recalculations in the Waste sector resulted in 18.9 per cent on average and 6,203.62 kt CO₂ eq. in total in the 1990-2012 trend. Additional information on recalculations is presented in CRF Table 8s3 for the relevant years. The main reasons for the change between the two submissions were; the application of the 2006 IPCC Guidelines, the revised GWPs in the UNFCCC Reporting guidelines and the estimation of new sources for the first time including, Composting and Open Burning of Waste. The impact of the recalculations in the Waste sector between annual Submissions in the 1990-2012 time series is outlined below in Figure 10.5.




Figure 10.5 Impact of Recalculations in Waste between annual Submissions 1990-2012

10.3 Effects on Emission Levels, Trends and Time-Series Consistency

Tables 10.2 and 10.3 outline the effect of the recalculations for the years 1990-2012 according to greenhouse gas and the IPCC sectors, respectively. The overall effect on the total emissions excluding LULUCF shows increases in estimates by 2.7 per cent (and 1,505.21 kt CO₂ eq.) in 1990 and by 1.8 per cent (and 1,068.10 kt CO₂ eq.) in 2012. There is no significant impact on the trend in total emissions (Chapter Two). Emissions increased in all years of the 1990-2012 trend by 1.7 per cent on average, Table 10.2 (c). The recalculations improve time-series consistency and comparability and they take account of the inventory review process by implementing the major outstanding inventory-specific recommendations of the latest annual review reports. It may be said that fully consistent greenhouse gas inventories are available for the years 1990-2012 and that these annual inventories are complete with respect to the coverage of the seven greenhouse gases and all IPCC source categories. The range of really important greenhouse gas emission sources in Ireland is quite small and the important elements of good practice are taken into account in the current approaches to estimating their emissions. The principal changes that that give rise to recalculated estimates for the years 1990-2012 included in the 2015 submission are outlined below (Tables 10.1 to 10.3 and Figure 10.6).

Table 10.1. Changes in Methodological Descriptions compared to 2014 NIR

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	DESCRIPTION OF METHODS	RECALCULATIONS	REFERENCE
	Categories where the 2015 NIR includes major changes in methodological descriptions compared to the 2014 NIR	Sub-categories where changes are reflected in recalculations of previous year estimates	Reference to sub-category, gas, pages in the NIR, Annex
Total (Net Emissions)			
1. Energy			
A. Fuel Combustion (Sectoral Approach)			
1. Energy Industries			
2. Manufacturing Ind & Const			
3. Transport	✓	✓	1A3b Road Transport CH ₄ and N ₂ O. Revised model; COPERT 4 v10.0 to COPERT 4 v11.0
4. Other Sectors			
5. Other			
B. Fugitive Emissions from Fuels			
1. Solid Fuels	✓	✓	Coal mining and post mining - new sector estimated in 2015 submission
2. Oil and Natural Gas			
C. CO ₂ Transport and Storage			
2. Industrial Processes and Product Use			
A. Mineral Industry			
B. Chemical Industry			
C. Metal Industry			
D. Non-Energy Products from Fuels and Solvent Use	✓	✓	Lubricants use and Paraffin wax use sectors estimated in 2015 submission, revised Solvents estimates
E. Electronics Industry	✓	✓	NF ₃ gas estimated in 2015 submission, revised F-gas GWPs
F. Product Uses as Substitutes for Ozone Depleting Substances	✓	✓	Revised F-gas GWPs in 2015 submission
G. Other Product Manufacture and Use	✓	✓	N ₂ O from Anaesthesia estimated in 2015 submission
H. Other	□	□	
3. Agriculture			
A. Enteric Fermentation			
B. Manure Management			
C. Rice Cultivation			
D. Agricultural Soils	✓	✓	N ₂ O from Organic Soils estimated in 2015 submission
E. Prescribed Burning of Savannas			
F. Field Burning of Agricultural Residues			
G. Liming	✓	✓	Liming estimated in 2015 submission
H. Urea Application	✓	✓	Urea Application estimated in 2015 submission
I. Other	□	□	
4. Land Use, Land-Use Change and Forestry			
A. Forest Land	□	□	
B. Cropland		□	
C. Grassland	□	□	
D. Wetlands			
E. Settlements	□	□	
F. Other Land	□	□	
G. Harvested Wood Products	✓	✓	New in 2015 submission
H. Other			
5. Waste			
A. Solid Waste Disposal			
B. Biological Treatment of Solid Waste	✓	✓	Composting estimated in 2015 submission
C. Incineration and Open Burning of Waste	✓	✓	Open Burning of Waste estimated in 2015 submission
D. Wastewater Treatment and Discharge			
E. Other			
6. Other			
Memo Items:			

Table 10.2. Recalculations by Gas 1990-2012

(a) Emissions by Gas 1990 –2012 reported in 2014 Submission (kt CO₂ eq)

GAS	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
CO ₂ emissions without net CO ₂ from LULUCF	32,424.0	35,232.5	44,689.2	47,098.3	45,676.1	45,151.2	45,805.5	47,626.9	47,227.2	47,213.6	47,005.7	41,749.6	41,292.1	37,716.3	38,011.4
CH ₄ emissions without CH ₄ from LULUCF	13,674.1	13,920.0	13,411.8	13,457.5	13,395.1	13,937.3	13,152.6	12,821.1	12,890.7	12,364.1	12,238.0	11,947.7	11,720.6	11,692.0	12,074.0
N ₂ O emissions without N ₂ O from LULUCF	9,112.1	9,555.2	9,484.8	8,974.2	8,604.4	8,519.4	8,339.6	8,123.9	7,991.5	7,745.4	7,640.1	7,551.1	7,837.0	7,288.4	7,416.6
HFCs	0.5	37.1	270.8	313.7	382.0	515.0	635.4	813.5	844.6	850.9	973.1	957.1	973.4	992.3	982.0
PFCs	0.1	75.4	305.4	296.0	212.4	228.8	182.4	168.3	148.3	130.6	106.2	65.6	37.0	13.2	8.0
SF ₆	35.5	82.9	54.3	67.8	67.7	115.4	68.8	102.0	63.4	66.2	57.5	41.2	34.7	47.7	39.2
Total (without LULUCF, with indirect)	55,246.3	58,903.2	68,216.3	70,207.5	68,337.7	68,467.1	68,184.5	69,655.7	69,165.7	68,370.7	68,020.5	62,312.3	61,894.9	57,750.0	58,531.2

(b) Recalculated Emissions by Gas 1990 –2012 reported in 2015 Submission (kt CO₂ eq)

GAS	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
CO ₂ emissions without net CO ₂ from LULUCF	32,742.4	35,693.9	45,094.5	47,470.7	45,952.7	45,554.3	46,042.4	47,952.2	47,403.1	47,489.8	47,170.9	41,981.6	41,553.8	37,898.3	38,066.6
CH ₄ emissions without CH ₄ from LULUCF	14,881.9	15,129.9	14,532.2	14,586.0	14,525.9	15,178.7	14,248.7	13,980.7	13,992.0	13,363.0	13,206.4	12,870.5	12,632.3	12,593.4	12,890.3
N ₂ O emissions without N ₂ O from LULUCF	9,013.2	8,597.9	8,585.9	8,843.9	8,171.4	8,221.3	7,926.4	8,134.8	7,806.2	6,931.5	7,103.2	7,318.9	7,476.1	7,061.3	7,395.6
HFCs	0.6	41.1	303.6	353.6	434.1	589.4	732.8	939.3	975.7	984.0	1,128.3	1,108.1	1,127.6	1,148.0	1,135.5
PFCs	0.1	97.6	397.8	379.5	267.9	286.0	234.8	216.4	191.0	168.1	136.1	83.6	46.6	15.9	9.6
SF ₆	33.9	79.1	51.8	64.6	64.5	110.0	65.3	96.8	60.2	62.9	54.7	39.2	33.1	45.5	37.4
NF ₃	NO	4.4	49.2	21.8	46.6	46.6	18.1	28.4	28.2	37.7	NO	NO	NO	NO	0.8
Total (without LULUCF, with indirect)	56,751.5	59,728.0	69,091.3	71,802.6	69,543.8	70,065.9	69,347.4	71,427.6	70,540.5	69,128.5	68,879.3	63,476.0	62,936.7	58,827.9	59,599.3

(c) Percentage Change in Emissions by Gas 1990-2012

GAS	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
CO ₂ emissions without net CO ₂ from LULUCF	1.0	1.3	0.9	0.8	0.6	0.9	0.5	0.7	0.4	0.6	0.4	0.6	0.6	0.5	0.1
CH ₄ emissions without CH ₄ from LULUCF	8.8	8.7	8.4	8.4	8.4	8.9	8.3	9.0	8.5	8.1	7.9	7.7	7.8	7.7	6.8
N ₂ O emissions without N ₂ O from LULUCF	-1.1	-10.0	-9.5	-1.5	-5.0	-3.5	-5.0	0.1	-2.3	-10.5	-7.0	-3.1	-4.6	-3.1	-0.3
HFCs	26.5	10.8	12.1	12.8	13.6	14.4	15.3	15.5	15.5	15.6	16.0	15.8	15.8	15.7	15.6
PFCs	28.6	29.5	30.2	28.2	26.1	25.0	28.7	28.5	28.7	28.7	28.2	27.5	25.8	20.3	19.0
SF ₆	-4.6	-4.6	-4.8	-4.7	-4.8	-4.7	-5.1	-5.1	-5.0	-4.9	-4.9	-4.8	-4.8	-4.6	-4.6
NF ₃	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total (without LULUCF, with indirect)	2.7	1.4	1.3	2.3	1.8	2.3	1.7	2.5	2.0	1.1	1.3	1.9	1.7	1.9	1.8

Table 10.3 Recalculations by IPCC Sector 1990-2012

(a) Emissions by IPCC Sector 1990 –2012 reported in 2014 Submission (kt CO₂ eq)

SOURCE AND SINK CATEGORIES	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
1. Energy	30,970.5	33,779.9	42,459.4	44,572.6	43,360.1	43,893.2	43,792.2	45,586.6	45,193.1	45,159.5	45,240.1	40,761.6	40,482.3	37,008.9	37,062.6
2. Industrial Processes and Product Use	3,258.5	3,150.9	4,313.1	4,442.3	3,904.1	3,276.1	3,467.5	3,710.5	3,670.0	3,703.5	3,512.3	2,620.6	2,415.3	2,292.5	2,493.9
3. Agriculture	19,634.1	20,314.4	19,970.4	19,594.8	19,378.8	19,511.0	19,315.0	18,857.3	18,721.9	18,283.2	18,150.6	17,937.4	18,004.5	17,380.6	17,967.4
5. Waste	1,383.3	1,658.0	1,473.5	1,597.9	1,694.7	1,786.8	1,609.7	1,501.2	1,580.8	1,224.5	1,117.6	992.8	992.7	1,068.0	1,007.3
6. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total (excl. LULUCF, with indirect)	55,246.3	58,903.2	68,216.3	70,207.5	68,337.7	68,467.1	68,184.5	69,655.7	69,165.7	68,370.7	68,020.5	62,312.3	61,894.9	57,750.0	58,531.2

(b) Recalculated Emissions by IPCC Sector 1990 –2012 reported in 2015 Submission (kt CO₂ eq)

SOURCE AND SINK CATEGORIES	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
1. Energy	31,117.8	33,892.1	42,525.1	44,626.3	43,402.6	44,034.5	43,826.4	45,649.0	45,152.1	45,115.7	45,210.7	40,731.3	40,359.1	36,879.6	36,950.0
2. Industrial Processes and Product Use	3,173.3	3,112.3	4,491.4	4,553.5	4,033.3	3,442.5	3,630.7	3,954.3	3,869.9	3,919.3	3,697.2	2,795.5	2,578.0	2,452.1	2,650.4
3. Agriculture	20,735.4	20,663.6	20,246.8	20,635.5	19,974.0	20,349.2	19,891.7	19,959.0	19,551.3	18,565.7	18,576.1	18,705.0	18,758.0	18,165.3	18,750.3
5. Waste	1,645.7	1,976.0	1,751.6	1,904.8	2,053.2	2,160.0	1,919.6	1,786.2	1,883.0	1,436.3	1,315.5	1,170.1	1,174.4	1,265.3	1,184.9
6. Other	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Total (excl. LULUCF, with indirect)	56,751.5	59,728.0	69,091.3	71,802.6	69,543.8	70,065.9	69,347.4	71,427.6	70,540.5	69,128.5	68,879.3	63,476.0	62,936.7	58,827.9	59,599.3

(c) Percentage Change in Emissions by Sector 1990-2012

SOURCE AND SINK CATEGORIES	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
1. Energy	0.5	0.3	0.2	0.1	0.1	0.3	0.1	0.1	-0.1	-0.1	-0.1	-0.1	-0.3	-0.3	-0.3
2. Industrial Processes and Product Use	-2.6	-1.2	4.1	2.5	3.3	5.1	4.7	6.6	5.4	5.8	5.3	6.7	6.7	7.0	6.3
3. Agriculture	5.6	1.7	1.4	5.3	3.1	4.3	3.0	5.8	4.4	1.5	2.3	4.3	4.2	4.5	4.4
5. Waste	19.0	19.2	18.9	19.2	21.2	20.9	19.3	19.0	19.1	17.3	17.7	17.9	18.3	18.5	17.6
6. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total (excl. LULUCF, with indirect)	2.7	1.4	1.3	2.3	1.8	2.3	1.7	2.5	2.0	1.1	1.3	1.9	1.7	1.9	1.8




Figure 10.6 Total Impact of Recalculations between annual Submissions 1990-2012

10.4 Response to the Review Process and Planned Improvements

Ireland recognises the need to deliver annual submissions in close conformity with the UNFCCC reporting guidelines on annual inventories to facilitate the work of expert review teams in conducting productive and efficient technical reviews of greenhouse gas inventories. Every attempt is made to participate in the UNFCCC review process and to facilitate the work of the UNFCCC secretariat, especially insofar as it impacts on the quality and transparency of the Irish estimates of emissions. The in-country review of Ireland's 2006 and 2013 submissions (UNFCCC, 2007, 2013) were an important development in this regard. The majority of the recommendations in the 2013 in-country review were implemented in the previous 2014 submission while further recommendations from the 2008 to 2013 centralised reviews of Ireland's inventory have also been addressed where feasible in the present submission. This involved greater application of country-specific information in a number of areas of the inventory and improved explanations and clarifications have been included in the 2015 NIR relating to the trend analysis and providing more circumstantial information where applicable. This submission is the first submission under the new UNFCCC Reporting guidelines and is prepared using the methodological guidance provided in the 2006 IPCC guidelines regarding revised nomenclature, new GWPs and sectoral disaggregation as well as the inclusion of new categories and gases. Annex 5.1 summarises the issues raised in the 2011 to 2013 annual inventory review reports and Ireland's response to those issues through the 2014 submission and the current submission. It may be stated therefore that the inventory material being submitted in 2015 broadly meets the principles of transparency, completeness, consistency, comparability and accuracy laid down in the UNFCCC reporting guidelines.

Further general improvements to greenhouse gas inventories are taking place through consolidation and implementation of the national system, which has been fully operational since 2007, and through application of formal QA/QC procedures that have been put into effect as an integral part of the national system. Memoranda of Understanding (MOU) which define the data inputs between the inventory agency and all key data providers and which outline the responsibilities that are conferred to the data providers under the national system (Table 1.1) underpin the national system in Ireland and have improved the quality and timely delivery of the activity data. Their application has identified where additional MOUs may be useful, including some secondary MOUs incorporated in 2009. An updated national climate change strategy was published in 2007 providing a framework in which internal review of annual inventories takes place among all stakeholders to monitor progress on the strategy, thereby fulfilling another important requirement of national system implementation.

The implementation of comprehensive QA/QC procedures in this reporting cycle according to the plan supporting the national inventory system maintains and enhances the general improvement in quality of Irish greenhouse gas inventories. The QA/QC elements include a plan and procedures for QA/QC in data selection and acquisition, data processing and reporting to comply with international requirements under Regulation No. 525/2013 of the European Parliament and of the Council and the Kyoto Protocol. The plan provides guidance on and templates for appropriate quality checking, documentation and traceability, the selection of appropriate source data and calculation methodologies. It extends to peer review and expert review of inventory data and outlines the annual requirements of a continuous improvement programme for the inventory. Participation in the internal review mechanisms within the EU as part of the QA/QC plan developed for the EU inventory under Regulation No. 525/2013 and its Implementing Regulation No. 749/2014 provides an opportunity to engage with other Member States in the examination and assessment of individual IPCC sectors and particular issues relating to methodologies and country-specific approaches that could bring mutual benefits to their greenhouse gas inventories.

PART II
SUPPLEMENTARY INFORMATION REQUIRED UNDER
ARTICLE 7.1 OF THE KYOTO PROTOCOL

Chapter 11 Emissions and Removals from LULUCF Activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol

11.1 General Information

11.1.1 Introduction

The major item of supplementary information required under Article 7.1 of the Kyoto Protocol, as outlined in annex II of the decision 2/CMP.8, is the estimation of anthropogenic emissions by sources and removals by sinks from land use land-use change and forestry activities under Article 3 paragraph 3, forest management (FM) under Article 3 paragraph 4 and any activities that a Party has elected under Article 3 paragraph 4 of the Kyoto Protocol.

Ireland **has elected** to account for optional activities under Cropland Management (CM) and Grazing land Management (GM) under Article 3 paragraph 4. Ireland has **not** elected to account for optional activities under Wetland Drainage and Rewetting, or Revegetation under Article 3.4. The approaches employed for data collection and the methodologies used to derive the estimates for Article 3.3 and FM, CM and GM 3.4 activities are described in Chapter 6, since the same approaches and time series are used for both Convention and KP reporting. The estimates of emissions and removals for these activities are compiled in supplementary CRF tables similar to those used for submitting the GHG inventory under the Convention as described in Chapter 6 of this NIR. Ireland has elected to account for KP activities at the end of the commitment period.

The reported net removals of CO₂ in 2013 on 299.07 kha of lands subject to afforestation/reforestation since 1990 is estimated at 3,710.19 kt CO₂eq while there were net emissions of 189.60 kt CO₂eq on a deforested area of 16.64 kha (Table 11.2 and 11.4). The overall forest sink for Article 3.3 forest increased from 2,335.44 kt CO₂eq in 2008 to 3,520.60 kt CO₂eq in 2013, primarily due to an increase in the area under afforestation and a decrease in emissions associated with deforestation.

Reported removals in 2013 on 449.53 kha of land under forest management is estimated to be 413.00 kt CO₂eq, most of which are associated with long term storage of C in harvested wood products.

Reported net-net emissions in 2013, relative to the 1990 base year, on 673.6 kha of land under cropland management is estimated to be +1,168 kt CO₂eq. This is mainly due to apparent changes in the patterns of utilisation of cropland and temporary grassland.

Reported net-net removals, relative to the 1990 base year, on 3,838.6 kha of land in 2013 under grazing land management is estimated to be -1,168.0 kt CO₂eq. This is mainly due to apparent changes in the patterns of utilisation of improved grassland and rough grazing.

11.1.2 Institutional Arrangements

The inventory for Article 3.3 and FM 3.4 activities is prepared by FERS Ltd, a consultant working to COFORD/DAFM (Council for Forest Research and Development) which in turn delivers the

information to the inventory agency under an agreed Memorandum of Understanding (Table 1.1). The reporting system adopts an activity based approach using the tier 3 CARBWARE national model that is applied specifically to report on Article 3.3 activities and forest management activities under article 3.4. To ensure consistency in reporting for *Lands converted to Forest Land* in the LULUCF inventory under the Convention (Chapter 6) and Afforestation and Reforestation under Kyoto Protocol, the same time series and methodological approach using the CARBWARE model has been developed and reported (see Chapter 6).

The inventory for Article 3.4 CM and GM activities are prepared by the EPA. The reporting system adopts an activity based approach using Tier 1 assessment of changes in land use on lands associated with CM and GM. To ensure consistency with reporting under Cropland and Grassland land use categories in LULUCF under the Convention, the same time series and methodological approach as reported in Chapter 6.

Table 11.1 shows the reported activities and pools. The definition of carbon pools are presented in Table 6.5.1 in section 6.3.2.6 of Chapter 6.

Table 11.1 Reported Activities and Pools (CRF Table NIR 1)

Activity	CHANGE IN CARBON POOL REPORTED ⁽¹⁾						
	Above-ground biomass	Below-ground biomass	Litter	Dead wood	Soil		HWP
					Mineral	Organic	
Article 3.3 activities							
Afforestation and reforestation	R	R	R	R	NA	R	R
Deforestation	R	R	R	R	NA	R	IO
Article 3.4 activities							
Forest management	R	R	R	R	NA	R	R
Cropland management	R	IE	NO	NO	R	NO	
Grazing land management	R	IE	NO	NO	R	R	
Revegetation	NA	NA	NA	NA	NA	NA	
Wetland drainage and rewetting	NA	NA	NA	NA	NA	NA	

Activity	GREENHOUSE GAS SOURCES REPORTED							
	Fertilization	Drained, rewetted and other soils		Nitrogen mineralization in mineral soils	Indirect N ₂ O emissions from managed soil	Biomass burning		
		N ₂ O	CH ₄	N ₂ O	N ₂ O	CO ₂	CH ₄	N ₂ O
Article 3.3 activities								
Afforestation and reforestation	IE	R	R	NO	IE	R	R	R
Deforestation	IE	R	R	R	IE	NO	NO	NO
Article 3.4 activities								
Forest management	IE	R	R	NO	IE	R	R	R
Cropland management		NO		IE		NO	R	R
Grazing land management		R		IE		NO	R	R
Revegetation	NA	NA	NA	NA	NA	NA	NA	NA
Wetland drainage and rewetting	NA	NA	NA		NA	NA	NA	NA

R indicates the reported carbon pools and emissions from biomass burning;

IE (included elsewhere) is used to show that emissions from fertilisation of soils and indirect N₂O emissions are included under Agriculture

HWP from lands reported under deforestation, which originated from the deforestation event at the time of the land-use change are accounted for on the basis of instantaneous oxidation (IO).

NA : Mineral soils are shown not to be "a source", so are not reported.

NO: Mineralisation losses in mineral soils do not occur since there are no changes in mineral soil carbon stocks from AR and FM, CO₂ emissions from Biomass burning under Cm and GM are assumed to be transient, and taken up in subsequent regrowth of vegetation.

Table 11.2 Land Transition Matrix (CRF Table NIR 2) for inventory year 2013

	ARTICLE 3.3 ACTIVITIES		ARTICLE 3.4 ACTIVITIES					Other ⁽⁶⁾	Total area at the end of the previous inventory year ⁽⁷⁾
	Afforestation and reforestation	Deforestation	Forest management ⁽⁵⁾	Cropland management (if elected)	Grazing land management (if elected)	Revegetation (if elected)	Wetland drainage and rewetting (if elected)		
	(kha)								
Article 3.3 activities									
Afforestation and reforestation	292.91	0.10							293.01
Deforestation		16.51							16.51
Article 3.4 activities									
Forest management		0.03	NO						0.03
Cropland management ⁽³⁾ (if elected)	0.19		NO	673.82	NO	NO	NO		674.01
Grazing land management ⁽³⁾ (if elected)	2.80		NO	NO	3,847.68	NO	NO		3,850.48
Revegetation ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Wetland drainage and rewetting ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Other ⁽⁴⁾	3.26	NO	449.53	NO	0.47	NA	NA	1,824.49	2,277.74
Total area at the end of the current inventory year	299.17	16.64	449.53	673.82	3,848.14	NA.NO	NA.NO	1,824.49	7,111.79

Areas and changes in areas between the previous and the current inventory year

Table 11.3 Key Categories for Article 3.3 and A.4 Activities (CRF Table NIR 3)

KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			Comments ⁽⁴⁾
		Associated category in UNFCCC inventory ⁽¹⁾ is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory ⁽²⁾ (including LULUCF)	Other ⁽³⁾	
Specify key categories according to the national level of disaggregation used ⁽¹⁾					
CM Mineral Soils	CO2	Cropland Remaining Croplands Mineral Soil	Yes	Approach 3, spatially explicit cropland management at parcel scale	High inter-annual variation in areas of land switching between crops and temporary grassland.
GM Organic Soil	CO2	Grassland Remaining Grassland Organic Soils	Yes	Approach 3 has been used to determine soil type. However, Approach 1 has been used to determine change in areas	The management of areas of drained organic soils gradually decreasing, leading to a net-net decrease in carbon loss. Need verification of this finding.

Table 11.4 Information Table on Reporting of emission/removals for Activities under Article 3, paragraphs 3 and 4 of the Kyoto Protocol

Year	A.1. KP_AR			A.2.KP_D			B.1.KP_FM			B.1.KP CM			B.1.KP GM		
	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
2008	-3021.69	0.92	0.26	461.91	0.17	0.41									
2009	-3272.73	0.91	0.27	387.21	0.18	0.41									
2010	-3502.54	1.15	0.28	205.21	0.18	0.40									
2011	-3627.42	1.01	0.28	318.55	0.23	0.40									
2012	-3678.07	0.96	0.29	240.57	0.23	0.40									
2013	-3823.19	1.04	0.29	61.89	0.23	0.41	-548.93	2.05	0.28	-344.76	0.003	7.9x10 ⁻⁵	4,901.1	15.97	6.8x10 ⁻⁴

11.1.3 Definitions and Application

For the definition of different carbon (C) pools please refer to Ch 6 section 6.2.3.6. Also see Table 11.1 for reporting of different C pools.

Forest definition

The definition of forest is the same as that adopted for the LULUCF inventory under the Convention. Forest land has a minimum area of 0.1 hectare, a minimum width of 20 m, trees higher than 5 m and a canopy cover of more than 20 per cent within the forest boundary, or trees able to reach these thresholds *in situ*. This is consistent with the forest definition contained in decision 16/CMP.1. The following attributes are also relevant to the definition:

- A tree is a woody perennial of a species forming a single main stem or several stems, and having a definitive crown;
- A forest includes windbreaks, shelterbelts and corridors of trees with an area of more than 0.1 ha and minimum width of 20 m;
- Forest is determined both by the presence of trees/stumps and the absence of other predominant land-uses. Areas under re-establishment (following clearfell) that have not yet reached but are expected to reach a canopy cover of 20 per cent and a minimum tree height of 5 m are included, as are temporarily un-stocked areas, resulting from human intervention, which are expected to be restocked;
- The forest area is determined by the forest boundary. The term forest boundary is defined by any man-made boundary enclosing the forest area or, in the absence of such boundary feature, the boundary of the forest is determined by extending out 1 m from the position of the pith-line of the outermost trees (NFI, 2007a);
- The forest area includes forest roads other open areas on forest land; forest in national parks, nature reserves and other protected areas such as those of specific scientific, historical, cultural or spiritual interest;
- The forest area excludes tree stands in agricultural production systems, for example in fruit plantations and Christmas tree plantations since these generally do not reach 5m in height;
- The term forest also includes trees in urban parks and gardens, provided these areas satisfy the forest definition.

The forest definition is applied to the NFI when land cover and use is determined (see Ch 6 Section 6.3.2.3). The classification forest roads, open forest areas within forest boundaries are done at the plot level when the NFI team established permanent sample plots.

Forest management

Ireland considers that all areas meeting the forest definition are managed through forestry operations (timber resource utilisation) or for other reasons such as conservation, control of invasive species, pests or diseases. Therefore, activities under FM include all areas which meet the forest definition and were first established before the 1st of January 1990, or are pre-existing semi-natural forests before 1st of January 1990.

Natural disturbances

Ireland applies the same definition for natural disturbance as that outlined in the annex of decision 2/CMP.7 paragraph 1a. Ireland considers that wildfires, insect and disease outbreaks, extreme weather and or geological disturbances outside the control of or not influenced by policy in Ireland. Ireland may wish to apply the provisions to exclude emissions from natural disturbances for the accounting for afforestation and reforestation (AR) under Article 3, paragraph 3, of the Kyoto Protocol and FM under Article 3, paragraph 4, of the Kyoto Protocol during the second commitment period in accordance with decision 2/CMP.7, annex, paragraph 33 (see section 11.6).

Carbon equivalent forest conversion (CEFC)

Ireland has not identified any land which qualifies for CEFC as outlined in the decision 2/CMP.7, annex, paragraph 37.

Cropland Management

The definition of cropland is the same as that adopted for the LULUCF inventory under the Convention. This is consistent with the definition contained in decision 16/CMP.1. *“Cropland management” is the system of practices on land on which agricultural crops are grown and on land that is set aside or temporarily not being used for crop production* The following activities are also relevant to the CM:

- Non-forest woodland and hedgerows, including boundary features, associated with identified croplands are included in the definition of CM. However assessment of these has not been included in this submission;
- CM includes areas identified as temporary grassland, with a history of cultivation for crops in the period 1990-2013;
- The CM area includes tree stands (not classified as forests) in agricultural production systems;
- CM excludes recreational areas in urban parks and gardens, which have been used for recreational/private vegetable growing.

Ireland considers all areas which have been identified as being utilised for crop cultivation in the period 1990-2013 as subject to Cropland Management activity. In any given year these include areas under crops and areas under temporary grasslands. Those areas previously reported under CM which have been converted to other land uses not reportable under mandatory 3.3 or 3.4 activities, and those not reportable under the elected activities will also continue to be reported under CM. However, as 2013 is the first reporting period, this does not arise in the 2015 submission.

Grazing land Management

The definition of Grazing land Management is the same as that adopted for Grassland for the LULUCF inventory under the Convention, with necessary caveats regarding semi-natural grasslands. This is consistent with the definition contained in decision 16/CMP.1, *“Grazing land management” is the system of practices on land used for livestock production aimed at manipulating the amount and type of vegetation and livestock produced*”. The following caveats are relevant to the definition:

- Non-forest woodland and hedgerows, including boundary features, associated with identified grasslands are included in the definition of GM, however, assessment of these has not been included in this submission;
- GM excludes areas identified as temporary grassland, with a history of cultivation for crops in the period 1990-2013;
- GM excludes semi-natural grasslands which have been unmanaged before the 1st Jan 1990. GM includes semi-natural grasslands which have fallen into this classification since 1st Jan 1990.
- GM excludes recreational areas such as urban parks, sporting facilities and gardens.

Ireland considers grassland areas which have identified as utilised as agricultural land in the period since 1st Jan 1990 as areas under Grazing land management, unless explicitly identified under another reported Art 3.3 or Art 3.4 activities. Therefore areas of land which had a history of crop cultivation, but are currently under grass are reported under CM and excluded from GM. Semi-natural grasslands identified prior to 1st Jan 1990, and continuing as semi-natural grassland are excluded. Semi-natural grasslands which have developed from managed grassland types since 1990 continue to be included as their current status is a direct result of a management decision.

Areas of grassland for recreational and settlement use are not included as grazing land, in the first instance. However, those areas previously reported under GM which have been converted to other land uses not reportable under mandatory 3.3 or 3.4 activities, and those not reportable under the elected activities will continue to be reported under GM. However, as 2013 is the first reporting period, this does not arise in the 2015 submission.

11.1.4 The geographical location of the boundaries of the areas that encompass activities under Article 3, paragraphs 3 and 4

The following information is provided to assist in review for compliance under annex 2 to Decision 2/CMP.8. The definition of reporting boundaries and their geographical locations for afforestation, deforestation and forest management areas are reported within the entire territory of Ireland, with further sub division of species strata within internal national boundaries. The national boundary is used as the basis for the random systematic grid sample used in the National forest inventory (NFI, see Section 6.3.2.3 in Ch 6).

The definition of reporting boundaries and their geographical locations for cropland management and grazing land management areas are reported within the entire territory of Ireland, with further sub division of management type and soil type strata within internal national boundaries. The national boundary is used as the basis for data collation for the Land Parcel Information System and Central Statistics Office analysis of Utilised Agricultural Area (see Section 6.5.1 in Ch 6).

11.1.5 Classification hierarchy and continuity of accounting reported activities over time

Activities under forest management (FM Art 3.4) are distinguished from AR lands based on the year of afforestation as derived from the IFORIS system and the NFI (sections 6.2.3.2 and 6.2.3.3 in Ch 6). This system identifies units of land subject to activities under Article 3, paragraph 3, of the Kyoto Protocol which would otherwise be included in land subject to forest management or elected activities under Article 3, paragraph 4, of the Kyoto Protocol under the provisions of decision

2/CMP.7, annex, paragraph 9. The hierarchy used for assigning land areas to specific activities ensures that all areas reported under afforestation or forest management activities cannot decrease unless converted to deforestation land (see KP NIR3).

Activities under cropland management (CM Art 3.4) and grazing land management (GM Art 3.4) are distinguished on the basis of land parcel histories derived from the Land Parcel Information System, and the CORINE land cover database (section 6.4.4 in Chapter 6). The hierarchy used for assigning land areas to specific activities ensures that all areas reported under CM or GM cannot decrease unless converted to Art 3.3 Afforestation land, or swapping between CM and GM. (see KP NIR3). From Figure 6.1, it can be seen that the hierarchy ensures no double counting areas of elected or mandatory activities

This means that in all cases once land is accounted for under activities under Article 3, paragraph 3 and forest management, cropland management or grazing land management under Article 3, paragraph 4 reporting shall continue throughout subsequent and contiguous commitment periods (see CRF table NIR 2, para 2(d) in annex II or decision 2/CMP.8).

11.1.6.1 The information on identifiable units of land under mandatory activities and spatial assessment units.

The NFI is the primary data source used to identify areas under ARD and FM (Ch 6 section 6.3.2.3). The primary classification of forest land in the NFI uses approach 2 as defined in Chapter 3, [Section 3.3](#) of the 2006 IPCC Guidelines for AFOLU for the representation of land areas for Article 3.3 activities and FM. The spatial assessment unit used in the NFI and for reporting is the area of permanent sample plots (0.05ha). The same assessment unit is used to determine AR, D and FM areas and these are not larger than 1ha in accordance with paragraph 3 in the annex to decision 2/CMP.7.

The NFI used information from the IFORIS system (Ch 6, 6.3.2.2) to identify forest lands afforested after the 31st of December 1989. This enables the inventory to distinguish between activities under AR and those occurring under FM.

A secondary classification of identified forest land is carried out at the plot inventory phase in the NFI. This secondary classification is done to verify that the forest definition was correctly applied and interpreted during the photo interpretation phase 1 and to further classify forest areas into areas under planted forest areas, temporary un-stocked forest areas, open areas with forest boundaries (e.g., rides, roads etc.). Forest stand attributes from the NFI were also collected to classify forest age, rotation stage (i.e. thicket, pre thinning, thinning cycle or rotation cycle), and management status so that inventories plots could be disaggregated into appropriate KP forest categories.

The hierarchical classification system used to define land use areas (see Ch 6 section 6.2.2.1) and the level 2 classification of forest lands ensured that land areas are not double counted and that CSC are calculated based on the spatial assessment unit for all activities.

Activities under ARD or FM are reported within the entire territory of Ireland, with further sub division into species/forest type strata (Table 11.5) within internal national boundaries.

In years when NFI inventory data is not available (i.e. before the 1st or after the last NFI) afforestation areas are tracked on a spatially explicit basis (IPCC Approach 3, see section 6.3.2.5) using while deforestation areas are identifiable but not spatially explicit (IPCC Approach 2, see section 6.3.2.4). Both approaches can detect a land use change at a resolution consistent with the

forest definition area of 0.1 ha. For deforestation activities, (5(KP-I)A.1.2) areas are stratified according to land use activities converted from forest area. This is consistent with forests converted to non-forest lands in LULUCF Convention reporting (CRF 4 B, C, D and E).

Table 11.5 Forest category codes used in CRF Tables 5 (KPI)

Forest_Category_Code	Forest_Category_Description
1	Spruce (Pure). Mainly Sitka and Norway spruce
2	Pine (Pure). Predominantly Scots and lodgepole pine
3	Larch (Pure)
4	Other conifers (Pure)
5	Fast growing broadleaves (Pure) such as ash, Alder, Sycamore, Birch
6	Slow growing broadleaves (Pure) such as Oak and Beech
7	Conifer mixes
8	Broadleaf mix
9	Conifer/Broadleaf mix
10	Open areas including biodiversity areas, roads within the forest boundary
11	Blown areas subjected to windthrow
12	Scrub, felled or failed areas (planted and unplanted)
13	New afforestation after 2006
14	Natural succession and regenerating land
101 to 115	Harvested areas. E.g 101 are harvested spruce areas
200	Burned areas

Forest stands were considered to be pure if one species represents 80 % or more of the canopy

Note: Categories 12 and 14 do not qualify as afforestation or reforestation under Article 3.3 of the KP, so are reported as NO in the CRF tables. Open areas are planned open areas in afforested areas for extraction roads of biodiversity enhancement. These are, however, assumed to be in steady state and reported as NO.

11.1.6.2 The information on identifiable units of land under elected activities and spatial assessment units.

The Land Parcel Information System is the primary data source used to identify areas under CM and GM (Ch 6, section 6.4.4).

The primary classification of cropland and grazing land uses approach 3 as defined in Chapter 3, [section 3.3](#) of the IPPC good practice guidance for AFOLU for the representation of land areas. The classification is based on the explicit spatial boundaries of land parcels with a tolerance of 0.1ha for each parcel. The same assessment unit is used to determine AR, D and FM areas and these are not larger than 1ha in accordance with paragraph 3 in the annex to decision 2/CMP.7.

The Land Parcel Information Systems is maintained by the DAFM, and is compatible and spatially consistent with the IFORIS system (Ch 6, 6.3.2.2). This enables the inventory to distinguish between activities under AR and conversion of CM and GM lands to AR. It also allows tracking of deforestation D to cropland and grassland land uses, which are reportable under Art 3.3 D.

The hierarchical classification system used to define land use areas (see Ch 6, section 6.2.2.1) and the level 2 classification of forest lands ensured that land areas are not double counted and that CSC are calculated based on the spatial assessment unit for all activities.

Activities under CM or GM are reported within the entire territory of Ireland.

At present, the analysis of the conversion of CM and GM to non-Art 3.4 activities, for example Settlement, is not spatially explicit. However, whilst the conversion to an alternative land use is captured, through adjustments in the land parcel data, the new-land use is not spatially explicit. Additional research is planned to explore planning and urban databases to address this.

11.1.7 Information that demonstrated elected activities are directly human induced and have occurred since 1990

Article 3.3 activities

Reforestation activities do not occur in Ireland and the relevant activities under Article 3.3 are limited to afforestation and deforestation. All afforested areas are a result of direct planting and establishing forest areas under guidelines of the Forest Service Grant and Premiums Scheme since the beginning of 1990 (Forest Service, 2003). The afforestation grant and premiums scheme was introduced under European Commission Council Regulation 2080/92 to support afforestation of agricultural land as part of accompanying measures to reform the Common Agricultural Policy. The afforestation grant and premiums dataset captures all areas afforested following successful grant application. All afforestation areas recorded by the Forest Service are verified using a strict control and referrals process, following a post establishment site visit by a forestry inspector (Forest Service 2003). Afforestation areas recorded by the Forest Service are verified using a strict control and referrals process, following a post-establishment site visit by a forestry inspector (Forest Service, 2003).

All deforestation areas are derived from legally-binding licence applications under the Forestry Act. These provisions fulfil the requirement to demonstrate that afforestation (i.e. planting of non-forest land with trees for development of the forest sector) and deforestation began on or after 1 January 1990 and are directly human-induced, which is necessary for the accounting of emissions and removals for activities under Article 3.3. These datasets were primarily digitised using the 1:12560 and 1:2500 Ordnance Survey Ireland (OSI) raster maps (see section 6.3.2 in Chapter 6).

Article 3.4 Forest management

All areas subject to FM activities are managed (see definition) and these are distinguished from ARD activities based on year of initial establishment (see 11.1.5). Areas under forest management can be categorised into the following:

1. *Land owned by the State forestry company- Coillte.* These areas [ca. 89 per cent of the total FM area] are subject to management plans (FMPs), forest inventory and routing application for felling licences under the Forestry Bill 2013. These management plans are updated every 5 years to ensure that timber future demands are met. This information is used for timber forecasting and securing timber sales. Other management activities carried out include pest and disease control, crop nutrition, biodiversity management, nutrient loading and runoff management, riparian woodland management etc. The Coillte FMP 2011-2015 for each forest region can be downloaded from:
http://www.coillte.ie/coillteforest/plans/forest_management_plans/forest_management_plans_2011_2015/.
2. *Private grant aided afforestation from 1985-1989:* These areas of forest were established before 1990 under the Forest Service Grant and Premiums Scheme. The application procedure for the scheme requires a detailed forest management plan for a 20 year period. In addition, the Forestry act requires management plans for clearfell, thinning and replanting operations, which are submitted when applying for felling licences, required under the new Forestry Bill 2013.

3. *Private grant aided forests since 1920*: These represent very small areas of existing forests which were grand aided under various other small schemes before 1985. Evidence of forest management activities for these lands include:
 - a) The 1975 private forest survey provided detailed management plans for all private forests.
 - b) The Forestry Bill 2013 requires management plans for clearfell, thinning and replanting operations, which are submitted when applying for felling licences.
 - c) The Forestry Bill 2013 aims to promote the development of forests and forest-related activities and industries in such a way that forests provide an economically, environmentally and socially sustainable yield of forest goods and services, while maintaining and enhancing their biological diversity. There are numerous sections in the act, which refer to specific management plans. For example section 10(1) “The Minister may, by notice in writing, require an owner of a forest to submit a forest management plan in respect of his or her forest to ensure that afforestation, felling, restocking, forest road works and other forestry related activities (including amenity and recreation uses of forestry) are being carried out in accordance with good forest practice”.
4. *Existing forests before 1920*: These include old estate forests and semi-natural forests/woodlands in existence before 1920. Specific forest management activities include those outlined under item 3 above and include:
 - a) Management guidelines from the ‘Woodlands of Ireland’ (<http://www.woodlandsofireland.com/>) with the objective to generate awareness of native woodlands amongst policy makers and the general public and to develop projects and sustainable management strategies aimed at ensuring the future viability of native woodlands.
 - b) Existing semi natural woodland grants also require submission of detailed management plans on application for grant aid.

Article 3.4 Cropland Management

All areas subject to CM activities are managed (see definition), in private ownership, and these are distinguished from GM activities based on history of land parcel use since 2000. The data which forms the basis of the analysis of CM is collected as an integral and mandated component of the Farm payments systems under the EU Common Agriculture Policy, which is under the administration of the Department of Agriculture, Food and Marine. In order to satisfy the regulatory controls under the CAP, all aspects of the farm payments system is subjected to rigorous auditing, including the mapping and spatial attribute data.

Areas under cropland management can be categorised into the following:

1. *Croplands under continuous tillage*

These are land parcels which are identified as having declared as tillage crops in all years since 2000, and with the assumption that these parcels were managed in a similar manner since 1990. It is reasonable to assume these lands are in long term equilibrium with respect to carbon pools. These lands tend to be in the direct ownership of the land manager/farmer.

2. Croplands under regular rotation with temporary grassland

These are land parcels which are identified as having declared as tillage crops in many of the years since 2000, and with the assumption that these parcels were managed in a similar manner since 1990. These lands tend to be in the direct ownership of the land manager/farmer, either in mixed farming systems or on relatively poor soils which need of frequent fallow periods to maintain fertility. However, a significant proportion may be subject to short term leasing arrangements, whereby the land manager can vary, with switching in land use or short periods, and little consideration of the long term sustainable management of the land. These lands are not in long term equilibrium with respect to carbon pools.

3. Lands under crops occasionally or infrequently, generally under grassland

These lands have spent far longer periods under grass than under crops during their recent histories. It is very likely that these lands are subject to very short term leasing arrangements between livestock and tillage farmers. Their carbon pools are likely close to the long term equilibrium condition of permanent grasslands.

4. Croplands converted to Settlement

Conversion of agricultural land to Settlement is relatively common in rural and suburban settings. These lands remain reportable under Art 3.4 activities.

Article 3.4 Grazing land Management

All areas subject to GM activities are managed (see definition) and these are distinguished from Cm activities based on a history of permanent grassland. Areas under grazing land management can be categorised into the following:

1. Grasslands identified as permanent grasslands

Grazing land is subdivided into improved grasslands, consisting of lands reported as managed as pasture, silage and hay.

Unimproved grasslands, consisting of lands reported as managed as rough grazing.

Semi-natural grasslands, consisting of unmanaged but accessible grasslands, which have been identified as in agricultural use at some stage since 1990.

2. Grasslands converted to Settlement

Conversion of agricultural land to Settlement is relatively common in rural and suburban settings. These lands remain reportable under Art 3.4 activities.

11.2 Methodologies and description of data

For detailed description of sources of activity a data and methods used please refer see sections 6.3.1 to 6.3.2 in Chapter 6).

11.2.1 Afforestation

See Chapter 6, sections 6.3.2.1 to 6.3.2.3, 6.3.2.5. For detailed data on areas over the time series see Tables 6.5.12, 6.5.13 and 6.5.14.

A more detailed description of removals and emissions associated with different forest types in 2013 (defined in table 11.5 above) is shown in table 11.6.

11.2.2 Forest management

See Chapter 6, sections 6.3.2.1 to 6.3.2.4. For detailed data on areas over the time series see Tables 6.5.4, 6.5.5 and 6.5.7.

A more detailed description of removals and emissions associated with different forest types in 2013 (defined in table 11.5 above) is shown in table 11.7.

Table 11.6 Detailed areas and CSC for level 3 forest categories under AR activities in 2013

Forest_Cat	Soil_Cat	kHa	kt C												kt_CO ₂
		Area	AG_GAin	AG_Loss	AG_Net	BG_GAin	BG_Loss	BG_Net	Litter_Net	DW_Net	Org Soils	DOC soils	Soils_Net	Total C	
1	Org	53.103	332.323	-85.487	246.835	65.240	-2.946	62.294	34.038	6.259	-31.331	-16.462	-47.792	301.634	-1105.990
1	Org_Min	29.630	156.408	-41.369	115.039	30.878	-1.343	29.535	13.514	4.853	-8.417	-9.185	-17.602	145.339	-532.911
1	Min	49.699	394.241	-131.449	262.792	81.166	-7.316	73.850	40.201	15.078	NO	NO	NO	391.922	-1437.046
2	Org	4.718	17.217	-4.391	12.826	3.868	-0.300	3.568	1.046	-0.043	-2.784	-1.463	-4.246	13.151	-48.220
2	Org_Min	0.629	0.628	-0.055	0.573	0.067	0.000	0.067	0.010	-0.019	-0.221	-0.195	-0.416	0.215	-0.788
2	Min	1.573	6.079	-1.971	4.109	1.376	0.000	1.376	1.192	0.000	NO	NO	NO	6.677	-24.481
3	Org	3.460	6.890	-2.568	4.322	1.907	-0.231	1.676	-0.527	0.676	-2.041	-1.073	-3.114	3.034	-11.124
3	Org_Min	2.831	1.084	-0.183	0.901	0.197	0.000	0.197	-0.037	-0.004	-0.852	-0.878	-1.730	-0.672	2.465
3	Min	7.235	17.235	-3.777	13.458	4.783	0.000	4.783	0.164	-0.182	NO	NO	NO	18.223	-66.817
4	Org	0.315	1.656	-0.158	1.498	0.199	0.000	0.199	0.068	-0.068	-0.186	-0.098	-0.283	1.414	-5.184
4	Org_Min	0.629	0.001	0.000	0.000	0.000	0.000	0.000	0.000	-0.002	-0.252	-0.195	-0.447	-0.447	1.640
4	Min	1.887	10.837	-2.320	8.517	4.244	0.000	4.244	1.122	-0.001	NO	NO	NO	13.883	-50.903
5	Org	2.089	1.497	-1.265	0.232	0.160	0.000	0.160	0.669	-0.217	-1.233	-0.648	-1.880	-1.038	3.805
5	Org_Min	3.258	0.272	-0.026	0.246	0.199	0.000	0.199	1.027	-0.193	-0.586	-1.010	-1.596	-0.318	1.166
5	Min	9.751	0.030	-41.681	-41.650	0.341	0.000	0.341	20.358	-0.206	NO	NO	NO	-21.157	77.574
6	Org	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
6	Org_Min	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
6	Min	5.033	5.883	-4.256	1.627	1.314	-0.025	1.289	0.293	0.366	NO	NO	NO	3.575	-13.109
7	Org	11.280	60.876	-29.515	31.361	12.943	-3.688	9.255	8.573	13.086	-6.655	-3.497	-10.152	52.123	-191.118
7	Org_Min	4.133	20.499	-5.156	15.343	4.170	-0.329	3.841	1.455	0.660	-1.070	-1.281	-2.351	18.948	-69.477
7	Min	6.606	31.422	-8.754	22.667	6.949	-0.622	6.327	3.113	2.262	NO	NO	NO	34.370	-126.022
8	Org	0.629	0.737	-0.566	0.170	0.022	0.000	0.022	0.284	-0.053	-0.371	-0.195	-0.566	-0.143	0.523
8	Org_Min	0.629	1.192	-1.127	0.065	0.170	0.000	0.170	0.577	-0.087	-0.189	-0.195	-0.384	0.341	-1.252
8	Min	1.887	1.832	-4.984	-3.152	0.538	0.000	0.538	2.458	-0.560	NO	NO	0.000	-0.716	2.624
9	Org	8.066	19.937	-21.494	-1.557	3.800	-0.292	3.508	9.928	0.638	-4.759	-2.500	-7.259	5.258	-19.280
9	Org_Min	3.887	6.772	-3.467	3.305	1.487	-0.136	1.351	1.145	0.222	-1.741	-1.205	-2.946	3.076	-11.278
9	Min	10.066	33.548	-44.979	-11.431	8.012	-2.508	5.504	17.576	2.735	NO	NO	NO	14.384	-52.742
11	Org	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
11	Org_Min	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
11	Min	0.315	0.096	-0.014	0.082	0.028	0.000	0.028	-0.023	-0.034	0.000	0.000	0.000	0.053	-0.194
10	Org	13.717	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
10	Org_Min	2.005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
10	Min	19.066	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
13	Org	14.403	9.202	-0.875	8.326	8.094	0.000	8.094	0.655	0.000	-8.498	-4.465	-12.963	4.112	-15.078
13	Org_Min	4.895	1.624	-0.154	1.469	1.428	0.000	1.428	0.116		-2.398	-1.517	-3.916	-0.903	3.309
13	Min	21.454	12.035	-1.145	10.890	10.586	0.000	10.586	0.856		0.000		0.000	22.333	-81.886
200	Org	0.194	NO	NO	NO	NO	NO	NO	NO	NO	-0.115	-0.060	-0.175	-0.175	0.642
200	Org_Min	IE	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
200	Min	IE	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
		299.070	1152.051	-443.187	708.865	254.167	-19.736	234.431	159.852	45.167	-73.697	-46.121	-119.818	1028.496	-3771.152

Table 11.7 Detailed areas and CSC for level 3 forest categories under FM activities in 2013

Forest_Cat	Soil_Cat	Kha	kt C												Net_kt_CO ₂
		ADJ_Area	AG_GAin	AG_Loss	AG_Net	BG_GAin	BG_Loss	BG_Net	Litter_Net	DW_Net	Org soil	DOC	Soils_Net	Net_kt_C	
1	Org	93.461	688.079	-391.776	296.302	164.164	-42.900	121.265	64.011	11.924	-19.162	-28.973	-48.135	445.367	-1633.012
1	Org/Min	14.436	221.358	-293.762	-72.404	55.787	-52.063	3.724	40.469	12.891	-3.572	-4.475	-8.047	-23.367	85.680
1	Min	38.368	233.596	-86.930	146.666	58.344	-3.689	54.654	-8.907	-4.243			0.000	188.170	-689.957
2	Org	38.905	175.765	-86.222	89.543	46.065	-9.678	36.388	3.046	0.069	-9.400	-12.061	-21.461	107.584	-394.476
2	Org/Min	3.985	30.685	-12.496	18.189	7.926	-0.741	7.185	2.300	-0.199	-1.034	-1.235	-2.269	25.206	-92.421
2	Min	6.979	37.308	-30.164	7.143	9.971	-5.016	4.955	4.789	0.734			0.000	17.621	-64.611
3	Org	1.498	0.344	-7.402	-7.058	0.049	0.000	0.049	0.568	-0.202	-0.374	-0.465	-0.839	-7.482	27.434
3	Org/Min	0.498	0.730	-7.837	-7.106	0.256	-2.848	-2.592	-0.053	0.292	-0.113	-0.155	-0.267	-9.726	35.663
3	Min	1.495	2.824	-6.158	-3.333	0.950	-1.398	-0.448	-0.648	0.181			0.000	-4.249	15.580
4	Org	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000	0.000	0.000
4	Org/Min	0.498	0.176	-10.183	-10.006	0.038	-3.579	-3.541	0.401	1.041	-0.107	-0.155	-0.261	-12.366	45.343
4	Min	1.495	9.997	-8.495	1.502	3.310	-2.159	1.151	0.397	1.857			0.000	4.907	-17.992
5	Org	2.495	3.076	-17.135	-14.059	0.500	0.000	0.500	1.762	0.000	-0.673	-0.774	-1.446	-13.243	48.557
5	Org/Min	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000	0.000	0.000
5	Min	9.471	0.347	-139.211	-138.864	0.102	-0.168	-0.066	13.836	0.721			0.000	-124.373	456.033
6	Org	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000	0.000	0.000
6	Org/Min	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000	0.000	0.000
6	Min	5.982	28.870	-25.893	2.977	8.481	0.000	8.481	2.854	-1.309			0.000	13.003	-47.676
7	Org	20.437	137.871	-72.156	65.715	34.592	-7.973	26.619	3.319	-1.592	-4.897	-6.336	-11.232	82.828	-303.703
7	Org/Min	2.976	58.333	-20.535	37.799	14.686	-1.174	13.513	-1.473	-0.774	-0.627	-0.922	-1.549	47.514	-174.219
7	Min	10.969	64.910	-96.616	-31.706	17.091	-20.392	-3.300	4.134	4.685			0.000	-26.188	96.024
8	Org	1.994	1.121	-18.783	-17.662	0.329	0.000	0.329	1.861	-0.106	-0.529	-0.618	-1.147	-16.724	61.323
8	Org/Min	0.997	0.869	-10.539	-9.670	0.255	-0.001	0.254	0.812	-0.037	-0.212	-0.309	-0.521	-9.161	33.590
8	Min	22.911	50.623	-314.959	-264.335	14.871	-0.478	14.393	30.867	-3.097			0.000	-222.172	814.631
9	Org	24.456	72.745	-89.663	-16.918	18.070	-2.012	16.058	7.622	0.081	-6.015	-7.581	-13.597	-6.754	24.765
9	Org/Min	1.333	19.933	-38.035	-18.102	5.026	-0.586	4.440	3.192	0.369	-0.277	-0.413	-0.690	-10.792	39.569
9	Min	28.637	141.164	-323.489	-182.326	38.826	-26.408	12.418	21.794	6.971			0.000	-141.143	517.526
10	Org	22.569	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
10	Org/Min	3.244	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			0.000	0.000	0.000
10	Min	17.836	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			0.000	0.000	0.000
11	Org	3.489	17.219	-5.714	11.505	4.667	-0.391	4.276	-2.243	-0.608	-0.893	-1.082	-1.975	10.955	-40.168
11	Org/Min	0.498	3.800	-0.614	3.186	1.071	0.000	1.071	0.181	-0.167	-0.099	-0.155	-0.253	4.018	-14.731
11	Min	1.495	8.464	-3.404	5.059	2.330	0.000	2.330	1.220	-0.526			0.000	8.084	-29.640
12	Org	13.957	1.325	-135.150	-133.825	0.423	-0.092	0.331	8.975	0.172	-3.484	-4.327	-7.810	-132.159	484.582
12	Org/Min	1.495	0.187	-23.340	-23.152	0.044	-0.049	-0.005	2.326	0.214	-0.321	-0.464	-0.785	-21.402	78.475
12	Min	19.440	5.540	-188.477	-182.937	2.041	-1.581	0.461	7.844	-0.429			0.000	-175.062	641.893
13	Org	3.200	0.358	0.000	0.358	0.062	0.000	0.062	0.000	0.000	0.000	-0.992	-0.992	-0.572	2.099
13	Org/Min	0.800	0.128	0.000	0.128	0.031	0.000	0.031	0.000	0.000	0.000	-0.248	-0.248	-0.090	0.328
13	Min	4.000	0.479	0.000	0.479	0.082	0.000	0.082	0.000	0.000			0.000	0.562	-2.059
14	Org	7.976	8.506	-1.989	6.517	1.107	0.000	1.107	-5.925	-0.345	-1.923	-2.472	-4.396	-3.042	11.153
14	Org/Min	2.979	6.448	-2.715	3.734	1.268	0.000	1.268	-3.129	-0.423	-0.661	-0.923	-1.584	-0.135	0.497
14	Min	11.969	14.805	-12.146	2.659	2.615	0.000	2.615	-2.244	-0.840			0.000	2.189	-8.028
200	Org	0.309	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.182	-0.096	-0.278	-0.278	1.020
200	Org/Min		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			0.000	0.000	0.000
200	Min		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			0.000	0.000	0.000
		449.535	2047.983	-2481.989	-434.005	515.431	-185.376	330.056	203.957	27.302	-54.555	-75.229	-129.784	-2.474	9.071

11.2.3 Deforestation

11.2.3.1 Information on how harvesting or forest disturbance that is followed by the re-establishment of a forest is distinguished from deforestation.

Ireland provides information on how lands subject to harvest or disturbance followed by re-establishment is distinguished from deforestation as required under paragraph 4b of the annex II to the decision 2/CMP.8. A forest area is classified as deforested when there is clear indication of a specific land use change for that area or if clearfelled areas have not been replanted within a period of 5 years. Whilst different methodologies have been used to detect deforestation over time (Sections 6.3.2.4, and 6.3.6 in Chapter 6), this definition of deforestation has been applied consistently in developing the 1990 to 2013 area time-series.

The new NFI 2012 enabled the detection of all deforestation events including illegal deforestation and failure to replant felled areas within 5 years. The NFI completes an inventory of all forest areas every 5 years on a rotation basis. The 3rd NFI cycle is due to commence in 2016. ***If a clearfelled area has not been planted with a successive crop within one NFI cycle (i.e. 5 years), the area is classified under deforestation.*** These areas are to be reported for the year deforestation is detected. Under the felling licence rules all replanted crop must be inspected after 5 years to ensure a 95 per cent survival rate. Crop is then considered to be successfully established for the next rotation.

11.2.3.2 Deforestation Information

See Chapter 6, sections 6.3.2.1 to 6.3.2.3, 6.3.2.4. For detailed data on areas over the time series see Tables 6.5.16 and 6.5.17.

Information for deforested areas supplied with the limited felling license application/or from the NFI provides details of the species, areas, volume of timber clear felled and an indication of the applicable land use transition category (see CRF 5(KP-I)A.2) into the following (see Table 11.8):

Forest land to Grassland (**F-GL**)

Forest land to Cropland (**F-CL**), this does not occur.

Forest land to Wetland (**F-WL**)

Forest land to Settlement (**F-S**)

Forest land to Other land (**F-OL**), these include land not classified above such as quarries, windfarms

Biomass, litter and deadwood pools for deforestation land were assumed to be immediately oxidised in the year deforestation occurs (see Chapter 6). The changes in biomass and deadwood C pools stock for these deforested lands converted to other land uses in the subsequent years is assumed to be zero and reported as NO. This is because all forest C pools have been oxidised in the previous year.

Table 11.8 CSC for forestland converted to other land categories

(also see information item in CRF table 4(KP-1)A.2)

	2013						
Deforestation		FL	CL	GL	WL	S	OL
Activity data							
Area subject to the activity	kha	NO	NO	6.64	2.22	4.74	3.03
Area of mineral soils	kha	NO	NO	6.27	0.34	1.93	1.49
Area of organic soils	kha	NO	NO	0.37	1.88	2.81	1.54
Change in carbon stock							
CSC above-ground biomass							
Gains	kt C	NO	NO	0.08	IE	NO	NO
Losses	kt C	NO	NO	-2.62	NO	-2.06	NO
Net change	kt C	NO	NO	-2.54	NO,IE	-2.06	NO
CSC below-ground biomass							
Gains	kt C	NO	NO	0.36	IE	NO	NO
Losses	kt C	NO	NO	-0.63	NO	-0.57	NO
Net change	kt C	NO	NO	-0.27	NO,IE	-0.57	NO
Net CSC in litter	kt C	NO	NO	-0.32	NO	-0.35	NO
Net CSC in dead wood	kt C	NO	NO	-0.08	NO	-0.04	NO
Net CSC in soils							
Mineral soils	kt C	NO	NO	NO	NO	-2.08	-1.63
Organic soils	kt C	NO	NO	-1.46	-1.94	-2.14	-1.39
Net emissions/removals							
CO2	kt	NO	NO	17.16	7.12	26.54	11.06

IE for wetland biomass gains reported under soils as specified in 2013 Wetland supplement for rewetted soils CH 4.

NO for mineral soils, not occurring because soils are demonstrated not to be a source. For emissions from mineral settlement and other land soils see section 6.3.6.1.1 in Ch 6.

NO not occurring

Biomass gains are only reported for grasslands based on 2006 AFOLU GPG Table 6.4 Ch 6.

11.2.4 Cropland management

See Chapter 6, sections 6.4.1 to 6.4.4. For detailed data on areas over the time series see Figures 6.4.1, 6.4.2.

11.2.5 Grazing land management

See Chapter 6, sections 6.5.1 to 6.5.4. For detailed data on areas over the time series see Figures 6.5.1, 6.5.2.

The difference in total area under GM and those within the Grassland land use category relate to the exclusion of semi-natural grasslands existing prior to 1st Jan 1990, and any subsequent land use and land management activities which change the status of these lands.

In general, the trend has been towards an increase in semi-natural grasslands, in response to policy measures to reduce over grazing of uplands areas, and in response to market drivers and reduction in the profitability of livestock production on marginal lands.

The trend is mitigated by conversion of semi-natural grasslands to forest. Figure 11.2.1 Time series of grasslands classified according to GM definition and Other grassland (unmanaged semi natural). "Grazing land remaining Grazing land", and "Agricultural land to Semi-natural" sum to the total GM area. "Other Grassland" is unmanaged grasslands throughout the period, and are excluded from the definition of GM.




Figure 11.2.1 Time series of grasslands classified according to GM defintion and Other grassland (unmanaged semi natural)

11.2.6 Direct and indirect emissions from N fertilisation

Indicated elsewhere (IE) under Agriculture emissions.

11.2.7 N₂O and CH₄ from drained and rewetted organic soils

See sections 6.3.4.5, 6.3.5.6 and 6.3.6.12 in Ch 6. Note that CO₂ emissions from drained organic soils are reported as IE in tables 4(KP-I) A.1, A.2 and B.1

11.2.8 N₂O losses from mineralization of soils due C loss associated with land use change

For afforestation activities this does not occur (NO) because we demonstrate that CSC in mineral soils is not a source. This also applies for forest to grassland conversions (see Section 11.4 below).

For deforestation to settlement and other land, CSC in mineral soils are reported and N₂O emission are reported using tier 1 approaches (see 6.3.6.1.1, Ch 6)

11.2.9 Biomass burning from Forest ARD and FM Fires

Areas of forest subjected to wild fires were obtained from Forest Service statistics (see section 6.3.4.4 and Table 6.5.6 in Chapter 6). These areas were assumed to be proportionally distributed between the Kyoto Protocol forestry categories afforestation/reforestation and forest management. For example, in 2008 the AR area in Table 11.1 represented 36 per cent of the total forest area, so it was assumed that 36 per cent of areas experiencing wild fires in 2008 are in the AR category. This determines the area for estimating biomass burned under AR and FM (see Table 6.5.6 in Ch 6), reported in CRF Table 4(KP II)4. The same assumptions are applied to years subsequent to 2008.

11.2.10 Biomass burning from CM Fires

Areas of cropland subjected to wild fires were extrapolated from Forest Service statistics (see section 6.4.9 and Table 6.5.6 in Chapter 6), and remote sensor detection of fires from the NASA FIRMS database, (see section 6.4.9 in Chapter 6).

11.2.11 Biomass burning from GM Fires

Areas of grazing land subjected to controlled fires were extrapolated from Forest Service statistics (see section 6.5.4 and Table 6.5.3 in Chapter 6), and remote sensor detection of fires from the NASA FIRMS database, (see section 6.5.4 in Chapter 6).

11.3 Justification for Omitting a Carbon Pool

This section provides detailed information that demonstrates that the mineral soil pools under afforestation, forest management and forest land conversions to grasslands are not a net source of anthropogenic GHG emissions; therefore, the mineral soil pool is not accounted for and reported as NO. If a pool is not reported and demonstrated that it is not a net source then the approach is consistent with requirements under paragraph 2(e) of the annex to the decision 2/CMP.8.

It also outlines the justification for omitting Litter and Dead Litter pools from CM and GM activities.


11.3.1 Afforestation: Mineral Soils for grassland, cropland and forest land transitions

Demonstration that soils are not a source is based on numerous research data, which have been subject to rigorous statistical analyses using different approaches. **The working null hypothesis is that mineral soils are a source when lands are converted to forest land, and visa-versa. The null hypothesis can be rejected if statistical tests prove that mineral soils are not a source. Therefore, we demonstrate that mineral soils for cropland and grassland conversion to and from forestry by rejecting the null hypothesis.**

Changes in mineral soil C pools over time (ΔC_{so}) are not reported because of all current information confirms that it is not a source. These findings are based on three research approaches, but this is re-evaluated every submission year following review of national research. There is also a new project ForCRep (2012-2016) specifically initiated to further develop a method for reporting mineral soils stock changes.

a) A chronosequence approach:

National research information does suggest that mineral soils are a sink for a minimum of 50 years following afforestation (Black et al., 2009b). These authors show that SOC is higher when a 9 year old stand was compared to year 0 (i.e. un-forested grassland in the chronosequence). Other information from 30 different afforested sites, previously located on grasslands, suggest that there is *no significant* change ($P>0.1$) of mineral soil C stocks over time following afforestation (Black, 2008 see Figure 11.1). If anything, the data suggest an increase in SOC stocks over time, suggesting that afforestation of mineral soils results in a net sink, albeit not significant. Therefore, we opt not to report stock changes for mineral soils because we can **reject the null hypothesis and demonstrate that the pool is not a source**. However, this can only be applied to grassland/forestland conversions, so additional data and analysis was required.



The solid line represents the linear change on C stock over time. The dashed and dotted lines represent the 95% confidence and prediction intervals

Figure 11.1 Variation in mineral soil carbon stocks and estimation of ΔC_{so} using the nationally derived data ($n = 30$).

b) A paired plot approach:

A National forest research (FORESTSOIL C and CARBiFOR2 projects 2007-2012) designed a soil carbon monitoring system for Ireland using country-specific land use and soil carbon information. The system is based on a stratified NFI sample of the country by soil type and land use. This element of the work concentrated on a paired plot approach to assess soil C stock changes due to afforestation and deforestation activities. The sampling strategy was designed to augment NFI plot measurements, but included an additional, paired plot, samples from adjacent non-forest land uses. The overall concept applies the assumption that changes in soil C stocks, due to transition from one land use to another, is a function of the difference between the forest and non-forest plot soil carbon pools and that both soil pools are in steady state. For this reason, all land use samples were assessed to have not undergone land use transitions in the past 20 to 50 years.

The following assumptions and conditions were applied:

- The analysis only applies to mineral soils. Organic soils stock changes are determined using emission factors. This includes organo-mineral soils, such as peaty-gley soils (see equations 6.3.23 and 6.3.24 in section 6.3.3.1.2(d) in Chapter 6);
- Carbon stock changes in mineral soils for all sample plots were at steady state when sampled;
- The age at steady state (i.e. the mean age of the land use or soil type) is equivalent to the transition time for soil C stocks to reach steady state;
- The land use transitions did not include wetlands or croplands because transitions between forestry and these land uses (and vice versa) because they were not detected in the random stratified sample grid of 60 out of 1762 NFI sample plots.(section 6.3.2.4 and Annex 3.4.A.2). In addition there are very few mineral soils in wetlands;

- Settlement and other land soils were not sampled because of technical difficulties in obtaining soil samples. Deforestation and transition to settlements and other land does occur in Ireland but the soils stocks are assumed to be a source and are reported (see section 11.4.1.1). No paired settlement or other land plots were identified in the random stratified sample taken in this study;
- This analysis is primarily concerned with transitions between forestry, scrub, un-managed grassland and managed grassland. Scrub in this case refers to land uses dominated by non-tree species such as gorse or bramble. These in effect are degraded or disused grasslands (un-managed grasslands), previously used for rough grazing;
- Changes in soil C stocks due to land use change is assumed to occur only if the difference between the forest and non-forest pair, within a given soil group is found to be significantly different following statistical analysis.

To quantify the relative importance of the different factors on Ireland's soil C stocks, and to test the null hypothesis for different land use and soil strata, we carried out a hierarchical analysis of variance and multiple regression analysis using SPSS statistical package.

Table 11.9 Record sample plots taken from mineral soils sites

Soil	Land use	Transition time (years)	Number of plots	Number of soil profiles
Gleys	Forest	30	10	50
	Un-managed grassland	30	4	20
	Managed grassland	30	4	20
	Scrub grassland	30	2	10
Brown earths	Forest	35	10	50
	Un-managed grassland	35	6	30
	Managed grassland	35	4	20
	Scrub grassland			
Brown podsols	Forest	50	10	50
	Un-managed grassland	50	6	30
	Managed grassland	50	4	20
	Scrub grassland	50		
Podsols	Forest	39	10	50
	Un-managed grassland	39	5	25
	Managed grassland	39	4	20
	Scrub grassland	39	1	5
Total			80	400


Land uses were first categorized as shown in Table 11.9, and then re-classified either as forest or grassland (Grassland/Forest pair). The forest grassland pair was categorised because there no significant difference soils stock or changes in the different grassland types.

Table 11.10 Results from the hierarchical analysis of variance on soils C at a depth of 0-30cm

Source	SS	MS	F	P
Between soils	763902	254634	3.49	<0.01
Land use within soils	29663	2963	1.24	0.34
Grassland/Forest within soils	20215	4043	0.81	0.48

Based the hierarchical analysis of variance on soils C at a depth of 0-30 cm, it was evident that there was a significant difference in soils C stock when soil types were compared. However, there was no difference in the soil C stock when the different land use classes were compared (Table 11.10). Therefore, the null hypothesis is rejected (i.e. the p value for the Land-use source of variance is >0.05). **This confirms that mineral soils are not a source since there is no difference in the soil C stock when forests, un-managed grasslands, managed grasslands and scrublands are compared.**

To further illustrate this point, Figure 11.2 shows that the mean soil C stock was significantly different within soil types for combined data from forests and grasslands (left panel Figure 11.2). Although there were marginal differences between the mean soil C stocks when forest and grassland plots are compared within the soil categories, these were not significantly different (right panel in Figure 11.2).



Histogram bars with different letters are significantly different at $P < 0.05$

Figure 11.2 Mean soil C reference values for forest and grasslands at steady state across different mineral soil types

Based on these analyses and the chronosequence soil stock changes (Black et al., 2009; NIR, 2010) it is evident that there is no significant change in soil C stocks for up to 30 years following transitions between all grassland types and forest land. Therefore **the null hypothesis is rejected and we clearly demonstrate that this pool is not a source for grassland/forest land transitions.** Similar results have been reported by Davis et al (2002) and Scott et al (2002) for studies conducted in New Zealand, where many more plots were sampled.

Based on these conclusions and related publications (Wellock et al., 2011) Ireland has elected not to account for mineral soil C stock changes following afforestation and deforestation from and into grassland uses, because we demonstrate that this pool is not a source across different mineral soil types. However, more research was required to demonstrate this for cropland/forest land conversions.

c) A new SOC database from the For CRep project:

In 2013, the ForCRep project (2012-2016) compiled a national database using all available SOC survey data sampled from mineral soils representing major land use categories in the republic of Ireland. The mineral soil database comprises of a total of 227 sample sites obtained from the Soil C project (Wellock et al., 2011), the Irish national soil database of Ireland (NCD, see Xu et al., 2011), the An Foras Taluntas project (Creamer, R., unpublished data), and the CARBiFOR project (Black et al 2009a). For this study it is not yet possible to make comparisons across different soil types because detailed soil profile data is still being processed and collected. However, a preliminary analysis was conducted to test the null hypothesis if mineral soils are a source across all cropland/ grassland and forest land transitions.

Analysis of variance confirmed that the **null hypothesis can be rejected for cropland to forest land conversions**. The SOC in forest mineral soils is significantly higher (37 per cent), when compared to croplands, confirming that these soils are a sink (Figure 11.3). In the **case of forest land conversions to cropland, the null hypothesis can be accepted** since mineral soils could represent a significant loss of SOC (i.e. a source). However, there are no deforestation activities and conversion to cropland in Ireland (see CRF table KP.A.2). This is primarily due to the poorer quality of land under forestry, which is not suitable of cropland production. Cropland conversions to forestry have declined in recent years because previous transitions represented marginal land not suitable for cropland production.

Based on a comparison of mean SOC values for all grasslands and forest land, there was no statistical difference in the mean SOC for these land categories (Figure 11.3). Therefore, **the null hypothesis is rejected and SOC in following grassland/forest land transitions are demonstrated not to be a source**.




Figure 11.3 Comparison of mean SOC values (histograms) and standard deviations (error bars) across different land use categories. Mean values with different alphabetical characters indicate a significant difference ($p < 0.05$)


11.3.2 Forest management: Mineral Soils

Mineral soils emissions were assumed to be zero for the Forest Management Reference Level (FMRL), so the same assumption is applied to reporting of CSC in mineral soils under FM in the 2nd commitment period.

Although there was no supporting information for this assumption under the FMRL review under 2/CMP.6, we now provide additional information supporting this assumption based on the new SOC database from the For CRep project.

a) First rotation forests older than 20 years old:

The ForCRep database was used to investigate the SOC stock changes of different mineral soils associated with 1st rotation crops older than 20 years old. Data were stratified into 4 basic soils groups due to significant differences in SOC between soil types. Based on regression analysis it was confirmed that CSC mineral soils in 1st rotation forests older than 20 are not a source (Figure 11.4). In contrast, the results suggest that CSC in gley and brown podsolic soils increase over the first rotation. This result is expected since an increased transfer of deadwood and litter carbon to soils as a result of thinning residues and mortality would result in an accumulation of recalcitrant C in soils (see Liski et al. 2005. Carbon and decomposition model YASSO for forest soils. Ecological Modelling 189, 168-182).



*Coefficients of determination (r²) values were significant when p values were > 0.05

Figure 11.4 Regression analysis showing trends in SOC stocks for different soils types as a function of forest age

b) Second rotation crops:

The national forest soil database only contains SOC data for one 2nd rotation site, so there is no reliable national information for 2nd rotation crops. Verifiable information supporting the assumption that 2nd rotation crops are not a source comes from literature reviews from regions with similar forest types and climatic conditions:

1. Zerva et al 2005. Soil carbon dynamics in a Sitka spruce (*Picea sitchensis* (Bong.) Carr.) chronosequence on a peaty gley. *Forest Ecology and Management*. 205: 227–240:- This study was conducted in Scotland in forests similar to those occurring in Ireland. The results show that SOC carbon stocks increased from 140 t C ha⁻¹ at the end of the 1st rotation to 249 tC ha⁻¹ at 30 years in the second rotation, so they are not a net source.
2. The effect of forest management on SOC varies considerably depending on management regime, species, climate harvest intensity etc. (see Waldchen et al 2013. The influence of changes in forest management over the past 200 years on present soil organic carbon stocks. *Forest Ecology and Management* 289(1):243–254). These authors report that there are no detectable effects of forest management on SOC. This study was based on 190 inventory plots taken from forests in Europe varying in age from 100-200 years old and with different management interventions from coppicing to selective harvest.
3. Many tier 3 modelling approaches (e.g. YASSO see Laiski et al., 2005) show that SOC generally accumulates in successive rotations due to increased inputs of C from harvest residues. In Ireland, there has been an age class shift from mature 1st rotation age forests to 2nd rotation forest (see Figure 6.3.9 Ch 6). This has resulted in an increase allocation of harvest residue to the dead organic matter pool, some of which is emitted due to decomposition. However, some of the recalcitrant C will accumulate in soils. The current hypothesis represented by the literature and the current age class structure of Irish forests under forest management suggest that mineral soils cannot be a net emission of CO₂. A study on organic soils in Ireland, soils which are more likely to result in emissions due to forestry, show that emissions from soils decline after the 1st rotation Byrne and Farrell, 2005).

It should be noted that the same approach for mineral soils are taken for reporting the FMRL and emissions/removals for FM in the 2nd commitment period. Therefore, this approach does not result in an underestimation of emissions associated with FM activities.

11.3.3 Cropland management Matter Carbon Pools

Based on the decision tree in Section 2.9.4.1 of the 2013 KP Supplement to the 2006 IPCC Guidelines, and Section 5.2.2.4, Vol 4 of the 2006 Guidelines, changes in Litter and Dead Matter carbon pools are assumed to be stable.

Changes in biomass associated with transitions between grassland and croplands within the CM cohort are estimated. Changes in the biomass of hedgerows, and other non-forest wood features, have not been estimated.

Biomass changes due to changes in the area of perennial woody crops are based on the analysis of the dominant crops, apple orchards and Christmas trees. In the case of Christmas, there is evidence that the market for trees is stable or increasing over time, and as such the biomass associated with this crop is stable or increasing, See section 6.4.7.

The area of apple orchard decreased in the early 1990s, but has been in near equilibrium over the last three census of the crop, see Figure 6.4.7.

Hedgerows are an integral part of the CM landscape. However, there is very limited long term monitoring data as to conditions and extent of these features. The EPA has funded a research project to pilot an analysis of historic and contemporary remote sensing data to establish a robust time series of changes in these landscapes. There is conflicting evidence as to the current trends in hedgerow and wooded area management within CM. Measures under planning guidelines, the Rural

Environment Payment Scheme, its antecedents and other policies, the maintenance, of existing hedgerows and establishment of new hedgerow has been encouraged. For example, under REPS 3 and 4 and AEOS 1 and 2 rural environmental protection schemes (see Figure 11.5), support was provided for establishment of approximately 10,000 km of new hedgerow (Teagasc Newsletter, Sep 2013¹⁰). However, the National Forest Inventory detected a decrease in hedgerow area of 4,548 ha between 2006 and 2012, albeit with a very large uncertainty (DAFM, 2013). At present it is not possible to provide a robust time series of hedgerows, and therefore it has not been possible to produce an estimate of biomass changes associated with their management.




Figure 41.5 Hedgerow established under a series of rural protection schemes

11.3.4 Grazing land management Carbon Pools

Based on the decision tree in Section 2.10.4.1 of the 2013 KP Supplement to the 2006 IPCC Guidelines, and Section 6.2.2.1, Vol 4 of the 2006 2006 Guidelines, changes in Litter and Dead Matter carbon pools are assumed to be in equilibrium and there is no requirement to estimate carbon stock changes in these pools, as a Tier 1 approach.

Biomass changes due to the transition between difference grazing land management types are interpreted as “not a source”. The main transitions which have occurred are:

- An increase in the intensive management of improved grasslands, which will tend to increase the productivity of the grazing land. However, this additional productivity is removed by harvesting for fodder or direct grazing. As such, any apparent increase in biomass is short lived. Average over the year, it is more likely than not to result in a slight increase in biomass, due to increased productivity. Therefore, the transition can reasonably be considered “not a source”.
- An increase in the area of land reverting from rough grazing to semi-natural grassland is inferred from the activity data. With the removals of grazing animals from rough grazing areas, there is a

¹⁰ http://www.teagasc.ie/publications/2013/2865/Environment_Newsletter_September2013.pdf
 Environmental Protection Agency 361

likely an increase in the biomass on these lands. However, this has not been quantified. However, it is reasonable to consider the transition to be “not a source” with respect to biomass.

Hedgerows are an integral part of the GM landscape. However, there is very limited long term monitoring data as to condition and extent of these features. The EPA has funded a research project to pilot an analysis of historic and contemporary remote sensing data to establish a robust time series of changes in these landscapes. There is conflicting evidence as to the current trends in hedgerow and wooded area management within GM. See section 11.4.2 for a more detailed discussion of available data. At present it is not possible to provide a robust time series of hedgerows, and therefore it has not been possible to produce an estimate of biomass changes associated with management of these

Factoring out Indirect and non-human induced emission/reductions

Ireland considers that all emissions/removals from Article 3.3 activities are directly human induced, since they are activities resulting from silvicultural intervention. No factoring out of indirect human-induced activities is considered in this submission due to a cited poor understanding of these influences (see Ainsworth and Long, 2005).

For FM, it is considered that the use of a forward looking baseline or FMRL factors out any non-human induced induction of emissions or removals such as N deposition, CO₂ fertilisation or age class legacies (see reviews carried out under 2.CMP6 and the Supplementary guidance on LULUCF 2013).

For CM and GM, Ireland considers emissions and removals to be dominated by, and indistinguishable from, the influence of management decisions and therefore all changes in emissions and removals to be directly human induced. By using Tier 1, emission factors, possible changes in the emissions and removals associated with direct impact of climate change or the fertilization effect of increased CO₂ levels are not included in the estimates. At this time, climate change impacts have not invalidated the use of temperate zone Tier 1 emission factors for Ireland¹¹, with an observed increase in temperature of 0.5°C between climate average for 1961-1990 and 1981 and 2010. Precipitation has been observed to increase slightly over the same period. Therefore Ireland remains in the “temperate, moist” climate zone. Therefore, “factoring out” is inherent, and estimates of emissions and removals are only driven by evaluation of human impacts on the activities.

11.4 Natural disturbances

Ireland indicates that we may wish to apply the provisions to exclude emissions from natural disturbances for the accounting for afforestation and reforestation under Article 3, paragraph 3, of the Kyoto Protocol and forest management under Article 3, paragraph 4, of the Kyoto Protocol during the second commitment period in accordance with decision 2/CMP.7, annex, paragraph 33a.

11.4.1 Calculation of background and margin

Emissions for wild fires may be excluded, if “triggered”, under the natural disturbance provision for both Article 3, paragraph 3, and Article 3, paragraph 4 activities. The calibration data to calculate the background level and margin includes the period 1990-2009, but has been extended to 2012 to account for most recent data (Table 11.11a and 11.11b). Wildfires have only been recorded in AR lands since 2008 and are assumed not to occur before 2008.

¹¹ Met Eireann, May 2012, Meteorological Note No. 14, A Summary of Climate Averages for Ireland 1981-2010
<http://www.met.ie/climate-ireland/SummaryClimAvgs.pdf>

Table 11.11a Total emissions form wild fires and area specific emissions from disturbances for the calibration period for FM

Year	Wildfires (kt CO ₂ eq.)	Insect /diseases	extreme weather	geological disturbances	other	Area (ha)
1990	113					389
1991	73					250
1992	47					161
1993	94					324
1994	108					372
1995	148					508
1996	164					565
1997	90					309
1998	47					163
1999	39					133
2000	97					334
2001	194					666
2002	45					153
2003	275					944
2004	160					550
2005	58					200
2006	58					200
2007	65					225
2008	51					175
2009	28					98
2010	185					636
2011	69					237
2012	17					60

Table 11.11b Total and area specific emissions from disturbances for the calibration period for AR

Year	Wildfires (kt CO ₂ eq.)	Insect /diseases	extreme weather	geological disturbances	other	Area (ha)
2008	17					99
2009	10					57
2010	64					377
2011	24					139
2012	6					35

The default method for estimating the background and margin for the natural disturbance provision is applied to the calibration data for both FM and AR disturbances using default approaches (Box 2.3.6 IPCC 2013 GPG KP-LULUCF). The final background and margin is derived at step VI, as outlined in the IPCC 2013 GPG KP-LULUCF (see table 8 below).

Table 11.12 Calculation steps used to derive the background and harvest for FM and AR for the second commitment period

(see Box 2.3.6 of IPCC 2013 GPG KP-LULUCF for detailed methods)

<i>kt CO₂ eq. for FM</i>						
Background/margin	I step	II step	III step	IV step	V step	VI step
Arithmetic mean	97	89	84	79	74	69
standard deviation	64	52	48	43	39	33
background+margin	225	193	179	164	151	136

<i>kt CO₂ eq. for AR</i>						
Background/margin	I step	II step	III step	IV step	V step	VI step
Arithmetic mean	0.09	0.09	0.09	0.09	0.09	0.09
standard deviation	0.08	0.08	0.08	0.08	0.08	0.08
background+margin	0.25	0.25	0.25	0.25	0.25	0.25

11.4.2 Trigger test for implementation of the natural disturbance provision

The reported emissions from wild fires under AR lands for 2013 were 26.99 kt CO₂eq, which is higher than the background plus margin (0.25 kt CO₂) presented in table 11.12. This means that Ireland can elect to exclude these emissions under the provisions set out under para 33 and 34 of the annex to 2/CMP.7 (i.e. trigger test in CRF table 4(KP-1) A1.1 should be “yes”). However, Ireland will not elect to avail of this provision for AR land disturbances at this time because the method used to track and record forest fires (approach 2) does not facilitate the ability to geo-reference the locations of wild fire events as required under para 2f(i) of annex II to the decision 2/CMP.8.

Ireland can show that these emissions from disturbances and subsequent removals following rehabilitation of lands can be excluded from accounting and fulfil all other required conditions outlined in the relevant decisions 2/CMP.7 and 2/CMP.8. Since Ireland has elected end of period accounting, possible approaches are being explored to georeferenced the location of wild fire events and Ireland may elect to exclude natural disturbance events from accounting, should this be triggered and appropriate methods are available.

The reported emissions from wild fires in lands under FM activities are lower than the background and margin presented in table 11.12. Therefore, Ireland cannot trigger the election of the natural disturbance provision in 2013 (i.e. trigger test in CRF table 4(KP-1) A1.1 should be “no”).

11.4.3 Exclusion of emissions from salvage logging

Salvage logging does not occur in lands subjected to forest fires. This is consistent with the assumption that all biomass and DOM is immediately oxidised when subjected to wild fire (see section 6.3.4.4 in Ch 6).

Ireland does not include disturbance events emissions associated with windthrow damage under the natural disturbance provision because all timber in windthrown areas is assumed to be recovered by salvage logging. Therefore, salvage logging will be excluded from natural disturbance emissions, should the provision be triggered (see para 2f(vi) in annex II of 2.CMP/8.). Emissions associated with windthrow (i.e. biomass, litter, deadwood etc.) are captured in the NFI inventory and included under

ARD or FM CSCs. Insect and disease infestations currently cause minimal emissions and are assumed to be captured by the inventory as reflected by the NFI permanent sample plot data.

11.5 FMRL and technical corrections

Ireland's forest management reference level (FMRL) as inscribed in the appendix to the annex to decision 2/CMP.7 is -142.07 kt CO₂ equivalent, see table 11.14 below.

Ireland has performed recalculations for the historic time series and 2013 and will apply a technical correction when accounting for the second commitment period. The requirement to apply a recalculation is based on conditions as outlined in the IPCC 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol (IPCC 2013 GPG KP-LULUCF):

- Use of new models to derive the reported carbon stock (CSC) changes in the inventory 2013. The same new model has been applied to the activity data used for the forest management reference level (FMRL) inscribed in the annex to 2/CMP.7. The new version of CARBWARE v5 is now used for the basis for CSC changes in biomass, litter and deadwood pools. The FMRL submission (see 2/CMP7) used CARBWARE version 4.5.
- There have been a range of methodological changes for estimation CO₂, N₂O and CH₄ emissions from organic and mineral soils as outlined in the 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands (IPCC Wetland supplement 2013). These include offsite DOC emissions of C, N₂O and CH₄ emissions due to drainage of organic soils.
- In accordance with Decision 2/CMP.6, Ireland's FMRL submission included a description of the domestic policies adopted and implemented no later than December 2009 and explain how these policies have been considered in the construction of the FMRL. Ireland confirms that the construction of the FMRL does not include assumptions about changes to domestic policies adopted and implemented after December 2009. However, Ireland has obtained new historical data for:
 - FM areas due to new deforestation data based on the repeat NFI in 2012 (See Table 4 below).
 - New historical harvest rates from FM areas before 2009 based on new national forest inventory (NFI) data (highlighted in bold in Table 11.13).

These data have been used to recalculate the historical and projected time series and a technical correction will be applied **when accounting** for FM activities.

- New historical activity data (prior to 2009) and new methods for HWP estimations have been applied. Therefore HWP pools emission reductions have been recalculated for the whole time series. The new methods include differentiation of domestic harvests from deforestation and those originating from FM lands remaining as FM land (IPCC 2013 GPG KP-LULUCF).

Table 11.13 A comparison of previous activity data for 1990-2009 used for the FMRL submission and the new data used in the 2013 inventory

Year	Area (Kha)		Harvests (M m ³)	
	FM 2013 inventory	FMRL submission	FM 2013 (this submission)	FMRL submission
1990	465.26	481.35	1.676	1.787
1991	465.24	479.91	1.767	1.837
1992	465.22	476.96	2.082	2.156
1993	465.20	479.40	2.097	2.003
1994	465.18	480.70	2.283	2.220
1995	464.84	474.14	2.377	2.424
1996	464.51	463.20	2.461	2.520
1997	464.18	461.33	2.313	2.398
1998	463.84	463.32	2.632	2.493
1999	463.51	466.75	2.765	2.842
2000	462.65	462.14	3.002	2.940
2001	461.79	456.02	2.822	2.700
2002	460.94	450.85	2.899	2.911
2003	460.08	458.81	2.986	2.951
2004	459.22	456.45	2.829	2.818
2005	458.37	454.35	2.925	2.775
2006	456.37	477.21	2.947	2.803
2007	454.77	444.52	2.864	2.160
2008	452.77	446.94	2.209	2.056
2009	451.97	447.29	2.682	2.392
2010	451.17		3.036	1.833
2011	449.57		2.729	2.061
2012	449.57		2.735	2.314
2013	449.53		2.827	2.390

Major reasons for the difference in the FMRLcorr and FMRL include:

- The new version of the CARBWARE model (v5) is a dynamic model which provides more accurate estimates of biomass, litter and deadwood CSC over time. The older version 4.5 of the model assumed that younger age classes (i.e. those less than 7 years old) exhibited a zero CSC.
- Higher CO₂ and non-CO₂ emissions from soils due to the new Wetland supplement methodology.
- Larger removals from the HWP pool in the FMRLcorr due to new methods applied and the higher historical harvest up to 2009, based on the new data. The allocation of HWP feedstock to FMRLcorr was also higher than the FMRL submission due to a change in methodology used.

Table 11.14 Summary of the FMRL recalculated (FMRLcorr) values for different pools in 2013, compared to the FMRL and reported FM (reported FM) emission/removal for 2013

Note that the FMRL shown is the value for 2013 not the annual average over the period 2013-2020 as indicated in the annex to 2/CMP7

	FMRLcorr 2013	FMRL 2013	Reported FM 2013
Biomass (kt C)	53.99	-75.88	-103.95
Litter	194.28	66.05	203.96
Deadwood	21.50	95.28	27.30
Soils	-130.08	-95.41	-129.78
Sub-total C-CO₂	-512.21	36.54	9.07
Fires CO₂ eq	12.00	12.00	72.50
N₂O and CH₄ drainage CO₂eq	128.44	10.95	128.36
HWP CO₂ eq	-420.85	207.50	-622.93
Total	-792.62	266.99	-413.00
FMRL 2013-2020		-142.07	
Indicative accountable amount for 2013 under FM (kt CO ₂ eq)			108.69

Reasons for the differences between the FMRLcorr and reported FM emission/removals are due to:

- Higher losses of biomass stock for the reported FM areas for 2013 due to a higher harvest than what was used in the projected FMRLcorr, based on policies implemented up to the end of 2009.
- The higher HWP removal for the reported FM for 2013, compared to the FMRLcorr, is due to the higher reported harvest of 2.827 Mm³, compared to a projected 2.39 Mm³ in the FMRL submission.

11.6 Harvested wood products

Ireland reports and accounts for domestically produced harvested wood products (HWP) using the first order decay approach (tier 2). The same approach is used for reporting under the conventions (see section, see 6.3.7.1 in Ch 6). However, these estimates exclude harvest emissions accounted AR activities in the 1st commitment period as set out under see para 2g(iv) of annex II to decision 2.CMP.8 (see Table 11.15). The HWP estimated are not adjusted for FM because this was not elected for the 1st commitment period (see Table 6.5.23 in Ch 6). All harvests from deforestation are accounted for on the basis of instantaneous oxidation.

Table 11.15 Adjusted HWP stock and stock changes for AR activities since 2007

Year	inflow tC			stock tC			Δ stock tC			Δ stock kt CO ₂	Solid wood	kt C			Yrs
	SW	WBP	PPB	SW	WBP	PPB	SW	WBP	PPB	Total	Initial stock	Cgain	C loss	Net	Half- life
2007	6269.3	6502.7	NO	6207.6	6413.4	NO	6207.6	6413.4	0.0	-46.3	12.8	12.8	-0.2	12.6	29.9
2008	IE	IE	NO	6085.9	6238.0	NO	-121.7	-175.4	0.0	1.1	12.1	0.0	-0.3	-0.3	IO
2009	IE	IE	NO	5966.6	6067.5	NO	-119.3	-170.6	0.0	1.1	11.8	0.0	-0.3	-0.3	IO
2010	IE	IE	NO	5849.6	5901.5	NO	-117.0	-165.9	0.0	1.0	11.5	0.0	-0.3	-0.3	IO
2011	IE	IE	NO	5734.9	5740.2	NO	-114.7	-161.4	0.0	1.0	11.3	0.0	-0.3	-0.3	IO
2012	IE	IE	NO	5622.4	5583.2	NO	-112.5	-157.0	0.0	1.0	11.0	0.0	-0.3	-0.3	IO
2013	10544.2	10753.8	NO	15952.7	16036.6	NO	10330.3	10453.4	0.0	-76.2	31.4	21.3	-0.5	20.8	30.0

IE indicated as emissions in 1st commitment period

IO harvests in the 1st commitment period are accounted on the basis of instantaneous oxidation

11.6.1 Information on activity data for the harvested wood products categories used for estimating the harvested wood products pool removed from domestic forests, for domestic consumption and for export, (para 2g(i and vii) of annex II of 2/CMP.8

Ireland derives HWP feedstock from domestically produced products (excluding imported HWP), such as sawnwood (SW), wood based panel (WBP), paper and paper board (PPB) from FAO/EUROSTAT data using Eq 2.8.1 and 2.8.2 of the IPCC Supplementary guidance 2013. This uses the data shown in CRF Gs4 and f_{irw} and f_{pulp} ratios to derive the volume of SW, WBP and PPB (see Table 6.3.21 in section 6.3.71 of Ch 6).

11.6.2 Information on half-lives used in estimating the emissions and removals for these categories in accordance with decision 2/CMP.7, annex, paragraph 29 or 30

The half-lives uses for HWP are the default values as indicated in section 6.3.7.1 f of Ch 6.

11.6.3 Information on whether emissions from harvested wood products originating from forests prior to the start of the second commitment period have been included in the accounting (see para 2g(iii) of annex II to decision 2.CMP.8).

Ireland uses HWP C inflows dating back to 1900 based on methods outlined in section 6.3.7.1 of Ch 6.

11.6.4 Information on how emissions from the harvested wood products pool that have been accounted for during the first commitment period on the basis of instantaneous oxidation have been excluded from the accounting for the second commitment period; (see para 2g(iv) of annex II to decision 2.CMP.8).

Harvests accounted on the basis of instantaneous oxidation under AR activities in the first commitment period represented 1 to 4 per cent of the total harvest over the period 2008-2012 (see Table 11.16). The emissions are excluded from accounting for the second commitment period by applying a corrected AR fraction (f_{ARcorr}) of zero to calculate HWP feedstocks under AR activities for the period 2007 to 2013. This means that only harvest for 2007 and from 2013 onwards are used as HWP feedstock under AR activities.

11.6.5 Information showing that harvested wood products resulting from deforestation have been accounted on the basis of instantaneous oxidation; (see para 2g(v) of annex II to decision 2.CMP.8).

The estimation of the annual fraction of harvest originating from the different forest activities (i.e. forest remaining forest (fFM), land converted to forest (fAR) and deforested (fD) harvest) are derived using Eq. 2.8.3 in Ch 2 of the IPCC supplementary guidance (see Table 11.16). The input information for the different activities (j) are derived from harvest data shown in Tables 6.3.4, 6.3.12 and 6.3.17 in Ch 6. All harvests for deforested land are assumed to be immediately oxidised by applying a fD corrected ratio of zero (fDcorr, Table 11.16), so CSC in the CRF under HWP are reported as IO under in CRF table 4(KP-1)C.

Table 11.16 The estimated fractions of HWP feedstock originating from different forest activities under article 3.3 and 3.4

The fDcorr fraction are the adjusted fraction to zero to account for harvest as instantaneous oxidation (2g(v) of annex II to 2/CMP.8)

	fAR	fARcorr	fD	fDcorr	fFM
1990	0.00	0.00	0.00	0.00	1.00
1991	0.00	0.00	0.00	0.00	1.00
1992	0.00	0.00	0.00	0.00	1.00
1993	0.00	0.00	0.00	0.00	1.00
1994	0.00	0.00	0.00	0.00	1.00
1995	0.00	0.00	0.03	0.00	0.97
1996	0.00	0.00	0.03	0.00	0.97
1997	0.00	0.00	0.03	0.00	0.97
1998	0.00	0.00	0.02	0.00	0.98
1999	0.00	0.00	0.02	0.00	0.98
2000	0.00	0.00	0.06	0.00	0.94
2001	0.00	0.00	0.06	0.00	0.94
2002	0.00	0.00	0.06	0.00	0.94
2003	0.00	0.00	0.06	0.00	0.94
2004	0.00	0.00	0.06	0.00	0.94
2005	0.00	0.00	0.06	0.00	0.94
2006	0.00	0.00	0.32	0.00	0.68
2007	0.03	0.03	0.14	0.00	0.83
2008	0.01	0.00	0.10	0.00	0.89
2009	0.09	0.00	0.09	0.00	0.83
2010	0.05	0.00	0.04	0.00	0.91
2011	0.06	0.00	0.04	0.00	0.90
2012	0.04	0.00	0.04	0.00	0.93
2013	0.06	0.06	0.00	0.00	0.93

11.6.6 Information showing that carbon dioxide emissions from harvested wood products in solid waste disposal sites, where these emissions are separately accounted for, and from wood harvested for energy purposes have been accounted on the basis of instantaneous oxidation;

Ireland does not report emissions of CO₂ from biogenic sources from landfills in the waste sector. The emissions associated with instantaneous oxidation of wood used for energy have been excluded from the HWP estimates. This is done using the term f_{IRW} in Eq 2.8.1 and 2.8.2 of the IPCC Supplementary guidance 2013 (see table 6.3.21 and section 6.3.7.1 c in Ch 6). The f_{IRW} estimate is based on industrial roundwood inflows from domestic harvest, which exclude wood used for energy purposes. The ratio of the industrial roundwood to total roundwood volumes, published in FAO/EUROSTAT, decreases from 0.99 in 2005 to 0.84 by 2013. This decrease is consistent with an increase utilisation of timber for energy purposes in Ireland.

11.7 Uncertainty Analysis

The IPCC tier 1 approach is applied to estimate uncertainties for the Article 3.3 activities described in this chapter using the methods for combining uncertainties given in section 6.3 of the IPCC good practice guidance for LULUCF (see equations 6.3.34 and 6.3.35 in section 6.3.4.7 in Chapter 6).

For detailed characterisation of individual uncertainties refer to sections 6.3.4.9 (FM), 6.3.5.9 (AR), 6.3.6.1.3 (D) and 6.3.7.1 (HWP) in CH 6. Note that the uncertainty estimates for HWP in AR lands is different due to the exclusion of feedstock for 2008-2012 (see section 11.7.4).

Table 11.17 Uncertainty estimates of forest activity estimates provides in CRF tables

Activity	CRF table	Combined uncertainty in year (±%)					
		2008	2009	2010	2011	2012	2013
AR	4(KP-I)A.1	21.69	20.86	20.36	20.27	20.35	20.28
	4(KP-I)C	24.24	24.63	24.61	24.60	24.58	24.57
	4(KP-II)2	93.41	93.75	94.00	94.12	94.16	94.19
	4(KP-II)4	52.20	52.20	52.20	52.20	52.20	52.20
	Total	22.76	21.80	21.60	21.23	21.21	20.81
D	4(KP-I)A.2	51.41	55.33	55.48	45.70	51.51	120.10
	4(KP-II)2	148.45	151.56	151.56	164.05	154.56	154.43
	4(KP-II)3	66.00	66.00	66.00	66.00	66.00	66.00
	Total	58.51	62.65	83.98	68.20	78.04	140.13
FM	4(KP-I)A.1	NA	NA	NA	NA	NA	5687.51
	4(KP-I)C	NA	NA	NA	NA	NA	24.28
	4(KP-II)2	NA	NA	NA	NA	NA	78.52
	4(KP-II)4	NA	NA	NA	NA	NA	59.36
	Total	NA	NA	NA	NA	NA	132.85
CM	4(KP-I) B.2	NA	NA	NA	NA	NA	70%
	4(KP-II)2	NA	NA	NA	NA	NA	NA
	4(KP-II)4	NA	NA	NA	NA	NA	105%
	Total	NA	NA	NA	NA	NA	70%
GM	4(KP-I) B.3	NA	NA	NA	NA	NA	33%
	4(KP-II)2	NA	NA	NA	NA	NA	133%
	4(KP-II)4	NA	NA	NA	NA	NA	105%
	Total	NA	NA	NA	NA	NA	75%

11.8 Quality Assurance and Quality Control (QA\QC)

The same QA/QC procedures were carried out for KP LULUCF as reported for forest lands under section 6.10.1. The entire compilation for this submission for both LULUCF (Chapter 6) and forest activities under Article 3.3 and 3.4 of the Kyoto Protocol (Chapter 11) were reviewed externally by an independent consultant, qualified as a UNFCCC expert reviewer for LULUCF/KP-LULUCF in March 2012. This provides an important element of quality assurance for this 2015 submission. Following the findings of this independent peer review, both chapter 6 and 11 of this report have been substantially improved to provide additional transparency and consistency between Convention and KP reporting for LULUCF.

The QA/QC could not be completed in time for submission due to problems with the release of a bug free CRF reporter, particularly for the KP LULUCF sector.

11.9 Recalculations in KP LULUCF

Numerous changes have been implemented to the KP LULUCF for the 2nd commitment period due to new methodological guidelines i.e. 2006 IPCC Guidelines, the 2013 Revised Supplementary Methods and IPCC 2013 GPG KP-LULUCF. All of these recalculations are outlined in section 6.11 of Ch 6.

The 2015 submission is the first time Ireland is reporting of CM and GM. No recalculations with respect to KP LULUCF.

Chapter 12 Information on Accounting of Kyoto Units

12.1 Background Information

Ireland's Standard Electronic Format report for the 2014 reported year, which contains the information required in paragraph 11 of the annex to decision 15/CMP.1 and adhering to the guidelines of the SEF has been submitted to the UNFCCC Secretariat electronically –

RREG1_IE_2014.xlsx

The contents of the SEF report (R1) can also be found in Appendix 1 – *SIAR Supplementary Information* of this document.

The contents of the report can also be found in annex 6 of this document.

12.2 Summary of Information Reported in the SEF Tables

There was 292,244,072 AAUs in Ireland's domain of the Union Registry at the end of the year 2014, of which 223,713,226 units were in the Party holding accounts; 1,810,375 units in the entity holding accounts; 735 units in the other cancellation accounts and 66,719,736 units in the retirement account.

There was 9,002,110 CERs in the registry at the end of 2014: 903,808 CERS were in the Party holding accounts; 5,315,249 CERs were held in the entity holding accounts; 2059 in the other cancellation accounts and 2,780,994 CERs were held in the retirement account.

There was 2,949,515 ERUs in the registry at the end of 2014; 1,635,668 ERUs were in the Party holding accounts; 74,964 ERUs were held on the entity holding accounts and 1,238,883 ERUs were held in the retirement account.

There was 1,193,621 t-CERs in the registry at the end of 2014; all of which were held on the entity holding accounts. The registry did not contain any RMUs or I-CERs.

There were no units in the Article 6 issuance and conversion accounts; no units in the Article 3.3 and Article 3.4 issuance or cancellation accounts and no units in the Article 12 afforestation and reforestation accounts.

The total amount of the units in the registry corresponded to 305,389,318 tonnes CO₂ eq.

Ireland's assigned amount is 314,184,272 tonnes CO₂eq.

Table 12.1 Information on the SEF tables

Annual Submission Item	Reported for 2014
15/CMP.1 annex I.E paragraph 11: Standard electronic format (SEF)	<p>Ireland's Standard Electronic Format report for the 2014 reported year, which contains the information required in paragraph 11 of the annex to decision 15/CMP.1 and adhering to the guidelines of the SEF has been submitted to the UNFCCC Secretariat electronically –</p> <p>RREG1_IE_2014.xlsx</p> <p>The contents of the SEF report (R1) can also be found in Appendix 1 – <i>SIAR Supplementary Information</i> of this document.</p>

12.3 Discrepancies and notifications

There were no discrepant transactions, no CDM notifications, no non-replacements and no invalid units in 2014. Accordingly, no actions were taken or changes made to address discrepancies for the 2014 reporting period (Table 12.2).

Table 12.2 Discrepancies and notifications

Annual Submission Item	Reported for 2014
15/CMP.1 annex I.E paragraph 12: List of discrepant transactions	<p>No discrepant transactions, pursuant of 15/CMP.1 annex I.E paragraph 12, occurred in the 2014 reporting period.</p> <p>The contents of the report R2 can also be found in the Appendix 1 – <i>SIAR Supplementary Information</i> of this document.</p> <p>Refer to Separate Electronic Attachment "SIAR Reports 2015-IE v 1.0.xls" Worksheet R2.</p>
15/CMP.1 annex I.E paragraph 13 & 14: List of CDM notifications	<p>No CDM notifications were received during the 2014 reporting period, pursuant of 15/CMP.1 annex I.E paragraphs 13 & 14.</p> <p>The contents of the Report R3 can also be found in Appendix 1 – <i>SIAR Supplementary Information</i> of this document.</p> <p>Refer to Separate Electronic Attachment "SIAR Reports 2015-IE v 1.0.xls" Worksheet R3.</p>

Annual Submission Item	Reported for 2014
15/CMP.1 annex I.E paragraph 15: List of non-replacements	No non-replacements occurred during the 2014 reporting period, pursuant of 15/CMP.1 annex I.E paragraph 15. The contents of the Report R4 can be found in Appendix 1 – <i>SIAR Supplementary Information</i> of this document. Refer to Separate Electronic Attachment “SIAR Reports 2015-IE v 1.0.xls” Worksheet R4.
15/CMP.1 annex I.E paragraph 16: List of invalid units	No invalid units exist as at 31 December 2014, pursuant of 15/CMP.1 annex I.E paragraph 16. The contents of the Report R5 can also be found in Appendix 1 – <i>SIAR Supplementary Information</i> of this document. Refer to Separate Electronic Attachment “SIAR Reports 2015-IE v 1.0.xls” Worksheet R5.
15/CMP.1 annex I.E paragraph 17 Actions and changes to address discrepancies	No actions were taken or changes made to address discrepancies for the 2014 reporting period.

12.4 Publicly Accessible Information

The public has access via the registry website to information on registry account types and account holders, information regarding Article 6 projects (currently no Article 6 projects in Ireland), information on transactions and the list of account holders authorised to hold Kyoto units in their account (Table 12.3).

Table 12.3 Publicly Accessible Information

Annual Submission Item	Reported for 2014
15/CMP.1 annex I.E Publicly accessible information	There was no change regarding publicly accessible information during 2014. The following information is publicly accessible and is available via the homepage of Ireland’s domain on the Union Registry – https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml

Annual Submission Item	Reported for 2014
	<p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p> <p>In accordance with the requirements of Annex E to Decision 13/CMP.1, all required information for a Party with an active Kyoto registry is provided with the exceptions as outlined below.</p>
	<p>Account Information (Paragraph 45)</p> <p>In line with the data protection requirements of Regulation (EC) No 45/2001 and Directive 95/46/EC and in accordance with Article 110 and Annex XIV of Commission Regulation (EU) No 389/2013, the information on account representatives, account holdings, account numbers, all transactions made and carbon unit identifiers, held in the EUTL, the Union Registry and any other KP registry (required by paragraph 45) is considered confidential.</p> <p>The most up-to-date account information may be accessed via the homepage of Ireland's domain on the Union Registry https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p>
	<p>JI projects in Ireland (Paragraph 46)</p> <p>Note that no Article 6 (Joint Implementation) projects are reported as conversion to an ERU under an Article 6 project, as this did not occur in the specified period.</p>
	<p>Holding and transaction information of units (Paragraph 47)</p> <p>Holding and transaction information is provided on a holding type level, due to more detailed information being declared confidential.</p> <p>The detailed information on transactions is considered confidential according to Article 110 of Commission Regulation (EU) No 389/2013:</p> <p><i>Information, including the holdings of all accounts, all</i></p>

Annual Submission Item	Reported for 2014
	<p><i>transactions made, the unique unit identification code of the allowances and the unique numeric value of the unit serial number of the Kyoto units held or affected by a transaction, held in the EUTL, the Union Registry and any other KP registry shall be considered confidential except as otherwise required by Union law, or by provisions of national law that pursue a legitimate objective compatible with this Regulation and are proportionate.</i></p> <p>The most up-to-date account information may be accessed via the homepage of Ireland's domain on the Union Registry https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p>
<p>15/CMP.1 annex I.E</p> <p>Publicly accessible information</p>	<p><u>Paragraph 47c</u></p> <p>Ireland does not host JI projects</p> <p><u>Paragraph 47e</u></p> <p>Ireland does not perform LULUCF activities and therefore does not issue RMUs</p> <p><u>Paragraph 47g</u></p> <p>No ERUs, CERs, AAUs and RMUs have been cancelled on the basis of activities under Article 3, paragraphs 3 and 4 to date.</p> <p><u>Paragraph 47h</u></p> <p>No ERUs, CERs, AAUs and RMUs have been cancelled following determination by the Compliance Committee that the Party is not in compliance with its commitment under Article 3, paragraph 1 to date.</p> <p><u>Paragraph 47j</u></p> <p>No ERUs, CERs, AAUs and RMUs have been retired to date</p> <p><u>Paragraph 47k</u></p> <p>There is no previous commitment period to carry ERUs, CERs, and AAUs over from.</p>
	<p>Entities authorised to hold Kyoto Units (Paragraph 48)</p>

Annual Submission Item	Reported for 2014
	<p>In line with the data protection requirements of Regulation (EC) No 45/2001 and Directive 95/46/EC and in accordance with Article 110 and Annex III of the Commission Regulation (EU) no 389/2013, the legal entity contact information (required by paragraph 48) is considered confidential.</p> <p>The most up-to-date account information may be accessed via the homepage of Ireland's domain on the Union Registry https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p>

12.5 Calculation of the Commitment Period Reserve

The commitment period reserve (CPR) is the lower of the two values given by 90 per cent of the assigned amount and five times the estimate of total emissions in the most recently reviewed inventory. The inventory for 2012 submitted on 15 May 2014 is the most recently reviewed inventory for Ireland ([FCCC/ARR/2014/IRL](#)). The total emissions in 2012 amounted to 58,531,238 tonnes *CO₂ equivalent* and five times this estimate is 292,656,190 tonnes *CO₂ equivalent*. This value is greater than 90 per cent of the assigned amount (282,765,845 tonnes *CO₂ equivalent*) determined in the review of Ireland's initial report ([FCCC/IRR/2007/IRL](#)) and therefore the commitment period reserve is 282,765,845 tonnes *CO₂ equivalent*.

12.6 Accounting for Activities under Article 3.3

In the initial report under the Kyoto Protocol ([FCCC/IRR/2007/IRL](#)), Ireland elected to account for all activities under Article 3, paragraph 3, of the Kyoto Protocol on the basis of commitment period accounting. No information on the accounting of KP-LULUCF is therefore included in the SEF tables.

Chapter 13 Changes in National System

13.1 Changes in National System since previous submission

Ireland's national system is described in section 1.2 of Chapter 1. There were no changes in the institutions or resources involved in the national system during the current reporting cycle.

Chapter 14 Changes in National Registry

14.1 Introduction

The national registry of Ireland is described in the initial report under the Kyoto Protocol ([FCCC/IRR/2007/IRL](#)). Ireland's national registry was established initially for the implementation of Directive 2003/87/EC (EP and CEU, 2003) on emissions trading. The registry software was purchased from the Department of the Environment, Food and Rural Affairs in the UK and has been developed in consultation with other Member States that also purchased this software as part of the GRETA group. 2012 saw the transition from the national registry using the GRETA registry software to the Consolidated System of EU Registries (CSEUR).

The following changes to the national registry of IRELAND have occurred in 2014.

These changes are summarised in this chapter and further details are provided in electronic form as Appendix 1 *SIAR Supplementary Information* to the NIR.

14.2 Information on Changes in National Registry

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(a) Change of name or contact	Addition of contact details: Mr Damien FAHEY Environmental Protection Agency PO Box 3000, Johnstown Castle Estate, Co Wexford, IRELAND. Email: d.fahey@epa.ie / etradmin@epa.ie Telephone: +353 (0)53 91 70750 Fax: +353 (0)53 91 60699
15/CMP.1 annex II.E paragraph 32.(b) Change regarding cooperation arrangement	No change of cooperation arrangement occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(c) Change to database structure or the capacity of national registry	An updated diagram of the database structure is attached as Annex A in electronic form as Appendix 1 <i>SIAR Supplementary Information</i> to the NIR. Versions of the CSEUR released after 6.1.7.1 (the production version at the time of the last Chapter 14 submission) introduced changes in the structure of the database. These changes were limited and only affected EU ETS

Reporting Item	Description
	<p>functionality.</p> <p>No change was required to the database and application backup plan or to the disaster recovery plan.</p> <p>No change to the capacity of the national registry occurred during the reported period.</p>
<p>15/CMP.1 annex II.E paragraph 32.(d)</p> <p>Change regarding conformance to technical standards</p>	<p>Changes introduced since Version 6.1.7.1 of the national registry were limited and only affected EU ETS functionality.</p> <p>However, each release of the registry is subject to both regression testing and tests related to new functionality. These tests also include thorough testing against the DES and were successfully carried out prior to the relevant major release of the version to Production (see Annex B) in electronic form as Appendix 1 <i>SIAR Supplementary Information to the NIR</i>. Annex H testing was carried out in February 2015 and the successful test report has been attached in electronic form as Appendix 1 <i>SIAR Supplementary Information to the NIR</i>.</p> <p>No other change in the registry's conformance to the technical standards occurred for the reported period.</p>
<p>15/CMP.1 annex II.E paragraph 32.(e)</p> <p>Change to discrepancies procedures</p>	<p>No change of discrepancies procedures occurred during the reporting period.</p>
<p>15/CMP.1 annex II.E paragraph 32.(f)</p> <p>Change regarding security</p>	<p>No change of security measures occurred during the reporting period</p>
<p>15/CMP.1 annex II.E paragraph 32.(g)</p> <p>Change to list of publicly available information</p>	<p>The following information is publicly accessible and is available via the homepage of Ireland's domain on the Union Registry – https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p> <p>In accordance with the requirements of Annex E to Decision 13/CMP.1, all required information for a Party with an active</p>

Reporting Item	Description
	Kyoto registry is provided with the exceptions as outlined below.
	<p>Account Information (Paragraph 45)</p> <p>In line with the data protection requirements of Regulation (EC) No 45/2001 and Directive 95/46/EC and in accordance with Article 110 and Annex XIV of Commission Regulation (EU) No 389/2013, the information on account representatives, account holdings, account numbers, all transactions made and carbon unit identifiers, held in the EUTL, the Union Registry and any other KP registry (required by paragraph 45) is considered confidential.</p> <p>The most up-to-date account information may be accessed via the homepage of Ireland's domain on the Union Registry – https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p>
	<p>JI projects in Ireland (Paragraph 46)</p> <p>Note that no Article 6 (Joint Implementation) projects are reported as conversion to an ERU under an Article 6 project, as this did not occur in the specified period.</p>
	<p>Holding and transaction information of units (Paragraph 47)</p> <p>Holding and transaction information is provided on a holding type level, due to more detailed information being declared confidential.</p> <p>The detailed information on transactions is considered confidential according to Article 110 of Commission Regulation (EU) No 389/2013:</p> <p><i>Information, including the holdings of all accounts, all transactions made, the unique unit identification code of the allowances and the unique numeric value of the unit serial number of the Kyoto units held or affected by a transaction, held in the EUTL, the Union Registry and any other KP registry shall be considered confidential except as otherwise required by Union law, or by provisions of national law that pursue a legitimate objective compatible with this Regulation and are</i></p>

Reporting Item	Description
	<p><i>proportionate.</i></p> <p>The most up-to-date information may be accessed via the homepage of Ireland's domain on the Union Registry – https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website: http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports</p>
	<p><u>Paragraph 47c</u></p> <p>Ireland does not host JI projects.</p> <p><u>Paragraph 47e</u></p> <p>Ireland does not perform LULUCF activities and therefore does not issue RMUs</p> <p><u>Paragraph 47g</u></p> <p>No ERUs, CERs, AAUs and RMUs have been cancelled on the basis of activities under Article 3, paragraphs 3 and 4 to date.</p> <p><u>Paragraph 47h</u></p> <p>No ERUs, CERs, AAUs and RMUs have been cancelled following determination by the Compliance Committee that the Party is not in compliance with its commitment under Article 3, paragraph 1 to date.</p> <p><u>Paragraph 47j</u></p> <p>No ERUs, CERs, AAUs and RMUs have been retired to date</p> <p><u>Paragraph 47k</u></p> <p>There is no previous commitment period to carry ERUs, CERs, and AAUs over from.</p>
	<p>Entities authorised to hold Kyoto Units (Paragraph 48)</p> <p>In line with the data protection requirements of Regulation (EC) No 45/2001 and Directive 95/46/EC and in accordance with Article 110 and Annex III of the Commission Regulation (EU) no 389/2013, the legal entity contact information (required by paragraph 48) is considered confidential.</p> <p>The most up-to-date account information may be accessed via the homepage of Ireland's domain on the Union Registry – https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml</p> <p>See also the EPA website:</p>

Reporting Item	Description
	http://www.epa.ie/climate/emissionstradingoverview/union%20registry/publicreports
15/CMP.1 annex II.E paragraph 32.(h) Change of Internet address	No change of the registry internet address occurred during the reporting period. Ireland's domain of the Union Registry can be found at this link: https://ets-registry.webgate.ec.europa.eu/euregistry/IE/public/reports/publicReports.xhtml
15/CMP.1 annex II.E paragraph 32.(i) Change regarding data integrity measures	No change of data integrity measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(j) Change regarding test results	Changes introduced since version 6.1.7.1 of the national registry were limited and only affected EU ETS functionality. Both regression testing and tests on the new functionality were successfully carried out prior to release of the version to Production. The site acceptance test was carried out by quality assurance consultants on behalf of and assisted by the European Commission; the report is attached as Annex B . Annex H testing was carried out in February 2015 and the successful test report has been attached in electronic form as Appendix 1 <i>SIAR Supplementary Information</i> to the NIR.
The previous Annual Review Recommendations	No recommendations relevant to registry operations in FCCC/ARR/2014/IRL of 08 April 2015.

Chapter 15 Minimisation of Adverse Impacts under Article 3, paragraph 14

15.1 Introduction

Article 3, paragraph 14, of the Kyoto Protocol requires that Annex I Parties shall strive to meet their commitments under Article 3, paragraph 1 of the Kyoto Protocol in such a way as to minimize adverse social environmental and economic impacts on developing country Parties, particularly those Parties identified in Article 4, paragraphs 8 and 9, of the Convention. Information on how commitments under Article 3, paragraph 14, are being implemented is to be prioritised under a number of actions as set down in section H of the annex to guidelines for the preparation of supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol (Decision 15/CMP.1). These requirements are addressed in this chapter. There has been no change to the information provided since the previous inventory submission.

15.2 Context

As a Member State of the European Union, Ireland's commitments under the Kyoto Protocol are being implemented under Decision 2005/166/EC, governing joint fulfilment under Article 4, and Decision 280/2004/EC, which covers specific emissions monitoring and reporting requirements. In this context, the minimization of adverse impacts on developing countries is also largely dictated by the European Commission's policy on climate change and by its policies and programmes affecting developing countries. Regulation at the European level also controls or influences market conditions, fiscal incentives, tax and duty exemptions and subsidies in all economic sectors in Member States.

The impact assessment of new policy initiatives has been established in the European Union, which allows their potential adverse social, environmental and economic impacts on various stakeholders, including developing country Parties, to be identified and limited at an early stage within the legislative process. Impact Assessment Guidelines specifically address impacts on third countries and also issues related to international relations. This provides a framework in which Member States like Ireland can also ensure a high level of protection of the environment and contribute to the integration of environmental considerations into the preparation and adoption of specified plans and programmes with a view to promoting sustainable development.

15.3 Specific Elements

a) The progressive reduction or phasing out of market imperfections, fiscal incentives, tax and duty exemptions and subsidies in all greenhouse-gas-emitting sectors, taking into account the need for energy price reforms to reflect market prices and externalities

Ireland's electricity market has been deregulated and the levy supporting the use of peat for electricity generation under a Public Service Agreement has been discontinued. Tax incentives contributed to the development of Ireland's most recent gas field off the west coast but such incentives will be severely curtailed for any similar developments in the future under new legislation.

Reforms of the Common Agricultural Policy have resulted in changes to subsidies in agriculture, which are now linked to environmental, food safety and animal welfare standards. The EU Emissions Trading Scheme is a market-based emissions control measure which applies to major combustion and process emission sources of CO₂ and a carbon tax is being introduced for fossil fuel use outside the ETS.

b) Removing subsidies associated with the use of environmentally unsound and unsafe technologies

Environmentally unsound and unsafe technologies may be regarded as technologies that would not conform to the concept of sustainable development and the objective and principles of the UNFCCC. The EC has addressed this issue by developing legislation to ensure that the price for coal produced in Member States is not lower than the price of coal of similar quality available from third countries and by phasing out subsidies on fossil fuel production and consumption by 2010. No environmentally unsound or unsafe technologies are in operation in Ireland.

c) Cooperating in the technological development of non-energy uses of fossil fuels, and supporting developing country Parties to this end

The Irish Government is represented on the energy and environment strands of the Seventh Framework Programme (FP7) for Research and Technological Development (RTD). This representation includes the FP7 Energy Programme Committees that focuses on developing and agreeing the annual work programme and strategic vision for the FP7 Energy Work programme 2007–2013. Much of the focus of this (energy theme) initiative is on energy mitigation through supporting technological development and transfer through joint collaborations and calls with emerging economies including India, Russia and Brazil.

The International Energy Agency (IEA) is the energy forum and think-tank for 26 OECD countries. The Irish Government is a Party to four Renewable Energy Implementing Agreements of the IEA on Bioenergy, Ocean, Wind and RE Technology Deployment (RETD). Ireland provides national delegates to the executive committees of the Implementing Agreements and nominates and supports country experts to a number of tasks. The Government also sits on the Committee for Energy research and technology (CERT). Ireland is a member of the EU Expert Group on Technology, which supports the EC in climate negotiations. This expert group is focused on the transfer of technology to reduce the impacts of climate change and on supporting developing countries to this end.

d) Cooperating in the development, diffusion, and transfer of less-greenhouse-gas-emitting advanced fossil-fuel technologies, and/or technologies, relating to fossil fuels, that capture and store greenhouse gases, and encouraging their wider use; and facilitating the participation of the least developed countries and other non-Annex I Parties in this effort

The EU collaborates with other Annex I and Non-Annex I Parties (Brazil, Saudi Arabia, China, Colombia, India, Korea, Mexico and South Africa) in the Carbon Sequestration Leadership Forum (CSLF). The CSLF is a ministerial-level international climate change initiative that is focused on the development of improved cost-effective technologies for the capture transport and long-term safe storage of CO₂. The mission of the CSLF is to facilitate the development and deployment of such technologies via collaborative efforts that address key technical, economic, and environmental obstacles. The CSLF will also promote awareness and champion legal, regulatory, financial, and institutional environments conducive to such technologies.

Ireland began its support to the Renewable Energy and Energy Efficiency Partnership (REEEP) in 2005. Following the decision by the Irish Government in 2007 to offset all its carbon emissions from official travel, REEEP was chosen as its implementing partner. REEEP is a Public-Private partnership and was launched by the United Kingdom along with other partners at the Johannesburg World Summit on Sustainable Development in August 2002. By providing opportunities for concerted collaboration among its partners, REEEP aims to accelerate the marketplace for renewable energy and energy efficiency. Funding from Ireland is being prioritised for projects in its programme countries of Ethiopia, Lesotho, Mozambique, Tanzania, Uganda, Zambia and Malawi.

Ireland provides development assistance in line with the priorities expressed by partner countries. To date requests for assistance in the area of technology are primarily in connection with water supply, transport infrastructure and agriculture. An innovative programme in Ethiopia carries out operational participatory research with farmers, extension workers and government officials to identify, develop, and disseminate new agricultural technologies. Some of the successful technologies are based on traditional practices, for example soil conservation techniques. Other new technologies are related to new crop varieties and irrigation. In addition to ODA, private companies also provide technology and advice to developing countries, particularly in the energy sector. Due to the range of funding sources no precise figure is available for funding attributed to technology development and transfer. Ireland's support to REEEP is worth mentioning again here as an example of Ireland's support for technology transfer. REEEP brings the private and public sectors together to facilitate the financing, development and transfer of renewable energy technologies. Ireland believes that this type of public-private collaboration is essential for the development of appropriate and environmentally sound technologies and to facilitate their application and use in developing countries.

e) Strengthening the capacity of developing country Parties identified in Article 4, paragraphs 8 and 9, of the Convention for improving efficiency in upstream and downstream activities relating to fossil fuels, taking into consideration the need to improve the environmental efficiency of these activities

The EU contributes to strengthening the capacities of countries engaged in the export of fossil fuels through the work of the Energy Expert Group of the Gulf Cooperation Council (GCC), in particular under the working sub-group on energy efficiency. As part of the EU's research programme, a project called "EUROGULF" was launched with the objective of to analyse The European Commission's planned e-network on clean energy technologies, is aiming to promote research and technical development of clean energy technologies in the GCC countries.

Ireland currently holds the Programme Chair of Renewable Energy and Energy Efficiency Partnership, a Type 2 International NGO. The Renewable Energy and Energy Efficiency Partnership (REEEP) is a global partnership that works to reduce the barriers in policy, regulatory and financial structures that bar and limit the uptake of renewable-energy and energy-efficiency technologies and projects. This Partnership focuses on deployment of projects in sub-Saharan Africa, Asia and Latin America. Ireland is actively involved in the partnership, alongside energy-related organisations from Australia, Austria, Canada, Germany, Italy, Spain, the Netherlands, New Zealand, Norway, the UK, the USA and the European Commission.

Ireland is a founding member of the UNEP SEFI Public Finance Alliance, or 'SEF Alliance'. This is a member-driven coalition of public and publicly backed organisations that finance sustainable-energy

markets in various countries, including emerging and developing economies. . Members use the platform to exchange best practices, pool resources, launch joint projects and assist other governments in establishing new or similar financing models. The SEF Alliance is under the remit of the Sustainable Energy Finance Initiative (SEFI) of the United Nations Environment Programme (UNEP) but is governed directly by its members and pursues activities according to their interests. In 2008, the Alliance published Public Finance for Climate Change Mitigation, which provided an overview of mechanisms being used by the public sector to help scale up the climate mitigation markets, with a particular focus on the clean energy sector. In 2008, the SEF Alliance also published a Public Venture Capital Study which examined current clean-energy venture financing, focusing on the role of public sector-sponsored venture capital.

f) Assisting developing country Parties which are highly dependent on the export and consumption of fossil fuels in diversifying their economies

Ireland supports a range of EU activities aimed at reducing dependence on the consumption of fossil fuels, in particular those EU support programmes for the promotion of renewable energies and energy efficiency in developing countries. Renewable energy cooperation with Mediterranean and Gulf countries which led to the Mediterranean Solar Plan, endorsed in 2008 with the objective of installing 20 GW of new generation capacity in solar and other renewable energy sources around the Mediterranean Sea by 2020. Another objective is to create a sub-regional electricity market between Morocco, Tunisia and Algeria and to progressively integrate it with the electricity market of the EU. Important initiatives which target energy efficiency and renewable energy projects in South America, Africa and Asia include the Africa, Caribbean and the Pacific (ACP-E) Energy Facility, the Latin America Investment Facility (LAIF), the Euro-Solar Programme in Latin America and the Global Energy Efficiency and Renewable Energy Fund (GEEREF).

Glossary

Annex 1 Parties	Countries listed in Annex I to the United Nations Framework Convention on Climate Change
Base year	The year or period under the Kyoto Protocol on which quantified emission limitation or reduction commitments in the commitment period are based.
BOD	Biochemical Oxygen Demand
CARBWARE	A forest model to calculate carbon stock change and growth increment for Irish forests
CFCs	Chlorofluorocarbons
CH₄	Methane
CHP	Combined Heat and Power.
CMMS	Cattle Movement and Monitoring System
CO	Carbon Monoxide
CO₂	Carbon Dioxide
CO₂ equivalent	The equivalent mass as CO ₂ of other greenhouse gases converted on the basis of their global warming potential (GWP)
COFORD	National Council for Forest Research and Development
Commitment Period	The years 2008 to 2012 (first CP) or 2013 to 2020 (second CP) inclusive for which quantified emission limitation or reduction commitments are established under the Kyoto Protocol
COP	Conference of the Parties
CORINAIR	Co-ordinated Information on the environment in the European Community-AIR. CORINAIR was one of several collaborative exercises initiated under the CORINE programme to harmonise the collection and dissemination of information on the environment in the EU.
CRF	Common Reporting Format
DAF	Department of Agriculture and Food
DAFM	Department of Agriculture, Food and the Marine
DCENR	Department of Communications, Energy and Natural Resources
DEHLG	Department of Environment Heritage and Local Government
DNDC	DeNitrification-DeComposition, is a computer simulation model of carbon and nitrogen biogeochemistry in agri-ecosystems
EMEP	European Monitoring and Evaluation Programme, a co-operative programme for monitoring and evaluation of the long-range transmissions of air pollutants in Europe
Emission	(of a greenhouse gas). The release of greenhouse gases into the atmosphere.
Enteric Fermentation	The digestive process in ruminant animals (e.g cattle and sheep) where bacteria convert the feed to a usable form of energy for the animal, producing CH ₄ as a by product
EUROSTAT	Statistical Agency of the European Union
FAO	Food and Agriculture Organisation of the United Nations
FFS	Farm Facilities Survey
FIPS	Forest Inventory and Planning System
Fluorinated Gases	HFCs, PFCs, SF ₆ and NF ₃

Fossil Fuel	Peat, coal, oil and natural gas and associated derivatives
FTA	Fraction of BOD in sludge that degrades anaerobically
GDP	Gross Domestic Product
Gg	Gigagram (10^9 g) = kilo tonne = 1,000 tonnes
Greenhouse Gas	A gas in the atmosphere that allows solar radiation through to the earth's surface, but traps some of the heat radiated back from the earth's surface
GWP	The cumulative warming over a specified time period, e.g. 100 years, resulting from a unit mass of a greenhouse gas emitted at the beginning of that time period, expressed relative to an absolute GWP of 1 for CO ₂
HCFCs	Hydrochlorofluorocarbon
HFCs	Hydrofluorocarbons
HGV	Heavy Goods Vehicle
IEA	International Energy Agency
IEF	Implied Emission Factor
IPC	Integrated Pollution Control
IPCC	Intergovernmental Panel on Climate Change
IUCC	Information Unit on Climate Change
kt	kilo tonne (1,000 tonnes)
Kyoto Protocol	The Protocol to the UNFCCC adopted by Decision 1/CP.3 under which industrialised countries agreed to reduce their combined greenhouse gas emissions in 1990 by at least 5 per cent by the period 2008-2012
LTO	Landing and Take-off cycle
MMS	Manure Management System
Montreal Protocol	Protocol on substances that deplete the ozone layer
Mt	million tonnes or mega tonnes
N₂O	Nitrous Oxide
NBP	Net Biome Productivity
NEE	Net Ecosystem Exchange
NF₃	Nitrogen trifluoride
NIR	National Inventory Report
NMVOC	Non Methane Volatile Organic Compounds
NO_x	Nitrogen Oxides
NRA	National Roads Authority
OSPAR	Oslo and Paris Convention for the Protection of the Marine Environment
PFCs	Perfluorocarbons
SBSTA	Subsidiary Body for Scientific and Technological Advice
SEAI	Sustainable Energy Authority of Ireland
SF₆	Sulphur Hexafluoride
Sink	The reservoir or pool in which sequestered carbon is stored; the process of sequestration
SO₂	Sulphur Dioxide
Teagasc	Irish Agriculture and Food Development Authority
TPER	Total Primary Energy Requirement
UNECE	United Nations Economic Commission for Europe
UNFCCC	United Nations Framework Convention on Climate Change
VOC	Volatile Organic Compounds

References

- Adams, M., Wagner, A., Goodwin, J., O'Leary, E., Creedon, M., McHugh, M and Gibson, C. 2005. Compiling Emission Inventories of HFCs, PFCs and SF6 for inclusion in Ireland's greenhouse gas submissions to the EU under Decision 280/2004/EC under the United National Framework Convention on Climate Change. Netcen/ ED48411/R1, AEA Technology UK.
- AEA. 2011. HFC consumption and emissions forecasting. Containing an update to the June 2008 HFC projections. Report to Defra.
- Alam S, Duffy P, Hyde B, McNabola A., Estimation and back extrapolation of CO2 emissions from the road transport sector: Emissions in Ireland, 1990 to 2013, 23rd International Conference on Modelling, Monitoring and Management of Air Pollution, Valencia, Spain, 1st-3rd June , 2015, pp1-13.
- B. Baesens, T. van Gestel, S. Viaene, M. Stepanova, J. Suykens, and J. Van-thienen. Benchmarking state-of-the-art classification algorithms for credit scoring. *Journal of the Operational Research Society*, 54:627-635, 2003.
- Black, K., Hendrick, E., Gallagher, G., Farrington, P., 2012. Establishment of Ireland's projected reference level for Forest Management for the period 2013-2020 under Article 3.4 of the Kyoto Protocol. *Irish Forestry* 69: 7-33.
- Black, K., Tobin, B., Saiz, G., Byrne, K. and Osbourne, B. 2004. Improved estimates of biomass expansion factors for Sitka spruce. *Irish Forestry* 61: 50-65
- Black, K.G. 2008. Scaling up from the stand to regional level: an analysis based on the major forest species in Ireland. Conference proceedings IIASA GHG Uncertainty workshop, Vienna, Sept 2008
- Black, K.G., Bolger, T., Davis, P., Nieuwenhuis, M., Reidy, B., Saiz, G., Tobin, B. and Osbourne, B. 2007. Inventory and Eddy Covariance Based Estimates of Annual Carbon Sequestration in a Sitka spruce (*Picea sitchensis* (Bong.) Carr.) Forest Ecosystem. *J Eur For Res* 126: 167-178
- Black, K.G., Byrne, K.A., Mencuccini, M., Tobin, B., Nieuwenhuis, M., Reidy, B., Bolger, T., Saiz, G., Green, C., Farrell, E.P. and Osborne, B. 2009b. Carbon stock and stock changes across a Sitka spruce chronosequence on surface water gley soils. *Forestry*, 85(3), 255-271
- Black, K.G., Farrell, E.P. (eds) 2006. Carbon Sequestration in Irish Forest Ecosystems. Council for Forest Research and Development (COFORD), Dublin, ISBN 1 902696 48 4
- Black, K.G., O'Brien, P., Redmond, J., Barrett, F. and Twomey M. 2009a. The extent of peatland afforestation in Ireland. *Irish Forestry* 65, 61-71.
- Bossard, M., Feranec, J. and Otahel, J. 2000. CORINE land cover technical guide – Addendum, 2000. Technical Report No. 40, EEA.
- Boyle, O. C., 1987. National Database on Waste. An Foras Forbartha, Dublin.
- Broad, L, Lynch, T. 2006. Panel data validation using cross-sectional methods. *Irish Forestry* 63: 80-95.

Büttner, G., Feranec, J., Jaffrain, G., Mari, L., Maucha, G. and Soukup, T. 2004. The CORINE land cover 2000 project. In Reuter, R. (Ed.), EARSel eProceedings 3(3). Paris (EARSel), pp. 331–346. CEC. 1993. CORINE Land Cover Technical Guide. European Union, Directorate-Environment, Luxembourg.

C. Bigler and H. Bugmann. Predicting the time of tree death using dendrochronological data. *Ecological Applications*, 14(3):902{914, June 2004.

Carey, P., Carty, G., Crowe, M., and Rudden, P.J. 1996. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

CEC (Commission of the European Communities), 2004. Commission Decision 2004/156/EC of 29/01/2004 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC. O.J. L59, 26 February 2004.

CEC (Council of the European Communities), 1991. Council Directive 91/676/EEC of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources O.J. L167, July 1993.

CEC (Council of the European Communities), 1999. Council Directive 99/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations. O.J. L85, 29 March 1999.

CEU (Council of the European Union), 1999. Council Directive 1999/31/EC on the landfill of waste. O.J. L182, July 1999.

Collins, C., Le Bolloch, O. and Meaney, B. 2005. National Waste Report 2004. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Collins, C., Meaney, B., Nolan, K., Maher, H. and Corish, C. 2004b. National Waste Database Interim Report 2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Collins, C., Meaney, B., Nolan, K., Maher, H. and Murphy, D. 2004a. National Waste Database Interim Report 2002. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Commission Decision 2013/162/EU on determining Member States' annual emission allocations for the period from 2013 to 2020 pursuant to Decision No 406/2009/EC of the European Parliament and of the Council

Commission Implementing Decision 2013/634/EU on the adjustments to Member States' annual emission allocations for the period from 2013 to 2020 pursuant to Decision No 406/2009/EC of the European Parliament and of the Council

Cossu, R., Andreottola, G. and Muntoni, A., 1996. Modelling Landfill Gas Production. In *Landfilling of Waste Biogas*. Edited by T. H. Christensen, R. Cossu and R. Stegman. E and FN Spon, London.

Council Decision EU 2015/1339 on the conclusion, on behalf of the European Union, of the Doha Amendment to the Kyoto Protocol to the United Nations Framework Convention on Climate Change and the joint fulfilment of commitments thereunder

Council Decision EU 2015/1340 on the conclusion, on behalf of the European Union, of the Agreement between the European Union and its Member States, of the one part, and Iceland, of the other part, concerning Iceland's participation in the joint fulfilment of commitments of the European

Union, its Member States and Iceland for the second commitment period of the Kyoto Protocol to the United Nations Framework Convention on Climate Change

Crowe, M., Fanning, A., Nolan, K., Carty, G. and Howley, D. 2000. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Crowley, A. M. 2001. Prediction of net energy requirements of beef cattle. M.Agr.Sc. Thesis, National University of Ireland, University College Dublin, November 2001, 145 pp.

Crowley, A. M., Keane, M.G., Agabriel, J. and O'Mara F. P. (2002). Prediction of net energy requirements of beef cattle. Proceedings of Agricultural Research Forum, 11 and 12 March 2002, Tullamore, p 19.

Cruickshank, M.M. and Tomlinson, R.W. 1996. Application of CORINE Land Cover Methodology to the U.K.-Some Issues Raised from Northern Ireland. Global Ecology and Biogeography Letters 5 (4/5) Remote Sensing and GIS in the Service of Ecology and Biogeography: A Series of Case Studies.

Cruickshank, M.M., Tomlinson, R.W. and Trew, S. 2000. Application of CORINE land-cover mapping to estimate the carbon stored in the vegetation of Ireland. Journal of Environmental Management, 58, 269-287.

CSO. Transport Omnibus 2009-2010 & 2011. 2010 & 2012

<http://www.cso.ie/en/releasesandpublications/transport/archive/>

CTC (Clean Technology Centre), 2005. NMVOC Inventory for Ireland (SNAP Sector 6 Solvent and Other Product Use). Report to the Environmental Protection Agency.

DAFF (Department of Agriculture, Fisheries and Food), 2004. CMMS Statistics Report 2003. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DAFF (Department of Agriculture, Fisheries and Food), 2005. CMMS Statistics Report 2004. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DAFF (Department of Agriculture, Fisheries and Food), 2006. CMMS Statistics Report 2005. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DAFF (Department of Agriculture, Fisheries and Food), 2007. CMMS Statistics Report 2006. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DAFF (Department of Agriculture, Fisheries and Food), 2008. CMMS Statistics Report 2007. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DAFF (Department of Agriculture, Fisheries and Food), 2009. AIM Bovine Statistics Report 2008. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie

DAFF (Department of Agriculture, Fisheries and Food), 2010. AIM Bovine Statistics Report 2009. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie

DAFM (Department of Agriculture, Food and the Marine), 2011. AIM Bovine Statistics Report 2010. Department of Agriculture, Food and the Marine, Dublin. www.agriculture.gov.ie

DAFM (Department of Agriculture, Food and the Marine), 2012. AIM Bovine Statistics Report 2011. Department of Agriculture, Food and the Marine, Dublin. www.agriculture.gov.ie

DAFM (Department of Agriculture, Food and the Marine), 2013. AIM Bovine Statistics Report 2012. Department of Agriculture, Food and the Marine, Dublin. www.agriculture.gov.ie

DAFM (Department of Agriculture, Food and the Marine), 2014. AIM Bovine Statistics Report 2013. Department of Agriculture, Food and the Marine, Dublin. www.agriculture.gov.ie

Decision No. 406/2009/EC of the European Parliament and of the Council on the effort of Member States to reduce their greenhouse gas emissions to meet the Community's greenhouse gas emission reduction commitments up to 2020

DEHLG (Department of the Environment Heritage and Local Government), 2005. European Communities(Good Agricultural Practice for Protection of Waters) Regulations 2005. SI No 788 of 2005. Government Publications Office, Dublin.

DEHLG (Department of the Environment Heritage and Local Government), 2006. European Communities(Good Agricultural Practice for Protection of Waters) Regulations 2006. SI No 378 of 2006. Government Publications Office, Dublin.

DEHLG (Department of the Environment Heritage and Local Government), 2007. Limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints, varnishes and vehicle refinishing products regulations 2007. S.I. No. 199 of 2007. Government Publications Office, Dublin.

DEHLG (Department of the Environment Heritage and Local Government), 2008. Waste Management (Certification of Historic Unlicensed Waste Disposal and Recovery Activity) Regulations 2008. S.I. No. 524 of 2008. Government Publications Office, Dublin.

DEHLG (Department of the Environment Heritage and Local Government), 2009. European Communities(Good Agricultural Practice for Protection of Waters) Regulations 2009. SI No 101 of 2009. Government Publications Office, Dublin.

DEHLG (Department of the Environment Heritage and Local Government), 2010. European Communities(Good Agricultural Practice for Protection of Waters) Regulations 2010. SI No 610 of 2010. Government Publications Office, Dublin.

Del Prado, A., Brown, L., Schulte, R., Ryan, M. and Scholefield. 2006. Principles of development of a mass balance N cycle model for temperate grasslands: an Irish case study Nutrient Cycling in Agroecosystems (2006) 74:115–131.

DELG (Department of the Environment and Local Government), 1997. Sustainable Development. A Strategy for Ireland. Department of the Environment and Local Government, Custom House, Dublin.

DELG (Department of the Environment and Local Government), 1998. A Policy Statement. Waste Management. Changing our Ways. Department of the Environment and Local Government, Custom House, Dublin.

DOE (Department of the Environment), 1992. The Environmental Protection Agency Act, 1992. SI No 7 of 1992. Department of the Environment, Custom House, Dublin.

DOE (Department of the Environment), 1994. Recycling for Ireland. A Strategy for Recycling Domestic and Commercial Waste. Department of the Environment, Custom House, Dublin.

DOE (Department of the Environment), 1997. IRELAND : Second National Communication under the UN Framework Convention on Climate Change. Department of the Environment, Custom House, Dublin.

DOE,DELG,DEHLG,DOT,DOTTS 1990 to 2013, Irish Bulletin of Vehicle and Driver Statistics (26 reports).

E. R. DeLong, D. M. DeLong, and D. L. Clarke-Pearson. Comparing the areas under two or more correlated receiver operating characteristic curves: A nonparametric approach. *Biometrics*, 44:837-845, 1988.

Edwards, P.N., Christie, J.M. 1981. Yield models for forest management. Forestry Commission Booklet No. 48. HMSO, London.

EP and CEU (European Parliament and Council of the European Union), 2001. Directive 2001/81/EC of 23 October 2001 on national emissions ceilings for certain atmospheric pollutants. O.J. L309, 27 November 2001.

EP and CEU (European Parliament and Council of the European Union), 2003. Directive 2003/87/EC establishing a scheme for greenhouse gas emission allowance trading within the Community and amending Council Directive 96/61/EC. O.J. L275, 25 October 2003.

EP and CEU (European Parliament and Council of the European Union), 2004a. Decision 280/2004/EC of the European Parliament and of the Council concerning a mechanism for monitoring Community greenhouse gas emissions and for implementing the Kyoto Protocol. O.J. L49, 19 February 2004.

EP and CEU (European Parliament and Council of the European Union), 2004b. Directive 2004/42/CE on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC. O.J. L143, 30 April 2004.

EP and CEU (European Parliament and Council of the European Union). Directive 2009/30/EC amending Directive 98/70/EC as regards the specification of petrol, diesel and gas-oil and introducing a mechanism to monitor and reduce greenhouse gas emissions and amending Council Directive 1999/32/EC as regards the specification of fuel used by inland waterway vessels and repealing Directive 93/12/EC.

ERL (Environmental Resources Limited), 1993. Towards a Recycling Strategy for Ireland. Department of the Environment, Custom House, Dublin.

European Semiconductor Industry Association PFC Voluntary Agreement Final Report, 2011

Farrelly, N., NiDhubhain, A., Nieuwenhuis, M., 2011. Productivity models for Sitka spruce in Ireland based on GIS data and readily accessible site quality factors. *Irish Forestry* 68: 6-23.

Fealy, R., and Green, S., "Teagasc-EPA Soils and Subsoils Mapping Project, Final report." .EPA, 2009

Fealy, R., Loftus, M. and Meehan, G. 2006. EPA Soil and Subsoil Mapping Project: Summary, Methodology, Description for Subsoils, Land Cover, Habitat and Soils Mapping/Modelling. EPA project report, Environmental Protection Agency, Johnstown Castle, Wexford.

Fehily Timoney and Company, 1985. Code of Good Practice for the Use of Biosolids in Agriculture. Guidelines for Farmers. Department of Environment and Local Government, Dublin.

Finn, J. et al, 2001. Estimation of Emissions of Non-Methane Volatile Organic Compounds from SNAP Sector 6: Solvent and Other Product Use for Ireland in 1998. Prepared for the Environmental Protection Agency by the Clean Technology Centre, Cork Institute of Technology. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Fogarty, G., Coggins, K and Gallagher, G., 1999, Forest Inventory and Planning System (FIPS). Irish Timber Growers Association Forestry Yearbook 2000. ITGA, Dublin.

Forest Service, 2000. Department of the Marine and Natural Resources Forestry Statistics. Internal publication.

Forestry Scheme Manual (2003). Forest Service, Department of Agriculture Food and the Marine. Dublin.

FSAI (Food Safety Authority of Ireland), 1999. Recommended Dietary Allowances for Ireland. Food Safety Authority of Ireland, Dublin. <http://www.fsai.ie/>

Gallagher, G., Hendrick, E. and Byrne, K. 2004 Preliminary Estimates of Biomass Carbon Stock Changes in Managed Forests in the Republic of Ireland over the period 1990-2000. In Irish Forestry Vol. 61 No 1, pages 16-35.

Gardiner, M. J. and Radford, T., 1980. Soil Associations of Ireland and their Land Use Potential. Explanatory Bulletin to Soil Map of Ireland 1980, Soil Survey Bulletin No 36, An Foras Taluntais, Dublin.

"Gkatzoflias D. (Emisia S.A.), C. Kouridis (Emisia S.A.), L. Ntziachristos (LAT/AUTH), Z. Samaras (LAT/AUTH). February 2012. COPERT 4 Computer programme to calculate emissions from road transport. User manual (version 9.0).

"

Gkatzoflias D., Kouridis C., Ntziachristos L. and Samaras Z., 2007. Copert 4. Computer programme to calculate emissions from road transport. Laboratory of Applied Thermodynamics, Mechanical Engineering Department, Aristotle University Thessaloniki.

Goodwin J., Gschrey B., Salisbury E., 2013, Emission Inventories for HFCs, PFCs and SF6 for inclusion in Ireland's submission to the EU under Decision 280/2004/EC and the United Nations Framework Convention on Climate Change

Hamilton, G.J. 1975. Forest Mensuration Handbook. Forestry Commission Booklet 39, HMSO, London.

Hargreaves, K., Milne, R. and Cannell, M.G.R. 2003. Carbon balance of afforested peatland in Scotland. Forestry, 76, 299-317.

Hawkins M and Black K. 2012. Validating generalised diameter-height models for several species and heterogeneous crop structures. Irish Forestry 69:58.73

Hazeu, G.W. and De Wit, A.J.W. 2004. CORINE land cover database of the Netherlands: monitoring land cover changes between 1986 and 2000. EARSel eProceedings 3 (3). Paris (EARSel)

<http://unfccc.int/resource/docs/2007/arr/irl.pdf>

Hyde, B., Carton, O.T. and Murphy, W.E. (2008). Farm Facilities Survey – Ireland 2003. Report prepared for the Department of Agriculture by Teagasc, Johnstown Castle, Co. Wexford. 150 pages

ICRL, 1990. Notes on the development and after-use of landfill sites. Guidance Note 17/78. Eight Edition, December, 1990. Interdepartmental Committee on the Redevelopment of Contaminated Land.

INRA, 1989. Ruminant Nutrition. Recommended Allowances and Feed Tables. R. Jarrige, Ed. John Libbey Eurotext, London and Paris.

IPCC (Intergovernmental Panel on Climate Change), 1997. Revised 1996 Guidelines for National Greenhouse Gas Inventories. OECD, Paris.

IPCC (Intergovernmental Panel on Climate Change), 2000. IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IGES (Institute for Global Environmental Strategies), Kanagawa, Japan.

IPCC (Intergovernmental Panel on Climate Change), 2003. Good Practice Guidance for Land Use, Land-Use Change and Forestry. IGES (Institute for Global Environmental Strategies), Kanagawa, Japan.

IUCC (Information Unit on Climate Change), 1998. The Kyoto Protocol to the Framework Convention on Climate Change. UNEP, Geneva.

J. Wunder, B. Brzeziecki, H. Zybura, B. Reineking, C. Bigler, and H. Bugmann. Growth-mortality relationships as indicators of life-history strategies: a comparison of nine tree species in unmanaged European forests. *Oikos*, 117:815-828, 2008.

J. Wunder, B. Reineking, C. Bigler, and H. Bugmann. Predicting tree mortality from growth data: how virtual ecologists can help real ecologists. *Journal of Ecology*, 96:174-187, 2006b.

J. Wunder, C. Bigler, R. Bjrn, L. Fahsse, and H. Bugmann. Optimisation of tree mortality models based on growth patterns. *Ecological Modelling*, 197:196-206, 2006a.

Katsis Petros, Leonidas Ntziachristos, Giorgos Mellios. December 3rd, 2012. Description of new elements in COPERT 4 v10.0.

Kenneth A. Byrne, Ger Kiely (2006). Partitioning of respiration in an intensively managed grassland. *Plant and Soil* 282: 281-289.

Kiely G., Xu X., Hsieh C.-I., Leahy P., Sottocornola M., Laine A., Mishurov M., Albertson J., Katul G., Oren R. and O. Carton (2009). CELTICFLUX: Measurement and Modeling of GHG Fluxes from Grasslands and a Peatland in Ireland. *AsiaFlux Newsletter* No 28, March 2009.

L. Broad and T. Lynch. Growth models for Sitka spruce in Ireland. *Irish Forestry*, 63(1-2), 2006a.

L. Broad and T. Lynch. Panel data validation using cross-sectional methods. *Irish Forestry*, 63(1-2), 2006b.

Le Bolloch, Meaney, B., Cope, J and Doyle, M., 2006. National Waste Report 2005: Data Update. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Le Bolloch, O., Cope, J., Kurz, I., Meaney, B. and Higgins, T. 2009. National Waste Report 2007. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Le Bolloch, O., Cope, J., Meaney, B and Kurz, I., 2007. National Waste Report 2006. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

- Li, C., Frolking, S. and Harriss, R. 1994. Modelling carbon biogeochemistry in agricultural soils. *Global Biogeochemical Cycles*, 8, 237-254.
- Li, C., Narayanan, V. and Harriss, R.C. 1996. Model estimates of nitrous oxide emissions from agricultural lands in the United States. *Global Biogeochemical Cycles* 10, 297-306.
- M.I. Khalil, G. Kiely, P. O'Brien and C. Müller . (2012) Organic carbon stocks in agricultural soils in Ireland using combined empirical and GIS approaches. *Geoderma*, 193-194 (2013) 222–235.
- Matthias Peichl, Paul Leahy, and Gerard Kiely (2011). Six-year Stable Annual Uptake of Carbon Dioxide in Intensively Managed Humid Temperate Grassland Ecosystems 14: 112–126 doi:10.1007/s10021-010-9398-2.
- McCoole, F., Derham, J., Kurz, I and McDonagh, M. 2011. National Waste Report 2009. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Derham, J., Kurz, I. and Higgins, T. 2009. National Waste Report 2008. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Kurz, I., McDonagh, M., Derham, J., O'Neill, D. 2012. National Waste Report 2010. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McCoole, F., Kurz, I., McDonagh, M., O'Neill, D. 2013. National Waste Report 2011. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- McGoff, N.M., Eaton, J.M., Byrne, K., Leahy, P. and Kiely, G. 2007. Measurement of soil carbon stocks in Irish soils. British Ecological Society Annual Meeting, September 10-12, University of Glasgow, UK.
- McInnes, G. (editor), 1996. CORINAIR/EMEP Atmospheric Emission Inventory Guidebook. First Edition. European Environment Agency, Copenhagen.
- MCOS (M. C. O'Sullivan), 1994. Report on Solid Waste Recycling (Packaging). Department of the Environment, Custom House, Dublin.
- Meaney, B., Collins, C., Nolan, K., Cahill, E., Delaney, J., Murray, B., Healy, J. and Carty, G. 2003. National Waste Database Report 2001, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Monaghan, S., Smith, D., Wall, B. and O'Leary, G. 2009. Urban Waste Water Discharges in Ireland for Population Equivalents Greater than 500 Persons. A Report for the years 2006 and 2007. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Mulqueen et al., 1999. Forest Drainage Engineering: A design Manual. COFORD, Dublin
- N. A. Obuchowski. Nonparametric analysis of clustered ROC curve data. *Biometrics*, 53:567{578, 1997.
- NEUT (Working Group on Nutrients and Eutrophication under the OSPAR Convention), 1999. Screening Procedure for Irish Coastal Waters with Regard to Eutrophication Status.
- NFI. 2007a. National Forest Inventory - Republic of Ireland - Results. Forest Service, Department of Agriculture, Fisheries and Food, Johnstown Castle Estate, Co Wexford.
- NFI. 2007b. National Forest Inventory - Republic of Ireland - Methodology. Forest Service, Department of Agriculture, Fisheries and Food, Johnstown Castle Estate, Co Wexford.

- O'Brien, P. 2007. Data Analysis and Estimation of Greenhouse Gas Emissions and Removals for the IPCC Sector Land Use Land-Use Change and Forestry in Ireland. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland (in preparation).
- O'Doherty, S. and McCulloch, A. 2002. Emission Inventories of HFC, PFC and SF6. RTDI Project L.S.-5.1.3(a) Final Report. Environmental Protection Agency, Johnstown Castle, Wexford.
- O'Leary, E., 2002. Emission Inventories for HFCs, PFCs and SF6 for Ireland 1995-2000. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Leary, E., Finn, J. and Cunningham, D., 2002. Emission Inventories for HFCs, PFCs and SF6 for Ireland 1998. RTDI Project L.S.-5.1.3(b) Final Report. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Leary, G. and Carty, G. Urban Waste Water Discharges in Ireland 1998. A Report for the years 1996 and 1997. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Leary, G., Fanning, A. and Carty, G. 2000. Urban Waste Water Discharges in Ireland. A Report for the years 1998 and 1999.
- O'Leary, G., Meaney, B. and Carty, G. 1997. Urban Waste Water Discharges in Ireland. A report for the years 1994 and 1995. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Mara, F., 2006. Development of Emission Factors for the Irish Cattle Herd. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Mara, F., P. Caffrey, P. J., and Drennan, M. J., 1997. Net energy values of grass silage determined from comparative feeding trials. Irish Journal of Agricultural and Food Research, 36, 110.
- P. Duffy, E. Hanley, S. Barry, B. Hyde and M.S. Alam. Ireland Informative Inventory Report 2013. Air pollutant emissions in Ireland 1990-2013. Reported to the secretariat of the UNECE convention on long-range transboundary air pollution. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- P. J. Heagerty, T. Lumley, and M. S. Pepe. Time-dependent ROC curves for censored survival data and a diagnostic marker. Biometrics, 56:337{344, 2000.
- P. Martin-Davila, J. Fortun, C. Gutierrez, P. Marti-Belda, A. Candelas, A. Honrubia, R. Barcena, A. Martinez, A. Puente, E. de Vicente, and S. Moreno. Analysis of a quantitative PCR assay for CMV infection in liver transplant recipients: an intent to find the optimal cut-off value. Journal of Clinical Virology, 33:138{144, 2005.
- Pastramas, N.; Samaras, Ch.; Mellios, G.; Ntziachristos, L. May 12, 2014. Update of the Air Emissions Inventory Guidebook - Road Transport 2014 Update.
- Purcell, T. 1979. Inventory of Private Woodlands – 1973. Forest and Wildlife Service, Dublin.
- R. A. Monserud and H. Sterba. Modeling individual tree mortality for Austrian forest species. Forest Ecology and Management, 113:109{123, 1999.
- R. Monserud. Simulation of forest tree mortality. Forest Science, 22:438{444,1976.)
- Regulation of the European Parliament and of the Council No. 525/2013 on a mechanism for monitoring and reporting greenhouse gas emissions and for reporting other information at national and Union level relevant to climate change and repealing Decision No 280/2004/EC

- Richards, K.G., Fenton, O., Khalil, M.I., Haria, A.H., Humphreys, J., Doody, D., Moels, R., Morgan, G. and Jordan, P. 2009. Good water status: The integration of sustainable grassland production and water resources in Ireland. *Tearmann: Irish journal of agri-environmental research*, 7, 143-160, 2009.
- Richardson, S. (editor), 1999. CORINAIR/EMEP Atmospheric Emission Inventory Guidebook. Second Edition. European Environment Agency, Copenhagen.
- RSA.Free speed survey 2008 & 2011 (Urban and Rural), Research Department. 2008 & 2011
- RSA.Survey of Free Speed 2006 &2007: Urban and Rural, Research Department. 2006 & 2007.
- Ryan, M. Brophy, C., Connolly, J., McNamara, K. and Carton, O.T. 2006. Monitoring of nitrogen leaching on a dairy farm during four drainage seasons. *Irish Journal of Agricultural and Food Research* 45: 115–134, 2006.
- S. A. Macskassy, F. Provost, and S. Rosset. ROC confidence bands: An empirical evaluation. In *Proceedings of the 22nd International Conference on Machine Learning*, Bonn, Germany, 2005.
- S. Mason and N. Graham. Areas beneath the relative operating characteristics (ROC) and relative operating levels (ROL) curves: Statistical significance and interpretation. *Quarterly Journal of the Royal Meteorological Society*, 128:2145{2166, 2002.
- Saiz, G., Black, K. and Reidy, B. 2007. Assessment of soil CO₂ efflux and its components using a process-based model in a young temperate forest site. *Goedermia* 139:79-89.
- SBSTA (Subsidiary Body for Scientific and Technological Advice), 1999. Guidelines for the Preparation of National Communications by Parties included in Annex I to the Convention. Part I: UNFCCC Reporting Guidelines on Annual Inventories. FCCC/SBSTA/1999/6/Add.1. UNFCCC Secretariat, Bonn.
- SBSTA (Subsidiary Body for Scientific and Technological Advice), 2002. Guidelines for the Preparation of National Communications by Parties included in Annex I to the Convention. Part I: UNFCCC Reporting Guidelines on Annual Inventories. FCCC/SBSTA/2002/L.5/Add.1, UNFCCC Secretariat, Bonn.
- Smith, D., O'Neill, N., Doris, Y. and Moriarty, J., 2004. Urban Waste Water Discharges in Ireland. A Report for the Years 2002/2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Smith, D., O'Neill, N., Wall, B and Crowe, M. 2007. Urban Waste Water Discharges in Ireland. A Report for the years 2004 and 2005. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Smith, D., O'Neill, N., Clinton, F. and Crowe, M. 2003. Urban Waste Water Discharges in Ireland. A Report for the years 2000 and 2001. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- T. Fawcett. An introduction to ROC analysis. *Pattern Recognition Letters*, 27:861{874, 2006.
- T. Sing, O. Sander, N. Beerenwinkel, and T. Lengauer. ROCR: Visualizing classifier performance in R. *Bioinformatics*, 21(20):3940{3941, 2005.
- Thistlethwaite, G., Baggott, S. L. and Goodwin, J., 2005. National Inventory System Scoping Report for the Greenhouse Gas Emission Inventory in Ireland. AEAT/ENV/R/1990 Issue 1, NETCEN, UK.

Tobin, B., Black, K., Osborne, B. 2006. Assessment of allometric algorithms for estimating leaf biomass, leaf area index and litter fall in different aged Sitka spruce forests. *Forestry* 79: 453-465

Tomlinson, R.W., Cruickshank, M.M., Cruickshank, J.G., Mallon, G., McStravick, A., Byrne, K., Collins, J., O'Toole, P., Farrell, E., Cunningham, D., Milne, R. and Walsh, M. 2005. Impact of land use and land-use change on carbon emission/fixation. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

UNECE (United Nations Economic Commission for Europe), 1999. Protocol to the 1979 Convention on Long-Range Transboundary Air Pollution to Abate Acidification, Eutrophication and Ground Level Ozone. UNECE Secretariat, Geneva.

"UNFCCC (United Nations Framework Convention on Climate Change). 2014.: Report of the individual review of the annual submission of Ireland submitted in 2013.

CCCC/ARR/2013/IRL (In-country review, Ireland)"

"UNFCCC (United Nations Framework Convention on Climate Change). 2015.: Report of the individual review of the annual submission of Ireland submitted in 2014.

CCCC/ARR/2014/IRL (UN Secretariat, Bonn)"

UNFCCC Decision 24/CP.19 Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention

UNFCCC 1/CMP.8 Amendment to the Kyoto Protocol pursuant to its Article 3, paragraph 9 (the Doha Amendment)

WB -World Bank. World Development Indicators (Ireland), 2013. <http://data.worldbank.org/country/ireland>.

Yamulki, S., Anderson, R., Peace, A. & Morison, J.I.L. (2013) Soil CO₂, CH₄ and N₂O fluxes from an afforested lowland raised peatbog in Scotland: implications for drainage and restoration. *Biogeosciences* 10: 1051-1065.

Yan, T., Agnew, R. E., Gordon, F. J. and Porter, M. G. (2000). Prediction of methane energy output in dairy and beef cattle offered grass silage-based diets. *Livestock Production Science*, 64: 253-263.

Annex 1

Key Category Analyses

1.A 2013 Key Category Analysis Level Assessment excluding LULUCF

1.B 2013 Key Category Analysis Level Assessment including LULUCF

1.C 2013 Key Category Analysis Trend Assessment excluding LULUCF

1.D 2013 Key Category Analysis Trend Assessment including LULUCF

1.E Information on the level of disaggregation

1.F Description of methodology used for identifying key categories

Table 1.A 2013 Key Category Analysis Level Assessment excluding LULUCF

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	10483.36	17.82	17.82
2	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6626.21	11.27	29.09
3	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	5633.17	9.58	38.66
4	1.A.1	Energy Industries - Gaseous Fuels	CO2	4373.09	7.43	46.10
5	1.A.1	Energy Industries - Solid Fuels	CO2	3797.89	6.46	52.56
6	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3131.86	5.32	57.88
7	1.A.4.b	Residential - Liquid Fuels	CO2	2753.85	4.68	62.56
8	1.A.1	Energy Industries - Peat Fuel	CO2	2579.98	4.39	66.95
9	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	2039.08	3.47	70.41
10	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	1709.71	2.91	73.32
11	1.A.4.b	Residential - Gaseous Fuels	CO2	1420.84	2.42	75.74
12	2.F.1	Product Uses as Substitutes for ODS -Refrigeration and air-con (incl. MAC)	HFC	1124.16	1.91	77.65
13	2.A.1	Cement Production	CO2	1111.75	1.89	79.54
14	5.A	Solid Waste Disposal	CH4	1106.03	1.88	81.42
15	1.A.4.b	Residential - Solid Fuels	CO2	1098.84	1.87	83.29
16	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1015.60	1.73	85.01
17	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	995.39	1.69	86.71
18	1.A.4.b	Residential - Peat Fuel	CO2	929.40	1.58	88.29
19	3.A.2	Enteric Fermentation - Sheep	CH4	696.09	1.18	89.47
20	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	632.26	1.07	90.54
21	1.A.4.c	Agriculture - Liquid Fuels	CO2	541.91	0.92	91.46
22	3.G.1	Liming - Limestone CaCO3	CO2	515.69	0.88	92.34
23	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	480.10	0.82	93.16
24	1.A.1	Energy Industries - Liquid Fuels	CO2	401.78	0.68	93.84
25	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	325.65	0.55	94.39
26	3.B.1.1	Manure Management - Dairy Cattle	CH4	286.76	0.49	94.88
27	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	225.44	0.38	95.27
28	3.B.2.1	Manure Management - Non-Dairy Cattle	N2O	217.03	0.37	95.63
29	3.B.1.3	Manure Management - Swine	CH4	190.41	0.32	95.96
30	2.A.2	Lime Production	CO2	189.64	0.32	96.28
31	1.A.3.d	Navigation - Liquid Fuels	CO2	175.55	0.30	96.58
32	1.A.3.e	Other Transport - Gaseous Fuels	CO2	148.53	0.25	96.83
33	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	132.53	0.23	97.06
34	5.D	Wastewater treatment and discharge	N2O	118.70	0.20	97.26
35	1.A.3.c	Rail Transport - Liquid Fuels	CO2	117.65	0.20	97.46

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
36	2.F.4	Product Uses as Substitutes for ODS -Aerosols (incl. MDIs)	HFC	114.66	0.19	97.65
37	1.A.3.b	Road Transport - Liquid Fuels	N2O	98.33	0.17	97.82
38	1.A.1	Energy Industries - Other Fuels	CO2	86.50	0.15	97.97
39	1.A.4.b	Residential - Solid Fuels	CH4	85.73	0.15	98.11
40	3.B.1.4	Manure Management - Other livestock	CH4	84.89	0.14	98.26
41	1.A.4.b	Residential - Peat Fuel	CH4	68.42	0.12	98.37
42	1.A.1	Energy Industries - Gaseous Fuels	N2O	67.46	0.11	98.49
43	2.D.3	Solvents	CO2	66.15	0.11	98.60
44	1.A.4.c	Agriculture - Liquid Fuels	N2O	56.84	0.10	98.70
45	5.D	Wastewater treatment and discharge	CH4	50.86	0.09	98.78
46	3.A.3	Enteric Fermentation - Swine	CH4	49.94	0.08	98.87
47	3.A.4	Enteric Fermentation - Other livestock	CH4	49.61	0.08	98.95
48	3.B.1.2	Manure Management - Sheep	CH4	48.01	0.08	99.04
49	1.A.1	Energy Industries - Peat Fuel	N2O	43.93	0.07	99.11
50	5.C.1 & 2	Incineration and open burning of waste - Incineration (fossil C)	CO2	43.05	0.07	99.18
51	2.G.3.a	Other Product Manufacture and Use - Other (Anaesthesia in medical applications)	CO2	41.06	0.07	99.25
52	3.B.2.1	Manure Management - Dairy Cattle	N2O	40.55	0.07	99.32
53	2.E.1	Electronics Industry - Integrated circuit or semiconductor	HFC, PFC, SF6, NF3	34.63	0.06	99.38
54	2.F.3	Product Uses as Substitutes for ODS -Fire protection	HFC	34.53	0.06	99.44
55	1.B.2.b	Fugitive emissions - Natural gas	CH4	23.26	0.04	99.48
56	3.H.	Urea Application	CO2	21.66	0.04	99.52
57	1.B.1.a	Fugitive Emissions - Coal Mining and Handling	CH4	20.15	0.03	99.55
58	2.G.1	Other Product Manufacture and Use - Electrical equipment	SF6	18.60	0.03	99.58
59	2.D.1	Lube oil from Transport	CO2	19.03	0.03	99.61
60	2.D.2	Paraffin wax use	CO2	16.88	0.03	99.64
61	1.A.3.b	Road Transport - Liquid Fuels	CH4	16.47	0.03	99.67
62	1.A.3.c	Rail Transport - Liquid Fuels	N2O	13.68	0.02	99.69
63	5.B.1	Biological treatment of solid waste - Composting	CH4	13.25	0.02	99.72
64	5.B.1	Biological treatment of solid waste - Composting	N2O	11.85	0.02	99.74
65	3.B.2.3	Manure Management - Swine	N2O	11.79	0.02	99.76
66	3.B.2.4	Manure Management - Other livestock	N2O	11.63	0.02	99.78
67	3.B.2.2	Manure Management - Sheep	N2O	11.50	0.02	99.80
68	1.A.3.a	Domestic Aviation - Liquid Fuels	CO2	10.01	0.02	99.81
69	1.A.4.b	Residential - Liquid Fuels	CH4	9.40	0.02	99.83
70	1.A.4.b	Residential - Biomass	CH4	9.06	0.02	99.84
71	1.A.4.a	Commercial/Institutional - Biomass	CH4	7.90	0.01	99.86

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
72	1.A.2	Manufacturing Industries & Construction - Biomass	N2O	7.49	0.01	99.87
73	1.A.4.b	Residential - Liquid Fuels	N2O	6.62	0.01	99.88
74	1.A.1	Energy Industries - Solid Fuels	N2O	6.05	0.01	99.89
75	1.A.1	Energy Industries - Biomass	N2O	5.45	0.01	99.90
76	1.A.4.b	Residential - Solid Fuels	N2O	5.11	0.01	99.91
77	1.A.2	Manufacturing Industries & Construction - Biomass	CH4	4.71	0.01	99.92
78	1.A.4.b	Residential - Peat Fuel	N2O	3.81	0.01	99.92
79	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	N2O	3.36	0.01	99.93
80	1.A.4.a	Commercial/Institutional - Liquid Fuels	CH4	3.35	0.01	99.94
81	1.A.4.b	Residential - Gaseous Fuels	CH4	3.17	0.01	99.94
82	2.G.2	Other Product Manufacture and Use - SF6 and PFCs from other product use	SF6	2.92	0.00	99.95
83	1.A.4.a	Commercial/Institutional - Liquid Fuels	N2O	2.85	0.00	99.95
84	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CH4	2.22	0.00	99.95
85	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CO2	2.16	0.00	99.96
86	1.A.1	Energy Industries - Gaseous Fuels	CH4	2.11	0.00	99.96
87	1.A.1	Energy Industries - Peat Fuel	CH4	1.62	0.00	99.96
88	1.A.3.e	Other Transport - Gaseous Fuels	N2O	1.58	0.00	99.97
89	1.A.2	Manufacturing Industries & Construction - Solid Fuels	N2O	1.54	0.00	99.97
90	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CH4	1.47	0.00	99.97
91	1.A.3.d	Navigation - Liquid Fuels	N2O	1.43	0.00	99.98
92	1.A.4.b	Residential - Biomass	N2O	1.43	0.00	99.98
93	1.A.1	Energy Industries - Biomass	CH4	1.40	0.00	99.98
94	1.A.4.a	Commercial/Institutional - Biomass	N2O	1.26	0.00	99.98
95	1.A.1	Energy Industries - Other Fuels	N2O	1.14	0.00	99.98
96	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	N2O	1.09	0.00	99.99
97	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CH4	0.91	0.00	99.99
98	1.A.4.c	Agriculture - Liquid Fuels	CH4	0.88	0.00	99.99
99	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CH4	0.86	0.00	99.99
100	1.A.4.b	Residential - Gaseous Fuels	N2O	0.76	0.00	99.99
101	1.A.1	Energy Industries - Other Fuels	CH4	0.72	0.00	99.99
102	1.A.1	Energy Industries - Solid Fuels	CH4	0.71	0.00	99.99
103	1.A.4.a	Commercial/Institutional - Gaseous Fuels	N2O	0.53	0.00	99.99
104	5.C.1 & 2	Incineration and open burning of waste - Incineration	N2O	0.45	0.00	100.00
105	1.A.3.d	Navigation - Liquid Fuels	CH4	0.42	0.00	100.00
106	1.B.2.a	Fugitive Emissions - Oil Refining	CH4	0.33	0.00	100.00
107	1.A.3.e	Other Transport - Gaseous Fuels	CH4	0.33	0.00	100.00
108	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	N2O	0.26	0.00	100.00

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
109	1.A.1	Energy Industries - Liquid Fuels	N2O	0.24	0.00	100.00
110	2.A.4.d	Other process uses of carbonates - Limestone use (Other)	CO2	0.21	0.00	100.00
111	1.A.3.c	Rail Transport - Liquid Fuels	CH4	0.17	0.00	100.00
112	1.A.3.a	Domestic Aviation - Liquid Fuels	N2O	0.16	0.00	100.00
113	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CH4	0.11	0.00	100.00
114	1.A.1	Energy Industries - Liquid Fuels	CH4	0.11	0.00	100.00
115	5.C.1 & 2	Incineration and open burning of waste - Incineration	CH4	0.07	0.00	100.00
116	2.A.4.b	Other process uses of carbonates - Soda Ash (Other use)	CO2	0.06	0.00	100.00
117	2.A.4.d	Other process uses of carbonates - Bricks (Other)	CO2	0.03	0.00	100.00
118	1.A.2	Manufacturing Industries & Construction - Peat Fuel	N2O	0.01	0.00	100.00
119	1.A.3.a	Domestic Aviation - Liquid Fuels	CH4	0.01	0.00	100.00
120	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CH4	0.00	0.00	100.00
121	1.A.4.a	Commercial/Institutional - Solid Fuels	CO2	0.00	0.00	100.00
122	1.A.4.a	Commercial/Institutional - Peat Fuel	CO2	0.00	0.00	100.00
123	1.A.4.a	Commercial/Institutional - Solid Fuels	CH4	0.00	0.00	100.00
124	1.A.4.a	Commercial/Institutional - Peat Fuel	CH4	0.00	0.00	100.00
125	1.A.4.a	Commercial/Institutional - Solid Fuels	N2O	0.00	0.00	100.00
126	1.A.4.a	Commercial/Institutional - Peat Fuel	N2O	0.00	0.00	100.00
127	2.A.3	Glass Production	CO2	0.00	0.00	100.00
128	2.B.1	Chemical Industry - Ammonia Production	CO2	0.00	0.00	100.00
129	2.B.2	Chemical Industry - Nitric Acid Production	N2O	0.00	0.00	100.00

Table 1.B 2013 Key Category Analysis Level Assessment including LULUCF

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Absolute Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	10483.36	14.66	14.66
2	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6626.21	9.27	23.93
3	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	5633.17	7.88	31.81
4	4.C.1	LULUCF - Grassland Remaining Grassland	CO2	4763.15	6.66	38.48
5	1.A.1	Energy Industries - Gaseous Fuels	CO2	4373.09	6.12	44.59
6	1.A.1	Energy Industries - Solid Fuels	CO2	3797.89	5.31	49.91
7	4.A.2	LULUCF - Land Converted to Forest Land	CO2	3746.98	5.24	55.15
8	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3131.86	4.38	59.53
9	1.A.4.b	Residential - Liquid Fuels	CO2	2753.85	3.85	63.38
10	4.D.1	LULUCF - Wetlands Remaining wetlands	CO2	2598.00	3.63	67.02
11	1.A.1	Energy Industries - Peat Fuel	CO2	2579.98	3.61	70.63
12	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	2039.08	2.85	73.48
13	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	1709.71	2.39	75.87
14	1.A.4.b	Residential - Gaseous Fuels	CO2	1420.84	1.99	77.86
15	2.F.1	Product Uses as Substitutes for ODS -Refrigeration and air-con (incl. MAC)	HFC	1124.16	1.57	79.43
16	2.A.1	Cement Production	CO2	1111.75	1.56	80.99
17	5.A	Solid Waste Disposal	CH4	1106.03	1.55	82.53
18	1.A.4.b	Residential - Solid Fuels	CO2	1098.84	1.54	84.07
19	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1015.60	1.42	85.49
20	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	995.39	1.39	86.88
21	1.A.4.b	Residential - Peat Fuel	CO2	929.40	1.30	88.18
22	3.A.2	Enteric Fermentation - Sheep	CH4	696.09	0.97	89.16
23	4.G	LULUCF - Harvested wood products	CO2	691.95	0.97	90.12
24	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	632.26	0.88	91.01
25	1.A.4.c	Agriculture - Liquid Fuels	CO2	541.91	0.76	91.77
26	3.G.1	Liming - Limestone CaCO3	CO2	515.69	0.72	92.49
27	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	480.10	0.67	93.16
28	1.A.1	Energy Industries - Liquid Fuels	CO2	401.78	0.56	93.72
29	4.B.1	LULUCF - Cropland Remaining Cropland	CO2	364.38	0.51	94.23
30	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	325.65	0.46	94.69
31	3.B.1.1	Manure Management - Dairy Cattle	CH4	286.76	0.40	95.09
32	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	225.44	0.32	95.40
33	3.B.2.1	Manure Management - Non-Dairy Cattle	N2O	217.03	0.30	95.71
34	3.B.1.3	Manure Management - Swine	CH4	190.41	0.27	95.97
35	2.A.2	Lime Production	CO2	189.64	0.27	96.24
36	1.A.3.d	Navigation - Liquid Fuels	CO2	175.55	0.25	96.48
37	1.A.3.e	Other Transport - Gaseous Fuels	CO2	148.53	0.21	96.69
38	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	132.53	0.19	96.88
39	5.D	Wastewater treatment and discharge	N2O	118.70	0.17	97.04
40	1.A.3.c	Rail Transport - Liquid Fuels	CO2	117.65	0.16	97.21
41	2.F.4	Product Uses as Substitutes for ODS -Aerosols (incl. MDIs)	HFC	114.66	0.16	97.37

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Absolute Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
42	1.A.3.b	Road Transport - Liquid Fuels	N2O	98.33	0.14	97.51
43	1.A.1	Energy Industries - Other Fuels	CO2	86.50	0.12	97.63
44	1.A.4.b	Residential - Solid Fuels	CH4	85.73	0.12	97.75
45	3.B.1.4	Manure Management - Other livestock	CH4	84.89	0.12	97.87
46	4.D.1	LULUCF - Wetlands Remaining wetlands	CH4	78.36	0.11	97.98
47	4.A.1	LULUCF - Forest land Remaining Forest Land	CO2	74.00	0.10	98.08
48	1.A.4.b	Residential - Peat Fuel	CH4	68.42	0.10	98.18
49	1.A.1	Energy Industries - Gaseous Fuels	N2O	67.46	0.09	98.27
50	4.E.2	LULUCF - Land Converted to Settlements	N2O	67.14	0.09	98.36
51	2.D.3	Solvents	CO2	66.15	0.09	98.46
52	4.C.2	LULUCF - Land Converted to Grassland	CO2	59.33	0.08	98.54
53	1.A.4.c	Agriculture - Liquid Fuels	N2O	56.84	0.08	98.62
54	4.E.2	LULUCF - Land Converted to Settlements	CO2	54.96	0.08	98.70
55	5.D	Wastewater treatment and discharge	CH4	50.86	0.07	98.77
56	4.F.2	LULUCF - Land Converted to Other Land	N2O	50.81	0.07	98.84
57	3.A.3	Enteric Fermentation - Swine	CH4	49.94	0.07	98.91
58	3.A.4	Enteric Fermentation - Other livestock	CH4	49.61	0.07	98.98
59	3.B.1.2	Manure Management - Sheep	CH4	48.01	0.07	99.04
60	4.C.1	LULUCF - Grassland Remaining Grassland	CH4	44.09	0.06	99.11
61	1.A.1	Energy Industries - Peat Fuel	N2O	43.93	0.06	99.17
62	5.C.1 & 2	Incineration and open burning of waste - Incineration (fossil C)	CO2	43.05	0.06	99.23
63	2.G.3.a	Other Product Manufacture and Use - Other (Anaesthesia in medical applications)	CO2	41.06	0.06	99.28
64	3.B.2.1	Manure Management - Dairy Cattle	N2O	40.55	0.06	99.34
65	2.E.1	Electronics Industry - Integrated circuit or semiconductor	HFC, PFC, SF6, NF3	34.63	0.05	99.39
66	2.F.3	Product Uses as Substitutes for ODS -Fire protection	HFC	34.53	0.05	99.44
67	4.D.1	LULUCF - Wetlands Remaining wetlands	N2O	32.75	0.05	99.48
68	1.B.2.b	Fugitive emissions - Natural gas	CH4	23.26	0.03	99.52
69	3.H.	Urea Application	CO2	21.66	0.03	99.55
70	1.B.1.a	Fugitive Emissions - Coal Mining and Handling	CH4	20.15	0.03	99.58
71	2.G.1	Other Product Manufacture and Use - Electrical equipment	SF6	18.60	0.03	99.60
72	2.D.1	Lube oil from Transport	CO2	19.03	0.03	99.63
73	2.D.2	Paraffin wax use	CO2	16.88	0.02	99.65
74	1.A.3.b	Road Transport - Liquid Fuels	CH4	16.47	0.02	99.67
75	1.A.3.c	Rail Transport - Liquid Fuels	N2O	13.68	0.02	99.69
76	5.B.1	Biological treatment of solid waste - Composting	CH4	13.25	0.02	99.71
77	5.B.1	Biological treatment of solid waste - Composting	N2O	11.85	0.02	99.73
78	3.B.2.3	Manure Management - Swine	N2O	11.79	0.02	99.75
79	3.B.2.4	Manure Management - Other livestock	N2O	11.63	0.02	99.76
80	3.B.2.2	Manure Management - Sheep	N2O	11.50	0.02	99.78
81	4.F.2	LULUCF - Land Converted to Other Land	CO2	11.06	0.02	99.79
82	1.A.3.a	Domestic Aviation - Liquid Fuels	CO2	10.01	0.01	99.81
83	1.A.4.b	Residential - Liquid Fuels	CH4	9.40	0.01	99.82

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Absolute Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
84	1.A.4.b	Residential - Biomass	CH4	9.06	0.01	99.83
85	1.A.4.a	Commercial/Institutional - Biomass	CH4	7.90	0.01	99.84
86	1.A.2	Manufacturing Industries & Construction - Biomass	N2O	7.49	0.01	99.85
87	4.D.2	LULUCF - Land converted to wetlands	CO2	7.12	0.01	99.86
88	4.A.1	LULUCF - Forest land Remaining Forest Land	CH4	7.08	0.01	99.87
89	4.C.1	LULUCF - Grassland Remaining Grassland	N2O	6.66	0.01	99.88
90	1.A.4.b	Residential - Liquid Fuels	N2O	6.62	0.01	99.89
91	1.A.1	Energy Industries - Solid Fuels	N2O	6.05	0.01	99.90
92	1.A.1	Energy Industries - Biomass	N2O	5.45	0.01	99.91
93	1.A.4.b	Residential - Solid Fuels	N2O	5.11	0.01	99.92
94	1.A.2	Manufacturing Industries & Construction - Biomass	CH4	4.71	0.01	99.92
95	1.A.4.b	Residential - Peat Fuel	N2O	3.81	0.01	99.93
96	4.C.2	LULUCF - Land Converted to Grassland	N2O	3.74	0.01	99.93
97	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	N2O	3.36	0.00	99.94
98	1.A.4.a	Commercial/Institutional - Liquid Fuels	CH4	3.35	0.00	99.94
99	1.A.4.b	Residential - Gaseous Fuels	CH4	3.17	0.00	99.95
100	2.G.2	Other Product Manufacture and Use - SF6 and PFCs from other product use	SF6	2.92	0.00	99.95
101	1.A.4.a	Commercial/Institutional - Liquid Fuels	N2O	2.85	0.00	99.95
102	4.A.2	LULUCF - Land converted to Forest Land	CH4	2.64	0.00	99.96
103	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CH4	2.22	0.00	99.96
104	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CO2	2.16	0.00	99.96
105	1.A.1	Energy Industries - Gaseous Fuels	CH4	2.11	0.00	99.97
106	1.A.1	Energy Industries - Peat Fuel	CH4	1.62	0.00	99.97
107	1.A.3.e	Other Transport - Gaseous Fuels	N2O	1.58	0.00	99.97
108	1.A.2	Manufacturing Industries & Construction - Solid Fuels	N2O	1.54	0.00	99.97
109	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CH4	1.47	0.00	99.98
110	1.A.3.d	Navigation - Liquid Fuels	N2O	1.43	0.00	99.98
111	1.A.4.b	Residential - Biomass	N2O	1.43	0.00	99.98
112	1.A.1	Energy Industries - Biomass	CH4	1.40	0.00	99.98
113	1.A.4.a	Commercial/Institutional - Biomass	N2O	1.26	0.00	99.98
114	1.A.1	Energy Industries - Other Fuels	N2O	1.14	0.00	99.99
115	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	N2O	1.09	0.00	99.99
116	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CH4	0.91	0.00	99.99
117	1.A.4.c	Agriculture - Liquid Fuels	CH4	0.88	0.00	99.99
118	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CH4	0.86	0.00	99.99
119	1.A.4.b	Residential - Gaseous Fuels	N2O	0.76	0.00	99.99
120	1.A.1	Energy Industries - Other Fuels	CH4	0.72	0.00	99.99
121	1.A.1	Energy Industries - Solid Fuels	CH4	0.71	0.00	99.99
122	1.A.4.a	Commercial/Institutional - Gaseous Fuels	N2O	0.53	0.00	99.99
123	4.A.1	LULUCF - Forest land Remaining Forest Land	N2O	0.49	0.00	100.00
124	5.C.1 & 2	Incineration and open burning of waste - Incineration	N2O	0.45	0.00	100.00
125	1.A.3.d	Navigation - Liquid Fuels	CH4	0.42	0.00	100.00
126	1.B.2.a	Fugitive Emissions - Oil Refining	CH4	0.33	0.00	100.00

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	Absolute Values (kt CO ₂ eq)	2013 Level assessment %	Cumulative Total %
127	1.A.3.e	Other Transport - Gaseous Fuels	CH4	0.33	0.00	100.00
128	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	N2O	0.26	0.00	100.00
129	1.A.1	Energy Industries - Liquid Fuels	N2O	0.24	0.00	100.00
130	2.A.4.d	Other process uses of carbonates - Limestone use (Other)	CO2	0.21	0.00	100.00
131	4.A.2	LULUCF - Land converted to Forest Land	N2O	0.18	0.00	100.00
132	1.A.3.c	Rail Transport - Liquid Fuels	CH4	0.17	0.00	100.00
133	1.A.3.a	Domestic Aviation - Liquid Fuels	N2O	0.16	0.00	100.00
134	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CH4	0.11	0.00	100.00
135	1.A.1	Energy Industries - Liquid Fuels	CH4	0.11	0.00	100.00
136	4.B.1	LULUCF - Cropland Remaining Cropland	CH4	0.08	0.00	100.00
137	5.C.1 & 2	Incineration and open burning of waste - Incineration	CH4	0.07	0.00	100.00
138	2.A.4.b	Other process uses of carbonates - Soda Ash (Other use)	CO2	0.06	0.00	100.00
139	2.A.4.d	Other process uses of carbonates - Bricks (Other)	CO2	0.03	0.00	100.00
140	4.B.1	LULUCF - Cropland Remaining Cropland	N2O	0.02	0.00	100.00
141	1.A.2	Manufacturing Industries & Construction - Peat Fuel	N2O	0.01	0.00	100.00
142	1.A.3.a	Domestic Aviation - Liquid Fuels	CH4	0.01	0.00	100.00
143	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CH4	0.00	0.00	100.00
144	1.A.4.a	Commercial/Institutional - Solid Fuels	CO2	0.00	0.00	100.00
145	1.A.4.a	Commercial/Institutional - Peat Fuel	CO2	0.00	0.00	100.00
146	1.A.4.a	Commercial/Institutional - Solid Fuels	CH4	0.00	0.00	100.00
147	1.A.4.a	Commercial/Institutional - Peat Fuel	CH4	0.00	0.00	100.00
148	1.A.4.a	Commercial/Institutional - Solid Fuels	N2O	0.00	0.00	100.00
149	1.A.4.a	Commercial/Institutional - Peat Fuel	N2O	0.00	0.00	100.00
150	2.A.3	Glass Production	CO2	0.00	0.00	100.00
151	2.B.1	Chemical Industry - Ammonia Production	CO2	0.00	0.00	100.00
152	2.B.2	Chemical Industry - Nitric Acid Production	N2O	0.00	0.00	100.00

Table 1.C 2013 Key Category Analysis Trend Assessment excluding LULUCF

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	4690.42	10483.36	17.82	20.26	20.26
2	1.A.1	Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	7.43	8.73	28.99
3	1.A.4.b	Residential - Peat Fuel	CO2	3123.37	929.40	1.58	8.32	37.31
4	1.A.4.b	Residential - Liquid Fuels	CO2	1175.35	2753.85	4.68	5.53	42.84
5	1.A.4.b	Residential - Solid Fuels	CO2	2483.42	1098.84	1.87	5.32	48.16
6	1.A.1	Energy Industries - Solid Fuels	CO2	4844.66	3797.89	6.46	4.41	52.57
7	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6581.38	5633.17	9.58	4.28	56.85
8	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	1420.84	2.42	4.11	60.96
9	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	873.02	2039.08	3.47	4.09	65.05
10	2.F.1	Product Uses as Substitutes for ODS -Refrigeration and air-con (incl. MAC)	HFC	0.00	1124.16	1.91	4.05	69.10
11	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1957.00	1015.60	1.73	3.65	72.75
12	1.A.1	Energy Industries - Liquid Fuels	CO2	1254.90	401.78	0.68	3.24	75.99
13	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	995.39	1.69	2.75	78.74
14	1.A.1	Energy Industries - Peat Fuel	CO2	3164.78	2579.98	4.39	2.52	81.26
15	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	325.65	0.55	2.08	83.35
16	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2198.38	1709.71	2.91	2.05	85.40
17	3.A.2	Enteric Fermentation - Sheep	CH4	1176.34	696.09	1.18	1.89	87.28
18	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3398.80	3131.86	5.32	1.41	88.69
19	5.A	Solid Waste Disposal	CH4	1396.49	1106.03	1.88	1.23	89.92
20	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6702.59	6626.21	11.27	1.16	91.08
21	2.A.1	Cement Production	CO2	884.00	1111.75	1.89	0.70	91.78
22	1.A.4.b	Residential - Peat Fuel	CH4	227.65	68.42	0.12	0.60	92.38
23	3.G.1	Liming - Limestone CaCO3	CO2	355.04	515.69	0.88	0.53	92.92
24	1.A.4.c	Agriculture - Liquid Fuels	CO2	660.30	541.91	0.92	0.51	93.43
25	1.B.2.b	Fugitive emissions - Natural gas	CH4	156.05	23.26	0.04	0.50	93.93
26	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	0.00	132.53	0.23	0.48	94.41
27	1.A.4.b	Residential - Solid Fuels	CH4	196.51	85.73	0.15	0.42	94.83
28	2.F.4	Product Uses as Substitutes for ODS -Aerosols (incl. MDIs)	HFC	0.00	114.66	0.19	0.41	95.24
29	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	559.22	480.10	0.82	0.36	95.60
30	1.A.3.d	Navigation - Liquid Fuels	CO2	84.90	175.55	0.30	0.32	95.92
31	1.A.1	Energy Industries - Other Fuels	CO2	0.00	86.50	0.15	0.31	96.23
32	1.A.3.e	Other Transport - Gaseous Fuels	CO2	62.04	148.53	0.25	0.30	96.53
33	3.B.1.1	Manure Management - Dairy Cattle	CH4	354.22	286.76	0.49	0.29	96.82
34	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	684.58	632.26	1.07	0.28	97.10
35	1.A.1	Energy Industries - Gaseous Fuels	N2O	10.21	67.46	0.11	0.20	97.31
36	5.C.1 & 2	Incineration and open burning of waste - Incineration (fossil C)	CO2	90.61	43.05	0.07	0.18	97.49
37	1.A.3.b	Road Transport - Liquid Fuels	N2O	48.09	98.33	0.17	0.17	97.66
38	1.A.3.a	Domestic Aviation - Liquid Fuels	CO2	51.13	10.01	0.02	0.15	97.82
39	1.B.1.a	Fugitive Emissions - Coal Mining and Handling	CH4	55.56	20.15	0.03	0.13	97.95

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
40	3.B.1.2	Manure Management - Sheep	CH4	81.95	48.01	0.08	0.13	98.09
41	2.F.3	Product Uses as Substitutes for ODS - Fire protection	HFC	0.00	34.53	0.06	0.12	98.21
42	2.A.2	Lime Production	CO2	214.08	189.64	0.32	0.12	98.33
43	2.E.1	Electronics Industry - Integrated circuit or semiconductor	HFC, PFC, SF6, NF3	1.17	34.63	0.06	0.12	98.45
44	1.A.3.b	Road Transport - Liquid Fuels	CH4	47.47	16.47	0.03	0.12	98.57
45	3.H.	Urea Application	CO2	44.47	21.66	0.04	0.09	98.65
46	3.B.1.3	Manure Management - Swine	CH4	160.23	190.41	0.32	0.09	98.74
47	3.B.1.4	Manure Management - Other livestock	CH4	61.32	84.89	0.14	0.08	98.82
48	1.A.3.c	Rail Transport - Liquid Fuels	CO2	133.19	117.65	0.20	0.07	98.89
49	3.B.2.1	Manure Management - Non-Dairy Cattle	N2O	189.89	217.03	0.37	0.07	98.96
50	5.D	Wastewater treatment and discharge	N2O	95.64	118.70	0.20	0.07	99.04
51	2.D.1	Lube oil from Transport	CO2	0.00	19.03	0.03	0.07	99.10
52	2.D.3	Solvents	CO2	79.33	66.15	0.11	0.06	99.16
53	1.A.4.c	Agriculture - Liquid Fuels	N2O	69.26	56.84	0.10	0.05	99.22
54	5.B.1	Biological treatment of solid waste - Composting	CH4	0.00	13.25	0.02	0.05	99.26
55	3.B.2.1	Manure Management - Dairy Cattle	N2O	51.18	40.55	0.07	0.05	99.31
56	5.D	Wastewater treatment and discharge	CH4	61.10	50.86	0.09	0.04	99.35
57	5.B.1	Biological treatment of solid waste - Composting	N2O	0.00	11.85	0.02	0.04	99.40
58	2.D.2	Paraffin wax use	CO2	5.09	16.88	0.03	0.04	99.44
59	2.G.2	Other Product Manufacture and Use - SF6 and PFCs from other product use	SF6	12.90	2.92	0.00	0.04	99.48
60	3.A.4	Enteric Fermentation - Other livestock	CH4	37.87	49.61	0.08	0.04	99.51
61	1.A.1	Energy Industries - Peat Fuel	N2O	52.07	43.93	0.07	0.04	99.55
62	1.A.4.b	Residential - Peat Fuel	N2O	12.66	3.81	0.01	0.03	99.58
63	2.G.3.a	Other Product Manufacture and Use - Other (Anaesthesia in medical applications)	CO2	31.34	41.06	0.07	0.03	99.61
64	1.A.4.a	Commercial/Institutional - Biomass	CH4	0.00	7.90	0.01	0.03	99.64
65	3.B.2.2	Manure Management - Sheep	N2O	18.33	11.50	0.02	0.03	99.67
66	3.A.3	Enteric Fermentation - Swine	CH4	41.37	49.94	0.08	0.03	99.69
67	1.A.4.b	Residential - Solid Fuels	N2O	11.71	5.11	0.01	0.03	99.72
68	1.A.4.b	Residential - Liquid Fuels	CH4	3.71	9.40	0.02	0.02	99.74
69	1.A.4.b	Residential - Biomass	CH4	14.08	9.06	0.02	0.02	99.76
70	1.A.1	Energy Industries - Biomass	N2O	0.00	5.45	0.01	0.02	99.78
71	2.A.4.d	Other process uses of carbonates - Bricks (Other)	CO2	5.07	0.03	0.00	0.02	99.80
72	1.A.2	Manufacturing Industries & Construction - Biomass	N2O	3.04	7.49	0.01	0.02	99.81
73	1.A.4.b	Residential - Liquid Fuels	N2O	2.48	6.62	0.01	0.01	99.83
74	1.A.4.a	Commercial/Institutional - Liquid Fuels	CH4	6.58	3.35	0.01	0.01	99.84
75	1.A.2	Manufacturing Industries & Construction - Biomass	CH4	1.91	4.71	0.01	0.01	99.85
76	1.A.2	Manufacturing Industries & Construction - Solid Fuels	N2O	4.12	1.54	0.00	0.01	99.86
77	2.G.1	Other Product Manufacture and Use - Electrical equipment	SF6	20.52	18.60	0.03	0.01	99.87
78	1.A.4.b	Residential - Gaseous Fuels	CH4	0.61	3.17	0.01	0.01	99.88
79	1.A.3.c	Rail Transport - Liquid Fuels	N2O	15.49	13.68	0.02	0.01	99.89
80	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CO2	0.00	2.16	0.00	0.01	99.90
81	1.A.4.a	Commercial/Institutional - Liquid Fuels	N2O	4.68	2.85	0.00	0.01	99.90

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
82	1.A.1	Energy Industries - Solid Fuels	N2O	7.74	6.05	0.01	0.01	99.91
83	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CH4	0.51	2.22	0.00	0.01	99.92
84	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	N2O	4.83	3.36	0.01	0.01	99.92
85	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CH4	2.30	0.86	0.00	0.01	99.93
86	1.A.1	Energy Industries - Gaseous Fuels	CH4	3.43	2.11	0.00	0.01	99.93
87	1.A.1	Energy Industries - Biomass	CH4	0.00	1.40	0.00	0.01	99.94
88	3.B.2.3	Manure Management - Swine	N2O	10.08	11.79	0.02	0.00	99.94
89	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	216.24	225.44	0.38	0.00	99.95
90	1.A.1	Energy Industries - Liquid Fuels	N2O	1.47	0.24	0.00	0.00	99.95
91	1.A.4.a	Commercial/Institutional - Biomass	N2O	0.00	1.26	0.00	0.00	99.96
92	1.A.1	Energy Industries - Other Fuels	N2O	0.00	1.14	0.00	0.00	99.96
93	3.B.2.4	Manure Management - Other livestock	N2O	10.31	11.63	0.02	0.00	99.96
94	1.A.4.b	Residential - Biomass	N2O	2.23	1.43	0.00	0.00	99.97
95	1.A.3.e	Other Transport - Gaseous Fuels	N2O	0.67	1.58	0.00	0.00	99.97
96	5.C.1 & 2	Incineration and open burning of waste - Incineration	CH4	0.83	0.07	0.00	0.00	99.97
97	1.A.3.d	Navigation - Liquid Fuels	N2O	0.67	1.43	0.00	0.00	99.98
98	1.A.1	Energy Industries - Other Fuels	CH4	0.00	0.72	0.00	0.00	99.98
99	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CH4	2.06	1.47	0.00	0.00	99.98
100	5.C.1 & 2	Incineration and open burning of waste - Incineration	N2O	1.04	0.45	0.00	0.00	99.98
101	1.A.4.b	Residential - Gaseous Fuels	N2O	0.15	0.76	0.00	0.00	99.99
102	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	N2O	0.47	1.09	0.00	0.00	99.99
103	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CH4	0.40	0.91	0.00	0.00	99.99
104	1.A.3.a	Domestic Aviation - Liquid Fuels	N2O	0.55	0.16	0.00	0.00	99.99
105	1.A.4.a	Commercial/Institutional - Gaseous Fuels	N2O	0.12	0.53	0.00	0.00	99.99
106	1.A.1	Energy Industries - Peat Fuel	CH4	1.90	1.62	0.00	0.00	99.99
107	1.A.1	Energy Industries - Liquid Fuels	CH4	0.39	0.11	0.00	0.00	99.99
108	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	N2O	0.00	0.26	0.00	0.00	100.00
109	1.A.1	Energy Industries - Solid Fuels	CH4	0.91	0.71	0.00	0.00	100.00
110	1.A.4.c	Agriculture - Liquid Fuels	CH4	1.07	0.88	0.00	0.00	100.00
111	1.A.3.d	Navigation - Liquid Fuels	CH4	0.20	0.42	0.00	0.00	100.00
112	1.A.3.e	Other Transport - Gaseous Fuels	CH4	0.14	0.33	0.00	0.00	100.00
113	1.B.2.a	Fugitive Emissions - Oil Refining	CH4	0.21	0.33	0.00	0.00	100.00
114	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CH4	0.00	0.11	0.00	0.00	100.00
115	2.A.4.d	Other process uses of carbonates - Limestone use (Other)	CO2	0.15	0.21	0.00	0.00	100.00
116	2.A.4.b	Other process uses of carbonates - Soda Ash (Other use)	CO2	0.10	0.06	0.00	0.00	100.00
117	1.A.3.c	Rail Transport - Liquid Fuels	CH4	0.19	0.17	0.00	0.00	100.00
118	1.A.3.a	Domestic Aviation - Liquid Fuels	CH4	0.02	0.01	0.00	0.00	100.00
119	1.A.2	Manufacturing Industries & Construction - Peat Fuel	N2O	0.00	0.01	0.00	0.00	100.00
120	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CH4	0.00	0.00	0.00	0.00	100.00
121	1.A.4.a	Commercial/Institutional - Solid Fuels	CO2	2.56	0.00	0.00	0.00	100.00
122	1.A.4.a	Commercial/Institutional - Peat Fuel	CO2	135.73	0.00	0.00	0.00	100.00
123	1.A.4.a	Commercial/Institutional - Solid Fuels	CH4	0.01	0.00	0.00	0.00	100.00

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
124	1.A.4.a	Commercial/Institutional - Peat Fuel	CH4	0.33	0.00	0.00	0.00	100.00
125	1.A.4.a	Commercial/Institutional - Solid Fuels	N2O	0.01	0.00	0.00	0.00	100.00
126	1.A.4.a	Commercial/Institutional - Peat Fuel	N2O	0.56	0.00	0.00	0.00	100.00
127	2.A.3	Glass Production	CO2	13.33	0.00	0.00	0.00	100.00
128	2.B.1	Chemical Industry - Ammonia Production	CO2	990.23	0.00	0.00	0.00	100.00
129	2.B.2	Chemical Industry - Nitric Acid Production	N2O	995.32	0.00	0.00	0.00	100.00

Table 1.D 2013 Key Category Analysis Trend Assessment including LULUCF

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
1	1.A.3.b	Road Transport - Liquid Fuels	CO2	4690.42	10483.36	14.66	15.29	15.29
2	4.A.2	LULUCF - Land Converted to Forest Land	CO2	27.26	3746.98	5.24	10.00	25.28
3	4.A.1	LULUCF - Forest land Remaining Forest Land	CO2	2719.66	74.00	0.10	7.27	32.56
4	1.A.1	Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	6.12	6.59	39.14
5	1.A.4.b	Residential - Peat Fuel	CO2	3123.37	929.40	1.30	6.08	45.22
6	1.A.4.b	Residential - Liquid Fuels	CO2	1175.35	2753.85	3.85	4.17	49.40
7	1.A.4.b	Residential - Solid Fuels	CO2	2483.42	1098.84	1.54	3.87	53.27
8	4.C.1	LULUCF - Grassland Remaining Grassland	CO2	6067.33	4763.15	6.66	3.87	57.13
9	4.B.1	LULUCF - Cropland Remaining Cropland	CO2	1524.52	364.38	0.51	3.21	60.34
10	1.A.1	Energy Industries - Solid Fuels	CO2	4844.66	3797.89	5.31	3.10	63.44
11	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CO2	873.02	2039.08	2.85	3.08	66.52
12	1.A.4.b	Residential - Gaseous Fuels	CO2	269.73	1420.84	1.99	3.08	69.60
13	2.F.1	Product Uses as Substitutes for ODS -Refrigeration and air-con (incl. MAC)	HFC	0.00	1124.16	1.57	3.02	72.62
14	3.D.1	Agricultural Soils - Direct Soil Emissions	N2O	6581.38	5633.17	7.88	2.94	75.56
15	1.A.4.a	Commercial/Institutional - Liquid Fuels	CO2	1957.00	1015.60	1.42	2.65	78.21
16	1.A.1	Energy Industries - Liquid Fuels	CO2	1254.90	401.78	0.56	2.37	80.57
17	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CO2	223.49	995.39	1.39	2.06	82.63
18	1.A.1	Energy Industries - Peat Fuel	CO2	3164.78	2579.98	3.61	1.76	84.39
19	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CO2	871.24	325.65	0.46	1.52	85.91
20	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CO2	2198.38	1709.71	2.39	1.44	87.36
21	3.A.2	Enteric Fermentation - Sheep	CH4	1176.34	696.09	0.97	1.36	88.72
22	4.D.1	LULUCF - Wetlands Remaining wetlands	CO2	2189.38	2598.00	3.63	0.97	89.68
23	3.A.1	Enteric Fermentation - Dairy Cattle	CH4	3398.80	3131.86	4.38	0.92	90.60
24	5.A	Solid Waste Disposal	CH4	1396.49	1106.03	1.55	0.86	91.47
25	4.G	LULUCF - Harvested wood products	CO2	410.59	691.95	0.97	0.73	92.20
26	3.A.1	Enteric Fermentation - Non-Dairy Cattle	CH4	6702.59	6626.21	9.27	0.60	92.80
27	2.A.1	Cement Production	CO2	884.00	1111.75	1.56	0.56	93.36
28	1.A.4.b	Residential - Peat Fuel	CH4	227.65	68.42	0.10	0.44	93.80
29	3.G.1	Liming - Limestone CaCO3	CO2	355.04	515.69	0.72	0.41	94.21
30	1.B.2.b	Fugitive emissions - Natural gas	CH4	156.05	23.26	0.03	0.37	94.58
31	1.A.4.c	Agriculture - Liquid Fuels	CO2	660.30	541.91	0.76	0.36	94.94
32	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CO2	0.00	132.53	0.19	0.36	95.29
33	1.A.4.b	Residential - Solid Fuels	CH4	196.51	85.73	0.12	0.31	95.60
34	2.F.4	Product Uses as Substitutes for ODS -Aerosols (incl. MDIs)	HFC	0.00	114.66	0.16	0.31	95.91
35	3.D.2	Agricultural Soils - Indirect Soil Emissions	N2O	559.22	480.10	0.67	0.25	96.16
36	1.A.3.d	Navigation - Liquid Fuels	CO2	84.90	175.55	0.25	0.24	96.40
37	1.A.1	Energy Industries - Other Fuels	CO2	0.00	86.50	0.12	0.23	96.63
38	1.A.3.e	Other Transport - Gaseous Fuels	CO2	62.04	148.53	0.21	0.23	96.86
39	3.B.1.1	Manure Management - Dairy Cattle	CH4	354.22	286.76	0.40	0.20	97.06
40	3.B.1.1	Manure Management - Non-Dairy Cattle	CH4	684.58	632.26	0.88	0.18	97.24

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
41	4.E.2	LULUCF - Land Converted to Settlements	N2O	5.76	67.14	0.09	0.16	97.41
42	1.A.1	Energy Industries - Gaseous Fuels	N2O	10.21	67.46	0.09	0.15	97.56
43	4.C.2	LULUCF - Land Converted to Grassland	CO2	2.71	59.33	0.08	0.15	97.71
44	4.F.2	LULUCF - Land Converted to Other Land	N2O	0.08	50.81	0.07	0.14	97.85
45	5.C.1 & 2	Incineration and open burning of waste - Incineration (fossil C)	CO2	90.61	43.05	0.06	0.13	97.98
46	1.A.3.b	Road Transport - Liquid Fuels	N2O	48.09	98.33	0.14	0.13	98.11
47	1.A.3.a	Domestic Aviation - Liquid Fuels	CO2	51.13	10.01	0.01	0.11	98.23
48	1.B.1.a	Fugitive Emissions - Coal Mining and Handling	CH4	55.56	20.15	0.03	0.10	98.32
49	3.B.1.2	Manure Management - Sheep	CH4	81.95	48.01	0.07	0.10	98.42
50	2.F.3	Product Uses as Substitutes for ODS -Fire protection	HFC	0.00	34.53	0.05	0.09	98.51
51	2.E.1	Electronics Industry - Integrated circuit or semiconductor	HFC, PFC, SF6, NF3	1.17	34.63	0.05	0.09	98.60
52	1.A.3.b	Road Transport - Liquid Fuels	CH4	47.47	16.47	0.02	0.09	98.69
53	2.A.2	Lime Production	CO2	214.08	189.64	0.27	0.08	98.77
54	3.B.1.3	Manure Management - Swine	CH4	160.23	190.41	0.27	0.07	98.84
55	3.H.	Urea Application	CO2	44.47	21.66	0.03	0.06	98.90
56	3.B.2.1	Manure Management - Non-Dairy Cattle	N2O	189.89	217.03	0.30	0.06	98.96
57	3.B.1.4	Manure Management - Other livestock	CH4	61.32	84.89	0.12	0.06	99.02
58	5.D	Wastewater treatment and discharge	N2O	95.64	118.70	0.17	0.06	99.08
59	4.E.2	LULUCF - Land Converted to Settlements	CO2	73.88	54.96	0.08	0.06	99.14
60	1.A.3.c	Rail Transport - Liquid Fuels	CO2	133.19	117.65	0.16	0.05	99.19
61	2.D.1	Lube oil from Transport	CO2	0.00	19.03	0.03	0.05	99.24
62	2.D.3	Solvents	CO2	79.33	66.15	0.09	0.04	99.28
63	1.A.4.c	Agriculture - Liquid Fuels	N2O	69.26	56.84	0.08	0.04	99.31
64	5.B.1	Biological treatment of solid waste - Composting	CH4	0.00	13.25	0.02	0.04	99.35
65	5.B.1	Biological treatment of solid waste - Composting	N2O	0.00	11.85	0.02	0.03	99.38
66	3.B.2.1	Manure Management - Dairy Cattle	N2O	51.18	40.55	0.06	0.03	99.41
67	5.D	Wastewater treatment and discharge	CH4	61.10	50.86	0.07	0.03	99.44
68	2.D.2	Paraffin wax use	CO2	5.09	16.88	0.02	0.03	99.48
69	3.A.4	Enteric Fermentation - Other livestock	CH4	37.87	49.61	0.07	0.03	99.51
70	4.F.2	LULUCF - Land Converted to Other Land	CO2	0.55	11.06	0.02	0.03	99.53
71	2.G.2	Other Product Manufacture and Use - SF6 and PFCs from other product use	SF6	12.90	2.92	0.00	0.03	99.56
72	1.A.1	Energy Industries - Peat Fuel	N2O	52.07	43.93	0.06	0.02	99.59
73	1.A.4.b	Residential - Peat Fuel	N2O	12.66	3.81	0.01	0.02	99.61
74	2.G.3.a	Other Product Manufacture and Use - Other (Anaesthesia in medical applications)	CO2	31.34	41.06	0.06	0.02	99.63
75	1.A.4.a	Commercial/Institutional - Biomass	CH4	0.00	7.90	0.01	0.02	99.66
76	3.A.3	Enteric Fermentation - Swine	CH4	41.37	49.94	0.07	0.02	99.68
77	3.B.2.2	Manure Management - Sheep	N2O	18.33	11.50	0.02	0.02	99.70
78	4.D.2	LULUCF - Land converted to wetlands	CO2	0.00	7.12	0.01	0.02	99.72
79	1.A.4.b	Residential - Solid Fuels	N2O	11.71	5.11	0.01	0.02	99.73
80	4.C.1	LULUCF - Grassland Remaining Grassland	N2O	0.90	6.66	0.01	0.02	99.75
81	1.A.4.b	Residential - Liquid Fuels	CH4	3.71	9.40	0.01	0.02	99.76
82	1.A.1	Energy Industries - Biomass	N2O	0.00	5.45	0.01	0.01	99.78

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
83	1.A.4.b	Residential - Biomass	CH4	14.08	9.06	0.01	0.01	99.79
84	2.A.4.d	Other process uses of carbonates - Bricks (Other)	CO2	5.07	0.03	0.00	0.01	99.81
85	3.B.2.5	Manure Management - Indirect N2O emissions	N2O	216.24	225.44	0.32	0.01	99.82
86	1.A.2	Manufacturing Industries & Construction - Biomass	N2O	3.04	7.49	0.01	0.01	99.83
87	4.A.1	LULUCF - Forest land Remaining Forest Land	CH4	11.06	7.08	0.01	0.01	99.84
88	1.A.4.b	Residential - Liquid Fuels	N2O	2.48	6.62	0.01	0.01	99.85
89	4.C.2	LULUCF - Land Converted to Grassland	N2O	0.08	3.74	0.01	0.01	99.86
90	1.A.4.a	Commercial/Institutional - Liquid Fuels	CH4	6.58	3.35	0.00	0.01	99.87
91	1.A.2	Manufacturing Industries & Construction - Biomass	CH4	1.91	4.71	0.01	0.01	99.88
92	1.A.2	Manufacturing Industries & Construction - Solid Fuels	N2O	4.12	1.54	0.00	0.01	99.89
93	4.A.2	LULUCF - Land converted to Forest Land	CH4	0.00	2.64	0.00	0.01	99.89
94	1.A.4.b	Residential - Gaseous Fuels	CH4	0.61	3.17	0.00	0.01	99.90
95	2.G.1	Other Product Manufacture and Use - Electrical equipment	SF6	20.52	18.60	0.03	0.01	99.91
96	4.D.1	LULUCF - Wetlands Remaining wetlands	N2O	34.26	32.75	0.05	0.01	99.91
97	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CO2	0.00	2.16	0.00	0.01	99.92
98	1.A.3.c	Rail Transport - Liquid Fuels	N2O	15.49	13.68	0.02	0.01	99.92
99	1.A.4.a	Commercial/Institutional - Liquid Fuels	N2O	4.68	2.85	0.00	0.01	99.93
100	1.A.1	Energy Industries - Solid Fuels	N2O	7.74	6.05	0.01	0.01	99.93
101	1.A.4.a	Commercial/Institutional - Gaseous Fuels	CH4	0.51	2.22	0.00	0.00	99.94
102	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	N2O	4.83	3.36	0.00	0.00	99.94
103	1.A.2	Manufacturing Industries & Construction - Solid Fuels	CH4	2.30	0.86	0.00	0.00	99.95
104	3.B.2.3	Manure Management - Swine	N2O	10.08	11.79	0.02	0.00	99.95
105	1.A.1	Energy Industries - Biomass	CH4	0.00	1.40	0.00	0.00	99.96
106	1.A.1	Energy Industries - Gaseous Fuels	CH4	3.43	2.11	0.00	0.00	99.96
107	1.A.1	Energy Industries - Liquid Fuels	N2O	1.47	0.24	0.00	0.00	99.96
108	1.A.4.a	Commercial/Institutional - Biomass	N2O	0.00	1.26	0.00	0.00	99.97
109	1.A.1	Energy Industries - Other Fuels	N2O	0.00	1.14	0.00	0.00	99.97
110	3.B.2.4	Manure Management - Other livestock	N2O	10.31	11.63	0.02	0.00	99.97
111	1.A.3.e	Other Transport - Gaseous Fuels	N2O	0.67	1.58	0.00	0.00	99.97
112	1.A.4.b	Residential - Biomass	N2O	2.23	1.43	0.00	0.00	99.98
113	5.C.1 & 2	Incineration and open burning of waste - Incineration	CH4	0.83	0.07	0.00	0.00	99.98
114	1.A.3.d	Navigation - Liquid Fuels	N2O	0.67	1.43	0.00	0.00	99.98
115	1.A.1	Energy Industries - Other Fuels	CH4	0.00	0.72	0.00	0.00	99.98
116	1.A.2	Manufacturing Industries & Construction - Liquid Fuels	CH4	2.06	1.47	0.00	0.00	99.98
117	5.C.1 & 2	Incineration and open burning of waste - Incineration	N2O	1.04	0.45	0.00	0.00	99.99
118	1.A.4.b	Residential - Gaseous Fuels	N2O	0.15	0.76	0.00	0.00	99.99
119	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	N2O	0.47	1.09	0.00	0.00	99.99
120	1.A.2	Manufacturing Industries & Construction - Gaseous Fuels	CH4	0.40	0.91	0.00	0.00	99.99
121	1.A.4.a	Commercial/Institutional - Gaseous Fuels	N2O	0.12	0.53	0.00	0.00	99.99
122	1.A.3.a	Domestic Aviation - Liquid Fuels	N2O	0.55	0.16	0.00	0.00	99.99
123	1.A.1	Energy Industries - Peat Fuel	CH4	1.90	1.62	0.00	0.00	99.99
124	4.A.1	LULUCF - Forest land Remaining Forest Land	N2O	0.77	0.49	0.00	0.00	99.99

Ranking	IPCC Sub-category	Emission Source / Activity	Direct GHG	1990 Emissions (kt CO ₂ eq)	2013 Emissions (kt CO ₂ eq)	2013 Level assessment (%)	2013 Trend assessment (%)	Cumulative Total (%)
125	1.A.1	Energy Industries - Liquid Fuels	CH4	0.39	0.11	0.00	0.00	99.99
126	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	N2O	0.00	0.26	0.00	0.00	100.00
127	1.A.1	Energy Industries - Solid Fuels	CH4	0.91	0.71	0.00	0.00	100.00
128	1.A.3.d	Navigation - Liquid Fuels	CH4	0.20	0.42	0.00	0.00	100.00
129	1.A.4.c	Agriculture - Liquid Fuels	CH4	1.07	0.88	0.00	0.00	100.00
130	1.A.3.e	Other Transport - Gaseous Fuels	CH4	0.14	0.33	0.00	0.00	100.00
131	4.A.2	LULUCF - Land converted to Forest Land	N2O	0.00	0.18	0.00	0.00	100.00
132	4.D.1	LULUCF - Wetlands Remaining wetlands	CH4	76.80	78.36	0.11	0.00	100.00
133	1.B.2.a	Fugitive Emissions - Oil Refining	CH4	0.21	0.33	0.00	0.00	100.00
134	1.A.2	Manufacturing Industries & Construction - Non-Renewable waste	CH4	0.00	0.11	0.00	0.00	100.00
135	4.C.1	LULUCF - Grassland Remaining Grassland	CH4	43.06	44.09	0.06	0.00	100.00
136	2.A.4.d	Other process uses of carbonates - Limestone use (Other)	CO2	0.15	0.21	0.00	0.00	100.00
137	2.A.4.b	Other process uses of carbonates - Soda Ash (Other use)	CO2	0.10	0.06	0.00	0.00	100.00
138	1.A.3.c	Rail Transport - Liquid Fuels	CH4	0.19	0.17	0.00	0.00	100.00
139	1.A.3.a	Domestic Aviation - Liquid Fuels	CH4	0.02	0.01	0.00	0.00	100.00
140	1.A.2	Manufacturing Industries & Construction - Peat Fuel	N2O	0.00	0.01	0.00	0.00	100.00
141	1.A.2	Manufacturing Industries & Construction - Peat Fuel	CH4	0.00	0.00	0.00	0.00	100.00
142	4.B.1	LULUCF - Cropland Remaining Cropland	CH4	0.07	0.08	0.00	0.00	100.00
143	4.B.1	LULUCF - Cropland Remaining Cropland	N2O	0.02	0.02	0.00	0.00	100.00
144	1.A.4.a	Commercial/Institutional - Solid Fuels	CO2	2.56	0.00	0.00	0.00	100.00
145	1.A.4.a	Commercial/Institutional - Peat Fuel	CO2	135.73	0.00	0.00	0.00	100.00
146	1.A.4.a	Commercial/Institutional - Solid Fuels	CH4	0.01	0.00	0.00	0.00	100.00
147	1.A.4.a	Commercial/Institutional - Peat Fuel	CH4	0.33	0.00	0.00	0.00	100.00
148	1.A.4.a	Commercial/Institutional - Solid Fuels	N2O	0.01	0.00	0.00	0.00	100.00
149	1.A.4.a	Commercial/Institutional - Peat Fuel	N2O	0.56	0.00	0.00	0.00	100.00
150	2.A.3	Glass Production	CO2	13.33	0.00	0.00	0.00	100.00
151	2.B.1	Chemical Industry - Ammonia Production	CO2	990.23	0.00	0.00	0.00	100.00
152	2.B.2	Chemical Industry - Nitric Acid Production	N2O	995.32	0.00	0.00	0.00	100.00

1.E Information on the level of disaggregation

The disaggregation approach found in Chapter 4.2 of the 2006 IPCC Guidelines has been followed.

1.F Description of methodology used for identifying key categories

Approach 1 in the 2006 IPCC Guidelines has been followed; key categories have been identified using a pre-determined cumulative emissions threshold. Key categories are those that, when summed together in descending order of magnitude, add up to 95 per cent of the total level.

Annex 2

Assessment of Uncertainty

2.A 2013 Uncertainty Assessment excluding LULUCF

2.B 2013 Uncertainty Assessment including LULUCF

2.C Description of methodology used for identifying uncertainties

Table 2.A.i 2013 Uncertainty Assessment excluding LULUCF, CO₂

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combine d Uncertain ty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared	
1	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO2	1880.66	4373.09	1.00	2.50	2.69	0.04	0.00	0.11	0.11	0.15	0.02	
2	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO2	1254.90	401.78	1.00	2.50	2.69	0.00	0.00	0.01	-0.04	0.04	0.00	
3	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO2	0.00	86.50	1.00	5.00	5.10	0.00	0.00	0.00	0.01	0.01	0.00	
4	1.A.1 Fuel combustion - Energy Industries - Peat	CO2	3164.78	2579.98	1.00	5.00	5.10	0.05	0.00	0.06	-0.06	0.09	0.01	
5	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO2	4844.66	3797.89	1.00	5.00	5.10	0.11	0.01	0.09	-0.11	0.14	0.02	
6	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO2	873.02	2039.08	7.00	3.00	7.62	0.07	0.00	0.36	0.06	0.36	0.13	
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO2	2198.38	1709.71	10.00	2.50	10.31	0.09	0.01	0.43	-0.03	0.43	0.18	
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO2	0.00	132.53	1.00	5.00	5.10	0.00	0.00	0.00	0.01	0.01	0.00	
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CO2	0.00	2.16	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00	
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO2	871.24	325.65	2.00	5.00	5.39	0.00	0.00	0.02	-0.05	0.05	0.00	
11	1.A.3.a Domestic Aviation	CO2	51.13	10.01	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00	
12	1.A.3.b Road Transportation	CO2	4690.42	10483.36	1.25	3.00	3.25	0.34	0.11	0.33	0.30	0.44	0.19	
13	1.A.3.c Railways	CO2	133.19	117.65	1.00	1.00	1.41	0.00	0.00	0.00	0.00	0.00	0.00	
14	1.A.3.d Domestic Navigation - Liquid Fuels	CO2	84.90	175.55	1.00	2.00	2.24	0.00	0.00	0.00	0.00	0.01	0.00	
15	1.A.3.e Other Transportation	CO2	62.04	148.53	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.01	0.00	
16	1.A.4 Other Sectors - Gaseous Fuels	CO2	493.22	2416.22	2.50	2.50	3.54	0.02	0.00	0.15	0.08	0.17	0.03	
17	1.A.4 Other Sectors - Liquid Fuels	CO2	3792.65	4311.36	10.00	5.00	11.18	0.67	0.45	1.07	0.03	1.07	1.16	
18	1.A.4 Other Sectors - Peat	CO2	3259.11	929.40	10.00	20.00	22.36	0.12	0.02	0.23	-0.86	0.89	0.80	
19	1.A.4 Other Sectors - Solid Fuels	CO2	2485.97	1098.84	5.00	10.00	11.18	0.04	0.00	0.14	-0.26	0.29	0.09	
20	2.A.1 Cement Production	CO2	884.00	1111.75	1.50	1.50	2.12	0.00	0.00	0.04	0.01	0.04	0.00	
21	2.A.2 Lime Production	CO2	214.08	189.64	5.00	5.00	7.07	0.00	0.00	0.02	0.00	0.02	0.00	
22	2.A.3 Glass Production	CO2	13.33	0.00	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00	
23	2.A.4 Other Process Uses of Carbonates	CO2	5.32	0.31	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00	
24	2.B.1 Ammonia Production	CO2	990.23	0.00	1.00	5.00	5.10	0.00	0.00	0.00	-0.09	0.09	0.01	
25	2.D Non-energy Products from Fuels and Solvent Use	CO2	84.42	102.06	30.00	5.00	30.41	0.00	0.00	0.08	0.00	0.08	0.01	
26	3.G Liming	CO2	355.04	515.69	5.00	50.00	50.25	0.19	0.04	0.06	0.13	0.15	0.02	
27	3.H Urea Application	CO2	44.47	21.66	5.00	50.00	50.25	0.00	0.00	0.00	-0.02	0.02	0.00	
28	5.C Incineration and Open Burning of Waste	CO2	90.61	43.05	10.00	5.00	11.18	0.00	0.00	0.01	0.00	0.01	0.00	
Total CO2			32821.78	37123.45				1.76					2.67	
Level								1.32	Trend					1.63

Table 2.A.ii 2013 Uncertainty Assessment excluding LULUCF, CH₄

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	CH ₄	0.00	1.40	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CH ₄	3.43	2.11	1.00	70.00	70.01	0.00	0.00	0.00	0.00	0.00	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CH ₄	0.39	0.11	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CH ₄	0.00	0.72	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	CH ₄	1.86	1.56	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CH ₄	0.91	0.71	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	CH ₄	1.91	4.71	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH ₄	0.40	0.91	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH ₄	2.06	1.47	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH ₄	0.00	0.11	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CH ₄	0.00	0.00	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH ₄	2.30	0.86	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	CH ₄	0.02	0.01	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	CH ₄	47.47	16.47	1.25	71.00	71.01	0.00	0.00	0.00	-0.04	0.04	0.00
15	1.A.3.c Railways	CH ₄	0.19	0.17	1.00	60.00	60.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	CH ₄	0.20	0.42	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	CH ₄	0.14	0.33	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	CH ₄	14.08	16.96	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	CH ₄	1.12	5.40	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	CH ₄	11.36	13.62	10.00	66.00	66.75	0.00	0.00	0.00	0.00	0.00	0.00
21	1.A.4 Other Sectors - Peat	CH ₄	227.98	68.42	10.00	50.00	50.99	0.00	0.00	0.02	-0.15	0.15	0.02
22	1.A.4 Other Sectors - Solid Fuels	CH ₄	196.51	85.73	5.00	50.00	50.25	0.01	0.00	0.01	-0.10	0.10	0.01
23	1.B.1 Fugitive emissions from Solid Fuels	CH ₄	55.56	20.15	10.00	50.00	50.99	0.00	0.00	0.01	-0.03	0.03	0.00
24	1.B.2.a Fugitive Emissions from Fuels - Oil and Natural Gas - Oil	CH ₄	0.21	0.33	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
25	1.B.2.b Fugitive Emissions from Fuels - Oil and Natural Gas - Natural Gas	CH ₄	156.05	23.26	10.00	50.00	50.99	0.00	0.00	0.01	-0.12	0.12	0.01
26	3.A Enteric Fermentation	CH ₄	11356.97	10553.71	1.00	17.00	17.03	9.34	87.15	0.26	-0.36	0.45	0.20
27	3.B Manure Management	CH ₄	1342.29	1242.33	1.00	19.00	19.03	0.16	0.03	0.03	-0.05	0.06	0.00
28	5.A Solid Waste Disposal	CH ₄	1396.49	1119.29	34.64	34.64	48.99	0.87	0.76	0.97	-0.20	0.99	0.97
29	5.C Incineration and Open Burning of Waste	CH ₄	0.83	0.07	10.00	30.00	31.62	0.00	0.00	0.00	0.00	0.00	0.00
30	5.D Wastewater Treatment and Discharge	CH ₄	61.10	50.86	10.00	30.00	31.62	0.00	0.00	0.01	-0.01	0.01	0.00
Total CH₄			14881.83	13232.21				10.38					1.23
						Level uncertainty, CH₄			3.22	Trend uncertainty, CH₄			1.11
Combined CO₂ and CH₄			47703.60	50355.66				12.13					3.90
						Level uncertainty, CO₂ and CH₄			3.48	Trend uncertainty, CO₂ & CH₄			1.97

Table 2.A.iii 2013 Uncertainty Assessment excluding LULUCF, N₂O

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	N ₂ O	0.00	5.45	1.00	63.00	63.01	0.00	0.00	0.00	0.01	0.01	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	N ₂ O	10.21	67.46	1.00	50.00	50.01	0.00	0.00	0.00	0.05	0.05	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	N ₂ O	1.47	0.24	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	N ₂ O	0.00	1.14	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	N ₂ O	52.07	43.93	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.01	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	N ₂ O	7.74	6.05	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	N ₂ O	3.04	7.49	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N ₂ O	0.47	1.09	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N ₂ O	4.83	3.36	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N ₂ O	0.00	0.26	1.00	20.00	20.02	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	N ₂ O	0.00	0.01	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N ₂ O	4.12	1.54	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	N ₂ O	0.55	0.16	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	N ₂ O	48.09	98.33	1.25	68.00	68.01	0.01	0.00	0.00	0.06	0.06	0.00
15	1.A.3.c Railways	N ₂ O	15.49	13.68	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	N ₂ O	0.67	1.43	1.00	90.00	90.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	N ₂ O	0.67	1.58	1.00	25.00	25.02	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	N ₂ O	2.23	2.68	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	N ₂ O	0.27	1.29	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	N ₂ O	76.42	66.32	10.00	50.00	50.99	0.00	0.00	0.02	-0.01	0.02	0.00
21	1.A.4 Other Sectors - Peat	N ₂ O	13.22	3.81	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
22	1.A.4 Other Sectors - Solid Fuels	N ₂ O	11.72	5.11	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
23	2.B.2 Nitric Acid Production	N ₂ O	995.32	0.00	1.00	10.00	10.05	0.00	0.00	0.00	-0.18	0.18	0.03
24	2.G Other Product Manufacture and Use	N ₂ O	31.34	41.06	5.00	5.00	7.07	0.00	0.00	0.01	0.00	0.01	0.00
25	3.B Manure Management	N ₂ O	496.01	517.94	11.22	50.00	51.24	0.20	0.04	0.14	0.00	0.14	0.02
26	3.D.1 Direct N ₂ O Emissions From Managed Soils	N ₂ O	6581.38	5633.17	7.80	100.00	100.30	92.27	8514.43	1.09	-2.09	2.36	5.57
27	3.D.2 Indirect N ₂ O Emissions From Managed Soils	N ₂ O	559.22	480.10	11.18	50.00	51.23	0.17	0.03	0.13	-0.09	0.16	0.03
28	5.C Incineration and Open Burning of Waste	N ₂ O	1.04	12.30	10.00	10.00	14.14	0.00	0.00	0.00	0.00	0.00	0.00
29	5.D Wastewater Treatment and Discharge	N ₂ O	95.64	118.70	10.00	10.00	14.14	0.00	0.00	0.03	0.00	0.03	0.00
Total N₂O			9013.24	7135.66				92.67					5.66
			Level uncertainty, N₂O					9.63	Trend uncertainty, N₂O				
Combined CO₂, CH₄ and N₂O			56716.85	57491.33				104.81					9.56
			Level uncertainty, CO₂, CH₄ & N₂O					10.24	Trend uncertainty, CO₂, CH₄ & N₂O				

Table 2.A.iv 2013 Uncertainty Assessment excluding LULUCF – aggregate F-gases and Total for all gases

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO2eq)	Emissions in 2013 (kt CO2eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	2.E Electronics Industry	Aggregate F-gases	1.17	1307.97	20.00	10.00	22.36	0.25	0.06	0.65	0.23	0.69	0.48
2	2.G Other Product Manufacture and Use	Aggregate F-gases	33.42	21.52	10.00	0.00	10.00	0.00	0.00	0.01	0.00	0.01	0.00
Total F-gases			34.59	1329.50				0.25	0.48				
Level uncertainty, F-gases								0.50	Trend uncertainty, F-gases				0.69
TOTAL for all gases			56751.44	58820.83				105.05	10.04				
Total level uncertainty for all GHGs								10.25	Total trend uncertainty for all GHGs				3.17

Table 2B.i 2013 Uncertainty Assessment including LULUCF CO₂

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO ₂	1880.66	4373.09	1.00	2.50	2.69	0.04	0.00	0.11	0.11	0.15	0.02
2	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO ₂	1254.90	401.78	1.00	2.50	2.69	0.00	0.00	0.01	-0.04	0.04	0.00
3	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO ₂	0.00	86.50	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Peat	CO ₂	3164.78	2579.98	1.00	5.00	5.10	0.04	0.00	0.06	-0.06	0.09	0.01
5	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO ₂	4844.66	3797.89	1.00	5.00	5.10	0.10	0.01	0.09	-0.11	0.14	0.02
6	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO ₂	873.02	2039.08	7.00	3.00	7.62	0.06	0.00	0.36	0.06	0.36	0.13
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO ₂	2198.38	1709.71	10.00	2.50	10.31	0.08	0.01	0.43	-0.03	0.43	0.18
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO ₂	0.00	132.53	1.00	5.00	5.10	0.00	0.00	0.00	0.01	0.01	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CO ₂	0.00	2.16	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO ₂	871.24	325.65	2.00	5.00	5.39	0.00	0.00	0.02	-0.05	0.05	0.00
11	1.A.3.a Domestic Aviation	CO ₂	51.13	10.01	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.3.b Road Transportation	CO ₂	4690.42	10483.36	1.25	3.00	3.25	0.30	0.09	0.33	0.30	0.44	0.19
13	1.A.3.c Railways	CO ₂	133.19	117.65	1.00	1.00	1.41	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.d Domestic Navigation - Liquid Fuels	CO ₂	84.90	175.55	1.00	2.00	2.24	0.00	0.00	0.00	0.00	0.01	0.00
15	1.A.3.e Other Transportation	CO ₂	62.04	148.53	1.00	2.50	2.69	0.00	0.00	0.00	0.00	0.01	0.00
16	1.A.4 Other Sectors - Gaseous Fuels	CO ₂	493.22	2416.22	2.50	2.50	3.54	0.02	0.00	0.15	0.08	0.17	0.03
17	1.A.4 Other Sectors - Liquid Fuels	CO ₂	3792.65	4311.36	10.00	5.00	11.18	0.59	0.35	1.07	0.03	1.07	1.16
18	1.A.4 Other Sectors - Peat	CO ₂	3259.11	929.40	10.00	20.00	22.36	0.11	0.01	0.23	-0.86	0.89	0.80
19	1.A.4 Other Sectors - Solid Fuels	CO ₂	2485.97	1098.84	5.00	10.00	11.18	0.04	0.00	0.14	-0.26	0.29	0.09
20	2.A.1 Cement Production	CO ₂	884.00	1111.75	1.50	1.50	2.12	0.00	0.00	0.04	0.01	0.04	0.00
21	2.A.2 Lime Production	CO ₂	214.08	189.64	5.00	5.00	7.07	0.00	0.00	0.02	0.00	0.02	0.00
22	2.A.3 Glass Production	CO ₂	13.33	0.00	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
23	2.A.4 Other Process Uses of Carbonates	CO ₂	5.32	0.31	5.00	2.50	5.59	0.00	0.00	0.00	0.00	0.00	0.00
24	2.B.1 Ammonia Production	CO ₂	990.23	0.00	1.00	5.00	5.10	0.00	0.00	0.00	-0.09	0.09	0.01
25	2.D Non-energy Products from Fuels and Solvent Use	CO ₂	84.42	102.06	30.00	5.00	30.41	0.00	0.00	0.08	0.00	0.08	0.01
26	3.G Liming	CO ₂	355.04	515.69	5.00	50.00	50.25	0.17	0.03	0.06	0.13	0.15	0.02
27	3.H Urea Application	CO ₂	44.47	21.66	5.00	50.00	50.25	0.00	0.00	0.00	-0.02	0.02	0.00
28	4.A.1 Forest Land Remaining Forest Land	CO ₂	-2719.66	74.00	51.00	114.00	124.89	0.15	0.02	0.09	5.32	5.32	28.26
29	4.B.1 Cropland Remaining Cropland	CO ₂	-1524.52	-364.38	20.59	69.15	72.15	-0.42	0.18	-0.17	1.35	1.36	1.85
30	4.C.1 Grassland Remaining Grassland	CO ₂	6067.33	4763.15	12.22	90.00	90.83	6.90	47.60	1.34	-2.12	2.51	6.29
31	4.C.2 Land Converted to Grassland	CO ₂	97.76	138.31	43.95	101.11	110.24	0.24	0.06	0.14	0.06	0.15	0.02
32	4.D.1 Wetlands Remaining Wetland	CO ₂	2291.19	2684.47	21.49	101.45	103.70	4.44	19.71	1.33	0.56	1.45	2.09
33	4.D.2 Land Converted to Wetlands	CO ₂	0.00	7.12	2.50	50.00	50.06	0.01	0.00	0.00	0.01	0.01	0.00
34	4.E.2 Land Converted to Settlements	CO ₂	73.88	54.96	39.97	81.83	91.07	0.08	0.01	0.05	-0.03	0.06	0.00
35	4.F.2 Land Converted to Other Land	CO ₂	0.55	11.06	51.93	75.00	91.23	0.02	0.00	0.01	0.01	0.02	0.00
36	4.G Harvested Wood Products	CO ₂	-410.59	-691.95	25.00	26.92	36.74	-0.41	0.16	-0.40	-0.12	0.42	0.17
37	5.C Incineration and Open Burning of Waste	CO ₂	90.61	43.05	10.00	5.00	11.18	0.00	0.00	0.01	0.00	0.01	0.00
Total CO₂			36724.97	40053.22				5.09				110.11	
						Level uncertainty, CO₂			2.26	Trend uncertainty, CO₂			10.49

Table 2B.ii 2013 Uncertainty Assessment including LULUCF, CH₄

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	CH ₄	0.00	1.40	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CH ₄	3.43	2.11	1.00	70.00	70.01	0.00	0.00	0.00	0.00	0.00	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CH ₄	0.39	0.11	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CH ₄	0.00	0.72	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	CH ₄	1.86	1.56	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CH ₄	0.91	0.71	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	CH ₄	1.91	4.71	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH ₄	0.40	0.91	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH ₄	2.06	1.47	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH ₄	0.00	0.11	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	CH ₄	0.00	0.00	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH ₄	2.30	0.86	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	CH ₄	0.02	0.01	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	CH ₄	47.47	16.47	1.25	71.00	71.01	0.00	0.00	0.00	-0.04	0.04	0.00
15	1.A.3.c Railways	CH ₄	0.19	0.17	1.00	60.00	60.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	CH ₄	0.20	0.42	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	CH ₄	0.14	0.33	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	CH ₄	14.08	16.96	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	CH ₄	1.12	5.40	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	CH ₄	11.36	13.62	10.00	66.00	66.75	0.00	0.00	0.00	0.00	0.00	0.00
21	1.A.4 Other Sectors - Peat	CH ₄	227.98	68.42	10.00	50.00	50.99	0.00	0.00	0.02	-0.15	0.15	0.02
22	1.A.4 Other Sectors - Solid Fuels	CH ₄	196.51	85.73	5.00	50.00	50.25	0.00	0.00	0.01	-0.10	0.10	0.01
23	1.B.1 Fugitive emissions from Solid Fuels	CH ₄	55.56	20.15	10.00	50.00	50.99	0.00	0.00	0.01	-0.03	0.03	0.00
24	1.B.2.a Fugitive Emissions from Fuels - Oil	CH ₄	0.21	0.33	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
25	1.B.2.b Fugitive Emissions from Fuels - Natural Gas	CH ₄	156.05	23.26	10.00	50.00	50.99	0.00	0.00	0.01	-0.12	0.12	0.01
26	3.A Enteric Fermentation	CH ₄	11356.97	10553.71	1.00	17.00	17.03	8.21	67.49	0.26	-0.36	0.45	0.20
27	3.B Manure Management	CH ₄	1342.29	1242.33	1.00	19.00	19.03	0.14	0.02	0.03	-0.05	0.06	0.00
28	4.A LULUCF - Forest Land	CH ₄	58.58	77.27	30.00	100.00	104.40	0.13	0.02	0.05	0.03	0.06	0.00
29	4.B.1 LULUCF - Cropland remaining Cropland	CH ₄	0.07	0.08	100.00	39.10	107.37	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	CH ₄	263.55	374.97	96.40	91.20	132.70	0.79	0.63	0.83	0.16	0.85	0.72
31	4.D.1 LULUCF - Wetlands remaining Wetlands	CH ₄	152.31	147.37	86.00	66.50	108.71	0.26	0.07	0.29	-0.01	0.29	0.09
32	5.A Solid Waste Disposal	CH ₄	1396.49	1119.29	34.64	34.64	48.99	0.76	0.58	0.97	-0.20	0.99	0.97
33	5.C Incineration and Open Burning of Waste	CH ₄	0.83	0.07	10.00	30.00	31.62	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	CH ₄	61.10	50.86	10.00	30.00	31.62	0.00	0.00	0.01	-0.01	0.01	0.00
Total CH₄			15356.35	13831.90				10.31				2.04	
						Level uncertainty, CH₄			3.21	Trend uncertainty, CH₄			1.43
Combined CO₂ and CH₄			52081.32	53885.12				15.40				112.15	
						Level uncertainty, CO₂ and CH₄			3.92	Trend uncertainty, CO₂ & CH₄			10.59

Table 2B.iii 2013 Uncertainty Assessment including LULUCF, N₂O

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	1.A.1 Fuel combustion - Energy Industries - Biomass	N ₂ O	0.00	5.45	1.00	63.00	63.01	0.00	0.00	0.00	0.01	0.01	0.00
2	1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	N ₂ O	10.21	67.46	1.00	50.00	50.01	0.00	0.00	0.00	0.05	0.05	0.00
3	1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	N ₂ O	1.47	0.24	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
4	1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	N ₂ O	0.00	1.14	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
5	1.A.1 Fuel combustion - Energy Industries - Peat	N ₂ O	52.07	43.93	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.01	0.00
6	1.A.1 Fuel combustion - Energy Industries - Solid Fuels	N ₂ O	7.74	6.05	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
7	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Biomass	N ₂ O	3.04	7.49	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
8	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N ₂ O	0.47	1.09	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
9	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N ₂ O	4.83	3.36	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
10	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N ₂ O	0.00	0.26	1.00	20.00	20.02	0.00	0.00	0.00	0.00	0.00	0.00
11	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Peat	N ₂ O	0.00	0.01	2.00	50.00	50.04	0.00	0.00	0.00	0.00	0.00	0.00
12	1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N ₂ O	4.12	1.54	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
13	1.A.3.a Domestic Aviation	N ₂ O	0.55	0.16	1.00	66.00	66.01	0.00	0.00	0.00	0.00	0.00	0.00
14	1.A.3.b Road Transportation	N ₂ O	48.09	98.33	1.25	68.00	68.01	0.01	0.00	0.00	0.06	0.06	0.00
15	1.A.3.c Railways	N ₂ O	15.49	13.68	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
16	1.A.3.d Domestic Navigation - Liquid Fuels	N ₂ O	0.67	1.43	1.00	90.00	90.01	0.00	0.00	0.00	0.00	0.00	0.00
17	1.A.3.e Other Transportation	N ₂ O	0.67	1.58	1.00	25.00	25.02	0.00	0.00	0.00	0.00	0.00	0.00
18	1.A.4 Other Sectors - Biomass	N ₂ O	2.23	2.68	10.00	50.00	50.99	0.00	0.00	0.00	0.00	0.00	0.00
19	1.A.4 Other Sectors - Gaseous Fuels	N ₂ O	0.27	1.29	2.50	50.00	50.06	0.00	0.00	0.00	0.00	0.00	0.00
20	1.A.4 Other Sectors - Liquid Fuels	N ₂ O	76.42	66.32	10.00	50.00	50.99	0.00	0.00	0.02	-0.01	0.02	0.00
21	1.A.4 Other Sectors - Peat	N ₂ O	13.22	3.81	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
22	1.A.4 Other Sectors - Solid Fuels	N ₂ O	11.72	5.11	5.00	50.00	50.25	0.00	0.00	0.00	-0.01	0.01	0.00
23	2.B.2 Nitric Acid Production	N ₂ O	995.32	0.00	1.00	10.00	10.05	0.00	0.00	0.00	-0.18	0.18	0.03
24	2.G Other Product Manufacture and Use	N ₂ O	31.34	41.06	5.00	5.00	7.07	0.00	0.00	0.01	0.00	0.01	0.00
25	3.B Manure Management	N ₂ O	496.01	517.94	11.22	50.00	51.24	0.18	0.03	0.14	0.00	0.14	0.02
26	3.D.1 Direct N ₂ O Emissions From Managed Soils	N ₂ O	6581.38	5633.17	7.80	100.00	100.30	81.20	6592.83	1.09	-2.09	2.36	5.57
27	3.D.2 Indirect N ₂ O Emissions From Managed Soils	N ₂ O	559.22	480.10	11.18	50.00	51.23	0.15	0.02	0.13	-0.09	0.16	0.03
28	4.A LULUCF - Forest Land	N ₂ O	92.86	171.66	30.00	100.00	104.40	0.29	0.08	0.12	0.13	0.17	0.03
29	4.B.1 LULUCF - Cropland remaining Cropland	N ₂ O	0.02	0.02	100.00	100.00	141.42	0.00	0.00	0.00	0.00	0.00	0.00
30	4.C.1 LULUCF - Grassland Remaining Grassland	N ₂ O	0.90	6.66	91.02	100.00	135.22	0.01	0.00	0.01	0.01	0.02	0.00
31	4.C.2 LULUCF - Land converted to Grassland	N ₂ O	0.08	3.74	90.80	100.00	135.07	0.01	0.00	0.01	0.01	0.01	0.00
32	4.D.1 LULUCF - Wetlands remaining Wetlands	N ₂ O	47.16	43.71	86.00	100.00	131.89	0.09	0.01	0.09	-0.01	0.09	0.01
33	5.C Incineration and Open Burning of Waste	N ₂ O	1.04	12.30	10.00	10.00	14.14	0.00	0.00	0.00	0.00	0.00	0.00
34	5.D Wastewater Treatment and Discharge	N ₂ O	95.64	118.70	10.00	10.00	14.14	0.00	0.00	0.03	0.00	0.03	0.00
Total N₂O			9173.01	7490.36				82.13				5.71	
						Level uncertainty, N₂O			9.06	Trend uncertainty, N₂O			2.39
Combined CO₂, CH₄ and N₂O			61254.32	61375.48				97.52				117.86	
						Level uncertainty, CO₂, CH₄ & N₂O			9.88	Trend uncertainty, CO₂, CH₄ & N₂O			10.86

Table 2B.iv 2013 Uncertainty Assessment including LULUCF – aggregate F-gases and Total for all gases

	KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Emissions in 1990 (kt CO ₂ eq)	Emissions in 2013 (kt CO ₂ eq)	Activity Data (AD) Uncertainty (%)	Emission Factor (EF) Uncertainty (%)	Combined Uncertainty (%)	Contribution to Variance by Category in Year 2013	Combined Emissions Uncertainty Squared	Uncertainty in Trend in Total Emissions due to AD (%)	Uncertainty in Trend in Total Emissions due to EF (%)	Combined Uncertainty in Trend in Total Emissions (%)	Combined Trend Uncertainty Squared
1	2.E Electronics Industry	Aggregate F-gases	1.17	1307.97	20.00	10.00	22.36	0.22	0.05	0.65	0.23	0.69	0.48
2	2.G Other Product Manufacture and Use	Aggregate F-gases	33.42	21.52	10.00	0.00	10.00	0.00	0.00	0.01	0.00	0.01	0.00
Total F-gases			34.59	1329.50				0.22					0.48
Level uncertainty, F-gases								0.47	Trend uncertainty, F-gases				0.69
TOTAL for all gases			61288.92	62704.98				97.74					118.34
Total level uncertainty for all GHGs								9.89	Total trend uncertainty for all GHGs				10.88

2.C Description of methodology used for estimating uncertainties

Approach 1 in 2006 IPCC Guidelines has been followed. Uncertainties for each category and gas have been estimated using equations 3.1 and 3.2 in Volume 1 of the 2006 IPCC Guidelines. A combination of these uncertainties by category has been used to estimate overall uncertainty for 2013 and the uncertainty in the trend.

$$U_{total} = \sqrt{(U_1^2 + U_2^2 + \dots + U_n^2)} \quad \text{Equation 3.1, Vol 1, 2006 IPCC Guidelines}$$

Where:

U_{total} = the percentage uncertainty in the product of the quantities (half the 95 per cent confidence interval divided by the total and expressed as a percentage);

U_i = the percentage uncertainties associated with each of the quantities.

$$U_{total} = \frac{\sqrt{((U_1 \times x_1) + (U_2 \times x_2))^2 + \dots + (U_n \times x_n)^2}}{|x_1 + x_2 + \dots + x_n|} \quad \text{Equation 3.2, Vol 1, 2006 IPCC Guidelines}$$

Where:

U_{total} = the percentage uncertainty in the sum of the quantities (half the 95 percent confidence interval divided by the total (i.e., mean) and expressed as a percentage). This term ‘uncertainty’ is thus based upon the 95 percent confidence interval;

x_i and U_i = the uncertain quantities and the percentage uncertainties associated with them, respectively.

Annex 3.1.A

Energy - Combustion (IPCC Sector 1.A)

3.1.1 – 3.1.2 Calculation Sheets for Energy 2013

3.1.3 – 3.1.5 Comparison of Reference and Sectoral Approach

3.1.6 – 3.1.8 Time-Series of Implied Emission Factors (IEFs) in Categories 1.A.1 and 1.A.2

Table 3.1.1 Calculation Sheet for Emissions from Fuel Combustion 2013 (continued on following pages)

	Sectoral Disaggregation of Fuel Combustion from National Energy Balance			Emission Factors			Emissions		
	Sector/Fuel	kTOE	TJ	CO ₂ kg/TJ	CH ₄ kg/TJ	N ₂ O kg/TJ	CO ₂ kt	CH ₄ Mg	N ₂ O Mg
	1A1a Public Electricity								
1	Coal	970.32	40625.48	93486	0.7	0.5	3797.89	28.44	20.31
2	Peat	497.53	20830.40	117991	3.0	7.0	2457.81	62.49	145.81
3	Fuel Oil and Gas Oil	37.78	1581.98	78537	0.8	0.3	124.24	1.28	0.49
4	Natural Gas	1829.95	76616.17	56860	1.1	3.0	4356.37	82.20	226.13
5	Biomass (LFG, Wood & MSW biomass)	109.75	4594.87	100901	12.1	4.0	463.63	55.83	18.28
6	MSW (non-renewable, fossil)	22.86	956.98	90393	30.0	4.0	86.50	28.71	3.83
	Public Electricity Total	3468.18	145205.88				10822.82	258.95	414.86
	1A1b Refinery Fuel								
7	Refinery Gas	69.09	2892.53	95129	1.0	0.1	275.16	2.89	0.29
8	Natural Gas	56.68	2373.04	7045	1.0	0.1	16.72	2.37	0.24
9	LPG	0.61	25.62	2	1.0	0.1	0.000043	0.03	0.00
10	Gasoil/Diesel/DERV	0.14	5.90	402205	3.0	0.6	2.37	0.02	0.00
	Refinery Total	126.52	5297.09				294.26	5.31	0.53
	1A1c Manufacture of Briquettes								
11	Peat	25.74	1077.52	113381	2.0	1.5	122.17	2.16	1.62
	1A2a-1A2g Industry Fuel								
12	Bituminous Coals	82.22	3442.36	94600	10.0	1.5	325.65	34.42	5.16
13	Briquettes	0.52	21.87	98860	2.0	1.5	2.16	0.04	0.03
14	Kerosene	78.44	3284.28	71400	3.0	0.6	234.50	9.85	1.97
15	Fuel Oil	137.81	5769.81	76000	3.0	0.6	438.51	17.31	3.46
16	LPG	116.05	4858.72	63700	1.0	0.1	309.50	4.86	0.49
17	Gasoil/Diesel/DERV	124.84	5226.69	73300	3.0	0.6	383.12	15.68	3.14
18	Pet Coke	88.43	3702.53	92934	3.0	0.6	344.09	11.11	2.22
19	Naphta	-	-	75330	3.0	0.6	-	-	-
20	Natural Gas	870.00	36425.04	55980	1.0	0.1	2039.08	36.43	3.64
21	Biomass (solid)	149.92	6276.79	112000	30.0	4.0	703.00	188.30	25.11
22	Biomass (gas)	2.64	110.74	54600	1.0	0.1	6.05	0.11	0.01
23	Non Renewable wastes (fossil)	34.89	1460.67	90732	3.0	0.6	132.53	4.38	0.88
	Industry Total	1685.76	70579.51				4209.13	322.50	46.11

Table 3.1.1 Calculation Sheet for Emissions from Fuel Combustion 2013 (continued from previous page)

	Sectoral Disaggregation of Fuel Combustion from National Energy Balance			Emission Factors			Emissions		
	Sector/Fuel	kTOE	TJ	CO ₂ kg/TJ	CH ₄ kg/TJ	N ₂ O kg/TJ	CO ₂ kt	CH ₄ Mg	N ₂ O Mg
	1A3a Aviation								
24	Domestic Aviation Kerosene & Gasoline	3.37	140.93	71015	2.1	3.9	10.01	0.30	0.55
	1A3b Road Transport Fuel								
25	Gasoline	1197.17	50123.01	69960	11.2	1.4	3506.61	562.43	70.99
26	Gasoil/Diesel/DERV	2272.22	95133.40	73300	1.0	2.7	6973.28	95.80	258.76
27	LPG	1.30	54.57	63700	12.8	3.8	3.48	0.70	0.21
28	Liquid Biofuels	102.24	4280.77	70253	69.0	41.9	300.74	16.11	9.78
	Road Transport Total	3572.94	149591.76				10483.36	658.93	329.96
	1A3c-1A3e Other Transport Fuel								
29	Railway Diesel	38.34	1605.04	73300	4.2	28.6	117.65	6.66	45.90
30	Navigation Fuel Oil	-	-	76000	7.0	2.0	-	-	-
31	Navigation Gasoil	57.20	2394.90	73300	7.0	2.0	175.55	16.76	4.79
32	Gas Distribution Use (Natural Gas)	63.37	2653.34	55980	5.0	2.0	148.53	13.27	5.31
33	Railway Biofuel	-	-	70800	4.2	28.6	-	-	-
	Other Transport Total	158.91	6653.28				441.73	36.69	56.00
	1A4a Commercial/Institutional Fuel								
34	Bituminous Coal	-	-	94600	10.0	1.5	-	-	-
35	Anthracite + Manufactured Ovoids	-	-	98300	10.0	1.5	-	-	-
36	Lignite	-	-	101000	10.0	1.5	-	-	-
37	Briquettes	-	-	98860	10.0	1.4	-	-	-
38	Fuel Oil	9.85	412.36	76000	10.0	0.6	31.34	4.12	0.25
39	LPG	8.65	362.04	63700	5.0	0.1	23.06	1.81	0.04
40	Gasoil / Diesel/ DERV	288.92	12096.41	73300	10.0	0.6	886.67	120.96	7.26
41	Gasoil (Fishing)	24.29	1016.84	73300	7.0	2.0	74.53	7.12	2.03
42	Pet Coke	0.0002	0.01	92934	10.0	0.6	0.0007	0.0001	0.000005
43	Natural Gas	424.69	17781.04	55980	5.0	0.1	995.39	88.91	1.78
44	Biomass	25.09	1050.28	112000	300.0	4.0	117.63	315.08	4.20
45	Biogas	4.42	185.05	54600	5.0	0.1	10.10	0.93	0.02
	Commercial/Institutional Total	785.90	32904.02				2010.99	538.93	15.57

Table 3.1.1 Calculation Sheet for Emissions from Fuel Combustion 2013 (continued from previous page)

	Sectoral Disaggregation of Fuel Combustion from National Energy Balance			Emission Factors			Emissions		
				CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
	Sector/Fuel	kTOE	TJ	kg/TJ	kg/TJ	kg/TJ	kt	Mg	Mg
	1A4b Residential Fuel								
46	Bituminous Coal	172.99	7242.79	94600	300.0	1.5	685.17	2172.84	10.86
47	Anthracite + Manufactured Ovoids	82.74	3463.99	98300	300.0	1.5	340.51	1039.20	5.20
48	Lignite	17.30	724.34	101000	300.0	1.5	73.16	217.30	1.09
49	Sod Peat	127.70	5346.71	104000	300.0	1.4	556.06	1604.01	7.49
50	Briquettes	90.20	3776.48	98860	300.0	1.4	373.34	1132.94	5.29
51	Kerosene	705.99	29558.53	71400	10.0	0.6	2110.48	295.59	17.74
52	LPG	40.04	1676.36	63700	5.0	0.1	106.78	8.38	0.17
53	Gasoil / Diesel/ DERV	160.97	6739.68	73300	10.0	0.6	494.02	67.40	4.04
54	Petroleum Coke	10.94	458.01	92934	10.0	0.6	42.56	4.58	0.27
55	Natural Gas	606.22	25381.01	55980	5.0	0.1	1420.84	126.91	2.54
56	Biomass	29.06	1216.59	112000	297.9	3.9	136.26	362.37	4.79
	Residential Total	2044.15	85584.50				6202.92	7031.51	59.47
	1A4c Agriculture Fuel								
57	Gasoil	176.58	7393.02	73300	4.7	25.8	541.91	35.01	190.74
58	Biomass	-	-	110000	300.0	4.0	-	-	-
	Agriculture Total	176.58	7393.02				541.91	35.01	190.74
	Total Energy	12048.04	504427.51				35139.30	8890.28	1115.41

Table 3.1.2 Emissions from Fuel Combustion Allocated by IPCC Source Category 2013

	GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA Consumption (TJ)	CO ₂ (t/TJ)	IMPLIED EMISSION FACTORS CH ₄ (kg/TJ)	N ₂ O (kg/TJ)	EMISSIONS CO ₂ (kt)	CH ₄	N ₂ O
A	1.A.1. Energy Industries	151,580.4933				8,659.2716	0.2664	0.4170
B	Solid Fuels	40,625.4830	93.4855	0.7000	0.5000	3,797.8944	0.0284	0.0203
C	Liquid Fuels	4,506.0315	89.1653	0.9366	0.1750	401.7818	0.0042	0.0008
D	Gaseous Fuels	78,989.2074	55.3631	1.0706	2.8658	4,373.0910	0.0846	0.2264
E	Peat Fuels	21,907.9150	117.7646	2.9508	6.7295	2,579.9779	0.0646	0.1474
F	Biomass	4,594.8741	100.9013	12.1500	3.9788	463.6289	0.0558	0.0183
F	Other Fuels	956.9823	90.3929	30.0000	4.0000	86.5044	0.0287	0.0038
G	1.A.2 Manufacturing Industries and Construction	70,579.5127				4,206.9688	0.3225	0.0461
H	Solid Fuels	3,442.3560	94.6000	10.0000	1.5000	325.6469	0.0344	0.0052
I	Liquid Fuels	22,842.0414	74.8493	2.5746	0.4936	1,709.7103	0.0588	0.0113
J	Gaseous Fuels	36,425.0449	55.9802	1.0000	0.1000	2,039.0830	0.0364	0.0036
	Peat Fuels	21.8741	98.8600	2.0000	1.5000	2.1625	0.0000	0.0000
K	Biomass	6,387.5300	111.0049	29.4973	3.9324	709.0471	0.1884	0.0251
L	Other Fuels	1,460.6663	90.7316	3.0000	0.6000	132.5286	0.0044	0.0009
M	1.A.3 Transport	156,385.9681				10,935.0987	0.7120	0.3963
N	Solid Fuels	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
O	Liquid Fuels	149,451.8541	72.1742	4.5677	2.5507	10,786.5641	0.6827	0.3812
P	Gaseous Fuels	2,653.3393	55.9802	5.0000	2.0000	148.5346	0.0133	0.0053
Q	Biomass	4,280.7746	70.2527	3.7633	2.2852	300.7358	0.0161	0.0098
R	1.A.4 Other Sectors	124,864.6865				8,681.2829	7.5983	0.2637
S	Solid Fuels	20,554.3141	98.6770	300.0000	1.4556	2,028.2376	6.1663	0.0299
T	Liquid Fuels	58,696.4154	72.1820	9.1632	3.7567	4,236.8232	0.5378	0.2205
U	Gaseous Fuels	43,162.0478	55.9802	5.0000	0.1000	2,416.2221	0.2158	0.0043
V	Biomass	2,451.9093	107.6680	276.6727	3.6738	263.9921	0.6784	0.0090
W	1.A.5 Other (Not specified elsewhere)⁽⁶⁾	NO	NO	NO	NO	NO	NO	NO
X	1.A. Fuel Combustion	503,410.6607				32,482.6220	8.8993	1.1232
	Memo Items							
Y	Aviation Bunkers	25,255.1500	71.4000	0.3192	2.3524	1,803.2177	0.0081	0.0594
Z	Marine Bunkers	5,978.7317	73.6426	7.0000	2.0000	440.2891	0.0419	0.0120
AA	CO ₂ from Biomass	17,715.0880	98.0748			1,737.4041	NA	NA

Table 3.1.3 Correspondence between National Disaggregation of Sources and IPCC Combustion Source Categories

IPCC Source Category/Fuel Groups from Table 3.1.2	National Disaggregated Sources from Table 3.1.1
A 1.A.1 Energy Industries (A = B+C+D+E+F)	
B (a) Solid Fuels	1+2+11
C (b) Liquid Fuels	3+7+9+10
D (c) Gaseous Fuels	4+8
E (d) Biomass	5
F (e) Other Fuels	6
G 1.A.2 Manufacturing Industries (G = H+I+J+K+L)	
H (a) Solid Fuels	12+13
I (b) Liquid Fuels	14+15+16+17+18+19
J (c) Gaseous Fuels	20
K (d) Biomass	21+22
L (e) Other Fuels	23
M 1.A.3 Transport (M = N+O+P+Q)	
N (a) Solid Fuels	NO
O (b) Liquid Fuels	24+25+26+27+29+30+31
P (c) Gaseous Fuels	32
Q (d) Biomass	28+33
R 1.A.4 Other Sectors (R = S+T+U+V)	
S (a) Solid Fuels	34+35+36+37+46+47+48+49+50
T (b) Liquid Fuels	38+39+40+41+42+51+52+53+54+57
U (c) Gaseous Fuels	43+55
V (d) Biomass	44+45+56+58
W 1.A.5 Other	NO
X 1.A Fuel Combustion (X = A+G+M+R+W)	

Table 3.1.4 Emissions of CO₂ from the Reference Approach in 2013 [CRF 2012 Table 1.A(b)]

FUEL TYPES			Unit	Production	Imports	Exports	International bunkers	Stock change	Apparent consumption	Conversion factor (TJ/Unit) ¹	NCV/ GCV ⁽²⁾	Apparent consumption (TJ)	Carbon emission factor (t C/TJ)	Carbon content (kt)	Carbon stored (kt C)	Net carbon emissions ((kt) C)	Fraction of carbon oxidized	Actual CO ₂ emissions ((kt) CO ₂)	
Liquid fossil	Primary fuels	Crude oil	Mg	NO	3,153,470.92	222,239.84		95,000.00	2,836,231.08	42.81	NCV	121,430.96	20.00	2,428.62	NO	2,428.62	1.00	8,904.94	
		Orimulsion	Mg	NO		NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO
		Natural gas liquids	Mg	NO		NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO
	Secondary fuels	Gasoline	Mg		631,404.57	11,841.70	NO	5,162.69	NO	44.59	NCV	27,395.74	19.08	522.71	NO	522.71	1.00	1,916.61	
		Jet kerosene	Mg		841,756.11	396.14	572,700.76	-4,987.51	273,646.73	44.10	NCV	12,067.71	19.47	234.99	NO	234.99	1.00	861.65	
		Other kerosene	Mg		404,143.02	3,878.58	NO	-17,653.00	417,917.44	44.20	NCV	18,470.24	19.47	359.67	NO	359.67	1.00	1,318.77	
		Shale oil	Mg			NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	NO
		Gas/diesel oil	Mg		2,008,042.79	338.41	120,535.60	-3,508.29	1,890,677.07	43.31	NCV	81,882.01	19.99	1,636.90	NO	1,636.90	1.00	6,001.95	
		Residual fuel oil	Mg		184,209.31	914,506.58	18,395.30	1,708.54	-750,401.11	41.24	NCV	-30,943.39	20.73	-641.37	NO	-641.37	1.00	-2,351.69	
		Liquefied petroleum gases (LPG)	Mg		91,117.08	27,864.61		2,051.97	61,200.50	47.16	NCV	2,885.96	17.37	50.14	NO	50.14	1.00	183.84	
		Ethane	Mg			NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	NO
		Naphtha	Mg			NO	27,023.34		-1,340.84	-25,682.49	44.00	NCV	-1,130.11	20.00	-22.60	NO	-22.60	1.00	-82.88
		Bitumen	Mg		278,417.63	73.83			278,343.81	37.70	NCV	10,492.98	22.00	230.85	230.85	0.00	1.00	0.00	
		Lubricants	Mg		37,326.00	6,646.00	NO		30,680.00	42.29	NCV	1,297.36	20.00	25.95	25.95	0.00	1.00	0.00	
		Petroleum coke	Mg		145,748.03	109.11		-14,925.35	160,564.27	32.22	NCV	5,173.99	25.43	131.57	NO	131.57	1.00	482.43	
		Refinery feedstocks	Mg			NO	20,686.51		4,658.53	-25,345.04	44.59	NCV	-1,130.12	20.00	-22.60	NO	-22.60	1.00	-82.88
		Other oil	Mg			NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	NO
Other liquid fossil												42.77		0.86	0.86	0.00		0.00	
		White Spirit	Mg	NO	999.00	27.00	NO	NO	972.00	44.00	NCV	42.77	20.00	0.86	0.86	0.00	1.00	0.00	
		Aviation Gasoline	Mg	NO		NO	NO	NO	NO	44.59	NCV	NO	19.08	NO	NO	NO	1.00	NO	
		Other Petroleum products	Mg	NO		NO	NO	NO	NO	40.20	NCV	NO	20.00	NO	NO	NO	1.00	NO	
		Paraffin Wax	Mg	NO		NO	NO	NO	NO	40.20	NCV	NO	20.00	NO	NO	NO	1.00	NO	
Liquid fossil totals												247,936.09		4,935.67	257.65	4,678.02		17,152.74	
Solid fossil	Primary fuels	Anthracite ⁽⁵⁾	Mg	NO	65,609.18	219.48		5,694.58	59,695.12	27.84	NCV	1,662.04	26.81	44.56	NO	44.56	1.00	163.38	
		Coking coal	Mg	NO	NO	NO			NO	29.10	NCV	NO	25.80	NO	NO	NO	1.00	NO	
		Other bituminous coal	Mg	NO	2,304,942.72	18,401.63	NO	220,964.64	2,065,576.45	25.54	NCV	52,764.56	25.80	1,361.33	NO	1,361.33	1.00	4,991.53	
		Sub-bituminous coal	Mg	NO	NO	NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	NO
		Lignite	Mg	NO	46,066.44	1,346.48		8,072.37	36,647.59	19.82	NCV	726.22	27.55	20.00	NO	20.00	1.00	73.35	
		Oil shale and tar sand	Mg	NO		NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	NO
	Secondary fuels	BKB ⁽⁴⁾ and patent fuel	Mg			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO
		Coke oven/gas coke	Mg			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO	NO	1.00	NO
		Coal tar	Mg				NO	NO		NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO
Other solid fossil												264.67		7.10	NO	7.10		26.02	
		Manufactured Ovoids	Mg	NO	30,144.30	12,403.20	NO	9,470.28	8,270.82	32.00	NCV	264.67	26.81	7.10	NO	7.10	1.00	26.02	
Solid fossil totals												55,417.49		1,432.98	NO	1,432.98		5,254.27	
Gaseous fossil		Natural gas (dry)	TJ	6,319.90	155,638.78	NO		-150.82	162,109.50	1.00	NCV	162,109.50	15.55	2,521.29	NO	2,521.29	1.00	9,244.73	
Other gaseous fossil												NO		NO	NO	NO		NO	
Gaseous fossil totals												162,109.50		2,521.29	NO	2,521.29		9,244.73	
Waste (non-biomass fraction)			TJ	2,418.11		NO	NO	NO	2,418.11	1.00	NCV	2,418.11	24.71	59.75	NO	59.75	1.00	219.08	
Other fossil fuels												NO		NO	NO	NO		NO	
Other fossil			TJ	NO		NO	NO	NO	NO	NO	NCV	NO	NO	NO	NO	NO	NO	NO	
Peat ^(5,6)			TJ	54,073.57		NO	357.50	23,440.25	30,275.81	1.00	NCV	30,275.81	30.95	937.00	NO	937.00	1.00	3,435.68	
Total												498,157.00		9,886.69	257.65	9,629.04		35,306.49	
Biomass total												17,845.05		505.28	NO	505.28		1,852.70	
		Solid biomass	TJ	9,112.77	1,496.84	0.21		39.93	10,569.47	1.00	NCV	10,569.47	29.09	307.49	NO	307.49	1.00	1,127.45	
		Liquid biomass	TJ	913.46	3,293.21	NO		35.10	4,171.57	1.00	NCV	4,171.57	33.10	138.06	NO	138.06	1.00	506.24	
		Gas biomass	TJ	2,016.78		NO	NO	NO	2,016.78	1.00	NCV	2,016.78	14.90	30.05	NO	30.05	1.00	110.18	
		Other non-fossil fuels (biogenic waste)	TJ	1,087.23		NO	NO	NO	1,087.23	1.00	NCV	1,087.23	27.30	29.68	NO	29.68	1.00	108.83	

Table 3.1.5 Comparison of Results from Sectoral Approach and Reference Approach for 2013 (CRF 2012 Table 1.A(c))

FUEL TYPES	REFERENCE APPROACH			SECTORAL APPROACH ⁽¹⁾		DIFFERENCE ⁽²⁾	
	Apparent energy consumption ⁽³⁾ (PJ)	Apparent energy consumption (excluding non-energy use, reductants and feedstocks) ⁽⁴⁾ (PJ)	CO ₂ emissions (kt)	Energy consumption (PJ)	CO ₂ emissions ⁽⁵⁾ (kt)	Energy consumption (%)	CO ₂ emissions ⁽⁶⁾ (%)
Liquid fuels (excluding international bunkers)	247.94	236.51	17,152.74	236.51	17,209.41	0.00	-0.33
Solid fuels (excluding international bunkers)	55.42	55.50	5,254.27	55.50	5,222.38	0.00	0.61
Gaseous fuels	162.11	161.23	9,244.73	161.23	8,976.93	0.00	2.98
Other fossil fuels	2.42	2.42	219.08	2.42	219.03	0.00	0.02
Peat	30.28	31.05	3,435.68	31.05	3,511.54	0.00	-2.16
Total⁽⁵⁾	498.16	486.71	35,306.49	486.71	35,139.30	0.00	0.48

Table 3.1.6 (a) Implied emission factors (IEFs) for CO₂ – Liquid Fuels in Sector 1.A.1.a

Energy (TJ)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Heavy Fuel Oil	13978.93	23298.22	23793.05	23504.40	26473.38	25318.78	25772.37	33483.46	45689.26	57070.34	41771.86	50019.02	36436.44	23912.76	30201.29	30040.79	26180.99	15749.12	14187.59	8382.58	4308.13	1695.56	1648.98	1393.89	Energy balance data
Gasoil	303.16	259.85	346.47	216.54	779.55	649.62	389.77	476.39	606.32	1082.71	1212.63	1082.71	822.86	1169.32	1645.71	2841.39	2259.13	580.00	457.38	355.46	1095.69	325.16	304.27	188.08	Energy balance data
total	14282.09	23558.07	24139.52	23720.94	27252.93	25968.40	26162.15	33959.85	46295.57	58153.04	42984.49	51101.72	37259.30	25082.08	31847.00	32882.18	28440.12	16329.13	14644.96	8738.04	5403.82	2020.72	1953.25	1581.98	Liquid Fuels CRFReporter
Emission Factors (t CO₂/TJ)																									
Heavy Fuel Oil	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	
Gasoil	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	
CO₂ Emissions National Approach Tier 1)																									
Heavy Fuel Oil	1062.40	1770.66	1808.27	1786.33	2011.98	1924.23	1958.70	2544.74	3472.38	4337.35	3174.66	3801.45	2769.17	1817.37	2295.30	2283.10	1989.76	1196.93	1078.26	637.08	327.42	128.86	125.32	105.94	
Gasoil	22.22	19.05	25.40	15.87	57.14	47.62	28.57	34.92	44.44	79.36	88.89	79.36	60.32	85.71	120.63	208.27	165.59	42.51	33.53	26.06	80.31	23.83	22.30	13.79	
total	1084.62	1789.71	1833.67	1802.21	2069.12	1971.84	1987.27	2579.66	3516.83	4416.71	3263.55	3880.81	2829.48	1903.08	2415.93	2491.37	2155.35	1239.45	1111.78	663.13	407.73	152.70	147.63	119.72	
IEF calculated	75.94	75.97	75.96	75.98	75.92	75.93	75.96	75.96	75.96	75.95	75.92	75.94	75.94	75.87	75.86	75.77	75.79	75.90	75.92	75.89	75.45	75.57	75.58	75.68	IEF National Approach
CO₂ emissions from ETS (Tier 3 bottom up)	1086.52	1788.62	1834.97	1803.59	2067.85	1985.81	1986.13	2617.23	3519.32	4533.60	3484.39	3925.34	2905.74	1992.62	2539.76	2562.77	2222.00	1284.23	1143.44	680.41	423.79	158.24	152.61	124.24	kt CO₂ CRFReporter
difference	0.17%	-0.06%	0.07%	0.08%	-0.06%	0.70%	-0.06%	1.44%	0.07%	2.58%	6.34%	1.13%	2.62%	4.49%	4.88%	2.79%	3.00%	3.49%	2.77%	2.54%	3.79%	3.50%	3.27%	3.64%	
IEF reported	76.08	75.92	76.02	76.03	75.88	76.47	75.92	77.07	76.02	77.96	81.06	76.81	77.99	79.44	79.75	77.94	78.13	78.65	78.08	77.87	78.42	78.31	78.13	78.54	IEF CRFReporter




Figure 3.1.1 Implied emission factors (IEFs) for CO₂ – Liquid Fuels in Sector 1.A.1.a

Table 3.1.7 (a) Implied emission factors (IEFs) for CO₂ – Solid Fuels in Sector 1.A.1.a

Energy (TJ)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Coal	51972.08	51137.65	58287.50	56810.07	58544.91	62584.78	62177.52	60202.93	61174.75	52951.80	59728.50	63375.92	61343.32	55395.13	57003.60	59307.55	50953.01	49049.52	41505.71	32444.13	36320.56	38228.46	48571.51	40625.48	Energy balance data
Milled Peat	23463.58	24374.71	25363.72	23222.17	23821.80	23385.71	22832.80	22871.73	21516.72	21516.72	20021.53	22466.79	22529.09	21010.53	13816.83	20488.09	18970.00	18806.62	23856.88	23314.27	20144.16	19799.83	23027.37	20830.40	Energy balance data
Sod Peat	1323.57	1218.74	904.22	445.56	314.51	314.51	982.85	353.83	183.47	183.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	Energy balance data
total	76759.23	76731.10	84555.44	80477.79	82681.22	86285.00	85993.17	83428.49	82874.94	74651.99	79750.03	85842.71	83872.41	76405.67	70820.42	79795.64	69923.01	67856.14	65362.59	55758.40	56464.72	58028.29	71598.87	61455.88	Solid Fuels CRFReporter
Emission Factors (t CO₂/TJ)																									
Coal	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	94.60	
Milled Peat	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	115.00	
Sod Peat	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	104.00	
CO₂ Emissions National Approach (Tier 1)																									
Coal	4916.56	4837.62	5514.00	5374.23	5538.35	5920.52	5881.99	5695.20	5787.13	5009.24	5650.32	5995.36	5803.08	5240.38	5392.54	5610.49	4820.15	4640.08	3926.44	3069.21	3435.93	3616.41	4594.86	3843.17	
Milled Peat	2698.31	2803.09	2916.83	2670.55	2739.51	2689.36	2625.77	2630.25	2474.42	2474.42	2302.48	2583.68	2590.85	2416.21	1588.93	2356.13	2181.55	2162.76	2743.54	2681.14	2316.58	2276.98	2648.15	2395.50	
Sod Peat	137.65	126.75	94.04	46.34	32.71	32.71	102.22	36.80	19.08	19.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
total	7752.52	7767.46	8524.86	8091.12	8310.56	8642.59	8609.98	8362.24	8280.63	7502.74	7952.79	8579.04	8393.92	7656.59	6981.48	7966.62	7001.71	6802.85	6669.98	5750.36	5752.50	5893.39	7243.01	6238.67	
IEF calculated	101.00	101.23	100.82	100.54	100.51	100.16	100.12	100.23	99.92	100.50	99.72	99.94	100.08	100.21	98.58	99.84	100.13	100.25	102.05	103.13	101.88	101.56	101.16	101.51	IEF National Approach
CO₂ emissions from ETS (Tier 3 bottom up)	7909.31	7795.76	8477.38	7955.29	8205.33	8645.06	8857.12	8606.14	8145.16	7454.31	8084.48	8688.84	8397.65	7731.52	7078.28	7909.68	6966.22	6703.73	6630.99	5765.58	5688.15	5857.32	7228.47	6255.70	kt CO₂ CRFReporter
difference	1.98%	0.36%	-0.56%	-1.71%	-1.28%	0.03%	2.79%	2.83%	-1.66%	-0.65%	1.63%	1.26%	0.04%	0.97%	1.37%	-0.72%	-0.51%	-1.48%	-0.59%	0.26%	-1.13%	-0.62%	-0.20%	0.27%	
IEF reported	103.04	101.60	100.26	98.85	99.24	100.19	103.00	103.16	98.28	99.85	101.37	101.22	100.12	101.19	99.95	99.12	99.63	98.79	101.45	103.40	100.74	100.94	100.96	101.79	IEF CRFReporter

IEF CO₂ - Solid Fuels 1A1a




Figure 3.1.2 Implied emission factors (IEFs) for CO₂ – Solid Fuels in Sector 1A.1.a

Table 3.1.8 Implied emission factors (IEFs) for CO₂ – Liquid Fuels in Sector 1.A.2.f

Energy (TJ)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Energy balance data
Kerosene	144.70	90.47	85.15	95.80	164.98	212.88	287.39	313.99	372.54	473.65	452.37	474.04	456.99	493.83	452.85	2326.07	1695.03	1390.03	1574.29	1573.05	1232.17	973.91	832.34	860.94	Energy balance data
Fuel Oil	3611.38	2007.31	2415.72	2466.61	3153.10	2935.85	2748.41	3012.82	2979.30	3180.05	3390.73	2668.90	2255.59	1951.78	1818.20	2367.06	1689.75	1534.88	1362.43	1347.66	1048.21	300.58	246.50	551.05	Energy balance data
LPG	1918.60	1467.29	1423.87	1498.04	1601.82	1578.86	1537.29	1551.88	1705.02	1799.71	1978.32	1862.06	2206.72	2839.82	3369.70	2960.05	3022.38	3401.18	3642.11	2143.33	2748.68	2658.72	2564.18	3105.81	Energy balance data
Gasoil	3119.23	4295.10	4379.59	4295.10	4970.99	5675.05	4435.91	4633.05	4689.37	4999.16	5139.97	5419.73	5321.05	5288.70	4858.54	5767.26	5527.08	5419.28	5008.75	4715.99	4460.62	4273.74	4075.87	3549.72	Energy balance data
Petroleum Coke	1971.87	1994.03	1462.29	1329.35	1971.87	2193.43	1019.17	2747.33	2414.99	2614.39	4608.42	5277.33	5951.78	7586.37	9018.88	9924.14	9829.63	10015.55	9196.13	4818.39	3053.84	2832.63	3799.72	3702.53	Energy balance data
Naphta	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	44.00	0.00	0.00	0.00	Energy balance data
total	10809.78	9898.20	9810.62	9728.91	11906.77	12640.08	10072.17	12303.07	12205.22	13110.97	15613.80	15746.06	16236.14	18204.52	19562.16	23388.58	21807.88	21804.92	20827.71	14642.43	12587.52	11039.57	11518.61	11770.04	Energy balance data
Emission Factors (t CO₂/TJ)																									
Kerosene	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40	71.40
Fuel Oil	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00
LPG	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70	63.70
Gasoil	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30	73.30
Petroleum Coke	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00	76.00
Naphta	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33	73.33
CO₂ Emissions National Approach (Tier 1)																									
Kerosene	10.33	6.46	6.08	6.84	11.78	15.20	20.52	22.42	26.60	33.82	32.30	33.85	32.63	35.26	32.33	166.08	121.03	99.25	112.40	112.32	87.98	69.54	59.43	61.47	
Fuel Oil	274.46	152.56	183.59	187.46	239.64	223.12	208.88	228.97	226.43	241.68	257.70	202.84	171.42	148.34	138.18	179.90	128.42	116.65	103.54	102.42	79.66	22.84	18.73	41.88	
LPG	122.21	93.47	90.70	95.43	102.04	100.57	97.93	98.85	108.61	114.64	126.02	118.61	140.57	180.90	214.65	188.56	192.53	216.66	232.00	136.53	175.09	169.36	163.34	197.84	
Gasoil	228.64	314.83	321.02	314.83	364.37	415.98	325.15	339.60	343.73	366.44	376.76	397.27	390.03	387.66	356.13	422.74	405.14	397.23	367.14	345.68	326.96	313.26	298.76	260.19	
Petroleum Coke	149.86	151.55	111.13	101.03	149.86	166.70	77.46	208.80	183.54	198.69	350.24	401.08	452.34	576.56	685.44	754.23	747.05	761.18	698.91	366.20	232.09	215.28	288.78	281.39	
Naphta	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	3.23	0.00	0.00	0.00	
total	788.74	722.09	715.76	708.82	870.91	924.81	733.16	901.87	892.13	958.50	1146.24	1156.87	1190.22	1331.94	1429.96	1714.73	1597.39	1594.20	1517.22	1066.37	905.01	790.29	829.04	842.78	
IEF calculated	72.97	72.95	72.96	72.86	73.14	73.16	72.79	73.30	73.09	73.11	73.41	73.47	73.31	73.17	73.10	73.32	73.25	73.11	72.85	72.83	71.90	71.59	71.97	71.60	IEF National Approach kt CO₂
CO₂ emissions difference	823.55	757.28	741.57	732.28	905.72	963.53	751.15	950.37	934.76	1004.65	1227.59	1250.02	1295.28	1465.86	1589.16	1904.62	1768.76	1766.53	1672.93	1150.95	958.75	839.13	893.15	905.47	CRFReporter
IEF reported	76.19	76.51	75.59	75.27	76.07	76.23	74.58	77.25	76.59	76.63	78.62	79.39	79.78	80.52	81.24	81.43	81.11	81.01	80.32	78.60	76.17	76.01	77.54	76.93	IEF CRFReporter
% Share of Fuels																									
Kerosene	1.34%	0.91%	0.87%	0.98%	1.39%	1.68%	2.85%	2.55%	3.05%	3.61%	2.90%	3.01%	2.81%	2.71%	2.31%	9.95%	7.77%	6.37%	7.56%	10.74%	9.79%	8.82%	7.23%	7.31%	
Fuel Oil	33.41%	20.28%	24.62%	25.35%	26.48%	23.23%	27.29%	24.49%	24.41%	24.25%	21.72%	16.95%	13.89%	10.72%	9.29%	10.12%	7.75%	7.04%	6.54%	9.20%	8.33%	2.72%	2.14%	4.68%	
LPG	17.75%	14.82%	14.51%	15.40%	13.45%	12.49%	15.26%	12.61%	13.97%	13.73%	12.67%	11.83%	13.59%	15.60%	17.23%	12.66%	13.86%	15.60%	17.49%	14.64%	21.84%	24.08%	22.26%	26.39%	
Gasoil	28.86%	43.39%	44.64%	44.15%	41.75%	44.90%	44.04%	37.66%	38.42%	38.13%	32.92%	34.42%	32.77%	29.05%	24.84%	24.66%	25.34%	24.85%	24.05%	32.21%	35.44%	38.71%	35.39%	30.16%	
Petroleum Coke	18.24%	20.15%	14.91%	13.66%	16.56%	17.35%	10.12%	22.33%	19.79%	19.94%	29.52%	33.52%	36.66%	41.67%	46.10%	42.43%	45.07%	45.93%	44.15%	32.91%	24.26%	25.66%	32.99%	31.46%	
Naphta	0.41%	0.44%	0.45%	0.45%	0.37%	0.35%	0.44%	0.36%	0.36%	0.34%	0.28%	0.28%	0.27%	0.24%	0.22%	0.19%	0.20%	0.20%	0.21%	0.30%	0.35%	0.00%	0.00%	0.00%	
total	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	

Annex 3.1.B

Energy - Transport (IPCC Sector 1.A.3)

3.1.9 – 3.1.12 Civil aviation data 1990-2013

3.1.13 Vehicle population data 1990-2013

Table 3.1.9 Number of Domestic LTOs by departure airport 1990-2013

Domestic LTOs No.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
ex Dublin	7657	6944	6885	5921	6153	7235	7742	8134	8991	10183	11018	10947	10849	11261	12143	9976	10392	10803	9611	7844	6074	3331	2190	2101
ex Cork	2872	2604	2582	2221	2307	2713	2903	3050	3372	3819	4132	4106	4069	4223	4438	3649	3721	4608	3919	2872	1861	809	445	441
ex Shannon	1737	1576	1562	1343	1396	1641	1757	1845	2040	2310	2500	2484	2462	2555	2865	2809	2892	2277	1897	1349	1077	834	764	800
ex Galway	1425	1293	1282	1102	1145	1347	1441	1514	1674	1895	2051	2038	2019	2096	2224	1631	1615	1815	1848	1563	1746	1252	51	31
ex Sligo	620	562	557	479	498	586	627	658	728	824	892	886	878	912	946	759	748	754	785	741	678	381	35	25
ex Donegal	581	527	523	449	467	549	588	617	682	773	836	831	824	855	717	684	747	736	754	739	697	721	733	723
ex Knock	445	404	400	344	358	421	450	473	523	592	641	637	631	655	753	565	557	568	481	510	454	253	79	83
ex Kerry	1133	1027	1019	876	910	1070	1145	1203	1330	1506	1630	1620	1605	1666	1755	1477	1515	1506	1418	1170	1048	460	781	776
ex Waterford	236	214	213	183	190	223	239	251	278	314	340	338	335	348	254	181	191	279	456	231	472	707	175	155
ex Other	347	314	312	268	279	328	350	368	407	461	499	496	491	510	539	495	518	411	476	305	282	277	241	282
Total	17053	15465	15334	13187	13703	16113	17242	18115	20024	22679	24538	24381	24164	25080	26634	22226	22896	23757	21645	17324	14389	9025	5494	5417
Domestic LTOs (%)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
ex Dublin	45%	45%	45%	45%	45%	45%	45%	45%	45%	45%	45%	45%	45%	45%	46%	45%	45%	45%	44%	45%	42%	37%	40%	39%
ex Cork	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	17%	16%	16%	19%	18%	17%	13%	9%	8%	8%
ex Shannon	10%	10%	10%	10%	10%	10%	10%	10%	10%	10%	10%	10%	10%	10%	11%	13%	13%	10%	9%	8%	7%	9%	14%	15%
ex Galway	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	8%	7%	7%	8%	9%	9%	12%	14%	1%	1%
ex Sligo	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	4%	3%	3%	3%	4%	4%	5%	4%	1%	0%
ex Donegal	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	4%	5%	8%	13%	13%
ex Knock	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	2%	2%	2%	3%	3%	3%	1%	2%
ex Kerry	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	7%	6%	7%	7%	7%	5%	14%	14%
ex Waterford	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	2%	1%	3%	8%	3%	3%
ex Other	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	2%	3%	4%	5%
Total	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%

Table 3.1.10 Distances between airport pairs used to estimate fuel consumption for cruise phase

Nautical Miles		Cork	Galway	Donegal	Dublin	Knock	Kerry	Shannon	Sligo	Waterford	Other
		EICK	EICM	EIDL	EIDW	EIKN	EIKY	EINN	EISG	EIWF	
EICK	Cork		89.18	192.52	124.89	124.88	43.37	54.12	146.58	56.04	89.18
EICM	Galway	89.18		106.92	96.28	36.93	70.51	35.94	60.13	95.09	89.18
EIDL	Donegal	192.52	106.92		121.75	70.16	177.15	142.26	46.80	177.42	89.18
EIDW	Dublin	124.89	96.28	121.75		95.56	139.93	105.34	97.52	79.89	89.18
EIKN	Knock	124.88	36.93	70.16	95.56		106.99	72.70	23.53	121.02	89.18
EIKY	Kerry	43.37	70.51	177.15	139.93	106.99		38.25	130.45	89.97	89.18
EINN	Shannon	54.12	35.94	142.26	105.34	72.70	38.25		95.53	74.21	89.18
EISG	Sligo	146.58	60.13	46.80	97.52	23.53	130.45	95.53		137.05	89.18
EIWF	Waterford	56.04	95.09	177.42	79.89	121.02	89.97	74.21	137.05		89.18
	Other	89.18	89.18	89.18	89.18	89.18	89.18	89.18	89.18	89.18	

Table 3.1.11 LTO emissions factors by aircraft type

Aircraft Type	kg of fuel per LTO	CH ₄ kg/ LTO	N ₂ O kg/ LTO
A30B	1540.55	0.12	0.20
A310	1540.55	0.63	0.20
A320	802.33	0.06	0.10
A321	802.33	0.14	0.10
A332	2231.52	0.13	0.20
A333	2231.52	0.13	0.20
A343	2231.52	0.39	0.20
AT43	115.20	0.02	0.02
AT72	137.00	0.03	0.02
ATP	569.51	0.10	0.10
B462	569.51	0.14	0.10
B463	569.51	0.14	0.10
B733	825.39	0.08	0.10
B734	825.39	0.08	0.10
B737	784.12	0.09	0.10
B738	763.48	0.07	0.10
B752	1253.00	0.02	0.10
B762	1617.09	0.33	0.10
B763	1617.09	0.12	0.20
B764	1617.09	0.10	0.20
BE20	51.80	0.06	0.01
BE40	58.30	0.06	0.01
CL30	569.51	0.10	0.10
CL60	569.51	0.10	0.10
DC10	2381.18	0.24	0.20
GLF2	569.51	0.14	0.10
GLF4	569.51	0.14	0.10
GLF5	569.51	0.03	0.10
H25B	569.51	0.14	0.10
LJ31	569.51	0.14	0.10
LJ45	569.51	0.14	0.10
LJ60	569.51	0.14	0.10
MD11	1003.06	0.24	0.20
MD82	1003.06	0.19	0.10
MD83	1003.06	0.19	0.10
T154	2190.00	7.59	0.20
Other	49.57	0.02	0.10

Table 3.1.12 Weighted Cruise fuel use per flight (IEF) by departure airport 1990-2013

kg Fuel	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
ex Dublin	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	494.8	442.3	459.8	537.6	496.4	510.5	526.2	490.8	409.1	390.2	359.3
ex Cork	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	394.2	245.3	266.9	399.9	410.7	439.8	494.8	501.9	394.4	173.2	168.7
ex Shannon	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	979.8	1010.3	1055.9	1059.3	978.1	990.0	938.6	826.5	625.6	655.3	549.0
ex Galway	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	167.3	159.4	160.3	158.5	176.9	196.1	160.5	159.2	147.9	124.7	102.8
ex Sligo	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	165.7	164.2	164.1	166.9	165.6	163.6	167.6	168.0	143.7	108.2	97.6
ex Donegal	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	213.7	212.1	210.6	213.8	215.2	212.4	216.0	215.6	191.0	126.4	127.0
ex Knock	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	214.5	192.6	202.7	242.8	244.8	230.1	201.6	186.8	176.0	240.2	232.3
ex Kerry	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	421.1	246.7	247.8	247.0	242.4	452.1	757.9	753.8	533.6	242.7	240.6
ex Waterford	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	158.9	104.8	109.8	105.5	210.1	287.0	130.4	164.8	151.2	101.9	72.7
ex Other	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	150.6	160.2	157.6	148.5	165.4	139.9	140.7	141.8	159.7	138.0	128.0
Total	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	454.5	402.0	433.5	492.9	451.1	467.0	485.6	440.6	334.9	326.7	298.9

Table 3.1.13 Vehicle numbers, by technology class 1990, 2000-2013

Sector	Subsector	Technology	1990	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	38455	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	62490	37	29	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	187469	6941	5063	2955	1471	393	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	192276	124459	101411	67394	51709	33473	25246	15904	11489	3758	3111	2349	2016	1901	949
Passenger Cars	Gasoline 0,8 - 1,4 l	Open Loop	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 1 - 91/441/EEC	0	229196	211281	202575	179992	161058	130173	107123	77929	81303	56760	41730	28922	22019	7739
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 2 - 94/12/EEC	0	333506	405547	396295	389764	379093	359730	336491	307589	365013	336509	286154	253098	216941	201061
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 3 - 98/69/EC Stage2000	0	0	0	71457	137486	204417	273777	266772	258354	295682	299096	287708	285219	280034	276634
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	73236	143395	204450	223224	240298	243904	248206	251944
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	14814	27035	39297
Passenger Cars	Gasoline 0,8 - 1,4 l	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	18263	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	29677	22	18	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	89030	4191	3129	1900	967	269	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	91313	75149	62671	43318	34002	22886	18190	12265	9511	2199	1786	1387	1146	1046	504
Passenger Cars	Gasoline 1,4 - 2,0 l	Open Loop	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 1 - 91/441/EEC	0	138389	130570	130207	118358	110118	93792	82616	64507	47570	32588	24352	16717	12425	4434
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 2 - 94/12/EEC	0	201371	250625	254722	256298	259193	259194	259509	254611	213566	193205	166989	143831	119396	106786
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 3 - 98/69/EC Stage2000	0	0	0	45930	90407	139764	197262	205741	213856	173002	171724	168168	162071	154101	146907
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	56481	118698	119622	128163	143356	140996	138768	136307
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	8750	15439	21812
Passenger Cars	Gasoline 1,4 - 2,0 l	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1319	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline >2,0 l	ECE 15/02	2144	2	2	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline >2,0 l	ECE 15/03	6431	377	294	188	101	30	0	0	0	0	0	0	0	0	0
Passenger Cars	Gasoline >2,0 l	ECE 15/04	6596	6767	5879	4282	3564	2584	2192	1621	1353	188	149	118	86	73	33
Passenger Cars	Gasoline >2,0 l	PC Euro 1 - 91/441/EEC	0	12463	12249	12872	12406	12432	11303	10921	9174	4072	2713	1912	1408	1012	425
Passenger Cars	Gasoline >2,0 l	PC Euro 2 - 94/12/EEC	0	18134	23512	25181	26865	29261	31235	34305	36208	18283	16082	13114	10753	8375	7055
Passenger Cars	Gasoline >2,0 l	PC Euro 3 - 98/69/EC Stage2000	0	0	0	4540	9476	15778	23772	27197	30413	14810	14294	13367	12108	10800	9699
Passenger Cars	Gasoline >2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	7466	16880	10241	10668	11967	11303	10464	9733
Passenger Cars	Gasoline >2,0 l	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	732	1248	1723
Passenger Cars	Gasoline >2,0 l	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	69656	30950	24735	17092	13167	8901	7170	5187	4309	780	659	543	501	430	182
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 1 - 91/441/EEC	0	53969	49069	49217	44564	42329	36971	34938	29228	17728	12052	9011	6615	4943	1452
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 2 - 94/12/EEC	0	78532	94187	96282	96502	99633	102168	109746	115363	66463	62641	54765	50321	42158	38544
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 3 - 98/69/EC Stage2000	0	0	0	17361	34040	53725	77756	87007	96897	115476	127136	127691	133178	131765	132268
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	23886	53781	121117	163082	221145	245016	264644	288016
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	53087	106094	160045
Passenger Cars	Diesel 1,4 - 2,0 l	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Diesel >2,0 l	Conventional	6057	3439	3057	2112	1795	1214	1071	844	760	189	135	109	88	68	26
Passenger Cars	Diesel >2,0 l	PC Euro 1 - 91/441/EEC	0	5997	6065	6083	6077	5772	5525	5688	5158	4295	2472	1814	1167	783	209
Passenger Cars	Diesel >2,0 l	PC Euro 2 - 94/12/EEC	0	8726	11641	11900	13159	13586	15267	17866	20358	16100	12847	11023	8877	6678	5545
Passenger Cars	Diesel >2,0 l	PC Euro 3 - 98/69/EC Stage2000	0	0	0	2146	4642	7326	11619	14164	17100	27973	26074	25702	23493	20873	19029
Passenger Cars	Diesel >2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	3888	9491	29339	33446	44513	43222	41923	41435

Sector	Subsector	Technology	1990	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Passenger Cars	Diesel >2,0 l	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	9365	16807	23025
Passenger Cars	Diesel >2,0 l	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	LPG	Conventional	2830	220	220	150	153	153	0	0	0	0	0	0	0	0	0
Passenger Cars	LPG	PC Euro 1 - 91/441/EEC	0	259	253	245	236	225	211	196	183	164	144	125	106	88	72
Passenger Cars	LPG	PC Euro 2 - 94/12/EEC	0	200	195	189	182	173	163	151	137	123	113	97	83	69	57
Passenger Cars	LPG	PC Euro 3 - 98/69/EC Stage2000	0	0	0	0	0	0	90	83	81	79	76	72	68	63	57
Passenger Cars	LPG	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	90	83	81	79	76	72	68	63	57
Passenger Cars	LPG	PC Euro 5 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	LPG	PC Euro 6 - EC 715/2007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Hybrid Gasoline <1,4 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Hybrid Gasoline 1,4 - 2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Passenger Cars	Hybrid Gasoline >2,0 l	PC Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Light Commercial Vehicles	Gasoline <3,5t	Conventional	22695	1579	1026	664	448	294	141	89	49	33	25	17	11	5	4
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 1 - 93/59/EEC	0	1423	1146	900	718	549	357	217	134	99	79	60	45	34	28
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 2 - 96/69/EEC	0	1872	2104	1658	1366	1117	797	550	381	307	263	207	160	120	101
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 3 - 98/69/EC Stage2000	0	0	0	406	689	869	937	661	496	442	425	372	312	256	236
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	223	337	404	419	414	357	306	292
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 5 - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	30	53	78
Light Commercial Vehicles	Gasoline <3,5t	LD Euro 6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Light Commercial Vehicles	Diesel <3,5 t	Conventional	99834	54858	43847	35979	29768	23966	15614	14153	10621	8121	6516	4747	3498	1972	1441
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 1 - 93/59/EEC	0	49440	48962	48758	47757	44707	39654	34690	29131	24051	20170	16613	14285	12392	10951
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 2 - 96/69/EEC	0	65017	89886	89849	90802	91026	88477	87696	82841	74651	67336	57256	51017	43655	39482
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 3 - 98/69/EC Stage2000	0	0	0	22020	45829	70747	104091	105457	107724	107448	108917	103239	99411	93224	92220
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 4 - 98/69/EC Stage2005	0	0	0	0	0	0	0	35522	73131	98078	107365	114809	113696	111249	113834
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 5 - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	9620	19152	30260
Light Commercial Vehicles	Diesel <3,5 t	LD Euro 6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Gasoline >3,5 t	Conventional	294	167	140	119	102	87	71	53	44	42	38	30	25	23	24
Heavy Duty Trucks	Rigid <=7,5 t	Conventional	9628	3892	3046	2396	1912	1491	952	785	589	432	336	247	200	132	108
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro I - 91/542/EEC Stage I	0	2335	2199	2055	1891	1706	1423	1205	986	759	596	454	390	318	287
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro II - 91/542/EEC Stage II	0	3882	5080	4720	4453	4238	3820	3593	3282	2790	2361	1904	1667	1359	1228
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro III - 2000 Standards	0	0	0	1157	2248	3294	4504	4321	4267	4016	3802	3432	3247	2920	2868
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1467	2897	3677	3737	3827	3714	3485	3541
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	305	609	932
Heavy Duty Trucks	Rigid <=7,5 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 7,5 - 12 t	Conventional	9219	5889	4587	3588	2843	2199	1434	1159	830	550	387	281	225	150	129
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro I - 91/542/EEC Stage I	0	3533	3312	3077	2812	2515	2143	1781	1389	967	687	517	440	359	345

Sector	Subsector	Technology	1990	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro II - 91/542/EEC Stage II	0	5874	7651	7067	6623	6249	5753	5308	4626	3554	2724	2171	1878	1535	1477
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro III - 2000 Standards	0	0	0	1732	3343	4857	6784	6384	6015	5116	4386	3914	3660	3299	3450
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	2167	4084	4685	4311	4364	4185	3937	4258
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	343	688	1121
Heavy Duty Trucks	Rigid 7,5 - 12 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 12 - 14 t	Conventional	1414	1791	1494	1233	1032	839	612	542	401	265	176	124	103	71	66
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro I - 91/542/EEC Stage I	0	1075	1079	1057	1021	960	915	832	671	465	313	229	202	171	176
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro II - 91/542/EEC Stage II	0	1787	2492	2428	2404	2385	2457	2481	2234	1710	1240	960	860	730	754
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro III - 2000 Standards	0	0	0	595	1213	1854	2897	2983	2905	2462	1996	1731	1677	1570	1761
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1013	1972	2254	1962	1930	1918	1873	2173
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	157	327	572
Heavy Duty Trucks	Rigid 12 - 14 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 14 - 20 t	Conventional	273	511	469	396	351	304	245	226	187	145	104	80	68	48	43
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro I - 91/542/EEC Stage I	0	307	338	339	347	348	366	347	314	254	184	147	134	114	114
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro II - 91/542/EEC Stage II	0	510	782	779	818	864	982	1036	1044	934	728	618	571	488	487
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro III - 2000 Standards	0	0	0	191	413	672	1158	1245	1358	1344	1172	1114	1112	1049	1137
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	423	922	1231	1152	1242	1272	1251	1403
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	104	219	370
Heavy Duty Trucks	Rigid 14 - 20 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 20 - 26 t	Conventional	0	0	0	1	1	1	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro I - 91/542/EEC Stage I	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro II - 91/542/EEC Stage II	0	0	0	2	2	2	2	3	2	2	2	2	2	1	1
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro III - 2000 Standards	0	0	0	0	1	2	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Rigid 20 - 26 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 26 - 28 t	Conventional	0	0	0	1	1	1	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro I - 91/542/EEC Stage I	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro II - 91/542/EEC Stage II	0	0	0	2	2	2	2	3	2	2	2	2	2	1	1
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro III - 2000 Standards	0	0	0	0	1	2	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Rigid 26 - 28 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 28 - 32 t	Conventional	0	0	0	1	1	1	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro I - 91/542/EEC Stage I	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro II - 91/542/EEC Stage II	0	0	0	2	2	2	2	3	2	2	2	2	2	1	1

Sector	Subsector	Technology	1990	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
		Stage II															
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro III - 2000 Standards	0	0	0	0	1	2	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Rigid 28 - 32 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid >32 t	Conventional	0	0	0	1	1	1	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Rigid >32 t	HD Euro I - 91/542/EEC Stage I	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Rigid >32 t	HD Euro II - 91/542/EEC Stage II	0	0	0	2	2	2	2	3	2	2	2	2	2	1	1
Heavy Duty Trucks	Rigid >32 t	HD Euro III - 2000 Standards	0	0	0	0	1	2	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Rigid >32 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Rigid >32 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Rigid >32 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Articulated 40 - 50 t	Conventional	0	0	0	1	1	1	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro I - 91/542/EEC Stage I	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro II - 91/542/EEC Stage II	0	0	0	2	2	2	2	3	2	2	2	2	2	1	1
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro III - 2000 Standards	0	0	0	0	1	2	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Articulated 40 - 50 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heavy Duty Trucks	Articulated 50 - 60 t	Conventional	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro I - 91/542/EEC Stage I	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro II - 91/542/EEC Stage II	0	0	0	0	0	0	2	3	2	2	2	2	2	1	1
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro III - 2000 Standards	0	0	0	0	0	0	3	3	3	3	3	3	3	3	3
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	1	2	3	3	3	4	3	4
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Heavy Duty Trucks	Articulated 50 - 60 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Buses	Urban Buses Standard 15 - 18 t	Conventional	1590	1347	1243	1107	968	924	839	631	593	554	387	329	295	298	273
Buses	Urban Buses Standard 15 - 18 t	HD Euro I - 91/542/EEC Stage I	0	343	344	342	306	338	331	271	260	251	247	234	217	205	190
Buses	Urban Buses Standard 15 - 18 t	HD Euro II - 91/542/EEC Stage II	0	887	1091	1107	1014	1146	1142	950	927	913	914	880	829	800	752
Buses	Urban Buses Standard 15 - 18 t	HD Euro III - 2000 Standards	0	0	0	167	325	582	792	915	914	923	946	905	897	883	849
Buses	Urban Buses Standard 15 - 18 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	0	233	490	617	616	606	609	596
Buses	Urban Buses Standard 15 - 18 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	30	110	184	369
Buses	Urban Buses Standard 15 - 18 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Buses	Coaches Standard <=18 t	Conventional	3468	2877	2619	2294	2226	1823	1624	1563	1445	1315	893	760	691	693	639
Buses	Coaches Standard	HD Euro I - 91/542/EEC	0	733	726	709	705	668	640	672	634	597	571	540	507	477	445

Sector	Subsector	Technology	1990	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
	<=18 t	Stage I															
Buses	Coaches Standard <=18 t	HD Euro II - 91/542/EEC Stage II	0	1894	2299	2294	2334	2262	2209	2354	2262	2167	2112	2030	1940	1859	1765
Buses	Coaches Standard <=18 t	HD Euro III - 2000 Standards	0	0	0	347	746	1150	1532	2266	2230	2190	2188	2089	2099	2052	1990
Buses	Coaches Standard <=18 t	HD Euro IV - 2005 Standards	0	0	0	0	0	0	0	0	569	1163	1426	1422	1417	1416	1397
Buses	Coaches Standard <=18 t	HD Euro V - 2008 Standards	0	0	0	0	0	0	0	0	0	0	0	69	257	429	866
Buses	Coaches Standard <=18 t	HD Euro VI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mopeds	2-stroke <50 cm³	Conventional	637	686	609	520	492	439	394	362	344	331	377	297	241	208	188
Mopeds	2-stroke <50 cm³	Mop - Euro I	0	172	313	408	492	459	423	401	406	408	471	400	341	321	296
Mopeds	2-stroke <50 cm³	Mop - Euro II	0	0	0	0	0	78	144	215	208	210	256	206	181	170	161
Mopeds	2-stroke <50 cm³	Mop - Euro III	0	0	0	0	0	0	0	0	83	155	242	240	241	246	251
Mopeds	4-stroke <50 cm³	Conventional	637	686	608	520	492	439	394	362	344	331	377	297	240	208	188
Mopeds	4-stroke <50 cm³	Mop - Euro I	0	172	313	408	492	459	422	401	406	408	471	400	341	321	295
Mopeds	4-stroke <50 cm³	Mop - Euro II	0	0	0	0	0	78	144	215	208	210	256	206	180	170	161
Mopeds	4-stroke <50 cm³	Mop - Euro III	0	0	0	0	0	0	0	0	83	154	242	240	240	246	251
Motorcycles	2-stroke >50 cm³	Conventional	2320	3125	3358	3381	3580	3555	3499	3563	3792	4020	4335	4024	3714	3438	3329
Motorcycles	4-stroke <250 cm³	Conventional	2320	2500	2216	1893	1790	1600	1434	1318	1251	1206	1214	1046	891	756	699
Motorcycles	4-stroke <250 cm³	Mot - Euro I	0	625	1141	1488	1790	1671	1539	1461	1479	1487	1517	1408	1263	1169	1099
Motorcycles	4-stroke <250 cm³	Mot - Euro II	0	0	0	0	0	284	525	784	758	764	823	724	669	619	599
Motorcycles	4-stroke <250 cm³	Mot - Euro III	0	0	0	0	0	0	0	0	303	563	780	845	891	894	932
Motorcycles	4-stroke 250 - 750 cm³	Conventional	15148	16324	14467	12363	11687	10446	9366	8607	8171	7874	7104	6508	5864	5215	5325
Motorcycles	4-stroke 250 - 750 cm³	Mot - Euro I	0	4081	7453	9713	11687	10910	10051	9537	9657	9711	8880	8761	8308	8060	8368
Motorcycles	4-stroke 250 - 750 cm³	Mot - Euro II	0	0	0	0	0	1857	3427	5117	4952	4987	4820	4505	4398	4267	4564
Motorcycles	4-stroke 250 - 750 cm³	Mot - Euro III	0	0	0	0	0	0	0	0	1981	3675	4567	5256	5864	6164	7100
Motorcycles	4-stroke >750 cm³	Conventional	1683	1814	1608	1374	1299	1161	1041	956	908	875	789	723	652	579	592
Motorcycles	4-stroke >750 cm³	Mot - Euro I	0	453	828	1079	1299	1212	1117	1060	1073	1079	987	973	923	896	930
Motorcycles	4-stroke >750 cm³	Mot - Euro II	0	0	0	0	0	206	381	569	550	554	536	501	489	474	507
Motorcycles	4-stroke >750 cm³	Mot - Euro III	0	0	0	0	0	0	0	0	220	408	507	584	652	685	789

Annex 3.2

Industrial Processes (IPCC Sector 2)

3.2.A Cement production (IPCC sector 2.A.1)

3.2.B Lime production (IPCC sector 2.A.2)

3.2.C Glass Production (IPCC sector 2.A.3)

3.2.D Other process uses of carbonates (IPCC sector 2.A.4.a & 2.A.4.d)

3.2.E Soda ash use (IPCC sector 2.A.4.b)

Table 3.2.A Cement production 1990-2013

Activity data, emission factors and emissions

IPCC Sector 2A1 Cement	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Clinker production (kilotonnes)																
Cement Plant 1	925	921	1,399	1,462	1,496	1,566	1,691	1,669	1,665	1,685	1,424	706	579	564	853	904
Cement Plant 2	685	680	908	915	902	905	977	957	900	934	902	501	362	283	254	274
Cement Plant 3	NO	NO	802	1,010	1,015	1,115	1,204	1,228	1,227	1,214	1,010	790	745	545	615	407
Cement Plant 4	NO	NO	NO	NO	NO	381	411	547	608	609	557	441	367	413	466	479
Total	1,610	1,601	3,109	3,386	3,413	3,967	4,283	4,400	4,400	4,441	3,893	2,438	2,053	1,805	2,189	2,065
Emission Factor t CO₂/t Clinker Produced																
Cement Plant 1	0.546	0.546	0.546	0.546	0.542	0.534	0.504	0.536	0.537	0.534	0.536	0.537	0.533	0.534	0.537	0.535
Cement Plant 2	0.553	0.553	0.553	0.553	0.553	0.538	0.518	0.533	0.535	0.536	0.544	0.542	0.534	0.536	0.533	0.532
Cement Plant 3	NA	NA	0.542	0.542	0.542	0.542	0.544	0.536	0.531	0.537	0.550	0.558	0.552	0.546	0.546	0.542
Cement Plant 4	NA	NA	NA	NA	NA	0.525	0.686	0.540	0.528	0.529	0.535	0.533	0.523	0.523	0.530	0.546
IEF t CO₂/t Clinker	0.549	0.549	0.547	0.547	0.545	0.536	0.536	0.536	0.534	0.535	0.541	0.544	0.538	0.535	0.538	0.538
Emissions CO₂ (kilotonnes)																
Cement Plant 1	505	503	764	798	810	836	853	894	894	900	763	379	309	301	458	483
Cement Plant 2	379	376	502	506	499	487	506	510	481	500	491	272	193	152	136	146
Cement Plant 3	NO	NO	435	547	550	604	654	658	651	652	555	441	411	297	336	221
Cement Plant 4	NO	NO	NO	NO	NO	200	282	295	321	322	298	235	192	216	247	262
Total	884	879	1,701	1,851	1,860	2,127	2,295	2,357	2,348	2,374	2,107	1,327	1,105	966	1,177	1,112

Table 3.2.B Lime Production 1990-2013

Activity data, emission factors and emissions

IPCC Sector 2A2 Lime	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Lime production (kilotonnes)																
Lime Plant 1	84.4	97.9	90.5	97.5	102.8	112.8	116.0	100.9	102.2	116.6	125.9	94.7	114.1	119.7	132.9	127.9
Lime Plant 2	170.8	125.6	157.8	146.0	139.6	126.9	130.1	135.7	129.0	136.5	121.3	111.0	138.5	143.0	149.5	123.5
Total	255.2	223.5	248.3	243.5	242.4	239.7	246.1	236.7	231.2	253.0	247.1	205.7	252.6	262.7	282.4	251.4
Emission Factor t CO₂/t Lime Produced																
Lime Plant 1	0.757	0.827	0.801	0.826	0.808	0.813	0.808	0.791	0.791	0.818	0.773	0.764	0.761	0.767	0.761	0.752
Lime Plant 2	0.879	0.849	0.747	0.746	0.769	0.903	0.829	0.764	0.771	0.760	0.759	0.764	0.769	0.760	0.767	0.756
IEF t CO₂/t Lime	0.839	0.839	0.767	0.778	0.785	0.861	0.819	0.775	0.780	0.787	0.766	0.764	0.765	0.763	0.764	0.754
Emissions CO₂ (kilotonnes)																
Lime Plant 1	63.9	80.9	72.5	80.5	83.0	91.7	93.7	79.8	80.8	95.3	97.3	72.4	86.9	91.8	101.2	96.2
Lime Plant 2	150.2	106.6	117.9	108.9	107.3	114.6	107.8	103.6	99.5	103.7	92.1	84.9	106.5	108.7	114.7	93.4
Total	214.1	187.5	190.4	189.4	190.3	206.3	201.5	183.5	180.3	199.1	189.3	157.2	193.4	200.5	215.9	189.6

Table 3.2.C Glass Production 1990-2013

Activity data, emission factors and emissions

IPCC Sector 2A3 Glass production	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Carbonate use (kilotonnes)																
Glass plant 1	0.412	0.412	0.412	0.412	0.363	0.438	0.438	0.328	NO	NO	NO	NO	NO	NO	NO	NO
Glass plant 2	1.720	1.549	1.273	0.537	0.440	0.440	0.581	0.472	0.701	0.600	0.422	0.063	NO	NO	NO	NO
Glass bottle	60.000	60.000	60.000	60.000	30.000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Glass wool	1.746	1.746	2.057	1.809	1.161	0.734	0.709	0.628	0.708	0.699	0.461	NO	NO	NO	NO	NO
Total	63.878	63.707	63.742	62.758	31.964	1.612	1.727	1.428	1.409	1.299	0.882	0.063	NO	NO	NO	NO
Emission Factor t CO₂/t Carbonate Use																
Glass plant 1	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	NA	NA	NA	NA	NA	NA	NA	NA
Glass plant 2	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	0.275	NA	NA	NA	NA
Glass bottle	0.200	0.178	0.156	0.151	0.147	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Glass wool	0.441	0.441	0.441	0.438	0.431	0.425	0.423	0.415	0.415	0.415	0.415	NA	NA	NA	NA	NA
IEF t CO₂/t Carbonate Use	0.209	0.188	0.168	0.162	0.161	0.343	0.336	0.337	0.345	0.350	0.348	0.275	NA	NA	NA	NA
Emissions CO₂ (kilotonnes)																
Glass plant 1	0.113	0.113	0.113	0.113	0.100	0.120	0.120	0.090	NO	NO	NO	NO	NO	NO	NO	NO
Glass plant 2	0.473	0.426	0.350	0.148	0.121	0.121	0.160	0.130	0.193	0.165	0.116	0.017	NO	NO	NO	NO
Glass bottle	11.970	10.658	9.345	9.083	4.410	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Glass wool	0.769	0.769	0.906	0.793	0.500	0.312	0.300	0.261	0.294	0.290	0.191	NO	NO	NO	NO	NO
Total	13.325	11.966	10.714	10.136	5.131	0.553	0.580	0.481	0.487	0.455	0.307	0.017	NO	NO	NO	NO

Table 3.2.D Bricks, Ceramics, Limestone 1990-2013

Activity data, emission factors and emissions

IPCC Sector 2A4 Other process uses of carbonates	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Carbonate uses (includes clays, shale, bricks, tiles, flues and limestone) (kilotonnes)																
2A4a Ceramics:	110.73	118.73	140.20	138.84	128.07	133.38	133.50	158.60	152.13	137.45	83.21	16.35	15.83	22.54	0.75	0.62
Ceramics Plant 1	30.75	30.75	30.75	30.75	32.64	35.59	37.17	41.21	34.32	33.81	16.52	2.73	0.17	0.57	0.75	0.62
Ceramics Plant 2	40.00	40.00	43.14	44.54	45.90	46.12	47.58	46.06	47.71	45.06	26.22	13.62	15.66	21.96	NO	NO
Ceramics Plant 3	39.98	47.98	66.30	63.55	49.54	51.67	48.75	52.51	48.85	38.83	21.27	NO	NO	NO	NO	NO
Ceramics Plant 4	NO	NO	NO	NO	NO	NO	NO	18.83	21.25	19.74	19.19	NO	NO	NO	NO	NO
2A4d Limestone use:	NO	NO	NO	9.93	4.67	5.13	7.75	9.54	5.46	4.84	5.77	3.55	2.40	2.41	1.01	0.49
Power plant 1	NO	NO	NO	9.93	4.67	5.13	7.75	8.32	5.28	4.76	4.81	2.20	2.32	1.88	0.82	0.33
Power plant 2	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.09	0.96	1.35	0.08	0.53	0.19	0.16
Sugar processing	NE	NE	NE	NE	NE	NE	NE	1.22	0.18	NO	NO	NO	NO	NO	NO	NO
Total	110.73	118.73	140.20	148.77	132.74	138.52	141.25	168.14	157.59	142.29	88.98	19.90	18.22	24.95	1.76	1.11
Emission Factor t CO2/t Raw Material Use																
2A4a Ceramics:	0.047	0.048	0.048	0.045	0.047	0.047	0.048	0.048	0.050	0.051	0.050	0.033	0.027	0.037	0.041	0.053
Ceramics Plant 1	0.045	0.045	0.045	0.036	0.046	0.045	0.048	0.046	0.048	0.048	0.048	0.055	0.056	0.054	0.041	0.053
Ceramics Plant 2	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.034	0.028	0.026	0.036	NA	NA
Ceramics Plant 3	0.052	0.052	0.051	0.051	0.051	0.051	0.051	0.051	0.053	0.053	0.056	NA	NA	NA	NA	NA
Ceramics Plant 4	NA	NA	NA	NA	NA	NA	NA	0.051	0.062	0.069	0.068	NA	NA	NA	NA	NA
2A4d Limestone use:	NA	NA	NA	0.436	0.436	0.436	0.441	0.437	0.436	0.436	0.437	0.434	0.430	0.433	0.432	0.436
Power plant 1	NA	NA	NA	0.436	0.436	0.436	0.441	0.436	0.436	0.436	0.436	0.430	0.430	0.431	0.431	0.436
Power plant 2	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.440	0.440	0.440	0.440	0.440	0.440	0.437
Sugar processing	NA	NA	NA	NA	NA	NA	NA	0.440	0.438	NA	NA	NA	NA	NA	NA	NA
Emissions CO2 (kilotonnes)																
2A4a Ceramics:	5.226	5.64	6.664	6.296	6.045	6.256	6.361	7.534	7.658	7.039	4.184	0.531	0.42	0.826	0.031	0.033
Ceramics Plant 1	1.3935	1.3935	1.3935	1.104	1.506	1.598	1.79	1.883	1.6548	1.62981	0.8009	0.1515	0.00942	0.0309	0.0307	0.0331
Ceramics Plant 2	1.7597	1.7597	1.898	1.9595	2.0192	2.0291	2.0932	2.0263	2.099	1.98285	0.8916	0.3799	0.411	0.7951	NO	NO
Ceramics Plant 3	2.0725	2.487	3.3728	3.2328	2.5199	2.6284	2.478	2.656	2.585	2.0542	1.1866	NO	NO	NO	NO	NO
Ceramics Plant 4	NO	NO	NO	NO	NO	NO	NO	0.9689	1.3193	1.372	1.3048	NO	NO	NO	NO	NO
2A4d Limestone use:	NO	NO	NO	4.33	2.04	2.24	3.42	4.17	2.38	2.11	2.52	1.54	1.03	1.04	0.44	0.21
Power plant 1	NO	NO	NO	4.33	2.04	2.24	3.42	3.63	2.30	2.07	2.10	0.94	1.00	0.81	0.36	0.15
Power plant 2	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.04	0.42	0.60	0.04	0.23	0.08	0.07
Sugar processing	NE	NE	NE	NE	NE	NE	NE	0.54	0.08	NO	NO	NO	NO	NO	NO	NO
Total	5.23	5.64	6.66	10.63	8.08	8.49	9.78	11.70	10.04	9.15	6.70	2.07	1.45	1.87	0.47	0.25

Table 3.2.E Soda ash use 1990-2013

Activity data, emission factors and emissions

IPCC Sector 2A4 Soda ash use	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Soda ash used (kilotonnes)																
Lime Plant 2	0.237	0.168	0.171	0.216	0.132	0.154	0.198	0.202	0.150	0.136	0.106	0.132	0.177	0.166	0.211	0.153
Emission Factor t CO₂/t Soda ash used																
IEF t CO₂/t Soda ash use	0.411	0.411	0.411	0.411	0.411	0.411	0.411	0.411	0.413	0.411	0.411	0.411	0.411	0.411	0.415	0.415
Emissions CO₂ (kilotonnes)																
Lime Plant 2	0.097	0.069	0.070	0.089	0.054	0.063	0.081	0.083	0.062	0.056	0.044	0.054	0.073	0.068	0.087	0.063

Annex 3.3

Agriculture (IPCC Sector 3)

3.3.A Animal Populations

3.3.B Methane Emission Factors for Enteric Fermentation

3.3.C Methane Emission Factors for Manure Management

3.3.D.1 Allocation of Animal Wastes to Manure Management Systems – Cattle

3.3.D.2 Allocation of Animal Wastes to Manure Management Systems – Other Livestock

3.3.E Nitrogen excretion values for Livestock 1990-2013

3.3.F Input Parameters for the calculation of N₂O Emissions from Agricultural Soils

3.3.G Nitrogen application to agricultural soils from sewage sludge (3.D.1.2.b) 1990-2013

3.3.H Activity data, parameters and emission factors for Crop Residue (3.D.1.4) 1990-2013

Table 3.3.A Animal Populations 1990-2013

1000 head	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Total Cattle	6822	6921	6973	6959	6965	7009	7282	7491	7592	7348	7012	7022	6961	6971	6973	6951	6925	6827	6828	6813	6543	6428	6691	6835
Dairy Cows	1341	1309	1262	1256	1247	1239	1241	1227	1216	1187	1165	1165	1146	1146	1139	1025	1054	1054	1060	1060	1039	1076	1101	1123
All Other Cattle	5481	5612	5711	5703	5718	5770	6041	6264	6376	6160	5847	5857	5814	5826	5834	5926	5872	5773	5768	5753	5505	5352	5590	5712
Other Cows	730	801	903	958	990	1022	1098	1183	1222	1192	1171	1178	1153	1166	1179	1121	1171	1185	1198	1169	1125	1103	1138	1118
Dairy Heifers	172	156	187	191	207	230	238	244	226	212	205	202	223	221	234	214	204	197	195	196	234	252	268	271
Other Heifers	80	71	106	116	104	123	134	149	128	121	133	140	142	139	142	191	193	212	180	156	170	202	181	155
Cattle < 1 yrs	1716	1765	1695	1738	1736	1746	1852	1938	1965	1821	1752	1824	1799	1761	1771	1962	1953	1941	1959	1889	1761	1846	2036	1969
Cattle < 1 yrs - male	903	919	889	914	904	915	974	1023	1055	965	919	955	953	922	930	958	951	947	969	918	827	892	1023	959
Cattle < 1 yrs - female	813	846	806	824	832	831	878	915	910	856	833	869	846	839	842	1005	1002	994	990	971	935	954	1013	1009
Cattle 1 - 2 yrs	1663	1692	1638	1587	1586	1586	1639	1717	1783	1706	1517	1515	1593	1577	1535	1642	1506	1466	1496	1542	1407	1270	1376	1551
Cattle 1 - 2 yrs - male	986	981	982	958	952	964	996	1055	1086	1039	912	913	992	983	950	972	845	818	832	851	760	673	770	873
Cattle 1 - 2 yrs - female	677	711	656	630	634	622	643	662	697	667	605	602	601	594	585	670	661	648	664	690	647	597	606	679
Cattle > 2 yrs	1093	1099	1152	1078	1058	1023	1036	986	1002	1058	1016	941	845	902	911	734	782	715	687	750	760	640	554	609
Cattle > 2 yrs - male	826	798	830	773	740	712	732	690	708	737	722	642	560	599	605	537	565	510	476	501	506	426	361	388
Cattle > 2 yrs - female	266	301	322	305	318	311	304	296	294	321	295	299	284	303	305	197	217	206	211	249	254	214	193	221
Bulls	27	29	32	36	38	40	44	48	50	51	53	56	59	60	63	61	63	57	54	52	47	38	37	39
Total Sheep	8021	8484	8736	8977	8559	8364	8329	8051	8572	8547	7957	7455	6682	6481	6703	6431	6187	5656	5105	4727	4328	4429	4843	4917
Ewes Lowland	2397	2543	2622	2576	2511	2427	2369	2390	3056	2936	2814	2704	2637	2552	2464	2627	2414	2207	2057	1928	1920	1954	2036	2015
Ewes Upland	1961	2080	2145	2108	2055	1986	1938	1955	1310	1258	1206	1159	1130	1094	1056	657	604	552	514	482	480	489	509	504
Rams Lowland	64	67	70	69	67	66	62	64	81	79	77	75	73	72	70	77	74	69	63	58	59	59	61	62
Rams Upland	53	55	57	56	55	54	51	52	35	34	33	32	31	31	30	19	19	17	16	14	15	15	15	15
Other Sheep>1 - Lowland	164	96	89	99	107	113	106	118	172	153	143	128	129	144	140	124	122	109	112	103	96	101	116	112
Other Sheep>1 - Upland	134	79	73	81	88	92	86	97	74	66	61	55	55	62	60	31	31	27	28	26	24	25	29	28
Lambs - Lowland	1787	1960	2024	2194	2022	1994	2044	1856	2692	2815	2535	2312	1838	1768	2019	2317	2339	2140	1853	1693	1387	1429	1661	1745
Lambs - Upland	1462	1604	1656	1795	1654	1632	1672	1519	1154	1206	1086	991	788	758	865	579	585	535	463	423	347	357	415	436
Total Pigs	1222	1325	1404	1504	1514	1546	1643	1708	1810	1775	1727	1760	1791	1729	1704	1679	1632	1544	1486	1444	1508	1551	1532	1510
Gilts in Pig	21	22	25	23	22	24	25	27	26	25	21	23	20	20	22	20	22	21	21	20	19	19	20	19
Gilts not yet Served	12	14	15	14	15	18	17	18	19	16	18	19	20	18	19	20	19	16	16	17	15	15	15	15
Sows in Pig	83	90	96	101	99	100	103	108	109	109	110	107	110	104	102	100	96	96	91	89	92	90	84	82
Other Sows for Breeding	31	31	33	33	30	31	36	37	38	38	32	37	33	32	30	34	31	28	25	27	29	27	25	29
Boars	6	7	7	6	6	5	5	5	5	4	4	4	3	3	3	2	2	2	2	2	2	1	1	1
Pigs 20 Kg +	749	803	837	905	918	952	1016	1064	1144	1094	1038	1036	1062	1043	1028	1010	1034	939	932	911	953	965	960	926
Pigs Under 20 Kg	319	358	392	422	425	417	442	450	469	489	504	535	543	508	500	494	429	443	400	378	400	434	426	438
Total Poultry	11772	12698	13272	13072	14034	14438	15375	15548	15686	15490	15680	16024	15545	16152	17190	16573	15934	13324	13258	15277	15212	14947	15631	15005
Layer	1868	1800	2231	1832	1730	1371	1701	1580	1559	1537	1572	1676	1613	1907	1906	1950	1970	1813	1813	2145	2145	2060	2600	2828
Broiler	8035	8905	9067	9522	10393	11092	11730	12096	12287	12200	12426	12629	12322	12672	13375	12818	12360	9696	9696	11904	11904	11520	11520	10764
Turkey	1509	1633	1615	1358	1552	1616	1585	1513	1482	1393	1322	1358	1248	1209	1461	1274	1097	1330	1330	874	874	1078	1222	1125
Ducks	347	347	347	347	347	347	347	347	347	347	347	347	348	349	350	435	520	497	475	409	344	279	279	279
Geese	12	12	12	12	12	12	12	12	12	12	12	12	13	13	13	12	11	10	10	10	10	10	10	10
Horses	62	63	65	66	67	68	70	72	73	76	70	71	73	70	73	80	87	89	96	98	106	106	111	102
Mules	8	7	8	9	8	7	8	7	8	7	5	5	5	6	6	6	7	7	9	9	8	9	10	8
Goats	17	17	18	18	16	16	15	15	15	14	8	8	8	8	8	7	7	7	9	10	11	11	10	9
Farmed Deer	12	12	13	15	15	16	16	18	17	16	12	12	12	11	11	10	9	10	10	9	5	3	2	2
Mink	185	143	130	124	124	124	133	143	143	143	146	146	146	146	146	149	149	149	149	190	183	183	198	198
Fox	26	22	13	9	7	7	8	8	9	9	4	4	4	4	4	2	2	3	1	0	0	NO	NO	NO
Fertiliser (1000's tonnes/N)	379.3	370.1	358.3	378.0	404.8	428.8	416.9	380.4	432.0	442.9	407.6	368.7	363.5	388.1	362.5	352.2	342.1	321.6	309.0	306.8	362.4	295.8	296.5	353.0

Table 3.3.B CH₄ Emission Factors for Enteric Fermentation (kg/head/year)

Animal Category	Animal Liveweight (kg)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Cattle																									
Dairy cows	535.0	101.4	101.9	102.5	103.0	103.6	104.1	104.6	105.2	105.7	106.3	106.8	107.4	107.9	108.5	108.4	111.3	111.4	111.3	110.0	108.5	113.1	113.2	110.8	111.6
Beef cows (Suckler Cows)	500.0	74.0	74.0	74.1	74.1	74.1	74.1	74.1	74.1	74.1	74.1	74.2	74.2	74.2	74.2	74.5	75.5	74.3	73.2	74.9	72.8	72.9	74.1	75.5	73.1
Dairy heifers	388.0	51.8	51.7	51.6	51.4	51.3	51.2	51.1	50.9	50.8	50.7	50.5	50.4	50.3	50.2	50.2	50.2	50.2	50.2	50.2	50.2	50.2	50.2	50.2	50.2
Beef heifers	450.0	55.4	55.3	55.2	55.0	54.9	54.8	54.6	54.5	54.4	54.2	54.1	53.9	53.8	53.7	53.7	53.7	53.7	53.7	53.7	53.7	53.7	53.7	53.7	53.7
Bulls for breeding	500.0	86.4	86.0	85.6	85.3	84.9	84.5	84.2	83.8	83.4	83.0	82.7	82.3	81.9	81.5	81.5	81.5	81.5	81.5	81.5	81.5	81.5	81.5	81.5	81.5
Cattle 1 - 2 yrs - male																									
< 1 year	140.0	30.5	30.4	30.3	30.2	30.2	30.1	30.0	29.9	29.9	29.8	29.7	29.7	29.6	29.5	29.7	29.7	29.6	29.7	29.7	29.8	29.1	29.8	30.1	30.3
1 - 2 years	388.0	62.2	62.1	62.0	61.8	61.7	61.6	61.4	61.3	61.2	61.0	60.9	60.8	60.6	60.5	59.3	58.9	59.9	59.2	59.1	58.6	60.0	58.0	56.6	56.2
> 2 years*	500.0	55.1	53.5	51.9	50.2	48.6	47.0	45.4	43.8	42.2	40.6	38.9	37.3	35.7	34.1	35.2	37.7	37.8	38.6	37.0	38.8	39.8	38.3	37.2	37.3
Cattle > 2 yrs - female																									
< 1 year	140.0	27.0	27.1	27.2	27.2	27.3	27.3	27.4	27.5	27.5	27.6	27.6	27.7	27.7	27.8	27.9	27.7	27.7	27.6	27.6	27.6	27.5	27.6	27.7	27.7
1 - 2 years	388.0	53.5	52.9	52.2	51.5	50.8	50.1	49.4	48.7	48.0	47.4	46.7	46.0	45.3	44.6	44.5	45.6	46.4	46.6	47.0	47.7	48.6	47.9	48.0	48.1
> 2 years*	500.0	21.7	21.7	21.8	21.8	21.9	22.0	22.0	22.1	22.1	22.2	22.3	22.3	22.4	22.5	22.5	22.4	22.4	22.4	22.5	22.6	22.6	22.7	22.7	22.6
Sheep																									
Lowland Sheep		8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0
Upland Sheep		8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0
Rams		8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0
Sheep > 1 yrs		8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0	8.0
Lambs		2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7
Horses		18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0
Mules		10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
Goats		5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Deer		20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0
Pigs																									
Gilts in Pig	160.0	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
Gilts not yet Served	120.0	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Sows in Pig	200.0	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Other Sows for Breeding	210.0	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
Boars	225.0	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7
Pigs > 20 Kg	58.0	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7
Pigs < 20 Kg	13.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Poultry	2.4	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

* Note: This value is low because this category of animal only live part of their third year

Table 3.3.C CH₄ Emission Factors for Manure Management (kg/head/year)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Cattle																								
Dairy cows	10.6	10.5	10.5	10.5	10.4	10.4	10.4	10.3	10.3	10.3	10.2	10.2	10.1	10.1	10.1	10.3	10.3	10.2	10.2	10.1	10.3	10.3	10.2	10.2
Beef cows(Suckler Cows)	6.6	6.6	6.6	6.6	6.6	6.6	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.6	6.5	6.4	6.6	6.4	6.4	6.5	6.7	6.4
Dairy heifers	4.6	4.5	4.5	4.4	4.3	4.3	4.2	4.2	4.1	4.0	4.0	3.9	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8
Beef heifers	5.3	5.3	5.2	5.1	5.0	5.0	4.9	4.8	4.8	4.7	4.6	4.6	4.5	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4	4.4
Bulls for breeding	10.5	10.3	10.2	10.0	9.8	9.7	9.5	9.3	9.2	9.0	8.9	8.7	8.5	8.4	8.4	8.4	8.4	8.4	8.4	8.4	8.4	8.4	8.4	8.4
Male cattle																								
< 1 year	4.5	4.4	4.4	4.3	4.3	4.3	4.2	4.2	4.1	4.1	4.0	4.0	4.0	3.9	4.0	4.0	3.9	4.0	4.0	4.0	4.1	4.0	4.1	4.1
1 - 2 years	7.1	7.0	6.9	6.8	6.8	6.7	6.6	6.5	6.4	6.4	6.3	6.2	6.1	6.0	5.9	5.9	6.0	5.9	5.9	5.9	6.0	5.7	5.5	5.4
> 2 years*	3.0	2.8	2.7	2.5	2.3	2.2	2.0	1.9	1.7	1.6	1.4	1.3	1.1	1.0	1.1	1.3	1.2	1.3	1.2	1.4	1.4	1.3	1.2	1.2
Female cattle																								
< 1 year	4.0	4.0	4.0	4.0	4.0	4.0	3.9	3.9	3.9	3.9	3.9	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8
1 - 2 years	6.3	6.1	5.9	5.7	5.6	5.4	5.2	5.1	4.9	4.7	4.5	4.4	4.2	4.0	3.9	4.1	4.2	4.2	4.3	4.4	4.6	4.4	4.4	4.5
> 2 years*	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Sheep																								
Lowland Ewes	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Upland Ewes	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Rams	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Sheep >1 yrs	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Lambs	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Horses	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
Mules	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Goats	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Deer	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Mink	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Fox	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Pigs																								
Gilts in Pig	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0
Gilts not yet Served	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1
Sows in Pig	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0	8.0	8.0	8.1	8.0
Other Sows for Breeding	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8	18.8
Boars	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4	9.4
Pigs > 20 Kg	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1
Pigs < 20 Kg	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4
Poultry																								
Layers	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Broilers	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Turkeys	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Ducks	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.6	0.6	0.6
Geese	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1

* Note: This value is low because this category of animal only live part of their third year.

Table 3.3.D.1 Allocation of Animal Wastes to Animal Waste Management Systems – Cattle

Cattle	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Number of days housed																								
Dairy Cows	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	118	117	117	117	117	117	117	117
Suckler Cows	141	141	141	141	141	141	141	141	141	141	141	141	141	141	141	142	141	141	141	141	141	141	142	141
Dairy Heifer	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128	128
Other Heifer	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139	139
Under1yr	221	221	221	221	221	221	221	221	221	221	221	221	221	221	221	222	222	222	223	223	227	224	224	223
Oneto2yrs	156	156	156	156	156	156	156	156	156	156	156	156	156	156	154	155	154	154	154	156	157	154	153	153
Over2yrs	23	23	23	23	23	23	23	23	23	23	23	23	23	23	25	26	24	26	23	26	26	23	21	22
Bulls	156	156	156	156	156	156	156	156	156	156	156	156	156	156	154	155	154	154	154	156	157	154	153	153
Number of days grazing																								
Dairy Cows	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	247	248	248	248	248	248	248	248
Suckler Cows	224	224	224	224	224	224	224	224	224	224	224	224	224	224	224	223	224	224	224	224	224	224	223	224
Dairy Heifer	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237	237
Other Heifer	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226	226
Under1yr	144	144	144	144	144	144	144	144	144	144	144	144	144	144	144	143	143	143	142	142	138	141	141	142
Oneto2yrs	209	209	209	209	209	209	209	209	209	209	209	209	209	209	211	210	211	211	211	209	208	211	212	212
Over2yrs	342	342	342	342	342	342	342	342	342	342	342	342	342	342	340	339	341	339	342	339	339	342	344	343
Bulls	209	209	209	209	209	209	209	209	209	209	209	209	209	209	211	210	211	211	211	209	208	211	212	212
Proportion to each AWMS																								
Pit Storage																								
Dairy Cows	0.32	0.32	0.32	0.32	0.31	0.31	0.31	0.31	0.31	0.30	0.30	0.30	0.30	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29	0.29
Suckler Cows	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.27	0.26	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27
Dairy Heifer	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
Other Heifer	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38
Under1yr	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.42	0.41	0.42	0.42	0.42	0.42	0.42	0.42
Oneto2yrs	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.33	0.34	0.34	0.34	0.34	0.33	0.33	0.33
Over2yrs	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.03
Bulls	0.42	0.42	0.42	0.41	0.41	0.41	0.40	0.40	0.40	0.39	0.39	0.39	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38
Deep Bedding																								
Dairy Cows	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Suckler Cows	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Dairy Heifer	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Other Heifer	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Under1yr	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.18	0.20	0.20	0.20	0.20	0.20	0.20	0.20
Oneto2yrs	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09
Over2yrs	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02
Bulls	0.14	0.14	0.14	0.14	0.14	0.14	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
Pasture																								
Dairy Cows	0.66	0.66	0.66	0.66	0.67	0.67	0.67	0.67	0.68	0.68	0.68	0.68	0.69	0.69	0.69	0.69	0.69	0.69	0.69	0.69	0.70	0.70	0.69	0.69
Suckler Cows	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.64	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63
Dairy Heifer	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65
Other Heifer	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62
Under1yr	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.38	0.39	0.39	0.39
Oneto2yrs	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.58	0.58	0.58	0.58	0.58	0.57	0.57	0.58	0.58	0.58
Over2yrs	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.94	0.93	0.93	0.93	0.93	0.94	0.93	0.93	0.94	0.94	0.94
Bulls	0.44	0.44	0.45	0.45	0.45	0.46	0.46	0.47	0.47	0.48	0.48	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49	0.49

Table 3.3.D.2 Allocation of Animal Wastes to Animal Waste Management Systems – Other Livestock

Animal Category	Days housed	% housed	% outwintered	Manure Management System											
				Liquid system		Solid storage & dry lot		Pit-storage		Deep bedding		Litter		Pasture	
				portion	MCF	portion	MCF	portion	MCF	portion	MCF	portion	MCF	portion	MCF
Sheep															
Lowland Ewes	61.00	47.07	52.93	NA	NA	NA	NA	NA	NA	0.08	17.0%	NA	NA	0.92	1.0%
Upland Ewes	85.00	44.34	55.66	NA	NA	NA	NA	NA	NA	0.10	17.0%	NA	NA	0.90	1.0%
Lowland Rams	85.00	22.34	77.66	NA	NA	NA	NA	NA	NA	0.05	17.0%	NA	NA	0.95	1.0%
Upland Rams	85.00	22.34	77.66	NA	NA	NA	NA	NA	NA	0.05	17.0%	NA	NA	0.95	1.0%
Lowland Other Sheep>1yrs	61.00	47.07	52.93	NA	NA	NA	NA	NA	NA	0.08	17.0%	NA	NA	0.92	1.0%
Upland Other Sheep>1yrs	85.00	44.34	55.66	NA	NA	NA	NA	NA	NA	0.08	17.0%	NA	NA	0.92	1.0%
Lowland lambs	58.00	16.88	83.12	NA	NA	NA	NA	NA	NA	0.03	17.0%	NA	NA	0.97	1.0%
Upland lambs	58.00	16.88	83.12	NA	NA	NA	NA	NA	NA	0.03	17.0%	NA	NA	0.97	1.0%
Pigs															
Gilts in pig	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Gilts not yet served	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Sows in pig	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Other sows for breeding	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Boars	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Pigs < 20 kg	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Pigs > 20 kg	365.00	100.00	0.00	NA	NA	NA	NA	1.00	17.0%	NA	NA	NA	NA	NA	NA
Poultry															
Laying hens	365.00	88.00	12.00	NA	NA	NA	NA	0.74	17.0%	NA	NA	0.14	1.5%	0.12	1.0%
Broilers	365.00	100.00	0.00	NA	NA	NA	NA	NA	NA	NA	NA	1.00	1.5%	NA	NA
Turkeys	365.00	100.00	0.00	NA	NA	NA	NA	NA	NA	NA	NA	1.00	1.5%	NA	NA
Ducks	365.00	100.00	0.00	NA	NA	NA	NA	0.54	17.0%	NA	NA	0.46	1.5%	NA	NA
Geese	365.00	100.00	0.00	NA	NA	NA	NA	NA	17.0%	NA	NA	1.00	1.5%	NA	NA
Horses	143.00	100.00	0.00	NA	NA	0.39	2.0%	NA	NA	NA	NA	NA	NA	0.61	1.0%
Mule	143.00	100.00	0.00	NA	NA	0.39	2.0%	NA	NA	NA	NA	NA	NA	0.61	1.0%
Goat	61.00	100.00	0.00	NA	NA	NA	NA	NA	NA	0.58	17.0%	NA	NA	0.42	1.0%
Deer	61.00	100.00	0.00	NA	NA	NA	NA	NA	NA	0.17	17.0%	NA	NA	0.83	1.0%
Fur animals	365.00	100.00	0.00	0.40	17.0%	0.48	2.0%	0.12	17.0%	NA	NA	NA	NA	NA	NA

Table 3.3.E Nitrogen excretion values for Livestock 1990-2013

N excretion (kg/head/year)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
	N excretion values																							
Dairy Cows	95.5	95.9	96.3	96.6	97.0	97.3	97.7	98.0	98.4	98.7	99.1	99.5	99.8	100.2	100.0	102.1	102.0	101.7	100.7	99.7	102.6	102.6	101.1	101.2
Suckler Cows	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8
Dairy Heifer	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4
Other Heifer	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4	74.4
Under1yr	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6	27.6
One to 2yrs	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4
Over 2yrs	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2	37.2
Bulls	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8	73.8
Ewes Lowland	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3	12.3
Ewes Upland	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5
Rams - lowland	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
Rams - upland	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
Other Sheep>1 - lowland	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5
Other Sheep>1 - upland	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Lambs - lowland	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
Lambs - upland	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Gilts in pig	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0
Gilts not yet served	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2
Sows in pig	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0
Other breeding sows	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0	20.0
Boars	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0
Fatteners > 20 kg	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2	9.2
Fatteners < 20 kg	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
Laying hen per bird place	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Broiler per bird place	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Turkey per bird place	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Ducks	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Geese	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Horses	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4	48.4
Mules	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0
Goats	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9	12.9
Deer (red) 6 months - 2 years	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3
Deer (red) > 2 years	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5	27.5
Deer (fallow) 6 months-2 years	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7
Deer (fallow) > 2 years	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3	14.3
Deer (sika) 6 months - 2 years	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1
Deer (sika) > 2 years	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0
Mink	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6
Fox	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1	12.1

Table 3.3.F Input Parameters for the calculation of N₂O Emissions from Agricultural Soils

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
FracGASF	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.02	0.02
FracGRAZ	0.63	0.63	0.64	0.63	0.63	0.63	0.63	0.62	0.63	0.63	0.63	0.63	0.62	0.63	0.63	0.62	0.62	0.62	0.61	0.61	0.61	0.62	0.61	0.61
FracGASM1	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
FracGASM2	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
FracLEACH	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
FSN (tonnes/year)	379311	370121	358302	377985	404811	428826	416918	380350	431999	442916	407598	368667	363513	388080	362525	352165	342137	321553	308960	306806	362395	295795	296536	353044
FAM (tonnes/year)	127756	130109	131147	131790	132465	134025	139426	144508	146745	141046	134461	135603	135955	135114	135012	136646	134587	132707	132819	132310	127998	126184	132252	133947
FS (tonnes/year)	103	103	104	105	105	97	91	212	365	543	765	934	1127	1299	1688	1819	1690	1826	2426	2616	1939	1949	1941	1729
FCR (tonnes/year)	79897	79806	81785	66482	72504	78082	84532	81973	76465	79249	81125	72300	65211	63389	75616	58138	29850	28764	35179	30148	29677	34881	29395	32700
FPRP (tonnes/year)	294636	299505	305512	304830	304365	305314	313247	320101	328076	320391	307197	304509	301184	301833	300973	292780	291163	284776	281378	278683	272889	270077	276483	283462

Table 3.3.G Nitrogen application to agricultural soils from sewage sludge (3.D.1.2.B) 1990-2013

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
N applied (t/year)	102.73	103.31	104.16	104.73	105.08	96.95	90.53	212.32	364.97	543.10	764.53	933.83	1127.09	1298.57	1687.61	1819.00	1690.15	1826.39	2425.89	2615.83	1938.74	1949.09	1940.68	1728.67

Table 3.3.H Activity data, parameters and emission factors for Crop Residue (3.D.1.4) 1990-2013

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Crop T	kg d.m. ha ⁻¹																							
Maize	0	0	0	0	0	0	0	0	0	0	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071	13,071
Winter Wheat	8,010	7,298	7,120	5,963	7,209	7,743	8,366	7,120	7,209	8,366	8,811	8,722	7,832	7,743	9,345	7,832	8,722	7,743	8,544	7,654	7,921	9,078	6,586	8,277
Spring Wheat	5,963	5,963	6,586	6,319	5,696	6,230	6,675	6,052	6,853	6,942	7,209	7,120	6,408	6,764	7,654	6,764	6,942	6,853	5,874	6,052	6,764	7,387	5,429	7,209
Oats	5,874	6,230	6,052	5,696	5,429	5,785	6,230	5,696	5,429	5,963	6,675	6,319	6,319	6,586	6,942	5,963	6,230	6,675	6,764	6,319	6,675	7,031	5,874	6,408
Barley	5,251	5,251	5,607	4,717	4,806	5,429	6,052	5,073	4,984	5,963	6,408	6,230	4,895	5,785	6,497	5,518	6,052	5,963	6,141	5,607	6,230	6,942	5,785	6,764
Beans and Peas	4,459	4,277	4,550	3,913	4,095	3,640	4,095	3,640	4,550	4,095	4,641	4,277	4,095	4,368	4,914	4,641	4,277	4,004	4,095	4,823	5,005	5,187	4,368	4,732
Potatoes	5,478	6,138	6,358	5,786	6,622	6,072	6,644	5,698	5,742	7,040	7,392	7,370	7,414	7,590	9,108	7,634	7,348	7,480	6,754	6,072	7,568	7,568	5,676	8,404
Turnips	51,512	49,350	52,170	49,162	49,914	50,854	54,332	51,700	42,582	39,480	56,400	51,136	51,136	0	0	0	0	0	0	0	0	0	0	0
Sugarbeet	43,052	39,762	41,924	32,618	36,942	41,454	42,958	47,940	46,812	47,658	53,392	45,308	39,104	44,838	56,212	41,830	0	0	0	0	0	0	0	0
Fodder Beat	61,100	64,766	66,364	51,606	55,460	62,792	69,278	62,228	60,912	66,176	66,082	61,570	57,246	0	0	0	0	0	0	0	0	0	0	0
R _{actm}	kg d.m																							
Maize	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08
Winter Wheat	1.66	1.66	1.67	1.68	1.67	1.66	1.66	1.67	1.67	1.66	1.66	1.66	1.66	1.66	1.65	1.66	1.66	1.66	1.66	1.66	1.66	1.65	1.67	1.66
Spring Wheat	1.42	1.42	1.40	1.41	1.42	1.41	1.40	1.41	1.40	1.40	1.39	1.40	1.41	1.40	1.39	1.40	1.40	1.40	1.40	1.42	1.41	1.40	1.39	1.43
Oats	1.06	1.05	1.06	1.07	1.07	1.06	1.05	1.07	1.07	1.06	1.04	1.05	1.05	1.05	1.04	1.06	1.05	1.04	1.04	1.05	1.04	1.04	1.06	1.05
Barley	1.09	1.09	1.09	1.11	1.10	1.09	1.08	1.10	1.10	1.08	1.07	1.07	1.10	1.08	1.07	1.09	1.08	1.08	1.08	1.09	1.07	1.06	1.08	1.07
Beans and Peas	1.32	1.33	1.32	1.35	1.34	1.36	1.34	1.36	1.32	1.34	1.31	1.33	1.34	1.32	1.30	1.31	1.33	1.34	1.34	1.31	1.30	1.29	1.32	1.31
Potatoes	0.29	0.27	0.27	0.28	0.26	0.27	0.26	0.29	0.28	0.25	0.24	0.24	0.24	0.24	0.22	0.24	0.24	0.24	0.26	0.27	0.24	0.24	0.29	0.23
Turnips	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.11	1.11	1.10	1.10	1.10	1.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sugarbeet	1.11	1.11	1.11	1.12	1.11	1.11	1.11	1.10	1.10	1.10	1.10	1.10	1.11	1.10	1.10	1.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Fodder Beat	1.10	1.09	1.09	1.10	1.10	1.09	1.09	1.09	1.10	1.09	1.09	1.10	1.10	1.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
R _{actm}	kg d.m																							
Maize	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46
Winter Wheat	0.61	0.61	0.61	0.62	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61
Spring Wheat	0.68	0.68	0.67	0.67	0.68	0.67	0.67	0.68	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.68	0.68	0.67	0.67	0.68	0.67
Oats	0.52	0.51	0.51	0.52	0.52	0.52	0.51	0.52	0.52	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.52	0.51
Barley	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.46	0.45	0.46	0.45
Beans and Peas	0.44	0.44	0.44	0.45	0.44	0.45	0.44	0.45	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.45	0.44	0.44	0.44	0.44	0.44
Potatoes	0.26	0.25	0.25	0.26	0.25	0.25	0.25	0.26	0.26	0.25	0.25	0.25	0.25	0.25	0.24	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.26	0.25
Turnips	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sugarbeet	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Fodder Beat	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Crop Residues (F _{CR})	tonnes N ₂ O-N/yr																							
Maize	0	0	0	0	0	0	0	0	0	0	1,767	2,487	2,436	1,969	1,704	1,843	2,550	2,638	3,093	3,118	2,878	2,399	1,730	1,831
Winter Wheat	6,999	7,570	8,098	5,322	6,284	6,209	8,685	7,755	7,324	5,247	7,994	6,717	9,696	7,263	10,295	7,878	7,969	7,755	11,543	7,621	7,328	10,875	8,666	5,807
Spring Wheat	1,183	1,635	1,650	2,019	1,492	1,712	1,787	2,089	1,821	2,745	1,982	3,591	2,110	3,442	3,426	2,953	2,835	1,940	1,989	1,781	1,760	1,742	1,076	1,578
Oats	1,479	1,473	1,401	1,334	1,324	1,332	1,494	1,361	1,229	1,389	1,277	1,216	1,361	1,578	1,575	1,155	1,458	1,619	1,762	1,477	1,498	1,705	1,608	1,957
Barley	15,287	14,310	14,495	12,127	11,578	13,627	15,305	13,606	13,450	15,978	16,222	15,775	12,211	14,815	16,558	12,733	14,090	13,939	16,010	15,218	15,151	17,320	15,600	20,535
Beans and Peas	189	170	173	342	327	253	280	253	416	239	98	115	99	179	185	241	261	166	146	283	320	208	253	298
Potatoes	1,224	1,100	1,210	1,122	1,200	1,195	1,366	937	957	1,018	809	855	924	865	911	721	687	706	636	651	742	632	462	695
Turnips	8,587	6,959	6,735	6,009	6,568	6,330	6,880	6,311	4,622	3,921	2,775	2,161	1,921	0	0	0	0	0	0	0	0	0	0	0
Sugarbeet	32,802	31,307	30,977	25,001	30,993	34,360	32,732	36,420	36,246	37,893	40,330	33,191	28,954	33,278	40,962	30,613	0	0	0	0	0	0	0	0
Fodder Beat	12,148	15,283	17,047	13,206	12,739	13,064	16,004	13,240	10,401	10,818	7,871	6,192	5,497	0	0	0	0	0	0	0	0	0	0	0
EF ₁	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Emissions	kt N ₂ O																							
Maize	-	-	-	-	-	-	-	-	-	-	0.028	0.039	0.038	0.031	0.027	0.029	0.040	0.041	0.049	0.049	0.045	0.038	0.027	0.029
Winter Wheat	0.110	0.119	0.127	0.084	0.099	0.098	0.136	0.122	0.115	0.082	0.126	0.106	0.152	0.114	0.162	0.124	0.125	0.122	0.181	0.120	0.115	0.171	0.136	0.091
Spring Wheat	0.019	0.026	0.026	0.032	0.023	0.027	0.028	0.033	0.029	0.043	0.031	0.056	0.033	0.054	0.054	0.046	0.045	0.030	0.031	0.028	0.028	0.027	0.017	0.025
Oats	0.023	0.023	0.022	0.021	0.021	0.021	0.023	0.021	0.019	0.022	0.020	0.019	0.021	0.025	0.025	0.018	0.023	0.025	0.028	0.023	0.024	0.027	0.025	0.031
Barley	0.240	0.225	0.228	0.191	0.182	0.214	0.																	

Annex 3.4

Activity Data for LULUCF (IPCC Sector 4.A) Forest Land

3.4.A Derivation of Historic Deforestation Areas for LULUCF and KP LULUCF

3.4.A.1 Tracking Deforestation using CORINE Land Cover Datasets (GPG approach 3)

3.4.A.2 Sampling approach: NFI grid points and aerial photography (modified GPG approach 3)

3.4.A.3 Modification to deforestation records from 2006 onwards

3.4.A.4 Allometric Equations for Biomass

3.4.A.5 Growth Models and Pre-processing Functions for CARBWARE v5

3.4.A.6 Adjustment of CSO statistics, affecting 5.B Cropland and 5.C Grassland

3.4.B Detailed Non-Forest Land Use Change Matrice

3.4.A Derivation of Historic Deforestation Areas for LULUCF and KP LULUCF

Lack of a method to record historic land use change is a significant gap in the LULUCF inventory. Ireland has attempted to improve the methodology to track deforestation, in particular, but this has only been implemented since 2006.

There are currently two data sources available to transparently report historic deforestation. However, both methods are limited and are not fully in accordance with IPCC good practice guidance for LULUCF because they do not accurately represent forest area changes, which are consistent with the forest definition (minimum area of 0.1 ha).

3.4.A.1 Tracking Deforestation using CORINE Land Cover Datasets (GPG approach 3)

The reporting of LUC matrices in Table 7.4 of chapter 7 show deforestation areas since 1990 (KP_CRF, Chapter 11) and have been estimated using CLC 1990-2000 and CLC 2000-2006.

3.4.A.1.a Background Information

Coordination of Information on the Environment, CORINE, is an EU initiative established in 1985. The CORINE methodology for indicating *Change in Land Cover* (CLC) between 1990 and 2006 is complex (CEC 1993). Computer aided visual interpretation of satellite images (Büttner et al. 2004) was applied in the process of updating the 1990 European Land Cover to 2000 (± 1 year) and the Land Cover change detection for the interval of 1990–2000, and 2000–2006 using Landsat MSS and TM satellite images. The smallest unit identified in CLC 2000 is 25 ha, and the minimum width of a linear feature is 100 m. Changes detected in the CORINE CLC were incorporated in CORINE 2000/6 only if the final CORINE polygon met the minimum mapping unit criterion of 25 ha. This means that a newly afforested area can only be detected by CORINE if it is larger than 25 ha. Clearly this is unlikely to accurately represent afforestation or deforestation since 1990, because the average size of newly established private forest parcels is 8 ha, and they are highly disperse and fragmented (Black et al, 2009 previously supplied to ERT).

The forest definition used by CORINE Land over (Bossard et al. 2000) is: “Areas occupied by forest and woodlands with a vegetation pattern composed of native or exotic coniferous and/or deciduous trees and which can be used for the production of timber or other forest products. The forest trees are under normal climatic conditions higher than 5 m with a canopy closure of 30 per cent at least”. Codes 311 representing deciduous forests, 312 for coniferous forests and 313 for mixed forests were used to interpret the change in forest area. The class, CLC 324, was excluded from the analysis, based on the assumption that this would represent recently felled/replanted and afforested areas, which are less than 10 years old. CLC 324 areas also include some semi-natural woodlands and scrub colonisation (not defined as forest land in the NFI), including: a) birch scrub on cutaway peatland; b) hazel encroachment in the Burren landscape and gorse colonisation on rough grassing land. This reclassification of land areas without ground truthing is one of our main concerns with the CLC 1990 to 2006 analysis.

Comparison of more recent high resolution datasets and CORINE clearly show that there is a mismatch in land cover classification in Ireland (Black et al., 2009). Therefore, we suggest that the misrepresentation of the CORINE afforested and deforestation area between 1990 and 2006 in Ireland may be associated with:

- a) statistical misrepresentation of Irish forest land parcels in CORINE (i.e. low resolution of CORINE);
- b) aggregation of classified categories, which may not reflect forest area change. This may be particularly relevant for CLC 324 (transitional woodland and scrub land, which may also include areas subjected to encroachment by hazel on the Burren, birch colonisation of cutaway midland peat and gorse on grazed upland, all of which may not be defined as forest land according the national definition (chapter 11).

CORINE classification and resolution problems have been highlighted in other comparative studies across northern Europe (Hazeu and de Wit 2004, Cruickshank and Tomlinson 1996).

3.4.A.1.b Methodology

Despite the above mentioned inappropriateness of CLC for reporting areas under LULUCF in a consistent, representative and accurate manner, this methodology uses the only data currently available to track historic land use change (see chapter 7).

For this exercise, the following codes were extracted; CLC 311, 312 and 313 to represent forest land area that were present in 1990, but were converted to land cover other than forest in the 2000 and 2006 time series. The resulting polygons were then intersected with a national soils map using ARCGIS to derive a land use change and soil type matrix to the periods 1990 to 2000 and 2000-2006. The resulting forest and soils GIS layers were then sampled using the NFI sample grid as discussed in H1.2 below.

3.4.A.2 Sampling approach: NFI grid points and aerial photography (modified GPG approach 3)

This sampling approach is a modification of approach 3, where the grids or centroids are sampled using a systematic sampling procedure adopted in the NFI. Note:

- The NFI was not designed to track land use change because the systematic grid (2 x 2 km) sample weighting factor used to derive total areas statistics in 400 ha (i.e. 1 sample point represents 400 ha). For small changes in forest areas, such as deforestation the sampling error is very large. For example if 10 PSP grid point are identified to be deforested then the total area represented is 4,000 ha with a lower and upper confidence limit of 945 and 7,055 ha, respectively. This represents a sampling error of 76 per cent;
- Another problem with this method is that it does not represent forest area change in a manner that is consistent with the forest area definition (0.1 ha), so is in conflict with IPCC GPG for LULUCF. This is why the NFI afforested areas are statistically adjusted using the IFORIS spatial data to consistently represent afforestation areas (see Chapter 11). However, there is at present no data available to adjust the NFI estimates of deforested land.




Figure 3.4.A2-1: The NFI systematic sample approach used to classify land use for each permanent sample plot (PSP)

The use of the NFI stratified sample 2x2 km grid of PSP described in chapter 11. Assessment of ca.18,000 point intersects with aerial photographs from 2000 and 2006 provides the opportunity to assess deforestation for this period. This method identified 15 NFI PSP grid samples which were deemed to be deforested between 2000 and 2006. The current land uses of these previously deforested lands were determined from photo interpretation using the 2006 images. Figure 3.4.A2-1 shows 2 examples of the GIS analysis and photo interpretation.



Figure 3.4.A2-2 Examples of NFI PSP (as indicated by the red cross) which were classified forests in 2000 but have since been converted to other land uses in 2006

Assessments of deforestation from 1995 to 2000 were based on a GIS intersection of the 18,000 NFI plots with the FIP95 forest parcel polygon layer. This exercise produced 105 forest parcels which were classified as forest in the FIPS 95 dataset but where classified as non-forest land in the NFI aerial photography interpretation. These 105 polygons were cross-checked with 1995 black and white aerial photographs to verify that they were forests in 1995. However, most of the sampled forest polygons were deemed to not be deforested or were originally other land uses in 1995. This was due to interpretation inconsistencies of photographs and mapping errors in the FIPS95 layer. Only 5 NFI sample points were identified to be deforested between 1995 and 2000.

The final deforestation-land use change-soils matrices for 1995-2000 and 2000-2006 were obtained by intersecting identified deforested PSP points with the national soils map database (Table 2, Figure 3.4.A2-3), see database for detailed information.




Figure 3.4.A2-3 The Irish soils map showing intersection with NFI PSP plots determined to be deforested between 1995 and 2006

3.4.A.3 Modification to deforestation records from 2006 onwards

The current methods for recording deforestation from 2006 onwards included the use of felling licence records. However, a QA exercise conducted in 2010 highlighted that 134 to 268 ha per year of land deforested since 2006 was not included in the felling licence records, if forests were less than 10 years old. These areas are not subject to the forestry act felling licence application. However, records were kept because these areas were previously subject to premium payments under the afforestation scheme. Owners in receipt of these payments are obliged to notify the Forest service if these areas are taken out ('lands taken out') of the premiums payment due to deforestation. A data base of these records is being compiled to capture the land use change and soil categories if the information is available. However, until this information does become available, the land use and soil type matrices from the felling record data for corresponding years will be used. The biomass, litter and DOM losses associated with deforestation will be based on the NFI, PSP average of all 10 year old forest areas.

Combination of the three different approaches was used to produce deforestation data for the entire time series (Table 3.4.A3).

Table 3.4.A3 The new deforestation, land use change and soil type matrix (continued on following pages):

Period	Source	Land use	Soil category	Area (ha) per year	% for period
1990-1994	CLC1990-2000			20.6¹²	100
		Grassland	Mineral	2.5	12.2
		Grassland	Peat		
		Grassland	Peaty mineral	5.7	27.9
		Settlement	Mineral	10.2	49.4
		Settlement	Peat		
		Settlement	Peaty mineral		
		Wetland	Mineral		
		Wetland	Peat		
		Wetland	Peaty mineral		
		Other	Mineral	2.2	10.5
		Other	Peat		
		Other	Peaty mineral		
1995-1999	NFI-FIPs 95			333.3¹³	100

¹² The CLC 1990-1994 area was calculated using the values show in table 1a to be, where annual deforestation area 1990-1994 = $\frac{area_{1990 \rightarrow 2000}}{10 \times 5} \times 5$

Period	Source	Land use	Soil category	Area (ha) per year		% for period
		Grassland	Mineral	266.7	80	
		Grassland	Peat			
		Grassland	Peaty mineral			
		Settlement	Mineral			
		Settlement	Peat			
		Settlement	Peaty mineral			
		Wetland	Mineral			
		Wetland	Peat			
		Wetland	Peaty mineral			
		Other	Mineral	66.6	20	
2000-2005	NFI-2000-2006			857.1¹⁴	100	
		Grassland	Mineral	342.8	40	
		Grassland	Peat		0	
		Grassland	Peaty mineral	57.4	6.7	
		Settlement	Mineral	171.4	20	
		Settlement	Peat		0	
		Settlement	Peaty mineral		0	
		Wetland	Mineral	56.6	6.6	
		Wetland	Peat	114.0	13.3	
		Wetland	Peaty mineral		0	
		Other	Mineral	57.4	6.7	
		Other	Peat	57.4	6.7	
		Other	Peaty mineral		0	
2006	Felling licence and land taken out			376.44	100	
	242.34+134.1	Grassland	Mineral	5.3	1.4	
	(LFL+LTO) ¹⁵	Grassland	Peat		0	
		Grassland	Peaty mineral	19.7	5.2	
		Settlement	Mineral	17.1	4.5	
		Settlement	Peat		0	

¹³ NFI 1995-1999 area was calculated using the values show in table 2a to be, where the annual deforested area 1995-1999 = $\frac{area_1995 \rightarrow 2000}{6 \times 5} \times 5$

¹⁴ NFI 2000-2005 area was calculated using the values show in table 2b to be, where the annual deforested area 2000-2005 = $\frac{area_2000 \rightarrow 2006}{7 \times 6} \times 6$

¹⁵ LFL is areas from limited felling licence records and LTO is the areas from lands taken out

Period	Source	Land use	Soil category	Area (ha) per year		% for period
		Settlement	Peaty mineral	0.6	0.2	
		Wetland	Mineral		0	
		Wetland	Peat	299.9	79.7	
		Wetland	Peaty mineral	30.8	8.2	
		Other	Mineral	3.1	0.8	
		Other	Peat		0	
		Other	Peaty mineral		0	
2007	Felling licence and land taken out			338.7	100	
	174.83+163.9	Grassland	Mineral	0.6	0.2	
	(LFL+LTO) ⁴	Grassland	Peat	14.5	4.3	
		Grassland	Peaty mineral		0	
		Settlement	Mineral	4.7	1.4	
		Settlement	Peat	0.8	0.3	
		Settlement	Peaty mineral		0	
		Wetland	Mineral		0	
		Wetland	Peat	297.2	87.7	
		Wetland	Peaty mineral		0	
		Other	Mineral	8.6	2.5	
		Other	Peat	12.4	3.6	
		Other	Peaty mineral			
2008	Felling licence and land taken out			294.5	100	
	26.42+268	Grassland	Mineral	80.2	27.2	
	(LFL+LTO) ⁴	Grassland	Peat	0.04	0.01	
		Grassland	Peaty mineral		0	
		Settlement	Mineral	66.4	22.6	
		Settlement	Peat		0	
		Settlement	Peaty mineral		0	
		Wetland	Mineral		0	
		Wetland	Peat	24.5	8.3	
		Wetland	Peaty mineral	21.2	7.2	
		Other	Mineral	100.9	34.3	
		Other	Peat		0	
		Other	Peaty mineral	1.1	0.4	
2009	Felling licence and land taken out			196.9	100	
	49.9+147	Grassland	Mineral	5.1	2.6	
	(LFL+LTO) ⁴	Grassland	Peat			
		Grassland	Peaty mineral			

Period	Source	Land use	Soil category	Area (ha) per year	% for period
		Settlement	Mineral	15.4	7.8
		Settlement	Peat	1.5	0.7
		Settlement	Peaty mineral	1.5	0.8
		Wetland	Mineral		0
		Wetland	Peat		0
		Wetland	Peaty mineral		0
		Other	Mineral	121.1	61.5
		Other	Peat	19.9	10.1
		Other	Peaty mineral	32.4	16.4
2010	Felling licence and land taken out			124	100
	26+98	Grassland	Mineral	39.7	39.1
	(LFL+LTO) ⁴	Grassland	Peat		
		Grassland	Peaty mineral		
		Settlement	Mineral	7.9	6.3
		Settlement	Peat		0.7
		Settlement	Peaty mineral	47.2	37.9
		Wetland	Mineral		0
		Wetland	Peat	0.5	0.4
		Wetland	Peaty mineral		0
		Other	Mineral	18.5	14.8
		Other	Peat	4.5	3.6
		Other	Peaty mineral	6.1	6.9

3.4.A.4 Allometric Equations for Biomass

Table 3.4.A.4.a: Allometric equations used to calculate biomass component for individual trees (kg d.wt tree⁻¹)

Similar species are grouped into 6 different cohorts based on available research information (Spruces, Pines, Larches, Other conifers, fast growing broadleaves and slow growing broadleaves). Abbreviations: AB-above ground, TB-total biomass, BB-below ground, FB-foliage, SB-stem (i.e. timber >7cm diameter), L_{HR}= lop and top from harvest residues, DBH diameter at breast height (1.3 m) in cm, H –height in m.

Eq	Function	Range	Equation	Coefficients				r ²	RMSE	Slope	Source
				a	b	c	d				
Spruce											
1	AB	H>4.5m	$a \times DBH^b + c \times H^d$	0.23	2.12	5×10^{-7}	4.99	0.91	0.29	1.01	i, ii
2	AB	H<4.5m	$a \times H^b \times c$	1.32	1.7	1.38		0.86	0.2	1.1	i, ii
3	TB		$\exp[Ln(a) + b \times Ln(AG)]$	1.02	1.033			0.91	0.08	1.03	ii, iii
4	BB		TB-AB								
5	FB		$AB \times a + b \times \exp[-c \times AB]$	0.025	0.089	0.003		0.68	3.4	0.98	i, ii
6	SB		$\exp[Ln(a) + b \times Ln(AG)]$	0.405	1.09			0.99	2.99	1.03	ii, iii
7	L _{HR}		AB-SB								
Pines											
8	AB	H>3.8m	$a \times DBH^b + c \times H^d$	0.07	2.42	0.039	2.51	0.93	0.13	0.94	ii, iii
9	AB	H<3.8m	$a \times H^b$	0.12	3.91			0.95	0.74	0.95	i, ii
10	TB		$\exp[Ln(a) + b \times Ln(AG)]$	1.15	1.01			0.96	0.4	1.01	ii, iii
4	BB		TB-AB								
5	FB		$AB \times a + b \times \exp[-c \times AB]$	0.025	0.089	0.003		0.68	3.4	0.98	i, ii
11	SB		$\exp[Ln(a) + b \times Ln(AG)]$	0.71	1.005			0.97	0.27	0.96	ii, iii
7	L _{HR}		AB-SB								
Larch											
12	AB	H>2m	$a \times DBH^b + c \times H^d$	0.11	2.31	0.001	3.29	0.94	0.27	0.94	ii, iii
13	AB	H<2m	$a \times H^b$	0.03	1.91			0.67	0.44	1.2	i, ii
14	TB		$\exp[Ln(a) + b \times Ln(AG)]$	1.43	0.98			0.99	0.25	0.99	ii, iii
4	BB		TB-AB								

Eq	Function	Range	Equation	Coefficients				r^2	RMSE	Slope	Source
				a	b	c	d				
5	FB		$AB \times a + b \times \exp[-c \times AB]$	0.025	0.089	0.003		0.68	3.4	0.98	i, ii
15	SB		$\exp[Ln(a) + b \times Ln(AG)]$	0.903	0.972			0.98	0.28	0.96	ii, iii
7	L _{HR}		AB-SB								
Other conifers											
16	AB	H>3.8m	$a \times DBH^b + c \times H^d$	0.022	2.73	0.19	2.06	0.96	0.46	1.008	ii, iii
17	AB	H<3.8m	$a \times H^b \times c$	0.005	1.58	1.12		0.86	0.28	1.02	i, ii
18	TB		$\exp[Ln(a) + b \times Ln(AG)]$	1.59	0.96			0.99	0.28	1.005	ii, iii
4	BB		TB-AB								
5	FB		$AB \times a + b \times \exp[-c \times AB]$	0.025	0.089	0.003		0.68	3.4	0.98	i, ii
19	SB		$\exp[Ln(a) + b \times Ln(AG)]$	0.89	0.96			0.98	0.57	1.055	ii, iii
7	L _{HR}		AB-SB								
Slow growing broadleaves											
20	AB	H>3.0m	$a + \left[\frac{b \times DBH^c}{DBH^c + 246872} \right]$	0.08	25000	2.5	246872				iv
21	AB	H<3.0m	$a \times H^b$	0.031	1.72			0.84	0.88	0.91	i, ii
22	BB		$\exp(-a + Ln(DBH) + b)$	1.509	0.284						iv
23	FB	DBH>10cm	$a \times (DBH \times 10)^b$	0.009	1.47			0.96			v
24	FB	DBH<10cm	$AB \times 0.3$					0.78	1.2	0.79	i, ii
25	SB	DBH>19cm	$a \times (DBH \times 10)^b$	0.0002	2.5			0.97			v
26	SB	DBH<9cm	$\frac{AB + BB}{1.4}$								BEF
7	L _{HR}		AB-SB								
Slow growing broadleaves											
20	AB	H>3.0m	$a + \left[\frac{b \times DBH^c}{DBH^c + 246872} \right]$	0.06	25000	2.5	246872				iv

Eq	Function	Range	Equation	Coefficients				r^2	RMSE	Slope	Source
				a	b	c	d				
21	AB	H<3.0m	$a \times H^b$	0.031	1.72			0.84	0.88	0.91	i, ii
22	BB		$\exp(-a + \ln(DBH) + b)$	1.509	0.284						iv
27	FB	DBH>3cm	$a + b \times DBH^c$	0.375	0.0024	2.517		0.90			vi
28	FB	DBH<3cm	$AB \times 0.3$					0.78	1.2	0.79	i, ii
29	SB	DBH>35cm	$a \times DBH^b$	0.0001	2.535			0.97			v
30	SB	DBH<9cm	$\frac{AB + BB}{1.4}$								BEF, vii
7	L _{HR}		AB-SB								

i National research harvested tree database (COFORD funded project CARBiFOR)

ii Black et al., Biomass equations for modelling C dynamics in Irish forests (in prep)

iii Forest Research pulled tree database (Brice Nicholl, NRS, Forest Research, UK)

iv Brown S (2002) . Measuring carbon in forests: current status and future challenges. Environmental Pollution 116: 363-372.

v Johansson, T. Dry matter amounts and increment in 21-to 91-year-old common alder and grey alder some practical implicatons. Canadian Journal of Forest Research 29 1679-1690.

vi Bartelink, H.H., Allometric relationship for biomass and leaf area of beech (Fagus sylvatica L). Annals of Forest Science, 1997. 54: p. 39-50.

vii Black K., Tobin B., Saiz G., Byrne K. & Osborne B. (2004). Improved estimates of biomass expansion factors for Sitka spruce. Irish Forestry 61:50-65.

3.4.A.5 Growth Models and Pre-processing Functions for CARBWARE v5

CARBWARE pre-processing functions and growth models

The NFI permanent plot sampling procedure does not sample all trees in a plot. Therefore, it is not possible to derive productivity index information, such as Height index or Yield class, which can be used to drive conventional stand based productivity models. The alternative and most statistically valid procedure adopted was the use of single tree models, to simulate tree growth between NFI cycles. These models can be cross-validated and re-parameterised once a repeat NFI cycle is completed. This section discussed the development of the CARBWARE growth model from draft versions for submission to International, peer reviewed Scientific Journals.

3.4.A.5.1: Pre-processing functions

Height-Diameter And Crown Ratio Modelling For Six Species Cohorts.

It is common among forestry datasets that tree height (H) or crown ratio (CR) is not measured on every tree. This creates interest in estimating the height of such trees.

A common forest inventory approach used to derive missing H and CR values involves the use of single parameter (DBH) models based on species and plot specific predictions (NFI, 2007; Wykoff et al., 1982). However, it has been suggested that these Chapman-Richards functions, or derivations thereof, are problematic because the function approaches the asymptote too rapidly, particularly when there is a weak relationship between DBH and H in larger trees. In addition, individual plot DBH-H data is sometimes too sparse to parameterise plot specific functions. Generalised DBH-H functions avoid the need to parameterise relationship for every stand. Since the relationship between DBH and H is influenced by the relative competitive position of trees within a stand and management interventions, site-level stand-density information is often incorporated (Temesgen and Gadow, 2004). Taking their results as a starting point, we address here several issues that arise in the context of our modelling dataset. These include the application of nonlinear mixed effects models which successfully borrow strength across all permanent plots, thereby facilitating imputation in plots where data is sparse or unevenly distributed. The permanent sample plot data, taken from a range of spacing and thinning experiments, used in this study is well suited, albeit not arising by design, to evaluate these stand-density parameters to describe variations in H and CR across different silvicultural conditions.

Materials and methods

Data

Data used were obtained from Coillte Teoranta's (the Irish Forestry Board state commercial forestry company) permanent sample plot record system. The dataset contains records from many silvicultural and thinning trials established during the period 1963 to 2001. The trials were initially established as replicated experimental designs with repeated measurements typically undertaken every five years. The dataset is described in Broad and Lynch (2007).

Incorporating competition covariates

The modelling here follows Temesgen and Gadow (2004) who based their work on Yang et al. (1978) and incorporated competition covariates into the Yang/Weibull function (Table 1, Model 2). We evaluate that model and also use test for differences between management regimes conditional on the DBH-H model by incorporating dummy indicator variables in the linear regression models of the

model parameters. Our aim in this section was to test if the inclusion of certain covariates, typically relating to the competition in a forest stand/plot, improved the baseline DBH-H model (3.4.A.5a, Model 1). We also investigated whether the model was improved by including random effects on the level of the plot (Table 3.4.A.5a, Model 3).

The competition covariates are plot basal area (BA, $\text{m}^2 \text{ha}^{-1}$), basal area in larger trees (BAL, $\text{m}^2 \text{ha}^{-1}$) which is the integral of the empirical frequency distribution of the BA variable from the subject tree to the largest diameter tree in the plot and plot density (DENS, trees ha^{-1}). Models were fitted in NLMixed procedure in SAS using the Trust-Region algorithm. Grids were specified as starting values for parameters where sensible.




Figure 3.4.A.5a. Model 2 Height estimates (red) and actual heights (black)

The estimates presented here depict a “cloud” because they are conditioned on covariates that vary between trees (BAL) and plots (Density, Basal Area) and over time (BAL, Density, BA).

Table 3.4.A.5a

	Model	-2l	BIC
1	$H = a(1 - \exp(b.DBH^c))$	65185	65223
2	$H = (a_1 + a_2BAL + a_3BA)(1 - \exp(b.DBH^{(c_1 - c_2BAL)}))$	58341	58417
3	$H = (Ui + a_1 + a_2BAL + a_3DENS + a_4BA)(1 - \exp(b.DBH^{(c_1 - c_2BAL)}))$	44980	45034

Table 3.4.A.5b Likelihood statistics for different forms of the DBH-H model

Model 2 is the model used in CARBWARE for the 6 different cohorts. If dependent variables had no significant influence on the H model prediction, these variables were excluded from the model.

Cohort	Model (2 variation)	a1	a2	a3	b	c1	c2
Spruce	$H = (a_1 + a_2 BAL + a_3 BA)(1 - \exp(b.DBH^{(c_1 - c_2 BAL)}))$	33.69	-0.274	0.1603	0.024	0.8846	0.0064
Pine	$H = (a_1 + a_2 BAL + a_3 BA)(1 - \exp(-b.BAL))$	16.905	0.083	0.0803	0.042		
Larch	$H = (a_1 + a_2 BAL + a_3 BA)(1 - \exp(-b.BAL))$	32.59	0.1052	0.1229	0.023		
Conifers	$H = (a_1 + a_2 BAL + a_3 BA)(1 - \exp(-b.DBH^{c_1}))$	23.226	0.1381	0.0703	0.027	1.1021	
FGB	$H = (a_1 + a_2 BAL + a_3 BA)(1 - \exp(-b.DBH))$	14.661	0.1167	0.0187	0.076		
SGB	$H = (a_1 + a_2 BAL)(1 - \exp(-b.DBH^c))$	29.677	0.1034		0.044	0.7813	

BAL is the sum of the basal area of all individual trees larger than the subject tree (m² per ha)

BA is the basal area of all trees in the plot (normalised to a ha)

DBH is the diameter at breast height (cm)

Table 3.4.A.5c CR models used in CARBWARE for the 6 different cohorts

If dependent variables had no significant influence on the H model prediction, these variables were excluded from the model.

The CR model takes the form of:

$$CR = \frac{\exp(ICR)}{1 + \exp(ICR)}$$

where *ICR* is derived from the non-linear equations, which may vary for different cohorts.

Cohort	Model (<i>ICR</i> variations)	a1	a2	a3	a4	a5	b	c
Spruce	$ICR = (a_1 + a_2 BAL + a_3 \ln(CCF) + a_4 H + a_5 \left[\frac{H}{BAL} \right] + b DBH^c$	4.8705	-0.017	-0.397	-0.119	-0.296	0.0003	2
Pine	$ICR = (a_1 + a_2 BAL + a_3 \ln(CCF) + a_4 H + b DBH^c$	3.8478	-0.024	-0.213	-0.137		0.0002	2
Larch	$ICR = (a_1 + a_2 BAL + a_3 \ln(CCF) + a_4 H$	5.8306	-0.018	-0.794	-0.039			
Conifers	$ICR = (a_1 + a_2 BAL + a_3 \ln(CCF) + a_4 H + b DBH^c$	4.1759	-0.019	-0.394	-0.965		0.0004	2

Cohort	Model (<i>ICR</i> variations)	a1	a2	a3	a4	a5	b	c
FGB	$ICR = (a_1 + a_2 BAL + a_3 Ln(CCF) + a_4 H + a_5 \left[\frac{H}{DBH} \right] + b DBH)^c$	2.4539	-0.009	-0.145	-0.045	-0.591	0.0001	2
SGB	$ICR = (a_1 + a_2 BAL + a_3 H + a_5 \left[\frac{H}{BAL} \right])$	1.477	-0.005	-0.017	-0.578			

BAL is the sum of the basal area of all individual trees larger than the subject tree (m² per ha)

CCF is the crown competition factor, which is a measure of the crown areas of the subject tree relative to a open grown tree that would not be subjected to crown competition (taken from Hassenhaur, see section B of this appendix)

DBH is the diameter at breast height (cm)

H is height (m) form actual or predicted H estimates (Table 3.4.A.5b)




Figure 3.4.A.5b. Fitted and actual height plotted (all cohorts model 2) against actual height




Figure 3.4.A.5c. Raw residuals from the fitted model plotted against the fitted height value

External validation

Based on the data presented above, model 2 was selected for validation against external data sets. In this section we compare model predictions against data from PSP non-research plots.




Figure 3.4.A.5d. Estimated and observed validation heights versus DBH

Generalised and plot-specific models

In this section we discuss the implications of using a generalised DBH-H model (i.e. one whose parameters are fitted to the entire dataset) with a plot-specific model (i.e. one whose parameters are estimated for each plot separately). We compare a mixed effects model and a plot-specific model. The former is plot-specific by the inclusion of a random residual plot effect. In what follows, by mixed model we mean the random asymptote model (Table 3.4.A.5d, Model 3). To get an idea of the difference between plot-specific and mixed-model results, we extract a plot from the dataset that exhibits a wide range of DBH and H values and then compare the models for that plot. This makes sense because the context of the comparison is how well a given model will perform for a given plot, primarily. In particular we will compare the standard error of prediction for a new tree height for both models. In the case of the mixed model, this standard error of prediction is derived as conditional on the estimated random plot effect.

A plot-specific Yang/Weibull model gives a smaller standard error of prediction than the same model estimated from the entire dataset, because residual variability for any given model will always increase from a subset of the data (plot specific) to the entire dataset (generalised). In other words, the generalised model predictions are less precise than the plot-specific predictions for any given plot, and the model mean estimate tends *towards* the overall mean and *away* from the plot-specific mean.




Figure 3.4.A.5e Model predictions for a single plot with various models, all based on the Yang/Weibull function (cf. Table 5.2b-A.1)

Green, red and black are data, estimates, and single standard error of prediction, respectively. Competition denotes a generalised model with competition covariates (cf. Model 2, Table 3.4.A.5a), Random denotes a plot-specific random asymptote (cf. Model 3, Table 5.2b-A.1), Generalised denotes parameters are estimated from the entire dataset. The smallest standard error of prediction is associated with the Plot-specific model, followed by the + Comp. + Random model. Average s.e.p. for these models are 1.39 and 1.25 respectively.

Thinning effects

All observations in the dataset were categorised by us as “thinned” or “non-thinned” depending on the general management regime for the plot. We estimated the following model to test for a residual thinning effect, having conditioned on other effects. :

$$H = (U_i + a + a_1BAL + a_2DENS + a_3BA + a_4I(Thinned))(1 - \exp(b.DBH^{(c_1 - c_2BAL)}))$$

where $I(Thinned)$ is an indicator function valued 1 if the plot was thinned and 0 otherwise. The BIC of this model was 45037, and the Wald test for the a_4 parameter ($p = 0.08$) indicated that the thinning effect was not statistically significant at the 5% level. The a_4 estimate was greater than zero, perhaps reflecting the longer tail in the height distribution for trees in thinned plots (Figure 3.4.A.5f).




Figure 3.4.A.5a. Strip-plot of Heights in the calibration dataset

Discussion

We have shown that it is possible to derive a generalised model that performs well and which by its nature deals with the data sparseness issue by estimating the “typical” parameter value and modifying this value as a function of the plot- and tree-level characteristics. The BIC results and the graphical results suggest that the inclusion of covariates in the model improves the DBH-H model (i.e Model 2), as was shown by Temesgen and von Gadow.

The inclusion of covariates in the model is a move away from the baseline model, which is a generalised approach that presumes that competition (as measured on the scale of the plot by DENS, and BA, and on the scale of the tree by BAL) does not affect the allometric relationship between DBH and H over the tree’s lifetime, when subjected to different competition pressure introduced by spacing or thinning. In the next section we address the issue of generalised vs plot specific modelling. However, our results at this point suggest that the Temesgen and von Gadow model that models plot differences through competition variables is a unified single-step approach. By contrast, the plot-specific approach can be seen as a multi-step approach, whereby the DBH-H relationship for each subject is modelled individually, and competition effects are at best implicitly described by the plot-specific fitted parameters. We might suspect that datasets that are heterogeneous across plots might be more accurately modelled using plot-specific approaches. Similarly, a generalised model might perform well on plots that are nearer the centre of the sample space than plots where management conditions are more atypical for a given dataset.

In conclusion, we adopt the use of generalised competition based models in the CARBWARE software because this performs better across all data.

3.4.A.5.2: Growth Modelling

(a) Modelling diameter increments in Irish Forests

Introduction

The modelling approach adapted in this version of CARBWARE v5 is the use of diameter increment models for all trees with a DBH greater than 5cm. This model is a distance independent individual tree growth model parameterised on Coillte permanent plot data recorded every 4 to 6 years since 1954 to 2003. These include pure and mixed species stands at establishment planting densities of 5000 to 1000 trees per ha and with different thinning treatments. The advantage of using a single tree growth model and the nature of the parameterisation data set is that different silvicultural regimes and species mixtures can be handled by one generalised modelling framework. In addition, the application data set, i.e. the data from which models will be run, does not contain explicit complete longitudinal data representing stand variables, which are used in conventional growth models.

Data operations

Two datasets are referred to, Coillte permanent sample plot (PSP) and NFI. Some of the data operations referred to below differ between these because the former has complete enumeration on a plot and is longitudinal, the latter samples from the plot and is cross-sectional.

In general, the modelling framework that we base our work on, PrognAus (see various references below), informed the types of data operations required. The framework involves, using their terminology, site, competition and size variables. Our focus was on the latter variables, and site or plot effects were accounted for using mixed model methods, whereby plot or site effects are random, blocking, effects, rather than effects whose levels have physical dimension. In any case, site or plot effects are not a feature of the growth simulator. Furthermore, incomplete enumeration of certain independent variables meant that random effects were difficult to estimate because of the sparse data. We can illustrate that elsewhere but such detail is not relevant to the CARBWARE software manual.

The variables described here are those that feature in the diameter increment model that we aim towards calibrating:

$$\text{Dinc(cm)} = \exp(a_0 + a_1 \ln \text{DBH} + a_2 \text{DBH}^2 + a_3 \ln \text{CR} + a_4 \ln \text{CCF} + a_5 \cdot \text{BAL})$$

See Table 3.4.A.5.2a. below and the text for explanation of symbols.

Data operations were concerned with assembling datasets of the variables used in the growth model, insofar as was feasible. Below, we describe any substantive data operations that were performed on the variables of interest. We exclude from this description any operations related to “data cleaning”. The main data cleaning result was to omit negative diameter increments from the dataset. Such omissions were made after such derived variables as BAL, BA and plot density were calculated. That decision was based on the fact that the omission did not have a significant impact on the results, which suggested that no further modelling was necessary to compensate for the omission. Also, if the trees involved were omitted prior to the calculation of derived variables, those variables would have been subject to an even greater bias.

Table 3.4.A.5.2a. Explanation of some symbols used in the text

Variable	Formula	Scale of measurement
CR	Crown length/height	Range (0,1)
DBH	Diameter at 1.3 m	Cm
Crown competition factor (CCF)	The “open-grown” (e.g. if every tree had zero competitors) crown area of all trees in a plot expressed as a percentage of plot area.	Percent
BAL	A function for each plot that takes as its argument any tree’s rank in the diameter distribution ordered from smallest to largest and returns the combined basal area of all trees with higher rank.	M ² ha ⁻¹
BA	Plot basal area	M ² ha ⁻¹
Annualised diameter increment (Dinc)	(DBH(t+1)-DBH(t))/([t+1] – [t]). DBH(t) stands for “DBH on the occasion of the t th measurement”. Since measurement intervals vary, this implies that [t +1] – [t] = 1 is not necessarily true, hence the use of the term “annualised”.	cm

Open-grown crown width (cw), is an intermediary variable in the calculation CCF. We estimated cw using equations derived by Hasenauer (1997). These equations return open-grown crown width in *metres*. Hasenauer (1997) derived species-specific equations that we apply in approximation to cohorts,

Spruce : $cw = \exp(-0.3232) * ((DBH)^{0.6441})$

Other conifers : $cw = \exp(0.092) * ((DBH)^{0.538})$

Pine : $cw = \exp(-0.1797) * ((DBH)^{0.6267})$

Larch : $cw = \exp(-0.3396) * ((DBH)^{0.6823})$

Slow-growing broadleaves : $cw = \exp(-0.3973) * ((DBH)^{0.7328})$

Fast-growing broadleaves : $cw = \exp(0.1366) * ((DBH)^{0.6183})$

(where a circumflex denotes exponentiation.)

Open grown crown area (m²) = $(0.25) * (3.141593 * cw^2)$

NFI and PSP datasets differed primarily in the fact that PSP plots were fully enumerated, whereas NFI plots were sampled. The sampling method, in conjunction with an assumption of homogeneous spatial diameter distribution, informs the calculation of a sampling weight or *expansion factor* which is used to allow for the possibility that some trees on a given plot were not sampled. The expansion factor is inversely proportional to the prior probability of a given tree’s inclusion in the sample, based on the trees diameter class. Each tree in the sample is thus duplicated by a number of times equal to its expansion factor. This duplication is allowed for when calculating plot-level derived variables, e.g. Density, by incorporating the expansion factor into the equations. For example, the estimated number of trees on a plot with a single sampled tree of 8cm is (12.62/3)². See Figure 3.4.A.5.2a. for an explanation.




Figure 3.4.A.5.2a. The NFI sampling scheme at the plot-level

The expansion factor for a tree in the i th diameter class is $(R_3/R_i)^2$

Diameter increment

The diameter increment model for each cohort was calibrated by fitting to data from the PSP dataset.

$$Dinc = \exp(a_0 + a_1 \ln DBH + a_2 DBH^2 + a_3 \ln CR + a_4 \ln CCF + a_5 \cdot BAL) + e$$

Where a_i , $i = 1 \dots 5$ are coefficients and e is a residual that was autocorrelated between measurements on the same tree and independent otherwise. The fitting was done in the Glimmix procedure in SAS, and the model is a GLM with Gaussian variance function and a log link. This is slightly different from Monserud and Sterba (1997), who log-transformed the response, where we log-transform the expected value of $Dinc$, and didn't model autocorrelation.

Where fitting was unsatisfactory, i.e. because of parameter instability or data sparseness, a submodel was selected. A criterion of model selection was that the parameters should be qualitatively similar to those estimated by Monserud and Sterba (1997). In this respect, the fitting of the increment models is better described as model calibration than model selection.

The parameters for the fitted models were:

FGB

$$E(Dinc) = \exp(-2.8528 + \ln(DBH) \cdot 1.1729 - 0.00012 \cdot DBH^2 + \ln(CR) \cdot 0.8241 - 0.000015 \cdot CCF)$$

Larch

$$E(Dinc) = \exp(-2.2969 + \ln(DBH) \cdot 0.6338 - 0.00096 \cdot CCF)$$

OC

$$E(Dinc) = \exp(-1.4191 + \ln(DBH) \cdot 0.554 - 0.00025 \cdot DBH^2 + \ln(CR) \cdot 0.5549 - 0.00052 \cdot CCF - 0.00646 \cdot BAL)$$

Pine

$$E(\text{Dinc}) = \text{EXP}(-1.3466 + \text{LN}(\text{DBH}) * 0.741 - 0.001 * \text{DBH}^2 + \text{LN}(\text{CR}) * 0.998 - 0.00066 * \text{CCF} - 0.00417 * \text{BAL})$$

SGB

$$E(\text{Dinc}) = \text{EXP}(-2.5897 + \text{LN}(\text{DBH}) * 0.7534 - 0.00068 * \text{DBH}^2 - 0.0006 * \text{CCF} - 0.00979 * \text{BAL})$$

Spruce

$$E(\text{Dinc}) = \text{EXP}(-1.8628 + \text{LN}(\text{DBH}) * 0.9456 - 0.0005 * \text{DBH}^2 + \text{LN}(\text{CR}) * 1.1639 - 0.000638 * \text{CCF} - 0.00273 * \text{BAL})$$

Uncertainty:

In this section we look at various measures of the performance for the different models discussed above. The performance measures quoted give rough ideas about how the models perform. It should be noted that performance can be improved somewhat by including plot and site effects but since these are problematic for extrapolation from PSP to NFI, they were omitted from the Dinc model. They were also omitted from within NFI imputation models, by which we mean imputation models calibrated on NFI data, for similar considerations. They were not omitted from PSP-specific models.

We looked at the performance of the various models – DBH-H, CR, Dinc – for the two datasets. Some measures we could have used, that are used by Thurig et al (2005), for example, are *accuracy*, *precision*, and *excess error*, calculated as follows.

Accuracy : $((\sum(\text{predicted-observed})/n) * 100)/m$. Where m is $E(\text{obs})$, and n is the number of observations.

Precision : $\text{SD}(\text{pred-obs})$

Empirical Excess error (%): $((1 - \text{Sec})/\text{Sei}) * 100$. Where Sec is the *precision* of the calibration data, and Sei the *precision* of the independent data.

Theoretical Excess error : $(1/n)[\sum(\text{pred}_{(-1)} - \text{obs})^2 - \sum(\text{pred-obs})^2]$. Where $\text{pred}_{(-1)}$ is the leave one out prediction error

Note that *empirical excess error* is only viable when doing external validation.

Temesgen and von Gadow (2004), for example, use *root mean squared error* (RMSE) and Bias to evaluate their models.

Bias : $(\sum(\text{pred-obs})/n)$

RMSE : $\sqrt{(\sum(\text{pred-obs})^2 / (n-p))}$. Where p is the number of parameters in the model.

Another measure is *mean absolute error* (MAE).

MAE : $\sum |\text{pred-obs}| / n$

A certain amount of model selection was done, as noted above, when fitting the models to the data in the first place. This ensures that the fitted models are the most parsimonious to minimise residual error. However, model performance is best evaluated by external validation or, failing that, some cross-validation. We conduct leave-k-out cross validation on the Dinc calibration data. *MAE* and *RMSE* are calculated for each cross-validation dataset replicate. External validation data was only available for the PSP DBH-H model, and that is discussed in another document.




Figure 3.4.A.5.2b Within-sample Precision (upper panel) and Bias (lower panel) for imputation

Values are plotted for each dataset, for cohorts, and for models of Height and Crown ratio.




Figure 3.4.A.5.2c Leave k -out cross validation results, precision (top) and mean absolute error (bottom) for the Dinc model

The probability of inclusion in the validation dataset is 0.33. 20 cross-validation replicates are displayed.

Discussion and conclusions

The lines joining the points in Figure 3.4.A.5.2b. are only included to facilitate a comparison between panels. The interpolating lines in Figure 3.4.A.5.2c are indicative of variability between the different cross-validation runs. This variability is partly a function of data resources, i.e. the number of cases,

and the size of the validation sample as a proportion of the number of cases. The low variability of Pine and Spruce, the cohorts with by far the most number of cases, reflects this.

In Figure 3.4.A.5.2b, the better performance of PSP versus NFI is partly a result of including such blocking effects as site and plot. This idea is also illustrated with more detail in the document on DBH-H modelling.

From Figure 3.4.A.5.2b, bias levels are low for both NFI and PSP. Pine and Spruce, the most important cohorts, are among the top performers. This partly reflects the better data resources for those cohorts.

(b) Modelling height increments for small trees

Introduction

Height growth for small trees is a driving developmental force as trees compete for light and vertical growing space. Because of this, the small-tree portion of CARBWARE is a height-growth driven model; height growth is estimated first, and then diameter growth is predicted from height growth. Equations used to predict small-tree height increment vary by species, variant, silvicultural practice and site type. Most single tree based models for young growth, generally use the same the same predictors as described for DBH increment models. However, the NFI data set provides little or no information on predictors for young tree height. The development of a H growth model for trees less than 1.3 m to a maximum H of 2.3 to 5.1 m (i.e. the diameter at breast H, DBH) is described here. The model uses a empirical Chapman-Richards approach for different species with sub models for different height index ratios (i.e. mean H over age as proxy's for young stand productivity and site factors).

Methodology

Modelling framework

The model uses a empirical Chapman-Richards approach for different species with sub models for different height index ratios (i.e. mean H over age as proxy's for young stand productivity and site factors).

$$xHinc_{i,j}^{n+1} = \frac{a_1}{1 + \exp(-1 \times \left[\frac{age^n - a_2}{a_3} \right])}$$

where, xH is mean height of all trees in the NFI plot for the i th species and j th H index ratio at the determined age ($n+1$). The age of the forest (n) is obtained from the NFI stand attribute data. The partial coefficients (a) for each species and productivity class and goodness of f

Once the new mean tree H ($xH_n + xHinc_{n+1}$) is computed, the individual tree H is recalculated based on a scaling function:

$$H_{n+1} = \frac{H_n}{xH_n} \times H_n$$

where, H_{n+1} is the individual H of the tree in the plot in the year following the NFI , H_n is the individual H in the year the last NFI was completed (2005), and xH_n is the mean H of trees in the plot in the year the last NFI was completed.

The Productivity class (H over age) categories were defined to match conventional yield class productivity indices (YCe_q) as described by (Christy and Edwards, 1981). This was derived by comparison of Chapman Richard outputs from each H index ratio (HI) with static age-H tables at ca. 10 to 20 year old crops.

$$YCe_q = HI = \min, \{YCH_{ij} - xH_{ij}\}^2$$

where, YC_{eq} is the HI equivalent to YC at the lowest least-squares different between the yield table H values (YCH) and the predicted mean height (xH see equation 1) for the *i*th cohort and *j*th HI.

Selection of tree for H increment model

All trees with no measurable DBH are selected for growth increment using the H model. The CARBWARE model also selects eligible trees to be grown using the H growth model based on cohort-specific threshold DBH values (Table 5.2B-B.2). These are derived from analysis of the minimum DBH ranges suitable for the DBH increment model. The transition from the H to DBH increment model is based on the threshold H value which corresponds to the minimum allowable DBH value to be used in the DBH increment model (Table 5.2b-B.2). If a tree has a larger corresponding DBH than the threshold value, it is grown using the DBH increment model.

Table 3.4.A.5.2b. Threshold minimum DBH values suitable for use in DBH increment model and corresponding cut-off H values used for H growth in small trees

Cohort	DBH threshold (cm)	Corresponding H (m)
Spruce	4	2.7
Pine	4	5.1
Larch	2	3.6
Other conifers	4	3.1
Slow growing Broadleaves (SGB)	2	4.2
Fast growing Broadleaves (FGB)	2	3.2

Datasets and measure of goodness of fit

We used both the Coillte PSP and NFI individual tree data base to develop H-age curves (range 0.1 to 12 m). Data operations were concerned with assembling datasets of the variables used in the H model, insofar as was feasible.

We looked at the performance of the various models –H-Age for different cohort for the combined datasets. Some measures we could have used, that are used by Thurig et al (2005), for example, are accuracy, precision, and excess error, calculated as follows.

Fitted model parameters

Table 3.4.A.5.2c shows the partial coefficients for each species and productivity class for the Chapman-Richards H-Age functions.

Table 3.4.A.5.2c Spruce cohort

HI range	YCeq				Precision	RMSE	Bias
		a_1	a_2	a_3			
>1.2	>24	1.02	5.59	2.04	1.8	4.69	0.32
1-1.2	24	1.05	7.05	2.32	1.42	4.23	-0.23
0.8-1	22	0.76	5.98	1.63	1.33	3.21	0.11
0.6-0.8	20	0.66	5.51	1.33	0.66	2.55	0.56
0.5-0.6	18	0.57	5.26	1.12	0.89	1.69	0.45
0.4-0.5	16	0.53	5.35	1.47	1.11	3.66	0.32
0.3-0.4	14	0.48	5.32	0.54	0.74	3.54	0.62
0.2-0.3	12	0.44	6.59	2.20	1.53	4.53	0.24
0.1-0.2	10	0.35	6.93	2.27	0.69	1.77	-0.43
<0.1	<10	0.28	8.02	0.35	1.9	4.23	-0.7

3.4.A.5.2: CARBWARE stand modification functions

The NFI permanent plots structure is modified at each growth cycle iteration to simulate the losses associated with natural mortality and harvest. This section discusses the development of the CARBWARE modification functions from draft versions for submission to International, peer reviewed Scientific Journals.

3.4.A.5.2.1: Mortality models

Introduction

In the general context of forest growth models, and at the most basic level, the tree mortality module's role at each iteration is to classify a particular tree in the dataset as being either dead or alive. This paper approaches this problem in the context of an individual-tree model of mortality that is both age- and distance-independent. The specific modelling framework within which the mortality module will be applied, is a framework similar to the PrognAus framework, with the goal of estimating annual forest dynamics for Ireland.

Literature review

There are two areas of interest for the literature context of this paper: *tree-mortality* modelling, and *threshold-based* classification. (Note that this paper is not focussed on a survival analysis, as one might perhaps expect, because such models are time-dependent.)

1. Mortality modelling in Forest Succession.

Wunder et al. [2006a] compared the use of classical stress-thresholds in mortality modules of forest succession (\gap") models. They conclude that logistica1 regression-based models are superior to stress-threshold models with regard to predicting time of tree death.

Baesens et al. [2003] review threshold-based classifiers in the context of credit-scoring. They examine logistic regression, discriminant analysis, k-nearest neighbour, neural networks and

decision trees, advanced kernel-based classification algorithms such as support vector machines and least-squares support vector machines (LS-SVM). They assess performance using the classification accuracy and the area under the receiver operating characteristic curve. They found that both the LS-SVM and neural network classifiers yield a very good performance, but also simple classifiers such as logistic regression and linear discriminant analysis perform very well for credit scoring.

Bigler and Bugmann [2004] introduced a new approach to modelling tree mortality based on different growth patterns of entire tree-ring series. They were interested in predicting time of tree death. In their study, dendrochronological data from *Picea abies* (Norway spruce) in the Swiss Alps were used to calibrate mortality models using logistic regression. They introduced a mortality threshold and classified a tree as dead if its modelled mortality probability curve plotted over time went above that threshold. They ignored autocorrelation at the modelling stage, and applied a jackknife method to correct for the resulting biased variance estimates. They found that the most reliable models were those that included relative growth rate and a short-term growth trend as explanatory variables.

Focussing on the role played by life-history strategies in determining tree mortality Wunder et al. [2008] investigated whether the relationship between growth and mortality differs among tree species and site conditions. This carries on from Monserud [1976] who showed that reduced growth generally accompanies a higher mortality risk. For each of nine species, they modelled mortality probability as a function of relative basal area increment, tree size and site. They selected the species-specific model with the highest goodness-of-fit and calculated the area under the receiver operating characteristic curve and calibration measures. The discriminatory power as measured by AUC ranged from 0.62 to 0.87. They found that most growth-mortality relationships differed among species and sites, i.e. there is no universal growth-mortality relationship.

It has been noted that a lack of long-term growth/mortality data has made it difficult to evaluate the performance of mortality models. Wunder et al. [2006b] adopt a "virtual ecology" approach to this problem, simulating forests with either of two a priori specified growth-mortality relationships. They simulate different sampling regimes in these virtual forests, thereby generating virtual tree-ring data, forest inventory data, or a combination of both. They compare eight existing or newly developed models of different structural flexibility by their ability to model the growth-mortality relationship in the simulated data, and quantify the deviation from the *a priori* specified growth-mortality relationships with the Kullback-Leibler distance. Of the models they evaluated, the highest accuracies were obtained with tree-ring based models, which required only small (approx. 60) numbers of dead trees. For larger sample sizes (approx 500 dead trees) forest inventory based models were also seen to be accurate. They also showed that flexible statistical approaches were superior to less flexible models only for large sample sizes (totally 2000 trees) and that the additional use of Bayesian statistics, model accuracies only when model flexibility was constrained. They also provided guidelines for sufficient sampling schemes in real forests.

In the PrognAus framework, Monserud and Sterba [1999] modelled mortality in Austrian forests for six major species based on 5-year re-measurements of the permanent plot network of the Austrian National Forest Inventory. Their general results, varying slightly between species, was that inverse of tree diameter, crown ratio and BAL were respectively the three most closely correlated factors in their model with 5-year mortality rates. They compared mortality rates across tree diameter class, thereby identifying a classic U-shape in mortality rates as diameter class increased. They modelled mortality rates rather than individual tree mortality probability, and validated the model with the

chi-square statistic calculated between observed and estimated. Because the explanatory variables in their model were measured on the scale of the individual tree, they were able to calculate the classification success rate using the complement of the overall proportion of mortality (i.e., approximately 93%, although it is not clear from the text) as the threshold. On this basis, their model correctly classified between 81 and 92%, of live trees, and between 25 and 44%, of dead trees. However, their treatment of the threshold is very brief, and may not be a typical interpretation, e.g. in their interpretation, a tree is classified as dead if the threshold exceeds the modelled probability. Also, they derive a total correct classification accuracy of 86%.

Materials and Methods

We fitted logistic regression models to the growth dataset. We investigated model performance in the case of separate models for each cohort. (Principal issue here was the lack of data for some cohorts). The response variable was a binary indicator of mortality (arbitrarily, 1 = tree dead at time of DBH measurement, 0 = tree alive). We only included trees whose cause of death was natural mortality, e.g. such causes as windblown, diseased, were excluded. Explanatory variables were as such that were selected by Monserud and Sterba [1999] {DBH and transformations thereof, CR, BAL, CCF}, but we also investigated relative growth indicators that Bigler and Bugmann [2004] noted as being useful correlates. Site and plot effects were modelled as random, and consecutive observations on the same tree were modelled as being correlated. Conditional on this correlation structure the fixed effects parameters were selected by backward selection starting with the candidate set of covariates just listed.

Models were fitted by maximum likelihood and individual fixed effects were identified as non-significant on the basis of asymptotic Wald-tests. This was done for each cohort separately. Performance of candidate models was then evaluated by cross-validation and external validation (comparing fitted to observed mortality in NFI dataset) and with threshold-based classification tools like the ROC and ROL curves and related measures and hypothesis tests. Cross-validation was done on a leave-k out basis, where the data \left-out" was selected at random. Up to twenty independent cross-validation runs were performed, and up to 33% of the data was left-out as cross-validation data for each run.

Other performance measures were consulted, and the ROC convex hull played a role in our chosen classifier. We used threshold-averaging to investigate the performance of the classifier in cross-validation and bootstrap scenarios. We derived confidence bands for the ROC curve of the chosen classifier following the approach of Macskassy et al. [2005]. (Note, the authors have also developed techniques for point interval estimation also, the reference appearing in that paper.)

Performance measures in ROC space and their role in uncertainty analysis

The AUC of the ROC curve is the estimated probability that the classifier will give a higher score to positive cases than negative cases. (In our application, the estimated probability of mortality is higher for dead trees than live trees.) We envisage that an uncertainty analysis of the forest growth model of which the mortality classifier is a component part could utilise this probability and its standard error in monte-carlo simulation assessments of overall uncertainty and sensitivity.

The AUC is equivalent to the Mann-Whitney U-statistic, and methods for comparing AUCs have been developed as a result, e.g. Heagerty et al. [2000]. The principal complicating factor here is the underlying correlation structure of the comparison, which can be influenced by details pertaining to the derivation of the classification forecasts, the setup of the calibration datasets, or whether the

forecasts are clustered in some way, e.g. DeLong et al. [1988], Obuchowski [1997], Heagerty et al. [2000], Mason and Graham [2002].

The convex hull of a classifier, or group of classifiers, in ROC space, can be seen as the optimal attainable classification performance. Fawcett [2006] notes that candidate classifiers that do not attain the convex hull can be discarded, on the grounds that a better classifier in ROC space exists. He suggests a method for interpolating between candidate classifiers to better approach the limit of performance estimated by the convex hull based on mis-classification costs and the prior class distribution.

When comparing ROC curves, per se, a complicating factor when it is of interest to compare different classifiers crops up if the classifiers in question are of a different "class", e.g. a probabilistic classifier versus a discrete classifier, or, more generally, comparisons across model classes, whose scoring systems are incommensurate Fawcett [2006].

Datasets

Permanent Sample Plot

The mortality model is calibrated on data extracted from the permanent sample plot record system of Coillte Teoranta (the Irish Forestry Board state commercial forestry company). Broad and Lynch [2006b] provide details of the dataset in the context of modelling plot volume. The database consists of records of many silvicultural and thinning trials. These longitudinal trials were established from the 1950s onwards, and were initially established as replicated and blocked experimental designs Broad and Lynch [2006a].

Although there are several categories of disease or mortality causes in the PSP database {including, Windblown, Uprooted, Diseased, Broken and Dead}, we modelled only the binary response Dead/Alive for the initial model. In this way, after derived variables {basal area, plot density, etc.} were calculated, only data points that could be classified as Dead/Alive, were kept in the calibration dataset

National Forest Inventory Plot data

We validated the ROC curve for the chosen model on the NFI data. In the NFI sample, the probability that a tree's status as dead or alive will be recorded {more generally, the probability that any feature of the tree is measured} is a function of its diameter class at the time of survey, and its distance from the centre of the plot. The expansion factor concept is a weight that varies between each tree in the dataset that estimates the prior probability of the tree's inclusion in the dataset (see Ch6).

The question to address is whether we can arrive at a sensible definition of representative mortality. At issue is how to derive a binary individual-tree-level mortality rule based on information in the NFI dataset, given the fact that there is missing information due to the sampling scheme. With this in mind, Figure 3.4.A.5.2.1a classifies all dead trees in the PSP database by cohort, and describes the empirical distribution of diameter classes conditional on mortality status. (We have included the diameter class (0,7] for completeness, even though there is no equivalent in the NFI dataset.) Note that the left-hand column is very similar to the unconditional distribution of diameter classes, so it does not need to be displayed. On those grounds, a comparison of the columns of Figure 5.2c-A.2 shows the dramatic extent to which the chance of mortality declines if a tree does not die while in the lowest diameter class. For example, the global fraction of trees in the Spruce cohort in the

lowest diameter class is very small, but this class represents 50% of dead trees in the cohort. Similarly for Pine, OC and FGB.

The right-hand column of Figure 3.4.A.5.2.1a, at least for the cohorts with enough observations, suggests a way to make the operation of a binary mortality rule more accurate in the context of the NFI sampling scheme. The basic idea would be to use the column heights as weights in a finite mixture function whose components would be the outcome of the mortality rule. Rather than reducing the expansion factor by one unit when death is predicted (which, we can show, can lead to an unrealistically high global mortality rate), the actual reduction would be a function of the weight for the given diameter class. This method could be stochastic or deterministic. Other information might be used to inform the values of the weights, including a forester's rule of thumb about global mortality (i.e., ~6%), or information from the NFI or a meta-analysis.

A similar approach would be to mix the outcome of the mortality rule with the diameter class mortality weights. It might be possible to iteratively tune the weights and/or the rule's cut-off parameter.




Figure 3.4.A.5.2.1a The empirical distribution in the PSP dataset of diameter classes of dead/alive trees classified by cohorts

Results

Candidate model Number 1. Candidate model 1 was a fixed effects model. A logistic GLM was fitted in Glimmix. The _fixed effects were DBH, BAL, and

$$\text{RelDiamInc} \left(\frac{\text{growth}(t - t - 1)}{\text{DBH}(t)} \right)$$

Part of the reason for looking at this model was that it was not subject to additional uncertainty due to imputation of missing X data, as would have been the case with the model put forward by Monserud and Sterba [1999], which also conditioned mortality on CR, a variable that was not measured on every tree in our dataset.

There are several points of interest to the results of this model fitting:

1. The characteristics of the parameters.
2. The cross-validation exercise.
3. The out-of-sample/deployment performance. E.g. how well the model described NFI mortality.

Estimated parameters Candidate Model 1 (Used in CARBWARE models)

The fitted parameters and their standard errors are presented in Table 3.4.A.5.2.1a. We supply parameter estimates for cohort-wise fits and the fit to the entire dataset, with no cohort-effect parameter.

Table 3.4.A.5.2.1a Candidate model 1 parameters

Fast-growing broadleaves cohort

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	-2.9295	0.1510	11784	-19.41	0.0001
DBH	-0.4307	0.02508	11784	-17.17	0.0001
BAL	0.06816	0.004384	11784	15.55	0.0001
RelDiamInc	-1.6783	1.2147	11784	-1.38	0.1671

Larch cohort

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	3.0526	0.1691	6544	18.06	0.0001
DBH	-0.4373	0.01276	6544	-34.27	0.0001
BAL	0.05688	0.003066	6544	18.56	0.0001
RelDiamInc	-14.7793	2.5794	6544	-5.73	0.0001

Other conifers

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	4.3636	0.1090	21239	40.02	0.0001
DBH	-0.8384	0.01447	21239	-57.95	0.0001
BAL	0.05970	0.002078	21239	28.72	0.0001
RelDiamInc	-29.2957	1.0322	21239	-28.38	0.0001

Pine cohort

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	2.3952	0.04531	187E3	52.86	0.0001
DBH	-0.8127	0.007225	187E3	-112.49	0.0001
BAL	0.08083	0.000999	187E3	80.91	0.0001
RelDiamInc	-23.0015	0.3995	187E3	-57.57	0.0001

Slow growing broadleaves

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	29.6029	7.1305	1027	4.15	0.0001
DBH	-2.1970	0.4873	1027	-4.51	0.0001
BAL	-0.1225	0.01754	1027	-6.98	0.0001
RelDiamInc	-2199.90	521.36	1027	-4.22	0.0001

Spruce cohort

Parameter	Estimate	s.e.	df	Wald statistic	Wald p-value
Intercept	1.2286	0.02747	298E3	44.72	0.0001
DBH	-0.6640	0.003840	298E3	-172.93	0.0001
BAL	0.05051	0.000529	298E3	95.57	0.0001
RelDiamInc	-13.0524	0.2544	298E3	-51.30	0.0001

Candidate Model 2

The fixed effects in Candidate model 2 were those in Monserud and Sterba [1999], and diameter increment as a proportion of diameter (RelDiamInc).

Cross-validation and deployment performance

We performed plot-wise and case-wise leave k-out cross-validation of the chosen models. The case-wise deletion algorithm was very slow for the Pine and Spruce cohorts, in which case we opted to use only plot-wise deletion. The algorithm selected plots for deletion from the fitting dataset using a

Bernoulli mechanism with parameter p , which we sometimes changed depending on the number of plots in the cohort dataset. Details are provided with each graphical representation of the results in Figures below. Twenty "leave-outs" were performed and the variability in these twenty runs is represented by the dotted curves.




Figure 3.4.A.5.2.1b The Receiver operating characteristic curve for Candidate model 1 (panels (a),(b)) and model 2 (panel (c),(d)) in the Fast-growing broadleaves cohort

20-fold cross-validation plotwise with average leave-out probability $p = 0.3$. Curves for each cross-validation run and a threshold-averaged curve are shown.

We estimated the ROC curve for each cohort model's out-of-sample performance by comparing model predictions with the actual NFI mortality data (Figures 3.4.A.5.2.1b). The cross-validation and deployment performance plots are presented pair-wise in the Figures below. In all cases model candidate outperformed candidate based on false positives and fit. For example we show the results for Fast growing cohorts in Figure 3.4.A.5.2.1c. Note that Slow-growing broadleaves cohort did not have enough data for the cross-validation to be feasible, the ROC curve for that cohort depicts in-sample performance.

Models fitted to NFI data

When fitting models to the NFI data we used backward elimination, starting with the parameters in the Monserud and Sterba [1999] model. Relative diameter was not used, because the dataset is cross-sectional. In Figure 3.4.A.5.2.1c we present an example of the out-of-sample performance (i.e. their performance in predicting NFI data) of the two PSP-calibrated models, and the in-sample performance of the NFI-calibrated model.




Figure 3.4.A.5.2.1c The Receiver operating characteristic curve for Fast-growing broadleaves cohort Candidate models 1 and 2 fitted on PSP, and for the NFI-fitted model.

The selected CARBWARE models based on NFI data fits

Fast-growing broadleaves cohort

$$P_{mort} = IL(12.93 + 0.068 \times BAL - 2.868 \times CR - 0.962 \times DBH - 72.28 \times \frac{1}{DBH} + 0.009 \times DBH^2)$$

Larch cohort

$$P_{mort} = IL(-4.9266 + 0.04273 \times DBH)$$

Other conifers

$$P_{mort} = IL(-4.5226 + 0.067 \times BAL - 6.05 \times CR + 0.066 \times DBH)$$

Pine cohort

$$P_{mort} = IL(2.395 + 0.0408 \times BAL - 3.0036 \times CR - 0.2263 \times DBH - 24.21 \times \frac{1}{DBH})$$

Slow growing broadleaves

$$P_{mort} = IL(15.78 + 0.0109 \times BAL - 2.2807 \times CR - 0.771 \times DBH - 94.002 \times \frac{1}{DBH} + 0.00449 \times DBH^2)$$

Spruce cohort

$$P_{mort} = IL(6.8976 + 0.0912 \times BAL - 21.3795 \times CR - 0.8287 \times DBH - 49.15 \times \frac{1}{DBH} + 0.008 \times DBH^2)$$

Where ($0 < P_{mort} < 1$) is the probability the tree is dead. We map then this estimated probability onto the binary (Dead, Alive) outcome using a cutoff, which may differ between cohorts. More details on this is give elsewhere. $IL(.)$ is the inverse logit, e.g. $IL(x) = \exp(x)/(1+\exp(x))$.

Choosing the operational cut-off

To identify a cut-off level to use for the mortality probability estimate, we plotted the True positive rate (TPR) and (FPR) on the same axis versus the cut-off (e.g. Figure 5.2c-A.5). In forest mortality, the number of positive cases (dead trees) is usually greatly outnumbered by the number of negative cases. This suggests that, all mis-classification costs being equal, the cut-off should be chosen with a view to keeping as small as feasible the rate of false positives predicted by the resulting rule, even though the rate of true positives is reduced as an unavoidable consequence. When combining individual cohort results to make an aggregate prediction the issue of false positive rate is of particular importance for large cohorts, because they have a greater weight in the aggregate estimate. In Figure 5.2c-A.5 we represent an FPR of not greater than 0.001 with a blue vertical line, and an FPR of not greater than 0.01 with a green vertical line, to illustrate the trade-off involved in each particular case.

Table 3.4.A.5.2.1b Formulae for some standard performance measures used in the text

Performance Measure	Formula
Accuracy	$\frac{TP+TN}{TP+FP+TN+FN}$
Rate of positive predictions	$\frac{TP+FP}{TP+FP+TN+FN}$
Correlation Coefficient	$\frac{(TP \cdot TN) - (FP \cdot FN)}{\sqrt{(TP+FN) \cdot (TN+FP) \cdot (TP+FP) \cdot (TN+FN)}}$

Note TP, TN, FP, and FN are the numbers of true positives, true negatives, false positives and false negatives, which are tallied by comparing the predictions with the data.

Figures 3.4.A.5.2.1d illustrate some other considerations for choosing cut-off points, accuracy, rate of positive predictions and a correlation coefficient are plotted for a range of cut-offs.

The graphs illustrate why the accuracy measure should not be used in isolation when choosing a cut-off. For example, in Figure 3.4.A.5.2.1d a high accuracy is obtained despite the correlation coefficient indicating that the correlation between correct predictions and the data is worse than random, i.e. a negative correlation coefficient.




Figure 3.4.A.5.2.1d TPR (Black) and FPR (Red) versus cut-off for Fast-growing broadleaves

The vertical green line shows the cut-off where $FPR < 0.01$, the blue vertical line shows the cut-off where $FPR < 0.001$

Discussion

In binary classification, a common approach is to visualise the parameterised curve described by plotting two performance measures as a parametric curve parameterised by the threshold value. Comparing models based on classification and mis-classification rate (precision, recall, etc.) make more sense when there is some hierarchy of misclassification errors. That is, that we can quantify the relative importance of gains from correct classification and losses from incorrect classification. Such a loss function is particularly useful when the number of objects to be classified is not equal, because then the trade-off curves are much more likely to be nonlinear and the concept of trade-off between competing performance measures is not easy to understand. The problem is how to specify losses/gains, in other words, how to quantify Trade-off, how to measure gains and losses in the same units so a net trade-off can be calculated. Otherwise, it is not always clear, even for commonly presented parameterised curves, in what sense the trade-off is occurring, particularly when a “good” classifier, e.g. one that exhibits desirable tendencies in threshold-space, can a priori exhibit a number of different “shapes” when presented as a “trade-off” curve.

For example, the class ROC trade-off curve has *a priori* a sense in which a classifier is good or bad. This is when the majority of the ROC curve lies below the line of equality. However, the precision-recall curve is not so easily understood. We know that the best classifier from a group is the one with the largest area between the curve and the line of equality. However, because the value of the precision at zero threshold is a function of the number of objects in each class to be classified, it is

possible to have a “good” classifier for which that area is zero. However, such a classifier is probably not statistically better than the naive, 50:50 classifier. We propose that for a classifier to be demonstrably better than the naive classifier, it should at the minimum describe a positive region between the curve and the line of equality. We conclude that the precision-recall curve does not describe a trade-off, and that in fact, a trade-off should have a point of equilibrium and the gains and losses should be incurred when the threshold moves from that point in either direction. In other words, the gains and losses as quantified by the two performance measures should be negatively correlated, for the parameterised graph to truly describe a trade-off. The precision-recall performance measures, for example, are positively correlated (both have TP in the numerator), and so their parameterised curve representation does not describe a true trade-off situation in every region of threshold space. If we overlay the two graphs with precision and recall on the y and y' axes, and threshold on the x axis, we can see more clearly where a true trade-off may occur. It is likely that should a true trade-off occur, that the region between the parameterised curve and the line of equality will have to be positive. As external corroboration, DeLong et al. [1988] note that the cost or loss function is essential to deciding the optimal cutpoint/threshold for a ROC curve. In summary, there are therefore two issues: comparing classifiers and, given a classifier, choosing a cut-off point.




Figure 3.4.A.5.2.1e illustrating some other performance measures of the NFI-calibrated model for the Larch cohort across the cut-off range and in particular the 0.01 (green vertical) and 0.001 (blue vertical) cut-off points

This latter can only be done in conjunction with some kind of loss function describing costs of the different types of classification error. The kind of classifier we are using, based on multiple correlation/regression, and therefore wholly empirical, is easier to select than other types of classifier. We can use model selection criteria based on correlation/regression, or minimization of errors, or some other abstract modelling concepts. Then, the classifier selected, we can choose the cut-off. In what we call mechanistic classifiers, such as described in Martin-Davila et al. [2005], where the classifier is predicated first and foremost on an understood pathway, not naive correlation, the threshold has a physical dimension, and the choice of cut-off has a defined purpose in a physical system. Note that a logistic regression with a single explanatory variable can be made to fit such a schema. In fact, it might be possible to define a convex hull of the multiple explanatory variables to take the place of single-variable classifier in that schema. Also, some variables might be better at defining the threshold than others and this can also be examined. A convex hull defined by cut-off points in each explanatory variable might be envisaged to play the role of a kind of "syncretized" cut-off point. In such an instance, it would be relevant to assess the cross-correlations among the explanatory variables.

Conclusions

We set out to determine a logistic regression model of mortality that could be used to describe mortality in the NFI data. This was the ultimate goal of the model. We investigated the possibility of calibrating this model on the permanent sample plot longitudinal data but found that we could improve the result by simply calibrating the parameters on the NFI data alone. In the absence of a mis-classification cost function we chose the cut-off for transforming predictions on the logit scale to the binary (dead, alive) scale based on the false positive rate (the rate at which the model predicted mortality incorrectly). Specifically, we chose the cut-off to keep this as small as reasonably possible.

3.4.A.5.2.2: Other modifications in the growth simulator

Thinning/Harvest

We assume that all thinning occur randomly. Random thinning can be implemented on an individual plot level. The CARBWARE user sets a basal area (BA) to be removed as stipulated in the harvest activity data (in the 'Eventstable') so thinning of trees are selected at random from the plot until this target BA is achieved. The thinned or harvested trees in a given plot are removed from the growth database and populated in a modifier table within the CarwKP_xx database. These data are then called up in the allocation module (see Fig 6.3.8 Ch6).

Although it is common practice that clear felled stands are replanted within 2 years, the CARBWARE model does not re-populate clear felled plots due to uncertainty of re-establishment success and species choice. This is a conservative approach and is consistent with the rules applied, which differentiate between deforestation and clear fell with re-establishment.

3.4.A.6 Adjustment of CSO statistics, affecting 5.B Cropland and 5.C Grassland

This Annex relates to adjustments made to the CSO crop area due to an inconsistency in the time series between 1996 and 1997. The text extracted from the NIR 2007 is relevant to the 2015 submission only insofar as it refers to areas of crops. The adjustment no longer has knock on implications for the Grassland land use category.

Extract from NIR 2007 Section 7.10 a

Adjustment of CSO statistics, affecting 5.B Cropland and 5.C Grassland

A review of the Central Statistics Office (CSO) data for areas under agricultural use identified an apparent mismatch between total farmed area reported for the period 1990-1996 and that for the period 1997-2004 (

Figure). The total net increase in area reported in agricultural usage in 1997 was approximately 89,000 ha. The increase was shared across most crops and grassland types and was contrary to the contemporary trends in the agricultural sector.

There was no change in the data compilation strategy or analysis methodology employed by the CSO at this time that would explain this increase. It is difficult to assign a single cause to the reported increase in agricultural area, however a number of developments in methods of farm subsidy payments and administration at this time may have encouraged certain farmers to revise the area of utilised agricultural land that they reported to the CSO. The assumption made in this analysis is that the additional lands reported as utilised agricultural land reflect a change in perception by the farmers of what constituted utilised lands rather than an actual change in land management. In effect, the additional lands were always available, and were already managed as agricultural lands. The CSO statistics available since 1997 are interpreted as a more complete record of agricultural lands, with the proviso that there is some agricultural grade lands unutilised and therefore unreported. In the 2006 analysis of LULUCF, these areas of agricultural land were included in the “Other Land” category up to 1997. This was inappropriate as the lands are impacted by agricultural activities, albeit on an irregular basis. It is therefore appropriate to adjust the CSO data for the years up to 1997 to include these less utilised agricultural lands.




Figure 3.4.A.6 Extracted from NIR 2007: Figure 7.1: Estimated area of Agricultural Land in Ireland 1991-2005

The increase in farmed area reported for 2007 was reported against a background of a downward trend reported in years up to 1997 and again in the years afterwards (

Figure). The average annual decrease in total farmed area from 1991-1996 was 20,000 ha (+/- 18,000 ha), or 16,800 ha annually from a linear fit trend. If farmers had used the older perception of land utilisation, the reported area for 1997 would have been of the order of 107,000 ha lower than given by CSO, showing a decrease of 16,000 ha from 1996. This represents an adjustment to reported areas of approximately 2.5 per cent. This adjustment factor is applied *pro rata* backwards to 1990, as shown on

Figure . The adjustment is provisional on the findings of future research analysis of the survey data collated at the time. Another effect of the adjustment is to create a greater continuity between area of land converted to forest lands and settlements and that lost from wetlands (and peatlands) and agricultural lands.

3.4.B Detailed Non-Forest Land Use Change Matrices

This annex supplements the material in Table 6.4 Summary Land Use Matrices 1990-2013. It provides detailed tables for annual estimates of area for the non-Forest Land, land use categories. Shown are estimates of gains and losses related to each land use type on an annual basis, the subsequent annual net change in area and the resultant cumulative total area under each category.

Also shown in the tables are the summary values for Remaining Land and In Transition which appear in the Common Reporting Format submission.

The tables in the annex demonstrate the consistency between Table 6.4 and the CRF figures. These tables are an important tool for QA/QC.

Table 3.4.B-1 Cropland Matrix

	Year	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
CRF	Cropland Remaining Cropland	700.01	698.40	696.46	694.76	693.13	691.15	688.75	686.60	685.41	684.07	682.74	681.13	679.61
CRF	Cropland in Transition Cumulative	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CRF	Total Cropland	424.65	404.36	401.39	402.62	413.89	410.20	409.28	416.35	414.30	408.10	401.00	401.10	415.50
	Forest to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Grassland to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Wetland to Cropland													
	Settlement to Cropland													
	Other land to Cropland													
	Total Gain in Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Cropland to Forest	1.51	1.58	1.91	1.67	1.60	1.95	2.37	2.10	1.14	1.29	1.27	1.57	1.46
	Cropland to Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Cropland to Wetland													
	Cropland to Settlement	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.05	0.05	0.06	0.05	0.07	0.06
	Cropland to Other land													
	Total Loss from Cropland	1.54	1.61	1.94	1.70	1.63	1.98	2.42	2.15	1.20	1.35	1.31	1.64	1.52
	Net Change Forest/Cropland	-1.51	-1.58	-1.91	-1.67	-1.60	-1.95	-2.37	-2.10	-1.14	-1.29	-1.27	-1.57	-1.46
	Net Change Grassland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Wetland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Settlement/Cropland	-0.03	-0.03	-0.03	-0.03	-0.03	-0.04	-0.04	-0.05	-0.05	-0.06	-0.05	-0.07	-0.06
	Net Change Otherland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Matrix Increment	Total Annual net Change	-1.54	-1.61	-1.94	-1.70	-1.63	-1.98	-2.42	-2.15	-1.20	-1.35	-1.31	-1.64	-1.52
Total Matrix	Total Cropland	700.01	698.40	696.46	694.76	693.13	691.15	688.75	686.60	685.41	684.07	682.74	681.13	679.61

Table 3.4.B-1 Cropland Matrix (continued)

	Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
CRF	Cropland Remaining Cropland	678.22	677.39	676.56	675.75	675.42	675.15	674.89	674.62	674.29	674.07	673.87	673.68
CRF	Cropland in Transition Cumulative	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CRF	Total Cropland	423.70	436.40	423.90	401.10	381.00	379.40	385.00	369.40	353.90	365.40	382.10	377.20
		678.22	677.39	676.56	675.75	675.42	675.15	674.89	674.62	674.29	674.07	673.87	673.68
	Forest to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Grassland to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Wetland to Cropland												
	Settlement to Cropland												
	Other land to Cropland												
	Total Gain in Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Cropland to Forest	1.33	0.75	0.75	0.72	0.24	0.22	0.19	0.20	0.25	0.20	0.18	0.18
	Cropland to Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Cropland to Wetland												
	Cropland to Settlement	0.08	0.08	0.10	0.09	0.06	0.07	0.07	0.09	0.02	0.02	0.01	0.00
	Cropland to Other land												
	Total Loss from Cropland	1.41	0.83	0.84	0.80	0.30	0.28	0.26	0.29	0.27	0.22	0.19	0.18
	Net Change Forest/Cropland	-1.33	-0.75	-0.75	-0.72	-0.24	-0.22	-0.19	-0.20	-0.25	-0.20	-0.18	-0.18
	Net Change Grassland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Wetland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Settlement/Cropland	-0.08	-0.08	-0.10	-0.09	-0.06	-0.07	-0.07	-0.09	-0.02	-0.02	-0.01	0.00
	Net Change Otherland/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Matrix Increment	Total Annual net Change	-1.41	-0.83	-0.84	-0.80	-0.30	-0.28	-0.26	-0.29	-0.27	-0.22	-0.19	-0.18
Annual Matrix	Total Cropland	678.22	677.39	676.56	675.75	675.42	675.15	674.89	674.62	674.29	674.07	673.87	673.68

Table 3.4.B-2 Grassland Matrix

	Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
CRF	Grassland Remaining Grassland	4144.2	4138.156	4132.860	4127.75	4121.5	4114.0	4107.2	4103.3	4098.8	4094.4	4089.1	4083.6
CRF	Grassland in Transition	0.069	0.138	0.207	0.276	0.345	0.642	0.938	1.235	1.532	1.829	2.304	2.779
CRF	Total Grassland	4144.3	4138.294	4133.0671	4128.030	4121.9	4114.7	4108.2	4104.5	4100.3	4096.2	4091.5	4086.4
	Forest to Grassland	8	8	8	8	8	267	267	267	267	267	400	400
	Cropland to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Wetland to Grassland	61	61	61	61	61	30	30	30	30	30	75	75
	Settlement to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Other land to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Total Gain in Grassland	69	69	69	69	69	297	297	297	297	297	475	475
	Grassland to Forest	4.75	5.74	5.01	4.80	5.84	7.11	6.29	3.43	3.88	3.80	4.71	4.83
	Grassland to Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Grassland to Wetland	0	0	0	0	0	0	0	0	0	0	0	0
	Grassland to Settlement	0.31	0.28	0.29	0.31	0.38	0.39	0.50	0.54	0.59	0.63	0.52	0.74
	Grassland to Other land	0	0	0	0	0	0	0	0	0	0	0	0
	Total Loss from Grassland	5	6	5	5	6	8	7	4	4	4	5	6
	Net Change Forest/Grassland	3	2	3	3	2	260	260	263	263	263	395	395
	Net Change Grassland/Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Wetland/Grassland	61	61	61	61	61	30	30	30	30	30	75	75
	Net Change Settlement/Grassland	0	0	0	0	0	0	0	-1	-1	-1	-1	-1
	Net Change Other land/Grassland	0	0	0	0	0	0	0	0	0	0	0	0
Matrix Increment	Total Annual net Change	64	63	64	64	63	289	290	293	292	292	470	469
Annual Matrix	Total Grassland Input	4144.3	4138.3	4133.1	4128.0	4121.9	4114.7	4108.2	4104.5	4100.3	4096.2	4091.5	4086.4

Table 3.4.B-2 Grassland Matrix (continued)

	Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
CRF	Grassland Remaining Grassland	4078.0	4074.1	4069.8	4065.0	4060.5	4056.6	4053.0	4049.3	4044.5	4041.3	4038.5	4035.6
CRF	Grassland in Transition	3.254	3.729	4.204	4.692	4.780	4.869	5.357	5.943	6.928	8.314	8.807	9.337
CRF	Total Grassland	4081.3	4077.8	4074.0	4069.7	4065.3	4061.5	4058.4	4055.2	4051.5	4049.6	4047.3	4044.9
	Forest to Grassland	400	400	400	400	0	0	400	400	800	1200	0	64
	Cropland to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Wetland to Grassland	75	75	75	88	88	88	88	186	186	186	493	465
	Settlement to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Other land to Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Total Gain in Grassland	475	475	475	489	88	88	488	586	986	1386	493	529
	Grassland to Forest	4.89	3.07	3.41	3.66	3.62	3.23	2.81	2.99	3.74	2.99	2.63	2.77
	Grassland to Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Grassland to Wetland	0	0	0	0	0	0	0	0	0	0	0	0
	Grassland to Settlement	0.66	0.85	0.90	1.09	0.95	0.64	0.76	0.78	0.97	0.22	0.19	0.13
	Grassland to Other land	0	0	0	0	0	0	0	0	0	0	0	0
	Total Loss from Grassland	6	4	4	5	5	4	4	4	5	3	3	3
	Net Change Forest/Grassland	395	397	397	397	-4	-3	397	397	796	1197	-3	61
	Net Change Grassland/Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Wetland/Grassland	75	75	75	88	88	88	88	186	186	186	493	465
	Net Change Settlement/Grassland	-1	-1	-1	-1	-1	-1	-1	-1	-1	0	0	0
	Net Change Otherland/Grassland	0	0	0	0	0	0	0	0	0	0	0	0
Matrix Increment	Total Annual net Change	469	471	471	484	84	84	485	582	981	1382	490	527
Annual Matrix	Total Grassland Input	4081.3	4077.8	4074.0	4069.7	4065.3	4061.5	4058.4	4055.2	4051.5	4049.6	4047.3	4044.9

Table 3.4.B-3 Wetland Matrix

	Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
CRF	Wetland Remaining Wetland	1,298.08	1,285.80	1,278.67	1,268.31	1,260.74	1,247.04	1,238.65	1,232.66	1,227.04	1,220.81	1,219.76	1,213.20
CRF	Wetland in Transition	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.34
CRF	Total Wetland	1,298.08	1,285.80	1,278.67	1,268.31	1,260.74	1,247.04	1,238.65	1,232.66	1,227.04	1,220.81	1,219.93	1,213.55
	Forest to Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.17
	Cropland to Wetland												
	Grassland to Wetland												
	Settlement to Wetland												
	Other land to Wetland												
	Total Gain in Wetland	0	0	0	0	0	0	0	0	0	0	0.1706	0.1706
	Wetland to Forest	9.02	10.91	9.52	9.12	11.09	13.51	11.96	6.52	7.37	7.22	8.95	8.66
	Wetland to Cropland												
	Wetland to Grassland	0.06	0.06	0.06	0.06	0.06	0.03	0.03	0.03	0.03	0.03	0.07	0.07
	Wetland to Settlement												
	Wetland to Other land												
	Total Loss from Wetland	9.08	10.97	9.58	9.18	11.15	13.54	11.99	6.55	7.40	7.25	9.02	8.74
	Net Change Forest/Wetland	-9	-11	-10	-9	-11	-14	-12	-7	-7	-7	-9	-8
	Net Change Wetland/Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Wetland/Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Settlement/Wetland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Otherland/Wetland	0	0	0	0	0	0	0	0	0	0	0	0
Matrix Increment	Total Annual net Change	-9	-11	-10	-9	-11	-14	-12	-7	-7	-7	-9	-9
Annual Matrix	Matrix Total Wetland Annual	1299	1288	1279	1270	1258	1245	1233	1226	1219	1212	1203	1194

Table 3.4.B-3 Wetland Matrix (continued)

	Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
CRF	Wetland Remaining Wetland	1,206.93	1,204.91	1,214.24	1,209.09	1,207.23	1,204.05	1,201.60	1,201.54	1,199.16	1,197.57	1,195.19	1,192.75
CRF	Wetland in Transition	0.51	0.68	0.85	1.02	1.02	1.42	1.42	1.82	1.82	2.22	2.22	2.22
CRF	Total Wetland	1,207.44	1,205.60	1,215.10	1,210.11	1,208.25	1,205.47	1,203.03	1,203.36	1,200.99	1,199.80	1,197.42	1,194.97
	Forest to Wetland	0.17	0.17	0.17	0.17	0.00	0.40	0.00	0.40	0.00	0.40	0.00	0.00
	Cropland to Wetland												
	Grassland to Wetland												
	Settlement to Wetland												
	Other land to Wetland												
	Total Gain in Wetland	0.1706	0.1706	0.1706	0.1706	0	0.4	0	0.4	0	0.4	0	0
	Wetland to Forest	8.28	4.91	5.16	5.25	3.62	3.23	2.81	2.99	3.74	2.99	2.63	2.77
	Wetland to Cropland												
	Wetland to Grassland	0.07	0.07	0.07	0.09	0.09	0.09	0.09	0.19	0.19	0.19	0.49	0.47
	Wetland to Settlement												
	Wetland to Other land												
	Total Loss from Wetland	8.35	4.99	5.24	5.34	3.70	3.32	2.90	3.18	3.93	3.18	3.13	3.24
	Net Change Forest/Wetland	-8	-5	-5	-5	-4	-3	-3	-3	-4	-3	-3	-3
	Net Change Wetland/Cropland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Wetland/Grassland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Settlement/Wetland	0	0	0	0	0	0	0	0	0	0	0	0
	Net Change Otherland/Wetland	0	0	0	0	0	0	0	0	0	0	0	0
Matrix Increment	Total Annual net Change	-8	-5	-5	-5	-4	-3	-3	-3	-4	-3	-3	-3
Annual Matrix	Matrix Total Wetland Annual	1186	1182	1177	1173	1170	1168	1166	1165	1162	1161	1162	1159

Table 3.4.B-4 Settlement Matrix

	Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
CRF	Settlement Remaining Settlement	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78
CRF	Settlement in Transition	0.37	0.70	1.04	1.41	1.86	2.31	2.88	3.51	4.19	4.92	5.69	6.71
CRF	Total Settlement	98.14	98.48	98.82	99.19	99.63	100.09	100.66	101.29	101.97	102.69	103.46	104.49
	Forest to Settlement	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.17	0.17
	Cropland to Settlement	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.05	0.05	0.06	0.05	0.07
	Grassland to Settlement	0.31	0.28	0.29	0.31	0.38	0.39	0.50	0.54	0.59	0.63	0.52	0.74
	Wetland to Settlement												
	Other land to Settlement	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.04	0.04	0.04	0.03	0.05
	Total Gain in Settlement	0.37	0.34	0.34	0.36	0.45	0.45	0.57	0.63	0.68	0.73	0.77	1.02
	Settlement to Forest												
	Settlement to Cropland												
	Settlement to Grassland												
	Settlement to Wetland												
	Settlement to Other land												
	Total Loss from Settlement	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Forest/Settlement	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.17	0.17
	Net Change Settlement/Cropland	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.05	0.05	0.06	0.05	0.07
	Net Change Settlement/Grassland	0.31	0.28	0.29	0.31	0.38	0.39	0.50	0.54	0.59	0.63	0.52	0.74
	Net Change Settlement/Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other land/Settlement	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.04	0.04	0.04	0.03	0.05
Matrix Increment	Total Annual net Change	0.37	0.34	0.34	0.36	0.45	0.45	0.57	0.63	0.68	0.73	0.77	1.02
Annual Matrix	Total Settlement Input	98.14	98.48	98.82	99.19	99.63	100.09	100.66	101.29	101.97	102.69	103.46	104.49

Table 3.4.B-4 Settlement Matrix (continued)

	Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
CRF	Settlement Remaining Settlement	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78	97.78
CRF	Settlement in Transition	7.65	8.80	10.01	11.44	12.94	14.88	16.96	17.86	18.99	19.25	20.26	20.47
CRF	Total Settlement	105.42	106.57	107.79	109.21	110.71	112.65	114.74	115.64	116.77	117.02	118.04	118.25
	Forest to Settlement	0.17	0.17	0.17	0.17	0.40	1.20	1.20	0.00	0.00	0.00	0.80	0.06
	Cropland to Settlement	0.06	0.08	0.08	0.10	0.09	0.06	0.07	0.07	0.09	0.02	0.02	0.01
	Grassland to Settlement	0.66	0.85	0.90	1.09	0.95	0.64	0.76	0.78	0.97	0.22	0.19	0.13
	Wetland to Settlement												
	Other land to Settlement	0.04	0.06	0.06	0.07	0.06	0.04	0.05	0.05	0.06	0.01	0.01	0.01
	Total Gain in Settlement	0.94	1.15	1.21	1.43	1.50	1.94	2.08	0.91	1.13	0.25	1.02	0.21
	Settlement to Forest												
	Settlement to Cropland												
	Settlement to Grassland												
	Settlement to Wetland												
	Settlement to Other land												
	Total Loss from Settlement	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Forest/Settlement	0.17	0.17	0.17	0.17	0.40	1.20	1.20	0.00	0.00	0.00	0.80	0.06
	Net Change Settlement/Cropland	0.06	0.08	0.08	0.10	0.09	0.06	0.07	0.07	0.09	0.02	0.02	0.01
	Net Change Settlement/Grassland	0.66	0.85	0.90	1.09	0.95	0.64	0.76	0.78	0.97	0.22	0.19	0.13
	Net Change Settlement/Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Otherland/Settlement	0.04	0.06	0.06	0.07	0.06	0.04	0.05	0.05	0.06	0.01	0.01	0.01
Matrix Increment	Total Annual net Change	0.94	1.15	1.21	1.43	1.50	1.94	2.08	0.91	1.13	0.25	1.02	0.21
Annual Matrix	Total Settlement Input	105.42	106.57	107.79	109.21	110.71	112.65	114.74	115.64	116.77	117.02	118.04	118.25

Table 3.4.B-5 Other Land Matrix

	Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
CRF	Other Land Remaining Other Land	390.57	389.97	389.45	388.95	388.35	387.61	386.95	386.57	386.14	385.72	385.22	384.48
CRF	Other Land in Transition	0.00	0.00	0.01	0.01	0.01	0.08	0.14	0.21	0.28	0.34	0.46	0.57
CRF	Total Other Land	390568.7	389978.1	389460.7	388962.9	388356.60	387686.34	387091.11	386779.17	386419.60	386065.11	385675.45	385056.10
	Forest to Other Land	0.00	0.00	0.00	0.00	0.00	0.07	0.07	0.07	0.07	0.07	0.11	0.11
	Cropland to Other Land												
	Grassland to Other Land												
	Wetland to Other Land												
	Settlement to Other Land												
	Total Gain in Other Land	0.00	0.00	0.00	0.00	0.00	0.07	0.07	0.07	0.07	0.07	0.11	0.11
	Other Land to Forest	0.47	0.57	0.50	0.48	0.58	0.71	0.63	0.34	0.39	0.38	0.47	0.52
	Other Land to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Settlement	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.04	0.04	0.04	0.03	0.05
	Total Loss from Other Land	0.49	0.59	0.52	0.50	0.61	0.74	0.66	0.38	0.43	0.42	0.50	0.56
	Net Change Forest/Other Land	-0.47	-0.57	-0.50	-0.48	-0.58	-0.64	-0.56	-0.28	-0.32	-0.31	-0.36	-0.40
	Net Change Other Land/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other Land/Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other land/Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other land/Settlement	-0.02	-0.02	-0.02	-0.02	-0.02	-0.03	-0.03	-0.04	-0.04	-0.04	-0.03	-0.05
Matrix Increment	Total Annual net Change	-0.49	-0.59	-0.52	-0.50	-0.61	-0.67	-0.60	-0.31	-0.36	-0.35	-0.39	-0.45
Annual Matrix	Total Other Land	390.57	389.98	389.46	388.96	388.36	387.69	387.09	386.78	386.42	386.07	385.68	385.23

Table 3.4.B-5 Other Land Matrix (continued)

	Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
CRF	Other Land Remaining Other Land	383.55	382.62	381.45	380.06	378.41	376.84	374.93	372.99	370.52	368.22	364.78	362.02
CRF	Other Land in Transition	0.69	0.80	0.92	1.03	2.63	2.63	3.03	3.03	3.03	3.03	3.03	3.02
CRF	Total Other Land	384234.5	383418.4	382370.0	381089.86	381041.68	379473.96	377963.3	376022.60	373553.41	371249.74	367804.69	365044.55
	Forest to Other Land	0.11	0.11	0.11	0.11	1.60	0.00	0.40	0.00	0.00	0.00	0.00	0.00
	Cropland to Other Land												
	Grassland to Other Land												
	Wetland to Other Land												
	Settlement to Other Land												
	Total Gain in Other Land	0.11	0.11	0.11	0.11	1.60	0.00	0.40	0.00	0.00	0.00	0.00	0.00
	Other Land to Forest	0.55	0.36	0.42	0.47	0.56	0.50	0.44	0.47	0.58	0.47	0.41	0.43
	Other Land to Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Other Land to Settlement	0.04	0.06	0.06	0.07	0.06	0.04	0.05	0.05	0.06	0.01	0.01	0.01
	Total Loss from Other Land	0.60	0.42	0.48	0.54	0.62	0.54	0.49	0.52	0.65	0.48	0.42	0.44
	Net Change Forest/Other Land	-0.44	-0.25	-0.31	-0.36	1.04	-0.50	-0.04	-0.47	-0.58	-0.47	-0.41	-0.43
	Net Change Other Land/Cropland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other Land/Grassland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other land/Wetland	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	Net Change Other land/Settlement	-0.04	-0.06	-0.06	-0.07	-0.06	-0.04	-0.05	-0.05	-0.06	-0.01	-0.01	-0.01
Matrix Increment	Total Annual net Change	-0.48	-0.30	-0.37	-0.43	0.98	-0.54	-0.09	-0.52	-0.65	-0.48	-0.42	-0.44
Annual Matrix	Total Other Land	384.75	384.44	384.08	383.65	384.62	384.08	383.99	383.48	382.83	382.35	381.93	381.49

Annex 3.5

Waste (IPCC Sector 5)

3.5.A Summary of Parameter Input Values to Estimate Methane Generation using 2006 IPCC Guidelines Model

3.5.B Time Series of Solid Waste Disposal and Composition 1990-2013

3.5.C Parameters, EFs for Clinical Waste Incineration 1990-2013

3.5.D Parameters, EFs for Solvent (Liquid/Vapour destruction) Waste Incineration 1990-2013

Table 3.5.A Summary of Parameter Input Values to Estimate Methane Generation using 2006 IPCC Guidelines Model

Model Run	Grouping or IPCC licence number	Number of Sites	Active Period	Status in 2013	MSW Total (t)	MSW 2012 (t)	^a DOC Fraction	DOC _f	^b Decay Rate k	^c MCF
1	From 1969	13	1956-2012	Open	7,547,065	106,724	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.40-0.67, 1.0
2	From 1979	10	1972-2012	Open	7,275,047	144,705	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.40-0.67, 1.0
3	1985-2002	5	1983-2002	Closed	2,190,371	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.40-0.67, 1.0
4	Small Closed	9	1957-2003	Closed	2,602,723	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.40-0.67, 1.0
5	Recent Closed	16	1975-2008	Closed	7,486,891	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.40-0.67, 1.0
6	W0004	1	1997-2010	Closed	4,790,944	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.75	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	1.0
7	W0127	1	1976-2001	Closed	4,812,569	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.8,1.0
8	W0009	1	1971-2012	Open	3,595,721	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.8,1.0
9	W0012	1	1965-2009	Closed	2,675,434	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.8,1.0
10	W0081	1	1999-2011	Closed	2,015,205	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.75	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	1.0
11	W0047	1	1996-2010	Closed	2,123,265	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.6
12	W0201	1	2008-2012	Open	1,573,181	330,544	0.20, 0.22, 0.45, 0.46, 0.40, 0.32	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	1.0
13	New Sites	8	1995-2012	Open	5,044,835	187,100	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	1.0
14	Town Dumps	~250	1956-1998	Closed	15,372,064	NO	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.4
15	Sewage Sludge	1	1990-2012	Open	322,651	2,721	0.05	0.60	0.20	0.8,1.0
16	Street Cleanings	1	1990-2012	Open	1,694,181	41,379	0.15, 0.20, 0.40, 0.43, 0.24, 0.24	0.60	0.20, 0.10, 0.07, 0.04, 0.07, 0.10	0.8,1.0

a The six values are for food, garden, paper, wood and straw, textiles and disposable nappies

b The six values are for food, garden, paper, wood and straw, textiles, disposable nappies

c Where two values are given, the first is for years up to 1998 (pre landfill licensing) and the second is for subsequent years

Table 3.5.B Time Series of Solid Waste Disposal and Composition 1990-2013

Year	Pop	MSW Prod Rate kg/cap/day	MSW Managed tonnes	MSW to SWDS %	MSW to SWDS tonnes	Street Cleansing tonnes	Sewage Sludge tonnes	MSW Food %	MSW Garden %	MSW Paper %	MSW Wood %	MSW Textiles %	MSW Nappies %	MSW Other %	MSW Organic tonnes	MSW Garden tonnes	MSW Paper tonnes	MSW Wood tonnes	MSW Textiles tonnes	MSW Nappies ^a tonnes	MSW Other tonnes
1990	3,505,800	1.8	2,037,978	0.92	1,878,358	46,959	16,307	39.3%	0.0%	29.5%	5.2%	9.8%	0.0%	16.3%	756,805		567,893	99,436	187,836		313,348
1991	3,525,700	1.7	2,020,416	0.92	1,862,172	46,554	16,400	37.7%	0.0%	29.6%	5.2%	8.0%	0.0%	19.5%	720,201		565,619	98,579	152,700		371,628
1992	3,554,500	1.7	2,000,684	0.92	1,843,986	46,100	16,533	36.2%	0.0%	29.8%	5.2%	6.2%	0.0%	22.7%	683,379		562,688	97,616	118,019		428,384
1993	3,574,100	1.7	2,026,906	0.92	1,868,154	46,704	16,625	34.6%	0.0%	29.9%	5.2%	4.5%	0.0%	25.9%	662,157		572,689	98,895	85,941		495,175
1994	3,585,900	1.7	2,136,712	0.92	1,969,360	49,234	16,680	33.0%	0.0%	30.0%	5.2%	2.7%	0.0%	29.1%	666,215		606,484	104,253	55,151		586,491
1995	3,601,300	1.7	2,067,106	0.92	1,905,206	46,791	16,878	31.4%	0.0%	30.2%	4.5%	1.0%	1.0%	31.9%	613,637		589,328	88,157	19,063	19,063	622,750
1996	3,626,100	1.8	2,469,888	0.92	2,266,842	58,194	16,739	28.8%	0.0%	27.9%	3.9%	0.9%	0.9%	37.6%	669,951		648,655	90,924	21,594	20,209	873,702
1997	3,664,300	1.9	2,538,442	0.91	2,319,897	69,596	18,243	27.3%	0.0%	28.4%	3.4%	1.0%	0.9%	39.1%	653,010		678,856	80,368	23,365	20,522	933,372
1998	3,703,100	1.9	2,323,692	0.91	2,114,605	80,999	24,213	25.7%	0.0%	28.5%	2.9%	1.0%	0.8%	41.1%	564,568		625,733	63,028	22,164	18,318	901,793
1999	3,741,600	1.9	2,039,195	0.90	1,826,474	80,156	26,710	25.8%	0.0%	27.8%	2.4%	1.1%	0.9%	42.1%	491,443		529,580	45,562	21,303	16,564	802,178
2000	3,789,500	1.9	2,395,557	0.88	2,111,319	79,312	31,811	24.1%	0.0%	30.3%	1.7%	1.3%	0.9%	41.6%	528,734		664,771	37,818	28,065	20,740	910,503
2001	3,847,200	1.9	2,558,726	0.87	2,218,447	78,469	26,253	24.8%	0.0%	30.1%	1.1%	1.4%	1.0%	41.6%	569,674		690,623	25,648	32,799	23,215	954,957
2002	3,917,200	1.9	2,684,314	0.79	2,128,259	65,573	21,605	25.6%	0.0%	28.9%	0.9%	1.5%	1.1%	42.1%	561,679		634,604	18,701	31,836	24,188	922,824
2003	3,979,900	2.0	2,717,740	0.72	1,946,012	71,779	16,111	25.9%	0.0%	30.7%	0.8%	1.5%	1.1%	40.0%	522,038		618,804	15,439	30,580	23,008	807,922
2004	4,045,200	2.0	2,823,173	0.66	1,875,428	26,344	10,696	29.3%	3.5%	26.9%	2.2%	4.2%	3.5%	30.3%	557,429	67,351	512,230	41,958	80,332	66,674	575,799
2005	4,133,800	2.0	2,875,399	0.65	1,880,955	23,875	5,161	29.2%	3.5%	26.8%	4.9%	3.8%	3.5%	28.3%	556,859	66,423	511,214	93,091	71,690	67,383	538,170
2006	4,232,900	2.2	3,202,322	0.64	2,045,788	30,366	2,672	29.5%	3.8%	25.6%	1.3%	4.0%	3.3%	32.4%	613,413	79,162	531,462	26,737	83,558	69,237	672,586
2007	4,375,800	2.1	3,316,376	0.63	2,104,800	70,334	2,897	35.5%	3.2%	20.6%	1.8%	6.1%	4.1%	28.7%	771,665	70,182	449,152	39,439	133,109	88,219	623,368
2008	4,485,100	2.0	3,182,048	0.62	1,987,575	24,969	3,909	29.7%	3.3%	23.3%	1.9%	5.6%	4.4%	31.7%	598,525	66,578	469,169	37,873	113,599	88,632	638,167
2009	4,533,400	1.8	2,895,940	0.61	1,767,004	26,701	4,229	30.9%	3.8%	20.8%	1.5%	6.2%	4.9%	31.9%	553,485	68,386	373,199	26,883	111,295	88,504	571,952
2010	4,554,800	1.7	2,548,148	0.58	1,476,852	18,713	3,079	31.8%	3.7%	21.7%	0.9%	6.2%	4.7%	31.0%	475,230	54,650	324,252	14,016	92,463	70,663	464,291
2011	4,574,900	1.7	2,498,884	0.53	1,318,837	22,291	3,096	31.4%	3.4%	21.8%	1.1%	6.7%	4.5%	31.0%	421,250	45,743	292,830	14,841	90,130	60,065	415,284
2012	4,585,400	1.6	2,584,112	0.38	986,198	41,379	3,081	31.1%	3.5%	21.2%	0.9%	6.0%	4.6%	32.6%	319,687	36,457	217,805	9,241	61,785	47,132	335,470
2013	4,593,100	1.6	2,015,183	0.38	769,072	41,379	2,721	30.8%	3.5%	21.1%	0.9%	5.9%	4.5%	33.2%	249,687	28,431	170,702	7,266	48,182	36,757	269,426

^a Nappies are assumed to be included in the textiles proportion during the period 1990-1995 inclusive.

Table 3.5.C Parameters, EFs for Clinical Waste Incineration 1990-2013

	Quantity of Clinical Waste (SW _{CW})	Dry Matter content of Clinical Waste (dm _{CW})	Fraction of Carbon in the dry matter as % (CF _{CW})	Fraction of fossil carbon in total carbon as % (FCF _{CW})	Oxidation Factor (OF _{CW})	Emissions CO ₂ (Fossil)
	kt					kt CO ₂
	A	B	C	D	E	= A * B * C * D * E * 44/12
1990	4.0	NA	60.00	40.00	1.000	3.52
1991	4.0	NA	60.00	40.00	1.000	3.52
1992	4.0	NA	60.00	40.00	1.000	3.52
1993	4.0	NA	60.00	40.00	1.000	3.52
1994	4.0	NA	60.00	40.00	1.000	3.52
1995	4.0	NA	60.00	40.00	1.000	3.52
1996	4.0	NA	60.00	40.00	1.000	3.52
1997	4.0	NA	60.00	40.00	1.000	3.52
1998	NO	NA	60.00	40.00	1.000	NO
1999	NO	NA	60.00	40.00	1.000	NO
2000	NO	NA	60.00	40.00	1.000	NO
2001	NO	NA	60.00	40.00	1.000	NO
2002	NO	NA	60.00	40.00	1.000	NO
2003	NO	NA	60.00	40.00	1.000	NO
2004	NO	NA	60.00	40.00	1.000	NO
2005	NO	NA	60.00	40.00	1.000	NO
2006	NO	NA	60.00	40.00	1.000	NO
2007	NO	NA	60.00	40.00	1.000	NO
2008	NO	NA	60.00	40.00	1.000	NO
2009	NO	NA	60.00	40.00	1.000	NO
2010	NO	NA	60.00	40.00	1.000	NO
2011	NO	NA	60.00	40.00	1.000	NO
2012	NO	NA	60.00	40.00	1.000	NO
2013	NO	NA	60.00	40.00	1.000	NO

$$\text{Equation 5.1, 2006 IPCC Guidelines: CO}_2 \text{ emissions} = \sum i (SW_i * dm_i * CF_i * FCF_i * OF_i) * \frac{44}{12}$$

i, type of waste incinerated (CW: Clinical Waste)

Table 3.5.D Parameters, EFs for Solvent (Liquid/Vapour destruction) Waste Incineration 1990-2013

	Quantity of Fossil Liquid Waste (SW _i)	Dry Matter content of Fossil Liquid Waste (dm _i)	Fraction of Carbon in the dry matter as % (CF _i)	Fraction of fossil carbon in total carbon as % (FCF _i)	Oxidation Factor (OF _i)	Emissions CO ₂
	kt					kt CO ₂
	A	B	C	D	E	= A * B * C * D * E * 44/12
1990	27.084	NA	80.00	100.00	1.000	79.446
1991	27.084	NA	80.00	100.00	1.000	79.446
1992	27.084	NA	80.00	100.00	1.000	79.446
1993	27.084	NA	80.00	100.00	1.000	79.446
1994	27.084	NA	80.00	100.00	1.000	79.446
1995	27.084	NA	80.00	100.00	1.000	79.446
1996	27.084	NA	80.00	100.00	1.000	79.446
1997	22.482	NA	80.00	100.00	1.000	65.947
1998	17.880	NA	80.00	100.00	1.000	52.448
1999	18.940	NA	80.00	100.00	1.000	55.557
2000	20.000	NA	80.00	100.00	1.000	58.667
2001	21.491	NA	80.00	100.00	1.000	63.040
2002	21.830	NA	80.00	100.00	1.000	64.035
2003	32.821	NA	80.00	100.00	1.000	96.275
2004	37.415	NA	80.00	100.00	1.000	109.751
2005	36.229	NA	80.00	100.00	1.000	106.270
2006	35.042	NA	80.00	100.00	1.000	102.790
2007	27.970	NA	80.00	100.00	1.000	82.045
2008	20.898	NA	80.00	100.00	1.000	61.301
2009	21.298	NA	80.00	100.00	1.000	62.474
2010	18.177	NA	80.00	100.00	1.000	53.319
2011	12.615	NA	80.00	100.00	1.000	37.004
2012	15.129	NA	80.00	100.00	1.000	44.378
2013	14.445	NA	80.00	100.00	1.000	42.372

$$\text{Equation 5.1, 2006 IPCC Guidelines: CO}_2 \text{ emissions} = \sum i (SW_i * dm_i * CF_i * FCF_i * OF_i) * \frac{44}{12}$$

Annex 4

Ireland's Energy Balance 1990-2013

4.A Ireland's Energy Balance - Stakeholders, Surveys and Sources

4.B Expanded Energy Balance sheets for 2013

4.C Country specific carbon emission factors – fossil fuels

4.A Ireland's Energy Balance - Stakeholders, Surveys and Sources

Sustainable Energy Authority of Ireland (SEAI)

The Sustainable Energy Authority of Ireland was established as Ireland's national energy authority under the Sustainable Energy Act 2002. SEAI's mission is to play a leading role in transforming Ireland into a society based on sustainable energy structures, technologies and practices. To fulfil this mission SEAI aims to provide well-timed and informed advice to Government, and deliver a range of programmes efficiently and effectively, while engaging and motivating a wide range of stakeholders and showing continuing flexibility and innovation in all activities.

SEAI has a lead role in developing and maintaining comprehensive national and sectoral statistics for energy production, transformation and end use. This data is a vital input in meeting international reporting obligations, for advising policy makers and informing investment decisions. The Energy Policy Statistical Support Unit (EPSSU) is SEAI's specialist statistics team.

Its core functions are to:

- Collect, process and publish energy statistics to support policy analysis and development in line with national needs and international obligations;
- Conduct statistical and economic analyses of energy services sectors and sustainable energy options;
- Contribute to the development and promulgation of appropriate sustainability indicators.

National Legislation

- Sustainable Energy Act 2002.
<http://www.irishstatutebook.ie/pdf/2002/en.act.2002.0002.pdf>
- European Communities (Energy End-use Efficiency and Energy Services) Regulations 2009, (S.I. No. 542 of 2009).
<http://www.irishstatutebook.ie/pdf/2009/en.si.2009.0542.pdf>
- Sulphur Content of Heavy Fuel, Gas Oil, and Marine Fuel Regulations 2008, (S.I. 119 of 2008).
<http://www.irishstatutebook.ie/pdf/2008/en.si.2008.0119.pdf>

EU Legislative Requirements

- Under the European Energy Statistics Regulation of 2008, no.1099, Ireland is legally obliged to submit energy statistics to Eurostat. The Regulation came into force on 1st January 2009 and SEAI are collecting data on behalf of Ireland from this date. SEAI submit annual and monthly energy statistics to Eurostat on energy supply, transformation and end-use for solid fuels, natural gas, electricity & heat and renewables & wastes (Oil statistics are supplied by DCENR). This data is also used for Ireland's Energy Balance.
<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32008R1099:EN:NOT>
- Information regarding gas and electricity prices is sent to Eurostat twice a year under the EU Gas and Electricity Price Transparency Directive 90/377/EEC.
http://europa.eu/legislation_summaries/other/l27002_en.htm

Department of Communications, Energy and Natural Resources (DCENR) - Oil Security Division

The Oil Security Division of the Department of Communications, Energy and Natural Resources is responsible for the development and implementation of Ireland's strategic oil supply policy, with particular regard to the areas of contingency planning and Ireland's obligations under the EU and International Energy Agency (IEA), in order to ensure an effective system of security of supply at times of physical oil supply disruption.

The National Oil Reserves Agency (NORA) is responsible for holding Ireland's strategic oil stocks for use in the event of a supply disruption. NORA is funded by a levy on disposals of petroleum products currently 2 cents per litre.

Oil Security Division collects monthly returns from oil companies and consumers on disposals of petroleum products and calculates the levy liability of each company. This is done under the NORA Act 2007 and associated returns and levy Regulations. A full list of legislation is available on the NORA website at www.nora.ie

Oil Security Division also provides Monthly Oil Statistics to the IEA and Eurostat.

Solid Fuels & Petroleum Coke

Fuels

- Bituminous Coal
- Anthracite
- Manufactured Ovoids
- Lignite
- Milled & Sod Peat
- Peat Briquettes
- Petroleum Coke

Frequency

- Monthly solid fuel survey
- Annual CHP Survey

Data Sources

This data collection is a monthly survey of solid fuel imports and producers to obtain solid fuel statistics as required under the European Energy Statistics Regulation of 2008, no.1099. This data is aggregated for the annual Energy Balance.

In the Energy Balance, anthracite and manufactured ovoids are combined to protect confidentiality.

Estimations

Smaller solid fuel distributors are not surveyed. To cover these smaller units, a technique was developed in 2009 to estimate the total data for the smaller units. The estimation method used data from Ireland's National Statistical Institute (CSO) monthly trade statistics publication to identify overall solid fuel imports which in turn highlighted areas that were missing.

Sod Peat is currently estimated.

Validation

These data are validated against the Emissions Trading Scheme data once it becomes available and there is a data point match. If there is a discrepancy between the ETS figure and the survey figure, the ETS figure is used as this has been audited by the Environmental Protection Agency.

Sectoral Breakdown

The sectoral breakdown is sourced from the monthly solid fuel surveys, except for industry where this is sourced from ETS data for bituminous coal, milled peat and petroleum coke.

Oil & Biofuels

Fuels

- Crude Oil
- Refinery Gas
- Gasoline
- Kerosene
- Jet Kerosene
- Fueloil
- LPG
- Gasoil/Diesel/DERV
- Liquid Biofuel
 - Bioethanol
 - Biodiesel

Frequency

- Monthly oil & biofuels survey
- Annual CHP Survey

Data Sources

Oil data is collected monthly by the Department of Communications, Environment and Natural Resources (DCENR). Oil companies are required to report to DCENR under the National Oil Reserves Agency Act 2007 (No. 7 of 2007) and the National Oil Reserves Agency Act 2007 (Returns and Levy) Regulations 2007 (S.I. 567 of 2007). Each oil company sends their monthly return to DCENR in a prescribed Excel format. This data is then analysed and manually transferred to a single monthly Excel sheet called the OCS system. All fuels are collected in litres, except for LPG which is collected in tonnes. In March each year, DCENR provide SEAI with all twelve OCS Excel sheets from the previous year. From the 2013 data collection a new online database, OLA, was rolled out to all companies. Data will be drawn from the OLA system rather than the OCS Excel spread sheets and will be provided to SEAI on this basis.

Liquid Biofuel data is collected monthly by DCENR under the Biofuel Obligation Scheme introduced in the Energy (Biofuel Obligation and Miscellaneous Provisions) Act 2010 and is provided to SEAI on an annual basis.

Revenue excise data on oil (litres) is provided on a monthly basis to SEAI.

Validation

Oil data are validated against the Emissions Trading Scheme data once it becomes available and there is a data point match. If there is a discrepancy between the ETS figure and the survey figure, the ETS figure is used as this has been audited by the Environmental Protection Agency.

Sectoral Breakdown

Census of Industrial Production data may be used for the industry breakdown. A joint CSO/SEAI Business Energy Use Survey (BEUS) was introduced recently in order to address energy consumption in the commercial and industry sectors. The survey results are not available yet but will eventually replace the CIP when calculating the breakdown of the industry sub-sectors.

The TFC is split further as follows:

Gasoline

- Total TFC sourced from Revenue excise data
- Transport
 - Road Private Car – calculated from vehicle stock and average annual consumption
 - Public Passenger Services – calculated from vehicle stock and average annual consumption
 - Domestic Aviation – sourced from Revenue excise data
 - Fuel Tourism - estimates provided by the Department of the Environment, Community & Local Government
 - Unspecified - remainder

Kerosene

- Total TFC sourced from Revenue excise data
- Industry
 - Estimated as 10% of TFC
 - Industry sub sectors are calculated using the Heavy Fuel Oil split from the 2009 CIP
 - The new BEUS will improve on this estimate
- Residential
 - Estimated as 90% of TFC
 - The new BEUS will improve on this estimate

Jet Kerosene

- Total TFC sourced from OCS system
- Transport
 - Domestic Aviation – Split based on EPA modelling
 - International Aviation – Split based on EPA modelling

Fueloil

- Total TFC sourced from Revenue excise data plus ETS data
- Industry – total less 10 kilotonnes used in Commercial/Public Services sector
 - Total Basic metals and fabricated metal comes from ETS data.
 - The remaining sub sectors are calculated using the Heavy Fuel Oil split from the 2009 CIP after subtracting Basic metals

- The new BEUS will improve on this estimate
- Commercial/Public Services – estimated as 10 kilotonnes

LPG

- Total TFC sourced from OCS system
- Industry
 - Total industry based on supplier split from OCS sheets
 - Industry sub sectors calculated using the Derived Gas split from the 2009 CIP
- Transport
 - Sourced from Revenue data as a differential excise duty is charged
- Residential
 - Total residential based on supplier split from OCS sheets
- Commercial/Public Services
 - Total commercial/public services based on supplier split from OCS sheets

Gasoil/Diesel/Derv

- Industry
 - Total industry estimated based on 1990 sector split (15% of Revenue gasoil less fuel input for electricity generation)
 - Industry sub sectors are calculated using the Gas Oil split from the 2009 CIP
 - The new BEUS will improve on this estimate
- Transport
 - Road Freight – calculated from CSO tonne kilometres and European data on energy use per tonne kilometre
 - Road Private Car – calculated from vehicle stock and average annual consumption
 - Public Passenger Services – calculated from vehicle stock and average annual consumption plus Revenue excise data for buses
 - Rail – Provided by the rail network operator
 - Fuel Tourism - estimates provided by the Department of the Environment, Community & Local Government
 - Sourced from Revenue data as a differential excise duty is charged
 - Unspecified - remainder
- Residential
 - Total residential estimated based on 1990 sector split (19% of Revenue gasoil)
 - The new BEUS will improve on this estimate
- Commercial/Public Services
 - Total commercial/public services estimated based on 1990 sector split (44% of Revenue gasoil), less navigation and fisheries
 - The new BEUS will improve on this estimate
- Agriculture
 - Total agriculture estimated based on 1990 sector split (21% of Revenue gasoil)
- Fisheries
 - Sourced from Revenue data as a differential excise duty is charged

Natural Gas

Frequency

Monthly and annual surveys of Bord Gáis Networks

Annual CHP Survey

Data Sources

Natural gas data is collected monthly and annually from Bord Gáis Networks. Bord Gáis Networks own, operate, build and maintain the natural gas network in Ireland and connect all customers to the network.

Supply data are collected in cubic metres and in gross energy units (Terajoules). Data on the demand side are received in TJ only.

Validation

Data are validated against the Emissions Trading Scheme data once it becomes available and there is a data point match. If there is a discrepancy between the ETS figure and the survey figure, the ETS figure is used as this has been audited by the Environmental Protection Agency.

Sectoral Breakdown

Census of Industrial Production data are used for the industry breakdown. A joint CSO/SEAI Business Energy Use Survey was introduced recently in order to address energy consumption in the commercial and industry sectors. The survey results are not available yet but will eventually replace the CIP when calculating the breakdown of the industry sub-sectors.

Renewables & Non-Renewable Waste

Fuels

- Wind
- Hydro
- Biomass
- Renewable Waste
- Landfill Gas
- Biogas
- Solar
 - Thermal
 - Photovoltaic
- Geothermal
- Non-Renewable Waste

Frequency

Annual renewable surveys

Annual CHP survey

Data Sources

Wind & Hydro

- Sourced from monthly electricity surveys – see electricity data collection

Wind (auto generation)

- Wind auto production data is sourced from annual surveys of the auto producers.

Biomass, Renewable Waste & Non-Renewable Waste

- Wood suppliers are surveyed annually; however there is usually a high non-response rate.
- Residential non-traded wood is calculated using estimation techniques.
- Boardmills and the major sawmills that use wood waste for energy are also surveyed. The remaining smaller sawmills are estimated.
- The Environmental Protection Agency provides administrative data on some renewable and non-renewable waste.
- The Department of Agriculture, Food and the Marine provide administrative data on tallow used for energy purposes.
- Biomass and waste data are validated against the Emissions Trading Scheme data once it becomes available and there is a data point match. If there is a discrepancy between the ETS figure and the survey figure, the ETS figure is used as this has been audited by the Environmental Protection Agency.

Landfill Gas

- Landfill Gas data is sourced from annual surveys of landfill gas operators and from administrative data provided by the Environmental Protection Agency.

Biogas

- Biogas data is sourced from annual surveys of sludge biogas installations and other biogas installations in Ireland

Solar

- Solar thermal contribution to energy in Ireland is calculated on an annual basis from administrative data. Data for retrofits on older buildings comes from government grant schemes administered by SEAI since 2006 both for residential and commercial properties. Solar statistics on new residential buildings in Ireland is sourced from the Building Energy Rating system which is also administered by SEAI.

Geothermal

- Geothermal contribution to energy in Ireland is calculated on an annual basis from administrative data. Data for retrofits on older buildings comes from government grant schemes administered by SEAI since 2006 both for residential and commercial properties. However, the residential grants ended in 2011 and SEAI have been working on a new source for this data.

Electricity

Frequency

Monthly electricity generator survey

Monthly TSO survey

Quarterly electricity retail market reports

Data Sources

Electricity Supply data is collected through a monthly survey of all electricity generators and the Transmission System Operator (Eirgrid) as required under the European Energy Statistics Regulation of 2008, no.1099. This data is aggregated for the annual Energy Balance.

The electricity generator survey is a business survey of all of the main activity electricity producers in Ireland. The Transmission System Operator survey is an administrative survey as this is data collected or generated by the TSO.

Since 2012, electricity consumption data is sourced from the quarterly Electricity and Gas Retail Markets Annual Report which is published by the Commission for Energy Regulation. Prior to this, each electricity supplier was surveyed annually for consumption data.

Validation

The electricity generator data are validated against the Emissions Trading Scheme data on fuel inputs once available. If there is a discrepancy between the ETS figure and the survey figure, the ETS figure is used as this has been audited by the Environmental Protection Agency.

Sectoral Breakdown

Census of Industrial Production data are used for the industry breakdown. A joint CSO/SEAI Business Energy Use Survey was introduced recently in order to address energy consumption in the commercial and industry sectors. The survey results are not available yet but will eventually replace the CIP when calculating the breakdown of the industry sub-sectors.

Table 4.B Expanded Energy Balance Sheet 2013

2013	Units = ktoe	Coal	Bituminous Coal	Anthracite + Manufactured Ovoids	Coke	Lignite	Peat	Milled Peat	Sod Peat	Briquettes	Oil	Crude	Refinery Gas	Gasoline	Kerosene	Jet Kerosene	Fueloil	LPG	Gasoil / Diesel /DERV	Petroleum Coke	Naphta	Bitumen	White Spirit	Lubricants	Natural Gas	Renewables	Hydro	Wind	Biomass & Renewable Waste	Landfill Gas	Biogas	Liquid Biofuel	Solar	Geothermal	Non-Renewable Waste	Electricity	Heat	TOTAL	
Indigenous Production		0	0				1,292	1,164	128		0														151	798	50	391	244	37	11	22	11	33	58			2,299	
Imports		1,496	1,407	67		22	0				7,973	3,225	0	672	427	887	181	103	2,077	112	0	251	1	38	3,717	114			36			79				216		13,516	
Exports		10	0	10		1	9			9	1,212	227		13	4	0	901	31	0	0	28	0	0	7	0	0			0			0				33		1,264	
Mar. Bunkers		0					0				143						18		125							0												143	
Stock Change		-162	-147	-11		-4	-560	-547	0	-13	-73	-97		-5	19	5	-2	-2	-4	11	1	0	0	0	4	-2			-1			-1						-793	
Primary Energy Supply (incl non-energy)		1,324	1,260	46	0	17	723	617	128	-21	6,545	2,900	0	654	441	891	-739	69	1,948	124	-27	251	1	31	3,872	911	50	391	279	37	11	99	11	33	58	182	0	13,615	
Primary Energy Requirement (excl. non-energy)		1,324	1,260	46	0	17	723	617	128	-21	6,262	2,900	0	654	441	891	-739	69	1,948	124	-27	0	0	0	3,872	911	50	391	279	37	11	99	11	33	58	182	0	13,332	
Transformation Input		970	970	0	0	0	639	639	0	0	2,946	2,902	5	0	0	0	33	1	4	0	0	0	0	0	2,155	118	0	0	77	37	4	0	0	0	23	68	0	6,918	
Public Thermal Power Plants		970	970				498	498	0		38						33		4						1,830	110			73	37					23			3,468	
Combined Heat and Power Plants		0	0				9	9			6		5				0	1	0						268	8			4		4							292	
Pumped Storage Consumption																																				50		50	
Briquetting Plants		0					132	132			0															0													132
Oil Refineries & other energy sector		0					0				2,902	2,902													57	0											17		2,976
Transformation Output		0	0	0	0	0	106	0	0	106	2,932	0	69	563	111	0	917	71	1,179	0	22	0	0	0	0	41	0	0	26	14	2	0	0	0	5	1,806	0	4,891	
Public Thermal Power Plants		0					0				0															38			24	14					5	1,599		1,599	
Combined Heat and Power Plants - Electricity		0					0				0															3			1		2						177		177
Combined Heat and Power Plants - Heat																										0												0	
Pumped Storage Generation																																					30		30
Briquetting Plants							106			106	0															0													106
Oil Refineries							0				2,932		69	563	111	0	917	71	1,179		22					0													2,932

Table 4.B Expanded Energy Balance Sheet 2013 (continued)

	Coal	Bituminous Coal	Manufactured	Coke	Lignite	Peat	Milled Peat	Sod Peat	Briquettes	Oil	Crude	Refinery Gas	Gasoline	Kerosene	Jet Kerosene	Fueloil	LPG	Gasoil / Diesel /DERV	Petroleum Coke	Naphta	Bitumen	White Spirit	Lubricants	Natural Gas	Renewables	Hydro	Wind	Biomass & Renewable Waste	Landfill Gas	Biogas	Liquid Biofuel	Solar	Geothermal Non-Renewable Waste	Electricity	Heat	TOTAL	
Exchanges and transfers	24	-14	38	0	0	0	0	0	0	-26	0	0	0	218	-218	2	0	-4	-24	0	0	0	0	0	-440	-50	-391	0	0	0	0	0	0	0	440	0	-2
Electricity																									-440	-50	-391								440		0
Heat																																				0	
Other	24	-14	38							-26			0	218	-218	2		-4	-24						0												-2
Own Use and Distribution Losses	0					16	16			64		64				0	0	0						65	0										262		407
Available Final Energy Consumption	377	276	84	0	17	174	-38	128	85	6,441	-2	0	1,218	771	674	146	139	3,119	100	-5	251	1	31	1,653	353	0	0	202	0	7	99	11	33	35	2,098	0	11,131
Non-Energy Consumption	0	0	0	0	0	0	0	0	0	283	0	0	0	0	0	0	0	0	0		251	1	31	0	0	0	0	0	0	0	0	0	0	0	0	0	283
Final non-Energy Consumption (Feedstocks)	0					0				283											251	1	31	0	0											283	
Total Final Energy Consumption	355	255	83	0	17	218	0	128	90	6,145	0	0	1,198	784	606	148	166	3,143	99	0	0	0	0	1,633	357	0	0	203	0	7	102	11	33	35	2,081	0	10,825
Industry*	82	82	0			1	0	0	0	546	0	0	0	78	0	138	116	125	88	0	0	0	0	620	153	0	0	150		3	0	0	0	35	799	0	2,234
Non-Energy Mining	0	0				0				34				2		2	0	30	0					10	0									57		102	
Food & beverages	21	21				0	0			137				46		29	33	28	0					90	28			25		3				170		447	
Textiles and textile products	0	0				0				3				1		1	0	1	0					1	0									10		14	
Wood and wood products	0	0				0				3				0		0	0	2	0					2	100			100						34		139	
Pulp, paper, publishing and printing	0	0				0				3				1		1	0	1	0					3	0									19		25	
Chemicals & man-made fibres	0	0				0				29				11		7	3	8	0					56	0			0						145		230	
Rubber and plastic products	0	0				0				10				0		0	6	3	0					4	0									35		49	
Other non-metallic mineral products	61	61				0				141				7		5	1	39	88					15	25			25					35	51		327	
Basic metals and fabricated metal products	0	0				0				96				0		88	6	3	0					321	0									64		480	
Machinery and equipment n.e.c.	0	0				0				6				0		0	2	3	0					5	0									20		31	
Electrical and optical equipment	0	0				0				42				0		0	40	2	0					106	0									99		247	
Transport equipment manufacture	0	0				0				5				0		0	4	1	0					2	0									17		23	
Other manufacturing	0	0				0		0		39				8		5	20	5	0					6	0									78		122	

Table 4.B Expanded Energy Balance Sheet 2013 (continued)

	Coal	Bituminous Coal	Anthracite + Manufactured Oxoids	Coke	Lignite	Peat	Milled Peat	Sod Peat	Briquettes	Oil	Crude	Refinery Gas	Gasoline	Kerosene	Jet Kerosene	Fueloil	LPG	Gasoil / Diesel /DERV	Petroleum Coke	Naphta	Bitumen	White Spirit	Lubricants	Natural Gas	Renewables	Hydro	Wind	Biomass & Renewable Waste	Landfill Gas	Biogas	Liquid Biofuel	Solar	Geothermal	Non-Renewable Waste	Electricity	Heat	TOTAL
Transport	0	0	0	0	0	0	0	0	0	4,173	0	0	1,198	0	606	0	1	2,368	0	0	0	0	0	0	102	0	0	0	0	0	102	0	0	0	4	0	4,279
Road Freight	0					0				562								562							18						18						581
Road Light Goods Vehicle										309								309							10						10						319
Road Private Car	0					0				1,791			918				1	872							50						50						1,842
Public Passenger Services	0					0				147			20					127							5						5						151
Rail	0					0				38								38							0						0				4		42
Domestic Aviation	0					0				3			1		2										0												3
International Aviation	0					0				603					603										0												603
Fuel Tourism	0					0				243			109					134							7						7						250
Navigation	0					0				57								57							0												57
Unspecified	0					0				418			150			0		268							12						12						430
Residential	273	173	83		17	218		128	90	918			0	706		0	40	161	11					606	64			28				11	25		684		2,763
Commercial/Public Services	0	0	0	0	0	0	0	0	0	307	0	0	0	0	0	10	9	289	0	0	0	0	0	407	38	0	0	25	0	4	0	0	9	0	547	0	1,300
Commercial Services	0	0	0		0	0				198				0		1	6	191	0					178	34			25				0	9		393		803
Public Services	0					0		0	0	109						9	2	98						229	4					4					155		497
Agricultural	0	0				0				177			0	0			0	177						0	0										48		225
Fisheries	0					0				24								24							0												24
Statistical Difference	22	21	1	0	0	-44	-39	0	-5	14	-2	0	19	-14	68	-1	-27	-24	0	-5	0	0	0	20	-5	0	0	-2	0	0	-3	0	0	0	17	0	24

Table 4.C Country specific carbon emission factors – fossil fuels

Fuel	NCV (toe/t)	NCV (MJ kg ⁻¹)	CO2 (t/TJ)	Density (kg l ⁻¹)
Liquid fuels				
Crude Oil	1.0226	42.814	73.333	
Refinery Gas	1.1912	49.871	52.652	
LPG	1.1263	47.156	63.700	
Naphtha	1.0510	44.003	73.333	
Motor Gasoline	1.0650	44.589	69.960	0.7550
Aviation Gasoline	1.0650	44.589	69.960	0.7550
Jet Kerosene	1.0533	44.100	71.401	0.8000
Other Kerosene	1.0556	44.196	71.400	0.8000
DERV (Road Gasoil)	1.0344	43.308	73.300	0.8450
Heating and Other Gasoil	1.0344	43.308	73.300	0.8450
Residual Fuel Oil	0.9849	41.236	76.000	0.9416
Residual Fuel Oil (Electricity Generation)	0.9630	40.320	79.134	
Fuel Oil – Low Sulphur content	0.9849	41.236	76.000	0.9416
Fuel Oil – High Sulphur content	0.9849	41.236	76.000	0.9416
White Spirits	1.0510	44.003	73.333	
Lubricants	1.0100	42.287	73.333	
Petroleum Coke	0.7697	32.224	92.934	
Bitumen (including Orimulsion)	0.9004	37.698	80.667	
Solid Fuels				
Coal (electricity generation)	0.5888	24.653	93.486	
Other Bituminous Coal (imports)	0.6649	25.199	94.600	
Other Bituminous Coal (default)	0.6650	25.199	94.600	
Anthracite	0.6650	27.842	98.260	
Lignite/Brown Coal	0.4733	19.816	101.200	
Patent Fuels (Manufactured Ovoids)	0.7643	32.000	98.260	
Milled Peat	0.1798	7.527	117.991	
Sod Peat	0.3130	13.105	104.000	
BKB/Peat Briquettes	0.4430	18.548	98.860	
	NCV (MJ m ⁻³)	NCV (MJ kg ⁻¹)	CO2 (t/TJ)	Density (kg m ⁻³)
Gaseous fuels				
Natural Gas-Indigenous	34.8108	47.111	56.198	0.7389
Natural Gas-Imported	35.7367	47.055	55.971	0.7595
Natural Gas- weighted average	35.2738		56.085	

Annex 5.1

Ireland's Response to the Recommendations in the UNFCCC Annual Review Reports

5.1a for Submission 2013

5.1b for Submission 2014

Table 5.1a Ireland's Response to the recommendations in the UNFCCC Annual Review Report for Submission 2013

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
Overview	Inventory planning	13	During the review, the ERT noted that the limited resources allocated to inventory work lead to the need for considerable prioritization in relation to inventory preparation, management and improvement. The ERT commends Ireland for the work carried out, but also recommends that Ireland maintain the quality of the inventory and facilitate further implementation of inventory improvements, if necessary by allocating sufficient resources.	Incomplete	
General	Key category analysis and uncertainty assessment	14	The ERT encourages Ireland to move to tier 2 in future submissions.	Incomplete	
General	QA/QC activities and documentation	15	The ERT recommends that Ireland ensure sufficient resources are available for the QA/QC activities and documentation.	Incomplete	
General	Inventory management	16	The ERT recommends that Ireland further improve the transparency of the NIR regarding AD, EFs and other parameters. Ireland uses emission estimates from EU ETS reports; however, the NIR does not provide information on the methods used to produce those data, or on the QA/QC carried out for those data used in the inventory. The ERT also noted that the description of the models used for the estimation of fluorinated gas (F-gas) consumption in the NIR is not fully transparent.		
General	Follow-up to previous reviews	17 & 72	The ERT found that some inconsistencies (i.e. mostly to do with typing and transcription errors) remain between the NIR (Submitted on 13 August 2013) and the CRF tables, as well as within the NIR and between the reporting under the Convention and under the Kyoto Protocol.		
General	Completeness	18	The ERT reiterates the encouragement made in the previous review report to investigate inclusion of non-mandatory categories reported as "NE" (not estimated), such as N2O use in anaesthesia.	N2O emissions from anaesthesia have been estimated for the first time in 2015 submission.	Chapter 4
Energy	Transparency	22, 23, 24	(22) The ERT noted that Ireland has improved the description of its QA/QC procedures related to the use of EU ETS data for the estimation of emissions from public electricity and heat production. However, the ERT further noted that in its NIR the Party does not provide information on category-specific QA/QC measures applied for other EU ETS data used in the inventory. The ERT considers that this lack of information on QA/QC measures applied in relation to the use of EU ETS data reduces the transparency of the Party's reporting. The ERT therefore recommends that Ireland provide more information on the category-specific QA/QC measures applied in relation to the use of EU ETS data in the NIR. (23) Ireland's inventory for the energy sector is, as referred to in several paragraphs in the NIR, based on emission data reported by plants. As the NIR does not provide transparent information on these data, the ERT met with difficulties in reviewing them. The ERT recommends that Ireland improve the transparency of the emission estimates in the energy sector, for example by providing a table in the energy chapter of the NIR for CO2 that includes the percentage distribution of emissions based on EU ETS data and calculated emissions, in order to improve the transparency of the	Additional information has been added to the NIR.	Chapter 3, Section 3.2.1

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			reporting and to facilitate future review activities. (24) Plant-specific data (on emissions and energy consumption) play an important role in the compilation of the inventory for fuel combustion for stationary combustion, particularly for the categories energy industries and manufacturing industries and construction, but also in the compilation of the energy balance. The estimates of CO ₂ emissions from energy industries for 2005–2011 are derived from EU ETS data. The emission estimates for 1990–2004 are also based on data from the plants, but are not as detailed as those based on EU ETS data. The ERT considers that this approach gives a consistent time series. However, it is not always clear from the NIR for which sectors and to what extent data from plants are included directly in the inventory. The ERT recommends that Ireland clarify this issue in its NIR.		
Energy	Transparency	26	The national energy balance produced by SEAI is a fundamental data source for the compilation of the Irish GHG inventory for the energy sector. The ERT notes that it is crucial that the energy balance is of high quality and that it is transparently described in the NIR. However, the ERT did not find the description of the energy balance to be sufficiently transparent. During the review week, SEAI gave a presentation on the production of the energy balance and the oil balance, which is compiled by the Department of Communications, Energy and Natural Resources. The ERT strongly recommends that Ireland include in the NIR a detailed and transparent description, including information on the data sources on which the balance is based, and whether the data sources are derived using a top-down and/or a bottom-up approach or based on surveys. The ERT also recommends that Ireland include in the NIR a transparent description of how the oil balance is produced, the basis for dividing diesel oil consumption between road traffic and non-road traffic and the allocation of other gas oils (marine diesel oil and light fuel oil) between the user categories.	Additional information has been added to the NIR.	Annex C
Energy	Completeness	27, 28	(27) The choice of EFs is not always well documented and justified in the NIR. For instance, the CO ₂ EF for petroleum coke for the earlier years of the time series (1990–2004) is based on average EU ETS data reported for the years 2005–2009, while for the years 2005–2011 Ireland uses an annual country-specific CO ₂ EF. This is also the case for the N ₂ O EF of 28.6 kg/TJ for agricultural mobile machinery, which is taken from the Intergovernmental Panel on Climate Change (IPCC) 2006 IPCC Guidelines for National Greenhouse Gas Inventories (hereinafter referred to as the 2006 IPCC Guidelines). The ERT notes that the corresponding EF in the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (hereinafter referred to as the Revised 1996 IPCC Guidelines) is 0.6 kg/TJ. The ERT recommends that Ireland document and justify the choice of EFs in the energy sector in the NIR. (28) The NIR does not provide information on CO ₂ EFs applied, net calorific values or densities for all energy carriers for which emissions are calculated. The ERT recommends that Ireland provide this information, if possible in a table, to increase the transparency of the inventory.	Additional information has been added to the NIR.	Annex C
Energy	Transparency	29	The ERT found that category-specific QA/QC procedures are not described adequately in the NIR for the use of plant-specific data. The ERT recommends that Ireland include this information in the NIR.		
Energy	Completeness, Key Categories	30, 38	(30) Since emissions from all non-road vehicles are not estimated explicitly in the Irish inventory, the ERT concludes that emissions from non-road vehicles can, in principle, be included in the Irish inventory, assuming that the energy balance is complete. The ERT has no information that indicates that this assumption is incorrect. However, the NIR only explicitly explains that the inventory includes emission estimates for tractors and other machinery used in agriculture and		

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			forestry. It is not clear from the NIR whether fuels used for other non-road vehicles (e.g. motorized equipment in the construction and building industries and lawn mowers in the residential sector) are included in the inventory. Emissions for these categories should be included in the total for stationary and mobile consumption under the appropriate categories. If the assumption of the ERT is correct, this would mean that, with the exception of machinery used in the agriculture sector, Ireland uses the same EFs for both stationary and mobile equipment. The ERT recommends that Ireland justify this assumption in the NIR. The ERT also recommends that Ireland justify why fuel consumption for non-road vehicles is included in the inventory and further recommends that Ireland improve the reporting on non-road vehicles by preparing a more accurate and transparent inventory for that category. (38) CO ₂ emissions from liquid fuels used for the category agriculture/forestry/fisheries include stationary emissions and mobile emissions from tractors and other agricultural machinery. The total consumption of diesel in the sector is divided into 90 per cent for mobile sources and 10 per cent for stationary consumption. The reference for the assumption is not given in the NIR. The ERT recommends that Ireland improve the description of the methodology and assumptions used in the NIR.		
Energy	Consistency	33	The ERT considers that the data for apparent energy consumption (excluding non-energy use and feedstock in CRF table 1.A(c)) should be consistent with the figures shown in CRF table 1.A(d). There are some discrepancies between the figures in the two tables, for example, for 1990, 2010 and 2011. The ERT recommends that Ireland correct these discrepancies, or explain why these discrepancies occur, in the NIR.		
Energy	Transparency	34	The NIR (page 72) states that the reference approach can be used to report national total emissions in cases where the detailed AD required for the sectoral approach are not available. The ERT considers the point of this description is unclear. The ERT encourages Ireland to clarify the statement in the NIR.		
Energy	Comparability	35	The ERT identified discrepancies between the energy consumption reported in the International Energy Agency (IEA) statistics and that reported in the CRF tables for domestic aviation, as the figures are generally 20–60 per cent lower in the CRF tables than the international statistics, while energy consumption for domestic navigation reported in the CRF tables is up to twice as high as reported in the IEA statistics for 2000 onwards, owing to a sharp drop in the IEA figures for gas and diesel oil consumption after 1999. The ERT encourages Ireland to provide a discussion on this issue in its NIR.	The fuel consumption data reflected in the IEA data for Ireland are reported by the Department of Communications, Energy and Natural Resources (DCENR), while the data reflected in the CRF tables are taken from the energy balances prepared by SEAI. The correct AD are those from the national energy balances and SEAI is currently in discussion with DCENR to provide all statistical information to IEA in the future on all types of	Annex C

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
				fuels consumed in the country, including any revisions to historical data, in an effort to improve the consistency of all energy data sets.	
Energy	Key categories, Stationary combustion: solid, liquid and gaseous fuels – CO2	37	According to information provided in the NIR, emissions from manufacturing industries and construction are estimated by multiplying energy consumption from the energy balance by country-specific CO2 EFs and default CH4 and N2O EFs for stationary combustion from the 2006 IPCC Guidelines. It is not clear from the NIR what reference is used in the inventory for the country-specific CO2 EFs, except for the CO2 EF for petroleum coke, which is based on EU ETS data. The ERT recommends that Ireland describe the references used for the country-specific CO2 EFs in the NIR.	It is all ETS data for CO2 in manufacturing industries sector (1.A.2).	
Energy	Key categories, Road transportation: liquid fuels – CO2, CH4 and N2O	39	Ireland uses the COPERT IV model, version 9.1, to estimate CH4 and N2O emissions from road transportation. The COPERT model is not described in the NIR. The ERT recommends that Ireland include in the NIR a general description of the COPERT model, for example, that the model is fuel driven (or kilometres driven) and that Ireland describe the relevant national data included in the model.	Additional information has been added to the NIR.	Chapter 3, Section 3.2.1.3.2
Energy	Key categories, Road transportation: liquid fuels – CO2, CH4 and N2O	40	The total consumption of gasoline and diesel oil used in the calculation of CO2 emissions from road transportation is from the energy balance, but there is no description in the NIR of how the amount of fuel is compiled in the balance. The ERT considers that the lack of this information makes the amount of fuel used in the calculation of CO2 emissions unclear to some extent. The ERT recommends that Ireland describe in the NIR how fuel consumption for road transportation is estimated.	Additional information has been added to the NIR.	Annex C
Energy	Non-key categories Oil and natural gas – CH4	41	Fugitive emissions of CH4 from offshore gas production are reported under the category fugitive emissions from oil and natural gas, and other sources and emissions from fuel combustion for energy purposes, for example pipeline transportation, are reported under the category other transportation. The ERT notes that it is not clear where emissions from fuel combustion for energy purposes at offshore platforms are reported in the CRF tables, other than for pipeline transportation at the platforms. The ERT recommends that Ireland explain in the NIR the different emission sources in the offshore sector (gas production), the compiling of the energy balance and where in the CRF tables the different emission sources are reported.		
IP and Solvent and other product use	Transparency	45	Information in the NIR lacks transparency regarding AD, EFs and other parameters. In particular, this relates to emission estimates taken directly from EU ETS reports, which are taken as good-quality data, but which are not followed by justification of their use and further explanations in the NIR regarding the methodology used. In addition, the inventory is not fully transparent, owing to a lack of description of the models used for the estimation of F-gas consumption in the NIR. The ERT recommends that Ireland provide more information to ensure transparency with regard to the EU ETS data used for mineral products.	Additional information has been added to the NIR.	Chapter 4
IP and	Consistency	46	Regarding QA/QC, the ERT identified minor issues related to the use of notation keys, for	Emissions from this	

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
Solvent and other product use			instance indirect GHG emissions from glass production are reported as "NE", while GHG emissions for that subcategory are reported as not occurring ("NO") for the years after production ceases. The ERT encourages Ireland to use consistent notation keys for the same subcategory.	subcategory are "NO" (not occurring) for the years after production ceases and will be corrected in the next submission.	
IP and Solvent and other product use	Completeness	47	Ireland reported "NE" for N2O emissions from anaesthesia and aerosol cans in the solvent and other product use sector, for which there is no methodology available in the Revised 1996 IPCC Guidelines or the IPCC good practice guidance; however, many Parties report these emissions. The ERT encourages Ireland to investigate these categories and, if necessary, report the corresponding emissions in the NIR.	N2O emissions from anaesthesia have been estimated for the first time in 2015 submission	Chapter 4
IP and Solvent and other product use	Key categories Cement production – CO2	48	Ireland uses plant-specific AD and EFs from the four plants operating in the country. Since 2004, emission data from EU ETS reports have been used; however, the methodology is not fully explained in the NIR, except that it takes into account the cement kiln dust (CKD) and calcium oxide (CaO) content of the clinker. Following recommendations made in previous review reports, information on the confidentiality of the CaO and magnesium oxide (MgO) contents of the clinker has been added to the NIR. During the review, Ireland provided AD to the ERT for 2008–2012 from each of the four cement plant operators. Using stoichiometric calculations for CaO and MgO and this information, the ERT found minor discrepancies in the CO2 emission estimates reported in the CRF tables. To enhance transparency, the ERT recommends that Ireland better explain in the NIR the methodology used to produce the EU ETS data and correct the minor discrepancies.	Information on calcium oxide (CaO) and magnesium oxide (MgO) content of clinker has been provided to the inventory agency by the plant operators, however, this information is not published in this inventory report as the cement producers deem it to be confidential. The data are available to the expert review teams for annual GHG inventory reviews upon request.	Chapter 4, Section 4.2.1
IP and Solvent and other product use	Key categories, 2.F Consumption of Halocarbons and SF6 – mobile air conditioning	49	Ireland reports both actual and potential emissions for two subcategories – mobile air conditioning (MAC) and domestic, commercial, transport and industrial refrigeration, and stationary air conditioning. The emission estimates are calculated using models that were provided to the ERT during the review week. The ERT was also provided with two background reports (Adams et al. (2005) and O'Doherty and McCulloch (2002)), on which the models are based. The ERT considers that the models are not transparently described in the NIR and recommends that Ireland enhance its description of the models employed in its NIR.	Following an independent study, estimates from most categories in sector 2.F have been revised and additional information about those recalculations has been added to the NIR.	Chapter 4, Section 4.4
IP and Solvent and other product use	Key categories, 2.F Consumption of Halocarbons and SF6 – mobile air conditioning	50	For MAC, the ERT found that the tier 2a bottom-up methodology was applied instead of the tier 3b bottom-up method, as stated in the NIR. The NIR also states that the new fleet entering each year can be derived from table E-5 of the NIR, which is not the case. The ERT recommends that Ireland provide information on the number of new vehicles that is used for the estimation, as well as confirm that the correct tier is used, in its NIR.	Following an independent study, estimates from most categories in sector 2.F have been revised and additional information about those recalculations has been added to the NIR.	Chapter 4, Section 4.4
IP and Solvent and	Key categories, 2.F	51	During the review week, the ERT found that the model used for estimating emissions from MAC only included the scrapping of vehicles at the end of their lifetime, which is not in line with the	Following an independent study, estimates from most	Chapter 4, Section 4.4

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
other product use	Consumption of Halocarbons and SF6 – mobile air conditioning		IPCC good practice guidance method, and that there was no consideration of the remaining charge at end of lifetime and no indication of whether the new regulations (starting from 2009) on recovering gas at decommissioning are being followed in Ireland. The ERT also found an error in the input data of the 2011 EF for MAC, where the amount of fluid remaining in products at decommission was reported as “NO” although emissions from disposal were estimated. The ERT recommends that Ireland correct the errors in the method by using the methodology provided in the IPCC good practice guidance (section 3.7.5), including default values for those parameters that are not country specific.	categories in sector 2.F have been revised and additional information about those recalculations has been added to the NIR.	
IP and Solvent and other product use	Key categories, 2.F Consumption of Halocarbons and SF6 – mobile air conditioning	52	Regarding HFC consumption in domestic, commercial, transport and industrial refrigeration and stationary air conditioning, Ireland estimates actual emissions using the tier 2 approach, using a proportion of the sales as an indication of the annual leakage. During the review week, with access to the model in Excel spread sheets, the ERT noticed that one of the gases used in transport refrigeration – refrigerant R-134a – was not included in the calculation of potential emissions of HFC-134a, leading to an underestimation of the potential emissions. The ERT recommends that Ireland estimate the missing potential emissions and include them in its annual submission. The ERT also recommends that Ireland improve the QA/QC procedures, documentation and data collection for the category, and encourages Ireland to investigate the availability of market surveys and industrial association information to ensure that no data are omitted.	Following an independent study, estimates from most categories in sector 2.F have been revised and additional information about those recalculations has been added to the NIR.	Chapter 4, Section 4.4
IP and Solvent and other product use	Key categories, 2.F Consumption of Halocarbons and SF6 – fire protection	53	The ERT noted that Ireland is possibly the only country in Europe estimating HFC-227ea and HFC-23 emissions for the fire protection subcategory for 1990 to 1995, on the basis of a time series since 1984, when HFCs were rarely used. The ERT considers that Ireland might have used other gases instead of these HFC species. The ERT recommends that Ireland revise the estimation for the whole time series in order to check the proper identification of the gases used in this category and provide further explanations in the NIR.	Following an independent study, estimates have been revised and the years 1990 to 1995 are now reported as Not Occurring.	Chapter 4, Section 4.4.2.3
IP and Solvent and other product use	Non-key categories Lime production – CO2	54	The NIR indicates that CO2 emissions from lime production are estimated by lime producers for the development of the first national action plan (NAP1) under European Union directive 2003/87/EC regarding the EU ETS, and that the emissions were calculated in accordance with the methods described in the supporting European Union decision 2004/156/EC. Nevertheless, the NIR does not indicate how this methodology adheres to the IPCC good practice guidance. During the review, Ireland explained that the fluctuation in the annual implied emission factor (IEF) depends on the amount of lime produced at each plant. The ERT considers this explanation to be insufficient and recommends that Ireland provide more detailed information about this methodology in the NIR.		
IP and Solvent and other product use	Non-key categories Limestone and dolomite use – CO2	55	The NIR indicates that CO2 emissions from limestone and dolomite use are estimated on the basis of the quantity of limestone used by the companies and an EF of 0.44 t CO2/t limestone, which is the stoichiometric ratio of CO2 to calcium carbonate. However, the CRF tables include such a value for up to 2000 only and after that the IEF fluctuates. The ERT recommends that Ireland provide more detailed information about this fluctuation in the NIR.	The EF values will be updated in the next NIR.	Chapter 4

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
Agriculture	Transparency	57	The ERT noted that the methods used in the calculations are not fully described in the NIR and that the publications referred to in the NIR are not publicly available. The ERT therefore recommends that the Party improve the description of the methods used for the estimation of emissions, especially in relation to the reports by O'Mara (2006) and Hyde et al. (2008), because both publications contain detailed background information on country specific parameters for livestock, which have an influence on all categories reported by the Party and are frequently referred to in the NIR.	The hyperlink in the references section of the NIR will be updated. Redesign of the EPA website has led to issues with hyperlinks previously referenced. The Hyde et al report was never officially published and is available to review teams on request.	Chapter 5
Agriculture	Key categories Enteric fermentation – CH ₄	61	The EFs for non-dairy cattle were generally higher in the 1990s than in recent years. Ireland provided in the NIR justifications only for male beef cattle more than two years old, but not for the other age classes. The Party explained that this was because of a higher proportion of late maturing breeds, a lower quality of forage and a lower usage of concentrates during the 1990s. This information was also found in O'Mara (2006). Therefore, the ERT recommends that the Party include this information in the NIR or make the publication readily available to readers in order to increase transparency.	Information will be provided in the NIR.	Chapter 6 section 6.2.5
Agriculture	Key categories Enteric fermentation – CH ₄	62	Ireland has been using the tier 1 method for estimating CH ₄ emissions from sheep, although it has been recommended to use tier 2 for estimating these emissions in several previous review reports, taking into account that enteric fermentation has been a key category. The ERT considered that the use of the tier 1 method is still in line with the IPCC good practice guidance, taking into account the significance of emissions from sheep in this subcategory (accounting for 10.8 per cent and 6.8 per cent of the total emissions from enteric fermentation in 1990 and 2011, respectively) and the decision tree in figure 4.2 of the IPCC good practice guidance. Nevertheless, the ERT reiterates the recommendation made in the previous review report that the Party move up to a tier 2 method when reliable data and information become available, because enteric fermentation is a key category, in both the level and trend analyses, and because the Party explained that an increase in CH ₄ emissions from sheep is expected in the near future owing to the potential rise in market demand for lamb. The ERT also recommends that the Party correct the description of how the default EF was adjusted for each subcategory of sheep, if the Party continues to use the tier 1 methodology for estimating these emissions.	Information will be provided in the NIR.	Chapter 6 section 6.2.6 and 6.9
Agriculture	Key categories Manure management – N ₂ O	63	Fixed N excretion rates have been used for all animals except dairy cattle for the whole time series. In the category manure management, dairy cattle and non-dairy cattle were found to be significant subcategories, accounting for 25.5 per cent and 61.7 per cent, respectively, of the total N excreted by animals in 2011, while the excretion rates for other animals were not significant. The Party explained that the inventory agency was in on-going discussions with relevant stakeholders to develop dynamic N excretion rates for cattle. The ERT recommends that the Party continue the investigation and apply the results, in particular for dairy and non-dairy cattle taking into account their significance, when the data become available.	The inventory agency will continue to engage with the agricultural research community in developing dynamic N excretion rates for cattle.	
Agriculture	Key categories Agricultural soils – N ₂ O	64	Ireland has reported direct N ₂ O emissions and indirect N ₂ O emissions (leaching and runoff) from sewage sludge application. However, the Party did not report indirect N ₂ O emissions from atmospheric deposition for the whole time series, even though the estimation methodologies for indirect emissions from sewage sludge are provided in the IPCC good practice guidance. The	Emissions from this source category will be estimated and included in future submissions.	Chapter 6 section 6.5.3 and 6.8

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			ERT considered that this could lead to a potential underestimation of emissions; therefore, the ERT included this issue in the list of potential problems and further questions raised by the ERT during the review week. In response to the list of potential problems and further questions, the Party submitted revised estimates for the entire time series and provided a description of the method used to calculate the emissions, including all of the parameters used for the year 2011. Ireland calculated the indirect N ₂ O emissions from atmospheric deposition based on the AD (the amount of N in sewage sludge) reported for the direct N ₂ O emissions from sewage sludge in its original submission, and using the IPCC default fraction of livestock N excretion that volatilizes as ammonia (NH ₃) and nitrous oxides (NO _x) (FracGASM) and the IPCC default EF for N ₂ O emissions from atmospheric deposition of N. The ERT confirmed that the calculation was conducted in accordance with the IPCC good practice guidance, and the ERT therefore considers that the potential underestimation has been resolved. As a result, the indirect N ₂ O emissions from atmospheric deposition have increased by 0.10 Gg CO ₂ eq or 0.02 per cent for 1990 and 2.05 Gg CO ₂ eq or 0.48 per cent for 2011 compared with the original submission. The Party explained that Ireland was in the process of investigating the applicability of estimating ammonia emissions from the spreading of sewage sludge on agricultural land. The ERT welcomes the effort and recommends that the Party replace the default FracGASM data with country-specific data when they become available.		
LULUCF	Completeness	67	The key driver for the rise in net removals over the reported period is a steadily increasing growth in living biomass on land converted to forest land owing to a continuous programme of afforestation over more than 20 years. The contribution to the trend from categories other than forest land is small in absolute numbers but substantial in relation to the emissions in the category. Emissions from cropland increased by 1,788.3 per cent over the period owing to an increase in cropland areas that involves the conversion of improved grassland. Emissions from grassland decreased by 55.4 per cent over the period owing to a rapid decline in extensive grazing of sheep in the early 1990s, on-going afforestation activities and intensification of livestock practices. Emissions from wetlands decreased by 29.5 per cent owing to a decrease in demand for peat. Emissions from settlements decreased by 7.1 per cent owing to an abrupt halt of infrastructure planning. Removals from other land increased by 12,980.0 per cent because grasslands that are no longer reported under agriculture are reverting to natural grasslands and are reported under other land. Ireland provided additional information on the drivers for the trends during the review and the ERT recommends that Ireland include this information in its NIRs of future submissions.	Discussion of the key drivers of change in these categories is provided in the text in relevant sections of the NIR for Cropland and Grasslands. Discussion of drivers to changes in wetlands is restricted to Peat extraction activities. Further analysis is required to assess the underlying drivers of changes in settlements, due to the very complex economic and social interactions which influence behaviour in this sector.	6.4.1, 6.4.5, 6.4.9, 6.5.1, 6.5.4, 6.5.7, 6.6.3, 6.6.6, 6.6.7, 6.9
LULUCF	Consistency	68	As indicated in previous review reports, the Party used different versions of the CARBWARE model to estimate emissions and removals from LULUCF sinks and sources under the Convention and under the Kyoto Protocol. In response to recommendations made in previous review reports, Ireland developed an approach to overcome consistency problems. For reporting for 2007 onwards, Ireland uses the same version of the CARBWARE model for reporting under the Convention and under the Kyoto Protocol. The reporting for 1990 to 2006 is based on the FORCARB model. To correct for biases introduced by using different models, the historical time series calculated with the FORCARB model was rescaled using data for 2007 to 2011 for the two models. The ERT commends Ireland for its efforts to make the reporting transparent and consistent, but also notes that the correction made to the historical time series for living biomass	Completed	6.3.1

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			and dead organic matter was only based on the relationship between the two methods (CARBWARE and FORCARE) for five years. The ERT recommends that Ireland continue to assess the requirement for the correction of the historical time series when new modelled data and data from the second NFI and other sources become available.		
LULUCF	Comparability	69	The ERT noted the efforts made by the Party to verify the estimates by comparing IEFs for net removals from forest land remaining forest land and land converted to forest land with net removals reported by other Parties and by independent techniques (eddyflux). However, comparing IEFs for LULUCF may not always be relevant since Parties use different approaches for area accumulation, and factors other than the ones of interest may affect the comparison. For instance, net removals are affected by very country-specific parameters (harvest levels). The ERT noted that the IEF for gains in living biomass for land converted to forest land was higher, and that the IEF for losses in living biomass for forest land converted to other land-use categories was lower, compared with other Parties. The ERT recommends that Ireland continuously verify estimates for gains and losses separately for land converted to forest land and forest land converted to other land-use categories, and include information in the NIR justifying the deviation in IEFs from other countries.	Completed	6.10.3
LULUCF		70	In addition to the major revision by implementing a new approach to calculate carbon stock changes in forest land, Ireland also reclassified the 20-year transition period used for land converted to forest land and forest land converted to other land-use categories to comply with the approach used for afforestation, reforestation and deforestation under the Kyoto Protocol. Forests are now classified as pre- and post-1990 forests, while other land-use change categories continue to apply the 20-year transition period. This resulted in a significant change in land-use change trends. The ERT acknowledges this initiative, but also notes that the use of different transition periods and the effects on the accumulated areas in different land-use change categories require further clarification in the NIR. The ERT recommends that Ireland add more information which clearly describes the use of different transition times for different land-use change categories, including the relationship to the activities reported under the Kyoto Protocol, in the NIR. To further enhance the transparency related to the methods to estimate land-use changes, the ERT encourages Ireland to improve the description on how annual land-use changes (which is now included in the category sections of the NIR) are estimated by merging it into the section explaining how different data sources are used to estimate the land use.	Detailed analysis of agricultural land parcel has indicated that there is very limited occurrence of permanent conversion of land between grassland and cropland. It is challenging therefore to assign a transition period for these two land use categories.	
LULUCF	LULUCF for NIR	71	The information in the NIR for LULUCF has also undergone major revisions, mainly in order to align with the changes in methods for forest-related categories, but also because of the results of a peer review conducted in 2012. The ERT commends Ireland for these efforts. The NIR is generally well written and includes most of the necessary information for a proper review. However, the NIR could be further improved by including: (a) a table defining the carbon pools reported for each category; (b) a clear description of how notation keys have been used throughout the reporting of the LULUCF sector; (c) more information on the temporal and spatial coverage of the different data sources used for the sector; and (d) a better description of the land-use change matrix.	Completed	6.3.2.6 and 6.3.2

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			In addition, the numbering and references to equations and tables needs to be checked. The ERT encourages Ireland to improve the NIR by addressing the issues listed above.		
LULUCF	Accuracy, Transparency	72, 73	<p>(72) During the review, the ERT identified inconsistencies between the NIR and the CRF tables, as well as within the NIR and between the reporting under the Convention and under the Kyoto Protocol. As explained to the Party during the review, the identified inconsistencies were mostly to do with typing and transcription errors and included:</p> <p>(a) For 2011, net CO₂ removals are 3,821.95 Gg for land converted to forest land and 3,759.62 Gg for afforestation and reforestation (the reported areas are the same);</p> <p>(b) For 2011, net CO₂ emissions are 29.76 Gg for deforestation and net CO₂ removals are 29.81 Gg for forest land converted to other land reported information items in CRF table 5 (the reported areas are the same);</p> <p>(c) According to the NIR, the EF used for limestone is 0.12 Mg CO₂-C/Mg, but in table 5(IV), the IEF for cropland is 0.11 Mg CO₂-C/Mg;</p> <p>(d) The CO₂ emissions from biomass burning for afforestation and reforestation in 2011 (83.42 Gg) are four times higher than the emissions from biomass burning for land converted to forest land (21.09 Gg);</p> <p>(e) The area of managed wetland (peatland) in CRF table 5D (52.48 kha) is smaller than the area reported under peatland in CRF table 5(II) (52.95 kha) for 2011.</p> <p>(73) The ERT recommends that Ireland strengthen the QA/QC procedure to avoid transcription and typing errors in the NIR and the CRF tables. The ERT also recommends that Ireland explain real differences between related categories in the LULUCF sector under the Convention and activities under the Kyoto Protocol.</p>	Completed	6.10.0
LULUCF	Key categories Land converted to forest land – CO ₂	74	As also noted in the previous review report, Ireland does not report carbon stock changes in the soil organic carbon (SOC) pool (mineral soils) for land converted to forest land. Ireland states and verifies in chapter 11 of the NIR that the pool is not a source according to the requirements of the annex to decision 16/CMP.1, although this provision is only applicable to activities under the Kyoto Protocol. Since default methods to estimate carbon stock changes in mineral soils for this category are available in the Good Practice Guidance for Land Use, Land-Use Change and Forestry (hereinafter referred to as the IPCC good practice guidance for LULUCF), and considering that land converted to forest land is a key category for Ireland, the ERT recommends that Ireland report carbon stock changes in the SOC pool (mineral soils for land converted to forest land).	Completed	6.3.6.1.1
LULUCF	Non-key categories Forest land converted to other land-use categories – CO ₂	75	Ireland does not report carbon stock changes in the SOC pool (mineral soils for grassland) for forest land converted to other land-use categories (to grassland, settlements and other land), mentioning that the mineral SOC pool is verified not to be a source (grassland) and that there is no documentation of methods in the IPCC good practice guidance for LULUCF (settlements, other land). As mentioned in paragraph 74 above, the “not a source” provision is only applicable for activities under the Kyoto Protocol. The ERT reiterates the recommendation made in the previous review report that Ireland report carbon stock changes in mineral soils for forest land converted to other land-use categories for categories where methods are provided in the IPCC good practice guidance for LULUCF.	Completed	6.3.6.1.1
LULUCF	Non-key categories	76	As indicated in previous review reports, the ERT noted that the areas of natural grassland that are not grazed in the inventory year were reported under the land-use category other land.	Completed	6.4

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
	Other land – CO ₂		During the review, Ireland informed the ERT that existing data were explored to establish a robust methodology for determination of the extent of natural grassland for inclusion within the land-use category grassland. The ERT reiterates the recommendation made in the previous review report that Ireland introduce natural grassland areas as a subdivision of the land-use category grassland.		
LULUCF	Non-key categories Biomass burning – CH ₄ and N ₂ O	77	The ERT found that Ireland uses different units for the reporting of AD for biomass burning under the Convention (area burned) and under the Kyoto Protocol (mass burned). Using different units makes the assessment of consistency between the Convention and the Kyoto Protocol reporting of LULUCF difficult. Therefore, the ERT recommends that Ireland use and report the same units for AD for biomass burning under the Convention and under the Kyoto Protocol (see also para. 105 below).	Completed	6.3.6
Waste	Completeness	81	The ERT encourages Ireland to maintain an on-going watch on commercial composting and consider introducing the calculation of corresponding emission estimates when appropriate.	CH ₄ and N ₂ O emissions from sector 5.B.1 Composting have been estimated for the first time in 2015 submission.	Chapter 7, Section 7.3, NIR 2015
Waste	Completeness	82	Ireland has provided an aggregate uncertainty for emissions from solid waste disposal on land but not for wastewater handling or incineration in the waste section of the NIR. While the overall inventory uncertainty does include incineration and wastewater uncertainty, the ERT encourages Ireland to expand the discussion of uncertainty in the waste chapter to include aggregate wastewater and incineration uncertainty in its next annual submission.	Revised and more detailed uncertainty analysis has been developed and it includes estimates for wastewater and waste incineration and open burning.	Annex 2, NIR 2015
Waste	Completeness	84	QA/QC procedures are generally well documented in the NIR. However, discussions during the review week confirmed that some recent sector-specific QA activities have not been documented. The ERT encourages Ireland to fully document these activities in its NIR of the next annual submission.	Further information will be provided in the NIR.	Chapter 8 section 8.6
Waste	Key categories Solid waste disposal on land – CH ₃	85	Ireland uses the tier 2 method of the IPCC good practice guidance, first-order decay (FOD) model, to estimate emissions from managed waste disposal on land, which is consistent with the 2006 IPCC Guidelines. Ireland provided justification for the use of the 2006 FOD model in the NIR. AD to support this model are obtained from the national waste database published by EPA on a yearly basis since 2001 and periodically prior to 2001. The model is run separately for larger landfills and aggregations of smaller landfills, as well as for sewage sludge and street sweepings. This provides a good degree of flexibility to reflect operational circumstances at individual landfills and groupings of similar landfills. The ERT encourages Ireland to continue to review structural changes in the landfill sector.	The inventory agency will continue to review structural changes in the landfill sector and discuss such changes in the NIR.	Chapter 8 section 8.7
Waste	Key categories Solid waste disposal on land – CH ₄	87	Ireland provides information on waste composition in annex I of the NIR. The ERT noted the recommendation made in previous review reports that Ireland provide further information on the composition of “organic waste” in the NIR to improve the transparency of the inventory. This information has not been provided in this submission. Therefore, the ERT reiterates this recommendation and also recommends that Ireland report on the composition of street sweepings and other municipal solid waste (MSW) to support the degradable organic carbon (DOC) values reported in the NIR.	The inventory agency will provide further information in the NIR.	Chapter 8 section 8.2 and 8.7 and Annex I

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
Waste	Key categories Solid waste disposal on land – CH ₄	88	The ERT noted that some model parameters used in the FOD model are not discussed in the NIR. For example, the time lag between disposal and methanogenesis, oxidation and fraction of CH ₄ in landfill gas are not explicitly discussed. The ERT recommends that Ireland include a discussion of these model parameters in its NIR, including the values used and the justification for their use.	The inventory agency will provide further information in the NIR.	
Waste	Non-key categories Wastewater handling – CH ₄ and N ₂ O	90	The ERT finds that the information in the NIR on industrial wastewater is limited and should be expanded to better describe the source and derivation of AD and the industrial sectors contributing to the BOD load. The ERT encourages Ireland to include AD from wastewater in annex I of the NIR.		
Waste	Non-key categories Wastewater handling – CH ₄ and N ₂ O	92	Ireland reports that all wastewater treatment plants employ aerobic processes, but that a small proportion of sludge is treated anaerobically. The fraction of BOD in sludge that is treated anaerobically is applied consistently throughout the whole time series and is 3 per cent based upon data in O'Leary and Carty (1998). Publications provided to the ERT which pre- and post-date this publication, show some variation in this value (e.g. O'Leary et al. (1997) suggest a value of 8 per cent and O'Leary et al. (2000) suggest a value of 6 per cent). The ERT recommends that Ireland review this assumption in the light of structural changes in the wastewater treatment network that have occurred since the publication in 1998 and report on this in the NIR.		
Waste	Non-key categories Wastewater handling – CH ₄ and N ₂ O	93	Ireland reports in the NIR that one wastewater treatment plant in Ringsend processes sludge for reuse and captures the resulting CH ₄ emissions for use on site. However, this site is not accounted for in the AD data from 2005 onwards when CH ₄ collection commenced. Rather, a quantity of BOD is deducted from the total BOD to effectively remove this plant from the system. The ERT concludes that this is not transparent and recommends that Ireland include BOD from Ringsend in the total BOD and provide a deduction as CH ₄ captured in CRF table 6.B. Furthermore, the ERT recommends that Ireland review and provide additional justification for the estimate of population equivalent covered by this plant to enhance transparency.	The inventory agency will review the status of anaerobic digestion of sludge in wastewater treatment plants in Ireland and include further detailed information in the NIR.	Chapter 8 section 8.3.1 and 8.7
Waste	Non-key categories Wastewater handling – CH ₄ and N ₂ O	94	During the review, Ireland confirmed that the quantity of CH ₄ consumed at Ringsend for production of on-site electricity is captured in the energy balance and reported appropriately in the energy sector. The ERT encourages Ireland to document this in the next NIR and to cross-check this value with the CH ₄ capture reported in table 6.B.	The inventory agency will review the status of anaerobic digestion of sludge in wastewater treatment plants in Ireland and include further detailed information in the NIR.	Chapter 8 section 8.3.1 and 8.7
Waste	Non-key categories Waste	95	Ireland has included emissions from solvent and clinical waste incineration in its 2011 submission. These emissions are estimated according to the tier 1 method in the 2006 IPCC Guidelines. AD are sourced from the national waste database. While there are no appropriate	Emission factor information with references has been included in 2014	Tables 8.4 and 8.5, NIR 2014

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
	incineration – CO ₂ , CH ₄ and N ₂ O		EFs for the incineration of solvents and clinical waste in the Revised 1996 IPCC Guidelines or the IPCC good practice guidance, the ERT encourages Ireland to include a discussion in its NIR on the applicability of EFs from the 2006 IPCC Guidelines to its incinerator units.	submission.	
Waste	Non-key categories Waste incineration – CO ₂ , CH ₄ and N ₂ O	96	The ERT found a small instance of double counting in clinical waste incinerated in CRF table 6.C. This double counting has occurred because the total value of clinical waste incinerated is reported against both biogenic and non-biogenic waste. The ERT recommends that Ireland correct this double counting in the CRF tables by disaggregating the AD into biogenic and non-biogenic components.		
KP-LULUCF	Completeness	99	Ireland has identified CO ₂ emissions from afforestation and reforestation under Article 3, paragraph 3, of the Kyoto Protocol as a key category for 2011. The ERT recommends that Ireland include a paragraph explaining the assessment of key categories for the KP-LULUCF activities in its NIR (chapter 11 on supplementary information under Article 7, paragraph 1, of the Kyoto Protocol).		
KP-LULUCF	Completeness	100	The ERT notes that the Party's IEF for gains in living biomass for afforestation and reforestation was very high and that the IEF for losses in living biomass for forest land converted to other land-use categories was low compared with those of other Parties. During the review, Ireland provided information justifying these differences, including evidence for the high yields in Sitka spruce plantations. The ERT assessed and accepted the justification. The ERT recommends that Ireland continuously verify estimates for gains and losses separately for afforestation, reforestation and deforestation and include information on the justification of the deviation in IEFs from those of other countries in its NIR.	Completed	6.10.6
KP-LULUCF	Consistency	101	During the review, the ERT identified inconsistencies between the NIR and the CRF tables, as well as within the NIR and between the reporting under the Convention and under the Kyoto Protocol in the CRF tables. The identified inconsistencies were mostly because of typing and transcription errors. In addition to errors listed in paragraph 72 above, Ireland reports different areas for deforestation in CRF table for 2011 (8.50 kha) and in the NIR (p. 235) (8.49 kha). The ERT recommends that Ireland strengthen the QA/QC process to avoid transcription errors and typing errors in the NIR and the CRF tables, and explain the real differences between related categories and activities, if necessary, in future submissions.	Completed	6.10.1
KP-LULUCF	transparency, completeness	102	As previous annual review reports pointed out, the ERT noted that the areas for afforestation and reforestation at the end of a reported year do not match the areas for afforestation and reforestation at the beginning of the next reporting year, as shown in table CRF table NIR-2. During the review, the problem relates to how the table is interpreted, and exclusively concerns Parties that include lands under deforestation which were previously reported under afforestation	Completed	NIR2, Tab3 11.2 NIR Ch 11

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
			and reforestation. The problem was solved through discussion and consideration between the Party and the ERT. The ERT recommends that Ireland correct the reported areas in CRF table NIR-2 and, in the upper left cell of CRF table NIR-2, include the part of the existing afforestation and reforestation area at the beginning of the reported year that is not deforested during the reported year, and in the upper second left cell, include the deforested area during the reported year previously reported as afforestation and reforestation. Deforested areas during the reported year that have not previously been reported under afforestation and reforestation shall be reported as "other" in the column deforestation. Land areas afforested during the reported year should be included under "other" in the column afforestation and reforestation.		
KP-LULUCF	Afforestation and reforestation – CO ₂ , CH ₄ , N ₂ O	104	Ireland provides information in the NIR verifying that the SOC pool (mineral soils) is not a net source. The ERT found the information to be mostly relevant for justifying the exclusion of the SOC pool from reporting. However, some of the information provided was not relevant to justifying the exclusion of the SOC pool from reporting. The ERT strongly recommends that Ireland strengthen the arguments which justify that the mineral SOC pool is not a source under afforestation and reforestation by only including very specific information in the NIR.	Completed	11.3
KP-LULUCF	Afforestation and reforestation – CO ₂ , CH ₄ , N ₂ O	105	The ERT found that Ireland uses different units for the reporting of AD for biomass burning under the Convention (area burned) and under the Kyoto Protocol (mass burned). Using different units makes the assessment of consistency difficult and the ERT therefore recommends that Ireland use the same units for reporting AD for biomass burning under the Convention as used under the Kyoto Protocol (see also para. 77 above).	Completed	6.3.6
KP-LULUCF	Deforestation – CO ₂	106	Deforestation areas are identified using the IPCC approach 2 based on legally binding licence applications for harvest under the Forestry Act. These provisions fulfil the requirement to demonstrate that deforestation is directly human induced. Deforestation is reported using the boundaries of the entire territory of Ireland. The Party explained that, since AD for deforestation is considered very uncertain, the deforested area will be verified using the second NFI, which ended in 2012. The ERT acknowledges this initiative and recommends that Ireland provide information on this verification.	Completed	11.1.7
KP-LULUCF	Deforestation – CO ₂	107	Ireland does not report SOC changes in mineral soils for deforestation. The ERT considers that, if no evidence can be provided to justify that the pool is not a source following conversion, SOC changes need to be reported. Ireland includes some information justifying that conversions to grassland may not result in emissions, but there is no justification for omitting conversions to settlement and other land. The ERT strongly recommends that Ireland report carbon stock changes in the SOC pool (mineral soils) for deforestation if it cannot be justified that the pool is not a source.	Completed	6.3.6.1
Changes to the national registry		114	The SIAR, part II, states that Ireland is not fully reporting changes in the national registry related to the description of database structure. Although the Party resubmitted a simplified data model during the assessment cycle, the information contained within the model is not sufficient. This is evidenced by the lack of descriptions of each entity in the diagram and the omission of some diagram entities mandated in the DES. The recommendation contained in the SIAR, part II, is that, following major changes, the Party provide a data model that contains all entities required by the DES complete with descriptions in its NIR. The ERT recommends that Ireland include this information.		

Table 5.1b Ireland's Response to the recommendations in the UNFCCC Annual Review Report for Submission 2014

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
Cross-cutting	General	4	Include the information on the key drivers of emission/removal trends on cropland, grassland, wetlands, settlements and other land in the next NIR	More discussion on trends included in NIR 2015.	Chapter 2, Chapter 6, Chapter 11, Annex 3.4
Cross-cutting	General	77	Include a paragraph explaining the assessment of key categories for the KP-LULUCF activities in chapter 11 in its NIR in order to enhance the transparency of its NIR		
Energy	Reference approach	21, 22	Further investigate the difference between reference approach and sectoral approach, and report accordingly in the next NIR	Resolved, no difference between two approaches in 2015 submission.	Chapter 3, Annex 3.1.A
Energy	Feedstocks and non-energy use of fuels	25	Investigate the emissions related to the non-energy use of lubricants, other than road transportation, and report accordingly in the next submission	Lubricants (and Paraffin wax) emissions are reported for the first time in 2015 submission.	Chapter 4
Energy	Stationary combustion: liquid and gaseous fuels – CO ₂	26	Improve the transparency of the reporting of emission estimates of this category by providing more information in relation to the use of EU ETS data in the NIR	Additional information provided.	Chapter 3, Annex 3.1.A, MMR IR Article 10
Energy	Stationary combustion: liquid and gaseous fuels – CO ₂	27	Provide information on AD and CO ₂ EFs for the different types of fuel and industrial activities reported under other (manufacturing industries and construction)		Chapter 3, Annex 3.1.A
Energy	Stationary combustion: liquid and gaseous fuels – CO ₂	28	Investigate further the issue on high IEF for gaseous fuels in petroleum refining and report accordingly in the next NIR	Ireland uses both national statistics fuel data and EU ETS emissions data for reporting of CO ₂ emissions from energy combustion in refinery. Unusual IEFs are a result of refinery gases/natural gas proportion that is reported in Energy Balance where activity data is derived from and different mix reported in EU ETS. Ireland is working closely with Energy Balance provider	Chapter 3, Annex 3.1.A, Annex 4

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
				on harmonising the fuel mix with that reported in EU ETS.	
Energy	Other transportation: liquid fuels – CO2	30	Reviews the notation key used to report liquid fuels and, as appropriate, change the notation key from “NO” to “IE”, and provide a transparent description on the basis for dividing fuel consumption between road and non-road traffic	Notation key IE in sector 1A5 Other applies to all other types of transportation i.e. non-road traffic that could not be further disaggregated and separated from stationary sources. The energy balance provider does not offer separate statistics for road and non-road transport.	Chapter 3
Energy	Oil and natural gas: gaseous fuels – CO2, CH4	31	Provide the explanation on where fugitive emissions of CH4 and CO2 from natural gas exploration and transmission are reported both in the CRF tables and in the NIR, and provide a detailed description of how the emissions from each activity are estimated in the NIR	Fugitive emissions of CH4 from natural gas for all activities are reported in two categories: category 1B2b2 Production and 1B2b5 Distribution. Other activities CH4 emissions from natural gas are reported as IE. CO2 emissions from natural gas are reported in pipeline compressors sector 1A3e other transportation.	
Energy	Oil and natural gas: gaseous fuels – CO2, CH4	32	Explain where fugitive CO2 emissions from natural gas and fugitive CH4 emissions from venting and flaring are allocated in the CRF tables	Fugitive CO2 emissions from natural gas are included together with combustion CO2 emissions in sector 1A3e other (pipeline compressors). Gas venting CH4 emissions could not be separated and are included in two fugitive categories gas is reported under: 1B2b2 Production and 1B2b5 Distribution. Gas flaring did not take place in	

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
				Ireland apart from two instances in two years (1999 and 2001) and CO ₂ emissions were reported accordingly for those years in category 1B2c2ii gas flaring.	
Energy	Oil and natural gas: gaseous fuels – CO ₂ , CH ₄	32	Use notation keys consistently between the NIR and the CRF tables	Notation keys have been applied consistently in CRF and NIR in 2015 submission.	
Energy	Oil and natural gas: gaseous fuels – CO ₂ , CH ₄	33	Use the appropriate notation keys and provide a detailed description of how the emissions from each activity are estimated in the NIR. Change fugitive emissions from other leakages from natural gas to IE (from NO).	Category 1B2b6 Other (i.e. leakages from natural gas) is not occurring NO as potential leakages from natural gas are considered negligible.	
Energy	Oil and natural gas: gaseous fuels – CO ₂ , CH ₄	34	Include the information on a mobile drilling unit in the Kinsale field in 2001 in the next NIR	Note added to NIR.	Chapter 3
Industrial processes and solvent and other product use	Consumption of halocarbons and SF ₆ – HFCs and SF ₆	40	Provide additional information of how the potential sources (e.g. from imported products) are considered in the emission estimates from this category to ensure a complete and accurate inventory	Potential sources are not estimated in 2015 submission (according to the 2006 IPCC Guidelines).	
Industrial processes and solvent and other product use	Limestone and dolomite use – CO ₂	41	Ensure consistency within the NIR and between the NIR and CRF tables in future submissions re stoichiometric ratio of CO ₂ to calcium carbonate (CaCO ₃) used as an EF. It fluctuates instead of being constant as stated in NIR (0.44 t CO ₂ /t limestone).	NIR text corrected.	Chapter 4
Agriculture	Manure management – CH ₄ and N ₂ O	50	Develop dynamic N excretion rates for non-dairy cattle and use the related data in the inventory, when the data become available	Inventory agency continues to engage with the agricultural research community to develop dynamic N excretion rates for non-dairy cattle.	
Agriculture	Agricultural soils – N ₂ O	51	Replace the default FracGASM data with country- specific data when they become available	The default FracGASM value will be replaced when country specific data is available.	

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
LULUCF	General	54	Follow the structure of NIR shown in the annex to decision 24/CP.19	Completed	
LULUCF	General	55	Include the information on key drivers of emission/removal trends on cropland, grassland, wetlands, settlements and other land in its NIR	Discussion of the key drivers of change in these categories is provided in the text in relevant sections of the NIR for Cropland and Grasslands. Discussion of drivers for changes in wetlands is restricted to Peat extraction activities. Further analysis is required to assess the underlying drivers of changes in settlements, due to the very complex economic and social interactions which influence behaviour in this sector.	6.4.1, 6.4.5, 6.4.9, 6.5.1, 6.5.4, 6.5.7, 6.6.3, 6.6.6, 6.6.7, 6.9
LULUCF	Forest land – CO ₂	56	Correct this typographical error on value of country-specific EF for organic forest soils	Completed	6.3.3.1.2
LULUCF	Forest land – CO ₂	57	Report the removals for the pool or report the pool as “NE” instead of “NO” or report the carbon stock changes as “NA” if the carbon stock changes in the pool are assumed to be zero because losses are balanced out by gains	Clarification in text, do not agree. Any notation can be used as long as it is clarified in text	6.3.3.1.2
LULUCF	Forest land – CO ₂	58	Delete the sentence “emissions from soils due to biomass burning resulting from forest wildfires are assumed to be negligible and do not occur (NO)” from its NIR in order to avoid confusion	Completed	6.3.4.4
LULUCF	Forest land – N ₂ O	59	Use the same tier to estimate the carbon stock changes in and the N ₂ O emissions from soils in the same category	Not valid there is new methodology under AFOLU 2006	
LULUCF	Wetlands remaining wetlands – CO ₂	60	Include the information on the carbon losses in DOM removed from managed wetlands in its NIR and the documentation box in CRF table in order to enhance transparency	Text has been added in the NIR. Will consider options for inclusion of information in CRF.	6.6.1
LULUCF	Land converted to wetlands – CO ₂	61	Include the information on mineral soils in wetlands in the next annual submission in order to clarify what kind of soils are included in wetland areas	To be considered in 2015 submission	

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
LULUCF	Settlements remaining settlements – CO2	62	Report the carbon stock changes in this category as “NA” instead of as “NO” and include an explanation on the use of the notation key to its NIR	To be considered in 2015 submission	
Waste	General	65	Correct inconsistencies between NIR table 8.6 and table I.2 in annex I to the NIR	Completed	
Waste	General	66	Fully document the sector-specific QA activities in the NIR.		
Waste	General	67	Expand the discussion of uncertainty in the waste chapter to include the uncertainty estimates for wastewater and incineration	Revised and more detailed uncertainty analysis has been developed and it includes estimates for wastewater and waste incineration and open burning.	Annex 2, NIR 2015
Waste	Solid waste disposal on land – CH4	68	Disaggregate the AD for the years up to 2003 in order to ensure time-series consistency		
Waste	Solid waste disposal on land – CH4	69	Update the information on MSW generation in the NIR and the CRF tables (error due to the non-inclusion of town dumps and the landfill site W047 in the national waste reports and information on waste composition in annex I to the NIR)		
Waste	Solid waste disposal on land – CH4	70	Include a discussion of these model parameters in FOD model in its next NIR, including the values used and justification for their use		
Waste	Wastewater handling – CH4 and N2O	71	Provide a discussion of the methodology used in the NIR in order to increase the transparency of its reporting		
Waste	Wastewater handling – CH4 and N2O	72	Describe the source and derivation of AD and the industrial sectors contributing to the BOD load		
Waste	Waste incineration – CO2, CH4 and N2O	73	Include a discussion in the NIR on the applicability of the EFs from the 2006 IPCC Guidelines to the incinerator units.		
Waste	Waste incineration – CO2, CH4 and N2O	74	Correct this double counting of AD in the quantity of clinical waste incinerated in the CRF tables by disaggregating the AD into biogenic and non- biogenic components	Waste incineration has been estimated separately for biogenic and non-biogenic components in 2015 submission.	
KP-LULUCF	General	77	Include a paragraph explaining the assessment of the key category analysis for the KP-LULUCF activities in chapter 11 of its NIR.		

Sector	Issue	ARR Paragraph	Recommendation	Party response	NIR Section
KP-LULUCF	Afforestation and reforestation – CO ₂ , CH ₄ and N ₂ O	79	Include this information on implied carbon stock change factors for organic soils in afforestation and reforestation in its NIR	Completed	Reported in ch 6
KP-LULUCF	Deforestation – CO ₂ , CH ₄ and N ₂ O	80	Include this information on implied carbon stock change factors for organic soils under deforestation in its NIR	Completed	reported in Ch 6 and cross reference in ch 11
KP-LULUCF	Deforestation – CO ₂ , CH ₄ and N ₂ O	81	Include this information that the sampling plot for measuring carbon stocks in above- and belowground biomass contained regenerating young broadleaf forest/scrub, and that stump and root biomass is greater than stems and branch biomass in the regenerating young broadleaf forest/scrub in its NIR in order to enhance transparency		

Annex 5.3

Standard Electronic Format (SEF) 2014

Party IE
 Submission Year 2015
 Reported Year 2014
 Commitment Period 1

Table 1. Total quantities of Kyoto Protocol units by account type at beginning of reported year						
Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	223,701,906	1,635,668	NO	903,808	NO	NO
Entity holding accounts	1,810,375	78,734	NO	5,277,156	1,051,429	NO
Article 3.3/3.4 net source cancellation accounts	NO	NO	NO	NO		
Non-compliance cancellation account	NO	NO	NO	NO		
Other cancellation accounts	735	NO	NO	2,059	NO	NO
Retirement account	66,719,736	1,238,883	NO	2,780,994	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	292,232,752	2,953,285	NO	8,964,017	1,051,429	NO

Party	IE											
Submission Year	2015											
Reported Year	2014											
Commitment Period	1											
Table 2a. Annual internal transactions												
	Additions						Subtractions					
Transaction type	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Art6 issuance and conversion												
Party verified projects		NO					NO		NO			
Independently verified projects		NO					NO		NO			
Art3.3 and 3.4 issuance or cancellation												
3.3 Afforestation reforestation			NO				NO	NO	NO	NO		
3.3 Deforestation			NO				NO	NO	NO	NO		
3.4 Forest management			NO				NO	NO	NO	NO		
3.4 Cropland management			NO				NO	NO	NO	NO		
3.4 Grazing land management			NO				NO	NO	NO	NO		
3.4 Revegetation			NO				NO	NO	NO	NO		
Art 12 afforestation and reforestation												
Replacement of expired tCERs							NO	NO	NO	NO	NO	
Replacement of expired ICERs							NO	NO	NO	NO		
Replacement for reversal of storage							NO	NO	NO	NO		NO
Replacement for non-submission of certification report							NO	NO	NO	NO		NO
Other cancellation							NO	NO	NO	NO	NO	NO
Subtotal		NO	NO				NO	NO	NO	NO	NO	NO

Transaction type	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Retirement	NO	NO	NO	NO	NO	NO

Party	IE											
Submission Year	2015											
Reported Year	2014											
Commitment Period	1											
Table 2b. Annual external transactions												
	Additions						Subtractions					
Transfers and acquisitions	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
CDM	NO	NO	NO	NO	142,192	NO	NO	NO	NO	NO	NO	NO
EU	11,320	NO	NO	38,093	NO	NO	NO	15,470	NO	372,267	NO	NO
GB	NO	11,700	NO	372,267	NO	NO	NO	NO	NO	NO	NO	NO
Subtotal	11,320	11,700	NO	410,360	142,192	NO	NO	15,470	NO	372,267	NO	NO

Additional Information												
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Independently verified ERU								NO				

Table 2c. Total annual transactions												
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Total (Sum of table 2(a) and 2(b))	11,320	11,700	NO	410,360	142,192	NO	NO	15,470	NO	372,267	NO	NO

Party	IE
Submission Year	2015
Reported Year	2014
Commitment Period	1

Table 3. Expiry, cancellation and replacement

	Expiry, cancellation and requirement to replace		Replacement					
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Transaction or event type								
Temporary CERs (tCERs)								
Expired in retirement and replacement accounts	NO							
Replacement of expired tCERs			NO	NO	NO	NO	NO	
Expired in holding accounts	NO							
Cancellation of tCERs expired in holding accounts	NO							
Long-term CERs (ICERs)								
Expired in retirement and replacement accounts		NO						
Replacement of expired ICERs			NO	NO	NO	NO		
Expired in holding accounts		NO						
Cancellation of ICERs expired in holding accounts		NO						
Subject to replacement for reversal of storage		NO						
Replacement for reversal of storage			NO	NO	NO	NO		NO
Subject to replacement for non-submission of certification report		NO						
Replacement for non-submission of certification report			NO	NO	NO	NO		NO
Total			NO	NO	NO	NO	NO	NO

Party	IE
Submission Year	2015
Reported Year	2014
Commitment Period	1

Table 4. Total quantities of Kyoto Protocol units by account type at end of reported year

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	223,713,226	1,635,668	NO	903,808	NO	NO
Entity holding accounts	1,810,375	74,964	NO	5,315,249	1,193,621	NO
Article 3.3/3.4 net source cancellation accounts	NO	NO	NO	NO		
Non-compliance cancellation account	NO	NO	NO	NO		
Other cancellation accounts	735	NO	NO	2,059	NO	NO
Retirement account	66,719,736	1,238,883	NO	2,780,994	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	292,244,072	2,949,515	NO	9,002,110	1,193,621	NO

Party	IE
Submission Year	2015
Reported Year	2014
Commitment Period	1

Table 5 a. Summary information on additions and subtractions												
	Additions						Subtractions					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Starting Values												
Issuance pursuant to Article 3.7 and 3.8	314,184,272											
Non-compliance cancellation							NO	NO	NO	NO		
Carry-over	NO	NO		NO								
Subtotal	314,184,272	NO		NO			NO	NO	NO	NO		
Annual Transactions												
Year 0 (2007)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 1 (2008)	2,029,090	NO	NO	3,778,980	NO	NO	1,245,569	NO	NO	108,000	NO	NO
Year 2 (2009)	3,237,702	NO	NO	9,285,054	NO	NO	3,269,878	NO	NO	5,762,220	NO	NO
Year 3 (2010)	5,439,324	722,440	NO	9,241,948	NO	NO	9,339,288	NO	NO	9,111,174	NO	NO
Year 4 (2011)	5,560,581	237,295	NO	13,946,988	NO	NO	18,004,498	435,000	NO	10,465,530	NO	NO
Year 5 (2012)	2,175,913	1,290,326	NO	6,237,051	503,489	NO	4,680,827	NO	NO	8,790,419	NO	NO
Year 6 (2013)	NO	1,693,196	NO	1,309,858	547,940	NO	3,854,805	554,972	NO	600,578	NO	NO
Year 7 (2014)	11,320	11,700	NO	410,360	142,192	NO	NO	15,470	NO	372,267	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Subtotal	18,453,930	3,954,957	NO	44,210,239	1,193,621	NO	40,394,865	1,005,442	NO	35,210,188	NO	NO
Total	332,638,202	3,954,957	NO	44,210,239	1,193,621	NO	40,394,865	1,005,442	NO	35,210,188	NO	NO

Table 5 b. Summary information on replacement								
	Expiry, cancellation and requirement to replace		Replacement					
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Previous CPs			NO	NO	NO	NO	NO	NO
Year 1 (2008)		NO	NO	NO	NO	NO	NO	NO
Year 2 (2009)		NO	NO	NO	NO	NO	NO	NO
Year 3 (2010)		NO	NO	NO	NO	NO	NO	NO
Year 4 (2011)		NO	NO	NO	NO	NO	NO	NO
Year 5 (2012)	NO	NO	NO	NO	NO	NO	NO	NO
Year 6 (2013)	NO	NO	NO	NO	NO	NO	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO
Total	NO	NO	NO	NO	NO	NO	NO	NO

Table 5 c. Summary information on retirement						
Year	Retirement					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Year 1 (2008)	NO	NO	NO	NO	NO	NO
Year 2 (2009)	19,668,515	NO	NO	713,192	NO	NO
Year 3 (2010)	16,991,714	NO	NO	223,643	NO	NO
Year 4 (2011)	16,230,638	394,883	NO	730,497	NO	NO
Year 5 (2012)	13,828,869	844,000	NO	1,113,662	NO	NO
Year 6 (2013)	NO	NO	NO	NO	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO
Total	66,719,736	1,238,883	NO	2,780,994	NO	NO

Party	IE											
Submission Year	2015											
Reported Year	2014											
Commitment Period	1											
Table 6 a. Memo item: corrective transactions relating to additions and subtractions												
Additions							Subtractions					
AAUs	ERUs	RMUs	CERs	tCERs	ICERs		AAUs	ERUs	RMUs	CERs	tCERs	ICERs

Table 6 b. Memo item: corrective transactions relating to replacement							
Expiry, cancellation and requirement to replace				Replacement			
tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs

Table 6 c. Memo item: corrective transactions relating to retirement						
Retirement						
AAUs	ERUs	RMUs	CERs	tCERs	ICERs	

Annex 5.4

Greenhouse Gases GWP and IPCC Reporting Format

Table 5.4.1 Greenhouse Gases and GWP Values

Greenhouse Gas	Chemical Formula	IPCC GWP (100-yr horizon) ^a
Carbon Dioxide	CO ₂	1
Methane	CH ₄	25
Nitrous Oxide	N ₂ O	298
Hydrofluorocarbons (HFCs)		
HFC-23	CHF ₃	14800
HFC-32	CH ₂ F ₂	675
HFC-41	CH ₃ F	92
HFC-43-10mee	CF ₃ CHFCHFCF ₂ CF ₃	1640
HFC-125	CHF ₂ CF ₃	3500
HFC-134	CHF ₂ CHF ₂	1100
HFC-134a	CH ₂ FCF ₃	1430
HFC-143	CH ₂ FCHF ₂	353
HFC-143a	CH ₃ CF ₃	4470
HFC-152	CH ₂ FCH ₂ F	53
HFC-152a	CH ₃ CHF ₂	124
HFC-161	CH ₃ CH ₂ F	12
HFC-227ea	CF ₃ CHF ₂ CF ₃	3220
HFC-236cb	CH ₂ FCF ₂ CF ₃	1340
HFC-236ea	CHF ₂ CHF ₂ CF ₃	1370
HFC-236fa	CF ₃ CH ₂ CF ₃	9810
HFC-245ca	CH ₂ FCF ₂ CHF ₂	693
HFC-245fa	CHF ₂ CH ₂ CF ₃	1030
HFC-365mfc	CH ₃ CF ₂ CH ₂ CF ₃	794
Perfluorocarbons		
PFC-14	CF ₄	7,390
PFC-116	C ₂ F ₆	12,200
PFC-218	C ₃ F ₈	8,830
PFC-3-1-10	C ₄ F ₁₀	8,860
PFC-318	c-C ₄ F ₈	10,300
PFC-4-1-12	C ₅ F ₁₂	9,160
PFC-5-1-14	C ₆ F ₁₄	9,300
PFC-9-1-18	C ₁₀ F ₁₈	>7,500
Perfluorocyclopropane	c-C ₃ F ₆	>17,340
Sulphur Hexafluoride	SF ₆	22,800
Nitrogen trifluoride	NF ₃	17,200

(a) GWP (global warming potential) as provided by the IPCC in its Fourth Assessment Report (WG1 errata to Table 2.14)

Table 5.4.2 IPCC Reporting Format (Level 1 and Level 2)

IPCC SOURCE and SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆	NF ₃
1. Energy							
A. Fuel Combustion (Sectoral Approach)							
1. Energy Industries							
2. Manufacturing Industries and Construction							
3. Transport							
4. Other Sectors							
5. Other							
B. Fugitive Emissions from Fuels							
1. Solid Fuels							
2. Oil and Natural Gas							
C. Carbon Dioxide Transport and Storage							
2. Industrial Processes and Product Use							
A. Mineral Industry							
B. Chemical Industry							
C. Metal Production							
D. Non-Energy Products from Fuels and Solvent Use							
E. Electronic Industry							
F. Product Uses as Substitutes for ODS							
G. Other Product Manufacture and Use							
H. Other							
3. Agriculture							
A. Enteric Fermentation							
B. Manure Management							
C. Rice Cultivation							
D. Agricultural Soils							
E. Prescribed Burning of Savannas							
F. Field Burning of Agricultural Residues							
G. Liming							
H. Urea Application							
I. Other							
4. Land-Use Land-Use Change and Forestry							
A. Forest Land							
B. Cropland							
C. Grassland							
D. Wetland							
E. Settlements							
F. Other Land							
G. Harvested Wood Products							
H. Other							
5. Waste							
A. Solid Waste Disposal							
B. Biological Treatment of Solid Waste							
C. Waste Incineration and Open Burning of Waste							
D. Wastewater Treatment and Discharge							
E. Other							
6. Other							
Memo Items:							
International Bunkers							
Aviation							
Navigation							
Multilateral Operations							
CO₂ Emissions from Biomass							
CO₂ captured							
Long-term storage of C in waste disposal sites							
Indirect N₂O							
Indirect CO₂							

The grey cells indicate sources/sinks where no emissions/removals of the various gases are expected

Appendix 1

Standard Independent Assessment Report (Electronic Appendix)

AN GHNÍOMHAIREACHT UM CHAOMHNÚ COMHSHAOIL

Tá an Gníomhaireacht um Chaomhnú Comhshaoil (GCC) freagrach as an gcomhshaoil a chaomhnú agus a fheabhsú mar shócmhainn luachmhar do mhuintir na hÉireann. Táimid tiomanta do dhaoine agus don chomhshaoil a chosaint ó éifeachtaí díobhálacha na radaíochta agus an truaillithe.

Is féidir obair na Gníomhaireachta a roinnt ina trí phríomhréimse:

Rialú: *Déanaimid córais éifeachtacha rialaithe agus comhlíonta comhshaoil a chur i bhfeidhm chun torthaí maithe comhshaoil a sholáthar agus chun díriú orthu siúd nach gcloíonn leis na córais sin.*

Eolas: *Soláthraímid sonraí, faisnéis agus measúnú comhshaoil atá ar ardchaighdeán, spriocdhírthe agus tráthúil chun bonn eolais a chur faoin gcinnteoireacht ar gach leibhéal.*

Tacaíocht: *Bímid ag saothrú i gcomhar le grúpaí eile chun tacú le comhshaoil atá glan, táirgiúil agus cosanta go maith, agus le hiompar a chuirfidh le comhshaoil inbhuanaithe.*

Ár bhFreagrachtaí

Ceadúnú

- Déanaimid na gníomhaíochtaí seo a leanas a rialú ionas nach ndéanann siad dochar do shláinte an phobail ná don chomhshaoil:
- saoráidí dramhaíola (m.sh. láithreáin líonta talún, loisceoirí, stáisiúin aistrithe dramhaíola);
- gníomhaíochtaí tionsclaíocha ar scála mór (m.sh. déantúsaíocht cógaisíochta, déantúsaíocht stroighne, stáisiúin chumhachta);
- an dionalmhaíocht (m.sh. muca, éanlaith);
- úsáid shrianta agus scaoileadh rialaithe Orgánach Géinmhodhnaithe (OGM);
- foinsí radaíochta ianúcháin (m.sh. trealamh x-gha agus radaiteiripe, foinsí tionsclaíocha);
- áiseanna móra stórála peitiril;
- scardadh dramhuisce;
- gníomhaíochtaí dumpála ar farraige.

Forfheidhmiú Náisiúnta i leith Cúrsaí Comhshaoil

- Clár náisiúnta iniúchtaí agus cigireachtaí a dhéanamh gach bliain ar shaoráidí a bhfuil ceadúnas ón nGníomhaireacht acu.
- Maoirseacht a dhéanamh ar fhreagrachtaí cosanta comhshaoil na n-údarás áitiúil.
- Caighdeán an uisce óil, arna sholáthar ag soláthraithe uisce phoiblí, a mhaoirsiú.
- Obair le húdaráis áitiúla agus le gníomhaireachtaí eile chun dul i ngleic le coireanna comhshaoil trí chomhordú a dhéanamh ar líonra forfheidhmiúcháin náisiúnta, trí dhírú ar chiontóirí, agus trí mhaoirsiú a dhéanamh ar leasúcháin.
- Cur i bhfeidhm rialachán ar nós na Rialachán um Dhramhthrealamh Leictreach agus Leictreonach (DTLL), um Shrian ar Shubstaintí Guaiseacha agus na Rialachán um rialú ar shubstaintí a idíonn an ciseal ózóin.
- An dlí a chur orthu siúd a bhriseann dlí an chomhshaoil agus a dhéanann dochar don chomhshaoil.

Bainistíocht Uisce

- Monatóireacht agus tuairisciú a dhéanamh ar cháilíocht aibhneacha, lochanna, uisce idirchriosacha agus cósta na hÉireann, agus screamhuisce; leibhéal uisce agus sruthanna aibhneacha a thomhas.
- Comhordú náisiúnta agus maoirsiú a dhéanamh ar an gCreat-Treoir Uisce.
- Monatóireacht agus tuairisciú a dhéanamh ar Cháilíocht an Uisce Snámha.

Monatóireacht, Anailís agus Tuairisciú ar an gComhshaoil

- Monatóireacht a dhéanamh ar cháilíocht an aeir agus Treoir an AE maidir le hAer Glan don Eoraip (CAFÉ) a chur chun feidhme.
- Tuairisciú neamhspleách le cabhrú le cinnteoireacht an rialtais náisiúnta agus na n-údarás áitiúil (m.sh. tuairisciú tréimhsiúil ar staid Chomhshaoil na hÉireann agus Tuarascálacha ar Tháscairí).

Rialú Astaíochtaí na nGás Ceaptha Teasa in Éirinn

- Fardail agus réamh-mheastacháin na hÉireann maidir le gáis ceaptha teasa a ullmhú.
- An Treoir maidir le Trádáil Astaíochtaí a chur chun feidhme i gcomhair breis agus 100 de na táirgeoirí dé-ocsaíde carbóin is mó in Éirinn

Taighde agus Forbairt Comhshaoil

- Taighde comhshaoil a chistiú chun brúnna a shainathint, bonn eolais a chur faoi bheartais, agus réitigh a sholáthar i réimsí na haeráide, an uisce agus na hinbhuanaitheachta.

Measúnacht Straitéiseach Timpeallachta

- Measúnacht a dhéanamh ar thionchar pleananna agus clár beartaithe ar an gcomhshaoil in Éirinn (m.sh. mórfhleananna forbartha).

Cosaint Raideolaíoch

- Monatóireacht a dhéanamh ar leibhéal radaíochta, measúnacht a dhéanamh ar nochtadh mhuintir na hÉireann don radaíocht ianúcháin.
- Cabhrú le pleananna náisiúnta a fhorbairt le haghaidh éigeandálaí ag eascairt as taismí núicléacha.
- Monatóireacht a dhéanamh ar fhorbairtí thar lear a bhaineann le saoráidí núicléacha agus leis an tsábháilteacht raideolaíochta.
- Sainseirbhísí cosanta ar an radaíocht a sholáthar, nó maoirsiú a dhéanamh ar sholáthar na seirbhísí sin.

Treoir, Faisnéis Inrochtana agus Oideachas

- Comhairle agus treoir a chur ar fáil d'earnáil na tionsclaíochta agus don phobal maidir le hábhair a bhaineann le caomhnú an chomhshaoil agus leis an gcosaint raideolaíoch.
- Faisnéis thráthúil ar an gcomhshaoil ar a bhfuil fáil éasca a chur ar fáil chun rannpháirtíocht an phobail a spreagadh sa chinnteoireacht i ndáil leis an gcomhshaoil (m.sh. Timpeall an Tí, léarscáileanna radóin).
- Comhairle a chur ar fáil don Rialtas maidir le hábhair a bhaineann leis an tsábháilteacht raideolaíoch agus le cúrsaí práinnfhreagartha.
- Plean Náisiúnta Bainistíochta Dramhaíola Guaisí a fhorbairt chun dramhaíl ghuaiseach a chosc agus a bhainistiú.

Múscailt Feasachta agus Athrú Iompraíochta

- Feasacht chomhshaoil níos fearr a ghiniúint agus dul i bhfeidhm ar athrú iompraíochta dearfach trí thacú le gnóthais, le pobail agus le teaghlacha a bheith níos éifeachtúla ar acmhainní.
- Tástáil le haghaidh radóin a chur chun cinn i dtithe agus in ionaid oibre, agus gníomhartha leasúcháin a spreagadh nuair is gá.

Bainistíocht agus struchtúr na Gníomhaireachta um Chaomhnú Comhshaoil

Tá an ghníomhaíocht á bainistiú ag Bord lánaimseartha, ar a bhfuil Ard-Stiúrthóir agus cúigear Stiúrthóirí. Déantar an obair ar fud cúig cinn d'Oifigí:

- An Oifig Aeráide, Ceadúnaithe agus Úsáide Acmhainní
- An Oifig Forfheidhmithe i leith cúrsaí Comhshaoil
- An Oifig um Measúnú Comhshaoil
- An Oifig um Cosaint Raideolaíoch
- An Oifig Cumarsáide agus Seirbhísí Corparáideacha

Tá Coiste Comhairleach ag an nGníomhaireacht le cabhrú léi. Tá dáréag comhaltaí air agus tagann siad le chéile go rialta le plé a dhéanamh ar ábhair imní agus le comhairle a chur ar an mBord.

Headquarters
PO Box 3000, Johnstown Castle Estate
County Wexford, Ireland
Bosca Poist 3000, Eastát Chaisleán Bhaile Sheáin
Contae Loch Garman Éire

T: +353 53 9160600
F: +353 53 9160699
E: info@epa.ie
W: www.epa.ie
Lo Call: 1890 33 55 99

EPA Regional Inspectorate Dublin
McCumiskey House
Richview
Clonskeagh Road
Dublin 14
Tel: 01-268 0100
Fax: 01-268 0199

EPA Regional Inspectorate Cork
Inniscarra
Co. Cork
Tel: 021-4875540
Fax: 021-4875545

EPA Regional Inspectorate Castlebar
John Moore Road
Castlebar
Co. Mayo
Tel: 094-9048400
Fax: 094-9021934

EPA Regional Inspectorate Kilkenny
Seville Lodge
Callan Road
Kilkenny
Tel: 056-7796700
Fax: 056-7796798

EPA Regional Inspectorate Monaghan
The Glen
Monaghan
Tel: 047-77600
Fax: 047-84987

E: info@epa.ie
W: www.epa.ie
LoCall: 1890 33 55 99

