

Switzerland's Greenhouse Gas Inventory 1990–2015

National Inventory Report

Including reporting elements under the Kyoto Protocol

Submission of April 2017
under the United Nations Framework Convention on Climate Change
and under the Kyoto Protocol



Schweizerische Eidgenossenschaft
Confédération suisse
Confederazione Svizzera
Confederaziun svizra

Swiss Confederation

Federal Office for the Environment FOEN

Publisher

Federal Office for the Environment FOEN, Climate Division, 3003 Bern, Switzerland
www.bafu.admin.ch/climate
www.climatereporting.ch

Inventory Process Manager

Regine Röthlisberger FOEN, Climate Division, Bern

National Inventory Compiler

Anouk-Aimée Bass FOEN, Air Pollution Control and Chemicals Division, Bern

NIR Lead Authors

Daniel Bretscher Agroscope, Zürich

Jürg Heldstab INFRAS, Zürich

Beat Rihm Meteotest, Bern

Nele Rogiers FOEN, Forest Division, Bern

Bettina Schächli INFRAS, Zürich

Markus Sommerhalder CSD, Bern

Cornelia Stettler Carbotech, Basel

Felix Weber INFRAS, Zürich

QA/QC Officer

Michael Bock FOEN, Climate Division, Bern

National Registry Administrator

Stefan Meier FOEN, Climate Division, Bern

Bern, April 2017

Switzerland's Greenhouse Gas Inventory 1990–2015

National Inventory Report 2017

Including reporting elements under the Kyoto Protocol

Submission of April 2017
under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol

NIR Chapter / Sector	Responsible Persons
Executive Summary	Author: Jürg Heldstab (INFRAS)
1. Introduction	Lead Author: Jürg Heldstab (INFRAS) Authors: Regine Röhliberger (FOEN; QA/QC), Bettina Schächli (INFRAS; Uncertainty analyses), Adrian Schilt (FOEN; KCA), Felix Weber (INFRAS; Key categories, Uncertainty analyses)
2. Trends in GHG emissions and removals	Lead Author: Jürg Heldstab (INFRAS) Author: Felix Weber (INFRAS)
3. Energy	Lead Author: Jürg Heldstab (INFRAS) Authors: Jürg Heldstab (INFRAS; Overview, RA/SA, Bunker fuels, Feedstocks, Country-specific issues, Mobile sources in 1A2-1A5), Bettina Schächli (INFRAS; Stationary sources in 1A1, 1A2, 1A4) Sector expert for Civil Aviation: Theo Rindlisbacher (FOCA) Technical contributors: Anouk-Aimée Bass (FOEN; EMIS data base operation, QA/QC checks), Benedict Notter (INFRAS; non-road), Sabine Schenker (FOEN; Reference Approach), Philipp Wüthrich (INFRAS; Road transportation)
4. Industrial processes and product use	Lead Authors: Bettina Schächli (INFRAS), Cornelia Stettler (Carbotech; F-gases) Authors: Bettina Schächli (INFRAS), Sabine Schenker (FOEN; EMIS data base operation, QA/QC checks), Cornelia Stettler (Carbotech; F-gases) Technical contributor (F-gases): Cornelia Stettler (Carbotech)
5. Agriculture	Lead Author: Daniel Bretscher (Agroscope) Author and technical contributor: Daniel Bretscher (Agroscope)
6. LULUCF	Project Leader: Andreas Schellenberger (FOEN) Lead Author: Beat Rihm (Meteotest) Authors and Sector experts: Daniel Bretscher (Agroscope; Cropland, Grassland), Paolo Camin (FOEN; HWP), Markus Didion (WSL; Yasso07), Andreas Gubler (Agroscope; NABO), Armin Keller (Agroscope: NABO), Jens Leifeld (Agroscope; non-Forest Land), Nele Rogiers (FOEN; Forest Land), Andreas Schellenberger (FOEN; QA/QC, Cross-cutting issues) Sector experts: Sonja Keel (Agroscope), Lukas Mathys (Nategra), Sonja Paul (University of Basel), Esther Thürig (WSL), Chloé Wüst (Agroscope) Technical contributors: Christoph Könitzer (Sigmaphan), Martin Lindenmann (Sigmaphan), Stefan Müller (Meteotest), Felix Weibel (SFSO)

NIR Chapter / Sector	Responsible Persons (continued)
7. Waste	Lead Author: Markus Sommerhalder (CSD) Author: Rainer Kegel (FOEN; EMIS data base operation, QA/QC checks), Markus Sommerhalder (CSD)
8. Other (IPCC sector 7)	Lead Author: Markus Sommerhalder (CSD) Authors: Markus Sommerhalder (CSD), Felix Weber (INFRAS)
9. Indirect CO ₂ and N ₂ O emissions	Lead Author: Bettina Schächli (INFRAS) Authors: Bettina Schächli (INFRAS); Adrian Schilt (FOEN) Sector experts: Sabine Schenker (FOEN), Adrian Schilt (FOEN)
10. Recalculations	Lead Author: Bettina Schächli (INFRAS) Authors: Daniel Bretscher (Agroscope), Jürg Heldstab (INFRAS), Beat Rihm (Meteotest), Nele Rogiers, (FOEN), Bettina Schächli (INFRAS), Andreas Schellenberger (FOEN), Cornelia Stettler (Carbotech), Felix Weber (INFRAS)
11. KP-LULUCF	Author: Nele Rogiers (FOEN) Sector experts: Paolo Camin (FOEN), Markus Didion (WSL), Esther Thürig (WSL)
12. Accounting of Kyoto units	Authors: Stefan Meier (FOEN), Regine Röthlisberger (FOEN) Sector experts: Susanne Riedener (FOEN)
13. Changes in National Registry	Author: Stefan Meier (FOEN)
14. Minimization of adverse effects	Authors: Regine Röthlisberger (FOEN)
Annex 1 KCA	Author: Felix Weber (INFRAS) Sector expert: Anouk-Aimée Bass (FOEN), Adrian Schilt (FOEN)
Annex 2 Assessment of uncertainty	Authors: Bettina Schächli (INFRAS), Felix Weber (INFRAS)
Annex 3 Other detailed information...	Authors: Jürg Heldstab (INFRAS), Bettina Schächli (INFRAS), Cornelia Stettler (Carbotech)
Annex 4 National energy balance	Authors: Felix Weber (INFRAS)
Annex 5 Additional information	Authors: Stefan Reimann (EMPA; F-gases), Dominik Brunner (EMPA; CH ₄), Stephan Henne (EMPA; CH ₄)
Annex 6 Information on the CRF Reporter	Author: Regine Röthlisberger (FOEN)
Annex 7 Information on the UNFCCC review process	Author: Felix Weber (INFRAS)

Table of Contents

Table of Contents	5
Glossary.....	9
Executive summary	13
Acknowledgements	26
PART 1	27
1 Introduction.....	27
1.1 Background information on Swiss greenhouse gas inventories, climate change and supplementary information of the Kyoto Protocol (KP).....	27
1.2 National inventory arrangements.....	29
1.3 Inventory preparation and data collection, processing, and storage.....	37
1.4 Methodologies and data sources	38
1.5 Description of key categories	40
1.6 General uncertainty evaluation.....	48
1.7 General assessment of completeness	56
2 Trends in greenhouse gas emissions and removals.....	58
2.1 Aggregated greenhouse gas emissions 2015 (UNFCCC)	58
2.2 Emission trends by gas	60
2.3 Emission trends by sources and sinks	62
2.4 Emission trends for precursor gases and SO ₂	68
2.5 Emission trends (Kyoto Protocol)	71
3 Energy	73
3.1 Overview.....	73
3.2 Source category 1A – Fuel combustion activities.....	77
3.3 Source category 1B - Fugitive emissions from fuels	181
3.4 Source category 1C – CO ₂ transport and storage.....	192
4 Industrial processes and product use	193
4.1 Overview.....	193
4.2 Source category 2A – Mineral industry	196
4.3 Source category 2B – Chemical industry	211
4.4 Source category 2C – Metal industry	221
4.5 Source category 2D – Non-energy products from fuels and solvent use.....	228
4.6 Source category 2E – Electronics industry.....	238

4.7	Source category 2F – Product uses as substitutes for ozone depleting substances.....	240
4.8	Source category 2G – Other product manufacture and use	253
4.9	Source category 2H – Other.....	263
5	Agriculture	267
5.1	Overview.....	267
5.2	Source category 3A – Enteric fermentation.....	270
5.3	Source category 3B – Manure management.....	283
5.4	Source category 3C – Rice cultivation	302
5.5	Source category 3D – Agricultural soils	302
5.6	Source category 3E – Prescribed burning of savannahs	320
5.7	Source category 3F – Field burning of agricultural residues.....	320
5.8	Source category 3G – Liming.....	320
5.9	Source category 3H – Urea application.....	322
6	LULUCF	324
6.1	Overview of LULUCF.....	324
6.2	Land-use definitions and classification systems.....	338
6.3	Approaches used for representing land areas, land-use databases.....	345
6.4	Category 4A – Forest land.....	349
6.5	Category 4B – Cropland	383
6.6	Category 4C – Grassland	389
6.7	Category 4D – Wetlands	400
6.8	Category 4E – Settlements.....	403
6.9	Category 4F – Other land	406
6.10	Categories 4(III) and 4(IV) – N ₂ O emissions	407
6.11	Category 4G – Harvested wood products (HWP)	410
7	Waste	415
7.1	Overview.....	415
7.2	Source category 5A – Solid waste disposal	420
7.3	Source category 5B – Biological treatment of solid waste	426
7.4	Source category 5C – Incineration and open burning of waste	430
7.5	Source category 5D – Wastewater treatment and discharge.....	435
7.6	Source category 5E – Other	444
8	Other.....	447
8.1	Overview.....	447

8.2	Source category 6 – Other	448
9	Indirect CO ₂ and N ₂ O emissions	451
9.1	Overview	451
9.2	Methodological issues	455
10	Recalculations and improvements	460
10.1	Explanations and justifications for recalculations, including in response to the review process	460
10.2	Implications for emission levels	478
10.3	Implications for emissions trends, including time series consistency.....	479
10.4	Planned improvements, including in response to the review process.....	480
PART 2	482
11	KP-LULUCF	482
11.1	General information	485
11.2	Land-related information.....	487
11.3	Activity-specific information	490
11.4	Article 3.3.....	501
11.5	Article 3.4.....	505
11.6	Other information	511
11.7	Information Relating to Article 6	512
12	Information on accounting of Kyoto Units	513
12.1	Background information.....	513
12.2	Summary of information reported in the SEF tables	513
12.3	Discrepancies and notifications	514
12.4	Publicly accessible information.....	514
12.5	Calculation of the Commitment Period Reserve (CPR)	515
12.6	KP-LULUCF Accounting.....	515
13	Information on changes in National Registry	516
14	Information on minimization of adverse impacts in accordance with Article 3, Paragraph 14.....	517
Annexes	521
Annex 1: Key category analysis (KCA)	521
Annex 2: Assessment of uncertainty	523	
Annex 3: Other detailed methodological descriptions for individual source or sink categories	536	
Annex 4: National energy balance and reference approach	560	
Annex 5: Additional information.....	566	

Annex 6: Information on the CRF Reporter 578

Annex 7: Information on the UNFCCC review process 582

References 590

References to EMIS database comments 623

Glossary

AD	Activity data
AFOLU	Agriculture, Forestry and Other Land Use
AREA1	Swiss Land Use Statistics 1979/85 (ASCH1 data re-evaluated according to the AREA set of land-use and land-cover categories)
AREA2	Swiss Land Use Statistics 1992/97 (ASCH2 data re-evaluated according to the AREA set of land-use and land-cover categories)
AREA3	Swiss Land Use Statistics, third survey 2004/09
AREA4	Swiss Land Use Statistics, fourth survey 2013/18
ART	Agroscope Reckenholz-Tänikon Research Station (formerly FAL) since 2014 Agroscope
ASCH1	Swiss Land Use Statistics, first survey 1979/85
ASCH2	Swiss Land Use Statistics, second survey 1992/97
AAU	Assigned Amount Unit (under the Kyoto Protocol)
BCEF, BEF	Biomass conversion and expansion factor, biomass expansion factor
Carbura	Swiss organisation for the compulsory stockpiling of oil products
Cemuisse	Association of the Swiss Cement Industry
CER	Certified Emission Reduction (under the Kyoto Protocol)
CC	Combination category
CDM	Clean Development Mechanism (under the Kyoto Protocol)
CFC	Chlorofluorocarbon (organic compound: refrigerant, propellant)
CH ₄	Methane, 2006 IPCC GWP: 25 (UNFCCC 2014a, Annex III)
CHP	Combined heat and power
chp.	Chapter
CNG	Compressed natural gas
CLRTAP	UNECE Convention on Long-Range Transboundary Air Pollution
CO	Carbon monoxide
CO ₂ , CO ₂ eq	Carbon dioxide, carbon dioxide equivalent
CORINAIR	CORe INventory of AIR emissions (under the European Topic Centre on Air Emissions and under the European Environment Agency)
CRF	Common reporting format
DBH	Diameter (of trees) at breast height
DDPS	Federal Department of Defence, Civil Protection and Sport
DETEC	Dept. of the Environment, Transport, Energy and Communications
EF	Emission factor
EMEP	European Monitoring and Evaluation Programme (under the Convention on Long-range Transboundary Air Pollution)
EMIS	Swiss Emission Information System

EMPA	Swiss Federal Laboratories for Material Testing and Research
ERT	Expert review team (under the UNFCCC and the Kyoto Protocol)
ERU	Emission Reduction Unit (under the Kyoto Protocol)
EV	Erdöl-Vereinigung (Swiss Petroleum Association)
FAL	Swiss Federal Research Station for Agroecology and Agriculture (since 2006: ART; since 2014 Agroscope)
FCA	Federal Customs Administration
FEDRO	Swiss Federal Roads Office
FiBL	Research Institute of Organic Agriculture
FMRL	Forest management reference level
FOAG	Federal Office for Agriculture
FOCA	Federal Office of Civil Aviation
FOD	First order decay (model)
FOEN	Federal Office for the Environment (former name SAEFL until 2005)
FOITT	Federal Office of Information Technology, Systems and Telecommunication
GHG	Greenhouse gas
GL	Guidelines
g	Gramme
GVS	Swiss Foundry Association
GWP	Global Warming Potential
ha	hectare
HFC	Hydrofluorocarbons (e.g. HFC-32 difluoromethane)
HWP	Harvested Wood Products
ICAO	International Civil Aviation Organization
IDM	FOEN Internal Document Management System
IDP	Inventory Development Plan
IPCC	Intergovernmental Panel on Climate Change
IPPU	Industrial processes and product use
JI	Joint Implementation (under the Kyoto Protocol)
KCA	Key category analysis
kha	Kilo hectare
kt	Kilo tonne (1'000 tonnes)
LPG	Liquefied Petroleum Gas (Propane/Butane)
LTO	Landing-Take-off-Cycle (Aviation)
LULUCF	Land Use, Land-Use Change and Forestry
MOFIS	Swiss federal vehicle registration database
MSW	Municipal solid waste
NABO	Swiss Soil Monitoring Network

NCV	Net calorific value
NEU	Non-energy use of fuels
NF ₃	Nitrogen trifluoride 2006 IPCC GWP: 17'200 (UNFCCC 2014a, Annex III)
NFI1, NFI2	First (1983–1985), Second (1993–1995), Third (2004–2006)
NFI3, NFI4	and Fourth (2009–2017) National Forest Inventory
NIR	National Inventory Report
NIS	National Inventory System
NFR	Nomenclature for Reporting (under the UNECE)
NMVOC	Non-methane volatile organic compounds
N ₂ O	Nitrous oxide; 2006 IPCC GWP: 298 (UNFCCC 2014a, Annex III)
NO _x	Nitrogen oxides
ODS	Ozone-depleting substances (CFCs, halons etc.)
PFC	Perfluorinated carbon compounds (e.g. Tetrafluoromethane)
SAEFL	Swiss Agency for the Environment, Forests and Landscape (since 2006: Federal Office for the Environment FOEN)
SEF	Standard electronic format (under the Kyoto Protocol)
SBV	Schweizerischer Bauernverband; Swiss Farmers Union
SF ₆	Sulphur hexafluoride, 2006 IPCC GWP: 22800 (UNFCCC 2014a, Annex III)
SFOE	Swiss Federal Office of Energy
SFSO	Swiss Federal Statistical Office
SGWA	Swiss Gas and Water Industry Association (see SVGW / SSIGE)
SKW	Schweizerischer Kosmetik- und Waschmittelverband (Swiss association of cosmetics and detergents)
SO ₂	Sulphur dioxide
SOC	Soil organic carbon
SOLV	Swiss Organisation for the Solvent Recovery of Industrial Enterprises in the Packaging Sector
SVGW / SSIGE	Schweizerischer Verein des Gas- und Wasserfaches / Société Suisse de l'Industrie du Gaz et des Eaux (Swiss Gas and Water Industry Association)
SWISSMEM	Swiss Mechanical and Electrical Engineering Industries (Schweizer Maschinen-, Elektro- und Metallindustrie)
tCER	Temporary Certified Emission Reduction (under the Kyoto Protocol)
QA/QC	Quality assurance/Quality control
QMS	Quality management system
UNECE	United Nations Economic Commission for Europe
UNFCCC	United Nations Framework Convention on Climate Change
VOC	Volatile organic compounds

VSG	Verband der Schweizerischen Gasindustrie / Association Suisse de l'Industrie Gazière (ASIG) (Swiss gas industry association)
VSZ	Verband Schweizerische Ziegelindustrie (Swiss association of brick and tile industry)
VSLF	Swiss association for coating and paint applications
VSTB	Swiss Association of Grass Drying Plants
WSL	Swiss Federal Institute for Forest, Snow and Landscape Research
WWT	Wastewater treatment
ZPK	Verband der Schweizerischen Zellstoff-, Papier- und Kartonindustrie

Executive summary

ES 1 Background information on greenhouse gas inventories, climate change and supplementary information required under Art. 7.1. KP

ES 1.1 Background information on climate change

In 2016 a comprehensive overview on Swiss climate – basics, consequences and perspectives – has been published by the Swiss Academies of Sciences (SCNAT 2016). Recent data confirms a warming trend in Switzerland with an observed increase in the mean annual temperature of 1.75°C between 1864 and 2012 (FOEN 2014d). Over the last 30 years Swiss temperature has increased with an annual average warming rate of 0.35°C/decade (CH2011 2011). The most visible change in the Alps resulting from global warming is the retreat of glaciers with a volume loss of 12% since 1999 (FOEN 2014d).

The observed trends in precipitation are less pronounced than in temperature. They generally show an increase in winter and spring, whereas for summer and autumn no significant trends are detectable. Regional scenarios predict an increase in mean winter precipitation and a decrease in summer, which will have a marked impact on the hydrological cycle. Furthermore, higher intensity of storms as well as reduced snowfall and snow cover duration are expected, increasing the risk and frequency of floods, landslides and debris flows.

Concerning biodiversity, climate change is expected to affect species composition, distribution, their cycles, synchronicity, the overall genetic diversity and the provision of ecosystem services. This in turn would raise the vulnerability of forests and potentially impair their protective, productive, and social functions.

For agriculture, a moderate warming of 2°C to 3°C might increase productivity; however, if the temperature rises beyond that level, the increase in heat waves and drought periods would prove problematic for the cultivation of land and for livestock husbandry.

Various sectors of the Swiss economy are likely to be adversely affected by progressing climate change: in particular, winter tourism would suffer from increased scarcity of snow, hydroelectric power stations are confronted with altered runoff and sediment transport regimes, and insurance companies may face increased losses due to winter storms and floods. Natural hazards and extreme weather events potentially pose a growing risk to infrastructure and human health. Heat waves in combination with elevated tropospheric ozone levels are cause for serious concern. Finally, it remains to be seen to what extent vector borne diseases spread due to changing climatic conditions. Recently Switzerland analysed these challenges in detail and developed an effective adaptation strategy in order to hedge against negative effects resulting from climate change in Switzerland (FOEN 2012b).

ES.1.2 Background information on greenhouse gas inventories

On 10 December 1993, Switzerland ratified the United Nations Framework Convention on Climate Change (UNFCCC). Since 1996, the submission of its national greenhouse gas inventory has been based on the 2006 IPCC Guidelines. From 1998 onwards, the inventories

have been submitted in the Common Reporting Format (CRF). In 2004, Switzerland started submitting a yearly National Inventory Report (NIR) under the UNFCCC.

On 9 July 2003, Switzerland ratified the Kyoto Protocol (KP) under the UNFCCC. The Swiss National Inventory System (NIS) according to Article 5.1 of the Kyoto Protocol was implemented and is fully operational.

The 2017 inventory submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol includes the NIR on hand, the greenhouse gas inventory 1990–2015 including also the Kyoto Protocol LULUCF tables 2008–2015 in the common reporting format as well as the standard electronic format (SEF) tables and the standard independent assessment report (SIAR) from the National Registry.

The Federal Office for the Environment (FOEN) is in charge of compiling the emission data and bears overall responsibility for Switzerland's national greenhouse gas inventory and the national registry. In addition to the FOEN, the Swiss Federal Office of Energy (SFOE), Agroscope, the Swiss centre of excellence for agricultural research, and the Federal Office of Civil Aviation (FOCA) participate directly in the compilation of the inventory. Several other administrative offices and research institutions are involved in the preparation of the inventory.

In preparing the national greenhouse gas inventory, Switzerland took into account some of the recommendations and encouragements of the "Draft Report on the individual review of the annual submission of Switzerland submitted in 2016" (UNFCCC 2017). The changes in response to the review process are documented in chp. 10.1.1). Furthermore, improvements addressed already in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016 are presented in the sectoral chapters and chp. 10.1. Note, these issues were considered to be resolved by the expert review team (ERT) and were implemented in the reporting tables submitted on 7 November 2016. For completeness, the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

The structure of Switzerland's NIR corresponds to the UNFCCC annotated outline (UNFCCC 2014a) and it contains three parts:

PART 1 reports the obligations under the UNFCCC,

PART 2 shows the additional obligations under the Kyoto Protocol and several

Annexes provide detailed information on selected issues of Part 1 and Part 2.

Chapter 1 of the NIR, the introduction, provides an overview of Switzerland's National System including institutional arrangements for producing the inventory, the process and methodologies used for inventory preparation, and the QA/QC procedures.

- The data sources used to compile the national inventory and to estimate greenhouse gas emissions and removals are the Swiss Emission Information System (EMIS), national energy statistics, data from industry associations, as well as further statistics and models for road transportation, off-road vehicles and machinery, agriculture, land use, land-use change and forestry (LULUCF), and waste. Emissions are calculated according to methodologies recommended by the 2006 IPCC Guidelines (IPCC 2006) including the recommended nomenclature and methodologies concerning uncertainty and QA/QC activities. The data in the EMIS database are pre-processed in order to enable transfers to the CRF Reporter required for reporting under the UNFCCC and under the Kyoto Protocol.

- All inventory data are assembled and prepared for input into the CRF Reporter by the GHG Inventory Core Group, which is responsible for ensuring the conformity of the inventory with the Updated UNFCCC Reporting Guidelines on Annual Inventories (UNFCCC 2014a) and the Guidance for reporting information on activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (UNFCCC 2014b). In the preparation of this report, the Inventory Core Group was supported by consultants. Their mandate included editing of the NIR, data quality controls, and an analysis of the consistency between the emission modelling and the recommendations of the IPCC Good Practice Guidance. Furthermore, the consultants contributed to the key category analyses and carried out the uncertainty analyses. They were also involved in improving the inventory, e.g. by performing tasks defined in the Inventory Development Plan (IDP).
- The inventory quality management system (QMS) is designed to comply with the objectives of good practice guidance, i.e. to ensure and improve transparency, consistency, comparability, completeness, accuracy and confidence in national GHG emission and removal estimates. The QA/QC Officer is responsible for the enforcement of the defined quality standards. The National Inventory System complies with the ISO 9001:2008 standard (Quality Management System) and is certified by the Swiss TS Technical Services AG (Swiss-TS 2016).
- A National Inventory System Supervisory Board was established by decision of the FOEN Directorate in summer 2006. The Board oversees activities related to the GHG Inventory and to the National Registry.

Chapter 1 provides information on key categories and uncertainties.

Chapter 2 contains an analysis of trends in Switzerland's greenhouse gas emissions by sources and removals by sinks for all sectors.

Chapters 3 to 9 provide principal source and sink category estimates.

Chapter 10 justifies, explains and summarises the recalculations and improvements. It also contains an overview of the planned improvements.

In **PART 2, Chapter 11** reports KP LULUCF data, **Chapter 12** presents information on accounting of Kyoto Units, **Chapter 13** lists changes in the National Registry, and **Chapter 14** includes information on minimization of adverse impacts in accordance with Article 3, paragraph 14.

ES.1.3 Background information on supplementary information required under article 7.1. of the Kyoto Protocol (KP)

As described above, Chapter 11 of PART 2 provides information on KP-LULUCF.

Switzerland only accounts for the mandatory activity Forest management under Article 3, paragraph 4 of the Kyoto Protocol (FOEN 2016c). In accordance with Annex I to Decision 2/CMP.7 (Annex I, Para 13), credits from Forest management are capped in the second commitment period. Thus for Switzerland the cap is set at 3.5% of the 1990 emissions (excluding LULUCF).

Switzerland chooses to account for emissions and removals from activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (FOEN 2016c, FOEN 2016d) over the entire second commitment period. In addition to the mandatory submission of the inventory years 2013–2015, data for the years 1999–2012 are available and shown in Switzerland's NIR.

ES.2 Summary of national emission and removal related trends, and emissions and removals from KP-LULUCF activities

ES.2.1 GHG Inventory 2015

In 2015, Switzerland emitted 48'038 kt (kilo tonnes) of CO₂ equivalent (CO₂ eq), corresponding to 5.8 tonnes of CO₂ equivalent per capita (CO₂: 4.7 tonnes per capita), to the atmosphere, excluding emissions from international bunkers (aviation and marine), excluding indirect greenhouse gas emissions and excluding emissions and removals from the sector Land use, land-use change, and forestry (LULUCF). For the emissions that are relevant under the Kyoto Protocol see chapter ES.3.3.

Key category analysis (KCA)

Several key category analyses are carried out by level (years 1990 and 2015) and trend assessment (period 1990–2015), both including LULUCF categories (see details in chp. 1.5.1.2 and IPCC (2006)).

- Approach 1: in 2015, 31 categories among a total of 167 are identified as level key categories. About half of these categories are part of sector 1 Energy, accounting for the largest share of total national emissions.
- Approach 2: in 2015, 23 categories among a total of 167 categories are identified as level key categories. Under Approach 2, the most important categories stem from sectors 3 Agriculture and 4 LULUCF.

Key category analyses are also performed excluding LULUCF categories. They are not represented in the NIR but are available on request.

Switzerland's GHG emissions by gases

Table E- 1 shows Switzerland's annual GHG emissions by individual gases from 1990 (base year) to 2015. Total emissions excluding LULUCF reach a minimum in 2014, which is 8.8% below base year emissions in 1990.

Table E- 1 Switzerland's GHG emissions in CO₂ equivalent (kt) by gas; 1990–2015. The column below on the far right indicates the percentage change in emissions in 2015 as compared to the base year 1990. HFCs increased by 6'198'125% when compared to 1990 levels (1990 = 0.025 kt CO₂ equivalent).

Greenhouse Gas Emissions	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
CO ₂ emissions including net CO ₂ from LULUCF	43'776	39'647	48'532	43'474
CO ₂ emissions excluding net CO ₂ from LULUCF	44'171	43'423	43'607	45'799
CH ₄ emissions including CH ₄ from LULUCF	6'132	5'802	5'411	5'321
CH ₄ emissions excluding CH ₄ from LULUCF	6'102	5'782	5'396	5'307
N ₂ O emissions including N ₂ O from LULUCF	2'916	2'771	2'624	2'516
N ₂ O emissions excluding N ₂ O from LULUCF	2'829	2'689	2'547	2'438
HFCs	0	245	622	1'064
PFCs	117	17	50	44
SF ₆	137	93	144	203
NF ₃	0	0	0	0
Total (including LULUCF)	53'078	48'575	57'383	52'622
Total (excluding LULUCF)	53'357	52'251	52'365	54'856

Greenhouse Gas Emissions	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Change base year to 2015 (%)
	CO ₂ equivalent (kt)										
CO ₂ emissions including net CO ₂ from LULUCF	45'928	43'123	43'594	41'745	43'713	40'049	40'913	42'040	38'346	37'771	-13.7%
CO ₂ emissions excluding net CO ₂ from LULUCF	45'380	43'388	44'708	43'538	45'053	40'993	42'259	43'202	39'269	38'751	-12.3%
CH ₄ emissions including CH ₄ from LULUCF	5'336	5'317	5'392	5'309	5'289	5'232	5'192	5'131	5'134	5'099	-16.8%
CH ₄ emissions excluding CH ₄ from LULUCF	5'321	5'301	5'379	5'295	5'276	5'217	5'179	5'118	5'121	5'085	-16.7%
N ₂ O emissions including N ₂ O from LULUCF	2'517	2'541	2'561	2'523	2'570	2'515	2'496	2'460	2'475	2'425	-16.8%
N ₂ O emissions excluding N ₂ O from LULUCF	2'437	2'461	2'484	2'448	2'496	2'440	2'423	2'385	2'400	2'352	-16.9%
HFCs	1'111	1'186	1'236	1'247	1'324	1'406	1'486	1'514	1'527	1'536	see caption
PFCs	51	49	58	63	64.50	67.72	71.27	51.93	44.03	57.21	-50.9%
SF ₆	186	172	222	180	148	160	209	252	259	256	86.7%
NF ₃	0.0	0.0	0.1	5	8	6	0.4	0.1	0.4	0.5	-
Total (including LULUCF)	55'129	52'388	53'063	51'071	53'117	49'436	50'368	51'450	47'784	47'144	-11.2%
Total (excluding LULUCF)	54'486	52'556	54'087	52'776	54'370	50'289	51'628	52'523	48'620	48'038	-10.0%

With regard to the distribution of emissions by individual greenhouse gases, CO₂ is the largest single contributor to emissions, accounting for 80.7% of total gross GHG emissions (excluding LULUCF). The shares of CH₄ and N₂O are about 10.6% and 4.9%, respectively. The shares of the three gases show slightly decreasing trends in the period 1990–2015, whereas aggregated F-gases, which contributed with only 0.5% in 1990, increased to reach 3.8% in 2015 (Table E- 2).

Table E- 2 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (kt) and shares of different GHG (%), selected years.

Greenhouse Gas Emissions (excluding LULUCF)	1990		1995		2000		2005		2010	
	kt CO ₂ eq	%								
CO ₂	44'171	82.8%	43'423	83.1%	43'607	83.3%	45'799	83.5%	45'053	82.9%
CH ₄	6'102	11.4%	5'782	11.1%	5'396	10.3%	5'307	9.7%	5'276	9.7%
N ₂ O	2'829	5.3%	2'689	5.1%	2'547	4.9%	2'438	4.4%	2'496	4.6%
HFCs	0	0.0%	245	0.5%	622	1.2%	1'064	1.9%	1'324	2.4%
PFCs	117	0.2%	17	0.0%	50	0.1%	44	0.1%	64	0.1%
SF ₆	137	0.3%	93	0.2%	144	0.3%	203	0.4%	148	0.3%
NF ₃	0	0.0%	0	0.0%	0	0.0%	0	0.0%	8	0.0%
Total (excluding LULUCF)	53'357	100%	52'251	100%	52'365	100%	54'856	100%	54'370	100%

Greenhouse Gas Emissions (excluding LULUCF)	2011		2012		2013		2014		2015	
	kt CO ₂ eq	%								
CO ₂	40'993	81.5%	42'259	81.9%	43'202	82.3%	39'269	80.8%	38'751	80.7%
CH ₄	5'217	10.4%	5'179	10.0%	5'118	9.7%	5'121	10.5%	5'085	10.6%
N ₂ O	2'440	4.9%	2'423	4.7%	2'385	4.5%	2'400	4.9%	2'352	4.9%
HFCs	1'406	2.8%	1'486	2.9%	1'514	2.9%	1'527	3.1%	1'536	3.2%
PFCs	68	0.1%	71	0.1%	52	0.1%	44	0.1%	57	0.1%
SF ₆	160	0.3%	209	0.4%	252	0.5%	259	0.5%	256	0.5%
NF ₃	6	0.0%	0.4	0.0%	0.1	0.0%	0.4	0.0%	0.5	0.0%
Total (excluding LULUCF)	50'289	100%	51'628	100%	52'523	100%	48'620	100%	48'038	100%




Figure E- 1 Contribution of individual gases to Switzerland's GHG emissions (excluding LULUCF) in 2015.

Uncertainty Analyses

Uncertainties were assessed with Approach 1 and 2 for Switzerland's GHG inventory including and excluding LULUCF categories for the years 1990 and 2015 (level) and for the period 1990–2015 (trend) (see details in chp. 1.6 and IPCC (2006)). The uncertainty results for Approach 2 are displayed in Table E- 3. When excluding LULUCF, Approach 2 level uncertainty amounts to 3.56% and trend uncertainty to 2.62%. Due to high uncertainties in sector 4 LULUCF, overall uncertainties are generally higher for the analyses including LULUCF categories (level: 6.14%, trend: 5.52%).

Table E- 3 Switzerland's relative uncertainties for national total GHG emissions excluding and including the LULUCF sector – Approach 2: Level uncertainties 2015 and trend uncertainties 1990–2015.

Approach 2 Uncertainty Analysis		
Inventory	Level uncertainty	Trend uncertainty
	2015	1990-2015
excl. LULUCF	3.56%	2.62%
incl. LULUCF	6.14%	5.52%

Recalculations

For the latest recalculated year, 2014, the total national emissions (excluding LULUCF) increased from 48'617 kt CO₂ eq (Subm. 2016) to 48'620 kt CO₂ eq (Subm. 2017). See detailed explanations of the recalculations in the sectoral chapters and the summary in chp. 10.

ES.2.2 KP-LULUCF Activities

Switzerland reports the mandatory LULUCF activities Afforestation and Deforestation (Reforestation is not occurring in Switzerland) under Article 3, paragraph 3, of the Kyoto Protocol, and Forest management as a mandatory activity under Article 3, paragraph 4, of the Kyoto Protocol. The total contribution of these activities is shown in Table E- 4 and corresponds with values of the KP reporting tables. All activities include emissions and removals of all GHG (i.e. CO₂, CH₄, N₂O) from Harvested wood products (HWP), biomass burning, drainage and N mineralization, where appropriate (see chp. 11.3).

Table E- 4 Net CO₂ eq emissions (positive sign) and removals (negative sign) for activities accounted for under Article 3, paragraph 3 and Forest management under Article 3, paragraph 4, of the Kyoto Protocol, 1990, 1995, 2000, 2005, and 2008–2015 (in kt CO₂ eq)..

Greenhouse gas source and sink activities	1990	1995	2000	2005
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)			
A. Article 3.3 activities	90.28	107.91	123.99	118.66
B. Article 3.4 Forest management	-1'554.35	-4'152.86	4'526.25	-3'113.96

Greenhouse gas source and sink activities	2008	2009	2010	2011	2012	2013	2014	2015
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)							
A. Article 3.3 activities	73.31	117.86	130.75	132.95	133.98	132.95	133.19	135.71
B. Article 3.4 Forest management	-1'693.07	-2'539.30	-2'514.69	-1'184.36	-2'520.72	-2'484.03	-1'077.73	-2'536.44

ES.3. Overview of source and sink category estimates and trends, including KP-LULUCF activities

ES.3.1 GHG inventory (Convention on Climate Change)

Table E- 5 shows the GHG emissions and removals by the main source and sink categories. Sector 1 Energy clearly dominates national emissions, accounting for more than three quarters of the total GHG emissions (excluding LULUCF), as shown in Table E- 6. Sectors 2 Industrial processes and product use (IPPU) and 3 Agriculture contribute a considerable share of GHG emissions as well, while sectors 5 Waste and 6 Other are of minor importance. LULUCF categories from sector 4 are a net GHG sink.

Overall, Switzerland's GHG emissions decreased in 2015 compared to 1990. This effect is mainly driven by decreases in the sectors Energy and Agriculture, which outweigh the increase in the sector Industrial processes and product use.

Table E- 5 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (kt) and the contribution of individual source (positive numbers) and sink (negative numbers) categories for selected years.

Source and Sink Categories	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
1. Energy	41'846	41'878	42'171	44'006
1A1 Energy industries	2'519	2'643	3'172	3'816
1A2 Manufacturing industries and construction	6'453	6'206	5'928	6'000
1A3 Transport	14'660	14'266	15'930	15'860
1A4 Other sectors	17'632	18'171	16'630	17'879
1A5 Other	220	163	151	139
1B Fugitive emissions from fuels	362	430	359	313
2. Industrial processes and product use	3'585	2'922	3'139	3'795
3. Agriculture	6'780	6'489	6'108	6'078
5. Waste	1'133	950	934	964
6. Other	12	12	13	14
Total (excluding LULUCF)	53'357	52'251	52'365	54'856
4. Land use, land-use change and forestry	-279	-3'676	5'018	-2'234
Total (including LULUCF)	53'078	48'575	57'383	52'622

Source and Sink Categories	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
	CO ₂ equivalent (kt)										%
1. Energy	43'612	41'578	42'943	41'844	43'218	39'162	40'552	41'489	37'464	37'113	-11.3%
1A1 Energy industries	4'032	3'719	3'837	3'674	3'847	3'598	3'641	3'737	3'609	3'279	30.1%
1A2 Manufacturing industries and construction	6'169	5'981	6'025	5'717	5'832	5'389	5'397	5'500	5'108	4'989	-22.7%
1A3 Transport	15'975	16'299	16'650	16'446	16'336	16'155	16'273	16'184	16'075	15'338	4.6%
1A4 Other sectors	17'002	15'163	16'025	15'607	16'785	13'610	14'850	15'697	12'303	13'151	-25.4%
1A5 Other	143	136	131	133	138	125	133	134	139	135	-38.4%
1B Fugitive emissions from fuels	291	280	275	269	282	285	260	239	230	221	-38.9%
2. Industrial processes and product use	3'785	3'848	3'924	3'811	4'022	4'064	4'066	4'096	4'140	3'992	11.3%
3. Agriculture	6'112	6'168	6'273	6'194	6'213	6'159	6'126	6'060	6'150	6'074	-10.4%
5. Waste	965	949	933	915	904	890	869	864	855	846	-25.3%
6. Other	12	14	13	13	12	13	14	14	12	12	2.0%
Total (excluding LULUCF)	54'486	52'556	54'087	52'776	54'370	50'289	51'628	52'523	48'620	48'038	-10.0%
4. Land use, land-use change and forestry	643	-169	-1'024	-1'705	-1'253	-854	-1'260	-1'073	-836	-894	220.7%
Total (including LULUCF)	55'129	52'388	53'063	51'071	53'117	49'436	50'368	51'450	47'784	47'144	-11.2%

It becomes apparent in Figure E- 2 that the GHG emission trend in the period 1990–2015 is subject to fluctuations but with a decreasing trend starting in 2005. The fluctuations emerge from the year-to-year variability of the energy sector emissions caused by changing winter temperatures (and hence, changing demand of heating). In addition, since 2005 a growing decoupling of fuel combustion emissions and winter temperature conditions is visible. That is, the emission reductions are not only caused by weather conditions, but are also the result of emission reduction measures.




Figure E- 2 Switzerland's GHG emissions in CO₂ equivalent (kt) by sectors, 1990–2015 (excluding LULUCF).

Table E- 6 gives a more detailed impression of individual sectors' contributions to total emissions for selected years (excluding LULUCF). In general, the relative contributions of the different sectors have been rather stable between 1990 and 2015. When comparing the contributions in 2015 to 1990, the following development can be observed:

- Smaller relative contributions in sectors 1 Energy and 5 Waste.
- Larger relative contributions in sector 2 Industrial processes and product use.
- Almost equal relative contribution in sectors 3 Agriculture (1990: 12.7%, 2015: 12.6%) and 6 Other (1990: 0.023%, 2015: 0.026%).

Table E- 6 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (kt) and the contribution of individual source categories for selected years.

Source and Sink Categories	1990		1995		2000		2005		2010	
	kt CO ₂ eq	%								
1. Energy	41'846	78.4%	41'878	80.1%	42'171	80.5%	44'006	80.2%	43'218	79.5%
1A1 Energy industries	2'519	4.7%	2'643	5.1%	3'172	6.1%	3'816	7.0%	3'847	7.1%
1A2 Manufacturing industries and construction	6'453	12.1%	6'206	11.9%	5'928	11.3%	6'000	10.9%	5'832	10.7%
1A3 Transport	14'660	27.5%	14'266	27.3%	15'930	30.4%	15'860	28.9%	16'336	30.0%
1A4 Other sectors	17'632	33.0%	18'171	34.8%	16'630	31.8%	17'879	32.6%	16'785	30.9%
1A5 Other	220	0.4%	163	0.3%	151	0.3%	139	0.3%	138	0.3%
1B Fugitive emissions from fuels	362	0.7%	430	0.8%	359	0.7%	313	0.6%	282	0.5%
2. Industrial processes and product use	3'585	6.7%	2'922	5.6%	3'139	6.0%	3'795	6.9%	4'022	7.4%
3. Agriculture	6'780	12.7%	6'489	12.4%	6'108	11.7%	6'078	11.1%	6'213	11.4%
5. Waste	1'133	2.1%	950	1.8%	934	1.8%	964	1.8%	904	1.7%
6. Other	12	0.0%	12	0.0%	13	0.0%	14	0.0%	12	0.0%
Total (excluding LULUCF)	53'357	100.0%	52'251	100.0%	52'365	100.0%	54'856	100.0%	54'370	100.0%

Source and Sink Categories	2011		2012		2013		2014		2015	
	kt CO ₂ eq	%								
1. Energy	39'162	77.9%	40'552	78.5%	41'489	79.0%	37'464	77.1%	37'113	77.3%
1A1 Energy industries	3'598	7.2%	3'641	7.1%	3'737	7.1%	3'609	7.4%	3'279	6.8%
1A2 Manufacturing industries and construction	5'389	10.7%	5'397	10.5%	5'500	10.5%	5'108	10.5%	4'989	10.4%
1A3 Transport	16'155	32.1%	16'273	31.5%	16'184	30.8%	16'075	33.1%	15'338	31.9%
1A4 Other sectors	13'610	27.1%	14'850	28.8%	15'697	29.9%	12'303	25.3%	13'151	27.4%
1A5 Other	125	0.2%	133	0.3%	134	0.3%	139	0.3%	135	0.3%
1B Fugitive emissions from fuels	285	0.6%	260	0.5%	239	0.5%	230	0.5%	221	0.5%
2. Industrial processes and product use	4'064	8.1%	4'066	7.9%	4'096	7.8%	4'140	8.5%	3'992	8.3%
3. Agriculture	6'159	12.2%	6'126	11.9%	6'060	11.5%	6'150	12.7%	6'074	12.6%
5. Waste	890	1.8%	869	1.7%	864	1.6%	855	1.8%	846	1.8%
6. Other	13	0.0%	14	0.0%	14	0.0%	12	0.0%	12	0.0%
Total (excluding LULUCF)	50'289	100.0%	51'628	100.0%	52'523	100.0%	48'620	100.0%	48'038	100.0%

ES.3.2 KP-LULUCF activities

An overview of net CO₂ equivalent emissions and removals of activities under Article 3, paragraph 3 and Forest management under paragraph 4 of the Kyoto Protocol is shown in Table E- 7 and in Figure E- 3.

Detailed quantitative information of the years 1990–2015 is reported in chp. 11.4, chp. 11.5, and displayed in Table 11-1. Annual changes in the emissions from Afforestation and Deforestation can directly be attributed to the changes in the area of Deforestations. The relative changes in the area of managed forest are comparatively low and fluctuations in the annual net carbon stock changes in Forest management can primarily be explained by changes in the losses from the living biomass, dead wood and litter pools (Table 11-2). The reason for the extraordinary high emissions in 2000 and the small removals in the following year 2001 for Forest management is the winter storm “Lothar” end of 1999, which caused large-scale damages in the forest stands and increased losses of living biomass due to salvage logging. Harvesting rates in Swiss forests gradually increased since 1991 until 2007, reaching peak values in 2006 and 2007 and thus also resulting in small removals from Forest management in those years. Because harvesting rates started to decline in 2008 (Table 6-16) due to the international and domestic economic framework conditions, removals from Forest management increased since 2008, still showing high year-to-year variability. Fluctuations in the HWP pool can mainly be attributed to changes in the production of sawnwood (see chp. 6.11), which is strongly linked to the domestic harvesting rate in Swiss forests.

Table E- 7 Net CO₂ eq emissions (positive sign) and removals (negative sign) of activities accounted for under Article 3, paragraph 3 (Afforestation, Deforestation) and paragraph 4 (Forest management and Harvested wood products HWP) of the Kyoto Protocol, 1990, 1995, 2000, 2005, and 2008–2015 (in kt CO₂ eq).

Greenhouse gas source and sink activities	1990	1995	2000	2005
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)			
A. Article 3.3 activities	90.28	107.91	123.99	118.66
Afforestation	-2.48	-12.20	-15.82	-19.12
Deforestation	92.76	120.11	139.81	137.78
B. Article 3.4 Forest Management	-1'554.35	-4'152.86	4'526.25	-3'113.96
Forest management excl. HWP	-323.56	-3'591.98	5'360.80	-2'350.90
HWP	-1'230.78	-560.89	-834.55	-763.06

Greenhouse gas source and sink activities	2008	2009	2010	2011	2012	2013	2014	2015
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)							
A. Article 3.3 activities	73.31	117.86	130.75	132.95	133.98	132.95	133.19	135.71
Afforestation	-21.35	-21.97	-20.61	-18.81	-18.35	-17.41	-15.31	-16.73
Deforestation	94.66	139.83	151.36	151.77	152.33	150.36	148.50	152.43
B. Article 3.4 Forest Management	-1'693.07	-2'539.30	-2'514.69	-1'184.36	-2'520.72	-2'484.03	-1'077.73	-2'536.44
Forest management excl. HWP	-1'164.90	-2'168.91	-2'103.36	-926.62	-2'352.99	-2'323.15	-971.71	-2'466.91
HWP	-528.17	-370.39	-411.33	-257.74	-167.74	-160.87	-106.02	-69.53




Figure E- 3 GHG emissions (positive sign) and removals (negative sign), 1990–2015 (in kt CO₂ eq). Shown are data for Afforestation (too small to be distinguishable) and Deforestation under Article 3, paragraph 3, Forest management (excluding HWP) and HWP under Article 3, paragraph 4.

ES.3.3 GHG inventory (Kyoto Protocol)

Relevant emissions and removals under the Kyoto Protocol are shown in Table E- 8 and Table E- 9 sorted by sectors and GHG respectively. The reported total emissions differ from those reported under the UNFCCC, because sector 6 Other as well as 4 LULUCF and international bunkers are not accounted for under the Kyoto Protocol. On the other hand, activities under Article 3, paragraph 3 (Afforestation, Reforestation and Deforestation) and Article 3, paragraph 4 (Forest management, Cropland management, Grazing land management, and Revegetation) as well as indirect CO₂ emissions are included in the tables. Under the activities of Article 3, paragraph 4, of the Kyoto Protocol, Switzerland only accounts for Forest management Base year emissions (as shown in Table E- 8 and Table E- 9), which are relevant for calculating the cap on activities under Art. 3.4 (see decision

2/CMP.7, paragraph 13) are reported in Switzerland's Second Initial Report (FOEN 2016c) and the update to the report following the review (FOEN 2016d).

Table E- 8 Summary of Switzerland's GHG emissions in CO₂ equivalent (kt) as well as emissions and removals under KP-LULUCF 2008–2015, 1990–2015 by sectors. Excluded are emissions and removals from sectors 4 LULUCF, 6 Other and from International bunkers.

Annex A sources	Sector	Base year initial report	1990	1991	1992	1993	1994	1995	1996	1997	1998
		CO ₂ equivalent (kt)									
Annex A sources	1 Energy + indirect CO ₂ from this sector	41'881	41'890	44'325	44'314	42'126	40'997	41'905	42'819	41'851	43'407
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'893	3'950	3'549	3'359	3'020	3'196	3'166	3'016	2'912	3'009
	3 Agriculture	6'804	6'780	6'744	6'629	6'527	6'513	6'489	6'446	6'261	6'218
	5 Waste + indirect CO ₂ from this sector	1'142	1'135	1'050	1'053	1'005	948	951	947	938	925
	Total (Annex A sources)	53'719	53'755	55'668	55'355	52'678	51'654	52'512	53'229	51'962	53'558

Annex A sources	Sector	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
		CO ₂ equivalent (kt)									
Annex A sources	1 Energy + indirect CO ₂ from this sector	43'189	42'188	43'604	41'993	43'214	43'575	44'019	43'626	41'590	42'956
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'063	3'307	3'396	3'419	3'497	3'777	3'909	3'897	3'956	4'033
	3 Agriculture	6'124	6'108	6'170	6'136	6'059	6'036	6'078	6'112	6'168	6'273
	5 Waste + indirect CO ₂ from this sector	914	936	951	975	954	973	965	966	951	934
	Total (Annex A sources)	53'289	52'539	54'121	52'522	53'724	54'362	54'971	54'600	52'665	54'197

KP-LULUCF	Sector	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
		CO ₂ equivalent (kt)							
KP-LULUCF	Art.3.3 Afforestation & reforestation	-22	-21	-19	-18	-17	-15	-17	
	Art.3.3 Deforestation	140	151	152	152	150	149	152.4	
	Art.3.4 Forest management	-2'539	-2'515	-1'184	-2'521	-2'484	-1'078	-2'536	
	Art.3.4 Cropland management	NA	NA	NA	NA	NA	NA	NA	
	Art.3.4 Grazing land management	NA	NA	NA	NA	NA	NA	NA	
	Revegetation	NA	NA	NA	NA	NA	NA	NA	

Annex A sources	Sector	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
		CO ₂ equivalent (kt)							
Annex A sources	1 Energy + indirect CO ₂ from this sector	41'856	43'229	39'173	40'562	41'500	37'475	37'118	-11%
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'920	4'132	4'173	4'175	4'202	4'246	4'098	4%
	3 Agriculture	6'194	6'213	6'159	6'126	6'060	6'150	6'074	-10%
	5 Waste + indirect CO ₂ from this sector	916	905	892	870	865	856	848	-25%
	Total (Annex A sources)	52'886	54'480	50'397	51'733	52'628	48'727	48'138	-10%

Table E- 9 Switzerland's total GHG emissions (excluding 4 LULUCF, 6 Other and International bunkers) and the contribution of individual gases in CO₂ equivalent (kt), 1990–2015 as well as emissions and removals under KP-LULUCF 2008–2015.

Annex A sources	GHG	Base year initial report	1990	1991	1992	1993	1994	1995	1996	1997	1998
		CO ₂ equivalent (kt)									
		CO ₂ + indirect CO ₂	44'521	44'571	46'569	46'370	43'917	42'953	43'686	44'394	43'266
CH ₄	6'091	6'102	6'035	5'953	5'843	5'791	5'782	5'743	5'601	5'531	
N ₂ O	2'853	2'829	2'824	2'794	2'730	2'702	2'688	2'685	2'591	2'587	
HFCs	0.0	0.0	1.6	16	33	81	245	296	360	454	
PFCs	117	117	99	81	35	21	17	20	21	24	
SF ₆	137	137	139	141	121	107	93	90	124	153	
NF ₃	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Total (Annex A sources)	53'719	53'755	55'668	55'355	52'678	51'654	52'512	53'229	51'962	53'558	

Annex A sources	GHG	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
		CO ₂ equivalent (kt)									
		CO ₂ + indirect CO ₂	44'616	43'782	45'234	43'596	44'794	45'360	45'915	45'496	43'498
CH ₄	5'429	5'395	5'435	5'400	5'322	5'290	5'307	5'320	5'300	5'378	
N ₂ O	2'551	2'546	2'561	2'537	2'489	2'446	2'438	2'436	2'460	2'484	
HFCs	528	622	720	798	893	1'014	1'064	1'111	1'186	1'236	
PFCs	26	50	28	33	62	65	44	51	49	58	
SF ₆	140	144	145	158	165	186	203	186	172	222	
NF ₃	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	
Total (Annex A sources)	53'289	52'539	54'121	52'522	53'724	54'362	54'971	54'600	52'665	54'197	
KP-LULUCF	Art.3.3	CO ₂									70
		CH ₄									NO
		N ₂ O									2.2
KP-LULUCF	Art.3.4	CO ₂									-1'702
		CH ₄									2.7
		N ₂ O									6.1

Annex A sources	GHG	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990	
		CO ₂ equivalent (kt)								%
		CO ₂ + indirect CO ₂	43'649	45'164	41'102	42'366	43'308	39'377	38'852	-13%
CH ₄	5'295	5'276	5'217	5'179	5'118	5'120	5'085	-17%		
N ₂ O	2'448	2'495	2'439	2'422	2'384	2'400	2'351	-17%		
HFCs	1'247	1'324	1'406	1'486	1'514	1'527	1'536	see caption		
PFCs	63	64	68	71	52	44	57	-51%		
SF ₆	180	148	160	209	252	259	256	87%		
NF ₃	5.1	8.5	6.2	0.4	0.1	0.4	0.5	NA		
Total (Annex A sources)	52'886	54'480	50'397	51'733	52'628	48'727	48'138	-10%		
KP-LULUCF	Art.3.3	CO ₂	114	127	130	131	130	133		
		CH ₄	NO	NO	NO	NO	NO	NO		
		N ₂ O	2.4	2.4	2.4	2.4	2.4	2.4	2.4	
KP-LULUCF	Art.3.4	CO ₂	-2'548	-2'523	-1'196	-2'529	-2'492	-1'086	-2'545	
		CH ₄	2.7	2.3	4.1	2.3	2.3	2.5	2.5	
		N ₂ O	6.2	6.0	7.1	6.0	6.0	6.2	6.2	

ES.4. Other information

Emission trends for precursor gases show a very pronounced decline (see Table 2-6 and Figure 2-9). A strict air pollution control policy led to strong decreases of slightly 56% (NO_x) up to 83% (SO₂) in the period 1990–2015 in emissions of precursor gases and SO₂. The main reduction measures were abatement of exhaust emissions from road vehicles and stationary combustion equipment, taxation of solvents and sulphured fuels, and voluntary agreements with industry sectors (FOEN 2010i, Swiss Confederation 1985, 1997).

Acknowledgements

The GHG inventory preparation is a joint effort which is based on input from many federal agencies, institutions, associations, companies and individuals. Their effort was essential for the successful completion of the present inventory report.

The Federal Office for the Environment would like to acknowledge the valuable support it has received from the many contributors to this document. In particular, it would like to thank all the data suppliers, including the Office of the Environment of the Principality of Liechtenstein for providing its fossil fuel consumption data, as well as experts, authors and both national and international reviewers.

PART 1

1 Introduction

1.1 Background information on Swiss greenhouse gas inventories, climate change and supplementary information of the Kyoto Protocol (KP)

1.1.1 Information on climate change

The report of the Swiss Advisory Body on Climate Change (OcCC) provides an assessment of the observed and expected impacts of climate change on Switzerland and the vulnerability of various ecological and socio-economic systems (OcCC 2008). In the course of the 21st century, Swiss climate is projected to depart significantly from present and past conditions (CH2011 2011). Recent data confirms a warming trend with an observed increase in mean annual temperature of 1.75°C between 1864 and 2012 (FOEN 2014d). Over the last 100 years, mean annual temperatures increased by 0.13–0.20°C per decade, with a substantially accelerated warming in recent decades. According to the non-intervention scenarios (A2, A1B), seasonal mean temperatures will rise by another 2.7–4.8°C by the end of this century compared to the period 1980–2009. Under the climate stabilization scenario (RCP3PD), Swiss climate would still change over the next decades, but is projected to stabilize at a mean warming of 1.2–1.8°C compared to the period 1980–2009 (FOEN 2014d). In 2016 a comprehensive overview on Swiss climate – basics, consequences and perspectives – has been published by the Swiss Academies of Sciences (SCNAT 2016).

The most visible change in the Alps resulting from global warming is the retreat of glaciers, which showed a volume loss of 12% since 1999 (FOEN 2014d). The area covered by alpine glaciers continuously diminishes. From about 2'900 km² of Alpine glacier area in the mid-1970s, only 2'100 km² remained in 2003 and an estimated 1'900 km² in 2013. A dramatic future loss in glacier covered area of 50–90% by 2100 has recently been modelled for a temperature increase of 2–6°C for Switzerland (FOEN 2014d).

The observed trends in precipitation are less distinct than in temperature. Compared to the last 30 years, and depending on the scenario considered, the best estimates of summer mean precipitation for all Swiss regions is projected to decrease by 8%–28% over the 21st century. Uncertainties due to climate model imperfections and natural variability typically amount to 15% in precipitation (CH2011 2011). The change in summer mean precipitation will have a marked impact on the hydrological cycle: on the Central Plateau and in the very south of Switzerland, small and medium watercourses will dry up more frequently and natural replenishment of groundwater will decrease accordingly. Apart from changes to the mean temperature and precipitation, the nature of extreme events is also expected to change (CH2011 2011). More frequent, intense and longer-lasting summer warm spells and heat waves are expected, while the number of cold winter days and nights decrease in the projections for future climate in Switzerland. This is particularly relevant for alpine areas, tourism and forestry due to the risk of more frequent floods, landslides and debris flows.

The warming trend and changing precipitation patterns are expected to have significant effects on ecosystems. The Biodiversity Monitoring Switzerland reports that impacts of climate change are already being observed. A report about climate change in Switzerland

summarizes several climate change affected indicators such as the phenological spring phases, flowering indices and animal specific indices (FOEN 2014d). The indicators show significant changes in a wide range of ecosystems during the last decades. The report also emphasizes that typical alpine vascular plants have shifted uphill over the past century. Generally, climate change is expected to affect species composition, distribution, their cycles, synchronicity, the overall genetic diversity and the provision of ecosystem services. It will raise the vulnerability of forests and impair their protective, productive and social functions. For agriculture, a moderate warming of 2–3°C might increase productivity, however, if temperatures rise beyond that level, the increase in heat waves and drought periods would prove problematic for the cultivation of land and for livestock husbandry.

Various sectors of the Swiss economy are likely to be affected by progressing climate change. In particular, the tourism industry will be hit, as the potentially beneficial effects for summer tourism will not compensate for the loss of income in mountain resorts during winter due to scarcity of snow. Cable car stations may lose their stability due to instabilities of permafrost soils. Hydroelectric power stations may be affected by altered runoff and sediment transport regimes, and insurance companies may face increased losses due to winter storms and floods. Natural hazards and extreme weather events potentially pose a growing risk to infrastructure and human health. Heat waves and elevated tropospheric ozone levels are cause for serious concern, as evidenced by the impacts of the heat wave in 2003. Finally, it remains to be seen to what extent vector borne diseases spread due to changing climatic conditions. Switzerland has recently analysed these challenges in detail and developed an effective adaptation strategy in order to hedge against negative effects resulting from climate change in Switzerland (FOEN 2012b).

1.1.2 Information on the greenhouse gas inventory

On 10 December 1993, Switzerland ratified the United Nations Framework Convention on Climate Change (UNFCCC). Since 1996, the submission of its national greenhouse gas inventory has been based on IPCC guidelines. From 1998 onwards, the inventories have been submitted in the Common Reporting Format (CRF). In 2004, Switzerland started submitting an annual National Inventory Report (NIR) under the UNFCCC.

On 9 July 2003, Switzerland ratified the Kyoto Protocol under the UNFCCC. In November 2006 Switzerland submitted its Initial Report under Article 7, paragraph 4 of the Kyoto Protocol (FOEN 2006h). The Swiss National Inventory System (NIS) according to Article 5.1 of the Kyoto Protocol has been implemented in 2006 and is fully operational. On 6 December 2007, the NIS quality management system was certified to comply with ISO 9001:2000 requirements (SQS 2008); it has been audited and recertified several times with the latest audit on 28th November 2016 (ISO 9001:2008, Swiss-TS 2016). The quality management system includes the accounting and reporting of the National Registry as well. The April 2008 submission of the Swiss GHG inventory (FOEN 2008) has been Switzerland's first submission under both the UNFCCC and the Kyoto Protocol.

On 28 August 2015, Switzerland submitted its instrument of acceptance of the Doha amendment to the Kyoto Protocol to the United Nations Framework Convention on Climate Change. The Initial Report for the second commitment period (FOEN 2016c) was submitted simultaneously with the inventory 2016. An update following the in-country review by an expert review team was submitted on 7th November 2016 to the UNFCCC secretariat (FOEN 2016d).

The 2015 inventory submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol was restructured in accordance with the Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention (UNFCCC 2014a) and the Guidance for reporting information on activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (UNFCCC 2014b). The submission could not be completed in 2015 due to the problems with the UN software (CRF Reporter). The first part of the submission included the NIR submitted on 27 April 2015 and the standard electronic format (SEF) tables (CP2 on 27 April 2015, CP1 on 27 May 2015). The second part consisting of the reporting tables (CRF tables and Kyoto Protocol LULUCF tables) is submitted in April 2016.

The 2017 inventory submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol includes the NIR on hand, the Greenhouse gas inventory 1990 to 2015, the Kyoto Protocol LULUCF tables 2008 to 2015 in the common reporting format and the SEF tables as well as the standard independent assessment report (SIAR) from the National Registry.

1.1.3 Supplementary information required under art. 7.1. KP

Information on KP-LULUCF is provided in chp. 11 of PART 2.

Switzerland only accounts for the mandatory activity Forest management under Article 3, paragraph 4 of the Kyoto Protocol (FOEN 2016c). In accordance with Annex I to Decision 2/CMP.7 (Annex I, Para 13), credits from Forest management are capped in the second commitment period. Thus, for Switzerland the cap amounts to 3.5% of the 1990 emissions (excluding LULUCF).

Switzerland has chosen to account over the entire commitment period for emissions and removals from activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (FOEN 2016c, FOEN 2016d). In addition to the mandatory submission of the inventory years 2013–2015, data for the years 1999–2012 are available and shown in Switzerland's NIR.

1.2 National inventory arrangements

1.2.1 Institutional, legal and procedural arrangements

Based on the Organisation Ordinance for the Federal Department of the Environment, Transport, Energy and Communications (DETEC), the Federal Office for the Environment (FOEN) is the designated national authority for climate policy and environmental monitoring. According to the decree of the Federal Council of 8 November 2006, the FOEN is in charge of the National Inventory System (NIS) (Figure 1-1). The Swiss National Inventory System was formally set up in 2006 in compliance with the requirements of the UNFCCC and the Kyoto Protocol (FOEN 2006h). In this context, the FOEN established the process "Climate Reporting", which covers maintaining the National Inventory System and fulfilling all reporting obligations under the UNFCCC and the Kyoto Protocol. The process, led and managed by the Climate division of the FOEN, is fully operational ever since and ensures timely fulfilment of Switzerland's reporting obligations.




Figure 1-1 Institutional arrangements of the National Inventory System. Colours refer to divisions at FOEN. yellow: Climate division, red: Air Pollution Control and Chemicals division, beige: Forest division.

Legal arrangements

The CO₂ act (Swiss Confederation 2011) and the CO₂ ordinance (Swiss Confederation 2012) are the main legal instruments regarding climate policies. They also define the implementing bodies and, for all measures that are regulated at the national level, sanctions for non-compliance to climate policies and measures. The FOEN plays a central role in the development, evaluation and implementation of policies and measures.

With regard to statistical investigations, the legal basis is laid down in the Federal Statistics Act (Swiss Confederation 1992a) and the corresponding Ordinance on the Conduct of Federal Statistical Surveys (Swiss Confederation 1993). The greenhouse gas inventory, the institution responsible for it and the institutions contributing to it are explicitly listed in the ordinance.

Institutional arrangements

There are well-established agreements and long-standing collaborations with institutions of the federal administration and private entities (Table 1-1) that guarantee the continuity of the National Inventory System (Figure 1-1). While agreements with institutions of the federal administration are normally open-ended, several large contracts with private entities are on a

four-year basis, with an option for renewal for another four-year term. This enables continuous collaboration and ensures the technical competence and experience of the staff involved.

The overall responsibility for the greenhouse gas inventory lies with the Climate division of the FOEN. The Air Pollution Control and Chemicals division of the FOEN maintains and updates the emissions database (greenhouse gases and air pollutants) in very close collaboration with the Climate division. The national energy statistics from the Federal Office of Energy (SFOE) provides the basis for the Energy sector. The Federal Office for Civil Aviation (FOCA) delivers the domestic and international aviation emissions. A consultancy (Carbotech) is mandated to survey and model fluorinated gases use and emissions and to provide an annual update thereof. Agriculture emissions are compiled by the federal research institute Agroscope. For LULUCF, detailed area survey data are provided by the Swiss Federal Statistical Office (SFSO). Two consultancies (Sigmoplan/Meteotest) are mandated to process the area survey data to derive land-use and land-use change data and related emissions. The Swiss Federal Institute for Forest, Snow and Landscape Research (WSL) is in charge of the national forest inventory and forestry-related modelling, providing the relevant input for the Forest division of the FOEN, who is compiling forestry emissions and removals. The LULUCF sector is coordinated by a member of the Climate division of the FOEN. A collaboration between two consultancies (INFRAS/CSD) is mandated to update the National Inventory Report (NIR).

Single national entity with overall responsibility for the inventory:

Federal Office for the Environment (FOEN)
 Climate Division, Climate Reporting and Adaptation Section
 Dr. Regine Röthlisberger, process manager
 CH-3003 Bern, Switzerland
 +41 58 462 92 59
climate@bafu.admin.ch
www.climatereporting.ch

Table 1-1 Overview of the institutional arrangements and tasks

Institutions of the federal administration	
FOEN Climate division	Overall responsibility for the greenhouse gas inventory
FOEN Air Pollution Control and Chemicals division	EMIS data base and data archiving
FOEN Forest division	Forestry emissions and removals
Swiss Federal Office of Energy (SFOE)	Energy statistics
Federal Office of Civil Aviation (FOCA)	Aviation emissions
Swiss Federal Statistical Office (SFSO)	Area surveys for (KP-) LULUCF
Swiss Federal Institute for Forest, Snow and Landscape Research WSL	National forest inventory, forestry related modelling
Agroscope Federal Research Institute	Agriculture emissions and removals
Private entities	
Carbotech	Fluorinated gases emissions
Sigmoplan/Meteotest	(KP-) LULUCF
Infras/CSD	NIR editing, uncertainty analyses

1.2.2 Overview of inventory planning, preparation and management

The process of inventory planning, preparation and management in Switzerland is well-established. Responsibilities and decision-making power are assigned to specific people or groups of people (Figure 1-1). The management responsibility for the NIS lies with the **National Inventory System Supervisory Board (NISSB)**. The board consists of a member of the FOEN directorate and FOEN division heads of the relevant divisions (Climate, Forest, Air Pollution Control and Chemicals, International Affairs). At the operational level, the process of planning, preparation and management of the greenhouse gas inventory is led by the **process manager**. The **QA/QC officer** oversees design, development, and operation of the quality management system and is the primary contact point during the UN review process. The **GHG inventory core group** is the committee that combines all technical expertise required for greenhouse gas inventory planning, preparation and management. It consists of the process manager, the QA/QC officer, the inventory compiler, sectoral experts, as well as the NIR compiler. Additional experts join the core group as required. The GHG inventory core group ensures conformity of the inventory with the relevant UNFCCC reporting guidelines (UNFCCC 2014a), timely inventory preparation, and consideration and approval of methodological changes, choice of data and recalculations. The **GHG inventory working group** encompasses all technical personnel involved in the inventory preparation process or representing institutions that play a significant role as suppliers of data.

Inventory planning, preparation, and management follow an annual cycle according to a plan-do-check-act cycle (Table 1-2). Planning of the inventory cycle starts with the first meeting of the GHG inventory core group in May, where work is scheduled, priorities with regard to inventory development are set and decisions regarding planned improvements are taken. Data compilation usually starts in June with the first data sets for the preceding year becoming available. Quality control activities form part of the data acquisition process. They are routinely carried out by the EMIS (Swiss Emission Information System) experts and the sectoral experts. The UN review process in September provides further input to the inventory development plan (IDP). Recommendations and suggestions are discussed in the core group and future work is prioritized. The supervisory board (NISSB) is provided with the management review in October and asked for formal approval of the planned way of proceeding. An important stage in inventory preparation is the preparation and quality control of the reporting tables (CRF) in December and January and the key category and uncertainty analyses towards end of January. The editing of the National Inventory Report (NIR) progresses alongside data compilation, with a draft of the NIR going into internal review in March. Suggestions from the internal review are dealt with before submission as far as possible. If the internal review suggests large revisions, they are taken up in the IDP for future improvements. The inventory is presented to the NISSB for official consideration and approval around end of March. Submission is coordinated by the process manager and carried out by the national inventory compiler. Archiving of inventory material is performed after submission by the EMIS and sectoral experts, by the contributing authors and by the QA/QC officer.

Table 1-2 Annual cycle of inventory planning, preparation and management

	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr
Data compilation												
QC EMIS Experts												
QC Sectoral Experts												
UN Review												
Inventory Development Plan												
CRF Tables												
QC CRF Tables												
KCA / Uncertainties												
NIR												
Internal review NIR and CRF Tables												
Official consideration and approval												
Submission												x
Archiving												
Meeting of Core Group	x				x			x		x		
Meeting of Working Group											x	
Meeting of NIS Supervisory Board						x					x	

1.2.3 Quality assurance, quality control and verification plan

The national inventory system has an established quality management system (QMS) that complies with the requirements of ISO 9001:2008. Certification has been obtained in 2007 and is upheld since through annual audits (Swiss TS 2016). The QMS is designed to comply with the UNFCCC reporting guidelines (UNFCCC 2014a) to ensure and continuously improve transparency, consistency, comparability, completeness, accuracy, and confidence in national GHG emission and removal estimates. The quality manual (FOEN, 2017a) – as required by ISO 9001:2008 – contains all relevant information regarding the QMS. It is updated annually and available to all members of the GHG inventory core group.

General QC procedures

The general QC activities as described in Table 6.1 of the IPCC reporting guidelines (IPCC 2006) are implemented in the annual cycle of inventory compilation (Table 1-2). Routine annual quality control procedures comprise checks related to new data and database operations, spot-checks for transcription errors, correct use of conversion factors and units, and correct calculations. There are checklists for the most important sectoral data suppliers and EMIS database experts.

Integrity of the database is ensured by creating a new database for every single submission and comparing the results from the new database with those from the previous version. Consistency of data between categories is to a large extent ensured by the design of the database, where specific emission factors and activity data that apply to various categories are used jointly by all categories to calculate emissions.

Checks regarding the correct aggregation are done on initial set-up of the various aggregations. There are also automated checks implemented in the database in order to identify incorrect internal aggregation processes.

Recalculations are compiled in a document and made available to the data compilers and the members of the GHG inventory core group including the NIR authors. The recalculations file

is of great importance in the QC procedures regarding the reporting tables (CRF) and in the preparation of the NIR. QC procedures regarding the reporting tables (CRF) comprise a detailed comparison of the reporting tables (CRF) of the previous submission with those of the current submission for the base year and the latest common year. In addition, the time-series consistency is incrementally checked by comparing the latest inventory year with the preceding year. Any exceptional deviations are investigated by the sectoral or the EMIS database experts. These checks are performed in a multi-step process, first by collaborators of the Climate division of FOEN and the EMIS database experts. Then, after the required changes were implemented by the EMIS experts, also by the NIR authors.

The NIR is subject to an internal review prior to submission. The review of every section is carried out by personnel not involved in the preparation of the reviewed section, but who is familiar with the reporting under the UNFCCC. Archiving of the database and related internal documentation is carried-out by the inventory compiler, while any other material is archived on the internal data management system by the QA/QC officer. Publicly available material is published after submission on the website owned by the FOEN (www.climatereporting.ch).

Category-specific QC procedures

Whenever new emission factors are considered, they are compared to the IPCC default values and to the values used in previous years. If the values are based on better or more appropriate data and compare reasonably well with the IPCC default values (or if differences can be explained), the new values are presented to the core group for adoption in future inventories. Similarly, if new activity data have become available for a particular category, a comparison between existing and new activity data is made and if the new data provide a more consistent or more reliable basis for the inventory, they are again presented to the core group for inclusion in future inventories. Quite often, sectoral and/or EMIS experts commission research to look into a particular topic in more detail. Results from these mid- to long-term projects are presented to the inventory core group. The core group decides on how to best implement the results and documents the agreed procedure in the inventory development plan. The general procedures regarding category-specific QC is also described in the quality manual (FOEN, 2017a), while specific activities are documented in the corresponding sectoral chapters.

Quality assurance procedures

As required by ISO 9001 there are periodic internal audits covering all processes. In addition, an external organisation is mandated to do the annual audit of the ISO 9001 quality management system.

Apart from these audits, there are expert peer reviews for specific sectors commissioned on a case-by-case basis. The results and suggestions for improvements from these reviews are discussed in the core group and specific tasks for future implementation are taken up into the inventory development plan. The most recent expert peer reviews covered the Industrial Processes (CSD 2013), LULUCF (VTI 2011) and Waste sector (Rytec 2010).

Likewise, recommendations and encouragements from the UNFCCC expert review teams (ERT) are also added to the inventory development plan, discussed in the core group and implementation in future submissions planned. Specific actions resulting from suggestions from the ERT are listed in chp. 10 Recalculations.

Verification activities

In the energy sector, the standard verification activity carried out on an annual basis is the reference approach, as documented in chp. 3.2.1 of the NIR and CRF Table 1.A(b).

In addition, the FOEN supports a long-term monitoring programme carried out by the Swiss Federal Laboratories for Materials Science and Technology (EMPA). In the frame of this programme continuous measurements of atmospheric concentrations of various halogenated gases are made at the high-Alpine research station Jungfraujoch (3580 m asl), from which Swiss emissions of some fluorinated greenhouse gases can be estimated. These data are compared with the emissions reported in the greenhouse gas inventory. The results are briefly summarized in Annex 5.1.

Furthermore, a research project is developing an independent estimate of CH₄ emissions in Switzerland based on atmospheric CH₄ measurements and inverse modelling of atmospheric transport. The initial research project showed very promising results leading to a follow-up project which is currently on-going. The first results show a very good agreement between modelled emissions and emission estimates according to the greenhouse gas inventory. A summary is provided in Annex 5.2.

Treatment of Confidentiality Issues

Nearly all of the data necessary to compile the Swiss GHG inventory are publicly available. There are, however, a few exceptions:

- (i) Emission data that refer to a single enterprise are in general confidential.
- (ii) The reporting of disaggregated emissions from F-gases is confidential (not confidential as aggregated data).
- (iii) In the civil aviation sub-sector one data source (FOCA 1991) has been marked confidential by the Federal Office of Civil Aviation (FOCA).
- (iv) Unpublished AREA land use statistics raw data have been temporarily classified confidential by the Swiss Federal Statistical Office (SFSO).

The FOEN collects the data needed for calculating emissions of HFCs, PFCs, NF₃ and SF₆ from private companies or industry associations. In the National Inventory Report, the activity data underlying emission estimates of HFCs, PFCs, NF₃ and SF₆ are only partly presented at the most disaggregated level for reasons of confidentiality. However, complete emissions are reported in aggregated tables.

Confidential data will be made available by the FOEN in line with the procedures agreed under the UNFCCC for the technical review of GHG inventories (UNFCCC 2015).

Public access to the Swiss Greenhouse Gas Inventory

FOEN operates a website (www.climatereporting.ch) where the Swiss GHG inventories (NIR, reporting tables, UNFCCC review reports), the Swiss National Communications and other reports submitted to the UNFCCC and the Kyoto Protocol may be downloaded. On this website, most papers, internal reports, domestic reviews, and other difficult-to-access

materials ('grey literature') quoted in the Swiss GHG inventory are provided online. The climate reporting homepage thus provides the option for public review.

1.2.4 Changes in the national inventory arrangements since previous submission

Changes to institutional, legal and procedural arrangements (24/CP.19, 22. (a)):

There are no changes to arrangements with other institutions of the federal administration. The agreements regarding responsibilities and deliverables are maintained.

For the second commitment period, the contracts with private companies were put out for tender. The tenders were won by the contractors that have already collaborated in inventory preparation during the first commitment period, except for the editing of the NIR, where the previous contractor teamed up with a new partner (see Table 1-1). The contracts are running for four years with an option to extend by another four years in order to guarantee continuity in inventory preparation. The contractor responsible for the fluorinated gases emissions is now also a member of the GHG inventory core group.

Furthermore, the National Inventory System Supervisory Board (NISSB), which originally not only covered the National Inventory System but also the National Registry, has been formally split in two separate boards with separate mandates and responsibilities in 2014. The NISSB is now overseeing all aspects related to reporting obligations under the UNFCCC (including reporting of the National Registry in the NIR). The Emission Registry Supervisory Board ERSB on the other hand deals with management issues related to the National Registry.

Changes in staff and capacity (24/CP.19, 22. (b)):

The inventory process manager retired on 30. June 2016. The former deputy process manager, Regine Röthlisberger, has taken over as of 1. July 2016. Since 1 September 2016, Michael Bock has taken over the role of QA/QC officer from Regine Röthlisberger.

Changes to national entity with overall responsibility for the inventory (24/CP.19, 22. (c)):

No changes.

Changes to the process of inventory planning (24/CP.19, 22.(d,e)/23./24.):

No changes.

Changes to the process of inventory preparation (24/CP.19, 25./26.):

No changes.

Changes to the process of inventory management (24/CP.19, 27.):

No changes.

1.3 Inventory preparation and data collection, processing, and storage

An overview over the inventory preparation is given above and is schematically shown in Figure 1-1. Each sector has an assigned sectoral expert who is responsible for conformity with the relevant reporting guidelines, selection of appropriate methods and data sources, and collection, processing and updating of data (Figure 1-2).

For the sectors Energy, IPPU (excl. fluorinated gases) and Waste, data collection and processing is done by the Air Pollution Control and Chemicals division of the FOEN. Emissions of road and non-road transportation are provided by INFRAS, a consultancy mandated by Traffic Section of FOEN. The use of fluorinated gases and related emissions in the corresponding source categories of the IPPU sector are provided by Carbotech, a consultancy mandated by FOEN to collect and process relevant data. For Agriculture, data collection and processing is provided by Agroscope, the Federal Research Institute for Agriculture. Land-use and land use change data from the Swiss Federal Statistical Office is compiled by Meteotest/SigmaPlan, in close collaboration with the Forest division of the FOEN. The Swiss Federal Institute for Forest, Snow and Landscape Research WSL provides further input, which is processed by the Forest Division. Emission and removal estimates from forest land are calculated by the Forest division of the FOEN.

All people responsible for data collection and processing in a particular sector are preparing their data for import into the National Air Pollution Database EMIS, which compiles all inventory data, including activity data and emission factors. EMIS was originally established in the late 1980s in order to record and monitor emissions of air pollutants, but it has since been extended to cover greenhouse gases and additional emission sources. The original EMIS database underwent a full redesign and a migration to a new software platform in 2005/2006. In preparation for the submission in 2015, all processes relevant to the GHG inventory have been restructured according to the 2006 IPCC Guidelines (IPCC 2006) and the revised reporting tables (CRF).

The EMIS database as well as background information on activity data and emission factors are archived by the national inventory compiler for each submission. In the sectors where data collection is made by EMIS experts (e.g. Energy, IPPU, Waste), additional background information is compiled as appropriate (e.g. interim worksheets; references; rationale for choice of methods, data sources, activity data, emission factors). Whenever such documents are cited, they are labelled as "EMIS 2017/NFR-Code" in this report.




Figure 1-2 Schematic overview: Data collection and processing, compilation in EMIS database, export to CRF Reporter and National Inventory Report (NIR). Abbreviations: see glossary.

1.4 Methodologies and data sources

According to the revised reporting guidelines under the UNFCCC (UNFCCC 2014a) and the Kyoto Protocol (UNFCCC 2014b), emissions are calculated based on standard methods and procedures provided in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006), the 2013 KP supplement (IPCC 2014), and the 2013 wetlands supplement (IPCC 2014a). All key categories are estimated using Approach 2 or higher or country-specific methods. The methodological tier used is described in detail in the sectoral chapters of the NIR and compiled in the CRF reporting table Summary 3.

Various data suppliers contribute to the greenhouse gas inventory (Table 1-3). While most data stem from official statistics either from the FOEN or from other federal offices, some data is drawn from national associations or consultancies that maintain well-established models or data-bases. Details on activity data and emission factors are provided in the sectoral chapters of the NIR.

Table 1-3 Primary data providers for the various inventory categories. Generally, statistics are updated annually. However, the on-road and non-road emission models of INFRAS, the complete area survey by the SFSO as well as the national forest inventory by the WSL require large efforts and are therefore updated every couple of years. Coloured boxes mark those sectors to which each data provider contributes. Abbreviations: see glossary.

Institution	Subject	Inventory category (numbering according to reporting tables)											
		1A1	1A2	1A3	1A4	1A5	1B	2	3	4 / KP	5	6	indir. CO ₂ / N ₂ O
FOEN, Air Pollution Control and Chemicals division	EMIS database												
FOEN, Climate division	monitoring data due to ordinance of reduction of CO ₂ (Confederation 2012)												
FOEN, Waste division	Waste statistics												
INFRAS	Road transportation emission model												
INFRAS	Non-road emission model												
SFOE	Swiss overall energy statistics												
SFOE	Swiss statistics of renewable energies												
SFOE	Swiss wood energy statistics												
SFOE	Energy consumption in industry (helbling statistics)												
FOCA	Civil aviation												
Swiss Air Force Administration (DDPS)	Military aviation												
SGWA	Gas distribution losses												
Carbotech	F-gases, post-combustion of NMVOC												
Swissmem	National SF ₆ balance												
SFSO	Agriculture, LULUCF												
Agroscope	Agriculture, LULUCF												
SBV	Agriculture												
FOEN, Forest division	Forest statistics												
WSL	National Forest Inventory												
SigmaPlan, Meteotest	LULUCF												

1.5 Description of key categories

The aim of the key category analysis (KCA) is to identify relevant categories that have a strong influence on Switzerland's GHG inventory in terms of absolute emission and removal levels, trends and uncertainties (IPCC 2006, chp. 4). Data collection as well as quality assurance and control are prioritised for key categories during the inventory resource allocation.

1.5.1 GHG inventory

1.5.1.1 Methodology

The key category analysis is performed according to the 2006 IPCC Guidelines (IPCC 2006, chp. 4) and Decision 24/CP.19 (UNFCCC 2014a, Annex 1, Para. 39) for 1990 and the latest reported year (2015) including all GHG (CO₂, CH₄, N₂O, HFCs, PFCs, SF₆ and NF₃). A total of 167 categories are used to disaggregate Switzerland's total GHG emissions for the purpose of this key category analysis. The disaggregation level of the categories is selected based on country-specific relevance, i.e. the most important sources in Switzerland are disaggregated on a more detailed level.

Both, Approach 1 (with a proposed threshold of 95%) and Approach 2 (with a proposed threshold of 90%) level and trend assessments are applied, including emissions from sector 4 LULUCF. Indirect N₂O emissions are not considered for the key category analysis. However, indirect CO₂ emissions are considered for the key category analysis in this submission.

Uncertainty data for key category analysis Approach 2 stems from the uncertainty analysis Approach 1 (see chp. 1.6.1.2) and therefore does not incorporate correlations.

1.5.1.2 KCA (including LULUCF categories)

Approach 1

For 2015, among the total of 167 categories, 31 are identified as **level key categories** under Approach 1 (Table 1-4).

Fifteen of the key categories belong to sector 1 Energy, accounting for the largest share of CO₂ equivalent emissions in 2015. The other key categories are more or less equally distributed between sectors 2 Industrial processes and product use, 3 Agriculture, 4 LULUCF and 5 Waste.

There are three major key sources, each contributing more than 10% to the level assessment, all being part of sector 1 Energy:

- 1A3b Fuel Combustion, Road Transportation, Gasoline, CO₂
- 1A3b Fuel Combustion, Road Transportation, Diesel, CO₂
- 1A4b Fuel Combustion, Other Sectors, Residential, Liquid Fuels, CO₂

Within the ten most relevant key categories (level contribution), only 3A Enteric fermentation and 4A1 Forest land remaining forest land are not part of sector 1 Energy.

Table 1-4 Switzerland's Approach 1 level key categories for the year 2015 including LULUCF categories, sorted by emission contribution to the national total.

APPROACH 1 LEVEL ASSESSMENT FOR 2015						
A	B	C	D	E	F	G
Code	IPCC Category	GHG	Ex,t (kt CO ₂ eq)	Ex,t (kt CO ₂ eq)	Lx,t	Cumulative Total
1A3b	Road Transportation: Gasoline	CO ₂	7669	7669	14.4%	14.4%
1A3b	Road Transportation: Diesel	CO ₂	7182	7182	13.5%	27.9%
1A4b	Residential: Liquid Fuels	CO ₂	5864	5864	11.0%	38.9%
3A	Enteric Fermentation	CH ₄	3345	3345	6.3%	45.2%
1A4a	Commercial: Liquid Fuels	CO ₂	2702	2702	5.1%	50.3%
1A4b	Residential: Gaseous Fuels	CO ₂	2600	2600	4.9%	55.2%
4A1	Forest land remaining Forest land	CO ₂	-2396	2396	4.5%	59.7%
1A1	Energy Industries: Other Fuels	CO ₂	2382	2382	4.5%	64.1%
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO ₂	2220	2220	4.2%	68.3%
1A2	Manufacturing Industry and Construction: Liquid Fuels	CO ₂	1888	1888	3.5%	71.9%
2A1	Cement production	CO ₂	1715	1715	3.2%	75.1%
2F1	Refrigeration and air conditioning	HFC	1432	1432	2.7%	77.8%
1A4a	Commercial: Gaseous Fuels	CO ₂	1401	1401	2.6%	80.4%
3Da	Direct emissions from managed soils	N ₂ O	1070	1070	2.0%	82.4%
4B1	Cropland remaining Cropland	CO ₂	1029	1029	1.9%	84.4%
3B1-3B4	Manure management	CH ₄	808	808	1.5%	85.9%
4A2	Land converted to Forest	CO ₂	-517	517	1.0%	86.8%
1A2	Manufacturing Industry and Construction: Solid Fuels	CO ₂	474	474	0.9%	87.7%
1A1	Energy Industries: Liquid Fuels	CO ₂	471	471	0.9%	88.6%
3Db	Indirect emissions from managed soils	N ₂ O	448	448	0.8%	89.5%
1A4c	Agriculture and Forestry: Liquid Fuels	CO ₂	407	407	0.8%	90.2%
1A1	Energy Industries: Gaseous Fuels	CO ₂	399	399	0.7%	91.0%
1A2	Manufacturing Industry and Construction: Other Fuels	CO ₂	367	367	0.7%	91.7%
5A	Solid Waste Disposal	CH ₄	351	351	0.7%	92.3%
4E2	Land converted to Settlements	CO ₂	326	326	0.6%	92.9%
3B5	Indirect N ₂ O emissions from manure management	N ₂ O	282	282	0.5%	93.5%
4C2	Land converted to Grassland	CO ₂	216	216	0.4%	93.9%
2G	Other Product Manufacture and Use	SF ₆	210	210	0.4%	94.3%
1B2	Oil and Natural Gas Energy Production	CH ₄	194	194	0.4%	94.6%
5D	Wastewater treatment and discharge	CH ₄	177	177	0.3%	95.0%
5D	Wastewater treatment and discharge	N ₂ O	144	144	0.3%	95.2%

For the **base year 1990**, 32 categories are identified as **level key categories** under Approach 1 (Table 1-5). The following categories are key according to level in 1990, but not in 2015:

- 1A3a Civil Aviation, Liquid Fuels, CO₂
- 1A5 Other (Military), Liquid fuels, CO₂

- 2A4 Other process uses of carbonates, CO₂
- 2 Indirect CO₂ emissions, CO₂
- 2C3 Aluminium production, CO₂
- 4G Harvested wood products, CO₂

On the other hand, the following categories are key according to level in 2015, but not in 1990:

- 2F1 Refrigeration and air conditioning, HFC
- 2G Other product manufacture and use, SF₆
- 4C2 Land converted to grassland, CO₂
- 5D Wastewater treatment and discharge, CH₄
- 5D Wastewater treatment and discharge, N₂O

Table 1-5 Switzerland's Approach 1 level key categories for the base year 1990 including LULUCF categories, sorted by emission contribution to the national total.

APPROACH 1 LEVEL ASSESSMENT FOR BASE YEAR						
A	B	C	D	E	F	G
Code	IPCC Category	GHG	Ex,0 (kt CO2 eq)	Ex,0 (kt CO2 eq)	Lx,0	Cumulative Total
1A1	Energy Industries: Other Fuels	CO2	1492	1492	2.6%	69.1%
1A1	Energy Industries: Liquid Fuels	CO2	686	686	1.2%	87.3%
1A1	Energy Industries: Gaseous Fuels	CO2	243	243	0.4%	93.5%
1A2	Manufacturing Industry and Construction: Liquid Fuels	CO2	3889	3889	6.7%	51.2%
1A2	Manufacturing Industry and Construction: Solid Fuels	CO2	1275	1275	2.2%	73.8%
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO2	1060	1060	1.8%	79.9%
1A2	Manufacturing Industry and Construction: Other Fuels	CO2	192	192	0.3%	94.7%
1A3a	Civil Aviation: Liquid Fuels	CO2	253	253	0.4%	93.1%
1A3b	Road Transportation: Gasoline	CO2	11334	11334	19.6%	19.6%
1A3b	Road Transportation: Diesel	CO2	2633	2633	4.6%	62.0%
1A4a	Commercial: Liquid Fuels	CO2	4261	4261	7.4%	44.5%
1A4a	Commercial: Gaseous Fuels	CO2	981	981	1.7%	81.6%
1A4b	Residential: Liquid Fuels	CO2	10099	10099	17.5%	37.1%
1A4b	Residential: Gaseous Fuels	CO2	1450	1450	2.5%	71.6%
1A4c	Agriculture and Forestry: Liquid Fuels	CO2	485	485	0.8%	90.2%
1A5	Other (Military): Liquid Fuels	CO2	218	218	0.4%	94.3%
1B2	Oil and Natural Gas Energy Production	CH4	336	336	0.6%	92.1%
2	Indirect CO2 emissions	CO2	365	365	0.6%	91.5%
2A1	Cement production	CO2	2581	2581	4.5%	66.5%
2A4	Other process uses of carbonates	CO2	164	164	0.3%	94.9%
2C3	Aluminium production	CO2	139	139	0.2%	95.2%
3A	Enteric Fermentation	CH4	3585	3585	6.2%	57.5%
3B1-3B4	Manure management	CH4	924	924	1.6%	83.2%
3B5	Indirect N2O emissions from manure management	N2O	242	242	0.4%	94.0%
3Da	Direct emissions from managed soils	N2O	1239	1239	2.1%	75.9%
3Db	Indirect emissions from managed soils	N2O	635	635	1.1%	88.4%
4A1	Forest land remaining Forest land	CO2	-319	319	0.6%	92.7%
4A2	Land converted to Forest	CO2	-571	571	1.0%	89.4%
4B1	Cropland remaining Cropland	CO2	913	913	1.6%	84.8%
4E2	Land converted to Settlements	CO2	394	394	0.7%	90.9%
4G	HWP Harvest Wood Prod	CO2 biog.	-1231	1231	2.1%	78.1%
5A	Solid Waste Disposal	CH4	763	763	1.3%	86.1%

Regarding the **trend assessment** between the base year 1990 and the most recent year 2015, 33 categories are identified as trend key categories under Approach 1 (Table 1-6).

Table 1-6 Switzerland's Approach 1 trend key categories between 1990 and 2015 including LULUCF categories, sorted by contribution to the trend assessment.

APPROACH 1 TREND ASSESSMENT FOR 2015							
A	B	C	D	E	F	G	H
Code	IPCC Category	GHG	Ex,0 (kt CO2 eq)	Ex,t (kt CO2 eq)	Trend Assessm.	Contr. to Trend	Cumulative Total
1A3b	Road Transportation: Diesel	CO2	2633	7182	0.084%	17.9%	17.9%
1A4b	Residential: Liquid Fuels	CO2	10099	5864	0.053%	11.3%	29.2%
1A3b	Road Transportation: Gasoline	CO2	11334	7669	0.041%	8.7%	37.9%
4A1	Forest land remaining Forest land	CO2	-319	-2396	0.035%	7.5%	45.4%
1A2	Manufacturing Industry and Construction: Liquid Fuels	CO2	3889	1888	0.027%	5.7%	51.1%
2F1	Refrigeration and air conditioning	HFC	0	1432	0.025%	5.3%	56.4%
1A4b	Residential: Gaseous Fuels	CO2	1450	2600	0.023%	4.9%	61.2%
4G	HWP Harvest Wood Prod	CO2 biog.	-1231	-70	0.023%	4.8%	66.1%
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO2	1060	2220	0.022%	4.7%	70.8%
1A1	Energy Industries: Other Fuels	CO2	1492	2382	0.018%	3.9%	74.7%
1A4a	Commercial: Liquid Fuels	CO2	4261	2702	0.018%	3.9%	78.6%
1A2	Manufacturing Industry and Construction: Solid Fuels	CO2	1275	474	0.011%	2.4%	81.0%
2A1	Cement production	CO2	2581	1715	0.010%	2.1%	83.1%
1A4a	Commercial: Gaseous Fuels	CO2	981	1401	0.009%	2.0%	85.1%
5A	Solid Waste Disposal	CH4	763	351	0.006%	1.2%	86.3%
4B1	Cropland remaining Cropland	CO2	913	1029	0.004%	0.8%	87.1%
2	Indirect CO2 emissions	CO2	365	106	0.004%	0.8%	87.9%
1A2	Manufacturing Industry and Construction: Other Fuels	CO2	192	367	0.003%	0.7%	88.6%
1A1	Energy Industries: Gaseous Fuels	CO2	243	399	0.003%	0.7%	89.3%
3A	Enteric Fermentation	CH4	3585	3345	0.003%	0.7%	90.0%
4C2	Land converted to Grassland	CO2	91	216	0.002%	0.5%	90.5%
1A1	Energy Industries: Liquid Fuels	CO2	686	471	0.002%	0.5%	91.0%
2C3	Aluminium production	CO2	139	0	0.002%	0.5%	91.4%
4A2	Land converted to Forest	CO2	-571	-517	0.002%	0.4%	91.9%
3Db	Indirect emissions from managed soils	N2O	635	448	0.002%	0.4%	92.3%
2C3	Aluminium production	PFC	116	0	0.002%	0.4%	92.7%
1B2	Oil and Natural Gas Energy Production	CH4	336	194	0.002%	0.4%	93.0%
1A3b	Road Transportation: Gasoline	N2O	137	25	0.002%	0.4%	93.4%
2G	Other Product Manufacture and Use	SF6	137	210	0.002%	0.3%	93.7%
1A3a	Civil Aviation: Liquid Fuels	CO2	253	137	0.001%	0.3%	94.0%
1A3b	Road Transportation: Gasoline	CH4	116	18	0.001%	0.3%	94.4%
5B	Biological Treatment of Solid Waste	CH4	36	110	0.001%	0.3%	94.6%
1A3b	Road Transportation: Diesel	N2O	6	79	0.001%	0.3%	94.9%
1A4b	Residential: Biomass	CH4	110	26	0.001%	0.3%	95.2%

Approach 2

Given that the threshold is set at 90%, the number of key categories is smaller under Approach 2 compared to Approach 1 for both, level and trend assessment.

Concerning the **level assessment**, 23 out of 157 categories are identified as key categories for the current reporting year 2015 (Table 1-7, Table 1-8).

The most relevant sources in 2015 contributing more than 10% to the level assessment are the following:

- 3Da Direct emissions from managed soils, N₂O
- 4A1 Forest land remaining forest land, CO₂

Several categories being key categories in Approach 1 are not key anymore when assessed with Approach 2. Nevertheless, there are 3 categories only being key under Approach 2 (year 2015):

- 4C1 Grassland remaining grassland, CO₂
- 4D1 Wetlands remaining wetlands, CO₂
- 4 III N mineralization, N₂O

Table 1-7 Switzerland's Approach 2 level key categories for the year 2015 including LULUCF categories, sorted by contribution to the uncertainty of the level assessment.

Code	IPCC Category	GHG	Ex,t (kt CO ₂ eq)	Ex,t (kt CO ₂ eq)	Lx,t	Cumulative Total
4A1	Forest land remaining Forest land	CO ₂	-2396	2396	21.8%	21.8%
3Da	Direct emissions from managed soils	N ₂ O	1070	1070	11.0%	32.8%
4C1	Grassland Remaining	CO ₂	112	112	9.3%	42.1%
3Db	Indirect emissions from managed soils	N ₂ O	448	448	7.6%	49.7%
3B5	Indirect N ₂ O emissions from manure management	N ₂ O	282	282	7.0%	56.7%
3A	Enteric Fermentation	CH ₄	3345	3345	5.5%	62.2%
4A2	Land converted to Forest	CO ₂	-517	517	4.3%	66.5%
3B1-3B4	Manure management	CH ₄	808	808	4.2%	70.7%
4B1	Cropland remaining Cropland	CO ₂	1029	1029	2.7%	73.4%
2F1	Refrigeration and air conditioning	HFC	1432	1432	2.6%	76.0%
1A1	Energy Industries: Other Fuels	CO ₂	2382	2382	2.4%	78.4%
5D	Wastewater treatment and discharge	N ₂ O	144	144	2.1%	80.6%
4E2	Land converted to Settlements	CO ₂	326	326	1.5%	82.1%
1A4b	Residential: Gaseous Fuels	CO ₂	2600	2600	1.2%	83.3%
5A	Solid Waste Disposal	CH ₄	351	351	1.1%	84.4%
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO ₂	2220	2220	1.1%	85.4%
5D	Wastewater treatment and discharge	CH ₄	177	177	0.9%	86.3%
4III	Direct N ₂ O from Disturbance	N ₂ O	56	56	0.7%	87.0%
4D1	Wetland remaining Wetland	CO ₂	68	68	0.7%	87.7%
1A4a	Commercial: Gaseous Fuels	CO ₂	1401	1401	0.6%	88.3%
4C2	Land converted to Grassland	CO ₂	216	216	0.6%	89.0%
2A1	Cement production	CO ₂	1715	1715	0.6%	89.6%
1A3b	Road Transportation: Diesel	CO ₂	7182	7182	0.6%	90.2%

Regarding the **trend assessment** between the base year 1990 and the most recent year 2015, 26 categories are identified as trend key categories under Approach 2 (Table 1-8).

Table 1-8 Switzerland's Approach 2 trend key categories between 1990 and 2015 including LULUCF categories, sorted by contribution to the trend assessment.

APPROACH 2 TREND ASSESSMENT WITH UNCERTAINTIES FOR 2015							
A	B	C	D	E	F	G	H
Code	IPCC Category	GHG	Ex,0 (kt CO2 eq)	Ex,t (kt CO2 eq)	Trend Assessm.	Contr. to Trend	Cumulative Total
4A1	Forest land remaining Forest land	CO2	-313	-2369	3.494%	35.1%	35.1%
4G	HWP Harvest Wood Prod	CO2 biog.	671	38	1.333%	13.4%	48.4%
4C1	Grassland Remaining	CO2	84	92	0.789%	7.9%	56.3%
2F1	Refrigeration and air conditioning	HFC	0	1428	0.493%	4.9%	61.3%
3Db	Indirect emissions from managed soils	N2O	636	418	0.385%	3.9%	65.2%
3B5	Indirect N2O emissions from manure management	N2O	231	300	0.293%	2.9%	68.1%
2C3	Aluminium production	CO2	139	0	0.213%	2.1%	70.2%
1A1	Energy Industries: Other Fuels	CO2	1496	2379	0.198%	2.0%	72.2%
4A2	Land converted to Forest	CO2	-581	-508	0.191%	1.9%	74.1%
5A	Solid Waste Disposal	CH4	768	350	0.182%	1.8%	76.0%
2C3	Aluminium production	PFC	116	0	0.178%	1.8%	77.7%
1A4b	Residential: Gaseous Fuels	CO2	1449	2599	0.114%	1.1%	78.9%
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO2	1059	2219	0.114%	1.1%	80.0%
5D	Wastewater treatment and discharge	N2O	116	147	0.109%	1.1%	81.1%
2	Indirect CO2 emissions	CO2	366	107	0.108%	1.1%	82.2%
4B1	Cropland remaining Cropland	CO2	914	1027	0.108%	1.1%	83.3%
1A2	Manufacturing Industry and Construction: Solid Fuels	CO2	1276	474	0.083%	0.8%	84.1%
1A3b	Road Transportation: Gasoline	N2O	138	25	0.081%	0.8%	84.9%
1A3b	Road Transportation: Diesel	CO2	2632	7183	0.075%	0.8%	85.7%
4C2	Land converted to Grassland	CO2	91	217	0.075%	0.8%	86.4%
1A1	Energy Industries: Solid Fuels	CO2	49	0	0.075%	0.8%	87.2%
2G	Other Product Manufacture and Use	N2O	104	44	0.070%	0.7%	87.9%
5D	Wastewater treatment and discharge	CH4	132	178	0.058%	0.6%	88.5%
5B	Biological Treatment of Solid Waste	CH4	35	110	0.057%	0.6%	89.1%
1B2	Oil and Natural Gas Energy Production	CH4	337	194	0.056%	0.6%	89.6%
1A3b	Road Transportation: Gasoline	CH4	116	18	0.055%	0.6%	90.2%

1.5.1.3 Summary of combined KCA including LULUCF categories

A summary of the key category analysis for 2015 is shown in Table 1-9, considering level and trend assessments for both, Approach 1 and Approach 2.

Table 1-9 Summary of Switzerland's combined KCA Approach for 2015 including LULUCF categories sorted by NFR code (first column). The abbreviations used in the last column indicate both, the approach and whether a certain category is identified as key because of the level or trend assessment (L1 = level according to Approach 1, L2 = level according to Approach 2) or the trend assessment (T1 = trend according to Approach 1, T2 = trend according to Approach 2).

Code	IPCC Category	GHG	Identification Criteria
1A1	Energy Industries: Gaseous Fuels	CO2	L1, T1
1A1	Energy Industries: Liquid Fuels	CO2	L1, T1
1A1	Energy Industries: Other Fuels	CO2	L1, L2, T1, T2
1A1	Energy Industries: Solid Fuels	CO2	T2
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO2	L1, L2, T1, T2
1A2	Manufacturing Industry and Construction: Liquid Fuels	CO2	L1, T1
1A2	Manufacturing Industry and Construction: Other Fuels	CO2	L1, T1
1A2	Manufacturing Industry and Construction: Solid Fuels	CO2	L1, T1, T2
1A3a	Civil Aviation: Liquid Fuels	CO2	T1
1A3b	Road Transportation: Gasoline	CH4	T1, T2
1A3b	Road Transportation: Gasoline	CO2	L1, T1
1A3b	Road Transportation: Gasoline	N2O	T1, T2
1A3b	Road Transportation: Diesel	CO2	L1, L2, T1, T2
1A3b	Road Transportation: Diesel	N2O	T1
1A4a	Commercial: Gaseous Fuels	CO2	L1, L2, T1
1A4a	Commercial: Liquid Fuels	CO2	L1, T1
1A4b	Residential: Biomass	CH4	T1
1A4b	Residential: Gaseous Fuels	CO2	L1, L2, T1, T2
1A4b	Residential: Liquid Fuels	CO2	L1, T1
1A4c	Agriculture and Forestry: Liquid Fuels	CO2	L1
1B2	Oil and Natural Gas Energy Production	CH4	L1, T1, T2
2A1	Cement production	CO2	L1, L2, T1
2C3	Aluminium production	CO2	T1, T2
2C3	Aluminium production	PFC	T1, T2
2F1	Refrigeration and air conditioning	HFC	L1, L2, T1, T2
2G	Other Product Manufacture and Use	N2O	T2
2G	Other Product Manufacture and Use	SF6	L1, T1
2	Indirect CO2 emissions	CO2	T1, T2
3A	Enteric Fermentation	CH4	L1, L2, T1
3B1-3B4	Manure management	CH4	L1, L2
3B5	Indirect N2O emissions from manure management	N2O	L1, L2, T2
3Da	Direct emissions from managed soils	N2O	L1, L2
3Db	Indirect emissions from managed soils	N2O	L1, L2, T1, T2
4A1	Forest land remaining Forest land	CO2	L1, L2, T1, T2
4A2	Land converted to Forest	CO2	L1, L2, T1, T2
4B1	Cropland remaining Cropland	CO2	L1, L2, T1, T2
4C1	Grassland Remaining	CO2	L2, T2
4C2	Land converted to Grassland	CO2	L1, L2, T1, T2
4D1	Wetland remaining Wetland	CO2	L2
4E2	Land converted to Settlements	CO2	L1, L2
4G	HWP Harvest Wood Prod	CO2 biog.	T1, T2
4III	Direct N2O from Disturbance	N2O	L2
5A	Solid Waste Disposal	CH4	L1, L2, T1, T2
5B	Biological Treatment of Solid Waste	CH4	T1, T2
5D	Wastewater treatment and discharge	CH4	L1, L2, T2
5D	Wastewater treatment and discharge	N2O	L1, L2, T2

1.5.2 KP-LULUCF inventory

Switzerland identified four key categories for activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (Forest management, Afforestation and Reforestation, Deforestation, Harvested wood products). The approach relies on full inventory KCA (with LULUCF), KP - CRF association and qualitative assessment. A detailed description is presented in chp. 11.6.1.

1.6 General uncertainty evaluation

1.6.1 GHG inventory

1.6.1.1 Approach 1 and Approach 2 analysis

This chapter presents the main results of the uncertainty evaluation Approach 1 and Approach 2 in accordance with the 2006 IPCC Guidelines (IPCC 2006/Chapter 3 Uncertainties). Concerning key assumptions and requirements for both approaches we refer to the guidelines, here we only recap the clues of both approaches:

- Approach 1: based on propagation of error, uncertainty in the emission level in 2015 and in the trend between the reporting year (2015) and the base year (1990) is estimated for the inventory total and for the single source categories and gases using uncertainty ranges of corresponding activity data and emission factors.
- Approach 2: is based on Monte Carlo analysis (IPCC 2006, UNFCCC 2014a). This approach provides a detailed category-by-category assessment of uncertainty, particularly where uncertainties are large, distribution is non-normal, the algorithms are complex functions and/or there are correlations between some of the activity data, emission factors or both. The principle of Monte Carlo analysis is to select random values for emission factor and activity data from within their individual probability distributions, and to calculate the corresponding emission values. This procedure is repeated until an adequately stable result has been found. The results of all iterations yield the overall emission probability distribution.

All uncertainties are given as half of the 95% confidence interval divided by the mean and expressed as a percentage (approximately two standard deviations) as suggested by the 2006 IPCC Guidelines (IPCC 2006).

The uncertainty analyses Approach 1 and Approach 2 are updated on a yearly basis.

The following chapters present the overall results of the uncertainty evaluation. Specific information about the uncertainty estimation for activity data, emission factors or emissions of each source category is included in the respective sectoral chapters (chp. 3 to 9) below. Detailed results of both approaches as well as further methodological information concerning Approach 2 are presented in Annex A2.

1.6.1.2 Data used

The evaluation includes uncertainties regarding activity data, emission factors and – in a few cases – emissions. Uncertainties in the GWP values are not taken into account.

Uncertainties are estimated for the reporting year. For some categories uncertainties in activity data and emission factors are considerably lower as compared to 1990. For example, uncertainty of N₂O emissions in 2B2 Nitric acid production used to be much higher than in the current reporting year, for which emissions are derived from continuous N₂O measurements. Continuous N₂O measurements started in 2013.

Uncertainty distributions are assumed to be symmetric for Approach 1. For the Monte Carlo simulation, some asymmetric distributions (triangular) were adopted.

For categories with quantitative uncertainty data available, the input information from studies or from the data suppliers is used for the uncertainty evaluation. This is mainly the case for key categories. However, no explicit information on uncertainties is available for a few key categories. For these cases, authors of the NIR chapters, FOEN experts involved and several data suppliers derived estimates of uncertainties based on the 2006 IPCC Guidelines (IPCC 2006) default values and on information concerning the process of data collection for activity data and emission factors (import or sales statistics, surveys or modelling). Several experts from data suppliers were contacted for further information on some of the uncertainties. Industry associations/sources also provided published or unpublished uncertainty estimates for their data. The data sources can be found in the relevant sub-sections on “Uncertainties and time-series consistency” in each of the sectoral chapters (chp. 3 to 9) below.

For categories with no quantitative uncertainty data available, the NIR provides qualitative estimates of uncertainties. The elaboration of a quantitative uncertainty assessment for these categories would present a large effort with only limited effect on the overall uncertainty and therefore it has been decided to realize a semi-quantitative assessment. This includes the definition of a list of the combined uncertainties for all gases and three uncertainty levels: low, medium and high (see Table 1-10). These values are motivated by the comparison of uncertainty analyses of several countries carried out by de Keizer et al. (2007), as presented at the 2nd Internat. Workshop on Uncertainty in Greenhouse Gas Inventories (Vienna 27-28 September 2007), and by expert judgement from sectoral experts and authors.

Table 1-10 Semi-quantitative (combined) uncertainties (U) for the emission of categories with no quantitative uncertainty data available. Note that there is no source of NF₃, for which a semi-quantitative uncertainty value is required.

Gas	Uncertainty Category	Combined Uncertainty
CO ₂	low	2%
	medium	10%
	high	40%
CH ₄	low	15%
	medium	30%
	high	60%
N ₂ O	low	40%
	medium	80%
	high	150%
HFC	medium	20%
PFC	medium	20%
SF ₆	medium	20%

The uncertainties of sector 1 Energy were broadly updated for the last submission in 2016. They are described in detail in chp. 3.2.4.7. Main result of the new evaluation were a lowering of the uncertainty of CO₂ for this sector.

Despite the investigation carried out for the current uncertainty analyses it will be necessary to further motivate institutions to supply not only average data but also estimates of associated uncertainties.

1.6.1.3 Results of Approach 1 uncertainty evaluation

Level and trend uncertainty analyses are carried out excluding and including the LULUCF sector. Table 1-11 gives a summary for the Approach 1 uncertainties for the national total emissions and removals.

Table 1-11 Switzerland's relative uncertainties for national total GHG emission excluding and including the LULUCF sector – Approach 1: Level uncertainties 2015 and trend uncertainties 1990–2015. The uncertainty analysis is based on emissions including indirect CO₂ emissions.

Approach 1 Uncertainty Analysis		
Inventory	Level uncertainty	Trend uncertainty
	2015	1990-2015
excl. LULUCF	3.75%	1.51%
incl. LULUCF	6.36%	4.03%

Uncertainty analysis results for the Switzerland's GHG inventory 2015 and including indirect CO₂ emissions with Approach 1:

- The level uncertainty 2015 is **3.75%** excluding LULUCF and **6.36%** including LULUCF. Explanations are given below in the chapter on Approach 2 results.
- The trend uncertainty 1990–2015 is **1.51%** excluding LULUCF and **4.03%** including LULUCF.
- Compared to the results of the previous inventory 2014 and excluding indirect CO₂ emissions (FOEN 2016):
- In 2014, level uncertainty was **3.81%** excl. LULUCF and **4.76%** incl. LULUCF
- In the period 1990–2014, trend uncertainty was **1.44%** excl. LULUCF and **2.31%** incl. LULUCF.

The level uncertainty excluding LULUCF slightly decreased (from 3.81% to 3.75%) compared to the previous uncertainty analysis since several uncertainties in sector 1 Energy and 2 Industrial processes and product use have been reduced in the current submission. The level uncertainty when including LULUCF, however, has clearly increased in 2015 compared to 2014 (from 4.76% to 6.36%). This is mainly a result of an increase in uncertainty values in category 4A1 Forest land remaining forest land (see sectoral chapters on uncertainties). Trend uncertainties including LULUCF increased for the current submission, which can be

explained by an increase in uncertainties in the LULUCF sector. Excluding LULUCF, there is only a slight increase in trend uncertainty. Indirect CO₂ emissions are included for the first time in Submission 2017. This contributes to an increase in trend uncertainty.

Detailed results of the Approach 1 uncertainty analysis for GHG emissions 2015 per category are shown in Table A – 2. Details of the uncertainty estimates for specific source categories are provided in the sub-sections on “Uncertainties and time-series consistency” in each of the chapters on source categories below.

It should be noted that the results of the Approach 1 uncertainty analysis for GHG emissions do not, or not fully, take into account the following factors that may further increase uncertainties:

- correlations between source categories that are not considered by Approach 1 (e.g. production data used for industry emissions in both categories 1A2 Manufacturing industries and 2 Industrial processes and product use, or cattle numbers used for emissions related to enteric fermentation and animal manure production);
- errors due to neglected temporal variability when assuming constant parameters over time (e.g. emission factors).
- errors due to non-normal, asymmetric distribution of the uncertainties;
- errors due to methodological shortcomings, i.e. simplified approaches;
- errors due to sources not reported (these are assumed to be very small).

On the other hand, the Approach 2 uncertainty evaluation described below explicitly takes into account correlations between sources and asymmetric distributions.

1.6.1.4 Results of Approach 2 uncertainty evaluation (Monte Carlo)

In the present analysis, Monte Carlo simulations were performed to estimate uncertainties both in emissions 2015 and in emission trends 1990–2015, at the source category level as well as for the inventory as a whole (excluding and including LULUCF). The simulations were run with the commercial software package Crystal Ball¹ (© Decisioneering, Release 11.1.2.4.400). This tool generates random numbers within user defined probability ranges and probability distributions. As a result, selected statistics are produced for the forecast variables.

Assumptions for the Monte Carlo simulations are given in Annex 2. In this chapter, only the main results of the simulations for level and trend analyses are presented.

¹ https://docs.oracle.com/cd/E40248_01/epm.1112/cb_statistical/frameset.htm?index.html

Table 1-12 Switzerland's relative uncertainties for national total GHG emission excluding and including the LULUCF sector – Approach 2: Level uncertainties 2015 and trend uncertainties 1990–2015. The emission trend 1990–2015 is -10.45% excl. LULUCF and -11.65% incl. LULUCF. The emissions shown in this table include indirect CO₂ emissions.

Approach 2 (Monte Carlo) Uncertainty Analysis						
Version	Level uncertainty 2015			Trend uncertainty 1990-2015		
	2.5 percentile	97.5 percentile	mean	2.5 percentile	97.5 percentile	mean
excl. LULUCF	-3.39%	3.73%	3.56%	-13.14%	-7.90%	2.62%
incl. LULUCF	-6.08%	6.21%	6.14%	-17.18%	-6.14%	5.52%

Approach 2 uncertainties excluding LULUCF and including indirect CO₂ emissions:

- The total uncertainty level of Switzerland's 2015 national total GHG emissions excluding LULUCF is **3.56%** with a slightly asymmetric 95% confidence interval between 96.61% and 103.73%.
- The trend in national total emissions excluding LULUCF between 1990 and 2015 is -10.45%. With a probability of 95% the trend lies within the range of -13.14% to -7.90%, which corresponds to a **mean trend uncertainty of 2.62%**.

Approach 2 uncertainties including LULUCF and including indirect CO₂ emissions:

- The total uncertainty level of Switzerland's 2015 national total GHG emissions including LULUCF is **6.14%** with a slightly asymmetric 95% confidence interval between 93.92% and 106.21%.
- The trend in national total emissions including LULUCF between 1990 and 2015 is -11.65%. With a probability of 95%, the trend lies within the range of -17.18% to -6.14%, which corresponds to a **mean trend uncertainty of 5.52%**.

That means that level and trend uncertainty are considerably higher if LULUCF categories are included in the uncertainty analysis, which is caused by large contributions, large uncertainties and strong trends of several LULUCF categories.




Figure 1-3 Probability distributions of the simulated total emissions excluding LULUCF for the base year 1990 (in orange) and year 2015 (in green). The vertical lines show simulated mean and percentile values (black for 2015, red for 1990). The number of Monte Carlo runs is 100'000. X-axis: total emissions (excl. LULUCF) [kt CO₂ eq]. Y-axis: Probability (left) and Frequency (right). The simulated values deviate from the reported inventory values (see Table A – 8 for detailed deviations: E.g. the simulated value including indirect CO₂ emissions 2015 is 49'414 kt CO₂ eq, whereas the reported value is 48'151 kt CO₂ eq, 2.6% lower than the simulated value). The emissions shown in this figure include indirect CO₂ emissions.

In the course of Monte Carlo simulation, the uncertainties are also evaluated by gas (see Table 1-13). As expected, CO₂ emissions have the highest precision or the lowest uncertainties among the Kyoto gases.

Table 1-13 Approach 2 level uncertainties by gas for the total national emissions 2015 excluding LULUCF. The emissions shown in this table include indirect CO₂ emissions.

Gas	Emission 2015 (excl. LULUCF) kt CO ₂ eq	Lower bound 2.5 percentile kt CO ₂ eq	Upper bound 97.5 percentile kt CO ₂ eq	Mean absolute uncertainty kt CO ₂ eq	Mean relative uncertainty %
CO ₂	38'865	38'574	39'160	293	0.75%
CH ₄	5'087	4'162	6'013	926	18.2%
N ₂ O	3'612	2'320	5'153	1'416	39.2%
HFC	1'536	1'249	1'823	287	18.7%
PFC	57	49	66	8	14.7%
SF ₆	256	204	307	52	20.2%
NF ₃	0.49	-0.46	1.43	0.95	195%
Total	49'414	47'739	51'255	1'758	3.56%

Detailed results per category of the Monte Carlo simulation are presented in Table A – 7, inputs on probability distributions and correlation coefficients in Tables Table A – 4 to Table A – 6.

The following chart – called Tornado plot – shows the results of a sensitivity analysis, depicting the most important uncertainties. These can either be emission factors, activity data or emissions. The bars depict the amount of uncertainty introduced compared to total emissions (on x-axis). On the left-hand side, the variable is indicated containing the information of type (EM emission, EF emission factor, AD activity data), NFR number and gas. The letter “t” refers to year 2015.

Categories 4A1 Forest land remaining forest land (CO₂), 4C1 Grassland remaining grassland (CO₂) and 3Da. Direct N₂O emissions from managed soils are the most important contributors to level uncertainty. The fact that two out of the three most important contributors stem from the LULUCF sector explains the result given above that the level uncertainty including LULUCF is higher than without LULUCF.

Based on the analysis of the predominant contributions to the uncertainty of the Swiss greenhouse gas inventory, the FOEN commissions and/or supports various projects. In line with UNFCCC (2017/ID#L.12) the soil carbon model Yasso07 will be further developed to improve the accuracy in the estimates of carbon changes in mineral soil, litter and dead wood in forest land (see chp. 6.4.6 for details). A Tier 3 methodological approach for the quantification of carbon stock changes in agricultural soils (cropland remaining cropland and grassland remaining grassland) is currently under development at Agroscope research station (see chp. 6.5.6 and chp. 6.6.6, respectively). In the agriculture sector, a research project run by Agroscope looks into CH₄ and N₂O emissions from current pasture practices using micrometeorological measurements and modelling approaches. With regard to HFC emissions, the FOEN supports a long-term monitoring initiative at Jungfrauoch to derive independent emission estimates from atmospheric concentrations (Annex 5.1). Another, more recent verification activity addresses national total CH₄ and N₂O emissions (Annex 5.2).




Figure 1-4 Tornado plot of the uncertainties by category. Abbrev.: “t” refers to 2015, “EF” emission factor, “AD” activity data, “EM” emissions. X-axis: Simulated national total of CO₂ eq emissions (in kt) including LULUCF in 2015 (the simulated values deviate from the reported inventory values, see Table A – 8 for details). The width of the bar shows the combined uncertainty introduced by the corresponding uncertainty. Uncertainty analysis includes indirect CO₂ emissions.

Further results of the Monte Carlo simulations are shown in Annex 2.2.

1.6.1.5 Comparison of Approach 1 and Approach 2

In the GHG inventory, the variability of the uncertainties is high, their statistical distribution may clearly deviate from normal distributions and they can be correlated. Approach 1 is based on simple error propagation, which assumes only small, normally distributed and uncorrelated uncertainties. The application of the Approach 1 is therefore not the optimal method for determining the uncertainties of a GHG inventory. The more appropriate choice, which is recommended by the IPCC 2006 Guidelines (IPCC 2006), is the Monte Carlo simulation (Approach 2), which is designed for uncertainties of any extent, any statistical distribution and any correlated parameters. The results of the Monte Carlo simulation are therefore considered to provide a more realistic picture of the uncertainties than the results of Approach 1.

Level uncertainty

Approach 2 excl. LULUCF leads to an overall level uncertainty of 3.56%, which is slightly lower than the result of Approach 1 (3.75%). The correct treatment of large uncertainties, asymmetric distributions for agricultural sources and accounting for relevant correlations lead all together to a slight decrease in the level uncertainty.

The same holds for the level uncertainty incl. LULUCF. With Approach 2, an overall level uncertainty of 6.14% is computed, which is also slightly lower than the result of Approach 1 6.36%.

Trend uncertainty

In terms of trend uncertainty, the results of Approach 2 show considerably higher uncertainties than the results of Approach 1. If LULUCF categories are excluded, Approach 2 leads to an uncertainty of 2.62% and Approach 1 to 1.51%, whereas when LULUCF categories are included, the numbers are 5.52% and 4.03%, respectively. Positive correlations for activity data and emission factors between the base year and 2015 tend to increase trend uncertainty. This effect is enforced for the analysis including LULUCF due to the high uncertainty values in this sector.

1.6.2 KP-LULUCF inventory

Uncertainty estimates for KP-LULUCF activities are presented in chp. 11.3.1.5.

1.7 General assessment of completeness

1.7.1 GHG inventory

Source and sink categories that are not estimated or included elsewhere are listed in CRF Table9.

For the following categories, for which reporting is “not mandatory”, the notation key “not estimated” is used

- CH₄ emissions from 4.A Forest Land/4(II), 4.C Grassland/4(II), 4.E Settlements/4(V) Biomass Burning/Settlements burning,
- CO₂ emissions from 4.E Settlements/4(V) Biomass Burning/Settlements burning,
- N₂O emissions from 4.D Wetlands/4(II), 4.E Settlements/4(V) Biomass Burning/Settlements burning.

Notation key “included elsewhere” is used in 1A2c and 2B1 for reasons of confidentiality, and in 1A3a / aviation gasoline, 4(I)E, 4(IV), 4(V), 5C1 and 5D2 because data for further disaggregation are not available.

1.7.2 KP-LULUCF inventory

For all known sources and sinks, complete estimates are accomplished for the current submission. Notation keys for the activity coverage and the reported pools are displayed in CRF table NIR-1. A detailed justification for the reported method is given in chapter 11.3.1.2.

2 Trends in greenhouse gas emissions and removals

This chapter provides an overview of Switzerland's GHG emissions and removals in 2015 as well as trends for the period 1990–2015. Numbers in chp. 2.1–2.4 are relevant for reporting under the UNFCCC, whereas numbers in chp. 2.5 refer to accounting under the Kyoto Protocol.

2.1 Aggregated greenhouse gas emissions 2015 (UNFCCC)

Table 2-1 shows the aggregated emissions of all GHG 2015 for each sector and the relative shares of the sectors. Furthermore, emission data on international aviation and marine bunkers are provided. As the table indicates, CO₂ is the main contributor to total greenhouse gas emissions followed by CH₄, N₂O and F-gases. Sector 1 Energy is the main source concerning climate-related emissions followed by sectors 3 Agriculture, 2 IPPU and 5 Waste. In contrast, sector 4 LULUCF is a net sink regarding GHG emissions in 2015.

A breakdown of Switzerland's total emissions by gas (excluding LULUCF) is given in Figure 2-1. Figure 2-2 charts the relative contributions of the individual sectors (excluding LULUCF) to the emissions of each GHG. Trends in GHG emissions are given in chp. 2.2 to 2.5.

The national total of 48'038 kt of CO₂ equivalent (excluding LULUCF) corresponds to 5.8 tonnes of CO₂ equivalent per capita² (CO₂: 4.7 tonnes per capita) emitted to the atmosphere in 2015 (Table 2-1).

Table 2-1 Switzerland's GHG emissions in CO₂ equivalent (kt) by gas and sector in 2015.

Sectors	CO ₂	CH ₄	N ₂ O	HFCs	PFCs	SF ₆	NF ₃	Total	Share
	CO ₂ equivalent (kt)								
1 Energy	36'595	287	231					37'113	77.3%
2 IPPU	2'091	2	49	1'536	57	256	0.5	3'992	8.3%
3 Agriculture	44	4'153	1'877					6'074	12.6%
5 Waste	10	643	193					846	1.8%
6 Other	11	1	1					12	0.0%
Total (excluding LULUCF)	38'751	5'085	2'352	1'536	57	256	0.5	48'038	100.0%
4 LULUCF	-981	13	74					-894	-1.9%
Total (including LULUCF)	37'771	5'099	2'425	1'536	57	256	0.5	47'144	98.1%
<i>International aviation bunkers</i>	4'902	1	40					4'943	
<i>International marine bunkers</i>	25	0.01	0.23					25	

² Population statistics taken from SFSO 2016a.




Figure 2-1 Contribution of individual gases to Switzerland's GHG emissions (excluding LULUCF) in 2015.




Figure 2-2 Relative contributions of the individual sectors (excluding LULUCF) to GHG emissions in 2015.

A clear dominance of CO₂ emissions in 2015 is related to source category 1A Fuel combustion within sector 1 Energy. CH₄ and N₂O emissions mainly originate from agriculture, while F-gas emissions by definition only stem from industrial processes.

2.2 Emission trends by gas

Emission trends by gas for the period 1990–2015 are summarized in Table 2-2.

Table 2-2 Switzerland's GHG emissions in CO₂ equivalent (kt) by gas; 1990–2015. The column below on the far right indicates the percentage change in emissions in 2015 as compared to the base year 1990. HFCs increased by 6'198'125% when compared to 1990 levels (1990 = 0.025 kt CO₂ equivalent).

Greenhouse Gas Emissions	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
CO ₂ emissions including net CO ₂ from LULUCF	43'776	39'647	48'532	43'474
CO ₂ emissions excluding net CO ₂ from LULUCF	44'171	43'423	43'607	45'799
CH ₄ emissions including CH ₄ from LULUCF	6'132	5'802	5'411	5'321
CH ₄ emissions excluding CH ₄ from LULUCF	6'102	5'782	5'396	5'307
N ₂ O emissions including N ₂ O from LULUCF	2'916	2'771	2'624	2'516
N ₂ O emissions excluding N ₂ O from LULUCF	2'829	2'689	2'547	2'438
HFCs	0	245	622	1'064
PFCs	117	17	50	44
SF ₆	137	93	144	203
NF ₃	0	0	0	0
Total (including LULUCF)	53'078	48'575	57'383	52'622
Total (excluding LULUCF)	53'357	52'251	52'365	54'856

Greenhouse Gas Emissions	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Change base year to 2015 (%)
	CO ₂ equivalent (kt)										
CO ₂ emissions including net CO ₂ from LULUCF	45'928	43'123	43'594	41'745	43'713	40'049	40'913	42'040	38'346	37'771	-13.7%
CO ₂ emissions excluding net CO ₂ from LULUCF	45'380	43'388	44'708	43'538	45'053	40'993	42'259	43'202	39'269	38'751	-12.3%
CH ₄ emissions including CH ₄ from LULUCF	5'336	5'317	5'392	5'309	5'289	5'232	5'192	5'131	5'134	5'099	-16.8%
CH ₄ emissions excluding CH ₄ from LULUCF	5'321	5'301	5'379	5'295	5'276	5'217	5'179	5'118	5'121	5'085	-16.7%
N ₂ O emissions including N ₂ O from LULUCF	2'517	2'541	2'561	2'523	2'570	2'515	2'496	2'460	2'475	2'425	-16.8%
N ₂ O emissions excluding N ₂ O from LULUCF	2'437	2'461	2'484	2'448	2'496	2'440	2'423	2'385	2'400	2'352	-16.9%
HFCs	1'111	1'186	1'236	1'247	1'324	1'406	1'486	1'514	1'527	1'536	see caption
PFCs	51	49	58	63	64.50	67.72	71.27	51.93	44.03	57.21	-50.9%
SF ₆	186	172	222	180	148	160	209	252	259	256	86.7%
NF ₃	0.0	0.0	0.1	5	8	6	0.4	0.1	0.4	0.5	-
Total (including LULUCF)	55'129	52'388	53'063	51'071	53'117	49'436	50'368	51'450	47'784	47'144	-11.2%
Total (excluding LULUCF)	54'486	52'556	54'087	52'776	54'370	50'289	51'628	52'523	48'620	48'038	-10.0%

As shown in Table 2-2, Table 2-3, and Figure 2-3, total emissions excluding LULUCF reach a minimum in 2015, being clearly below base year emissions. There is no discernible trend in the period 1990–2005. Only from 2005 onwards, a decreasing trend starts to develop. The emission maximum occurred in 1991. The decrease to a minimum of total emissions in 2015 compared to the base year 1990 is also visible including LULUCF, although the net CO₂ sink generated by LULUCF categories was generally smaller after 1997.

There is a strong correlation between CO₂ emissions and winter climatic conditions (number of heating degree days; see footnote 3, page 66 for further information) in the period 1990–2015 (see Figure 2-7). However, the relative developments of heating degree days and CO₂ emissions are clearly drifting apart in the years since 2003, which indicates that additional effects like reduction measures contribute to emission reductions (Figure 2-7).

Between 1990 and 2015, CH₄ (excluding LULUCF) decreased mainly due to a reduction of livestock that led to a reduction of emissions from enteric fermentation in the agricultural sector. Moreover, from 2000 onwards, a change in waste legislation banning the disposal of municipal solid waste in landfills contributed to this trend.

As a consequence of the declining livestock population and reduced input of synthetic fertilizers, N₂O emissions mainly stemming from manure management and agricultural soils decrease between 1990 und 2015 as well.

HFC emissions increased significantly due to their application as substitutes for CFCs, while PFC emissions declined. SF₆ emissions show relatively large fluctuations between 1990 and 2015. This is the effect of annual fluctuations of the market volumes in the production of electrical equipment and on the other hand the effect of changes in other applications. The increase of SF₆ emissions in recent years is due the disposal of sound proof windows. NF₃ has been used only short-term in the photovoltaic industry.

Table 2-3 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (kt) and the contribution of individual gases (as percentage values) for selected years.

Greenhouse Gas Emissions (excluding LULUCF)	1990		1995		2000		2005		2010	
	kt CO ₂ eq	%								
CO ₂	44'171	82.8%	43'423	83.1%	43'607	83.3%	45'799	83.5%	45'053	82.9%
CH ₄	6'102	11.4%	5'782	11.1%	5'396	10.3%	5'307	9.7%	5'276	9.7%
N ₂ O	2'829	5.3%	2'689	5.1%	2'547	4.9%	2'438	4.4%	2'496	4.6%
HFCs	0	0.0%	245	0.5%	622	1.2%	1'064	1.9%	1'324	2.4%
PFCs	117	0.2%	17	0.0%	50	0.1%	44	0.1%	64	0.1%
SF ₆	137	0.3%	93	0.2%	144	0.3%	203	0.4%	148	0.3%
NF ₃	0	0.0%	0	0.0%	0	0.0%	0	0.0%	8	0.0%
Total (excluding LULUCF)	53'357	100%	52'251	100%	52'365	100%	54'856	100%	54'370	100%

Greenhouse Gas Emissions (excluding LULUCF)	2011		2012		2013		2014		2015	
	kt CO ₂ eq	%								
CO ₂	40'993	81.5%	42'259	81.9%	43'202	82.3%	39'269	80.8%	38'751	80.7%
CH ₄	5'217	10.4%	5'179	10.0%	5'118	9.7%	5'121	10.5%	5'085	10.6%
N ₂ O	2'440	4.9%	2'423	4.7%	2'385	4.5%	2'400	4.9%	2'352	4.9%
HFCs	1'406	2.8%	1'486	2.9%	1'514	2.9%	1'527	3.1%	1'536	3.2%
PFCs	68	0.1%	71	0.1%	52	0.1%	44	0.1%	57	0.1%
SF ₆	160	0.3%	209	0.4%	252	0.5%	259	0.5%	256	0.5%
NF ₃	6	0.0%	0.4	0.0%	0.1	0.0%	0.4	0.0%	0.5	0.0%
Total (excluding LULUCF)	50'289	100%	51'628	100%	52'523	100%	48'620	100%	48'038	100%

Figure 2-3 shows Switzerland's relative GHG emission trends by gas. The base year 1990 is set to 100%.




Figure 2-3 Relative trend of Switzerland's main GHG emissions by gas (excluding LULUCF), 1990–2015 (base year 1990 = 100%). F-gases are not illustrated (see Figure 4-3).

2.3 Emission trends by sources and sinks

Table 2-4 shows the emission trends for all major sources and sink categories. As the largest share of emissions originates from sector 1 Energy, the table includes further information concerning the contributions of energy-related source categories.

2.3.1 Overview

In order to understand trends within the sector 1 Energy, the individual source categories are considered separately. See “Emission trends in sector 1 Energy” and Figure 2-6 below.

In line with economic development, overall emissions in sector 2 Industrial processes and other product use (IPPU) show a decreasing trend in the early 90s and a gradual increase between 1998 and 2015, except for the economically difficult year 2009. Since 2005, the Ordinance on Chemical Risk Reduction (Swiss Confederation 2005) is in place and regulates the use of F-gases. The dominant source category of sector 2 is 2A Mineral industry although the emissions decreased by approximately 1/3 since 1990. If sources are analysed in more detail 2A1 Cement production is the most relevant emitter in this category. Emissions of 2F Product uses as substitutes for ozone-depleting substances (ODS), the second most important source in sector 2 increased by some orders of magnitude since 1990 due to the replacement of CFCs by HFCs. Source category 2G Other product manufacture and use with SF₆ and PFC emissions from electrical equipment and other product use, as well as N₂O emissions from the application in households and hospitals has increased by approximately

factor 2 since 1990. Other source categories in sector 2 are of minor importance with regard to the overall greenhouse gas emissions.

GHG emissions in sector 3 Agriculture are driven by populations of cattle and swine and by fertilizer use. Both factors have been declining, thus leading to a decrease in CH₄ and N₂O emissions until 2004. Subsequently, emissions increased slightly until 2008 and decreased again afterwards mainly due to the evolution of the cattle population.

Total emissions from the source category 5 Waste continuously decrease between 1990 and 2015, with a short increasing phase from 2000 until 2003. The main driver of the decreasing trend is the emission reduction in solid waste disposal, which was reinforced through a change of legislation in 2000 that banned disposal of combustible waste in landfills. Therefore, an increasing amount of municipal solid waste is being incinerated, with emissions reported under source 1A1 Energy industries rather than sector 5 Waste. Altogether, “waste-related” emissions (including emissions from all waste management activities reported in 1 Energy, 3 Agriculture, and 5 Waste) are increasing since 1990 and show a stagnation since 2006 (see Figure 7-3 in chp. 7.1).

The total emissions from sector 6 Other (fire damages) show fluctuations on a very low level. Emissions from sector 6 Other are not accounted for under the Kyoto Protocol and are of minor importance.

Table 2-4 Switzerland's GHG emissions in CO₂ equivalent (kt) by sources and sinks, 1990–2015. The column below on the far right (digits in italic) indicates the percentage change in emissions in 2015 as compared to the base year 1990.

Source and Sink Categories	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
1. Energy	41'846	41'878	42'171	44'006
1A1 Energy industries	2'519	2'643	3'172	3'816
1A2 Manufacturing industries and construction	6'453	6'206	5'928	6'000
1A3 Transport	14'660	14'266	15'930	15'860
1A4 Other sectors	17'632	18'171	16'630	17'879
1A5 Other	220	163	151	139
1B Fugitive emissions from fuels	362	430	359	313
2. Industrial processes and product use	3'585	2'922	3'139	3'795
3. Agriculture	6'780	6'489	6'108	6'078
5. Waste	1'133	950	934	964
6. Other	12	12	13	14
Total (excluding LULUCF)	53'357	52'251	52'365	54'856
4. Land use, land-use change and forestry	-279	-3'676	5'018	-2'234
Total (including LULUCF)	53'078	48'575	57'383	52'622

Source and Sink Categories	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
	CO ₂ equivalent (kt)										%
1. Energy	43'612	41'578	42'943	41'844	43'218	39'162	40'552	41'489	37'464	37'113	-11.3%
1A1 Energy industries	4'032	3'719	3'837	3'674	3'847	3'598	3'641	3'737	3'609	3'279	30.1%
1A2 Manufacturing industries and construction	6'169	5'981	6'025	5'717	5'832	5'389	5'397	5'500	5'108	4'989	-22.7%
1A3 Transport	15'975	16'299	16'650	16'446	16'336	16'155	16'273	16'184	16'075	15'338	4.6%
1A4 Other sectors	17'002	15'163	16'025	15'607	16'785	13'610	14'850	15'697	12'303	13'151	-25.4%
1A5 Other	143	136	131	133	138	125	133	134	139	135	-38.4%
1B Fugitive emissions from fuels	291	280	275	269	282	285	260	239	230	221	-38.9%
2. Industrial processes and product use	3'785	3'848	3'924	3'811	4'022	4'064	4'066	4'096	4'140	3'992	11.3%
3. Agriculture	6'112	6'168	6'273	6'194	6'213	6'159	6'126	6'060	6'150	6'074	-10.4%
5. Waste	965	949	933	915	904	890	869	864	855	846	-25.3%
6. Other	12	14	13	13	12	13	14	14	12	12	2.0%
Total (excluding LULUCF)	54'486	52'556	54'087	52'776	54'370	50'289	51'628	52'523	48'620	48'038	-10.0%
4. Land use, land-use change and forestry	643	-169	-1'024	-1'705	-1'253	-854	-1'260	-1'073	-836	-894	220.7%
Total (including LULUCF)	55'129	52'388	53'063	51'071	53'117	49'436	50'368	51'450	47'784	47'144	-11.2%

The percentage shares of source categories are shown for selected years in Table 2-5, whereas Figure 2-4 to Figure 2-6 are graphical representations of the data in Table 2-4. For the time series of the source categories of sector 1 Energy see chp. 3.

Table 2-5 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (kt) and the contribution of individual source categories (as percentage values) for selected years.

Source and Sink Categories	1990		1995		2000		2005		2010	
	kt CO ₂ eq	%								
1. Energy	41'846	78.4%	41'878	80.1%	42'171	80.5%	44'006	80.2%	43'218	79.5%
1A1 Energy industries	2'519	4.7%	2'643	5.1%	3'172	6.1%	3'816	7.0%	3'847	7.1%
1A2 Manufacturing industries and construction	6'453	12.1%	6'206	11.9%	5'928	11.3%	6'000	10.9%	5'832	10.7%
1A3 Transport	14'660	27.5%	14'266	27.3%	15'930	30.4%	15'860	28.9%	16'336	30.0%
1A4 Other sectors	17'632	33.0%	18'171	34.8%	16'630	31.8%	17'879	32.6%	16'785	30.9%
1A5 Other	220	0.4%	163	0.3%	151	0.3%	139	0.3%	138	0.3%
1B Fugitive emissions from fuels	362	0.7%	430	0.8%	359	0.7%	313	0.6%	282	0.5%
2. Industrial processes and product use	3'585	6.7%	2'922	5.6%	3'139	6.0%	3'795	6.9%	4'022	7.4%
3. Agriculture	6'780	12.7%	6'489	12.4%	6'108	11.7%	6'078	11.1%	6'213	11.4%
5. Waste	1'133	2.1%	950	1.8%	934	1.8%	964	1.8%	904	1.7%
6. Other	12	0.0%	12	0.0%	13	0.0%	14	0.0%	12	0.0%
Total (excluding LULUCF)	53'357	100.0%	52'251	100.0%	52'365	100.0%	54'856	100.0%	54'370	100.0%

Source and Sink Categories	2011		2012		2013		2014		2015	
	kt CO ₂ eq	%								
1. Energy	39'162	77.9%	40'552	78.5%	41'489	79.0%	37'464	77.1%	37'113	77.3%
1A1 Energy industries	3'598	7.2%	3'641	7.1%	3'737	7.1%	3'609	7.4%	3'279	6.8%
1A2 Manufacturing industries and construction	5'389	10.7%	5'397	10.5%	5'500	10.5%	5'108	10.5%	4'989	10.4%
1A3 Transport	16'155	32.1%	16'273	31.5%	16'184	30.8%	16'075	33.1%	15'338	31.9%
1A4 Other sectors	13'610	27.1%	14'850	28.8%	15'697	29.9%	12'303	25.3%	13'151	27.4%
1A5 Other	125	0.2%	133	0.3%	134	0.3%	139	0.3%	135	0.3%
1B Fugitive emissions from fuels	285	0.6%	260	0.5%	239	0.5%	230	0.5%	221	0.5%
2. Industrial processes and product use	4'064	8.1%	4'066	7.9%	4'096	7.8%	4'140	8.5%	3'992	8.3%
3. Agriculture	6'159	12.2%	6'126	11.9%	6'060	11.5%	6'150	12.7%	6'074	12.6%
5. Waste	890	1.8%	869	1.7%	864	1.6%	855	1.8%	846	1.8%
6. Other	13	0.0%	14	0.0%	14	0.0%	12	0.0%	12	0.0%
Total (excluding LULUCF)	50'289	100.0%	51'628	100.0%	52'523	100.0%	48'620	100.0%	48'038	100.0%




Figure 2-4 Switzerland's GHG emissions in CO₂ equivalent (kt) by sectors, 1990–2015 (excluding LULUCF).




Figure 2-5 Relative emission trends (CO₂ eq.) by main source categories (base year 1990 = 100%).

2.3.2 Emission trends in sector 1 Energy

The main source categories within sector 1 Energy – representing the major sources of Switzerland’s GHG emissions – are shown in Figure 2-6. Values are given in Table 2-5.

It is noteworthy that due to Switzerland’s electricity production structure (the most of it is generated by hydroelectric and nuclear power plants in 2015; see SFOE (2016): Table 24), category 1A1 Energy industries plays only a minor role. It does not represent thermal power stations as in many other countries, but primarily waste incineration plants. The following emission trends emerge within the sector 1 Energy:

- Despite differing trends of individual source categories, the overall emissions from the sector 1 Energy remain at a relatively constant level (orange/bold line in Figure 2-6) in the period 1990–2005. From 2005–2015 the combination of effective reduction measures and warm winters (see Figure 2-7) led to decreasing emissions (see details below under 1A4 Other sectors).
- Overall emissions 2015 from source category 1A1 Energy industry are higher than in 1990. The time series shows an increase until 2006 and a decreasing trend from then on, fluctuations being caused by varying combustion activities in the petroleum refinery industry, waste-to-energy, new installations of district heating and weather related forcing of heating activities (see Figure 2-6 and values in Table 2-5).
- The trend for category 1A3 Transport is quite stable with minor fluctuations. They are representing the overall economic development in Switzerland fairly well (gross domestic product) (SFSO 2009a, SFSO 2016e). The slight decrease of transport emissions since 2008 as well as the drop from 2014 to 2015 is largely caused by decreasing fuel tourism (EV 2015a) (see chp. 3.2.9.2.2).

- The trend for source category 1A4 Other sectors reflects the impact of climatic variations on energy demand for heating. The strong correlation with the number of “heating degree days”³ – used as an index of cold weather conditions – is apparent from Figure 2-7, which shows CO₂ emissions from source category 1A4 Fuel combustion – Other sectors (only stationary sources) and the number of heating degree days. The number of heating degree days in 2015 increased compared to 2014. CO₂ emissions caused by fuel combustion in source category 1A4 Other sectors – stationary sources increased simultaneously (see Figure 2-7). In the period 1990–2015, the number of buildings and apartments increased as well as the average floor space per person and workplace. Both phenomena result in an increase in the total area heated by more than one third. Over the same period, however, higher standards were specified for insulation and for combustion equipment efficiency for both new and renovated buildings, compensating for the emissions from the additional area heated.




Figure 2-6 Emission trends (CO₂ eq.) for the source categories in sector 1 Energy. The trend for the entire sector 1 Energy is represented by the bold line with orange squares.

³ Heating degree days: Number of degrees per day calculated as the difference between 20°C (room temperature) and the daily average outdoor temperature for such days where the daily average temperature is below 12°C (e.g. daily outdoor average equals 7°C, then for that day 20–7 = 13). The number of degrees per day are summed up for a year t to yield the heating degree days of year t.




Figure 2-7 Relative trend for CO₂ emissions from 1A4 Fuel Combustion - Other Sectors (stationary sources only) compared with the number of heating degree days.

2.3.3 Emission trends in sector 4 LULUCF

Figure 2-8 illustrates the net emissions and removals of sector 4 LULUCF in Switzerland. Specific values are given in Table 2-4. LULUCF emissions are dominated by biomass dynamics in forests. Except for the years 2000 and 2006, the removals in sector 4 LULUCF were higher than the emissions throughout the period 1990–2015. A strong year to year variation is evident over the whole period. The reasons for the positive value for LULUCF in 2000 (and the small removals in 1990 and 2001) are the storms Vivian (February 1990) and Lothar (December 1999), respectively, which caused great damages in the forest stands and increased harvesting. Harvesting rates in Swiss forests tended to increase since 1991 with maxima in 2006 and 2007 resulting in a net emission in 2006 and a minor removal in 2007. Because harvesting rates started to decline in 2008 due to the international and domestic economic framework conditions, removals from LULUCF increased again since 2007 and showed a moderate year-to-year variability over the last inventory years. The annual contributions of CH₄ and N₂O emissions from LULUCF in the period 1990–2015 are relatively small compared to the CO₂ emissions and removals.




Figure 2-8 Net GHG emissions and removals of sector 4 Land use, land-use change and forestry (LULUCF), 1990–2015 (in kt CO₂ eq). Positive values refer to emissions, negative values refer to removals.

2.4 Emission trends for precursor gases and SO₂

Emission trends for precursor gases (IPCC 2006, Volume 1, Chapter 7) show a very pronounced decline (see Table 2-6 and Figure 2-9). A strict air pollution control policy and the implementation of a large number of emission reduction measures led to a decrease between 56% and 83% in emissions of precursor gases and SO₂ over the period 1990–2015. The main reduction measures were abatement of exhaust emissions from road vehicles and stationary combustion equipment, taxation of solvents and sulphured fuels, and voluntary agreements with the industry (FOEN 2010i, Swiss Confederation 1985, 1997).

Table 2-6 Switzerland's precursor gases and SO₂ emissions (kt), 1990–2015 (without NMVOC from LULUCF).

Precursor gases and SO ₂	1990	1995	2000	2005
	kt			
NO _x	143	118	107	93
CO	798	533	426	331
NMVOC	298	197	142	102
SO ₂	40	26	15	15

Precursor gases and SO ₂	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	kt									
NO _x	90	86	85	79	77	72	72	71	67	63
CO	306	287	276	260	249	227	220	213	195	185
NMVOC	98	94	92	90	88	85	83	82	80	77
SO ₂	14	12	12	10	10	9	9	9	8	7




Figure 2-9 Relative trends for precursor and SO₂ emissions (without NMVOC from LULUCF), 1990–2015 (base year 1990 = 100%).

Sector 1 Energy is by far the largest source of precursor gas emissions (see Table 2-7), with the only exception being NMVOC, where sector 2 Industrial processes and product use is the dominant source (see Figure 2-10). The total shown in Table 2-7 includes NMVOC emissions from LULUCF, which are estimated at almost 100 kt per year according to SAEFL (1996a).

Table 2-7 Precursor and SO₂ emissions (kt) by source, 2015. (NE = not estimated, NO = not occurring)

Sectors	NO _x		CO		NMVOC		SO ₂	
	kt	%	kt	%	kt	%	kt	%
1 Energy	59.23	93.8%	176.98	95.2%	22.20	12.8%	5.89	88.6%
2 IPPU	0.30	0.5%	5.55	3.0%	48.71	28.2%	0.65	9.7%
3 Agriculture	3.15	5.0%	NO	NO	3.90	2.3%	NO	NO
4 LULUCF	0.02	0.0%	0.68	0.4%	95.52	55.3%	NE	NE
5 Waste	0.38	0.6%	2.00	1.1%	2.42	1.4%	0.10	1.5%
6 Other sources	0.08	0.1%	0.68	0.4%	0.11	0.1%	0.01	0.2%
Total	63.17	100.0%	185.89	100.0%	172.86	100.0%	6.64	100.0%

Figure 2-10 shows the relative contributions of the various sectors for each individual gas excluding LULUCF (data from Table 2-7).




Figure 2-10 Relative contributions of individual sectors to precursor and SO₂ emissions in 2015 (without LULUCF).

2.5 Emission trends (Kyoto Protocol)

Relevant emission and removals as accounted for under the Kyoto Protocol are shown in Table 2-8 and Table 2-9, sorted by sectors and gases, respectively. Base year emissions for the second commitment period are reported in Switzerland's second Initial Report (FOEN 2016c) and the update to the report following the review (FOEN 2016d).

The reported total emissions differ from those reported under the UNFCCC, as sector 6 Other – in addition to LULUCF and international bunkers – is not accounted for under the Kyoto Protocol. On the other hand, activities under Article 3, paragraph 3 (Afforestation, Reforestation and Deforestation) and Article 3, paragraph 4 (Forest management, Cropland management, Grazing land management, and Revegetation) are taken into account. Under the activities of Article 3, paragraph 4 of the Kyoto Protocol, Switzerland only accounts for the mandatory activity Forest management.

Table 2-8 Summary of Switzerland's GHG emissions in CO₂ equivalent (kt) as well as emissions and removals under KP-LULUCF 2008–2015, 1990–2015 by sectors. Excluded are emissions and removals from sectors 4 LULUCF, 6 Other and from International bunkers.

Annex A sources	Sector	Base year initial report	1990	1991	1992	1993	1994	1995	1996	1997	1998
		CO ₂ equivalent (kt)									
Annex A sources	1 Energy + indirect CO ₂ from this sector	41'881	41'890	44'325	44'314	42'126	40'997	41'905	42'819	41'851	43'407
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'893	3'950	3'549	3'359	3'020	3'196	3'166	3'016	2'912	3'009
	3 Agriculture	6'804	6'780	6'744	6'629	6'527	6'513	6'489	6'446	6'261	6'218
	5 Waste + indirect CO ₂ from this sector	1'142	1'135	1'050	1'053	1'005	948	951	947	938	925
	Total (Annex A sources)	53'719	53'755	55'668	55'355	52'678	51'654	52'512	53'229	51'962	53'558

Annex A sources	Sector	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
		CO ₂ equivalent (kt)									
Annex A sources	1 Energy + indirect CO ₂ from this sector	43'189	42'188	43'604	41'993	43'214	43'575	44'019	43'626	41'590	42'956
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'063	3'307	3'396	3'419	3'497	3'777	3'909	3'897	3'956	4'033
	3 Agriculture	6'124	6'108	6'170	6'136	6'059	6'036	6'078	6'112	6'168	6'273
	5 Waste + indirect CO ₂ from this sector	914	936	951	975	954	973	965	966	951	934
	Total (Annex A sources)	53'289	52'539	54'121	52'522	53'724	54'362	54'971	54'600	52'665	54'197

KP-LULUCF	Art.3.3	2009–2015										
		2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990			
KP-LULUCF	Art.3.3	Afforestation & reforestation										-21
		Deforestation										95
	Art.3.4	Forest management										-1'693
		Cropland management										NA
		Grazing land management										NA
		Revegetation										NA

Annex A sources	Sector	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
		CO ₂ equivalent (kt)							
Annex A sources	1 Energy + indirect CO ₂ from this sector	41'856	43'229	39'173	40'562	41'500	37'475	37'118	-11%
	2 Industrial processes and product use + indirect CO ₂ from this sector	3'920	4'132	4'173	4'175	4'202	4'246	4'098	4%
	3 Agriculture	6'194	6'213	6'159	6'126	6'060	6'150	6'074	-10%
	5 Waste + indirect CO ₂ from this sector	916	905	892	870	865	856	848	-25%
	Total (Annex A sources)	52'886	54'480	50'397	51'733	52'628	48'727	48'138	-10%

KP-LULUCF	Art.3.3	2009–2015								
		2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990	
KP-LULUCF	Art.3.3	Afforestation & reforestation	-22	-21	-19	-18	-17	-15	-17	
		Deforestation	140	151	152	152	150	149	152.4	
	Art.3.4	Forest management	-2'539	-2'515	-1'184	-2'521	-2'484	-1'078	-2'536	
		Cropland management	NA							
		Grazing land management	NA							
		Revegetation	NA							

Table 2-9 Switzerland's total GHG emissions (excluding 4 LULUCF, 6 Other and International bunkers) and the contribution of individual gases in CO₂ equivalent (kt), 1990–2015 as well as emissions and removals under KP-LULUCF 2008–2015.

Annex A sources	GHG	Base year initial report	1990	1991	1992	1993	1994	1995	1996	1997	1998
		CO ₂ equivalent (kt)									
	CO ₂ + indirect CO ₂	44'521	44'571	46'569	46'370	43'917	42'953	43'686	44'394	43'266	44'810
	CH ₄	6'091	6'102	6'035	5'953	5'843	5'791	5'782	5'743	5'601	5'531
	N ₂ O	2'853	2'829	2'824	2'794	2'730	2'702	2'688	2'685	2'591	2'587
	HFCs	0.0	0.0	1.6	16	33	81	245	296	360	454
	PFCs	117	117	99	81	35	21	17	20	21	24
	SF ₆	137	137	139	141	121	107	93	90	124	153
	NF ₃	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Total (Annex A sources)	53'719	53'755	55'668	55'355	52'678	51'654	52'512	53'229	51'962	53'558

Annex A sources	GHG	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
		CO ₂ equivalent (kt)									
	CO ₂ + indirect CO ₂	44'616	43'782	45'234	43'596	44'794	45'360	45'915	45'496	43'498	44'819
	CH ₄	5'429	5'395	5'435	5'400	5'322	5'290	5'307	5'320	5'300	5'378
	N ₂ O	2'551	2'546	2'561	2'537	2'489	2'446	2'438	2'436	2'460	2'484
	HFCs	528	622	720	798	893	1'014	1'064	1'111	1'186	1'236
	PFCs	26	50	28	33	62	65	44	51	49	58
	SF ₆	140	144	145	158	165	186	203	186	172	222
	NF ₃	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
	Total (Annex A sources)	53'289	52'539	54'121	52'522	53'724	54'362	54'971	54'600	52'665	54'197

KP-LULUCF	GHG	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
		CO ₂ equivalent (kt)							
Art.3.3	CO ₂								
Art.3.3	CH ₄								NO
Art.3.3	N ₂ O								2.2
Art.3.4	CO ₂								-1'702
Art.3.4	CH ₄								2.7
Art.3.4	N ₂ O								6.1

Annex A sources	GHG	2009	2010	2011	2012	2013	2014	2015	2015 vs. 1990
		CO ₂ equivalent (kt)							
	CO ₂ + indirect CO ₂	43'649	45'164	41'102	42'366	43'308	39'377	38'852	-13%
	CH ₄	5'295	5'276	5'217	5'179	5'118	5'120	5'085	-17%
	N ₂ O	2'448	2'495	2'439	2'422	2'384	2'400	2'351	-17%
	HFCs	1'247	1'324	1'406	1'486	1'514	1'527	1'536	see caption
	PFCs	63	64	68	71	52	44	57	-51%
	SF ₆	180	148	160	209	252	259	256	87%
	NF ₃	5.1	8.5	6.2	0.4	0.1	0.4	0.5	NA
	Total (Annex A sources)	52'886	54'480	50'397	51'733	52'628	48'727	48'138	-10%

KP-LULUCF	GHG	2009	2010	2011	2012	2013	2014	2015
		Art.3.3	CO ₂	116	128	131	132	131
Art.3.3	CH ₄	NO	NO	NO	NO	NO	NO	NO
Art.3.3	N ₂ O	2.4	2.4	2.4	2.4	2.4	2.4	2.35
Art.3.4	CO ₂	-2'548	-2'523	-1'196	-2'529	-2'492	-1'086	-2'545
Art.3.4	CH ₄	2.7	2.3	4.1	2.3	2.3	2.5	2.5
Art.3.4	N ₂ O	6.2	6.0	7.1	6.0	6.0	6.2	6.2

3 Energy

3.1 Overview

This chapter provides information on the estimation of the greenhouse gas emissions from the sector 1 Energy. The following source categories are reported:

- 1A Fuel combustion
- 1B Fugitive emissions from fuels

In Switzerland, the sector 1 Energy is the most relevant source of greenhouse gases. The emissions of the period 1990–2015 are depicted in Figure 3-1 and Table 3-1.




Figure 3-1 Switzerland's GHG emissions of sector 1 Energy 1990–2015 in CO₂ equivalent (kt).

Considering total emissions of sector 1 Energy, fluctuations with no trend are observed in the period 1990–2005. From 2006 onwards, a decreasing trend can be identified, again superposed by fluctuations. 2014 and 2015 show the lowest values of the entire period 1990–2015. Four source categories dominate the emissions:

- 1A3 Transport and 1A4 Other sectors are the main sources of the sector 1 Energy.
- 1A1 Energy industries and 1A2 Manufacturing industries and construction are less important.
- 1A5 Other (Military) and 1B Fugitive emissions play only a minor role.

The trends of the individual gases are given in Table 3-1 and Figure 3-2:

- By far the most important gas emitted from sector 1 Energy is CO₂. Fluctuations reflect inter alia the climatic variability in Switzerland (see Figure 2-7 and related comments).
- The decreasing trend of CH₄ emissions since 1990 is the result of improved gas transmission and distribution networks, resulting in substantially lower fugitive emissions, and reduced emissions from gasoline passenger cars due to catalytic converters. Furthermore, improved combustion technologies in 1A4 Other sectors also contribute to the decreasing trend.
- The changes in N₂O emissions can mainly be explained by changes in the emission of road transportation due to changes in emission factors for diesel and gasoline combustion. The first generation of catalytic converters generated N₂O as an unintended by-product in the exhaust gases, leading to an increase in N₂O emissions until 2000. With new converter materials being used, the emission factors are decreasing since 2001 with strongest reduction during 2003 and 2004. For further details, see chp. 3.2.9.2.2.

Table 3-1 GHG emissions of source category 1 Energy by gas in CO₂ equivalent (kt)

Gas	1990	1995	2000	2005
CO ₂ equivalent (kt)				
CO ₂	40'899	40'916	41'331	43'352
CH ₄	652	642	515	420
N ₂ O	295	320	326	234
Sum	41'846	41'878	42'171	44'006

Gas	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2014 to 2015	1990 to 2015
CO ₂ equivalent (kt)											%	
CO ₂	42'984	40'981	42'344	41'259	42'614	38'599	39'992	40'931	36'944	36'595	-1%	-11%
CH ₄	400	374	368	355	367	338	324	312	289	287	-1%	-56%
N ₂ O	228	222	231	231	238	225	237	245	232	231	0%	-22%
Sum	43'612	41'578	42'943	41'844	43'218	39'162	40'552	41'489	37'464	37'113	-1%	-11%




Figure 3-2 Relative trends of the greenhouse gases of sector 1 Energy in the period 1990–2015. The base year 1990 represents 100%.

The following table summarises the emissions of sector 1 Energy in 2015. The table also includes emissions from international bunkers (aviation and marine) as well as CO₂ emissions from biomass burning, which both are not accounted for under the Kyoto Protocol but are included in the reporting tables.

Table 3-2 Summary of sector 1 Energy, emissions in 2015 in kt CO₂ equivalent (Total: rounded values). For full biomass CO₂ emissions see Table 3-20. Note that in reporting table CRF Table10s2 biogene CO₂ emissions from 5C are missing (for an overview of errors of the CRF Reporter see Annex 6).

Sector Energy	CO ₂	CH ₄	N ₂ O	Total
	CO ₂ equivalent (kt)			
1 Energy	36'595	286.5	231.3	37'113
1A Fuel combustion	36'568	92.5	231.2	36'892
1A1 Energy industries	3'251	0.7	26.9	3'279
1A2 Manufacturing industries and construction	4'950	5.0	33.8	4'989
1A3 Transport	15'209	19.2	109.8	15'338
1A4 Other sectors	13'024	67.3	59.6	13'151
1A5b Other (mobile)	134	0.2	1.2	135
1B Fugitive emissions from fuels	27	194.1	0.1	221
International bunkers	4'926	0.6	40.4	4'967
CO ₂ emissions from biomass	6'971	-	-	6'971

In 2015, 51 key source categories are identified in the Swiss greenhouse gas inventory according to level or trend (Table 1-9). Amongst these, 21 belong to the sector 1 Energy. The key categories (according to level and trend) from sector 1 Energy are depicted in Figure 3-3 (Approach 1) and Figure 3-4 (Approach 2).




Figure 3-3 Key categories in the Swiss GHG inventory from sector 1 Energy determined by Approach 1 (L1 = key category according to Approach 1 level in 2015; T1 = key category according to Approach 1 trend 1990–2015).




Figure 3-4 Key categories in the Swiss GHG inventory from sector 1 energy determined by Approach 2 (L2 = key category according to Approach 2 level in 2015; T2 = key category according to Approach 2 trend 1990–2015).

3.2 Source category 1A – Fuel combustion activities

3.2.1 Comparison of the Sectoral Approach with the Reference Approach

Two methods are applied for modelling CO₂ emissions from the sector 1 Energy, the Sectoral Approach and the Reference Approach. For the inventory under the Framework Convention on Climate Change and the Kyoto Protocol the Sectoral Approach is used. The Reference Approach is only used for verification purposes (quality control activity).

Figure 3-5 depicts the two approaches, showing the input data used and the disaggregation of fuel types that ultimately allows for comparing the two approaches.




Figure 3-5 Calculation of Reference and Sectoral Approach. The input data for both approaches stem from the Swiss overall energy statistics (SFOE 2016). While the Reference Approach considers the net import/export balance, the Sectoral Approach considers the fuel consumption. The dark grey arrows depict fuel deductions where occurring. The dashed arrow from the Feedstock use to Other fossil fuels stands for the CO₂ emissions from cracker by-products (originating from feedstock use of LPG and naphtha) which are accounted for under Other fossil fuels. The graphic box of the import of Residual fuel oil is dashed since there is no more import of residual fuel oil.

The Sectoral Approach is based on sectoral energy consumption data from the Swiss overall energy statistics (SFOE 2016) and additional source-specific information. In the Sectoral Approach, fossil fuel consumption statistics are combined with bottom-up data and modelling of fuel consumption. A detailed description of the Sectoral Approach is provided in chp. 3.2.4.

The Reference Approach on the other hand corresponds to a top-down approach based on net quantities of fuel imported into Switzerland as listed in the energy supply statistics of the

Swiss overall energy statistics (SFOE 2016). Apparent consumption (in tonnes) is derived from imports and exports of primary fuels (crude oil, natural gas, coal⁴), secondary fuels (gasoline, diesel oil etc.) and stock changes. For crude oil, a constant value for carbon content and net calorific value is applied for the entire time period, although these properties may vary depending on origin. For solid, gaseous, secondary liquid and other fuels, the same carbon content values and net calorific values are applied as in the Sectoral Approach (see Table 3-9 and Table 3-10, Table 3-12 and Table 3-13 in chp. 3.2.4.2). After the exclusion of feedstocks and non-energy use of fuels (see chp. 3.2.3), the net carbon emissions and the effective CO₂ emissions are calculated for the Reference Approach as shown in the reporting tables 1A(b)–1A(d). The oxidation factor is set to 1 (see chp. 3.2.4.4.1). The Reference Approach covers the CO₂ emissions of all net imported primary fuels and emissions of imported secondary fuels. In 2014, 44% of all liquid fossil fuels sold in Switzerland (without kerosene) were produced in Swiss refineries, in 2015 after closing of one refinery, the share dropped down to 30% (EV 2016). In addition the reporting tables 1.A(b) provide information of the Reference Approach of total biomass use in Switzerland as well including now also the so far missing consumption of so-called other non-fossil fuels (biogenic waste).

All necessary data for calculating the Reference Approach are implemented in the EMIS database and all the data on import, export, bunkers, stock changes, apparent consumption, carbon emission factors, carbon stored and actual emissions are calculated in the EMIS database under the following conditions:

- For the Reference Approach, gas oil and diesel oil are reported together, since the reporting table template structure requires this aggregation. Accordingly, a weighted average NCV is calculated based on values given in Table 3-9. In contrast, marine bunkers consist of diesel oil only and are reported using the country-specific NCV of 43.0 TJ/kt.
- Liechtenstein's liquid fossil fuel consumption is subtracted from the input figures in SFOE (2016), as the Swiss overall energy statistics includes Liechtenstein's liquid fuel consumption (see also chp. 3.2.4).

The differences in energy consumption and CO₂ emissions between Reference and Sectoral Approach are calculated within the EMIS database. For the entire period, they are below 1% and in the range of about 1% for the energy consumption and the CO₂ emissions, respectively, as shown in Table 3-3 and in Figure 3-6. Various effects influence the difference between Reference and Sectoral Approach. On the one hand, energy and carbon contents of crude oil may vary over time. However, no data are available to quantify this effect. On the other hand, the efficiency of Swiss refineries and the market share of secondary fuel imports potentially influence the difference between the Reference and Sectoral Approach. Apparent differences between the Reference Approach and the IEA energy statistics (IEA 2012) are discussed in Annex 4.2.

⁴ Coking coal is included under other bituminous coal.

Table 3-3 Differences in energy consumption and CO₂ emissions between the Reference and the Sectoral Approach. The difference is calculated according to $[(RA-SA)/SA] \cdot 100\%$ with RA = Reference Approach, SA = Sectoral Approach.

	1990	1995	2000	2005
	%			
Energy consumption	0.6	0.8	0.4	0.5
CO ₂ emissions	0.8	0.9	0.8	0.9

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	%									
Energy consumption	0.6	0.5	0.4	0.6	0.6	0.6	0.5	0.4	0.4	0.2
CO ₂ emissions	1.0	0.8	0.9	1.1	1.0	1.1	0.9	0.9	0.9	0.6




Figure 3-6 Time series for the differences between Reference and Sectoral Approach. Numbers are taken from Table 3-3. See caption there for further information.

3.2.2 International bunker fuels (1D)

3.2.2.1 Source category description for 1D

With Switzerland being a landlocked country, international aviation dominates emissions from bunker fuels by far. International navigation is limited to activities on the river Rhine (Basel – Rotterdam) and navigation on Lake Geneva (bordering France) and Lake Constance (bordering Germany and Austria).

Table 3-4 Source category description of International bunkers.

1D	Source	Specification
1D1	International aviation (aviation bunkers)	Bunker fuels include fuel used for international aviation only.
1D2	International navigation (marine bunkers)	Marine bunkers of the Rhine river and crossborder navigation on Lake Geneva and Lake Constance.

3.2.2.2 Methodological issues for 1D

3.2.2.2.1 International aviation / aviation bunkers (1D1)

Following the decision tree of the 2006 IPCC Guidelines (IPCC 2006, Volume 2 Energy, chp. 3 Mobile Combustion, Figure 3.6.1), the emissions from aviation bunkers are calculated with a Tier 3A method because of availability of data on the origin and destination of flights and also on air traffic movements delivered by the Federal Office of Civil Aviation (FOCA).

The Tier 3A method follows standard modelling procedures at the level of single aircraft movements based on detailed movement statistics. For international aviation (aviation bunkers), the flights departing from Switzerland to a destination abroad are selected. The emission factors are country-specific based on measurement and analyses of fuel samples. The activity data of the bunker is summarised in Table 3-5 (see also Table 3-74). Given that detailed information about activity data is available, the resulting fuel consumption is considered complete. In spite of this, there remain small differences between the fuel consumption modelled bottom-up and the total fuel sold (SFOE 2016, FOCA 2016). In 1990, the modelled consumption adds up to 1.01 million tonnes, whereas 1.05 million tonnes of fuel was sold. Such difference of 4% is considered acceptable, because discrepancies up to 10% can easily result from fuelling strategies of airlines (FOCA 2006a). Investigation showed, that airlines are calculating whether it is economically beneficial to refuel at a place with lower fuel price. In order to match the bottom-up calculation with the fuel quantity sold, any occurring difference is attributed to international bunker emissions. The factor between calculated international fuel consumption and adjusted international fuel consumption is used to scale the bunker emissions linearly. For instance in 2015, the bunker fuel consumption and the emissions had to be expanded by the factor 1.020, the correction factor was 0.98 (FOCA 2016). For the more recent years, the modelled and actual total fuel sales are listed in Table 3-5. Table 3-6 provides an overview of total fuel consumption of international aviation (bunker) in energy units.

Table 3-5 Comparison between modelled and actual fuel sales in bunker fuel consumption for aviation

Modelled and actual fuel sales	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	Fuel consumption in t										
Modelled domestic fuel sales	38'754	38'550	43'968	37'627	39'626	39'252	42'047	43'414	42'064	44'462	43'680
Modelled international fuel sales	1'152'614	1'196'731	1'287'062	1'391'656	1'345'919	1'395'428	1'511'279	1'527'522	1'528'863	1'561'678	1'590'013
Total modelled fuel sales (FOCA)	1'191'368	1'235'281	1'331'030	1'429'283	1'385'545	1'434'680	1'553'326	1'570'936	1'570'927	1'606'140	1'633'693
Actual fuel sales (GEST)	1'148'131	1'203'868	1'289'152	1'382'835	1'324'224	1'390'824	1'488'805	1'523'116	1'539'963	1'549'228	1'602'319
Difference between FOCA and GEST	3.8%	2.6%	3.2%	3.4%	4.6%	3.2%	4.3%	3.1%	2.0%	3.7%	2.0%
Correction factor	0.962	0.974	0.968	0.966	0.954	0.969	0.957	0.969	0.980	0.963	0.980

Table 3-6 International bunker fuels (1D1): aviation bunkers. Consumption of kerosene in TJ (Liechtenstein's kerosene consumption is subtracted, see chp. 3.2.4).

1D1 International aviation	1990	1995	2000	2005
	Fuel consumption in TJ			
1D1 International aviation	41'884	49'918	63'687	47'671
1990 = 100%	100%	119%	152%	114%

1D1 International aviation	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	Fuel consumption in TJ									
1D1 International aviation	50'109	53'543	57'844	55'238	58'118	62'211	63'627	64'709	65'006	67'333
1990 = 100%	120%	128%	138%	132%	139%	149%	152%	154%	155%	161%

3.2.2.2 International navigation / navigation bunkers (1D2)

According to the decision tree concerning navigation bunkers (IPCC 2006, Volume 2 Energy, chp. 3 Mobile Combustion, Figure 3.5.1), emissions from international navigation are calculated with a Tier 2 approach for CO₂ (with country-specific carbon contents) and with a Tier 1 approach for CH₄ and N₂O using IPCC default emission factors. On the river Rhine and on the lakes of Geneva and Konstanz, some of the boats cross the border and go abroad (Germany, France). Fuels bought in Switzerland will therefore become bunker fuel. Accordingly, the amount of bunker diesel oils is reported as a memo item "International bunker / navigation".

- Only diesel oil is relevant for navigation on the river Rhine. Since there is an exemption from fuel taxation, activity data on marine river bunkers on the Rhine are well documented by the customs administration for the years 1997–2015 (FCA 2016a).
- For navigation on two border lakes (Lake Constance, Lake Geneva), bunker fuel consumption was reported in INFRAS (2011a) after having performed surveys among the shipping companies involved. Activity data of these bunkers is summarised in Table 3-7. Data from 1995–2012 have been provided by the three navigation companies concerned as documented in INFRAS 2011a. For older data, proxies such as passenger data on a national basis had to be consulted. As marine lake bunkers provide only a minor share of the total international navigation (between 6% for the year 1990 and 23% for 2015) this approach is justified. The emission factor for CO₂ is country-specific and in accordance with Table 3-12.

Table 3-7 International bunker fuels (1D2): Navigation. Consumption of diesel oil in TJ.

1D2 International navigation	1990	1995	2000	2005
	Fuel consumption in TJ			
1D2 International navigation	813	756	526	500
1990 = 100%	100%	93%	65%	62%

1D2 International navigation	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	Fuel consumption in TJ									
1D2 International navigation	462	477	459	435	472	421	379	356	311	335
1990 = 100%	57%	59%	57%	54%	58%	52%	47%	44%	38%	41%

3.2.2.3 Uncertainties and time-series consistency for 1D

International aviation: see general remarks in chp. 3.2.4.7.

Consistency: Time series of 1D are all considered consistent.

3.2.2.4 Category-specific QA/QC and verification for 1D

The general QA/QC procedures are described in chp. 1.2.3. Furthermore QA/QC procedures conducted for all 1A source categories are listed in chp. 3.2.4.8.

3.2.2.5 Category-specific recalculations for 1D

Based on a question raised by the energy expert during the review 2016, there are no more CH₄ emissions for cruise activities. This implies higher NMVOC emissions for those activities. All years 1990–2014 have been recalculated.

Small recalculation due to change in the NCV of diesel used in international navigation to the same as for other diesel processes. Therefore small changes occur in international navigation and in fuel tourism and statistical difference for diesel (which is integrated in 1A3biii).

3.2.2.6 Category-specific planned improvements for 1D

No category-specific improvements are planned.

3.2.3 Feedstocks and non-energy use of fuels

The Swiss overall energy statistics (SFOE 2016) reports feedstocks and non-energy fuel use on an aggregated level only. Some disaggregation is provided by the petroleum balance of the annual report of the Swiss petroleum association (EV 2016). To complement this source, bottom-up data from annual monitoring reports of the Swiss emissions trading scheme (ETS) and from surveys of individual companies are used to provide a detailed breakdown into specific petroleum products and coal type. For submission 2015, a new and more differentiated breakdown of feedstocks and non-energy use of fuels was developed. The reassessment of feedstocks and other non-energy use of fuels is documented in an internal documentation (FOEN 2015g).

- Feedstocks and non-energy use of fuels is reported in reporting tables 1.A(d) and differentiated in the following fuel types:
- Liquefied petroleum gas and naphtha are exclusively used in one single Swiss plant as feedstocks in the thermal cracking process for the production of ammonia and ethylene (see source categories 2B1 and 2B8b under 4.3.2.1 and 4.3.2.4, respectively). Accordingly, activity data for liquefied petroleum gas and naphtha are confidential and included in fuel type Other oil in reporting table 1.A(d).
- Bitumen is the most important petroleum product, which is used as a feedstock in Switzerland. It is mainly used for road paving with asphalt and to a lower extent in asphalt roofing (see source category 2D3 under 4.5.2.2).
- Lubricants are used in a variety of processes, including the blending with motorcycle fuel. Use of lubricants is considered partly emissive (see source category 2D1 under 4.5.2.1). According to the 2006 IPCC guidelines (IPCC 2006), 20% of lubricants are oxidized during use (ODU).

- Petroleum coke is used as a feedstock by two consumers only, i.e. for the production of silicium carbide and graphite as well as of anodes in primary aluminium production (up to 2006) in source categories 2B5 and 2C3, respectively (see chp. 4.3.2.3 and 4.4.2.2). Apart from bottom-up information from these two consumers, top-down information is provided by the Swiss petroleum association (EV 2016). Accordingly, activity data are confidential and included in fuel type Other oil in reporting table 1.A(d).
- Paraffin waxes for non-energy use are reported under Other oil, since there is no separate category for paraffin waxes in reporting table 1.A(d). The information used stems from the statistics of the Swiss petroleum association (EV 2016). Use of paraffin waxes is considered partly emissive (see source category 2D2 under 4.5.2.1). According to the 2006 IPCC guidelines (IPCC 2006), 20% of paraffin waxes are oxidized during use (ODU).
- Other oil comprises all other unspecified petroleum products for non-energy use. Please note that the net consumption of non-energy use of fuels reported in Swiss overall energy statistics includes sulphur produced by the refineries as well. This amount of sulphur is now subtracted resulting in lower fuel quantities for non-energy use of other oil for the entire time series.
- Other bituminous coal (anthracite) is also used as feedstock in the Swiss production plant for silicium carbide and graphite in source category 2B5 (chp. 4.3.2.3). Accordingly, activity data for other bituminous coal are confidential and thus denoted as “C” in reporting tables 1.A(d). Please note that in column “Carbon stored” of the reporting table 1.A(b) notation key “NA” is given since the template cannot cope with notation key “C”. After consulting the responsible for the Swiss overall energy statistics the feedstock use of other bituminous coal has to be included in its stock changes.

Table 3-8 This table is only available in the confidential version of this chapter. It provides a complete time series of the fuel quantity, carbon excluded and the reported CO₂ emissions from feedstocks and non-energy use of fuels.

Category-specific recalculations

The net consumption of non-energy use of fuels reported in Swiss overall energy statistics includes sulphur produced by the refineries as well. This amount of sulphur is now subtracted resulting in lower fuel quantities for non-energy use of other oil for the entire time series reported in CRF table 1.A(d). See also chp. 3.2.4.9.

3.2.4 Country-specific issues of 1A Fuel combustion

3.2.4.1 System boundaries: Differences between UNFCCC and CLRTAP reporting

Switzerland reports its greenhouse gas emissions according to the requirements of the UNFCCC as well as other air pollutants according to the requirements of the CLRTAP. The nomenclature for both reportings is (almost) the same (NFR), but there are differences concerning the system boundaries. Under the UNFCCC, the national total for assessing compliance is based on fuel sold within the national territory, whereas under the CLRTAP, the national total for assessing compliance is based on fuel used within the territory. Thus, fuel sold in Switzerland but consumed abroad (“fuel tourism”) is accounted for in Switzerland’s GHG inventory, but not in the reporting under the CLRTAP. The difference

between the two approaches amounts to several percent, with considerable variation from year to year due to fluctuating fuel price differences between Switzerland and its neighbouring countries.

Also emissions from civil aviation are differentiated under the UNFCCC and the CLRTAP: Only emissions from domestic flights are accounted for in the GHG inventory, while emissions from international flights are reported as memo items. For the reporting under the CLRTAP, landing and takeoff (LTO) emissions of domestic and international flights are accounted for, while emissions of international and domestic cruise flights are reported under memo items only (see Figure 3-7).

Differences between reporting under CLRTAP and UNFCCC concerning the accounting to the national total			CLRTAP / NFR-Templates			UNFCCC / CRF-Tables	
			accounted to				
			National total	National total for compliance	Memo item	National total	Bunker 1 D
Road transportation 1 A 3 b	Fuel sold in 1 A 3 b	Fuel used 1 A 3 b i-vii	Yes	Yes	Yes	Yes	No
		Fuel tourism and statistical difference 1 A 3 b viii	Yes	No	No	Yes	No
Aviation 1 A 3 a	Civil/Domestic aviation	Landing and Take-Off (LTO)	Yes	Yes	No	Yes	No
		Cruise	No	No	Yes	Yes	No
	International aviation	Landing and Take-Off (LTO)	Yes	Yes	No	No	Yes
		Cruise	No	No	Yes	No	Yes

Figure 3-7 Accounting rules for emissions from 1A3a Civil aviation and 1A3b Road transportation for CLRTAP and UNFCCC

3.2.4.2 Net calorific values (NCV)

Table 3-9 summarizes the net calorific values (NCV) which are used in order to convert from energy amounts in tonnes into energy quantities in gigajoules (GJ).

- For gasoline, jet kerosene, diesel oil and gas oil, values for 1998 and 2013 are based on measurements. Constant values are used for the period 1990 to 1998 and from 2013 onwards.
- For residual fuel oil measurements for 1998 are available.
- For liquefied petroleum gas, petroleum coke, other bituminous coal, lignite and wood, NCVs are given by Swiss Federal Office for Energy (SFOE 2016, 2016b) and partly based on measurements from the cement industry (Cemsuisse 2010a).
- NCV of natural gas is annually reported by the Swiss Gas and Water Industry Association (SGWA), see Table 3-10.

More detailed explanations including information about the origin of the NCV for individual energy sources are given below.

Table 3-9 Net calorific values of fuels (NCV) 1990–1998 and from 2013 onwards. For years between 1998 and 2013, the NCVs are linearly interpolated. Natural gas see Table 3-10. Data source SGWA stands for annually updated reports of the Swiss Gas and Water Industry Association, latest report stems from SGWA (2016).

Net calorific values (NCV)		1990-1998	2013-2015
Fuel	Data Sources	NCV [GJ/t]	NCV [GJ/t]
Gasoline	EMPA (1999), SFOE/FOEN (2014)	42.5	42.6
Jet kerosene	EMPA (1999), SFOE/FOEN (2014)	43.0	43.2
Diesel oil	EMPA (1999), SFOE/FOEN (2014)	42.8	43.0
Gas oil	EMPA (1999), SFOE/FOEN (2014)	42.6	42.9
Residual fuel oil	EMPA (1999)	41.2	41.2
Liquefied petroleum gas	SFOE (2016)	46.0	46.0
Petroleum coke	SFOE (2016), Cemsuisse (2010a)	35.0	31.8
Other bituminous coal	SFOE (2016), Cemsuisse (2010a)	28.1	25.5
Lignite	SFOE (2016), Cemsuisse (2010a)	20.1	23.6
Natural gas	SGWA	<i>see table below</i>	
Biofuel	Data Sources	NCV [GJ/t]	NCV [GJ/t]
Biodiesel	assumed equal to diesel oil	42.8	43.0
Bioethanol	assumed equal to gasoline	42.5	42.6
Biogas	assumed equal to natural gas	<i>see table below</i>	
Wood	SFOE (2016b)	9.4-10.4	9.4-10.4

Gasoline, jet kerosene, diesel oil and gas oil

The net calorific values for gasoline, jet kerosene, diesel oil and gas oil are provided by a national measurement campaign. The campaign was realized by the EMPA (Swiss Federal Laboratories for Materials Science and Technology) in 1998 for the first time (EMPA 1999). Previous data are not available. The values for 1990–1998 are therefore assumed to be constant at the 1998 levels. An updated study, commissioned by the Federal Office for the Environment (FOEN) and the Federal Office for Energy (SFOE) was conducted in 2013 (SFOE/FOEN 2014). The study is based on a representative sample covering summer and winter fuel qualities from the main import streams. The sampling started in July 2013 for a duration of six months. Samples were taken fortnightly from nine different sites (large-scale storage facilities and the two Swiss refineries) and analysed for carbon content and calorific value amongst other. These updated values are used from 2013 onwards, while the NCVs 1999 – 2012 are linearly interpolated between the measured values of 1998 and 2013.

Residual fuel oil

Residual fuel oil plays only a minor role in energy supply. Therefore, this fuel type was not analysed in the most recent measurement campaign in 2013. Thus, respective NCVs refer to the measurement campaign by EMPA (1999) in 1998. The NCV for residual fuel oil is assumed to be constant for the entire time series. The same approach is applied for the CO₂ emission factor (see Table 3-12).

Liquefied petroleum gas

The net calorific value (NCV) attributed to liquefied petroleum gas is taken from the Swiss overall energy statistics (SFOE 2016)⁵ and is therefore country-specific.

Petroleum coke, other bituminous coal, lignite

NCVs of petroleum coke, other bituminous coal and lignite are based on data from the SFOE and on measurements of samples taken from Switzerland's cement plants. Cement plants are the largest consumers of these fuels in Switzerland. The samples from the individual plants were compiled over nine months in 2009 and analysed for calorific value by an independent analytical laboratory. The original data is collected in an internal documentation provided by the Swiss Association of the Swiss Cement Industry – Cemsuisse (Cemsuisse 2010a). For each fuel type, the measurements from the individual plants were weighted according to the relative consumption of each plant. Between 1999 and 2010 the values are linearly interpolated (see SFOE 2016, p. 61). This approach is also used in order to derive CO₂ emission factors for the three fuels (see chp. 3.2.4.3.3).

Natural gas

The net calorific value of natural gas (and also the CO₂ emission factor of natural gas, see Table 3-13) is calculated based on measurements of gas properties and corresponding import shares of individual gas import stations. Measurements of gas properties are available on an annual basis since 2009 and for selected years before. Import shares are available for 1991, 1995, 2000, 2005, 2007 and from 2009 onwards on an annual basis. Estimated import shares for the years 1991, 1995 and 2000 are taken from Quantis (2014). This study focused on gas imports of the Swiss gas grid for the years 1991, 1995 and 2000. Missing values for the years in between are interpolated. The calculation procedure is documented in FOEN (2016h).

⁵ It is assumed that LPG consists of 50% propane and 50% butane.

Table 3-10 Net calorific values of natural gas and biogas for selected years. Years in-between are linearly interpolated. Data source annual reports of the Swiss Gas and Water Industry Association SGWA. Spreadsheet to determine national averages: FOEN 2016h.

Year	NCV [GJ/t]
1990	46.5
1991	46.5
1995	47.5
2000	47.2
2005	46.6
2007	46.3
2009	46.4
2010	46.3
2011	46.1
2012	45.8
2013	45.7
2014	45.7
2015	46.6

Wood

The net calorific value of wood depends on the type of wood fuel (for e.g. log wood, wood chips, pellets) and are based on the Swiss wood energy statistics (SFOE 2016b). Table 3-9 illustrates the range of the NCV for all wood fuel types.

Biofuels

Regarding the small amount of biofuels used in Switzerland, the NCVs are assumed to be equal to the corresponding values of the fossil fuels substituted (i.e. biodiesel – diesel oil, bioethanol – gasoline, biogas – natural gas).

3.2.4.3 Swiss energy model and final energy consumption

3.2.4.3.1 Swiss overall energy statistics

The fundamental data on final energy consumption is provided by the Swiss overall energy statistics (SFOE 2016). However, since Switzerland and Liechtenstein form a customs and monetary union governed by a customs treaty, data regarding liquid fuels in the Swiss overall energy statistics also cover liquid fuel consumption in Liechtenstein. In order to calculate the correct Swiss fuel consumption, Liechtenstein's liquid fossil fuel consumption, given by Liechtenstein's energy statistics (OS 2016), is subtracted from the numbers provided by the Swiss overall energy statistics. In all years of the reporting period the sum of liquid fossil fuels used in Liechtenstein was less than half a percent of the Swiss consumption.

The energy-related activity data in the energy model and thus in the GHG inventory correspond to the energy balance provided in the Swiss overall energy statistics (SFOE 2016). The energy statistics are updated annually and contain all relevant information about primary and final energy consumption. This includes annual aggregated consumption data

for various fuels and main consumers such as households, transport, energy industries, industry, and services (see energy balance in Annex 4).

The main data sources of the Swiss overall energy statistics are:

- The Swiss organisation for the compulsory stockpiling of oil products (Carbura) and the Swiss petroleum association (EV) for data on import, export, sales, stocks of oil products and for processing of crude oil in refineries.
- Annual import data for natural gas from the Swiss gas industry association (VSG).
- Annual import data for petroleum products and coal from the Federal Customs Administration (FCA).
- Data provided by industry associations (GVS, SGWA, Cemsuisse, VSG, VSTB etc.).
- Swiss renewable energy statistics (SFOE).
- Swiss wood energy statistics (SFOE)
- Swiss statistics on combined heat and power generation (SFOE)

As can be seen in Figure 3-8, fossil fuels amount to slightly more than half of primary energy consumption. The main end-users of fossil fuels are the transport and the housing sector, as electricity generation is predominantly based on hydro- and nuclear power stations. The most recent energy balance is given in Annex 4.




Figure 3-8 Switzerland's primary energy consumption in 2015 by fuels (see corresponding data in SFOE 2016).

Table 3-11 shows primary energy consumption excluding nuclear fuel and water power. Liquid fossil fuel consumption changed only little since 1990. This is the combined effect of a

marked increase of the consumption in the transport sector and a substantial decrease of gas oil use in the residential and industry sector. Natural gas consumption increased since 1990, compensating to some extent the decreasing use of gas oil.

Table 3-11 Switzerland's energy consumption in 1990–2015 by fuel type. Only those fuels are shown that are implemented in the EMIS database (no water or nuclear power), The numbers are based on the fuels sold principle, thus they include consumption from fuel tourism, all fuels sold for domestic and international aviation as well as liquid fuels consumed in Liechtenstein.

Year	Gasoline	Kerosene	Diesel	Gas oil	Residual fuel oil	Refinery gas & LPG	Petroleum coke	Solid fuels	Natural gas	Other fuels	Bio fuels	Total
	TJ	TJ	TJ	TJ	TJ	TJ	TJ	TJ	TJ	TJ	TJ	TJ
1990	155'785	48'067	47'557	218'510	23'342	8'890	1'400	14'901	68'599	19'160	46'684	652'895
1991	162'225	46'562	48'154	238'602	23'590	12'437	980	12'662	76'902	18'596	48'662	689'372
1992	168'100	49'099	46'706	236'809	24'170	11'492	315	8'758	80'808	19'009	47'562	692'828
1993	155'897	50'776	44'978	225'920	17'165	12'388	1'120	7'342	84'758	19'158	47'837	667'338
1994	156'087	52'109	47'748	207'141	17'860	13'455	1'470	7'432	83'587	19'154	45'785	651'828
1995	151'290	54'947	48'604	217'523	17'278	12'756	1'260	7'962	92'123	19'687	47'772	671'204
1996	155'209	56'753	45'597	226'289	15'097	13'939	1'015	5'956	99'710	20'584	51'240	691'388
1997	161'171	58'774	47'385	212'223	12'581	14'236	280	4'590	96'260	21'655	48'162	677'317
1998	162'477	61'268	49'209	222'407	15'882	15'259	455	3'810	99'065	23'803	49'717	703'352
1999	168'025	65'224	52'184	212'349	11'058	15'805	521	3'875	102'588	24'403	50'421	706'454
2000	168'165	68'019	55'677	196'137	7'923	13'649	551	5'970	101'970	26'536	50'087	694'684
2001	163'543	64'150	56'709	213'089	9'942	14'069	410	6'073	106'132	27'068	53'410	714'596
2002	160'375	59'335	58'721	196'655	6'446	15'584	679	5'325	104'170	27'877	53'010	688'178
2003	159'636	53'358	62'251	208'040	7'061	13'642	202	5'713	110'116	27'643	55'456	703'118
2004	156'812	50'350	66'893	203'370	7'561	16'429	1'819	5'420	113'615	28'845	56'345	707'460
2005	152'062	50'994	73'065	205'729	5'805	16'432	2'906	6'040	116'646	29'236	58'416	717'331
2006	147'436	53'443	79'063	195'926	6'419	18'578	3'324	6'517	113'412	31'233	61'381	716'732
2007	146'012	57'010	84'885	171'313	5'179	15'587	2'830	7'296	110'395	30'015	60'335	690'859
2008	142'801	60'967	93'143	178'833	4'606	16'288	3'516	6'562	117'589	30'854	63'979	719'138
2009	138'968	58'471	94'569	173'219	3'575	16'301	3'254	6'193	112'807	29'811	64'005	701'172
2010	134'043	61'397	98'247	182'305	3'027	15'463	3'498	6'208	126'013	31'185	68'519	729'905
2011	128'856	65'438	100'876	143'760	2'292	14'856	2'957	5'842	111'774	30'882	64'875	672'409
2012	124'301	67'021	106'996	154'448	2'780	12'247	3'148	5'269	122'521	31'145	70'773	700'651
2013	118'634	68'068	111'824	162'532	1'959	15'053	2'735	5'667	129'027	30'925	74'346	720'769
2014	113'875	68'541	114'688	122'704	1'701	14'473	3'148	5'904	111'770	31'320	70'021	658'145
2015	105'592	70'788	113'161	129'159	892	9'822	1'145	5'406	119'420	32'084	73'706	661'174

3.2.4.3.2 Energy model – Conceptual overview

For the elaboration of the greenhouse gas and air pollutants inventories, information about energy consumption is needed at a much more detailed level than provided by the Swiss overall energy statistics (SFOE 2016). Activity data in sector 1 Energy are therefore calculated and disaggregated by the Swiss energy model, which is an integral part of the emission database EMIS. The model is developed and updated annually by the Swiss Federal Office for the Environment (FOEN). It relies on the Swiss overall energy statistics and is complemented with further data sources, e.g. Liechtenstein's liquid fuel sales (OS 2016), the Swiss renewable energy statistics (SFOE 2016a), the energy consumption statistics in the industry and services sectors (SFOE 2016d) as well as additional information from the industry and the Swiss wood energy statistics (SFOE 2016b).

The Swiss overall energy statistics are not only the main data input into the energy model, but also serve as calibration and quality control instrument: The total energy consumption given by the Swiss overall energy statistics has to be equal to the sum of the disaggregated

activity data of all source categories within the energy sector (including memo items/bunker). Differences are explicitly taken into account as “statistical differences” (see chp. 3.2.4.1).

As shown in Figure 3-9, the energy model consists of several sub-models, such as the industry model, the civil aviation model, the road transportation model, the non-road transportation model, and the energy model for wood combustion. A brief overview of each of these models is given below. However, depending on the scope of these sub-models, they are either described in the corresponding source category chapter or in an overarching chapter preceding the detailed description of the individual source categories. In chp. 3.2.4.3.3, the resulting sectoral disaggregation is shown separately for each fuel type.




Figure 3-9 Overview of Switzerland's energy model. In the abbreviation SFOE 201X the “X” refers to the latest edition of the Swiss overall energy statistics.

Industry model (Details are given in chp. 3.2.6.2.1)

In order to produce consistent time-series, the industry model is a composite of the energy consumption statistics in the industry and services sectors (SFOE 2016d), which is based on a comprehensive annual survey, and a bottom-up industry model (Prognos 2013), which is periodically calibrated to the Swiss overall energy statistics. The resulting industry model provides a split of energy consumption by source category and fuel type. Further

disaggregation is then achieved by using plant-level industry data for specific processes, as far as available.

Civil aviation model (Details are given in chp. 3.2.9.2.1)

The civil aviation model is developed and updated by the Federal Office for Civil Aviation FOCA. It aggregates single aircraft movements according to detailed movement statistics of the Swiss airports. Differentiation of domestic and international aviation is based on the information on departure and destination of each flight in the movement database.

Road transportation model (Details are given in chp. 3.2.9.2.2)

The road transportation model is a territorial model, accounting for traffic on Swiss territory only. The model is based on detailed vehicle stock data (from the vehicle registration database of the Federal Roads Office FEDRO), mileage per vehicle category differentiated into different driving patterns and specific consumption and emission factors. The difference between fuel sales and the territorial model (road and non-road models combined) is reported under fuel tourism and statistical difference.

Non-road transportation model (Details are given in chp. 3.2.4.5.1)

The non-road transportation model covers all remaining mobile sources, i.e. industrial vehicles, construction machinery, agricultural and forestry machinery, gardening machinery as well as railways, navigation and military vehicles (except for military aviation, which is considered separately (see chp. 3.2.11.2.1)). The model combines vehicle numbers, their operation hours, engine power, and load factors to derive specific fuel consumption, emission factors and resulting emissions. Data stem from surveys among producers, various user associations, and the national database of non-road vehicles run by FEDRO.

Energy model for wood combustion (Details are given in chp. 3.2.4.5.2)

Based on the Swiss wood energy statistics (SFOE 2016b), total wood consumption is disaggregated into source categories (public electricity and heat production, industry, commercial/institutional, residential, agriculture/forestry/fisheries) and into 24 different combustion installations (ranging from open fireplaces to large-scale automatic boiler or heat and power plants). Where available, industry data on wood combustion is taken into account to allocate parts of the wood consumption as given by the Swiss wood energy statistics to a specific source category.

3.2.4.3.3 Disaggregation of the energy consumption by source category and fuel types

The energy model as outlined above disaggregates total energy consumption as provided by the Swiss overall energy statistics (SFOE 2016) into the relevant source categories 1A1-1A5. For each fuel type, the disaggregation process of the energy model as shown schematically

in Figure 3-9, the interaction between the different sub-models and additional data sources are visualized separately in Figure 3-11 to Figure 3-19.




Figure 3-10 Switzerland's energy consumption 1990–2015 by source categories 1A1-1A5 as based on the Swiss energy model. In the same period population increased by about 20%, industrial production by almost 70% and the motor vehicle fleet by 50% (SFOE 2016, table 43b).

Starting from the total energy consumption from the Swiss energy statistics for each fuel type, the energy is assigned to the relevant source categories based on the various sub-models of the energy model, mentioned in chp. 3.2.4.3.2 above. In addition the following assignments are considered as well.

Within source categories 1A4a and 1A4b, the amount of gas oil and natural gas used for co-generation in turbines and engines is derived from a model of stationary engines developed by Eicher + Pauli (Kaufmann 2015) for the statistics on combined heat and power generation (SFOE 2016c). The residual energy is then assigned to boilers which are not further specified.

For source category 1A4c Other sectors – Agriculture/forestry/fishing, specific bottom-up industry information is available for grass drying. Its fuel consumption is determined by the Swiss association of grass drying plants (VSTB) and is subtracted from the total fuel consumption of 1A2.

In order to report all energy consumption, the statistical differences as reported in the Swiss overall energy statistics are allocated to source category 1A4a Other sectors – Commercial/institutional (stationary combustion) and 1A3b viii Fuel tourism and statistical differences.




Figure 3-11 Schematic disaggregation of 1A Fuel consumption for gasoline and diesel oil. Marine bunker fuel consumption is based on the national customs statistics (see chp. 3.2.2.2.2 on memo items)




Figure 3-12 Schematic disaggregation of 1A Fuel consumption for kerosene. Fuel consumption for military aircraft is provided by the Federal Department of Defence, Civil Protection and Sport. The differentiation between domestic and international aviation as well as between CR and LTO is provided by the civil aviation model (see chp. 3.2.2.2.1)




Figure 3-13 Schematic disaggregation of 1A Fuel consumption for gas oil. The Swiss overall energy statistics provide gas oil use for energy conversion and the amount thereof being used for district heating. Based on this information, gas oil use is split into 1A1a i Electricity generation and 1A1a iii Heat plants. According to the non-road model, a small amount of gas oil is consumed in source category 1A3d navigation (steam-powered vessels).




Figure 3-14 Schematic disaggregation of 1A Fuel consumption for residual fuel oil. The Swiss overall energy statistics report residual fuel oil use in energy conversion and the amount thereof consumed in electricity production (one single fossil fuel power station, operational from 1985 to 1994), district heating, and in petroleum refineries. Based on this information, residual fuel oil use in Energy industries is split into 1A1a i Electricity generation, 1A1a iii Heat plants and 1A1b Petroleum refining.




Figure 3-15 Schematic disaggregation of 1A Fuel consumption for liquefied petroleum gas (LPG).




Figure 3-16 Schematic disaggregation of 1A Fuel consumption for petroleum coke.




Figure 3-17 Schematic disaggregation of 1A Fuel consumption for lignite and bituminous coal. The Swiss overall energy statistics provide bituminous coal use for energy conversion and the amount thereof being used for district heating. Based on this information, use of bituminous coal in energy industries is split into 1A1a i Electricity generation and 1A1a iii Heat plants up to 1995. Coal consumption for Public electricity and heat production ceased thereafter.




Figure 3-18 Schematic disaggregation of 1A Fuel consumption for natural gas. The Swiss overall energy statistics (SFOE 2016) provide gas use in the transformation sector (energy conversion and distribution losses). Distribution losses as estimated by the Swiss Gas and Water Industry Association SVGW are subtracted and reported under source category 1B2 Fugitive emissions from fuels. The remaining fuel consumption for natural gas is reported under 1A1a Public electricity and heat production.




Figure 3-19 Schematic disaggregation of 1A Fuel consumption for wood. For a detailed description of the energy model for wood combustion see chp. 3.2.4.5.2.

3.2.4.4 Emission factors of 1A Fuel combustion

3.2.4.4.1 Oxidation factor for 1A Fuel combustion

For the emission calculation, an oxidation factor of 100% is assumed for all fossil fuel combustion processes, since the technical standards for combustion installations in Switzerland are high and the small fraction of originally non-oxidised carbon retained in ash, particulates or soot is likely to be oxidized later. This is consistent with the 2006 IPCC Guidelines and the EU and Swiss guidelines for the Emissions Trading Scheme (ETS), where also a default oxidation factor of 100% was applied.

Because an oxidation factor of 100% is assumed, indirect CO₂ emissions from CO and NMVOC are implicitly reported as direct CO₂ emissions in sector 1A Energy and no indirect emissions are reported for sector 1A in chp. 9.

3.2.4.4.2 CO₂ Emission factors for 1A Fuel combustion

General CO₂ emission factors

The CO₂ emission factors applied for the time series 1990–2015 are given in Table 3-12. Detailed information regarding the underlying data and assumptions are provided in chp. 3.2.4.2 Net calorific values (NCV), since in most cases, NCVs and carbon content were determined jointly.

Table 3-12 CO₂ emission factors 1990–1998 and years from 2013 onwards. For years between 1998 and 2013, the factors are linearly interpolated. Data source SGWA stands for annually updated reports of the Swiss Gas and Water Industry Association.

CO ₂ emission factors			1990-1998	2013-2015
Fossil fuel	CS/D	Data sources	t CO ₂ / TJ	t CO ₂ / TJ
Gasoline	CS	EMPA (1999), SFOE/FOEN (2014)	73.9	73.8
Jet kerosene	CS	EMPA (1999), SFOE/FOEN (2014)	73.2	72.8
Diesel oil	CS	EMPA (1999), SFOE/FOEN (2014)	73.6	73.3
Gas oil	CS	EMPA (1999), SFOE/FOEN (2014)	73.7	73.7
Residual fuel oil	CS	EMPA (1999)	77.0	77.0
Liquefied petroleum gas	CS	FOEN (2015d)	65.5	65.5
Petroleum coke	CS	Cemsuisse (2010a)	91.4	91.4
Other bituminous coal	CS	Cemsuisse (2010a)	92.7	92.7
Lignite	CS	Cemsuisse (2010a)	96.1	96.1
Natural gas	CS	SGWA	<i>see table below</i>	
Biofuel	CS/D	Data sources		
Biodiesel	CS	assumed equal to diesel oil	73.6	73.3
Bioethanol	CS	assumed equal to gasoline	73.9	73.8
Biogas	CS	assumed equal to natural gas	<i>see table below</i>	
Wood	CS	Cemsuisse (2010a)	99.9	99.9

CO₂ emission factors for natural gas

Table 3-13 Time series of CO₂ emission factors of natural gas. SGWA refers to annual updates of properties of natural gas that are provided by the Swiss Gas and Water Industry Association.

CO ₂ emission factors			1990	1995	2000	2005
Fuel	CS/D	Data sources	t CO ₂ / TJ			
Natural gas/Biogas	CS	SGWA	56.1	55.7	56.2	56.4

CO ₂ emission factors			2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Fuel	CS/D	Data sources	t CO ₂ / TJ									
Natural gas/Biogas	CS	SGWA	56.5	56.5	56.5	56.5	56.5	56.6	56.5	56.4	56.5	56.4

CO₂ emission factors for wood

The CO₂ emission factor for wood combustion activities is taken from Cemsuisse (2010a).

3.2.4.4.3 CH₄ Emission factors for 1A Fuel combustion

General CH₄ emission factors

An overview of the general CH₄ emission factors is given in Table 3-14. These emission factors are used for most stationary combustion processes (exceptions are discussed in the detailed sectoral chapters where they occur). For stationary combustion, mainly IPCC default emission factors are used for the entire time period. For wood, country-specific factors are used. Details are given below in Table 3-15. CH₄ emission factors related to transport activities (aviation, road and non-road transportation) are category specific and given in the corresponding chapters.

Table 3-14 CH₄ emission factors for stationary combustion for the whole time period.

CH₄ emission factors			1990-2015
Fuel	CS/D	Data sources	g CH₄ / GJ
Gas oil	D	IPCC (2006)	3
Residual fuel oil	D	IPCC (2006)	3
Liquefied petroleum gas	D	IPCC (2006)	1
Petroleum coke	D	IPCC (2006)	3
Other bituminous coal	D	IPCC (2006)	10
Lignite	D	IPCC (2006)	10
Natural gas	D	IPCC (2006)	1
Biofuel	CS/D	Data Sources	
Biogas	D	IPCC (2006)	1
Wood	CS	Nussbaumer and Hälg (2015)	1.3 - 240

CH₄ emission factors for wood

There are many different combustion installations in use which have very different CH₄ emission factors. A detailed overview of all applied wood related CH₄ emission factors for the entire time series is given in Table 3-15.

The CH₄ emission factor for each combustion type is modelled based on VOC measurements at wood combustion installations (Nussbaumer and Hälg 2015), assuming a CH₄ to VOC ratio of 0.4.

The EF for the different combustion installations varies between 1.3 and 240 g CH₄/GJ depending on the year, rated thermal input and technology used. The EF value of a single category represents the emission characteristics of a large number of combustion installations with a range of technology types, maintenance and operating conditions at a given time. According to their lifespan, existing combustion installations are gradually replaced by installations of new technology with better combustion, resulting in gradually decreasing emission factors.

Table 3-15 CH₄ emission factors for wood combustion installations.

1A Wood combustion	Unit	CH ₄			
		1990	1995	2000	2005
Open fireplaces	g/GJ	160	149	138	127
Closed fireplaces, log wood stoves	g/GJ	160	149	138	127
Pellet stoves	g/GJ	16	15	14	13
Log wood hearths	g/GJ	240	229	218	207
Log wood boilers	g/GJ	200	161	122	83
Log wood dual chamber boilers	g/GJ	240	229	218	207
Automatic chip boilers < 50 kW	g/GJ	20	17	13	10
Automatic pellet boilers < 50 kW	g/GJ	6.7	5.6	4.5	3.4
Automatic chip boilers 50-500 kW w/o wood proc. companies	g/GJ	20	16	13	8.9
Automatic pellet boilers 50-500 kW	g/GJ	6.7	5.4	4.1	2.8
Automatic chip boilers 50-500 kW within wood proc. companies	g/GJ	20	16	13	8.9
Automatic chip boilers > 500 kW w/o wood proc. companies	g/GJ	13	11	8.1	5.6
Automatic pellet boilers > 500 kW	g/GJ	6.7	5.4	4.1	2.8
Automatic chip boilers > 500 kW within wood proc. companies	g/GJ	13	11	8.1	5.6
Combined chip heat and power plants	g/GJ	13	11	8.1	5.6
Plants for renewable waste from wood products	g/GJ	13	11	8.1	5.6

1A Wood combustion	Unit	CH ₄									
		2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Open fireplaces	g/GJ	124	122	120	120	120	120	120	120	120	119
Closed fireplaces, log wood stoves	g/GJ	124	122	120	117	113	110	107	103	100	98
Pellet stoves	g/GJ	12	12	12	12	12	12	12	12	12	12
Log wood hearths	g/GJ	204	202	200	193	187	180	173	167	160	157
Log wood boilers	g/GJ	76	68	60	58	57	55	53	52	50	49
Log wood dual chamber boilers	g/GJ	204	202	200	193	187	180	173	167	160	154
Automatic chip boilers < 50 kW	g/GJ	9.3	8.7	8.0	8.0	8.0	8.0	8.0	8.0	8.0	7.7
Automatic pellet boilers < 50 kW	g/GJ	3.1	2.9	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.6
Automatic chip boilers 50-500 kW w/o wood proc. companies	g/GJ	8.2	7.4	6.7	6.7	6.7	6.7	6.7	6.7	6.7	6.5
Automatic pellet boilers 50-500 kW	g/GJ	2.5	2.3	2.0	2.0	2.0	2.0	2.0	2.0	2.0	1.9
Automatic chip boilers 50-500 kW within wood proc. companies	g/GJ	8.2	7.4	6.7	6.7	6.7	6.7	6.7	6.7	6.7	6.5
Automatic chip boilers > 500 kW w/o wood proc. companies	g/GJ	5.0	4.5	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.9
Automatic pellet boilers > 500 kW	g/GJ	2.5	2.3	2.0	2.0	2.0	2.0	2.0	2.0	2.0	1.9
Automatic chip boilers > 500 kW within wood proc. companies	g/GJ	5.0	4.5	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.9
Combined chip heat and power plants	g/GJ	5.0	4.5	4.0	3.6	3.1	2.7	2.2	1.8	1.3	1.3
Plants for renewable waste from wood products	g/GJ	5.0	4.5	4.0	3.6	3.1	2.7	2.2	1.8	1.3	1.3

3.2.4.4 N₂O Emission factors for 1A Fuel combustion

Table 3-16 shows the general N₂O emission factors in source category 1A which are based on default values from the 2006 IPCC Guidelines (IPCC 2006) and kept constant over the whole period. N₂O emission factors related to transport activities (aviation, road and non-road transportation) and for waste incineration are category specific and given in the corresponding chapters.

Table 3-16 N₂O emission factors. Default emission factors are used for all fuels for the whole time period.

N ₂ O emission factors			1990-2015
Fuel	CS/D	Data sources	g N ₂ O / GJ
Jet Kerosene	D	IPCC (2006)	2
Gas oil	D	IPCC (2006)	0.6
Residual fuel oil	D	IPCC (2006)	0.6
Liquefied petroleum gas	D	IPCC (2006)	0.1
Petroleum coke	D	IPCC (2006)	0.6
Other bituminous coal	D	IPCC (2006)	1.5
Lignite	D	IPCC (2006)	1.5
Natural gas	D	IPCC (2006)	0.1
Biofuel	CS/D	Data sources	
Biogas	D	IPCC (2006)	0.1
Wood	D	IPCC (2006)	4

3.2.4.5 Models overlapping more than one source category

3.2.4.5.1 Non-road transportation model (excl. aviation)

Choice of method

- The GHG emissions are calculated by a Tier 3 method based on the decision tree Fig. 3.3.1 in chp. 3. Mobile Combustion in IPCC (2006), complemented with
- Tier 2 for railways CO₂, Fig. 3.4.1 in IPCC (2006)
- Tier 3 for railways CH₄, N₂O and precursors / SO₂, Fig. 3.4.2 in IPCC (2006)
- Tier 2 for navigation, Fig. 3.5.1 (Box 1) in IPCC (2006)

Methodology

The emissions of the non-road sector underwent an extended revision in 2014/2015, resulting in an update of GHG emissions including precursors and SO₂. Results are documented in FOEN (2015j). The non-road categories considered are listed in Table 3-17. All of them include several technologies (diesel oil, 2- or 4-stroke gasoline, natural gas, gas oil), and emission standards according to the classification shown in Figure 3-20.

Table 3-17 Non-road categories as specified in FOEN (2015j) and the corresponding nomenclature in the CRF.

Non-road categories (by Corinair)	Nomenclature CRF
Construction machinery	1.A.2.g.vii Off-road vehicles and other machinery
Industrial machinery	1.A.2.g.vii Off-road vehicles and other machinery
Railway machinery	1.A.3.c. Railways
Navigation machinery	1.A.3.d. Domestic Navigation
Garden-care/professional appliances	1.A.4.a.ii Commercial/institutional, Off-road vehicles and other machinery
Garden-care/hobby appliances	1.A.4.b.ii Residential, Off-road vehicles and other machinery
Agricultural machinery	1.A.4.c.ii Agriculture/forestry/fishing, Off-road vehicles and other machinery
Forestry machinery	1A.4.c.ii Agriculture/forestry/fishing, Off-road vehicles and other machinery
Military machinery (excl. aviation)	1.A.5.b Other, mobile, Military




Figure 3-20 Each non-road vehicle is classified by its engine-power class, engine type, machine type and machine category (FOEN (2015j), INFRAS 2015a).

The emission modelling is based on activity data and emission factors by means of the following equation, which holds on the most disaggregated level of engine power class (Figure 3-20):

$$Em = N \cdot H \cdot P \cdot \lambda \cdot \varepsilon \cdot CF_1 \cdot CF_2 \cdot CF_3 \cdot$$

with

Em	=	emission per engine type (in g/a)
N	=	number of vehicles (--)
H	=	number of operation hours per year (h/a)
P	=	engine power output (kW)
λ	=	effective load factor (--)
ε	=	emission factor (g/kWh)
CF_1	=	correction factor for the effective load (--)
CF_2	=	correction factor for dynamical engine use (--)
CF_3	=	degradation factor due to aging (--)

With this equation, the emissions of the following gases are calculated:

- GHG: CH₄, N₂O
- precursor gases: NO_x, CO
- air pollutant: VOC
- fuel consumption: in this case, ε represents the consumption instead of emission factor (in g/kWh)
- For other gases, the following method is applied:
- CO₂ is calculated as product of fuel consumption and CO₂ emission factors (Table 3-12)
- SO₂ is calculated as product of fuel consumption and SO₂ emission factors (Table A – 19)
- NMVOC is calculated as the difference between VOC and CH₄

The total emission and consumption per non-road category is calculated by taking the sum over all engine-power classes, engine types, and machine types.

Emissions are only calculated in steps of 5 years 1980, 1985, 1990, ... 2050. Emissions for years in-between (1981, 1982 etc.) are interpolated linearly. A more detailed description of the analytical details is given in the Annex of FOEN (2015j).

Emission factors

Emission factors are taken from various sources based on measurement, modelling and literature.

- CO₂ and SO₂ emission factors are country-specific, see Table 3-12 and Table A – 19
- For other gases, the main data sources are EPA (2010), IFEU (2010), EMEP/EEA (2013) and Integer (2013).

For a detailed description of emission factors and their origin, see tables in the annex of FOEN (2015j) and online in the database belonging to INFRAS (2015a)⁶.

Activity data

Activity data were collected by surveys among producers and several user associations in Switzerland (FOEN 2015j), and by evaluating information from the national database of non-road vehicles (MOFIS) run by the the Federal Roads Office (FEDRO 2014). In addition, several publications serve as further data sources:

- SBV (2013) for agricultural machinery
- SFSO (2013a) for agricultural machinery
- Jardin Suisse (2012) for garden care /hobby and professional appliances
- KWF (2012) for forestry machinery
- The national statistics on imports/exports of non-road vehicles was assessed by FCA (2015c)
- Off-Highway Research (2005, 2008, 2012) provided information on the number of non-road vehicles.
- Federal Department of Defence, Civil Protection and Sport: List of military machinery with vehicle stock, engine-power classes and operating hours (DDPS 2014a).

From these data sources, all necessary information was developed like size distributions, modelling of the fleets, annual operating hours (age-dependent), load factors, year of placing on the market and age distribution. Details are documented in FOEN (2015j). All activity data (vehicle stocks, operating hours, consumption factors) can be downloaded by query from the public part of the non-road database INFRAS (2015a), which is the data pool of FOEN (2015j). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels.

⁶ <http://www.bafu.admin.ch/luft/00596/06906/offroad-daten/index.html?lang=en>

3.2.4.5.2 Energy model for wood combustion

Choice of method

The emissions from wood combustion in 1A Fuel combustion activities are calculated by a Tier 2 method based on the decision tree for stationary fuel combustion (IPCC 2006, Volume 2 Energy, chp. 2 Stationary Combustion, Figure 2.1 on page 2.15).

Methodology

The Swiss wood energy statistics (SFOE 2016b) provide both the annual wood consumption for specified categories of combustion installations (table K, categories 1-19), see Table 3-18 below, and the allocations of the combustion categories to the sectoral consumer categories (table N, household, agriculture/forestry, industry, services, electricity and district heating). This allows for assigning the annual wood consumption at the level of combustion installation categories directly to the source categories 1A1a Public electricity and heat production, 1A2gviii Other, 1A4ai Commercial/Institutional, 1A4bi Residential and 1A4ci Agriculture/forestry/fishing (EMIS 2017/1A Holzfeuerungen).

Table 3-18 Categories of wood combustion installations based on SFOE 2016b.

1A Wood combustion, categories
Open fireplaces
Closed fireplaces, log wood stoves
Pellet stoves
Log wood hearths
Log wood boilers
Log wood dual chamber boilers
Automatic chip boilers < 50 kW
Automatic pellet boilers < 50 kW
Automatic chip boilers 50-500 kW w/o wood processing companies
Automatic pellet boilers 50-500 kW
Automatic chip boilers 50-500 kW within wood processing companies
Automatic chip boilers > 500 kW w/o wood processing companies
Automatic pellet boilers > 500 kW
Automatic chip boilers > 500 kW within wood processing companies
Combined chip heat and power plants
Plants for renewable waste from wood products

Emission Factors

Emission factors are described in chp. 3.2.4.4.2 for CO₂, 3.2.4.4.3 for CH₄, and 3.2.4.4.4 for N₂O.

Activity Data

Total activity data are based on the Swiss wood energy statistics (SFOE 2016b). As additional data source, specific bottom-up information from the industry is used in order to allocate wood combustion emissions directly to a particular source category. Thus, activity data of wood combustion within 1A2f, 1A2giv and 1A4ci are allocated on the basis of industry information (see Figure 3-19 and EMIS 2017/1A Holzfeuerungen):

- Wood energy consumption in source categories 1A2f Brick and tile production, 1A2f Cement production and 1A2giv Fibreboard are subtracted from the activity data of 1A2gviii Automatic chip boiler >500 kW without wood processing companies and 1A2gviii Plants for renewable waste from wood products, respectively.
- Since 2013, also the wood energy consumption in 1A4ci Grass drying is available and has been subtracted from the activity data in 1A4ci Automatic chip boiler >500 kW without wood processing companies.

Table 3-19 Wood energy consumption in 1A Fuel combustion.

1A Wood combustion	Unit	1990	1995	2000	2005
Total	TJ	28'165	29'419	27'041	30'785
Open fireplaces	TJ	227	271	196	181
Closed fireplaces, log wood stoves	TJ	7'275	7'178	6'493	7'047
Pellet stoves	TJ	0	0	7	48
Log wood hearths	TJ	8'524	7'030	4'744	4'029
Log wood boilers	TJ	5'308	5'571	5'109	5'366
Log wood dual chamber boilers	TJ	1'964	1'779	978	481
Automatic chip boilers < 50 kW	TJ	239	434	550	754
Automatic pellet boilers < 50 kW	TJ	0	0	56	805
Automatic chip boilers 50-500 kW w/o wood proc. companies	TJ	688	1'332	1'793	2'707
Automatic pellet boilers 50-500 kW	TJ	0	0	2	99
Automatic chip boilers 50-500 kW within wood proc. companies	TJ	1'287	1'720	1'755	1'918
Automatic chip boilers > 500 kW w/o wood proc. companies	TJ	327	992	1'596	2'244
Automatic pellet boilers > 500 kW	TJ	0	0	0	9
Automatic chip boilers > 500 kW within wood proc. companies	TJ	1'347	2'048	2'232	2'531
Combined chip heat and power plants	TJ	0	3	186	127
Plants for renewable waste from wood products	TJ	979	1'060	1'345	2'438

1A Wood combustion	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Total	TJ	31'163	29'879	33'879	35'274	37'634	33'283	37'595	41'360	35'412	36'922
Open fireplaces	TJ	171	150	150	137	123	87	84	83	62	64
Closed fireplaces, log wood stoves	TJ	6'939	6'282	6'827	7'011	7'913	6'586	7'468	8'333	6'737	7'451
Pellet stoves	TJ	65	72	93	109	140	127	156	184	157	182
Log wood hearths	TJ	3'604	2'914	2'839	2'491	2'268	1'576	1'485	1'351	900	918
Log wood boilers	TJ	5'162	4'586	4'892	4'751	4'905	3'691	3'826	3'884	2'822	2'972
Log wood dual chamber boilers	TJ	423	348	338	289	272	195	190	181	125	119
Automatic chip boilers < 50 kW	TJ	775	727	833	861	1'008	801	867	943	740	787
Automatic pellet boilers < 50 kW	TJ	1'193	1'248	1'562	1'731	2'104	1'814	2'153	2'499	2'102	2'378
Automatic chip boilers 50-500 kW w/o wood proc. companies	TJ	2'914	2'828	3'224	3'288	3'780	3'251	3'821	4'329	3'561	4'100
Automatic pellet boilers 50-500 kW	TJ	184	259	365	434	540	515	624	726	692	871
Automatic chip boilers 50-500 kW within wood proc. companies	TJ	1'882	1'751	1'872	1'874	2'012	1'744	1'885	2'026	1'676	1'832
Automatic chip boilers > 500 kW w/o wood proc. companies	TJ	2'443	2'511	3'122	3'400	3'963	3'631	4'350	5'065	4'425	5'200
Automatic pellet boilers > 500 kW	TJ	39	56	80	84	92	139	161	186	192	214
Automatic chip boilers > 500 kW within wood proc. companies	TJ	2'470	2'348	2'451	2'459	2'689	2'348	2'510	2'684	2'250	2'470
Combined chip heat and power plants	TJ	242	1'058	2'467	3'423	2'756	3'900	5'010	5'421	5'325	3'792
Plants for renewable waste from wood products	TJ	2'657	2'740	2'764	2'933	3'071	2'877	3'005	3'465	3'647	3'573

3.2.4.6 Emissions from Biomass (memo item)

CO₂ emissions from biomass do not count for the national total emissions and therefore are a memo item only. The CO₂ emissions from biomass as reported in the reporting tables are incomplete as the following CO₂ emissions are not foreseen for reporting in the reporting tables: 2G4 Use of tobacco, 2H2 Food and beverages, 5A Solid waste disposal, 5B Biological treatment of solid waste and 5D Wastewater treatment and discharge.

Table 3-20 provides an overview of effective CO₂ emissions from biomass in 2015 and their reporting in the reporting tables (without land-use, land-use change and forestry). For further information on the biomass CO₂ emissions refer to the respective source category chapters.

Table 3-20 Effective biomass CO₂ emissions in 2015 and their representation in the reporting tables. Note that in reporting table CRF Table10s2 biogene CO₂ emissions from 5C are missing (for an overview of errors of the CRF Reporter see Annex 6).

Biomass CO ₂ emissions	Unit	2015	Note
1A1 Energy industries (without MSW incineration)	kt	488	Included in CRF
1A1 Energy generation from MSW Incineration	kt	2'188	Included in CRF
1A2 Manufacturing industry and construction	kt	1'340	Included in CRF
thereof use of waste derived fuels in cement production	kt	51	
thereof use of bio fuels (1A2gvii)	kt	6	
1A3 Transport	kt	170	Included in CRF
1A4 Other sectors (Commercial/institutional, residential)	kt	2'785	Included in CRF
1A5 Other	kt	0.21	Included in CRF
2H2 Food and beverages industry	kt	15	Not included in CRF
2G Other product use (Consumption of tobacco)	kt	10	Not included in CRF
5A Solid waste disposal on land	kt	46	Not included in CRF
5B Biological treatment of waste (composting and anaerobic digestion)	kt	375	Not included in CRF
5C Waste incineration (without MSW incineration)	kt	143	Included in CRF
5D Wastewater handling	kt	128	Not included in CRF
Total biomass combustion CO ₂ emissions included in CRF	kt	7'115	
Total energy related biomass combustion CO ₂ emissions included in CRF 1A	kt	6'971	See tab "Summary2" in CRF
Total biomass CO ₂ emissions in Switzerland in 2015	kt	7'689	

3.2.4.7 Uncertainty and time series consistency for source category 1A

Basic uncertainties of AD and EF CO₂ by fuel type

Table 3-21 Uncertainties of activity data and CO₂ emission factors for 1A Fuel combustion.

Fuel type	Uncertainties		
	Activity data	CO ₂ emission factors	CO ₂ emissions
	%		
kerosene	0.96	0.16	0.97
gasoline	0.69	0.13	0.70
diesel oil	0.88	0.07	0.88
liquid fuels	0.69	0.08	0.69
solid fuels	5.00	5.06	7.12
gaseous fuels	5.00	1.04	5.11
other fuels	5.00	9.22	10.5
biomass	10.0	--	--

Liquid fuels

Uncertainty of the CO₂ emission factors: In 2013, a large measurement campaign was carried out to determine the CO₂ emission factors of the dominant liquid fuels (SFOE/FOEN 2014). From the standard deviation presented in this study, the 95% uncertainties are derived and shown in Table 3-22 as lower and upper values as well as relative uncertainties.

The uncertainties were updated for submission 2016. The new values are applied for the current submission. The former values were also based on measurements but were carried out in 1998 (EMPA 1999), but with a smaller number of samples (between 10 and 30 samples per fuel type). The former uncertainties were then 1.16% for kerosene, 1.36% for gasoline and 0.47% for diesel oil and 0.61% for gas oil.

For mobile combustion, the 2006 IPCC Guidelines provide default uncertainties for the CO₂ emission factor of kerosene as 2%, gasoline 4% and diesel oil 1% (IPCC 2006, vol. 2, TABLE 3.2.1). Switzerland's measurements indicate much lower uncertainties. For stationary combustion, the 2006 IPCC Guidelines give no default values but show instead a summary of an uncertainty assessment of CO₂ emission factors for stationary combustion for selected countries (IPCC 2006, vol 2, TABLE 2.13). The values lie in the range between 0.5% and 3% and are again higher than the values derived from the Swiss measurements.

Table 3-22 Uncertainties of aggregated results of measurements of the CO₂ emission factors of selected liquid fuels (SFOE/FOEN 2014).

Fuel type	CO ₂ emission factors (measurements)			95% uncertainties EF(CO ₂)		no. samples
	mean t/TJ	lower t/TJ	upper t/TJ	absolute t/TJ	relative	
kerosene	72.81	72.70	72.93	0.12	0.16%	24
gasoline	73.80	73.71	73.90	0.10	0.13%	138
diesel oil	73.30	73.25	73.35	0.05	0.07%	75
gas oil	73.67	73.61	73.73	0.06	0.08%	138

Uncertainties of activity data: The values shown in Table 3-23 are based on a written message of SFOE to FOEN (SFOE 2012a). It lists two kinds of relevant errors: errors of measurements and errors of the conversion from mass to energy units. For gasoline and diesel oil, a third source of errors stems from the transformation of products. These errors are multiplicative, therefore the relative uncertainties have to be summed up.

Up to submission in 2015, a single expert estimate has been used, being the same for all liquid fuels (1%). The updated and more sophisticated uncertainties are equal (kerosene) or slightly lower.

Table 3-23 Sources of errors contributing to the total uncertainty of the activity data of selected liquid fuels (SFOE 2012a).

source of uncertainty	kerosene	gasoline	diesel oil	gasoil
	activity data uncertainty in %			
Measurement	0.39%	0.39%	0.39%	0.39%
Conversion mass to energy	0.57%	0.29%	0.29%	0.29%
Product transformation	0.00%	0.00%	0.20%	0.00%
Total uncertainty	0.96%	0.69%	0.88%	0.69%

Gaseous fuels

Uncertainty of the CO₂ emission factor: The composition of the imported gas is analysed in detail at the import stations. From this information, the FOEN annually calculates the CO₂

emission factor for each import station and the weighted mean. To estimate the uncertainty of the emission factor, the weighted standard deviation is calculated and is multiplied by the factor 1.96 to extend the standard deviation to 95% uncertainty interval. This calculation has been carried out for 13 years within the period 1990 and 2015. The uncertainties fluctuate between 0.50% and 1.48% with a mean of 1.04%, which is used as the uncertainty of the CO₂ emission factor for gaseous fuels.

Uncertainty of activity data: There is no country-specific estimate of the uncertainty for the consumption of natural gas. It is taken from the 2006 IPCC Guidelines (IPCC 2006, vol. 2, TABLE 2.15), which give a range of 2%-5% for industrial combustion and 3%-5% for commercial, institutional and residential combustion. For Switzerland, an overall value of 5% is used.

Solid fuels

Uncertainty of the CO₂ emission factor: There is no country-specific uncertainty available. The 2006 IPCC Guidelines suggest a range from 0.5% to 10% (IPCC 2006, vol. 2 TABLE 2.13) . For Switzerland, an uncertainty of 5% is chosen (medium of suggested range).

Uncertainty of activity data: There is no country-specific estimate of the uncertainty for the consumption of coal. It is taken from the 2006 IPCC Guidelines (IPCC 2006, vol. 2, TABLE 2.15, which give a range of 2%-5% for industrial combustion and 3%-5% for commercial, institutional and residential combustion. For Switzerland, an overall value of 5% is used (as for natural gas).

Other fuels (waste to energy)

Uncertainty of the CO₂ emission factor: There are two factors influencing the uncertainty of CO₂ emissions from municipal solid waste incineration (1A1): the carbon content of waste and the fossil carbon fraction of the carbon content.

- The carbon content is determined in a study by Fellner et al. (2007). A relation between the calorific value of waste and the carbon content is derived therein, which provides upper and lower limits. The relation is tested by measurements. The difference between upper and lower limits (5.9%) is interpreted as 95% confidence interval for the carbon content.
- The fossil fraction of the carbon content is determined in another study by Mohn et al. (2011). A field application of the radio carbon (¹⁴C) method was applied to calculate the ratio of biogenic versus fossil CO₂ emissions from five waste-to-energy plants. Gas samples for ¹⁴CO₂ analysis were taken at the plants during miscellaneous seasons. Six measuring campaigns of three weeks periods were carried out for three plants and three campaigns, again of three weeks periods, were carried out for two plants. That means that the measurements lasted 72 weeks in total. The 95% confidence interval of the campaigns result in a biogenic fraction of 52.3% ± 3.8% (Table 3-24), which corresponds to an uncertainty of 7.1%. The results fit well to a former measurement campaign on three plants which yield 52.0% ± 3.7% for the biogenic fraction (Mohn 2008). For the uncertainty analysis the latest result is used.

Table 3-24 Measures shares of fossil and biogenic share in five MSW plants campaign.

Plant	Shares		Uncertainty		Measurement campaigns (duration) weeks
	fossil %	biogenic %	absolute %	relative %	
Buchs	47.7	52.3	3.6	7.5	6 x 3 = 18
Winterthur	43.4	56.6	3.9	9.0	6 x 3 = 18
Linthgebiet	50.6	49.4	3.4	6.7	6 x 3 = 18
Fribourg	54.5	45.5	3.1	5.7	3 x 3 = 18
Zuchwil	45.9	54.1	3.7	8.1	3 x 3 = 18
Median / sum	47.7	52.3	1.8	3.8	72

- The fossil-CO₂ emission factor results from a multiplication of the carbon content and the fossil fraction. The uncertainty of the CO₂ emission factor is thus the addition of the corresponding uncertainties by error propagation: $[(5.9\%)^2 + (3.8\%)^2]^{0.5} = 7.0\%$. (For the previous submission 9.2% had been applied based on the former results).

Uncertainty of activity data: There is no country-specific estimate of the uncertainty for the consumption of waste. It is taken from the 2006 IPCC Guidelines (IPCC 2006, vol. 2, p. 2.40), which states: “Experts believe that the uncertainty resulting from the two errors (*systematic, random*) combined is probably in the range of $\pm 5\%$ for most developed countries.” In accordance with that statement, the value of 5% is used.

Biomass

Uncertainty of the CO₂ emission factor: For CO₂ emissions of biomass burning, no uncertainty is estimated (memo item).

Uncertainty of activity data: no country-specific uncertainty of the activity data is available. The 2006 IPCC Guidelines suggest 2% to 5% for industrial, institutional and residential combustion and 10% to 30% for biomass burning in small sources (IPCC 2006, vol. 2, TABLE 2.15). About 80% of the CO₂ emissions from biomass burning stem from industrial, institutional and residential combustion and 20% from small residential installations with less developed statistical systems (see Table 3-19). An average uncertainty of 10% is applied for biomass burning in all source categories.

Uncertainty of CH₄ and N₂O emission factors

Since the CO₂ emissions vastly dominate the GHG emissions of source category 1A (almost 99%), the uncertainty evaluation of the non-CO₂ emissions is carried out on a semi-quantitative level (see Table 3-25).

Only for **1A3b Road transportation** a quantitative analysis has been performed. Following a study for the road transportation in Germany (IFEU/INFRAS 2009), where the same handbook of emission factors is used as in Switzerland (INFRAS 2010), the uncertainties for the CH₄ and N₂O emission factors have been determined (see lines 1A3b gasoline and diesel oil in Table 3-25). The uncertainties of CH₄ and N₂O emissions of CNG (1A3b), which were not investigated in IFEU/INFRAS (2009), have been estimated qualitatively as “medium” according to Table 1-10. For **1A1, 1A2, 1A3a, 1A3c, 1A3d, 1A3e, 1A4a, 1A4b,**

1A4c, 1A5 the uncertainties of CH₄ and N₂O emissions have similarly been estimated qualitatively (see Table 3-25).

Summary

Table 3-25 below provides a summary of the uncertainties of 1A Fuel combustion as derived in the preceding sections. The uncertainty of the CO₂ emissions (“combined uncertainty”) are calculated from the uncertainties of the activity data and the emission factors by Approach 1 error propagation.

Table 3-25 Uncertainties of 1A Fuel combustion categories for activity data, emission factors and combined uncertainties. The latter are calculated by Approach 1. (For 1A2/Other Fuels a mean uncertainty is assumed based on semi-quantitative estimations from Table 1-10. The emission factor uncertainty is calculated “backward”⁷ from the combined and the activity data uncertainty). CH₄ and N₂O: semi-quantitative uncertainties (see Table 1-10).

1A Fuel Combustion Categories	Fuel type	Uncertainties				
		Activity data	CO ₂ em. factors	CO ₂ emissions	CH ₄ emissions	N ₂ O emissions
		%	%	%	--	--
1. Energy industries	liquid fuels	0.7	0.1	0.7	medium	medium
1. Energy industries	solid fuels	5.0	5.1	7.1	medium	medium
1. Energy industries	gaseous fuels	5.0	1.0	5.1	medium	medium
1. Energy industries	other fuels	5.0	9.2	10.5	medium	medium
2. Manufacturing industries and construction	liquid fuels	0.7	0.1	0.7	medium	medium
2. Manufacturing industries and construction	solid fuels	5.0	5.1	7.1	medium	medium
2. Manufacturing industries and construction	gaseous fuels	5.0	1.0	5.1	medium	medium
2. Manufacturing industries and construction	other fuels	5.0	9.2	10.5	medium	medium
3a. Transport; Domestic aviation	kerosene	1.0	0.2	1.0	high	high
3b. Transport; Road transportation	gasoline	0.7	0.1	0.7	37.0	50.0
3b. Transport; Road transportation	diesel oil	0.9	0.1	0.9	20.0	22.0
3b. Transport; Road transportation	gaseous fuels	5.0	1.0	5.1	medium	medium
3c. Transport; Railways	diesel oil	0.9	0.1	0.9	medium	medium
3d. Transport; Domestic navigation	liquid fuels	0.7	0.1	0.7	medium	high
3e. Transport; Other transportation	gaseous fuels	5.0	1.0	5.1	medium	medium
4a. Other sectors; Commercial/institutional	liquid fuels	0.7	0.1	0.7	medium	medium
4a. Other sectors; Commercial/institutional	gaseous fuels	5.0	1.0	5.1	medium	medium
4b. Other sectors; Residential	liquid fuels	0.7	0.1	0.7	medium	medium
4b. Other sectors; Residential	solid fuels	5.0	5.1	7.1	medium	medium
4b. Other sectors; Residential	gaseous fuels	5.0	1.0	5.1	medium	medium
4c. Other sectors; Agriculture/forestry/fishing	liquid fuels	0.7	0.1	0.7	medium	medium
4c. Other sectors; Agriculture/forestry/fishing	gaseous fuels	5.0	1.0	5.1	medium	medium
5. Other	liquid fuels	0.7	0.1	0.7	medium	high
1A Stationary sources	biomass	10.0	--	--	medium	medium
1A Mobile sources	biomass	10.0	--	--	high	high

Time series consistency 1A

Time series for 1A Fuel combustion are all considered consistent.

$${}^7 U(EF) = \sqrt{U(EM)^2 - U(AD)^2}$$

3.2.4.8 Category-specific QA/QC and verification for source category 1A

Various QA/QC activities are relevant for all source categories in 1A. Therefore, they are briefly described here and not repeated again in the chapters dealing with source categories 1A1 to 1A5.

Comparison of emission estimates using different approaches

At the level of total energy-related CO₂ emissions, a quality control consists in the comparison of emissions modelled using the sectoral approach with emissions calculated based directly on fuel consumption according to the Swiss overall energy statistics (SFOE 2016). The differences in total CO₂ emissions for the entire time period are negligible, indicating the completeness of the inventory.

The cross-check of the Reference and Sectoral Approach is also used for an assessment of emissions related to the consumption of fuels in the energy sector. Again, a good agreement between the two approaches is found (see chp. 3.2.1).

Activity data checks

The SFOE constructs a national commodity balance expressed in mass and in energy units including mass balances of fuel conversion industries.

The gross carbon supply in the Reference Approach has been adjusted for fossil fuel carbon destined for non-energy use. The numbers in the Swiss overall energy statistics (SFOE 2016) are consistent with those provided by international organisations, e.g. IEA.

Emission factor check and review

Emission factors for the main fossil fuels have been reassessed for submission 2015. In 2013, the Federal Office for the Environment (FOEN) and the Swiss Federal Office for Energy (SFOE) launched an in-depth investigation into the NCV and CO₂ emission factors of gas oil, diesel oil, gasoline, and kerosene (SFOE/FOEN 2014), see description under 3.2.4.2 Net calorific values (NCV). The values differ only marginally from previously used values. The CO₂ emission factors compare well with the IPCC default values (see Table 3-26).

Table 3-26 Comparison of default CO₂ emission factors from IPCC 2006 with current country-specific values for selected fuels.

CO ₂ Emission Factors	IPCC 2006			Switzerland
	Lower	Upper	Default	CS
	t CO ₂ / TJ			
Gasoline	67.5	73.0	69.3	73.8
Jet kerosene	69.7	74.4	71.5	72.8
Diesel oil	72.6	74.8	74.1	73.3
Gas oil	72.6	74.8	74.1	73.7

The CO₂ emission factor for gasoline is higher than the IPCC range. However, as the value from earlier measurements was confirmed and the new value is based on more than 100 fuel samples taken from July to December 2013, the value is considered to correctly represent national circumstances.

For natural gas, the CO₂ emission factor has been reassessed for submission 2017. A country-specific CO₂ emission factor, based on measurements of gas properties and corresponding import shares of individual gas import stations is calculated (see also chp. 3.2.4.2). The resulting values are largely consistent with the CO₂ EF used by the countries from which gas is imported (Germany (IEF: 55.5–56.0 t CO₂/TJ), the Netherlands (IEF: 56.5–56.8 t CO₂/TJ), Norway (IEF: 56.1 t CO₂/TJ), France (IEF: 56.0–57.0 t CO₂/TJ), Italy (IEF: 55.3 to 56.9 t CO₂/TJ) and Denmark (IEF: 56.9–57.5 t CO₂/TJ). It lies within the range given by the IPCC (lower 54.3 t CO₂/TJ, upper 58.3 t CO₂/TJ, compared to the country-specific value of 56.4 t CO₂/TJ for 2015).

The CH₄ emission factors from combustion of wood have been scrutinized and revised based on Nussbaumer and Hälg (2015). The range of country-specific values is not entirely consistent with the upper and lower IPCC default values (Table 3-27). However, as the country-specific emission factors are based on an extensive measurement campaign, they are considered representative for Swiss circumstances.

Table 3-27 Comparison of default CH₄ emission factors from the 2006 IPCC Guidelines (IPCC 2006) with country-specific values

CH ₄ Emission factors	IPCC 2006			Switzerland CS
	Lower	Upper	Default	
kg CH ₄ / TJ				
Wood	10	100	30	1.3 - 240

Expert review

As described in chp. 1.2.3, data from source category 1A and the initial draft of the NIR were scrutinized in an external review involving national experts and stakeholders in the different fields related to emissions from stationary sources.

3.2.4.9 Category-specific recalculations for source category 1A in general

- Reference Approach: Consumption of so-called other non-fossil fuels (biogenic waste) was missing so far. It is now also included in the Reference Approach in CRF table 1.A(b).
- 1A: The CO₂ emission factor for natural gas is linearly interpolated between 1995–2000. (Before it was kept constant between 1995–1998).
- 1A: The CO₂ emission factor for natural gas has changed. There was a minor error in the calculations for the years as follows:
2009: from 56'400g/GJ to 56'500 g/GJ,
2010: from 56'400g/GJ to 56'500 g/GJ,
2014: from 56'700g/GJ to 56'500 g/GJ

- 1A: Small recalculations due to rounding of activity data 2013, 2014 in the Swiss overall energy statistics (SFOE 2016) concerning bituminous coal and natural gas.
- 1A: Non-energy use of fuels (NEU): The net consumption of non-energy use of fuels reported in Swiss overall energy statistics includes sulphur produced by the refineries as well. This amount of sulphur is now subtracted resulting in lower fuel quantities for NEU of other oil for the entire time series reported in CRF table 1.A(d).

3.2.4.10 Planned improvements for source category 1A in general

No improvements for 1A general are planned

3.2.5 Source category 1A1 - Energy industries (stationary)

3.2.5.1 Source category description for 1A1 (stationary)

Table 3-28 Key categories (KCA incl. LULUCF) of 1A1 Energy industries

A	B	C	D
Code	IPCC Category	GHG	Identification Criteria
1A1	Energy Industries: Gaseous Fuels	CO2	L1, T1
1A1	Energy Industries: Liquid Fuels	CO2	L1, T1
1A1	Energy Industries: Other Fuels	CO2	L1, L2, T1, T2
1A1	Energy Industries: Solid Fuels	CO2	T2

Source category 1A1 Energy industries comprises emissions from fuels combusted by the fuel extraction and energy-producing industries. The most important source category is 1A1a Public electricity and heat production, followed by 1A1b Petroleum refining. Activities in source category 1A1c Manufacture of Solid Fuels and other energy industries is virtually not occurring in Switzerland (apart from a tiny charcoal production activity in historic trade).

Within source category 1A1a, heat and electricity production in waste incineration plants cause the largest emissions, as electricity production in Switzerland is dominated by hydroelectric power plants and nuclear power stations (SFOE 2016). Emissions from industries producing heat and/or electricity (CHP) for their own use are included in category 1A2 Manufacturing Industries and Construction.

Table 3-29 Specification of source category 1A1 Energy Industries.

1A1	Source	Specification
1A1a	Public electricity and heat production	Main source are waste incineration plants with heat and power generation (Other fuels) and public district heating systems. The only fossil fuelled public electricity generation unit "Vouvry" (300 MWe; no public heat production) ceased operation in 1999.
1A1b	Petroleum refining	Combustion activities supporting the refining of petroleum products, excluding evaporative emissions. Emissions of SO ₂ from Claus units in refineries.
1A1c	Manufacture of solid fuels and other energy industries	Charcoal production

3.2.5.2 Methodological issues for 1A1 (stationary)

3.2.5.2.1 Public electricity and heat production (1A1a)

Public electricity and heat production in Switzerland encompasses different plant types where various fuels are used (Table 3-30) Energy recovery from municipal solid waste and special waste incineration is mandatory in Switzerland and plants are equipped with energy recovery systems. The emissions from municipal solid waste and special waste incineration plants are therefore reported under category 1A1a. There was a single fossil fuel power station operating with residual fuel oil in Vouvry. However, the power station closed down in 1999.

Table 3-30 Plant type and fuels used in source category 1A1a.

Plant type	Fuel type
Heat plants for renewable wastes	wood waste (biomass)
Heating boilers >300 MW (Vouvry)	residual fuel oil
Heating boilers <300 MW	gas oil, residual fuel oil, bituminous coal
Central heating boilers for district heating	natural gas, gas oil, residual fuel oil, bituminous coal
Wood combined heat and power generation	wood, wood waste (biomass)
Engines and boilers at fermentation plants	digestion gas (biogas)
Engines on landfill sites	landfill gas (biogas)
Municipal solid waste incineration plants	municipal solid wastes (other, waste-to-energy)
Special waste incineration plants	special wastes (other, waste-to-energy)

Methodology (1A1a)

For Public electricity and heat production (1A1a) a country-specific approach, as explained in chp. 3.2.4.5 is used combining Tier 2 and Tier 3 methods (IPCC 2006, Volume 2 Energy, chp. 2 Stationary Combustion, Figure 2.1).

Emission factors (1A1a)

The following table presents the emission factors used in 1A1a. Emission factors for gas oil, residual fuel oil and natural gas are further explained in chp. 3.2.4.4.

Table 3-31 Emission Factors for 1A1a Public Electricity and Heat Production in 2015.

1A1a Public electricity and heat production	CO ₂	CO ₂ bio.	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
	t/TJ	t/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ
Gas oil	73.7	NA	3.0	0.6	34	2	18	6.5
Residual fuel oil	NO	NA	NO	NO	NO	NO	NO	NO
Petroleum coke	NO	NA	NO	NO	NO	NO	NO	NO
Natural gas	56.4	NA	1.0	0.1	18	2	0.5	10
Other (waste-to-energy), fossil	88.8	NA	NA	1.6				
Other (waste-to-energy), biogenic	NA	92.3	NA	1.4	32	2.4	3.7	8.5
Biomass (wood, renewable waste)	NA	99.9	1.3	4.0	116	2	11	98
Biogas (co-generation from landfills, fermentation engines)	NA	100.5	2.2	0.1	47	3	15	64

Emission factors for waste incineration and biogas use

Specific emission factors within 1A1a Public electricity and heat production apply for municipal solid waste incineration, special waste incineration and for biogas use (landfill gas and digestion gas). The emission factors for CO₂, CH₄, N₂O, NO_x, CO, NMVOC and SO₂ are country-specific and based on measurements and expert estimates, as documented in EMIS 2017/1A1a Kehrichtverbrennungsanlagen, EMIS 2017/1A1a Sondermüllverbrennungsanlagen, EMIS 1A1a Vergärung IG, EMIS 2017/1A1a Vergärung LW und EMIS 2017/1A1a Kehrichtdeponien.

Source-specific CO₂ emission factors for municipal solid waste incineration plants

C-content of waste is calculated based on the net calorific value (NCV), which is deduced by a standard method and published on a yearly basis since 2009 by SFOE for each MSWIP and as a Swiss average (FOEN/SFOE/VBSA, 2016). In deviation from the general description of oxidation factors in 3.2.4.4.1 an oxidation factor of 0.99 is assumed here. The assumption is based on measurements in two MSWIPs in Zurich (AWEL 2009) and on a study in Austria (Zeschmar-Lahl 2004), where the MSWIP have the same standards as in Switzerland. The measurements in Zurich showed transfer coefficients into air of 0.96–0.99 and the ones in Austria stated a transfer coefficient into air of 0.989.

The fossil fraction of waste incinerated in MSWIP is based on a study conducted in the year 2014 (Rytec 2014). The study uses data from three measurement campaigns during which the waste composition has been analysed (FOEN 2014o) and measurements of the radioactive isotope carbon-14 (¹⁴C) in the flue gas for calibration (Mohn 2011). The CO₂ emission factor in MSWIPs fluctuates over the reporting period because of gradual changes in the net calorific values of the waste. Please refer to Table 3-32 for data.

Table 3-32 Emission factor CO₂ total, share of CO₂ fossil and net calorific value (NCV) in municipal solid waste incineration plants (MSWIP) from 1990–2015.

1A1a Public electricity and heat production, Other fossil fuels	Unit	1990	1995	2000	2005
CO ₂ total (MSWIP)	t/TJ	92.80	91.86	91.09	91.49
Share of CO ₂ fossil (MSWIP)	1	0.497	0.505	0.513	0.505
NCV of waste (MSWIP)	TJ/t	0.0114	0.0119	0.0124	0.0121

1A1a Public electricity and heat production, Other fossil fuels	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
CO ₂ total (MSWIP)	t/TJ	91.62	91.90	92.20	92.50	92.32	92.25	92.62	92.76	92.43	92.28
Share of CO ₂ fossil (MSWIP)	1	0.501	0.497	0.493	0.489	0.486	0.482	0.478	0.478	0.478	0.478
NCV of waste (MSWIP)	TJ/t	0.0121	0.0119	0.0117	0.0116	0.0117	0.0117	0.0115	0.0114	0.0116	0.0117

Sodium bicarbonate and calcium carbonate are used in MSWIP for flue gas treatment. Sodium bicarbonate is used since 2013 and calcium carbonate was used between 1990 and 2005. According to IPCC 2006 the corresponding emission are reported in source category 2A4d.

Source-specific CO₂ emission factors for special waste incineration plants

Based on detailed information regarding waste composition and estimated emission factors in the years 1992–2004 a weighted average emission factor for special waste incineration was calculated. Special waste is assumed to be of entirely fossil origin. Overall, a specific emission factor of 1.45 t CO₂/t waste results for special waste. This value is considerably higher than the one reported in SAEFL (2000). As there is no newer data on the special waste composition the emission factor deduced as described above is used for the whole period from 1990 until today. See documentation in EMIS 2017/1A1a Sondermüllverbrennungsanlagen.

Source-specific CH₄ emission factors in municipal and special waste incineration plants

Emissions of CH₄ are not occurring in waste incineration plants because of the high temperatures and the long dwell time in the combustion chamber as confirmed by Mohn (2013). In the year 2013 EMPA assessed the N₂O and CH₄ emission factors for MSWIP (Mohn 2013). In this study EMPA evaluated measurements that were performed in 2011 in five Swiss MSWIP with different Denox techniques (SCR, SNCR). For most of the measurements CH₄ concentrations were below the detection limit of 0.3 ppm. The study concluded that "CH₄ emission concentrations were very low and below the background concentration of 1.8 ppm". CH₄ emissions are considered to be negligible for municipal waste incineration and are therefore set to zero. The same fact applies for special waste incineration.

Source-specific N₂O emission factors for municipal solid waste incineration

In 2013, a study evaluated N₂O measurements that have been performed in the years 2010–2011 in the flue gas of five Swiss municipal waste incineration plants (Mohn 2013) and derived plant-specific emission factors for Selective Catalytic Reduction (SCR) and Selective Non-Catalytic Reduction (SNCR) equipped installations.

Average Swiss emission factors have been calculated according to the state of equipment of all Swiss waste incineration plants (with two types of Denox-equipment (SCR, SNCR) and without Denox-equipment). For installations without Denox-equipment the emission factor comes from (SAEFL 2000). According to the state of equipment of all Swiss waste incineration plants in the years 1990, 1994, 1998, 2004, 2008 and 2012, weighted average N₂O emission factors have been calculated, based on the amounts of waste burnt in every plant. For the years in between, the N₂O emission factors were linearly interpolated. Since 2012 the emission factor is assumed to be constant (however the emission factor related to energy changes by reason of the conversion with the net calorific value of waste). It is planned to calculate a new value periodically, depending on data available. See documentation in EMIS 2017/1A1a Kehricht- und Sondermüllverbrennungsanlagen. The emission factor is therefore not constant over time.

Source-specific N₂O emission factors for special waste incineration

The emission factor of special waste for the year 1990 is based on SAEFL (2000). It is assumed that this value (3.1 g/GJ) then increases until 2003 (6.1g/GJ) due to the installation of Denox-equipment and thereafter declines as a result of optimized installations.

Table 3-33 N₂O emission factors of 1A1a Municipal solid and special waste incineration.

1A1a Public electricity and heat production, Other fossil fuels	Unit	1990	1995	2000	2005
N ₂ O (MSWIP)	kg/TJ	5.26	2.96	2.06	1.44
N ₂ O (SWIP)	kg/TJ	3.06	4.23	5.41	5.48

1A1a Public electricity and heat production, Other fossil fuels	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
N ₂ O (MSWIP)	kg/TJ	1.41	1.40	1.38	1.41	1.40	1.40	1.43	1.44	1.42	1.41
N ₂ O (SWIP)	kg/TJ	5.16	4.84	4.53	4.21	3.89	3.57	3.25	2.94	2.62	2.30

Activity data (1A1a)

Activity data for liquid, gaseous, solid fuels and wood are based on the Swiss overall energy statistics (SFOE 2016) and additional data sources as described in 3.2.4.3. Activity data for Other fuels are based on the amount of waste incinerated in MSWIPs and SWIPs (FOEN 2016i, see Table 3-35). Activity data for combined heat and power generation in landfills and in biogas facilities are taken from the Swiss renewable energy statistics (SFOE 2016a).

Please note that waste-to-energy activities in CRF Table 1.A(a)s1 are allocated to fuel types 'Other fossil fuels' and 'Biomass'. 'Other fossil fuels' encompasses emissions from fossil share of MSWIP and from SWIP. Whereas 'Biomass' covers emissions from wood, waste wood, landfill gas use in co-generation, digestion gas and biogenic share from MSWIP.

Table 3-34 Activity data in 1A1a Public Electricity and Heat Production.

1A1a Public electricity and heat production	Unit	1990	1995	2000	2005
Total fuel consumption	TJ	40'414	39'216	50'018	57'230
Gas oil	TJ	980	554	790	1'300
Residual fuel oil	TJ	3'214	1'813	340	290
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	530	46	NO	NO
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	4'339	5'422	8'292	9'827
Other (waste-to-energy), fossil	TJ	16'605	16'870	22'482	24'711
Biomass	TJ	14'747	14'511	18'114	21'103
Other (waste-to-energy), biogenic	TJ	14'163	13'394	16'889	19'797
Biomass (wood, renewable waste)	TJ	301	466	547	844
Biogas (co-generation from landfills, fermentation engines)	TJ	282	651	679	462

1A1a Public electricity and heat production	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Total fuel consumption	TJ	60'574	58'220	59'487	58'518	62'638	60'837	64'728	64'861	61'013	62'958
Gas oil	TJ	1'280	800	490	540	500	400	800	670	780	470
Residual fuel oil	TJ	300	220	180	130	40	10	NO	NO	NO	NO
Petroleum coke	TJ	NO									
Other bituminous coal	TJ	NO									
Lignite	TJ	NO									
Natural gas	TJ	8'663	7'910	8'468	8'073	9'926	7'512	8'213	8'460	5'092	7'070
Other (waste-to-energy), fossil	TJ	26'940	25'791	25'880	24'853	26'002	25'575	26'262	25'738	26'049	26'832
Biomass	TJ	23'390	23'498	24'470	24'922	26'169	27'340	29'452	29'993	29'092	28'586
Other (waste-to-energy), biogenic	TJ	21'940	21'415	21'464	21'249	22'275	22'272	23'051	22'489	23'112	23'716
Biomass (wood, renewable waste)	TJ	939	1'458	2'311	2'877	2'958	3'982	5'032	5'948	4'324	3'072
Biogas (co-generation from landfills, fermentation engines)	TJ	511	625	695	796	937	1'086	1'370	1'556	1'657	1'798

Since 1990 the use of waste-derived fuels increased considerably. This is due to the fact that since 1st of January 2000, disposal of combustible wastes in landfill sites is prohibited by law (TVA Art. 32). The increase is also partly due to municipal solid waste imported from neighbouring countries to optimize the load factor of MSWIPs. During the reporting period the consumption of natural gas increased, and the consumption of liquid fuels decreased. This is due to a fuel shift in combined heat and power generation and the closure of the only power station located in Vouvry that has been operated with residual fuel oil in the 1990ies.

Municipal solid waste incineration and special waste incineration

Figure 7-4 in Sector 5 Waste gives an overview over the waste amounts, their treatment and their reporting in the Swiss greenhouse gas inventory. Municipal solid waste includes waste generated in households and waste of similar composition from other sources.

The amount of municipal solid waste in kt reported in Table 3-35 is the total amount of waste burned (it includes fossil and biogenic shares). The fossil and biogenic share in TJ are given as well.

Special waste is composed of special wastes with high calorific value, wastewater and sludge with organic load, inorganic solids and dusts, inorganic sludge containing heavy metals, acids and alkalis, PCB-containing wastes, non-metallic shredder residues, contaminated soil, filter materials and chemicals residues and others.

Table 3-35 Activity data for 1A1a iv Other: Municipal solid waste and special waste incinerated with heat and/or power generation 1990–2015. The amount of municipal solid waste in kt is the total amount of waste burned.

1A1a iv Public electricity and heat production, Other	Unit	1990	1995	2000	2005
Total fuels	TJ	30'768	30'264	39'371	44'508
Municipal solid waste fossil	TJ	13'995	13'664	17'790	20'197
Municipal solid waste biogenic	TJ	14'163	13'394	16'889	19'797
Special waste	TJ	2'610	3'206	4'692	4'514
Total fuels	kt	2'603	2'433	3'040	3'527
Municipal solid waste (fossil and biogenic)	kt	2'470	2'270	2'801	3'297
Special waste	kt	133	163	239	230

1A1a iv Public electricity and heat production, Other	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Total fuels	TJ	48'880	47'206	47'344	46'102	48'277	47'847	49'313	48'228	49'161	50'548
Municipal solid waste fossil	TJ	22'028	21'159	20'871	20'334	21'062	20'724	21'108	20'593	21'163	21'717
Municipal solid waste biogenic	TJ	21'940	21'415	21'464	21'249	22'275	22'272	23'051	22'489	23'112	23'716
Special waste	TJ	4'912	4'632	5'009	4'519	4'941	4'851	5'155	5'145	4'886	5'115
Total fuels	kt	3'896	3'816	3'865	3'827	3'968	3'924	4'104	4'035	4'066	4'150
Municipal solid waste (fossil and biogenic)	kt	3'646	3'580	3'610	3'597	3'717	3'676	3'841	3'773	3'817	3'889
Special waste	kt	250	236	255	230	252	247	263	262	249	261

3.2.5.2.2 Petroleum refining (1A1b)

Methodology (1A1b)

Up to 2015, two refineries were in operation in Switzerland. Since one of the refineries ceased operation in 2015, the data are considered confidential since 2014. Data are available to reviewers on request. Based on the generalised decision tree Fig. 2.1 for stationary combustion (IPCC Guidelines 2006, vol.2, chp. 2), Switzerland applies a Tier 2 approach with country-specific emission factors for CO₂ emissions and a Tier 1 approach with 2006 IPCC default emission factors for CH₄ and N₂O emissions. The calculations are based on measurements and data from the refining industry as documented in the EMIS database (EMIS 2017/1A1b Heizkessel Raffinerien).

Emission factors (1A1b)

Emission factors of residual fuel oil, petroleum coke and refinery gas are estimated based on measurements from the two refineries for the years 2005 – 2011 and 2013 – 2015 provided in the framework of the Swiss emissions trading system. From 2005 onwards, the measured emission factors are applied. The emission factors for 2012 are interpolated between 2011 and 2013. In years before 2005, the emission factors of residual fuel oil and petroleum coke are based on the weighted mean of the available data (2005 – 2011 and 2013 – 2015). The CO₂ emission factor of refinery gas is based on an estimate provided by one of the two refining plants for the years 1990-2004, which is assumed to be constant. Since 2013 the annual emission factor is derived from annual monitoring reports and the allocation report (2005-2011), which provide plant-specific data.

The resulting CO₂ emission factors are all within the given ranges in IPCC (2006). For emission factors of N₂O and CH₄ default values according to IPCC 2006 are used.

Table 3-36 Emission factors for 1A1b Petroleum refining in 2015.

1A1b Petroleum refining	CO ₂	CH ₄	N ₂ O	NO _x	NMVOC	SO ₂	CO
	t/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ
Residual fuel oil	C	C	C	C	C	C	C
Refinery gas	C	C	C	C	C	C	C
Petroleum coke	C	C	C	C	C	C	C

Activity data (1A1b)

Activity data on fuel combustion for petroleum refining (1A1b) is provided by the Swiss overall energy statistics (SFOE 2016) and by the industry (bottom-up data). The data from the industry is collected by Carburia and forwarded to the Swiss Federal Office of Energy for inclusion in the Swiss overall energy statistics (SFOE 2016).

Refinery gas is the most important fuel used in source category 1A1b. Energy consumption, in particular use of refinery gas, has increased substantially since 1990 because one of the two Swiss refineries operated at reduced capacity in 1990 and resumed full production in later years. In 2012, one of the refineries was closed over six months due to insolvency and the search for a new buyer (EV 2014).

Table 3-37 Activity data for 1A1b Petroleum refining.

1A1b Petroleum refining	Unit	1990	1995	2000	2005
Total fuel consumption	TJ	5'629	9'836	9'636	14'548
Residual fuel oil	TJ	1'259	1'786	1'908	902
Refinery gas	TJ	4'370	8'050	7'728	11'833
Petroleum coke	TJ	NO	NO	NO	1'813

1A1b Petroleum refining	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Total fuel consumption	TJ	16'013	13'774	15'118	14'473	14'176	13'169	11'242	13'834	14'173	7'232
Residual fuel oil	TJ	692	1'182	692	733	891	764	1'212	1'094	C	C
Refinery gas	TJ	13'508	11'033	11'978	11'706	11'282	10'720	8'249	11'055	C	C
Petroleum coke	TJ	1'813	1'558	2'449	2'035	2'003	1'685	1'781	1'685	C	C

3.2.5.2.3 Manufacture of solid fuels and other energy industries (1A1c)

Methodology (1A1c)

In source category 1A1c Manufacture of Solid Fuels and other energy industries, only the emissions from charcoal production are reported as no other activities occur in Switzerland.

Based on the generalised decision tree in Fig. 2.1 for stationary combustion (IPCC Guidelines 2006, vol.2, chp. 2), emissions are estimated using a Tier 2 approach.

Emission factors (1A1c)

The CO₂ emission factor is based on literature (USEPA 1995) and CH₄, NO_x, CO and NMVOC emission factors are taken from the revised 1996 IPCC Guidelines (EMIS 2017/1A1c).

Table 3-38 Emission factors for 1A1c Manufacture of Solid Fuels and other energy industries in 2015. The CO₂ emission factor refers to CO₂ of biogenic origin.

1A1c Charcoal	Unit	CO ₂ biog.	CH ₄	NO _x	NM VOC	SO ₂	CO
Charcoal production	kg/TJ	16'900	1'000	10	1'700	NA	7'000

Activity data (1A1c)

The annual amount of charcoal produced is based on detailed queries with the few remaining sites where charcoal is produced. The main producer is the Köhlerverein Romoos, small quantities are produced at individual traditional local trade shows (Karthause Ittingen, Freilichtmuseum Ballenberg), as documented in EMIS 2017/1A1c. The FAO database contained values that differ substantially from these detailed bottom-up data. FAO has been informed about the discrepancy and was provided with the data used in the greenhouse gas inventory.

The charcoal is not used in the industry anymore but mainly for barbecues. Production has increased between 1990 and 2015 due to two regular charcoal production sites starting operation in 2004, low wood prices and increased demand for local charcoal in Switzerland (Koehlerei 2014).

Table 3-39 Activity data for 1A1c Manufacture of Solid Fuels and other energy industries.

1A1c Charcoal	Unit	1990	1995	2000	2005
Charcoal production	TJ	1.25	1.43	2.20	3.37

1A1c Charcoal	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Charcoal production	TJ	3.33	3.48	3.52	3.48	3.19	3.30	3.62	2.82	3.82	3.30

3.2.5.3 Uncertainties and time-series consistency for 1A1 (stationary)

The uncertainty of CO₂ emission factors is described in chp. 1.6.1.2 and 3.2.4.7. The uncertainty in emissions of non-CO₂ gases are estimated to be medium, i.e. 30% for CH₄ and 80% for N₂O (see also chp. 1.6, Table 1-10).

Consistency: Time series for 1A1 Energy industries are all considered consistent.

3.2.5.4 Category-specific QA/QC and verification for 1A1 (stationary)

The general QA/QC procedures are described in chp. 1.2.3. Furthermore QA/QC procedures conducted for all 1A source categories are listed in chp. 3.2.4.8.

Concerning activity data and emission factors in the refinery sector, collections of emissions and fuel combustion statistics at large combustion plants for pollution legislation purposes exist. This plant-level data was used to cross-check national energy statistics from this sector for representativeness.

3.2.5.5 Category-specific recalculations for 1A1 (stationary)

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 or 2014 of at least 1 kt CO₂ eq are quantified. All the other recalculations have an impact of less than 1 kt CO₂ eq in the years 1990 and 2014.

- 1A1a: Recalculation concerning boilers using natural gas due to recalculation in losses of natural gas in distribution network (2013, 2014).
- 1A1a: The time series value of waste generation rate for 2013 has slightly changed due to the correction of an error in waste statistics.
- 1A1a: The use of carbonates for sulphur oxide removal in 1A1a Municipal solid waste incineration plants has been moved to 2A4d Other process uses of carbonates (1990–2006). In 1990, this recalculation leads to a decrease in emissions in category 1A1a by 0.3 kt CO₂ eq and in 2014 the emissions decrease by 1.4 kt CO₂ eq.
- 1A1a: Recalculation in residual fuel oil boilers due to mistake in calculations in the energy model for the entire time series (1990–2014). This recalculation reduces emissions in 2014 by about 2 kt CO₂ eq and in 1990 by about 1 kt CO₂ eq.
- 1A1b: In last submission the emission factor of CO₂ of refinery gas and residual fuel oil used in refineries was linear interpolated from 1990–2005. It was changed to be constant from 1990–2004. This leads to a recalculation between 1991 and 2004.
- 1A1b: There was a transcription error in the calculation of the CO₂ emission factor of refinery gas in 2014. The correction of the error results in an increase of the emissions in 2014 by about 11 kt CO₂ eq.

3.2.5.6 Category-specific planned improvements for 1A1 (stationary)

The N₂O emission factor of MSWIP is assessed periodically. It is planned to calculate a new weighted average N₂O emission factor for the year 2016 to be used in the next submission.

3.2.6 Source category 1A2 - Manufacturing industry and construction (stationary without 1A2g vii)

3.2.6.1 Source category description for 1A2 (stationary)

Table 3-40 Key categories (KCA incl. LULUCF) of 1A2 Manufacturing industries and construction.

Code	IPCC Category	GHG	Identification Criteria
1A2	Manufacturing Industry and Construction: Gaseous Fuels	CO ₂	L1, L2, T1, T2
1A2	Manufacturing Industry and Construction: Liquid Fuels	CO ₂	L1, T1
1A2	Manufacturing Industry and Construction: Other Fuels	CO ₂	L1, T1
1A2	Manufacturing Industry and Construction: Solid Fuels	CO ₂	L1, T1, T2

[Source category 1A2 contains the sum of emissions of stationary and mobile sources – the statement on key categories holds for the aggregated emission only. However, the CO₂ emissions of 1A2 from Liquid Fuels are dominated by the stationary sources, 87%, which means that the emissions of 1A2g vii only play a minor role within category 1A2.]

The source category 1A2 Manufacturing industries and construction comprises all emissions from the combustion of fuels in stationary boilers and cogeneration facilities within manufacturing industries and construction. This includes use of conventional fossil fuels as well as waste derived fuels and biomass. Use of fossil fuels as feedstocks or other non-energy use of fuels as for example bitumen and lubricants are reported in CRF-table 1.A(d) and described in chp. 3.2.3.

Table 3-41 Specification of source category 1A2 Manufacturing industries and construction in Switzerland.

1A2	Source	Specification
1A2a	Iron and steel	Iron and steel industry: boilers, cupola furnaces in iron foundries and heating furnaces in steel production
1A2b	Non-ferrous metals	Non-ferrous metals industry: secondary aluminium production, copper alloys production
1A2c	Chemicals	Chemical industry: production of chemicals such as ammonia, nitric acid, ethylene, acetic acid and sulphuric acid as well as silicon carbide (amongst others)
1A2d	Pulp, paper and print	Pulp, paper and print industry
1A2e	Food processing, beverages and tobacco	Food processing, beverages and tobacco industry: meat production, milk products, convenience food, chocolate, sugar and baby food (amongst others).
1A2f	Non-metallic minerals	Fine ceramics, container glass, glass, glass wool, lime, rock wool, mixed goods, cement, brick and tile
1A2giv	Wood and wood products	Fibreboard production
1A2gviii	Other	Industrial fossil fuel and biomass boilers and engines that do not provide heat or electricity to the public.

3.2.6.2 Methodological issues for 1A2 (stationary)

3.2.6.2.1 Methodology (1A2) and Industry model

For fuel combustion in source category 1A2 Manufacturing industries and construction, a country-specific approach is used combining Tier 2 and Tier 3 methods (IPCC 2006, Volume 2 Energy, chp. 2 Stationary Combustion, Figure 2.1).

Overview Industry Model

The industry model is one sub-model of the Swiss energy model (see chp. 3.2.4.3). The industry model disaggregates the stationary fuel consumption into the source categories and processes under 1A2 Manufacturing industries and construction. The following figure visualizes the disaggregation process.




Figure 3-21 Schematic presentation of the data sources used for the industrial sectors 1A2a – 1A2g. The reference SFOE 201X refers to the 2016 edition of the corresponding energy statistics. For each fuel type, the Swiss overall energy statistics provide the total consumption for industry. The total consumption is then distributed to the different source categories based on information from industry surveys (SFOE 2016d) and the Prognos industry model. The grey boxes on the right show the specific bottom-up industry information.

The total fuel consumption regarding each fuel type in the industry sector is provided by the Swiss overall energy statistics (see description of the Swiss overall energy statistics in chp. 3.2.4.3). The energy disaggregation into the source categories 1A2a to 1A2g is carried out for each fuel type individually based on the energy consumption statistics in the industry and services sectors (SFOE 2016d). These statistics are available since 1999 for gas oil and natural gas. For all other fossil fuels (i.e. residual fuel oil, liquefied petroleum gas, petroleum coke, other bituminous and lignite) data are available since 2002. In order to generate a consistent time series since 1990, additional data from an industry model is applied (Prognos 2013) as described in the following sub-section.

In addition, the share of fuel used for co-generation in turbines and engines within 1A2 is derived from a model of stationary engines developed by Eicher + Pauli (Kaufmann 2015) for the statistics on combined heat and power generation (SFOE 2016c).

Energy consumption statistics in the industry and services sectors

The energy consumption statistics in the industry and services sectors (SFOE 2016d) refers to representative surveys with about 12'000 workplaces in the industry and services sectors that are then grossed up or extrapolated to the entire industry branch. For certain sectors and fuel types (i.e. industrial waste, residual fuel oil, other bituminous coal and lignite) the surveys represents a census covering all fuel consumed. The surveys are available for all years since 1999 or 2002, depending on the fuel type.

In 2015, a change in the survey method of the energy consumption statistics in the industry and services sectors was implemented (SFOE 2015d). In brief, the business and enterprise

register, which forms the basis for the samples of the surveys, was revised. While previously the business and enterprise register was based on direct surveys with work places, it is now based on annual investigations of registry data (e.g. from the old-age and survivors' insurance). In the course of this revision, a comparative assessment was conducted for the year 2013. This comparison shows that the energy consumption in the source categories of 1A2 stationary are modified by less than 1 percent, but also that the differences between the new and the old results for 2013 are not statistically significant (SFOE 2015d). As these statistics are only used for allocation of total energy consumption to different source categories, the impact on the different source categories consists only of a reallocation of the energy consumption and does not affect the total of the sector. Moreover, only consumption of gas oil and natural gas is affected. For all these reasons, the time series consisting of data based on the old (1990–2012) and new (2013–2015) survey method are therefore considered consistent.

Modelling of industry categories

The energy consumption statistics in the industry and services sectors (SFOE 2016d) are complemented by a bottom-up industry model (Prognos 2013). The model is based on 164 individual industrial processes and further 64 processes related to infrastructure in industry. Fuel consumption of a specific process is calculated as the product of the process activity data and the process specific fuel consumption factor.

It provides data on the disaggregation of total fuel consumption according to different industries and services between 1990 and 2012. For the time period, where the two disaggregation methods overlap, systematic differences between the two time series can be detected. These two data sets have been combined in order to obtain consistent time series of the shares of each source category 1A2a–1A2g for each fuel type. For this purpose, the approach to “generate consistent time series from overlapping time series” is used according to the 2006 IPCC Guidelines, Volume 1, chp. 5, consistent overlap (IPCC 2006). To illustrate the approach, an example for gas oil attributed to source category 1A2c is provided in the following figure. A detailed description for all fuel types and source categories (1A2a–1A2g), including further assumptions, is provided in the underlying documentation of the EMIS database (EMIS 2017/1A2 Sektorgliederung Industrie).




Figure 3-22 Illustrative example for combining time series with consistent overlap according to the 2006 IPCC Guidelines (IPCC 2006). The y-axis indicates the share of source category 1A2c of total gas oil consumption in the industry sector. The green line, which is based on the combination of the shares from the energy consumption statistics in the industry and services sectors (SFOE 2016d, red line from 1999 to 2015) and the bottom-up industry model (Prognos 2013, blue line from 1990 to 2012), corresponds to the share finally used to calculate the fuel consumption in 1A2c. Similar calculations are performed for each source category and fuel type, see also EMIS database documentation (EMIS 2017/1A2 Sektorgliederung Industrie).

Bottom-up industry data

Grey colored boxes in Figure 3-21 represent source categories, i.e. 1A2a–d, 1A2f and 1A2g for which bottom-up data from the industry are used in order to disaggregate the fuel consumption within a particular source category. These data consist of validated and verified monitoring data from the Swiss emissions trading scheme implemented under the Ordinance for the Reduction of CO₂ Emissions (Swiss Confederation 2012) and are discussed in depth in the following chapters 3.2.6.2.2 to 3.2.6.2.8.

The bottom-up information provides activity data for specific industrial production processes and forms a subset of the total fuel consumption allocated to each source category by the approach described above. Therefore, the fuel consumptions of the bottom-up industry processes are subtracted from the total fuel consumption of the respective source category and the remaining fuel consumptions are considered as fuels used in boilers of each source category. This method ensures that the sum of fuel consumption over all processes of a source category corresponds to the total fuel consumption as documented in the energy consumption statistics in the industry and services sectors (SFOE 2016d).

There is a difference in calculating the emissions of precursors from boilers and bottom-up industry processes. For boilers, fuel consumption is used as activity data whereas for bottom-up processes production data is used.

Further specific statistical data

Fuel consumption of wood, wood waste, biogas and sewage gas in manufacturing industries is based on the Swiss wood energy statistics (SFOE 2016b) as well as on data from the Swiss renewable energy statistics (SFOE 2016a) and the Statistics on combined heat and power generation in Switzerland (SFOE 2016c), respectively. Emissions from these sources

are reported under 1A2gviii Other due to insufficient information regarding sectoral disaggregation.

Emission factors (1A2)

The following table presents the emission factors of fuel consumption in source category 1A2 Manufacturing industry and construction (see also chp. 3.2.4.4).

Table 3-42 Emission factors for 1A2 Manufacturing industries and construction in 2015. Values that are highlighted in green are described in chp. 3.2.4.4.

1A2 Emission factors. Mix of bottom-up and top-down approach (modelling) for GHG	CO ₂ fossil	CO ₂ bio.	CH ₄	N ₂ O
	t/TJ	t/TJ	kg/TJ	kg/TJ
Gas oil	73.7		3	0.6
Residual fuel oil	77.0		<3 (lower IEF than default emission factor)	0.6
Liquefied petroleum gas	65.5		1	0.1
Petroleum coke	91.4		<3 (lower IEF than default emission factor)	0.6
Other bituminous coal	92.7		<10 (lower IEF than default emission factor)	1.5
Lignite	96.1		<10 (lower IEF than default emission factor)	1.5
Natural gas	56.4		1	0.1
Other fossil fuels (including solvents, plastics, waste tyres and rubber (see 1A2f))	71.9	3.6	2.5	3.4
Biomass (wood, biogas, biodiesel, bioethanol and other biogenic waste)		90.9	5.2	3.3

Other fossil fuels comprise various fossil waste derived fuels used in 1A2f Cement production as well as cracker by-products, i.e. gasolio and heating gas used for steam production in a chemical plant in source category 1A2c. The emission factors of CO₂, CH₄ and N₂O are implied emission factors based on the fossil waste fuel mix. In addition the CH₄ emission factor includes the total CH₄ emissions of the cement industry based on direct exhaust measurements at the chimneys of the cement plants (see documentation in EMIS 2017/1A2fi Zementwerke_Feuerung) based on industry data and emission declarations according to the Ordinance on Air Pollution Control. Implied CH₄ emission factors of source category 1A2 for residual fuel oil, petroleum coke, other bituminous coal and lignite are thus lower than the default emission factors of source category 1A documented in chp. 3.2.4.4.3 (see detailed description below in chapter Cement (1A2f i)).

The emission factors of the precursors NO_x, CO, NMVOC and SO₂ for all fuels in source category 1A2 are provided in Annex A3.1.1. The emission factors for NO_x and CO for natural gas and gas oil used in boilers are derived from a large number of air pollution control

measurements of combustion installations (Leupro 2012). This study analysed a large dataset from various cantons in Switzerland that was collected between 2000 and 2011. The emission factors for NO_x and CO for residual fuel oil, petroleum coke, other bituminous coal and lignite used in boilers are country-specific and documented in the Handbook on emission factors for stationary sources (SAEFL 2000) The implied emission factors for NO_x decreased significantly. NMVOC and SO₂ emission factors are country-specific and documented in SAEFL (2000).

In contrast to combustion in boilers, emission factors of precursors and SO₂ for fuel combustion in bottom-up industry processes are based on bottom-up industry data. Production-weighted emission factors based on various air pollution control measurements under the Ordinance on Air Pollution Control (Swiss Confederation 1985) are used to derive the corresponding process-specific emission factors.

Activity data (1A2)

The following table shows the total fuel consumption reported in source category 1A2 as described above in the industry model and displays the fuel switch within Swiss industry. Since 1990, the use of residual fuel oil and other bituminous coal has decreased. In the same period, natural gas consumption has more than doubled. Regarding the fuels used within Swiss industry in 2015, natural gas consumption accounts for the largest share followed by biomass and gas oil.

Source category 1A2gviii Other comprising emissions from boilers and engines is the most important category within source category 1A2 Manufacturing Industries and construction in 2015. 1A2f Non-metallic minerals and 1A2c Chemicals are the second and third most important fuel consumers.

Table 3-43 Activity data fuel consumption in 1A2 Manufacturing industries and construction.

Source	Unit	1990	1995	2000	2005
1A2 Manufacturing industries and constr. (stationary sources)	TJ	88'183	89'133	87'861	91'679
Gas oil	TJ	21'754	23'529	25'145	24'711
Residual fuel oil	TJ	18'870	13'678	5'675	4'613
Liquefied petroleum gas	TJ	4'520	4'706	5'921	4'599
Petroleum coke	TJ	1'400	1'260	551	1'093
Other bituminous coal	TJ	13'476	7'303	5'716	4'899
Lignite	TJ	265	153	124	742
Natural gas	TJ	18'721	27'898	31'383	34'372
Other fossil fuels	TJ	2'555	2'817	4'054	4'525
Biomass	TJ	6'622	7'788	9'292	12'126

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2 Manufacturing industries and constr. (stationary sources)	TJ	94'261	92'370	93'523	87'935	90'678	84'257	85'523	87'527	82'231	80'734
Gas oil	TJ	23'539	21'602	21'386	21'005	20'686	16'771	17'157	17'902	12'340	12'636
Residual fuel oil	TJ	5'427	3'776	3'734	2'713	2'096	1'518	1'568	848	351	226
Liquefied petroleum gas	TJ	5'070	4'554	4'310	4'595	4'181	4'136	3'998	3'998	3'538	3'584
Petroleum coke	TJ	1'512	1'271	1'067	1'219	1'495	1'272	1'367	1'049	1'240	795
Other bituminous coal	TJ	4'186	4'959	4'445	4'263	4'348	3'818	3'694	3'910	2'403	1'946
Lignite	TJ	1'931	1'937	1'717	1'531	1'460	1'624	1'175	1'357	3'102	3'060
Natural gas	TJ	35'840	36'910	38'719	35'126	38'042	36'903	38'013	39'400	39'946	39'137
Other fossil fuels	TJ	4'293	4'224	4'975	4'958	5'183	5'307	4'883	5'186	5'270	5'252
Biomass	TJ	12'462	13'135	13'170	12'527	13'188	12'909	13'667	13'877	14'042	14'098

The following chapters describe the fuel consumption of the different source categories 1A2a-1A2gviii, the specific industrial production processes based directly on bottom-up industry data and additional source-specific emission factors. Further information is documented in the respective EMIS documentation (EMIS 2017/1A2a-g).

3.2.6.2.2 Iron and steel (1A2a)

The source category 1A2a Iron and steel consists both of fuels used in boilers and specific industrial production processes, i.e. reheating furnaces in steel plants and cupola furnaces in iron foundries.

There is no primary iron and steel production in Switzerland. Only secondary steel and iron production using recycled steel scrap occurs. Iron is produced in 14 iron foundries. About 75% of the iron is processed in induction furnaces and 25% in cupola furnaces using other bituminous coal. Part of the other bituminous coal is also used to increase the carbon content of the raw material steel scrap to produce iron alloys with higher carbon content as well as reducing agent. Since other bituminous coal first of all acts as fuel in cupola furnaces it was decided to report its CO₂ emissions in source category 1A2a. Additionally, also limestone is used as flux in cupola furnaces yielding geogenic CO₂ emissions. These emissions are newly reported in source category 2A4d Other carbonate uses. The share of induction furnaces increased since 1990 with a sharp increase in 2009 due to the closure of at least one cupola furnace. Induction furnaces use electricity for the melting process and therefore only process emissions occur, which are reported in source category 2C1 Iron and steel production. Due to the reduced iron production and the switch from cupola to induction furnaces in iron foundries the consumption of other bituminous coal has decreased.

Today, steel is only produced in two steel production plants after closure of two plants in 1994. Both plants use electric arc furnaces (EAF) with carbon electrodes for melting the steel scrap. Therefore, only emissions from the reheating furnaces are reported in source category 1A2a. Since 1995, these furnaces use natural gas only for reheating the ingot moulds prior to the rolling mills. Process emissions from steel production are included in source category 2C1 Iron and steel production. Steel production and the related natural gas consumption was significantly reduced in 1995 and the use of residual fuel oil ceased with the closure of two steel companies. Since 1995, steel production increased continuously until 2004 to reach the same production level as 1990. Since then, steel production is about constant. Only in 2009, the production was considerably lower due to the economic crisis. One steel producer switched its production to high quality steel and therefore the specific energy use per tonne of steel produced increased between 1995 and 2000. This led to higher natural gas consumption.

Today fuel consumption of source category 1A2a consists mainly of natural gas but also liquefied petroleum gas and gas oil and small amounts of other bituminous coal are used.

Table 3-44 Activity data fuel consumption in 1A2a Iron and steel.

Source	Unit	1990	1995	2000	2005
1A2a Iron and steel	TJ	3'310	2'570	3'351	3'389
Gas oil	TJ	480	262	338	401
Residual fuel oil	TJ	346	131	20	39
Liquefied petroleum gas	TJ	408	193	286	217
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	433	289	266	154
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	1'643	1'695	2'440	2'578

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2a Iron and steel	TJ	4'208	4'415	4'156	3'190	3'773	3'805	3'634	3'530	3'653	3'976
Gas oil	TJ	311	326	307	279	315	271	172	139	86	221
Residual fuel oil	TJ	52	36	51	39	51	2	NO	NO	NO	NO
Liquefied petroleum gas	TJ	313	295	246	214	219	226	438	438	388	393
Petroleum coke	TJ	NO									
Other bituminous coal	TJ	150	160	177	70	64	73	55	55	52	44
Lignite	TJ	NO									
Natural gas	TJ	3'382	3'598	3'374	2'588	3'125	3'233	2'969	2'898	3'127	3'317

3.2.6.2.3 Non-ferrous metals (1A2b)

The source category 1A2b Non-ferrous metals consists both of fuels used in boilers and specific industrial production processes, i.e. secondary aluminium production and non-ferrous metal foundries, producing mainly copper alloys.

Until 1993, secondary aluminium production plants have been in operation using gas oil. Emissions from primary aluminium production in Switzerland are reported in source category 2C3 as induction furnaces have been used. Its last production site closed down in April 2006.

Regarding non-ferrous metal industry in Switzerland, only casting and no production of non-ferrous metals occur. There is one large company and several small foundries, which are organized within the Swiss foundries association (Schweizerischer Giessereiverband, GVS) providing production data.

Fuel consumption of source category 1A2b represents only a small amount of the total fuel consumption in source category 1A2 in 2015. Fuels consumed in 2015 are mainly natural gas as well as gas oil and small amounts of residual fuel oil and liquefied petroleum gas. Fuel consumption within this source category decreased since 1990 due to the closing down of the secondary aluminium production and the strong reduction of the non-ferrous metal production since 2000.

Table 3-45 Activity data fuel consumption in 1A2b Non-ferrous metals.

Source	Unit	1990	1995	2000	2005
1A2b Non-ferrous metals	TJ	2'379	1'969	1'560	977
Gas oil	TJ	587	347	236	125
Residual fuel oil	TJ	NO	NO	NO	NO
Liquefied petroleum gas	TJ	27	17	15	7
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	1'765	1'605	1'309	845

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2b Non-ferrous metals	TJ	1'162	1'022	1'042	1'006	1'218	1'177	1'746	1'593	1'917	1'765
Gas oil	TJ	72	94	112	167	112	76	153	128	90	75
Residual fuel oil	TJ	NO	NO	0	0	0	0	1	23	NO	44
Liquefied petroleum gas	TJ	10	8	7	7	8	8	11	11	10	10
Petroleum coke	TJ	NO									
Other bituminous coal	TJ	NO									
Lignite	TJ	NO									
Natural gas	TJ	1'080	920	924	833	1'098	1'093	1'581	1'430	1'817	1'636

3.2.6.2.4 Chemicals (1A2c)

In Switzerland, there are more than thirty chemical companies mainly producing fine chemicals and pharmaceuticals. Fossil fuels are mostly used for steam production and process heat. The process emissions from the production of chemicals such as ammonia, nitric acid, ethylene, acetic acid and sulphuric acid as well as silicon carbide are reported in source category 2B, see chp. 4.3.

There is one large company producing ammonia and ethylene by thermal cracking of liquefied petroleum gas and light virgin naphtha (see also descriptions in chp. 3.2.3 for feedstock use). As by-products from the cracking process, so-called heating gas and gasolio are produced, which are used thermally for steam production within the same plant. For reasons of confidentiality, fuel consumption and emissions of these by-products are included in Other fossil fuels of 1A2f in the reporting tables. Data are available to reviewers on request.

The CO₂ emission factors of gasolio and heating gas are plant-specific based on monitoring reports of the Swiss ETS.

Since the fuel quality of gasolio and heating gas are of similar quality as residual fuel oil and gas oil, respectively, the same default IPCC emission factors are assumed for CH₄ and N₂O (see Table 3-42 and Table 3-14 (CH₄ EF of residual fuel oil)).

Table 3-46 Emission factors for 1A2c Chemicals are documented in the confidential NIR, which is available to reviewers on request.

The fuels consumed in 2015 include mainly natural gas as well as minor amounts of gas oil. Fuel consumption in this source category has slightly decreased between 1990 and 2015. Consumption of gas oil and residual fuel oil have decreased in that period, while natural gas consumption has increased.

Table 3-47 Activity data fuel consumption in 1A2c Chemicals.

Source	Unit	1990	1995	2000	2005
1A2c Chemicals	TJ	14'436	15'158	13'500	15'477
Gas oil	TJ	3'942	3'313	3'215	3'345
Residual fuel oil	TJ	1'434	693	252	36
Liquefied petroleum gas	TJ	15	13	12	10
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	9'044	11'138	10'020	12'086
Other fossil fuels	TJ	IE	IE	IE	IE

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2c Chemicals	TJ	14'995	14'810	14'610	12'611	11'814	12'167	13'909	14'125	12'128	12'951
Gas oil	TJ	3'210	2'556	2'261	2'498	2'103	1'847	2'055	1'797	1'321	1'226
Residual fuel oil	TJ	71	6	79	91	66	0.2	0.2	1	NO	NO
Liquefied petroleum gas	TJ	11	10	9	9	8	7	10	10	9	9
Petroleum coke	TJ	NO									
Other bituminous coal	TJ	NO									
Lignite	TJ	NO									
Natural gas	TJ	11'704	12'239	12'261	10'014	9'637	10'312	11'845	12'317	10'798	11'716
Other fossil fuels	TJ	IE									

3.2.6.2.5 Pulp, paper and print (1A2d)

Around half a dozen paper producers and several printing facilities exist in Switzerland. The only cellulose production plant was closed in 2008. Thermal energy is mainly used for provision of steam used in the drying process within paper production. Emissions from use of carbonate in flue gas treatment in cellulose production is reported in 2A4d Other process use of carbonates.

Fuel consumption in 1A2d consists both of fuels used in boilers and specific industrial production processes. In this source category only biomass from cellulose production (until 2008) is included, based on data from the only production site. Biomass used in paper production is reported in source category 1A2gviii, because no comprehensive information exists to distribute biomass consumption to the specific industry sectors within 1A2 as explained in chapter 3.2.4.6.

The overall fuel consumption within the Swiss pulp and paper industry has considerably decreased since 1990, due to the closure of the cellulose production plant in 2008 and of several paper producers in the last years. The fuels used in 2015 are mainly natural gas as well as gas oil. Since 1990 residual fuel oil and gas oil have decreased, while natural gas consumption increased.

Table 3-48 Activity data of fuel consumption in 1A2d Pulp, paper and print.

Source	Unit	1990	1995	2000	2005
1A2d Pulp, paper and print	TJ	11'762	13'700	11'580	11'379
Gas oil	TJ	1'188	1'751	1'403	1'456
Residual fuel oil	TJ	5'250	3'061	1'417	2'092
Liquefied petroleum gas	TJ	86	141	148	100
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	3'153	7'389	6'918	5'678
Biomass	TJ	2'085	1'358	1'694	2'053

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2d Pulp, paper and print	TJ	11'493	10'236	9'455	6'124	6'773	6'051	5'374	5'474	4'644	3'579
Gas oil	TJ	1'291	1'096	1'019	948	852	561	623	711	297	372
Residual fuel oil	TJ	3'305	1'885	1'887	1'084	279	4	3	0	22	19
Liquefied petroleum gas	TJ	79	71	60	62	61	62	67	67	60	60
Petroleum coke	TJ	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Other bituminous coal	TJ	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Natural gas	TJ	4'742	5'085	5'164	4'030	5'581	5'424	4'681	4'696	4'265	3'128
Biomass	TJ	2'076	2'099	1'324	NO						

3.2.6.2.6 Food processing, beverages and tobacco (1A2e)

In Switzerland, the source category 1A2e Food, beverages and tobacco includes around 200 companies. According to the national food industry association, the major part of revenues is provided by meat production, milk products and convenience food. Further productions comprise chocolate, sugar or baby food (Fial 2013). Fossil fuels are used for steam production and drying processes. Fuel consumption in 1A2e is exclusively based on information from the energy consumption statistics in the industry and services sectors (SFOE 2016d) and Prognos (2013).

In 2015, the fuels used in this category were mainly natural gas as well as gas oil and small amounts of liquefied petroleum gas. There was an increase in fuel consumption between 1990 and 2015. This is due to the increased production in this sector. The consumption of residual fuel oil and gas oil ceased and has decreased, respectively, whereas natural gas and liquefied petroleum gas consumption has increased significantly.

Table 3-49 Activity data fuel consumption in 1A2e Food processing, beverages and tobacco.

Source	Unit	1990	1995	2000	2005
1A2e Food processing, beverages and tobacco	TJ	9'859	8'784	10'439	10'239
Gas oil	TJ	7'410	5'511	5'515	4'070
Residual fuel oil	TJ	1'160	466	137	NO
Liquefied petroleum gas	TJ	204	308	535	534
Petroleum coke	TJ	NO	NO	NO	NO
Other bituminous coal	TJ	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO
Natural gas	TJ	1'085	2'500	4'251	5'635

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2e Food processing, beverages and tobacco	TJ	11'519	11'221	10'975	12'558	13'161	11'374	11'310	13'079	12'440	11'600
Gas oil	TJ	3'811	3'500	3'376	3'687	3'778	3'197	3'237	3'681	2'395	2'413
Residual fuel oil	TJ	NO									
Liquefied petroleum gas	TJ	678	596	535	736	659	675	935	935	828	838
Petroleum coke	TJ	NO									
Other bituminous coal	TJ	NO									
Lignite	TJ	NO									
Natural gas	TJ	7'031	7'126	7'064	8'135	8'723	7'502	7'138	8'463	9'218	8'348

3.2.6.2.7 Non-metallic minerals (1A2f)

The source category 1A2f Non-metallic minerals includes several large fuel consumers within mineral industry, e.g. cement, brick and tile, glass and rock wool production. All fuel consumption of these specific industrial production processes are based on bottom-up industry data.

The fuels consumed in this source category are very diverse, depending on the fuel use within the specific industry process (see detailed documentation below). Fuel consumption in 2015 comprises mainly other fossil fuels, natural gas, lignite, biomass and other bituminous coal.

Between 1990 and 2015 there has been a switch in fuel consumption from other bituminous coal and residual fuel oil to other fossil fuels, natural gas, lignite and biomass. The most important emission source within this category is cement production. Information on bottom-up data of fuel consumption and some source-specific emission factors are described in the following. Detailed data at process level cannot be provided, since they are mostly confidential. Therefore, aggregated data for 1A2f are shown in the following table.

Table 3-50 Activity data fuel consumption in 1A2f Non-metallic minerals.

Source	Unit	1990	1995	2000	2005
1A2f Non-metallic minerals	TJ	25'613	19'884	18'056	17'832
Gas oil	TJ	1'871	1'629	1'642	1'389
Residual fuel oil	TJ	5'382	5'578	3'649	2'420
Liquefied petroleum gas	TJ	523	498	468	324
Petroleum coke	TJ	550	300	480	638
Other bituminous coal	TJ	12'665	6'758	5'415	4'364
Lignite	TJ	265	153	124	737
Natural gas	TJ	1'769	1'566	1'496	1'861
Other fossil fuels	TJ	2'555	2'817	4'054	4'525
Biomass	TJ	33	585	728	1'575

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2f Non-metallic minerals	TJ	17'770	18'036	17'902	17'102	18'196	17'801	16'956	17'119	17'785	16'599
Gas oil	TJ	1'483	1'343	1'299	1'260	1'269	1'238	1'097	1'174	1'276	1'182
Residual fuel oil	TJ	1'704	1'744	1'598	1'374	1'519	1'403	1'456	801	209	130
Liquefied petroleum gas	TJ	227	181	160	95	102	127	108	113	45	52
Petroleum coke	TJ	903	912	1'036	994	1'130	1'081	920	815	1'052	622
Other bituminous coal	TJ	3'661	4'348	3'912	3'940	3'992	3'474	3'403	3'478	1'973	1'498
Lignite	TJ	1'834	1'790	1'596	1'379	1'348	1'493	1'081	1'283	2'912	2'856
Natural gas	TJ	2'057	2'017	1'919	1'731	2'048	1'938	2'085	2'506	3'147	3'151
Other fossil fuels	TJ	4'293	4'224	4'975	4'958	5'183	5'307	4'883	5'186	5'270	5'252
Biomass	TJ	1'609	1'476	1'406	1'371	1'604	1'739	1'923	1'764	1'901	1'856

Cement (1A2f)

Methodology

In Switzerland, there are six plants producing clinker and cement. The Swiss plants are rather small and do not exceed a production capacity of 3'000 tonnes of clinker per day. All of them use modern dry process technology. Cement industry emissions stem from incineration of a wide variety of fossil and waste derived fuels used to generate high temperatures needed for the calcination process.

Emission factors

The CH₄ emission factor includes the overall CH₄ emissions of the cement industry based on direct exhaust measurements at the chimneys of the cement plants. Therefore, these CH₄ emissions are reported under the fuel type other fossil fuels in the CRF-tables.

Table 3-51 Emission factors for cement industry in 2015. Emission factors for CO₂ and N₂O are fuel-specific (see Table 3-43).

Cement industry (part of 1A2f)	CO ₂	N ₂ O	CH ₄	NO _x	NM VOC	SO ₂	CO
	t/TJ		g/t clinker				
Cement	fuel specific		6	930	59	270	1'900

The NCVs and CO₂ emission factors for waste oil, solvents and residues from distillation, plastics, mix of special waste with saw dust (CSS), sewage sludge, wood waste, animal meal and saw dust are based on a study of Cemsuisse (Cemsuisse 2010a). The values for waste tyres are taken from Hackl and Mauschitz (2003). The biogenic fraction of waste tyres is based on an Austrian study and published by the German Ministry of Environment (UBA 2006). The emission factor of N₂O for all waste derived fuels is the same and is taken from IPCC 2006 guidelines (IPCC 2006, vol 2, chp.2 table 2.3 industrial wastes).

Table 3-52 NCV, CO₂ and N₂O emission factors as well as biomass fraction of waste derived fuels (Other fossil fuels and Biomass) used in the cement industry in 2015.

Cement industry (part of 1A2f) Waste derived fuel	NCV	EF CO ₂ Tot.	EF N ₂ O	Fraction biomass-C
	MJ/kg	kg CO ₂ /GJ	g/GJ	%
Waste oil	32.48	74.35	4	0
Waste coke from coke filters	23.7	97	4	0
Mixed industrial waste	18.34	74	4	0
Other fossil waste fuels	20.85	97	4	0
Solvents and residues from distillation	23.63	73.99	4	0.9
Waste tyres and rubber	26.4	84	4	27
Plastics	25.24	84.66	4	27.7
Mix of special waste with saw dust (CSS)	9.22	102.4	4	78.5
Sewage sludge (dried)	9.39	94.52	4	100
Wood waste	16.26	99.9	4	100
Animal meal	16.81	86.66	4	100
Sawdust	16.26	99.9	4	100
Agricultural waste / other biomass	12.72	110	4	100

Activity data

Data on fuel consumption is provided by the industry, for recent years based on monitoring reports of the Swiss ETS as documented in the EMIS database (EMIS 2017/1A2f Zementwerke Feuerung).

In 2015, the Swiss cement industry used about similar amounts of standard fossil and waste derived fuels. Fossil fuels used in cement industry are mainly lignite, other bituminous coal

and petroleum coke. In addition, also fossil and biogenic waste derived fuels are used. Fossil wastes comprise plastics, solvents and residues from distillation, waste tyres and rubbers, and waste oil whereas biogenic wastes contain mainly wood waste, animal residues and sewage sludge. The main fossil fuels used in 1990 were other bituminous coal, residual fuel oil and other fossil fuels.

Fuel consumption in cement plants has decreased between 1990 and 2015. This is partly due to a decrease in production since 1990 and an increase in energy efficiency. In the same period the fuel mix has changed significantly from mainly fossil fuels to the above mentioned mix of fuels, including biogenic fractions of waste derived fuels.

In the reporting tables, the mainly biogenic waste derived fuels are reported under fuel type Biomass, whereas mainly fossil waste derived fuels are reported under fuel type Other fossil fuels (however, both fuel types also contain a fossil and a biogenic fraction, respectively, see Table 3-52).

Table 3-53 Activity data: Overview on fuel use in 1A2f cement industry.

Cement industry (part of 1A2f)	Unit	1990	1995	2000	2005
Cement, total incl. waste	TJ	17'193	12'772	11'018	11'623
Cement fossil without waste	TJ	15'319	9'993	7'332	6'208
Gas oil	TJ	NO	NO	NO	72
Residual fuel oil	TJ	1'907	2'825	1'530	637
Petroleum coke	TJ	550	300	480	638
Other bituminous coal	TJ	12'235	6'547	5'176	4'120
Lignite	TJ	265	153	124	737
Gas	TJ	362	168	22	4
Cement, waste derived fuel	TJ	1'874	2'780	3'686	5'415
Other fossil fuels	TJ	1'841	2'195	2'998	3'931
Waste oil	TJ	1'169	1'485	1'519	1'411
Waste coke from coke filters	TJ	59	59	59	58
Mixed industrial waste	TJ	NO	NO	NO	NO
Other fossil waste fuels	TJ	NO	NO	NO	NO
Solvents and residues from distillation	TJ	284	181	427	976
Waste tyres and rubber	TJ	330	415	421	645
Plastics	TJ	NO	55	572	841
Biomass	TJ	33	585	688	1'484
Mix of special waste with saw dust (CSS)	TJ	23	136	158	133
Sewage sludge (dried)	TJ	9	128	332	494
Wood waste	TJ	NO	321	NO	NO
Animal meal	TJ	NO	NO	198	856
Sawdust	TJ	NO	NO	NO	NO
Agricultural waste / other biomass	TJ	NO	NO	NO	NO

Cement industry (part of 1A2f)	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Cement, total incl. waste	TJ	11'719	12'022	11'954	11'816	12'388	12'187	11'462	11'866	12'339	11'348
Cement fossil without waste	TJ	6'401	6'914	6'389	6'127	6'278	5'859	5'406	5'512	5'847	4'917
Gas oil	TJ	57	NO	NO	NO	5	1	0.1	88	75	87
Residual fuel oil	TJ	220	175	135	100	112	101	297	86	58	45
Petroleum coke	TJ	903	912	1'036	994	1'130	1'081	920	815	1'052	622
Other bituminous coal	TJ	3'383	4'033	3'618	3'650	3'662	3'167	3'097	3'203	1'713	1'267
Lignite	TJ	1'834	1'790	1'596	1'379	1'348	1'493	1'081	1'283	2'912	2'856
Gas	TJ	4	4	4	4	21	16	11	38	37	41
Cement, waste derived fuel	TJ	5'319	5'108	5'565	5'689	6'109	6'329	6'056	6'354	6'492	6'431
Other fossil fuels	TJ	3'814	3'727	4'237	4'394	4'580	4'685	4'225	4'599	4'596	4'582
Waste oil	TJ	1'279	844	866	1'278	1'253	1'170	839	876	923	1'142
Waste coke from coke filters	TJ	60	NO								
Mixed industrial waste	TJ	NO	2	1	1	NO	NO	NO	NO	NO	NO
Other fossil waste fuels	TJ	NO	48	105	137	45	55	36	25	19	12
Solvents and residues from distillation	TJ	981	1'295	1'476	1'032	1'189	1'264	1'294	1'414	1'273	1'292
Waste tyres and rubber	TJ	568	525	794	828	842	1'033	964	985	1'021	958
Plastics	TJ	926	1'013	995	1'119	1'252	1'163	1'092	1'299	1'360	1'177
Biomass	TJ	1'504	1'381	1'328	1'295	1'530	1'644	1'831	1'756	1'896	1'850
Mix of special waste with saw dust (CSS)	TJ	146	164	157	131	123	96	100	96	103	80
Sewage sludge (dried)	TJ	560	549	511	475	477	483	527	418	428	420
Wood waste	TJ	NO	NO	NO	61	292	409	586	732	886	896
Animal meal	TJ	799	664	658	621	624	614	572	479	457	412
Sawdust	TJ	NO	NO	NO	NO	6	24	17	32	21	42
Agricultural waste / other biomass	TJ	NO	5	2	7	7	18	28	NO	NO	NO

Lime (1A2f)

In Switzerland there is only one plant producing lime. Fossil fuels are used for the burning process (calcination) of limestone. Between 1994 and 2012 fuel consumption in lime production was dominated by residual fuel oil. However in 2013, the main kiln has been switched to natural gas. Since 1995, no other bituminous coal is used anymore as it was replaced by residual fuel oil.

Container Glass (1A2f)

Today, there exists only one production plant for container glass in Switzerland. In 2014, fuel consumption for container glass production includes only natural gas. Since 1990, fuel consumption has drastically decreased due to reduction in production. Until 2003 only residual fuel oil was used and since 2004 the share of natural gas has increased to reach a stable share between 2006 and 2012. The large increase in natural gas between 2012 and 2013 is due to the fact that the plant has switched its glass kiln completely to natural gas in autumn 2013.

Tableware Glass (1A2f)

Today, there exists only one production plant for tableware glass in Switzerland. Fuel consumption for tableware glass currently includes only liquefied petroleum gas as residual fuel oil was eliminated in 1995. Since 1990, fuel consumption has strongly decreased because of the closure of one production plant in 2006.

Glass wool (1A2f)

In Switzerland, Glass wool is produced in two plants. Currently, fuel consumption for glass wool production includes only natural gas. Production of glass wool has increased since 1990, but the natural gas consumption decreased. This can be explained by an increase in energy efficiency in the production process between 1990 and 2015.

Fine ceramics (1A2f)

In Switzerland, the main production of fine ceramics is sanitary ware produced by one big and some small companies. In earlier years, also other ceramics were produced as for example glazed ceramic tiles, electrical porcelain and earthenware. Since 2001, only sanitary ware is produced.

Since 2010, fuel consumption within fine ceramics production is natural gas only. In 2001 the fuel mix consisted of natural gas and gas oil. Since then, it has continuously shifted to natural gas. Compared to the production of other fine ceramics, the production of sanitary ware is more energy-intensive. Therefore, the specific energy use per tonne of produced fine

ceramics has increased since 1990. This results in a lower reduction of fuel consumption compared to the reduction in production between 1990 and 2015.

Brick and tile (1A2f)

Methodology

In Switzerland there are about 20 plants producing bricks and tiles. Mainly fossil fuels but also paper production residues, animal grease and wood are used for drying and burning of the clay blanks.

Emission factors

The CO₂ emission factors for wood and animal grease are based on a study of Cemsuisse (Cemsuisse 2010a), see Table 3-52, whereas the one for paper production residues is taken from a German study on secondary fuels (UBA 2006) as documented in the EMIS database (EMIS 2017/1A2f Ziegeleien).

For CH₄ and N₂O emission factors of paper production residues and animal grease default values for wood waste and other liquid fuels, respectively, according to IPCC 2006 are used. For wood the CH₄ and N₂O emission factors according to the energy model for wood combustion (automatic chip boiler >500 kW, w/o wood processing companies), see chp. 3.2.4.5.2, are taken.

Activity data

Since 2013, plant-specific activity data are available from monitoring reports of the Swiss ETS. Fuels used in the brick and tile production in 2015 are mainly natural gas but also residual fuel oil and gas oil. Apart from a production recovery in the years around 2004, the production has gradually decreased since 1990, which is also represented in the overall fuel consumption decrease. Regarding the fuels used, there has been a considerable shift from residual fuel oil to natural gas from 1990 onwards as well as to a lesser extent, a shift from liquefied petroleum gas and gas oil to natural gas from 2004 onwards. Small amounts of paper production residues, wood and animal grease are used since 2000.

Rock wool (1A2f)

In Switzerland there is one single producer of rock wool. Cupola furnaces are used for the melting of rocks at a temperature of 1500°C.

Currently, other bituminous coal and natural gas are used in the production process. Until 2004 also gas oil and liquefied petroleum gas were used. In 2005, these fuels were substituted by natural gas.

Mixed goods (1A2f)

The production of mixed goods mainly includes the production of bitumen for road paving. A total of 110 production sites are producing mixed goods at stationary production sites.

The main fuels used in 2015 are gas oil and natural gas. Since 1990, there has been a fuel switch from gas oil to natural gas.

3.2.6.2.8 Other (1A2g stationary)

Methodology (1A2g stationary)

Source category 1A2g_{iv} Wood and wood products includes fuel consumption of fibreboard production. Fibreboards are produced in two companies in Switzerland, where thermal energy is used for heating and drying processes.

Source category 1A2g_{viii} Other covers fossil fuel combustion in boilers of manufacturing industries and construction mainly within non-metallic mineral industries as well as combustion of wood, wood waste, biogas and sewage gas in all manufacturing industries.

This source category accounts for about one third of the overall fuel consumption in 2015 of 1A2 Manufacturing industries and construction.

Methodologically, the fossil fuel consumption in boilers of 1A2g_{viii} comprises also all the residual entities of the industry installations that could not be allocated to any other source categories 1A2a-f.

Emission factors (1A2g stationary)

The CO₂ emission factors for wood waste and animal grease in 1A2g_{iv} Fibreboard production are based on a study of Cemsuisse (Cemsuisse 2010a), see Table 3-52. For wood waste the respective CH₄ and N₂O emission factors of the energy model for wood combustion, see chp. 3.2.4.5.2, are taken whereas for animal grease the default values of IPCC 2006 for other liquid biofuels are used. For biogas and sewage gas in 1A2g_{viii} Other boilers and engines industry the same emission factors as for natural gas are assumed.

Activity data (1A2g stationary)

1A2g_{iv} Fibreboard production

In source category fibreboard production, mainly wood waste as well as natural gas are used. Since 1990, the production of fibreboard and thus the fuel consumption have increased significantly. The fuel mix has strongly shifted between 1990 and 2015 from fossil fuels to biomass (wood waste). Between 2001 and 2013, also animal grease was used for fibreboard production. Since 2004, data on annual fuel consumption is taken from monitoring reports of the industry as documented in the EMIS database (EMIS 2017/1A2g_{iv}).

1A2gviii Other boilers and engines industry

Activity data for wood combustion is based on Swiss wood energy statistics (SFOE 2016b) whereas sewage and biogas consumption is based on data from the Swiss renewable energy statistics (SFOE 2016a) and the Statistics on combined heat and power generation in Switzerland (SFOE 2016c). Further information on wood energy consumption is provided in chapter 3.2.4.5.2.

Since 1990, the consumption of residual fuel oil and liquefied petroleum gas decreased. Solid fossil fuel consumption also decreased, whereas biomass and natural gas consumption increased.

Table 3-54 Activity data fuel consumption in 1A2g iv Wood and wood products and 1A2g viii Other (stationary).

Source	Unit	1990	1995	2000	2005
1A2g iv: Wood and wood products, 1A2g viii: Other (stationary)	TJ	20'825	27'068	29'376	32'386
Gas oil	TJ	6'276	10'716	12'796	13'925
Residual fuel oil	TJ	5'298	3'749	199	26
Liquefied petroleum gas	TJ	3'256	3'536	4'457	3'407
Petroleum coke	TJ	850	960	71	456
Other bituminous coal	TJ	378	256	34	380
Lignite	TJ	NO	NO	NO	5
Natural gas	TJ	262	2'005	4'948	5'689
Other fossil fuels	TJ	NO	NO	NO	NO
Biomass	TJ	4'505	5'845	6'869	8'499

Source	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2g iv: Wood and wood products, 1A2g viii: Other (stationary)	TJ	33'112	32'629	35'383	35'343	35'743	31'883	32'592	32'607	29'665	30'265
Gas oil	TJ	13'362	12'688	13'012	12'166	12'256	9'581	9'821	10'272	6'875	7'147
Residual fuel oil	TJ	295	105	119	124	182	109	109	22	120	33
Liquefied petroleum gas	TJ	3'753	3'392	3'293	3'473	3'124	3'031	2'428	2'424	2'199	2'222
Petroleum coke	TJ	609	360	31	224	365	191	447	234	189	173
Other bituminous coal	TJ	375	452	355	252	293	270	236	376	378	403
Lignite	TJ	97	147	121	152	111	131	95	75	189	204
Natural gas	TJ	5'845	5'926	8'012	7'796	7'828	7'400	7'713	7'090	7'575	7'841
Other fossil fuels	TJ	NO									
Biomass	TJ	8'777	9'560	10'440	11'156	11'584	11'170	11'744	12'113	12'141	12'242

3.2.6.3 Uncertainties and time-series consistency for 1A2 (stationary)

The uncertainty of CO₂ emissions from fuel combustion is described in the uncertainty analysis of source category 1A Fuel combustion in chp. 3.2.4.7. Uncertainty in emissions of other non-CO₂ gases is estimated to be medium, i.e. 30% for CH₄ and 80% for N₂O (see Table 1-10).

Consistency: Time series for 1A2 Manufacturing industries and construction are all considered consistent.

3.2.6.4 Category-specific QA/QC and verification for 1A2 (stationary)

The general QA/QC procedures are described in chp. 1.2.3. Furthermore QA/QC procedures conducted for all 1A source categories are listed in chp. 3.2.4.8.

3.2.6.5 Category-specific recalculations for 1A2 (stationary)

Recalculations at the aggregation level of 1A2 amount to -8 kt CO₂ eq for natural gas and about -1 kt CO₂ eq for petroleum coke in 2014. Reallocation of consumption of natural gas and gas oil at the level of the source categories 1A2a-1A2g viii causes differences, which however do not have any influence on total emissions reported in 1A2.

- 1A2: Recalculations due to new available statistics (SFOE2016d) (2012, 2013), causing a reallocation of energy to the different categories 1A2a-1A2g. Total energy consumption in 1A2 is not affected by this recalculation.
- 1A2: Amount of used gas oil in households, industry and commercial sector in Liechtenstein has been redistributed. Therefore the amount of used gas oil in boilers in households, industry and the commercial sector for Switzerland has changed too (1990–2014).
- 1A2d: The use of limestone for sulphur oxide removal in 1A2d Cellulose production has been moved to 2A4d Other process uses of carbonates (1990–2008). This reallocation leads to a decrease in emissions by 4 kt CO₂ eq in 1990 in 1A2d (and a corresponding increase in 2A4d).
- 1A2d: The so far missing emissions of CH₄ and N₂O from biogenic waste used in 1A2d Cellulose production are now reported for 1990–2008 based on default emission factors from the 2006 IPCC Guidelines for sulphite lyes. This recalculation leads to an increase by about 1 kt CO₂ eq in 1990.
- 1A2f: Activity data for 2014 of other bituminous coal and petroleum coke in cement production has changed due to the correction of an assignment error between the two fuels. This results in a recalculation of about +18 kt CO₂ eq for bituminous coal and -18 kt CO₂ eq for petroleum coke.
- 1A2f: AD (gas oil, natural gas) of 1A2f Production of mixed goods have been revised for 2014 based on corrected data from industry association.
- 1A2f: The emission factors of NO_x, NMVOC and CO of 1A2f Production of mixed goods have been revised from 1991 onwards based on air pollution control measurements (2001–2015).
- 1A2f: Revised interpolated CO emission factors of 1A2f Rockwool production in 2014 due to new plant-specific data for 2015.
- 1A2f: The emission factors of NMVOC and SO₂ as well as CO from 1A2f Glass production (speciality tableware) have been revised from 1991 and 1996, respectively, onwards.
- 1A2f: Fuel mix and consumption of 1A2f Lime production have been revised for 1990–1994 and 1990–2008 based on industry data and current net calorific values of gas oil, respectively, resulting in minor changes of AD of gas oil and residual fuel oil for 1990–2008 and 1994, respectively. In addition, it was found that not petroleum coke was used up to 1994 but other bituminous coal. All these changes induce recalculations in AD (gas oil, residual fuel oil, petroleum coke and other bituminous coal) of 1A2gviii Industrial boilers in the respective years. In 1990, this recalculation leads to a shift in emissions from bituminous coal from 1A2gviii to 1A2f (23 kt CO₂ eq).
- 1A2f: The conversion factor used for calculation of NMVOC emissions from total carbon based on air pollution control measurements has been revised resulting in adjusted NMVOC emission factors of 1A2f Brick and tile production and 1A2f Fine ceramics production for the entire time series

- 1A2g viii: Recalculations in AD (1990–2008: gas oil, residual fuel oil, 1990–1994: petroleum coke and other bituminous coal) of 1A2gviii Industrial boilers due to revised fuel mix and consumption in 1A2f Lime production. In 1990, this recalculation leads to an shift in emissions from bituminous coal from 1A2gviii to 1A2f (23 kt CO₂ eq) and a reduction of emissions from petroleum coke by 21 kt CO₂ eq and gas oil by 1 kt CO₂ eq.

3.2.6.6 Category-specific planned improvements for 1A2 (stationary)

No category-specific improvements are planned.

3.2.7 Source category 1A4 – Other sectors (stationary 1A4 ai/bi/ci)

3.2.7.1 Source category description for 1A4 (stationary)

Table 3-55 Key categories (KCA incl. LULUCF) of 1A4 Other sectors.

Code	IPCC Category	GHG	Identification Criteria
1A4a	Commercial: Gaseous Fuels	CO ₂	L1, L2, T1
1A4a	Commercial: Liquid Fuels	CO ₂	L1, T1
1A4b	Residential: Biomass	CH ₄	T1
1A4b	Residential: Gaseous Fuels	CO ₂	L1, L2, T1, T2
1A4b	Residential: Liquid Fuels	CO ₂	L1, T1
1A4c	Agriculture and Forestry: Liquid Fuels	CO ₂	L1

[Each of the source categories 1A4a, 1A4b, 1A4c contain the sum of emissions of stationary and mobile sources – the above statements on key categories hold for the aggregated emissions of 1A4a etc. only. The CO₂ emissions of 1A4a and 1A4b from Liquid Fuels are vastly dominated by the stationary sources, which means that the emissions of 1A4aii and 1A4bii only play a minor role within category 1A4a and 1 A4b. For 1A4c, however, the emissions of 1A4cii are dominating those of 1A4ci. See chp. 3.2.10.1]

Table 3-56 Specification of source category 1A4 Other sectors.

1A4	Source	Specification
1A4ai	Commercial/institutional	Emissions from stationary combustion in commercial and institutional buildings.
1A4bi	Residential	Emissions from stationary fuel combustion in households.
1A4ci	Agriculture/forestry/fishing	Emissions from stationary fuel combustion of agriculture and grass drying.

3.2.7.2 Methodological issues for 1A4 (stationary)

Methodology (1A4 stationary)

Emissions from stationary combustion in source categories 1A4ai, 1A4bi and 1A4ci are estimated based on country-specific CO₂ emission factors using a Tier 2 approach according to the decision tree for stationary combustion of the 2006 IPCC Guidelines (IPCC 2006, Volume 2 Energy, chp. 2 Stationary Combustion, Figure 2.1). Direct emission measurements are not available. A Tier 1 approach is applied with 2006 IPCC defaults EFs for CH₄ emissions of natural gas and gasoil for boilers and N₂O emissions of all fuels and technologies.

For the calculation of the emissions from the use of gas oil and natural gas the following sources are differentiated: (a) heat only boilers, (b) combined heat and power production in turbines and (c) combined heat and power production in engines.

Emissions from 1A4ci originate from fuel combustion for grass drying and wood combustion for heating in agriculture and forestry. For grass drying, information is provided by the grass drying association.

Emission factors (1A4 stationary)

Table 3-57 Emission factors for stationary combustion in 1A4ai Other sectors commercial/institutional in 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

Source/fuel	CO ₂	CO ₂ biog.	CH ₄	N ₂ O	NO _x	CO	NMVOC	SO ₂
	t/TJ		kg/TJ					
1A4a Other sectors: Commercial/institutional								
Gas oil (weighted average)	73.7		10	0.6	34	6.6	6	17.5
Gas oil (heat only boilers)	73.7		10	0.6	34	6.5	6	17.6
Gas oil (engines)	73.7		10	0.6	40	30.0	8	16.6
Natural gas (weighted average)	56.4		2.1	0.1	22.1	12.7	1.9	0.5
NG (heat only boilers)	56.4		1	0.1	17.5	10.0	2	0.5
NG (turbines)	56.4		2.0	0.1	60.0	15.0	0.1	0.5
NG (engines)	56.4		20	0.1	96.7	57.1	1	0.5
Other bituminous coal	NO		NO	NO	NO	NO	NO	NO
Lignite	NO		NO	NO	NO	NO	NO	NO
Biomass (weighted average)		86.8	19.5	3.1	93.9	622.4	29.3	7.8
Biomass (wood)		92.9	23.2	3.7	109.3	745.6	34.8	9.3
Biomass (biogas)		56.4	1	0.1	17.5	10.0	2.0	0.5

Table 3-58 Emission factors for stationary combustion in 1A4bi Other sectors residential in 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

Source/fuel	CO ₂	CO ₂ biog.	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂
	t/TJ		kg/TJ					
1A4b Other sectors: Residential								
Gas oil (weighted average)	73.7		10	0.6	35.5	12	6	17.5
Gas oil (heat only boilers)	73.7		10	0.6	35.5	12	6	17.6
Gas oil (engines)	73.7		2	0.6	40	30	8	16.6
Natural gas (weighted average)	56.4		1.2	0.1	16.7	13.4	4	0.5
NG (heat only boilers)	56.4		1	0.1	16.5	13	4	0.5
NG (turbines)	56.4		2	0.1	60	15	0.1	0.5
NG (engines)	56.4		20	0.1	34.2	57.1	1	0.5
Other bituminous coal	92.7		300	1.5	65	2000	100	350
Lignite	NO		NO	NO	NO	NO	NO	NO
Biomass (Wood)		95.2	49.7	3.8	89.1	1332.7	74.5	9.5

Table 3-59 Emission factors for stationary combustion in 1A4ci Agriculture/forestry/fishing in 2015.

1A4c Agriculture/forestry/fishing	Unit	CO ₂ fossil	CO ₂ biog.	CH ₄	N ₂ O
Grass drying (fossil, biogenic)	kg/TJ	54'978	65'145	6.1	0.8
Biomass	kg/TJ	NA	137'864	21	5.5

Charcoal and bonfires

Emission factors concerning CO₂, CH₄ and N₂O emissions of charcoal use in the residential source categories (1A4bi) are taken from the 2006 IPCC Guidelines (IPCC 2006). Default emission factors according to the guidelines are also applied for CH₄ and N₂O emissions resulting from bonfires. The CO₂ emission factor for bonfires in the residential category (1A4bi) is based on literature (SAEFL 2000). Emission factors of precursors are taken from the EMEP/EEA Guidebook (2013).

Table 3-60 Emission factors for use of charcoal and bonfires in 1A4bi Other sectors residential in 2015.

1A4bi Other sectors: residential stationary combustion	Unit	CO ₂ biog.	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
Use of charcoal	kg/TJ	112'000	200	1	50	600	11	6'000
Bonfires	kg/TJ	99'900	300	4	50	600	11	6'000

Activity data (1A4 stationary)

General energy sources

Activity data about the energy sources gas oil, residual fuel oil, natural gas and biomass are calculated by the Swiss energy model (see 3.2.4.3 for further information). For other energy sources such as other bituminous coal, activity data is provided directly by the Swiss overall energy statistics (SFOE 2016). Grass drying activities for source category 1A4ci are reported by the Swiss association of grass drying plants (VSTB) (as standard tonne of dried grass) as documented in the EMIS database (EMIS 2017/1A4ci Grastrocknung). Since submission 2015, the actual fuel consumption for grass drying is available and used for emission calculations.

Table 3-61 Activity data in 1A4a Commercial/Institutional (stationary).

Source/Fuel	Unit	1990	1995	2000	2005
1A4a Other sectors:	TJ	78'077	85'328	81'618	87'754
Commercial/institutional					
Gas oil	TJ	57'622	58'811	53'013	54'937
Gas oil heat only boilers	TJ	57'599	58'635	52'662	54'620
Gas oil engines	TJ	24	175	351	318
Natural gas	TJ	17'495	22'715	24'234	27'440
NG heat only boilers	TJ	17'219	21'544	22'498	25'407
NG turbines	TJ	85	78	NO	28
NG engines	TJ	192	1'093	1'737	2'004
Other bituminous coal	TJ	NO	NO	NO	NO
Lignite	TJ	NO	NO	NO	NO
Biomass (total)	TJ	2'959	3'802	4'371	5'377
Biomass (wood)	TJ	2'935	3'780	4'313	5'231
Biomass (biogas)	TJ	24	23	58	146

Source/Fuel	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A4a Other sectors:	TJ	82'577	74'772	79'010	77'017	83'372	69'084	75'946	80'829	64'454	70'715
Commercial/institutional											
Gas oil	TJ	51'554	45'450	47'585	45'699	48'778	38'900	41'814	44'328	34'191	36'406
Gas oil heat only boilers	TJ	51'260	45'269	47'416	45'545	48'660	38'796	41'720	44'242	34'109	36'324
Gas oil engines	TJ	293	181	169	154	119	105	94	86	82	82
Natural gas	TJ	25'034	23'511	24'987	24'496	27'207	23'805	26'783	28'213	21'946	24'837
NG heat only boilers	TJ	23'082	21'585	23'130	22'683	25'503	22'224	25'231	26'715	20'510	23'401
NG turbines	TJ	23	28	29	26	23	17	5	7	7	7
NG engines	TJ	1'929	1'898	1'829	1'787	1'681	1'564	1'548	1'490	1'429	1'429
Other bituminous coal	TJ	NO									
Lignite	TJ	NO									
Biomass (total)	TJ	5'990	5'810	6'437	6'822	7'387	6'379	7'348	8'288	8'317	9'472
Biomass (wood)	TJ	5'773	5'497	6'007	6'323	6'742	5'647	6'521	7'238	6'975	7'885
Biomass (biogas)	TJ	216	313	431	499	644	732	827	1'050	1'341	1'587

Table 3-62 Activity data in 1A4b Residential (stationary).

Source/Fuel	Unit	1990	1995	2000	2005
1A4b Other sectors: Residential	TJ	185'285	189'244	170'409	185'917
Gas oil	TJ	136'887	133'548	116'295	124'024
Gas oil heat only boilers	TJ	136'887	133'544	116'242	123'961
Gas oil engines	TJ	1	4	53	63
Natural gas	TJ	25'841	34'074	36'256	42'623
NG heat only boilers	TJ	25'781	33'815	35'817	42'093
NG turbines	TJ	NO	NO	NO	NO
NG engines	TJ	60	258	439	530
Other bituminous coal	TJ	630	460	130	400
Lignite	TJ	NO	NO	NO	NO
Biomass	TJ	21'926	21'162	17'728	18'869

Source/Fuel	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A4b Other sectors: Residential	TJ	178'757	159'078	170'197	166'819	180'901	145'330	160'343	171'385	134'415	144'108
Gas oil	TJ	118'885	102'729	108'715	105'296	111'731	86'989	94'103	99'373	75'136	79'406
Gas oil heat only boilers	TJ	118'823	102'663	108'663	105'254	111'695	86'955	94'072	99'344	75'109	79'379
Gas oil engines	TJ	63	65	52	42	36	34	32	29	27	27
Natural gas	TJ	40'914	39'151	42'381	42'462	48'222	40'903	47'036	50'946	42'367	46'107
NG heat only boilers	TJ	40'372	38'605	41'840	41'924	47'717	40'433	46'570	50'498	41'937	45'677
NG turbines	TJ	NO	3	3	NO						
NG engines	TJ	542	542	537	538	506	470	466	448	430	430
Other bituminous coal	TJ	400	400	400	400	400	400	400	400	400	400
Lignite	TJ	NO									
Biomass	TJ	18'557	16'798	18'701	18'661	20'547	17'037	18'804	20'665	16'513	18'195

Table 3-63 Activity data in 1A4ci Agriculture/forestry/fishing (stationary).

Source/Fuel	Unit	1990	1995	2000	2005
1A4c Agriculture/forestry/fishing	TJ	2'323	2'032	1'703	1'644
Grass drying (fossil, biogenic)	TJ	1'895	1'544	1'223	994
Biomass	TJ	428	488	480	649
1990=100%		100%	87%	73%	71%

Source/Fuel	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A4c Agriculture/forestry/fishing	TJ	1'437	1'507	1'445	1'486	1'435	1'461	1'398	1'040	1'151	1'152
Grass drying (fossil, biogenic)	TJ	845	948	822	856	739	891	685	458	524	431
Biomass	TJ	592	558	622	630	697	570	713	582	627	721
1990=100%		62%	65%	62%	64%	62%	63%	60%	45%	50%	50%

Charcoal and bonfires

Besides the main energy sources, also charcoal use and bonfires are accounted for in source category 1A4bi. The energy source charcoal is only used for charcoal grills. The total charcoal consumption under 1A4bi is very small compared to other fuels used for heating purposes. The activity data are the sum of charcoal production under 1A1c and net imports provided by the Swiss overall energy statistics (SFOE 2016).

The total wood demand for bonfires is assumed to be constant over time. Per capita wood demand is decreasing since 1990 due to an increasing number of inhabitants (for further details see documentation in EMIS 2017/1A4bi Lagerfeuer).

Table 3-64 Activity data in 1A4bi Charcoal and bonfires.

1A4bi Other sectors: residential stationary combustion	Unit	1990	1995	2000	2005
Use of charcoal	TJ	311	291	292	313
Bonfires	TJ	160	160	160	160

1A4bi Other sectors: residential stationary combustion	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Use of charcoal	TJ	303	313	354	343	343	343	344	343	354	353
Bonfires	TJ	160	160	160	160	160	160	160	160	160	160

3.2.7.3 Uncertainties and time-series consistency for 1A4 (stationary)

The uncertainty of CO₂ emissions from fuel combustions is described in the uncertainty analysis in chp. 3.2.4.7. Uncertainty in emissions of other non-CO₂ gases is estimated to be medium: 30% for CH₄ and 80% for N₂O (see Table 1-10).

Consistency: Time series for 1A4 Other sectors are all considered to be consistent.

3.2.7.4 Category-specific QA/QC and verification for 1A4 (stationary)

The general QA/QC procedures are described in chp. 1.2.3. Furthermore QA/QC procedures conducted for all 1A source categories are listed in chp. 3.2.4.8.

3.2.7.5 Category-specific recalculations for 1A4 (stationary)

- 1A4: Recalculations of the CO₂ emission factor of natural gas (see chp. 3.2.4.9) result in a change in the emission level of -13 kt CO₂ eq in 2014 and 2 kt CO₂ eq in 1990.
- 1A4ci: The missing emission factors in last year's submission of NO_x, CO, NMVOC and SO₂ for 1A4ci Plants for renewable waste from wood products in 2014 are now included in the inventory.
- 1A4ci: The emission factors of NMVOC as well as NO_x and CO from 1A4ci Drying of grass have been revised from 1990 and 1991, respectively, onwards based on air pollution control measurements (2005–2015).
- 1A4ai, 1A4bi and 1A4ci Wood combustion AD of automatic boilers and stoves have been revised for 1990–2014 and 2011–2014, respectively due to minor recalculations in Swiss wood energy statistics (SFOE 2016b).

3.2.7.6 Category-specific planned improvements for 1A4 (stationary)

No category-specific improvements are planned.

3.2.8 Source category 1A2 – Manufacturing industry and construction (mobile 1A2g vii)

3.2.8.1 Source category description for 1A2 Manufacturing industry and construction (mobile 1A2g vii)

Note for Key categories 1A2:

See chp. 3.2.6 and note that source category 1A2 contains the sum of emissions of stationary and mobile sources – the statement on key categories holds for the aggregated emission only. However, the CO₂ emissions of 1A2 from fuel consumption are dominated by the stationary sources, 87%, which means that the emissions of 1A2g vii only play a minor role within category 1A2].

Table 3-65 Specification of source category 1A2 Manufacturing industries and construction (mobile).

Source	Specification
Mobile Combustion in manufacturing industries and construction	industry sector: forklifts and snow groomers etc. construction machines: excavators, loaders, dump trucks, mobile compressors etc.

3.2.8.2 Methodological issues for 1A2 Manufacturing industry and construction (mobile 1A2g vii)

Methodology (1A2g vii)

Based on the decision tree Fig. 3.3.1 in chp. “3. Mobile Combustion” in IPCC (2006) the emissions of industry and construction vehicles and machinery are calculated by a Tier 3 method with the non-road transportation model described in chp. 3.2.4.5.1.

Emission factors (1A2g vii)

- The CO₂ emission factors applied for the time series 1990–2015 for diesel oil, gasoline and compressed natural gas are country-specific and are given in Table 3-12.
- The CH₄ and N₂O emission factors are country-specific and are shown in Table 3-66 to Table 3-68 for diesel oil, gasoline and CNG engines for all emission standards.
- For SO₂ the emission factors are country-specific. See also Table A – 19 in Annex 3.1.5 for diesel oil, gasoline, gas oil.
- The emission factors for precursors are country-specific and are given in FOEN (2015j).
- NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference between VOC and CH₄ emissions.
- The implied emission factors 2015 are shown in Table 3-69.

All emission factors (GHG, precursors, SO₂) can be downloaded by query from the public part of the non-road database INFRAS (2015a)⁸. They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels.

Table 3-66 Emission factors CH₄ and N₂O for industry and construction vehicles with diesel engines by emission standards including the year of enforcement.

Gas	Power class kW	PreEU-A	PreEU-B	EU-I	EU-II	EU-III A	EU-III B	EU-IV
		<1996	1996	2002/03	2002/04	2006/08	2011/12	2014
		g/kWh						
CH ₄	<18	0,0547	0,0547	0,0384	0,0240	0,0142	0,0142	0,0142
CH ₄	18–37	0,0578	0,0578	0,0221	0,0134	0,0089	0,0089	0,0089
CH ₄	37–56	0,0319	0,0319	0,0156	0,0110	0,0079	0,0055	0,0058
CH ₄	56–75	0,0319	0,0319	0,0156	0,0110	0,0079	0,0031	0,0031
CH ₄	75–130	0,0218	0,0218	0,0108	0,0084	0,0067	0,0031	0,0031
CH ₄	130–560	0,0218	0,0218	0,0103	0,0072	0,0053	0,0031	0,0031
CH ₄	>560	0,0218	0,0218	0,0103	0,0072	0,0053	0,0031	0,0031
N ₂ O	0–3000	0,035	0,035	0,035	0,035	0,035	0,035	0,035

⁸ <https://www.bafu.admin.ch/bafu/en/home/topics/air/state/non-road-datenbank.html>

Table 3-67 Emission factors CH₄ and N₂O for industry and construction vehicles with gasoline engines by emission standards including the year of enforcement.

Gas	Power class ccm	PreEU-A	PreEU-B	PreEU-C	EU-I	EU-II
		<1996	1996	2000	2004	2005/09
		g/kWh				
CH ₄	<66	2,04	2,04	2,04	1,394	1,394
CH ₄	66–100	1,36	1,36	1,36	1,088	1,088
CH ₄	100–225	0,68	0,68	0,68	0,408	0,408
CH ₄	>225	0,68	0,68	0,68	0,34	0,306
N ₂ O	0–3000	0,03	0,03	0,03	0,03	0,03

Table 3-68 Emission factors CH₄ and N₂O for industry and construction vehicles with CNG engines.

Gas	without catalyst	with catalyst
	g/kWh	
CH ₄	0,552	0,035
N ₂ O	0,05	0,05

Table 3-69 Implied emission factors 2015 for industry and construction vehicles.

1A2gvii Non-road vehicles and other machinery	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
	t/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ	kg/TJ
Gasoline	73.8	41.3	1.1	107	769	1.8	19'589
Diesel oil	73.3	0.7	3.3	372	31	0.5	156
Natural gas	56.4	NA	NA	NA	NA	NA	NA

Activity data (1A2g vii)

Activity data for non-road (1A2gvii) are described in chp. 3.2.4.5.1 (non-road transportation model). Values are taken from FOEN (2015j). Data on biofuels are provided by the statistics of renewable energies (SFOE 2016a). Activity data are shown in Table 3-70 and in Annex A3.1.4. Detailed data can be downloaded from the online database of INFRAS (2015a).

Table 3-70 Activity data for industry and construction vehicles.

Source/Fuel	Unit	1990	1995	2000	2005
1A2gvii Non-road vehicles and other machinery	TJ	5'722	6'861	7'644	8'170
Gasoline	TJ	196	224	227	225
Diesel oil	TJ	5'359	6'380	7'108	7'629
Natural gas	TJ	167	257	301	292
Biodiesel	TJ	NO	NO	8	24
Bioethanol	TJ	NO	NO	NO	0.02

Source/Fuel	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A2gvii Non-road vehicles and other machinery	TJ	8'292	8'414	8'536	8'657	8'779	8'810	8'841	8'873	8'904	8'935
Gasoline	TJ	224	223	222	221	220	212	205	197	189	182
Diesel oil	TJ	7'755	7'881	8'007	8'133	8'259	8'295	8'331	8'367	8'403	8'440
Natural gas	TJ	288	283	279	274	270	261	252	243	234	225
Biodiesel	TJ	25	27	28	29	31	42	53	63	74	85
Bioethanol	TJ	0.03	0.04	0.05	0.07	0.08	0.72	1.36	2.00	2.64	3.29

Underlying activity data (vehicle stock, operating hours) of mobile non-road sources can also be downloaded by query from the public part of the non-road database INFRAS (2015a), see footnote 8. They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels.

3.2.8.3 Uncertainties and time-series consistency for 1A2g vii (mobile)

Uncertainties by fuel type are given in Table 3-25.

3.2.8.4 Category-specific QA/QC and verification for 1A2g vii (mobile)

The general QA/QC procedures are described in chp. 1.2.3. Furthermore, QA/QC procedures conducted for all 1A source categories are listed in 3.2.4.8.

3.2.8.5 Category-specific recalculations for 1A2g vii (mobile)

No recalculations were carried out for source category 1A2g vii.

3.2.8.6 Category-specific planned improvements for 1A2g vii (mobile)

No category-specific improvements are planned.

3.2.9 Source category 1A3 - Transport

3.2.9.1 Source category description for 1A3

Table 3-71 Key categories (KCA incl. LULUCF) of 1A3 Transport.

Code	IPCC Category	GHG	Identification Criteria
1A3a	Civil Aviation: Liquid Fuels	CO ₂	T1
1A3b	Road Transportation: Gasoline	CH ₄	T1, T2
1A3b	Road Transportation: Gasoline	CO ₂	L1, T1
1A3b	Road Transportation: Gasoline	N ₂ O	T1, T2
1A3b	Road Transportation: Diesel	CO ₂	L1, L2, T1, T2
1A3b	Road Transportation: Diesel	N ₂ O	T1

Table 3-72 Specification of source category 1A3 Transport.

1A3	Source	Specification
1A3a	Domestic aviation	Large (jet, turboprop) and small (piston) aircrafts, helicopters
1A3b i	Road Transportation	Passenger cars
1A3b ii		Light duty trucks
1A3b iii		Heavy duty trucks and buses
1A3b iv		Motorcycles
1A3b v		Other
1A3c	Railways	Diesel locomotives
1A3d	Domestic navigation	Passenger ships, motor and sailing boats on the Swiss lakes and the river Rhine
1A3e	Other transportation - Pipeline compressors	Compressor station in Ruswil, Lucerne

For information on international bunker fuel emissions from international aviation and navigation, see chp. 3.2.2.

3.2.9.2 Methodological issues for 1A3

3.2.9.2.1 Domestic aviation (1A3a)

Methodology (1A3a)

The emissions of domestic aviation are modelled by a Tier 3A method (IPCC 2006, Volume 2, chp. 3 Mobile Combustion, Table 3.6.2 and figure 3.6.2) developed by FOCA (2006) and based on origin and destination of single movements by aircraft type according to detailed movement statistics. LTO emissions are modelled based on the individual engine type. The emissions of domestic aviation are modelled together with the international aviation reported in 1D1 (aviation bunker, see chp. 3.2.2.2.1).

FOCA is represented in the emissions technical working group (CAEP WG3) and in the modelling and database group (CAEP MDG) of the International Civil Aviation Organisation (ICAO). FOCA is directly involved in the development of ICAO guidance material for the calculation of aircraft emissions and in the update of the IPCC Guidelines (via the secretariat of ICAO CAEP (Committee on Aviation Environmental Protection)). The Tier 3A method applied for the emission modelling is in line with the methods developed in the working groups mentioned. The modelling scheme for domestic aviation refers to aircraft basic data, activity data and emission factors that result in calculated emissions. Respective values are ultimately imported into the EMIS database as shown in Figure 3-23.

The Tier 3A method follows standard modelling procedures on the level of single movements based on detailed movement statistics. The primary key for all calculations is the aircraft tail number, which allows to calculate on the most precise level, namely on the level of the individual aircraft and engine type. Every aircraft is linked to the FOCA engine data base containing emission factors for more than 700 individual engine types with different power settings. Emissions in the landing and take-off cycle (LTO) are calculated with aircraft category dependent flight times and corresponding power settings. Cruise emissions are calculated based on the individual aircraft type and the trip distance for every flight. For piston-engine powered aircraft and helicopters, to the knowledge of FOCA, it has been the only provider of publicly available engine data and a full methodology. All piston engine data

and study results have been published in 2007 (FOCA 2007a). The guidance on the determination of helicopter emissions has been published in 2009 (FOCA 2009a) and updated in 2015 (FOCA 2015a).

The movement database from Swiss airports registers the departure and destination airports of each flight. With this information, all flights from and to Swiss airports are differentiated into domestic and international flights prior to the emission calculation. The emissions of domestic flights are reported under 1A3a Domestic Aviation, the emissions of international flights are reported under 1D1 international aviation (international bunkers).

The emission factors used are either country-specific or taken from the ICAO engine emissions databank, from EMEP/CORINAIR databases (EMEP/EEA 2013), Swedish Defence Research Agency (FOI) and Swiss FOCA measurements (precursors). Cruise emission factors are generally calculated from the values of the ICAO engine emissions databank, aircraft performance tables and from confidential airline data. Pollutant emission factors are adjusted to cruise conditions by using the Boeing Fuel Flow Method 2. For N_2O , the IPCC default emission factor of 2 kg/TJ is used. For the methane split of unburned hydrocarbons, the 10% methane share for the LTO, given in IPCC 2006 is used. For cruise emissions, no methane is reported. Activity data are derived from a detailed movement statistics.




Figure 3-23 Modelling scheme (activity data, emission factors, emissions) for domestic aviation.

A complete emission modelling (LTO and cruise emissions for domestic and international flights) has been carried out by FOCA for 1990, 1995, 2000, 2002, 2004–2015. The results of

the emission modelling have been transmitted from FOCA to FOEN in an aggregated form. FOEN calculated the implied emission factors 1990, 1995, 2000, 2002, 2004 and carried out a linear interpolation for the years in-between. The interpolated implied emission factors were multiplied with the annual fuel sold from Swiss overall energy statistics (SFOE in respective years), providing the missing emissions of domestic aviation for the years 1991–1994, 1996–1999, 2001 and 2003.

Details of emission factors and activity data follow below. Further tables containing more information are also given in Annex A3.1.2, more detailed descriptions of the emission modelling may be found in FOCA (2006).

Emission factors (1A3a)

LTO

The FOCA engine emissions database consists of more than 700 individual engine data sets. Jet engine factors for engines above 26.7 kN thrust (emission certificated) are identical to the ICAO engine emissions database. Emission factors for lower thrust engines, piston engines and helicopters were taken from manufacturers or from own measurements. Emission factors for turboprops could be obtained in collaboration with the Swedish Defence Research Agency (FOI).

Cruise

The fuel flows of the whole Airbus fleet (which produces a great portion of the Swiss inventory) have been modelled on the basis of real operational aircraft data from flight data recorders (FDR) of Swiss International Airlines. GHG emission factors have been modelled on the basis of the ICAO engine databank and corrected to cruise conditions using FDR engine parameters and the Boeing Fuel Flow Method 2. For older aircraft types (pre 2003), part of the cruise emission factors were taken from EMEP/CORINAIR (EMEP/EEA 2013) and from former CROSSAIR (FOCA 1991). For new aircraft type entries, the FOCA models the cruise emission factors based on the aircraft type characteristics and the engine models fitted to the aircraft. The model uses proprietary aircraft information as well as public information from the ICAO engine database. For those aircraft types, which dominate the fuel consumption in Switzerland, flight data recorder information has been used to calibrate emission factors. The factors are updated periodically to take account of flight operational improvements, as well. Calculation results for international aviation emissions are periodically compared to Eurocontrol results. For piston engine aircraft and helicopters, Swiss FOCA has produced its own data, which were taken under real flight conditions (2005 data, FOCA 2009a, FOCA 2015a).

In 2016, the FOCA Helicopter Emissions Calculation Guidance has been updated and implemented into the emissions calculation for the 2015 emission inventory (FOCA 2015a). FOCA now uses engine power specific emission factors for most helicopters, taking into account lower power requirement per engine, if engines are installed in a twin engine configuration. On top of the few non-public manufacturer data sources, FOCA introduced 80 individual engine models replacing most of the generic engine assignments.

Kyoto gases

- CO₂: the emission factor of 72.8 t/TJ in 2014 is country-specific and is based on measurements and analyses of fuel samples (see Table 3-12 and Table 3-73)
- CH₄, NMVOC (country-specific; CORINAIR): VOC emissions (see Precursors below) are split into CH₄ and NMVOC by a constant share of 0.1 (CH₄) and 0.9 (NMVOC) for LTO. For cruise flights the VOC emissions do not consist of CH₄ emissions. The implied emission factor for CH₄ is shown in Table 3-73.
- The N₂O emission factor regarding jet kerosene is default given by the 2006 IPCC Guidelines (IPCC 2006). It is assumed that the emission factor for international cruise is sufficient for all kind of flight periods (LTO and cruise) and remains stable over the entire time period 1990–2014 (see Table 3-73).

Precursors

- Assignment of emission factors for 1990 and 1995: The fleet that operated in and from Switzerland during those years has been analysed. The corresponding most frequent engines within an aircraft category (ICAO Code) have been assigned to every aircraft type.
- Assignment of emission factors for the year 2000, 2002 and 2004 to 2015: the actual engine of every single aircraft operating in and from Switzerland has been assigned. FOCA uses the aircraft tail number as the key variable which links activity data and individual aircraft engine information (see Annex A 3.1.2 Table A – 12 Aircraft Engine Combinations).

FOCA determines the emission factors of different gases as follows given in Table 3-73.

Table 3-73 Implied emission factors of 1A3a in 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A3a Aviation	Unit	CO ₂ fossil	CH ₄	N ₂ O	NO _x	NMVOC	SO ₂	CO
Kerosene, domestic, LTO	kg/TJ	72'800	13.5	2.0	193	121.5	20.7	2'890
Kerosene, domestic, CR	kg/TJ	72'800	NA	2.0	282	48.1	22.0	684
Kerosene, international, LTO	kg/TJ	72'800	3.5	2.0	292	31.2	23.2	304
Kerosene, international, CR	kg/TJ	72'800	NA	2.0	310	9.0	23.2	44

Activity data (1A3a)

The statistical basis has been extended after 1996. Therefore, the modelling details are not exactly the same for the years 1990/1995 as for the subsequent years. The source for the 1990 and 1995 modelling are the movement statistics, which records information for every movement on airline, number of seats, Swiss airport, arrival/departure, origin/destination, number of passengers, distance. From 1996 onwards, every movement in the FOCA statistics also contains the individual aircraft tail number (aircraft registration). This is the key variable to connect airport data and aircraft data. The statistics may contain more than one million records with individual tail numbers. All annual aircraft movements recorded are split

into domestic and international flights (there are 460'978 aircraft movements in the total of scheduled and charter traffic in 2015 as provided by FOCA 2016).

Non-scheduled, non-charter and general aviation (including Helicopters)

- Airports and most of the airfields report individual aircraft data (aircraft registration). FOCA may therefore compute the inventory for small aircraft with Tier 3A method, too. However, for 1990 and 1995, the emissions data for non-scheduled, non-charter and General Aviation (helicopters etc.) could not be calculated with a Tier 3A method. Its fuel consumption is estimated to be 10% of the domestic fuel consumption. Data were taken from two FOCA studies (FOCA 1991, FOCA 1991a). For 2000–2007, all movements from airfields are known, which allows a more detailed modelling of the emissions (FOCA 2007a).
- Helicopter flights which do not take off from an official airport or airfield such as transport flights, flights for lumbering, animal transports, supply of alpine huts, heli-skiing and flight trainings in alpine regions cannot be recorded with the movement data base from airports and airfields. These emissions are taken into account using the statistics of the Swiss Helicopter Association (Unternehmensstatistik der Schweizer Helikopterunternehmen). These statistics are officially collected by FOCA and updated annually (see FOCA 2004 as illustrative example for all subsequent years). In this case, emissions are calculated based on operating hours of the helicopters, with emission factors taken from the helicopter study (see FOCA 2015a).
- Since 2007, the data of these helicopter statistics are included electronically in the data warehouse of the model and undergo first some plausibility checks (E-plaus software). In order to distinguish between single engine helicopters and twin engine helicopters a fix split of 87% for single engine helicopters and 13% for twin engine helicopters has been applied for the entire commitment period until 2014 based on investigations in 2004 (FOCA 2004). From 2015, the statistics allowed to assign the individual helicopters from the helicopter companies. All emissions from helicopter flights without using an official airport or an official airfield are considered domestic emissions.

Fuel consumption: Table 3-74 summarises the activity data for domestic aviation (1A3a). It also includes international aviation, which belongs to the memo items, international bunkers/aviation (see also chp. 3.2.2). In order to split the fuel consumption for domestic and international flights, the FOCA calculates the fuel for each domestic and international flight bottom up. A first validation of this calculation can be done top down for the sum of all flights: The total annual aviation fuel sold known from robust energy statistics in a country should correspond very closely (within a few percent) to the modelled total fuel consumption of domestic and international flights together. For 2015, the modelled total fuel consumption in Switzerland was 2.0% higher than the fuel sold value, so the model showed a slight overestimation. The total fuel sold as reported in the Swiss overall energy statistics is considered the most robust value for reporting, so the modelled total fuel consumption is scaled downwards such that the sum of domestic and international fuel consumption becomes identical with the fuel sold. The scaling is only done on the international fuel for the following reasons: Eurocontrol calculations for Switzerland's international flights fuel consumption are usually a few percent lower than the FOCA result. For domestic flights, the FOCA takes every movement including the smallest aircraft into account and applies conservative emission factors. An indication of this is the fact that Eurocontrol calculations for

Switzerland's domestic flight fuel consumption is usually only around half the value reported by Switzerland. In summary, Switzerland reports the domestic fuel consumption according to the modelled value (conservative estimation), whereas the international fuel consumption (bunker) is scaled downwards so that the sum of domestic and international fuel consumption becomes identical with the fuel sold, as reported in the Swiss overall energy statistics.

Table 3-74 Fuel consumption of civil aviation in TJ for separated for domestic / international and LTO / cruise. Domestic consumption and the corresponding emissions are reported under 1A3a, international consumption is reported under Memo items, international bunkers (FOCA 2007, 2007a, 2008–2015).

1A3a/1D1 Civil aviation	1990	1995	2000	2005
	Fuel consumption in TJ			
Kerosene, domestic, LTO	1'050	935	772	517
Kerosene, domestic, CR	2'401	2'139	1'767	1'182
Kerosene, international, LTO (not part of national total)	4'277	5'097	6'503	4'868
Kerosene, international, CR (not part of national total)	37'608	44'821	57'184	42'804
Total Civil aviation	45'334	52'993	66'225	49'370
1990 = 100%	100%	117%	146%	109%

1A3a/1D1 Civil aviation	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	Fuel consumption in TJ									
Kerosene, domestic, LTO	506	544	512	497	463	507	502	494	525	387
Kerosene, domestic, CR	1'152	1'347	1'106	1'207	1'225	1'301	1'365	1'323	1'396	1'500
Kerosene, international, LTO (not part of national total)	5'095	5'401	5'737	5'449	5'622	6'017	6'199	6'208	6'142	6'459
Kerosene, international, CR (not part of national total)	45'013	48'142	52'107	49'789	52'496	56'194	57'428	58'501	58'864	60'874
Total Civil aviation	51'766	55'434	59'462	56'942	59'805	64'019	65'494	66'526	66'927	69'220
1990 = 100%	114%	122%	131%	126%	132%	141%	144%	147%	148%	153%

3.2.9.2.2 Road transportation (1A3b)

Methodology (1A3b)

Choice of method

- The CO₂ emissions are calculated by a Tier 2 method based on the decision tree Fig. 3.2.2 in chp. 3. Mobile Combustion in IPCC (2006).
- The CH₄ and the N₂O emissions are calculated by a Tier 3 method based on the decision tree Fig. 3.2.3 in chp. 3. Mobile Combustion in IPCC (2006).
- The use of urea in urea-based catalysts is reported in chp. 4.5.2.2 under 2D3d as recommended in the reporting table's footnotes.

Connections between road model, non-road model and Swiss overall energy statistics

For the source categories related to transport, INFRAS developed a territorial emission model for road transportation (1A3b, INFRAS 2010 / FOEN 2010i; for details refer to Annex A3.1.3) and a model for non-road transportation (mobile sources in 1A2g vii, 1A3c, 1A3d, 1A4a ii, 1A4b ii, 1A4c ii, 1A5b (excl. military aviation); (FOEN 2015j, see also "non-road transportation model" in chp. 3.2.4.5.1).

Due to fuel price differences in the vicinity of the national borders, gasoline stations used to sell relevant amounts of fuels to foreign car owners. This amount of fuel was mainly

consumed abroad, called **fuel tourism**, was not captured by the territorial road model, but had to be included in the GHG inventory for the UNFCCC reporting. No fuel tourism is assumed for the non-road model.

The Swiss overall energy statistics provide information on the amounts of fuel sold, which contains territorial consumption **and** fuel tourism. From the amounts of fuel sold, the consumptions modelled by the territorial road and non-road models – i.e. fuel used – are subtracted. The resulting differences to the amount of fuels sold represent the amount of fuel tourism plus statistical differences⁹. Figure 3-24 shows how the models and the Swiss overall energy statistics are linked to determine the GHG emissions from road and non-road transportation:

- CO₂ emissions are calculated by using fuel sales and country-specific CO₂ emission factors.
- CH₄ and N₂O emissions are calculated in three steps:
- From fuel used and country-specific CH₄ and N₂O emission factors, the territorial emissions are calculated.
- The differences between fuels sold and fuels used (territorial) are interpreted as fuel tourism plus and statistical differences. These amounts of gasoline and diesel oil are multiplied with implied CH₄ and N₂O emission factors, which are deduced from the road transportation, to form the CH₄ and N₂O emissions resulting from fuel tourism plus statistical difference.
- CH₄ and N₂O emissions from the territorial model plus CH₄ and N₂O from fuel tourism and statistical difference are added to the total CH₄ and N₂O emissions reported to the UNFCCC.

⁹ The amount of fuel tourism has been estimated by SFOE (2010d). The result showed that the difference between fuels sales and fuels determined by the traffic model tend to overestimate the “true” fuel tourism. It was concluded that the difference also contains potential underestimation of the mileage and other statistical errors. Therefore, the difference between fuel sales and fuel used in the traffic model is therefore indicated in the NIR as “fuel tourism and statistical difference”.




Figure 3-24 Connections between fuel sold and fuel used for road and non-road transportation. Fuel sold is provided by the Swiss overall energy statistics (minus Liechtenstein's gasoline and diesel oil consumption and bunker fuels for navigation). Fuel used results from the territorial road and non-road models. The residual fuel consists of fuel tourism and a statistical difference. Its emissions are calculated by means of implied emission factors deduced from the territorial road model. The diagram holds separately for gasoline and diesel oil.

Methodology of the territorial road transportation model

The emission computation is based on two sets of data (FOEN 2010i):

- Emission factors: specific emissions in grams per activity data unit.
- Traffic activity data: vehicle kilometres travelled (hot emissions), number of starts/stops and vehicle stock (cold start, evaporation emissions and running losses) or fuel consumption per vehicle category.

Emission modelling consists of three parts:

hot emissions: $E_{hot} = VKT \cdot EF_{hot}$

start emissions: $E_{start} = N_{start} \cdot EF_{start}$

evaporative emissions: $E_{evap,i} = N_{evap,i} \cdot EF_{evap,i}$

with

- EF_{hot} , EF_{start} , EF_{evap} : Emission factors for ordinary driving conditions (hot motor), cold start and evaporative (VOC) emissions (after stops, running losses, diurnal losses)
- VKT : Vehicle km travelled
- N_{start} : Number of starts
- i runs over three evaporation categories: stops, running losses, diurnal losses

- $N_{\text{evap},i}$: Number of stops (i = "after stops") or number of vehicles (i = "running losses" and "diurnal losses")

Note that cold start emissions for CH₄ and N₂O are not accounted for in the model described. During the in-country review in 2016, the ERT identified a potential underestimation. Switzerland therefore estimated N₂O cold start excess emissions for PC and LDV for 1990, 2014 and 2015 by means of emission factors of the Copert model, as recommended by the ERT. For the years 1991–2013 the emissions were interpolated linearly between 1990 and 2014. The emission factors per Euro class are documented in the EMEP/EEA air pollutant emission inventory guidebook - 2013 on p. 91 ff. (EMEP/EEA 2013). Note that the ERT confirmed that this approach complies with the 2006 IPCC Guidelines.

Emission factors (1A3b)

CO₂

- The country-specific CO₂ emission factors are described in chp.3.2.4.4.2. Values are shown in Table 3-12 (gasoline, diesel oil) and in Table 3-13 (natural gas). The values in 2014 are also shown in Table 3-75.
- The same emission factors are also applied for the calculation of the emissions resulting from fuel tourism.

CH₄

- Country-specific emission factors are applied. Details see below ("*Country-specific emission factors*").
- CH₄ emissions from fuel tourism: From the territorial model, implied emission factors for CH₄ are derived per vehicle category and per fuel type corresponding to mean emission factors for Switzerland (see Figure 3-24). These factors are then applied to calculate the emissions resulting from fuel tourism. To verify this approach, a comparison of these emission factors with implied emission factors of the neighbouring countries have been carried out. The differences are small between Switzerland, Austria, and Germany because all three countries use the same emission factors (INFRAS 2010, TUG 2009), whereas there are some differences compared to France and Italy that use other emission factors (COPERT). Nevertheless, the use of the mean Swiss emission factors seems to be the consistent approach.
- For biofuels, no country-specific EFs for CH₄ are available. Therefore, emissions have been estimated using the EFs for alternative fuel vehicles provided in table 3.2.4 on page 3.23 of Volume 2 of the 2006 IPCC Guidelines (IPCC 2006).

N₂O

- Country-specific emission factors are applied. Details see below ("*Country-specific emission factors*").
- N₂O emissions from fuel tourism: The same approach as for CH₄ is applied (see paragraph above)
- For biofuels no country-specific EFs for N₂O are available. Therefore, emissions have been estimated using the EFs for alternative fuel vehicles provided in table 3.2.4 on page

3.23 of Volume 2 of the 2006 IPCC Guidelines (IPCC 2006). The value of 101 mg/km from the 2006 IPCC Guidelines was used for urban buses running on CNG only. For the bi-fuel passenger cars, it is assumed that they use gasoline mainly during the start but otherwise run on CNG; therefore the respective CNG emission factor for light duty vehicles of 27 mg/km from the same source was applied.

- Cold start emission factors 2015 are for PC (gasoline) 0.011 kg/TJ, and for LDV 0.025 kg/TJ.

Country-specific emission factors

Emission factors for other gases are country-specific derived from “emission functions” which are determined from a compilation of measurements from various European countries with programs using similar driving cycles (legislative as well as standardized real-world cycles, like “Common Artemis Driving Cycle” (CADC)). The method has been developed in 1990–1995 and has been extended and updated in 2000, 2004 and 2010. These emission factors are compiled in a “Handbook of Emission Factors for Road Transport” (seer INFRAS 2010, TUG 2009). Version 3.1 is presented and documented on the website <http://www.hbefa.net/>. A later version has recently been made available and will be used in future submissions. The general emission factor methodology is documented in TUG (2009). The resulting emission factors are published on CD ROM (“Handbook of emission factors for Road Transport”, INFRAS 2010). They refer to the so-called “traffic situations”, which represents characteristic patterns of driving behaviour and which serve as a key to the disaggregation of the activity data. The underlying database contains a dynamic fleet compositions model simulating the release of new exhaust technologies and the fading out of old technologies. Corrective factors are provided to account for future technologies. Further details are shown in Annex A3.1.3.

Implied emission factors for GHG, precursors and SO₂

The following Table 3-75 presents mean emission factors for the GHG, the precursors and SO₂ in 2015. More or less pronounced decreases of the emission factors have occurred in the last years due to new emission regulations and subsequent new exhaust technologies (mandatory use of catalytic converters for gasoline cars and lower limits for sulphur content in diesel fuels). Early models of catalytic converters have been substantial sources of N₂O, leading to an emission increase until 1998. Recent converter technologies have overcome this problem resulting in a decrease of the (mean) emission factor.

Note that an inconsistency in the attribution of natural gas to the vehicle categories leads to an error in the implied emission factors for gas-driven light duty vehicles. The error will be corrected for the next submission.

Table 3-75 Implied emission factors in 2015 for road transportation. For more details see Annex A3.1.3.

1A3b Road Transportation Gasoline / Bioethanol	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
	kg/TJ						
Passenger cars	73'800	5.0	0.8	54	82	0.36	642
Light duty vehicles	73'800	10	2.8	160	163	0.38	2211
Heavy duty vehicles	NO	NO	NO	NO	NO	NO	NO
Motorcycles	73'800	64	1.4	135	487	0.38	4344
Fuel tourism and statistical differences	73'800	6.9	0.5	59	104	0.36	786

1A3b Road Transportation Diesel / Biodiesel	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
	kg/TJ						
Passenger cars	73'300	0.29	2.5	224	12	0.45	58
Light duty vehicles	73'300	0.27	2.0	328	11	0.48	64
Heavy duty vehicles	73'300	0.18	3.4	329	7	0.47	117
Motorcycles	NO	NO	NO	NO	NO	NO	NO
Fuel tourism and statistical difference	73'300	0.25	2.5	275	10	0.46	80

1A3b Road Transportation Gas / Biogas	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
	kg/TJ						
Passenger cars	56'400	6.9	2.7	29	0.6	0.02	204
Light duty vehicles	56'400	0.10	0.004	0.02	0.01	0.000008	0.7
Heavy duty vehicles	56'400	2.4	7.0	154	0.2	0.01	108
Motorcycles	NO	NO	NO	NO	NO	NO	NO
Fuel tourism and statistical difference	56'400	2.8	4.0	175	0.2	0.01	123.2

Activity data (1A3b)

Energy-related activity data (basis for modelling the CO₂ emissions)

The amount of gasoline and diesel fuel sold in Switzerland serves as the activity data for the calculation of the CO₂ emissions. The Swiss overall energy statistics provides the amount of gasoline and diesel oil sold in Switzerland and the Principality of Liechtenstein (SFOE 2016). From these numbers, Liechtenstein's sales, the non-road consumption, the bunker fuels and the fugitive emissions from transmission, storage and fuelling of gasoline (reported under 1B2av Distribution of oil products) are subtracted. The result gives the inventory-relevant consumption for estimating the CO₂ emissions. It contains the fuel consumption of the traffic model plus the amount of fuel tourism and statistical differences.

The consumption of biofuels is based on the Swiss overall energy statistics (SFOE 2016), the Swiss renewable energy statistics (SFOE 2016a) and the Federal Customs Administrations (FCA 2016).

Table 3-76 shows the split of fuel sales into territorial road transportation model, the territorial non-road transportation model and fuel tourism (incl. statistical difference).

- The relevant numbers for road transportation are given as two different contributions in the rows "on road fuel consumption (model)" and "fuel tourism and statistical difference".
- The emissions from natural gas combustion for road transportation originate from activity data of two vehicle categories: biofuel CNG/petrol passenger cars and urban buses running purely on CNG. Data source is FCA (2016).

- Consumption of biofuels for road transportation (biodiesel, bioethanol and biogas) starts in Switzerland in 1997.

Table 3-76 Split of fuel sales between territorial “on-road consumption (model)”, “non-road consumption (models)” and “fuel tourism and statistical differences” (residual value to sales amounts) for gasoline, diesel oil, natural gas (CNG) and biofuels (Vegetable/Waste oil is included in the numbers of Biodiesel) in PJ. Numbers may not add to totals due to rounding.

Activity data for fossil fuels on-road and non-road categories	Source category	1990	1995	2000	2005
		PJ			
Gasoline					
on-road consumption (model)	1A3b	135.6	140.8	146.9	136.1
fuel tourism and statistical difference	1A3b	17.8	8.1	19.0	13.9
non-road consumption (models)	1A2gvii; 1A3dii; 1A4aii,bii,cii; 1A5b	2.4	2.4	2.3	2.1
Gasoline sold in Switzerland		155.8	151.3	168.2	152.1
Diesel oil					
on-road consumption (model)	1A3b	36.5	39.8	44.9	57.4
fuel tourism and statistical difference	1A3b	-0.7	-4.4	-3.2	1.2
non-road consumption (models)	1A2gvii; 1A3c,dii; 1A4cii; 1A5b	11.0	12.4	13.5	13.9
Diesel oil sold in Switzerland		46.7	47.8	55.2	72.6
Natural gas					
on-road consumption (model)	1A3b	NO	NO	NO	0.1
fuel tourism and statistical difference	1A3b	NO	NO	NO	0.1
non-road consumption (models)	1A2gvii	0.2	0.3	0.3	0.3
Natural gas sold in on- and non-road categories		0.2	0.3	0.3	0.5
Biodiesel					
on-road consumption (model)	1A3b	NO	NO	0.1	0.2
non-road consumption (models)	1A2gvii; 1A3c,dii; 1A4cii; 1A5b	NO	NO	0.01	0.04
Biodiesel sold in Switzerland		NO	NO	0.1	0.3
Bioethanol					
on-road consumption (model)	1A3b	NO	NO	NO	0.02
non-road consumption (models)	1A2gvii; 1A3dii; 1A4bii; cii; 1A5b	NO	NO	NO	0.0002
Bioethanol sold in Switzerland		NO	NO	NO	0.02
Biogas					
on-road consumption (model)	1A3b	NO	NO	NO	NO
non-road consumption (models)		NO	NO	NO	NO
Biogas sold in Switzerland		NO	NO	NO	NO

Activity data on-road and non-road categories	Source category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
		PJ									
Gasoline											
on-road consumption (model)	1A3b	132.1	129.1	126.4	122.8	119.3	116.0	112.7	109.4	106.1	102.6
fuel tourism and statistical difference	1A3b	13.2	14.9	14.4	14.2	12.9	11.0	9.8	7.5	6.1	1.3
non-road consumption (models)	1A2gvii; 1A3dii; 1A4aii,bii,cii; 1A5b	2.1	2.0	2.0	1.9	1.9	1.9	1.8	1.8	1.7	1.7
Gasoline sold in Switzerland		147.4	146.0	142.8	139.0	134.0	128.9	124.3	118.6	113.9	105.6
Diesel oil											
on-road consumption (model)	1A3b	61.1	65.3	68.2	70.8	73.9	75.1	76.3	77.5	78.8	78.7
fuel tourism and statistical difference	1A3b	3.4	4.9	10.1	8.7	9.1	10.6	15.5	19.1	20.8	19.3
non-road consumption (models)	1A2gvii; 1A3c,dii; 1A4cii; 1A5b	14.1	14.3	14.4	14.6	14.8	14.8	14.8	14.8	14.8	14.8
Diesel oil sold in Switzerland		78.6	84.4	92.7	94.1	97.8	100.5	106.6	111.5	114.4	112.8
Natural gas											
on-road consumption (model)	1A3b	NO	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.3	0.3
fuel tourism and statistical difference	1A3b	0.1	0.2	0.3	0.5	0.5	0.5	0.4	0.4	0.3	0.3
non-road consumption (models)	1A2gvii	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Natural gas sold in on- and non-road categories		0.4	0.6	0.7	0.9	1.0	1.0	0.9	0.9	0.9	0.9
Biodiesel											
on-road consumption (model)	1A3b	0.3	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.7	1.5
non-road consumption (models)	1A2gvii; 1A3c,dii; 1A4cii; 1A5b	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Biodiesel sold in Switzerland		0.4	0.4	0.5	0.4	0.4	0.4	0.5	0.5	0.8	1.6
Bioethanol											
on-road consumption (model)	1A3b	0.02	0.1	0.1	0.03	0.1	0.1	0.1	0.1	0.2	0.6
non-road consumption (models)	1A2gvii; 1A3dii; 1A4bii; cii; 1A5b	0.0003	0.0003	0.0004	0.001	0.001	0.01	0.01	0.02	0.02	0.03
Bioethanol sold in Switzerland		0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.2	0.6
Biogas											
on-road consumption (model)	1A3b	NO	0.03	0.03	0.03	0.1	0.1	0.2	0.2	0.3	0.3
non-road consumption (models)		NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Biogas sold in Switzerland		NO	0.03	0.03	0.03	0.05	0.1	0.2	0.2	0.3	0.3

Mileage-related activity data (basis for modelling of the non-CO₂ emissions by means of a traffic model)

The activity data are derived from different data sources:

- **Vehicle stock:** The federal vehicle registration database MOFIS (run by the Federal Roads Office FEDRO) contains vehicle stock data including all parameters needed for the emission modelling (vehicle category, engine capacity, fuel type, total weight, vehicle age and exhaust technology). The data are not public, but the ordinary vehicle stock numbers are published by the Swiss Federal Statistical Office (SFSO 2016c). The stock numbers from MOFIS are used for 1990–2010, whereas for 2011–2014 numbers are provided from a vehicle fleet projection by Prognos (2012a). With the help of a fleet turnover model the vehicle categories are split up into “sub-segments”, which are used to link with the specific emission factors of the same categorisation (vehicle category, size class, fuel type, emission standard [“Euro classes”], see also INFRAS 2010, TUG 2009).
- **The transport performance, i.e. the mileage** is calculated from the specific mileage per vehicle category (based on surveys/Mikrozensus ARE/SFSO 2005) times the number of vehicles. This figure is calibrated to the official statistics of traffic performance (SFSO 2009c and SFSO 2010c)¹⁰. For the period 2010–2015 the mileages are modelling results from Prognos (2012a) and ARE (2012).
- **Numbers of starts/stops:** Derived from vehicles stock, with data on trip length distributions and parking time distributions (ARE/SFSO 2005).

The transport performance is attributed to “traffic situations” (characteristic patterns of driving behaviour) which serve as a key to select the appropriate emission factor which are also available per traffic situation. The relative shares of the traffic situations are derived from a national road traffic model (operated by the Federal Office of Spatial Development, see ARE 2010). The traffic model is based on an origin-destination matrix that is assigned to a network of about 20'000 road segments. The model is calibrated partly bottom-up and partly top-down: bottom-up by a number of traffic counts from the national traffic-counter network (395 stations all over Switzerland, FEDRO 2010), and top-down by the total of the mileage per vehicle category. Furthermore, it supplies all the attributes needed for assigning a “traffic situation” to each road segment. The traffic model in combination with consumption factors (per vehicle category, size class, fuel type, emissions standard and per traffic situation) allows to calculate the **territorial** road traffic consumption of gasoline and diesel oil.

Table 3-77 shows the time series of the mileage per vehicle category. The total mileage has constantly been growing by about 1 per cent per year. The major part of vehicle kilometres was driven by passenger cars over the whole period. In the same period, on-road fuel consumption increased less strongly indicating improved fuel efficiency. This effect is also

¹⁰ As reported in the NIR of submission 2015 (FOEN 2015), a recalibration of the mileage per vehicle category has been performed based on the latest figures on population growth and economy (Prognos 2012a, ARE 2012). As a result vehicle kilometres from 1993 onwards were slightly lower in total; fleet compositions have changed, with slight impacts on implied emission factors; also fuel consumption in fuel tourism was slightly reduced; the modelled share of biofuels has been reduced to be consistent with real-world developments. However the overall impacts of these revisions on emissions have been minor.

reflected in Table 3-78 that depicts the specific fuel consumption per vehicle-km. For most vehicle categories, the specific consumption has decreased in the period 1990–2015.

Table 3-77 Mileages in millions of vehicle kilometres. PC: passenger cars, LDV: light duty vehicles, HDV: heavy duty vehicles).

Veh. category	1990	1995	2000	2005
	million vehicle-km			
PC	42'650	43'824	48'063	50'465
LDV	2'758	2'746	2'978	3'300
HDV	1'992	2'107	2'273	2'127
Coaches	108	110	99	106
Urban Bus	174	192	200	229
2-Wheelers	2'025	1'744	1'999	2'204
Sum	49'707	50'724	55'612	58'432
(1990=100%)	100%	102%	112%	118%

Veh. category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	million vehicle-km									
PC	50'812	51'208	51'949	52'852	53'341	54'000	54'730	55'424	56'085	56'731
LDV	3'374	3'473	3'529	3'584	3'621	3'663	3'701	3'735	3'765	3'793
HDV	2'189	2'203	2'223	2'172	2'210	2'250	2'290	2'329	2'369	2'407
Coaches	118	120	114	119	119	119	118	118	118	118
Urban Bus	233	240	245	249	251	254	257	261	264	267
2-Wheelers	2'262	2'300	2'366	2'385	2'407	2'436	2'465	2'494	2'523	2'552
Sum	58'989	59'544	60'426	61'361	61'950	62'722	63'562	64'362	65'123	65'868
(1990=100%)	119%	120%	122%	123%	125%	126%	128%	129%	131%	133%

Table 3-78 Specific fuel consumption of road transport, not including fuel tourism and statistical differences (PC: passenger cars, LDV: light duty vehicles, HDV: heavy duty vehicles).

Veh. cat.	Fuel	1990	1995	2000	2005
		MJ/veh-km			
PC	Gasoline	3.18	3.23	3.14	3.04
	Diesel	2.91	2.90	2.80	2.46
	CNG	0.00	0.00	0.00	0.00
LDV	Gasoline	3.17	3.18	3.18	3.19
	Diesel	3.86	3.86	3.75	3.42
HDV	Diesel	10.91	10.85	10.33	10.77
Coach	Diesel	11.84	11.69	11.33	11.22
Urban Bus	Diesel	16.22	16.29	15.80	15.37
	CNG	0.00	0.00	0.00	0.00
2-Wheeler	Gasoline	1.11	1.22	1.25	1.28
Average		3.46	3.54	3.42	3.24
		100%	102%	99%	94%

Veh. cat.	Fuel	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
		MJ/veh-km									
PC	Gasoline	2.99	2.97	2.93	2.90	2.86	2.81	2.77	2.71	2.66	2.61
	Diesel	2.46	2.41	2.40	2.35	2.33	2.30	2.28	2.24	2.21	2.17
	CNG	0.00	0.00	2.91	2.88	2.85	2.83	2.53	2.51	2.49	2.46
LDV	Gasoline	3.19	3.21	3.21	3.20	3.19	3.18	3.17	3.15	3.13	3.10
	Diesel	3.42	3.37	3.34	3.32	3.31	3.31	3.30	3.29	3.26	3.22
HDV	Diesel	10.77	10.71	10.73	10.65	10.59	10.55	10.50	10.46	10.41	10.37
Coach	Diesel	11.22	11.23	11.22	11.18	11.16	11.16	11.15	11.14	11.12	11.11
Urban Bus	Diesel	15.37	15.24	15.23	15.05	14.94	14.81	14.76	14.72	14.68	14.64
	CNG	0.00	0.00	20.34	20.32	20.36	20.58	20.52	20.46	20.38	20.31
2-Wheeler	Gasoline	1.28	1.29	1.31	1.33	1.35	1.34	1.34	1.34	1.34	1.33
Average		3.24	3.20	3.17	3.12	3.07	3.03	2.99	2.94	2.89	2.85
		94%	92%	91%	90%	89%	87%	86%	85%	84%	82%

For modelling of cold start and evaporative emissions of passenger cars and light duty vehicles, also vehicle stock and start numbers are used for activity data. The corresponding numbers are summarised in Table 3-79. Vehicle stock figures correspond to registration data. The starts per vehicle are based on specific surveys (ARE/SFSO 2005).

Table 3-79 Vehicle stock numbers and average number of starts per vehicle per day (PC: passenger cars, LDV: light duty vehicles).

Veh. Category	1990	1995	2000	2005
	stock in 1000 vehicles			
PC	2'985	3'229	3'545	3'862
LDV	221	238	260	291
2-Wheelers	764	704	732	770
starts per vehicle per day				
PC	2.61	2.53	2.46	2.40
LDV	1.97	1.97	1.96	1.96
2-Wheelers	1.59	1.54	1.50	1.54

Veh. Category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	stock in 1000 vehicles									
PC	3'894	3'956	3'990	4'010	4'076	4'195	4'302	4'396	4'477	4'548
LDV	298	307	312	317	326	328	331	334	337	339
2-Wheelers	784	789	804	807	816	815	815	816	818	820
starts per vehicle per day										
PC	2.39	2.38	2.37	2.35	2.34	2.34	2.33	2.33	2.32	2.32
LDV	1.96	1.96	1.96	1.96	1.96	1.96	1.96	1.96	1.96	1.96
2-Wheelers	1.54	1.55	1.56	1.56	1.57	1.57	1.57	1.58	1.58	1.58

Further details are given in Annex A3.1.3.

3.2.9.2.3 Railways (1A3c)

Methodology (1A3c)

As mentioned in chp. 3.2.4.5.1, the emissions are calculated by the non-road transportation model, using approach at

- Tier 2 for CO₂ (based on decision tree Fig. 3.4.1 in IPCC 2006)
- Tier 3 for CH₄, N₂O and precursors/SO₂ (based on decision tree Fig. 3.4.2 in IPCC 2006).

The entire Swiss railway system is electrified. Electric locomotives are used in passenger as well as freight railway traffic. Diesel locomotives are used for shunting purposes in marshalling yards and for construction activities only.

Emissions are calculated for the years 1990, 1995, 2000, 2005 etc. up to 2015 based on fuel used. For the years in-between, the emissions are interpolated linearly.

Emission factors (1A3c)

Only diesel oil is being used as fuel, therefore all emission factors refer to diesel oil.

- The CO₂ emission factor applied for the time series 1990–2015 for diesel oil is country-specific and is given in Table 3-12.
- The CH₄ and N₂O emission factors of diesel locomotives are shown in Table 3-80.
- For SO₂ the emission factors are country-specific. See also Table A – 19 (row diesel oil).
- The emission factors for precursors are country-specific and are given in FOEN (2015j).
- NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference between VOC and CH₄ emissions.
- Implied emission factors 2015 are shown in Table 3-81.

All emission factors (GHG, precursors, SO₂) can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

Table 3-80 CH₄ and N₂O emission factors for rail vehicles

Gas	Power class	Rail vehicles with diesel oil engines				
		PreEU <2000	UIC1 2000	UIC2 2003	EU3a 2006	EU3b 2012
	kW	g/kWh				
CH ₄	<18	0.0547	0.0384	0.024	0.0142	0.0142
CH ₄	18–37	0.0578	0.0221	0.0134	0.0089	0.0089
CH ₄	37–56	0.0319	0.0156	0.011	0.0079	0.0055
CH ₄	56–75	0.0319	0.0156	0.011	0.0079	0.0031
CH ₄	75–130	0.0218	0.0108	0.0084	0.0067	0.0031
CH ₄	>130	0.0218	0.0103	0.0072	0.0053	0.0031
N ₂ O	all	0.035	0.035	0.035	0.035	0.035

Table 3-81 Implied emission factors 2015 for rail vehicles. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A3c Railways	Unit	CO ₂ fossil	CO ₂ biogen	CH ₄	N ₂ O	NO _x	NMVOC	SO ₂	CO
Diesel oil	kg/TJ	73'300	NA	1.3	3.6	1'005	116	0.47	531
Biodiesel	kg/TJ	NA	73'300	1.1	3.1	874	101	0.40	461

Activity data (1A3c)

Activity data for non-road (1A3c) are described in chp. 3.2.4.5.1 (non-road transportation model). Values are taken from FOEN (2015j). Data on biofuels are provided by the statistics of renewable energies (SFOE 2016a). Activity data are shown in Table 3-82 and in Annex A3.1.4.

Table 3-82 Activity data (diesel oil consumption) for railways.

1A3c Railways	Unit	1990	1995	2000	2005
Diesel oil	TJ	390	441	455	472
Biodiesel	TJ	NO	NO	0.5	1.5
Total Railways	TJ	390	441	456	474
1990 = 100%		100%	113%	117%	121%

1A3c Railways	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Diesel oil	TJ	476	480	484	488	492	472	452	432	412	392
Biodiesel	TJ	1.5	1.6	1.7	1.8	1.8	2.3	2.7	3.1	3.5	4.0
Total Railways	TJ	478	482	486	490	494	474	455	435	415	396
1990 = 100%		123%	124%	125%	126%	127%	122%	117%	112%	107%	102%

Underlying activity data (vehicle stock, operating hours) of mobile non-road sources can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

3.2.9.2.4 Domestic navigation (1A3d)

Methodology (1A3d)

Based on the decision tree Fig. 3.5.1 Box 1 of the 2006 IPCC Guidelines (IPCC 2006) the emissions of navigation are calculated by a Tier 3 method with the non-road transportation model described in chp. 3.2.4.5.1.

There are passenger ships, dredgers, fishing boats, motor and sailing boats on the lakes and rivers of Switzerland. The emissions are calculated for the years 1990, 1995, 2000, 2005 etc. up to 2015 based on fuel used. For the years in-between, the emissions are linearly interpolated.

On the river Rhine as well as on the lakes of Geneva and of Constance, some of the boats cross the border. Fuels bought in Switzerland but used for international navigation are therefore reported as bunker fuels (memo items, chp. 3.2.2.)

Emission factors (1A3d)

- The CO₂ emission factor applied for the time series 1990–2015 for diesel oil, gasoline and gas oil are country-specific and are given in Table 3-12.
- The CH₄ and N₂O emission factors are country-specific and are shown below in the Table 3-83 to Table 3-85 for all fuel types and emission standards.
- For SO₂ the emission factors are country-specific. See also Table A – 19 in Annex A3.1.5 rows diesel oil, gasoline, gas oil.
- The emission factors for precursors are country-specific and are given in FOEN (2015j).
- NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference between VOC and CH₄ emissions.
- The implied emission factors 2015 are shown in Table 3-86.

All emission factors (GHG, precursors, SO₂) can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

Table 3-83 CH₄ and N₂O emission factors for ships with diesel engines.

Gas	Power class	Ships with diesel oil engines				
		PreSAV (<1995)	SAV 1995	EU-I 2003	EU-II 2008	EU-III A 2009
		g/kWh				
CH ₄	<18	0.0547	0.0547	0.0384	0.024	0.0142
CH ₄	18–37	0.0578	0.0578	0.0221	0.0134	0.0089
CH ₄	37–56	0.0319	0.0319	0.0156	0.011	0.0079
CH ₄	56–75	0.0319	0.0319	0.0156	0.011	0.0079
CH ₄	75–130	0.0218	0.0218	0.0108	0.0084	0.0067
CH ₄	>130	0.0218	0.0218	0.0103	0.0072	0.0053
N ₂ O	all	0.035	0.035	0.035	0.035	0.035

Table 3-84 CH₄ and N₂O emission factors for ships with gasoline engines by emission standards including the year of enforcement.

Gas	Power class	boats with 2-stroke gasoline engines			boats with 4-stroke gasoline engines		
		PreSAV <1995	SAV 1995	SAV/EU 2007	PreSAV <1995	SAV 1995	EU 2007
		g/kWh					
CH ₄	<4.4	18.2	1.54	1.75	1.25	1.10	1.25
CH ₄	4.4–7.4	18.2	0.84	0.91	1.00	0.60	0.65
CH ₄	7.4–37	18.2	0.42	0.56	1.00	0.30	0.40
CH ₄	37–74	18.2	0.42	0.56	1.00	0.20	0.30
CH ₄	74–100	18.2	0.42	0.56	1.00	0.17	0.25
CH ₄	>100	18.2	0.42	0.56	1.00	0.10	0.25
N ₂ O	0–300	0.01	0.01	0.01	0.03	0.03	0.03

Table 3-85 CH₄ and N₂O emission factors for steamboats by the year of enforcement.

Gas	steamboats		
	<2000	2000-2004	>2004
	g/kWh		
CH ₄	0.0218	0.0103	0.0072
N ₂ O	0.035	0.035	0.035

Table 3-86 Implied emission factors 2015 for navigation. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A3d Navigation	Unit	CO ₂ fossil	CO ₂ biog.	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
Gasoline	kg/TJ	73'800	NA	20.0	1.9	542	368	0.7	7'471
Diesel oil	kg/TJ	73'300	NA	1.8	3.4	933	293	0.5	533
Gas oil	kg/TJ	73'700	NA	0.2	0.7	26	1.6	12	6.9
Biodiesel	kg/TJ	NA	73'300	1.6	3.0	811	255	0.4	463
Bioethanol	kg/TJ	NA	73'800	11.6	1.3	343	221	0.2	4'619

Activity data (1A3d)

Activity data for navigation (1A3d) are described in chp. 3.2.4.5.1 (non-road transportation model). Values are taken from FOEN (2015j). Data on biofuels is provided by the statistics of renewable energies (SFOE 2016a). Activity data are shown in Table 3-87 and in Annex A3.1.4.

Table 3-87 Fuel consumption of (domestic) navigation.

1A3d Navigation	Unit	1990	1995	2000	2005
Gasoline	TJ	701	654	616	565
Diesel oil	TJ	738	724	792	800
Gas oil	TJ	110	139	147	150
Biodiesel	TJ	NO	NO	0.9	2.5
Bioethanol	TJ	NO	NO	NO	0.1
Total Navigation	TJ	1'550	1'517	1'556	1'518
1990 = 100%		100%	98%	100%	98%

1A3d Navigation	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Gasoline	TJ	814	827	841	855	868	871	874	877	879	882
Diesel oil	TJ	559	553	547	541	534	529	524	518	513	508
Gas oil	TJ	152	154	155	157	159	157	156	154	153	151
Biodiesel	TJ	2.6	2.8	2.9	3.1	3.2	4.4	5.5	6.6	7.8	8.9
Bioethanol	TJ	0.1	0.1	0.1	0.2	0.2	2.2	4.2	6.2	8.1	10.1
Total Navigation	TJ	1'527	1'537	1'546	1'556	1'565	1'564	1'563	1'562	1'561	1'560
1990 = 100%		99%	99%	100%	100%	101%	101%	101%	101%	101%	101%

Underlying activity data (vehicle stock, operating hours) of mobile non-road sources can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

3.2.9.2.5 Other transportation (1A3e)

Methodology (1A3e)

The emissions are calculated with a Tier 2 method (the 2006 IPCC Guidelines (IPCC 2006) do not contain a decision tree to determine the Tier level specifically).

Source 1A3e includes only pipeline transportation (1A3e i) from a compressor station located in Ruswil. Emissions of CO₂, CH₄, N₂O, NO_x, CO, NMVOC and SO₂ are reported. The

compressor station uses a centrifugal compressor according to Transitgas AG (the company operating the compressor station and the pipeline network).

Emission factors (1A3e)

- The CO₂ emission factor applied for the time series 1990–2015 for natural gas is country-specific and is given in Table 3-12.
- The CH₄ emission factor corresponds to the one used for gas turbines in Switzerland (SAEFL 2000) as suggested by expert judgement. The CH₄ EF is assumed to be 5 g/GJ up to 1995 and 2 g/GJ from 2000 onwards, with linear interpolation in between. This corresponds with the fact that a catalyst was fitted to the system, which reduced the CH₄ emissions of the gas turbine. For the value 2014 see Table 3-88.
- For N₂O emission factors the IPCC 2006 default value is used as displayed in Table 3-88.
- For SO₂ the emission factors are country-specific. The emission factor 2015 is shown in Table 3-88. See also Table A – 19 in Annex A3.1.5 row natural gas.
- The emission factors for precursors are country-specific and are given in SAEFL (2000); see also EMIS 2015/1A3e “Gasturbinen; Erdgas”.

Table 3-88 Emission factors of 1A3e i Pipeline transportation / compressor station located in Ruswil in 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A3e Other transportation	Unit	CO ₂ fossil	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
Gas	kg/TJ	56'400	2	0.1	60	0.1	0.5	15

Activity data (1A3e)

The data on fuel consumption for the operation of the compressor station in Ruswil is based on the Swiss overall energy statistics (SFOE 2016; Table 17).

Table 3-89 Activity data of 1A3e.

1A3ei Pipeline transport	Unit	1990	1995	2000	2005
Natural gas	TJ	560	310	340	1'070
1990=100%		100%	55%	61%	191%

1A3ei Pipeline transport	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Natural gas	TJ	1'700	1'430	1'460	950	830	840	810	410	830	760
1990=100%		304%	255%	261%	170%	148%	150%	145%	73%	148%	136%

3.2.9.3 Uncertainties and time-series consistency for 1A3

For a general description of the uncertainty analysis and time series consistency of the Energy sector see chp. 3.2.4.7 where uncertainties of activity data and emission factors of fuels are shown (Table 3-25) and explained in detail.

Consistency: Time series for 1A3 Transport are all considered consistent.

3.2.9.4 Category-specific QA/QC and verification for 1A3

General

The general QA/QC measures are described in chp. 1.2.3 and partly also in chp. 3.2.4.8.

Specific: Domestic aviation (1A3a)

Emissions

Total calculated emissions for domestic and international flights have been compared between different years. The development of total emissions with time is consistent with a fleet renewal of former Swissair in the early nineties, the technological improvements and changes in fleet composition.

Emission factors

- From total fuel burnt, total distance, number of passenger (without freight) per aircraft type, the fuel consumption per 100 passenger km has been calculated (backward calculation). The result of 2 to 10 kg fuel/100 passenger km is in line with expectations for 1990 passenger fleets.
- The implied emission factors were calculated for 2015 and compared with previous years.

Activity data

- In an independent Tier 3B calculation, EUROCONTROL performed a fuel calculation for Switzerland's international flights, based on collected flight plan data and single movements. The results for the years 2004, 2005 and 2007 matched the FOCA calculations by more than 97.4%. The FOCA results were generally 1% to 2% higher but included the total number of actual flight movements of all flights, including VFR (visual flight rules) and non-scheduled flights such as helicopter movements in alpine regions.
- Comparison between total movement numbers in the calculation and in the corresponding published statistics. Example: In 1990 calculation, FOCA considered all flights for which there was a form 'Traffic report to the airport authorities' filled in (total heavy aircraft). The total number of movements in 1990 is 263'951 (without Basel). The published number of movements for scheduled and charter flights in 1990 is: 263'952 (without Basel).
- The bottom-up calculation of total fuel matches the total fuel sold within a few percents.
- Real-world fuel consumption was compared with modelled consumption for selected aircraft of four Swiss airlines. The difference between the two methods was smaller than 1%.

Specific: Road transportation (1A3b)

Comparison between the 2006 IPCC Guideline's default and Switzerland's emission factors

- CO₂ (see also Table 3-26): IPCC default value for gasoline is 69.3 t/TJ and for diesel oil 74.1 t/TJ (IPCC 2006, Table 3.2.1). Switzerland's emission factors vary between 73.8 and 73.9 t/TJ for gasoline – 6% higher than IPCC – and between 73.3 and 73.6 t/TJ for diesel oil – about 1% below IPCC default value.
- CH₄: The IPCC default emission factor for gasoline motors with oxidation catalysts is 25 kg/TJ with an uncertainty range from 7.5 to 86 kg/TJ (IPCC 2006, Table 3.2.2). Switzerland's emission factor for gasoline passenger cars dropped from 26.6 kg/TJ (1990) to 5.0 kg/TJ (2015) and is therefore in the lower part of IPCC's uncertainty range. For diesel oil the IPCC default emission factors lies in the range of 1.6-9.5 kg/TJ, whereas Switzerland's range is lower 0.3-1.5 kg/TJ.
- N₂O: The IPCC default emission factor for gasoline motors with oxidation catalysts lies in the uncertainty range 2.6-24 kg/TJ (IPCC 2006, Table 3.2.2). Switzerland's emission factor for gasoline passenger cars dropped from 4.2 kg/TJ (1997) to 0.8 kg/TJ (2015) and is therefore in the lower part of and below IPCC's uncertainty range. For diesel oil the IPCC default emission factors lies in the range of 1.3-12 kg/TJ, whereas Switzerland's range is lower between 0.2 kg/TJ (1990) and 2.5 kg/TJ (2015).

The international project for the update of the emission factors for road vehicles is overseen by a group of external national and international experts that guarantees an independent quality control. For the update of the modelling of Switzerland's road transport emissions, which has been carried out between 2008 and 2010, several experts from the federal administration have conducted the project. The results have undergone extensive plausibility checks and comparisons with earlier estimates.

The emission factors CH₄ and N₂O used for the modelling of 1A3b Road Transportation are taken from the handbook of emission factors (INFRAS 2010), which is also applied in Germany, Austria, Netherlands, and Sweden. The Swiss emission factors for CH₄ and N₂O used in 1A3b were additionally compared with those depicted in the CRF from Germany and a good match was found. Possible small differences might result from a varying fleet composition.

For gasoline, the activity data is easily verified due to the fact, that 98% of the gasoline sold 2015 in Switzerland is consumed by 1A3b Road Transportation itself. Therefore, the amount of gasoline reported in the Swiss overall energy statistics is a strong control and verification parameter for the activity data of 1A3b.

3.2.9.5 Category-specific recalculations for 1A3

- 1A3a: As recommended by the ERT and in line with the IPCC reporting GLs there are no more CH₄ emissions for cruise activities. This implies higher NMVOC emissions for cruise activities. All years 1990–2014 have been recalculated.
- 1A3b: The CO₂ emission factor for biogas used in road transportation was out-dated and has now been adapted 1990–2014 to the one of natural gas.
- 1A3b/1A3d: Small recalculation due to a change in the NCV of diesel used in international navigation to equalise with other diesel processes. Therefore, small changes

occurred 1990–2014 in fuel tourism and statistical difference for diesel which is integrated in 1A3biii.

- 1A3bi cars and 1A3bii LDT: N₂O cold start excess emissions for PC and LDV by means of emission factors of the Copert model have been added for 1990, 2013 and 2014. For the years 1991–2012 the emissions were linearly interpolated between 1990 and 2013. Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

3.2.9.6 Category-specific planned improvements for 1A3

- 1A3a Civil aviation: No category-specific improvements are planned.
- 1A3b Road Transportation: A general update of the emission factors and the activity data is on-going. The results will be presented in future submissions.
- 1A3b: Note that an inconsistency in the attribution of natural gas to the vehicle categories leads to an error in the implied emission factors for gas-driven light duty vehicles. The error will be corrected within the general update mentioned above.
- 1A3c, 1A3d, 1A3e: No category-specific improvements are planned.

3.2.10 Source category 1A4 - Other sectors (mobile)

3.2.10.1 Source category description for 1A4 Other sectors (mobile)

Key categories 1A4

See key categories mentioned in chp. 3.2.7.1, which are vastly dominated by the emissions of the stationary sources.

CO₂ from the combustion of Liquid Fuels in 1A4c Agriculture/Forestry/Fishing (level).

Table 3-90 Specification of source category 1A4 Other sectors (mobile 1A4 aii/bii/cii).

1A4	Source	Specification
1A4a ii	Commercial/ institutional	Emission from non-road vehicles (professional gardening) and motorised equipment
1A4b ii	Residential	Emissions from mobile machinery (hobby, gardening) and motorised equipment
1A4c ii	Agriculture/forestry	Emissions from non-road vehicles and machinery in agriculture and forestry

3.2.10.2 Methodological issues for 1A4 Other sectors (mobile)

Methodology (1A4 Other sectors, mobile)

Based on the decision tree Fig. 3.3.1 in chp. “3. Mobile Combustion” in the 2006 IPCC Guidelines (IPCC 2006), the emissions of vehicles and machinery in 1A4 are calculated by a Tier 3 method with the non-road transportation model described in chp. 3.2.4.5.1.

Emission factors (1A4 Other sectors, mobile)

In the categories 1A4a ii and 1A4b ii only gasoline is being used as fuel. In category 1A4c ii mainly diesel oil is consumed and only a small amount of gasoline (e.g. chainsaws).

- The CO₂ emission factors applied for the time series 1990–2015 are country-specific and are given in Table 3-12.
- The CH₄ and N₂O emission factors are country-specific and are shown in Table 3-66 and Table 3-67 for diesel oil and gasoline engines for all emission standards.
- For SO₂ the emission factors are country-specific. See also Table A – 19 in Annex A3.1.5 for diesel oil, gasoline, gas oil.
- The emission factors for precursors are country-specific and are given in FOEN (2015j).
- NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference between VOC and CH₄ emissions.
- Implied emission factors 2015 are shown in Table 3-91.

All emission factors (GHG, precursors, SO₂) can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

Table 3-91 Implied emission factors 2015 for 1A4 Other sectors (1A4a ii – 1A4c ii mobile). Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A4 Non-road machinery	Unit	CO ₂ fossil	CH ₄	N ₂ O	NO _x	NMVOC	SO ₂	CO
1A4a ii Gardening professional, gasoline	kg/TJ	73'800	92	1.1	185	1'468	5.2	26'422
1A4b ii Gardening, gasoline	kg/TJ	73'800	50	1.4	159	1'025	2.0	25'023
1A4c ii Forestry and agriculture, gasoline	kg/TJ	73'800	88	1.1	174	1'544	3.6	23'900
1A4c ii Forestry and agriculture, diesel oil	kg/TJ	73'300	1.6	3	518	61	0.5	307

Activity data (1A4 Other sectors, mobile)

Activity data are described in chp. 3.2.4.5.1 (non-road transportation model) and are shown in Table 3-92 and in Annex A3.1.4.

Table 3-92 Activity data for non-road vehicles and machinery in 1A4 Other sectors (mobile).

1A4 Non-road machinery	Unit	1990	1995	2000	2005
1A4a ii Gardening professional, gasoline	TJ	191	245	295	295
1A4b ii Gardening, gasoline	TJ	142	155	165	165
1A4c ii Forestry and agriculture, gasoline	TJ	1'160	1'070	963	823
1A4c ii Forestry and agriculture, diesel oil	TJ	4'269	4'604	4'921	4'804
Total 1A4 non-road machinery	TJ	5'761	6'073	6'343	6'088
1990=100%		100%	105%	110%	106%

1A4 Non-road machinery	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1A4a ii Gardening professional, gasoline	TJ	293	292	290	288	287	280	272	265	258	251
1A4b ii Gardening, gasoline	TJ	165	164	164	163	163	161	160	158	156	155
1A4c ii Forestry and agriculture, gasoline	TJ	797	770	743	716	689	664	638	613	588	562
1A4c ii Forestry and agriculture, diesel oil	TJ	4'820	4'836	4'852	4'868	4'884	4'883	4'881	4'880	4'878	4'876
Total 1A4 non-road machinery	TJ	6'075	6'062	6'049	6'036	6'023	5'987	5'952	5'916	5'880	5'845
1990=100%		105%	105%	105%	105%	105%	104%	103%	103%	102%	101%

Underlying activity data (vehicle stock, operating hours) of mobile non-road sources can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

3.2.10.3 Uncertainties and time-series consistency for 1A4 Other sectors, mobile

Uncertainties by fuel type are given in Table 3-25.

3.2.10.4 Category-specific QA/QC and verification for 1A4 Other sectors, mobile

The general QA/QC procedures are described in chp. 1.2.3. Furthermore QA/QC procedures conducted for all 1A source categories are listed in chp. 3.2.4.8.

3.2.10.5 Category-specific recalculations for 1A4 Other sectors, mobile

No category-specific recalculations were carried out.

3.2.10.6 Category-specific planned improvements for 1A4 Other sectors, mobile

No category-specific improvements are planned.

3.2.11 Source category 1A5b - Other (mobile)

3.2.11.1 Source category description for 1A5b (mobile)

Source category 1A5b – Other (mobile) is not a key category.

All of the Swiss source categories of 1A5 refer to mobile sources of military activities (1A5b). Stationary activities are not occurring.

Table 3-93 Specification of Swiss source category 1A5 Other.

1A5	Source	Specification
1A5bi	Military aviation	Emissions from military aircrafts
1A5bii	Military non-road vehicles and machines	Emissions from machines like power generators, tanks, bulldozers, boats etc.

3.2.11.2 Methodological issues for 1A5b Other (mobile)

3.2.11.2.1 Military aviation (1A5b i)

Methodology (1A5b i Other, military aviation)

To calculate the emissions from military aviation, a Tier 2 method is used.

Emission factors (1A5b i Other, military aviation)

Emissions of NO_x, CO and VOC have been modelled in detail by the Federal Office for Military Aviation (Bundesamt für Betriebe der Luftwaffe) for 1990 and 1995. From these inputs, FOEN determined average emission factors 1990 and 1995. For 1991–1994 the emission factors are linearly interpolated. For 1996–2014, the factors for 1995 are used. The emissions are then calculated yearly based on average emission factors.

The extension of the emission modelling to CO₂, CH₄, N₂O, NMVOC and SO₂ is also accomplished by FOEN.

- The CO₂ emission factor applied for the time series 1990–2015 for kerosene is country-specific and is given in Table 3-12.
- CH₄: The division of VOC into CH₄ and NMVOC is carried out as described for civil aviation (see chp. 3.2.9.2.1).
- N₂O: The emission factor is set equal to the emission factor of 1A3a Civil aviation (FOCA 2015).
- NO_x, VOC, CO: Engine producer information is used (CORINAIR, for details see SAEFL 1996: p. 202) for calculation of the emission factors in 1990 and 2000. For 1991–1999 the values are linearly interpolated between 1990 and 2000. For 2001–2015, the values 2000 are used.
- SO₂: The emission factor is taken from the EMEP/EEA Guidebook (EMEP/EEA 2013) and is assumed to be constant over the period 1990–2015.

Table 3-94 Implied emission factors 1A5b i military aviation in 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A5bi Military aviation	Unit	CO ₂ fossil	CH ₄	N ₂ O	NO _x	NMVOC	SO ₂	CO
Jet kerosene	kg/TJ	72'800	3.6	2	133	32	23	672

Activity data (1A5b i Other, military aviation)

The fuel consumption 1990–2015 is known on an annual basis (DDPS 2016). A very small fraction of fuel is consumed for training abroad and might be allocated under “International aviation” (less than 3% of total military aviation consumption). Since the exact numbers for the fuels used abroad is not known, it is not subtracted from the total consumption but included under national military aviation, as recommended by the IPCC Guidelines (2006, chp. 3.6.1.4).

Table 3-95 Activity data (fuel consumption) for military aviation.

1A5 Other Military aviation	1990	1995	2000	2005
	fuel consumption in TJ			
Jet kerosene	2'733	1'955	1'794	1'624

1A5 Other Military aviation	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	fuel consumption in TJ									
Jet kerosene	1'676	1'577	1'505	1'529	1'592	1'420	1'527	1'542	1'615	1'567

3.2.11.2 Military non-road vehicles (1A5b ii Other, military machinery)

Methodology (1A5b ii Other, military machinery)

Emissions are calculated as part of the non-road transportation model (chp. 3.2.4.5.1) corresponding to a Tier 3 according to the the decision tree Fig. 3.3.1 in chp. 3. Mobile Combustion in IPCC (2006).

Emission factors (1A5b ii Other, military machinery)

- The CO₂ emission factors applied for the time series 1990–2015 for diesel oil, gasoline and compressed natural gas are country-specific as shown in Table 3-12.
- The CH₄ and N₂O emission factors are country-specific and are shown in Table 3-66 to Table 3-68 for diesel oil and gasoline engines for all emission standards.
- For SO₂ the emission factors are country-specific. See also Table A – 19 in Annex A3.1.5, rows diesel oil, gasoline.
- The emission factors for precursors are country-specific and are given in FOEN (2015j).
- NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference between VOC and CH₄ emissions.
- Implied emission factors are shown in Table 3-96.

All emission factors (GHG, precursors) can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

Table 3-96 Implied emission factors 1A5b ii military non-road vehicles 2015. Emission factors that are highlighted in green are described in chp. 3.2.4.4.

1A5bii Military non-road	Unit	CO ₂ fossil	CO ₂ biog.	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
Gasoline	kg/TJ	73'800	NA	44.8	1.5	134	817	1.8	24077
Diesel	kg/TJ	73'300	NA	1	3	465	37	0.5	194
Biodiesel	kg/TJ	NA	73'300	0.8	2.6	404	32	0.4	169
Bioethanol	kg/TJ	NA	73'800	10.9	1	72	305	0.2	15052

Activity data (1A5b ii Other, military machinery)

Activity data for military non-road vehicles (1A5b ii) are described in chp. 3.2.4.5.1 (non-road transportation model). Values are taken from FOEN (2015j). Data on biofuels are provided by the statistics of renewable energies (SFOE 2016a). Activity data are shown in Table 3-97 and in Annex A3.1.4.

Table 3-97 Activity data (fuel consumption) for military non-road vehicles.

1A5bii Military non-road	1990	1995	2000	2005
	fuel consumption in TJ			
Military non-road	239	248	252	257
Gasoline	19	19	19	19
Diesel	220	228	233	238
Biodiesel	NO	NO	0.3	0.7
Bioethanol	NO	NO	NO	0.0

1A5bii Military non-road	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	fuel consumption in TJ									
Military non-road	261	265	268	272	275	275	275	275	275	275
Gasoline	19	19	18	18	18	18	17	17	17	17
Diesel	241	245	249	253	256	256	256	256	255	255
Biodiesel	0.8	0.8	0.9	0.9	1.0	1.3	1.6	1.9	2.3	2.6
Bioethanol	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.3

Underlying activity data (vehicle stock, operating hours) of mobile non-road sources can be downloaded by query from the public part of the non-road database INFRAS (2015a). They can be queried by vehicle type, fuel type, power class and emission standard either at aggregated or disaggregated levels (see footnote 8).

3.2.11.3 Uncertainties and time-series consistency for 1A5b Other (mobile)

For a general description of the uncertainty analysis and time series consistency of the Energy sector see chp. 3.2.4.7 where uncertainties of activity data and emission factors of fuels are shown (Table 3-25) and explained in detail.

Consistency: Time series for 1A5b Other are all considered consistent.

3.2.11.4 Category-specific QA/QC and verification for 1A5b Other (mobile)

The general QA/QC measures are described in chp. 1.2.3 and partly also in chp. 3.2.4.8.

The activity data of military aviation (1A5b), kerosene consumption, is provided by the Federal Department of Defence, Civil Protection and Sport. For a compatibility check with the emission data base of civil aviation, they are sent to the FOCA (office of the Federal Department of the Environment, Transport, Energy and Communications).

3.2.11.5 Category-specific recalculations for 1A5b Other (mobile)

No category-specific recalculations were carried out.

3.2.11.6 Category-specific planned improvements for 1A5b Other (mobile)

No category-specific improvements are planned.

3.3 Source category 1B - Fugitive emissions from fuels

3.3.1 Source category description for 1B

Table 3-98 Key categories (KCA incl. LULUCF) of 1B Fugitive emissions from fuels.

Code	IPCC Category	GHG	Identification Criteria
1B2	Oil and Natural Gas Energy Production	CH4	L1, T1, T2

The only relevant source categories of fugitive emissions in Switzerland are:

- Oil (1B2a)
- Natural gas (1B2b)
- Venting and flaring (1B2c)

3.3.2 Source category 1B1 – Solid Fuels

Coal mining is not occurring in Switzerland. There are no greenhouse gas emissions from coal handling.

3.3.3 Source category fugitive emissions from 1B2a – Oil

3.3.3.1 Source category description for 1B2a

In Switzerland, oil production is not occurring. Fugitive emissions in the oil industry result exclusively from the refineries and several fuel handling stations. At the beginning of 2015, one of the two refineries ceased operation. The extents of the two existing oil pipelines in Switzerland are approximately 40 km and 70 km, respectively. The pipelines are mainly laid underground.

Table 3-99 Specification of source category fugitive emissions from 1B2a Oil in Switzerland.

1B2	Source	Specification
1B2a	Fugitive emissions attributed to oil	Emissions from refining/storage of oil and the distribution of oil products: transport of crude oil in pipelines.

3.3.3.2 Methodological issues for 1B2a

Methodology (1B2a)

According to the decision tree for crude oil transport, refining and upgrading, Switzerland estimates 1B2a fugitive emissions from oil based on a Tier 1 (1B2a iii) and a Tier 2 (1B2a iv, v) approach (IPCC 2006, Volume 2 Energy, chp. 4 Fugitive Emissions, Figure 4.2.3). For emission calculations plant-specific activity data are available.

For source 1B2a fugitive emissions from oil, fugitive emissions of CH₄ are reported, which occur only in 1B2a iii Transport and 1B2a iv Refining/storage. Indirect CO₂ emissions resulting from NMVOC emissions in this source category are reported in chp. 9.

Emission factors (1B2a)

For oil transport (1B2a iii), the default emission factors from the 2006 IPCC Guidelines for pipeline transportation are used to calculate emissions. Values provided in Table 3-100 are converted using a crude oil density of 0.82 t/m³.

For oil refining and storage (1B2a iv), plant-specific emission factors for CH₄ and NMVOC are used. The emission factors for CH₄ are delineated from an emission estimation project in one of the refineries in 1992 called CRISTAL. The estimation from the other refinery is assumed to be twice as high, because the technology of the plant is older. Then a weighted mean based on the quantity of crude oil used in both refineries was calculated (for further details see the internal documentation of the EMIS database, EMIS 2017/1B2a iv). The emission factors for SO₂ emissions from Claus units in refineries are country-specific and based on measurements and data from industry and expert estimates.

For oil distribution from storage tanks and gasoline stations (1B2a v), the NMVOC emission factor for oil distribution from tanks and gasoline stations is country-specific, based on a model which takes annual gasoline sales and technical equipment of gasoline stations and storage tanks into account (see internal database documentation in EMIS 2017/1B2a v Benzinumschlag Tanklager and EMIS 2017/1B2a v Benzinumschlag Tankstellen). An expert team (Weyer and Partner AG) is in charge of providing annual updates of the modelled NMVOC emissions based on their own database of Swiss storage tanks and gasoline vapour recovery systems. The model is calibrated with spot checks of the gas recovery systems of gas stations.

Table 3-100 Emission factors for fugitive emissions of source category 1B2a Oil in 2015.



Source/fuel	Unit	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
1B2a Oil								
Exploration	g/t	NO	NO	NO	NO	NO	NO	NO
Production	g/t	NO	NO	NO	NO	NO	NO	NO
Transport	g/t	NA	6.59	NA	NA	65.9	NA	NA
Refining/Storage	g/t	NA	45	NA	NA	430	38	NA
Distribution of oil products:								
Gasoline storage tank	g/GJ	NA	NA	NA	NA	7	NA	NA
Distribution of oil products:								
Gasoline station	g/GJ	NA	NA	NA	NA	9	NA	NA
Source/fuel	Unit	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
1B2a Oil								
Exploration	g/t	NO	NO	NO	NO	NO	NO	NO
Production	g/t	NO	NO	NO	NO	NO	NO	NO
Transport	g/t	NA	6.59	NA	NA	65.9	NA	NA
Refining/Storage	g/t	NA	45	NA	NA	430	38	NA
Distribution of oil products:								
Gasoline storage tank	g/GJ	NA	NA	NA	NA	7	NA	NA
Distribution of oil products:								
Gasoline station	g/GJ	NA	NA	NA	NA	9	NA	NA

Activity data (1B2a)

For oil transport (1B2a iii) and oil refining and storage (1B2a iv), activity data (crude oil use in the two refineries) are based on annual statistics of the Swiss petroleum association (EV 2016). The annual amount of processed crude oil in Claus units is based on the Swiss overall energy statistics (SFOE 2016).

For oil distribution from storage tanks and gasoline stations (1B2a v), gasoline sales based on the Swiss overall energy statistics (SFOE 2016), corrected for consumption of Liechtenstein, are used as activity data.

In analogy to Figure 3-11 to Figure 3-19, Figure 3-25 shows how the energy model (chp. 3.2.4.3.3) attributes imported crude oil (as provided by the Swiss overall energy statistics, SFOE 2016), losses of gasoline and of natural gas to source category 1B.






Figure 3-25 Attribution of fugitive emissions to source categories of 1B Fugitive emissions from fuels.

Table 3-101 Activity data for fugitive emissions from 1B2a Oil.

1B2a Oil products	Unit	1990	1995	2000	2005
Crude oil	kt	3'127	4'657	4'649	4'877
Gasoline transport	TJ	157'335	152'575	169'331	152'955

1B2a Oil products	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Crude oil	kt	5'563	4'720	5'133	4'833	4'546	4'452	3'455	4'935	4'975	2'836
Gasoline transport	TJ	148'256	146'832	143'620	139'726	134'723	129'506	124'969	119'280	114'466	106'074

3.3.3.3 Uncertainties and time-series consistency for 1B2a

Based on expert judgement, a preliminary uncertainty assessment of all sources in source category 1B2 results in medium confidence in the emissions estimate (see Table 1-10).

Consistency: Time series for 1B2a Oil are all considered consistent.

3.3.3.4 Category-specific QA/QC and verification for 1B2a

The general QA/QC measures are described in chp. 1.2.3 and partly also in chp. 3.2.4.8. No further source specific activities undertaken for fugitive emissions from oil (1B2a).

3.3.3.5 Category-specific recalculations for 1B2a

The following recalculations were implemented in submission 2017. These recalculations cause a change in emission levels in 1990 or 2014 by less than 1 kt CO₂.

- 1B2a: Small recalculations in activity data due to change of units (1990–2014).

- 1B2a: SO₂ emissions from Claus units were previously reported in source category 2H3 and are now reported in 1B2a iv.

3.3.3.6 Category-specific planned improvements for 1B2a

No category-specific improvements are planned.

3.3.4 Source category fugitive emissions from 1B2b – Natural gas

3.3.4.1 Source category description for 1B2b

Emissions from natural gas production (1B2b ii) are only occurring for the years of operation of the single production plant in Switzerland from 1985–1994. Other emissions in this source category occur from natural gas transmission (1B2b iv) and distribution (1B2b v). Emissions from accidents in the gas pipeline system are reported under source category 1B2b vi Other Leakage.

Table 3-102 Specification of source category fugitive emissions from 1B2b Natural gas in Switzerland.

1B2	Source	Specification
1B2b	Fugitive emissions attributed to natural gas	Emissions from gas network

3.3.4.2 Methodological issues for 1B2b

Methodology (1B2b)

According to the decision tree for natural gas systems (IPCC 2006, Volume 2 Energy, chp. 4 Fugitive Emissions, Figure 4.2.1), Switzerland follows a Tier 1 approach for fugitive emissions concerning 1B2b ii Production and a Tier 2 approach for fugitive emissions attributed to 1B2b iv Transmission and storage as well as 1B2b v Distribution.

Emissions from source category 1B2 are key. However, the contribution from 1B2b ii is small and therefore the use of a Tier 1 method for this source category is justified. The emissions from source category 1B2b ii are calculated based on annual production data and default emission factors (IPCC Tier 1 approach). Production data under 1B2b ii are only available for the years 1990–1994 because the single production site was closed in 1994.

For emission calculations from source category 1B2b iv, 1B2b v and 1B2b vi country-specific emission factors and activity data are available. Emissions are calculated with a country-specific method which first assesses the losses of natural gas in the gas network including pipelines, fittings and gas devices, as these data represent the activity data. Based on the gas losses, CO₂, CH₄ and NMVOC emissions are calculated with country-specific emission factors which reflect the composition of the gas lost.

Emissions from gas transmission (source category 1B2b iv) include emissions from transport pipelines including the transit pipeline and the single compressor station. Emissions

comprise leakages from gas pipelines, small-scale damages, maintenance work and leakages of pipeline fittings. Gas storages are considered as components of the distribution network and the respective emissions are included in source category 1B2b v.

Source category 1B2b v Distribution covers emissions from the gas distribution pipelines and network components (e.g. control units, fittings and gas meters) as well as fugitive emissions at the end users. Emission calculations for the gas distribution network are based on the length, material and pressure of the gas pipelines. Fugitive emissions at the end users arise from on-site and indoor pipelines and the permanent leakiness of the different gas appliances in households, industry and natural gas fuelling stations. In the calculations, the number and kind of end users and connected gas appliances are considered.

Emission factors (1B2b)

For natural gas production, CO₂, CH₄ and NMVOC default emission factors are taken from the 2006 IPCC Guidelines (IPCC 2006) as documented in the internal emission database documentation (EMIS 2017/1B2b Diffuse Emissionen Erdgas).

Emission factors for transmission, distribution and other leakages (source category 1B2b iv 1B2b v and 1B2b vi) are calculated based on the average CO₂, CH₄, and NMVOC concentrations of natural gas and its average lower calorific value in Switzerland as described in Quantis (2014) and EMIS 2017/1B2b Diffuse Emissionen Erdgas. Since 2012, the annual average CH₄ concentration and lower calorific value of natural gas in Switzerland are provided by the Swiss Gas and Water Industry Association (SGWA).

Table 3-103- Emission factors for fugitive emissions of source category 1B2b Natural gas in 2015.

1B2b Natural gas	CO₂	CH₄	N₂O	NO_x	NMVOC	SO₂	CO
	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ
1B2b ii Production	NO	NO	NO	NO	NO	NO	NO
1B2b iv Transmission	309	18'082	NA	NA	1'434	NA	NA
1B2b v Distribution	309	18'082	NA	NA	1'434	NA	NA
1B2b vi Other Leakage	309	18'082	NA	NA	1'434	NA	NA

Activity data (1B2b)

Activity data for fugitive emissions from gas production (1B2b ii) are the actual gas production data for the years 1990–1994 (SFOE 2016).

For gas transmission (1B2b iv), distribution (1B2b v), and other leakage (1B2b vi), the activity data have been reassessed in a recent study by Quantis (2014) and updated in 2016 (EMIS 2017/1B2b Diffuse Emissionen Erdgas). The activity data represent the amount of natural gas lost from the gas network and are shown in Table 3-104.

For source categories 1B2b iv and 1B2b v, information regarding the gas transport and distribution network from the Swiss Gas and Water Industry Association (SGWA) is used to derive the activity data (see Quantis 2014 and EMIS 2017/1B2b Diffuse Emissionen Erdgas).

For transmission pipelines a constant emission factor per pipeline length is applied accounting for losses from purging and cleaning flows, pipeline damages and leaky fittings and mountings. For the one compressor station a constant emission rate based on the physical power of the turbines is employed including emissions due to shutting down and starting of the gas turbines, leakages at regulating valves and fittings, maintenance and gasometry work.

The calculation of losses from source category 1B2b v Distribution follows a detailed country-specific approach that considers losses from the pipeline network as well as losses at the end users.

The calculated gas losses from the pipeline network depend on the length, material and pressure of the pipelines. Gas losses due to permanent leakiness, small-scale damages, network maintenance and the network components are evaluated separately. As no applicable loss rates are available for the network compounds in Switzerland (installed control units, fittings, storage systems and gas meters), a fixed percentage is applied to the permanent gas losses.

Regarding the end users, gas losses from on-site and indoor pipelines as well as gas losses due to the permanent leakiness of gas appliances are evaluated. Pipeline loss rates apply to the number of households, industrial users and gas fuelling stations separately. Regarding the gas appliances, different loss rates are assigned to the number of gas heating systems, gas cooking stoves and gas fuelling stations.

For some (earlier) years in the time series, sufficient input data are not available to calculate the gas losses. For these years, polynomial interpolations are applied to assess the activity data.

For significant emission events due to accidents the Swiss Pollutant Release and Transfer Register is considered, and emissions are attributed to source category 1B2b vi Other Leakage. So far, two events have been reported by the transit pipeline operator, one in 2010 and one in 2011.

Fugitive emissions from pipelines are the major emission source in source category 1B2b. Fugitive emissions from damages and ruptures of the pipelines, maintenance of the pipelines and the components are very small (Quantis 2014). Total CH₄ emissions from gas transmission and distribution decreased due to gradual replacement of cast-iron pipes with polyethylene pipes.

Table 3-104 Activity data (amount of gas lost) for fugitive emissions from 1B2b Natural gas

1B2b Natural Gas	Unit	1990	1995	2000	2005
1B2b Natural gas	GJ	866'811	846'239	686'190	543'896
1B2b ii Production	GJ	130'000	NO	NO	NO
1B2b iv Transmission	GJ	26'565	29'211	30'923	31'860
1B2b v Distribution	GJ	710'246	817'028	655'267	512'036
1B2b vi Other Leakag	GJ	NO	NO	NO	NO

1B2b Natural Gas	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
1B2b Natural Gas	GJ	523'734	519'852	504'640	492'119	482'349	474'762	466'414	434'622	421'123	421'123
1B2b ii Production	GJ	NO									
1B2b iv Transmission	GJ	31'949	32'786	32'906	32'921	32'930	32'904	32'681	33'309	33'338	33'338
1B2b v Distribution	GJ	491'785	487'066	471'734	459'198	449'419	441'858	433'733	401'313	387'785	387'785
1B2b vi Other Leakag	GJ	NO	NO	NO	NO	35'444	28'114	NO	NO	NO	NO

3.3.4.3 Uncertainties and time-series consistency for 1B2b

According to the assessment by Quantis (2014), an uncertainty of 30% is estimated for fugitive CH₄ emissions from natural gas pipelines in Switzerland.

A preliminary uncertainty assessment of all other sources in source category 1B2 based on expert judgement results in medium confidence in the emissions estimate (see Table 1-10).

Consistency: Time series for 1B2b Natural gas are all considered consistent.

3.3.4.4 Category-specific QA/QC and verification for 1B2b

The general QA/QC measures are described in chp. 1.2.3.

As suggested by the 2006 IPCC Guidelines (IPCC 2006) the gas industry was involved in the reassessment of fugitive emissions from the natural gas system in 2014 (Quantis 2014) and 2016 (EMIS 2017/1B2b Diffuse Emissionen Erdgas).

3.3.4.5 Category-specific recalculations for 1B2b

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 or 2014 of at least 1 kt CO₂ eq are quantified.

- 1B2b ii: Recalculation in the Swiss overall energy statistics (SFOE 2016) concerning production of natural gas between 1985 and 1994.
- 1B2b ii: Reassessment of the emission factors for source category 1B2b ii.
- 1B2b iv and 1B2b v: Recalculations were carried out due to the update of the employed calculation tool with minor corrections of individual natural gas loss rates, minor corrections of individual data regarding the Swiss gas network as well as minor changes in the polynomial interpolations for years with no sufficient data of the gas network available (EMIS 2017/1B2b Diffuse Emissionen Erdgas). This recalculation leads to an increase of 15 kt CO₂ eq in 1990.

3.3.4.6 Category-specific planned improvements for 1B2b

No category-specific improvements are planned.

3.3.5 Source category 1B2c – Venting and flaring

3.3.5.1 Source category description for 1B2c

In Switzerland, oil production is not occurring, and only one production site for natural gas production was operational from 1985 – 1994. Therefore, emissions from flaring result primarily from the torches, which were operational at the two refineries (1B2c i Flaring). Since 2015, there is only one refinery in operation. In addition, CO₂ emissions from H₂ production in one of the two refineries are also reported under 1B2c.

Table 3-105 Specification of source category 1B2c Venting and flaring in Switzerland.

1B2	Source	Specification
1B2 c	Fugitive emissions attributed to venting and flaring.	The combustion of excess gas at the oil refinery (flaring) only. Emissions from H ₂ production Emissions from gas production (1990-1994 only)

3.3.5.2 Methodological issues for 1B2c

Methodology (1B2c)

According to the decision tree for crude oil transport, refining and upgrading, Switzerland follows a Tier 2 method for emissions attributed to 1B2c i Flaring, Oil in order to estimate fugitive emissions under 1B2c fugitive emissions from venting and flaring (IPCC 2006, Volume 2 Energy, chp. 4 Fugitive Emissions, Figure 4.2.3). For emission calculations plant-specific emission factors and activity data are available from the refining industry.

For source category 1B2c i Flaring, Oil, emissions of CO₂ as well as CH₄, N₂O, NO_x, CO and NMVOC are considered. For source category 1B2c ii Flaring, Gas emissions of CO₂ as well as CH₄, N₂O and NMVOC are considered. Since the CO₂ emission factors assume an oxidation of 100%, no indirect emissions need to be accounted for. Therefore, from this source category no indirect emissions are reported in chp. 9.

One of the refining plants produces H₂ from butane leading to process emissions of CO₂. Emissions are estimated based on plant-specific data.

Emission factors (1B2c)

Emission factors are based on data from the refining industry as documented in the internal emission database documentation (EMIS 2017/1B2c Raffinerie Abfackelung). Since 2005 (with the exception of 2012), the refining industry provides annual data on the emissions from flaring under the Federal Act on the Reduction of CO₂ Emissions (Swiss Confederation 2011) based on daily measurements of CO₂ emission factors of the flared gases. From these data annual emission factors are derived. Since 2005, the evolution of the other emission factors (CH₄, N₂O, NO_x, CO and NMVOC) is assumed to vary proportionally to the CO₂ emission factor. Emission factors are considered confidential and are available to reviewers on request.

The CO₂ emission factor of H₂ production from butane, which is produced in one of the refineries since the end of 2004, is equal to the stoichiometric emission factor.

The emissions from flaring in the gas production facility are calculated based on default emission factors provided in the 2006 IPCC guidelines.

Table 3-106 Emission factors for 1B2c Venting and flaring.

Source/fuel	Unit	CO ₂	CH ₄	N ₂ O	NO _x	NM VOC	SO ₂	CO
1B2c Venting and flaring	g/t	C	C	C	C	C	C	C
1B2cii Flaring from gas production	g/GJ	33	0.0021	0.00058	NA	0.017	NA	NA
Other: H2 production refinery (butane)	g/GJ	C	NA	NA	NA	NA	NA	NA

Activity data (1B2c)

Before 2005, the amount of flared gas is assumed to be proportional to the amount of crude oil processed in the refineries. The Swiss petroleum association provides data on the use of crude oil on an annual basis (EV 2016). Between 2001 and 2004, one of the two refineries made major changes to their installations (new cracker, new flaring installation) and their standard operation process. Therefore, emissions from flaring decreased significantly thereafter. Since 2005, the industry provides data on the amount of gas flared.

For gas production, the amount flared is estimated based on the amount of gas produced.

For H₂ production in one of the refining plants, annual data on butane consumption are provided by the industry since 2005, when the H₂ production unit was installed. Data are confidential and they are available to reviewers on request.

Table 3-107 Activity data for 1B2c Venting/flaring.

1B2c Venting and flaring	Unit	1990	1995	2000	2005
Crude oil used	kt	3'127	4'657	4'649	4'877
Flaring from gas production	GJ	130'000	NO	NO	NO
Other: H2 production refinery (butane)	GJ	NO	NO	NO	C

1B2c Venting and flaring	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Crude oil used	kt	5'563	4'720	5'133	4'833	4'546	4'452	3'455	4'935	4'975	2'836
Flaring from gas production	GJ	NO									
Other: H2 production refinery (butane)	GJ	C	C	C	C	C	C	C	C	C	C

3.3.5.3 Uncertainties and time-series consistency for 1B2c

A preliminary uncertainty assessment of all sources in source category 1B2 based on expert judgement results in medium confidence in the emissions estimate (see Table 1-10).

Consistency: Time series for 1B2c Venting and flaring are all considered consistent.

3.3.5.4 Category-specific QA/QC and verification for 1B2c

The general QA/QC measures are described in chp. 1.2.3. No category-specific QA/QC activities were undertaken.

3.3.5.5 Category-specific recalculations for 1B2c

The following recalculations were implemented in submission 2017. Recalculations cause a change in emission levels in 1990 or 2014 by less than 1 kt CO₂.

- 1B2c: CH₄ emission factor of flaring activity in refineries changed due to less rounding (2005-2014).
- 1B2c: Emissions of N₂O , CO₂ and CH₄ from flaring in gas production were missing before. Now emission factors from the 2006 IPCC Guidelines are used (1990-1994).

3.3.5.6 Category-specific planned improvements for 1B2c

Emissions from flaring in gas production (1990-2004) were assessed for the first time in Submission 2017. They amount to 4t CO₂ eq in 1990. The data are not yet implemented in the CRF tables. The emissions will be integrated in Submission 2018.

3.4 Source category 1C – CO₂ transport and storage

CO₂ transport and CO₂ storage is not occurring in Switzerland.

4 Industrial processes and product use

4.1 Overview

This chapter provides information on the estimation of the GHG emissions from sector 2 Industrial processes and product use. The following source categories are reported:

- 2A Mineral industry
- 2B Chemical industry
- 2C Metal industry
- 2D Non-energy products from fuels and solvent use
- 2E Electronics industry
- 2F Product uses as substitutes for ozone-depleting substances (ODS)
- 2G Other product manufacture and use
- 2H Other

Emissions within this sector comprise GHG emissions as by-products from industrial processes and also emissions of F-gases during production, use and disposal. Emissions from fuel combustion in industry are reported in source category 1A2 under sector 1 Energy.

According to the 2006 IPCC guidelines this sector provides also information on the GHG emissions from solvent and product use. CO₂ emissions from solvent and partly from product use are due to post-combustion of NMVOC in order to reduce NMVOC in exhaust gases. The disposal of solvents is reported in the waste sector (chp. 7).

Indirect emissions CO₂ and N₂O from fossil CO and NMVOC as well as NO_x and NH₃ emissions, respectively, are reported in chapter 9. Since the CO₂ emissions from 2C1 Secondary steel production, electric arc furnace and 2C3 Primary aluminium production are based on carbon mass balances their emissions of CO and NMVOC, and of CO, respectively, are not accounted for calculation of the indirect CO₂ emissions. Biogenic NMVOC and CO emissions occur in source category 2H2 Food and beverages and 2G4 tobacco consumption and are not reported as indirect CO₂ emissions.

For several industrial processes within source categories 2A Mineral industry, 2B Chemical industry and 2C Metal industry data and information on emission factors and activity data are classified as confidential (C). For reviewers there is an additional version of chapter 4 Industrial processes and product use available, including all confidential data and information.

Figure 4-1 shows the evolution of GHG emissions in sector 2 between 1990 and 2015.




Figure 4-1 Switzerland's greenhouse gas emissions of sector 2 Industrial processes and product use 1990–2015.

2A Mineral industry remains the dominant source of sector 2 accounting for around half of the GHG emissions in 2015 although absolute emissions have decreased since 1990. 2B Chemical industry accounts for a small share and shows a decreasing trend since 1990. 2C Metal industry shows a strong decreasing trend and accounts only for a small share in 2015. 2D Non-energy products also only have a minor contribution in 2015.

2F Product uses as substitutes for ozone depleting substances (ODS) is of increasing importance: The emissions have increased since 1990 and account for almost half of total GHG emissions in sector 2 in 2015. This is primarily due to the replacement of CFCs and other ODS by HFCs in many technical applications. 2G Other product manufacture and use shows no clear trend since 1990. 2E Electronic industry and 2H Other are of little importance with regard to the overall GHG emissions of sector 2.

In Table 4-1, the development of GHG emissions in sector 2 Industrial processes and product use is given by gases. Dominant gases are CO₂ and F-gases in 2015 whereas N₂O and CH₄ have only a minor contribution. The relative trend of these gases referring to the base year 1990 is shown in Figure 4-2 and Figure 4-3.

Table 4-1 GHG emissions of sector 2 Industrial processes and product use by gases in kt CO₂ equivalent.

Gas	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
CO ₂	3'158	2'424	2'206	2'381
CH ₄	1.8	1.8	1.7	2.6
N ₂ O	171	140	115	100
F-gases	254	356	816	1'312
Sum	3'585	2'922	3'139	3'795

Gas	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	CO ₂ equivalent (kt)									
CO ₂	2'331	2'337	2'297	2'215	2'372	2'327	2'203	2'205	2'259	2'091
CH ₄	2.3	2.6	2.9	1.9	2.7	2.8	2.8	2.1	2.1	2.1
N ₂ O	104	101	109	99	102	94	94	70	48	49
F-gases	1'348	1'407	1'516	1'495	1'545	1'640	1'766	1'818	1'830	1'849
Sum	3'785	3'848	3'924	3'811	4'022	4'064	4'066	4'096	4'140	3'992




Figure 4-2 Relative trends of the GHG (without F-gases, see Figure 4-3) of sector 2 Industrial processes and product use in the period 1990–2015. The base year 1990 represents 100%.

Figure 4-2 shows that since 1990 the emissions of CO₂ and N₂O from sector 2 Industrial processes and product use have both decreased. Emissions of CH₄ have increased in the same time period with considerable interannual fluctuation. However, absolute emissions are small compared to CO₂ and N₂O.

Figure 4-3 shows a large increase in emissions of F-gases compared to the year 1990. Main contributions in the inventory 1990 result from the PFC emissions in the smelting process of aluminium production (chp. 4.4.2.2) and from the use of SF₆ in electrical equipment and sound proof windows (chp. 4.8.2.1 and chp. 4.8.2.2). The increase from 1995 onwards is due

to the increasing product uses of HFCs as substitutes for ODS (chp. 4.7) in refrigeration and air conditioning. Most relevant and main source of F-gases emissions in 2015 is the use of HFC in refrigeration and air conditioning.




Figure 4-3 Relative trends of the F-gases of sector 2 Industrial processes and product use in the period 1990–2015. The base year 1990 represents 100%.

4.2 Source category 2A – Mineral industry

4.2.1 Source category description

Table 4-2 Key categories (KCA incl. LULUCF) of 2A Mineral industries.

Code	IPCC Category	GHG	Identification Criteria
2A1	Cement production	CO ₂	L1, L2, T1

Table 4-3 Specification of source category 2A Mineral industry in Switzerland.

2A	Source	Specification
2A1	Cement production	Geogenic CO ₂ emissions from calcination process in cement production; Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting operations
2A2	Lime production	Geogenic CO ₂ emissions from calcination process in lime production; Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting operations
2A3	Glass production	Geogenic CO ₂ emissions from production of container and tableware glass, and glass wool
2A4	Other process uses of carbonates	Geogenic CO ₂ emissions from fine ceramics, brick and tile and rock wool production as well as from use of sodium bicarbonate; Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting operations in plaster production Geogenic CO ₂ emissions from carbonate use for fluegas purification in cellulose production and waste incineration plants

4.2.2 Methodological issues

4.2.2.1 Cement production (2A1)

In Switzerland, there are six plants producing clinker and cement. The Swiss plants are rather small and do not exceed a capacity of 3'000 tonnes of clinker per day. All of them use modern dry process technology.

Emissions of geogenic CO₂ occur during the production of clinker, which is an intermediate component in the cement manufacturing process. During the production of clinker, limestone, which is mainly calcium carbonate (CaCO₃), is heated (calcined) to produce lime (CaO) and CO₂ as by-product. The CaO reacts subsequently with minerals in the raw materials and yields clinker. During this reaction step no further CO₂ is emitted. Clinker is then mixed with other components such as gypsum to make cement.

Blasting operations in the limestone quarries are another source of emissions for both CO₂ and precursor greenhouse gases such as NO_x, CO, NMVOC and SO₂.

Methodology

Calcination process

The geogenic CO₂ emissions from the calcination process in cement production are determined by a Tier 2 method according to the decision tree Fig. 2.1. of 2006 IPCC Guidelines (vol. 3, chp. 2.1 Cement production).

In Switzerland, no long wet or long dry kilns are used. Only modern preheater or precalciner kilns are used and also no so-called low-alkali cement is produced. Therefore, there is no land-filling of calcined cement dust (cement kiln dust, CKD) in Switzerland. In the cement

plants all the filter dust is collected in high performance electrostatic precipitator or bag filters (having an efficiency of more than 99.999%) and being recycled to the kiln feed. In some cases small portions of the CKD are added directly to the cement as filler. Due to the kiln technology used in Switzerland the degree of decarbonization of the CKD is almost equal to that of the kiln feed, meaning, that this CKD has not been decarbonised yet.

Blasting operations

Emissions resulting from blasting operations during the digging of limestone are calculated by a Tier 2 method according to EMEP/EEA Guidebook 2013 (EMEP/EEA 2013, chp. 2A1, Fig. 3.1) using country-specific emission factors. The CO₂ emissions from "blasting" are related only to the usage of explosives in the quarries and not to the fuel consumption of construction machinery such as bulldozers etc. The amount of used explosives is reported to be 0.13 kg/t cement¹¹ (EMIS 2017/2A1 Zementwerke übriger Betrieb).

Total emissions reported for the production of cement are the sum of emissions from calcination process and blasting operations. The share of CO₂ emissions from blasting operations in limestone quarries is well below one tenth of a per cent of the geogenic CO₂ emissions from the calcination process.

Emission factors

Calcination process

The emission factor of CO₂ from calcination is provided per tonne of clinker. It accounts for geogenic emissions from the carbonate containing raw material, emissions from organic carbon content of the raw material and from cement kiln dust (CKD).

For emissions from calcination of the carbonate containing raw material a value of 525 kg CO₂/t clinker is used according to the guidance document of the Swiss ETS (FOEN 2013q, chp. G.7). The emissions from the organic carbon content of the raw material are assumed to be a constant share of 0.2% of the raw material (i.e. 11.37 kg CO₂/t clinker). The emission factor of CKD is estimated based on plant-specific data available for 2013 – 2015 that are provided by the cement industry association (cemsuisse). From this data, an average emission factor of CO₂ from CKD is calculated (0.35 kg CO₂/t clinker).

Based on these emission factors a total emission factor per ton of clinker is calculated. The emission factor is assumed constant for the entire time period.

¹¹ The CO₂ emission factor for the use of blasting agents amounts to 600 kg CO₂/t of blasting agent. For the average amount on blasting agent used per kg cement measurement data for the year 2002 were taken. Measurement data were available for four Swiss cement plants, covering more than 60% of the Swiss cement production. Therefore, this information is considered representative for cement plants in Switzerland.

Table 4-4 CO₂ emission factor for calcination in 2A1 Cement Production 1990 to 2015.

2A1 Cement production	Unit	1990 - 2015
Calcination, CO ₂	kg/t clinker	536.7

Blasting operations

The emission factors are country-specific based on emission factors of civil explosives and information on the specific consumption of explosives in the quarries as documented in the Handbook on emission factors for stationary sources (SAEFL 2000) as documented in the EMIS database (EMIS 2017/2A1 Zementwerke übriger Betrieb). They are assumed to be constant over the entire time period and are given per tonne of clinker.

Table 4-5 Emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting operations in g/t clinker from source category 2A1 Cement Production in 2015

2A1 Cement production	Unit	CO ₂	NO _x	CO	NMVOC	SO ₂
Blasting operations	g/t clinker	34.1	3.3	3.3	8.6	0.1

Activity data

Since 1990, data on annual clinker production are provided by the industry association cemsuisse as documented in the EMIS database (EMIS 2017/2A1_Zementwerke Rohmaterial). From 2008 onwards they are based on plant-specific annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

Table 4-6 Activity data of clinker production

2A1 Cement production	Unit	1990	1995	2000	2005
Clinker production	kt	4'808	3'706	3'214	3'442

2A1 Cement production	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Clinker production	kt	3'452	3'512	3'461	3'443	3'642	3'587	3'368	3'415	3'502	3'195

4.2.2.2 Lime production (2A2)

During the production of lime calcium carbonate (CaCO₃) is heated (calcined) yielding burnt lime (CaO) and CO₂ as by-product. In Switzerland, there is only one plant producing lime. There is no industry in Switzerland producing lime for its own requirements, except for sugar production. A request to both sugar producing plants confirmed that indeed they produce lime from limestone in own shaft kilns. However, the CO₂ is re-captured in the sugar production process and thus no CO₂ emissions occur.

Blasting operations in quarries are another source of emissions for both CO₂ and precursor emissions such as NO_x, CO, NMVOC and SO₂.

Methodology

Calcination process

The geogenic CO₂ emissions from the calcination process in lime production are determined by a country-specific method according to the decision tree Fig. 2.2. of 2006 IPCC guidelines (vol. 3, chp. 2.2 Lime production).

Blasting operations

Emissions resulting from blasting operations during the digging of limestone are calculated by a Tier 2 method according to EMEP/EEA Guidebook 2013 (EMEP/EEA 2013, chp. 2A2, Fig. 3.1) using country-specific emission factors. The CO₂ emissions from "blasting" are related only to the usage of explosives in the quarries and not to fuel consumption of e.g. bulldozers etc.

Total emissions reported for the production of lime are the sum of emissions from calcination process and blasting operations. CO₂ emissions from blasting operations in limestone quarries account only for a small share of the total emissions.

Emission factors

Calcination process

The emission factor for CO₂ from calcination of limestone depends both on the purity of the limestone and the degree of calcination (i.e. amount of rest CO₂ remaining in the lime produced). A plant-specific value has been calculated based on industry declaration and it is assumed to be constant for the years 1990–2012 (EMIS 2017/2A2 Kalkproduktion, Rohmaterial). The value is confidential and is available to reviewers on request. Since 2013, emission factors are derived from annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

Table 4-7 CO₂ emission factor for calcination process in lime production in kg/t lime for 1990–2015 are documented in the confidential NIR, which is available to reviewers on request.

Blasting operations

The emission factors are country-specific as documented in EMIS 2017/2A2 Kalkproduktion, übriger Betrieb. The values are confidential and they are available to reviewers on request.

Table 4-8 CO₂ emission factor for the calcination process in lime production in kg/t lime and emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting operations in g/t lime in 2015

2A2 Lime production	Unit	CO ₂	NO _x	CO	NMVOC	SO ₂
Calcination	kg/t	C	NA	NA	NA	NA
Blasting operations	g/t	C	C	C	C	C

Activity data

Activity data on annual lime production are provided by the only existing plant in Switzerland, as documented in the EMIS database (EMIS 2017/2A2 Kalkproduktion, Rohmaterial and EMIS 2017/2A2 Kalkproduktion übriger Betrieb). Since 2009 they are based on plant-specific annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

Detailed activity data are not reported since they are considered confidential.

Table 4-9 In the confidential NIR, the respective table with activity data on lime production are separately reported and available to reviewers.

4.2.2.3 Glass production (2A3)

Source category 2A3 Glass production comprises geogenic CO₂ emissions from the carbonate containing raw materials, i.e. soda ash, limestone and dolomite. In Switzerland, the following three glass types are produced: container glass, tableware glass and glass wool. Today, there is only one production plant remaining for container glass and tableware glass after the other plants closed in 2002 and 2006, respectively. Glass wool is produced in two plants.

Methodology

For determination of geogenic CO₂ emissions from glass production, a Tier 2 method according to the decision tree Fig. 2.3 of 2006 IPCC guidelines (vol. 3, chp. 2.4 Glass production) is used. For glass production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = M_{\text{Glass type}} \cdot \text{EF}_{\text{Glass type}} \cdot (1 - \text{cullet ratio})$$

The cullet ratio describes the share of recycled glass material which is used in the production. The melting of cullet causes no geogenic CO₂ emissions.

From 2005 onwards, the geogenic CO₂ emissions from 2A3 Container glass production is determined according to a Tier 3 method based on the amount of carbonate containing raw materials used, i.e. soda, dolomite and limestone and their effective carbonate content.

Emission factors

The emission factors for glass production in Switzerland are taken from IPCC 2006 (vol.3, chp. 2.4 Glass production, Table 2.6). For the production of container glass (1990-2004), tableware glass (1990-2015) and glass wool (1990-2015) the values for glass type container, tableware and fibreglass are taken, respectively. As the emission factors are material properties, they remain constant over time.

From 2005 onwards, effective amounts of carbonate containing raw materials (soda ash, dolomite and limestone) are available from ETS monitoring reports for the container glass

production and thus the corresponding CO₂ emission factors are taken from IPCC 2006 (vol. 3, chp. 2.1, Table 2.1). As these emission factors are material properties, they remain constant over time.

Table 4-10 Geogenic CO₂ emission factor for glass production in g/t glass and g/t carbonate containing raw material (IPCC 2006).

2A3 Glass production	Unit	CO₂ geogenic	
Glass wool (fibre glass insulation)	g/t	250'000	
Glass (speciality tableware)	g/t	100'000	
		1990–2004	2005–2015
Container glass	g/t	210'000	
Soda use	g/t soda		414'920
Dolomite use	g/t dolomite		477'320
Limestone use	g/t limestone		439'710

Table 4-11 In the confidential NIR, a comparison of implied CO₂ emission factors based on Tier 2 and Tier 3 approaches is provided for container glass production in g/t glass for the time period 2005-2011.

Activity data and cullet ratios

Source category 2A3 Glass production is dominated by the emissions from the production of container glass and glass wool.

For glass wool production, activity data are based on data from the two glass wool production plants in Switzerland. Since 2008, activity data are based on plant-specific annual monitoring reports.

Activity data of tableware and container glass production are based on data from Swiss glass producers.

Detailed information on activity data for container glass production and tableware production is confidential as there is only one production plant for container glass and tableware glass, respectively. Data are available to the reviewers on request (EMIS 2017/2A3 Hohlglass Produktion, EMIS 2017/2A3 Glas übrige Produktion and EMIS 2017/2A3 Glaswolle Produktion Rohprodukt).

Table 4-12 Activity data of glass production in Switzerland and cullet ratio in % as well as consumption of carbonate containing raw materials in container glass production

2A3 Glass production	Unit	1990	1995	2000	2005
Container glass					
Production	kt	C	C	C	C
Cullet ratio	%	C	C	C	NA
Soda use	kt	NA	NA	NA	C
Dolomite use	kt	NA	NA	NA	C
Limestone use	kt	NA	NA	NA	C
Glass (speciality tableware)					
Production	kt	C	C	C	C
Cullet ratio	%	C	C	C	C
Glass wool					
Production	kt	24.3	24.2	31.1	37.5
Cullet ratio	%	21	45	69	65

2A3 Glass production	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Container glass											
Production	kt	C	C	C	C	C	C	C	C	C	C
Cullet ratio	%	NA									
Soda use	kt	C	C	C	C	C	C	C	C	C	C
Dolomite use	kt	C	C	C	C	C	C	C	C	C	C
Limestone use	kt	C	C	C	C	C	C	C	C	C	C
Glass (speciality tableware)											
Production	kt	C	C	C	C	C	C	C	C	C	C
Cullet ratio	%	C	C	C	C	C	C	C	C	C	C
Glass wool											
Production	kt	38.1	44.5	44.4	33.5	35.7	41.4	38.7	33.4	32.3	31.4
Cullet ratio	%	73	71	69	69	71	72	61	67	67	67

4.2.2.4 Other process uses of carbonates (2A4)

Source category 2A4 Other process uses of carbonates comprises geogenic CO₂ emissions from production of fine ceramics (2A4a), bricks and tiles (2A4a) and rockwool (2A4d), from use of carbonates for sulphur oxide removal in municipal solid waste incineration plants (2A4d) and cellulose production (ceased in 2008) (2A4d), from use of limestone in iron-foundries (cupola furnaces) (2A4d) and from use of sodium bicarbonate (2A4d) as well as emissions of CO₂, NO_x, CO, NMVOC and SO₂ from blasting operations in plaster production (2A4d).

Ceramics (2A4a)

Source category 2A4a Ceramics consists of the production of fine ceramics and brick and tile.

Fine ceramics (2A4a)

In Switzerland, the main production of fine ceramics is sanitary ware. The carbonate containing raw materials limestone and dolomite as well as small amounts of soda ash are used in product glazes only. All information on the fine ceramics production is documented in EMIS 2017/2A4a Feinkeramik Produktion.

Methodology

The geogenic CO₂ emissions from fine ceramics production are determined by a Tier 2 method according to the decision tree Fig. 2.4 of 2006 IPCC guidelines (IPCC 2006, vol. 3, chp. 2.5 Other process uses of carbonates).

For fine ceramics production in Switzerland, this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = (M_{\text{Limestone}} \cdot \text{EF}_{\text{Limestone}}) + (M_{\text{Dolomite}} \cdot \text{EF}_{\text{Dolomite}}) + (M_{\text{Soda Ash}} \cdot \text{EF}_{\text{Soda Ash}})$$

Emission factors

The CO₂ emission factors of limestone, dolomite and soda ash are taken from IPCC 2006 (vol. 3, chp. 2.1, Table 2.1). As these emission factors are material properties, they remain constant over time.

Table 4-13 Geogenic CO₂ emission factors used for fine ceramics and the production of brick and tile in g/t carbonate containing raw material and g/t product, respectively.

2A4a Ceramics	Unit	CO₂ geogenic			
Fine ceramics		1990–2015			
Limestone use	g/t limestone	439'710			
Dolomite use	g/t dolomite	477'320			
Soda use	g/t soda	414'920			
		1990–2012	2013	2014	2015
Brick and tile production	g/t	117'000	100'000	110'000	103'000

Activity data

Activity data for carbonate containing raw materials (i.e. limestone, dolomite and soda ash) used in the glazes of the fine ceramics production are extrapolated values based on industry data from the largest production plant in Switzerland. Detailed activity data are considered confidential. They are available to the reviewers on request.

Brick and tile production (2A4a)

In Switzerland, there are about 20 plants producing bricks and tiles. The manufacturing process uses limestone containing clay as main raw material.

Methodology

The brickearth used in Switzerland for the production of bricks and tiles does not consist of pure and defined contents of clay minerals but its clay content is varying depending on the individual pit, comprising other minerals such as calcite, dolomite and quartz. Compared to other countries, the fraction of carbonate containing raw material is relatively high. Detailed data on the composition of carbonate containing raw materials from the Swiss brick and tile industry were not available before 2013. Therefore, for the period 1990 until 2012 data from a comparison of geogenic CO₂ emissions based on representative analyses of the carbonate content of the clay used for brick and tile production in a number of plants in Switzerland and the European Union are applied. This study was carried out by the Swiss association of brick and tile industry (Verband Schweizerische Ziegelindustrie, VSZ) in 2012 (see EMIS 2017/2A4a Ziegeleien).

Since 2013, the Swiss brick and tile production plants are legally obliged to report geogenic emissions from carbonate containing raw materials annually (Federal Act on the Reduction of CO₂ Emissions, Swiss Confederation 2011 and Ordinance for the Reduction of CO₂ Emissions, Swiss Confederation 2012). The emissions are estimated from analyses of the carbonate content of the raw materials and an assumed calcination factor of 100%. This procedure corresponds to a Tier 3 method according to the decision tree Fig. 2.4 of 2006 IPCC guidelines (IPCC 2006, vol. 3, chp. 2.5 Other process uses of carbonates). Between 1990 and 2012 a Tier 2 method is applied.

Emission factors

According to the above mentioned study, bricks emit a weighted average of 13.2% of geogenic CO₂ (variation range 5.4% - 24%) and roof tiles have a weighted average of 8.6% (variation range 5.6% - 13%). Based on the production shares of the largest Swiss brick producer, a production ratio for bricks to tiles of 2:1 was assumed for the whole period from 1990 to 2012. This resulted in an average geogenic CO₂ emission factor of 117 kg CO₂/t brick and tile, which was assumed constant for the time period 1990 to 2012.

Since 2013, a production weighted emission factor is derived based on the plant-specific monitoring data of the geogenic CO₂ emissions from the carbonate containing raw materials. For emission factors see Table 4-13.

Activity data

Activity data are based on production data from the Swiss association of brick and tile industry (VSZ). Since 2011 they are based on plant-specific annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

Table 4-14 Activity data for the production of fine ceramics including the use of limestone, soda and dolomite in the glazes, brick and tile, rock wool and plaster as well as other use of carbonates (sodium bicarbonate) in Switzerland in kt.

2A4a Ceramics	Unit	1990	1995	2000	2005
Fine ceramics production	kt	C	C	C	C
Limestone use	kt	C	C	C	C
Dolomite use	kt	C	C	C	C
Soda use	kt	C	C	C	C
Brick and tile production	kt	1'271	1'115	959	1'086
2A4d Other					
Rock wool production	kt	C	C	C	C
Carbonate use in waste incineration plants	kt	0.7	0.8	0.8	0.6
Limestone use in cellulose	kt	8.5	9.4	9.3	8.3
Limestone use in iron foundries	kt	8.6	5.8	5.4	2.7
Other use of carbonates	kt	5.9	5.4	7.0	7.3
Plaster production	kt	319	304	288	327

2A4a Ceramics	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Fine ceramics production	kt	C	C	C	C	C	C	C	C	C	C
Limestone use	kt	C	C	C	C	C	C	C	C	C	C
Dolomite use	kt	C	C	C	C	C	C	C	C	C	C
Soda use	kt	C	C	C	C	C	C	C	C	C	C
Brick and tile production	kt	1'065	975	865	701	879	800	792	785	765	726
2A4d Other											
Rock wool production	kt	C	C	C	C	C	C	C	C	C	C
Carbonate use in waste incineration plants	kt	0.1	NO	NO	NO	NO	NO	NO	2.7	2.7	2.7
Limestone use in cellulose	kt	8.0	7.8	6.5	NO						
Limestone use in iron foundries	kt	2.6	2.8	3.1	1.3	1.0	1.2	0.9	0.9	0.8	0.7
Other use of carbonates	kt	7.0	7.0	7.4	6.6	6.9	6.4	7.6	6.1	6.7	10.5
Plaster production	kt	323	314	295	293	335	293	271	213	166	140

Other uses of soda ash (2A4b)

Soda ash is mainly used in the glass production, which is reported separately in source category 2A3 Glass production. A very small amount of soda ash is also applied in glazes of fine ceramics and is thus accounted for in source category 2A4a Ceramics (see Table 4-13).

Other (2A4d)

Rock wool production (2A4d)

In Switzerland, there is one single producer of rock wool. The plant uses carbonate containing raw materials like dolomite, basalt, cement and further additives as documented in the EMIS database (EMIS 2017/2A4d Steinwolle Produktion).

Methodology

Since 2013, rock wool manufacturers are legally obliged to report geogenic CO₂ emissions from carbonate containing raw material annually. For the years 2005-2011 and 2013 plant-specific data on raw material consumption and emission factors is available from monitoring reports of the Swiss ETS. From this information, data for the other years are interpolated for calculating an implied emission factor.

The geogenic CO₂ emissions from rock wool production are determined by a Tier 3 method according to IPCC 2006 (vol. 3, chp. 2.5 Other process uses of carbonates).

Emission factors

For rock wool production in Switzerland, the CO₂ emission factor is based on measurements of the oxides (CaO, MgO, Na₂O, K₂O, MnO) of the carbonate containing raw materials and the product for the years 2005 to 2011 as well as since 2013. Based on the difference in the oxide content in the raw material and the products, the total geogenic CO₂ emissions are determined. Consequently, the emission factor is specified as g/t rock wool. Since data on the carbonate content are missing for the years 1990 to 2004 and 2012 the mean value of the years 2005-2011 and 2013 is applied for these years.

The CO₂ emission factors are confidential. They are available to reviewers on request.

Table 4-15 Geogenic CO₂ emission factors used for rock wool production and other carbonate uses, CO₂ fossil, NO_x, CO, NMVOC and SO₂ emission factors for plaster production in g/t carbonate containing raw material and g/t product, respectively for 2015.

2A4d Other	Unit	CO ₂ geogenic	CO ₂ fossil	NO _x	CO	NMVOC	SO ₂
Rock wool production	g/t	C	NA	NA	NA	NA	NA
Carbonate use in waste incineration plants	g/t	523'880	NA	NA	NA	NA	NA
Limestone use in iron foundries	g/t	439'710	NA	NA	NA	NA	NA
Other carbonate uses	g/t	523'880	NA	NA	NA	NA	NA
Plaster production	g/t rocks	NA	144	5.6	33	14.4	0.24

Table 4-16 In the confidential NIR, the respective table with geogenic CO₂ emission factors used for rock wool production is separately reported and available to reviewers.

Activity data

Activity data are based on industry data from the single rock wool production plant in Switzerland (monitoring reports of the Swiss ETS) and are therefore confidential. They are available to reviewers on request.

Other carbonate uses (2A4d)

Methodology

In 2014, an assessment was carried out in order to identify sources of CO₂ emissions from carbonate use for sulphur oxide removal and acid neutralization, which were not considered in the Swiss greenhouse gas inventory so far (INFRAS 2015). The survey among selected potentially relevant industrial plants, industry associations, Swiss cantons and the Swiss customs administration (EZV) comprised the following substances: limestone (CaCO₃), dolomite (CaMg(CO₃)₂), sodium bicarbonate (NaHCO₃) and soda ash (Na₂CO₃).

Besides applications of calcium hydroxide and sodium hydroxide in flue gas treatment also a few applications of limestone and sodium bicarbonate for sulphur oxide removal could be

identified in Switzerland. Limestone had been used in the cellulose production up to 2008, when the plant was closed, and in one municipal solid waste incineration plant up to 2005. Another waste incineration plant has used sodium bicarbonate since 2013.

In cupola furnaces of iron foundries limestone is also used as flux as documented in the EMIS database (EMIS 2017/1A2a Eisengiessereien Kupolöfen).

Limestone is also used to neutralize acid waste water in one chemical production plant. These emissions are reported in source category 2B10 Limestone pit.

Additionally, it is assumed, that all applications of sodium bicarbonate result in a complete conversion to CO₂. Since there is no production of sodium bicarbonate in Switzerland, the annual emissions can be estimated based on the net import.

The method for calculating the geogenic CO₂ emissions from the use of limestone and sodium bicarbonate in all these source categories corresponds to a Tier 2 method according to the decision tree Fig. 2.4 of 2006 IPCC guidelines (IPCC 2006, vol. 3, chp. 2.5 Other process uses of carbonates).

Emission factors

The emission factors of limestone and sodium bicarbonate are based on the stoichiometry of CaCO₃ (2006 IPCC Guidelines, vol. 3 chp. 2.1, table 2.1, IPCC 2006) and NaHCO₃ (CRC 2004), respectively, see Table 4-15. A conversion factor of 100% is assumed for all applications of both carbonates.

Activity data

Activity data on limestone use in flue gas treatment in cellulose production are based on expert estimates on the specific consumption of limestone per tonne of cellulose as documented in the EMIS database (EMIS 2017/1A2d Zellulose Produktion)

The activity data of limestone and sodium bicarbonate use in waste incineration plants are provided by the industry as documented in the EMIS database (EMIS 2017/1A1a Kehrichtverbrennungsanlagen).

The amount of limestone used as flux in iron foundries (cupola furnaces) is estimated by the Swiss foundry association to be in the range of 30 – 50% of the coal consumed. Therefore, an average share of 40% is assumed to calculate the activity data of limestone use (EMIS 2017/1A2a Eisengiessereien Kupolöfen).

The activity data of sodium bicarbonate correspond to the net import of sodium bicarbonate. These data are provided by the Swiss customs administration (EZV, FCA 2016b).

For activity data see Table 4-14.

Plaster production (2A4d)

Methodology

There are two plaster production sites in Switzerland. The emissions stem mainly from blasting operations.

Emissions from blasting operations are determined by a country-specific method analogous to a Tier 2 method of EMEP/EEA Guidebook 2013 (EMEP/EEA 2013).

Emission factors

As there are no specific emission factors for gypsum mining, the emission factors for cement raw material mining are taken instead (with a rough estimate that 1.5 t of raw material are required for production of 1 t of cement). This method is documented in EMIS 2017/2A4d Gips-Produktion übriger Betrieb.. For emission factors see Table 4-15.

Activity data

The activity data of the annual amount of raw material processed in the plaster production are based on industry data and expert estimates as documented in EMIS 2017/2A4d Gips-Produktion übriger Betrieb (see Table 4-14).

4.2.3 Uncertainties and time-series consistency

The uncertainty for CO₂ emissions in 2A1 Cement production, which is a key category regarding level and trend, amounts to 4%. The uncertainty of CO₂ emissions was calculated following the steps in Table 3.2 in 2006 IPCC Guidelines (IPCC 2006, vol. 1, chp. 3, p. 3.30-3.31). An uncertainty of 2% is assumed for activity data and 3% for the emission factor, which consists of an average emission factor per tonne of clinker for calcination of the carbonate containing raw material (FOEN 2013q, chp. G.7) and a correction for the content of organic carbon and cement kiln dust.

Combined uncertainty is estimated to be 3% for emissions from 2A2 Lime production and 4% for emissions from 2A3 Glass production (expert estimate).

For CO₂ emissions in source category 2A4 Other process uses of carbonates, an overall uncertainty of 3% is assumed. Most of the data stems from industrial plants participating in the Swiss ETS, which requires that the uncertainty in the emissions does not exceed a given limit (1.5%-7.5%, depending on the amount of emissions resulting from a given source) and from the Swiss Federal Customs Administration.

Consistency: Time series for 2A Mineral industry are all considered consistent.

4.2.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

For submission 2017, implied emission factors of 2A3 container glass production were assessed by both a Tier 2 and Tier 3 method for the years 2005 – 2011. This comparison provides an indication of the differences caused by the switch in the Tier level from Tier 2 (1990-2004) to Tier 3 (2005-2015).

4.2.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 and 2014 of at least 0.3 kt CO₂ eq are quantified. All the other recalculations have an impact of less than 0.3 kt CO₂ eq in the years 1990 and 2014.

- 2A1: The EF for CO₂ geog. has been changed to a constant value of 536.718 kg/t clinker for the whole time period which is higher than the EF used before. It now corresponds to the calculation in the Swiss emissions trading system. This results in an emission increase of 860 kt over the entire period 1990-2015. In 2014 the increase amounts to 21.6 kt CO₂ eq and in 1990 the increase amounts to 56 kt CO₂ eq.
- 2A3: AD of 2A3 Container glass have been revised for 2003-2006 based on monitoring reports of the Swiss emissions trading scheme.
- 2A4d: The use of carbonates for sulphur oxide removal in municipal solid waste incineration plants has been moved from 1A1a to 2A4d Other process uses of carbonates (entire time series). In 2014, this reallocation leads to an increase in emissions by 1.4 kt CO₂ eq and in 1990 the emissions increase by 0.3 kt CO₂ eq.
- 2A4d: The use of limestone for sulphur oxide removal in cellulose production has been moved from 1A2d to 2A4d Other process uses of carbonates (1990-2008). In 1990, this reallocation leads to an increase in emissions by 4 kt CO₂ eq.
- 2A4d: The emissions of geogenic CO₂ from limestone use in cupola furnaces of iron foundries are newly reported in the inventory for the entire time series. This leads to an increase by 0.4 kt CO₂ eq in 2014 and 4 kt CO₂ eq in 1990.
- 2A4d: Negligible rounding changes in AD of 2A4d Other use of carbonates for 1990-2013.
- 2A4d: AD of 2A4d Carbonate use in municipal solid waste incineration has been adjusted for 2014 yielding revised AD for 2014 of 2A4d Other use of carbonates as well.

4.2.6 Category-specific planned improvements

There are no category-specific planned improvements

4.3 Source category 2B – Chemical industry

4.3.1 Source category description

Approach 1 and 2 key category 2B

Source category 2B Chemical industry is not a key category.

Table 4-17 Specification of source category 2B Chemical industry in Switzerland.

2B	Source	Specification
2B1	Ammonia production	Emissions of CO ₂ and NMVOC are reported in 2B8b Ethylene production
2B2	Nitric acid production	Emissions of N ₂ O and NO _x from the production of nitric acid
2B5	Carbide production	Emissions of CO ₂ , CH ₄ and SO ₂ from the production of silicon carbide
2B8	Petrochemical and carbon black production	Emissions of CO ₂ and NMVOC from ethylene production. In Switzerland there is only ethylene production under this source category
2B10	Other	Emissions of CO ₂ , CH ₄ , CO and NMVOC from acetic acid production; CO ₂ emissions from limestone pit and niacin production; NMVOC emissions from PVC production (ceased in 1996); SO ₂ emissions from sulphuric acid production

4.3.2 Methodological issues

4.3.2.1 Ammonia production (2B1)

Ammonia (NH₃) is produced in one single plant in Switzerland by catalytic reaction of nitrogen and synthetic hydrogen (see Figure 4-4). Ammonia is not produced in an isolated reaction plant but is part of an integrated production chain (see Figure 4-5).

The starting production process is the thermal cracking of liquefied petroleum gas (LPG) and light virgin naphtha yielding ethylene (ethene, C₂H₄), and a series of by-products such as e.g. synthetic hydrogen and methane, which are used as educts for further production steps. According to the Swiss ammonia producer it is not possible to split and allocate the emissions of the cracking process (CO₂ and NMVOC) to every single product such as, e.g., ethylene, acetylene (ethyne, C₂H₂), cyanic acid or ammonia. Therefore, all CO₂ and NMVOC emissions of the cracking process are allocated to the ethylene production and are reported under the category 2B8b Ethylene production. Thus, for source category 2B1 Ammonia production, CO₂ and NMVOC emissions are reported as included elsewhere (IE). All information on the ammonia production and the cracking process is documented in EMIS 2017/2B1 Ammoniak-Produktion and EMIS 2017/2B8b Ethen-Produktion, respectively.




Figure 4-4 Process flow chart for the production of ammonia (NH₃) from nitrogen (N₂) and hydrogen (H₂, synth.). Hydrogen is derived from the thermal cracking process in the same plant (see Figure 4-5).




Figure 4-5 Process flow chart for the production of ethylene (C₂H₄) and acetylene (C₂H₂) by thermal cracking of liquefied petroleum gas (LPG) and light virgin naphtha. The intermediate product H₂, synth. is used as educt in the ammonia production in the same plant (see Figure 4-4).

Table 4-18 Activity data for ammonia production in Switzerland are documented in the confidential NIR, which is available to reviewers on request.

4.3.2.2 Nitric acid production (2B2)

In Switzerland, there is one single plant producing nitric acid (HNO₃). Nitric acid is produced by catalytic oxidation of ammonia (NH₃) with air. At temperatures of 800°C nitric monoxide (NO) is formed. During cooling, nitrogen monoxide reacts with excess oxygen to form nitrogen dioxide (NO₂). The nitrogen dioxide reacts with water to form 60% nitric acid (HNO₃). Today, two types of processes are used for nitric acid production: single pressure or dual pressure plants. In Switzerland a dual pressure plant is installed.

During this process, nitrous oxide (N_2O) can be formed as an unintentional by-product. In addition, also some nitrogen oxide (NO_x) is produced. In the Swiss production plant abatement of NO_x is done by selective catalytic reduction (SCR, installed in 1988), which reduces NO_x to N_2 and O_2 (the SCR in this plant is also used for treatment of other flue gases and was not installed for the HNO_3 production specially). In 1990, an automatic control system for the dosing of ammonia to the SCR process was installed. A new catalyst installed in 2013 reduced the N_2O emissions.

No additional abatement technique is installed to destroy N_2O . A decomposition of N_2O occurs, to some extent, simultaneously in the NO_x reduction process.

Methodology

According to decision tree Fig. 3.2 of the IPCC 2006 guidelines (vol. 3, chp. 3.3 Nitric acid production), the N_2O emissions from nitric acid production are determined by a Tier 2 method during the time period 1990-2012 and by a Tier 3 method since 2013, based on direct measurements. The NO_x emissions are calculated by a Tier 2 method according to the decision tree Fig. 3.1 in EMEP/EEA (2013) (chp. 2B Chemical industry) using a plant-specific emission factor.

Emission factors

The N_2O and NO_x emission factors for nitric acid production in Switzerland are based on measurements from the single nitric acid production plant.

The measurement of N_2O was carried out in 2009 according to the guideline VDI-Richtlinie 2469/Blatt 1 (Messen gasförmiger Emissionen - Messen von Distickstoffmonoxid - Manuelles gaschromatographisches Verfahren) and is the only plant-specific measurement of N_2O emissions. The test gas is sucked in via a heated titanium sensor and then treated with a solution of potassium permanganate and hydrogen peroxide in order to remove nitrogen oxides and further disturbing components. The N_2O concentration is then measured using a gas chromatograph with an electron capture detector. The measurement uncertainty is $\pm 20\%$ (minimum $\pm 0.5 \text{ mg/m}^3$). On repeated enquires the plant confirmed that since a denitrification system and an automatic control system for the ammonia addition was installed in 1988 and 1990, respectively, no modifications were made in the production line until 2012. Therefore, a constant N_2O -emission factor is assumed for this time period. A new catalyst installed in 2013 reduced the N_2O emissions, which are measured online by NDIR photometry from 2013 onwards.

The NO_x emission factor is the mean value based on three plant-specific measurements in 2007, 2009 and 2012. Since no modifications were made in the production line between 1990 and 2012 a constant emission factor is assumed for this time period. In 2013, the volume of the SCR-plant was duplicated. This modification together with the new catalyst in the production line slightly reduced the NO_x emission factor. The values are documented in EMIS 2017/2B2 Salpetersäure Produktion.

Table 4-19 Emission factors for N₂O and NO_x for nitric acid production in Switzerland in kg/t nitric acid for 2015. Data refers to 100% nitric acid

2B2 Nitric acid production	Unit	N ₂ O	NO _x
	kg/t	C	C

Activity data

Activity data on annual production of nitric acid (100%) are provided annually by the Swiss production plant for the entire time period 1990-2015. Since 2013, activity data of the annual nitric acid production is taken from annual monitoring reports from the Swiss Emissions Trading Scheme (ETS). The data are confidential but available to reviewers (see EMIS 2017/2B2 Salpetersäure Produktion).

Table 4-20 Activity data for the production of nitric acid (100%) in Switzerland are documented in the confidential NIR, which is available to reviewers on request.

4.3.2.3 Carbide production (2B5)

In Switzerland, there is one single plant producing carbide. The plant produces silicon carbide, which is used in abrasives, refractories, metallurgy and anti-skid flooring. The Swiss silicon carbide is produced in an electric furnace at temperatures above 2000°C using the Acheson process. The starting materials are quartz sand (SiO₂), petroleum coke and anthracite (C) which yield silicon carbide (SiC) and carbon monoxide (CO). The CO is converted to CO₂ in excess oxygen and released to the atmosphere. Petroleum coke and anthracite – although to a lower portion – may contain volatile organic compounds, which can form methane (CH₄) as an unintended by-product. There is no abatement techniques installed which could capture the CO₂ or CH₄ emissions.

Methodology

According to decision tree Fig. 3.5 of the IPCC 2006 guidelines (vol. 3, chp. 3.6 Carbide production), the CO₂ and CH₄ emissions from silicon carbide production are determined by a Tier 2 method. The SO₂ emissions are calculated by a Tier 2 method according to the decision tree Fig. 3.1 in EMEP/EEA (2013) (chp. 2B Chemical industry) using plant-specific emission factors.

Emission factors

The CO₂, CH₄ and SO₂ emission factors are confidential and available to reviewers on request. The values are partly based on measurements from the single silicon carbide production plant and are documented in EMIS 2017/2B5 Graphit und Siliziumkarbid Produktion.

Table 4-21 In the confidential NIR, a respective table with emission factors of fossil CO₂ in kg/t silicon carbide are provided. Data are available to reviewers on request.

Table 4-22 Emission factors for CO₂, CH₄ and SO₂ for carbide production in kg/t silicon carbide in Switzerland in kg/t for 2014.

2B5 Carbide production	Unit	CO₂	CH₄	SO₂
2B5a Silicon carbide	kg/t	C	C	C

Activity data

Activity data on annual production of silicon carbide are provided annually from 1997 onwards by the Swiss production plant. For the time period 1990-1996 activity data are based on industry data for 1990 and 1995 and interpolated values in between. For 2015, the plant did not provide any data. Therefore activity data are assumed to remain at the same level as in 2014.

The data are confidential but available to reviewers on request (see EMIS 2017/2B5 Graphit und Siliziumkarbid Produktion).

Table 4-23 In the confidential NIR, the respective table with activity data on silicon carbide production in Switzerland is separately reported and available to reviewers.

4.3.2.4 Petrochemical and carbon black production (2B8)

Ethylene (2B8b)

Ethylene (ethene, C₂H₄) is produced by a single plant in Switzerland by thermal cracking of liquefied petroleum gas (LPG) and virgin naphtha. Ethylene is not produced in an isolated process but is co-processed together with several other products such as H₂, CH₄, and C₂H₂ (see flow chart in Figure 4-5 in chp. 4.3.2.1). From the thermal cracking process, emissions of CO₂ and NMVOC are released. They are both allocated entirely to the production of ethylene, which is the first product within the integrated production chain. CH₄ emissions to atmosphere do not occur since CH₄ is completely used as an educt in the downstream production of cyanic acid (HCN) in the same facility (see Figure 4-5 and for further information see EMIS 2017/2B8b Ethen-Produktion). Therefore, CH₄ emissions are reported as NA for ethylene production and only CO₂ and NMVOC emissions are reported.

Methodology

According to decision trees Fig. 3.8 of the IPCC 2006 guidelines (vol. 3, chp. 3.9 Petrochemical and carbon black production) and Fig. 3.1 of EMEP/EEA (2013) (chp. 2B Chemical industry), the CO₂ and NMVOC emissions, respectively, from ethylene production are determined by a Tier 2 method using plant-specific emission factors (EMIS 2017/2B8b Ethylene production).

Emission factors

The CO₂ and NMVOC emission factors for ethylene production are based on industry data from the single ethylene production plant in Switzerland. Annual emission data were only available from the year 2000 onwards. For the period 1990-1999 a constant value, i.e. the mean value of the years 2000-2009 was assumed.

The emission factors for ethylene production are considered confidential; however, they are available to reviewers on request.

Table 4-24 Emission factors for CO₂ and NMVOC in ethylene production, NMVOC in acetic acid production, CO₂ in limestone pit and niacin production and SO₂ in sulphuric acid production for 2015 in kg/t product.

	Unit	CO ₂	NMVOC	SO ₂
2B8 Petrochemical and carbon black production				
2B8b Ethylene	kg/t	C	C	NA
2B10 Other				
Acetic acid production	kg/t	NA	C	NA
Limestone pit	kg/t	C	NA	NA
Niacin production	kg/t	C	NA	NA
Sulphuric acid production	kg/t	NA	NA	C

Table 4-25 CO₂ fossil emission factors in 2B8b Ethylene are documented in the confidential NIR, which is available to reviewers on request.

Activity data

Activity data on the annual production of ethylene are provided annually by the single ethylene production plant in Switzerland. Since 2013, activity data of the annual nitric acid production is taken from annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

The data are considered confidential but available to reviewers on request.

Table 4-26 Activity data for the production of ethylene, acetic acid, niacin, PVC and sulphuric acid as well as for limestone pit in Switzerland in kt.

	Unit	1990	1995	2000	2005
2B8 Petrochemical and carbon black production					
2B8b Ethylene	kt	C	C	C	C
2B10 Other					
Acetic acid production	kt	30	27	24	8
Limestone pit	kt	C	C	C	C
Niacin production	kt	C	C	C	C
PVC production	kt	43	43	NO	NO
Sulphuric acid production	kt	C	C	C	C

	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2B8 Petrochemical and carbon black production											
2B8b Ethylene	kt	C	C	C	C	C	C	C	C	C	C
2B10 Other											
Acetic acid production	kt	8	9	18	28	20	18	12	C	C	C
Limestone pit	kt	C	C	C	C	C	C	C	C	C	C
Niacin production	kt	C	C	C	C	C	C	C	C	C	C
PVC production	kt	NO									
Sulphuric acid production	kt	C	C	C	C	C	C	C	C	C	C

4.3.2.5 Other (2B10)

Source category 2B10 Other comprises emissions from production of acetic acid, sulphuric acid, niacin and PVC (ceased in 1996) as well as from limestone pits.

Acetic acid production (2B10)

In Switzerland, there is only one plant producing acetic acid (CH₃COOH) remaining after the other one stopped its production by the end of 2012. The still existing plant emits NMVOC only whereas from the latter one also emissions of CO₂, CH₄ and CO occur.

Methodology

In order to determine emissions of CO₂ and CH₄ from acetic acid a country-specific method analogous to a Tier 2 method according to the IPCC 2006 guidelines (vol. 3) is used. The CO and NMVOC emissions are calculated by a Tier 2 method according to the decision tree Fig. 3.1 in EMEP/EEA (2013) (chp. 2B Chemical industry).

Emission factors

The emission factors for CO₂, CH₄, CO and NMVOC from acetic acid production in Switzerland are plant-specific and based on data from industry and expert estimates documented in EMIS 2017/2B10 Essigsäure-Produktion.

In the plant which ceased production by the end of 2012 process emissions had been treated in a flue gas incineration. Thus, the reported emissions of CH₄, CO and NMVOC only

occurred in case of malfunction, which resulted in strongly fluctuating plant-specific emission factors. In addition, the resulting implied emission factors based on the emissions of both plants are modulated by considerable production fluctuations of one of the plants from 2000 onwards.

The emission factors for acetic acid production are confidential but available to reviewers on request.

Table 4-27 In the confidential NIR, the respective table with emission factors for CO₂ and CH₄ in acetic acid production are separately reported and available to reviewers.

Activity data

The annual amount of produced acetic acid is based on data from industry and from the Swiss industry association for the chemical, pharmaceutical and biotech industry (scienceindustries) documented in EMIS 2017/2B10 Essigsäure-Produktion (see Table 4-26).

The data for acetic acid production since 2013 are confidential, since there is only one manufacturer remaining. The data are available for reviewers on request.

Limestone pit (2B10)

In one chemical plant acids are neutralized in a so-called limestone pit yielding geogenic CO₂ emissions.

Methodology

According to decision tree Fig. 2.4 of the IPCC 2006 guidelines (vol. 3, chp. 2.5 Other process uses of carbonates), the CO₂ emissions from the limestone pit are determined by Tier 2 method.

Emission factors

The CO₂ emission factor is considered confidential but available to reviewers on request.

Activity data

Activity data of annual consumption of calcium carbonate are provided by the chemical plant from 1999 onwards as documented in EMIS 2017/2B10 Kalksteingrube. For the years 2005-2011 and since 2013 they are based on monitoring reports of the Swiss ETS. Since no data are available of the limestone pit for the time period 1990-1998, the annual activity is derived from the average annual consumption between 1999 and 2015.

Activity data is considered confidential but available to reviewers on request.

Niacin production (2B10)

CO₂ emissions from niacin production of the single manufacturer in Switzerland are reported since submission 2014. CO₂ is released in the last reaction step of the niacin production.

Methodology

In order to determine emissions of CO₂ from niacin production, a country-specific method analogous to a Tier 2 method according to the IPCC 2006 guidelines (vol. 3) is used.

Emission factors

The CO₂ emission factor is plant-specific based on monitoring reports of the Swiss ETS and is assumed to be constant as documented in the EMIS database (EMIS 2017/2B10 Niacin-Produktion). The emission factor is considered confidential but available to reviewers on request.

Activity data

Activity data of annual niacin production were provided by the Swiss production plant for the entire time period as documented in EMIS 2017/2B10 Niacin-Produktion. For the years 2005-2011 and since 2013 they are based on monitoring reports of the Swiss ETS.

Activity data are considered confidential but available to reviewers on request.

PVC and sulphuric acid production (2B10)

Sulphuric acid (H₂SO₄) is produced by one plant only in Switzerland. From this production process SO₂ is emitted. Until 1996, also PVC was produced in Switzerland releasing NMVOC emissions.

Methodology

In order to determine SO₂ and NMVOC emissions from sulphuric acid and PVC production, respectively, a Tier 2 method according to the decision tree Fig. 3.1 of EMEP/EEA (2013) (chp. 2B Chemical industry) with plant-specific emission factors is used.

Emission factors

The emission factor for SO₂ from sulphuric acid production in Switzerland is plant-specific and based on measurement data from industry and expert estimates documented in the EMIS database (EMIS 2017/2B10 Schwefelsäure-Produktion).

The SO₂ emission factor is confidential but available to reviewers on request.

For PVC production the NMVOC emission factor was based on industry information and expert estimates (EMIS 2017/2B10 PVC-Produktion).

Activity data

The annual amount of sulphuric acid and PVC produced is based on data from industry and expert estimates documented in EMIS 2017/2B10 Schwefelsäure-Produktion and EMIS 2017/2B10 PVC-Produktion (see Table 4-26). The activity data for sulphuric acid production are confidential but available to reviewers on request.

4.3.3 Uncertainties and time-series consistency

The uncertainties for CO₂ in source category 2B are estimated to be medium, (see Table 1-11 Semi-quantitative uncertainties for non-key categories) resulting in a relative uncertainty of 10%. For CH₄ a combined uncertainty of 20% is estimated.

For N₂O emissions from 2B2 Nitric acid production, the uncertainty is assumed to be 7.5% since the Swiss ETS requires that an uncertainty of 7.5% is not exceeded for continuous N₂O measurements.

Consistency: Time series for 2B Chemical industry are all considered consistent.

4.3.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

4.3.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. The impact of these recalculations is less than 0.3 kt CO₂ eq.

- 2B10: Minor changes in CO₂ emissions from 2B10 Limestone pit due to rounding differences of AD for 1990-1998.
- 2B10: The extrapolated values of the SO₂ emission factor of 2B10 Sulphuric acid production have been revised (1990-2008).

4.3.6 Category-specific planned improvements

No category-specific improvements are planned.

4.4 Source category 2C – Metal industry

4.4.1 Source category description

Table 4-28 Key categories (KCA incl. LULUCF) of 2C Metal industry.

Code	IPCC Category	GHG	Identification Criteria
2C3	Aluminium production	CO ₂	T1, T2
2C3	Aluminium production	PFC	T1, T2

Table 4-29 Specification of source category 2C Metal industry in Switzerland.

2C	Source	Specification
2C1	Iron and steel production	Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from the production of iron and steel
2C2	Ferroalloys production	Production is not occurring in Switzerland
2C3	Aluminium production	Emissions of PFC, CO ₂ , NO _x , CO, NMVOC, and SO ₂ from the production of primary aluminium (ceased in 2006). Emissions from use of SF ₆ in aluminium foundries.
2C4	Magnesium production	Emissions from use of SF ₆ in magnesium foundries
2C7	Other	Emissions of CO and NMVOC from non-ferrous metal foundries Emissions of CO ₂ , NO _x , CO and SO ₂ from battery recycling

4.4.2 Methodological issues

4.4.2.1 Iron and steel production (2C1)

There is no primary iron and steel production in Switzerland. Only secondary steel production occurs, which is steel production from recycled steel scrap. After closing down of two steel plants in 1994, there remain two plants in Switzerland. Both plants use electric arc furnaces (EAF) with a carbon electrode for melting the steel scrap. During the melting process CO₂ emissions occur mainly from scrap, electrodes and carburization coal whereas the produced steel, filter dust and slag act as carbon sinks. Emissions of precursors such as NO_x, CO, NMVOC and SO₂ occur as well.

In Switzerland, no production of pig iron occurs but iron is processed in foundries only. Today, there exist about 14 iron foundries in Switzerland. About 75% of the iron is processed in induction furnaces and 25% in cupola furnaces. From induction furnaces only precursors are emitted. In cupola furnaces also CO₂ emissions from other bituminous coal occur. Other bituminous coal acts first of all as fuel but also as carburization material and reductant. Therefore it was decided to report those CO₂ emissions in source category 1A2a. Geogenic CO₂ emissions from use of limestone in cupola furnaces are reported in 2A4d Other process uses of carbonates.

The CO₂ emissions from 2C1 Secondary steel production, electric arc furnace are based on a carbon mass balance considering all carbon sources and sinks of the process. Therefore,

the emissions of CO and NMVOC are no longer included in the calculation of the indirect CO₂ emissions from sector 2 IPPU in order to avoid a double counting.

Methodology

For determination of CO₂ emission from iron and steel production a mixture of a Tier 2 and a Tier 3 method according to decision tree Fig. 4.7 IPCC 2006 (vol. 3, chp. 4.2 Iron & steel and metallurgical coke production) is used. For the years 2005-2011 and 2013 plant-specific data on the carbon mass balance is available from monitoring reports of the Swiss ETS, since under the Ordinance for the Reduction of CO₂ Emissions (Swiss Confederation 2012) the plants are required to report their emissions annually (Tier 3). From this information, data for the other years are interpolated for calculating an implied emission factor. In Switzerland, no CH₄ emissions occur in the EAF process.

Emissions of all precursors are determined by a Tier 2 method based on the decision tree Fig. 3.1 in chapter 2C1 in EMEP/EEA (2013) using country-specific emission factors (EMIS 2017/2C1).

Emission factors

The emission factors for iron and steel production in Switzerland are country-specific and are based on measurements from industry and expert estimates documented in the EMIS database (EMIS 2017/2C1 Eisengiessereien Elektroschmelzofen/übriger Betrieb, EMIS 2017/2C1 Stahl-Produktion Elektroschmelzöfen and EMIS 2017/2C1 Stahlwerke Walzwerke).

The electrode consumption in the two Swiss plants differs. For the calculations all carbon sources (graphite electrodes, steel scrap, alloy coal, etc.) and carbon sinks (steel, filter dust and slag) for the years 2005–2011 and 2013–2014 were taken into account. Based on these carbon mass balances, a mean plant-specific CO₂ emission factor results. The reported CO₂ emission factor for Swiss steel industry is the production-weighted average. Consequently, there are no indirect CO₂ emissions to be accounted for in chp. 9 Indirect CO₂ and N₂O emissions for source category 2C1.

The plant-specific data are confidential but available to reviewers on request.

Table 4-30 CO₂ emission factor of electric arc furnaces in 2C1 Steel production in kg/t.

2C1 Steel production	Unit	1990	1995	2000	2005
CO ₂	kg/t	8.3	8.0	7.7	8.8

2C1 Steel production	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
CO ₂	kg/t	9.1	8.5	6.8	6.8	7.6	7.1	7.9	8.5	8.2	8.6

Emission factors for all precursors emitted from steel production are based on air pollution control measurements of the steel plants. For submission 2016, emission factors of NO_x, NMVOC, SO₂, and CO have been revised based on air pollution control measurements at

the electric arc furnaces of the two plants in 1999, 2005 and 2010 and in 1998, 2009 and 2014, respectively. The emission factors from iron production in foundries are provided by the Swiss foundry association (GVS).

Table 4-31 Emission factors for NO_x, CO and NMVOC in iron production, for CO₂, NO_x, CO, NMVOC and SO₂ in steel production, for CO and NMVOC in non-ferrous metal production and for CO₂, NO_x, CO and SO₂ in battery recycling for 2015.

2C Metal industry	Unit	CO ₂	NO _x	CO	NMVOC	SO ₂
2C1 Iron production	kg/t	IE	0.01	4.1	4	NA
2C1 Steel production	kg/t	8.6	0.14	0.7	0.1	0.014
2C7a Non-ferrous metals	kg/t	NA	NA	0.24	0.05	NA
2C7c Battery recycling	kg/t	570	0.62	0.9	0.20	0.01

Activity data

Activity data on annual production of iron and steel are provided annually by the Swiss foundry association (Giesserei-Verband Schweiz, GVS) and the steel plants, respectively. Since 2009, activity data of the annual steel production is taken from annual monitoring reports from the Swiss Emissions Trading Scheme (ETS).

Table 4-32 Production of iron, steel, aluminium and non-ferrous metals as well as amount of batteries recycled in Switzerland in kt.

2C Metal industry	Unit	1990	1995	2000	2005
2C1 Iron production	kt	170	130	120	67
2C1 Steel production	kt	1'108	716	1'022	1'159
2C3 Aluminium production	kt	87	21	36	45
2C7a Non-ferrous metals	kt	55	60	70	33
2C7c Battery recycling	kt	3.0	3.0	3.0	2.8

2C Metal industry	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2C1 Iron production	kt	67	72	78	49	53	61	46	45	43	37
2C1 Steel production	kt	1'254	1'267	1'315	935	1'218	1'322	1'252	1'231	1'315	1'296
2C3 Aluminium production	kt	12	NO	NO	NO	NO	NO	NO	NO	NO	NO
2C7a Non-ferrous metals	kt	30	28	21	15	20	12	18	7	7	7
2C7c Battery recycling	kt	2.4	2.4	2.5	3.4	3.3	2.4	2.4	2.3	2.6	2.5

4.4.2.2 Aluminium production (2C3)

Methodology

The last production site for primary aluminium in Switzerland closed down in April 2006. According to the 2006 IPCC Guidelines (IPCC 2006, vol.3, chp. 4.4, fig. 4.11) CO₂ emissions were calculated by a Tier 2 method using a country-specific emission factor. For PFC emissions, a more specific Tier 3 method with facility-specific data according to the 2006 IPCC Guidelines (IPCC 2006) was used. Operating smelter emissions have been monitored periodically by the industry for selected years.

FOEN import statistics indicate in the year 2003 part of the SF₆ imports to be related to the aluminium industry, referring to cleaning process in foundries. The 2006 IPCC Guidelines

mention use of SF₆ in aluminium production for magnesium alloys on a low scale but do not provide further information for evaluation. Accordingly, the same evaluation methodology as for magnesium foundries with an emission factor based on a Tier 2 method is applied.

Emission factors

The emission factor for CO₂ of 1.6 tonnes per tonne of aluminium is country-specific. It is based on measurements and data from industry and expert estimates, documented in the EMIS database (EMIS 2016/2C3 Aluminium Produktion). CO₂ emissions from aluminium production stem from the oxidation of the anode in the electrolysis process. In Switzerland, only prebake anode technology was used. For the anode consumption, a constant mean value of 0.43 tonnes per tonne of aluminium was applied. It is assumed that the anode consisted completely of carbon and that it was fully oxidized during the process. Therefore, there are no indirect CO₂ emissions to be accounted for in chp. 9 Indirect CO₂ and N₂O emissions from CO emissions of primary aluminium production. But as the NMVOC emissions originate solely from the production of the electrodes at the plants they have to be considered for the calculation of the indirect CO₂ emissions in chp. 9.

Before the close down of the only Swiss primary aluminium factory in 2006, PFC emission factors of operating smelters have been monitored periodically. The factory provided own measurements for 1990, 1999 and 2000 yielding smaller EFs than the European average (Alcan 2003). The comparison of these data with data from IAI (2005) on global PFC emissions from aluminium production showed that the monitored emissions from the smelter in Switzerland were lower by a factor of about 4. This seems to be plausible because they used point feed prebake (PFPB) technology which is known for the lowest emissions per tonne of aluminium. Therefore, for the years with no measured emission data available, the average European emission factors as reported by the European Aluminium Association (Alcan 2002) are applied, reduced by a 'general reduction factor' of four for both PFC gases (CF₄ and C₂F₆). The resulting emission factors for Switzerland are within the uncertainty range according to the 2006 IPCC Guidelines (variations by a factor of 10 using same technologies). It is assumed that the PFC emissions consist to 90% of CF₄ and to 10% of C₂F₆. Because aluminium production was closed in 2006, it is not possible to redo any measurements or to collect any information about the process details retroactively. The emission factors, measured and reported by the industry back then, have decreased between 1990 and 2006 due to technical efforts to reduce emissions (Alcan 2003).

Table 4-33 PFC emission factors for aluminium production in Switzerland. Aluminium production in Switzerland ceased in 2006.

Gas	Unit	1990	1995	2000	2005
CF ₄	kg/t	0.1530	0.0833	0.0360	0.0315
C ₂ F ₆	kg/t	0.0170	0.0093	0.0040	0.0035

Gas	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
CF ₄	kg/t	0.0315	NO								
C ₂ F ₆	kg/t	0.0035	NO								

There are no measurements of SF₆ emissions available from aluminium foundries to identify the fraction of SF₆ destroyed or transformed in the cleaning process. For SF₆ used in aluminium foundries (2C3) it is therefore assumed that the total imported amount is emitted, in accordance with the default emission factor (1000 kg per tonne of imported substance) of the 2006 IPCC Guidelines (IPCC 2006).

Activity data

In 2006, the last aluminium production site in Switzerland was closed. Activity data on aluminium production from 1997 to 2006 are based on annual data published by the Swiss Aluminium Association. For earlier years, data were provided directly by the aluminium industry. Activity data for aluminium production in Switzerland are given in Table 4-32.

Activity data on SF₆ used in aluminium foundries (2C3) is derived from import data from FOEN statistics. Import companies indicated in the year 2003 a portion of SF₆ imports for foundries to be used for aluminium cleaning. For the activity data of any particular year, the mean value of the imports in the present and the previous year is used to account for possible time lag between import and consumption (e.g. for 2004 the mean value of 2003 and 2004 import data are used). It is assumed that the total imported amount is emitted within one year. In 2011, a study was carried out among members of the Swiss Foundry Association (GVS), confirming that SF₆ is not used any more in aluminium foundries. As no details on the imported amount are available for the time period 2003–2011, a steady decrease of the import amount of SF₆ is assumed from 2003 until the final elimination of SF₆ for aluminium cleaning in 2011. This assumption is based on the above-mentioned survey and on information obtained on applications within the category 'others' from FOEN import statistics.

4.4.2.3 Magnesium production (2C4)

Use of SF₆ in magnesium foundries (2C4)

SF₆ is used in Swiss magnesium foundries since 1997. There have been two magnesium foundries known to be using SF₆. In 2007 one of them closed down.

Methodology

SF₆ is used in magnesium foundries in the cleaning process as inert gas to fill casting forms. The Swiss Foundry Association (GVS) has not provided information on emission factors and hence a Tier 2 method is used.

Emission factors

There are no measurements of SF₆ emissions available to identify the fraction of SF₆ destroyed or transformed in the process. For SF₆ used in magnesium foundries (2C4) it is therefore assumed that the total imported amount is emitted, in accordance with the default emission factor (1000 kg per tonne of imported substance) of the 2006 IPCC Guidelines (IPCC 2006).

Activity data

Activity data on SF₆ used in magnesium foundries (2C4) are based on import data from FOEN statistics. For the activity data of any particular year, the mean value of the imports in the present and the previous year is used to account for possible time lag between import and consumption (e.g. for 2015 the mean value of 2014 and 2015 import data are used). It is assumed that the total imported amount is emitted within one year. The remaining magnesium foundry reported activity data for 2008 to 2015 to the SWISSMEM statistics, the information is in accordance with import data from FOEN statistics. The fact that only one magnesium foundry uses SF₆ was confirmed by a survey, which has been carried out in 2011 among members of the Swiss Foundry Association (GVS).

4.4.2.4 Other (2C7)

Battery recycling and non-ferrous metal foundries (2C7)

There is one battery recycling plant in Switzerland. The recycling is done by applying the Sumitomo process. The batteries are first pyrolysed at temperatures of 700°C in a reducing atmosphere in a shaft kiln. The gas with the carbonised components then goes to a post-combustion step where it is completely oxidised at temperatures of 1000°C. The flue gas is then directed to a flue gas treatment installation. The metal fraction from the pyrolysis goes to a melting furnace where it is reduced by addition of coal and magnesium oxide. As reducing agent coke and Carburit is used.

In Switzerland, there are one large company and several small plants operating non-ferrous metal foundries producing mainly copper alloys. During the melting process emissions of CO and NMVOC occur.

Methodology

To determine emissions of CO₂, NO_x, CO and SO₂ from battery recycling and of CO and NMVOC from non-ferrous metal foundries, Tier 2 methods according to EMEP/EEA Guidebook 2013 (EMEP/EEA 2013, chp. 2C7c and 2C7a) with country-specific emission factors are used.

Emission factors

The emission factors of CO₂, NO_x, SO₂, CO from battery recycling between 1990 and 2002 are based on measurements in 2003 as well as mass balances of the single recycling site and are assumed constant. Since 2003 they are based on air pollution control measurements from 2003 and 2012 and are assumed constant during this time period. Emission factors of NMVOC are also based on air pollution control measurements from 2003 and 2012 and are reported for the first time in submission 2017. They are assumed constant for the entire time period (EMIS 2017/2C7 Batterie-Recycling).

Emission factors of CO and NMVOC from non-ferrous metal foundries in Switzerland are country-specific and based on measurements from industry and expert estimates

documented in the EMIS database (2017/2C7 Buntmetallgiessereien Elektroöfen) (see Table 4-31).

Activity data

The annual amount of recycled batteries and produced non-ferrous metals in Switzerland is reported from industry and the foundry association as documented in the EMIS database (EMIS 2017/2C7 Batterie-Recycling and 2017/2C7 Buntmetallgiessereien Elektroöfen), (see Table 4-32).

4.4.3 Uncertainties and time-series consistency

The uncertainty of CO₂ emissions in 2C1 Iron and steel production amounts to 7.1%. Production data of the steel industry have a high confidence and its uncertainty is estimated at 5%. The uncertainty for the CO₂ emission factor is estimated at 5%.

For the emission of CO₂ and PFC from 2C3 Aluminium production, which is a key category for both gases, combined uncertainties of 20.6 % and 9%, respectively, are determined. The emission factor uncertainty for CO₂ and PFC are estimated to be 20% and 6.4%, respectively. The uncertainty in the activity data is estimated to be 5%.

For the emissions of SF₆ from the use in 2C4 Magnesium the combined uncertainty is estimated at 25.5%.

The uncertainty of CO₂ emissions from source category 2C7 Other is estimated to be 20% (expert estimate).

Consistency: Time series for 2C Metal industry are all considered consistent.

4.4.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

For source category 2C4 Magnesium production, the data received from SWISSMEM and import firms have been checked for double counting.

4.4.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. These recalculations cause a change in emission levels 1990 or 2014 by less than 0.3 kt CO₂ eq.

- 2C1: The conversion factor used for calculation of NMVOC emissions from total carbon based on air pollution control measurements has been revised resulting in an adjusted NMVOC emission factor of 2C1 Secondary steel production, electric arc furnace from 1995 onwards.

- 2C7c: The NMVOC emission factor of 2C7c Battery recycling is newly reported in the inventory based on air pollution control measurements (2003 and 2012).
- 2C7c: The emission factors of CO₂, NO_x, CO and SO₂ from 2C7c Battery recycling have been revised from 2002 onwards based on air pollution control measurements (2003 and 2012).
- No recalculations reported in the submission 2017 for F-gases in source category 2C.

4.4.6 Category-specific planned improvements

No category-specific improvements are planned.

4.5 Source category 2D – Non-energy products from fuels and solvent use

4.5.1 Source category description

Source category 2D – Non-energy products from fuels and solvent use is not a key category.

Table 4-34 Specification of source category 2D Non-energy products from fuels and solvent use in Switzerland.

2D	Source	Specification
2D1	Lubricant use	Emissions of CO ₂ from primary usage of lubricants in machinery and vehicles
2D2	Paraffin wax use	Emissions of CO ₂ from primary usage of paraffin waxes
2D3a	Solvent use	Emissions of NMVOC from coating applications, degreasing, dry cleaning and chemical products as well as emissions of CO ₂ resulting from post-combustion of NMVOC in exhaust gases of these sources
2D3b	Road paving with asphalt	Emissions of NMVOC from road paving with asphalt
2D3c	Asphalt roofing	Emissions of CO and NMVOC from asphalt roofing;
2D3d	Urea use in SCR catalysts of diesel engines	Emissions of CO ₂ from urea use in SCR catalysts of diesel engines

4.5.2 Methodological issues

4.5.2.1 Lubricant use (2D1) and Paraffin wax use (2D2)

Lubricants are mostly used in industrial and transportation applications. They can be subdivided into motor oils, industrial oils and greases, which differ in terms of physical characteristics, commercial applications and environmental fate. Lubricants in engines are primarily used for their lubricating properties and associated CO₂ emissions are therefore considered as non-combustion emissions reported in 2D1 Lubricant use.

The source category 2D2 Paraffin wax use includes products such as petroleum jelly, paraffin waxes and other waxes, including mixtures of saturated hydrocarbons, solid at ambient temperature. Paraffin waxes are separated from crude oil during the production of

light (distillate) lubricating oils. Emissions from the use of waxes occur primarily when the waxes or derivatives of paraffins are combusted during use (e.g. candles).

Methodology

CO₂ emissions from oxidation of lubricants and paraffin wax are calculated by a Tier 1 method according to the 2006 IPCC Guidelines (IPCC 2006, vol. 3, chp. 5.2 and 5.3) applying the IPCC default oxidation fraction of 0.2.

Emission factors

The emission factors of CO₂ from lubricant and paraffin wax use in Switzerland are based on default IPCC values for NCV, carbon content and oxidation fraction documented in vol. 2, chp.1 and vol. 3, chp. 5.2 and 5.3, respectively, of IPCC 2006, see also EMIS 2017/2D1 Lubricant use and EMIS 2017/2D2 Paraffin wax use.

Table 4-35 CO₂ emission factor of 2D1 Lubricant use and 2D2 Paraffin wax use for 2015 in kg/t.

	Unit	CO ₂
2D1 Lubricant use	kg/t	590
2D2 Paraffin wax use	kg/t	590

Activity data

The annual amount of lubricant and paraffin wax used in Switzerland is derived from the Swiss petroleum association (EV 2016).

Table 4-36 Use of lubricants in Switzerland.

	Unit	1990	1995	2000	2005
2D1 Lubricant use	kt	80	61	63	72
2D2 Paraffin wax use	kt	11	10	12	10

	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2D1 Lubricant use	kt	68	71	66	52	55	54	51	53	54	52
2D2 Paraffin wax use	kt	11	9	9	6	5	5	3	4	4	4

4.5.2.2 Other (2D3)

Solvent use (2D3a)

Since the 2006 IPCC Guidelines (vol. 3, chp. 5.5) refer to the EMEP/EEA Guidebook 2013 regarding methodologies for estimating NMVOC emissions from solvent use, the respective NFR codes are indicated as reference as well. In the following sections, the NMVOC emissions from coating applications (2D3d NFR), degreasing (2D3e NFR), dry cleaning

(2D3f NFR) as well as production and processing of chemical products (2D3g NFR) are reported. Due to the obligations of the Ordinance on Air Pollution Control (Swiss Confederation 1985) and Ordinance on the Incentive Tax on Volatile Organic Compounds (Swiss Confederation 1997) several industrial plants use facilities and equipment to reduce NMVOC in exhaust gases and room ventilation output. Often this implies the feeding of air with high NMVOC content into the burning chamber of boilers or other facilities to incinerate NMVOC. These CO₂ emissions from post-combustion of NMVOC are estimated based on industry data and expert estimates (Carbotech 2016a).

The source categories paint application in construction, paint application on wood, industrial and non-industrial paint application, production of fine chemicals and cleaning of parts in metal processing account for the largest share of NMVOC emissions from 2D3a in 2015.

Coating applications (2D3d NFR)

Methodology

For the determination of NMVOC emissions from coating applications a Tier 2 method according to the EMEP/EEA Guidebook 2013 is used based on the consumption of paints, lacquers, thinners etc. and their solvent content. Switzerland's Informative Inventory Report 2017 contains a detailed description of the methods and country-specific data used for estimating the NMVOC emissions from 2D3d NFR Coating applications (FOEN 2017f).

Emission factors

Emission factors for NMVOC are based on data from VSLF and retailers as documented in the EMIS database (EMIS 2017/2D3d NFR). In recent years, values of all emission factors for coating applications declined as a result of both a reduction of the solvent content and replacing of solvent based paint by water based paint due to increasingly strict NMVOC regulations by the EU directive (EC 2004). In addition, powder coatings, which are far more efficient, replaced in this time period the conventional paint (rough estimate: 1 t of powder coating replaces 3 t of conventional paint).

For 2D3d NFR Paint application in construction the emission factor of NMVOC is based on a case study by VSLF in 2005 and expert estimates.

Table 4-37 NMVOC emission factors of coating applications, degreasing, dry cleaning, chemical products, manufacture and processing in 2D3a Solvent use for 2015.

2D3a Solvent use	Unit	NMVOC
Coating applications (2D3d NFR)		
Paint application, construction	kg/t paint	55
Paint application, households	kg/t paint	86
Paint application, industrial & non-industrial	kg/t paint	345
Paint application, wood	kg/t paint	291
Paint application, car repair	kg/t paint	400
Degreasing (2D3e NFR)		
Cleaning of electronic components	kg/t solvent	500
Degreasing of metal	kg/t solvent	460
Other industrial cleaning	kg/t solvent	610
Dry cleaning (2D3f NFR)	kg/t solvent	500
Chemical products, manufacture and processing (2D3g NFR)		
Fine chemicals production	t/production index	3.6
Glue production	kg/t glue	0.8
Handling and storing of solvents	t/production index	1.9
Ink production	kg/t ink	8.5
Paint production	kg/t paint	3.5
Pharmaceutical production	kg/t pharmaceutical	7.7
Polyester processing	kg/t polyester	50
Polystyrene processing	kg/t polystyrene	16
Polyurethane processing	kg/t polyurethane	3.6
PVC processing	kg/t PVC	4.0
Rubber processing	kg/tyres	0.14
Tanning of leather	kg/employee	0.68

Activity data

The activity data correspond to the annual consumption of paints. The consumption and solvent content are estimated according to information from the Swiss association for coating and paint applications (VSLF) and in addition from relevant retailers for paint applications in households (EMIS 2017/2D3d NFR). Between 1990 and 1998, the total consumption of paint decreased considerably and increases continuously again since 2001. This trend results from the opposing trends in the different source categories:

- 2D3d NFR Paint application, construction: Activity data of paint application in construction show a substantial reduction compared to 1990 levels. The increasing tendency in paint application since 2000 can be explained by an increase in the construction activity in Switzerland. Since 2000, the expenditures on construction have increased and are thus contributing to an increase in paint application in construction. Before 2000, there was a decline in construction activity, which explains the decreasing tendency in paint application.
- 2D3d NFR Paint application, industrial & non industrial: Between 1990 and 2015, the activity of industrial and non-industrial paint application decreased significantly. Due to structural changes in the industrial sectors, there is a decreasing tendency in emissions between 2000 and 2005. Slight annual fluctuations can be explained by the development of the economic situation, e.g. slight increase in 2007 and decrease in 2008.
- 2D3d NFR Paint application, households: Activity data of paint application in households has more than doubled between 1990 and 2015 due to an increase in demand. The

number of private households increased since 1990, thus leading to an increasing tendency in paint application in the household sector.

Table 4-38 Activity data of coating applications, degreasing, dry cleaning and chemical products, manufacture and processing in Switzerland.

2D3a Solvent use	Unit	1990	1995	2000	2005
Coating applications (2D3d NFR)					
Paint application, construction	kt	122	66	33	42
Paint application, households	kt	12	13	13	20
Paint application, industrial & non-industrial	kt	20	21	21	8.8
Paint application, wood	kt	6.0	6.3	6.5	7.7
Paint application, car repair	kt	2.7	2.2	2.0	1.9
Degreasing (2D3e NFR)					
Cleaning of electronic components	kt	0.90	0.56	0.35	0.64
Degreasing of metal	kt	16	10	5.9	2.6
Other industrial cleaning	kt	0.6	0.6	0.6	1.4
Dry cleaning (2D3f NFR)					
Dry cleaning	kt	1.30	1.01	0.72	0.43
Chemical products, manufacture and processing (2D3g NFR)					
Fine chemicals production	prod. index	70	100	163	224
Glue production	kt	19	32	44	60
Handling and storing of solvents	prod. index	70	100	163	224
Ink production	kt	20	18	18	18
Paint production	kt	138	122	117	122
Pharmaceutical production	kt	16	21	20	28
Polyester processing	kt	11	7	6	7
Polystyrene processing	kt	20	19	19	24
Polyurethane processing	kt	17	35	45	54
Production of adhesive tape	kt	2	NO	NO	NO
PVC processing	kt	94	94	78	64
Rubber processing	tyres	120'000	119'375	103'667	67'000
Tanning of leather	employees	110	108	102	88

2D3a Solvent use	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Coating applications (2D3d NFR)											
Paint application, construction	kt	44	45	48	51	54	56	59	61	61	61
Paint application, households	kt	20	20	23	25	28	28	28	28	28	29
Paint application, industrial & non-industrial	kt	8.9	9.1	8.8	8.6	8.3	8.2	8.0	7.9	7.8	7.7
Paint application, wood	kt	7.8	8.0	8.7	9.3	10.0	10.0	10.0	10.0	10.1	10.3
Paint application, car repair	kt	1.8	1.8	1.8	1.7	1.7	1.5	1.4	1.2	1.2	1.3
Degreasing (2D3e NFR)											
Cleaning of electronic components	kt	0.61	0.57	0.60	0.63	0.67	0.70	0.73	0.73	0.73	0.73
Degreasing of metal	kt	2.5	2.4	2.3	2.2	2.1	2.0	1.9	1.9	1.9	1.9
Other industrial cleaning	kt	1.3	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
Dry cleaning (2D3f NFR)											
Dry cleaning	kt	0.37	0.32	0.26	0.20	0.14	0.09	0.03	0.03	0.03	0.03
Chemical products, manufacture and processing (2D3g NFR)											
Fine chemicals production	prod. index	246	283	280	295	314	299	302	305	308	311
Glue production	kt	62	64	64	64	63	63	63	62	62	62
Handling and storing of solvents	prod. index	246	283	280	295	314	299	302	305	308	311
Ink production	kt	18	19	19	19	19	21	24	26	26	25
Paint production	kt	124	125	125	126	126	126	126	126	125	124
Pharmaceutical production	kt	28	29	29	30	30	30	30	30	30	30
Polyester processing	kt	7.2	7.6	6.2	4.8	3.4	3.5	3.7	3.7	3.7	3.7
Polystyrene processing	kt	26	26	29	31	34	36	31	32	32	33
Polyurethane processing	kt	59	70	67	52	54	40	40	38	38	37
Production of adhesive tape	kt	NO									
PVC processing	kt	69	78	73	62	52	55	40	38	37	36
Rubber processing	tyres	70'000	70'000	72'500	75'000	77'500	80'000	80'000	81'000	82'000	83'000
Tanning of leather	employees	88	87	87	87	87	87	86	85	85	84

Degreasing and dry cleaning (2D3e NFR, 2D3f NFR)

Methodology

Source category 2D3e NFR comprises emissions from degreasing of electronic components, metal and other industrial cleaning. For the determination of NMVOC emissions from degreasing and dry cleaning a Tier 2 method according to the EMEP/EEA Guidebook 2013 is used based on the consumption of solvents. Switzerland's Informative Inventory Report 2017 contains a detailed description of the methods and country-specific data used for

estimating the NMVOC emissions from 2D3e NFR Degreasing and 2D3f NFR dry cleaning (FOEN 2017f).

Emission factors

Emission factors for NMVOC emissions from degreasing are based on data from the association of Swiss mechanical and electric engineering industries (swissmem) including VOC balance evaluations in 2004, 2007 and 2012 and expert estimates as documented in the EMIS database (EMIS 2017/2D3e NFR). For emission factors in 2015 see Table 4-37.

NMVOC emission factors for dry cleaning are estimated based on data and information from a survey of selected dry cleaning facilities that are representative for Swiss dry cleaning facilities and import statistics as documented in the EMIS database (EMIS 2017/2D3f NFR).

Activity data

Activity data of degreasing correspond to the annual consumption of solvents used for degreasing. Data are based on data from the association of Swiss mechanical and electric engineering industries (swissmem) in 2004, 2007 and 2012, VOC balances, import statistics and expert estimates, documented in the EMIS database (EMIS 2017/2D3e NFR)). A comparison between the surveys and the evaluations of VOC balances showed an underestimation of the survey data by about 6%. Thus, the emissions based on survey data from the industry association (swissmem) have been corrected by +10%. Activity data is provided in Table 4-38.

For dry cleaning, activity data is based on the amount of tetrachloroethylene (PER) and non-halogenated solvents imported and estimates of the share used for dry cleaning. Activity data for 2012 are based on the most recent survey at cantons and cleaning facilities as well as data from the Swiss supervising association of textile cleaning (VKTS). Activity data for 1990 are based on net imports of PER. For the years in between, data are interpolated linearly and after 2012, the activity data are assumed to remain constant, as documented in the EMIS database (EMIS 2017/2D3f NFR).

Chemical products, manufacture and processing (2D3g NFR)

Methodology

Based on the decision tree Fig. 3.1 in chapter 2D3g in EMEP/EEA (2013), for source category 2D3 Chemical products a Tier 2 method using country-specific emission factors is used for calculating the NMVOC emissions. Switzerland's Informative Inventory Report 2017 contains a detailed description of the methods and country-specific data used for estimating the NMVOC emissions from 2D3g NFR Chemical products, manufacture and processing (FOEN 2017f).

Emission factors

Emission factors for NMVOC are mainly provided by industry associations, i.e. for

- fine chemicals production, pharmaceutical production and handling and storing of solvents: Swiss business association for the chemical, pharmaceutical and biotech industry (scienceindustries)
- paint and ink production: Swiss association for coating and paint applications (VSLF) and the Swiss Organisation for the Solvent Recovery of Industrial Enterprises in the Packaging Sector (SOLV)
- polyurethane processing: Swiss plastics association
- polyester processing: Swiss polyester association
- tanning of leather: Swiss leather tanning association.

For the other processes in source category 2D3 (2D3g NFR) data are based on information from the industry and expert estimates as documented in the EMIS database.(EMIS 2017/2D3g). For emission factors see Table 4-37.

Activity data

The activity data are mainly production or consumption data provided by industry associations and by the Swiss Federal Office of Statistics, i.e. for

- fine chemicals production and handling and storing of solvents: Swiss Federal Office of Statistics
- pharmaceutical production: Swiss business association for the chemical, pharmaceutical and biotech industry (scienceindustries)
- paint and ink production: Swiss association for coating and paint applications (VSLF) and Swiss Organisation for the Solvent Recovery of Industrial Enterprises in the Packaging Sector (SOLV)
- polyurethane processing: Swiss plastics association
- polyester processing: Swiss polyester association
- tanning of leather: Swiss leather tanning association.

For the other processes in source category 2D3 (2D3g NFR) data are based on information of from the industry and expert estimates as documented in the EMIS database. Since 1994 no production of adhesive tape is occurring in Switzerland anymore.

For activity data see Table 4-38.

Road paving with asphalt (2D3b)

Methodology

Asphalt road surfaces are composed of compacted aggregate and asphalt binder. From road surfacing operations only NMVOC emissions occur. Based on the decision tree Fig. 3.1 in

chapter 2D3b in EMEP/EEA (2013), the NMVOC emissions from 2D3b Road paving with asphalt are determined by a Tier 2 method based on country-specific emission factors as documented in EMIS 2017/2D3b NFR.

Emission factors

The emission factor for NMVOC emissions from 2D3b Road paving with asphalt comprises NMVOC emissions from the use of prime coatings and from the bitumen content in asphalt products (about 5%). The NMVOC content in the bitumen has decreased considerably between 1990 and 2010. The values are based on industry data from 1990, 1998, 2007, 2010 and 2013. All other years are interpolated and complemented with expert estimates documented in the EMIS database.

Table 4-39 Emission factors of 2D3b Road paving with asphalt and 2D3c Asphalt roofing for 2015.

	Unit	CO	NMVOC
2D3b Road paving	kg/t asphalt concrete	NA	0.54
2D3c Asphalt roofing	kg/t asphalt sealing sheeting	124	21

Activity data

Activity data on the amount of asphalt products (so-called mixed goods) used for road paving is based on annual data from the association of asphalt production industry (SMI) for 1990 and from 1998 onwards and expert estimates for the years between.

Table 4-40 Activity data for road paving with asphalt, asphalt roofing and urea use in SCR catalysts.

	Unit	1990	1995	2000	2005
2D3b Road paving with asphalt					
Asphalt concrete	kt	5'500	4'800	5'170	4'780
2D3c Asphalt roofing					
Asphalt sealing sheeting	kt	50.0	44.6	41.4	30.1
2D3d Urea use in SCR catalysts					
AdBlue	kt	NO	NO	NO	0.3

	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2D3b Road paving with asphalt											
Asphalt concrete	kt	5'400	5'100	5'160	5'200	5'250	5'300	4'770	4'770	5'260	4'850
2D3c Asphalt roofing											
Asphalt sealing sheeting	kt	28.0	25.9	25.6	25.3	25.0	24.8	24.5	24.2	23.9	23.6
2D3d Urea use in SCR catalysts											
AdBlue	kt	2.6	5.6	10.0	14.1	17.0	18.9	20.8	22.7	23.5	23.5

Asphalt roofing (2D3c)

Methodology

This source category comprises emissions from production and use of asphalt roofing materials (saturated felt, roofing and siding shingles, roll roofing and sidings). These products are used in roofing and other building applications. From 2D3c Asphalt roofing only precursors such as CO and NMVOC arise. CO is emitted during the production process of asphalt roofing materials whereas NMVOC emissions are released during the entire production and laying processes (primers included). Based on the decision tree Fig. 3.1 in chapter 2D3c in EMEP/EEA (2013), the emissions of NMVOC, particulate matter and CO from Asphalt roofing are determined by a Tier 2 method based on country-specific emission factors as documented in the EMIS database (EMIS 2017/2D3c Dachpappen Produktion und Verlegung).

Emission factors

The emission factors from Asphalt roofing are based on information from the industry association, literature and expert estimates as documented in the EMIS database (see Table 4-39).

Activity data

Activity data is based on data from industry and expert estimates as documented in the EMIS database (see Table 4-40).

Urea use in SCR catalysts of diesel engines (2D3d)

This source category encompasses CO₂ emissions from the use of urea containing AdBlue in diesel engines with SCR-catalysts in road transportation (Euro V/VI and Euro 5/6).

Methodology

In accordance with the 2006 IPCC Guidelines the consumption of Ad Blue is reported in this submission following a methodology suggested in the EMEP/EEA guidebook 2013 (EMEP/EEA 2013; part B, chp. 1.A.3.b.i-iv, page 48). A specific percentage of the fuel consumption of SCR-vehicles in road transportation according to their Euro class is applied for Ad Blue consumption estimates. Emissions are calculated according to following formula:

$$\text{CO}_2 \text{ Emissions} = \text{EF} \cdot \text{FC} \cdot \text{Share of SCR vehicles mileage} \cdot \text{Specific urea share}$$

“FC” relates to the fuel consumption in tonnes of the entire vehicle category. “Share of SCR vehicles mileage” implies the mileage share of SCR-vehicles in the entire vehicle category and “Specific urea share” comprises the percentage of fuel consumption, which relates to AdBlue (urea solution) consumption.

Emission factors

The emission factor for CO₂ emissions from urea use in SCR-catalysts in vehicles is a default value (EMEP/EEA 2013) considering the molecular mass conversion of urea into CO₂ during the reaction with water and the content of 32.5% of the aqueous AdBlue urea solution.

Activity data

Activity data on AdBlue consumption as well as annual mileage are provided by INFRAS (INFRAS 2010) on a yearly basis as documented in EMIS 2017/2D3d NFR Urea (AdBlue) Einsatz Strassenverkehr. For activity data see Table 4-40.

4.5.3 Uncertainties and time-series consistency

The uncertainty of total CO₂ emissions from the entire source category 2D – Non-energy products from fuels and solvent use is estimated to be 50% (expert estimate).

Consistency: Time series for 2D Non-energy products from fuels and solvent use are all considered consistent.

4.5.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3 and partly also in chp. 3.2.4.8.

4.5.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 and 2014 of at least 0.3 kt CO₂ eq are quantified.

- 2D3b: The AD of 2D3b Road paving has been revised for 2014 based on corrected data from industry association.
- 2D3a: The survey on post-combustion of NMVOC emissions from coating applications, degreasing, dry-cleaning and manufacture and processing of chemical products has been revised resulting in changes of CO₂ emissions of 2D3a Post-combustion of NMVOC from solvent use for the entire time series. In 1990, this recalculation leads to a decrease in emissions by 1 kt CO₂ eq and in 2014, this recalculation leads to an increase by 1 kt CO₂ eq.

4.5.6 Category-specific planned improvements

No category-specific improvements are planned.

4.6 Source category 2E – Electronics industry

4.6.1 Source category description

Source category 2E Electronics industry is not a key category.

Source category 2E Electronics industry comprises HFC, PFC, NF₃ and SF₆ emissions from consumption of the applications listed below in Table 4-41.

Table 4-41 Specification of source category 2E Electronics industry in Switzerland.

2E	Source	Specification
2E1	Integrated circuit or semiconductor	Etching and cleaning processes in the production of IC and semiconductors (similar cleaning services for printed wiring boards included in the evaluation)
2E2	TFT flat panel display	No production of TFT flat panel displays in Switzerland, activities contained in the production of displays for watches
2E3	Photovoltaics	Emissions from photovoltaic manufacturing
2E4	Heat transfer fluids	No application in Switzerland assumed*
2E5	Other	Test activities (for example related to printed wiring boards), research activities

* Heat transfer fluids subject of research, for example ORC systems. Alternative products available with low GWP as for example Novec 649 and 7000

4.6.2 Methodological issues

Emission calculations are based on import data from FOEN statistics for etching and cleaning processes of the electronics industry, covering different source categories as listed in Table 4-41 (until 2010 import declarations for electronic industry under solvents). Process-specific transformation and emission rates are used. A survey within the electronics industry was carried out for the submission in 2015 to distribute the imported substances to the different source categories of electronic industry and to obtain information on waste air treatment. More information are available from Carbotech (2015).

Methodology

A Tier 2a approach with process gas-specific parameters is used for emission calculations. IPCC default values for the gas-specific transformation rate of different processes and general values for the exhaust treatment efficiency are applied.

Imports of electronics industry were included in FOEN statistics under solvents until 2010. For the inventory report 2011 (FOEN 2011) interviews were conducted with the industry to get in-depth information on allocation of imported PFC volumes to different applications and to obtain process-specific information from consumers. Until 2010, most PFC imports declared as 2F5 Solvents or 2F6 Other were related to the electronics industry (2E). Since 2011, PFC import declarations have been improved and information is provided for the source category 2E separately. A survey was carried out for the submission in 2015 to

determine contributions of different source categories 2E1–2E5 (Table 4-41). As a result, the peak of NF_3 imports (and corresponding emissions) between 2009 and 2011 was found to be related to photovoltaic manufacture.

Emission factors

Default emission factors according to the 2006 IPCC Guidelines are used for production and waste-air treatment. An exhaust treatment is assumed probable for most applications due to the Chemical Risk Reduction Ordinance (Swiss Confederation 2005), which limits emissions for industrial applications, such as semiconductor manufacturing, to 5% of the total substance used. For some large users the presence of exhaust treatment was confirmed in a survey.

Activity data

Activity data are based on FOEN import statistics and industry information.

4.6.3 Uncertainties and time-series consistency

The uncertainty for the emissions from the use of HFC, PFC, SF_6 and NF_3 in 2E Electronics industry is estimated at 51% (HFC), 81.5% (PFC), 61.5% (SF_6), 195% (NF_3) based on a Monte Carlo simulation.

Consistency: Time series for 2E Electronics industry are all considered consistent.

4.6.4 Category-specific QA/QC and verification

The entire time series are compared between the current and the previous submissions. The general QA/QC measures are described in chp. 1.2.3.

4.6.5 Category-specific recalculations

No category-specific recalculations were carried out, but all data on the recovery were replaced by the notation key "NA". The definition of recovery in the CRF tables and the use of recovery in the calculation of the implied emission factor are not coherent in the submission 2016 and led to false results for the implied emission factor.

4.6.6 Category-specific planned improvements

No category-specific improvements are planned.

4.7 Source category 2F – Product uses as substitutes for ozone depleting substances

4.7.1 Source Category Description

Table 4-42 Key categories (KCA incl. LULUCF) of 2F Product uses as substitutes for ozone depleting substances.

Code	IPCC Category	GHG	Identification Criteria
2F1	Refrigeration and air conditioning	HFC	L1, L2, T1, T2

Source category 2F Product uses as substitutes for ozone depleting substances comprises HFC and PFC emissions from consumption of the applications listed in Table 4-43.

Table 4-43 Specification of source category 2F Product uses as substitutes for ozone depleting substances in Switzerland.

2F	Source	Specification
2F1	Refrigeration and air conditioning	Emissions from refrigeration and air conditioning (incl. heat pumps and tumble dryers)
2F2	Foam blowing agents	Emissions from foam blowing, incl. polyurethane spray
2F4	Aerosols	Emissions from use as aerosols, incl. metered dose inhalers
2F5	Solvents	Emissions from use as solvents

The following graph shows HFC and PFC emissions from different applications in source category 2F. In 2015 refrigeration and air conditioning equipment account by far for the highest emissions with a share of 97% of the total emissions in source category 2F. Further, emissions are dominated by HFCs and only a minor contribution comes from PFCs (generally less than 1%).




Figure 4-6 Development of emissions under source category 2F Product uses as substitutes for ozone depleting substances (1990 to 2015). HFC and small amounts of PFC are used as substitutes for ozone depleting substances. Most relevant today are emissions from the built up refrigerant stock in equipment.

4.7.2 Methodological issues

The data models used for source category 2F are complex and therefore a comprehensive documentation of all relevant model parameters is not possible within the NIR. Most relevant is the contribution of 2F1 refrigeration and air conditioning. Calculations are carried out for different applications separately.

- 2F1a Stationary refrigeration and air conditioning
 - Domestic refrigeration
 - Commercial and industrial refrigeration
 - Stationary air conditioning, heat pumps and tumble dryers
- 2F1a Mobile refrigeration and air conditioning
 - Mobile air conditioning in different vehicle types
 - Transport refrigeration for different vehicle types

Annex A3.2 shows an illustrative example of the model structure and parameters used for calculating emissions from mobile air conditioning in cars. Where possible, the most important assumptions for the data model are documented in Table 4-44. More information of the individual data and models is available from Carbotech (2015) as well as related

background documents. This information is FOEN internal due to confidentiality of data, but is open for consultation by reviewers.

4.7.2.1 Refrigeration and air conditioning (2F1)

Methodology

The inventory under source category 2F1 includes different applications and equipment types. For each individual emission, models are used for calculating actual emissions as per the 2006 IPCC Guideline's Tier 2a approach (emission factor approach). In order to obtain the most reliable data for the calculations, two different approaches are applied to get the stock data needed for the model calculations. For the following applications a 'bottom up' approach is applied relying on statistics, product informations and expert estimations:

- Domestic refrigeration
- Mobile air conditioning for different vehicle types
- Transport refrigeration for different vehicles types
- Stationary air conditioning (direct and indirect systems)
- Heat pumps
- Tumble dryers

On the other hand, a 'top down' approach is applied for the calculation of the stock in commercial and industrial equipment, starting with the total imported amount of refrigerant. To determine the portion used for commercial and industrial refrigeration, the refrigerant consumption of other applications is subtracted from the import amount (consumption for the production and maintenance based on the bottom up calculations of stock as given in the example of mobile air conditioning in Annex A3.2). The evaluation of commercial and industrial refrigeration is carried out together, and the portion of industrial refrigeration is included under commercial refrigeration in the reporting tables.

The combination of 'bottom up' with 'top down' calculations leads to more comprehensive results than using just one approach. Noteworthy, in the hypothetical but possible case of incomplete 'bottom up' evaluations, remaining imported refrigerant would be attributed to the production and maintenance of industrial and commercial refrigeration equipment. This might be a reason why the resulting refrigerant stock of commercial and industrial refrigeration, which serves as the residual, tends to be higher than in neighbouring countries.

The import data as reported to FOEN are adjusted for imported substances to be used in Liechtenstein. This is to eliminate double counting with the inventory data of Liechtenstein. The split factor is based on the proportion of employees in the industrial and service sector (share of import for Liechtenstein <1%). The adjustment does not affect the bottom up calculations and leads to an adjustment of commercial and industrial refrigeration mainly.

Figure 4-7 shows the required data for the model calculation of refrigeration and air conditioning.




Figure 4-7 Required data for the model to calculate emissions from refrigeration and air conditioning in Switzerland.

In 2008, the revised Chemical Risk Reduction Ordinance (Swiss Confederation 2005) was introduced. As part of this revision, an obligation for operators handling equipment containing more than 3 kg of HFCs to provide information to FOEN on the date of operation start, type of equipment, type and amount of refrigerant and date of disposal was introduced. This data source provides valuable information and has been used to improve the estimates used for modelling emissions under source category 2F. However, it did not allow to directly draw the stock data or emission factors for the national inventory.

Emission factors

Emission factors for manufacturing, product life and disposal as well as average product lifetime are established on the basis of expert judgement and literature. Direct monitoring of the product life emission factors is only done at the company level for internal use and has been used partly for the verification of quality (confidential data from retailers and other type of industry). The product life factors and further parameters (i.e. re-filling frequency, handling losses and reuse of refrigerant) are used to allocate imported F-gases to new products and maintenance activities.

Table 4-44 displays the detailed model parameters used for the present submission. Changes of model parameters within the period 1990 to 2015 are indicated with values in brackets. The parameters in brackets are applied for the inventory 2015. For product life emission factors of some equipment types, a dynamic model which implies that emission

decrease linearly between 1995 and 2015 due to improved production technologies and the continuous sensitisation of service technicians is applied. The start/end values are based on expert statements (UBA 2005, UBA 2007, Schwarz 2001, Schwarz and Wartmann 2005). The charge at end of life for different applications has been analysed considering the technical minimal charge of equipment and the expected frequency of maintenance (UBA/Ökorecherche 2012). Disposal losses are calculated based on expert assumptions on the portion of broken equipment (100% loss) and on assumptions on disposal losses for professional recovery at site or waste treatment by specialized companies.

Table 4-44 Typical values of lifetime, charge and emission factors used in the model calculations for 1990 to 2015 for refrigeration and air conditioning equipment. Changes of model parameters within this time period are indicated with the starting year of the application in brackets (for example a charge of 4.7–7.5 kg was applied for heat pumps until 2000 and a lower charge of 2.8–4.5 kg from 2000 onwards. A linear interpolation is applied for the product life emission factor of commercial and industrial refrigeration, stationary air conditioning and for the emission factor of mobile air conditioning between the starting year and 2015.

Equipment type	Product life time	Initial charge of new product	Manufacturing emission factor	Product life emission factor	Charge at end of life *)	Export of retiring equipment **)	Disposal loss emission factor ***)
	[a]	[kg]	[% of initial charge]	[% per annum]	[% of initial charge of new product]	[% of retiring equipment]	[% of remaining charge]
Domestic refrigeration	16	0.1	NO	0.5	92	0-5 (2015:3)	19 ****)
Commercial and industrial refrigeration	10	NR	0.5	Sinking from 12 in 1990 to 7 in 2015	78-90	NE	19
Transport refrigeration: trucks/vans	10	1.8-7.8	1.5	15	86	90	28
Transport refrigeration: wagons	16	NR	NO	10	100	NE	28
Stationary air conditioning: direct cooling systems	15	NR	3 (2005: 1)	Sinking from 10 in 1995 to 4 in 2010	74-89	NE	28
Stationary air conditioning: indirect cooling systems	15	NR	1	Sinking from 6 in 1995 to 4 in 2010	85-89	NE	19
Stationary air conditioning: heat pumps	15	4.7-7.5 (2000: 2.8-4.5)	3 (2005: 1)	2	86	NE	19
Stationary air conditioning: tumble dryers	15	0.4	0.5	2	74	NE	19
Mobile air conditioning: cars	15	Sinking from 0.84 1990 to 0.55 in 2014	NO	8.5	58	31-72 (2015: 44)	50
Mobile air conditioning: truck/van cabins	12	1.1	NO	10 (2010: 8.5)	69-73	90 trucks (50 vans)	50
Mobile air conditioning: buses	12	7.5	NO	20 (2001: 15)	100	50	50
Mobile air conditioning: trains	16	20	NO	5.5	100	NE	20

*) Calculated value taking into account annual loss and portion refilled over the whole product life where applicable.

**) Allocation of disposal losses to export country

***) Calculated value taking into account share of total refrigerant loss and emission factor of professional disposal. Disposal losses of HFC and PFC occur from 2000 onwards (introduction of HFCs and PFCs starting 1991 and 10 to 16 years lifetime of equipment). The value of 50% for mobile air conditioning is based on UBA 2005 and expert assumptions on share of total refrigerant loss, e.g. due to road accident.

****) Takes into account HFC-134a content in foams, based on information from the recycling organisation SENS.

NR = Not relevant as only aggregate data is used

NO = Not occurring (only import of charged units)

NE = Not estimated

Activity data

Activity data are taken from industry information and national statistics such as for admission of new cars, buses, vans and trucks. Stock data is modelled dynamically. Due to the large number of sub-models used for modelling the total emissions for source category 2F1, no table on time series of activity data is provided here. For illustration, Annex A3.2 shows the detailed calculation model for car air conditioning including the time series for the activity data for this particular sub-model. Mobile air conditioning accounts for approx. 30% of the total emissions (CO₂ eq) of source category 2F1 Refrigeration and air conditioning in the inventory 2015.

For the inventory report 2012 (FOEN 2012) a cross check has been performed for results from model calculation and FOEN statistics on disposal and recycling of HFCs. This has indicated a significant gap with higher disposal values in model calculations compared to the FOEN disposal statistics. Some of the gap is explained by the onsite reuse and recycling of refrigerants, which is not reflected in the FOEN statistics and by other factors as e.g. the not accounted export of refrigeration equipment. Export rates used in model calculations are given in Table 4-44.

The registered refrigerant import is assumed to cover the consumption of Switzerland and Liechtenstein. To avoid double counting with the inventory data of Liechtenstein, the activity data for the equipment type commercial and industrial refrigeration is reduced by 0.9%, based on the share of imports of substances to be used in Liechtenstein. The reduction factor is based on the proportion of employees in the industrial and service sector in these two countries. For other equipment types no scope for double counting with the inventory of Liechtenstein was identified and therefore no correction factor is applied.

4.7.2.2 Foam blowing agents (2F2)

Methodology

In Switzerland no production of open cell foam based on HFCs is reported by the industry. Therefore, only closed cell PU and XPS foams, PU spray applications and further closed cell applications as sandwich elements are relevant under source category 2F2.

The emission model (Tier 2a) for foam blowing has been developed 'top down' based on import statistics for products, industry information and expert assumptions for market volumes and emission factors. Emissions from further not specified have been calculated (Tier 1a) as residual balance between FOEN import statistics and consumption in PU spray, PU and XPS foams.

Emission factors

For emission factors and lifetime of XPS and PU foam, expert estimates and default values according to the 2006 IPCC Guidelines (IPCC 2006, Volume 3, p. 7.37) are used. For PU spray, expert estimates and specific default values according to the 2006 IPCC Guidelines (IPCC 2006, Volume 3, p. 7.37) are used. Unknown applications are evaluated following the

Gamlen model recommended in the 2006 IPCC Guidelines (IPCC 2006). First-year losses are allocated to the country of production.

Table 4-45 Typical values on lifetime, charge and emission factors used in model calculations for foam blowing

Product	Product lifetime	Charge of new product	Manufacturing emission factor	Product life emission factor	Charge at end of life
Foam type	years	% of product weight	% of initial charge	% per annum	% charge of new product
PU foam	50	4.5	NR	NR	Calculated charge minus emissions over lifetime (so far not relevant, products still in use)
XPS foam HFC-134a	50	6.5	NR	NR / 0.7**	
XPS foam HFC-152a				100 / 0**	
PU spray all HFC	50	13.6 / 0 *	<1%	95 / 2.5 **	
Unknown use:					
HFC 134a, HFC 227ea, HFC 365 mfc	20	NR	10	10 / 4.5 **	
HFC 152a			100	100 / 0 **	

* The first value represents the charge of HFC 1995 (start of HFC use as substitutes for ozone depleting substances). The HFC amount was reduced continuously between 1995 and 2008. Since 2009 the production of PU spray is HFC free in Switzerland.

** Data for 1st year / following years (HFC-152a all emissions allocated to production)

NR Not relevant (PU foam: no substances according to this protocol have been used; XPS foam: emissions occur outside Switzerland; unknown use: calculations are based on the remaining propellant import amount).

Activity data

HFCs have been used until 2008 in the Swiss production of PU spray. The export rate of PU spray from Swiss production was about 96.5% of total production volume in the time period of HFC use. About one third of PU spray sold in Switzerland originates from local production, the rest is imported. For PU rigid foams no HFCs are used as foam blowing agent (only pentane and CO₂). There has been no production of XPS in Switzerland with HFCs. XPS foams were 100% imported until 2010. In 2011 a new production facility was started which, however, does not use HFCs. The HFC import not related to the main applications above has been allocated to further unknown applications (possible use in the production of sandwich elements mentioned by an import company of foam blowing agents has not been confirmed).

Detailed activity data for this source category are available from Carbotech (2015) as well as related background documents at FOEN, but not reported due to confidentiality.

4.7.2.3 Fire protection (2F3)

No emissions occur in source category 2F3 within Switzerland. The application of HFCs, PFCs and SF₆ in fire extinguishers is prohibited by law.

4.7.2.4 Aerosols (2F4)

Methodology

The Tier 2a emission model for Aerosol / MDI is based on a 'top-down' approach using import statistics for HFCs.

Emission factors

A manufacturing emission factor of 1% is applied. The model then assumes prompt emissions, i.e. 50% of the remaining substance is emitted in the first year and the rest in the second year, in line with the 2006 IPCC Guidelines.

Activity data

In most aerosol applications, HFCs have been replaced already in the past years. According to the information of companies filling aerosol bottles for use in households, e.g. cosmetics, cloth care and paint, no HFC is being used. For special technical applications – especially metered dose inhalers (MDI) – HFC is still in use. Compared to the total amount of aerosol applied, the HFC use for MDI is considered to be irrelevant.

Activity data are based on import statistics. The export and import of filled products is unknown, but assumed to be in a similar range.

4.7.2.5 Solvents (2F5)

Methodology

HFCs and PFCs are used as solvents. Emissions are calculated according to a Tier 1a method according to the 2006 IPCC Guidelines on basis of a 'top-down' approach using import statistics and industry information on allocation of the imported HFC and PFC amounts to different applications.

The import data as reported to FOEN cover imported substances to be used in Switzerland and Liechtenstein, and are therefore split in proportion of inhabitants of the two countries to avoid double counting.

Emission factors

In line with the 2006 IPCC Guidelines prompt emissions are assumed, i.e. half of the initial amount is emitted in the first year, the other half in the second year.

Activity data

Activity data are based on import statistics. Since the inventory report of the year 2011 (FOEN 2011), interviews were made with industry to get in-depth information on allocation of imported HFC and PFC volumes to different applications. These interviews revealed that most imported PFCs declared as Solvents (2F5) or Other (2F9) before 2011 are actually related to the electronics industry. Therefore, the model for allocation of imported PFC volumes was adjusted accordingly. Since 2011 imports for semiconductors manufacturing and further etching processes of electronics industry are registered as separate category in FOEN import statistics.

To avoid double counting with the inventory data of Liechtenstein, the import data reported to FOEN which is assigned to source category 2F5 in the inventory of Switzerland is reduced by 0.5%. The reduction factor is based on the proportion of inhabitants in these two countries.

4.7.2.6 Other applications (2F6)

There are no further applications of substitutes for ozone depleting substances in Switzerland.

4.7.3 Uncertainties and time-series consistency

For refrigeration equipment, air conditioning equipment as well as for foam blowing, a Monte Carlo analysis according to IPCC Good Practice Guidance for the evaluation of uncertainties of model calculations according to Tier 1 and 2 has been carried out. The Monte Carlo analysis was performed on the inventory data of the current GHG inventory (submission 2016). For the purpose of the Monte Carlo analysis, the uncertainty of all relevant parameters (e.g. initial appliance charge, product life emission factor, import and export volumes, etc.) used in the emission models for the applications as per Table 4-46 below has been characterised using the following statistical distributions:

- Triangular distribution (defined by the three parameters minimum, maximum and most likely value)
- Uniform distribution (same probability for the whole spectrum)
- Normal or lognormal distribution

The analysis was carried out with 10'000 cycles. Details on the distributions of parameters used (i.e. type of distribution, minimum, maximum, most likely value) are available from background documents at FOEN.

For the submission 2006 the uncertainty for the import statistic data had been estimated for the first time. Discussions with the persons responsible for data collection in the years 1997–2015 led to the estimations of standard deviation and minimal and maximal values given in Table 4-46. A normal distribution is used in the Monte Carlo analysis and the standard deviation, minimal and maximal values applied to define the probability ranges.

Table 4-46 Estimated uncertainty for the data of the imported substances.

Year	Std. Dev.	Minimal	Maximal	Remarks
Up to 1999	20%	15%	50%	Assumed that the data is not complete
2000 – 2003	20%	20%	20%	Data can be incomplete or possible double declaration
2004 – 2015	10%	20%	20%	Data can be incomplete or possible double declaration

Table 4-47 summarises the results for the application-specific emission models. The “value 2015” represents the reported emissions in kt CO₂ equivalent for the specific application for the year 2015. The uncertainty values stem from the Monte Carlo analysis. The uncertainty of the resulting total emissions from source category 2F Product uses as substitutes for ODS is about 20%. Higher values result for the contributions of single applications.

Uncertainties of more than 20% have been calculated for the following applications:

- Stationary air conditioning
- Commercial and industrial refrigeration
- Domestic refrigeration
- Foam blowing
- Aerosols
- Solvents

Uncertainties of 15% to 20% have been calculated for the following applications:

- Mobile air conditioning
- Transport refrigeration

For the model calculation of domestic refrigeration no uncertainty value is given due to a very asymmetric distribution. Calculation of refrigerant stocks is not reported in detail here because the uncertainties for stock and new filled refrigerant related to the split of refrigerant on different applications is of small importance for the overall emissions (different applications show similar characteristics for the building of stocks and related emissions). Detailed data are available from background documents at FOEN (excel calculations). For refrigerant stocks maximum uncertainties of over 30% result for HFC-134a in commercial and industrial refrigeration.

Relevant parameters for the building of stock in PU foam are the PU foam import and export rate of past years and the PU spray first year emission factor. The data base for PU sprays has been significantly improved with effect from the 2007 submission (FOEN 2007). This is attributed to improved models which have been elaborated by the main producer and its blowing agent import firm. However, the following three factors lead to a small amount remaining in the stock with a relative high uncertainty: high import and export rate of PU spray, lacking information on import of PU spray and on propellant used in import products and high emission factor of the first year.

Table 4-47 Summary of results for model parameter “emissions” from Monte Carlo analysis for 2015 data on selected emission sources.

Application	Model parameter	Value 2015 kt CO ₂ eq.	Average kt CO ₂ eq.	Median kt CO ₂ eq.	min. kt CO ₂ eq.	max. kt CO ₂ eq.	Uncertainty %
2F1 Refrigeration and air conditioning	Emissions in kt CO ₂ eq.	1'438	1'533	1'521	1'122	2'090	21
- Commercial / Industrial refrigeration		758	857	844	489	1'427	38
- Mobile air conditioning		435	411	408	301	597	17
- Stationary air conditioning		214	229	228	145	331	25
- Transport refrigeration		30.4	33.0	32.8	24.9	43.5	17
- Domestic refrigeration		0.4	3.1	0.9	0.1	12.9	*)
2F2 Foam blowing agents		32.6	50.2	48.9	26.2	212.2	61
2F4 Aerosols		14.1	14.6	14.6	5.7	24.2	54
2F5 Solvents		1.0	1.0	1.0	0.3	1.9	61
Total 2F Product use as substitutes for ODS		1'485	1'598	1'588	1'184	2'148	20

*) very asymmetric distribution, therefore standard deviation not indicated.

Consistency: Time series for 2F are all considered consistent.

4.7.4 Category-specific QA/QC and verification

The entire time series are compared between the current and the previous submission.

Recalculations were identified and explained. All modelling results produced by Carbotech (2015) have been checked in detail by FOEN specialists.

The assumptions of decreasing emission factors for the different equipment types under source category 2F1 Refrigeration and air conditioning have been cross-checked with the inventories of Austria and Germany and have been found to be in line with the assumptions made for these inventories.

The emission factor of category 2F used in the Swiss inventory was compared to the corresponding emission factors of other countries (UNFCCC: <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (INFRAS 2012). Concerning ODS substitutes the following sources of emissions are deemed most relevant: HFC-125, HFC-134a and HFC-143a from stationary and commercial refrigeration

as well as mobile air conditioning. The product life factor is relevant, since there is no production of halocarbons in Switzerland. For all these sources Switzerland's emission factors lie in the midfield of the range of other countries except for the life factor in mobile air conditioning and commercial and industrial refrigeration. However, when compared to neighbouring countries such as Germany, very similar values are used. The Swiss product life factors are often lower than the average for the following reasons. First, since 2005 the Chemical Risk Reduction Ordinance (Swiss Confederation 2005) is in place that ensures the proper handling and disposal of halocarbons and SF₆. Second, the decommissioning sector is well-organized by the SENS foundation and recycling is taxed in advance. Third, servicing staff is well-trained to proper handling and disposal of respective appliances. And fourth, the good economic conditions allow a higher renewal frequency and higher equipment standard.

The FOEN supports a monitoring campaign at the high-altitude research station Jungfrauoch, where various greenhouse gases are measured continuously. The location of the research station normally provides for analysis of tropospheric background concentrations. However, under special meteorological conditions, an estimate of Swiss emissions can be derived from the measurements. For HFC-134a, HFC-125, HFC-152a, HFC-143a, HFC-336mfc and HFC-32 a comparison of the inventory data with the inferred emissions is presented in Annex A5.1. Estimated emissions based on measurements at Jungfrauoch agree fairly well with the emission estimates HFC-134a, HFC-125, HFC-143 and HFC-32 of the Swiss greenhouse gas inventory. Larger differences result for less relevant contributions of HFC-152a. The allocations of first year emissions of foam blowing agents to the country of production might be the reason for the observed differences.

4.7.5 Category-specific recalculations

Recalculations reported in submission 2017

- 2F1: Product life emission factor of commercial and industrial refrigeration is now assumed to decrease slower, i.e. linearly from 12% in 1995 to 7% in 2015.
- 2F1: The correction factor for refrigerant import to avoid double countings with Liechtenstein has been applied for the whole time period (so far only considered for the time period 2007 to 2014).
- 2F5: The correction factor for solvents import to avoid double countings with Liechtenstein has been applied for the whole time period (so far only considered for the time period 2007 to 2014).

4.7.6 Category-specific planned improvements

Gradual improvement of the data quality in co-operation with industry is ongoing. As in the past years, methodologies and emission models will be updated during the yearly process of F-gas inquiry. The focus will be on:

- Split of commercial and industrial refrigeration.
- Improvements of HFC emission calculations from refrigeration and air conditioning equipment.

- Changes are expected and will be analysed in this area due to the revision of the Chemical Risk Reduction Ordinance and CO₂ compensation programmes (share of products with HFC, recycling of HFC, early replacement of HFC).

4.8 Source category 2G – Other product manufacture and use

4.8.1 Source category description

Table 4-48 Key categories (KCA incl. LULUCF) of 2G Other product manufacture and use.

Code	IPCC Category	GHG	Identification Criteria
2G	Other Product Manufacture and Use	N2O	T2
2G	Other Product Manufacture and Use	SF6	L1, T1

Table 4-49 Specification of source category 2G Other product manufacture and use in Switzerland.

2G	Source	Specification
2G1	Electrical equipment	Emissions of SF ₆ from use in electrical equipment
2G2	SF ₆ and PFCs from other product use	Emissions of SF ₆ and PFC not accounted in other source categories (i.e. for particle accelerators, soundproof windows, leakage detection, research and laboratory use)
2G3	N ₂ O from product uses	Emissions of N ₂ O from the use of N ₂ O in hospitals; Emissions of N ₂ O from the use of aerosol cans
2G4	Other	Emissions of NMVOC from domestic solvent use, printing, other solvent and product use as well as emissions of CO ₂ resulting from post-combustion of NMVOC in exhaust gases of these sources Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ as well as CO from use of fireworks and tobacco, respectively; Emissions of HFC not accounted in other source categories

4.8.2 Methodological issues

4.8.2.1 Electrical equipment (2G1)

Methodology

Under an agreement with FOEN, the industry association SWISSMEM is reporting actual emissions of SF₆ on basis of a mass-balance approach (Tier 3a). The mass balance includes mainly data for the production, installation, operation and disposal of electrical equipment, but included in past years also small amounts of SF₆ for other applications (i.e. research, magnesium foundry). SWISSMEM is collecting data from its members and is crosschecking the reported SF₆ consumption data with data from importers of SF₆. Installations in operation with electrical equipment containing SF₆ are periodically inspected for leakage, and losses are refilled (topping up). The refilled quantities and any SF₆ charge required during repair are reported as emissions at the time of filling. A product lifetime of 35 years is assumed.

Emission factors

Emission factors for source category 2G1 are based on industry information and are calculated values based on the mass-balance data. The discontinuity in emission factor from 2005 to 2006 data is due to the inspection intervals, optimised data collection and technical optimisation of equipment. The trend for reduced emission factors can be linked to the existing agreement of SWISSMEM and FOEN on the reduction of SF₆ emissions.

Activity data

Activity data are based on industry information. The wide annual fluctuation of SF₆ emissions from electrical equipment is related to the annual fluctuation of market volumes for such equipment as well as variations in inspection intervals and equipment break-down requiring topping up of SF₆ charge in the equipment. Import declarations obtained for FOEN import statistics are cross-checked regularly in order to eliminate double counting between SWISSMEM data and other import declarations.

4.8.2.2 SF₆ and PFCs from other product use (2G2)

Methodology

The emissions reported under 2G2 are related to the use of SF₆ for industrial particle accelerators (2G2b), the use of SF₆ for soundproof windows (2G2c) and other PFC and SF₆ use (2G2e). 2G2e summarizes research/analytics and further applications (including the unallocated difference in SF₆ emissions based on the FOEN import statistics and the SWISSMEM mass balance).

Under an agreement with FOEN, the industry association SWISSMEM is reporting actual emissions of SF₆ from industrial particle accelerators on the basis of a mass-balance approach (Tier 3a).

For 2G2c soundproof windows and 2G2e Other a Tier 2 approach is applied. Therefore, the unallocated amount of SF₆ under 2G2e has been assigned as application of cables and electrical control systems. Further evaluations of applications under 2G2e are based on FOEN import statistics and industry data, including applications with direct emissions and applications with banks. No further details are provided due to confidentiality. Data are available from Carbotech (2015).

Emission factors

For the unallocated amount of SF₆ assigned to cables and electrical control systems, the emission factor is assumed to be 4% for manufacturing and 1% per year during the product life. 100% of the remaining charge is emitted at the time of disposal after a lifetime of 40 years. Because of the long lifetime, the disposal emissions are not relevant for the results.

For soundproof windows an emission rate of 1% per year is assumed, including the portion of broken windows. Since 2008, there is no production of windows with SF₆ in Switzerland. For the manufacturing in former years the emission factor is assumed to be 33%.

Activity data

Activity data are based on import statistics and industry information. For the unallocated amount of SF₆ assigned to cables and electrical control systems an export rate of 80% is assumed similar to electrical equipment 2G1. Also for the inventory report 2013 (FOEN 2015), the split factors for allocation of imported amounts to different applications was checked through industry interviews and in-depth analysis in order to eliminate double counting between SWISSMEM data and other import declarations. Interviews with industry were carried out for the present inventory to identify applications of substances related to research under source category 2G2e Other.

4.8.2.3 N₂O from product uses (2G3)

Methodology

Emissions of N₂O from the source category 2G3 occur from the anaesthesia use in hospitals (2G3a Medical applications) and from the use of aerosol cans in households (2G3b Other). For both categories a Tier 2 method based on the production/consumption of N₂O is used (IPCC 2006 (vol. 3 chp. 8.4)).

Emission factors

For source category 2G3a Medical applications the emission factor is calculated based on the amount of N₂O sold in Switzerland divided by the number of inhabitants. The amount of N₂O sold for anaesthesia purpose is derived from sales data from the companies concerned based on annual data from 2005–2011 (EMIS 2017/2G3a Lachgasanwendung Spitäler).

Source category 2G3b Other includes N₂O emissions from whipped-cream makers using gas capsules for private households and restaurants. The emission factor is calculated based on sales data and N₂O content of gas capsules sold in Switzerland divided by the number of inhabitants (EMIS 2017/2G3b Lachgasanwendung Haushalt).

Table 4-50 N₂O emission factors for the source categories 2G3a Medical applications and 2G3b Other in 2015.

2G3a Use of N ₂ O for anaesthesia	Unit	1990	1995	2000	2005
N ₂ O	g/inhabitant	43	29.8	16.6	12.0
2G3b N ₂ O from aerosol cans					
N ₂ O	g/inhabitant	9.3	9.6	9.8	10.5

2G3a Use of N ₂ O for anaesthesia	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
N ₂ O	g/inhabitant	10.0	9.0	8.0	7.0	7.0	6.0	5.7	5.5	5.2	5.0
2G3b N ₂ O from aerosol cans											
N ₂ O	g/inhabitant	10.8	11.0	11.3	11.5	11.8	12.0	12.2	12.4	12.7	12.9

Activity data

For the source categories 2G3a Medical applications and 2G3b Other the activity data correspond to the Swiss population (SFSO 2016a) (EMIS 2017/2G3a Lachgasanwendung Spitaler and EMIS 2017/2G3b Lachgasanwendung Haushalt).

Table 4-51 Activity data for the source categories 2G3a Use of N₂O for anaesthesia and 2G3b N₂O from aerosol cans.

2G3 N ₂ O from product uses	Unit	1990	1995	2000	2005
2G3a, 2G3b	inhabitants	6'796'000	7'081'000	7'209'000	7'501'000

2G3 N ₂ O from product uses	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2G3a, 2G3b	inhabitants	7'558'000	7'619'000	7'711'000	7'801'000	7'878'000	7'912'000	7'997'000	8'089'000	8'189'000	8'282'000

4.8.2.4 Other (2G4)

Since the 2006 IPCC Guidelines (vol. 3, chp. 5.5) refer to the EMEP/EEA Guidebook 2013 regarding methodologies for estimating NMVOC emissions from solvent use, the respective NFR codes are indicated as reference as well. In the following sections the NMVOC emissions from domestic solvent use (2D3a NFR), printing (2D3h NFR), other solvent use (2D3i NFR) as well as other product use (2G NFR) are reported. From other product use (2G NFR) also CO₂, NO_x, CO and SO₂ as well as CO from the use of fireworks and tobacco, respectively, are emitted.

Due to the obligations of the Ordinance on Air Pollution Control (Swiss Confederation 1985) and Ordinance on the Incentive Tax on Volatile Organic Compounds (Swiss Confederation 1997) several industrial plants use facilities and equipment to reduce NMVOC in exhaust gases and room ventilation output. Often this implies the feeding of air with high NMVOC content into the burning chamber of boilers or other facilities to incinerate NMVOC. These CO₂ emissions from post-combustion of NMVOC are estimated based on industry data and expert estimates (Carbotech 2016a).

Switzerland's Informative Inventory Report 2017 contains a detailed description of the methods and country-specific data used for estimating the NMVOC emissions from the most important sources within source category 2G4 Other (FOEN 2017f).

Domestic solvent use (2D3a NFR)

Methodology

The source category 2G4 Domestic solvent use (2D3a NFR) Domestic solvent use including fungicides comprises mainly the use of cleaning agents and solvents in private households for building and furniture cleaning and cosmetics and toiletries but also the use of spray cans and pharmaceuticals. These products contain solvents, which evaporate during use or after the application. Among the numerous NMVOC emission sources, the use of household cleaning agents is the largest single source in source category 2D3.

Based on the decision tree Fig. 3.1 in chapter 2D3a in EMEP/EEA (2013), the emissions are calculated by a Tier 2 method (EMIS 2017/2D3a) using country-specific emission factors. All emissions related to domestic solvent use are calculated proportional to the Swiss population.

Emission factors

- *Household cleaning agents:* The source category 2D3a Use of cleaning agents includes the use of cosmetics, toiletries, cleaning agents and care products. Its resulting emission factor bases thus on a multitude of products, their NMVOC contents, emission fractions and consumption numbers. About 80% of the NMVOC emissions stem from the use of cosmetics and toiletries whereas the rest arises from the use of cleaning agents and care products. Available data sources consist of surveys of the use of household cleaning agents, cosmetics and toiletries in Switzerland (1990) and information from the Swiss association of cosmetics and detergents (SKW 2010) as well as surveys from Germany (1998, 2005). From 2001 until 2010 a constant EF is assumed for domestic use of cleaning agents. The value is based both on information from the Swiss association of cosmetics and detergents (SKW 2010) and from a German study on NMVOC emissions from solvent use and abatement possibilities by Theloke J. (2005). There were no significant improvements in the solvent compositions of the employed detergents. In a study conducted in 2013/2014 in Switzerland more accurate data of household cleaning agents, cosmetics and toiletries was collected based on comprehensive surveys at retailers, producers, industry associations and experts as well as analysis of import statistics (Hubschmid 2014). As a result of this study, the emission factor of household cleaning agents was adjusted in 2013. The study indicates again an increase in the NMVOC emission factor in 2013.
- *Domestic use of spray cans:* Emission factors of domestic use of spray cans are based on surveys in Switzerland (1990) and information from the Swiss Association for Aerosols (ASA) for years 1998, 2001, 2007 and 2010 and from a German study on NMVOC emissions from solvent use and abatement possibilities by Theloke J. (2005). In a study conducted in 2013/2014 in Switzerland more accurate data of domestic spray cans were collected based on comprehensive surveys at retailers, producers, industry associations and experts as well as analysis of import statistics (Hubschmid 2014). As a result of this study, the emission factor of spray cans was adjusted in 2013. The study indicates again an increase in the NMVOC emission factor in 2013.
- *Domestic use of pharmaceutical products:* Emission factors of domestic use of pharmaceutical products are available from surveys in Switzerland (1990) and Germany (1998) and from the Swiss business association for the chemical, pharmaceutical and biotech industry (scienceindustries) for 2011, as documented in the EMIS database. For years with no survey data, emission factors are interpolated.

Table 4-52 Emission factors for NMVOC for source category 2G4 Other: domestic solvent use, printing, other solvent use, other product use in 2015.

2G4 Other	Unit	NMVOC
Domestic solvent use (2D3a NFR)		
Domestic use of spray cans	g/inhabitant	329
Domestic use of pharmaceuticals	g/inhabitant	30
Household cleaning agents	g/inhabitant	989
Printing (2D3h NFR)		
Printing	kg/t ink	280
Package printing	kg/t ink	181
Other solvent use (2D3i NFR)		
Production of cosmetics	kg/employee	64
Production of paper and paperboard	g/t	35
Production of perfume and flavour	kg/employee	38
Production of textiles	kg/employee	8
Production of tobacco	kg/employee	12
Removal of paint and lacquer	g/inhabitant	34
Scientific laboratories	kg/employee	15
Other product use (2G NFR)		
Application of glues and adhesives	kg/t solvent	732
Commercial & industrial use of cleaning agents	g/employee	454
Cosmetic institutions	kg/employee	28
De-icing of airplanes	kg/t de-icing agent	280
Glass wool enduction	g/t glass wool	190
Hairdressers	kg/employee	14
Health care, other	kg/employee	8
Medical practices	kg/employee	8
Preservation of wood	kg/t preservative	110
Rock wool enduction	g/t rock wool	380
Underseal treatment & conservation of vehicles	kg/t underseal agent	400
Use of concrete additives	g/t additive	740
Use of cooling lubricants	kg/t lubricant	6
Use of lubricants	kg/t lubricant	340
Use of pesticides	kg/t pesticide	33
Use of tobacco	kg/Mio cigarette eq.	9

Activity data

As described in the methodology chapter, the activity data used for calculating the NMVOC emissions from domestic solvent use corresponds to the Swiss population (SFSO 2016a), see Table 4-53.

Table 4-53 Activity data for source category 2G4 domestic solvent use, printing, other solvent use, other product use.

2G4 Other	Unit	1990	1995	2000	2005
Domestic solvent use (2D3a NFR)	inhabitants	6'796'000	7'081'000	7'209'000	7'501'000
Printing (2D3h NFR)					
Printing	kt ink	13	13	14	5.5
Package printing	kt ink	5.9	5.9	5.5	9.1
Other solvent use (2D3i NFR)					
Fat, edible & non-edible oil extraction	kt	40	38	12	NO
Production of cosmetics	employees	2'200	2'200	2'267	2'100
Production of paper and paperboard	kt	1'510	1'560	1'780	1'750
Production of perfume and flavour	employees	2'200	2'325	2'567	3'200
Production of textiles	employees	25'200	26'763	24'300	17'067
Production of tobacco	employees	3'300	2'988	2'729	2'710
Removal of paint and lacquer	inhabitants	6'796'000	7'081'000	7'209'000	7'501'000
Scientific laboratories	employees	10'194	18'604	23'217	23'000
Vehicles dewaxing	vehicles	200'000	166'250	72'667	NO
Other product use (2G NFR)					
Application of glues and adhesives	kt	4.0	3.0	2.0	1.5
Commercial & industrial use of cleaning agents	employees	3'950'000	3'867'500	3'954'667	4'133'667
Cosmetic institutions	employees	2'600	3'100	3'533	3'800
De-icing of airplanes	kt	1.3	1.3	1.3	2.5
Fireworks	kt	0.8	1.0	1.5	1.4
Glass wool enduction	kt	24	24	31	37
Hairdressers	employees	20'553	22'826	23'530	22'200
Health care, other	employees	113'000	129'250	145'667	161'667
Medical practices	employees	27'625	42'047	50'833	55'357
Preservation of wood	kt paint	6.0	7.9	8.7	7.2
Rock wool enduction	kt	38	40	51	46
Underseal treatment & conservation of vehicles	kt	0.06	0.06	0.08	0.12
Use of concrete additives	kt	24	25	29	36
Use of cooling lubricants	kt	5.0	5.2	5.8	6.4
Use of lubricants	kt	1.3	1.3	1.3	3.7
Use of pesticides	kt	2.4	2.0	1.7	1.5
Use of tobacco	Mio cigarette eq.	16'192	15'774	14'751	13'369

2G4 Other	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Domestic solvent use (2D3a NFR)	inhabitants	7'558'000	7'619'000	7'711'000	7'801'000	7'878'000	7'912'000	7'997'000	8'089'000	8'189'000	8'282'000
Printing (2D3h NFR)											
Printing	kt ink	4.0	2.6	2.5	2.5	2.4	2.2	2.0	1.9	1.9	2.0
Package printing	kt ink	11	13	13	13	13	13	13	13	13	13
Other solvent use (2D3i NFR)											
Fat, edible & non-edible oil extraction	kt	NO									
Production of cosmetics	employees	2'100	2'100	2'100	2'100	2'100	2'100	2'100	2'100	2'100	2'100
Production of paper and paperboard	kt	1'690	1'734	1'700	1'540	1'540	1'380	1'390	1'400	1'410	1'420
Production of perfume and flavour	employees	3'300	3'400	3'425	3'450	3'475	3'500	3'521	3'542	3'563	3'583
Production of textiles	employees	16'733	16'400	16'200	14'200	13'800	14'800	14'789	14'778	14'767	14'756
Production of tobacco	employees	2'705	2'700	2'825	2'950	3'075	3'200	3'200	3'200	3'200	3'200
Removal of paint and lacquer	inhabitants	7'558'000	7'619'000	7'711'000	7'801'000	7'878'000	7'912'000	7'997'000	8'089'000	8'189'000	8'282'000
Scientific laboratories	employees	23'000	23'000	23'000	23'000	23'000	23'000	23'083	23'167	23'250	23'333
Vehicles dewaxing	vehicles	NO									
Other product use (2G NFR)											
Application of glues and adhesives	kt	1.4	1.3	1.2	1.2	1.1	1.0	1.9	1.0	1.0	1.0
Commercial & industrial use of cleaning agents	employees	4'208'333	4'283'000	4'323'333	4'363'667	4'404'000	4'333'333	4'262'667	4'192'000	4'236'000	4'280'000
Cosmetic institutions	employees	4'000	4'200	4'400	4'600	4'800	5'000	5'111	5'222	5'333	5'444
De-icing of airplanes	kt	2.1	1.7	2.4	3.1	3.7	4.4	5.1	5.1	5.1	5.1
Fireworks	kt	1.5	1.7	2.0	2.0	1.7	2.0	1.9	2.3	1.8	1.6
Glass wool enduction	kt	38	44	44	33	36	41	39	33	32	31
Hairdressers	employees	22'200	22'200	23'000	23'000	23'000	23'000	23'000	23'000	23'000	23'000
Health care, other	employees	163'333	165'000	163'000	163'000	163'000	163'000	163'000	163'000	163'000	163'000
Medical practices	employees	56'471	57'586	58'700	58'700	58'700	58'700	58'700	58'700	58'700	58'700
Preservation of wood	kt paint	7.0	6.9	6.1	5.3	4.5	3.6	2.8	2.0	2.0	2.0
Rock wool enduction	kt	53	63	58	53	56	57	57	54	53	47
Underseal treatment & conservation of vehicles	kt	0.13	0.13	0.14	0.15	0.16	0.17	0.18	0.18	0.18	0.18
Use of concrete additives	kt	35	34	34	34	41	44	38	38	38	38
Use of cooling lubricants	kt	5.7	4.9	4.2	2.8	3.6	4.0	3.7	3.7	3.7	3.7
Use of lubricants	kt	2.6	1.5	0.4	0.3	0.4	0.5	0.4	0.4	0.4	0.4
Use of pesticides	kt	1.5	1.5	1.7	1.9	2.0	2.2	2.2	2.2	2.2	2.2
Use of tobacco	Mio cigarette eq.	13'808	13'072	13'310	13'667	12'443	11'856	12'705	12'162	10'628	10'284

Printing (2D3h NFR)

Methodology

The source category 2G4 Printing (2D3h NFR) has been split into 2G4 Other printing industry and 2G4 Package printing. A Tier 2 method according to the EMEP/EEA Guidebook 2013 is applied using country-specific emission factor for calculating the NMVOC emissions from ink applications.

Emission factors

Emission factors for NMVOC are based on data from industry associations (Swiss Organisation for the Solvent Recovery of Industrial Enterprises in the Packaging Sector (SOLV)), VOC balances in the printing industry, German studies on NMVOC emissions from solvent use (Theloke 2005) and expert estimates, as documented in the EMIS database (EMIS 2017/2G4). For emission factors of 2015 see Table 4-52.

Activity data

The activity data used for calculating the NMVOC emissions correspond to the annual consumption of printing ink. This data stem from industry associations (SOLV), VOC balances in the printing industry and expert estimates, documented in the EMIS database (EMIS 2017/2G4). For activity data see Table 4-53.

Other solvent use (2D3i NFR)

Methodology

Source category 2G4 Other solvent use (NFR 2D3i) consists of a number of solvent uses in various production processes and services. Based on the decision tree Fig. 3.1 in chapter 2D3i in EMEP/EEA (2013), a Tier 2 method using country-specific emission factors is applied for calculating the NMVOC emissions from the different solvent applications in source category 2D3i Other solvent use (EMIS 2017/2G4). For the source category 2G4 Other solvent use (NFR 2D3i) Not-attributable solvent emissions so-called direct emission data is available only.

Emission factors

Emission factors for NMVOC are country-specific based on data from industry and services, industry associations, German studies on NMVOC emissions from solvent use (Theloke et al. 2000 and Theloke 2005) and expert estimates, as documented in the EMIS database (EMIS 2017/2G4). For emission factors see Table 4-52.

Activity data

For the majority of production processes and services – such as production of perfume and flavour and production of textiles – the activity data correspond to the respective number of employees (SFSO 2016e). The quantity of NMVOC emission per employee originates from the bottom-up approach in these industrial sectors and the decentralized political structure in Switzerland. The determined NMVOC emissions of representative production sites or service institutions are referred to the number of employees in order to calculate the Swiss total.

For production of paper and paperboard and fat, edible and non-edible oil extraction, the activity data are based on production volumes. Annual production volumes of paper and paperboard are provided by the Swiss association of pulp, paper and paperboard industry

(ZPK). For the removal of paint and lacquer the activity data correspond to the number of inhabitants (SFSO 2016a).

For activity data see Table 4-53.

Other product use (2G NFR)

Methodology

Within source category 2G Other product use, the major NMVOC emission sources in 2015 are 2G Commercial and industrial use of cleaning agents, 2G De-icing of airplanes and 2G Health care, other.

Based on the decision tree Fig. 3.1 in chapter 2G in EMEP/EEA (2013), for source category 2G Other product use Tier 2 methods using country-specific emission factors are applied for calculating the emissions from the different product applications and the use of fireworks and tobacco (EMIS 2017/2G).

For the source categories 2G Renovation of corrosion inhibiting coatings and 2G Use of aerosol cans in commerce and industry so-called direct emission data is available only.

Indirect CO₂ emissions from CO and NMVOC emissions from tobacco use are of biogenic origin and are therefore not reported in chp. 9 Indirect CO₂ and N₂O emissions.

Emission factors

Emission factors for NMVOC are based on data from industry and services, industry associations, German studies on NMVOC emissions from solvent use (Theloke et al. 2000 and Theloke 2005) and expert estimates, as documented in the EMIS database (EMIS 2017/2G4). For emission factors see Table 4-52.

Emission factors of CO₂, NO_x, CO and SO₂ as well as CO from use of fireworks and tobacco (EMIS 2017/2G), respectively, are displayed in the following table. Emission factors of fireworks are documented in FOEN (2014p).

Table 4-54 Emission factors for CO₂, NO_x, CO, SO₂ for source category 2G4 Fireworks in 2015

2G4 Other	Unit	CO ₂	NO _x	CO	SO ₂
Other product use (2G NFR)					
Fireworks	kg/t	43	0.26	7.4	4.1
Use of tobacco	kg/Mio cigarette eq.	1'000	NA	80	NA

Activity data

For the production processes, such as enduction of glass and rock wool and part of the applications in services, such as preservation of wood and application of glues and adhesives the activity data are based on production volume or employed agents. For the other part of applications in services, such as house cleaning in services, commerce and

industry and medical practices the activity data correspond to the respective number of employees. The quantity of NMVOC emission per employee originates from the bottom-up approach in these service sectors and the decentralized political structure in Switzerland. The determined NMVOC emissions of representative production sites or service institutions are referenced to the number of employees in order to calculate the Swiss total.

The activity data stem from industry, services, industry associations, Swiss federal statistical office and expert estimates and are documented in the EMIS database. Activity data for annual tobacco consumption and the annual firework sales are provided by the Swiss addiction prevention foundation ("Sucht Schweiz") and the statistics of the Swiss federal office for police (FEDPOL 2016), respectively.

For activity data see Table 4-53.

HFC not accounted in other source categories

Emissions of HFC not accounted for in any other source categories are reported under 2G4 Other. For confidentiality reasons, no further details are provided. Information is available from FOEN (confidential report Carbotech, 2015).

Methodology

A Tier 2 approach is applied for HFCs with prompt emissive applications based on import statistics and industry data.

Emission factors

Prompt emissions of HFC are calculated following the 2006 IPCC Guidelines assuming a total loss of product within two years (50% loss in the the first and 50% in the second year).

Activity data

HFC activity data under 2G4 are based on FOEN import statistic and industry data.

4.8.3 Uncertainties and time-series consistency

The uncertainty of total CO₂ emissions from the entire source category 2G is estimated at 50% (expert estimate).

The uncertainty of N₂O emissions from source category 2G3, which is a key category for this gas regarding trend according to Approach 2, is estimated at 80% (expert estimate, see Table 1-10).

The uncertainty of SF₆, HFC and PFC emissions in source category 2G is estimated at 24%, 12% and 15% respectively based on a Monte Carlo analysis (further details available from

background documents). Source category 2G is a key category for SF₆ regarding level and trend according to Approach 1.

Time series is consistent, with exception of the source category 2G2 Electrical equipment where from 2000 onwards the data are based on a Tier 3a approach instead of model calculations according to Tier 2 as applied for data before 2000. Due to lack of basic information it is not possible to provide a consistent time series for category 2G2 Electrical equipment retroactively.

4.8.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

4.8.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 and 2014 of at least 0.3 kt CO₂ eq are quantified.

- 2G: Unfortunately there is a double-counting in the NMVOC emissions from 2G De-icing of airplanes for the years 1990–2006. This mistake will be corrected in submission 2018.
- 2G3a: The N₂O emission factor for 2020 of 2G3a Use of N₂O in hospitals has been adapted to the current scenario of Swiss population development yielding revised interpolated values for 2012–2014.
- 2G4: The survey on post-combustion of NMVOC emissions from printing and other solvent and product use has been revised resulting in changes of CO₂ emissions of 2G4 Post-combustion of NMVOC from other solvent and product use for the entire time series. In 2014, this recalculation leads to an increase in emissions by 1 kt CO₂ eq.

4.8.6 Category-specific planned improvements

No category-specific improvements are planned.

4.9 Source category 2H – Other

4.9.1 Source category description

Source category 2H Other is not a key category.

Table 4-55 Specification of source category 2H Other in Switzerland.

2H	Source	Specification
2H1	Pulp and paper	Emissions from NMVOC from pulp and paper including chipboard, fibreboard and cellulose production (ceased in 2008)
2H2	Food and beverages industry	Emissions of CO and NMVOC from production of food and drink
2H3	Other	Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting and shooting;

4.9.2 Methodological Issues

4.9.2.1 Pulp and paper (2H1)

Methodology

In 2014, the production of chipboard and fibreboard are the relevant industrial processes in the source category 2H1 Pulp and paper. In Switzerland, chipboard and fibreboard are produced in one and two plants, respectively. The cellulose production was closed down in 2008 and is not occurring anymore in Switzerland. The NMVOC emissions are calculated by a Tier 2 method according to EMEP/EEA (2013) using country-specific emission factors.

Emission factors

The emission factor for NMVOC emissions from pulp and paper production in Switzerland is country-specific and based on measurements and data from industry and expert estimates documented in EMIS 2017/2H1. The implied emission factor given in Table 4-48 is production-weighted and related to chipboard and fibreboard production.

Table 4-56 Emission factors for CO and NMVOC in pulp and paper production and food and beverages industry, CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting for 2015.

2H Other	Unit	CO ₂	NO _x	CO	NMVOC	SO ₂
2H1 Pulp and paper	g/t	NA	NA	NA	550	NA
2H2 Food and beverage industry (exc. beer, wine, spirits)	g/t	NA	NA	250	3'480	NA
2H2 Food and beverage industry (beer, wine, spirits)	g/m ³	NA	NA	NA	360	NA
2H3 Blasting and shooting	kg/t	400	35	310	60	0.5

Activity data

The annual amount of pulp and paper produced in Switzerland is based on data from industry and expert estimates documented in EMIS 2017/2H1.

Table 4-57 Pulp and paper production, food and beverages production, amount of used explosives and processed crude oil in Switzerland.

2H Other	Unit	1990	1995	2000	2005
2H1 Pulp and paper	kt	604	593	641	693
2H2 Food and beverage industry (exc. beer, wine, spirits)	kt	2'254	2'116	2'301	2'138
2H2 Food and beverage industry (beer, wine, spirits)	m3	560'972	516'519	492'208	452'877
2H3 Blasting and shooting; blasting agent and powder	kt	2.6	1.3	1.9	0.8

2H Other	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
2H1 Pulp and paper	kt	731	790	765	544	602	564	533	510	516	519
2H2 Food and beverage industry (exc. beer, wine, spirits)	kt	2'167	2'343	2'428	2'437	2'400	2'487	2'422	2'378	2'540	2'387
2H2 Food and beverage industry (beer, wine, spirits)	m3	451'924	462'141	479'293	465'753	467'699	461'453	454'903	449'070	446'567	447'709
2H3 Blasting and shooting; blasting agent and powder	kt	1.5	1.1	1.4	2.1	2.4	2.9	2.3	2.2	2.1	2.2

4.9.2.2 Food and beverages industry (2H2)

Methodology

In Switzerland, production of beverages comprises wine, beer and spirits and food industry comprises production of bread, sugar, smoked meat, roasting of coffee and the milling industry. The CO and NMVOC emissions from food and beverages industry are calculated by a Tier 2 method according to EMEP/EEA (2013) using country-specific emission factors.

Emission factors

The emission factors for CO and NMVOC emissions from food and beverages industry in Switzerland are country-specific and based on measurements and data from industry and expert estimates as documented in the EMIS database (EMIS 2017/2H2). The implied emission factors are production-weighted (Table 4-56).

Indirect CO₂ emissions from CO and NMVOC emissions from food and beverages industry are of biogenic origin and are therefore, not reported in chp. 9 Indirect CO₂ and N₂O emissions.

Activity data

The annual amount of food and beverages produced in Switzerland is based on data from industry and the farmers' association (SBV) and expert estimates as documented in EMIS 2017/2H2 (Table 4-57).

4.9.2.3 Other (2H3)

Methodology

For determination of emissions of CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting, an analogous Tier 2 method with country-specific emission factors is used as documented in the EMIS database (EMIS 2017/2H3 Sprengen und Schiessen).

Emission factors

The emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting activities in Switzerland are country-specific and based on measurements and data from industry and expert estimates (see Table 4-56).

Activity data

The annual amount of used explosives is based on the Federal statistics on explosives (FEDPOL 2016) (Table 4-57).

4.9.3 Uncertainties and time-series consistency

The uncertainty for CO₂ emissions from 2H3 Other is estimated to be 8% (expert judgement) since activity data are taken from customs statistics.

Consistency: Time series for 2H Other are all considered consistent.

4.9.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

4.9.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. These recalculations cause a change in emission levels 1990 and 2014 of less than 0.3 kt CO₂ eq.

- 2H1: AD of 2H1 Chipboard production have been revised from 2005 onwards based on a changed density value of the chipboard produced.
- 2H2: Activity data of wine production has slightly decreased in the year 2011 due to the correction in the underlying statistics from the Swiss Alcohol Administration.
- 2H2: Activity data of sugar production has slightly changed for the years 2011, 2012 and 2014 due to corrections in the statistics provided by the Swiss Sugar Association.
- 2H2: Activity data of meat smokehouses has slightly changed for the years 2007, 2013 and 2014 due to corrections in the statistics provided by the Swiss Federal Office for Statistics.
- 2H3: SO₂ emissions from Claus units were previously reported in source category 2H3 and are now reported in 1B2aiv.

4.9.6 Category-specific planned improvements

No category-specific improvements are planned.

5 Agriculture

5.1 Overview

This chapter provides information on the estimation of the greenhouse gas emissions from the sector Agriculture. The following source categories are reported:

- 3A Enteric fermentation, CH₄ emissions from domestic livestock,
- 3B Manure management, emissions of CH₄, N₂O and NO_x,
- 3D Agricultural soils, emissions of N₂O, NO_x and NMVOC,
- 3G Liming, emissions of CO₂
- 3H Urea application; emissions of CO₂

Categories 3C Rice cultivation, 3E Prescribed burning of savannahs and 3F Field burning of agricultural residues do not occur in Switzerland and are therefore not reported.

CO₂ emissions from soils are reported under Land use, land-use change and forestry. CO₂ emissions from energy use in agriculture are reported under 1A4c Agriculture/forestry/fishing.

Total greenhouse gas emissions from the agriculture sector in 2015 were 6'074 kt CO₂ equivalents which is a contribution of 12.6% to the total of Swiss greenhouse gas emissions (including indirect CO₂, excluding LULUCF, Table 2-1, Table 5-1). Main agricultural sources of greenhouse gases in 2015 were 3A Enteric fermentation, emitting 55% of all agricultural greenhouse gases, followed by 3D Agricultural soils with 25% and 3B Manure management with 19% (Figure 5-1). 3G Liming and 3H Urea application contributed 0.5% and 0.2% respectively.




Figure 5-1 Greenhouse gas emissions of the agricultural sector in kt CO₂ equivalents 1990–2015.

Table 5-1 Greenhouse gas emissions of the agricultural sector in kt CO₂ equivalents 1990–2015.

Gas	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	kt CO ₂ equivalent									
CO ₂	49	42	42	42	42	42	42	38	36	37
CH ₄	4'509	4'496	4'410	4'353	4'374	4'363	4'333	4'235	4'201	4'135
N ₂ O	2'222	2'206	2'177	2'132	2'098	2'083	2'072	1'988	1'981	1'951
Sum	6'780	6'744	6'629	6'527	6'513	6'489	6'446	6'261	6'218	6'124

Gas	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	kt CO ₂ equivalent									
CO ₂	39	40	41	39	44	42	42	46	44	42
CH ₄	4'120	4'156	4'130	4'092	4'061	4'099	4'136	4'159	4'262	4'214
N ₂ O	1'949	1'975	1'965	1'928	1'932	1'936	1'934	1'963	1'967	1'938
Sum	6'108	6'170	6'136	6'059	6'036	6'078	6'112	6'168	6'273	6'194

Gas	2010	2011	2012	2013	2014	2015
	kt CO ₂ equivalent					
CO ₂	44	44	42	42	46	44
CH ₄	4'197	4'184	4'175	4'139	4'174	4'153
N ₂ O	1'972	1'932	1'910	1'879	1'931	1'877
Sum	6'213	6'159	6'126	6'060	6'150	6'074

CH₄ and N₂O emissions generally declined from 1990 until 2004 (Figure 5-2). Subsequently emissions increased slightly until 2008 and decreased again until 2015. This general development can be explained by the development of the cattle population and the input of mineral fertilisers. Use of mineral fertiliser declined due to the introduction of the “Proof of Ecological Performance (PEP)” in the early 1990s (Agroscope 2016b, Leifeld and Fuhrer 2005), while the cattle population was influenced by the market situation, the milk quotation system (suspended in 2009) and the general agricultural policy- and subsidy-system (OECD 2013). Most emission factors did not change significantly over the inventory years. CO₂ emissions display high year to year variability due to variability of urea application, which depends among others on the relative price levels of different industrial fertilisers.




Figure 5-2 Trend of the greenhouse gases of the agricultural sector 1990–2015. The base year 1990 represents 100%.

Among the key categories of the Swiss inventory, five are from the agricultural sector (Figure 5-3).




Figure 5-3 Key categories (Approach 1 and Approach 2) in the agricultural sector 1990 and 2015.

5.2 Source category 3A – Enteric fermentation

5.2.1 Source category description

Table 5-2 Key categories (KCA incl. LULUCF) of 3A Enteric fermentation

Code	IPCC Category	GHG	Identification Criteria
3A	Enteric Fermentation	CH4	L1, L2, T1

This emission source comprises the domestic livestock population broken down into 3 cattle categories (mature dairy cattle, other mature cattle, growing cattle), sheep, swine, buffalo, camels, deer, goats, horses, mules and asses, poultry, rabbits and livestock not covered by the agricultural census (livestock NCAC)(Table 5-3).

Emissions from 3A Enteric fermentation declined from 1990 until 2004, mainly due to a reduction in the number of cattle. However, between 2004 and 2008 cattle livestock numbers and subsequently CH₄ emissions increased, whereas since 2008 they were decreasing again.

Cattle contribute over 94% to the overall emissions from 3A Enteric fermentation and the contribution of mature dairy cattle is more than 62%.

Table 5-3 Specification of source category 3A Enteric fermentation.

3A	Source	Specification
3A1	Cattle	Mature Dairy Cattle
		Other Mature Cattle
		Growing Cattle (Fattening Calves, Pre-Weaned Calves, Breeding Cattle 1st year (Breeding Calves + Breeding Cattle 4-12 months), Breeding Cattle > 1 year, Fattening Cattle (Fattening Calves 0-4 months, Fattening Cattle 4-12 months))
3A2	Sheep	Lambs < 1 year Mature Sheep
3A3	Swine	
3A4a	Buffalo	Bisons < 3 years Bisons > 3 years
3A4b	Camels	Llamas < 2 years Llamas > 2 years
		Alpacas < 2 years Alpacas > 2 years
3A4c	Deer	Fallow Deer Red Deer
3A4d	Goats	
3A4e	Horses	Horses < 3 years Horses > 3 years
3A4f	Mules and Asses	Mules Asses
3A4g	Poultry	
3A4h i	Rabbits	
3A4h ii	Livestock NCAC	Sheep
		Goats
		Horses < 3 years
		Horses > 3 years
		Mules Asses

5.2.2 Methodological issues

5.2.2.1 Methodology

For mature dairy cattle a detailed Tier 3 model approach is applied, predicting gross energy intake by the means of a feeding model that takes into account animal performance and diet bio-chemical composition. A country-specific methane conversion rate (Y_m) was derived from a series of studies representing Swiss specific feeding conditions.

Emission estimation for all other cattle categories follows a Tier 2 approach. This means that detailed country-specific data on nutrient requirements and feed intake were used. CH_4 conversion rates were taken from the 2006 IPCC Guidelines.

Methods for all other animal categories are based on a Tier 2 approach, estimating country-specific energy intake rates. Methane conversion rates were taken from the 2006 IPCC Guidelines or from published peer reviewed literature.

The calculation of CH_4 emissions is conducted in Agroscope 2017.

5.2.2.2 Emission factors

All emission factors for 3A Enteric fermentation are country-specific, based on IPCC equation 10.21 (IPCC 2006):

$$EF = \frac{GE * (Y_m \div 100) * 365 \text{ days / y}}{55.65 \text{ MJ / kg } CH_4}$$

EF = annual CH_4 emission factor (kg/head/year)

GE = gross energy intake (MJ/head/day)

Y_m = methane conversion rate, which is the fraction of gross energy in feed converted to methane (%)

55.65 MJ/kg = energy content of methane.

5.2.2.2.1 Gross energy intake (GE)

For calculating the gross energy intake (GE), country-specific methods based on available data on requirements of net energy, digestible energy and metabolisable energy were used. The different energy levels used for energy conversion from energy required for maintenance and production to GE intake are illustrated in Figure 5-4. The respective conversion factors are given in Table 5-4.

For the **cattle categories** detailed estimations for energy requirements are necessary. As the Swiss Farmers Union (SBV) does not provide these estimates on a detailed cattle sub-category level, requirements for each cattle source category were calculated individually following the feeding recommendations for Switzerland provided in RAP (1999) and Morel et al. (2015). These RAP recommendations are also used by the Swiss farmers as the basis for their cattle feeding regimes and for filling in application forms for direct payments; they are therefore highly appropriate.




Figure 5-4 Levels of feed energy conversion (Soliva 2006).

Table 5-4 Conversion factors used for calculation of energy requirements of individual livestock categories (Soliva 2006). GE: Gross energy; DE: Digestible energy; ME: Metabolisable energy; NEL: Net energy for lactation; NEV: Net energy for growth. Blue: annually changing parameters, value for 2015.

Livestock Category		Conversion Factors	
Mature Dairy Cattle		NEL to GE	0.341
Other Mature Cattle		NEL to GE	0.265
Growing Cattle	<i>Fattening Calves</i>	<i>ME to GE</i>	<i>0.939</i>
	<i>Pre-Weaned Calves</i>	<i>NEL to GE</i>	<i>0.299</i>
	<i>Breeding Calves</i>	<i>NEL to GE</i>	<i>0.358</i>
	<i>Breeding Cattle (4-12 months)</i>	<i>NEL to GE</i>	<i>0.319</i>
	<i>Breeding Cattle (> 1 year)</i>	<i>NEL to GE</i>	<i>0.313</i>
	<i>Fattening Calves (0-4 months)</i>	<i>NEV to GE</i>	<i>0.355</i>
	<i>Fattening Cattle (4-12 months)</i>	<i>NEV to GE</i>	<i>0.397</i>
	Sheep	<i>Fattening Sheep</i>	<i>NEV to GE</i>
<i>Milksheep</i>		<i>NEL to GE</i>	<i>0.287</i>
Swine		DE to GE	0.682
Buffalo		NA	NA
Camels		NA	NA
Deer		NA	NA
Goats		NEL to GE	0.283
Horses		DE to GE	0.700
Mules and Asses		DE to GE	0.700
Poultry		ME to GE	0.700
Rabbits		NA	NA
Livestock NCAC		NA	NA

For **mature dairy cattle** a detailed feeding model from the Agroscope Institute for Livestock Sciences was used to predict gross energy intake (Agroscope 2014c).

Energy and protein requirements were estimated based on animal performance (body weight, milk production, pregnancy) following the standard feeding recommendations for Switzerland (RAP 1999). An average body weight of 650 kg was assumed for the typical Swiss mature dairy cattle. Statistics of annual milk production are provided by the Swiss Farmers Union (SBV 2016, Table 5-5). Milk production includes marketed milk, milk consumed by calves on farms and milk sold outside the commercial industry (MISTA 2016). It should be noted that daily milk yield refers to milk production during lactation (305 days) and not during the whole year (365 days). Accordingly, milk production and energy requirement for lactation was zero during the two remaining months when the cows are dry. During the dry months additional energy requirements for pregnancy were accounted for.

To cover total animal energy and protein requirements, typical Swiss specific basic feed rations were defined as model inputs. The average basic feed ration in summer consisted of 92% fresh grass and 8% maize cubes. In winter the feed ration consisted of 10% maize silage, 13% grass silage, 72% hay and 5% fodder beet. Concentrates are automatically supplemented in the model according to additional energy and protein requirements not covered by the basic feed ration. Concentrates consisted of a varying mixture of barley grains, wheat grains, maize grains, maize gluten, soybean meal and rapeseed meal according to specific animal requirements. Subsequently, average bio-chemical composition and properties of the total feed ration (e.g. energy content, protein content, digestibility) were derived, weighing the respective values of the individual feed ingredients given in the Swiss Feed Database (Agroscope 2014b). Finally, gross energy intake was estimated based on the total feed intake and the gross energy content of the total ration that was 18.26 MJ/kg on average for the years 1990–2015.

In the year 2003 yearly milk yield surpassed 6000 kg. To achieve yearly milk yields higher than 6000 kg, cows have to be fed with an increasing share of feed concentrates that have a substantially higher net energy (NE) density than the basic feed ration. The model reproduces this behaviour. Due to the increasing ratio of net energy to gross energy the increase of gross energy intake is slower after the year 2003 although milk yield increases more or less at the same rate (Table 5-6).

A more exhaustive model description is contained in Agroscope 2016b.

Table 5-5 Average daily milk production during 305 days of lactation in Switzerland 1990–2015.

Milk Production Cattle		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Population Size Mature Dairy Cattle	head	783'100	780'500	763'500	744'450	749'700	739'641	736'043	711'613	701'343	683'545
Lactation Period	day	305	305	305	305	305	305	305	305	305	305
Milk Yield Mature Dairy Cattle	kg/head/day	16.06	16.35	16.39	16.78	16.75	17.09	16.96	17.48	17.97	18.40
Milk Yield Other Mature Cattle	kg/head/day	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20

Milk Production Cattle		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Population Size Mature Dairy Cattle	head	669'410	669'410	657'924	638'288	621'008	620'708	618'065	614'795	628'516	599'361
Lactation Period	day	305	305	305	305	305	305	305	305	305	305
Milk Yield Mature Dairy Cattle	kg/head/day	18.75	18.97	19.34	19.77	20.43	20.45	20.57	21.21	21.66	22.27
Milk Yield Other Mature Cattle	kg/head/day	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20	8.20

Milk Production Cattle		2010	2011	2012	2013	2014	2015
Population Size Mature Dairy Cattle	head	589'024	589'239	591'212	586'609	587'385	583'277
Lactation Period	day	305	305	305	305	305	305
Milk Yield Mature Dairy Cattle	kg/head/day	22.46	22.63	22.57	22.27	22.87	23.10
Milk Yield Other Mature Cattle	kg/head/day	8.20	8.20	8.20	8.20	8.20	8.20

For **other mature cattle** and **growing cattle**, data on energy intake were based on the feeding requirements according to RAP (1999) and Morel et al. (2015). In the calculation of the NE data, the animal's weight, daily growth rate, daily feed intake (dry matter), daily feed energy intake, and energy required for milk production and pregnancy for the respective sub-categories were considered. The method is described in detail in Soliva (2006) but has been revised slightly since. NE is further subdivided into NE for lactation (NEL) and NE for growth (NEV)(Table 5-4). For some of the growing cattle categories NEL is used, rather than NEV that would seem logical. However, cattle-raising is often coupled with dairy cattle activities and therefore the same energy unit (NEL) is used in these cases. Exceptions are the fattening calves (milk-fed calves), whose requirement for energy is expressed as metabolisable energy (ME).

Table 5-6 Gross energy intake per head of different livestock groups. Sub-categories not contained in the reporting tables (CRF) are displayed in italic. Whole time series on a livestock sub-category level are provided in Annex 3 A.3.3.

Gross Energy Intake	1990-2011									
	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011
	MJ/head/day									
Cattle										
Mature Dairy Cattle	259.3	267.4	280.1	291.6	292.2	295.2	297.2	299.8	300.7	301.4
Other Mature Cattle	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6
Growing Cattle (weighted average)	101.0	101.5	101.0	97.7	97.7	97.4	97.3	97.0	96.9	96.6
<i>Fattening Calves</i>	<i>47.1</i>									
<i>Pre-Weaned Calves</i>	<i>38.1</i>									
<i>Breeding Calves</i>	<i>27.8</i>									
<i>Breeding Cattle (4-12 months)</i>	<i>90.1</i>									
<i>Breeding Cattle (> 1 year)</i>	<i>143.6</i>									
<i>Fattening Calves (0-4 months)</i>	<i>50.0</i>									
<i>Fattening Cattle (4-12 months)</i>	<i>126.3</i>									
Sheep	21.2	24.0	22.4	22.8	22.6	22.2	22.0	22.7	22.6	22.6
Swine	28.3	31.9	28.0	26.6	26.3	26.9	26.7	27.0	27.2	26.9
Buffalo (weighted average)	NA	136.6	146.9	140.6	138.9	130.4	129.1	134.8	136.9	139.8
Camels (weighted average)	NA	NA	34.8	31.7	31.7	31.7	31.6	31.5	31.0	31.4
Deer (weighted average) ¹⁾	50.5	55.3	56.4	55.4	55.8	55.9	56.5	56.8	56.5	56.7
Goats	25.0	27.9	25.7	25.4	25.3	25.0	25.0	25.3	25.1	25.6
Horses (weighted average)	107.3	106.9	107.4	107.7	107.7	107.7	107.7	107.8	107.9	107.9
Mules and Asses (weighted average)	39.2	39.7	39.5	39.4	39.5	39.3	39.2	40.0	40.2	39.9
Poultry ²⁾	1.5	1.3	1.4	1.2	1.2	1.3	1.3	1.3	1.2	1.2
Rabbits	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
Livestock NCAC (weighted average)	95.6	83.4	37.7	33.3	33.2	32.9	31.7	34.3	36.9	37.2

Gross Energy Intake	2012-2015			
	2012	2013	2014	2015
	MJ/head/day			
Cattle				
Mature Dairy Cattle	301.2	299.8	302.5	303.6
Other Mature Cattle	250.6	250.6	250.6	250.6
Growing Cattle (weighted average)	96.1	96.0	96.0	96.4
<i>Fattening Calves</i>	<i>47.1</i>	<i>47.1</i>	<i>47.1</i>	<i>47.1</i>
<i>Pre-Weaned Calves</i>	<i>38.1</i>	<i>38.1</i>	<i>38.1</i>	<i>38.1</i>
<i>Breeding Calves</i>	<i>27.8</i>	<i>27.8</i>	<i>27.8</i>	<i>27.8</i>
<i>Breeding Cattle (4-12 months)</i>	<i>90.1</i>	<i>90.1</i>	<i>90.1</i>	<i>90.1</i>
<i>Breeding Cattle (> 1 year)</i>	<i>143.6</i>	<i>143.6</i>	<i>143.6</i>	<i>143.6</i>
<i>Fattening Calves (0-4 months)</i>	<i>50.0</i>	<i>50.0</i>	<i>50.0</i>	<i>50.0</i>
<i>Fattening Cattle (4-12 months)</i>	<i>126.3</i>	<i>126.3</i>	<i>126.3</i>	<i>126.3</i>
Sheep	22.5	22.4	22.5	22.5
Swine	26.5	27.3	27.4	27.4
Buffalo (weighted average)	135.9	136.0	134.6	134.5
Camels (weighted average)	31.6	31.9	31.8	31.6
Deer (weighted average) ¹⁾	57.0	58.1	58.0	58.0
Goats	25.6	25.6	25.0	25.0
Horses (weighted average)	107.9	108.0	108.1	108.3
Mules and Asses (weighted average)	39.9	39.6	39.6	39.6
Poultry ²⁾	1.2	1.2	1.2	1.2
Rabbits	1.2	1.2	1.2	1.2
Livestock NCAC (weighted average)	37.9	38.2	38.5	38.7

¹⁾ Deer: Gross energy intake per animal place (mother with offspring)

²⁾ Poultry data is not gross energy intake (GE) but metabolizable energy intake (ME)

The gross energy intake for other mature cattle is significantly higher than IPCC default values, since the category “other mature cattle” in Switzerland only includes mature cows that produce offspring for meat (so-called “suckler cows” or “mother cows”). Milk production of other mature cattle is 2500 kg per head and year (305 days of lactation) and has not changed over the inventory time period (Morel et al. 2015).

The gross energy intake of growing cattle was calculated separately for all sub-categories displayed in Table 5-6 (in italics) and subsequently averaged (weighted average). No

methane is generated from milk. Therefore, the energy intake from milk or milk products is not considered when estimating methane emission factors from enteric fermentation of calves. The values for all 7 sub-categories are constant over time. Since the composition of the growing cattle category changed over time (e.g. more pre-weaned calves, fewer fattening calves, see Table 5-8), the average gross energy intake for growing cattle also changes slightly. To calculate an annual emission factor, the categories breeding calves and breeding cattle 4-12 months were combined in the category breeding cattle 1st year (not shown in Table 5-6 and Table 5-8). Accordingly, the respective animals have two separate gross energy intake values, i.e. 27.8 MJ/head/day for the first 4 months and 90.1 MJ/head/day for the last 8 months. The same procedure is applied for fattening calves 0–4 months (50.0 MJ/head/day) and fattening cattle 4–12 months (126.3 MJ/head/day) summing up to the category fattening cattle.

Energy requirements and gross energy intake of **sheep, swine, goats and poultry** were obtained from the respective estimates of the Swiss Farmers Union (SBV 2016, Giuliani 2016). These estimates are not officially published anymore in the statistical yearbooks (e.g. SBV 2016) but are still available from background data and are based on the same method as earlier published energy requirement statistics (e.g. SBV 2007).

Gross energy intake for **horses, mules and asses** were estimated by Stricker (2012), mainly based on Meyer and Coenen (2002).

Table 5-7 Dry matter and gross energy requirements for buffalo, camels and deer according to Flisch et al. (2009).

		DM Intake	GE Intake
		kg DM/head/day	MJ/head/day
Buffalo	Bisons < 3 years	4.93	90.99
	Bisons > 3 years	10.68	197.14
Camels	Llamas < 2 years	1.34	24.77
	Llamas > 2 years	2.33	42.97
	Alpacas < 2 years	0.82	15.16
	Alpacas > 2 years	1.51	27.80
Deer ¹⁾	Fallow Deer	2.74	50.55
	Red Deer	5.48	101.10

1) Requirements for deer are assessed per animal place i.e. mother with offspring.

For **buffalo, camels and deer**, energy intake was derived from data on dry matter intake provided in Flisch et al. (2009) (Table 5-7). According to the 2006 IPCC Guidelines an energy density of 18.45 MJ*kg⁻¹ was used to convert dry matter to gross energy.

Energy intake of **rabbits** was estimated by Menzi (2014) based on Schlegel and Menzi (2013).

Finally for **livestock NCAC** the same energy intakes as the respective animal categories in the official census were used.

Final compilation of livestock gross energy intake was conducted in Agroscope (2017). Resulting estimates are provided in Table 5-6 (main categories) and in Annex 3 A3.3 (all years and all sub-categories).

5.2.2.2.2 Methane conversion rate (Y_m)

For the methane conversion rate (Y_m), few country-specific data exist. Accordingly, for most animal categories default or literature values were used. Due to its great importance a country-specific Y_m was used for **mature dairy cattle**. A value of 6.9% was derived from a series of measurements conducted under Swiss specific feeding and husbandry conditions at the Federal Institute of Technology in Zürich (based on data compiled in Zeitz et al. (2012) and additional measurements described in Estermann et al. (2001), Külling et al. (2002) and Staerfl et al. (2012)).

For all **other cattle categories, sheep and buffalo** default values recommended by the IPCC for developed countries in Western Europe were used (IPCC 2006: Table 10.12, 10.13, 10A.2, 10A.3). For all juvenile cattle consuming only milk (i.e. fattening calves) the methane conversion rate is assumed to be zero.

According to table 10.13 in IPCC (2006) two different Y_m were used for **sheep**, namely 4.5% for lambs <1 year and 6.5% for mature sheep. Overall Y_m was subsequently weighted according to the population structure. For **camels and deers** the same overall methane conversion rate as for sheep was applied, assuming the same relationship between adult and juvenile animals.

For **Swine** a methane conversion rate of 0.6% was used. This value was suggested by Crutzen et al. (1986) and was confirmed by the compilation of references in Minonzio et al. (1998). Since the 2006 IPCC Guidelines do not provide a default value for **goats**, an Y_m of 6% was adopted based on the work of Martínez-Fernández et al. (2014) and Fernández et al. (2013). For **Horses, mules and asses** an Y_m of 2.45% was used, which corresponds to a methane energy loss of 3.5% of digestible energy (Vermorel et al. 1997, Minonzio et al. 1998) and a feed digestibility of 70% (Stricker 2012). For **poultry** a country-specific value (0.16% of metabolisable energy) was used. This value was evaluated in an in vivo trial with broilers (Hadorn and Wenk 1996). For **rabbits** an Y_m of 0.6% was applied as suggested in the national GHG inventory of Italy (ISPRA 2014). Finally, as for gross energy intake, the same methane conversion rates as the respective animals in the official census were used for **livestock NCAC**.

5.2.2.3 Activity data

Livestock population data was obtained from statistics published by the Swiss Farmers Union (SBV 2016) and the Swiss Federal Statistical Office (SFSO 2016)(Table 5-8). All activity data was revised and harmonized during a joint effort of the Agroscope Reckenholz-Tänikon Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012).

The category other mature cattle only includes mature cows used to produce offspring for meat.

Emission estimation for growing cattle was conducted at a more disaggregated level than the one displayed in the reporting tables (CRF). The livestock category growing cattle in the reporting tables includes the sub-categories fattening calves, pre-weaned calves, breeding

calves, breeding cattle 4–12 months, breeding cattle >1 year, fattening calves 0–4 months and fattening cattle 4–12 months. Although not growing cattle in the proper sense, bulls are contained in the categories breeding cattle (>1 year) and fattening cattle (4–12 months) according to their purposes. This disaggregation of the cattle category enhances the accuracy of the emission estimation procedure from livestock activities (also refer to chp. 5.3.2.1).

Emission estimation for buffalo, camels, horses, mules and asses and deers was also conducted on a more disaggregated level than displayed in the reporting tables (CRF). Additional data on a livestock sub-category level is contained in Annex 3 A3.3.

Additionally to official statistical data, population data of livestock not covered by the agricultural census of the Swiss Federal Statistical Office was assessed. The respective category “Livestock NCAC” (livestock not covered by agricultural census) consists of sheep, goats, horses and mules and asses held for non-agricultural purposes (e.g. horses for sports and leisure) and/or livestock held by private persons or enterprises that do not fulfil the criteria of an agricultural enterprise. Data for the respective horses, mules and asses were derived from Poncet et al. (2007, 2009) and Schmidlin et al. (2013). For sheep and goats, data from individual cantons having full livestock census was used to estimate the relative share for the whole of Switzerland. The respective estimates were conducted in the course of the elaboration of the gross nutrient balance of the Swiss Federal Statistical Office (SFSO 2016b).

Table 5-8 Activity data for calculating methane emissions from 3A Enteric fermentation (ART/SHL 2012, SBV 2016, SFSO 2016, SFSO 2016b). The complete time series on a livestock sub-category level are provided in Annex 3 A3.3.

Population Size	1990-2011									
	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011
	1'000 head									
Cattle	1'855	1'748	1'588	1'555	1'567	1'572	1'604	1'597	1'591	1'577
Mature Dairy Cattle	783	740	669	621	618	615	629	599	589	589
Other Mature Cattle	12	23	45	78	87	94	98	108	111	111
Growing Cattle	1'060	986	874	856	862	863	877	890	891	877
<i>Fattening Calves</i>	112	102	103	106	101	100	95	101	99	101
<i>Pre-Weaned Calves</i>	10	18	36	62	67	72	76	86	88	88
<i>Breeding Cattle 1st Year</i>										
<i>Breeding Calves</i>	214	166	76	75	77	76	80	77	77	75
<i>Breeding Cattle (4-12 months)</i>	132	129	161	147	147	147	152	149	149	145
<i>Breeding Cattle (> 1 year)</i>										
<i>Breeding Cattle 2nd Year</i>	253	239	222	205	210	210	213	212	213	207
<i>Breeding Cattle 3rd Year</i>	151	139	130	113	110	109	110	119	119	116
<i>Fattening Cattle</i>										
<i>Fattening Calves (0-4 months)</i>	88	82	43	35	35	34	36	35	34	34
<i>Fattening Cattle (4-12 months)</i>	100	110	105	112	114	114	116	112	111	111
Sheep	395	387	421	446	448	444	446	432	434	424
Swine	1'787	1'446	1'498	1'609	1'635	1'573	1'540	1'557	1'589	1'579
Buffalo	0	0	0	0	0	0	0	1	1	1
Camels	0	0	1	3	3	4	4	5	6	6
Deer ¹⁾	0	1	3	4	4	4	5	5	6	6
Goats	68	53	62	74	76	79	81	81	83	83
Horses	28	41	50	55	56	58	59	60	62	57
Mules and Asses	6	8	12	16	16	17	18	19	20	19
Poultry	5'938	6'251	6'983	8'260	7'670	8'228	8'543	8'809	9'025	9'478
Rabbits	61	41	28	25	24	27	25	28	35	34
Livestock NCAC	28	30	94	86	82	80	86	90	91	99

Population Size	2012-2015			
	2012	2013	2014	2015
	1'000 head			
Cattle	1'565	1'557	1'563	1'554
Mature Dairy Cattle	591	587	587	583
Other Mature Cattle	114	117	118	118
Growing Cattle	859	854	857	853
<i>Fattening Calves</i>	99	97	97	91
<i>Pre-Weaned Calves</i>	91	93	93	93
<i>Breeding Cattle 1st Year</i>				
<i>Breeding Calves</i>	73	72	72	73
<i>Breeding Cattle (4-12 months)</i>	140	138	139	140
<i>Breeding Cattle (> 1 year)</i>				
<i>Breeding Cattle 2nd Year</i>	200	198	199	200
<i>Breeding Cattle 3rd Year</i>	112	111	111	112
<i>Fattening Cattle</i>				
<i>Fattening Calves (0-4 months)</i>	34	34	34	34
<i>Fattening Cattle (4-12 months)</i>	112	111	112	111
Sheep	417	409	403	347
Swine	1'544	1'485	1'498	1'496
Buffalo	1	0	1	1
Camels	6	6	6	6
Deer ¹⁾	6	6	6	6
Goats	85	85	85	71
Horses	58	57	57	55
Mules and Asses	20	20	20	20
Poultry	9'955	10'079	10'723	10'825
Rabbits	28	28	27	25
Livestock NCAC	112	107	106	101

¹⁾ Deer: numbers correspond to animal places i.e. mother with offspring.




Figure 5-5 Relative development of main animal categories 1990–2015. The category with the strongest increase, i.e. other mature cattle, is not displayed, as it increases to over 980% of the 1990 value by 2015.

Livestock populations in Switzerland are primarily influenced by the general agricultural policy, i.e. the subsidy system, the milk quotation system and the development of the economic framework conditions. The number of cattle declined slightly until the year 2004. However, cattle livestock numbers increased between 2004 and 2008, mainly due to an increase of the number of growing cattle. Since 2008 the cattle population was decreasing again, possibly due to the suspension of the milk quotation system in 2009.

After a decrease until 1996, the number of swine increased until 2006 – a process that has been observed in many other European countries (SBV 2004: p.69). Since then, the number of swine has fluctuated slightly below the level of 2006. The number of poultry shows a rapid increase between 1990 and 2015 with a distinct dip only between 2005 and 2006, a consequence of changed human consumption patterns as a result of the avian flu in 2006.

The number of sheep was more or less constant while the number of goats increased following a decline between 1990 and 1995.

5.2.3 Uncertainties and time-series consistency

For the uncertainty analysis the input data from ART (2008a) was used and was updated with current activity and emission data as well as with new default uncertainties of the 2006 IPCC Guidelines. The arithmetic mean of the lower and upper bound uncertainty was used for activity data (6.4%) and for emission factors (16.9%), resulting in a combined uncertainty of 18.1% for Approach 1 analysis.

For the Approach 2 analysis, asymmetric probability distributions as well as possible correlations of input data were considered. The uncertainty interval of the Approach 2 analysis lies between -18% and +18% (Annex A2, Table A – 7).

For further uncertainty-results also consult chp. 1.6.1 and Annex A2.

The time series 1990–2014 are all considered consistent, although the following issues should be considered:

- Gross energy intake of some of the aggregated animal categories reveals some fluctuations during the inventory period due to varying shares of the sub-categories.
- Gross energy intake as well as the implied emission factor for mature dairy cattle increase, mainly as a result of higher milk production (Table 5-5).
- Between 1998 and 1999 the questionnaire for the collection of livestock data was modified. In some animal categories this led to minor ruptures in the time series. Consequences for overall emissions are, however, of minor importance. An analysis conducted in 2012 revealed, that while the average annual change for the years 1990–2011 over all animal categories (excluding other mature cattle) was 3.3% points, the annual change for the years 1998–1999 was 3.8% points (ART/SHL 2012).
- For the last seven inventory years the population statistics of growing cattle were not available in the usual format. Data for 2009 to 2015 is based on the animal traffic database. Aggregation was adapted to the format necessary for the AGRAMMON-model and the GHG inventory by the Swiss College of Agriculture SHL (SHL 2010). Data in the animal traffic database are considered more complete than the data from the survey of the SFSO because the latter includes also animals held outside agricultural enterprises.

5.2.4 Category-specific QA/QC and verification

General QA/QC measures are described in NIR chp. 1.2.3.

All further category-specific QA/QC activities are described in a separate document (Agroscope 2016b). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed. Furthermore, comparisons with data from other countries were conducted and discussed where possible. Agroscope 2016b is continuously updated with the most recent inventory data.

Livestock data were compared with the livestock data provided by the FAO and checked for plausibility. In all cases the new recalculated data according to ART/SHL (2012) are considered more reliable than the FAO data. Small inconsistencies (usually in the order of $\pm 2\%$) are due to updates of provisional data that are not considered by the FAO. For horses, mules and asses disagreements might be due to the different accounting of agricultural and non-agricultural horses. The Swiss inventory system accounts for all animals and differentiates between animals captured by the official agricultural census and livestock not covered by agricultural census. Moreover, the numbers of mules and asses is higher in the Swiss GHG inventory because unlike the FAO, Switzerland accounts also for ponies and lesser horses. The total numbers of poultry in the GHG inventory and the FAO data also show minor discrepancies due to different accounting of turkeys, geese, ducks and quails.

Seasonal fluctuation of the cattle population was analysed for the years 2005–2007 based on detailed information from the Swiss Farmers Union (SBV 2007a). Seasonal fluctuations are usually in the order of $\pm 3\%$ with census data (April) always slightly above the annual mean. Data from the animal traffic database (i.e. cattle populations for the years 2009–2015) refer to annual mean population.

IPCC tables with data for estimating emission factors for cattle (such as weight, weight gain, milk production) were filled in, checked for consistency and confidence and compared with IPCC default values (refer to Annex 3 A3.3).

Country-specific energy-intake rates for all cattle categories were compared to intake rates estimated with the IPCC Tier 2 default methodology (see Agroscope 2016b for details). Both approaches are comparable in the assessment of net energy requirements. However, the IPCC approach resulted in higher estimates of GE-intake. Further analyses suggest that the IPCC conversion rates of net energy into gross energy are unrealistic for conditions in Switzerland. Given the experimentally verified high feed quality standards in Switzerland, the results of the country-specific inventory method are thus much more plausible than the estimates using the unaltered IPCC default method. Moreover, a discrepancy of approximately 8.5% was found when comparing the overall GE-intake of the cattle population with the respective estimate of the Swiss Farmers Union (Giuliani 2016). As found for the comparison with the IPCC approach, different assumptions on net energy densities of the feed might explain the divergence.

During the past years a couple of studies were conducted to verify methane emissions at the regional scale, comparing bottom-up estimates with atmospheric measurements. While virtually all these measurements are subjects to great uncertainties, the overall picture support the bottom-up approach in the Swiss GHG inventory or at least does not indicate the omission of a significant methane source. Hiller et al. (2014a) found that methane emissions could be underestimated by the inventory method when they measured atmospheric CH₄ concentrations over the Reuss-valley with an airplane. However, the methodological approach applied by Hiller et al. (2014a) still relies on a number of rather uncertain basic assumptions and is therefore not beyond doubts. Additionally, it should be noted, that methane emission estimates from the agriculture sector in the Swiss GHG inventory were revised since, and currently lie approximately 10% above the estimates used in this study. Stieger (2013) and Stieger et al. (2015) reported a very good accordance of bottom-up estimates and flux measurements with a tethered balloon system. Bamberger et al. (2014) conducted regional CH₄ measurements with a measurement device mounted on a car. Measurement precision and duration was not sufficient to validate bottom-up inventory estimates. Nonetheless, they concluded that a locally relevant emission source considered negligible in the emission inventory would have been identified. Finally Henne et al. (2015) utilised continuous CH₄ observations originating from four sites of the CarboCount CH network on the Swiss Plateau and two additional sites. These observations were combined with atmospheric transport simulations and an inverse modelling framework in order to deduce the spatial distribution of CH₄ emissions in Switzerland and the adjacent countries. Total Swiss emissions were estimated to 200 ± 20 kt CH₄/yr as mean and standard deviation. This value is in very good agreement with the value of 206 ± 33 kt CH₄/yr given in the revised 2015 submission for the year 2013 (FOEN 2015, see also Annex 5 A 5.2 therein).

5.2.5 Category-specific recalculations

General information on recalculations is provided in chp. 10.

- CH₄ emission factors for the year 2012 and 2014 of sheep, goats, swine and poultry and "other" (livestock NCAC) were revised due to updates of provisional net energy intake data by the Swiss Farmers Union (Giuliani 2016). The effect of the recalculation on overall greenhouse gas emissions is negligible (<1 kt CO₂ equivalents).

5.2.6 Category-specific planned improvements

Planned improvements for future submissions are the further development, adaptation and verification of the dairy cattle feeding model (GE, Y_m).

5.3 Source category 3B – Manure management

5.3.1 Source category description

Table 5-9 Key categories (KCA incl. LULUCF) of 3B Manure management

Code	IPCC Category	GHG	Identification Criteria
3B1-3B4	Manure management	CH ₄	L1, L2
3B5	Indirect N ₂ O emissions from manure management	N ₂ O	L1, L2, T2

The emission source is the domestic livestock population broken down into 3 cattle categories (mature dairy cattle, other mature cattle, growing cattle), sheep, swine, buffalo, camels, deer, goats, horses, mules and asses, poultry, rabbits and livestock not covered by agricultural census (Livestock NCAC) (Table 5-10). Six (CH₄) respectively five (N₂O) different manure management systems are considered as well as indirect N₂O emissions from 3B Manure management (Table 5-11). Additionally, NO_x emissions from manure management are estimated.

The total emissions from 3B Manure management closely follow the development of the cattle population. Emissions declined from 1990 until 2004, increased until 2008 and subsequently decreased again.

Significant contributors to CH₄ emissions from 3B Manure management are cattle with approximately 75% and swine with approximately 22%. Likewise, cattle and swine contribute significantly to N₂O emissions with 71% and 14% respectively.

Table 5-10 Specification of source category 3B Manure management by livestock categories.

3B	Source	Specification
3B1	Cattle	Mature Dairy Cattle
		Other Mature Cattle
		Growing Cattle (Fattening Calves, Pre-Weaned Calves, Breeding Cattle 1 st year (Breeding Calves + Breeding Cattle 4-12 months), Breeding Cattle > 1 year (Breeding Cattle 2 nd year + Breeding Cattle 3 rd year), Fattening Cattle (Fattening Calves 0-4 months + Fattening Cattle 4-12 months))
3B2	Sheep	Lambs < 1 year Mature Sheep Fattening Sheep Milk Sheep
3B3	Swine	Piglets Fattening Pig over 25 kg Dry Sows Nursing Sows Boars
3B4a	Buffalo	Bisons < 3 years Bisons > 3 years
3B4b	Camels	Llamas < 2 years Llamas > 2 years Alpacas < 2 years Alpacas > 2 years
3B4c	Deer	Fallow Deer Red Deer
3B4d	Goats	Goat Places
3B4e	Horses	Horses < 3 years Horses > 3 years
3B4f	Mules and Asses	Mules Asses
3B4g	Poultry	Growers Layers Broilers Turkey Other Poultry
3B4h i	Rabbits	
3B4h ii	Livestock NCAC	Sheep Goats Horses < 3 years Horses > 3 years Mules Asses

Table 5-11 Specification of source category 3B Manure management by manure management systems.

3B	Source	Specification
3B6a	Direct Emissions	Liquid systems
3B6b		Solid storage and dry lot
3Bc / 3D		Pasture, range and paddock
3B6d		Digesters (anaerobic digestion)
3B6e		Other
3B5a	Indirect Emissions	Atmospherical deposition
3B5b		Leaching and run-off

5.3.2 Methodological issues

5.3.2.1 Methodology

The calculation is based on methods described in the 2006 IPCC Guidelines (CH₄: IPCC 2006 equation 10.23; N₂O: IPCC 2006 equation 10.25).

CH₄ emissions from 3B Manure management were generally estimated using a Tier 2 methodology. For cattle a more detailed method was applied, estimating volatile solids (VS) excretion based on gross energy intake estimates as used for enteric fermentation. Due to lacking default values, VS excretion from buffalo, camels, horses and deer was equally estimated based on gross energy intake. For the remaining livestock categories default parameters were used. Methane conversion factors (MCF) are from IPCC (2006; solid storage, pasture range and paddock, anaerobic digesters, deep litter, poultry manure) or were modelled according to Mangino et al. (2001) (liquid systems, digesters).

N₂O emissions from 3B Manure management were estimated using a Tier 2 methodology. Activity data were adjusted to the particular situation of Switzerland in coordination with the Swiss ammonia model AGRAMMON (Kupper et al. 2013). Detailed country-specific data on nitrogen excretion rates, manure management system distribution and nitrogen volatilisation were applied. Emission factors for direct N₂O emissions (EF₃) are based on IPCC (2006) whereas the emission factor for indirect emissions from atmospheric deposition is country-specific (Bühlmann et al. 2015 and Bühlmann 2014). Leaching of NO₃⁻ from manure management systems was considered negligible and is thus not included in the estimates. The N₂O emissions from pasture, range and paddock are reported under 3D Agricultural soils, source category 3Da3 (Urine and dung deposited by grazing animals).

For calculation of CH₄ and N₂O emissions, slightly different livestock sub-categories were used (Table 5-12). The livestock categories reported in the reporting tables (CRF) are the same, but the respective sub-categories as a basis for the calculation are different. The categorization for the estimation of CH₄ emissions had to be adapted to data availability for energy requirements, while the categorisation for the estimation of N₂O emissions is determined by the respective categorisation of the Swiss ammonia inventory (AGRAMMON, Kupper et al. 2013, Flisch et al. 2009). Nevertheless, there is no inconsistency in the total number of animals as they are the same both for CH₄ and N₂O emissions. Note that although not growing cattle in the proper sense, bulls are contained in the categories breeding cattle >1 year, breeding cattle 3rd year and/or fattening cattle according to their purposes.

The calculation of CH₄ and N₂O emissions is conducted in Agroscope 2017.

Table 5-12 Livestock categories for estimating CH₄ and N₂O emissions from 3B Manure management.

3B	CH ₄	N ₂ O
Cattle	Mature Dairy Cattle	Mature Dairy Cattle
	Other Mature Cattle	Other Mature Cattle
	Growing Cattle Fattening Calves Pre-Weaned Calves Breeding Cattle 1 st year (Breeding Calves + Breeding Cattle 4-12 months) Breeding Cattle > 1 year Fattening Cattle (Fattening Calves 0-4 months + Fattening Cattle 4-12 months)	Growing Cattle Fattening Calves Pre-Weaned Calves Breeding Cattle 1 st year Breeding Cattle 2 nd year Breeding Cattle 3 rd year Fattening Cattle
Sheep	Lambs < 1 year Mature Sheep	Fattening Sheep Milk Sheep
Swine	Swine	Piglets Fattening Pig over 25 kg Dry Sows Nursing Sows Boars
Buffalo	Bisons < 3 years Bisons > 3 years	Bisons < 3 years Bisons > 3 years
Camels	Llamas < 2 years Llamas > 2 years Alpacas < 2 years Alpacas > 2 years	Llamas < 2 years Llamas > 2 years Alpacas < 2 years Alpacas > 2 years
Deer	Fallow Deer Red Deer	Fallow Deer Red Deer
Goats	Goats	Goat places
Horses	Horses < 3 years Horses > 3 years	Horses < 3 years Horses > 3 years
Mules and Asses	Mules Asses	Mules Asses
Poultry	Poultry	Growers Layers Broilers Turkey Other Poultry
Rabbits	Rabbits	Rabbits
Livestock NCAC	Sheep Goats Horses < 3 years Horses > 3 years Mules Asses	Sheep Goats Horses < 3 years Horses > 3 years Mules Asses

5.3.2.2 Emission factors CH₄

Calculation of CH₄ emissions from 3B Manure management is based on methods described in the 2006 IPCC Guidelines (IPCC 2006, equation 10.23):

$$EF_T = VS_T \cdot 365 \text{ days / year} \cdot B_{0T} \cdot 0.67 \text{ kg / m}^3 \cdot \sum_S MCF_S \cdot MS_{TS}$$

EF_T = annual CH₄ emission factor for livestock category T (kg/head/year)

VS_T = daily volatile solids (VS) excreted for livestock category T (kg/head/day)

B_{0T} = maximum CH₄ producing capacity for manure produced by livestock category T (m³/kg)

0.67 kg/m³ = conversion factor of m³ CH₄ to kilograms CH₄

MCF_S = CH₄ conversion factors for each manure management system S (%)

MS_{TS} = fraction of livestock category T 's manure handled using manure management system S (dimensionless)

5.3.2.2.1 Volatile solids excretion (VS)

The daily excretions of volatile solids (VS) for **all cattle sub-categories** were estimated according to equation 10.24 in the 2006 IPCC Guidelines (IPCC 2006):

$$VS = \left[GE \cdot \left(1 - \frac{DE\%}{100} \right) + (UE \cdot GE) \right] \cdot \left[\frac{1 - ASH}{EDF} \right]$$

VS = volatile solids excretion per day on a dry-organic matter basis (kg/day)

GE = gross energy intake (MJ/head/day)

DE = digestibility of the feed (%)

$(UE \cdot GE)$ = urinary energy expressed as fraction of GE

ASH = ash content of manure calculated as a fraction of the dry matter feed intake

EDF = energy density of feed, conversion factor for dietary GE per kg of dry matter (MJ/kg)

Gross energy intake was calculated according to the method described in chp. 5.2.2.2.1. In the case of **mature dairy cattle** the same model was used as for the estimation of CH₄ emissions from 3A Enteric fermentation. Content of net energy, gross energy and ash in feed dry matter as well as feed digestibility were also estimated using the Agroscope feeding model (Agroscope 2014c). The digestibility of feed is of crucial importance for the calculation of volatile solids. The modelled values for dairy cattle are somewhat higher than the IPCC default and were compared to measurements from feeding trials in Switzerland. The comparison revealed that modelled values are on average slightly higher than measurements. Accordingly, an adjustment was made in order to take account of the high feeding level that is usually above maintenance (Ramin and Huhtanen 2012). High feeding

levels may lead to an increase in rumen passage rate and subsequently to lower feed digestibility (Nousiainen et al. 2009). The correction decreased the feed digestibility on average by 2.5 per cent points. Resulting feed digestibility was 72.2% on average, gross energy content (EDF) was 18.26 MJ/kg and ash content was 9.0% each with very small fluctuations along the time series. Urinary energy expressed as fraction of gross energy was 0.04 (IPCC 2006).

For **calves and other growing cattle** IPCC default values of 65% respectively 60% were taken for the feed digestibility. For the urinary energy expressed as fraction of gross energy and for the energy density of the feed (EDF) the IPCC default values, i.e. 0.04 and 18.45 MJ/kg were adopted. Furthermore, an ash content of 8.0% was used for all these categories.

For VS excretion of the livestock categories **sheep, swine, goats, mules and asses, poultry, rabbits and livestock NCAC**, default values from IPCC were taken (IPCC 2006, Tables 10A-7, 10A-8, 10A-9).

For **buffalo, camels, horses and deer** VS excretion was again estimated using equation 10.24 in the 2006 IPCC Guidelines with default values for feed digestibility and ash content (IPCC 2006). Feed digestibility was 55% for buffalos, 60% for camels and deer (assuming similar feed composition as for sheep) and 70% for horses. The urinary energy as fraction of the gross energy was 0.04, the energy density of the feed (EDF) was 18.45 MJ/kg. The ash content of manure was 8.0% for buffalo, camels and deer and 4.0% for horses (IPCC 2006).

Finally for **livestock NCAC** the same VS excretion rates as for the respective animal categories in the official census were used.

5.3.2.2.2 Maximum CH₄ producing capacity (B₀)

For the methane producing capacity (B₀) default values were used (IPCC 2006). For deer the same value as for sheep was applied as no default value was available (i.e. 0.19 m³/kg).

5.3.2.2.3 Methane conversion factor (MCF)

For estimating CH₄ emissions from manure management, six different manure management systems are distinguished. Switzerland has an average annual temperature below 15°C (MeteoSwiss 2014) and was therefore allocated to the cool climate region without differentiation.

In the case of **solid manure** and **pasture range and paddock** the default MCF values from table 10.17 of the 2006 IPCC Guidelines were used (Table 5-13).

Liquid/slurry systems are responsible for the major part of methane emissions from Manure management (90% on average). Accordingly a more detailed model was used to determine the respective MCF. For this purpose the model developed by Mangino et al. (2001), that is also used to derive the 2006 IPCC default values, was adapted to the specific conditions of Switzerland. On a monthly time step, loading of a virtual liquid/slurry manure system was simulated according to the VS excretion of the total livestock herd and the manure management system distribution (MS) in the respective inventory year. Thereby it was assumed that excretion on pasture, range and paddock takes only place during summer

months, i.e. from April to September. Subsequently, monthly manure degradation was forecast using the temperature-dependent van't Hoff-Arrhenius equation with the parametrization as suggested by Mangino et al. (2001). Monthly mean air temperatures for the Swiss central plateau during the 1981-2010 time period were obtained from the Federal Office of Meteorology and Climatology (MeteoSwiss 2014). Minimum temperature in the liquid/slurry system was allowed to drop to 1°C instead of 5°C as proposed in the original model (see e.g. Vergé et al. 2007, Van der Zaag et al. 2013). Any carry-over effect of undegraded manure from one month to the next was neglected (see e.g. Park et al. 2006, Van der Zaag et al. 2013). Finally, an annual methane conversion factor was calculated by dividing the total VS degraded by the total load of VS.

Several authors have found that the simulated MCF-values according to the model described above are unrealistically high (Park et al. 2006, Van der Zaag et al. 2013). Consequently they propose to use a management and design practice factor (MDP factor) to bring the modelled factors into accordance with measurements. Accordingly a MDP factor of 0.8 was applied here as suggested by Mangino et al. (2001). The resulting MCF-values for liquid/slurry systems range from 13.5% to 14.5%. The variation of the MCF along the time series is due to varying shares of manure dropped on pasture, range and paddock. The higher the share of manure dropped on pasture, range and paddock, the lower is the overall MCF (as livestock is only grazing during summer, the relative share of low methane conversion factors during the cold winter month increases when summer grazing time increases).

Anaerobic digestion of animal manure is increasing in Switzerland since the 1990s but is still not widespread (3.7% of all animal manure in 2015). Emissions from the digestion plant itself are reported under source category 5B2 (Anaerobic digestion at biogas facilities) and described in chp. 7.3. However, emissions from manure storage before alimentation into the digester are reported under agriculture. The amount of manure digested anaerobically was estimated based on total energy production (SFOE 2016a) and eight monitoring protocols of agricultural biogas plants (Genossenschaft Ökostrom Schweiz 2014, GES Biogas GmbH 2014). According to the data in the monitoring protocols the total amount of manure entering the plant originated mainly from cattle manure stored as liquid/slurry (57%) and solid storage (23%) and from swine manure stored as liquid/slurry (20%). It is assumed that 22.5% of the liquid/slurry manure is coming from the farm where the biogas plant is located and is hence directly fed into the digester on a daily basis without being stored (Koehli 2014). The respective MCF was thus set to zero. As solid manure usually has a low MCF and is stored for only a short period before being fed into the digester, the respective MCF was also set to zero. The MCF for the remaining liquid/slurry manure that is delivered from neighbouring farms to the biogas plant was estimated with the methodology described in the "Standard method for compensating projects of the type "agricultural biogas plants"" (FOEN 2014n). This method is based on the "Approved small scale baseline and monitoring methodology AMS-III.D./Version 19.0. Methane recovery in animal manure management systems" and relies thus on a generally accepted foundation (UNFCCC 2013c).

According to this methodology the MCF value for conventional liquid/slurry systems given in Table 5-13 is reduced according to the duration of pre-storage before the manure is delivered to the digester:

$$MCF_{PSAD} = MCF_{LS} * \left(\frac{14.49 * (e^{-k*AI_j} - 1)}{AI_j} + 1 \right)$$

MCF_{PSAD} = CH₄ conversion factor for pre-storage of liquid manure before delivery to biogas plants (%)

MCF_{LS} = CH₄ conversion factor for liquid/slurry systems (%)

k = Degradation rate constant (0.069)

AI_j = average pre-storage time period (day)

The average pre-storage time was estimated to be 12 days (Koehli 2014). The resulting weighted average MCF-value for anaerobic digestion varies between 2.6% and 2.8%. Variation is due to the variation of the underlying MCF of liquid/slurry systems.

Fattening calves, sheep, camels, deer and goats are kept in **deep litter systems**. A MCF of 10% was adopted, which is the mean value between the IPCC default values for cattle and swine deep bedding <1 month and >1 month at 10°C (IPCC 2006). The choice of a MCF of 10% for deep litter is supported by the specific feeding and manure management regime in Switzerland (especially cold winter temperatures) and confirmed by a number of studies representative for the country-specific management conditions (Amon et al. 2001, Külling et al. 2002, Külling et al. 2003, Moller et al. 2004, Hindrichsen et al. 2006, Park et al. 2006 and Sommer et al. 2007, Zeitz et al. 2012). For further details see FOEN 2011 (16.5 attachment E).

For all poultry categories a MCF value of 1.5% was used according to the default value for **poultry manure systems** in the 2006 IPCC Guidelines.

Table 5-13 Manure management systems and methane conversion factors (MCFs). Blue: annually changing parameters, value for 2015.

Manure management system		Description	MCF (%)
Pasture		Manure is allowed to lie as it is, and is not managed (distributed, etc.).	1.0
Solid storage		Dung and urine are excreted in a barn. The solids (with and without litter) are collected and stored in bulk for a long time (months) before disposal.	2.0
Liquid/slurry		Combined storage of dung and urine under animal confinements for longer than 1 month.	13.7
Digesters		Storage before alimentation into anaerobic digester. Storage system can be liquid/slurry or solid storage.	2.6
Other	Deep litter	Dung and urine is excreted in a barn with lots of litter and is not removed for a long time (months).	10.0
	Poultry system	Manure is excreted on the floor with or without bedding.	1.5

5.3.2.2.4 Manure management system distribution (MS)

The fraction of animal manure handled using different manure management systems (MS) as well as the percentages of urine and dung deposited on pasture, range and paddock was separately assessed for each livestock category (Table 5-14). The fractions are determined by the livestock husbandry system (e.g. tie stall or loose housing system) as defined in Flisch et al. (2009). Estimation is conducted within the Swiss ammonium model AGRAMMON (Kupper et al. 2013) based on expert judgement and values from the literature (1990, 1995) and on extensive farm surveys (2002, 2007, 2010 and 2015). The data clearly reproduce the shift towards an increased use of pasture, range and paddocks and a decrease in solid storage. The changes of the manure management system distribution reflect the shift to a more animal-friendly livestock husbandry in the course of the agricultural policy reforms during the 1990s and the early 20th century. One of the most important voluntary programs in this context is called "RAUS" and implies at least 156 days of pasture per year (Swiss Confederation, 2008). Accordingly, the share of mature dairy cattle (and other animals) going to pastures increased substantially and the length of stay on the pasture increased by 50%. In the year 2007 78% of the dairy cattle were held on farms participating in the RAUS program. The average number of pasture days (including all farms) in that year was 181, and it was 177 in 2010. It can thus be assumed, that already in the early years of the new millennium most farms accomplished the transition to RAUS and that a new management standard was reached at this point of time, which did not change significantly afterwards.

The amount of manure digested anaerobically was estimated based on total energy production (SFOE 2016a) and eight monitoring protocols of agricultural biogas plants (Genossenschaft Ökostrom Schweiz 2014, GES Biogas GmbH 2014) as described under 5.3.2.2.3.

5.3.2.3 Activity data CH₄

Activity data of all livestock categories covered by the official census was obtained from SBV (2016) and the SFSO (2016). The respective data was revised and harmonized during a joint effort of the Agroscope Reckenholz Tänikon Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012). Additionally to official statistical data, population data of livestock not covered by the agricultural census of the Swiss Federal Statistical Office was assessed (Poncet et al. 2007, Poncet et al. 2009, Schmidlin et al. 2013, SFSO 2016b). For further details and additional data on a livestock sub-category level refer to chp. 5.2.2.3, Table 5-8 as well as Annex 3 A3.3.

Table 5-14 Manure management system distribution (MS) according to the AGRAMMON model. Detailed data on livestock sub-category levels are provided in Annex 3 A3.3.

MS Distribution	1990					1995					2002					
	%					%					%					
	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	
Mature Dairy Cattle	63.7	27.6	8.3	0.4	0.0	65.7	24.4	9.5	0.4	0.0	65.3	16.2	18.0	0.5	0.0	
Other Mature Cattle	41.2	32.1	26.3	0.4	0.0	39.3	34.1	26.2	0.4	0.0	39.8	20.7	39.1	0.5	0.0	
Growing Cattle (weighted average)	47.5	31.7	15.8	0.4	4.5	48.4	30.9	15.9	0.4	4.5	42.1	25.4	27.5	0.5	4.5	
Cattle	Fattening Calves	14.3	0.0	0.0	0.4	85.2	15.0	0.0	0.0	0.4	84.6	21.6	0.0	0.3	0.5	77.6
	Pre-Weaned Calves	41.2	32.1	26.3	0.4	0.0	39.3	34.1	26.2	0.4	0.0	41.2	21.1	37.3	0.5	0.0
	Breeding Cattle 1st Year	37.0	48.5	14.1	0.4	0.0	38.0	47.4	14.2	0.4	0.0	33.7	38.8	27.0	0.5	0.0
	Breeding Cattle 2nd Year	45.3	28.9	25.4	0.4	0.0	47.3	26.7	25.6	0.4	0.0	37.8	23.4	38.4	0.5	0.0
	Breeding Cattle 3rd Year	50.5	29.1	20.0	0.4	0.0	51.4	27.9	20.3	0.4	0.0	42.2	22.5	34.8	0.5	0.0
	Fattening Cattle	70.1	24.1	0.0	0.4	5.5	66.4	27.7	0.0	0.4	5.6	67.3	26.8	2.2	0.5	3.2
Sheep (weighted average)	0.0	0.0	30.1	0.0	69.9	0.0	0.0	30.3	0.0	69.7	0.0	0.0	33.2	0.0	66.8	
Swine (weighted average)	98.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	98.0	0.3	0.1	1.5	0.0	
Buffalo (weighted average)	NA	NA	NA	NA	NA	47.5	26.8	25.6	0.0	0.0	38.1	23.5	38.4	0.0	0.0	
Camels (weighted average)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.0	0.0	33.5	0.0	66.5	
Deer (weighted average)	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	
Goats	0.0	0.0	13.6	0.0	86.4	0.0	0.0	13.6	0.0	86.4	0.0	0.0	12.2	0.0	87.8	
Horses (weighted average)	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	76.1	23.9	0.0	0.0	
Mules and Asses (weighted average)	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	76.9	23.1	0.0	0.0	
Poultry (weighted average)	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.5	0.0	99.5	0.0	0.0	2.6	0.0	97.4	
Rabbits	0.0	100.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	
Livestock NCAC (weighted average)	0.0	93.2	6.8	0.0	0.0	0.0	87.2	7.3	0.0	5.6	0.0	38.5	26.9	0.0	34.5	

MS Distribution	2007					2010					2015					
	%					%					%					
	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid storage	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	
Mature Dairy Cattle	67.5	13.6	17.7	1.3	0.0	67.0	14.3	16.9	1.7	0.0	69.8	10.7	16.2	3.3	0.0	
Other Mature Cattle	49.5	20.2	29.0	1.3	0.0	47.9	17.9	32.4	1.7	0.0	50.6	14.7	31.4	3.3	0.0	
Growing Cattle (weighted average)	45.6	23.9	25.3	1.3	4.0	45.2	25.6	23.4	1.7	4.1	46.9	22.6	23.7	3.3	3.5	
Cattle	Fattening Calves	21.8	0.0	0.2	1.3	76.7	16.9	0.0	0.2	1.7	81.1	22.0	0.0	1.2	3.3	73.4
	Pre-Weaned Calves	50.0	18.6	30.1	1.3	0.0	44.6	32.8	20.9	1.7	0.0	35.8	27.7	33.2	3.3	0.0
	Breeding Cattle 1st Year	41.0	34.5	23.3	1.3	0.0	43.4	33.4	21.5	1.7	0.0	44.7	30.9	21.1	3.3	0.0
	Breeding Cattle 2nd Year	41.4	20.8	36.5	1.3	0.0	43.2	20.8	34.3	1.7	0.0	42.3	19.5	34.8	3.3	0.0
	Breeding Cattle 3rd Year	45.6	21.3	31.8	1.3	0.0	46.3	21.4	30.6	1.7	0.0	54.1	17.0	25.5	3.3	0.0
	Fattening Cattle	62.3	29.2	4.3	1.3	2.9	57.7	33.2	4.0	1.7	3.4	61.4	27.7	4.8	3.3	2.8
Sheep (weighted average)	0.0	0.0	39.2	0.0	60.8	0.0	0.0	33.7	0.0	66.3	0.0	0.0	33.6	0.0	66.4	
Swine (weighted average)	94.5	0.1	1.2	4.1	0.0	94.0	0.3	0.1	5.6	0.0	88.7	0.0	0.0	11.2	0.0	
Buffalo (weighted average)	42.3	21.1	36.5	0.0	0.0	44.5	21.2	34.3	0.0	0.0	44.7	20.5	34.8	0.0	0.0	
Camels (weighted average)	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Deer (weighted average)	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Goats	0.0	0.0	7.1	0.0	92.9	0.0	0.0	10.0	0.0	90.0	0.0	0.0	11.6	0.0	88.4	
Horses (weighted average)	0.0	78.7	21.3	0.0	0.0	0.0	74.4	25.6	0.0	0.0	0.0	80.5	19.5	0.0	0.0	
Mules and Asses (weighted average)	0.0	75.2	24.8	0.0	0.0	0.0	79.3	20.7	0.0	0.0	0.0	77.5	22.5	0.0	0.0	
Poultry (weighted average)	0.0	0.0	3.7	0.0	96.3	0.0	0.0	2.7	0.0	97.3	0.0	0.0	3.0	0.0	97.0	
Rabbits	0.0	100.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	
Livestock NCAC (weighted average)	0.0	34.7	28.7	0.0	36.6	0.0	39.6	27.5	0.0	32.9	0.0	41.6	24.1	0.0	34.2	

5.3.2.4 Emission factors N₂O

Estimation of direct N₂O emissions from manure management relies basically on the same animal waste management systems as the estimation of CH₄ emissions (compare chp. 5.3.2.2). All emission factors are based on default values given in table 10.21 of the 2006 IPCC Guidelines (Table 5-15). For liquid/slurry systems a weighted emission factor was

calculated based on the share of systems with and without natural crust cover. Data on occurrence of natural crusts on slurry tanks of Swiss farms was raised in the census conducted for the Swiss ammonia inventory AGRAMMON (Kupper et al. 2013). Results suggest that formation of thick and permanent natural crusts on slurry tanks is not widespread in Switzerland. The share of systems with crust formation ranges from 0.0 to 7.1% and leads to a N₂O emission factor that ranges from 0.0000 to 0.0004 kg N₂O-N/kg N, respectively.

Table 5-15 Emission factors for calculating N₂O emissions from manure management. Blue: annually changing parameters, value for 2015.

Animal waste management system	Emission factor
	kg N ₂ O-N / kg N
Liquid/Slurry: with natural crust cover	0.005
Liquid/Slurry: without natural crust cover	0.000
Solid storage	0.005
Anaerobic digester	0.000
Cattle and swine deep bedding: no mixing	0.010
Poultry manure	0.001
Indirect emissions due to volatilisation	0.026

The emission factor for indirect N₂O emissions after volatilisation of NH₃ and NO_x from manure management systems was reassessed during a literature review by Bühlmann et al. 2015 and Bühlmann 2014. Due to the fragmented land use in Switzerland, where agricultural land use alternates with natural and semi-natural ecosystems over short distances, the share of volatilised nitrogen that is re-deposited in (semi-)natural habitats is on average higher than 55%. Thus, the assumption made in the 2006 IPCC Guidelines that “a substantial fraction of the indirect emissions will in fact originate from managed land”, cannot be applied to Switzerland. Accordingly, the overall emission factor for indirect emissions was estimated by calculating an area-weighted mean of the indirect emission factor for managed land (i.e. 0.01 based on IPCC 2006) and the indirect emission factor for (semi-)natural land (as provided in Bühlmann 2014). Due to slightly changing land use over the inventory time period, the resulting emission factor shows some small temporal variation around a mean value of 2.55%. Note that the emission factor in cell R37 of CRF Table3.B(b) refers to kg N₂O/kg N instead of kg N₂O-N/kg N.

5.3.2.5 Activity data N₂O

Activity data for N₂O emissions from 3B Manure management was estimated according to equation 10.25 of the 2006 IPCC Guidelines:

$$N_2O_{D(mm)} = \left[\sum_S \left[\sum_T (N_{(T)} \cdot Nex_{(T)} \cdot MS_{(T,S)}) \right] \cdot EF_{3(S)} \right] \cdot \frac{44}{28}$$

N₂O_{D(mm)} = direct N₂O emissions from manure management (kg N₂O/year)

N_(T) = number of head of livestock species/category *T* (head)

N_{ex(T)} = annual average N excretion per head of species/category *T* (kg N/head/year)

MS_(T,S) = fraction of total annual nitrogen excretion for each livestock species/category *T* that is managed in manure management system *S*

EF_{3(S)} = emission factor for direct N₂O emissions from manure management system *S* (kg N₂O-N/kg N)

44/28 = conversion of (N₂O-N)_(mm) emissions to N₂O_(mm) emissions

5.3.2.5.1 Livestock population

Activity data of all livestock categories covered by the official census was obtained from SBV (2016) and the SFSO (2016). The respective data set was revised and harmonized during a joint effort of the Agroscope Reckenholz Tänikon Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012). Additionally to official statistical data, population data of livestock not covered by the agricultural census of the Swiss Federal Statistical Office was assessed (Poncet et al. 2007, Poncet et al. 2009, Schmidlin et al. 2013, SFSO 2016b). For further details and additional data on a livestock sub-category level refer to chp. 5.2.2.3, Table 5-8 as well as Annex 3 A3.3.

5.3.2.5.2 Nitrogen excretion (N_{ex})

Data on nitrogen excretion per animal category (kg N/head/year) is country-specific and was obtained from Kupper et al. (2013) (Table 5-16). These values are based on the "Principles of Fertilisation in Arable and Forage Crop Production" (Flisch et al. 2009). Unlike to the method in the IPCC Guidelines, the age structure of the animals and the different use of the animals (e.g. fattening and breeding) are considered. Standard nitrogen excretion rates are modified within the AGRAMMON model in order to account for changing agricultural structures and production techniques over the years (e.g. milk yield, use of feed concentrates, protein reduced animal feed etc.; Kupper et al. (2013)). This more disaggregated approach leads to considerable lower calculated nitrogen excretion rates compared to IPCC mainly because lower N_{ex}-rates of young animals are considered explicitly.

The nitrogen excretion rates are given on an annual basis, considering replacement of animals (growing cattle, swine, poultry, rabbits) and including excretions from corresponding offspring and other associated animals (sheep, deer, goats, swine, rabbits) (ART/SHL 2012).

As an exception, nitrogen excretion of **mature dairy cattle** was separately calculated within the same feeding model that was used for CH₄ emissions from 3A Enteric fermentation and from 3B Manure management (Agroscope 2014c, see also chp. 5.2.2.2). Nitrogen excretion of mature dairy cattle is dependent on milk production and feed properties. In the year 2003 yearly milk yield surpassed 6000 kg. To achieve yearly milk yields higher than 6000 kg the cows have to be fed with an increasing share of feed concentrates. Due to the energy dense feed concentrates, the ratio between net energy content and protein content increases. Since 2003 the increase in nitrogen excretion rate is thus slower than before, although milk yield increased more or less at the same rate from 1990 to 2015.

Sheep in Switzerland are fed mainly on roughage from extensive pasture and meadows (Flisch et al. 2009) and are estimated to excrete approximately 8.0 kg N per head and year. This is considerably lower than IPCC default. However, nitrogen excretion is averaged over the whole population, of which roughly 40% are lambs and other immature animals. **Swine** show a significant decrease in nitrogen excretion rates until 2006, which can be explained by the increasing use of protein-reduced fodder (Kupper et al. 2013).

Table 5-16 Nitrogen excretion rates of Swiss livestock, 1990–2015. The complete time series on a livestock sub-category level are provided in Annex 3 A3.3.

Nitrogen Excretion	1990-2011									
	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011
	kg N/head/year									
Mature Dairy Cattle	102.7	105.5	109.7	113.2	113.3	113.9	114.2	114.7	114.9	115.0
Other Mature Cattle	80.0	80.0	80.0	80.0	80.0	80.0	80.0	80.0	80.0	80.0
Growing Cattle (weighted average)	33.1	33.4	33.6	33.1	33.2	33.2	33.2	33.4	33.4	33.4
<i>Fattening Calves</i>	13.0	13.0	13.0	13.0	13.0	13.0	13.0	13.0	13.0	13.0
<i>Pre-Weaned Calves</i>	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0	34.0
<i>Breeding Cattle 1st Year</i>	25.0	25.0	25.0	25.0	25.0	25.0	25.0	25.0	25.0	25.0
<i>Breeding Cattle 2nd Year</i>	40.0	40.0	40.0	40.0	40.0	40.0	40.0	40.0	40.0	40.0
<i>Breeding Cattle 3rd Year</i>	55.0	55.0	55.0	55.0	55.0	55.0	55.0	55.0	55.0	55.0
<i>Fattening Cattle</i>	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0	33.0
Sheep (weighted average)	7.5	7.6	8.1	8.1	8.2	8.3	8.2	8.5	8.5	8.5
Swine (weighted average)	13.4	12.8	10.5	9.4	9.2	9.1	9.2	9.2	9.2	9.1
Buffalo (weighted average)	NA	37.2	41.1	38.7	38.0	34.9	34.4	36.5	37.3	38.4
Camels (weighted average)	NA	NA	14.1	12.8	12.8	12.8	12.8	12.8	12.6	12.7
Deer (weighted average) ¹⁾	20.0	21.9	22.3	21.9	22.1	22.1	22.4	22.5	22.4	22.4
Goats	10.5	10.4	10.6	10.5	10.6	10.5	10.5	10.7	10.6	10.8
Horses (weighted average)	43.6	43.5	43.6	43.7	43.7	43.7	43.7	43.7	43.7	43.7
Mules and Asses (weighted average)	15.7	15.7	15.7	15.7	15.7	15.7	15.7	15.7	15.7	15.7
Poultry (weighted average)	0.6	0.5	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Rabbits	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Livestock NCAC (weighted average)	38.8	33.8	14.5	12.5	12.4	12.5	12.2	13.1	14.3	14.8

Nitrogen Excretion	2012-2015			
	2012	2013	2014	2015
	kg N/head/year			
Mature Dairy Cattle	114.9	114.7	115.2	115.3
Other Mature Cattle	80.0	80.0	80.0	80.0
Growing Cattle (weighted average)	33.3	33.3	33.4	33.5
<i>Fattening Calves</i>	13.0	13.0	13.0	13.0
<i>Pre-Weaned Calves</i>	34.0	34.0	34.0	34.0
<i>Breeding Cattle 1st Year</i>	25.0	25.0	25.0	25.0
<i>Breeding Cattle 2nd Year</i>	40.0	40.0	40.0	40.0
<i>Breeding Cattle 3rd Year</i>	55.0	55.0	55.0	55.0
<i>Fattening Cattle</i>	33.0	33.0	33.0	33.0
Sheep (weighted average)	8.5	8.6	8.5	9.6
Swine (weighted average)	9.0	9.0	8.9	8.9
Buffalo (weighted average)	36.9	36.9	36.4	36.4
Camels (weighted average)	12.8	12.9	12.8	12.7
Deer (weighted average) ¹⁾	22.5	23.0	23.0	23.0
Goats	10.8	10.9	10.9	12.6
Horses (weighted average)	43.7	43.8	43.8	43.8
Mules and Asses (weighted average)	15.7	15.7	15.7	15.7
Poultry (weighted average)	0.5	0.5	0.5	0.5
Rabbits	1.0	1.0	1.0	1.0
Livestock NCAC (weighted average)	15.3	15.5	15.6	16.5

¹⁾ Deer: Excretion per animal place

5.3.2.5.3 Manure management system distribution (MS)

The split of nitrogen flows into the different animal waste management systems and its temporal dynamics are based on the respective analysis in the AGRAMMON model (Kupper et al. 2013). The distribution is consistent with the allocation of volatile solids used for the calculation of CH₄ emissions from 3B Manure management (for further information refer to chp. 5.3.2.2.4).

5.3.2.5.4 Volatilisation of NH₃ and NO_x from manure management systems

Indirect N₂O emissions from the deposition of NH₃ and NO_x volatilised from manure management are considered. Losses of ammonia from stables and manure storage systems to the atmosphere are calculated according to the Swiss ammonia model AGRAMMON (Kupper et al. 2013). Specific loss-rates for all major livestock categories are estimated based on agricultural structures and techniques (e.g. stable type, manure management system, measures to reduce NH₃ emissions). Accordingly, the overall fraction of nitrogen volatilised underlies certain temporal dynamics that can be explained by changes in agricultural management practices (e.g. the transition to more animal friendly housing systems). It ranges from 15.3 to 20.8%.

For the volatilisation of NO_x default values from the EMEP/EEA air pollutant emission inventory guidebook 2013 were used, assuming that 50% and 25% of the nitrogen is present in the form of TAN (total ammonia nitrogen) in liquid/slurry and solid storage systems respectively (EMEP/EEA 2013). Accordingly, it is estimated that 0.005% and 0.25% of the total nitrogen in liquid/slurry and solid storage systems are lost to the atmosphere. In this context the management systems “anaerobic digestion” and “deep litter” are treated as liquid/slurry- and solid storage systems respectively.

Note that volatilisation from pasture, range and paddock manure is included under 3Db (Indirect N₂O emissions from managed soils). A graphical overview of the nitrogen flow system is given in Figure 5-6 and respective numbers are provided in Table 5-20.

5.3.3 Uncertainties and time-series consistency

For the uncertainty analysis the input data from ART (2008a) were used and were updated with current activity and emission data as well as with new default uncertainties from the 2006 IPCC Guidelines. The arithmetic mean of the lower and upper bound was used for activity data and for emission factors in the Approach 1 analysis (Table 5-17).

For the Approach 2 analysis, asymmetric probability distributions as well as possible correlations of input data were considered.

For further results also consult chp. 1.6.1 and Table A – 7.

Table 5-17 Uncertainties for 3B Manure management 2015. (AD: Activity data; EF: Emission factor; CO: Combined).

Uncertainty 3B		Approach 1			Approach 2		
		AD	EF	CO	low	high	mean
		%			%		
CH ₄		6.4	54.1	54.4	-55	54	55
N ₂ O direct	Liquid/slurry / Anaerobic digester	31.9	75.0	81.5	-77	88	83
N ₂ O direct	Solid storage / Deep bedding	31.9	75.0	81.5	-76	89	83
N ₂ O indirect	Indirect emissions	46.5	240.0	244.5	-77	126	102

The time series 1990–2015 are all considered consistent, although the following issues should be considered:

- For time series consistency of livestock population data and gross energy intake see chp. 5.2.3.
- The MCF for liquid/slurry systems varies according to the development of the grazing management over the years as described under chp. 5.3.2.2.3.
- Input data from the AGRAMMON-model is available for the years 1990 and 1995 (expert judgement and literature) as well as for 2002, 2007, 2010 and 2015 (extensive surveys on approximately 3000 farms). Values in-between the assessment years were interpolated linearly.
- The emission factor for indirect N₂O emissions after volatilisation of NH₃ and NO_x from manure management systems varies according to varying land use as described in chp. 5.3.2.4.

5.3.4 Category-specific QA/QC and verification

General QA/QC measures are described in NIR chp. 1.2.3.

All further category-specific QA/QC activities are described in a separate document (Agroscope 2016b). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed. Furthermore, comparisons with data from other countries were conducted and discussed where possible. Agroscope 2016b is continuously updated with the most recent inventory data.

For quality of livestock population data and livestock energy intake consult chp. 5.2.4.

5.3.4.1 QA/QC and verification CH₄

IPCC tables with data for estimating emission factors of all livestock categories (such as weight, feed digestibility, maximum CH₄ producing capacity (B₀) or daily excretion of volatile solids) were filled in, checked for consistency and confidence and compared with IPCC default values (refer to Annex 3 A3.3).

VS excretion of various animal categories is based on IPCC default values. A cross check of these estimate was conducted during the 2016 submission. VS excretion of the total livestock population was estimated by using exclusively equation 10.24 of the 2006 IPCC Guidelines and GEI data for all animal categories. Using this approach, total VS excretion for the year 2014 was 4.1% higher than reported in the Swiss GHG inventory. Most of the discrepancy can be attributed to swine, for which the default value for VS excretion is rather low (i.e. 0.31 kg/head/day as weighted mean for 2014 compared to 0.43 kg/head/day from the approach based on equation 10.24). However, Minonzio et al. 1998 also suggest a low VS-excretion of 0.30 kg/head/day on average, based on the Swiss typical feeding recommendations. They assume a digestibility of the organic matter of 83%. Using this value in IPCC equation 10.24 would also yield a VS-excretion of 0.31 kg/head/day. This finding supports the adoption of the IPCC default VS-excretion for swine. As for swine, equation 10.24 yields higher VS-excretion values for sheep and goats. Also in these cases the default

values for feed digestibility (i.e. 60%) might be too low for Swiss specific conditions. In summary there is no clear indication that the approach using exclusively equation 10.24 would result in a better estimate of overall VS excretion. As for some of the parameters used in equation 10.24 (such as e.g. feed digestibility for swine) no reliable country-specific data was available, it was thus decided to still use the IPCC default values for VS excretion of the animal categories concerned.

Factors for methane conversion (MCF) and manure management system distribution (MS) were analysed considering the national agricultural context. The estimated MCF-values for liquid/slurry systems in Switzerland are lower than the IPCC default value for liquid/slurry system, without natural crust cover, at a temperature $\leq 10^{\circ}\text{C}$. However, a relatively low MCF is supported by the fact that more than 80% of all liquid/slurry storage tanks are covered and approximately one third of the remaining tanks have a surface crust (Kupper et al. 2013). Furthermore, a series of laboratory measurements of MCF-values by the group of animal nutrition from the Swiss Federal Institute of Technology in Zürich yielded consistently low MCF-values (Zeit et al. 2012).

During the past years studies were conducted to verify methane emissions at the regional scale comparing bottom-up estimates with atmospheric measurements (Bamberger et al. 2014, Henne et al. 2015, Hiller et al. 2014, Hiller et al. 2014a, Stieger 2013, Stieger et al. 2015). For further information on these studies see chp. 5.2.4.

5.3.4.2 QA/QC and verification N₂O

N₂O estimation is based on the Swiss ammonium emission model AGRAMMON that is documented in Kupper et al. (2013).

All relevant data needed for the calculation of N₂O emissions such as nitrogen excretion rates, manure management system distribution and N₂O emission factors were checked for consistency and were compared to the corresponding values of other countries and to the IPCC default value if available (Agroscope 2016b).

As one of the most important parameters, nitrogen excretion rates were analysed in more detail. For mature dairy cattle, all model inputs were compared to available data on feeding regimes as applied in the field (e.g. rations composition, amount of feed concentrates and silage, nitrogen content of the feed). Furthermore, modelled values were compared to measurements of feeding trials of the animal nutrition group of the Swiss Federal Institute of Technology in Zürich. Measurements were on average almost 30 kg/head/year lower than modelled values. However, nitrogen intake as well as nitrogen losses through milk were in very good accordance supporting the excretion rates. It is thus most likely, that some of the nitrogen excreted was lost (e.g. volatilised) before manure could be collected and nitrogen could be stabilised for measurements (see e.g. Van Dorland et al. 2007). In order to validate the total nitrogen excretion of the whole livestock population a cross check was conducted comparing the bottom up inventory estimates with an independent top down approach. Thereby, the total amount of nitrogen contained in animal livestock products such as meat, milk or eggs (output) was subtracted from the total amount of nitrogen in animal feedstuff produced in or imported to the country (input). Under the condition that the nitrogen pool in the animal population remains constant, the result should be equal to the amount of nitrogen excreted in the manure (e.g. Spiess 2011). Accordance was good (average discrepancy of

±2%) for the years 1990 – 2005. However, for later years the top down estimates were on average 10% higher than the bottom up estimates. Reasons for this behaviour are not yet clear and this finding will be subject to further analysis. N_{ex} -values for the most important animal categories (mature dairy cattle and swine, being responsible for 65% of total nitrogen excretion) were compared to the values of the alternative gross energy approach suggested in equation 10.32 in the 2006 IPCC Guidelines. For swine, the IPCC approach estimated on average 18% lower N_{ex} values for the years 1990–2004. This is probably due to an underestimation of the feed protein content in this model calculation and the inventory estimates are considered more realistic. Differences were smaller than 3% for years after 2005. All QA/QC checks of the N_{ex} values are further elaborated in Agroscope 2016b.

5.3.5 Category-specific recalculations

General information on recalculations is provided in chp. 10.

- The excretion of VS of horses was revised due to a respective recommendation during the annual review in 2016 (UNFCCC 2017; A6). New values are based on equation 10.24 in the 2006 IPCC Guidelines. New emission estimates are slightly higher (+ 0.44 kt CO₂ eq. on average, 1990: +0.36 kt CO₂ eq., 2014: +0.48 kt CO₂ eq.)
- The amount of VS in the manure management system anaerobic digesters was revised for the year 2014 due to updated estimates of the respective AD in the waste sector. Overall CH₄ and N₂O emissions increased by 0.7 kt CO₂ equivalents.
- All estimates based on AGRAMMON data were recalculated for the years 2011–2014 due to a new interpolation after new survey results for 2015 became available. The impact on overall emissions in kt CO₂ equivalents is: 2011: 0.4; 2012: 0.9; 2013: 1.4; 2014: 1.9.
- For the years 2011–2014, the MCF for liquid/slurry systems was slightly revised due to new model runs based on the new AGRAMMON data. Effects on overall CH₄ emissions in kt CO₂ equivalents are: 2011: 0.5; 2012: 0.9; 2013: 1.3; 2014: 1.7.

5.3.6 Category-specific planned improvements

Planned improvements for future submissions are the further development, adaptation and verification of the dairy cow feeding model (GE, DE, VS-excretion, N-excretion).

New country-specific values for N_{ex} for most livestock categories will become available in the next years due to a revision of the “Principles of Fertilization in Arable and Forage Crop Production (Flisch et al. 2009)”.

A release of a new version of the AGRAMMON-Model is planned for 2017. If possible the respective AGRAMMON projections will be included during the next GHG inventory submission.

NMVOC emissions will be revised in submission 2018 and separately reported for 3B Manure management and 3D Agricultural soils.

5.4 Source category 3C – Rice cultivation

Rice cultivation is of minor importance in Switzerland. The agricultural land used for rice cultivation and the annual yield of rice are not estimated by the Swiss Farmers Union (SBV 2016). There is only some insignificant upland rice cultivation in the southern part of Switzerland. CH₄ emissions are assumed to be zero. They are therefore not considered in the emission calculation.

5.5 Source category 3D – Agricultural soils

5.5.1 Source category description

Table 5-18 Key categories (KCA incl. LULUCF) of 3B Manure management

Code	IPCC Category	GHG	Identification Criteria
3Da	Direct emissions from managed soils	N ₂ O	L1, L2
3Db	Indirect emissions from managed soils	N ₂ O	L1, L2, T1, T2

The source category 3D includes direct and indirect N₂O emissions from managed soils (Table 5-19). Direct emissions are further subdivided in emissions from 1. Inorganic N fertilisers, 2. Organic N fertilisers, 3. Urine and dung deposited by grazing animals, 4. Crop residues, 5. Mineralisation/immobilisation associated with loss/gain of soil organic matter, 6. Cultivation of organic soils (i.e. histosols) and 7. Other (i.e. Domestic synthetic fertilisers). Indirect N₂O emissions are further subdivided in 1. Atmospheric deposition and 2. Nitrogen leaching and run-off. All indirect N₂O emissions after deposition of NO_x and NH₃ or after leaching of NO₃⁻ are reported under source category 3Db Indirect N₂O Emissions from managed soils. This includes indirect N₂O emissions after NO₃⁻ leaching from N mineralisation in cropland remaining cropland and grassland remaining grassland. To avoid double counting the respective emissions are not reported under source category 4(IV) Indirect N₂O emissions from managed soils or in CRF Table6 “Indirect emissions of N₂O and CO₂” (see also chp. 9).

Table 5-19 Specification of source category 3D Agricultural soils.

3D	Source	Specification
3Da	Direct N ₂ O emissions from managed soils	<ol style="list-style-type: none"> 1. Inorganic N fertilisers 2. Organic N fertilisers (animal manure applied to soils, sewage sludge applied to soils, other organic fertilisers applied to soils) 3. Urine and dung deposited by grazing animals 4. Crop residues (incl. residues from meadows and pasture) 5. Mineralisation/immobilisation associated with loss/gain of soil organic matter 6. Cultivation of organic soils (i.e. histosols) 7. Other (domestic synthetic fertilisers)
3Db	Indirect N ₂ O emissions from managed soils	<ol style="list-style-type: none"> 1. Atmospheric deposition 2. Nitrogen leaching and run-off

Furthermore, NO_x emissions from managed soils as well as NMVOC emissions are estimated.

Direct and indirect N₂O emissions from managed soils have decreased since 1990 in almost all major sub-categories. Only N₂O emissions from 3Da3 (Urine and dung deposited by grazing animals) increased due to a higher share of manure excreted on pasture, range and paddock. NO_x emissions have declined by 21% since 1990. The general trends can be explained by a reduction in the number of cattle and a reduced input of mineral fertilisers due to the introduction of the "Proof of Ecological Performance (PEP)" requiring a balanced fertiliser management (Agroscope 2016b, Leifeld and Fuhrer 2005). Major changes occurred mainly in the 1990's while most emissions were more or less stable after the year 2000.

The most significant N₂O emission sources are animal manure applied to soils (Ø 27%), nitrogen input from atmospheric deposition (Ø 23%), inorganic nitrogen fertilisers (Ø 15%) and urine and dung deposited by grazing animals (Ø 11%).

5.5.2 Methodological issues

5.5.2.1 Methodology

For the calculation of N₂O emissions from 3D Agricultural soils a country-specific Tier 2 method was applied that is based on the IULIA model from Schmid et al. (2000). IULIA is an IPCC-derived method for the calculation of N₂O emissions from agriculture that basically uses the same emission factors, but adjusts the activity data to the particular situation of Switzerland. IULIA is continuously updated. New values for nitrogen excretion rates, manure management system distribution and ammonium emission factors from the Swiss ammonium model AGRAMMON were adopted (Kupper et al. 2013). Furthermore, the updated version of the "Principles of Fertilisation in Arable and Forage Crop Production" (GruDAF; Flisch et al. 2009) was used instead of obsolete data from FAL/RAC 2001 and Walther et al. 1994. Most recently, additional livestock categories, new emission factors for indirect N₂O emissions from atmospheric deposition, new estimates for nitrogen leaching and run-off as well as new NO_x emission factors were implemented.

The modelling of the N₂O emissions is conducted in Agroscope (2017) and is consistent with source category 3B N₂O emissions from manure management. The model structure is displayed in Figure 5-6 and the corresponding amounts of nitrogen are given in Table 5-20.




Figure 5-6 Diagram depicting the methodology of the approach to calculate the N₂O emissions in agriculture (red arrows). Black frames and the respective letters refer to the nitrogen flows in Table 5-20. Note that the figure shows explicitly the methodology of the approach and not necessarily the physical nitrogen flows.

Table 5-20 Nitrogen flows of the N-flow-model for Swiss agriculture. Letters refer to the letters in Figure 5-6. Processes refer to the nitrogen flows in the black frames in Figure 5-6 from left to right or from top to bottom.

	Process	Amount of N			CRF table
		1990	2015		
		tN			
A	1 Pasture, range and paddock	13'578	23'062	= B	3.Da3
	2 Liquid/slurry systems	91'925	77'008		3.B(b)
	3 Solid storage	35'913	17'931		3.B(b)
	4 Other AWMS	8'430	15'313		3.B(b)
	5 Commercial fertiliser	75'339	50'577		= F
B	1 Pasture, range and paddock	13'578	23'062	= A1-A4	3.Da3
	2 NH ₃ volatilisation housing	11'347	15'428		3.B(b)5
	3 N ₂ O emission liquid/slurry	0	23		3.B(b)
	4 NO _x volatilisation liquid/slurry and digester	5	4		3.B(b)5
	5 Leaching manure management	0	0		3.B(b)5
	6 Manure applied to soils	115'195	87'335		3.Da2
	7 N ₂ O emission solid storage	180	90		3.B(b)
	8 N ₂ O emission other AWMS	46	53		3.B(b)
	9 NO _x volatilisation solid storage and deep litter	109	71		3.B(b)5
	10 NH ₃ volatilisation storage	9'388	7'248		3.B(b)5
C	1 NH ₃ deposition manure management	20'734	22'676	= B2+B10	3.B(b)5
	2 NO _x deposition manure management	114	75	= B4+B9	
	3 Leaching manure management	0	0	= B5	
D	1 Plant available N PR&P	9'771	17'190	= B1	3.Da3
	2 N ₂ O emission PR&P	260	439		
	3 NO _x volatilisation PR&P	75	127		
	4 NH ₃ volatilisation PR&P	674	1'192		
	5 Leaching and run-off PR&P	2'798	4'114		
E	1 Plant available N animal manure	58'637	50'873	= B6	3.Da2
	2 N ₂ O emission application animal manure	1'152	873		
	3 NO _x volatilisation application animal manure	634	480		
	4 NH ₃ volatilisation application animal manure	31'034	19'528		
	5 Leaching and run-off application animal manure	23'740	15'580		
F	1 Plant available N com. fertiliser	54'050	38'337	= A5	3.Da1,2,7
	2 N ₂ O emission application com. fertiliser	753	506		
	3 NO _x volatilisation application com. fertiliser	414	278		
	4 NH ₃ volatilisation application com. fertiliser	4'595	2'433		
	5 Leaching and run-off application com. fertiliser	15'526	9'023		
G	1 Cultivation of organic soils (ha)	18'039	17'363		3.Da6
H	1 Mineralisation/immobilisation soil organic matter	620	748		3.Da5
I	1 N in crop residues pasture, range and paddock	21'689	21'604		3.Da4
	2 N in crop residues arable crops	11'347	10'479		
J	1 NH ₃ volatilisation agricultural area	2'134	2'099		
K	1 NH ₃ volatilisation alpine area	269	237		
L	1 NH ₃ deposition fertiliser appl. and PR&P	36'303	23'153	= D4+E4+F4	3.Db1
	2 NO _x deposition fertiliser appl. and PR&P	1'123	885	= D3+E3+F3	
	3 NH ₃ deposition agricultural and alpine area	2'403	2'336	= J+K	
	4 Leaching and run-off fertiliser appl. and PR&P	42'064	28'717	= D5+E5+F5	3.Db2
	5 Leaching and run-off mineralisation SOM	128	133		
	6 Leaching and run-off crop residues	6'808	5'723		

5.5.2.2 Direct N₂O emissions from managed soils (3Da)

Calculation of Direct N₂O emissions from managed soils is based on IPCC 2006 equation 11.2 including six terms for activity data and three different emission factors:

$$N_2O_{Direct-N} = (F_{SN} + F_{ON} + F_{CR} + F_{SOM}) \cdot EF_1 + F_{OS} \cdot EF_2 + F_{PRP} \cdot EF_3$$

N₂O_{Direct-N} = annual direct N₂O–N emissions produced from managed soils (kg N₂O–N/year)

F_{SN} = annual amount of synthetic fertiliser N applied to soils (kg N/year)

F_{ON} = annual amount of animal manure, compost, sewage sludge and other organic N additions applied to soils (kg N/year)

F_{CR} = annual amount of N in crop residues, including N-fixing crops, returned to soils (kg N/year)

F_{SOM} = annual amount of N in mineral soils that is mineralised, in association with loss of soil C from soil organic matter as a result of changes of land use or management (kg N/year)

F_{OS} = annual area of managed/drained organic soils (ha)

F_{PRP} = annual amount of urine and dung N deposited by grazing animals on pasture, range and paddock (kg N/year)

EF₁ = emission factor for N₂O emissions from N inputs (kg N₂O–N/kg N input)

EF₂ = emission factor for N₂O emissions from drained/managed organic soils (kg N₂O–N/ha/year)

EF₃ = emission factor for N₂O emissions from urine and dung N deposited on pasture, range and paddock by grazing animals (kg N₂O–N/kg N input)

5.5.2.2.1 Emission factors

Emission factors for calculating 3Da Direct N₂O emissions from managed soils are all based on default values as provided in the 2006 IPCC Guidelines (Table 5-21). Due to the lack of data no fertiliser specific emission factors were applied for EF₁. The emission factor for urine and dung deposited by grazing animals was calculated as the weighted mean between the emission factor for cattle, poultry and pigs (EF_{3PRP, CPP} = 0.02) and the emission factor for sheep and “other animals” (EF_{3PRP, SO} = 0.01) according to the shares of nitrogen excreted by the respective animals.

Table 5-21 Emission factors for calculating direct N₂O emissions from managed soils (IPCC 2006). Blue: annually changing parameters, value for 2015.

Emission Source	Emission factor
EF ₁ Inorganic N fertilisers (kg N ₂ O-N/kg)	0.0100
EF ₁ Organic N fertilisers (kg N ₂ O-N/kg)	0.0100
EF ₁ Crop residue (kg N ₂ O-N/kg)	0.0100
EF ₁ Mineralisation/immobilisation soil organic matter (kg N ₂ O-N/kg)	0.0100
EF ₁ Other (domestic synthetic fertilisers) (kg N ₂ O-N/kg)	0.0100
EF ₂ Cultivation of organic soils (kg N ₂ O-N/ha)	8.0000
EF ₃ Urine and dung deposited by grazing animals (kg N ₂ O-N/kg)	0.0190

5.5.2.2.2 Activity data

Activity data for calculation of 3Da Direct soil emissions includes 1. Inorganic N fertilisers, 2. Organic N fertilisers, 3. Urine and dung deposited by grazing animals, 4. Crop residues, 5. Nitrogen from mineralisation/immobilisation associated with loss/gain of soil organic matter 6. Area of organic soils (i.e. histosols) and 7. Other (i.e. Domestic inorganic fertilisers).

Emissions from **inorganic nitrogen fertilisers** include urea and other mineral fertilisers (mainly ammonium-nitrate). The amount of nitrogen input due to these fertilisers is obtained from Agricura (2015). Fertiliser statistics are based on sales statistics of the compulsory storekeepers of fertilisers (Pflichtlagerhalter) and small importers. Agricura conducts plausibility checks with import-data received by the Directorate General of Customs (Oberzolldirektion). It is estimated that 4% of the mineral fertilisers are used for non-agricultural purposes (Kupper et al. 2013). These fertilisers are used in public green areas, sports grounds and home gardens. In the reporting tables (CRF) they are reported under 3Da7 **Other (Domestic inorganic fertilisers)** while emission calculation is conducted together with 3Da1. In some occasions, as for instance for the estimation of indirect N₂O emissions from managed soils, the sum of urea, other mineral fertilisers, sewage sludge, other organic fertilisers and domestic fertilisers is referred to as “commercial fertilisers” (see also Figure 5-6 and Table 5-20).

Organic nitrogen fertilisers include animal manure, sewage sludge and other organic fertilisers. The amount of nitrogen in **animal manure applied to soils** is calculated according to the methods described in chp. 5.3.2.5. As suggested in chp. 10.5.4. and equation 10.34 of the 2006 IPCC Guidelines, all nitrogen excreted on pasture, range and paddock as well as all nitrogen volatilised prior to final application to managed soils is subtracted from the total excreted manure (for the estimation of the respective N-volatilisation during manure management see chp. 5.3.2.5, compare also Figure 5-6 and Table 5-23). Fra_{CGASM} in CRF Table3.D represents the amount of nitrogen volatilised as NH₃, NO_x and N₂O from housing and manure storage divided by the manure excreted in the stable (liquid/slurry, solid storage, digesters, deep litter and poultry manure). The nitrogen input from manure applied to soils under 3Da2a in CRF Table3.D can thus be calculated with the numbers given in CRF Table3.B(b) and 3.D. Nitrogen from bedding material was not accounted for under animal manure applied to soils. The respective nitrogen is included in the nitrogen returned to soils as crop residues.

The amount of **sewage sludge** applied to agricultural soils was estimated according to Kupper et al. (2013). Since 2003 the use of sewage sludge as fertiliser is prohibited in Switzerland. However, a transition period applies for some areas. Cantons could therefore prolong this period until 2008 in individual cases (UVEK 2003). **Other organic fertilisers** include compost as well as liquid and solid digestates from biogas plants and are also estimated according to Kupper et al. (2013). Additionally nitrogen input through co-substrats in agricultural biogas plants is accounted for under this sub-category.

Calculation of emissions from **urine and dung deposited by grazing animals** is based on equation 11.5 of the 2006 IPCC Guidelines. Estimation of total livestock nitrogen excretion was described under 5.3.2.5. The share of manure nitrogen excreted on pasture, range and paddock was estimated according to the AGRAMMON-model (Kupper et al. 2013; Table 5-14). For each livestock category the share of animals that have access to grazing, the number of days per year they are actually grazing as well as the number of hours per day grazing takes place was assessed. Estimates are based on values from the literature and expert judgement (1990, 1995) and on surveys on approximately 3000 farms (2000, 2007, 2010, 2015).

N₂O emissions from **crop residues** are based on the amount of nitrogen in crop residues returned to soil. For **arable crops**, data on total annual crop yields were adopted from the statistical yearbooks of the Swiss Farmers Union (SBV 2016). Subsequently the relationship between nitrogen returned in crop residues and fresh matter crop yield was determined for each crop and hereafter the overall amount of nitrogen returned to soils was calculated as follows:

$$F_{CR,AC} = \sum_T \left(Y_T \cdot \frac{NR_T}{SY_T} \right)$$

$F_{CR,AC}$ = amount of nitrogen in crop residues from arable crops returned to soils (t N)

Y_T = amount of fresh matter crop yield for crop T (t)

NR_T = standard amount of nitrogen in crop residues for crop T (dt/ha)

SY_T = standard amount of fresh matter crop yield for crop T (dt/ha)

Standard values for fresh matter crop yields and nitrogen contained in crop residues are given in the "Principles of Fertilisation in Arable and Forage Crop Production" (FAL/RAC 2001 and Flisch et al. 2009). For sugar beet and fodder beet it is assumed that 10% of the crop residues are removed from the fields for animal fodder.

Crop residues from **meadows and pastures** were also assessed. Two third of the agricultural land consists of grassland which underscores the importance of this source for Switzerland. According to the 2006 IPCC Guidelines (chp. 11.2.1.3) crop residues on pastures should be included in the estimation of N₂O emission from agricultural soils only for years when renewal of pastures happened. However, the area of meadows and pastures applied here refers to permanent grassland (in contrast to leys and intensive meadows). Renewal of these grasslands is not common practice in Switzerland. Crop residues from meadows and pasure therefore refer here only to field losses from feed not eaten by the animals and feed losses due to trampling effects.

$$F_{CR,MP} = \sum_P \left(A_P \cdot \frac{SY_{DM,P}}{10} \cdot N_{DM,P} \div 1000 \cdot R_P \right)$$

$F_{CR,MP}$ = amount of nitrogen in crop residues from meadows and pastures returned to soils (t N)

A_P = area of meadow and pasture of type P (ha)

$SY_{DM,P}$ = standard dry matter yield per area of meadow and pasture of type P (dt/ha)

$N_{DM,P}$ = dry matter nitrogen content of meadow and pasture of type P (kg/t)

R_P = ratio of residues to harvested yield for meadows and pasture of type P (kg/kg)

Areas of intensive meadows, natural meadows, pasture and alpine and Jurassic pasture were obtained from SBV (2016) and from the SFSO (2016). Standard dry matter yields per area, nitrogen content of dry matter as well as % yield losses were based on the original IULIA model (Schmid et al. 2000).

Estimated values of total crop production, nitrogen incorporated with crop residues $F_{(CR)}$, residue/crop ratio, dry matter fraction of residues and nitrogen content of residues are provided in Annex 3 A3.3.

Assessment of nitrogen **mineralisation/immobilisation associated with loss/gain of soil organic matter** was conducted based on data from the LULUCF sector. For reasons of consistency, losses and gains of soil organic matter on cropland and grasslands were accounted for. The same methodology as described under 6.10.2 was applied. Nitrogen mineralisation was estimated by dividing the carbon loss on cropland remaining cropland and grassland remaining grassland with a C/N-ratio of 9.8 according to Leifeld et al. (2007). It should be noted that the carbon losses were assessed based on land use changes on a sub-category level. Only land use changes that led to a net carbon stock loss were considered, excluding land use changes that led to a net carbon stock increase. Consequently, the carbon losses used here are not identical with the net carbon stock changes reported in the reporting tables (CRF). N₂O emissions from nitrogen mineralisation of land converted to cropland or land converted to grassland are reported under source category 4(III) "Direct nitrous oxide (N₂O) emissions from nitrogen (N) mineralisation/immobilisation associated with loss/gain of soil organic matter resulting from change of land use or management of mineral soils".

Estimates of N₂O emissions from **cultivated organic soils** are based on the area of cultivated organic soils and the IPCC default emission factor for N₂O emissions from cultivated organic soils (IPCC 2006). The area of cultivated organic soils corresponds to the total area of organic soils under cropland and grassland as reported in CRF Table 4.B and 4.C (see also 6.2.2).

The relevant activity data for calculating N₂O emissions from soils is displayed in Table 5-22. Additional information is given in Annex 3 A3.3.

Table 5-22 Activity data for calculating 3Da Direct N₂O emissions from managed soils (1990–2015).

Activity Data		1990-1999									
		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		t N/yr									
1. Inorganic N fertilisers	Urea	17'000	12'500	12'171	11'842	11'514	11'185	10'856	7'900	6'698	6'996
	Other mineral fertilisers	49'912	54'604	54'741	50'366	47'046	47'375	45'592	40'964	42'358	44'364
2. Organic N fertilisers	a. Animal manure	115'195	113'859	111'200	109'516	108'361	106'426	103'795	99'670	97'604	93'591
	b. Sewage sludge	4'815	4'840	4'866	4'891	4'916	4'942	4'624	4'307	3'990	3'673
	c. Other organic fertilisers	824	935	1'046	1'157	1'268	1'380	1'421	1'457	1'494	1'531
3. Urine and dung deposited by grazing animals		13'578	13'860	13'937	13'899	14'147	14'333	16'222	17'532	18'871	19'729
4. Crop residues	Arable crops	11'347	11'181	11'067	11'263	10'649	10'854	12'161	11'759	11'820	10'572
	Residues PR&P	21'689	21'717	21'809	21'700	21'372	21'744	21'837	21'911	21'949	21'898
5. Min./imm. associated with loss/gain of SOM		620	615	616	595	676	660	682	667	741	747
6. Cultivation of organic soils (ha)		18'039	18'014	17'989	17'964	17'941	17'912	17'884	17'853	17'823	17'792
7. Other (domestic inorganic fertilisers)		2'788	2'796	2'788	2'592	2'440	2'440	2'352	2'036	2'044	2'140

Activity Data		2000-2009									
		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		t N/yr									
1. Inorganic N fertilisers	Urea	7'978	8'169	8'385	7'066	8'232	6'910	6'254	8'680	6'905	5'551
	Other mineral fertilisers	42'902	46'647	45'087	44'006	43'224	43'394	43'090	43'064	41'863	40'433
2. Organic N fertilisers	a. Animal manure	90'885	89'568	87'107	86'105	85'333	86'284	86'510	86'650	88'539	87'679
	b. Sewage sludge	3'356	2'934	2'513	2'091	1'670	1'248	1'054	859	573	286
	c. Other organic fertilisers	1'569	1'785	1'949	2'116	2'332	2'525	2'789	3'096	3'503	3'888
3. Urine and dung deposited by grazing animals		21'913	23'689	25'276	24'967	24'522	24'552	24'626	24'594	24'802	24'372
4. Crop residues	Arable crops	11'907	10'415	11'479	9'771	11'846	11'546	10'567	11'544	11'523	11'907
	Residues PR&P	21'900	21'931	21'922	22'020	22'029	21'845	21'849	21'928	21'904	21'916
5. Min./imm. associated with loss/gain of SOM		747	747	747	747	747	756	712	625	502	727
6. Cultivation of organic soils (ha)		17'761	17'731	17'700	17'669	17'638	17'608	17'585	17'562	17'545	17'518
7. Other (domestic inorganic fertilisers)		2'120	2'284	2'228	2'128	2'144	2'096	2'056	2'156	2'032	1'916

Activity Data		2010-2015					
		2010	2011	2012	2013	2014	2015
		t N/yr					
1. Inorganic N fertilisers	Urea	7'424	6'788	5'589	6'015	8'245	7'232
	Other mineral fertilisers	45'856	40'156	39'723	37'857	41'260	36'703
2. Organic N fertilisers	a. Animal manure	87'648	87'458	87'538	86'805	87'491	87'335
	b. Sewage sludge	0	0	0	0	0	0
	c. Other organic fertilisers	4'326	4'369	4'472	4'588	4'701	4'811
3. Urine and dung deposited by grazing animals		23'979	23'648	23'528	23'286	23'255	23'062
4. Crop residues	Arable crops	10'473	12'209	11'272	10'113	12'388	10'479
	Residues PR&P	21'918	21'964	21'919	21'786	21'709	21'604
5. Min./imm. associated with loss/gain of SOM		760	760	760	757	747	748
6. Cultivation of organic soils (ha)		17'492	17'466	17'440	17'416	17'387	17'363
7. Other (domestic inorganic fertilisers)		2'220	1'956	1'888	1'828	2'063	1'831




Figure 5-7 Relative development of the most important activity data for 3Da Direct N₂O emissions from managed soils 1990–2015.

Figure 5-7 depicts the development of the most important activity data for 3Da Direct N₂O emissions from managed soils. The use of inorganic N-fertiliser declined mainly during the 1990s due to the agricultural policy reforms and the introduction of the “Proof of Ecological Performance (PEP)” that requires a balanced fertiliser management. Simultaneously, nitrogen input from animal manure declined due to declining livestock populations (mainly cattle). Urine and dung deposited by grazing animals increased substantially due to the shift to more animal-friendly livestock husbandry in the course of the agricultural policy reforms during the 1990s and the early 21st century (see also chp. 5.3.2.2.4). N inputs from crop residues remained more or less constant during the inventory time period due to more or less stable crop production.

5.5.2.3 Indirect N₂O emissions from atmospheric deposition of N volatilised from managed soils (3Db1)

N₂O emissions from atmospheric deposition of N volatilised from managed soil were estimated based on equations 11.9 and 11.11 of the 2006 IPCC Guidelines. However, the method was adapted to the far more detailed approach of Switzerland:

$$N_2O_{(ATD)} - N = \left[\begin{array}{l} \sum_i (F_{CN_i} * Frac_{GASF_i}) \\ + \sum_T (F_{AM_T} * Frac_{GASM_T}) \\ + \sum_T (F_{PRP_T} * Frac_{GASP_T}) \\ + NH3_{AS} + NH3_{AA} \end{array} \right] + \left[\begin{array}{l} (F_{CN} + F_{AM}) * Frac_{NOXA} \\ + F_{PRP} * Frac_{NOXP} \end{array} \right] * EF_4$$

$N_2O_{(ATD)}-N$ = annual amount of N₂O–N produced from atmospheric deposition of N volatilised from managed soils (kg N₂O–N/year)

F_{CN_i} = annual amount of commercial fertiliser N of type i applied to soils (kg N/year)

F_{CGASFI} = fraction of commercial fertiliser N of type i that volatilises as NH_3 (kg N/kg N)

F_{AMT} = annual amount of managed animal manure N of livestock category T applied to soils (kg N/year)

F_{CGASMT} = fraction of applied animal manure N of livestock category T that volatilises as NH_3 (kg N/kg N)

F_{PRPT} = annual amount of urine and dung N deposited on pasture, range and paddock by grazing animals of livestock category T (kg N/year)

F_{CGASPT} = fraction of urine and dung N deposited on pasture, range and paddock by grazing animals of livestock category T that volatilises as NH_3 (kg N/kg of N)

$\text{NH}_{3\text{AS}}$ = ammonia volatilised from the vegetation cover on agricultural soils (kg N/ha)

$\text{NH}_{3\text{AA}}$ = ammonia volatilised from the vegetation cover from the alpine area (kg N/ha)

F_{CN} = total amount of commercial fertiliser N applied to soils (kg N/year)

F_{AM} = total amount of managed animal manure N applied to soils (kg N/year)

F_{CNOXA} = fraction of applied N (commercial fertilisers and animal manure) that volatilises as NO_x (kg N/kg N)

F_{PRP} = total amount of urine and dung N deposited on pasture, range and paddock by grazing animals (kg N/year)

F_{CNOXP} = fraction of urine and dung N deposited on pasture, range and paddock that volatilises as NO_x (kg N/kg of N)

EF_4 = emission factor for N_2O emissions from atmospheric deposition of N on soils and water surfaces (kg $\text{N}_2\text{O-N}$ / kg N volatilised).

5.5.2.3.1 Emission factor

The emission factor for indirect N_2O emissions from atmospheric deposition of N volatilised from managed soils is the same as used for the assessment of indirect N_2O emissions after volatilisation of NH_3 and NO_x from manure management systems. The emission factor was reassessed by a literature review by Bühlmann et al. 2015 and Bühlmann 2014. Due to slightly changing land use, the resulting emission factor shows some small variations around a mean value of 2.55%. For further information see chp. 5.3.2.4.

5.5.2.3.2 Activity data

The estimation of volatilisation of ammonia and NO_x was harmonized with the Swiss ammonia model AGRAMMON using the same emission factors and basic parameters (Table 5-23). Losses of commercial fertiliser nitrogen, animal manure N applied to soils, urine and dung N deposited on pasture, range and paddock by grazing animals as well as ammonia losses from agricultural soils and alpine areas due to processes in the vegetation cover were considered. For the calculation of NH_3 emissions, changes of agricultural structures (changes

to more animal friendly housing systems) and techniques (manure management, measures to reduce NH₃ emissions) are considered and explain temporal dynamics.

Ammonia volatilisation from **commercial fertiliser N** was estimated separately for synthetic fertilisers (urea and other synthetic fertilisers), sewage sludge, and other organic fertilisers (compost, liquid and solid digestates from biogas plants). Ammonia volatilisation of nitrogen in synthetic fertilisers is 15% for urea and 2% for other synthetic fertilisers. These estimates are based on a literature review by van der Weerden and Jarvis (1997) who examined ammonia emission factors for ammonium nitrate and urea for grassland and cropland soils. The emission factors for all other synthetic fertilisers (as straight and compound fertilisers) were assumed to be similar to that for ammonium nitrate. Ammonia emission factors for sewage sludge range from 20% to 26% depending on the composition of the sludge (Kupper et al. 2013). Other organic fertilisers include compost as well as liquid and solid digestates. Ammonia emission factors are 3.4% for compost, 21% - 30% for liquid digestate and 4.0% for solid digestate. The ammonia loss rate for liquid digestates decreased from 2007 until 2010 due to the increasing use of trailing hoses during field application.

Total $Frac_{GASF}$ as reported in CRF Table3.D declined considerably from 6.6% in 1990 to 5.4% in 2015 due to a change in the shares of the different commercial fertilisers: the use of urea and sewage sludge (which both have high NH₃ emission factors) has declined since 1990.

Different ammonia loss factors were used for **animal manure N applied to soils** from different livestock categories according to the detailed approach of the AGRAMMON model (Kupper et al. 2013). Overall weighted $Frac_{GASMT}$ for animal manure applied to soils slightly declined from 27% in the early 1990s to 22% in 2015.

Ammonia volatilisation from **urine and dung N deposited on pasture, range and paddock by grazing animals** was also assessed individually for each livestock category. Weighted mean loss rates ($Frac_{GASPT}$) range from 5.0% to 5.2%.

As an additional source, **volatilisation of ammonia from the vegetation cover** on agricultural soils and from alpine areas was accounted for (Kupper et al. 2013), assuming that 2.0 kg NH₃-N/ha and 0.5 kg NH₃-N/ha are emitted from agricultural land and the alpine area, respectively (Schjoerring and Mattsson 2001).

NO_x emissions were estimated separately for applied fertiliser N (commercial fertilisers, animal manure) and for urine and dung N deposited on pasture, range and paddock by grazing animals. NO_x emission factors for applied fertilisers and for urine and dung N deposited on pasture, range and paddock are 0.55% each, based on Stehfest and Bouwman (2006).

Nitrogen pools and flows for calculating 3Db Indirect N₂O emissions from managed soils are displayed in Table 5-24. Additional information is given in Annex 3 A3.3.

Table 5-23 Overview of NH₃ and NO_x emission factors used for the assessment of 3Db Indirect N₂O emissions from atmospheric deposition (1990–2015). Complete time series on a livestock sub-category level are provided in Annex 3 A3.3.

Emission factors volatilisation		1990-2011									
		1990	1995	2000	2005	2006	2007	2008	2009	2010	2011
		%									
NH ₃ from commercial fertiliser N (Frac _{GASFI})		6.10	5.83	5.35	4.41	4.30	4.76	4.48	4.21	4.26	4.40
	Urea	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00
	Other Mineral Fertilisers	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00
	Recycling Fertilisers (weighted average)	17.58	19.74	20.29	14.14	13.91	13.86	13.35	12.31	11.00	11.10
	Sewage Sludge	20.00	23.94	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07
	Compost	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43
	Digestate Liquid	30.00	30.00	30.00	30.00	30.00	30.00	27.00	24.00	21.00	21.00
	Digestate Solid	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00
NH ₃ from application of animal manure N (Frac _{GASMT})		26.94	27.07	25.29	25.06	25.31	25.53	25.01	24.40	23.80	23.51
	Mature Dairy Cattle	29.41	29.53	28.05	27.76	27.89	28.02	27.39	26.76	26.13	25.76
	Other Mature Cattle	27.35	27.05	25.61	26.71	27.25	27.76	27.26	26.74	26.21	26.00
	Growing Cattle (weighted average)	27.67	27.81	26.07	26.45	26.88	27.30	26.75	26.27	25.76	25.48
	Sheep (weighted average)	8.81	9.36	9.34	10.48	10.89	11.32	11.19	11.07	10.96	10.85
	Swine (weighted average)	22.85	22.43	20.62	20.55	20.79	21.06	20.41	19.77	19.12	18.98
	Other Livestock (weighted average)	11.46	12.24	11.50	11.50	11.55	11.59	11.78	12.03	12.28	12.37
NH ₃ from urine and dung N deposited on PR&P (Frac _{GASPT})		4.96	5.00	5.07	5.16	5.17	5.20	5.16	5.13	5.10	5.11
	Mature Dairy Cattle	4.95	4.93	4.87	4.82	4.81	4.80	4.80	4.80	4.80	4.80
	Other Mature Cattle	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98
	Growing Cattle (weighted average)	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98
	Sheep (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
	Swine (weighted average)	NA	NA	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00
	Other Livestock (weighted average)	5.00	6.97	8.04	9.56	9.75	10.14	9.51	8.91	8.41	8.63
NH ₃ from Agricultural Soils (kg/ha/year)		2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00
NH ₃ from Alpine Area (kg/ha/year)		0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
NO _x from applied fertilisers (Frac _{NOXA})		0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55
NO _x from urine and dung N deposited on PR&P (Frac _{NOXP})		0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55

Emission factors volatilisation		2012-2015			
		2012	2013	2014	2015
		%			
NH ₃ from commercial fertiliser N (Frac _{GASFI})		4.21	4.43	4.72	4.81
	Urea	15.00	15.00	15.00	15.00
	Other Mineral Fertilisers	2.00	2.00	2.00	2.00
	Recycling Fertilisers (weighted average)	11.33	11.57	11.80	12.01
	Sewage Sludge	26.07	26.07	26.07	26.07
	Compost	3.43	3.43	3.43	3.43
	Digestate Liquid	21.00	21.00	21.00	21.00
	Digestate Solid	4.00	4.00	4.00	4.00
NH ₃ from application of animal manure N (Frac _{GASMT})		23.20	22.93	22.64	22.36
	Mature Dairy Cattle	25.39	25.02	24.65	24.28
	Other Mature Cattle	25.79	25.57	25.36	25.15
	Growing Cattle (weighted average)	25.21	24.95	24.69	24.42
	Sheep (weighted average)	10.74	10.63	10.52	10.40
	Swine (weighted average)	18.85	18.70	18.56	18.42
	Other Livestock (weighted average)	12.40	12.47	12.55	12.63
NH ₃ from urine and dung N deposited on PR&P (Frac _{GASPT})		5.12	5.14	5.15	5.17
	Mature Dairy Cattle	4.80	4.80	4.79	4.79
	Other Mature Cattle	4.98	4.98	4.98	4.98
	Growing Cattle (weighted average)	4.98	4.98	4.98	4.98
	Sheep (weighted average)	5.00	5.00	5.00	5.00
	Swine (weighted average)	14.00	14.00	14.00	14.00
	Other Livestock (weighted average)	8.70	9.05	9.39	9.84
NH ₃ from Agricultural Soils (kg/ha/year)		2.00	2.00	2.00	2.00
NH ₃ from Alpine Area (kg/ha/year)		0.50	0.50	0.50	0.50
NO _x from applied fertilisers (Frac _{NOXA})		0.55	0.55	0.55	0.55
NO _x from urine and dung N deposited on PR&P (Frac _{NOXP})		0.55	0.55	0.55	0.55

Table 5-24 Overview of N pools and flows for calculating 3Db Indirect N₂O emission from managed soils (1990–2015). Complete time series on a livestock sub-category level are provided in Annex 3 A3.3.

Nitrogen pools and flows		1990-2011									
		1990	1995	2000	2005	2006	2007	2008	2009	2010	2011
		t N/yr									
	Animals manure N applied to soils	115'195	106'426	90'885	86'284	86'510	86'650	88'539	87'679	87'648	87'458
	Commercial fertiliser	75'339	67'321	57'925	56'173	55'243	57'855	54'876	52'074	59'826	53'269
	Area of agricultural soils (ha)	1'066'981	1'080'226	1'072'492	1'065'118	1'065'199	1'060'242	1'058'100	1'055'648	1'051'748	1'051'866
	Alpine area (ha)	538'676	499'774	496'667	487'956	484'816	486'686	485'812	485'330	486'383	483'414
Deposition	Sum volatilised N (NH ₃ and NO _x)	39'829	36'888	30'520	28'658	28'833	29'447	29'162	28'091	27'927	27'367
	NH ₃ emissions from commercial fertilisers	4'595	3'922	3'096	2'480	2'375	2'755	2'458	2'194	2'551	2'345
	NH ₃ emissions from applied animal manure	31'034	28'805	22'981	21'619	21'896	22'119	22'140	21'390	20'862	20'563
	NH ₃ emissions from pasture, range and paddock	674	717	1'110	1'267	1'274	1'279	1'280	1'251	1'224	1'209
	NH ₃ emissions from agricultural soils	2'403	2'410	2'393	2'374	2'373	2'364	2'359	2'354	2'347	2'345
	NO _x emissions from commercial fertilisers	414	370	319	309	304	318	302	286	329	293
	NO _x emissions from applied animal manure	634	585	500	475	476	477	487	482	482	481
	NO _x emissions from PR&P	75	79	121	135	135	135	136	134	132	130
Leaching and run-off	Sum leaching and run-off	49'000	45'614	40'409	37'742	37'064	37'374	36'808	35'809	36'500	35'555
	Leaching and run-off from commercial fertilisers	15'526	13'874	11'403	10'539	10'263	10'641	9'992	9'386	10'672	9'503
	Leaching and run-off from applied animal manure	23'740	21'933	17'891	16'189	16'072	15'938	16'122	15'803	15'636	15'602
	Leaching and run-off from pasture, range and paddock	2'798	2'954	4'314	4'606	4'575	4'524	4'516	4'393	4'278	4'219
	Leaching and run-off from crop residues	6'808	6'718	6'655	6'265	6'022	6'157	6'087	6'096	5'778	6'096
	Leaching and run-off from mineralisation of SOM	128	136	147	142	132	115	91	131	136	136

Nitrogen pools and flows		2012-2015			
		2012	2013	2014	2015
		t N/yr			
	Animals manure N applied to soils	87'538	86'805	87'491	87'335
	Commercial fertiliser	51'672	50'288	56'268	50'577
	Area of agricultural soils (ha)	1'051'063	1'049'923	1'051'183	1'049'478
	Alpine area (ha)	481'379	479'745	475'773	474'821
Deposition	Sum volatilised N (NH ₃ and NO _x)	26'929	26'549	26'924	26'375
	NH ₃ emissions from commercial fertilisers	2'177	2'227	2'658	2'433
	NH ₃ emissions from applied animal manure	20'308	19'905	19'810	19'528
	NH ₃ emissions from pasture, range and paddock	1'205	1'196	1'197	1'192
	NH ₃ emissions from agricultural soils	2'343	2'340	2'340	2'336
	NO _x emissions from commercial fertilisers	284	277	309	278
	NO _x emissions from applied animal manure	481	477	481	480
	NO _x emissions from PR&P	129	128	128	127
Leaching and run-off	Sum leaching and run-off	35'088	34'436	36'010	34'573
	Leaching and run-off from commercial fertilisers	9'218	8'971	10'038	9'023
	Leaching and run-off from applied animal manure	15'616	15'485	15'608	15'580
	Leaching and run-off from pasture, range and paddock	4'197	4'154	4'149	4'114
	Leaching and run-off from crop residues	5'921	5'691	6'083	5'723
	Leaching and run-off from mineralisation of SOM	136	135	133	133

Figure 5-8 depicts the development of the most important activity data for 3Db Indirect N₂O emissions from managed soils. Ammonia emissions from application of commercial fertilisers declined mainly due to reduced fertiliser use and due to the decreasing share of fertilisers with high ammonia emission rates (i.e. urea and sewage sludge). Ammonia emissions from applied animal manure declined mainly due to declining livestock populations and hence due to the reductions of available manure N. The fraction of applied animal manure N that volatilises as NH₃ (Frac_{GASMT}) declined slightly and also contributed to the decreasing trend.




Figure 5-8 Relative development of the most important activity data for 3Db Indirect N₂O emissions from managed soils 1990–2015.

5.5.2.4 Indirect N₂O emissions from leaching and run-off from managed soils (3Db2)

N₂O emissions from leaching and run-off from managed soils are estimated based on equation 11.10 of the 2006 IPCC Guidelines:

$$N_2O_{(L)-N} = (F_{CN} + F_{AM} + F_{PRP} + F_{CR} + F_{SOM}) \cdot \text{Frac}_{LEACH-(H)} \cdot EF_5$$

$N_2O_{(L)-N}$ = annual amount of N₂O–N produced from leaching and run-off of N additions to managed soils (kg N₂O–N/year)

F_{CN} = annual amount of commercial fertiliser N applied to soils (kg N/year)

F_{AM} = annual amount of managed animal manure N applied to soils (kg N/year)

F_{PRP} = annual amount of urine and dung N deposited by grazing animals (kg N/year)

F_{CR} = annual amount of N in crop residues, including N-fixing crops, returned to soils (kg N/year)

F_{SOM} = annual amount of N in mineral soils that is mineralised, in association with loss of soil C from soil organic matter as a result of changes of land use or management (kg N/year)

$\text{Frac}_{LEACH-(H)}$ = fraction of all N added to/mineralised in managed soils that is lost through leaching and runoff (kg N/kg of N additions)

EF_5 = emission factor for N₂O emissions from N leaching and run-off (kg N₂O–N/kg N leached and run-off)

5.5.2.4.1 Emission factor

The emission factor for indirect N₂O emissions from leaching and run-off from managed soils is 0.0075 kg N₂O–N/kg N according to the 2006 IPCC Guidelines (IPCC 2006).

5.5.2.4.2 Activity data

For the calculation of N₂O emissions from leaching and run-off from managed soils, N-leaching from commercial fertilisers (including synthetic fertilisers, sewage sludge, compost, and liquid and solid digestates from biogas plants) (F_{CN}), managed animal manure N applied to soils (F_{AM}), urine and dung N deposited by grazing animals (F_{PRP}), N in crop residues returned to soils (F_{CR}) and N mineralised in mineral soils (F_{SOM}) were accounted for. The method for the assessment of the respective amounts of nitrogen is described in chp. 5.5.2.2 and numbers are contained in Table 5-22.

Fra_{CLEACH} was estimated for the years 1990 and 2010 by dividing the available amount of nitrogen by the amount of nitrogen that is lost due to leaching and run-off in Switzerland according to model estimates of Prasuhn 2016. The respective loss rates are 20.6% for 1990 and 17.8% for 2010. Spiess and Prasuhn (2006), confirm that the loss rates were somewhat higher in the early 1990s and then declined due to the agricultural policy reforms. Accordingly, the reduction in the nitrate loss rate was implemented between 1995 and 2010 with constant loss rates after 2010. The same loss rates were applied to all nitrogen pools independent of their origin and composition. The resulting amount of nitrogen that is lost through leaching and run-off is given in Table 5-24.

Figure 5-8 depicts the development of the most important activity data for 3Bb Indirect N₂O emissions from managed soils. Both leaching and run-off from commercial fertiliser and animal manure N declined during the inventory time period due to the reduced nitrogen inputs and the decreasing nitrate loss rates (Fra_{CLEACH}).

5.5.2.5 NMVOC emissions

Estimation of NMVOC emissions of meadows and arable land is based on Spirig and Neftel (2002). VOC flows are estimated in Warneke et al. (2002) (for meadows) and König et al. (1995) (for arable land). Emissions were measured in a field trial in Austria (Karl et al. 2001).

5.5.3 Uncertainties and time-series consistency

For the uncertainty analysis the input data from ART (2008a) were used and were updated with current activity and emission data as well as with new default uncertainties of the 2006 IPCC Guidelines. The arithmetic mean of the lower and upper bound uncertainty is used for activity data and for emission factors, resulting in combined Approach 1 uncertainties as shown in Table 5-25. For 3Da (Direct N₂O emissions – Fertilisers) the sub-positions 3Da 1, 2, 4, 5 and 7 were combined according to Approach 1 error propagation.

For the Approach 2 analysis, asymmetric probability distributions as well as possible correlations of input data were considered (see Table A – 7).

For further results also consult chp. 1.6.1.

Table 5-25 Uncertainties for 3D Agricultural soils. (AD: Activity data; EF: Emission factor; CO: Combined).

Uncertainty 3D		Approach 1			Approach 2		
		AD	EF	CO	low	high	mean
		%			%		
Direct soil emissions	Fertilisers	14.8	135.0	135.8	-64	87	76
	Organic soils	29.4	137.5	140.6	-68	98	83
	Urine and dung deposited on PR&P	67.8	132.5	148.8	-68	113	91
Indirect soil emissions	Atmospheric deposition	39.1	240.0	243.2	-76	121	99
	Leaching and run-off	22.3	163.3	164.8	-77	102	90

The time series 1990–2015 are all considered consistent, although the following issues should be considered:

- Input data from the AGRAMMON-model is available for the years 1990 and 1995 (expert judgement and literature) as well as for 2002, 2007, 2010 and 2015 (extensive surveys on approximately 3000 farms). Values in-between the assessment years were interpolated linearly.
- Estimated amounts of sewage sludge and compost are available for the years 1990, 1995, 2000, 2005, 2007 and 2010. Years in-between were interpolated linearly. Beyond 2010, the 2010 value was used and will be updated as further survey results become available.
- The emission factor for indirect N₂O emissions following volatilisation of NH₃ and NO_x from applied fertilisers and urine and dung excreted on PR&P varies according to varying land use as described in chp. 5.3.2.4.

For more details on time-series consistency see also chp. 5.2.3 and 5.3.3.

5.5.4 Category-specific QA/QC and verification

General QA/QC measures are described in NIR chp. 1.2.3.

All further category-specific QA/QC activities are described in a separate document (Agroscope 2016b). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed.

The Swiss ammonium emission model AGRAMMON is documented in Kupper et al. (2013) and Agrammon (2010). Generally the Reporting of N₂O emissions in the Swiss national GHG inventory is consistent with the reporting of other nitrogen compounds (NH₃, NO_x) under the CLRTAP.

All relevant parameters needed for the calculation of direct and indirect nitrogen inputs to agricultural soils (e.g. F_{CN}, MS-distribution, Frac_{GAS}F, N_{ex}, Frac_{GAS}MT, F_{ON}, F_{CR}, Frac_{LEACH}) were checked for consistency and confidence and were compared (where possible) to IPCC default values, values of other countries as well as values in the literature. As one of the most important parameters, nitrogen excretion was analysed in more detail as described in chp. 5.3.4.2.

For quality of livestock population data consult chp. 5.2.4.

The estimate for the area of cultivated histosols in the agricultural sector is consistent with the estimates reported under cropland and grassland in the LULUCF sector. A literature study conducted by Leifeld et al. (2003) estimates $17'000 \pm 5'000$ ha which is close to the numbers reported in the LULUCF sector (17'700 ha on average).

The country-specific value of $Frac_{LEACH}$ is based on a very detailed model for the assessment of leaching and run-off in Switzerland (Hürdler et al. 2015, Prasuhn 2016) that takes into account regional parameters such as topography, different crop species as well as fertiliser application levels.

N_2O emission factors were compared to values in the literature to ensure plausibility. Implied emission factors are similar to measured values from the literature representative for Swiss conditions (Agroscope 2016b).

5.5.5 Category-specific recalculations

General information on recalculations is provided in chp. 10.

- The AD for direct N_2O emissions from N in crop residues returned to soils was revised for the year 2014 due to data updates by the Swiss Farmers Union. Overall emissions increased by 0.02 kt CO_2 equivalents.
- The alpine area used to estimate NH_3 volatilisation from the vegetation cover and N_2O emissions from crop residues on pasture and meadows was slightly revised with negligible effects on overall emissions.
- All estimates based on AGRAMMON data were recalculated for the years 2011–2014 due to a new interpolation after new survey results for 2015 became available. The impact on overall emissions in kt CO_2 equivalents is: 2011: 0.4; 2012: 0.9; 2013: 1.4; 2014: 1.9.
- N-input from co-substrates in agricultural biogas plants for 2014 was revised due to an error correction in the background file of the waste sector. The impact on overall emissions is negligible.
- AD for N_2O emissions from the cultivation of organic soils was revised due to new AREA-projections in the LULUCF sector for all inventory years. Overall emissions decreased by 0.03 kt CO_2 equivalent on average (1990: no change; 2014: -0.08 kt CO_2 equivalents).
- The AD for direct N_2O emissions from N in mineral soils that is mineralized/immobilized in association with loss of soil C was revised due to new projections in the LULUCF sector for all inventory years. Overall emissions increased by 0.13 kt CO_2 equivalents on average (1990: +0.00 kt CO_2 equivalents; 2014: +0.20 kt CO_2 equivalents).
- $Frac_{LEACH}$ was recalculated due to new model estimates of NO_3 losses on agricultural soils from Prasuhn (2016). Overall emissions decreased by 18.67 kt CO_2 equivalents on average (1990–2014).

5.5.6 Category-specific planned improvements

New country-specific values for N_{ex} for most livestock categories will become available in the next years due to a revision of the “Principles of Fertilization in Arable and Forage Crop Production (Flisch et al. 2009)”.

A release of a new version of the AGRAMMON-Model is planned for 2017. If possible the respective AGRAMMON projections will be included during the next GHG inventory submission.

NMVOC emissions will be revised in submission 2018 and separately reported for 3B Manure management and 3D Agricultural soils.

5.6 Source category 3E – Prescribed burning of savannahs

Burning of savannahs does not occur (NO) in Switzerland.

5.7 Source category 3F – Field burning of agricultural residues

Field burning of agricultural residues does not occur (NO) in Switzerland.

Emissions from open burning of branches in agriculture and forestry were reported here in the past. However, the respective emissions were moved to the LULUCF and the waste sector based on recommendations from the UNFCCC expert review teams (e.g. UNFCCC 2017; W12 and W13). Respective information can be found under source category 4V “Biomass Burning” (see chp. 6.4.2.13) and source category 5C “Incineration and open burning of waste” (see chp. 7.4).

5.8 Source category 3G – Liming

5.8.1 Source category description

CO₂ emission from 3G Liming is not a key category.

Emissions from the application of lime (Ca(CO₃)) and dolomite (CaMg(CO₃)₂) to agricultural soils are reported.

The emissions due to liming of agricultural soils range from 22.2 to 32.9 kt CO₂ per year.

5.8.2 Methodological issues

A simple Tier 1 approach was adopted using estimated amounts of lime and dolomite applied and IPCC default emission factors.

5.8.2.1 Emission factor

The availability of a country-specific emission factors for agricultural lime and dolomite application was investigated, but no domestic measurement data could be found. Consequently, the IPCC default carbon conversion factors for carbonate containing lime (0.12 t C per t Ca(CO₃)) and for dolomite (0.13 t C per t CaMg(CO₃)₂) were used (IPCC 2006).

5.8.2.2 Activity data

The total annual amount of lime and dolomite applied to agricultural soils is between 51'300 Mg (1990) and 74'050 Mg (2008–2015). It was estimated by Agroscope in 2009 for the period 1990–2008. For 2009–2015 the same value as for 2008 was used: An inquiry in 2013 including the most important production and trading companies of lime products suggests that the consumption of limestone remained constant in this period (Agroscope 2014a). The split of lime into calcium carbonate and dolomite is based on the following assumptions and data:

- $\text{Ca}(\text{CO}_3)$ contained in mixed compound fertilizers as reported by Agricura (2015),
- All material originating from nuclear power plants and from the sugar beet industry is $\text{Ca}(\text{CO}_3)$,
- The remaining lime not covered under the points above was divided fifty fifty into $\text{Ca}(\text{CO}_3)$ and $\text{CaMg}(\text{CO}_3)_2$.

5.8.3 Uncertainties and time-series consistency

The amount of total lime applied in agriculture is mainly based on expert judgement; the resulting number is uncertain. A relative uncertainty of $\pm 40\%$ was used as an approximation (Agroscope 2014a). For the emission factor of lime a lower uncertainty of $\pm 5\%$ was chosen, because it is a simple chemical process. The combined Approach 1 uncertainty is thus $\pm 40.3\%$. Approach 2 uncertainties do not differ significantly from Approach 1 uncertainties.

For further results also consult chp. 1.6.1.

Consistency: Time series for 3G Liming are all considered consistent.

5.8.4 Category-specific QA/QC and verification

General QA/QC measures are described in NIR chp. 1.2.3.

No further category-specific quality assurance activities were conducted.

5.8.5 Category-specific recalculations

General information on recalculations is provided in chp. 10.

All lime applied in agriculture was previously reported as $\text{Ca}(\text{CO}_3)$. During the 2017 submission the total amount of lime was split into calcium carbonate and dolomite. The new specific EF for dolomite (i.e. 0.13) is slightly higher than the old EF that was used previously for both limestone and dolomite (i.e. 0.12). Accordingly, overall emissions increased by 0.15 kt CO_2 equivalents on average (1990: +0.12 kt CO_2 equivalents; 2014: +0.23 kt CO_2 equivalents).

5.8.6 Category-specific planned improvements

No category-specific improvements are planned.

5.9 Source category 3H – Urea application

5.9.1 Source category description

CO₂ emission from 3H Urea application is not a key category.

Adding urea to soils during fertilisation leads to a loss of CO₂ that was fixed during the industrial production process of the fertilizer. Emissions in Switzerland range from 8.7 to 26.7 kt CO₂ per year with a general decreasing trend from 1990 to 2014.

5.9.2 Methodological issues

A simple Tier 1 approach was adopted using estimated amounts of urea applied and IPCC default emission factors.

5.9.2.1 Emission factor

No country-specific emission factors are available. Consequently, the IPCC default emission factor of 0.20 t of C per t of urea was applied.

5.9.2.2 Activity data

The amount of urea applied to agricultural soils was obtained from Agricura (2015). Fertiliser statistics are based on sales statistics by the compulsory storekeepers of fertilisers (Pflichtlagerhalter) and small importers. Agricura conducts plausibility checks with import-data received by the Directorate General of Customs (Oberzolldirektion).

5.9.3 Uncertainties and time-series consistency

An uncertainty of $\pm 5\%$ for the activity data was estimated according to ART (2008a). An uncertainty of $\pm 5\%$ was assumed for the emission factor since it is a simple chemical process. The combined Approach 1 uncertainty is hence $\pm 7.1\%$. Approach 2 uncertainties do not differ significantly from Approach 1 uncertainties.

For further results also consult chp. 1.6.1.

Consistency: Time series for 3H Urea application are all considered consistent.

5.9.4 Category-specific QA/QC and verification

General QA/QC measures are described in NIR chp. 1.2.3.

No further category-specific quality assurance activities were conducted.

5.9.5 Category-specific recalculations

General information on recalculations is provided in chp. 10.

No category-specific recalculations were carried out.

5.9.6 Category-specific planned improvements

No category-specific improvements are planned.

6 LULUCF

6.1 Overview of LULUCF

6.1.1 Methodology

Chapter 6 presents estimates of greenhouse gas emissions by sources and removals by sinks from land use, land-use change and forestry (LULUCF). The sector LULUCF includes emissions and removals from the carbon pool in Harvested wood products (HWP). Data acquisition and calculations are based on the Guidelines for National Greenhouse Gas Inventories (IPCC 2006), Volume 4 "Agriculture, Forestry and Other Land Use" (AFOLU). In many subcategories country-specific emission factors were used.

The land areas in the period 1990–2015 are represented by geographically explicit land use data with a resolution of one hectare (following approach 3 for representing land areas; IPCC 2006). Direct and repeated assessment of land use with full spatial coverage also enables to calculate spatially explicit land-use change matrices. In 2004, the Swiss Land Use Statistics AREA was launched. Simultaneously, aerial photos from two earlier Swiss Land Use Statistics (1979/85 and 1992/97) were re-evaluated, applying the same approach. The AREA surveys 1, 2 and 3 was completed in 2013 and the interpretation of the entire Swiss territory is available for three time slices. In this submission, results of the ongoing AREA4 survey for the western part of Switzerland were included.

The six main land-use categories required by IPCC (2006) are: A. Forest land, B. Cropland, C. Grassland, D. Wetlands, E. Settlements and F. Other land. These categories were divided in 18 sub-divisions of land use. A further spatial stratification reflects "elevation" (3 zones), "geomorphologic and climatic conditions" (adopting the five production regions of the National Forest Inventory; NFI) and "soil type" (mineral, organic).

Country-specific emission factors and carbon stocks for Forest Land were derived from three completed National Forest Inventories (NFI1, NFI2, NFI3, finalized in 1985, 1995, and 2006, respectively) and the first five annual tranches of the continuous NFI4 (2011–2015). The inventories comprised ca. 3'400 (5-years interval from NFI4), 6'500 (NFI2 and NFI3) and 11'000 (NFI1) terrestrial sampling plots (see Table 6-12), where biomass stock, growth, cut and mortality were measured.

For the remaining land-use categories, carbon stocks and GHG emissions and removals were derived from particular research activities, domestic surveys and measurements in the fields of agriculture (Cropland, Grassland) and nature conservation (Wetlands). Partially, also IPCC default values and expert estimates were used.

6.1.2 Emissions and removals

Table 6-1 and Figure 6-1 summarize the CO₂ emissions and removals as a result of carbon losses and gains for the years 1990–2015. The total net emissions and removals of CO₂ from 1990 to 2015 varied between -5'082 kt (1996) and 4'925 kt (2000).

Table 6-1 and Figure 6-1 show a breakdown of Switzerland's CO₂ balance in the LULUCF sector. Five components were differentiated:

- Gains in carbon stock of living biomass on all land uses and due to land-use changes; this component represents the largest sink of carbon.
- Losses in carbon stock of living biomass on all land uses and due to land-use changes; this component represents the largest source of carbon. The highest losses were observed in the year 2000 after a heavy storm with windfall in December 1999.
- Net carbon stock changes in dead organic matter (DOM; consisting of dead wood and litter) on Forest land remaining forest land as well as on Forest land converted to non-forest land: this component represents a sink of carbon in most years.
- Net carbon stock changes (1) in soils due to the use of soils (especially of organic soils) and due to land-use changes, and (2) by wildfires. Taken together, both components persistently represent a source of carbon in the period under investigation.
- Net carbon stock changes in Harvested wood products (HWP). In the period under investigation this component persistently represents a sink of carbon, i.e. the carbon stock stored in wood products was increasing.

The largest part of gains and losses in carbon stocks of biomass occurred in forests, where growth of biomass exceeded cut and mortality, except for the years 2000 and 2006 (see also chp. 2.3.3). Overall, the LULUCF sector was a sink of on average $-1'745 \text{ kt CO}_2 \text{ yr}^{-1}$ between 1990 and 2015 (see Table 6-1 and Figure 6-3).

Table 6-1 CO₂ emissions and removals in the LULUCF sector, 1990–2015. See main text for the respective components. Positive values refer to emissions; negative values refer to removals. In this data set, emissions of CH₄ and N₂O are not included; GHG (i.e. CO₂, CH₄, N₂O) emissions and removals in the LULUCF sector (in CO₂ eq) are shown in Figure 6-3.

LULUCF	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	kt CO ₂ yr ⁻¹									
Gains of living biomass	-12'696	-12'712	-12'857	-12'815	-12'756	-13'001	-12'954	-12'942	-12'634	-12'645
Losses of living biomass	12'479	9'232	8'878	8'677	9'656	9'252	9'083	9'871	11'020	10'926
Net change in dead organic matter	292	-412	175	-73	536	-232	-1'584	-766	-816	-508
Net change in organic and mineral soils and wildfires	761	761	760	757	764	765	768	763	777	775
LULUCF (excluding HWP)	836	-3131	-3044	-3453	-1800	-3216	-4686	-3074	-1653	-1453
Net change in Harvested Wood Products (HWP)	-1'231	-942	-774	-631	-477	-561	-396	-314	-424	-479
Total LULUCF	-395	-4072	-3818	-4084	-2277	-3777	-5082	-3388	-2076	-1932

LULUCF	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	kt CO ₂ yr ⁻¹									
Gains of living biomass	-13'058	-12'661	-12'715	-12'677	-13'371	-12'791	-13'250	-13'860	-13'448	-13'473
Losses of living biomass	18'337	12'362	10'085	11'296	11'072	11'404	14'000	13'548	12'552	11'899
Net change in dead organic matter	-293	-517	-63	-933	-589	-958	-383	-22	-450	-597
Net change in organic and mineral soils and wildfires	773	771	769	767	765	783	806	800	760	748
LULUCF (excluding HWP)	5760	-44	-1923	-1547	-2123	-1562	1174	465	-586	-1423
Net change in Harvested Wood Products (HWP)	-835	-577	-443	-441	-637	-763	-626	-731	-528	-370
Total LULUCF	4925	-621	-2366	-1988	-2760	-2325	548	-265	-1114	-1793

LULUCF	2010	2011	2012	2013	2014	2015	Mean
	kt CO ₂ yr ⁻¹						
Gains of living biomass	-13'401	-14'027	-13'415	-13'422	-14'143	-13'442	-13'122
Losses of living biomass	12'824	12'459	11'805	12'276	12'173	12'038	11'508
Net change in dead organic matter	-1'109	125	-325	-625	389	-263	-385
Net change in organic and mineral soils and wildfires	757	757	757	770	763	756	768
LULUCF (excluding HWP)	-929	-686	-1179	-1001	-817	-911	-1'231
Net change in Harvested Wood Products (HWP)	-411	-258	-168	-161	-106	-70	-513
Total LULUCF	-1340	-944	-1347	-1162	-923	-981	-1'745




Figure 6-1 CO₂ emissions and removals in the LULUCF sector (in kt CO₂), 1990–2015, broken down for: (i) CO₂ removals due to the gain (growth) of living biomass, (ii) CO₂ emissions due to the loss (cut and mortality) of living biomass, (iii) net CO₂ emissions and removals due to changes in dead organic matter, (iv) net CO₂ emissions from soils and wildfires, and (v) net CO₂ removals from Harvested wood products. Positive values indicate emissions, negative values indicate removals.

The non-CO₂ emissions associated with land use, land-use change and forestry were relatively small. Between 1990 and 2015 maximum annual CH₄ emissions were 1.29 kt yr⁻¹, and maximum annual N₂O emissions were 0.30 kt yr⁻¹ (32 kt CO₂ eq and 90 kt CO₂ eq, respectively; see year 1997 in Figure 6-2). The emissions arose from drained organic soils (N₂O; CRF Table4(II)), flooded land/reservoirs (CH₄; CRF Table4(II)), nitrogen mineralization associated with loss of soil organic matter resulting from land use and land-use change (N₂O; CRF Table4(III)), nitrogen leaching and run-off on non-agricultural soils (indirect N₂O emissions; CRF Table4(IV)), wildfires on Forest land and Grassland (CH₄ and N₂O; CRF Table4(V)), and controlled burning of residues from forestry (CH₄ and N₂O; CRF Table4(V)). The calculation methods are based on default procedures of IPCC (2006, Volume 4).




Figure 6-2 N₂O and CH₄ emissions in the LULUCF sector (in kt CO₂ eq), 1990–2015.

Figure 6-3 shows the resulting net GHG emissions and removals in the LULUCF sector 1990–2015, including both CO₂ and non-CO₂ (i.e. CH₄, N₂O) fluxes. Further explanatory notes on LULUCF data can be found in chp. 2.3.3 “Emission trends in sector 4 LULUCF”.




Figure 6-3 Net GHG emissions and removals in the LULUCF sector, 1990–2015 (in kt CO₂ eq). Positive values refer to emissions, negative values refer to removals.

6.1.3 Approach for calculating carbon emissions and removals

6.1.3.1 Work steps

The selected procedure for calculating carbon emissions and removals in the LULUCF sector corresponds to a Tier 2 approach as described in IPCC 2006 (Volume 4, chp. 3). It can be summarised as follows:

- Define managed and unmanaged land: In Switzerland, all land is considered to be managed. Other land (CC61, see Table 6-2) is defined as the residual area without any relevant human activities.
- Define land-use categories and sub-divisions with respect to available land use data (see Table 6-2). Combination categories (CC) were defined on the basis of the AREA land-use and land-cover categories (Table 6-6; SFSO 2006a).
- Define criteria and collect data for the spatial stratification of the land-use categories.
- For Forest land: Measure or estimate the carbon stocks in living biomass ($stockC_l$), in dead wood ($stockC_d$), in litter ($stockC_h$), and in soil ($stockC_s$) for each spatial stratum of the combination categories (CC).
For non-Forest land: Measure or estimate the carbon stocks in living biomass ($stockC_l$), in dead organic matter ($stockC_{dom}$), and in soil ($stockC_s$) for each spatial stratum of the combination categories (CC).
- For Forest land: Measure or estimate the gain of carbon in living biomass ($gainC_l$), the loss of carbon in living biomass ($lossC_l$), the net carbon stock change in dead wood ($changeC_d$), in litter ($changeC_h$), and in soil ($changeC_s$) for each spatial stratum of the combination categories (CC).
For non-Forest land: Measure or estimate the gain of carbon in living biomass ($gainC_l$), the loss of carbon in living biomass ($lossC_l$), the net carbon stock change in dead organic matter ($changeC_{dom}$), and in soil ($changeC_s$) for each spatial stratum of the combination categories (CC).
- Calculate the land use and the land-use change matrix for each spatial stratum.
- For Forest land: Calculate the net carbon stock changes in living biomass (ΔC_l), in dead wood (ΔC_d), in litter (ΔC_h), and in soil (ΔC_s) for all cells of the land-use change matrix for each year under consideration.
For non-Forest land: Calculate the net carbon stock changes in living biomass (ΔC_l), in dead organic matter (ΔC_{dom}), and in soil (ΔC_s) for all cells of the land-use change matrix for each year under consideration.
- Finally, aggregate the results by summarising the carbon stock changes over combination categories and spatial strata according to the level of disaggregation displayed in the reporting tables.
- Calculate emissions and removals of the carbon pool in Harvested wood products (HWP).

The combination category CC11 (see Table 6-2) refers a conversion from land to forest land that corresponds to the Swiss definition for afforestation activities under Article 3, paragraph 3, of the Kyoto Protocol as defined in the Initial Report for the first commitment period (FOEN 2006h). For the reporting under the UNFCCC, afforested areas were allocated to category 4A2 (Land converted to forest land), where they were reported in an individual subdivision afforestation (no capitalisation, first letter in lowercase; see chp. 6.4.1). The same afforested areas were reported as Afforestations (with capitalisation, first letter in uppercase) under the

Kyoto-Protocol (see chp. 11.1.3). In a nutshell, the diction Afforestation was consistently used to indicate the Kyoto Protocol Article 3, paragraph 3 activity.

Table 6-2 Land-use categories used in this report (combination categories CC): 6 main land-use categories (identical to the UNFCCC land-use categories) and 18 sub-divisions. Additionally, descriptive remarks, abbreviations used in the reporting tables, and CC codes are given. For a detailed definition of the combination categories see Table 6-6 and SFSO (2006a).

CC Main category	CC Sub-division	Remarks	Terminology in CRF tables	CC code
A. Forest Land	afforestation	areas converted to forest by active measures, e.g. planting	afforestation	11
	productive forest	dense and open forest meeting the criteria of forest land	productive	12
	unproductive forest	brush forest and forest on unproductive areas meeting the criteria of forest land	unproductive	13
B. Cropland		arable and tillage land (annual crops and leys in arable rotations)		21
C. Grassland	permanent grassland	meadows, pastures (low-land and alpine)	permanent	31
	shrub vegetation	agricultural and unproductive areas predominantly covered by shrubs	woody, shrub	32
	vineyard, low-stem orchard, tree nursery	perennial agricultural plants with woody biomass (no trees)	woody, vine	33
	copse	agricultural and unproductive areas covered by perennial woody biomass including trees	woody, copse	34
	orchard	permanent grassland with fruit trees	woody, orchard	35
	stony grassland	grass, herbs and shrubs on stony surfaces	unproductive, stony	36
	unproductive grassland	unmanaged grass vegetation	unproductive	37
D. Wetlands	surface water	lakes and rivers	surface water	41
	unproductive wetland	reed, extensively managed wetland	unprod wetland	42
E. Settlements	buildings and constructions	areas without vegetation such as houses, roads, construction sites, dumps	building	51
	herbaceous biomass in settlements	areas with low vegetation, e.g. lawns	herb	52
	shrubs in settlements	areas with perennial woody biomass (no trees)	shrub	53
	trees in settlements	areas with perennial woody biomass including trees	tree	54
F. Other Land		areas without soil and vegetation: rocks, sand, scree, glaciers		61

6.1.3.2 Calculating carbon stock changes

For calculating carbon stock changes, the following input parameters (mean values per hectare) were quantified for all combination categories (CC) and spatial strata (i):

stockC _{l,i,CC}	carbon stock in living biomass (t C ha ⁻¹)
stockC _{d,i,CC}	carbon stock in dead wood (t C ha ⁻¹)
stockC _{h,i,CC}	carbon stock litter (organic soil horizons) (t C ha ⁻¹)
stockC _{s,i,CC}	carbon stock in soil (t C ha ⁻¹)
gainC _{l,i,CC}	annual gain (gross growth) of carbon in living biomass (t C ha ⁻¹ yr ⁻¹)
lossC _{l,i,CC}	annual loss (cut and mortality) of carbon in living biomass (t C ha ⁻¹ yr ⁻¹)
changeC _{d,i,CC}	annual net carbon stock change in dead wood (t C ha ⁻¹ yr ⁻¹)
changeC _{h,i,CC}	annual net carbon stock change in litter (t C ha ⁻¹ yr ⁻¹)
changeC _{s,i,CC}	annual net carbon stock change in soil (t C ha ⁻¹ yr ⁻¹)

In the reporting tables on non-forest land under the UNFCCC (Table4.B to Table4.F), the carbon stocks and carbon stock changes of litter and dead wood are merged into "dead organic matter" (DOM):

$$\text{stockC}_{\text{dom},i,\text{CC}} = \text{stockC}_{\text{d},i,\text{CC}} + \text{stockC}_{\text{h},i,\text{CC}}$$

$$\text{changeC}_{\text{dom},i,\text{CC}} = \text{changeC}_{\text{d},i,\text{CC}} + \text{changeC}_{\text{h},i,\text{CC}}$$

On this basis, the total changes in carbon stocks (t C yr⁻¹) in living biomass (deltaC_l), in dead wood (deltaC_d), in litter (deltaC_h), and in soils (deltaC_s) were calculated for all cells of the land-use change matrix for each year under consideration. Each cell is characterized by a land-use category before the conversion (b), a land-use category after the conversion (a), and the area of converted land within the spatial stratum (i). This approach includes cases without any land-use change (a = b).

Equations 6.1–6.8 show, according to the AFOLU-Guidelines (IPCC 2006, Volume 4), two approaches and their application for calculating carbon gains and losses: (1) the gain-loss approach (Equation 2.4; IPCC 2006, Volume 4) and (2) the stock-difference approach (Equation 2.5; IPCC 2006, Volume 4).

The gain-loss approach is defined as:

$$\text{deltaC}_{\text{l},i,\text{ba}} = (\text{gainC}_{\text{l},i,\text{a}} - \text{lossC}_{\text{l},i,\text{a}}) * A_{i,\text{ba}} \quad (6.1)$$

$$\text{deltaC}_{\text{d},i,\text{ba}} = \text{changeC}_{\text{d},i,\text{a}} * A_{i,\text{ba}} \quad (6.2)$$

$$\text{deltaC}_{\text{h},i,\text{ba}} = \text{changeC}_{\text{h},i,\text{a}} * A_{i,\text{ba}} \quad (6.3)$$

$$\text{deltaC}_{\text{s},i,\text{ba}} = \text{changeC}_{\text{s},i,\text{a}} * A_{i,\text{ba}} \quad (6.4)$$

The stock-difference approach is defined as:

$$\text{deltaC}_{l,i,ba} = [(\text{stockC}_{l,i,a} - \text{stockC}_{l,i,b}) / \text{CT}] * A_{i,ba} \quad (6.5)$$

$$\text{deltaC}_{d,i,ba} = [(\text{stockC}_{d,i,a} - \text{stockC}_{d,i,b}) / \text{CT}] * A_{i,ba} \quad (6.6)$$

$$\text{deltaC}_{h,i,ba} = [(\text{stockC}_{h,i,a} - \text{stockC}_{h,i,b}) / \text{CT}] * A_{i,ba} \quad (6.7)$$

$$\text{deltaC}_{s,i,ba} = [(\text{stockC}_{s,i,a} - \text{stockC}_{s,i,b}) / \text{CT}] * A_{i,ba} \quad (6.8)$$

where:

a	land-use category after conversion (CC = a)
b	land-use category before conversion (CC = b)
ba	land-use conversion from b to a
i	spatial stratum
$A_{i,ba}$	area of land (ha) converted from b to a in the spatial stratum i (area converted in the inventory year if CT=1 year, or the sum of the areas converted within the last 20 years if CT=20 years)
CT	conversion time (yr), see chp. 6.1.3.3.

Table 6-3 pinpoints which approach was used for calculating the carbon stock changes for the various types of land-use conversion and carbon pools (living biomass, dead wood/litter, mineral soil and organic soil).

The gain-loss approach was used in cases of no land-use change and generally for continuous transitions, e.g. the growth of living biomass on land converted to forest land. The stock-difference approach was used for abrupt changes following discrete events (e.g. loss of biomass by deforestation, CT = 1 year) as well as for slow processes such as the change in soil carbon content (CT = 20 years, see chp. 6.1.3.3).

For the conversions between different forest combination categories the approach was chosen in such a way that potential carbon losses cannot be underestimated: e.g. for CC12 to CC13 stock-difference is used, for CC13 to CC12 gain-loss is used, see Table 6-3.

In case of land-use changes involving "Buildings and Constructions" (CC51), 50% of the difference between soil carbon stocks before and after the conversion was reported as a source or sink, respectively; for a detailed documentation see chp. 6.8.2.2.

Table 6-3 Calculation approach (gain-loss or stock-difference) and conversion time periods (CT in years) applied for different land-use changes and carbon pools. KP = corresponding activity under the Kyoto Protocol; NF = non-forest combination category. Combination categories CC11 to CC61 were introduced in Table 6-2.

Change in main land-use category or sub-division	Living biomass	Dead wood, litter	Mineral soil	Organic soil	Remarks
no change in category KP and UNFCCC	gain-loss	gain-loss	gain-loss	gain-loss	
CC13 to CC12 UNFCCC: 4A1 KP: forest management	gain-loss	stock-diff., 20	stock-diff., 20	gain-loss	
CC12 to CC13 UNFCCC: 4A1 KP: forest management	stock-diff., 20	stock-diff., 20	stock-diff., 20	gain-loss	
CC11 to CC12 UNFCCC: 4A1 KP: afforestation >20 yr	gain-loss	gain-loss	gain-loss	gain-loss	
change to CC11 UNFCCC: 4A2 KP: afforestation ≤20 yr	gain-loss	stock-diff., 20	stock-diff., 20	gain-loss	Dead organic matter is 0 in CC11 and in NF; direct human-induced
NF to CC12/CC13 UNFCCC: 4A2 KP: forest management	gain-loss	stock-diff., 20	stock-diff., 20	gain-loss	
change to CC51 UNFCCC: 4E2 KP: deforestation	stock-diff., 1	stock-diff., 1	stock-diff., 20 (50%)	stock-diff., 20 (50%)	Buildings/constructions; soil: stock-difference reduced by 50% in changes from/to CC51
change to CC52-54 UNFCCC: 4E2	stock-diff., 1	stock-diff., 1	stock-diff., 20	gain-loss	Unsealed settlement areas
change to CC21 UNFCCC: 4B2	stock-diff., 1	stock-diff., 1	stock-diff., 20	gain-loss	Cropland
change to CC31-37 UNFCCC: 4C2	stock-diff., 1	stock-diff., 1	stock-diff., 20	gain-loss	Grassland
change to CC41 UNFCCC: 4D2	stock-diff., 1	stock-diff., 1	stock-diff., 1	gain-loss	Surface water
change to CC42 UNFCCC: 4D2	stock-diff., 1	stock-diff., 1	stock-diff., 20	gain-loss	Unproductive wetland
change to CC61 UNFCCC: 4F2	stock-diff., 1	stock-diff., 1	stock-diff., 20	stock-diff., 20	Other land

6.1.3.3 Considering the conversion time (CT)

Changes in the soil carbon stock, and this is also true for the increase of woody biomass, as a result of land-use changes are slow processes that might take decades. Therefore, IPCC (2006, Volume 4, chp. 2) suggests implementing a conversion time (CT). Following the IPCC default value (CT = 20 years), carbon emissions or removals due to a soil carbon stock difference (stockCs,i,a – stockCs,i,b) do not occur in one year but are distributed evenly over the 20 years following the land-use conversion.

A conversion time of 20 years was applied to all soil carbon stock changes (except land converted to surface water). Accordingly, the category 2 in reporting tables Table4.A to Table4.F contain the cumulative area remaining in the respective category in the reporting year.

The combination category afforestations (CC11) is a transitional category by definition in the land-use survey. Areas converted to afforestations are reported in category 2 in the CRF Table 4.A with the same conversion time as for other forest subcategories (20 years). However, after 20 years, afforestations remaining afforestations (according to the land-use survey) are reported in category 1 of CRF Table 4.A and are merged with productive forests (CC12). Note: Under the Kyoto Protocol Afforestations are processed differently (see chp. 11.2.3).

Table 6-3 shows the conversion times applied to carbon stock changes in living biomass, in dead organic matter (dead wood, litter), and in soils for all types of land-use changes.

There are no consistent data sources on land-use changes before 1990, but it is well known (ARE/FOEN 2007, FOEN 2016k) that the main trends of the Swiss land-use dynamics (e.g. increase of forests and settlements) did arise before 1972. Therefore, it was assumed that between 1971 and 1989 the annual rate of all land-use changes was the same as in 1990. Based on this assumption it was possible to produce the land-use data required for the consideration of the conversion time in that period and to consider it in the years 1990 to 2009 in accordance with the 20 years conversion period.

6.1.3.4 Displaying results in the Common Reporting Format (CRF)

In the reporting tables Table 4.A to Table 4.F, a part of the combination categories (CC) and associated spatial strata are shown at an aggregated level for optimal documentation and overview. The values of ΔC are accordingly summarised. Positive values of $\Delta C_{l,i,ba}$ were inserted in the column "Gains" and negative values in the column "Losses", respectively. The values of $\Delta C_{d,i,ba}$, $\Delta C_{h,i,ba}$ were inserted into columns "Net carbon stock change in dead wood" and "Net carbon stock change in litter" in CRF Table 4.A, and the values of $\Delta C_{dom,i,ba}$ were inserted into columns "Net carbon stock change in dead organic matter" in the reporting tables Table 4.B to Table 4.F. The values of $\Delta C_{s,i,ba}$ were inserted in column "Net carbon stock change in soils" in the reporting tables Table 4.A to Table 4.F.

The reporting tables Table 4.B to Table 4.F are subdivided in two parts: (1) X land remaining X land and (2) Land converted to X land. Changes of areas from one combination category to another within the same main land-use category are reported in part (1) of the reporting tables. For example, the area of "shrub vegetation" (CC32) converted to "permanent grassland" (CC31) would be reported in CRF Table 4.C under 4C1 in the sub-division "permanent". As CC31 and CC32 do have different carbon stocks in biomass, a carbon stock change would be calculated according to the equations presented in chp. 6.1.3.2.

The CRF Reporter generated errors or inconsistent content in several reporting tables related to the LULUCF sector (see Annex 6):

- CRF Table 4: N₂O emissions from 4(IV) (indirect emissions) are not separated in the categories 4A, 4B, 4C, 4D, 4E, and 4F but just shown in CRF Table 4(IV). But they are summarized in the total sum of N₂O emissions in CRF Table 4 (cell D7) without being displayed in this CRF Table 4. The sum is correct (considering the inclusion of the indirect emissions).
- CRF Table 4(V): Notation "IEF" for CO₂, and values for CH₄ and N₂O in line 9 and 10 are missing.

- CRF Table10s4: Similar problem as in CRF Table4 above; in sector 4 LULUCF the sum (as displayed in row 38) is not just the sum of N₂O emissions from the categories 4A+4B+4C+4D+4E+4F (rows 39 to 46), but includes indirect N₂O emissions from 4(IV). The sum is correct (considering the inclusion of the indirect emissions).

6.1.4 Carbon stocks, emission factors, and net changes at a glance

Table 6-4 lists carbon stocks, gains, losses and net changes of carbon for the pools living biomass, dead wood, litter and soil stratified by combination category (CC) and spatial strata for the year 1990. These values remain constant during the inventory period 1990–2015 with the following exceptions (highlighted cells):

- Carbon stock, gain and loss of living biomass, carbon stock and net change in dead wood, net change in litter, and net change in mineral soils of productive forest (CC12): Derivation of the data and the annual values are described in chp. 6.4.2.5, chp. 6.4.2.6 and chp. 6.4.2.7.
- Carbon stock, gain and loss of living biomass of cropland (CC21): Annual data of CC21 are listed in chp. 6.5.2.

The derivation of the individual carbon stocks and emission factors is explained in detail in chapters 6.4 to 6.9. Positive values refer to gains in carbon stock; negative values refer to losses in carbon stock.

Table 6-4 Carbon stocks and changes in living biomass, in dead wood, in litter and in soils for the combination categories (CC), stratified by elevation zone, NFI region, and soil type. The values are valid for the period 1990–2015 with the exception of the values in the highlighted cells, which change annually (numbers given here are for the year 1990); cf. main text.

Combination category (CC)	NFI region	Elevation zone z	Carbon stock in living biomass (stockC _{l,i})	Carbon stock in dead wood (stockC _{d,i})	Carbon stock in litter (stockC _{h,i})	Carbon stock in mineral soil (stockC _{s,i})	Carbon stock in organic soil (stockC _{s,i})	Gain of living biomass (gainC _{l,i})	Loss of living biomass (lossC _{l,i})	Net change in dead wood (changeC _{d,i})	Net change in litter (changeC _{h,i})	Net change in mineral soil (changeC _{s,i})	Net change in organic soil (changeC _{s,i})
			[t C ha ⁻¹]					[t C ha ⁻¹ yr ⁻¹]					
11 Afforestations	1	1	10.00	0	0	82.65	145.6	2.39	-0.21	0	0	0	-2.6
	1	2	10.00	0	0	102.03	145.6	2.39	-0.21	0	0	0	-2.6
	1	3	7.50	0	0	121.34	145.6	1.35	-0.1	0	0	0	-2.6
	2	1	10.00	0	0	55.40	145.6	2.39	-0.21	0	0	0	-2.6
	2	2	10.00	0	0	62.12	145.6	2.39	-0.21	0	0	0	-2.6
	2	3	7.50	0	0	122.00	145.6	1.35	-0.1	0	0	0	-2.6
	3	1	10.00	0	0	66.10	145.6	2.39	-0.21	0	0	0	-2.6
	3	2	10.00	0	0	75.91	145.6	2.39	-0.21	0	0	0	-2.6
	3	3	7.50	0	0	95.78	145.6	1.35	-0.1	0	0	0	-2.6
	4	1	10.00	0	0	66.47	145.6	2.39	-0.21	0	0	0	-2.6
	4	2	10.00	0	0	74.39	145.6	2.39	-0.21	0	0	0	-2.6
	4	3	7.50	0	0	69.48	145.6	1.35	-0.1	0	0	0	-2.6
	5	1	10.00	0	0	102.37	145.6	2.39	-0.21	0	0	0	-2.6
	5	2	10.00	0	0	108.99	145.6	2.39	-0.21	0	0	0	-2.6
	5	3	7.50	0	0	107.08	145.6	1.35	-0.1	0	0	0	-2.6
12 Productive forest	1	1	126.76	5.84	9.51	82.65	145.6	3.60	-2.38	0.00	-0.14	0.00	-2.6
	1	2	124.81	5.67	7.53	102.03	145.6	3.21	-2.28	-0.15	-0.09	0.00	-2.6
	1	3	84.75	6.02	7.76	121.34	145.6	1.95	-1.36	-0.04	-0.17	0.00	-2.6
	2	1	133.95	9.14	8.70	55.40	145.6	4.63	-4.77	-0.04	-0.12	0.00	-2.6
	2	2	146.61	8.80	11.42	62.12	145.6	4.63	-4.61	-0.06	-0.09	0.00	-2.6
	2	3	101.51	8.80	11.42	122.00	145.6	1.60	-1.05	-0.06	-0.09	0.00	-2.6
	3	1	135.29	9.52	7.51	66.10	145.6	4.56	-3.35	0.00	-0.05	0.00	-2.6
	3	2	147.51	8.46	16.29	75.91	145.6	4.15	-3.78	-0.07	-0.06	0.00	-2.6
	3	3	119.40	9.30	26.21	95.78	145.6	2.48	-2.75	-0.02	-0.13	0.00	-2.6
	4	1	94.68	6.88	3.15	66.47	145.6	3.24	-3.19	0.02	-0.20	0.00	-2.6
	4	2	104.44	7.80	19.99	74.39	145.6	2.49	-2.59	0.00	-0.15	0.00	-2.6
	4	3	96.36	8.15	33.37	69.48	145.6	1.81	-2.47	-0.06	-0.06	0.00	-2.6
	5	1	71.66	2.34	8.22	102.37	145.6	2.74	-0.92	-0.10	-0.13	0.00	-2.6
	5	2	76.87	3.06	11.03	108.99	145.6	2.20	-0.61	-0.01	-0.17	0.00	-2.6
	5	3	76.48	3.28	30.77	107.08	145.6	1.61	-0.30	0.08	-0.02	0.00	-2.6
13 Unproductive forest	1	1	38.53	0	9.51	82.65	145.6	0	0	0	0	0	-2.6
	1	2	51.10	0	7.53	102.03	145.6	0	0	0	0	0	-2.6
	1	3	51.34	0	7.76	121.34	145.6	0	0	0	0	0	-2.6
	2	1	20.45	0	8.70	55.40	145.6	0	0	0	0	0	-2.6
	2	2	35.83	0	11.42	62.12	145.6	0	0	0	0	0	-2.6
	2	3	51.33	0	11.42	122.00	145.6	0	0	0	0	0	-2.6
	3	1	20.45	0	7.51	66.10	145.6	0	0	0	0	0	-2.6
	3	2	47.53	0	16.29	75.91	145.6	0	0	0	0	0	-2.6
	3	3	42.36	0	26.21	95.78	145.6	0	0	0	0	0	-2.6
	4	1	21.60	0	3.15	66.47	145.6	0	0	0	0	0	-2.6
	4	2	31.48	0	19.99	74.39	145.6	0	0	0	0	0	-2.6
	4	3	29.88	0	33.37	69.48	145.6	0	0	0	0	0	-2.6
	5	1	20.83	0	8.22	102.37	145.6	0	0	0	0	0	-2.6
	5	2	23.82	0	11.03	108.99	145.6	0	0	0	0	0	-2.6
	5	3	24.35	0	30.77	107.08	145.6	0	0	0	0	0	-2.6

(Table 6-4 continued)

Combination category (CC)	NFI region	Elevation zone z	Carbon stock in living biomass (stockCl,i)	Carbon stock in dead wood (stockCd,i)	Carbon stock in litter (stockCh,i)	Carbon stock in mineral soil (stockCs,i)	Carbon stock in organic soil (stockCs,i)	Gain of living biomass (gainCl,i)	Loss of living biomass (lossCl,i)	Net change in dead wood (changeCd,i)	Net change in litter (changeCh,i)	Net change in mineral soil (changeCs,i)	Net change in organic soil (changeCs,i)
21 Cropland	n.s.	n.s.	4.34	0	0	53.40	240	0.00	-0.34	0	0	0	-9.52
31 Permanent Grassland	n.s.	1	7.08	0	0	62.02	240	0	0	0	0	0	-9.52
	n.s.	2	6.00	0	0	67.50	240	0	0	0	0	0	-9.52
	n.s.	3	7.95	0	0	75.18	240	0	0	0	0	0	-9.52
32 Shrub Vegetation	n.s.	1	20.45	0	0	62.02	240	0	0	0	0	0	-5.3
	n.s.	2	20.45	0	0	67.50	240	0	0	0	0	0	-5.3
	n.s.	3	20.45	0	0	75.18	240	0	0	0	0	0	-5.3
33 Vineyards et al.	n.s.	n.s.	3.74	0	0	53.40	240	0	0	0	0	0	-9.52
34 Copse	n.s.	1	20.45	0	0	62.02	240	0	0	0	0	0	-5.3
	n.s.	2	20.45	0	0	67.50	240	0	0	0	0	0	-5.3
	n.s.	3	20.45	0	0	75.18	240	0	0	0	0	0	-5.3
35 Orchards	n.s.	n.s.	24.62	0	0	64.76	240	0	0	0	0	0	-9.52
36 Stony Grassland	n.s.	n.s.	7.16	0	0	26.31	240	0	0	0	0	0	-5.3
37 Unproductive Grassland	n.s.	n.s.	7.87	0	0	68.23	240	0	0	0	0	0	-5.3
41 Surface Waters	n.s.	n.s.	0	0	0	0	240	0	0	0	0	0	0
42 Unproductive Wetland	n.s.	n.s.	6.50	0	0	68.23	240	0	0	0	0	0	-5.3
51 Buildings, Constructions	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0
52 Herbaceous Biomass in S.	n.s.	n.s.	9.54	0	0	53.40	240	0	0	0	0	0	-9.52
53 Shrubs in Settlements	n.s.	n.s.	15.43	0	0	53.40	240	0	0	0	0	0	-5.3
54 Trees in Settlements	n.s.	n.s.	20.72	0	0	53.40	240	0	0	0	0	0	-5.3
61 Other Land	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0

Legend			
Elevation zones:		NFI regions:	
1	< 601 m	1	Jura
2	601 - 1200 m	2	Central Plateau
3	> 1200 m	3	Pre-Alps
		4	Alps
		5	Southern Alps
			n.s. = no stratification
			Annual data

6.1.5 Uncertainty estimates

Table 6-5 gives an overview of uncertainty estimates of activity data (AD) and of emission factors (EF). For categories 4A–4F (highlighted in yellow), the uncertainties of AD mainly depend on the uncertainty of the AREA survey data (see chp. 6.3.3, Table 6-10). For categories 4D1, 4(II)–4(V) and 4G other data sources are relevant; they are presented in detail in the respective chapters (6.X.3) of the LULUCF categories, along with the uncertainty estimates of EF.

In general, AD uncertainty is lower than EF uncertainty, because AD are mostly based on a systematic survey with high spatial resolution (such as AREA), while EFs include parameters that are difficult to measure or to model such as carbon stocks in biomass, growth rates and biogeochemical processes.

Table 6-5 Uncertainty estimates in the LULUCF sector, expressed as half of the 95% confidence intervals. Highlighted AD uncertainties depend mainly on the uncertainty of the AREA survey.

IPCC category		Gas	Activity data uncertainty	Emission factor uncertainty
			%	%
4A1	Forest land remaining forest land	CO ₂	1.1	88.6
4A2	Land converted to forest land	CO ₂	1.6	88.6
4B1	Cropland remaining cropland	CO ₂	4.9	26.4
4B2	Land converted to cropland	CO ₂	5.2	31.6
4C1	Grassland remaining grassland	CO ₂	5.2	867.9
4C2	Land converted to grassland	CO ₂	5.3	30.7
4D1	Wetlands remaining wetlands	CO ₂	61.9	72.2
4D2	Land converted to wetlands	CO ₂	4.0	21.4
4E1	Settlements remaining settlements	CO ₂	4.4	50.0
4E2	Land converted to settlements	CO ₂	4.5	50.0
4F1	Other land remaining other land	CO ₂	NA	NA
4F2	Land converted to other land	CO ₂	3.1	50.0
4(II)	Drained organic soils	N ₂ O	34.2	137.5
4(II)D2	Flooded land	CH ₄	10.0	70.0
4(III)	N mineralization	N ₂ O	5.3	135.0
4(IV)2	Leaching (indirect emissions)	N ₂ O	20.0	161.5
4(V)	Wildfires	CO ₂	NA	NA
4(V)	Wildfires	CH ₄	30.0	70.0
4(V)	Wildfires	N ₂ O	30.0	70.0
4G	Harvested wood products	CO ₂	4.5	57.0

6.2 Land-use definitions and classification systems

6.2.1 Combination Categories (CC) as derived from AREA Land Use Statistics

The nomenclature of the Swiss Land Use Statistics (AREA) evaluated by the Swiss Federal Statistical Office (SFSO 2006a) is the basis for the land-use categories and subcategories used for land area representation in the LULUCF sector. In the course of the AREA surveys (see chp. 6.3.1), every hectare of Switzerland's territory was assigned to a land-use category (NOLU04) and to a land-cover category (NOLC04) (SFSO nomenclature version 2004). The 46 NOLU04 categories and 27 NOLC04 categories of AREA were aggregated to 18 combination categories (CC) as shown in Table 6-6, thus implementing the main categories proposed by IPCC as well as country-specific sub-divisions (see Table 6-2). The first digit of the CC code represents the main land-use category according to IPCC, whereas the second digit stands for respective sub-divisions.

All subcategories of Forest land, Grassland and Wetlands are defined as managed and reported under managed land in CRF Table 4.1 (Cropland and Settlements are regarded to be managed by default; Other land (4F) is regarded to be unmanaged by default). In a

nutshell, the entire land area of Switzerland – except for 4F Other land – is reported to be managed.

The sub-divisions were defined with respect to possible differentiation of biomass densities, carbon turnover, and soil carbon contents. They were defined in 2006 in an evaluation process involving experts from the FOEN, the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), the Swiss Federal Statistical Office and Agroscope as well as private consultants. The evaluation process resulted in the elaboration of Table 6-6. The definition of the CCs was strongly influenced by the NOLC04/NOLU04 classification of the Swiss Land Use Statistics AREA (SFSO 2006a). Most criteria and thresholds as defined therein were adopted. Some examples are:

- For Forest land, the criteria correspond to NFI thresholds with respect to minimum area, width, crown cover, and tree height.
- For NOLC04 31 (land cover shrub), the criteria include: vegetation height <3 m, degree of coverage >80%, dominated by shrubs, dwarf-shrubs, and bushes.
- For NOLC04 32 (land cover brush meadows), the criteria include vegetation height <3 m, degree of coverage 50–80%, dominated by shrubs, dwarf-shrubs, and bushes.

With regard to carbon content in biomass, there is a strong relation to the vegetation type (i.e. land cover in most cases). This is exemplarily reflected by the mainly horizontal arrangement of the individual CCs in Table 6-6. With regard to carbon turnover and soil organic carbon the CC definition was driven by the consideration that most vegetation units are subject to a similar management that leads to comparable carbon fluxes in biomass and soil.

For individual CCs (especially Forest Land, i.e. CC11, CC12, CC13) further spatial stratifications were introduced (cf. chp. 6.2.2) with the intent to approximate the real/natural differences in carbon stock, carbon turnover and soil conditions as good as possible.

The underlying criteria to include land use subcategories such as shrub vegetation, vineyards, low-stem orchards, tree nurseries, copse and orchards under Grassland with woody biomass are: (1) They do not fulfil the criteria for forests; (2) There is an agricultural management in general; (3) They all have woody biomass (i.e. perennial vegetation) with grass understory. Under Cropland, in contrast, there are no perennial crops, but annual crops and leys in arable rotations. All perennial crops are included in the Grassland subcategories.

6.2.2 Spatial stratification

In order to quantify carbon stocks and GHG emissions and removals in the LULUCF sector as accurately as possible, Switzerland's territory was stratified by means of three site criteria: soil type (mineral or organic), elevation and forest production region.

Soil Type

Most soils in Switzerland are mineral soil types. A digital map showing estimates of the surface of organic soils in Switzerland was elaborated by Wüst-Galley et al. (2015). As there is no single data set from which the location of organic soils across the country could be adequately deduced, the authors evaluated numerous spatial and non-spatial data sets providing information on geology, soils, forest habitats and vegetation. According to Wüst-Galley et al. (2015) the total area of organic soils is 28 kha (0.8% of the total area covered by soils).

The definition of organic soils in the GHG inventory is as follows:

Intact or degraded peaty soils are considered organic soils. Where information on soil organic carbon (SOC) is known, the definition of organic soils from the IPCC (IPCC 2006, Volume 4, chp. 3, Annex 3A.5) was used to classify soils as mineral / organic (see Wüst-Galley et al. 2015: 11). Thus, this definition was used for the ground-truthing of forest habitat maps and fen inventories. This definition also formed the basis of the classification of soil types from the soils maps, as organic or mineral. Here however, two soils types ("anmoorig" and "antorfig" soils) could not be classified; these have a ranges of SOC and peat depth that are wider than those given in the IPCC definition, meaning they cannot be classified as either mineral or organic soils. Due to lack of information regarding their distribution, they were not explicitly considered in the estimate of organic soils (see Wüst-Galley et al. 2015: 14-15 and 61); including these additional soil types would lead to inconsistency of the definition of organic soils across the country, because their distribution is known for a small area in Switzerland.

For the other data sets used in the construction of the organic soils map (geology maps, hydrogeology maps and other habitat maps), no information on SOC is available, and the presence of peat was used as evidence of organic soils. The carbon content of peat meets the IPCC definition of organic soils.

Consistency: A single map of organic soils is applied to all years (1990 to present), meaning the classifications used are consistent through time. The same definition of organic soils was used across the whole country.

Elevation

For Forest land (CC11-CC13) and permanent grassland (CC31), three elevation zones were differentiated: <601 m a.s.l. (meters above sea level), 601-1200 m a.s.l., and >1200 m a.s.l. (Figure 6-4). Elevation data from the Federal Office of Topography (swisstopo.ch) on a 25x25 m raster (product DHM25) were used to map the three zones.

Forest production region

Forest land was furthermore differentiated into the five production regions of the National Forest Inventory (EAFV/BFL 1988; Brassel and Brändli 1999; Brändli 2010). The NFI regions were adopted from EAFV/BFL (1988) as shown in Figure 6-4:

1. Jura, 2. Central Plateau, 3. Pre-Alps, 4. Alps, 5. Southern Alps.

Applying all spatial stratifications, 30 different strata (referred to as subscript i in chp. 6.1.3.2) would be theoretically possible. Not all of them, but altogether 29 have been actually realised and applied for the calculation of LULUCF-associated carbon emissions and removals.




Figure 6-4 Map showing the spatial stratification according to NFI production region, elevation zone, and soil type.

6.2.3 The land-use tables and change matrices

In Table 6-7 the land-use statistics resulting from spatial stratification (chp. 6.2.2) and interpolation in time (chp. 6.3.2) are exemplarily shown for the year 1990. The table gives also the size of the individual spatial strata.

Table 6-7 Land use (in terms of combination categories CC) projection by the end of 1990, stratified separately for elevation (3 zones), soil type (mineral or organic) and NFI region (1-5), in kha.

CC:	11	12	13	21	31	32	33	34	35	36	37	41	42	51	52	53	54	61	Sum
Altitude																			
<601 m	1.1	224.8	6.2	299.8	153.8	2.6	22.5	32.4	1.2	0.5	2.9	138.6	5.2	116.7	47.5	2.8	18.6	2.0	1079.1
601-1200 m	1.4	504.2	18.1	131.7	358.2	8.7	3.9	29.9	0.3	2.5	1.5	9.7	5.7	46.4	17.0	0.9	5.3	8.1	1153.9
>1200 m	1.4	377.8	79.9	0.4	425.4	144.4	0.0	27.1	0.0	148.7	61.9	13.3	14.3	11.4	3.7	0.2	1.0	585.2	1896.1
	3.9	1106.9	104.2	432.0	937.4	155.7	26.5	89.3	1.6	151.6	66.3	161.6	25.1	174.6	68.2	3.9	24.9	595.3	4129.0
Soil																			
mineral	3.9	1103.3	104.1	420.3	931.8	155.6	26.4	89.0	1.6	151.6	66.1	161.3	21.4	173.3	67.7	3.9	24.8	595.3	4101.3
organic	0.0	3.6	0.2	11.7	5.6	0.1	0.0	0.4	0.0	0.0	0.2	0.3	3.7	1.2	0.5	0.0	0.1	0.025	27.7
	3.9	1106.9	104.2	432.0	937.4	155.7	26.5	89.3	1.6	151.6	66.3	161.6	25.1	174.6	68.2	3.9	24.9	595.3	4129.0
NFI region																			
1	0.7	197.2	8.3	78.0	122.6	0.9	4.7	11.9	0.3	0.2	0.6	23.6	1.2	26.8	10.9	0.5	4.7	0.5	493.5
2	0.8	227.2	4.1	307.0	152.4	0.9	9.9	27.4	1.0	0.2	1.6	70.4	4.1	84.9	34.7	1.6	12.6	0.7	941.5
3	1.0	214.3	13.0	30.2	261.3	10.4	0.8	17.8	0.1	8.5	6.8	30.6	12.0	26.8	9.2	0.5	2.9	15.0	661.2
4	1.1	331.6	56.1	13.8	365.4	110.2	9.5	24.5	0.2	118.1	49.2	26.2	7.2	26.9	9.8	0.8	3.0	524.8	1678.2
5	0.3	136.6	22.6	3.0	35.7	33.3	1.5	7.8	0.0	24.6	8.1	10.7	0.7	9.2	3.7	0.6	1.9	54.2	354.6
	3.9	1106.9	104.2	432.0	937.4	155.7	26.5	89.3	1.6	151.6	66.3	161.6	25.1	174.6	68.2	3.9	24.9	595.3	4129.0

Table 6-8 shows the overall trends of land-use changes between 1990 and 2015. For example, the area of afforestations (CC11) decreased by 77% during this period, while the area of unproductive forests (CC13) increased by 3%. CC11 is decreasing because the area of new afforestations has been decreasing during this period and because most of the afforestations turn to productive forests after a certain time period.

Table 6-8 Statistics of land use (in terms of combination categories CC) and relative change (%) between 1990 and 2015, in kha.

CC:	11	12	13	21	31	32	33	34	35	36	37	41	42	51	52	53	54	61	Sum
Year:																			
1990	3.9	1106.9	104.2	432	937.4	155.7	26.5	89.34	1.56	151.6	66.3	161.6	25.1	174.6	68.2	3.93	24.9	595.3	4129.0
1991	3.8	1109.1	104.5	431.2	935.6	155.2	26.5	88.3	1.5	151.4	66.2	161.6	25.1	176.2	68.7	4.0	25.3	595.0	4129.0
1992	3.7	1111.3	104.7	430.4	933.8	154.7	26.6	87.3	1.4	151.1	66.0	161.6	25.1	177.9	69.3	4.0	25.6	594.6	4129.0
1993	3.5	1113.4	104.9	429.4	932.3	154.2	26.5	86.2	1.4	150.9	65.8	161.6	25.1	179.5	69.9	4.1	25.9	594.2	4129.0
1994	3.4	1115.3	105.1	428.0	931.6	153.7	26.5	85.2	1.3	150.7	65.7	161.6	25.1	181.1	70.4	4.2	26.2	593.9	4129.0
1995	3.2	1117.1	105.3	426.1	931.5	153.1	26.5	84.3	1.3	150.6	65.5	161.6	25.1	182.7	71.1	4.2	26.2	593.5	4129.0
1996	3.0	1118.7	105.4	424.2	931.6	152.6	26.4	83.4	1.3	150.5	65.4	161.6	25.2	184.3	71.9	4.2	26.2	593.1	4129.0
1997	2.8	1120.2	105.6	422.2	931.9	152.2	26.3	82.5	1.2	150.4	65.3	161.7	25.2	185.8	72.7	4.2	26.1	592.8	4129.0
1998	2.6	1121.5	105.7	420.2	932.2	152.0	26.2	81.6	1.2	150.4	65.1	161.7	25.2	187.3	73.5	4.2	25.9	592.4	4129.0
1999	2.4	1122.8	105.8	418.2	932.5	151.8	26.1	80.7	1.2	150.5	65.0	161.7	25.2	188.8	74.3	4.2	25.8	592.1	4129.0
2000	2.2	1124.1	105.9	416.2	932.7	151.6	26.1	79.8	1.1	150.5	64.8	161.8	25.2	190.4	75.2	4.2	25.7	591.8	4129.0
2001	2.0	1125.4	105.9	414.2	933.0	151.4	26.0	78.8	1.1	150.5	64.7	161.8	25.2	191.9	76.0	4.2	25.5	591.4	4129.0
2002	1.8	1126.7	106.0	412.2	933.3	151.1	25.9	77.9	1.1	150.6	64.5	161.8	25.2	193.4	76.8	4.2	25.4	591.1	4129.0
2003	1.5	1128.0	106.1	410.2	933.6	150.9	25.8	77.0	1.0	150.6	64.4	161.9	25.3	194.9	77.6	4.2	25.3	590.7	4129.0
2004	1.3	1129.3	106.2	408.2	933.8	150.7	25.7	76.1	1.0	150.6	64.2	161.9	25.3	196.5	78.4	4.2	25.1	590.4	4129.0
2005	1.2	1130.8	106.2	406.7	933.6	150.4	25.6	75.0	1.0	150.6	64.0	162.0	25.3	198.0	79.2	4.2	24.9	590.1	4129.0
2006	1.1	1132.3	106.3	405.6	933.4	150.2	25.5	73.8	1.0	150.7	63.9	162.0	25.3	199.6	79.6	4.2	24.7	589.8	4129.0
2007	1.0	1133.9	106.4	404.6	933.3	150.1	25.4	72.4	0.9	150.7	63.7	162.1	25.3	201.1	80.0	4.1	24.5	589.5	4129.0
2008	1.0	1136.0	106.7	404.0	933.3	149.6	25.2	71.2	0.9	150.7	63.4	162.2	25.2	202.1	80.2	4.1	24.2	588.8	4129.0
2009	1.0	1137.5	106.8	402.3	933.0	149.3	25.2	70.4	0.9	150.8	63.3	162.2	25.3	203.7	80.8	4.1	24.2	588.3	4129.0
2010	0.9	1138.6	106.9	400.5	932.6	149.2	25.1	69.7	0.9	150.8	63.1	162.3	25.3	205.3	81.4	4.1	24.2	588.1	4129.0
2011	0.9	1139.7	107.0	398.8	932.3	149.1	25.0	69.1	0.9	150.8	63.0	162.3	25.3	207.0	82.0	4.1	24.2	587.8	4129.0
2012	0.9	1140.8	107.0	397.0	931.9	148.9	25.0	68.4	0.8	150.8	62.9	162.3	25.3	208.6	82.6	4.1	24.2	587.6	4129.0
2013	0.9	1142.0	107.2	395.8	931.1	148.8	24.9	67.7	0.8	150.8	62.7	162.4	25.3	210.2	83.1	4.1	24.0	587.3	4129.0
2014	0.9	1143.2	107.3	394.2	930.8	148.7	24.8	67.1	0.8	150.8	62.6	162.4	25.3	211.7	83.5	4.1	23.9	587.1	4129.0
2015	0.9	1144.3	107.3	392.5	930.3	148.5	24.7	66.4	0.8	150.8	62.5	162.4	25.3	213.3	84.1	4.1	23.9	586.8	4129.0
Change:	-77	3	3	-9	-1	-5	-7	-26	-50	-1	-6	1	1	22	23	4	-4	-1	0

The annual land-use changes across the entire territory of Switzerland (change-matrices, see examples for 1990 and 2015 in Table 6-9) were obtained by adding up the annual changes on a hectare basis per combination category (CC). For calculating the carbon stock

changes, fully stratified (cf. chp. 6.2.2) land-use change tables were used for each year (Meteotest 2017). More aggregated change-matrices are reported in CRF Table4.1 for each year in the period 1990–2015.

It is worth noting that in general the numbers given in the change-matrices (Table 6-9) cannot be directly compared with the figures of category 2 in CRF Table4.A, Table4.B, Table4.C, Table4.D, Table4.E, and Table4.F (Land converted to X), where the cumulative area remaining in the respective category in the reporting year is recorded (cf. the description of conversion time of 20 years in chp. 6.1.3.3). In contrast, the change matrices present the land-use changes occurring in the specified year only.

Table 6-9 Annual land-use changes in 1990 and in 2015 (change matrices). Units: ha/year, rounded values. Empty cells indicate that no change occurred.

1990		change to CC																			decrease
		11	12	13	21	31	32	33	34	35	36	37	41	42	51	52	53	54	61		
change from CC	11		369	1	0	0	1		0						0	0	0		0	372	
	12			158	5	125	86	6	59		12	19	11	7	117	27	11	17	49	709	
	13		678		8	354	48	5	89	0	3	3	1	3	41	20	3	15	10	1280	
	21	8	1	5		663	6	181	35	1	4	4	4	4	632	317	21	18	22	1926	
	31	136	166	480	717		1007	123	311	4	46	43	9	11	870	490	27	44	67	4554	
	32	24	1022	715	2	126		9	309		14	15	6	0	24	8	5	3	30	2313	
	33	1	2	4	126	65	4	28		2	0	1	0		50	26	4	3	5	323	
	34	20	536	63	143	866	49	35		11	9	23	4	3	171	94	6	41	14	2087	
	35		0	0	8	13	0	4	46						4	2	0	1	0	80	
	36	3	27	26	2	162	243	1	41			89	4	0	8	1	0		45	652	
	37	7	26	6	1	8	234	1	68		10		3	0	6	2		0	13	384	
	41	0	4	1	2	2	6	0	4		4	1		17	11	2	1	0	99	156	
	42	5	27	6	1	3	2	0	2		0	0	6		4	1	0	0	1	59	
	51	38	18	4	86	158	11	5	7		3	5	6	4		271	58	46	5	726	
	52	7	4	1	16	32	3	1	1		0	1	1	2	349		68	387	0	874	
	53	5	9	0	6	7	2	0	2				0	2	45	28		46	0	150	
54	2	6	0	1	2	0	0	3			0	0	1	78	152	8		0	253		
61	4	41	17	16	67	93	8	31		287	33	96	2	13	1	0	1		709		
increase	261	2936	1489	1140	2653	1794	381	1036	18	394	236	152	55	2425	1443	211	621	361	17607		
2015		change to CC																			decrease
		11	12	13	21	31	32	33	34	35	36	37	41	42	51	52	53	54	61		
change from CC	11		64	0		0								0					65		
	12			318	1	232	139	2	78		33	20	16	15	95	26	11	11	74	1069	
	13		677		2	398	76	2	68		5	2	1	2	31	16	1	9	11	1301	
	21	2	1	1		1780	6	141	26	1	6	11	6	8	500	260	10	4	16	2779	
	31	16	115	415	877		793	73	222	2	84	33	8	12	806	446	17	12	87	4016	
	32	3	661	525	2	143		3	288		20	12	7	1	13	4	2	1	33	1719	
	33	0	2	2	119	93	4		20	1	1	0		0	34	26	2	2	5	309	
	34	2	490	63	39	576	58	11		4	10	21	6	1	94	57	1	21	16	1470	
	35				1	6		1	16						1	0				25	
	36	0	17	22	2	90	204	0	44			56	4		5	1			44	490	
	37	2	15	3	1	5	196		50		16		3	1	4	1			13	310	
	41	0	3	0	0	1	6		2		3	3		11	5	1	0		102	138	
	42	0	30	3	0	1	1		1		0	0	8		1	1		0	1	48	
	51	17	11	2	55	137	8	2	4		7	7	6	2		313	50	26	6	651	
	52	7	5	1	14	43	3	1	2		1	2	1	2	465		62	309	0	918	
	53	3	14	0	4	10	3	0	1			1		0	53	43		42		174	
54	1	6	1	0	1	0		2			0	0	0	115	314	22			463		
61	1	27	12	14	44	74	5	30		316	16	103	1	6	1	0			648		
increase	54	2139	1366	1131	3559	1569	242	852	8	501	186	170	55	2227	1509	179	436	409	16592		

6.3 Approaches used for representing land areas, land-use databases

6.3.1 Swiss Land Use Statistics (AREA)

Data of the Swiss Land Use Statistics (AREA) evaluated by the Swiss Federal Statistical Office (SFSO 2016d) form the basis of activity data. In the course of the AREA surveys, every hectare of Switzerland's territory (4'129 kha) is assigned to one of 46 land-use categories and to one of 27 land-cover categories by means of stereographic interpretation of aerial photos (SFSO 2006a).

For the reconstruction of the land use conditions in Switzerland during the period 1990–2015 four datasets are used:

- Land Use Statistics "1979/85" (AREA1), status: completed
- Land Use Statistics "1992/97" (AREA2), status: completed
- Land Use Statistics "2004/09" (AREA3), status: completed
- Land Use Statistics "2013/18" (AREA4), status: 25% of territory processed.

The aerial photos for AREA1, AREA2 and AREA3 were taken 1977–1986, 1990–1998 and 2004–2009, respectively. In the course of AREA3 all photos were simultaneously (re-) interpreted according to the newly designed AREA set of land-use and land-cover categories based on the nomenclature 'NOAS04' (SFSO 2006a). The AREA4 survey was started in 2014; in this submission data based on aerial photos 2012–2014 were included.

The inter-survey period is not identical throughout the Swiss territory, but varies regionally. It averages approximately 12 years for AREA1, AREA2 and AREA3; for AREA4 the period will be shorter, approximately 9 years. This methodical characteristic needs to be considered when reconstructing the annual country-wide status of land use or when calculating annual rates of land-use change.

6.3.2 Interpolation of the status for each year

The exact dates of aerial photo shootings are known for each hectare. However, the exact occurrence date (year) of a land-use change on a specific hectare is unknown. The actual change can have taken place in any year between two AREA surveys. In this study, it was assumed that the probability of a land-use change from AREA1 to AREA2, from AREA2 to AREA3 and from AREA3 to AREA4 is uniformly distributed over the respective interim period between two surveys. Therefore, the land-use change of each hectare has to be equally distributed over its specific interim period.

Thus, the land-use status for the years between two data collection dates can be calculated by linear interpolation. Dates of aerial photo shootings (i.e. starting and ending year of the inter-survey period) and the land-use categories of AREA1, AREA2, AREA3 and AREA4 for every hectare were used for these calculations. An example is shown in Figure 6-5: A hectare had been assigned to the land-use category Cropland in AREA1 (aerial photo in 1980). A land-use change to 'Surrounding of Buildings' was discovered 10 years later (1990) in AREA2.




Figure 6-5 Hypothetical development of land use for a supposed survey period 1980–2020. The linear land-use changes between AREA1, AREA2, AREA3 and AREA4 considering as example a hectare changing from “Cropland” to “Surrounding of Buildings” and later from “Surrounding of Buildings” to “Buildings”. For 2020, a linear interpolation has been carried out between AREA4 and a virtual fifth survey (AREA5v) that was modelled for the year 2025 (here resulting in no change of land use).

The “state 1988” of that hectare is determined by calculating the fractions of the two land-use categories for the year 1988. A linear development from “Cropland” to “Surrounding of Buildings” during the whole interim period was assumed. Thus, in 1988 the hectare was split up in two fractions: 80% is “Surrounding of Buildings” and 20% is “Cropland”. The same procedure can be applied for two survey dates between AREA2 and AREA3 (here exemplarily shown for the period 1990–2004, highlighting “state 2003”) or between AREA3 and AREA4 (here exemplarily shown for the period 2004–2016, highlighting “state 2009”).

After completion, AREA4 will comprehend aerial photos from 2013–2018.

To obtain consistent and complete nationwide data for each year during the ongoing AREA4 survey two analyses are required:

1) For those hectares that currently have no data in AREA4, the land-use states after AREA3 were interpolated between AREA3 and a “virtual” 4th survey (AREA4v). AREA4v was modelled for each sample point using a Markov-chain approach, where transition probabilities between AREA3 and AREA4v were assessed based on the transition distribution between AREA2 and AREA3 within each spatial stratum (Sigmaplan 2017).

2) For those hectares that were already covered by AREA4, the land-use states after the flight year of AREA4 were interpolated between AREA4 and a "virtual" 5th survey (AREA5v). AREA5v was modeled for each sample point using a Markov-chain approach, where transition probabilities between AREA4 and AREA5v were assessed based on the transition distribution between AREA3 and AREA4 within each spatial stratum (Sigmaplan 2017). Therefore, the land-use changes occurring after the flight year of AREA4 (i.e. when it was covered) were calculated from the linear development detected between AREA4 and the virtual 5th survey AREA5v for this type of hectare (regarding CC and spatial strata) (see Figure 6-5: example "state 2020").

The "virtual survey" approach used in 1) and 2) was evaluated successfully by modelling a "virtual" AREA3 from transition probabilities between AREA1 and AREA2 and comparing the results to the published AREA3 data (Sigmaplan 2017).

The status for each individual year in the period 1990–2015 for the whole Swiss territory results from the summation of the fractions of all hectares per combination category CC, additionally considering the spatial strata where appropriate.

6.3.3 Uncertainties and time-series consistency of activity data

An overview of uncertainty estimates for activity data (AD) and emission factors (or biomass parameters) is shown in Table 6-5. Details related to uncertainties of AREA data are presented in this chapter, while uncertainties of other AD (such as wild fires and consumption of Harvested wood products) and emission factors are presented in the respective chapters (6.X.3) of the LULUCF categories.

In most cases, the uncertainty of AD for categories 4A–4F depends on the quality of the AREA survey data. For categories with relevant emissions from drained organic soils, also the uncertainty of the spatial allocation of organic soils (see chp. 6.2.2 and below) was considered.

The uncertainty of AREA-based activity data has two main sources (Table 6-10). They were quantified on the basis of the AREA data (SFSO 2016d) as follows:

1) Interpretation error: In the AREA survey, the first classification of the aerial photos is checked by a second independent interpreter. The portion of sampling points with a mismatch of the first and the second interpretation was used as the uncertainty of the interpretation. This uncertainty of interpretation integrates all errors related to the manual interpretation of land-use and land-cover classes on aerial photographs. While it is clear that this is rather an estimate of the maximum potential interpretation error than of the actual interpretation error, it is reported hereafter unless more accurate information is available.

2) Statistical sampling error: In the AREA survey, the land-use types are interpreted on points situated on a regular 100x100 m grid. Thus, the uncertainty of the surface area covered by a certain land-use type or land-use change decreases with increasing numbers of sampling points. Assuming a binomial distribution of the errors, this uncertainty was calculated as

$$U_{\text{sampling}} = 100 * 1.96 * (\text{number of points})^{-0.5}$$

The number of sampling points lies between 2'506 (for 4D2) and 1'356'570 (for 4C1) leading to values of U_{sampling} between 3.9% and 0.2%.

The overall uncertainty was calculated as:

$$U_{\text{overall}} = (U_{\text{interpret}}^2 + U_{\text{sampling}}^2)^{0.5}$$

Table 6-10 Sources of AD uncertainty and overall uncertainties in the allocation of land-use categories, expressed as half of the 95% confidence intervals. Calculations are based on AREA data from SFSO (2016d).

Category	Description	Interpretation uncertainty	Sampling uncertainty	Overall uncertainty
4A1	Forest land remaining forest land	1.1	0.2	1.1
4A2	Land converted to forest land	1.1	1.1	1.6
4B1	Cropland remaining cropland	4.9	0.3	4.9
4B2	Land converted to cropland	4.9	1.8	5.2
4C1	Grassland remaining grassland	5.2	0.2	5.2
4C2	Land converted to grassland	5.2	1.0	5.3
4D1	Wetlands remaining wetlands	0.9	0.5	1.0
4D2	Land converted to wetlands	0.9	3.9	4.0
4E1	Settlements remaining settlements	4.4	0.4	4.4
4E2	Land converted to settlements	4.4	1.1	4.5
4F1	Other land remaining other land	1.4	0.3	1.4
4F2	Land converted to other land	1.4	2.8	3.1

An analysis of the uncertainty of the spatial allocation of organic soils published by Wüst-Galley et al. (2015) resulted in: 37.8% for Cropland, 55.9% for Grassland and 61.9% for Wetlands. For Forest land and Settlements, the emission from organic soils were not considered in the calculation of the overall uncertainty (Meteotest 2017).

Activity data for wildfires were taken from the Swissfire database (see chapters 6.4.2.12 and 6.6.2.5). The uncertainty of those areas was estimated as 30% by expert judgment.

Consistency: Time series for activity data are all considered consistent; they were calculated based on consistent methods for interpolation and extrapolation and homogenous databases.

6.3.4 QA/QC and verification of activity data

The general QA/QC measures are described in chp. 1.2.3.

The AREA survey is a well-defined and controlled, long-term process in the responsibility of the Swiss Federal Statistical Office (SFSO 2006a). The data supplied by SFSO (2016d) were checked for consistency (Sigmaplan 2017).

The temporal interpolation and extrapolation of the AREA sample is quite a complex procedure, whose internal consistency was checked systematically as described in Sigmaplan (2017). Further checks (interannual comparisons, plausibility) were carried out after producing the land-use change tables presented in chp. 6.2.3.

A systematic cross-check between the activity data reported under LULUCF category 4A Forest land and under the Kyoto Protocol activity Forest management was carried out (see chp. 11.3.2.2).

It was checked and confirmed that the total country area remains constant over the inventory period.

6.3.5 Recalculations of activity data

The AD time series 1991–2014 was updated as a result of the following activities:

The most recent land-use data from the fourth area survey (AREA4) were included (SFSSO 2016d). They are based on aerial photographs from 2012 and 2014. The interpolation and projection procedures were adapted accordingly (see chp. 6.3.2 and Sigmaplan 2017).

Along with the AREA4 survey the SFSSO continuously performs consistency checks and corrections in the data of AREA3 (2004–2009). As a consequence, there was a decrease in the area of cropland in favour of grassland (see Sigmaplan 2017 for details).

6.3.6 Planned improvements for activity data

The uncertainty of Switzerland's activity data for land areas will decrease gradually with the increase in the sample size of the current survey AREA4. Interpretation and further processing of AREA4 is expected to be completed in 2020.

In response to UNFCCC (2017/ID#L.7), the description of the identification of the country-specific combination categories and subdivisions (cf. Table 6-2) will be improved.

6.4 Category 4A – Forest land

6.4.1 Description

Table 6-11 Key categories (KCA including LULUCF) in category 4A.

Code	IPCC Category	GHG	Identification Criteria
4A1	Forest land remaining Forest land	CO ₂	L1, L2, T1, T2
4A2	Land converted to Forest	CO ₂	L1, L2, T1, T2

Only temperate forests are occurring in Switzerland. Forest is defined as a minimum area of land of 0.0625 ha with crown cover of at least 20% and a minimum width of 25 m. The minimum height of the dominant trees must be 3 m or have the potential to reach 3 m at maturity in situ (FOEN 2006h). The following forest areas are not subject of the criteria of minimum stand height and minimum crown cover, but must have the potential to achieve it: afforested, regenerated, as well as burned, cut or damaged areas. Although orchards, parks,

camping grounds, open tree formations in settlements, gardens, cemeteries, sports and parking fields may fulfil the (quantitative) forest definition, they were not considered as forests (FOEN 2006h).

According to the Federal Act on Forest, it is one objective to “conserve the forest in its area and spatial distribution” (Swiss Confederation 1991: Art. 1a). Any change of the forested area has to be authorized. Therefore, all forests in Switzerland are considered to be under management.

For reporting purposes, the different forest types were allocated to afforestations (CC11), productive forest (CC12) and unproductive forest (CC13) based on AREA categories (see Table 6-2 and Table 6-6; SFSO 2006a; Didion and Thürig 2013).

Note for afforested areas: The diction afforestation is consistently used for reporting under the UNFCCC, and the diction Afforestation for reporting the activity under the Kyoto Protocol (see chp. 6.1.3.1 and chp. 11.1.3).

A detailed description of the category unproductive forest CC13 can be found in chp. 6.4.2.8.

6.4.2 Methodological issues

6.4.2.1 Choice of method and National Forest Inventories

The calculation approach and the applied conversion time periods for different land-use changes within, from and to forest land and the respective carbon pools are shown in Table 6-3.

Data for growing stock, gross growth (gain in carbon stock of living biomass), cut and mortality (losses in carbon stock of living biomass) were derived from four available Swiss National Forest Inventories (NFIs; see Table 6-12). A description of NFI1 and NFI2 methodologies can be found in EAFV/BFL (1988) and in Brassel and Brändli (1999). Data and methodology of NFI3 are described in Brändli (2010). The inventories NFI1, NFI2 and NFI3 were based on full surveys that were repeated in intervals of approximately 10 years. The fourth inventory (NFI4, 2009–2017) is carried out as a continuous survey where annually a nationally representative subsample of approximately 12% of the Swiss forests is surveyed and evaluated. Otherwise, the methodology remained identical to Brändli (2010).

For assembling the results for the GHG inventory 1990–2015, NFI4 data for the years 2011–2015 were available in addition to the data from NFI1–3. Results of the 5-year subset of the NFI4 were summarised in the data release NFI4 2011–2015 (Thürig et al. 2017; Rigling and Schaffer 2015).

Table 6-12 Characteristics of the National Forest Inventories 1, 2, and 3 and NFI4 2011–2015, accessible forest sample plots without brush forest.

	NFI 1	NFI 2	NFI 3	NFI 4 2011-2015
Inventory cycle	1983-1985	1993-1995	2004-2006	2011-2015
Grid size	1 x 1 km	1.4 x 1.4 km	1.4 x 1.4 km	1.4 x 1.4 km
Terrestrial sample plots	10'981	6'412	6'608	3'403
Measured single trees	128'441	76'394	77'959	39'955

6.4.2.2 Stratification

Spatial strata

Forests in Switzerland reveal a high heterogeneity in terms of elevation, growth conditions, tree species composition, and interannual growth variability. To account for the heterogeneity, the Swiss NFI uses a spatial stratification based on five production regions and three elevation belts (Brändli 2010). To find explanatory variables that significantly reduce the variance of gross growth, an analysis of variance was done (Table 6-13).

Table 6-13 Analysis of variance of gross growth of data from NFI2 and NFI3. Explanatory variables: Tree species, NFI production region, and elevation.

	Gross growth	
	F-value	p-value
Coniferous / Deciduous	421	<0.0001
Production region	45	<0.0001
Elevation	34	<0.0001

The analysis of variance indicated that production region, elevation, and tree species all significantly explain differences in gross growth. Therefore, the explanatory variables considered here were:

- tree species: coniferous and deciduous species
- the five NFI production regions:
 1. Jura, 2. Central Plateau, 3. Pre-Alps, 4. Alps, 5. Southern Alps
- elevation: <601 m, 601–1200 m, >1200 m.

Values for growing stock, gross growth, harvesting and mortality were calculated for each of these 30 strata.

Additional stratification: eastern and western Alps

In the Swiss Alps (NFI production region 4) below an elevation of 1200 m, climate between the eastern and the western part differs substantially. An additional stratification for the eastern and the western part of the Alps below 1200 m (Alps <601 m east, Alps <601 m

west, Alps 601-1200 m east, Alps 601-1200 m west was included; see Thürig et al. 2005a for details). This additional stratification resulted in very small datasets per stratum.

Gains and losses in carbon stock of living biomass were estimated for the eastern and western Alps separately. The emission factors for the Alps below 1200 m were then calculated as a weighted mean of the percentage of forest biomass situated in the western and in the eastern Alps. The ratios for the pooled emission factors correspond to the biomass ratios for the two regions, which were calculated for the three periods between NFI1 and NFI2, NFI2 and NFI3, and NFI3 and NFI4 2011–2015 (Table 6-14).

Table 6-14 Ratio of biomass in the eastern and western Alps (NFI production region 4) for 1985–1994 (derived from NFI1 and NFI2; source: Brassel and Brändli 1999), for 1995–2005 (derived from NFI2 and NFI3 data; source: Brändli 2010) and for 2006–2015 (derived from NFI3 and NFI4 2011–2015 data; source: Thürig et al. 2017). For NFI4 2011–2015, NFI region 4 <601 m and 601–1200 m were aggregated (Thürig et al. 2017).

Elevation [m]	1985 - 1994		1995 - 2005		2006-2015	
	NFI 1-2 Eastern	NFI 1-2 Western	NFI 2-3 Eastern	NFI 2-3 Western	NFI 3-4 Eastern	NFI 3-4 Western
<601	0.56	0.44	0.53	0.47	0.58	0.42
601-1200	0.62	0.38	0.61	0.39	0.58	0.42

Aggregation of strata

Since only the data for the years 2011 to 2015 of the continuously sampled NFI4 (see chp. 6.4.2.1) were used in this submission, several spatial strata were represented by a low number of plots (Thürig et al. 2017). Due to the large variability between sample plots a minimum number of sample plots is needed to obtain reliable estimates of means and sampling errors. Smaller strata were thus merged with neighbouring strata for this submission. For NFI4 2011–2015, the following strata were aggregated and treated as single strata:

- NFI region 2 Central Plateau 601-1200 m and >1200 m:
new stratum NFI region 2 Central Plateau >600 m (276 plots)
- NFI region 3 Pre-Alps ≤600 m and 601-1200 m:
new stratum NFI region 3 Pre-Alps ≤1200 m (400 plots)
- NFI region 4 Alps West ≤600 m and 601-1200 m:
new stratum NFI region 4 Alps West ≤1200 m (142 plots)
- NFI region 4 Alps East ≤600 m and 601-1200 m:
new stratum NFI region 4 Alps East ≤1200 m (162 plots)

6.4.2.3 Estimation of growing stock in biomass

The biomass of all tree compartments (stem-wood over bark including stock, coarse and small branches, needles/leaves, and roots) were estimated based on established allometries to tree-dimensions (Table 6-15; Thürig and Herold 2013). Estimates for branches, foliage and roots were derived from tree diameter at breast height (DBH). For stem-wood over bark

including stock, additionally, diameter at tree height 7 m (D7) and total tree height were required. Except for roots, the biomass functions were empirically derived from a large number of single-tree data from Swiss forest sites (see references in Table 6-15).

Table 6-15 Applied allometric biomass functions, dependencies and references. DBH: tree diameter at breast height; D7: diameter at tree height 7 m.

Tree parts	Input parameter	Nr. of trees	References
Stem-wood over bark incl. stock	DBH, D7, height	12'000	Kaufmann et al. 2001
Coarse branches (≥ 7 cm)	DBH	40'000	Kaufmann et al. 2001
Small branches (< 7 cm)	DBH	40'000	Kaufmann et al. 2001
Needles, Leaves	DBH	400	Perruchoud et al. 1999
Broadleaved Roots	DBH	443	Wutzler et al. 2008
Coniferous Roots	DBH	80	Zell and Thürig 2013

The biomass of all individual trees was calculated and, in a second step, single-tree estimates of gains and losses were obtained as the difference in tree biomass between subsequent NFIs (Thürig and Herold 2013).

6.4.2.4 Carbon content

A mean carbon content of 50% was used to convert the volume of alive trees to biomass. The carbon content estimate represents an approximation which was based on carbon fractions for coniferous and broadleaved trees in temperate forests provided in Tab. 4.3 in Volume 4 of IPCC (2006), and on the fact that in Switzerland coniferous trees are more abundant than broadleaved trees (see Table 051 in Brändli 2010).

6.4.2.5 Productive forests (CC12): growing stock, gains and losses of living biomass

Values for growing stock, gains (gross growth) and losses (cut and mortality) of living biomass for productive forests (CC12, without afforestations) were derived from 5'456 common sample plots measured during NFI1 and NFI2 (Kaufmann 2001), 5'581 samples measured during NFI2 and NFI3 (Brändli 2010) and 3'280 samples measured during NFI3 and NFI4 2011–2015 (Thürig et al. 2017). All values derived from the national forest inventories refer to above- and below-ground biomass in mass units (t C ha^{-1}) per spatial stratum.

Annual gain of living biomass – gross growth

Annual values of gross growth were derived from the NFI1 and NFI2 datasets for the period 1985–1994, from the NFI2 and NFI3 datasets for the period 1995–2005 and from the NFI3 and NFI4 2011–2015 datasets for the period 2006–2015. Annual values of gross growth were assumed to remain constant in the intersurvey periods of NFI1 to NFI2, NFI2 to NFI3 and of NFI3 to NFI4 2011–2015, respectively (Table 6-17).

Annual loss of living biomass – cut and mortality

An average value for cut and mortality (CM) was derived from the NFI1 and NFI2 dataset for the period 1985–1994, from the NFI2 and NFI3 datasets for the period 1995–2005 and from the NFI3 and NFI4 2011–2015 datasets for the period 2006–2015. To calculate annual values of cut and mortality (CM_y) for the years 1985 to 1994, 1995 to 2005 and 2006 to 2015, respectively, the average amount of cut and mortality from NFI was weighted by the percentage of the relative harvesting amounts taken from the forest statistics (Table 6-16; FOEN 2016k and former editions; Swiss Federal Statistical Office: Wood production in Switzerland 1975–2015, <https://www.pxweb.bfs.admin.ch/>). These relative harvesting amounts, used as weighting factors, were calculated for each year per NFI intersurvey period.

Data from the forest statistics (Table 6-16) show that harvesting rates were extraordinary high in 1990 after storm Vivian (February 1990) and in 2000 after the storm Lothar (December 1999). Harvesting rates in Swiss forests tended to increase since 1991. In 2008 harvesting rates started to decline due to the international and domestic economic framework conditions.

Table 6-16 Annual harvesting amount in m³ merchantable timber specified for five NFI production region as well as for coniferous and deciduous tree species for the period 1990–2015 (FOEN 2016k and former editions; <https://www.pxweb.bfs.admin.ch/>).

Year	1. Jura		2. Central plateau		3. Pre-Alps		4. Alps		5. Southern Alps		Total [m ³]
	Conif. [m ³]	Dec. [m ³]									
1990	687'327	358'647	1'769'813	606'718	1'285'639	138'126	1'301'313	70'064	21'575	22'456	6'261'678
1991	476'956	354'002	1'017'232	489'742	877'851	133'155	1'064'650	72'229	24'356	26'736	4'536'909
1992	555'523	372'249	1'199'596	571'610	735'680	128'934	736'230	70'706	47'388	28'637	4'446'553
1993	550'536	373'298	1'206'294	562'232	723'565	132'676	649'938	63'940	42'511	32'785	4'337'775
1994	621'726	392'967	1'270'296	530'906	798'449	136'103	717'840	66'896	40'986	33'746	4'609'915
1995	650'572	407'119	1'388'932	570'552	774'040	154'108	590'859	56'714	51'643	33'869	4'678'408
1996	520'335	381'365	1'066'770	567'769	654'554	151'164	506'107	59'674	48'288	38'889	3'994'915
1997	599'981	394'846	1'176'333	576'415	742'830	153'719	574'152	63'650	61'043	40'189	4'383'158
1998	604'703	422'216	1'330'973	627'633	836'806	164'348	657'409	108'848	50'626	41'485	4'845'047
1999	602'652	398'648	1'342'905	639'150	824'142	173'845	593'844	68'786	44'556	39'181	4'727'709
2000	994'262	387'183	3'916'680	934'372	2'241'486	213'858	436'743	57'105	21'236	35'049	9'237'974
2001	443'612	338'751	2'020'561	594'616	1'477'489	157'710	510'730	60'152	22'237	35'722	5'661'580
2002	442'519	329'480	1'406'758	493'905	1'090'875	134'603	528'144	63'303	31'236	35'794	4'556'617
2003	557'454	315'096	1'669'605	518'273	1'195'090	142'055	588'062	62'739	37'111	35'486	5'120'971
2004	655'757	305'681	1'774'841	515'877	1'119'243	164'745	488'722	70'090	29'995	35'571	5'160'522
2005	653'049	359'808	1'810'839	614'845	1'010'979	180'546	514'905	70'603	35'462	33'614	5'284'650
2006	735'256	405'850	1'779'973	687'428	1'116'868	229'781	569'673	84'656	43'443	48'599	5'701'527
2007	793'459	425'790	1'587'494	699'076	1'144'370	230'284	621'234	82'414	62'799	43'638	5'690'558
2008	705'815	459'994	1'281'782	727'581	1'018'497	224'634	664'086	82'623	53'064	44'123	5'262'199
2009	598'292	461'055	1'149'202	701'188	878'565	224'490	678'212	90'001	56'375	42'316	4'879'696
2010	647'176	494'739	1'090'994	722'644	992'435	248'151	720'659	99'773	60'391	52'037	5'128'999
2011	617'887	513'720	1'061'986	741'587	983'040	253'300	686'797	101'644	61'822	53'305	5'075'088
2012	566'782	488'626	970'748	719'003	825'019	225'988	665'506	94'480	51'475	50'757	4'658'384
2013	576'744	521'122	948'706	739'180	834'166	254'726	670'170	117'841	64'745	50'928	4'778'328
2014	619'002	539'721	945'695	777'852	863'150	259'888	654'300	110'816	95'192	47'603	4'913'219
2015	528'202	505'431	916'020	766'645	753'783	244'149	625'555	96'230	62'233	53'649	4'551'897

Growing stock: calculation of time series

In order to develop a consistent time series, annual growing stocks of living biomass (stockC_i) were calculated per spatial strata (i) for productive forests (CC12) backward or forward starting from the growing stock 2005, determined from NF13 (abbreviations are explained in chp. 6.1.3.2):

$$\text{stockC}_{i,i,12,iy} = \text{stockC}_{i,i,12,2005} - \sum_{n=2005}^{iy} [\text{gainC}_{i,i,12,n}] + \sum_{n=2005}^{iy} [\text{lossC}_{i,i,12,n}] \quad \text{for } iy < 2005$$

$$\text{stockC}_{i,i,12,iy} = \text{stockC}_{i,i,12,2005} \quad \text{for } iy = 2005$$

$$\text{stockC}_{i,i,12,iy} = \text{stockC}_{i,i,12,2005} + \sum_{n=2006}^{iy} [\text{gainC}_{i,i,12,n}] - \sum_{n=2006}^{iy} [\text{lossC}_{i,i,12,n}] \quad \text{for } iy > 2005$$

where “iy” indicates the inventory year (here: 1985-2015), “n” the years between 2005 and the inventory year iy.

The backward calculation was used for the time period 1985–2004 (iy <2005), where the annual growing stock equals the growing stock 2005 minus the net change based on the gains due to the annual gross growth (gainC_{i,i,12,iy}) and the losses due to the annual amounts of cut and mortality (lossC_{i,i,12,iy}).

The forward calculation was used for the time period after 2005 (iy >2005), where the annual growing stock equals the growing stock 2005 plus the net change based on the gains due to the annual gross growth (gainC_{i,i,12,iy}) and the losses due the annual amounts of cut and mortality (lossC_{i,i,12,iy}).

Annual values of gross growth (gains in carbon stock of living biomass), cut and mortality (losses in carbon stock of living biomass) and calculated growing stocks (carbon stocks in living biomass) for the period 1990 to 2015 specified for all spatial strata are displayed in Table 6-17.

All working steps and data required to reproduce the calculation of emission factors for productive forests (CC12) in the period 1990–2015 are summarized in FOEN (2017b).

Table 6-17 Annual carbon data of living biomass for productive forest (CC12) disaggregated for NFI region (NFI) and elevation zone (Elev.), 1990–2015, for stocks (stockCl), gains (gross growth, gainCl) and losses (cut and mortality, lossCl). Highlighted data for 1990 are displayed in Table 6-4.

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹]									
1	1	126.76	127.98	129.46	130.77	132.09	133.23	134.28	133.76	132.96	131.95
1	2	124.81	125.74	127.12	128.29	129.47	130.45	131.34	132.06	132.55	132.96
1	3	84.75	85.34	86.31	87.13	87.96	88.65	89.29	90.06	90.71	91.33
2	1	133.95	133.81	135.28	136.19	137.13	138.05	138.65	139.44	140.01	140.13
2	2	146.61	146.63	148.28	149.41	150.55	151.66	152.45	153.52	154.34	154.73
2	3	101.51	102.05	103.03	103.89	104.75	105.58	106.34	106.76	107.11	107.35
3	1	135.29	136.50	138.24	140.19	142.11	143.89	145.45	147.60	149.61	151.41
3	2	147.51	147.87	149.24	150.97	152.71	154.25	155.76	157.31	158.59	159.55
3	3	119.40	119.13	119.71	120.60	121.51	122.26	123.05	124.14	125.04	125.75
4	1	94.68	94.73	94.88	95.29	95.99	96.54	97.67	99.62	101.43	102.44
4	2	104.44	104.35	104.63	105.46	106.47	107.36	108.43	109.50	110.42	110.82
4	3	96.36	95.70	95.48	95.87	96.43	96.86	97.53	98.28	98.88	99.26
5	1	71.66	73.49	75.13	76.69	78.08	79.43	80.77	81.37	81.93	82.44
5	2	76.87	78.46	79.93	81.29	82.57	83.82	85.05	86.55	88.01	89.47
5	3	76.48	77.78	79.04	80.08	81.15	82.22	83.20	84.41	85.49	86.67

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹]									
1	1	131.12	129.46	129.41	129.44	129.30	128.99	129.11	128.99	128.92	129.10
1	2	133.44	133.04	134.06	135.11	135.94	136.57	136.55	136.30	136.24	136.52
1	3	91.97	92.07	92.98	93.89	94.66	95.30	96.47	97.56	98.72	99.99
2	1	140.20	134.70	133.69	134.12	133.99	133.68	131.28	129.32	128.03	127.17
2	2	155.06	149.42	148.47	149.00	148.95	148.67	145.61	144.42	144.02	144.04
2	3	107.58	106.09	105.90	106.14	106.20	106.19	145.61	144.42	144.02	144.04
3	1	153.15	152.66	153.65	155.34	156.83	158.25	156.43	155.72	155.45	155.64
3	2	160.50	157.22	156.38	156.77	156.82	156.97	156.43	155.72	155.45	155.64
3	3	126.48	124.26	123.64	123.83	123.80	123.92	124.99	126.03	127.23	128.64
4	1	104.16	106.22	108.16	110.03	111.84	113.65	117.18	117.61	117.94	118.18
4	2	111.66	112.89	113.95	114.96	115.85	116.88	117.18	117.61	117.94	118.18
4	3	99.81	100.71	101.45	102.14	102.71	103.49	104.66	105.74	106.75	107.73
5	1	83.03	83.78	84.51	85.23	85.97	86.70	87.24	87.93	88.62	89.36
5	2	90.97	92.59	94.19	95.77	97.34	98.93	100.17	101.37	102.61	103.86
5	3	87.91	89.41	90.89	92.28	93.61	95.02	96.32	97.48	98.72	99.94

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹]									
1	1	129.02	128.91	129.04	129.00	128.77	128.92				
1	2	136.55	136.63	136.94	137.14	137.15	137.53				
1	3	101.19	102.41	103.70	104.95	106.16	107.47				
2	1	126.38	125.60	125.12	124.63	124.02	123.51				
2	2	144.16	144.33	144.79	145.27	145.67	146.17				
2	3	144.16	144.33	144.79	145.27	145.67	146.17				
3	1	155.34	155.05	155.41	155.60	155.68	156.19				
3	2	155.34	155.05	155.41	155.60	155.68	156.19				
3	3	129.87	131.12	132.60	134.06	135.47	137.04				
4	1	118.23	118.34	118.56	118.57	118.67	118.96				
4	2	118.23	118.34	118.56	118.57	118.67	118.96				
4	3	108.64	109.61	110.61	111.59	112.61	113.67				
5	1	89.73	90.05	90.49	90.89	91.33	91.63				
5	2	104.95	106.02	107.18	108.26	109.20	110.26				
5	3	101.09	102.22	103.45	104.56	105.43	106.56				

(Table 6-17 continued)

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	3.60	3.60	3.60	3.60	3.60	3.60	3.37	3.37	3.37	3.37
1	2	3.21	3.21	3.21	3.21	3.21	3.21	3.04	3.04	3.04	3.04
1	3	1.95	1.95	1.95	1.95	1.95	1.95	1.80	1.80	1.80	1.80
2	1	4.63	4.63	4.63	4.63	4.63	4.63	4.54	4.54	4.54	4.54
2	2	4.63	4.63	4.63	4.63	4.63	4.63	4.56	4.56	4.56	4.56
2	3	1.60	1.60	1.60	1.60	1.60	1.60	1.28	1.28	1.28	1.28
3	1	4.56	4.56	4.56	4.56	4.56	4.56	4.23	4.23	4.23	4.23
3	2	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15
3	3	2.48	2.48	2.48	2.48	2.48	2.48	2.50	2.50	2.50	2.50
4	1	3.24	3.24	3.24	3.24	3.24	3.24	3.44	3.44	3.44	3.44
4	2	2.49	2.49	2.49	2.49	2.49	2.49	2.50	2.50	2.50	2.50
4	3	1.81	1.81	1.81	1.81	1.81	1.81	1.90	1.90	1.90	1.90
5	1	2.74	2.74	2.74	2.74	2.74	2.74	2.04	2.04	2.04	2.04
5	2	2.20	2.20	2.20	2.20	2.20	2.20	2.18	2.18	2.18	2.18
5	3	1.61	1.61	1.61	1.61	1.61	1.61	1.79	1.79	1.79	1.79

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	3.37	3.37	3.37	3.37	3.37	3.37	3.77	3.77	3.77	3.77
1	2	3.04	3.04	3.04	3.04	3.04	3.04	3.37	3.37	3.37	3.37
1	3	1.80	1.80	1.80	1.80	1.80	1.80	2.11	2.11	2.11	2.11
2	1	4.54	4.54	4.54	4.54	4.54	4.54	4.52	4.52	4.52	4.52
2	2	4.56	4.56	4.56	4.56	4.56	4.56	4.89	4.89	4.89	4.89
2	3	1.28	1.28	1.28	1.28	1.28	1.28	4.89	4.89	4.89	4.89
3	1	4.23	4.23	4.23	4.23	4.23	4.23	4.11	4.11	4.11	4.11
3	2	4.15	4.15	4.15	4.15	4.15	4.15	4.11	4.11	4.11	4.11
3	3	2.50	2.50	2.50	2.50	2.50	2.50	2.67	2.67	2.67	2.67
4	1	3.44	3.44	3.44	3.44	3.44	3.44	2.29	2.52	2.52	2.52
4	2	2.50	2.50	2.50	2.50	2.50	2.50	2.29	2.52	2.52	2.52
4	3	1.90	1.90	1.90	1.90	1.90	1.90	2.16	2.16	2.16	2.16
5	1	2.04	2.04	2.04	2.04	2.04	2.04	2.52	2.52	2.52	2.52
5	2	2.18	2.18	2.18	2.18	2.18	2.18	2.17	2.17	2.17	2.17
5	3	1.79	1.79	1.79	1.79	1.79	1.79	1.85	1.85	1.85	1.85

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	3.77	3.77	3.77	3.77	3.77	3.77				
1	2	3.37	3.37	3.37	3.37	3.37	3.37				
1	3	2.11	2.11	2.11	2.11	2.11	2.11				
2	1	4.52	4.52	4.52	4.52	4.52	4.52				
2	2	4.89	4.89	4.89	4.89	4.89	4.89				
2	3	4.89	4.89	4.89	4.89	4.89	4.89				
3	1	4.11	4.11	4.11	4.11	4.11	4.11				
3	2	4.11	4.11	4.11	4.11	4.11	4.11				
3	3	2.67	2.67	2.67	2.67	2.67	2.67				
4	1	2.52	2.52	2.52	2.52	2.52	2.52				
4	2	2.52	2.52	2.52	2.52	2.52	2.52				
4	3	2.16	2.16	2.16	2.16	2.16	2.16				
5	1	2.52	2.52	2.52	2.52	2.52	2.52				
5	2	2.17	2.17	2.17	2.17	2.17	2.17				
5	3	1.85	1.85	1.85	1.85	1.85	1.85				

(Table 6-17 continued)

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	-2.38	-2.12	-2.29	-2.29	-2.46	-2.55	-3.89	-4.17	-4.37	-4.20
1	2	-2.28	-1.83	-2.04	-2.03	-2.23	-2.32	-2.32	-2.54	-2.63	-2.56
1	3	-1.36	-0.99	-1.13	-1.12	-1.26	-1.32	-1.03	-1.15	-1.18	-1.16
2	1	-4.77	-3.17	-3.72	-3.70	-3.71	-4.03	-3.75	-3.97	-4.41	-4.47
2	2	-4.61	-2.98	-3.50	-3.49	-3.53	-3.84	-3.50	-3.74	-4.18	-4.23
2	3	-1.05	-0.62	-0.74	-0.74	-0.77	-0.84	-0.86	-0.93	-1.04	-1.05
3	1	-3.35	-2.82	-2.60	-2.64	-2.77	-2.99	-2.09	-2.22	-2.44	-2.49
3	2	-3.78	-2.78	-2.42	-2.41	-2.60	-2.63	-2.61	-2.87	-3.19	-3.20
3	3	-2.75	-1.90	-1.60	-1.57	-1.73	-1.69	-1.41	-1.59	-1.79	-1.77
4	1	-3.19	-3.10	-2.83	-2.55	-2.69	-2.27	-1.49	-1.62	-2.42	-1.72
4	2	-2.59	-2.21	-1.66	-1.48	-1.61	-1.33	-1.43	-1.59	-2.10	-1.67
4	3	-2.47	-2.03	-1.41	-1.25	-1.38	-1.14	-1.15	-1.30	-1.51	-1.35
5	1	-0.92	-1.09	-1.18	-1.35	-1.38	-1.40	-1.44	-1.49	-1.54	-1.45
5	2	-0.61	-0.72	-0.84	-0.92	-0.94	-0.98	-0.67	-0.72	-0.72	-0.67
5	3	-0.30	-0.35	-0.57	-0.54	-0.53	-0.63	-0.58	-0.71	-0.61	-0.54

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	-5.03	-3.41	-3.35	-3.51	-3.67	-4.05	-3.65	-3.88	-3.83	-3.58
1	2	-3.43	-2.02	-1.99	-2.21	-2.41	-2.56	-3.38	-3.62	-3.43	-3.09
1	3	-1.70	-0.89	-0.88	-1.03	-1.16	-1.20	-0.95	-1.01	-0.95	-0.84
2	1	-10.04	-5.54	-4.12	-4.67	-4.85	-5.23	-6.92	-6.47	-5.80	-5.38
2	2	-10.20	-5.52	-4.03	-4.62	-4.83	-5.14	-6.57	-6.07	-5.30	-4.87
2	3	-2.77	-1.47	-1.05	-1.22	-1.29	-1.34	-6.57	-6.07	-5.30	-4.87
3	1	-4.72	-3.24	-2.54	-2.74	-2.82	-2.80	-4.73	-4.82	-4.39	-3.93
3	2	-7.44	-4.99	-3.77	-4.10	-4.00	-3.77	-4.73	-4.82	-4.39	-3.93
3	3	-4.72	-3.12	-2.31	-2.53	-2.38	-2.16	-1.60	-1.64	-1.46	-1.27
4	1	-1.37	-1.50	-1.57	-1.62	-1.63	-1.67	-2.00	-2.09	-2.19	-2.29
4	2	-1.28	-1.44	-1.50	-1.61	-1.48	-1.53	-2.00	-2.09	-2.19	-2.29
4	3	-1.00	-1.16	-1.20	-1.33	-1.12	-1.18	-0.99	-1.08	-1.15	-1.17
5	1	-1.29	-1.32	-1.32	-1.31	-1.31	-1.24	-1.97	-1.83	-1.83	-1.77
5	2	-0.56	-0.58	-0.59	-0.60	-0.59	-0.57	-0.92	-0.97	-0.92	-0.92
5	3	-0.29	-0.31	-0.40	-0.46	-0.38	-0.44	-0.54	-0.69	-0.61	-0.63

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	-3.85	-3.87	-3.63	-3.81	-4.00	-3.62				
1	2	-3.33	-3.28	-3.06	-3.17	-3.35	-2.98				
1	3	-0.91	-0.89	-0.83	-0.85	-0.91	-0.80				
2	1	-5.30	-5.30	-4.99	-5.00	-5.13	-5.02				
2	2	-4.76	-4.73	-4.42	-4.41	-4.50	-4.39				
2	3	-4.76	-4.73	-4.42	-4.41	-4.50	-4.39				
3	1	-4.41	-4.40	-3.76	-3.92	-4.04	-3.61				
3	2	-4.41	-4.40	-3.76	-3.92	-4.04	-3.61				
3	3	-1.43	-1.42	-1.19	-1.21	-1.25	-1.10				
4	1	-2.47	-2.41	-2.30	-2.52	-2.42	-2.23				
4	2	-2.47	-2.41	-2.30	-2.52	-2.42	-2.23				
4	3	-1.25	-1.19	-1.16	-1.17	-1.14	-1.09				
5	1	-2.15	-2.20	-2.07	-2.12	-2.07	-2.21				
5	2	-1.07	-1.10	-1.00	-1.08	-1.22	-1.11				
5	3	-0.70	-0.72	-0.62	-0.73	-0.98	-0.72				

6.4.2.6 Productive forests (CC12): carbon stocks in dead wood, litter and in mineral soils

Carbon stocks in dead wood

Carbon stock in dead wood depends on the available volume in different decay stages, and on the associated wood density and carbon content. The influence of wood decay on wood density and on carbon content of dead wood was investigated by Dobbertin and Jüngling (2009). For the two dominant tree species in Swiss forests, Norway spruce (*Picea abies*) and European beech (*Fagus sylvatica*), a significant decrease in wood density from alive trees to wood in advanced decay (duff wood) was found, i.e., Norway spruce (0.39 to 0.247 g cm⁻³) and beech (0.56 to 0.233 g cm⁻³; cf. Tab. 1 in Didion et al. 2014a). Carbon content varied very little (± 1.2 to 1.4%) between alive trees and dead wood in different decay stages. The findings by Dobbertin and Jüngling (2009) are consistent with recent data from Switzerland for Norway spruce (Wunder and Bont 2015).

The total amount of carbon in the dead wood pool (henceforth dead wood) in Switzerland is estimated as the sum of carbon in

- stemwood of standing dead trees ≥ 12 cm DBH
- lying dead trees ≥ 7 cm DBH
- branchwood ≥ 7 cm in diameter
- coarse roots > ca. 5 mm in diameter of dead trees ≥ 12 cm DBH.

A time series of carbon stocks in dead wood was simulated with the soil carbon model Yasso07 (Didion and Thürig 2016; see description in chp. 6.4.2.7). Stratified estimated dead wood stocks for 1990 are shown in Table 6-18. Annual values for dead wood stocks since 1990 are displayed in Table 6-19.

Carbon stocks in litter (organic soil horizons) and in mineral soils

Soil carbon stocks were estimated by Nussbaum et al. (2012, 2014) based on soil profiles and robust geostatistical methods. The soil profiles are part of a database maintained at the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL). The data were collected over the past 30 years (N=1033 sites) distributed among different forest types throughout Switzerland. The same dataset was re-analysed (N=1030; Nussbaum et al. 2012) to update estimates of carbon stocks in litter (organic soil horizons L - litter, F - fermentation and H - humus) prepared by Moeri (2007).

The data for litter and soil carbon stocks were stratified by the five NFI production regions and three elevation levels (Table 6-18). The estimated national average carbon stocks in mineral forest soils are 79.9 t C ha⁻¹ (0–30 cm topsoil) and 125.8 t C ha⁻¹ (0–100 cm), respectively (Nussbaum et al. 2014), and 16.7 t C ha⁻¹ in the organic soil horizons (litter) of mineral forest soils (Table 3 in Nussbaum et al 2012).

The sites in the WSL soil database which were used by Nussbaum et al. (2012, 2014) and by Moeri (2007) were visited mostly between 1990 and 2005. Hence, it is not possible to attribute the national estimates of carbon stocks in mineral forest soils and in litter to one single year. Consequently, a combination of these carbon stocks and the carbon stock

changes derived from the Yasso07 model (Didion and Thürig 2016; see chp. 6.4.2.7) would not result in a consistent time series for litter and soilcarbon stocks. Thus, it was assumed that the values from Nussbaum et al. (2012, 2014) are representative for the period 1990 until the inventory year.

Table 6-18 Dead wood stocks (stockC_d) in Swiss productive forests (CC12) by spatial stratum in t C ha⁻¹ for 1990 (Tables A-22 (means) and A-24 (SE) in Didion and Thürig 2016). Carbon stocks in organic soil horizons (litter; stockC_h; Table 3 in Nussbaum et al. 2012; used for CC12, CC13) and carbon stocks in mineral soil (0–30 cm; stockC_s; Table 5 in Nussbaum et al. 2012, Nussbaum et al. 2014; used for CC11, CC12, CC13) were assumed to be representative for 1990–2015. The data were stratified by five NFI production regions and three elevation zones. Dead wood and litter stocks in NFI region 2, 601-1200 and >1200 m were aggregated due to the low number of samples >1200 m. Average values ± single standard errors are given.

NFI region	Elevation [m]	Carbon stock in dead wood 1990 (stockC _{d,i,12}) [t C ha ⁻¹]	Carbon stock in litter (stockC _{h,i,12} , stockC _{h,i,13}) [t C ha ⁻¹]	Carbon stock in mineral topsoil 0-30 cm (stockC _{s,i,11} , stockC _{s,i,12} , stockC _{s,i,13}) [t C ha ⁻¹]
1	<601	5.84 ± 0.09	9.51 ± 1.57	82.65 ± 3.34
1	601-1200	5.67 ± 0.15	7.53 ± 0.70	102.03 ± 3.56
1	>1200	6.02 ± 0.06	7.76 ± 1.74	121.34 ± 5.39
2	<601	9.14 ± 0.09	8.70 ± 0.68	55.40 ± 1.55
2	601-1200	8.8 ± 0.1	11.42 ± 1.45	62.12 ± 1.68
2	>1200	8.8 ± 0.1	11.42 ± 1.45	122.00 ± 7.07
3	<601	9.52 ± 0.32	7.51 ± 1.25	66.10 ± 2.06
3	601-1200	8.46 ± 0.13	16.29 ± 1.55	57.91 ± 2.00
3	>1200	9.3 ± 0.09	26.21 ± 4.77	95.78 ± 3.27
4	<601	6.88 ± 0.32	3.15 ± 0.47	66.47 ± 2.44
4	601-1200	7.8 ± 0.07	19.99 ± 2.64	74.39 ± 2.42
4	>1200	8.15 ± 0.09	33.37 ± 3.53	69.48 ± 1.85
5	<601	2.34 ± 0.05	8.22 ± 1.62	102.37 ± 4.07
5	601-1200	3.06 ± 0.05	11.03 ± 2.11	108.99 ± 4.09
5	>1200	3.28 ± 0.08	30.77 ± 5.43	107.08 ± 4.11
Switzerland		7.44 ± 0.03	16.73 ± 0.83	79.93 ± 1.52

Table 6-19 Carbon stock in dead wood for CC12, 1990–2015. Highlighted data for 1990 are displayed in Table 6-4 and Table 6-18.

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: carbon stock in dead wood (stockCd,i) [t C ha ⁻¹]									
1	1	5.84	5.88	5.89	5.91	5.89	5.89	6.13	6.33	6.48	6.61
1	2	5.67	5.55	5.42	5.31	5.18	5.09	5.06	5.01	4.97	4.92
1	3	6.02	6.02	5.99	5.97	5.93	5.91	6.00	6.06	6.12	6.16
2	1	9.14	9.17	9.15	9.13	9.07	9.05	9.42	9.73	9.99	10.19
2	2	8.80	8.80	8.76	8.73	8.66	8.63	9.01	9.33	9.60	9.84
2	3	8.80	8.80	8.76	8.73	8.66	8.63	9.01	9.33	9.60	9.84
3	1	9.52	9.56	9.57	9.57	9.53	9.54	9.70	9.82	9.91	9.99
3	2	8.46	8.43	8.37	8.33	8.26	8.23	8.28	8.27	8.29	8.31
3	3	9.30	9.32	9.31	9.31	9.26	9.26	9.73	10.10	10.46	10.78
4	1	6.88	6.92	6.94	6.96	6.94	6.96	7.12	7.26	7.40	7.51
4	2	7.80	7.82	7.81	7.81	7.79	7.80	7.76	7.69	7.64	7.59
4	3	8.15	8.12	8.06	8.02	7.95	7.92	7.93	7.91	7.89	7.87
5	1	2.34	2.25	2.17	2.09	2.01	1.95	1.91	1.86	1.82	1.79
5	2	3.06	3.06	3.05	3.04	3.03	3.03	3.01	2.98	2.95	2.93
5	3	3.28	3.35	3.42	3.48	3.52	3.58	3.33	3.10	2.90	2.73

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: carbon stock in dead wood (stockCd,i) [t C ha ⁻¹]									
1	1	6.72	6.83	6.91	7.07	7.16	7.26	7.43	7.56	7.70	7.84
1	2	4.87	4.84	4.79	4.76	4.73	4.71	4.60	4.48	4.39	4.31
1	3	6.19	6.22	6.24	6.30	6.33	6.38	6.60	6.77	6.96	7.14
2	1	10.38	10.54	10.68	10.92	11.06	11.23	11.43	11.56	11.71	11.88
2	2	10.04	10.23	10.39	10.61	10.77	10.95	11.06	11.11	11.19	11.29
2	3	10.04	10.23	10.39	10.61	10.77	10.95	11.06	11.11	11.19	11.29
3	1	10.05	10.11	10.14	10.24	10.30	10.37	10.51	10.57	10.65	10.73
3	2	8.31	8.33	8.31	8.31	8.33	8.36	8.12	7.89	7.70	7.52
3	3	11.06	11.33	11.54	11.79	12.01	12.24	12.43	12.57	12.74	12.90
4	1	7.62	7.72	7.79	7.90	8.00	8.11	8.36	8.57	8.78	8.99
4	2	7.52	7.48	7.41	7.38	7.36	7.36	7.40	7.42	7.45	7.50
4	3	7.84	7.83	7.78	7.78	7.78	7.80	8.03	8.21	8.39	8.57
5	1	1.75	1.72	1.69	1.67	1.65	1.64	2.02	2.36	2.66	2.93
5	2	2.90	2.89	2.86	2.85	2.84	2.85	2.95	3.04	3.13	3.21
5	3	2.56	2.42	2.28	2.17	2.06	1.96	2.07	2.15	2.23	2.31

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: carbon stock in dead wood (stockCd,i) [t C ha ⁻¹]									
1	1	8.00	8.10	8.20	8.31	8.35	8.45				
1	2	4.25	4.16	4.09	4.03	3.95	3.89				
1	3	7.33	7.45	7.58	7.71	7.78	7.89				
2	1	12.06	12.18	12.30	12.43	12.46	12.58				
2	2	11.42	11.48	11.54	11.63	11.62	11.68				
2	3	11.42	11.48	11.54	11.63	11.62	11.68				
3	1	10.83	10.86	10.92	10.99	10.96	10.99				
3	2	7.40	7.21	7.08	6.96	6.81	6.68				
3	3	13.10	13.18	13.31	13.45	13.48	13.56				
4	1	9.22	9.37	9.53	9.69	9.78	9.90				
4	2	7.56	7.57	7.59	7.62	7.63	7.64				
4	3	8.77	8.88	9.01	9.15	9.23	9.32				
5	1	3.19	3.41	3.61	3.80	3.96	4.11				
5	2	3.30	3.37	3.43	3.50	3.55	3.60				
5	3	2.39	2.45	2.50	2.56	2.61	2.65				

6.4.2.7 Productive forests (CC12): changes in carbon stocks in dead wood, in litter and in mineral soils

Switzerland used the soil carbon model Yasso07 to estimate temporal changes in carbon stocks in mineral forest soil (0–100 cm), organic soil horizons (LFH; litter) and in dead wood for productive forests (CC12). The implementation of Yasso07 (Tuomi et al. 2009, 2011) in the Swiss GHG inventory is described in detail in Didion et al. (2012) and Didion and Thürig (2016). Didion et al. (2014a) demonstrated the validity of the model for application in Swiss forests.

Yasso07 is a model of carbon cycling in mineral soil, litter and dead wood. For estimating stocks of organic carbon in mineral soil up to a depth of ca. 100 cm and the temporal dynamics of the carbon stocks, Yasso07 requires information on carbon inputs from dead organic matter (i.e. non-woody inputs, including foliage and fine roots, woody inputs, including standing and lying dead wood and dead roots) and climate (temperature, temperature amplitude and precipitation).

By default, Yasso07 does not provide separate estimates of carbon pool sizes for dead wood, litter and soil. In order to report estimates for each pool, the structure of Yasso07 was examined for deriving separate estimates. Dead wood, litter and soil pools could be correlated with modeled data based on the category of carbon input, i.e., non-woody and woody material, and the five carbon compartments in Yasso07, i.e. four chemical partitions (insoluble, soluble in ethanol, soluble in water or in acid and humus). The approach was validated using independent, measured data (see Didion et al. 2012).

Using annual data for climate and for carbon inputs obtained from the Swiss NFIs, Yasso07 was used for estimating the annual carbon stock changes in mineral soil, litter and dead wood. For an overview Table 6-18 shows the carbon stocks in Swiss forests by stratum for the pools mineral soil, litter (based on Nussbaum et al 2012, 2014; see 6.4.2.6) and dead wood (based on Didion and Thürig 2016; see 6.4.2.6). Annual stratified values of carbon stock changes for dead wood, litter and mineral soils can be found in Table 6-20. Carbon stocks and carbon stock changes were validated as described in Didion and Thürig (2016).

Table 6-20 Net carbon stock change in dead wood, in litter and in mineral soils for productive forest (CC12), 1990–2015. Highlighted data for 1990 are displayed in Table 6-4. Positive values refer to gains in carbon stock, negative values refer to losses in carbon stock.

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: net change in dead wood (changeCd,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.00	0.04	0.01	0.01	-0.02	0.00	0.24	0.20	0.16	0.13
1	2	-0.15	-0.12	-0.13	-0.11	-0.13	-0.09	-0.03	-0.05	-0.04	-0.04
1	3	-0.04	0.00	-0.03	-0.02	-0.05	-0.02	0.09	0.06	0.06	0.04
2	1	-0.04	0.02	-0.02	-0.02	-0.06	-0.02	0.36	0.31	0.26	0.21
2	2	-0.06	0.00	-0.03	-0.03	-0.07	-0.03	0.38	0.32	0.28	0.23
2	3	-0.06	0.00	-0.03	-0.03	-0.07	-0.03	0.38	0.32	0.28	0.23
3	1	0.00	0.04	0.01	0.00	-0.04	0.01	0.16	0.12	0.09	0.08
3	2	-0.07	-0.03	-0.06	-0.04	-0.07	-0.03	0.05	0.00	0.02	0.02
3	3	-0.02	0.02	-0.01	0.00	-0.05	0.00	0.46	0.38	0.36	0.32
4	1	0.02	0.05	0.02	0.02	-0.01	0.01	0.17	0.14	0.14	0.11
4	2	0.00	0.02	-0.01	0.01	-0.02	0.01	-0.04	-0.07	-0.05	-0.06
4	3	-0.06	-0.03	-0.06	-0.04	-0.07	-0.03	0.01	-0.03	-0.01	-0.02
5	1	-0.10	-0.09	-0.09	-0.08	-0.08	-0.06	-0.04	-0.04	-0.04	-0.04
5	2	-0.01	0.00	-0.01	0.00	-0.01	0.00	-0.02	-0.03	-0.02	-0.02
5	3	0.08	0.08	0.06	0.06	0.05	0.06	-0.24	-0.23	-0.20	-0.18

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: net change in dead wood (changeCd,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.12	0.10	0.08	0.16	0.09	0.10	0.17	0.12	0.14	0.14
1	2	-0.05	-0.04	-0.05	-0.03	-0.03	-0.01	-0.12	-0.11	-0.09	-0.08
1	3	0.03	0.03	0.02	0.06	0.03	0.05	0.22	0.18	0.18	0.18
2	1	0.18	0.17	0.14	0.23	0.15	0.17	0.20	0.13	0.15	0.16
2	2	0.20	0.19	0.16	0.22	0.16	0.18	0.10	0.05	0.08	0.10
2	3	0.20	0.19	0.16	0.22	0.16	0.18	0.10	0.05	0.08	0.10
3	1	0.05	0.06	0.03	0.10	0.06	0.07	0.14	0.06	0.08	0.08
3	2	0.00	0.01	-0.01	0.00	0.02	0.03	-0.24	-0.23	-0.19	-0.17
3	3	0.27	0.27	0.22	0.24	0.22	0.23	0.19	0.14	0.16	0.16
4	1	0.11	0.10	0.07	0.11	0.09	0.11	0.26	0.20	0.21	0.21
4	2	-0.07	-0.04	-0.06	-0.03	-0.02	0.00	0.04	0.03	0.03	0.04
4	3	-0.03	-0.01	-0.04	0.00	0.00	0.02	0.23	0.19	0.18	0.18
5	1	-0.04	-0.03	-0.03	-0.02	-0.02	-0.01	0.39	0.34	0.30	0.27
5	2	-0.03	-0.02	-0.02	-0.01	-0.01	0.01	0.11	0.09	0.09	0.08
5	3	-0.16	-0.14	-0.14	-0.12	-0.11	-0.09	0.10	0.09	0.08	0.08

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: net change in dead wood (changeCd,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.16	0.10	0.10	0.11	0.04	0.11				
1	2	-0.06	-0.09	-0.07	-0.06	-0.08	-0.06				
1	3	0.19	0.12	0.13	0.13	0.07	0.10				
2	1	0.19	0.12	0.12	0.13	0.03	0.12				
2	2	0.13	0.06	0.07	0.08	-0.01	0.07				
2	3	0.13	0.06	0.07	0.08	-0.01	0.07				
3	1	0.10	0.03	0.06	0.07	-0.03	0.03				
3	2	-0.13	-0.18	-0.14	-0.11	-0.15	-0.13				
3	3	0.20	0.09	0.13	0.14	0.03	0.07				
4	1	0.23	0.16	0.16	0.16	0.09	0.12				
4	2	0.07	0.01	0.02	0.04	0.00	0.01				
4	3	0.20	0.11	0.13	0.14	0.07	0.09				
5	1	0.26	0.22	0.20	0.19	0.16	0.16				
5	2	0.09	0.06	0.06	0.07	0.05	0.05				
5	3	0.08	0.06	0.06	0.06	0.04	0.04				

(Table 6-20 continued)

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: net change in litter (changeCh,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	-0.14	0.06	-0.08	-0.04	-0.16	-0.05	0.30	0.16	0.04	-0.01
1	2	-0.09	0.07	-0.05	-0.01	-0.13	-0.01	0.25	0.09	0.07	0.02
1	3	-0.17	-0.05	-0.12	-0.06	-0.15	-0.02	0.17	0.02	0.06	0.03
2	1	-0.12	0.10	-0.05	-0.04	-0.16	-0.02	0.18	0.09	0.01	-0.05
2	2	-0.09	0.10	-0.03	-0.02	-0.15	0.00	0.18	0.05	0.03	-0.02
2	3	-0.09	0.10	-0.03	-0.02	-0.15	0.00	0.18	0.05	0.03	-0.02
3	1	-0.05	0.10	-0.01	-0.02	-0.14	0.02	0.05	-0.04	-0.06	-0.05
3	2	-0.06	0.08	-0.05	-0.01	-0.15	0.03	0.23	0.02	0.07	0.04
3	3	-0.13	0.00	-0.11	-0.03	-0.16	0.02	0.15	-0.07	0.06	0.03
4	1	-0.20	-0.05	-0.14	-0.09	-0.16	-0.05	0.18	0.08	0.10	0.02
4	2	-0.15	-0.02	-0.13	-0.06	-0.16	0.00	0.19	0.02	0.08	0.01
4	3	-0.06	0.06	-0.10	0.02	-0.15	0.06	0.24	0.04	0.13	0.01
5	1	-0.13	-0.07	-0.10	-0.08	-0.09	0.00	0.38	0.22	0.21	0.15
5	2	-0.17	-0.10	-0.13	-0.09	-0.11	-0.01	0.29	0.13	0.14	0.11
5	3	-0.02	0.06	-0.06	0.01	-0.10	0.09	0.44	0.17	0.20	0.16

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: net change in litter (changeCh,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.01	0.00	-0.05	0.24	0.00	0.08	-0.11	-0.18	-0.05	0.00
1	2	0.00	0.02	-0.05	0.15	0.02	0.10	0.00	-0.07	0.03	0.06
1	3	-0.01	0.03	-0.04	0.04	0.03	0.08	0.03	0.00	0.06	0.06
2	1	-0.05	-0.04	-0.06	0.20	-0.01	0.08	-0.20	-0.26	-0.12	-0.05
2	2	-0.05	-0.01	-0.06	0.15	0.01	0.09	-0.11	-0.18	-0.05	-0.01
2	3	-0.05	-0.01	-0.06	0.15	0.01	0.09	-0.11	-0.18	-0.05	-0.01
3	1	-0.09	-0.03	-0.09	0.10	0.00	0.05	0.07	-0.09	-0.01	0.03
3	2	-0.02	0.04	-0.07	0.07	0.05	0.09	-0.12	-0.16	-0.05	-0.02
3	3	-0.03	0.04	-0.07	-0.01	0.06	0.10	-0.20	-0.16	-0.05	-0.05
4	1	0.04	0.01	-0.05	0.10	0.04	0.10	-0.05	-0.11	-0.02	0.01
4	2	-0.03	0.04	-0.08	0.09	0.08	0.14	-0.04	-0.10	-0.03	0.02
4	3	-0.05	0.06	-0.09	0.12	0.13	0.19	0.04	-0.04	0.01	0.09
5	1	0.08	0.12	0.04	0.14	0.04	0.17	0.11	-0.01	0.00	0.02
5	2	0.05	0.09	0.02	0.10	0.03	0.15	0.15	0.04	0.06	0.05
5	3	0.07	0.13	0.01	0.14	0.09	0.28	0.15	0.04	0.08	0.07

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: net change in litter (changeCh,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.07	-0.06	-0.04	0.02	-0.15	0.08				
1	2	0.15	-0.08	0.00	0.05	-0.15	0.02				
1	3	0.15	-0.09	0.03	0.06	-0.11	-0.03				
2	1	0.04	-0.08	-0.05	0.00	-0.17	0.05				
2	2	0.09	-0.09	-0.01	0.03	-0.18	0.03				
2	3	0.09	-0.09	-0.01	0.03	-0.18	0.03				
3	1	0.11	-0.09	0.03	0.07	-0.18	0.02				
3	2	0.10	-0.16	0.00	0.05	-0.20	-0.05				
3	3	0.10	-0.21	-0.02	0.04	-0.17	-0.10				
4	1	0.10	-0.06	-0.01	0.03	-0.14	0.01				
4	2	0.13	-0.13	-0.01	0.06	-0.14	-0.04				
4	3	0.22	-0.14	-0.04	0.08	-0.13	-0.07				
5	1	0.10	-0.03	0.01	0.02	-0.09	0.02				
5	2	0.14	-0.03	0.02	0.04	-0.06	-0.01				
5	3	0.22	-0.09	0.00	0.07	-0.07	-0.04				

(Table 6-20 continued)

NFI	Elev.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
		CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001
1	2	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.001	0.001
1	3	-0.001	-0.001	-0.001	-0.001	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002
2	1	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.001
2	2	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.001	0.001	0.001
2	3	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.001	0.001	0.001
3	1	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
3	2	0.001	0.001	0.001	0.001	0.001	0.000	0.001	0.001	0.001	0.002
3	3	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
4	1	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	0.000	0.000
4	2	0.000	0.000	0.000	0.000	-0.001	-0.001	0.000	0.000	0.000	0.000
4	3	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.002	0.002	0.002
5	1	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002	-0.002	-0.001	-0.001
5	2	-0.001	-0.001	-0.002	-0.002	-0.002	-0.002	-0.002	-0.001	-0.001	-0.001
5	3	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.002	0.002	0.002

NFI	Elev.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
		CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
1	2	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
1	3	-0.002	-0.002	-0.002	-0.002	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001
2	1	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
2	2	0.001	0.001	0.001	0.001	0.001	0.002	0.002	0.001	0.001	0.001
2	3	0.001	0.001	0.001	0.001	0.001	0.002	0.002	0.001	0.001	0.001
3	1	0.001	0.001	0.000	0.000	0.001	0.001	0.001	0.001	0.001	0.001
3	2	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
3	3	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.001	-0.001
4	1	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
4	2	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
4	3	0.002	0.002	0.002	0.001	0.002	0.002	0.002	0.002	0.002	0.002
5	1	-0.001	-0.001	-0.001	-0.001	-0.001	0.000	0.000	0.000	0.000	0.000
5	2	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001
5	3	0.002	0.002	0.002	0.002	0.002	0.002	0.003	0.003	0.003	0.003

NFI	Elev.	2010	2011	2012	2013	2014	2015				
		CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹]									
1	1	0.001	0.001	0.001	0.001	0.001	0.001				
1	2	0.002	0.002	0.002	0.002	0.002	0.002				
1	3	-0.001	-0.001	-0.001	-0.001	-0.002	-0.002				
2	1	0.001	0.001	0.001	0.001	0.001	0.001				
2	2	0.001	0.001	0.001	0.001	0.001	0.001				
2	3	0.001	0.001	0.001	0.001	0.001	0.001				
3	1	0.001	0.001	0.001	0.001	0.001	0.001				
3	2	0.002	0.002	0.002	0.002	0.002	0.002				
3	3	-0.001	-0.001	-0.001	-0.001	-0.002	-0.002				
4	1	0.001	0.001	0.001	0.001	0.001	0.000				
4	2	0.000	0.000	0.000	0.000	0.000	0.000				
4	3	0.002	0.002	0.002	0.002	0.002	0.002				
5	1	0.001	0.001	0.001	0.001	0.001	0.001				
5	2	0.000	0.000	0.000	0.000	0.000	0.000				
5	3	0.003	0.004	0.004	0.004	0.004	0.004				

Carbon stock changes in the soil pool are small (Table 6-20). The Yasso07 data are supported by measurements of the Swiss Soil Monitoring Network (see chp. 6.4.4). Carbon

stock changes in litter are higher and more erratic than changes in the dead wood and soil pools (Fig. 6-6). This is expected since non-woody material decomposes faster than dead wood (Tuomi et al. 2011) and there is a higher interannual variability in the production of foliage (Etzold et al. 2011). The carbon stock change in the dead wood pool after 2000 is to a large extent driven by the increase in the dead wood stocks following the hurricane Lothar (1999). As Lothar occurred between the NFI2 (1993–1995) and NFI3 (2004–2006), it strongly affects the results of the change analysis for dead wood volume in the period NFI2 to NFI3. Although the majority of the windthrown trees were removed from the forest, the dead wood stock increased significantly. As particularly the larger sized felled trees decay slowly (Didion et al. 2014a), the storm resulted in a sustained carbon sink. The additional dead wood pool which was created by the storm will slowly release the stored carbon over the coming decades. The trend of decreasing harvest rates for several years after NFI3 (Table 6-16) further sustained the carbon sink of dead wood as mature trees, which could be harvested remain in the forest to potentially contribute to the dead wood pool. Large-scale disturbance events like Lothar that occur between two consecutive NFIs strongly affected the estimates of annually accumulating mass of carbon in dead wood that drives the Yasso07 simulation. This bias is expected to disappear following the switch to a continuous sampling approach in the NFI4 (Brändli and Speich 2011).




Figure 6-6 Mean carbon stock change (CSC) for three pools soil (0–100 cm), litter, dead wood and their sum (ALL), 1990–2015, in $t\ C\ ha^{-1}$. Note the difference of the y-axis scale between $Soil_{csc}$ and $Litter_{csc}$, $Dead\ wood_{csc}$ and ALL_{csc} , respectively. Negative values indicate losses in carbon stock, positive values gains in carbon stock. The error bars indicate the double standard error.

6.4.2.8 Unproductive forests (CC13)

Unproductive forests consist of brush forests, inaccessible stands and unproductive forest not covered by the NFI. Unproductive forests exhibit a high variability (see examples of unproductive forests in Switzerland in FOEN 2014f).

For transparency reasons, productive and unproductive forest areas are reported separately. However, there is only scarce information available on unproductive forests. In unproductive forests, wood is not harvested for economic reasons. Only in exceptional cases (e.g. wood log blocks a hiking trail) there can be an intervention where the log is moved, but not removed from the stand. Moreover, since yearly harvesting amounts from forest statistics (FOEN 2016k) are distributed over the productive forests, total harvesting in Swiss forests

was accounted for under productive forests (CC12), and thus all harvesting amounts were accounted for.

The NFI does not include unproductive stands CC13 in its regular inventory scheme because (1) the plots are difficult to access or it is not possible to carry out precise measurements (brush forests), (2) the plots are inaccessible or (3) the NFI forest definition is not fulfilled (forest not covered by the NFI).

- **Brush forests:** Since brush forests have no direct economic value in terms of wood harvest, an inventory of these stands has not been attributed high priority. During NFI3, some plots in brush forests were visited for the first time, but only a limited number of attributes such as tree species, stem diameter and crown cover were collected.
- **Inaccessible stands:** Inaccessible stands are forests which cannot be visited because of safety reasons (see description in Brändli 2010: 89). They are mainly located in the Alps and often grow on sites of low productivity, including rocky sites and sites at high elevation near the tree line with a short vegetation period and low biological activity.
- **Unproductive forests not covered by NFI:** After the review of its first Initial Report (FOEN 2006h), Switzerland had to apply a forest definition for reporting activities under the Kyoto Protocol Art. 3.3 and Art. 3.4, which is different from the definition applied by the Swiss NFI and the Land Use Statistics AREA. The same definition is used for reporting under the UNFCCC and under the Kyoto Protocol. Because the country definition (NFI and AREA) was not in line with the specific requirements of the Kyoto Protocol forest definition, Switzerland had to develop an approach to classify certain AREA categories as forest. Those areas are not covered by the regular NFI and are situated in the threshold range between forests and alpine pastures with woody biomass of very low productivity. More specifically, it concerns combination categories of “pastures or grasslands with clusters of trees” (NOLC04 47/NOLU04 222, NOLC04 47/NOLU04 223, NOLC04 47/NOLU04 242) and “alpine sheep grazing pastures, in general with open forest” with “clusters of trees”) NOLC04 44/NOLU04 243; cf. Table 6-6).

Carbon stocks in living biomass

- **Brush forest:** Brush forests in Switzerland mainly consist of *Alnus viridis*, horizontal *Pinus mugo* var. *prostrata* with a percentage cover of 65% and 16%, respectively (Table 1 in Düggelein and Abegg 2011). Following the NFI definition, brush forests are dominated by more than two thirds by shrubs. For brush forests, no NFI data are available to derive their growing stock. In a case study, Düggelein and Abegg (2011) analysed the carbon stock of total living biomass in Swiss brush forests and found an average value of 20.45 t C ha⁻¹.
- **Inaccessible stands:** Inaccessible stands are considered similar to brush forest regarding biomass and carbon stock. Their area is determined based on land cover ‘tree vegetation’ in typically remote and high-elevation land uses such as avalanche chutes (NOLU04 403 and 422; Table 6-6).
- **Unproductive forests not covered by NFI:** These forests are mainly associated with extensively pastured land where sparse tree vegetation (NOLC04 44 and 47; Table 6-6) is found. As those forests are assumed to grow preferably on bad site conditions, an average growing stock (>7 cm diameter) of 150 m³ ha⁻¹ was assumed. Multiplied by the mean BCEF of 0.69 (i.e. weighted mean based on the quotient of stemwood volume and total tree biomass of coniferous and broadleaved trees as described in Thürig and Herold 2013), an average biomass for these forests of 102.75 t ha⁻¹ was estimated, which corresponds to 51.38 t C ha⁻¹ (using a carbon content of 50%; see chp. 6.4.2.4).

The carbon stock of living biomass in unproductive forest was calculated as a weighted average of brush forest, inaccessible stands and unproductive forest not covered by NFI per spatial stratum:

$$\text{stockC}_{i,i,13} = F_i * \text{stockC}_{i,i,13bi} + (1 - F_i) * \text{stockC}_{i,i,13u}$$

where F_i is the fraction of the brush and inaccessible forest per spatial stratum i ,

$\text{stockC}_{i,i,13bi}$ is the carbon stock of brush and inaccessible forest (20.45 t C ha⁻¹),

$\text{stockC}_{i,i,13u}$ is the carbon stock of forest on unproductive areas (51.38 t C ha⁻¹).

Table 6-21 shows the resulting carbon stocks in living biomass of unproductive forest per spatial stratum in t C ha⁻¹.

Table 6-21 Area of brush forest, inaccessible forest and unproductive forest not covered by NFI, their areal fractions (F_i : fraction of brush and inaccessible forest per stratum i) and the resulting weighted carbon stocks in living biomass in t C ha⁻¹ of unproductive forests (CC13) specified for all spatial strata ($\text{stockC}_{i,i,13}$).

NFI region	Altitude [m]	Brush forest [ha]	Inaccesssible forest [ha]	Forest not covered by NFI [ha]	Fraction of brush and inaccessible forest (F_i)	Fraction of forest not covered by NFI ($1-F_i$)	Carbon stock in living biomass ($\text{stockC}_{i,i,13}$) [t C ha ⁻¹]
1	<601	49	0	69	0.42	0.58	38.53
	601-1200	44	0	4'841	0.01	0.99	51.10
	>1200	6	0	4'648	0.00	1.00	51.34
2	<601	188	0	0	1.00	0.00	20.45
	601-1200	94	0	93	0.50	0.50	35.83
	>1200	1	0	633	0.00	1.00	51.33
3	<601	11	0	0	1.00	0.00	20.45
	601-1200	172	0	1'210	0.12	0.88	47.53
	>1200	3'486	5	8'482	0.29	0.71	42.36
4	<601	26	0	1	0.96	0.04	21.60
	601-1200	1'058	5	589	0.64	0.36	31.48
	>1200	42'795	50	18'808	0.69	0.31	29.88
5	<601	243	1	3	0.99	0.01	20.83
	601-1200	2'249	0	275	0.89	0.11	23.82
	>1200	17'776	7	2'568	0.87	0.13	24.35

Carbon stocks in dead wood, litter, and mineral soil

As stated above, CC13 consists of different types of forests and data are hardly available. So far, there are no data available for carbon stocks in dead wood in unproductive forests (CC13). Dead wood on CC13 forest stands was assumed to be zero.

Carbon stocks in litter and in mineral soil under unproductive forests reveal a high spatial heterogeneity, and specific data are not available. Both carbon stocks of soil carbon and litter were assumed to be the same as for productive forests, which were derived from Nussbaum et al. (2012, 2014) (see Table 6-18).

Values for carbon stocks in dead wood, litter, and in mineral soil for CC13 are listed in Table 6-4.

Changes in carbon stocks of living biomass

There are a few case studies on carbon stocks, but similarly to neighbouring countries with forests in mountainous regions, there are no repeated forest inventory data available for these unproductive forests (also known as “mountain forest without harvest”). As no harvesting is conducted in unproductive forests, gross growth and cut and mortality of unproductive forest were assumed to be in balance. This approach is confirmed by three studies in which basal area and crown cover were used as a proxy for the stock of living biomass (Huber and Thürig 2014; Ginzler 2014; Huber and Frehner 2013). An increase in basal area or crown cover, respectively, was positively correlated with an increase in living biomass (e.g. Nowak and Crane 2002). Living biomass in brush forests was increasing during the stage of establishment: the stand developed from a stand with grasses, herbs and some shrubs towards a stand dominated by shrubs and with a denser crown cover. A decrease in crown cover in unproductive forests was observed when natural disturbances like avalanches or rock fall partially damaged the stand. The following studies provide evidence that living biomass in unproductive forests is not a source of carbon:

- Huber and Thürig (2014) analysed the available data on diameters of the terrestrial inventories NFI3 and NFI4 2009-2012. The authors found that the number of trees had increased over the approximately 6 year period between the two inventories. Since no allometric functions were available for these stands, it was not possible to calculate stocks from these data. The authors estimated an increase in the mean basal area from 4.59 m² ha⁻¹ in 2006 to 5.47 m² ha⁻¹ in 2012.
- Ginzler (2014) analysed the crown cover density of 135 aerial photographs between 2006 (NFI3) and 2011 (NFI4) and found no statistical change in crown cover density of well-established, existing brush forests. The terrestrial NFI data, however, showed a slight increase in the basal area of trees in brush forests.
- Huber and Frehner (2013) showed that the expansion of Green Alder (*Alnus viridis*) in eastern Switzerland has doubled in the past 75 years. Especially in the Alps or at unproductive sites, brush forests were expanding as summer pastures were abandoned. At these sites, an increase in crown cover was observed which correlates with an increment in carbon stocks. A literature review by Huber and Frehner (2012; for an overview see FOEN 2014f) showed that Green Alder has in general a strong annual gross growth, not only in very young stands, and that stands of Green Alder can be very vital at an age of over 100 years.

Considering the observed dynamics in Swiss brush forests, it was concluded that living biomass in unproductive forests was not a net source of carbon over the last decades. Applying a Tier 1 approach, living biomass is reported to be in equilibrium. In Table 6-4 and in CRF Table4.A, this approach is transcribed into “gains (gain_{C_{i,i,13}}) = losses (loss_{C_{i,i,13}}) = 0”.

Changes in carbon stocks of dead wood, litter, and mineral soil

There are no repeated measurements of carbon stocks in dead wood, in litter, and in mineral soil.

Above, transparent and verifiable information is given that in Switzerland living biomass in brush forest is increasing. An increase in biomass leads to an increase in dead wood

production and in litter, which in turn can lead to an accumulation in soil carbon. Based on these conceptual considerations, it was concluded that dead wood, litter, and mineral soil in unproductive forests were not a net source of carbon over the last decades. Applying a Tier 1 approach, thus, dead wood, litter, and mineral soil are reported to be in equilibrium. In Table 6-4 and in CRF Table 4.A, this approach is transcribed into “ $\text{change}C_{d,i,11} = \text{change}C_{h,i,11} = \text{change}C_{s,i,13} = 0$ ”. The Tier 1 approach is supported by the following evidences:

- Unproductive forest stands occur on higher elevation where microbiological processes in soils are slow (Hagedorn et al. 2010; Davidson and Janssens 2006).
- Unproductive forests grow on poor or rocky sites with thin or no organic layer. Brush forest protect the soils; in particular Alder brush is not even destroyed by avalanches or small-to-medium rock fall (Huber and Frehner 2014). By stabilizing soils, brush forests act as a good protection against soil erosion (Richard 1995; Stangl 2004).
- Green Alder has an ameliorative effect on the soil with its nitrogen-fixing root nodules (Huber and Frehner 2014). Amelioration of soils enables an increase in biomass production which on the other hand increases the amount of litter and dead wood and finally leads to accumulation of soil carbon.
- No active logging occurs on unproductive stands and consequently, there is no human impact on the soils, litter and dead wood.

By providing this transparent and verifiable information (survey of peer-reviewed literature and reasoning based on sound knowledge of likely system responses), the requirements for an application of the Tier 1 approach are considered to be fulfilled.

For conversions within Forest land (CC13 to CC12 and CC12 to CC13), no changes in carbon stocks of litter and soil carbon of mineral soils were calculated because carbon stocks of litter and mineral soil are the same for CC12 and CC13.

With the exception of brush forests, it is very likely that carbon stocks in litter and in mineral soil are smaller under unproductive forests than under productive forests. As the area changing from CC13 to CC12 is larger than from CC12 to CC13 (see Table 6-9), by applying the stock-difference method (see Table 6-3) with the same carbon stocks for litter and mineral soil under productive and unproductive forest, the resulting emissions are not underestimated.

6.4.2.9 Afforestations (CC11)

Carbon stock and changes in carbon stock changes of living biomass

Thürig and Traub (2015: Table 6) estimated the average carbon stock and gains and losses in living biomass of afforestations in Switzerland. Data are shown in Table 6-4.

In Switzerland, land-use change from non-forest to forest is usually not caused by plantation but by abandonment of agricultural land-use (Rutherford et al. 2008). These newly forested areas are often characterized by continuously growing trees with a large diversity in diameter at breast height (DBH) and tree age. Afforested stands established by plantation or even-aged young forest stands, however, are generally characterized by a large number of trees in small DBH classes and few trees in large DBH classes. Thürig and Traub (2015) selected

NFI plots to represent both types of afforestation. Young stands were defined as stands that changed from non-forest to forest between two consecutive NFIs with at least 85% of the trees with a DBH smaller or equal to 20 cm. As there is almost no land-use change from non-forest to forest below 600 m above sea level, results were stratified for below 1200 m above sea level and above 1200 m. As a consequence of the plot selection, small losses caused by natural mortality or cut of single trees occur.

Carbon stock and changes in carbon stock of dead wood and litter

On afforestations, carbon stocks in litter and dead wood were assumed to be zero (IPCC 2006, Volume 4, chp. 4.3.2). Applying the stock-difference calculation approach (Table 6-3), calculated changes in the litter and dead wood pool after a afforestation were rather small since the major part of afforestations (CC11) in Switzerland occur on grasslands and in settlements (see Table 6-9) where there is no litter and no dead wood (Table 6-4).

Carbon stock and changes in carbon stock of mineral soil

The estimates for soil carbon stocks from Nussbaum et al. (2012, 2014) were used for afforestations (see Table 6-4 and Table 6-18). Carbon stock changes were calculated with the stock-difference method (see Table 6-3).

6.4.2.10 Organic soils

Carbon stock in organic soils

The mean soil organic carbon stock (0–30 cm) for organic soils under forest land is $145.6 \pm 24.1 \text{ t C ha}^{-1}$ (Wüst-Galley et al. 2016). This value was used for CC11, CC12, and CC13 (cf. Table 6-4).

Changes in carbon stocks of organic soils

Drainage of forests is not a permitted practice in Switzerland (Swiss Confederation 1991). There are no nation-wide survey data available. It is possible that parts of the Swiss forest have been drained before 1990 or have been established on drained areas. In order not to underestimate the emissions, all organic forest soils were assumed to be drained.

For the calculation of changes in carbon stocks of organic soils, the default emission factor of $2.6 \text{ t C ha}^{-1} \text{ yr}^{-1}$ was applied for all forest stands (CC11, CC12, and CC13; cf. Table 6-4) according to the Wetlands Supplement (IPCC 2014a: Table 2.1).

6.4.2.11 N₂O emissions from Forest land

Fertilization of forests is prohibited by the Federal Act on Forest and the adherent ordinance (Swiss Confederation 1991, 1992). The Federal Act on Forest (Art. 18) states: “The use of

environmentally hazardous substances in the forest is prohibited” with a direct reference to the Federal Act on the Protection of the Environment (Swiss Confederation 1983). Details of the Federal Act on Forest Art. 18 had initially been regulated in the Ordinance on Forest (Art. 27). Since 2005, the Ordinance on Chemical Risk Reduction (Swiss Confederation 2005: Art. 4) prohibits the application of fertilizers, including liming, in forests. Hence, the application of fertilizers, including liming in forests was prohibited since 1991 in Switzerland. Furthermore, these management practices have never been common practice in Swiss forestry. There is thus considerable evidence to justify the assumption that this situation is valid since 1990. Therefore, no emissions were reported in category A. in CRF Table4(I) (notation key “NO”).

N₂O emissions from drainage of organic soils was calculated for Forest land with an emission factor of 2.8 kg N₂O-N ha⁻¹ and reported in category A. in CRF Table4(II). This is the default value given in the Wetlands Supplement (IPCC 2014a, Table 2.5) for temperate forest land.

The calculation of emissions reported in CRF Table4(III) and Table4(IV), i.e. direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils, is described in chp. 6.10.

6.4.2.12 Emissions from wildfires

Data on wildfires affecting Swiss forest land were obtained from cantonal authorities and were compiled by the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL, Swissfire database, <http://www.wsl.ch/swissfire>). Table 6-22 shows the time series 1990 to 2015 of the area burnt and the associated emissions.

As controlled burning of forest stands is not allowed in Switzerland all fires in forests were considered “wildfires”. All fires were assigned to productive forests. In this way, emissions are not underestimated, since the “available fuel” of productive forests is higher than the carbon stocks of afforestations and unproductive forests. Moreover, this approach reflects reality quite well, since fires on afforestations or in unproductive forests are rather unlikely to occur for the following reasons:

- Non-Forest land to Forest land (or Afforestations under the Kyoto Protocol Art. 3.3) and unproductive forest: the “available fuel” is small, there is very little dead woody material on the surface which can catch fire (Zumbrunnen et al. 2012).
- Unproductive forests: the “available fuel” is small since tree cover is not very dense (Zumbrunnen et al. 2012). Moreover, in remote areas the cause of fire is restricted to lightning strikes.

CO₂ emissions from wildfires were encompassed (“IE”) in the data in CRF Table4.A. Losses in living biomass are reflected in the NFI dataset. Carbon changes in dead wood, litter and soil carbon calculated with Yasso07 also cover the influence of forest fires and other disturbances by using NFI data as an input (see chp. 2.3.3 in Didion and Thürig 2016).

CH₄ and N₂O emissions from wildfires (Table 6-22) were calculated using Equation 2.27 in Volume 4 of IPCC (2006) with the following parameters:

- For CH₄ the default emission factor of 4.7 g kg⁻¹ dry matter burned and for N₂O, the default emission factor of 0.26 g (kg combusted biomass)⁻¹ was applied (IPCC 2006 Volume 4, Table 2.5).
- The mass of “available fuel” encompasses carbon stocks of living biomass, dead wood, and litter. On average, the amount of living biomass amounts to 92.84 t C ha⁻¹ or 185.67 t biomass ha⁻¹. This value was derived from the mean growing stock in NFI1, NFI2, NFI3 and NFI4 2009–2013 (Brassel and Brändli 1999; Brändli 2010; Abegg et al. 2014) as a weighted value of the regions affected by forest fires (82% of the fires occur in the southern Alps, 15% in the Central Alps; FOEN 2016k). The average amount of litter in Swiss forests was 16.73 t C ha⁻¹ (Nussbaum et al. 2012; see) or 33.46 t biomass ha⁻¹ and average stocks of dead wood were calculated per NFI-period based on data from Didion and Thürig 2016 (see Table 6-19).
- The fraction of the biomass combusted was 0.45 (IPCC 2006, Volume 4, Table 2.6).

CH₄ and N₂O emissions caused by wildfires were reported in CRF Table4(V). CO₂ emissions caused by wildfires were included in CRF Table4.A and thus identified as “IE” in CRF Table4(V). Further, in CRF Table4(V), CH₄ and N₂O emissions from wildfires of all types of forests were reported under 4(V)A1, because it is not known which fires occur on Forest land remaining forest land and which on Land converted to forest land. Consequently, category 4(V)A2 has the notation key "IE".

Table 6-22 Productive forest land affected by wildfires (WSL, Swissfire database) and resulting CH₄ and N₂O emissions 1990–2015.

Forest land	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Area burnt	ha	1'060.6	70.5	26.9	17.7	233.2	362.8	232.0	1'389.5	197.7	11.2
CH ₄	t	529.4	35.2	13.4	8.8	116.4	181.1	115.8	693.6	98.670	5.590
N ₂ O	t	29.3	1.9	0.7	0.5	6.4	10.0	6.4	38.4	5.458	0.309

Forest land	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Area burnt	ha	47.4	12.9	418.0	527.0	24.6	39.1	110.9	234.9	37.4	48.5
CH ₄	t	23.7	6.5	208.7	263.1	12.3	19.5	55.3	117.2	18.673	24.188
N ₂ O	t	1.3	0.4	11.5	14.6	0.7	1.1	3.1	6.5	1.033	1.338

Forest land	Unit	2010	2011	2012	2013	2014	2015
Area burnt	ha	25.9	168.7	23.7	24.2	43.2	42.3
CH ₄	t	12.9	84.2	11.8	12.1	21.6	21.1
N ₂ O	t	0.7	4.7	0.7	0.7	1.2	1.2

6.4.2.13 Emissions from controlled burning

CH₄ and N₂O emissions from open burning of residues from forests were reallocated from category 5C2 (open burning of waste) in the waste sector to category 4(V)A1 (controlled burning in forest land remaining forest land) as requested by the ERT during the in-country review 2016. The rationale was provided in UNFCCC (2017/ID#W.13), see also UNFCCC (2017/ID#L.13) and UNFCCC (2017a/ID#1).

Emissions from controlled burning covers the burning of residues in forestry; controlled burning of forest stands is not allowed in Switzerland.

Emissions were calculated by a Tier 2b approach based on chp. 5.2. in Volume 5 of IPCC (2006). The emissions of burning of residues in forestry are calculated by multiplying the annual estimate of branches burnt (in kt) by emission factors (IPCC default values).

The amount of natural residues burnt openly was estimated by INFRAS (2014). Open burning of such residues is regulated in the Ordinance on Air Pollution Control OAPC, (Swiss Confederation 1985: Art. 26b). In Switzerland cantonal authorities are responsible for surveilling that the regulations of the OAPC are respected. For INFRAS (2014) an inquiry of some cantonal authorities has been performed in order to assess the activity data for these processes.

The emission factors of burning of branches in forestry were calculated based on EMEP/EEA (2013) except for CH₄ und N₂O which were based on EMEP/CORINAIR (EMEP/EEA 2002), see also documentation in EMIS (2016/5C2 *Abfallverbrennung in der Land- und Forstwirtschaft*).

6.4.2.14 NMVOC emissions

Estimates for annual biogenic emissions of NMVOC in Switzerland for forests (and natural grassland) are available in SAEFL (1996a): The values are 92.0 kt for coniferous forests, 2.4 kt for deciduous forests and 0.61 kt for forest fires. These numbers are based on a study by Andreani-Aksoyoglu and Keller (1995). Approximately 97% of the total emissions were monoterpene and the rest consisted of isoprene (Keller et al. 1995).

6.4.3 Uncertainties and time-series consistency

Uncertainties

For living biomass, the uncertainty was estimated based on the following information:

- Stem wood of growth (gains of living biomass) and cut & mortality (losses of living biomass) in NFI4 2009–2013 and differences between NFI3 and NFI4 2009–2013 (Abegg et al. 2014):
 - mean gain 8.95 m³ ha⁻¹ year⁻¹, mean loss -7.64 m³ ha⁻¹ year⁻¹, resulting mean net change in stem volume 1.31 m³ ha⁻¹ year⁻¹
 - relative uncertainty of mean net change in volume: 54% (double standard error (2SE); for calculation see Thürig et al. 2015)
- Carbon content in solid wood: The uncertainty was estimated to be 2% (2SE) based on Monni et al. (2007) (2% relative standard deviation; RSD), and Lamtom and Savidge (2003) (4-8% RSD).
- Biomass expansion function (for the Swiss GHG inventory, allometric functions for individual trees are applied) and conversion into mass with wood density: The uncertainty related to the expansion and conversion of stem volume to whole tree biomass was based on Lehtonen and Heikkinen (2016) including 21.2% (2SE) sampling uncertainty and 22.2% (2SE) model uncertainty.

Thus, the total uncertainty of net carbon stock change in living biomass ($U_{liv.biom}$) in terms of carbon per unit area can be calculated following equation 3.1 in chp. 'Quantifying Uncertainties' (Volume 1 of IPCC 2006):

$$U_{liv.biom} = \sqrt{54^2 + 2^2 + 21.2^2 + 22.2^2} = 62.15\%$$

The uncertainty in the estimates of annual carbon stock changes derived with the Yasso07 model originates from the following sources as described in Didion and Thürig (2016):

- carbon input estimates obtained from the NFI (measurement errors, allometries, etc.) (chp. 2.3.3 in Didion and Thürig 2016);
- decomposition parameters used in the Yasso07 model (chp. 2.3.1 in Didion and Thürig 2016).

No data are available yet to estimate the uncertainty associated with the spatially interpolated temperature and precipitation data. MeteoSwiss is currently developing ensembles of the gridded climate data that will allow to quantify the effect of this source of uncertainty on the annual carbon stocks; the ensembles are expected to become available in 2017. Based on preliminary results from MeteoSwiss (Vogel 2013), the additional uncertainty for carbon stock change estimates is expected to be minor.

The uncertainty associated with carbon inputs (dead wood production and litterfall) was estimated based on estimates of uncertainty in (a) litter turnover rates (Wutzler and Mund 2007), (b) wood densities of deadwood in different decay stages (Dobbertin and Jüngling 2009), and (c) spatial uncertainty in the NFI data approximated based on the estimation error for tree volume reported for the NFI (see chp. 1.4 in Brändli 2010). Based on the mean carbon inputs and the estimated uncertainty, a distribution of possible values was obtained. Finally, the combined uncertainty from these sources was calculated. The uncertainty in the Yasso07 parameters was estimated based on a Markov Chain Monte Carlo approach (see also Tuomi et al. 2011). A distribution of possible parameter values was provided by A. Lehtonen, Natural Resources Institute Finland. The uncertainty of Yasso07 estimates on carbon stocks and carbon stock changes in different pools, resulting from the uncertainty of carbon inputs and of model parameters, was obtained through Monte Carlo simulations: 10 values for carbon inputs and 10 parameter combinations were selected randomly and the combined uncertainty in Yasso07 estimates of carbon stocks and carbon stock changes in the soil, litter, and dead wood pools was calculated as described in Didion and Thürig (2016).

Based on this approach, the absolute uncertainty (double standard error) of the estimates of C stock changes are:

- $U_{Soil.abs} = 0.00014 \text{ t C ha}^{-1} \text{ yr}^{-1}$,
- $U_{Litter.abs} = 0.00371 \text{ t C ha}^{-1} \text{ yr}^{-1}$, and
- $U_{Deadwood.abs} = 0.00708 \text{ t C ha}^{-1} \text{ yr}^{-1}$,

which correspond to the relative uncertainties (2SE) of

- $U_{\text{Soil}} = 13.18\%$,
- $U_{\text{Litter}} = 2.31\%$, and
- $U_{\text{Deadwood}} = 679.46\%$.

The total uncertainty associated with carbon stock change in all four pools was estimated using equation 3.2 in chp. 'Quantifying Uncertainties' (Volume 1 of IPCC 2006):

$$U_{\text{tot}} = \frac{\sqrt{(U_{\text{liv.biom}} * X_{\text{liv.biom}})^2 + (U_{\text{soil}} * X_{\text{soil}})^2 + (U_{\text{Litter}} * X_{\text{Litter}})^2 + (U_{\text{Deadwood}} * X_{\text{Deadwd}})^2}}{|X_{\text{liv.biom}} + X_{\text{soil}} + X_{\text{Litter}} + X_{\text{Deadwood}}|}$$

With mean carbon stock changes in 2014 in

living biomass ($X_{\text{liv.biom}}$): $0.53 \text{ t C ha}^{-1} \text{ yr}^{-1}$,

soil (X_{Soil}): $0.001 \text{ t C ha}^{-1} \text{ yr}^{-1}$,

litter (X_{Litter}): $-0.160 \text{ t C ha}^{-1} \text{ yr}^{-1}$, and

dead wood (X_{Deadwood}): $0.001 \text{ t C ha}^{-1} \text{ yr}^{-1}$,

where positive values refer to gains in carbon stock; negative values refer to losses in carbon stock. Thus,

$$U_{\text{tot}} = \frac{\sqrt{(U_{\text{liv.biom}} * X_{\text{liv.biom}})^2 + (U_{\text{soil}} * X_{\text{soil}})^2 + (U_{\text{Litter}} * X_{\text{Litter}})^2 + (U_{\text{Deadwood}} * X_{\text{Deadwd}})^2}}{|X_{\text{liv.biom}} + X_{\text{soil}} + X_{\text{Litter}} + X_{\text{Deadwood}}|}$$

$$U_{\text{tot}} = \frac{\sqrt{(62.15 * -0.53)^2 + (13.18 * -0.001)^2 + (2.31 * 0.160)^2 + (679.46 * -0.001)^2}}{|(-0.53) + (-0.001) + 0.160 + (-0.001)|}$$

Thus, the resulting relative uncertainty of the total carbon stock change for forest land is 88.6%. It should be noted that this value is an overestimation. Responsible for the high value is the high uncertainty estimate for the mean net change in volume of living biomass (54%) which is the result of the calculation of the relative uncertainty for a small net change in volume living biomass of $1.31 \text{ m}^3 \text{ ha}^{-1} \text{ year}^{-1}$. In fact, the uncertainties of mean gains and losses in living biomass from which the net change is calculated are only $\pm 2\%$ and $\pm 4\%$, respectively. For the combined stock change from soil, litter and dead wood the uncertainty is 5%.

The emission factor uncertainty for wildfires is 70%. This is the default value given for non- CO_2 emissions in the Good Practice Guidance (IPCC 2003, chp. 3.2.1.4.2.4) and also corresponds to the uncertainty of the combustion factor from IPCC 2006 (Volume 4, Table 2.6, mean = 0.45, 2SE = 0.32).

Uncertainties of activity data of category 4A Forest land are described in chp. 6.3.3. Table 6-5 lists the relative uncertainties in the LULUCF sector.

Time-series consistency

Consistent time series of annual carbon stocks of living biomass were calculated backward or forward starting from the growing stock 2005, as derived from NFI3 (see chp. 6.4.2.5).

Consistent time series of dead wood, litter and soil carbon were calculated with the model Yasso07 (see Didion and Thürig 2016 and chp. 6.4.2.7).

6.4.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

Suitability of the soil carbon model Yasso07 for application for forests in Switzerland

The validity of the Yasso07 model in Swiss forests was examined by Didion et al. (2014a). The study analyzed, among other, the accuracy of Yasso07 for reproducing observed carbon decomposition in litter and dead wood in Swiss forests. The authors found that no significant differences existed between simulated and observed remaining carbon in foliage and fine root litter after 10 years and in lying dead trees after 14 to 21 years.

Afforestation – Growing stock and changes in growing stock

A comparison with IPCC default values and NFI data from neighbouring countries is included in Thürig and Traub (2015). The study supports the plausibility of the Swiss estimates: they are well within the range of the IPCC default values as well as the Austrian and German estimates.

Swiss estimates were also compared with literature values. Based on data of the German forest inventory (*Bundeswaldinventur II*), Paul et al. (2009) reported a carbon sequestration rate of $2.8 \text{ t C ha}^{-1} \text{ a}^{-1}$ in the first 20 years following an afforestation.

Afforestation – Litter

In an experiment by Zimmermann and Hiltbrunner (2012; COST E639-project “Turnover and stabilization of soil organic matter: effect of land-use change in alpine regions”), litter accumulation in a 40 year old afforestation with Norway Spruce was determined. The authors found accumulation rates of $0.17\text{-}0.20 \text{ t C ha}^{-1} \text{ yr}^{-1}$. Further relevant studies are discussed in chp. 11.3.1.2.

Carbon balance of two mountain forest ecosystems in Switzerland – Net ecosystem exchange and soil respiration

Measurements of the net ecosystem exchange (NEE) and of soil respiration were conducted at a montane mixed forest over 5 years (Lägeren; 2005–2009; NFI production region 2), and

at a subalpine coniferous forest over 12 years (Davos; 1997–2009; Swiss Plateau, NFI production region 4).

(1) Etzold et al. (2011) determined the net ecosystem exchange (NEE) by eddy covariance (EC) measurements. EC measurements, as well as biometric estimates indicate that both sites with two different mountain forest types were significant carbon sinks in the respective periods. During 2005 to 2009 NEE of the Lägeren forest ranged from -366 to -662 g C m⁻² yr⁻¹ (mean: -415 g C m⁻² yr⁻¹), and of the Davos forest from -47 to -274 g C m⁻² yr⁻¹ (mean: -154 g C m⁻² yr⁻¹).

(2) Rühr and Eugster (2009) measured soil respiration rates at these two Swiss forest sites. Modelled changes in soil carbon storage with the dynamic soil carbon model Yasso07 gave comparable results with measured soil respiration. Rühr and Eugster (2009) found that soils at the alpine site Davos acted as a significant carbon sink. Soils at the Lägeren site were neither a significant carbon sink nor a significant carbon source. This domestic study confirms the broadly spread knowledge that it is very difficult to detect short term changes in soil carbon stocks, since the uncertainty of the measurement is often higher than the actual change of the annual estimates (e.g. Falloon and Smith 2003).

Changes in soil carbon stocks – Soil organic carbon (SOC) dataset of the Swiss Soil Monitoring Network

The objective of the Swiss Soil Monitoring Network (<http://www.nabo.ch>; NABO) is to assess soil quality in the long term and to validate appropriate soil protection measures. NABO operates about 110 long-term monitoring sites throughout Switzerland. Most of them were sampled for the first time between 1985 and 1989 and resampled every five years ever since. 28 sites are located in forests (SAEFL 1993).

The long-term soil monitoring sites are resampled every 5 years. At each site, four replicate bulked soil samples from the upper soil layer 0–20 cm are taken within an area of 10m*10m. Each bulked sample consists of 25 single cores taken according to a stratified random sampling scheme. Further details are provided by SAEFL (2000a) and FOEN (2015p). Currently, results of sampling campaigns 1 to 5 are available for Forest Land and Grassland. For Cropland, additionally the results of sampling campaign 6 were already produced.

The spatial variation of bulk density was included in calculating the carbon pools. Bulk density and soil skeleton (>2 mm) were measured repeatedly for all monitoring sites at the occasion of sampling campaigns 4 to 6 (2000–2014), but not in the previous campaigns. The mass of fine earth (<2 mm; M_FE) per total soil volume (V_tot, including skeleton and pores) was determined for four volumetric samples 0–20 cm per site and campaign to derive the so-called apparent density of fine earth ($D = M_FE / V_tot$). Subsequently, SOC pools 0–20 cm [t/ha] were calculated by $D [g/cm^3] * SOC [\% w./w.] * 20 [cm]$. For each site, the site-specific apparent density was used; repeated apparent density measurements per site were used to account for the variability of the bulk density.

The SOC pools for the forest top soils (0–20 cm) ranged between 35.4 t C ha⁻¹ (min) and 135.8 t C ha⁻¹ (max) and were on average 70.6 t C ha⁻¹. In these numbers one coniferous forest site was excluded as it revealed large SOC pools up to 191 t C ha⁻¹. Figure 6-7 shows that on average, SOC pools did not change monotonously during the measurement period

between 1989 and 2009 in the sampled forest soils. At some of the forest monitoring sites higher values were found in the third resampling campaign.




Figure 6-7 Time series of measured SOC pool changes in the top soil (0–20 cm) at the 28 NABO forest sites from the 1st to the 5th resampling campaigns. SOC pools were centred by the median SOC pool of all resamplings of the monitoring site. Each pool value presents the median of four bulked soil samples per campaign with measured SOC and bulk density. The error bars indicate the 25% and 75% percentiles resulting from the spatial variation of the sites and the errors along the measurement chain. The elevation of the forest sites ranges between 380 and 1690 m a.s.l.

Detailed studies at monitoring sites showed that short-term temporal variation of soil properties can result from different site conditions at the sampling date, e.g. regarding soil moisture, soil temperature and bulk density (Keller et al. 2006). For instance, at two forest sites six resamplings within three years revealed short-term variation of the SOC content between $\pm 1.8\%$ and $\pm 0.6\%$ (simple standard error). Therefore, the majority of the measured temporal variation for all forest sites was interpreted as natural variation (noise) and not as real SOC changes (signal). This hypothesis is also supported by the fact that the soil samples in the third resampling campaign were taken earlier in spring time as in the other sampling campaigns and hence, soil moisture content of the samples was higher on average. This might explain the large temporal variation, in particular at coniferous forest sites with a pronounced organic layer. Using a robust linear regression approach for the SOC pool data of the forest soils, the 95% confidence interval for the SOC pool was $\pm 1.5 \text{ t C ha}^{-1}$. In order to capture as good as possible the natural variation of these site-specific characteristics, standard operation procedures and quality assurance were implemented since the 4th soil campaign. Further work will focus on the correction of the measured carbon pools to equivalent mass of the fine earth <2 mm. In this way, the 95% confidence interval of the mean SOC pool can probably be reduced to some degree.

In comparison, the mean change in SOC which was obtained with Yasso07 for the period 1991–2010 was $-0.00075 \pm 0.00053 \text{ t C ha}^{-1} \text{ yr}^{-1}$ (2SE; based on data in Didion and Thürig 2016). This value would correspond to $-0.015 \text{ t C ha}^{-1}$ for the monitoring period of the NABO network (20 years). The analysis strongly suggests that modelled SOC pool changes are not inconsistent with the repeated soil inventories in the NABO network. As indicated by the 95%

confidence interval for the 28 forest monitoring sites, the noise is two orders of magnitude higher than the modelled signal.

Uncertainty Estimates

The uncertainty for carbon stock changes in dead wood, litter and soil organic matter reported by Finland, where the Yasso07 model is also applied, was 31.5% for the year 2014 (Statistics Finland 2016: chp. 6.4.3.2). For the total uncertainty in the change of living biomass, Finland reported 20% (Statistics Finland 2016: chp. 6.4.3.1).

6.4.5 Category-specific recalculations

Activity data 1990–2014 were updated (see chp. 6.3.5).

The most recent NFI data (NFI4 2011–2015) were applied for productive forests (Thürig et al. 2017; see chp. 6.4.2.1).

Modelling carbon stock changes in dead wood, litter and mineral soil with Yasso07 for productive forests: the methodological change in the National Forest Inventory (NFI) from a periodic (i.e. NFI1, NFI2, and NFI3) to a continuous sampling starting with the NFI4 required modifications to the approach for estimating dead wood and litter inputs. To derive estimates of carbon inputs between NFI3 and NFI4, plots which were visited within the latest 5 years of the NFI4 were considered, i.e. currently 5-year tranche NFI4 2011–2015. Thus, the sites used in the Yasso07 simulations (cf. chp. 2.4.1 in Didion and Thürig 2016) changed in comparison with the GHGI inventory submission 2016 (FOEN 2016); N is now 2616 compared to 2607 in FOEN (2016) of which 1591 are common sites. Further, the accuracy of the estimates of dead wood and litter production was improved. Besides minor corrections, an inconsistency was removed in the NFI1 regarding the classification of laying trees as alive or dead. This resulted in more accurate estimates of the dead wood and litter production in the period NFI1–2.

In response to issue ID#W.13 in UNFCCC (2017) (see also UNFCCC 2017/ID#L.13 and UNFCCC 2017a/ID#1) CH₄ and N₂O emissions from open burning of residues from forests were reallocated from category 5C2 (open burning of waste) in the waste sector to category 4(V)A1 (controlled burning in forest land remaining forest land) (see chp. 6.4.2.13). Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

6.4.6 Category-specific planned improvements

Further research is underway to estimate annual values for carbon gains in living biomass. First results building on a study on the relationship between climate and gains in living biomass (Thürig et al. 2009) are expected in the next years.

The implementation of the soil model Yasso07 to improve the accuracy in the estimates of temporal changes in soil carbon, litter and dead wood will be further developed. Planned improvements (in line with UNFCCC (2017/ID#L.12) include:

- Investigating the validity of the further development of Yasso07 for application in Switzerland. The new model version Yasso15 includes, among other, a new parameter set that improves the sensitivity of the simulated decomposition to temperature and precipitation. Preliminary results from Switzerland using a beta-version of Yasso15 indicate improvements over Yasso07, particularly with regards to soil carbon stocks.
- Improving the completeness of the litter inputs by accounting for the contribution of (a) fine-woody litter <7 cm and (b) litter from the herb- and shrub layer. A field study was recently completed which will provide estimates on biomass turnover of plants in the herb- and shrub layer of NFI sites. These activities will improve the accuracy of the simulated estimates.
- Quantifying the effect of uncertainty associated with the spatially interpolated temperature and precipitation data on CSC estimates. MeteoSwiss is currently developing ensembles of the gridded precipitation data that will allow to address this source of uncertainty in the simulations with Yasso07. The ensembles are expected to become available in 2017.

Improvements to the application of Yasso07 in Switzerland are foreseen at the earliest for the GHG inventory 1990–2016 (to be submitted in 2018).

Projects in a new national research programme ([Sustainable Use of Soil as a Resource: "SOM control"](#)) aim at identifying the drivers of soil organic matter storage in Swiss forest soils. The objectives are to assess how forest productivity and tree species composition affect soil organic matter storage, to investigate if and how land-use history affects carbon pools in soils, to estimate the influence of climate, temperature and precipitation on soil organic matter stocks, to link soil organic matter stocks to physico-chemical parameters controlling soil organic matter stabilization and to model soil organic matter and evaluate the residuals to measured soil organic matter stocks. In the long term, results from these studies are expected to improve the reporting of emissions by sources and removals by sinks in category 4A. In a follow-up project financed by FOEN, the implications for carbon stabilization in mineral forest soils is investigated and the application to further development of the Yasso model is examined.

In response to issue ID#L.11 in UNFCCC (2017), data on the share of organic soils affected by past draining activities under Forest land will be collected for the next submission, using additional descriptive information from the NFI surveys (see chp. 6.4.2.10).

6.5 Category 4B – Cropland

6.5.1 Description

Table 6-23 Key categories (KCA including LULUCF) in category 4B.

Code	IPCC Category	GHG	Identification Criteria
4B1	Cropland remaining Cropland	CO2	L1, L2, T1, T2

Swiss croplands belong to the cold temperate wet climatic zone.

Carbon stocks in above-ground living biomass and carbon stocks in mineral and organic soils were considered.

Croplands (CC21) include annual crops and leys in arable rotations (see Table 6-2 and Table 6-6). Because arable cropping mainly occurs in the temperate Swiss Central Plateau and no elevation-dependent soil carbon stocks are available for Swiss croplands (Leifeld et al. 2005), no stratification of carbon stocks was applied.

In 2015, category 4B1 Cropland remaining cropland was a net source of 1028.78 kt CO₂ due to losses in living biomass and emissions from organic soils. Average living biomass was increasing slightly over the period 1990–2015. However, annual fluctuations in carbon stocks of biomass are considerable (see Table 6-24). Carbon stocks on mineral soils were assumed to be in balance (i.e. no carbon stock changes occur in mineral soils). Thus, all soil emissions in category 4B1 are considered to originate from carbon mineralization in organic soils, mainly in the lowest elevation zone (z1: 86%). Overall, organic soils accounted for 2.7% of cropland area in Switzerland (Table 6-7).

Category 4B2 Land converted to cropland was a small net source of 41.22 kt CO₂ in 2015 mainly due to carbon losses in mineral soils under Grassland converted to cropland.

6.5.2 Methodological issues

6.5.2.1 Carbon in living biomass

Annual biomass carbon stocks are shown in Table 6-24. They were calculated as area-weighted means of standing stocks at harvest for the seven most important annual crops (barley, wheat, maize, silage maize, sugar beet, fodder beet, potatoes) and as cumulated annual harvested biomass for leys.

The annual mean standing biomass carbon stock per hectare was calculated as:

$$\text{Biomass cropland} = \sum_f (A_f / A_t) * C_f$$

where A_f = Area of crop type f , A_t = total cropping area and C_f = yield (annual crops, leys) for the particular crop (t C ha⁻¹). Annual values for A_f , A_t and C_f were published by the Swiss Farmers Union (SBV 2016).

The resulting mean biomass stock for Swiss cropland over the inventory time period was 4.74 ± 0.36 (1 SD) t C ha⁻¹.

Table 6-24 Annual values for arable crop yields (SBV 2016), resulting area-weighted carbon stock means (t C ha⁻¹) and carbon stock changes (gain/loss) (t C ha⁻¹ yr⁻¹), assuming a carbon fraction of 0.5 (IPCC default). Highlighted data for 1990 are displayed in Table 6-4.

Crop	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	CC21: yield (stock) [t C ha ⁻¹] and gain/loss [t C ha ⁻¹ yr ⁻¹] in living biomass									
Barley	2.36	2.47	2.48	2.54	2.19	2.29	2.69	2.69	2.86	2.18
Wheat	2.36	2.54	2.35	2.53	2.34	2.56	2.84	2.55	2.63	2.24
Maize	3.51	3.42	3.53	3.78	3.72	3.55	3.64	3.94	3.87	3.78
Silage maize	7.37	6.59	7.15	6.72	6.11	6.03	4.98	7.08	6.88	6.49
Sugar beet	7.41	6.91	7.04	7.63	6.72	6.78	7.83	7.76	7.42	7.48
Fodder beet	6.70	6.51	6.64	6.77	5.66	5.49	6.41	6.53	6.06	5.79
Potatoes	4.47	4.39	4.65	4.83	3.65	3.88	5.36	5.05	4.44	3.87
Leys	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11
Mean yield	4.34	4.30	4.39	4.44	4.11	4.28	4.51	4.72	4.67	4.43
Gain/loss	-0.34	-0.03	0.09	0.05	-0.32	0.17	0.23	0.21	-0.05	-0.24

Crop	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	CC21: yield (stock) [t C ha ⁻¹] and gain/loss [t C ha ⁻¹ yr ⁻¹] in living biomass									
Barley	2.55	2.38	2.68	2.35	2.92	2.61	2.64	2.57	2.58	2.73
Wheat	2.53	2.35	2.42	2.16	2.62	2.46	2.49	2.57	2.58	2.60
Maize	4.10	3.80	3.92	1.83	4.09	4.10	3.30	4.32	4.12	4.42
Silage maize	6.68	6.45	4.93	5.96	6.52	8.23	7.01	8.02	8.09	7.60
Sugar beet	8.74	6.51	8.52	7.89	8.60	8.50	7.30	8.37	8.73	9.37
Fodder beet	6.71	5.75	5.95	5.67	6.13	6.15	6.25	6.21	6.30	6.72
Potatoes	4.67	4.13	4.30	3.71	4.34	4.26	3.60	4.59	4.71	5.12
Leys	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11	6.11
Mean	4.71	4.51	4.54	4.41	4.89	4.97	4.70	5.07	5.13	5.19
Gain/loss	0.28	-0.21	0.03	-0.13	0.49	0.08	-0.27	0.37	0.06	0.06

Crop	2010	2011	2012	2013	2014	2015	mean 1990-2015		
	CC21: yield (stock) [t C ha ⁻¹] and gain/loss [t C ha ⁻¹ yr ⁻¹] in living biomass								
Barley	2.56	2.75	2.75	2.45	3.19	3.00	2.59		
Wheat	2.48	2.72	2.49	2.32	2.73	2.65	2.50		
Maize	3.61	4.13	3.85	3.12	3.75	2.65	3.69		
Silage maize	7.47	8.42	8.06	7.79	8.46	7.20	7.01		
Sugar beet	8.03	10.38	9.58	7.61	10.06	7.55	8.03		
Fodder beet	6.49	6.65	5.58	4.67	5.52	4.95	6.09		
Potatoes	4.26	5.04	4.52	3.59	4.89	3.69	4.38		
Leys	6.11	6.11	6.11	6.11	6.11	6.09	6.11		
Mean	4.99	5.44	5.23	4.93	5.46	4.96	4.74		
Gain/loss	-0.20	0.45	-0.21	-0.30	0.52	-0.50	0.01		

6.5.2.2 Carbon in soils

Soil carbon stocks in mineral soils under cropland were calculated based on Leifeld et al. (2003, 2005). The approach correlated measured soil organic carbon stocks (t ha⁻¹) for arable land and leys with soil texture after correction for soil depth and stone content. Area upscaling used the Swiss digital soil map (SFSO 2000a), and average stocks were calculated as weighted means using the area of arable land and leys. The mean soil organic carbon stock (0–30 cm) for cropland was 53.40 ± 5 t C ha⁻¹ (uncertainty 9%).

It should be noted that current carbon stocks are not only the result of the conditions for productivity and carbon turnover under different land-use types, but are also determined by farmers' decisions to use a site in a specific way due to the demands of a crop or the suitability of a site, e.g. regarding machine use (see Leifeld et al. 2003: 65).

Soil carbon stocks in organic soils under cropland were calculated based on Leifeld et al. (2003, 2005). The approach used measured carbon stocks in Swiss organic soils. The mean soil organic carbon stock (0–30 cm) for cultivated organic soils was $240 \pm 48 \text{ t C ha}^{-1}$ (uncertainty 20%).

6.5.2.3 Changes in carbon stocks

Carbon stocks in living biomass intermittently increased from 4.34 t C ha^{-1} in 1990 to 4.96 t C ha^{-1} in 2015 (Table 6-24; SBV 2016). The difference in biomass stock between a specific year and the preceding year was reported as gain or loss of carbon (see Table 6-24). The resulting values are in the range between -0.50 and $0.52 \text{ t C ha}^{-1} \text{ yr}^{-1}$ with an average of $0.01 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for the inventory time period.

Applying a Tier 1 approach, changes in carbon stocks, in dead organic matter, and in mineral soils were assumed to be in equilibrium for Cropland remaining cropland.

The annual net carbon stock change in organic soils was estimated to $-9.52 \text{ t C ha}^{-1}$ according to measurements in Europe including Switzerland as compiled by Leifeld et al. (2003, 2005) and verified by ART (2009b).

In the case of land-use change, the net carbon changes in biomass and soil were calculated as described in chp. 6.1.3.

6.5.2.4 N₂O emissions from cropland

N₂O emissions from drainage of organic soils (category 4(II)) on cropland are reported in the agriculture sector (CRF Table3.D).

The calculation of emissions for categories 4(III) and 4(IV) (direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils) is described in chp. 6.10.

6.5.3 Uncertainties and time-series consistency

Uncertainties of activity data of category 4B Cropland are described in chp. 6.3.3. For calculating the overall uncertainty of category 4B, the relevant emissions from living biomass, mineral soils and organic soils were considered (Metecost 2017).

- Living biomass: The relative uncertainty in yield determination was estimated as 13% for biomass carbon from agricultural land (Leifeld and Fuhrer 2005). Data on biomass yields for different elevations and management intensities as published by FAL/RAC (2001) were based on many agricultural field experiments and have a high reliability. The absolute uncertainties per hectare, calculated with the implied emission factors of 2015, are: $0.065 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for 4B1 and $0.010 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for 4B2.
- Mineral soils: A range of possible carbon stock changes in mineral soils was determined by the Swiss Soil Monitoring Network (NABO). The upper and lower margin of the 95% confidence interval for carbon stock changes under cropland was $0 \pm 0.15 \text{ t C ha}^{-1} \text{ yr}^{-1}$ (see chp. 6.5.4). $0.15 \text{ t C ha}^{-1} \text{ yr}^{-1}$ was used as absolute uncertainty for 4B1 and 4B2.

- Organic soils: The uncertainty of the carbon stock change (emission factor) in organic soils is 23% as reported by Leifeld et al. (2003: 56) and the uncertainty of the activity data (area of organic soil) is 37.8% (see chp. 6.3.3), resulting in a combined uncertainty of 44.2%. Thus, the absolute uncertainties of the total organic soil emissions in 2015 are 42.93 kt C for 4B1 and 1.03 kt C for 4B2. By dividing those uncertainties with the total area of 4B1 and 4B2, respectively, the absolute uncertainties per hectare result in 0.116 t C ha⁻¹ yr⁻¹ for 4B1 and 0.046 t C ha⁻¹ yr⁻¹ for 4B2.

The root sum squares of the above-mentioned three absolute uncertainties are 0.200 t C ha⁻¹ yr⁻¹ for 4B1 and 0.157 t C ha⁻¹ yr⁻¹ for 4B2. These absolute uncertainties were used to calculate relative emission factor uncertainties for 4B1 and 4B2 by dividing with the mean net carbon stock change per hectare of 4B1 and 4B2, respectively. In 2015, the mean net carbon stock changes were -0.758 t C ha⁻¹ for 4B1 and -0.498 t C ha⁻¹ for 4B2 (calculated from CRF Table 4.B). The resulting relative uncertainties are 26.4% for 4B1 and 31.6% for 4B2, respectively (see Table 6-5).

Consistency: Time series for category 4B Cropland are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.5.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

Changes in living biomass

In 2012 an assessment of the appropriateness of the estimated pools of carbon in living biomass was conducted (ART 2012a). It came to the conclusion that almost all carbon stocks and carbon stock changes are in the range of the respective default values provided in the guidelines (IPCC 2006). Nevertheless, there is room for improvements. However, given the relatively low significance of the respective emissions a major effort in this area is hardly justified. Consequently, the biomass carbon pools will eventually be recalculated only in the course of the new planned Tier 3 approach for quantification of carbon stocks and carbon stock changes in agricultural soils (see chp. 6.5.6).

Changes in soil carbon stocks

The Tier 1 assumption that changes of carbon stocks in mineral soils are zero for Cropland remaining cropland was applied. In response to the ERT recommendation to develop appropriate methods to support this assumption (UNFCCC 2011: §94), the following evidence is provided: The SOC pools measured at 29 cropland monitoring sites of the Swiss Soil Monitoring Network (NABO; see chp. 6.4.4) featuring mineral soils indicate no significant changes from 1990 to 2014 (Figure 6-8). The decline from the first to the second sampling campaign was identified as artefact introduced by the date of sampling; in the first campaign, samplings were conducted substantially later in the year compared with the remaining campaigns, which induced higher SOC contents and thus SOC pools. The range of the calculated SOC pools was large (20.6–88.4 t C ha⁻¹) with a mean of 46.7 t C ha⁻¹.

The SOC pools for three additional cropland sites featuring organic soils ranged from 205 to 269 t C ha⁻¹ in the first and from 141 to 236 t C ha⁻¹ for the sixth sampling campaign (not included in Figure 6-8). Thus, SOC pools 0–20 cm of these sites declined by 14–63 t C ha⁻¹ over a period of 30 years (however, the effective losses over the whole soil profiles are even higher due to decreasing depths of soil layers). The monitoring scheme applied by NABO is able to detect relative changes in SOC contents of less than 5% per 10 years for mineral cropland soils (minimum detectable change for about 30 monitoring sites including three or more sampling campaigns). This corresponds to a minimum detectable change of roughly 0.15 t C ha⁻¹ yr⁻¹ for SOC pools.

The results provide evidence that Swiss cropland mineral soils did not act as a net carbon source or sink during the last 30 years.




Figure 6-8 Measured SOC pools for topsoils (0–20 cm) and their changes for 29 NABO long-term monitoring sites featuring mineral soils and used as cropland during the time period 1985–2014. The elevation of the sites ranges from 324 to 945 m a.s.l. Top panel: SOC pools 0–20 cm per site and sampling; the dots indicate the mean and the bars the range of 5% and 95% percentiles of bootstrap samples taking into account the variability in SOC contents of the individual replicates per site and sampling as well as the variations of the bulk density. Bottom panel: boxplot of changes in SOC pools since first sampling (boxes indicate the lower and the upper quartiles with the median indicated; lines include all observations inside the range of 1.5 times the interquartile distance, observations beyond that range are indicated as dots).

Short-term land-use changes in arable rotations

Short-term land-use changes between "grassland" and cropland are to be expected for leys in arable rotations. However, leys were allocated to cropland by the Swiss Land Use Statistics (AREA) and were thus not considered grasslands in the common sense (i.e. permanent grassland). Furthermore, only long-term changes between cropland and grassland are considered relevant for carbon stock changes in soils. Since only long-term

land-use changes are registered by the Swiss Land Use Statistics (AREA), carbon stock changes in soils associated with land-use changes between cropland and grassland and vice versa were adequately reported in the Swiss GHG inventory.

6.5.5 Category-specific recalculations

Activity data 1991–2014 were updated (see chp. 6.3.5).

Provisional carbon stocks and carbon stock changes in living biomass for the years 2013 and 2014 were replaced by definitive values from SBV (2016).

The provisional carbon stock change in living biomass for 1990 was replaced by the definitive value calculated from the difference 1989–1990 (cf. Table 6-24).

6.5.6 Category-specific planned improvements

Recently, Price et al. (2017) created a nationwide model for tree biomass in Switzerland (both inside and outside of forest), using structural information available from airborne laser scanning. The model offers significant opportunity for improved estimates of carbon stocks in living biomass on land use combination categories where tree biomass has either not been included or only roughly estimated until now. The suitability of the obtained data for reporting in category 4B will be evaluated.

Information on carbon stock changes in soils for cropland remaining cropland will become available from Agroscope research activities. A pilot study to evaluate possible Tier 3 methodological approaches for quantification of carbon stocks and carbon stock changes in agricultural soils was completed (Köck et al. 2013). Based on this, an evaluation of four soil carbon models was started in 2015, using data from Swiss long-term experiments. Model evaluation is scheduled to be finalized by the end of 2017.

Projects in the national research programme "Sustainable Use of Soil as a Resource" (<http://www.nfp68.ch/en/Pages/Home.aspx>) focus on (1) sustainable management of organic soils, (2) agricultural management and below ground carbon inputs, and (3) an integrated modelling framework to monitor and predict trends of agricultural management and their impact on soil functions at multiple scales. In the long term, results from these studies are expected to improve the reporting of emissions by sources and removals by sinks in category 4B.

6.6 Category 4C – Grassland

6.6.1 Description

Table 6-25 Key categories (KCA including LULUCF) in category 4C.

Code	IPCC Category	GHG	Identification Criteria
4C1	Grassland Remaining	CO ₂	L2, T2
4C2	Land converted to Grassland	CO ₂	L1, L2, T1, T2

Swiss grasslands belong to the cold temperate wet climatic zone.

Carbon stocks in living biomass and carbon stocks in mineral and organic soils were considered.

Grasslands were subdivided into permanent grassland (CC31), shrub vegetation (CC32), vineyards, low-stem orchards ('Niederstammobst') and tree nurseries (CC33), copse (CC34), orchards ('Hochstammobst'; CC35), stony grassland (CC36), and unproductive grassland (CC37) (see Table 6-2 and Table 6-6).

In category 2 in CRF Table 4.C, the land-use types CC32, CC33, CC34 and CC35 were merged under the notation 'woody' and CC36 and CC37 were merged under 'unproductive' (see Table 6-2).

In 2015, category 4C1 Grassland remaining grassland was a net source of 112.04 kt CO₂. Carbon stocks in living biomass and carbon stocks in mineral soils are assumed to be in balance (i.e. no carbon stock changes occur). The highest contribution to the CO₂ emissions of category 4C1 was thus generated by carbon mineralization in organic soils under permanent grasslands (CC31), although only 0.6% of CC31 soils in Switzerland are organic soils. Contributions of other Grassland remaining grassland categories (CC32-CC37) were of minor importance.

Category 4C2 Land converted to grassland was a net source of 216.34 kt CO₂ in 2015. The highest individual contribution came from subcategory 4C2.1 Forest land converted to grassland being responsible for a net source of 399.07 kt CO₂. Most of this source was due to net changes in living biomass from deforestation. Subcategories 4C2.2 to 4C2.5 were net sinks due to sequestration of CO₂ in mineral soils and biomass in the course of the conversion to grassland.

6.6.2 Methodological issues

6.6.2.1 Carbon in living biomass

Permanent grassland (CC31)

Permanent grasslands range in elevation from <300 m to 3000 m above sea level. Because both biomass productivity and soil carbon rely on the prevailing climatic and pedogenic conditions, grassland stocks were calculated separately for three elevation zones (corresponding to those used in category 4A Forest Land).

Standing stocks for permanent grasslands (t C ha⁻¹) were calculated as the annual cumulative yield of differentially managed grasslands (meadows, pastures, alpine pastures and meadows) based on FAL/RAC (2001; Table 6-26), assuming a carbon fraction of 0.5. Mean standing above-ground biomass stocks were taken for each of the altitudinal zones (see Table 6-27) because the spatial distribution of grassland management types is not known.

Table 6-26 Annual yields of different types of managed permanent grassland (CC31). Each value represents the mean of two fertilization levels (based on FAL/RAC 2001).

Management	Elevation [m]	Annual yield [t C ha ⁻¹]
Meadow	<601	5.88
	601-1200	4.38
	>1200	3.25
Pasture	<601	4.63
	601-1200	3.75
	>1200	2.75
Alpine pasture and meadow	601-1200	3.75
	>1200	0.75

Data for root biomass C were compiled by ART (2011a) based on published data of Swiss grassland. Carbon stocks in roots are in the range of 1.82–5.70 t C ha⁻¹ depending on elevation. Root biomass was added to above-ground biomass to derive the total living biomass for CC31. Table 6-27 shows the living biomass of permanent grassland for the three elevation zones as the the sum of cumulated annual yield (above-ground biomass) and roots.

Table 6-27 Root biomass (C_{root}), above ground biomass (C_{yield}) and total living biomass C_l of permanent grassland (CC31).

Elevation [m]	C _{root}	C _{yield}	C _l
	[t C ha ⁻¹]		
<601	1.82	5.26	7.08
601-1200	2.04	3.96	6.00
>1200	5.70	2.25	7.95

Shrub vegetation (CC32) and Copse (CC34)

Due to the lack of accurate data, the living biomass of shrub vegetation and copse was assumed to be equal to the living biomass of brush forest as described in chp. 6.4.2.8, where brush forest is assumed to contain 20.45 t C ha⁻¹ (Düggelin and Abegg 2011).

Vineyards, low-stem orchards and tree nurseries (CC33)

Low-stem orchards are small fruit trees distinguished from CC35 ('orchards') by a maximum stem-height of 1 m and a much higher stand density. Only low-stem orchards and vineyards are considered in the following because no stand densities for tree nurseries are available.

This is justified because tree nurseries comprise only ca. 8% (1'378 ha tree nurseries, SFSO 2002) of the total area of CC33, i.e., 17'054 ha, 15'436 ha vineyards (SFSO 2005) and 240 ha low-stem orchards (Widmer 2006).

The standing carbon stock of living biomass per ha (CI) for CC33 was therefore calculated as:

$$CI = [(CI \text{ vineyards} * \text{area vineyards}) + (CI \text{ low-stem orchards} * \text{area low-stem orchards})] / (\text{area vineyards} + \text{area low-stem orchards})$$

CI of vineyards is 3.61 t C ha⁻¹, calculated based on the mean stand density (5'556 vines ha⁻¹) and the mean carbon content in the woody biomass of one plant including roots (0.65 kg C; Ruffner 2005).

For small fruit trees on low-stem orchards, no literature value was found for biomass expansion factors. Therefore, the following assumptions were made: Diameter at breast height (DBH) of such trees was assumed to be 10 cm and the stem height was assumed to be 1 m. The bole shape of low-stem apple trees can be approximated by a cylinder shape.

$$\text{Stem wood volume} = r^2 * \pi * \text{height} = (5 \text{ cm})^2 * 3.1 * 100 \text{ cm} = 7.75 \text{ dm}^3$$

Based on expert knowledge (Kaufmann 2005), the percentage of branches was estimated as 100%, and the percentage of roots was estimated as 30% of the stem wood volume. This results in a BEF of 2.3. A wood density of 0.55 kg dm⁻³ (Vorreiter 1949) and the default IPCC carbon content of 50% (IPCC 2006, Volume 4, chp. 5.2.2.2) were assumed. With these assumptions the carbon content of a tree of the type low-stem ('Niederstamm') was calculated as follows:

$$\begin{aligned} \text{C low-stem} &= \text{stem wood volume} * \text{BEF} * \text{wood density} * \text{carbon content} \\ &= 7.75 \text{ dm}^3 * 2.3 * 0.55 \text{ kg/dm}^3 * 0.5 = 4.9 \text{ kg C} \end{aligned}$$

The mean stand density of low-stem orchards was estimated as 2500 ha⁻¹ (Widmer 2006), resulting in a CI of 12.25 t C ha⁻¹.

The resulting CI for CC33 is 3.74 t C ha⁻¹.

Orchards (CC35)

Orchards consists of larger fruit trees ('Hochstammobst') planted at a low density with grass understory. CI of orchards trees was calculated as:

$$\begin{aligned} \text{CI biomass} &= (\text{carbon per fruit tree [t C]} * \text{number of fruit trees} / \text{area orchards [ha]}) \\ &+ \text{carbon in grass [t C ha}^{-1}] \end{aligned}$$

The carbon content of a large fruit tree with a DBH of 25–35 cm was calculated as follows:

$$\text{C (Hochstamm)} = \text{Stem wood volume} * \text{KE-Factor} = 225 \text{ kg C}$$

where:

- Stem wood volume of an apple tree assuming a cylindrical stem with mean DBH of 30 cm and a stem height of 7 m amounts to 0.5 m³, and

- KE-Factor [t C m^{-3}] = BEF * Density * C-content = 0.45 (Wirth et al. 2004: 68, Table 16).

From the total fruit-growing area of 41'480 ha (SFSO 2005), the area of low-stem trees (240 ha, see CC33) was subtracted, and the remaining area of 41'240 ha was divided by the number of large fruit trees calculated as the mean of the counts in 1991 (3'616'301 trees) and 2001 (2'900'000 trees; SFSO 2002). This resulted in a mean stand density of 79 trees ha^{-1} . The resulting woody biomass of CC35 is thus 17.78 t C ha^{-1} . Because orchards typically have a grass understory, the biomass of CC31 was added to the woody biomass. The biomass of CC31 (cf. Table 6-27) was weighted with the area of CC35 in the three elevation zones (i.e. 6.84 t C ha^{-1}) and added to the woody biomass to obtain a total biomass stock of 24.62 t C ha^{-1} for CC35.

Stony grassland (CC36)

Approximately 35% of the surface of CC36 (herbs and shrubs on stony surfaces) is covered by vegetation. No accurate data were available for this category. Therefore, the carbon stock of brush forest (20.45 t C ha^{-1} ; cf. chp. 6.4.2.8; Düggelein and Abegg 2011) was multiplied by 0.35 to account for the 35% vegetation coverage. This results in a carbon stock for stony grassland of 7.16 t C ha^{-1} .

Unproductive grassland (CC37)

The category CC37 includes grass and herbaceous plants at watersides of lakes and rivers including dams and other flood protection structures, constructions to protect against avalanches and rock slides, and alpine infrastructure (e.g. for skiing). For none of these land-use types, biomass data are currently available. Therefore, the area-weighted mean of permanent grasslands biomass in the three elevation zones (cf. Table 6-27), 7.87 t C ha^{-1} , was assumed to be representative for the biomass on unproductive grassland CC37.

6.6.2.2 Carbon in soils

Permanent grassland (CC31)

Carbon stocks in grassland soil refer to a depth of 0–30 cm.

Soil carbon stocks in mineral soils under permanent grassland CC31 were calculated based on Leifeld et al. (2003, 2005). The approach correlates measured soil organic carbon stocks (t ha^{-1}) for permanent grasslands with soil texture and elevation after correction for soil depth and stone content. Area upscaling made use of the Swiss digital soil map (SFSO 2000a) and topography. Mean carbon stock values calculated for grasslands CC31 are given in Table 6-28. Their uncertainty is between 12% and 21%.

It should be noted that the current carbon stocks are not only the result of the conditions for productivity and carbon turnover under different land-use types, but are also determined by farmers' decisions to use a site in a specific way due to the demands of a crop or the suitability of a site, e.g. regarding machine use (see Leifeld et al. 2003: 65).

Table 6-28 Mean carbon stocks and their uncertainties under permanent grassland on mineral soils (0–30 cm).

Elevation [m]	Cs [t C ha ⁻¹]
<601	62.02 ± 13
601-1200	67.50 ± 12
>1200	75.18 ± 9

Soil carbon stocks in organic soils under permanent grassland were calculated based on Leifeld et al. (2003, 2005). The approach used measured carbon stocks in Swiss organic soils without differentiation between cropland and grassland. The mean soil organic carbon stock (0–30 cm) for organic soils is 240 ± 48 t C ha⁻¹ (uncertainty 20%).

Shrub vegetation (CC32)

Due to the lack of data, the values of CC31 (Table 6-28) were used as the mineral soil carbon stocks for this category (0–30 cm).

The mean soil organic carbon stock (0–30 cm) for organic soils is 240 t C ha⁻¹.

Vineyards, low-stem orchards and tree nurseries (CC33)

The category includes carbon stocks in soils of vineyards, low-stem orchards and tree nurseries. In accordance with the estimate of carbon stocks in biomass, only vineyards and low-stem orchards were considered. Both land-use types were assumed to have grass undercover. Therefore, the soil carbon stock may range between the values for grassland and cropland. It was decided to take the soil carbon content values of cropland, i.e. 53.40 t C ha⁻¹ (mineral soils, 0–30 cm) and 240 t ha⁻¹ (organic soils, 0–30 cm) for CC33 (see chp. 6.5.2.2).

Copse (CC34)

Due to the lack of data, the values of CC31 (Table 6-28) were used as the mineral soil carbon stocks for this category (0–30 cm).

The mean soil organic carbon stock (0–30 cm) for organic soils is 240 t C ha⁻¹.

Orchards (CC35)

The carbon stock in soils under orchards was calculated in accordance to the biomass calculation. No specific value for orchards was available, and the mean value of grassland mineral soil carbon stocks from the two lower elevation zones (i.e. 64.76 t C ha⁻¹; cf. Table 6-28) was taken for mineral soils (0–30 cm), and the value of 240 t ha⁻¹ for organic soils (0–30 cm).

Stony grassland (CC36)

Soil organic carbon stocks under herbs and shrubs on stony surfaces were calculated according to the procedure described in chp. 6.6.2.1, i.e. it was assumed that not more than 35% of the area of CC36 is covered with vegetation and thus only 35% of the area bears a mineral soil while the remainder is bare rock. Land use of this category mostly belongs to 'grassland' and 'unproductive land' and likely includes many of the former alpine grasslands (SFSO 2005). These grasslands are mainly located at elevations >1200m a.s.l. Thus, using the respective value from Table 6-28, the carbon stock of CC36 was calculated as:

$$Cs \text{ of CC36} = 0.35 * Cs \text{ permanent grassland } >1200 \text{ m} = 26.31 \text{ t C ha}^{-1} \text{ (0–30 cm)}$$

The mean soil organic carbon stock (0–30 cm) for organic soils is 240 t C ha⁻¹. It was assumed that the small area covered by organic soils in CC36 (cf. category 1 in CRF Table 4.C 'stony'), albeit entitled 'stony grassland', does not contain significant contributions from stones because bogs are free of stones as a matter of nature and fens usually contain, if any, only fine mineral sediments.

Unproductive grassland (CC37)

The combination category CC37 'unproductive grassland' includes grass and herbaceous plants at watersides of lakes and rivers including dams and other flood protection structures, constructions to protect against avalanches and rock slides, and alpine infrastructure (e.g. for skiing). For none of these land-use types, soil carbon stock data are currently available. In accordance with the procedure followed for biomass, the soil carbon stock of CC37 was assumed to correspond to the mean value of carbon stocks under permanent grassland on mineral soils for all elevation zones (Table 6-28). The carbon stock in soils under CC37 is thus 68.23 t C ha⁻¹.

The mean soil organic carbon stock (0–30 cm) for organic soils is 240 t C ha⁻¹.

6.6.2.3 Changes in carbon stocks

Applying a Tier 1 approach, changes in carbon stocks in biomass, in dead organic matter, and in mineral soils were assumed to be in equilibrium for Grassland remaining grassland.

The annual net carbon stock change in organic soils on managed grassland (CC31, CC33 and CC35) was estimated as -9.52 t C ha⁻¹ according to measurements in Europe including Switzerland as compiled by Leifeld et al. (2003, 2005) and verified by ART (2009b). For extensively managed grasslands (CC32, CC34, CC36 and CC37) the emission from organic soils was estimated as 5.30 t C ha⁻¹ yr⁻¹ according to available domestic data (ART 2011b).

In the case of land-use change, the net carbon changes in biomass and soil of CC31, CC32, CC33, CC34, CC35, CC36, and CC37 were calculated as described in chp. 6.1.3.

6.6.2.4 N₂O emissions from grassland

N₂O emissions from drainage of organic soils (category 4(II)) on grassland were reported in the agriculture sector (CRF Table3.D).

The calculation of emissions for categories 4(III) and 4(IV) (direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils) is described in chp. 6.10.

6.6.2.5 Emissions from wild fires

Data on wildfires affecting Swiss grassland are obtained from cantonal authorities and are compiled by the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL, Swissfire database, <http://www.wsl.ch/swissfire>). Table 6-29 shows the annual burnt area from 1990 to 2015. The Swissfire database differentiates between 'grassland' and 'unproductive land'. As 'unproductive land' can partially cover the grassland categories CC32, CC34, CC36 and CC37 the sum of both categories was reported. Controlled burning is not a common practice in Switzerland. Therefore, all fires were assigned to "wildfires".

The CH₄ and N₂O emissions were calculated using Equation 2.27 in Volume 4 of IPCC (2006) with the following parameters:

- For CH₄ the default emission factor of 2.3 g (kg combusted biomass)⁻¹ and for N₂O, the default emission factor of 0.21 g (kg combusted biomass)⁻¹ was applied (IPCC 2006, Volume 4, Table 2.5, Savanna and Grassland).
- The mass of "available fuel" encompasses the carbon stock of living biomass (litter and dead wood carbon stocks are zero for grassland). On average, the amount of living biomass amounted to 18.26 t biomass ha⁻¹ or, assuming a carbon fraction of 0.5, to a carbon stock of 9.13 t C ha⁻¹. This value was derived from the carbon stocks of all grassland categories (CC31 to CC37) as an area-weighted mean using the geographical extensions in 2015.
- The fraction of the biomass combusted was 0.74 (IPCC 2006 Volume 4, Table 2.6, Savanna and Grassland).

The resulting annual CH₄ and N₂O emissions 1990–2015 on burnt areas in category 4C Grassland are shown in Table 6-29 and are reported in CRF Table4(V).

Table 6-29 Area of 4C Grassland affected by wildfires (WSL, Swissfire database) and resulting CH₄ and N₂O emissions, 1990–2015.

Grassland	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Area burnt	ha	637.1	22.3	6.0	18.9	174.5	82.0	42.8	371.8	72.0	18.9
CH ₄	t	19.7	0.7	0.2	0.6	5.4	2.5	1.3	11.5	2.229	0.586
N ₂ O	t	1.8	0.1	0.0	0.1	0.5	0.2	0.1	1.1	0.203	0.054

Grassland	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Area burnt	ha	21.8	7.5	257.2	137.9	4.3	3.8	5.7	87.6	29.2	2.7
CH ₄	t	0.7	0.2	8.0	4.3	0.1	0.1	0.2	2.7	0.904	0.085
N ₂ O	t	0.1	0.0	0.7	0.4	0.0	0.0	0.0	0.2	0.083	0.008

Grassland	Unit	2010	2011	2012	2013	2014	2015
Area burnt	ha	1.3	56.3	4.4	3.4	2.3	3.3
CH ₄	t	0.0	1.7	0.1	0.1	0.1	0.1
N ₂ O	t	0.0	0.2	0.0	0.0	0.0	0.0

6.6.2.6 NMVOC emissions

Estimates for annual biogenic emissions of NMVOC (CRF Table4) for forests and natural grassland in Switzerland are available in SAEFL (1996a): the value for natural grassland (unproductive vegetation) is 0.51 kt.

6.6.3 Uncertainties and time-series consistency

Uncertainties of activity data of category 4C Grassland are described in chp. 6.3.3. For calculating the overall uncertainty of category 4C, the relevant emissions from living biomass, mineral soils and organic soils were considered (Meteotest 2017).

- Living biomass: The relative uncertainty in yield determination was estimated as 13% for biomass carbon from both cropland and grassland (Leifeld and Fuhrer 2005). Data on biomass yields for different elevations and management intensities as published by FAL/RAC (2001) were based on many agricultural field experiments and have a high reliability. The absolute uncertainties per hectare, calculated with the implied emission factors of 2015, are: 0.001 t C ha⁻¹ yr⁻¹ for 4C1 and 0.104 t C ha⁻¹ yr⁻¹ for 4C2.
- Mineral soils: A range of possible carbon stock changes in mineral soils was determined by the Swiss Soil Monitoring Network (NABO). The upper and lower margin of the 95% confidence interval for carbon stock changes under grassland was 0 ± 0.9 t C ha⁻¹ yr⁻¹ for a 20-years period (see chp. 6.6.4) or 0 ± 0.20 t C ha⁻¹ for one year (0.9/20^{0.5}=0.20). 0.20 t C ha⁻¹ yr⁻¹ was used as absolute uncertainty for 4C1 and 4C2.
- Organic soils: The uncertainty of the carbon stock change (emission factor) in organic soils is 23% as reported by Leifeld et al. (2003: 56) and the uncertainty of the activity data (area of organic soil) is 55.9% (see chp. 6.3.3), resulting in a combined uncertainty of 60.4%. Thus, the absolute uncertainties of the total organic soil emissions in 2015 are 32.22 kt C for 4C1 and 5.66 kt C for 4C2. By dividing those uncertainties with the total area of 4C1 and 4C2, respectively, the absolute uncertainties per hectare result in 0.025 t C ha⁻¹ yr⁻¹ for 4C1 and 0.074 t C ha⁻¹ yr⁻¹ for 4C2.

The root sum squares of the above-mentioned three absolute uncertainties are 0.203 t C ha⁻¹ yr⁻¹ for 4C1 and 0.238 t C ha⁻¹ yr⁻¹ for 4C2. These absolute uncertainties were used to

calculate relative emission factor uncertainties for 4C1 and 4C2 by dividing with the mean net carbon stock change per hectare of 4C1 and 4C2, respectively. In 2015, the mean net carbon stock changes were $-0.023 \text{ t C ha}^{-1}$ for 4C1 and $-0.776 \text{ t C ha}^{-1}$ for 4C2 (calculated from CRF Table 4.C). The resulting relative uncertainties are 867.9% for 4C1 and 30.7% for 4C2, respectively (see Table 6-5).

For wildfires, the emission factor uncertainties of CH_4 and N_2O were set to 70% (identical to forest land, see chp. 6.4.3).

Consistency: Time series for category 4C Grassland are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.6.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

Changes in living biomass

The assumption of a constant carbon stock in living biomass was reconsidered (UNFCCC 2007: §97). According to Schneider (2010) yields on meadows and pastures did not increase since 1990. Neither management nor the share of clover did significantly change over the past 20 years. Consequently, the current approach was reconfirmed.

In 2012 an assessment of the appropriateness of the estimated pools of carbon in living biomass was conducted (ART 2012a). It came to the conclusion that almost all carbon stocks and carbon stock changes are in the range of the respective default values provided in the guidelines (IPCC 2006). Nevertheless, there is room for improvements. However, given the relatively low significance of the respective emissions a major effort in this area is hardly justified. Consequently, the biomass carbon pools will eventually be recalculated only in the course of the new planned Tier 3 approach for quantification of carbon stocks and carbon stock changes in agricultural soils (see chp. 6.6.6).

Changes in soil carbon stocks

In response to the ERT recommendation to improve the documentation for the procedure to estimate changes in the SOC pool (UNFCCC 2007: §97), the following evidence is provided: The SOC pool measured at 33 grassland monitoring sites in the Swiss Soil Monitoring Network (NABO; see chp. 6.4.4) showed, on average, a slight increase during the period 1985 to 2000 and a slight decrease thereafter (Figure 6-9). SOC pools ranged between 20.9 and $183.2 \text{ t C ha}^{-1}$, the average SOC pool for the 33 grassland monitoring sites was 77.9 t C ha^{-1} (0–20 cm). Two alpine grassland sites above 1200 m a.s.l showed remarkable SOC pools of about 120 and 173 t C ha^{-1} (0–20 cm). The confidence interval of the mean SOC pool changes over time was $\pm 0.9 \text{ t C ha}^{-1}$. In total, the results of the soil monitoring data indicate that Swiss grassland mineral soils did not act as a net source or sink of carbon during the last 20 years.




Figure 6-9 Time series of measured SOC pool changes in the top soil (0–20 cm) at the 33 NABO grassland sites from the 1st to the 5th resampling campaigns. SOC pools were centred by the median SOC pool of all resamplings of the monitoring site. Each pool value presents the median of four bulked soil samples per campaign with measured SOC and bulk density. The error bars indicate the 25% and 75% percentiles resulting from the spatial variation of the sites and the errors along the measurement chain. The elevation of the grassland sites ranges between 265 and 2340 m a.s.l.

The slight increase and decrease will be subject of further detailed analysis. Partly, it may be attributed to natural variation of soil sampling (see chp. 6.4.4). The temporal change of the SOC content at grassland sites with intensive management and large manure application may also be related to the nitrogen input fluxes and nitrogen content in soil. Therefore, the total nitrogen content of the archived soil samples will be measured and the correlation to the SOC content analysed. Moreover, management data of the monitoring sites gathered directly from the farmers since 1985 permitted the calculation of annual nutrient fluxes of the sites (Keller et al. 2005). In this way temporal changes in nutrient management of the grassland sites can be separated from other effects that may cause temporal changes of SOC contents in grassland soils.

Short-term land-use changes between Grassland and Cropland

See chp. 6.5.4.

6.6.5 Category-specific recalculations

Activity data 1991–2014 were updated (see chp. 6.3.5).

The carbon stock in living biomass for orchards (CC35) and unproductive grassland (CC37) were recalculated for the time series 1990–2014 using area-weighted means instead of arithmetic means.

The amount of “fuel” for wildfires was recalculated for the time series 1990–2014 with the updated biomass carbon stocks.

6.6.6 Category-specific planned improvements

Recently, Price et al. (2017) created a nationwide model for tree biomass in Switzerland (both inside and outside of forest), using structural information available from airborne laser scanning. The model offers significant opportunity for improved estimates of carbon stocks in living biomass on land use combination categories where tree biomass has either not been included or only roughly estimated until now. The suitability of the obtained data for reporting in category 4C will be evaluated.

Information on carbon stock changes in soils for grassland remaining grassland will become available from Agroscope research activities. A pilot study to evaluate possible Tier 3 methodological approaches for quantification of carbon stocks and carbon stock changes in agricultural soils (including meadows and pastures) was completed (Köck et al. 2013). Based on this, an evaluation of four soil carbon models was started in 2015, using data from Swiss long-term experiments. Model evaluation is scheduled to be finalized by the end of 2017.

A study on GHG emissions from an extensively used fen under grassland management (Agroscope in collaboration with the University of Basel, 2013–2019, financed by FOEN) will improve the robustness of country-specific emission factor estimates for grassland soils rich in organic matter in the medium term. A further objective of the study is to estimate emission reductions by rewetting.

Projects in the national research programme ("[Sustainable Use of Soil as a Resource](#)") focus on (1) sustainable management of organic soils, (2) agricultural management and below ground carbon inputs, and (3) an integrated modelling framework to monitor and predict trends of agricultural management and their impact on soil functions at multiple scales. In the long term, results from these studies are expected to improve the reporting of emissions by sources and removals by sinks in category 4C.

6.7 Category 4D – Wetlands

6.7.1 Description

Table 6-30 Key categories (KCA including LULUCF) in category 4D.

Code	IPCC Category	GHG	Identification Criteria
4D1	Wetland remaining Wetland	CO2	L2

Carbon stocks in living biomass and carbon stocks in mineral and organic soils were considered.

Wetlands were subdivided into surface waters (CC41) and unproductive wetlands such as shore vegetation, fens or (raised) bogs (CC42) (see Table 6-2 and Table 6-6).

6.7.2 Methodological issues

6.7.2.1 Carbon in living biomass

Surface waters (CC41)

Surface waters have no carbon stocks by definition.

Unproductive wetland (CC42)

CC42 consists of (very) extensively managed grassland, bushes or tree groups. The pool of living biomass was estimated as 6.50 t C ha⁻¹ (Mathys and Thürig 2010).

6.7.2.2 Carbon in soils

The soil carbon stock for surface waters (CC41) is zero. However, for CC41 situated in areas with organic soil (see chp. 6.2.2 and Table 6-7), a soil carbon stock of 240 t C ha⁻¹ (0–30 cm) was assumed. These surface waters were assumed to be shallow ponds as integrated parts of fens or bogs.

Land cover in CC42 includes bogs and fens protected by Federal Legislation (Swiss Confederation 1991a, 1994) as well as reed. More than 10% of the unproductive wetlands are located on organic soils (cf. Table 6-7). In this case the carbon stock in soils is 240 t C ha⁻¹ (0–30 cm). Currently, no specific soil data are available for CC42 on mineral soils. As a first approximation, it was assumed that the soil carbon stock of unproductive wetlands is similar to unproductive grassland (CC37) on mineral soils (mean value: 68.23 t C ha⁻¹; 0–30 cm).

6.7.2.3 Changes in carbon stocks

Applying a Tier 1 approach, changes in carbon stocks in biomass, in dead organic matter, and in mineral soils were assumed to be in equilibrium for Wetlands remaining wetlands.

The emission from organic soils under CC41 was assumed to be zero because the respective areas are not drained.

The emission from organic soils under CC42 was estimated to be 5.30 t C ha⁻¹ yr⁻¹ according to domestic data (ART 2011b). This value was used for weakly managed ecosystems such as fens and (very) extensively managed ecosystems such as raised bogs. Bogs and fens are protected to a large part by Federal Ordinances (Swiss Confederation 1991a, 1994) and drainage is not allowed. However, the impact of old drainages constructed before 1990 probably leads to a certain emission.

In the case of land-use change, the net carbon changes in biomass and soil of both CC41 and CC42 were calculated as described in chp. 6.1.3.

For land converted to unproductive wetland (CC42) a conversion time of one year was chosen for the carbon stock change in living biomass and in dead organic matter (see Table 6-3). This assumption was reviewed as recommended by the ERT (UNFCCC 2009: §82), and

it was concluded that it is justified because the exact year of the land use change is not known (cf. chp. 6.3.2). For carbon stock changes in soil the conversion time is 20 years.

6.7.2.4 Non-CO₂ emissions from wetlands

An estimate of 0.43 kt CH₄ yr⁻¹ emitted by reservoirs (flooded lands) was given by Hiller et al. (2014). The estimate encompasses 97 artificial lakes covering a total area of 10.6 kha. This emission is reported in category D.2 in CRF Table4(II).

N₂O emissions from drainage of organic soils was calculated for unproductive wetlands (CC42) and reported in category D.3 in CRF Table4(II). The emission factor of 1.6 kg N₂O-N ha⁻¹ used is the default value given in the IPCC Wetlands Supplement (IPCC 2014a, Table 2.5) for shallow drained, nutrient-rich grassland.

The calculation of emissions for categories 4(III) and 4(IV) (direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils) is described in chp. 6.10.

6.7.3 Uncertainties and time-series consistency

Uncertainties of activity data of category 4D Wetlands are described in chp. 6.3.3.

For calculating the overall uncertainty of 4D1, only the relevant emissions from organic soils were considered (Meteotest 2017).

- Organic soils: The uncertainty of the carbon stock change (emission factor) in organic soils is 72.2% calculated on the basis of measurement data compiled by ART (2011b). The uncertainty of the activity data (area of organic soil) is 61.9% (see chp. 6.3.3).
- For calculating the overall uncertainty of 4D2, the relevant emissions from living biomass, mineral soils and organic soils were considered (Meteotest 2017).
- Living biomass: The relative uncertainty in yield determination was estimated as 13% for biomass carbon from both cropland and grassland (Leifeld and Fuhrer 2005). Data on biomass yields for different elevations and management intensities as published by FAL/RAC (2001) were based on many agricultural field experiments and have a high reliability. The absolute uncertainty per hectare, calculated with the implied emission factors of 2015, is 0.132 t C ha⁻¹ yr⁻¹ for 4D2.
- Mineral soils: Based on expert judgement, a value of 50% was chosen for the emission factor uncertainty in 4D2. The absolute uncertainty per hectare, calculated with the implied emission factor of 2015, is 0.394 t C ha⁻¹ yr⁻¹ for 4D2.
- Organic soils: The uncertainty of the carbon stock change (emission factor) in organic soils is 72.2% calculated on the basis of measurement data compiled in ART (2011b) and the uncertainty of the activity data (area of organic soil) is 61.9% (see chp. 6.3.3), resulting in a combined uncertainty of 95.1%. Thus, the absolute uncertainties of the total organic soil emissions in 2015 are 0.857 kt C for 4D2. By dividing this uncertainty with the total area of 4D2, the absolute uncertainty per hectare results in 0.198 t C ha⁻¹ yr⁻¹ for 4D2.

The root sum squares of the above-mentioned three absolute uncertainties is $0.460 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for 4D2. This absolute uncertainty was used to calculate the relative emission factor uncertainty for 4D2 by dividing with the mean net carbon stock change per hectare of 4D2. In 2015, the mean net carbon stock change was $-2.148 \text{ t C ha}^{-1}$ for 4D2 (calculated from CRF Table 4.D). The resulting relative uncertainties is 21.4% for 4D2 (see Table 6-5).

The uncertainty for CH_4 emitted by flooded lands can be very high (IPCC 2006, Volume 4, Appendix 3). As a best guess, a value of 70% was chosen for the emission factor of 4(II)D2 (cf. Table 6-5).

For N_2O emissions from drainage of organic soils (4(II)D3) the uncertainties for agricultural soils given in the 2006 IPCC Guidelines was used. The arithmetic mean of the lower and upper bound uncertainty (137.5%) was used for the emission factor (see also chp. 5.5.3).

Consistency: Time series for category 4D Wetlands are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.7.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

No category-specific QA/QC activities were carried out.

6.7.5 Category-specific recalculations

- Activity data 1991–2014 were updated (see chp. 6.3.5).

6.7.6 Category-specific planned improvements

Recently, Price et al. (2017) created a nationwide model for tree biomass in Switzerland (both inside and outside of forest), using structural information available from airborne laser scanning. The model offers significant opportunity for improved estimates of carbon stocks in living biomass on land use combination categories where tree biomass has either not been included or only roughly estimated until now. The suitability of the obtained data for reporting in category 4D will be evaluated.

6.8 Category 4E – Settlements

6.8.1 Description

Table 6-31 Key categories (KCA including LULUCF) in category 4E.

Code	IPCC Category	GHG	Identification Criteria
4E2	Land converted to Settlements	CO ₂	L1, L2

Carbon stocks in living biomass and carbon stocks in mineral and organic soils were considered.

Settlements were subdivided into buildings/constructions (CC51), herbaceous biomass in settlements (CC52), shrubs in settlements (CC53), and trees in settlements (CC54) (see Table 6-2 and Table 6-6).

Switzerland decided not to prepare estimates of GHG emissions by biomass burning in settlements (4(V)E) (not mandatory).

N₂O emissions associated with inputs from N (fertilizers, crop residues) in Settlements (4(I)E; see chp. 5.5) were included in the Agriculture sector.

6.8.2 Methodological issues

6.8.2.1 Carbon in living biomass

Buildings and constructions (CC51)

By default, buildings/constructions have no carbon stocks.

Herbaceous biomass, shrubs and trees in settlements (CC52, CC53, CC54)

Carbon stocks in living biomass are: 9.54 t C ha⁻¹ for CC52, 15.43 t C ha⁻¹ for CC53, and 20.72 t C ha⁻¹ for CC54 (Mathys and Thürig 2010: Table 7).

6.8.2.2 Carbon in soils

The carbon stocks in mineral and organic soils for CC51 (buildings and construction) were assumed to be zero.

In case of land-use changes to CC51 or from CC51 on mineral or organic soils, 50% of the difference between carbon stocks before and after the change was reported as emission and removal, respectively. The reason for this is that the soil organic matter on construction sites is stored temporarily and is later used for replanting the surroundings or for vegetating dumps, for example. According to paragraph 7 of the "Ordinance against deterioration of soils" (Swiss Confederation 1998) the soil material excavated on a construction site must be treated in such a way that it can be used as a soil again. When the material is re-used (e.g. for re-cultivations) the fertility of the soil must not be affected. This regulation ensures that a large part of the soil organic matter is preserved on land converted to and from CC51.

Prior to 1998 there were federal acts and good practice guidance for engineers focusing on physical soil protection and on preserving soil fertility. The Ordinance against pollution of soils (Swiss Confederation 1986) was in force between 1986 and 1998 (without a specific focus on the soil material on construction sites). As a legal basis the Federal Act on the Protection of the Environment (Swiss Confederation 1983) formulates the aim of preventing soil pollution ("physical, chemical and biological modification of the natural condition of the soil") and of preserving the natural foundations of life sustainably, in particular biological diversity and the fertility of the soil. The act and concomitant ordinances form a legal framework for prosecuting violators. It is very likely that also prior to the latest environmental legislation the protection measures at construction sites were applied in an appropriate way as the awareness for soil fertility has been traditionally high in Switzerland. A good practice

guidance of the Swiss Society of Engineers and Architects on gardening and landscaping operations may serve as example (SIA 1988).

Switzerland chose the factor 0.5 (i.e. 50% stabilised SOC fraction which is not likely to be oxidised in the medium term) to reflect this domestic soil protection measure (see discussion in Leifeld et al. 2003: 67). Thus, the equation 6.8 presented in chp. 6.1.3.2 was adjusted as follows if $a=CC51$ or $b=CC51$:

$$\Delta C_{s,i,ba} = [0.5 * (\text{stock}_{Cs,i,a} - \text{stock}_{Cs,i,b}) / CT] * A_{i,ba}$$

where:

a	land-use category after conversion (CC = a)
b	land-use category before conversion (CC = b)
ba	land use conversion from b to a
i	spatial stratum
$A_{i,ba}$	area of land (ha) converted from b to a in the spatial stratum i (the sum of the areas converted within the last 20 years)
CT	conversion time (20 years), see Table 6-3.

The carbon stock in mineral soils for CC52, CC53, and CC54 is 53.40 t C ha⁻¹ (0–30 cm). This is the same value as for cropland.

For organic soils the carbon stock for CC52, CC53, and CC54 was assumed as 240 t C ha⁻¹ (0–30 cm).

6.8.2.3 Changes in carbon stocks

Applying a Tier 1 approach, changes in carbon stocks in biomass, in dead organic matter, and in mineral soils were assumed to be in equilibrium for Settlements remaining settlements.

On organic soils, the following emission factors were applied:

- 9.52 t C ha⁻¹ yr⁻¹ for CC52. This corresponds to the value used for cropland because CC52 areas are managed (gardens, parks) (Leifeld et al. 2003, 2005 and verified by ART 2009b).
- 5.30 t C ha⁻¹ yr⁻¹ for CC53 and CC54. This corresponds to the value used for extensively managed grasslands (ART 2011b).

In the case of land-use change, the net carbon changes in biomass and soil of CC51, CC52, CC53, and CC54 were calculated as described in chp. 6.1.3 and 6.8.2.2.

6.8.2.4 N₂O emissions from settlements

The calculation of emissions for categories 4(III) and 4(IV) (direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils) is described in chp. 6.10.

6.8.3 Uncertainties and time-series consistency

Based on expert judgement, a value of 50% was chosen for the emission factor uncertainty in category 4E (Table 6-5).

Uncertainties of activity data of category 4E Settlements are described in chp. 6.3.3.

Consistency: Time series for category 4E Settlements are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.8.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

No category-specific QA/QC activities were carried out.

6.8.5 Category-specific recalculations

- Activity data 1991–2014 were updated (see chp. 6.3.5)

6.8.6 Category-specific planned improvements

Recently, Price et al. (2017) created a nationwide model for tree biomass in Switzerland (both inside and outside of forest), using structural information available from airborne laser scanning. The model offers significant opportunity for improved estimates of carbon stocks in living biomass on land use combination categories where tree biomass has either not been included or only roughly estimated until now. The suitability of the obtained data for reporting in category 4E will be evaluated.

In response to UNFCCC (2017/ID#KL.4), the assumption that only 50% of the difference between the carbon stocks before and after the change is reported as a source or sink, respectively, for afforestation and deforestation will be reviewed.

6.9 Category 4F – Other land

6.9.1 Description

Category 4F – Other land is not a key category.

As shown in Table 6-2 and in Table 6-6 Other land (CC61) covers non-vegetated areas such as glaciers, rocks and shores.

6.9.2 Methodological issues

By definition, Other land has no carbon stocks. Coherently, changes in carbon stock in biomass, in dead organic matter, and in soils were assumed to be zero for Other land remaining other land.

In the case of land converted to other land, the net carbon changes in biomass and soil were calculated as described in chp. 6.1.3.

The calculation of emissions on land converted to other land for categories 4(III) and 4(IV) (direct N₂O emissions from nitrogen mineralization in mineral soils and indirect N₂O emissions from managed soils) is described in chp. 6.10.

6.9.3 Uncertainties and time-series consistency

Based on expert judgement, a value of 50% was chosen for the emission factor uncertainty in subcategory 4F2 (Table 6-5).

Uncertainties of activity data of category 4F Other Land are described in chp. 6.3.3.

Consistency: Time series for category 4F Other land are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.9.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

No category-specific QA/QC activities were carried out.

6.9.5 Category-specific recalculations

- Activity data 1991–2014 were updated (see chp. 6.3.5).

6.9.6 Category-specific planned improvements

No category-specific improvements are planned.

6.10 Categories 4(III) and 4(IV) – N₂O emissions

6.10.1 Description

Table 6-32 Key categories (KCA including LULUCF) in category 4F.

Code	IPCC Category	GHG	Identification Criteria
4III	Direct N ₂ O from Disturbance	N ₂ O	L2

This chapter presents the methods for calculating direct (4(III)) and indirect (4(IV)) N₂O emissions from nitrogen (N) mineralization in mineral soils. The source of nitrogen is N mineralization associated with loss of soil organic matter resulting from change of land use or management of mineral soils. As the approaches applied were not Tier 3, no N₂O immobilization is reported.

- In category 4(III), direct N₂O emissions from nitrogen mineralization associated with loss of soil organic matter are reported.
- In category 4(IV)2, indirect emissions of N₂O due to nitrogen leaching and run-off after mineralization of soil organic matter are reported.

The following N₂O emissions were included in the Agriculture sector:

- N₂O emissions on grassland remaining grassland (4(III)C1, see chp. 5.5). In Switzerland, grassland is regarded to be under agricultural management.
- Indirect N₂O emissions due to atmospheric deposition (4(IV)1; all land uses) and leaching and run-off on agricultural land (4(IV)2; Cropland remaining cropland and Grassland remaining grassland) (see chp. 5.3).

6.10.2 Methodological issues

Direct N₂O emissions

Direct N₂O emissions (category 4(III)) as a result of the disturbance of mineral soils associated with change of land use or management of mineral soils were calculated according to IPCC (2006, Vol. 4, chp. 11):

$$\text{Emission(N}_2\text{O)} = -\text{deltaCs} * 1 / (\text{C:N}) * \text{EF1} * 44 / 28, \text{ if deltaCs} < 0 \quad [\text{kt N}_2\text{O}]$$

where:

- deltaCs: soil carbon change induced by land-use change [kt C]
- C:N: C to N ratio of the soil before the land-use change
- EF1: default emission factor = 0.01 kg N₂O-N (kg N)⁻¹, IPCC 2006 (Volume 4, Table 11.1)

deltaCs was calculated according to the methodology described in chp. 6.1.3.2. If deltaCs is zero or positive (carbon gain) there are no N₂O emissions provoked by a land-use change.

The value of the C:N ratio is related to the land-use category before the change. For cropland and grassland the ratio is 9.8 according to Leifeld et al. (2007). This value was also used for the mineral soils in wetlands (CC42) and unsealed settlement areas (CC 52, CC53, CC54). For forest land, the default value of C:N=15 was used (IPCC 2006, Volume 4, Equation 11.8).

Indirect N₂O emissions

The indirect N₂O emissions (4(IV)) as a result of N leaching and run-off after mineralization of soil organic matter were calculated as follows using default emission factors (IPCC 2006, Volume 4, Table 11.3):

$$\text{Emission(N}_2\text{O)} = -\text{deltaCs} * \text{Fra}_{\text{CLEACH}} / (\text{C:N}) * \text{EF5} * 44 / 28, \text{ if deltaCs} < 0 \quad [\text{kt N}_2\text{O}]$$

where:

- Fra_{CLEACH} : fraction of mineralized N lost by leaching or run-off, $Fra_{CLEACH}=21.8\%$. (The corresponding value in the agriculture sector was recently updated (17.8%), see chp. 5.5.2.4.2).
- EF5: default emission factor = $0.0075 \text{ kg N}_2\text{O-N (kg N)}^{-1}$, IPCC 2006 (Volume 4, Table 11.3)

If ΔC_s is zero or positive (carbon gain) there are no N_2O emissions.

For calculating ΔC_s , all land-use changes and conversions between land-use subcategories were taken into account.

On productive forest land (CC12), small annual changes in carbon contents of mineral soils are reported that were calculated with the Yasso07 model (chp. 6.4.2.7). These changes were deliberately not considered for the calculation of N_2O emissions in categories 4(III) and 4(IV) as they are not associated with a land-use change or any change in management.

6.10.3 Uncertainties and time-series consistency

Relative uncertainties for the emission factors were estimated as the mean of the upper and the lower limit of the uncertainty ranges listed in IPCC (2006), Vol 4, Tables 11.1 and 11.3 (Table 6-5):

- Uncertainty (EF1): 135%
- Uncertainty (EF5): 162%

The uncertainty of the activity data for category 4(III) corresponds to the uncertainty of land converted to cropland or land converted to grassland, i.e. 5.3% (Table 6-10).

The uncertainty of the activity data for category 4(IV)2 was adopted from ART (2008a): 20%.

Consistency: Time series for categories 4(III) and 4(IV) N_2O emissions from nitrogen mineralization are all considered consistent; they were calculated based on consistent methods and homogenous databases.

6.10.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

No category-specific QA/QC activities were carried out.

6.10.5 Category-specific recalculations

- Activity data 1991–2014 were updated (see chp. 6.3.5).

6.10.6 Category-specific planned improvements

The fraction of mineralized N lost by leaching or run-off (Fra_{CLEACH}) used for calculating N_2O emissions in category 4(IV) will be harmonised with the updated value in the agricultural sector (see chp. 5.5.2.4.2).

6.11 Category 4G – Harvested wood products (HWP)

6.11.1 Description

Table 6-33 Key categories (KCA including LULUCF) in category 4G.

Code	IPCC Category	GHG	Identification Criteria
4G	HWP Harvest Wood Prod		T1, T2

The data presented in this chapter are estimates of net emissions and removals from HWP due to changes in the HWP carbon pool.

The approach to calculate carbon stock changes in HWP can be generally characterized as a production approach as described in chp. 12, Volume 4 of IPCC (2006). Changes in carbon stocks in Swiss forests are presented in chp. 6.4. Changes in the wood products pool containing products made from wood harvested in Switzerland are described in this chapter. The wood products pool also includes products made from domestic harvest that were exported and are in use in other countries.

To calculate carbon stock changes in HWP, product categories and half-lives were used following the methodologies described in IPCC (2006) and IPCC (2014).

6.11.2 Methodological issues

The same methodology was used for reporting under the UNFCCC and accounting under the Kyoto Protocol for HWPs in Switzerland consistent with paragraph 29 in the Annex to Decision 2/CMP.7, which states that “transparent and verifiable activity data for harvested wood products categories are available, and accounting is based on the change in the harvested wood products pool of the second commitment period, estimated using the first-order decay function”. Therefore, in this chapter the terminology of the Kyoto Protocol is used, i.e. it is referred to the activities Afforestation, Deforestation and Forest management (as defined in FOEN 2006h; see chp. 11.1.3).

For the estimation of carbon stocks and carbon stock change, the equations described in IPCC (2014) were used.

A Tier 1 approach (instantaneous oxidation) was applied to the product category paper because the available statistical data are poor due to uncertainties in i) the estimate of the proportion of the national harvest used for producing paper, ii) the use of recycled products, and due to the fact that the share of this product was considered too negligible to justify the (financial) effort to collect higher quality data (see Figure 4.1 in Volume 1 of IPCC 2006).

A Tier 2 approach, first order decay, was applied to the product categories panels and sawnwood, according to equation 2.8.5 in IPCC (2014).

- Emissions occurring during the second commitment period from HWPs removed from forests prior to the start of the second commitment period were also accounted for. The starting year used to estimate the delayed emissions from the existing pools is 1900.
- Emissions from the HWP pool accounted for in the first commitment period on the basis of instantaneous oxidation were excluded from the accounting for the second commitment period (FOEN 2016j).
- Emissions from HWP in solid waste disposals and from wood harvested from energy purposes were accounted on the basis of instantaneous oxidation (FOEN 2016j).
- Exported HWP were included in the calculation (notation key "IE" in CRF table 4(KP-I)C).
- The feedstock from domestic harvest was calculated according to equation 2.8.1 in IPCC (2014). Exported round wood was not included in the calculation of HWP.
- Based on the available datasets it was not possible to differentiate between HWP from Afforestation and HWP from Forest management. Since tree diameters (DBH) of the majority of trees on areas afforested since 1990 are small, the amount of HWP from Afforestation is not existing or negligibly small. Therefore, all carbon stock changes in HWPs were reported under Forest management.
- The change in carbon stocks of HWPs was estimated separately for each product category and differentiating HWPs from Deforestation and from Forest management including HWP from Afforestations by applying equation 2.8.4 in IPCC (2014). Applying instantaneous oxidation to HWPs originating from Deforestation, the same results were obtained for changes in carbon stocks of HWP reported under the UNFCCC (CRF Table4.Gs1) and under the Kyoto Protocol (CRF table 4(KP-I)C).

6.11.2.1 Activity data

Activity data for production of HWPs for the product categories sawnwood and wood panels were based on national statistics (FOEN 2016k) and FAO forest product statistics (Forestry Production and Trade; <http://faostat3.fao.org/download/F/FO/E>). They are described in detail in FOEN (2016j). The time series are shown in the CRF Table4.Gs2. For the second commitment period, activity data were restricted exclusively to Swiss national statistics.

Data on production of sawnwood was derived from national saw mill statistics (FOEN 2016k). Data on production of wood panels was derived from surveys of the processing industries (FOEN 2016k). In order to estimate the share of industrial roundwood originating from domestic forests, as feedstock for HWP production, data from National Wood Use statistic and National Foreign Trade statistics were used (FOEN 2016k).

In order to estimate carbon amounts in each HWP category and subcategory, default conversion factors were taken from IPCC (2014; Table 2.8.1).

6.11.2.2 Emission factors

Emission factors for specific product categories were calculated with default half-lives of 25 years for wood panels and 35 years for sawn wood (IPCC 2014; eq. 2.8.5).

6.11.2.3 Results

Emissions and removals per product category are listed in Table 6-34. Figure 6-10 shows the resulting CO₂ emissions and removals.

Table 6-34 Total CO₂ emissions and removals from Harvested wood products (changeC_{HWP}) derived from sawnwood (changeC_{HWP-sawnwood}) and panels (changeC_{HWP-panels}), 1990–2015, originating from Forest management, in kt CO₂ yr⁻¹ (positive values refer to emissions, negative values refer to removals). HWPs originating from Deforestation were calculated using instantaneous oxidation.

Greenhouse gas emissions and removals	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	Net CO ₂ emissions/removals (kt CO ₂ yr ⁻¹)									
changeC _{HWP}	-1'230.8	-941.7	-774.3	-630.5	-477.3	-560.9	-396.1	-313.8	-423.5	-479.0
changeC _{HWP-sawnwood}	-751.1	-526.1	-326.9	-224.9	-153.8	-312.0	-191.2	-135.2	-226.7	-297.8
changeC _{HWP-panels}	-479.7	-415.5	-447.4	-405.6	-323.5	-248.9	-204.9	-178.6	-196.8	-181.1

Greenhouse gas emissions and removals	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	Net CO ₂ emissions/removals (kt CO ₂ yr ⁻¹)									
changeC _{HWP}	-834.5	-577.0	-442.5	-440.7	-636.7	-763.1	-625.8	-730.7	-528.2	-370.4
changeC _{HWP-sawnwood}	-427.5	-213.4	-149.0	-133.0	-272.4	-342.4	-327.7	-287.1	-233.8	-179.4
changeC _{HWP-panels}	-407.0	-363.6	-293.6	-307.7	-364.3	-420.7	-298.1	-443.6	-294.4	-191.0

Greenhouse gas emissions and removals	2010	2011	2012	2013	2014	2015
	Net CO ₂ emissions/removals (kt CO ₂ yr ⁻¹)					
changeC _{HWP}	-411.3	-257.7	-167.7	-160.9	-106.0	-69.5
changeC _{HWP-sawnwood}	-170.8	-70.0	45.0	141.1	36.0	62.5
changeC _{HWP-panels}	-240.5	-187.8	-212.7	-302.0	-142.0	-132.0

Fluctuations in the HWP pool can mainly be attributed to changes in the production of sawnwood (see Table 6-34), which correlates strongly with the domestic harvesting rate.




Figure 6-10 CO₂ emissions and removals from category 4G Harvested wood products originating from Forest management (KP), 1990–2015 (in kt CO₂; positive values refer to emissions, negative values refer to removals).

6.11.3 Uncertainties and time-series consistency

For category 4G HWP, the following information on relative uncertainty was used:

- Activity data:
 - Roundwood harvest: 5% (national activity data from the Swiss Forestry Statistics, annual complete survey) HWP Production:
 - Sawnwood: 5% for activity data prior to 1990 and 2% for activity data since 1990 (national activity from survey on wood processing in sawmills, combined survey, FOEN 2016k)
 - Wood Panels: 4% for activity data prior to 1990 and 2% for activity data since 1990 (national activity from survey in the wood industry, FOEN 2016k).
- Conversion factors:
 - Wood density: 25% (default from IPCC 2006, Volume 4, Table 12.6)
 - Carbon contents in wood products: 10% (Lamlom and Savidge 2003, assessment of carbon content in wood); IPCC 2006, Volume 4, Table 12.6)
 - Emission factors (half-life estimates): 50% (default from IPCC 2006, Volume 4, Table 12.6).

The relative uncertainty of the emission factors for HWP amounts to 57%:

$$U_{\text{HWP EF}} = \sqrt{25^2 + 10^2 + 50^2} = 57\%$$

The total relative uncertainty of carbon losses and gains in HWP was thus calculated as:

$$U_{\text{HWP Sawnwood}} = \sqrt{5^2 + 5^2 + 25^2 + 10^2 + 50^2} = 57\%$$

$$U_{\text{HWP Panels}} = \sqrt{5^2 + 4^2 + 25^2 + 10^2 + 50^2} = 57\%$$

Consistency: Time series for category 4G Harvested wood products are all considered consistent; they were calculated based on consistent methods and homogenous databases (FOEN 2016j).

6.11.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

No category-specific QA/QC activities was carried out.

6.11.5 Category-specific recalculations

The activity data were updated for the whole time series. More specifically, the share of domestic to total HWP was revised.

Net CO₂ emissions and removals from Harvested wood products (HWP) Deforestations (KP) were accounted for on the basis of instantaneous oxidation.

6.11.6 Category-specific planned improvements

In response to UNFCCC (2017/ID#KL.8), the available data from the FAOSTAT database will be used to calculate the contribution of paper and paperboard to the HWP pool and report in the next submission. However, for the calculation there are some methodological challenges, such as how to determine the amount of domestic pulp wood contained in recycled products. Therefore, a study will be carried out to obtain additional national activity data on paper and paperboard to improve the calculation of the contribution of this pool.

7 Waste

7.1 Overview

7.1.1 Greenhouse gas emissions

Within sector 5 Waste, emissions from five source categories are considered:

- 5A Solid waste disposal
- 5B Biological treatment of solid waste
- 5C Incineration and open burning of waste
- 5D Wastewater treatment and discharge
- 5E Other (no direct GHG emissions, but indirect GHG emissions reported in chp. 9)




Figure 7-1 Switzerland's greenhouse gas emissions from sector 5 Waste 1990–2015. There are no direct greenhouse gas emissions from sector 5 E Other.

The total greenhouse gas emissions from sector 5 Waste show a decrease within the reporting period. 5A Solid waste disposal and 5D Wastewater treatment and discharge are the two dominant source categories.

Table 7-1 Trend of total GHG emissions from sector 5 Waste in Switzerland 1990–2015.

Gas	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
CO ₂	54	30	19	12
CH ₄	939	775	759	785
N ₂ O	140	144	156	167
Sum	1133	950	934	964

Gas	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	CO ₂ equivalent (kt)									
CO ₂	13	12	12	11	11	11	10	10	10	10
CH ₄	782	764	745	724	709	692	677	665	655	643
N ₂ O	170	173	176	180	184	187	181	189	190	193
Sum	965	949	933	915	904	890	869	864	855	846

CH₄ is the most important greenhouse gas in sector 5 Waste over the whole reporting period. Nevertheless, CH₄ emissions have decreased. Two processes determine the trend: a dominating decreasing trend from 5A Solid waste disposal and a distinct increasing trend from 5B Biological treatment of solid waste (refer to

Figure 7-3 and Table 7-14).

The relative trends of the gases are shown in Figure 7-2.




Figure 7-2 Relative trends of total emissions of CO₂, CH₄ and N₂O from sector 5 Waste in Switzerland 1990–2015.

According to the 2006 IPCC Guidelines (IPCC 2006) all emissions from waste-to-energy, i.e. emissions resulting when waste material is used directly as fuel or converted into a fuel, are reported under sector 1 Energy (see also Figure 7-4). Therefore, the largest share of waste-related emissions in Switzerland is not reported under sector 5 Waste. This is illustrated in

Figure 7-3 which provides an overview of all waste-related GHG emissions in Switzerland reported in chp. 7 and elsewhere in the NIR (see also Figure 7-4).




Figure 7-3 Total waste-related GHG emissions from 1990–2015, reported in different sectors.

7.1.2 Overview of waste management in Switzerland

The goals and principles regarding waste management in Switzerland are stated in the Guidelines on Swiss Waste Management (BUS 1986), in the Waste Concept for Switzerland (SAEFL 1992) and in the ordinance regarding waste avoidance and waste management (Swiss Confederation 2015). The four principles are:

- The generation of waste shall be avoided as far as possible.
- Pollutants from manufacturing processes and in products shall be reduced as far as possible.
- Waste shall be recycled wherever this is environmentally beneficial and economically feasible.
- Waste shall be treated in an environmentally sound way. In the long term only materials of final storage quality shall be disposed of in landfills.

Figure 7-4 gives a general overview of the type of treatment and amounts of waste treated in the respective sectors in Switzerland, including imports and exports. Only waste fractions that are emission-relevant are shown. The figure further illustrates where the processes related to the waste management system are reported in the NIR. The following details can be provided regarding the different sectors:

- **1 Energy:** In accordance with the 2006 IPCC Guidelines (IPCC 2006) emissions from waste-to-energy activities, where waste is used as an alternative fuel for energy production, are reported in 1A Fuel combustion activities. This applies to municipal solid waste incineration plants (MSWIP) and special waste incineration plants (SWIP), where energy is recovered (1A1a). MSWIP treat burnable municipal solid waste as well as sewage sludge, burnable construction waste and some special wastes. Cement industry uses conventional fossil fuels but also alternative fuels, which are special waste, dried sewage sludge, biomass as well as plastics collected separately or segregated from solid waste streams (1A2fi). The digestion of biomass and the use of landfill gas is also reported in sector 1 Energy, as such biogas is used for combined heat and power generation. The energy production from renewable goods, such as the use of wood waste in wood-fired power stations, is reported under 1A2, 1A4ai and 1A4bi and 1A4ci.
- **3 Agriculture:** Since 2003 it is forbidden in Switzerland to use sewage sludge as a fertilizer. In 2014, within sector 3 Agriculture only compost used as fertilizer was emission-relevant (N₂O emissions as described in chp. 5.5.2.2, Table 5-22).
- **5 Waste:** Only emissions from waste management activities not used for energy production are reported under sector 5 Waste. Solid waste disposal does not occur anymore in Switzerland as incineration is mandatory for disposal of combustible waste since 2000. Emissions from composting are described under 5B1. Emissions related to digestion, but not directly related to energy production (such as the storage of digested biomass), are reported under 5B2. 5C Waste incineration and open burning of waste accounts for a small fraction only, consisting of illegal waste incineration, sewage sludge incineration, burning of residues in agriculture and private households as well as cremations. Special waste incineration without energy recovery, such as cable incineration or hospital waste incineration plants, no longer take place in Switzerland and is thus crossed out in the figure. These waste fractions are nowadays incinerated in MSWIP and are therefore reported under sector 1 Energy. Emissions related to wastewater treatment are reported under 5D.




Figure 7-4 Overview on the type of treatment and amounts of waste treated in the respective sectors in Switzerland in 2015. Abbreviations: MSWIP: Municipal Solid Waste Incineration Plant, SWIP: Special Waste Incineration Plant, SSIP: Sewage Sludge Incineration Plant, D: Germany, F: France, A: Austria, I: Italy.

Regarding the treatment and amounts of relevant waste types the following details can be provided (state in 2015, recycled amounts are not shown in Figure 7-4 because they are not emission-relevant):

- Municipal solid waste:** In Switzerland more than 50% of the municipal solid waste is collected separately and recycled (FOEN 2016i). The amount of waste incinerated includes imported MSW, mainly from neighbouring countries such as Germany, France, Austria and Italy. The import of waste into Switzerland needs to be authorized by the Federal Office for the Environment. A part of the separately collected plastic fractions from households and industry which cannot be recycled, is used as an alternative fuel in the cement industry.
- Construction waste:** More than 50% of the construction waste is recycled. About half of the recycling takes place at the construction sites, e.g. by reusing material left after breaking up the road cover. The other half is separated at the construction sites and recycled individually, e.g. used glass, metals, concrete etc. A minor amount of combustible construction waste is incinerated in MSWIP. The remaining, inert

construction waste is disposed of in landfills for inert waste (ERM 2016; Wüest&Partner 2015).

- **Special waste:** Special waste refers to a highly diverse waste fraction containing hospital wastes, batteries, electronic waste, hazardous industrial sludge, contaminated soils, solvents, chemicals etc. The special waste is either recycled, biologically treated, landfilled, burnt or exported for landfilling in foreign countries (FOEN 2016i). Only the amount of incinerated special waste is emission-relevant (EMIS 2017/1A1a Kehrichtverbrennungsanlagen and EMIS 2017/1A1a Sondermüllverbrennungsanlagen). Some special waste is also used as an alternative fuel in the cement production (EMIS 2017/1A2f i Zementwerke Feuerung).
- **Sewage sludge:** Since 2010 sewage sludge may not be used anymore as a fertilizer in agriculture due to the content of organic contaminants, heavy metals and other substances. Therefore, all sewage sludge is incinerated, either in MSWIP or in SSIP without energy recovery (internal information provided by the waste section of FOEN). Dried sewage sludge is also used as an alternative fuel in the cement industry (EMIS 2017/1A2fi Zementwerke Feuerung).
- **Biomass:** The term biomass refers to a broad range of materials such as garden waste, grass, wood waste, liquid manure and production remains from the food industry or further fractions, depending on the process concerned. Biomass from agriculture, forestry and private gardens are burned without energy recovery (EMIS 2017/5C2 Abfallverbrennung Land- und Forstwirtschaft). Biomass is also digested or composted (in large-scale composting facilities or backyards). Quantities of biomass refer to wet matter. Biomass such as used wood or animal fat is used as an alternative fuel in the cement industry (EMIS 2017/1A2fi Zementwerke Feuerung). Compost is used as a fertilizer (see Table 5-22 “Other organic fertilisers”).

7.2 Source category 5A – Solid waste disposal

7.2.1 Source category description

Table 7-2 Key categories (KCA incl. LULUCF) of 5A Solid waste disposal.

Code	IPCC Category	GHG	Identification Criteria
5A	Solid Waste Disposal	CH ₄	L1, L2, T1, T2

The source category 5A1 Managed waste disposal sites comprises all emissions from managed solid waste landfill sites. As incineration is mandatory for combustible waste since 2000, inputs into managed solid waste landfill sites have dropped to zero. Remaining emissions thus stem from landfilling before 2000. Emissions from the source category 5A2 Unmanaged waste disposal sites are included in source category 5A1 Managed waste disposal sites. This is motivated by the fact that in Switzerland to date no official unmanaged waste disposal sites exist. Although no reliable data is available, the effective quantity of waste not properly treated in landfills is estimated to be very small.

In Switzerland, six managed biogenic active landfill sites were equipped to recover landfill gas in 2015 (SFOE 2016a). While some landfill gas is used to generate heat only, the landfill gas is generally used in co-generation plants in order to produce electricity and heat. A small amount of the landfill gas is flared (Consaba 2016).

Table 7-3 Specification of source category 5A Solid waste disposal in Switzerland.

5A	Source	Specification
5A1	Managed waste disposal sites	Emissions from managed solid waste landfill sites.
5A2	Unmanaged waste disposal sites	Officially no unmanaged waste disposal sites exist (included in 5A1)

7.2.2 Methodological issues

Methodology (5A)

Emissions are calculated by a Tier 2 method based on the decision tree in Fig. 3.1 of chp. 3. Solid waste disposal in IPCC (2006). The spreadsheet for the First Order Decay (FOD) model provided by IPCC (2006) has been applied and parametrised for Swiss conditions (FOEN 2016g).

The values for the parameter degradable organic carbon (DOC) are provided for each waste fraction (Table 7-4). For all waste types the IPCC default values are used, except for industrial waste. For industrial waste the default value for wood and straw is used, as most of the industrial waste deposited in Switzerland is assumed to be wood waste.

Table 7-4 Degradable organic carbon (DOC) values for fractions of different waste compositions (weight fraction, wet basis).

Waste composition (weight fraction, wet basis)	IPCC default value		Country-specific parameters	
	Range	Default	Swiss Value	Reference and remarks
Food waste	0.08-0.20	0.15	0.15	
Garden	0.18-0.22	0.2	0.2	
Paper	0.36-0.45	0.4	0.4	
Wood and straw	0.39-0.46	0.43	0.43	
Textiles	0.20-0.40	0.24	0.24	
Disposable nappies	0.18-0.32	0.24	NO	not relevant/no activity data
Sewage sludge	0.04-0.05	0.05	0.05	
Industrial waste	0.00-0.54	0.15	0.43	all waste wood

The methane generation rate [1/yr] is chosen according to wet temperate conditions (Table 7-5). For all waste types the IPCC default values are used, except for industrial waste. For industrial waste the default value for wood and straw is used, again based on the fact that most of it is assumed to be wood waste.

Table 7-5 Methane generation rate [1/yr] according to waste by composition for wet temperature conditions.

Waste composition (weight fraction, wet basis)	IPCC default value		Country-specific parameters	
	Range	Default	Swiss Value	Reference and remarks
Food waste	0.1–0.2	0.185	0.185	
Garden	0.06–0.1	0.1	0.1	
Paper	0.05–0.07	0.06	0.06	
Wood and straw	0.02–0.04	0.03	0.03	
Textiles	0.05–0.07	0.06	0.06	
Disposable nappies	0.06–0.1	0.1	NO	not relevant/no activity data
Sewage sludge	0.1–0.2	0.185	0.185	
Industrial waste	0.08–0.1	0.09	0.03	all waste wood

The general parameters are set as follows:

- DOCf (fraction of DOC dissimilated) = 0.5 (IPCC default value)
- Delay time (months) = 6 (IPCC default value)
- Fraction of methane (F) in developed landfill gas = 0.5 (IPCC default value)
- Conversion factor, C to CH₄ = 1.33 (IPCC default value)
- Oxidation factor (OX) = 0.1

The oxidation factor OX has been set to 0.1 according to the 2006 IPCC Guidelines, chp. 3 (IPCC 2006), since it is standard practice in Switzerland to cover the landfills, e.g. with soil.

For the methane correction factors (MCF) for the different solid waste disposal site types IPCC default values are used. Between 1990 and 2015 (the IPCC spreadsheet has to be parametrised from 1950 to 2030/2050) waste distribution to the following three solid waste disposal site types has taken place (for both MSW and IW):

- Methane correction factor (MCF) for unmanaged, shallow SWDS = 0.4 (IPCC default value)
- Methane correction factor (MCF) for unmanaged, deep SWDS = 0.8 (IPCC default value)
- Methane correction factor (MCF) for managed SWDS = 1 (IPCC default value)
- The other two MCF (managed, semi-aerobic and uncategorised) are not relevant because such SWDS are not occurring in Switzerland, i.e. no waste has been distributed to such sites.

The waste composition of MSW deposited has changed during the last 60 years (see Table 7-6).

Table 7-6 Composition of MSW going to solid waste disposal sites (BUS 1978, BUS 1984, FOEN 2014o).

Food	Garden	Paper	Wood	Textile	Nappies	Plastics, other inert	Time period
23.8%	4.2%	36.0%	4.0%	4.0%	0.0%	28.0%	1950-1979
26.5%	2.9%	30.6%	4.3%	3.1%	0.0%	32.6%	1980-1989
21.4%	1.6%	28.0%	5.0%	3.0%	0.0%	41.0%	1990-1999
26.6%	1.4%	21.0%	2.0%	3.0%	0.0%	46.0%	2000-2009
31.5%	1.7%	17.2%	1.8%	3.2%	0.0%	44.6%	2010 -

With these parametrisations and the activity data for municipal solid waste (MSW), industrial waste (IW) and sewage sludge (SS) the amount of CH₄ generated in landfills is calculated. The amount of CH₄ recovered and used as fuel for combined heat and power generation or flared is then subtracted.

For combined heat and power generation and flaring, the emissions of other gases are considered to be proportional to the amount of CH₄ burnt (Table 7-7).

Emission factors (5A)

Emission factors for CO₂, CH₄, CO, NMVOC and SO₂ are country-specific based on measurements and expert estimates, as documented in EMIS 2017/1A1 & 5A Kehrichtdeponien. CO₂ emissions from non-biogenic waste are included, while CO₂ emissions from biogenic waste are excluded from total emissions. Table 7-7 presents the emission factors used in 5A1.

Table 7-7 Emission factors for 5A1 Managed waste disposal sites in 2015.

5A1 Managed waste disposal sites	Unit	CO ₂ biogen	CO ₂ fossil	CH ₄	NO _x	CO	NMVOC	SO ₂
Direct emissions from landfill	t / t CH ₄ produced	3.0	NA	1.0	NA	NA	NA	NA
Flaring	kg / t CH ₄ burned	2750	NA	NA	1.0	17	NA	NA

Activity data (5A)

There are three kinds of activity data for 5A1 Managed waste disposal sites: Waste quantities disposed on landfills, direct CH₄ emissions and CH₄ flared.

For the calculation of these three kinds of activity data the amounts of MSW, IW and SS deposited on SWDS are relevant.

Table 7-8 Activity data in 5A1: Waste disposed on managed waste disposal sites from 1990 to 2015 (documented in EMIS 2017/1A1a & 5A Kehrichtdeponien).

5A1 Managed waste disposal sites	Unit	1990	1995	2000	2005
Municipal solid waste (MSW)	kt	650.0	540.0	291.7	13.7
Construction waste (CW)	kt	150.0	60.0	53.9	1.4
Sewage sludge (SS)	kt (dry)	60.0	28.1	4.2	1.0
Open burned waste	kt	NO	NO	NO	NO
Total waste quantity	kt	860.0	628.1	349.7	16.1

5A1 Managed waste disposal sites	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Municipal solid waste (MSW)	kt	3.6	1.5	1.2	NO						
Construction waste (CW)	kt	0.8	NO								
Sewage sludge (SS)	kt (dry)	0.3	NO								
Open burned waste	kt	NO									
Total waste quantity	kt	4.7	1.5	1.2	NO						

Table 7-8 documents the amounts of municipal solid waste, construction waste and sewage sludge disposed of on managed waste disposal sites over the time period 1990–2015 (as documented in EMIS 2017/1A1a & 5A Kehrichtdeponien). The continuous decline of the waste amounts landfilled happened due to changes in the legislative framework, making incineration mandatory for disposal of combustible waste and banning the disposal of combustible waste on landfills from 1 January 2000. The amounts of combustible waste disposed of on managed waste disposal sites reached zero in 2009. Open burning of waste on SWDS has occurred in the past but is assumed to have ceased since 1990 (expert judgement) and is therefore NO in Table 7-8.

With these primary activity data total CH₄ emissions generated are calculated using the spreadsheet FOD model provided by IPCC (2006). For the calculation of direct CH₄ emissions, CH₄ flared and used in co-generation units is determined and subtracted from total CH₄ emissions (

Table 7-9). The landfill gas recovered and used as fuel for co-generation units is reported under 1A1 Energy in accordance with the 2006 IPCC Guidelines (IPCC 2006). The sum of landfill gas flared and landfill gas used in co-generation units is reported as being recovered in CRF table 5.A.

The amount of CH₄ used in co-generation stems from the Swiss statistics of renewable energies (SFOE 2016a). The amount of landfill gas flared has been assessed in a separate investigation in the year 2015 (Consaba 2016). The CH₄ flared has been estimated as follows:

1. A list of all managed SWDS that are still operated or have been closed since 1990 was compiled.
2. Their technical equipment was assessed and deduced (motors, torches, gas drainage, etc.).
3. Four types of managed landfill sites according to their equipment and CH₄ management were distinguished:
 - a) landfills with gas recovery in combined heat and power generation, boiler and torch
 - b) landfills with gas recovery or thermal treatment (boiler, torch, non-catalytic oxidation, flameless oxidation)
 - c) landfill gas recovery without methane elimination (biofilter, aerobiosation)

d) landfills without gas treatment (direct release)

4. A survey was conducted in 14 managed SWDS and data on their operation mode has been collected.
5. With these data the amounts flared in SWDS category a) and b) were estimated.
6. The amount flared on all managed SWDS has been extrapolated considering the waste amounts deposited.
7. A time series for the amount of methane torched relative to the total amount of CH₄ estimated with the Swiss FOD IPCC 2006 model (managed and unmanaged sites) has been calculated.

The amount flared is expressed as a percentage of CH₄ produced in all SWDS in Switzerland. The percentage flared varies between 5 and 15% since 1990.

Table 7-9 Activity data in 5A1: Direct CH₄ emissions, CH₄ flared and CH₄ used in co-generation units (CHP) from 1990 to 2015 (as documented in EMIS 2017/1A1a & 5A Kehrichtdeponien).

5A1 Managed waste disposal sites	Unit	1990	1995	2000	2005
CH ₄ direct emissions	kt	30.5	22.9	20.9	21.1
CH ₄ flared	kt	1.8	5.2	5.6	3.4
CH ₄ used in co-generation units (reported under 1A1a)	kt	4.9	12.1	11.3	4.1

5A1 Managed waste disposal sites	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
CH ₄ direct emissions	kt	20.9	20.1	19.2	18.3	17.5	16.7	16.0	15.3	14.7	14.0
CH ₄ flared	kt	3.2	2.9	2.6	2.4	2.4	2.1	1.8	1.6	1.4	1.4
CH ₄ used in co-generation units (reported under 1A1a)	kt	2.7	2.1	1.8	1.5	1.0	0.9	0.9	0.8	0.6	0.4

The CH₄ generated in landfill sites decreased since 1990 because waste quantities disposed of in landfills have been decreasing. Together with the relative increase of CH₄ recovery from 1990 until 2015 this is the reason for CH₄ emissions from the source category 5A being a key source regarding trend.

7.2.3 Uncertainties and time-series consistency

Uncertainty in CH₄ and CO₂ emissions from 5A Solid waste disposal

For lack of a detailed uncertainty analysis with the new FOD model, a combined uncertainty of 30% is assumed for the CH₄ emissions (EMIS 2017/1A1a & 5A Kehrichtdeponien).

Consistency: Time series for 5A Solid waste disposal are all considered consistent.

7.2.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

7.2.5 Category-specific recalculations

No category-specific recalculations were carried out.

7.2.6 Category-specific planned improvements

In response to UNFCCC (2017/ID#W.6) the reporting of long-term storage of C in waste disposal sites in the CRF tables will be considered.

7.3 Source category 5B – Biological treatment of solid waste

7.3.1 Source category description

Table 7-10 Key categories (KCA incl. LULUCF) of 5B Biological treatment of solid waste.

Code	IPCC Category	GHG	Identification Criteria
5B	Biological Treatment of Solid Waste	CH ₄	T1, T2

Source category 5B Biological treatment of solid waste comprises the process-related GHG emissions from composting and from digesting of organic waste.

5B1 Composting covers the GHG emissions from centralized composting plants with a capacity of more than 100 tonnes of organic matter per year. Backyard composting is also common practice in Switzerland and taken into account.

In 5B2 Anaerobic digestion at biogas facilities emissions from gas leakages occur as well as from digested matter (solid leftovers after completion of anaerobic microbial degradation of organic matter) which is being composted. The biogas is used for combined heat and power generation or upgraded and used as fuel.

In 5B Biological treatment of solid waste the emissions from the composting of digested matter as well as the CH₄ losses from biogas facilities and emerging from biogas upgrading are included. The emissions related to the use of biogas for combined heat and power generation as well as emissions from biogas-upgrading are reported in sector 1 Energy.

Table 7-11 Specification of source category 5B Biological treatment of solid waste.

5B	Source	Specification
5B1	Composting	Process-related emissions from composting of organic waste
5B2	Anaerobic digestion at biogas facilities	Process-related emissions from digesting of organic waste

7.3.2 Methodological issues

7.3.2.1 Composting (5B1)

Methodology (5B1)

Emissions are calculated by a Tier 2 method based on chp. 4.1.1 Biological treatment of solid waste in IPCC (2006).

Emission factors (5B1)

The emission factors used for source category 5B1 Composting are summarized in Table 7-12 and documented in detail in EMIS 2017/5B1 Kompostierung Industrie. Emission factors are country-specific and encompass CO₂, CH₄, NH₃, N₂O and NMVOC based on measured or estimated values reported in the literature. CH₄ emission factors are taken from SFOE (1999), N₂O emission factors from Schenk et al. (1997) and SFOE (1999).

Activity data (5B1)

Activity data for source category 5B1 Composting are shown in Table 7-13 and documented in detail in EMIS 2017/5B1 Kompostierung Industrie. The activity data for industrial composting from 1990 to 2002 and for 2007 are based on waste statistics. For the year 2020 a value is projected by scaling the activity data for 2007 by the increase of the population from 2007 until 2020. The quantities from 2003 to 2006 and from 2008 to 2019 are interpolated. However, the waste statistics only covers centralized composting plants with a capacity of more than 100 tonnes of organic matter per year. Therefore, the quantities composted in smaller sites, in particular neighbourhood and backyard composting, are estimated to be approx. 10% of the quantities of centralized composting.

7.3.2.2 Anaerobic digestion at biogas facilities (5B2)

Methodology (5B2)

In source category 5B2 Anaerobic digestion at industrial and agricultural biogas facilities are considered. The produced biogas is used for combined heat and power generation or upgraded to natural gas quality. Accordingly, biogas upgrading is considered as a separate process in 5B2. However, emissions from the use of biogas as fuel for combined heat and power generation are reported under sector 1 Energy, in accordance with the 2006 IPCC Guidelines (IPCC 2006).

For the emissions from 5B2 Anaerobic digestion at biogas facilities, a Tier 2 method is used. While industrial and agricultural biogas facilities are separately considered, the same emission factors are used (see below). As mentioned above, emissions from biogas upgrading are estimated separately, based on the amount of biogas upgraded.

Emissions of greenhouse gases from industrial and agricultural biogas facilities are estimated to be a constant emission factor for each biogas facility. This is based on an evaluation of measurement data for methane losses that has shown that those losses are not dependent on the amount of substrate processed in a particular facility. Therefore, CH₄

emissions are calculated based on an emission factor per plant multiplied by the number of industrial and agricultural biogas facilities, respectively.

In contrast, emissions of air pollutants are calculated based on estimates from up to seven different process steps (such as pre-storage, primary and secondary digester, interim storage, maturing, handling of biogas etc.), as documented in EMIS 2017/1A1a & 5 B 2_Vergärung LW and EMIS 2017/1A1a & 5 B 2_Vergärung IG. However, as NMVOC and CO emissions from source category 5B are of biogenic origin, they are not considered for the calculation of indirect CO₂ emissions in chp. 9 Indirect CO₂ and N₂O emissions.

N₂O emissions from source category 5B2 are considered to be negligible according to the 2006 IPCC Guidelines (IPCC 2006), and are therefore set to zero.

Emission factors (5B2)

Table 7-12 presents the emission factors used in 5B2 Anaerobic digestion at biogas facilities. As documented in FOEN (2015n), the emission factor for CH₄ for anaerobic digestion at industrial and agricultural biogas facilities is based on investigations performed in the framework of the GHG emission compensation projects. Field measurements indicate that there is no correlation between the produced amount of biogas and the amount of biogas lost to the atmosphere. The investigated data show that on average each biogas facility loses 1.23 t CH₄ per year to the atmosphere. This value is used to estimate the emissions from industrial and agricultural biogas facilities in Switzerland.

The emission factor for losses of CH₄ from biogas upgrading is based on official regulations regarding maximal CH₄ leakage, as well as studies focussing on CH₄ emissions from biogas upgrading. Accordingly, regulations by the Swiss Gas and Water Association (SGWA 2016a) set an emission limit value (ELV) for CH₄ losses from biogas upgrading. In 1990, such losses were allowed to be 5% of the upgraded amount, in 2014 the limit was lowered to 2.5%. Measurements in a few biogas upgrade installations in 2007, 2013 and 2014 showed the following losses: 2007 one plant: 2.6%, 2013 one plant: 1%, 2014 three plants: 1.3%, 1.8%, and 3.5%. The measurements showed that the emission limits were respected (with the exception of one plant in 2014) and therefore Switzerland decided to set the losses from biogas upgrading to the ELV with the assumption of a linear improvement between the 1990 and the 2014 value. The continuous improvement seems plausible, as newer plants show fewer losses and values of less than 1%–2.5% are state of the art.

Activity data (5B2)

Activity data for 5B2 Anaerobic digestion at biogas facilities, as shown in Table 7-13, are based on data from the Swiss renewable energy statistics (SFOE 2016a). Relevant are the number of industrial and agricultural biogas facilities, as well as the total amount of biogas upgraded.

Table 7-12 Emission factors for 5B Biological treatment of solid waste in 2015.

5B Biological treatment of solide waste	Unit	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂
Composting	g/t composted waste	5'000	70	NA	NA	1'700	NA
Digestion (industrial biogas facilities)	t/facility	1.23	NA	NA	NA	NA	NA
Digestion (agricultural biogas facilities)	t/facility	1.23	NA	NA	NA	NA	NA
Biogas up-grade	g/GJ	500	NA	NA	NA	NA	NA

Table 7-13 Activity data in 5B Biological treatment of solid waste.

5B Biological treatment of solide waste	Unit	1990	1995	2000	2005
Composting	kt wet	260	400	640	735
Digestion (industrial biogas facilities)	number	NO	4	11	14
Digestion (agricultural biogas facilities)	number	98	76	68	72
Biogas up-grade	GJ	NO	NO	19'866	40'637

5B Biological treatment of solide waste	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Composting	kt wet	737	739	747	755	763	771	779	787	795	803
Digestion (industrial biogas facilities)	number	13	16	16	21	22	28	26	26	25	26
Digestion (agricultural biogas facilities)	number	80	77	75	75	72	80	89	97	98	99
Biogas up-grade	GJ	41'710	50'966	71'721	85'008	121'627	168'170	236'074	283'503	368'862	447'877

To improve transparency the CH₄ and N₂O emissions of source category 5B Biological treatment of solid waste are shown on a completely disaggregated level in Table 7-14.

Table 7-14 CH₄ and N₂O emissions of 5B Biological treatment of solid waste.

5B Biological treatment of solide waste	Gas	Unit	1990	1995	2000	2005
Composting	CH ₄	t	1'300	2'000	3'200	3'676
	N ₂ O	t	18.2	28.0	44.8	51.5
Digestion (industrial)	CH ₄	t	NO	4.9	13.6	17.2
Digestion (agricultural)	CH ₄	t	120.5	93.5	83.6	88.6
Biogas up-grade	CH ₄	t	NO	NO	15.7	28.0

5B Biological treatment of solide waste	Gas	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Composting	CH ₄	t	3'685	3'693	3'734	3'774	3'814	3'855	3'895	3'935	3'975	4'016
	N ₂ O	t	51.6	51.7	52.3	52.8	53.4	54.0	54.5	55.1	55.7	56.2
Digestion (industrial)	CH ₄	t	16.0	19.7	19.7	25.9	27.1	34.5	32.0	32.0	30.8	32.0
Digestion (agricultural)	CH ₄	t	98.4	94.7	92.3	92.3	88.6	98.4	109.5	119.4	120.6	121.8
Biogas up-grade	CH ₄	t	27.8	32.9	44.8	51.3	70.9	94.7	128.0	147.7	184.4	223.9

7.3.3 Uncertainties and time-series consistency

Uncertainty in CH₄ emissions from composting and digestion

The uncertainty of the CH₄ emission factors in source category 5B1 Composting is estimated at 40% (EMIS 2017/5B1 Kompostierung Industrie). The contribution of 5B1 to the CH₄ emissions of 5B is about 75%. For 5B2 Anaerobic digestion at biogas facilities the same uncertainty is assumed.

The uncertainty of the related activity data is estimated at 10% (EMIS 2017/5B1 Kompostierung Industrie) due to the high reliability of the waste statistics.

The uncertainty of the N₂O emission factor for 5B1 Composting is unknown. Therefore, the uncertainty of N₂O emissions is estimated to be medium, which is attributed by Table 1-10 to a combined uncertainty of 80%.

Consistency: Time series for 5B Biological treatment of solid waste are all considered consistent.

7.3.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

7.3.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 or 2014 of at least 0.2 kt CO₂ eq are quantified. All the other recalculations have an impact of less than 0.2 kt CO₂ eq in the years 1990 and 2014.

- 5B2: The CH₄ emission factor for biogas upgrading has been recalculated for the whole time series from 1990–2014. The net calorific value of methane has been adjusted to 50 GJ/t (from 46.5) to be in accordance with the fact sheet with CO₂-emission factors (FOEN 2015d) and the rest of the inventory. This recalculation leads to a decrease by 0.3 kt CO₂ eq in 2014.

7.3.6 Category-specific planned improvements

Activity data for backyard composting is assumed to be approx. 10% of the amount of waste composted in industrial plants in the year 2007 and later. This value is based on expert judgements. It is planned for a subsequent submission to assess composting activities, i.e. activity data and emission factors. .

7.4 Source category 5C – Incineration and open burning of waste

7.4.1 Source category description

Source category 5C – Incineration and open burning of waste is not a key category.

There is a long tradition in Switzerland to incinerate waste. The heat generated during the incineration has to be recovered if technically and economically feasible. In accordance with the 2006 IPCC Guidelines (IPCC 2006) emissions from waste-to-energy activities are dealt with in 1A1a Public electricity and heat production.

5C1 encompasses incineration of hospital wastes, illegal waste incineration, incineration of insulation of materials from cables, of sewage sludge and in crematoria.

5C2 consists of emissions from open burning of branches in agriculture and gardening. Natural agricultural and gardening residues consist of fallen fruit trees, part of diseased residue which are cut up, collected and burned off-site. Field burning of agricultural residues

does not occur in Switzerland. Emissions from open burning of natural residues in forestry are reported in LULUCF sector 4 V (chapter 6.4.2.13).

Table 7-15 Overview of waste incineration sources reported under 5C.

5C	Waste incineration	Specification
5C1	Hospital waste incineration	Emissions from incinerating hospital waste in hospital incinerators
	Illegal waste incineration	Emissions from illegal incineration of municipal solid wastes at home. Emissions from waste incineration at construction sites (open burning)
	Insulation material from cables	Emissions from incinerating cable insulation materials
	Sewage sludge	Emissions from sewage sludge incineration plants
	Crematoria	Emissions from the burning of bodies in crematoria
5C2	Open burning of branches	Open burning of branches in agriculture and gardening.

7.4.2 Methodological issues

Methodology (5C)

Emissions are calculated using Tier 2 methods based on chp. 5.2. in IPCC (2006). In general, the GHG emissions are calculated by multiplying the waste quantity incinerated by emission factors. For crematoria, the GHG emissions are calculated by multiplying the number of cremations by emission factors.

For sewage sludge incineration plants the respective waste quantities are based on reliable statistical data (updated every two years until 2006). The emission factors are based on emission declarations from an incineration plant in 2002 that covered approximately one third of the Swiss capacities. Due to the lack of better or newer data these emission factors are kept constant since then and no improvement in flue gas cleaning standards is assumed.

For hospital waste incineration, illegal waste incineration and incineration of insulation material, the waste quantities used are based on expert estimates.

The emissions from burning of residues in agriculture and gardening are calculated by multiplying the annual estimate of branches burnt (in kt of wood equivalent) by emission factors (IPCC default method).

Emission factors (5C)

Emission factors of all categories within 5C1 for CO₂, CH₄, N₂O, CO, NMVOC and SO₂ are country-specific based on measurements and expert estimates, as documented in EMIS 2017/5C1. Table 7-16 presents an overview of the emission factors for 5C in 2015.

Table 7-16 Emission factors for 5C Waste incineration and open burning of waste in 2015.
Documentation/sources: EMIS 2017/5C1 and EMIS 2017/5C2 (details see Table 7-15), EMEP/EEA (2013).

5C Waste incineration and open burning of waste	CO₂ t/t	CH₄ kg/t	N₂O g/t	NO_x kg/t	CO kg/t	NMVOC kg/t	SO₂ kg/t
Hospital waste incineration	0.9	NA	60	1.5	1.4	0.3	1.3
Municipal waste incineration (illegal)	0.5	6.0	150	2.5	50	16	0.75
Industrial waste incineration	1.3	NA	NA	1.3	2.5	0.5	6.0
Sewage sludge incineration	NA	0.1	800	0.7	0.19	0.005	0.47
Open burning of natural residues in agriculture	NA	6.8	180	1.4	49	1.5	0.03
Open burning of natural residues in private households	NA	6.8	180	1.4	49	1.5	0.03
	CO₂ t/crem.	CH₄ kg/crem.	N₂O g/crem.	NO_x kg/crem.	CO kg/crem.	NMVOC kg/crem.	SO₂ kg/crem.
Crementation	NA	NA	NA	0.21	0.06	0.007	NA

Comments on the CO₂ emission factors:

- For all waste incineration categories, only CO₂ emissions from non-biogenic waste are taken into account.
- Hospital waste incineration: The waste is mainly of fossil origin. The default value for the CO₂ emission factor is taken from SAEFL (2000). Since 2002, no emissions from hospital waste incineration occur, as all hospital waste incinerator plants have been closed and hospital waste is incinerated in municipal solid waste incineration plants (accounted for in 1A1a).
- Illegal waste incineration: The CO₂ emission factor is estimated by using the same assumption as in case of MSW incineration: The C-content is based on the study by FOEN (2014I) and the fossil carbon fraction was determined by Ryttec (2014). See also chp. 3.2.5.2 and detailed information in EMIS 2017/1A1a Kehrlichtverbrennungsanlagen (pp. 5–7).
- Industrial waste (consists of cable insulation materials): The CO₂ emission factor is based on measurements of the flue gas treatment of a cable disassembling site where O₂ was measured in the flue gas. Assuming that the ratio of CO₂/O₂ is the same as in municipal solid waste incineration plants, a fraction of 7% of CO₂ results. Based on these assumptions, an emission factor of 1.3 kg/kg cable can be derived. Since 1995, no emissions from incinerating cable insulation materials occurred.
- Sewage sludge plants: As sewage sludge is biogenic waste, the emission factor for CO₂ is zero. It is assumed that the share of fossil fuel used during the start-ups is negligible.

Additional information on emission factors of all other (non-CO₂) gases:

- Hospital waste incineration: All emission factors are taken from SAEFL (2000).
- Illegal waste incineration: The emission factor for N₂O is taken from the 2006 IPCC Guidelines (IPCC 2006, vol. Waste), the emission factors for CH₄, SO₂, NO_x, NMVOC from SAEFL (2000) and USEPA (1995a).
- Industrial waste (cable insulation materials): All emission factors are adopted from SAEFL (2000).

- Sewage sludge plants: For 1990 emission factors are taken from SAEFL (2000). From 2002 onwards constant emission factors are used, which are deduced from measurements (LHA 2004) taken at the largest sewage sludge incineration plant incinerating one third of Switzerland's sewage sludge. Between 1990 and 2002 the emission factors are interpolated. They show reductions in NMVOC, CO, SO₂ and CH₄ emissions due to gradual technical improvements.
- Crematoria: NMVOC and CO emissions were reduced by technical improvements. A large number of measurements were analysed (crematoria as well as other types of installations are obliged to monitor their emissions by the Swiss Federal Ordinance on Air Pollution Control (Swiss Confederation 1985) such that plant-specific emission factors are available for installations with retrofitted flue gas treatment as well as non-retrofitted installations. The emission factors are calculated as weighted averages of cremations taking place in retrofitted and non-retrofitted cremation plants (EMIS 2017/5C1 Krematorien).
- The emission factors of burning of branches in agriculture and gardening are calculated based on EMEP/EEA (2013) except for CH₄ und N₂O which are based on EMEP/CORINAIR (EMEP/EEA 2002), see also documentation in EMIS 2017/5C2 Abfallverbrennung in der Land- und Forstwirtschaft.
- General remark: In years with no specific data for activity data or emission factors the respective data are interpolated.

General remark: Indirect CO₂ emissions from fossil CO and NMVOC emissions are reported in chp. 9 Indirect CO₂ and N₂O emissions.

Activity data (5C)

The activity data for 5C Waste incineration are the quantities of waste incinerated, see Table 7-17.

Table 7-17 Activity data for the different emission sources within source category 5C Waste incineration and open burning of waste.

5C Incineration and open burning of waste	Unit	1990	1995	2000	2005
Hospital waste incineration	kt	30.0	17.5	5.0	NO
Municipal waste incineration (illegal)	kt	32.3	26.2	24.9	21.7
Industrial waste incineration	kt	7.5	NO	NO	NO
Sewage sludge incineration	kt dry	57.0	50.2	64.3	94.9
Open burning of natural residues in agriculture	kt	16.5	15.2	14.0	12.8
Open burning of natural residues in private households	kt	6.1	4.9	3.6	2.4
Total	kt	149.3	114.0	111.8	131.7
Cremation	Numb.	37'513	40'968	44'821	48'169

5C Incineration and open burning of waste	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Hospital waste incineration	kt	NO									
Municipal waste incineration (illegal)	kt	22.6	22.1	22.4	20.7	21.0	20.3	20.3	19.9	19.3	19.3
Industrial waste incineration	kt	NO									
Sewage sludge incineration	kt dry	92.7	95.2	97.7	100.1	102.6	102.4	100.8	120.9	121.0	127.6
Open burning of natural residues in agriculture	kt	12.5	12.3	12.0	11.8	11.5	11.5	11.6	11.6	11.6	11.6
Open burning of natural residues in private households	kt	2.2	1.9	1.7	1.5	1.2	1.2	1.2	1.3	1.3	1.3
Total	kt	130.0	131.5	133.8	134.1	136.4	135.5	133.9	153.6	153.2	159.8
Cremation	Numb.	48'083	49'413	51'116	52'402	52'813	52'530	50'567	53'205	55'616	59'664

Hospital waste incineration: Does not occur anymore in specific hospital waste incineration plants since 2002. The amount of hospital waste burnt in 1990 stems from a study (BUS 1988).

Illegal municipal waste incineration: As waste incineration outside incineration plants is forbidden in Switzerland, no data is available. Illegal incineration of waste e.g. in wood stoves, garden fires, construction sites etc. is decreasing due to surveillance by authorities but also by citizens that would report open burning. However there still are cases of illegal waste incineration. It is assumed that 1% of all waste in Switzerland has been burnt illegally in 1990 and that this value decreases to 0.25% in 2030 and then remains constant.

Industrial waste incineration (cable insulation): Does not occur anymore since 1995. The amount burnt in 1990 is estimated by the amount reported by a company that was supposed to burn approx. 1/3 of all insulation material in Switzerland.

Sewage sludge incineration: Activity data for sewage sludge incineration is calculated as follows. Total amount of sewage sludge (according to waste statistics) minus sewage sludge burnt in MSWIP minus sewage sludge used as alternative fuel in cement industry.

Open burning of natural residues: The amount of natural residues burnt openly have been estimated in a study (INFRAS 2014). Open burning of such residues is regulated in the Ordinance on Air Pollution Control OAPC, Article 26b. In Switzerland cantonal authorities are responsible for surveilling that the regulations of the OAPC are respected. For the study an inquiry of some cantonal authorities has been performed in order to assess the activity data for these processes. Emissions from open burning of natural residues in forestry (5 C 2 ii) are reported in LULUCF sector 4 V (chapter 6.4.2.13).

Cremations: AD are reported by the Swiss Cremation Association. These statistics provide data for each year and are updated regularly.

7.4.3 Uncertainties and time-series consistency

The uncertainty assessment is based on expert judgment results in high uncertainties for CO₂ and CH₄ of 40% and 60% of emission estimates, respectively, and for N₂O in low uncertainty of 40% of emission estimates (see Table 1-10 for quantification of “low” and “high”).

Consistency: Time series for 5C Waste incineration and open burning of waste are all considered consistent.

7.4.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

7.4.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. These recalculations cause a change in emission levels 1990 or 2014 by less than 0.2 kt CO₂ eq.

- 5C1: Activity data for illegal waste incineration for the year 2013 has slightly increased due to the correction of an error in waste statistics.
- 5C1: Activity data for sewage sludge incineration for the year 2013 has slightly increased due to the correction of an error in waste statistics.
- 5C2: As a consequence of the greenhouse gas inventory UNFCCC in-country review 2016 greenhouse gas emissions from open burning of natural residues in forestry (5 C 2 ii) were moved to LULUCF sector 4 V (new in chapter 6.4.2.13). This improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

7.4.6 Category-specific planned improvements

No category-specific improvements are planned.

7.5 Source category 5D – Wastewater treatment and discharge

7.5.1 Source category description

Table 7-18 Key categories (KCA incl. LULUCF) of 5D Wastewater treatment and discharge.

Code	IPCC Category	GHG	Identification Criteria
5D	Wastewater treatment and discharge	CH4	L1, L2, T2
5D	Wastewater treatment and discharge	N2O	L1, L2, T2

The source category 5D1 Domestic wastewater comprises all emissions from liquid waste handling and sludge from housing and commercial sources (including grey water and night soil). In Switzerland, municipal wastewater treatment (WWT) plants treat wastewater from single cities or several cities/municipalities together. Wastewater in general is treated in three steps: 1. Mechanical treatment, 2. Biological treatment, and 3. Chemical treatment. The treated wastewater flows into a receiving system (lake, river or stream). Pre-treated effluents from industries are also handled for final treatment in municipal WWT plants (see below). Switzerland's wastewater management infrastructure – comprising about 850 WWT plants and 40'000–50'000 km of public sewers – is now practically complete (FOEN 2012h). The vast majority of WWT plants apply an anaerobic sludge treatment with sewage gas recovery, and use the sewage gas for heat production. About 290 WWT plants also apply combined heat and power (CHP) units.

The source category 5D2 Industrial wastewater comprises all emissions from liquid waste handling and sludge from industrial processes such as food processing, textiles, car-washing places, electroplating plants, and pulp/paper production. These processes may result in effluents with a high load of organics. Depending on the contaminants, an on-site pre-treatment is necessary in order to reduce the load of pollutants in the wastewater, to meet the regulatory standards (which are in place to preclude disruptions of the municipal WWT plants), and to reduce discharge fees. The on-site pre-treatment is generally anaerobic, in order to use the sewage gas as source for heat and power production. Currently, about 20 industrial WWT plants pre-treat wastewater before its discharge to the domestic sewage system, where the industrial wastewater is additionally treated together with domestic wastewater in municipal WWT plants. Due to this strong connection with domestic wastewater treatment, industrial wastewater is not identified as separate wastewater stream for the calculation of GHG emissions, but joined to the domestic wastewater treatment. For the calculation of emissions of other gases (NO_x, CO, NMVOC, SO₂), domestic and industrial wastewater streams are distinguished (i.e. different emission factors relative to population, see below).

Table 7-19 Specification of source category 5D Wastewater treatment and discharge.

5D	Source	Specification
5D1	Domestic wastewater	Emissions from liquid waste handling and sludge from housing and commercial sources
5D2	Industrial wastewater	Emissions of precursors from handling of liquid wastes and sludge from industrial processes (emissions of CH ₄ and N ₂ O are implemented in 5D1)

Category 5D contains all direct emissions from wastewater handling, including direct emissions of sewage gas (leakage), torching, combined heat and power (CHP) units, furnaces (only heat production), and upgrading of sewage gas to natural gas quality (which can then be fed into the natural gas network and/or used as fuel). However, wastewater treatment also leads to further emissions reported in other categories, as illustrated in Figure 7-5 below.

Emissions associated with sewage sludge drying are assumed to be negligible. The discharge of sewage sludge on agricultural soils has been phased out since 2003 and is generally forbidden since 2008. Therefore, this process is crossed out in Figure 7-5. The same applies to solid waste disposal on land (5A). All sewage sludge is incinerated either in MSW incineration plants (1A1a), SS incineration plants (5C) or used as alternative fuel in the cement industry (1A2f).




Figure 7-5 System boundaries of processes related to wastewater treatment. CHP= Combined heat and power generation. MSWIP = Municipal solid waste incineration plant. SSIP = Sewage sludge incineration plant.

7.5.2 Methodological issues

Emissions are calculated by a Tier 2 method based on the decision tree in Fig. 6.2 and Fig. 6.3 in chp. 6. Wastewater treatment and discharge in IPCC (2006). Details regarding the calculation of CH₄ and N₂O emissions are provided in the following.

7.5.2.1 CH₄ emissions

Methodology (5D, CH₄)

CH₄ emissions from wastewater treatment and discharge take into account emissions stemming from organically degradable material in wastewater and emissions related to sewage gas production (and recovery) from sewage sludge (in domestic as well as industrial installations). As noted above most industrial wastewater may be treated together with domestic wastewater in municipal WWT plants. The industrial/commercial contribution to total CH₄ emissions from wastewater is thus taken into account in the calculation of CH₄ emissions from domestic wastewater by means of a correction factor for additional industrial

biochemical oxygen demand (BOD) discharged into the domestic sewer system. However, industries handling wastewater with high BOD usually use anaerobic digesters to produce sewage gas. The emissions related to sewage gas production (and recovery) during industrial pre-treatment of wastewater are also taken into account in the calculation of emissions from municipal WWT plants, because the underlying Swiss renewable energy statistics in Switzerland (see below) does not differentiate between sewage gas production in domestic and industrial WWT plants.

Accordingly, total CH₄ emissions from domestic and industrial wastewater treatment and discharge are calculated as the sum of two terms:

$$CH_{4,total} = CH_{4,wastewater} + CH_{4,sewage\ gas}$$

(i) Wastewater

In accordance with the 2006 IPCC Guidelines (IPCC 2006) the contribution of wastewater seweraged to WWT plants is determined by:

$$CH_{4,wastewater} = EF_{wastewater} * T_{Plant} * TOW$$

$EF_{wastewater}$ corresponds to the emission factor (see below), T_{Plant} to the fraction of population connected to municipal WWT plants in each year and TOW to the total organically degradable material in the wastewater per year.

From all inhabitants (urban and rural) 90% were connected to WWT plants in 1990, and this percentage reached 97% in 2006, remaining constant thereafter. Switzerland reports emissions only from wastewater discharged to the public sewer system, without taking into account potential emissions from wastewater of unconnected inhabitants. However, emissions from the small fraction of wastewater not treated in WWT plants (since 2006 the wastewater from 3% of the population) are negligible. Federal law only permits alternative treatment systems in remote and sparsely populated regions. Some of such alternative systems treat wastewater very similar to centralized WWT plants, often under aerobic conditions. The sewage sludge from these small scale treatment installations is either dealt with by centralized WWT plants or MWIP (municipal waste incineration plants). Simpler systems are e.g. septic tanks with at least three chambers. However, the production of CH₄ in an anaerobic environment is strongly temperature dependent and significant CH₄ production is unlikely below 15°C due to the inactivity of methanogens (IPCC 2006). As in Switzerland alternative systems are typically buried, the wastewater reaches the rather constant temperature of the surrounding soil, approximately corresponding to the mean annual air temperature. At Grono, the warmest place in Switzerland, the mean annual temperature is 12.4°C. Accordingly, in alternative treatment systems the temperature of the wastewater is too low to produce substantial CH₄ emissions. CH₄ emissions from wastewater produced by inhabitants not connected to municipal WWT plants are thus considered insignificant and set to zero in the Swiss greenhouse gas inventory.

ii) Sewage gas

The CH₄ emissions resulting from sewage gas treatment (aiming at stabilizing the sewage sludge and producing sewage gas) are calculated based on a country-specific implied emission factor ($EF_{sewage\ gas}$, see below), which is normalized with population (P):

$$CH_{4,sewage\ gas} = EF_{sewage\ gas} * P$$

Emission factors (5D, CH₄)

(i) Wastewater

The wastewater of all connected inhabitants, i.e. virtually all wastewater generated in Switzerland, is seweraged to WWT plants using closed sewer systems. The emission factor according to the 2006 IPCC Guidelines (IPCC 2006), Equation 6.2, is represented by the product of the maximum CH₄ producing potential (B_0 , default value 0.60 kgCH₄/kgBOD) and the methane correction factor (MCF) for the wastewater treatment and discharge system. For the wastewater seweraged to centralized WWT plants, the 2006 IPCC Guidelines (IPCC 2006) propose that the MCF is zero (range 0.0–0.1) for well managed aerobic WWT plants. While WWT plants are generally well managed in Switzerland and mostly operated aerobically (with the exception of sewage sludge treatment, which is considered separately, see below), some CH₄ emissions may still occur. Therefore, the MCF is set to 0.05 (corresponding to the mid-value of the range of well managed aerobic WWT plants), which also brings total CH₄ emissions from WWT in Switzerland to similar values as estimated by Hiller et al. (2014) in their peer-reviewed study. This leads to the following constant emission factor:

$$EF_{wastewater} = B_0 * MCF = 0.60 \frac{kgCH_4}{kgBOD} * 0.05 = 0.03 \frac{kgCH_4}{kgBOD}$$

As mentioned above the maximum CH₄ producing capacity of the wastewater not treated in WWT plants is zero, as the wastewater has a temperature most likely too low to produce significant amounts of CH₄. Accordingly, the emission factor for wastewater not treated in WWT plants is zero and the corresponding emissions are zero, too.

(ii) Sewage gas

To calculate the country-specific implied emission factor $EF_{sewage\ gas}$ for CH₄ emissions from sewage gas treatment the total sewage gas production (in domestic and industrial systems) is taken into account based on detailed Swiss renewable energy statistics in Switzerland (SFOE 2016a). These statistics provide the amount of sewage gas used in furnaces and CHP installations, as well as the amount of sewage gas upgraded to natural gas quality. It is assumed that 2% of the total amount of sewage gas is flared and 0.75% of the total amount is leaking. It is further assumed that the leakage of upgraded gas linearly decreases from 5% in 1990 to 2.5% in 2014, remaining constant thereafter. The emission factor is adapted on a yearly basis due to the respective annual changes in population and the total production of sewage gas.

(iii) Values of emission factors referred to the number of inhabitants

The CH₄ emission factors for 5D Wastewater treatment and discharge are summarized in Table 7-20.

Table 7-20 Country-specific CH₄ emission factors for source category 5D Wastewater treatment and discharge in 2015 referred to the number of inhabitants. Detailed information is given in EMIS 2017/5D1 5D2 Kläranlagen GHG (Wastewater Handling - Emissions of Nitrous Oxide (N₂O) and Methane (CH₄), Update to the 2006 IPCC Guidelines).

5D Wastewater treatment and discharge	Unit	2015
Population	in 1000	8'282
Emissions from WW sewered to WWT plants	kg CH ₄ /person/a	0.80
Emissions from WW not sewered to WWT plants	kg CH ₄ /person/a	NA
Emissions from losses during sludge treatment	kg CH ₄ /person/a	0.06

Activity data (5D, CH₄)

(i) Wastewater

In correspondence with the emission factor $EF_{wastewater}$ given above, the activity data is the fraction of population connected to municipal WWT plants (T_{plant}), as well as the total organically degradable material (TOW) in domestic and industrial/commercial wastewater. According to the 2006 IPCC Guidelines (IPCC 2006), TOW is calculated by

$$TOW = P * BOD * 0.001 * I * 365$$

TOW is given in kg BOD/yr (BOD: biochemical oxygen demand) and P is the population (see Table 7-21). For BOD the default value for Europe given by the 2006 IPCC Guidelines (IPCC 2006) is used for Switzerland (60 g/inhabitant/day). The parameter I corresponds to the correction factor for additional industrial/commercial BOD discharged into domestic sewers with default value 1.25. While the amount of sewage sludge removed from WWT plants is known, detailed information about its BOD content is not available. Therefore, the amount of BOD removed with sewage sludge is set to zero, in accordance with the default value given by the 2006 IPCC Guidelines (IPCC 2006).

Time series of the activity data are shown in Table 7-21.

(ii) Sewage gas

As elaborated above, a per capita CH₄ emission factor ($EF_{sewage\ gas}$) is calculated for CH₄ emissions from separate sewage sludge treatment, and the respective activity data is population (Table 7-21).

7.5.2.2 N₂O emissions

Methodology (5D, N₂O)

Direct N₂O emissions from centralized WWT plants and N₂O emissions from wastewater effluent are calculated in accordance with the 2006 IPCC Guidelines (IPCC 2006).

(i) N₂O emissions from WWT plants

Direct N₂O emissions from WWT plants are determined with equation 6.9 of the 2006 IPCC Guidelines (IPCC 2006):

$$N_2O_{PLANTS} = EF_{PLANT} * P * T_{PLANT} * F_{IND-COM}$$

N_2O_{PLANTS} corresponds to the total N₂O emissions from WWT plants in kg N₂O/yr, P to the population, T_{PLANT} to the degree of utilization of modern, centralized WWT plants (%), $F_{IND-COM}$ to the correction factor for industrial and commercial co-discharged protein, and EF_{PLANT} to the emission factor from the plants.

(ii) N₂O emissions from wastewater effluents

The following equation from the 2006 IPCC Guidelines (IPCC 2006) for the N₂O emissions from wastewater effluent is used:

$$N_2O_{EFFLUENT} = EF_{EFFLUENT} * N_{EFFLUENT} * 44/28$$

$N_2O_{Effluent}$ corresponds to the total N₂O emissions from effluents (kg N₂O/yr), $N_{EFFLUENT}$ to the total amount of nitrogen discharged to the aquatic environment (kg N/yr), and $EF_{EFFLUENT}$ to the emission factor for N₂O emissions from discharged wastewater (kg N-N₂O/kg N). The following equation allows for the calculation of the total amount of nitrogen in the wastewater ($N_{EFFLUENT}$, kg N/yr, IPCC 2006):

$$N_{EFFLUENT} = (P * Protein * F_{NPR} * F_{NON-COM} * F_{IND-COM}) - N_{SLUDGE} - N_{WWT}$$

P corresponds to the population, $Protein$ to the annual per capita protein consumption (kg protein/inhabitant/yr), and F_{NPR} to the fraction of nitrogen in protein. $F_{NON-COM}$ is a factor accounting for non-consumed protein added to the wastewater. $F_{IND-COM}$ is a factor accounting for industrial and commercial co-discharged protein into the sewer system. N_{SLUDGE} is the amount of nitrogen removed with sewage sludge (kg N/yr), calculated as the product of sludge amount per year and its nitrogen concentration. The default value according to the 2006 IPCC Guidelines would be zero, but detailed data about sewage sludge removal as well as the nitrogen content of the sewage sludge is available for Switzerland (Külling et al. 2002a). In Switzerland sewage sludge is mostly burnt today in waste incineration plants and (cement) industry, previously it has also been used as fertilizer (now forbidden). N_{WWT} corresponds to the amount of nitrogen directly emitted by WWT plants in form of N₂O (N_2O_{Plants} , see calculation above).

Emission factors (5D, N₂O)

(i) N₂O emissions from WWT plants

The IPCC default emission factor is applied: $EF_{PLANT} = 3.2 \text{ g N}_2\text{O/inhabitant/yr}$ (IPCC 2006).

(ii) N₂O emissions from wastewater effluents

The IPCC default emission factor is applied: $EF_{EFFLUENT} = 0.005 \text{ kg N}_2\text{O/kg N}$ (IPCC 2006).

Activity Data (5D, N₂O)

(i) N₂O emissions from WWT plants

The needed time-dependent and country-specific activity data are summarized in Table 7-21:

- Population (P)
- Degree of utilization of modern, centralized WWT plants (T_{PLANT})
- In addition, the following constant factor is used:
- Industrial/commercial co-discharged protein, IPCC default value: $F_{IND-COM} = 1.25$ (IPCC 2006)

(ii) N₂O emissions from wastewater effluents

The time-dependent and country-specific activity data are also summarized in Table 7-21:

- Population (P)
- Annual per capita protein consumption (Protein) (SBV 2015)
- Mass of nitrogen contained in the removed sludge (N_{SLUDGE})

In addition, the following constant factors are used:

- Fraction of nitrogen in protein, IPCC default value: $FNPR = 0.16 \text{ kg N/kg protein}$ (IPCC 2006).
- Factor accounting for non-consumed protein added to the wastewater, IPCC default value: $F_{NON-CON} = 1.1$ (IPCC 2006). This value is recommended for countries without garbage disposal (which holds for Switzerland as it is illegal to discharge solid and liquid garbage with the wastewater, see Article 10 in the Waters Protection Ordinance, Swiss Confederation 1998a).
- Industrial/commercial co-discharged protein, IPCC default value: $F_{IND-COM} = 1.25$ (IPCC 2006).

Table 7-21 Activity data for source category 5D Wastewater treatment and discharge.

5D Wastewater treatment and discharge	Unit	1990	1995	2000	2005
Population	persons in 1000	6'796	7'081	7'209	7'501
Fraction of population connected to wastewater treatment plants	%	90.0	93.5	95.4	96.8
Connected persons	persons in 1000	6'116	6'621	6'877	7'261
Protein consumption	kg/inhab./a	38	37	37	36
N removed with sludge (N _{sludge})	N in t/a	9'465	9'009	8'831	9'026
N directly emitted (N _{WWT})	N in t/a	15.6	16.9	17.5	18.5
Total org. degr. material (TOW)	t/a	186'041	193'842	197'346	205'340

5D Wastewater treatment and discharge	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Population	persons in 1000	7'558	7'619	7'711	7'801	7'878	7'912	7'997	8'089	8'189	8'282
Fraction of population connected to wastewater treatment plants	%	97.0	97.0	97.0	97.0	97.0	97.0	97.0	97.0	97.0	97.0
Connected persons	persons in 1000	7'331	7'390	7'480	7'567	7'642	7'675	7'757	7'846	7'943	8'034
Protein consumption	kg/inhab./a	37	37	37	38	38	39	37	37	37	37
N removed with sludge (N _{sludge})	N in t/a	9'135	9'135	9'135	9'135	9'135	9'135	9'135	9'135	9'135	9'135
N directly emitted (N _{WWT})	N in t/a	18.7	18.8	19.0	19.3	19.5	19.5	19.7	20.0	20.2	20.4
Total org. degr. material (TOW)	t/a	206'900	208'570	211'089	213'552	215'660	216'591	218'918	221'436	224'174	226'720

7.5.2.3 Other gases

The sewage gas production generates emissions of further gases from flaring: CO₂ (biogenic), NO_x, CO, NMVOC, and SO₂. The emissions are calculated by multiplying population (as activity data, see Table 7-21) with country-specific emission factors based on measurements and expert estimates, documented in EMIS 2017/5D1 Wastewater Treatment Plants and EMIS 2017/5D2 Pre-treatment of industrial wastewater. The emission factors used are summarized in Table 7-22.

Table 7-22 Emission factors of CO₂ (biogenic), CH₄, N₂O, NO_x, CO, NMVOC and SO₂ for 5D Wastewater treatment and discharge in 2015.

5D Wastewater treatment and discharge	CO ₂ biog.	N ₂ O	CH ₄	NO _x	CO	NMVOC	SO ₂
	kg/person	g/person					
5D1 Domestic wastewater	13.5	59	856	22	38	0.5	2.4
5D2 Industrial wastewater	2.0	IE	IE	2.9	4.7	0.1	0.3

7.5.3 Uncertainties and time-series consistency

Uncertainty in CH₄ and N₂O emissions from 5D

7.5.3.1 CH₄ emissions

The default values of the 2006 IPCC Guidelines (IPCC 2006) are adopted to estimate the uncertainty of CH₄ emissions. The following specifications are given:

- Activity data: Uncertainties of the single factors U(population) = 5%, U(BOD) = 30%, U(I) = 20% lead to an aggregated uncertainty of U(AD) = 36%.
- CH₄ emission factor: Uncertainties of the single factors U(B₀) = 30%, U(MCF) = 10% (well managed plants) lead to an aggregated uncertainty of U(EF) = 32%.
- Combined uncertainty U(Em CH₄) = 48%.

7.5.3.2 N₂O emissions

By applying the default uncertainties of the 2006 IPCC Guidelines (IPCC 2006, chp. 6, table 6.11) for the activity data (population, protein consumption etc.) a total uncertainty of 32% results.

For the emission factor the 2006 IPCC Guidelines provide default values, too. However, the range for $EF_{EFFLUENT}$ covers an interval of 0.0005–0.25 (with default value 0.005). If this range is interpreted as the 95% uncertainty interval, a symmetrised uncertainty of 2500% would result, which is not considered appropriate. The 2006 IPCC Guidelines (IPCC 2006) do not explain how to apply the range, wherefore the default uncertainty is not adopted. Instead, the uncertainty is based on expert judgments assuming a high uncertainty of N₂O emissions from 5D Wastewater treatment and discharge in Switzerland. By means of Table 1-10 this qualitative estimation corresponds to 150% for the combined uncertainty. This value is used for the uncertainty analyses in chp. 1.7.

Consistency: Time series for 5D Wastewater treatment and discharge are all considered consistent.

7.5.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

7.5.5 Category-specific recalculations

The following recalculations were implemented in submission 2017. Recalculations that cause a change in emission levels 1990 or 2014 of at least 0.2 kt CO₂ eq are quantified. All the other recalculations have an impact of less than 0.2 kt CO₂ eq in the years 1990 and 2014.

- 5D1: Activity data of energy produced in combined heat and power generation engines (CHP) has been recalculated for 2011–2014 in the Swiss Statistics for Renewable Energies by SFOE. This leads to changed CH₄ and CO₂ biog. emission factors (emissions per capita).
- 5D: The N₂O emission factor for wastewater handling for the year 2014 has slightly decreased because it was updated according to the last published value in the Agristat 2015 statistics published by the Swiss Farmers Association. This recalculation leads to a decrease in emissions in 2014 by 2 kt CO₂ eq.

7.5.6 Category-specific planned improvements

No category-specific improvements are planned.

7.6 Source category 5E – Other

7.6.1 Source category description

Source category 5E Other is not a key category.

The source category 5E Other comprises NMVOC and CO emissions from car shredding stemming from residues of fuels (gasoline, diesel) and motor oil in the tanks and motors of the shredded vehicles. GHG emissions do not occur.

Table 7-23 Specification of source category 5E Other (car shredding)

5E	Source	Specification
5E	Car shredding plants	Emissions from car shredding plants

7.6.2 Methodological issues

Methodology (5E)

For the emissions from car shredding a Tier 2 method is used.

Indirect CO₂ emissions from fossil CO and NMVOC emissions are described in chp. 9 Indirect CO₂ and N₂O emissions.

Emission factors (5E)

An emission factor of 100 g NMVOC per tonne of shredded vehicle is applied for the period 1990–1995. From 2000 onward, 200 g/t are used. Between 1995 and 2000 the values are linearly interpolated. The NMVOC emission factor are based on measurements at four plants in the years from 2002 to 2008 (EMIS 2017/5E Shredder Anlagen). For CO a constant emission factor is applied over the whole reporting period.

Table 7-24 CO and NMVOC emission factors for 5E Other (car shredding) in 2015.

5E Other waste	Unit	CH ₄	N ₂ O	NO _x	CO	NMVOC	SO ₂
Shredding	g/t scrap	NA	NA	NA	5	200	NA

Activity data (5E)

The waste quantities from 1990 to 1999 are provided by the Swiss Shredding Association. The data from 2000 to 2007 are taken from Swiss waste statistics. From then onwards the quantities are assumed to remain constant due to the lack of data (see also EMIS 2017/5E Shredder Anlagen).

Table 7-25 Activity data 5E Other (car shredding).

5E Other waste	Unit	1990	1995	2000	2005
Shredding	kt	280	300	300	300

5E Other waste	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Shredding	kt	300	300	300	300	300	300	300	300	300	300

7.6.3 Uncertainties and time-series consistency

Uncertainties of 20% for the emission factor and 10% for the activity data are assumed.

Consistency: Time series for 5E Other are all considered consistent.

7.6.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

7.6.5 Category-specific recalculations

No category-specific recalculations were carried out.

7.6.6 Category-specific planned improvements

No category-specific improvements are planned.

8 Other

8.1 Overview

8.1.1 Greenhouse gas emissions

Within the sector 6 Other emissions from two sources are considered:

- Fire damage estates
- Fire damage motor vehicles




Figure 8-1 Switzerland's greenhouse gas emissions in the sector 6 Other 1990–2015.

Table 8-1 Trend of total GHG emissions from sector 6 Other in Switzerland 1990–2015.

Gas	1990	1995	2000	2005
	CO ₂ equivalent (kt)			
CO ₂	11.0	10.9	11.8	12.3
CH ₄	0.7	0.6	0.6	0.6
N ₂ O	0.6	0.5	0.5	0.6
Sum	12.2	12.1	12.9	13.5

Gas	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	CO ₂ equivalent (kt)									
CO ₂	11.4	12.6	11.8	11.5	11.3	12.2	12.7	13.1	10.5	11.3
CH ₄	0.6	0.7	0.6	0.6	0.6	0.6	0.7	0.7	0.6	0.6
N ₂ O	0.5	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.5	0.5
Sum	12.5	13.8	13.0	12.6	12.4	13.4	14.0	14.4	11.5	12.5

In sector 6 Other “fire damage estates” account for most of the emissions, the rest stems from “fire damage motor vehicles”. The total greenhouse gas emissions of this sector show

variations around 12 kt CO₂ eq during the reporting period. Consequently, sector 6 Other is an emission source of minor importance for the national total.

8.2 Source category 6 – Other

8.2.1 Source category description

Source category 6 Other is not a key category.

The sources reported in source category 6 Other are depicted in Table 8-2.

Table 8-2 Specification of source category 6 Other.

6	Source	Specification
6Ad	Fire damage estates	Emissions from fires in buildings.
6Ad	Fire damage motor vehicles	Emissions from fires in motor vehicles.

8.2.2 Methodological issues

Methodology (6 Other)

a) Fire damage estates

This source category describes the emissions from fire damages in estates and its furniture. For the emissions from fire damages in estates, a Tier 2 method is used. The AD and EF are country-specific. The estimation of GHG emissions is based on damage sums and fires reported from insurance companies.

b) Fire damage motor vehicles

This source category describes the emissions from fire damages in motor vehicles. For the emissions from fire damages in motor vehicles, a Tier 2 method is used. The AD and EF are country-specific. The estimation of GHG emissions is based on damage sums and fires reported from insurance companies.

Emission factors (6 Other)

a) Fire damage estates

Emission factors for CO₂, CO, NO_x and SO₂ are country-specific based on measurements and expert estimates originally completed for illegal waste incineration. It is assumed that for fire damage in estates emissions are similar (EMIS 2017/6Ad Brand- und Feuerschäden Immobilien).

The fraction between fossil and biogenic CO₂ emissions is assumed to remain constant since 2000 with 80% being fossil and 20% biogenic CO₂ emissions. Before 2000, it is assumed

that the fraction of fossil CO₂ emissions from burnt goods has been increasing linearly from 20% in 1950 to 80% in 2000.

Indirect CO₂ emissions from fossil CO and NMVOC emissions are documented in chp. 9 Indirect CO₂ and N₂O emissions.

b) Fire damage motor vehicles

Emission factors for CO₂, CO, NO_x and SO₂ are country-specific based on measurements and expert estimates originally gained from the combustion of cable insulation materials, documented in EMIS 2017/6Ad Brand- und Feuerschäden Motorfahrzeuge.

The emission factor for CH₄ from fire damage in motor vehicles is based on EPA (1992), while N₂O emissions have not been estimated for this source category.

Indirect CO₂ emissions from fossil CO and NMVOC emissions are documented in chp. 9 Indirect CO₂ and N₂O emissions.

Table 8-3 Emission factors for fire damages in 2015 (EMIS 2017/6Ad).

6A Other	Unit	CO ₂ biogenic	CO ₂ fossil	CH ₄	N ₂ O	NO _x	CO	NMVOC	SO ₂
6Ad Fire damage estates	t / kt burned good	400	1'500	3	0.25	2	100	16	1
6Ad Fire damage motor vehicles	t / kt burned good	NO	1'500	5	NE	1.3	2	2	5

Activity data (6 Other)

a) Fire damage estates

The fire insurance association of the cantons (Vereinigung kantonaler Feuerversicherungen, VKF) publishes the number of fire incidents in buildings each year and the total sum of monetary damage. Data from 1992 to 2001 shows that the average damage sum per fire incident in buildings amounts to approx. CHF 20'000. It is assumed that this corresponds to 780 kg of flammable material per case. It is further assumed that in average only 50% of the material actually burns down during an incident because of the intervention of the fire brigade. Thus, an amount of 400 kg of burnt material per fire case is estimated. With these assumptions, the amount of burnt material for each year can be calculated from the total sum of monetary damage published by VKF (EMIS 2017/6Ad Brand- und Feuerschäden Immobilien).

b) Fire damage motor vehicles

Based on data from a Swiss insurance company with 25% market share in 2002, the number of reported cases of fire damage to vehicles was extrapolated to the total vehicle number in Switzerland. It was estimated that one fire case per 790 vehicles occurs per year, remaining constant within the reporting period. Applying this ratio to the actual vehicle number, which is annually published by the Swiss Federal Statistical Office, the total number of fire incidents with vehicles in Switzerland is obtained for each year (EMIS 2017/6Ad Brand- und

Feuerschäden Motorfahrzeuge). During a car fire incident, a car burns down only partially. It is assumed that approx. 100 kg of material burns down during a car fire. With these assumptions, the total number of material burnt can be calculated from the total number of cars in Switzerland.

Table 8-4 Activity data: Burnt goods from 1990 to 2015 (documented in EMIS 2017/6Ad).

6A	Unit	1990	1995	2000	2005
6Ad Fire damage estates	kt	8.0	7.3	7.3	7.6
6Ad Fire damage motor vehicles	kt	0.48	0.52	0.58	0.64

6A	Unit	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
6Ad Fire damage estates	kt	6.9	7.7	7.2	7.0	6.8	7.4	7.8	8.0	6.3	6.8
6Ad Fire damage motor vehicles	kt	0.65	0.66	0.66	0.67	0.68	0.69	0.71	0.72	0.73	0.75

8.2.3 Uncertainties and time series consistency

Uncertainties of CO₂, CH₄ and N₂O emissions are estimated to be high (according to Table 1-10).

Consistency: Time series for 6Ad Fire damages are all considered consistent.

8.2.4 Category-specific QA/QC and verification

The general QA/QC measures are described in chp. 1.2.3.

8.2.5 Category-specific recalculations

- 6Ad: Activity data for fire damages motor vehicles has changed for the years 2003–2014 because the model has been updated with vehicle data provided by the SFOS.
- 6Ad: Activity data for fire damages estates has changed for the years 1996–2014 because the model has been updated with damage sums provided by the Swiss Building Insurance Association.

8.2.6 Category-specific planned improvements

No category-specific improvements are planned.

9 Indirect CO₂ and N₂O emissions

9.1 Overview


In this chapter, indirect CO₂ emissions that result from the atmospheric oxidation of NMVOC and CO as well as indirect N₂O emissions that are induced by the deposition of NO_x and NH₃ are documented. While indirect CO₂ emissions reported in this chapter are accounted for in the national total, indirect N₂O emissions are not.

Indirect emissions of CO₂ and N₂O are shown in CRF table 6, together with the emissions of the precursor gases CH₄, CO, NMVOC, NO_x and NH₃. While all emissions of precursor gases are shown in both CRF table 6 and in the respective sectors, the indirect emissions of CO₂ and N₂O shown in CRF table 6 only represent emissions not already included together with direct emissions in other sectors (in order to avoid double counting). Further, in the case of indirect CO₂ emissions, only carbon of fossil origin is considered. Accordingly, while e.g. NMVOC and CO of biogenic origin are shown as precursor gases in CRF table 6, they are not included for the calculation of indirect CO₂ emissions. Consequently, the implied emission factors may vary from sector to sector and also from year to year. Indirect CO₂ emissions resulting from the atmospheric oxidation of CH₄ are generally not considered.

Chapter 9.2 explains in detail the methodological issues to derive indirect CO₂ and N₂O emissions based on the emissions of the precursor gases NMVOC and CO, as well as NO_x and NH₃ from the different sectors. As an overview, the resulting indirect CO₂ emissions are shown in Table 9-1, as well as in Figure 9-1 and Figure 9-2. The resulting indirect N₂O emissions are shown in Table 9-2, as well as in Figure 9-3 and Figure 9-4.

Indirect CO₂ emissions are considered for both the uncertainty analysis (see chp. 1.6) and the key category analysis (see chp. 1.5).

Indirect N₂O emissions are not considered for the uncertainty analysis (see chp. 1.6), nor for the key category analysis (see chp. 1.5).

Figure 9-1 Switzerland's indirect fossil CO₂ emissions 1990–2015.Table 9-1 Indirect fossil CO₂ emissions from 1990 to 2015.

Indirect fossil CO ₂ emissions by source category	1990	1995	2000	2005
	(kt CO ₂)			
1 Energy	43.74	27.64	17.04	13.24
1B Fugitive emissions from fuels	43.74	27.64	17.04	13.24
2 Industrial processes and product use	365.12	243.91	167.94	113.87
2A Mineral industry	0.15	0.12	0.11	0.11
2B Chemical industry	1.34	0.39	0.05	0.06
2C Metal industry	3.36	2.41	2.21	1.26
2D Non-energy products from fuels and solvent use	231.91	140.12	93.54	55.90
2G Other product manufacture and use	125.53	98.86	70.11	55.23
2H Other	2.83	2.01	1.93	1.30
5 Waste	2.01	1.66	1.60	1.38
5A Solid waste disposal	NO	NO	NO	NO
5C Waste incineration and open burning of waste	1.95	1.56	1.47	1.25
5E Other	0.06	0.10	0.13	0.13
6 Other	1.04	1.03	1.11	1.16
6Ad Fire damages	1.04	1.03	1.11	1.16
Total	411.92	274.23	187.69	129.65

Indirect fossil CO ₂ emissions by source category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	(kt CO ₂)									
1 Energy	13.51	12.38	12.54	11.97	11.05	10.81	9.60	11.03	10.82	5.14
1B Fugitive emissions from fuels	13.51	12.38	12.54	11.97	11.05	10.81	9.60	11.03	10.82	5.14
2 Industrial processes and product use	111.87	108.23	108.95	109.01	110.05	109.38	108.65	106.39	106.63	105.82
2A Mineral industry	0.11	0.12	0.11	0.11	0.12	0.12	0.11	0.10	0.10	0.09
2B Chemical industry	0.06	0.06	0.08	0.06	0.09	0.06	0.06	0.06	0.05	0.05
2C Metal industry	1.22	1.26	1.35	0.85	0.93	1.05	0.81	0.81	0.78	0.68
2D Non-energy products from fuels and solvent use	54.94	53.14	53.45	53.77	54.23	53.12	51.38	50.81	51.01	50.14
2G Other product manufacture and use	53.75	52.03	52.15	52.22	52.41	52.52	54.21	52.60	52.74	52.86
2H Other	1.79	1.62	1.81	2.01	2.27	2.51	2.09	2.02	1.95	2.01
5 Waste	1.42	1.38	1.39	1.29	1.30	1.25	1.24	1.22	1.18	1.18
5A Solid waste disposal	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
5C Waste incineration and open burning of waste	1.29	1.25	1.26	1.15	1.16	1.11	1.10	1.08	1.05	1.05
5E Other	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
6 Other	1.06	1.18	1.10	1.07	1.04	1.13	1.19	1.22	0.96	1.04
6Ad Fire damages	1.06	1.18	1.10	1.07	1.04	1.13	1.19	1.22	0.96	1.04
Total	127.86	123.17	123.97	123.34	123.44	122.57	120.68	119.86	119.59	113.18




Figure 9-2 Relative trends of the indirect fossil CO₂ emissions in the period 1990–2015 by sector. The base year 1990 represents 100%.




Figure 9-3 Switzerland's indirect N₂O emissions 1990–2015.

Table 9-2 Indirect N₂O emissions from 1990 to 2015.

Indirect N ₂ O emissions by source category	1990	1995	2000	2005
	(kt)			
1 Energy	1.70	1.44	1.40	1.21
1A Fuel combustion activities	1.70	1.44	1.39	1.21
1B Fugitive emissions from fuels	0.00	0.00	0.00	0.00
2 Industrial processes and product use	0.02	0.01	0.02	0.01
2A Mineral industry	0.00	0.00	0.00	0.00
2B Chemical industry	0.00	0.00	0.00	0.00
2C Metal industry	0.00	0.00	0.00	0.00
2G Other product manufacture and use	0.01	0.01	0.01	0.01
2H Other	0.01	0.00	0.01	0.00
5 Waste	0.03	0.03	0.03	0.03
5A Solid waste disposal	0.02	0.01	0.01	0.01
5B Biological treatment of solid waste	0.00	0.00	0.00	0.00
5C Waste incineration and open burning of waste	0.00	0.00	0.00	0.00
5D Wastewater handling and discharge	0.00	0.01	0.01	0.01
6 Other	0.00	0.00	0.00	0.00
6Ad Fire damages	0.00	0.00	0.00	0.00
Total	1.75	1.48	1.44	1.25

Indirect N ₂ O emissions by source category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
	(kt)									
1 Energy	1.17	1.13	1.11	1.05	1.03	0.96	0.96	0.95	0.89	0.83
1A Fuel combustion activities	1.17	1.13	1.11	1.05	1.03	0.96	0.96	0.95	0.89	0.83
1B Fugitive emissions from fuels	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2 Industrial processes and product use	0.01									
2A Mineral industry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2B Chemical industry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2C Metal industry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2G Other product manufacture and use	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2H Other	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.01	0.00
5 Waste	0.03									
5A Solid waste disposal	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5B Biological treatment of solid waste	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
5C Waste incineration and open burning of waste	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5D Wastewater handling and discharge	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
6 Other	0.00									
6Ad Fire damages	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	1.21	1.17	1.15	1.09	1.07	1.00	1.00	0.98	0.93	0.87




Figure 9-4 Relative trends of the indirect N₂O emissions in the period 1990–2015 by sector. The base year 1990 represents 100%.

9.2 Methodological issues

9.2.1 Methodological issues to derive indirect CO₂ emissions

Indirect CO₂ emissions from the atmospheric oxidation of NMVOC are calculated based on the stoichiometric conversion (carbon content fraction * molecular weight of carbon dioxide / molecular weight of carbon). Thereby, a constant carbon content of NMVOC of 60% is assumed, based on the 2006 IPCC Guidelines (IPCC 2006). Indirect CO₂ emissions from the atmospheric oxidation of CO are also calculated based on the stoichiometric conversion (molecular weight of carbon dioxide / molecular weight of carbon monoxide). Thus, indirect CO₂ emissions (Em) result from the following equations:

$$Em_{CO_2, \text{ indirect from NMVOC}} = Em_{NMVOC, \text{ fossil}} * 0.6 * 44/12$$

$$Em_{CO_2, \text{ indirect from CO}} = Em_{CO, \text{ fossil}} * 44/28$$

Activity data for the calculation of indirect CO₂ emissions

Activity data to calculate indirect CO₂ emissions consists of NMVOC and CO emissions as reported in each individual sector and source category, carefully excluding NMVOC and CO emissions of biogenic origin and emissions already included as direct (CO₂) emissions (e.g. when using an oxidation factor of 100%). For the different sectors and source categories, the situation is as follows:

1A Energy: Since according to the 2006 IPCC Guidelines (IPCC 2006) emission factors in source category 1A Energy are based on the assumption of complete oxidation (100%), CO₂ resulting from the atmospheric oxidation of CO and NMVOC emitted from this source category is already accounted for in the corresponding emission factors for direct CO₂ emissions. The respective emissions are thus implicitly reported as direct CO₂ emissions in 1A and no indirect CO₂ emissions from 1A Energy are reported (see chp. 3.2.4.4.1).

1B Fugitive emissions from fuels: CO₂ resulting from the atmospheric oxidation of NMVOC and CO emitted from source category 1B is reported as indirect CO₂ emissions unless it is already accounted for implicitly as direct CO₂ emissions in 1B (chp. 3.3). For 1B, Table 9-3 illustrates in which processes CO and NMVOC emissions occur, and whether the related CO₂ emissions are reported as indirect CO₂ emissions or implicitly as direct CO₂ emissions. In summary, all CO₂ resulting from the atmospheric oxidation of CO emitted from 1B is implicitly included in 1B as direct CO₂ and is therefore not reported as indirect CO₂. CO₂ resulting from the atmospheric oxidation of NMVOC emitted from 1B is reported as indirect CO₂, except for CO₂ from source category 1B2c, where an oxidation factor of 100% is applied to calculate direct CO₂ emissions.

Table 9-3 Sources of indirect CO₂ emissions from source category 1B Fugitive emissions from fuels.

Source category name	CO emissions	NMVOC emissions
1B1a Coal mining and handling	-	-
1B2a Oil		
1B2a iii Oil – Transport	-	Indirect CO ₂ emissions reported in chp. 9
1B2a iv Oil – Refining/storage:	<i>Leakage:</i> -	<i>Leakage:</i> Indirect CO ₂ emissions reported in chp. 9
	<i>Crude oil:</i> -	<i>Crude oil:</i> -
1B2a v, Oil - Distribution of oil products:	<i>Gasoline storage tank:</i> -	<i>Gasoline storage tank:</i> Indirect CO ₂ emissions reported in chp. 9
	<i>Gasoline station:</i> -	<i>Gasoline station:</i> Indirect CO ₂ emissions reported in chp. 9
1B2b Natural gas		
1B2b ii, Natural gas - Production	-	Indirect CO ₂ emissions reported in chp. 9
1B2b iv, Natural gas – Transmission and storage	-	Indirect CO ₂ emissions reported in chp. 9
1B2b v, Natural gas - Distribution	<i>Distribution:</i> -	<i>Distribution:</i> -
	<i>Leakage gas pipeline:</i> -	<i>Leakage gas pipeline:</i> Indirect CO ₂ emissions reported in chp. 9
	<i>Other leakage:</i> -	<i>Other leakage:</i> Indirect CO ₂ emissions reported in chp. 9
1B2b vi, Other (losses due to major accidents)	-	Indirect CO ₂ emissions reported in chp. 9 (Emissions occur only in 2010 and 2011)
1B2c Venting and flaring	<i>Flaring:</i> CO ₂ emissions from CO are reported in 1B2c a as direct CO ₂ emissions (CO ₂ EF assumes an oxidation of 100%)	<i>Flaring:</i> CO ₂ emissions from NMVOC are reported in 1B2c a as direct CO ₂ emissions (CO ₂ EF assumes an oxidation of 100%)
	<i>H₂ Production:</i> -	<i>H₂ Production:</i> -

2 Industrial processes and product use: Except for CO and NMVOC emissions from 2C1 Secondary steel production, electric arc furnaces and CO emissions from anodes in source category 2C3 Primary aluminium production, none of the CO₂ emissions resulting from the atmospheric oxidation of NMVOC and CO emitted from sector 2 are already considered under the direct CO₂ emissions of this sector. The CO₂ emissions from 2C1 Secondary steel production, electric arc furnace are based on a carbon mass balance considering all carbon

sources and sinks of the process. Therefore, the emissions of both CO and NMVOC are not accounted for as indirect CO₂ emissions from sector 2 IPPU. For CO emissions from 2C3 Primary aluminium production full oxidation of the anodes is assumed and these emissions are therefore not included in the indirect emissions (see chp. 4.4.2.2). On the other hand the NMVOC emissions from 2C3 Primary aluminium production originate solely from the production of the electrodes at the plants therefore, they have to be considered for the calculation of indirect CO₂ emissions.

In addition, before indirect CO₂ emissions are calculated, biogenic NMVOC and CO need to be subtracted from the total NMVOC and CO emissions from sector 2. Biogenic NMVOC and CO emissions occur in the source categories 2H2 Food and beverages industry and 2G4 Other (tobacco consumption only), see chp 4.2.

3 Agriculture and 4 LULUCF: NMVOC and CO emissions from the sectors 3 Agriculture and 4 LULUCF are of biogenic origin. Accordingly, no indirect CO₂ emissions are reported for these sectors.

5 Waste: NMVOC and CO emissions from sector 5 Waste contain fossil and biogenic shares. Only CO₂ resulting from the atmospheric oxidation of fossil NMVOC and CO is reported as indirect CO₂. Emissions of fossil NMVOC and CO stem from the following processes:

- Landfills with open burning: Partly fossil, 0 since 1990.
- Illegal waste incineration: Partly fossil, fossil share is assumed to be the same as for waste incinerated in MWIP, see chp. 7.4 Incineration and open burning of waste (5C1).
- Insulation material from cables: See chp. 7.4 Incineration and open burning of waste (5C1).
- Hospital waste incineration: See chp. 7.4 Incineration and open burning of waste (5C1).
- Shredding: See chp. 7.6 Other.

6 Other: NMVOC and CO emissions from sector 6 Other contain fossil and biogenic shares. Only CO₂ resulting from the atmospheric oxidation of fossil NMVOC and CO is reported as indirect CO₂. Emissions of fossil NMVOC and CO stem from the following processes:

- Fire damage estate: Partly fossil, see chp. 8.2 (6Ad).
- Fire damage motor vehicles: See chp. 8.2 (6Ad).

9.2.2 Methodological issues to derive indirect N₂O emissions

Indirect N₂O emissions are estimated using a country-specific method according to a study of indirect N₂O emissions induced by nitrogen deposition in Switzerland (Bühlmann 2014, Bühlmann et al. 2015). In this study, ecosystem-specific emission factors for indirect N₂O resulting from nitrogen deposition were developed, based on a comprehensive literature survey. Thereby, the land cover types forests, grassland and wetlands were distinguished. In

a next step, the ecosystem-specific emission factors were combined with a highly-resolved nitrogen deposition map of Switzerland as well with the geo-referenced dataset of the Swiss Land Use Statistics (allowing for the localisation and estimation of spatial extent of the different ecosystems). This resulted in detailed and spatially resolved indirect N₂O emissions for Switzerland. To facilitate a simple application in the greenhouse gas inventory, the resulting total emissions were used to come up with a total emission factor expressed as indirect N₂O-N per N-deposition (deposited in form of NO_x or NH₃, see also chp. 5.3.2.4). The resulting total emission factor is in the order of 2.5% and slightly varies with time as the shares of the different ecosystems are not constant over time. Based on this country-specific emission factor, higher indirect N₂O emissions result compared to the emissions that would result by applying the 2006 IPCC Guidelines (IPCC 2006, see also Bühlmann et al. 2015).

To calculate indirect N₂O emissions induced by the deposition of NO_x and NH₃ according to Bühlmann (2014) and Bühlmann et al. (2015), total N-deposition is needed. It is derived from NO_x (which is always reported in NO₂ equivalents) and NH₃ emissions using the stoichiometric conversion according to the following equation:

$$\text{Mass-N} = \text{Mass-NO}_{2,\text{eq}} * 14/46 + \text{Mass-NH}_3 * 14/17$$

Activity data for the calculation of indirect N₂O emissions

The activity data to calculate indirect N₂O emissions from a specific sector corresponds to the NO_x and NH₃ emissions reported in the respective source categories. However, the following exceptions need to be considered:

- Indirect N₂O emissions from sector 3 Agriculture are reported in the respective sector together with direct N₂O emissions (chp. 5.3.2.5.4 Volatilisation of NH₃ and NO_x from manure management systems, chp. 5.5.2.3 Indirect N₂O emissions from atmospheric deposition of N volatilised from managed soils (3Db1) and chp. 5.5.2.4 Indirect N₂O emissions from leaching and run-off from managed soils (3Db2)).
- For sector 6, the only indirect N₂O emissions to be considered are those resulting from NO_x emissions in source category 6Ad Fire damages. All other indirect N₂O emissions are included in sector 3 Agriculture together with direct N₂O emissions.

9.2.3 Uncertainties and time series consistency

Indirect CO₂ emissions are included in the uncertainty analysis, but indirect N₂O emissions are not included.

Uncertainties of indirect CO₂ emissions are based on respective uncertainties of NMVOC and CO emissions. Uncertainties of NMVOC emissions are estimated based on AD and EF uncertainties as documented in Switzerland's Informative Inventory Report 2017 (FOEN 2017f). The estimated uncertainties do not distinguish between fossil and biogenic shares. Uncertainties of CO emissions are estimated based on AD uncertainties documented in Switzerland's Informative Inventory Report 2017 (FOEN 2017f) and EF uncertainties are based on expert judgements and uncertainties provided by the EMEP/EEA Guidebook.

Combined uncertainties of indirect CO₂ emissions amount to 22% for indirect CO₂ emissions from sector 1 Energy, 28% for indirect CO₂ emissions from sector 2 IPPU, 48% for indirect CO₂ emissions from sector 5 Waste and 60% for indirect CO₂ emissions from sector 6 Other.

Consistency: Time series for indirect CO₂ and N₂O emissions are all considered consistent.

9.2.4 Category-specific QA/QC and verification

The same QA/QC and verification procedures are conducted as for NMVOC, CO, NO_x and NH₃ related source categories in chp. 3 Energy, 4 Industrial processes and product use, 7 Waste and 8 Other.

9.2.5 Category-specific recalculations

- See NMVOC related recalculations reported in chp. 3 Energy, 4 Industrial processes and product use, 7 Waste and 8 Other.
- See CO related recalculations reported in chp. 4 Industrial processes and product use, 7 Waste and 8 Other.
- See NO_x and NH₃ related recalculations reported in chp. 3 Energy, 4 Industrial processes and product use, 7 Waste and 8 Other.
- The CO₂ emissions from 2C1 Secondary steel production, electric arc furnace are based on a carbon mass balance considering all carbon sources and sinks of the process. Therefore, the emissions of CO and NMVOC are no longer included in the calculation of the indirect CO₂ emissions from sector 2 IPPU in order to avoid a double counting.

Recalculations implemented in Switzerland's Saturday paper 2016:

- 2C3: The ERT found that the CO₂ emissions from aluminium production (direct and indirect) are potentially overestimated for the years 1990-2006. Switzerland eliminated any double-counting of CO₂ by removing the indirect CO₂ emissions from oxidation of CO from 2C3 Primary aluminium production. Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

9.2.6 Category-specific planned improvements

No category-specific improvements are planned.

10 Recalculations and improvements

10.1 Explanations and justifications for recalculations, including in response to the review process

10.1.1 Recommendations and encouragements from ERT and implementation

The Inventory Development Plan (IDP) is regularly updated, based on the “Reports of the individual review of the greenhouse gas inventory of Switzerland” (e.g. UNFCCC 2017), the outcome of domestic reviews, and other suggestions and feedback from the QA/QC procedures implemented in the inventory preparation process. The IDP represents the main instrument for continuous improvement of the Swiss GHG inventory in subsequent inventory cycles. It includes suggestions and recommendations for recalculations that have an impact on emission levels in the corresponding sectors.

The processing of the most recent expert review team’s recommendations and encouragements in the course of inventory preparation and compilation led to several recalculations and improvements (Table 10-1 and Table 10-2). Further recalculations had to be carried out due to improvements in some sectors. The details are explained below. An extensive list with all detailed recalculations and specifics of the recalculations is compiled by the EMIS experts and available to the reviewers on demand (in German/French only).

Due to unresolved technical issues around the CRF Reporter Switzerland was not in a position to provide correct reporting tables CRF Summary3s1 and Summary3s2 recommended by the ERT in issue ID#G.1 in UNFCCC (2017). To address the issue manually corrected reporting tables are presented in Annex 6 (Table A – 35 and Table A – 36).

Table 10-1 Recommendations from the ERT based on the Draft Review Report (UNFCCC 2017) and explanations for implementation 2016/2017.

ID	classification	recommendation	Detail = Answer including ref to chapter in NIR
Unresolved issues according to Table 3: Status of implementation of issues and/or problems raised in the previous review report of Switzerland			
G.1	General	Report the same and correct information in the CRF table summary table 3 and the NIR and improve the QC procedures at the final stage of the inventory compilation process of the annual submission	Some discrepancies between the NIR and the CRF tables are still detected and shown in a non-exhaustive list in Annex 6. To address issue #G.1 manually filled reporting tables CRF Summary 3s1 and Summary 3s2 are also presented in Annex 6.
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
General			
G.5	General - Key category analysis	The ERT recommends that Switzerland include indirect CO2 emissions in its key category analysis	chapter 1.5. Indirect CO2 emissions are taken into account in the KCA.
G.6	General - Uncertainty analysis	The ERT recommends that Switzerland include indirect CO2 emissions in its uncertainty analysis	chapter 1.6 Indirect CO2 emissions are taken into account in the uncertainty analysis.
G.7	General - Uncertainty analysis	The ERT recommends that Switzerland improve the transparency of its use of the uncertainty analysis to improve the inventory	The use of the uncertainty analyses for improving the inventory is described in chp. 1.6.1.4.
Energy			
E.8	International bunkers and multilateral operations – liquid fuels – CO2, CH4 and N2O	The ERT welcomes Switzerland's efforts in addressing this issue and recommends that it transparently report the recalculations of liquid fuel consumption and associated GHG emissions from international bunkers	Has been implemented for jet kerosene of international aviation bunkers, they are now identical in both CRF-tables (1.A(b) abd 1.D).
E.9	1.A.1.a Public electricity and heat production – other fuels – CO2	The ERT recommends that Switzerland include in its NIR the additional information to justify the application of 0.99 as the oxidation factor of the combustion of MSW in waste incineration power plants	Justification has been added to chapter 3.2.5.2.1
E.10	1.A.1.a Public electricity and heat production – other fuels – CH4	The ERT recommends that Switzerland either estimate and include in the inventory CH4 emissions from waste incineration based on the above-mentioned study, or report emissions as "NE" instead of "NA" and provide a justification in the NIR, consistent with the UNFCCC Annex I inventory reporting guidelines, that these emissions are considered insignificant	As explained in chapter 3.2.5.2.1, CH4 measurements in the exhaust gas of the MSWIP showed that they were mostly below the detection limit of 0.3 ppm. Based on this evidence, CH4 emissions from waste incineration are considered not applicable (NA).
E.11	1.A.1.b Petroleum refining, 1.A.4 Other sectors, 1.B.2.a Oil all fuels – CO2, CH4 and N2O	The ERT recommends that Switzerland improve the reporting of the level of the tier approach that is applied for petroleum refining, other sectors and oil transport in the NIR	chp. 3.2.5.2.2.: The description of the tier level for 1A1b has been changed accordingly. chp. 3.2.7.2.: The description of the tier level for 1A4 (stationary) has been changed accordingly. chp. 3.3.3.2.: The description of the tier level for 1B2a, Oil, has been changed accordingly.
E.12	1.A.2.a Iron and steel – limestone use – CO2	The ERT recommends that Switzerland either estimate and include in the inventory the CO2 emissions associated with limestone use in cupola furnaces, or report these emissions as "NE", indicate in the documentation box that they are considered insignificant and provide a justification in the NIR, consistent with the UNFCCC Annex I inventory reporting guidelines, that these emissions are considered insignificant	Chapter 4.2.2.4. CO2 emissions from limestone use in cupola furnaces are included in 2A4d Other use of carbonates.
E.13	1.A.2.d Pulp, paper and print – biomass – CH4 and N2O	The ERT recommends that Switzerland estimate and report CH4 and N2O emissions from biomass used as fuel in cellulose production in the period 1990–2008	CH4 and N2O emissions from biomass as fuel used in cellulose production have been estimated and reported in CRF Table 1.A(a)s2. See documentation of recalculation in chapter 3.2.6.5.
E.14	1.A.2.f Non-metallic minerals – biomass – CH4	The ERT recommends that Switzerland change the reported notation key for CH4 emissions from biomass used as fuel in non-metallic minerals from "NO" to "IE" for the years 1990–1999 and explain where the emissions are reported	The reported notation key for CH4 emissions from biomass used as fuel in non-metallic minerals has been changed from "NO" to "IE" for the years 1990–1999. In chp. 3.2.6.2.7, it is stated that the CH4 emission factor includes the overall CH4 emissions of the cement industry based on direct exhaust measurements at the chimneys of the cement plants. Therefore, these CH4 emissions are reported under the fuel type other fossil fuels in the CRF-tables.
E.15	1.A.3.b Road transportation – biomass – CO2, CH4 and N2O	The ERT welcomes Switzerland's efforts in addressing this issue and recommends that the Party estimate accurately CO2, CH4 and N2O emissions from biodiesel used in road transportation	As stated in chapter 3.2.9.6, the road transportation model is undergoing a re-assessment. The updated model will be used to estimate emissions from 1A3b in future submissions.

Table 10-1 continued.

ID	classification	recommendation	Detail = Answer including ref to chapter in NIR
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
Energy			
E.16	1.A.3.b.i Cars – 1.A.3.b.ii Light duty trucks gasoline and diesel – N2O	The ERT recommends that Switzerland explain the calculation of N2O emissions from cold start in road transportation in its NIR	A brief description has been added to chapter 3.2.9.2.2.
E.17	1.A.3.b.ii Light duty trucks – diesel – N2O	The ERT recommends that Switzerland change the reported notation key for N2O emissions of diesel from “NO” to “NA” for the years 1990–1995	A numerical value is provided in the CRF tables for N2O emissions from light duty trucks - diesel.
E.18	1.B.2.b Natural gas – gaseous fuels – CO2 and CH4	The ERT recommends that Switzerland recalculate CO2 and CH4 emissions of natural gas production for the years 1990–1994 by using EFs in line with the 2006 IPCC Guidelines	The default emission factors from 2006 IPCC Guidelines have been implemented in the inventory as described in chapter 3.3.4.2.
E.19	1.B.2.c Venting and flaring – natural gas – CO2, CH4 and N2O	The ERT recommends that Switzerland estimate and report flaring CO2, CH4 and N2O emissions from flaring of natural gas by using a methodology consistent with the 2006 IPCC Guidelines.	The default emission factors from 2006 IPCC Guidelines have been implemented in the database. Unfortunately they are not included in the CRF tables of this submission. This will be corrected in next submission. A methodological description can be found in chapter 3.3.5.2.
IPPU			
I.3	2. General (IPPU) – CO2	The ERT recommends that Switzerland improve the transparency of the reporting of indirect CO2 emissions from the IPPU sector by including detailed information on the AD and methodology used for the estimation	chp. 9.2.1 The description of indirect emissions from IPPU was improved. see also I.5 and I.6
I.4	2. General (IPPU) – HFCs, PFCs and SF6	The ERT recommends that the Party improve the description of the role of the data from Jungfrauoch station as a provider of verification data, not the input data for the inventory	The description of the project at Jungfrauoch has been revised to better reflect the fact, that the data is used as independent verification of the emission estimates used in the greenhouse gas inventory (Annex 5.1)
I.5	2.C.1 Iron and steel production – CO2	The ERT recommends that Switzerland either reallocate process emissions from iron processing in cupola furnaces from category 1.A.2.a to category 2.C.1 iron and steel production or, if that split is not possible, report these emissions as “IE” under category 2.C.1 and explain where they are reported	Chapter 3.2.6.2.2., Chapter 4.4.2.1. Since other bituminous coal first of all acts as fuel in cupola furnaces it was decided to report its CO2 emissions in source category 1A2a.
I.6	2.C.3 Aluminium production – CO2	The ERT recommends that the Party explain how indirect CO2 emissions from aluminium production are estimated and how it ensures that there is no double counting of emissions between the direct and indirect CO2 in its NIR	I.6 was addressed in chp. 4.4.2.2. and the Saturday Paper 2016, see Annex 7
I.7	2.C.3 Aluminium production – PFCs	2.C.3 Aluminium production – PFCs The ERT recommends that the Party include in its NIR more detailed information on the analysis of the measurements resulting in a lower EF for PFC emissions from aluminium production	The text in chapter 4.4.2.2 has been revised to clarify the issue as much as possible. Given that the company closed down in 2007 and that no additional information was found in the archives, it will not be possible to provide any additional information regarding this issue.
I.8	2.C.3 Aluminium production – HFCs, PFCs and SF6	The ERT recommends that Switzerland correct the description of the allocation of SF6 emissions from aluminium foundries in 2005 in the NIR to ensure consistency with the CRF tables	The description of the SF6 emissions from aluminium foundries are adapted and placed under 2.C.3 Aluminium production in the NIR (chp. 4.4.2.2)
I.9	2.E.1 Integrated circuit or semi- conductor – PFCs	The ERT recommends that Switzerland explain in detail how PFC emissions (especially CF4 emissions) from integrated circuits or semiconductors originate, including which species are converted into other species	In CRF table 2(II)B-Hs1, there is an error in the calculation of the implied emissions factor. Recovery is considered part of the emissions, as a result all substances show an implied emissions factor of 100% and higher for CF4 due to additional amount of CF4 from the transformation of other substances. The definition of recovery in CRF tables and its use for the calculation of the implied emissions factor are not coherent. To correct this, recovery was set NA.
I.10	2.F.1 Refrigeration and air conditioning – HFCs, PFCs	The ERT recommends that Switzerland exclude Liechtenstein when estimating HFC and PFC emissions from commercial and industrial refrigeration in the period 1991–2007	The import amount of refrigerants was adjusted for the period 1991 to 2007 to account for refrigerant use in Liechtenstein (chapter 4.7.2.1 and 4.7.5)
I.11	2.F.1 Refrigeration and air conditioning – HFCs, PFCs	The ERT recommends that Switzerland continue its efforts to acquire statistical data to allow the reporting of emissions to be split between industrial and commercial refrigeration or, if this is not possible, report the appropriate notation key “IE” for HFC and PFC emissions from industrial refrigeration with the information that emissions from that category are reported under commercial refrigeration	Efforts are ongoing to acquire statistical data to allow the reporting of emissions to be split between industrial and commercial refrigeration. So far the data quality is not sufficient. In CRF Table 2(II)B-Hs2, information is provided in the documentation box 2.F.1 that industrial refrigeration is included under commercial refrigeration.
I.12	2.F.1 Refrigeration and air conditioning – HFCs, PFCs	The ERT recommends that Switzerland improve the description of the assumptions made in the estimates for HFC and PFC emissions from refrigeration and air conditioning (2.F.1), especially for parameters that are not within the range given by the 2006 IPCC Guidelines	In chp. 4.7.2 the description of the model approach was improved and plausibility of lower emissions factors given a focus.

Table 10-1 continued.

ID	classification	recommendation	Detail = Answer including ref to chapter in NIR
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
Agriculture			
A.3	3. General (agriculture)	The ERT recommends that Switzerland correct the information on methodologies and EFs for N ₂ O emissions from manure management and agricultural soils in CRF table summary 3s2 to make it consistent with the EFs and methodologies actually used in the estimations	CRF table Summary3s2 has been updated accordingly.
A.6	3.B.4 Other livestock – CH ₄	The ERT recommends that Switzerland provide relevant supporting information in the NIR on the choice of the VS value used to estimate CH ₄ emissions from manure management of horses	VS excretion of horses has been changed to 1.90 kg/head/day based on equation 10.24 of the 2006 IPCC Guidelines. See NIR chp. 5.3.2.2.1.
A.7	3.D.a.3 Crop residues – N ₂ O	The ERT recommends that Switzerland provide relevant explanations on the assumptions used to estimate nitrogen input from crop residues on pastures in the NIR	An explanation is provided in NIR chp. 5.5.2.2.2.
A.8	3.D.a.5 Mineralization/immobilization associated with loss/gain of soil organic matter – N ₂ O	The ERT recommends that Switzerland provide in the NIR a clear indication on the usage of net carbon losses to estimate direct N ₂ O emissions from mineralization of soil organic matter	An explanation is provided in NIR chp. 5.5.2.2.2.
A.9	3.G Liming – CO ₂	The ERT recommends that Switzerland estimate CO ₂ emissions from liming taking into account the limestone and dolomite used	Total lime application was split into calcium carbonate and dolomite based on available data on Ca(CO ₃) application and the assumption that the remaining lime consists of 50% calcium carbonate and 50% dolomite. See NIR chp. 5.8.
LULUCF			
L.6	Land representation	The ERT recommends that Switzerland clarify in the NIR that all its lands are managed, or provide its definition of managed and unmanaged land and their areas over time	L.6 was addressed in chp. 6.1.3.1, chp. 6.2.1, and in chp. 6.4.1.
L.7	Land representation	In order to increase the transparency of the information on the identification of IPCC land use categories, the ERT recommends that the Party improve the description of the identification of the country-specific combination categories (i.e. land use and land-use change categories that are more detailed than those defined by the 2006 IPCC Guidelines)	L.7 was listed under planned improvements in chp. 6.3.6.
L.8	Land representation	In order to enhance the transparency of the reporting, the ERT recommends that Switzerland use the terms “afforestation” and “reforestation” in the NIR only when referring to activities under Article 3, paragraph 3, of the Kyoto Protocol	L.8 was addressed in chp. 6.1.3.1.
L.9	4. General (LULUCF) – CO ₂	The ERT recommends that the Party transparently report its definition of organic soils to estimate and report the carbon stock changes in organic soils	L.9 was addressed in chp. 6.2.2.
L.10	4. General (LULUCF) – CO ₂	The ERT recommends that Switzerland improve the transparency of the description of the equations used for calculations in the NIR by clarifying the meaning of areas (A _i ,b _a) used in the equations	L.10 was addressed in chp. 6.1.3.2.
L.11	4.A Forest land – CO ₂	In order to enhance the accuracy of the estimates of emissions and removals from forest land, the ERT recommends that Switzerland identify the areas of drained organic soils in forests accurately by collecting data on areas of organic soils under forest land affected by past draining activities	L.11 was listed under planned improvements in chp. 6.4.6.
L.13	4.A.1 Forest land remaining forest land – CH ₄ and N ₂ O	The ERT recommends that Switzerland explain the reallocation of CH ₄ and N ₂ O emissions from open burning of residues from forests from category 5.C.2 (open burning of waste) in the waste sector to the category 4(V).A.1 controlled burning in forest land remaining forest land (CRF table 4(V) biomass burning under the LULUCF sector) in its NIR	L.13 was addressed in chp. 6.4.2.13.

Table 10-1 continued.

ID	classification	recommendation	Detail = Answer including ref to chapter in NIR
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
Waste			
W.7	5.A Solid waste disposal on land – CH4	The ERT recommends that Switzerland report the correct tier for the methodology used to estimate CH4 emissions from solid waste disposal on land in CRF table summary 3s2 and in its NIR	Chapter 7.2.2: Methodical issues corrected to Tier 2.
W.8	5.A Solid waste disposal on land – CH4	The ERT recommends that Switzerland explain in more detail the assumptions, AD and methodologies used to estimate landfill with gas recovery both for electricity production and other purposes. <<This issue is not applicable to ARR 2015 as the study did not yet report.>>	Detailed explanations are given in chp. 7.2.2
W.9	5.B.1 Composting – CH4	Since emissions from the biological treatment of solid waste (5.B) is a key category, the ERT recommends that Switzerland review and, if necessary, revise, the AD for composting and demonstrate that they are accurate by providing supporting documentation in the NIR	W.9 is listed under planned improvements in chp. 7.3.6.
W.10	5.B.2 Anaerobic digestion at biogas facilities – CH4	However, to increase transparency in the NIR, the ERT recommends that Switzerland explain in more detail how it obtained the country-specific EF for CH4 losses from biogas facilities in its NIR	A detailed explanation is given in chp. 7.3.2.2
W.11	5.C Incineration and open burning of waste – CH4 and N2O	In order to improve transparency of the AD, the ERT recommends that Switzerland provide a more detailed explanation of the source, data acquisition and references of the AD, by type of waste, used to estimate CH4 and N2O emissions from incineration and open burning of waste	Detailed explanations are given in chp 7.4.2
W.12	5.C.2 Open burning of waste – CH4 and N2O	The ERT recommends that Switzerland identify in its NIR the definition of natural agricultural residue waste as a country-specific type of waste in Switzerland or national waste, as allowed by the definition of MSW in the 2006 IPCC Guidelines	In chapter 7.4.1 (additional text): Natural agricultural and gardening residues consist of fallen fruit trees, part of diseased residue which are cut up, collected and burned off-site.
W.13	5.C.2 Open burning of waste – CH4 and N2O	The ERT recommends that Switzerland explain the reallocation of CH4 and N2O emissions from open burning of residues from forests in the NIR	W.13 was addressed in Switzerland's Saturday paper 2016 (see Annex 7).
W.14	5.C.2 Open burning of waste (biogenic)	To increase consistency between the NIR and the CRF tables, the ERT recommends that Switzerland correct the AD reported in table 5.C for open burning of waste for natural residues and ensure consistency between the NIR and the CRF tables on these AD	Activity data in the CRF tables has been corrected.
KP-LULUCF			
KL.3	General (KP-LULUCF)	The ERT recommends that Switzerland address the transparency issues ID#L.6, ID#L.7, ID#L.9, ID#L.10 and ID#L.12 and provide the necessary information in relation to KP-LULUCF activities	For L.6, L.7, L.9, L.10 and L.12 see the respective issues above. When these issues are resolved in the LULUCF sector they are inherently addressed and resolved in the KP-LULUCF sector.
KL.4	Afforestation and reforestation, deforestation – CO2	The ERT recommends that the Party review the assumption that only 50 per cent of the difference between the carbon stocks before and after the change is reported as a source or sink, respectively, for afforestation (from settlements to forest land) and deforestation (from forest land to settlements) and, if necessary, revise its estimates for these KP-LULUCF activities	KL.4 was listed under planned improvements in chp. 6.8.6. and chp. 11.5.2.3.
KL.5	Deforestation	The ERT recommends that Switzerland address issue ID#L.11 and, if necessary, revise its estimates for deforestation	KL.5 (L.11) was listed under planned improvements in chp. 6.4.6.
KL.6	Forest management – CH4 and N2O	The ERT recommends that Switzerland explain in its NIR the estimation of CH4 and N2O emissions from open burning of residues from forests and its allocation to the category controlled burning in CRF table 4(KP-II)4 (GHG emissions from biomass burning for forest management) The ERT also recommends that Switzerland include the reallocated values in its FMRL, applying a technical correction if necessary	L.13 was addressed in chp. 6.4.2.13. KL.6 was listed under planned technical corrections for the FMRL in chp. 11.5.2.3. A technical correction of the FMRL is planned for the submission in 2019.
KL.7	Forest management	The ERT recommends that the Party report the correct values for both FMRL and the technical corrections in CRF table accounting	KL.7 was addressed by reporting the correct values for both FMRL and the technical correction in CRF table 4(KP-I)B.1.1.

Table 10-1 continued.

ID	classification	recommendation	Detail = Answer including ref to chapter in NIR
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
KP-LULUCF			
KL.8	Harvested wood products – CO2	The ERT recommends that Switzerland estimate and report carbon stock changes for the product category paper based on either the national or the internationally available data, or provide transparent justification in the NIR as to why the available information on AD for paper is not transparent and verifiable	ID#KP.L8 was listed under planned improvements in chp. 6.11.6.
KL.9	Harvested wood products – CO2	The ERT recommends that Switzerland increase the transparency of its reporting by correctly reporting the amount of exported HWP in CRF table 4(KP-I)C instead of using "NA" or by entering the notation key "IE" if exported HWP are included in the total HWP production	KL.9 was addressed by inserting "IE" in the CRF table 4(KP-I)C and was documented in chp. 6.11.2.
KL.10	Harvested wood products – CO3	The ERT recommends that the Party increase the transparency of the explanation in the NIR to clarify that exports of round wood are excluded from the calculations following equation 2.8.1	KL.10 was adressed in chp. 6.11.2.

Table 10-2 Encouragements from the ERT based on the Draft Review Report (UNFCCC 2017) and explanations for implementation 2016/2017.

ID	classification	encouragement	Details
Unresolved issues according to Table 5: Additional findings made during the 2016 technical review of the annual submission of Switzerland			
Energy			
E.6	Fuel combustion – reference approach – comparison with international data – other fossil fuels – CO ₂	The ERT encourages Switzerland to document in the NIR explanations to justify the deviations between the CRF tables and IEA energy consumption data of other fossil fuels	The allocation of liquid fossil, waste (non-biomass fraction) and biomass total in the reference approach has been reassessed. The reasons for the discrepancy between IEA and the CRF tables are provided in Annex 4 of the NIR.
E.7	Fuel combustion – reference approach – liquid fuels – CO ₂	The ERT encourages Switzerland to report apparent consumption of other kerosene and refinery feedstock separately in CRF table 1.A(b) or change the reported notation key for other kerosene and refinery feedstock in CRF table 1.A(b) from “NO” to “IE”	Notation key has been set to “IE” for “Import” of both fuels in CRF table 1.A(b) (incl. Information in documentation boxes)
IPPU			
not mentioned in ARR (only in preliminary findings)	IPPU 2A3	Improve description of table 4-8 to ensure that emission factors are presented in transparent manner.	The comparison between the Tier 3 and Tier 2 approach is described in the confidential NIR in chp. 4.2.2.3.
Agriculture			
A.4	3.A.4 Other livestock – CH ₄	The ERT encourages Switzerland to provide short relevant explanations of the methods used to estimate dry matter intake and gross energy for buffalo, camels and deer	An explanation and a table (Table 5-7) is provided in NIR chp. 5.2.2.2.1.
A.5	3.B Manure management – N ₂ O	The ERT encourages Switzerland to provide a short comparison of country-specific Nex rates for all animal categories to default values estimated in accordance with tier 2 of the 2006 IPCC Guidelines in the QA/QC section of the NIR with relevant explanations of deviations	The comparison was conducted and is documented in the QA/QC document. However, the QA/QC document is not yet updated completely due to the lack of some further analysis in other source categories.
LULUCF			
L.12	4.A Forest land – CO ₂	The ERT encourages the Party to continue its efforts to validate (and if possible, better adapt) the parameter sets applied, and to include this information in the NIR to increase the transparency of the description of the above-mentioned model. The ERT notes that if the new version of the model produces more accurate results than the previous version and the Party decides to use it in its estimates, the new version should be applied consistently over time	L.12 was listed under planned improvements in chp. 6.4.6.
Waste			
W.6	5. General (waste)	The ERT encourages Switzerland to estimate and report long-term storage of C in waste disposal sites, annual change in total long-term C storage and annual change in total long-term C storage in HWP waste in CRF table 5 If Switzerland continues to report long-term storage of C in waste disposal sites, annual change in total long-term C storage and annual change in total long-term C storage in HWP waste as “NE” in CRF table 5, the ERT encourages the Party to provide appropriate information in the documentation box of CRF table 5 and in its NIR	W.6 was listed under planned improvements in chp. 7.2.6.

10.1.2 Recalculations and improvements reported in Submission 2017

10.1.2.1 General recalculations and improvements

10.1.2.1.1 Feedstocks and non-energy use of fuels

- The net consumption of non-energy use of fuels reported in Swiss overall energy statistics includes sulphur produced by the refineries as well. This amount of sulphur is now subtracted resulting in lower fuel quantities for non-energy use of other oil for the entire time series reported in CRF table 1.A(d). See also chp. 3.2.4.9.

10.1.2.1.2 Category-specific recalculations for 1A

- Reference Approach: Consumption of so-called other non-fossil fuels (biogenic waste) was missing so far. It is now also included in the Reference Approach in CRF table 1.A(b).
- 1A: The CO₂ emission factor for natural gas is linearly interpolated between 1995–2000. (Before it was kept constant between 1995–1998).
- 1A: The CO₂ emission factor for natural gas has changed. There was a minor error in the calculations for the years as follows:
2009: from 56'400g/GJ to 56'500 g/GJ,
2010: from 56'400g/GJ to 56'500 g/GJ,
2014: from 56'700g/GJ to 56'500 g/GJ
- 1A: Small recalculations due to rounding of activity data 2013, 2014 in the Swiss overall energy statistics (SFOE 2016) concerning bituminous coal and natural gas.
- 1A: Non-energy use of fuels (NEU): The net consumption of non-energy use of fuels reported in Swiss overall energy statistics includes sulphur produced by the refineries as well. This amount of sulphur is now subtracted resulting in lower fuel quantities for NEU of other oil for the entire time series reported in CRF table 1.A(d).

10.1.2.2 Energy

Recalculations for sector 1 Energy are expressed quantitatively for changes of 1 kt CO₂ eq or larger, whereas smaller changes are described qualitatively. The description of recalculations in the following chapters follows the structure in chp. 3, i.e. all stationary sources are described first (chp. 10.1.2.2.1– 10.1.2.2.3), followed by mobile sources (chp. 10.1.2.2.4– 10.1.2.2.6).

10.1.2.2.1 Category-specific recalculations for 1A1 (stationary)

- 1A1a: Recalculation concerning boilers using natural gas due to recalculation in losses of natural gas in distribution network (2013, 2014).
- 1A1a: The time series value of waste generation rate for 2013 has slightly changed due to the correction of an error in waste statistics.
- 1A1a: The use of carbonates for sulphur oxide removal in 1A1a Municipal solid waste incineration plants has been moved to 2A4d Other process uses of carbonates (1990–

2006). In 1990, this recalculation leads to a decrease in emissions in category 1A1a by 0.3 kt CO₂ eq and in 2014 the emissions decrease by 1.4 kt CO₂ eq (and a corresponding increase in 2A4d).

- 1A1a: Recalculation in residual fuel oil boilers due to mistake in calculations in the energy model for the entire time series (1990–2014). This recalculation reduces emissions in 2014 by about 2 kt CO₂ eq and in 1990 by about 1 kt CO₂ eq.
- 1A1b: In last submission the emission factor of CO₂ of refinery gas and residual fuel oil used in refineries was linear interpolated from 1990–2005. It was changed to be constant from 1990–2004. This leads to a recalculation between 1991 and 2004.
- 1A1b: There was a transcription error in the calculation of the CO₂ emission factor of refinery gas in 2014. The correction of the error results in an increase of the emissions increased in 2014 by about 11 kt CO₂ eq.

10.1.2.2.2 Category-specific recalculations for 1A2 (stationary)

Recalculations at the aggregation level of 1A2 amount to -8 kt CO₂ eq for natural gas and about -1 kt CO₂ eq for petroleum coke in 2014. Reallocation of consumption of natural gas and gas oil at the level of the source categories 1A2a-1A2g viii causes differences, which however do not have any influence on total emissions reported in 1A2.

- 1A2: Recalculations due to new available statistics (SFOE2016d) (2012, 2013), causing a reallocation of energy to the different categories 1A2a-1A2g. Total energy consumption in 1A2 is not affected by this recalculation.
- 1A2: Amount of used gas oil in households, industry and commercial sector in Liechtenstein has been redistributed. Therefore the amount of used gas oil in boilers in households, industry and the commercial sector for Switzerland has changed too (1990–2014).
- 1A2d: The use of limestone for sulphur oxide removal in 1A2d Cellulose production has been moved to 2A4d Other process uses of carbonates (1990–2008). This recalculation leads to a decrease in emissions by 4 kt CO₂ eq in 1990 in 1A2d (and a corresponding increase in 2A4d).
- 1A2d: The so far missing emissions of CH₄ and N₂O from biogenic waste used in 1A2d Cellulose production are now reported for 1990–2008 based on default emission factors from the 2006 IPCC Guidelines for sulphite lyes. This recalculation leads to an increase by about 1 kt CO₂ eq in 1990.
- 1A2f: Activity data for 2014 of other bituminous coal and petroleum coke in cement production has changed due to the correction of an assignment error between the two fuels. This results in a recalculation of about +18 kt CO₂ eq for bituminous coal and -18 kt CO₂ eq for petroleum coke.
- 1A2f: AD (gas oil, natural gas) of 1A2f Production of mixed goods have been revised for 2014 based on corrected data from industry association.
- 1A2f: The emission factors of NO_x, NMVOC and CO of 1A2f Production of mixed goods have been revised from 1991 onwards based on air pollution control measurements (2001–2015).
- 1A2f: Revised interpolated CO emission factors of 1A2f Rockwool production in 2014 due to new plant-specific data for 2015.

- 1A2f: The emission factors of NMVOC and SO₂ as well as CO from 1A2f Glass production (speciality tableware) have been revised from 1991 and 1996, respectively, onwards.
- 1A2f: Fuel mix and consumption of 1A2f Lime production have been revised for 1990–1994 and 1990–2008 based on industry data and current net calorific values of gas oil, respectively, resulting in minor changes of AD of gas oil and residual fuel oil for 1990–2008 and 1994, respectively. In addition, it was found that not petroleum coke was used up to 1994 but other bituminous coal. All these changes induce recalculations in AD (gas oil, residual fuel oil, petroleum coke and other bituminous coal) of 1A2g viii Industrial boilers in the respective years. In 1990, this recalculation leads to a shift in emissions from bituminous coal from 1A2g viii to 1A2f (23 kt CO₂ eq).
- 1A2f: The conversion factor used for calculation of NMVOC emissions from total carbon based on air pollution control measurements has been revised resulting in adjusted NMVOC emission factors of 1A2f Brick and tile production and 1A2f Fine ceramics production for the entire time series
- 1A2g viii: Recalculations in AD (1990–2008: gas oil, residual fuel oil, 1990–1994: petroleum coke and other bituminous coal) of 1A2g viii Industrial boilers due to revised fuel mix and consumption in 1A2f Lime production. In 1990, this recalculation leads to a shift in emissions from bituminous coal from 1A2g viii to 1A2f (23 kt CO₂ eq) and a reduction of emissions from petroleum coke by 21 kt CO₂ eq and gas oil by 1 kt CO₂ eq.

10.1.2.2.3 Category-specific recalculations for 1A4 (stationary)

- 1A4: Recalculations of the CO₂ emission factor of natural gas (see chp. 3.2.4.9) result in a change in the emission level of -13 kt CO₂ eq in 2014 and 2 kt CO₂ eq in 1990.
- 1A4ci: The missing emission factors in last year's submission of NO_x, CO, NMVOC and SO₂ for 1A4ci Plants for renewable waste from wood products in 2014 are now included in the inventory.
- 1A4ci: The emission factors of NMVOC as well as NO_x and CO from 1A4ci Drying of grass have been revised from 1990 and 1991, respectively, onwards based on air pollution control measurements (2005–2015).
- 1A4ai, 1A4bi and 1A4ci Wood combustion AD of automatic boilers and stoves have been revised for 1990–2014 and 2011–2014, respectively due to minor recalculations in Swiss wood energy statistics (SFOE 2016b).

10.1.2.2.4 Category-specific recalculations for 1A3 (mobile)

- 1A3a: As recommended by the ERT and in line with the IPCC reporting GLs there are no more CH₄ emissions for cruise activities. This implies higher NMVOC emissions for cruise activities. All years 1990–2014 have been recalculated.
- 1A3b: The CO₂ emission factor for biogas used in road transportation was out-dated and has now been adapted 1990–2014 to the one of natural gas.
- 1A3b/1A3d: Small recalculation due to a change in the NCV of diesel used in international navigation to equalise with other diesel processes. Therefore small changes occurred 1990–2014 in fuel tourism and statistical difference for diesel which is integrated in 1A3biii.
- 1A3bi cars and 1A3bii LDT: N₂O cold start excess emissions for PC and LDV by means of emission factors of the Copert model have been added for 1990, 2013 and 2014. For

the years 1991–2012 the emissions were linearly interpolated between 1990 and 2013. Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

10.1.2.2.5 Category-specific recalculations for 1A4 aii/bii/cii mobile

- No category-specific recalculations were carried out.

10.1.2.2.6 Category-specific recalculations for 1A5b (mobile)

- No category-specific recalculations were carried out.

10.1.2.2.7 Category-specific recalculations for 1B2a

- 1B2a: Small recalculations in activity data due to change of units (1990–2014).
- 1B2a: SO₂ emissions from Claus units were previously reported in source category 2H3 and are now reported in 1B2a iv.

10.1.2.2.8 Category-specific recalculations for 1B2b

- 1B2b ii: Recalculation in the Swiss overall energy statistics (SFOE 2016) concerning production of natural gas between 1985 and 1994.
- 1B2b ii: Reassessment of the emission factors for source category 1B2b ii.
- 1B2b iv and 1B2b v: Recalculations were carried out due to the update of the employed calculation tool with minor corrections of individual natural gas loss rates, minor corrections of individual data regarding the Swiss gas network as well as minor changes in the polynomial interpolations for years with no sufficient data of the gas network available (EMIS 2017/1B2b Diffuse Emissionen Erdgas). This recalculation leads to an increase of 15 kt CO₂ eq in 1990.

10.1.2.2.9 Category-specific recalculations for 1B2c

- 1B2c: CH₄ emission factor of flaring activity in refineries changed due to less rounding (2005–2014).
- 1B2c: Emissions of N₂O, CO₂ and CH₄ from flaring in gas production were missing before. Now emission factors from the 2006 IPCC Guidelines are used (1990–1994).

10.1.2.3 Industrial processes and other product use

Recalculations for sector 2 Industrial processes and other product use are expressed quantitatively for changes amounting to 0.3 kt CO₂ eq or larger, whereas smaller changes are described qualitatively.

10.1.2.3.1 Category-specific recalculations for 2A

- 2A1: The EF for CO₂ geog. has been changed to a constant value of 536.718 kg/t clinker for the whole time period which is higher than the EF used before. It now corresponds to the calculation in the Swiss emissions trading system. This results in an emission increase of 860 kt over the entire period 1990–2015. In 2014 the increase amounts to 21.6 kt CO₂ eq and in 1990 the increase amounts to 56 kt CO₂ eq.
- 2A3: AD of 2A3 Container glass have been revised for 2003–2006 based on monitoring reports of the Swiss emissions trading scheme.
- 2A4d: The use of carbonates for sulphur oxide removal in municipal solid waste incineration plants has been moved from 1A1a to 2A4d Other process uses of carbonates (entire time series). In 1990, this recalculation leads to an increase in emissions by 0.3 kt CO₂ eq and in 2014 the emissions increase by 1.4 kt CO₂ eq.
- 2A4d: The use of limestone for sulphur oxide removal in cellulose production has been moved from 1A2d to 2A4d Other process uses of carbonates (1990–2008). In 1990, this recalculation leads to an increase in emissions by 4 kt CO₂ eq.
- 2A4d: The emissions of geogenic CO₂ from limestone use in cupola furnaces of iron foundries are newly reported in the inventory for the entire time series. This leads to an increase by 0.4 kt CO₂ eq in 2014.
- 2A4d: Negligible rounding changes in AD of 2A4d Other use of carbonates for 1990–2013.
- 2A4d: AD of 2A4d Carbonate use in municipal solid waste incineration has been adjusted for 2014 yielding revised AD for 2014 of 2A4d Other use of carbonates as well.

10.1.2.3.2 Category-specific recalculations for 2B

- 2B10: Minor changes in CO₂ emissions from 2B10 Limestone pit due to rounding differences of AD for 1990–1998.
- 2B10: The extrapolated values of the SO₂ emission factor of 2B10a Sulphuric acid production have been revised (1990–2008).

10.1.2.3.3 Category-specific recalculations for 2C

- 2C1: The conversion factor used for calculation of NMVOC emissions from total carbon based on air pollution control measurements has been revised resulting in an adjusted NMVOC emission factor of 2C1 Secondary steel production, electric arc furnace from 1995 onwards.
- 2C7c: The NMVOC emission factor of 2C7c Battery recycling is newly reported in the inventory based on air pollution control measurements (2003 and 2012).
- 2C7c: The emission factors of CO₂, NO_x, CO and SO₂ from 2C7c Battery recycling have been revised from 2002 onwards based on air pollution control measurements (2003 and 2012).

- No recalculations reported in the submission 2017 for F-gases in source category 2C.

10.1.2.3.4 Category-specific recalculations for 2D

- 2D3b: The AD of 2D3b Road paving has been revised for 2014 based on corrected data from industry association.
- 2D3a: The survey on post-combustion of NMVOC emissions from coating applications, degreasing, dry-cleaning and manufacture and processing of chemical products has been revised resulting in changes of CO₂ emissions of 2D3a Post-combustion of NMVOC from solvent use for the entire time series. In 1990, this recalculation leads to a decrease in emissions by 1 kt CO₂ eq and in 2014, this recalculation leads to an increase by 1 kt CO₂ eq.

10.1.2.3.5 Category-specific recalculations for 2E

- No category-specific recalculations were carried out, but all data on the recovery were replaced by the notation key "NA". The definition of recovery in the CRF tables and the use of recovery in the calculation of the implied emission factor are not coherent in the submission 2016 and led to false results for the implied emission factor.

10.1.2.3.6 Category-specific recalculations for 2F

- 2F1: Product life emission factor of commercial and industrial refrigeration is now assumed to decrease slower, i.e. linearly from 12% in 1995 to 7% in 2015.
- 2F1 The correction factor for refrigerant import to avoid double countings with Liechtenstein has been applied for the whole time period (so far only considered for the time period 2007 to 2014).
- 2F5 .The correction factor for solvents import to avoid double countings with Liechtenstein has been applied for the whole time period (so far only considered for the time period 2007 to 2014).

10.1.2.3.7 Category-specific recalculations for 2G

- 2G: Unfortunately there is a double-counting in the NMVOC emissions from 2G De-icing of airplanes for the years 1990–2006. This mistake will be corrected in submission 2018.
- 2G3a: The N₂O emission factor for 2020 of 2G3a Use of N₂O in hospitals has been adapted to the current scenario of Swiss population development yielding revised interpolated values for 2012–2014.
- 2G4: The survey on post-combustion of NMVOC emissions from printing and other solvent and product use has been revised resulting in changes of CO₂ emissions of 2G4 Post-combustion of NMVOC from other solvent and product use for the entire time series. In 2014, this recalculation leads to an increase in emissions by 1 kt CO₂ eq.

10.1.2.3.8 Category-specific recalculations for 2H

- 2H1: AD of 2H1 Chipboard production have been revised from 2005 onwards based on a changed density value of the chipboard produced.
- 2H2: Activity data of wine production has slightly decreased in the year 2011 due to the correction in the underlying statistics from the Swiss Alcohol Administration.
- 2H2: Activity data of sugar production has slightly changed for the years 2011, 2012 and 2014 due to corrections in the statistics provided by the Swiss Sugar Association.
- 2H2: Activity data of meat smokehouses has slightly changed for the years 2007, 2013 and 2014 due to corrections in the statistics provided by the Swiss Federal Office for Statistics.
- 2H3: SO₂ emissions from Claus units were previously reported in source category 2H3 and are now reported in 1B2a iv.

10.1.2.4 Agriculture

10.1.2.4.1 Category-specific recalculations for 3A

- CH₄ emission factors for the year 2012 and 2014 of sheep, goats, swine and poultry and "other" (livestock NCAC) were revised due to updates of provisional net energy intake data by the Swiss Farmers Union (Giuliani 2016). The effect of the recalculation on overall greenhouse gas emissions is negligible (<1 kt CO₂ equivalents).

10.1.2.4.2 Category-specific recalculations for 3B

- The excretion of VS of horses was revised due to a respective recommendation during the annual review in 2016 (UNFCCC 2017; A.6). New values are based on equation 10.24 in the 2006 IPCC Guidelines. New emission estimates are slightly higher (+ 0.44 kt CO₂ eq. on average, 1990: +0.36 kt CO₂ eq., 2014: +0.48 kt CO₂ eq.)
- The amount of VS in the manure management system anaerobic digesters was revised for the year 2014 due to updated estimates of the respective AD in the waste sector. Overall CH₄ and N₂O emissions increased by 0.7 kt CO₂ equivalents.
- All estimates based on AGRAMMON data were recalculated for the years 2011–2014 due to a new interpolation after new survey results for 2015 became available. The impact on overall emissions in kt CO₂ equivalents is: 2011: 0.4; 2012: 0.9; 2013: 1.4; 2014: 1.9.
- For the years 2011–2014, the MCF for liquid/slurry systems was slightly revised due to new model runs based on the new AGRAMMON data. Effects on overall CH₄ emissions in kt CO₂ equivalents are: 2011: 0.5; 2012: 0.9; 2013: 1.3; 2014: 1.7.

10.1.2.4.3 Category-specific recalculations for 3D

- The AD for direct N₂O emissions from N in crop residues returned to soils was revised for the year 2014 due to data updates by the Swiss Farmers Union. Overall emissions increased by 0.02 kt CO₂ equivalents.

- The alpine area used to estimate NH₃ volatilisation from the vegetation cover and N₂O emissions from crop residues on pasture and meadows was slightly revised with negligible effects on overall emissions.
- All estimates based on AGRAMMON data were recalculated for the years 2011–2014 due to a new interpolation after new survey results for 2015 became available. The impact on overall emissions in kt CO₂ equivalents is: 2011: 0.4; 2012: 0.9; 2013: 1.4; 2014: 1.9.
- N-input from co-substrates in agricultural biogas plants for 2014 was revised due to an error correction in the background file of the waste sector. The impact on overall emissions is negligible.
- AD for N₂O emissions from the cultivation of organic soils was revised due to new AREA-projections in the LULUCF sector for all inventory years. Overall emissions decreased by 0.03 kt CO₂ equivalent on average (1990: no change; 2014: -0.08 kt CO₂ equivalents).
- The AD for direct N₂O emissions from N in mineral soils that is mineralized/immobilized in association with loss of soil C was revised due to new projections in the LULUCF sector for all inventory years. Overall emissions increased by 0.13 kt CO₂ equivalents on average (1990: +0.00 kt CO₂ equivalents; 2014: +0.20 kt CO₂ equivalents).
- Fra_{CLEACH} was recalculated due to new model estimates of NO₃ losses on agricultural soils from Prasuhn (2016). Overall emissions decreased by 18.67 kt CO₂ equivalents on average (1990–2014).

10.1.2.4.4 Category-specific recalculations for 3G

- All lime applied in agriculture was previously reported as Ca(CO₃). During the 2017 submission the total amount of lime was split into calcium carbonate and dolomite. The new specific EF for dolomite (i.e. 0.13) is slightly higher than the old EF that was used previously for both limestone and dolomite (i.e. 0.12). Accordingly, overall emissions increased by 0.15 kt CO₂ equivalents on average (1990: +0.12 kt CO₂ equivalents; 2014: +0.23 kt CO₂ equivalents).

10.1.2.4.5 Category-specific recalculations for 3H

- No category-specific recalculations were carried out.

10.1.2.5 Land Use, Land-Use change and Forestry

10.1.2.5.1 Category-specific recalculations for 4A

- Activity data 1990–2014 were updated (see chp. 6.3.5).
- The most recent NFI data (NFI4 2011–2015) were applied for productive forests (CC12; see Thürig et al. 2017; see chp. 6.4.2.1).
- Modelling carbon stock changes in dead wood, litter and mineral soil with Yasso07 for productive forests: the methodological change in the National Forest Inventory (NFI) from a periodic (i.e. NFI1, NFI2, and NFI3) to a continuous sampling starting with the NFI4 required modifications to the approach for estimating dead wood and litter inputs. To derive estimates of carbon inputs between NFI3 and NFI4, plots which were visited within the latest 5 years of the NFI4 were considered, i.e. currently 5-year tranche NFI4 2011–2015. Thus, the sites used in the Yasso07 simulations (cf. chp. 2.4.1 in Didion and Thürig

2016) changed in comparison with the GHGI inventory submission 2016 (FOEN 2016); N is now 2616 compared to 2607 in FOEN (2016) of which 1591 are common sites. Further, the accuracy of the estimates of dead wood and litter production was improved. Besides minor corrections, an inconsistency was removed in the NFI1 regarding the classification of laying trees as alive or dead. This resulted in more accurate estimates of the dead wood and litter production in the period NFI1–2.

- In response to issue ID#W.13 in UNFCCC (2017) (see also UNFCCC 2017/ID#L.13 and UNFCCC 2017a/ID#1) CH₄ and N₂O emissions from open burning of residues from forests were reallocated from category 5C2 (open burning of waste) in the waste sector to category 4(V)A1 (controlled burning in forest land remaining forest land) (see chp. 6.4.2.13). Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

10.1.2.5.2 Category-specific recalculations for 4B

- Activity data 1991–2014 were updated (see chp. 6.3.5).
- Provisional carbon stocks and carbon stock changes in living biomass for the years 2013 and 2014 were replaced by definitive values from SBV (2016).
- The provisional carbon stock change in living biomass for 1990 was replaced by the definitive value calculated from the difference 1989–1990 (cf. Table 6-24).

10.1.2.5.3 Category-specific recalculations for 4C

- Activity data 1991–2014 were updated (see chp. 6.3.5).
- The carbon stock in living biomass for orchards (CC35) and unproductive grassland (CC37) were recalculated for the time series 1990–2014 using area-weighted means instead of arithmetic means.
- The amount of “fuel” for wildfires was recalculated for the time series 1990–2014 with the updated biomass carbon stocks.

10.1.2.5.4 Category-specific recalculations for 4D

- Activity data 1991–2014 were updated (see chp. 6.3.5).

10.1.2.5.5 Category-specific recalculations for 4E

- Activity data 1991–2014 were updated (see chp. 6.3.5)

10.1.2.5.6 Category-specific recalculations for 4F

- Activity data 1991–2014 were updated (see chp. 6.3.5)

10.1.2.5.7 Category-specific recalculations for 4(III), 4(IV)

- Activity data 1991–2014 were updated (see chp. 6.3.5).

10.1.2.5.8 Category-specific recalculations for 4G

- The activity data were updated for the whole time-series. More specifically, the share of domestic to total HWP was revised.
- Net CO₂ emissions and removals from Harvested wood products (HWP) Deforestations (KP) were accounted for on the basis of instantaneous oxidation.

10.1.2.6 Waste

- Recalculations for sector 5 Waste are expressed quantitatively for changes amounting to 0.2 kt CO₂ eq or larger, whereas smaller changes are described qualitatively.

10.1.2.6.1 Category-specific recalculations for 5A

- No category-specific recalculations were carried out.

10.1.2.6.2 Category-specific recalculations for 5B

- 5B2 The CH₄ emission factor for biogas upgrading has been recalculated for the whole time series from 1990–2014. The net calorific value of methane has been adjusted to 50 GJ/t (from 46.5) to be in accordance with the fact sheet with CO₂-emission factors (FOEN 2015d) and the rest of the inventory. This recalculation leads to a decrease by 0.3 kt CO₂ eq in 2014.

10.1.2.6.3 Category-specific recalculations for 5C

- 5C1 Activity data for illegal waste incineration for the year 2013 has slightly increased due to the correction of an error in waste statistics.
- 5C1 Activity data for sewage sludge incineration for the year 2013 has slightly increased due to the correction of an error in waste statistics.
- 5C2 As a consequence of the greenhouse gas inventory UNFCCC in-country review 2016 greenhouse gas emissions from open burning of natural residues in forestry (5 C 2 ii) were moved to LULUCF sector 4 V (new in chapter 6.4.2.13). Note, this improvement is not a recalculation in the strict sense because it was addressed in Switzerland's answers in the Saturday paper emerging from the in-country review process in 2016. The issue was considered to be resolved by the ERT and was implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

10.1.2.6.4 Category-specific recalculations for 5D

- 5D1 Activity data of energy produced in combined heat and power generation engines (CHP) has been recalculated for 2011–2014 in the Swiss Statistics for Renewable

Energies by SFOE. This leads to changed CH₄ and CO₂ biog. emission factors (emissions per capita).

- 5D The N₂O emission factor for wastewater handling for the year 2014 has slightly decreased because it was updated according to the last published value in the Agristat 2015 statistics published by the Swiss Farmers Association. This recalculation leads to a decrease in emissions in 2014 by 2 kt CO₂ eq.

10.1.2.6.5 Category-specific recalculations for 5E

- No category-specific recalculations were carried out.

10.1.2.7 Other

- 6Ad Activity data for fire damages motor vehicles has changed for the years 2003–2014 because the model has been updated with vehicle data provided by the SFOS.
- 6Ad Activity data for fire damages estates has changed for the years 1996–2014 because the model has been updated with damage sums provided by the Swiss Building Insurance Association.

10.1.2.8 Indirect CO₂ and N₂O Emissions

- See NMVOC related recalculations reported in chp. 3 Energy, 4 Industrial processes and product use, 7 Waste and 8 Other.
- See CO related recalculations reported in chp. 4 Industrial processes and product use, 7 Waste and 8 Other.
- See NO_x and NH₃ related recalculations reported in chp. 3 Energy, 4 Industrial processes and product use, 7 Waste and 8 Other.
- The CO₂ emissions from 2C1 Secondary steel production, electric arc furnace are based on a carbon mass balance considering all carbon sources and sinks of the process. Therefore, the emissions of CO and NMVOC are no longer included in the calculation of the indirect CO₂ emissions from sector 2 IPPU in order to avoid a double counting.

Recalculations implemented in Switzerland's Saturday paper 2016:

- 2C3: The ERT found that the CO₂ emissions from aluminium production (direct and indirect) are potentially overestimated for the years 1990-2006. Switzerland eliminated any double-counting of CO₂ by removing the indirect CO₂ emissions from oxidation of CO from 2C3 Primary aluminium production. Note, this improvement is not a recalculation in the strict sense because it has been addressed in Switzerland's answers in the Saturday paper emerging from the in house review process in 2016. The issue is considered to be resolved by the ERT and has been implemented in the reporting tables submitted on 7 November 2016. For completeness the Saturday paper (Attachment C) including comprehensive answers to the ERT is presented in Annex 7.

10.1.2.9 KP- LULUCF Inventory

- Net CO₂ emissions and removals from Harvested wood products (HWP) from deforestations are accounted for on the basis of instantaneous oxidation.
- Emissions from controlled burning are for the first time reported under Forest management. In previous submissions those emissions were reported under 5C2 “incineration and open burning of waste – natural residues”.

10.2 Implications for emission levels

Table 10-3 shows the effect of the recalculations on the results for the base year 1990. The recalculations resulted in an increase of the total emissions in CO₂ equivalents (without emissions/removals from CO₂ from LULUCF) of 49 kt CO₂ eq (0.09%) in submission 2017 as compared to submission 2016 of November 7. If the LULUCF sector is included, there is an increase of an increase of 648 kt CO₂ eq (1.24%) for submission 2017 as compared to submission 2016.

Table 10-3 Overview of implications of recalculations on 1990 data. Emissions are shown before the recalculation according to the previous submission of 7 November 2016 “Prev.” (FOEN 2016) and after the recalculation according to the present submission 2017 “Latest”. The difference refers to the absolute values (Latest - Previous).

Recalculation	CO ₂			CH ₄			N ₂ O			Sum (CO ₂ , CH ₄ and N ₂ O)		
	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.
Emissions for 1990												
Source and Sink Categories	CO ₂ equivalent (kt)									CO ₂ equivalent (kt)		
1. Energy	40'907	40'899	-7.49	636.1	651.9	15.80	294.2	294.8	0.63	41'837	41'846	8.94
2. IPPU	3'095	3'158	63.55	1.8	1.8	0.00	171.4	171.4	0.00	3'268	3'331	63.55
3. Agriculture	49	49	-0.32	4'508.8	4'509.2	0.36	2'245.8	2'222.2	-23.53	6'804	6'780	-23.49
4 LULUCF excl. HWP	-994	-395	599.0	29.4	29.4	0.00	86.9	86.9	0.00	-878	-279	598.97
5. Waste	54	54	0.00	938.8	938.7	-0.12	140.5	140.5	0.00	1'133	1'133	-0.12
6 Other	11	11	0.00	0.7	0.7	0.00	0.6	0.6	0.00	12	12	0.00
Sum (without F-gases)	43'122	43'776	654.7	6'116	6'132	16.04	2'939	2'916	-22.90	52'177	52'824	647.85

Recalculation	HFC			PFC			SF ₆			Sum (synthetic gases)		
	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.
Emissions for 1990												
Source and Sink Categories	CO ₂ equivalent (kt)									CO ₂ equivalent (kt)		
2 Ind. Processes (only syn. gases)	0.02	0.02	0.00	116.5	116.5	0.00	137.0	137.0	0.00	253.55	253.55	0.00

Recalculation	Sum (all gases)		
	Prev.	Latest	Differ.
Emissions for 1990			
Source and Sink Categories	CO ₂ equivalent (kt)		
Total CO₂ eq Em. with LULUCF	52'430	53'078	647.85
	100%	101.24%	1.24%
Total CO₂ eq Em. without LULUCF	53'308	53'357	48.88
	100%	100.09%	0.09%

Table 10-4 shows the effect of the recalculations on the results for the year 2014. The recalculations resulted in an increase of the total emissions in CO₂ equivalents (without emissions/removals from CO₂ from LULUCF) of 3.1 kt CO₂ eq (0.01%) submission 2017 compared to submission 2016. If the LULUCF sector is included, there is an increase of 114 kt CO₂ eq (0.24%) submission 2017.

Table 10-4 Overview of implications of recalculations on 2014 data. Emissions are shown before the recalculation according to the previous submission of 7 November 2016 "Prev." (FOEN 2016) and after the recalculation according to the present submission 2017 "Latest". The difference refers to the absolute values (Latest - Previous).

Recalculation	CO ₂			CH ₄			N ₂ O			Sum (CO ₂ , CH ₄ and N ₂ O)					
	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.			
Emissions for 2014															
Source and Sink Categories	CO ₂ equivalent (kt)									CO ₂ equivalent (kt)					
1. Energy	36'960	36'944	-16.75	289.1	288.5	-0.60	231.9	231.5	-0.33						
2. IPPU	2'236	2'259	22.99	2.1	2.1	0.00	48.3	48.0	-0.24						
3. Agriculture	46	46	0.23	4'147.5	4'174.1	26.66	1'980.7	1'930.6	-50.12						
4 LULUCF excl. HWP	-1'029	-923	106.04	13.3	13.3	0.00	69.4	74.5	5.02						
5. Waste	10	10	0.00	655.7	655.4	-0.22	191.4	189.5	-1.92						
6 Other	13	10.5	-2.42	0.7	0.6	-0.11	0.6	0.5	-0.13						
Sum (without F-gases)	38'236	38'346	110.1	5'108	5'134	25.73	2'522	2'475	-47.72						
										45'866	45'954	88.10			
Recalculation	HFC			PFC			SF6			NF3			Sum (synthetic gases)		
Emissions for 2014	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.	Prev.	Latest	Differ.
Source and Sink Categories	CO ₂ equivalent (kt)														
2 Ind. Processes (only syn. gases)	1'500.98	1'526.90	25.92	43.9	44.0	0.15	258.8	258.8	0.00	0.40	0.40	0.00	1'804.11	1'830.18	26.07
Recalculation													Sum (all gases)		
Emissions for 2014													Prev.	Latest	Differ.
Source and Sink Categories													CO ₂ equivalent (kt)		
Total CO₂ eq Em. with LULUCF													47'670	47'784	114.17
													100%	100.24%	0.24%
Total CO₂ eq Em. without LULUCF													48'617	48'620	3.11
													100%	100.01%	0.01%

10.3 Implications for emissions trends, including time series consistency

Due to recalculations, the emission trend 1990–2014 reported in the present 2017 submission has slightly changed. Compared to 1990, 2014 emissions (national total excl. LULUCF) showed a decrease of 8.80% before recalculation (Submission 2016). After recalculation in 2017, the decrease is slightly larger with a change 1990–2014 of 8.88%.

Table 10-5 Change of the emission trend 1990–2014 due to recalculation.

Recalculation	1990		2014		change 2014/1990	
	previous	latest	previous	latest	previous	latest
Submission						
Unit	CO ₂ eq (kt)				%	
Total excl. LULUCF	53'308	53'357	48'617	48'620	-8.80%	-8.88%

All time series in the present submission are consistent.

10.4 Planned improvements, including in response to the review process

- 1A1a: The N₂O emission factor of MSWIP is assessed periodically. It is planned to calculate a new weighted average N₂O emission factor for the year 2016 to be used in the next submission.
- 1A3b Road Transportation: A general update of the emission factors and the activity data is on-going. The update results will be presented in submission 2018 or latest in 2019.
- 1A3b: Note that an inconsistency in the attribution of natural gas to the vehicle categories leads to an error in the implied emission factors for gas-driven light duty vehicles. The error will be corrected within the general update mentioned above.
- 2F1: Calculation of commercial and industrial refrigeration are carried out together and will be split in response to the recommendations of the review process. As in the past years, all emission models will be updated during the yearly process of F-gas inquiry. The focus will be on HFC-emission calculations from refrigeration and air conditioning equipment, changes are expected in this area due to the revision of the relevant ordinance (ChemRRV) and CO₂ compensation programmes (share of products with HFC, recycling of HFC, early replacement of HFC).
- 3A: Planned improvements for future submissions are the further development, adaptation and verification of the dairy cattle feeding model (GE, Ym).
- 3B: Planned improvements for future submissions are the further development, adaptation and verification of the dairy cow feeding model (GE, DE, VS-excretion, N-excretion (Nex)).
New country-specific values for Nex for most livestock categories will become available in the next years due to a revision of the "Principles of Fertilization in Arable and Forage Crop Production (Flisch et al. 2009)".
A release of a new version of the AGRAMMON-Model is planned for 2017. If possible the respective AGRAMMON projections will be included during the next GHG inventory submission.
- NMVOC emissions will be revised in submission 2018 and separately reported for 3B Manure management and 3D Agricultural soils.
- 3D: New country-specific values for Nex for most livestock categories will become available in the next years due to a revision of the "Principles of Fertilization in Arable and Forage Crop Production (Flisch et al. 2009)".
A release of a new version of the AGRAMMON-Model is planned for 2017. If possible the respective AGRAMMON projections will be included during the next GHG inventory submission.
- NMVOC emissions will be revised in submission 2018 and separately reported for 3B Manure management and 3D Agricultural soils.
- 4A: Further research is underway to estimate annual values for carbon gains in living biomass. First results building on a study on the relationship between climate and gains in living biomass (Thürig et al. 2009) are expected in the next years.
- 4A: The implementation of the soil model Yasso07 to improve the accuracy in the estimates of temporal changes in soil carbon, litter and dead wood will be further developed. Planned improvements are in line with UNFCCC (2017/ID#L.12) and are foreseen at the earliest for the GHG inventory 1990–2016 (to be submitted in 2018).
- 4A: Projects in a new national research programme ([Sustainable Use of Soil as a Resource](#): "SOM control") aim at identifying the drivers of soil organic matter storage in Swiss forest soils.

- 4A: In response to issue ID#L.11 in UNFCCC (2017) data on the share of organic soils affected by past draining activities under Forest land will be collected for the next submission, using additional descriptive information from the NFI surveys (see chp. 6.4.2.10).
- 4B and 4C: Country-specific emission factor estimates (changes in carbon stocks) for cropland and grassland mineral soils will be improved by results of several research projects in the medium term.
- 4B, 4C, 4D, 4E: Recently, Price et al. (2017) created a nationwide model for tree biomass in Switzerland (both inside and outside of forest), using structural information available from airborne laser scanning. The model offers significant opportunity for improved estimates of carbon stocks in living biomass on land use combination categories where tree biomass has either not been included or only roughly estimated until now. The suitability of the obtained data for reporting in categories 4B, 4C, 4D, and 4E, respectively, will be evaluated.
- 4C: A study on GHG emissions from an extensively used fen under grassland management (Agroscope in collaboration with the University of Basel, 2013–2019, financed by FOEN) will improve the robustness of country-specific emission factor estimates for grassland soils rich in organic matter in the medium term.
- 4E: In response to UNFCCC (2017/ID#KL.4), the assumption that only 50% of the difference between the carbon stocks before and after the change is reported as a source or sink, respectively, for afforestation and deforestation will be reviewed.
- 4G: In response to UNFCCC (2017/ID#KL.8), the available data from the FAOSTAT database will be used to calculate the contribution of paper and paperboard to the HWP pool and report in the next submission. However, for the calculation there are some methodological challenges, such as how to determine the amount of domestic pulp wood contained in recycled products. Therefore, a study will be carried out to obtain additional national activity data on paper and paperboard to improve the calculation of the contribution of this pool.
- 4(IV): The fraction of mineralized N lost by leaching or run-off (Fra_{CLEACH}) used for calculating N_2O emissions in category 4(IV) will be harmonised with the updated value in the agricultural sector (see chp. 5.5.2.4.2).
- 5B1 Activity data for backyard composting is assumed to be approx. 10% of the amount of waste composted in industrial plants in the year 2007 and later. This share is not constant over time, but assumed to be approx. 3% in 1990 and 13% in 2000. These values are based on an expert judgement. It is planned for a subsequent submission to assess and verify the activity data. In the same course emission factors will be verified and completed if necessary.
- For further planned improvements please refer to respective chp. “planned improvements” for each category.

PART 2

11 KP-LULUCF

Switzerland has chosen to account over the entire second commitment period for emissions and removals from activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (FOEN 2016c, FOEN 2016d). In addition to the mandatory submission of the inventory years 2013, 2014 and 2015, data for the years 1990–2012 are available. Switzerland accounts for the mandatory activity Forest management under Article 3, paragraph 4, of the Kyoto Protocol (FOEN 2016c). Switzerland applies the condition of direct human-induced in relation to Afforestation and Deforestation very strictly for both activities (see chp. 11.1.3, FOEN 2010d, FOEN 2010h). CRF table NIR-1 shows the activity coverage and the carbon pools reported for the mandatory activities under Article 3, paragraph 3 and for Forest management under paragraph 4, of the Kyoto Protocol. Detailed information on completeness of the activity coverage and reported pools is given in chp. 11.3.1.2. The areas and change in areas between the previous and the current inventory year are shown in CRF table NIR-2. CRF table NIR-3 summarizes the results of the KCA for LULUCF activities under the Kyoto Protocol.

An overview of net annual CO₂ eq emissions and removals of activities under Article 3, paragraph 3 and Forest management under Article 3, paragraph 4, of the Kyoto Protocol is shown in Figure 11-1 and Table 11-1. Annual removals from Afforestation and Deforestation fluctuate (Figure 11-2), which can mainly be attributed to the changes in their respective areas (Table 11-2). The relative changes in the area of managed forest are comparatively low and fluctuations in the annual net changes in Forest management can primarily be explained by changes in the losses from the living biomass, dead wood and litter pools (Table 11-2). The reason for the high emissions in 2000 and the small removals in the following year 2001 for Forest management is the winter storm “Lothar” end of 1999, which caused great damages in the forest stands and increased losses of living biomass due to salvage logging. Harvesting rates in Swiss forests gradually increased since 1991 until 2007, reaching peak values in 2006 and 2007 and thus also resulting in small removals from Forest management in those years. Because harvesting rates started to decline in 2008 (Table 6-16) due to the international and domestic economic framework conditions, removals from Forest management increased since 2008, still showing high year-to-year variability. Fluctuations in the HWP pool can mainly be attributed to changes in the production of sawnwood (see chp. 6.11 and Table 6-34), which is strongly linked to the domestic harvesting rate.




Figure 11-1 GHG emissions (positive sign) and removals (negative sign), 1999–2015 (in kt CO₂). Shown are data for Afforestation (too small to be distinguishable) and Deforestation under Article 3, paragraph 3, Forest management (excluding HWP) and HWP under Article 3, paragraph 4.

Table 11-1 Net CO₂ eq emissions (positive sign) and removals (negative sign) for activities accounted for under Article 3, paragraph 3 and Forest management under Article 3, paragraph 4, of the Kyoto Protocol, 1990, 1995, 2000, 2005, and 2008–2015. Abbreviations are explained in chp. 6.1.3.2; except for loss_{N₂O}: N₂O emissions associated with land-use change (LUC) or from drainage; C_{L,ag}: carbon in above-ground living biomass; C_{L,bg}: carbon in below-ground living biomass; C_{s,m}: carbon in mineral soil; C_{s,o}: carbon in organic soil; C_{HWP}: carbon in HWP; loss_{biomburn}: CH₄ and N₂O emissions from biomass burning. Figures correspond with CRF table 4(KP) and CRF table 4(KP-I)B.1.

Greenhouse gas source and sink activities	1990	1995	2000	2005
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)			
A. Article 3.3 activities	90.28	107.91	123.99	118.66
A.1. Afforestation and Reforestation incl. N ₂ O; 4(KP)	-2.48	-12.20	-15.82	-19.12
Afforestation <= 20 yr	-2.49	-12.26	-15.90	-19.21
Afforestation > 20 yr	0.00	0.00	0.00	0.00
lossN ₂ O (from drainage and LUC)	0.01	0.06	0.08	0.09
A.2. Deforestation incl. N ₂ O; 4(KP)	92.76	120.11	139.81	137.78
Deforestation excl. N ₂ O	92.65	119.43	138.54	135.92
lossN ₂ O (from LUC)	0.11	0.68	1.27	1.86
B. Article 3.4 activities	-1'554.35	-4'152.86	4'526.25	-3'113.96
B.1. Forest management; 4(KP)	-1'554.35	-4'152.86	4'526.25	-3'113.96
gainC _{L,ag}	-9'759.85	-9'697.92	-9'728.97	-9'764.94
gainC _{L,bg}	-2'749.41	-2'751.06	-2'763.42	-2'776.95
lossC _{L,ag}	8'870.59	6'792.19	13'984.45	8'468.59
lossC _{L,bg}	2'682.91	2'005.86	3'902.33	2'461.04
changeC _h	387.10	-69.38	74.64	-534.03
changeC _d	177.31	74.39	-152.06	-246.16
changeC _{s,m}	-1.43	-0.26	-3.14	-4.56
changeC _{s,o}	35.88	36.23	36.45	36.69
changeC _{HWP}	-1'230.78	-560.89	-834.55	-763.06
Forest management excl. CH ₄ and N ₂ O; 4(KP-I)B.1	-1'587.70	-4'170.85	4'515.73	-3'123.37
loss _{biomburn} (CH ₄ and N ₂ O)	28.41	12.99	5.49	4.36
lossN ₂ O (from drainage)	4.94	4.99	5.02	5.06
B.2. Cropland management	NA	NA	NA	NA
B.3. Grazing land management	NA	NA	NA	NA
B.4. Revegetation	NA	NA	NA	NA

Greenhouse gas source and sink activities	2008	2009	2010	2011	2012	2013	2014	2015
	Net CO ₂ equivalent emissions/removals (kt CO ₂ eq)							
A. Article 3.3 activities	73.31	117.86	130.75	132.95	133.98	132.95	133.19	135.71
A.1. Afforestation and Reforestation incl. N ₂ O; 4(KP)	-21.35	-21.97	-20.61	-18.81	-18.35	-17.41	-15.31	-16.73
Afforestation <= 20 yr	-21.45	-22.07	-20.16	-18.35	-16.55	-15.10	-14.05	-13.46
Afforestation > 20 yr	0.00	0.00	-0.55	-0.55	-1.88	-2.38	-1.32	-3.32
lossN ₂ O (from drainage and LUC)	0.10	0.10	0.09	0.08	0.08	0.07	0.06	0.06
A.2. Deforestation incl. N ₂ O; 4(KP)	94.66	139.83	151.36	151.77	152.33	150.36	148.50	152.43
Deforestation excl. N ₂ O	92.51	137.57	149.09	149.49	150.04	148.07	146.21	150.14
lossN ₂ O (from LUC)	2.15	2.26	2.27	2.28	2.29	2.30	2.29	2.29
B. Article 3.4 activities	-1'693.07	-2'539.30	-2'514.69	-1'184.36	-2'520.72	-2'484.03	-1'077.73	-2'536.44
B.1. Forest management; 4(KP)	-1'693.07	-2'539.30	-2'514.69	-1'184.36	-2'520.72	-2'484.03	-1'077.73	-2'536.44
gainC _{L,ag}	-10'294.55	-10'306.39	-10'311.04	-10'316.05	-10'321.07	-10'327.22	-10'333.43	-10'337.72
gainC _{L,bg}	-2'947.67	-2'951.88	-2'953.99	-2'956.07	-2'958.14	-2'960.56	-2'962.97	-2'964.82
lossC _{L,ag}	9'492.18	8'861.42	9'330.45	9'260.44	8'521.49	8'785.24	9'033.29	8'399.62
lossC _{L,bg}	2'765.07	2'601.04	2'753.86	2'733.14	2'522.87	2'607.39	2'680.79	2'491.54
changeC _h	75.40	-97.58	-548.41	469.18	67.25	-193.72	604.86	72.14
changeC _d	-296.69	-316.89	-414.57	-160.23	-225.77	-274.70	-35.21	-169.13
changeC _{s,m}	-4.53	-4.67	-5.13	-5.56	-5.26	-5.39	-5.40	-4.95
changeC _{s,o}	37.04	37.15	37.23	37.31	37.39	37.52	37.65	37.72
changeC _{HWP}	-528.17	-370.39	-411.33	-257.74	-167.74	-160.87	-106.02	-69.53
Forest management excl. CH ₄ and N ₂ O; 4(KP-I)B.1	-1'701.92	-2'548.19	-2'522.94	-1'195.58	-2'528.97	-2'492.31	-1'086.44	-2'545.15
loss _{biomburn} (CH ₄ and N ₂ O)	3.74	3.78	3.12	6.08	3.09	3.12	3.52	3.52
lossN ₂ O (from drainage)	5.10	5.12	5.13	5.14	5.15	5.17	5.19	5.20
B.2. Cropland management	NA	NA	NA	NA	NA	NA	NA	NA
B.3. Grazing land management	NA	NA	NA	NA	NA	NA	NA	NA
B.4. Revegetation	NA	NA	NA	NA	NA	NA	NA	NA

The CRF table accounting (“Information table on accounting for activities under Articles 3.3 and 3.4 of the Kyoto Protocol”) gives an overview of the CO₂ eq emissions and removals from Afforestation and Deforestation under Article 3, paragraph 3 and Forest management under Article 3, paragraph 4.

In 2015, Forest management (including HWP, biomass burning, and drainage of organic soils) in Switzerland caused removals of -2536.44 kt CO₂ eq. The debit incurred from activities under Article 3.3 is 135.71 kt CO₂ eq.

11.1 General information

The inventory datasets on which the calculations are based (Swiss Land Use Statistics AREA and National Forest Inventory NFI) are described in chp. 6.2, 6.3 and 6.4.2.1, respectively.

Methodological issues and assumptions concerning the calculation of activity data and emission factors used for the reporting under Article 3, paragraphs 3 and 4, of the Kyoto Protocol follow the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006) as described in chp. 6.4.2 and the KP-Supplement (IPCC 2014).

11.1.1 Definition of forest and any other criteria

The forest definition used under the Kyoto Protocol is defined in Switzerland's first Initial Report (FOEN 2006h, Sect. E) and it is still valid for the second commitment period (FOEN 2016c; see also chp. 6.4.1 in this submission). Forest is defined as a minimum area of land of 0.0625 ha with crown cover of at least 20% and a minimum width of 25 m. The minimum height of the dominant trees must be 3 m or have the potential to reach 3 m at maturity in situ. The selected parameters are listed in CRF table NIR1.

Some source categories were explicitly excluded from the land-use category Forest land, although they may partly fulfil the requirements of the Swiss forest definition used under the Kyoto Protocol (see chp. 6.2.1, Table 6-6; chp. 6.4.1 and FOEN 2006h section E). Those are:

- Vineyards, Low-Stem Orchards, Tree nurseries, Copses and Orchards in the land-use category Grassland;
- Cemeteries and public parks in the land-use category Settlements.

11.1.2 Elected activities under article 3, paragraph 4, of the Kyoto Protocol

Switzerland only accounts for the mandatory activity Forest management under Article 3, paragraph 4, of the Kyoto Protocol (FOEN 2016c). In accordance with Annex I to Decision 2/CMP.7 (Annex I, Para 13), additions to the assigned amount resulting from Forest management under Article 3, paragraph 4, are capped. This cap for the second commitment period amounts to 15'037'884 t CO₂ for the entire commitment period 2013–2020 (FOEN 2016d).

11.1.3 Description of how the definitions of each activity under article 3.3 and each elected activity under article 3.4 have been implemented and applied consistently over time

The Swiss definitions of Afforestation, Deforestation and Forest management are published in Switzerland's Initial Report (see FOEN 2006h, Sect. E and F). These definitions are still valid for the second commitment period (FOEN 2016c). Switzerland applies the condition of direct human-induced in relation to Afforestation and Deforestation very strictly for both activities (see FOEN 2010d, FOEN 2010h).

For the notation of activities under article 3.3 and article 3.4 of the Kyoto Protocol, the first character is capitalised (see chp. 6.1.3.1): Afforestation, Deforestation and Forest management.

Afforestation

Afforestation is the conversion to forest of an area not fulfilling the definition of forest for a period of at least 50 years if the definition of forest in terms of minimum area (625 m²) is fulfilled, and the conversion is a direct human-induced activity (FOEN 2006h).

Natural forest regeneration following the abandonment of subalpine pastures is not considered to be a direct human-induced activity. Only conversions to forest land which can clearly be attributed as direct human-induced from aerial photographs (SFSO 2016d; see also chp. 6.3.1) are considered as Afforestation under the Kyoto Protocol. Examples of direct human-induced conversions to forest land (Afforestations) are shown in FOEN (2010h).

Deforestation

Deforestation is the permanent conversion of areas fulfilling the definition of forest in terms of minimum forest area (625 m²) to areas not fulfilling the definition of forest as a consequence of direct human influence (FOEN 2006h).

Temporary removals of (cluster of) trees (e.g. for the construction of high-voltage power lines, cable-car and powerlines, maintenance roads along railway lines and highways) are not reported as Deforestation under the Kyoto Protocol because in those cases the forest stand has to be re-established. In the NFI methodology (Brändli 2010: 91) "forest aisles" under power lines are explicitly classified as forests. These forest aisles underlie however a specific management, i.e. maximum tree height is limited to a certain height. The NFI dataset thus covers such areas with a specific forest management practice.

After approximately 12 years (see chp. 6.3.1) it is possible to check if deforestations or other land-use changes have been correctly classified. Sigmaplan (2012a) screened the classification of all land-use changes classified as Deforestation under the Kyoto Protocol. They found that 86% of all these Kyoto Deforestations were still deforested after 20 years, whereas 14% were temporary removals of tree cover, which should be classified as "management interventions" rather than as real land-use changes. As no reclassification was done, the area of Deforestations reported under the Kyoto Protocol Art. 3.3 is in fact an overestimation. Accordingly, emissions are overestimated since implied emission factors for Deforestations are higher than for Forest management (see CRF table 4(KP-I)A.2 for Deforestations and CRF table 4(KP-I)B.1 for Forest management). The area of the current land-use after Deforestation is given as information item in CRF table 4(KP-I)A.2. Since no

additional activities besides Forest management are elected under Art. 3.4, only the activity data are given and no information on changes in carbon stocks is provided.

Reforestation

Reforestation does not occur in Switzerland (FOEN 2006h, Sect. E; see also chp. 11.4.1).

Forest management

Forest management includes all activities serving the purpose of fulfilling the Federal Law on Forests (Swiss Confederation 1991, Art. 1c), i.e. the obligation to conserve forests and to ensure forest functions – such as wood production, protection against natural hazards, preservation of biodiversity, purification of drinking water, and maintenance of recreational value – in a sustainable manner.

11.1.4 Description of precedence conditions and/or hierarchy among 3.4 activities and how they have been consistently applied in determining how land was classified.

Since Switzerland only accounts for Forest management from the activities of Article 3, paragraph 4, of the Kyoto Protocol, the hierarchy among 3.4 activities does not affect Swiss reporting.

11.2 Land-related information

11.2.1 Spatial assessment unit used for determining the area of the units of land

The spatial assessment unit for the submission of the KP-reporting tables covers the entire territory of Switzerland, i.e. 4'129.03 kha (see chp. 6.3.1; Table 6-8).

All activity data for reporting the activities under the Kyoto Protocol are retrieved from the Swiss Land Use Statistics (SFSO 2016d; see also chp. 6.3.1). The Swiss Land Use Statistics AREA (SFSO 2006a) uses a georeferenced sample grid with a grid size of 100 m by 100 m. To each grid point a specific combination category (see Table 6-2) is assigned.

11.2.2 Methodology used to develop the land transition matrix

The methodology used to develop the land transition matrix is described in detail in chp. 6.2.3.

11.2.3 Maps and/or database to identify the geographical locations and the system of identification codes for the geographical locations

All Afforestations and Deforestations are accounted for under Article 3, paragraph 3 and are not reported under Forest management under Article 3, paragraph 4. Afforestations older

than the conversion period of 20 years are still reported under Afforestation: “Total for Activity A.1” in CRF Table 4(KP-I)A.1 equals the cumulated afforested areas since 1990 as shown in Table 11-2 and CRF table NIR-2. The area of deforestations displayed under “Total for activity A.2” in CRF table 4(KP-I)A.2 encompasses the cumulated area of deforestations since 1990 (see also Table 11-2 and CRF table NIR-2). However, only the cumulated area of deforestations of the last 20 years are relevant to calculate changes in carbon stocks (Table 6-3).

The calculation of changes in carbon stocks is described in chp. 11.3.1.1. The changes in areas between the activities under Article 3, paragraph 3 and Article 3, paragraph 4 are listed in CRF table NIR-2.

The area under Forest management is subdivided into productive forests (CC12) and unproductive forests (CC13; for a description see chp. 6.4.2.8). Productive forests in Switzerland reveal a high heterogeneity in terms of elevation, growth conditions and tree species composition (see chp. 6.2.2 and Figure 6-4). Therefore, Switzerland has been stratified into five National Forestry Inventory production regions (L1: Jura, L2: Central Plateau, L3: Pre-Alps, L4: Alps, L5: Southern Alps), three elevation zones (Z1: <601 m, Z2: 601-1200 m, Z3: >1200 m) and two soil types (mineral soils and organic soils). In the reporting tables, the stratification of the activity data into production region (L) and elevation level (Z) is indicated in the column “Subdivision”.

Area reported under Afforestation, Deforestation and Forest management

Land Use Statistics (AREA) data allow to clearly separate between the land areas subject to a specific activity. Absolute and cumulated activity data since 1990 of Afforestations and Deforestations and forest under Forest management are listed in Table 11-2. The total country area remains constant and amounts to 4'129.03 kha (Table 6-8).

Table 11-2 Activity data for activities under Article 3, paragraphs 3 and 4, 1990, 1995, 2000, 2005, and 2008–2015. Data for Afforestation and Deforestation and values depicting the area under Forest management were derived from the Swiss Land Use Statistics (AREA) (SFSO 2006a, 2016d). See also CRF table NIR-2.

	Unit	1990	1995	2000	2005
Afforestation area	kha	0.26	0.12	0.06	0.08
Cumulated area of Afforestation since 1990 [kha]	kha	0.26	1.28	1.66	1.98
Deforestation area	kha	0.30	0.34	0.36	0.33
Cumulated area of Deforestation since 1990 [kha]	kha	0.30	1.90	3.67	5.44
Forest management area	kha	1'210.65	1'221.59	1'229.11	1'236.22

	Unit	2008	2009	2010	2011	2012	2013	2014	2015
Afforestation area	kha	0.07	0.06	0.06	0.06	0.06	0.06	0.06	0.05
Cumulated area of Afforestation since 1990	kha	2.20	2.26	2.31	2.37	2.42	2.49	2.55	2.60
Deforestation area	kha	0.21	0.32	0.35	0.35	0.35	0.34	0.34	0.35
Cumulated area of Deforestation since 1990	kha	6.24	6.56	6.91	7.26	7.61	7.95	8.28	8.63
Forest management area	kha	1'241.89	1'244.03	1'245.20	1'246.26	1'247.33	1'248.59	1'249.84	1'250.93

Afforestation

Activity data for Afforestations are derived from the Swiss Land Use Statistics (AREA; SFSO 2006a, 2016d; see also chp. 6.3.1). A detailed description of the identification of Afforestations fulfilling the Kyoto definition is provided in FOEN (2010h).

Deforestation

Data for Deforestations are derived from the Swiss Land Use Statistics (AREA; SFSO 2006a, 2016d; see also chp. 6.3.1). A detailed description of the identification of Deforestations under the Kyoto Protocol from the AREA dataset is given in FOEN (2010d) and Sigmaphan (2010a).

Not all changes from a forest combination category (afforestation CC11, productive forest CC12 and unproductive forest CC13) to a non-forest combination category correspond to the definition of Deforestation according to the Kyoto Protocol Art. 3.3. The following criteria are used to identify conversions from a forest combination category to a non-forest combination category, which are not classified as Deforestations under the Kyoto Protocol Art. 3.3 (FOEN 2010d):

1. Non-permanent conversions are due to common forest management practices, natural dynamics or hazards:

- Tree loss is temporally limited, i.e., areas with loss of tree biomass, but where a change in land use cannot be identified: Natural regeneration, which is a common practice in Swiss forest management, is expected, but could not yet be recognized on the aerial photograph at the time the AREA survey (see chp. 6.3.1) was conducted. Also, in the NFI methodology (Brändli 2010: 91) "forest aisles" under power lines are explicitly classified as forests (see also chp. 11.1.3). Further, a study by Sigmaphan (2012a) showed that, although the aspect of "temporal limitation" was considered when classifying Deforestations, at the end still 14% of these Kyoto Deforestations were in fact "short-term reduction of crown coverage" and should be classified as "management interventions" rather than as real land-use changes (see chp. 11.1.3).
- Tree loss is spatially limited, i.e., conversion is caused by an alteration of the surrounding stand, but the change does not affect the tree cover at the sample point: this criterion applies also to the case of a Swiss-specific silvo-pastoral system of grasslands with tree cover. It is very difficult for interpreters of aerial photographs to determine this land use/land cover correctly. In fact, these points could be attributed to two coequal land-use types: agricultural area (NOLU04 2XX) and forest area (NOLU04 3XX; cf. Table 6-6). Land cover on these points is in general open forest (NOLC04 44), linear woods (NOLC04 46) or cluster of trees (NOLC04 47; cf. Table 6-6). When tree vegetation on these grasslands becomes denser over time, land owners remove single trees every now and then. This management practice can lead to the fact that an interpreter of aerial photographs reclassifies the sample point into a different land-use type during a later survey (i.e. change from forest area NOLU04 3XX to agricultural area NOLU04 2XX), although in reality no LUC took place on these sites; and, moreover, all elements of the Kyoto forest definition are still fulfilled (see Table 2 in FOEN 2010d).

2. Conversions of combination categories (see Table 6-2 and Table 6-6) which do not meet the definition of Deforestation as defined under the Kyoto Protocol and in Switzerland's Initial Report (FOEN 2006h):

- Areas smaller than the minimum area of 625 m².

- Areas with a reduction in forest cover on the grid point but still fulfilling the Kyoto definition of forest, i.e. having the potential to reach 3 m at maturity in situ.

3. No change in land use took place: reduction of tree cover without land-use change; former land use was mainly pasture

4. Tree loss is not direct human-induced: Conversion due to natural hazards and dynamics.

The four criteria were applied to the the land-use change data of the AREA survey (see chp. 6.2.2) for calculating annual values of the respective area (e.g. 1.030 kha in 2015, see Table 11-6).

It was ensured that the criteria and the application to identify conversions which do not correspond to Deforestations under the Kyoto Protocol do not result in inconsistencies in the estimates of changes in carbon stocks on the converted areas. If a sample point in the AREA dataset is not classified as Kyoto Deforestation, it remains classified as Forest management. The classification under Forest management implies that carbon stocks on these areas are based on NFI data (see NIR chp. 6.4.2.1). Thus, carbon stock changes are reflected in the Implied Emission Factors in the reporting tables and are completely accounted for.

Forest management

Since all forests in Switzerland are subject to certain forest management practices, the area under the activity Forest management corresponds to the forest area (see FOEN 2006h, Sect. E; FOEN 2016c) as derived from the Swiss Land Use Statistics (AREA; SFSO 2006a, 2016d; see also chp. 6.3). Changes in pools for the following geographical locations are reported:

- productive forest remaining productive forests (CC12 remaining)
- productive forest converted to unproductive forests (CC12 to CC13)
- unproductive forest remaining unproductive forests (CC13 remaining)
- unproductive forest converted to productive forests (CC13 to CC12).

Forest land is expanding in Switzerland (Table 11-2). The land-use change matrix (Table 6-9) shows that these conversions are mainly occurring on former grasslands (CC3X to CC12 or to CC13). The main reason is natural forest regeneration in the Alpine area due to the abandonment of land. Evidence for this process is provided, for example, by Rigling and Schaffer (2015), Brändli (2014), SWI (2009), and Gehrig-Fasel et al. (2007).

11.3 Activity-specific information

11.3.1 Methods for carbon stock change and GHG emission and removal estimates

11.3.1.1 Description of the methodologies and the underlying assumptions used

Emission factors for Afforestations, Deforestations and Forest management were accounted for following the methodology described in chp. 6.1.3.2. The methodological approach is

based on the details provided in Table 6-3, and on equations 6.1-6.8 and it is displayed in detail for each carbon pool in Table 11-4. Annual values for carbon stocks and carbon stock changes in the pools of living biomass, dead wood, litter and soil carbon of Afforestations (CC11), productive forests (CC12) and unproductive forests (CC13) are displayed in Table 6-4, Table 6-17, and Table 6-20. All working steps and data required to reproduce the calculation of emission factors the reporting tables are summarized in FOEN (2017c).

Separation of above- and below-ground living biomass

Carbon stock of total living biomass can be separated into above- and below-ground components using the ratios listed in Table 11-3. Carbon stock of total living biomass can be separated into above- and below-ground components using the ratios listed in Table 11-3. Under the UNFCCC both pools were reported in an aggregated way, under the Kyoto Protocol the pools were reported separately. For Forest management, the stratified ratios shown in Table 11-3 were used. For Afforestation and Deforestation the domestic mean value (0.30) was used.

Table 11-3 Root-to-shoot ratios to separate total living biomass into above- and below-ground living biomass. The ratios are retrieved from the NFI (Brändli 2010: Table 95).

NFI region	Elevation [m]	Root-to-shoot ratios for living trees
1	<601	0.22
	601-1200	0.27
	>1200	0.35
2	<601	0.22
	601-1200	0.24
	>1200	0.40
3	<601	0.23
	601-1200	0.28
	>1200	0.37
4	<601	0.25
	601-1200	0.30
	>1200	0.40
5	<601	0.28
	601-1200	0.32
	>1200	0.40
Switzerland	<601	0.23
	601-1200	0.27
	>1200	0.39
	average	0.30

Table 11-4 Application of the methodology described in equations 6.1-6.8 in chp. 6.1.3.2 and in Table 6-3 for calculating carbon stock changes for the Kyoto activities Afforestations (CC11) younger than 20 years (≤ 20 yr) and older than 20 years (>20 yr), Deforestations and Forest management with four “geographic locations” (CC12 remaining, CC13 remaining, CC12 to CC13, i.e. conversions from CC12 to CC13, and CC13 to CC12, i.e. conversions from CC13 to CC12). In the case of Afforestations (CC51 to CC11) and Deforestation (C1X to CC51), changes in carbon of mineral soils and of organic soils are accounted for by reducing the difference in soil carbon pools by 50% (see chp. 6.8.2.2). A conversion time CT of 20 years is applied for all pools except for the loss of living biomass, litter and dead wood after Deforestation (CT=1 year). Subscripts used: l = living biomass, h = litter, d = dead wood, s_m = mineral soil, s_o = organic soil, i = spatial stratum, a = land-use-type after the conversion, b = land-use-type before the conversion. CC11 (Afforestation), CC12 (productive forests) and CC13 (unproductive forests) refer to the specific combination category (see Table 6-2).

	Living Biomass	Litter	Dead Wood	Mineral Soil	Organic Soil
Afforestation CC11 ≤ 20 yr	gain-loss $\text{gain}C_{l,i,CC11} - \text{loss}C_{l,i,CC11}$	stock-difference, CT=20 $(\text{stock}C_{h,i,CC11} - \text{stock}C_{h,i,b})/CT$	stock-difference, CT=20 $(\text{stock}C_{d,i,CC11} - \text{stock}C_{d,i,b})/CT$	stock-difference, CT=20 C51 to CC11: $0.5(\text{stock}C_{s,m,i,CC11} - \text{stock}C_{s,m,i,b})/CT$ Other to CC11: $(\text{stock}C_{s,m,i,CC11} - \text{stock}C_{s,m,i,b})/CT$	gain-loss
Afforestation CC11 > 20 yr	gain-loss $\text{gain}C_{l,i,CC12} - \text{loss}C_{l,i,CC12}$	gain-loss $\text{change}C_{h,i,CC12}$	gain-loss $\text{change}C_{d,i,CC12}$	gain-loss $\text{change}C_{s,m,i,CC12}$	gain-loss
Deforestation	stock-difference, CT=1 $(\text{stock}C_{l,i,a} - \text{stock}C_{l,i,CC12})/CT$	stock-difference, CT=1 $(\text{stock}C_{h,i,a} - \text{stock}C_{h,i,CC12})/CT$	stock-difference, CT=1 $(\text{stock}C_{d,i,a} - \text{stock}C_{d,i,CC12})/CT$	stock-difference, CT=20 C1X to CC51: $0.5(\text{stock}C_{s,m,i,CC51} - \text{stock}C_{s,m,i,CC1X})/CT$ C1X to other: $(\text{stock}C_{s,m,i,a} - \text{stock}C_{s,m,i,CC1X})/CT$	C1X to CC51: stock-difference, CT=20 $0.5(\text{stock}C_{s,o,i,CC51} - \text{stock}C_{s,o,i,CC1X})/CT$ C1X to other: gain-loss
CC12 remaining	gain-loss $\text{gain}C_{l,i,CC12} - \text{loss}C_{l,i,CC12}$	gain-loss $\text{change}C_{h,i,CC12}$	gain-loss $\text{change}C_{d,i,CC12}$	gain-loss $\text{change}C_{s,m,i,CC12}$	gain-loss
CC13 remaining	gain-loss $\text{gain}C_{l,i,CC13} - \text{loss}C_{l,i,CC13} = 0$	gain-loss $\text{change}C_{h,i,CC13} = 0$	gain-loss $\text{change}C_{d,i,CC13} = 0$	gain-loss $\text{change}C_{s,m,i,CC13} = 0$	gain-loss
CC12 to CC13	stock-difference, CT=20 $(\text{stock}C_{l,i,CC13} - \text{stock}C_{l,i,CC12})/CT$	stock-difference, CT=20 $(\text{stock}C_{h,i,CC13} - \text{stock}C_{h,i,CC12})/CT = 0$	stock-difference, CT=20 $(\text{stock}C_{d,i,CC13} - \text{stock}C_{d,i,CC12})/CT = (0 - \text{stock}C_{d,i,CC12})/20$	stock-difference, CT=20 $(\text{stock}C_{s,m,i,CC13} - \text{stock}C_{s,m,i,CC12})/CT = 0$	gain-loss
CC13 to CC12	gain-loss $\text{gain}C_{l,i,CC12} - \text{loss}C_{l,i,CC12}$	stock-difference, CT=20 $(\text{stock}C_{h,i,CC12} - \text{stock}C_{h,i,CC13})/CT = 0$	stock-difference, CT=20 $(\text{stock}C_{d,i,CC12} - \text{stock}C_{d,i,CC13})/CT = \text{stock}C_{d,i,CC12}/20$	stock-difference, CT=20 $(\text{stock}C_{s,m,i,CC12} - \text{stock}C_{s,m,i,CC13})/CT = 0$	gain-loss

Reforestation

Reforestation does not occur in Switzerland (FOEN 2006h, Sect. E).

Afforestation ≤ 20 years: units of land not harvested since the beginning of the commitment period

Living biomass

- Gain and loss in living biomass of Afforestations (gross growth and cut and mortality) was taken from the study by Thürig and Traub (2015). Values are available for two elevation levels (Table 6-4).

Litter and dead wood

- On Afforestations, carbon stocks in litter and dead wood were assumed to be zero (IPCC 2006, chp. 4.3.2; see also chp. 11.3.1.2 for details). Applying the stock-difference calculation approach (Table 6-3), calculated changes in the litter and dead wood pool after an afforestation event were rather small since the major part of Afforestations (CC11) in Switzerland occur on grasslands CC31 and settlements CC5X (see Table 6-9), where there is no litter and no dead wood (Table 6-4).

Soil carbon

- Organic soils: In the case of organic soils, emissions due to drainage were calculated as described in chp. 6.4.2.10 and chp. 11.3.1.2.
- Mineral soils: In the case of land-use conversions to Afforestations, the difference in soil carbon stocks between land use before the conversion and after the afforestation event (CC11) was considered.
- In case of land-use changes to CC51 or from CC51 on mineral or organic soils, 50% of the difference between carbon stocks before and after the change were reported as emission and removal, respectively (see chp. 6.8.2.2).

Afforestation >20 years: units of land harvested since the beginning of the commitment period

After 20 years, afforested areas are subject to common forest management practices and the first thinnings and treatments are conducted. These afforested areas are, however, not reclassified to the activity Forest management: all afforestations after 1990 are consistently reported under Afforestation under Article 3.3 (CRF table 4(KP-I)A.1; see chp. 11.2.3). Emissions and removals for the carbon pools of Afforestations older than 20 years were calculated using the emission factors of productive (CC12) forests (see methodological description under Forest management), since nearly all of the afforestations (99.9%) develop into productive forests.

Deforestation

The differences in carbon stock of living biomass, litter and dead wood between Forest land and the land-use type after the conversion was immediately accounted for after Deforestation (conversion time = 1 year). Losses in soil carbon due to disturbance caused by Deforestation and conversion to CC51 (buildings and constructions) were accounted for by reducing the difference in soil carbon stocks by 50% (Covington 1981; Rusch et al. 2009; see also chp. 6.1.3.2) over a conversion period of 20 years (see Table 6-3; chp. 6.8.2.2).

Forest management

Living biomass

- Gain in living biomass (gross growth) of productive forests was used for “CC12 remaining” (Table 6-17). Gain of unproductive forests was used for “CC13 remaining” and amounts to zero (see chp. 6.4.2.8; Table 6-4).
- Losses in living biomass reflects yearly cut and mortality in productive forests “CC12 remaining” (Table 6-17). Unproductive forests are not systematically harvested (description chp. 6.4.2.8; Table 6-4). Thus losses of unproductive forests “CC13 remaining” are zero. Moreover, since yearly harvesting amounts from forest statistics (FOEN 2016k) are distributed over the productive forests, total harvesting in forests was accounted for under “CC12 remaining”.
- For the conversions between different forest combination categories (“CC13 to CC12” and “CC12 to CC13”) the method is chosen in such a way that no potential carbon losses are underestimated: For areas which changed from “CC12 to CC13” the difference in carbon stocks of living biomass was considered and a net loss in carbon stock of living

biomass is reported; in the case of a conversion from “CC13 to CC12” a gain-loss approach was applied, since applying a stock-difference approach would lead to a considerable sink in living biomass in this category.

Litter, dead wood and soil

- For productive forests “CC12 remaining”, values for yearly changes in carbon stock of litter, dead wood and soil were used (Table 6-20). Estimates of those yearly changes were obtained from simulations with Yasso07 (see chp. 6.4.2.7). For unproductive forests “CC13 remaining”, yearly changes in litter and dead wood and soil carbon stock were assumed to be zero (chp. 6.4.2.8).
- For the conversions between different forest combination categories (“CC13 to CC12” and “CC12 to CC13”) the difference in carbon stock of dead wood, litter and soil carbon was taken into account. For dead wood, the conversion “CC12 to CC13” leads to a net loss in carbon stock, in the case of a conversion “CC13 to CC12”, a net gain is reported (Table 6-4).
- In the case of organic soils, emissions due to drainage were calculated as described in chp. 6.4.2.10.

Differences in accounting for “Forest categories 4A1 and 4A2” under the UNFCCC and Forest management under the Kyoto Protocol Art. 3.4

Under the Kyoto Protocol Art. 3.4, natural forest regeneration is reported under Forest management as productive forest (CC12) or unproductive forest (CC13) as soon as the KP definition of forest is fulfilled (CRF table NIR-1). Changes within the activity Forest management are reported under the Kyoto Protocol in the combination categories “CC12 to CC13” and “CC13 to CC12”.

Under the UNFCCC, all changes in land use from non-forest land to forest land are reported in the land-use category 4A2 for a conversion time of 20 years. For further details and a quantitative comparison see chp. 11.3.2.2.

11.3.1.2 Justification when omitting any carbon pool or GHG emissions/removals from activities under article 3.3 and elected activities under article 3.4

CRF table NIR-1 summarizes the activity coverage and the carbon pools reported. When using the Tier 1 approach (IPCC 2006 Volume 4, chp. 1.3) assuming a specific carbon pool to be in balance, the carbon pool is indicated as not reported (NR). This is the case for litter and dead wood under Afforestation. Also for all pools of unproductive forests, no changes are reported.

Changes in carbon pools not reported - Afforestation ≤ 20 years: litter and dead wood

Applying the stock-difference calculation approach (cf. Table 11-4), calculated changes in the litter and dead wood pool after Afforestation under 20 years are zero (see chp. 6.4.2.9), since on Afforestations carbon stock in litter and dead wood is assumed to be zero (IPCC 2006, Volume 4, chp. 4.3.2).

Because Afforestation is not a key category (see chp. 11.6.1) and carbon stock changes in litter and dead wood were considered too negligible to justify the (financial) effort to collect higher quality data (see Figure 4.1 in Volume 1 of IPCC 2006), a Tier 1 estimate for these pools for Afforestations under 20 years was applied. Verifiable information to justify this approach is provided here:

- Changes in litter after afforestation: Under the Kyoto Protocol, changes in the litter pool after Afforestations were not reported. In an experiment by Zimmermann and Hiltbrunner (2012) litter accumulation of an afforestation with Norway Spruce was determined 40 years after afforestation. The authors found accumulation rates of 0.17-0.20 t C ha⁻¹ yr⁻¹. Other studies show even higher accumulation rates, e.g., 0.24-0.34 t C ha⁻¹ yr⁻¹ for afforestations with Norway spruce in the Southern Alps (Thuille and Schulze 2006), 0.24 t C ha⁻¹ yr⁻¹ for afforestation with ash and maple (Alberti et al. 2008) and 0.36 t C ha⁻¹ yr⁻¹ for Scotch pine (Vesterdal et al. 2002). In Finnish forests, Karhu et al. (2011) found that over 18 years the mean annual rate of carbon accumulation in the litter was 0.28 Mg ha⁻¹ for Scots pine and 0.15 Mg ha⁻¹ for birch.
- Based on a literature overview, Jandl et al. (2007) argued that the accumulation of a forest floor layer in, e.g., a conifer forest, results in a carbon sink. The authors concluded that after afforestation, forest floors accumulate carbon quickly. A long-term consequence of afforestation is the gradual incorporation of carbon in the carbon pool of the mineral soil. Guidi et al. (2014) found that the carbon stocks in the organic layers were affected by land-use change, with more carbon stored under early-stage forest compared with grassland abandoned 10 years ago, and highest carbon stocks were found under the old forest dominated by *Fagus sylvatica* and *Picea abies*.
- Changes in dead wood after afforestation: no changes in the dead wood pool after afforestations were reported. Zimmermann and Hiltbrunner (2012) showed that 40 years after afforestation with Norway Spruce, dead wood volume amounted to 10.4 t C ha⁻¹. This corresponds to an annual increase of carbon stored in the dead wood of 0.26 t C ha⁻¹ yr⁻¹ for afforestations with Norway Spruce.
- Besides the results of the case studies listed above, a reasoning based on sound knowledge of likely system responses (Grassi and Blujdea 2011) was provided: At stand level, the pools dead wood and litter of afforestation on cropland and grassland cannot be a source, especially if the previous land use did not have perennial woody biomass. On afforestations, tree growth is assumed to follow an exponential pattern, which can also be assumed for the accumulation of litter and dead wood.

Note that for Afforestations older than 20 years, estimates of carbon stock changes in dead wood, litter and soil were reported.

Changes in carbon pools not reported - Unproductive forests

A description of unproductive forests and the reasoning why the living biomass, litter, dead wood and soil pools were reported to be in equilibrium and thus not a source is given in detail in chp. 6.4.2.8.

Based on the fact that unproductive forest land only covers 8-9% of the area under Forest management (CRF table 4(KP.I)B) and based on the description of these stands in chp. 6.4.2.8, emissions or removals of any of the pools of unproductive forests cannot account for more than 25% of the activity Forest management. According to Fig 1.2 note 4 in IPCC (2006), Volume 4, 25% is the threshold that would require a higher Tier. Because of limited financial resources (IPCC 2006 Volume 1, chp. 4.1.2, Fig. 4.1), Switzerland decided to use the Tier 1 approach and reported no changes in the carbon pools of living biomass, litter,

dead wood and mineral soil of unproductive forest areas. Emissions from organic soils were accounted for using default factors from IPCC 2014 (Tier 1).

Greenhouse gas sources reported

- Fertilization of forests is prohibited by the Swiss forest law and adherent ordinances (Swiss Confederation 1991, 1992). Additionally, the “Ordinance on Chemical Risk Reduction” (Swiss Confederation 2005) prohibits the application of fertilizers, including liming, in forests. Thus, emissions from fertilization were reported as “not occurring”.
- Drainage of forests is not a permitted practice in Switzerland and since 1991 not a permitted practice in Switzerland (Swiss Confederation 1991). In order not to underestimate the emissions, all organic forest soils were reported to be drained. CO₂ emissions due to drainage are calculated as described in chp. 6.4.2.10. N₂O emissions from drainage of organic soils was calculated as described in chp. 6.4.2.11.
- Biomass burning: emissions of CO₂ were given the notation “IE”; emissions of CH₄ and N₂O were reported. The calculation of these emissions is described in chp. 6.4.2.12.

Greenhouse gas sources reported as “included elsewhere (IE)”

- Emissions from Biomass burning on Afforestations were reported under Forest management. In this way, emissions were not underestimated, since the carbon stock (“available fuel”) in productive forests is higher than the carbon stocks in afforestations and unproductive forests. Moreover, this approach reflects reality well, since fires in afforestations or in unproductive forests are rather unlikely to occur (see chp. 6.4.2.12).
- Biomass burning on areas under Forest management: CO₂ emissions were reported as “IE”. The reported losses of living biomass and dead wood are covered by NFI data and thus the values reported in the reporting tables Table4.A and 4(KP-I)B.1 include these losses. Emissions of CH₄ and N₂O were reported. The calculation of these emissions is described in chp. 6.4.2.12.

Greenhouse gas sources reported as “not occurring”

- HWP from Afforestation: based on the available datasets it was not possible to differentiate between HWP from Afforestation and HWP from Forest management. The amount of HWP from Afforestation is not existing or negligibly small. Therefore, all carbon stock changes in HWPs from Afforestation were reported under Forest management (see chp. 6.11.2). HWP from Deforestation was accounted for on the basis of instantaneous oxidation “IO”.

11.3.1.3 Information on whether or not indirect and natural GHG emissions and removals were factored out

No anthropogenic GHG emissions and removals from elevated carbon dioxide concentrations, indirect nitrogen deposition or the dynamic effects of the age structure resulting from LULUCF activities under Article 3, paragraphs 3 and 4 prior to 01 January 1990 were factored out.

The IPCC does not give specific methods for factoring out these effects. Besides this, there are no reliable country-specific data available. Investigations on elevated CO₂ concentrations on growth showed complex relationships in the mid-term. Some species showed an increase

others a decrease and some no change in growth (Bader et al. 2013). Opposing patterns are also reported regarding the effect of nitrogen deposition: A positive effect of N deposition on growth was found by e.g. Spiecker (1999) and Jarvis and Linder (2000). Other studies (e.g., Hyvönen et al. 2008; Högberg et al. 2006; Braun et al. 2010; Gschwantner 2006; Meining et al. 2008) indicate that N-deposition, while leading to soil acidification, can cause a reduction in growth. Such acidification processes are widely detected in Swiss forest soils (Braun and Flückiger 2012).

11.3.1.4 Changes in data and methods since the previous submission (recalculations)

Due to unresolved technical issues around the CRF Reporter Switzerland was not in a position to provide correct reporting tables CRF Summary3s1 and Summary3s2 recommended by the ERT in issue ID#G.1 in UNFCCC (2017). To address the issue manually corrected reporting tables are presented in Annex 6 (Table A – 35 and Table A – 36).

Table 10-1 (recommendations) and Table 10-2 (encouragements) list the improvements and planned improvements made since the last submission (FOEN 2016) in response to the questions, recommendations and encouragements of the UNFCCC Expert Review Team during the in-country review 2016.

Methodological improvements for Forest land valid for the LULUCF and the KP-LULUCF sector are listed in chp. 6.4.5. The following Kyoto-specific methodological modification was made.

- Net CO₂ emissions and removals from Harvested wood products (HWP) from deforestations were accounted for on the basis of instantaneous oxidation.

11.3.1.5 Uncertainty estimates

Uncertainty estimates of activity data are discussed in detail in chp. 6.1.5 and are shown in Table 6-10. Uncertainty estimates of emission factors for the reported activities under the Kyoto Protocol are shown in Table 6-5, overall uncertainties in Table 11-5.

A detailed description of the determination of the emission factor uncertainty of Forest management can be found in chp. 6.4.3. Table 6-5 lists the relative uncertainties in the LULUCF sector: an uncertainty of 88.6% was calculated for Afforestations, 50.0% for Deforestations and 88.6% for Forest management.

Lands fulfilling the definition of forest (see chp. 11.1.1) were accounted for under Forest management. This means, that the area under Forest management resulting from natural regeneration, is attributed the uncertainty of Forest management.

Table 11-5 Uncertainty estimates of activity data and emission factors and the overall uncertainty of activities reported under the Kyoto Protocol Article 3.3 and Article 3.4.

Activity under KP	Associated category in UNFCCC inventory (chp. 6.3)	Activity data uncertainty	Emission factor	Combined uncertainty
		%	%	%
Afforestation	4A2 Land converted to forest land	1.6	88.6	88.6
Deforestation	mainly 4E2 Land converted to settlements	4.5	50.0	50.2
Forest management	4A1 Forest land remaining forest land	1.1	88.6	88.6

11.3.1.6 Information on other methodological issues

N₂O emissions as a result of the disturbance associated with land-use conversion (Deforestation) were reported in CRF table 4(KP-II)3. The emissions were calculated according to the methodology described in chp. 6.10.

11.3.1.7 The year of the onset of an activity, if after 2013

The starting year of the activities reported can directly be derived from the land-use change matrix (Table 6-9), from which a continuous time series was derived (Table 11-2).

11.3.2 Category-specific QA/QC and verification

In chp. 6.4.4 category-specific QA/QC and verification items for forest land are described. The general QA/QC measures are described in chp. 1.2.3.

11.3.2.1 Changes in soil carbon stock under Afforestation

The assumption that soils are acting as small sinks under Afforestation is supported by Jandl et al. (2007) who reviewed several studies on the effect of different forest management systems (including afforestations) on soil carbon sequestration and concluded that a long-term consequence of afforestation is the gradual incorporation of carbon in the mineral associated soil carbon pool.

11.3.2.2 Comparison of the forest areas reported in the reporting tables

A direct comparison of the areas reported in the reporting tables under the Convention Forest land remaining forest land (CRF Table 4.A) and under Forest management under the Kyoto Protocol (CRF table 4(KP-I)B.1) is not possible due to the different structure of these reporting tables and due to different reporting requirements:

- Conversions to Forest land which are not human-induced (natural regeneration) were not accounted for as Afforestations under the Kyoto Protocol. These areas were reported under KP Art. 3.4 Forest management in CRF table 4(KP-I)B.1 as soon as the definition of Forest was fulfilled. Under the Convention, these afforestations were reported under land-use category 4A2 with a conversion time of 20 years.
- Afforestations under the Kyoto Protocol which are older than 20 years were consistently reported under Art. 3.3 (sub-division >20 years in CRF table 4(KP-I)A.1: units of land harvested since the beginning of the commitment period). Thus, there is no

reclassification of the units of lands reported under Art. 3.3. In contrast, under the UNFCCC, afforestations older than 20 years were reallocated to the land-use category 4A1 Forest land remaining forest land.

- Not all changes from a forest combination category (CC11, CC12, CC13) to a non-forest combination category correspond to the definition of Deforestation according to the Kyoto Protocol Art. 3.3. (see above). These areas remained under the Kyoto Protocol Art. 3.4 activity Forest management and were included in the areas as reported in CRF table 4(KP-I)B.1.
- Reporting of land-use changes LUC: Since only the KP activity Forest management is accounted for under KP Art. 3.4, changes from other KP activities to Forest land were not reported as land-use change, but were reported as CC12 or CC13 as soon as the KP definition of forest was fulfilled. Only conversions within the activity Forest management were reported under the Kyoto Protocol, i.e. CC12 to CC13 and CC13 to CC12. Under the UNFCCC, land-use change to forest land were reported in category 4A2.

In Meteotest (2017) the reported activity data for the inventory year 2015 were examined and compared (Table 11-6). The differences in the reporting tables Table4.A, 4(KP-I)A.1 and 4(KP-I)B.1 can be explained and the resulting budget of areas reported under the Convention and the Kyoto Protocol are identical.

Table 11-6 Area budget (in kha) of KP-LULUCF and LULUCF under the UNFCCC in the year 2015 for forest land. The references in the table are valid for the reporting tables used until 2015 (Meteotest 2017).

Activity	Table, Cells	Area UNFCCC 2015 kha	Area KP 2015 kha	Check Difference kha	Remarks
All Forest Land					
Forest management	4(KP-I)B.1, D11		1'250.932		a)
Afforestations <= 20 years	4(KP-I)A.1, C29		1.318		b)
Afforestations > 20 years	4(KP-I)A.1, C13		1.284		c)
Total area KP			1'253.534		
Non-Kyoto loss of forest cover			-1.030		d)
Forest land UNFCCC	4.A, C10	1'252.504			e)
Total		1'252.504	1'252.504	0.000	
Afforestation, CC11					
UNFCCC	4.A, C32+C36+C40 +C44+C48	1.318			f)
KP (<= 20 years)	4(KP-I)A.1, C29		1.318	0.000	g)

Remarks:

a) KP Forest management consists of CC12 and CC13 areas fulfilling the criteria of the KP.

b) KP Afforestations are afforested areas since 1990 cumulated over 20 years at most.

c) KP Afforestations "older than 20 years" (>20 years) comprise the area that has been afforested since more than 20 years. In the UNFCCC reporting tables these areas belong to 4A1 (CC12 or CC13).

d) The non-Kyoto loss of forest cover is the part of the total area of forest loss (reported under the UNFCCC) not fulfilling the definition of deforestations according to the Kyoto Protocol (see chp. 11.2.3). For the comparison this area must be subtracted from the KP forest area. It is an annual value (not cumulated) calculated on the basis of the AREA survey data (chp. 6.2.2).

e) The total Forest land in CRF Table4.A covers productive forests (CC12), unproductive forests (CC13) and afforestations (CC11). It is congruent with the forest area derived from the aerial photos of the AREA survey (chp. 6.2.2).

f) The CC11 area under the UNFCCC can be derived from CRF Table4.A.2 by summing up the afforestation categories.

g) The cumulated (20 years) CC11 area of KP and UNFCCC are congruent.

11.3.2.3 Impact of forest management on changes in carbon stocks in soil and in litter

Accounting for forest management impacts on carbon storage in litter and soil in Swiss productive forests with Yasso07

To estimate carbon stocks and carbon stock changes in the reported litter and soil pools, Switzerland uses the carbon cycling model Yasso07 (Didion et al. 2012, Didion and Thürig 2016; chp. 6.4.2.7). Yasso07 requires information on carbon inputs from dead organic matter (i.e. non-woody inputs, including foliage and fine roots, woody inputs, including standing and lying dead wood and dead roots) and climate (temperature, temperature amplitude and precipitation). The carbon inputs are obtained for each plot in the NFI that is simulated with Yasso07. The NFI plots were repeatedly measured since the first inventory in 1985 and, hence, observed changes in the volume of living and dead biomass reflect, among other, the site-specific impact of forest management practices. Based on harvesting statistics and allometric relationships, the production of dead wood (incl. dead roots, stems, stumps and branches) and litter from living trees (i.e. controlled by forest management) and as harvest residues were estimated.

Thus, the Yasso07-model reflects the impact of forest management practices: effects of common forest management on carbon stocks in litter (including non-woody and woody material) and soil were fully accounted for in the GHG inventory (Didion 2014).

Literature Review

A detailed screening of the available scientific literature on the impact of forest management on carbon stock changes in litter and soils is provided in Didion (2014). The majority of studies indicated no significant effect of forest management on soil carbon stocks with the exception of clearcutting (e.g. Jandl et al. 2007). Since silvicultural practices in Switzerland are regulated by law and exclude intensive management options such as clearcuts, fertilization or liming (Swiss Confederation 1991, 1992), no or only minor forest management impacts on soil carbon stocks can be expected. The production of litter is directly affected by silvicultural practices since the removal of trees results in harvest residues and in a decrease in the amount of remaining foliage (e.g. Van Miegroet and Olsson 2011). Generally, the impact of forest management on litter production is temporary and losses of litter carbon can be rapidly replaced (Nave et al. 2010).

11.4 Article 3.3.

Net removals from Afforestation and emissions from Deforestation under Article 3, paragraph 3 differ by one order of magnitude (Figure 11-2, Figure 11-3). Since carbon from living biomass is immediately removed after clear-cutting, deforestation can be seen as a process where carbon is lost over a very short time. In contrast, afforestation is a slow process where carbon is sequestered and accumulated over decades. CO₂ emissions on organic soils under Afforestation and Deforestation are due to former drainage (see chp. 11.3.1.2). Figure 11-2 shows removals of CO₂ eq from Afforestation and emissions of CO₂ eq from Deforestation for the years 1999–2015. Corresponding data for 2000, 2005, and 2008-2015 are listed in Table 11-1.




Figure 11-2 Removals (negative sign) and emissions (positive sign) from Afforestation under Article 3, paragraph 3 (upper panel) and emissions from Deforestation under Article 3, paragraph 3 (lower panel) shown per carbon pool, 1999–2015 (in kt CO₂ eq). For abbreviations see Table 11-11.




Figure 11-3 Net removals (negative sign) of Afforestations under Article 3, paragraph 3 and emissions (positive sign) of Deforestations under Article 3, paragraph 3, 1999–2015 (in kt CO₂ eq).

11.4.1 Information that demonstrates that activities under Article 3.3. began on or after 01 January 1990 and before December 2020 and are direct human-induced.

The Swiss definitions of Afforestation and Deforestation only consider direct human-induced activities (see FOEN 2006h, Sect. E and FOEN 2010d).

Reforestation

For more than 100 years, the area of forest in Switzerland has been increasing (see chp. 11.5.2). A decrease in forest area as a result of deforestation is not possible, since deforestation is strongly regulated by the Federal Law on Forests (Swiss Confederation 1991). Therefore, reforestation of areas not forested for a period of at least 50 years does not occur in Switzerland (FOEN 2006h, Sect. E). Switzerland only considers Afforestation and Deforestation under Article 3, paragraph 3.

Afforestation

Switzerland is very restrictive in reporting Afforestations under the Kyoto Protocol and only reports planted forests under Afforestation (see chp. 11.1.3; FOEN 2010h).

The annual rate of all afforested areas since 1990 is assessed based on AREA data (chp. 6.3). For reporting under the Kyoto Protocol, afforested areas since 1990 always remain in the Afforestation category. Therefore, the area in this category has been increasing since 1990 (see Table 11-2).

Afforestations older than 20 years are subject to common forest management practices including harvesting (see chp. 11.3.1.1). These areas are reported in CRF table 4(KP-I)A.1.

Deforestation

In Switzerland, direct human-induced Deforestation is subject to authorization (Swiss Confederation 1991, Art. 5). Deforestation is only allowed for projects with public interests and in these cases, the deforestation has to be compensated by an afforestation of equal area.

For details concerning the classification of Deforestations under the Kyoto Protocol see chp. 11.2.3). Only deforestation events carried out after 01 January 1990 are considered. For reporting under the Kyoto Protocol, deforested areas since 1990 remain in the Deforestation category. Therefore, the area in this category has been increasing since 1990 (see Table 11-2). Since Switzerland only accounts for KP Art. 3.4 activity Forest management, these deforested areas are not accounted for under another KP Art. 3.4 activity.

11.4.2 Information on how harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from Deforestation

The Swiss definition of Deforestation only covers permanent conversions from Forest land to non-forest land. In the process of the interpretation of land conversions based on AREA data (chp. 11.2.3), the definition is implemented by applying the criteria discussed in chp. 11.2.3. This approach was verified by Sigmaplan (2012a).

The criteria distinguish between permanent conversions and transient situations like harvesting or forest disturbance followed by forest re-establishment. Construction of e.g. pipelines and power supply lines within a forest area are transient situations (see chp. 11.1.3 and 11.2.3; Brändli 2010). As described in FOEN (2010d), these non-permanent conversions are not classified as Deforestation under the Kyoto Protocol.

11.4.3 Information on the size and geographical location of forest areas that have lost forest cover but which are not yet classified as deforested

The AREA survey provides a detailed overview of land-use changes with regard to land cover and land use (see chp. 6.2 and 6.3). Temporal changes of land cover can lead to a reclassification in AREA from a forest combination category to a non-forest combination category. However, not all changes from Afforestation CC11, productive forest CC12, or unproductive forest CC13 to a non-forest combination category correspond to the definition of Deforestation according to the Kyoto Protocol Art. 3.3. Explicit criteria were developed (cf. FOEN 2010d and chp. 11.2.3) to identify which conversions from a forest combination category to a non-forest combination category do not correspond to Kyoto Deforestation under the Kyoto Protocol.

11.4.4 Information related to the natural disturbances provision under Article 3.3

Switzerland does not apply the provision of exclusion of natural disturbances for Afforestation and reforestation under Article 3, paragraph 3, of the Kyoto Protocol (FOEN 2016c).

11.4.5 Information on Harvested wood products under Article 3.3

The calculation of carbon stock changes in Harvested wood products HWP is described in detail in chp. 6.11. The change in carbon stocks was estimated differentiating HWPs originating from Afforestation, Deforestation and from Forest management.

Applying instantaneous oxidation to HWPs originating from Deforestations, the same results are obtained for changes in carbon stocks of HWP reported under the UNFCCC (CRF Table 4.Gs1) and under the Kyoto Protocol (CRF table 4(KP-I)C) as shown in Table 6-34.

11.5 Article 3.4

GHG emissions and removals differentiated for the reported carbon pools and net GHG emissions and removals of KP Article 3, paragraph 4 activity Forest management for the years 1999–2015 are shown in Figure 11-4. Corresponding data for 2000, 2005, and 2008–2015 are listed in Table 11-1. The annual fluctuations in the GHG emissions and removals from Forest management are described in the context of Figure 11-1.




Figure 11-4 Emissions (positive sign) and removals (negative sign) broken down for the reported carbon pools under Forest management (upper panel) and the net emissions and removals from Forest management (lower panel), 1999–2015 (in kt CO₂ eq). For abbreviations see Table 11-1.

11.5.1 Information that demonstrates that activities under Article 3.4. have occurred since 1 January 1990 and are human-induced

According to the Federal Act on Forest, the extent and the spatial distribution of the total forest area in Switzerland has to be preserved (Swiss Confederation 1991, Art. 1) and thus, any change of the forested area has to be authorized. All Swiss forests are under continuous observation of the Swiss Forest Service and monitored by the NFI. Therefore, all forests in Switzerland are subject to forest management and reported under Forest management under Article 3, paragraph 4 of the Kyoto Protocol (FOEN 2006h, Sect. F).

11.5.2 Information relating to Forest management

There is a long tradition of forest protection in Switzerland. The first federal Forest Act came into force in 1876, but it only covered the higher-elevation regions. Its aims were to put a halt to the depletion of forests, to manage the remaining forest areas in a sustainable way, and to promote afforestation. The Forest Act of 1902 covered the whole country. The Forest Act as well as an increasing economic development resulted in an increase of the forested area in Switzerland by nearly 50% today compared to the mid-19th century (Figure 11-5). Also the growing stock increased significantly due to changes in forest management practices. The Forest Act (Swiss Confederation 1991) that came into force in 1993 reaffirmed the long-standing Swiss tradition of preserving both forest area and forest as a natural ecosystem. It prescribes sustainable forest management, prohibits clearing, and bans deforestation unless it is replaced by an equal area of afforested land or an equivalent measure to improve biodiversity.




Figure 11-5 Historical mean growing stock ($\text{m}^3 \text{ha}^{-1}$) and forest area (kha) in Switzerland since 1880.

In 2004, the Swiss national forest programme was published, outlining an action plan for the period 2004–2015 (SAEFL 2004b). It specifies five priority objectives: (1) the forests' protective function is guaranteed, (2) the economic viability of the forestry sector is improved, (3) the value-added chain for wood is strengthened, (4) biodiversity is conserved and (5) forest soils, trees and drinking water are not threatened. These objectives encompass that CO_2 removals by sinks and emissions by sources in the forests shall be recognized in terms of compliance with the Kyoto Protocol while making better use of the potential of forests for timber production and fuel wood through economic incentives and implementing new technologies.

In November 2006, the Swiss government communicated in its Initial Report to the UNFCCC that Switzerland will be accounting for Forest management under Article 3.4 of the Kyoto Protocol (FOEN 2006h).

To implement the objectives of the national forest programme (SAEFL 2004b), FOEN has formulated its wood resource policy (FOEN 2008h) which is coordinated with the other relevant sectoral policies (e.g. energy policy, regional development policy). This wood resource policy defines, among other things, the direction to be taken by federal policy in relation to wood promotion on completion of the “Wood 21” wood promotion programme which was finalized at the end of 2008. Under this programme, a wood action plan was started in 2009. The main focus in the implementation of the action plan lies on the ecologically and economically effective use of wood. Regarding the efficient use of wood, cascade use is prioritized, i.e. wood is used as material prior to its use for energy. In the case of energy use, greater overall efficiency of the conversion technology should be targeted.

11.5.2.1 Conversion of natural forest to planted forest

Not applicable.

Switzerland did not choose to apply the concept of carbon equivalent forests (see CRF table 4(KP-I)B.1.2).

11.5.2.2 Forest management reference level (FMRL)

Switzerland's Forest management reference level (FMRL) is documented in FOEN (2011). The Swiss FMRL is inscribed in the appendix to the annex to Decision 2/CMP.7 and amounts to +0.220 Mt CO₂ eq. yr⁻¹. The FMRL was subject to a technical assessment. Based on the technical assessment report (UNFCCC 2011a) and applying guidance of IPCC (2014), a technical correction of Switzerland's FMRL was made as described in FOEN (2016: chp. 11.5.2.3).

11.5.2.3 Technical correction Forest management reference level

The technical corrections were implemented for the submission in 2015 (FOEN 2015) and were described in detail in FOEN (2016). For the submission in 2017, no improvements leading to a technical correction of the FMRL were implemented.

Switzerland decided not to provide technical corrections of the FMRL on an annual basis, but to correct the FMRL periodically.

In order to reflect the most recent scientific knowledge, data availability and model versions and to ensure methodological consistency between the reference level and reporting for Forest management (corrections are listed in chp. 10.1.2.5.1), the next technical correction of the FMRL will be reported in Switzerland's submission in 2019 including the following improvements:

To ensure ensure methodological consistency between the reference level and reporting for Forest management, the technical corrections listed in chp. 10.1.2.5.1 will be applied.

In response to UNFCCC (2017/ID#KL.4), the assumption that only 50% of the difference between the carbon stocks before and after the change is reported as a source or sink, respectively, for afforestation (from settlements to forest land) and deforestation (from forest land to settlements) will be reviewed.

For the Massimo model, it is planned to improve the algorithms and assumptions, the code, the usability and the documentation. The major aspects of the revision, which will be implemented over the coming years, include:

- Improved representation of natural mortality to more accurately reflect the background level for natural disturbances (see chp. 11.5.2.4.2).
- Verification of allometries used to obtain estimates of whole tree volume and biomass, incl. branches, foliage, and roots.
- To ensure methodological consistency between the reference level and reporting for Forest management, the most recent NFI data will be used (see chp. 10.1.2.5.1).

Planned improvements for Yasso07 are described in chp. 6.4.6.

- Investigating the validity of the further development of Yasso07 for application in Switzerland.
- Improving the completeness of the litter inputs by accounting for the contribution of a) fine-woody litter <7 cm and b) litter from the herb- and shrub layer.
- Investigating the causes of carbon stabilization in mineral forest soils and examining the application to further development of the Yasso model.
- Quantifying the effect of uncertainty associated with the spatially interpolated temperature and precipitation data on CSC estimates.

Harvested wood products

- The activity data will be updated for the whole time-series. More specifically, the share of domestic to total HWP will be revised for the calculation of the technical correction.
- Paper and paperboard are not yet included in the current calculations of HWP for the FMRL. Since these products have a half-life of 2 years, the contribution of this pool to total HWP will be small. Currently, the quality of this activity data needs to be improved. See chp. 6.11.6 (UNFCCC 2017/ID#KL.8).

Pools or emissions and removals to be removed from the FMRL:

- CH₄ and N₂O emissions from wildfires were considered for the calculation of the background level and the margin with respect to the application of excluding natural disturbances from the accounting (see UNFCCC2017/ID#KL.2; FOEN 2016d). By moving these emissions to the background level, CH₄ and N₂O emissions from wildfires must be excluded from the contributing pools of the FMRL accordingly (see Table 11-14 in FOEN 2016).

Pools or emissions and removals not included in the FMRL so far:

- Emissions from controlled burning covers the emissions from burning of residues in forestry. In this submission, these emissions are for the first time reported in the category Forest land within the LULUCF sector. Before, these emissions have been reported under 5C2 "incineration and open burning of waste – natural residues". The emissions from controlled burning should be added as a pool to the FMRL (UNFCCC 2017/ID#KL.6).

11.5.2.4 Information related to the natural disturbance provision under Article 3.4

11.5.2.4.1 Application of the provision of natural disturbances

As indicated in Switzerland's Initial Report (FOEN 2016c, FOEN 2016d), Switzerland intends to apply, in the case of significant magnitude events, the provision of natural disturbances for units of lands under Forest management during the second commitment period in accordance with decision 2/CMP.7. In cases or events in which emissions from natural disturbances are higher than the nationally established threshold value and in which all other requirements defined in 2/CMP.7 and IPCC (2014) are met, Switzerland will evaluate and decide whether the effort would be justified to exclude them.

In the inventory years 2013, 2014 and 2015, no natural disturbances causing emissions exceeding the upper confidence interval (background level plus margin) occurred. Thus, no emissions from natural disturbances were excluded for 2013, 2014 and 2015.

11.5.2.4.2 Technical correction of the background level and margin

The background level and margin have been reviewed and are reported in the update of Switzerland's Second Initial Report under the Kyoto Protocol (FOEN 2016d).

There is no technical correction of the background level and margin in this submission.

11.5.2.5 Information on Harvested wood products under Article 3.4

Methodology, estimates and uncertainties of carbon stock changes in the HWP pools are described in chp. 6.11. The same methodology was applied for reporting HWP from Forest land under the UNFCCC and accounting for HWP from Forest management under the Kyoto Protocol. A time series for changes in the HWP pool is shown in Table 6-34 and Figure 6-10. An overview of emissions and removals resulting from the HWP pool is presented in Table 11-1 and Figure 11-1.

11.5.3 Information relating to Cropland management, Grazing Land management, Revegetation and Wetland drainage and rewetting if elected, for the Base Year

Not applicable.

11.5.4 Information that demonstrates that emissions and removals resulting from elected Article 3, paragraph 4 activities are not accounted for under activities under Article 3, paragraph 3

This information is requested in the Annex to 15/CMP.1 paragraph 9(c). The reporting of Forest management under Article 3, paragraph 4 is clearly separated from the reporting of the activities under Article 3, paragraph 3.

Units of lands with ARD (Afforestation, Reforestation and Deforestation) activities, are reported under Article 3, paragraph 3. These areas always remain under Article 3, paragraph 3. Afforestations older than 20 years are accounted for based on emission factors of mature forests under Forest management. These units of lands are reported in CRF table 4(KP-I)A.1

and not under Forest management. Thus, there is no double counting of units of lands under article 3, paragraph 3 to Article 3, paragraph 4.

11.5.5 Information that indicates to what extent removals from Forest management offset the debit incurred under Article 3, Paragraph 3

This information will only be available at the end of the commitment period.

11.6 Other information

11.6.1 Key category analysis for Article 3.3 and 3.4 activities

The results of the Approach 1 key category analysis including LULUCF for the year 2015 are shown in Table 1-4 (by emissions) and summarized in Table 1-9. The method is explained in chp. 1.5. The smallest UNFCCC category, considered key based on an Approach 1 level assessment is "5D Wastewater treatment and discharge, N₂O" with a contribution of 144.38 kt CO₂ eq.

The following LULUCF activities under the Kyoto Protocol are listed in CRF table NIR-3 because their associated LULUCF categories in the UNFCCC inventory are key categories under the level or trend assessment:

- Forest management (-2475.62 kt CO₂) encompasses CO₂ emissions from Forest management excluding HWP, biomass burning and drainage (see Table 11-1) and is a key category under the Kyoto Protocol because its absolute contribution is higher than the smallest category considered key in the UNFCCC inventory. This activity is associated with the UNFCCC category Forest land remaining forest land. Since the total Swiss forest is considered as managed, there is a good agreement between the activity under the Kyoto Protocol and the UNFCCC LU category. According to Table 1-9 the UNFCCC category "Forest land remaining forest land" is both level and trend key category under Approaches 1 and 2 assessments in 2015.
- Afforestation and Reforestation (-16.79 kt CO₂; encompasses CO₂ emissions; CRF table 4(KP)) is not a key category under the Kyoto Protocol because its absolute contribution is substantially lower than the smallest category considered key in the UNFCCC inventory. The associated UNFCCC category Land converted to Forest Land includes converted areas after natural regenerations due to abandonment of land, which are not reported as Afforestation under the Kyoto Protocol. The UNFCCC category Land converted to forest Land is both level and trend key category under Approaches 1 and 2 assessments in 2015 (Table 1-9).
- Deforestation (150.14 kt CO₂; Deforestation excluding HWP encompasses CO₂ emissions; CRF table 4(KP)) is a key category under the Kyoto Protocol because its contribution is higher than the smallest UNFCCC category considered key. The associated UNFCCC category is Land converted to settlements, but only a part of this UNFCCC category represents the activity Deforestation under the Kyoto Protocol (see chp. 11.2.3). The UNFCCC category Land converted to settlements is level key category under Approaches 1 and 2 assessments in 2015 (Table 1-9).
- Harvested wood products (-69.53 kt CO₂; CRF table 4(KP-I)C) is not a level key category under the Approach 1 assessment because its contribution is lower than the smallest UNFCCC category considered key. The same method is used for the calculation of HWP under the UNFCCC and the KP. Using Approach 2, the UNFCCC category HWP is also not a level key category. It is trend key category under Approaches 1 and 2 assessments in 2015 (Table 1-9).

11.7 Information Relating to Article 6

Switzerland does not host Joint Implementation projects.

12 Information on accounting of Kyoto Units

12.1 Background information

The Swiss Emissions Trading Registry completed the go-live process and got fully operational with the International Transaction Log (ITL) on December 4, 2007. As part of the go-live process the entire Assigned Amount of 242'838'402 has been issued as Assigned Amount Units (AAUs) for the first commitment period.

The user interface is located on the Swiss Emissions Trading Registry website (<https://www.emissionsregistry.admin.ch>). Switzerland uses the CR registry software, which has been developed by Lippke & Wagner GmbH. Switzerland cooperates with Monaco regarding registry issues.

The following registry systems' reporting includes the standard electronic format (SEF) tables and the standard independent assessment report (SIAR) tables in accordance with sections E and G of the annex to decision 15/CMP.1.

12.2 Summary of information reported in the SEF tables

The Standard Electronic Format reports for units with applicable commitment period 1 (CP1), and with applicable commitment period 2 (CP2) for 2016, have been submitted to the UNFCCC Secretariat electronically.

Overview of CP1 units

The total balances of CP1 units remained unchanged in the reporting year 2016. By the end of 2016, 247,620,769 AAUs were held in the Swiss Emissions Trading Registry (Table 12-1). In addition, 4,210,465 Emission Reduction Units (ERUs), 25,756,722 Certified Emission Reductions (CERs), and 114,793 Temporary Certified Emission Reductions (tCERs) were held in the Swiss Emissions Trading Registry. A total of 4,796,312 AAUs, 3,651,820 ERUs, 7,896,871 CERs, and all of the 114,793 tCERs have been voluntarily cancelled in the period from 2008 to 2016.

Table 12-1 Total quantities of CP1 Kyoto Protocol units by account type at the end of 2016 (SEF table 4)

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	5'794'523	NO	NO	1'640'531	NO	NO
Entity holding accounts	NO	NO	NO	181'123	NO	NO
Article 3.3/3.4 net source cancellation accounts	172'587	NO	1'013'340	NO		
Non-compliance cancellation account	NO	NO	NO	NO		
Other cancellation accounts	4'796'312	3'651'820	NO	7'896'871	114'793	NO
Retirement account	236'857'347	558'645	8'267'540	16'038'197	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	247'620'769	4'210'465	9'280'880	25'756'722	114'793	NO

Overview of CP2 units

By the end of the reporting year 2016 a total balance of 17,145,594 CERs were held in the Swiss Emissions Trading Registry, of which 854,839 CERs have been voluntarily cancelled (Table 12-2).

Table 12-2 Total quantities of CP2 Kyoto Protocol units by account type at the end of 2016 (SEF table 4)

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	NO	NO	NO	15'297	NO	NO
Entity holding accounts	NO	NO	NO	16'275'458	NO	NO
Retirement account	NO	NO	NO	NO	NO	NO
Previous period surplus reserve account	NO					
Article 3.3/3.4 net source cancellation accounts	NO	NO	NO	NO		
Non-compliance cancellation account	NO	NO	NO	NO		
Voluntary cancellation account	NO	NO	NO	854'839	NO	NO
Cancellation account for remaining units after carry-over	NO	NO	NO	NO	NO	NO
Article 3.1 ter and quarter ambition increase cancellation account	NO					
Article 3.7 ter cancellation account	NO					
tCER cancellation account for expiry					NO	
ICER cancellation account for expiry						NO
ICER cancellation account for reversal of storage						NO
ICER cancellation account for non-submission of certification report						NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	NO	NO	NO	17'145'594	NO	NO

12.3 Discrepancies and notifications

Switzerland's reports on discrepancies (R-2), Clean development mechanism (CDM) notifications (R-3), non-replacements (R-4) including reversal of storage and failure of certification and invalid units (R-5) have been uploaded on the UNFCCC Submission Portal.

During the reported year 2016, the Swiss Emissions Trading Registry had no discrepancies, no CDM notifications, no non-replacements including reversal of storage and failure of certification and no invalid units. Therefore, the SIAR tables R-2, R-3, R-4 and R-5 are empty and no actions and changes have been taken to address discrepancies.

12.4 Publicly accessible information

In accordance to section E of the annex to decision 13/CMP.1 the Swiss Emissions Trading Registry makes non-confidential information available to the public via webpage or user-interface.

Non-confidential information is publicly available on the Swiss Emissions Trading Registry website <https://www.emissionsregistry.admin.ch>. The national allocation plan is accessible under 'Allocation' in the Public Information menu. The report 'Accounts' provides a list of open accounts in the national registry. The 'Surrendering Obligation', and 'Surrendered units' per operator are also publicly accessible.

Data of transfers and holdings of individual accounts are considered as business secrets and the disclosure may prejudice their competitiveness. Information on acquiring and transferring units of companies (as legal persons) is therefore regarded as personal data. Article 19 of the Federal Act on Data Protection (FADP, SR 235.1 Bundesgesetz vom 19. Juni 1992 über den Datenschutz (DSG) 2) enacts that federal bodies may disclose personal data if there is a

legal basis for doing so or if there is an overriding public interest. In the present case these conditions are not fulfilled. Therefore, the registry of Switzerland cannot make the information on acquiring and transferring accounts publicly available and considers them as confidential. The Representative identifier (13/CMP.1 Annex paragraph 45 (d)), as well as all information according to 13/CMP.1 Annex paragraph 45 (e) are also considered as confidential. Therefore, this information is not publicly available. A statement on which information is considered as confidential can be found on the public website <https://www.emissionsregistry.admin.ch>.

All other information referred to in paragraphs 44 to 48 to the annex to decision 13/CMP.1 are made publicly available by the Swiss Emissions Trading Registry, if they are not covered by the above mentioned articles.

Information related to Article 6 projects is publicly accessible on the website <http://www.bafu.admin.ch/ji-e>. Switzerland does not host Joint Implementation (JI)-projects and therefore no issuance of ERUs has taken place.

12.5 Calculation of the Commitment Period Reserve (CPR)

The commitment period reserve and the assigned amount for the second commitment period is defined in the *Report to facilitate the calculation of the assigned amount pursuant to Article 3, paragraphs 7bis, 8 and 8bis, of the Kyoto Protocol for the second commitment period 2013–2020 (Switzerland's Initial Report under the Kyoto Protocol, 2nd CP)* (FOEN 2016c), and the update to the report following the review (FOEN 2016d). Switzerland's assigned amount for the second commitment period is 361'768.527 kt CO₂ equivalent. The commitment period reserve is 325'591.674 kt CO₂ equivalent.

12.6 KP-LULUCF Accounting

Switzerland chose to account over the entire commitment period for emissions and removals from activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol.

13 Information on changes in National Registry

Table 13-1 Changes in the national registry in accordance with §32 decision 15/CMP.1

Annual Submission Item	Reporting
15/CMP.1 annex II.E paragraph 32.(a): Change of name or contact	No change in the name or contact information of the registry administrator occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(b): Change of cooperation arrangement	No change of cooperation arrangement occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(c): Change of the database or the capacity of National Registry	No change to the database or to the capacity of the national registry occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(d): Change of conformance to technical standards	No change in the registry's conformance to technical standards occurred for the reported period.
15/CMP.1 annex II.E paragraph 32.(e): Change of discrepancies procedures	No change of discrepancies procedures occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(f): Change of Security	No change of security measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(g): Change of list of publicly available information	No change to the list of publicly available information occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(h): Change of Internet address	No change of the registry Internet address occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(i): Change of data integrity measures	No change of data integrity measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(j): Change of test results	No change of test results occurred during the reporting period.

14 Information on minimization of adverse impacts in accordance with Article 3, Paragraph 14

The Convention (Art. 4 §8 and §10) and its Kyoto Protocol (Art. 2 §3 and Art. 3 §14) commit Parties to strive to implement climate policies and measures in such a way as to minimize adverse economic, social and environmental impacts on developing countries when responding to climate change.

Context

Switzerland strives to design climate change policies and measures in a way as to ensure a balanced distribution of mitigation efforts by implementing climate change response measures in all sectors and for different gases. Indirectly, this approach is deemed to minimize also the scope of potential adverse impacts on concerned actors (including developing countries). Though, due to Switzerland's size and share related to international trade – mainly concentrated on the EU – and greenhouse gas emissions, it is not assumed that Swiss climate change policies have any significant adverse economic, social and environmental impacts in developing countries. Additionally, the policies and measures are very much compatible and consistent with those of the European Union in order to avoid trade distortion, non-tariff barriers to trade and to set similar incentives. All major projects of law in Switzerland are accompanied by impact assessments, inter alia including evaluation of trade-related issues. In accordance with international law, this approach strives at ensuring that Switzerland is implementing those climate change response measures, which are least trade distortive and do not create unnecessary barriers to trade. Consistently, Switzerland notifies all proposed non-tariff measures having a potential impact on trade to the WTO, where specific concerns can be raised by other parties. Moreover, Switzerland belongs to the most important donors in the area of Aid for Trade.

The impact assessment is accompanied by a broad internal and external consultation process, inter alia inviting competent actors to provide advice on international economic, social and environmental aspects of proposed policies and measures. The open public consultation process, together with regular policy dialogues with other countries guarantee that all domestic and foreign stakeholders can raise concerns and issues about new policy initiatives, i.e. including those concerns about possible adverse impacts on other countries.

Progressive reduction or phasing out of market imperfections, fiscal incentives, tax and duty exemptions and subsidies in all greenhouse-gas-emitting sectors, taking into account the need for energy price reforms to reflect market prices and externalities

Environmental policy in Switzerland, including climate change policies, are guided by the "polluter pays" principles, as enshrined in the Federal Law on the Protection of the Environment. Accordingly, the internalization of external costs and adequate price signals are key aspects of Switzerland's climate change policy. Regarding greenhouse gas emissions, market-based instruments, such as the Swiss Emissions Trading Scheme, the supplemental use of Certified Emission Reductions from the Clean Development Mechanism or the levy for heating and process fuels are important measures to put a price on emissions of greenhouse

gases (see Sixth National Communication for more details), that are then reflected in market prices and thus internalizing externalities.

Fiscal incentives, tax and duty exemptions and subsidies

Price-based measures are recognized as essential instruments for promoting the efficient use of resources and to reduce market imperfections. In 2001 Switzerland introduced a heavy vehicle fee (HVF). It is applied to passenger and freight transport vehicles of more than 3.5 tonnes gross weight. The impact of the HVF introduction was most clearly reflected by changes in traffic volume (truck-kilometres) but also in reduced air pollution, a renewal of the heavy vehicle fleet and an increase of load per vehicle, fewer trucks having transported more goods. Two thirds of the revenues are used to finance major railway infrastructure projects (such as the two base tunnels through the Alps), and one third is transferred to the cantons.

In 2008 Switzerland introduced a CO₂ levy on heating and process fuel to set an incentive for a more efficient use of fossil fuels, promote investment in energy-efficient technologies and the use of low-carbon or carbon-free energy sources. The 2013 amendment to the CO₂ act (Swiss Confederation 2012) still encompasses the imposition of a CO₂ levy on heating and process fuel. Companies, especially those industries with substantial CO₂ emissions from use of heating fuels, may apply for exemption from the CO₂ levy, provided the company commits to emission reductions. The company has to elaborate an emission reduction target, based on the technological potential and economic viability of various measures within the company. While the proceeds from the CO₂ levy were initially to be fully and equally refunded to the Swiss population and to the business community in proportion of wages paid, a parliamentary decision of June 2009 earmarked a third of the revenues from the CO₂ levy to CO₂ relevant measures in the building sector (Building refurbishment programme). The partial earmarking of revenues from the CO₂ tax is limited in the revised CO₂ act to a maximum of 300 million Swiss francs per year.

The economic impact of the Swiss climate policy was analysed in two studies¹². The impact is considered to be very small.

Switzerland does not subsidize fossil fuels in general. There are some minor schemes in place though that may be regarded as fossil fuel subsidies. In international comparison, however, these schemes are limited: At the federal level, a few tax exemptions and reductions provide some form of support to users of fossil fuels. Farmers, foresters, fishermen and the fuel use of snow cats are exempt from the mineral oil tax that is normally levied on sales of mineral oils, while public transport companies benefit from a reduced rate. Some vehicles are also exempt from the performance-related Heavy Vehicle Fee (HFV), e.g. agricultural vehicles, vehicles used for the concessionary transport of persons or vehicles for police, fire brigade, oil and chemical emergency unit, civil protection and ambulances.

¹² Ecoplan (2009): Volkswirtschaftliche Auswirkungen der Schweizer Post-Kyoto-Politik, im Auftrag des BAFU. BAFU (2010): Synthesebericht zur Volkswirtschaftlichen Beurteilung der Schweizer Klimapolitik nach 2012.

The need for energy prices reforms

World-wide subsidies for fossil fuels are estimated at 300–500 billion USD per annum, depending on the level of energy prices. This huge market distortion does not only produce severe fiscal problems for the countries concerned, it is also a major obstacle for enhanced investments in energy efficiency measures and renewable energies.

Switzerland as a member of the Friends of Fossil Fuels Subsidies Reform group supports the gradual and sustained reduction of unnecessary market-distortions. Switzerland under its Economic Development Cooperation supports partner countries in the design and implementation of energy tariff reforms, as an element of infrastructure financing programs. Switzerland has been an initiator of specialized international programs, including the World Bank's Energy Sector Management Program ESMAP. The Energy Efficiency Governance Handbook has been produced with Swiss financing (IEA/EBRD 2010).

Removing subsidies associated with the use of environmentally unsound and unsafe technologies

Switzerland doesn't subsidize the use of environmentally unsound and unsafe technologies.

Strengthening the capacity of developing country Parties for improving efficiency in upstream and downstream activities relating to fossil fuels, taking into consideration the need to improve the environmental efficiency of these activities

Switzerland supports through different projects the enhancement of efficiency in industrial production, i.e. "cleaner production". These cleaner production projects promote eco-efficient means of production and better working conditions attained through technical improvements and behavioural changes in both management and staff in industrial companies and services. The resulting rise of economic and environmental efficiency and improved competitiveness is gained through the systematic optimisation of energy use, processing of raw material, more efficient use of resources and thus better protection of the environment.

Furthermore, there is a rising awareness and demand by consumers for environmentally sound products. In order to alleviate potential adverse economic impacts of corresponding national measures Switzerland promotes and supports the development of international standards, especially with regard to the sustainable use of natural resources (including agricultural commodities), e.g. through the creation of sustainability standards, financial incentives and favourable framework conditions in developing countries by consultancy services and technology transfer. Further information is contained in chp. 7 of Switzerland's Sixth National Communication (FOEN 2014d).

Assisting developing country Parties which are highly dependent on the export and consumption of fossil fuels in diversifying their economies

Most developing and transition countries have, in recent years, taken important steps towards trade liberalisation, in order to align their trade policies with multilateral trade agreements. The Swiss State Secretariat for Economic Affairs (SECO) supports these efforts, because a multilaterally acknowledged and respected set of regulations for

international transactions not only strengthens trade as such, but also creates more potent and legally secure markets to the benefit of all players.

The measures taken by SECO are aimed at creating the necessary conditions for earning additional income in the beneficiary countries and thereby contribute directly to the alleviation of poverty. SECO is focusing on three areas of intervention along the value chain: (i) International competitiveness (ii) Enabling framework conditions for trade (iii) Improving market access.

For example market access: Trade between developing and industrial countries is still insufficiently developed respectively not diversified enough. On one hand, the developing countries lack the necessary production capacities, transport infrastructure and know-how; on the other hand, tariff and non-tariff barriers to trade make direct access to markets more difficult.

Switzerland promotes access to Swiss markets by granting preferential tariffs on products from developing and emerging countries. In addition, SECO runs programmes for promoting imports to Switzerland and the rest of Europe. The easing of market entry for products from disadvantaged countries is an important contribution to the promotion and diversification of trade, the increase of export revenues and thus to the economic development of the partner countries. Switzerland supports developing and transition countries in the following areas:

- Generalized system of preferences (GSP)
- Swiss Import Promotion Program (www.sippo.ch)
- Development of new private voluntary social and environmental standards based on international multi-stakeholder approaches: private sustainability standards Better Cotton, 4C (Common Code for the Coffee Community), Roundtable for Sustainable Biofuels, etc.

Finally, Switzerland is a strong supporter of the EITI (Extractive Industries Transparency Initiative). We share a belief that the rational use of natural resource wealth is an important driving force for sustainable economic growth that contributes to sustainable development and poverty reduction. The sustainable management of natural resource wealth – as supported by EITI principle and criteria incl. regular publication and audit of revenues – is key to mobilize the funds for diversification strategies.

Changes compared to the latest submission

There are no changes compared to the 2016 submission.

Annexes

Annex 1: Key category analysis (KCA)

Table A – 1 Overview over the total of 167 categories included in the KCA (level of disaggregation)

No.	NFR	IPCC source and sink categories			Fuel	Gas
1	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Biomass	CH4
2	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Gaseous Fuels	CH4
3	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Liquid Fuels	CH4
4	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Solid Fuels	CH4
5	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Gaseous Fuels	CO2
6	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Liquid Fuels	CO2
7	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Other Fuels	CO2
8	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Solid Fuels	CO2
9	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Biomass	N2O
10	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Gaseous Fuels	N2O
11	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Liquid Fuels	N2O
12	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Other Fuels	N2O
13	1A1	1. Energy	A. Fuel combustion activities	1. Energy industries	Solid Fuels	N2O
14	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Biomass	CH4
15	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Gaseous Fuels	CH4
16	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Liquid Fuels	CH4
17	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Other Fuels	CH4
18	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Solid Fuels	CH4
19	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Gaseous Fuels	CO2
20	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Liquid Fuels	CO2
21	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Other Fuels	CO2
22	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Solid Fuels	CO2
23	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Biomass	N2O
24	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Gaseous Fuels	N2O
25	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Liquid Fuels	N2O
26	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Other Fuels	N2O
27	1A2	1. Energy	A. Fuel combustion activities	2. Manufacturing industries and construction	Solid Fuels	N2O
28	1A3a	1. Energy	A. Fuel combustion activities	3. Transport; Domestic aviation	kerosene	CH4
29	1A3a	1. Energy	A. Fuel combustion activities	3. Transport; Domestic aviation	kerosene	CO2
30	1A3a	1. Energy	A. Fuel combustion activities	3. Transport; Domestic aviation	kerosene	N2O
31	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Biomass	CH4
32	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Diesel	CH4
33	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gaseous Fuels	CH4
34	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gasoline	CH4
35	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Diesel	CO2
36	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gaseous Fuels	CO2
37	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gasoline	CO2
38	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Biomass	N2O
39	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Diesel	N2O
40	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gaseous Fuels	N2O
41	1A3b	1. Energy	A. Fuel combustion activities	3. Transport; Road transportation	Gasoline	N2O
42	1A3c	1. Energy	A. Fuel combustion activities	3. Transport; Railways	Biomass	CH4
43	1A3c	1. Energy	A. Fuel combustion activities	3. Transport; Railways	Diesel	CH4
44	1A3c	1. Energy	A. Fuel combustion activities	3. Transport; Railways	Diesel	CO2
45	1A3c	1. Energy	A. Fuel combustion activities	3. Transport; Railways	Biomass	N2O
46	1A3c	1. Energy	A. Fuel combustion activities	3. Transport; Railways	Diesel	N2O
47	1A3d	1. Energy	A. Fuel combustion activities	3. Transport; Domestic navigation	Biomass	CH4
48	1A3d	1. Energy	A. Fuel combustion activities	3. Transport; Domestic navigation	Liquid Fuels	CH4
49	1A3d	1. Energy	A. Fuel combustion activities	3. Transport; Domestic navigation	Liquid Fuels	CO2
50	1A3d	1. Energy	A. Fuel combustion activities	3. Transport; Domestic navigation	Biomass	N2O
51	1A3d	1. Energy	A. Fuel combustion activities	3. Transport; Domestic navigation	Liquid Fuels	N2O
52	1A3e	1. Energy	A. Fuel combustion activities	3. Transport; Other transportation	Gaseous Fuels	CH4
53	1A3e	1. Energy	A. Fuel combustion activities	3. Transport; Other transportation	Gaseous Fuels	CO2
54	1A3e	1. Energy	A. Fuel combustion activities	3. Transport; Other transportation	Gaseous Fuels	N2O
55	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Biomass	CH4
56	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Gaseous Fuels	CH4
57	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Liquid Fuels	CH4
58	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Gaseous Fuels	CO2
59	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Liquid Fuels	CO2
60	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Biomass	N2O
61	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Gaseous Fuels	N2O
62	1A4a	1. Energy	A. Fuel combustion activities	4. Other sectors; Commercial/institutional	Liquid Fuels	N2O
63	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Biomass	CH4
64	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Gaseous Fuels	CH4
65	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Liquid Fuels	CH4
66	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Solid Fuels	CH4
67	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Gaseous Fuels	CO2
68	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Liquid Fuels	CO2
69	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Solid Fuels	CO2
70	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Biomass	N2O
71	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Gaseous Fuels	N2O
72	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Liquid Fuels	N2O
73	1A4b	1. Energy	A. Fuel combustion activities	4. Other sectors; Residential	Solid Fuels	N2O
74	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Biomass	CH4
75	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Gaseous Fuels	CH4
76	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Liquid Fuels	CH4
77	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Gaseous Fuels	CO2
78	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Liquid Fuels	CO2
79	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Biomass	N2O
80	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Gaseous Fuels	N2O
81	1A4c	1. Energy	A. Fuel combustion activities	4. Other sectors; Agriculture/forestry/fishing	Liquid Fuels	N2O

No.	NFR	IPCC source and sink categories			Fuel	Gas
82	1A5	1. Energy	A. Fuel combustion activities	5. Other	Biomass	CH4
83	1A5	1. Energy	A. Fuel combustion activities	5. Other	Liquid Fuels	CH4
84	1A5	1. Energy	A. Fuel combustion activities	5. Other	Liquid Fuels	CO2
85	1A5	1. Energy	A. Fuel combustion activities	5. Other	Biomass	N2O
86	1A5	1. Energy	A. Fuel combustion activities	5. Other	Liquid Fuels	N2O
87	1B2	1. Energy	B. Fugitive emissions from fuels	2. Oil & nat. gas & other em. from energy prod.		CH4
88	1B2	1. Energy	B. Fugitive emissions from fuels	2. Oil & nat. gas & other em. from energy prod.		CO2
89	1B2	1. Energy	B. Fugitive emissions from fuels	2. Oil & nat. gas & other em. from energy prod.		N2O
90	1ind	1. Energy	Indirect emissions			CO2
91	2A1	2. IPPU	A. Mineral industry	1. Cement production		CO2
92	2A2	2. IPPU	A. Mineral industry	2. Lime production		CO2
93	2A3	2. IPPU	A. Mineral industry	3. Glass production		CO2
94	2A4	2. IPPU	A. Mineral industry	4. Other process uses of carbonates		CO2
95	2B5	2. IPPU	B. Chemical industry	5. Carbide production		CH4
96	2B5	2. IPPU	B. Chemical industry	5. Carbide production		CO2
97	2B8	2. IPPU	B. Chemical industry	8. Petrochemical and carbon black production		CO2
98	2B10	2. IPPU	B. Chemical industry	10. Other (Niacin, Limestone pit)		CO2
99	2B2	2. IPPU	B. Chemical industry	2. Nitric acid production		N2O
100	2C1	2. IPPU	C. Metal industry	1. Iron and steel production		CO2
101	2C3	2. IPPU	C. Metal industry	3. Aluminium production		CO2
102	2C3	2. IPPU	C. Metal industry	3. Aluminium production		PFC
103	2C4	2. IPPU	C. Metal industry	4. Magnesium production		SF6
104	2C7	2. IPPU	C. Metal industry	7. Other		CO2
105	2D	2. IPPU	D. Non-energy products from fuels and solvent use			CO2
106	2E1	2. IPPU	E. Electronics industry	1. Integrated circuit or semiconductor		HFC
107	2E1	2. IPPU	E. Electronics industry	1. Integrated circuit or semiconductor		PFC
108	2E1	2. IPPU	E. Electronics industry	1. Integrated circuit or semiconductor		SF6
109	2E3	2. IPPU	E. Electronics industry	3. Photovoltaics		NF3
110	2E4	2. IPPU	F. Product uses as subst. for ODS	4. Heat transfer fluid		PFC
111	2F1	2. IPPU	F. Product uses as subst. for ODS	1. Refrigeration and air conditioning		HFC
112	2F1	2. IPPU	F. Product uses as subst. for ODS	1. Refrigeration and air conditioning		PFC
113	2F2	2. IPPU	F. Product uses as subst. for ODS	2. Foam blowing agents		HFC
114	2F4	2. IPPU	F. Product uses as subst. for ODS	4. Aerosols		HFC
115	2F5	2. IPPU	F. Product uses as subst. for ODS	5. Solvents		HFC
116	2G	2. IPPU	G. Other product manufacture and use			CO2
117	2G	2. IPPU	G. Other product manufacture and use			HFC
118	2G	2. IPPU	G. Other product manufacture and use			PFC
119	2G	2. IPPU	G. Other product manufacture and use			SF6
120	2G	2. IPPU	G. Other product manufacture and use			N2O
121	2H	2. IPPU	H. Other			CO2
122	2ind	2. IPPU	Indirect emissions			CO2
123	3A	Agriculture	A. Enteric Fermentation			CH4
124	3B	Agriculture	B. Manure Management			CH4
125	3B	Agriculture	B. Manure Management	liquid		N2O
126	3B	Agriculture	B. Manure Management	solid		N2O
127	3B5	Agriculture	B. Manure Management	indirect		N2O
128	3Da1/2/4/5/7	Agriculture	D. Agricultural Soils; Direct Soil Em.	fertilizer		N2O
129	3Da6	Agriculture	D. Agricultural Soils; Direct Soil Em.	organic soils		N2O
130	3Da3	Agriculture	D. Agricultural Soils; Pasture, Range and Paddock Manure			N2O
131	3Db1	Agriculture	D. Agricultural Soils; Indirect Em.	deposition		N2O
132	3Db2	Agriculture	D. Agricultural Soils; Indirect Em.	leaching and runoff		N2O
133	3G	Agriculture	G. Limestone			CO2
134	3H	Agriculture	H. Urea application			CO2
135	4 II	4. LULUCF	Drainage, rewetting and other management of organic and mineral soils			CH4
136	4 II	4. LULUCF	Drainage, rewetting and other management of organic and mineral soils			N2O
137	4 III	4. LULUCF	N mineralization			N2O
138	4 IV	4. LULUCF	Indirect emissions			N2O
139	4 V	4. LULUCF	Biomass burning			CH4
140	4 V	4. LULUCF	Biomass burning			CO2
141	4 V	4. LULUCF	Biomass burning			N2O
142	4A1	4. LULUCF	A. Forest land	1. Forest land remaining forest land		CO2
143	4A2	4. LULUCF	A. Forest land	2. Land converted to forest land		CO2
144	4B1	4. LULUCF	B. Cropland	1. Cropland remaining cropland		CO2
145	4B2	4. LULUCF	B. Cropland	2. Land converted to cropland		CO2
146	4C1	4. LULUCF	C. Grassland	1. Grassland remaining grassland		CO2
147	4C2	4. LULUCF	C. Grassland	2. Land converted to grassland		CO2
148	4D1	4. LULUCF	D. Wetlands	1. Wetlands remaining wetlands		CO2
149	4D2	4. LULUCF	D. Wetlands	2. Land converted to wetlands		CO2
150	4E1	4. LULUCF	E. Settlements	1. Settlements remaining settlements		CO2
151	4E2	4. LULUCF	E. Settlements	2. Land converted to settlements		CO2
152	4F2	4. LULUCF	F. Other land	2. Land converted to other land		CO2
153	4G	4. LULUCF	G. Harvested wood products			CO2
154	5A	5. Waste	A. Solid waste disposal			CH4
155	5A	5. Waste	A. Solid waste disposal			CO2
156	5B	5. Waste	B. Biological treatment of solid waste			CH4
157	5B	5. Waste	B. Biological treatment of solid waste			N2O
158	5C	5. Waste	C. Incineration and open burning of waste			CH4
159	5C	5. Waste	C. Incineration and open burning of waste			CO2
160	5C	5. Waste	C. Incineration and open burning of waste			N2O
161	5D	5. Waste	D. Wastewater treatment and discharge			CH4
162	5D	5. Waste	D. Wastewater treatment and discharge			N2O
163	5ind	5. Waste	Indirect emissions			CO2
164	6	6. Other				CH4
165	6	6. Other				CO2
166	6	6. Other				N2O
167	6ind	6. Other	Indirect emissions			CO2

Annex 2: Assessment of uncertainty

A2.1 Detailed results of Approach 1 uncertainty analysis

The table on the next pages shows the detailed results of Approach 1 uncertainty analysis. The structure of the table is identical to Table 3.2 of the 2006 IPCC Guidelines (IPCC 2006). For explanations to the columns see pp. 3.30–3.31 in vol. 1 of IPCC (2006).

Table A – 2 Results of Approach 1 uncertainty analysis, overall results are presented at the bottom of the last part of the table (Table 3.2 of IPCC 2006 Guidelines). The uncertainty analysis includes indirect CO₂ emissions.

IPCC Source category	Gas	Base year emissions or removals	Year 2015 emissions or removals	AD uncertainty	EF uncertainty	Combined uncertainty	Contribution to variance by Category in 2015	Type A sensitivity	Type B sensitivity	Uncertainty in trend in nat. emissions introduced by EF uncertainty	Uncertainty in trend in nat. emissions introduced by AD uncertainty	Uncertainty introduced into the trend in total national emissions	
		kt CO ₂ eq	kt CO ₂ eq	%	%	%	-	%	%	%	%	-	
1A1	1. Energy industries	Biomass	0.45	0.28	10.0	28.3	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Gaseous F.	0.11	0.18	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Liquid F.	0.52	0.27	0.7	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Solid F.	0.13	-	5.0	29.6	30.0	-	0.0000	-	0.00	-	0.00
		Gaseous F.	243.40	388.76	5.0	1.0	5.1	0.002	0.0034	0.0075	0.00	0.05	0.00
		Liquid F.	685.81	470.76	0.7	0.1	0.7	0.000	0.0025	0.0088	0.00	0.01	0.00
		Other F.	1'491.55	2'381.90	5.0	9.2	10.5	0.279	0.0199	0.0445	0.18	0.31	0.13
		Solid F.	49.13	-	5.0	5.1	7.1	-	0.0008	-	0.0008	-	0.00
		Biomass	22.58	13.67	10.0	79.4	80.0	0.001	0.0001	0.0003	0.01	0.00	0.00
		Gaseous F.	0.13	0.21	5.0	79.8	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Liquid F.	1.11	0.45	0.7	80.0	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Other F.	24.33	12.62	5.0	79.8	80.0	0.000	0.0002	0.0002	0.01	0.00	0.00
		Solid F.	0.24	-	5.0	79.8	80.0	-	0.0000	-	0.00	-	0.00
		Biomass	4.28	1.86	10.0	28.3	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
Gaseous F.	0.49	0.98	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Liquid F.	4.48	1.40	0.7	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Other F.	0.77	0.53	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Solid F.	0.31	0.22	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Gaseous F.	1'059.63	2'220.03	5.0	1.0	5.1	0.058	0.0240	0.0415	0.03	0.29	0.09		
Liquid F.	3'889.15	1'888.12	0.7	0.1	0.7	0.001	0.0289	0.0353	0.00	0.03	0.00		
Other F.	192.29	367.22	5.0	9.2	10.5	0.007	0.0037	0.0069	0.03	0.05	0.00		
Solid F.	1'274.70	474.46	5.0	5.1	7.1	0.005	0.0122	0.0089	0.06	0.06	0.01		
Biomass	5.14	13.76	10.0	79.4	80.0	0.001	0.0002	0.0003	0.01	0.00	0.00		
Gaseous F.	0.68	1.33	5.0	79.8	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Liquid F.	13.03	10.85	0.7	80.0	80.0	0.000	0.0000	0.0002	0.00	0.00	0.00		
Other F.	2.32	5.58	5.0	79.8	80.0	0.000	0.0001	0.0001	0.01	0.00	0.00		
Solid F.	6.14	2.24	5.0	79.8	80.0	0.000	0.0001	0.0001	0.00	0.00	0.00		
kerosene	0.17	0.13	1.0	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
kerosene	252.55	137.37	1.0	0.2	1.0	0.000	0.0016	0.0026	0.00	0.00	0.00		
kerosene	2.06	1.12	1.0	150.0	150.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
1A2	2. Manufacturing industries and construction	Biomass	0.45	0.28	10.0	28.3	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Gaseous F.	0.11	0.18	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Liquid F.	0.52	0.27	0.7	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Solid F.	0.13	-	5.0	29.6	30.0	-	0.0000	-	0.00	-	0.00
		Gaseous F.	243.40	388.76	5.0	1.0	5.1	0.002	0.0034	0.0075	0.00	0.05	0.00
		Liquid F.	685.81	470.76	0.7	0.1	0.7	0.000	0.0025	0.0088	0.00	0.01	0.00
		Other F.	1'491.55	2'381.90	5.0	9.2	10.5	0.279	0.0199	0.0445	0.18	0.31	0.13
		Solid F.	49.13	-	5.0	5.1	7.1	-	0.0008	-	0.0008	-	0.00
		Biomass	22.58	13.67	10.0	79.4	80.0	0.001	0.0001	0.0003	0.01	0.00	0.00
		Gaseous F.	0.13	0.21	5.0	79.8	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Liquid F.	1.11	0.45	0.7	80.0	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Other F.	24.33	12.62	5.0	79.8	80.0	0.000	0.0002	0.0002	0.01	0.00	0.00
		Solid F.	0.24	-	5.0	79.8	80.0	-	0.0000	-	0.00	-	0.00
		Biomass	4.28	1.86	10.0	28.3	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
Gaseous F.	0.49	0.98	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Liquid F.	4.48	1.40	0.7	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Other F.	0.77	0.53	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Solid F.	0.31	0.22	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Gaseous F.	1'059.63	2'220.03	5.0	1.0	5.1	0.058	0.0240	0.0415	0.03	0.29	0.09		
Liquid F.	3'889.15	1'888.12	0.7	0.1	0.7	0.001	0.0289	0.0353	0.00	0.03	0.00		
Other F.	192.29	367.22	5.0	9.2	10.5	0.007	0.0037	0.0069	0.03	0.05	0.00		
Solid F.	1'274.70	474.46	5.0	5.1	7.1	0.005	0.0122	0.0089	0.06	0.06	0.01		
Biomass	5.14	13.76	10.0	79.4	80.0	0.001	0.0002	0.0003	0.01	0.00	0.00		
Gaseous F.	0.68	1.33	5.0	79.8	80.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
Liquid F.	13.03	10.85	0.7	80.0	80.0	0.000	0.0000	0.0002	0.00	0.00	0.00		
Other F.	2.32	5.58	5.0	79.8	80.0	0.000	0.0001	0.0001	0.01	0.00	0.00		
Solid F.	6.14	2.24	5.0	79.8	80.0	0.000	0.0001	0.0001	0.00	0.00	0.00		
kerosene	0.17	0.13	1.0	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
kerosene	252.55	137.37	1.0	0.2	1.0	0.000	0.0016	0.0026	0.00	0.00	0.00		
kerosene	2.06	1.12	1.0	150.0	150.0	0.000	0.0000	0.0000	0.00	0.00	0.00		
1A3a	3. Transport: Domestic aviation	Biomass	0.45	0.28	10.0	28.3	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Gaseous F.	0.11	0.18	5.0	29.6	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00
		Liquid F.	0.52	0.27	0.7	30.0	30.0	0.000	0.0000	0.0000	0.00	0.00	0.00

IPCC Source category	Gas	Base year emissions or removals	Year 2015 emissions or removals	AD uncertainty	EF uncertainty	Combined uncertainty	Contribution to variance by Category in 2015	Type A sensitivity	Type B sensitivity	Uncertainty in trend in nat. emissions introduced by EF uncertainty	Uncertainty in trend in nat. emissions introduced by AD uncertainty	Uncertainty introduced into the trend in total national emissions
1A3b	3. Transport; Road transportation	Biomass	-	0.12	10.0	59.2	60.0	0.000	0.0000	0.00	0.00	0.00
		Diesel	1.67	0.60	0.9	20.0	20.0	0.000	0.0000	0.00	0.00	0.00
		Gaseous F.	-	0.04	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00
		Gasoline	116.04	18.00	0.7	37.0	37.0	0.000	0.0016	0.06	0.00	0.00
		Diesel	2'632.59	7'181.99	0.9	0.1	0.9	0.018	0.0907	0.1343	0.01	0.17
		Gaseous F.	-	35.53	5.0	1.0	5.1	0.000	0.0007	0.00	0.00	0.00
		Gasoline	11'334.49	7'669.15	0.7	0.1	0.7	0.013	0.0437	0.1434	0.01	0.14
		Biomass	-	1.93	10.0	149.7	150.0	0.000	0.0000	0.01	0.00	0.00
		Diesel	5.63	79.19	0.9	22.0	22.0	0.001	0.0014	0.0015	0.03	0.00
		Gaseous F.	-	0.74	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00
		Gasoline	136.88	25.09	0.7	50.0	50.0	0.001	0.0018	0.0005	0.00	0.00
		Biomass	-	0.00	10.0	59.2	60.0	0.000	0.0000	0.00	0.00	0.00
Diesel	0.03	0.01	0.9	30.0	30.0	0.000	0.0000	0.00	0.00	0.00		
Diesel	28.69	28.72	0.9	0.1	0.1	0.000	0.0001	0.0005	0.00	0.00		
Biomass	-	0.00	10.0	149.7	150.0	0.000	0.0000	0.00	0.00	0.00		
Diesel	0.43	0.42	0.7	80.0	80.0	0.000	0.0000	0.00	0.00	0.00		
Biomass	-	0.00	10.0	59.2	60.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	1.68	0.29	0.7	30.0	30.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	114.27	113.28	0.7	0.1	0.1	0.000	0.0002	0.0021	0.00	0.00		
Biomass	-	0.01	10.0	149.7	150.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	1.16	1.23	0.7	150.0	150.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	0.07	0.04	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00		
Gaseous F.	31.42	42.86	5.0	1.0	5.1	0.000	0.0003	0.0008	0.00	0.01		
Gaseous F.	0.02	0.02	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00		
Biomass	9.22	4.63	10.0	28.3	30.0	0.000	0.0001	0.0001	0.00	0.00		
Gaseous F.	0.70	1.30	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	16.34	9.66	0.7	30.0	30.0	0.000	0.0001	0.0002	0.00	0.00		
Gaseous F.	981.49	1'400.80	5.0	1.0	5.1	0.023	0.0100	0.0262	0.01	0.19		
Liquid F.	4'260.86	2'701.66	0.7	0.1	0.7	0.002	0.0199	0.0505	0.00	0.05		
Biomass	3.50	9.45	10.0	79.4	80.0	0.000	0.0001	0.0002	0.01	0.00		
Gaseous F.	0.52	0.74	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	10.35	6.59	0.7	80.0	80.0	0.000	0.0000	0.00	0.00	0.00		
Biomass	109.97	25.57	10.0	28.3	30.0	0.000	0.0000	0.00	0.00	0.00		
Gaseous F.	0.70	1.36	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	34.83	20.04	0.7	30.0	30.0	0.000	0.0002	0.0004	0.01	0.00		
Solid F.	4.73	3.00	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00		
Gaseous F.	1'449.69	2'600.43	5.0	1.0	5.1	0.079	0.0247	0.0486	0.03	0.34		
Liquid F.	10'099.07	5'863.66	0.7	0.1	0.7	0.007	0.0571	0.1096	0.00	0.11		
Solid F.	58.40	37.08	5.0	5.1	7.1	0.000	0.0003	0.0007	0.00	0.00		
Biomass	25.86	21.37	10.0	79.4	80.0	0.001	0.0000	0.0004	0.00	0.01		
Gaseous F.	24.53	1.37	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00		
Liquid F.	0.28	0.18	5.0	80.0	80.0	0.001	0.0001	0.0003	0.01	0.00		
Solid F.	-	-	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00		

IPCC Source category	Gas	Base year emissions or removals kt CO2 eq	Year 2015 emissions or removals kt CO2 eq	AD uncertainty %	EF uncertainty %	Combined uncertainty %	Contribution to variance by Category in 2015	Type A sensitivity %	Type B sensitivity %	Uncertainty in trend in nat. emissions introduced by EF uncertainty %	Uncertainty in trend in nat. emissions introduced by AD uncertainty %	Uncertainty introduced into the trend in total national emissions	
1. Energy	A. Fuel combustion activities	Biomass	1.97	0.35	10.0	28.3	30.0	0.000	0.0000	0.00	0.00	0.00	
		Gaseous F.	0.02	0.01	5.0	29.6	30.0	0.000	0.0000	0.00	0.00	0.00	
		Liquid F.	6.45	1.43	0.7	30.0	30.0	0.000	0.0001	0.00	0.00	0.00	
		4. Other sectors; Agriculture / forestry / fishing	41.45	13.12	5.0	1.0	5.1	0.000	0.0004	0.00	0.00	0.00	
		Liquid F.	485.09	407.17	0.7	0.1	0.7	0.000	0.0004	0.00	0.01	0.00	
		Biomass	0.51	1.01	10.0	79.4	80.0	0.000	0.0000	0.00	0.00	0.00	
		Gaseous F.	0.02	0.01	5.0	79.8	80.0	0.000	0.0000	0.00	0.00	0.00	
		Liquid F.	4.20	4.60	0.7	80.0	80.0	0.000	0.0001	0.00	0.00	0.00	
		Biomass	-	0.00	10.0	59.2	60.0	0.000	0.0000	0.00	0.00	0.00	
		Liquid F.	0.27	0.16	0.7	60.0	60.0	0.000	0.0000	0.00	0.00	0.00	
2. Industrial processes and product use	B. Fugitive emissions from fuels	Liquid F.	217.65	134.04	0.7	0.1	0.7	0.000	0.0011	0.00	0.00	0.00	
		Biomass	-	0.00	106.1	106.1	150.0	0.000	0.0000	0.00	0.00	0.00	
		Liquid F.	1.84	1.17	106.1	106.1	150.0	0.000	0.0000	0.00	0.00	0.00	
		2. Oil and natural gas and other emissions from energy production	335.53	194.09	5.0	29.6	30.0	0.015	0.0019	0.0036	0.03	0.00	
		CO2	25.96	27.06	5.0	8.7	10.0	0.000	0.0001	0.0005	0.00	0.00	
		N2O	0.60	0.14	5.0	79.8	80.0	0.000	0.0000	0.0000	0.00	0.00	
		CO2	43.74	5.14	15.5	15.5	21.9	0.000	0.0006	0.0001	0.01	0.00	
		Indirect emissions	2580.79	1714.79	2.0	3.0	3.6	0.017	0.0106	0.0321	0.09	0.00	
		1. Cement production	53.35	43.32	2.0	2.0	2.8	0.000	0.0001	0.0008	0.00	0.00	
		2. Lime production	15.25	7.31	2.0	3.0	3.6	0.000	0.0001	0.0001	0.00	0.00	
3. Metal industry	C. Metal industry	3. Glass production	163.95	85.24	2.0	2.0	2.8	0.000	0.0011	0.0016	0.00	0.00	
		4. Other process uses of carbonates	65.49	5.39	2.0	7.2	7.5	0.000	0.0010	0.0001	0.01	0.00	
		2. Nitric acid production	15.72	19.61	2.0	10.0	10.2	0.000	0.0001	0.0004	0.00	0.00	
		5. Carbide production	1.74	2.12	2.0	20.0	20.1	0.000	0.0000	0.0000	0.00	0.00	
		5. Carbide production	94.08	98.49	2.0	10.0	10.2	0.000	0.0003	0.0018	0.01	0.00	
		8. Petrochemical and carbon black production	11.87	12.81	2.0	10.0	10.2	0.000	0.0000	0.0002	0.00	0.00	
		10. Other (Niacin, Limestone pit)	9.20	11.15	2.0	5.0	5.4	0.000	0.0001	0.0002	0.00	0.00	
		1. Iron and steel production	139.26	-	5.0	20.0	20.6	-	0.0023	-	0.05	-	0.00
		3. Aluminium production	116.46	-	6.4	6.4	9.0	-	0.0019	-	0.01	-	0.00
		4. Non-energy products from fuels and solvent use	D. Non-energy products from fuels and solvent use	3. Aluminium production	-	39.51	18.0	18.0	25.5	0.000	0.0007	0.0007	0.02
4. Magnesium production	65.52			61.39	35.4	35.4	50.0	0.000	0.0001	0.0000	0.00	0.00	
7. Other	-			0.17	36.1	36.1	51.0	0.000	0.0000	0.0000	0.00	0.00	
1. Integrated circuit or semiconductor	-			4.69	57.6	57.6	81.5	0.000	0.0001	0.0001	0.01	0.00	
3. Photovoltaics	-			6.51	43.5	43.5	61.5	0.000	0.0001	0.0001	0.01	0.00	
4. Heat transfer fluid	-			0.49	137.9	137.9	195.0	0.000	0.0000	0.0000	0.00	0.00	
1. Refrigeration and air conditioning	0.02			1'431.58	14.1	14.1	20.0	-	-	-	-	-	0.43
2. Foam blowing agents	0.05			5.89	31.1	31.1	44.0	0.000	0.0001	0.0001	0.00	0.00	
4. Aerosols	-			32.65	32.2	32.2	45.5	0.001	0.0006	0.0006	0.02	0.03	
5. Solvents	-			14.09	29.7	29.7	42.0	0.000	0.0003	0.0003	0.01	0.01	

	IPCC Source category	Gas	Base year emissions or removals	Year 2015 emissions or removals	AD uncertainty	EF uncertainty	Combined uncertainty	Contribution to variance by Category in 2015	Type A sensitivity	Type B sensitivity	Uncertainty in trend in nat. emissions introduced by EF uncertainty	Uncertainty in trend in nat. emissions introduced by AD uncertainty	Uncertainty introduced into the trend in total national emissions
2G	2. Ind. Processes	CO ₂	6.60	34.28	35.4	35.4	50.0	-	0.0005	0.0006	0.02	0.03	0.00
		HFC	-	56.53	8.5	8.5	12.0	0.000	0.0011	0.0011	0.01	0.01	0.00
2H	2. Ind. Processes	PFC	-	46.63	10.6	10.6	15.0	0.000	0.0009	0.0009	0.01	0.01	0.00
		SF ₆	137.01	209.74	17.0	17.0	24.0	0.011	0.0017	0.0039	0.03	0.09	0.01
2I	2. Ind. Processes	N ₂ O	105.87	44.10	1.0	80.0	80.0	0.006	0.0009	0.0008	0.07	0.00	0.01
		CO ₂	1.04	0.89	3.0	7.1	7.7	0.000	0.0000	0.0000	0.00	0.00	0.00
3A	3. Agriculture	CO ₂	365.12	105.82	12.5	25.0	27.9	0.004	0.0041	0.0020	0.10	0.03	0.01
		CH ₄	3584.87	3344.78	6.4	16.9	18.1	1.643	0.0033	0.0625	0.06	0.57	0.33
3B	3. Agriculture	CH ₄	924.34	808.11	6.4	54.1	54.4	0.867	0.0002	0.0151	0.01	0.14	0.02
		N ₂ O	-	11.00	31.9	75.0	81.5	0.000	0.0002	0.0002	0.02	0.01	0.00
3D	3. Agriculture	N ₂ O	105.65	66.77	31.9	75.0	81.5	0.013	0.0005	0.0012	0.04	0.06	0.00
		N ₂ O	242.32	281.59	46.5	240.0	244.5	2.122	0.0013	0.0053	0.30	0.35	0.21
3E	3. Agriculture	N ₂ O	1'049.85	799.56	14.8	135.0	135.8	5.280	0.0024	0.0149	0.32	0.31	0.20
		N ₂ O	67.58	65.05	29.4	137.5	140.6	0.037	0.0001	0.0012	0.01	0.05	0.00
3D	3. Agriculture	N ₂ O	121.78	205.46	67.8	132.5	148.8	0.419	0.0018	0.0038	0.24	0.37	0.19
		N ₂ O	462.94	326.44	39.1	240.0	243.2	2.821	0.0015	0.0061	0.37	0.34	0.25
3G	3. Agriculture	N ₂ O	172.10	121.43	22.3	163.3	164.8	0.179	0.0006	0.0023	0.09	0.07	0.01
		CO ₂	22.25	32.86	40.0	5.0	40.3	0.001	0.0002	0.0006	0.00	0.03	0.00
3H	3. Agriculture	CO ₂	26.71	11.36	5.0	5.0	7.1	0.000	0.0002	0.0002	0.00	0.00	0.00
		CH ₄	763.39	350.90	21.2	21.2	30.0	0.050	0.0060	0.0066	0.13	0.20	0.06
5A	5. Waste	CO ₂	-	-	7.1	7.1	10.0	-	-	-	-	-	-
		CH ₄	35.51	109.84	10.0	40.0	41.2	0.009	0.0015	0.0021	0.06	0.03	0.00
5B	5. Waste	N ₂ O	5.42	16.75	10.0	79.4	80.0	0.001	0.0002	0.0003	0.02	0.00	0.00
		CH ₄	8.89	5.39	10.0	59.2	60.0	0.000	0.0000	0.0001	0.00	0.00	0.00
5C	5. Waste	CO ₂	53.73	9.94	10.0	38.7	40.0	0.000	0.0007	0.0002	0.03	0.00	0.00
		N ₂ O	16.78	31.98	10.0	38.7	40.0	0.001	0.0003	0.0006	0.01	0.01	0.00
5D	5. Waste	CH ₄	130.92	177.28	32.0	36.0	48.2	0.033	0.0012	0.0033	0.04	0.15	0.02
		N ₂ O	118.27	144.38	32.0	146.5	150.0	0.210	0.0007	0.0027	0.11	0.12	0.03
5I	5. Waste	CO ₂	2.01	1.18	33.6	33.6	47.6	0.000	0.0000	0.0000	0.00	0.00	0.00
		CH ₄	0.66	0.60	42.4	42.4	60.0	0.000	0.0000	0.0000	0.00	0.00	0.00
6	6. Other	CO ₂	10.96	11.35	28.3	28.3	40.0	0.000	0.0000	0.0002	0.00	0.01	0.00
		N ₂ O	0.60	0.51	106.1	106.1	150.0	0.000	0.0000	0.0000	0.00	0.00	0.00
6I	6. Other	CO ₂	1.04	1.04	17.2	57.8	60.3	0.000	0.0000	0.0000	0.00	0.00	0.00
		Indirect emissions	-	-	-	-	-	-	-	-	-	-	-

IPCC Source category		Gas	Base year emissions or removals	Year 2015 emissions or removals	AD uncertainty	EF uncertainty	Combined uncertainty	Contribution to variance by Category in 2015	Type A sensitivity	Type B sensitivity	Uncertainty in trend in nat. emissions introduced by EF uncertainty	Uncertainty in trend in nat. emissions introduced by AD uncertainty	Uncertainty introduced into the trend in total national emissions
			kt CO2 eq	kt CO2 eq	%	%	%		%	%	%	%	
4 II	Drainage, rewetting and other management of organic and mineral soils	CH4	10.75	10.75	10.0	70.0	70.7	0.000	0.0000	0.0002	0.00	0.00	0.00
4 III		N2O	7.77	8.00	34.2	137.5	141.7	0.001	0.0000	0.0001	0.00	0.01	0.00
4 IV		N2O	58.71	55.57	5.3	135.0	135.1	0.025	0.0001	0.0010	0.01	0.01	0.00
4 V		N2O	9.60	9.09	20.0	161.5	162.7	0.001	0.0000	0.0002	0.00	0.00	0.00
	Biomass burning	CH4	18.63	2.54	30.0	70.0	76.2	0.000	0.0003	0.0000	0.02	0.00	0.00
		CO2	-	-	10.0	70.0	70.7	-	-	-	-	-	-
4A1	1. Forest land remaining forest land	N2O	10.81	0.98	30.0	70.0	76.2	0.000	0.0002	0.0000	0.01	0.00	0.00
4A2		CO2	-318.64	-2'395.51	1.1	88.6	88.6	20.152	0.0395	0.0448	3.50	0.07	12.25
	2. Land converted to forest land	CO2	-571.07	-516.82	1.6	88.6	88.6	0.938	0.0002	0.0097	0.02	0.02	0.00
4B1		CO2	912.56	1'028.78	4.9	26.4	26.9	0.342	0.0042	0.0192	0.11	0.13	0.03
4B2	1. Cropland remaining cropland	CO2	46.58	41.22	5.2	31.6	32.0	0.001	0.0000	0.0008	0.00	0.01	0.00
4C1		CO2	79.58	112.04	5.2	867.9	867.9	4.234	0.0008	0.0021	0.68	0.02	0.46
4C2	2. Land converted to grassland	CO2	91.25	216.34	5.3	30.7	31.2	0.020	0.0025	0.0040	0.08	0.03	0.01
4D1		CO2	67.56	68.10	61.9	72.2	95.1	0.019	0.0002	0.0013	0.01	0.11	0.01
4D2	1. Wetlands remaining wetlands	CO2	21.87	34.13	4.0	21.4	21.8	0.000	0.0003	0.0006	0.01	0.00	0.00
4E1		CO2	17.27	54.61	4.4	50.0	50.2	0.003	0.0007	0.0010	0.04	0.01	0.00
4E2	2. Land converted to settlements	CO2	394.46	325.71	4.5	50.0	50.2	0.120	0.0004	0.0061	0.02	0.04	0.00
4F2		CO2	94.36	120.21	3.1	50.0	50.1	0.016	0.0007	0.0022	0.03	0.01	0.00
4G	G. Harvested wood products	CO2	-1'230.78	-69.53	4.5	57.0	57.2	0.007	0.0190	0.0013	1.08	0.01	1.18
Total Uncertainty including LULUCF			53'490	47'258	Percentage uncertainty in total inventory:			40.50			Trend uncertainty:		16.23
								6.36					4.03

A2.2 Detailed description of Approach 2 uncertainty analysis by Monte Carlo simulation

A2.2.1 Work steps

As a first step, the probability distributions need to be selected and their parameters need to be defined for the activity data and emission factors, based on measured data, literature or expert judgement. The mean values of the probability distributions are set equal to the values of the GHG inventory. In most cases, normal distributions are assumed. For some agricultural categories, triangular distributions are applied (see below).

In a second step, correlation coefficients for activity data, CO₂ emission factors and emission levels are chosen. Correlations may have a significant effect on the overall inventory uncertainty. Depending on whether correlations are negative or positive, they can lead to a decrease or increase in level uncertainty, respectively. Regrading trend uncertainty, positive correlations lead to a decrease and negative correlations to an increase in the trend uncertainty. Correlations were defined only for categories with relevant contributions to total uncertainty. If a large set of parameters is correlated between each other, the resulting correlation matrix might be mathematically inconsistent. In this case, the software Crystal Ball adjusts correlation coefficients iteratively such that the resulting correlation matrix is mathematically consistent. The modification of the correlation coefficients amounts in average to 0.10.

In the third step, Monte Carlo simulations are carried out to produce uncertainty results (see below). Several runs were performed to study the sensitivity to the choice of correlation strengths.

A2.2.2 Assumptions for the probability distributions

For almost all source and sink categories, normal distributions are chosen. The important exceptions are agricultural source categories as indicated in Table A – 3, where triangular distributions are applied.

Table A – 3 Probability distribution assigned to activity data and emission factors (1990 and 2015) of categories that are not considered normally distributed. For all other categories normal probability distributions have been assigned.

IPCC Source Category				Gas	Probability distribution	
					AD	EF
3B5	Agriculture	B. Manure Management	indirect	N2O	triangular	triangular
3Da1/2/4/5/7	Agriculture	D. Agricultural Soils; Direct Soil Emissions	fertilizer	N2O	normal	triangular
3Da6	Agriculture	D. Agricultural Soils; Direct Soil Emissions	organic soils	N2O	normal	triangular
3Da3	Agriculture	D. Agricultural Soils; Pasture, Range and Paddock Manure		N2O	triangular	triangular
3Db1	Agriculture	D. Agricultural Soils; Indirect Emissions	deposition	N2O	triangular	triangular
3Db2	Agriculture	D. Agricultural Soils; Indirect Emissions	leaching and runoff	N2O	triangular	triangular

A2.2.3 Assumptions for the correlation coefficients

Since there are no quantitative correlation coefficients available, only the following values have been used (if any are assumed):

- “strong” positive correlations are set to $r = 1.0$ (like perfect correlations),

- “medium” correlations are set to $r = \pm 0.5$.
- “weak” correlations are set to $r = \pm 0.25$.

The following assumptions are made for the level uncertainty:

- Activity data of liquid and gaseous fuels from the categories 1A2, 1A4a and 1A4b are negatively correlated ($r = -0.5$), since the total amount is well known but the partitioning into the different categories is less precisely known. By choosing negative correlations, overestimations in a category during the simulations are compensated by underestimations in one or more of the other categories.
- Activity data of 3A (Enteric Fermentation) and 3B (Manure Management) are positively correlated ($r = 0.5$) since they are both based on the same livestock numbers.
- Activity data 3B liquid and 3Db Indirect N₂O emissions (3Db1 deposition and 3Db2 leaching and runoff) are medium and positively correlated.

The following assumptions are made for the trend uncertainty:

- The CO₂ emission factor of category 1A1 (Other fuels) is correlated between 1990 and 2015. A medium positive correlation is assumed ($r = 0.5$).
- Activity data/emissions of the major sources (1A2: CO₂, 1A3: CO₂, 1A4: CO₂, 3A: CH₄, 3B: CH₄, 2F: HFC) are correlated between 1990 and 2015. A medium positive correlation is assumed ($r = 0.5$).
- The emission factors of agricultural categories are correlated between 1990 and 2015. A strong positive correlation is assumed.

Table A – 4 Correlation coefficients for correlated activity data.

b_AD_1A2_Gaseous_Fuels_CO2	b_AD_1A2_Gaseous_Fuels_CO2	b_AD_1A4a_Gaseous_Fuels_CO2	b_AD_1A4b_Gaseous_Fuels_CO2	t_AD_1A2_Liquid_Fuels_CO2	t_AD_1A2_Gaseous_Fuels_CO2	t_AD_1A4a_Gaseous_Fuels_CO2	t_AD_1A4b_Gaseous_Fuels_CO2
b_AD_1A2_Gaseous_Fuels_CO2	1.0	-0.5	1.0				
b_AD_1A4a_Gaseous_Fuels_CO2	-0.5	1.0	-0.5	1.0			
b_AD_1A4b_Gaseous_Fuels_CO2	0.5	0.5	0.5	0.5	1.0		
t_AD_1A2_Liquid_Fuels_CO2				1.0	-0.5	1.0	
t_AD_1A4a_Gaseous_Fuels_CO2				0.5	-0.5	-0.5	1.0
t_AD_1A4b_Gaseous_Fuels_CO2				0.5	-0.5	-0.5	1.0
b_AD_1A2_Liquid_Fuels_CO2	b_AD_1A2_Liquid_Fuels_CO2	b_AD_1A4a_Liquid_Fuels_CO2	b_AD_1A4b_Liquid_Fuels_CO2	t_AD_1A2_Liquid_Fuels_CO2	t_AD_1A2_Gaseous_Fuels_CO2	t_AD_1A4a_Liquid_Fuels_CO2	t_AD_1A4b_Liquid_Fuels_CO2
b_AD_1A2_Liquid_Fuels_CO2	1.0	-0.5	1.0				
b_AD_1A4a_Liquid_Fuels_CO2	-0.5	1.0	-0.5	1.0			
b_AD_1A4b_Liquid_Fuels_CO2	0.5	0.5	0.5	0.5	1.0		
t_AD_1A2_Liquid_Fuels_CO2				1.0	-0.5	1.0	
t_AD_1A4a_Liquid_Fuels_CO2				0.5	-0.5	-0.5	1.0
t_AD_1A4b_Liquid_Fuels_CO2				0.5	-0.5	-0.5	1.0
t_AD_3B_liquid_N2O	t_AD_3B_liquid_N2O	t_AD_3B_solid_N2O	t_AD_3B5_indirect_N2O	t_AD_3Db1_deposition_N2O	t_AD_3Db2_leaching and runoff_N2O		
t_AD_3B_liquid_N2O	1.0	-0.4	1.0				
t_AD_3B_solid_N2O	-0.4	1.0	1.0				
t_AD_3B5_indirect_N2O			1.0	1.0			
t_AD_3Db1_deposition_N2O	0.4	0.4	0.4	0.5	1.0		
t_AD_3Db2_leaching and runoff_N2O	0.4	0.4	0.4	0.5	0.5	1.0	
b_AD_3B_solid_N2O	b_AD_3B_solid_N2O	b_AD_3B5_indirect_N2O	b_AD_3Db1_deposition_N2O	b_AD_3Db2_leaching and runoff_N2O			
b_AD_3B_solid_N2O	1.0	1.0	1.0				
b_AD_3B5_indirect_N2O	0.5	0.5	0.5	1.0			
b_AD_3Db1_deposition_N2O	0.5	0.5	0.5	0.5	1.0		
b_AD_3Db2_leaching and runoff_N2O	0.5	0.5	0.5	0.5	0.5	1.0	
b_AD_1A3b_Gasoline_CO2	t_AD_1A3b_Gasoline_CO2						
b_AD_1A3b_Gasoline_CO2	1.0						
t_AD_1A3b_Gasoline_CO2	0.5	1.0					
b_AD_1A3b_Diesel_CO2	t_AD_1A3b_Diesel_CO2						
b_AD_1A3b_Diesel_CO2	1.0						
t_AD_1A3b_Diesel_CO2	0.5	1.0					

Table A – 5 Correlation coefficients for correlated CO₂ emission factors.

	b_EF_1A1_Other Fuels_CO2	t_EF_1A1_Other Fuels_CO2
b_EF_1A1_Other Fuels_CO2	1.0	
t_EF_1A1_Other Fuels_CO2	0.5	1.0
	b_EF_3B_solid_N2O	t_EF_3B_solid_N2O
b_EF_3B_solid_N2O	1.0	
t_EF_3B_solid_N2O	1.0	1.0
	b_EF_3Da1/2/4/5/7_fertilizer_N2O	t_EF_3Da1/2/4/5/7_fertilizer_N2O
b_EF_3Da1/2/4/5/7_fertilizer_N2O	1.0	
t_EF_3Da1/2/4/5/7_fertilizer_N2O	1.0	1.0
	b_EF_3Da3_0_N2O	t_EF_3Da3_N2O
b_EF_3Da3_0_N2O	1.0	
t_EF_3Da3_N2O	1.0	1.0
	b_EF_3Da6_organic soils_N2O	t_EF_3Da6_organic soils_N2O
b_EF_3Da6_organic soils_N2O	1.0	
t_EF_3Da6_organic soils_N2O	1.0	1.0
	b_EF_3Db2_leaching and runoff_N2O	t_EF_3Db2_leaching and runoff_N2O
b_EF_3Db2_leaching and runoff_N2O	1.0	
t_EF_3Db2_leaching and runoff_N2O	1.0	1.0
	b_EF_3Db1_deposition_N2O	t_EF_3Db1_deposition_N2O
b_EF_3Db1_deposition_N2O	1.0	
t_EF_3Db1_deposition_N2O	1.0	1.0
	b_EF_3B5_indirect_N2O	t_EF_3B5_indirect_N2O
b_EF_3B5_indirect_N2O	1.0	
t_EF_3B5_indirect_N2O	1.0	1.0

Table A – 6 Correlation coefficients for correlated emission levels.

	b_EM_2F1_HFC	t_EM_2F1_HFC		
b_EM_2F1_HFC	1.0			
t_EM_2F1_HFC	0.5	1.0		
	b_EM_3A_CH4	b_EM_3B_CH4	t_EM_3A_CH4	t_EM_3B_CH4
b_EM_3A_CH4	1.0			
b_EM_3B_CH4	0.5	1.0		
t_EM_3A_CH4	0.5		1.0	
t_EM_3B_CH4		0.5	0.5	1.0
	b_EM_3G_CO2	b_EM_3H_CO2	t_EM_3G_CO2	t_EM_3H_CO2
b_EM_3G_CO2	1.0			
b_EM_3H_CO2		1.0		
t_EM_3G_CO2	0.5		1.0	
t_EM_3H_CO2		0.5		1.0

A2.2.4 Detailed results of Monte Carlo simulations

Table A – 7 Results of Approach 2 uncertainty analysis, Monte Carlo simulation (Table 3.3 of the 2006 IPCC Guidelines, see also explanations therein on pp. 3.42-3.43 for each column). The uncertainty analysis includes indirect CO₂ emissions. If year 2015 emissions are zero, combined uncertainties are indicated as NO.

Categories (NFR, fuel, gas)	Base year (1990) emissions/ removals	Year 2015 emissions/ removals	Activity data uncertainty		Emission factor/ estimation parameter uncertainty		Combined uncertainty		Contribution to variance in year 2015	Inventory trend in national emissions 1990-2015	Uncertainty introduced into the trend in total national emissions with respect to base year	
	kt CO ₂ eq	kt CO ₂ eq	(-) %	(+) %	(-) %	(+) %	(-) %	(+) %	(fraction)	(% of base year)	(-) %	(+) %
1A1 Biomass CH4	0.45	0.28	-10	10	-28	28	-29	31	0.0000	-38	-36	35
1A1 Gaseous Fuels CH4	0.11	0.18	-5	5	-30	30	-30	30	0.0000	63	-57	57
1A1 Liquid Fuels CH4	0.52	0.27	-1	1	-30	30	-30	30	0.0000	-49	-34	34
1A1 Solid Fuels CH4	0.13	-	-5	5	-30	30	NO	NO	-	-100	-30	30
1A1 Gaseous Fuels CO2	243.40	398.76	-5	5	-1	1	-5	5	0.0000	64	-10	10
1A1 Liquid Fuels CO2	685.81	470.76	-1	1	-0	0	-1	1	0.0000	-31	-1	1
1A1 Other Fuels CO2	1'491.55	2'381.90	-5	5	-9	9	-10	11	0.0070	60	-16	16
1A1 Solid Fuels CO2	49.13	-	-5	5	-5	5	NO	NO	-	-100	-7	7
1A1 Biomass N2O	22.58	13.67	-10	10	-79	79	-80	82	0.0000	-40	-94	94
1A1 Gaseous Fuels N2O	0.13	0.21	-5	5	-80	80	-79	80	0.0000	63	-153	153
1A1 Liquid Fuels N2O	1.11	0.45	-1	1	-80	80	-80	80	0.0000	-59	-86	86
1A1 Other Fuels N2O	24.33	12.62	-5	5	-80	80	-80	80	0.0000	-48	-90	90
1A1 Solid Fuels N2O	0.24	-	-5	5	-80	80	NO	NO	-	-100	-80	80
1A2 Biomass CH4	4.28	1.86	-10	10	-28	28	-29	31	0.0000	-57	-33	32
1A2 Gaseous Fuels CH4	0.49	0.98	-5	5	-30	30	-30	30	0.0000	102	-68	68
1A2 Liquid Fuels CH4	4.48	1.40	-1	1	-30	30	-30	30	0.0000	-69	-32	31
1A2 Other Fuels CH4	0.77	0.53	-5	5	-30	30	-30	30	0.0000	-32	-36	36
1A2 Solid Fuels CH4	0.31	0.22	-5	5	-30	30	-30	30	0.0000	-29	-37	37
1A2 Gaseous Fuels CO2	1'059.63	2'220.03	-5	5	-1	1	-5	5	0.0014	109	-9	9
1A2 Liquid Fuels CO2	3'889.15	1'888.12	-1	1	-0	0	-1	1	0.0000	-51	-1	1
1A2 Other Fuels CO2	192.29	367.22	-5	5	-9	9	-10	11	0.0002	91	-22	23
1A2 Solid Fuels CO2	1'274.70	474.46	-5	5	-5	5	-7	7	0.0001	-63	-8	8
1A2 Biomass N2O	5.14	13.76	-10	10	-79	79	-79	81	0.0000	167	-228	229
1A2 Gaseous Fuels N2O	0.68	1.33	-5	5	-80	80	-79	81	0.0000	96	-175	177
1A2 Liquid Fuels N2O	13.03	10.85	-1	1	-80	80	-80	80	0.0000	-17	-104	104
1A2 Other Fuels N2O	2.32	5.58	-5	5	-80	80	-80	81	0.0000	141	-209	209
1A2 Solid Fuels N2O	6.14	2.24	-5	5	-80	80	-80	81	0.0000	-64	-86	85
1A3a kerosene CH4	0.17	0.13	-1	1	-30	30	-30	30	0.0000	-24	-37	37
1A3a kerosene CO2	252.55	137.37	-1	1	-0	0	-1	1	0.0000	-46	-1	1
1A3a kerosene N2O	2.06	1.12	-1	1	-150	150	-150	150	0.0000	-45	-173	172
1A3b Biomass CH4	-	0.12	-10	10	-59	59	-59	61	0.0000	-	-	-
1A3b Diesel CH4	1.67	0.60	-1	1	-20	20	-20	20	0.0000	-64	-21	21
1A3b Gaseous Fuels CH4	-	0.04	-5	5	-30	30	-30	30	0.0000	-	-	-
1A3b Gasoline CH4	116.04	18.00	-1	1	-37	37	-37	37	0.0000	-85	-37	37
1A3b Diesel CO2	2'632.59	7'181.99	-1	1	-0	0	-1	1	0.0005	173	-2	2
1A3b Gaseous Fuels CO2	-	35.53	-5	5	-1	1	-5	5	0.0000	-	-	-
1A3b Gasoline CO2	11'334.49	7'669.15	-1	1	-0	0	-1	1	0.0003	-32	-1	1
1A3b Biomass N2O	-	1.93	-10	10	-150	150	-150	152	0.0000	-	-	-
1A3b Diesel N2O	5.63	79.19	-1	1	-22	22	-22	22	0.0000	1'309	-312	313
1A3b Gaseous Fuels N2O	-	0.74	-5	5	-80	80	-80	80	0.0000	-	-	-
1A3b Gasoline N2O	136.88	25.09	-1	1	-50	50	-50	50	0.0000	-82	-51	51
1A3c Biomass CH4	-	0.00	-10	10	-59	59	-60	61	0.0000	-	-	-
1A3c Diesel CH4	0.03	0.01	-1	1	-30	30	-30	30	0.0000	-57	-33	33
1A3c Diesel CO2	28.69	28.72	-1	1	-0	0	-1	1	0.0000	0	-1	1
1A3c Biomass N2O	-	0.00	-10	10	-150	150	-150	150	0.0000	-	-	-
1A3c Diesel N2O	0.43	0.42	-1	1	-80	80	-80	80	0.0000	-3	-111	111
1A3d Biomass CH4	-	0.00	-10	10	-59	59	-59	61	0.0000	-	-	-
1A3d Liquid Fuels CH4	1.68	0.29	-1	1	-30	30	-30	30	0.0000	-82	-30	30
1A3d Liquid Fuels CO2	114.27	113.28	-1	1	-0	0	-1	1	0.0000	-1	-1	1
1A3d Biomass N2O	-	0.01	-10	10	-150	150	-151	152	0.0000	-	-	-
1A3d Liquid Fuels N2O	1.16	1.23	-1	1	-150	150	-150	149	0.0000	6	-219	218
1A3e Gaseous Fuels CH4	0.07	0.04	-5	5	-30	30	-30	30	0.0000	-46	-34	34
1A3e Gaseous Fuels CO2	31.42	42.86	-5	5	-1	1	-5	5	0.0000	36	-9	9
1A3e Gaseous Fuels N2O	0.02	0.02	-5	5	-80	80	-80	80	0.0000	36	-135	135
1A4a Biomass CH4	9.22	4.63	-10	10	-28	28	-29	31	0.0000	-50	-34	33
1A4a Gaseous Fuels CH4	0.70	1.30	-5	5	-30	30	-30	30	0.0000	84	-62	63
1A4a Liquid Fuels CH4	16.34	9.66	-1	1	-30	30	-30	30	0.0000	-41	-35	35
1A4a Gaseous Fuels CO2	981.49	1'400.80	-5	5	-1	1	-5	5	0.0006	43	-7	7
1A4a Liquid Fuels CO2	4'260.86	2'701.66	-1	1	-0	0	-1	1	0.0000	-37	-1	1
1A4a Biomass N2O	3.50	9.45	-10	10	-79	79	-79	81	0.0000	170	-229	233
1A4a Gaseous Fuels N2O	0.52	0.74	-5	5	-80	80	-80	81	0.0000	42	-140	140
1A4a Liquid Fuels N2O	10.35	6.59	-1	1	-80	80	-80	80	0.0000	-36	-95	95

Categories (NFR, fuel, gas)	Base year (1990) emissions/ removals	Year 2015 emissions/ removals	Activity data uncertainty		Emission factor/ estimation parameter uncertainty		Combined uncertainty		Contribution to variance in year 2015	Inventory trend in national emissions 1990-2015	Uncertainty introduced into the trend in total national emissions with respect to base year	
			kt CO2 eq	kt CO2 eq	(-) %	(+) %	(-) %	(+) %			(-) %	(+) %
1A4b Biomass CH4	109.97	25.57	-10	10	-28	28	-29	31	0.0000	-77	-31	30
1A4b Gaseous Fuels CH4	0.70	1.36	-5	5	-30	30	-30	30	0.0000	94	-65	66
1A4b Liquid Fuels CH4	34.83	20.04	-1	1	-30	30	-30	30	0.0000	-43	-34	35
1A4b Solid Fuels CH4	4.73	3.00	-5	5	-30	30	-30	30	0.0000	-36	-36	35
1A4b Gaseous Fuels CO2	1'449.69	2'600.43	-5	5	-1	1	-5	5	0.0020	79	-8	8
1A4b Liquid Fuels CO2	10'099.07	5'863.66	-1	1	-0	0	-1	1	0.0002	-42	-1	1
1A4b Solid Fuels CO2	58.40	37.08	-5	5	-5	5	-7	7	0.0000	-37	-9	8
1A4b Biomass N2O	25.86	21.37	-10	10	-79	79	-80	81	0.0000	-17	-104	104
1A4b Gaseous Fuels N2O	0.77	1.37	-5	5	-80	80	-79	80	0.0000	78	-163	164
1A4b Liquid Fuels N2O	24.53	14.26	-1	1	-80	80	-80	80	0.0000	-42	-93	92
1A4b Solid Fuels N2O	0.28	0.18	-5	5	-80	80	-80	81	0.0000	-37	-94	95
1A4c Biomass CH4	1.97	0.35	-10	10	-28	28	-29	31	0.0000	-82	-31	30
1A4c Gaseous Fuels CH4	0.02	0.01	-5	5	-30	30	-30	30	0.0000	-69	-32	31
1A4c Liquid Fuels CH4	6.45	1.43	-1	1	-30	30	-30	30	0.0000	-78	-31	31
1A4c Gaseous Fuels CO2	41.45	13.12	-5	5	-1	1	-5	5	0.0000	-68	-5	5
1A4c Liquid Fuels CO2	485.09	407.17	-1	1	-0	0	-1	1	0.0000	-16	-1	1
1A4c Biomass N2O	0.51	1.01	-10	10	-79	79	-79	81	0.0000	97	-176	179
1A4c Gaseous Fuels N2O	0.02	0.01	-5	5	-80	80	-80	80	0.0000	-69	-84	83
1A4c Liquid Fuels N2O	4.20	4.60	-1	1	-80	80	-80	80	0.0000	9	-119	118
1A5 Biomass CH4	-	0.00	-10	10	-59	59	-59	61	0.0000	-	-	-
1A5 Liquid Fuels CH4	0.27	0.16	-1	1	-60	60	-60	60	0.0000	-40	-70	70
1A5 Liquid Fuels CO2	217.65	134.04	-1	1	-0	0	-1	1	0.0000	-38	-1	1
1A5 Biomass N2O	-	0.00	-106	106	-106	106	-119	195	0.0000	-	-	-
1A5 Liquid Fuels N2O	1.84	1.17	-106	106	-106	106	-120	195	0.0000	-37	-210	177
1B2 CH4	335.53	194.09	-5	5	-30	30	-30	30	0.0004	-42	-35	35
1B2 CO2	25.96	27.06	-5	5	-9	9	-10	10	0.0000	4	-15	15
1B2 N2O	0.60	0.14	-5	5	-80	80	-80	80	0.0000	-77	-82	82
1ind CO2	43.74	5.14	-15	15	-15	15	-22	22	0.0000	-88	-22	22
2A1 CO2	2'580.79	1'714.79	-2	2	-3	3	-4	4	0.0004	-34	-4	4
2A2 CO2	53.35	43.32	-2	2	-2	2	-3	3	0.0000	-19	-4	4
2A3 CO2	15.25	7.31	-2	2	-3	3	-4	4	0.0000	-52	-4	4
2A4 CO2	163.95	85.24	-2	2	-2	2	-3	3	0.0000	-48	-3	3
2B2 N2O	65.49	5.39	-2	2	-7	7	-8	7	0.0000	-92	-8	7
2B5 CO2	15.72	19.61	-2	2	-10	10	-10	10	0.0000	25	-16	16
2B5 CH4	1.74	2.12	-2	2	-20	20	-20	20	0.0000	22	-32	32
2B8 CO2	94.08	98.49	-2	2	-10	10	-10	10	0.0000	5	-15	15
2B10 CO2	11.87	12.81	-2	2	-10	10	-10	10	0.0000	8	-15	15
2C1 CO2	9.20	11.15	-2	2	-5	5	-5	5	0.0000	21	-8	8
2C3 CO2	139.26	-	-5	5	-20	20	NO	NO	-	-100	-21	21
2C3 PFC	116.46	-	-6	6	-6	6	NO	NO	-	-100	-9	9
2C4 SF6	-	39.51	-18	18	-18	18	-25	26	0.0000	-	-	-
2C7 CO2	1.65	1.41	-2	2	-20	20	-20	20	0.0000	-15	-26	26
2D CO2	65.52	61.39	-35	35	-35	35	-50	50	0.0001	-6	-68	68
2E1 HFC	-	0.17	-36	36	-36	36	-51	51	0.0000	-	-	-
2E1 PFC	-	4.69	-58	58	-58	58	-81	82	0.0000	-	-	-
2E1 SF6	-	6.51	-43	43	-43	43	-62	61	0.0000	-	-	-
2E3 NF3	-	0.49	-138	138	-138	138	-195	195	0.0000	-	-	-
2E4 PFC	-	-	-51	51	-51	51	NO	NO	-	-	-	-
2F1 HFC	0.02	1'431.58	-14	14	-14	14	-20	20	0.0092	5'779'464	-1'154'502	1'158'737
2F1 PFC	0.05	5.89	-31	31	-31	31	-44	44	0.0000	10'878	-4'822	4'819
2F2 HFC	-	32.65	-32	32	-32	32	-46	46	0.0000	-	-	-
2F4 HFC	-	14.09	-30	30	-30	30	-42	42	0.0000	-	-	-
2F5 HFC	-	0.98	-43	43	-43	43	-62	62	0.0000	-	-	-
2G CO2	6.60	34.28	-35	35	-35	35	-50	50	0.0000	420	-265	266
2G HFC	-	56.53	-8	8	-8	8	-12	12	0.0000	-	-	-
2G PFC	-	46.63	-11	11	-11	11	-15	15	0.0000	-	-	-
2G SF6	137.01	209.74	-17	17	-17	17	-24	24	0.0003	53	-44	44
2G N2O	105.87	44.10	-1	1	-80	80	-80	80	0.0001	-58	-87	87
2H CO2	1.04	0.89	-3	3	-7	7	-8	8	0.0000	-15	-10	10
2ind CO2	365.12	105.82	-13	13	-25	25	-28	28	0.0001	-71	-29	29
3A CH4	3'584.87	3'344.78	-6	6	-17	17	-18.0	18.0	0.0410	-7	-17	17
3B CH4	924.34	808.11	-6	6	-54	54	-55	54	0.0218	-13	-51	50
3B N2O	-	11.00	-34	30	-50	100	-76	89	0.0000	-	-	-
3B N2O	105.65	66.77	-34	30	-50	100	-76	89	0.0003	-35	-59	39
3B5 N2O	242.32	281.59	-38	55	-80	400	-76	126	0.0484	10	-59	70

Categories (NFR, fuel, gas)	Base year (1990) emissions/ removals	Year 2015 emissions/ removals	Activity data uncertainty		Emission factor/ estimation parameter uncertainty		Combined uncertainty		Contribution to variance in year 2015	Inventory trend in national emissions 1990-2015	Uncertainty introduced into the trend in total national emissions with respect to base year	
			(-) %	(+) %	(-) %	(+) %	(-) %	(+) %			(fraction)	(% of base year)
	kt CO2 eq	kt CO2 eq										
3Da1/2/4/5/7 N2O	1'049.85	799.56	-16	13	-70	200	-64	87	0.0941	-24	-34	20
3Da6 N2O	67.58	65.05	-29	29	-75	200	-68	98	0.0007	-4	-48	45
3Da3 N2O	121.78	205.46	-53	83	-65	200	-68	114	0.0104	74	-93	149
3Db1 N2O	462.94	326.44	-33	45	-80	400	-76	121	0.0616	-34	-68	35
3Db2 N2O	172.10	121.43	-22	22	-93	233	-77	102	0.0031	-29	-45	25
3G CO2	22.25	32.86	-40	40	-5	5	-40	40	0.0000	48	-52	52
3H CO2	26.71	11.36	-5	5	-5	5	-7	7	0.0000	-57	-6	6
5A CH4	763.39	350.90	-21	21	-21	21	-30	30	0.0012	-54	-33	33
5A CO2	-	-	-7	7	-7	7	NO	NO	-	-	-	-
5B CH4	35.51	109.84	-10	10	-40	40	-41	41	0.0002	209	-134	134
5B N2O	5.42	16.75	-10	10	-79	79	-80	80	0.0000	210	-261	260
5C CH4	8.89	5.39	-10	10	-59	59	-60	60	0.0000	-39	-70	70
5C CO2	53.73	9.94	-10	10	-39	39	-40	40	0.0000	-81	-41	41
5C N2O	16.78	31.98	-10	10	-39	39	-40	40	0.0000	91	-87	86
5D CH4	130.92	177.28	-32	32	-36	36	-48	48	0.0008	35	-81	81
5D N2O	118.27	144.38	-32	32	-147	147	-150	149	0.0053	23	-237	237
5ind CO2	2.01	1.18	-34	34	-34	34	-48	48	0.0000	-41	-55	56
6 CH4	0.66	0.60	-42	42	-42	42	-60	60	0.0000	-8	-82	82
6 CO2	10.96	11.35	-28	28	-28	28	-40	40	0.0000	3	-58	58
6 N2O	0.60	0.51	-106	106	-106	106	-150	150	0.0000	0	-214	212
6ind CO2	1.04	1.04	-17	17	-58	58	-60	60	0.0000	0	-86	85
4 II CH4	10.75	10.75	-10	10	-70	70	-70	70	0.0000	0	-101	100
4 II N2O	7.77	8.00	-34	34	-138	138	-142	142	0.0000	3	-202	203
4 III N2O	58.71	55.57	-5	5	-135	135	-135	134	0.0006	-5	-185	185
4 IV N2O	9.60	9.09	-20	20	-162	162	-164	163	0.0000	-6	-223	223
4 V CH4	18.63	2.54	-30	30	-70	70	-77	76	0.0000	-86	-77	77
4 V CO2	-	-	-10	10	-70	70	NO	NO	-	-	-	-
4 V N2O	10.81	0.98	-30	30	-70	70	-76	75	0.0000	-91	-76	76
4A1 CO2	-318.64	-2'395.51	-1	1	-89	89	88	-88	0.5039	651	670	-670
4A2 CO2	-571.07	-516.82	-2	2	-89	89	89	-88	0.0235	-10	119	-119
4B1 CO2	912.56	1'028.78	-5	5	-26	26	-27	27	0.0086	13	-41	40
4B2 CO2	46.58	41.22	-5	5	-32	32	-32	32	0.0000	-11	-43	43
4C1 CO2	79.58	112.04	-5	5	-868	868	-870	864	0.1066	39	-1'482	1'478
4C2 CO2	91.25	216.34	-5	5	-31	31	-31	31	0.0005	137	-81	80
4D1 CO2	67.56	68.10	-62	62	-72	72	-95	95	0.0005	1	-135	135
4D2 CO2	21.87	34.13	-4	4	-21	21	-22	22	0.0000	56	-41	40
4E1 CO2	17.27	54.61	-4	4	-50	50	-50	50	0.0001	215	-166	165
4E2 CO2	394.46	325.71	-5	5	-50	50	-50	50	0.0030	-17	-65	66
4F2 CO2	94.36	120.21	-3	3	-50	50	-50	50	0.0004	27	-81	82
4G CO2	-1'230.78	-69.53	-5	5	-57	57	57	-58	0.0002	-94	57	-58
incl LULUCF	53'490	47'258					-6.08	6.21		-11.83	-5.53	5.51
excl. LULUCF	53'768	48'151					-3.39	3.73		-10.66	-2.69	2.55

A2.2.5 Relation between simulated and inventory values

The Monte Carlo method simulates a probability distribution of the Swiss greenhouse gas emissions from which all relevant statistical parameters can be derived (mean, standard deviation and percentiles). The simulated mean value may slightly differ from the reported value.

The discrepancy between simulated and reported values becomes apparent when mean numbers in Figure 1-3 are compared to reported numbers in the summary tables. It is not a relevant issue for the uncertainty analysis but can be confusing for readers and reviewers who carefully study the numbers. For transparency reasons, the numbers are explained in Table A – 8.

The absolute percentiles generated by the simulation are firstly expressed as relative numbers (the simulated mean is set to 100%). Then the relative numbers are transferred to the numbers reported in the summary tables, then they are applied to derive the absolute uncertainties.

Table A – 8 Mean values, 2.5 and 97.5 percentiles of the Monte Carlo simulation and corresponding values of the reported emissions (as listed in summary tables). The uncertainty analysis includes indirect CO₂ emissions.

Year	Parameters	Unit	Emission (excl. LULUCF)	Lower bound 2.5 percentile	Upper bound 97.5 percentile	Lower uncertainty	Upper uncertainty
2015	simulated values						
	absolute	kt CO ₂ eq	49'414	47'739	51'255	-1'675	1'841
	relative	%	100.00%	96.61%	103.73%	-3.39%	3.73%
	reported values						
	absolute	kt CO ₂ eq	48'151	46'519	49'945	-1'632	1'794
	relative	%	100.00%	96.61%	103.73%	-3.39%	3.73%
1990	simulated values						
	absolute	kt CO ₂ eq	55'308	53'337	57'539	-1'971	2'230
	relative	%	100.00%	96.44%	104.03%	-3.56%	4.03%
	reported values						
	absolute	kt CO ₂ eq	53'768	51'852	55'937	-1'916	2'168
	relative	%	100.00%	96.44%	104.03%	-3.56%	4.03%

Annex 3: Other detailed methodological descriptions for individual source or sink categories

A3.1 Sector Energy

A3.1.1 Emission from manufacturing industries and construction

The emission factors of precursors in the manufacturing industries and construction sector are given below. Further and more detailed emission factors can be found in Switzerland's Informative Inventory Report (FOEN 2017f).

Emission factors for greenhouse gases are given in 3.2.6.2.

Table A – 9 Emission factors 2015 of precursors from boiler in 1A2 Manufacturing industries and construction.

1A2 Boiler	NO _x	NM VOC	SO ₂	CO
	kg/TJ			
Boiler gas oil	32	2	18	7
Boiler residual fuel oil	125	4	291	10
Boiler liquefied petroleum gas	19	2	0.5	9
Boiler petroleum coke	125	4	291	10
Boiler other bituminous coal	200	10	500	100
Boiler lignite	208	10	500	100
Boiler natural gas	19	2	0.5	9

A3.1.2 Civil aviation

This paragraph contains further information on the emission modelling. More complete information is provided in FOCA (2006–2014) and on request for reviewers by FOCA.

Emission factors (1A3a)

Table A – 10 Aircraft cruise factors, used for cruise emission calculation (extract of list of 881 aircraft) GKL_ICAO = ICAO seat categories. Mass emissions are given in kilograms or grams per nautical mile (NM).

Aircraft Cruise_Factors						
Aircraft_ICAO	GKL_ICAO	Cruise_D_Source	kg_fuel_NM	kg_NOx_NM	g_VOC_NM	g_CO_NM
AA1	0	P002FOCA	0.21	0.0098	1.79	61.7
AA5	0	P002FOCA	0.21	0.0098	1.79	61.7
AC11	0	P002FOCA	0.21	0.0098	1.79	61.7
AC14	0	P002FOCA	0.21	0.0098	1.79	61.7
AC50	0	P001FOCA	0.77	0.021	4.14	364.17
AC68	0	P001FOCA	0.77	0.0075	4.14	364.17
AC6T	1	FOCAINV95-03.2T	1.58	0.021	0.87	2.9
AC90	1	FOCAINV95-03.2T	1.58	0.021	0.87	2.9
AC95	1	FOCAINV95-03.2T	1.58	0.021	0.87	2.9
AEST	0	P001FOCA	0.77	0.021	4.14	364.17
AJET	0	FOCAEDBJ014	2.92	0.0146	8.53	63
ALO2	0	FOCAHeli	1.91	0.024	0.42	2.1
ALO3	0	FOCAHeli	1.91	0.024	0.42	2.1
AN12	0	AN26*2	5.36	0.0062	143	348
AN2	0	FOCA/91/DC3	0.82	0.0002	13.7	1000
AN22	6	FOCAINV95-03.2T*2	3.16	0.042	1.74	5.8
AN24	2	AN26	2.68	0.0031	71.7	174
AN26	1	500	2.68	0.0031	71.7	174
AN72	2	FOCAINV95-03.2J	6.4	0.1	0.83	10
AR7	0	P002FOCA	0.21	0.0098	1.79	61.7
AR7A	0	P002FOCA	0.21	0.0098	1.79	61.7
AS02	0	P002FOCA	0.21	0.0098	1.79	61.7
AS16	0	P002FOCA	0.21	0.0098	1.79	61.7
AS20	0	P002FOCA	0.21	0.0098	1.79	61.7
AS24	0	P002FOCA	0.21	0.0098	1.79	61.7
AS25	0	P002FOCA	0.21	0.0098	1.79	61.7
AS26	0	P002FOCA	0.21	0.0098	1.79	61.7
AS2T	0	FOCAEDBT758	0.95	0.005	1.8	12
AS30	0	FOCAHeli*2	3.82	0.048	0.82	4.2
AS32	1	FOCAHeli*2	3.82	0.048	0.82	4.2
AS33	0	FOCAHeli*2	3.82	0.048	0.82	4.2
AS35	0	FOCAHeli	1.91	0.024	0.42	2.1
AS50	0	FOCAHeli*2	3.82	0.048	0.82	4.2
AS55	0	FOCAHeli*2	3.82	0.048	0.82	4.2
AS65	0	FOCAHeli*2	3.82	0.048	0.82	4.2
ASK1	0	P002FOCA	0.21	0.0098	1.79	61.7
ASTA	0	FOCAINV95-03.B	3.016	0.046	0.3	2.8
ASTR	0	FOCAINV95-03.B	3.016	0.046	0.3	2.8
ASTRA	0	FOCAINV95-03.B	3.016	0.046	0.3	2.8
AT42	1	FOCAINV95-03.2T	1.58	0.021	0.87	2.9
AT43	1	500	1.6	0.013	0	15

Activity data (1A3a)

LTO-cycle times (minutes). ICAO standard cycle times were originally designed for emissions certification, not for emissions modelling. Today, they do generally not match real world aircraft LTO operations. Swiss FOCA has therefore adjusted some of the ICAO standard cycle times for different aircraft categories. For jets, the mean time for taxi-in and taxi-out at Swiss airports has been determined 20 minutes instead of the standard 26 minutes.

Table A – 11 For jets, business jets, turboprops, piston engines and helicopters, the times in mode are shown and are based on ICAO, US EPA and Swiss FOCA data. “Type” is a classification variable. J = Jet, T = Turboprop, P = Piston, H = Helicopter, HP = Helicopter with Piston Engine, B = Business jet, SJ = Supersonic Jet, E = Electric Aircraft. The number in “Type” stands for the number of engines. For Jet Aircraft, the cycle times and associated thrust settings still lead to an overestimation of LTO emissions (FOCA 2007b).

LTO Cycle				
Type	Time_Take_Off	Time_Climbout	Time_Approach	Time_Taxi
1J	0.7	2.2	4	20
1T	0.5	2.5	4.5	13
1P	0.3	2.5	3	12
1H	0	3	5.5	5
2B	0.4	0.5	1.6	13
3B	0.4	0.5	1.6	13
2T	0.5	2.5	4.5	13
4T	0.5	2.5	4.5	13
2J	0.7	2.2	4	20
3J	0.7	2.2	4	20
4J	0.7	2.2	4	20
2P	0.3	2.5	3	12
3P	0.3	2.5	3	12
4P	0.3	2.5	3	12
2H	0	3	5.5	5
4SJ	1.2	2	2.3	20
3H	0	3	5.5	5
4H	0	3	5.5	5
4B	0.4	0.5	1.6	13
1HP	0	4	5.5	5
2HP	0	4	5.5	5
3HP	0	4	5.5	5
4HP	0	4	5.5	5
1B	0.4	0.5	1.6	13
1E	0.7	10	5	13
4E	0.3	10	5	13
6J	0.7	2.2	4	20

Table A – 12 Aircraft-Engine Combinations and associated codes for SWISS FOCA emissions database. (Extract from list of more than 40'000 individual aircraft)

Aircraft Engine Combinations							
Engine Name	Aircraft Name	Aircraft Registr.	No. Eng.	Code	Type	Aircr. ICAO	Source
V2527-A5	AIRBUS A320-232	ECHXA	2	J220	2J	A320	1IA003
CF34-3B1	BOMBARDIER CRJ200ER (CL-600-2B19)	ECHXM	2	J090	2J	CRJ2	1GE034
CFM56-3C1	BOEING 737-4K5	ECHXT	2	J022	2J	B734	1CM007
TPE331-11U-611G	FAIRCHILD (SWEARIN-GEN) SA227AC METR	ECHXY	2	T310	2T	SW4	FOI
CFM56-5B4/P	AIRBUS A320-214	ECHYC	2	J067	2J	A320	3CM026
CFM56-5B4/P	AIRBUS A320-214	ECHYD	2	J067	2J	A320	3CM026
CF34-3B1	BOMBARDIER CRJ200ER (CL-600-2B19)	ECHYG	2	J090	2J	CRJ2	1GE034
CFEC-FE738-1-1B	DASSAULT FALCON 2000	ECHYI	2	B130	2B	F2TH	FOI-Honeywell
GA TPE331-11U-612G		ECHZH	2	T310	2T	FA3	FOI
CF34-3B1	BOMBARDIER CRJ200ER (CL-600-2B19)	ECHZR	2	J090	2J	CRJ2	1GE034
CFM56-7B27B1	BOEING 737-86Q (WINGLETS)	ECHZS	2	J075	2J	B738	3CM034
CFM56-5B4/P	AIRBUS A320-214	ECHZU	2	J067	2J	A320	3CM026
CF34-3B1	BOMBARDIER CRJ200ER (CL-600-2B19)	ECIAA	2	J090	2J	CRJ2	1GE034
FJ44-1A	CESSNA 525 CITATIONJET	ECIAB	2	B001	2B	C525	FOCA
CFM56-5B4/P	AIRBUS A320-214	ECIAG	2	J067	2J	A320	3CM026
V2527-A5	AIRBUS A320-232	ECIAZ	2	J220	2J	A320	1IA003
BRBR700-710A2-20	BOMBARDIER BD-700-1A10 GLOBAL EX-PRE	ECIBD	2	J854	2J	GLEX	4BR009
PT6A-60A	BEECH-CRAFT KING AIR 350 (RAYTHEON B)	ECIBK	2	T738	2T	B350	FOI
CF34-3B1	BOMBARDIER CRJ200ER (CL-600-2B19)	ECIBM	2	J090	2J	CRJ2	1GE034
CFM56-7B27B1	BOEING 737-81Q (WINGLETS)	ECICD	2	J075	2J	B738	3CM034
CFM56-5B4/P	AIRBUS A320-214	ECICK	2	J067	2J	A320	3CM026

Emissions (1A3a)

The output of the FOCA emission modelling consists of tables with the following structure:

Table A – 13 Extract of the output file of FOCA emission and fuel consumption modelling. Upper part: LTO, lower part: cruise (example for 2004). Emissions and fuel consumption in tonnes.

Airport	Distance	Type Traffic	Movements	Type	Aircraft ICAO	Engine Name	Fuel (LTO) tons	Emissions (LTO) in tons					
	Km		No.					CO ₂	H ₂ O	SO ₂	NO _x	VOC	CO
LSGG	181501.69	Taxi	165	2B	C550	JT15D-4	5673.492	17871.5	6978.395	5.673	26.04	139	359.2
LSGG	164165.197	Taxi	77	2J	B752	RB211-535E4	47470.5	149532.1	58388.72	47.47	554.91	0	361.47
LSGG	133166.837	Taxi	118	2B	F2TH	CFE738-1-1B	6164.2728	19417.46	7582.056	6.164	87.539	40.59	185.53
LSGG	117228.943	Taxi	99	3B	F900	TFE731-60-1C	5668.542	17855.91	6972.307	5.669	46.937	28.13	163.44
LSGG	114258.902	Taxi	134	2B	LJ45	TFE731-20R	4725.108	14884.09	5811.883	4.725	31.31	53.62	169.01
LSGG	112510.267	Taxi	100	2B	F2TH	CFE738-1-1B	5223.96	16455.47	6425.471	5.224	74.186	34.4	157.23
LSGG	107945.477	Taxi	96	2B	C560	JT15D-5D	3795.3216	11955.26	4668.246	3.795	16.959	271.6	287.98
LSGG	181501.69	Taxi	165	2B	C550	JT15D-4	307732.68	969357.9	378511.2	307.7	4513	29.43	274.71
LSGG	164165.197	Taxi	77	2J	B752	RB211-535E4	673698.47	2122150	828649.1	673.7	7986.4	647.8	1038.2
LSGG	133166.837	Taxi	118	2B	F2TH	CFE738-1-1B	225781.85	711212.8	277711.7	225.8	3311.2	21.59	201.55
LSGG	117228.943	Taxi	99	3B	F900	TFE731-60-1C	298139.18	939138.4	366711.2	298.1	4372.3	28.52	266.14
LSGG	114258.902	Taxi	134	2B	LJ45	TFE731-20R	193723.81	610230	238280.3	193.7	2841	18.53	172.93
LSGG	106761.289	Taxi	100	2B	F2TH	CFE738-1-1B	181011.75	570187	222644.4	181	2654.6	17.31	161.58
LSGG	103217.159	Taxi	96	2B	C560	JT15D-5D	175002.74	551258.6	215253.4	175	2566.5	16.74	156.22

A3.1.3 Road transportation

Emission factors (1A3b)

The derivation of the emission factors for road vehicles is described in detail in TUG (2009). Some important features of the emission factor methodologies are summarised in this paragraph.

The emission factors have to be differentiated according to vehicle categories. Each category contains a number of vehicle classes, which differ by emission concepts. The following table illustrates the classes of the passenger cars. Similar “segmentations” hold for the other vehicle categories too. Emission factors for vehicle classes are combined to average emission factors for vehicles categories weighted according to the fleet composition, which varies from year to year (see below).

Table A – 14 Vehicle segmentation of the passenger cars. Each segment is subdivided into three cubic capacities: <1.4 litre, 1.4-2.0 litres, >2.0 litres (INFRAS 2010).

Fuel type	Vehicle segment
Gasoline	<ECE
	AGV82 (CH)
	PreEuro 3WayCat <1987
	PreEuro 3WayCat 1987-90
	ECE-15'00
	ECE-15'01/02
	ECE-15'03
	Euro-1
	Euro-2
	Euro-3
	Euro-4
	Euro-5
	Euro-6
Diesel	<1986
	1986-1988
	Euro-1
	Euro-2
	Euro-3
	Euro-4
	Euro-5 Diesel Particle Filter
	Euro-6 Diesel Particle Filter

The emission factors published in the handbook (CD ROM, INFRAS 2010) are classified by “traffic situations”. The scheme (see table below) distinguishes the traffic situations along 4 dimensions: urban/rural areas, 5 functional road types, speed limit and 4 levels of service. This leads to the definition of 276 different traffic situations in total. A traffic situation is primarily characterised by the type of road which induces a typical driving behaviour. (Because driving behaviour is not independent of the amount of traffic on that particular road, on the same segment different driving patterns may exist.) For the handbook several typical traffic situations have been defined, based on driving behaviour studies in Germany and in Switzerland (see e.g. SAEFL 1995, chp. 4).

Table A – 15 Traffic situation-scheme in HBEFA 3.1 (INFRAS 2010). Every traffic situation is characterised by a typical driving pattern (i.e. a speed-time curve)

Area	Road type	Levels of service	Speed Limit [km/h]													
			30	40	50	60	70	80	90	100	110	120	130	>130		
Rural	Motorway-Nat.	4 levels of service														
	Semi-Motorway	4 levels of service														
	TrunkRoad/Primary-Nat.	4 levels of service														
	Distributor/Secondary	4 levels of service														
	Distributor/Secondary(sinuous)	4 levels of service														
	Local/Collector	4 levels of service														
	Local/Collector(sinuous)	4 levels of service														
	Access-residential	4 levels of service														
Urban	Motorway-Nat.	4 levels of service														
	Motorway-City	4 levels of service														
	TrunkRoad/Primary-Nat.	4 levels of service														
	TrunkRoad/Primary-City	4 levels of service														
	Distributor/Secondary	4 levels of service														
	Local/Collector	4 levels of service														
	Local/Collector	4 levels of service														
	Access-residential	4 levels of service														

Traffic situations are defined independently of vehicle categories (LDV, HDV, 2-wheelers). But behind the same traffic situation each vehicle category may know its own “driving pattern” which may be expressed as a speed curve (i.e. speed time series). Emission factors originally are derived for these underlying driving patterns based on measurements performed on laboratory test benches. Emission factors per traffic situation are then calculated by attributing the driving patterns to different traffic situations (based on statistical analyses).

Emission factors for Switzerland are shown in the next table (FOEN 2010i, updated based on Prognos 2012a). They represent weighted averages over all traffic situations. The year indicates the date when the corresponding vehicle class appears in the market. E.g. “Euro-3” standard came into force on 1 Jan, 2001, but the first vehicles with Euro-3 standard already appeared in 1999.

Table A – 16 Mean emission factors of passenger cars (PC), light duty vehicles (LDV), heavy duty vehicles (HDV), coaches, urban buses (Bus) and Motorcycles (MC) in grams per kilometre, incl. cold starts and evaporation. (FOEN 2010i, updated based on Prognos 2012a). CO₂ (rep.) refers to the fossil part, CO₂ (total) includes fossil and biomass. HC and NMHC corresponds to VOC and NMVOC.

Pollutant	Year	PC	LDV	HDV	Coach	Bus	MC
grams per vehicle kilometre, incl. cold starts and evaporation							
CH ₄	1990	0.080	0.090	0.020	0.017	0.053	0.236
CH ₄	1995	0.050	0.065	0.018	0.016	0.046	0.159
CH ₄	2000	0.033	0.039	0.013	0.014	0.034	0.120
CH ₄	2005	0.021	0.020	0.009	0.011	0.018	0.103
CH ₄	2010	0.013	0.012	0.004	0.007	0.005	0.092
CH ₄	2015	0.009	0.006	0.002	0.004	0.003	0.082
CO	1990	9.58	20.16	2.39	2.09	5.99	14.70
CO	1995	5.46	14.60	2.18	2.01	5.68	14.14
CO	2000	3.59	8.86	1.77	1.84	4.64	13.62
CO	2005	2.53	4.39	1.61	1.73	2.92	11.68
CO	2010	1.64	2.09	1.44	1.70	1.26	8.09
CO	2015	1.19	1.28	1.27	1.63	1.03	5.63
CO ₂ (rep.)	1990	234	249	803	871	1'194	82
CO ₂ (rep.)	1995	237	252	799	860	1'199	90
CO ₂ (rep.)	2000	230	254	759	833	1'162	92
CO ₂ (rep.)	2005	217	246	790	823	1'127	94
CO ₂ (rep.)	2010	195	237	768	812	1'078	96
CO ₂ (rep.)	2015	169	224	739	794	1'045	91
CO ₂ (total)	1990	234	249	803	871	1'194	82
CO ₂ (total)	1995	237	252	799	860	1'199	90
CO ₂ (total)	2000	230	255	760	834	1'163	92
CO ₂ (total)	2005	217	246	793	826	1'131	94
CO ₂ (total)	2010	199	240	777	821	1'094	99
CO ₂ (total)	2015	180	232	760	817	1'079	98
HC	1990	1.58	2.02	0.85	0.70	2.20	3.69
HC	1995	0.92	1.38	0.74	0.66	1.93	2.65
HC	2000	0.57	0.77	0.56	0.60	1.42	2.08
HC	2005	0.36	0.38	0.38	0.47	0.73	1.64
HC	2010	0.23	0.19	0.16	0.27	0.22	1.19
HC	2015	0.17	0.12	0.07	0.16	0.12	0.85
N ₂ O	1990	0.009	0.005	0.008	0.008	0.003	0.002
N ₂ O	1995	0.012	0.007	0.009	0.008	0.003	0.002
N ₂ O	2000	0.011	0.009	0.009	0.008	0.003	0.002
N ₂ O	2005	0.005	0.007	0.008	0.007	0.002	0.002
N ₂ O	2010	0.003	0.006	0.030	0.014	0.001	0.002
N ₂ O	2015	0.002	0.005	0.041	0.023	0.002	0.002
N ₂ O	2020	0.002	0.005	0.043	0.029	0.005	0.002
N ₂ O	2025	0.002	0.005	0.043	0.032	0.008	0.002
N ₂ O	2030	0.002	0.004	0.042	0.033	0.009	0.002
N ₂ O	2035	0.002	0.004	0.042	0.033	0.010	0.002
N ₂ O	2040	0.002	0.004	0.041	0.033	0.010	0.002
N ₂ O	2045	0.002	0.004	0.041	0.033	0.010	0.002
N ₂ O	2050	0.002	0.003	0.041	0.032	0.010	0.002
NMHC	1990	1.504	1.930	0.827	0.681	2.151	3.451
NMHC	1995	0.871	1.320	0.721	0.640	1.880	2.489
NMHC	2000	0.538	0.735	0.543	0.582	1.383	1.964
NMHC	2005	0.343	0.362	0.373	0.459	0.714	1.538
NMHC	2010	0.217	0.180	0.155	0.265	0.209	1.096
NMHC	2015	0.163	0.116	0.069	0.156	0.110	0.773
NO _x	1990	2.147	4.167	22.457	22.929	33.896	0.294
NO _x	1995	1.608	3.485	20.751	21.648	32.841	0.392
NO _x	2000	1.301	3.067	18.148	19.938	29.997	0.423
NO _x	2005	0.968	2.595	15.051	17.360	24.703	0.444
NO _x	2010	0.679	2.117	9.512	13.381	18.000	0.400
NO _x	2015	0.535	1.857	6.213	10.142	13.576	0.352
SO ₂	1990	0.040	0.093	0.714	0.774	1.061	0.010
SO ₂	1995	0.031	0.041	0.173	0.186	0.260	0.011
SO ₂	2000	0.022	0.034	0.131	0.144	0.201	0.008
SO ₂	2005	0.001	0.001	0.005	0.005	0.007	0.000
SO ₂	2010	0.001	0.001	0.005	0.005	0.007	0.001
SO ₂	2015	0.001	0.001	0.005	0.005	0.006	0.001

Table A – 17 Cold start emission factors of passenger cars (PC) and light duty vehicles (LDV). The factors are adopted from Copert and recommended in the EMEP/EEA Gudibook (EMEP/EEA 2013).

Cold start emission factors N ₂ O				
Euro class	PC		LDV	
	gasoline mg/km	diesel oil mg/km	gasoline mg/km	diesel oil mg/km
EURO0	10.0	0	10.0	0
EURO1	19.0	0	47.3	0
EURO2	12.9	3	86.9	3
EURO3	8.4	15	17.1	15
EURO4	5.5	15	13.7	15
EURO5	2.0	15	1.8	15
EURO6	1.7	9	1.7	9

Activity data (1A3b)

Activity data for the emission model are the mileages of the vehicle categories per traffic situation. To that aim, three steps must be carried out.

1. Vehicle turnover: The vehicle fleet is built up for each year accounting for the stock changes. This vehicle turnover is modelled on the basis of new registrations and by applying survival probabilities. Trends in traffic volume per vehicle category, including structural changes (size distributions, shares of diesel vehicles) are then combined to draw the continual substitution of older technologies by new ones constantly altering the fleet composition or mileage by emission concepts in all vehicle categories (see following Figure A-1).

2. The total mileage is calculated by vehicle stock multiplied with the specific mileage per vehicle and annum. The latter data are derived from surveys and from specific odometer readings during vehicle inspections (ARE 2002).

3. Assignment of the mileage to the traffic situations for all vehicle categories. This step requires the adoption of the traffic model: Each road segment carries its mileage and its traffic, which allows the assignment sought.




Figure A – 1 Mileage composition by emission concept (in million vehicle kilometres per year) for passenger cars, light duty trucks (“light commercial vehicles”), motorcycles and heavy duty trucks (“heavy goods vehicles”), FOEN 2010i.

Modelling hot exhaust emissions (1A3b)

As a next step in the modelling process, the mileage classified by vehicle segments and traffic situations is multiplied with the emission factors resulting in hot exhaust emissions.

The results do not yet contain the emissions from fuel tourism and statistical differences. For this purpose a special procedure is carried out (described in chp. 3.2.9.2.2), providing the fuel consumption of fuel tourism and statistical differences. From that, the emissions are calculated by multiplication with mean emission factors.

Cold start and evaporative emissions (1A3b)

The handbook also contains emission factors for modelling cold start excess emissions and evaporative emissions (diurnal and hot/warm soak and running losses). For a technical description the reader may refer to TUG (2009).

Results show that for CO₂ the hot exhaust emissions contribute to 97% of the total. Only 3% stem from cold start excess emissions. For CH₄, however, the picture is different. Only about 40% of the emission total is hot exhaust. More than 59% are cold start excess emissions, the rest results evaporative emissions. For N₂O, no cold start emissions or evaporative emissions are taken into account due to lack of data. However, the missing N₂O emissions have been estimated by means of emission factors provided by the Copert model and which are provided in the EMEP Guidebook (EMEP/EEA 2013). For details see chp. 3.2.9.2.2.

A3.1.4 Non-road vehicles: supplementary activity data

The following table shows some aggregated information on stock numbers and annual operation hours of non-road vehicles. Detailed information is available in the report FOEN (2015j) and most disaggregated information is available by query from the online non-road database INFRAS (2015a):

<https://www.bafu.admin.ch/bafu/en/home/topics/air/state/non-road-datenbank.html>

Table A – 18 Overview over stock and operating hours of non-road vehicles (FOEN 2015j):

Upper table: Number of vehicles,

middle table Specific operating hours per year

lower table: Total operating hours per year (in million hours)

Category	1980	1990	2000	2010	2020	2030
	number of vehicles					
Construction machinery	63'364	58'816	52'729	57'102	60'384	62,726
Industrial machinery	26'714	43'244	70'671	69'786	69'757	70,083
Agricultural machinery	292'773	324'567	337'869	318'876	309'825	305,235
Forestry machinery	11'815	13'844	13'055	11'857	10'831	10,170
Garden-care / hobby appliances	1'198'841	1'539'624	1'944'373	2'322'737	2'464'323	2,499,627
Navigation machinery	94'866	103'383	93'912	95'055	97'522	99,104
Railway machinery	529	1'300	1'255	697	640	640
Military machinery	13'092	13'373	14'272	13'083	12'853	12,856

Category	1980	1990	2000	2010	2020	2030
	Specific operating hours per year					
Construction machinery	247	322	406	417	424	429
Industrial machinery	666	670	684	680	675	671
Agricultural machinery	136	119	112	103	99	95
Forestry machinery	203	199	203	193	188	182
Garden-care / hobby appliances	12	17	20	64	77	81
Navigation machinery	39	38	38	36	35	35
Railway machinery	877	613	617	783	719	719

Category	1980	1990	2000	2010	2020	2030
	million operating hours per year					
Construction machinery	15.7	19.0	21.4	23.8	25.6	26.9
Industrial machinery	17.8	29.0	48.4	47.5	47.1	47.0
Agricultural machinery	39.9	38.8	37.7	33.0	30.6	29.0
Forestry machinery	2.4	2.8	2.6	2.3	2.0	1.9
Garden-care / hobby appliances	14.6	25.7	39.3	149.7	190.8	201.3
Navigation machinery	3.7	3.9	3.5	3.4	3.4	3.4
Railway machinery	0.5	0.8	0.8	0.5	0.5	0.5
Military machinery	0.8	0.9	0.9	0.9	0.9	0.9
Total	95	121	155	261	301	311

A3.1.5 Sulphur dioxide (SO₂)

Table A – 19 shows sulphur contents and SO₂ emission factors per fuel type. Explanations:

- For liquid and solid fuels the SO₂ emission factors are determined by the sulphur content. The upmost lines in Table A – 19 “maximum legal limit on sulphur content” show the maximum values as defined in the Federal Ordinance on Air Pollution Control OAPC (Swiss Confederation 1985).
- The lines in the middle part of Table A – 19 contain the effective sulphur contents. They are based on measurements: Summary and annual reports of the Swiss Petroleum Association (EV), reports by the Federal Customs Administration (FCA) since 2000.
- The lines at the bottom part of Table A – 19 give the emission factors in kg/TJ. They are calculated from the sulphur content *S*, the net calorific value NCV and the quotient of the molar masses of S and SO₂

$$EF_{SO_2} = \frac{M_{SO_2}}{M_S} \cdot \frac{S}{NCV} = 2 \cdot \frac{S}{NCV}$$

- Coal: The legal limit of sulphur content depends on the size of the heat capacity of the combustion system. The value shown in the table above (1%, 350 kg/TJ SO₂) holds for heat capacity below 1 MW; see OAPC Annex 4, §513 (Swiss Confederation 1985). For larger capacities the value is 3% (OAPC Annex 5, §2, Swiss Confederation 1985). For industrial combustion plants, the limit for the exhaust emissions actually sets the corresponding maximum sulphur content to 1.4% (500 kg/TJ).
- Residual fuel oil: OAPC Annex 5, §11, lit.2 sets 2.8% for the legal limit. Simultaneously, OAPC dispenses from emission control measurements if residual fuel oil is used with sulphur content of maximum 1% (see OAPC Annex 3, §421, lit. 2, Swiss Confederation 1985), which holds for most combustion plants.

Table A – 19 Sulphur content and SO₂ emission factors.

year	maximum legal limit of sulphur content					
	Diesel oil ppm	Gasoline ppm	Gas oil ppm	Natural gas ppm	Res. fuel oil %	Coal %
1990	1400	200	2000	190	1.0	1.0
1991	1300	200	2000	190	1.0	1.0
1992	1200	200	2000	190	1.0	1.0
1993	1000	200	2000	190	1.0	1.0
1994	500	200	2000	190	1.0	1.0
2000	350	150	2000	190	1.0	1.0
2005	50	50	2000	190	1.0	1.0
2008	50	50	1000	190	1.0	1.0
2009	10	50	1000	190	1.0	1.0
2010-2015	10	10	1000	190	1.0	1.0

year	Effective sulphur content		
	Diesel oil ppm	Gasoline ppm	Gas oil ppm
1990	1400	200	1600
1991	1300	200	1300
1992	1200	200	1200
1993	1000	200	1000
1994	434	200	1350
1995	341	200	1170
1996	372	200	1160
1997	353	200	1250
1998	402	200	926
1999	443	200	650
2000	272	142	680
2001	250	121	830
2002	235	101	798
2003	200	81	700
2004	10	8.0	700
2005	10	8.0	799
2006	10	8.0	699
2007	10	8.0	630
2008	10	8.0	641
2009	7.2	5.2	603
2010	9	6	548
2011	5	8	116
2012	7	6	617
2013	8	5	253
2014	7	3	385
2015	7	3	384

year	Effective SO ₂ emission factor							
	Diesel oil (average in road transportation)	Gasoline	Gas oil	Natural gas	Res. fuel oil	Lignite	Bituminous coal Boiler 1A1, 1A2, 1A3 / Boiler 1A4	Kerosene (average)
kg/TJ								
1990	65.4	9.4	75.1	0.5	473.2	500.0	500/350	25.6
1991	60.8	9.4	61.0	0.5	432.0	500.0	500/351	25.1
1992	56.1	9.5	56.3	0.5	417.5	500.0	500/352	25.3
1993	46.8	9.5	46.9	0.5	422.3	500.0	500/353	25.3
1994	20.3	9.5	63.4	0.5	374.3	500.0	500/354	25.2
1995	15.9	9.3	54.9	0.5	377.2	500.0	500/355	25.3
1996	17.3	9.2	54.5	0.5	378.6	500.0	500/356	25.2
1997	16.4	9.3	58.7	0.5	339.8	500.0	500/357	25.0
1998	18.7	9.3	43.5	0.5	402.9	500.0	500/358	24.9
1999	20.6	9.3	30.5	0.5	301.0	500.0	500/359	25.0
2000	12.6	6.6	31.9	0.5	320.4	500.0	500/360	24.9
2001	11.6	5.7	39.0	0.5	398.1	500.0	500/361	24.3
2002	10.9	4.7	37.5	0.5	398.1	500.0	500/362	23.6
2003	9.3	3.7	32.9	0.5	383.5	500.0	500/363	23.2
2004	0.5	0.4	32.9	0.5	368.9	500.0	500/364	23.2
2005	0.5	0.4	37.5	0.5	378.6	500.0	500/365	22.8
2006	0.5	0.4	32.8	0.5	361.2	500.0	500/366	21.4
2007	0.5	0.4	29.6	0.5	343.7	500.0	500/367	21.6
2008	0.5	0.4	30.1	0.5	326.2	500.0	500/368	21.5
2009	0.5	0.4	25.3	0.5	308.7	500.0	500/369	21.4
2010	0.5	0.4	25.7	0.5	291.3	500.0	500/370	21.4
2011	0.5	0.4	24.1	0.5	291.1	500.0	500/371	21.5
2012	0.5	0.4	22.4	0.5	291.0	500.0	500/372	21.6
2013	0.5	0.4	20.8	0.5	290.9	500.0	500/373	21.7
2014	0.5	0.4	19.2	0.5	290.9	500.0	500/374	21.7
2015	0.5	0.4	17.6	0.5	290.9	500.0	500/375	21.7

A3.2 Industrial processes and product use (illustrative example of mobile air conditioning)

The use of HFC as substitutes of ODS in 2F1 refrigeration and air conditioning is the main factor for the increase of HFC emission 1990 to 2015. Refrigerants contained in installed equipment lead to a considerable stock with annual losses depending from equipment type between 0.5% to 20% (see Table 4-44). Emissions are calculated for the production, operation, service and disposal of equipment. The following illustrative example shows the calculations for the example of mobile air conditioning (HFC-134a use as refrigerant). The example is calculated bottom up based on vehicle statistics and informations on air conditioning equipment. There is no production of air conditioning equipment for cars in Switzerland, equipment is imported already charged.

Table A – 20 Applied model parameters and assumption for mobile air conditioning of cars

Characteristic values			
Initial charge in kg HFC per unit AC	1994	0.8	kg
	2002	0.7	kg
	2014	0.6	kg
	Extrapolation of other years		
Lifetime		15	years
Production			
Import of precharged equipment		100	%
Operation			
Annual losses		8.5	%
Recharge of losses (6.8% of 8.5%)		70	%
Additional service losses over lifetime		10	%
Disposal			
Export rate		31-72	%
Share with total loss of refrigerant		40	%
Disposal loss of professional recovery		15	%
Assumed reuse of recovered chemical		80	%

Table A – 21 Bottom up calculations to identify number of air conditioning equipment and amount of HFC-134a

Year	New registered vehicles	Vehicles in use	Disposed vehicles	New equipment: number of air conditioning units with HFC-134a in new registered cars			Equipment stock: Number of air conditioning units with HFC-134a in use		Equipment disposal	Initial equipment charge
				Portion of vehicles with AC [%]	HFC-134a as refrigerant [%]	AC units with HFC-134 [units]	Portion of vehicles with HFC-134a [%]	AC units with HFC-134 [units]		
1989	335'094	2'895'842		5	0	0	0	0	0	0.85
1990	327'456	2'985'399	237'899	6	0	0	0	0	0	0.84
1991	314'824	3'057'800	242'423	7	10	2'204	0	2'204	0	0.83
1992	296'009	3'091'230	262'579	9	30	7'992	0	10'196	0	0.83
1993	262'814	3'109'524	244'520	14	66	24'284	1	34'480	0	0.82
1994	270'009	3'165'043	214'490	19	90	46'172	3	80'652	0	0.81
1995	272'897	3'229'169	208'771	24	100	65'495	5	146'147	0	0.78
1996	269'529	3'268'073	230'625	38	100	102'421	8	248'568	0	0.77
1997	272'441	3'323'421	217'093	52	100	141'669	12	390'237	0	0.76
1998	297'336	3'383'275	237'482	68	100	202'188	18	592'426	0	0.75
1999	317'985	3'467'275	233'985	75	100	238'489	24	830'914	0	0.73
2000	315'398	3'545'247	237'426	77	100	242'856	30	1'073'771	0	0.72
2001	317'126	3'629'713	232'660	85	100	269'557	37	1'343'328	0	0.71
2002	295'109	3'704'822	220'000	87	100	256'745	43	1'600'073	0	0.70
2003	271'541	3'754'000	222'363	89	100	241'671	49	1'841'744	0	0.69
2004	269'211	3'811'351	211'860	91	100	244'982	55	2'086'726	0	0.68
2005	259'426	3'863'807	206'970	92	100	238'672	60	2'325'398	0	0.66
2006	269'421	3'899'917	233'311	96	100	258'644	66	2'581'839	2'204	0.65
2007	284'674	3'955'787	228'804	96	100	273'287	72	2'847'133	7'992	0.64
2008	288'525	4'030'965	213'347	96	100	276'984	77	3'099'833	24'284	0.63
2009	266'018	4'051'569	245'414	96	100	255'377	82	3'309'039	46'172	0.61
2010	294'239	4'119'370	226'438	96	100	282'469	86	3'526'013	65'495	0.60
2011	327'896	4'209'300	237'966	96	100	314'780	90	3'738'372	102'421	0.59
2012	328'139	4'254'725	282'714	96	100	315'013	92	3'911'717	141'669	0.58
2013	310'154	4'320'885	243'994	96	92	273'928	92	3'983'456	202'188	0.56
2014	304'083	4'384'490	240'478	96	85	248'132	91	3'993'099	238'489	0.55
2015	327'143	4'458'069	253'564	96	77	241'510	90	3'991'753	242'856	0.55

Table A – 22 Results and structure of emission calculations of HFC-134a from mobile air conditioning of cars for the inventories 1990 to 2015.

HFC-134a	Activity			Emissions				Recharge	
	Input with vehicles	Stock	Disposed	Production	Stock incl. Recharge	Disposal	Total	import in bulk	recovered and reused
	[t]	[t]	[t]	[t]	[t]	[t]	[t]	[t]	[t]
1990	0	0	0	NO	0	0	0	0	0
1991	2	2	0	NO	0	0	0	0	0
1992	7	8	0	NO	1	0	1	0	0
1993	20	28	0	NO	3	0	3	1	0
1994	37	65	0	NO	6	0	6	2	0
1995	51	114	0	NO	10	0	10	4	0
1996	79	190	0	NO	17	0	17	7	0
1997	107	292	0	NO	27	0	27	12	0
1998	151	434	0	NO	40	0	40	19	0
1999	175	597	0	NO	55	0	55	28	0
2000	175	756	0	NO	69	0	69	38	0
2001	191	927	0	NO	85	0	85	49	0
2002	180	1'081	0	NO	99	0	99	59	0
2003	166	1'217	0	NO	112	0	112	69	0
2004	165	1'349	0	NO	124	0	124	78	0
2005	158	1'470	0	NO	135	0	135	87	0
2006	168	1'597	1	NO	146	0	147	94	0
2007	174	1'724	3	NO	158	0	158	101	2
2008	173	1'841	9	NO	169	3	171	107	5
2009	156	1'927	20	NO	177	6	183	110	10
2010	169	2'019	25	NO	185	7	192	114	13
2011	185	2'108	40	NO	193	11	205	113	21
2012	181	2'167	64	NO	199	15	213	109	33
2013	155	2'190	72	NO	201	19	220	110	36
2014	136	2'189	77	NO	201	21	222	109	39
2015	133	2'181	82	NO	200	22	222	107	42

A3.3 Agriculture

Additional data for estimating CH₄ emission from 3A Enteric fermentation

Table A – 23 Data for estimating enteric fermentation emission factors for cattle (Table according to outline in IPCC 1997c, p 4.31 – 4.33).

Type	Age ^a	Weight ^a kg	Weight Gain ^a kg/day	Feeding Situation / Further Specification ^a	Milk ^b kg/day	Work hrs/day	Pregnant ^a % 305 days of lactation	Digestibility of Feed % ^d	Y _m ^d %	Em. Factor kg/head/year ^e
Mature Dairy Cattle	NA	650	0		16.1-23.1 ^c	0		72	6.90	117.3 - 137.4
Other Mature Cattle	NA	650	0		8.2	0		60	6.50	106.8
Fattening Calves	0-98 days	124	1.43	Rations of unskimmed milk and supplementary milk feed when life weight exceeds 100 kg. Rations are apportioned on two servings per day.	0	0	0	65	0.00	0.0
Pre-Weaned Calves	0-300 days	195	0.88	"Natura beef" production, milk from mother cow and additional feed.	0	0	0	65	6.50	16.3
Breeding Calves	0-105 days	85	0.67	Feeding plan for a dismission with 14 to 15 weeks. Milk feed concentrate (100kg in total), hay (80 kg in total).	0	0	0	65	6.50	30.0
Breeding Cattle (4-12 months)	4-12 month	210	0.80	Premature race (Milk-race)	0	0	0	60	6.50	
Breeding Cattle (> 1 year)	12-28/30 month	450	0.80	Premature race (Milk-race)	0	0	0	60	6.50	61.2
Fattening Calves (0-4 months)	0-132 days	115	0.83	Diet based on milk or milk-powder and feed concentrate, hay and/or silage	0	0	0	65	6.50	
Fattening Cattle (4-12 months)	4-12 month	361	1.37	Feeding recommendations for fattening steers, concentrate based	0	0	0	60	6.50	43.2

^a Data source: RAP 1999 and calculations according to Soliva 2006.

^b Milk production in kg/day is calculated by dividing the average annual milk production per head by 305 days (lactation period).

^c data source: Swiss farmers union (MISTA 2015).

^d data source: IPCC 2006 and Zeliz et al. 2012.

^e For better comparability emission factors of young cattle were converted to kg/head/year although the time span of most of the individual categories is less than 365 days.

Table A – 24 Gross energy intake of Swiss livestock 1990–2015.

Gross Energy Intake	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
	MJ/head/day																										
Cattle																											
Mature Dairy Cattle	259.3	261.6	261.9	265.0	264.7	267.4	266.4	270.4	274.2	277.4	280.1	281.8	284.5	287.7	291.5	291.6	292.2	295.2	297.2	298.8	300.7	301.4	301.2	301.2	299.8	302.5	303.6
Other Mature Cattle	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6	250.6
Growing Cattle (weighted average)	101.0	101.1	101.3	100.8	101.3	101.5	101.2	101.7	100.5	98.8	101.0	99.8	98.4	98.0	98.4	97.7	97.7	97.4	97.3	97.0	96.8	96.6	96.1	96.1	96.0	96.0	96.4
Fattening Calves	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1	47.1
Pre-Weaned Calves	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1	38.1
Breeding Calves	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8
Breeding Cattle (>1 year)	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	90.1	
Breeding Cattle (>1 year)	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	143.6	
Fattening Calves (<4 months)	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	
Fattening Cattle (<4 months)	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	126.3	
Fattening Cattle (<4 months)	21.2	21.7	22.2	22.4	23.8	24.0	22.0	22.1	21.5	22.9	22.4	22.9	22.8	22.7	23.3	22.8	22.6	22.2	22.0	22.7	22.6	22.6	22.5	22.4	22.5	22.5	
Sheep	28.3	28.9	29.0	29.1	28.5	31.9	29.8	29.9	27.9	29.0	28.0	27.7	27.0	27.2	26.6	26.3	26.9	26.7	27.0	27.2	26.9	26.5	27.3	27.4	27.4	27.4	
Swine	NA	NA	134.7	138.2	137.5	136.6	137.0	136.5	137.0	130.6	146.9	137.5	140.8	137.2	140.0	140.6	138.9	130.4	129.1	134.8	136.9	139.8	135.9	138.0	134.6	134.5	
Buffalo (weighted average)	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	91.0	
Bisons < 3 years	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	192.1	
Bisons > 3 years	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
Camels (weighted average)	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	24.8	
Llamas < 2 years	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	43.0	
Llamas > 2 years	16.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	15.2	
Alpacas < 2 years	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	27.8	
Alpacas > 2 years	50.5	51.6	53.6	54.5	55.0	55.3	59.7	55.4	54.8	55.5	56.4	55.8	56.7	55.2	55.6	55.4	55.8	55.9	56.5	56.8	56.5	56.7	57.0	58.1	58.0	58.0	
Deer (weighted average)	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	50.5	
Fallow Deer ¹⁾	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	101.1	
Red Deer ²⁾	25.0	24.6	25.0	25.4	25.5	27.9	25.3	25.6	26.9	25.8	25.7	26.0	25.2	25.4	25.2	25.4	25.3	25.0	25.0	25.3	25.1	25.6	25.6	25.6	25.0	25.0	
Horses (weighted average)	107.3	107.3	107.3	107.3	107.1	106.9	107.1	107.3	107.3	107.3	107.2	107.4	107.5	107.6	107.6	107.7	107.7	107.7	107.7	107.8	107.9	107.9	107.9	107.9	108.0	108.3	
Horses < 3 years	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	
Horses > 3 years	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	
Mules and Asses (weighted average)	39.2	39.2	39.2	39.2	39.5	39.7	39.7	39.8	39.6	39.8	39.5	39.5	39.5	39.5	39.5	39.4	39.5	39.3	39.2	39.2	40.0	40.2	39.9	39.8	39.6	39.6	
Mules	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	
Asses	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	
Poultry ²⁾	1.5	1.5	1.6	1.3	1.4	1.3	1.4	1.4	1.3	1.4	1.4	1.4	1.4	1.3	1.3	1.3	1.2	1.2	1.3	1.3	1.2	1.2	1.2	1.2	1.2	1.2	
Rabbits	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	
Livestock NCAG (weighted average)	95.6	95.6	95.6	95.6	96.1	93.4	43.2	40.1	38.6	38.6	37.7	36.5	35.7	34.6	34.6	33.3	33.2	32.9	31.7	34.3	36.9	37.2	37.9	38.2	38.5	38.7	
Sheep Non-Agr.	21.2	21.7	22.2	22.4	23.8	24.0	22.0	22.1	21.5	22.9	22.4	22.9	22.8	22.7	23.3	22.8	22.6	22.2	22.0	22.7	22.6	22.6	22.6	22.6	22.5	22.5	
Goats Non-Agr.	25.0	24.6	25.0	25.4	25.5	27.9	25.3	25.6	26.9	25.8	25.7	26.0	25.2	25.4	25.2	25.4	25.3	25.0	25.0	25.3	25.1	25.6	25.6	25.6	25.0	25.0	
Horses < 3 years Non-Agr.	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	101.4	
Horses > 3 years Non-Agr.	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	109.0	
Mules Non-Agr.	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	86.0	
Asses Non-Agr.	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	37.9	

¹⁾ Deer: Gross energy intake per animal (see: review web offering)
²⁾ Poultry data not Gross Energy Intake (GEI) but Metabolizable Energy Intake (MEI)

Table A – 25 Livestock population 1990–2015.

Population Size	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
Cattle	1855.2	1828.9	1782.6	1745.1	1755.4	1748.3	1747.1	1672.9	1640.9	1608.7	1588.0	1611.4	1597.7	1570.2	1544.5	1554.7	1566.9	1571.8	1604.3	1597.5	1591.2	1574.4	1564.6	1567.5	1562.8	1564.3	1564.3
Mature Dairy Cattle	783.1	780.5	763.5	744.5	749.7	739.6	736.0	711.6	701.3	683.5	669.4	669.4	657.9	638.3	621.0	620.7	618.1	614.8	628.5	628.5	599.0	589.2	589.2	591.2	586.6	587.4	583.3
Other Mature Cattle	12.0	14.0	17.0	18.0	20.0	23.0	28.0	32.0	36.0	41.2	44.9	50.6	58.1	65.1	70.0	78.5	87.3	93.5	98.4	108.4	111.3	110.7	114.4	118.9	118.0	117.9	117.9
Other Mature Cattle	1060.1	1034.4	1002.1	982.6	985.7	985.6	983.0	929.3	903.5	884.0	873.7	891.3	877.7	866.7	853.6	855.5	861.5	863.4	887.4	889.7	880.9	877.5	889.0	894.0	854.0	857.4	853.1
Calving Cattle	112.3	113.4	109.5	111.1	101.4	101.7	112.0	106.0	108.1	116.4	103.3	114.7	114.4	113.9	111.3	105.6	101.2	100.5	95.0	100.5	99.4	100.5	96.6	97.0	96.7	90.9	90.9
Pre-Weaned Calves	9.6	11.2	13.6	14.4	16.0	18.4	22.4	25.6	28.8	33.2	35.7	40.4	46.9	52.3	56.6	62.5	67.3	72.2	76.1	85.8	88.1	87.6	90.6	92.5	93.4	93.4	93.3
Breeding Cattle 1st Year	346.4	336.7	324.0	308.2	306.2	294.7	286.1	260.1	253.5	218.7	236.0	238.1	229.5	219.8	214.7	222.0	223.3	223.3	232.4	232.4	226.4	220.7	212.3	212.3	211.3	211.3	211.3
Breeding Cattle 2nd Year	253.3	251.9	250.5	238.7	237.2	236.6	243.0	232.9	217.4	187.5	221.9	219.3	212.7	212.7	205.4	204.7	210.2	210.5	212.7	212.7	212.8	207.5	199.5	197.7	198.6	198.6	198.6
Breeding Cattle 3rd Year	150.7	148.4	146.7	142.3	141.3	139.4	139.3	132.7	117.9	129.8	130.4	126.0	124.0	120.9	115.3	110.1	109.1	109.1	109.6	118.8	119.2	116.2	111.8	111.8	111.2	111.2	111.8
Fattening Cattle	187.8	174.8	167.8	168.0	183.5	192.9	178.9	165.4	163.1	210.2	147.1	148.5	141.7	144.1	144.7	147.5	146.6	145.1	151.6	146.6	145.1	145.0	146.2	146.2	145.7	146.1	145.3
Fattening Cattle	395.2	409.4	414.7	424.0	405.4	386.7	418.6	420.4	422.3	423.5	420.7	420.0	429.5	444.8	440.5	446.4	447.5	443.6	446.2	443.9	434.1	424.0	417.3	409.5	402.8	404.0	347.0
Fattening Sheep	190.6	200.8	201.0	211.1	201.2	191.4	207.6	208.0	208.7	221.7	216.6	216.6	219.9	228.6	227.5	228.4	230.6	230.0	229.4	223.9	228.2	221.8	219.3	216.2	208.5	204.0	204.0
Milk Sheep	4.3	4.0	3.8	3.5	3.3	3.0	2.8	3.1	4.4	5.8	6.7	7.2	8.0	8.1	8.9	9.5	10.2	10.7	11.7	12.4	12.4	12.4	12.8	13.3	13.7	13.6	13.6
Piglets	299.4	282.5	290.8	299.6	297.2	274.8	240.9	252.2	261.8	281.0	296.6	316.8	326.6	322.8	327.8	337.6	366.5	344.8	336.1	338.4	350.9	352.7	344.7	333.3	332.2	330.0	330.0
Fattening Pig over 25 kg	1024.6	989.6	972.8	943.0	855.5	787.9	778.7	779.6	837.4	734.4	750.9	762.5	767.9	751.7	753.2	786.7	786.1	763.9	763.2	778.5	788.1	787.2	775.5	748.1	750.8	777.3	777.3
Dry Sows	129.3	126.0	124.9	125.3	117.1	108.9	98.8	104.3	110.9	101.2	104.8	108.0	108.6	105.3	107.9	112.7	105.7	105.4	104.7	106.1	103.4	103.4	97.4	95.8	94.2	93.3	93.3
Nursing Sows	37.4	36.8	36.8	37.3	35.1	33.0	30.2	29.9	31.4	35.0	36.7	37.5	36.5	36.8	35.3	36.3	36.5	34.9	32.6	33.1	33.5	32.3	31.0	29.4	29.4	29.4	29.4
Bovars	8.4	8.1	8.0	8.2	7.7	7.1	6.3	6.4	6.4	6.2	6.2	6.2	5.8	5.3	5.2	5.1	4.9	4.2	4.0	3.8	3.7	3.3	3.0	3.2	2.9	2.7	2.7
Buffalo	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.4	0.5	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.6
Bisons < 3 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Bisons > 3 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Camels	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Lamas < 2 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Lamas > 2 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Alpacas < 2 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Alpacas > 2 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Deer	0.2	0.4	0.7	0.9	1.2	1.4	1.9	1.9	2.2	2.6	2.8	2.9	3.1	3.2	3.5	3.8	4.2	4.4	4.8	5.1	5.5	5.7	5.7	5.7	5.7	5.7	6.0
Fallow Deer	0.2	0.4	0.6	0.8	1.1	1.3	1.6	1.8	2.0	2.4	2.5	2.6	2.7	2.9	3.2	3.5	3.7	4.0	4.4	4.9	5.0	5.0	4.9	4.9	4.9	5.1	5.1
Red Deer	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.6	0.7	0.7	0.7	0.8	0.8	0.9	0.9
Goats	68.3	65.2	58.2	56.7	54.9	53.2	56.8	58.0	60.1	61.6	62.5	63.0	66.0	67.4	70.6	74.0	76.3	78.1	81.4	81.2	82.8	83.0	84.7	84.5	84.7	84.5	84.7
Goat Places	44.8	43.1	38.4	37.3	35.9	34.6	37.1	37.7	39.8	40.8	41.4	42.1	43.0	44.9	46.2	48.5	50.5	51.9	53.4	54.3	54.7	55.9	57.4	57.4	57.9	56.1	56.1
Horses	28.2	30.2	32.3	34.5	37.9	41.4	43.0	45.8	46.3	48.5	50.3	51.1	52.7	53.7	55.1	56.4	57.7	59.0	60.2	62.1	62.1	57.2	58.0	57.2	57.2	55.5	55.5
Horses < 3 years	6.1	6.5	7.0	7.4	9.2	11.0	10.7	10.0	10.0	11.0	10.1	9.7	9.5	9.4	9.4	9.4	9.5	9.6	9.6	9.6	9.6	8.7	8.3	8.0	7.1	6.5	4.7
Horses > 3 years	22.1	23.7	25.4	27.1	28.7	30.4	32.3	35.8	36.3	37.5	40.2	40.4	41.7	43.3	44.3	46.8	46.9	48.1	49.4	51.1	53.4	49.0	50.1	50.2	50.7	50.8	50.8
Mules and Asses	5.9	6.3	6.7	7.2	7.4	7.6	8.5	9.4	9.9	11.3	11.8	12.5	13.2	14.1	14.8	16.0	16.5	17.2	17.8	19.2	20.4	19.0	20.1	19.6	18.6	19.7	19.7
Mules	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Asses	5.7	6.1	6.6	7.0	7.1	7.3	8.1	9.0	9.6	10.9	11.4	12.0	12.8	13.6	14.4	15.5	15.9	16.7	17.3	18.4	19.4	16.2	19.3	18.9	18.9	19.0	
Poultry ¹⁾	5938.2	5646.8	5501.6	6409.8	6330.3	6250.7	6440.5	6552.5	6739.6	6907.5	6938.0	6938.0	7338.6	7597.3	8060.7	8260.4	8228.5	8542.8	8808.4	9024.9	9477.7	9584.6	10079.0	10723.4	10824.6	10824.6	
Layers	718.9	664.2	709.6	719.2	716.8	714.4	732.1	732.9	793.5	760.9	831.7	745.3	753.9	809.0	853.1	867.7	888.4	901.8	919.0	966.7	925.5	969.7	1075.6	1054.5	1195.6	1195.6	
Broilers	2019.9	2198.6	2095.5	2990.2	3110.6	3231.0	3283.2	3342.0	3502.3	3747.4	3607.8	3993.2	4298.2	4519.4	4970.8	5060.4	4481.3	5002.4	5456.2	5984.1	6282.1	5984.1	6282.1	6580.3	6763.7	6897.8	
Turkey	94.7	117.4	140.1	162.8	166.5	170.2	173.8	184.4	157.8	154.6	172.6	123.0	123.9	133.8	138.8	132.3	137.1	112.5	53.8	52.4	58.1	57.8	51.2	55.1	57.2	49.3	
Other Poultry	21.8	21.2	20.6	20.0	18.4	16.9																					

Additional data for estimating CH₄ and N₂O emission from 3B Manure management

Table A – 26 Data for estimating manure management CH₄ emission factors (Table according to outline in IPCC 1997c, Tables B-1 to B-7).

Type	Weight kg ^a	Digestibility of Feed % ^b	Energy Intake MJ/day	Feed Intake kg/day	% Ash Dry Basis ^b	VS kg/head/day	B ₀ m ³ CH ₄ /kg VS ^b
Mature Dairy Cattle	650	72	259 - 304	15.89 ^c	8.8 - 9.1	4.08 - 4.82	0.24
Other Mature Cattle	650	60	251	10.96 ^c	8	5.50	0.18
Fattening Calves	124	65	47	2.02 ^a	8	0.92	0.18
Pre-Weaned Calves	195	65	38	2.99 ^a	8	0.74	0.18
Breeding Calves	85	65	28	2.19 ^a	8	0.54	0.18
Breeding Cattle (4-12 months)	210	60	90	4.88 ^a	8	1.98	0.18
Breeding Cattle (> 1 year)	450	60	144	7.78 ^a	8	3.15	0.18
Fattening Calves (0-4 months)	115	65	50	3.00 ^a	8	0.97	0.18
Fattening Cattle (4-12 months)	361	60	126	6.84 ^a	8	2.77	0.18
Sheep	NA	60	21 - 24	1.09-1.41 ^c	8	0.40 ^b	0.19
Swine	NA	75	26 - 32	NA	2	0.31 ^b	0.45
Buffalo	NA	55	129 - 147	7.00-7.96 ^c	8	3.29	0.10
Camels	NA	60	31 - 38	1.68-2.05 ^c	8	0.69	0.26
Deer	NA	60	51 - 60	2.74-3.24 ^c	8	1.27	0.19
Goats	NA	60	25 - 28	1.21-1.47 ^c	8	0.30 ^b	0.18
Horses	NA	70	107 - 108	7.73-7.88 ^c	4	1.9 ^b	0.33
Mules and Asses	NA	70	39 - 40	NA	4	0.94 ^b	0.33
Poultry	NA	NA	1.2 - 1.6 ^d	NA	NA	0.01 ^b	0.37
Rabbits	NA	NA	1.2	NA	NA	0.10	0.32
Livestock NCAC	NA	NA	NA	NA	NA	0.68	0.27

^a RAP 1999

^b IPCC 1997c and IPCC 2006

^c Flisch et al. 2009

^d metabolizable energy (ME)

Table A – 27 Manure management system distribution in Switzerland 1990–2015.

MS Distribution	1990					2002					2007					2010					2015										
	Liquid / Slurry	Solid manure	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid manure	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid manure	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid manure	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)	Liquid / Slurry	Solid manure	Pasture range and paddock	Digesters	Other (Deep litter, Poultry manure)						
Mature Dairy Cattle	63.7	27.6	8.3	0.4	0.0	65.7	24.4	9.5	0.4	0.0	63.3	16.2	18.0	0.5	0.0	67.5	13.6	17.7	1.3	0.0	67.0	14.3	16.9	1.7	0.0	69.8	10.7	16.2	3.3	0.0	
Other Mature Cattle	41.2	32.1	28.3	0.4	0.0	39.8	34.1	28.2	0.4	0.0	38.8	20.7	39.1	0.5	0.0	49.5	20.2	28.3	1.3	0.0	47.9	17.9	32.4	1.7	0.0	50.6	14.7	31.4	3.3	0.0	
Growing Cattle (weighted average)	47.5	31.7	15.8	0.4	4.5	48.4	30.9	15.9	0.4	4.5	42.1	25.4	27.5	0.5	4.5	45.6	23.9	25.3	1.3	4.0	45.2	25.6	23.4	1.7	4.1	46.9	22.6	23.7	3.3	3.5	
Featuring Calves	14.3	0.0	0.0	0.4	85.2	15.0	0.0	0.0	0.4	84.6	21.6	0.0	0.3	0.5	77.6	21.8	0.0	0.2	1.3	76.7	16.9	0.0	0.2	1.7	81.1	0.0	0.0	1.2	3.3	73.4	
Pre-Weaned Calves	41.2	32.1	26.3	0.4	0.0	39.3	34.1	26.2	0.4	0.0	41.2	21.1	37.3	0.5	0.0	50.0	18.6	30.1	1.3	0.0	44.6	32.8	20.9	1.7	0.0	35.8	27.7	32.3	3.3	0.0	
Breeding Cattle 1st Year	37.0	48.5	14.1	0.4	0.0	38.0	47.4	14.2	0.4	0.0	33.7	38.8	27.0	0.5	0.0	41.0	34.5	23.3	1.3	0.0	43.4	33.4	21.5	1.7	0.0	44.7	30.9	21.1	3.3	0.0	
Breeding Cattle 2nd Year	46.3	26.9	25.4	0.4	0.0	47.3	26.7	25.6	0.4	0.0	37.8	23.4	38.4	0.5	0.0	41.4	20.8	36.5	1.3	0.0	43.2	20.8	30.4	1.7	0.0	42.3	19.5	34.8	3.3	0.0	
Breeding Cattle 3rd Year	50.5	29.1	20.0	0.4	0.0	51.4	27.9	20.3	0.4	0.0	42.2	22.5	34.8	0.5	0.0	45.6	21.3	31.8	1.3	0.0	46.3	20.8	30.6	1.7	0.0	54.1	17.0	25.5	3.3	0.0	
Featuring Cattle	70.1	24.1	0.0	0.4	5.5	66.4	27.7	0.0	0.4	5.6	67.3	23.8	2.2	0.5	3.2	62.3	29.2	4.3	1.3	2.9	57.7	33.2	4.0	1.7	3.4	61.4	27.7	4.8	3.3	2.8	
Sheep (weighted average)	0.0	0.0	30.1	0.0	69.9	0.0	0.0	30.3	0.0	69.7	0.0	0.0	33.2	0.0	66.8	0.0	0.0	39.2	0.0	60.8	0.0	0.0	33.7	0.0	66.3	0.0	0.0	33.6	0.0	66.4	
Featuring Sheep	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Milksheep	0.0	0.0	11.4	0.0	88.6	0.0	0.0	11.4	0.0	88.6	0.0	0.0	26.1	0.0	73.9	0.0	0.0	24.1	0.0	75.9	0.0	0.0	22.8	0.0	77.2	0.0	0.0	40.9	0.0	59.1	
Pigs	98.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	98.0	0.3	0.1	1.5	0.0	94.5	0.1	1.2	4.1	0.0	94.0	0.3	0.1	5.6	0.0	88.7	0.0	0.0	11.2	0.0	
Featuring Pig over 25 kg	96.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	97.6	0.8	0.0	1.5	0.0	94.8	0.7	0.4	4.1	0.0	92.0	2.3	0.0	0.2	5.6	0.0	86.8	0.0	0.0	11.2	0.0
Dry Sows	98.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	98.4	0.0	0.1	1.5	0.0	94.8	0.1	1.0	4.1	0.0	94.2	0.0	0.2	5.6	0.0	88.7	0.0	0.1	11.2	0.0	
Nursing Sows	96.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	97.8	0.7	0.0	1.5	0.0	95.0	0.5	0.3	4.1	0.0	94.2	0.2	0.0	94.2	0.2	86.8	0.0	0.0	11.2	0.0	
Boars	96.8	0.0	0.0	1.2	0.0	98.7	0.0	0.0	1.3	0.0	97.7	0.5	0.2	1.5	0.0	94.7	0.0	1.2	4.1	0.0	92.5	1.2	0.6	5.6	0.0	89.5	0.0	0.2	11.2	0.0	
Buffalo (weighted average)	NA	NA	NA	NA	NA	47.5	26.8	25.6	0.0	0.0	38.1	23.5	38.4	0.0	0.0	42.3	21.1	36.5	0.0	0.0	44.5	21.2	34.3	0.0	0.0	44.7	20.5	34.8	0.0	0.0	
Bisons < 3 years	46.6	29.0	26.4	0.0	0.0	47.5	26.8	25.6	0.0	0.0	38.1	23.5	38.4	0.0	0.0	42.3	21.1	36.5	0.0	0.0	44.5	21.2	34.3	0.0	0.0	44.7	20.5	34.8	0.0	0.0	
Bisons > 3 years	46.6	29.0	26.4	0.0	0.0	47.5	26.8	25.6	0.0	0.0	38.1	23.5	38.4	0.0	0.0	42.3	21.1	36.5	0.0	0.0	44.5	21.2	34.3	0.0	0.0	44.7	20.5	34.8	0.0	0.0	
Camels (weighted average)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
Llamas < 2 years	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Llamas > 2 years	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Alpacas < 2 years	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Alpacas > 2 years	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Deer (weighted average)	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Fallow Deer	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Red Deer	0.0	0.0	30.7	0.0	69.3	0.0	0.0	30.7	0.0	69.3	0.0	0.0	33.5	0.0	66.5	0.0	0.0	40.2	0.0	59.8	0.0	0.0	34.5	0.0	65.5	0.0	0.0	33.0	0.0	67.0	
Goats	0.0	0.0	13.6	0.0	86.4	0.0	0.0	13.6	0.0	86.4	0.0	0.0	12.2	0.0	87.8	0.0	0.0	7.1	0.0	92.9	0.0	0.0	10.0	0.0	90.0	0.0	0.0	11.6	0.0	88.4	
Goat Places	0.0	0.0	13.6	0.0	86.4	0.0	0.0	13.6	0.0	86.4	0.0	0.0	12.2	0.0	87.8	0.0	0.0	7.1	0.0	92.9	0.0	0.0	10.0	0.0	90.0	0.0	0.0	11.6	0.0	88.4	
Horses (weighted average)	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Horses < 3 years	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Horses > 3 years	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Mules and Asses (weighted average)	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Mules	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Asses	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	6.8	0.0	0.0	0.0	93.2	
Poultry (weighted average)	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.5	0.0	99.5	0.0	0.0	2.6	0.0	97.4	0.0	0.0	3.7	0.0	96.3	0.0	0.0	0.0	2.7	0.0	97.3	0.0	0.0	3.0	0.0	
Growers	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.6	0.0	99.4	0.0	0.0	0.2	0.0	99.8	0.0	0.0	1.5	0.0	98.5	0.0	0.0	0.0	1.2	0.0	98.8	0.0	0.0	0.4	0.0	
Layers	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.6	0.0	99.4	0.0	0.0	0.2	0.0	99.8	0.0	0.0	1.5	0.0	98.5	0.0	0.0	0.0	1.2	0.0	98.8	0.0	0.0	0.4	0.0	
Broilers	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.4	0.0	99.6	0.0	0.0	0.6	0.0	99.4	0.0	0.0	1.2	0.0	98.8	0.0	0.0	0.0	0.3							

Additional data for estimating N₂O emissions from 3D Agricultural soils

Table A – 29 Additional data for estimating N₂O emission from crop residues (2015).

2015		Total crop production	Nitrogen incorporated with	N ₂ O emissions from crop
		kg DM	crop residues F _(CR)	residues
			t N	t N ₂ O
1. Cereals	Wheat	435'605'450	1'825	28.68
	Barley	168'165'700	857	13.47
	Maize	81'112'950	773	12.15
	Oats	6'606'200	41	0.64
	Rye	10'130'300	43	0.67
	Other:			
	Triticale	42'424'350	208	3.27
	Spelt	12'416'800	114	1.79
	Mix of Fodder Cereals	957'950	5	0.08
	Mix of Bread Cereals	112'200	0	0.01
	Millet	154'800	0	0.00
2. Pulse	Dry Beans	1'251'200	50	0.78
	Peas (Eiweisserbsen)	12'778'050	376	5.91
	Soybeans	3'445'900	142	2.23
	Leguminous Vegetables	2'779'238	403	6.34
	Lupines	273'680	11	0.17
3. Tuber and Root	Potatoes	78'256'000	286	4.49
	Other:			
	Fodder Beet	7'728'000	62	0.97
	Sugar Beet	298'256'200	2'684	42.18
5. Other	Fruit	48'109'150	192	3.02
	Grass	6'052'976'044	21'604	339.49
	Green Corn	96'037'795	44	0.69
	Non-Leguminous Vegetables	55'320'071	515	8.10
	Rape	78'303'600	1'143	17.97
	Renewable Energy Crops	1'224'000	18	0.28
	Silage Corn	564'928'205	324	5.10
	Sunflowers	8'314'700	176	2.77
	Tobacco	1'060'000	28	0.43
	Berries	2'578'800	15	0.24
	Vine	21'557'200	129	2.03
	Oil Squash	5'280	0	0.00
	Oil Hemp	23'400	1	0.02
	Oil Flax	267'300	2	0.04
	Hops	23'000	0	0.00
Medicinal Plants and Herbs	360'000	9	0.15	
Total Non-leguminous		2'020'039'401	9'497	149.24
Total Leguminous		20'528'068	982	15.43
Total excluding Grass		2'040'567'469	10'479	164.67
Total including Grass		8'093'543'512	32'083	504.17

Table A – 30 Additional data for estimating N₂O emission from crop residues (fractions)(2015).

2015		Residue/ Crop ratio kg/kg	Dry matter fraction of residue kg/kg	Nitrogen content of residues kg/kg
1. Cereals	Wheat	1.15	0.85	0.0037
	Barley	1.00	0.85	0.0051
	Maize	1.11	0.85	0.0086
	Oats	1.27	0.85	0.0049
	Rye	1.17	0.85	0.0036
	Other :			
	Triticale	1.25	0.85	0.0039
	Spelt	1.56	0.85	0.0059
	Mix of Fodder Cereals	1.00	0.85	0.0051
	Mix of Bread Cereals	1.17	0.85	0.0037
	Millet	0.20	0.90	0.0070
2. Pulse	Dry Beans	1.13	0.85	0.0353
	Peas (Eiweisserbsen)	1.25	0.85	0.0235
	Soybeans	1.00	0.85	0.0412
	Other:	3.87	0.16	0.0328
	Leguminous Vegetables	1.00	0.85	0.0412
	Lupines	0.47	0.13	0.0127
3. Tuber and Root	Potatoes			
	Other :	0.41	0.15	0.0233
	Fodder Beet	0.67	0.15	0.0220
	Sugar Beet	NA	0.17	0.0040
5. Other	Fruit	0.27	NA	0.0214
	Grass	0.05	0.32	0.0091
	Green Corn	0.46	0.13	0.0230
	Non-Leguminous Vegetables	1.86	0.85	0.0083
	Rape	1.86	0.85	0.0083
	Renewable Energy Crops	0.05	0.32	0.0115
	Silage Corn	2.00	0.60	0.0150
	Sunflowers	1.18	NA	0.0221
	Tobacco	NA	0.20	0.0060
	Berries	NA	0.20	0.0060
	Vine	0.46	0.13	0.0230
	Oil Squash	4.62	0.85	0.0106
	Oil Hemp	1.25	0.85	0.0071
	Oil Flax	NA	1.00	NA
	Hops	1.18	NA	0.0221
	Medicinal Plants and Herbs	1.18	NA	0.0221

Table A – 31 NH₃ and NO_x emission factors for 3Db Indirect N₂O emissions from managed soils 1990–2015.

Emission Factors Volatilisation	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
	%																										
NH ₃ from application of animal manure N (F _{rac,co,sm})	26.94	26.97	26.96	26.91	27.00	27.07	26.71	26.35	25.98	25.63	25.29	24.94	24.54	24.71	24.88	25.06	25.31	25.53	25.01	24.40	23.80	23.51	23.20	22.93	22.64	22.36	
Mature Dairy Cattle	28.41	29.43	29.46	29.48	29.51	29.53	29.26	28.97	28.67	28.37	28.05	27.71	27.37	27.50	27.63	27.76	27.89	28.02	27.39	26.76	26.13	25.76	25.39	25.02	24.65	24.28	
Other Mature Cattle	27.35	27.29	27.23	27.17	27.11	27.05	26.80	26.53	26.24	25.94	25.61	25.25	24.87	25.52	26.03	26.71	27.25	27.76	27.26	26.74	26.21	26.00	25.79	25.57	25.36	25.15	
Growing Cattle (weighted average)	27.67	27.67	27.68	27.71	27.77	27.81	27.47	27.16	26.78	26.33	26.07	25.82	25.20	25.63	26.05	26.45	26.98	27.30	26.75	26.27	25.76	25.48	25.21	24.95	24.69	24.42	
Sheep (weighted average)	8.81	8.92	9.03	9.14	9.25	9.36	9.36	9.37	9.36	9.36	9.34	9.33	9.32	9.69	10.08	10.48	10.89	11.27	11.19	11.07	10.96	10.85	10.74	10.63	10.52	10.40	
Swine (weighted average)	22.85	22.77	22.68	22.60	22.52	22.43	22.10	21.76	21.40	21.02	20.62	20.21	19.77	20.03	20.59	20.79	20.55	20.79	21.06	20.41	19.77	19.12	18.98	18.85	18.70	18.56	
Buffalo (weighted average)	NA	NA	28.38	28.43	28.49	28.54	28.30	28.05	27.78	27.49	27.17	26.84	26.48	26.86	27.23	27.61	27.98	28.35	27.61	26.89	26.18	25.93	25.68	25.43	25.18	24.93	
Camels (weighted average)	NA	NA	NA	NA	NA	NA	9.36	9.37	9.39	9.40	9.42	9.43	9.44	9.82	10.20	10.60	11.01	11.44	11.28	11.13	10.98	10.89	10.79	10.70	10.61	10.52	
Deer (weighted average)	8.79	8.90	9.01	9.13	9.24	9.35	9.36	9.37	9.39	9.40	9.42	9.43	9.44	9.82	10.20	10.60	11.01	11.44	11.28	11.13	10.98	10.89	10.79	10.70	10.61	10.52	
Goats	9.05	9.17	9.28	9.40	9.51	9.62	9.45	9.28	9.11	8.93	8.76	8.59	8.42	9.42	10.37	11.28	12.15	12.98	11.68	10.30	8.85	9.09	9.33	9.57	9.82	10.06	
Horses (weighted average)	9.23	9.35	9.47	9.58	9.70	9.82	9.64	9.26	8.98	8.65	8.22	7.87	8.31	8.63	8.95	9.26	9.57	9.88	9.82	9.77	9.72	9.88	10.04	10.21	10.37	10.53	
Mules and Asses (weighted average)	9.23	9.35	9.47	9.58	9.70	9.82	9.65	9.26	8.98	8.65	8.24	7.99	7.61	8.03	8.46	8.86	9.31	9.73	10.47	11.18	11.85	11.55	11.24	10.92	10.60	10.27	
Poultry (weighted average)	13.76	14.00	14.25	14.48	14.72	14.96	14.79	14.64	14.47	14.30	14.11	13.94	13.78	13.47	13.14	12.88	12.67	12.50	12.91	13.64	14.38	14.36	14.34	14.32	14.29	14.28	
Rabbits	9.33	9.33	9.33	9.33	9.33	9.33	9.26	9.19	9.11	9.04	8.97	8.90	8.83	8.80	8.77	8.74	8.72	8.69	8.75	8.82	8.89	8.89	8.89	8.89	8.89	8.89	
Livestock NCAC (weighted average)	9.23	9.35	9.47	9.58	9.70	9.81	9.56	9.41	9.27	9.13	8.97	8.84	8.68	9.12	9.54	10.03	10.44	10.86	10.71	10.44	10.21	10.22	10.30	10.34	10.39	10.45	
NH ₃ from urine and dung N deposited on PR&P (F _{rac,co,sm})	4.96	4.97	4.98	4.99	4.99	5.00	5.02	5.03	5.05	5.06	5.07	5.07	5.08	5.10	5.13	5.16	5.17	5.20	5.16	5.13	5.10	5.11	5.12	5.14	5.15	5.17	
Mature Dairy Cattle	4.95	4.94	4.94	4.94	4.94	4.93	4.91	4.90	4.89	4.88	4.87	4.86	4.85	4.84	4.82	4.81	4.80	4.80	4.80	4.80	4.80	4.80	4.80	4.80	4.79	4.79	
Other Mature Cattle	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	
Growing Cattle (weighted average)	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	
Sheep (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Swine (weighted average)	NA	NA	NA	NA	NA	NA	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00	14.00
Buffalo (weighted average)	NA	NA	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	4.98	
Camels (weighted average)	NA	NA	NA	NA	NA	NA	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Deer (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Goats	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Horses (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Mules and Asses (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
Poultry (weighted average)	NA	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	42.00	
Rabbits	NA																										
Livestock NCAC (weighted average)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
NH ₃ from commercial fertilizer N (F _{rac,co,sm})	6.10	5.37	5.38	5.62	5.82	5.83	5.85	5.65	5.30	5.17	5.35	5.04	5.01	4.69	4.81	4.41	4.30	4.76	4.48	4.21	4.26	4.40	4.21	4.43	4.72	4.81	
Urea	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	15.00	
Other Mineral Fertilisers	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	
Recycling Fertilisers (weighted average)	17.58	18.04	18.48	18.91	19.33	19.74	19.92	20.07	20.18	20.26	20.29	19.35	18.20	16.91	15.64	14.14	13.91	13.86	13.35	12.31	11.00	11.10	11.33	11.57	11.80	12.01	
Sewage Sludge	20.00	20.79	21.58	22.37	23.15	23.94	24.37	24.79	25.22	25.64	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	26.07	
Compost	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	3.43	
Digestate Liquid	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	30.00	
Digestate Solid	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	
NH ₃ from Agricultural Soils (kg/ha/year)	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	
NH ₃ from Alpine Area (kg/ha/year)	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	
NO _x from applied fertilisers (F _{rac,nox})	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	
NO _x from urine and dung N deposited on PR&P (F _{rac,nox})	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	0.55	

Annex 4: National energy balance and reference approach

A4.1 Swiss energy balance: energy flows

The diagram shows a summary of the Swiss energy flow 2015 as published by the Swiss Federal Office of Energy (SFOE 2016). Diagram languages are German and French. The energy balance is also provided in tabular form in Table A – 32.




Figure A – 2 Energy flow in Switzerland 2015 in TJ (SFOE 2016)

Table A – 32 Switzerland's energy balance 2015 (SFOE 2016) in TJ¹³.Tabelle 4
Tableau 4Energiebilanz der Schweiz für das Jahr 2015 (in TJ)
Bilan énergétique de la Suisse pour 2015 (en TJ)

	Holzenergie	Kohle	Müll und Industrieabfälle	Rohöl	Erdölprodukte	Gas	Wasserkraft	Kernbrennstoffe	Übrige erneuerbare Energien	Elektrizität	Fernwärme	Total
	Énergie du bois	Charbon	Ord. mén. et déchets ind.	Pétrole brut	Produits pétroliers	Gaz	Énergie hydraulique	Combustibles nucléaires	Autres énergies renouvelables	Électricité	Chaleur à distance	Total
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
Inlandproduktion	38 060	–	56 630	–	–	0	142 150	–	26 000	–	–	262 840
+ Import	2 100	5 440	–	121 340	337 620	119 420	–	241 040	1 840	152 300	–	981 100
+ Export	– 110	0	–	–	– 17 930	–	–	–	–	– 156 030	–	– 174 070
+ Lagerveränderung!	–	– 30	–	860	10 900	–	–	–	–	–	–	– 11 730
= Bruttoverbrauch	40 050	5 410	56 630	122 200	330 590	119 420	142 150	241 040	27 840	– 3 730	0	1 081 600
+ Energieumwandlung:												
• Wasserkraftwerke	–	–	–	–	–	–	– 142 150	–	–	142 150	–	0
• Kernkraftwerke	–	–	–	–	–	–	–	– 241 040	–	79 540	1 100	– 160 400
• konventionell-thermische Kraft-, Fernheiz- und Fernheizkraftwerke	– 2 190	–	– 46 430	–	– 470	– 6 950	–	–	–	9 580	18 950	– 27 510
• Gaswerke	–	–	–	–	–	0	–	–	–	–	–	0
• Raffinerien	–	–	–	– 122 200	122 240	–	–	–	–	–	–	– 40
• Diverse Erneuerbare	– 1 010	–	–	–	–	850	–	–	– 7 270	6 180	0	– 1 250
+ Eigenverbrauch des Energiesektors, Netzverluste, Verbrauch der Speicherungen	–	–	–	–	– 7 040	– 390	–	–	–	– 24 030	– 1 760	– 33 220
+ Nichtenergetischer Verbrauch	–	–	–	–	– 20 900	–	–	–	–	–	–	– 20 900
= Endverbrauch	36 850	5 410	10 200	0	424 420	112 930	0	0	20 570	209 690	18 290	838 360
Haushalte	17 970	400	–	–	79 520	46 260	–	–	13 500	67 540	7 200	232 390
Industrie	10 430	5 010	10 200	–	16 270	39 370	–	–	1 560	64 760	7 080	154 680
Dienstleistungen	7 860	0	–	–	35 030	25 530	–	–	3 190	62 550	4 010	138 170
Verkehr	–	–	–	–	290 530	1 390	–	–	2 070	11 290	–	305 280
Statistische Differenz inkl. Landwirtschaft	590	0	–	–	3 070	380	–	–	250	3 550	0	7 840

1 + Lagerabnahme
– Lagerzunahme
1 + diminution de stock
– augmentation de stock¹³ Liechtenstein's consumption of liquid fuels is included in the numbers (see chapter below on Final Swiss energy consumption).

A4.2 Differences between IEA data and the reference approach

Reviewers have repeatedly asked for explanations of the apparent differences between the energy data held by the International Energy Agency (IEA) and the data reported in the reference approach. In order to clarify the pertaining issues, the reasons for the major differences are given below. Data for the year 2010 are used to illustrate the description.

General remarks

The net calorific values used by IEA differ from those used in the GHG inventory. In order to avoid differences caused by the conversion with different NCV, the comparison between IEA and the Reference Approach is made in kt.

Stock changes as reported by IEA are only including primary stocks (IEA 2005), while the reporting in the Reference Approach includes secondary and tertiary stocks. This results in a particularly large difference for gas oil, as retailers and end-consumers hold considerable amounts of heating fuel on stock. The IEA subsumes secondary and tertiary stock changes under statistical difference.

All data regarding liquid fuel consumption reported by the IEA includes fuel consumption in Liechtenstein (Geographical coverage in IEA 2012). For reporting purposes under the UNFCCC, consumption of Liechtenstein is subtracted.

Data sources used for the comparison shown in Table A – 33 below are:

- Switzerland's greenhouse gas inventory 1990–2011, submission of 15. April 2013, CRF-table 1.A(b), (FOEN 2013).
- Energy statistics of OECD countries (2012 Edition), (IEA 2012).

Liquid fuels

The total amount of liquid fuel consumption as reported in the greenhouse gas inventory is 11'052 kt. There is a difference of 13 kt (0.1%) between CRF and IEA. This difference is primarily caused by the different methodology used for aviation bunkers (see below).

Crude oil

Crude oil in the reference approach contains additives, while IEA lists them separately (data in italics in Table A – 33). The difference between CRF and IEA is smaller than 0.1% if the sum of additives, refinery feedstocks and crude oil is considered.

Gasoline

The comparison is made for motor gasoline only. Aviation gasoline is included under aviation fuels. Gasoline reported by IEA includes gasoline used in Liechtenstein (LIE), which is subtracted for reporting under the UNFCCC. The difference between CRF and IEA is approximately 0.1%, if the consumption of LIE is taken into account.

Aviation fuels

The different aviation fuels are aggregated in the greenhouse gas inventory. For comparison of IEA and reference approach, all aviation fuels are summed up. The difference between IEA and reference approach if considering the apparent final consumption is 12 kt (approximately 1% of imports). This difference is largely due to a different methodology used to estimate international bunker. Aviation bunkers have to be reported monthly to the IEA. As the tier 3 approach used for the greenhouse gas inventory is not available on a monthly basis, the international bunker fuel estimate of IEA consists of the total consumption at the two international airports in Zurich and Geneva, while all remaining fuel use is considered domestic. The reporting in the national greenhouse gas inventory is based on a much more detailed approach, where information on single flights is taken into account. Due to the different approach, the numbers are somewhat different. However, the order of magnitude is the same, and the information in the inventory is based on a higher-tier method and presumably more accurate.

Diesel and gas oil

The IEA numbers include diesel and gas oil used in Liechtenstein. Furthermore, stock changes are reported differently in the CRF and by the IEA. Secondary and tertiary stock changes are subsumed under statistical difference by the IEA, while they are included in the stock change reported in the reference approach. If the statistical difference is taken into account, the difference in the apparent consumption is less than 0.1%.

Residual fuel oil

Data agree between IEA and UNFCCC. It seems as if there is a rounding error in the imported amounts, leading to an apparent difference of 1 kt. According to the foreign trade statistics, 33'693 t of residual fuel oil had been imported in 2010.

Bitumen

Bitumen is a main feedstock in the greenhouse gas inventory. Data between IEA and the reference approach compare well. Again, small differences are likely due to the use of rounded values, leading to apparent differences of the order of 1-2 kt.

Petroleum coke

There are considerable differences (26 kt) in the reported numbers for petroleum coke import. The reason for this apparent difference is that for IEA, all petroleum coke is reported together. In the greenhouse gas inventory submitted in 2013, however, only the petroleum coke used as a fuel was reported under petroleum coke, while calcined petroleum coke was reported together with "other oil" as feedstocks. This is largely a consequence of the treatment of fuels and feedstocks in the Swiss overall energy statistics (SFOE 2012).

Lubricants

There are small differences between IEA and the reference approach, as the data reported to the IEA comprises a slightly different set of customs tariff headings for lubricants to the one used for the Swiss energy statistic. The substances not reported under lubricants in the reference approach are reported under other oil.

Liquefied petroleum gas (LPG)

The reporting of liquefied petroleum gas in the greenhouse gas inventory includes white spirit and lamp oil. As for petroleum coke, IEA numbers include fuels that are used as feedstocks, while in the reference approach, only liquefied petroleum gas, white spirit and lamp oil used as fuels are reported under liquefied petroleum gas. The difference in apparent consumption between IEA and the reference approach is 3 kt (0.03% of total liquid fuel consumption).

Other oil products

In the greenhouse gas inventory, all other oil products are reported together, while IEA has a finer degree of disaggregation. As already mentioned above, the share of petroleum coke that is used as a feedstock is reported under other oil in the greenhouse gas inventory. Therefore, the difference between IEA and the reference approach corresponds largely to the difference in apparent consumption of petroleum coke.

Solid fuels

Solid fuels play only a minor role in Switzerland (246 kt) and are reported in good agreement.

Gaseous fuels

In the greenhouse gas inventory, the amount of gas reported under 1B2b Fugitive emissions is subtracted from the total gas import as reported by IEA, as this gas is not used for energy purposes. Taking this into account the difference is of the order of 2 TJ.

Table A – 33 Comparison of the IEA energy statistic with the Reference Approach for the year 2010. Numbers in italics are fuels that are reported in a finer disaggregation in the IEA energy statistic than in the Reference Approach. Numbers in bold aggregate the data to the level of disaggregation used in the Reference Approach.

CRF vs. IEA (2010) Gg	Import		Export		Bunker		Stock change		Stat. diff.		LIE		Consumption	
	IEA	CRF	IEA	CRF	IEA	CRF	IEA	CRF	IEA	CRF	IEA	CRF	IEA	CRF
Crude oil	<i>4'488</i>	<i>4'546</i>					0	1	0				<i>4'488</i>	<i>4'547</i>
Refinery feedstocks	3						1		2				6	
Additives/blending components	51						-1		2				52	
													4'546	4'547
Motor gasoline	1'850	1'838					-9	-6	4	15			1'830	1'832
Aviation gasoline	7						-2		-1				4	
Kerosene type jet fuel	<i>1'354</i>	<i>1'362</i>			-1'367	-1'352			2	6			-7	12
Other Kerosene	3												3	
													0	12
Gas/diesel oil	3'510	3'485	-21	-39	-10	-11	38	1'072	1'020	27			4'510	4'507
Fuel oil	33	34	-323	-316			-17	-17	7				-300	-299
Liquefied petroleum gases (LPG)	50	54	-24	-25						0.1			26	29
White spirit & SBP	7									-1			6	
													32	29
Bitumen	317	318	-2	-2									315	317
Lubricants	86	72	-38	-16					7				55	56
Petroleum coke	73	47											73	47
Naphtha	1							5		-1			5	
Paraffin waxes	1												1	
Non-specified oil products / other oil	4	63	-	-23			-	-6					4	33
													10	33
Liquid fuels													11'039	11'052
Anthracite	7												7	
Other bituminous coal	123	152					36	32					159	184
Lignite	66	62					-4						62	62
Coke oven coke	18												18	
Solid fuels													246	246
Natural gas (TJ, NCV)	126'014	125'627											126'014	125'627
Fugitive emissions (TJ, NCV)		389												389
Gaseous fuels													126'014	126'016

Additional information regarding reporting of waste-derived fuels

During the in-country review in 2016, the ERT identified that the apparent consumption of non-biomass fraction of waste in the CRF table 1.A(b) was systematically smaller than the consumption reported to IEA. The difference stems from the assumptions made with regard to the fossil and renewable fractions. The SFOE, which is responsible for reporting to the IEA, allocates total wastes to 50% fossil and 50% renewable. For the greenhouse gas inventory, a more sophisticated method based on a detailed analysis of waste composition and measurements in the flue gas of waste incineration plants is used to estimate fossil and renewable fractions (see chp. 3.2.5.2.1).

Annex 5: Additional information

A 5.1 Independent verification of the National Swiss Inventory for F-gases

Introduction

Since 2000 the Swiss Federal Laboratories for Materials Science and Technology (Empa) performs continuous measurements of halogenated greenhouse gases at the high-Alpine site of Jungfrauoch (3580 m asl). These measurements are used for estimating emissions of fluorinated greenhouse gases (HFCs, SF₆) from Switzerland and neighbouring countries. This information can be used for an independent assessment of Swiss inventory data of these greenhouse gases. The independent emission estimate is not used directly for deriving data for the inventory. Data is used, however, to identify discrepancies, which in turn lead to a reassessment of the corresponding part of the inventory and to evaluate options for further improvements of the inventory.

For this independent assessment the so-called tracer-ratio method is applied, where Swiss pollution events of HFC and SF₆, arriving at Jungfrauoch, are scaled to concurrent pollution events of carbon monoxide (CO) and then multiplied by the Swiss CO emission inventory (see Figure A – 3 for a graphical description of the method). Other methods that rely on atmospheric observations are also being developed at Empa for future usage. Similar approaches are also used for independent verification of greenhouse gas emissions in the United Kingdom (UK MetOffice – using measurements from Mace Head, Ireland) and in Australia (CSIRO – using measurements from Cape Grim, Tasmania).

Method description

For yearly estimates of Swiss emissions of HFCs and SF₆ only periods are used when the air masses at the high-Alpine station of Jungfrauoch are predominantly influenced by emissions from Switzerland. The number of events which can be used each year depends on the meteorological conditions and is between 7-15 days per year (mostly in the summer). The process to select these periods is shown in Figure A – 3 and is shortly described here. First, the trajectories from the COSMO-model from MeteoSwiss are screened for periods when the Jungfrauoch site has been under the influence of air masses which were within the Swiss boundary layer for the last 48 hours. Second, for these periods mixing ratios of HFCs and SF₆ are compared with those of CO. Periods which show a concurrent increase for both groups of compounds are selected for the independent assessment of Swiss emissions, as this is taken as an indication of thorough mixing of Swiss emissions during the transport to the height of Jungfrauoch. Third, the emissions are calculated for each case/day using the formula in Figure A – 3. The resulting emissions are only used for the annual emission estimate if they are within three standard deviations of the average (Grubbs test). This criterion is met by approximately 90% of the selected data. Finally, annual emissions are estimated as the median from these individual cases. These annual estimates are merged to a 3-year annual average centered over a 3-year period (e.g. the estimate for the 2015 emissions is calculated by using data from 2014–2016). Since 2009 the error of the estimates for HFCs has been assessed by using the range of the 25%-75% percentiles of the estimates from single pollution events. For estimates between 2001–2008 the average of the 2009–2011 errors has been taken. For SF₆, with comparably low emissions and a higher degree of uncertainty, a general uncertainty of 50% is estimated, based on the long-term average of the 25%-75% percentiles. An additional absolute error could occur if the Swiss

emissions of CO are over/underestimated by the inventory. This would linearly be transmitted to the emissions of the fluorinated greenhouse gases.




Figure A – 3 Description of the procedure to estimate annual emissions of HFCs from Switzerland by using continuous measurements of HFCs at Jungfraujoch (Switzerland).

Results and discussion:

In the following, Swiss emissions of five HFCs (HFC-134a, HFC-125, HFC-152a, HFC-143a, HFC-32) and of SF₆ are estimated based on data from Jungfraujoch and are compared to the emission estimate of the Swiss greenhouse gas inventory. Further emission estimates of other fluorinated greenhouse gases will be added in future National Inventory Reports (NIR) upon availability.

HFC-134a

HFC-134a is the most important anthropogenic HFC. Its main source is the diffuse emission from its usage as cooling agent in mobile air conditioners (MACs). Further relevant applications are the usage as propellant and in cooling mixtures in the industrial and commercial refrigeration as well as in stationary air conditioners and heat pumps. The stock of HFC-134a in MACs and the related emissions have been steadily increasing over the past years. The stabilization of the total emissions between 2005 and 2009 is related to the decreasing HFC used in propellants and to optimizations in the industrial and commercial refrigeration. Increasing tendencies are found again afterwards until around 2013 in the inventory and to a lesser degree in the measurement-based estimates, due to the still growing stock of HFC-134a in refrigeration and air conditioning equipment and due to new applications using HFC-134a for research (i.e. as tracer gas). In most recent years emissions are stable in both the inventory and in the measurement-based estimates. Estimated emissions based on measurements at Jungfraujoch agree fairly well with the emission estimates of the Swiss greenhouse gas inventory. Until 2007 the emissions according to the inventory were slightly higher than the ones based on measurements although data for both methods often agreed within the estimated uncertainty of 25%. Afterwards, the gap got larger but in recent years the relative difference remained constant.




Figure A – 4 Comparison of HFC-134a emissions from Switzerland: Inventory and estimates from measurements at Jungfraujoch.

HFC-125

HFC-125 is mainly used in cooling mixtures in air conditioners and commercial refrigeration equipment. Estimated emissions from Jungfrauoch measurement data are in fairly good agreement with emissions provided by the inventory. Until 2013, emissions from the inventory consistently exceeded emissions based on measurements. However, since 2014 estimated emissions from the measurement-based method approach the emissions from the inventory.




Figure A – 5 Comparison of HFC-125 emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

HFC-152a

HFC-152a is mainly used as a blowing agent in open-cell polyurethane (PU) foams, in closed cell PU sprays and closed-cell extruded polystyrene (XPS) foams. In open cell foams, 100% of emissions are related to the blowing process. In closed cell foams a portion of the blowing agent remains in the product and emissions occur continuously over the lifetime, depending on the cell- and molecular-structure of the blowing agent. Unlike for other blowing agents, experts assume that within the first year of the lifetime of the foam 95–100% of HFC-152a is emitted. The emissions of the first year are commonly allocated to the country of production (according to UNFCCC good practice guidance). These assumptions and allocation are also applied for the model used in the Swiss inventory for estimating HFC-152a emissions under the source category 2F2 (Foam Blowing).

HFC-152a emissions from foams in the inventory are mainly related to the production and consumption of PU spray. Most of other foam products are imported and consequently these emissions are allocated to the country of origin. The reported decrease in the inventory since 2003 reflects the replacement of HFC-152a in PU spray.

Up to the year 2002 estimated emissions from Jungfrauoch measurement data are lower than reported in the inventory and from then onwards they are higher. This can be explained

by the UNFCCC practice to allocate HFC-152a emissions of the first year to the country of production of foams (which is except for PU spray mainly outside Switzerland). However, in reality a fraction of these first year emissions actually occur during usage of the products (e.g. for insulation) in Switzerland and therefore are reflected in the measurements but are not reflected by definition in the inventory¹⁴. Emissions estimated from Jungfrauoch show a consistent negative trend related to the partial phase-out of HFC-152a from the foam-blowing applications.




Figure A – 6 Comparison of HFC-152a emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

HFC-143a and HFC-32

HFC-143a and HFC-32 are mainly used in cooling agent mixtures in commercial refrigeration and stationary air conditioners (together with HFC-134a and/or HFC-125). Until 2013 HFC-143a emissions estimated from Jungfrauoch measurement data were consistently slightly lower than emissions provided by the inventory. However, since 2014 they agree well with the inventory data, due to a rise in emissions estimated by the measurement-based method. A similar development can also be seen in estimated HFC-125 emissions (see above).

Measurement-based estimates of HFC-32 are consistently lower than the data from the inventory but show a parallel constant increase in emissions.

¹⁴ Nonetheless it is important to apply the UNFCCC approach in the inventory as otherwise double counting may occur when allocating the total emissions to the country of origin and the country of product use.




Figure A – 7 Comparison of HFC-143a emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.




Figure A – 8 Comparison of HFC-32 emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

Sulfur hexafluoride (SF₆)

Emissions of SF₆ in Switzerland are mainly due to its use as an insulator of electrical equipment, as for example in gas insulated switchgears and in gas circuit breakers. Additional minor emissions arise from magnesium smelters and its former use from insulating windows and various other applications. Emissions for both methods show a remarkable similarity in the trend, except for the period of 2001–2003, when slightly higher emissions were estimated from the measurement-based method. The data quality of the inventory has been improved in this period. Since 2003, a mass balance approach based on industry data is applied for the use of SF₆ in electrical equipment. Increasing SF₆ emissions 2012–2014 result in the inventory from the disposal of insulating windows (all SF₆ released).




Figure A – 9 Comparison of SF₆ emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

A 5.2 Independent verification of methane emissions

Introduction

In 2013 the Swiss Federal Laboratories for Material Science and Technology (Empa), ETH Zürich and University of Bern established a greenhouse gas (GHG) observing network in Switzerland as part of the CarboCount-CH project (www.carbocount.ch). The network consists of four sites that continuously measure the atmospheric concentration of carbon dioxide (CO₂) and methane (CH₄). The sites were chosen to cover most of the densely populated and agriculturally used Swiss Plateau (see Figure A – 10). Atmospheric transport simulations confirm that the measurements at these sites are sensitive to emissions from a large part of Switzerland (Oney et al., 2015). The aim of CarboCount-CH was to better understand and quantify the exchange of the abovementioned GHGs between the atmosphere and the land surface by inverse modelling. Here, the results of such an inverse modelling approach to validate total Swiss CH₄ emissions are reported. The analysis was carried out for the measurement period March 2013 to February 2014, the first year with data available from all four CarboCount-CH sites. In addition, CH₄ observations from the high altitude site Jungfrauojoch (Empa) and the mountaintop site Schauinsland (Germany, Umweltbundesamt) were used to further constrain the emission estimates. The results summarized here are excerpts from the publication by Henne et al. (2015). An update of this estimate for the full period 2013–2015 is currently being prepared in a follow-up study and is planned to be integrated in next years' NIR.

Methods

The applied approach is based on source sensitivities calculated for each measurement site with the Lagrangian atmospheric transport model FLEXPART. FLEXPART is driven with high resolution meteorological input data from the numerical weather prediction model COSMO (7 km by 7 km horizontal resolution) provided by the Swiss national weather service (MeteoSwiss). For each site 3-hourly source sensitivities were calculated by running the model in time-inverted mode, releasing in each 3-hour time interval 50'000 air parcels and following them 4 days backward in time.

When combining source sensitivities with a map of emission fluxes, atmospheric concentrations at the location of the observations can be obtained. These simulated concentrations can be compared with the measurements, and through “inverse modeling” an optimized (*a posteriori*) emission distribution within and around Switzerland can be estimated that minimizes the differences between simulated and observed concentrations while also considering the uncertainties of the initial (*a priori*) emission distribution.

Two different inverse methods were applied: a Bayesian approach and an extended Kalman filter. Technical details of these approaches can be found in Henne et al. (2015) and Brunner et al. (2012), respectively.

As *a priori* emissions for Switzerland the MAIOLICA CH₄ inventory was used (Hiller et al., 2014), which disaggregates the emissions reported by the Swiss National Inventory Report (NIR) onto a regular spatial grid. For emissions outside of Switzerland the European TNO/MACC-2 inventory was employed (Kuenen et al., 2014). The total anthropogenic emissions of the MAIOLICA inventory was 176 kt yr⁻¹, which corresponds to the Swiss CH₄ emissions in 2012 as reported by the NIR in 2014. The MAIOLICA inventory also includes a small contribution from natural sources of 3 kt yr⁻¹ (<2%).

Next to a reference inversion a number of sensitivity inversions (30) were set up by systematically varying different aspects of the inversion system (inverse algorithm, transport model, particle release height, *a priori* emissions, boundary conditions, uncertainty covariance, selection of observational data). In our reference case the measurements at the two CarboCount-CH sites Beromünster and Lägern-Hochwacht were used together with measurements at Schauinsland and Jungfrauoch. The CarboCount-CH sites Gimmiz and Fruebuel, which were more difficult to simulate due to their proximity to local CH₄ sources, were only treated as validation sites.




Figure A – 10 : Map of Switzerland illustrating the location of CarboCount-CH sites (blue), supplementary sites (black), cantonal borders (gray), major cities (dark blue), and selected canton names (red). The sites are: Beromünster (tall tower, BEO), Lägern Hochwacht (mountain top, tower, LHW), Gimmiz (flat, tower, GIM), Fruebuel (mountain, near surface, FRU), Schauinsland (mountain top, SSL), and Jungfrauoch (high Alpine, JFJ).

Results and Discussion

The inversion results are presented in terms of national total emissions and spatial distribution. The inversion system was not set-up to optimize emissions by category separately, but to estimate the spatial distribution of total emissions. Nevertheless, through the spatial information the results can provide qualitative insights into the contribution from specific source categories that dominate in a given region. Further details and discussion on the inversion performance and results can be found in Henne et al. (2015).

National Totals

The best inverse estimate of total Swiss CH₄ emissions for the observation period March 2013 to February 2014 is 196 ± 18^{15} kt yr⁻¹. This number represents the average and standard deviation over the reference and all sensitivity inversions (Figure A – 11). This is in close agreement with the NIR value reported in 2015 for the years 2012 and 2013 of 206 ± 33 kt yr⁻¹. The uncertainty given in the NIR amounts to 16% of the national total and was also used as the uncertainty of our *a priori* emissions. The inversion results suggest a reduced uncertainty of 9%.

In conclusion, the independent atmospheric observations of the CarboCount-CH network largely confirm the estimate of the NIR and further suggest a reduced uncertainty on the reported value.




Figure A – 11: Histogram of total Swiss CH₄ emissions taken from all individual sensitivity inversions: low (light green) and high (light orange) particle releases. The base inversion prior (green) and posterior (blue) estimate as well as the average over all sensitivity inversions (red) and the NIR 2015 estimate (purple) are indicated by their Gaussian probability density functions. Taken from Henne et al. (2015).

Spatial and Temporal Distribution and Source Processes

In Figure A – 12 the spatial distribution of the *a priori* emissions (a), the *a posteriori* emissions (b), their absolute (c) and relative (d) difference (a posteriori minus a priori) is shown. An irregular inversion grid was used that exhibits high spatial resolution close to the observations and gets coarser with distance to these. In the *a priori* emissions the dominating role of agricultural emissions in the rural areas of the Cantons Lucerne and Thurgau/Appenzell is clearly seen. In contrast, the densely populated areas of Zurich, Basel and Geneva do not show up as emission hot-spots, as expected from the small contribution of emissions from natural gas distribution and waste water treatment reported in the NIR.

A posteriori emissions estimated by the inversion were smaller than *a priori* emissions in the agricultural areas of Canton Lucerne and Canton Thurgau, but were increased in the north-

¹⁵ All uncertainties given here represent 1- σ confidence levels (66% confidence interval around the mean).

eastern part of Switzerland (Cantons Appenzell and Saint Gallen). Smaller differences were seen in other parts of the country. When allowing the inversion to derive seasonal mean instead of annual mean emissions, a clear seasonal cycle with reduced winter time and increased spring to summer emissions was detected. This is in line with the temperature dependent seasonality expected from manure handling and storage and the productivity-dependent seasonality of milk cows. We conclude that the emission reductions as estimated by the inversion are largely due to an overestimation in agricultural emissions in the *a priori* estimate by 10% to 20% that can mainly be attributed to the winter season. A strong seasonality can be expected for emissions from manure handling and storage (lower winter temperatures). Hence, the observed overestimation in the *a priori* may be attributed to this emission process.




Figure A – 12: (a) *a priori* (MAIOLICA) and (b) *a posteriori* surface fluxes of CH₄ in the base inversion with low particle release heights. (c) absolute and (d) relative (to *a priori*) difference between *a posteriori* and *a priori* fluxes. For panels c and d red (blue) colors indicate higher (lower) *a posteriori* than *a priori* emissions. Taken from Henne et al. (2015).

An alternative sensitivity inversion used *a priori* emissions from the EDGAR inventory (v4.2 FT2010, EC JRC 2009). In EDGAR, the Swiss country total amounts to 228 kt yr⁻¹, mostly due to about 25 kt yr⁻¹ larger emissions from the natural gas distribution network (IPCC category 1B2: fugitive emissions from oil and gas). *A posteriori* emissions in this sensitivity inversion were very similar to those of the reference inversion, which in turn required large reductions from the *a priori* distribution especially in densely populated areas (see Figure A – 13). From this we conclude, that the natural gas emissions as given in the NIR (8 kt yr⁻¹) are in much better agreement with our atmospheric observations than the emission in the EDGAR inventory (32 kt yr⁻¹).




Figure A – 13: Annual average (left) *a priori* CH₄ fluxes and (right) absolute difference between *a posteriori* and *a priori* fluxes when using EDGAR *a priori* information

Most sensitivity inversions yielded a similar spatial distribution of the *a posteriori* emissions as the reference inversion. This robust pattern suggests that the increased emissions estimated for north-eastern Switzerland are a true feature, but it remains unclear what source is responsible for this result. A possible candidate is differences in farming practices between this area and the rest of the country, resulting in different per head emissions from livestock. Other potential anthropogenic sources are waste water handling, composting and anaerobic digestion in biogas production and fugitive emissions from the natural gas distribution network. However, there is no indication that these processes occur disproportionately in north-eastern Switzerland. Natural emissions, e.g. from the “Alter Rhein” wetland area, may also contribute to the unexpectedly high emissions from this area. Additional observations at a new site established in this area were conducted during a few months in 2016 to better characterize this source. These observations will be included in the update planned for next year's NIR.

Annex 6: Information on the CRF Reporter

The CRF Reporter still seems to generate errors in the numerical output of some reporting tables. A non-exhaustive list of issues and errors identified so far is provided in the table Table A – 34. Where numbers are not identical in the reporting tables (CRF) and in the NIR, a remark was added in the NIR. This aspect should be taken into consideration when comparing information in the NIR with the reporting tables (CRF).

Additionally, in response to the findings of the ERT, which noted some discrepancies between the NIR and the information included in CRF Summary 3s1 and Summary 3s2 (UNFCCC2017/ID#G.1), we present manually filled reporting tables in Table A – 35 and Table A – 36.

Table A – 34: Identified errors in the output of some reporting tables (CRF).

CRF table	Problem	Solution
Table1A(b)	Template cannot cope with notation key "C" in column "Carbon stored", (resulting in no "net carbon emissions" and "Actual CO2 emissions").	Workaround: use of "NA" instead of "C".
Table1A(d)	The column "Reported under ..." should include the NFR code in the Text.	Actually there is no possibility to change it, because a drop-down-menu determines the text.
Table2(I)s1-s2	In all empty cells should be written „NO“, but the effort would be very big to generate in the navigation tree of the CRF Reporter for every cell a new nod and then to adapt all the import files to these changes.	Because the effort would be quite big, and there is no real benefit, we passed on it.
Table2(II)	In all empty cells should be written „NO“, but the effort would be very big to generate in the navigation tree of the CRF Reporter for every cell a new nod and then to adapt all the import files to these changes.	Because the effort would be quite big, and there is no real benefit, we indicate in the documentation box: „2.B.9, 2.C, 2.E, 2.F.1-2.F6: "NO" for all empty cells".
Table2(II)B-Hs1	In all empty cells should be written „NO“, but the effort would be very big to generate in the navigation tree of the CRF Reporter for every cell a new nod and then to adapt all the import files to these changes.	Because the effort would be quite big, and there is no real benefit, we passed on it.
Table3.B(a)s2	Cell I55: MCF for Pasture for Rabbits: Cell is empty. Should be either "NO" or 1.00.	Problem in the CRF Reporter.
Table3.B(a)s2	Cell L103: MCF for Burned for Fuel for Poultry: Cell is empty. Should be "NO".	Problem in the CRF Reporter.
Table3.B(b)	The IEF for "indirect emissions from atmospheric deposition" (cell R37): Wrong molecular weight. The IEF is displayed in kg N ₂ O per kg N handled instead of kg N ₂ O-N per kg N handled.	Problem in the CRF Reporter.
Table4	N ₂ O from 4(IV) (indirect emissions) are not separated in the categories 4A/B/C/D/E/F and just shown in table 4(IV). But they are summarized in the total sum in table 4 without been displayed in this table 4. The sum is correct (with indirect emissions).	Problem in the CRF Reporter.
Table4	There are empty cells in the CH ₄ - and N ₂ O-columns; numbers and/or notation keys are not inserted	Problem in the CRF Reporter.
Table4.A	All the different documentation-box texts are in the same cell.	Problem in the CRF Reporter.
Table4.B	All the different documentation-box texts are in the same cell.	Problem in the CRF Reporter.
Table4(V)	IEF and Values in Line 9 and 10 are missing.	Problem in the CRF Reporter.
Summary3s1	Wrong information in Method applied and emission factor for several sectors and gases: There should be indicated „NO“ instead of empty cells.	Problem in the CRF Reporter. „NO“ is not imported and there is no possibility to change it, because „NO“ or „NA“ are not in the drop-down menu.
Table10s2	CO ₂ emissions from biomass (row 60) are wrong. Biogenic CO ₂ emissions from 5C are missing.	Problem in the CRF Reporter.
Table10s4	For N ₂ O in sector 4 LULUCF the sum is not just the displayed sum of N ₂ O emissions from the categories 4A+4B+4C+4D+4E+4F (rows 39 to 46) but including indirect N ₂ O from 4(IV). The sum is correct (with indirect emissions).	Problem in the CRF Reporter.
KP-CRF-tables	Text in documentation boxes is right-aligned.	Problem in the CRF Reporter.
4(KP-I)A.2	Information items: every row is shown twice. This is not necessary.	Problem in the CRF Reporter.
4(KP-I)B.1	Description of category and sub-category is shown twice under „identification code“ and „subdivision“. This is not wrong but not necessary.	Problem in the CRF Reporter.

Table A – 35 Manually filled reporting table CRF Summary3s1.

Inventory 2015
Submission 2017
SWITZERLAND

SUMMARY 3 SUMMARY REPORT FOR METHODS AND EMISSION FACTORS USED
(Sheet 1 of 2)

CATEGORIES	CO ₂		CH ₄		N ₂ O		HFC _s		PFC _s		SF ₆		Unspecified mix of HFC _s and PFC _s		NF ₃	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
1. Energy	CS.D.T1.T2.T3	CS.D	CS.T1.D.T2.T3	CR.CS.D	CS.D.T1.T2.T3	CS.D										
A. Fuel combustion	CS.T1.T2.T3	CS	CS.T1.T2.T3	CR.CS.D	CS.D.T1.T2.T3	CS.D										
1. Energy industries	CS.T2.T3	CS	CS.T1.T2.T3	CR.CS.D	CS.D.T1.T2.T3	CS.D										
2. Manufacturing industries and construction	T1.T2.T3	CS	T2.T3	CS.D	T2.T3	CS.D										
3. Transport	T1.T2.T3	CS	T2.T3	CR.CS	CS.D.T2.T3	CS.D										
4. Other sectors	T1.T2.T3	CS	CS.T2.T3	CS.D	CS.T2.T3	CS.D										
5. Other	T2.T3	CS	CS.CR	T2	CS.D	CS.D										
B. Fugitive emissions from fields	CS.D.T1.T2	CS.D	CS.D.T1.T2	CS.D	D.T2	D										
1. Solid fuels	CS.D.T1.T2	CS.D	CS.D.T1.T2	CS.D	D.T2	D										
2. Oil and natural gas	CS.D.T1.T2	CS.D	CS.D.T1.T2	CS.D	D.T2	D										
C. CO ₂ transport and storage	CR.CS.T1.T2.T3	CR.CS.D.PS	T2	D	CS.T2	CS.PS	T1.T2	CS.D	T1.T2.T3	CS.D	T2.T3	CS.D	T2	T2	D	
2. Industrial processes	CS.T2.T3	CS.D.PS	T2	D	CS.T2	CS.PS	T1.T2	CS.D	T1.T2.T3	CS.D	T2.T3	CS.D	T2	T2	D	
A. Mineral industry	T2	PS	T2	D	T3	PS					T2.D	T2.D				
B. Chemical industry	CR.T3	CS.PS														
C. Metal industry	CR.CS.T1	CR.CS.D														
D. Non-energy products from fields and solvent use																
E. Electronic industry																
F. Product uses as ODS substitutes	CR.CS	CS			T2	CS	T2	D	T2	D	T2	CS.D	T1.T2	CS.D	T2	D
G. Other product manufacture and use	CS	CS					T2	D	T2	D	T2	CS.D	T1.T2	CS.D	T2	D
H. Other																

Use the following notation keys to specify the method applied:

- D (IPCC default)
- RA (Reference Approach)
- T1 (IPCC Tier 1)
- T1a, T1b, T1c (IPCC Tier 1a, Tier 1b and Tier 1c, respectively)
- T2 (IPCC Tier 2)
- T3 (IPCC Tier 3)

If using more than one method within one source category, list all the relevant methods. Explanations regarding country-specific methods, other methods or any modifications to the default IPCC methods, as well as information regarding the use of different methods per source category where more than one method is indicated, should be provided in the documentation box. Also use the documentation box to explain the use of notation OTH.

Use the following notation keys to specify the emission factor used:

- D (IPCC default)
- CR (CORINAIR)
- CS (Country Specific)
- PS (Plant Specific)
- OTH (Other)
- M (model)

Where a mix of emission factors has been used, list all the methods in the relevant cells and give further explanations in the documentation box. Also use the documentation box to explain the use of notation OTH.

Table A – 36 Manually filled reporting table CRF Summary3s2.

Inventory 2015
Submission 2017
SWITZERLAND

SUMMARY 3 SUMMARY REPORT FOR METHODS AND EMISSION FACTORS USED
(Sheet 2 of 2)

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO ₂		CH ₄		N ₂ O		HFC ₃		PFC ₃		SF ₆		Unspecified mix of HFC ₃ and PFC ₃		NF ₃	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
3. Agriculture	T1	D	T2	CS,D	CS,T1	CS,D										
A. Enteric fermentation			T2	CS												
B. Manure management			T2	CS,D												
C. Rice cultivation																
D. Agricultural soils ⁽³⁾					CS,T1	CS,D										
E. Prescribed burning of savannas																
F. Field burning of agricultural residues																
G. Liming	T1	D														
H. Urea application	T1	D														
I. Other carbon-containing fertilizers																
J. Other																
4. Land use, land-use change and forestry	T2,T3	CS,D	T1	CS,D	T1	D										
A. Forest land	T2,T3	CS	T1,T2	CS,D	T1,T2	CS,D										
B. Cropland	T2	CS														
C. Grassland	T2	CS	T1	D	T1	D										
D. Wetlands	T2	CS	T1	D	T1	D										
E. Settlements	T2	CS														
F. Other land	T2	CS														
G. Harvested wood products	T2	D														
H. Other																
5. Waste	T2	CS	T2,CS	CS,D	T2	CS,D										
A. Solid waste disposal			T2,CS	CS												
B. Biological treatment of solid waste			T2	CS												
C. Incineration and open burning of waste			T2	CS												
D. Wastewater treatment and discharge	T2	CS	T2	CS,D	T2	CS,D										
E. Other																
6. Other (as specified in summary I-A)	T1	CS	T1	CS	CS,T1b	CS,D										

Use the following notation keys to specify the method applied.

D (IPCC default)

RA (Reference Approach)

T1 (IPCC Tier 1)

T2 (IPCC Tier 2)

T3 (IPCC Tier 3)

When more than one method is indicated, should be provided in the documentation box. Also use the documentation box to explain the use of different methods per source category.

Use the following notation keys to specify the emission factor used:

D (IPCC default)

CR (CORNAIR)

CS (Country Specific)

PS (Plant Specific)

OTH (Other)

M (model)

CR (CORNAIR)

CS (Country Specific)

OTH (Other)

M (model)

T1a, T1b, T1c (IPCC Tier 1a, Tier 1b and Tier 1c, respectively)

T2 (IPCC Tier 2)

T3 (IPCC Tier 3)

When more than one method is indicated, should be provided in the documentation box. Also use the documentation box to explain the use of different methods per source category.

Use the following notation keys to specify the emission factor used:

D (IPCC default)

CR (CORNAIR)

CS (Country Specific)

PS (Plant Specific)

OTH (Other)

M (model)

CR (CORNAIR)

CS (Country Specific)

OTH (Other)

M (model)

T1a, T1b, T1c (IPCC Tier 1a, Tier 1b and Tier 1c, respectively)

T2 (IPCC Tier 2)

T3 (IPCC Tier 3)

When more than one method is indicated, should be provided in the documentation box. Also use the documentation box to explain the use of different methods per source category.

Use the following notation keys to specify the emission factor used:

D (IPCC default)

CR (CORNAIR)

CS (Country Specific)

PS (Plant Specific)

OTH (Other)

M (model)

Where a mix of emission factors has been used, list all the methods in the relevant cells and give further explanations in the documentation box. Also use the documentation box to explain the use of notation OTH.

Documentation box:

- Parties should provide the full information on methodological issues, such as methods and emission factors used, in the relevant sections of chapters 3 to 8 (see section 2.2 of each of Chapters 3 - 8) of the national inventory report (NIR). Use this documentation box to provide references to relevant sections of the NIR if any additional information and further details are needed to understand the content of this table.
- Where a mix of methods/emission factors has been used within one source category, use this documentation box to specify those methods/emission factors for the various sub-sources where they have been applied.
- Where the notation OTH (Other) has been entered in this table, use this documentation box to specify those other methods/emission factors.

Documentation box

Annex 7: Information on the UNFCCC review process

A 7.1 Attachment C of the Saturday paper from the 10 September 2016

**Potential Problems formulated in the course of the review of the 2015
and 2016 annual submissions of Switzerland and of the report to
facilitate the calculation of the assigned amount for the second
commitment period**

Bern, Switzerland. 10 September 2016

For the ERT,

Ms. Laura Dawidowski, lead reviewer Mr. Ioannis Sempos, lead reviewer

ATTACHMENT C

Overview of inventory potential problems identified for the base year that could lead to an adjustment

Annex A sources

Switzerland

Abbreviations:

2006 IPCC GL: 2006 IPCC Guidelines for National Greenhouse Gas Inventories

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

Sector, category, sub-category (with code)	Gas	KC / non-KC	Identified inventory problem in terms of:		
			Missing estimate	Estimate provided but not in line with the 2006 IPCC GL	Estimate provided but lack of transparency
2. Industrial processes and product use 2.C. Metal industry 2.C.3. Aluminium production	CO ₂	KC		X	
<p>Description of problem identified:</p> <p>Switzerland presents, in the NIR, Chapter 4, direct CO₂ emissions from aluminium production estimated using a country specific emission factor that has been obtained on the basis of the anode consumption in the electrolysis process. Moreover, as it was reported in the NIR / page 220: "It is assumed that the anode consisted completely of carbon and that it was fully oxidized during the process." On the other hand, in Chapter 9 (and in CRF table 6), the Party reports indirect CO₂ emission estimates from aluminium production, based on CO and NMVOC monitoring data obtained in the stacks of the aluminium foundries.</p> <p>The ERT notes that according to the 2006 IPCC GLs, in making indirect CO₂ emission estimates, inventory compilers should assess each category to ensure that the carbon is not already covered in the estimates of CO₂ direct emissions taking into consideration the assumptions and approximations made.</p> <p>During the review week, the Party indicated that it could not provide any additional information about the estimation of emissions from aluminium production, given that the last production site for primary aluminium in Switzerland closed down in April 2006.</p> <p>The ERT is of the view that the carbon content of the CO and NMVOCs measured in the stack of the aluminium foundries have been accounted for in the estimation of the direct CO₂ emissions, and for this reason the inclusion of indirect CO₂ emissions represents a double counting.</p> <p>Therefore the ERT concluded that the CO₂ emissions from aluminium production (direct and indirect) are potentially overestimated for the years 1990-2006 and, as a result, the Annex A emissions for the base year are potentially overestimated.</p>					
<p>Recommendation by ERT:</p> <p>The ERT recommends that the Party either:</p> <p>(a) Demonstrate that there is no double counting between the CO₂ direct and indirect emissions reported for aluminium production in 1990 – 2006</p> <p style="text-align: center;">Or</p> <p>(b) Submit revised estimates for CO₂ emissions from aluminium production (category 2.C.3) excluding indirect CO₂ emissions from aluminium production for the entire time series.</p>					

Response / Information by Party:

As already explained during the review week, the NMVOC emissions from source category 2C3 Primary aluminium production originate solely from the **production of the electrodes at the plants** as documented in the Handbook on emission factors for stationary sources (SAEFL 2000). Thus, the resulting indirect CO₂ emissions from NMVOC are not already reported as direct CO₂ emissions. On the other hand, the CO emissions are based on the literature and on information from the plant (measurements at the chimney), and thus predominantly stem from the anodes. Therefore, indirect CO₂ from these CO emissions are already accounted for in the direct CO₂ emissions since it is assumed that the anodes are fully oxidized during the process. This indeed represented a double counting in the inventory.

The double-counting of CO₂ is eliminated by removing the indirect CO₂ emissions from oxidation of CO from 2C3 Primary aluminium production. The CO₂ emissions reported as direct emissions in category 2C3 remain unchanged. The indirect CO₂ emissions reported in CRF table 6 and table 9-1 of the NIR are now excluding indirect CO₂ emissions from oxidation of CO from 2C3. This approach is in line with the reporting of the CO₂ emissions from 1A Fossil fuel consumption with an oxidation factor of 100% and the 2006 IPCC guidelines, vol.1 chap. 7.2.1.5, where it is stated that also for a few IPPU categories, CO₂ estimates can be based on carbon mass balances.

Table: Emissions of 2C3 Primary aluminium production and indirect CO₂ from 2C

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	kt									
2C3 Primary aluminium production										
CO ₂	139.3	131.0	120.6	58.3	38.7	33.2	42.5	43.7	51.7	55.1
CO	3.5	3.3	3.0	1.5	1.0	0.8	1.1	1.1	1.3	1.4
indirect CO ₂	5.5	5.1	4.7	2.3	1.5	1.3	1.7	1.7	2.0	2.2
<i>from anode production only:</i>										
NMVOC	0.06	0.05	0.05	0.02	0.02	0.01	0.02	0.02	0.02	0.02
Indirect CO ₂ emissions										
2C	10.8	9.9	9.6	6.7	5.9	4.9	4.8	4.9	5.4	5.5
2C, excluding CO from 2C3	5.3	4.8	4.9	4.4	4.4	3.6	3.2	3.2	3.3	3.4

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	kt									
2C3 Primary aluminium production										
CO ₂	56.9	58.1	64.3	70.2	71.8	71.7	19.2	NO	NO	NO
CO	1.4	1.5	1.6	1.8	1.8	1.8	0.5	NO	NO	NO
indirect CO ₂	2.2	2.3	2.5	2.8	2.8	2.8	0.8	NO	NO	NO
<i>from anode production only:</i>										
NMVOC	0.02	0.02	0.03	0.03	0.03	0.03	0.01	NO	NO	NO
Indirect CO ₂ emissions										
2C	5.7	5.5	5.4	5.5	5.6	5.5	3.5	2.8	2.9	2.0
2C, excluding CO from 2C3	3.4	3.2	2.9	2.8	2.8	2.7	2.7	2.8	2.9	2.0

	2010	2011	2012	2013	2014
	kt				
2C3 Primary aluminium production					
CO ₂	NO	NO	NO	NO	NO
CO	NO	NO	NO	NO	NO
indirect CO ₂	NO	NO	NO	NO	NO
<i>from anode production only:</i>					
NMVOC	NO	NO	NO	NO	NO
Indirect CO ₂ emissions					
2C	2.4	2.7	2.3	2.3	2.4
2C, excluding CO from 2C3	2.4	2.7	2.3	2.3	2.4

Potential problem unsolved? Rationale:

Overview of inventory potential problems identified for the base year that could lead to an adjustment

Annex A sources

Switzerland

Abbreviations:

2006 IPCC GL: 2006 IPCC Guidelines for National Greenhouse Gas Inventories

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

Sector, category, sub-category (with code)	Gas	KC / non-KC	Identified inventory problem in terms of:		
			Missing estimate	Estimate provided but not in line with the 2006 IPCC GL	Estimate provided but lack of transparency
5.C.2 Open burning of waste	CH ₄ , N ₂ O	Non-KC		x	
<p>Description of problem identified: Switzerland has reported CH₄ and N₂O emissions from open burning of residues from forests under the waste sector (category 5.C.2 Open burning of waste, subcategory Other (natural residues)).</p> <p>During the review, Switzerland explained that the residues in forest are the residues collected from forest activities which are then burnt off-site.</p> <p>The ERT notes that the CO₂ emissions from both on-site and off-site burning of biomass in forest land have been correctly included in the carbon stock changes in the biomass pool and thus reported under the category 4.A: forest land in the LULUCF sector (in accordance with the 2006 IPCC Guidelines).</p> <p>The ERT also notes that CO₂ as well as non-CO₂ emissions from burning of forest biomass in both controlled burning and in wildfires are to be reported in the LULUCF sector, CRF table 4(V) 'Biomass burning'.</p> <p>The ERT further notes that the definition of 'other waste' in section 2.2.4 of Chapter 2 in Volume 5 of the 2006 IPCC GLs does not include forest residues.</p> <p>The ERT considers that the CO₂ and non-CO₂ emissions arising from burning of the biomass should be allocated to the same category unless specific provisions apply.</p> <p>The ERT thus considers that CH₄ and N₂O emissions from the open burning of residues from forest should be reported in the LULUCF sector, category Forest Land, controlled burning, in CRF table 4(V) 'Biomass burning'. The ERT considers that, by reporting these emissions in the waste sector, Switzerland may be overestimating its Annex A base year emissions and, as a result, its assigned amount.</p>					
<p>Recommendation by ERT:</p> <p>The ERT recommends that Switzerland either:</p> <p>(a) Demonstrate that the CH₄ and N₂O emissions from open burning of forest residues are correctly allocated to category 5.C.2 and, as a result, there is no overestimation of the Annex A emissions in the base year</p> <p>OR</p> <p>(b) recalculate the emissions from the LULUCF and waste sectors for the whole time series by allocating the CH₄ and N₂O emissions from open burning of residues from forests from category 5.C.2 open burning of waste to the category 4(V).A.1 controlled burning in Forest land remaining forest land (CRF table 4(V) biomass burning under the LULUCF sector).</p>					

Response / Information by Party:

Switzerland followed recommendation (b) and moved open burning of forest residues from the waste sector 5.C.2 to LULUCF 4(V).A.1 "Controlled burning, residues from forestry". The recalculation leads to a decrease of CH₄ and N₂O emissions in 5.C.2 and to a corresponding increase in 4(V).A.1 in the amount as shown in the following table.

Table: Activity data and emissions moved from 5.C.2 to 4(V).A.1

4 V A 1 Controlled burning of residues from forestry	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄	kt	0.196	0.190	0.184	0.178	0.172	0.167	0.161	0.155	0.149	0.143
N ₂ O	kt	0.0052	0.0050	0.0049	0.0047	0.0046	0.0044	0.0043	0.0041	0.0039	0.0038
Activity data	kt	28.83	27.96	27.10	26.23	25.37	24.50	23.64	22.77	21.91	21.04

4 V A 1 Controlled burning of residues from forestry	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄	kt	0.137	0.131	0.125	0.120	0.114	0.108	0.102	0.096	0.090	0.084
N ₂ O	kt	0.0036	0.0035	0.0033	0.0032	0.0030	0.0029	0.0027	0.0025	0.0024	0.0022
Activity data	kt	20.18	19.31	18.45	17.58	16.72	15.85	14.99	14.12	13.26	12.39

4 V A 1 Controlled burning of residues from forestry	Unit	2010	2011	2012	2013	2014
CH ₄	kt	0.078	0.079	0.079	0.080	0.080
N ₂ O	kt	0.0021	0.0021	0.0021	0.0021	0.0021
Activity data	kt	11.53	11.58	11.64	11.69	11.75

Accordingly, emissions from controlled burning of residues from forestry are reported under the activity "Forest Management" in Table 4(KP-II)4.

CO₂ emissions from controlled fires, as well as those from wildfires, are encompassed ("IE") in the data in CRF Table 4.A and Table 4(KP-I)B.1. Carbon losses in living biomass are reflected in the NFI data-set. Carbon stock changes in dead wood, litter and soil calculated with Yasso07 also cover the influence of forest fires and other disturbances by using NFI-data as an input.

Potential problem unsolved? Rationale:

Overview of inventory potential problems identified for 2013 and 2014 that could lead to an adjustment

Annex A sources

Switzerland

Abbreviations:

2006 IPCC GL: 2006 IPCC Guidelines for National Greenhouse Gas Inventories

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

LDT: light duty trucks, PC: passenger cars

Sector, category, sub-category (with code)	Gas	KC / non-KC	Identified inventory problem in terms of:		
			Missing estimate	Estimate provided but not in line with the 2006 IPCC GL	Estimate provided but lack of transparency
1. Transport, 1.A.3.Transport 1.A.3.b. Road transportation 1.A.3.bi Cars and 1.A.3.bii LDT Gasoline, diesel	N2O	KC		X	
<p>Description of problem identified:</p> <p>Switzerland reported in the NIR (p561) that: "for N2O, no cold start emissions or evaporative emissions are taken into account due to lack of data." (CH4 emissions during cold start are included in the estimates). Switzerland estimates N2O emissions from road transportation for cars and LDT (both gasoline and diesel) by a Tier 3 method: a country-specific territorial emission model with EFs from "The Handbook of Emission Factors, HBEFA" http://www.hbefa.net/e/index.html.</p> <p>During the review, Switzerland indicated that "The Handbook of Emission Factors" does not contain specific N2O EF for cold start. The Party also indicated that the ongoing update of the road transportation model (planned to be incorporated in the 2018 submission) will:</p> <ol style="list-style-type: none"> 1) integrate a new model for cold start emissions 2) review the EF for N2O <p>The ERT considered that 2006 IPCC GLs provides a methodology for estimating N2O cold start emissions for the urban mode (Table 3.2.5 / page 3.24 / volume 2 / 2006 IPCC GLs).</p> <p>In response to a question raised by the ERT during the review week, the Party provided, informally, revised estimations for N2O cold start up emissions for PC and LDT, by applying the latest version of COPERT EFs from EMEP/EEA air pollutant emission inventory guidebook – 2013¹. The ERT considers that this approach would resolve the potential underestimation.</p>					
<p>Recommendation by ERT:</p> <p>The ERT recommends that Switzerland either</p> <ol style="list-style-type: none"> (a) demonstrate that N2O emissions from cold start are not underestimated in the official estimates OR (b) submit officially the revised N2O estimates provided during the review (or those estimated with any methodology consistent with the IPCC 2006 Guidelines) for the whole time-series 1990-2014. 					

¹ <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>

Response / Information by Party:

We implemented recommendation (b)

During the review we presented provisional estimations for N₂O cold start excess emissions for PC and LDT for 1990 and 2014 by means of emission factors of the Copert model, which were recommended by the ERT. The factors are documented in the EMEP/EEA air pollutant emission inventory guidebook - 2013 on p. 91 ff. (<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>)

The ERT confirmed that this approach complies with the 2006 IPCC Guidelines. We corrected the emissions (for PC 1990) and extended the calculation of 2014 for 2013. For the years 1991–2012 we interpolated the emissions linearly between 1990 and 2013.

The final results for N₂O cold start excess emissions of 1A3b are shown in the following table:

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PC	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806
LDT	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806
Sum	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806
PC	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806
LDT	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806
Sum	441920	432002	423084	414166	405248	396330	387412	378494	369576	360658	351740	342822	333904	324986	316068	307150	298232	289314	280396	271478	262560	253642	244724	235806

The results are implemented in the EMIS database and are added to the other emissions of category 1A3b in the re-submitted CRF tables.

Potential problem unsolved? Rationale:

References

Abegg, M., Brändli, U.-B., Cioldi, F., Fischer, C., Herold-Bonardi, A., Huber, M., Keller, M., Meile, R., Rösler, E., Speich, S., Traub, B., Vidondo, B. 2014: Fourth national forest inventory – result tables and maps on the Internet for the NFI 2009–2013 (NFI 4b). Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.

<http://www.lfi.ch/resultate> [queries from database].

Agrammon 2010: The Swiss ammonia model, Federal Office for the Environment FOEN. Internet version in German, French and English: <http://www.agrammon.ch/documents-to-download>, Technical description for download (updated periodically):

<http://www.agrammon.ch/technical-description>

Agricura 2015: Geschäftsbericht 2014/2015. Agricura, Bern [confidential/internal].

Agroscope 2014a: Lime application in Swiss Agriculture. Internal documentation by Bretscher, D., Agroscope Reckenholz-Tänikon Research Station, Zürich [confidential/internal].

Agroscope 2014b: Futtermitteldatenbank. Agroscope, Institut für Nutztierwissenschaften (INT). Posieux, Schweiz. <http://www.feedbase.ch> [queries from database].

Agroscope 2014c: Fütterungstool zur Ermittlung des Energie- und Stickstoffumsatzes von Milchkühen in der Schweiz. Agroscope, Institut für Nutztierwissenschaften (INT). Posieux, Schweiz. <http://www.feedbase.ch> [queries from database].

Agroscope 2016b: Agricultural CH₄ and N₂O emissions in Switzerland: QA/QC. Internal documentation (with continual update) by Bretscher, D. Agroscope Research Station; Zürich, Switzerland [confidential/internal].

Agroscope 2017: Treibhausgasinventar Sektor Landwirtschaft, Tabellen zur Berechnung der Treibhausgasemissionen der schweizerischen Landwirtschaft. Internes Dokument. [GHG inventory, sector Agriculture. Spreadsheets to calculate the GHG emissions of Switzerland's agriculture. Internal document]. Agroscope Reckenholz-Tänikon Research Station, Zürich [confidential/internal].

Alberti, G., Peressotti, A., Piussi, P., Zerbi, G. 2008: Forest ecosystem carbon accumulation during a secondary succession in the Eastern Prealps of Italy. *Forestry* 81: 1–11. <http://dx.doi.org/10.1093/forestry/cpm026>

Alcan 2002: Written communication from Kurt Buxmann (ALCAN) to Carbotech, 30.1.2002 [confidential/internal].

Alcan 2003: Written communication from François Veuthey (ALCAN) to Carbotech [confidential/internal].

Amon, B., Amon, T., Boxberger, J., Alt, C. 2001: Emissions of NH₃, N₂O and CH₄ from dairy cows housed in a farmyard manure tying stall (housing, manure storage, manure spreading). *Nutrient Cycling in Agroecosystems* 60 (1–3): 103–113.

<http://dx.doi.org/10.1023/A:1012649028772>

Andreani-Aksoyoglu, S., Keller, J. 1995: Estimates of monoterpene and isoprene emissions from the forests in Switzerland. *Journal of Atmospheric Chemistry* 20: 71–87.

<http://dx.doi.org/10.1007/BF01099919>

ARE 2002: Fahrleistungen der Schweizer Fahrzeuge. Ergebnisse der periodischen Erhebung Fahrleistungen (PEFA) 2000. Federal Office for Spatial Development, Bern.
<http://www.climatereporting.ch>

ARE 2010: Nationales Personenverkehrsmodell – Basismodell 2005, Bern.
<https://www.aren.admin.ch/aren/de/home/medien-und-publikationen/publikationen/verkehr/nationales-personenverkehrsmodell-des-uvek-basismodell-2005.html> [08.02.2017]

ARE 2012: Ergänzungen zu den schweizerischen Verkehrsperspektiven bis 2030. Bundesamt für Raumentwicklung (ARE), Bern.
<https://www.aren.admin.ch/aren/de/home/medien-und-publikationen/publikationen/verkehr/ergaenzungen-zu-den-schweizerischen-verkehrsperspektiven-bis-203.html>

ARE/BAFU 2007: Le paysage sous pression, suite 3. [Landschaft unter Druck, 3. Fortschreibung]. Federal Office for Spatial Development, Bern and Federal Office for the Environment, Bern. <https://www.aren.admin.ch/aren/de/home/raumentwicklung-und-raumplanung/grundlagen-und-daten/raumb Beobachtung/landschaft-unter-druck.html>

ART 2008a: Uncertainty in agricultural CH₄ and N₂O emissions of Switzerland. Internal documentation by Bretscher, D. and Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich. <http://www.climatereporting.ch>

ART 2009b: Emission factor drained peatlands Switzerland – A brief analysis of recent studies and comparison to EF used in the Swiss GHG Inventory. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich.
<http://www.climatereporting.ch>

ART 2011a: Summary of the available published data on root biomass and root carbon in Swiss grasslands. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich. <http://www.climatereporting.ch>

ART 2011b: First estimate on CO₂ emission factor of organic soils under unproductive wetland. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich. <http://www.climatereporting.ch>

ART 2012a: Carbon in Living Biomass and Dead Organic Matter in Croplands and Grasslands: Current state of the Swiss GHG-Inventory. Internal documentation by Bretscher, D., Agroscope Reckenholz-Tänikon Research Station, Zürich [confidential/internal].

ART/SHL 2012: Categorization of livestock animals in Switzerland. D. Bretscher and T. Kupper. Agroscope Research Station Zürich (ART), Schweizerische Hochschule für Landwirtschaft Zollikofen (SHL). March 2012. <http://www.climatereporting.ch>

AWEL 2009: Qualitätsentwicklung konventionell ausgetragener Schlacke. Zurich, April 2009.
http://www.awel.zh.ch/internet/audirektion/awel/de/abfall_rohstoffe_altlasten/veroeffentlichungen/jcr_content/contentPar/publication_2/publicationitems/schlacke_qualitaetsentwicklung/download.spooler.download.1320054462148.pdf/qualitaetsentwicklung_austrag_schlacke_2009.pdf

Bader, M.K., Leuzinger, S., Keel, S.G., Siegwolf, T.W., Hagedorn, F., Schleppei, P., Körner, C. 2013: Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project. *Journal of Ecology* 101 (6): 1509–1519.
<http://dx.doi.org/10.1111/1365-2745.12149>

Bamberger, I., Stieger, J., Buchmann, N., Eugster, W. 2014: Spatial variability of methane: Attributing atmospheric concentrations to emissions. *Environmental Pollution* 190: 65–74. <http://dx.doi.org/10.1016/j.envpol.2014.03.028>

Brändli, U.-B. (2014): Mehr Laubbäume und weniger Fichtenholz im Mittelland. Medienmitteilung 06.11.2014, Birmensdorf. http://www.wsl.ch/medien/news/landesforstinventar_2014/index_DE

Brändli, U.-B. (Red.) 2010: Schweizerisches Landesforstinventar. Ergebnisse der dritten Aufnahme 2004–2006. [Results of the third Swiss national forest inventory 2004–2006]. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft, Birmensdorf. Bundesamt für Umwelt, Bern, 312 S. <http://www.lfi.ch/publikationen/publ/lfi3-fr.php>

Brändli, U.-B., Speich, S. 2011: Das LFI4 – eine kontinuierliche Erhebung. *LFI-News* 13. <http://www.lfi.ch/publikationen/publ/lfi-info/LFIinfo13.pdf>

Brassel, P., Brändli, U.-B. 1999: Schweizerisches Landesforstinventar. Ergebnisse der Zweitaufnahme 1993–1995. [Results of the second Swiss national forest inventory 1993–1995]. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft, Birmensdorf. Bundesamt für Umwelt, Wald und Landschaft, Bern. Haupt, Bern, Stuttgart, Wien [available in German, French and Italian]. <http://www.lfi.ch/publikationen/publ/lfi2-en.php>

Braun, S., Flückiger, W. 2012: Bodenversauerung in den Flächen des Interkantonalen Walddauerbeobachtungsprogramms. *Schweizerische Zeitschrift für Forstwesen* 163: 374–382. <http://dx.doi.org/10.3188/szf.2012.0374>http://www.iap.ch/publikationen/szf163_braun_versauerung.pdf

Braun, S., Thomas, V.F., Quiring, R., Flückiger, W. 2010: Does nitrogen deposition increase forest production? The role of phosphorus. *Environmental Pollution* 158: 2043–2052. <http://dx.doi.org/10.1016/j.envpol.2009.11.030>

Bühlmann, T. 2014: Durch Stickstoffdeposition induzierte Emissionen von Stickoxiden und Lachgas aus (semi-)natürlichen Ökosystemen. Analyse der schweizerischen Situation mit Vorschlägen zur Umsetzung im Schweizerischen Emissionsinventar (EMIS). Bundesamt für Umwelt (BAFU), Juli 2014. Bern; Schweiz [confidential/internal].

Bühlmann, T., Hiltbrunner, E., Körner, Ch., Rihm, B., Achermann, B. 2015: Induction of indirect N₂O and NO emissions by atmospheric nitrogen deposition in (semi-)natural ecosystems in Switzerland. *Atmospheric Environment* 103: 94–101. <http://dx.doi.org/10.1016/j.atmosenv.2014.12.037>

BUS 1978: Stand der kommunalen Abfallbeseitigung in der Schweiz. Bern, 1. Januar 1978, Eidg. Amt für Umweltschutz, 1978 [hardcopy].

BUS 1984: Abfallerhebung, Schriftenreihe Umweltschutz, Nr. 27, 1984 (ergänzt 1987). <http://www.climatereporting.ch>

BUS 1986: Leitbild für die Schweizerische Abfallwirtschaft. [Lignes directrices pour la gestion des déchets en Suisse]. Schriftenreihe Umweltschutz Nr. 51. Bundesamt für Umweltschutz BUS. [Les cahiers de l'environnement No 51. Office fédéral de la protection de l'environnement], Bern. <http://www.climatereporting.ch>

BUS 1988: Luftreinhaltung beim Verbrennen von Spitalabfällen, Bulletin 1/88. Bern, 1988.

Carbotech 2015: Swiss Greenhouse Gas Inventory 2013: PFCs, HFCs, NF₃ and SF₆ Emissions. Confidential report no. 251.16. For internal use on behalf of the Federal Office for the Environment, Bern. Carbotech, Basel [confidential/internal].

Carbotech 2016a: CO₂ aus VOC-Verbrennung Confidential report no. 129.15. For internal use on behalf of the Federal Office for the Environment, Bern. Carbotech Basel [confidential/internal].

Cemsuisse 2010a: Bestimmung der Emissionsfaktoren an Mischproben, Interner Bericht von Wessling im Auftrag von cemsuisse, 2010 [confidential/internal].

CH2011 2011: Swiss Climate Change Scenarios CH2011. Published by C2SM, MeteoSwiss, ETH, NCCR Climate, and OcCC, Zurich, Switzerland: 1–88. <http://www.ch2011.ch>

Consaba 2016: Erhebung Verwertung Deponiegas über Fackelanlagen in der Schweiz 1990 bis 2014. Consaba GmbH (A. Bauen) im Auftrag des Bundesamtes für Umwelt (BAFU), Bern, Schlussbericht, 23.09.2016. <http://www.climatereporting.ch>

Covington, W.W. 1981: Changes in forest floor organic matter and nutrient content following clear cutting in northern hardwoods. Ecology 62(1): 41–48. <http://www.jstor.org/pss/1936666>

CRC Handbook of Chemistry and Physics 2008: Chemical Rubber Company. CRC Handbook of Chemistry and Physics : A Ready-reference Book of Chemical and Physical Data. D. R. Lide: 84th edition 2003-2004.

Crutzen, P.J., Aselmann, I., Seiler, W. 1986: Methane production by domestic animals, wild ruminants, other herbivorous fauna, and humans. Tellus 38B(3–4): 271–284. <http://dx.doi.org/10.1111/j.1600-0889.1986.tb00193.x>

CSD 2013: Swiss Greenhouse Gas Inventory Submission 2012. National Review of Sector Industrial Processes. On behalf of the Federal Office for the Environment, Bern. Basel. <http://www.climatereporting.ch>

Davidson, E.A., Janssens, I.A. 2006: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. Nature 440: 165–173. <http://dx.doi.org/10.1038/nature04514>

DDPS 2014a: List of military non-road machinery (stocks, engine-power, annual operating hours). E-Mail C. Stucki (Federal Department of Defence, Civil Protection and Sport) to B. Notter (INFRAS) 27.05.2014 [confidential/internal].

DDPS 2016: Consumption of aviation fuel and jet kerosene of Swiss military aircraft 1990–2015. Didier Castella (Federal Department of Defence, Civil Protection and Sport) to Anouk-Aimée Bass (FOEN, Bern), 21.09.2016 [confidential/internal].

de Keizer, C., Ramirez, A., van der Sluijs, J. 2007: Uncertainty Ranges and Correlations Assumed in Tier 2 Studies of Several European Countries. In: Proceedings of the 2nd International Workshop on Uncertainties in Greenhouse Gas Inventories, 27–28 September 2007, Laxenburg, Austria: 35–39. <http://www.ibspan.waw.pl/ghg2007/GHG-total.pdf>

Didion, M. 2014: Impacts of forest management on carbon stock changes in litter and soil in Swiss forests. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern. <http://www.climatereporting.ch>

- Didion, M., Frey, B., Rogiers, N., Thürig, E. 2014a:** Validating tree litter decomposition in the Yasso07 carbon model. *Ecological Modelling* 291: 58–68.
<http://dx.doi.org/10.1016/j.ecolmodel.2014.07.028>
- Didion, M., Kaufmann, E., Thürig, E. 2012:** Estimation of carbon stocks and stock changes in soil, LFH layer and deadwood in Swiss forests with Yasso07. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern. <http://www.climatereporting.ch>
- Didion, M., Kaufmann, E., Thürig, E. 2013:** Data on soil carbon stock change, carbon stock and stock change in surface litter and in coarse deadwood prepared for the Swiss GHGI 1990–2012. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
<http://www.climatereporting.ch>
- Didion, M., Thürig, E. 2013:** Revision der CC13 (Harmonizing activity data from the Swiss land use statistics (AREA) with the Kyoto Protocol guidelines). Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern. <http://www.climatereporting.ch>
- Didion, M., Thürig, E. 2016:** Data on soil carbon stock change, carbon stock and stock change in surface litter and in coarse dead wood prepared for the Swiss GHGI 2017 (1990–2015). Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
<http://www.climatereporting.ch>
- Dobbertin, M., Jüngling, E. 2009:** Totholzverwitterung und C-Gehalt. Zwischenergebnisse. [Wood density and carbon content with changing degree of dead wood decay: First results]. Swiss Federal Research Institute for Forest, Snow and Landscape Research, Birmensdorf.
<http://www.climatereporting.ch>
- Düggelin, C., Abegg, M. 2011:** Modelle zur Biomasse- und Holzvolumenschätzung im Schweizer Gebüschwald. *Schweizerische Zeitschrift für Forstwesen* 162 (2): 32–40.
<http://dx.doi.org/10.3188/szf.2011.0032>
- EAFV (Eidg. Anstalt für das forstliche Versuchswesen) / BFL (Bundesamt für Forstwesen und Landschaftsschutz) (eds.) 1988:** Schweizerisches Landesforstinventar. Ergebnisse der Erstaufnahme 1982–1986. [Results of the first Swiss national forest inventory 1982–1986]. Ber. Eidgenöss. Forsch.anst. Wald Schnee Landsch. 305.
<http://www.lfi.ch/publikationen/publ/lfi1-en/php>
- EC 2004:** Directive 2004/42/CE of the European Parliament and of the Council of 21 April 2004 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC. Official Journal of the European Union. <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32004L0042&qid=1455622565462&from=EN>
- EMEP/EEA 2002:** EMEP/CORINAIR Emission Inventory Guidebook. European Environment Agency. 3rd edition October 2002 update.
<http://www.eea.europa.eu/publications/EMEPCORINAIR3/page002.html>
- EMEP/EEA 2013:** Air Pollutant Emission Inventory Guidebook 2013, European Environment Agency. <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> [1.02.2017]

EMIS 2017/(NFR-Code): Comments to EMIS (Swiss Emission Information System) database. Internal documents. Federal Office for the Environment, Bern [confidential/internal].

EMPA 1999: Written communication from Dr. H.W. Jäckle (EMPA, Dübendorf) to Andreas Liechti (FOEN, Bern), 09.03.1999. <http://www.climatereporting.ch>

EPA 1992: Emission factors for automobile body incineration. EPA - 1992 - AP-42 - Vol. I - CH 2.6. October 1992. <https://www3.epa.gov/ttn/chieff/ap42/ch02/final/c02s06.pdf>

EPA 2010: Exhaust and Crankcase Emission Factors for Nonroad Engine Modelling – Compression Ignition. EPA-420-R-10-018 / NR-009d. Assessment and Standards Division Office of Transportation and Air Quality U.S. Environmental Protection Agency. July 2010. <http://www.epa.gov/otaq/models/nonrdmdl/nonrdmdl2010/420r10018.pdf> [1.02.2017]

ERM 2016: Bauabfälle in der Schweiz – Tiefbau. Aktualisierung 2015. Schlussbericht. Energie- und Ressourcen-Management GmbH, Freienbach, Mai 2016. https://www.bafu.admin.ch/dam/bafu/de/dokumente/abfall/externe-studien-berichte/studie_bauabfaelle_tiefbau_2015.pdf

Estermann, B. L., Wettstein, H.-R., Sutter, F., Kreuzer, M. 2001: Nutrient and energy conversion of grass-fed dairy and suckler beef cattle kept indoors and on high altitude pasture. Animal Research 50(6): 477–493. <http://dx.doi.org/10.1051/animres:2001109>

Etzold, S., Ruehr, N.K., Zweifel, R., Dobbertin, M., Zingg, A., Pluess, P., Häsler, R., Eugster, W. Buchmann, N. 2011: The Carbon Balance of Two Contrasting Mountain Forest Ecosystems in Switzerland: Similar Annual Trends, but Seasonal Differences. Ecosystems 14: 1289–1309. <http://dx.doi.org/10.1007/s10021-011-9481-3>

EV 2014: Jahresbericht 2013. Erdöl-Vereinigung [Rapports annuel 2013. L'Union Pétrolière]. Zürich. https://www.erdoel.ch/images/com_evdocs/1621_ev_ib13_d.pdf

EV 2016: Jahresbericht 2015. Erdöl-Vereinigung [Rapports annuel 2015. L'Union Pétrolière]. Zürich https://www.erdoel.ch/images/com_evdocs/ev_ib_2015_de.pdf [09.02.2017]

EV2015a: Tanktourismus und Eurokurs. Mario Keller, MK Consulting GmbH im Auftrag der Erdöl-Vereinigung. Bern, 18. Dez. 2015. https://www.erdoel.ch/images/pdf/Studie_Tanktourismus_d.pdf

FAL/RAC 2001: Grundlagen für die Düngung im Acker- und Futterbau 2001. [Principles of fertilization in crop and feed production]. Eidgenössische Forschungsanstalt für Agrarökologie und Landbau / Eidgenössische Forschungsanstalt für Pflanzenbau, Agrarforschung, June 2001, Zürich-Reckenholz, Nyon. <http://www.climatereporting.ch> [available in German and French]

Falloon, P., Smith, P. 2003: Accounting for changes in soil carbon under the Kyoto protocol: need for improved long-term data sets to reduce uncertainty in model projections. Soil Use and Management 19(3): 265–269. <http://dx.doi.org/10.1079/SUM2003201>

FCA 2016: Mineralölsteuer: Versteuerte Mengen 2014. Federal Customs Administration (FCA)

FCA 2016a: Bunkers of diesel oil 1990–2015. Written communication Wolfgang Kobler, Federal Customs Administration (FCA) to Anouk Bass (FOEN), 30.09.2016 [confidential/internal]

FCA 2016b: Import- und Exportstatistik Swiss-Impex. Tarifnummer 2836 – carbonate <https://www.swiss-impex.admin.ch> [24.02.2017]

FEDPOL 2016: Bundesamt für Polizei fedpol, Statistik Pyrotechnikumsatz in der Schweiz. https://www.fedpol.admin.ch/fedpol/de/home/sicherheit/sprengestoff_pyrotechnik/pyrotechnische_gegenstaende/pyrotechnikumsatz.html

FEDRO 2010: Swiss Automatic Road Traffic Counts (SARTC). Federal Roads Office, Bern. System description: <https://www.astra.admin.ch/astra/en/home/documentation/traffic-data/data-and-publication/swiss-automatic-road-traffic-counts--sartc-.html> [08.02.2017].

Data 2009:

https://www.astra.admin.ch/dam/astra/en/dokumente/verkehrsdaten/jahresergebnisse2009.xls.download.xls/annual_results_2009.xls

FEDRO 2014: Extract 2AA of MOFIS database (reference date 30.09.2013) of the Federal Roads Office (FEDRO) containing vehicle stock numbers. E-mail from A. Lehmann (FEDRO) to P. Wüthrich, INFRAS on 14.11.2014 [hardcopy].

Fellner et al. 2007: A New Method to Determine the ratio of Electricity Production from Fossil and Biogenic Sources in Waste-to-Energy Plants. Environ. Sci. Technol. 2007, 41, 2579–2586. <http://dx.doi.org/10.1021/es0617587>

Fernández, C., Espinós, F.J., López, M.C., García-Diego, F.J., Cervera, C. 2013: Representation of a mathematical model to predict methane output in dairy goats. Computers and Electronics in Agriculture 91(0): 1–9. <http://dx.doi.org/10.1016/j.compag.2012.11.013>

Fial 2013: Die Schweizer Nahrungsmittel-Industrie im Jahr 2012. http://fial.ch/de/statistics/fial_Statistik%202012_de.pdf

Flisch, R., Sinaj, S., Charles, R., Richner, W. 2009: Grundlagen für die Düngung im Acker- und Futterbau 2009. Forschungsanstalt Agroscope Changins-Wädenswil ACW und Agroscope Reckenholz-Tänikon ART, Agrarforschung 16 (2). http://www.agrarforschungschweiz.ch/archiv_11de.php?jahr=2009&band=16&heft=02

FOCA 1991: Crossair confidential data 1991. Federal Office of Civil Aviation, Bern [confidential/internal].

FOCA 1991a: L'aviation civile Suisse en 1990. Federal Office of Civil Aviation, Bern. <http://www.climatereporting.ch>

FOCA 2004: Unternehmensstatistik der Schweizerischen Helikopterunternehmen. Federal Office of Civil Aviation, Bern. <http://www.climatereporting.ch>

FOCA 2006: GHG emissions of Swiss civil aircraft in 1990 and 2004: data, proceeding and description of the calculations. Written communication from Theo Rindlisbacher and Paul Stulz (FOCA, Bern) to Andreas Liechti (FOEN, Bern), 20./22.02.2006. <http://www.climatereporting.ch>

FOCA 2006a: GHG emissions of Swiss civil aircraft in 1990, 1995, 2000, 2002, 2004 and 2005: data, proceeding and description of the calculations. Written communication from Theo Rindlisbacher (FOCA, Bern) to Paul Filliger (FOEN, Bern), 17.11.2006.

<http://www.climatereporting.ch>

FOCA 2007: GHG emissions of Swiss civil aircraft in 2006. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 03.12.2007.

<http://www.climatereporting.ch>

FOCA 2007a: Aircraft Piston Engine Emissions. Summary Report. Federal Office of Civil Aviation. Report Ref. 0 / 3/33/33-05-003 ECERT. Bern. <http://www.climatereporting.ch>

FOCA 2007b: Validation of ADAECAM (Advanced Aircraft Emission Calculation Method). Report on fuel calculation. Federal Office of Civil Aviation. Report Ref. 0/3/33/33-05-007.021. Bern, 10.08.2007. <http://www.climatereporting.ch>

FOCA 2008: GHG emissions of Swiss civil aircraft in 2007. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 01.12.2008.

<http://www.climatereporting.ch>

FOCA 2009: GHG emissions of Swiss civil aircraft in 2008. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 04.11.2009.

<http://www.climatereporting.ch>

FOCA 2009a: Guidance on the determination of helicopter emissions. Federal Office of Civil Aviation. Report Ref. 0/3/33/ 33-05-020. Edition 1. Bern. March 2009.

<http://www.climatereporting.ch>

FOCA 2010: GHG emissions of Swiss civil aircraft in 2009. Written communication from Theo Rindlisbacher (FOCA, Bern) to Sophie Hoehn (FOEN, Bern), 18.11.2010.

<http://www.climatereporting.ch>

FOCA 2011: GHG emissions of Swiss civil aircraft in 2010. Written communication from Theo Rindlisbacher (FOCA, Bern) to Sophie Hoehn (FOEN, Bern), 09.11.2011.

<http://www.climatereporting.ch>

FOCA 2012: GHG emissions of Swiss civil aircraft in 2011. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 07.09.2012.

<http://www.climatereporting.ch>

FOCA 2013: GHG emissions of Swiss civil aircraft in 2012. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 07.11.2013.

<http://www.climatereporting.ch>

FOCA 2014: GHG emissions of Swiss civil aircraft in 2013 and corrections for the year 2011. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 30.10.2014 [confidential/internal].

FOCA 2015: GHG emissions of Swiss civil aircraft in 2014. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 29.10.2015 [confidential/internal].

FOCA 2015a: Guidance on the Determination of Helicopter Emissions. Federal Office of Civil Aviation. Reference: COO.2207.111.2.2015750. Edition 2. Bern. December 2015.

<https://www.bazl.admin.ch/bazl/en/home/specialists/regulations-and-guidelines/environment/pollutant-emissions/aircraft-engine-emissions/guidance-on-the-determination-of-helicopter-emissions.html> [28.03.2017]

FOCA 2016: GHG emissions of Swiss civil aircraft in 2015. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 6.12.2016 [confidential/internal].

FOEN 2006h: Switzerland's Initial Report under Article 7, paragraph 4 of the Kyoto Protocol. Federal Office for the Environment, Bern. Including the Update following the UNFCCC review (FCCC/IRR/2007/CHE). <http://www.climatereporting.ch>

FOEN 2008: Switzerland's Greenhouse Gas Inventory 1990–2006, National Inventory Report and CRF-tables 2008. Submission of 15 April 2008 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2008h: Wood Resource Policy: Strategy, Objectives and Action Plan for the Resource Wood. Federal Office for the Environment, Bern. <https://www.bafu.admin.ch/ud-1001-e>

FOEN 2010d: Deforestations in Switzerland as reported under the Kyoto Protocol Art. 3.3. Forest Division, Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2010h: Afforestations in Switzerland as reported under the Kyoto Protocol Art. 3.3. Forest Division, Federal Office for the Environment, Bern. <http://www.bafu.admin.ch/ghginv-ref> <http://www.climatereporting.ch>

FOEN 2010i: Pollutant Emissions from Road Transport, 1990 to 2035. Updated in 2010. Environmental studies no. 1021. Federal Office for the Environment, Bern. <https://www.bafu.admin.ch/uw-1021-e>

FOEN 2011: Switzerland's Greenhouse Gas Inventory 1990–2009: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008–2009, SEF and SIAR tables from the National Registry. Submission of 15 April 2011 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2011i: Switzerland's Submission on Reference Levels as an accounting approach for Forest Management under the Kyoto Protocol. Submission of 28 February 2011 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. http://unfccc.int/files/meetings/ad_hoc_working_groups/kp/application/pdf/awqkp_switzerland_2011.pdf

FOEN 2012: Switzerland's Greenhouse Gas Inventory 1990–2010: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008–2010, SEF and SIAR tables from the National Registry. Submission of 13 April 2012 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2012b: Anpassung an den Klimawandel in der Schweiz – Ziele, Herausforderungen und Handlungsfelder. Erster Teil der Strategie des Bundesrates. Federal Office for the Environment, Bern. <http://www.bafu.admin.ch/klimaanpassung/11529>

FOEN 2012h: Adressliste der Schweizer Kläranlagen mit Angaben zur Ausbaugrösse. Swiss Federal Office of Energy, Bern.

https://www.bafu.admin.ch/dam/bafu/de/dokumente/wasser/fachinfo-daten/adressliste_der_schweizerklaeranlagenmitangabenzurausbaugroesse.pdf

FOEN 2013: Switzerland's Greenhouse Gas Inventory 1990–2011: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008–2010, SEF and SIAR tables from the National Registry. Submission of April 2013 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2013q: Guidance document on activity specific monitoring methods, Appendix G, Federal Office for the Environment, Bern.

FOEN 2014d: Switzerland's Sixth National Communication under the UNFCCC; Third National Communication under the Kyoto Protocol to the UNFCCC. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2014f: Illustration of unproductive forests in Switzerland. Internal report by the Swiss Federal Office for the Environment. <http://www.climatereporting.ch>

FOEN 2014i: Kohlenstoffgehalt Abfall auf Basis des Heizwerts – Berechnungsgrundlage, FOEN, Bern. 31.10.2014 [confidential/internal].

FOEN 2014n: Standardmethode für Kompensationsprojekte des Typs „Landwirtschaftliche Biogasanlagen“. Version für informelle Konsultation, 1.5.2014. Federal Office for the Environment, Geschäftsstelle Kompensation, Bern [confidential/internal].

FOEN 2014o: Erhebung der Kehrrichtzusammensetzung 2012. Federal Office for the Environment, Bern.

<http://www.news.admin.ch/NSBSubscriber/message/attachments/33597.pdf>
<http://www.news.admin.ch/NSBSubscriber/message/attachments/33601.pdf>

FOEN 2014p: Von Arx U.: Feuerwerkskörper. Umweltauswirkungen und Sicherheitsaspekte. Umwelt-Wissen Nr.1423. Federal Office for the Environment, Bern.
<https://www.bafu.admin.ch/uw-1423-d>

FOEN 2015: Switzerland's Greenhouse Gas Inventory 1990–2013: National Inventory Report, CRF-tables. Submission of 15 April 2015 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2015d: Faktenblatt. CO₂-Emissionsfaktoren des Treibhausgasinventars der Schweiz. Federal Office for the Environment, Bern.
<https://www.bafu.admin.ch/dam/bafu/en/dokumente/klima/fachinfo-daten/co2-emissionsfaktorendesschweizerischentreibhausgasinventars.pdf>

FOEN 2015g: Zusammenstellung der Feedstocks basierend auf GEST-, EV- und bottom-up Industrie-Daten, BAFU, internal documentation (confidential, Excelfile), Federal Office for the Environment, Bern [confidential/internal].

FOEN 2015j: Non-road energy consumption and pollutant emissions. Study for the period from 1980 to 2050 (Original in German: Energieverbrauch und Schadstoffemissionen des Nonroad-Sektors. Studie für die Jahre 1980–2050. Schlussbericht). INFRAS. Ed. by the Federal Office for the Environment. <https://www.bafu.admin.ch/uw-1519-e> [8.02.2017]

FOEN 2015n: Biological Treatment of Solid Waste (5B) – Revision 2015 (internal documentation) [confidential/internal].

FOEN 2015p: Ergebnisse der Nationalen Bodenbeobachtung (NABO) 1985-2009. Zustand und Veränderungen der anorganischen Schadstoffe und Bodenbegleitparameter. Umwelt-Zustand Nr. 1507. Federal Office for the Environment, Bern. <http://www.bafu.admin.ch/uz-1507-f> [French] <https://www.bafu.admin.ch/uz-1507-d> [German]

FOEN 2016: Switzerland's Greenhouse Gas Inventory 1990–2014: National Inventory Report, CRF-tables. Submission of 15 April 2016 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2016c: Switzerland's Second Initial Report under the Kyoto Protocol. Report to facilitate the calculation of the assigned amount pursuant to Article 3, paragraphs 7bis, 8 and 8bis, of the Kyoto Protocol for the second commitment period 2013–2020. <http://www.climatereporting.ch><http://www.climatereporting.ch/>

FOEN 2016d: Switzerland's Second Initial Report under the Kyoto Protocol. Update following the in-country review by an expert review team coordinated by the UNFCCC secretariat. <http://www.climatereporting.ch>

FOEN 2016g: UNFCCC FOD spreadsheet model Switzerland for estimating methane emissions from landfill sites [confidential/internal].

FOEN 2016h: CO₂-Emissionsfaktor Erdgas, Zeitreihe für das Treibhausgasinventar der Schweiz, internal documentation, Federal Office for the Environment, Bern [confidential/internal].

FOEN 2016i: Abfallstatistiken des Jahres 2015 sowie Zeitreihen. Diverse Dateien. https://www.bafu.admin.ch/bafu/de/home/themen/abfall/zustand/daten/Abfallstatistiken_2015.html [14.02.2017] <https://www.bafu.admin.ch/bafu/de/home/themen/abfall/zustand/daten/abfallstatistiken--zeitreihen.html> [14.02.2017]

FOEN 2016j: Harvested wood products (HWP) in the Swiss GHG inventory – Description of time series and modeling methodology for accounting under the Kyoto Protocol and results. Internal report, Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

FOEN 2016k: Jahrbuch Wald und Holz 2015. Umwelt-Zustand Nr. 1520 [Annuaire La forêt et le bois 2015. Etat de l'environnement no 1520]. Federal Office for the Environment, Bern. <http://www.bafu.admin.ch/uz-1520-d> [German] <http://www.bafu.admin.ch/uz-1520-f> [French]

FOEN 2017a: Qualitätsmanagement Treibhausgasinventar – Handbuch. BAFU, Bern [confidential/internal].

FOEN 2017b: Handbuch Berechnung der Wald-Emissionsfaktoren fürs Inventarjahr 2015 [Manual: Calculation of EFs for Forest Land]. Including 1 Excel data file. Internal document. Federal Office for the Environment, Bern [confidential/internal].

FOEN 2017c: Handbuch Anleitung zum Ausfüllen der Kyoto Tabellen Wald fürs Inventarjahr 2015. [Instruction manual for completing data of forest related activities in Kyoto Tables]. Internal document, Federal Office for the Environment, Bern [confidential/internal].

FOEN 2017f: Switzerland's Informative Inventory Report 2017 (IIR), Submission under the UNECE Convention on Long-range Transboundary Air Pollution, Submission of March 2017 to the United Nations ECE Secretariat. Federal Office for the Environment, Bern.

https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/fachinfo-daten/switzerland_s_informativeinventoryreport2016iir.pdf

FOEN/SFOE/VBSA 2016: Einheitliche Heizwert- und Energiekennzahlenberechnung der Schweizer KVA nach europäischem Standardverfahren. Resultate 2015. Bern. 6. April 2016.

http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_355_210852.pdf&endung=Einheitliche

Gehrig-Fasel, J., Guisan, A., Zimmermann, N.E. 2007: Treeline shifts in the Swiss Alps: climate change or land abandonment? *Journal of Vegetation Science* 18: 571–582.

<http://dx.doi.org/10.1111/j.1654-1103.2007.tb02571.x>

Genossenschaft Ökostrom Schweiz 2014: Projekte zur Emissionsverminderung im Inland – Methanemissionsreduktion bei landwirtschaftlichen Biogasanlagen in der Schweiz (Bündel I) – Monitoringberichte zu Handen BAFU für die Jahre 2010, 2011 und 2012. Frauenfeld, Schweiz [confidential/internal].

GES Biogas GmbH 2014: Projekte zur Emissionsverminderung im Inland – Methanemissionsreduktion bei landwirtschaftlichen Biogasanlagen in der Schweiz (Bündel II) – Monitoringberichte zu Handen BAFU für die Jahre 2011 und 2012. Zürich, Schweiz [confidential/internal].

Ginzler, C. 2014: Crown cover and basal area in unproductive forest: special analysis of plots covered by aerial photographs and terrestrial NFI. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern [confidential/internal].

Giuliani, S. 2016: Energiebedarfswerte der Nutztiere für die Periode 2010–2013. Written communication from Silvano Giuliani (SBV, Swiss Farmers Union) to Daniel Bretscher (ART, Reckenholz), 14.09.2015 [confidential/internal].

Grassi, G., Blujdea, V. 2011: Pools to be reported under KP-LULUCF. Harmonized guidance and decision tree on the application of "not a source" principle. Joint Research Centre, Ispra. Presentation at "JRC technical workshop on LULUCF issues under the Kyoto Protocol", 21 November 2011. <http://slideplayer.com/slide/743707/>

Gschwantner, T. 2006: Zuwachsänderungen nach den Daten der Österreichischen Waldinventur und ihre klimatischen Ursachen. [Growth changes according to the data of the Austrian National Forest Inventory and their climatic causes]. BFW-Berichte 133, Wien, Bundesamt für Wald [hardcopy].

Guidi, C., Vesterdal, L., Gianelle, D., Rodeghiero, M. 2014: Changes in soil organic carbon and nitrogen following forest expansion on grassland in the Southern Alps. *Forest Ecology and Management* 328: 103–116. <http://dx.doi.org/10.1016/j.foreco.2014.05.025>

Hackl, A., Mauschwitz, G. 2003: Emissionen aus Anlagen der Österreichischen Zementindustrie IV. Jahresreihe 2000–2002. Weitra/Wien. <http://www.climatereporting.ch>

Hadorn, R., Wenk, C. 1996: Effect of different sources of dietary fibre on nutrient and energy utilization in broilers. 2. Energy and N-balance as well as whole body composition. *Archiv für Geflügelkunde* 60: 22–29 [hardcopy].

- Hagedorn, F., Martin, M., Rixen, C., Rusch, S., Bebi, P., Zürcher, A., Siegwolf, R.T.W., Wipf, W., Escape, C., Roy, J., Hättenschwiler, S. 2010:** Short-term responses of ecosystem carbon fluxes to experimental soil warming at the Swiss alpine treeline. *Biogeochemistry*: 97: 7–19. <http://dx.doi.org/10.1007/s10533-009-9297-9>
- Henne, S., Brunner, D., Steinbacher, M., Emmenegger, L. 2015:** Verifikation des Schweizer Methan-Emissionsinventars. Eidgenössische Materialprüfungs- und Forschungsanstalt (EMPA), Abteilung Luftfremdstoffe / Umwelttechnik. Dübendorf; Schweiz. <http://www.climatereporting.ch>
- Hiller, R.V., Bretscher, D., DeISontro, T., Diem, T., Eugster, W., Henneberger, R., Hobi, S., Hodson, E., Imer, D., Kreuzer, M., Künzle, T., Merbold, L., Niklaus, P.A., Rihm, B., Schellenberger, A., Schroth, M.H., Schubert, C.J., Sigrist, H., Stieger, J., Buchmann, N., Brunner, D. 2014:** Anthropogenic and natural methane fluxes in Switzerland synthesized within a spatially explicit inventory. *Biogeosciences* 11: 1941–1959. <http://dx.doi.org/10.5194/bg-11-1941-2014>
- Hiller, R.V., Neininger, B., Brunner, D., Gerbig, Ch., Bretscher, D., Künzle, T. 2014a:** Airborne CH₄ flux measurements for validation of emissions from an agriculturally dominated area in Switzerland. *Journal of Geophysical Research*. 119(8): 4874–4887. <http://dx.doi.org/10.1002/2013JD020918>
- Hindrichsen, I.K., Wettstein, H.-R., Machmüller, A., Bach Knudsen, K.E., Madsen, J., Kreuzer, M. 2006:** Methane emission, nutrient degradation and nitrogen turnover in dairy cows and their slurry at different milk production scenarios with and without concentrate supplementation. *Agricultural Ecosystems and Environment* 113: 150–161. <http://dx.doi.org/10.1016/j.agee.2005.09.004>
- Högberg, P., Fan, H., Quist, M., Binkley, D., Tamm, C.O. 2006:** Tree growth and soil acidification in response to 30 years of experimental nitrogen loading on boreal forest. *Global Change Biology* 12: 489–499. <http://dx.doi.org/10.1111/j.1365-2486.2006.01102.x>
- Huber, B., Frehner M. 2012:** Forschungsprojekt Grünerle. Bericht erstellt im Auftrag des Bundesamtes für Umwelt (BAFU), Bern. Abenis AG Chur und Forstingenieurbüro Monika Frehner, Sargans. <http://www.climatereporting.ch>
- Huber, B., Frehner M. 2013:** Die Verbreitung und Entwicklung der Grünerlenbestände in der Ostschweiz. *Schweizerische Zeitschrift für Forstwesen* 164: 87–94. <http://www.szfi.org/toc/swif/164/4>
- Huber, T., Thürig, E. 2014:** Special analysis of stocks of living biomass in non-productive forest (CC13) on terrestrial NFI plots. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern. <http://www.climatereporting.ch>
- Hubschmid, S. 2014:** VOC-Emissionen aus lösungsmittelhaltigen Haushaltsprodukten, Master Thesis, MAS Umwelttechnik und -management, FHNW [confidential/internal].
- Hürdler, J., Prasuhn, V., Spiess, E. 2015:** Abschätzung diffuser Stickstoff- und Phosphoreinträge in die Oberflächengewässer der Schweiz. MODIFFUS 3.0. – Bericht im Auftrag des Bundesamtes für Umwelt BAFU, Agroscope, Zürich, ca. 115 S. <https://www.agroscope.admin.ch/dam/agroscope/de/dokumente/themen/umwelt-ressourcen/gewaesserschutz/huerdler-prasuhn-spiess-endbericht-modiffus.pdf>

Hyvönen, R., Persson, T., Andersson, S., Olsson, B. Ågren, G.I., Linder, S. 2008: Impact of long-term nitrogen addition on carbon stocks in trees and soils in northern Europe. *Biogeochemistry* 89: 121–137. <http://dx.doi.org/10.1007/s10533-007-9121-3>

IAI 2005: Aluminium for future Generations. Sustainability Update 2004. International Aluminium Institute, London. <http://www.climatereporting.ch>

IEA 2005: Energy statistics manual, IEA/OECD Paris. http://www.iea.org/publications/freepublications/publication/statistics_manual-1.pdf

IEA 2012: Energy statistics of OECD countries, 2012 edition, IEA/OECD Paris. <http://www.iea.org/stats/index.asp>

IFEU 2010: Aktualisierung des Modells TREMOD – Mobile Machinery (TRE-MOD-MM). Endbericht im Auftrag des Umweltbundesamtes. UBA Texte 28/2010. Institut für Energie- und Umweltforschung (IFEU), Heidelberg. <https://www.umweltbundesamt.de/sites/default/files/medien/461/publikationen/3944.pdf> [1.02.2017]

INFRAS 2010: The Handbook Emission Factors for Road Transport (HBEFA), version 3.1 (MS AccessXP runtime application and report). INFRAS in cooperation with further editors: FOEN/Switzerland; Umweltbundesamt Dessau/Germany; Umweltbundesamt Wien/Austria; Swedish Road Administration, ADEME/France; SFT/Norway. Bern. 30.01.2010. <http://www.hbefa.net/e/index.html> [queries from database].

INFRAS 2011a: Bunker fuels Bodensee/Genfersee, Datenverfügbarkeit und Berechnung der Bunker Fuels. Internal document. Federal Office for the Environment. September 2011 [confidential/internal].

INFRAS 2012: Verification of Swiss implied emission factors. INFRAS, Zürich. <http://www.climatereporting.ch>

INFRAS 2014: Verbrennung natürlicher Wald-, Feld- und Gartenabfälle im Freien. Datengrundlagen zur Aktualisierung der Jahresleistungen. Zürich, 29. Januar 2014.

INFRAS 2015: CO₂-Emissions from Carbonate, Assessment for FOEN. Internal documentation FOEN [confidential/internal]. <http://www.bafu.admin.ch/ghginv-CP2-ref>

INFRAS 2015a: Online database of emission factors, activity data and emissions of non-road vehicles (corresponding report is FOEN 2015j) <https://www.bafu.admin.ch/bafu/en/home/topics/air/state/non-road-datenbank.html> [queries from database].

Integer 2013: Emissions Control in Non-Road Mobile Machinery (NRMM) Markets: 2013 Edition. Integer Research Limited, London. <https://www.integer-research.com/wp-content/uploads/2014/07/NRMM-markets-2013-Brochure-amended.pdf> [1.02.2017]

IPCC 1997c: Greenhouse Gas Inventory Reference Manual, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Reference Manual (Volume 3). Intergovernmental Panel on Climate Change. <http://www.ipcc-nggip.iges.or.jp/public/ql/invs6.htm>

IPCC 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change. <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm> [11.01.2017]

- IPCC 2014:** 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol (KP Supplement). Intergovernmental Panel on Climate Change. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds). Published: IPCC, Switzerland. <http://www.ipcc-nggip.iges.or.jp/public/kpsqg/index.html> [1.02.2017]
- IPCC 2014a:** 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands, Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds). Published: IPCC, Switzerland. <http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html> [1.02.2017]
- ISPRA 2014:** Italian Greenhouse Gas Inventory 1990–2012. Inventory Report and CRF-tables. Submission to the UNFCCC Secretariat 2014. Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA). Rome, Italy. http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8108.php
- Jandl, R., Lindner, M., Vesterdal, L., Bauwens, B., Baritz, R., Hagedorn, F., Johnson, D. W., Minkinen, K., Byrne, K. A. 2007:** How strongly can forest management influence soil carbon sequestration? *Geoderma* 137: 253–268. <http://dx.doi.org/10.1016/j.geoderma.2006.09.003>
- Jardin Suisse 2012:** Zahlen zum Schweizerischen Gartenbau. Unternehmerverband Gärtner Schweiz. <http://www.climatereporting.ch>
- Jarvis, P.G., Linder, S. 2000:** Constraints to growth of boreal forests. *Nature* 405: 904–905. <http://dx.doi.org/10.1038/35016154>
- Karhu, K., Wall, A., Vanhala, P., Liski, J., Esala, M., Regina, K. 2011:** Effects of afforestation and deforestation on boreal soil carbon stocks – Comparison of measured C stocks with Yass07 model results. *Geoderma* 164: 33–45. <http://dx.doi.org/10.1016/j.geoderma.2011.05.008>
- Karl, T., Guenther, A., Lindinger, C., Jordan, A., Fall, R., Lindinger, W. 2001:** Eddy covariance measurements of oxygenated volatile organic compound fluxes from crop harvesting using a redesigned proton-transfer-reaction mass spectrometer. *Journal of Geophysical Research* 106 D20: 24157–24167. <http://dx.doi.org/10.1029/2000JD000112>
- Kaufmann, E. 2001:** Estimation of Standing Timber, Growth and Cut. In: Brassel, P.; Lischke, H. (eds). *Swiss National Forest Inventory: Methods and Models of the Second Assessment*. Birmensdorf, Swiss Federal Research Institute WSL. 162–196. http://www.lfi.ch/publikationen/publ/LFI2_Methoden.pdf
- Kaufmann, E. 2005:** Personal communication from Edgar Kaufmann (WSL, Zürich) to Esther Thürig (FOEN, Bern), 12.12.2005. <http://www.climatereporting.ch>
- Kaufmann, U. 2015:** Energieverbrauch stationäre Motoren und Gasturbinen ab 1990. Excel data file for internal use on behalf of the Federal Office for the Environment, Bern. Eicher + Pauli, Liestal [confidential/internal].
- Keller, A., Desaulles, A., Schwab, P., Weisskopf, P., Scheid, S., Oberholzer, H.-R. 2006:** Monitoring soil quality in the long term: examples from the Swiss Soil Monitoring Network. *Mitteilungen der Österreichischen Bodenkundlichen Gesellschaft*, Heft 73: 5–12. <http://www.climatereporting.ch>

- Keller, A., Rossier, N., Desales, A. 2005:** Schwermetallbilanzen von Landwirtschaftsparzellen der Nationalen Bodenbeobachtung. NABO – Nationales Bodenbeobachtungsnetz der Schweiz. [Heavy-metal balances of agricultural soil monitoring sites. NABO – Swiss Soil Monitoring Network]. Schriftenreihe der FAL 54, Zürich-Reckenholz. <http://www.climatereporting.ch>
- Keller, J., Andreani-Aksoyoglu, S., Joss, U. 1995:** Inventory of natural emissions in Switzerland. In: Power, H., Moussiopoulos, N., Brebbia C.A. (eds.): Air Pollution III, Volume 2, Air Pollution Engineering and Management: 339–346. Computational Mechanics Publications. <http://www.climatereporting.ch>
- Köck, K., Leifeld, J., Fuhrer, J. 2013:** A model-based inventory of sinks and sources of CO₂ in agricultural soils in Switzerland: development of a concept. Agroscope, Zürich. <http://www.climatereporting.ch>
- Koehlerei, 2014:** Geschichte der Holzköhlerei in Romoos. <http://www.koehlerei.ch>
- Koehli L. 2014:** Landwirtschaftliche Biogasanlagen. Written communication from Lorenz Koehli (Oekostrom Schweiz) to Daniel Bretscher (Agroscope), 27.08.2014 [confidential/internal].
- König, G., Brunda, M., Puxbaum, H., Hewitt, C.N., Duckham, S.C., Rudolph, J. 1995:** Relative contribution of oxygenated hydrocarbons to the total biogenic VOC emissions of selected mid-European agricultural and natural plant species. Atmospheric Environment 29: 861–874. [http://dx.doi.org/10.1016/1352-2310\(95\)00026-U](http://dx.doi.org/10.1016/1352-2310(95)00026-U)
- Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier van der Gon, H. A. C. 2014:** TNO-MACC_II emission inventory; a multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, Atmos. Chem. Phys., 14, 10963–10976, <http://dx.doi.org/10.5194/acp-14-10963-2014>
- Külling D. R., Stadelmann F. X., Candinas T. 2002a:** Nährstoffe und Schwermetalle im Klärschlamm 1975–1999, Agrarforschung 9(5): 200–205. http://www.agrarforschungschweiz.ch/artikel/2002_05_27.pdf
- Külling, D. R., Dohme, F., Menzi, H., Sutter, F., Lischer, P., Kreuzer, M. 2002:** Methane emissions of differently fed dairy cows and corresponding methane and nitrogen emissions from their manure during storage. Environmental Monitoring and Assessment 79 (2): 129–150. <http://dx.doi.org/10.1023/A:1020248700255>
- Külling, D. R., Menzi, H., Sutter, F., Lischer, P., Kreuzer, M. 2003:** Ammonia, nitrous oxide and methane emissions from differently stored dairy manure derived from grass- and hay-based rations. Nutrient Cycling in Agroecosystems 65 (1): 13–22. <http://dx.doi.org/10.1023/A:1021857122265>
- Kupper, T., Bonjour, C., Achermann, B., Zaucker, F., Rihm, B., Menzi, H. 2013:** Ammoniakemissionen in der Schweiz 1990–2010 und Prognose bis 2020. Hochschule für Agrar-, Forst- und Lebensmittelwissenschaften, Zollikofen. <http://www.agrammon.ch/dokumente-zum-download>
- KWF 2012:** Forstmaschinenstatistik zeigt deutliche Stabilisierung des Marktes. 17. KWF-Tagung. <http://www.kwf-tagung.org/fr/aktuelles/news-detailanzeige/eintrag/39.html>
- Lamlom, S.H., Savidge, R.A. 2003:** A reassessment of carbon content in wood: variation within and between 41 North American species. Biomass and Bioenergy 25: 381–388. [http://dx.doi.org/10.1016/S0961-9534\(03\)00033-3](http://dx.doi.org/10.1016/S0961-9534(03)00033-3)

- Lehtonen, A. , Heikkinen, J. 2016:** Uncertainty of upland soil carbon sink estimate for Finland. Canadian Journal of Forest Research 46: 310-322. <http://dx.doi.org/10.1139/cjfr-2015-0171>
- Leifeld, J., Bassin, S., Fuhrer, J. 2003:** Carbon stocks and carbon sequestration potentials in agricultural soils in Switzerland. Schriftenreihe der FAL 44. Zürich-Reckenholz. <http://www.climatereporting.ch>
- Leifeld, J., Bassin, S., Fuhrer, J. 2005:** Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. Agriculture, Ecosystems & Environment 105 (1/2): 255–266. <http://dx.doi.org/10.1016/j.agee.2004.03.006>
- Leifeld, J., Fuhrer, J. 2005:** Greenhouse gas emissions from Swiss agriculture since 1990: Implications for environmental policies to mitigate global warming. Environmental Science & Policy 8: 410–417. <http://dx.doi.org/10.1016/j.envsci.2005.04.001>
- Leifeld, J., Zimmermann, M., Fuhrer, J. 2007:** Characterization of soil carbon stocks and site-specific sequestration potentials of agricultural soils. Extended Summary. Final Report, Contract Number: 810.03.0716 / 2003.C.04. Agroscope Reckenholz-Tänikon Research Station ART. <http://www.climatereporting.ch>
- Leupro 2012:** Entwicklung und Prognose der Emissionsfaktoren Feuerungen für den Zeitraum 1990–2035. <http://www.climatereporting.ch>
- LHA 2004:** Letter of „Lufthygieneamt beider Basel“ to FOEN, Basel, 04.05.2004 [confidential/internal].
- Mangino, J., Bartram, D., Brazy, A. 2001:** Development of a methane factor to estimate emissions from animal waste lagoons. U.S.EPA's 17th Annual Emission Inventory Conference, Atlanta GA; U.S. EPA. <http://www.climatereporting.ch>
- Martínez-Fernández, G., Abecia, L., Ramos-Morales, E., Martín-García, A.I., Molina-Alcaide, E., Yáñez-Ruiz, D.R. 2014:** Effects of propyl propane thiosulfinate on nutrient utilization, ruminal fermentation, microbial population and methane emissions in goats. Animal Feed Science and Technology 191:16–25. <http://dx.doi.org/10.1016/j.anifeedsci.2014.01.008>
- Mathys, L., Thürig E. 2010:** Baumbiomasse in der Landschaft. [Living biomass of trees in Non-Forest Land]. Final report, Sigmaphan and WSL on behalf of the Federal Office for the Environment, Bern. <http://www.climatereporting.ch>
- Meining, S., V.Wilpert, K., Schäffer, J., Schröter, H. 2008:** Waldzustandsbericht 2008 der Forstlichen Versuchs- und Forschungsanstalt Baden-Württemberg. Freiburg <http://www.climatereporting.ch>
- Menzi, H. 2014:** Stickstoffausscheidung Kaninchen. Written communication from Harald Menzi (Agroscope INT) to Daniel Bretscher (Agroscope INH), 05.12.2014 [confidential/internal].
- MeteoSwiss (Federal Office of Meteorology and Climatology) 2014:** Normwerte 1981–2010: Lufttemperatur 2m. Federal Office of Meteorology and Climatology MeteoSwiss, Zurich. <http://www.meteoschweiz.admin.ch/home/klima/vergangenheit/klimanormwerte/normwerte-pro-messgroesse.html>

Meteotest 2017: Treibhausgasinventar Sektor LULUCF, Tabellen zur Berechnung der Treibhausgasemissionen im Sektor LULUCF in der Schweiz inklusive Begleitdokumentation. Internes Dokument. [GHG inventory, sector LULUCF. Spreadsheets to calculate the GHG emissions of the LULUCF sector in Switzerland inclusive accompanying documents. Internal document]. Meteotest, Bern.

Meyer, H., Coenen, M., 2002: Pferdefütterung. Parey Buchverlag Berlin, 4. Auflage; ISBN 3-8263-3398-5 [hardcopy].

Minonzo, G., Grub, A., Fuhrer, J. 1998: Methanemissionen der schweizerischen Landwirtschaft. Schriftenreihe Umwelt Nr. 298. FOEN / BUWAL, Bern [hardcopy].

MISTA 2016: Milchstatistik der Schweiz 2015. Schweizerischer Bauernverband (SBV), TSM Treuhand GmbH, Schweizer Milchproduzenten (SMP).

Moeri, A.C. 2007: Kohlenstoffvorräte in Schweizer Waldböden mit besonderer Berücksichtigung der organischen Auflage. Diplomarbeit bei der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft (WSL). Geographisches Institut der Universität Zürich. <http://www.climatereporting.ch>

Mohn, J. 2011: Bestimmung des Anteils biogener und fossiler CO₂ Emissionen aus Schweizer KVAs. Eidg. Materialprüfungsanstalt und Forschungsanstalt. Schlussbericht. Im Auftrag des Bundesamts für Umwelt. <http://www.climatereporting.ch>

Mohn, J. 2013: Bestimmung der N₂O und CH₄ Emissionen aus Schweizer KVAs. Eidg. Materialprüfungs- und Forschungsanstalt. Schlussbericht. Im Auftrag des Bundesamts für Umwelt. <http://www.climatereporting.ch>

Mohn, J., Szidat, S., Fellner, J., Rechberger, H., Quartier, R., Buchmann, B., Emmenegger, L. 2008: Determination of biogenic and fossil CO₂ emitted by waste incineration based on ¹⁴CO₂ and mass balances. Bioresource Technology 99: 6471–6479 <http://dx.doi.org/10.1016/j.biortech.2007.11.042>

Moller, H.B., Sommer, S.G., Ahring, B.K. 2004: Biological degradation and greenhouse gas emissions during pre-storage of liquid animal manure. Journal of Environmental Quality 33 (1): 27–36. <http://dx.doi.org/10.2134/jeq2004.0027>

Monni, S., Peltoniemi, M., Palosuo, T., Lehtonen, A., Mäkipää, R., Savolainen, I. 2007: Uncertainty of forest carbon stock changes – implications to the total uncertainty of GHG inventory of Finland. Climatic Change 81: 391–413. <http://dx.doi.org/10.1007/s10584-006-9140-4>

Morel, I., Chassot, A., Schlegel, P., Jans, F., Kessler, J. 2015: Fütterungsempfehlungen für die Mutterkuh. Fütterungsempfehlungen für Wiederkäuer (Grünes Buch). Agroscope Posieux. Posieux; Schweiz.

Nave, L.E., Vance, E.D., Swanston, C. W., Curtis, P.S. 2010: Harvest impacts on soil carbon storage in temperate forests. Forest Ecology and Management 259: 857–866. <http://dx.doi.org/10.1016/j.foreco.2009.12.009>

Nousiainen, J., Rinne, M., Huhtanen, P. 2009: A meta-analysis of feed digestion in dairy cows. 1. The effects of forage and concentrate factors on total diet digestibility. Journal of dairy science 92(10): 5019–5030. <http://dx.doi.org/10.3168/jds.2008-1833>

- Nowak, D., Crane, D. 2002:** Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution* 116: 381–389. [http://dx.doi.org/10.1016/S0269-7491\(01\)00214-7](http://dx.doi.org/10.1016/S0269-7491(01)00214-7)
- Nussbaum, M., Papritz, A., Baltensweiler, A., Walthert, L. 2012:** Organic carbon stocks of Swiss forest soils. Final report, including CD ROM. Institute of Terrestrial Ecosystems, ETH Zürich and Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. <http://dx.doi.org/10.3929/ethz-a-007555133>
- Nussbaum, M., Papritz, A., Baltensweiler, A., Walthert, L. 2014:** Estimating soil organic carbon stocks of Swiss forest soils by robust external-drift kriging. *Geoscientific Model Development* 7: 1197–1210. <http://dx.doi.org/10.5194/gmd-7-1197-2014>
- Nussbaumer, T., Hälgl, L. 2015:** Emissionsfaktoren von Holzfeuerungen – Aktualisierung und Ergänzung 2014. Verenum im Auftrag des Bundesamtes für Umwelt (BAFU) inkl. Excelfile [confidential/internal].
- OcCC 2008:** Das Klima ändert – was nun? Der neue UN-Klimabericht (IPCC 2007) und die wichtigsten Ergebnisse aus Sicht der Schweiz. The Advisory Body on Climate Change, Bern. <http://proclimweb.scnat.ch/Products/OcCC-IPCC/OcCC-IPCC-lowres.pdf> [08.02.2017]
- OECD 2013:** Agricultural policy monitoring and evaluation 2013: OECD countries and emerging economies. Organisation for Economic Co-operation and Development (OECD); Paris, France. http://www.oecd-ilibrary.org/agriculture-and-food/agricultural-policy-monitoring-and-evaluation-2013_agr_pol-2013-en
- Off-Highway Research 2005:** The Market for Construction Equipment and Agricultural Tractors in Switzerland. Internal document INFRAS Bern. September 2005 [confidential/hardcopy].
- Off-Highway Research 2008:** The Market for Construction Equipment and Agricultural Tractors in Switzerland. Internal document FOEN. November 2008 [confidential/internal].
- Off-Highway Research 2012:** The Market for Construction Equipment and Agricultural Tractors in Switzerland. Internal document FOEN. April 2012 [confidential/internal].
- Oney, B., Henne, S., Gruber, N., Leuenberger, M., Bamberger, I., Eugster, W., and Brunner, D. 2015:** The CarboCount CH sites: characterization of a dense greenhouse gas observation network, *Atmos. Chem. Phys.*, 15, 11147–11164. <http://dx.doi.org/10.5194/acp-15-11147-2015>
- OS 2016:** Liechtenstein's energy statistics 2016. Office of statistics Liechtenstein. <http://www.llv.li/files/as/energiestatistik-2015.pdf>
- Park, K.H., Thompson, A.G., Marinier, M., Clark, K., Wagner-Riddle, C. 2006:** Greenhouse gas emissions from stored liquid swine manure in a cold climate. *Atmospheric Environment* 40 (4): 618–627. <http://dx.doi.org/10.1016/j.atmosenv.2005.09.075>
- Paul, C., Weber, M., Mosandl, R. 2009:** Kohlenstoffbindung junger Aufforstungsflächen. Literaturstudie. Karl Gayer Institut und Lehrstuhl für Waldbau der Technischen Universität München, Freising, 70 p. <https://mediatum.ub.tum.de/doc/1219390/1219390.pdf>
- Perruchoud, D., Kienast, F., Kaufmann, E., Bräker, O.U 1999:** 20th Century Carbon Budget of Forest Soils in the Alps. *Ecosystems* 2: 320–337. <http://dx.doi.org/10.1007/s100219900083>

Poncet, P.-A., Boessinger, M., Guillet, A., Klopfenstein, S., König-Bürgi, D., Lüth, A., Martin, R., Montavon, S., Obexer-Ruff, G., Rieder, S., Rubli, S., Rüegg, P., Trolliet, Ch. F. 2009: Wirtschafts-, gesellschafts- und umweltpolitische Bedeutung des Pferdes in der Schweiz – Was gibt es Neues seit 2007? Bericht der Arbeitsgruppe Pferdebranche. Avenches, Schweiz. (ID 21284)

https://www.agroscope.admin.ch/agroscope/de/home/publikationen/suchen/webcode/_jcr_content/par/externalcontent.external.exturl.pdf/aHR0cHM6Ly9pcmEuYWdyb3Njb3BILmNoL2RILUNIL0VpbnplbH/B1YmXpa2F0aW9uL0Rvd25sb2FkRXh0ZXJuP2VpbnplbHB1YmXp/a2F0aW9uSWQ9MjlxMDA=.pdf

Poncet, P.-A., Guillet, A., Jallon, L., Lüth, A., Martin, R., Montavon, S., Sunier, E., Trolliet, Ch. F., Wohlfender, K. 2007: Wirtschafts-, gesellschafts- und umweltpolitische Bedeutung des Pferdes in der Schweiz. Bericht der Arbeitsgruppe Pferdebranche. Avenches, Schweiz. (ID 21285)

https://www.agroscope.admin.ch/agroscope/de/home/publikationen/suchen/webcode/_jcr_content/par/externalcontent.external.exturl.pdf/aHR0cHM6Ly9pcmEuYWdyb3Njb3BILmNoL2RILUNIL0VpbnplbH/B1YmXpa2F0aW9uL0Rvd25sb2FkRXh0ZXJuP2VpbnplbHB1YmXp/a2F0aW9uSWQ9MjlxMDI=.pdf

Prasuhn, V. 2016: Abklärung zum Umweltziel Landwirtschaft: Reduktion der landwirtschaftsbedingten Stickstoffeinträge in die Gewässer um 50% gegenüber 1985. Bericht im Auftrag des Bundesamtes für Umwelt (BAFU). Agroscope. Zürich; Schweiz.

<https://www.agroscope.admin.ch/agroscope/de/home/publikationen.html>

Price, B., Gomez, A., Mathys, L., Gardi, O., Schellenberger, A., Ginzler, C., Thürig, E. 2017: Tree biomass in the Swiss landscape: nationwide modelling for improved accounting for forest and non-forest trees. *Environmental Monitoring and Assessment*, 189: 106.

<http://doi.org/10.1007/s10661-017-5816-7>

Prognos 2012a: Die Energieperspektiven für die Schweiz bis 2050 – Energienachfrage und Elektrizitätsangebot in der Schweiz 2000–2050. Prognos AG im Auftrag des Bundesamtes für Energie, Basel.

[http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_564869151.pdf&endung=Die Energieperspektiven für die Schweiz bis 2050](http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_564869151.pdf&endung=Die+Energieperspektiven+für+die+Schweiz+bis+2050)

Prognos 2013: CO₂-Emissionen 1990–2012 von Industrie- und Dienstleistungen, Endbericht / Kurzdokumentation zuhanden Bundesamt für Umwelt, Bern. Prognos, Basel [confidential/internal].

Quantis 2014: Methanemissionen der Schweizer Gaswirtschaft. Zeitreihe 1990 bis 2012. Schlussbericht. Quantis im Auftrag des Schweizerischen Vereins des Gas- und Wasserfaches SVGW und des Bundesamts für Umwelt BAFU.

<http://www.climatereporting.ch>

Ramin, M., Huhtanen, P. 2012: Development of non-linear models for predicting enteric methane production. *Acta Agriculturae Scandinavica, Section A — Animal Science* 62(4): 254–258. <http://dx.doi.org/10.1080/09064702.2013.765908>

RAP 1999: Fütterungsempfehlungen und Nährwerttabellen für Wiederkäuer [Apports alimentaires recommandés et tables de la valeur nutritive des aliments pour les ruminants]. Landwirtschaftliche Lehrmittelzentrale, Zollikofen. Vierte Auflage [hardcopy].

Richard, L. 1995: Ecologie des mégaphorbiaies subalpines à aune vert de la Vanoise et des régions environnantes (seconde partie) – Phytoécologie. Trav Sci Parc Nat Vanoise 19: 131–160.

Rigling, A., Schaffer, H.P. (Eds.) 2015: Forest Report 2015. Condition and Use of Swiss Forests. Federal Office for the Environment (FOEN), Bern, Swiss Federal Institute (WSL), Birmensdorf. 144 pages. <https://www.bafu.admin.ch/uz-1512-e> [English], <https://www.bafu.admin.ch/uz-1512-d> [German]

Ruffner, H.P. 2005: Written communication from Hans Peter Ruffner (Forschungsanstalt Wädenswil) to Jens Leifeld (ART, Reckenholz), 19.12.2005. <http://www.climatereporting.ch>

Rühr, N., Eugster, W. 2009: Soil respiration fluxes and carbon sequestration of two mountain forests in Switzerland. Report on behalf of the Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

Rusch, S., Hagedorn, F., Zimmermann, S., Lüscher, P. 2009: Bodenkohlenstoff nach Windwurf – eine CO₂-Quelle? [Soil carbon after wind throw – a source of CO₂?]. Report on behalf of the Federal Office for the Environment. <http://www.climatereporting.ch> [German]

Rutherford, G.N., Bebi, P., Edwards, P.J., Zimmermann, N.E. 2008: Assessing land-use statistics to model land cover change in a mountainous landscape in the European Alps. Ecological Modelling 212(3–4): 460–471. <http://dx.doi.org/10.1016/j.ecolmodel.2007.10.050>

Rytec 2010: Review of the waste sector of the national greenhouse gas inventory of Switzerland. Final report. Rytec AG, Münsingen. <http://www.climatereporting.ch>

Rytec 2014: Fossiler Kohlenstoffanteil im verbrannten Abfall 1992 – 2012. Errechnung einer Zeitreihe aus der Verknüpfung der Rauchgasmessung 2010/11 und der erhobenen Kehrrechtzusammensetzung 1992, 2002, 2012. Rytec AG, Münsingen. <http://www.climatereporting.ch>

SAEFL 1992: Abfallkonzept für die Schweiz, Schriftenreihe Umwelt Nr. 173. Swiss Agency for the Environment, Forests and Landscape, Bern [hardcopy].

SAEFL 1993: NABO – Nationales Bodenbeobachtungsnetz. Messresultate 1985–1991. [NABO – Réseau d'observation des sols: période d'observation 1985–1991]. Schriftenreihe Umwelt Nr. 200. [Cahier de l'environnement No 200]. Swiss Agency for the Environment, Forests and Landscape, Bern. <https://www.bafu.admin.ch/bafu/de/home/themen/boden/publikationen-studien/publikationen/nationales-bodenbeobachtungsnetz-nabo.html>

SAEFL 1995: Emissionen des Strassenverkehrs 1950–2010. [Emissions polluantes du trafic routier de 1950 à 2010]. Schriftenreihe Umwelt Nr. 255. [Cahier de l'environnement No 255]. Swiss Agency for the Environment, Forests and Landscape, Bern. [hardcopy, available in German and French].

SAEFL 1996: Schadstoffemissionen und Treibstoffverbrauch des Offroad Sektors, Umwelt-Materialien Nr. 49. Elektrowatt Ingenieurunternehmung AG, Zürich; Technik Thermische Maschinen, Niederrohrdorf; Swiss Agency for the Environment, Forests and Landscape, Bern [hardcopy].

SAEFL 1996a: Luftschadstoff-Emissionen aus natürlichen Quellen der Schweiz. [Emissions polluantes dues aux sources naturelles en Suisse]. Schriftenreihe Umwelt Nr. 257. [Cahier de l'environnement No 257]. Swiss Agency for the Environment, Forests and Landscape, Bern. <http://www.climatereporting.ch>

SAEFL 2000: Handbuch Emissionsfaktoren für stationäre Quellen. Ausgabe 2000, Reihe Vollzug Umwelt. Swiss Agency for the Environment, Forests and Landscape, Bern [hardcopy].

SAEFL 2000a: Nationales Boden-Beobachtungsnetz – Veränderungen von Schadstoffgehalten nach 5 und 10 Jahren. [Réseau national d'observation des sols – Variations des teneurs en polluants après 5 et 10 ans de suivi]. Schriftenreihe Umwelt Nr. 320. [Cahier de l'environnement No 320]. Swiss Agency for the Environment, Forests and Landscape, Bern. https://www.bafu.admin.ch/dam/bafu/de/dokumente/boden/uz-umwelt-zustand/nationales_boden-beobachtungsnetzabveraenderungenvonschadstoff.pdf

SAEFL 2004b: Swiss national forest programme (Swiss NFP), Environmental documentation No. 363, Swiss Agency for the Environment, Forests and Landscape, Bern. 117 pp. https://www.bafu.admin.ch/dam/bafu/en/dokumente/wald-holz/ud-umwelt-diverses/waldprogramm_schweizwap-chhandlungsprogramm2004-2015.pdf

SBV 2004: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2003. [Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung 2003]. Swiss Farmers Union, Brugg [hardcopy, available in German and French].

SBV 2007: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2006. [Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung, 2006]. Swiss Farmers Union, Brugg [hardcopy, available in German and French].

SBV 2007a: Données mensuelles sur l'agriculture. 67.8. August 2007. [Landwirtschaftliche Monatszahlen 67.8. August 2007]. Swiss Farmers Union, Brugg. <http://www.climatereporting.ch> [available in German and French]

SBV 2013: Betriebsinterne Verrechnungsansätze und Inventar-Grunddaten (BIV) 2013. Schweizerischer Baumeisterverband. Verlag SBV [confidential/internal].

SBV 2016: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2015. [Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung, 2015]. Swiss Farmers Union, Brugg. [available in German and French, partly online, partly hardcopy] <http://www.sbv-usp.ch/de/statistik>

Schenk, M. .K., Appel, S. and Daum, D. 1997: N₂O emissions during composting of organic waste. Acta Hort. (ISHS) 450: 253–262. http://www.actahort.org/books/450/450_30.htm

Schjoerring, J.K., Mattsson, M. 2001: Quantification of ammonia exchange between agricultural cropland and the atmosphere: measurements over two complete growth cycles of oilseed rape, wheat, barley and pea. Plant and Soil 228(1): 105–115. <http://www.jstor.org/stable/42950948>

Schlegel, P., Menzi, H. 2013: Schweizer Kaninchenproduktion und deren Stickstoff-, Phosphor- und Kaliumausscheidung. Agrarforschung Schweiz 4(2): 60–67. http://www.agrarforschungschweiz.ch/artikel/2013_02_1839.pdf

Schmid, M., Neftel, A., Fuhrer, J. 2000: Lachgasemissionen aus der Schweizer Landwirtschaft. [Emissions de protoxyde d'azote de l'agriculture Suisse]. Schriftenreihe der FAL 33. Zürich-Reckenholz. <http://www.climatereporting.ch>

Schmidlin, L., Bachmann, I., Flierl, S., Schwarz, A., Roesch, A., Rieder S., von Niederhäusern R. 2013: Wirtschafts-, Gesellschafts- und Umweltpolitische Bedeutung des Pferdes in der Schweiz – Stand 2013. Agroscope Forschungsanstalt Liebefeld- Posieux ALP-Haras, Schweizerisches Nationalgestüt. Avenches, Schweiz.
<http://www.cofichev.ch/Htdocs/Files/v/5885.pdf/Publicationsautres/SCHMIDLINLetalBerichtPferdebranche2013AGROSCOPE20131220.pdf>

Schneider, M. 2010: Personal communication from Manuel Schneider (ART, Reckenholz) to Daniel Bretscher (ART, Reckenholz), 21.09.2010. <http://www.climatereporting.ch>

Schwarz, W. 2001: Emissionen des Kältemittels R-134a aus mobilen Klimaanlageen. Jährliche Emissionsraten von bis zu sieben Jahre alten Pkw-Klimaanlagen. [Emission of Refrigerant R-134a from Mobile Air-Conditioning Systems Annual Rate of Emission from Passenger-Car Air-Conditioning Systems up to Seven Years Old]. Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 360 09 006), Frankfurt.
<http://www.oekorecherche.de/sites/default/files/publikationen/vollr134a.pdf>

Schwarz, W., Wartmann, S. 2005: Emissionen und Emissionsprognosen von H-FKW, FKW und SF₆ in Deutschland – Aktueller Stand und Entwicklung eines Systems zur jährlichen Ermittlung. Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 202 41 356), Frankfurt.
<http://www.umweltbundesamt.de/sites/default/files/medien/publikation/long/3000.pdf>

SCNAT 2016: Akademien der Wissenschaften Schweiz (2016), Brennpunkt Klima Schweiz. Grundlagen, Folgen und Perspektiven. Swiss Academies Reports 11 (5),
<http://www.naturwissenschaften.ch/organisations/proclim/activities/brennpunkt/downloads/81637-brennpunkt-klima-schweiz>

SFOE 1999: Ökologischer, energetischer und ökonomischer Vergleich von Vergärung, Kompostierung und Verbrennung fester biogener Abfallstoffe, Edelmann W., Schleiss K.
<http://www.bfe.admin.ch/php/modules/enet/streamfile.php?file=000000007031.pdf>

SFOE 2010d: Tanktourismus. Swiss Federal Office of Energy, Bern. In German, with Summary in French.
<http://www.news.admin.ch/NSBSubscriber/message/attachments/19578.pdf>
[\[1.02.2017\]](#)

SFOE 2012: Schweizerische Gesamtenergiestatistik 2011. Statistique globale suisse de l'énergie 2011. Swiss Federal Office of Energy, Bern. In German and French.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_872595201.pdf

SFOE 2012a: Frage BAFU zu den Unsicherheiten der fossilen Energieträger GEST. Written message of the Swiss Federal Office of Energy (SFOE) to the Federal Office for the Environment (FOEN) [confidential/internal].

SFOE 2015d: Energieverbrauch in der Industrie und im Dienstleistungssektor. Resultate 2014 (German/French). Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_567447663.pdf

SFOE 2016: Schweizerische Gesamtenergiestatistik 2015. Statistique globale suisse de l'énergie 2015. Swiss Federal Office of Energy, Bern. In German and French.

[http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_83041321.pdf&endung=Schweizerische Gesamtenergiestatistik 2015](http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_83041321.pdf&endung=Schweizerische%20Gesamtenergiestatistik%202015)

SFOE 2016a: Schweizerische Statistik der erneuerbaren Energien (Swiss renewable energy statistics). Ausgabe 2015. Swiss Federal Office of Energy, Bern.

http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_818961010.pdf&endung=Schweizerische%20Statistik%20der%20erneuerbaren%20Energien

SFOE 2016b: Schweizerische Holzenergiestatistik. Erhebung für das Jahr 2015. Swiss Federal Office of Energy, Bern.

http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_373702251.pdf&endung=Schweizerische%20Holzenergiestatistik

SFOE 2016c: Thermische Stromproduktion inklusive Wärmekraftkopplung (WKK) in der Schweiz. Ausgabe 2015. Swiss Federal Office of Energy, Bern.

http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_481124506.pdf.

SFOE 2016d: Energieverbrauch in der Industrie und im Dienstleistungssektor. Resultate 2015 (German/French). Swiss Federal Office of Energy, Bern.

http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_14044384.pdf

SFOE/FOEN 2014: Messung von Heizwerten und CO₂-Emissionsfaktoren von Erdölprodukten 2013. Statistische Analyse der Messresultate, Bern [confidential/internal].

SFSO 2000a: Digital soil map 1:200'000 („Bodeneignungskarte“, BEK). Swiss Federal Statistical Office, GEOSTAT, Neuchâtel.

<https://www.bfs.admin.ch/bfs/de/home/dienstleistungen/geostat/geodaten-bundesstatistik/boden-nutzung-bedeckung-eignung/abgeleitete-und-andere-daten/bodeneignungskarte-schweiz.html> [queries from database].

SFSO 2002: Einblicke in die schweizerische Landwirtschaft. [Insights into Swiss Agriculture]. Swiss Federal Statistical Office, Neuchâtel. [hardcopy, available in German and French]

SFSO 2005: Swiss Land Use Statistics (Arealstatistik Schweiz). Supply of hectare-based data of the first survey (Arealstatistik 1979/85, ASCH1) and second survey (Arealstatistik 1992/97, ASCH2). Swiss Federal Statistical Office, Neuchâtel.

http://www.bfs.admin.ch/bfs/portal/fr/index/infothek/erhebungen_quellen/blank/blank/arealstatistik/01.html [queries from database]

SFSO 2006a: Arealstatistik 2004/09 – Kategorienkatalog. Bodenbedeckung, Bodennutzung. Sektion Geoinformatik, Swiss Federal Statistical Office, Neuchâtel.

<http://www.climatereporting.ch>

SFSO 2009a: Volkswirtschaftliche Gesamtrechnung der Schweiz im Jahr 2008 (National Accounting 2008) Bruttoinlandprodukt (BIP), Swiss Federal Statistical Office, Neuchâtel, November 2009.

<http://www.bfs.admin.ch/bfs/portal/de/index/themen/04/22/publ.html?publicationID=3777>

SFSO 2013a: Landwirtschaftliche Betriebszählung – Zusatzerhebung. Swiss Federal Statistical Office (SFSO), Neuchâtel. [internal/confidential].

SFSO 2016: STAT-TAB: Die interaktive Statistikdatenbank; Datenwürfel für Thema 07.2 – Landwirtschaft. Swiss Federal Statistical Office, Neuchâtel.

https://www.pxweb.bfs.admin.ch/Default.aspx?px_language=de [queries from database].

SFSO 2016a: Die Bevölkerung der Schweiz 2015. Swiss Federal Office. Neuchâtel.

<https://www.bfs.admin.ch/bfsstatic/dam/assets/1401562/master>

SFSO 2016b: Background data of the Gross Nutrient Balance of the Swiss Federal Statistical Office. Swiss Federal Statistical Office, Neuchâtel [confidential/internal, queries from database].

SFSO 2016c: Switzerland's road vehicle stock. Swiss Federal Statistical Office

<https://www.bfs.admin.ch/bfs/de/home/aktuell/neue-veroeffentlichungen.gnpdetail.2017-0321.html>

SFSO 2016d: Supply of provisional data of the AREA Land Use Statistics. Written communication from Felix Weibel and Jürg Burkhalter (SFSO, Neuchâtel) to Christoph Könitzer (Sigmaplan, Bern), 25 July 2016 [confidential/internal, queries from database].

SFSO 2016e: Die Beschäftigungsstatistik der Schweiz 2014. Swiss Federal Statistical Office, Neuchâtel. <http://www.bfs.admin.ch/bfs/portal/de/index/themen/06/02/blank/data.html>

SFSO 2016f: Bruttoinlandprodukt pro Einwohner 1990–2015. Swiss Federal Statistical Office, Neuchâtel.

<https://www.bfs.admin.ch/bfs/de/home/statistiken/volkswirtschaft.assetdetail.214640.html>

[22.02.2017]

SGWA 2016: Erdgaseigenschaften. Written communication R. Huber (Swiss Gas and Water Industry Association) to R. Röthlisberger (FOEN). 23.06.2016 [confidential/internal]

SGWA 2016a: G13d Richtlinie für die Einspeisung von erneuerbaren Gasen G13d, Ausgabe März 2016. Swiss Gas and Water Industry Association

http://www.svgw.ch/index.php?id=137&no_cache=1&tx_commerce_pi1%5BshowUid%5D=291&tx_commerce_pi1%5BcatUid%5D=11&cHash=7c480e936b2a3f323dabe5199c7a1c5c

SHL 2010: Bestimmung Tiere der Rindviehkategorien ab 2009 für die Berechnung des Ammoniakinventars. Schweizerische Hochschule für Landwirtschaft SHL, Zollikofen.

<http://www.climatereporting.ch>

SIA 1988: Norm Nr. 318, Garten- und Landschaftsbau (Practice Guidance No. 318 on Gardening and Landscaping Operations). Swiss Society of Engineers and Architects (SIA).

<http://www.webnorm.ch/481573fd-6a93-4890-9aa6-96b16753f4c2/D/DownloadAnhang>

Sigmaplan 2010a: Deforestation under Kyoto Protocol. Documentation of implementation. Internal documentation by Mathys, L., Sigmaplan, Bern on behalf of the Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

Sigmaplan 2012a: LULUCF and KP-LULUCF. Comparison of deforestation data. Final Report. Internal document. Sigmaplan, Bern. <http://www.climatereporting.ch>

SigmaPlan 2017: Treibhausgasinventar Sektor LULUCF, Tabellen zur Berechnung der Aktivitätsdaten im Sektor LULUCF in der Schweiz inklusive Begleitdokumentation. Internes Dokument. [GHG inventory, sector LULUCF. Spreadsheets to calculate the activity data of the LULUCF sector in Switzerland inclusive accompanying documents. Internal document]. SigmaPlan, Bern.

SKW 2010: Annual reports of the Swiss Cosmetic and Detergent Association. Zurich.
http://www.skw-cds.ch/fileadmin/skwcdsch/01_Verband/JB2010D.pdf

Soliva, C.R. 2006: Report to the attention of IPCC about the data set and calculation method used to estimate methane formation from enteric fermentation of agricultural livestock population and manure management in Swiss agriculture. On behalf of the Federal Office for the Environment, Bern. ETH Zurich, Institute of Animal Science.
<http://www.climatereporting.ch>

Sommer, S.G., Petersen, S.O., Sorensen, P., Poulsen, H.D., Moller, H.B. 2007: Methane and carbon dioxide emissions and nitrogen turnover during liquid manure storage. Nutrient Cycling in Agroecosystems 78 (1): 27–36. <http://dx.doi.org/10.1007/s10705-006-9072-4>

Spiecker, H. 1999: Growth trends in European forests, Do we have sufficient knowledge? In Karjalainen T, Spiecker H, Laroussinie, O. (Eds) 1999: Causes and consequences of accelerating tree growth in Europe. Proceedings of the international seminar held in Nancy, France 14–16. May 1998. European Forest Institute Proceedings No. 27: 157–169.

Spiess, E. 2011: Nitrogen, phosphorus and potassium balances and cycles of Swiss agriculture from 1975 to 2008. Nutrient Cycling in Agroecosystems 91 (3): 351–365.
<http://dx.doi.org/10.1007/s10705-011-9466-9>

Spiess, E., Prasuhn, V. 2006: Weniger Nitrat im Grundwasser dank Ökomassnahmen. AGRARForschung 13(1): 4–9.
http://www.agrarforschungschweiz.ch/artikel/2006_01_1145.pdf

Spirig, C., Neftel, A. 2002: Biogene VOC und Aerosole. Bedeutung der biogenen flüchtigen organischen Verbindungen für die Aerosolbildung. Schriftenreihe der FAL 42. Zürich-Reckenholz [hardcopy].

SQS 2008: IQNet and SQS ISO 9001:2000 Certificate. Certified area: National Inventory System. Field of activity: Greenhouse Gas Inventory and CO₂ Statistics. Registration Number: CH-34433. Issued on 04.01.2008. <http://www.climatereporting.ch>

Staerfl, S. M., Amelchanka, S. L., Kälber, T., Soliva, C. R., Kreuzer, M., Zeitz, J. O. 2012: Effect of feeding dried high-sugar ryegrass ('AberMagic') on methane and urinary nitrogen emissions of primiparous cows. Livestock Science 150(1–3): 293–301.
<http://dx.doi.org/10.1016/j.livsci.2012.09.019>

Stangl, R. 2004: Ingenieurbiologische Hangsicherungsmassnahmen in Wildbacheinzugsgebieten – Bestandesentwicklung und langfristige Wirksamkeit. Klagenfurt: Internationale Forschungsgesellschaft Interpraevent, Tagungspublikation Internationales Symposium Interpraevent 2004, Band 1, pp. 295–305. <http://www.climatereporting.ch>

Statistics Finland 2016: Greenhouse gas emissions in Finland 1990–2014. National Inventory Report under the UNFCCC and the Kyoto Protocol. Statistics Finland, Helsinki.
http://unfccc.int/files/national_reports/annex_i_ghg_inventories/national_inventories_submissions/application/zip/fin-2016-nir-15jun16.zip

Stehfest, E., Bouwman, L. 2006: N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. *Nutrient Cycling in Agroecosystems* 74(3): 207–228.

<http://dx.doi.org/10.1007/s10705-006-9000-7>

Stieger, J. 2013: Methane emissions at farm and regional scales: origin, magnitude and spatio-temporal variability. PhD Thesis ETHZ, No. 21538; Zürich, Switzerland.

<http://www.climatereporting.ch>

Stieger, J., Bamberger, I., Buchmann, N., Eugster, W. 2015: Validation of farm-scale methane emissions using nocturnal boundary layer budgets. *Atmos. Chem. Phys.* 15 (24): 14055–14069. <http://dx.doi.org/10.5194/acp-15-14055-2015>

Stricker, B. 2012: Energiebedarfswerte der Pferde, Esel und Ponys. Written communication from Brigitte Stricker (ALP-Haras, Agroscope Liebefeld-Posieux research station) to Daniel Bretscher (ART, Reckenholz), August-October 2012. <http://www.climatereporting.ch>

SWI 2009: Forests continue conquest of countryside. Swissinfo.ch, published online 28.07.2009. <http://www.swissinfo.ch/eng/forests-continue-conquest-of-countryside/990884>

Swiss Civil Aviation Emissions Inventory 2015. Federal Office of Civil Aviation FOCA. Reference: FOCA / 047.1-00017/00002. 29.11.2016 [confidential/internal].

Swiss Confederation 1983: Loi fédérale du 7 octobre 1983 sur la protection de l'environnement (Loi sur la protection de l'environnement, LPE). Status as of 01 January 2017. http://www.admin.ch/ch/f/rs/c814_01.html [official text in German, French and Italian]; Federal Act on the Protection of the Environment [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19830267/index.html> [05.04.2017]

Swiss Confederation 1985: Ordonnance du 16 décembre 1985 sur la protection de l'air (OPair). Status as of 1 April 2017 http://www.admin.ch/ch/f/rs/c814_318_142_1.html [official text in German, French and Italian]; Ordinance on Air Pollution Control [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19850321/index.html> [05.04.2017]

Swiss Confederation 1986: Ordonnance du 9 juin 1986 sur les polluants du sol (Ordinance against pollution of soils). In force until 1998; replaced by Swiss Confederation, 1998 [hardcopy].

Swiss Confederation 1991: Loi fédérale du 4 octobre 1991 sur les forêts (Loi sur les forêts, LFo). Status as of 01 January 2017. http://www.admin.ch/ch/f/rs/c921_0.html [official text in German, French and Italian]; Federal Act on Forest [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19910255/index.html> [05.04.2017]

Swiss Confederation 1991a: Ordonnance du 21 janvier 1991 sur la protection des hauts-marais et des marais de transition d'importance nationale (Ordonnance sur les hauts-marais). Status as of 01 July 2015. http://www.admin.ch/ch/f/rs/c451_32.html [official text in German, French and Italian] [05.04.2017]

Swiss Confederation 1992: Ordonnance du 30 novembre 1992 sur les forêts (OFo). Status as of 01 January 2017. http://www.admin.ch/ch/f/rs/c921_01.html [official text in German, French and Italian]; Ordinance on Forest [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19920310/index.html> [05.04.2017]

Swiss Confederation 1992a: Bundesstatistikgesetz vom 9. Oktober 1992. As of 1. January 2016. http://www.admin.ch/ch/d/sr/c431_01.html [official text in German, French and Italian]; Federal Statistics Act [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19920252/index.html> [05.04.2017]

Swiss Confederation 1993: Verordnung über die Durchführung von statistischen Erhebungen des Bundes vom 30. Juni 1993. Status as of 1. December 2016. http://www.admin.ch/ch/d/sr/c431_012_1.html [official text in German, French and Italian] [05.04.2017]

Swiss Confederation 1994: Ordonnance du 7 septembre 1994 sur la protection des bas-marais d'importance nationale (Ordonnance sur les bas-marais). Status as of 1. July 2015. http://www.admin.ch/ch/f/rs/c451_33.html [official text in German, French and Italian] [05.04.2017]

Swiss Confederation 1997: Ordonnance du 12 novembre 1997 sur la taxe d'incitation sur les composés organiques volatils (OCOV). As at 01 January 2009. http://www.admin.ch/ch/f/rs/c814_018.html [official text in German, French and Italian]; Ordinance on the Incentive Tax on Volatile Organic Compounds [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19970460/index.html> [05.04.2017]

Swiss Confederation 1998: Ordonnance du 1er juillet 1998 sur les atteintes portées aux sols (OSol) (Ordinance against deterioration of soils). Status as of 12. April 2012. http://www.admin.ch/ch/f/rs/c814_12.html [official text in German, French and Italian] [05.04.2017]

Swiss Confederation 1998a: Gewässerschutzverordnung vom 28 October 1998. Status as of 7. February 2017. http://www.admin.ch/ch/d/sr/c814_201.html [official text in German, French and Italian]; Ordinance on the Incentive Tax on Volatile Organic Compounds [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/19983281/index.html> [05.04.2017]

Swiss Confederation 2005: Ordinance of 18 May 2005 on Risk Reduction related to the Use of certain particularly dangerous Substances, Preparations and Articles (Ordinance on Chemical Risk Reduction, ORRChem). SR Number 814.81. Status as of 1 February 2017. http://www.admin.ch/ch/d/sr/814_81/index.html [official text in German, French and Italian]; Ordinance on the Reduction of Risks relating to the Use of Certain Particularly Dangerous Substances, Preparations and Articles [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/20021520/index.html> [05.04.2017]

Swiss Confederation 2008: Verordnung des EVD über Ethoprogramme (Ethoprogrammverordnung), 25. Juni 2008 (Stand am 1. Oktober 2008). Der Schweiz. Bundesrat, Bern, Schweiz. <http://www.admin.ch/opc/de/official-compilation/2008/3785.pdf>

Swiss Confederation 2011: Bundesgesetz über die Reduktion der CO₂-Emissionen (CO₂-Gesetz). Status as of 1. January 2013. http://www.admin.ch/ch/d/sr/c641_71.html [official text in German, French and Italian]; Federal Act on the Reduction of CO₂ Emissions [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/20091310/index.html> [05.04.2017]

Swiss Confederation 2012: Verordnung über die Reduktion der CO₂-Emissionen (CO₂-Verordnung). Status as of 12.2.2013. http://www.admin.ch/ch/d/sr/c641_711.html [official text in German, French and Italian]; Ordinance on the Reduction of CO₂ Emissions [English translation for information purposes only]: <https://www.admin.ch/opc/en/classified-compilation/20120090/index.html> [05.04.2017]

Swiss Confederation 2015: Verordnung über die Vermeidung und die Entsorgung von Abfällen (Abfallverordnung, VVEA). Status as of 19. July 2016. <https://www.admin.ch/opc/de/classified-compilation/20141858/> [official text in German, French and Italian]. [05.04.2017]

Swiss-TS 2016: Quality management according to ISO 9001:2008. Certificate. Certified area: National Inventory System and Reporting under the UNFCCC and the Kyoto Protocol. Registration Number: 13-299-113. Issued on 28.11.2016. <http://www.climatereporting.ch>

Theloke, J. 2005: NMVOC-Emissionen aus der Lösemittelanwendung und Möglichkeiten zu ihrer Minderung. Fortschritt-Berichte VDI Reihe 15 Nr. 252. Düsseldorf: VDI-Verlag. <http://elib.uni-stuttgart.de/opus/volltexte/2005/2339>

Theloke, J. et al. 2000: Ermittlung der Lösemittlemissionen 1994 in Deutschland und Methoden zur Fortschreibung. Institut für Energiewirtschaft und Rationelle Energieanwendung (IER) der Universität Stuttgart. Forschungsbericht 295 42 628. Im Auftrag des Umweltbundesamtes. <https://www.umweltbundesamt.de/sites/default/files/medien/publikation/long/2484.pdf>

Thuille, A., Schulze, E.D. 2006: Carbon dynamics in successional and afforested spruce stands in Thuringia and the Alps. Global Change Biology 12: 325–342. <http://dx.doi.org/10.1111/j.1365-2486.2005.01078.x>

Thürig, E., Herold, A. 2013: Recalculation of emission factors in Swiss forests for the Swiss GHGI. Internal documentation of technical adjustments of data delivery and more recent data. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. <http://www.climatereporting.ch>

Thürig, E., Rösler, E., Didion, M. 2015: NFI 2009–2013: Calculation of emission factors for living biomass in Swiss forests for the Swiss GHGI 2015. Internal documentation of data delivery and more recent data. Report commissioned by the Swiss Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

Thürig, E., Rösler, E., Didion, M. 2015: NFI4 2011–2015: Calculation of emission factors for living biomass in Swiss forests for the Swiss GHGI 2017. Internal documentation of data delivery and more recent data. Report commissioned by the Swiss Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

Thürig, E., Traub, B. 2015: Non-forest areas converted to forest: standing stock, gains and losses in biomass. Report commissioned by the Swiss Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

Thürig, E., Weber, P., Lischke, H., Schmatz, D., Kaufmann, E., Zandt, H., Dobbertin, M., Zingg, A., Cherubini, P., Waldner, P. 2009: Verbesserung der Klimaabhängigkeit der Wachstumsfunktion für Szenarienanalysen der Waldentwicklung (Holznutzungspotential, Nachhaltigkeit) KLIWAWA. [Improvement of climate dependency of growth curves for analyzing forest development scenarios (sustainable Potential Wood Supply)]. Forschungsanstalt für Wald, Schnee und Landschaft WSL. Research project on behalf of the Federal Office for the Environment, Bern. <http://www.climatereporting.ch>

TUG 2009: Emission Factors from the Model PHEM for the HBEFA Version 3, Institute for Internal Combustion Engines and Thermodynamics, Report Nr. I-20/2009 Haus-Em 33/08/679, University of Technology, Graz.
http://www.hbefa.net/e/documents/HBEFA_31_Docu_hot_emissionfactors_PC_LCV_HDV.pdf [1.02.2017]

Tuomi, M., Laiho, R., Repo, A., Liski, J. 2011: Wood decomposition model for boreal forests. Ecological Modelling 222: 709–718.
<http://dx.doi.org/10.1016/j.ecolmodel.2010.10.025>

Tuomi, M., Thum, T., Järvinen, H., Fronzek, S., Berg, B., Harmon, M., Trofymow, J.A., Sevanto, S., Liski, J. 2009: Leaf litter decomposition – Estimates of global variability based on Yasso07 model. Ecological Modelling 220: 3362–3371.
<http://dx.doi.org/10.1016/j.ecolmodel.2009.05.016>

UBA 2005: Emissionen, Aktivitätsraten und Emissionsfaktoren von fluoridierten Treibhausgasen (F-Gasen) in Deutschland für die Jahre 1995–2002. Texte: 14 / 05. Umwelt Bundes Amt. Berlin.
<https://www.umweltbundesamt.de/sites/default/files/medien/publikation/long/2902.pdf>

UBA 2006: Einsatz von Sekundärbrennstoffen, Forschungsbericht 204 42 203/02, UBA-FB 000893; http://www.eefa.de/pdf/UBA-Texte_07_06.pdf

UBA 2007: Daten von H-FKW, FKW und SF6 für die nationale Emissionsberichterstattung gemäß Klimarahmenkonvention für die Berichtsjahre 2004 und 2005“ "Fluorierte Treibhausgase". Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 205 41 217/01), Frankfurt.
<http://www.umweltbundesamt.de/sites/default/files/medien/publikation/long/3439.pdf>

UBA/Ökorecherche 2012: Entsorgung von Kältemitteln aus stationären Anlagen, Emissionsfaktor der Entsorgung, Emissionsberichterstattung von F-Gasen in der Kälte- und Klimatechnik. UBA Umweltbundesamt Berlin – 24.10.12. Öko-Recherche Büro für Umweltforschung und -beratung GmbH, W. Schwarz, Barbara Gschrey. Umweltbundesamt Berlin, 24. Oktober 2012, Ergebnisse des Forschungsvorhabens 1. Gewerbekälte und 2. Industriekälte.
http://www.umweltbundesamt.de/sites/default/files/medien/419/dokumente/4_entsorgung_stanlagen1.pdf

UNFCCC 2007: Report of the individual review of the greenhouse gas inventory of Switzerland submitted in 2006. FCCC/ARR/2006/CHE, 31 July 2007.
<http://unfccc.int/resource/docs/2007/arr/che.pdf>

UNFCCC 2009: Report of the individual review of the greenhouse gas inventories of Switzerland submitted in 2007 and 2008. FCCC/ARR/2008/CHE, 29 April 2009.
<http://unfccc.int/resource/docs/2009/arr/che.pdf>

UNFCCC 2011: Report of the individual review of the annual submission of Switzerland submitted in 2010. FCCC/ARR/2010/CHE, 23 May 2011.

<http://unfccc.int/resource/docs/2011/arr/che.pdf>

UNFCCC 2011a: Report of the technical assessment of the forest management reference level submission of Switzerland submitted in 2011 (FCCC/TAR/2011/CHE). Published online on 21 September 2011. <http://unfccc.int/resource/docs/2011/tar/che01.pdf>

UNFCCC 2013c: Approved small scale baseline and monitoring methodology AMS-III.D./Version 19.0. Methane recovery in animal manure management systems.

<https://cdm.unfccc.int/methodologies/SSCmethodologies/approved>

UNFCCC 2014a: Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention. Decision 24/CP.19 (FCCC/CP/2013/10/Add.3)

<http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf> [11.01.2017]

UNFCCC 2014b: Guidance for reporting information on activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol. Decision 6/CMP.9 (FCCC/KP/CMP/2013/9/Add.1)

<http://unfccc.int/resource/docs/2013/cmp9/eng/09a01.pdf> [20.02.2016]

UNFCCC 2015: Guidelines for the technical review of information reported under the Convention related to greenhouse gas inventories, biennial reports and national communications by Parties included in Annex I to the Convention. Decision 13/CP.20 (FCCC/CP/2014/10/Add.3) <http://unfccc.int/resource/docs/2014/cop20/eng/10a03.pdf>

UNFCCC 2017: Draft Report on the individual review of the annual submission of Switzerland submitted in 2016. FCCC/ARR/2016/CHE, 3 March 2017 [confidential/internal].

UNFCCC 2017a: Draft Report on the review of the report to facilitate the calculation of the assigned amount for the second commitment period of the Kyoto Protocol of Switzerland, FCCC/IRR/2016/CHE, 3 March 2017 [confidential/internal].

USEPA 1995: Compilation of air pollutant emission factors, Chapter 10.7 Charcoal.

<https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emission-factors>

USEPA 1995a: US EPA 1995, "Emission factors for open burning of municipal refuse".

<http://www.epa.gov/ttn/chief/ap42/ch02/final/c02s05.pdf>

UVEK 2003: Düngen mit Klärschlamm wird verboten, UVEK Eidgenössisches Departement für Umwelt, Verkehr, Energie, Kommunikation.

http://www.admin.ch/cp/d/3e816ebe_1@presse1.admin.ch.html

Van der Weerden, T.J., Jarvis, S.C. 1997: Ammonia emission factors for N fertilizers applied to two contrasting grassland soils. Environmental Pollution 95(2): 205–211.

[http://dx.doi.org/10.1016/S0269-7491\(96\)00099-1](http://dx.doi.org/10.1016/S0269-7491(96)00099-1)

Van der Zaag, A. C., MacDonald, J. D., Evans, L., Vergé, X. P. C., Desjardins, R. L. 2013: Towards an inventory of methane emissions from manure management that is responsive to changes on Canadian farms. Environmental Research Letters 8(3): 035008.

<http://dx.doi.org/10.1088/1748-9326/8/3/035008>

- Van Dorland, H. A., Wettstein, H. R., Leuenberger, H., Kreuzer, M. 2007:** Effect of supplementation of fresh and ensiled clovers to ryegrass on nitrogen loss and methane emission of dairy cows. *Livestock Science* 111(1–2): 57–69.
<http://dx.doi.org/10.1016/j.livsci.2006.11.015>
- Van Miegroet, H., Olsson, M. 2011:** Ecosystem Disturbance and Soil Organic Carbon – A Review. In: *Soil Carbon in Sensitive European Ecosystems*. John Wiley & Sons, Ltd, Chichester: 85–117. <http://dx.doi.org/10.1002/9781119970255.ch5>
- Vergé, x. P. C., De Kimpe, C., Desjardins, R. L. 2007:** Agricultural production, greenhouse gas emissions and mitigation potential. *Agricultural and Forest Meteorology* 142(2–4): 255–269. <http://dx.doi.org/10.1016/j.agrformet.2006.06.011>
- Vermorel, M., Martin-Rosset, W., Vernet, J. 1997:** Energy utilization of twelve forages or mixed diets for maintenance by sport horses. *Livestock Production Science* 47(2): 157–167. [http://dx.doi.org/10.1016/S0301-6226\(96\)01402-9](http://dx.doi.org/10.1016/S0301-6226(96)01402-9)
- Vesterdal, L., Ritter, E., Gundersen, P. 2002:** Change in soil organic carbon following afforestation of former arable land. *Forest Ecology and Management* 169: 137–147. [http://dx.doi.org/10.1016/S0378-1127\(02\)00304-3](http://dx.doi.org/10.1016/S0378-1127(02)00304-3)
- Vogel, R. 2013:** Quantifying the uncertainty of spatial precipitation analyses with radar-gauge observation ensembles, Zürich: Federal Office of Meteorology and Climatology, MeteoSwiss. <http://www.meteoswiss.admin.ch/content/dam/meteoswiss/en/Ungebundene-Seiten/Publikationen/Scientific-Reports/doc/sr95vogel.pdf>
- Vorreiter, L. 1949:** Holztechnologisches Handbuch; Band I: Allgemeines, Holzkunde, Holzschutz und Holzvergütung. Verlag Georg Fromme & Co., Wien [hardcopy].
- VTI 2011:** Review of the LULUCF Sector of Switzerland's Greenhouse Gas Inventory 1990–2008. Johann Heinrich von Thünen-Institut, Braunschweig. <http://www.climatereporting.ch>
- Walther U., Menzi, H., Ryser, J.-P., Fleisch, R., Jeangros, B., Maillard, A., Siegenthaler, A., Vuilloud, P.A. 1994:** Grundlagen für die Düngung im Acker- und Futterbau. *Agrarforschung* 1(7): 1–40 [hardcopy].
- Warneke C., Luxembourg, S.L., de Gouw, J.A., Rinne, H.J.I., Guenther, A.B., Fall, R. 2002:** Disjunct eddy covariance measurements of oxygenated VOC fluxes from an Alfalfa field before and after cutting. *Journal of Geophysical Research* 107 D8: 4067. <http://dx.doi.org/10.1029/2001JD000594>
- Widmer, A. 2006:** Written communication from Albert Widmer (Forschungsanstalt Wädenswil) to Jens Leifeld (ART, Reckenholz), 20.06.2006. <http://www.climatereporting.ch>
- Wirth, C., Schulze, E. D., Schwalbe, G., Tomczyk, I., Weber, G.-E., Weller, E. 2004:** Dynamik der Kohlenstoffvorräte in den Wäldern Thüringens: Abschlussbericht zur 1. Phase des BMBF-Projektes "Modelluntersuchung zur Umsetzung des Kyoto-Protokolls". *Mitteilungen der Thüringer Landesanstalt für Wald, Jagd und Fischerei* 23 [hardcopy].
- Wüest & Partner 2015:** Bauabfälle in der Schweiz – Hochbau. Studie 2015 Schlussbericht. Wüest & Partner, Zürich, 28. September 2015 <https://www.bafu.admin.ch/bafu/de/home/themen/abfall/zustand/daten.html>
- Wunder, J., Bont, Z. 2015:** Einfluss von Kernfäulen auf den Kohlenstoffspeicher Wald. Wunder Consulting. Projekt im Auftrag des Bundesamt für Umwelt. <http://www.climatereporting.ch>

- Wüst-Galley, C., Grünig, A., Leifeld, J. 2015:** Locating organic soils for the Swiss Greenhouse Gas Inventory. Agroscope Science No. 26, Zürich.
<http://www.climatereporting.ch>
- Wüst-Galley, C., Mössinger, E., Leifeld, J. 2016:** Loss of the soil carbon storage function of drained forested peatlands. *Mires and Peat*, 18(07), 1–22. <http://mires-and-peat.net/pages/volumes/map18/map1807.php>
- Wutzler, T., Mund, M. 2007:** Modelling mean above and below ground litter production based on yield tables. *Silva Fennica* 41(3): 559–574. <https://doi.org/10.14214/sf.289>
- Wutzler, T., Wirth, C., Schumacher, J., 2008:** Generic biomass functions for common beech (*Fagus sylvatica*) in Central Europe: predictions and components of uncertainty. *Canadian Journal of Forest Research* 38: 1661–1675. <http://dx.doi.org/10.1139/X07-194>
- Zeitz, J. O., Soliva, C. R., Kreuzer, M. 2012:** Swiss diet types for cattle: how accurately are they reflected by the Intergovernmental Panel on Climate Change default values? *Journal of Integrative Environmental Sciences*, 9:sup1,199–216.
<http://dx.doi.org/10.1080/1943815X.2012.709253>
- Zell, J., Thürig, E. 2013:** Root biomass functions for for the GHG reporting under the UNFCCC and under the KP in Switzerland. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Internal report to FOEN [confidential/internal].
- Zeschmar-Lahl 2004:** Schadstoffverbringung in Senken durch Abfallbehandlung. Dr. Barbara Zeschmar-Lahl, BZL Kommunikation und Projektsteuerung GmbH, Oyten
http://www.bzl.info/de/sites/default/files/SIDAF_04_BZL_end.PDF
- Zimmermann, S., Hiltbrunner, D. 2012:** Turnover and stabilization of soil organic matter: effect of land-use change in alpine regions. Schlussbericht [German]. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.
<http://www.climatereporting.ch>
- Zumbrunnen, T., Menéndez, P., Bugmann, H., Conedera, M., Gimmi, U., Bürgi, M. 2012:** Human impacts on fire occurrence: a case study of hundred years of forest fires in a dry alpine valley in Switzerland. *Regional Environmental Change* 12: 935–949.
<http://dx.doi.org/10.1007/s10113-012-0307-4>

References to EMIS database comments

Table A – 37 Assignments of NFR Codes to titles of EMIS (Swiss Emission Information System) database comments. These internal documents will be made available to reviewers on request. green cell: NEW comment for the submission at hand.

NFR Code CRF [UNECE]	EMIS Title	NFR Code CRF [UNECE]	EMIS Title
1 A 1 a & 2 A 4 d	Kehrichtverbrennungsanlagen	2 D 3 a [2 D 3 g]	Klebstoff-Produktion
1 A 1 a	Sondermüllverbrennungsanlagen	2 D 3 a [2 D 3 g]	Lösungsmittel-Umschlag und -Lager
1 A 1 a & 5 A	Kehrichtdeponien	2 D 3 a [2 D 3 g]	Pharmazeutische Produktion**
1 A 1 a & 5 B 2	Vergärung IG (industriell-gewerblich)	2 D 3 a [2 D 3 g]	Polyester-Verarbeitung
1 A 1 a & 5 B 2	Vergärung LW (landwirtschaftlich)	2 D 3 a [2 D 3 g]	Polystyrol-Verarbeitung
1 A 1 c	Holzkohle Produktion	2 D 3 a [2 D 3 g]	Polyurethan-Verarbeitung
1 A 2 a & 2 A 4 d	Eisengiessereien Kupolöfen	2 D 3 a [2 D 3 g]	PVC-Verarbeitung
1 A 2 a	Stahl-Produktion Wärmeöfen**	2 D 3 a [2 D 3 g]	Gerben von Ledermaterialien
1 A 2 b	Buntmetallgiessereien übriger Betrieb**	2 D 3 b	Strassenbelagsarbeiten**
1 A 2 b & 2 C 3	Aluminium Produktion	2 D 3 c	Dachpappen Produktion Emissionen aus Bitumen
1 A 2 c & 2 B 8 b [2 B 10 a]	Ethen-Produktion*	2 D 3 c	Dachpappen Produktion Voranstrich
1 A 2 d & 2 A 4 d	Zellulose-Produktion Feuerung*	2 D 3 c	Dachpappen Verlegung Bitumen
1 A 2 f	Kalkproduktion, Feuerung*	2 D 3 c	Dachpappen Verlegung Voranstrich
1 A 2 f	Mischgut Produktion	2 D 3 d	Urea (AdBlue) Einsatz Strassenverkehr
1 A 2 f	Zementwerke Feuerung	2 G 3 a	Lachgasanwendung Spitäler**
1 A 2 f & 2 A 3	Glas übrige Produktion*	2 G 3 b	Lachgasanwendung Haushalt**
1 A 2 f & 2 A 3	Glaswolle Produktion Rohprodukt*	2 G 4 [2 D 3 a]	Pharma-Produkte im Haushalt
1 A 2 f & 2 A 3	Hohlglas Produktion*	2 G 4 [2 D 3 a]	Reinigungs- und Lösemittel; Haushalte
1 A 2 f & 2 A 4 a	Feinkeramik Produktion*	2 G 4 [2 D 3 a]	Spraydosen Haushalte**
1 A 2 f & 2 A 4 a	Ziegeleien**	2 G 4 [2 D 3 h]	Verpackungsdruckereien**
1 A 2 f & 2 A 4 d	Steinwolle Produktion*	2 G 4 [2 D 3 h]	Druckereien uebrige
1 A 2 g iv	Faserplatten Produktion**	2 G 4 [2 D 3 i]	Entfernung von Farben und Lacken
1 A 3 a & 1 A 5	Flugverkehr	2 G 4 [2 D 3 i]	Entwachsung von Fahrzeugen
1 A 3 b i-viii	Strassenverkehr	2 G 4 [2 D 3 i]	Kosmetika-Produktion**
1 A 3 c	Schienerverkehr	2 G 4 [2 D 3 i]	Lösungsmittel-Emissionen IG nicht zugeordnet
1 A 3 e	Gastransport Kompressorstation	2 G 4 [2 D 3 i]	Öl- und Fettgewinnung
1 A 4 b i	Holzkohle-Verbrauch	2 G 4 [2 D 3 i]	Papier- und Karton-Produktion**
1 A 4 b i	Lagerfeuer	2 G 4 [2 D 3 i]	Parfum- und Aromen-Produktion**
1 A 4 c i	Gastrocknung**	2 G 4 [2 D 3 i]	Tabakwaren Produktion**
1 B 2 a iv	Raffinerie, Leckverluste	2 G 4 [2 D 3 i]	Textilien-Produktion
1 B 2 a v	Benzinumschlag Tanklager	2 G 4 [2 D 3 i]	Wissenschaftliche Laboratorien
1 B 2 a v	Benzinumschlag Tankstellen	2 G 4 [2 G]	Korrosionsschutz im Freien
1 B 2 b ii	Gasproduktion	2 G 4 [2 G]	Betonzusatzmittel-Anwendung
1 B 2 b iv-vi	Netzverluste Erdgas	2 G 4 [2 G]	Coiffeursalons
1 B 2 c	Raffinerie, Abfackelung	2 G 4 [2 G]	Fahrzeug-Unterbodenschutz**
1 Energy Model***	Energie New	2 G 4 [2 G]	Feuerwerke
1A	Holzfeuerungen	2 G 4 [2 G]	Flugzeug-Enteisung
1A2g vii, 1A3c, 1A3e, 1A5b (c)	Off-Road	2 G 4 [2 G]	Gas-Anwendung
2 A 1	Zementwerke Rohmaterial	2 G 4 [2 G]	Gesundheitswesen, übrige**
2 A 1	Zementwerke übriger Betrieb	2 G 4 [2 G]	Glaswolle Imprägnierung*
2 A 2	Kalkproduktion, Rohmaterial*	2 G 4 [2 G]	Holzschutzmittel-Anwendung
2 A 2	Kalkproduktion, übriger Betrieb*	2 G 4 [2 G]	Klebstoff-Anwendung
2 A 4 d	Karbonatanwendung weitere	2 G 4 [2 G]	Kosmetik-Institute
2 A 5 a	Gips-Produktion übriger Betrieb**	2 G 4 [2 G]	Kühlschmiermittel-Verwendung
2 A 5 a	Kieswerke	2 G 4 [2 G]	Medizinische Praxen**
2 B 1	Ammoniak-Produktion*	2 G 4 [2 G]	Pflanzenschutzmittel-Verwendung
2 B 10 [2 B 10 a]	Ammoniumnitrat-Produktion*	2 G 4 [2 G]	Reinigung Gebäude IGD**
2 B 10 [2 B 10 a]	Chlorgas-Produktion*	2 G 4 [2 G]	Schmierstoff-Verwendung
2 B 10 [2 B 10 a]	Essigsäure-Produktion*	2 G 4 [2 G]	Spraydosen IndustrieGewerbe
2 B 10 [2 B 10 a]	Formaldehyd-Produktion	2 G 4 [2 G]	Tabakwaren Konsum
2 B 10 [2 B 10 a]	PVC-Produktion	2 G 4 [2 G]	Steinwolle-Imprägnierung*
2 B 10 [2 B 10 a]	Salzsäure-Produktion*	2 H 1	Faserplatten Produktion**
2 B 10 [2 B 10 a]	Schwefelsäure-Produktion*	2 H 1	Zellulose Produktion übriger Betrieb*
2 B 10	Kalksteingrube*	2 H 1	Spanplatten Produktion*
2 B 10	Niacin-Produktion*	2 H 2	Bierbrauereien
2 B 2	Salpetersäure Produktion*	2 H 2	Branntwein Produktion
2 B 5	Graphit und Siliziumkarbid Produktion*	2 H 2	Brot Produktion
2 C - 2 G	Synthetische Gase	2 H 2	Fleischräuchereien
2 C 1	Eisengiessereien Elektroschmelzöfen	2 H 2	Kaffeeröstereien
2 C 1	Eisengiessereien übriger Betrieb	2 H 2	Müllereien
2 C 1	Stahl-Produktion Elektroschmelzöfen**	2 H 2	Wein Produktion
2 C 1	Stahl-Produktion übriger Betrieb**	2 H 2	Zucker Produktion
2 C 1	Stahl-Produktion Walzwerke**	2 H 3	Sprengen und Schiessen
2 C 7 a	Buntmetallgiessereien Elektroöfen**	2 I	Holzbearbeitung
2 C 7 c	Verzinkereien	2 L	NH3 aus Kühlanlagen
2 C 7 c	Batterie-Recycling*	3	Landwirtschaft
2 D 1	Schmiermittel-Anwendung	3 C	Reisanbau
2 D 2	Paraffinwachs-Anwendung	5 B 1	Kompostierung Industrie
2 D 3 a [2 D 3 d]	Farben-Anwendung Bau	5 B 1	Kompostierung, Verbreitung als Dünger im Haushalt
2 D 3 a [2 D 3 d]	Farben-Anwendung andere	5 B 2	Biogasaufbereitung (Methanverlust)
2 D 3 a [2 D 3 d]	Farben-Anwendung Haushalte**	5 C 1 [5 C 1 a]	Abfallverbrennung illegal
2 D 3 a [2 D 3 d]	Farben-Anwendung Holz	5 C 1 [5 C 1 b i]	Kabelabbrand
2 D 3 a [2 D 3 d]	Farben-Anwendung Autoreparatur	5 C 1 [5 C 1 b iii]	Spitalabfallverbrennung
2 D 3 a [2 D 3 e]	Elektronik-Reinigung	5 C 1 [5 C 1 b iv]	Klärschlammverbrennung
2 D 3 a [2 D 3 e]	Metallreinigung	5 C 1 [5 C 1 b v]	Krematorien
2 D 3 a [2 D 3 e]	Reinigung Industrie übrige	5 C 2	Abfallverbrennung Land- und Forstwirtschaft
2 D 3 a [2 D 3 f]	Chemische Reinigung**	5 D 1 [5 D]	Kläranlagen kommunal (Luftschadstoffe)
2 D 3 a [2 D 3 g]	Druckfarben Produktion	5 D 2 [5 D]	Kläranlagen industriell (Luftschadstoffe)
2 D 3 a [2 D 3 g]	Farben-Produktion	5 D 1 / 5 D 2 [5 D]	Kläranlagen GHG
2 D 3 a [2 D 3 g]	Feinchemikalien-Produktion**	5 E	Shredder Anlagen
2 D 3 a [2 D 3 g]	Gummi-Verarbeitung**	6 A d	Brand- und Feuerschäden Immobilien
2 D 3 a [2 D 3 g]	Klebband-Produktion	6 A d	Brand- und Feuerschäden Motorfahrzeuge

* confidential process

** confidential EMIS comment

*** work in progress

Prozess nicht relevant für Inventar ab 1990

Neue Kommentare