



# Greenhouse Gas Emissions in the Netherlands 1990-2008 National Inventory Report 2010



# Greenhouse Gas Emissions in the Netherlands 1990-2008 National Inventory Report 2010

C.W.M. van der Maas  
P.W.H.G. Coenen<sup>1</sup>  
P.J. Zijlema<sup>2</sup>  
K. Baas<sup>3</sup>  
G. van den Berghe<sup>2</sup>  
G.J. van den Born  
A.T. Brandt<sup>4</sup>  
B. Guis<sup>3</sup>  
G. Geilenkirchen  
R. te Molder  
D.S. Nijdam  
C.J. Peek  
S.M. van der Sluis

<sup>1</sup> Netherlands Organization for Applied Scientific Research (TNO),  
P.O. Box 80015, NL-3508-TA Utrecht

<sup>2</sup> NL Agency, PO Box 8242, NL-3503 RE Utrecht

<sup>3</sup> Statistics Netherlands (CBS), P.O. Box 24500, NL-2490 HA Den Haag

<sup>4</sup> Dutch Emission Authority, P.O.Box 91503, IPC 652, 2509 EC Den Haag



### **Greenhouse Gas Emissions in the Netherlands 1990-2008**

© Netherlands Environmental Assessment Agency (PBL), Bilthoven, April 2010

PBL report 500080017 / 2010

#### **Contacts:**

Wim van der Maas (PBL/IMP) ([wim.van.der.maas@rivm.nl](mailto:wim.van.der.maas@rivm.nl))

Peter Zijlema (NIE/SenterNovem) ([p.zijlema@agentschapnl.nl](mailto:p.zijlema@agentschapnl.nl))

This investigation has been performed by order and for the account of the Directorate-General for Environmental Protection, Climate Change and Industry Division, of the Netherlands Ministry of Spatial Planning, Housing and the Environment, within the framework of project Emissieregistratie M/500080/NIR, 'Netherlands Pollutant Emission Register'.

Report prepared for submission in accordance with The UN Framework Convention on Climate Change (UNFCCC) and the European Union's Greenhouse Gas Monitoring Mechanism [Including electronic Excel spreadsheet files containing the Common Reporting Format (CRF) data for 1990 to 2008]

Netherlands Environmental Assessment Agency (PBL)

P.O. Box 303

3720 AH Bilthoven

the Netherlands

Tel: +31-30-274 274 5

Fax: +31-30-274 44 79

[www.pbl.nl/en](http://www.pbl.nl/en)

# Acknowledgements

Many colleagues from a number of organisations (CBS, EC-LNV, LEI, Alterra, NL Agency, PBL, RIVM and TNO) have been involved in the annual update of the Netherlands Pollutant Release & Transfer Register (PRTR), also called the Emission Registration (ER) system, which contains emissions data on about 350 pollutants. The emission calculations, including those for greenhouse gas emissions, are performed by members of the ER 'Task Forces'. This is a major task, since the Netherlands' inventory contains many detailed emission sources.

Subsequently, the emissions and activity data of the Netherlands' inventory is converted into the IPCC source categories contained in the CRF files, which form a supplement to this report.

We are very grateful to Mr Bert Leekstra who automated this process to create the CRF files. This way we can avoid a second labour intensive quality check on the data.

The description of sources, analysis of trends and uncertainty estimates in emissions (see Chapters 3 to 8) of the various sources has been made in cooperation with the following emission experts: Mr Guus van den Berghe (NL Agency) (waste), Mr Gert-Jan van der Born (land use), Mr Gerben Geilenkirchen (transport), Mr Romuald te Molder (trends, key sources), Mr Durk Nijdam (solvent and product use), Mr Jos Olivier (energy), Mr Kees Peek (fugitive, industrial processes, other waste), Mr Kees Baas (CBS) (wastewater handling), Mrs Marian van Schijndel and Ms Sietske van der Sluis (agriculture). In addition, Mr Bas Guis of CBS has provided pivotal information on CO<sub>2</sub> related to energy use. This group has also provided activity data and additional information for the CRF files in cases where these were not included in the data sheets submitted by the ER Task Forces. We are particularly grateful to Mrs Marian Abels, Mr Jack Pesik, Mr Dirk Wever and Mr Kees Peek, for their contribution to data processing, chart production and quality control.

We greatly appreciate the contributions of each of these groups and individuals to this National Inventory Report and supplemental CRF files, as well as the external reviewers that provided comments on the draft report.



# Rapport in het kort

## Broeikasgasemissies in Nederland 1990-2008

Dit rapport over de Nederlandse inventarisatie van broeikasgasemissies is op verzoek van het Ministerie van VROM opgesteld om te voldoen aan de nationale rapportageverplichtingen in 2010 van het Klimaatverdrag van de Verenigde Naties (UNFCCC) en van het Bewakingsmechanisme Broeikasgassen van de Europese Unie. Dit rapport bevat de volgende informatie:

- trendanalyses voor de emissies van broeikasgassen in de periode 1990-2008;
- een analyse van zogenaamde sleutelbronnen en de onzekerheid in hun emissies volgens de 'Tier 1'-methodiek van het IPCC-rapport over Good Practice guidance;
- documentatie van gebruikte berekeningsmethoden, databronnen en toegepaste emissiefactoren;
- een overzicht van het kwaliteitssysteem en de validatie van de emissiecijfers voor de Nederlandse Emissieregistratie.

Geconcludeerd wordt dat de emissies van de zes broeikasgassen, uitgedrukt in CO<sub>2</sub> equivalenten, in 2008 in totaal met bijna 3% gedaald zijn ten opzichte van het basisjaar [1990 voor CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O en 1995 voor HFK's, PFK's en SF<sub>6</sub> (F-gassen); exclusief het bos- en landgebruik (LULUCF)]. Emissie van CO<sub>2</sub> exclusief LULUCF is in de periode 1990-2008 met 10% gestegen, terwijl de emissies van CH<sub>4</sub> en N<sub>2</sub>O met respectievelijk 33% en 42% zijn gedaald over dezelfde periode. De emissies van F-gassen zijn in de periode 1995-2008 met gemiddeld 71% afgenomen. De emissies van HFK's en PFK's daalden in die periode met respectievelijk 68% en 87%. De SF<sub>6</sub> emissies daalden met 26%.

Ten opzichte van 2007 zijn in 2008 zowel de CO<sub>2</sub>-emissies als de CH<sub>4</sub>-emissies licht gestegen (circa 2 en 1%) en de N<sub>2</sub>O-emissies sterk gedaald (-24%). Dit laatste is het gevolg van maatregelen in de industrie. De totale broeikasgasemissies zijn in 2008 vrijwel gelijkgebleven ten opzichte van 2007.

Trefwoorden: broeikasgassen, emissies, trends, methodiek, klimaat





# Contents

- Acknowledgements 5
- Rapport in het kort 7
  - Broeikasgasemissies in Nederland 1990-2008 7
- Samenvatting 13
  - National Inventory Report (NIR) 13
  - Ontwikkeling van de broeikasgasemissies 13
  - Methoden 13
- Executive Summary 15
  - ES1 Background information on greenhouse gas inventories and climate change 15
  - ES2 Summary of national emission and removal related trends 16
  - ES3 Overview of source and sink category emission estimates and trends 16
  - ES4 Other information 17
- Part I: Annual Inventory Report 21
- 1 Introduction 23
  - 1.1 Background information on greenhouse gas inventories and climate change 23
  - 1.2 Institutional arrangements for inventory preparation 24
  - 1.3 Inventory preparation 25
  - 1.4 Brief description of methodologies and data sources used 26
  - 1.5 A brief description of the key categories 28
  - 1.6 Information on the QA/QC plan 28
  - 1.7 Evaluating general uncertainty 31
  - 1.8 General assessment of the completeness 33
- 2 Trends in greenhouse gas emissions 35
  - 2.1 Emission trends for aggregated greenhouse gas emissions 35
  - 2.2 Emission trends by gas 36
  - 2.3 Emission trends specified by source category 39
  - 2.4 Emission trends for indirect greenhouse gases and SO<sub>2</sub> 39
- 3 Energy [CRF Sector 1] 43
  - 3.1 Overview of sector 43
  - 3.2 Fuel Combustion [1A] 44
  - 3.3 Fugitive emissions from fuels [1B] 65
- 4 Industrial processes [CRF Sector 2] 69
  - 4.1 Overview of sector 69
  - 4.2 Mineral products [2A] 70
  - 4.3 Chemical industry [2B] 73
  - 4.4 Metal production [2C] 76
  - 4.5 Food and drink production [2D] 78
  - 4.6 Production of halocarbons and SF<sub>6</sub> [2E] 78
  - 4.7 Consumption of halocarbons and SF<sub>6</sub> [2F] 79
  - 4.8 Other industrial processes [2G] 81

■	5	Solvent and other product use [CRF Sector 3]	83
	5.1	Overview of sector	83
	5.2	Indirect CO <sub>2</sub> emissions from Solvents and product use (Paint application [3A], Degreasing and dry cleaning [3B] and Other [3D])	84
	5.3	Miscellaneous N <sub>2</sub> O emissions from solvents and product use (use of N <sub>2</sub> O for anaesthesia [3D1] and N <sub>2</sub> O from aerosol cans [3D3])	85
■	6	Agriculture [CRF Sector 4]	87
	6.1	Overview of the sector	87
	6.2	Enteric fermentation [4A]	90
	6.3	Manure management [4B]	92
	6.4	Agricultural soils [4D]	96
■	7	Land use, land use change and forestry [CRF Sector 5]	101
	7.1	Overview of sector	101
	7.2	Methods	101
	7.3	Data	102
	7.4	Recalculations	102
	7.5	Forest Land [5A]	102
	7.6	Cropland [5B]	106
	7.7	Grassland [5C]	107
	7.8	Wetland [5D]	108
	7.9	Settlement [5E]	108
	7.10	Other Land [5F]	109
	7.11	Other [5G]	110
■	8	Waste [CRF Sector 6]	111
	8.1	Overview of sector	111
	8.2	Solid waste disposal on land [6A]	113
	8.3	Wastewater handling [6B]	114
	8.4	Waste incineration [6C]	116
	8.5	Other waste handling [6D]	118
■	9	Other [CRF Sector 7]	119
■	10	Recalculations and improvements	121
	10.1	Explanation and justification for the recalculations	121
	10.2	Implications for emission levels	123
	10.3	Implications for emission trends, including time-series consistency	123
	10.4	Recalculations, response to the review process and planned improvements	123
■	Part II: Supplementary Information required under Article 7, Paragraph 1		127
■	11	KP-LULUCF	129
	11.1	General information	129
	11.2	Land-related information	129
	11.3	Activity-specific information	131
	11.4	Article 3.3	134
	11.5	Article 3.4	134
	11.6	Other information	134
	11.7	Information relating to Article 6	134
■	12	Information on accounting of Kyoto units	135
	12.1	Background information	135
	12.2	Summary of information reported in the SEF tables	135
	12.3	Discrepancies and notifications	135
	12.4	Publicly accessible information	136
	12.5	Calculation of the commitment period reserve (CPR)	136
	12.6	KP-LULUCF accounting	136
■	13	Information on changes in national system	137

■	14	Information on changes in national registry	139
	14.1	Previous Review Recommendations	139
	14.2	Changes to National Registry	141
■	15	Information on minimisation of adverse impacts in accordance with Article 3, paragraph 14	143
■	16	Other information	147
■		Annexes	149
■		Annex 1 Key sources	151
	A1.1	Introduction	151
	A1.2	Changes in key sources compared to previous submission	156
	A1.3	Tier 1 key source and uncertainty assessment	156
■		Annex 2 Detailed discussion of methodology and data for estimating CO <sub>2</sub> emissions from fossil fuel combustion	173
	A2.1	Introduction	173
	A2.2	Starting points for the Netherlands list	173
	A2.3	The Netherlands list	174
	A2.4	Fact sheets	176
	A2.5	Using the Netherlands list in national monitoring, European CO <sub>2</sub> emissions trade and in the e-MJV National monitoring	176
	A2.6	Defining and maintaining the Netherlands list	177
	A2.7	Application of the Netherlands standard and source-specific CO <sub>2</sub> emission factors in the national emission inventory	177
■		Annex 3 Other detailed methodological descriptions for individual source or sink categories	179
■		Annex 4 CO <sub>2</sub> Reference Approach and comparison with Sectoral Approach	181
	A4.1	Comparison of CO <sub>2</sub> emissions	181
	A4.2	Causes of differences between the two approaches	181
	A4.3	Other country-specific data used in the Reference Approach	182
	A4.4	Feedstock component in the CO <sub>2</sub> Reference Approach	183
■		Annex 5 Assessment of completeness and (potential) sources and sinks	185
■		Annex 6 Additional information to be considered as part of the NIR submission	187
	A6.1	List of protocols	187
	A6.2	Documentation of uncertainties used in IPCC Tier 1 uncertainty assessments and Tier 2 key source identification	188
	A6.3	Background documents and uncertainty discussion papers	188
	A6.4	Documentation of Quality Assurance and Quality Control for national greenhouse gas inventory compilation and reporting	188
	A6.5	Documentation of Changes to the National Registry	188
		Registry Information	189
■		Annex 7 Tables 6.1 and 6.2 of the IPCC Good Practice guidance	207
■		Annex 8 Emission Factors and Activity Data Agriculture	209
■		Annex 9 Chemical compounds, global warming potentials, units and conversion factors	223
	A9.1	Chemical compounds	223
	A9.2	Global Warming Potentials for selected greenhouse gases	223
	A9.3	Units	224
	A9.4	Other conversion factors for emissions	224
■		Annex 10 List of abbreviations	225
■		References	227



# Samenvatting

Het National Inventory Report (NIR) 2010 bevat de rapportage van broeikasgasemissies (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub> en de F-gassen) over de periode 1990 tot en met 2008.

De emissiecijfers in de NIR 2010 zijn berekend volgens de protocollen behorend bij het 'National System' dat is voorgeschreven in het Kyoto Protocol. In de protocollen zijn de methoden vastgelegd voor zowel het basisjaar (1990 voor CO<sub>2</sub>, CH<sub>4</sub> en N<sub>2</sub>O en 1995 voor de F-gassen) als voor de emissies in de periode tot en met 2012. De protocollen staan op de website [www.broeikasgassen.nl](http://www.broeikasgassen.nl). Alle emissiecijfers en bijbehorende documentatie worden ook, met een vertraging van enkele maanden, gepubliceerd op [www.emissieregistratie.nl](http://www.emissieregistratie.nl)

## National Inventory Report (NIR)

Dit rapport over de Nederlandse inventarisatie van broeikasgasemissies is op verzoek van het ministerie van VROM opgesteld om te voldoen aan de nationale rapportageverplichtingen in 2010 van het Klimaatverdrag van de Verenigde Naties (UNFCCC) het Kyoto protocol en het Bewakingsmechanisme Broeikasgassen van de Europese Unie. Dit rapport bevat de volgende informatie:

- trendanalyses voor de emissies van broeikasgassen in de periode 1990-2008
- een analyse van zogenaamde sleutelbronnen en de onzekerheid in hun emissies volgens de 'Tier 1'-methodiek van het IPCC-rapport over Good Practice guidance;
- documentatie van gebruikte berekeningsmethoden, databronnen en toegepaste emissiefactoren;
- een overzicht van het kwaliteitssysteem en de validatie van de emissiecijfers voor de Nederlandse Emissieregistratie;
- de wijzigingen die in de methoden voor het berekenen van broeikasgasemissies zijn aangebracht na de review van het Nationaal Systeem broeikasgassen vanuit het Klimaatverdrag. Op basis van de methoden die in de NIR en de Nederlands protocollen broeikasgassen zijn vastgelegd, is de basisjaaremisse bepaald, en de hoeveelheid broeikasgassen die Nederland in de periode 2008 t/m 2012 (volgens het Kyoto Protocol) mag uitstoten.

Vanaf dit jaar bevat de NIR ook de informatie die voorgeschreven is volgens artikel 7 van het Kyoto protocol. Door deze toevoeging voldoet Nederland aan de meest recente rapportagerichtlijnen van de UNFCCC.

Een losse annex bij dit rapport bevat elektronische data over emissies en activiteitsdata in het zogenaamde Common Reporting Format (CRF), waar door het secretariaat van het VN-Klimaatverdrag om wordt verzocht. In de bijlagen bij dit rapport zijn de samenvattende emissie- en trendtabellen '7A' en 10 op basis van het CRF opgenomen voor 1990-2008. Daarnaast bevatten de bijlagen ondermeer een overzicht van sleutelbronnen en onzekerheden.

De NIR gaat niet specifiek in op de invloed van het gevoerde overheidsbeleid met betrekking tot emissies van broeikasgassen; meer informatie hierover is te vinden in de Milieubalans en de vierde Nationale Communicatie onder het Klimaatverdrag, die eind 2005 is verschenen.

## Ontwikkeling van de broeikasgasemissies


De emissieontwikkeling in Nederland wordt beschreven en toegelicht in dit National Inventory Report (NIR 2010). Figuur ES.1 geeft het emissieverloop over de periode 1990-2008 weer. De totale emissies bedroegen in 2008 circa 206,9 Tg (Mton ofwel miljard kg) CO<sub>2</sub> equivalenten en waren daarmee bijna drie procent lager (Box ES.1) dan de emissies in het basisjaar (213,3 Tg CO<sub>2</sub> eq). De hier gepresenteerde emissies zijn exclusief de emissies van landgebruik en bossen (LULUCF); deze emissies tellen pas mee vanaf het emissiejaar 2008 (waarover in deze submitisie wordt gerapporteerd) onder het Kyoto Protocol. De emissie van CO<sub>2</sub> is sinds 1990 met circa 10% toegenomen, terwijl de emissies van de andere broeikasgassen met circa 42% zijn afgenomen ten opzichte van het basisjaar.

In 2008 daalde N<sub>2</sub>O emissie verder door de maatregelen bij de salpeterzuurproductie.

In 2008 bedroeg de N<sub>2</sub>O emissie 11.8 miljard kg ten opzichte van 15.4 miljard kg CO<sub>2</sub> eq in 2007. Deze emissiereductie wordt teniet gedaan door een stijging van de CO<sub>2</sub> emissie die thans weer op het niveau van 2005 ligt. De totale emissie van broeikasgassen in 2008 is op het zelfde niveau gebleven als in 2007.

## Methoden

De methoden die Nederland hanteert voor de berekening van de broeikasgasemissies zijn vastgelegd in protocollen,



te vinden op [www.broeikasgassen.nl](http://www.broeikasgassen.nl). De protocollen zijn opgesteld door Agentschap NL, in nauwe samenwerking met deskundigen van de Emissieregistratie (voor wat betreft de beschrijving en documentatie van de berekeningsmethoden). Na vaststelling van deze protocollen in de Stuurgroep EmissieRegistratie (december 2005), zijn de protocollen vastgelegd in een wettelijke regeling door het ministerie van VROM. De methoden maken onderdeel uit van het Nationaal Systeem (artikel 5.1 van het Kyoto Protocol) en zijn bedoeld voor de vaststelling van de emissies in zowel het basisjaar als in de jaren in de budgetperiode. Naar aanleiding van de review van het zogenaamde 'Initial Report' zijn de

methoden en protocollen aangepast. Deze zijn daarmee in overeenstemming met de IPCC Good Practice guidance and Uncertainty Management, dat als belangrijkste voorwaarde is gesteld aan de te hanteren methoden voor de berekening van broeikasgassen. Deze methoden zullen de komende jaren (tot 2014) worden gehanteerd; tenzij er grote veranderingen plaatsvinden in bijvoorbeeld de beschikbaarheid van basisdata of de implementatie van beleidsmaatregelen aanleiding geeft de methoden aan te passen.

### Box ES.1 Onzekerheden

De emissies van broeikasgassen kunnen niet exact worden gemeten of berekend. Onzekerheden zijn daarom onvermijdelijk. Het PBL schat de onzekerheid in de jaarlijkse totale broeikasgasemissies op circa 4%. Dit is geschat op basis van informatie van emissie-experts in een eenvoudige analyse van de onzekerheid (volgens IPCC Tier 1). De totale uitstoot van broeikasgassen

ligt daarmee met 95% betrouwbaarheid tussen de 197 en 218 Tg (Mton). De onzekerheid in de emissietrend tussen het basisjaar (1990/1995) en 2008 is geschat op circa 3%; dat wil zeggen dat de emissietrend in die periode met 95% betrouwbaarheid ligt tussen de -6 tot +0%.

# Executive Summary

## ES1 Background information on greenhouse gas inventories and climate change

This report documents the 2010 Netherlands' annual submission of its greenhouse gas emission inventory in accordance with the guidelines provided by the United Nations Framework Convention on Climate Change (UNFCCC), the Kyoto Protocol and the European Union's Greenhouse Gas Monitoring Mechanism. These guidelines, which also refer to Revised 1996 IPCC Guidelines and IPCC Good Practice guidance and Uncertainty Management reports, provide a format for the definition of source categories and for calculation, documentation and reporting of emissions. The guidelines aim at facilitating verification, technical assessment and expert review of the inventory information by independent Expert Review Teams of the UNFCCC. Therefore, the inventories should be transparent, consistent, comparable, complete and accurate as elaborated in the UNFCCC Guidelines for reporting and be prepared using good practice as described in the IPCC Good Practice Guidance. This National Inventory Report (NIR) 2010, therefore, provides explanations of the trends in greenhouse gas emissions, activity data and (implied) emission factors for the period 1990-2008. It also summarises descriptions of methods and data sources of Tier 1 assessments of the uncertainty in annual emissions and in emission trends; it presents an assessment of key sources following the Tier 1 and Tier 2 approaches of the IPCC Good Practice Guidance; and describes Quality Assurance and Quality Control activities. This report provides no specific information on the effectiveness of government policies for reducing greenhouse gas emissions. This information can be found in the annual Environmental Balance (in Dutch: 'Milieubalans') prepared by the Netherlands Environmental Assessment Agency (PBL) and the 4<sup>th</sup> National Communication (NC4) prepared by the Government of the Netherlands.

The Common Reporting Format (CRF) spreadsheet files, containing data on emissions, activity data and implied emission factors, accompany this report. The complete set of CRF files as well as the NIR in pdf format can be found at the website [www.greenhousegases.nl](http://www.greenhousegases.nl). From July 2010, the emissions and documentation can also be found on [www.prtr.nl](http://www.prtr.nl)

### Climate Convention and Kyoto Protocol

This NIR is prepared as a commitment under the UNFCCC and under the Kyoto Protocol. The NIR now also contains

a part 2 that focuses on supplementary information under article 7 of the Kyoto protocol. One of the commitments is the development of a National System for greenhouse gas emissions (art. 5.1 of the Protocol). This National System developed in the period 2000-2005 was reviewed by an Expert Review Team of the UNFCCC in April 2007 and found to be in compliance with the requirements.


### Key categories

For identification of the 'key categories' according to the IPCC Good Practice approach, national emissions are allocated according to the IPCC potential key category list wherever possible. The IPCC Tier 1 method consists of ranking this list of source category-gas combinations for the contribution to both the national total annual emissions and the national total trend. The results of these listings are presented in Annex 1: the largest sources, the total of which adds up to 95% of the national total, are 32 sources for annual level assessment and 30 sources for the trend assessment from a total of 70 sources. Both lists can be combined to give an overview of sources, which meet either of these two criteria. Next, the IPCC Tier 2 method for identification of key sources is used, which requires incorporating the uncertainty to each of these sources before ordering the list of shares. The result is a list of 43 source categories from a total of 70 that could be identified as 'key sources' according to the definition of the IPCC Good Practice Guidance report. Finally, four key categories are found in the LULUCF sector (Sector 5), after inclusion of 9 LULUCF subcategories in the key category analysis.

### Institutional arrangements for inventory preparation

The greenhouse gas inventory of the Netherlands is based on the national Pollutant Release & Transfer Register (PRTR). The general process of inventory preparation has existed many years and is organised as a project with an annual cycle. In 2000, an improvement programme was initiated under the lead of SenterNovem (now NL Agency) to transform the general process of the greenhouse gas inventory of the PRTR into a National System, according to the requirements under article 5.1 of the Kyoto Protocol.

The Netherlands Environmental Assessment Agency (PBL) has been contracted by the Ministry of Housing, Spatial Planning and the Environment (VROM) to compile and maintain the PRTR and to co-ordinate the preparation of the NIR and filling the CRF (see Figure ES.2). NL Agency is designated by law as the National Inventory Entity (NIE) and co-ordinates



the overall QA/QC activities and the support/response to the UNFCCC review process.

### Monitoring protocols

As part of the improvement programme, the methodologies for calculating greenhouse gas emission in the Netherlands were reassessed and compared with UNFCCC and IPCC requirements. For the key sources and for sinks, the methodologies and processes are elaborated, re-assessed and revised when required. The final revision was done after review of the National System (including the protocols). The present CRF/NIR is based on methodologies approved during/after the review of the National System and the calculation of the Assigned Amount of the Netherlands. Monitoring protocols describing methodologies, data sources and the rationale for their selection are available at [www.greenhousegases.nl](http://www.greenhousegases.nl).

### Organisation of the report

This report is in line with the prescribed NIR format, starting with an introductory Chapter 1, containing background information on the Netherlands' process of inventory preparation and reporting; key categories and their uncertainties; a description of methods, data sources and emission factors, and a description of the quality assurance system, along with verification activities applied to the data. Chapter 2 provides a summary of trends for aggregated greenhouse gas emissions by gas and by main source. Chapters 3 to 9 present detailed explanations for emissions in different sectors. Chapter 10 presents information on recalculations, improvements and response to issues raised in external reviews on the NIR 2009 and on the draft of the NIR 2010. In addition, the report provides detailed information on key categories, methodologies and other relevant reports in 10 Annexes.

### ES2 Summary of national emission and removal related trends

In 2008, total direct greenhouse gas emissions (excluding emissions from LULUCF) in the Netherlands were estimated

at 206.9 Tg CO<sub>2</sub> equivalents (CO<sub>2</sub> eq). This is 3% below the emissions in the base year (213.3 Tg CO<sub>2</sub> eq). In the Netherlands, the base year emissions are 1990 for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, and 1995 for fluorinated gases. CO<sub>2</sub> emissions (excluding LULUCF) increased by about 10% from 1990 to 2008, mainly due to the increase in the emissions in the 1A1a Public electricity sector and 1A3 Transport sector. CH<sub>4</sub> emissions decreased by 33% in 2008 compared to the 1990 level, mainly due to decrease in the waste sector, the agricultural sector and fugitive emissions in the energy sector. N<sub>2</sub>O emissions decreased by about 42% in 2008 compared to 1990, mainly due to decrease in emissions from agriculture and from industrial processes, which partly compensated emission increases from fossil fuel combustion (mainly from transport). Of the fluorinated greenhouse gases, HFC and PFC emissions decreased in 2008 by about 68% and 87%, respectively, while SF<sub>6</sub> emissions decreased by 26%. Total emissions of all F-gases decreased by about 71% compared to the 1995 level (chosen as the base year).


Between 2007 and 2008, total greenhouse gas emissions (excluding LULUCF) stabilised. CO<sub>2</sub> emissions increased (excluding LULUCF) by 3.4 Tg. The total greenhouse gas emission remain fairly constant because N<sub>2</sub>O emissions dropped by 3.7 Tg CO<sub>2</sub> eq. This is caused by nitric acid production the emission of which decreased by approximately 12.1 Gg N<sub>2</sub>O in the period 2007-2008.

### ES3 Overview of source and sink category emission estimates and trends

Tables ES.1 and ES.2 provide an overview of the emission trends (in CO<sub>2</sub> equivalents) per gas and per IPCC source category. The Energy sector (category 1) is by far the largest contributor to national total greenhouse gas emissions. The emissions of this sector increased substantially compared to 1990. In contrast, emissions from the other sectors decreased compared to the base year, the largest being Industrial Processes, Waste and Agriculture.

Sectors showing the largest growth in CO<sub>2</sub> equivalent emissions since 1990 are Transport (1A3) and Energy





industries (1A1) (+37% and +24%, respectively). Half the marked increase in the public electricity sector of almost 30% between 1990 and 1998 is caused by a shift of cogeneration plants from manufacturing industries to the public electricity and heat production sector due to a change of ownership (joint ventures), simultaneously causing a 15% decrease in industry emissions in the early 1990s (1A2).

#### ES4 Other information

##### General uncertainty evaluation

The results of the uncertainty estimation according to the IPCC Tier 1 uncertainty approach are summarised in Annex 1 of this report. The Tier 1 estimation of annual uncertainty in CO<sub>2</sub> eq emissions results in an overall uncertainty of 4%, based on calculated uncertainties of 2%, 16%, 60% and 36% for CO<sub>2</sub> (excluding LULUCF), CH<sub>4</sub>, N<sub>2</sub>O and F-gases, respectively. However, these figures do not include the correlation between source categories (e.g., cattle numbers for enteric fermentation and animal manure production), or a correction for not-reported sources. Therefore, the actual uncertainty of

total annual emissions per compound and of the total will be somewhat higher; it is currently estimated by PBL at:

CO <sub>2</sub>	±3%	HFCs	±50%
CH <sub>4</sub>	±25%	PFCs	±50%
N <sub>2</sub> O	±65%	SF <sub>6</sub>	±50%
Total greenhouse gas emissions			±5%

Table A1.4 of Annex 1 summarises the estimate of the trend uncertainty 1990-2008 calculated according to the IPCC Tier 1 approach in the IPCC Good Practice Guidance (IPCC, 2001). The result is a trend uncertainty in the total CO<sub>2</sub> eq emissions (including LULUCF) for 1990-2008 (1995 for F-gases) of ±3% points. This means that the decrease in total CO<sub>2</sub> eq emissions between 1990 and 2008 (including LULUCF), which is calculated to be 3%, will be between -6% and +0%. Per individual gas, the trend uncertainty in total emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and the total group of F-gases has been calculated at ±3%, ±9%, ±14% and ±10% points, respectively. More details on the level and trend uncertainty assessment can be found in Annex 7.

Summary of emission trend per gas (unit: Tg CO<sub>2</sub> equivalents)

Table ES.1

	CO <sub>2</sub> incl. LULUCF	CO <sub>2</sub> excl. LULUCF	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	SF <sub>6</sub>	Total (incl. LULUCF)	Total (excl. LULUCF)
<i>Base yr</i>	161.9	159.3	25.5	20.2	6.0	1.9	0.3	215.9	213.3
1990	161.9	159.3	25.5	20.2	4.4	2.3	0.2	214.6	212.0
1991	166.5	164.3	25.9	20.6	3.5	2.2	0.1	218.8	216.8
1992	164.8	162.3	25.5	21.5	4.4	2.0	0.1	218.5	216.0
1993	169.0	166.8	25.2	21.9	5.0	2.1	0.1	223.4	221.1
1994	168.9	166.7	24.4	21.4	6.5	2.0	0.2	223.4	221.2
1995	172.9	170.6	24.1	21.5	6.0	1.9	0.3	226.9	224.5
1996	179.8	177.7	23.5	21.4	7.7	2.2	0.3	234.8	232.7
1997	173.5	171.1	22.5	21.1	8.3	2.3	0.3	228.1	225.7
1998	175.6	173.2	21.7	20.7	9.3	1.8	0.3	229.5	227.2
1999	170.2	167.8	20.7	20.0	4.9	1.5	0.3	217.6	215.2
2000	172.2	169.7	19.8	19.3	3.9	1.6	0.3	217.1	214.6
2001	177.6	175.2	19.3	18.1	1.6	1.5	0.3	218.4	216.0
2002	178.1	175.7	18.4	17.3	1.7	2.2	0.3	217.9	215.6
2003	182.0	179.7	17.9	16.9	1.5	0.6	0.2	219.2	216.8
2004	183.3	181.0	17.6	17.4	1.6	0.3	0.3	220.5	218.2
2005	178.2	175.8	17.2	17.3	1.5	0.3	0.3	214.7	212.4
2006	174.9	172.5	16.8	17.1	1.7	0.3	0.2	211.1	208.7
2007	174.8	172.3	16.8	15.4	1.8	0.3	0.2	209.5	206.9
2008	178.1	175.7	17.1	11.8	1.9	0.3	0.2	209.4	206.9

Summary of emission trend per source category (unit: Tg CO<sub>2</sub> equivalents)

Table ES.2

	1. Energy	2. Ind. Proc.	3. Solvents	4. Agriculture	5. LULUCF	6. Waste	7. Other	Total (incl. LULUCF)	Total (excl. LULUCF)
<i>Base yr</i>	154.0	22.2	0.5	22.5	2.6	12.8	NA	215.9	213.3
1990	154.0	22.2	0.5	22.5	2.6	12.8	NA	214.6	212.0
1991	159.1	21.2	0.5	23.0	2.2	12.9	NA	218.8	216.6
1992	157.8	21.5	0.4	23.6	2.5	12.7	NA	218.5	216.0
1993	162.5	22.3	0.4	23.5	2.2	12.3	NA	223.4	221.1
1994	161.8	24.3	0.4	22.8	2.2	11.9	NA	223.4	221.2
1995	165.8	23.6	0.4	23.5	2.3	11.3	NA	226.9	224.5
1996	173.6	24.8	0.4	22.9	2.2	10.9	NA	234.8	232.7
1997	166.0	26.1	0.3	22.6	2.4	10.6	NA	228.1	225.7
1998	168.2	26.4	0.4	22.0	2.3	10.2	NA	229.5	227.2
1999	162.6	21.2	0.4	21.6	2.4	9.4	NA	217.5	215.2
2000	164.7	20.2	0.3	20.4	2.5	8.9	NA	217.1	214.6
2001	170.7	16.7	0.3	19.9	2.4	8.4	NA	218.4	216.0
2002	171.3	17.1	0.2	18.9	2.3	8.0	NA	217.9	215.6
2003	175.1	15.6	0.2	18.4	2.3	7.5	NA	219.2	216.8
2004	176.3	16.0	0.2	18.5	2.3	7.2	NA	220.5	218.2
2005	171.2	15.7	0.2	18.5	2.4	6.8	NA	214.7	212.4
2006	167.9	15.8	0.2	18.4	2.4	6.4	NA	211.1	208.7
2007	167.8	14.7	0.2	18.3	2.5	6.0	NA	209.5	206.9
2008	172.1	10.5	0.2	18.5	2.4	5.7	NA	209.4	206.9

### Completeness of the national inventory

The Netherlands' greenhouse gas emission inventory includes all sources identified by the Revised Intergovernmental Panel on Climate Change (IPCC) Guidelines (IPCC, 1996) – with the exception of the following very minor sources:

- oil transport (1B2a3), due to missing activity data
- charcoal production (1B2) and use (1A4), due to missing activity data
- CO<sub>2</sub> from asphalt roofing (2A5), due to missing activity data
- CO<sub>2</sub> from road paving (2A6), due to missing activity data
- CH<sub>4</sub> from enteric fermentation of poultry (4A9), due to missing emission factors

- N<sub>2</sub>O from industrial waste water (6B1), due to negligible amounts
- Precursor emissions (carbon monoxide (CO), nitrogen oxide (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO<sub>2</sub>)) from Memo item "International bunkers" (international transport) have not been included.

For more information on this issue, see Annex 5.

Differences between NIR 2009 and NIR 2010 due to recalculations (Unit: Tg CO<sub>2</sub> eq, F-gases: Gg CO<sub>2</sub> eq)

Table ES.3

Gas	Source	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
CO <sub>2</sub> Incl. LUCF	NIR09	161.9	173.0	172.1	177.6	178.1	182.0	183.3	178.2	174.9	175.2
	NIR10	<b>161.9</b>	172.9	172.2	177.6	178.1	182.0	183.3	178.2	174.9	174.8
	Diff.	0.0%	-0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	-0.2%
CO <sub>2</sub> Excl. LUCF	NIR09	159.3	170.7	169.6	175.2	175.7	179.6	181.0	175.8	172.5	172.7
	NIR10	<b>159.3</b>	170.6	169.7	175.3	175.8	179.7	181.0	175.8	172.5	172.3
	Diff.	0.0%	-0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	-0.2%
CH <sub>4</sub>	NIR09	25.5	24.2	19.8	19.3	18.4	17.9	17.6	17.2	16.8	17.0
	NIR10	25.5	24.1	19.8	19.3	18.4	17.9	17.6	17.2	16.8	16.8
	Diff.	0.0%	0.0%	-0.1%	-0.1%	-0.1%	-0.1%	0.0%	0.0%	0.0%	-0.7%
N <sub>2</sub> O	NIR09	20.2	21.5	19.3	18.1	17.3	16.9	17.4	17.3	17.1	15.6
	NIR10	20.2	21.5	19.3	18.1	17.3	16.9	17.4	17.3	17.1	15.4
	Diff.	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.2%
PFCs Gg	NIR09	2,264	1,938	1,582	1,489	2,187	621	286	266	257	327
	NIR10	2,264	<b>1,938</b>	1,582	1,489	2,187	621	286	266	257	323
	Diff.	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.2%
HFCs Gg	NIR09	4,432	6,020	3,829	1,469	1,541	1,377	1,507	1,358	1,566	1,738
	NIR10	4,432	<b>6,018</b>	3,891	1,561	1,653	1,511	1,650	1,514	1,728	1,845
	Diff.	0.0%	0.0%	1.6%	6.2%	7.2%	9.7%	9.5%	11.5%	10.3%	6.2%
SF <sub>6</sub> Gg	NIR09	217	301	319	323	283	243	246	238	202	214
	NIR10	217	<b>301</b>	318	322	281	242	265	254	217	226
	Diff.	0.0%	0.0%	-0.3%	-0.4%	-0.5%	-0.8%	7.5%	6.9%	7.1%	5.6%
Total Excl. LUCF	NIR09	212.0	224.7	214.4	215.9	215.5	216.7	218.0	212.2	208.5	207.5
	NIR10	212.0	224.5	214.6	216.0	215.6	216.8	218.2	212.4	208.7	206.9
	Diff.	0.0%	-0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	-0.3%
Total Incl. LUCF	NIR09	214.6	227.0	216.9	218.3	217.8	219.0	220.4	214.6	210.9	210.0
	NIR10	214.6	226.9	217.1	218.4	217.9	219.2	220.5	214.7	211.1	209.5
	Diff.	0.0%	-0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	-0.3%

Emission trends for indirect greenhouse gases and SO<sub>2</sub> (Unit: Gg)

Table ES.4

	1990	1995	2000	2005	2006	2007	2008
Total NO <sub>x</sub>	545	450	378	326	312	287	279
Total CO	1,108	895	757	665	656	636	643
Total NMVOC	459	322	230	175	166	163	163
Total SO <sub>2</sub>	188	128	71	63	63	59	50

### Methodological changes, recalculations and improvements

This NIR 2010 is based on the envisaged National System of the Netherlands under article 5.1 of the Kyoto Protocol, as developed in the last decade and finalised by December 2005. In past years the results of various improvement actions have been implemented in the methodologies and processes of the preparation of the greenhouse gas inventory of the Netherlands. Compared to the NIR/CRF 2009 and based on the results of the review of the National System by an Expert Review Team of the UNFCCC, some recalculations were undertaken in the last year. The major recalculations are the methodical change in the assessment of HFC emissions from stationary refrigeration and airconditioning, and an error correction in recent years for the oil and gas sector. The ratio behind the recalculations is documented in the Chapters 3-8 and 10.

Table ES.3 provides the results of recalculations in the NIR 2010 compared to the NIR 2009.

### Improving the QA/QC system

The QA/QC programme (quality assurance / quality control) is up to date and all procedures and processes have been

established to meet the National System requirements (as part of the annual activity programme of the Netherlands PRTR). QA/QC activities to be undertaken as part of the National System are described in Chapter 1. Some actions that remained since the NIR 2007 are now being implemented.

### Emission trends for indirect greenhouse gases and SO<sub>2</sub>

Compared to 1990, CO and NMVOC emissions were reduced in 2008 by 58% and 69%, respectively. For SO<sub>2</sub> this is 75%, and for NO<sub>x</sub>, the 2008 emissions are 50% lower than the 1990 level. Table ES.4 provides trend data.

In contrast to the direct greenhouse gases, emissions of precursors from road transport have not been corrected for fuel sales according to the national energy statistics but are directly related to transport statistics on vehicle-km, which differs to some extent from the IPCC approach.

Recalculations (due to changes in methodologies and/or allocation) have only been performed for 1990, 1995, 2000 and 2005 to 2008 for all sources.



# Part I: Annual Inventory Report





# Introduction

## 1.1 Background information on greenhouse gas inventories and climate change

### 1.1.1 Background information on climate change

The United Nations Framework Convention on Climate Change (UNFCCC) was ratified by the Netherlands in 1994 and entered into force in March of 1994. One of the commitments made by the ratifying Parties under the Convention is to develop, publish and regularly update national emission inventories of greenhouse gases. This national inventory report, together with the CRF, represent the 2010 national emission inventory of greenhouse gases under the UNFCCC (part 1 of this report) and under its Kyoto Protocol (part 2 of this report).

### Geographical coverage

The reported emissions include those that have to be allocated to the legal territory of the Netherlands. This includes a 12-mile zone from the coastline and also inland water bodies. It excludes Aruba and the Netherlands Antilles, which are self-governing dependencies of the Royal Kingdom of the Netherlands. Emissions from offshore oil and gas production on the Dutch part of the continental shelf are included.

### 1.1.2 Background information on greenhouse gas inventory

As indicated, this national inventory report documents the 2010 Greenhouse Gas Emission Inventory for the Netherlands under the UNFCCC and under the Kyoto Protocol. The estimates provided in the report are consistent with the Intergovernmental Panel on Climate Change (IPCC) 1996 Guidelines for National Greenhouse Gas Inventories (IPCC, 1997) and the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC, 2001) and the IPCC Good Practice Guidance for Land Use, Land Use Change and Forestry (LULUCF). The methodologies applied for the Netherlands' inventory are also consistent with the guidelines under the Kyoto Protocol and the European Union's Greenhouse Gas Monitoring Mechanism.

For detailed assessments of the extent to which changes in emissions are due to the implementation of policy measures, see the Environmental Balance (PBL, 2009; in Dutch), the Fourth and the Fifth Netherlands National Communication under the United Nations Framework Convention on Climate Change (VROM, 2005 resp. VROM, 2009) and the Netherlands Report on Demonstrable Progress under Article 3.2 of the Kyoto Protocol (VROM, 2006b).

The Netherlands also reports emissions under other international agreements, such as the United Nations Economic Commission for Europe (UNECE), Convention on Long Range Transboundary Air Pollutants (CLRTAP) and the EU National Emission Ceilings (NEC) Directive. All these estimates are provided by the Netherlands Pollutant Release and Transfer Register (PRTR), which is compiled by a special project in which various organisations cooperate. The greenhouse gas inventory and the PRTR share the same underlying data, which ensures consistency between the inventories and other internationally reported data. Several institutes are involved in the process of compiling the greenhouse gas inventory (see also Section 1.3).

The National Inventory Report (NIR) covers the six direct greenhouse gases included in the Kyoto Protocol: carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs) and sulphur hexafluoride (SF<sub>6</sub>) (the F-gases). Emissions of the following indirect greenhouse gases are also reported: nitrogen oxides (NO<sub>x</sub>), carbon monoxide (CO), and non-methane volatile organic compounds (NMVOC), as well as sulphur oxides (SO<sub>x</sub>).

This report provides explanations of the trends in greenhouse gas emissions per gas and per sector for the 1990–2008 period and summarises descriptions of methods and data sources for: (a) Tier 1 assessments of the uncertainty in annual emissions and in emission trends; (b) key source assessments following the Tier 1 and Tier 2 approaches of the IPCC Good Practice Guidance (IPCC, 2001); (c) quality assurance and quality control (QA/QC) activities.

Under the National System under Article 5.1 of the Kyoto Protocol, methodologies were established (and documented) in monitoring protocols. These protocols are annually re-assessed and revised, if needed, for example based on recommendations of UN reviews. The monitoring protocols and the general description of the National System are available on the website [www.greenhousegases.nl](http://www.greenhousegases.nl). The emissions reported in the NIR 2010 are based on these methodologies, which have been incorporated in the National System for greenhouse gases. The emissions and all the documentation are, with a delay of some months, also available on the website [www.prtr.nl](http://www.prtr.nl)

In 2007, the UN performed an in-country initial review under the Kyoto Protocol. The review concluded that

the Netherlands National System has been established in accordance with the guidelines and that it meets the requirements. This was confirmed also by later reviews such as the centralised review of the NIR 2009. The National System has remained unchanged with the exception of an organisational change per 1 January 2010. At that date, coordination of the above mentioned PRTR project (emissions registration project) shifted from PBL (Netherlands Environmental Assessment Agency) to RIVM (National Institute for Public Health and the Environment). The year 2010 is used as a transitional period. Arrangements are made to ensure that this transition has no consequences for the quality of the products of the PRTR project.

The structure of this report complies with the format required by the UNFCCC (FCCC/SBSTA/2004/8 and the latest annotated outline of the National Inventory report including reporting elements under the Kyoto protocol). It also includes supplementary information under Article 7 of the Kyoto Protocol. Part 2 gives an overview of this information.

Greenhouse gas emissions presented in this report are given in gigagrammes (Gg) and teragrammes (Tg). Global warming potential (GWP) weighed emissions of the greenhouse gases are also provided (in CO<sub>2</sub> equivalents), using the GWP values in accordance with the Kyoto Protocol and using the IPCC GWP for a time horizon of 100 years. The GWP of each individual greenhouse gas is provided individually in Annex 9.

The Common Reporting Format (CRF) spreadsheet files accompany this report as electronic annexes (the CRF files are included in the zip file for this submission: NETHERLANDS-2010-v1.3.zip). The CRF files contain detailed information on greenhouse gas emissions, activity data and (implied) emission factors specified by sector, source category and greenhouse gas. The complete set of CRF files as well as this report comprise the National Inventory Report (NIR) and are published on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Other information, such as protocols of the methods used to estimate emissions, is also available on this website. Section 10 provides details on the extent to which the CRF data files for 1990–2008 have been completed and on improvements made since the last submission.

### 1.1.3 Background information on supplementary information under Article 7 of the Kyoto Protocol

Part 2 of this report provides the supplementary information under (Article 7) of the Kyoto Protocol. As the Netherlands has not elected any activities to include under Article 3, paragraph 4 of the Kyoto Protocol, the supplementary information on KP LULUCF deals with activities under Article 3, paragraph 3. Information on the accounting of Kyoto units is also provided in the SEF file SEF\_NL\_2010\_2\_14-42-58 22-4-2010.xls and in the SIAR report file SIAR\_Reports\_2009\_NL v1.0.xls.

## 1.2 Institutional arrangements for inventory preparation

### 1.2.1 Overview of institutional arrangements for the inventory preparation

The Ministry of Housing, Spatial Planning and the Environment (VROM) has overall responsibility for climate change policy issues including the preparation of the inventory.

In August 2004, VROM assigned SenterNovem (now NL agency) executive tasks bearing on the National Inventory Entity (NIE), the single national entity required under the Kyoto Protocol. In December 2005, SenterNovem (now NL Agency) was designated by law as the NIE. In addition to coordinating the establishment and maintenance of a National System, the tasks of NL Agency include overall coordination of improved QA/QC activities as part of the National System and coordination of the support/response to the UNFCCC review process. The National System is described in more detail in the (Fourth and Fifth National Communication (VROM, 2009).

Since 1 January 2010, RIVM has been assigned by VROM to take over the role of PBL as coordinating institute for compiling and maintaining the pollutants emission register/inventory (PRTR system), containing about 350 pollutants including the greenhouse gases. The PRTR project system is used as basis for the NIR and for filling the CRF.

### 1.2.2 Overview of inventory planning


The Dutch Pollutant Release & Transfer Register (PRTR) has been in operation in the Netherlands since 1974. This system encompasses data collection, data processing and registering and reporting emission data for some 350 policy-relevant compounds and compound groups that are present in air, water and soil. The emission data is produced in an annual (project) cycle (MNP, 2006). This system is also the basis for the national greenhouse gas inventory. In April 2004, overall coordination of the PRTR was outsourced by VROM to the PBL. As per 1 January 2010, coordination has been assigned to RIVM.

The main objective of the PRTR is to produce an annual set of unequivocal emission data that is up-to-date, complete, transparent, comparable, consistent and accurate. In addition to RIVM, various external agencies contribute to the PRTR by performing calculations or submitting activity data. These include: CBS (Statistics Netherlands), PBL, TNO (Netherlands Organisation for Applied Scientific Research), NL Agency, Centre for Water Management, Deltares and several institutes related to the Wageningen University and Research Centre (WUR).

### Responsibility for reporting

The NIR part 1 is prepared by RIVM as part of the PRTR project. Most institutes involved in the PRTR also contribute to the NIR (including CBS and TNO).. In addition, NL Agency is involved in its role as NIE. NL Agency also prepares the NIR part 1 and takes care of integration and submission to the UNFCCC in its role as NIE. Submission to the UNFCCC only takes place after approval by VROM.





### 1.2.3 Overview of the inventory preparation and management under Article 7 of the Kyoto Protocol

Following the annotated outline, the supplementary information under Article 2 of the Kyoto Protocol is reported together in the NIR part 2. This information is prepared by NL Agency, using information from various other involved organisations, such as the NEa (the Netherlands Emissions Authority), the WUR and the Ministry (VROM).

## 1.3 Inventory preparation

### 1.3.1 GHG and KP-LULUCF inventory

The primary process of preparing the greenhouse gas inventory in the Netherlands is summarised in Figure 1.1. This process includes three major steps that are described in more detail in the following Sections.

For the KP-LULUCF inventory, the inventory preparation is combined with the work for reporting LULUCF by the unit Wettelijke Onderzoekstaken Natuur & Milieu, part of Wageningen UR. The project team LULUCF oversees data management, the preparation of the reports for land-use, land-use change and forestry and the QA/QC activities and decides on further improvements.

### 1.3.2 Data collection processing and storage

Various data suppliers provide the basic data for emission estimates. The most important data sources for greenhouse gas emissions include:

#### 1.3.2.1 Statistical data

Statistical data are provided under various (not specifically greenhouse-gas related) obligations and legal arrangements. These include national statistics from Statistics Netherlands (CBS) and a number of other sources of data on sinks, water and waste. The provision of relevant data for greenhouse gases is guaranteed through covenants and an Order in Decree, the latter of which is under preparation by VROM. For greenhouse gases, relevant agreements with respect to waste management are in place with CBS and NL Agency. An

agreement with the Ministry of Agriculture, Nature and Food Quality (LNV) and related institutions was established in 2005.

#### 1.3.2.2 Data from individual companies

Data from individual companies are provided in the form of annual environmental reports (MJVs). A large number of companies have a legal obligation to submit an MJV that includes – in addition to other pertinent information – emission data validated by the competent authorities (usually provincial and occasionally local authorities that also issue permits to these companies). A number of companies with large combustion plants are also required to report information under the BEES/A regulation. Some companies provide data voluntarily within the framework of environmental covenants. The data in these MJVs are used for verifying the calculated CO<sub>2</sub> emissions from energy statistics for industry, energy sector and refineries. If reports from major industries contain plant-specific information on activity data and emission factors of sufficient quality and transparency, this data is used in the calculation of CO<sub>2</sub> emission estimates for specific sectors.

The MJVs from individual companies provide essential information for calculating the emissions of substances other than CO<sub>2</sub>. The calculations of industrial process emissions of non-CO<sub>2</sub> greenhouse gases (e.g., N<sub>2</sub>O, HFC-23 and PFCs released as by-products) are mainly based on information from these MJVs, as are the calculated emissions from precursor gases (CO, NO<sub>x</sub>, NMVOC) and SO<sub>2</sub>. As reported in previous NIRs, only those MJVs with high-quality and transparent data are used as a basis for calculating total source emissions in the Netherlands.

#### 1.3.2.3 Additional greenhouse-gas-related data

Additional greenhouse gas related data are provided by other institutes and consultants that are specifically contracted to provide information on sectors not sufficiently covered by the above-mentioned data sources. For greenhouse gases, contracts and financial arrangements are made (by PBL/RIVM) with, for example, various agricultural institutes and TNO. In addition, NL Agency contracts out various tasks to consultants (such as collecting information on F-gas emissions from cooling and product use, on improvement actions).

During 2004, the Ministry of LNV also issued contracts to a number of agricultural institutes; these consisted of, in particular, contracts for developing a monitoring system and protocols for the LULUCF data set. Based on a written agreement between LNV and PBL/RIVM, these activities are also part of the PRTR.

#### 1.3.2.4 Processing and storage

Data processing and storage are coordinated by PBL/RIVM; these processes consist most notably of the elaboration of emission estimates and data preparation in the emissions data base and the CRF. The emission data are stored in a central database, thereby satisfying – in an efficient and effective manner – national and international criteria on emission reporting. This year, PBL automated the process to fill the CRF with emissions from the central database.

The actual emission calculations and estimates that are made using the input data are implemented in five task forces, each dealing with specific sectors:

- energy, industry and waste (combustion, process emissions, waste handling)
- agriculture (agriculture, sinks)
- consumers and services (non-industrial use of products)
- transport (including bunker emissions)
- water (less relevant for greenhouse gas emissions)

The task forces consist of experts from several institutes. In 2009, in addition to the PBL, these included TNO, CBS, Centre for Water Management, Deltares, FO-I (the Facilitating Organisation for Industry, which co-ordinates annual environmental reporting by companies), NL Agency (Waste Management division) and two agricultural research institutes: Alterra (sinks) and LEI. The task forces are responsible for assessing emission estimates based on the input data and emission factors provided. PBL commissioned TNO to assist compiling the CRF.

#### 1.3.3 Reporting, QA/QC, archiving and overall coordination

The NIR is prepared by PBL with input from the experts in the relevant PRTR task forces and from NL Agency. This step includes documentation and archiving. VROM formally approves the NIR before it is submitted; in some cases approval follows consultation with other ministries. NL Agency is responsible for coordinating QA/QC and responses to the EU and for providing additional information requested by the UNFCCC after the NIR and the CRF have been submitted. NL Agency is also responsible (in collaboration with RIVM) for coordinating the submission of supporting data to the UNFCCC review process.

For KP-LULUCF, consistency with the values submitted for the Convention was assured by using the same base data and calculation structure, and apply different calculations only where applicable based on the definitions and details of data for the KP LULUCF CRF Tables. The data and calculations were thus subject to the same QA/QC (Van den Wyngaert et al., 2009). The full time series since 1990 was calculated explicitly for re/afforestation and deforestation.

The calculated values were entered in the CRF reporting system at Alterra, and checked by the LULUCF sectoral

expert. They were then exported as an XML file and sent to the Dutch inventory, which imported the data in the CRF database for all sectors and again checked. Any strange or incomplete values were reported to the LULUCF sectoral expert, checked and if necessary corrected.

Verification with other international statistics was performed only with FAO. The area of forest is systematically lower for FAO. This may be due to a different methodology. For discussion on different outcomes of different estimates of forest cover in the Netherlands, see Nabuurs et al., 2005. The net increase in forest area in the FAO statistics is higher than reported for KP-LULUCF, and this may indicate that the 1990 estimate may be low in the FAO statistics. These values indicate a conservative estimate of the net forest increase in the Netherlands.

The mean C stock in Dutch forests (used as emission factor for deforestation under the KP) is slightly higher in the UNFCCC estimates than in the FAO estimates. Considering that different conversion factors were used, the estimates are close, while the difference has the tendency to increase. If this continues for the 2010 FAO estimate, this will be reason for investigation. These values indicate a conservative estimate of C emissions from deforestation.

No values from FAO are available on young forests. FAO statistics also provide no information on fires or disturbances for the Kyoto period, since at the national level these statistics are no longer kept. The same accounts for EFFIS, the European Forest Fires Information System.

## 1.4 Brief description of methodologies and data sources used

### 1.4.1 GHG inventory

#### 1.4.1.1 Methodologies

Table 1.1 provides an overview of the methods used to estimate greenhouse gas emissions. Monitoring protocols, documenting the methodologies and data sources used in the greenhouse gas inventory of the Netherlands as well as other key documents are listed in Annex 6. The protocols were elaborated, together with relevant experts and institutes, as part of the monitoring improvement program.

Explanation of notation keys used:

- Method applied: D, IPCC default; RA, reference approach; T, IPCC Tier; C, CORINAIR; CS, country-specific; M, model.
- Emission factor used: D, IPCC default; C, CORINAIR; CS, country-specific; PS, plant-specific; M, model.
- Other keys: NA, not applicable, NO, not occurring; NE, not estimated; IE, included elsewhere.

All key documents are electronically available in PDF format at [www.greenhousegases.nl](http://www.greenhousegases.nl). The monitoring protocols describe methodologies, data sources and QA/QC procedures for estimating greenhouse gas emissions in the Netherlands. The sector-specific chapters provide a brief description per key source of the methodologies applied for estimating the emissions.

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>		CH <sub>4</sub>		N <sub>2</sub> O		HFCs		PFCs		SF <sub>6</sub>	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
<b>1. Energy</b>	CS,D,T2,T3	CS,D,PS	CS,T1,T1b,T2,T3	CS,D,PS	CS,T1,T2	CS,D						
A. Fuel Combustion	CS,D,T2	CS,D	CS,T1,T2,T3	CS,D	CS,T1,T2	CS,D						
1. Energy Industries	T2	CS	T2	CS	T1,T2	CS,D						
2. Manufacturing Industries and Construction	T2	CS	T2	CS	T1,T2	CS,D						
3. Transport	CS,T2	CS	CS,T2,T3	CS,D	CS,T2	CS,D						
4. Other Sectors	T2	CS	T1,T2	CS,D	T1	D						
5. Other	D,T2	D	CS,T2	CS	CS,T2	CS						
B. Fugitive Emissions from Fuels	CS,T2,T3	CS,PS	T1b,T2,T3	CS,D,PS	NA	NA						
1. Solid Fuels	T2	CS	T1b	D	NA	NA						
2. Oil and Natural Gas	CS,T2,T3	CS,PS	T1b,T2,T3	CS,D,PS	NA	NA						
<b>2. Industrial Processes</b>	CS,T1,T1a,T1b,T2	CS,D,PS	CS,T1,T2	CS,D	CS,T2	CS,PS	T1,T2	CS,PS	CS,T2	PS	CS,T2	D,PS
A. Mineral Products	CS	CS,D,PS	NA	NA	NA	NA						
B. Chemical Industry	CS,T1,T1b	CS,D,PS	T1,T2	D	T2	PS	NA	NA	NA	NA	NA	NA
C. Metal Production	T1a,T2	CS	NA	NA	NA	NA	NA	NA	T2	PS	NA	NA
D. Other Production	T1b	CS										
E. Production of Halocarbons and SF <sub>6</sub>							T1,T2	PS	NA	NA	NA	NA
F. Consumption of Halocarbons and SF <sub>6</sub>							T2	CS	CS,T2	PS	CS,T2	D,PS
G. Other	CS,T1b	CS,D	CS	CS	CS	CS	NA	NA	NA	NA	NA	NA
<b>3. Solvent and Other Product Use</b>	CS	CS			CS	CS						
<b>4. Agriculture</b>			T1,T2	CS,D	T1,T1b,T2,T3	CS,D						
A. Enteric Fermentation			T1,T2	CS,D								
B. Manure Management			T2	CS	T2	D						
C. Rice Cultivation			NA	NA								
D. Agricultural Soils			NA	NA	T1,T1b,T2,T3	CS,D						
E. Prescribed Burning of Savannas			NA	NA	NA	NA						
F. Field Burning of Agricultural Residues			NA	NA	NA	NA						
G. Other			NA	NA	NA	NA						
<b>5. Land Use, Land-Use Change and Forestry</b>	CS,D	CS,D	NA	NA	NA	NA						
A. Forest Land	CS	CS	NA	NA	NA	NA						
B. Cropland			NA	NA	NA	NA						
C. Grassland			NA	NA	NA	NA						
D. Wetlands			NA	NA	NA	NA						
E. Settlements			NA	NA	NA	NA						
F. Other Land			NA	NA	NA	NA						
G. Other	D	D	NA	NA	NA	NA						
<b>6. Waste</b>	NA	NA	T2	CS	T2	CS,D						
A. Solid Waste Disposal on Land	NA	NA	T2	CS								
B. Waste-water Handling			T2	CS	T2	D						
C. Waste Incineration	NA	NA	NA	NA	NA	NA						
D. Other	NA	NA	T2	CS	T2	CS						
<b>7. Other (as specified in Summary 1.A)</b>	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

#### 1.4.1.2 Data sources

The monitoring protocols provide detailed information on activity data used for the inventory. In general, the following primary data sources supply the annual activity data used in the emission calculations:

- fossil fuel data: (1) national energy statistics from CBS (National Energy Statistics; Energy Monitor); (2) natural gas and diesel consumption in the agricultural sector (Agricultural Economics Institute, LEI)
- residential bio fuel data: (1) annual survey of residential woodstove and fireplace penetration from the Association for Comfortable Living (Vereniging Comfortabel Wonen); (2) a 1996 survey on wood consumption by owners of residential woodstoves and fireplaces from the Stove and Stack Association (Vereniging van Haard en Rookkanaal, VHR)
- transport statistics: monthly statistics for traffic and transportation
- industrial production statistics: (1) annual inventory reports from individual companies; (2) national statistics
- consumption of HFCs: annual reports from the accountancy firm PriceWaterhouseCoopers (only HFC data are used due to inconsistencies for PFCs and SF<sub>6</sub> with emissions reported elsewhere)
- consumption/emissions of PFCs and SF<sub>6</sub>: reported by individual firms
- anaesthetic gas: data provided by Hoekloos, the major supplier of this gas
- spray cans containing N<sub>2</sub>O: the Dutch Association of Aerosol Producers (Nederlandse Aerosol Vereniging, NAV)

- animal numbers: from the CBS/LEI agricultural database, plus data from the annual agricultural census
- manure production and handling: from the CBS/LEI national statistics
- fertiliser statistics: from the LEI agricultural statistics
- forest and wood statistics: (1) harvest data: FAO harvest statistics; (2) stem-volume, annual growth and fellings: Dirkse et al, (2003) (3) carbon balance: National Forestry Inventory data based on two inventories: HOSP (1988-1992) and MFV (2001-2005)
- land use and land use change: based on digitised and digital topographical maps of 1990 and 2004 (Kramer et al, 2009, in preparation)
- area of organic soils: De Vries (2004)
- soil maps: De Groot et al. (2005)
- waste production and handling: Working Group on Waste Registration (WAR), NL Agency and CBS
- CH<sub>4</sub> recovery from landfills: Association of Waste Handling Companies (VVAV).

Many recent statistics are available on the internet at CBS's statistical website [Statline](#) and in the CBS/PBL [environmental data compendium](#). However, it should be noted that the units and definitions used for domestic purposes on those websites occasionally differ from those used in this report (for instance: temperature corrected CO<sub>2</sub> emissions versus actual emissions in this report; in other cases, emissions are presented with or without the inclusion of organic CO<sub>2</sub> and with or without LULUCF sinks and sources).

## 1.4.2 KP-LULUCF inventory

### 1.4.2.1 Methodologies

The methods used to estimate data on sinks and sources as well as the units of land subject to Article 3.3 afforestation, reforestation and deforestation are additional to the methods used for LULUCF. The methodology of the Netherlands to assess the emission from LULUCF is based on a wall-to-wall approach for the estimation of area per category of land use. For the wall-to-wall map overlay approach were used harmonised and validated digital topographical maps of 1990 and 2004 (Kramer et al., 2009). The result was a national scale land use and land use change matrix.

To distinguish between mineral soils and peat soils, an overlay was made between two Basic Nature maps and the Dutch Soil Map (de Vries et al., 2003). The result is a map with national coverage that identifies for each pixel whether it was subject to RA or D between 1990 and 2004, and whether it is located on a mineral or on an organic soil.

### 1.4.2.2 Data sources

The changes in land use are based on comparing detailed maps that best represent land use in 1990 and 2004. Both datasets on land use were especially developed to support the temporal and spatial development in land use and especially designed to support policy in the field of nature conservation. Changes after 2004 have been obtained by linear extrapolation.

## 1.5 A brief description of the key categories

### 1.5.1 GHG inventory

The analysis of key sources is performed in accordance with the IPCC Good Practice Guidance (IPCC, 2001). To facilitate the identification of key sources, the contribution of source categories to emissions per gas are classified based on the IPCC potential key source list as presented in Table 7.1, Chapter 7 of the Good Practice Guidance.

A detailed description of the key source analysis is provided in Annex 1 of this report. Per sector, the key sources are also listed in the first Section of each of Chapters 3 to 8.

Compared to the key source analysis for the NIR 2009, one new key category has been identified:

- CH<sub>4</sub> emissions from Manure management, poultry (Tier 2 trend)

### 1.5.2 KP-LULUCF inventory

Deforestation is larger than the smallest key category in the Tier 1 key source analysis (excluding LULUCF). With 547 Gg CO<sub>2</sub> the annual contribution of re/afforestation under the KP is just below the smallest key category (Tier 1 analysis including LULUCF). Deforestation under the KP in 2008 causes emission of 780 Gg CO<sub>2</sub>, which is more than the smallest key category (Tier 1 analysis including LULUCF).

## 1.6 Information on the QA/QC plan

As one of the results of a comprehensive *inventory improvement program*, a National System fully in line with the Kyoto requirements was finalised and established at the end of 2005. As part of this system, an Act on the Monitoring of Greenhouse Gases also became effective in December 2005. This Act determines the establishment of the National System for monitoring of greenhouse gases and empowers the Minister for Housing, Spatial Planning and the Environment (VROM) to appoint an authority responsible for the National System and the National Inventory. The Act also determines that the National Inventory be based on methodologies and processes as laid down in the monitoring protocols. In a subsequent regulation the Minister has appointed NL Agency as NIE (National Inventory Entity, the single national entity under the Kyoto Protocol) and published a list of the protocols. Adjustments to the protocols will require official publication of the new protocols and announcement of publication in the official Government Gazette (Staatscourant).

As part of its National System, the Netherlands has developed and implemented a QA/QC program. This program is yearly assessed and updated, if needed. The key elements of the current program (SenterNovem, 2009) are briefly summarised in this chapter, notably those related to the current NIR.

### 1.6.1 QA/QC procedures for the CRF/NIR 2010

The Monitoring Protocols were elaborated and implemented in order to improve the transparency of the inventory (including methodologies, procedures, tasks, roles and responsibilities with regard to inventories of greenhouse gases). Transparent descriptions and procedures of these different aspects are described in the protocols for each gas and sector and in process descriptions for other relevant tasks in the National System. The protocols are assessed annually and updated if needed.

- Various QC issues
  - Inconsistencies in the key category analysis between CRF and NIR were analysed and removed. The key category analysis is updated in the NIR (Annex 1) as well as the CRF files.
  - The Expert Review Team (ERT) recommended providing more information in the NIR report and protocols, that was until now only included in background information. The Netherlands has updated the protocols; for various sectors this implies that more information is included in the protocols, as requested by the ERT.
  - The ERT recommended providing more specific information on sector specific QC activities. In 2009 and early 2010, a project was performed to re-assess and update both the information on uncertainties and on sector specific QC activities [Ecofys 2010/PBL/SenterNovem in print]. The results are used in the NIR 2010.
  - The Netherlands continues its efforts to include the correct notation keys in the CRF files.
- For the NIR 2010, changes were incorporated in and references were updated to the National System website ([www.greenhousegases.nl](http://www.greenhousegases.nl)), providing additional information on the protocols and relevant background documents;



- General QC checks were performed. To facilitate these general QC checks, a checklist was developed and implemented. A number of general QC checks have been introduced as part of the annual work plan of the PRTR and are also mentioned in the monitoring protocols. The QC checks included in the work plan aim at covering issues such as consistency, completeness and correctness of the CRF data. The general QC for the present inventory was largely performed in the institutes involved as integrated part of their PRTR work. The PRTR task forces fill in a standard-format database with emission data for 1990–2008. After a first check of the emission files by RIVM and TNO for completeness, the (corrected) data are available to the specific task force for checking consistency checks and trend analysis (comparability, accuracy). The task forces have access to information about the relevant emissions in the database. Several weeks before the dataset was fixed, a trend verification workshop was organised by PBL (December, 2009) See Box 1.1. The result of this workshop including actions for the taskforces to resolve the identified clarification issues are documented at PBL. Required changes to the database are then made by the taskforces.
- Quality Assurance for the current NIR includes the following activities:
  - A peer and public review on the basis of the draft NIR in January/February 2010. Results of this review are summarised in Chapter 10 and have been dealt with as far as possible in the present NIR.
  - In preparing this NIR, the results of former UNFCCC reviews, including the results of the recent reviews, including that of the NIR 2009 (in September 2009) have been taken into account wherever possible (see also Chapter 10.5 for an overview).
  - As part of the evaluation process of the previous cycle, internal audits were carried out by NL Agency on the use of the protocols and the implementation of QC checks. This year, the NIR process was given special attention. The audit resulted in some recommendations on transparency of the processes (e.g., improvement by drawing up manuals for key source analysis and data conversion from the central database to the CRF) and on tasks and responsibilities (e.g., capacity building, improvement of planning and communication).
- Up to and including 2008, PBL held ISO 9001/2000 certification. After 31 December 2008, PBL no longer applied for extension of this certificate, but uses its own quality management system, following the guidelines of the Dutch Institute for Quality Management (INK, a Dutch variety of the European Foundation for Quality Management (EFQM) Business Model). In practice, this did not have much impact on the quality checks and quality assurance within PBL. As part of this system, PBL will periodically contract consultants to assess the implementation of its quality system and the INK guidelines.
- *The annual activity program* of the PRTR (PBL, 2009). The work plan describes the tasks and responsibilities of the parties involved in the PRTR process, such as products, time schedules (planning) and emission estimation methods – including the monitoring protocols for the greenhouse gases – as well as those of the members of several task forces. The annual work plan also describes the general QC activities to be performed by the task forces before the annual database is fixed (see Section 1.6.2)
- The responsibility for the quality of data in *annual environmental reports* (MJVs) lies with the companies themselves, while validation of the data is the responsibility of the competent authorities. It is the responsibility of the institutes involved in the PRTR to judge whether or not to use the validated data of individual companies to assess the national total emissions. (CO<sub>2</sub> emissions, however, are based on energy statistics and standard emission factors, and only qualified specific emission factor from environmental reports are used).
- *Agreements/covenants* between PBL/RIVM and other institutes involved in the annual PRTR process. The general agreement is that by accepting the annual work plan, the institutes involved commit themselves to deliver capacity for the products specified in that work plan. The role and responsibility of each institute have been described (and agreed upon) within the framework of the PRTR work plan.
- *Specific procedures* that have been established to fulfil the QA/QC requirements as prescribed by the UNFCCC and Kyoto Protocol. General agreements on these procedures are described in the QA/QC program as part of the National System. The following specific procedures and agreements have been set out and described in the QA/QC plan and the annual PRTR work plan:
  - QC on data input and data processing, as part of the annual process towards trend analysis and fixation of the database following approval of the involved institutions.
  - Documentation of consistency, completeness and correctness of the CRF data (see also Section 1.6.2). Documentation is required for changes in the historical data set or in the emission trend that exceeds 5% at the sector level and 0.5% at the national total level.
  - Peer reviews of CRF and NIR by the NL Agency and institutions not fundamentally involved in the PRTR process.
  - Public review of the draft NIR: NL Agency organises every year a public review (by means of internet). Relevant comments are incorporated in the final NIR.

The QA/QC activities generally aim at a high-quality output of the emissions inventory and the National System; these are in line with international QA/QC requirements (IPCC Good Practice Guidance).

The QA/QC system should operate within the available means (capacity, finance). Within those boundaries, the main focal points of the QA/QC activities are:

- The *QA/QC programme* (SenterNovem, 2009) that has been developed and implemented as part of the National System. This program includes quality objectives for the National System, the QA/QC plan and a time schedule for implementation of the activities. It is updated annually as part of a yearly ‘evaluation and improvement cycle’ for the inventory and National System and held available for review.

- Audits: in the context of the annual work plan, it has been agreed that the involved institutions of the PRTR inform PBL concerning possible internal audits. Furthermore, NL Agency is assigned the task of organising audits, if needed, of relevant processes or organisational issues within the National System. In 2009, such an audit was performed for the NIR process (the previous Section).
- Archiving and documentation: internal procedures are agreed (amongst others in the PRTR annual activity program) for general data collection and the storage of fixed datasets in the PBL database, including the documentation/archiving of QC checks. The improved monitoring protocols have been documented and will be published on the website [www.greenhousegases.nl](http://www.greenhousegases.nl). To improve transparency, the implemented checklists for QC checks have been documented and archived. As part of the QA/QC plan the documentation and archiving system has been further upgraded. NL Agency (NIE) maintains the National System website and a central archive of relevant National System documents
- Each institution is responsible for QA/QC aspects related to reports based on the annually fixed database.
- *Evaluation and improvement*: those persons involved in the annual inventory tasks are invited once a year to evaluate the process. In this evaluation, the results of any internal and external review and evaluation are taken into account. The results are used for the annual update of the QA/QC program (including the improvement program) and the annual work plan. The (monitoring) improvement plan has been described in the previous sub-section;
- *Source-specific QC*: comparison of emissions with independent data sources was one of the study topics in the inventory improvement program. Because it did not seem possible to considerably reduce uncertainties through independent verification (measurements) – at least not on a national scale – this issue has received less priority. However, the theme is taken up in two projects. Following the UN review recommendations, NL Agency started a project together with Ecofys (Ecofys, 2010) and the PRTR project to re-assess and update the description of uncertainties and the sector specific QC activities. Furthermore, in the context of a longer term programme (including a large research program on climate change in the Netherlands, the National Research Program on Global Air Pollution and Climate Change (NOP-MLK) or by the Dutch Reduction Program on Other Greenhouse Gases (ROB)), the issue is also currently being studied.. To some extent (e.g., in the transport sector), comparisons can be made on the basis of independent data sets.
- In 2009, a quantitative assessment was made of the possible (in) consistencies in CO<sub>2</sub> emissions between data from ETS, NIR and National Energy Statistics. The figures that were analysed concerned about 40% of the CO<sub>2</sub> emissions in the Netherlands in 2008. The differences could reasonably be explained (e.g., different scope) within the given time available for this action [Zijlema, 2010].

### 1.6.2 Verification activities for the CRF/NIR 2010

Several weeks in advance of a trend analysis meeting, a snapshot from the database was made available by PBL in a

web-based application (Emission Explorer, EmEx) for checks by the institutes and experts involved (PRTR task forces). This allows the task forces to check for level errors and consistency in the algorithm/method used for calculations throughout the time series. The task forces perform checks such as for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from all sectors. The totals for the sectors are then compared with the previous year's data set. Where significant differences are found, the task forces evaluate the emission data in more detail. The results of these checks are then subject to discussion at the trend analysis workshop and subsequently documented.

Furthermore, the task forces were provided with the CRF Reporter software to check the time series of emissions per substance. The task forces examine these time series. During the trend analysis, the greenhouse gas emissions for all years between 1990 and 2008 were checked in two ways: (1) emissions from 1990 – 2008 should (with some exceptions) be identical to those reported last year; (2) the data for 2008 were compared with the trend development for each gas since 1990. Checks of outliers were carried out at a more detailed level for the sub-sources of all sector background tables:

- annual changes in emissions of all greenhouse gases
- annual changes in activity data
- annual changes in implied emission factors
- level values of implied emission factors

Exceptional trend changes and observed outliers are noted and discussed at the trend analysis workshop, resulting in an action list. Items on this list must either be processed within 2 weeks or be dealt with in next year's inventory.

The trend verification workshop held on 11 December 2009, showed the following results:

#### Issues per source category:

- Reallocation of CO<sub>2</sub> emissions from cat. 1.AA.4 a to 1.AA4 c based on new fuel consumption data from the agricultural sector.
- Because detailed data became available last year the historic emissions of F-gases changed
- Changes in emissions of CH<sub>4</sub> from cat. 4. Agriculture (whole time series) should be explained in Chapter 6
- Minor changes in activity data and emissions in most categories, caused by the new improved automated import procedure of data to the CRF Reporter, and should be explained in Chapter 10.

All above mentioned check are planned in the annual inventory preparation plan for 2010 (PBL, 2009). Furthermore, data checks (also for non greenhouse gases) are performed and a trend verification workshop is held. To facilitate the data checks and the trend verification workshop, three types of data sheets were prepared from the PRTR emission database:

- Based on the PRTR emission database, a table with a comparison of the emission years 2007 and 2008. In this table, differences >5% on sector level were marked for documenting trends;
- Based on the PRTR emission database, to check if no historical data have been accidentally changed, a table

CO <sub>2</sub>	±3%	HFCs	±50%
CH <sub>4</sub>	±25%	PFCs	±50%
N <sub>2</sub> O	±65%	SF <sub>6</sub>	±50%
Total greenhouse gases			±5%

with a comparison of the complete inventories of 2009 versus 2010;

- To check if no errors occur during transfer of data from the PRTR emission database to the CRF, a table with the comparison of data from both data sources.

The data checks are performed by the sector experts and others involved in preparing the emission database and the inventory. Communications (e-mail) between the participants in the data checks is centrally collected and analysed. This resulted in a checklist of actions to be taken. This checklist is used as input for the trend verification workshop and completed with the actions agreed in this workshop. Furthermore, in the trend verification workshop trends >5% on sector level are explained.

Completion of an action is reported in the checklist. Based on the completed checklist and the documentation of trends the dataset is formally agreed by the two most involved institutes: PBL and Statistics Netherlands (CBS). The acceptance of the dataset is, furthermore, a subject in the PRTR Executive body (WEM).

All documentation (e-mails, data sheets and checklist) are stored electronically on a server at PBL.

### 1.6.3 Treatment of confidentiality Issues

Some of the data used in the compilation of the inventory are confidential and cannot be published in print or electronic format. That is the reason the Netherlands has to use the notation key “C” in the CRF. Although this does impair the transparency of the inventory, all confidential data can be made available to the official review process of the UNFCCC.

## 1.7 Evaluating general uncertainty

The IPCC Tier 1 methodology for estimating uncertainty in annual emissions and trends has been applied to the list of possible key sources (see Annex 1) in order to obtain an estimate of the uncertainties in the annual emissions as well as in the trends. These uncertainty estimates have also been used for a first Tier 2 analysis to assess error propagation and to identify key sources as defined in the IPCC Good Practice Guidance (IPCC, 2001).

### 1.7.1 GHG inventory

The following information sources were used for estimating the uncertainty in activity data and emission factors (Olivier et al., 2009):

- estimates used for reporting uncertainty in greenhouse gas emissions in the Netherlands that were discussed at a national workshop in 1999 (Van Amstel et al., 2000a)
- default uncertainty estimates provided in the IPCC Good Practice Guidance report (IPCC, 2000)

- RIVM fact sheets on calculation methodology and data uncertainty (RIVM, 1999)
- other recent information on the quality of data (Boonekamp et al., 2001)
- a comparison with uncertainty ranges reported by other European countries have led to a number of improvements in (and increased underpinning of) the Netherlands’ assumptions for the present Tier 1 (Ramirez et al., 2006)

These data sources were supplemented with expert judgments from PBL and CBS emission experts (also for new key sources). This was followed by an estimation of the uncertainty in the emissions in 1990 and 2008 according to the IPCC Tier 1 methodology – for both the annual emissions and the emission trend for the Netherlands. All uncertainty figures should be interpreted as corresponding to a confidence interval of 2 standard deviations (2σ), or 95%. In cases where asymmetric uncertainty ranges were assumed, the largest percentage was used in the calculation.

The results of the uncertainty calculation according to the IPCC Tier 1 uncertainty approach are summarised in Annex 7 of this report. The Tier 1 calculation of annual uncertainty in CO<sub>2</sub> equivalent emissions results in an overall uncertainty of about 4% in 2008, based on calculated uncertainties of 2%, 16%, 60% and 36% for CO<sub>2</sub> (excluding LULUCF), CH<sub>4</sub>, N<sub>2</sub>O and F-gases, respectively. The uncertainty in CO<sub>2</sub> equivalent emissions including emissions from LULUCF is calculated to be 4% as well.

However, these figures do not include the correlation between source categories (e.g., cattle numbers for enteric fermentation and animal manure production) or a correction for not-reported sources. Therefore, the Tier 2 uncertainty of total annual emissions per compound and of the total will be somewhat higher; see Table 1.2 for the currently estimated values.

Table 1.3 shows the top ten sources (excluding LULUCF) contributing most to total annual uncertainty in 2008, after ranking the sources according to their calculated contribution to the uncertainty in total national emissions (using the column ‘Combined uncertainty as a percentage of total national emissions in 2008’ in Table A7.1).

Table A7.1 of Annex 7 summarises the estimate of the *trend uncertainty* 1990–2008 calculated according to the IPCC Tier 1 approach in the IPCC Good Practice Guidance (IPCC, 2001). The result is a trend uncertainty in the total CO<sub>2</sub> equivalent emissions (excluding LULUCF) for 1990–2008 (1995 for F-gases) of ±3% points. This means that the decrease in total CO<sub>2</sub> eq emissions between 1990 and 2008, which is calculated to be 3%, will be between 0% and 6%.

IPCC category	Category	Gas	Combined uncertainty as a percentage of total national emissions in 2008 <sup>a)</sup>
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3.1%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	1.4%
1A4a	Stationary combustion : Other sectors: Commercial/Institutional, gases	CO <sub>2</sub>	1.0%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	0.8%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	0.7%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	0.6%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	0.5%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	0.5%
4A!	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	N <sub>2</sub> O	0.5%
4B	Emissions from manure management	N <sub>2</sub> O	0.4%

a)calculated uncertainties, for ranking purposes not rounded off

IPCC cat.	Category	Gas	Uncertainty introduced into the trend in total national emissions
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	1.9%
1A4a	Stationary combustion : Other sectors: Commercial/Institutional, gases	CO <sub>2</sub>	1.3%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	1.1%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	0.7%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	0.6%
1A4b	Stationary combustion : Other sectors, Residential, gases	CO <sub>2</sub>	0.5%
2B2	Nitric Acid production	N <sub>2</sub> O	0.4%
1A4c	Stationary combustion: Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	0.4%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	0.3%
2F	Emissions from substitutes for ozone depleting substances	HFC	0.3%

Per individual gas, the *trend uncertainty* in total emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and the total group of F-gases has been calculated to be  $\pm 3\%$ ,  $\pm 9\%$ ,  $\pm 14\%$  and  $\pm 10\%$  points respectively. More details on the level and trend uncertainty assessment can be found in Annex 7. Table 1.4 shows the top ten sources (excluding LULUCF) contributing most to *trend uncertainty* (calculated) in the national total (using the column 'Uncertainty introduced into the trend in total national emissions' in Table A7.1).

Five of these key sources are included in both the list presented above and the list of the largest contributors to annual uncertainty.

The propagation of uncertainty in the emission calculations was assessed using the IPCC Tier 1 approach. In this method, uncertainty ranges are combined for all sectors or gases using the standard equations for error propagation. If sources are added, total error is the root of the sum of squares of the error in the underlying sources. Strictly speaking, this is only valid if the uncertainties meet the following conditions: (a) standard normal distribution ("Gaussian"); (b)  $2\sigma$  smaller than 60%; (c) independent (not-correlated) sector-to-sector and substance-to-substance. It is clear, however, for some sources that activity data or emission factors are correlated, which may change the overall uncertainty of the sum to an unknown extent. It is also known for some sources, that the uncertainty is not distributed normally; in particular, when uncertainties are very high (of an order of 100%), it is clear that the distribution will be positively skewed.

Even more important is the fact that although the uncertainty estimates have been based on the documented uncertainties mentioned above, uncertainty estimates are unavoidably – and ultimately – based on the judgment of the expert. On occasion, there is only limited reference to actual data for the Netherlands possible as support for these estimates. By focusing on the order of magnitude of the individual uncertainty estimates, it is expected that this data set provides a reasonable first assessment of the uncertainty of key source categories.

Furthermore, in 2006 a Tier 2 uncertainty assessment was carried out (Ramirez et al., 2006). This study used the same uncertainty assumption as the Tier 1 study but accounted for correlations and non-Gaussian distributions. Results reveal that the Tier 2 uncertainty in total Netherlands CO<sub>2</sub> equivalent emissions is in the same order of magnitude as that in the Tier 1 results, although a higher trend uncertainty is found (see Tables 1.5 and 1.6). Furthermore, the Tier 2 uncertainty for 1990 emissions is slightly higher (about 1.5%) than the uncertainty for the 2004 emissions. Finally, the resulting distribution for total Netherlands CO<sub>2</sub> equivalent emissions turns out to be clearly positively skewed.

As part of the above mentioned study, the expert judgments and assumptions made for uncertainty ranges in emission factors and activity data for the Netherlands have been compared to the uncertainty assumptions (and their underpinnings) used in Tier 2 studies carried out by other European countries, such as Finland, the United Kingdom,



Greenhouse gas	Tier 1 annual uncertainty <sup>1)</sup>	Tier 2 annual uncertainty <sup>2)</sup>
Carbon dioxide	1.9%	1.5%
Methane	18%	15%
Nitrous oxide	45%	42%
F-gases	27%	28%
Total	4.3%	3.9%

<sup>1)</sup> Calculated in NIR 2006. <sup>2)</sup> Source: Ramirez-Ramirez et al. (2006).

Effects of simplifying Tier 1 assumptions on the uncertainty in the emission trend for 1990–2004 (without LULUCF) Table 1.6

Greenhouse gas	Emission trend 1990-2004	Tier 1 trend uncertainty <sup>1)</sup>	Tier 2 trend uncertainty <sup>2)</sup>
Carbon dioxide	+13%	2.7%	2.1%
Methane	-32%	11%	15%
Nitrous oxide	-16%	15%	28%
F-gases	-75%	7.0%	9.1%
Total	+1.6%	3.2%	4.5%

Norway, Austria and Flanders (Belgium). The correlations that have been assumed in the various European Tier 2 studies have also been mapped and compared. The comparisons of assumed uncertainty ranges have already led to a number of improvements in (and increased underpinning of) the Netherlands' assumptions for the present Tier 1 approach. Although a straightforward comparison is somewhat blurred due to differences in the aggregation level at which the assumptions have been made, results show that for CO<sub>2</sub> the uncertainty estimates of the Netherlands are well within the range of European studies. For non-CO<sub>2</sub> gases, especially N<sub>2</sub>O from agriculture and soils, the Netherlands uses IPCC defaults which are on the high side compared to the assumptions used in some of the other European studies. This seems quite realistic in view of the state of knowledge on the processes that lead to N<sub>2</sub>O emission. Another finding is that correlations (covariance and dependencies in the emission calculation) seem somewhat under-addressed in most present-day European Tier 2 studies and may require more systematic attention in future Tier 2 studies.

In the assessments made above, only random errors have been estimated, assuming that the methodology used for the calculation does not include systematic errors. It is well known that, in practice, this may well be the case. Therefore, a more independent verification of the emission level and emission trends using, for example, comparisons with atmospheric concentration measurements is encouraged by the IPCC Good Practice Guidance. In the Netherlands, these approaches have been studied for several years, funded by the National Research Program on Global Air Pollution and Climate Change (NOP-MLK) or by the Dutch Reduction Program on Other Greenhouse Gases (ROB). The results of these studies can be found in Berdowski et al. (2001), Roemer and Tarasova (2002) and Roemer et al. (2003). In 2006, the research program '*Climate changes spatial planning*' started to strengthen knowledge on the relationship between greenhouse gas emissions and land-use and spatial planning.

### 1.7.2 KP-LULUCF inventory

The analysis combines uncertainty estimates of the forest statistics, land use and land use change data (topographical data) and the method used to calculate the yearly growth

in carbon increase and removals (Olivier et al., 2009). The uncertainty analysis is performed for Forests according to the Kyoto definition and is based on the same data and calculations as used for KP article 3.3 categories. Thus, the uncertainty for total net emissions from units of land under article 3.3 afforestation/reforestation are estimated at 63%, equal to the uncertainty in Land converted to Forest Land. Similarly, the uncertainty for total net emissions from units of land under article 3.3, deforestation is estimated at 66%, equal to the uncertainty in Land converted to Grassland (which includes for the sake of the uncertainty analysis all Forest land converted to any other type of land use).

## 1.8 General assessment of the completeness

### 1.8.1 GHG inventory

At present, the greenhouse gas emission inventory for the Netherlands includes all of the sources identified by the Revised IPCC Guidelines (IPCC, 1997). Except for a number of (very) minor sources Annex 5 presents the assessment of completeness and sources, potential sources and sinks for this submission of the NIR and the CRF.

### 1.8.2 KP-LULUCF inventory

As good data to relate carbon accumulation in litter and dead wood since the time of re/afforestation are lacking for The Netherlands, this carbon sink is conservatively estimated as zero. Carbon stock changes in mineral and organic soils are not reported. Research is ongoing for this topic and result is expected to be available prior to next submission. In the next submission, a recalculation will be made for the year 2008.

Forest fertilisation does not occur in the Netherlands and therefore fertilisation in re/afforested areas are reported as not occurring.

Nitrous oxide emissions associated with disturbance of soils when deforested areas are converted to cropland are currently not estimated. However, research is ongoing and results are expected to be included in one of the next submissions.

Greenhouse gas emissions (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) related to biomass burning are not estimated because biomass burning has not been monitored since 1996. Wildfire statistics indicate that forest fires rarely occurred in the two decades before 1996.

# Trends in greenhouse gas emissions


## 2.1 Emission trends for aggregated greenhouse gas emissions

Chapter 2 summarises the trends in greenhouse gas emissions during the period 1990–2008, by greenhouse gas and by sector. Detailed explanations of these trends are provided in

Chapters 3–8. In 2008, total direct greenhouse gas emissions (excluding emissions from Land Use, Land Use Change and Forestry, LULUCF) in the Netherlands are estimated at 206.9 Tg CO<sub>2</sub> eq. This is 3.0% lower than the 213.3 Tg CO<sub>2</sub> eq reported in the base year (1990; 1995 is the base year for fluorinated gases).

Greenhouse gas emissions (excl. LULUCF)

Figure 2.1






Figure 2.1 shows the trends and relative contributions of the different gases to the aggregated national greenhouse gas emissions. In the period 1990–2008, emissions of carbon dioxide (CO<sub>2</sub>) increased by 10% (excluding LULUCF), while emissions of non-CO<sub>2</sub> greenhouse gases decreased by 42% compared with base year emissions. Of the non CO<sub>2</sub> greenhouse gases, methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and fluorinated gases (F-gases) decreased by 33%, 42% and 71% respectively.

Emissions of LULUCF related sources decreased by 5.8% in 2008 compared to 2007. In 2008, total greenhouse gas emissions (including LULUCF) decreased by 0.1 Tg CO<sub>2</sub> eq compared to 2007 (209.4 Tg CO<sub>2</sub> eq in 2008).

## 2.2 Emission trends by gas


### 2.2.1 Carbon dioxide

Figure 2.2 presents the contribution of the most important sectors, as defined by the Intergovernmental Panel on Climate Change (IPCC) to the trend in total national CO<sub>2</sub> emissions (excluding LULUCF). In the period 1990–2008, the

national CO<sub>2</sub> emissions increased by 10% (from 159.3 to 175.7 Tg). The Energy sector is by far the largest contributor to CO<sub>2</sub> emissions in the Netherlands (96%), with the categories 1A1 “Energy industries” (37%) and 1A4 “Other sectors” (22%) as largest contributors in 2008.

The relatively high level of CO<sub>2</sub> emissions in 1996 is mainly explained by a very cold winter, which increased energy use for space heating in the residential sector. The resulting emissions are included in the category 1A4 “Other sectors”. The relatively low level of CO<sub>2</sub> emissions in the category 1A1 “Energy industries” in 1999 is explained by the marked increase in imported electricity and a shift from the use of coal to residual chemical gas and natural gas in 1999; the share of imported electricity almost doubled. However, this increased import of electricity led to only a temporary decrease in the CO<sub>2</sub> emissions. In the period 2000–2004, the pre-1999 annual increase in CO<sub>2</sub> emissions from this category – about 1–2% – was observed again. In 2007, import of electricity decreased.

In 2008, CO<sub>2</sub> emissions increased by 2% compared to 2007.



### 2.2.2 Methane

Figure 2.3 presents the contribution of the most important IPCC sectors to the trend in total CH<sub>4</sub> emissions. The national CH<sub>4</sub> emissions decreased by 33%, from 1.22 Tg in 1990 to 0.81 Tg in 2008 (25.5 to 17.1 Tg CO<sub>2</sub> eq). The Agriculture and Waste sector (54% and 30%) are the largest contributors in 2008.

Compared to 2007, national CH<sub>4</sub> emissions increased by 1% in 2008 (0.2 Tg CO<sub>2</sub> eq), due to the increase of CH<sub>4</sub> emissions mainly in the categories 1A, 4A and 4B.

### 2.2.3 Nitrous oxide

Figure 2.4 presents the contribution of the most important IPCC sectors to the trend in national total N<sub>2</sub>O emissions. The total national inventory of N<sub>2</sub>O emissions decreased by about 42%, from 65.2 Gg in 1990 to 37.9 Gg in 2008 (20.2 to 11.8 Tg CO<sub>2</sub> eq). The sector contributing the most to this decrease in N<sub>2</sub>O emissions is "Industrial Processes" (-85%).

Compared to 2007, total N<sub>2</sub>O emissions decreased by 24% in 2008 (-3.7 Tg CO<sub>2</sub> eq) due to the emission reduction measures in Nitric Acid production.


### 2.2.4 Fluorinated gases

Figure 2.5 shows the trend in F-gas emissions included in the national greenhouse gas inventory. The emission level of the total F-gases decreased by 71% between 1995 and 2008, from 8.3 Tg CO<sub>2</sub> eq in 1995 (base year for F-gases) to 2.4 Tg CO<sub>2</sub> eq in 2008. Emissions of hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) decreased by approximately 68% and 87% respectively during this same period, while sulphur hexafluoride (SF<sub>6</sub>) emissions decreased by 26%.


The aggregated emissions of F-gases decreased 0.2% between 2007 and 2008. HFC emissions showed an increase of 4%, PFC and SF<sub>6</sub> emissions decreased by 22% and 1% respectively.

### 2.2.5 Uncertainty in emissions specified by greenhouse gas

The uncertainty in the trend of CO<sub>2</sub> equivalent emissions of the six greenhouse gases together is estimated to be approximately 3% points, based on the IPCC Tier 1 Trend Uncertainty Assessment; see Section 1.7. Per individual gas, the *trend* uncertainty in total emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and the sum of the F-gases is estimated to be ±3%, ±9%, ±14% and ±10% points, respectively.



For all greenhouse gases together, the uncertainty estimate in *annual* emissions is  $\pm 4\%$  and for CO<sub>2</sub>  $\pm 2\%$ . The uncertainty estimates in *annual* emissions of CH<sub>4</sub> and N<sub>2</sub>O are  $\pm 25\%$  and  $\pm 60\%$  respectively, and for HFCs, PFCs and SF<sub>6</sub>,  $\pm 50\%$  (see Section 1.7).



### 2.3 Emission trends specified by source category

Figure 2.6 provides an overview of emission trends per IPCC sector in Tg CO<sub>2</sub> equivalents.

The IPCC Energy sector is by far the largest contributor to the total greenhouse gas emissions in the national inventory (contributing 74% in the base year and 82% in 2008). The relative share of the other sectors decreased correspondingly. The emission level of the Energy sector increased by approximately 12% in the period 1990–2008, and total greenhouse gas emissions from the Waste, Industrial Processes and Agriculture sectors decreased by 56%, 53%, and 18% respectively in 2008 compared to the base year.

Compared to 2007, greenhouse gas emissions in the Energy sector increased by about 4.3 Tg (mainly CO<sub>2</sub>) in 2008. The CO<sub>2</sub> emission from the combustion of fossil fuels in this category increased by approximately 4.5 Tg.



Trends in emissions by sub category are described in more detail in Chapters 3–8.

#### 2.3.1 Uncertainty in emissions by sector

The uncertainty estimates in annual CO<sub>2</sub> equivalent emissions of IPCC sectors Energy [1], Industry [2], Solvents and product use [3], Agriculture [4], and Waste [6] are about ±2%, ±10%, ±27%, ±40% and ±30% respectively; for sector 5 LULUCF it is ±100%. The uncertainty in the trend of CO<sub>2</sub> equivalent emissions per sector is calculated for sector 1 Energy at ±3% points in the 12% increase, for sector 2 Industry at ±5% points in the 56% decrease, for sector 4 Agriculture at ±15% points in the 17% decrease and for sector 6 Waste at ±8% points in the 56% decrease.

#### 2.4 Emission trends for indirect greenhouse gases and SO<sub>2</sub>

Figure 2.7 shows the trends in total emissions of carbon monoxide (CO), nitrogen oxides (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO<sub>2</sub>). Compared to 1990, CO and NMVOC emissions in 2008 were reduced by 42% and 64% respectively. For SO<sub>2</sub> this was as much as 75%, and for NO<sub>x</sub>, 2008 emissions are 50% lower than





the 1990 level. With the exception of NMVOC, most of the emissions stem from fuel combustion.

Because of the problems identified with annual environmental reporting (see Section 1.3.2.), emissions of CO from industrial sources are not verified. However, experts have suggested that possible errors will have a minor effect on total emission levels. Due to lack of data, the time series for 1991–1994 and 1996–1999 was interpolated between 1990 and 1995.

In contrast to direct greenhouse gases, calculations of emissions of precursors from road transport are not based on fuel sales according to the national energy statistics but are directly related to transport statistics on a vehicle-kilometre basis. To some extent, this is different from the IPCC approach (see Section 3.2.8.4.).

Uncertainty in the emission factors for NO<sub>x</sub>, CO and NMVOC from fuel combustion is estimated to be in the range of 10–50%. The uncertainty in the emission factors of SO<sub>2</sub> from fuel combustion (basically the sulphur content of the fuels) is estimated to be 5%. For most compounds, the uncertainty in the activity data is relatively small compared to the uncertainty in the emission factors. Therefore, the uncertainty in the overall total of sources included in the inventory is estimated to be in the order of 25% for CO, 15% for NO<sub>x</sub>, 5% for SO<sub>2</sub>, and approximately 25% for NMVOC (TNO, 2004).



# Energy [CRF Sector 1]

# 3

## ■ Major changes in the Energy sector compared to the National Inventory Report 2009

*Emissions:* Compared to the previous submission the GHG emissions increased by 2%

*Key sources:* No changes compared to previous submission.

*Methodologies:* No changes in methodologies compared to last submission

## 3.1 Overview of sector

### 3.1.1 The Dutch Energy System

#### Energy Supply and Energy Demand

As for most developed countries, the energy system in the Netherlands is largely driven by the combustion of fossil fuels (Figure 3.1). In 2008, natural gas is supplying about 43.6% of the total primary fuels used in the Netherlands, followed by liquid fuels (38.2%) and solid fossil fuels (10.1%). The contribution of non-fossil fuels, including renewables and waste streams is rather limited.

Part of the supply of energy is not used for energy purposes. It is either used as feed stocks in (petro) chemical or fertiliser industry (19.8%) or lost as waste heat in cooling towers and cooling water in power plants (14.6%).

Emissions from fuel combustion are consistent with the national energy statistics. The time-series of the energy statistics is not fully consistent at the detailed sector and detailed fuel-type levels for the years 1991 to 1994. This inconsistency is caused by revisions in the economic classification scheme implemented in 1993, a change from the “special trade” to “general trade” system to define the domestic use of oil products, some error corrections and the elimination of statistical differences. These changes were incorporated into the data sets for 1990, 1995 and subsequent years, thus creating the existing inconsistency with the 1991–1994 dataset. For the base year 1990, CBS has re-assessed the original statistics and made them compatible with the ‘new’ 1993 classification system and ECN (Energy Research Centre of the Netherlands) was commissioned to re-allocate the statistics of 1991–1994 at a higher level of detail (for both fuels and sectors). This is also visible in Figure 3.1, where fuel use is only shown as a total value.

#### Trends in fossil fuel use and fuel mix

Natural gas represents a very large share of the national energy consumption in all non-transport sectors: power generation, industry and other sectors (mainly for space heating). Oil products are primarily used in the transport

sector, refineries and in the petrochemical industry, while the use of coal is limited to power generation and steel production.

Although the combustion of fossil waste (reported under Other Fuels) has tripled since 1990, its share in total fossil fuel use is still only 1% at the present time. In the 1990–2008 period, total fossil fuel combustion increased by 12%, due to a 15% increase in gas consumption, while liquid fuel use increased by 7%. At the same time, the combustion of solid fuels decreased by 12%.

Total fossil fuel consumption for combustion increased by about 2% between 2007 and 2008, mainly due to a 5% increase in gas consumption, but less solid and liquid fuels were used.

### 3.1.2 GHG Emissions from the energy sector


During combustion, carbon and hydrogen from fossil fuels are converted mainly into carbon dioxide (CO<sub>2</sub>) and water (H<sub>2</sub>O), releasing the chemical energy in the fuel as heat. This heat is generally either used directly or used (with some conversion losses) to produce mechanical energy, often to generate electricity or for transportation.

The energy sector is the most important sector in the Dutch greenhouse gas emission inventory, and contributes approximately 96% of CO<sub>2</sub> emissions in the country. The contribution of the energy sector to total greenhouse gas emissions in the country increases from 73% in 1990 to 83% in 2008. The greenhouse gas emissions from this sector are for over 98% in the form of CO<sub>2</sub> (see the previous chapter, Figure 2.2).

The energy sector includes:

- exploration and exploitation of primary energy sources,
- conversion of primary energy sources into more useable energy forms in refineries and power plants
- transmission and distribution of fuels
- final use of fuels in stationary and mobile applications.

Emissions arise from these activities by combustion and as fugitive emissions, or escape without combustion. Emissions



from the energy sector are to be reported in the source category split as shown in Figure 3.2.

#### Overview of shares and trends in emissions

Table 3.1 and Figure 3.2 show the contribution of the source categories in the Energy sector to the total national greenhouse gas inventory. About 47% of CO<sub>2</sub> emissions from fuel combustion stems from the combustion of natural gas, 18% from solid fuels (coal) and 34% from liquid fuels. CH<sub>4</sub> and N<sub>2</sub>O emissions from fuel combustion contribute 1.4% to the total emissions from this sector.

#### Key sources

Table 3.1 presents the key categories in the Energy sector specified by both level and trend (see also Annex 1). The key categories in 1A1, 1A2, 1A3 and 1A4 are based on aggregated emissions by fuel type and category, which is in line with the IPCC Good Practice Guidance (see Table 7.1 in IPCC 2001). Since CO<sub>2</sub> emissions have the largest share in the total of national greenhouse gas emissions, it is not surprising that – with the exception of inland aviation and railways – almost

all CO<sub>2</sub> sources are identified as key category. The total CH<sub>4</sub> emissions from stationary combustion sources together are also identified as a key category.

No changes are found compared to the key source analysis for the NIR 2009.

### 3.2 Fuel Combustion [1A]

#### 3.2.1 Comparison of the sectoral approach with the reference approach

Emissions from fuel combustion are generally estimated by multiplying fuel quantities combusted at specific energy processes with fuel and, in case of non-CO<sub>2</sub> greenhouse gases, source category dependent emission factors. This Sectoral Approach (SA) is based on fuel demand statistics. The IPCC guidance requires –as a quality control activity–also estimating CO<sub>2</sub> emissions from fuel combustion on the basis of a national carbon balance, derived from the fuel supply statistics. This is the Reference Approach (RA). In Annex


Sector/category	Gas	Key	Emissions in base year	Emissions in 2007	Emissions in 2008	Change 2008 - 2007	Contribution to total in 2008 (%)		
			Level, Trend	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	By sector	Of total gas
1 Energy	CO <sub>2</sub>			151.1	165.0	168.8	3.8	98	96
	CH <sub>4</sub>			2.4	2.1	2.5	0.4	1.4	15
	N <sub>2</sub> O			0.5	0.7	0.8	0.0	0.4	6
	All			154.0	167.8	172.1	4.3	100	83
1A Fuel combustion	CO <sub>2</sub>			149.9	163.4	167.5	4.1	97	95
	CH <sub>4</sub>			0.7	1.3	1.7	0.4	1.0	10
	N <sub>2</sub> O			0.5		0.8	0.0	0.4	6
	All			151.2	165.4	169.9	4.5	99	82
1A Emissions from stationary combustion (excl. 1A3)	CH <sub>4</sub>	L,T		0.6	1.2	1.6	0.4	0.9	10
1A1 Energy Industries	CO <sub>2</sub>			52.5	65.1	65.2	0.1	38	37
1A1a. Public Electricity and Heat Production	CO <sub>2</sub>			39.9	52.7	52.4	-0.3	30	30
1A1a liquids	CO <sub>2</sub>	L1,T1		0.2	0.7	0.8	0.1	0.5	0.5
1A1a solids	CO <sub>2</sub>	L,T1		25.8	26.1	25.4	-0.7	15	14
1A1a gas	CO <sub>2</sub>	L,T		13.3	23.7	23.9	0.3	14	14
1A1a other fuels: waste incineration	CO <sub>2</sub>	L1,T		0.6	2.2	2.2	0.1	1	1
1A1b. Petroleum refining	CO <sub>2</sub>			11.0	10.6	10.9	0.3	6	6
1A1b liquids	CO <sub>2</sub>	L,T1		10.0	8.0	8.2	0.2	5	5
1A1b gases	CO <sub>2</sub>	L1,T1		1.0	2.6	2.7	0.1	2	2
1A1c Manufacture of Solid Fuels and Other Energy Industries	CO <sub>2</sub>			1.5	1.8	1.9	0.1	1	1
1A1c gases	CO <sub>2</sub>	L,T1		1.5	1.8	1.9	0.1	1	1
1A2 Manufacturing industries and construction	CO <sub>2</sub>			32.7	27.7	27.5	-0.3	16	16
1A2 liquids	CO <sub>2</sub>	L,T1		8.6	9.1	9.0	-0.1	5	5
1A2 solids	CO <sub>2</sub>	L		5.0	4.5	4.7	0.1	3	3
1A2 gases	CO <sub>2</sub>	L,T		19.0	14.1	13.8	-0.3	8	8
1A2a. Iron and steel	CO <sub>2</sub>			4.0	4.5	4.8	0.3	3	3
1A2b. Non-Ferrous Metals	CO <sub>2</sub>			0.2	0.3	0.2	0.0	0.1	0.1
1A2c. Chemicals	CO <sub>2</sub>			17.1	12.8	12.5	-0.4	7	7
1A2d. Pup, Paper and Print	CO <sub>2</sub>			1.7	1.4	1.2	-0.2	0.7	0.7
1A2e. Food Processing, Beverages and Tobacco	CO <sub>2</sub>			4.1	3.8	3.6	-0.2	2	2
1A2f. Other	CO <sub>2</sub>			5.5	4.9	5.2	0.3	3	3
1A3. Transport	CO <sub>2</sub>			26.0	35.2	35.5	0.3	21	20
	N <sub>2</sub> O			0.3	0.4	0.4	0.0	0.3	4
	All			26.4	35.7	36.0	0.3	21	17
1A3a. Civil aviation	CO <sub>2</sub>			0.04	0.04	0.04	0.0	0.0	0.0
1A3b. Road	CO <sub>2</sub>			25.5	34.5	34.7	0.3	20	20
1A3b gasoline	CO <sub>2</sub>	L,T1		10.9	13.0	12.9	-0.1	7	7
1A3b diesel oil	CO <sub>2</sub>	L,T		11.8	20.5	20.8	0.3	12	12
1A3b LPG	CO <sub>2</sub>	L1,T		2.7	1.0	1.0	0.0	0.6	0.6
1A3b. Road	N <sub>2</sub> O			0.3	0.4	0.4	0.0	0.3	4
1A3c. Railways	CO <sub>2</sub>			0.1	0.1	0.1	0.0	0.1	0.1
1A3d. Navigation	CO <sub>2</sub>	L1		0.4	0.6	0.6	0.0	0.4	0.4
1A4. Other sectors	CO <sub>2</sub>			38.2	35.0	38.8	3.8	23	22
	CH <sub>4</sub>			0.4	1.0	1.4	0.4	0.8	8.4
	All			38.7	36.0	40.2	3.8	23	19
1A4 liquids (excl. From 1A4c)	CO <sub>2</sub>	T		1.5	0.4	0.4	-0.1	0.2	0.2
1A4a. Commercial/Institutional	CO <sub>2</sub>			8.5	10.1	11.1	0.9	6	6
1A4a gases	CO <sub>2</sub>	L,T		7.6	9.9	10.9	1.1	6	6
1A4b. Residential	CO <sub>2</sub>	L,T1		19.5	16	18	1.9	10	10
	CH <sub>4</sub>			0.4	0.3	0.3	0.0	0.2	2
1A4b gases	CO <sub>2</sub>			18.7	15.7	17.6	1.9	10	10
1A4c. Agriculture/Forestry/Fisheries	CO <sub>2</sub>	L,T1		10.2	8.9	9.9	1.0	6	6
1A4c liquids	CO <sub>2</sub>	L,T1		2.9	2.2	2.4	0.2	1	1
1A4c gases	CO <sub>2</sub>	L,T1		7.3	6.7	7.5	0.8	4	4
1A5 Other	CO <sub>2</sub>			0.6	0.3	0.4	0.1	0.2	0.2
1B Fugitive emissions from fuels	CO <sub>2</sub>			1.2	1.6	1.3	-0.2	0.8	0.8

Sector/category	Gas	Key	Emissions in base year	Emissions in 2007	Emissions in 2008	Change 2008 - 2007	Contribution to total in 2008 (%)		
1B Fugitive emissions from fuels	CH <sub>4</sub>		1.7	0.8	0.8	0.0	0.5	5	0.4
	All		2.8	2.4	2.2	-0.2	1.3		1.0
1B1. Solid fuels transformation: coke production	CO <sub>2</sub>		0.4	0.4	0.4	0.0	0.2	0.2	0.2
1B2. venting/flaring	CO <sub>2</sub>	T	0.8	0.1	0.1	0.0	0.0	0.0	0.0
1B2. venting/flaring	CH <sub>4</sub>	T	1.3	0.4	0.4	0.0	0.2	2	0.2
National Total GHG emissions (excl. CO <sub>2</sub> LULUCF)	CO <sub>2</sub>		159.3	172.3	175.7	3.4		100	85
	CH <sub>4</sub>		25.5	16.8	17.1	0.2		100	8
	N <sub>2</sub> O		20.2	15.4	11.8	-3.7		100	6
	All		213.3	206.9	206.9	0.0			100

Note: Key sources in the 1A1, 1A2, and 1A4 categories are based on aggregated emissions of CO<sub>2</sub> by fuel type.

## 1. Energy Sector

Figure 3.2



4, a detailed comparison of the sectoral approach with the reference approach is shown.

### Energy Supply balance

The energy supply balance for the Netherlands in 1990 and 2008 is shown in Table 3.2 at a relatively high aggregation level. The Netherlands produces large amounts of natural

gas, both onshore (Groningen gas) and offshore; 74% of the gas produced is exported. Natural gas represents a very large share of the national energy supply.

Table 3.2 Energy Supply Balance for the Netherlands (PJ NCV/year)

Year	Role	Indicator Name	Solid fuels	Crude oil and petroleum	Gas
1990	Supply	Primary production	0	171	2,301
		Total imports	491	5,367	85
		Stock change	-22	2	0
		Total exports	-101	-4,076	-1,081
		Bunkers	0	-500	0
		Gross inland consumption	-368	-964	-1,305
	Demand	Final non-energy consumption	-11	-328	-101
2008	Supply	Primary production	0	93	2,511
		Total imports	562	7,564	794
		Stock change	-21	-61	1
		Total exports	-203	-5,511	-1,852
		Bunkers	0	-811	0
		Gross inland consumption	-338	-1,274	-1,454
	Demand	Final non-energy consumption	-13	-597	-82

With carbon contents of each specific fuel, a national carbon balance can be derived from the energy supply balance and from this the national CO<sub>2</sub> emissions can be estimated by determining how much of this carbon is oxidised in any process within the country. To allow this, international bunkers are to be considered as “exports” and subtracted from the gross national consumption.

### 3.2.2 International bunker fuels

The Rotterdam area has four large refineries, producing relatively large quantities of heavy fuel oils. An important fraction of these heavy fuel oils is sold as international bunkers. In addition, most marine fuel oil produced in Russia is transported to Rotterdam where it is sold on the market. Combined, this makes Rotterdam the world’s largest supplier of marine bunker oils. The quantities of this bunker fuel are presented in Figure 3.3.

The Dutch refineries also produce considerable amounts of aviation fuels delivered to air carriers at the airports of the countries. In addition, Schiphol Airport is Western Europe’s largest supplier of aviation bunker fuels (jet-fuel). Given the small size of the country, almost all of this aviation fuels is used in international aviation. Figure 3.3 presents the time series of the fuel quantities exported to marine and aviation bunkers.

### 3.2.3 Feed stocks and non-energy use of fuels

Table 3.2 shows that in 2008, 47% of the gross national consumption of petroleum products is used in non-energy applications, mainly as feedstock (naphta) in the petrochemical industry and in products in many applications (bitumen, lubricants, etc.). Also a fraction of the gross national consumption of natural gas (6%, mainly in the ammonia production) and coal (4%, mainly in the iron and steel production) is used for non-energy applications and hence not directly oxidised. In many cases, these products will finally be oxidised in waste incinerators or during use (lubricants in two stroke engines). In the Reference Approach these product flows are excluded from the calculation of CO<sub>2</sub> emissions.

### 3.2.4 CO<sub>2</sub> capture from flue gases and subsequent CO<sub>2</sub> storage, if applicable

Not applicable yet.

### 3.2.5 Country-specific issues

See above.


### 3.2.6 Energy Industries [1A1]

#### 3.2.6.1 Source category description

“Energy Industries” are the main source category contributing to the Energy sector, accounting for 38% of the greenhouse gas emissions from this sector in 2008. In this sector, three source categories are included: Public Electricity and Heat Production [1A1a], Petroleum Refining [1A1b] and Manufacture of Solid Fuels and Other Energy Industries [1A1c]. Within these categories, natural gas and coal combustion by public electricity and heat production and oil combustion by petroleum refining are the dominating key sources. However, other key sources are liquid fuels and other fuels (waste) in public electricity and heat production, and natural gas combustion in petroleum refining and in manufacturing of solid fuels and other energy industries. CH<sub>4</sub> and N<sub>2</sub>O emissions from 1A1 “Energy Industries” contribute relatively little to the total national inventory of greenhouse gas emissions. CH<sub>4</sub> from stationary combustion is a key source, due to an increase of the CH<sub>4</sub> emission factor from small CHP plants. N<sub>2</sub>O emissions from “Energy Industries” are not identified as a key source (see Table 3.1).

In 2008, CO<sub>2</sub> emissions from category 1A1 “Energy Industries” contributed 25% to the total national greenhouse gas emission inventory (excluding LULUCF), while CH<sub>4</sub> and N<sub>2</sub>O emissions from this same category contributed relatively little to the total national greenhouse gas emissions. The share contributed by 1A1 “Energy Industries” to the total greenhouse gas emissions from the Energy sector increased from 34% in 1990 to 38% in 2008 (see Figure 3.2), partly due to a change in ownership of CHP plants (joint ventures, which are allocated to this source category).

Between 1990 and 2008, total CO<sub>2</sub> emissions from 1A1 “Energy Industries” increased by 24% (see Figure 3.4). In 2008, CO<sub>2</sub>



emissions from 1A1 “Energy Industries” remained stable compared to the emission in 2007.

#### Public Electricity and Heat Production [1A1a]

The Dutch electricity sector has a few notable features: it has a large share of coal-fired power stations and a large fraction of gas-fired cogeneration plants, with many of the latter being operated as joint ventures with industries. Compared to other countries in the EU, nuclear energy and renewable energy provide very little of the total primary energy supply in the Netherlands. The two main renewable energy sources are biomass and wind.

This source category also includes all emissions from large-scale waste incineration, since all incineration facilities also produce heat and/or electricity and hence, the waste incinerated in these installations is regarded as a fuel. In addition, a large fraction of the blast furnace gas and a significant part of coke oven gas produced by the one iron and steel plant in the Netherlands is combusted in the public electricity sector.

In 2008, 1A1a “Public Electricity and Heat Production” was the largest source category within the 1A1 Energy industries, accounting for 80% of the total greenhouse gas emissions from this category (see Figure 3.4 and Table 3.1). CO<sub>2</sub> emissions from waste incineration of fossil carbon represent 4% of the total greenhouse gas emissions in 1A1a Public electricity and heat production. In 2008, the emissions of CO<sub>2</sub> from the combustion of fossil fuels in this source category decreased by 1%.

Between 1990 and 2008, total CO<sub>2</sub> emissions from “Public Electricity and Heat Production” increased by 31%. The increasing trend in electric power production corresponds to considerably increased CO<sub>2</sub> emissions from fossil fuel combustion by power plants, which are partly compensated for by a shift from coal to natural gas and the increased efficiency of power plants.

The CO<sub>2</sub> emission level from waste incineration of fossil carbon increased from 0.6 Tg CO<sub>2</sub> in 1990 to 2.2 Tg CO<sub>2</sub> in 2008



due to the increasing amounts of municipal waste that are combusted instead of being deposited in landfills, which is the result of environmental policy to reduce waste disposal in landfills (see Chapter 8). The increasing level of CO<sub>2</sub> emissions in this period is partly compensated by a shift from the use of coal to natural gas and the increased efficiency of power plants.

The increase in the CO<sub>2</sub> emission factor for “other fuels” since 2004 is due to the increase in the share of plastics (which have a high carbon fraction) in the combustible waste (see Table 8.6 on the composition of incinerated waste). The decrease in 2006 and 2007 in the implied emission factor for CO<sub>2</sub> from biomass is due to the increase of the share of pure biomass (co-combusted with coal-firing) as opposed to the organic carbon in waste combustion with energy recovery. For the former type a lower emission factor is applied than for the latter.

Between 1990 and 1998, changed ownership relations of plants (joint ventures) caused a shift of cogeneration plants from category 1A2 “Manufacturing Industries” to “Public Electricity and Heat Production”. Half of the almost 30% increase in natural gas combustion that occurred between 1990 and 1998 is largely explained by cogeneration plants and a few large chemical waste gas-fired steam boilers being shifted from “Manufacturing Industries” to the “Public Electricity and Heat Production” due to changed ownership (joint ventures). The corresponding CO<sub>2</sub> emissions allocated to the Energy sector increased from virtually zero in 1990 to 8.5 Tg in 1998 and 9.1 Tg in 2005. The same criterion applies for emissions from waste incineration, which are included in this category since they all are subject to heat or electricity recovery, albeit this is not their main activity. Most of the combustion of biogas recovered at landfill sites is in CHP operated by utilities; therefore, it is allocated in this category.

A remarkable drop is shown in the emissions from 1A1a ‘Electricity and heat production’ in 1999 (–6% compared to 1998), which is, however, associated to the increasing emission trend in the 1990–1998 period and 2000 and





thereafter. In fact, electricity consumption in the Netherlands was 2% higher in 1999 than in 1998. The relatively low emissions for 1999 are explained by the higher share of imported electricity in domestic electricity consumption in that year, which was almost double that in 1998 (10% in 1998 versus 20% in 1999), and to a relatively large shift from coal to chemical waste gas and natural gas in 1999. The high import of electricity corresponds to approximately 4 Tg CO<sub>2</sub>, while the shift from coal to natural gas and oil corresponds to approximately 1 Tg CO<sub>2</sub> in 1999. The net import of electricity decreased again in 2001, and this was compensated for by an increased production of electricity from gas and coal combustion in the public electricity sector. In 2004, CO<sub>2</sub> emissions increased by 3% as a direct result of the start-up in 2004 of a large new gas-fired 790 MW<sub>e</sub> cogeneration plant, and a 2% decrease in coal combustion.

The strong increase in liquid fuel use in 1994 and 1995, with a sharp increase in 1995, is due to chemical waste gas being used in joint venture electricity and heat production facilities. This also explains the somewhat lower IEF for CO<sub>2</sub> from liquids since 1995.

#### Petroleum Refining [1A1b]

The Rotterdam harbour area houses four major refineries (a fifth one is located at Vlissingen) which export about 50% of their products to the European market. Consequently, the Dutch petrochemical industry is relatively large.

The share of 1A1b “Petroleum Refining” in total greenhouse gas emissions from the category 1A1 “Energy Industries” is estimated to be 21% in 1990 and 16% in 2008. However, the combustion emissions from this category should be viewed in relation to the fugitive emissions reported under category 1B2. Between 1990 and 2008 total CO<sub>2</sub> emissions from the refineries (including fugitive CO<sub>2</sub> emissions from hydrogen production reported in 1B2a-iv Refining) fluctuated between 11 and 12 Tg.

For 1A1b “Petroleum Refining” the calculation of emissions from fuel combustion is based on the sectoral energy statistics, using the fuel consumption for energetic purposes as activity data (including the consumption of residual refinery gases). Although the same method is still used, the quality of the data used to calculate and report CO<sub>2</sub> emissions is now improved by incorporating the CO<sub>2</sub> emissions reported by the individual refineries for 2002 onwards. Since 1998, one refinery has operated the SGHP unit, supplying all the hydrogen for a large-scale hydrocracker. When producing hydrogen, CO<sub>2</sub> is also produced as a co-product from the chemical processes (CO<sub>2</sub> removal and a two stage CO shift reaction). Refinery data specifying these fugitive CO<sub>2</sub> emissions are available and used for 2002 onwards and are reported in the category 1B2. The fuel used to provide the carbon for this non-combustion process is subtracted from the fuel consumption used to calculate the combustion emissions reported in this category.

The use of plant-specific emission factors for refinery gas for 2002 onwards – arithmetically resulting from the reported CO<sub>2</sub> emissions and combustion emissions as calculated using the default data – also causes changes in the implied emission

factor for CO<sub>2</sub> for total liquid fuel compared to the years prior to 2002 (emission factor for refinery gas is adjusted to get exact correspondence between the total calculated CO<sub>2</sub> emissions and the total CO<sub>2</sub> emissions officially reported by the refineries). Besides this non-energy/feedstock use of fuel for hydrogen production, for years prior to 2002 the energy and carbon balance between the oil products produced does not match the total crude oil input and of fuel used for combustion. The conclusion drawn, therefore, is that not all residual refinery gases and other residual fuels are accounted for in the national energy statistics. The carbon difference is always a positive figure. As such, it is assumed that part of the residual refinery gases and other residual fuels are all combusted (or incinerated by flaring) but not monitored/ reported by the industry are thus unaccounted for. The CO<sub>2</sub> emissions from this varying fuel consumption are included in the fuel type ‘liquids’. This represents approximately 10% (5–20%) of the total fuel consumption accounted for in the statistics. For 1998–2001, the unspecified CO<sub>2</sub> process emissions from the hydrogen plant are also included.

The inter-annual variation in the IEFs for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from liquid fuels is explained both by the high and variable shares (between 40% and 55%) of refinery gas in total liquid fuel. This has a relatively low default emission factor compared to most other oil products and has variable emission factors for the years 2002 onward, and by the variable addition of ‘unaccounted for’ liquids that is only used for estimating otherwise missing CO<sub>2</sub> emissions (but not for CH<sub>4</sub> and N<sub>2</sub>O). However for 2002 onwards, the ‘unaccounted for’ amount has been reduced substantially due to the subtraction of fuel used for the non-combustion process of producing hydrogen (with CO<sub>2</sub> as by-product), of which the emissions are now reported under 1B2.

In fact, it is assumed that after the re-allocation of this fugitive CO<sub>2</sub> source and if more detailed CO<sub>2</sub> emissions reported by the individual refineries are used, which is presently not the case, no ‘unaccounted for’ liquid fuel would remain for these years. As the ‘unaccounted for’ amounts decreases over time, this causes the IEFs of CH<sub>4</sub> and N<sub>2</sub>O to increase over time because the ‘unaccounted for’ fuel use was determined solely to calculate CO<sub>2</sub> emissions due to net carbon losses only, not for other emissions. All remaining differences with the CO<sub>2</sub> calculation based on the national energy statistics and default emission factors are, therefore, shown up in the calculated carbon content of the combusted refinery gas and thus in the implied emission factor of CO<sub>2</sub> for liquid fuel.

#### Manufacture of Solid Fuels and Other Energy Industries [1A1c]

In accordance with IPCC classification guidelines emissions from fuel combustion for on-site coke production by the iron and steel company (Corus) are included in 1A2 “Manufacturing Industries and Construction” since this is an integrated coke, iron and steel plant (see Section 3.2.7). The emissions from the combustion of solid fuels of one independent coke production facility (Sluiskil), the operation of which discontinued in 1999 is also included in category 1A2.

Source category 1A1c also comprises:

- Combustion of “own” fuel use by the oil and gas production industry for heating purposes (the difference

Fuel	Amount of fuel used in 2008 (TJ NCV)	Emission factors (g/GJ)		
		CO <sub>2</sub>	N <sub>2</sub> O	CH <sub>4</sub>
Natural gas	498510	56.70	0.10	6.72
Coal	202840	96.23	1.40	0.44
Waste Gas	95526	68.77	0.10	3.60
Waste, biomass	30433	109.60	4.97	30.00
Blast Furnace Gas	23825	239.08	0.10	0.35
Solid biomass	20342	109.60	4.00	30.00

between the amounts of fuel produced and sold, minus the amounts of associated gas which is either flared or vented or otherwise lost by leakage).

- Fuel combustion for space heating and in use in compressors for gas and oil pipeline transmission by the gas, oil and electricity transport and distribution companies.

The share of 1A1c 'Manufacture of solid fuels (coke) and other energy industries (fuel production) in the total greenhouse gas emissions from the category 1A1 "Energy Industries" is approximately 3% in both 1990 and 2008. This category comprises mostly CO<sub>2</sub> emissions from the combustion of natural gas. The dominating source is the use for energy purposes in oil and gas production and in the transmission industry. The combustion emissions from oil and gas production refer to 'own use' of the gas and oil production industry, which is the difference between the amounts of fuel produced and sold, after subtraction the amounts of associated gas which is either flared or vented or otherwise lost by leakage. Production and sales data are based on the national energy statistics; amounts flared and vented are based on reports from the sector. CO<sub>2</sub> emissions from this source category increased from 1.5 Tg in 1990 to 1.9 Tg CO<sub>2</sub> in 2008 mainly due to the exploitation of less favourable production sites for oil and gas production compared with those exploited in the past. This fact explains the steady increase in time shown by this category with respect to gas consumption. The inter-annual variability in the emission factors for CH<sub>4</sub> from gas combustion is mainly due to the variable losses in the compressor stations of the gas transmission network, which are reported in the Annual Environmental Reports (MJVs) of the gas transport company are included here. The small amounts of solid fuel combustion by the one stand-alone coke production plant in Sluiskil, the operation of which was discontinued in 1999, are not separately recorded in the energy statistics but are included in the iron and steel industry (category 1A2a).

### 3.2.6.2 Methodological issues

The emissions from this source category are essentially estimated by multiplying fuel use statistics with country specific emission factors (Tier 2 method). Activity data are derived from the aggregated statistical data from the national energy statistics, which is published annually by the CBS (see [www.cbs.nl](http://www.cbs.nl)). The aggregated statistical data from the national energy statistics is based on confidential data from individual companies. When necessary, data from individual companies are also used, for example, when companies report a different emission factor for derived gases (see the following section).

For CO<sub>2</sub> and N<sub>2</sub>O, IPCC default emission factors are used, with the exception of CO<sub>2</sub> for natural gas and chemical waste gas and coal, for which country-specific emission factors are used. When available, company-specific or sector-specific emission factors have been used, in particular for derived gases such as refinery gas, chemical waste gas and blast furnace gas (see Annex 2, Table A2.1). If companies report different emission factors for derived gases, it is possible to deviate from the standard emission factor for estimating the emissions for these companies. The CH<sub>4</sub> emission factors are taken from Scheffer et al, 1997. An overview of the emission factors used for the most important fuels (up to 95% of the fuel use) in the sector Energy Industries [1A1] is provided in Table 3.3. Since some emission data in this sector originate from individual companies, the values (in Table 3.3) represent partly implied emission factors.

Notes to the source specific emission factors:

- The standard CH<sub>4</sub> emission factor for natural gas is 5.7 g/GJ. Only for gas fired CHP, a higher emission factor is used, which explains the higher emission factor for this sector.
- The CO<sub>2</sub> emissions from coal and blast furnace gas are based on emission data from the NEA. Therefore, the implied emission factor is slightly different from the standard country-specific emission factor.
- The CO<sub>2</sub> emissions from waste gas are CO<sub>2</sub> emissions occurring in the chemical industry and in refineries. The emissions are partly based on emission data from the NEA.
- The N<sub>2</sub>O emission factor from waste combustion is depending on the amount of waste incinerated in incinerators with or without a SNCR, which have emission factors of 9.43 g/GJ and 1.89 g/GJ respectively.

More details on emission factors, methodologies, the data sources used and country-specific source allocation issues are provided in the monitoring protocols (see [www.greenhousegases.nl](http://www.greenhousegases.nl), Protocol 0042: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from 'Stationary Combustion: Fossil Fuels' and Protocol 0078: Emissions from biomass combustion). According to the IPCC Guidelines, only fossil-fuel related CO<sub>2</sub> emissions are included in the total national inventory, thus excluding CO<sub>2</sub> from organic carbon sources from the combustion of biomass. The CO<sub>2</sub> from biomass from waste incineration is reported as memo item.

### 3.2.6.3 Uncertainties and time-series consistency

The uncertainty in CO<sub>2</sub> emissions of this category is estimated to be 2% (see Section 1.7 for more details). The accuracy of fuel consumption data in power generation and oil refineries is generally considered to be very accurate, with an estimated uncertainty of approximately 0.5%. The high accuracy in most

of these activity data is due to the limited number of utilities and refineries that report their large fuel consumption as part of the national energy statistics and which are verified as part of the European Emission Trading Scheme. The two exceptions are solids in power generation and liquids in refineries, which have a larger estimated uncertainty of 1% and 10%, respectively, based on the share of blast furnace gas in total solid consumption, the ‘unaccounted for’ liquids calculated for refineries and the recalculations made for 2002-2004 as presented in this report (Olivier et al., 2009). The high uncertainty in the liquids in refineries apply mainly to the years prior to 2002, for which accurate reported CO<sub>2</sub> emissions are not available at the required aggregation level. The consumption of gas and liquid fuels in the 1A1c category is mainly from the oil and gas production industry, where the split into own use and venting/flaring has proven to be quite difficult, and thus a high uncertainty of 20% is assigned. For other fuels a 10% uncertainty is used, which refers to the amount of fossil waste being incinerated and thus to the uncertainties in the total amount of waste and the fossil and biomass fractions.

For natural gas, the uncertainty in the CO<sub>2</sub> emission factor is now estimated to be 0.25% (instead of 1%) based on the recent fuel quality analysis reported by Heslinga and Van Harmelen (2007) and further discussed in Olivier et al. (2009). This value is not yet used in the uncertainty assessment in Section 1.7 and key source assessment in Annex 1. For hard coal (bituminous coal), an analysis was made of coal used in power generation (Van Harmelen and Koch, 2002). For the default power plant factor, 94.7 CO<sub>2</sub>/GJ is the mean value of 1270 samples taken in 2000, which is accurate within about 0.5%. However, in 1990 and 1998 the emission factor varies  $\pm 0.9$  CO<sub>2</sub>/GJ (see Table 4.1 in Van Harmelen and Koch, 2002); consequently when the default factor is applied to other years, the uncertainty is apparently larger, about 1%. Analysis of the default CO<sub>2</sub> emission factors for coke oven gas and blast furnace gas reveals uncertainties of about 10% and 15%, respectively (data reported by the steel plant). Since the share of BF/OX gas in total solid fuel emissions from power generation is about 15-20%, the overall uncertainty in the CO<sub>2</sub> emission factor of solids in power generation is estimated to be about 3%. The CO<sub>2</sub> emission factors of chemical waste gas and – to a lesser extent – of BF/OX gas are more uncertain than those of other fuels used by utilities. Thus, for liquid fuels in these sectors is a higher uncertainty of 10% assumed in view of the quite variable composition of the refinery gas used in both sectors. For natural gas and liquid fuels in ‘Oil and Gas Production’ (1A1c), uncertainties of 5% and 2% are assumed, respectively, which refer to the variable composition of the offshore gas and oil produced. For the CO<sub>2</sub> emission factor of other fuels (fossil waste), an uncertainty of 5% is assumed, which reflects the limited accuracy of the waste composition and of the carbon fraction per waste stream. The uncertainty in the emission factors of CH<sub>4</sub> and N<sub>2</sub>O from stationary combustion is estimated at about 50%, which is an aggregate for the various subcategories (Olivier et al., 2009).

#### 3.2.6.4 Source-specific QA/QC and verification

The trends in fuel combustion in the “Public Electricity and Heat Production” (1A1a) are compared to trends in domestic electricity consumption (production plus net imports).

First, large annual changes are identified and explained (e.g., changes in fuel consumption by joint ventures). For ‘Oil Refineries’ (1A1b), a carbon balance calculation is made to check completeness. Moreover the trend in total CO<sub>2</sub> reported as fuel combustion from refineries is compared to trends in activity indicators, such as total crude throughput. The IEF trend tables are then checked for changes, and inter-annual variations are explained in this NIR. More details on the validation of the energy data are to be found in the monitoring [protocol 0042: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from ‘Stationary Combustion: Fossil Fuels’](#)

#### 3.2.6.5 Source-specific recalculations

The emissions for category 1.A.1.C were recalculated as stated in the previous section. The emissions from natural gas are now in accordance with the final data from the oil and gas sector. The recalculation resulted in a reduction of the 2007 greenhouse gas emission of approximately 0.4 Tg CO<sub>2</sub> eq.

#### 3.2.6.6 Source-specific planned improvements,

From 2002 onwards, the data reported by the Dutch refineries are used to calculate plant-specific emission factors for CO<sub>2</sub> that represent an improvement compared to the use of the standard EF. This procedure will be continued. Analysis of the ETS data revealed that the use of these data would not improve the inventory,


### 3.2.7 Manufacturing Industries and Construction [1A2]

#### 3.2.7.1 Source category description

This source category consists of the six categories 1A2a ‘Iron and Steel’, 1A2b ‘Non-ferrous Metals’, 1A2c ‘Chemicals’, 1A2d ‘Pulp, Paper and Print’, 1A2e ‘Food Processing, Beverages and Tobacco’ and 1A2f ‘Other’. Within these categories, liquid fuel and natural gas combustion by the chemical industry, solid fuel combustion by the iron and steel industry and natural gas combustion by the food processing and other industries are the dominating emission sources. However, natural gas in the pulp and paper industries and liquid fuels (mainly for off-road machinery) in the other industries are also large emission sources. The shares of CH<sub>4</sub> and N<sub>2</sub>O emissions from industrial combustion are relatively small and these are no key sources. Natural gas is mostly used in the chemical, food and drinks and other industries; solid fuels (that means coal and coke-derived fuels, such as blast furnace/oxygen furnace gas) are mostly used in 1A2a ‘Iron and Steel’ industry; liquid fuels are mostly used in 1A2c ‘Chemicals’ industry and in 1A2f ‘Other’ industries.

Another feature of industry in the Netherlands is that it operates a large number of combined heat and power (CHP) facilities (and sometimes also steam boilers). Several of these facilities have changed ownership over time and are now operated as joint venture concerns with electrical utilities, the emissions of which are reported in “Energy Industries” (1A1a).

In the Netherlands the source category 1A2c Chemicals is the largest fuel user within 1A2 manufacturing industries and construction (see Figure 3.6). In this industry liquid fuel use is 109 PJ and natural gas use is 88 PJ in 2008. A second important industry is included in 1A2f other industries and




include emissions from mineral products (cement, bricks, glass, other building materials), textiles, wood, wood products and building construction industry. Solid fuels (35 PJ in 2008) are almost exclusively used in 1A2a Iron and Steel. In this industry, a limited amount of natural gas is also used. All other industries are almost completely run on natural gas.

In 2008, the share of CO<sub>2</sub> emissions from 1A2 “Manufacturing Industries and Construction” in the total national greenhouse gas emission inventory was estimated to be 16% compared to 21% in 1990. In contrast, the share of the other greenhouse gas emissions in this category is relatively small. Category 1A2c ‘Chemical industry’ is the largest contributor to CO<sub>2</sub> emissions, accounting for approximately 45% of the total emissions from the manufacturing industry in 2008.

In the period 1990–2008, CO<sub>2</sub> emissions from combustion in 1A2 “Manufacturing Industries and Construction” decreased by 16% (see Figure 3.7). The chemical industry contributes the

most to this decrease in emissions in this source category, with its contribution to CO<sub>2</sub> emissions decreasing by 4.7 Tg. Total CO<sub>2</sub> emissions from 1A2 “Manufacturing Industries and Construction” in 2008 decreased 1% compared to 2007.

The derivation of these figures, however, should also be viewed in the context of industrial process emissions of CO<sub>2</sub> since the separation of the source categories is not always fixed. Most industry process emissions of CO<sub>2</sub> are reported in CRF sector 2 (soda ash, ammonia, carbon electrodes and industrial gases such as hydrogen and carbon monoxide). However in manufacturing processes, this oxidation is accounted for in the energy statistics as the production and combustion of residual gases (e.g., in the chemical industry) – as is often the case in the Netherlands – then the corresponding CO<sub>2</sub> emissions are reported as combustion and not as an industrial process in sector 2.



### Iron and Steel [1A2a]

This sector refers mainly to the integrated steel plant Corus, which produces approximately 6000 kton crude steel (in addition to approximately 100 kton of electric steel production and iron foundries). Since Corus is an integrated plant, the category includes fuel combustion for on-site coke production as well as the emissions of the combustion of blast furnace gas and oxygen furnace gas in the steel industry.

The contribution of 1A2a 'Iron and steel' to the CO<sub>2</sub> emissions from 1A2 "Manufacturing Industries and Construction" was about 12% in 1990 and 17% in 2008. Inter-annual variations in CO<sub>2</sub> emissions from fuel combustion from the iron and steel industry can be explained as being mainly due to varying amounts of solid fuels used in this sector. In 2008 CO<sub>2</sub> emissions from solid fuel combustion of the iron and steel industry increased slightly (+0.3 Tg).

The 14% decrease in solid fuel use in 1999 and the 10% decrease in associated CO<sub>2</sub> emissions corresponds with the 8% decrease in crude steel production. When all CO<sub>2</sub> emissions from the sector are combined – including the net process emissions

reported under category 2C1 – total emissions closely follow the inter-annual variation in crude steel production. Total CO<sub>2</sub> emissions have remained rather constant in the period 1990–2008 even though production has increased by about 30%. This indicates a substantial energy efficiency improvement in the sector.

The inter-annual variation in the IEF for CO<sub>2</sub> from solid fuels is due to variable shares of BF/OX gas and coke oven gas, which have much higher and lower emission factors, respectively, than hard coal and coke have. The relative low IEFs in 1990–1994 compared to later years are due to the higher share of coke oven gas in the solid fuel mix in those years due to CO gas combustion by the independent coke manufacturer in Sluiskil, which was in these years not accounted for in the energy statistics separately but included in this category.

### Non-Ferrous Metals [1A2b]

This category consists mainly of two aluminium smelters. CO<sub>2</sub> emissions from anode consumption in the aluminium industry are included in 2C. This small source category only contributes about 0.2 Tg CO<sub>2</sub> to the total national greenhouse



gas inventory, predominantly from the combustion of natural gas. Energy use in the aluminium industry is largely based on electricity, the emissions of which are included in 1A1a 'Public electricity and heat production'.

The amounts of liquid and solid fuels vary considerably between years, but the differences in the amounts and related emissions are almost negligible. The inter-annual variation of the IEFs from liquid fuels is a result of changes in the mix of underlying fuels (e.g., the share of LPG which has a relatively low emission factor) and partly due to the small amounts used.

#### Chemicals [1A2c]

The share of 1A2c 'Chemical industry' to the total CO<sub>2</sub> emissions from 1A2 "Manufacturing Industries and Construction" decreased from 52% in 1990 to 45% in 2008. The combustion of natural gas and liquid fuels accounts for 40% respectively 58% in the CO<sub>2</sub> emissions from the chemical industry. CO<sub>2</sub> emissions from this source category have decreased by approximately 27% since 1990, which is mainly due to the 47% decrease in the consumption of natural gas during the same period.

The steady decreasing CO<sub>2</sub> emissions from the combustion of natural gas can be largely explained by the decreasing numbers of cogeneration facilities in this industrial sector. CO<sub>2</sub> emissions from liquid fuel combustion stem predominantly from the combustion of chemical waste gas. The marked decrease in liquid fuel consumption since 1995 is not due to a decrease in chemical production or data errors, but mainly to a large shift of ownership of a large cogeneration plant – one using chemical waste gas – into a joint venture, thus re-allocating it to energy industries. This also explains the 88% decrease in solid fuel combustion in 1994 and the 28% decrease in liquid fuel combustion in 1995. In these years, the then-existing coal-fired and oil-fired cogeneration plants, respectively, shifted to joint ventures and thus moved to the 'Energy Industry'.

Taking into account all CO<sub>2</sub> emissions, including the net process emissions included in category 2B and the re-allocation of CO<sub>2</sub> emissions to the energy industry, the total CO<sub>2</sub> emission level from the chemical industry was rather constant in the period 1990–2008. Given that since 1990 the production has increased significantly, the constant emission level indicates substantial improvements in the efficiency of energy use and/or structural changes within the chemical industry.

The increase in 2003 of the IEF for CO<sub>2</sub> from liquid fuels is also explained by the increase in the use of chemical waste gas and the change in composition. For CO<sub>2</sub> from chemical waste gas from liquid and solid fuels, source-specific emission factors are used for 1995 onwards based on data of selected years. For 16 individual plants, residual chemical gas from liquids is hydrogen, for which the specific CO<sub>2</sub> emission factor is 0. For CO<sub>2</sub> from phosphorous furnace gas, plant-specific values are used, with values around 149.5 kg/GJ. This gas is made from coke and therefore included in solid fuels. The operation of the phosphorous plant started around 2000, which explains the increase in the IEF for solid fuels to about

149.5 kg/GJ. For another 9 companies, plant-specific CO<sub>2</sub> emission factors were used based on annual reporting by the companies (most in the 50–55 range, with exceptional values of 23 and 95). The increased use of chemical waste gas (included in liquid fuels) since 2003 and the changes in the mix of compositions explain the increase in the IEF for liquid fuels from about 55 to about 67 kg/GJ. For 1990, an average sector-specific value for the chemical industry was calculated using the plant-specific factors for 1995 from the 4 largest companies and the amounts used per company in 1990. For more details, see Appendix 2 of the NIR 2005.

#### Pulp, Paper and Print [1A2d]

The contribution of 1A2d 'Pulp, paper and print' to CO<sub>2</sub> emissions from 1A2 "Manufacturing Industries and Construction" is estimated to be approximately 5% in 1990 and about 4% in 2008. In line with the decreased consumption of natural gas, CO<sub>2</sub> emissions have decreased by approximately 31% since 1990, of which a substantial fraction is used for co-generation. The relatively low CO<sub>2</sub> emissions in 1995 can be explained by re-allocation of emissions to the energy sector due to the above-mentioned formation of joint ventures. Between 1990 and 2008, CO<sub>2</sub> emissions from gaseous fuel combustion decreased by about 30%.

The amounts of liquid and solid fuel combustion vary considerably between years, but the amounts and related emissions are almost negligible. The inter-annual variation in the IEFs for liquid fuels is due to variable shares of derived gases and LPG in total liquid fuel combustion.

#### Food Processing, Beverages and Tobacco [1A2e]

The share of 1A2e 'Food processing, beverages and tobacco industries' in the CO<sub>2</sub> emission from 1A2 "Manufacturing Industries and Construction" was 12% in 1990 and 13% in 2008. CO<sub>2</sub> emissions decreased by almost 12% in the period 1990–2008. This is due to a decrease since 2003 of joint ventures of cogeneration plants located in the pulp and paper industry, of which the emissions were formerly allocated in 1A2e but are now reported under public electricity and heat production (1A1a). This shift in allocation corresponds with a CO<sub>2</sub> decrease of about 0.3 Tg. In 2008, CO<sub>2</sub> emissions from gaseous fuel combustion in this source category decreased by about 4% compared to last submission.

The amounts of liquid and solid fuels vary considerably between years, but the amounts and related emissions are verifiably small. The inter-annual variation in the IEFs for liquid fuels is due to variable shares of LPG in total liquid fuel combustion.

#### Other [1A2f]

This category includes all other industry branches, including mineral products (cement, bricks, other building materials, glass), textiles, wood and wood products. Also included are emissions from the building construction industry and from off-road vehicles (mobile machinery) for building construction and for the construction of roads and waterways and other off-road sources except agriculture (liquid fuels). The latter refers mainly to sand and gravel production.

Fuel	Amount of fuel used in 2008 (TJ NCV)	Emission factors (g/GJ)		
		CO <sub>2</sub>	N <sub>2</sub> O	CH <sub>4</sub>
Natural gas	243689	56.7	0.1	6.98
Waste Gas	106090	66.55	0.1	3.6
Gas / Diesel Oil	20809	74.3	0.6	5
Coke Oven Gas	15379	42.6	0.1	2.8
Blast Furnace Gas	14511	239.1	0.1	0.35
Solid biomass	8943	109.6	4	30
Other	12720	NA	NA	NA

The share of category 1A2f 'Other' (including construction and other off-road machinery) in CO<sub>2</sub> emissions from 1A2 "Manufacturing Industries and Construction" was approximately 17% in 1990 and 19% in 2008. Most of the 5.2 Tg CO<sub>2</sub> emissions from this source category in 2008 stem from gas combustion (3.3 Tg), while almost all of the remaining CO<sub>2</sub> emissions are associated with the combustion of liquid fuels (1.7 Tg CO<sub>2</sub>), of which off-road machinery accounts for 1.5 Tg CO<sub>2</sub>. CO<sub>2</sub> emissions from this source category have decreased by 6% since 1990. In 2008, total CO<sub>2</sub> emissions from the other manufacturing industries increased by 5% compared to 2007.

### 3.2.7.2 Methodological issues

The methods used for this source category are the same as those used for 1A1 Energy Industries. A country-specific top-down (Tier 2) method is used for calculating emissions for fuel combustion from 'Manufacturing Industries and Construction' (1A2). Fuel combustion emissions in this sector are calculated using fuel consumption data from national sectoral energy statistics and IPCC default emission factors for CO<sub>2</sub> and N<sub>2</sub>O, with the exception of CO<sub>2</sub> for natural gas and chemical waste gas and coal, for which country-specific emission factors are used. When available, company-specific or sector-specific emission factors have been used, in particular for derived gases such as chemical waste gas, blast furnace gas and coke oven gas (see Annex 2). More details on methodologies, data sources used and country-specific source allocation issues are provided in the monitoring protocols (see [www.greenhousegases.nl](http://www.greenhousegases.nl)).

An overview of the emission factors used for the most important fuels (up to 95% of the fuel use) in the sector Manufacturing Industries and Construction [1A2] is provided in Table 3.4. Since some emission data in this sector originate from individual companies, the values in the Table 3.4 represent partly implied emission factors.

Notes to the implied emission factors:

- The standard CH<sub>4</sub> emission factor for natural gas is 5.7 g/GJ. Only for gas powered CHP, a higher emission factor is used, which explains the higher emission factor for this sector.
- The CO<sub>2</sub> emissions from coke oven gas, blast furnace gas and waste gas are based on emission data from the NEA. Therefore, the implied emission factor is slightly different from the standard country-specific emission factor.
- Emission factor for CH<sub>4</sub> and N<sub>2</sub>O from gas/diesel oil used in Machinery are based on source specific estimation methods.

More details on emission factors methodologies, the data sources used and country-specific source allocation issues are provided in the monitoring protocols (see [Hwww.greenhousegases.nl](http://www.greenhousegases.nl)).

In 'Iron and Steel Industry', a substantial large fraction of total CO<sub>2</sub> emissions is reported as process emissions in CRF 2C1, based on net losses calculated from the carbon balance from the coke and coal inputs in the blast furnaces and the blast furnace gas produced. Since the fraction of BF/OX gas captured and used for energy varies over time, the trend in the combustion emissions of CO<sub>2</sub> accounted for by this source category should be viewed in association with the reported process emissions. The fuel combustion emissions from on-site coke production by the iron and steel company Corus are included here in 1A2a instead of in 1A1c, since these are reported in an integrated and aggregated manner. In addition to including the emission from Corus, this category also includes the combustion emissions of a small electric steel producer and – for the period 1990–1994 – of one small independent coke production facility for which the fuel consumption was not separately included in the national energy statistics during this period. The fugitive emissions, however, from all coke production sites are reported separately (see Section 3.2.7.1).

For the chemical industry, CO<sub>2</sub> emissions from the production of silicon carbide, carbon black, methanol and ethylene from the combustion of residual gas (produced as by-product from the non-energy use of fuels) are included in 1A2c 'Chemicals'. Although these CO<sub>2</sub> emissions are more or less process-related, they are included in 1A2 for practical purposes: consistency with Energy statistics that account for the combustion of residual gases. This inclusion in 1A2 is justified since there is no strict IPCC guidance on where to include those emissions.

The fuel consumption data in 1A2f 'Other Industries for Construction' and 'Other Off-road' are not based on large surveys. Therefore, the energy consumption data of this part of the Category 1A2f are the least accurate.

Details of the method for this source category are described in Protocol 0042: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from 'Stationary Combustion: Fossil Fuels'

### 3.2.7.3 Uncertainties and time-series consistency

The uncertainty in CO<sub>2</sub> emissions of this category is estimated to be about 3% (see Section 1.7 for more details). The accuracy of fuel consumption data in the manufacturing industries



is generally considered to be rather accurate, about 2%, with the exception of those for derived gases included in solids and liquids (Olivier et al., 2009). This includes the uncertainty in the subtraction of the amounts of gas and solids for non-energy/feedstock uses, including the uncertainty in the conversion from physical units to Joules, and the completeness of capturing blast furnace gas in total solid consumption and chemical waste gas in liquid fuel consumption.

For natural gas, the uncertainty in the CO<sub>2</sub> emission factor is now estimated to be 0.25% (instead of 1%) based on the recent fuel quality analysis reported by Heslinga and Van Harmelen (2007) and further discussed in Olivier et al. (2009), but not yet used in the uncertainty assessment in Section 1.7 and Annex 1. The 5% uncertainty estimate in the CO<sub>2</sub> emission factor for liquids is based on an uncertainty of 10% in the emission factor for chemical waste gas in order to account for the quite variable composition of the gas and its more than 50% share in the total liquid fuel use in the sector. An uncertainty of 10% is assigned for solids, which reflects the uncertainty in carbon contents of blast furnace gas/oxygen furnace gas based on the standard deviation in a 3-year average. BF/OX gas accounts for the majority of solid fuel use in this sector.

#### 3.2.7.4 Source-specific QA/QC and verification

The trends in CO<sub>2</sub> emissions from fuel combustion in the iron and steel industry, non-ferrous industry, food processing, pulp and paper and other industries are compared to trends in the associated activity data: crude steel and aluminium production, indices of food production, pulp and paper production and cement and bricks production. Large annual changes are identified and explained (e.g., changed fuel consumption by joint ventures). Moreover, for the iron and steel industry the trend in total CO<sub>2</sub> emissions reported as fuel combustion-related emissions (included in 1A2a) and industrial process emissions (included in 2C1) is compared to the trend in the activity data (crude steel production). A similar comparison is made for the total trend in CO<sub>2</sub> emissions from the chemical industry (sum of 1A2c and 2B) and trends split per main fuel type or specific process (chemical waste gas combustion and process emissions from ammonia production), IEF trend tables are checked for large changes and large inter-annual variations at different levels and explained in the NIR. More details on the validation of the energy data are found in the monitoring protocol 0042: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from 'Stationary Combustion: Fossil Fuels'.

#### 3.2.7.5 Source-specific recalculations

No source-specific re-calculations were performed during the compilation of this NIR.

#### 3.2.7.6 Source-specific planned improvements

There is no source-specific improvement planned.

### 3.2.8 Transport [1A3]

#### 3.2.8.1 Source category description

The source category 1A3 'Transport' comprises the following sources: 'Civil Aviation' (1A3a), 'Road Transportation' (1A3b), 'Railways' (1A3c), 'Water-borne Navigation' (1A3d) and 'Other

Transportation' (1A3e). The source category 'Civil Aviation' only includes emissions from domestic aviation - aviation with departure and arrival in the Netherlands. In the same manner, the source category 'Water-borne Navigation' only includes emissions from domestic inland navigation. Emissions from international aviation and navigation (aviation and marine bunkers) are reported separately; see Section 3.8. The source categories 'Road Transportation' and 'Railways' include all emissions from fuel sold to road transportation and railways in the Netherlands.

The source category 'Other Transportation' is not used, instead emissions from other mobile sources are reported in different source categories in the inventory. Emissions from national fisheries and off-road agricultural machinery, such as tractors, are included in 1A4c 'Agriculture, Forestry and Fisheries'; see Section 3.2.8, whereas emissions from other off-road machinery, such as road and building construction equipment, are reported under category 1A2f 'Other'. Emissions from fuel combustion by military aircraft and shipping activities are included in 1A5 'Other'; see Section 3.2.9. Energy consumption for pipeline transport is not recorded separately in the national energy statistics, but included in 1A1c for gas compressor stations and in 1A4a for pipelines for oil and other products.

#### Overview of shares and trends in emissions

The source category 1A3 'Transport' is responsible for 17% of total greenhouse gas emissions in the Netherlands. Between 1990 and 2008, the total greenhouse gas emission increased by 36% to 36.0 Tg CO<sub>2</sub> eq in 2008. This increase is mainly caused by an increase in fuel consumption and corresponding CO<sub>2</sub> emissions from road transportation. The greenhouse gas emissions from the transport sector are summarised in Figure 3.8. CO<sub>2</sub> emissions from 1A3b 'Road transport' are dominant in this source category, accounting for more than 95% of total emissions over the whole time-series.


After a decrease of greenhouse gas emissions by the transport sector in 2007, emissions increased again in 2008 (0.3 Tg) due to an increase in fuel consumption by road transportation (3 PJ) and a small decrease in the use of biofuels.

#### Civil Aviation [1A3a]

The share of 1A3a 'Civil Aviation' in total national greenhouse gas emissions was less than 0.1% in both 1990 and 2008. The reported energy use and greenhouse gas emissions by civil aviation in the Netherlands are based on a rough estimate of fuel consumption in 2000, which is applied to the whole time-series (see Section 3.5.2). Therefore, emissions remain constant over the whole time-series.

#### Road transport [1A3b]

The contribution of 1A3b 'Road transport' to the national inventory of CO<sub>2</sub> emissions was 16% in 1990 and 20% in 2008. Between 1990 and 2008, CO<sub>2</sub> emissions from road transport have increased from 25.5 Tg to 34.7 Tg. For the most part, this increase in CO<sub>2</sub> emissions is caused by a large increase in diesel use. In this time span, diesel use increased by 76% (121 PJ) due to a large growth in freight transport and the growing number of diesel passenger cars and light duty vehicles in



the Dutch car fleet. As a consequence, the share of diesel in fuel sales to road transport (PJ) has increased significantly between 1990 and 2008. The share of LPG has decreased significantly, while the share of gasoline has decreased slightly.


In 2007, CO<sub>2</sub> emissions from road transport decreased due to an increase in the use of biofuels in road transport: the share of biofuels in total energy use by road transport increased from 0.4% in 2006 to 2.8% in 2007. In 2008, the share of biofuels decreased to 2.6%. Combined with an increase in energy use by the transport sector, this resulted in an increase in CO<sub>2</sub> emissions.

CH<sub>4</sub> emissions from road transport fell from 7.5 Gg in 1990 to 2.2 Gg in 2008, which translates to a decrease of about 70%. This decrease results from a reduction in total VOC emissions resulting from the implementation of European Union emission legislation for new road vehicles. Total combustion and evaporative VOC emissions by road transport decreased by approximately 79% during the period 1990–2008, primarily due to the penetration of catalyst- and canister-equipped vehicles in the passenger car park. Between 2007 and 2008,

CH<sub>4</sub> emissions stayed at the same level due to a minor increase in emissions from biofuels (0.01 Gg). Even though the total use of biofuels decreased between 2007 and 2008, the use of ethanol in road transport increased. Therefore, the increase in ethanol use in road transportation led to an increase in CH<sub>4</sub> emissions from biofuels, even though the use of biodiesel and the total use of biofuels decreased.

N<sub>2</sub>O emissions from road transport increased from 0.9 Gg in 1990 to 1.6 Gg N<sub>2</sub>O in 1997, but decreased to 1.4 Gg in 2008. The increase in N<sub>2</sub>O emissions up to 1997 can be explained by the increased vehicle kilometres of petrol cars equipped with a catalytic converter, as the latter have higher emission factors than cars without this emission control technology. The subsequent decrease in N<sub>2</sub>O emissions between 1997 and 2008, despite an increase in vehicle-kilometres in this period, can be explained by a mix of developments:

- Subsequent generations of catalytic converters (the second was introduced in 1996) appear to have lower N<sub>2</sub>O emission factors (Gense, 2002);
- The share of diesel cars in the passenger car park, which are assumed to have lower emissions per vehicle kilometre



than catalyst-equipped petrol cars, has increased during the last few years.

Between 2007 and 2008, N<sub>2</sub>O emissions from road transport decreased slightly (0.02 Gg).

#### Rail transport [1A3c]

The share of 1A3c Rail transport in total national greenhouse gas emissions was only 0.1% in 1990 and 2008. Total emissions have remained fairly constant over the whole time-series (0.1 Tg CO<sub>2</sub> eq).

#### Navigation [1A3d]

The share of domestic water-borne navigation (1A3d) in total national greenhouse gas emissions is small. Emissions were about 0.4 Tg in 1990 and 0.6 Tg in 2008.

#### Key sources

CO<sub>2</sub> emissions from 1A3b 'Road Transport', all fuel types, and from Navigation are identified as key sources.

#### 3.2.8.2 Methodological issues

A detailed description of the methodology and data sources used to calculate transport emissions is provided in Klein et al. (2009) and in the monitoring protocols that can be found at [www.greenhousegases.nl](http://www.greenhousegases.nl) and are listed in Section 3.1.

#### Civil Aviation [1A3a]

An IPCC Tier 2 methodology is used for calculating the greenhouse gas emissions of 'Civil Aviation'. It is not possible to use fuel sales figures because there are no reliable data available on the distribution of these sales between national, international and military aviation. Therefore, the figures included in the national energy statistics (CBS Energy Balance) are not used. Instead, fuel consumption by domestic aviation has been roughly estimated based on the 2000 consumption figures of aviation petrol (avgas) and jet

kerosene for domestic flights in the Netherlands reported by the Civil Aviation Authority Netherlands (Pulles, 2000). Because of the very small amounts involved (342 TJ aviation petrol and 230 TJ jet kerosene) and since the number of domestic flights has decreased since 2000, there is no reason to expect an increase in energy use and emissions since 2000. Therefore, the figures for the year 2000 are used for the whole time-series. The emission factors used to calculate CO<sub>2</sub> emissions from kerosene and aviation petrol are derived from Vreuls and Zijlema (2009). Default IPCC emission factors for kerosene and aviation petrol are used to calculate emissions of CH<sub>4</sub> and N<sub>2</sub>O. Deliveries of bunkers to international aviation are not included in this source category.

Emissions of precursor gases (NO<sub>x</sub>, CO, NMVOC and SO<sub>2</sub>) reported in the NIR under domestic air traffic are the uncorrected emission values from the Netherlands Pollutant Emissions Register and refer to aircraft emissions associated with the Landing and Take-Off (LTO) cycles of Schiphol Airport. By far the most aircraft activities (>90%) in the Netherlands are related to Schiphol Airport; therefore emissions from other airports are ignored. No attempt has been made to estimate non-greenhouse gas emissions specifically related to domestic flights (including cruise emissions of these flights) since these emissions are almost negligible.

#### Road Transport [1A3b]

For national policy, air pollution from 'Road Transport' is, in general, calculated bottom-up from statistics collected on vehicle-kilometres and emission factors expressed in grams per vehicle-kilometre. The Revised IPCC Guidelines (IPCC, 1997) prescribe countries to report greenhouse gas emissions from combustion on the basis of fuel sales within the national territory. Thus, 'Road Transport' emissions of the direct greenhouse gases CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O are calculated using fuel sales data.

An IPCC Tier 2 methodology is used for calculating CO<sub>2</sub> emissions from 'Road Transport', using Dutch data on fuel sales to 'Road Transport' from Statistics Netherlands (CBS) and country-specific emission factors, as reported in Klein et al. (2009). See Annex 2 for more details.

An IPCC Tier 3 methodology is used for calculating CH<sub>4</sub> emissions from 'Road Transport', using data on the mass fractions of different compounds in total VOC emissions (Veldt, 1993). VOC emissions from 'Road Transport' are calculated using data on vehicle-kilometres from Statistics Netherlands (CBS), and VOC emission factors obtained from the Netherlands Organisation for Applied Scientific Research (TNO), as reported in Klein et al. (2009). The mass fraction is dependent on the fuel type and – for petrol vehicles – whether or not the vehicle is equipped with a catalyst. Petrol-fuelled vehicles equipped with a catalyst emit more CH<sub>4</sub> per unit of VOC than vehicles without a catalyst. In absolute terms, however, passenger cars with catalysts emit far less CH<sub>4</sub> than passenger cars without a catalyst because total VOC emissions are far lower.

Diesel-fuelled vehicles emit less CH<sub>4</sub> per unit of total VOC than petrol-fuelled vehicles without a catalyst. For each diesel-fuelled vehicle category, the calculation methodology distinguishes between several vehicle characteristics, such as age, fuel type and weight. In addition, the methodology also distinguishes three road types and takes into account cold starts.

N<sub>2</sub>O emissions from 'Road Transport' are also calculated using an IPCC Tier 3 (country-specific) methodology, using data on fuel deliveries from Statistics Netherlands and energy-specific emission factors. The emission factors for passenger cars and light-duty vehicles using petrol or LPG are based on country-specific data (Gense, 2002). Emission factors for diesel light-duty vehicles, heavy-duty vehicles, motorcycles and mopeds are based on Riemersma et al. (2003).

Since CO<sub>2</sub> emissions from 'Road Transport' are considered to be a key source (see Table 3.1), the present Tier 2 methodology complies with the IPCC Good Practice Guidance (IPCC, 2001). CH<sub>4</sub> and N<sub>2</sub>O emissions from 'Road Transport' are not a key source.

Emissions of all other compounds, including ozone precursors and SO<sub>2</sub>, which are more directly involved in air quality, are calculated bottom-up using vehicle-kilometre data (with fuel consumption figures that are somewhat different from the energy supply statistics. See Section 3.5.4 for more details).

Emissions from biodiesel and ethanol in gasoline are reported separately in the CRF. The emission calculation is comparable to the emission calculation for diesel/gasoline and is based on the sold amount of biodiesel and ethanol, as reported by Statistics Netherlands.

#### Rail Transport [1A3c]

Information on fuel use by diesel trains is obtained from the Dutch Railways (NS). For calculating CO<sub>2</sub> emissions, country-specific emission factors are used (Olivier, 2004); see Annex

2 for more details. For CH<sub>4</sub> and N<sub>2</sub>O emissions, IPCC default emission factors are used.

#### Navigation [1A3d]

An IPCC Tier 2 methodology is used for CO<sub>2</sub> emissions from domestic navigation. CO<sub>2</sub> emissions are calculated based on fuel deliveries to water-borne navigation in the Netherlands and country-specific emission factors (Klein Goldewijk et al., 2004). In the Netherlands, domestic commercial inland ships are allowed to use bunker fuels (sold without levies and VAT). Although the national energy statistics (CBS Energy Balance) distinguish between trips on the Rhine River and other inland shipping in the fuel consumption data for shipping, the sum of bunker fuel sales and domestic fuel sales to water-borne navigation in the national energy statistics includes fuel used for international navigation that should not be reported as part of domestic shipping according to IPCC Good Practice. Using the Emission Monitor Shipping (EMS) however, it is possible to distinguish between national and international navigation based on ton-kilometres travelled by ships (AVV, 2003). The share of fuel used by international navigation as calculated with the EMS is, therefore, subtracted from the total fuel sales to navigation in order to arrive at the fuel sales to national navigation, which is reported under 1A3d.

The present Tier 2 methodology level complies with the IPCC Good Practice Guidance (IPCC, 2001). Emissions from fisheries are allocated to the domestic source category 1A4c 'Commercial/ Institutional/Fisheries' as required by the IPCC Reporting Guidelines (see Section 3.2.5).

#### 3.2.8.3 Uncertainties and time-series consistency

The uncertainty in CO<sub>2</sub> emissions from 'Road Transport' is estimated to be about 4% in annual emissions (see Section 1.7 for more detailed information). The uncertainty in fuel sales to road transport is estimated to be 2% for petrol, 5% for diesel oil and 10% for LPG. The uncertainty in the CO<sub>2</sub> emission factor for petrol and diesel is calculated to be 0.4% and 0.2% respectively, while the uncertainty in the CO<sub>2</sub> emission factor for LPG is estimated to be 0.2%. For petrol and diesel fuel, the uncertainty in the emission factor for CO<sub>2</sub> is based on 50 samples of petrol and diesel fuel from petrol stations in the Netherlands in 2004 (Olivier, 2004). There are however indications that the carbon content of petrol and diesel fuels for road transport is changing due to tightening of European fuel quality standards. It is therefore unknown if these uncertainty figures are still valid. The uncertainty in CO<sub>2</sub> emissions from Road Transport might thus be underestimated.

The uncertainty in CH<sub>4</sub> emissions from 'Road Transport' is estimated to be approximately 60% in annual emissions. The share of CH<sub>4</sub> in VOC emissions is based on the report of Veldt and Van der Most (1993) and the composition of VOC emissions from 'Road Transport' has not been validated since. It is possible that the mass fraction of CH<sub>4</sub> has changed due to, for example, recent changes in the aromatic content of road transport fuels or improved exhaust after-treatment technology. The uncertainty in annual N<sub>2</sub>O emissions from 'Road Transport' is estimated to be 50%. N<sub>2</sub>O emission factors have not been updated recently and therefore are relatively uncertain.

The uncertainty in fuel used by (domestic) 'Civil Aviation' is presently estimated to be about +/- 50%. The high uncertainty is due to the lack of data on fuel sales for domestic flights. The uncertainty in emission factors for Civil Aviation is estimated to be +/- 0.5% for CO<sub>2</sub> and +/- 100% for CH<sub>4</sub> and N<sub>2</sub>O.

The uncertainty in fuel used by (domestic) 'Inland Navigation' is estimated to be approximately +/- 20%. The uncertainty in emission factors is estimated to be +/- 0.2% for CO<sub>2</sub> and +/- 100% for CH<sub>4</sub> and N<sub>2</sub>O.

The uncertainties of the CO<sub>2</sub> emission factors for 'Civil aviation' and 'Inland Navigation' (expressed as the standard error of the mean) are much lower than the uncertainties presented in the NIRs of other West European countries (Ramírez et al., 2007) and might be underestimated. For more details on the uncertainty estimates for 1A3 'Transport, see Oliver et al. (2009).

The methodologies used to estimate emissions from transport are consistent throughout the time-series.

### 3.2.8.4 Source-specific QA/QC and verification

#### Vehicle-kilometre approach versus IPCC approach

The Netherlands applies two methodologies to calculate CO<sub>2</sub> emissions from 'Road Transport': (1) the IPCC approach (based on fuel sales) and (2) the (informal) domestic approach, which is based on fuel consumed on national territory, calculated on the basis of transport statistics in terms of vehicle-kilometres travelled and fuel consumption per vehicle kilometre. A comparison between both approaches gives an indication of the validity of the fuel sales data.

The difference in fuel consumption inferred from transport statistics compared with data on fuel sales inferred from supply statistics on deliveries to fuelling stations is in the range of about 4–9% (data on fuel sales are higher). This is caused mostly by differences in diesel and LPG figures, which differ annually by up to 23% with an average of about 12% and 14% respectively (see NIR 2007, Figure 3.8). Differences in petrol sales and calculated petrol consumption are much smaller with an average of around 2%. The differences in fuel sales and the calculated fuel consumption on Dutch territory can partly be explained by the fact that part of the fuel sold in the Netherlands is consumed abroad and vice versa (Van Amstel et al., 2000a). Another explanation is the lack of a reliable fuel consumption figures per vehicle kilometres for most vehicle types. It can be concluded that roughly both methods show similar trends in fuel consumption by fuel type over the last 10 years.

### 3.2.8.5 Source-specific recalculations

The greenhouse gas emissions from 1A3d 'Navigation' for the year 2007 have been recalculated this year compared to last year's submission, based on new data on fuel use by domestic inland navigation. The data on fuel use by domestic aviation in 2007 were not available in time for last year's submission; therefore emissions in 2007 were held constant compared to 2006. This year, new data on fuel use in 2007 became

available. This recalculation has led to an increase in the 2007 greenhouse gas emissions by approximately 2 Gg CO<sub>2</sub> eq.

The greenhouse gas emissions from 1A3c 'Railways' for the year 2007 have also been recalculated using new data on fuel use by railways that became available after last year's submission.

### 3.2.8.6 Source-specific planned improvements

There is no source-specific improvement planned.

## 3.2.9 Other Sectors [1A4]

### 3.2.9.1 Source category description

Source category 1A4 "Other sectors" comprises the following categories:

- 1A4a 'Commercial and Institutional Services'. This sector comprises commercial and public services such as banks, schools and hospitals, and trade, retail and communication; it also includes the production of drinking water and miscellaneous combustion emissions from waste handling activities and from wastewater treatment plants.
- 1A4b 'Residential'. This sector refers to fuel consumption by households for space heating, water heating and cooking. Space heating requires about three-quarters of the total consumption of natural gas.
- 1A4c 'Agriculture, Forestry and Fisheries'. This sector comprises stationary combustion emissions from agriculture, horticulture, greenhouse horticulture, cattle breeding and forestry, and fuel combustion emissions from fisheries and from off-road machinery used in agriculture (mainly tractors).


CO<sub>2</sub> emissions of 1A4 "Other sectors" increased by 2% in the period 1990–2008. In 2008, CO<sub>2</sub> emissions from 1A4 "Other sectors" increased by 11% compared to the 2007 level. This increase is the result of a relative cold winter in 2008.

The share of CO<sub>2</sub> emissions from 1A4 "Other sectors" in total national CO<sub>2</sub> equivalent emissions (excluding LULUCF) was about 18% in 1990 and 19% in 2008, respectively. The share of CH<sub>4</sub> emissions from this source category in the national total greenhouse gas emissions is very small (0.7%); the share of N<sub>2</sub>O emissions is almost negligible; 1A4b 'Residential' is the main contributor, contributing approximately 10% to the total national CO<sub>2</sub> equivalent emissions.

About 13% of the total CH<sub>4</sub> emissions in the Energy sector originate from the 'Residential' sector (0.3 Tg CO<sub>2</sub> eq, see Table 3.1). Over 80% of these CH<sub>4</sub> emissions stem from gas combustion in particular from cooking losses; the remainder is from biofuel combustion.

The decreased emissions in 'Agricultural' are due to energy conservation measures in the category of greenhouse horticulture, CO<sub>2</sub> emissions from off-road machinery used in agriculture and from fisheries are included in the total emissions from category 1A4c (total CO<sub>2</sub> emissions from 1A4c: approximately 10 Tg CO<sub>2</sub>).





Within this source category, the combustion of gases and liquids form a key source for CO<sub>2</sub> emissions. See Table 3.1 for details.

#### Commercial / Institutional [1A4a]

The CO<sub>2</sub> emission in the 'Commercial/Institutional Services' sector has increased since 1990 by 30%. However, when a temperature correction is taken into account, the structural, anthropogenic trend shows a somewhat lower increase in this period. The 'Commercial/Institutional Services' sector has grown strongly during this period: the amount of manpower (in man-years) increased by 35% in the period 1990–2008. This increase is roughly comparable with the increase of fuel consumption (excluding electricity) of 35%, and thus of CO<sub>2</sub> emissions. It should be noted that about 0.8 Tg in 2008, are emissions from cogeneration facilities, which may also provide electricity to the public grid.

The emission trends should not be considered to be very robust. The fossil fuel consumption of natural gas and the small uses of liquid and solid fuels in this category show a very large inter-annual variation due to the relatively large inaccuracy of fuel consumption data in the energy statistics.

This large inaccuracy is a result of the calculation scheme used in the national energy statistics, which allocates all fossil fuel use remaining after subtraction of the amounts allocated to the previous source categories (1A1, 1A2, 1A3) and other categories (1A4b and 1A4c) to this category. Thus, all uncertainties in the other allocations accumulate in this remaining category, which also results in large inter-annual changes in the underlying fuel mix of solid and liquid fuels. This explains the relatively large inter-annual variation that can be observed in the IEFs of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O for solid and liquid fuels. As mentioned above, the strong decrease of CO<sub>2</sub> emissions in 2005, and of gas and solids consumption, must be an artefact of the very large uncertainty in the fuel consumption data of this category, which is for natural gas magnified in 2005 by the assumption of almost constant gas consumption in the agricultural category.

For 1991–1994, in particular, the detailed fuel mix assumed for liquid and solids fuels was different from the adjoining years 1990 and 1995 due to the revision of the energy statistics at a high aggregation level (discussed in Section 3.1.1). The biomass combustion reported here refers mainly to the combustion of biogas recovered by waste water treatment

plants (WWTP), which shows a rather smooth increasing trend, and biomass consumption by industrial companies, which are classified in this economic sector, for example landfill gas used as fuel. According to the renewable energy statistics, the latter increased substantially in 2005.

#### Residential [1A4b]

When corrected for the inter-annual variation in temperatures, the trend in total CO<sub>2</sub> – i.e. in gas consumption – becomes quite smooth, with inter-annual variations of less than 5%. The variations are much larger for liquid and solid fuels because of the much smaller figures. The biomass consumption is almost all wood (fuel wood, other wood: also less than 1% waste). The cause of the irregularity in biomass fuel use in 1999 is unknown but may be due to a small error in the survey procedures (for details see the monitoring protocol 0078 on biomass fuel combustion).

The IEF for CH<sub>4</sub> from national gas combustion is the aggregate of the standard emission factor for gas combustion of 5.7 g/GJ plus the 30 g/GJ of total residential gas combustion that represents start-up losses, which occur mostly in cooking but also in central heating and warm water production devices. This second component is neither accounted for in the IPCC default nor in emission factors used by most other countries.

In the ‘Residential’ sector, CO<sub>2</sub> emissions have remained almost constant since 1990. The structural anthropogenic trend including a temperature correction shows a significant decrease in this period. Although the number of households and residential dwellings increased since 1990, the average fuel consumption per household decreased more, mainly due to the improved insulation of dwellings and the increased efficiency of heating apparatus (increased use of high-efficient boilers for central heating).

#### Agriculture / Forestry / Fishing [1A4c]

Most of the energy in this source category is used for space heating and water heating; although some energy is used for cooling. The major fuel used in the categories is natural gas, which accounts for approximately 81% of total fossil fuel consumption; much less liquid fuel is used by off-road machinery and by fisheries. Almost no solid fuels are used in these sectors.

Total CO<sub>2</sub> emissions in the ‘Agriculture, Forestry and Fisheries’ category have decreased since 1990, mainly due to decrease in gas consumption for stationary combustion as a result of different energy conservation measures (amongst others in the greenhouse horticulture). The surface area of heated greenhouses has increased but their energy consumption has been reduced. It should be noted that about 1 Tg of the CO<sub>2</sub> emissions from the agricultural sector are emissions from cogeneration facilities, which may also provide electricity to the public grid.

In addition, since the fall of 2005 CO<sub>2</sub> from the hydrogen production plant in a refinery is starting to be used for crop fertilisation in greenhouse horticulture, thereby avoiding some CO<sub>2</sub> emissions otherwise generated by CHP facilities merely for producing CO<sub>2</sub> for horticulture. Total annual amounts, however, will be limited to a few tenths of Tg

CO<sub>2</sub>. In addition, in 2008 production and use of biogas from composting of manure in the ‘Agriculture/Forestry/Fisheries’ category increased from virtually zero to 0.5 PJ.

The CO<sub>2</sub> emissions from off-road machinery in agriculture in 2008 amount to 1.3 Tg, whereas the CO<sub>2</sub> emissions from fisheries amount to about 0.9 Tg. CO<sub>2</sub> emissions from fisheries have shown a decreasing trend in recent years, whereas CO<sub>2</sub> emissions from agricultural machinery have fluctuated in these years. In 2008, the CO<sub>2</sub> emissions from agricultural machinery increased by approximately 0.2 Tg compared to 2007.

#### 3.2.9.2 Methodological issues

In this category liquid and gaseous fossil fuels are key sources of CO<sub>2</sub> emissions (in particular, gaseous fossil fuels, which cover about 81% of the source category 1A4). Emissions from the combustion of gases in the categories 1A4a, 1A4b and 1A4c are identified as key sources, as are the emissions from the combustion of liquids in 1A4c. IPCC Tier 2 methodologies are used to calculate greenhouse gas emissions from stationary and mobile combustion in this category. More details on methodologies, the data sources used and country-specific source allocation issues are provided in the monitoring protocols ([www.greenhousegases.nl](http://www.greenhousegases.nl)).

The activity data for the ‘Residential’ sector (1A4b) and from stationary combustion in agriculture (1A4c-i) are compiled using data from separate surveys for these categories (‘HOME’ survey, formerly the ‘BAK’ and ‘BEK’ surveys, and LEI). However, due to late availability of the statistics on agricultural fuel use, preliminary data are often used for the most recent year in the national energy statistics. Also, it is likely that trends in agricultural fuel consumption are estimated using indicators that take no account of the varying heating demand due to changes in heating degree days. The fuel consumption data in 1A4a ‘Commercial/Institutional Services’ is determined by subtracting the energy consumption allocated to the other source categories (1A1, 1A2, 1A3) and other categories (1A4b and 1A4c) from the total energy consumption, which means that resulting activity data are the least accurate of all three categories. The emission factors for CO<sub>2</sub> from natural gas and from diesel fuel are based on country-specific data; for the CH<sub>4</sub> emission factors country specific values are also used, which for the residential gas combustion includes start-up losses, a factor mostly neglected by other countries. For other factors IPCC defaults were used (see Annex 2 and the monitoring protocols on [www.greenhousegases.nl](http://www.greenhousegases.nl)).

Emissions from ‘Off-road Machinery and Fisheries’ in this category (1A4c-ii) are calculated based on IPCC Tier 2 methodologies. The fuel use data is combined with country-specific emission factors for CO<sub>2</sub> and IPCC default emission factors for N<sub>2</sub>O and CH<sub>4</sub>. Fuel consumption by ‘Fisheries’ (1A4c-ii) is included in the Netherlands international bunker statistics, which are part of the national energy statistics. However, since the national energy statistics does not separately account for fisheries, it is not possible to use fuel sales figures in the national energy statistics. Instead, the fuel consumption of diesel oil and heavy fuel oil by fisheries is estimated based on statistics of the number of days at

Fuel	Amount of fuel used in 2008 (TJ NCV)	Emission factors (g/GJ)		
		CO <sub>2</sub>	N <sub>2</sub> O	CH <sub>4</sub>
Natural gas	635802	56.7	0.1	84.4
Gas / Diesel Oil	28079	74.3	0.6	5

sea ('hp-days') of four types of Dutch fishing ships. This information is compiled by LEI, and the estimate includes specific fuel consumption per ship [per day and per unit of power (hp) based on a study of TNO (Hulskotte, 2004b)]. This amount is reported as part of category 1A4c and subtracted from the amount of bunker fuel consumption in the national energy statistics. The modified bunker figures are reported as a Memo item. For more details, see the monitoring [protocol 0050 for Fisheries](#)

Fuel consumption by off-road agricultural machinery is derived from the EMMA model (Hulskotte, 2009). This model is based on sales data for different types of mobile machinery and assumptions on the average use (hours per year) and the fuel consumption (kilogram per hour) for different machine types. The results of the EMMA model are adjusted to make sure the model is consistent with the national energy statistics (see also 1A2f). It is assumed that only diesel fuel is used by mobile machinery. The use of gasoline and LPG is small and not specifically part of the national energy statistics. Instead it is part of the total use of gasoline and LPG in the transport sector.

An overview of the emission factors used for the most important fuels (up to 95% of the fuel use) in the Other Sectors [1A4] is provided in Table 3.5. Since some emission data is used for individual companies, some of these values represent implied emission factors.

Notes to the implied emission factors:

- The standard CH<sub>4</sub> emission factor for natural gas is 5.7 g/GJ. Only for gas engines, a higher emission factor is used, which explains the higher emission factor for this sector.
- Emission factor for CH<sub>4</sub> and N<sub>2</sub>O from gas/diesel oil used in Machinery are based on source specific estimation methods.

More details on emission factors, methodologies, the data sources used and country-specific source allocation issues are provided in the monitoring protocols (see [www.greenhousegases.nl](http://www.greenhousegases.nl)).

### 3.2.9.3 Uncertainties and time-series consistency

It should be noted that the energy consumption data for the total category 1A4 "Other sectors" are much more accurate than the data for the other categories of 1A4. In particular, energy consumption by the commercial/institutional and – to some extent – agricultural categories (in particular the latest year) is monitored less accurately than that by the 'Residential' sector. Trends of emissions and activity data of these categories should be treated with some caution when drawing conclusions. The uncertainty in total CO<sub>2</sub> emissions from this source category is about 6%, with an uncertainty of the composite parts of about 5% for the 'Residential' sector,

10% for the 'Agricultural' sector and 20% for the 'Service' sector (see Section 1.7 and Annex 1 for more details).

The uncertainty in gas consumption data is estimated at 5% for the 'Residential' sector, 10% for 'Agriculture' and 20% for the 'Commercial' sector. An uncertainty of 20% is assumed for liquid fuel use for 'Off-road Machinery and Fisheries' and in the 'Service' sector. Since the uncertainty in small figures in national statistics are generally larger than large numbers, as also indicated by the high inter-annual variability of the data, the uncertainty in solid fuel consumption is estimated to be even higher at 50%. However, the uncertainty of fuel statistics for the *total* "Other sectors" is somewhat smaller than the data for the sectors: consumption per fuel type is defined as the remainder of total national supply after subtraction of amount used in the 'Energy', 'Industry' and "Transport" sectors. Subsequently, energy consumption by the residential and agricultural sectors is estimated separately using a trend analysis of sectoral data ('HOME' survey of the 'Residential' sector and LEI data for 'Agriculture').

For natural gas the uncertainty in the CO<sub>2</sub> emission factor is now estimated at 0.25% (instead of 1%) based on the recent fuel quality analysis reported by Heslinga and Van Harmelen (2007) and further discussed in Olivier et al. (2009), but this has not yet been used in the uncertainty assessment in Section 1.7 and Annex 1. For the CO<sub>2</sub> emission factors for liquids and solids, uncertainties of 2% and 5% were assigned. The uncertainty in CH<sub>4</sub> and N<sub>2</sub>O emission factors is estimated to be much higher (about 50% and 100%, respectively).

If the changes made in earlier years are indicative of the quality of the data (see Table 3.22 of NIR 2004 and Table 3.26a of NIR 2005; Klein Goldewijk et al., 2004; 2005), then the uncertainty in total CO<sub>2</sub> emissions from this source category is about 7%, with an uncertainty of the composite parts of 3% for the 'Residential' sector, 15% for the 'Agricultural' sector and 20% for the 'Service' sector. This is in line with the results from the Tier 1 uncertainty analysis.

Since most of the fuel consumption in this source category is used for space heating, the gas consumption from the "Other sectors" varies considerably across years due to variations in winter temperatures over time. For trend analysis a method is used to correct the CO<sub>2</sub> emissions from gas combustion for the varying winter temperatures. This involves the use of the number of heating degree-days under normal climate conditions, which is determined by the long-term trend as explained in Visser (2005).

The deviating IEFs in the 1991–1994 period of CH<sub>4</sub> for liquids and gas and of N<sub>2</sub>O for liquids are due to the higher aggregation level used in the revised energy statistics.



Category		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Military ships	Emission factor	75.25	2.34	1.87
Military aviation	Emission factor	72.9	5.8	10
Total	Emissions in 2008 (Gg)	392.32	0.04	0.02

<sup>1)</sup> Source: Hulsakotte, 2004b.

### 3.2.9.4 Source-specific QA/QC and verification

The trends in CO<sub>2</sub> from the three categories were compared to trends in related activity data: the number of households, number of persons employed in the 'Service' sectors and the area of heated greenhouses. Large annual changes were identified in special trend tables and explanations were sought (e.g., inter-annual changes in CO<sub>2</sub> emissions by calculating temperature-corrected trends to identify the anthropogenic emission trends). The trend tables for the IEFs were then used to identify large changes and large inter-annual variations at the category level for which explanations were sought and included in the NIR. More details on the validation of the energy data can be found in the monitoring protocol 0042: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from 'Stationary Combustion: Fossil Fuels'.

### 3.2.9.5 Source-specific recalculations

This submission includes revised activity data for the gas consumption in categories 1AA4A and 1AA4C. Based on new data from agriculture fuel use (natural gas) which were allocated in this sector are now partly reallocated to 1AA4A.

### 3.2.9.6 Source-specific planned improvements,

There are no source-specific recalculations envisaged.

## 3.2.10 Other [1A5]

### 3.2.10.1 Source category description

Category 1A5 'Others' includes the emissions from military ships and aircraft (in 1A5b). This category is not a key source. CO<sub>2</sub> emissions from this source category are approximately 0.5 Tg, with some inter-annual variation caused by different levels of operations, including fuel use for multilateral operations, which are included here. Emissions of CH<sub>4</sub> and N<sub>2</sub>O are negligible.

The emission factors used are presented in Table 3.6.

### 3.2.10.2 Methodological issues

A country-specific top-down (Tier 2) method is used for calculating the emissions for fuel combustion from 1A5 'Others'. The fuel combustion emissions in this sector are calculated using fuel consumption data for both shipping and aviation that have been obtained from the Ministry of Defence and are the total emissions for domestic military shipping and aviation activities and the so-called multilateral operations. The fuel data for aviation consist of a mixture of jet kerosene, F65 and SFC. In the national energy statistics these activity data are included in the bunker fuel consumption. The sector-specific emission factors that are used are those reported by the Ministry of Defence. The methodology and data sources for the calculation of these emissions can be found on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 3.2.10.3 Uncertainties and time-series consistency

The uncertainty in CO<sub>2</sub> emissions from fuel combustion from 1A5 'Others' is estimated to be about 20% in annual emissions. The uncertainty for CH<sub>4</sub> and N<sub>2</sub>O emissions is estimated to be about 100%. The accuracy of fuel consumption data is tentatively estimated at 20%. For emission factors, the uncertainties were estimated at 2% for CO<sub>2</sub> and 100% for CH<sub>4</sub> and N<sub>2</sub>O.

A consistent methodology is used throughout the time-series. The time-series consistency of the activity data is good due to the continuity in the data provided.

### 3.2.10.4 Source-specific QA/QC and verification

This source category is covered by the general QA/QC procedures discussed in Chapter 1.

### 3.2.10.5 Source-specific recalculations

There have been no source-specific recalculations.

### 3.2.10.6 Source-specific planned improvements,

There are no source-specific planned improvements.

## 3.3 Fugitive emissions from fuels [1B]

This source category includes fuel-related emissions from non-combustion activities in the energy production and transformation industries:

- 1B1 'Solid Fuels' (coke manufacture)
- 1B2 'Oil and Gas' (production, gas processing, oil refining, transport, distribution).

The contribution of emissions from source category 1B to the total national greenhouse gas emissions inventory was 1.3% in 1990 and 1.0% in 2008.

Between 1990 and 2008, total greenhouse gas emissions in this category decreased from 2.8 Tg to 2.2 Tg.

### 3.3.1 Solid fuels [1B1]

#### 3.3.1.1 Source category description

Fugitive emissions from this category refer mainly to CO<sub>2</sub> from 1B1b 'Coke Manufacture' (see Table 3.1). The Netherlands currently has only one on-site coke production facility at the iron and steel plant of Corus. A second independent coke producer in Sluisil discontinued its activities in 1999. The fugitive emissions of CO<sub>2</sub> and CH<sub>4</sub> from both coke production sites are included here. We note that fugitive emissions from all coke production sites are included (in contrast with fuel combustion emissions from on-site coke production by the iron and steel industry, which are included in 1A2a

instead of 1A1c, since these are reported in an integrated and aggregated manner).

There are no fugitive emissions from coal mining and handling activities (1B1a) in the Netherlands; these activities ceased with the closing of the last coal mine in the early 1970s.

With respect to fugitive emissions from 'Charcoal Production', the Netherlands has one large state of the art production location that serves most of the Netherlands and also occupies a large share of the market of our neighbouring countries. These emissions are presently not accounted for. Recent research showed only minor CH<sub>4</sub> emissions of the plant due to the abatement technology used. Because no activity data is available on a regular basis we do not foresee inclusion of this minor source in the inventory.

### 3.3.1.2 Methodological issues

The CO<sub>2</sub> emissions related to transformation losses (1B1) from *coke ovens* are based on national energy statistics of coal inputs and coke and coke oven gas produced and a carbon balance of the losses. The completeness of the accounting in the energy statistics of the coke oven gas produced is not an issue, since the not-captured gas is by definition included in the net carbon loss calculation used for the process emissions.

Detailed information on activity data and emission factors can be found in the monitoring protocols on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 3.3.1.3 Uncertainties and time-series consistency

For emissions from 'Coke Production' (included in 1B1b) the uncertainty in annual CO<sub>2</sub> emissions from this source category is estimated to be about 50%. This uncertainty refers to the precision with which the mass balance calculation of carbon losses in the conversion from coking coal to coke and coke oven gas can be made (for details, see Olivier et al., 2009).

The methodology used to estimate emissions from solid fuel transformation is consistent throughout the time-series.

### 3.3.1.4 Source-specific QA/QC and verification

No source-specific QA/QC and verification.

### 3.3.1.5 Source-specific recalculations

There have been no source-specific recalculations in comparison to the previous submission.

### 3.3.1.6 Source-specific planned improvements

No source-specific improvements planned.

## 3.3.2 Oil and Natural Gas [1B2]

### 3.3.2.1 Source category description

The fugitive emissions – mostly CH<sub>4</sub> – from category 1B2 comprise non-fuel combustion emissions from flaring and venting, emissions from oil and gas production, emissions from gas transport (compressor stations) and gas distribution networks (pipelines for local transport) and oil refining.

The fugitive CO<sub>2</sub> emissions from refineries are included in the combustion emissions reported in category 1A1b. In addition, the combustion emissions from exploration and production are reported under 1A1c.

From the 2007 submission the Process emissions of CO<sub>2</sub> from a hydrogen plant of a refinery (about 0.9 Tg CO<sub>2</sub> per year) are reported in this category. Refinery data specifying these fugitive CO<sub>2</sub> emissions are available from 2002 onwards (environmental report from the plant) and re-allocated from 1A1b to 1B2a-iv for 2002 onwards.

CO<sub>2</sub> from gas flaring (including the venting of gas with high carbon dioxide content) and methane from gas venting/flaring are identified as key sources (see Table 3.1).

Gas production of which about 50% is exported, and gas transmission varies according to demand – in cold winters, more gas is produced – which explains the peak in 1996. The length of the gas distribution network is still gradually expanding as new neighbourhoods are being built; mostly using PVC and PE, which are also used to replace cast iron pipelines (see Table 3.44 in NIR 2005). The IEF for gas distribution gradually decreases as the share of grey cast iron pipelines decreases due to gradual replacement and expansion of the network. The present share is about 6%; in 1990 this was still 11%.

There is very little oil production in the Netherlands. The emission factors of CO<sub>2</sub> and CH<sub>4</sub> from oil and gas production, in particular for venting and flaring, have been reduced significantly and are now about 25% of the 1990 level. This is due to the implementation of environmental measures to reduce venting and flaring by optimizing the utilisation of energy purposes of produced gas that was formerly wasted.

### 3.3.2.2 Methodological issues

Country-specific methods comparable with the IPCC Tier 3 method are used to estimate the emission of fugitive CH<sub>4</sub> and CO<sub>2</sub> emissions from 'Oil and Gas Production and Processing' (1B2) (Grontmij, 2000). The emissions for CH<sub>4</sub> from gas venting and flaring are plant-specific.

The IPCC Tier 3 method for CH<sub>4</sub> from 'Gas Distribution' (1B2) is based on two country-specific emission factors of 610 m<sup>3</sup> (437 Gg) methane for grey cast iron and 120 m<sup>3</sup> (86 Gg) for other materials per 1000 km of pipeline due to leakages; the emission factors are based on seven measurements of leakage per hour on grey cast iron at one pressure level and on 18 measurements at three pressure levels for other materials (PVC, steel, nodular cast iron and PE) and subsequently aggregated to factors for the material mix in 2004. From 2004 onwards, the gas distribution sector annually records the number of leaks found per material, and any future possible trends in the emission factors are derived from these data. Fugitive emissions of methane from refineries in category 1B2 are based on a 4% share in total VOC emissions reported in the annual environmental reports of the Dutch companies (Spakman et al., 2003). For more information, see the monitoring protocols available on [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 3.3.2.3 Uncertainty and time-series consistency

The uncertainty in CO<sub>2</sub> emissions from gas flaring and venting is estimated to be about 50%, while the uncertainty in methane emissions from oil and gas production (venting) and gas transport and distribution (leakage) is estimated to be 25% and 25% in annual emissions, respectively. The uncertainty in the emission factor of CO<sub>2</sub> from gas flaring and venting (1B2) is estimated at 2%. This uncertainty takes the variability in the gas composition of the smaller gas fields into account for flaring. For venting, this uncertainty accounts for the high amounts of CO<sub>2</sub> gas produced at a few locations, which is then processed and the CO<sub>2</sub> extracted and subsequently vented. For CH<sub>4</sub> from fossil fuel production (gas venting) and distribution, the uncertainty in the emission factors is estimated to be 25% and 50%, respectively. This uncertainty refers to the changes in reported venting emissions by the oil and gas production industry over the past years and to the limited number of measurements made of gas leakage per leak for different types of materials and pressures, on which the Tier 2 methodology for methane emissions from gas distribution is based.

A consistent methodology is used to calculate emissions throughout the whole time-series.

### 3.3.2.4 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures which are discussed in Chapter 1.

### 3.3.2.5 Source-specific recalculations

There have been no source-specific recalculations in comparison to the previous submission.

### 3.3.2.6 Source-specific planned improvements

There is no source specific improvement planned.



# Industrial processes [CRF Sector 2]

# 4

## ■ Major changes in sector 2 Industrial Processes compared to the National Inventory Report 2009

*Emissions:* Mainly due to the reduction in N<sub>2</sub>O emissions from the Chemical Industry, the total greenhouse gas emissions in this sector decreased by 45% in 2008 compared to 2007.

*Key sources:* Nitric acid production is from now on no longer a Tier 2 level key source.

*Methodologies:* In response to the ERT report 2009, the method to estimate HFC emissions from Stationary refrigeration (2F1) has been changed. From now on, the complicated mass-balance method is replaced by a straightforward EF-based method.

Furthermore, information about pre-charged refrigeration equipment and new SF<sub>6</sub> emissions from the source Electrical equipment came available. As a result, the emissions of these sources have been changed for a number of years.

### 4.1 Overview of sector

Emissions of greenhouse gases in this sector include all non-energy-related emissions from industrial activities (including construction) and all emissions from the use of the F-gases HFCs, PFCs and SF<sub>6</sub> (including their use in other sectors). Please note that due to the specific Dutch estimation methods and absence of required activity data, only the actual emissions of F-gases can be estimated in detail. The potential emissions can not be calculated in the detail which is required in the CRF. In Table 4.1 aggregated estimates of the potential emissions from refrigeration and air conditioning (the main source of F-gases) are given.

The following protocols on [www.greenhousegases.nl](http://www.greenhousegases.nl) describe the methodologies applied for estimating emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and F-gases of Industrial processes in the Netherlands:

- Protocol 0043: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from Process emissions: fossil fuels;

- Protocol 0054: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from Process emissions and product use;
- Protocol 0055: N<sub>2</sub>O from Nitric acid production (2B2);
- Protocol 0056: N<sub>2</sub>O from Caprolactam production (2B5);
- Protocol 0057: PFCs from Aluminum production (2C3);
- Protocol 0058: HFC<sub>23</sub> from HCFC<sub>22</sub> production (2E1);
- Protocol 0059: HFCs from Handling (2E3);
- Protocol 0060: HFCs from Stationary refrigeration (2F1);
- Protocol 0061: HFCs from Mobile air conditioning (2F1);
- Protocol 0062: HFCs from Foams (2F2);
- Protocol 0063: HFCs from Aerosols (2F4);
- Protocol 0066: SF<sub>6</sub> from Electrical equipment (2F8);
- Protocol 0065: SF<sub>6</sub> and PFCs from Semiconductor manufacturing (2F7);
- Protocol 0064: SF<sub>6</sub> from Sound-proof windows (2F9).


#### Key sources

The key sources in this sector are presented in Table 4.2. Annex 1 presents all sources identified in the Industrial processes sector in the Netherlands. Nitric Acid production is from now on only a Tier 2 trend key source, due to the

Aggregated estimate of the potential emissions (Units: Gg CO<sub>2</sub> eq)

Table 4.1

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
HFC-134a	229	496	871	1,234	1,455	1,720	1,932	2,201	2,387	2,628	2,775	3,058	3,172	3,247	3,250
HFC-125		65	193	309	445	672	947	1,255	1,514	1,830	2,101	2,365	2,658	2,914	3,134
HFC-143a		44	149	260	400	598	855	1,101	1,340	1,619	1,870	2,106	2,365	2,605	2,812
HFC-32		30	101	155	209	222	319	348	370	404	435	473	517	537	520
Unspecified HFCs							1,238	2,153	3,049	3,981	4,881	5,747	6,574	7,358	8,135
Sum	229	634	1,314	1,958	2,508	3,213	5,291	7,058	8,660	10,462	12,063	13,749	15,286	16,660	17,850



reduction achieved in this category. Other key sources are CO<sub>2</sub> emissions from Ammonia production, CO<sub>2</sub> emissions from steel and aluminium production, HFC emissions from Substitutes for ozone-depleting substances and the production of HCFC-22. Also, the Manufacture of other chemical products is a key source for CO<sub>2</sub>.

#### Overview of shares and trends in emissions

Figure 4.1 and Table 4.2 show the trends in total greenhouse gas emissions from the sector Industrial processes.

In 2008, Industrial processes contributed 5% to the total national greenhouse gas emissions (without LULUCF) in comparison to 11% in the base year. The sector is a major source of N<sub>2</sub>O emissions in the Netherlands, accounting for 9% of the national total N<sub>2</sub>O emissions.

Category 2B Chemical industry contributes most to emissions from this sector. Compared to the base year, total CO<sub>2</sub> equivalent greenhouse gas emissions of the sector declined by 6.3 Tg to 4.8 Tg CO<sub>2</sub> eq in 2008 (-57%). CO<sub>2</sub> emissions from Industrial processes decreased 14% during the period 1990–

2008. N<sub>2</sub>O emissions decreased 85% in the same period. Total emissions of fluorinated gases (F-gases) have been strongly reduced.

In 2008, total greenhouse gas emissions in the sector decreased by 29% or 4.2 Tg CO<sub>2</sub> eq compared to 2007. CO<sub>2</sub> emissions decreased by 6%, HFC emissions showed an increase of 0.06 Tg CO<sub>2</sub> eq and PFC emissions decreased by 0.07 Tg CO<sub>2</sub> eq, while SF<sub>6</sub> emissions remained at the same level as the previous year. N<sub>2</sub>O emission decreased by 78% or 3.8 Tg CO<sub>2</sub> eq in 2008.

### 4.2 Mineral products [2A]

#### 4.2.1 Source category description

##### General description of the source categories

This category comprises emissions of greenhouse gases related to the production and use of non-metallic minerals in:

- 2A1 Cement clinker production: CO<sub>2</sub> emissions;
- 2A3 Limestone and dolomite use: CO<sub>2</sub> emissions;

Sector/category	Gas	Key	Emissions in base year	Emissions in 2007	Emissions in 2008	Change 2008 - 2007	Contribution to total in 2008 (%)		
			Level, Trend	Tg CO <sub>2</sub> eq	Tg CO2 eq	Tg CO <sub>2</sub> eq	Tg CO2 eq	By sector	Of total gas
2 Industry	CO <sub>2</sub>		7.9	7.2	6.7	-0.4		4%	3%
	CH <sub>4</sub>		0.3	0.3	0.3	0.0		1.7%	0.1%
	N <sub>2</sub> O		7.1	4.8	1.0	-3.8		9%	1%
	HFC		6.0	1.8	1.9	0.1		100%	0.9%
	PFC		1.9	0.3	0.3	-0.1		100%	0.1%
	SF6		0.3	0.2	0.2	0.0		100%	0.1%
	All		23.5	14.7	10.5	-4.2			5%
2A. Mineral Products	CO <sub>2</sub>		0.9	1.1	1.1	0.0	11%	0.6%	0.5%
2B. Chemical industry	CO <sub>2</sub>		3.7	3.6	3.5	-0.1	33%	2.0%	2%
	N <sub>2</sub> O		7.1	4.8	1.0	-3.8	10%	9%	0.5%
	All		11.1	8.7	4.8	-3.9	46%	2%	2%
2B1 Emissions from ammonia production	CO <sub>2</sub>	L1	3.1	3.0	2.9	-0.2	27%	2%	1%
2B2 Emissions from nitric acid production	N <sub>2</sub> O	T	6.3	4.3	0.6	-3.7	5%	5%	0.3%
2B5 Emissions from caprolactam production	N <sub>2</sub> O								
2B5 Other chemical product manufacture	CO <sub>2</sub>	L	0.6	0.6	0.6	0.0	6%	0.4%	0.3%
2C. Metal Production	CO <sub>2</sub>		2.9	2.1	1.8	-0.3	17%	1.0%	0.9%
	PFC		1.9	0.1	0.1	0.0	1%	29%	0.0%
	All		4.8	2.2	1.9	-0.3	18%		0.9%
2C1 Iron and steel production (carbon inputs)	CO <sub>2</sub>	L1,T1	2.5	1.6	1.3	-0.3	13%	0.8%	0.7%
2C3 PFC emissions from aluminium production	PFC	T	1.9	0.1	0.1	0.0	0.7%	29%	0.0%
2D. Other Production	CO <sub>2</sub>		0.1	0.0	0.0	0.0	0.0%	0.0%	0.0%
2E. Production of halocarbons and SF <sub>6</sub>	HFC		5.8	0.3	0.2	0.0	2%	12%	0.1%
2E1 HFC-23 emissions from HCFC-22 manufacture	HFC	T	5.8	0.2	0.2	0.0	2%	11%	0.1%
2F. Consumption of Halocarbons and SF6	HFC		0.2	1.6	1.7	0.1	16%	88%	0.8%
	PFC		0.0	0.2	0.2	0.0	2%	71%	0.1%
	SF6		0.3	0.2	0.0	-0.2	0%	0%	0.0%
	All		0.6	2.0	2.1	0.1	20%		1.0%
	CO <sub>2</sub>		0.2	0.3	0.3	0.0	3%	0.2%	0.2%
2G. Other	N <sub>2</sub> O		0.0	0.0	0.0	0.0	0%	0%	0.0%
	All			0.3	0.3	0.0	3%	0.2%	0.2%
Total National emissions (excl. CO <sub>2</sub> LUCF)	CO <sub>2</sub>		159.3	172.3	175.7	3.4			
	CH <sub>4</sub>		25.5	16.8	17.1	0.2			
	N <sub>2</sub> O		20.2	15.4	11.8	-3.7			
	HFCs		6.0	1.8	1.9	0.1			
	PFCs		1.9	0.3	0.3	-0.1			
	SF6		0.3	0.2	0.2	0.0			
	All		213.3	206.9	206.9	0.0			

- 2A4 Soda ash production and use: CO<sub>2</sub> emissions;
- 2A7 Other (the production of glass and other production and use of minerals): CO<sub>2</sub> emissions.

CO<sub>2</sub> emissions from 2A5 Asphalt roofing and 2A6 Road paving with asphalt are not estimated. For more information see Annex 5.

#### Overview of shares and trends in emissions

Total CO<sub>2</sub> emissions in category 2A increased from 0.9 Tg in 1990 to 1.1 Tg in 2008 (see Table 4.2). The increased emissions during the period 1990-2008 are related to the increased production levels during that period.

#### Activity data and (implied) emission factors

Detailed information on activity data and emission factors can be found in the monitoring protocols on the website [www.greenhousegases.nl](http://www.greenhousegases.nl)

Activity data are based on the following sources:

- Cement clinker production: the environmental reports (MJVs) of the single Dutch company are used.
- Limestone and dolomite use: environmental reports are used for emission data. Activity data on plaster production for use in desulphurising installation for power plants are based on the environmental reports of the coal-fired power plants. Data on the consumption of limestone and



dolomite are based on statistical information obtained from Statistics Netherlands (CBS) and can be found on the website [www.cbs.nl](http://www.cbs.nl).

- Soda ash production and use: the environmental reports for data on the non-energy use of coke are used. For activity data on soda use, see following bullet Glass production;
- Glass production: activity data are based on data from Statistics Netherlands (CBS) and the trade organisation.

The following emission factors (EF) are used to estimate the CO<sub>2</sub> emissions from the different source categories:

- Cement clinker production: Because of changes in raw material composition it is not possible to estimate reliable CO<sub>2</sub> process emissions by calculating the clinker production (as AD) by a default EF. For that reason the company has chosen to base the calculation of CO<sub>2</sub> emissions on the carbonate content of the process input. For more information, see Section 4.2.2
- Limestone use: EF= 0.440 t/t (IPCC default);
- Dolomite use: EF= 0.477 t/t (IPCC default);
- Soda ash production: EF= 0.415 t/t (IPCC default);
- Glass production: Plant-specific EFs have been used for the years 1990 (0.13 t CO<sub>2</sub> /t glass), 1995 (0.15 t CO<sub>2</sub> /t glass) and 1997 (0.18 t CO<sub>2</sub> /t glass). For other years in the time series, there were not enough data available to calculate plant-specific EFs. For the missing years 1991-1994 and 1996, EFs have been estimated by interpolation. Because no further measurement data are available, the emission factor for 1998– 008 is kept at the same level as the EF of 1997 (0.18 t CO<sub>2</sub> /t glass).

#### 4.2.2 Methodological issues

For all the source categories, country-specific methodologies are used to estimate emissions of CO<sub>2</sub>, in compliance with the IPCC Good Practice Guidance (IPCC, 2001). More detailed descriptions of the methods used and emission factors are found in Protocols 0043 and 0054 on [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 4.1.

- 2A1 Cement clinker production: the CO<sub>2</sub> emissions from this source category are based on (measured) data reported by the single company in the Netherlands that produces clinkers. The methodology for measurements and for calculating emissions can be described as follows:
- The first carbonate input in the kiln is the raw material. The CO<sub>2</sub> emission is calculated on a monthly basis by multiplying the amount of raw material by a derived process EF. From every batch in a month a sample is taken just before the raw material is fed to the kiln. The process EFs and composition data for batches of raw material are determined in a laboratory. The EF is determined by measuring the weight loss of the sample (excluded the amount of organic carbon). The monthly EF is set as the average of all sample EFs determined that month. The second carbonate input in the kiln is sewage sludge. Also the CO<sub>2</sub> emission from this source is calculated monthly by multiplying the amount of sewage sludge by the monthly derived process EF. Besides the CO<sub>2</sub> emissions resulting from calcination of the carbonate input in the kiln, the company considers the CO<sub>2</sub> emission from the burning off the small amount of organic carbon in the raw material as a process emission. As a result, the total yearly process

emissions of the company is the sum of all monthly emissions of the following sources:

- CO<sub>2</sub> from the calcination of the carbonate input of the raw material, marl;
- CO<sub>2</sub> from the calcination of the carbonate input of sewage sludge;
- CO<sub>2</sub> from the burning of organic carbon in the raw material.
- Emission data obtained from the environmental report related to clinker production figures give an implied emission factor of 0.48 – 0,54 t/t clinker (IPCC Default = 0.51 t/t clinker).
- There is a monitoring protocol applied for emissions trading (this protocol is approved by the Netherlands Emission authority (NEa); the Government organisation responsible for emission trading (ETS) in the Netherlands) This organisation is also responsible for the verification of the reported data from the company that produces clinkers.
- 2A3 Limestone and dolomite use: the CO<sub>2</sub> emissions from this source category are based on consumption figures for limestone use – derived from plaster production figures – for flue gas desulphurisation (FGD) with a wet process by coal-fired power plants and for apparent dolomite consumption (mostly used for road construction). No activity data are available to estimate other sources of limestone and dolomite use.
- 2A4 Soda ash production and use: only one company in the Netherlands is producing soda ash using the Solvay process. CO<sub>2</sub> emissions are calculated based on the non-energy use of coke, assuming the 100% oxidation of carbon.
- 2A7 Other: CO<sub>2</sub> emissions from this source category refer to Glass production. Emissions are estimated based on gross glass production data and a country-specific emission factors.

#### 4.2.3 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of uncertainties according to IPCC source category.

Uncertainty estimates used in the Tier 1 analysis are based on the judgment of experts since no detailed information is available for assessing the uncertainties of the emissions reported by the facilities (Cement clinker production, Limestone and dolomite use and Soda ash production). The uncertainty in CO<sub>2</sub> emissions from cement production is estimated to be approximately 10% in annual emissions; for Limestone/dolomite use and other sources the uncertainty is estimated to be 25%, based on the relatively high uncertainty in the activity data.

Activity data for Soda ash use, Glass production and Limestone and dolomite use are assumed to be relatively uncertain (25%). The uncertainties of the IPCC default emission factors used for some processes are not assessed. However, as these are, minor sources for CO<sub>2</sub> this was not given any further consideration.



#### Time-series consistency

Consistent methodologies have been applied for all source categories. The time series involve a certain amount of extrapolation with respect to the activity data for *Soda ash use*, thereby introducing further uncertainties in the first part of the time series of this source.

#### 4.2.4 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedure discussed in Chapter 1.

#### 4.2.5 Source-specific recalculations

No source-specific recalculations were performed in the preparation of this NIR.

#### 4.2.6 Source-specific planned improvements

There are no source-specific improvements planned.

### 4.3 Chemical industry [2B]

#### 4.3.1 Source category description

The national inventory of the Netherlands comprises emissions of greenhouse gases related to four source categories as belonging to this category:

- 2B1 Ammonia production: CO<sub>2</sub> emissions: in the Netherlands, natural gas is used as feedstock for ammonia production. CO<sub>2</sub> is produced as a by-product during the chemical separation of hydrogen from the natural gas. During the process of ammonia (NH<sub>3</sub>) production, hydrogen and nitrogen are combined to react together to manufacture ammonia. Only prompt process emissions from the ammonia/urea production are included in this source category. Emissions from the use of urea in domestic agricultural activities are included in category 5C (see Chapter 7).
- 2B2 Nitric acid production: N<sub>2</sub>O emissions: the production of nitric acid (HNO<sub>3</sub>) generates nitrous oxide (N<sub>2</sub>O) as a by-product of the high-temperature catalytic oxidation of ammonia.
- 2B4 Carbide production: CH<sub>4</sub> emissions: petrol cokes are used during the production of silicon carbide; the volatile compounds in the petrol cokes form CH<sub>4</sub>.
- 2B5 CO<sub>2</sub> and N<sub>2</sub>O emissions from Other chemical product manufacture:
  - Industrial gas production: hydrogen and carbon monoxide are produced mainly from natural gas used as chemical feedstock, but they can also be produced from petroleum coke and coke, during which processes CO<sub>2</sub> is produced.
  - Carbon electrode production: carbon electrodes are produced from petroleum coke and coke used as feedstock, during which processed CO<sub>2</sub> is produced.
  - Activated carbon production: Norit is one of world's largest manufacturers of activated carbon, for which peat is used as carbon source and CO<sub>2</sub> is produced as by-product.
  - Caprolactam production: N<sub>2</sub>O emissions result from the production of caprolactam.
  - Ethylene oxide production: CO<sub>2</sub> emissions result from the production of ethylene oxide.

Adapic acid (2B3) and calcium carbide (included in 2B4) are not produced in the Netherlands. CO<sub>2</sub> emissions resulting from the use of fossil fuels as feedstocks for the production of silicon carbide, carbon black, ethylene and methanol are included in the Energy sector (1A1a and 1A2c; see Sections 3.2.1. and 3.3.1. for more details).

#### 4.3.2 Key sources

Ammonia production and Other chemical product manufacture are identified as key-sources for CO<sub>2</sub> emissions. Nitric acid production is no longer a Tier 2 level key source for N<sub>2</sub>O emissions. Due to the emission reduction, it is devaluated to a trend key source (see Table 4.2).

#### 4.3.3 Overview of shares and trends in emissions

Figure 4.2 shows the trend in CO<sub>2</sub> equivalent emissions from 2B 'Chemical industry' in the period 1990–2008. Table 4.2 gives an overview of shares in emissions of the main categories.

Emissions from this category contributed 5% to the total national greenhouse gas emissions (without LULUCF) in the base year and 2% in 2008. Nitric acid production is the most important source of N<sub>2</sub>O emissions from industrial processes in the Netherlands. The contribution of N<sub>2</sub>O emissions from 2B 'Chemical industry' was 3% of the total national greenhouse gas emission inventory in the base year and 0.5% in 2008.

From 1990 to 2008, total greenhouse gas emissions in 2B 'Chemical industry' decreased by 57%, mainly due to reduction of N<sub>2</sub>O emissions from the production of nitric acid. In 2008, total greenhouse gas emissions in 2B 'Chemical industry' decreased by 45% or 3.9 Tg CO<sub>2</sub> eq.

Table 4.3 shows that N<sub>2</sub>O emissions from the chemical industry remained rather stable between 1990 and 2000 – when there was no policy aimed at controlling these emissions.

From the 2002 submissions onwards, the N<sub>2</sub>O emission from the nitric acid production is based on measurements.

Until 2002, N<sub>2</sub>O emissions from nitric acid production were based on default IPCC emission factors. N<sub>2</sub>O emission measurements made in 1998 and 1999 have resulted in new emission factors. Because no measures have been taken and the operation conditions did not change during the period 1990–1998, the emission factors obtained from the measurements have been used to recalculate the emissions for the period 1990–1998. Technical measures implemented at one of the nitric acid plants in 2001 resulted in an emission reduction of 9% compared to 2000. The decreased emission level in 2002 compared to 2001 is related to the decreased production level of nitric acid in that year. In 2003, emissions and production did not fluctuate, whereas in 2004 the increased emission level is once again related to the marked increase in production. In 2005 and 2006, the N<sub>2</sub>O emissions of the nitric acid plants remained almost at the same level as in 2004. Technical measures implemented at all nitric acid plants in the third quarter of 2007 resulted in an emission reduction of 23% compared to 2006. In 2008, the full effect of the measures is reflected in the low emission.


Trend in N<sub>2</sub>O emissions from Chemical industry processes (2B) (Units: Gg CO<sub>2</sub> eq)

Table 4.3

	1990	1995	2000	2005	2006	2007	2008
B2. Nitric acid production	6,330	6,278	5,898	5,659	5,597	4,305	558
B5. Other	766	805	936	705	662	497	481
Total	7,096	7,083	6,834	6,364	6,259	4,802	1,039

After 2002, more accurate measurements were performed to estimate N<sub>2</sub>O emissions from caprolactam production (2B5). Calculations of the pre-2003 emissions are based on a production-index series (real production data are confidential business information) over the period 1990-2004 and the 2003 and 2004 measurements from the company.

The decreased emission level of the caprolactam plant in 2005 compared to 2004 is related to the decreased production level in that year. In 2006, the N<sub>2</sub>O emissions of the caprolactam plant remained almost at the same level as in 2005. A better process control and a lower production level resulted in an emission reduction of 25% in 2007 compared to

2006. In 2008, the emissions remained at the same level as in 2007.

#### 4.3.4 Activity data and (implied) emission factors

Detailed information on activity data and emission factors can be found in monitoring protocols 0043, 0054, 0055 and 0056 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Activity data are based on the following sources:

- Ammonia production: activity data on use of natural gas are obtained from Statistics Netherlands (CBS).
- Nitric acid production: activity data are confidential. Emissions are reported by the companies.

- Carbide production: silicon carbide production figures are derived from the Environmental Report (MJV) of the relevant company.
- Other: activity data on caprolactam production are confidential. Only emissions are reported by the companies. This year a production-index series over the period 1990-2005 were received from the company. For Ethylene oxide production only capacity data are available; therefore, a default capacity utilisation rate of 86% is used to estimate CO<sub>2</sub> emissions (based on Neelis et al., 2005). Activity data for estimating CO<sub>2</sub> emissions are based on data for feedstock use of fuels provided by Statistics Netherlands (CBS).

The emission factors used to estimate greenhouse gas emissions from the different source categories are based on:

- Ammonia production: a country-specific CO<sub>2</sub> emission factor is used. This emission factor is based on a 17% fraction of the carbon in the gas-feedstock not being oxidised during the ammonia manufacture and was calculated from the carbon contained in the urea produced (based on Neelis et al., 2003).
- Nitric acid production: plant-specific N<sub>2</sub>O emission factors are used (which are confidential).
- Silicon carbide production: the IPCC default emission factor is used for CH<sub>4</sub>.
- Other: plant-specific N<sub>2</sub>O emission factors are used for Caprolactam production (confidential). A default emission factor of 0.45 tons CO<sub>2</sub> per ton of ethylene oxide production is used. Country-specific CO<sub>2</sub> emission factors are used to estimate the CO<sub>2</sub> emissions of the other source categories because no IPCC methodologies exist for these processes. For activated carbon an emission factor of 1 t/t Norit derived from the carbon losses from peat uses is used.

#### 4.3.5 Methodological issues

For all the source categories of the chemical industry, the methodologies used to estimate the greenhouse gas emissions are in compliance with the IPCC Good Practice Guidance (IPCC, 2001). Country-specific methodologies are used for the CO<sub>2</sub> process emissions from the chemical industry. More detailed descriptions of the methods used and emission factors can be found in the protocols (0043, 0054, 0055 and 0056) described on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 4.1:

- 2B1 Ammonia production: a method equivalent to IPCC Tier 1b; the amount of natural gas used as feedstock and a country-specific emission factor are used to estimate CO<sub>2</sub> emissions. This emission factor is based on the assumption that the fraction of carbon in the gas-feedstock oxidised during the ammonia manufacture is 17%. This figure is based on reported carbon losses from urea production (Neelis et al., 2003).
- 2B2 Nitric acid production: an IPCC Tier 2 method is used to estimate N<sub>2</sub>O emissions. The emission factors are based on plant-specific measured data which are confidential. The emissions are based on data reported by the nitric acid manufacturing industry and are included in the national Pollutant Release & Transfer Register (PRTR).
- 2B5 Other chemical products: N<sub>2</sub>O emissions from 2B5 Other chemical industry, which mainly originate from

Caprolactam production, are also based on emission data reported by the manufacturing industry (based on measurements). Emission factors and activity data are confidential. The aggregated CO<sub>2</sub> emissions included in this source category are identified as a key source and based on country-specific methods and emission factors. These refer to the production of:

- Industrial gases: CO<sub>2</sub> emissions are estimated based on use of fuels (mainly natural gas) as chemical feedstock. An oxidation fraction of 20% is assumed, based on reported data in environmental reports from the relevant facilities.
- Carbon electrodes: CO<sub>2</sub> emissions are estimated based on fuel use (mainly petroleum coke and coke). A small oxidation fraction – 5% – is assumed, based on reported data in the environmental reports.
- Activated carbon: CO<sub>2</sub> emissions are estimated on the basis of the production data for Norit and by applying an emission factor of 1 t/t Norit. The emission factor is derived from the carbon losses from peat uses reported in the environmental reports. As peat consumption is not included in the national energy statistics, the production data since 1990 have been estimated based on an extrapolation of production level of 33 Tg reported in 2002. This is considered to be justified because this source contributes relatively little to the national inventory of greenhouse gases.
- Ethylene oxide: CO<sub>2</sub> emissions are estimated based on capacity data by using a default capacity utilisation rate of 86% and applying an emission factor of 0.45 t/t ethylene oxide.

For the minor sources of CH<sub>4</sub> emissions included in this source category, IPCC Tier 1 methodologies and IPCC default emission factors are used.

#### 4.3.6 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Table A7.1 and A7.2 provides estimates of uncertainties according to IPCC source categories.

No accurate information is available for assessing the uncertainties of the emissions reported by the facilities (i.e. Ammonia, Nitric acid, Caprolactam production). Activity data are assumed to be relatively certain. The uncertainties in CO<sub>2</sub> emissions from Ammonia production and Other chemical products are estimated to be approximately 2% and 50%, respectively, in annual emissions. The uncertainty in the annual emissions of N<sub>2</sub>O from Nitric acid production and Caprolactam production is estimated to be approximately 20%.

##### Time-series consistency

Consistent methodologies are used throughout the time series for the sources in this category.

#### 4.3.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures as discussed in Chapter 1.

Implied emission factors for CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub> from aluminium production (Units: kg/Tg) (2C3)

Table 4.4

Gas	1990	1995	2000	2005	2006	2007	2008
CF <sub>4</sub>	1.02	1.10	0.53	0.03	0.03	0.04	0.03
C <sub>2</sub> F <sub>6</sub>	0.18	0.18	0.12	0.01	0.00	0.01	0.00

Although ammonia and urea production data are considered confidential, international statistics such as UN, IFA and USGS do report production data for the Netherlands.

#### 4.3.8 Source-specific recalculations

There have been no source-specific recalculations in comparison to the previous submission.

#### 4.3.9 Source-specific planned improvements

There are no source-specific improvements planned.

### 4.4 Metal production [2C]

#### 4.4.1 Source category description

The national inventory of the Netherlands comprises emissions of greenhouse gases related to three source categories as belonging to 2C Metal production:

- 2C1 Iron and steel production: CO<sub>2</sub> emissions: The Netherlands has one integrated iron and steel plant (Corus, previously Hoogovens). During the production of iron and steel, coke and coal are used as reducing agents in the blast and oxygen furnaces, resulting in the production of the by-products blast furnace gas and oxygen furnace gas. A small part of these gases are emitted (lost) and the rest are subsequently used as fuels for energy purposes. Only the carbon losses are reported in category 2C1. In addition, CO<sub>2</sub> is produced as by-product from the use of limestone and during the conversion from pig iron to steel. These emissions are also reported in this category.
- 2C3 Aluminium production: CO<sub>2</sub> and PFC emissions: in the Netherlands aluminium is produced at two primary aluminium smelters (Pechiney and Aludel). CO<sub>2</sub> is produced by the reaction of the carbon anodes with alumina and by the reaction of the anode with other sources of oxygen (especially air).
- The PFCs (CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>) from the Aluminium industry are formed during the phenomenon known as the 'anode effect' (AE), which occurs when the concentration of aluminium oxide in the reduction cell electrolyte drops below a certain level.

2C2 Ferroalloys production and 2C4 Magnesium and aluminium foundries, both of which use SF<sub>6</sub> as a cover gas, do not occur in the Netherlands. No other sources of metal production (2C5) are identified in the inventory.

#### 4.4.2 Key sources

Iron and steel production (carbon inputs) is identified as a key source for CO<sub>2</sub> emissions, Aluminium production as a key source for PFC emissions (see Table 4.2).

#### 4.4.3 Overview of shares and trends in emissions

Table 4.2 gives an overview of shares in emissions of the main categories.

Total CO<sub>2</sub> emissions from 2C1 'Iron and steel production' decreased by 1.1 Tg during the period 1990–2008. In 2008, CO<sub>2</sub> emissions remained at the same level as in 2007.

PFC emissions from primary 'Aluminium industry' (2C3) decreased by 1.8 Tg CO<sub>2</sub> eq between 1995 and 2008. Because in 2008 the number of anode-effects decreased the PFC emissions decreased compared to 2007.

Table 4.4 shows the trend in implied CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub> emission factors (IEF) for aluminium production during the period 1990–2008. The largest company produces approximately two-thirds of the national total production. The IEFs decreased by 97% between 1995 and 2008. In 1998, the smallest company switched from side feed to point feed; this switch was followed by the larger company in 2002/2003, thereby explaining the decreased IEF from this year onwards. The higher level of the IEF in 2002 is caused by specific process-related problems during the switching process by the larger producer.

#### 4.4.4 Activity data and (implied) emission factors

Detailed information on activity data and emission factors can be found in the monitoring protocols 0043, 0054 and 0057 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl)

Activity data are based on the following sources:

- Iron and steel production: data on coke production and coal input, limestone use and the carbon balance are reported by the relevant company (by means of an environmental report);
- Aluminium production: activity data and emissions are based on data reported in the environmental reports of both companies.

Emission factors used in the inventory to estimate greenhouse gas emissions are based on:

- Iron and steel production: EF (limestone use) = 0.440 tons CO<sub>2</sub> per ton (IPCC default); EF (blast furnace gas) = 0.21485 tons CO<sub>2</sub> per GJ (plant specific);
- Aluminium production: EF (consumption of anodes) = 0.00145 tons CO<sub>2</sub> per ton aluminium (plant specific; IPCC default = 0.0015 t/t aluminium).

EF for PFCs is plant-specific and confidential. Emissions of PFCs are obtained from the environmental reports of both companies.

#### 4.4.5 Methodological issues

The methodologies used to estimate the greenhouse gas emissions for all source categories of metal production are in compliance with the IPCC Good Practice Guidance (IPCC, 2001). More detailed descriptions of the methods used and emission factors are found in protocols 0043, 0054 and 0057 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 4.1.

### Iron and steel production (2C1):

CO<sub>2</sub> emissions are estimated using a Tier 2 IPCC method and country-specific value for the carbon contents of the fuels. Carbon losses are calculated from coke and coal input used as reducing agents in the blast and oxygen furnaces, including other carbon sources such as limestone and the carbon contents in the iron ore (corrected for the fraction that ultimately remains in the steel produced):

- CO<sub>2</sub> from coke/coal inputs = amount of coke \* EF<sub>coke</sub> + amount of coal \* EF<sub>coal</sub> – (blast furnace gas + oxygen oven gas produced) \* EF<sub>BFGas</sub> (1a)
- CO<sub>2</sub> from limestone use = limestone use \* ZF(limestone) \* EF<sub>limestone</sub> (1b)
- CO<sub>2</sub> from ore/steel = (C-mass in ore, scrap and raw iron purchased – C-mass in raw steel)\* 44/12 (1c)
- The same emission factors for blast furnace gas and oxygen furnace gas are used (see Annex 2).

As mentioned above, only the carbon losses are reported in category 2C1. The carbon contained in the blast furnace gas and oxygen furnace gas produced as by-products and subsequently used as fuels for energy purposes is subtracted from the carbon balance and included in the Energy sector (1A1a and 1A2a; see Sections 3.2.2 and 3.2.3).

Data reported in the annual environmental reports (2000–2005) of Corus are used to calculate the CO<sub>2</sub> emissions from limestone use and the conversion from pig iron to steel in the period 1990–2000. The amount of limestone stone was calculated from the average consumption of 107.9 kg (lower: 107,1 kg; upper: 109.0 kg) in 2000–2005 per ton of crude steel produced. A similar calculation was made for the CO<sub>2</sub> from the carbon fractions in ore and crude steel. The CO<sub>2</sub> emission was calculated from the average CO<sub>2</sub> emission of 8.3 kg (lower: 6.0 kg; upper: 11.9 kg) in 2000–2005 per ton of crude steel produced.

### Aluminium production (2C3)

A Tier 1a IPCC method (IPCC, 2001) is used to estimate CO<sub>2</sub> emissions from the anodes used in the primary production of aluminium, with aluminium production being as activity data. In order to calculate the IPCC default emission factor the stoichiometric ratio of carbon needed to reduce the aluminium ore to pure aluminium is based on the reaction  $Al_2O_3 + 3/2C \rightarrow 2Al + 3/2 CO_2$ . This factor is corrected to include additional CO<sub>2</sub> produced by the reaction of the carbon anode with oxygen in the air. A country-specific emission factor of 0.00145 tons CO<sub>2</sub> per ton aluminium is used to estimate CO<sub>2</sub> emissions, and it has been verified that this value is within the range of the IPCC factor of 0.0015 and the factor of 0.00143 calculated by the World Business Council for Sustainable Development (WBCSD) (WBCSD/WRI, 2004). PFC emissions from primary aluminium production reported by these two facilities are based on the IPCC Tier 2 method for the complete period 1990–2008. Emission factors are plant-specific and are based on measured data.

### 4.4.6 Uncertainties and time-series consistency

#### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Table A7.1 and A7.2 provides estimates of uncertainties according to IPCC source categories.

No accurate information is available for assessing the uncertainties of the emissions reported by the facilities. Activity data are assumed to be relatively certain. The uncertainties in CO<sub>2</sub> emissions from iron and steel production (2C1) and aluminium production (2C3) are estimated to be approximately 6% and 5%, respectively, in annual emissions.

#### Time-series consistency

Consistent methodologies are used throughout the time series for the sources in this category.

### 4.4.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures as discussed in Chapter 1.

### 4.4.8 Source-specific recalculations

There have been no source-specific recalculations in comparison to the previous submission.

### 4.4.9 Source-specific planned improvements

There are no source-specific improvements planned.

### 4.4.10 Uncertainties and time-series consistency

#### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of uncertainties according to IPCC source category. The uncertainty in annual CO<sub>2</sub> emissions is estimated to be approximately 6% and 5% for Iron and steel production and Aluminium production respectively, whereas the uncertainty in PFC emissions from Aluminium production is estimated to be 20%. The uncertainty in the activity data is estimated at 2% for Aluminium production and 3% for Iron and steel production. The uncertainty in the emission factors for CO<sub>2</sub> is estimated at 5% and for PFC from Aluminium production at 20%.

#### Time-series consistency

The time series are based on consistent methodologies for the sources in this category. PFC emissions from the production of aluminium by the main company during the period 1990–1998 are based on the extrapolation of measured data from 1999, thereby increasing the uncertainties of the emissions during that period. It is assumed, however, that the emission factors reflect the plant specific circumstances better than the default emission factors used in previous reporting.

### 4.4.11 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures discussed in Chapter 1.

### 4.4.12 Source-specific recalculations

There are no source-specific improvements planned.



#### 4.4.13 Source-specific planned improvements

There were no source-specific recalculations in this submission.

### 4.5 Food and drink production [2D]

#### 4.5.1 Source category description

This category comprises CO<sub>2</sub> emissions related to food and drink production in the Netherlands.

CO<sub>2</sub> emissions in this source category are related to the non-energy use of fuels. Carbon is oxidised during these processes, resulting in CO<sub>2</sub> emissions.

#### 4.5.2 Key sources

This minor source is no key source for CO<sub>2</sub>

#### 4.5.3 Overview of shares and trends in emissions

Emissions vary at around 0.05 Tg, and are rounded off to either 0.1 or 0.0 Tg (see Table 4.2).

#### 4.5.4 Activity data and (implied) emission factors

Detailed information on the activity data and emission factors can be found in monitoring protocol 0043 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

The activity data used to estimate CO<sub>2</sub> emissions from this source are based on national energy statistics from Statistics Netherlands (CBS) on Coke consumption. Emission factors are derived from the national default carbon content of coke (Corus, MJVs 2000-2008).

#### 4.5.5 Methodological issues

The methodology used to estimate the greenhouse gas emissions complies with the IPCC Good Practice Guidance (IPCC, 2001). More detailed descriptions of the method used and the emission factors can be found in protocol 0043 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 4.1.

CO<sub>2</sub> emissions are calculated based on the non-energy use of fuels by the food and drink industry as recorded in the national energy statistics, multiplied by an emission factor. The emission factor is based on the national default carbon contents of the fuels (see Annex 2), under the assumption that the carbon is fully oxidised to CO<sub>2</sub>.

#### 4.5.6 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of the uncertainties according to the IPCC source category. The uncertainty in the emissions of this category is estimated to be 5%. Since this is a very small emission source, the uncertainties in this category are not analysed further in more detail. Therefore, in the uncertainty analysis and the key source analysis the emissions in this category (2D) are combined with the emissions in category 2G (Other industrial emissions), see Section 4.8.

#### Time-series consistency

The time series is based on consistent methodologies and activity data for this source.

#### 4.5.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures which are discussed in Chapter 1.

#### 4.5.8 Source-specific recalculations

During the compilation of the CRF an error in the 1990 emission of CO<sub>2</sub> was detected and corrected.

#### 4.5.9 Source-specific planned improvements

There are no source-specific improvements planned.

### 4.6 Production of halocarbons and SF<sub>6</sub> [2E]

#### 4.6.1 Source category description

The national inventory of the Netherlands comprises emissions of greenhouse gases related to the following source categories in this category:

- 2E1 Production of HCFC-22: HFC-23 emissions.
- HCFC-22 is produced at one plant in the Netherlands. Tri-fluoromethane (HFC-23) is generated as a by-product during the production of chlorodifluoromethane (HCFC-22) and emitted through the plant condenser vent.
- 2E3 Handling activities: emissions of HFCs. There is one company in the Netherlands that repackages HFCs from large units (e.g., containers) into smaller units (e.g., cylinders) and in addition trading with HFCs. Besides this company there are a lot of companies in the Netherlands which are importing small units with HFCs and sell them in the trading areas.

#### 4.6.2 Key sources

Production of HCFC-22 (HFC-23 emission) is a key source; see Table 4.2.

#### 4.6.3 Overview of shares and trends in emissions

Table 4.2 gives an overview of shares in emissions of the main categories.

Total HFC emissions in category 2E were 5.8 Tg in 1995 and 0.2 Tg CO<sub>2</sub> eq in 2008, with HFC-23 emissions from HCFC-22 production being the major source of HFC emissions. HFC emissions from handling contributed 8% to the total HFC emissions from this category in 2008.

Table 4.5 shows the trend in HFC emissions from the categories HCFC-22 production and HFCs from handling activities for the period 1990–2008. The emissions of HFC-23 increased about by 35% in the period 1995–1998 due to the increased production of HCFC-22. However, in the period 1998–2000, the emissions of HFC-23 decreased by 69% following the installation of a thermal afterburner at the plant.

The operation time of the thermal afterburner (84% in 2000; 95% in 2001; 93.6% in 2002) is the primary factor explaining the variation in emission levels during the period 2000–2002. The decreased emission (33%) in 2003 can be explained mostly

	1990	1995	2000	2005	2006	2007	2008
2E1. HFC-23	4,432	5,759	2,421	196	281	243	212
2E3. HFCs	NO	12	418	39	37	24	18
HFC Total	4,432	5,771	2,838	235	318	267	230

by a lower production level. Despite a higher production level, emissions have remained stable because the operation time of the thermal afterburner increased from 92% in 2003 to 96% in 2004. The decreased emission (45%) in 2005 can be explained by a higher operation time of the thermal afterburner (97.1% in 2005) and a lower production level. Because of a higher production level, emission increased (40%) in 2006. The decreased emission (16%) in 2007 can be explained by a higher operation time of the thermal afterburner.

The significant emission fluctuations in category 2.E.3 during the period 1992-2008 can be explained by the large variety in handling activities, which depends on the demand of the costumers.

#### 4.6.4 Activity data and (implied) Emission factors

The activity data used to estimate emissions of F-gases from this category are based on confidential information provided by the manufacturers:

- Production of HCFC-22: production figures on HCFC-22 are confidential.
- Handling activities (HFCs): activity data used to estimate HFC emissions are confidential.

(Implied) emission factors used to estimate the emissions of F-gases from this category are based on the following:

- Production of HCFC-22: Destruction factor of the thermal afterburner used is 99.99%.
- Handling activities (HFCs): the emission factors used are plant-specific and confidential, and they are based on 1999 measurement data. More detailed information on the activity data and emission factors can be found in the monitoring protocols 0058 and 0059 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

#### 4.6.5 Methodological issues

The methodologies used to estimate the greenhouse gas emissions included in this category are in compliance with the IPCC *Good Practice Guidance* (IPCC, 2001). More detailed descriptions of the method used and emission factors can be found in the protocols 0058 and 0059 on website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 4.1:

- Production of HCFC-22 (2E1)*: this source category is identified as a key source for HFC-23 emissions. In order to comply with the IPCC Good Practice Guidance (IPCC, 2001), an IPCC Tier 2 method is used to estimate the emissions of this source category. HFC-23 emissions are calculated using both (measured) data obtained on the mass flow of HFC-23 produced in the process and a destruction factor to estimate the reduction of this HFC-23 flow by the afterburner.

- Handling activities (HFCs) (2E3)*: Tier 1 country-specific methodologies are used to estimate the handling emissions of HFCs. The estimations are based on emissions data reported by the manufacturing and sales companies.

#### 4.6.6 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of uncertainties according to the IPCC source category.

The uncertainty in HFC emissions from HCFC-22 production is estimated to be about 15%, while the uncertainty in HFC emissions from handling activities is estimated to be about 20%. The uncertainty in the activity data for these sources is estimated at 10%. The uncertainties in the emission factors for HFC-23 from HCFC-22 production and for HFC from Handling activities are estimated at 10% and 20%, respectively. These figures are all based on the judgments of experts.

##### Time-series consistency

The time series is based on consistent methodologies and activity data for this source.

#### 4.6.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures discussed in Chapter 1.

#### 4.6.8 Source specific recalculations

There have been no source-specific recalculations compared to the previous submission.

#### 4.6.9 Source-specific planned improvements

There are no source-specific improvements planned for this category.

### 4.7 Consumption of halocarbons and SF<sub>6</sub> [2F]

#### 4.7.1 Source category description

Halocarbons and SF<sub>6</sub> are released from the use of these compounds in different products. The national inventory of the Netherlands comprises emissions of greenhouse gases related to the following source category: 2F(1-9): Emissions from substitutes for Ozone-depleting substances.

The inventory comprises the following sources in this source category:

- 2F1. Stationary refrigeration: HFC emissions;
- 2F1. Mobile air conditioning: HFC emissions;
- 2F2. Foam Blowing: HFC emissions; (included in 2F6);



	1990	1995	2000	2005	2006	2007	2008
HFC-134a	NO	45	204	433	469	495	511
HFC-143a	NO	7	110	294	331	367	400
HFC-125	NO	8	90	243	274	303	329
HFC-152a	NO	NO	NO	NO	NO	NO	NO
HFC-32	IE,NO	0.80	7.00	11.44	12.48	13.30	13.35
Other HFCs	NO	187	642	297	327	399	439
HFC Total	IE,NO	247	1,053	1,279	1,412	1,578	1,693
PFC use	18	37	193	178	194	222	180
SF <sub>6</sub> use	217	301	318	254	217	226	224
Total HFC/PFC/SF <sub>6</sub>	236	585	1,564	1,711	1,823	2,026	2,097

- 2F4. Aerosols/Metered dose inhalers: HFC emissions; (included in 2F6);
- 2F6. Other applications using ODS substitutes
- 2F7. Semiconductor manufacture : PFC emissions (SF<sub>6</sub> emissions included in 2F9);
- 2F8. Electrical equipment: SF<sub>6</sub> emissions (included in 2F9);
- 2F9. Other: SF<sub>6</sub> emissions from Sound-proof windows and Electron microscopes

In the Netherlands, many processes related to the use of HFCs and SF<sub>6</sub> takes place in only one or two companies. Because of the sensitivity of data from that companies, only the sum of the HFC emissions of 2F2 and 2F4 (included in 2F6) and of the SF<sub>6</sub> emissions of 2F7 and 2F8 is reported (included in 2F9).

#### 4.7.2 Key sources

Emissions from Substitutes for ozone-depleting substances [2F] are identified as a key source (see Table 4.2).

#### 4.7.3 Overview of shares and trends in emissions

The contribution of F-gas emissions from category 2F to the total national inventory of F-gas emissions was 4% in the base year 1995 and 88% in 2008. This corresponds to 1.6 Tg CO<sub>2</sub> eq and accounts for 0.8% in the national total greenhouse gas emissions in 2008.

The level of HFC emissions increased by a factor of 7 in 2008 compared to 1995, mainly due to increased HFC consumption as a substitute for (H)CFC use. PFC emissions increased due to a higher production level of the Semiconductor manufacturing industry. And actual emissions of SF<sub>6</sub> remained rather stable during the period 1995–2008. Table 4.6 gives an overview of the trends in actual emissions from 1990–2008.

#### Activity data and (implied) emission factors

Detailed information on the activity data and emission factors can be found in the monitoring protocols 0060–0066 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

The activity data used to estimate the emissions of the F-gases are based on the following sources:

- Consumption data of HFCs (Stationary refrigeration, Mobile air-conditioning, Aerosols and Foams) are obtained from annual reports from PriceWaterhouseCoopers.
- Activity data on the use of PFCs in Semiconductor manufacturing and SF<sub>6</sub> in Sound-proof windows and electron microscopes are obtained from different individual companies (confidential information).

Emission factors used to estimate the emissions of the F-gases in this category are based on the following sources:

- Stationary refrigeration, Mobile air conditioning, Aerosols and Foams: annual leak rates are based on surveys (De Baedts et al., 2001) and other literature.
- Semiconductor manufacturing: emission factors which are confidential information of the only company.
- Sound-proof windows: EF used for production is 33% (IPCC default); EF (leak rate) used during the lifetime of the windows is 2% per year (IPCC default).
- Electron microscopes: emission factors are confidential information of the only company.

The source Electrical equipment comprises SF<sub>6</sub> emissions of users of high-voltage circuit breakers and the only international test laboratory for power switches. The emissions from the circuit breakers are obtained from EnergieNed, the Federation of Energy Companies in the Netherlands and the emissions from testing from the test laboratory.

#### 4.7.4 Methodological issues

To comply with the IPCC Good Practice Guidance (IPCC, 2001) IPCC Tier 2 methods are used to estimate emissions of the sub-sources Stationary refrigeration, Mobile air conditioning, Aerosols, Foams and Semiconductor manufacturing.

The country-specific methods for the sources Sound-proof windows and Electron microscopes are equivalent to IPCC Tier 2 methods and from 2007 onwards the country-specific method for the source Electrical equipment is equivalent to the IPCC Tier 3 method.

More detailed descriptions of the methods used and emission factors can be found in the protocols 0060–0066 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 4.1.

#### 4.7.5 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of the uncertainties according to the IPCC source category. The uncertainty in HFC emissions from HFC consumption is estimated to be 50%, and the uncertainties in PFC and SF<sub>6</sub> emissions are estimated to be about 25% and 55%, respectively. The uncertainty in the activity data for the HFC sources and for SF<sub>6</sub> and PFC sources is estimated at 10%, 50% and 5%, respectively. For the emission

		1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
HFCs	NIR 2009	NO	249	991	828	759	922	1,052	1,123	1,248	1,471
	NIR 2010	NA,NO	247	1,053	919	870	1,055	1,195	1,279	1,412	1,578
	Difference	NA,NO	-2	62	91	112	134	144	156	164	107
SF <sub>6</sub>	NIR 2009	217	301	319	323	283	243	246	238	202	214
	NIR 2010	217	301	318	322	281	242	265	254	217	226
	Difference	0	0	-1	-1	-2	-2	18	16	14	12

factors the uncertainties are estimated 50%, 25% and 25%. All of these figures are based on the judgments of experts.

#### Time series consistency

Consistent methodologies have been used to estimate emissions from these sources.

#### 4.7.6 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures discussed Chapter 1.

#### 4.7.7 Source-specific recalculations

In response to the ERT report 2009, the method to estimate HFC emissions from Stationary refrigeration (2F1) has been changed. From now on the complicated mass-balance method is replaced by a straightforward EF-based method. More detailed information about this change can be found in the protocol 0060 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Furthermore, information about pre-charged refrigeration equipment and new SF<sub>6</sub> emissions from the only electrical equipment testing company in the Netherlands came available.

The results of the recalculation and changes were corrected in this submission (see Table 4.7).

#### 4.7.8 Source-specific planned improvements

There are no source-specific improvements planned for this category.

### 4.8 Other industrial processes [2G]

#### 4.8.1 Source category description

The national inventory of the Netherlands comprises emissions of greenhouse gases related to four source categories in this category:

- Fireworks and candles: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions;
- Degassing of drinking water: CH<sub>4</sub> emissions;
- Miscellaneous non-energy fossil fuel product uses, (e.g., lubricants and waxes); CO<sub>2</sub> emissions (about 0.2 Tg).

The CO<sub>2</sub> emissions reported in category 2G stem from the direct use of specific fuels for non-energy purposes, which results in partially or fully 'oxidation during use (ODU)' of the carbon contained in the products – for example, lubricants, waxes and other fuels. With the exception of lubricants and waxes no other fuels are included in this category. Oxidation for mineral turpentine is included in Sector 3 (Indirect CO<sub>2</sub> of solvent use).

#### 4.8.2 Key sources

There are no key sources identified from these source category (see also Annex 1).

#### 4.8.3 Overview of shares and trends in emissions

The small CO<sub>2</sub> and CH<sub>4</sub> emissions remained rather constant between 1990 and 2008.

#### 4.8.4 Activity data and (implied) emission factors

Detailed information on the activity data and emission factors can be found in the monitoring protocols 0043 and 0054 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl)

The activity data used are based on the following sources:

- Fireworks: data on annual sales from branch organisation;
- Candles: average use of 3.3 kg per person ([www.bolsius.com](http://www.bolsius.com));
- Production of drinking water: Volume Statistics Netherlands (CBS);
- Fuel use: energy statistics obtained from Statistics Netherlands (CBS).

Emission factors:

- Fireworks: CO<sub>2</sub>: 43 t/t; CH<sub>4</sub>: 0.78 t/t; N<sub>2</sub>O: 1.96 t/t (Brouwer et al., 1995);
- Candles: 2.3 t/t (EPA, 2001);
- Production of drinking water: 2.47 tons CH<sub>4</sub>/10<sup>6</sup> m<sup>3</sup>;
- Use of fuels for production of lubricants: ODU factor of 50% (IPCC default);
- Production of waxes: ODU factor of 100% (IPCC default).

CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from Fireworks and candles showed a 'peak' in 1999 because of the millennium celebrations.

#### 4.8.5 Methodological issues

The methodologies used to estimate the greenhouse gas emissions included in this category are in compliance with the IPCC Good Practice Guidance (IPCC, 2001). More detailed descriptions of the methods used and the emission factors can be found in protocols 0043 and 0054 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 4.1:

- Fireworks and candles: country-specific methods and emission factors are used to estimate emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O.
- Degassing of drinking water: a country-specific methodology and emission factor are used to estimate the CH<sub>4</sub> emissions, which is the main source of CH<sub>4</sub> emissions in this category.
- Miscellaneous non-energy fossil fuel product uses (i.e. lubricants and waxes): a Tier 1 method is used to estimate

emissions from lubricants and waxes using IPCC default emission factors.

#### 4.8.6 Uncertainties and time-series consistency

##### Uncertainties

The Tier 1 uncertainty analysis in Annex 7 shown in Tables A7.1 and A7.2 provides estimates of the uncertainties according to IPCC source category.

The uncertainty in CO<sub>2</sub> emissions of other industrial processes is estimated to be approximately 20% (5% in activity data and 20% in emission factor), mainly due to the uncertainty in the ODU factor for lubricants. The uncertainty in the activity data (such as domestic consumption of these fuel types) is generally very large, since it is based on production-, import- and export figures.

The uncertainty in CH<sub>4</sub> emissions of other industrial processes is estimated to be 50% (10% in activity data and 50% in emission factor). The uncertainty in N<sub>2</sub>O emissions of other industrial processes is estimated at 70% (50% in activity data and 50% in emission factor). All figures are based on the judgments of experts, since no specific monitoring data or literature is available for the current situation in the Netherlands.

##### Time-series consistency

Consistent methodologies and activity data have been used to estimate the emissions of these sources.

#### 4.8.7 Source specific QA/QC and verification

The source categories are covered by the general QA/QC procedures discussed in Chapter 1.

#### 4.8.8 Source specific recalculations

There have been no source-specific recalculations in comparison to the previous submission.

#### 4.8.9 Source specific planned improvements

There are no source-specific improvements

# Solvent and other product use [CRF Sector 3]

- Major changes in Sector 3 Solvent and other product use compared to the National Inventory Report 2009

*Emissions:* No changes.

*Key sources:* There are no key sources in allocation in this sector.

*Methodologies:* There have been no methodological changes in this sector.

## 5.1 Overview of sector

Emissions of the greenhouse gases in this sector include indirect emissions of CO<sub>2</sub> related to the release of non-methane volatile organic compounds (NMVOC) with the use of solvents and a wide range of other fossil carbon-containing products (e.g., paints, cosmetics, cleaning agents). In addition, this sector includes N<sub>2</sub>O emissions originating from the use of N<sub>2</sub>O as anaesthesia and as a propelling agent in aerosol cans (for example cans with cream).

The Netherlands has three source categories in this IPPC Common Reporting Format (CRF) sector:

- 3A, 3B, 3D “Solvents and other product use”: indirect CO<sub>2</sub> emissions (related to NMVOC)
- 3D “Anaesthesia”: N<sub>2</sub>O emissions
- 3D “Aerosol cans”: N<sub>2</sub>O emissions.

This sector comprises non-combustion emissions from households, services, hospitals, research- and governmental institutions etc, except for the following emissions:

- Use of F-gases (HFCs, PFCs and SF<sub>6</sub>). In accordance with the IPCC Reporting Guidelines F-gases are included in 2 “Industrial processes” (thus including their use in the Residential and Commercial sectors)
- Direct non-energy use of mineral oil products (e.g., lubricants, waxes). These are included in 2G “Industrial processes”
- Several minor sources of CH<sub>4</sub> emissions from non-industrial, non-combustion sources. These are included in Sector 2G because the CRF does not permit methane emissions to be included in Sector 3

The following emission from the manufacturing industry is also included in this Chapter:

- Indirect CO<sub>2</sub> emissions from 3C “Chemical products, manufacture and processing”. These NMVOC emissions are included in categories 3A, 3B and 3D.

The following protocol, which can be accessed on [www.greenhousegases.nl](http://www.greenhousegases.nl), describes the methodologies applied for estimating CO<sub>2</sub> and N<sub>2</sub>O emissions from solvent and product use in the Netherlands:

- [Protocol 0054: CO<sub>2</sub>, N<sub>2</sub>O en CH<sub>4</sub> from Other process emissions and product use.](#)

### Overview of shares and trends in emissions

Table 5.1 shows the contribution of the emissions from Solvent and other product use in the Netherlands. Total greenhouse gas emissions from Solvent and product use in the Netherlands were 0.5 Tg CO<sub>2</sub> eq in 1990 and 0.2 Tg CO<sub>2</sub> eq in 2008.

Total emissions of the sector declined by 60% between 1990 and 2003, and remained quite stable between 2003 and 2008. CO<sub>2</sub> emissions from the sector decreased by 59% between 1990 and 2008, mainly due to decreasing indirect emissions from paints that resulted from the implementation of an emission reduction program for NMVOC (KWS, 2000). N<sub>2</sub>O emissions from anaesthesia fell by 87% from 1990 to 2008 due to the better dosing in hospitals and other medical institutions. The emissions of N<sub>2</sub>O from food aerosol cans increased 101% in this period. Total N<sub>2</sub>O emissions from this chapter have declined since 1990 by 67%.

### Key sources

Solvent and product use is a minor source of greenhouse gas emissions. No key sources are included in this sector. The most relevant sources are indirect CO<sub>2</sub> emissions from paint application and the use of N<sub>2</sub>O for anaesthesia in hospitals.

Sector/category	Gas	Key	Emissions base-year (1990)	Emissions 2007	Emissions 2008	Change 2008 - 2007	Contribution to total in 2008 (in %)		
			Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	By sector	Of total gas	Of total CO <sub>2</sub> eq
3 Solvent and other product use	CO <sub>2</sub>	Level, Trend	0.3	0.1	0.1	0.0		0.1	0.1
	N <sub>2</sub> O		0.2	0.1	0.1	0.0		0.6	0.0
	All		0.5	0.2	0.2	0.0			0.1
3A. Paint application	CO <sub>2</sub>		0.2	0.1	0.1	0.0	29	0.0	0.0
3A. Paint application	All		0.2	0.1	0.1	0.0	29		0.0
3B. Degreasing and dry cleaning	CO <sub>2</sub>		0.0	0.0	0.0	0.0	1	0.0	0.0
3B. Degreasing and dry cleaning	All		0.0	0.0	0.0	0.0	1		0.0
3D. Other	CO <sub>2</sub>		0.1	0.1	0.1	0.0	33	0.0	0.0
	N <sub>2</sub> O		0.2	0.1	0.1	0.0	36	0.6	0.0
3D1 Anaesthesia	N <sub>2</sub> O		0.2	0.0	0.0	0.0	13	0.2	0.0
3D3 Aerosol cans	N <sub>2</sub> O		0.0	0.0	0.0	0.0	23	0.4	0.0
3D. Other	All		0.3	0.1	0.1	0.0			0.1
National Total GHG emis- sions (excl. CO <sub>2</sub> LULUCF)	CO <sub>2</sub>		159.3	172.3	175.7	3.4		100.0	
	N <sub>2</sub> O		20.2	15.4	11.8	-3.7		100.0	
	All		213.3	206.9	206.9	0.0			

## 5.2 Indirect CO<sub>2</sub> emissions from Solvents and product use (Paint application [3A], Degreasing and dry cleaning [3B] and Other [3D])

### 5.2.1 Source category description

CRF source category 3A Paint application includes the indirect CO<sub>2</sub> emissions of solvents from the use of both industrial paints and paints used by households and professional painters. Indirect emissions from the use of solvents in degreasing and dry cleaning are included in CRF source category 3B, which covers the use of solvents for cleaning and degreasing of surfaces, the dry cleaning of clothing and textiles and the degreasing of leather.

### 5.2.2 Activity data and implied emission factors

Detailed information on the activity data and emission factors of NMVOC estimates can be found in the monitoring protocol 0054 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Activity data: consumption data and NMVOC contents of products are mainly provided by trade associations, such as the VVVF (for paints), the NCV (for cosmetics) and the NVZ (for detergents). Consumption of almost all solvent-containing products has increased since 1990. However, the general NMVOC content of products (especially paints) has decreased over the last years, resulting in a steady decline in NMVOC emissions since 1990 (see Section 2.4). Due to the increased sales of hairspray and deodorant sprays NMVOC emissions have increased slightly in recent years. It is assumed that the NMVOC contents of these products have remained stable.

Emission factors: it is assumed that all NMVOC in the product is emitted (with the exception of some cleaning products and methylated spirit, which are partly broken down in sewerage treatment plants after use, or used as fuel in BBQs or fondue sets (methylated spirit). The carbon contents of NMVOC

emissions are documented in the monitoring protocol on the website [Hwww.greenhousegases.nl](http://www.greenhousegases.nl).

### 5.2.3 Methodological issues

Country-specific carbon contents of the NMVOC emissions from 3A Paint application, 3B Degreasing and dry cleaning and 3D Other product use are used to calculate indirect CO<sub>2</sub> emissions. Monitoring of NMVOC emissions from these sources differs per source. Most of the emissions are reported by branch organisations (e.g., paints, detergents and cosmetics). The indirect CO<sub>2</sub> emissions from NMVOC are calculated from the average carbon contents of the NMVOC in the solvents.

Category	3A	3B	3D
C-content NMVOC (%)	0.72	0.16	0.69

The carbon content of degreasing and dry cleaning is very low due to the high share of chlorinated solvents (mainly tetrachloroethylene used for dry cleaning). The emissions are then calculated as follows:

$$\text{CO}_2 \text{ (in Gg)} = \sum \{ \text{NMVOC emission in subcategory } i \text{ (in Gg)} \times \text{C-fraction subcategory } i \} \times 44/12$$

The fraction of organic carbon (of natural origin) in the NMVOC emissions is assumed to be negligible.

### 5.2.4 Uncertainty and time-series consistency

#### Uncertainty

These sources do not affect the overall total or the trend in the direct greenhouse gas emissions. The uncertainty of indirect CO<sub>2</sub> emissions is not explicitly estimated for this category, but it is expected to be fairly low. Based on the expert judgment, the uncertainty in the NMVOC emissions is estimated to be 25%, and the uncertainty in the carbon

contents is estimated at 10%, resulting in an uncertainty in CO<sub>2</sub> emissions of approximately 27%.

#### Time-series consistency

Consistent methodologies have been applied for all source categories. As the quality of the activity data used was not uniform throughout the complete time-series, some extrapolation of the data was required. It is assumed that the accuracy of the estimates is not significantly affected by this. The emission estimates for the source categories are expected to be reasonably good.

#### 5.2.5 Source-specific QA/QC and verification

This source category is covered by the general QA/QC procedures discussed in Chapter 1.

#### 5.2.6 Source-specific recalculations

There were no recalculations in this sector.

#### 5.2.7 Source-specific planned improvements

There are no source-specific improvements planned.

### 5.3 Miscellaneous N<sub>2</sub>O emissions from solvents and product use (use of N<sub>2</sub>O for anaesthesia [3D1] and N<sub>2</sub>O from aerosol cans [3D3])

#### 5.3.1 Source category description

Emissions of N<sub>2</sub>O from the use of anaesthesia are included in 3D1. Emissions of N<sub>2</sub>O from aerosol cans are included in category 3D3.

#### 5.3.2 Activity data and implied emission factors

Detailed information on the activity data and emission factors of N<sub>2</sub>O estimates are found in the monitoring protocol 0054 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

**Activity data:** The major hospital supplier of N<sub>2</sub>O for aesthetic use reports the consumption data of aesthetic gas in the Netherlands annually. The Dutch Association of Aerosol Producers (NAV) reports data on the annual sales of N<sub>2</sub>O-containing spray cans. Missing years are then extrapolated on the basis of this data. Domestic sales of cream in aerosol cans have shown a strong increase since 1990. The increase is reflected in the increased emissions.

**Emission factors:** The emission factor used for N<sub>2</sub>O in anaesthesia is 1 kg/kg. Sales and consumption of N<sub>2</sub>O for anaesthesia are assumed to be equal each year. The emission factor for N<sub>2</sub>O from aerosol cans is estimated to be 7.6 g/can (based on data provided by one producer), and is assumed to be constant over time.

#### 5.3.3 Methodological issues

Country-specific methodologies are used for the N<sub>2</sub>O sources in Sector 3. Since the emissions in this source category are from non-key sources for N<sub>2</sub>O, the present methodology complies with the IPCC Good Practice Guidance (IPCC, 2001). A full description of the methodology is provided in the monitoring protocol 0054 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 5.3.4 Uncertainties and time-series consistency

#### Uncertainties

These sources do not affect the overall total or trend in the Dutch emissions of direct greenhouse gases. For N<sub>2</sub>O emissions, the uncertainty is estimated to be approximately 50% based on the judgment of experts. Uncertainty in the activity data of N<sub>2</sub>O use is estimated to be 50% and that of the emission factor to be 0% (all gas is released).

#### Time-series consistency

Consistent methodologies have been applied for all source categories. The quality of the activity data needed was not uniform for the complete time-series, requiring some extrapolation of data. This is not expected to introduce significant problems with the accuracy of the estimates. The estimates for the source categories are expected to be quite good.

#### 5.3.5 Source-specific QA/QC and verification

This source category is covered by the general QA/QC procedures discussed in Chapter 1.

#### 5.3.6 Source-specific recalculations

There are no source-specific recalculations compared to the previous submission.

#### 5.3.7 Source-specific planned improvements

There are no source-specific improvements planned.





# Agriculture [CRF Sector 4]

## ■ Major changes in the Agriculture sector compared to the National Inventory Report 2009

*Emissions:* Compared to 2007, nitrous oxide (N<sub>2</sub>O) emissions and methane (CH<sub>4</sub>) emissions increase as a result of increasing numbers of livestock.

*Key sources:* The CH<sub>4</sub> emission from “Manure management, poultry” is now key source.

*Methodologies:* As a result of error correction feed intake of female cattle has been recalculated in 2009. This recalculation also uncovered an error in volume-excretion for 1990-2000. This affects CH<sub>4</sub> from manure management for all animal categories. Due to this recalculation methane (CH<sub>4</sub>)

emissions for the years 1990-1996 are slightly higher and for the years 1997-2000 the emissions are lower.

Methane emissions from agricultural soils are regarded as natural, non-anthropogenic emissions and therefore are not included.

## 6.1 Overview of the sector

Emissions of greenhouse gases from “Agriculture” include all anthropogenic emissions from the agricultural sector, with the exception of emissions from fuel combustion and carbon dioxide (CO<sub>2</sub>) emissions by land use in agriculture. These emissions are included in 1A4c “Agriculture/forestry/fisheries” (Section 3.6) and in 5 “Land Use, Land Use Change and Forestry” (LULUCF Sections 7.3 and 7.4).

In the Netherlands, three source categories occur in the agricultural sector:

- 4A “Enteric fermentation”: CH<sub>4</sub> emissions
- 4B “Manure management”: CH<sub>4</sub> and N<sub>2</sub>O emissions
- 4D “Agricultural soils”: N<sub>2</sub>O emissions.

The other Intergovernmental Panel on Climate Change (IPCC) categories – 4C “Rice cultivation”, 4E “Prescribed burning of savannas”, 4F “Field burning of agricultural residues” and 4G “Other” – do not occur in the Netherlands. Open fires/burning in the field is prohibited by law and therefore negligible in practice.

Manure management (4B) includes all emissions from confined animal waste management systems (AWMS). CH<sub>4</sub> emissions from animal manure produced in the meadow during grazing are included in category 4B “Manure management”; N<sub>2</sub>O emissions from this source are included in category 4D2 “Animal production”. These different approaches are in accordance with IPCC Guidelines (IPCC, 2001).

The following protocols on [www.greenhousegases.nl](http://www.greenhousegases.nl) describe the methodologies, activity data and emission factors applied in estimating N<sub>2</sub>O and CH<sub>4</sub> emissions in the agricultural sector in the Netherlands:

- Protocol 0067: CH<sub>4</sub> from Enteric fermentation (4A)
- Protocol 0068: N<sub>2</sub>O from Manure management (4B)
- Protocol 0069: CH<sub>4</sub> from Manure management (4B)
- Protocol 0070: N<sub>2</sub>O from Agricultural soils: indirect emissions (4D)
- Protocol 0071: N<sub>2</sub>O from Agricultural soils: direct emissions and emissions from animal production (4D).

### Overview of shares and trends in emissions

Table 6.1 shows the contribution of the agricultural source categories to the total national greenhouse gas inventory. This table also presents the key sources identified in the agricultural sector as specified by trend or level, or both.

In 2008, CO<sub>2</sub> equivalent emissions from Sector 4 “Agriculture” contributed 8.9% to the total national emissions (without LULUCF) compared to 11% in 1990. In 2008, emissions of CH<sub>4</sub> and N<sub>2</sub>O from agricultural sources accounted for 54% and 80% of the national total CH<sub>4</sub> and N<sub>2</sub>O emissions. Category 4A “Enteric fermentation” is the main source of CH<sub>4</sub> emissions and category 4D “Agricultural soils” is the largest source of N<sub>2</sub>O emissions included in this sector.

Total greenhouse gas emissions from Agriculture decreased by approximately 13% between 1990 and 2008, from 22.5 Tg CO<sub>2</sub> eq in 1990 to 18.5 Tg CO<sub>2</sub> eq in 2008 (see also Figure 6.1). This decrease was largely the result of decreasing numbers

Sector/category	Gas	Key	Emissions base-year	Emissions 2007	Emissions 2008	Change 2008 – 2007	Contribution to total in 2008 (%)		
			Level, Trend	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	By sector	Of total gas	Of total CO <sub>2</sub> eq
4. Agriculture	CH <sub>4</sub>			10.5	9.0	9.1	49	54	4.4
	N <sub>2</sub> O			11.9	9.3	9.4	51	80	4.5
	All			22.5	18.3	18.5	100		8.9
4A. Enteric fermentation	CH <sub>4</sub>			7.5	6.3	6.5	35	38	3.1
4A1 Cattle	CH <sub>4</sub>	L,T		6.8	5.6	5.8	31	34	2.8
4A Swine	CH <sub>4</sub>			0.4	0.4	0.4	2	2	0.2
4A2-13 Other animals	CH <sub>4</sub>			0.3	0.3	0.3	2	2	0.1
4B. Manure management	CH <sub>4</sub>			3.0	2.6	2.7	14	16	1.3
	N <sub>2</sub> O	L,T2		0.8	0.9	0.9	5	8	0.4
	All			3.8	3.5	3.6	19		1.7
4B1 Cattle	CH <sub>4</sub>	L		1.6	1.5	1.5	8	9	0.7
4B8 Swine	CH <sub>4</sub>	L		1.1	1.1	1.1	6	7	0.5
4B9 Poultry	CH <sub>4</sub>	T2		0.3	0.1	0.1	0	0	0.0
4B2-7, 10-13 Other animals	CH <sub>4</sub>			0.0	0.0	0.0	0	0	0.0
4D Agriculture soils	N <sub>2</sub> O			11.1	8.4	8.5	46	72	4.1
4D1 Direct soil emissions	N <sub>2</sub> O	L,T		4.7	4.7	4.8	26	41	2.3
4D2 Animal production on agricultural soils	N <sub>2</sub> O	L,T		1.4	0.6	0.6	3	5	0.3
4D3 Indirect emissions	N <sub>2</sub> O	L,T		5.0	3.1	3.0	16	26	1.5
National Total GHG emissions (excl. CO <sub>2</sub> LUCF)	CH <sub>4</sub>			25.5	16.8	17.1		100	100
	N <sub>2</sub> O			20.2	15.4	11.8		100	
	All			213.3	206.9	206.9			

\*Key sources: L = Level; T = Trend; 1 = Tier 1; 2 = Tier 2.

Numbers of animals in 1990–2008 (1000 heads) (CBS, 2009)

Table 6.2

Animal type	1990	1995	2000	2005	2006	2007	2008
Cattle	4,926	4,654	4,070	3,799	3,745	3,763	3,890
- Adult dairy cattle	1,878	1,708	1,504	1,433	1,420	1,413	1,466
- Adult non-dairy cattle	120	146	163	152	143	144	127
- Young Cattle	2,929	2,800	2,403	2,214	2,182	2,206	2,297
Sheep	1,702	1,674	1,308	1,363	1,376	1,369	1,213
Goats	61	76	179	292	310	324	355
Horses	70	100	118	133	128	134	144
Pigs (*1000)	13.9	14.4	13.1	11.3	11.4	11.7	12.0
Poultry (*1000)	95.6	92.2	107.2	95.9	94.7	96.0	99.7

of livestock, a decreased application of animal manure and a decreased use of synthetic fertilisers.

Compared to 2007, nitrous oxide (N<sub>2</sub>O) emissions and methane (CH<sub>4</sub>) emissions increase as a result of increasing numbers of livestock.

#### Overview of trends in activity data

Livestock numbers are the primary activity data used in the calculation of CH<sub>4</sub> and N<sub>2</sub>O.


Activity data for the livestock numbers are based on the annual agricultural survey performed by Statistics Netherlands (CBS). Data can be found on the website [www.cbs.nl](http://www.cbs.nl), in Annex 8, Table A8.1 and in background documents (e.g., Van der Hoek and Van Schijndel, 2006). Table 6.2 presents an overview.

For cattle, three categories are distinguished:

- mature dairy cattle: adult cows for milk production
- mature non-dairy cattle: adult cows for meat production
- young cattle: mix of different age categories for breeding and meat production, including adult male cattle.

Between 1990 and 2008, (dairy) cattle, pig and sheep numbers decreased by 22, 14 and 29% respectively, while poultry numbers remained fairly constant. Goat numbers increased by a factor 5 and horse numbers nearly doubled in this period.

For mature dairy cattle, the decrease in numbers was associated with an increase in milk production per cow between 1990 and 2008. The increased milk production per cow is the a result of both genetic changes (due to breeding programmes for milk yield) as well as the increase in feed intake and the high feeding quality of cattle diets. Total milk production in the Netherlands is determined mainly by European Union (EU) policy on milk quotas, which remained



unchanged in the same period. In order to comply with the unchanged milk quota, animal numbers of mature dairy cattle had to decrease to counteract the effect of increased milk production per cow. Between 1990 and 2008 the numbers of young (dairy) cattle follow the same trends as those of adult female cattle – namely, a decrease.

The Netherlands' manure and fertilizer policy also influences livestock numbers. Young cattle, pig and poultry numbers in particular decreased as a result of the introduction of measures such as buying up part of the pig and poultry production rights (ceilings for total phosphate production by animals) by the government and lowering the maximum application standards for manure and artificial fertilizer. For pigs and young cattle the decreasing trend of the past has levelled off in the last couple of years. For pigs, it changed into a slight increase since 2004.

The increased number of swine in 1997 was a direct result of the outbreak of classical swine fever in that year (Annex 8 Table A8.1). In areas where this disease was present, the transportation of pigs, sows and piglets to

the slaughterhouse was not allowed, so the animals had to remain on the pig farms for a relatively long period (accumulation of pigs).

An increase in the number of poultry is observed between 1990 and 2002. In 2003, however, poultry numbers decreased by almost 30% as a direct result of the avian flu outbreak. In the years after 2003 the poultry population recovered to a large extent and reached a level of 4% below the 2002 level in 2008.

The increase in the number of goats might be explained as an effect of the milk quota for cattle. As result of the milk quota for cattle and the market development for goat milk products, farmers tend to change their management towards goats.

	1990	1995	2000	2005	2006	2007	2008
<i>Mature dairy cattle</i>	110	116	120	126	128	129	128
<i>Mature non-dairy cattle</i>	65	66	67	71	71	72	73
<i>Young cattle</i>	37	37	35	34	34	34	34

## 6.2 Enteric fermentation [4A]

### 6.2.1 Source category description

Methane emissions from Enteric fermentation are produced as a by-product of the digestive process in which organic matter (mainly carbohydrates) is degraded and utilized by micro-organisms under anaerobic conditions. Both ruminant (e.g. cattle, sheep and goats) and non-ruminant animals (e.g. pigs and horses) produce CH<sub>4</sub>, although ruminants produce far more CH<sub>4</sub> per unit of feed intake than non-ruminants due to the specialisation of the digestive system to digest fibrous material. Ruminants have a strongly expanded chamber, the rumen, at the front side of their digestive tract that allows the selective retention of feed particles and that supports intensive microbial fermentation of the feed. This yields several nutritional advantages including the capacity to digest fibrous material and the synthesis of microbial protein that can be digested in the intestine, but which is also accompanied by a high methane production by methanogens in the rumen.

Buffalo and camels do not occur in the Netherlands. The emissions from llamas, mules and donkeys are negligible and, therefore, not taken up in the inventory. Enteric fermentation methane emission from poultry is not estimated due to the negligible amount of CH<sub>4</sub> emission for this animal category. The IPCC Guidelines do not provide a default emission factor for this animal category. Other countries do not estimate emissions from poultry either.

### 6.2.2 Overview of shares and trends in emissions

In 2008 Enteric fermentation accounted for 35% of the total greenhouse gas emissions from the agricultural sector (see Table 6.1). In the Netherlands, cattle accounts for the majority of CH<sub>4</sub> emissions from Enteric fermentation (89%) in 2008. The second largest CH<sub>4</sub> emission source in category 4A is swine (6%). 4A Other consists of sheep, goats and horses, and accounts for 5%.

CH<sub>4</sub> emissions from Enteric fermentation decreased from 7.5 Tg CO<sub>2</sub> eq to 6.5 Tg (–14%) between 1990 and 2008, with CH<sub>4</sub> emissions from Enteric fermentation by cattle and swine decreasing by 15% and 14%, respectively. From 2007 to 2008 a rather small increase indicates a stabilisation.

### 6.2.3 Activity data and emission factors

Trends in CH<sub>4</sub> emission from enteric fermentation are explained by a change in animal numbers, a change in emission factor or both.

Detailed information on data sources for activity data and emission factors can be found in the following monitoring protocol:

- [Protocol 0067: CH<sub>4</sub> from Enteric fermentation \(4A\)](#)

All relevant documents concerning methodology, emission factors and activity data are published on the website [www.greenhousegases.nl](http://www.greenhousegases.nl). Table 6.2 (in Section 6.1) presents an overview of animal numbers. In Annex 8, Tables A8.1, A8.2 and A8.3 show the activity data for all animal categories and emission factors for cattle.

For swine, sheep, goat and horses, default IPCC emission factors are used (1.5, 8, 5 and 18 kg/animal, respectively). Changes in emissions for these animal categories are therefore explained entirely by changes in animal numbers. To a great extent this is also the case for cattle, but the total decrease in CH<sub>4</sub> emission is lower due to an increase in implied emission factor (IEF).

### Trends in cattle IEF

The emission factors for three cattle categories are calculated annually. For mature dairy cattle, a Tier 3 approach is used to calculate the CH<sub>4</sub> production per cow per year on the basis of data on the share of feed components and their chemical nutrient composition (Smink et al, 2005). For mature non-dairy and young cattle, a Tier 2 approach is used to calculate the CH<sub>4</sub> production per animal per year on the basis of data on the feed intake (Smink, 2005). For more information on the methods and the recalculation used, see Section 6.2.2 and Section 6.2.5.

Table 6.3 shows the implied emission factors (IEFs) of the different cattle categories reported. The implied emission factor for young cattle is an average of several subcategories (Annex 8, Table A8.3)

For both mature dairy cattle and mature non-dairy cattle, IEFs increased primarily as a result of an increase in total feed intake during the period 1990–2008. For dairy cattle, a change in the feed nutrient composition partly counteracted this effect (see Section 6.2.2). For young cattle the decrease of IEF between 1990 and 2008 can be explained by a decrease in the average total feed intake due to a shift towards relatively more meat calves in the population of young cattle (Annex 8 Table A8.1).

### Comparison of cattle IEF with IPCC defaults

Table 6.4 shows that the mature dairy cattle IEF follows the increasing trend in milk production. Compared to the default IPCC IEF of 118 kg CH<sub>4</sub> per cow for mature dairy cattle (at a milk production rate of 6700 kg / cow / yr), the IEF used in the Netherlands is slightly lower.

In 1997 for instance, a milk production of about 6800 kg per year per cow led to an emission factor of 117 kg/animal/yr, less than 1% lower than the default of 118 kg/animal/yr. An explanation of the difference can be found in the data on feed intake, dietary composition and nutrient composition

	1990	1995	2000	2005	2006	2007	2008
Milk production	6050	6613	7296	7554	7744	7945	7926
IEF for methane	110	116	120	126	128	129	128

of dietary components as an input to an alternative country specific model that predicts the methane emission factor for mature dairy cattle (Bannink, 2010). With increasing milk production per cow a decrease in the amount of CH<sub>4</sub> emission per unit of milk produced (from 0.018 to 0.016 kg CH<sub>4</sub> / kg milk) can be seen.

The higher IEF for mature non-dairy cattle, (compared to the IPCC default value of 48 per animal) can be explained by the higher total feed intake per adult non-dairy cow. The relatively large share of meat calves for white and rose veal production explains the relatively low IEF for young cattle compared to the IPCC default value (Annex 8 Table A8.1).

#### 6.2.4 Methodological issues

A detailed description of the method, data sources and emission factors is found in the protocol on [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 6.2.1. A specified description with more details on data and data sources on cattle used in calculations until 2006 can be found in Smink et al. (2005) and Smink (2005). In 2009, a recalculation was carried out for the whole time series (Bannink, 2009 and Van Bruggen 2009).

Emissions from enteric fermentation are calculated from activity data on animal numbers and the appropriate emission factors.

$\text{CH}_4 \text{ emission} = \sum \text{EF}_i (\text{kg CH}_4/\text{animal}_i) * [\text{number of animals for livestock category } i]$

#### Cattle

The emission factors for cattle are calculated annually for several subcategories of dairy and non-dairy cattle. For mature dairy cattle a country-specific method based on a Tier 3 methodology is followed; for the other cattle categories, the calculation is based on a country-specific Tier 2 methodology.

The feed intake of cattle, which is estimated from the energy requirement calculation used in the Netherlands, is the most important parameter in the calculation of the CH<sub>4</sub> emission factor for cattle. For instance for dairy cows the energy requirement expressed as net energy value of lactation (or VEM in Dutch) is calculated on the basis of total milk production and feed composition. For young cattle the energy requirement is calculated on the basis of total weight gain and feed composition.

The intake of grass silage, maize silage, wet by-products, concentrates and grass products is estimated from national statistics found at [www.cbs.nl](http://www.cbs.nl) (Van Bruggen, 2009). More information on the Netherlands VEM system is presented in Smink et al. (2005) and Tamminga et al. (2004).

#### Mature dairy cattle

The CH<sub>4</sub> emission from enteric fermentation by mature dairy cattle is calculated by a Tier 3 approach using dynamic modelling (Smink et al., 2005). The model of Mills et al. (2001) is employed, including updates (Bannink et al., 2005). This model is based on the mechanistic, dynamic model of rumen fermentation processes developed by Dijkstra et al. (1992). It has been developed for mature cattle and is therefore not suitable for other ruminant categories such as young cattle. The model calculates the gross energy (GE) intake and CH<sub>4</sub> emission factor (kg CH<sub>4</sub>/cow/yr) and the methane conversion factor (MCF; % of GE intake converted into CH<sub>4</sub>) on the basis of data on the share of feed components (grass silage, maize silage, wet by-products and concentrates), their chemical nutrient composition (soluble carbohydrates, starch, NDF, crude protein, ammonia, crude fat and ash) and the intrinsic degradation characteristics of starch, NDF and crude protein in the rumen. Data on the share of feed components in the diet are found at [www.cbs.nl](http://www.cbs.nl) (Van Bruggen, 2009). Data on the chemical nutrient composition of individual roughages are provided by Blgg (a leading laboratory in the Dutch agricultural and horticultural sector with roughage sampling, and analytical and advisory activities, and able to deliver data that can be taken is representative of the average Dutch farming conditions; [www.blgg.com](http://www.blgg.com)). Data used between 1990 and present are published by Bannink (2010) (via [www.prtr.nl](http://www.prtr.nl)).

#### Young cattle and non-dairy cattle

The methane emission factor (EF) for enteric fermentation by non-dairy and young cattle is calculated by multiplying the GE intake by a methane conversion factor (Smink, 2005). Changes in GE intake are based on changes in the total feed intake and on the share of feed components. Data on the amounts of feed components, expressed as dry matter (DM) intake are found at [www.cbs.nl](http://www.cbs.nl) (Van Bruggen, 2009) and in Annex 8 Table A8.2 Gross Energy intake can be found.

The equation for calculating the EF (in kg per animal per year) is:

$$\text{EF} = (\text{MCF} * \text{GE intake} * 365 \text{ d/yr}) / 55.65 \text{ MJ/kg CH}_4$$

Where:

EF: Emission factor (kg CH<sub>4</sub>/animal/yr)

MCF: Methane conversion factor; fraction of the gross energy of feed intake converted to CH<sub>4</sub>

GE intake: Gross energy intake (MJ/animal/d)

Where:

- GE intake = Dry Matter intake (kg DM/animal/d) × 18.45 MJ/kg DM (IPCC, 2001)

- MCF = 0.04 for white veal calves and 0.06 for the other categories of young cattle and mature non-dairy cattle (IPCC, 2001)



### Other livestock

Emission factors for the source categories swine, sheep, horses and goats are based on default IPCC Tier 1 EF (IPCC, 1997).

The share in total CH<sub>4</sub> enteric fermentation emissions by these other livestock categories (sheep, goats, horses and swine) is less than 10% of the total CH<sub>4</sub> enteric fermentation emissions. According to IPCC good practice guidance (GPG), no Tier 2 method is needed if the share of a source category is less than 25–30% of the total emission by a key source category.

As already mentioned in Section 6.2.1, enteric fermentation emission from poultry is not estimated due to a lack of data on CH<sub>4</sub> emission factors for this animal category.

### 6.2.5 Uncertainty and time-series consistency

#### Uncertainty

The Tier 1 uncertainty analysis shown in Annex 7 provides estimates of uncertainty according to IPCC source categories. The uncertainty of CH<sub>4</sub> emissions from enteric fermentation from cattle sources is based on the judgements of experts and is estimated to be about 20% in annual emissions, using a 5% uncertainty for animal numbers (Olivier et al., 2009) and 15% for the emission factor (Bannink, 2010). The uncertainty in the emission factor for swine and other animals is estimated to be 50% and 30%, respectively (Olivier et al., 2009)

#### Time-series consistency

A consistent methodology is used throughout the time-series; see also Section 6.2.5. Emissions are calculated from animal population data and emission factors. The animal population data are collected in an annual census and published by Statistics Netherlands over a long period of time (several decennia). Emission factors are either constant (default IPCC) or are calculated from feed intake data collected by an annual survey published by Statistics Netherlands.

The compilers of the activity data strive to use consistent methods to produce the activity data. The time-series consistency of these activity data is, therefore, very good due to the continuity in the data provided.

### 6.2.6 Source-specific QA/QC and verification

This source category is covered by the general QA/QC procedures discussed in Chapter 1.

### 6.2.7 Source-specific recalculations

During the compilation of the CRF an error in the 2007 CH<sub>4</sub> emission was detected and corrected. The 2007 emission increased by 11.9 Gg CO<sub>2</sub> eq.

### 6.2.8 Source-specific planned improvements

There are no source-specific improvements planned.

## 6.3 Manure management [4B]

### 6.3.1 Source category description

Both CH<sub>4</sub> and N<sub>2</sub>O are emitted during the handling or storage of manure from cattle, pigs, poultry, sheep, goats and horses. These emissions are related to the quantity and the composition of the manure, and to the manure management system types and conditions. For instance, in comparison to anaerobic conditions, aerobic conditions in the manure management system will in general increase N<sub>2</sub>O emissions and decrease CH<sub>4</sub> emissions. Furthermore, longer storage times and higher temperatures will increase CH<sub>4</sub> emissions compared to shorter storage times and lower temperatures.

Buffalo and camels do not occur in the Netherlands, and the numbers of llamas, mules and donkeys are negligible and therefore not estimated. Three animal manure management systems are distinguished for emission estimates of both CH<sub>4</sub> and N<sub>2</sub>O: liquid and solid manure management systems and manure produced in the meadow while grazing.


In accordance with IPCC Guidelines, N<sub>2</sub>O emissions from manure produced in the meadow during grazing are not taken into account in the source category Manure management (see Section 6.1), but are included in the source category Agricultural soils (Section 6.4).

### 6.3.2 Overview of shares and trends in emissions

In 2008, Manure management accounted for 19% (CH<sub>4</sub> and N<sub>2</sub>O) of the total greenhouse gas emissions from the agricultural sector (Table 6.1 and Figure 6.2). In the Netherlands CH<sub>4</sub> emissions from Manure management are particularly related to cattle and swine manure management, which, in 2008, contributed 8% and 6%, respectively, to the total greenhouse gas emissions in the agricultural sector. Poultry is a minor key source for CH<sub>4</sub> emissions by manure management. Furthermore, N<sub>2</sub>O emissions from Manure management contribute 5% of the total greenhouse gas emissions from the agricultural sector.

Between 1990 and 2008, the emission of CH<sub>4</sub> from Manure management decreased by 11%. Emissions from cattle, swine and poultry decreased by 5%, 2% and 81%, respectively during this period. From 2007 to 2008, the emission of CH<sub>4</sub> from Manure management slightly increased.

The emissions of N<sub>2</sub>O from Manure management increased 11% between 1990 and 2008, from 2.6 to 2.9 Gg N<sub>2</sub>O in 2008 (Table 6.1). The relatively large decrease in N<sub>2</sub>O emissions from solid manure in 2003 is a direct result of the decrease in poultry animal manure. This decrease was due to the reduction in the number of poultry animals that followed the avian flu epidemic. In 2004 and 2005, N<sub>2</sub>O emissions increased once again following the recovery of poultry animal numbers, while in 2006 the emission decreased as a consequence of lower poultry numbers. In 2007, emissions increased as a result of increasing animal population and higher N excretion per animal. In 2008, emissions increased as a result of increasing animal population.



### 6.3.3 Activity data and (implied) emission factors

Detailed information on data sources (for activity data and emission factors) can be found in the following monitoring protocols:

- Protocol 0069: CH<sub>4</sub> from Manure management (4B)
- Protocol 0068: N<sub>2</sub>O from Manure management (4B)

More details and specific data (activity data and emission factors), including data sources (emission factors), are documented in the background documents. All relevant documents concerning methodology, emission factors and activity data are published on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Activity data on animal numbers can be found on the website [www.cbs.nl](http://www.cbs.nl), in Annex 8, Table A8.1 and in a background document (Van der Hoek and Van Schijndel, 2006). Emission factor data can be found in Annex 8 Tables A8.3 to A8.8.

The decreased CH<sub>4</sub> emission from swine between 1990 and 2008 results from the decrease in animal numbers and manure production (Annex 8, Tables A8.1 and A8.8). The

decrease is less than expected due to an increase in emission factor (Annex 8, Table A8.7). For mature non-dairy cattle and young cattle, emissions decrease only as a result of animal numbers and a small increase in emission factor. For mature dairy cattle, the decrease in CH<sub>4</sub> emissions is much lower than the decrease in animal numbers as a consequence of a higher IEF. For poultry, the large decrease in CH<sub>4</sub> emissions between 1990 and 2008 can only be explained by the lower IEF.

The slightly increased N<sub>2</sub>O emission from Manure management between 1990 and 2008 is explained by an increase in IEF partly counteracted by a decrease in nitrogen (N) excretion in the stable.

### CH<sub>4</sub> implied emission factors (IEF) for Manure management

The CH<sub>4</sub> IEF for Manure management is calculated annually for all animal categories. A Tier 2 approach is used based on country-specific data on animal manure production per animal, on manure characteristics (such as organic matter (OM) content) and (liquid) manure storage conditions). For more information on methodology, see Section 6.3.2 and 6.3.5.



CH<sub>4</sub> implied emission factor (kg/head.year) for Manure management as specified by animal category, 1990–2008

Table 6.5

Animal type	1990	1995	2000	2005	2006	2007	2008
<i>Cattle</i>							
- mature dairy cattle	27.70	30.48	33.15	37.50	38.34	39.19	37.50
- mature non-dairy cattle	3.23	3.53	3.45	3.45	3.45	3.45	3.45
- young cattle	7.70	8.22	7.18	6.63	6.52	6.42	6.86
<i>Swine*</i>	3.90	4.43	4.61	4.50	4.46	4.42	4.43
<i>Swine excl piglets</i>	6.22	7.25	7.55	7.54	7.55	7.55	7.54
- fattening pigs	4.97	6.08	6.32	6.32	6.32	6.32	6.32
- breeding swine	11.39	12.24	12.86	12.96	12.99	12.95	13.38
<i>Poultry</i>	0.14	0.10	0.06	0.03	0.03	0.03	0.02

\* IEF is calculated on basis of total pig numbers, including piglets. However, manure production by piglets is accounted for in manure production by adult breeding swine.

Table 6.5 shows the implied emission factors for Manure management specified by the animal categories that contribute the most to CH<sub>4</sub> emissions.

### Trends in IEF

#### *Mature dairy cattle*

The IEF for manure management of mature dairy cattle increased between 1990 and 2008 because the increased milk production in that period (Table 6.4) is accompanied by an increase in manure production per cow and an increase in organic matter content of cattle manure. Both developments result from a higher feed intake. A third development concerns the shift in the proportion of the two dairy manure management systems (liquid manure in the stable and manure production in the meadow). The share of the amount of liquid stable manure increased between 1990 and 2008, while simultaneously the amount of manure produced in the meadow during grazing was reduced (Annex 8, Table A8.8). This is a consequence of the increase of the average time period dairy cattle are kept indoors. An explanation for this is the increase in average farm size. Since large herds are difficult to collect for indoor milking, farmers tend to keep the animals indoors for 365 days per year. With stable manure showing a 17-fold higher emission factor for CH<sub>4</sub> emissions, the shift to more stable manure increased the methane emission per cow (Annex 8, Table A8.7; Van der Hoek and Van Schijndel, 2006).

In short, between 1990 and 2008 the increase in the manure production per cow and in the organic matter content of dairy cattle manure combined with a shift to more stable manure resulted in an increased methane emission from manure management per cow.

#### *Poultry*

For poultry, the substantial decrease in CH<sub>4</sub> IEF of manure management between 1990 and 2008 mainly explains the CH<sub>4</sub> emission decrease. This decrease can be explained by a shift in the proportion of the two poultry manure management systems (solid and liquid manure) in this period. The proportion of the solid manure system increased between 1990 and 2008 from approximately 40% to more than 95%. So the liquid manure system was almost completely replaced by the solid manure system. Compared to the liquid manure system the CH<sub>4</sub> emission factor for the solid system is about

15-fold lower (Annex 8, Table A8.7). Overall, this leads to a substantially decreased IEF, which in combination with only a slight increase in animal numbers fully explains the decrease in CH<sub>4</sub> emissions (Van der Hoek and Van Schijndel, 2006).

#### *Swine*

Compared to 1990, the IEF of swine manure management (based on total swine numbers, including piglets), increased in 1993 and 1997 as a result of storage of manure under higher temperature (increased storage capacity below stable) and in 1995 due to increasing Volatile Solids (Annex 8, Table A8.4 and A8.5). There are inter-annual changes not explained by this. These changes can be explained by looking at emissions factors of underlying swine categories. The calculation method for CH<sub>4</sub> emissions from swine manure management is based on the liquid manure production of adult breeding swine (in which manure production by piglets is accounted for). So presenting the underlying IEFs gives a better understanding of the inter-annual changes.

For fattening pigs the 22% increase in IEF between 1990 and 1995 is explained by a 4% decrease in manure production per animal combined with a 20% increase in organic matter (OM) content of the manure and a higher storage temperature. The 4% increase in IEF between 1995 and 2000 is explained by an 8% increase due to higher storage temperature counteracted by a 4% decrease in manure production per animal. These manure volume changes are mainly the result of a change in liquid manure handling. In order to decrease the liquid manure volume, the mixing of rinsing water with manure was prevented as much as possible. As a consequence not only manure volume decreased, but also an increase in the OM concentration of manure occurred. A higher OM content results in a higher emission factor.

The inter-annual changes in the IEF for breeding pigs' manure are explained by inter-annual changes in the relative amount of different swine categories. Furthermore, between 1999 and 2000 a 2% decrease in manure production per animal occurred as a result of a change in liquid manure handling. In order to decrease the manure volume, the mixing of rinsing water with manure was prevented as much as possible.

For more details see Van der Hoek and Van Schijndel (2006) and Annex 8, Tables A8.4 to A8.8.

**N<sub>2</sub>O implied emission factor for Manure management and total N-excretion per animal manure management system, 1990-2008 (Units: mln kg/year and kg N<sub>2</sub>O/kg manure)**

**Table 6.6**

	1990	1995	2000	2005	2006	2007	2008
Total N-excretion	506.5	508.1	424.5	385.6	382.5	398.6	411.6
-liquid system	445.2	438.5	350.5	312.7	310.8	325.4	335.4
-solid storage	61.3	69.6	74.1	72.9	71.6	73.2	76.2
N <sub>2</sub> O emission manure management	2.63	2.88	2.88	2.78	2.74	2.81	2.92
N <sub>2</sub> O IEF manure management	0.0052	0.0057	0.0068	0.0072	0.0072	0.0071	0.0071

#### Comparison with IPCC default methane emission factor

The emission factors per animal type used by the Netherlands cannot be compared directly to the IPCC default values because of the assumptions on the share of the different animal manure management systems underlying the IPCC defaults.

Also the values of one of the underlying parameters per manure management system, Volatile Solids (VS), also called Organic Matter (OM) per animal type are not directly comparable. The Netherlands approach differs from the IPCC method in that the Netherlands uses the VS content of the manure (kg VS per kg manure) instead of volatile solids VS produced per animal per day (kg per head per day) in the IPCC calculation equations. By multiplying the VS per kg manure with the manure production per year, the annual VS production in manure in the Netherlands can be compared with the annual VS production underlying the default IPCC emission factors. More details are presented in Annex 8. Compared to the IPCC default MCF values, the Netherlands MCF values for liquid manure systems of swine (1990-1996) and cattle are slightly lower because part of the manure is stored under cooler conditions. For solid manure systems, the Netherlands uses a MCF of 1.5% for all animal categories (see Section 6.3.2); for manure production in the meadow, it uses the IPCC default MCF value.

#### N<sub>2</sub>O implied emission factor (IEF) for Manure management

Emission factors for N<sub>2</sub>O from Manure management represent the IPCC default values for liquid and solid management systems, 0.001 and 0.02 respectively.

Table 6.6 shows that the N<sub>2</sub>O emissions from Manure management increased between 1990 and 2008, mainly as a consequence of the increase in the N<sub>2</sub>O IEF. The explanation is that between 1990 and 2008 the proportion of the total solid manure N excretion increased. Compared to the liquid manure system, the N<sub>2</sub>O emission factor for the solid system is 20-times higher, which explains the increased overall IEF. This increased IEF was not fully counteracted by the decrease in the total N excretion and therefore has led to a small increase in N<sub>2</sub>O emissions.

The N<sub>2</sub>O emissions of solid manure decreased in 2003 as a direct result of the decrease in poultry manure production. This decrease was due to the reduction in poultry numbers that followed the avian flu epidemic (see also Section 6.1, Table 6.2).

#### 6.3.4 Methodological issues

##### Methane emissions from animal manure

A Tier 2 approach is followed for CH<sub>4</sub> emission calculations. The amounts of manure (in kg) produced are calculated annually for every manure management system per animal category. Country-specific CH<sub>4</sub> emission factors are calculated for all three manure management systems for every animal category on a Tier 2 level. These calculations are based on country-specific data on:

- manure characteristics: organic matter (OM) and maximum CH<sub>4</sub> producing potential (Bo)
- manure management system conditions (storage temperature and period) for liquid manure systems, which determine the methane conversion factor (MCF).

The amount of manure produced is calculated by multiplying manure production factors (in kg per head per year) by animal numbers. Detailed descriptions of the methods can be found on the website [www.greenhousegases.nl](http://www.greenhousegases.nl). More specified data on Manure management are based on statistical information on manure management systems found at [www.cbs.nl](http://www.cbs.nl) (Van Bruggen, 2009). These data are also documented in Van der Hoek and Van Schijndel (2006) and in Annex 8, Table A8.8.

For the methane conversion factor (MCF) for solid manure systems and manure produced in the meadow, IPCC default values are used. The IPCC guidelines recommend a MCF value of 0.01 for stored solid cattle manure and MCF = 0.015 for stored solid poultry manure. However, literature shows that CH<sub>4</sub> emissions from stored solid cattle manure are probably higher (see Van der Hoek and Van Schijndel, 2006). For this reason, the Netherlands set the MCF value for stored solid cattle manure equal to the MCF for stored solid poultry manure. The IPCC guidelines recommend a MCF value of 0.01 for manure produced in the meadow. This value is used in the CH<sub>4</sub> emission calculations.

Although the approach of the method applied by the Netherlands for CH<sub>4</sub> calculations differs slightly from the IPCC method, it is in accordance with the IPCC GPG. The Netherlands uses a country-specific emission factor for a specific animal category, which is expressed as the amount of CH<sub>4</sub> emitted per kg animal manure per year, whereas in the IPCC method the emission factor is expressed as the amount of methane (in kg) emitted per animal per year.

Since the CH<sub>4</sub> emissions from manure management from cattle, swine and poultry are key sources (see Table 6.1), the

present country-specific Tier 2 methodology fully complies with the IPCC Good Practice Guidance (IPCC, 2001).

#### Nitrous oxide emissions from animal manure

For the manure management systems and animal categories distinguished, the total N content of the manure produced – also called N excretion – (in kg N) is calculated by multiplying N excretion factors (kg .y<sup>-1</sup> per head) and animal numbers. Activity data are collected in compliance with a Tier 2 method. However, N<sub>2</sub>O emission factors used for liquid and solid manure management systems are IPCC defaults. The method used is fully in compliance with the IPCC Good Practice Guidance (IPCC, 2001), which is required for this key source. N<sub>2</sub>O emissions from manure produced in the meadow during grazing are not taken into account in the source category manure management. In accordance with the IPCC guidelines, this source is included in the source category agricultural soils (see Sections 6.1 and 6.4).

#### 6.3.5 Uncertainty and time-series consistency

##### Uncertainty

The Tier 1 uncertainty analysis shown in Annex 7 provides estimates of uncertainty according to IPCC source categories. The uncertainty in the annual CH<sub>4</sub> and N<sub>2</sub>O emissions from manure management from cattle and swine is estimated to be approximately 100%. The uncertainty in the amount of animal manure (10%) is based on a 5% uncertainty in animal numbers and a 5–10% uncertainty in excretion per animal. The resulting uncertainty of 7–11% was rounded off to 10%. The uncertainty in the CH<sub>4</sub> emission factors for Manure management, based on the judgments of experts, is estimated to be 100% (Olivier et al., 2009).

##### Time-series consistency

A consistent methodology is used throughout the time-series. The time-series consistency of the activity data is very good due to the continuity in the data provided.

#### 6.3.6 Source-specific QA/QC

This source category is covered by the general QA/QC procedures, discussed in Chapter 1.

#### 6.3.7 Source-specific recalculations

In 2005, an error was made while a new method to calculate the feed intake by cattle was applied. Feed intake is the basis for the calculation of the methane emission factor from enteric fermentation by cattle and also N excretion. All these errors have been corrected in the NIR 2009. Recalculation uncovered a few errors in volume excretion for the years 1990–2000. These errors are corrected in this NIR. The effect of the recalculation is minor and has resulted to a very small increase of the CH<sub>4</sub> emission in 1990 (0.2 Gg CH<sub>4</sub>).

#### 6.3.8 Source-specific planned improvements

A possible technical measure to prevent methane emissions due to Manure management is manure treatment in an anaerobic digester. In 2007, 0.6% of the total liquid stable manure has been treated in an anaerobic digester is ([www.cbs.nl](http://www.cbs.nl)). The Netherlands will examine future needs and possibilities in this area to include anaerobic treatment in the methodology and to extend calculations.

### 6.4 Agricultural soils [4D]

#### 6.4.1 Source category description

In the Netherlands, this source consists of the N<sub>2</sub>O source categories specified in Table 6.1:

- Direct soil emissions from the application of synthetic fertilisers, animal manure and sewage sludge to soils, and from N-fixing crops, crop residues and the cultivation of histosols (4D1);
- Animal production – animal manure produced in the meadow during grazing (4D2);
- Indirect emissions from N leaching and run-off, and from N deposition (4D3).

#### 6.4.2 Overview of shares and trends in emissions

In 2008, agricultural soils contributed 46% to the total greenhouse gas emissions in the agricultural sector. Direct and indirect N<sub>2</sub>O emissions and emissions from animal production in the meadow contributed 26%, 16% and 3%, respectively, to the total greenhouse gas emissions in the agricultural sector.

Total N<sub>2</sub>O emissions from Agricultural soils decreased by 24% between 1990 and 2008 (see Figure 6.3). Direct emissions increased by 3%, while indirect emissions and emissions from animal manure produced in the meadow decreased 39 and 56%, respectively.

This decrease is caused by a relatively high decrease in N-input to soil (from manure and chemical fertiliser application and animal production in the meadow), partly counteracted by the increased IEF in this period that resulted from a shift from the surface spreading of manure to the incorporation of manure into soil as a result of ammonia policy.

#### 6.4.3 Key sources

Both direct and indirect N<sub>2</sub>O soil emissions are level and/or trend key sources (see Table 6.1).


#### 6.4.4 Activity data and (implied) emission factors

Detailed information on data sources (for activity data and emission factors) can be found in the following monitoring protocols:

- Protocol 0070: N<sub>2</sub>O from Agricultural soils: indirect emissions (4D)
- Protocol 0071: N<sub>2</sub>O from Agricultural soils: direct emissions and emissions from animal production (4D)

More details and specific data (activity data and emission factors), including data sources (emission factors), are documented in background documents. All relevant documents concerning methodology, emission factors and activity data are published on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

The calculation of N<sub>2</sub>O emissions from Agricultural soils is based on various activity data, for example animal numbers (see Section 6.1) and nitrogen flows. For an overview of data sources, see the protocols or the background document (Van der Hoek et al., 2007). The activity data and emission factors can also be found in Annex 8, Tables A8.10, A8.11 and A8.12.



### Nitrogen flows

Table 6.7 present the N flows from synthetic fertilisers consumption and from animal manure production and application in the Netherlands (Annex 8, Table A8.10 gives N flows for all years). 80-85% of the manure N collected in the stable and in storage is applied to soils. A small portion of the manure N (approximately 1-7%) is exported; while approximately 12-15% is emitted as ammonia during storage.

The total amount of gross N available for soil (total manure production and fertiliser minus net export) applied to agricultural soils (including production of animal manure in the meadow) decreased by approximately 36% between 1990 and 2008. This is explained by the Netherlands manure and fertiliser policy, aimed at reducing N leaching and run-off. This policy regulates the amount of manure production and its application by the introduction of measures such as pig and poultry production rights and maximum nutrient application standards for manure and fertiliser.

Of the manure N applied to the soil between 1990 and 2008 the part emitted as ammonia (NH<sub>3</sub>) decreased from 23 to 8%, due to a change in the method of animal manure

application to agricultural soils. Before 1991 manure was applied to the soil by spreading on the surface of grasslands and arable land. Initiated by the Netherlands' policy to reduce ammonia emissions, this practice changed in 1991 into manure incorporation into the soil (e.g., shallow injection or ploughing in), resulting in lower NH<sub>3</sub> emissions. Ultimately, between 1990 and 2008 the part of the N in manure and synthetic fertiliser emitted as NH<sub>3</sub> (in the stable and during, storage, grazing and application to the field) decreased from approximately 18% to 13%.

About 30% of the total nitrogen flow to the soil is subject to leaching and run-off (default IPCC Frac<sub>leach</sub> factor).

The decrease in indirect N<sub>2</sub>O emissions is fully explained by the decrease in N lost by atmospheric deposition and by leaching and run-off. The decrease in N<sub>2</sub>O emissions from animal manure produced in the meadow is also entirely reflected in the decrease in N input to soil by this source. The 3% increase in direct N<sub>2</sub>O emissions can mainly be explained by the 36% decrease in the direct N input to soil by manure and chemical fertiliser application in combination with a 39% increase of the IEF.

	1990	1995	2000	2005	2006	2007	2008	Change 2008 - 1990
<i>Nitrogen fertiliser consumption</i>	412.4	405.8	339.5	279.2	287.8	257.5	238.1	-42%
<i>of which ammonium fertiliser</i>	3.6	11.2	6.6	30.6	42.9	17.2	12.8	258%
<i>NH<sub>3</sub>-N emission during application</i>	11.2	10.5	9.2	9.8	10.9	9.9	8.2	-27%
<i>Net fertiliser to soil</i>	401.1	395.3	330.3	269.4	276.9	247.6	229.9	-43%
<i>Nitrogen excretion by animals</i>	694.4	680.1	549.2	478.9	471.2	479.7	497.2	-28%
<i>Nitrogen excretion in animals houses</i>	506.5	508.1	424.5	385.6	382.5	398.6	411.6	-19%
<i>of which in solid form</i>	61.3	69.6	74.1	72.9	71.6	73.2	76.2	24%
<i>of which in liquid form</i>	445.2	438.5	350.5	312.7	310.8	325.4	335.4	-25%
<i>NH<sub>3</sub>-N emission in animal houses</i>	73.2	73.5	60.2	49.8	49.6	50.1	51.1	-30%
<i>Net available manure for application</i>	433.3	434.6	364.3	335.9	332.9	348.5	360.5	-17%
<i>Nitrogen in net manure exported abroad*</i>	6.4	22.1	14.7	14.9	15.8	28.8	30.5	377%
<i>NH<sub>3</sub>-N emission during application</i>	98.3	51.4	36.8	33.2	32.6	34.4	27.9	-72%
<i>Net animal manure to soil</i>	328.6	361.2	312.8	287.8	284.5	285.3	302.2	-8%
<i>Nitrogen excretion in meadow</i>	188.0	171.9	124.6	93.3	88.7	81.1	85.5	-54%
<i>NH<sub>3</sub>-N emission in meadow</i>	13.0	11.9	8.5	7.1	6.9	6.1	6.5	-50%
<i>Gross nitrogen available for soil (total manure prod. + fertiliser - export)</i>	1100.4	1063.8	873.9	743.2	743.2	708.4	704.8	-36%
<i>Nitrogen fixation in arable crops</i>	7.8	4.9	4.7	4.5	4.6	4.5	4.1	-48%
<i>Nitrogen in crop residues left in field</i>	36.4	34.9	34.1	32.1	30.1	29.6	26.8	-26%
<i>Nitrogen in histosols</i>	52.4	52.4	52.4	52.4	52.4	52.4	52.4	0%
<i>Nitrogen in sewage sludge on agric. land</i>	5.0	1.5	1.5	1.2	1.1	1.0	1.0	-80%
<i>Atmospheric deposition agr. NH<sub>3</sub>-N em</i>	195.9	147.5	115.1	100.3	100.3	100.9	94.0	-52%
<i>Nitrogen lost through leaching and run off</i>	330.1	319.1	262.2	223.0	223.0	212.5	211.4	-36%

\* export and import figures can be found in Annex 8, Table A8.10a

**Direct and animal production N<sub>2</sub>O implied emission factors for Agricultural soils by CRF category**  
(Units: kg N/kg N-input)

Table 6.8

	1990	1995	2000	2005	2006	2007	2008
<i>Nitrogen input from applic. of synthetic fertilisers</i>	0.011	0.011	0.011	0.010	0.010	0.010	0.010
<i>Nitrogen input from manure applied to soils</i>	0.011	0.020	0.020	0.020	0.020	0.020	0.020
<i>Nitrogen input from animal production</i>	0.016	0.016	0.015	0.015	0.015	0.015	0.015

#### Implied emission factor

Table 6.8 shows the implied emissions factors (IEF) for N<sub>2</sub>O emissions from Agricultural soils for the most important sources. For (direct) soil emissions by manure application to soil, a 77% increase in IEF occurred in the period 1990–2008, which is caused by an ammonia policy driven shift from the surface spreading of manure to the incorporation of manure into the soil. Combined with an 8% decrease in net N manure input to soil (see Table 6.7), this explains the 63% increase in N<sub>2</sub>O from manure application.

#### 6.4.5 Methodological issues

Direct and indirect N<sub>2</sub>O emissions from agricultural soils, as well as N<sub>2</sub>O emissions by animal production in the meadow are estimated using country-specific activity data on N-input to soil and NH<sub>3</sub> volatilisation during grazing, manure management (stable and storage) and manure application. Most of these data are estimated at a Tier 2 or Tier 3 level. The present methodologies fully comply with the IPCC Good Practice Guidance (IPCC, 2001). For a description of the

methodologies and data sources used, see the monitoring protocols on [www.greenhousegases.nl](http://www.greenhousegases.nl). A full description of the methodologies is provided in Van der Hoek et al. (2007), with more details in Kroeze (1994). An overview of the emission factors used is presented in Table 6.9. Default IPCC emission factors are included for comparison.

#### Direct N<sub>2</sub>O emissions

The IPCC Tier 1b/2 methodology is used to estimate direct N<sub>2</sub>O emissions for two soil types (organic and inorganic soils). Emissions from the application of synthetic fertiliser have been estimated for two types of synthetic fertilisers (ammonium phosphate/sulphate and other synthetic fertilisers). Emissions from animal manure application are estimated for two types of manure application methods (surface spreading and incorporation into soil).

The country-specific emission factors are lower for mineral soils (e.g., 0.01 kg N/kg N input) and higher for organic soils (0.02 kg N/kg N input) compared to the IPCC default of 0.0125



Source	Default IPCC	Mineral soils	Organic soils	Reference
Nitrogen fertiliser	0.0125			
Ammonium fertiliser		0.005	0.01	4
Other fertilisers		0.01	0.02	1,4
Animal manure application	0.0125			
Surface spreading		0.01	0.02	1
Incorporation into soil		0.02	0.02	1
Sewage sludge	0.0125	0.01		2
Biological nitrogen fixation crops	0.0125	0.01		1
Crop residues	0.0125	0.01		2
Cultivation of organic soils (histosols)			0.02	2,3
Animal manure during grazing	0.02			
Faeces		0.01	0.01	1
Urine		0.02	0.02	1

References 1 = Kroeze, 1994; 2 = Van der Hoek et al. (2007); 3 = Kuikman et al., 2005; 4 = Kuikman et al., 2006.

kg N/kg N -input. A higher emission factor of 0.02 kg N/kg N input is also used for manure incorporation into soil.

The higher value for incorporation is explained by two mechanisms. Incorporation of animal manure into the soil produces less ammonia emission and hence more reactive nitrogen enters the soil. Furthermore, the animal manure is more concentrated (e.g., hot spots) in comparison with surface spreading and hence the process conditions for nitrification and denitrification can be more suboptimal.

A review of the literature in 2005 showed that in most experiments with simultaneous surface spreading and incorporation the latter produces higher nitrous oxide emissions. It was, however, not possible to derive a new emission factor for incorporation or shallow (sod) injection (Kuikman et al., 2006). Therefore it was decided not to change the existing emission factors.

#### Animal production

The IPCC Tier 1b/2 methodology is used to estimate direct N<sub>2</sub>O emissions from animal production. The country specific method uses total animal production minus ammonia emission from pasture (Kroeze, 1994). For Animal production a distinction is made between N in urine and N in faeces. The country-specific emission factors are lower for faeces (0.01 kg N/kg net N input) compared to the IPCC default of 0.02 kg N/kg N-input. For urine, the emission factor is 0.02 kg N/kg net N input. This means that on gross N input, the emission factor is a little lower compared to the IPCC default of 0.02 kg N/kg gross N input. The emission factor for urine is higher than for faeces because the ratio mineral nitrogen/total nitrogen is higher in urine than in faeces, leading to faster nitrification and denitrification in urine-affected spots. Furthermore, urine penetrates faster into the soil than faeces, which enhances the lack of sufficient oxygen in the soil for the nitrification process. Together with the higher mineral nitrogen ratio in urine, it is clear that urine creates a higher potential for suboptimal conditions for nitrification and denitrification than faeces.

#### Indirect N<sub>2</sub>O emissions

The IPCC Tier 1 method is used to estimate indirect N<sub>2</sub>O emissions. Indirect N<sub>2</sub>O emissions resulting from atmospheric

deposition are estimated using country-specific data on NH<sub>3</sub> emissions (estimated at a Tier 3 level). IPCC default values are used for N<sub>2</sub>O emission factors because of the lack of country-specific data.

Indirect N<sub>2</sub>O emissions resulting from leaching and run-off N emissions are estimated using country-specific data on total N-input into soil (estimated at a Tier 2 level). IPCC default values are used for the fraction of N-input to soil that leaches from the soil and ends up partly as N<sub>2</sub>O emissions from groundwater and surface water (Fracleach) and for the N<sub>2</sub>O emission factors. The main reason to use IPCC defaults is that direct and indirect N<sub>2</sub>O emissions in the Netherlands partially originate from the same soils and sources. In the Netherlands, no experimental data are available to evaluate the value of the emission factor for indirect emissions.

#### 6.4.6 Uncertainty and time-series consistency

##### Uncertainty

The Tier 1 uncertainty analysis, shown in Annex 7, provides estimates of uncertainty according to IPCC source categories. The uncertainty in direct N<sub>2</sub>O emissions from Agricultural soils is estimated to be approximately 60%. The uncertainty in indirect N<sub>2</sub>O emissions from N used in agriculture is estimated to be more than a factor of 2 (Olivier et al., 2009).

##### Time-series consistency

Consistent methodologies are used throughout the time series. The time-series consistency of the activity data is very good due to the continuity in the data provided.

#### 6.4.7 Source-specific QA/QC

This source category is covered by the general QA/QC procedures discussed in Chapter 1.

#### 6.4.8 Source-specific recalculations

During the compilation of the CRF an error was detected in the N<sub>2</sub>O emission for 2007 was detected and removed. The emission decreased by 180.3 Gg CO<sub>2</sub> eq.

#### 6.4.9 Source-specific planned improvements

The specific characteristics of the Netherlands agricultural soils (with relatively high water tables) justify the calculation

of the “fracleach” and the emission factors on the basis of country-specific data. Therefore, the Netherlands will examine the needs and possibilities of extending calculations in the future in order to improve the methodology.

A higher emission factor than the IPCC default is used for the incorporation of manure into soil. However, the findings of a survey on N<sub>2</sub>O emission factors for the field-scale application of animal manure abroad did not provide the necessary underpinning for an update of long-term average N<sub>2</sub>O emission factors for this source in the Netherlands (Van der Hoek et al., 2007). Consequently, research has been carried out since 2007 to gain an insight into this. First results on the effect of manure application technique on N<sub>2</sub>O emission in field and laboratory experiments were recently presented (Velthof et al, 2010). In order to make a translation into adjustments of the emission factors further investigations are carried out now.



# Land use, land use change and forestry [CRF Sector 5]

## ■ Major changes in the LULUCF sector compared to the National Inventory Report 2009

*Emissions:* The emission data from LULUCF from 2008 are slightly lower than in 2007.

*Key sources:* No changes to NIR 2009.

*Methodologies:* Changes in methodology have not been made.

### 7.1 Overview of sector

This chapter describes the 2008 greenhouse gas inventory for the Land Use, Land Use Change and Forestry (LULUCF) sector. It covers both the sources and sinks of CO<sub>2</sub> greenhouse gases from land use, land use change and forestry. The emission of nitrous oxide (N<sub>2</sub>O) from land use is included in the “Agriculture” sector (category 4D) and the emission of methane (CH<sub>4</sub>) from wetlands is not estimated due to the lack of data. All other emissions from forestry and land use can be considered to be negligible.

Land use in the Netherlands is dominated by agriculture (57%), settlements (13%), forestry (10%, including trees outside forests) and 2% comprises dunes, nature reserves, wildlife areas and heather. The remaining area (19%) in the Netherlands is open water. The soils in the Netherlands are dominated by mineral soils, mainly sandy soils and clay soils (of fluvial or marine origin). Organic soils, used mainly as meadowland or hayfields, cover about 8% of the land area. The Netherlands has an intensive agricultural system with high inputs of nutrients and organic matter. The agricultural land is used as grassland (51%), arable (25%), fodder maize (12%) and the remaining agricultural land is used for horticulture, fallow, fruit trees, etc. Grassland and fodder maize are cultivated in rotation. About 80% of the grasslands are permanent grasslands (of which 5% are high nature value grasslands); the remaining 20% is temporary grassland. Since 1990, the agricultural land area has decreased by about 5%, mainly because of conversion to settlements/infrastructure and nature.

The LULUCF sector in the Netherlands is estimated to be a net source, amounting in 2008 to some 2.4 Tg CO<sub>2</sub> equivalents. The fact that the LULUCF sector is a net source is due to the large contribution of carbon emitted from drained peat soils,

which exceeds the sequestration of carbon in forestry. The LULUCF sector is responsible for 1.2% of total greenhouse gas emission in the Netherlands.

The structure of this Section and of the main submission for the National Inventory Report and Common Reporting Format (CRF) tables is based on the categories of the CRF tables at the 9th Conference of Parties to the United Nations Framework Convention on Climate Change (UNFCCC). The Sector 5 Report Tables in the CRF format have been submitted using the CRF Reporter.

### 7.2 Methods

The methodology of the Netherlands to assess the emission from LULUCF is based on the IPCC 1996 Revised Guidelines and its updates in the Good Practice Guidance: a carbon stock change approach based on inventory data subdivided into appropriate pools and land use types and a wall-to-wall approach for the estimation of area per category of land use. The information on the activities and land use categories used covers the entire territorial (land and water) surface area of the Netherlands.

The inventory comprises six classes: Forest Land; Cropland; Grassland; Wetlands; Settlements and Other Land. There is also a category “Other” which includes emissions from land use related activities such as liming. The changes in land use (“remaining” or “converted”) are presented in a 6 x 6 matrix, which is fully in accordance with the approach described in the IPCC guidelines. To better match available national maps and databases on land use, the category “Forest Land” is the aggregation of two main subdivisions: Forest (according to the Kyoto definition) and Trees outside Forest, and the category “Grassland” is the aggregation of two main

subdivisions: Grasslands and Nature. The latter subdivision includes heather, peat land and moors. All categories are relevant in the Netherlands.

The carbon cycle of a managed forest and wood production system is considered in the calculations of the relevant CO<sub>2</sub> emissions. The carbon stocks in soils from a single stratified measurement campaign for the various types of land use are used to calculate the emissions from land use categories. For the Netherlands, it is assumed that the impact of land use in terms of loss of soil carbon is likely to be relatively small. Simulation shows that large quantities of carbon accumulate in soil. We have assumed no changes in the carbon stocks due to land and soil management and cultivation practices over the period 1990–2008. This is a conservative approach. The research that is ongoing aims to underpin whether land conversions are a source of soil carbon or not, and is not directed to quantifying the size of the sinks in the Netherlands.

### 7.3 Data

In this NIR, the changes in land use are based on comparing detailed maps that best represent land use in 1990 and 2004. Both datasets on land use were especially developed to support the temporal and spatial development in land use and especially designed to support policy in the field of nature conservation (MNP, 2008). In the future, updates of the digital land use map will become available regularly and these will suite the future LULUCF process in their aim to present accurate information on land use changes. Changes in land use over the period 1990–2004 were checked in detail (Kramer, 2009). Omissions due to methodological reasons (e.g., legend, classification and gridding) were manually adjusted in favour of a correct presentation of the changes in land use over the period 1990–2004. The sum of all land use categories is constant over time. It is likely that the updated reference maps will also follow future updates of the land use change matrix. Changes after 2004 have been obtained by linear extrapolation.

### 7.4 Recalculations

This year, there were no recalculations.

The methodologies applied for estimating CO<sub>2</sub> emissions and removals of the land use change and forestry in the Netherlands are described in the updates of the two protocols (see also the website at [www.greenhousegases.nl](http://www.greenhousegases.nl)):

- Protocol 0072: CO<sub>2</sub> from forest (5A)
- Protocol 0073: CO<sub>2</sub> from total land use categories (5B-5G)

Table 7.1 shows the sources and sinks in the LULUCF sector in 1990 and 2008. For 1990 and 2008, the total net emissions are estimated to be approximately 2.6 Tg CO<sub>2</sub> and 2.4 Tg CO<sub>2</sub> respectively, with the major source being CO<sub>2</sub> emissions from the decrease in carbon (C) stored in organic soils and peat lands: 4.2 Tg CO<sub>2</sub>, included in 5C1 “Grassland remaining grassland”, resulting from agricultural and water management. The major sink is the storage of carbon in

forests: - 2.8 Tg CO<sub>2</sub>, this includes the emissions from “Forest Land remaining forest land” (5A1) and “Land converted to Forest Land” (5A2). Sector 5 “Land use, land use change and forestry” (LULUCF) accounted for 1.2% of the total national CO<sub>2</sub> emission in 2008.

## 7.5 Forest Land [5A]

### 7.5.1 Source category description

This category includes emissions and sinks of CO<sub>2</sub> caused by changes in forestry and other woody biomass stock. All forests in the Netherlands are classified as temperate forest, with 30% of the forests being coniferous, 22% broad-leaved and the remaining area a mix of both. The share of mixed and broad-leaved forests has grown in recent decades (Dirkse et al., 2003).

The category includes two subcategories: 5A1 “Forest Land remaining Forest Land” and 5A2 “Land converted to Forest Land”. The first category includes estimates of changes in the carbon stock from different carbon pools in the forest. The second category includes estimates of the changes in land use from mainly agricultural areas into forest land since 1990.

### 7.5.2 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

The methodology of the Netherlands to assess the emission from LULUCF is based on a wall-to-wall approach for the estimation of area per category of land use. For the wall-to-wall map overlay approach were used harmonised and validated digital topographical maps of 1990 and 2004 (Kramer et al., 2009). The result was a national scale land-use and land-use change matrix. The information on the activities and land-use categories used covers the entire territorial (land and water) surface area of the Netherlands.

### 7.5.3 Definition

The land use category “Forest Land” is defined as all land with woody vegetation consistent with thresholds used to defined forest land in the national GHG inventory, sub divided into managed and unmanaged units and also by ecosystem type as specified in IPCC Guidelines. It also includes systems with vegetation that currently fall below, but are expected to exceed the threshold of the forest land category (IPCC, 2003; 2006).

The Netherlands has chosen to define the land-use category “Forest Land” as all land with woody vegetation, now or expected in the near future (e.g., clear-cut areas to be replanted, young afforestations). This is further stratified in:

- “Forest” or “Forest according to the Kyoto definition” (FAD), - all forest land which complies to the following (more strict than IPCC) definition chosen by the Netherlands for the Kyoto protocol: forests are patches of land exceeding 0.5 ha with a minimum width of 30 m, with tree crown cover at least 20% and tree height at least 5 m, or, if this is not the case, these thresholds are likely to be achieved at the particular site. Roads in the forest less than 6 m wide are also considered to be forest. This definition

Sector/category	Gas	Key	Emissions base year (1990)	Emissions 2007	Emissions 2008	Change 2008-2007	Contribution to total in 2008 (%)		
			Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Gg CO <sub>2</sub> eq	By sector	Of total gas	Of total CO <sub>2</sub> eq
Key sources		Level, Trend							
5. Total land use categories	CO <sub>2</sub>		2.6	2.5	2.4	-0.1	100	1.4	1.2
5A. Forest land	CO <sub>2</sub>		-2.5	-2.7	-2.8	-0.1	-116		
5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	L,T	-2.5	-2.2	-2.2	0.0	-90		
5A2. Land converted to Forest Land	CO <sub>2</sub>	L,T	0.0	-0.6	-0.6	-0.1	-26		
5B. Cropland	CO <sub>2</sub>		0.0	0.0	0.0	0.0	2		
5B1. Cropland remaining Cropland	CO <sub>2</sub>					0.0	0		
5B2. Land converted to Cropland	CO <sub>2</sub>		0.0	0.0	0.0	0.0	2		
5C. Grassland	CO <sub>2</sub>		4.6	4.8	4.8	0.0	196		
5C1 Grassland remaining Grassland	CO <sub>2</sub>	L	4.2	4.2	4.2	0.0	174		
5C2. Land converted to Grassland	CO <sub>2</sub>	L2,T2	0.4	0.5	0.6	0.0	22		
5D. Wetlands	CO <sub>2</sub>		0.0	0.1	0.1	0.0	2		
5D1. Wetlands remaining Wetlands	CO <sub>2</sub>		NE	NE	NE				
5D2. Land converted to Wetlands	CO <sub>2</sub>		0.0	0.1	0.1	0.0	2		
5E. Settlements	CO <sub>2</sub>		0.2	0.3	0.3	0.0	12		
5E1. Settlements remaining Settlements	CO <sub>2</sub>					0.0			
5E2. Land converted to Settlements	CO <sub>2</sub>		0.2	0.3	0.3	0.0	12		
5F. Other Land	CO <sub>2</sub>		0.0	0.0	0.0	0.0	1		
5F1. Other Land remaining Other Land	CO <sub>2</sub>		0.0	0.0	0.0	0.0	0		
5F2. Land converted to Other Land	CO <sub>2</sub>		0.0	0.0	0.0	0.0	1		
5G. Other	CO <sub>2</sub>		0.2	0.1	0.1	0.0	3		
Total National CO <sub>2</sub> Emissions (incl. CO <sub>2</sub> LULUCF)	CO <sub>2</sub>		161.9	174.8	178.1	3.3			
	All		215.9	209.5	209.4	-0.1			

conforms to the FAO reporting and was chosen within the ranges set by the Kyoto protocol.

- “Trees outside Forests” (TOF), that is - wooded areas that comply with the previous forest definition except for their surface ( $\leq 0.5$  ha or less than 30 m width). These represent fragmented forest plots as well as groups of trees in parks and nature terrains and most woody vegetation lining roads and fields. These areas comply with the GPG-LULUCF definition of Forest Land (they have woody vegetation) but not to the strict forest definition that the Netherlands applies.

#### 7.5.4 Methodological issues

Removals and emissions of CO<sub>2</sub> from changes in forestry and woody biomass stock are estimated based on country-specific Tier 2 methodology. The approach chosen follows the IPCC 1996 Revised Guidelines and its updates in the Good Practice Guidance on Land Use, Land Use Change and Forestry (IPCC, 2003). The basis assumption is that the net flux can be derived from converting the change in growing stock volume in the forest into carbon. Detailed descriptions of the methods used and emission factors can be found in the protocol 0072 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 7.1.

The Netherlands' National System follows the carbon cycle of a managed forest and wood products system. The pools are distinguished by above-ground biomass, below-ground biomass, litter, dead wood, and soil organic carbon. Changes in the carbon stock are calculated for above-ground biomass, below-ground biomass and dead wood and litter in forests. Calculations for the living biomass carbon balance are carried out at plot level.

#### Living biomass

The following steps are taken to calculate the net carbon flux in living biomass. First, the age of the stand and the limit of dominant height are calculated, followed by a calculation of the height and expected volume in the next year. Based on the expected volume for the next year and from the number of trees, the average tree volume for the next year is derived. The next step is the calculation of the average diameter of the tree in the next year. The above-ground and below-ground total biomass is derived using the equations from the COST E21 database. The desired net flux is derived from the difference in tree mass between two years, the basic wood density and the carbon content of the dry mass. This last step is represented in the following equation:

$$\Delta C_{FFG} = \sum_i^n (A_i \cdot G_{TOTALi}) \cdot CF$$

$$G_{TOTALi} = (\overline{B_{i+1}} - \overline{B_{it}}) \cdot nt_{it}$$

where:

$\Delta C_{FFG}$  Total net carbon emission due to biomass increase for Forest

land remaining Forest land – FAD in The Netherlands kg C ha<sup>-1</sup>

$A_i$  Area represented per NFI plot ha

$CF$  Carbon fraction of living biomass 0.5

and

$G_{TOTALi}$	Biomass increase for NFI plot i	kg DW
$\overline{B}_{it}$	Average tree biomass of NFI plot i at time t	kg DW
$\overline{B}_{it+1}$	Average tree biomass of NFI plot i at time t+1	kg DW
$nt_{it}$	Living tree density of NFI plot i at time t	ha <sup>-1</sup>

### Thinning

Thinning was carried out in all plots that met the criteria for thinning (age > 110 years or growing stock more than 300 m3 ha<sup>-1</sup>). The number of trees thinned was based on the volume harvested, and the net carbon flux due to thinning is then calculated from the average biomass of a single tree and the carbon content of the dry mass.

### Deadwood

The net carbon flux to dead wood is calculated as the remainder of the input of dead wood due to mortality minus the decay of the dead wood. Leaves and roots were not taken into account for the build up of dead wood. The mortality rate was assumed to be a fixed fraction of the standing volume (0.4% year<sup>-1</sup>), and the current stock of dead wood volume is assumed to be 6.6% of the living wood volume (based on data from Timber Production Statistics and Forecast (HOSP) and the MFV). A net build up may exist, since Dutch forestry only began to pay attention to dead wood a decade ago.

The following equations are used to calculate the net carbon flux to dead wood:

$$\Delta C_{FFDW} = \sum (A_i \cdot (B_{DWint_{oi}} - B_{DWout_{oi}})) \cdot CF$$

$$B_{DWint_{oi}} = B_{it} \cdot f_{mort}$$

$$B_{DWout_{oi}} = \left( \frac{V_{SDi}}{L_{SDi}} + \frac{V_{LDi}}{L_{LDi}} \right) \cdot D_{DW}$$

$\Delta C_{FFDW}$  Total net carbon emission due to change in dead wood for Forest land remaining Forest land – FAD in the Netherlands

$B_{DWint_{oi}}$  Annual mass transfer into dead wood pool of NFI plot i

$B_{DWout_{oi}}$  Annual mass transfer out of dead wood pool of NFI plot i

$B_{it}$  Stand living biomass of NFI plot i at time t

$f_{mort}$  Mortality fraction (0.4% year<sup>-1</sup>)

$V_{SDi}$  Volume of standing dead wood of NFI plot i

$V_{LDi}$  Volume of lying dead wood of NFI plot i

$L_{SDi}$  Species specific longevity of standing dead wood

$L_{LDi}$  Species specific longevity of standing lying wood

$D_{DW}$  Species specific average wood density of dead wood

### Litter

The carbon stock change from changes in the litter layer was estimated using a stock change method at national level. Data for litter layer thickness and carbon in litter were available from five different datasets. None of these datasets could be used exclusively. Therefore, a stepwise approach was used to estimate the national litter carbon stock and change therein in a consistent way. Additional, selected forest stands, on poor and rich sands, were intensively sampled with the explicit purpose to provide conversion factors or functions.

None of the available datasets could be used exclusively. Therefore, a stepwise approach was used to estimate the national litter carbon stock and change therein in a consistent way. After which a hierarchy was developed to accord mean litter stock values to any of the sampled plots of the available forest inventories (HOSP and MFV). The difference between 2004 and 1990 was estimated and a mean annual rate of carbon accumulation was calculated. A Monte Carlo uncertainty analysis was carried out and showed that the result was considered the more conservative.

### 7.5.5 Activity data

Activity data on land use and land use change are derived from the land use maps and the land use change matrix. Carbon content is based on the soil map of the Netherlands (scale 1:50,000) combined with results of LSK, a national random check of map units that provides detailed descriptions of soil profiles. The random check was implemented both nationwide and on a stratified scale, combining main categories and/or symbol units in order to produce a more homogenous classification with respect to landscaping, soil formation or parent material. Within this framework, this random check was meant to provide further quantitative information for the existing soil maps.

Activity data on forests is based on forest inventories carried out in 1988–1992 (HOSP data) and in 2001–2002 and 2004–2005 (MFV data). HOSP data, which includes plot level data (in total 2007 plots, about 400 per year) for growing stock volume, increment, age, tree species, height, tree number and dead wood, was used for the 1990 situation. Forward calculation using this data was applied to the year 1999. Additional data on felling, final cut and thinning was used to complete the data set. MFV plot level data (in total 3622 plots, with same items as HOSP) was applied to the years 2000–2004. In addition, in order to assess the changes in activity data, databases with tree biomass information, with allometric equations to calculate above-ground and below-ground biomass and with forest litter, as well as wood harvest statistics, soil carbon estimations and high-resolution topographical maps of 1990 and 2004 were used. See the website at [www.greenhousegases.nl](http://www.greenhousegases.nl) for more details on activity data.

	1990	1995	2000	2005	2008
5A Forest Land	-2,532	-2,798	-2,670	-2,871	-2,847
5A1 Forest Land remaining Forest Land of which	-2,529	-2,739	-2,448	-2,414	-2,208
Live trees	-3,745	-3,509	-3,505	-3,308	-3,337
Harvest	1,745	1,257	1,528	1,337	1,440
Trees outside Forest	-211	-180	-160	-132	-116
Dead wood	-307	-307	-311	-312	-311
5A2 Land converted to Forest Land (Afforestation)	-3	-59	-222	-457	-639

### 7.5.6 Implied emission factors

The total emissions from the tree component after deforestation is calculated by multiplying the total area deforested with the average carbon stock in living biomass, above- as well as below ground (Nabuurs et al., 2005) and the average carbon stock in dead organic matter and litter. Thus it is assumed that with deforestation, all carbon stored in above- and below ground biomass as well as in dead wood and litter is lost to the atmosphere. National averages are used as there is no record of the spatial occurrence of specific forest types.

The average carbon stock in living biomass follows the calculations from the gap filled forest inventory data. The calculated emission factors show a progression over time. The EF for biomass is 60.4 Mg C ha<sup>-1</sup> in 1990 and increase to 85.3 Mg C ha<sup>-1</sup> in 2008. The EF for litter is 29.0 Mg C ha<sup>-1</sup> in 1990 and increase to 35.9 Mg C ha<sup>-1</sup> in 2008 and the EF for dead wood is 0.45 Mg C ha<sup>-1</sup> in 1990 and increase to 3.26 Mg C ha<sup>-1</sup> in 2008. The systematic increase in average standing carbon stock reflects the fact that annual increment exceeds annual harvests in the Netherlands.

The IEF for biomass increase in land converted to either FAD or TOF reflects the age distribution of the re/afforested areas and will attain a constant value 20 years after 1990.

### Non CO<sub>2</sub> emissions in forest land

N<sub>2</sub>O emissions might occur as a result of using fertiliser in forests or from drainage. Both management practices are rarely applied in forestry in the Netherlands. Thus, it is assumed that N<sub>2</sub>O emissions are irrelevant in forests. CH<sub>4</sub> emissions resulting from forest fires are considered to be negligible because fires seldom occur.

### 7.5.7 Uncertainty and time-series consistency

#### Uncertainties

The Tier 1 analysis in Annex 7 shown in Table A7.1 provides estimates of uncertainties according to IPCC source category. The Netherlands uses a Tier 1 analysis for the uncertainty assessment of the sector LULUCF. The analysis combines uncertainty estimates of the forest statistics, land use and land use change data (topographical data) and the method used to calculate the yearly growth in carbon increase and removals. The uncertainty in the CO<sub>2</sub> emissions from 5A1 “Forest Land remaining Forest Land” is calculated at 67%. The uncertainty in the CO<sub>2</sub> emission from 5A2 “Land converted to Forest Land” is calculated at 63%. See Olivier et al. (2009) for details.

The uncertainty in implied emission factors of 5A1 “Forest Land remaining Forest Land” concerns forest and trees outside the forest. As the methodology and data sets used are the same for both sources, the uncertainty calculation is performed for forests and the result is considered to be representative for trees outside forests as well.

The uncertainty in the implied emission factor of increment in living biomass is calculated at 13% (rounded off to 15% in the calculation spreadsheet). The uncertainty in the implied emission factor of decrease in living biomass is calculated at 30%. The uncertainty in the net carbon flux from dead wood is calculated at 30% (rounded off to 50% in the Tier 1 calculation spreadsheet)

#### Uncertainty in implied emission factor of 5A2 ‘Land converted to Forest Land’

For the increment in living biomass, the same data and calculations are used as for 5A1 “Forest Land remaining Forest Land” and, therefore, the same uncertainties are used in the Tier 1 calculation spreadsheet.

For soil carbon stock changes after land use change it is assumed that the average carbon stock in the soil under the new and old land use is the same (Groot et al., 2005). Therefore, the uncertainty is the uncertainty of the change in carbon content in mineral soil, which is calculated at 38% (rounded off to 50% in the Tier 1 calculation spreadsheet); see Section 7.3.3.

#### Uncertainty in activity data in categories 5A1 and 5A2

The activity data used is area change, calculated by comparing two topographic maps. The uncertainty of one topographic map is estimated at 5% (expert judgment). Therefore, the uncertainty in comparing two topographic maps is theoretically 5 × 5 = 25%. This is without doubt an overestimation, as not all land use may change over a decade.

#### Time-series consistency

The updated time series for category 5A shows an average of about 2,800 Gg CO<sub>2</sub> yr<sup>-1</sup> and with a range from 2,500 Gg CO<sub>2</sub> yr<sup>-1</sup> to 3,000 Gg CO<sub>2</sub> yr<sup>-1</sup> over the period 1990-2008 (see Table 7.2). The figures in category 5A1 show the net result of the sequestration in live trees, in trees outside forest, dead wood and litter and the emission from harvest. The figures for live trees and harvest only change slightly over time, with no clear direction. The figures for afforestation have steadily increased since 1990 and reached in 2007 a sequestration level of 575 Gg CO<sub>2</sub> yr<sup>-1</sup>.



### 7.5.8 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures as discussed in Chapter 1.

### 7.5.9 Source-specific recalculations

This year there were no recalculations.

### 7.5.10 Category-specific planned improvements

No improvements planned.

## 7.6 Cropland [5B]

### 7.6.1 Source category description

The source category 5B “Cropland” includes only the emissions of CO<sub>2</sub> from 5B2 “Land converted into Cropland”.

The land use category “Cropland” is defined as all arable and tillage land, including rice-fields, and agro-forestry systems where the vegetation structure falls below the thresholds used for the Forest Land category (IPCC, 2003).

### 7.6.2 Activity data and (implied) emission factors

The activity data is derived from the land use maps and the land use change matrix. Carbon content is based on the soil map of the Netherlands (scale 1:50,000) combined with results of LSK, a national random check of map units that provides detailed descriptions of soil profiles. The random check was implemented both nationwide and on a stratified scale, combining main categories and/or symbol units in order to produce a more homogenous classification with respect to landscaping, soil formation or parent material. Within this framework, this random check was meant to provide further quantitative information for the existing soil maps.

### 7.6.3 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

One consistent approach was used over all land use categories. See par 7.2.2.

### 7.6.4 Definitions

The Netherlands has chosen to define croplands as arable lands and nurseries (including tree nurseries). Intensive grasslands are not included in this category and are reported under Grasslands. For part of the agricultural land, rotation between cropland and grassland is frequent, but data on where exactly this is occurring are as yet lacking. Currently, the situation on the topographical map is leading, with lands under agricultural crops and classified as arable lands at the time of recording reported under Cropland and lands with grass vegetation at the time of recording classified as Grassland.

### 7.6.5 Methodological issues

The type of land use is determined using digitised and digital topographical maps (scale: 1:10,000), which allows the land-use matrix to be completed according to the recommendations in the Good Practice Guidance on Land Use, Land Use Change and Forestry (IPCC, 2003). The years 1990 and 2004 are based on observations of land use; the values for the period in between are obtained through linear

interpolations, and the values for the years after 2004 are obtained by means of extrapolation. For more information on the methodology, see the description on land use and the land use change matrix in Chapter 7.2.

More detailed descriptions of the methods used and emission factors can be found in the protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

## 7.6.6 Uncertainty and time-series consistency

### Uncertainties

The Tier 1 analysis in Annex 7 shown in Table A7.1 provides estimates of uncertainties according to IPCC source categories. The Netherlands uses a Tier 1 analysis for the uncertainty assessment of the sector LULUCF. The uncertainties in the Dutch analysis of carbon levels depend on the collective factors with which the calculations are implemented (calculation of the organic substances in the soil profile and the conversion to a national level) and data on land use and land use change (topographical data). The uncertainty in the CO<sub>2</sub> emissions from 5B2 “Land converted to Cropland” is calculated at 56%; see Olivier et al. (2009) for details.

### Uncertainty in the implied emission factor of 5B2 Land converted to Cropland

The uncertainty in the implied emission factor of 5B2 “Land converted to Cropland” refers to the change in carbon content of mineral soils. The uncertainty in the change in the carbon content of mineral soils is calculated to be 38% (rounded off to 50% in the Tier 1 calculation spreadsheet, since it is the order of magnitude that is important).

### Uncertainty in activity data

The activity data used is area change, calculated by comparing two topographic maps. The uncertainty of one topographic map is estimated to be 5% (expert judgment). Therefore, the uncertainty for comparing two topographic maps is theoretically  $5 \times 5 = 25\%$ . This is without doubt an overestimation as not all land use may change over a decade.

### Time-series consistency

The yearly emission of CO<sub>2</sub> due to the conversion of land converted to cropland shows a small yearly increase from 35 Gg CO<sub>2</sub> in 1990 to 48 Gg CO<sub>2</sub> in 2008.

### 7.6.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

### 7.6.8 Source-specific recalculations

This year there were no recalculations.

### 7.6.9 Category-specific planned improvements

For this land use category no improvements are planned in the direct future.

## 7.7 Grassland [5C]

### 7.7.1 Source category description

The source category 5C “Grassland” includes only the emissions of CO<sub>2</sub> from 5C1 “Grassland remaining Grassland” and 5C2 “Land converted into Grassland”. The source category 5C1 is by far the most important source of CO<sub>2</sub> within the sector LULUCF.

### 7.7.2 Activity data and (implied) emission factors

The activity data is derived from the land use maps and the land use change matrix. Carbon content is based on the soil map of the Netherlands in combination with results of LSK, a national random check of map units that provides detailed descriptions of soil profiles (see Section 7.3.1). The activity data for organic soils is based on soil maps (1:50,000 for the period 1960–1990), recent inventories on organic soils (2001–2003), profile information from LSK and data on field levels in 1990 and 2000.

### 7.7.3 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

One consistent approach was used over all land use categories. See par 7.2.2.

### 7.7.4 Definition

The land use category “Grassland” is defined as rangeland and pasture land that is not considered as croplands. It also includes vegetation that falls below the threshold used in the forest land category and are not expected to exceed, without human intervention, the threshold used in the forest land category. The category also includes all grassland from wild lands to recreational areas as well as agricultural and silvi-pastoral systems, subdivided into managed and unmanaged consistent with national definitions. (IPCC, 2003). It is stratified in:

- “Grasslands” - all areas predominantly covered by grass vegetation (whether natural, recreational or cultivated)
- “Nature” - all natural areas excluding grassland (natural grasslands and grasslands used for recreation purposes). It mainly consists of heath land, peat moors and other nature areas. Many have the occasional tree as part of the typical vegetation structure. This category was in the previous submissions a subcategory within Forest land

The Netherlands currently reports under grassland any type of terrain which is predominantly covered by grass vegetation. No distinction is made between agricultural intensively and extensively managed grasslands and natural grasslands. However, the potential and the need for this are currently under discussion. Apart from pure grasslands, all orchards (with standard fruit trees, dwarf varieties or shrubs) are included in the category grasslands. They do not conform to the forest definition, and while agro-forestry systems are mentioned in the definition of Croplands, this is motivated by the cultivation of soil under trees. However, in the Netherlands the main undergrowth of orchards is grass. We therefore chose to report them as grasslands. As for grasslands no change in above-ground biomass is reported, the carbon stored in these trees is not reported.

### 7.7.5 Methodological issues

For information on the methodology to assess land use and land use change see Chapter 7.2. A country-specific Tier 2 method is used to estimate CO<sub>2</sub> emissions from the drainage of organic soils (Grassland remaining Grassland).

For grassland, CO<sub>2</sub> emissions resulting from soil subsidence of peat land by oxidation of peat due to managed drainage are added. The CO<sub>2</sub> emission of 5C1 “Grassland remaining Grassland” is calculated and based on observations on yearly subsidence rates for various types of peat and available information on the extent of drainage and subsequent soil carbon losses through oxidation for each peat type and drainage level (Kuikman et al., 2005). The country-specific method used is based on the recommendations given in the IPCC 2003 Good Practice Guidance (IPCC, 2003). Uncertainty in the decrease in the area of organic soils in past decades – in particular, the estimate for 1990 – has led to the conclusion that the area can be considered to be relatively constant yet likely to be still decreasing at a low rate since 1990 (223,000 ha is the observed area of organic soils and thus a conservative estimate). The 2003 stated area of organic soils with the relevant water management conditions and measures and calculated loss of organic matter calculates an implied emission factor of on average 19.04 ton CO<sub>2</sub>/ha (Kuikman, 2005). For the period 1990-2008 the emissions from organic soils under grassland are based on the fixed area and implied emission factor value. Both are the result of analysis of the developments in a range of different peat lands (including water and soil management). The area used so far conflicts to some extent with the results for grassland on organics soils of the land use change matrix. The matrix shows a 4% smaller area and overtime a very slight decrease in area. As long as the loss of carbon cannot be verified and calculated on an annual bases (based on accurate condition data e.g., temperature and water management) the use of year specific area figures of the matrix introduces a pseudo accuracy. Therefore we have decided not to change the calculation methodology as outlined in Kuikman et al, 2005. More detailed descriptions of the methods used and emission factors can be found in protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 7.7.6 Uncertainty and time-series consistency

#### Uncertainties

The Tier 1 analysis in Annex 7 shown in Table A7.1 provides estimates of uncertainties according to the IPCC source category. The uncertainty for the CO<sub>2</sub> emissions in categories 5C1 Grassland remaining Grassland and 5C2 Land converted to Grassland is calculated to be 56%; see Olivier et al. (2009) for details.

#### Uncertainty in the implied emission factor of 5C1 Grassland remaining Grassland

The uncertainty for the oxidation of organic soils in category 5C1 is calculated at 55%. Combined with the 38% uncertainty of the change in carbon content of mineral soils (see Section 7.3.3), the overall uncertainty in the implied emission factor for category 5C1 will probably remain in the 50% range (50% used in the Tier 1 calculation spreadsheet).



### Uncertainty in the implied emission factor of 5C2 Land converted to grassland

For the uncertainty of 5C “Land converted to Grassland”, reference is made to the description of 5B2 “Land converted to Cropland” (Section 7.3.3). The calculation for “Land converted to Grassland” is based on the same assumptions as those made for 5B2 “Land converted to Cropland” and are, therefore, identical. The uncertainty is estimated to be 38% (50% used in the Tier 1 calculation spreadsheet).

### Uncertainty in activity data of categories 5C1 and 5C2

The activity data used is area change, calculated by comparing two topographic maps. The uncertainty of one topographic map is estimated to be 5% (expert judgment). Therefore, the uncertainty in comparing two topographic maps is theoretically  $5 \times 5 = 25\%$ . This is without doubt an overestimation as not all land use may change over a decade.

### Time-series consistency

The yearly source of CO<sub>2</sub> that results from the drainage of organic soils is 4.246 Gg CO<sub>2</sub>. The yearly emission of CO<sub>2</sub> due to the conversion of forest land to grassland shows a steady increase from 394 Gg CO<sub>2</sub> in 1990 to 550 Gg CO<sub>2</sub> in 2008.

#### 7.7.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

#### 7.7.8 Source-specific recalculations

This year there were no recalculations.

#### 7.7.9 Category-specific planned improvements

For this land use category no improvements are planned in the direct future

## 7.8 Wetland [5D]

### 7.8.1 Source category description

The source category 5D “Wetland” includes only CO<sub>2</sub> emissions from 5D1 “Wetland remaining Wetland” and 5D2 “Land converted to Wetland”.

### 7.8.2 Activity data and (implied) emission factors

The activity data is derived from the land use maps and the land use change matrix (see Section 7.3.2.). The carbon content of wetlands is not estimated and is set at zero in the land use change matrix.

### 7.8.3 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

One consistent approach was used over all land use categories. See par 7.2.2.

### 7.8.4 Definition

The land use category “Wetland” includes land that is covered or saturated with water for all or part of the year and does not fall into the forest land, cropland, grassland or settlements categories. It includes reservoirs as a managed sub-division and natural lakes and rivers as unmanaged sub-divisions (IPCC, 2003). Though the Netherlands is a country

with many wet areas by nature, many of these are covered by a grassy vegetation and those are included under grasslands. Some wetlands are covered by a more rough vegetation of wild grasses or shrubby vegetation, which is reported in the subcategory “Nature” of Grassland. Forested wetlands like willow coppice are reported in the subcategories FAD or TOF of Forest Land, depending on their surface.

In the Netherlands, only reed marshes and open water bodies are included in the Wetland land use category. This includes natural open water in rivers, but also man-made open water in channels, ditches and artificial lakes. It includes bare areas which are under water only part of the time as a result of tidal influences, and very wet areas without vegetation. It also includes “wet” infrastructure for boats, i.e. waterways but also the water in harbours and docks.

### 7.8.5 Methodological issues

For information on the methodology to assess land use and land use change see Chapter 7.2. The emission of CH<sub>4</sub> from wetlands is not estimated due to the lack of data. More detailed descriptions of the methods used and the emission factors can be found in protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

### 7.8.6 Uncertainty and time-series consistency

#### Uncertainties

For information on the uncertainty estimates, the reader is referred to Section 7.2.5, which discusses the uncertainty of soil carbon and changes in land use.

#### Time-series consistency

The time-series shows a consistent small increase from 40 Gg CO<sub>2</sub> in 1990 to 56 Gg CO<sub>2</sub> in 2008.

#### 7.8.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

#### 7.8.8 Source-specific recalculations

This year there are no recalculations.

#### 7.8.9 Category-specific planned improvements

For this land use category no improvements are planned in the direct future.

## 7.9 Settlement [5E]

### 7.9.1 Source category description

This source category 5E “Settlement” includes only those CO<sub>2</sub> emissions from 5E1 “Settlements remaining Settlements” and 5E2 “Land converted to Settlements”.

### 7.9.2 Activity data and (implied) emission factors

The activity data are derived from the land use maps and the land use change matrix. Estimates of carbon content are based on the soil map of the Netherlands in combination with results of LSK, a national random check of map units that provides detailed descriptions of soil profiles. There is a lack of information on the carbon content for most of the

settlement grid cells. Consequently, the carbon content was calculated using a weighed average over all carbon stock classes within each land use category.

### 7.9.3 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

One consistent approach was used over all land use categories. See par 7.2.2.

### 7.9.4 Definition

The land use category “Settlements” includes all developed land, including transportation infrastructure and human settlements of any size, unless they are already included under other categories (IPCC, 2003). In the Netherlands, the main classes included are urban areas and transportation infrastructure, and built-up areas. Built-up areas include any constructed item, independent of the type of construction material, which is (expected to be) permanent, fixed to the soil surface and serves as place for residence, trade, traffic and/or labour. Thus it includes houses, blocks of houses and apartments, office buildings, shops and warehouses but also fuel stations and greenhouses. Urban areas and transportation infrastructure include all roads, whether paved or not, are included in the land use category Settlements with exception of forest roads which are included in the official forest definition. It also includes train tracks, (paved) open spaces in urban areas, parking lots and graveyards. Though some of the last classes are actually covered by grass, the distinction cannot be made based on maps. As even the grass graveyards are not managed as grasslands, inclusion in the land use category “Settlements” conforms better to the rationale of the land use classification

### 7.9.5 Methodological issues

For information on the methodology to assess land use and land use change see Chapter 7.2. The reporting is considered to be a Tier 2 level (see protocol 0073). Because there has been no change in soil carbon and, in any case, no loss of soil carbon was expected for the period 1990–2004, the emissions from 5E1 “Settlement land remaining Settlement” are set at zero. The category 5E2 “Land converted to Settlement” includes the conversion from mainly grassland, cropland and other land to settlements. More detailed descriptions of the methods used and the emission factors can be found in the protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 7.1.

### 7.9.6 Uncertainty and time-series consistency

#### Uncertainties

Uncertainty estimates are provided in Section 7.2.5, which discusses the uncertainty of soil carbon and changes in land use.

#### Time-series consistency

The time-series shows a consistent increase from 212 Gg CO<sub>2</sub> in 1990 to 296 Gg CO<sub>2</sub> in 2008.

### 7.9.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

### 7.9.8 Source-specific recalculations

This year there are no recalculations.

### 7.9.9 Category-specific planned improvements

For this land use category no improvements are planned in the direct future

## 7.10 Other Land [5F]

### 7.10.1 Source category description

This source category 5F “Other Land” includes only CO<sub>2</sub> emissions from 5F1 “Other Land remaining Other Land” and 5F2 “Land converted to Other Land”.

### 7.10.2 Activity data and (implied) emission factors

The activity data are derived from the land use maps and the land use change matrix. Carbon content is based on the soil map of the Netherlands in combination with results of LSK, a national random check of map units that provides detailed descriptions of soil profiles.

### 7.10.3 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

One consistent approach was used over all land use categories. See par 7.2.2.

### 7.10.4 Definition

The land use category “Other Land” was included to allow the total of identified land to match the national area where data are available. It includes bare soil, rock, ice and all unmanaged land area that do not fall in any of the other five categories. (IPCC, 2003).

In general, Other Land does not have a substantial amount of carbon. The Netherlands uses this land use category to report the surfaces of bare soil which are not included in any other category. In the Netherlands this refers mostly to almost bare sands and the earliest stages of succession from sand in the coastal areas (beaches, dunes and sandy roads) or uncultivated land alongside rivers. It does not include bare areas that emerge from shrinking and expanding water surfaces (these “emerging surfaces” are included in wetlands).

### 7.10.5 Methodological issues

For information on the methodology to assess land use and land use change see Chapter 7.2. The land use category Other Land is introduced to allow wall-to-wall reporting of land areas even if not all land could be allocated to a land use category. The carbon stored in land allocated to Other Land need not be reported (as it is assumed that Other Land has no carbon). In previous submissions, a quite broad definition of Other Land was used, and the carbon in land converted to or from Other Land was assumed to change between reported and not reported. Therefore, large positive and negative emissions were reported which did not actually reflect changes in carbon in soil, but the reporting status of carbon in soil. This was deemed not realistic. In the 2009 submission a more narrow definition of Other Land is used, and the reporting of “administrative emissions” was stopped. More

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008
5G Other (liming of agricultural soils)	183	98	98	80	85	86	79	75	81	71	71

detailed descriptions of the methods used and the emission factors can be found in protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 7.1.

#### 7.10.6 Uncertainty and time-series consistency

##### Uncertainties

For information on the uncertainty estimation, the reader is referred to Section 7.2.5, which discusses the uncertainty of soil carbon and changes in land use.

##### Time-series consistency

The time-series shows a consistent small increase from 18 Gg CO<sub>2</sub> in 1990 to 25 Gg CO<sub>2</sub> in 2008.

#### 7.10.7 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

#### 7.10.8 Source-specific recalculations

This year there are no recalculations.

#### 7.10.9 Category-specific planned improvements

For this land use category no improvements are planned in the direct future.

### 7.11 Other [5G]

#### 7.11.1 Source category description

The source category 5G “Other” includes only the emissions of CO<sub>2</sub> from the liming of agricultural land with limestone and dolomite. Limestone and dolomite are used in the agricultural sector to increase the chalk content of the soil.

##### Activity data and (implied) emission factors

The activity data is derived from agricultural statistics for total lime fertilisers (period 1990–2008). Data available on the application of limestone and dolomite does not address its use on grassland and cropland separately.

#### 7.11.2 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

Information on liming was derived from national, yearly updated, statistics on fertiliser use. The yearly amount of limestone and dolomite are converted into carbon dioxide emissions conform the calculations in the guidelines.

##### Definition

The category “other land” was included to allow the total of identified land to match the national area for which data are available. It includes bare soils, rock and all unmanaged land area that do not fall in any other five categories.

#### 7.11.3 Methodological issues

The reporting is considered to be at the Tier 2 level (see protocol 0073). Limestone (“lime marl”) and dolomite (“carbonic magnesium lime”) amounts, reported in CaO equivalents, are multiplied with the emission factors for limestone (440 kg CO<sub>2</sub>/ton pure limestone) and for dolomite (0.477 tons CO<sub>2</sub> per ton pure dolomite). More detailed descriptions of the methods used and the emission factors can be found in protocols 0072 and 0073 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 7.1.

#### 7.11.4 Uncertainty and time-series consistency

##### Uncertainties

The Tier 1 analysis in Annex 7 shown in Table A7.1 provides estimates of uncertainties according to IPCC source category. The uncertainty in the CO<sub>2</sub> emissions from 5G “Liming of soils” is calculated to be 25%. The uncertainty in the activity data is estimated to be 25%, and the uncertainty in emission factors is 1%. When considered over a longer time span, all carbon that is applied through liming is emitted.

##### Time-series consistency

The methodology used to calculate CO<sub>2</sub> emissions from limestone and dolomite for the period 1990–2008 is consistent over time. The use of chalk containing fertiliser in the Netherlands decreased from 265 million kg in 1990 to 101 million kg in 2008. Over that period the proportion of limestone more than doubled, from about 12% in 1990 up to almost 30% in 2008 and the proportion of dolomite decreased from about 35 to 40% in 1990 to levels below 30% in 2008 (the remaining 38% is earth foam). Although the figures for 2008 show a relative increase in the use of limestone and dolomite, in absolute figures the supply was 10% below the supply in 2007. The CO<sub>2</sub> emissions related to these fertilisers is 70 Gg CO<sub>2</sub>, which is 10 Gg CO<sub>2</sub> less than in 2007 (see Table 7.3).

#### 7.11.5 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures, as discussed in Chapter 1.

#### 7.11.6 Source-specific recalculations

Not applicable.

#### 7.11.7 Category-specific planned improvements

A recalculation over 2008 will be carried out when data become available.

# Waste [CRF Sector 6]

## ■ Major changes in Waste sector compared to the National Inventory Report 2009

*Emissions:* In 2008, the total greenhouse gas emissions in this sector decreased further. Emissions in the period 1990-2007 did not change compared to the previous NIR.

*Key sources:* No changes in key sources compared to previous submission.

*Methodologies:* There were no methodological changes compared to last submission.

### 8.1 Overview of sector

The national inventory of the Netherlands comprises four source categories in the Waste sector:

- 6A Solid waste disposal: CH<sub>4</sub> (methane) emissions;
- 6B Wastewater handling: CH<sub>4</sub> and N<sub>2</sub>O emissions;
- 6C Waste incineration: CO<sub>2</sub> emissions (included in [1A1a]);
- 6D Other waste: CH<sub>4</sub> emissions.

Carbon dioxide emissions from the anaerobic decay of waste in land fill sites are not included, since this is considered to be part of the carbon cycle and is not a net source. The Netherlands does not report emissions from waste incineration facilities in the Waste sector because these facilities also produce electricity or heat used for energetic purposes and, as such, these emissions are included in category 1A1a (to comply with IPCC reporting guidelines). Methodological issues concerning this source category are briefly discussed in Section 8.4.

The following protocols, which can be found on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), describe the methodologies applied for estimating CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions of the Waste sector in the Netherlands (see also Annex 6):

- Protocol 0074: CH<sub>4</sub> from Waste disposal (6A1);
- Protocol 0075: CH<sub>4</sub>, N<sub>2</sub>O from Wastewater treatment (6B);
- Protocol 0076: CH<sub>4</sub>, N<sub>2</sub>O from Industrial composting (6D);
- Protocol 0078: CO<sub>2</sub> CH<sub>4</sub> N<sub>2</sub>O from Biomass (1A);

The Waste sector accounted for 3% of total national emissions (without LULUCF) in 2008 compared with 6% in 1990, with the emissions of CH<sub>4</sub> and N<sub>2</sub>O accounting for 91% and 9% of CO<sub>2</sub> equivalent emissions from the sector, respectively. Emissions of CH<sub>4</sub> from waste – almost all (87%) from Landfills (6A) – accounted for 29% of the national total CH<sub>4</sub> emissions in 2008. The N<sub>2</sub>O emissions from the Waste sector stem from domestic and commercial wastewater. The fossil-fuel related emissions from waste incineration, mainly CO<sub>2</sub>, are included in the fuel combustion emissions from the Energy Sector (1A1) since all large-scale incinerators also produce electricity and/or heat for energetic purposes.


Emissions from the Waste sector decreased by 53% between 1990 and 2008 (see Figure 8.1), mainly due to a 56% reduction in CH<sub>4</sub> from Landfills (6A1 'Managed waste disposal on land'). Between 2007 and 2008 the CH<sub>4</sub> emissions from landfills decreased by about 7%. The decreased methane emission from "Landfills" since 1990 is the result of:

- increasing recovery and recycling of waste for composting and/or incineration
- a considerable reduction in the amount of municipal solid waste (MSW) disposal at landfills
- a decreasing organic waste fraction in the waste disposed
- increasing methane recovery from the landfills (from 5% in 1990 to 23% in 2008) (see SenterNovem, 2009)

Table 8.1 shows the contribution of the emissions from the Waste sector to the total greenhouse gas emissions in the Netherlands and also presents the key sources in this sector specified by level, trend or both. The list of all (key- and non-key) sources in the Netherlands is shown in Annex 1. Total greenhouse gas emissions from the Waste sector decreased from 12.8 Tg CO<sub>2</sub> eq in 1990 to 5.7 Tg CO<sub>2</sub> eq in 2008. This decrease is mainly due to (SenterNovem, 2008):

- Increased recovery and recycling, resulting in a decreasing amount of solid waste disposed at landfills;
- A decreasing amount of organic waste disposed of at landfills;
- Increasing CH<sub>4</sub> recovery from landfills

CH<sub>4</sub> emissions from landfills contribute the largest share to the greenhouse gas emissions of this sector. Category 6A1 Solid waste disposal sites (SWDS) is a key source specified by both level and trend, category 6B N<sub>2</sub>O emissions from waste water handling is a minor key source (L2) when uncertainties are taken into account (see Annex 1).



Contribution of main categories and key sources in Sector 6 Waste

Table 8.1

Sector/category	Gas	Key	Emissions base-year	Emissions 2007	Emissions 2008	Change 2008–2007	Contribution to total in 2008 (%)		
			Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Tg CO <sub>2</sub> eq	Gg	By sector	Of total gas	Of total CO <sub>2</sub> eq
6 Waste	CH <sub>4</sub>		12.3	5.5	5.2	-0.4	91	30	2
	N <sub>2</sub> O		0.5	0.5	0.5	0.0	9	4	0.2
	All		12.8	6.0	5.7	-0.4	100		3
6A. Solid Waste Disposal on Land	CH <sub>4</sub>		12.0	5.3	4.9	-0.4	87	29	2
6A1. Managed Waste Disposal on Land	CH <sub>4</sub>	L.T	12.0	5.3	4.9	-0.4	87	29	2
6B Waste water handling	N <sub>2</sub> O	L2	0.5	0.5	0.5	0.0	8	4	0.2
	CH <sub>4</sub>		0.3	0.2	0.2	0.0	3	1	0.1
	All		0.8	0.7	0.7	0.0	12		0.3
6D. Other	CH <sub>4</sub>		0.0	0.1	0.1	0.0	1	0.4	0.0
National Total GHG emissions (excl. CO <sub>2</sub> LULUCF)	CH <sub>4</sub>		25.5	16.8	17.1	0.2		100	
	N <sub>2</sub> O		20.2	15.4	11.8	-3.7		100	
	All		213.3	206.9	206.9	0.0			100



## 8.2 Solid waste disposal on land [6A]

### 8.2.1 Source category description

In 2008 there were 22 operating landfill sites as well as a few thousand older sites that are still reactive. CH<sub>4</sub> recovery takes place at 51 sites in the Netherlands. As a result of anaerobic degradation of the organic material within the landfill body, all of these landfills produce CH<sub>4</sub> and CO<sub>2</sub>. Landfill gas comprises about 60% (vol.) CH<sub>4</sub> and 40% (vol.) CO<sub>2</sub>. Due to a light overpressure, the landfill gas migrates into the atmosphere. On several landfill sites the gas is extracted before it is released into the atmosphere and subsequently used as an energy source or flared off. In both of these cases the CH<sub>4</sub> in the extracted gas is not released into the atmosphere. The CH<sub>4</sub> may be degraded (oxidised) to some extent by bacteria when it passes through the landfill cover; this results in a lower CH<sub>4</sub> emission.

Anaerobic degradation of organic matter in landfills is a time-dependent process and may take many decades. Some of the factors influencing this process are known; some are not. Each landfill site has its own unique characteristics: concentration and type of organic matter, moisture and temperature, among others. The major factors determining the decreased net CH<sub>4</sub> emissions are lower quantities of organic carbon deposited into landfills (organic carbon content × total amount of land-filled waste) and higher methane recovery rates from landfills (see Sections 8.2.2 and 8.2.3).

The share of CH<sub>4</sub> emissions from landfills in the total national inventory of greenhouse gas emissions was 6% in 1990 and 3% in 2008. Between 1990 and 2008 CH<sub>4</sub> emissions have decreased by 55%. This decrease is due to the increase in recovered CH<sub>4</sub> – from about 5% in 1990 to 23% in 2008 – but also to the decrease in methane produced in solid waste disposal sites.

In 2008 solid waste disposal on land accounted for 87% of the total emissions in the Waste sector and 2% of the total national CO<sub>2</sub> equivalent emissions (see Table 8.1).

The policy that has been implemented in the Netherlands is one directly aimed at reducing the amount of waste used in land fill sites. This policy requires enhanced prevention of waste production and the increased recycling of waste, followed by incineration. As early as the 1990s the government introduced bans on the use of certain categories of waste for land-filling; for example, the organic fraction of household waste. Another method implemented to reduce land-filling was to raise the landfill tax to comply with the increased costs of incinerating waste. Depending on the capacity of incineration, the government can grant exemption from these ‘obligations’. Due to this policy the amount of waste used as landfill has decreased, thereby reducing emissions from this source category from more than 14 million tonnes in 1990 to 2 million tonnes in 2008.

### 8.2.2 Methodological issues

A more detailed description of the method used and emission factors can be found in the protocol 0074 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 8.1.

Activity data on the amount of waste disposed of at landfill sites are mainly based on the annual survey performed by the Working Group on Waste Registration at all the landfill sites in the Netherlands. This data can be found on the website [www.uitvoeringafvalbeheer.nl](http://www.uitvoeringafvalbeheer.nl) and are documented in SenterNovem (2007). This document also contains the amount of CH<sub>4</sub> recovered from landfill sites yearly.

The implied emission factors correspond with the IPCC default values.

In order to calculate the CH<sub>4</sub> emissions from all the landfill sites in the Netherlands, the simplifying assumption was made that all the wastes are assumed to be landfill on one landfill site, an action that started in 1945. However, as stated above, characteristics of individual sites vary substantially. CH<sub>4</sub> emissions from this ‘national landfill’ are then calculated using a first-order decomposition model (first-order decay function) with an annual input of the total amounts deposited and the characteristics of the land-filled waste and the amount of landfill gas extracted. This is equivalent to the IPCC Tier 2 methodology. Since the CH<sub>4</sub> emissions from landfills are a key source, the present methodology is in line with the IPCC Good Practice Guidance (IPCC, 2001).

Parameters used in the landfill emissions model are as follows:

- total amount of land-filled waste
- fraction of degradable organic carbon (DOC) (see Table 8.2 for a detailed time-series)
- CH<sub>4</sub> generation (decomposition) rate constant (k): 0.094 up to and including 1989, decreasing to 0.0693 in 1995 and constant thereafter; this corresponds to half-life times of 7.4 and 10.0 years respectively (see Table 8.2 for a detailed time-series);
- CH<sub>4</sub> oxidation factor: 10%;
- fraction of DOC actually dissimilated (DOCF): 0.58; (see also (Oonk, 1994))
- CH<sub>4</sub> conversion factor (IPCC parameter): 1.0.
- Fraction of methane in landfill gas is determined yearly from 2002 onwards, based on the composition of landfill gas at all sites with CH<sub>4</sub> recovery. For the years until 2001, the fraction of methane in landfill gas is set at 60%

Trend information on IPCC Tier 2 method parameters that change over time is provided in Table 8.2. The change in DOC values is due to such factors as the prohibition of land filling of combustible wastes, whereas the change in k-values (CH<sub>4</sub> generation rate constant) is caused by a sharp increase in the recycling of vegetable, fruit and garden waste in the early 1990's. The integration time for the emission calculation is defined as the period from 1945 to the year for which the calculation is made.

### 8.2.3 Uncertainty and time-series consistency

#### Uncertainty

The Tier 1 uncertainty analysis shown in Tables A7.1 and A7.2 of Annex 7 provides estimates of uncertainties according to IPCC source category and gas. The uncertainty in CH<sub>4</sub> emissions of solid waste disposal sites is estimated to be approximately 35% in annual emissions. The uncertainty in the

Parameter	1990	1995	2000	2005	2006	2007	2008
Waste generation rate <sup>1)</sup> (kg/cap/day)	1.52	1.50	1.69	1.70	1.70	1.70	1.74
Fraction MSW disposed to SWDS	0.38	0.29	0.09	0.01	0.01	0.01	0.01
Fraction DOC in MSW	0.13	0.13	0.11	0.08	0.07	0.07	0.07
Fraction of waste incinerated	0.08	0.09	0.12	0.12	0.12	0.12	0.12
Fraction of waste recycled	0.63	0.75	0.80	0.84	0.84	0.84	0.84
CH <sub>4</sub> generation rate constant (k)	0.09	0.07	0.07	0.07	0.07	0.07	0.07
Number of SWDS recovering CH <sub>4</sub>	45	50	55	50	50	51	52
Waste incineration <sup>2)</sup> (Tg)	3.9	4.7	7.1	7.1	7.1	7.2	7.2
Fraction CH <sub>4</sub> in landfill gas	0.6	0.6	0.6	0.522	0.55	0.521	0.51

1) Waste generation rate refers to MSW (municipal solid waste), excluding inorganic industrial waste such as construction or demolition waste.

2) Waste incineration refers to the total amount of waste incinerated: municipal solid waste, industrial waste, commercial waste, sewage sludge.

activity data and the emission factor are estimated to be 30% and 15%, respectively. For a more detailed analysis of these uncertainties, see Olivier et al. (2009).

#### Time-series consistency

The estimates for all years are calculated from the same model, which means that the methodology is consistent throughout the time-series. The time-series consistency of the activity data is very good due to the continuity in the data provided. Since 2002 the fraction of CH<sub>4</sub> in landfill gas is determined yearly based on the composition of the landfill gas of the sites recovering CH<sub>4</sub>. It is expected that this will reflect the average fraction of CH<sub>4</sub> in the landfill gas better than the default used in previous inventories and slightly reduces uncertainties in the emission estimations of the post-2001 period. This "new" CH<sub>4</sub> fraction is only used to estimate methane in the recovered biogas and not for the generation of methane within the landfill site.

#### 8.2.4 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures as discussed in Chapter 1.

#### 8.2.5 Source-specific recalculations

There are no source specific recalculations compared to the previous submission. For the year 2007 there was an error-correction in the emission. All landfill emissions were corrected due to a correction in the amount of waste disposed on land in 2007.

#### 8.2.6 Source specific planned improvements

During the review of the NIR 2006 by the ERT (in 2008) it was recommended to investigate the composition of soils in order to verify the fraction of organic carbon present and to include this fraction in the estimation of CH<sub>4</sub> emissions. In 2008 a project started studying, among others things, contaminated soils at landfill sites. In the first half of 2010 the results will be available and will be incorporated in the estimation method.

### 8.3 Wastewater handling [6B]

#### 8.3.1 Source category description

This source category covers emissions released from Wastewater handling and includes emissions from industrial, commercial and domestic wastewater and septic tanks.

The treatment of urban wastewaters and the resulting wastewater sludge is accomplished using aerobic and/or anaerobic processes. During the treatment, the biological breakdown of Degradable Organic Compounds (DOC) and nitrogen compounds can result in CH<sub>4</sub> and N<sub>2</sub>O emissions, respectively. The discharge of effluents subsequently results in indirect N<sub>2</sub>O emissions from surface waters due to the natural breakdown of residual nitrogen compounds. The source category also includes the CH<sub>4</sub> emissions from anaerobic industrial wastewater treatment plants (WWTP) and from septic tanks, but these are small compared to urban WWTP.

N<sub>2</sub>O emissions from waste water treatment (see Table 8.1) contributed about 4% to total N<sub>2</sub>O emissions in 2008 and 0.2% in total CO<sub>2</sub> eq. N<sub>2</sub>O emissions from waste water handling decreased by 2% during the period 1990–2008. This small decrease is the result of two counteracting trends. Improved biological breakdown of nitrogen compounds at urban WWTPs (see Table 8.4) leads to a gradual increase of N<sub>2</sub>O emissions. However, this improved nitrogen removal results in lower effluent loads (see Table 8.4) and a subsequent decrease in the (indirect) N<sub>2</sub>O emissions from human sewage.

The contribution of wastewater handling to the national total of CH<sub>4</sub> emissions in 2008 was 1%. Since 1994, CH<sub>4</sub> emissions from wastewater treatment plants have decreased due to the introduction in 1990 of a new sludge stabilisation system in one of the largest wastewater treatment plants. As the operation of the plant took a few years to optimise, venting emissions were higher in the introductory period (1991–1994) than under normal operating conditions.

The amount of wastewater and sludge being treated does not change much over time. Therefore, the inter-annual changes in methane emissions can be explained by varying fractions



Wastewater handling emissions of CH<sub>4</sub> and N<sub>2</sub>O (Units: Gg/year)

Table 8.3

	1990	1995	2000	2005	2006	2007	2008
CH <sub>4</sub> industrial wastewater	0.25	0.33	0.34	0.36	0.33	0.32	0.33
CH <sub>4</sub> domestic & commercial wastewater	9.07	7.90	7.96	8.20	8.12	8.37	8.23
CH <sub>4</sub> septic tanks	4.47	3.25	2.20	1.22	1.11	0.98	0.86
Net CH <sub>4</sub> emissions	13.79	11.48	10.50	9.78	9.56	9.68	9.42
CH <sub>4</sub> recovered and/or flared	33.0	39.2	40.4	41.9	43.8	42.6	43.7
Recovery/flared (% gross emission)	70.5	77.4	79.4	81.1	82.1	81.5	82.3
N <sub>2</sub> O domestic & commercial wastewater	0.66	0.75	0.88	0.99	1.05	1.10	1.14
N <sub>2</sub> O from human sewage	0.37	0.37	0.53	0.44	0.38	0.37	0.34
Total N <sub>2</sub> O emissions	1.50	1.40	1.41	1.43	1.43	1.47	1.48

Activity data of domestic and commercial wastewater handling (Gg/year), total volume of treated urban waste water (Mm<sup>3</sup>/year) and percentage of population connected to septic tanks (%)

Table 8.4

	1990	1995	2000	2005	2006	2007	2008
Wastewater DOC <sup>1)</sup>	933	921	921	943	938	942	946
Sludge DOC	254	269	281	298	318	294	316
Nitrogen removed in urban WWTP	42.0	47.7	55.8	63.1	66.8	70.0	72.5
Nitrogen in effluents <sup>2)</sup>	53.8	41.5	33.8	27.8	24.3	23.6	23.6
Treated volume	1,711	1,908	2,034	1,841	1,854	2,069	1,929
% Inhabitants with septic tanks	4.0	2.8	1.9	1.1	0.9	0.8	0.7

1) DOC, Degradable organic component.

2) Total of industrial, domestic and commercial effluents.

of methane being flared instead of vented or used for energy purposes.

Table 8.3 shows the trend in greenhouse gas emissions from the different sources of wastewater handling.

### 8.3.2 Methodological issues

#### Activity data and (implied) emission factors

Detailed information on activity data and emission factors can be found in the monitoring protocol 0075 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

Most of the activity data on wastewater treatment are collected by Statistics Netherlands in yearly questionnaires which cover all urban WWTPs as well as all anaerobic industrial WWTPs; see also [www.statline.cbs.nl](http://www.statline.cbs.nl) for detailed statistics on wastewater treatment. Table 8.4 shows the development in the key activity data with respect to urban (= domestic and commercial) wastewater treatment. Due to varying weather conditions the volumes of treated wastewater and of the total load of DOC can fluctuate from year to year, depending on how much run-off rainwater enters the sewer systems.

Table 8.3 shows that total N<sub>2</sub>O emissions from wastewater handling decreased only 2% between 1990 and 2008. This small decrease is the overall result of an increase in N<sub>2</sub>O emissions caused by improved nitrogen removal at urban WWTPs, and a decrease in the (indirect) N<sub>2</sub>O emissions from human sewage as a result of lower effluent loads (see Table 8.4).

From Table 8.4 it can be concluded that the DOC of treated wastewater and sludge does not significantly change over time. Therefore, the inter-annual changes in CH<sub>4</sub> emissions can be explained by varying fractions of CH<sub>4</sub> being vented instead of flared or used for energy purposes.

The source Septic Tanks has steadily decreased from 1990 onwards. This can be explained by the increased number of households connected to the sewer system in the Netherlands (and therefore no longer using septic tanks; see Table 8.4).

A full description of the methodology is provided in the monitoring protocol 0075 (see the website [www.greenhousegases.nl](http://www.greenhousegases.nl)) and in the background document (Oonk et al., 2004). In general, the emissions are calculated according to the IPCC guidelines, with country-specific parameters and emission factors being used for CH<sub>4</sub> emissions from wastewater handling (including sludge). The calculation methods are equivalent to the IPCC Tier 2 methods.

#### CH<sub>4</sub> emissions

For anaerobic industrial WWTPs, the CH<sub>4</sub> emission factor is expressed as 0.056 t/t DOC design capacity, assuming a utilisation rate of 80%, a CH<sub>4</sub>-producing potential (B<sub>0</sub>) of 0.22 t/t DOC and a methane recovery (MR) of 99%.

For Urban wastewater treatment and anaerobic sludge handling, the combined emission factor is defined as 0.0085 tons CH<sub>4</sub> per ton DOC<sub>influent</sub>. The following parameters are underlying the calculation of this emission factor (for further details, see also background document Oonk et al., 2004):

- Methane formation  $B_0 = 0.25 \text{ t CH}_4/\text{t DOC converted}$  (IPCC, 1996).
- $MCF_{stp}$  = Methane Correction Factor of Sewage Treatment Plants = 3.5% (Doorn et al, 1997, as referred to in IPCC-GPG, 2001).
- 37% of the  $DOC_{inflow}$  remains in the sludge (Country Specific Long Term Annual Average).
- MCF of anaerobic sludge treatment = 54% (Country Specific LTAA).
- In anaerobic sludge treatment 42% of the incoming DOC is digested (Country Specific LTAA).
- $CH_4$  recovery (MR) from anaerobic sludge treatment = 94% (Hobson & Palfrey, 1996, as referred to in IPCC-GPG, 2001).

Incidental venting of biogas at urban WWTPs is recorded by the plant operators and subsequently reported to Statistics Netherlands.

For septic tanks, the emission factor for  $CH_4$  is expressed as 0.0075 tons per year per person connected to a septic tank, assuming a methane correction factor (MCF) of 0.5 and a  $CH_4$ -producing potential ( $B_0$ ) of 0.25. In 2008, only 0.7% of the population was connected to a septic tank.

#### **N<sub>2</sub>O emissions**

The  $N_2O$  emissions from wastewater handling are not determined via the average per-capita protein intake – as many countries do – but on basis of data on the total Nitrogen loads removed in Urban Waste Water treatment plants (see also Table 8.4). Influent and effluent loads of Urban WWTP are monitored systematically in accordance with the rules of the EU Urban Waste Water Treatment Directive, by all of the Dutch Regional Water Authorities. Waste water treated at Urban WWTP is a mix of household waste water, run-off rainwater and waste water from industries and services, so the forthcoming  $N_2O$  emissions are reported under the category 6B2 ‘Domestic and commercial waste water’. Because of their insignificance compared to  $N_2O$  from urban wastewater treatment, no  $N_2O$  emissions were estimated for separate industrial wastewater treatment and from septic tanks.

$N_2O$  emissions from the biological N removal processes in Urban WWTP as well as indirect  $N_2O$  emission from effluents are calculated using the IPCC default emission factor of 0.01 tons  $N_2O$ -N per ton N removed or discharged. Since  $N_2O$  emissions from wastewater handling is identified in earlier NIRs as a key source, the present Tier 2 methodology complies with the IPCC Good Practice Guidance (IPCC, 2001).

#### **8.3.3 Uncertainties and time-series consistency**

##### **Uncertainties**

The Tier 1 uncertainty analysis in Tables A7.1 and A7.2 in Annex 7 provides estimates of uncertainties according to IPCC source category and gas. The uncertainty in annual  $CH_4$  and  $N_2O$  emissions from wastewater handling are estimated to be 30% and 50%, respectively. The uncertainty in activity data is based on the judgments of experts and estimated to be 20%. The uncertainty in emission factors for  $CH_4$  and  $N_2O$  are estimated to be 25% and 50% respectively.

#### **Time-series consistency**

The same methodology has been used to estimate emissions for all years, thereby providing a good time-series consistency. The time-series consistency of activity data is very good due to the continuity in the data provided by Statistics Netherlands.

#### **8.3.4 Source-specific QA/QC and verification**

The source categories are covered by the general QA/QC procedures as discussed in Chapter 1.

#### **8.3.5 Source-specific recalculations**

Compared to the previous submission no recalculations took place for this submission.

#### **8.3.6 Source-specific planned improvements**

There are no source specific planned improvements compared to the previous submission.

### **8.4 Waste incineration [6C]**

#### **8.4.1 Source category description**

The source category ‘Waste incineration’ is included in category 1A1 (Energy industries) as part of the source 1A1a Public electricity and heat production, since all waste incineration facilities in the Netherlands also produce electricity and/or heat used for energetic purposes. According to the IPCC Guidelines (IPCC, 2001), these are included in category 1A1a: Public electricity and heat production: other fuels (see Section 3.2.1).

#### **8.4.2 Methodological issues**

##### **Activity data and emission factors**

The activity data for the amount of waste incinerated are mainly based on the annual survey performed by the Working Group on Waste Registration at all 11 waste incinerators in the Netherlands. Data can be found on the website [www.uitvoeringafvalbeheer.nl](http://www.uitvoeringafvalbeheer.nl) and in a background document (SenterNovem, 2009).

A more detailed description of the method used and the emission factors can be found in the protocol 0078 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), as indicated in Section 8.1.

Total  $CO_2$  emissions – i.e. the sum of organic and fossil carbon – from waste incineration are reported per facility in the annual environmental reports and included in the ER-I data set. The fossil-based and organic  $CO_2$ ,  $CH_4$  en  $N_2O$  emissions from Waste incineration are calculated from the total amount of waste incinerated. The composition of the waste (the six types listed in Table 8.5) is determined per waste stream (residential and several others). An assumption is made for each of these six types of waste with respect to the specific carbon and fossil carbon fractions, which will subsequently yield the  $CO_2$  emissions. Table 8.6 shows the total amounts of waste incinerated, the fractions of the different waste components used for calculating the amounts of fossil and organic carbon in the waste (from their fossil and organic carbon fraction) and the corresponding amounts of fossil and organic carbon in total waste incinerated. The method is

Waste type	Non household waste		Household Waste	
	Carbon fraction	Fossil fraction	Carbon fraction	Fossil Fraction
WIP <sup>1)</sup> : paper/cardboard (%)	30	0	30	23
WIP: wood (%)	45	0	37	6
WIP: other organic (%)	20	0	22	6
WIP: plastics (%)	54	100	45	86
WIP: other combustible (%)	32	41	32	50
WIP: non-combustible (%)	1	100	1	100

1) WIP, Waste incineration plant; listed are the residential waste fractions; for waste fractions of other waste types (considered fixed in time), see Joosen and De Jager, 2003.

Composition of incinerated waste

Table 8.6

	1990	1995	2000	2005	2006	2007	2008
Total waste incinerated (Gg)	2,780	2,913	4,896	5,503	5,542	5,790	6,053
of which household waste (Gg)	2,310	2,083	3,115	4,413	4,086	3,738	3,681
of which							
paper/cardboard (weight %)	26%	33%	32%	25%	25%	25%	26%
wood (weight %)	1%	2%	2%	3%	3%	3%	3%
other organic (weight %)	51%	37%	35%	35%	35%	34%	33%
plastics (weight %)	8%	11%	13%	19%	20%	20%	19%
other combustible (weight %)	3%	5%	5%	6%	6%	6%	6%
non-combustible (weight %)	11%	13%	13%	13%	11%	12%	12%
Total waste incinerated (TJ)	22,746	27,903	51,904	55,058	56,087	59,678	62,341
Energy content (MJ/kg)	8.2	9.6	10.6	10.0	10.1	10.3	10.3
Fraction organic (energy %)	58.2%	55.2%	50.4%	47.8%	47.9%	47.9%	48.8%
Amount of fossil carbon (Gg)	164	221	433	561	578	596	611
Amount of organic carbon (Gg)	544	561	938	909	926	988	1,046

1) WIP, Waste incineration plant (Not included incineration plant for specific waste streams as sewage sludge.), listed are the residential waste fractions; for waste fractions of other waste types (considered fixed in time), see Joosen and De Jager, 2003).

described in detail in Joosen and De Jager (2003) and in the monitoring protocol. CH<sub>4</sub> emissions are based on the total amount of energy in incinerated waste and therefore an emission factor of 30 kg/TJ is applied. Based on measurement data (Spoelstra, 1993), an emission factor of 20 g/ton waste is applied for N<sub>2</sub>O from incineration with SCR. For Incineration with SNCR an emission of 100 g/ton is applied. The percentage SCR increased from 6% in 1990 to 39% in 2008.

In 2005 the carbon fraction of the household waste fraction and the percentage fossil of these fractions were determined. These values are still used for the calculation of the fossil and not fossil emissions from household waste. For the other fraction, the older values are still used (Bosselaar and Gerlagh, 2006).

### 8.4.3 Uncertainties and time-series consistency

#### Uncertainties

The Tier 1 uncertainty analysis is shown in Tables A7.1 and A7.2 in Annex 7 provides estimates of uncertainties according to IPCC source category and gas. The uncertainty in annual CO<sub>2</sub> emissions from Waste incineration is estimated at 11%. The main factors influencing these emissions are the total amount being incinerated. The fractions of different waste components used for calculating the amounts of fossil and organic carbon in the waste (from their fossil and organic

carbon fraction) and the corresponding amounts of fossil and organic carbon in the total waste incinerated. The uncertainty in the amounts of incinerated fossil waste and the uncertainty in the corresponding emission factor are estimated to be 10% and 5% respectively. These figures are based on expert judgment.

#### Time-series consistency

The time-series are based on consistent methodologies for this source category. The time-series consistency of the activity data is considered to be very good due to the continuity of the data provided by Statistics Netherlands.

#### 8.4.4 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures that are discussed in Chapter 1.

#### 8.4.5 Source-specific recalculations

All emissions prior to 2004 are recalculated. The emission calculation in the previous submissions was erroneous and not in line with the activity data. This has been corrected in this submission. The activity data have not been recalculated. Please note that the emissions are reported under Category 1.A.1.

#### 8.4.6 Source-specific planned improvements

There are no source-specific improvements planned for this category.

### 8.5 Other waste handling [6D]

#### 8.5.1 Source category description

This source category, which consists of the CH<sub>4</sub> and N<sub>2</sub>O emissions from composting separately collected organic waste from households, is not considered to be a key source. Emissions from small-scale composting of garden waste and food waste by households are not estimated as these are assumed to be negligible. It should be noted that non-CO<sub>2</sub> emissions from the combustion of biogas at wastewater treatment facilities are allocated to category 1A4 “Fuel combustion – Other sectors” because this combustion is partly used for heat or power generation at the treatment plants.

The amount of composted organic waste from households increased from nearly 0 million tons up to 1.3 million tons in 2008. In 2008 there were 23 industrial composting sites in operation; these accounted for 1% of the emissions in the Waste sector in that year (see Table 8.1).

#### 8.5.2 Methodological issues

##### Activity data and (implied) emission factors

Detailed information on activity data and emission factors can be found in the monitoring protocol 0076 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl).

The activity data for the amount of organic waste composted at industrial composting facilities are mainly based on the annual survey performed by the Working Group on Waste Registration at all industrial composting sites in the Netherlands. Data can be found on the website [www.uitvoeringafvalbeheer.nl](http://www.uitvoeringafvalbeheer.nl) and in a background document (SenterNovem, 2009). This document also contains the amount of compost produced on a yearly basis.

The emission factors are based on the average emissions (per metric ton of composted organic waste) of a number of facilities that were measured in the late 1990s (during a large-scale monitoring program in the Netherlands). Recently the emission factors have been measured again (at three facilities, one measurement per facility) in the Netherlands. The average of these three measurements for methane was much lower than the applied emission factor, with a wide range. Because of the small number of measurements and the wide range of values these new insights have not been used.

A more detailed description of the method used and the emission factors can be found in protocol 0076 on the website [www.greenhousegases.nl](http://www.greenhousegases.nl) as indicated in Section 8.1.

A country-specific methodology is used for estimating the industrial composting of organic food and garden waste from households. Since this source is not considered to be a key source, the present methodology level complies with the general IPCC Good Practice Guidance (IPCC, 2001). No

mention is made of a method for estimating the industrial composting of organic waste in the Good Practice Guidance.

#### 8.5.3 Uncertainties and time-series consistency

##### Uncertainty

The emissions of this source category are calculated using an average emission factor that has been obtained from the literature. Given the large scatter in reported emission factors the uncertainty is estimated to be more than 100%.

##### Time-series consistency

The time-series consistency of the activity data is very good due to the continuity in the data provided.

#### 8.5.4 Source-specific QA/QC and verification

The source categories are covered by the general QA/QC procedures that are discussed in Chapter 1.

#### 8.5.5 Source-specific recalculations

There are no source-specific recalculations compared to the previous submission.

#### 8.5.6 Source-specific planned improvements

In 2007 the NIR 2006 was reviewed by an ERT. As a result of the review, the ERT recommended to investigate the application of compost to land and report the emissions from this application. In 2008 a study began to collect this information. In this study also the wide range of applied emission factors was examined. The results will be (internationally) peer reviewed in the first half of 2010. This peer review might result in new emission factors for the next submission

After finalizing this study the emissions from the application of compost to land will be reported.

# Other [CRF Sector 7]

9

The Netherlands allocates all emissions in Sectors 1 to 6;  
there are no sources of greenhouse gas emissions included in  
Sector 7



# Recalculations and improvements

## ■ Major changes compared to the National Inventory Report 2009

This chapter addresses the changes in emissions compared to the previous submission.

For the submission of this NIR 2010, the data for the most recent year (2008) were added to the corresponding Common Reporting Format (CRF). This submission also includes recalculated data for CH<sub>4</sub> emissions from manure management in agriculture. Furthermore during the compilation of this NIR some errors from previous submissions were detected and corrected. These result in minor changes in emissions over the total period 1990-2008.

Responding to the recommendations of the ERT the method for estimation of F-gases from refrigeration and airconditioning was improved.

For more details on the effect and justification of the recalculations, the reader is referred to Chapters 3–8.

## 10.1 Explanation and justification for the recalculations

### 10.1.1 GHG inventory

For this submission (NIR 2010), the Netherlands uses the CRF reporter software 3.3.22. The present CRF tables are based on improved methodologies after the UNFCCC review in 2009. These improved methodologies are also described in the (updated) monitoring protocols 2010 (see Annex 6).

This chapter summarises the relevant changes in emission figures compared to the NIR 2009 (and CRF version 1.3).

A distinction is made between:

- methodological changes: new emission data are reported resulting from revised or new estimation methods; improved emission factors or activity data are also captured in recalculations as a result of methodological changes
- allocation: changes in the allocation of emissions to different sectors (only affecting the totals per category or sector)
- error corrections: correction of incorrect data

#### 10.1.1.1 Methodological changes

As part of the QA/QC activities in the Netherlands, the process of assessing and documenting methodological changes has been improved. This is now done using a brief check list for timely discussion with involved experts and users of information on likely changes. This process should improve peer review of and timely documentation on the background and justification of changes.

In response to the ERT report 2009, one methodological change was implemented in this submission: stationary cooling (2F1).

Furthermore information about pre-charged refrigeration equipment and new SF<sub>6</sub> emissions from the source Electrical equipment came available. As a result the emissions of these sources have been changed for a number of years.

#### 10.1.1.2 Source allocation

Due to new statistical information, the allocation of the fuel consumption of natural gas in the categories 1.AA.4 Commercial and institutional and 1.AA.4. B Agriculture could be improved. Emissions are now also allocated according to the new split. Because the emission factors for CH<sub>4</sub> and N<sub>2</sub>O in the two categories are different the total CO<sub>2</sub> for the two categories changed to a limited extent (-0.5 Gg CO<sub>2</sub> eq in 1990 and +2.2 Gg CO<sub>2</sub> eq in 2007) compared to the previous submission. Furthermore an allocation error related to the fuel consumption of liquid fuels in 1990 was detected, emissions shifted from category 1A.2 to 2B (42.9 CO<sub>2</sub> eq).

#### 10.1.1.3 Error correction

Following the improvements from last year, the Netherlands used a new automated system to transfer the activity (and emission) data from the National Inventory Database to the CRFReporter. Most of the activity data and emissions in this submission were loaded in the CRF Reporter using this system. In doing so, minor errors in the data in the CRF of last submission were eliminated. These error corrections (in the order of magnitude of max ± 50 Gg CO<sub>2</sub> eq per category) will not be explained in detail below. The automated transfer of



Differences between NIR 2009 and NIR 2010 for the period 1990–2007 due to recalculations  
(unit: Tg CO<sub>2</sub> eq; for F-gases: Gg CO<sub>2</sub> eq).

Table 10.1

Gas	Source	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
CO <sub>2</sub> Incl. LUCF	NIR09	161.9	173.0	172.1	177.6	178.1	182.0	183.3	178.2	174.9	175.2
	NIR10	<b>161.9</b>	172.9	172.2	177.6	178.1	182.0	183.3	178.2	174.9	174.8
	Diff.	0.0%	-0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	-0.2%
CO <sub>2</sub> Excl. LUCF	NIR09	159.3	170.7	169.6	175.2	175.7	179.6	181.0	175.8	172.5	172.7
	NIR10	<b>159.3</b>	170.6	169.7	175.3	175.8	179.7	181.0	175.8	172.5	172.3
	Diff.	0.0%	-0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	-0.2%
CH <sub>4</sub>	NIR09	25.5	24.2	19.8	19.3	18.4	17.9	17.6	17.2	16.8	17.0
	NIR10	<b>25.5</b>	24.1	19.8	19.3	18.4	17.9	17.6	17.2	16.8	16.8
	Diff.	0.0%	0.0%	-0.1%	-0.1%	-0.1%	-0.1%	0.0%	0.0%	0.0%	-0.7%
N <sub>2</sub> O	NIR09	20.2	21.5	19.3	18.1	17.3	16.9	17.4	17.3	17.1	15.6
	NIR10	<b>20.2</b>	21.5	19.3	18.1	17.3	16.9	17.4	17.3	17.1	15.4
	Diff.	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.2%
PFCs Gg	NIR09	2,264	1,938	1,582	1,489	2,187	621	286	266	257	327
	NIR10	2,264	<b>1,938</b>	1,582	1,489	2,187	621	286	266	257	323
	Diff.	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-1.2%
HFCs Gg	NIR09	4,432	6,020	3,829	1,469	1,541	1,377	1,507	1,358	1,566	1,738
	NIR10	4,432	<b>6,018</b>	3,891	1,561	1,653	1,511	1,650	1,514	1,728	1,845
	Diff.	0.0%	0.0%	1.6%	6.2%	7.2%	9.7%	9.5%	11.5%	10.3%	6.2%
SF <sub>6</sub> Gg	NIR09	217	301	319	323	283	243	246	238	202	214
	NIR10	217	<b>301</b>	318	322	281	242	265	254	217	226
	Diff.	0.0%	0.0%	-0.3%	-0.4%	-0.5%	-0.8%	7.5%	6.9%	7.1%	5.6%
Total Excl. LUCF	NIR09	212.0	224.7	214.4	215.9	215.5	216.7	218.0	212.2	208.5	207.5
	NIR10	212.0	224.5	214.6	216.0	215.6	216.8	218.2	212.4	208.7	206.9
	Diff.	0.0%	-0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	-0.3%
Total Incl. LUCF	NIR09	214.6	227.0	216.9	218.3	217.8	219.0	220.4	214.6	210.9	210.0
	NIR10	214.6	226.9	217.1	218.4	217.9	219.2	220.5	214.7	211.1	209.5
	Diff.	0.0%	-0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	-0.3%

Note: base year values as applied for the calculation of the Assigned Amount are indicated in bold.

the emissions of the precursor gases will be introduced in the next submission.

Furthermore the following error corrections and or improvements were made for this submission:

- As response to the review more detailed data on F-gas use and consumption were made available. Due to these improved data the emissions could be calculated in more detail according to the protocol. This resulted in changed emission figures for SF<sub>6</sub> and HFCs (2.F.9 and 2.F.1) in most recent years. For the base year (1995) the emissions of F-gases decreased by 1.6 Gg CO<sub>2</sub> eq and increased by 115.7 Gg CO<sub>2</sub> eq in 2007 compared to the latest submission.
- In general, the 2007 figures were updated whenever improved statistical data became available after the submission of 2009. The far most important is the removal of the erroneous emission of CO<sub>2</sub> and CH<sub>4</sub> in the categories 1.A.1.C and 1.B.2. C.2.3 Flaring. This error correction was already announced in the NIR 2009 and amounts to a reduction of 152 Gg CO<sub>2</sub> eq.
- In category 1.AA.1. A public electricity the basic activity data for waste combustion were analysed in detail. Errors in the emission calculations (CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>) prior to 2005 were discovered and corrected (+9.1 Gg CO<sub>2</sub> eq in 1990).
- An historic error in the CO<sub>2</sub> emission from category 2.D.2 Food and Drink in 1990 was removed (-0.06 Gg CO<sub>2</sub> eq in 1990).

- In 2005 an error was made while a new method to calculate the feed intake by cattle was applied. Feed intake is the basis for the calculation of the methane emission factor from enteric fermentation by cattle and also the N excretion. All these errors have been corrected in the NIR 2009. Recalculation uncovered a few errors in volume excretion for the years 1990-2000. These errors are corrected in this NIR. The effect of the recalculation is minor and has resulted to a very small increase of the CH<sub>4</sub> emission in 1990 (2.7 Gg CO<sub>2</sub> eq). Furthermore for 2007 an error was detected in the calculation of direct soil emissions (4.D). Correcting this error reduced the emission with 180 Gg CO<sub>2</sub> eq).
- An historic error in the CH<sub>4</sub> emission from category 4.B Manure Management in 1997 was removed (-164 Gg CO<sub>2</sub> eq in 1997).

Due to the above mentioned error corrections the national emissions in 1990 increased with 0.006 Tg CO<sub>2</sub> eq. For 1995 the corrections led to a decrease in emissions of 0.02 Tg CO<sub>2</sub> eq. The total national emission decreased for 2007 with 0.6 Tg CO<sub>2</sub> eq compared to the latest submission.

All above changes in previous data (methodological, allocation and error correction) are explained in the CRF.

#### 10.1.2 KP-LULUCF inventory

Not applicable, the Netherlands reports the KP data for the first time in the NIR 2010.

**Differences between NIR 2009 and NIR 2010 with respect to emission trends during the period 1990–2007 (Units: Gg CO<sub>2</sub> eq, rounded).**

**Table 10.2**

Gas	Trend (absolute)Ç			Trend (%age)		
	NIR 2009	NIR 2010	Difference	NIR 2009	NIR 2010	Difference
CO <sub>2</sub> eq [Gg] <sup>1)</sup>						
CO <sub>2</sub>	13,198	13,184	-14	8.4%	8.1%	-0.3%
CH <sub>4</sub>	-8,714	-8,713	2	-33.6%	-34.1%	-0.5%
N <sub>2</sub> O	-3,083	-3,081	2	-22.8%	-23.7%	-0.9%
HFCs	-2,866	-2,704	162	-60.8%	-59.5%	1.3%
PFCs	-2,008	-2,008	0	-85.6%	-85.7%	-0.2%
SF <sub>6</sub>	-15	-1	14	-1.6%	4.0%	5.5%
Total	-3,488	-3,322	166	-2.1%	-2.4%	-0.3%

<sup>1)</sup> Excluding LULUCF

## 10.2 Implications for emission levels

### 10.2.1 GHG inventory

This chapter outlines and summarises the implications of the changes described in Section 10.1, for the emission levels over time. Table 10.1 elaborates the differences between the submissions from last year and the current NIR with respect to the level of the different greenhouse gases. More detailed explanations are elaborated in the relevant Chapters 3–8.

#### 10.2.1.1 Effect of recalculations on base year and 2007 emission levels

Table 10.1 gives the changes due to the recalculations for the base year 1990, 1995, 2000 and 2005 to 2007 (compared to the NIR 2009).

### 10.2.2 KP- LULUF inventory

Not applicable, the Netherlands reports the KP data for the first time in the NIR 2010.

## 10.3 Implications for emission trends, including time-series consistency

### 10.3.1 GHG inventory

In general, the recalculations improve both the accuracy and time-series consistency of the estimated emissions. Table 10.2 presents the changed trends in the greenhouse gas emissions during this period due to the recalculations carried out.

### 10.3.2 KP- LULUF inventory

Not applicable, the Netherlands reports the KP data for the first time in the NIR 2010.

## 10.4 Recalculations, response to the review process and planned improvements

### 10.4.1 GHG inventory

#### 10.4.1.1 Recalculations

No recalculations are anticipated in the next submission of the CRF.

#### 10.4.1.2 Response to the review process

##### Public and peer review

Drafts of the NIR are subject to an annual process of general public review and a peer review.

One remark was received from the public on the draft NIR 2010 of January 2010. This issue (transfer of CO<sub>2</sub> from refinery to greenhouse horticulture and food industry) was clarified in the present NIR 2010.

The peer review includes a general check on all chapters. In addition, a special focus is given to a specific sector or topic each year. This year, a separate study (Hanschke et al, 2010) focused on the emissions in the transport sector. In general, the conclusion of this peer review report is that the used methodologies are in line with the IPCC Guidelines and that the transport chapter of the NIR and the protocols are written in a clear and consistent manner with good quality. For the various subsectors in transport, the report contains recommendations for improvement of transparency of methodological descriptions, correction of possible inconsistencies and inaccuracies in the text.

##### UNFCCC reviews

The NIR 2009 was reviewed in the fall of 2009 and the report (FCCC/ARR/2009/NLD) was published on 16 December 2009. In Table 10.3 contains the improvements made in response to UNFCCC review 2009

#### 10.4.1.3 Completeness of sources

The Netherlands' greenhouse gas emission inventory includes all sources identified by the Revised Intergovernmental Panel on Climate Change (IPCC) Guidelines (IPCC, 1996) with the exception of the following, very minor, sources:

- oil transport (1B2a3), due to missing activity data
- charcoal production (1B2) and use (1A4), due to missing activity data
- CO<sub>2</sub> from asphalt roofing (2A5), due to missing activity data
- CO<sub>2</sub> from road paving (2A6), due to missing activity data
- CH<sub>4</sub> from enteric fermentation of poultry (4A9), due to missing emission factors
- N<sub>2</sub>O from industrial waste water (6B1), due to negligible amounts
- Precursor emissions (i.e. carbon monoxide (CO), nitrogen oxide (NO<sub>x</sub>), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (S<sub>2</sub>)) from Memo item "International bunkers" (international transport) have not been included.

Recommendations ERT	Reference	Improvement made
<b>General</b>		
<i>Improve completeness</i>	10	First order estimates were made which proved the emissions are very minor. In addition, emissions from lime production have been reported under process emissions of sugar production. The notation key NE is replaced by IE. Finally, attempts are made to retrieve basic activity data (road paving and asphalt roofing, see Annex 5) to calculate the emissions on a regular basis which could be included in the CRF (NIR 2011).
<i>Use of annotated outline of the NIR</i>	15	Included in this submission
<i>KSA with and without LULUCF in NIR</i>	21	Annex 1 of the NIR includes the results of KSA including and excluding LULUCF
<i>Complete KS table 7 in CRF in latest inventory year</i>		Now improved and included in 2008 CRF. Please note that we were not able to indicate including and excluding LULUCF together in the CRF table. This is a bug in the CRF reporter. We choose to report all KS in the table (including LULUCF).
<i>Verify uncertainty estimates</i>	24	Based on the study by Ecofys on uncertainties and QA/QC procedures (Ecofys, 2010), improvements have been made in the present NIR by adding sections on uncertainties and QA/QC to the monitoring protocols. Other recommendations will be implemented in the NIR 2011.
<i>Improve consistency in documentation recalculations</i>	26	All changes shown in CRF table 8 are addressed in Chapter 10 of the NIR
<i>Provide information on ETS data use</i>	28	In 2009, a comparison of the ETS data and the data in the inventory was made. Conclusion was that due to definition differences the use of ETS data will not improve the emission inventory.
<i>Include information from protocols in NIR</i>	31/33	From 2009 onwards the Netherlands improved the 'balance' between NIR, protocols and background reports. This process started in 2009 and was finalised in 2010. For example, we included relevant emission factors in chapter 3. Furthermore, we improved the links between NIR and the protocols.
<i>Establish a centralised archiving system</i>	34	The Netherlands has not further centralised the archiving system. However, the archiving in the Netherlands is centrally accessible for the involved staff as well for NL Agency/NIE). "The review team recommended the further centralisation of the archiving of intermediate calculations. Most documentation and archiving are already centralised, with the exception of some intermediate/supporting data calculations archived at task force level. This recommendation will also be considered during the data process in the coming years." At the moment, we are discussing the need and options to further centralise these intermediate/supporting data calculations into the central database. The present arrangements will also be one of the special themes for the coming audit.
<b>Energy</b>		
<i>Improve completeness</i>	42	Based on figures from industry, we could assess that the emissions from lime production in the Netherlands were included in the figures of the food industry. Thus, the inventory is complete for lime production. The other sectors addressed have not yet been estimated, but efforts will be made to assess the availability of activity data to perform emission calculations.
<i>Increase use of plant specific EF</i>	45	The current inventory makes use of available plant specific EFs for Refineries, Chemical Industry, Iron and Steel and Electricity producers and will continue to do so.
<i>Documentation on use of ETS data in QA/QC</i>	46	A comprehensive note on this issue is available to UNFCCC ERT teams but not included in the NIR (see also recommendation 28)
<i>Include corrected data for jet kerosene in 2007</i>	51	Error correction performed
<i>Documentation on oxidation losses for chemical waste gases</i>	53	The Netherlands considers commissioning a study into this matter to update the data.
<i>Include corrected data for 1.AA.1.C 2007</i>	54	Error correction performed
<i>Improvement of notation keys</i>	55	Included in this submission
<b>Industrial processes and solvent and other product use</b>		
<i>Improve completeness</i>	58	See recommendation 42 Furthermore, the potential emissions of F-gases are included in the present NIR.
<i>Improve use of notation keys</i>	59	Recommendations of ERT on use of notation keys are followed up. The issue of confidentiality is elaborated in the NIR.
<i>Reorganise NIR information</i>	60	Structure of NIR is now in conformity with the latest recommendations of the UNFCCC
<i>Improve documentation of allocation of emissions</i>	64	Now improved in the NIR
<i>Improve documentation of methods for air-conditioning and refrigeration</i>	66	Now improved in the NIR, and recommended changes in method were made
<i>Improve documentation of EF for Cement Production</i>	67	Now improved in the NIR
<i>Improve documentation of method for SF<sub>6</sub> emissions</i>	68	Now improved in the NIR

Recommendations ERT	Reference	Improvement made
<b>Agriculture</b>		
<i>Include background data from NIR in CRF</i>	70	Not implemented completely because NIR and CRF are both available to reviewers
<i>Explain differences in animal statistics</i>	73	The animal statistics in the CRF are consistent with the national statistics at CBS. These data are also reported to the FAO. The observed differences may be due to definition differences or differences in reporting years.
<i>Include anaerobic treatment of animal waste in inventory</i>	75	The biogas from this activity is reported under 1AA4C
<i>Improve documentation on manure export</i>	76	Now improved in the NIR
<i>Improve documentation on N-excretion</i>	77	Now improved in the NIR
<i>Correct two inconsistencies</i>	78	Now improved in the NIR
<b>Waste</b>		
<i>Provide documentation on EF used in wastewater handling</i>	90	Reference to the relevant protocol is made in the NIR
<i>Provide documentation on recalculations and QA/QC</i>	91	The recalculations performed are described in Chapter 10 in the NIR. The source specific QA/QC for waste incineration was the cause of the recalculations.
<b>National Registry</b>		
<i>National registry improvement</i>	96	See NIR Part 2, Section 14.1.1
<i>DES (Data Exchange Standards) related measures</i>	96	See NIR Part 2, Section 14.1.2
<i>Public information</i>	96	See NIR Part 2, Section 14.1.3

For more extended information on this issue, see Annex 5.

#### 10.4.1.4 Completeness of the CRF files

For the years 1991–1994, energy data are less detailed for all industrial source categories than in both the preceding and following years, but they adequately cover all sectors and source categories. All emissions are specified per fuel type (solid, liquid and gaseous fossil fuels). Coal-derived gases (coke oven gas, blast furnace gas etc.) are included in Solid fuels and refinery gases and residual chemical gases are included in Liquid fuels (also LPG, except for Transport). The fuel category Other Fuels is used to report emissions from fossil waste in waste incineration (included in 1A1a).

Since the Industrial processes source categories in the Netherlands often comprise only a few companies, it is generally not possible to report detailed and disaggregated data. Activity data are confidential and not reported when a source category comprises three (or fewer) companies.

Potential emissions (total consumption data) for PFCs and SF<sub>6</sub> are not reported due to the confidentiality of the consumption data. A limited number of companies report emissions or consumption data, and actual estimates are made on the basis of these figures. Data to estimate potential emissions, however, are confidential (Confidential Business Information).

#### 10.4.1.5 Planned improvements

The Netherlands National System was established by the end of 2005, in line with the requirements under the Kyoto Protocol and under the EU Monitoring Mechanism. The establishment of the National System was a result of the implementation of a monitoring improvement programme (see Section 1.6). In 2007, the system was reviewed during the initial review. The review team concluded that the Netherlands' National System has been established in accordance with the guidelines for national systems under Article 5, Section 1, of the Kyoto Protocol (decision 19/CMP.1) and that it meets the

requirements for implementation of the general functions of the National System as well the specific functions of inventory planning, inventory preparation and inventory management.

#### Monitoring improvement

The National System includes an annual evaluation and improvement process. The evaluation is based on experience in previous years, and results of UN reviews, peer reviews and audits. Where needed, improvements are included in the annual update of the QA/QC program (SenterNovem, 2009).

One of the recent improvement actions relates to the emission factor (EF) for natural gas. This EF has been calculated on a yearly basis for a number of years, using detailed data from the gas supply companies. The annual EF was established in this way for the NIR 2006, for 2004 and the base year 1990. For both years, the emission factor proved to be 56.8. Given the time constraints, the EF for intermediate years was assumed to be constant. In 2009, a study analysed this further using two further sample years and the conclusion was that annual fluctuations in intermediate years were very minor. It was therefore decided not to carry out more detailed assessment for further intermediate years and to maintain the EF for these intermediate years at 56.8, especially since these years are neither base years nor commitment period years. Since 2007, the EF has been assessed annually. The value in both 2007 and 2008 was 56.7 (Zijlema, 2008 and Zijlema, 2009), see Annex 2.

#### Monitoring protocol and QA/QC program

The Netherlands uses monitoring protocols that describe the methodology, data sources (and the rationale for their selection). These protocols are available on the website [www.greenhousegases.nl](http://www.greenhousegases.nl). The protocols were given a legal basis in December 2005. The monitoring protocols are assessed annually and –when needed– updated. The initial review recommended that some of the protocols should include more details (inclusion of some additional information that is now only included in background documents). For

2009, the Netherlands has included this recommendation in its QA/QC program and to improve the 'balance' between NIR, protocols and background reports. This process started in 2009 and was finalised in 2010.

The QA/QC program for this year (SenterNovem, 2009) also continues the assessment of improvement options in the longer term, partly based on the consequences of the new 2006 IPCC guidelines. This will provide a basis for a possible improvement program for the longer term.

In response to the ERT report 2009, one methodological change was implemented in this submission: stationary cooling. The review team recommended the further centralisation of the archiving of intermediate calculations. Most documentation and archiving is already centralised, with the exception of some intermediate/supporting data calculations archived at task force level. This recommendation will also be considered during the data process in the coming years.

#### 10.4.2 KP-LULUCF inventory

Not applicable, The Netherlands reports the KP data for the first time in the NIR 2010.

## Part II: Supplementary Information required under Article 7, Paragraph 1





## 11.1 General information

### 11.1.1 Definition of forest and any other criteria

The Netherlands identified in its Initial Report the single minimum values under Article 3.3 of the Kyoto Protocol.

The complete forest definition the Netherlands uses for Kyoto reporting is: Forest is land with woody vegetation and with tree crown cover of more than 20% and area of more than 0.5 ha. The trees should be able to reach a minimum height of 5 m at maturity in situ. They may consist either of closed forest formations where trees of various storeys and undergrowth cover a high proportion of the ground, or open forest formations with a continuous vegetation cover in which tree crown cover exceeds 20%. Young natural stands and all plantations established for forestry purposes which have yet to reach a crown density of 20% or tree height of 5 m are included under forest, as areas normally forming part of the forest area which are temporally unstocked as a result of human intervention or natural causes but which are expected to revert to forest. Forest Land also includes:

- forest nurseries and seed orchards that constitute an integral part of the forest;
- forest road, cleared tracts, firebreaks and other small open areas, all smaller than 6 m. within the forest;
- forest in national parks, nature reserves and other protected areas such as those of special environmental, scientific, historical, cultural or spiritual interest, with an area of more than 0.5 ha and a width of more than 30m;
- windbreaks and shelterbelts of trees with an area of more than 0.5 ha and a width of more than 30 m.

This excludes tree stands in agricultural production systems, for example, in fruit plantations and agro forestry systems.

This definition is in line with the FAO reporting since 1984 and was chosen within the ranges set by the Kyoto Protocol. The definition matches the subcategory “Forests according to the Kyoto definition” (abbreviated as FAD) of Forest Land in the inventory under the Convention on Climate Change.

### 11.1.2 Elected activities under Article 3, paragraph 4, of the Kyoto Protocol

The Netherlands has not elected any activities to include under Article 3, paragraph 4, of the Kyoto Protocol.

### 11.1.3 Description of how the definitions of each activity under Article 3.3 and each elected activity under Article 3.4 have been implemented and applied consistently over time

Units of land subject to Article 3.3 afforestation and reforestation are reported jointly and are defined as units of land that did not comply with the forest definition on 1 January 1990 and do so at any time (that can be measured) before 31 December 2012. Land is classified as re/afforested as long as it complies with the forest definition.

Units of land subject to Article 3.3 deforestation are defined as units of land that did comply to the forest definition at in time on or after 1 January 1990, and again ceased to comply to this forest definition at any moment in time (that can be measured) after 1 January 1990. Once land is classified as deforested, it remains in this category, even if it is reforested and thus complies with the forest definition again later in time.

### 11.1.4 Description of precedence conditions and/or hierarchy among Article 3.4 activities, and how they have been consistently applied in determining how land was classified.

This is not applicable as no article 3.4 activities have been elected.

## 11.2 Land-related information

### 11.2.1 Spatial assessment unit used for determining the area of the units of land under Article 3.3

The Netherlands has complete and spatially explicit land use mapping that allows for geographical stratification at 25 m x 25 m (0.0625 ha) pixel resolution (Kramer et al., 2009). This corresponds with the wall-to-wall approach used for reporting under the Convention (approach 3 in GPG-LULUCF chapter 2) and is described as reporting method 2 in GPG-LULUCF for Kyoto (par. 4.2.2.2). ARD activities are recorded on a pixel basis. For each pixel individually, it is known whether it is part of a patch that complies to the forest definition or not.

Any pixel changing from non-compliance to compliance to the forest definition is treated as re/afforestation. This may be the result of a group of clustered pixels that together cover at least 0.5 ha of non-forest land changing land use into forest land. It may also occur when one or more pixels adjacent to

year	AR land remaining AR land	Other land converted to AR land	AR land converted to D land	D land remaining D land	Other land converted to D land	Other land remaining other land	Land in KP article 3.3 ARD
1990	0	2.56	0	0	1.99	4,146.95	4.55
1991	2.56	2.56	0	1.99	1.99	4,142.40	9.10
1992	5.12	2.56	0	3.98	1.99	4,137.85	13.65
1993	7.68	2.56	0	5.98	1.99	4,133.29	18.21
1994	10.24	2.56	0	7.97	1.99	4,128.74	22.76
1995	12.80	2.56	0	9.96	1.99	4,124.19	27.31
1996	15.36	2.56	0	11.95	1.99	4,119.64	31.86
1997	17.92	2.56	0	13.95	1.99	4,115.09	36.41
1998	20.47	2.56	0	15.94	1.99	4,110.54	40.96
1999	23.03	2.56	0	17.93	1.99	4,105.99	45.51
2000	25.59	2.56	0	19.92	1.99	4,101.43	50.07
2001	28.15	2.56	0	21.91	1.99	4,096.88	54.62
2002	30.71	2.56	0	23.91	1.99	4,092.33	59.17
2003	33.27	2.56	0	25.90	1.99	4,087.78	63.72
2004	35.83	2.56	0	27.90	1.99	4,083.23	68.27
2005	38.39	2.56	0	29.88	1.99	4,078.68	72.82
2006	40.95	2.56	0	31.87	1.99	4,074.13	77.37
2007	43.51	2.56	0	33.87	1.99	4,069.57	81.93
2008	46.07	2.56	0	35.86	1.99	4,065.02	86.48

a forest patch change land use. Similarly, any pixel changing from compliance with the Kyoto forest definition to non-compliance is treated as deforestation, whether it involves the whole group of clustered pixels or just a subgroup of them. Thus, the assessment unit of land subject to ARD is 25 m x 25 m (0.0625 ha).

### 11.2.2 Methodology used to develop the land transition matrix

The Netherlands has complete and spatially explicit land-use mapping with map dates on 1 January 1990 and 1 January 2004 (Kramer et al., 2009). An overlay was made between those two maps and this resulted in a land-use change matrix between January 1990 and January 2004. Mean annual rates of change for all land-use transitions between those years was calculated by linear interpolation, and after 2004 by extrapolation. The values based on extrapolation after 1 January 2004 will be subject to recalculation when a new land-use map of later date has been created. Our aim is to make land-use maps for 1 January 2008 and 1 January 2013, ensuring that we are able to capture land-use changes between 1990 and 2008, and between 2008 and 2012 (IPCC, 2003).

Thus, in table NIR-2 the transitions from “other land” to either AR or D activities during the reporting year 2008 (last row in table NIR-2) are extrapolated values based on the mean annual rate of land use change between 1990 and 2004, and will be subject to recalculation when updates of the land-use maps become available. Land subject to AR or D between 1990 and 2007 is based on the sum of

- the cumulative area under AR respectively under D for the (reporting) years 1990 to 2003, as derived from a land-use map overlay (these values can be considered as final); and
- the cumulative area under AR respectively under D for the (reporting) years 2004 to 2007, based on an extrapolation of the mean annual rate of land use change between 1990 and 2004 (these values will be subject to recalculation when updates of the land use maps become available).

Table 11.1 gives the annual values from 1990 on for the article 3.3 related cells in table NIR-2. Due to the use of extrapolation in the current submission, the values from 2004 on can be considered preliminary, with updates foreseen in the 2012 submission.

The summed values in Table 11.1 for AR (AR land remaining AR land + Other land converted to AR land) match with the sum of values reported under the Convention sector 5.A.2 land converted to Forest Land subcategory Forests according to the Kyoto definition (FAD), and Forest Land - Trees outside Forest converted to Kyoto Forest (included in Forest land – Kyoto Forest) for the respective years. The annual values for deforestation (Other land converted to D land) match with the sum of the values reported in sectors 5.B.2.1 Forest Land - FAD to 5.F.2.1 Forest Land – FAD, and Forest Land – Kyoto forest converted to Trees outside Forest (included in Forest land - Trees outside Forest) for the respective years.

### 11.2.3 Maps and/or database to identify the geographical locations, and the system of identification codes for the geographical locations

The land use information reported under both the Convention (see also par. 7.1.2) and the Kyoto Protocol is based on two maps for monitoring nature development in the Netherlands, “Basiskaart Natuur” (BN) for 1990 and 2004.

The source material for BN1990 consists of the paper topographic map 1:25,000 (Top25) and digital topographical map 1:10,000 (Top10Vector). Map sheets with exploration years in the period 1986-1994 were used. The source material for BN 2004 consists of the digital topographic map 1:10,000 (Top10Vector). All topographic maps have been explored in the period 1999-2003. For the BN 2004, information from the Top 10 vector is combined with four other sources, that is information from two subsidy regulations (information from 2004), a map with the geophysical regions of the Netherlands

Characteristics	BN 1990	BN 2004
Name	Historical Land use Netherlands 1990	Base map Nature 2004
Aim	Historical land use map for 1990	Base map for monitoring nature development
Resolution	25 m	25 m
Coverage	Netherlands	Netherlands
Base year source data	1986-1994	1999-2003
Source data	Hard copy topographical maps at 1:25,000 scale and digital topographical maps at 1:10,000	Digital topographical maps at 1:10,000 and additional sources to distinguish specific nature types
Number of classes	10	10
Distinguished classes	Grassland, Arable land, Heath land/peat moor, Forest, Buildings, Water, Reed marsh, Sand, Built-up area, Greenhouses	Grassland, Nature grassland, Arable land, Heath land, Forest, Built-up area and Infrastructure, Water, Reed marsh, Drifting sands, Dunes and beaches

(Fysisch Geografische Regio's) and a map with the land use in 2000 (Bestand BodemGebruik, 2000; Kramer et al., 2007). Table 11.2 summarises the characteristics of both maps (taken from Kramer et al., 2009).

In 2008, a series of improvements were made to the methodology for digitalisation, classification and aggregation. One of the main improvements for the 1990 map is a better distinction between built-up areas and agricultural lands. This was based on manually checking of all areas. If the source information was a paper map, it was converted to a digital high resolution raster map. Then both Top10Vector files and digitised Top25 maps were (re)classified to match the requirements set by UNFCCC reporting. In this process additional data sets were used, and the forest definition was applied to distinguish forests that comply with the minimum area and width chosen for the Kyoto Protocol (see Section 11.1.1) from other wooded areas ("Trees outside forests").

Simultaneously, harmonisation between the different source materials was applied to allow a sufficiently reliable overlay. Harmonisation included the use of road maps to check the representation of linear features, and correct for any artefact movement of roads due to differences in source material.

The final step in the creation of the land use maps was the aggregation to 25 m × 25 m raster maps. For the 1990 map (which to a large extent was based on information derived from paper maps), an additional validation step was applied to check on the digitising and classifying processes.

To distinguish between mineral soils and peat soils, an overlay was made between the two BN maps and the Dutch Soil Map (de Vries et al., 2003). The result is a map with national coverage that identifies for each pixel whether it was subject to AR or D between 1990 and 2004, and whether it is located on a mineral or on an organic soil.

Following this procedure, the status as re/afforested area or deforested area is certain for each of the individual locations on the map that were subject to ARD between 1990 and 2004. However, it is unknown for each individual location when exactly this occurred. A mean annual rate for the Netherlands as a whole is derived from this by interpolating. For ARD occurring after 1 January 2004 until the reporting year, the mean annual rate for ARD activities is derived by

extrapolating the mean annual rates between 1990 and 2004. As such, the exact location of ARD activities after 2004 is not known. The location will be specified as soon as a new land use map of later date is created. All ARD will be recalculated for the years where extrapolated data have been used.

### 11.3 Activity-specific information

#### 11.3.1 Methods for carbon stock change and GHG emission and removal estimates

##### Description of the methodologies and the underlying assumptions used

The linkage between AR the reporting based on land use (sub)categories for the Convention is as follows:

- 5.A.2.1 Cropland converted to Forest Land – Forests according to the Kyoto definition
- 5.A.2.2 Grassland converted to Forest Land – Forests according to the Kyoto definition
- 5.A.2.3 Wetland converted to Forest Land – Forests according to the Kyoto definition
- 5.A.2.4 Settlement converted to Forest Land – Forests according to the Kyoto definition
- 5.A.2.5 Other Land converted to Forest Land – Forests according to the Kyoto definition as well as the conversion from 5.1.1. Trees outside Forest to Forests according to the Kyoto definition, included in 5.1.1. Forests according to the Kyoto definition.

The methodologies used to calculate carbon stock changes due to AR activities are in accordance with those under the Convention as presented in Section 7.2.4. The carbon stock changes due to changes in biomass were attributed to above-respectively below-ground biomass using one average R value derived from the plots 0-20 years old (Van den Wyngaert et al., 2009). Carbon stock changes in dead wood and litter are not reported (see Section 11.3.1.2). Carbon stock changes in mineral and organic soils are not reported in this submission, but will be from next year onwards. A recalculation will be made for 2008.

The linkage between D and the reporting based on land use (sub)categories for the Convention is as follows:

- 5.B.2.1 Forest Land – Forests according to the Kyoto definition converted to Cropland

Year	AR - CSC in AB biomass	AR - CSC in BG biomass	D - CSC in AB biomass	D - CSC in BG biomass	D - CSC in litter	D - CSC in dead wood
1990	0.4	0.2	-90.1	-18.0	-51.3	-0.8
1991	1.3	0.6	-91.7	-18.3	-52.3	-1.3
1992	2.6	1.2	-93.8	-18.8	-53.3	-1.7
1993	4.3	1.9	-95.7	-19.1	-54.3	-2.2
1994	6.4	2.9	-97.9	-19.6	-55.3	-2.7
1995	9.0	4.1	-100.0	-20.0	-56.3	-3.2
1996	11.9	5.4	-102.1	-20.4	-57.3	-3.7
1997	15.4	7.0	-104.4	-20.8	-58.3	-4.1
1998	19.2	8.7	-106.4	-21.2	-59.3	-4.6
1999	23.5	10.7	-108.6	-21.6	-60.3	-5.1
2000	28.2	12.8	-107.8	-20.4	-61.3	-2.6
2001	39.7	18.0	-110.3	-20.9	-62.3	-3.1
2002	46.3	21.0	-112.8	-21.4	-63.3	-3.5
2003	53.4	24.3	-115.5	-21.9	-64.3	-4.0
2004	61.0	27.7	-118.2	-22.4	-64.3	-4.4
2005	69.2	31.4	-120.8	-22.9	-64.3	-4.9
2006	77.8	35.4	-123.4	-23.3	-64.3	-5.4
2007	87.0	39.5	-125.9	-23.8	-64.3	-5.8
2008	96.6	43.9	-128.3	-24.2	-64.3	-6.3

5.C.2.1 Forest Land – Forests according to the Kyoto definition converted to Grassland

5.D.2.1 Forest Land – Forests according to the Kyoto definition converted to Wetland

5.E.2.1 Forest Land – Forests according to the Kyoto definition converted to Settlement

5.F.2.1 Forest Land – Forests according to the Kyoto definition converted to Other Land

as well as the conversion from Forests according to the Kyoto definition to Trees outside Forest and, included in 5.1.1. Trees outside Forest

The methodologies used to calculate carbon stock changes due to D activities are in accordance with those under the Convention as presented in Section 7.2.4. The carbon stock changes due to changes in biomass change were differentiated in above respectively below ground biomass using data available from the simple bookkeeping model used (Van den Wyngaert et al., 2009). All emissions were attributed to the year of deforestation, and no emissions were reported for any other years. As under the Convention, emissions as well as areas under deforestation are reported on an annual basis, while under the KP areas are reported cumulative while all emissions are reported in the year of deforestation, emissions are equal under both reporting, but implied emission factors are different.

Carbon stock changes in mineral and organic soils are not reported in this submission, but will be from next year on. A recalculation will be made for 2008.

Liming of forest in the Netherlands might occur occasionally but no statistics are available. All liming based on quantities of product sold is attributed to agricultural land (Cropland, Grassland) which is the main sector where liming occurs. Liming is thus reported only for deforested land that is converted to any of these categories. The total amount of liming

is reported in sector 5G of the Convention and described in Section 7.8. There is no information how much of the total amount of lime is applied on croplands and grasslands that are reported under deforestation (as opposed to other croplands and grasslands). A mean per ha lime application was calculated based on the total amount of lime applied and the total area under grassland and cropland. This was multiplied with the total area of grassland and cropland reported under article 3.3 deforestation to calculate the amount of CO<sub>2</sub> emission due to liming.

#### 11.3.1.2 Justification when omitting any carbon pool or GHG emissions/removals from activities under Article 3.3 and elected activities under Article 3.4

##### Carbon stock change due to changes in dead wood and litter in units of land subject to article 3.3 ARD

The national forest inventory provides an estimate for the average amount of litter (in plots on sandy soils only) and the amount of dead wood (all plots). The data provide the age of the trees and assume that the plots are no older than the trees. As such, the age of the plot does not provide or predict any litter accumulation from previous forests. This is illustrated in a very weak relationship between plot age and carbon in litter, and a large variation in dead wood even for young plots. As good data to relate carbon accumulation in litter and dead wood since the time of re/afforestation are lacking for the Netherlands, we report this carbon sink conservatively as zero.

##### Carbon stock change due to changes in soil carbon in units of land subject to article 3.3 ARD

Carbon stock changes in mineral and organic soils are not reported in this submission. Research is ongoing on this topic and results are expected to be available prior to next submission. It is foreseen that for the next submission, a recalculation will be made for the year 2008.

Year	AR - CSC in AB biomass	AR - CSC in BG biomass	D - CSC in AB biomass	D - CSC in BG biomass	D - CSC in litter	D - CSC in dead wood
1990	0.4	0.1	0.5	0.1	0.0	0.0
1991	0.8	0.2	1.0	0.2	0.0	0.0
1992	1.2	0.2	1.4	0.3	0.0	0.0
1993	1.6	0.3	1.9	0.4	0.0	0.0
1994	1.9	0.4	2.3	0.5	0.0	0.0
1995	2.3	0.4	2.8	0.5	0.0	0.0
1996	2.7	0.5	3.2	0.6	0.0	0.0
1997	3.0	0.6	3.7	0.7	0.0	0.0
1998	3.4	0.7	4.1	0.8	0.0	0.0
1999	3.8	0.7	4.6	0.9	0.0	0.0
2000	4.4	0.8	5.3	1.0	0.0	0.0
2001	4.7	0.9	5.7	1.1	0.0	0.0
2002	5.1	0.9	6.2	1.1	0.0	0.0
2003	5.4	1.0	6.6	1.2	0.0	0.0
2004	5.8	1.1	7.0	1.3	0.0	0.0
2005	6.2	1.1	7.5	1.4	0.0	0.0
2006	6.5	1.2	7.9	1.4	0.0	0.0
2007	6.9	1.2	8.4	1.5	0.0	0.0
2008	7.3	1.3	8.8	1.6	0.0	0.0

CSC : carbon stock change

AR : afforestation and reforestation

AB : above ground

D : deforestation

BG : below ground

#### **N<sub>2</sub>O emission due to nitrogen fertilisation in units of land subject to article 3.3 AR**

Forest fertilisation does not occur in the Netherlands. Therefore, fertilisation in re/afforested areas is reported NO.

#### **N<sub>2</sub>O emission due to soil disturbance associated with conversion to cropland in units of land subject to article 3.3 D**

Nitrous oxide emissions associated with disturbance of soils when deforested areas are converted to cropland are currently not estimated, but research is ongoing. Results are expected to be included in one of the next submissions.

#### **GHG emission due to biomass burning in units of land subject to article 3.3 ARD**

Greenhouse gas emissions (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) related to biomass burning are not estimated because biomass burning has not been monitored since 1996. Wildfire statistics indicated that forest fires rarely occurred in the two decades before 1996 (Wijdeven et al., 2006).

#### **11.3.1.3 Information on whether or not indirect and natural GHG emissions and removals have been factored out**

For all article 3.3 AR activities, forests were created only after 1990 and factoring out of effects on age structure of practices and activities before 1990 is not relevant. For article 3.3 D activities, the increase in mean carbon stock since 1990 may be an effect of changes in management as well as a change in age structure resulting from activities and practices before 1990. However, it is not known which factor contributes to what extent. This increase in mean carbon stock results in a higher carbon emission due to deforestation. Thus, not factoring out the effect of age structure dynamics since 1990 results in a more conservative estimate of emissions due to article 3.3 D activities.

There has been no factoring out of indirect GHG emissions and removals due to effects of elevated carbon dioxide concentrations or nitrogen deposition. To our knowledge, there is no internationally agreed methodology to factor out the effects of these that could be applied to our data.

#### **11.3.1.4 changes in data and methods since the previous submission (recalculations)**

Not applicable.

#### **11.3.1.5 Uncertainty estimates**

The Tier 1 analysis in Annex 7, Table A7.3 provides estimates of uncertainties of LULUCF categories. The Netherlands uses a Tier 1 analysis for the uncertainty assessment of the sector LULUCF. The analysis combines uncertainty estimates of the forest statistics, land use and land use change data (topographical data) and the method used to calculate the yearly growth in carbon increase and removals (Olivier et al., 2009). The uncertainty analysis is performed for Forests according to the Kyoto definition (par. 7.2.5) and is based on the same data and calculations as used for KP article 3.3 categories.

Thus, the uncertainty for total net emissions from units of land under article 3.3 afforestation/reforestation are estimated at 63%, equal to the uncertainty in Land converted to Forest Land. Similarly, the uncertainty for total net emissions from units of land under article 3.3 deforestation is estimated at 66%, equal to the uncertainty in Land converted to Grassland (which includes for the sake of the uncertainty analysis all Forest land converted to any other type of land use. See Olivier et al., 2009). As a result of recent improvements in both maps and calculations (see NIR 2009), it is likely that the current estimate is an overestimate of the actual uncertainty. It is foreseen that new uncertainty estimates will be



calculated before the final accounting for the KP commitment period.

#### 11.3.1.6 Information on other methodological issues

There is no additional information on other methodological issues

#### 11.3.1.7 The year of the onset of an activity, if after 2008

Not relevant for this submission.

### 11.4 Article 3.3

#### 11.4.1 Information that demonstrates that activities under Article 3.3 began on or after 1 January 1990 and before 31 December 2012 and are direct human-induced

The land use map is dated on 1 January 1990. Only ARD activities relative to this map, that is after this date, are taken into account.

In The Netherlands, forests are protected under the Forest Law (1961), which stipulates that “The owner of ground, on which a forest stand, other than through pruning, has been harvested or otherwise destroyed, is obliged to replant the forest stand within a period of three years after the harvest or destruction of the stand (...)”. A system of permits is applied for deforestation, and compensation forests have been planted at other locations. This has in the past created problems for (local) nature agencies that wanted to restore the more highly valued heather and peat areas in the Netherlands and as a result will not allow forest regeneration on areas where it is not intended.

With the historic and current scarcity of land in the Netherlands (which has the highest population density of Europe), any land use is the result of deliberate human decisions.

#### 11.4.2 Information on how harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation

Following the forest definition and the mapping practice applied in the Netherlands, areas subject to harvesting or forest disturbance are still classified as forests and as such will not result in a change in land use in the overlay of the land use maps (Kramer et al., 2009).

#### 11.4.3 Information on the size and geographical location of forest areas that have lost forest cover but which are not yet classified as deforested

The land use maps do not provide information on forest areas that have lost forest cover if they are not classified as deforested. However, from the national forest inventory it can be estimated that about 0.3% of the forests was classified as clear cut area, that is without tree cover.

### 11.5 Article 3.4

This is not applicable as no article 3.4 activities have been elected

### 11.6 Other information

#### 11.6.1 Key category analysis for Article 3.3 activities and any elected activities under Article 3.4

Under the Convention, conversion to Forest Land (5A2) is a key category. Despite differences in definition between forests under the Convention and under the Kyoto Protocol, 5A2 is a corresponding category and as such re/afforestation is considered a key category under the KP.

Under the Convention, conversion of Forest Land to Grassland (5C2) is a key category. Despite differences in definition between forests under the Convention and under the Kyoto Protocol, 5C2 is a corresponding category and as such deforestation is considered a key category under the KP.

Additionally, deforestation is larger than the smallest key category in the Tier 1 key source analysis (excluding LULUCF). The smallest key source based on level for Tier 1 level analysis including LULUCF is 639 Gg CO<sub>2</sub> (5A2 Land converted to Forest Land, see Annex 1). With 546.68 Gg CO<sub>2</sub> the annual contribution of re/afforestation under the KP is just below the smallest key category (Tier 1 level analysis including LULUCF). Deforestation under the KP in 2008 causes an emission of 780.45 Gg CO<sub>2</sub>, which is more than the smallest key category (Tier 1 level analysis including LULUCF).

#### 11.7 Information relating to Article 6

The Netherlands is not buying or selling emission rights from JI projects related to land subject to a project under Article 6 of the Kyoto protocol.

# Information on accounting of Kyoto units

## 12.1 Background information

The Netherlands' Standard Electronic Format report for 2009 containing the information required in paragraph 11 of the annex to decision 15/CMP.1 and adhering to the guidelines of the SEF has been submitted to the UNFCCC Secretariat electronically - SEF\_NL\_2010\_2\_14-42-58 22-4-2010.xls.

## 12.2 Summary of information reported in the SEF tables

There were 1.011.775.096 AAUs in The Netherlands' National Emission Trading Registry at the end of the year 2009, of which 842.767.016 units were in the Party holding account; 85.495.450 units in the entity holding accounts; no units in the other cancellation accounts and 83.512.630 units in the retirement account.

There were 17.241.136 CERs in the registry at the end of 2009: 12.458.380 CERs were held in the Party holding account, 4.782.736 CERs were held in the entity holding accounts, 20 CERs in the other cancellation accounts and no CERs were held in the retirement account.

There were 1.037.208 ERUs in the registry at the end of 2009: 917.099 ERUs were held in the Party holding account, 120.109 ERUs were held in the entity holding accounts, no ERUs in the other cancellation accounts or retirement account.

The registry did not contain any RMUs, t-CERs or I-CERs.

There were no units in the Article 6 issuance and conversion accounts; no units in the Article 3.3 and Article 3.4 issuance or cancellation accounts and no units in the Article 12 afforestation and reforestation accounts.

The total amount of the units in the registry corresponded to 1.030.053.440 tonnes CO<sub>2</sub> eq.

The Netherlands' assigned amount is 1.001.262.141 tonnes CO<sub>2</sub>eq.

## 12.3 Discrepancies and notifications

Annual Submission Item	Submission
15/CMP.1 annex I.E paragraph 11: Standard electronic format (SEF)	The Standard Electronic Format report for 2009 has been submitted to the UNFCCC Secretariat electronically (SEF_NL_2010_2_14-42-58 22-4-2010.xls) and the contents of the report (R1) can also be found in annex 6 of this document.
15/CMP.1 annex I.E paragraph 12: List of discrepant transactions	The list of discrepant transactions for the year 2009, pursuant to 15/CMP.1 annex I.E paragraph 12, has been submitted to the UNFCCC Secretariat electronically. The contents of the Report R2 can also be found in Annex 6 of this document. Refer to Separate Electronic Attachment "SIAR Reports 2009-NL v 1.0.xls" Worksheet R2.
15/CMP.1 annex I.E paragraph 13 & 14: List of CDM notifications	No CDM notifications occurred in 2009.
15/CMP.1 annex I.E paragraph 15: List of non-replacements	No non-replacements occurred in 2009.
15/CMP.1 annex I.E paragraph 16: List of invalid units	No invalid units exist as at 31 December 2009.
15/CMP.1 annex I.E paragraph 17: Actions and changes to address discrepancies	Under paragraph 14.1 "Previous Review Recommendations" the actions and changes to address discrepancies are discussed.



## 12.4 Publicly accessible information

Annual Submission Item	Submission
15/CMP.1 annex I.E Publicly accessible information	<p>The information as described in 13/CMP.1 annex II.E paragraphs 44-48 is publicly available at the following internet address (URL); <a href="http://www.emissieautoriteit.nl/english/public-information-kyoto">http://www.emissieautoriteit.nl/english/public-information-kyoto</a></p> <p>All required information for a Party with an active Kyoto registry is provided with the following exceptions;</p> <p>paragraph 46 Article 6 Project Information. The Netherlands does not host JI projects as laid down in National legislation. This fact is stated on the mentioned internet address. That the Netherlands does not host JI projects is implied by article 16.46c of the Environment Act (Wet milieubeheer) and explicitly stated in the explanatory memorandum to the act implementing the EC linking Directive (Directive 2004/101/EC, the Directive that links the ETS to the project based activities under the Kyoto Protocol). As is explained in the memorandum, the government decided not to allow JI projects in the Netherlands since it would only increase the existing shortage of emission allowances / assigned amount units.) paragraph 47a/d/f/ in/out/current Holding and transaction information is provided on a holding type level, due to more detailed information being declared confidential by EU regulation. This follows from article 10 of EU Regulation 2216/2004/EC, that states that "All information, including the holdings of all accounts and all transactions made, held in the registries and the Community independent transaction log shall be considered confidential for any purpose other than the implementation of the requirements of this Regulation, Directive 2003/87/EC or national law."</p> <p>paragraph 47c The Netherlands does not host JI projects as laid down in National legislation. (ref. submission paragraph 46 above)</p> <p>paragraph 47e The Netherlands does not perform LULUCF activities and therefore does not issue RMUs</p> <p>paragraph 47g No ERUs, CERs, AAUs and RMUs have been cancelled on the basis of activities under Article 3, paragraphs 3 and 4 to date.</p> <p>paragraph 47h No ERUs, CERs, AAUs and RMUs have been cancelled following determination by the Compliance Committee that the Party is not in compliance with its commitment under Article 3, paragraph 1 to date.</p> <p>paragraph 47j No ERUs, CERs, AAUs and RMUs have been retired to date</p> <p>paragraph 47k There is no previous commitment period to carry ERUs, CERs, and AAUs over from.</p>

## 12.5 Calculation of the commitment period reserve (CPR)

In April 2008 the Netherlands became eligible under the Kyoto Protocol. Its assigned amount was fixed at 1,001,262,141 tonnes CO<sub>2</sub> equivalent. The CPR was calculated at that point in time at 901,135,927 tonnes CO<sub>2</sub> equivalent. The CPR has not been changed.

## 12.6 KP-LULUCF accounting

Not applicable, because The Netherlands has chosen for end-of-period accounting for KP-LULUCF

## Information on changes in national system

Extensive information on the national inventory system is described in this National Inventory Report under the appropriate Sections as required by the UNFCCC guidelines. More extensive background information on the National System is also included in the Netherlands 5th National Communication and in the Initial Report. The Initial Review in 2007 concluded that the Netherlands National System has been established in accordance with the guidelines.

There have been no changes in the National System since the last submission and since the initial report, with the exception of following two issues:

- The coordination of the Emission Registration Project, in which emissions of about 350 substances are annually calculated, was performed until 1 January 2010 by PBL. As of 1 January 2010, coordination has been assigned to RIVM. Processes, protocols and methods remain unchanged. Many of the former experts from PBL have also shifted to RIVM.
- The name of SenterNovem (single national entity/NIE) has changed as of 1 January 2010 to NL Agency.

Both changes do not have any impact on the functions of the National System.



# Information on changes in national registry

## 14.1 Previous Review Recommendations

### 14.1.1 First Recommendation: National Registry Improvement

#### 14.1.1.1 Recommendation

“The Netherlands should further improve the measures put in place in its national registry with a view to minimizing operator errors and ensuring efficient exchange of data with other registry systems, including the international transaction log, in accordance with paragraph 115 (b) and (e) of the annex to decision 22/CMP.1”

#### 14.1.1.2. Action taken

Detailed recommendation document IAR/2009/NL/2/1 under chapter 1.2. Summary of findings, paragraph 5

‘The Netherlands should further improve the measures put in place in its national registry with a view to ensuring minimal operator errors and reliable interoperability with other registry systems, including the ITL, in accordance with paragraph 115 of the annex to decision 22/CMP.1 and paragraph 25 of the annex to decision 24/CP.8. The key recommendations are listed in paragraph 6 below.’

The detailed recommendations refer to paragraph 6. The Second Recommendation refer to the same paragraph. Therefore the action taken to fulfil this recommendation is discussed together with the Second Recommendation.

### 14.1.2 Second Recommendation: DES Related measures

#### 14.1.2.1 Recommendation

“The Netherlands should implement measures listed in paragraph 6 of the summary of findings in the SIAR by the time of its next annual submission, and report in that submission on the changes made to the national registry following the successful implementation and testing of those measures, including any test plans and test reports”

#### 14.1.2.2 Action Taken

Detailed recommendation document IAR/2009/NL/2/1 under chapter 1.2. Summary of findings, paragraph 6

Following the SIAR recommendations, the following measures should be implemented in the registry by the end of 2009

at the latest. The Netherlands should report in its next annual submission on the changes made to its registry following the successful implementation and testing of those measures, including any relevant test plans and test reports.

- a. On Data Exchange Standards (DES) response code 2008, automated internal validations should be performed on account identifiers to ensure their format is valid before messages are sent to the ITL.
- b. On DES response codes 4003 and 4010, mitigation strategies should be implemented to minimize the reuse of recently used transaction unit blocks.
- c. On DES response codes 4008, automated internal validations should be performed to ensure that transaction proposals do not contain unit blocks involved in an inconsistency.
- d. In accordance with paragraph 25(e) of the annex to decision 24/CP.8, the Netherlands should review thoroughly the design and implementation of its registry to prevent inconsistencies from occurring and provide timely and accurate records of its unit holdings and relevant transactions when inconsistencies occur.

#### Ref (a)

To reduce chance of response code 2008 being triggered a change was implemented in version 4.2 of the used Greta registry software. This version has been implemented 10 November 2009. The change added additional automated internal validations on account identifier entered by the registry user. Validations now performed are for example:

- Check that the account number entered consists of all and no more than the 4 required elements; acquiring county code, account type, account unique identifier and commitment period.
- Check that the entered account type represents a holding account if a regular internal or external transfer is performed.
- Check that the account identifier exists if an internal transfer is performed.

Because the data exchange standard (DES) does not facilitate checking the existence of a particular account identifier in an external registry prior to performing an transfer it is not possible to completely prevent users from using an invalid account identifier causing response code 2008 being triggered.

#### Ref (b)

Document IAR/2009/NL/2/1 under P2.2.10.5 states that “The response code 4003/4010 is a common discrepancy among the registries because of a limitation in the DES message flow”. The Netherlands shares this view and has been working with the Secretariat to propose a change in the DES that removes this limitation. The Netherlands has been advocating this change since before the ITL was implemented. Recently this change known under number CR13 was approved by the Change Advisory Board and it is currently scheduled to be implemented in the first half of 2010.

Pending the outcome of this change request no other mitigation strategies have been extensively considered for implementation in the Registry software. There are two reasons for this:

1. the DES change would be much more efficient and effective than any mitigating change in the registry could ever be, excluding drastic trading impeding measures
2. mitigating changes in the registry could have a negative effects on another important issue with the DES, namely the unit-block fragmentation issue. The unit-block fragmentation issue is currently still under investigation.

This investigation first needs to be completed to determine which measures (in the registry, ITL or DES protocol) would be most effective and efficient when considering both DES problems.

As reported further on under ‘14.2.3 Change to the database or the capacity of National Registry’ implemented Greta software version 4.2 contains changes that improve the capacity of the Netherlands registry. This version is used or soon will be used by the registries of more than ten parties. This will increase the average capacity of registries. Although this has not been investigated in detail this should reduce across all registries the amount of 4003/4010 response code occurrences. This is because the discussed DES limitation is amplified when transactions take longer to finalise and the finalisation time decreases with registries capacity increasing.

#### Ref (c)

In 2008 there were twelve transactions in the Netherlands registry that contained unit blocks involved in an inconsistency identified in the Netherlands registry. These twelve transactions can be traced back to two identical cases with two separate incoming transactions that had not been finalised yet by the registry that initiated the transaction to the Netherlands registry. Once the previously discussed DES change # CR13 is implemented both cases would not occur. This would prevent inconsistencies from occurring in such cases and no unit blocks with inconsistencies would be contained in transaction proposals. Now that the change #CR13 is approved Greta version 5.0, planned for 2010, will contain this change.

#### Ref (d)

A newer Greta version 4.3 is currently in development that will contain several improvements to reconciliation messaging. This version is scheduled to be implemented in the registry in the first quarter of 2010. Additionally a new windows service has been introduced to improve and simplify the logical design of the system. This service is designed to

provide one single framework for the processing of incoming and outgoing messages, in time allowing to concentrate all logic concerning messaging in one part of the system. Hereby creating a robust basis to start improving messaging reliability, efficiency and the capacity of the registry as a result. Up to now improvements of messaging have been focussed on transaction messaging. If needed version 5.0, planned for mid 2010, will contain further improvements to bring these improvements to transaction messaging also to reconciliation messaging.

### 14.1.3 Public Information

#### 14.1.3.1 Recommendation

The Netherlands should enhance the user interface of its registry by the time of its next annual submission by providing the public information referred to in paragraphs 45, 46 and 48 of the annex to decision 13/CMP.1, and report, in the annual submission, on any changes to that public information.

#### 14.1.3.2 Action Taken

#### IAR/2009/NL/1/1 Detailed recommendations P2.4.2.1 & P2.4.2.3

‘Netherlands must either provide its account information in the user interface of its national registry or declare this information confidential as required by 13/CMP.1 Annex II paragraph 45.’

‘Netherlands must either provide the list of legal entities in the user interface of its national registry or declare the information confidential as required by 13/CMP.1 Annex II paragraph 48.’

The information on accounts and authorised legal entities was already provided on the registry website itself as per the requirement. This fact was stated on the website:

‘A list of active accounts and authorised Legal Entities in the Registry is also available for download in comma separated file format (.csv).’

In addition to this information being available on the registry website a link to the CITL was also provided.

To avoid confusion the link to the CITL has been removed. The information on accounts is now only provided through the registry website itself.

#### IAR/2009/NL/1/1 Detailed recommendations P2.4.2.2

‘Netherlands has declared that it won’t host article 6 projects and therefore won’t report its list of projects. In order to be compliant with 13/CMP.1 Annex II paragraph 46, Netherlands must either provide an empty list of article 6 project in the user interface of its national registry or declare this information confidential.’

This recommendation is in contradiction with the document ‘Publicly Available Information; Reporting Guidance for National Registries’ version 4.0 dated 22/09/2009 as released by the UNFCCC Secretariat. This document states on page six as a mandatory requirement 44.9:

‘ If data for a specific area do not exist because the underlying business event has not occurred, a note should be presented indicating the registry did not experience the business event creating the data for the requirement rather than displaying a blank list or report’

**Format:** Explanation (Paragraph Reference).

Example: No Article 6 (Joint Implementation) project is reported as conversion to an ERU under an Article 6 project did not occur in the specified period. (Paragraph 46).

On the registry website, no empty list is added. Instead, the wording has been changed to the suggested wording in the guidance.

## 14.2 Changes to National Registry

Reporting Item	Submission
15/CMP.1 annex II.E paragraph 32. (a) <i>Change of name or contact</i>	There is no change in name or contact information of the registry administrator designated by the Netherlands.
15/CMP.1 annex II.E paragraph 32. (b) <i>Change of cooperation arrangement</i>	There is no change in the cooperation arrangement itself. However the Greta collective of Registries decided to change IT supplier from Siemens to SFW Ltd. Development and support was taken over by SWF as of 9 February 2009.
15/CMP.1 annex II.E paragraph 32. (c) <i>Change to database or the capacity of national registry</i>	In 2009 two registry software version updates have been implemented, namely to the Greta version 4.1 on 23 June and version 4.2 on 10 November. Both upgrades have incorporated changes that increased the capacity of the Registry. The following capacity-improving measures have been implemented compared to the previously used version. Update from version 3.0.84 to version 4.1 Internal database procedures have been improved to increase resource efficiency. Update from version 4.1 to version 4.2 A new windows service has been introduced to improve and simplify the logical design of the system. This service is designed to provide one single framework for the processing of incoming and outgoing messages, in time allowing to concentrate all logic concerning messaging in one part of the system. This creates a robust basis to start improving messaging reliability, efficiency and thus the capacity of the registry as a result.
15/CMP.1 annex II.E paragraph 32. (c) <i>Change to database or the capacity of national registry (continued)</i>	Most checks on incoming messages are now performed asynchronously instead of synchronously, hereby considerably decreasing the time needed to create the synchronous response. This change has eliminated time-out errors in processing incoming messages. The asynchronous processing of incoming messages is now performed in sequence as opposed to in parallel. This change has increased the robustness of message processing and resource efficiency, thus further increasing the capacity of the registry. The functionality allowing initiation of transfers has been improved by using a smarter data integrity algorithm. This change increases robustness of the system when several users are trying to initiate transfers from the same account. Thus further increasing the capacity of the registry for outgoing transfers. Together with the above improvements to the registry system, automated testing was introduced for system testing. This makes it possible to test more and to better performance test the system. Testing has proven that the system is now able to process in and outgoing messages containing 2500 unit blocks without problems on mainstream hardware. Calculations have been made suggesting that the system is able to process messages containing up to 9000 unit blocks. This limit is likely imposed by network delays external to the system. Please consult in annex 6.5 the release notes for details on the changes compared to Greta version 3.0.84 used in 2008. Please note that the Greta version 4.1 included the changes made in version 4.0: - Release notes version 4.0.14 - Release notes version 4.1.16 - Release notes version 4.2.21
15/CMP.1 annex II.E paragraph 32. (d) <i>Change of conformance to technical standards</i>	No change in the registry's conformance to technical standards occurred for the reporting period.
15/CMP.1 annex II.E paragraph 32. (e) <i>Change of discrepancies procedures</i>	No changes were made to the procedures to prevent and/or resolve discrepancies.
15/CMP.1 annex II.E paragraph 32. (f) <i>Change of security</i>	No change of security measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32. (g) <i>Change of list of publicly available information</i>	Publication of public information has been brought in line with the document 'Publicly Available Information; Reporting Guidance for National Registries' version 4.0 dated 22/09/2009 as released by the UNFCCC Secretariat.
15/CMP.1 annex II.E paragraph 32. (h) <i>Change of Internet address</i>	No change of the registry Internet address occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32. (i) <i>Change of data integrity measures</i>	A registry software version update was implemented on 10 November. This Greta version 4.2 implemented the following data integrity improvement: The functionality allowing initiation of transfers has been improved by using a smarter data integrity algorithm making use of data check summing. This change increases robustness of the system when several users are trying to initiate transfers from the same account. It prevents data being changed concurrently that might lead to data integrity related problems. Netherlands have been working to review their disaster recovery plans and disaster recovery test plans with a view to testing disaster recovery shortly after finalising these plans. Due to a busy trading period in December a scheduled test for this month has been postponed to February 2010. It is anticipated to include the revised plans together with the test report will be part of the submission next year.
15/CMP.1 annex II.E paragraph 32. (j) <i>Change of test results</i>	The following test plans and test results for version 4.1 and 4.2 of the Greta registry software have been included in annex 6.5: - Greta test plan and report for version 4.1 - (C)ITL test plan and report for version 4.1 - Certification email from Trasy on behalf of UNFCCC Secretariat and European Commission - Greta test plan and report for version 4.2 - Localisation test plan used for versions 4.1 and 4.2





# Information on minimisation of adverse impacts in accordance with Article 3, paragraph 14

The Netherlands has reported information on minimisation of adverse impacts in its 5th National Communication, submitted to the UNFCCC in December 2009. The relevant information is repeated and where appropriate updated below.

The Kyoto Protocol was adopted in pursuit of the ultimate objective of the Convention, and hence its full implementation by the Netherlands is intended to contribute to preventing dangerous anthropogenic interference with the climate system. Ambitious mitigation goals are necessary to ascertain a future for all countries. In striving to develop policies and measures to reduce greenhouse gas emissions, Parties to the Kyoto Protocol should implement those policies and measures to minimise adverse effects, including the adverse effects of climate change, effects on international trade, and social, environmental and economic impacts on other Parties, especially developing country Parties identified in article 4, paragraphs 8 and 9 of the Convention.

Adverse impacts on developing countries are reduced if global temperature increase is limited to 2 degrees Celsius, if dependence on fossil fuels decreases, and if Annex I Parties are able to develop low-carbon energy systems and reduce fossil fuel consumption.

The European Community actively undertakes a large number of activities to have positive impacts on other countries and their ability to tackle climate change, specifically capacity building and technology transfer activities. During the climate conference in Bali in 2007, it was agreed that adaptation should also be part of a new climate convention. To achieve this, technologies and financing mechanisms need to be developed further and made available to developing countries that are seriously affected by climate change. These include low-lying areas such as Vietnam and Bangladesh, islands that barely rise above sea level (the Maldives and parts of

Comoros), and mountain states that are losing their glaciers (and therefore their sources of water). The Dutch Government, through the Development Minister has adopted 'the polluter pays' as the guiding principle in financing adaptation in developing countries. This means that those who emit high levels of CO<sub>2</sub> pay for the adaptation to climate change of others whose emissions are low but who suffer the consequences. The latter are mostly developing countries.

The discussions and negotiations on climate, and the Netherlands' contribution to them have so far largely had an international character. As already stated, in the 2007–2008 period adaptation became an important element in international climate negotiations. In 2008, a number of countries pledged € 6 billion for an international climate fund for mitigation and adaptation to be managed by the World Bank. As a result of Dutch efforts, this fund is to have a special facility for renewable energy programmes in developing countries. In addition, partly as a result of Dutch efforts, 'avoided deforestation' has been placed on the agenda for the negotiations on a new climate convention. Avoided deforestation means that countries are offered financial compensation for preserving their forests, thereby helping to avoid increases in CO<sub>2</sub> emissions.

For the ultimate impact in developing countries, one significant development is that an increasing number of actors are becoming aware of environmental degradation and the need to adapt to climate change. New forms of cooperation, modified technologies and financing mechanisms are being developed so that developing countries can adapt to the consequences of climate change. Attention to adaptation is also increasing at national level. In 2008, for example, the new Ministry of the Environment in Peru made adaptation one of its main policy priorities. A number of areas in the Peruvian Andes are of great importance for the climatologically condi-

tions throughout the entire Andes region. Peru has, therefore, decided to take on a leadership role at regional level in developing an adaptation policy for the international climate negotiations. The Ministry is working closely with local NGOs, such as the Asociación Especializada para el Desarrollo Sostenible (AEDES), on this. The Dutch organisation both ENDS and other NGOs are following this process closely, so that they can contribute to the policy.

In 2007, Dutch civil society organisations, united in the national HIER climate campaign, took the initiative to organise a national meeting on adaptation for representatives of environmental and development organisations. The latter are becoming increasingly convinced of the need to take account of the effects of climate change in their programmes and projects. An example of adaptation is a programme run by the Dutch Red Cross Climate Centre, which builds the capacity of national associations in developing countries to deal with increasingly frequent extreme weather systems. A start has also been made on developing new forms of cooperation between universities, insurance companies, Directorate General for International Cooperation (DGIS), the Ministry of Housing, Spatial Planning and the Environment, and environmental organisations.

The flexible mechanisms under the Protocol - Emissions Trading, Joint Implementation and the Clean Development Mechanism - are all tools incorporated into the Protocol in order to share efforts in reducing greenhouse gases. This ensures that investment is made where there is optimal greenhouse gas-reducing effects, thus ensuring minimal impact on the world economy. Furthermore, the Netherlands policy aims to achieve a balance in measures oriented to CO<sub>2</sub> reduction and measures oriented to reduction of non-CO<sub>2</sub> greenhouse gases.

Dutch policies and measures on the promotion of renewable energies further contribute to reduction of dependence on fossil fuels, meeting rural electricity needs, and improving air quality. Similarly, Dutch activities on the promotion of energy efficiency and CHP measures can reduce energy costs and contribute to the improvement of air quality.

#### Market imperfections, removing subsidies, etc. (action (a) and (b))

Changes to subsidies under the EC Common Agricultural Policy (CAP) now link payments to environmental, food safety and animal welfare standards, not to agricultural production volume. This encourages responsible agricultural practices.

As a member of the EU, the Dutch electricity market is also liberalised for all users. Also market oriented instruments are implemented as part of the climate change policies, such as the ETS system. In the Netherlands also, a significant part of the N<sub>2</sub>O emissions have been included in the system.

One of the packages of measures in the Netherlands is oriented to investigating, demonstrating and implementing the use of carbon capture and storage systems in the Netherlands. Pilot projects have started and/or are being prepared.

#### Technological development of non-energy use of fossil fuels (action (c))

The technological development of non-energy use of fossil fuels is not a policy or research priority in the Netherlands. Priority is given to improve energy efficiency.

#### Promotion of bio fuels (action (d) and (e))

Expectations are that the worldwide use of biomass in the energy supply will increase considerably in the coming decades. This will be accompanied by the large-scale planting of energy crops. New areas will be opened up for agriculture. Countries and producers will see opportunities for new activities. But there is growing concern that this must not be at the expense of other important values for nature, environment and society. To accommodate these feelings, criteria will be needed that indicate whether biomass has been produced in a responsible way.

At the request of the Government, the project group 'Sustainable production of biomass', under the chairmanship of Prof. Dr. Jacqueline Cramer, from the beginning of 2006 has been bringing together different views on sustainable production. On this basis, the project group has drawn up a framework for the testing of the sustainability of biomass production. This report describes this 'testing framework for sustainable produced biomass' and its elaboration in the form of criteria and indicators. The project group defines the sustainability of large-scale production of biomass on the basis of six relevant themes. These themes are for the greater part linked to the 'Triple P' of sustainable development: People, Planet and Profit, supplemented with specific themes for biomass.

The project group distinguishes six relevant themes:

- *Greenhouse gas emissions:* Calculated over the whole chain, the use of biomass must produce fewer emissions of greenhouse gases net than on average with fossil fuel. For electricity production, the emission reduction must now amount to at least 50-70%, for the application in transportation fuels at least 30%. These percentages must increase further by innovation in the future. The percentages are minimum requirements. Here the basic principle must be that policy instruments should promote a higher percentage above the minimum requirement by differentiating strongly on the basis of the emission reduction of greenhouse gases. The project group thinks it desirable to achieve, in about ten years time, at least 80 to 90% emission reduction in relation to the current fossil reference. This means that in 2010 it will have to be evaluated to what degree the minimum requirement will have to be tightened up in 2011 to attain the objective of 80 to 90% in ten years time. This aim can be achieved when innovative biofuels are applied and a much more efficient cultivation for the production of energy. The development of new acreage for the planting of biomass for energy must not lead in the longer term to the release of large quantities of carbon that had been stored there (in soil or vegetation).
- *Competition with food and other local applications:* The production of biomass for energy must not endanger the food supply and other local applications (such as for medicines or building materials). Criteria for this have not been deter-

mined yet; reporting on changes in land use in the region and in prices for food and land is of great importance here.

- *Biodiversity*: Biomass production must not affect protected or vulnerable biodiversity and will, where possible, have to strengthen biodiversity. Often, local laws and regulations have been grafted on international agreements about biodiversity. Vulnerable areas and areas with a high value for biodiversity must be spared, where possible restoration of biodiversity is desirable.
- *Environment*: In the production and processing of biomass, the quality of soil, surface and ground water and air must be retained or even increased. This makes demands, for example, on the use of fertilisers and pesticides, but it also requires the application of the 'best practices' for instance to prevent erosion or additional emission of harmful substances.
- *Prosperity*: The production of biomass must contribute to local prosperity. Criteria for this have not yet been developed. Reports that fit in with descriptions according to the Global Reporting Initiative can indicate if, for instance, the economic value of the biomass production will directly benefit the local community
- *Social Well-being*: The production of biomass must contribute to the social well-being of the employees and the local population. The production of biomass must at least comply with international principles that have been laid down by the International Labour Organisation, in the UN Universal Declaration of Human Rights and in other treaties. Reports must also bring to light any violations of property rights or corruption.

Strengthening capacity and assisting developing countries (15. CMP.1 para 2. Actions (e) and (f))

The Netherlands promotes the transfer of technology and capacity building through various channels, for example through:

- EU programmes and mechanisms;
- participation in IEA programmes;
- bilateral or multilateral programmes and schemes
- use of flexible mechanisms such as CDM.

These are extensively described in the 5th National Communication.

A recent impact evaluation by the World Bank shows that increased use of electricity leads to improved quality of air in homes (through better cooking facilities), better health provisions (including refrigeration for medicines, good light for operations), and greater availability of information (reading lights, radios, the Internet). This enables the poor to more easily meet their basic needs. The Netherlands has set itself the target of providing 10 million people in developing countries with modern energy by 2015. Between 2004 and 2007, through a variety of programmes funded by the Netherlands, 5.5 million people started to use of modern, sustainable forms of energy, including biogas installations, small-scale hydraulic power stations, solar panels etc. On the basis of the most recent calculations, the total number of people benefiting had reached 6.3 million by the end of 2008.

Multilaterally, the Netherlands cooperates in the Energy Fund for Africa (the World Bank, the IFC and the African Development Bank), the Asia Sustainable Technology and Alternative Energy (ASTAE) programme for Southeast Asia, and the Energy Sector Management Assistance Programme (ESMAP). The results of this cooperation include knowledge on the application of renewable energy, strengthening national organisations for renewable energy and access to renewable energy for the poor. For example, hydraulic power stations have been set up in Zambia, and solar energy programmes in Mongolia. In Southeast Asia, energy loans are provided to projects that use renewable energy.

Bilaterally, the Netherlands participates in a regional programme for the Great Lakes region, in the renovation of existing hydraulic power plants, the transmission of electricity, extending the electricity network and decentralising energy programmes. Methane excavation from Lake Kivu has a prominent place in these activities. In Rwanda, a national energy plan has been developed in which renewable energy has been integrated. In the DRC, a programme has been elaborated to renew the electricity network in Kisangani. In Indonesia, a number of Dutch energy partners, including NL Agency, are supporting a nationally operating energy programme, together with the World Bank and GTZ. As a result, thousands of households are now connected to the electricity network and solar panels and small-scale hydraulic power stations have been installed. Cleaner energy is now also available in the form of biogas and vegetable oils.

A large number of new initiatives have been developed in cooperation with civil society organisations and the private sector. The Netherlands Development Organisation (SNV) conducts biogas programmes in Asia and a biogas programme is being set up, together with SNV and Hivos, in six African countries. The Global Village Energy Partnership strengthens local small-scale private enterprises in developing countries. The Free Energy Foundation works to increase the use of solar power in Africa. Partnerships with Nuon, Philips and the FMO's Access to Energy fund collaborate with local companies that supply consumers with electricity on the basis of renewable energy. This helps to make low-energy products available in local markets. Here, too, the results contribute to poverty reduction and economic development, help reduce the growth of CO<sub>2</sub> emissions and strengthen the position of women.



## Other information

No other information



# Annexes





# Annex 1

## Key sources

### A1.1 Introduction

As explained in the Good Practice Guidance (IPCC, 2001), a key source category is prioritised within the national inventory system because its estimate has a significant influence on a country's total inventory of direct greenhouse gases in terms of the absolute level of emissions, the trend in emissions, or both.

For the identification of key sources in the Netherlands inventory, we allocated the national emissions according to the Intergovernmental Panel on Climate Change (IPCC) potential key source list, as presented in Table 7.1 in Chapter 7 of the Good Practice Guidance. As suggested in this table, the carbon dioxide (CO<sub>2</sub>) emissions from stationary combustion (1A1, 1A2 and 1A4) are aggregated by fuel type. CO<sub>2</sub>, methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions from “Mobile combustion: road vehicles” (1A3) are assessed separately. The CH<sub>4</sub> and N<sub>2</sub>O emissions from aircrafts and ships are relatively small (about 1–2 Gg CO<sub>2</sub> equivalents). Other mobile sources are not assessed separately by gas. “Fugitive emissions from oil and gas operations” (1B) is an important source of greenhouse gas emissions in the Netherlands. The most important gas/source combinations in this category are separately assessed. Emissions in other IPCC sectors are disaggregated, as suggested by IPCC.

The IPCC Tier 1 method consists of ranking the list of source category/gas combinations according to their contribution to the national total annual emissions and to the national total trend. The darker green areas at the top of the tables in this Annex are the largest sources, of which the total adds up to 95% of the national total (excluding LULUCF): 31 sources for annual level assessment (emissions in 2008) and 30 sources for the trend assessment out of a total of 70 sources. Both lists can be combined to obtain an overview of sources that meet any of these two criteria.

The IPCC Tier 2 method for identification of key sources requires the incorporation of the uncertainty in each of these sources before ordering the list of shares. This has been carried out using the uncertainty estimates presented in Annex 7 (for details on the Tier 1 uncertainty analysis see Olivier et al., 2009). The results of the Tier 1 and Tier 2 level

and trend assessments are summarised in Table A1.1 and show a total of 38 key sources (excluding LULUCF).

As expected, the Tier 2 level and trend assessment increases the importance of very uncertain sources. It can be concluded that in using the results of a Tier 2 key source assessment, 1 more source is added to the list of 35 Tier 1 level and trend key sources (excluding LULUCF) as were identified in the last submission:

- CH<sub>4</sub> emissions from “Manure management, poultry” (Tier 2 trend)

The share of these sources in the national annual total becomes more important when taking their uncertainty (50%–100%) into account (Table A1.4). We then include the most important Land Use, Land Use Change and Forestry (LULUCF) emission sinks and sources in the Tier 1 and Tier 2 key source calculations to identify the key sources in IPCC Sector 5. This results in 4 additional key sources, giving an overall total of 42 key sources (including LULUCF); see also Table A1.2. In this report, the key source assessment is based on emission figures from Common Reporting Format (CRF) 2010 version 1.1, submitted to the European Union (EU) in January 2010.

Key source list identified by the Tier 1 and 2 level and trend assessments (based on CRF tables 2010 version 1.1. Level assessment for 2008 emissions (excluding LULUCF sources)

Table A1.1

Category	Gas	Category name	Key source?	Tier 1 Level	Tier 1 Trend	Tier 2 Level	Tier 2 Trend
<i>ENERGY</i>							
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: liquids	Key(L1,T1)	1	1	0	0
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: solids	Key(L,T1)	1	1	1	0
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: gases	Key(L,T)	1	1	1	1
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: waste incineration	Key(L,T)	1	1	1	1
1A1b	CO <sub>2</sub>	Stationary combustion: Petroleum Refining: liquids	Key(L,T1)	1	1	1	0
1A1b	CO <sub>2</sub>	Stationary combustion: Petroleum Refining: gases	Key(L1,T1)	1	1	0	0
1A1c	CO <sub>2</sub>	Stationary combustion: Manuf. of Solid Fuels and Other En. Ind.: liquids	Non key	0	0	0	0
1A1c	CO <sub>2</sub>	Stationary combustion: Manuf. of Solid Fuels and Other En. Ind.: gases	Key(L,T1)	1	1	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, liquids	Key(L,T1)	1	1	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, solids	Key(L,)	1	0	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, gases	Key(L,T)	1	1	1	1
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: gasoline	Key(L,T1)	1	1	1	0
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: diesel oil	Key(L,T)	1	1	1	1
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: LPG	Key(L1,T)	1	1	0	1
1A3	CO <sub>2</sub>	Mobile combustion: water-borne navigation	Key(L1,)	1	0	0	0
1A3	CO <sub>2</sub>	Mobile combustion: aircraft	Non key	0	0	0	0
1A3	CO <sub>2</sub>	Mobile combustion: other (railways)	Non key	0	0	0	0
1A3	CH <sub>4</sub>	Mobile combustion: other (non-road)	Non key	0	0	0	0
1A3	N <sub>2</sub> O	Mobile combustion: other (non-road)	Non key	0	0	0	0
1A3	CH <sub>4</sub>	Mobile combustion: road vehicles	Non key	0	0	0	0
1A3	N <sub>2</sub> O	Mobile combustion: road vehicles	Non key	0	0	0	0
1A4	CO <sub>2</sub>	Stationary combustion: Other Sectors, solids	Non key	0	0	0	0
1A4a	CO <sub>2</sub>	Stationary combustion: Other Sectors: Commercial/Institutional, gases	Key(L,T)	1	1	1	1
1A4b	CO <sub>2</sub>	Stationary combustion: Other Sectors, Residential, gases	Key(L,T1)	1	1	1	0
1A4c	CO <sub>2</sub>	Stationary combustion: Other Sectors, Agriculture/Forestry/Fisheries, gases	Key(L,T1)	1	1	1	0
1A4c	CO <sub>2</sub>	Stationary combustion: Other Sectors, Agriculture/Forestry/Fisheries, liquids	Key(L,T1)	1	1	1	0
1A4	CO <sub>2</sub>	Stationary combustion: Other Sectors, liquids excl. From 1A4c	Key(,T)	0	1	0	1
1A5	CO <sub>2</sub>	Military use of fuels (1A5 Other)	Non key	0	0	0	0
1A	CH <sub>4</sub>	Emissions from stationary combustion: non-CO <sub>2</sub>	Key(L,T)	1	1	1	1
1A	N <sub>2</sub> O	Emissions from stationary combustion: non-CO <sub>2</sub>	Non key	0	0	0	0
1B1	CH <sub>4</sub>	Coal mining					
1B1b	CO <sub>2</sub>	Coke production	Non key	0	0	0	0
1B2	CO <sub>2</sub>	Fugitive emissions from venting/flaring: CO <sub>2</sub>	Key(,T)	0	1	0	1
1B2	CH <sub>4</sub>	Fugitive emissions venting/flaring	Key(,T)	0	1	0	1
1B2	CH <sub>4</sub>	Fugitive emissions from oil and gas: gas distribution	Non key	0	0	0	0
1B2	CH <sub>4</sub>	Fugitive emissions from oil and gas operations: other	Non key	0	0	0	0

Category	Gas	Category name	Key source?	Tier 1 Level	Tier 1 Trend	Tier 2 Level	Tier 2 Trend
<i>INDUSTRIAL PROCESSES</i>							
2A1	CO <sub>2</sub>	Cement production	Non key	0	0	0	0
2A3	CO <sub>2</sub>	Limestone and dolomite use	Non key	0	0	0	0
2A7	CO <sub>2</sub>	Other minerals	Non key	0	0	0	0
2B1	CO <sub>2</sub>	Ammonia production	Key(L1,)	1	0	0	0
2B2	N <sub>2</sub> O	Nitric acid production	Key(T)	0	1	0	1
2B5	N <sub>2</sub> O	Caprolactam production	Non key	0	0	0	0
2B5	CO <sub>2</sub>	Other chemical product manufacture	Key(L,)	1	0	1	0
2C1	CO <sub>2</sub>	Iron and steel production (carbon inputs)	Key(L1,T1)	1	1	0	0
2C3	CO <sub>2</sub>	CO <sub>2</sub> from aluminium production	Non key	0	0	0	0
2C3	PFC	PFC from aluminium production	Key(T)	0	1	0	1
2F	SF <sub>6</sub>	SF <sub>6</sub> emissions from SF6 use	Non key	0	0	0	0
2F	HFC	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	Key(L,T)	1	1	1	1
2E	HFC	HFC-23 emissions from HCFC-22 manufacture	Key(T)	0	1	0	1
2E	HFC	HFC by-product emissions from HFC manufacture	Non key	0	0	0	0
2F	PFC	PFC emissions from PFC use	Non key	0	0	0	0
2G	CO <sub>2</sub>	Other industrial: CO <sub>2</sub>	Non key	0	0	0	0
2G	CH <sub>4</sub>	Other industrial: CH <sub>4</sub>	Non key	0	0	0	0
2G	N <sub>2</sub> O	Other industrial: N <sub>2</sub> O	Non key	0	0	0	0
<i>SOLVENTS AND OTHER PRODUCT USE</i>							
3	CO <sub>2</sub>	Indirect CO2 from solvents/product use	Non key	0	0	0	0
3	CH <sub>4</sub>	Solvents and other product use	IE in 2G				
<i>AGRICULTURAL SECTOR</i>							
4A1	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	Key(L,T)	1	1	1	1
4A8	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	Non key	0	0	0	0
4A	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	Non key	0	0	0	0
4B	N <sub>2</sub> O	Emissions from manure management	Key(L,T2)	1	0	1	1
4B1	CH <sub>4</sub>	Emissions from manure management : cattle	Key(L,)	1	0	1	0
4B8	CH <sub>4</sub>	Emissions from manure management : swine	Key(L,)	1	0	1	0
4B9	CH <sub>4</sub>	Emissions from manure management: poultry	Key(T2)	0	0	0	1
4B	CH <sub>4</sub>	Emissions from manure management : other	Non key	0	0	0	0
4C	CH <sub>4</sub>	Rice cultivation					
4D1	N <sub>2</sub> O	Direct N <sub>2</sub> O emissions from agricultural soils	Key(L,T)	1	1	1	1
4D3	N <sub>2</sub> O	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	Key(L,T)	1	1	1	1
4D2	N <sub>2</sub> O	Animal production on agricultural soils	Key(L,T)	1	1	1	1
<i>WASTE</i>							
6A1	CH <sub>4</sub>	CH <sub>4</sub> emissions from solid waste disposal sites	Key(L,T)	1	1	1	1
6B	CH <sub>4</sub>	Emissions from wastewater handling	Non key	0	0	0	0
6B	N <sub>2</sub> O	Emissions from wastewater handling	Non key	0	0	0	0
6C	all	Emissions from waste incineration					
6D	CH <sub>4</sub>	OTHER CH <sub>4</sub>	Non key	0	0	0	0
3, 6D	N <sub>2</sub> O	OTHER N <sub>2</sub> O	Non key	0	0	0	0
<i>KEY SOURCE CATEGORIES (EXCL. LULUCF)</i>			38	31	30	25	21

Category	Gas	Category name	Key source?	Tier 1 Level	Tier 1 Trend	Tier 2 Level	Tier 2 Trend
<i>ENERGY</i>							
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: liquids	Key(L1,T1)	1	1	0	0
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: solids	Key(L,T1)	1	1	1	0
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: gases	Key(L,T)	1	1	1	1
1A1a	CO <sub>2</sub>	Stationary combustion: Public Electricity and Heat Production: waste incineration	Key(L,T)	1	1	1	1
1A1b	CO <sub>2</sub>	Stationary combustion: Petroleum Refining: liquids	Key(L,T1)	1	1	1	0
1A1b	CO <sub>2</sub>	Stationary combustion: Petroleum Refining: gases	Key(L1,T1)	1	1	0	0
1A1c	CO <sub>2</sub>	Stationary combustion: Manuf. of Solid Fuels and Other En. Ind.: liquids	Non key	0	0	0	0
1A1c	CO <sub>2</sub>	Stationary combustion: Manuf. of Solid Fuels and Other En. Ind.: gases	Key(L,T1)	1	1	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, liquids	Key(L,T1)	1	1	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, solids	Key(L,)	1	0	1	0
1A2	CO <sub>2</sub>	Emissions from stationary combustion: Manufacturing Industries and Construction, gases	Key(L,T)	1	1	1	1
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: gasoline	Key(L,T1)	1	1	1	0
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: diesel oil	Key(L,T)	1	1	1	1
1A3b	CO <sub>2</sub>	Mobile combustion: road vehicles: LPG	Key(L1,T)	1	1	0	1
1A3	CO <sub>2</sub>	Mobile combustion: water-borne navigation	Key(L1,)	1	0	0	0
1A3	CO <sub>2</sub>	Mobile combustion: aircraft	Non key	0	0	0	0
1A3	CO <sub>2</sub>	Mobile combustion: other (railways)	Non key	0	0	0	0
1A3	CH <sub>4</sub>	Mobile combustion: other (non-road)	Non key	0	0	0	0
1A3	N <sub>2</sub> O	Mobile combustion: other (non-road)	Non key	0	0	0	0
1A3	CH <sub>4</sub>	Mobile combustion: road vehicles	Non key	0	0	0	0
1A3	N <sub>2</sub> O	Mobile combustion: road vehicles	Non key	0	0	0	0
1A4	CO <sub>2</sub>	Stationary combustion: Other Sectors, solids	Non key	0	0	0	0
1A4a	CO <sub>2</sub>	Stationary combustion: Other Sectors: Commercial/Institutional, gases	Key(L,T)	1	1	1	1
1A4b	CO <sub>2</sub>	Stationary combustion: Other Sectors, Residential, gases	Key(L,T1)	1	1	1	0
1A4c	CO <sub>2</sub>	Stationary combustion: Other Sectors, Agriculture/Forestry/Fisheries, gases	Key(L,T1)	1	1	1	0
1A4c	CO <sub>2</sub>	Stationary combustion: Other Sectors, Agriculture/Forestry/Fisheries, liquids	Key(L,T1)	1	1	1	0
1A4	CO <sub>2</sub>	Stationary combustion: Other Sectors, liquids excl. From 1A4c	Key(,T)	0	1	0	1
1A5	CO <sub>2</sub>	Military use of fuels (1A5 Other)	Non key	0	0	0	0
1A	CH <sub>4</sub>	Emissions from stationary combustion: non-CO <sub>2</sub>	Key(L,T)	1	1	1	1
1A	N <sub>2</sub> O	Emissions from stationary combustion: non-CO <sub>2</sub>	Non key	0	0	0	0
1B1	CH <sub>4</sub>	Coal mining					
1B1b	CO <sub>2</sub>	Coke production	Non key	0	0	0	0
1B2	CO <sub>2</sub>	Fugitive emissions from venting/flaring: CO <sub>2</sub>	Key(,T)	0	1	0	1
1B2	CH <sub>4</sub>	Fugitive emissions venting/flaring	Key(,T)	0	1	0	1
1B2	CH <sub>4</sub>	Fugitive emissions from oil and gas: gas distribution	Non key	0	0	0	0
1B2	CH <sub>4</sub>	Fugitive emissions from oil and gas operations: other	Non key	0	0	0	0

Category	Gas	Category name	Key source?	Tier 1 Level	Tier 1 Trend	Tier 2 Level	Tier 2 Trend
<b>INDUSTRIAL PROCESSES</b>							
2A1	CO <sub>2</sub>	Cement production	Non key	0	0	0	0
2A3	CO <sub>2</sub>	Limestone and dolomite use	Non key	0	0	0	0
2A7	CO <sub>2</sub>	Other minerals	Non key	0	0	0	0
2B1	CO <sub>2</sub>	Ammonia production	Key(L1,)	1	0	0	0
2B2	N <sub>2</sub> O	Nitric acid production	Key(,T)	0	1	0	1
2B5	N <sub>2</sub> O	Caprolactam production	Non key	0	0	0	0
2B5	CO <sub>2</sub>	Other chemical product manufacture	Key(L,)	1	0	1	0
2C1	CO <sub>2</sub>	Iron and steel production (carbon inputs)	Key(L1,T1)	1	1	0	0
2C3	CO <sub>2</sub>	CO <sub>2</sub> from aluminium production	Non key	0	0	0	0
2C3	PFC	PFC from aluminium production	Key(,T)	0	1	0	1
2F	SF <sub>6</sub>	SF <sub>6</sub> emissions from SF <sub>6</sub> use	Non key	0	0	0	0
2F	HFC	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	Key(L,T)	1	1	1	1
2E	HFC	HFC-23 emissions from HCFC-22 manufacture	Key(,T)	0	1	0	1
2E	HFC	HFC by-product emissions from HFC manufacture	Non key	0	0	0	0
2F	PFC	PFC emissions from PFC use	Non key	0	0	0	0
2G	CO <sub>2</sub>	Other industrial: CO <sub>2</sub>	Non key	0	0	0	0
2G	CH <sub>4</sub>	Other industrial: CH <sub>4</sub>	Non key	0	0	0	0
2G	N <sub>2</sub> O	Other industrial: N <sub>2</sub> O	Non key	0	0	0	0
<b>SOLVENTS AND OTHER PRODUCT USE</b>							
3	CO <sub>2</sub>	Indirect CO <sub>2</sub> from solvents/product use	Non key	0	0	0	0
3	CH <sub>4</sub>	Solvents and other product use	IE in 2G				
<b>AGRICULTURAL SECTOR</b>							
4A1	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	Key(L,T)	1	1	1	1
4A8	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	Non key	0	0	0	0
4A	CH <sub>4</sub>	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	Non key	0	0	0	0
4B	N <sub>2</sub> O	Emissions from manure management	Key(L,T2)	1	0	1	1
4B1	CH <sub>4</sub>	Emissions from manure management : cattle	Key(L,)	1	0	1	0
4B8	CH <sub>4</sub>	Emissions from manure management : swine	Key(L,)	1	0	1	0
4B9	CH <sub>4</sub>	Emissions from manure management: poultry	Key(,T2)	0	0	0	1
4B	CH <sub>4</sub>	Emissions from manure management : other	Non key	0	0	0	0
4C	CH <sub>4</sub>	Rice cultivation					
4D1	N <sub>2</sub> O	Direct N <sub>2</sub> O emissions from agricultural soils	Key(L,T)	1	1	1	1
4D3	N <sub>2</sub> O	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	Key(L,T)	1	1	1	1
4D2	N <sub>2</sub> O	Animal production on agricultural soils	Key(L,T)	1	1	1	1
<b>LAND USE, LAND USE CHANGE AND FORESTRY</b>							
5A1	CO <sub>2</sub>	5A1. Forest Land remaining Forest Land	Key(L,T)	1	1	1	1
5A2	CO <sub>2</sub>	5A2. Land converted to Forest Land	Key(L,T)	1	1	1	1
5B2	CO <sub>2</sub>	5B2. Land converted to Cropland	Non key	0	0	0	0
5C1	CO <sub>2</sub>	5C1. Grassland remaining Grassland	Key(L,)	1	0	1	0
5C2	CO <sub>2</sub>	5C2. Land converted to Grassland	Key(L2,T2)	0	0	1	1
5D2	CO <sub>2</sub>	5D2. Land converted to Wetlands	Non key	0	0	0	0
5E2	CO <sub>2</sub>	5E2. Land converted to Settlements	Non key	0	0	0	0
5F2	CO <sub>2</sub>	5F2. Land converted to Other Land	Non key	0	0	0	0
5G	CO <sub>2</sub>	5G. Other (liming of soils)	Non key	0	0	0	0
<b>WASTE</b>							
6A1	CH <sub>4</sub>	CH <sub>4</sub> emissions from solid waste disposal sites	Key(L,T)	1	1	1	1
6B	CH <sub>4</sub>	Emissions from wastewater handling	Non key	0	0	0	0
6B	N <sub>2</sub> O	Emissions from wastewater handling	Non key	0	0	0	0
6C	all	Emissions from waste incineration					
6D	CH <sub>4</sub>	OTHER CH <sub>4</sub>	Non key	0	0	0	0
3, 6D	N <sub>2</sub> O	OTHER N <sub>2</sub> O	Non key	0	0	0	0
<b>TOTAL KEY SOURCE CATEGORIES (INCL. LULUCF)</b>			42	35	34	28	24

## A1.2 Changes in key sources compared to previous submission

Due to the use of emission data for 2008 in the key source analysis, the following changes have taken place compared to the previous NIR:

- CH<sub>4</sub> emissions from “Manure management, poultry” (Tier 2 trend)

## A1.3 Tier 1 key source and uncertainty assessment

In Tables A1.3 and A1.4, the source ranking is done according to the contribution to the 2008 annual emissions total and to the base year to 2008 trend respectively. This resulted in 31 level key sources and 30 trend key sources (indicated in the blue lines at the top, excluding LULUCF). Inclusion of LULUCF sources in the analysis adds 4 additional Tier 1 level and trend key sources (see Table A1.2).

Source ranking using IPCC Tier 1 level assessment 2008 excluding LULUCF (amounts in Gg CO<sub>2</sub> eq)

Table A1.3

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	12%	12%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	12%	24%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	10%	34%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	9%	42%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	7%	49%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	6%	55%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	61%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9059	4%	65%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	4%	69%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	4%	73%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	3%	76%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	2%	78%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	2%	81%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	2%	83%
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3112	2%	84%
2B1	Ammonia production	CO <sub>2</sub>	2850	1%	86%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	2723	1%	87%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	88%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	2242	1%	89%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	90%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	91%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1500	1%	92%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	1%	92%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	1%	93%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	1%	94%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	0%	94%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	95%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	837	0%	95%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	95%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	96%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	0%	96%
2B2	Nitric acid production	N <sub>2</sub> O	558	0%	96%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	96%
2A7	Other minerals	CO <sub>2</sub>	471	0%	97%
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	97%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	97%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	97%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	97%
2A1	Cement production	CO <sub>2</sub>	399	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	98%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	0%	98%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	98%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	363	0%	98%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	99%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	99%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	99%



IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	269	0%	99%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	99%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	99%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	0%	99%
2F	PFC emissions from PFC use	PFC	180	0%	99%
6B	Emissions from wastewater handling	CH <sub>4</sub>	180	0%	99%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	100%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	100%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	100%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	0%	100%
2C3	PFC from aluminium production	PFC	72	0%	100%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	100%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	100%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	100%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	12%	12%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	11%	23%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	10%	33%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	8%	41%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	6%	47%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	6%	53%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	58%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9059	4%	63%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	4%	67%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	3%	70%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	3%	73%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	2%	75%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	2%	77%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	2%	80%
5C1	5C1. Grassland remaining Grassland	CO <sub>2</sub>	4246	2%	82%
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3112	1%	83%
2B1	Ammonia production	CO <sub>2</sub>	2850	1%	84%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	2723	1%	86%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	87%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	2242	1%	88%
5A1	5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	2208	1%	89%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	90%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	90%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1623	1%	91%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	1%	92%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	1%	93%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	1%	93%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	0%	94%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	94%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	837	0%	94%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	95%
5A2	5A2. Land converted to Forest Land	CO <sub>2</sub>	639	0%	95%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	95%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	0%	96%
2B2	Nitric acid production	N <sub>2</sub> O	558	0%	96%
5C2	5C2. Land converted to Grassland	CO <sub>2</sub>	550	0%	96%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	96%
2A7	Other minerals	CO <sub>2</sub>	471	0%	97%
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	97%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	97%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	97%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	97%
2A1	Cement production	CO <sub>2</sub>	399	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	98%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	0%	98%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	98%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	363	0%	98%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	98%
5E2	5E2. Land converted to Settlements	CO <sub>2</sub>	296	0%	99%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	99%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	99%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	269	0%	99%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	99%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	99%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	0%	99%
6B	Emissions from wastewater handling	CH <sub>4</sub>	198	0%	99%
2F	PFC emissions from PFC use	PFC	180	0%	99%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	99%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	100%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	100%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	0%	100%
2C3	PFC from aluminium production	PFC	72	0%	100%
5G	5G. Other (liming of soils)	CO <sub>2</sub>	71	0%	100%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	100%
5D2	5D2. Land converted to Wetlands	CO <sub>2</sub>	56	0%	100%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	100%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	100%
5B2	5B2. Land converted to Cropland	CO <sub>2</sub>	48	0%	100%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	100%
5F2	5F2. Land converted to Other Land	CO <sub>2</sub>	25	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	15%	15%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	13%	28%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	9%	38%
2B2	Nitric acid production	N <sub>2</sub> O	558	8%	45%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	7%	53%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	6%	59%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	64%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	3%	67%
2C3	PFC from aluminium production	PFC	72	2%	70%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	3112	2%	72%
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	2723	2%	74%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	2242	2%	77%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	2%	79%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1623	2%	81%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	2%	82%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	363	1%	84%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	85%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	1%	86%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	1%	88%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	1%	89%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	1%	90%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	9059	1%	90%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	837	1%	91%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	1%	92%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	1%	93%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	94%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	94%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	1%	95%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	0%	95%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	0%	96%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	96%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	96%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	97%
2A7	Other minerals	CO <sub>2</sub>	471	0%	97%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	0%	97%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	98%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	98%
2B1	Ammonia production	CO <sub>2</sub>	2850	0%	98%
2F	PFC emissions from PFC use	PFC	180	0%	98%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	99%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	99%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	99%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	99%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	99%
6B	Emissions from wastewater handling	CH <sub>4</sub>	198	0%	99%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	99%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	99%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	100%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	100%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	100%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	100%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	100%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	0%	100%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	269	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	100%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	100%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	100%
2A1	Cement production	CO <sub>2</sub>	399	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	100%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	15%	15%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	13%	27%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	9%	37%
2B2	Nitric acid production	N <sub>2</sub> O	558	8%	44%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	7%	52%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	6%	58%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	63%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	3%	66%
2C3	PFC from aluminium production	PFC	72	2%	68%
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3112	2%	71%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	2723	2%	73%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	2242	2%	75%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	2%	77%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1623	2%	79%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	1%	81%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	363	1%	82%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	84%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	1%	85%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	1%	86%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	1%	87%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	1%	88%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9059	1%	89%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	837	1%	90%
5A2	5A2. Land converted to Forest Land	CO <sub>2</sub>	639	1%	91%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	1%	91%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	1%	92%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	93%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	93%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	0%	94%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	0%	94%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	0%	95%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	95%
5A1	5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	2208	0%	95%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	96%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	0%	96%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	96%
2A7	Other minerals	CO <sub>2</sub>	471	0%	97%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	97%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	97%
5C2	5C2. Land converted to Grassland	CO <sub>2</sub>	550	0%	97%
2B1	Ammonia production	CO <sub>2</sub>	2850	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	98%
2F	PFC emissions from PFC use	PFC	180	0%	98%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	98%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	98%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	98%
5C1	5C1. Grassland remaining Grassland	CO <sub>2</sub>	4246	0%	99%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	99%
5G	5G. Other (liming of soils)	CO <sub>2</sub>	71	0%	99%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	99%
5E2	5E2. Land converted to Settlements	CO <sub>2</sub>	296	0%	99%
6B	Emissions from wastewater handling	CH <sub>4</sub>	198	0%	99%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	99%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	99%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	100%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	100%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	100%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Cum. Share
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	0%	100%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	100%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	269	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	100%
5D2	5D2. Land converted to Wetlands	CO <sub>2</sub>	56	0%	100%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	100%
5B2	5B2. Land converted to Cropland	CO <sub>2</sub>	48	0%	100%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	0%	100%
5F2	5F2. Land converted to Other Land	CO <sub>2</sub>	25	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	100%
2A1	Cement production	CO <sub>2</sub>	399	0%	100%
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	100%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	100%



## A1.4 Tier 2 key source assessment

Using the uncertainty estimate for each key source as a weighting factor (see Annex 7), the key source assessment was performed again. This is called the Tier 2 key source assessment. The results of this assessment are presented in Tables A1.5 and A1.6 for the contribution to the 2008 annual

emissions total and to the trend respectively. Comparison with the Tier 1 assessment presented in Tables A1.3 and A1.4 show less level and trend key sources (25 and 21 respectively instead of 31 and 30). Inclusion of LULUCF sources in the analysis adds 3 additional Tier 2 level and trend key sources (see Table A1.2).

Source ranking using IPCC Tier 2 level assessment 2008 excluding LULUCF (in Gg CO<sub>2</sub> eq).

Table A1.5

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Uncertainty estimate	Level * Uncertainty	Share L*U	Cum. Share L*U
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3112	2%	206%	3%	20%	20%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	2%	61%	1%	9%	30%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	20%	1%	7%	37%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	2%	34%	1%	5%	42%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	1%	100%	1%	5%	47%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9059	4%	14%	1%	4%	51%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	1%	100%	1%	4%	55%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	10%	5%	1%	3%	58%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	3%	16%	0%	3%	61%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	100%	0%	3%	64%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	9%	5%	0%	3%	67%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1623	1%	51%	0%	3%	69%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	50%	0%	3%	72%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	12%	3%	0%	3%	74%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	4%	10%	0%	2%	77%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	0%	100%	0%	2%	79%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	2%	10%	0%	2%	80%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	20%	0%	2%	82%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	4%	5%	0%	1%	83%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	71%	0%	1%	85%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	21%	0%	1%	86%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	7%	2%	0%	1%	87%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	12%	1%	0%	1%	88%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	6%	2%	0%	1%	89%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	2242	1%	11%	0%	1%	90%
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	54%	0%	1%	90%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	50%	0%	1%	91%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	50%	0%	1%	92%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	50%	0%	1%	92%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	50%	0%	1%	93%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	51%	0%	0%	93%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	28%	0%	0%	94%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	20%	0%	0%	94%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	56%	0%	0%	95%
2B2	Nitric acid production	N <sub>2</sub> O	558	0%	22%	0%	0%	95%
2A7	Other minerals	CO <sub>2</sub>	471	0%	25%	0%	0%	95%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	0%	10%	0%	0%	96%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	0%	25%	0%	0%	96%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	30%	0%	0%	96%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Uncer- tainty estimate	Level * Uncer- tainty	Share L*U	Cum. Share L*U
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	837	0%	10%	0%	0%	96%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	54%	0%	0%	97%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	20%	0%	0%	97%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	1%	6%	0%	0%	97%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	363	0%	20%	0%	0%	97%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	269	0%	25%	0%	0%	98%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	25%	0%	0%	98%
2B1	Ammonia production	CO <sub>2</sub>	2850	1%	2%	0%	0%	98%
6B	Emissions from wastewater handling	CH <sub>4</sub>	198	0%	32%	0%	0%	98%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	54%	0%	0%	99%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	100%	0%	0%	99%
2F	PFC emissions from PFC use	PFC	180	0%	25%	0%	0%	99%
2A1	Cement production	CO <sub>2</sub>	399	0%	11%	0%	0%	99%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	0%	50%	0%	0%	99%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	27%	0%	0%	99%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	2723	1%	1%	0%	0%	99%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	0%	14%	0%	0%	99%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	50%	0%	0%	99%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	60%	0%	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	21%	0%	0%	100%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	5%	0%	0%	100%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	32%	0%	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	50%	0%	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%	0%	0%	100%
2C3	PFC from aluminium production	PFC	72	0%	20%	0%	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	71%	0%	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	5%	0%	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	22%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	112%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	112%	0%	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	20%	0%	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Uncertainty estimate	Level * Uncertainty	Share L*U	Cum. Share L*U
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	3112	1%	206%	2.98%	18%	18%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4883	2%	61%	1.38%	8%	26%
5C1	5C1. Grassland remaining Grassland	CO <sub>2</sub>	4246	2%	56%	1.10%	7%	32%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	10928	5%	20%	1.02%	6%	38%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	4896	2%	34%	0.76%	5%	43%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1494	1%	100%	0.70%	4%	47%
5A1	5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	2208	1%	67%	0.68%	4%	51%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9059	4%	14%	0.60%	4%	55%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1118	1%	100%	0.52%	3%	58%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	20841	10%	5%	0.49%	3%	61%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5782	3%	16%	0.43%	3%	63%
4B	Emissions from manure management	N <sub>2</sub> O	906	0%	100%	0.42%	3%	66%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	17619	8%	5%	0.42%	2%	68%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	1623	1%	51%	0.39%	2%	71%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	1624	1%	50%	0.38%	2%	73%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25365	12%	3%	0.37%	2%	75%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7503	3%	10%	0.35%	2%	77%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	636	0%	100%	0.30%	2%	79%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	4681	2%	10%	0.22%	1%	80%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2350	1%	20%	0.22%	1%	81%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	9001	4%	5%	0.21%	1%	83%
2B5	Other chemical product manufacture	CO <sub>2</sub>	649	0%	71%	0.21%	1%	84%
5A2	5A2. Land converted to Forest Land	CO <sub>2</sub>	639	0%	63%	0.19%	1%	85%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1940	1%	21%	0.19%	1%	86%
5C2	5C2. Land converted to Grassland	CO <sub>2</sub>	550	0%	66%	0.17%	1%	87%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	13817	6%	2%	0.14%	1%	88%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	23927	11%	1%	0.12%	1%	89%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12886	6%	2%	0.12%	1%	90%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	22,2	1%	11%	0.12%	1%	90%
6B	Emissions from wastewater handling	N <sub>2</sub> O	459	0%	54%	0.11%	1%	91%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	432	0%	50%	0.10%	1%	91%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	430	0%	50%	0.10%	1%	92%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	379	0%	50%	0.09%	1%	93%
5E2	5E2. Land converted to Settlements	CO <sub>2</sub>	296	0%	56%	0.08%	0%	93%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	327	0%	50%	0.08%	0%	94%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	293	0%	51%	0.07%	0%	94%
2B5	Caprolactam production	N <sub>2</sub> O	481	0%	28%	0.06%	0%	94%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	637	0%	20%	0.06%	0%	95%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	224	0%	56%	0.06%	0%	95%
2B2	Nitric acid production	N <sub>2</sub> O	558	0%	22%	0.06%	0%	95%
2A7	Other minerals	CO <sub>2</sub>	471	0%	25%	0.06%	0%	96%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	1010	0%	10%	0.05%	0%	96%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	390	0%	25%	0.05%	0%	96%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	296	0%	30%	0.04%	0%	96%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	837	0%	10%	0.04%	0%	97%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	155	0%	54%	0.04%	0%	97%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	392	0%	20%	0.04%	0%	97%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	1349	1%	6%	0.04%	0%	97%

IPCC	Category	Gas	CO <sub>2</sub> eq last year	Share	Uncer- tainty estimate	Level * Uncer- tainty	Share L*U	Cum. Share L*U
1A4	Stationary combustion : Other Sec- tors, liquids excl. From 1A4c	CO <sub>2</sub>	363	0%	20%	0.03%	0%	98%
1B2	Fugitive emissions from oil and gas ope- rations: gas distribution	CH <sub>4</sub>	269	0%	25%	0.03%	0%	98%
2A3	Limestone and dolomite use	CO <sub>2</sub>	261	0%	25%	0.03%	0%	98%
2B1	Ammonia production	CO <sub>2</sub>	2850	1%	2%	0.03%	0%	98%
6B	Emissions from wastewater handling	CH <sub>4</sub>	198	0%	32%	0.03%	0%	98%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	117	0%	54%	0.03%	0%	98%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	52	0%	100%	0.02%	0%	99%
2F	PFC emissions from PFC use	PFC	180	0%	25%	0.02%	0%	99%
2A1	Cement production	CO <sub>2</sub>	399	0%	11%	0.02%	0%	99%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	72	0%	50%	0.02%	0%	99%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	129	0%	27%	0.02%	0%	99%
5D2	5D2. Land converted to Wetlands	CO <sub>2</sub>	56	0%	56%	0.01%	0%	99%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	2723	1%	1%	0.01%	0%	99%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	212	0%	14%	0.01%	0%	99%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	56	0%	50%	0.01%	0%	99%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	45	0%	60%	0.01%	0%	99%
5B2	5B2. Land converted to Cropland	CO <sub>2</sub>	48	0%	56%	0.01%	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	127	0%	21%	0.01%	0%	100%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	447	0%	5%	0.01%	0%	100%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	64	0%	32%	0.01%	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	0%	50%	0.01%	0%	100%
5G	5G. Other (liming of soils)	CO <sub>2</sub>	71	0%	25%	0.01%	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	16	0%	100%	0.01%	0%	100%
2C3	PFC from aluminium production	PFC	72	0%	20%	0.01%	0%	100%
5F2	5F2. Land converted to Other Land	CO <sub>2</sub>	25	0%	56%	0.01%	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	6	0%	71%	0.00%	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	89	0%	5%	0.00%	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	18	0%	22%	0.00%	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	2	0%	112%	0.00%	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	0%	112%	0.00%	0%	100%
1A1c	Stationary combustion : Manuf. of So- lid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	1	0%	20%	0.00%	0%	100%

With respect to Tier 2 level key sources, and perhaps surprisingly, the energy industries, with the highest share (30%) in the national total are not number one when uncertainty estimates are included. As Table A1.5 shows, two large but quite uncertain N<sub>2</sub>O sources are now in the top five list of level key sources:

- indirect N<sub>2</sub>O emissions from nitrogen used in agriculture;
- direct N<sub>2</sub>O emissions from agricultural soils.

The uncertainty in these emissions is estimated at 50% to 200%, with indirect N<sub>2</sub>O emissions having an uncertainty of a factor of two; one or two orders of magnitude higher than the 4% uncertainty estimated for CO<sub>2</sub> from the energy industries.

Source ranking using IPCC Tier 2 trend assessment excluding LULUCF (in Gg CO<sub>2</sub> eq).

Table A1.6

IPCC	Category	Gas	CO <sub>2</sub> eq base year	CO <sub>2</sub> eq last year	level assessment last year	trend assessment	Uncertainty estimate	Trend * uncertainty	% Contr. to trend	Cumulative
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	4975	3112	2%	1%	206%	2%	24%	24%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	12011	4896	2%	3%	34%	1%	15%	39%
2B2	Nitric acid production	N <sub>2</sub> O	6330	558	0%	3%	22%	1%	8%	47%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	1449	636	0%	0%	100%	0%	5%	53%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	5759	212	0%	3%	14%	0%	5%	58%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	7632	10928	5%	2%	20%	0%	5%	63%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	247	1623	1%	1%	51%	0%	5%	67%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	584	1624	1%	1%	50%	0%	4%	71%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	11832	20841	10%	5%	5%	0%	3%	74%
2C3	PFC from aluminium production	PFC	1901	72	0%	1%	20%	0%	2%	76%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	775	72	0%	0%	50%	0%	2%	79%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	1476	363	0%	1%	20%	0%	1%	80%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	273	52	0%	0%	100%	0%	1%	82%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4674	4883	2%	0%	61%	0%	1%	83%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	1252	390	0%	0%	25%	0%	1%	84%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	601	2242	1%	1%	11%	0%	1%	86%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	2738	1010	0%	1%	10%	0%	1%	87%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	6783	5782	3%	0%	16%	0%	1%	88%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	13348	23927	12%	5%	1%	0%	1%	88%
4B	Emissions from manure management	N <sub>2</sub> O	814	906	0%	0%	100%	0%	1%	89%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	19020	13817	7%	2%	2%	0%	1%	90%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1526	1940	1%	0%	21%	0%	1%	91%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2893	2350	1%	0%	20%	0%	1%	91%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9999	9059	4%	0%	14%	0%	1%	92%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	271	432	0%	0%	50%	0%	1%	92%
2B5	Caprolactam production	N <sub>2</sub> O	766	481	0%	0%	28%	0%	0%	93%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	250	117	0%	0%	54%	0%	0%	93%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	189	56	0%	0%	50%	0%	0%	94%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	157	45	0%	0%	60%	0%	0%	94%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	207	837	0%	0%	10%	0%	0%	95%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	2514	1349	1%	1%	6%	0%	0%	95%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	226	327	0%	0%	50%	0%	0%	95%

IPCC	Category	Gas	CO <sub>2</sub> eq base year	CO <sub>2</sub> eq last year	level assessment last year	trend assessment	Uncertainty estimate	Trend * uncertainty	% Contr. to trend	Cumulative
2A7	Other minerals	CO <sub>2</sub>	275	471	0%	0%	25%	0%	0%	96%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	405	637	0%	0%	20%	0%	0%	96%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	316	129	0%	0%	27%	0%	0%	96%
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	10902	12886	6%	1%	2%	0%	0%	97%
2B5	Other chemical product manufacture	CO <sub>2</sub>	606	649	0%	0%	71%	0%	0%	97%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7330	7503	4%	0%	10%	0%	0%	97%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	301	224	0%	0%	56%	0%	0%	98%
2F	PFC emissions from PFC use	PFC	37	180	0%	0%	25%	0%	0%	98%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1574	1494	1%	0%	100%	0%	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	566	392	0%	0%	20%	0%	0%	98%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	8644	9001	4%	0%	5%	0%	0%	98%
6B	Emissions from wastewater handling	CH <sub>4</sub>	290	198	0%	0%	32%	0%	0%	99%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	18696	17619	9%	0%	5%	0%	0%	99%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	438	379	0%	0%	50%	0%	0%	99%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	5033	4681	2%	0%	10%	0%	0%	99%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	1	64	0%	0%	32%	0%	0%	99%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	403	430	0%	0%	50%	0%	0%	99%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	1042	2723	1%	1%	1%	0%	0%	100%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1140	1118	1%	0%	100%	0%	0%	100%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25776	25365	12%	0%	3%	0%	0%	100%
2A3	Limestone and dolomite use	CO <sub>2</sub>	232	261	0%	0%	25%	0%	0%	100%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	255	269	0%	0%	25%	0%	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	11	16	0%	0%	100%	0%	0%	100%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	319	296	0%	0%	30%	0%	0%	100%
6B	Emissions from wastewater handling	N <sub>2</sub> O	466	459	0%	0%	54%	0%	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	148	127	0%	0%	21%	0%	0%	100%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	395	447	0%	0%	5%	0%	0%	100%
2B1	Ammonia production	CO <sub>2</sub>	3096	2850	1%	0%	2%	0%	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	3	6	0%	0%	71%	0%	0%	100%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	297	293	0%	0%	51%	0%	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	12	18	0%	0%	22%	0%	0%	100%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	162	155	0%	0%	54%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	1	2	0%	0%	112%	0%	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	41	0%	0%	50%	0%	0%	100%
2A1	Cement production	CO <sub>2</sub>	416	399	0%	0%	11%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	1	0%	0%	112%	0%	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	2	1	0%	0%	20%	0%	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	91	89	0%	0%	5%	0%	0%	100%

IPCC	Category	Gas	CO <sub>2</sub> eq base year	CO <sub>2</sub> eq last year	level assessment last year	trend assessment	Uncertainty estimate	Trend * uncertainty	% Contr. to trend	Cumulative
4D3	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	4975	3112	1%	1%	206%	2%	23%	23%
6A1	CH <sub>4</sub> emissions from solid waste disposal sites	CH <sub>4</sub>	12011	4896	2%	3%	34%	1%	14%	37%
2B2	Nitric acid production	N <sub>2</sub> O	6330	558	0%	3%	22%	1%	8%	45%
4D2	Animal production on agricultural soils	N <sub>2</sub> O	1449	636	0%	0%	100%	0%	5%	50%
2E	HFC-23 emissions from HCFC-22 manufacture	HFC	5759	212	0%	3%	14%	0%	5%	55%
2F	Emissions from substitutes for ozone depleting substances (ODS substitutes): HFC	HFC	247	1623	1%	1%	51%	0%	4%	59%
1A4a	Stationary combustion : Other Sectors: Commercial/Institutional, gases	CO <sub>2</sub>	7632	10928	5%	2%	20%	0%	4%	64%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	584	1624	1%	1%	50%	0%	3%	67%
1A3b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	11832	20841	10%	4%	5%	0%	3%	70%
5A2	5A2. Land converted to Forest Land	CO <sub>2</sub>	3	639	0%	0%	63%	0%	3%	73%
2C3	PFC from aluminium production	PFC	1901	72	0%	1%	20%	0%	2%	75%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	775	72	0%	0%	50%	0%	2%	77%
1A4	Stationary combustion : Other Sectors, liquids excl. From 1A4c	CO <sub>2</sub>	1476	363	0%	1%	20%	0%	1%	79%
4B9	Emissions from manure management : poultry	CH <sub>4</sub>	273	52	0%	0%	100%	0%	1%	80%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	1252	390	0%	0%	25%	0%	1%	81%
4D1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4674	4883	2%	0%	61%	0%	1%	82%
1A1a	Stationary combustion : Public Electricity and Heat Production: waste incineration	CO <sub>2</sub>	601	2242	1%	1%	11%	0%	1%	84%
5A1	5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	2529	2208	1%	0%	67%	0%	1%	85%
1A3b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	2738	1010	0%	1%	10%	0%	1%	86%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	6783	5782	3%	0%	16%	0%	1%	87%
1A1a	Stationary combustion : Public Electricity and Heat Production: gases	CO <sub>2</sub>	13348	23927	11%	5%	1%	0%	1%	87%
4B	Emissions from manure management	N <sub>2</sub> O	814	906	0%	0%	100%	0%	1%	88%
5C2	5C2. Land converted to Grassland	CO <sub>2</sub>	394	550	0%	0%	66%	0%	1%	89%
1A2	Stationary combustion : Manufacturing Industries and Construction, gases	CO <sub>2</sub>	19020	13817	6%	2%	2%	0%	1%	89%
1A1b	Stationary combustion : Petroleum Refining: liquids	CO <sub>2</sub>	9999	9059	4%	0%	14%	0%	1%	90%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: gases	CO <sub>2</sub>	1526	1940	1%	0%	21%	0%	1%	91%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, liquids	CO <sub>2</sub>	2893	2350	1%	0%	20%	0%	1%	91%
1A3	Mobile combustion: road vehicles	N <sub>2</sub> O	271	432	0%	0%	50%	0%	1%	92%
2B5	Caprolactam production	N <sub>2</sub> O	766	481	0%	0%	28%	0%	0%	92%
3, 6D	OTHER N <sub>2</sub> O	N <sub>2</sub> O	250	117	0%	0%	54%	0%	0%	93%
1A4	Stationary combustion : Other Sectors, solids	CO <sub>2</sub>	189	56	0%	0%	50%	0%	0%	93%
1A3	Mobile combustion: road vehicles	CH <sub>4</sub>	157	45	0%	0%	60%	0%	0%	94%
2C1	Iron and steel production (carbon inputs)	CO <sub>2</sub>	2514	1349	1%	1%	6%	0%	0%	94%
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	207	837	0%	0%	10%	0%	0%	94%
5C1	5C1. Grassland remaining Grassland	CO <sub>2</sub>	4246	4246	2%	0%	56%	0%	0%	95%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	226	327	0%	0%	50%	0%	0%	95%
2A7	Other minerals	CO <sub>2</sub>	275	471	0%	0%	25%	0%	0%	95%
5E2	5E2. Land converted to Settlements	CO <sub>2</sub>	212	296	0%	0%	56%	0%	0%	96%
1A3	Mobile combustion: water-borne navigation	CO <sub>2</sub>	405	637	0%	0%	20%	0%	0%	96%
3	Indirect CO <sub>2</sub> from solvents/product use	CO <sub>2</sub>	316	129	0%	0%	27%	0%	0%	96%



IPCC	Category	Gas	CO <sub>2</sub> eq base year	CO <sub>2</sub> eq last year	level as- essment last year	trend assess- ment	Uncer- tainty estimate	Trend * uncer- tainty	% Contr. to trend	Cumu- lative
1A3b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	10902	12886	6%	1%	2%	0%	0%	97%
2B5	Other chemical product manufacture	CO <sub>2</sub>	606	649	0%	0%	71%	0%	0%	97%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	301	224	0%	0%	56%	0%	0%	97%
4B1	Emissions from manure management : cattle	CH <sub>4</sub>	1574	1494	1%	0%	100%	0%	0%	97%
1A4c	Stationary combustion : Other Sectors, Agriculture/Forestry/Fisheries, gases	CO <sub>2</sub>	7330	7503	3%	0%	10%	0%	0%	98%
2F	PFC emissions from PFC use	PFC	37	180	0%	0%	25%	0%	0%	98%
1A5	Military use of fuels (1A5 Other)	CO <sub>2</sub>	566	392	0%	0%	20%	0%	0%	98%
1A2	Stationary combustion : Manufacturing Industries and Construction, liquids	CO <sub>2</sub>	8644	9001	4%	0%	5%	0%	0%	98%
1A4b	Stationary combustion : Other Sectors, Residential, gases	CO <sub>2</sub>	18696	17619	8%	0%	5%	0%	0%	98%
6B	Emissions from wastewater handling	CH <sub>4</sub>	290	198	0%	0%	32%	0%	0%	99%
5G	5G. Other (liming of soils)	CO <sub>2</sub>	183	71	0%	0%	25%	0%	0%	99%
4A8	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	438	379	0%	0%	50%	0%	0%	99%
1A2	Stationary combustion : Manufacturing Industries and Construction, solids	CO <sub>2</sub>	5033	4681	2%	0%	10%	0%	0%	99%
6D	OTHER CH <sub>4</sub>	CH <sub>4</sub>	1	64	0%	0%	32%	0%	0%	99%
1A1b	Stationary combustion : Petroleum Refining: gases	CO <sub>2</sub>	1042	2723	1%	1%	1%	0%	0%	99%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	403	430	0%	0%	50%	0%	0%	99%
5D2	5D2. Land converted to Wetlands	CO <sub>2</sub>	40	56	0%	0%	56%	0%	0%	100%
2A3	Limestone and dolomite use	CO <sub>2</sub>	232	261	0%	0%	25%	0%	0%	100%
1A1a	Stationary combustion : Public Electricity and Heat Production: solids	CO <sub>2</sub>	25776	25365	12%	0%	3%	0%	0%	100%
4B8	Emissions from manure management : swine	CH <sub>4</sub>	1140	1118	1%	0%	100%	0%	0%	100%
5B2	5B2. Land converted to Cropland	CO <sub>2</sub>	35	48	0%	0%	56%	0%	0%	100%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	255	269	0%	0%	25%	0%	0%	100%
4B	Emissions from manure management : other	CH <sub>4</sub>	11	16	0%	0%	100%	0%	0%	100%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	319	296	0%	0%	30%	0%	0%	100%
5F2	5F2. Land converted to Other Land	CO <sub>2</sub>	18	25	0%	0%	56%	0%	0%	100%
2B1	Ammonia production	CO <sub>2</sub>	3096	2850	1%	0%	2%	0%	0%	100%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	148	127	0%	0%	21%	0%	0%	100%
2C3	CO <sub>2</sub> from aluminium production	CO <sub>2</sub>	395	447	0%	0%	5%	0%	0%	100%
6B	Emissions from wastewater handling	N <sub>2</sub> O	466	459	0%	0%	54%	0%	0%	100%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	3	6	0%	0%	71%	0%	0%	100%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	297	293	0%	0%	51%	0%	0%	100%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	162	155	0%	0%	54%	0%	0%	100%
2E	HFC by-product emissions from HFC manufacture	HFC	12	18	0%	0%	22%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	N <sub>2</sub> O	1	2	0%	0%	112%	0%	0%	100%
2A1	Cement production	CO <sub>2</sub>	416	399	0%	0%	11%	0%	0%	100%
1A3	Mobile combustion: aircraft	CO <sub>2</sub>	41	41	0%	0%	50%	0%	0%	100%
1A3	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	1	0%	0%	112%	0%	0%	100%
1A1c	Stationary combustion : Manuf. of Solid Fuels and Other En. Ind.: liquids	CO <sub>2</sub>	2	1	0%	0%	20%	0%	0%	100%
1A3	Mobile combustion: other (railways)	CO <sub>2</sub>	91	89	0%	0%	5%	0%	0%	100%



# Annex 2

## Detailed discussion of methodology and data for estimating CO<sub>2</sub> emissions from fossil fuel combustion

The Netherlands list of fuels and standard CO<sub>2</sub> emission factors was originally approved in 2004 by the Steering Committee Emission Registration, and is revised following decisions on the CO<sub>2</sub> emission factor for natural gas by this Steering Group in its meetings 25 April 2006 and 21 April 2009.

On 21 April 2009, Steering Committee Emission Registration delegated the authority to decide on revisions of the list to the Working Group Emission Monitoring (WEM). On 8 December 2009 the present document (version December 2009) was approved by the WEM.

For a description of the methodology and activity data used for the calculation of CO<sub>2</sub> emissions from fossil fuel combustion we refer to the monitoring protocols (see Annex 6, protocols 0042 for stationary sources and protocols 0044-0051 for mobile sources).

### A2.1 Introduction

For national monitoring of greenhouse gas emissions under the framework of the UN Climate Change Convention (UNFCCC) and monitoring at corporate level for the European CO<sub>2</sub> emissions trade, international agreements state that each country must draw up a national list of defined fuels and standard CO<sub>2</sub> emission factors. This is based on the IPCC list (with default CO<sub>2</sub> emission factors), but should include national values that reflect the specific national situation. This list will also be used by the Netherlands in the e-MJV (electronic annual environmental report), because these are used for the national monitoring.

The Netherlands list of energy carriers and standard CO<sub>2</sub> emission factors (further referred to as “the Netherlands list”) is now available in the form of:

1. a table containing the names (in Dutch and English) of the energy carrier and the accompanying standard energy content and CO<sub>2</sub> emissions factor;
2. a fact sheet per energy carrier, substantiating the values given, presenting similar names and possible specifications, and providing an overview of the codes that organizations use for the individual energy carriers.

This Annex is for people using the Netherlands list. It contains the starting points for this list and indicates how it should be used for various objectives, for example, national monitoring of greenhouse gas emissions, the European CO<sub>2</sub> emissions trade and the e-MJV. It also includes background information. The list, plus this document and the background documents for substantiating the specific Netherlands values can be found on the website: [www.greenhousegases.nl](http://www.greenhousegases.nl) / [www.broeikasgassen.nl](http://www.broeikasgassen.nl).

Based on new scientific knowledge in 2006 the CO<sub>2</sub> emission factor for natural gas has been changed for the period 1990-2006.

From 2007 onwards, the CO<sub>2</sub> emission factor for natural gas has been assessed annually. In this document, the CO<sub>2</sub> emission factor for natural gas for 2008 has been determined.

### A2.2 Starting points for the Netherlands list

The following starting points were used to draw up the Netherlands list:

1. The list contains all the fuels, as included in the IPCC guidelines (Revised 1996 Intergovernmental Panel on Climate Change (IPCC) for national greenhouse gas inventories, further known as the “1996 IPCC guidelines”), Table 1-1 (in Chapter 1 of the Reference Manual, Volume 3 of the 1996 IPCC guidelines) and the differentiation thereof in the Workbook Table 1.2 (Module 1 of the Workbook, Volume 2 of the 1996 IPCC guidelines). The 1996 IPCC guidelines are applicable to the national monitoring of greenhouse gas emissions under the UNFCCC framework.
2. The list contains all fuels, as included in European Commission (EC) Directive 2004/156/EG on reporting CO<sub>2</sub> emissions trading (“... defining guidelines for monitoring and reporting greenhouse gas emissions...”), Appendix 1, Chapter 8.
3. The definition of fuels is based on the definition used by the CBS (Statistics Netherlands) when collating energy statistics.
4. As a result of the 1996 IPCC guidelines and the EC Directive 2004/156/EG mentioned in 1 and 2 above, the CO<sub>2</sub> emission factors are accurate to one digit after the decimal point.
5. The list assumes the standard CO<sub>2</sub> emission factors as used in the 1996 IPCC guidelines and the EC Directive 2004/156/EG but, where the Netherlands’ situation deviates from this norm, specific standard values for the Netherlands are used, which are documented and substantiated.

### A2.3 The Netherlands list

A study was carried out in 2002 with respect to specific Netherlands CO<sub>2</sub> emission factors (TNO, 2002). This study showed that, for a limited number of Dutch fuels, their situations deviated such that national values needed to be determined. For a number of fuels, the previously defined national values (Emission Registration, 2002) could be updated but for others new values were required.

A specific Netherlands standard CO<sub>2</sub> emissions factor has been determined for the following fuels, which does not appear in the 1996 IPCC Guidelines or in the EC Directive 2007/589/EG, but has been added as specification for one of the following fuels in:

1. petrol/gasoline
2. gas and diesel oil
3. LPG
4. coke coals (coke ovens and blast furnaces)
5. other bituminous coal
6. coke ovens/gas cokes
7. coke oven gas
8. blast furnace gas
9. oxygen furnace gas
10. phosphorus furnace gas
11. natural gas.

For industrial gases, chemical waste gas is also differentiated from refinery gas. For the IPCC main group “other fuels”, only non-biogenic waste is differentiated.

#### Coking Coal

For coking coal the standard CO<sub>2</sub> emissions factor is also a weighed average, for example of coke coals used in coke ovens and in blast furnaces.

#### Natural gas

In 2006, a study was commissioned to research methods to determine the CO<sub>2</sub> emission factor for natural gas (TNO, 2006). This resulted in an advice to use natural gas a country specific factor from the year 1990 onwards (SenterNovem, 2006). In its meeting 25 April 2006, the Steering Committee Emission Registration agreed with this advice and approved an update of the National list for the period 1990-2006.

From 2007 onwards, the CO<sub>2</sub> emission factor for natural gas has been assessed annually. In the meeting of the Steering Committee Emission Registration of 21 April 2009, the procedure was approved for the annual update of the emission factor of natural gas. In this document (version December 2009), the emission factor of natural gas for 2008 was determined according to this procedure.

#### Waste

From 2009 onwards, on the Netherlands list the fuel Waste (non-biogenic) is replaced by the fuel Waste. This fuel concerns all waste that is incinerated in the Netherlands, both residential waste and other waste. In addition, from 2009 onwards the heating value and the emission factor will be determined annually on the Netherlands list. These values are not used as input for the calculation of greenhouse gas emissions under the framework of the UN Climate Change Convention (UNFCCC), but are the result of these calculations (see Renewable Energy Monitoring Protocol update 2006). In the e-MJV these value can be used by companies that incinerate waste.

In this document (version December 2009), the heating value and the emission factor of Waste are determined for 2008. The incinerated waste is a mix of biogenic and non-biogenic waste. Therefore, the percentage biogenic is given for both the heating value and the emission factor.

#### Biomass

The list also includes biomass as a fuel, with accompanying specific Netherlands CO<sub>2</sub> emission factors. Biomass emissions are reported separately in the national monitoring of greenhouse gas emissions under the UNFCCC framework (as memo element) and are not included in the national emissions figures. For the European CO<sub>2</sub> emissions trading, the emissions are not included because an emissions factor of zero is used for biomass.

The CO<sub>2</sub> emissions factor for wood is used for solid biomass, and that of palm oil is used for liquid biomass<sup>1</sup>. A weighed average of three specified biogases is used as the standard factor for gaseous biomass:

1. sewage treatment facility (WWTP) biogas

<sup>1</sup> The heating value and the emission factor of liquid biomass are not used in the calculations of the national transport emissions for biofuels. For an explanation, see Klein, 2009 (. Table 1.31)

# Netherlands fuels and standard CO<sub>2</sub> emission factors, version December 2009

Main group (Dutch)	Main group (English) IPCC (supplemented)	Unit	Heating value (MJ/unit)	CO <sub>2</sub> EF (kg/GJ)
	A. Liquid Fossil, Primary Fuels			
<i>Ruwe aardolie</i>	Crude oil	kg	42.7	73.3
<i>Orimulsion</i>	Orimulsion	kg	27.5	80.7
<i>Aardgascondensaat</i>	Natural Gas Liquids	kg	44.0	63.1
	Liquid Fossil, Secondary Fuels/ Products			
<i>Motorbenzine</i>	Petrol/gasoline	kg	44.0	72.0
<i>Kerosine luchtvaart</i>	Jet Kerosene	kg	43.5	71.5
<i>Petroleum</i>	Other Kerosene	kg	43.1	71.9
<i>Leisteenolie</i>	Shale oil	kg	36.0	73.3
<i>Gas-/dieselolie</i>	Gas/ Diesel oil	kg	42.7	74.3
<i>Zware stookolie</i>	Residual Fuel oil	kg	41.0	77.4
<i>LPG</i>	LPG	kg	45.2	66.7
<i>Ethaan</i>	Ethane	kg	45.2	61.6
<i>Nafta's</i>	Naphtha	kg	44.0	73.3
<i>Bitumen</i>	Bitumen	kg	41.9	80.7
<i>Smeeroliën</i>	Lubricants	kg	41.4	73.3
<i>Petroleumcokes</i>	Petroleum Coke	kg	35.2	100.8
<i>Raffinaderij grondstoffen</i>	Refinery Feedstocks	kg	44.8	73.3
<i>Raffinaderijgas</i>	Refinery Gas	kg	45.2	66.7
<i>Chemisch restgas</i>	Chemical Waste Gas	kg	45.2	66.7
<i>Overige oliën</i>	Other Oil	kg	40.2	73.3
	B. Solid Fossil, Primary Fuels			
<i>Antraciet</i>	Anthracite	kg	26.6	98.3
<i>Cokeskolen</i>	Coking Coal	kg	28.7	94.0
<i>Cokeskolen (cokeovens)</i>	Coking Coal (used in coke oven)	kg	28.7	95.4
<i>Cokeskolen (basismetale)</i>	Coking Coal (used in blast furnaces)	kg	28.7	89.8
<i>(Overige bitumineuze) steenkool</i>	Other Bituminous Coal	kg	24.5	94.7
<i>Sub-bitumineuze kool</i>	Sub-bituminous Coal	kg	20.7	96.1
<i>Bruinkool</i>	Lignite	kg	20.0	101.2
<i>Bitumineuze Leisteen</i>	Oil Shale	kg	9.4	106.7
<i>Turf</i>	Peat	kg	10.8	106.0
	Solid Fossil, Secondary Fuels			
<i>Steenkool- en bruinkoolbriketten</i>	BKB & Patent Fuel	kg	23.5	94.6
<i>Cokesoven/ gascokes</i>	Coke Oven/Gas Coke	kg	28.5	111.9
<i>Cokesovengas</i>	Coke Oven gas	MJ	1.0	41.2
<i>Hoogovengas</i>	Blast Furnace Gas	MJ	1.0	247.4
<i>Oxystaalovengas</i>	Oxy Gas	MJ	1.0	191.9
<i>Fosforovengas</i>	Phosphor Gas	Nm <sup>3</sup>	11.6	149.5
	C. Gaseous Fossil Fuels			
<i>Aardgas</i>	Natural Gas (dry)	Nm <sup>3</sup>	31.65	56.7 <sup>2</sup>
<i>Koolmonoxide</i>	Carbon Monoxide	Nm <sup>3</sup>	12.6	155.2
<i>Methaan</i>	Methane	Nm <sup>3</sup>	35.9	54.9
<i>Waterstof</i>	Hydrogen	Nm <sup>3</sup>	10.8	0.0
	Biomass <sup>3</sup>			
<i>Biomassa vast</i>	Solid Biomass	kg	15.1	109.6
<i>Biomassa vloeibaar</i>	Liquid Biomass	kg	39.4	71.2
<i>Biomassa gasvormig</i>	Gas Biomass	Nm <sup>3</sup>	21.8	90.8
<i>RWZI biogas</i>	Wastewater biogas	Nm <sup>3</sup>	23.3	84.2
<i>Stortgas</i>	Landfill gas	Nm <sup>3</sup>	19.5	100.7
<i>Industrieel fermentatiegas</i>	Industrial organic waste gas	Nm <sup>3</sup>	23.3	84.2
	D. Other fuels			
<i>Afval<sup>4</sup></i>	Waste	kg	10.3	97.5

2. landfill gas

3. industrial organic waste gas.

The heating values are the same as those used by the CBS for observed fuels in its surveys for collating energy statistics.

<sup>2</sup> The emission factor for natural gas in this table (56.7 kg CO<sub>2</sub>/GJ) is applicable for the calculation of the emissions in 2007 (Zijlema, 2008) and in 2008 (Zijlema, 2009). For the period 1990-2006, the emission factor remains unchanged (56.8 kg CO<sub>2</sub>/GJ). In the future, the emission factor for natural gas will be updated annually.

<sup>3</sup> Biomass: the value of the CO<sub>2</sub> emission factor is shown as a memo item in reports for the climate change convention; the value is zero for emissions trading and for the Kyoto Protocol.

<sup>4</sup> The percentage biogenic in the heating value is 49%. The percentage biogenic in the emission factor is 63% (based on 2008).

## A2.4 Fact sheets

A fact sheet (consisting of at least two sections) has been drawn up for each fuel:

1. General information
  - a. Name of the fuel, in Dutch and English
  - b. Other names used (Dutch and English)
  - c. Description
  - d. Codes (in Dutch) used to specify the fuel
  - e. Unit
2. Specific values and substantiation
  - a. Heating value
  - b. Carbon content
  - c. CO<sub>2</sub> emissions factor
  - d. Density (if relevant), converting from weight to volume or converting from gases to m<sup>3</sup> standard natural gas equivalents
  - e. Substantiating the choices, plus accurate referral to references and/or specific text sections within the reference
  - f. Year and/or period for which the specific values apply.

If a standard Dutch value for a fuel exists, this has been added to the fact sheet (as a third section containing the same information as that described under 1) and 2) above).

## A2.5 Using the Netherlands list in national monitoring, European CO<sub>2</sub> emissions trade and in the e-MJV National monitoring

### National monitoring

The 1996 IPCC Guidelines are among those valid for national monitoring under the UNFCCC framework, which is reported annually in the NIR (National Inventory Report). This includes the default CO<sub>2</sub> emission factors shown in Table 1-1 (Chapter 1 of the Reference Manual, volume 3 of the 1996 IPCC Guidelines) and Table 1-2 (Module 1 of the Workbook, volume 2 of the 1996 IPCC Guidelines). With respect to the specification at national level: ‘...default assumptions and data should be used only when national assumptions and data are not available.’ (Overview of the Reporting Instructions, volume 1 of the 1996 IPCC Guidelines) and ‘...because fuel qualities and emission factors may differ markedly between countries, sometimes by as much as 10% for nominally similar fuels, national inventories should be prepared using local emission factors and energy data where possible.’ (Chapter 1, Section 1.1 of the Reference Manual, volume 3 of the 1996 IPCC Guidelines).

With respect to documentation: ‘When countries use local values for the carbon emission factors they should note the differences from the default values and provide documentation supporting the values used in the national inventory cal-

culations’ (Chapter 1, Section 1.4.1.1 of the Reference Manual, volume 3 of the 1996 IPCC Guidelines). Exactly when and how the Netherlands list should be used in the national monitoring process is further described in the 1996 IPCC Guidelines. The Netherlands list is included in the country’s national report to the UNFCCC on greenhouse gas emissions.

### Monitoring European CO<sub>2</sub> emissions trade

The EC Directive 2007/589/EG covers monitoring under the framework of the European CO<sub>2</sub> emissions trade. This Directive serves as a starting point for the Netherlands monitoring system for trading in emission allowances. With respect to the CO<sub>2</sub> emission factors and the calculations of CO<sub>2</sub> emissions at level 2a, the Directive states: ‘The operator should use the relevant fuel calorific values that apply in that Member State, for example as indicated in the relevant Member State’s latest national inventory, which has been submitted to the secretariat of the UNFCCC (EC Directive 2007/589/EC, Appendix II, Section 2.1.1.1).

With respect to the reports, this states that: ‘Fuels, and the resulting emissions must be reported in accordance with the IPCC format for fuels (...) this is based on the definitions set out by the IEA (International Energy Agency). If the Member State (relevant to the operator) has already published a list of fuel categories, including definitions and emission factors, which is consistent with the latest national inventory such as submitted to the UNFCCC secretariat, these categories and the accompanying emission factors should be used, if these have been approved within the framework of the relevant monitoring methodology.’ (EC Directive 2007/589/EG, Appendix I, Section 5).

When and how the Netherlands list should be used in the monitoring process under the framework of the EU CO<sub>2</sub> emissions trading is further explained in EC Directive 2007/589/EG and the Netherlands system for monitoring the trade in emission allowances.

### e-MJV

Within the UNFCCC framework, the national monitoring of greenhouse gases is partly based on the information provided in the MJVs (annual environmental reports). Information on CO<sub>2</sub> emissions trading is (also) reported in the MJV, which is why the Netherlands list is also used in the e-MJV. Since the monitoring of the energy covenant known as MJA (long-term energy agreement) can be carried out via the e-MJV, the Netherlands list is also used to compile these reports. Exactly how the Netherlands list should be used in the e-MJV is further described in the e-MJV itself.

### Use of the Netherlands list by other stakeholders in the Netherlands

The Netherlands list can also be used for other purposes (e.g., monitoring energy covenants, predicting future CO<sub>2</sub> emissions). Selections can be taken from the list, depending on the application. This usage is not defined in the legislation, but offers the advantage of harmonising the national monitoring under the UNFCCC framework. Whenever CO<sub>2</sub> emissions are defined for the Government, the Netherlands list will be used wherever possible.

## A2.6 Defining and maintaining the Netherlands list

VROM (Spatial Planning, Housing and the Environment) initiated the compilation of the Netherlands list, as it is responsible for the national monitoring of greenhouse gas emissions under the UNFCCC framework. This list has been prepared in consultation with those national institutes involved in national monitoring activities, such as PBL, CBS and NL Agency, and other relevant organisations, such as the e-MJV, CO<sub>2</sub> emissions trade and ECN. The Steering Committee Emission Registration (the collaborative agencies implementing the national monitoring) compiled the list during its meeting in October 2004.

The list will be maintained within the National System, the organisational structure that coordinates national greenhouse gas monitoring under the UNFCCC framework. The Netherlands list, this document and the background documents are all publicly accessible from the Dutch website ([www.broeikasgassen.nl](http://www.broeikasgassen.nl) or the English version, [www.greenhousegases.nl](http://www.greenhousegases.nl)). As part of the quality monitoring system for national monitoring of greenhouse gases, this list will be evaluated every three years.

This document was updated in November 2005 with some editorial changes. This document and the Netherlands list were updated in 2006 based on research for methods to determine the CO<sub>2</sub> emission factor for natural gas in the Netherlands for the period 1990-2006.

From 2007 onwards, the CO<sub>2</sub> emission factor for natural gas has been assessed annually, based on measurement by Gasunie and Zebragas. On 21 April 2009, this procedure was approved by the Steering Committee Emission Registration.

On 21 April 2009, Steering Committee Emission Registration delegated the authority to decide on revisions of the list to the Working Group Emission Monitoring (WEM). On 8 December 2009, the present document (version December 2009) was approved by the WEM. In this document (version December 2009) the CO<sub>2</sub> emission factor for natural gas for 2008 has been determined. In addition, the fuel Waste (non-biogenic) has been replaced by the fuel Waste, for which the heating value and the emission factor will be determined annually.

## A2.7 Application of the Netherlands standard and source-specific CO<sub>2</sub> emission factors in the national emission inventory

For the most common fuels (natural gas, coal, coal products, diesel and petrol), country-specific standard CO<sub>2</sub> emission factors are used; otherwise default IPCC emission factors are used (see Table A2.1). However, for some of the derived fuels the chemical composition and thus the CO<sub>2</sub> emission factor is highly variable between source categories and over time.

Thus, for blast furnace gas and oxygen furnace gas, refinery gas, chemical waste gas (liquids and solids treated separately) and solid waste (the biogenic and fossil carbon part treated separately), mostly source-specific (or plant-specific) emission

factors have been used, that may also change over time. In addition, for raw natural gas combustion by the oil and gas production industry a source-specific (or company-specific) CO<sub>2</sub> emission factor is used. This refers to the “own use” of unprocessed natural gas used by the gas and oil production industry, of which the composition may differ significantly from that of treated standard natural gas supplied to end-users. These emission factors are based on data submitted by industries in their Annual Environmental Reports (MJVs). These fuels are used in the subcategories “Public electricity and heat production” (1A1a), “Refineries” (1A1b) and “Other energy industries” included in 1A1c.

Fossil-based CO<sub>2</sub> emissions from waste incineration are calculated from the total amount of waste that is incinerated, split into six waste types per waste stream, each with a specific carbon content and fraction of fossil carbon in total carbon (see Section 8.4.2 for more details).

More details on methodologies, data sources used and country-specific source allocation issues are provided in the monitoring protocols (see Annex 6).





# Annex 3

## Other detailed methodological descriptions for individual source or sink categories

A detailed description of methodologies per source/ sink category can be found in protocols on the website [www.greenhousegases.nl](http://www.greenhousegases.nl), including country-specific emission factors. Annex 6 provides an overview of the available monitoring protocols at this site.



# Annex 4

## CO<sub>2</sub> Reference Approach and comparison with Sectoral Approach

### A4.1 Comparison of CO<sub>2</sub> emissions

The IPCC Reference Approach (RA) for CO<sub>2</sub> from energy use uses apparent consumption data per fuel type to estimate CO<sub>2</sub> emissions from fossil fuel use. This has been used as a means of verifying the sectoral total CO<sub>2</sub> emissions from fuel combustion (IPCC, 2001). For the Reference Approach energy statistics (production, imports, export, stock changes) were provided by Statistics Netherlands (CBS); national default, partly country-specific, CO<sub>2</sub> emission factors (see Annex 2.1, Tables A2.1 and A2.2) and constant carbon storage fractions based on the average of annual carbon storage fractions calculated per fossil fuel type for 1995-2002 from reported CO<sub>2</sub> emissions in the sectoral approach. Also, bunker fuels were corrected for the modification made to include fisheries, internal navigation, military aviation and shipping in domestic consumption instead of included in the bunker total in the original national energy statistics (see Annex 2.1, Tables A2.1 - A2.3).

Table A4.1 presents the results of the Reference Approach calculation for 1990-2008 and compared with the official national total emissions reported as fuel combustion (source category 1A). The annual difference calculated from the direct comparison varies between 0.3% for 2008 and 3.8% for 1990.

The Reference Approach (RA) and National Approach (NA) data show a 13% RA vs. 14% NA increase in emissions from liquid fuels (1990-2008) and a 12% RA vs. 14% NA increase from gaseous fuels; CO<sub>2</sub> emissions from solid fuels decreased in this period by 8% in the RA vs. an decrease of 3% in the NA. The emissions from others (fossil carbon in waste), which is only included in the NA increased from 0.6 Tg in 1990 to 2.2 Tg CO<sub>2</sub> in 2008. However, as will be discussed below, these numbers cannot be compared well since the RA includes sources not included in the NA and vice versa. Therefore, a corrected comparison will be made below.

### A4.2 Causes of differences between the two approaches

There are five main reasons for differences in the two approaches, of which two are *inherent to the comparison method* itself (see Table A4.2):

1. The CO<sub>2</sub> from *incineration of waste* that contains fossil carbon (reported under 6C or 1A1a) is not included in the Reference Approach;
2. The fossil-fuel related emissions reported as *process emissions* (sector 2) and *fugitive emissions* (Sector 1B), which are not included in the Sectoral Approach total of Sector 1A. The most significant are gas used as feedstock in ammonia production (2B1) and losses from coke/coal inputs in blast furnaces (2C1);

and others are *country-specific*:

3. In addition, the country-specific *carbon storage factors* used in the Reference Approach are multi-annual averages, so the RA calculation for a specific year will deviate somewhat from the factors that could be calculated from the specific mix of feedstock/non-energy uses of different fuels;
4. The use of *plant-specific emission factors* in the NA vs. national defaults in the RA;
5. Other differences could – in principle – be due to the presence of *statistical differences* between apparent consumption and total sectoral fuel use and/or to differences between total sectoral fuel use as used in the emission inventory and as included in the national energy statistics in cases where *plant-specific fuel use data* have been used.

However, the latter is not applicable to the Netherlands. National statistics are compiled in such a way that no statistical difference occurs (initial differences are removed by shifting to the most uncertain fuel entry). Moreover, the calculations are all based on the official sectoral energy statistics from Statistics Netherlands (CBS), which guarantees that the activity data in the inventory are identical to the national energy statistics.

	1990	1995	2000	2005	2006	2007	2008
<b>Reference Approach</b>							
<i>Liquid fuels</i> <sup>1)</sup>	49.7	51.4	53.8	55.2	54.6	58.2	55.9
<i>Solid fuels</i> <sup>1)</sup>	34.0	34.7	30.5	32.2	30.2	33.2	31.4 <sub>4</sub>
<i>Gaseous fuels</i>	71.9	79.9	81.0	81.8	79.6	77.1	80.6
<i>Others</i>	NA	NA	NA	NA	NA	NA	NA
<b>Total RA</b>	155.6	166.0	165.3	169.2	164.4	168.5	167.9
<b>National Approach</b>							
<i>Liquid fuels</i>	49.8	52.3	54.4	56.5	56.2	56.0	56.7
<i>Solid fuels</i>	31.0	32.5	28.8	30.1	28.6	30.7	30.1
<i>Gaseous fuels</i>	68.6	76.0	76.7	78.5	77.0	74.5	78.5
<i>Others</i> <sup>2)</sup>	0.6	0.8	1.5	2.1	2.1	2.2	2.2
<b>Total NA</b>	149.9	161.6	161.4	167.2	163.8	163.4	167.5
<b>Difference</b> <sup>3)</sup> (%)							
<i>Liquid fuels</i>	-0.1%	-1.7%	-1.1%	-2.3%	-2.8%	3.9%	-1.3%
<i>Solid fuels</i>	9.9%	7.2%	6.1%	6.9%	5.8%	8.2%	4.2%
<i>Gaseous fuels</i>	4.8%	5.2%	5.5%	4.2%	3.4%	3.4%	2.8%
<i>Other</i>	-100%	-100%	-100%	-100%	-100%	-100%	-100%
<b>Total</b>	3.8%	2.8%	2.4%	1.2%	0.4%	3.1%	0.3%

<sup>1)</sup> Specification of national fuel types used in the IPCC fuel type categories:

*Gasoline*: jet fuel, gasoline basis; aviation gasoline; motor gasoline; *Other Kerosene*: petroleum; *Other Oil*: oil aromates; other light oils; other oil products; *Other Bituminous Coal*: all hard coal; lignite/brown coal; *BKB and Patent Fuel*: coal derivatives.

<sup>2)</sup> Fossil-fuel component of waste combustion in waste incineration that also produce heat and electricity for energy purposes.

<sup>3)</sup> Defined as: (RA-NA)/NA.

### Correction of inherent differences

The correction terms for the RA/NA total are for the Netherlands:

1. waste incineration (in the Netherlands included in 1A1a, as 'other fuels');
2. selected CRF Sector 2 components listed in Table A4.2 and selected fugitive CO<sub>2</sub> emissions included in CRF Sector 1B.

If the RA is corrected by including the fossil waste and the NA by including selected Sector 1B and Sector 2 emissions that should be added to the 1A total before the comparison is made (see Table A4.2), then a much smaller difference remains between the approaches. Remaining differences are generally below  $\pm 1\%$  between +1.2% in 2007 and -1.4% in 2006, with a direct average of  $(-0.2 \pm 0.3)\%$  vs.  $(-0.2 \pm 2.1)\%$  in the uncorrected comparison.

The corrected RA and NA comparison per fuel type is presented in Table A4.3. This shows that the largest differences do not concentrate in a particular corner of the period. The corrected 1990-2007 trends also differ only slightly: 8.3% for the corrected National Approach (NA) (=F sum of sectoral emissions in source category 1A plus selected 1B and 2 minus fossil waste) and 8.3% for the corrected Reference Approach. We conclude that in total annual emissions, the remaining differences are now all between about -1.5% and +0.5%, except for 2006 which shows a -2.1% difference.

The corrected approaches show differences in emissions from liquid fuels up to -5% for a single year vs. -3% for uncorrected comparisons. For solid fuels, differences are up to 2% vs. 11% and for gaseous fuels -1% vs. +5%, respectively, if corrections are made for 2G (non energy uses of lubricants and waxes) in NA-liquids, 1B (coke production), 2A ('Soda Ash'), 2B5, 2C1 (blast furnaces) and 2D in NA-solids; and 1B2 (gas flaring, refineries) and 2B1 (ammonia) in NA-gases (Table

A4.2). Remaining differences must be due to the use of one multi-annual average carbon storage factor per fuel type for all years (see Section A4.3) and plant-specific emission factors in some cases as discussed in Section A4.4 (for more details, see Annex 2, Table A2.2).

### A4.3 Other country-specific data used in the Reference Approach

Apart from different *storage fractions* of non-energy use of fuels as presented in Table A4.5, other country-specific information used in the RA is found in:

#### Carbon contents (. CO<sub>2</sub> emission factors) used

For the fuels used in the Reference Approach, the factors used are listed in Table A.2.1. These are the national defaults. For 'other bituminous coal' and 'BKB & Patent fuel' the values are used of bituminous coal and coal bitumen respectively;

#### Fuel consumption in international marine and aviation bunkers

Some changes were made in the national energy statistics of total apparent consumption, mainly for diesel, jet kerosene and residual fuel oil, due the reallocation for the emissions inventory of part of the bunker fuels to domestic consumption (e.g., fisheries and inland navigation). This explains the difference between the original bunker statistics in the national energy statistics (and as reported to international agencies such as the IEA) and the bunker fuel data used in the Reference Approach calculation.

RA,NA, correction term	1990	1995	2000	2005	2006	2007	2008
Difference RA-NA	5.7	4.5	3.8	2.0	0.6	5.1	0.5
Reference Approach:	155.6	166.0	165.3	169.2	164.4	168.5	167.9
Other: fossil waste cf. NA	0.6	0.8	1.6	2.1	2.1	2.2	2.2
RA incl. fossil waste:	156.2	166.9	166.9	171.3	166.6	170.7	170.2
Diff. RA incl. Waste-NA:	5.1	3.7	2.2	0.0	-1.5	2.9	-1.8
National Approach:	149.9	161.6	161.5	167.2	163.8	163.4	167.5
CO <sub>2</sub> fossil in Sector 1B:							
1B1b. Solid Fuel Transf.	0.4	0.5	0.4	0.5	0.4	0.4	0.4
1B2c Flaring	0.4	0.3	0.2	0.1	0.1	0.1	0.0
1B2a-iv Oil refining	0.0	0.0	0.0	0.9	0.9	1.0	0.8
CO <sub>2</sub> fossil in Sector 2:							
A. Mineral Products	5.7	5.5	5.0	4.4	4.5	4.7	4.2
Soda Ash Production	0.1	0.1	0.1	0.1	0.1	0.1	0.1
B. Chemical industry							
1. Ammonia production	3.1	3.6	3.6	3.1	3.1	3.0	2.9
5. Other, excl. act. carbon	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	0.1	0.1	0.1	0.1	0.1	0.1	0.1
C. Metal industry							
1. Inputs in blast furnace	2.2	1.5	1.0	0.8	1.1	1.3	1.0
D. Other Production							
2. Food and Drink	0.1	0.0	0.0	0.0	0.0	0.0	0.0
G. Other (please specify)							
Other economic sectors **	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Not in NA-1A:	6.5	6.3	5.6	4.9	5.1	5.2	4.7
NA+1B+Ind. Proc.	156.4	167.8	167.1	172.1	168.9	168.6	172.2
RA+Fossil waste:	156.2	166.9	166.9	171.3	166.6	170.7	170.2
New difference (abs)	-0.2	-1.0	-0.2	-0.8	-2.4	2.1	-2.0
New difference (%)	-0.1%	-0.6%	-0.1%	-0.5%	-1.4%	1.2%	-1.2%

\*\* Comprises lubricants and waxes.

**Table A.4-3 Comparison of CO<sub>2</sub> emissions: differences between corrected Reference Approach (RA) versus corrected National Approach [(RA-NA)/NA] (in %)**

Fuel type *	1990	1995	2000	2005	2006	2007	2008
Liquids	-0.4%	-2.0%	-1.4%	-2.7%	-3.1%	3.6%	-1.6%
Solids	0.8%	0.2%	0.6%	2.2%	0.2%	2.0%	-0.9%
Gas	-0.4%	0.0%	0.5%	0.0%	-0.8%	-0.8%	-1.0%
Total (incl. waste)	-0.1%	-0.6%	-0.1%	-0.5%	-1.4%	1.2%	-1.2%

\* Liquids incl. 2G; Solids incl. 1B1,2A,2B5,2C1,2D; Gaseous incl. 1B2, 2B1; Total incl. fossil waste.

#### A4.4 Feedstock component in the CO<sub>2</sub> Reference Approach

Feedstock/non-energy uses of fuels in the energy statistics are also part of the IPCC Reference Approach for CO<sub>2</sub> from fossil fuel use. The fraction of carbon not oxidised during the use of these fuels during product manufacture or other uses is subtracted from total carbon contained in total apparent fuel consumption by fuel type. The fractions stored/oxidised have been calculated as three average values, one for gas, liquid and solid fossil fuels:  
 77.7±2% for liquid fuels;  
 55.5±13% for solid fuels;  
 38.8±4% for natural gas.

These were calculated from all processes for which emissions are calculated in the NA, either by assuming a fraction oxidised, for example ammonia, or by accounting for by-

product gases (excluding emissions from blast furnaces and coke ovens). (In Table A.4.4 of the NIR 2005, the calculation of annual oxidation fractions for 1995-2002 are presented and the average values derived from them.) It shows indeed that the factors show significant inter-annual variation, in particular for solid fuels.

The use of one average oxidation factor per fuel type for all years, whereas in the derivation of the annual oxidation figures differences up to a few per cent points can be observed, are one reason for differences between the RA and the corrected NA.

In Table A.4.4, the total CO<sub>2</sub> calculated as emitted from the oxidation of the non-energy uses in the Reference Approach is presented per fuel type. According to the Reference Approach dataset, the CO<sub>2</sub> emissions of this group of sources increased by about 7%, of which most are due to changes

**Trends in CO<sub>2</sub> emitted by feedstock use of energy carriers (production and direct uses) according to the correction term in the IPCC Reference Approach for CO<sub>2</sub> from fossil fuel use (in Tg CO<sub>2</sub>)**

**Table A4.4**

<b>Fuel type</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>Trend</b>
<i>Liquids</i> <sup>1)2)</sup>	5.0	3.2	3.7	5.4	4.5	6.9	6.2	1.2
<i>Solids</i> <sup>3)</sup>	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.1
<i>Gaseous</i>	3.5	3.8	3.9	3.5	3.0	3.2	2.9	-0.6
<i>Total</i>	8.7	7.2	7.8	9.1	7.7	10.3	9.3	0.6
<i>As% of RA</i>	5.6%	4.4%	4.7%	5.4%	4.7%	6.1%	5.6%	

1) Using country-specific carbon Oxidation Factors (multi-year average, fuel type averaged).

2) Excluding refineries.

3) Coal oils and tars (from coking coal), coke and other bituminous coal only; excluding emissions from blast furnaces and coke ovens.

**Carbon storage in the IPCC Reference Approach for CO<sub>2</sub> from fossil fuel use (in Tg CO<sub>2</sub>)**

**Table A4.5**

<b>Fuel type</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>Trend</b>
<i>Liquids</i>	17.4	18.0	21.3	30.5	28.1	34.9	31.9	14.5
<i>Solids</i>	0.6	0.5	0.5	0.5	0.4	0.6	0.7	0.1
<i>Gaseous</i>	2.2	2.4	2.5	2.2	1.9	2.0	1.8	-0.4
<i>Total</i>	20.2	21.0	24.4	33.3	30.5	37.5	34.4	14.2
<i>As% of RA</i>	11%	13%	19%	24%	24%	24%	22%	

Expressed as part of total carbon in apparent consumption of fossil fuels (without subtracting the stored part).

in emissions from liquid fuels. In Table A.4.4 and A4.5, the carbon storage in the RA calculation is shown. It shows that in the Netherlands about 10 to 25% of all carbon in the apparent consumption of fossil fuels is stored.



# Annex 5

## Assessment of completeness and (potential) sources and sinks

The Netherlands emissions inventory focuses on completeness and improving accuracy in the most relevant sources. This means that for all 'NE' sources, it is investigated what information is available and whether it could be assumed that a source is a really (very) small/negligible. For those sources that were not small, during the improvement programme, methods for estimating the emissions were developed.

As a result of this process, it was decided to keep only for very few sources as 'NE', since data for estimating emissions are not available and the source is very small. Of course, on regular basis it is being checked/re-assessed whether there are developments in NE sources that indicate (major) increase in emissions or new data sources for estimating emissions. For all except biofuels, this is the case for the 'NE' sources the ERT is referring to. For biofuels we are planning to incorporate activity data and emissions.

Following the 2009 review, several NE sources has been reviewed for the potential magnitude and the three potential biggest researched for estimating. As result lime production (2A2) is no longer included in this Annex, as these emissions were already included in the reported emissions of sugar production industries.

The Netherlands greenhouse gas emission inventory includes all sources identified by the Revised IPCC Guidelines (IPCC, 1996) – with the exception of the following (very) minor sources:

- Oil transport (1B2a3), due to missing activity data;
- The pipeline network in the Netherlands is dominated by the transport of natural gas (about 11,600 km in 2007). It is estimated that there is about 400 km pipelines for the transport of crude oil and about 1,000 km pipelines for the transport of oil products, chemical products and industrial gasses. Almost all pipelines are in the ground. Most of the pipelines for transport other than natural gas are short distance connections within industrial companies, compa-

nies close to each other and plant sites. For example, some pipelines in the Rotterdam area within the refinery sector; the emissions of those pipelines are included in the emission estimates as reported in 1A.

- There is no information available on the annual transport of the different types of fluids and also not on the material used for the pipelines.
- So it estimated that the emissions from oil transport by pipelines are a very minor source. Due to missing activity data as well as missing information on the material of the pipeline system, it is not possible to estimate this very minor source.
- Charcoal production (1B2), due to missing activity data;
- As indicated in the NIR 2008, there is information indicating that one company might produce charcoal. The production started after 1990. At this time, the production capacity is about 11,000 ton/year. Production levels are not known. The company is using the "Twin-retort" carbonisation process to produce charcoal. Compared to traditional charcoal production processes, CH<sub>4</sub> emissions from the Twin-retort system are far lower (Reumerman and Frederiks, 2002). CH<sub>4</sub> emissions from charcoal production are not estimated since these emissions are negligible.
- Charcoal use (1A4), due to missing activity data;
- Emissions from charcoal are only expected from barbecuing in the residential en services sector during the summer period. As indicated in the NIR 2008, there is no information on activity data on the actual charcoal use in 1A4 (other sectors), but it is assumed that these emissions are negligible. For these reasons the very low emissions are not estimated.
- CO<sub>2</sub> from asphalt roofing (2A5) and CO<sub>2</sub> from road paving (2A6), due to missing activity data;
- Information on the use of bitumen is only available for two groups: the chemical industry and all others. There is no information on the amount of asphalt roofing production and also no information on road paving with asphalt. The

statistical information on sales (value) of asphalt roofing and asphalt for road paving was finalised by 2002.

- Based on this information it was assumed that emissions related to this two categories are very low/undetectable and that effort in generating activity data would, therefore, not be cost effective. So not only the missing activity data but also the very limited amount of emissions was the rationale of the decision to not estimate these emissions.
- As follow up on the 2009 review, information has been collected from the branch organisation of roofing indicating that the number of producers of asphalt roofing declined from about 15 in 1990 to less than 5 in 2008 and that the import of asphalt roofing increased. Also, information has been researched on asphalt production (for road paving) as reported in the progress of the voluntary agreements for energy efficiency. A first estimate indicates that the CO<sub>2</sub> emissions could be around 0.5 kton.
- CH<sub>4</sub> from Enteric fermentation poultry (4A9), due to missing emission factors;
- For this source category no IPCC default emission factor is available
- N<sub>2</sub>O from Industrial wastewater (6B1), due to negligible amounts.
- As presented in the NIR 2008, page 194, the annual source for activity data are yearly questionnaires which cover all urban WWTPs and all anaerobic industrial WWTPs. For these industrial WWTPs, CH<sub>4</sub> emissions are estimated based on the design capacity of the installations (47 plants) and reported. CH<sub>4</sub> emissions reported for 2006 were 0.33 Gg CH<sub>4</sub>. From this anaerobic pre-treatment, there is no N<sub>2</sub>O emission
- In 2000, the Netherlands investigated not previously estimated sources for non-CO<sub>2</sub> emissions. One of these sources was the wastewater handling (DHV, 2000). As a result of this study, emissions were estimated (Oonk, 2004) and the methods are presented in the protocols CH<sub>4</sub>, N<sub>2</sub>O from wastewater treatment (6B).
- We are not able to estimate N<sub>2</sub>O emissions from aerobic industrial WWTPs, as there is no information available on these installations. In the priority setting for allocation of budgets for improvements in emission estimates, we did consider this as a source for which it could not be argued that a new data collection process or a new statistic was a priority. The argumentations for this decision includes following:
- The majority of the small and medium enterprises are linked to the municipal wastewater treatment plants (for which we made emission estimates) and do not have an their own wastewater treatment;
  - The anaerobic pre-treatment reduces the N load to the aerobic final treatment;
  - Aerobic (post) treatment is done for several of the industrial companies in the municipal WWTPs;
  - The composition of the industrial wastewater is mainly process water and although we have no specific information on the N-content of the influent, it is assumed that it is low N content.

In addition, there are indications that the number of industrial wastewater treatment plants will reduce in the near

future and this will also further minimise the minor effect of not estimating this source.

- Precursor emissions (i.e. CO, NO<sub>x</sub>, NMVOC and SO<sub>2</sub>) from Memo item international bunkers (international transport) have not been included.

# Annex 6

## Additional information to be considered as part of the NIR submission

The following information should be considered as part of this NIR submission:

### A6.1 List of protocols

Methodological description (monitoring protocols 2010, from 15 April 2010 available at the website).

Table A6.1

Protocol	IPCC code	Description	Gases
0041	All	Reference approach	CO <sub>2</sub>
0042	1A1 1A2 1A4	Stationary combustion (fossil) *	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0043	1A1b 1B1b 1B2aiv 2A4i 2B1 2B4i 2B5i 2B5vii 2B5viii 2C1vi 2D2 2Giv	Process emissions (fossil)	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0044	1A2f 1A4c	Mobile equipment	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0045	1A3a	Inland aviation	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0046	1A3b	Road transport	CO <sub>2</sub>
0047	1A3b	Road transport	N <sub>2</sub> O CH <sub>4</sub>
0048	1A3c	Rail transport	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0049	1A3d	Inland navigation	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0050	1A4c	Fisheries	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0051	1A5	Defence	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0052	1B2	Oil & gas production	CO <sub>2</sub> CH <sub>4</sub>
0053	1B2	Oil & gas distribution/transport	CO <sub>2</sub> CH <sub>4</sub>
0054	2A1 2A2 2A3 2A4ii 2A7i 2B5ix 2C1i 2C1vii 2C3 2Gi 2Gii 2Giii 2Gv 3A 3B 3C 3D	Process emissions (non-fossil)	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0055	2B2	Nitric acid	N <sub>2</sub> O
0056	2B5	Caprolactam	N <sub>2</sub> O
0057	2C3	Aluminium production	PFC
0058	2E1	HCFC-22 production	HFC
0059	2E3	HFC byproduct emissions	HFC
0060	2F1	Stationary refrigeration	HFC
0061	2F1	Mobile refrigeration	HFC
0062	2F2	Hard foams	HFC
0063	2F4	Aerosols	HFC
0064	2F8	Sound proof windows	SF <sub>6</sub>
0065	2F8	Semi-conductors	SF <sub>6</sub> PFC
0066	2F8	Electrical equipment	SF <sub>6</sub>
0067	4A	Enteric fermentation	CH <sub>4</sub>
0068	4B	Manure management	N <sub>2</sub> O
0069	4B	Manure management	CH <sub>4</sub>
0070	4D	Agricultural soils, indirect	N <sub>2</sub> O
0071	4D	Agricultural soils, direct	N <sub>2</sub> O
0072	5A	Forest	CO <sub>2</sub>
0073	5D-5G	Soil	CO <sub>2</sub>
0074	6A1	Waste disposal	CH <sub>4</sub>
0075	6B	Waste water treatment	CH <sub>4</sub> N <sub>2</sub> O
0076	6D	Large-scale composting	CH <sub>4</sub> N <sub>2</sub> O
<i>In addition to the emissions described in the protocols, two memo items are included in the National System</i>			
0077	Memo item	International bunker emissions	CO <sub>2</sub> N <sub>2</sub> O CH <sub>4</sub>
0078	1A, (CO <sub>2</sub> memo item)	Biomass	CO <sub>2</sub> CH <sub>4</sub> N <sub>2</sub> O

#### A6.2 Documentation of uncertainties used in IPCC Tier 1 uncertainty assessments and Tier 2 key source identification

- Olivier, J.G.J., L.J. Brandes, R.A.B. te Molder, 2009: Estimate of annual and trend uncertainty for Dutch sources of greenhouse gas emissions using the IPCC Tier 1 approach. PBL Report 500080013, PBL, Bilthoven
- Olsthoorn, X. and A. Pielaat, 2003: Tier-2 uncertainty analysis of the Dutch greenhouse gas emissions 1999. Institute for Environmental Studies (IVM), Free University, Amsterdam. IVM Report no. R03-06.
- Ramírez-Ramírez, A., C. de Keizer and J.P. van der Sluijs, 2006: Monte Carlo Analysis of Uncertainties in the Netherlands Greenhouse Gas Emission Inventory for 1990–2004, report NWS-E-2006-58, Department of Science, Technology and Society, Copernicus Institute for Sustainable Development and Innovation, Utrecht University, Utrecht, the Netherlands.

#### A6.3 Background documents and uncertainty discussion papers

- Van Amstel, A.R., J.G.J. Olivier and P.G. Ruysenaars (eds.), 2000a: Monitoring of Greenhouse Gases in the Netherlands: Uncertainty and Priorities for Improvement. Proceedings of a National Workshop held in Bilthoven, the Netherlands, 1 September 1999. WIMEK report/RIVM report no. 773201 003. Bilthoven, May 2000.
- Kuikman, P.J., J.J.H. van den Akker and F. de Vries, 2005: Lachgasemissie uit organische landbouwbodems. Alterra, Wageningen. Alterra rapport 1035-II.
- Hoek, K. W. van der, 2002: Uitgangspunten voor de mest- en ammoniakberekeningen 1999 tot en met 2001 zoals gebruikt in de Milieubalans 2001 en 2002, inclusief dataset landbouwemissies 1980-2001. RIVM rapport 773004013. RIVM, Bilthoven.
- Hoek, K. W. van der and M. W. van Schijndel, 2006: Methane and nitrous oxide emissions from animal manure management, including an overview of emissions 1990-2003. Background document for the Dutch National Inventory Report. RIVM report 680.125.002, Bilthoven.
- Hoek, K.W. van der, M.W. van Schijndel, P.J. Kuikman, 2007. Direct and indirect nitrous oxide emissions from agricultural soils, 1990 - 2003. Background document on the calculation method for the Dutch National Inventory Report. RIVM Report No. 68012.003./2007 MNP Report No. 500080003/2007 Bilthoven, the Netherlands.
- Nabuurs, G.J., I.J. van den Wyngaert, W.D. Daamen, A.T.F. Helmink, W de Groot, W.C. Knol, H. Kramer, P Kuikman, 2005: National System of Greenhouse Gas Reporting for Forest and Nature Areas under UNFCCC in the Netherlands - version 1.0 for 1990–2002. Alterra, Wageningen. Alterra rapport 1035-I.
- Van den Wyngaert, I.J.J., Kramer, H., Kuikman, P., Nabuurs, G.J. (2009) Greenhouse gas reporting of the LULUCF sector, revisions and updates related to the Dutch NIR 2009. Alterra report 1035.7, Alterra, Wageningen.

#### A6.4 Documentation of Quality Assurance and Quality Control for national greenhouse gas inventory compilation and reporting

- DHV, 2002: Quality Assurance and Quality Control for the Dutch National Inventory Report; report on phase 1, January 2002, report no. ML-BB-20010367. DHV, Amersfoort.
- PBL, 2008. Werkplan EmissieRegistratie ronde 2009 – 2010. PBL, Bilthoven, 2009.
- SenterNovem, the Netherlands National System:QA/QC programme 2009/2010 Version 5.0 December 2009.

#### A6.5 Documentation of Changes to the National Registry

- Release Notes, Greta Emissions Trading Registry Version 4.0.14, Date: 30th March 2009, SFW Limited, Southern House Station Approach Woking Surrey GU22 7UY
- Release Notes, Greta Emissions Trading Registry Version 4.1.16, Date: 18th June 2009, SFW Limited, Southern House Station Approach Woking Surrey GU22 7UY
- Release Notes, Greta Emissions Trading Registry Version 4.2.21, Date: 6th October 2009, SFW Limited, Southern House Station Approach Woking Surrey GU22 7UY
- Test report, Greta Registry Release V4.1, QME201 Version 2.0, Date of creation: 20 Jul 2009, SFW Limited, Southern House Station Approach Woking Surrey GU22 7UY
- Test report, Greta Registry Release V4.2, QME201 Version 2.0, Date of creation: 11 Nov 2009, SFW Limited, Southern House Station Approach Woking Surrey GU22 7UY
- ETS Testing Plan, EUROPEAN COMMISSION, DG ENVIRONMENT, Directorate C - Climate Change & Air, ENV.C.2 - Market based instruments including Greenhouse gas emissions trading, Brussels, ENV C2/
- Test approval of a E new reconciliation for NL in the Registry Test Environment, E-mail Registry Test Environment. Application manager for the CITL, 17th June 2009

## Registry Information

### Report R1

Total quantities of Kyoto Protocol units by account type at beginning of reported year

Table 1

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	924505409	NO	NO	7230375	NO	NO
Entity holding accounts	80858465	NO	NO	9280513	NO	NO
Article 3.3/3.4 net source cancellation accounts	NO	NO	NO	NO		
Non-compliance cancellation accounts	NO	NO	NO	NO		
Other cancellation accounts	NO	NO	NO	NO	NO	NO
Retirement account	NO	NO	NO	NO	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
<b>Total</b>	1005363874	NO	NO	16510888	NO	NO

Annual internal transactions

Table 2a

Transaction type	Additions						Subtractions					
	Unit type						Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
<b>Article 6 issuance and conversion</b>												
Party-verified projects		NO					NO		NO			
Independently verified projects		NO					NO		NO			
<b>Article 3.3 and 3.4 issuance or cancellation</b>												
3.3 Afforestation and reforestation			NO				NO	NO	NO	NO		
3.3 Deforestation			NO				NO	NO	NO	NO		
3.4 Forest management			NO				NO	NO	NO	NO		
3.4 Cropland management			NO				NO	NO	NO	NO		
3.4 Grazing land management			NO				NO	NO	NO	NO		
3.4 Revegetation			NO				NO	NO	NO	NO		
<b>Article 12 afforestation and reforestation</b>												
Replacement of expired tCERs							NO	NO	NO	NO	NO	
Replacement of expired ICERs							NO	NO	NO	NO		
Replacement for reversal of storage							NO	NO	NO	NO		NO
Replacement for non-submission of certification report							NO	NO	NO	NO		NO
<b>Other cancellation</b>							NO	NO	NO	20	NO	NO
<b>Sub-total</b>		NO	NO				NO	NO	NO	20	NO	NO

Transaction type	Retirement					
	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
<b>Retirement</b>	83512630	NO	NO	NO	NO	NO

	Additions						Subtractions					
	Unit type						Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
<b>Transfers and acquisitions</b>												
CDM	NO	NO	NO	5254583	NO	NO	NO	NO	NO	NO	NO	NO
AT	3109151	NO	NO	56216	NO	NO	612757	3000	NO	271500	NO	NO
BE	3077228	108750	NO	71520	NO	NO	8983451	NO	NO	780576	NO	NO
CZ	3216268	100369	NO	44000	NO	NO	6444347	NO	NO	323000	NO	NO
DK	50686258	NO	NO	4029468	NO	NO	35530872	1000	NO	21808528	NO	NO
EE	73864	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
FI	142303	NO	NO	NO	NO	NO	37000	NO	NO	211500	NO	NO
FR	47298959	NO	NO	2877810	NO	NO	60443543	NO	NO	3496254	NO	NO
DE	40125559	111100	NO	1552666	NO	NO	38312596	302950	NO	10375617	NO	NO
GR	986000	NO	NO	15000	NO	NO	36000	NO	NO	379000	NO	NO
HU	1389249	76730	NO	NO	NO	NO	NO	NO	NO	45769	NO	NO
IE	NO	NO	NO	NO	NO	NO	110600	NO	NO	534306	NO	NO
IT	1174680	NO	NO	39000	NO	NO	686521	NO	NO	442140	NO	NO
JP	NO	NO	NO	NO	NO	NO	NO	NO	NO	93227	NO	NO
LV	3526580	NO	NO	NO	NO	NO	84	NO	NO	NO	NO	NO
LT	1626300	NO	NO	1	NO	NO	800000	NO	NO	8001	NO	NO
LU	13548	NO	NO	NO	NO	NO	322000	NO	NO	12548	NO	NO
NZ	NO	240000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
NO	78957	NO	NO	NO	NO	NO	101202	NO	NO	4000	NO	NO
PL	7884512	NO	NO	1	NO	NO	1381000	NO	NO	18501	NO	NO
PT	494410	NO	NO	198000	NO	NO	629000	NO	NO	1304853	NO	NO
SK	241289	NO	NO	NO	NO	NO	329000	NO	NO	NO	NO	NO
SI	253416	NO	NO	NO	NO	NO	60000	NO	NO	72564	NO	NO
ES	999398	NO	NO	1232513	NO	NO	527482	NO	NO	1093587	NO	NO
SE	232800	NO	NO	355163	NO	NO	265000	NO	NO	707055	NO	NO
CH	NO	63800	NO	29122914	NO	NO	NO	50950	NO	12417442	NO	NO
UA	500000	650109	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
GB	41938096	50000	NO	28381431	NO	NO	47045148	5750	NO	18100070	NO	NO
<b>Sub-total</b>	<b>2.09E+08</b>	<b>1400858</b>	<b>NO</b>	<b>73230286</b>	<b>NO</b>	<b>NO</b>	<b>202657603</b>	<b>363650</b>	<b>NO</b>	<b>72500038</b>	<b>NO</b>	<b>NO</b>

## Additional information

Independently verified ERUs								NO				
-----------------------------	--	--	--	--	--	--	--	----	--	--	--	--

## Total annual transactions

## Table 2c

<b>Total (Sum of tables 2a and 2b)</b>	<b>2.09E+08</b>	<b>1400858</b>	<b>NO</b>	<b>73230286</b>	<b>NO</b>	<b>NO</b>	<b>202657603</b>	<b>363650</b>	<b>NO</b>	<b>72500058</b>	<b>NO</b>	<b>NO</b>
--	-----------------	----------------	-----------	-----------------	-----------	-----------	------------------	---------------	-----------	-----------------	-----------	-----------

## Expiry, cancellation and replacement

## Table 3

Transaction or event type	Expiry, cancellation and requirement to replace		Replacement					
			Unit type					
	Unit type		Unit type					
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
<b>Temporary CERs (tCERs)</b>								
Expired in retirement and replacement accounts	NO							
Replacement of expired tCERs			NO	NO	NO	NO	NO	
Expired in holding accounts	NO							
Cancellation of tCERs expired in holding accounts	NO							
<b>Long-term CERs (ICERs)</b>								
Expired in retirement and replacement accounts		NO						
Replacement of expired ICERs			NO	NO	NO	NO		
Expired in holding accounts		NO						
Cancellation of ICERs expired in holding accounts		NO						
Subject to replacement for reversal of storage		NO						
Replacement for reversal of storage			NO	NO	NO	NO		NO
Subject to replacement for non-submission of certification report		NO						
Replacement for non-submission of certification report			NO	NO	NO	NO		NO
<b>Total</b>			NO	NO	NO	NO	NO	NO

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	842767016	917099	NO	12458380	NO	NO
Entity holding accounts	85495450	120109	NO	4782736	NO	NO
Article 3.3/3.4 net source cancellation accounts	NO	NO	NO	NO		
Non-compliance cancellation accounts	NO	NO	NO	NO		
Other cancellation accounts	NO	NO	NO	20	NO	NO
Retirement account	83512630	NO	NO	NO	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
<b>Total</b>	<b>1011775096</b>	<b>1037208</b>	<b>NO</b>	<b>17241136</b>	<b>NO</b>	<b>NO</b>

## Summary information on additions and subtractions

Table 5a

	Additions						Subtractions					
	Unit type						Unit type					
Starting values	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Issuance pursuant to Article 3.7 and 3.8	1001262141											
Non-compliance cancellation							NO	NO	NO	NO		
Carry-over	NO	NO		NO								
<b>Sub-total</b>	<b>1001262141</b>	<b>NO</b>	<b></b>	<b>NO</b>	<b></b>	<b></b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b></b>	<b></b>
<b>Annual transactions</b>												
Year 0 (2007)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 1 (2008)	87571284	NO	NO	39222701	NO	NO	83469551	NO	NO	22711813	NO	NO
Year 2 (2009)	209068825	1400858	NO	73230286	NO	NO	202657603	363650	NO	72500058	NO	NO
Year 3 (2010)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 4 (2011)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 5 (2012)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 6 (2013)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<b>Sub-total</b>	<b>296640109</b>	<b>1400858</b>	<b>NO</b>	<b>112452987</b>	<b>NO</b>	<b>NO</b>	<b>286127154</b>	<b>363650</b>	<b>NO</b>	<b>95211871</b>	<b>NO</b>	<b>NO</b>
<b>Total</b>	<b>1297902250</b>	<b>1400858</b>	<b>NO</b>	<b>112452987</b>	<b>NO</b>	<b>NO</b>	<b>286127154</b>	<b>363650</b>	<b>NO</b>	<b>95211871</b>	<b>NO</b>	<b>NO</b>

## Summary information on replacement

Table 5b

	Requirement for replacement		Replacement						
	Unit type		Unit type						
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	
<b>Previous CPs</b>			NO	NO	NO	NO	NO	NO	
Year 1 (2008)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 2 (2009)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 3 (2010)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 4 (2011)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 5 (2012)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 6 (2013)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 7 (2014)	NO	NO	NO	NO	NO	NO	NO	NO	
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO	
<b>Total</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	

## Summary information on retirement

Table 5c

Year	Retirement					
	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Year 1 (2008)	NO	NO	NO	NO	NO	NO
Year 2 (2009)	83512630	NO	NO	NO	NO	NO
Year 3 (2010)	NO	NO	NO	NO	NO	NO
Year 4 (2011)	NO	NO	NO	NO	NO	NO
Year 5 (2012)	NO	NO	NO	NO	NO	NO
Year 6 (2013)	NO	NO	NO	NO	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO
<b>Total</b>	<b>83512630</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>



	Additions						Subtractions					
	Unit type						Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs

	Requirement for replacement		Replacement					
	Unit type		Unit type					
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs

	Retirement					
	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs

## Report R2

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
4003	811.58	221.57	NL-8974	23/01/2009 09:45:39	External Transfer	Terminated			
							DE-3443213489-3443386093	AAU	172605
							DE-3445831381-3445849775	AAU	18395
			NL-8975	23/01/2009 09:45:39	External Transfer	Terminated	GR-284197997-284506996	AAU	309000
							DE-3443329489-3443386093	AAU	56605
							DE-3445831381-3445849775	AAU	18395
			NL-9661	10/02/2009 11:49:00	Internal Transfer	Terminated			
							DE-1753584569-1753589968	AAU	5400
							DE-3566072952-3566074234	AAU	1283
							DE-3566103144-3566109143	AAU	6000
							DE-3566109144-3566109951	AAU	808
							DE-3566124952-3566127951	AAU	3000
							DE-3566127952-3566132951	AAU	5000
							DE-3566151156-3566152155	AAU	1000
							DE-3566161952-3566168843	AAU	6892
							DE-3566171520-3566171536	AAU	17
							DE-3568772800-3568776715	AAU	3916
							DE-3568796716-3568802788	AAU	6073
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
			NL-9669	10/02/2009 12:03:47	Internal Transfer	Terminated			
							DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428837163-428839162	AAU	2000
							SE-428843478-428853162	AAU	9685
							SE-428853163-428855477	AAU	2315
							SE-428855478-428855717	AAU	240
							SK-428695421-428695459	AAU	39
							SK-428695460-428701397	AAU	5938
							SK-428701398-428703677	AAU	2280
							SK-428703678-428704439	AAU	762
							SK-428704440-428705201	AAU	762
							SK-428705202-428710439	AAU	5238
							SK-433949830-433952129	AAU	2300
			NL-9670	10/02/2009 12:06:50	Internal Transfer	Terminated			

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
							DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428848459-428853162	AAU	4704
							SE-428853163-428855477	AAU	2315
							SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
			NL-9671	10/02/2009 12:12:09	Internal Transfer	Terminated			
							DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428855459-428855477	AAU	19
							SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
			NL-9672	10/02/2009 12:19:07	External Transfer	Terminated			
							DE-3569316883-3569317882	AAU	1000
			NL-9673	10/02/2009 12:24:40	Internal Transfer	Terminated			
							DE-3569316883-3569317882	AAU	1000
			NL-9674	10/02/2009 13:02:26	Internal Transfer	Terminated			
							DE-3568916327-3568916715	AAU	389
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569316882	AAU	5000
							DE-3569316883-3569317882	AAU	1000
			NL-9675	10/02/2009 13:40:15	Internal Transfer	Terminated			
							DE-3569316883-3569317882	AAU	1000
			NL-9678	10/02/2009 15:04:03	Internal Transfer	Terminated			
							DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568914716-3568916326	AAU	1611
							DE-3568916327-3568916715	AAU	389
							DE-3569311883-3569316882	AAU	5000
							DE-3569316883-3569317882	AAU	1000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-9682	10/02/2009 15:36:05	Internal Transfer	Terminated	IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428843478-428848458	AAU	4981
							SE-428848459-428853162	AAU	4704
							SE-428853163-428855458	AAU	2296
							SE-428855459-428855477	AAU	19
							SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
							DE-1720088581-1720090580	AAU	2000
							DE-3442879853-3442880527	AAU	675
							DK-1704299017-1704299695	AAU	679
							DK-1704299696-1704301526	AAU	1831
							DK-1705939086-1705939191	AAU	106
							DK-1705945750-1705947154	AAU	1405
							EE-243253544-243255867	AAU	2324
							ES-1710213556-1710218555	AAU	5000
							ES-1718885201-1718885504	AAU	304
							FR-3308269838-3308272767	AAU	2930
							FR-3308273600-3308275599	AAU	2000
							FR-3308280600-3308284669	AAU	4070
							FR-3308284670-3308285669	AAU	1000
							IT-3027183833-3027200832	AAU	17000
							RO-2564396638-2564400742	AAU	4105
							RO-2565135067-2565138066	AAU	3000
							RO-2565138067-2565139066	AAU	1000
							RO-2565144067-2565148066	AAU	4000
							RO-2603777359-2603777709	AAU	351
							RO-2603840590-2603842500	AAU	1911
							RO-2603845590-2603845811	AAU	222
							RO-2603845812-2603847222	AAU	1411
			NL-9683	10/02/2009 15:53:22	Internal Transfer	Terminated	DE-3566103144-3566109143	AAU	6000
							DE-3566124952-3566127951	AAU	3000
							DE-3566127952-3566132951	AAU	5000
							DE-3566151156-3566152155	AAU	1000
							DE-3568773716-3568776715	AAU	3000
			NL-11323	26/05/2009 12:28:37	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
			NL-11324	26/05/2009 12:30:06	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
			NL-11325	26/05/2009 12:31:06	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000

[illegible]

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
							FR-3267254582-3267254609	AAU	28
							FR-3285642694-3285642824	AAU	131
							FR-3288455693-3288457648	AAU	1956
							FR-3288462267-3288462351	AAU	85
							FR-3289102635-3289105634	AAU	3000
							FR-3289300983-3289301087	AAU	105
							FR-3289534974-3289536329	AAU	1356
							FR-3289536330-3289537422	AAU	1093
							FR-3289537423-3289537634	AAU	212
							FR-3289545812-3289546077	AAU	266
							FR-3289558635-3289560479	AAU	1845
							FR-3293753383-3293753996	AAU	614
							FR-3294835946-3294844945	AAU	9000
							FR-3301197816-3301202815	AAU	5000
							FR-3301202816-3301203815	AAU	1000
							FR-3301403112-3301407111	AAU	4000
							FR-3302599112-3302601111	AAU	2000
							FR-3302602112-3302604111	AAU	2000
							FR-3303636408-3303641407	AAU	5000
							FR-3303668573-3303669572	AAU	1000
							PL-3249253579-3249254324	AAU	746
			NL-12793	01/12/2009 10:13:30	External Transfer	Terminated			
							GB-3892021446-3892021505	AAU	60
							GB-3901447620-3901447862	AAU	243
							GB-3901449666-3901449780	AAU	115
							GB-3901547792-3901548226	AAU	435
							GB-3910430963-3910431551	AAU	589
							GB-3913551143-3913551557	AAU	415
							GB-3976982109-3976982922	AAU	814
							GB-3980733265-3980733838	AAU	574
							GB-3986931627-3986932189	AAU	563
							GB-3987027190-3987027744	AAU	555
							GB-3991549420-3991550234	AAU	815
							GB-3992440284-3992441110	AAU	827
							GB-3993039216-3993040044	AAU	829
							GB-3993068216-3993068776	AAU	561
							GB-3995247246-3995247799	AAU	554
							GB-3995310094-3995310652	AAU	559
							GB-3995362725-3995363283	AAU	559
							GB-3995707164-3995707977	AAU	814
							GB-3995847185-3995848008	AAU	824
							GB-3996429812-3996430377	AAU	566
							GB-3999250712-3999251303	AAU	592
							GB-4000404168-4000404217	AAU	50
							GB-4000478042-4000478637	AAU	596
							GB-4002378797-4002379394	AAU	598
							GB-4002397035-4002397353	AAU	319
							GB-4003815679-4003816252	AAU	574
			NL-12794	01/12/2009 10:15:48	External Transfer	Terminated			
							DE-1643914844-1643922166	AAU	7323
							DE-2135936907-2135944583	AAU	7677
							DE-3441941939-3441942591	AAU	653
							DE-3442682583-3442682929	AAU	347
							DE-3558925244-3558926243	AAU	1000

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
							DE-3588697472-3588698471	AAU	1000
							ES-1829735803-1829742368	AAU	6566
							ES-1829742369-1829753368	AAU	11000
							ES-1829753369-1829755802	AAU	2434
							GB-3764984053-3764986052	AAU	2000
							GB-4000051622-4000086754	AAU	35133
							GB-4000374755-4000395962	AAU	21208
							GB-4000547359-4000569257	AAU	21899
							PL-3186805746-3186807505	AAU	1760
			NL-12796	01/12/2009 10:19:10	External Transfer	Terminated			
							ES-1829671153-1829679368	AAU	8216
							ES-1829679369-1829683368	AAU	4000
							ES-1829717369-1829718845	AAU	1477
							ES-1829718846-1829721650	AAU	2805
							ES-1829727301-1829727368	AAU	68
							ES-1829727369-1829735802	AAU	8434
			NL-12801	01/12/2009 11:14:20	External Transfer	Terminated			
							AT-1585046206-1585048205	AAU	2000
							AT-1585048206-1585055205	AAU	7000
							AT-1585070056-1585071055	AAU	1000
			NL-12803	01/12/2009 11:16:47	Internal Transfer	Terminated			
							AT-1557053298-1557057381	AAU	4084
							AT-1570623324-1570646104	AAU	22781
							AT-1574166444-1574169163	AAU	2720
							AT-1574212740-1574214922	AAU	2183
							AT-1583303601-1583304573	AAU	973
							AT-1583304950-1583307073	AAU	2124
							AT-1584137537-1584146671	AAU	9135
							AT-1585040206-1585046205	AAU	6000
			NL-12806	01/12/2009 11:20:54	External Transfer	Terminated			
							AT-1557053298-1557057381	AAU	4084
							AT-1570623324-1570646104	AAU	22781
							AT-1574166444-1574169163	AAU	2720
							AT-1574212740-1574214922	AAU	2183
							AT-1583303601-1583304573	AAU	973
							AT-1583304950-1583307073	AAU	2124
							AT-1584137537-1584146671	AAU	9135
							AT-1585040206-1585046205	AAU	6000
							GB-3686527678-3686547677	AAU	20000
			NL-12811	01/12/2009 11:24:42	External Transfer	Terminated			
							AT-1584142672-1584146671	AAU	4000
							AT-1585040206-1585046205	AAU	6000
			NL-12947	07/12/2009 16:13:08	External Transfer	Terminated			
							CN-86986984-86987861	CER	878
							CN-86989121-86990379	CER	1259
			NL-12949	07/12/2009 16:16:09	External Transfer	Terminated			
							CN-86986984-86987861	CER	878
							CN-86989121-86990379	CER	1259
			NL-12950	07/12/2009 16:19:09	External Transfer	Terminated			
							CN-86986984-86987861	CER	878
							CN-86989121-86990379	CER	1259



DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-12951	07/12/2009 16:22:29	External Transfer	Terminated			
							CN-86986984-86987861	CER	878
							CN-86989121-86990379	CER	1259
			NL-12952	07/12/2009 16:25:41	External Transfer	Terminated			
							CN-86986984-86987861	CER	878
							CN-86989121-86990379	CER	1259
			NL-13067	16/12/2009 13:09:26	Internal Transfer	Terminated			
							BR-22518777-22531294	CER	12518
			NL-13068	16/12/2009 13:34:34	Internal Transfer	Terminated			
							BR-22518777-22531294	CER	12518
			NL-13069	16/12/2009 14:24:05	Internal Transfer	Terminated			
							CZ-1322159770-1322160390	AAU	621
							CZ-1322160391-1322164304	AAU	3914
							CZ-1322461352-1322465351	AAU	4000
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383
			NL-13070	16/12/2009 14:40:34	Internal Transfer	Terminated			
							CZ-1322159770-1322160390	AAU	621
							CZ-1322160391-1322164304	AAU	3914
							CZ-1322461352-1322465351	AAU	4000
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383
			NL-13077	17/12/2009 10:35:22	Internal Transfer	Terminated			
							CZ-1322464333-1322465351	AAU	1019
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383
4010	811.58	147.71							

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-8974	23/01/2009 09:45:39	External Transfer	Terminated	DE-3443213489-3443386093	AAU	172605
							DE-3445831381-3445849775	AAU	18395
							GR-284197997-284506996	AAU	309000
			NL-9661	10/02/2009 11:49:00	Internal Transfer	Terminated	DE-1753584569-1753589968	AAU	5400
							DE-3566072952-3566074234	AAU	1283
							DE-3566103144-3566109143	AAU	6000
							DE-3566109144-3566109951	AAU	808
							DE-3566124952-3566127951	AAU	3000
							DE-3566127952-3566132951	AAU	5000
							DE-3566151156-3566152155	AAU	1000
							DE-3566161952-3566168843	AAU	6892
							DE-3566171520-3566171536	AAU	17
							DE-3568772800-3568776715	AAU	3916
							DE-3568796716-3568802788	AAU	6073
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
			NL-9669	10/02/2009 12:03:47	Internal Transfer	Terminated	DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428837163-428839162	AAU	2000
							SE-428843478-428853162	AAU	9685
							SE-428853163-428855477	AAU	2315
							SE-428855478-428855717	AAU	240
							SK-428695421-428695459	AAU	39
							SK-428695460-428701397	AAU	5938
							SK-428701398-428703677	AAU	2280
							SK-428703678-428704439	AAU	762
							SK-428704440-428705201	AAU	762
							SK-428705202-428710439	AAU	5238
							SK-433949830-433952129	AAU	2300
			NL-9670	10/02/2009 12:06:50	Internal Transfer	Terminated	DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-9671	10/02/2009 12:12:09	Internal Transfer	Terminated	DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428848459-428853162	AAU	4704
							SE-428853163-428855477	AAU	2315
							SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
							DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568802789-3568805266	AAU	2478
							DE-3568805267-3568805788	AAU	522
							DE-3568838716-3568840715	AAU	2000
							DE-3568914716-3568916715	AAU	2000
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569317882	AAU	6000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428855459-428855477	AAU	19
							SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
			NL-9672	10/02/2009 12:19:07	External Transfer	Terminated	DE-3569316883-3569317882	AAU	1000
			NL-9673	10/02/2009 12:24:40	Internal Transfer	Terminated	DE-3569316883-3569317882	AAU	1000
			NL-9674	10/02/2009 13:02:26	Internal Transfer	Terminated	DE-3568916327-3568916715	AAU	389
							DE-3568920075-3568922685	AAU	2611
							DE-3569311883-3569316882	AAU	5000
							DE-3569316883-3569317882	AAU	1000
			NL-9675	10/02/2009 13:40:15	Internal Transfer	Terminated	DE-3569316883-3569317882	AAU	1000
			NL-9678	10/02/2009 15:04:03	Internal Transfer	Terminated	DE-3566156458-3566159674	AAU	3217
							DE-3568799400-3568802788	AAU	3389
							DE-3568914716-3568916326	AAU	1611
							DE-3568916327-3568916715	AAU	389
							DE-3569311883-3569316882	AAU	5000
							DE-3569316883-3569317882	AAU	1000
							ES-1829724350-1829727300	AAU	2951
							FR-3308272768-3308273599	AAU	832
							IT-2916908402-2916910842	AAU	2441
							RO-2568225422-2568235421	AAU	10000
							SE-428843478-428848458	AAU	4981
							SE-428848459-428853162	AAU	4704
							SE-428853163-428855458	AAU	2296
							SE-428855459-428855477	AAU	19

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-9682	10/02/2009 15:36:05	Internal Transfer	Terminated	SE-428855478-428855717	AAU	240
							SK-433949830-433952129	AAU	2300
							DE-1720088581-1720090580	AAU	2000
							DE-3442879853-3442880527	AAU	675
							DK-1704299017-1704299695	AAU	679
							DK-1704299696-1704301526	AAU	1831
							DK-1705939086-1705939191	AAU	106
							DK-1705945750-1705947154	AAU	1405
							EE-243253544-243255867	AAU	2324
							ES-1710213556-1710218555	AAU	5000
							ES-1718885201-1718885504	AAU	304
							FR-3308269838-3308272767	AAU	2930
							FR-3308273600-3308275599	AAU	2000
							FR-3308280600-3308284669	AAU	4070
							FR-3308284670-3308285669	AAU	1000
							IT-3027183833-3027200832	AAU	17000
							RO-2564396638-2564400742	AAU	4105
							RO-2565135067-2565138066	AAU	3000
							RO-2565138067-2565139066	AAU	1000
							RO-2565144067-2565148066	AAU	4000
			RO-2603777359-2603777709	AAU	351				
			RO-2603840590-2603842500	AAU	1911				
			RO-2603845590-2603845811	AAU	222				
			RO-2603845812-2603847222	AAU	1411				
			NL-9683	10/02/2009 15:53:22	Internal Transfer	Terminated	DE-3566103144-3566109143	AAU	6000
							DE-3566124952-3566127951	AAU	3000
							DE-3566127952-3566132951	AAU	5000
							DE-3566151156-3566152155	AAU	1000
							DE-3568773716-3568776715	AAU	3000
			NL-11323	26/05/2009 12:28:37	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
			NL-11324	26/05/2009 12:30:06	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
NL-11325	26/05/2009 12:31:06	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677			
				GB-3701126446-3701127768	AAU	1323			
				GB-3703857688-3703858687	AAU	1000			
				GB-3910245949-3910270948	AAU	25000			
				GB-3998247363-3998251362	AAU	4000			
NL-11326	26/05/2009 12:31:55	External Transfer	Terminated	DE-3597161354-3597180030	AAU	18677			
				GB-3701126446-3701127768	AAU	1323			
				GB-3703857688-3703858687	AAU	1000			
				GB-3910245949-3910270948	AAU	25000			
				GB-3998247363-3998251362	AAU	4000			
							DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-11327	26/05/2009 12:32:48	External Transfer	Cancelled	GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
							DE-3597161354-3597180030	AAU	18677
							GB-3701126446-3701127768	AAU	1323
							GB-3703857688-3703858687	AAU	1000
							GB-3910245949-3910270948	AAU	25000
							GB-3998247363-3998251362	AAU	4000
			NL-11328	26/05/2009 12:33:43	External Transfer	Terminated	DE-3566255851-3566256340	AAU	490
							DE-3566256341-3566257340	AAU	1000
							DE-3566266803-3566268413	AAU	1611
							DE-3567620582-3567620781	AAU	200
							DE-3567629307-3567629542	AAU	236
							DE-3597160031-3597161353	AAU	1323
							FR-3257472676-3257474437	AAU	1762
							FR-3267247961-3267250901	AAU	2941
							FR-3267254582-3267254609	AAU	28
							FR-3285642694-3285642824	AAU	131
							FR-3288455693-3288457648	AAU	1956
							FR-3288462267-3288462351	AAU	85
							FR-3289102635-3289105634	AAU	3000
							FR-3289300983-3289301087	AAU	105
							FR-3289534974-3289536329	AAU	1356
							FR-3289536330-3289537422	AAU	1093
							FR-3289537423-3289537634	AAU	212
							FR-3289545812-3289546077	AAU	266
							FR-3289558635-3289560479	AAU	1845
							FR-3293753383-3293753996	AAU	614
							FR-3294835946-3294844945	AAU	9000
							FR-3301197816-3301202815	AAU	5000
							FR-3301202816-3301203815	AAU	1000
							FR-3301403112-3301407111	AAU	4000
							FR-3302599112-3302601111	AAU	2000
							FR-3302602112-3302604111	AAU	2000
							FR-3303636408-3303641407	AAU	5000
							FR-3303668573-3303669572	AAU	1000
			NL-11329	26/05/2009 12:49:55	External Transfer	Terminated	PL-3249253579-3249254324	AAU	746
							DE-3566255851-3566256340	AAU	490
							DE-3566256341-3566257340	AAU	1000
							DE-3566266803-3566268413	AAU	1611
							DE-3567620582-3567620781	AAU	200
							DE-3567629307-3567629542	AAU	236
							DE-3597160031-3597161353	AAU	1323
							FR-3257472676-3257474437	AAU	1762
							FR-3267247961-3267250901	AAU	2941
							FR-3267254582-3267254609	AAU	28
							FR-3285642694-3285642824	AAU	131
							FR-3288455693-3288457648	AAU	1956
							FR-3288462267-3288462351	AAU	85
							FR-3289102635-3289105634	AAU	3000
							FR-3289300983-3289301087	AAU	105

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
							FR-3289534974-3289536329	AAU	1356
							FR-3289536330-3289537422	AAU	1093
							FR-3289537423-3289537634	AAU	212
							FR-3289545812-3289546077	AAU	266
							FR-3289558635-3289560479	AAU	1845
							FR-3293753383-3293753996	AAU	614
							FR-3294835946-3294844945	AAU	9000
							FR-3301197816-3301202815	AAU	5000
							FR-3301202816-3301203815	AAU	1000
							FR-3301403112-3301407111	AAU	4000
							FR-3302599112-3302601111	AAU	2000
							FR-3302602112-3302604111	AAU	2000
							FR-3303636408-3303641407	AAU	5000
							FR-3303668573-3303669572	AAU	1000
							PL-3249253579-3249254324	AAU	746
			NL-12793	01/12/2009 10:13:30	External Transfer	Terminated			
							GB-3892021446-3892021505	AAU	60
							GB-3901447620-3901447862	AAU	243
							GB-3901449666-3901449780	AAU	115
							GB-3901547792-3901548226	AAU	435
							GB-3910430963-3910431551	AAU	589
							GB-3913551143-3913551557	AAU	415
							GB-3976982109-3976982922	AAU	814
							GB-3980733265-3980733838	AAU	574
							GB-3986931627-3986932189	AAU	563
							GB-3987027190-3987027744	AAU	555
							GB-3991549420-3991550234	AAU	815
							GB-3992440284-3992441110	AAU	827
							GB-3993039216-3993040044	AAU	829
							GB-3993068216-3993068776	AAU	561
							GB-3995247246-3995247799	AAU	554
							GB-3995310094-3995310652	AAU	559
							GB-3995362725-3995363283	AAU	559
							GB-3995707164-3995707977	AAU	814
							GB-3995847185-3995848008	AAU	824
							GB-3996429812-3996430377	AAU	566
							GB-3999250712-3999251303	AAU	592
							GB-4000404168-4000404217	AAU	50
							GB-4000478042-4000478637	AAU	596
							GB-4002378797-4002379394	AAU	598
							GB-4002397035-4002397353	AAU	319
							GB-4003815679-4003816252	AAU	574
			NL-12794	01/12/2009 10:15:48	External Transfer	Terminated			
							DE-1643914844-1643922166	AAU	7323
							DE-2135936907-2135944583	AAU	7677
							DE-3441941939-3441942591	AAU	653
							DE-3442682583-3442682929	AAU	347
							DE-3558925244-3558926243	AAU	1000
							DE-3588697472-3588698471	AAU	1000
							ES-1829735803-1829742368	AAU	6566
							ES-1829742369-1829753368	AAU	11000
							ES-1829753369-1829755802	AAU	2434
							GB-3764984053-3764986052	AAU	2000
							GB-4000051622-4000086754	AAU	35133

DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-12796	01/12/2009 10:19:10	External Transfer	Terminated	GB-4000374755-4000395962 GB-4000547359-4000569257 PL-3186805746-3186807505	AAU AAU AAU	21208 21899 1760
							ES-1829671153-1829679368 ES-1829679369-1829683368 ES-1829717369-1829718845 ES-1829718846-1829721650 ES-1829727301-1829727368 ES-1829727369-1829735802	AAU AAU AAU AAU AAU AAU	8216 4000 1477 2805 68 8434
			NL-12801	01/12/2009 11:14:20	External Transfer	Terminated	AT-1585046206-1585048205 AT-1585048206-1585055205 AT-1585070056-1585071055	AAU AAU AAU	2000 7000 1000
			NL-12803	01/12/2009 11:16:47	Internal Transfer	Terminated	AT-1557053298-1557057381 AT-1570623324-1570646104 AT-1574166444-1574169163 AT-1574212740-1574214922 AT-1583303601-1583304573 AT-1583304950-1583307073 AT-1584137537-1584146671 AT-1585040206-1585046205	AAU AAU AAU AAU AAU AAU AAU AAU	4084 22781 2720 2183 973 2124 9135 6000
			NL-12806	01/12/2009 11:20:54	External Transfer	Terminated	AT-1557053298-1557057381 AT-1570623324-1570646104 AT-1574166444-1574169163 AT-1574212740-1574214922 AT-1583303601-1583304573 AT-1583304950-1583307073 AT-1584137537-1584146671 AT-1585040206-1585046205 GB-3686527678-3686547677	AAU AAU AAU AAU AAU AAU AAU AAU AAU	4084 22781 2720 2183 973 2124 9135 6000 20000
			NL-12811	01/12/2009 11:24:42	External Transfer	Terminated	AT-1584142672-1584146671 AT-1585040206-1585046205	AAU AAU	4000 6000
			NL-12947	07/12/2009 16:13:08	External Transfer	Terminated	CN-86986984-86987861 CN-86989121-86990379	CER CER	878 1259
			NL-12949	07/12/2009 16:16:09	External Transfer	Terminated	CN-86986984-86987861 CN-86989121-86990379	CER CER	878 1259
			NL-12950	07/12/2009 16:19:09	External Transfer	Terminated	CN-86986984-86987861 CN-86989121-86990379	CER CER	878 1259
			NL-12951	07/12/2009 16:22:29	External Transfer	Terminated	CN-86986984-86987861 CN-86989121-86990379	CER CER	878 1259
			NL-12952	07/12/2009 16:25:41	External Transfer	Terminated	CN-86986984-86987861	CER	878



DES Response Code	Average number of occurrences per transaction (X 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Units Involved Abbreviated		
	Reported Year	Prior to the Reported Year					Serial Number	Unit Type	Quantity
			NL-13067	16/12/2009 13:09:26	Internal Transfer	Terminated	CN-86989121-86990379	CER	1259
			NL-13068	16/12/2009 13:34:34	Internal Transfer	Terminated	BR-22518777-22531294	CER	12518
			NL-13069	16/12/2009 14:24:05	Internal Transfer	Terminated	BR-22518777-22531294	CER	12518
							CZ-1322159770-1322160390	AAU	621
							CZ-1322160391-1322164304	AAU	3914
							CZ-1322461352-1322465351	AAU	4000
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383
			NL-13070	16/12/2009 14:40:34	Internal Transfer	Terminated	CZ-1322159770-1322160390	AAU	621
							CZ-1322160391-1322164304	AAU	3914
							CZ-1322461352-1322465351	AAU	4000
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383
			NL-13077	17/12/2009 10:35:22	Internal Transfer	Terminated	CZ-1322464333-1322465351	AAU	1019
							CZ-1325723117-1325724745	AAU	1629
							CZ-1325762764-1325765745	AAU	2982
							CZ-1325765746-1325767763	AAU	2018
							CZ-1325767764-1325770763	AAU	3000
							CZ-1325770764-1325771214	AAU	451
							CZ-1326000299-1326000540	AAU	242
							CZ-1331643733-1331646008	AAU	2276
							CZ-1332040733-1332045732	AAU	5000
							CZ-1332292335-1332296334	AAU	4000
							CZ-1332365773-1332366155	AAU	383

# Annex 7

## Tables 6.1 and 6.2 of the IPCC Good Practice guidance

As described in Section 1.7, a Tier 1 uncertainty assessment was made to estimate the uncertainty in total national greenhouse gas emissions and in their trend. Tier 1 here means that non-Gaussian uncertainty distributions and correlations between sources have been neglected<sup>5</sup>. The uncertainty estimates for activity data and emission factors as listed in Table A7.2 were also used for a Tier 1 trend uncertainty assessment, as shown in Table A7.1. Uncertainties for the activity data and emission factors are derived from a mixture of empirical data and expert judgment and presented here as half the 95% confidence interval. The reason for halving the 95% confidence interval is that the value then corresponds to the familiar plus or minus value when uncertainties are loosely quoted as “plus or minus x%”.

Details on this calculation can be found in Table A7.2 and in Olivier et al. (2009). It should be stressed that most uncertainty estimates are ultimately based on collective expert judgment and therefore also rather uncertain (usually of the order of 50%). However, the reason to make these estimates is to identify the relatively most important uncertain sources.

For this purpose, a reasonable order-of-magnitude estimate of the uncertainty in activity data and in emission factors is usually sufficient: uncertainty estimates are a means to identify and prioritize inventory improvement activities, rather than an objective in itself.

This result may be interpreted in two ways: part of the uncertainty is due to inherent lack of knowledge on the sources that cannot be improved. Another part, however, can be attributed to elements of the inventory of which the uncertainty could be reduced in the course of time. The latter may be a result of either dedicated research initiated by the Inventory Agency or by other researchers. When this type of uncertainty is in sources that are expected to be relevant for emission reduction policies, the effectiveness of the policy package could be in jeopardy if the unreduced emissions turn out to be much less than originally estimated.

The results of this uncertainty assessment for the list of potential key sources can also be used to refine the Tier 1 key source assessment discussed above.

Uncertainty estimates for Tier 1 trend.

Table A7.1

	Uncertainty in emission level	Uncertainty in emission trend
<i>CO<sub>2</sub>eq</i>	± 5%	± 3%-points of 3% decrease
<i>CO<sub>2</sub></i>	± 3%	± 3%-points of 10% increase
<i>CH<sub>4</sub></i>	± 25%	± 9%-points of 33% decrease
<i>N<sub>2</sub>O</i>	± 60%*	± 15%-points of 41% decrease
<i>F-gases</i>	± 50%	± 10%-points of 72% decrease

\* The uncertainty is higher compared to last submission as a result of the emission reduction in the nitric acid production which was a large source with a relative low uncertainty.

<sup>5</sup> We note that a Tier 2 uncertainty assessment and a comparison with a Tier 1 uncertainty estimate based on similar data showed that in the Dutch circumstances the errors made in the simplified Tier 1 approach for estimating uncertainties are quite small (Olsthoorn and Pielaat, 2003; and Ramírez-Ramírez et al., 2006). This conclusion holds for both annual uncertainties and the trend uncertainty (see Section 1.7 for more details).

Tier 1 level and trend uncertainty assessment 1990–2008 (for F-gases with base year 1995) with the categories of the IPCC potential key source list (without adjustment for correlation sources).

Table A7.2

IPCC	Category	Gas	CO <sub>2</sub> eq base year abt	CO <sub>2</sub> eq last year abt	AD unc.	EF unc.	EM uncertainty	Combined Uncertainty as % of total national emissions	Type A sensitivity	Type B sensitivity	Potentiality in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A1a	Stationary combustion: Public Electricity and Heat Production: liquids	CO <sub>2</sub>	207	937	1%	10%	10%	0.2%	0.2%	0%	0.0%	0.0%	0.0%
1A1a	Stationary combustion: Public Electricity and Heat Production: solids	CO <sub>2</sub>	25 776	25 385	1%	3%	3%	0.4%	0.1%	11%	0.0%	0.2%	0.2%
1A1a	Stationary combustion: Public Electricity and Heat Production: gases	CO <sub>2</sub>	13 348	23 937	1%	1%	1%	0.1%	4.3%	11%	0.0%	0.1%	0.1%
1A1a	Stationary combustion: Public Electricity and Heat Production: waste incinerator	CO <sub>2</sub>	601	2 242	10%	5%	11%	0.1%	0.7%	1%	0.0%	0.1%	0.1%
1A1b	Stationary combustion: Petroleum Refining: liquids	CO <sub>2</sub>	9 999	9 099	10%	10%	14%	0.4%	-0.2%	4%	0.0%	0.0%	0.0%
1A1b	Stationary combustion: Petroleum Refining: gases	CO <sub>2</sub>	1 042	2 723	1%	1%	1%	0.3%	0.0%	1%	0.0%	0.0%	0.0%
1A1c	Stationary combustion: Manufacture of Solid Fuels and Other Ex. Ind.: liquids	CO <sub>2</sub>	2	1	20%	2%	20%	0.2%	0.0%	0%	0.0%	0.0%	0.0%
1A1c	Stationary combustion: Manufacture of Solid Fuels and Other Ex. Ind.: gases	CO <sub>2</sub>	1 526	1 940	20%	5%	21%	0.2%	0.2%	1%	0.0%	0.2%	0.2%
1A2	Stationary combustion: Manufacturing Industries and Construction: liquids	CO <sub>2</sub>	9 844	9 021	1%	5%	9%	0.2%	0.2%	4%	0.0%	0.1%	0.1%
1A2	Stationary combustion: Manufacturing Industries and Construction: solids	CO <sub>2</sub>	5 033	4 691	2%	10%	10%	0.2%	-0.1%	2%	0.0%	0.1%	0.1%
1A2	Stationary combustion: Manufacturing Industries and Construction: gases	CO <sub>2</sub>	19 030	13 017	2%	1%	2%	0.1%	-2.1%	0%	0.0%	0.2%	0.2%
1A4	Stationary combustion: Other Sectors: solids	CO <sub>2</sub>	189	96	90%	9%	90%	0.2%	-0.1%	0%	0.0%	0.0%	0.0%
1A4a	Stationary combustion: Other Sectors: Commercial/Institutional: gases	CO <sub>2</sub>	7 632	10 326	20%	1%	20%	1.0%	1.6%	5%	0.0%	1.4%	1.4%
1A4b	Stationary combustion: Other Sectors: Residential: gases	CO <sub>2</sub>	15 696	17 818	5%	1%	5%	0.4%	-0.3%	0%	0.0%	0.6%	0.6%
1A4c	Stationary combustion: Other Sectors: Agriculture/Forestry/Fisheries: gases	CO <sub>2</sub>	7 330	7 933	10%	1%	10%	0.4%	0.2%	2%	0.0%	0.6%	0.6%
1A4c	Stationary combustion: Other Sectors: Agriculture/Forestry/Fisheries: liquids	CO <sub>2</sub>	2 893	2 360	20%	2%	20%	0.2%	-0.2%	1%	0.0%	0.3%	0.3%
1A4	Stationary combustion: Other Sectors: liquids and From 1A4b	CO <sub>2</sub>	1 476	363	20%	2%	20%	0.2%	-0.5%	0%	0.0%	0.0%	0.0%
1A5	Military use of fuels (IAD Other)	CO <sub>2</sub>	666	392	20%	2%	20%	0.3%	-0.1%	0%	0.0%	0.1%	0.1%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	CH <sub>4</sub>	984	1 624	2%	50%	50%	0.4%	0.5%	1%	0.2%	0.0%	0.2%
1A	Emissions from stationary combustion: non-CO <sub>2</sub>	N <sub>2</sub> O	226	327	2%	50%	50%	0.1%	0.0%	0%	0.0%	0.0%	0.0%
1A2b	Mobile combustion: road vehicles: gasoline	CO <sub>2</sub>	12 902	12 096	2%	0%	2%	0.1%	1.0%	0%	0.0%	0.2%	0.2%
1A2b	Mobile combustion: road vehicles: diesel oil	CO <sub>2</sub>	11 832	20 861	6%	0%	6%	0.1%	4.2%	9%	0.0%	0.7%	0.7%
1A2b	Mobile combustion: road vehicles: LPG	CO <sub>2</sub>	2 738	1 010	10%	0%	10%	0.2%	-0.7%	0%	0.0%	0.1%	0.1%
1A2	Mobile combustion: waterborne navigation	CO <sub>2</sub>	406	637	20%	0%	20%	0.1%	0.1%	0%	0.0%	0.1%	0.1%
1A2	Mobile combustion: aircraft	CO <sub>2</sub>	41	41	90%	1%	90%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
1A2	Mobile combustion: other (railways)	CO <sub>2</sub>	31	89	6%	0%	6%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
1A2	Mobile combustion: other (non-road)	CH <sub>4</sub>	1	1	90%	100%	112%	0.2%	0.0%	0%	0.0%	0.0%	0.0%
1A2	Mobile combustion: other (non-road)	N <sub>2</sub> O	1	2	90%	100%	112%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
1A2	Mobile combustion: road vehicles	CH <sub>4</sub>	157	46	2%	60%	60%	0.3%	0.0%	0%	0.0%	0.0%	0.0%
1A2	Mobile combustion: road vehicles	N <sub>2</sub> O	271	432	6%	50%	50%	0.1%	0.1%	0%	0.0%	0.0%	0.0%
1B2	Fugitive emissions venting/flaring	CH <sub>4</sub>	1 252	390	2%	25%	25%	0.3%	-0.4%	0%	-0.1%	0.0%	0.1%
1B2	Fugitive emissions from oil and gas operations: gas distribution	CH <sub>4</sub>	256	359	2%	25%	25%	0.3%	0.0%	0%	0.0%	0.0%	0.0%
1B2	Fugitive emissions from oil and gas operations: other	CH <sub>4</sub>	162	156	20%	50%	50%	0.3%	0.0%	0%	0.0%	0.0%	0.0%
1B1b	CO <sub>2</sub> from coke production	CO <sub>2</sub>	420	430	90%	2%	90%	0.1%	0.0%	0%	0.0%	0.1%	0.1%
1B2	Fugitive emissions venting/flaring: CO <sub>2</sub>	CO <sub>2</sub>	775	72	90%	2%	90%	0.3%	-0.3%	0%	0.0%	0.0%	0.0%
2A1	Cement production	CO <sub>2</sub>	416	399	9%	10%	11%	0.3%	0.0%	0%	0.0%	0.0%	0.0%
2A2	Lime/limestone and dolomite use	CO <sub>2</sub>	232	261	29%	6%	26%	0.3%	0.0%	0%	0.0%	0.0%	0.0%
2A2	Other materials	CO <sub>2</sub>	275	471	29%	6%	29%	0.1%	0.1%	0%	0.0%	0.1%	0.1%
2B1	Ammonia production	CO <sub>2</sub>	3 036	2 899	2%	1%	2%	0.3%	-0.1%	1%	0.0%	0.0%	0.0%
2B2	Nitric acid production	N <sub>2</sub> O	6 330	559	10%	20%	22%	0.1%	-0.0%	0%	-0.0%	0.0%	0.0%
2B5	Coproduction production	N <sub>2</sub> O	756	481	20%	20%	28%	0.1%	-0.1%	0%	0.0%	0.1%	0.1%
2B5	Other chemical product manufacturing	CO <sub>2</sub>	608	849	90%	50%	71%	0.2%	0.0%	0%	0.0%	0.2%	0.2%
2C1	Iron and steel production (blast iron)	CO <sub>2</sub>	2 514	1 349	2%	6%	6%	0.0%	-0.5%	1%	0.0%	0.0%	0.0%
2C3	CO <sub>2</sub> from syngas production	CO <sub>2</sub>	395	447	2%	6%	6%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
2C3	HFC from syngas production	HFC	1 801	72	2%	20%	20%	0.0%	-0.0%	0%	-0.0%	0.0%	0.0%
2F	SF <sub>6</sub> emissions from SF <sub>6</sub> use	SF <sub>6</sub>	301	224	90%	25%	96%	0.1%	0.0%	0%	0.0%	0.1%	0.1%
2F	Emissions from substitutes for ozone depleting substances (CO <sub>2</sub> substitutes: HFC	HFC	247	1 623	10%	60%	61%	0.4%	0.6%	1%	0.3%	0.1%	0.3%
2F	HFC-23 emissions from HFC-23 manufacture	HFC	5 759	212	10%	10%	14%	0.0%	-0.0%	0%	-0.2%	0.0%	0.2%
2F	HFC by-product emissions from HFC manufacture	HFC	12	16	10%	20%	23%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
2F	HFC emissions from HFC use	HFC	37	180	6%	25%	26%	0.0%	0.1%	0%	0.0%	0.0%	0.0%
2G	Other industrial: CO <sub>2</sub>	CO <sub>2</sub>	140	127	6%	20%	21%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
2G	Other industrial: CH <sub>4</sub>	CH <sub>4</sub>	287	393	10%	50%	61%	0.1%	0.0%	0%	0.0%	0.0%	0.0%
2G	Other industrial: N <sub>2</sub> O	N <sub>2</sub> O	3	6	90%	50%	71%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
3	Indirect CO <sub>2</sub> from energy production	CO <sub>2</sub>	316	129	29%	10%	27%	0.0%	-0.1%	0%	0.0%	0.0%	0.0%
4A1	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: cattle	CH <sub>4</sub>	5 793	5 792	6%	14%	14%	0.4%	-0.4%	0%	-0.1%	0.2%	0.2%
4A2	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: swine	CH <sub>4</sub>	438	379	6%	50%	50%	0.1%	0.0%	0%	0.0%	0.0%	0.0%
4A	CH <sub>4</sub> emissions from enteric fermentation in domestic livestock: other	CH <sub>4</sub>	319	296	6%	30%	30%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
4B	Emissions from manure management	N <sub>2</sub> O	514	506	10%	100%	100%	0.4%	0.1%	0%	0.1%	0.1%	0.1%
4B1	Emissions from manure management: cattle	CH <sub>4</sub>	1 674	1 484	10%	100%	100%	0.1%	0.0%	1%	0.0%	0.1%	0.1%
4B	Emissions from manure management: poultry	CH <sub>4</sub>	1 140	1 118	10%	100%	100%	0.1%	0.0%	1%	0.0%	0.1%	0.1%
4B	Emissions from manure management: other	CH <sub>4</sub>	273	52	10%	100%	100%	0.0%	-0.1%	0%	-0.1%	0.0%	0.1%
4B	Emissions from manure management: other	CH <sub>4</sub>	11	16	10%	100%	100%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
4C1	Direct N <sub>2</sub> O emissions from agricultural soils	N <sub>2</sub> O	4 674	4 083	10%	60%	61%	1.4%	0.7%	2%	0.1%	0.3%	0.3%
4C2	Indirect N <sub>2</sub> O emissions from nitrogen used in agriculture	N <sub>2</sub> O	4 975	3 112	50%	200%	200%	0.3%	-0.0%	1%	-1.0%	1.0%	1.0%
4C2	Animal production on agricultural soils	N <sub>2</sub> O	1 440	636	10%	100%	100%	0.2%	-0.4%	0%	-0.4%	0.0%	0.4%
4C41	CH <sub>4</sub> emissions from solid waste: disposal sites	CH <sub>4</sub>	12 011	4 096	30%	15%	34%	0.0%	-1.1%	2%	-0.9%	1.0%	1.0%
4B	Emissions from wastewater handling	CH <sub>4</sub>	200	190	20%	25%	22%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
4B	Emissions from wastewater handling	N <sub>2</sub> O	468	469	50%	50%	44%	0.1%	0.0%	0%	0.0%	0.1%	0.1%
4D	Other HFC: CH <sub>4</sub>	CH <sub>4</sub>	1	64	20%	25%	22%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
3, 4D	Other HFC: N <sub>2</sub> O	N <sub>2</sub> O	200	117	20%	50%	64%	0.0%	-0.1%	0%	0.0%	0.0%	0.0%
5A1	SA1: Forest Land remaining Forest Land	CO <sub>2</sub>	2 529	2 236	29%	32%	27%	0.7%	-0.1%	1%	-0.1%	0.4%	0.4%
5A2	SA2: Land converted to Forest Land	CO <sub>2</sub>	3	639	29%	50%	67%	0.2%	0.3%	0%	0.2%	0.1%	0.2%
5B2	SB2: Land converted to Cropland	CO <sub>2</sub>	35	48	25%	50%	50%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
5C1	SC1: Grassland remaining Grassland	CO <sub>2</sub>	4 246	4 246	25.0%	50.0%	56%	1.2%	0.1%	2%	0.0%	0.7%	0.7%
5C2	SC2: Land converted to Grassland	CO <sub>2</sub>	394	550	25.0%	61.2%	66%	0.2%	0.1%	0%	0.0%	0.1%	0.1%
5D2	SD2: Land converted to Wetlands	CO <sub>2</sub>	40	56	25.0%	50.0%	56%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
5E2	SE2: Land converted to Settlements	CO <sub>2</sub>	212	296	25.0%	50.0%	56%	0.1%	0.0%	0%	0.0%	0.0%	0.1%
5F2	SF2: Land converted to Other Land	CO <sub>2</sub>	18	25	25.0%	50.0%	56%	0.0%	0.0%	0%	0.0%	0.0%	0.0%
5G	SG: Other (liming of soils)	CO <sub>2</sub>	183	71	25%	1%	25%	0.0%	0.0%	0%	0.0%	0.0%	0.0%

Emissions (Gg) and uncertainty estimates for the subcategories of Sector 5 LULUCF, as used in the Tier 1 uncertainty analysis.

Table A7.3

IPCC	Category	Gas	CO <sub>2</sub> eq 1990	CO <sub>2</sub> eq 2008	AD unc.	EF unc.	EM uncertainty estimate
5A1	5A1. Forest Land remaining Forest Land	CO <sub>2</sub>	-2.529	-2.208	25.0%	61.8%	67%
5A2	5A2. Land converted to Forest Land	CO <sub>2</sub>	-3	-639	25.0%	57.9%	63%
5B2	5B2. Land converted to Cropland	CO <sub>2</sub>	35	48	25.0%	50.0%	56%
5C1	5C1. Grassland remaining Grassland	CO <sub>2</sub>	4.246	4.246	25.0%	50.0%	56%
5C2	5C2. Land converted to Grassland	CO <sub>2</sub>	394	550	25.0%	61.2%	66%
5D2	5D2. Land converted to Wetlands	CO <sub>2</sub>	40	56	25.0%	50.0%	56%
5E2	5E2. Land converted to Settlements	CO <sub>2</sub>	212	296	25.0%	50.0%	56%
5F2	5F2. Land converted to Other Land	CO <sub>2</sub>	18	25	25.0%	50.0%	56%
5G	5G. Other (liming of soils)	CO <sub>2</sub>	183	71	25.0%	10%	25%

# Annex 8

## Emission Factors and Activity Data Agriculture

### Animal numbers

Table A8.1

	1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>							
Female young stock under 1 yr	752,658	696,063	562,563	499,937	488,356	509,863	532,259
Male young stock under 1 yr	53,229	44,163	37,440	33,778	31,736	32,470	33,545
Female young stock, 1-2 yrs	734,078	682,888	594,100	515,972	513,238	494,853	509,763
Male young stock, 1-2 yrs	34,635	33,118	26,328	18,149	17,206	13,627	14,939
Female young stock, 2 yrs and over	145,648	124,970	104,633	74,180	66,331	69,110	79,489
Cows in milk and in calf	1,877,684	1,707,875	1,504,097	1,433,202	1,419,716	1,413,166	1,466,134
Bulls for service 2 yrs and over	8,762	8,674	10,410	12,391	8,200	10,028	7,718
<i>Cattle for fattening</i>							
Meat calves, for rosé veal production	28,876	85,803	145,828	204,227	221,710	261,620	272,117
Meat calves, for white veal production	572,709	583,516	636,907	624,513	622,015	598,252	626,596
Female young stock < 1 yr	53,021	57,218	41,300	43,313	40,718	44,671	42,657
Male young stock (incl. young bullocks) < 1 yr	255,375	188,193	83,447	66,655	55,069	55,008	53,875
Female young stock, 1-2 yrs	56,934	66,653	44,807	43,452	43,381	41,102	44,005
Male young stock (incl. young bullocks), 1-2 yrs	178,257	169,546	88,669	52,788	52,392	49,280	52,029
Female young stock, 2 yrs and over	42,555	48,365	16,917	15,260	14,428	16,056	18,755
Male young stock (incl. young bullocks) ≥ 2 yrs	12,073	10,969	9,397	9,346	7,515	9,713	9,334
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	119,529	146,181	163,397	151,641	143,082	143,965	126,980
<i>Pigs</i>							
Piglets	5,190,749	5,596,117	5,102,434	4,562,991	4,646,509	4,837,355	4,965,922
Fattening pigs	7,025,102	7,123,923	6,504,540	5,504,295	5,475,689	5,558,828	5,838,974
Gilts not yet in pig	385,502	357,520	339,570	274,085	273,120	285,361	231,500
Sows	1,272,215	1,287,224	1,129,174	946,466	946,105	966,439	977,983
Young boars 1	13,893	11,382	6,917	6,486	5,516	4,192	4,335
Boars for service	27,587	21,297	35,182	17,235	9,028	10,479	7,753
<i>Poultry</i>							
Broilers	41,172,110	43,827,286	50,936,625	44,496,116	41,913,979	43,351,898	44,357,773
Broilers parents under 18 weeks	2,882,250	3,065,170	3,644,120	2,191,650	2,852,760	2,808,924	2,386,073
Broilers parents 18 weeks and over	4,389,830	4,506,840	5,397,520	3,596,700	3,992,590	4,260,360	4,862,707
Laying hens < 18 weeks, liq. manure	7,339,708	4,889,555	2,865,850	1,035,581	1,052,463	963,881	586,885
Laying hens < 18 weeks, solid manure	3,781,062	4,000,545	8,597,550	9,751,719	9,910,697	9,076,541	10,920,673
Laying hens ≥ 18 weeks, liq. manure	19,919,466	12,294,122	7,166,060	2,292,654	2,308,303	2,325,546	806,067
Laying hens ≥ 18 weeks, solid manure	13,279,644	16,977,598	25,406,940	29,549,756	29,751,467	29,973,709	32,780,059
Ducks for slaughter	1,085,510	868,965	958,466	1,030,867	1,043,349	1,134,146	1,063,799
Turkeys for slaughter	1,003,350	1,175,527	1,543,830	1,245,420	1,139,840	1,232,354	1,044,315
Turkeys parents under 7 months	28,550	13,930					
Turkeys parents 7 months and over	20,460	17,290					

## Animal numbers

Table A8.1 (continued)

	1990	1995	2000	2005	2006	2007	2008
<i>Rabbits (mother animals)</i>	105,246	64,234	52,252	48,034	40,517	49,413	41,410
<i>Minks (mother animals)</i>	543,969	456,104	584,806	691,862	694,017	802,853	848,664
<i>Foxes (mother animals)</i>	10,029	7,102	3,816	5,240	4,489	4,860	
<i>Other grazing animals</i>							
<i>Sheep (ewes)</i>	789,691	770,730	681,441	648,235	647,691	644,799	583,408
<i>Goats (mothers)</i>	37,472	43,231	98,077	172,159	177,423	188,676	207,882
<i>Horses</i>	49,931	70,101	78,892	87,807	83,262	86,008	93,170
<i>Ponies</i>	19,661	29,903	39,352	45,514	44,478	47,516	50,908

## Gross energy intake (MJ/head.day) for cattle

Table A8.2

	1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>							
<i>Female young stock under 1 yr</i>	73.6	75.6	75.0	75.8	76.4	77.0	75.5
<i>Male young stock under 1 yr</i>	86.1	86.7	85.1	89.1	89.4	89.8	89.8
<i>Female young stock, 1-2 yrs</i>	139.5	142.5	139.5	144.6	145.6	149.1	147.1
<i>Male young stock, 1-2 yrs</i>	151.1	162.2	155.9	154.1	154.1	155.8	154.7
<i>Female young stock, 2 yrs and over</i>	139.4	142.5	139.5	144.6	145.6	149.1	147.1
<i>Cows in milk and in calf</i>	279.6	292.1	306.8	321.2	327.2	333.2	332.1
<i>Bulls for service 2 yrs and over</i>	151.1	162.2	155.9	154.1	154.1	155.8	154.7
<i>Cattle for fattening</i>							
<i>Meat calves, for rosé veal production</i>	77.9	77.9	95.5	82.8	82.8	82.8	82.8
<i>Meat calves, for white veal production</i>	30.9	32.7	35.6	34.8	37.2	37.2	37.2
<i>Female young stock &lt; 1 yr</i>	73.6	75.5	74.9	75.8	76.2	76.7	75.3
<i>Male young stock (incl. young bullocks) &lt; 1 yr</i>	82.3	87.6	88.8	86.7	86.7	85.7	86.2
<i>Female young stock, 1-2 yrs</i>	139.5	142.4	139.3	144.4	145.4	148.9	147.0
<i>Male young stock (incl. young bullocks), 1-2 yrs</i>	167.3	164.1	154.1	157.5	157.4	156.0	156.7
<i>Female young stock, 2 yrs and over</i>	139.5	142.5	139.4	144.5	145.5	149.0	147.1
<i>Male young stock (incl. young bullocks) ≥ 2 yrs</i>	167.3	164.1	154.1	157.5	157.4	156.0	156.7
<i>Suckling cows (incl. fattening/grazing ≥ 2 yrs)</i>	165.0	167.1	169.1	180.0	181.4	184.0	184.9

## Emission factors enteric fermentation for cattle kg/animal. year

Table A8.3

	1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>							
<i>Female young stock under 1 yr</i>	29.0	29.8	29.5	29.8	30.1	30.3	29.7
<i>Male young stock under 1 yr</i>	33.9	34.1	33.5	35.0	35.2	35.3	35.3
<i>Female young stock, 1-2 yrs</i>	54.9	56.1	54.9	56.9	57.3	58.7	57.9
<i>Male young stock, 1-2 yrs</i>	59.5	63.8	61.3	60.7	60.7	61.3	60.9
<i>Female young stock, 2 yrs and over</i>	54.9	56.1	54.9	56.9	57.3	58.7	57.9
<i>Cows in milk and in calf</i>	110.5	115.8	120.0	126.3	127.8	129.4	128.3
<i>Bulls for service 2 yrs and over</i>	59.5	63.8	61.3	60.7	60.7	61.3	60.9
<i>Cattle for fattening</i>							
<i>Meat calves, for rosé veal production</i>	30.6	30.6	37.6	32.6	32.6	32.6	32.6
<i>Meat calves, for white veal production</i>	8.1	8.6	9.3	9.1	9.8	9.8	9.8
<i>Female young stock &lt; 1 yr</i>	29.0	29.7	29.5	29.8	30.0	30.2	29.6
<i>Male young stock (incl. young bullocks) &lt; 1 yr</i>	32.4	34.5	34.9	34.1	34.1	33.7	33.9
<i>Female young stock, 1-2 yrs</i>	54.9	56.0	54.8	56.8	57.2	58.6	57.9
<i>Male young stock (incl. young bullocks), 1-2 yrs</i>	65.8	64.6	60.7	62.0	62.0	61.4	61.7
<i>Female young stock, 2 yrs and over</i>	54.9	56.1	54.9	56.9	57.2	58.6	57.9
<i>Male young stock (incl. young bullocks) ≥ 2 yrs</i>	65.8	64.6	60.7	62.0	62.0	61.4	61.7
<i>Suckling cows (incl. fattening/grazing ≥ 2 yrs)</i>	64.9	65.8	66.6	70.8	71.4	72.4	72.8

**Volatile Solids (= Organic Matter) kg per 1000 kg manure**
**Table A8.4**

		1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>								
Female young stock under 1 yr	liquid manure	60	66	64	64	64	64	64
Male young stock under 1 yr	liquid manure	60	66	64	64	64	64	64
Female young stock, 1-2 yrs	liquid manure	60	66	64	64	64	64	64
Male young stock, 1-2 yrs	liquid manure	60	66	64	64	64	64	64
Female young stock, 2 yrs and over	liquid manure	60	66	64	64	64	64	64
Cows in milk and in calf	liquid manure	60	66	64	64	64	64	64
Bulls for service 2 yrs and over	liquid manure	60	66	64	64	64	64	64
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production	liquid manure	22.5	22.5	22.5	22.5	22.5	22.5	22.5
Meat calves, for white veal production	liquid manure	15	15	15	15	15	15	15
Female young stock < 1 yr	liquid manure	60	66	64	64	64	64	64
Male young stock (incl. young bullocks) < 1 yr	liquid manure	60	66	64	64	64	64	64
Female young stock, 1-2 yrs	liquid manure	60	66	64	64	64	64	64
Male young stock (incl. young bullocks), 1-2 yrs	liquid manure	60	66	64	64	64	64	64
Female young stock, 2 yrs and over	liquid manure	60	66	64	64	64	64	64
Male young stock (incl. young bullocks) ≥ 2 yrs	liquid manure	60	66	64	64	64	64	64
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	solid manure	140	153	150	150	150	150	150
<i>Cattle for breeding</i>								
Female young stock under 1 yr	Pasture	60	66	64	64	64	64	64
Male young stock under 1 yr								
Female young stock, 1-2 yrs	Pasture	60	66	64	64	64	64	64
Male young stock, 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	60	66	64	64	64	64	64
Cows in milk and in calf	Pasture	60	66	64	64	64	64	64
Bulls for service 2 yrs and over								
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production								
Meat calves, for white veal production								
Female young stock < 1 yr	Pasture	60	66	64	64	64	64	64
Male young stock (incl. young bullocks) < 1 yr								
Female young stock, 1-2 yrs	Pasture	60	66	64	64	64	64	64
Male young stock (incl. young bullocks), 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	60	66	64	64	64	64	64
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	60	66	64	64	64	64	64
<i>Pigs</i>								
<i>Piglets</i>								
Fattening pigs	liquid manure	50	60	60	60	60	60	60
Gilts not yet in pig	liquid manure	35	35	35	35	35	35	35
Sows	liquid manure	35	35	35	35	35	35	35
Young boars 1	liquid manure	35	35	35	35	35	35	35
Boars for service	liquid manure	35	35	35	35	35	35	35
<i>Poultry</i>								
Broilers	solid manure	508	508	508	508	508	508	508
Broilers parents under 18 weeks	solid manure	423	423	423	423	423	423	423
Broilers parents 18 weeks and over	solid manure	423	423	423	423	423	423	423
Laying hens < 18 weeks, liq. manure	liquid manure	90	93	93	93	93	93	93
Laying hens < 18 weeks, solid manure	solid manure	350	350	350	350	350	350	350
Laying hens ≥ 18 weeks, liq. manure	liquid manure	90	93	93	93	93	93	93
Laying hens ≥ 18 weeks, solid manure	solid manure	350	350	350	350	350	350	350
Ducks for slaughter	solid manure	209	209	209	209	209	209	209
Turkeys for slaughter	solid manure	464	464	464	464	464	464	464
Turkeys parents under 7 months	solid manure	464	464	464	464	464	464	464
Turkeys parents 7 months and over	solid manure	464	464	464	464	464	464	464
Rabbits (mother animals)	solid manure	367	367	367	367	367	367	367
Minks (mother animals)	solid manure	185	185	185	185	185	185	185
Foxes (mother animals)	solid manure	185	185	185	185	185	185	185

# Volatile Solids (= Organic Matter) per 1000 kg manure

Table A8.4 (continued)

		1990	1995	2000	2005	2006	2007	2008
<i>Ruminants, not cattle</i>								
Sheep (ewes)	<i>solid manure</i>	205	205	205	205	205	205	205
Goats (mothers)	<i>solid manure</i>	182	182	182	182	182	182	182
Horses	<i>solid manure</i>	250	250	250	250	250	250	250
Ponies	<i>solid manure</i>	250	250	250	250	250	250	250
<i>Ruminants, not cattle</i>								
Sheep (ewes)	<i>Pasture</i>	60	66	64	64	64	64	64
Goats (mothers)								
Horses	<i>Pasture</i>	60	66	64	64	64	64	64
Ponies	<i>Pasture</i>	60	66	64	64	64	64	64

# Methane conversion factor for pigs and poultry

Table A8.5

		1990	1995	2000	2005	2006	2007	2008
<i>Pigs</i>								
<i>Piglets</i>								
Fattening pigs	<i>liquid manure</i>	0.34	0.36	0.39	0.39	0.39	0.39	0.39
Gilts not yet in pig	<i>liquid manure</i>	0.34	0.36	0.39	0.39	0.39	0.39	0.39
Sows	<i>liquid manure</i>	0.34	0.36	0.39	0.39	0.39	0.39	0.39
Young boars 1	<i>liquid manure</i>	0.34	0.36	0.39	0.39	0.39	0.39	0.39
Boars for service	<i>liquid manure</i>	0.34	0.36	0.39	0.39	0.39	0.39	0.39
<i>Poultry</i>								
Broilers	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Broilers parents under 18 weeks	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Broilers parents 18 weeks and over	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Laying hens < 18 weeks, liq. manure	<i>liquid manure</i>	0.39	0.39	0.39	0.39	0.39	0.39	0.39
Laying hens < 18 weeks, solid manure	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Laying hens ≥ 18 weeks, liq. manure	<i>liquid manure</i>	0.39	0.39	0.39	0.39	0.39	0.39	0.39
Laying hens ≥ 18 weeks, solid manure	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Ducks for slaughter	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Turkeys for slaughter	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Turkeys parents under 7 months	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Turkeys parents 7 months and over	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Rabbits (mother animals)	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Minks (mother animals)	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015
Foxes (mother animals)	<i>solid manure</i>	0.015	0.015	0.015	0.015	0.015	0.015	0.015



		MCF	B0			MCF	B0
<i>Cattle for breeding</i>				<i>Pigs</i>			
<i>Female young stock under 1 yr</i>	liquid manure	0.17	0.25	<i>Piglets</i>			
<i>Male young stock under 1 yr</i>	liquid manure	0.17	0.25	<i>Fattening pigs</i>	liquid manure	0.34	
<i>Female young stock, 1-2 yrs</i>	liquid manure	0.17	0.25	<i>Gilts not yet in pig</i>	liquid manure	0.34	
<i>Male young stock, 1-2 yrs</i>	liquid manure	0.17	0.25	<i>Sows</i>	liquid manure	0.34	
<i>Female young stock, 2 yrs and over</i>	liquid manure	0.17	0.25	<i>Young boars 1</i>	liquid manure	0.34	
<i>Cows in milk and in calf</i>	liquid manure	0.17	0.25	<i>Boars for service</i>	liquid manure	0.34	
<i>Bulls for service 2 yrs and over</i>	liquid manure	0.17	0.25				
<i>Cattle for fattening</i>	liquid manure			<i>Poultry</i>			
<i>Meat calves, for rosé veal production</i>	liquid manure	0.14	0.25	<i>Broilers</i>	solid manure	0.34	
<i>Meat calves, for white veal production</i>	liquid manure	0.14	0.25	<i>Broilers parents under 18 weeks</i>	solid manure	0.34	
<i>Female young stock &lt; 1 yr</i>	liquid manure	0.17	0.25	<i>Broilers parents 18 weeks and over</i>	solid manure	0.34	
<i>Male young stock (incl. young bullocks) &lt; 1 yr</i>	liquid manure	0.17	0.25	<i>Laying hens &lt; 18 weeks, liq. manure</i>	liquid manure	0.34	
<i>Female young stock, 1-2 yrs</i>	liquid manure	0.17	0.25	<i>Laying hens &lt; 18 weeks, solid manure</i>	solid manure	0.34	
<i>Male young stock (incl. young bullocks), 1-2 yrs</i>	liquid manure	0.17	0.25	<i>Laying hens ≥ 18 weeks, liq. manure</i>	liquid manure	0.34	
<i>Female young stock, 2 yrs and over</i>	liquid manure	0.17	0.25	<i>Laying hens ≥ 18 weeks, solid manure</i>	solid manure	0.34	
<i>Male young stock (incl. young bullocks) ≥ 2 yrs</i>	liquid manure	0.17	0.25	<i>Ducks for slaughter</i>	solid manure	0.34	
<i>Suckling cows (incl. fattening/grazing ≥ 2 yrs)</i>	solid manure	0.015	0.25	<i>Turkeys for slaughter</i>	solid manure	0.34	
				<i>Turkeys parents under 7 months</i>	solid manure	0.34	
<i>Cattle for breeding</i>				<i>Turkeys parents 7 months and over</i>	solid manure	0.34	
<i>Female young stock under 1 yr</i>	Pasture	0.01	0.25	<i>Rabbits (mother animals)</i>	solid manure	0.34	
<i>Male young stock under 1 yr</i>				<i>Minks (mother animals)</i>	solid manure	0.34	
<i>Female young stock, 1-2 yrs</i>	Pasture	0.01	0.25	<i>Foxes (mother animals)</i>	solid manure	0.34	
<i>Male young stock, 1-2 yrs</i>							
<i>Female young stock, 2 yrs and over</i>	Pasture	0.01	0.25	<i>Ruminants, not cattle</i>			
<i>Cows in milk and in calf</i>	Pasture	0.01	0.25	<i>Sheep (ewes)</i>	solid manure	0.015	0.25
<i>Bulls for service 2 yrs and over</i>				<i>Goats (mothers)</i>	solid manure	0.015	0.25
<i>Cattle for fattening</i>				<i>Horses</i>	solid manure	0.015	0.25
<i>Meat calves, for rosé veal production</i>				<i>Ponies</i>	solid manure	0.015	0.25
<i>Meat calves, for white veal production</i>							
<i>Female young stock &lt; 1 yr</i>	Pasture	0.01	0.25	<i>Ruminants, not cattle</i>			
<i>Male young stock (incl. young bullocks) &lt; 1 yr</i>				<i>Sheep (ewes)</i>	pasture	0.01	0.25
<i>Female young stock, 1-2 yrs</i>	Pasture	0.01	0.25	<i>Goats (mothers)</i>			
<i>Male young stock (incl. young bullocks), 1-2 yrs</i>				<i>Horses</i>	pasture	0.01	0.25
<i>Female young stock, 2 yrs and over</i>	Pasture	0.01	0.25	<i>Ponies</i>	pasture	0.01	0.25
<i>Male young stock (incl. young bullocks) ≥ 2 yrs</i>							
<i>Suckling cows (incl. fattening/grazing ≥ 2 yrs)</i>	Pasture	0.01	0.25				

		1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>								
Female young stock under 1 yr	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Male young stock under 1 yr	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Female young stock, 1-2 yrs	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Male young stock, 1-2 yrs	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Female young stock, 2 yrs and over	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Cows in milk and in calf	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Bulls for service 2 yrs and over	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production	liquid manure	0.00052	0.00052	0.00052	0.00052	0.00052	0.00052	0.00052
Meat calves, for white veal production	liquid manure	0.00035	0.00035	0.00035	0.00035	0.00035	0.00035	0.00035
Female young stock < 1 yr	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Male young stock (incl. young bullocks) < 1 yr	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Female young stock, 1-2 yrs	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Male young stock (incl. young bullocks), 1-2 yrs	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Female young stock, 2 yrs and over	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Male young stock (incl. young bullocks) ≥ 2 yrs	liquid manure	0.00169	0.00186	0.00180	0.00180	0.00180	0.00180	0.00180
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	solid manure	0.00035	0.00038	0.00037	0.00037	0.00037	0.00037	0.00037
<i>Cattle for breeding</i>								
Female young stock under 1 yr	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Male young stock under 1 yr								
Female young stock, 1-2 yrs	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Male young stock, 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Cows in milk and in calf	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Bulls for service 2 yrs and over								
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production								
Meat calves, for white veal production								
Female young stock < 1 yr	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Male young stock (incl. young bullocks) < 1 yr								
Female young stock, 1-2 yrs	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Male young stock (incl. young bullocks), 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<i>Pigs</i>								
<i>Piglets</i>								
Fattening pigs	liquid manure	0.00383	0.00486	0.00527	0.00527	0.00527	0.00527	0.00527
Gilts not yet in pig	liquid manure	0.00268	0.00284	0.00307	0.00307	0.00307	0.00307	0.00307
Sows	liquid manure	0.00268	0.00284	0.00307	0.00307	0.00307	0.00307	0.00307
Young boars 1	liquid manure	0.00268	0.00284	0.00307	0.00307	0.00307	0.00307	0.00307
Boars for service	liquid manure	0.00268	0.00284	0.00307	0.00307	0.00307	0.00307	0.00307
<i>Poultry</i>								
Broilers	solid manure	0.00172	0.00172	0.00172	0.00172	0.00172	0.00172	0.00172
Broilers parents under 18 weeks	solid manure	0.00143	0.00143	0.00143	0.00143	0.00143	0.00143	0.00143
Broilers parents 18 weeks and over	solid manure	0.00143	0.00143	0.00143	0.00143	0.00143	0.00143	0.00143
Laying hens < 18 weeks, liq. manure	liquid manure	0.00790	0.00816	0.00816	0.00816	0.00816	0.00816	0.00816
Laying hens < 18 weeks, solid manure	solid manure	0.00118	0.00118	0.00118	0.00118	0.00118	0.00118	0.00118
Laying hens ≥ 18 weeks, liq. manure	liquid manure	0.00790	0.00816	0.00816	0.00816	0.00816	0.00816	0.00816
Laying hens ≥ 18 weeks, solid manure	solid manure	0.00118	0.00118	0.00118	0.00118	0.00118	0.00118	0.00118
Ducks for slaughter	solid manure	0.00071	0.00071	0.00071	0.00071	0.00071	0.00071	0.00071
Turkeys for slaughter	solid manure	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157
Turkeys parents under 7 months	solid manure	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157
Turkeys parents 7 months and over	solid manure	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157	0.00157
Rabbits (mother animals)	solid manure	0.00124	0.00124	0.00124	0.00124	0.00124	0.00124	0.00124
Minks (mother animals)	solid manure	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062
Foxes (mother animals)	solid manure	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062

		1990	1995	2000	2005	2006	2007	2008
<i>Ruminants, not cattle</i>								
<i>Sheep (ewes)</i>	<i>solid manure</i>	0.00051	0.00051	0.00051	0.00051	0.00051	0.00051	0.00051
<i>Goats (mothers)</i>	<i>solid manure</i>	0.00045	0.00045	0.00045	0.00045	0.00045	0.00045	0.00045
<i>Horses</i>	<i>solid manure</i>	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062
<i>Ponies</i>	<i>solid manure</i>	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062	0.00062
<i>Ruminants, not cattle</i>								
<i>Sheep (ewes)</i>	<i>Pasture</i>	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<i>Goats (mothers)</i>								
<i>Horses</i>	<i>Pasture</i>	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<i>Ponies</i>	<i>Pasture</i>	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011

		1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>								
Female young stock under 1 yr	liquid manure	3,500	3,500	3,500	3,500	3,500	3,500	4,000
Male young stock under 1 yr	liquid manure	5,000	5,000	5,000	5,000	5,000	5,000	5,000
Female young stock, 1-2 yrs	liquid manure	6,000	6,000	6,000	6,000	6,000	6,000	6,500
Male young stock, 1-2 yrs	liquid manure	11,500	11,500	11,500	11,500	11,500	11,500	11,500
Female young stock, 2 yrs and over	liquid manure	6,000	6,000	6,000	6,000	6,000	6,000	6,500
Cows in milk and in calf	liquid manure	16,000	16,000	18,000	20,500	21,000	21,500	20,500
Bulls for service 2 yrs and over	liquid manure	11,500	11,500	11,500	11,500	11,500	11,500	11,500
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production	liquid manure	5,000	5,000	5,000	5,000	5,000	4,300	4,300
Meat calves, for white veal production	liquid manure	3,500	3,500	3,500	3,000	3,000	3,000	3,000
Female young stock < 1 yr	liquid manure	3,500	3,500	3,500	3,500	3,500	3,500	4,000
Male young stock (incl. young bullocks) < 1 yr	liquid manure	4,500	4,500	4,500	4,500	4,500	4,500	4,500
Female young stock, 1-2 yrs	liquid manure	6,000	6,000	6,000	6,000	6,000	6,000	6,500
Male young stock (incl. young bullocks), 1-2 yrs	liquid manure	10,000	10,000	10,000	10,000	10,000	10,000	10,000
Female young stock, 2 yrs and over	liquid manure	6,000	6,000	6,000	6,000	6,000	6,000	6,500
Male young stock (incl. young bullocks) ≥ 2 yrs	liquid manure	10,000	10,000	10,000	10,000	10,000	10,000	10,000
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	solid manure	7,000	7,000	7,000	7,000	7,000	7,000	7,000
<i>Cattle for breeding</i>								
Female young stock under 1 yr	Pasture	1,500	1,500	1,500	1,500	1,500	1,500	1,000
Male young stock under 1 yr								
Female young stock, 1-2 yrs	Pasture	5,500	5,500	5,500	5,500	5,500	5,500	5,000
Male young stock, 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	5,500	5,500	5,500	5,500	5,500	5,500	5,000
Cows in milk and in calf	Pasture	7,000	7,000	7,000	5,500	5,000	4,500	5,500
Bulls for service 2 yrs and over								
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production								
Meat calves, for white veal production								
Female young stock < 1 yr	Pasture	1,500	1,500	1,500	1,500	1,500	1,500	1,000
Male young stock (incl. young bullocks) < 1 yr								
Female young stock, 1-2 yrs	Pasture	5,500	5,500	5,500	5,500	5,500	5,500	5,000
Male young stock (incl. young bullocks), 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	5,500	5,500	5,500	5,500	5,500	5,500	5,000
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	8,000	8,000	8,000	8,000	8,000	8,000	8,000
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<i>Pigs</i>								
<i>Piglets</i>								
Fattening pigs	liquid manure	1,300	1,250	1,200	1,200	1,200	1,200	1,200
Gilts not yet in pig	liquid manure	1,300	1,300	1,300	1,300	1,300	1,300	1,300
Sows	liquid manure	5,200	5,200	5,100	5,100	5,100	5,100	5,100
Young boars 1	liquid manure	1,300	1,300	1,300	1,300	1,300	1,300	1,300
Boars for service	liquid manure	3,200	3,200	3,200	3,200	3,200	3,200	3,200
<i>Poultry</i>								
<i>Broilers</i>								
Broilers	solid manure	10	11	11	10.9	10.9	10.9	10.9
Broilers parents under 18 weeks	solid manure	15.4	13.4	13.4	8.2	8.2	8.2	8.2
Broilers parents 18 weeks and over	solid manure	25.3	23.0	23.0	20.6	20.6	20.6	20.6
Laying hens < 18 weeks, liq. Manure	liquid manure	25.4	25.4	25.4	22.5	22.5	22.5	22.5
Laying hens < 18 weeks, solid manure	solid manure	10.0	10.0	9.0	7.6	7.6	7.6	7.6
Laying hens ≥ 18 weeks, liq. Manure	liquid manure	63.5	63.5	63.5	53.4	53.4	53.4	53.4
Laying hens ≥ 18 weeks, solid manure	solid manure	22.5	23.5	24.0	18.9	18.9	18.9	18.9
Ducks for slaughter	solid manure	86.3	70.0	70.0	70.0	70.0	70.0	70.0
Turkeys for slaughter	solid manure	37.9	45.0	45.0	45.0	45.0	45.0	45.0
Turkeys parents under 7 months	solid manure	49.4	49.4					
Turkeys parents 7 months and over	solid manure	78.6	78.6					
Rabbits (mother animals)	solid manure	377	377	377	377	377	377	377
Minks (mother animals)	solid manure	103.7	103.7	103.7	103.7	103.7	103.7	103.7
Foxes (mother animals)	solid manure	272.2	272.2	272.2	272.2	272.2	272.2	

Emission factors for methane from manure in CH<sub>4</sub>/kg manure. year

## Table A8.8 (continued)

		1990	1995	2000	2005	2006	2007	2008
<i>Ruminants, not cattle</i>								
<i>Sheep (ewes)</i>	<i>solid manure</i>	325	325	325	325	325	325	325
<i>Goats (mothers)</i>	<i>solid manure</i>	1,300	1,300	1,300	1,300	1,300	1,300	1,300
<i>Horses</i>	<i>solid manure</i>	5,200	5,200	5,200	5,200	5,200	5,200	5,200
<i>Ponies</i>	<i>solid manure</i>	2,100	2,100	2,100	2,100	2,100	2,100	2,100
<i>Ruminants, not cattle</i>								
<i>Sheep (ewes)</i>	<i>Pasture</i>	2,000	2,000	2,000	2,000	2,000	2,000	2,000
<i>Goats (mothers)</i>								
<i>Horses</i>	<i>Pasture</i>	3,300	3,300	3,300	3,300	3,300	3,300	3,300
<i>Ponies</i>	<i>Pasture</i>	2,100	2,100	2,100	2,100	2,100	2,100	2,100

		1990	1995	2000	2005	2006	2007	2008
<i>Cattle for breeding</i>								
Female young stock under 1 yr	liquid manure	26.5	29.8	29.0	23.0	22.8	24.6	29.2
Male young stock under 1 yr	liquid manure	39.6	40.8	37.0	37.0	36.7	36.6	35.9
Female young stock, 1-2 yrs	liquid manure	43.1	48.4	46.4	42.7	40.1	42.5	45.8
Male young stock, 1-2 yrs	liquid manure	90.6	101.9	96.8	88.5	87.4	89.6	86.0
Female young stock, 2 yrs and over	liquid manure	43.0	48.4	46.3	42.7	40.1	42.5	45.8
Cows in milk and in calf	liquid manure	95.9	104.0	97.2	103.2	103.9	110.8	103.2
Bulls for service 2 yrs and over	liquid manure	90.6	101.9	96.8	88.5	87.4	89.6	86.0
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production	liquid manure	28.9	28.9	34.1	27.2	27.0	28.1	10.7
Meat calves, for white veal production	liquid manure	10.6	11.6	11.9	10.6	11.2	11.0	27.4
Female young stock < 1 yr	liquid manure	26.2	29.4	28.6	22.8	22.5	24.4	28.8
Male young stock (incl. young bullocks) < 1 yr	liquid manure	28.9	29.5	26.6	27.0	27.3	26.6	26.0
Female young stock, 1-2 yrs	liquid manure	43.0	48.2	46.0	42.4	39.8	42.4	45.0
Male young stock (incl. young bullocks), 1-2 yrs	liquid manure	72.6	64.7	56.1	56.8	57.3	54.5	53.8
Female young stock, 2 yrs and over	liquid manure	43.1	48.4	46.1	42.5	39.9	42.4	44.9
Male young stock (incl. young bullocks) ≥ 2 yrs	liquid manure	72.6	64.7	56.1	56.8	57.3	54.5	53.8
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	solid manure	42.3	48.0	42.4	39.1	38.7	39.4	38.7
<i>Cattle for breeding</i>								
Female young stock under 1 yr	Pasture	15.3	14.4	13.0	17.0	16.6	14.3	7.5
Male young stock under 1 yr								
Female young stock, 1-2 yrs	Pasture	51.2	47.5	42.9	33.1	34.1	32.2	29.0
Male young stock, 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	51.2	47.5	42.9	33.1	34.1	32.2	29.0
Cows in milk and in calf	Pasture	52.6	52.5	39.3	30.8	28.8	25.7	31.3
Bulls for service 2 yrs and over								
<i>Cattle for fattening</i>								
Meat calves, for rosé veal production								
Meat calves, for white veal production								
Female young stock < 1 yr	Pasture	15.2	14.3	12.8	16.9	16.5	14.0	7.4
Male young stock (incl. young bullocks) < 1 yr								
Female young stock, 1-2 yrs	Pasture	51.2	47.5	42.9	33.1	34.1	32.0	29.4
Male young stock (incl. young bullocks), 1-2 yrs								
Female young stock, 2 yrs and over	Pasture	51.2	47.5	42.9	33.1	34.1	32.0	29.4
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	68.4	63.1	52.7	45.8	44.5	43.4	46.2
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	8,000	8,000	8,000	8,000	8,000	8,000	8,000
Male young stock (incl. young bullocks) ≥ 2 yrs								
Suckling cows (incl. fattening/grazing ≥ 2 yrs)	Pasture	0.00010	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<i>Pigs</i>								
<i>Piglets</i>								
Fattening pigs	liquid manure	14.3	14.5	12.3	12.3	12.6	12.6	12.9
Gilts not yet in pig	liquid manure	14.0	14.4	14.2	14.3	14.6	14.2	13.5
Sows	liquid manure	33.8	31.4	30.9	30.7	30.8	31.5	30.8
Young boars 1	liquid manure	14.0	14.4	14.2	14.3	14.6	14.2	13.5
Boars for service	liquid manure	25.0	24.6	22.9	23.7	23.9	23.3	23.5

## N excretion (kg/animal. yr)

Table A8.9 (continued)

		1990	1995	2000	2005	2006	2007	2008
<i>Poultry</i>								
Broilers	<i>solid manure</i>	0.61	0.63	0.51	0.55	0.53	0.53	0.53
Broilers parents under 18 weeks	<i>solid manure</i>	0.52	0.45	0.37	0.32	0.33	0.33	0.33
Broilers parents 18 weeks and over	<i>solid manure</i>	1.33	1.29	1.13	1.10	1.10	1.13	1.12
Laying hens < 18 weeks, liq. manure	<i>liquid manure</i>	0.38	0.36	0.31	0.32	0.33	0.34	0.34
Laying hens < 18 weeks, solid manure	<i>solid manure</i>	0.38	0.36	0.31	0.32	0.33	0.34	0.34
Laying hens ≥ 18 weeks, liq. manure	<i>liquid manure</i>	0.75	0.81	0.67	0.71	0.74	0.74	0.75
Laying hens ≥ 18 weeks, solid manure	<i>solid manure</i>	0.75	0.81	0.67	0.71	0.74	0.74	0.75
Ducks for slaughter	<i>solid manure</i>	1.12	1.09	0.99	0.89	0.91	0.85	0.76
Turkeys for slaughter	<i>solid manure</i>	1.98	1.97	1.85	1.81	1.66	1.69	1.71
Turkeys parents under 7 months	<i>solid manure</i>	2.38	2.78					
Turkeys parents 7 months and over	<i>solid manure</i>	3.17	3.04					
Rabbits (mother animals)	<i>solid manure</i>	8.7	8.1	7.6	8.2	8.1	8.0	7.9
Minks (mother animals)	<i>solid manure</i>	4.08	4.08	3.5	2.7	2.6	2.5	2.4
Foxes (mother animals)	<i>solid manure</i>	13.9	13.9	8.3	6.9	6.5	6.4	
<i>Ruminants, not cattle</i>								
Sheep (ewes)	<i>solid manure</i>	3.9	4.0	3.9	2.6	2.6	2.6	2.5
Goats (mothers)	<i>solid manure</i>	19.9	21.5	19.4	17.7	17.7	15.8	16.0
Horses	<i>solid manure</i>	33.3	33.3	33.3	33.3	33.3	32.1	30.3
Ponies	<i>solid manure</i>	14.4	14.4	14.4	14.4	14.4	13.8	13.2
<i>Ruminants, not cattle</i>								
Sheep (ewes)	<i>Pasture</i>	21.1	20.3	19.5	12.2	11.7	11.1	11.9
Goats (mothers)								
Horses	<i>Pasture</i>	30.2	30.2	30.2	30.2	30.2	29.4	28.2
Ponies	<i>Pasture</i>	19.9	19.9	19.9	19.9	19.9	19.4	18.9

## Nitrogen flows (Gg)

Table A8.10

	1990	1995	2000	2005	2006	2007	2008
Nitrogen fertilizer consumption	412.4	405.8	339.5	279.2	287.8	257.5	257.5
of which ammonium fertilizer	3.6	11.2	6.6	30.6	42.9	17.2	17.2
NH <sub>3</sub> -N emission during application	11.2	10.5	9.2	9.8	10.9	9.9	9.9
Net fertilizer to soil	401.1	395.3	330.3	269.4	276.9	247.6	247.6
Nitrogen excretion by animals	694.4	680.1	549.2	478.9	471.2	479.7	497.2
Nitrogen excretion in animals houses	506.5	508.1	424.5	385.6	382.5	398.6	411.6
of which in solid form	61.3	69.6	74.1	72.9	71.6	73.2	76.2
of which in liquid form	445.2	438.5	350.5	312.7	310.8	325.4	335.4
NH <sub>3</sub> -N emission in animal houses	73.4	73.8	60.6	50.2	50.0	50.5	51.5
Net available manure for application	433.1	434.4	364.0	335.5	332.5	348.1	360.1
Nitrogen in net manure exported abroad	6.4	22.1	14.7	14.9	15.8	28.8	30.5
NH <sub>3</sub> -N emission during application	98.3	51.4	36.8	32.2	32.6	34.4	27.9
Net animal manure to soil	328.4	360.9	312.4	287.4	284.1	284.9	301.8
Nitrogen excretion in meadow	188.0	171.9	124.6	93.3	88.7	81.1	85.5
NH <sub>3</sub> -N emission in meadow	13.0	11.9	8.5	7.1	6.9	6.1	6.5
Net animal manure to soil	175.0	160.0	116.2	86.2	81.9	75.0	79.1
Total nitrogen supply to soil	14.4	14.4	14.4	14.4	14.4	13.8	13.2
(manure + fertilizer – net export)	1100.4	1063.8	873.9	743.3	743.2	708.4	724.2
Nitrogen fixation in arable crops	7.8	4.9	4.7	4.5	4.6	4.5	4.1
Nitrogen in crop residues left in field	36.4	34.9	34.1	32.1	30.1	29.6	26.8
Nitrogen in histosols	52.4	52.4	52.4	52.4	52.4	52.4	52.4
Nitrogen in sewage sludge on agric. Land	5.0	1.5	1.5	1.2	1.1	1.0	1.0
Atmospheric deposition agr. NH <sub>3</sub> -N em	195.9	147.5	115.1	100.3	100.3	100.9	95.7
Nitrogen lost through leaching and run off	330.1	319.1	262.2	223.0	223.0	212.5	217.2



## Export and import

Table A8.10a

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Import (mln kg N)					1.1	1.1	2.1	0	0.3	0.7	1.3	0.4	1.9	1.7	2.0	1.9	1.3	1.5	1.4
Export (mln kg N)					22.3	23.1	15.1	11.0	10.1	13.5	16.0	18.4	21.6	13.3	17.5	16.8	17.1	30.3	32.0
Netto export (mln kg N)	6.4	6.8	11.2	15.0	21.2	22.0	13.0	11.0	9.8	12.8	14.7	18.0	19.7	11.6	15.5	14.9	15.8	28.8	30.5

Crop Area (100 m<sup>2</sup>)

Table A8.11

	1990	1995	2000	2005	2006	2007	2008
Winter wheat	13,510,369	12,559,909	12,050,981	11,603,963	12,150,157	12,442,902	14,061,700
Spring wheat	549,904	981,302	1,617,586	2,067,009	1,962,151	1,689,184	1,589,326
Winter barley	994,082	309,977	363,547	296,950	348,824	426,303	466,894
Spring barley	3,044,693	3,248,038	4,353,676	4,761,972	4,109,069	4,172,914	4,556,486
Rye	860,386	817,514	596,058	253,457	238,553	284,508	211,740
Oats	340,128	291,431	240,390	169,744	161,350	170,278	149,107
Triticale		257,947	664,635	408,259	369,430	388,865	319,918
Dried and green peas	1,090,832	69,149	75,204	192,508	57,381	60,594	39,448
Peas (green to harvest)	766,724	713,143	586,657	509,139	530,173	602,662	596,872
Marrowfats	79,350	36,732	38,849	39,585	48,223	27,846	52,318
Kidney beans	373,005	222,094	112,590	109,903	113,854	109,374	91,081
Broad and field beans	316,912	53,220	67,916	44,111	30,702	27,508	31,435
Grass seed	2,631,440	2,189,274	2,196,001	2,763,858	2,614,662	2,010,683	1,566,102
Rape seed	841,501	149,268	85,416	209,640	341,082	335,790	246,658
Caraway seed	34,158	121,059	13,806	9,034	2,920	3,905	3,646
Pop seed	26,356	141,119	58,806	28,286	61,241	50,295	84,182
Flax seed	553,468	440,738	437,930	473,339	442,593	345,633	261,793
Seed potatoes on sand or peat	548,553	536,058	709,599	352,313	340,510	332,014	322,942
Seed potatoes on clay	3,010,113	3,243,815	3,470,553	3,573,898	3,402,330	3,340,892	3,330,423
Potatoes on sand or peat	1,602,484	1,845,122	2,563,153	1,926,935	1,839,859	2,042,278	1,903,856
Potatoes on clay	6,086,924	6,170,599	6,180,900	4,656,037	5,107,937	5,204,137	5,026,336
Industrial potatoes	6,283,773	6,134,453	5,095,818	5,069,191	4,959,220	4,798,038	4,603,383
Sugar beets	12,499,462	11,608,057	11,099,810	9,131,265	8,278,170	8,202,608	7,223,132
Fodder beets	302,286	157,602	89,094	53,195	35,837	33,121	35,320
Lucerne	596,017	583,627	661,606	587,842	644,052	589,808	491,808
Green maize	20,181,089	21,921,725	20,532,074	23,508,819	21,803,584	22,155,358	24,344,498
Green manure	728,159	1,224,765	261,452	3,101,990	1,814,294	1,655,421	757,969
Grain maize		900,542	2,029,838	2,074,849	1,977,153	1,934,033	2,213,225
Corn cob mix		500,473	721,918	667,841	750,828	719,952	759,763
Chicory			475,596	433,848	236,232	258,617	340,924
Hemp			79,197	10,043	2,675	13,515	27,832
Onions	1,282,770	1,608,194	1,997,942	2,252,034	2,463,418	2,617,775	2,614,035
Other horticultural crops	808,437	598,220	1,088,320	1,186,888	1,006,004	875,290	834,797

	1990	1995	2000	2005	2006	2007	2008
<i>Strawberry</i>	186,688	176,313	174,568	230,089	295,921	296,381	292,561
<i>Endive</i>	23,392	27,629	25,198	27,971	27,812	33,169	28,795
<i>Asparagus</i>	266,313	232,356	208,408	233,366	246,053	238,333	247,717
<i>Gherkin</i>	25,738				18,190	25,293	41,912
<i>Cabbage for preservation</i>	157,620	178,353	152,753	139,794			
<i>Cauliflower</i>	236,792	242,970	216,038	239,408	266,682	263,256	253,917
<i>Broccoli</i>		53,379	84,602	131,115	148,472	158,682	173,165
<i>Cabbage (spring and autumn)</i>	100,151	113,850	101,629	107,505	273,562	286,360	306,418
<i>Celeriac</i>	136,263	141,421	128,519	112,772	122,737	138,531	132,991
<i>Beetroot</i>		35,349	29,015	27,619	35,872	36,976	40,473
<i>Lettuce</i>	95,475	104,217	108,978	130,353	159,596	191,860	207,633
<i>Leeks</i>	287,307	385,356	318,448	272,537	304,710	306,271	301,226
<i>Scorzonera</i>	139,536	148,006	113,796	86,697	91,680	99,631	95,895
<i>Spinach</i>	115,291	96,500	120,827	91,431	117,221	130,226	117,525
<i>Brussels sprouts</i>	480,319	438,811	483,409	309,508	335,443	335,223	324,297
<i>Industrial French beans</i>	369,501	467,764	362,736	425,410	389,430	375,083	342,903
<i>Runner beans</i>	22,493				10,903	6,842	7,090
<i>Broad beans green</i>	117,770	87,716	69,416	78,984	151,700	154,794	183,795
<i>Carrot</i>	302,983	327,442	298,512	255,140	273,072	264,776	265,776
<i>Winter carrot (Danvers)</i>	295,050	467,490	472,875	470,043	593,583	547,765	528,567
<i>Witlof</i>	591,896	388,881	419,858	342,321	359,158	347,839	316,209
<i>Other outside horticultural crops</i>	277,358	286,665	317,125	431,248	379,975	317,252	311,478

	N content kg N/ha	Crop residue Frac	N fixation
Winter wheat	28	0.1	
Spring wheat	28	0.1	
Winter barley	19	0.1	
Spring barley	19	0.1	
Rye	16	0.1	
Oats	19	0.1	
Triticale	24	0.1	
Dried and green peas	74	1.0	164
Peas (green to harvest)	194	1.0	164
Marrowfats	74	1.0	164
Kidney beans	74	1.0	164
Broad and field beans	16	1.0	325
Grass seed	28	1.0	
Rape seed	42	1.0	
Caraway seed	37	1.0	
Pop seed	20	1.0	
Flax seed	23	1.0	
Seed potatoes on sand or peat	26	1.0	
Seed potatoes on clay	26	1.0	
Potatoes on sand or peat	26	1.0	
Potatoes on clay	26	1.0	
Industrial potatoes	26	1.0	
Sugar beets	174	1.0	
Fodder beets	92	1.0	
Lucerne	23	1.0	422
Green maize	22	0.1	
Green manure	80	1.0	
Grain maize	70	1.0	
Corn cob mix	70	1.0	
Chicory	40	1.0	
Hemp	40	1.0	
Onions	4	1.0	
Other horticultural crops	40	1.0	
Strawberry	23	1.0	
Endive	78	1.0	
Asparagus	24	1.0	
Gherkin	78	1.0	
Cabbage for preservation	206	1.0	
Cauliflower	89	1.0	
Broccoli	89	1.0	
Cabbage (spring and autumn)	206	1.0	
Celeriac	78	1.0	
Beetroot	78	1.0	
Lettuce	25	1.0	
Leeks	62	1.0	
Scorzonera	78	1.0	
Spinach	62	1.0	
Brussels sprouts	206	1.0	
Industrial French beans	61	1.0	75
Runner beans	61	1.0	75
Broad beans green	13	1.0	185
Carrot	99	1.0	
Winter carrot (Danvers)	99	1.0	
Witloof	78	1.0	
Other outside horticultural crops	78	1.0	

# Annex 9

## Chemical compounds, global warming potentials, units and conversion factors

### A9.1 Chemical compounds

$CF_4$	Perfluoromethane (tetrafluoromethane)
$C_2F_6$	Perfluoroethane (hexafluoroethane)
$CH_4$	Methane
$CO$	Carbon monoxide
$CO_2$	Carbon dioxide
$HCFCs$	Hydrochlorofluorocarbons
$HFCs$	Hydrofluorocarbons
$HNO_3$	Nitric Acid
$NH_3$	Ammonia
$NO_x$	Nitrogen oxide ( $NO$ and $NO_2$ ), expressed as $NO_2$
$N_2O$	Nitrous oxide
$NMVOC$	Non-Methane Volatile Organic Compounds
$PFCs$	Perfluorocarbons
$SF_6$	Sulphur hexafluoride
<sup>2</sup>	Sulphur dioxide
$VOC$	Volatile Organic Compounds (may include or exclude methane)

### A9.2 Global Warming Potentials for selected greenhouse gases

Gas	Atmospheric lifetime	20-year GWP	100-year GWP <sup>1)</sup>	500-year GWP
$CO_2$	Variable (50-200)	1	1	1
$CH_4$ <sup>2)</sup>	12±3	56	21	6.5
$N_2O$	120	280	310	170
<i>HFCs</i> <sup>3)</sup> :				
<i>HFC-23</i>	264	9100	11700	9800
<i>HFC-32</i>	5.6	2100	650	200
<i>HFC-125</i>	32.6	4600	2800	920
<i>HFC-134a</i>	10.6	3400	1300	420
<i>HFC-143a</i>	48.3	5000	3800	1400
<i>HFC-152a</i>	1.5	460	140	42
<i>HFC-227ea</i>	36.5	4300	2900	950
<i>HFC-236fa</i>	209	5100	6300	4700
<i>HFC-245ca</i>	6.6	1800	560	170
<i>PFCs</i> <sup>3)</sup> :				
$CF_4$	50000	4400	6500	10000
$C_2F_6$	10000	6200	9200	14000
$C_3F_8$	2600	4800	7000	10100
$C_4F_{10}$	2600	4800	7000	10100
$C_6F_{14}$	3200	5000	7400	10700
$SF_6$	3200	16300	23900	34900

Source: IPCC (1996)

<sup>1)</sup> GWPs calculated with a 100-year time horizon (indicated in the shaded column) and from the SAR are used in this report (thus not of the Third Assessment Report), in compliance with the UNFCCC Guidelines for reporting (UNFCCC, 1999). Gases indicated in italics are not emitted in the Netherlands.

<sup>2)</sup> The GWP of methane includes the direct effects and the indirect effects due to the production of tropospheric ozone and stratospheric water vapour; the indirect effect due to the production of  $CO_2$  is not included.

### A9.3 Units

<i>MJ</i>	Mega Joule ( $10^6$ Joule)
<i>GJ</i>	Giga Joule ( $10^9$ Joule)
<i>TJ</i>	Tera Joule ( $10^{12}$ Joule)
<i>PJ</i>	Peta Joule ( $10^{15}$ Joule)
<i>Mg</i>	Mega gramme ( $10^6$ gramme)
<i>Gg</i>	Giga gramme ( $10^9$ gramme)
<i>Tg</i>	Tera gramme ( $10^{12}$ gramme)
<i>Pg</i>	Peta gramme ( $10^{15}$ gramme)
<i>ton</i>	metric ton (= 1 000 kilogramme = 1 Mg)
<i>kton</i>	kiloton (= 1 000 metric ton = 1 Gg)
<i>Mton</i>	Megaton (= 1 000 000 metric ton = 1 Tg)
<i>ha</i>	hectare (= $10^4$ m <sup>2</sup> )
<i>kha</i>	kilo hectare (= 1 000 hectare = $10^7$ m <sup>2</sup> = 10 km <sup>2</sup> )
<i>mln</i>	million (= $10^6$ )
<i>mld</i>	milliard (= $10^9$ )

### A9.4 Other conversion factors for emissions

From element basis to full molecular mass:		From full molecular mass to element basis	
$C \rightarrow CO_2$ :	$\times 44/12 = 3.67$	$CO_2 \rightarrow C$ :	$\times 12/44 = 0.27$
$C \rightarrow CH_4$ :	$\times 16/12 = 1.33$	$CH_4 \rightarrow C$ :	$\times 12/16 = 0.75$
$C \rightarrow CO$ :	$\times 28/12 = 2.33$	$CO \rightarrow C$ :	$\times 12/28 = 0.43$
$N \rightarrow N_2O$ :	$\times 44/28 = 1.57$	$N_2O \rightarrow N$ :	$\times 28/44 = 0.64$
$N \rightarrow NO$ :	$\times 30/14 = 2.14$	$NO \rightarrow N$ :	$\times 14/30 = 0.47$
$N \rightarrow NO_2$ :	$\times 46/14 = 3.29$	$NO_2 \rightarrow N$ :	$\times 14/46 = 0.30$
$N \rightarrow NH_3$ :	$\times 17/14 = 1.21$	$NH_3 \rightarrow N$ :	$\times 14/17 = 0.82$
$N \rightarrow HNO_3$ :	$\times 63/14 = 4.50$	$HNO_3 \rightarrow N$ :	$\times 14/63 = 0.22$
$S \rightarrow S_2$ :	$\times 64/32 = 2.00$	$S_2 \rightarrow S$ :	$\times 32/64 = 0.50$

# Annex 10

## List of abbreviations

TAD	Activity Data	EPA	US Environmental Protection Agency
AE	Anode Effect	ER-I	Emission Registration-Individual firms
AWMS	Animal Waste Management Systems	ERT	Expert Review Team
BAK	Monitoring report of gas consumption of small users	ET	Emissions Trading
BEES	Order governing combustion plant emissions requirements (1992) (in Dutch: “Besluit Emissie-Eisen Stookinstallaties”)	ETC/ACC	European Topic Centre on Air and Climate Change
BEK	Monitoring report of electricity consumption of small users	ETS	Emission Trading System
BF	Blast Furnace (gas)	EU	European Union
BOD	Biological Oxygen Demand	EZ	Ministry of Economic Affairs
C	Confidential (notation key in CRF)	FAD	Forest According to Definition
CO	Coke Oven (gas)	FAO	Food and Agricultural Organization (UN)
CS	Country-Specific (notation key in CRF)	F-gases	Group of fluorinated compounds comprising HFCs, PFCs and SF6
Cap	capita (person)	FGD	Flue Gas Desulphurization
CBS	Statistics Netherlands	FO-I	Dutch Facilitating Organization for Industry
CDM	Clean Development Mechanism (one of three mechanisms of the Kyoto Protocol)	GE	Gross Energy
CHP	Combined Heat and Power	GHG	Greenhouse Gas
CLRTAP	Convention on Long-range Transboundary Air Pollution (UN-ECE)	GPG	Good Practice Guidance
CORINAIR	CORe INventory AIR emissions	GIS	Gas Insulated Switchgear
CRF	Common Reporting Format (of emission data files, annexed to an NIR)	GWP	Global Warming Potential
CRT	Continuous Regeneration Trap	HBO	Heating Oil
DLO	Legal name of Wageningen University and Research Centre (Wageningen UR)	HDD	Heating-Degree Day
DM	Dry Matter	HFO	Heavy Fuel Oil
DOC	Degradable Organic Carbon	HOSP	Timber Production Statistics and Forecast (in Dutch: “Hout Oogst Statistiek en Prognose oogstbaar hout”)
DOCF	Degradable Organic Carbon Fraction	IE	Included Elsewhere (notation key in CRF)
EC-LNV	National Reference Centre for Agriculture	IEA	International Energy Agency
ECE	Economic Commission for Europe (UN)	IEF	Implied Emission Factor
ECN	Energy Research Centre of the Netherlands	INK	Dutch Institute for Quality Management
EEA	European Environment Agency	IPCC	Intergovernmental Panel on Climate Change
EF	Emission Factor	KNMI	Royal Netherlands Meteorological Institute
EGR	Exhaust Gas Recirculation	LEI	Agricultural Economics Institute
EIT	Economies-In-Transition (countries from the former SU and Eastern Europe)	LHV	Lower Heating Value
EMEP	European program for Monitoring and Evaluation of long-range transmission of air Pollutants	LNV	Ministry of Agriculture, Nature and Food Quality
EMS	Emission Monitor Shipping	LPG	Liquefied Petroleum Gas
EMSG	Emissions Registration Steering Group	LTO	Landing and Take-Off
ENINA	Task Group Energy, Industry and Waste Handling	LUCF	Land Use Change and Forestry
		LULUCF	Land Use, Land Use Change and Forestry
		MCF	Methane Conversion Factor
		MFV	Measuring Network Functions (in Dutch: “Meetnet Functievervulling”)
		MJV	Annual Environmental Report

MNP	Netherlands Environmental Assessment Agency (in Dutch: “Milieu- en Natuur Planbureau”)	TNO	Netherlands Organization for Applied Scientific Research
MR	Methane Recovery	TBFRA	Temperate and Boreal Forest Resources Assessment (ECE-FAO)
MSW	Municipal Solid Waste	TOF	Trees outside Forests
MW	Mega Watt	UN	United Nations
NA	Not Available; Not Applicable (notation key in CRF); also: National Approach	UNECE	United Nations Economic Commission for Europe
NACE	Statistical Classification of Economic Activities from the European Union: Nomenclature générale des Activités économiques dans les Communautés Européennes.	UNEP	United Nations Environment Program
NAM	Nederlandse Aardolie Maatschappij	UNFCCC	United Nation’s Framework Convention on Climate Change
NAV	Dutch Association of Aerosol Producers	VOC	Volatile Organic Compound
ND	No Data	VROM	Ministry of Housing, Spatial Planning and the Environment
NDF	Neutral Detergent Fiber	VS	Volatile Solids
NE	Not Estimated (notation key in CRF)	V&W	Ministry of Transport, Public Works and Water Management
NEAT	Non-Energy CO <sub>2</sub> emissions Accounting Tables (model of NEU-CO <sub>2</sub> Group)	WBCSD	World Business Council for Sustainable Development
NEC	National Emission Ceilings	WEB	Working Group Emission Monitoring of Greenhouse Gases
NGL	Natural Gas Liquids	WEM	Working Group Emission Monitoring
NIE	National Inventory Entity	WIP	Waste Incineration Plant
NIR	National Inventory Report (annual greenhouse gas inventory report to UNFCCC)	WUR	Wageningen University and Research Centre (or: Wageningen UR)
NLR	National Aerospace Laboratory	WWTP	Wastewater Treatment Plant
NOGEPa	Netherlands Oil and Gas Exploration and Production Association		
NOP-MLK	National Research Program on Global Air Pollution and Climate Change		
NS	Dutch Railways		
ODS	Ozone Depleting Substances		
ODU	Oxidized During Use (of direct non-energy use of fuels or of petrochemical product)		
OECD	Organization for Economic Cooperation and Development		
OM	Organic Matter		
OX	Oxygen Furnace (gas)		
PBL	Netherlands Environmental Assessment Agency		
PRTR	Pollutant Release and Transfer Register		
QA	Quality Assurance		
QC	Quality Control		
RA	Reference Approach (vs. Sectoral or National Approach)		
RIVM	National Institute for Public Health and the Environment		
RIZA	National Institute of Water Management and Waste Treatment		
ROB	Reduction Program on Other Greenhouse Gases		
SA	Sectoral Approach; also: National Approach (vs. Reference Approach)		
SBI	Standaard bedrijven indeling (NACE)		
SCR	Selective Catalytic Reduction		
SBSTA	Subsidiary Body for Scientific and Technological Advice (of Parties to the UNFCCC)		
SGHP	Shell Gasification and Hydrogen Production		
SNCR	Selective Non-Catalytic Reduction		
SW	Streefwaarde (Dutch for “target value”)		
SWDS	Solid Waste Disposal Site		



# References

- Amstel, A.R. van, J.G.J. Olivier, P.G. Ruysenaars (eds.), 2000a: Monitoring of Greenhouse Gases in the Netherlands: Uncertainty and Priorities for Improvement. Proceedings of a National Workshop held in Bilthoven, the Netherlands, 1 September 1999. WIMEK report/ RIVM report No. 773201003. Bilthoven.
- AVV, 2003: Emissie Monitoring Scheepvaart. Protocolen voor de berekening van emissies door scheepvaart. AVV Rotterdam.
- De Baedts, E.E.A. et al., 2001: Koudemiddelgebruik in Nederland. STEK, Baarn.
- Bank, M.P. van de, en H.M. Venderbosch, 1997: Sectorstudie bouwmaterialen, Adromi, NDS—96-013 NEEDIS-studie.
- Bannink, A., J. Dijkstra, J.A.N. Mills, E. Kebreab, J. France, 2005a: Nutritional strategies to reduce enteric methane formation in dairy cows. pp. 367-376. In: Emissions from European Agriculture. Eds. T. Kuczynski, U. Dämmgen, J. Webb and A. Myczko. Wageningen Academic Publishers, Wageningen, the Netherlands
- Bannink, A., J. Kogut, J. Dijkstra, E. Kebreab, J. France, A.M. Van Vuuren, S. Tamminga, 2005b: Estimation of the stoichiometry of volatile fatty acid production in the rumen of lactating cows. *Journal of Theoretical Biology*, in press.
- Bannink, A. 2008. Methane emission from Dutch dairy cows in 2006; estimate of the national average and its uncertainty. ASG rapport, Lelystad.
- Bannink, A., 2010: Methane emissions from enteric fermentation by dairy cows, 1990-2008. Background document on the calculation method and uncertainty analysis for the Dutch National Inventory Report on Greenhouse Gas emissions. ASG Report, Lelystad
- Berdowski, J.J.M., G.P.J. Draaijers, L.H.J.M. Janssen, J.C.Th. Hollander, M. van Loon, M.G.M. Roemer, A.T. Vermeulen, M. Vosbeek, H. Visser, 2001: Sources, Regional Scaling and Validation of Methane Emissions from the Netherlands and Northwest Europe. NOP, NOP-MLK Series, RIVM Report No. 410200084, Bilthoven.
- Boonekamp, P.G.M., H. Mannaerts, H.H.J. Vreuls, B. Wesseling, 2001: Protocol Monitoring Energiebesparing. ECN, ECN Report No. ECN-C-01-129, Petten; RIVM Report No. 408137005, Bilthoven.
- Brouwer J.G.H., J.H.J. Hulskotte, J.A. Annema, 1995: Afsteken van vuurwerk, WESP Report No. C3. RIVM Report No. 772414005, Bilthoven.
- Bruggen, C. van, 2006: Dierlijke mest en mineralen 2004. CBS, Voorburg.
- CBS, 2005a: Herziening duurzame energie 1990-2004. CBS, Voorburg/Heerlen.
- Coenen, P.W.H.G., Memorandum on recalculations as presented in the CRF submission 2006. TNO, Apeldoorn.
- Denier van der Gon, H.A.C., J.H.J. Hulskotte, 2002: Emissiefactoren van methaan en di-stikstofoxide uit luchtvaart en zeescheepvaart. TNO-MEP, TNO-Report No. R2002/294, Apeldoorn.
- DHV, 2000: Overview and explanation of results of project 'Identification of unknown sources of other greenhouse gases' (in Dutch). NOVEM, Utrecht, December 2000. Project No. 3743990070. DHV, DHV-Report No. ML-TB2000 0178, Amersfoort.
- DHV, 2002: Quality Assurance and Quality Control for the Dutch National Inventory Report. Report Phase 1. Report No. ML-BB-20010367, DHV, Amersfoort.
- Dirkse, G.M., W.P. Daamen, H. Schoonderwoerd, J.M. Paasman, 2003: Meetnet Functievervulling bos - Het Nederlandse bos 2001-2002. Expertisecentrum LNV, Rapport EC-LNV No. 2003/231, Ede.
- Dijkstra, J., H.D.St.C. Neal, D.E. Beever, J. France, 1992: Simulation of nutrient digestion, absorption and outflow in the rumen: model description. *Journal of Nutrition*, No. 122, p.2239-2256.
- Emission registration, Methodological report 37, 1997, update in Methodological report 37b, 2002
- Ecofys, Assessment of uncertainties and QA/QC procedures in the Dutch GHG Inventory Report, Utrecht, 2010 (in preparation)
- EPA, 2001: Candles and incense as potential sources of indoor air pollution: Market analysis and Literature review. EPA, Washington DC, Rapport No. EPA-600/R-01-001.
- EZ, 2000: Meerjarenafspraken Energie-efficiency, Resultaten 1999. Ministry of Economic Affairs, The Hague.
- Gense, N.J.L. and R.J. Vermeulen, 2002: N<sub>2</sub>O emissions from passenger cars. TNO report 02.OR.VM.016.1/NG, TNO Automotive, Delft.
- Grontmij 2000: Industrieplan-2. Uitvoering Milieubeleid Olie- en gaswinningsindustrie(NOGEPa) 1999-2002, IMP-2 definitief., 15 november 2000, SJ/CdV, De Bilt.
- Groot, W.J.M.de, R. Visschers, E. Kiestra, P.J. Kuikman and G.J. Nabuurs, 2005: National system to report to the UNFCCC on carbon stock and change of carbon stock related to land use and changes in land use in the Netherlands (in Dutch). Alterra, Alterra-rapport 1035-3.r, Wageningen.
- Groot, W.J.M. de, E. Kiestra, F. de Vries, P.J. Kuikman, 2005: National system of Greenhouse Gas Reporting for Land Use and Land Use Change: Carbon stock changes in the Netherlands due to land use changes 1990-2000. Alterra, Alterra report 1035-III, Wageningen.
- Guis, B., R. de Ridder, P.J. Zijlema, 2009: Verklaring verschillen tussen CO<sub>2</sub>-emissies in EU-ETS en andere rapportages, available at SenterNovem, Utrecht.
- Hanschke C.B. ea Peer review of the Dutch NIR 2010 for the category transport, ECN, ECN\_BS—10-06, February 2010, Petten
- Harmelen, A.K. van, W.R.R. Koch, 2002: CO<sub>2</sub> emission factors for fuels in the Netherlands. TNO, Apeldoorn.
- Heslinga, D.C. and A.K. van Harmelen, 2006. Vaststellingsmethodieken CO<sub>2</sub> emissiefactoren voor aardgas in Nederland. TNO, Rapport no. R.2006/ Project no. 64101, Apeldoorn.
- Hoek, K. W. van der, 2002: Uitgangspunten voor de mest- en ammoniakberekeningen 1999 tot en met 2001 zoals gebruikt in de Milieubalans 2001 en 2002, inclusief dataset landbouwemissies 1980-2001. RIVM rapport 773004013, Bilthoven.
- Hoek, K.W. van der and M.W. van Schijndel, 2006. Methane and nitrous oxide emissions from animal manure management, 1990-2003. Background document on the calculation method for the Dutch NIR. RIVM Report No. 680125002, MNP report 500080002, Bilthoven, the Netherlands.
- Hoek, K.W. van der, M.W. van Schijndel, P.J. Kuikman, 2007. Direct and indirect nitrous oxide emissions from agricultural soils, 1990-2003. Background document on the calculation method for the Dutch NIR. RIVM Report No. 680125003, MNP report 500080003, Bilthoven, the Netherlands.
- Hulskotte, J., 2004a: Protocol voor de jaarlijkse bepaling van de emissies van specifieke defensie-activiteiten conform de IPCC-richtlijnen. TNO-MEP, Apeldoorn.
- Hulskotte, J., 2004b: Protocol voor de vaststelling van de broeikasgasemissies van de visserij in Nederland conform de IPCC-richtlijnen. TNO-MEP, Apeldoorn.
- IPCC, 1996: IPCC Second Assessment. Climate Change 1995. IPCC, Geneva
- IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Office, Bracknell, UK.
- IPCC, 2001: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, IPCC-TSU NGGIP, Japan.
- IPCC, 2003: LUCF Sector Good Practice Guidance. Penman et al. (Eds.), IPCC Good practice Guidance for Land Use, Land Use Change and Forestry. IPCC NGGIP Programme. Publ. by IGES for IPCC. Japan.
- Joosen and De Jager: Afvalverbrandingsinstallaties, Notitie in het kader van de Marktmonitoring Duurzame Energie, Ecofys, EEPo2011, Augustus, 2003.
- Klein J., G. Geilenkirchen (PBL), A. Hoen, J. Hulskotte, N. van Duynhoven, R. de Lange, A. Hensema, D. Broekhuizen, H. Molnár-in 't Veld, Methoden voor de berekening van de emissies door mobiele bronnen in Nederland, november 2009
- Klein Goldewijk, K., J.G.J. Olivier, J.A.H.W. Peters, P.W.H.W. Coenen, H.H.J. Vreuls, 2004: Greenhouse Gas Emissions in the Netherlands 1990-2002, National Inventory Report 2004. RIVM Report 773201008, Bilthoven.
- Klein Goldewijk, K., J.G.J. Olivier, J.A.H.W. Peters, P.W.H.G. Coenen, H.H.J. Vreuls, 2005: Greenhouse Gas Emissions in the Netherlands 1990-2003. National Inventory Report 2005, RIVM Report No. 773201009, Bilthoven.
- Kramer, H. W. and Knol, 2005: Historisch grondgebruik in Nederland: een landelijk reconstructie van het grondgebruik 1990. Alterra, Alterra rapport 1035-4, Wageningen.

- Kramer, H., J. Oldengarm, G. van Dorland, I.J.J. van den Wyngaert and G.J. van den Born, (in preparation): Kyoto Land use and Land use Change. Alterra, Wageningen.
- Kroeze, C., 1994: Nitrous oxide. Emission inventory and options for control in the Netherlands. RIVM Report No. 773001004, Bilthoven.
- Kuikman, P.J., W.J.M. de Groot, R. Hendriks, A. Verhagen and F.J. de Vries (2002) Stocks of C in soils and emissions of CO<sub>2</sub> from agricultural soils in the Netherlands. Alterra, Wageningen-UR, Wageningen. Alterra rapport 561.
- Kuikman, P.J., van den Akker J.J.H., F. de Vries, 2005: Emissions of N<sub>2</sub>O and CO<sub>2</sub> from organic agricultural soils, Alterra, Wageningen UR. Alterra rapport 1035-2, Wageningen.
- Kuikman, P.J., K.W. van der Hoek, A. Smit, K. Zwart, 2006. Update of emission factors for direct emissions of nitrous oxide from agricultural soils on the basis of measurements in the Netherlands. Alterra rapport 1217, Alterra, Wageningen
- LEI/CBS, 2000: Land- en tuinbouwcijfers (Agriculture and horticulture statistics). On [www.lei.wur.nl](http://www.lei.wur.nl) LEI, Den Haag and on [www.CBS.nl](http://www.CBS.nl) CBS, Voorburg.
- Mills, J.A.N., J. Dijkstra, A. Bannink, S.B. Cammell, E. Kebreab, J. France, 2001: A mechanistic model of whole-tract digestion and methanogenesis in the lactating dairy cow: Model development, evaluation and application. *Journal of Animal Science* 79, 1584-1597.
- Monteny G.J., 2009: Audit en review van de emissieregistratie voor de Taakgroep Landbouw en Landgebruik. Met focus op de niet-CO<sub>2</sub>-broeikasgassen methaan en lachgas, Monteny Milieu Advies, Wageningen.
- Nabuurs, G.J., I.J. van den Wyngaert, W.D. Daamen, A.T.F. Helmink, W. de Groot, W.C. Knol, H. Kramer, P. Kuikman. 2005. National system of greenhouse gas reporting for forest and nature areas under UNFCCC in the Netherlands. Alterra report 1035-1. Alterra, Wageningen.
- Neelis, M., M. Patel, M. de Feber, 2003: Improvement of CO<sub>2</sub> emission estimates from the non-energy use of fossil fuels in the Netherlands, report commissioned by the Netherlands' agency for Energy and the Environment (NOVEM) and the Netherlands' Ministry of Housing, Spatial Planning and the Environment (VROM). Utrecht University, Copernicus Institute/Dept. of Science, Technology and Society, Utrecht.
- Neelis, M.L., M.K. Patel, P.W. Bach, W.G. Haije, 2005: Analysis of energy use and carbon losses in the chemical and refinery industries, report ECN-I-05-008, Energy research Centre of the Netherlands, Unit Energy Efficiency in Industry, Petten, the Netherlands, August 2005, 82 pp.
- Neelis, M., 2006: Peer review of the 'Industrial Processes' and 'Solvent and other product use' chapters of the draft National Inventory Report 2006. Copernicus Institute for Sustainable Development and Innovation, Utrecht University. Report NWS-E-2006-6.
- Neelis, M., P. Blinde, 2009: Emission from industrial processes: expert review of the draft Dutch National Inventory Report 2009. Ecofys International BV, Utrecht.
- Olivier, J.G.J., 2004: Note on Netherlands' CO<sub>2</sub> emission factors for petrol, diesel and LPG. Version 2, December 2004. MNP document No. M/773201/01/NI, Bilthoven.
- Olivier, J.G.J., L.J. Brandes, R.A.B. te Molder, 2009 (in print): Estimate of annual and trend uncertainty for Dutch sources of greenhouse gas emissions using the IPCC Tier 1 approach. PBL Report 500080013, PBL, Bilthoven.
- Olsthoorn, X. and A. Pielaat, 2003: Tier-2 uncertainty analysis of the Dutch greenhouse gas emissions 1999. Institute for Environmental Studies (IVM), Free University, Amsterdam. IVM Report No. Ro3 06.
- Oonk, 1994: Validation of landfill gas formation models, Oonk, H., A. Weenk, O. Coops, L. Luning, TNO Institute of Environmental and Energy Technology, December 1994, reference number 94-315
- Oonk et al., 2004: Methaan- en lachgasemissies uit afvalwater, TNO, Apeldoorn, TNO report R2004/486
- PBL, 2008: Werkplan 2007-2008. PBL, Bilthoven.
- PBL, 2009: Werkplan ER ronde 2010. PBL, Bilthoven.
- Pulles, H., 2000: Structuurschema Regionale en Kleine Luchthavens. Deel 1, CO<sub>2</sub>-berekeningswijze en -resultaten, Rijksluchtvaartdienst, Den Haag.
- Ramírez-Ramírez, A., C. de Keizer, J.P. van der Sluijs, 2006: Monte Carlo Analysis of Uncertainties in the Netherlands Greenhouse Gas Emission Inventory for 1990-2004. Department of Science, Technology and Society, Copernicus Institute for Sustainable Development and Innovation, Utrecht University, report NWS-E-2006-58, Utrecht, the Netherlands; July 2006.
- Reumerman, P.J. and Frederiks, B., 2002: Charcoal production with reduced emissions. 12th European Conference on Biomass for Energy, Industry and Climate Protection, Amsterdam.
- Riemersma, I.J., K. Jordaan, J. Oonk, 2003: N<sub>2</sub>O emissions of Heavy Duty vehicles. Netherlands Organisation for Applied Scientific Research (TNO), Delft.
- RIVM, 1999: Meten, rekenen en onzekerheden. De werkwijze van het RIVM Milieuonderzoek, RIVM Report No. 408129 005 (main report and addendum), Bilthoven.
- Roemer M., Th. Thijsse, T. van der Meulen, 2003: Verificatie van methaan emissies. ArenA, Journal of the Netherlands Association of Environmental Professionals (VVM), Den Bosch.
- Roemer M. and O. Tarasova, 2002: Methane in the Netherlands - an exploratory study to separate time scales. TNO report R2002/215. TNO, Apeldoorn.
- Segers, R., 2005: Herziening duurzame energie 1990-2004. On [www.cbs.nl](http://www.cbs.nl), June, 27th, 2005, Voorburg.
- Segers, R. and M. Wilmer, 2007: Renewable energy 2006. Explanation of updated preliminary figures (in Dutch). CBS, Voorburg, July 2007.
- SenterNovem, 2005c, Description of the National System (elaborated in collaboration with involved institutes. Also included in the Initial Report of the Netherlands for the calculation of its assigned amount under the Kyoto and the UNFCCC [VROM, 2006])
- SenterNovem 2006 Advies nieuwe emissiefactor voor aardgas vanaf 1990
- SenterNovem, 2008, QA/QC Programme 2008/2009 (elaborated in cooperation with ER).
- SenterNovem, 2008: Afvalverwerking in Nederland, gegevens 2007, SenterNovem Uitvoering Afvalbeheer, november 2008, 3UA0819, isbn978-90-5748-069-0..
- SenterNovem 2009, The Netherlands National System: QA/QC programme 2009/2010 Version 5.0 December 2009. SenterNovem, Utrecht.
- Smink, W., 2005: Calculated methane production from enteric fermentation in cattle excluding dairy cows. FIS background document. SenterNovem, Utrecht.
- Smink, W., K.W. van der Hoek, A. Bannink, J. Dijkstra, 2005: Calculation of methane production from enteric fermentation in dairy cows. SenterNovem, Utrecht.
- Spakman, J., Van Loon, M.M.J., Van der Auweraert, R.J.K., Gielen, D.J., Olivier, J.G.J., E.A. Zonneveld, 2003: Method for calculating greenhouse gas emissions. VROM-HIMH, The Hague. Report Emission Registration No. 37b, March 2003. Electronic update of original report No. 37 of July 1997. Only electronically available both in Dutch and in English at: [www.greenhousegases.nl](http://www.greenhousegases.nl).
- Spoelstra, H., 1993: N<sub>2</sub>O-emissions from combustion processes used in the generation of electricity. KEMA, Arnhem/RIVM, NOP report no. 410100049, Bilthoven.
- Tamminga, S., F. Aarts, A. Bannink, O. Oenema, G.J. Monteny, 2004: Actualisering van geschatte N en P excreties door rundvee. Reeks Milieu en Landelijk gebied 25. Wageningen.
- TNO, 2002: CO<sub>2</sub> emission factors for fuels in the Netherlands, TNO, Apeldoorn. TNO Report No. R2002/174.
- TNO 2006 Vaststellingsmethodieken voor CO<sub>2</sub> emissiefactoren van aardgas in Nederland, rapport 2006-A-R0079/B
- TNO, 2004: Uncertainty assessment of NO<sub>x</sub>, 2 and NH<sub>3</sub> emissions in the Netherlands. TNO Environment, Energy and Process Innovation, Apeldoorn.
- UNFCCC, 1999: UNFCCC Guidelines for reporting and review. UNFCCC Secretariat, Bonn. Doc. No FCCC/CP/1999/7. January 2000.
- UNFCCC, 2004: Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, part I: UNFCCC reporting guidelines on annual inventories (following incorporation of the provisions of Decision 13/CP.9). UNFCCC Secretariat, Bonn. Doc. No. FCCC/SBSTA/2004/8.
- Veldt, C. and P.F.J. van der Most, 1993: Emissiefactoren vluchtige organische stoffen uit verbrandingsmotoren. VROM, The Hague, Publikatiereeks Emissieregistratie No. 10.
- Velthof, G.L., J. Mosquera, J. Huis in 't Veld, E. Hummelink, 2010: Effect of manure application technique on nitrous oxide emission from agricultural soils. Wageningen UR, Livestock Research
- Visser, H., 2005: The significance of climate change in the Netherlands An analysis of historical and future trends (1901-2020) in weather conditions, weather extremes and temperature-related impacts. MNP Report No. 550002007, Bilthoven.
- Volkers, C.H. en K.E.L. Smekens, 2008; Kan het beter? Review van de taakgroep ENINA van de Emissieregistratie. ECN report ECN-E-08-047, ECN, Petten.
- Vreuls, H.H.J., 2006: Advies voor het gebruik van een nieuwe emissiefactor voor aardgas vanaf 1990. SenterNovem, Notitie 22 Maart 2006.
- Vreuls, H.H.J., 2006: The Netherlands: list of fuels and standard CO<sub>2</sub> emission factors, SenterNovem.
- Vreuls, H.H.J., and P.J. Zijlema, 2009: The Netherlands: list of fuels and standard CO<sub>2</sub> emission factors, version December 2009, SenterNovem.
- De Vries, F., 2004: The expansion of peat soils (In Dutch: De verbreiding van veengronden). In: Kekem, A.J. van (red.). Veengronden en stikstofleverend vermogen. Alterra, Wageningen. Alterra report 965.

- VROM, 2005: Fourth Netherlands' National Communication under the United Nations Convention on Climate Change, Ministry of VROM, The Hague.
- VROM 2006b: The Netherlands' Report on demonstrable Progress under article 3.2 of the Kyoto Protocol, Ministry of VROM, The Hague.
- VROM, 2006: Initial Report of the Netherlands for the calculation of its assigned amount under the Kyoto and the UNFCCC.
- WBCSD/WRI (World Business Council for Sustainable Development/World Resources Institute), 2004: Calculating Direct GHG Emissions from Primary Aluminium Metal Production. Guide to calculation worksheets. Available at 'GHG Protocol Initiative' website: <http://www.ghgprotocol.org/standard/tools.htm>.
- Wyngaert, I.J.J. van den, W. de Groot, P. Kuikman, G.J. Nabuurs, 2006: Updates of the Dutch National System for greenhouse gas reporting of the LULUCF sector. Alterra report 1035-5. Alterra, Wageningen.
- Wyngaert, I.J.J. van den, H. Kramer, P. Kuikman, and G.J. Nabuurs, (in preparation): Greenhouse gas reporting of the LULUCF sector, revisions and updates related to the Dutch NIR 2009, Alterra, Wageningen.
- Zijlema, P.J., 2008: Berekening van de standaard CO<sub>2</sub>-emissiefactor aardgas t.b.v. kalenderjaar 2007, SenterNovem, UEMBo804422, 28, Utrecht.
- Zijlema, P.J., 2009: Berekening van de standaard CO<sub>2</sub>-emissiefactor aardgas t.b.v. kalenderjaar 2008, SenterNovem, UEMBo917552, Utrecht.
- Zijlema P.J., 2010: Verklaring verschillen tussen CO<sub>2</sub>-emissies in EU-ETS en andere rapportages t.b.v. NIR 2010, Utrecht, 2010, available at NL Agency

### Emissions of greenhouse gases stabilized in 2008

The total greenhouse gas emission (excluding LULUCF) in 2008 stabilized at the level of 2007 although the emission of N<sub>2</sub>O dropped significantly (- 3.7 Tg CO<sub>2</sub> equivalents) due to measures in the nitric acid production. This decline was counteracted by an increase in the emissions of CO<sub>2</sub> and CH<sub>4</sub>.

In 2008, total direct greenhouse gas emissions (excluding emissions from LULUCF) in the Netherlands amount to 206.9 Tg CO<sub>2</sub> eq. This is three percent below the emissions in the base year (213.3 Tg CO<sub>2</sub> eq).

This report documents the 2009 Netherlands' annual submission of its greenhouse gas emission inventory in accordance with the guidelines provided by the United Nations Framework Convention on Climate Change (UNFCCC), the Kyoto Protocol and the European Union's Greenhouse Gas Monitoring Mechanism.

The report comprises explanations of observed trends in emissions; a description of an assessment of key sources and their uncertainty; documentation of methods, data sources and emission factors applied; and a description of the quality assurance system and the verification activities performed on the data.