

Annexes

ANNEX 1: Key Categories 572

| | | |
|-------|---|-----|
| A 1.1 | GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES | 572 |
| A 1.2 | QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES | 573 |
| A 1.3 | QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES | 573 |
| A 1.4 | QUANTITATIVE APPROACH 2 KCA FOLLOWING IPCC 2006 GUIDELINES | 616 |
| A 1.5 | KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM | 638 |
| A 1.6 | APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES | 640 |
| A 1.7 | USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY | 640 |
| A 1.8 | TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3 | 641 |

ANNEX 2: Assessment of Uncertainty 643

| | | |
|---------|--|-----|
| A 2.1 | ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1) | 643 |
| A 2.1.1 | Key Categories | 643 |
| A 2.1.2 | Tables of uncertainty estimates from the error propagation approach | 644 |
| A 2.2 | ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2) | 652 |
| A 2.2.1 | Overview of the Method | 652 |
| A 2.2.2 | Methodological details of the Monte Carlo model | 652 |
| A 2.2.3 | Quality Control Checks on the Monte Carlo Model Output | 653 |
| A 2.3 | UNCERTAINTY PARAMETERS | 654 |
| A 2.3.1 | Uncertainty Parameters used | 654 |
| A 2.3.2 | General Considerations | 670 |
| A 2.3.3 | Uncertainty in the Trend | 670 |
| A 2.4 | UNCERTAINTIES IN GWP WEIGHTED EMISSIONS | 671 |
| A 2.4.1 | Uncertainty in the emissions | 671 |
| A 2.4.2 | Uncertainty in the Trend | 671 |
| A 2.5 | SECTORAL UNCERTAINTIES | 673 |
| A 2.5.1 | Overview of the Method | 673 |

| | | |
|-------|---|-----|
| A 2.6 | COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES | 678 |
|-------|---|-----|

ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities. 680

| | | |
|----------|--|-----|
| A 3.1 | ENERGY 680 | |
| A 3.1.1 | Emission factors | 680 |
| A 3.1.2 | Commonly used references | 680 |
| A 3.1.3 | Feedstocks and Non-Energy Use (NEU) of fuels | 683 |
| A 3.1.4 | Aviation (MS 7) | 691 |
| A 3.1.5 | Gas leakage | 694 |
| A 3.2 | INDUSTRIAL PROCESSES (CRF SECTOR 2) | 695 |
| A 3.3 | AGRICULTURE (CRF SECTOR 3) | 695 |
| A 3.3.1 | Enteric Fermentation (3A) | 696 |
| A 3.3.2 | Manure Management (3B) | 697 |
| A 3.3.3 | Agricultural Soils (3D) | 702 |
| A 3.4 | LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4) | 711 |
| A 3.4.1 | Carbon stock changes due to afforestation and forest management (4A) | 717 |
| A 3.4.2 | Land Use Change and Soils (4B, 4C, 4E) | 728 |
| A 3.4.3 | Changes in stocks of carbon in non-forest biomass due to management and land use change (4B2, 4C2, 4E2) | 737 |
| A 3.4.4 | Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G) | 742 |
| A 3.4.5 | Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C) | 745 |
| A 3.4.6 | Emissions from drainage (organic soils) (4B1, 4C1) | 749 |
| A 3.4.7 | Emissions of N2O due to disturbance associated with land use conversion and land management changes (4(III)) | 749 |
| A 3.4.8 | On-site and off-site emissions from peat extraction (4D) | 750 |
| A 3.4.9 | Flooded Lands (4D) | 753 |
| A 3.4.10 | Harvested Wood Products (4G) | 754 |
| A 3.4.11 | Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs) | 755 |

| | | |
|-----------------|--|------------|
| | A 3.4.12 Uncertainty analysis of the LULUCF sector | 762 |
| A 3.5 | WASTE (CRF SECTOR 5) | 763 |
| | A 3.5.1 Solid Waste Disposal on Land (5A) | 763 |
| | A 3.5.2 Biological Treatment of Solid Waste (5B) | 772 |
| | A 3.5.3 Waste Incineration (5C) | 774 |
| | A 3.5.4 Wastewater Handling (5D) | 777 |
| A 3.6 | UK CROWN DEPENDENCIES AND OVERSEAS TERRITORIES | 781 |
| | A 3.6.1 Overview of Data Sources | 781 |
| | A 3.6.2 Activity and Emissions Data | 781 |
| ANNEX 4: | National Energy Balance | 788 |
| A 4.1 | UK ENERGY BALANCE | 788 |
| A 4.2 | FUELS DATA | 793 |
| A 4.2.1 | Reallocations of energy data and differences from UK energy statistics | 794 |
| ANNEX 5: | Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information. | 810 |
| A 5.1 | ANNUAL INVENTORY SUBMISSION | 810 |
| A 5.2 | SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1 | 810 |
| ANNEX 6: | Comparison of Emission Estimates Using Atmospheric Observations | 811 |
| A 6.1 | MODELLING APPROACH USED FOR COMPARISON WITH THE UK GHGI | 811 |
| A 6.2 | METHANE | 812 |
| A 6.3 | NITROUS OXIDE | 814 |
| A 6.4 | HYDROFLUOROCARBONS | 816 |
| | A 6.4.1 HFC-134a | 816 |
| | A 6.4.2 HFC-152a | 817 |
| | A 6.4.3 HFC-125 | 818 |
| | A 6.4.4 HFC-143a | 819 |

| | | |
|-----------------|--|------------|
| A 6.4.5 | HFC-23 | 820 |
| A 6.4.6 | HFC-32 | 822 |
| A 6.4.7 | HFC-43-10mee | 823 |
| A 6.4.8 | HFC-227ea | 824 |
| A 6.4.9 | HFC-365mfc | 825 |
| A 6.4.10 | HFC-245fa | 826 |
| A 6.5 | PERFLUOROCARBONS | 827 |
| A 6.5.1 | PFC-14 | 827 |
| A 6.5.2 | PFC-116 | 828 |
| A 6.5.3 | PFC-218 | 829 |
| A 6.5.4 | PFC-318 | 830 |
| A 6.6 | SULPHUR HEXAFLUORIDE | 831 |
| A 6.7 | NITROGEN TRIFLUORIDE | 832 |
| A 6.8 | CARBON DIOXIDE | 833 |
| ANNEX 7: | Analysis of EU ETS Data | 834 |
| A 7.1 | INTRODUCTION | 834 |
| A 7.2 | BACKGROUND | 837 |
| A 7.2.1 | EU ETS Data and GHG Inventories | 837 |
| A 7.2.2 | Scope of the UK EU ETS and Implications for the GHG Inventory | 838 |
| A 7.2.3 | Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration | 840 |
| A 7.3 | DATA PROCESSING | 841 |
| A 7.4 | EU ETS DATA COVERAGE | 843 |
| A 7.5 | EU ETS DATA USE IN THE UK GHGI | 845 |
| A 7.5.1 | Activity and Emissions Data | 845 |
| A 7.5.2 | Implied Emission Factors | 849 |
| ANNEX 8: | UK Domestic Emissions Reporting Requirements | 861 |
| A 8.1 | NATIONAL STATISTICS | 862 |

ANNEX 9: End User Emissions 863

| | | |
|-------|--|-----|
| A 9.1 | INTRODUCTION | 863 |
| A 9.2 | DEFINITION OF END USERS | 863 |
| A 9.3 | OVERVIEW OF THE END USER CALCULATIONS | 864 |
| A 9.4 | EXAMPLE END USER CALCULATION | 865 |
| A 9.5 | END USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY | 868 |
| A 9.6 | DETAILED EMISSIONS ACCORDING TO END USER CATEGORIES | 919 |

Table of Tables:

| | | |
|---------------|---|-----|
| Table A 1.3.1 | Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF) | 575 |
| Table A 1.3.2 | Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF) | 581 |
| Table A 1.3.3 | Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF) | 587 |
| Table A 1.3.4 | Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF) | 593 |
| Table A 1.3.5 | Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF) | 599 |
| Table A 1.3.6 | Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF) | 608 |
| Table A 1.5.1 | KCA Ranking..... | 638 |
| Table A 1.8.1 | Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol | 641 |
| Table A 2.1.1 | Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year | 644 |
| Table A 2.3.1 | Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO ₂) inventory | 655 |
| Table A 2.3.2 | Estimated uncertainties in the activity data and emission factors used in the methane (CH ₄) inventory | 663 |
| Table A 2.3.3 | Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N ₂ O) inventory..... | 666 |
| Table A 2.3.4 | Estimated uncertainties in the activity data and emission factors used in the F-gas inventory..... | 668 |
| Table A 2.4.1 | Summary of Monte Carlo Uncertainty Estimates | 672 |

| | | |
|----------------|--|-----|
| Table A 2.5.1 | Sectoral Uncertainty Estimates..... | 674 |
| Table A 2.6.1 | Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses | 679 |
| Table A 3.1.1 | Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics..... | 684 |
| Table A 3.1.2 | CAA aircraft types assigned to EMEP-EEA Emissions Inventory Guidebook aircraft types..... | 691 |
| Table A 3.1.3 | Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers | 695 |
| Table A 3.3.1 | Livestock Population Data by Animal Type ('000 animal places) | 696 |
| Table A 3.3.2 | Methane Emission Factors for Livestock Emissions for 2017 | 696 |
| Table A 3.3.3 | Methane conversion factors for Manure Management Systems in the UK | 697 |
| Table A 3.3.4 | Nitrogen Excretion Factors, kg N animal place ⁻¹ year ⁻¹ for livestock in the UK (1990-2017)..... | 698 |
| Table A 3.3.5 | Distribution of Animal Waste Management Systems (%) used for Different Animal types, 2017 | 699 |
| Table A 3.3.6 | Nitrous Oxide Emission Factors for Animal Waste Handling Systems | 701 |
| Table A 3.3.7 | EF for direct N ₂ O emissions from managed soils in the UK inventory | 702 |
| Table A 3.3.8 | Areas of UK Crops and quantities of fertiliser applied for 2017 | 702 |
| Table A 3.3.9 | Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2017..... | 704 |
| Table A 3.3.10 | Parameter values for crop residue management. | 706 |
| Table A 3.3.11 | N concentrations in above and below ground biomass | 707 |
| Table A 3.3.12 | Mineralised N from soils | 709 |
| Table A 3.4.1 | Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce (<i>P. Sitchensis</i>), yield class 12. | 718 |
| Table A 3.4.2 | Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 based on estimates of woodland area by age from the NFI and administrative records. | 725 |
| Table A 3.4.3 | Grouping of MLC land cover types for soil carbon change modelling..... | 728 |
| Table A 3.4.4 | Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling..... | 729 |
| Table A 3.4.5 | Sources of land use change data used to estimate changes in soil carbon in Great Britain for different periods..... | 729 |

| | | |
|----------------|--|-----|
| Table A 3.4.6 | Sources of land use change data used to estimate changes in soil carbon in Northern Ireland for different periods. | 730 |
| Table A 3.4.7 | Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK | 731 |
| Table A 3.4.8 | Weighted average change in equilibrium soil carbon density (t ha^{-1}) to 1 m deep for changes between different land types in England | 733 |
| Table A 3.4.9 | Weighted average change in equilibrium soil carbon density (t ha^{-1}) to 1 m deep for changes between different land types in Scotland | 733 |
| Table A 3.4.10 | Weighted average change in equilibrium soil carbon density (t ha^{-1}) to 1 m deep for changes between different land types in Wales | 733 |
| Table A 3.4.11 | Weighted average change in equilibrium soil carbon density (t ha^{-1}) to 1 m deep for changes between different land types in Northern Ireland | 733 |
| Table A 3.4.12 | Rates of change of soil carbon for land use change transitions | 734 |
| Table A 3.4.13 | Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W)..... | 734 |
| Table A 3.4.14 | Aggregation of Agricultural Census crop types for estimating biomass carbon stock changes from Cropland Management | 738 |
| Table A 3.4.15 | Biomass stock factors for UK Cropland types | 739 |
| Table A 3.4.16 | Aggregation of Countryside Survey Broad Habitats for estimating biomass carbon stock changes from Grassland Management | 739 |
| Table A 3.4.17 | Biomass stock factors for UK Grassland types | 739 |
| Table A 3.4.18 | Mean biomass carbon stock densities, tC/ha | 740 |
| Table A 3.4.19 | Weighted average change in equilibrium biomass carbon density (kg m^{-2}) for changes between different land types in England..... | 741 |
| Table A 3.4.20 | Weighted average change in equilibrium biomass carbon density (kg m^{-2}) for changes between different land types in Scotland. | 741 |
| Table A 3.4.21 | Weighted average change in equilibrium biomass carbon density(kg m^{-2}) for changes between different land types in Wales. | 741 |
| Table A 3.4.22 | Weighted average change in equilibrium biomass carbon density (kg m^{-2}) for changes between different land types in Northern Ireland. | 741 |
| Table A 3.4.23 | Countryside Survey data for Forest conversion | 743 |
| Table A 3.4.24 | Corrected Forest conversion rates..... | 744 |
| Table A 3.4.25 | IRS database property type descriptions by LULUCF sub-category | 747 |
| Table A 3.4.26 | Activity data for peat extraction sites in Northern Ireland | 750 |
| Table A 3.4.27 | Annual peat production, m^3 for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports)..... | 751 |

| | | |
|----------------|---|-----|
| Table A 3.4.28 | Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies | 757 |
| Table A 3.4.29 | Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies | 757 |
| Table A 3.4.30 | IPCC Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies | 761 |
| Table A 3.5.1 | Composition of waste sorting residues and mixed municipal waste | 764 |
| Table A 3.5.2 | DOC, DOCf and composition of waste materials | 765 |
| Table A 3.5.3 | Amount of waste landfilled (1945 to 2017) | 767 |
| Table A 3.5.4 | Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted..... | 768 |
| Table A 3.5.5 | Parameters used in landfill emission estimates for overseas territories and crown dependencies..... | 770 |
| Table A 3.5.6 | Activity Data: Inputs in the composting process..... | 772 |
| Table A 3.5.7 | Activity Data: Inputs in the anaerobic digestion process | 773 |
| Table A 3.5.8 | Activity Data: UK Waste Incineration | 774 |
| Table A 3.5.9 | Emissions Data: UK Waste Incineration | 775 |
| Table A 3.5.10 | UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data | 777 |
| Table A 3.5.11 | UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors..... | 778 |
| Table A 3.5.12 | UK Private Waste Water Management System Emission Estimates Parameters (5D1)..... | 779 |
| Table A 3.5.13 | UK Industrial Waste Water Treatment Activity Data (5D2) | 780 |
| Table A 3.6.1 | Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO ₂ equivalent)..... | 781 |
| Table A 3.6.2 | Isle of Man, Guernsey and Jersey – Combustion activity data | 782 |
| Table A 3.6.3 | Isle of Man, Guernsey and Jersey – Animal numbers..... | 782 |
| Table A 3.6.4 | Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N ₂ O) | 783 |
| Table A 3.6.5 | Cayman Islands, Falklands Islands, and Bermuda - Emissions of Direct GHGs (Mt CO ₂ equivalent) | 783 |
| Table A 3.6.6 | Cayman Islands, Falklands Islands, and Bermuda – Combustion activity data | 784 |
| Table A 3.6.7 | Cayman Islands, Falklands Islands, and Bermuda – Animal numbers..... | 785 |

| | | |
|----------------|---|-----|
| Table A 3.6.8 | Cayman Islands, Falklands Islands, and Bermuda – Total emissions from Agricultural Soils (kg N ₂ O)..... | 785 |
| Table A 3.6.9 | Cayman Islands, Falklands Islands, and Bermuda - Amount of synthetic fertilizer applied | 785 |
| Table A 3.6.10 | Gibraltar – Emissions of Direct GHGs (Mt CO ₂ equivalent)..... | 786 |
| Table A 3.6.11 | Gibraltar – Combustion activity data | 786 |
| Table A 4.1.1 | UK Energy Balance for 2017 (thousand tonnes of oil equivalent, gross energy basis) | 789 |
| Table A 4.2.1 | Mapping of fuels used in IPCC and the NAEI | 793 |
| Table A 4.2.2 | Fuel reconciliation - coal use in the latest year (Mtonnes)..... | 795 |
| Table A 4.2.3 | Fuel reconciliation – natural gas use in the latest year (Mtherms)..... | 798 |
| Table A 4.2.4 | Fuel reconciliation – Fuel oils use in latest year (Mtonnes) | 801 |
| Table A 4.2.5 | Fuel reconciliation – Gas oil use in latest year (Mtonnes) | 804 |
| Table A 4.2.6 | Fuel reconciliation – Use of petroleum gases in the latest year (Mtherms) | 807 |
| Table A 7.1.1 | Summary of the use of EU ETS data in the UK inventory | 836 |
| Table A 7.5.1 | Refinery Emissions Data Comparison and Revision to OPG Activity | 846 |
| Table A 7.5.2 | EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases)..... | 849 |
| Table A 7.5.3 | Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas) | 851 |
| Table A 7.5.4 | EU ETS data for fuels used at integrated steelworks & coke ovens (Emission Factors in kt/Mt for solid & liquid fuels, kt/Mth for gases) | 853 |
| Table A 7.5.5 | EU ETS data for Fuels used at Cement Kilns (kt / Mt) | 855 |
| Table A 7.5.6 | Comparison of Cement Sector Carbon Dioxide Emissions* within the UK GHGI and the EU ETS | 856 |
| Table A 7.5.7 | EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases)..... | 857 |
| Table A 7.5.8 | EU ETS emission factor data for production of lime (kt / Mt lime produced) | 858 |
| Table A 7.5.9 | EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas)..... | 859 |
| Table A 8.1.1 | Summary table of GHG emissions by NC Category, including net emissions/removals from LULUCF (Mt CO ₂ eq) – National Statistics coverage (UK only) | 862 |

| | | |
|---------------|--|-----|
| Table A 8.1.2 | Summary table of GHG emissions by Gas, including net emissions/removals from LULUCF (Mt CO ₂ eq) – National Statistics coverage (UK only) | 862 |
| Table A 9.4.1 | Example of the outputs during an end user calculation | 867 |
| Table A 9.5.1 | Sources reallocated to end users and the fuels used..... | 869 |
| Table A 9.5.2 | End user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation..... | 872 |
| Table A 9.6.1 | End user emissions from all National Communication categories, MtCO ₂ equivalent..... | 920 |
| Table A 9.6.2 | End user CO ₂ emissions from all National Communication categories, MtCO ₂ equivalent..... | 920 |
| Table A 9.6.3 | End user CH ₄ emissions from all National Communication categories, MtCO ₂ equivalent..... | 921 |
| Table A 9.6.4 | End user N ₂ O emissions from all National Communication categories, MtCO ₂ equivalent..... | 922 |

Table of Figures:

| | | |
|--------------|---|-----|
| Figure A 3.1 | Data flow diagrams for each land use sub-category, showing cross-linkages between sectors: (i) 4A and 4G, (ii) 4B, (iii) 4C, (iv) 4D, (v) 4E | 712 |
| Figure A 3.2 | The CARBINE-SCA model | 722 |
| Figure A 3.3 | Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes). | 731 |
| Figure A 3.4 | Decision tree for assessing the effects of cropland management activities on soil carbon stocks. | 736 |
| Figure A 3.5 | Crop type area in the UK 1989-2017 | 737 |
| Figure A 3.6 | Annual area of FIRMS thermal anomalies for GB for 2001 to the current inventory year (thermal anomalies were filtered to exclude those recorded over urban/industrial areas). | 748 |
| Figure A 3.7 | Time series of wildfire burnt areas in the UK 1990 to the current inventory year | 748 |
| Figure A 3.8 | The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions..... | 763 |
| Figure A 6.1 | Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 812 |
| Figure A 6.2 | Verification of the UK emission inventory estimates for methane in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD+CBW, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ)..... | 813 |

| | | |
|---------------|--|-----|
| Figure A 6.3 | Verification of the UK emission inventory estimates for methane in Gg yr ⁻¹ from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ($\pm 1\sigma$)..... | 814 |
| Figure A 6.4 | Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified..... | 814 |
| Figure A 6.5 | Verification of the UK emission inventory estimates for nitrous oxide in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 815 |
| Figure A 6.6 | Verification of the UK emission inventory estimates for nitrous oxide in Gg yr ⁻¹ from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ($\pm 1\sigma$)..... | 816 |
| Figure A 6.7 | Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 816 |
| Figure A 6.8 | Verification of the UK emission inventory estimates for HFC-134a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 817 |
| Figure A 6.9 | Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 817 |
| Figure A 6.10 | Verification of the UK emission inventory estimates for HFC-152a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 818 |
| Figure A 6.11 | Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 818 |
| Figure A 6.12 | Verification of the UK emission inventory estimates for HFC-125 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 819 |
| Figure A 6.13 | Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 820 |
| Figure A 6.14 | Verification of the UK emission inventory estimates for HFC-143a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 820 |

| | | |
|---------------|---|-----|
| Figure A 6.15 | Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 821 |
| Figure A 6.16 | Verification of the UK emission inventory estimates for HFC-23 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). The second plot is plotted from the year 2000 on an expanded y-axis. | 821 |
| Figure A 6.17 | Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 822 |
| Figure A 6.18 | Verification of the UK emission inventory estimates for HFC-32 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). | 823 |
| Figure A 6.19 | Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 823 |
| Figure A 6.20 | Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). | 824 |
| Figure A 6.21 | Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 824 |
| Figure A 6.22 | Verification of the UK emission inventory estimates for HFC-227ea in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). | 825 |
| Figure A 6.23 | Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 825 |
| Figure A 6.24 | Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). | 826 |
| Figure A 6.25 | Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 826 |
| Figure A 6.26 | Verification of the UK emission inventory estimates for HFC-245fa in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) | |

| | | |
|---------------|--|-----|
| | estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 827 |
| Figure A 6.27 | Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 827 |
| Figure A 6.28 | Verification of the UK emission inventory estimates for PFC-14 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 828 |
| Figure A 6.29 | Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 828 |
| Figure A 6.30 | Verification of the UK emission inventory estimates for PFC-116 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 829 |
| Figure A 6.31 | Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 829 |
| Figure A 6.32 | Verification of the UK emission inventory estimates for PFC-218 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 830 |
| Figure A 6.33 | Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 830 |
| Figure A 6.34 | Verification of the UK emission inventory estimates for PFC-318 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 831 |
| Figure A 6.35 | Monthly Northern Hemisphere trend in SF ₆ estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 831 |
| Figure A 6.36 | Verification of the UK emission inventory estimates for SF ₆ in Gg yr ⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$)..... | 832 |
| Figure A 6.37 | Monthly Northern Hemisphere trend in NF ₃ estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. | 832 |

| | | |
|---------------|---|-----|
| Figure A 6.38 | Monthly Northern Hemisphere trend in CO ₂ estimated from Mace Head observations (blue line). Red line denotes annual trend. Data under grey shading are not yet ratified. | 833 |
| Figure A 9.1 | Simplified fuel flows for an end user calculation..... | 864 |
| Figure A 9.2 | Fuel use in the example calculation..... | 866 |
| Figure A 9.3 | Comparison of 'direct' and end user emissions of carbon according the sectors considered in the end user example..... | 868 |

ANNEX 1: Key Categories

This annex contains the key category analysis for the latest GHG inventory¹. It contains:

- A description of the methodology used for identifying key categories
- Information on the level of disaggregation
- Information to fulfil the reporting requirements of Tables 4.2 and 4.3 of Volume 1 of the 2006 IPCC Guidelines, including and excluding land use, land-use change and forestry (LULUCF).

The annex also contains information relevant to the requirements of reporting under the Kyoto Protocol. The table below contains the additional KP information that Annex 1 needs to contain, and the locations of this information in the Annex².

| Requirements | Locations of the relevant information in this Annex |
|---|---|
| Description of methodology used for identifying key categories, <i>including KP-LULUCF</i> | See sections immediately below including “ <i>General approach used to identify Key Categories</i> ” and “ <i>Approach used to identify KP-LULUCF Key Categories</i> ”. |
| Reference to the key category tables in the CRF | This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category tables (Table 7) in the CRF. |
| Reference to the key category tables in the CRF, <i>including in the KP-LULUCF CRF tables</i> | This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category KP-LULUCF tables (Table NIR 3) in the CRF. |
| Information on the level of disaggregation | The tables in this Annex contain information on the level of disaggregation used. The level of disaggregation follows IPCC 2006 Guidelines. |
| Tables 4.2 to 4.4 of Volume 4 the 2006 IPCC guidelines | The data requested in the 2006 Guidelines tables, including and excluding LULUCF, are provided in Table A 1.3.1 to Table A 1.4.6 and Table 1.7 to Table 1.10 . |
| <i>Table NIR.3, as contained in the annex to decision 6/CMP.3</i> | A facsimile of Table NIR 3, provided in the CRF, is given in Table A 1.8.1 . |

A 1.1 GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES

In the UK inventory, certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. These key source categories have been

¹ Following the requirements to report information about uncertainties as set out in FCCC/CP/2013/10/Add.3. Report of the Conference of the Parties on its nineteenth session, held in Warsaw from 11 to 23 November 2013. Addendum Part two: Action taken by the Conference of the Parties at its nineteenth session.

² The information in this table has been taken directly from the UNFCCC document “Annotated outline of the National Inventory Report including reporting elements under the Kyoto Protocol”.

identified so that the resources available for inventory preparation may be prioritised, and the best possible estimates prepared for the most significant source categories.

The UK completes both quantitative and qualitative Key Category Analyses (KCAs).

The UK has used the method set out in Section 4.3.1 and Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (*Approach 1 to identify key categories*, and *Approach 2 to identify key categories* respectively) to quantitatively determine the key source categories.

The method used in the qualitative KCA is described below, and further descriptions of the methods the UK uses to quantitatively determine key categories are given later in this section.

A 1.2 QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES

Following IPCC good practice, a qualitative analysis of the inventory has been made to identify any additional key source categories, which may not have been identified using the quantitative analysis. The approach set out in Section 4.3.3 of the IPCC 2006 Guidelines has been applied, using the four criteria set out in the guidance, to judge whether a category is a key category. The criteria are:

1. (Use of) mitigation techniques and technologies;
2. Emissions growth (increase or decrease);
3. No quantitative assessment of uncertainties performed;
4. Completeness (examine qualitatively potential key categories that are not yet estimated quantitatively by applying the qualitative considerations above).

In addition, additional criteria have also been taken in account

5. High uncertainty (links to point 3 above);
6. Unexpectedly low or high emissions;
7. External recommendation has also been used as an additional criterion to identify key categories.

The results of the qualitative analysis did not identify any categories that were not already identified by the quantitative key category analysis.

A 1.3 QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES

A key category analysis has been completed for both level and trend. This KCA has been created using the 2006 IPCC Guidelines Approach 1 methodology. The factors that make a source a key category are:

- A high contribution to the level of emissions; and
- A high contribution to the trend;

For example, transport fuel (1A3b) is a key category for carbon dioxide because it is a large source of emissions and nitric acid production (2B2) because it shows a significant trend.

The category groupings are largely aligned to those suggested in Tables 4.5 and 4.6 in Volume 1, Chapter 4 of the 2006 IPCC guidelines, although we deviate in a number of cases, in particular:

- **Agriculture and LULUCF.** In the 2006 guidelines a different nomenclature for categorising agriculture and LULUCF sources and sinks was used compared to the adopted nomenclature, which means that it would be challenging and confusing to retain this categorisation when sources are grouped differently in the adopted nomenclature. The agriculture categories are analysed at the level of mostly 3 digit IPCC codes, whilst analysis for LULUCF sources and sinks is more aggregated. The UK inventory agency considers that the level of aggregation used in the UK method for the KCA is sufficiently detailed to target inventory improvements (e.g. consideration of the more important livestock types under agriculture) whilst not introducing unnecessary computational difficulties (e.g. use of “miscellaneous” categories to mop up the remainder within a sector). Further, the level of source/sink category aggregations in the KCA are aligned to how individual methods or models are used to derive the UK inventory estimates, and are therefore at an appropriate level of detail for the UK inventory
- **Fugitive Emissions.** The suggested categories are at a much more granular level (e.g. 1B2aii) than other sectors. We considered that this would lead to an undue diminishing of these sectors, decreasing their likelihood of being considered key, so have adopted a level of aggregation more consistent with other sectors
- **Miscellaneous emissions.** The suggested approach was to group a large number of small sources into one category. We considered that this would lead to an undue increase in the significance of these sources, increasing their likelihood of being considered key, so have adopted a level of aggregation more consistent with other sectors

The results of the key category analysis with and without LULUCF, for the base year and the latest reported year and for both Approaches 1 and 2 KCA, are summarised by sector and gas in **Section 1.5.1**. The tables indicate whether a key category arises from the level (L1) assessment or the trend (T1) assessment.

The results of the **level assessment** (based on Approach 1) with and without LULUCF for the base year and the latest reported year are shown **Table A 1.3.1** to **Table A 1.3.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the “Level Parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the level parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in level.

The results of the **trend assessment** (based on Approach 1) with and without LULUCF for the base year to the latest reported year are shown in **Table A 1.3.5** and **Table A 1.3.6**. The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use, Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of the “Trend parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the trend parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in trend.

Note that the tables in chapter 1 of the NIR summarise the key categories from both the approach 1 and approach 2 key categories analyses and the aggregations used are slightly different for the

two approaches. The category "3A" is therefore total emissions from category 3A, whilst categories 3A1 and 3A2 have also been identified as key categories in their own right. Category "2B Chemical industries - CO₂" is total CO₂ emissions from category 2B.

Table A 1.3.1 Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF)

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A1 | Energy industries: solid fuels | CO ₂ | 185,492.23 | 185,492.23 | 0.2184 | 0.2184 |
| 1A3b | Road transportation: liquid fuels | CO ₂ | 108,570.16 | 108,570.16 | 0.1278 | 0.3462 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 70,373.19 | 70,373.19 | 0.0829 | 0.4290 |
| 5A | Solid waste disposal | CH ₄ | 60,432.98 | 60,432.98 | 0.0711 | 0.5002 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 40,883.76 | 40,883.76 | 0.0481 | 0.5483 |
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 38,484.20 | 38,484.20 | 0.0453 | 0.5936 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 29,832.00 | 29,832.00 | 0.0351 | 0.6288 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 27,330.36 | 27,330.36 | 0.0322 | 0.6609 |
| 1B1 | Coal mining and handling | CH ₄ | 21,826.68 | 21,826.68 | 0.0257 | 0.6866 |
| 2B3 | Adipic acid production | N ₂ O | 19,934.61 | 19,934.61 | 0.0235 | 0.7101 |
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 19,881.28 | 19,881.28 | 0.0234 | 0.7335 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 19,868.96 | 19,868.96 | 0.0234 | 0.7569 |
| 1A4 | Other sectors: liquid fuels | CO ₂ | 19,607.73 | 19,607.73 | 0.0231 | 0.7800 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 17,784.67 | 17,784.67 | 0.0209 | 0.8009 |
| 4A | Forest land | CO ₂ | -15,026.26 | 15,026.26 | 0.0177 | 0.8186 |
| 4B | Cropland | CO ₂ | 14,265.93 | 14,265.93 | 0.0168 | 0.8354 |
| 3D | Agricultural soils | N ₂ O | 13,610.39 | 13,610.39 | 0.0160 | 0.8514 |
| 1B2 | Oil and gas extraction | CH ₄ | 12,345.07 | 12,345.07 | 0.0145 | 0.8660 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 9,229.34 | 9,229.34 | 0.0109 | 0.8768 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 7,611.13 | 7,611.13 | 0.0090 | 0.8858 |
| 2A1 | Cement production | CO ₂ | 7,295.26 | 7,295.26 | 0.0086 | 0.8944 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|--|--|------------------|------------------|
| 4C | Grassland | CO ₂ | -7,111.03 | 7,111.03 | 0.0084 | 0.9028 |
| 4E | Settlements | CO ₂ | 7,011.30 | 7,011.30 | 0.0083 | 0.9110 |
| 1B2 | Oil and gas extraction | CO ₂ | 5,777.92 | 5,777.92 | 0.0068 | 0.9178 |
| 2C1 | Iron and steel production | CO ₂ | 5,591.54 | 5,591.54 | 0.0066 | 0.9244 |
| 1A5 | Other: liquid fuels | CO ₂ | 5,293.44 | 5,293.44 | 0.0062 | 0.9306 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,935.90 | 4,935.90 | 0.0058 | 0.9364 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,732.53 | 4,732.53 | 0.0056 | 0.9420 |
| 2B8 | Petrochemical and carbon black production | CO ₂ | 4,538.91 | 4,538.91 | 0.0053 | 0.9473 |
| 5D | Wastewater treatment and discharge | CH ₄ | 4,218.90 | 4,218.90 | 0.0050 | 0.9523 |
| 2B2 | Nitric acid production | N ₂ O | 3,860.26 | 3,860.26 | 0.0045 | 0.9569 |
| 3B2 | Manure management from Sheep | N ₂ O | 3,442.70 | 3,442.70 | 0.0041 | 0.9609 |
| 2B1 | Ammonia production | CO ₂ | 1,895.00 | 1,895.00 | 0.0022 | 0.9631 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,881.31 | 1,881.31 | 0.0022 | 0.9654 |
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 1,698.56 | 1,698.56 | 0.0020 | 0.9674 |
| 4G | Harvested wood products | CO ₂ | -1,639.08 | 1,639.08 | 0.0019 | 0.9693 |
| 2A2 | Lime production | CO ₂ | 1,462.05 | 1,462.05 | 0.0017 | 0.9710 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,455.18 | 1,455.18 | 0.0017 | 0.9727 |
| 2C6 | Zinc production | CO ₂ | 1,358.83 | 1,358.83 | 0.0016 | 0.9743 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,312.28 | 1,312.28 | 0.0015 | 0.9759 |
| 5C | Incineration and open burning of waste | CO ₂ | 1,300.71 | 1,300.71 | 0.0015 | 0.9774 |
| 1A4 | Other sectors: solid fuels | CH ₄ | 1,265.22 | 1,265.22 | 0.0015 | 0.9789 |
| 1A3b | Road transportation: liquid fuels | CH ₄ | 1,246.70 | 1,246.70 | 0.0015 | 0.9804 |
| 1A1 | Energy industries: solid fuels | N ₂ O | 1,023.72 | 1,023.72 | 0.0012 | 0.9816 |
| 4B | Cropland | N ₂ O | 1,019.85 | 1,019.85 | 0.0012 | 0.9828 |
| 3G | Liming | CO ₂ | 1,015.18 | 1,015.18 | 0.0012 | 0.9840 |
| 5D | Wastewater treatment and discharge | N ₂ O | 783.59 | 783.59 | 0.0009 | 0.9849 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|--|--|------------------|------------------|
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 783.25 | 783.25 | 0.0009 | 0.9858 |
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 663.00 | 663.00 | 0.0008 | 0.9866 |
| 2A4 | Other process uses of carbonates | CO ₂ | 640.93 | 640.93 | 0.0008 | 0.9873 |
| 4E | Settlements | N ₂ O | 584.68 | 584.68 | 0.0007 | 0.9880 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 552.81 | 552.81 | 0.0007 | 0.9887 |
| 2G3 | N ₂ O from product uses | N ₂ O | 552.57 | 552.57 | 0.0007 | 0.9893 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 531.30 | 531.30 | 0.0006 | 0.9900 |
| 4D | Wetlands | CO ₂ | 486.95 | 486.95 | 0.0006 | 0.9905 |
| 2C3 | Aluminium production | CO ₂ | 450.32 | 450.32 | 0.0005 | 0.9911 |
| 2A3 | Glass production | CO ₂ | 405.54 | 405.54 | 0.0005 | 0.9915 |
| 4 | Indirect N ₂ O emissions from LULUCF | N ₂ O | 401.49 | 401.49 | 0.0005 | 0.9920 |
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 387.17 | 387.17 | 0.0005 | 0.9925 |
| 1A4 | Other sectors: peat | CO ₂ | 372.48 | 372.48 | 0.0004 | 0.9929 |
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 333.43 | 333.43 | 0.0004 | 0.9933 |
| 3H | Urea application to land | CO ₂ | 327.60 | 327.60 | 0.0004 | 0.9937 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 292.27 | 292.27 | 0.0003 | 0.9940 |
| 3A3 | Enteric fermentation from Swine | CH ₄ | 283.06 | 283.06 | 0.0003 | 0.9944 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|--|--|------------------|------------------|
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 278.70 | 278.70 | 0.0003 | 0.9947 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 276.73 | 276.73 | 0.0003 | 0.9950 |
| 1A1 | Energy industries: other fuels | CO ₂ | 246.13 | 246.13 | 0.0003 | 0.9953 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 241.68 | 241.68 | 0.0003 | 0.9956 |
| 2B7 | Soda ash production | CO ₂ | 231.55 | 231.55 | 0.0003 | 0.9959 |
| 4A | Forest land | N ₂ O | 231.22 | 231.22 | 0.0003 | 0.9961 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 224.74 | 224.74 | 0.0003 | 0.9964 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 198.91 | 198.91 | 0.0002 | 0.9966 |
| 3F | Field burning of agricultural residues | CH ₄ | 186.57 | 186.57 | 0.0002 | 0.9968 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 184.49 | 184.49 | 0.0002 | 0.9971 |
| 2B10 | Other Chemical Industry | CH ₄ | 174.65 | 174.65 | 0.0002 | 0.9973 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 157.18 | 157.18 | 0.0002 | 0.9975 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 148.72 | 148.72 | 0.0002 | 0.9976 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 140.41 | 140.41 | 0.0002 | 0.9978 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 138.62 | 138.62 | 0.0002 | 0.9980 |
| 5C | Incineration and open burning of waste | CH ₄ | 134.83 | 134.83 | 0.0002 | 0.9981 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 132.00 | 132.00 | 0.0002 | 0.9983 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 105.00 | 105.00 | 0.0001 | 0.9984 |
| 2B6 | Titanium dioxide production | CO ₂ | 104.63 | 104.63 | 0.0001 | 0.9985 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 103.32 | 103.32 | 0.0001 | 0.9986 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 92.30 | 92.30 | 0.0001 | 0.9987 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 76.86 | 76.86 | 0.0001 | 0.9988 |
| 1A4 | Other sectors: biomass | CH ₄ | 62.37 | 62.37 | 0.0001 | 0.9989 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 57.79 | 57.79 | 0.0001 | 0.9990 |
| 3F | Field burning of agricultural residues | N ₂ O | 57.66 | 57.66 | 0.0001 | 0.9990 |
| 1A5 | Other: liquid fuels | N ₂ O | 56.12 | 56.12 | 0.0001 | 0.9991 |
| 1A1 | Energy industries: solid fuels | CH ₄ | 49.32 | 49.32 | 0.0001 | 0.9992 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 47.59 | 47.59 | 0.0001 | 0.9992 |
| 5C | Incineration and open burning of waste | N ₂ O | 46.66 | 46.66 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 44.43 | 44.43 | 0.0001 | 0.9993 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 43.36 | 43.36 | 0.0001 | 0.9994 |
| 2G4 | Other product manufacture and use | N ₂ O | 41.00 | 41.00 | 0.0000 | 0.9994 |
| 1B2 | Oil and gas extraction | N ₂ O | 40.75 | 40.75 | 0.0000 | 0.9995 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 37.47 | 37.47 | 0.0000 | 0.9995 |
| 2C1 | Iron and steel production | CH ₄ | 36.90 | 36.90 | 0.0000 | 0.9996 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 36.88 | 36.88 | 0.0000 | 0.9996 |
| 2A4 | Other process uses of carbonates | CH ₄ | 31.10 | 31.10 | 0.0000 | 0.9997 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 29.98 | 29.98 | 0.0000 | 0.9997 |
| 1A4 | Other sectors: peat | CH ₄ | 26.35 | 26.35 | 0.0000 | 0.9997 |
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 19.54 | 19.54 | 0.0000 | 0.9997 |
| 1A1 | Energy industries: other fuels | CH ₄ | 18.55 | 18.55 | 0.0000 | 0.9998 |
| 5B | Biological treatment of solid waste | CH ₄ | 18.13 | 18.13 | 0.0000 | 0.9998 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.80 | 17.80 | 0.0000 | 0.9998 |
| 2C1 | Iron and steel production | N ₂ O | 17.74 | 17.74 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 14.54 | 14.54 | 0.0000 | 0.9998 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A3c | Railways: liquid fuels | N ₂ O | 13.75 | 13.75 | 0.0000 | 0.9999 |
| 5B | Biological treatment of solid waste | N ₂ O | 12.97 | 12.97 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 12.29 | 12.29 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 12.20 | 12.20 | 0.0000 | 0.9999 |
| 4C | Grassland | N ₂ O | 10.33 | 10.33 | 0.0000 | 0.9999 |
| 4C | Grassland | CH ₄ | 9.97 | 9.97 | 0.0000 | 0.9999 |
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 9.56 | 9.56 | 0.0000 | 0.9999 |
| 1A4 | Other sectors: biomass | N ₂ O | 9.14 | 9.14 | 0.0000 | 0.9999 |
| 1A1 | Energy industries: other fuels | N ₂ O | 6.72 | 6.72 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 6.37 | 6.37 | 0.0000 | 1.0000 |
| 4D | Wetlands | N ₂ O | 4.13 | 4.13 | 0.0000 | 1.0000 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 3.66 | 3.66 | 0.0000 | 1.0000 |
| 1A5 | Other: liquid fuels | CH ₄ | 3.56 | 3.56 | 0.0000 | 1.0000 |
| 4E | Settlements | CH ₄ | 3.00 | 3.00 | 0.0000 | 1.0000 |
| 4A | Forest land | CH ₄ | 2.89 | 2.89 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 2.79 | 2.79 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 2.46 | 2.46 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 2.10 | 2.10 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 1.47 | 1.47 | 0.0000 | 1.0000 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 1.41 | 1.41 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: biomass | CH ₄ | 0.47 | 0.47 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 0.38 | 0.38 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.31 | 0.31 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|--------------|--|------------------|--|--|------------------|------------------|
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.29 | 0.29 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.26 | 0.26 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: biomass | N ₂ O | 0.25 | 0.25 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 0.16 | 0.16 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling biomass | CH ₄ | 0.10 | 0.10 | 0.0000 | 1.0000 |
| 4B | Cropland | CH ₄ | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.08 | 0.08 | 0.0000 | 1.0000 |
| Total | | | 801,848.54 | 849,401.28 | 1.0000 | |

Table A 1.3.2 Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF)

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|------------------|--|--|------------------|------------------|
| 1A1 | Energy industries: solid fuels | CO ₂ | 185,492.23 | 185,492.23 | 0.2314 | 0.3668 |
| 1A3b | Road transportation: liquid fuels | CO ₂ | 108,570.16 | 108,570.16 | 0.1354 | 0.3668 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 70,373.19 | 70,373.19 | 0.0878 | 0.4546 |
| 5A | Solid waste disposal | CH ₄ | 60,432.98 | 60,432.98 | 0.0754 | 0.5300 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 40,883.76 | 40,883.76 | 0.0510 | 0.5810 |
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 38,484.20 | 38,484.20 | 0.0480 | 0.6290 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 29,832.00 | 29,832.00 | 0.0372 | 0.6663 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 27,330.36 | 27,330.36 | 0.0341 | 0.7004 |
| 1B1 | Coal mining and handling | CH ₄ | 21,826.68 | 21,826.68 | 0.0272 | 0.7276 |
| 2B3 | Adipic acid production | N ₂ O | 19,934.61 | 19,934.61 | 0.0249 | 0.7525 |
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 19,881.28 | 19,881.28 | 0.0248 | 0.7773 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 19,868.96 | 19,868.96 | 0.0248 | 0.8020 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|--|--|------------------|------------------|
| 1A4 | Other sectors: liquid fuels | CO ₂ | 19,607.73 | 19,607.73 | 0.0245 | 0.8265 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 17,784.67 | 17,784.67 | 0.0222 | 0.8487 |
| 3D | Agricultural soils | N ₂ O | 13,610.39 | 13,610.39 | 0.0170 | 0.8657 |
| 1B2 | Oil and gas extraction | CH ₄ | 12,345.07 | 12,345.07 | 0.0154 | 0.8811 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 9,229.34 | 9,229.34 | 0.0115 | 0.8926 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 7,611.13 | 7,611.13 | 0.0095 | 0.9021 |
| 2A1 | Cement production | CO ₂ | 7,295.26 | 7,295.26 | 0.0091 | 0.9112 |
| 1B2 | Oil and gas extraction | CO ₂ | 5,777.92 | 5,777.92 | 0.0072 | 0.9184 |
| 2C1 | Iron and steel production | CO ₂ | 5,591.54 | 5,591.54 | 0.0070 | 0.9254 |
| 1A5 | Other: liquid fuels | CO ₂ | 5,293.44 | 5,293.44 | 0.0066 | 0.9320 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,935.90 | 4,935.90 | 0.0062 | 0.9381 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,732.53 | 4,732.53 | 0.0059 | 0.9440 |
| 2B8 | Petrochemical and carbon black production | CO ₂ | 4,538.91 | 4,538.91 | 0.0057 | 0.9497 |
| 5D | Wastewater treatment and discharge | CH ₄ | 4,218.90 | 4,218.90 | 0.0053 | 0.9550 |
| 2B2 | Nitric acid production | N ₂ O | 3,860.26 | 3,860.26 | 0.0048 | 0.9598 |
| 3B2 | Manure management from Sheep | N ₂ O | 3,442.70 | 3,442.70 | 0.0043 | 0.9641 |
| 2B1 | Ammonia production | CO ₂ | 1,895.00 | 1,895.00 | 0.0024 | 0.9664 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,881.31 | 1,881.31 | 0.0023 | 0.9688 |
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 1,698.56 | 1,698.56 | 0.0021 | 0.9709 |
| 2A2 | Lime production | CO ₂ | 1,462.05 | 1,462.05 | 0.0018 | 0.9727 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,455.18 | 1,455.18 | 0.0018 | 0.9745 |
| 2C6 | Zinc production | CO ₂ | 1,358.83 | 1,358.83 | 0.0017 | 0.9762 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,312.28 | 1,312.28 | 0.0016 | 0.9779 |
| 5C | Incineration and open burning of waste | CO ₂ | 1,300.71 | 1,300.71 | 0.0016 | 0.9795 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A4 | Other sectors: solid fuels | CH ₄ | 1,265.22 | 1,265.22 | 0.0016 | 0.9811 |
| 1A3b | Road transportation: liquid fuels | CH ₄ | 1,246.70 | 1,246.70 | 0.0016 | 0.9826 |
| 1A1 | Energy industries: solid fuels | N ₂ O | 1,023.72 | 1,023.72 | 0.0013 | 0.9839 |
| 3G | Liming | CO ₂ | 1,015.18 | 1,015.18 | 0.0013 | 0.9852 |
| 5D | Wastewater treatment and discharge | N ₂ O | 783.59 | 783.59 | 0.0010 | 0.9861 |
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 783.25 | 783.25 | 0.0010 | 0.9871 |
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 663.00 | 663.00 | 0.0008 | 0.9879 |
| 2A4 | Other process uses of carbonates | CO ₂ | 640.93 | 640.93 | 0.0008 | 0.9887 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 552.81 | 552.81 | 0.0007 | 0.9894 |
| 2G3 | N ₂ O from product uses | N ₂ O | 552.57 | 552.57 | 0.0007 | 0.9901 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 531.30 | 531.30 | 0.0007 | 0.9908 |
| 2C3 | Aluminium production | CO ₂ | 450.32 | 450.32 | 0.0006 | 0.9913 |
| 2A3 | Glass production | CO ₂ | 405.54 | 405.54 | 0.0005 | 0.9919 |
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 387.17 | 387.17 | 0.0005 | 0.9923 |
| 1A4 | Other sectors: peat | CO ₂ | 372.48 | 372.48 | 0.0005 | 0.9928 |
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 333.43 | 333.43 | 0.0004 | 0.9932 |
| 3H | Urea application to land | CO ₂ | 327.60 | 327.60 | 0.0004 | 0.9936 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 292.27 | 292.27 | 0.0004 | 0.9940 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|--|--|------------------|------------------|
| 3A3 | Enteric fermentation from Swine | CH ₄ | 283.06 | 283.06 | 0.0004 | 0.9943 |
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 278.70 | 278.70 | 0.0003 | 0.9947 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 276.73 | 276.73 | 0.0003 | 0.9950 |
| 1A1 | Energy industries: other fuels | CO ₂ | 246.13 | 246.13 | 0.0003 | 0.9953 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 241.68 | 241.68 | 0.0003 | 0.9956 |
| 2B7 | Soda ash production | CO ₂ | 231.55 | 231.55 | 0.0003 | 0.9959 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 224.74 | 224.74 | 0.0003 | 0.9962 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 198.91 | 198.91 | 0.0002 | 0.9965 |
| 3F | Field burning of agricultural residues | CH ₄ | 186.57 | 186.57 | 0.0002 | 0.9967 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 184.49 | 184.49 | 0.0002 | 0.9969 |
| 2B10 | Other Chemical Industry | CH ₄ | 174.65 | 174.65 | 0.0002 | 0.9971 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 157.18 | 157.18 | 0.0002 | 0.9973 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 148.72 | 148.72 | 0.0002 | 0.9975 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 140.41 | 140.41 | 0.0002 | 0.9977 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 138.62 | 138.62 | 0.0002 | 0.9979 |
| 5C | Incineration and open burning of waste | CH ₄ | 134.83 | 134.83 | 0.0002 | 0.9980 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 132.00 | 132.00 | 0.0002 | 0.9982 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 105.00 | 105.00 | 0.0001 | 0.9983 |
| 2B6 | Titanium dioxide production | CO ₂ | 104.63 | 104.63 | 0.0001 | 0.9985 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 103.32 | 103.32 | 0.0001 | 0.9986 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 92.30 | 92.30 | 0.0001 | 0.9987 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 76.86 | 76.86 | 0.0001 | 0.9988 |
| 1A4 | Other sectors: biomass | CH ₄ | 62.37 | 62.37 | 0.0001 | 0.9989 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 57.79 | 57.79 | 0.0001 | 0.9990 |
| 3F | Field burning of agricultural residues | N ₂ O | 57.66 | 57.66 | 0.0001 | 0.9990 |
| 1A5 | Other: liquid fuels | N ₂ O | 56.12 | 56.12 | 0.0001 | 0.9991 |
| 1A1 | Energy industries: solid fuels | CH ₄ | 49.32 | 49.32 | 0.0001 | 0.9992 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 47.59 | 47.59 | 0.0001 | 0.9992 |
| 5C | Incineration and open burning of waste | N ₂ O | 46.66 | 46.66 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 44.43 | 44.43 | 0.0001 | 0.9993 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 43.36 | 43.36 | 0.0001 | 0.9994 |
| 2G4 | Other product manufacture and use | N ₂ O | 41.00 | 41.00 | 0.0001 | 0.9994 |
| 1B2 | Oil and gas extraction | N ₂ O | 40.75 | 40.75 | 0.0001 | 0.9995 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 37.47 | 37.47 | 0.0000 | 0.9995 |
| 2C1 | Iron and steel production | CH ₄ | 36.90 | 36.90 | 0.0000 | 0.9996 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 36.88 | 36.88 | 0.0000 | 0.9996 |
| 2A4 | Other process uses of carbonates | CH ₄ | 31.10 | 31.10 | 0.0000 | 0.9997 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 29.98 | 29.98 | 0.0000 | 0.9997 |
| 1A4 | Other sectors: peat | CH ₄ | 26.35 | 26.35 | 0.0000 | 0.9997 |
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 19.54 | 19.54 | 0.0000 | 0.9998 |
| 1A1 | Energy industries: other fuels | CH ₄ | 18.55 | 18.55 | 0.0000 | 0.9998 |
| 5B | Biological treatment of solid waste | CH ₄ | 18.13 | 18.13 | 0.0000 | 0.9998 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.80 | 17.80 | 0.0000 | 0.9998 |
| 2C1 | Iron and steel production | N ₂ O | 17.74 | 17.74 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 14.54 | 14.54 | 0.0000 | 0.9999 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|--|--|------------------|------------------|
| 1A3c | Railways: liquid fuels | N ₂ O | 13.75 | 13.75 | 0.0000 | 0.9999 |
| 5B | Biological treatment of solid waste | N ₂ O | 12.97 | 12.97 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 12.29 | 12.29 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 12.20 | 12.20 | 0.0000 | 0.9999 |
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 9.56 | 9.56 | 0.0000 | 0.9999 |
| 1A4 | Other sectors: biomass | N ₂ O | 9.14 | 9.14 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: other fuels | N ₂ O | 6.72 | 6.72 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 6.37 | 6.37 | 0.0000 | 1.0000 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 3.66 | 3.66 | 0.0000 | 1.0000 |
| 1A5 | Other: liquid fuels | CH ₄ | 3.56 | 3.56 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 2.79 | 2.79 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 2.46 | 2.46 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 2.10 | 2.10 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 1.47 | 1.47 | 0.0000 | 1.0000 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 1.41 | 1.41 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: biomass | CH ₄ | 0.47 | 0.47 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 0.38 | 0.38 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.31 | 0.31 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.29 | 0.29 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.26 | 0.26 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: biomass | N ₂ O | 0.25 | 0.25 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 0.16 | 0.16 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|--------------|--------------------------------------|------------------|--|--|------------------|------------------|
| 1B1 | Coal mining and handling biomass | CH ₄ | 0.10 | 0.10 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.08 | 0.08 | 0.0000 | 1.0000 |
| Total | | | 801,593.08 | 801,593.08 | 1.0000 | |

Table A 1.3.3 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF)

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|---------------------------------------|---|------------------|------------------|
| 1A3b | Road transportation: liquid fuels | CO ₂ | 113,364.64 | 113,364.64 | 0.2169 | 0.2169 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 73,421.79 | 73,421.79 | 0.1405 | 0.3574 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 61,476.35 | 61,476.35 | 0.1176 | 0.4750 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 23,723.43 | 23,723.43 | 0.0454 | 0.5204 |
| 1A1 | Energy industries: solid fuels | CO ₂ | 20,363.72 | 20,363.72 | 0.0390 | 0.5594 |
| 4A | Forest land | CO ₂ | -18,211.77 | 18,211.77 | 0.0348 | 0.5942 |
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 16,770.98 | 16,770.98 | 0.0321 | 0.6263 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 15,767.40 | 15,767.40 | 0.0302 | 0.6565 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 14,676.48 | 14,676.48 | 0.0281 | 0.6846 |
| 5A | Solid waste disposal | CH ₄ | 14,261.40 | 14,261.40 | 0.0273 | 0.7119 |
| 1A4 | Other sectors: liquid fuels | CO ₂ | 13,284.20 | 13,284.20 | 0.0254 | 0.7373 |
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 12,133.91 | 12,133.91 | 0.0232 | 0.7605 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 11,653.11 | 11,653.11 | 0.0223 | 0.7828 |
| 3D | Agricultural soils | N ₂ O | 11,466.79 | 11,466.79 | 0.0219 | 0.8047 |
| 4B | Cropland | CO ₂ | 10,971.23 | 10,971.23 | 0.0210 | 0.8257 |
| 4C | Grassland | CO ₂ | -8,861.80 | 8,861.80 | 0.0170 | 0.8427 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 4E | Settlements | CO ₂ | 6,451.34 | 6,451.34 | 0.0123 | 0.8550 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 5,443.73 | 5,443.73 | 0.0104 | 0.8654 |
| 1A1 | Energy industries: other fuels | CO ₂ | 5,021.63 | 5,021.63 | 0.0096 | 0.8750 |
| 1B2 | Oil and gas extraction | CH ₄ | 4,935.25 | 4,935.25 | 0.0094 | 0.8845 |
| 2A1 | Cement production | CO ₂ | 4,409.79 | 4,409.79 | 0.0084 | 0.8929 |
| 1B2 | Oil and gas extraction | CO ₂ | 4,230.64 | 4,230.64 | 0.0081 | 0.9010 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,226.81 | 4,226.81 | 0.0081 | 0.9091 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,043.20 | 4,043.20 | 0.0077 | 0.9168 |
| 5D | Wastewater treatment and discharge | CH ₄ | 3,445.75 | 3,445.75 | 0.0066 | 0.9234 |
| 2B8 | Petrochemical and carbon black production | CO ₂ | 2,846.51 | 2,846.51 | 0.0054 | 0.9289 |
| 3B2 | Manure management from Sheep | N ₂ O | 2,814.79 | 2,814.79 | 0.0054 | 0.9343 |
| 2C1 | Iron and steel production | CO ₂ | 2,501.24 | 2,501.24 | 0.0048 | 0.9391 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 2,276.82 | 2,276.82 | 0.0044 | 0.9434 |
| 4G | Harvested wood products | CO ₂ | -2,015.60 | 2,015.60 | 0.0039 | 0.9473 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,939.08 | 1,939.08 | 0.0037 | 0.9510 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,898.39 | 1,898.39 | 0.0036 | 0.9546 |
| 2B1 | Ammonia production | CO ₂ | 1,763.86 | 1,763.86 | 0.0034 | 0.9580 |
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 1,658.10 | 1,658.10 | 0.0032 | 0.9612 |
| 1A5 | Other: liquid fuels | CO ₂ | 1,558.20 | 1,558.20 | 0.0030 | 0.9641 |
| 5B | Biological treatment of solid waste | CH ₄ | 1,193.62 | 1,193.62 | 0.0023 | 0.9664 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,095.65 | 1,095.65 | 0.0021 | 0.9685 |
| 2A2 | Lime production | CO ₂ | 1,052.06 | 1,052.06 | 0.0020 | 0.9705 |
| 3G | Liming | CO ₂ | 939.13 | 939.13 | 0.0018 | 0.9723 |
| 5B | Biological treatment of solid waste | N ₂ O | 728.03 | 728.03 | 0.0014 | 0.9737 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|---------------------------------------|---|------------------|------------------|
| 5D | Wastewater treatment and discharge | N ₂ O | 725.93 | 725.93 | 0.0014 | 0.9751 |
| 2G3 | N ₂ O from product uses | N ₂ O | 645.41 | 645.41 | 0.0012 | 0.9763 |
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 641.74 | 641.74 | 0.0012 | 0.9776 |
| 1A4 | Other sectors: biomass | CH ₄ | 601.47 | 601.47 | 0.0012 | 0.9787 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 565.53 | 565.53 | 0.0011 | 0.9798 |
| 4E | Settlements | N ₂ O | 524.99 | 524.99 | 0.0010 | 0.9808 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 519.22 | 519.22 | 0.0010 | 0.9818 |
| 1B1 | Coal mining and handling | CH ₄ | 483.98 | 483.98 | 0.0009 | 0.9827 |
| 4B | Cropland | N ₂ O | 461.97 | 461.97 | 0.0009 | 0.9836 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 457.93 | 457.93 | 0.0009 | 0.9845 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 450.52 | 450.52 | 0.0009 | 0.9854 |
| 2A4 | Other process uses of carbonates | CO ₂ | 419.65 | 419.65 | 0.0008 | 0.9862 |
| 2A3 | Glass production | CO ₂ | 367.98 | 367.98 | 0.0007 | 0.9869 |
| 3H | Urea application to land | CO ₂ | 343.95 | 343.95 | 0.0007 | 0.9875 |
| 4D | Wetlands | CO ₂ | 336.78 | 336.78 | 0.0006 | 0.9882 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 324.69 | 324.69 | 0.0006 | 0.9888 |
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 318.00 | 318.00 | 0.0006 | 0.9894 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 289.24 | 289.24 | 0.0006 | 0.9899 |
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 281.17 | 281.17 | 0.0005 | 0.9905 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 268.38 | 268.38 | 0.0005 | 0.9910 |
| 5C | Incineration and open burning of waste | CO ₂ | 258.91 | 258.91 | 0.0005 | 0.9915 |
| 4 | Indirect N ₂ O emissions from LULUCF | N ₂ O | 243.62 | 243.62 | 0.0005 | 0.9920 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 206.19 | 206.19 | 0.0004 | 0.9924 |
| 2G4 | Other product manufacture and use | N ₂ O | 198.47 | 198.47 | 0.0004 | 0.9927 |
| 3A3 | Enteric fermentation from Swine | CH ₄ | 186.33 | 186.33 | 0.0004 | 0.9931 |
| 2B6 | Titanium dioxide production | CO ₂ | 181.02 | 181.02 | 0.0003 | 0.9934 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 175.44 | 175.44 | 0.0003 | 0.9938 |
| 1A4 | Other sectors: solid fuels | CH ₄ | 173.62 | 173.62 | 0.0003 | 0.9941 |
| 1A1 | Energy industries: biomass | N ₂ O | 171.90 | 171.90 | 0.0003 | 0.9944 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 161.99 | 161.99 | 0.0003 | 0.9947 |
| 4A | Forest land | N ₂ O | 144.65 | 144.65 | 0.0003 | 0.9950 |
| 2B7 | Soda ash production | CO ₂ | 140.18 | 140.18 | 0.0003 | 0.9953 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 137.70 | 137.70 | 0.0003 | 0.9956 |
| 1A1 | Energy industries: solid fuels | N ₂ O | 115.89 | 115.89 | 0.0002 | 0.9958 |
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 112.04 | 112.04 | 0.0002 | 0.9960 |
| 1A1 | Energy industries: other fuels | CH ₄ | 110.63 | 110.63 | 0.0002 | 0.9962 |
| 1A1 | Energy industries: biomass | CH ₄ | 109.01 | 109.01 | 0.0002 | 0.9964 |
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 102.56 | 102.56 | 0.0002 | 0.9966 |
| 1A4 | Other sectors: biomass | N ₂ O | 94.30 | 94.30 | 0.0002 | 0.9968 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 93.64 | 93.64 | 0.0002 | 0.9970 |
| 1A1 | Energy industries: other fuels | N ₂ O | 93.36 | 93.36 | 0.0002 | 0.9971 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 1A3b | Road transportation: liquid fuels | CH ₄ | 93.16 | 93.16 | 0.0002 | 0.9973 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 93.01 | 93.01 | 0.0002 | 0.9975 |
| 1A3b | Road transportation: biomass | CO ₂ | 92.80 | 92.80 | 0.0002 | 0.9977 |
| 1B1 | Coal mining and handling liquid fuels | CO ₂ | 91.16 | 91.16 | 0.0002 | 0.9979 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 82.03 | 82.03 | 0.0002 | 0.9980 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 80.97 | 80.97 | 0.0002 | 0.9982 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 72.08 | 72.08 | 0.0001 | 0.9983 |
| 2C3 | Aluminium production | CO ₂ | 71.62 | 71.62 | 0.0001 | 0.9984 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 64.53 | 64.53 | 0.0001 | 0.9986 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 61.71 | 61.71 | 0.0001 | 0.9987 |
| 2B10 | Other Chemical Industry | CH ₄ | 48.97 | 48.97 | 0.0001 | 0.9988 |
| 1A3c | Railways: solid fuels | CO ₂ | 41.15 | 41.15 | 0.0001 | 0.9989 |
| 5C | Incineration and open burning of waste | N ₂ O | 39.63 | 39.63 | 0.0001 | 0.9989 |
| 1B2 | Oil and gas extraction | N ₂ O | 39.36 | 39.36 | 0.0001 | 0.9990 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 38.62 | 38.62 | 0.0001 | 0.9991 |
| 2B2 | Nitric acid production | N ₂ O | 37.25 | 37.25 | 0.0001 | 0.9991 |
| 4C | Grassland | N ₂ O | 33.24 | 33.24 | 0.0001 | 0.9992 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 32.75 | 32.75 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 32.04 | 32.04 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 30.91 | 30.91 | 0.0001 | 0.9994 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 30.40 | 30.40 | 0.0001 | 0.9995 |
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 21.71 | 21.71 | 0.0000 | 0.9995 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|---------------------------------------|---|------------------|------------------|
| 2F5 | Solvents | HFCs, PFCs, SF ₆ and NF ₃ | 18.67 | 18.67 | 0.0000 | 0.9995 |
| 1A3c | Railways: liquid fuels | N ₂ O | 18.30 | 18.30 | 0.0000 | 0.9996 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 18.08 | 18.08 | 0.0000 | 0.9996 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.96 | 17.96 | 0.0000 | 0.9996 |
| 4C | Grassland | CH ₄ | 17.78 | 17.78 | 0.0000 | 0.9997 |
| 1A5 | Other: liquid fuels | N ₂ O | 16.48 | 16.48 | 0.0000 | 0.9997 |
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 15.07 | 15.07 | 0.0000 | 0.9997 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 12.48 | 12.48 | 0.0000 | 0.9998 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 12.16 | 12.16 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 10.86 | 10.86 | 0.0000 | 0.9998 |
| 2C1 | Iron and steel production | CH ₄ | 10.68 | 10.68 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 10.47 | 10.47 | 0.0000 | 0.9998 |
| 5C | Incineration and open burning of waste | CH ₄ | 10.45 | 10.45 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 10.05 | 10.05 | 0.0000 | 0.9999 |
| 4E | Settlements | CH ₄ | 8.25 | 8.25 | 0.0000 | 0.9999 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 7.00 | 7.00 | 0.0000 | 0.9999 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 6.28 | 6.28 | 0.0000 | 0.9999 |
| 2C1 | Iron and steel production | N ₂ O | 6.19 | 6.19 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 6.12 | 6.12 | 0.0000 | 0.9999 |
| 2A4 | Other process uses of carbonates | CH ₄ | 5.89 | 5.89 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: solid fuels | CH ₄ | 5.69 | 5.69 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | CO ₂ | 4.83 | 4.83 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|--------------|---|------------------|---------------------------------------|---|------------------|------------------|
| 1B1 | Coal mining and handling biomass | CH ₄ | 3.76 | 3.76 | 0.0000 | 1.0000 |
| 4A | Forest land | CH ₄ | 2.31 | 2.31 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 1.92 | 1.92 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 1.55 | 1.55 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 1.51 | 1.51 | 0.0000 | 1.0000 |
| 1A5 | Other: liquid fuels | CH ₄ | 1.03 | 1.03 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | CH ₄ | 0.94 | 0.94 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.35 | 0.35 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | CH ₄ | 0.34 | 0.34 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.33 | 0.33 | 0.0000 | 1.0000 |
| 4D | Wetlands | N ₂ O | 0.30 | 0.30 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.28 | 0.28 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 4B | Cropland | CH ₄ | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.01 | 0.01 | 0.0000 | 1.0000 |
| Total | | | 464,453.44 | 522,631.79 | 1.0000 | |

Table A 1.3.4 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF)

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|-----------------|---------------------------------------|---|------------------|------------------|
| 1A3b | Road transportation: liquid fuels | CO ₂ | 113,364.64 | 113,364.64 | 0.2390 | 0.2390 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 73,421.79 | 73,421.79 | 0.1548 | 0.3938 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 61,476.35 | 61,476.35 | 0.1296 | 0.5234 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 23,723.43 | 23,723.43 | 0.0500 | 0.5734 |
| 1A1 | Energy industries: solid fuels | CO ₂ | 20,363.72 | 20,363.72 | 0.0429 | 0.6163 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 16,770.98 | 16,770.98 | 0.0354 | 0.6517 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 15,767.40 | 15,767.40 | 0.0332 | 0.6849 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 14,676.48 | 14,676.48 | 0.0309 | 0.7159 |
| 5A | Solid waste disposal | CH ₄ | 14,261.40 | 14,261.40 | 0.0301 | 0.7459 |
| 1A4 | Other sectors: liquid fuels | CO ₂ | 13,284.20 | 13,284.20 | 0.0280 | 0.7739 |
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 12,133.91 | 12,133.91 | 0.0256 | 0.7995 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 11,653.11 | 11,653.11 | 0.0246 | 0.8241 |
| 3D | Agricultural soils | N ₂ O | 11,466.79 | 11,466.79 | 0.0242 | 0.8483 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 5,443.73 | 5,443.73 | 0.0115 | 0.8597 |
| 1A1 | Energy industries: other fuels | CO ₂ | 5,021.63 | 5,021.63 | 0.0106 | 0.8703 |
| 1B2 | Oil and gas extraction | CH ₄ | 4,935.25 | 4,935.25 | 0.0104 | 0.8807 |
| 2A1 | Cement production | CO ₂ | 4,409.79 | 4,409.79 | 0.0093 | 0.8900 |
| 1B2 | Oil and gas extraction | CO ₂ | 4,230.64 | 4,230.64 | 0.0089 | 0.8989 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,226.81 | 4,226.81 | 0.0089 | 0.9078 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,043.20 | 4,043.20 | 0.0085 | 0.9164 |
| 5D | Wastewater treatment and discharge | CH ₄ | 3,445.75 | 3,445.75 | 0.0073 | 0.9236 |
| 2B8 | Petrochemical and carbon black production | CO ₂ | 2,846.51 | 2,846.51 | 0.0060 | 0.9296 |
| 3B2 | Manure management from Sheep | N ₂ O | 2,814.79 | 2,814.79 | 0.0059 | 0.9356 |
| 2C1 | Iron and steel production | CO ₂ | 2,501.24 | 2,501.24 | 0.0053 | 0.9408 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 2,276.82 | 2,276.82 | 0.0048 | 0.9456 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,939.08 | 1,939.08 | 0.0041 | 0.9497 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,898.39 | 1,898.39 | 0.0040 | 0.9537 |
| 2B1 | Ammonia production | CO ₂ | 1,763.86 | 1,763.86 | 0.0037 | 0.9574 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|---------------------------------------|---|------------------|------------------|
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 1,658.10 | 1,658.10 | 0.0035 | 0.9609 |
| 1A5 | Other: liquid fuels | CO ₂ | 1,558.20 | 1,558.20 | 0.0033 | 0.9642 |
| 5B | Biological treatment of solid waste | CH ₄ | 1,193.62 | 1,193.62 | 0.0025 | 0.9667 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,095.65 | 1,095.65 | 0.0023 | 0.9691 |
| 2A2 | Lime production | CO ₂ | 1,052.06 | 1,052.06 | 0.0022 | 0.9713 |
| 3G | Liming | CO ₂ | 939.13 | 939.13 | 0.0020 | 0.9733 |
| 5B | Biological treatment of solid waste | N ₂ O | 728.03 | 728.03 | 0.0015 | 0.9748 |
| 5D | Wastewater treatment and discharge | N ₂ O | 725.93 | 725.93 | 0.0015 | 0.9763 |
| 2G3 | N ₂ O from product uses | N ₂ O | 645.41 | 645.41 | 0.0014 | 0.9777 |
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 641.74 | 641.74 | 0.0014 | 0.9790 |
| 1A4 | Other sectors: biomass | CH ₄ | 601.47 | 601.47 | 0.0013 | 0.9803 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 565.53 | 565.53 | 0.0012 | 0.9815 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 519.22 | 519.22 | 0.0011 | 0.9826 |
| 1B1 | Coal mining and handling | CH ₄ | 483.98 | 483.98 | 0.0010 | 0.9836 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 457.93 | 457.93 | 0.0010 | 0.9846 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 450.52 | 450.52 | 0.0009 | 0.9855 |
| 2A4 | Other process uses of carbonates | CO ₂ | 419.65 | 419.65 | 0.0009 | 0.9864 |
| 2A3 | Glass production | CO ₂ | 367.98 | 367.98 | 0.0008 | 0.9872 |
| 3H | Urea application to land | CO ₂ | 343.95 | 343.95 | 0.0007 | 0.9879 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 324.69 | 324.69 | 0.0007 | 0.9886 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 318.00 | 318.00 | 0.0007 | 0.9893 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 289.24 | 289.24 | 0.0006 | 0.9899 |
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 281.17 | 281.17 | 0.0006 | 0.9905 |
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 268.38 | 268.38 | 0.0006 | 0.9910 |
| 5C | Incineration and open burning of waste | CO ₂ | 258.91 | 258.91 | 0.0005 | 0.9916 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 206.19 | 206.19 | 0.0004 | 0.9920 |
| 2G4 | Other product manufacture and use | N ₂ O | 198.47 | 198.47 | 0.0004 | 0.9924 |
| 3A3 | Enteric fermentation from Swine | CH ₄ | 186.33 | 186.33 | 0.0004 | 0.9928 |
| 2B6 | Titanium dioxide production | CO ₂ | 181.02 | 181.02 | 0.0004 | 0.9932 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 175.44 | 175.44 | 0.0004 | 0.9936 |
| 1A4 | Other sectors: solid fuels | CH ₄ | 173.62 | 173.62 | 0.0004 | 0.9939 |
| 1A1 | Energy industries: biomass | N ₂ O | 171.90 | 171.90 | 0.0004 | 0.9943 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 161.99 | 161.99 | 0.0003 | 0.9946 |
| 2B7 | Soda ash production | CO ₂ | 140.18 | 140.18 | 0.0003 | 0.9949 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 137.70 | 137.70 | 0.0003 | 0.9952 |
| 1A1 | Energy industries: solid fuels | N ₂ O | 115.89 | 115.89 | 0.0002 | 0.9955 |
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 112.04 | 112.04 | 0.0002 | 0.9957 |
| 1A1 | Energy industries: other fuels | CH ₄ | 110.63 | 110.63 | 0.0002 | 0.9959 |
| 1A1 | Energy industries: biomass | CH ₄ | 109.01 | 109.01 | 0.0002 | 0.9962 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|---|---------------------------------------|---|------------------|------------------|
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 102.56 | 102.56 | 0.0002 | 0.9964 |
| 1A4 | Other sectors: biomass | N ₂ O | 94.30 | 94.30 | 0.0002 | 0.9966 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 93.64 | 93.64 | 0.0002 | 0.9968 |
| 1A1 | Energy industries: other fuels | N ₂ O | 93.36 | 93.36 | 0.0002 | 0.9970 |
| 1A3b | Road transportation: liquid fuels | CH ₄ | 93.16 | 93.16 | 0.0002 | 0.9972 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 93.01 | 93.01 | 0.0002 | 0.9974 |
| 1A3b | Road transportation: biomass | CO ₂ | 92.80 | 92.80 | 0.0002 | 0.9976 |
| 1B1 | Coal mining and handling liquid fuels | CO ₂ | 91.16 | 91.16 | 0.0002 | 0.9978 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 82.03 | 82.03 | 0.0002 | 0.9979 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 80.97 | 80.97 | 0.0002 | 0.9981 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 72.08 | 72.08 | 0.0002 | 0.9983 |
| 2C3 | Aluminium production | CO ₂ | 71.62 | 71.62 | 0.0002 | 0.9984 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 64.53 | 64.53 | 0.0001 | 0.9985 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 61.71 | 61.71 | 0.0001 | 0.9987 |
| 2B10 | Other Chemical Industry | CH ₄ | 48.97 | 48.97 | 0.0001 | 0.9988 |
| 1A3c | Railways: solid fuels | CO ₂ | 41.15 | 41.15 | 0.0001 | 0.9989 |
| 5C | Incineration and open burning of waste | N ₂ O | 39.63 | 39.63 | 0.0001 | 0.9989 |
| 1B2 | Oil and gas extraction | N ₂ O | 39.36 | 39.36 | 0.0001 | 0.9990 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 38.62 | 38.62 | 0.0001 | 0.9991 |
| 2B2 | Nitric acid production | N ₂ O | 37.25 | 37.25 | 0.0001 | 0.9992 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 32.75 | 32.75 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 32.04 | 32.04 | 0.0001 | 0.9993 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 30.91 | 30.91 | 0.0001 | 0.9994 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 30.40 | 30.40 | 0.0001 | 0.9995 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|--|---|---------------------------------------|---|------------------|------------------|
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 21.71 | 21.71 | 0.0000 | 0.9995 |
| 2F5 | Solvents | HFCs, PFCs, SF ₆ and NF ₃ | 18.67 | 18.67 | 0.0000 | 0.9995 |
| 1A3c | Railways: liquid fuels | N ₂ O | 18.30 | 18.30 | 0.0000 | 0.9996 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 18.08 | 18.08 | 0.0000 | 0.9996 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.96 | 17.96 | 0.0000 | 0.9997 |
| 1A5 | Other: liquid fuels | N ₂ O | 16.48 | 16.48 | 0.0000 | 0.9997 |
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 15.07 | 15.07 | 0.0000 | 0.9997 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 12.48 | 12.48 | 0.0000 | 0.9997 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 12.16 | 12.16 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 10.86 | 10.86 | 0.0000 | 0.9998 |
| 2C1 | Iron and steel production | CH ₄ | 10.68 | 10.68 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 10.47 | 10.47 | 0.0000 | 0.9998 |
| 5C | Incineration and open burning of waste | CH ₄ | 10.45 | 10.45 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 10.05 | 10.05 | 0.0000 | 0.9999 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 7.00 | 7.00 | 0.0000 | 0.9999 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 6.28 | 6.28 | 0.0000 | 0.9999 |
| 2C1 | Iron and steel production | N ₂ O | 6.19 | 6.19 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 6.12 | 6.12 | 0.0000 | 0.9999 |
| 2A4 | Other process uses of carbonates | CH ₄ | 5.89 | 5.89 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: solid fuels | CH ₄ | 5.69 | 5.69 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|--------------|---|------------------|---------------------------------------|---|------------------|------------------|
| 1A4 | Other sectors: peat | CO ₂ | 4.83 | 4.83 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling biomass | CH ₄ | 3.76 | 3.76 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 1.92 | 1.92 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 1.55 | 1.55 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 1.51 | 1.51 | 0.0000 | 1.0000 |
| 1A5 | Other: liquid fuels | CH ₄ | 1.03 | 1.03 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | CH ₄ | 0.94 | 0.94 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.35 | 0.35 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | CH ₄ | 0.34 | 0.34 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.33 | 0.33 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.28 | 0.28 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.01 | 0.01 | 0.0000 | 1.0000 |
| Total | | | 474,346.12 | 474,346.12 | 1.0000 | |

Table A 1.3.5 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF)

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|-----------------------------------|-----------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A1 | Energy industries: solid fuels | CO ₂ | 185,492.23 | 20,363.72 | 0.1025 | 0.2251 | 0.2251 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 9,229.34 | 61,476.35 | 0.0661 | 0.1451 | 0.3702 |
| 1A3b | Road transportation: liquid fuels | CO ₂ | 108,570.16 | 113,364.64 | 0.0594 | 0.1305 | 0.5007 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 70,373.19 | 73,421.79 | 0.0385 | 0.0844 | 0.5851 |
| 5A | Solid waste disposal | CH ₄ | 60,432.98 | 14,261.40 | 0.0244 | 0.0536 | 0.6387 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1B1 | Coal mining and handling | CH ₄ | 21,826.68 | 483.98 | 0.0143 | 0.0314 | 0.6702 |
| 2B3 | Adipic acid production | N ₂ O | 19,934.61 | - | 0.0136 | 0.0298 | 0.7000 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 531.30 | 11,653.11 | 0.0134 | 0.0293 | 0.7293 |
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 38,484.20 | 12,133.91 | 0.0120 | 0.0263 | 0.7556 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 17,784.67 | 175.44 | 0.0119 | 0.0262 | 0.7818 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 19,868.96 | 2,276.82 | 0.0109 | 0.0239 | 0.8056 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 40,883.76 | 15,767.40 | 0.0093 | 0.0205 | 0.8261 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 27,330.36 | 23,723.43 | 0.0093 | 0.0204 | 0.8465 |
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 19,881.28 | 16,770.98 | 0.0062 | 0.0136 | 0.8601 |
| 1A1 | Energy industries: other fuels | CO ₂ | 246.13 | 5,021.63 | 0.0057 | 0.0126 | 0.8727 |
| 3D | Agricultural soils | N ₂ O | 13,610.39 | 11,466.79 | 0.0042 | 0.0093 | 0.8820 |
| 4A | Forest land | CO ₂ | -15,026.26 | -18,211.77 | 0.0037 | 0.0081 | 0.8901 |
| 4B | Cropland | CO ₂ | 14,265.93 | 10,971.23 | 0.0032 | 0.0070 | 0.8971 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 29,832.00 | 14,676.48 | 0.0031 | 0.0067 | 0.9038 |
| 4E | Settlements | CO ₂ | 7,011.30 | 6,451.34 | 0.0028 | 0.0062 | 0.9100 |
| 1B2 | Oil and gas extraction | CH ₄ | 12,345.07 | 4,935.25 | 0.0026 | 0.0057 | 0.9157 |
| 2B2 | Nitric acid production | N ₂ O | 3,860.26 | 37.25 | 0.0026 | 0.0057 | 0.9214 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--------------------------------------|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A4 | Other sectors: liquid fuels | CO ₂ | 19,607.73 | 13,284.20 | 0.0023 | 0.0050 | 0.9264 |
| 1A5 | Other: liquid fuels | CO ₂ | 5,293.44 | 1,558.20 | 0.0018 | 0.0039 | 0.9303 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,732.53 | 4,226.81 | 0.0017 | 0.0038 | 0.9341 |
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 663.00 | 1,658.10 | 0.0015 | 0.0033 | 0.9374 |
| 4C | Grassland | CO ₂ | -7,111.03 | -8,861.80 | 0.0015 | 0.0032 | 0.9406 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,935.90 | 4,043.20 | 0.0014 | 0.0031 | 0.9437 |
| 5B | Biological treatment of solid waste | CH ₄ | 18.13 | 1,193.62 | 0.0014 | 0.0031 | 0.9467 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,455.18 | 1,939.08 | 0.0013 | 0.0028 | 0.9496 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 7,611.13 | 5,443.73 | 0.0012 | 0.0027 | 0.9522 |
| 5D | Wastewater treatment and discharge | CH ₄ | 4,218.90 | 3,445.75 | 0.0012 | 0.0026 | 0.9548 |
| 1B2 | Oil and gas extraction | CO ₂ | 5,777.92 | 4,230.64 | 0.0010 | 0.0023 | 0.9571 |
| 3B2 | Manure management from Sheep | N ₂ O | 3,442.70 | 2,814.79 | 0.0010 | 0.0021 | 0.9592 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,881.31 | 1,898.39 | 0.0010 | 0.0021 | 0.9613 |
| 2C6 | Zinc production | CO ₂ | 1,358.83 | - | 0.0009 | 0.0020 | 0.9634 |
| 2C1 | Iron and steel production | CO ₂ | 5,591.54 | 2,501.24 | 0.0009 | 0.0019 | 0.9653 |
| 5B | Biological treatment of solid waste | N ₂ O | 12.97 | 728.03 | 0.0008 | 0.0019 | 0.9671 |
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 1,698.56 | 268.38 | 0.0008 | 0.0018 | 0.9690 |
| 2B1 | Ammonia production | CO ₂ | 1,895.00 | 1,763.86 | 0.0008 | 0.0017 | 0.9707 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A3b | Road transportation: liquid fuels | CH ₄ | 1,246.70 | 93.16 | 0.0007 | 0.0016 | 0.9723 |
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 76.86 | 641.74 | 0.0007 | 0.0015 | 0.9739 |
| 1A4 | Other sectors: biomass | CH ₄ | 62.37 | 601.47 | 0.0007 | 0.0015 | 0.9753 |
| 1A4 | Other sectors: solid fuels | CH ₄ | 1,265.22 | 173.62 | 0.0007 | 0.0014 | 0.9768 |
| 5C | Incineration and open burning of waste | CO ₂ | 1,300.71 | 258.91 | 0.0006 | 0.0013 | 0.9781 |
| 1A1 | Energy industries: solid fuels | N ₂ O | 1,023.72 | 115.89 | 0.0006 | 0.0012 | 0.9793 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 224.74 | 565.53 | 0.0005 | 0.0011 | 0.9804 |
| 3G | Liming | CO ₂ | 1,015.18 | 939.13 | 0.0004 | 0.0009 | 0.9813 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 184.49 | 450.52 | 0.0004 | 0.0009 | 0.9822 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,312.28 | 1,095.65 | 0.0004 | 0.0009 | 0.9831 |
| 2G3 | N ₂ O from product uses | N ₂ O | 552.57 | 645.41 | 0.0004 | 0.0008 | 0.9839 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 1.41 | 324.69 | 0.0004 | 0.0008 | 0.9848 |
| 4G | Harvested wood products | CO ₂ | -1,639.08 | -2,015.60 | 0.0004 | 0.0008 | 0.9856 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 292.27 | 457.93 | 0.0003 | 0.0007 | 0.9863 |
| 5D | Wastewater treatment and discharge | N ₂ O | 783.59 | 725.93 | 0.0003 | 0.0007 | 0.9870 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|---|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 2B8 | Petrochemical and carbon black production | CO ₂ | 4,538.91 | 2,846.51 | 0.0003 | 0.0006 | 0.9876 |
| 1A4 | Other sectors: peat | CO ₂ | 372.48 | 4.83 | 0.0002 | 0.0005 | 0.9881 |
| 2A2 | Lime production | CO ₂ | 1,462.05 | 1,052.06 | 0.0002 | 0.0005 | 0.9887 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 552.81 | 519.22 | 0.0002 | 0.0005 | 0.9892 |
| 2C3 | Aluminium production | CO ₂ | 450.32 | 71.62 | 0.0002 | 0.0005 | 0.9897 |
| 4E | Settlements | N ₂ O | 584.68 | 524.99 | 0.0002 | 0.0005 | 0.9901 |
| 2A1 | Cement production | CO ₂ | 7,295.26 | 4,409.79 | 0.0002 | 0.0005 | 0.9906 |
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 333.43 | 15.07 | 0.0002 | 0.0005 | 0.9911 |
| 2G4 | Other product manufacture and use | N ₂ O | 41.00 | 198.47 | 0.0002 | 0.0005 | 0.9915 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 198.91 | 289.24 | 0.0002 | 0.0004 | 0.9920 |
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 783.25 | 281.17 | 0.0002 | 0.0004 | 0.9924 |
| 1A1 | Energy industries: biomass | N ₂ O | 0.25 | 171.90 | 0.0002 | 0.0004 | 0.9929 |
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 278.70 | 318.00 | 0.0002 | 0.0004 | 0.9933 |
| 3H | Urea application to land | CO ₂ | 327.60 | 343.95 | 0.0002 | 0.0004 | 0.9937 |
| 2A3 | Glass production | CO ₂ | 405.54 | 367.98 | 0.0002 | 0.0003 | 0.9940 |
| 4B | Cropland | N ₂ O | 1,019.85 | 461.97 | 0.0002 | 0.0003 | 0.9944 |
| 2B6 | Titanium dioxide production | CO ₂ | 104.63 | 181.02 | 0.0001 | 0.0003 | 0.9947 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 387.17 | 112.04 | 0.0001 | 0.0003 | 0.9950 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 241.68 | 30.40 | 0.0001 | 0.0003 | 0.9952 |
| 1A1 | Energy industries: biomass | CH ₄ | 0.47 | 109.01 | 0.0001 | 0.0003 | 0.9955 |
| 3F | Field burning of agricultural residues | CH ₄ | 186.57 | - | 0.0001 | 0.0003 | 0.9958 |
| 1A1 | Energy industries: other fuels | CH ₄ | 18.55 | 110.63 | 0.0001 | 0.0003 | 0.9961 |
| 1A3b | Road transportation: biomass | CO ₂ | - | 92.80 | 0.0001 | 0.0002 | 0.9963 |
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 19.54 | 102.56 | 0.0001 | 0.0002 | 0.9965 |
| 1B1 | Coal mining and handling liquid fuels | CO ₂ | - | 91.16 | 0.0001 | 0.0002 | 0.9968 |
| 1A1 | Energy industries: other fuels | N ₂ O | 6.72 | 93.36 | 0.0001 | 0.0002 | 0.9970 |
| 1A4 | Other sectors: biomass | N ₂ O | 9.14 | 94.30 | 0.0001 | 0.0002 | 0.9972 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 92.30 | 137.70 | 0.0001 | 0.0002 | 0.9974 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 157.18 | 161.99 | 0.0001 | 0.0002 | 0.9976 |
| 5C | Incineration and open burning of waste | CH ₄ | 134.83 | 10.45 | 0.0001 | 0.0002 | 0.9978 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 12.29 | 64.53 | 0.0001 | 0.0001 | 0.9980 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 148.72 | 30.91 | 0.0001 | 0.0001 | 0.9981 |
| 4D | Wetlands | CO ₂ | 486.95 | 336.78 | 0.0001 | 0.0001 | 0.9982 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 2B10 | Other Chemical Industry | CH ₄ | 174.65 | 48.97 | 0.0001 | 0.0001 | 0.9984 |
| 2A4 | Other process uses of carbonates | CO ₂ | 640.93 | 419.65 | 0.0001 | 0.0001 | 0.9985 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 276.73 | 206.19 | 0.0001 | 0.0001 | 0.9986 |
| 1A3c | Railways: solid fuels | CO ₂ | - | 41.15 | 0.0000 | 0.0001 | 0.9987 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 36.88 | 61.71 | 0.0000 | 0.0001 | 0.9988 |
| 3F | Field burning of agricultural residues | N ₂ O | 57.66 | - | 0.0000 | 0.0001 | 0.9989 |
| 4C | Grassland | N ₂ O | 10.33 | 33.24 | 0.0000 | 0.0001 | 0.9990 |
| 1A1 | Energy industries: solid fuels | CH ₄ | 49.32 | 5.69 | 0.0000 | 0.0001 | 0.9990 |
| 3A3 | Enteric fermentation from Swine | CH ₄ | 283.06 | 186.33 | 0.0000 | 0.0001 | 0.9991 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 103.32 | 82.03 | 0.0000 | 0.0001 | 0.9992 |
| 2F5 | Solvents | HFCs, PFCs, SF ₆ and NF ₃ | - | 18.67 | 0.0000 | 0.0000 | 0.9992 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 37.47 | 38.62 | 0.0000 | 0.0000 | 0.9993 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 47.59 | 10.86 | 0.0000 | 0.0000 | 0.9993 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 132.00 | 93.01 | 0.0000 | 0.0000 | 0.9993 |
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 9.56 | 21.71 | 0.0000 | 0.0000 | 0.9994 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|---|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A5 | Other: liquid fuels | N ₂ O | 56.12 | 16.48 | 0.0000 | 0.0000 | 0.9994 |
| 1B2 | Oil and gas extraction | N ₂ O | 40.75 | 39.36 | 0.0000 | 0.0000 | 0.9995 |
| 1A4 | Other sectors: peat | CH ₄ | 26.35 | 0.34 | 0.0000 | 0.0000 | 0.9995 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 138.62 | 93.64 | 0.0000 | 0.0000 | 0.9995 |
| 5C | Incineration and open burning of waste | N ₂ O | 46.66 | 39.63 | 0.0000 | 0.0000 | 0.9996 |
| 2A4 | Other process uses of carbonates | CH ₄ | 31.10 | 5.89 | 0.0000 | 0.0000 | 0.9996 |
| 4C | Grassland | CH ₄ | 9.97 | 17.78 | 0.0000 | 0.0000 | 0.9996 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 105.00 | 72.08 | 0.0000 | 0.0000 | 0.9997 |
| 4 | Indirect N ₂ O emissions from LULUCF | N ₂ O | 401.49 | 243.62 | 0.0000 | 0.0000 | 0.9997 |
| 4A | Forest land | N ₂ O | 231.22 | 144.65 | 0.0000 | 0.0000 | 0.9997 |
| 2C1 | Iron and steel production | CH ₄ | 36.90 | 10.68 | 0.0000 | 0.0000 | 0.9997 |
| 1A3c | Railways: liquid fuels | N ₂ O | 13.75 | 18.30 | 0.0000 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 0.38 | 10.05 | 0.0000 | 0.0000 | 0.9998 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.80 | 17.96 | 0.0000 | 0.0000 | 0.9998 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 43.36 | 18.08 | 0.0000 | 0.0000 | 0.9998 |
| 4E | Settlements | CH ₄ | 3.00 | 8.25 | 0.0000 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 44.43 | 32.04 | 0.0000 | 0.0000 | 0.9999 |
| 2B7 | Soda ash production | CO ₂ | 231.55 | 140.18 | 0.0000 | 0.0000 | 0.9999 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 0.16 | 6.12 | 0.0000 | 0.0000 | 0.9999 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 2.79 | 7.00 | 0.0000 | 0.0000 | 0.9999 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 29.98 | 12.16 | 0.0000 | 0.0000 | 0.9999 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 3.66 | 6.28 | 0.0000 | 0.0000 | 0.9999 |
| 2C1 | Iron and steel production | N ₂ O | 17.74 | 6.19 | 0.0000 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 14.54 | 12.48 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling biomass | CH ₄ | 0.10 | 3.76 | 0.0000 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 12.20 | 10.47 | 0.0000 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 6.37 | 1.55 | 0.0000 | 0.0000 | 1.0000 |
| 4D | Wetlands | N ₂ O | 4.13 | 0.30 | 0.0000 | 0.0000 | 1.0000 |
| 1A5 | Other: liquid fuels | CH ₄ | 3.56 | 1.03 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | CH ₄ | - | 0.94 | 0.0000 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 1.47 | 0.02 | 0.0000 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 57.79 | 32.75 | 0.0000 | 0.0000 | 1.0000 |
| 4A | Forest land | CH ₄ | 2.89 | 2.31 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 2.46 | 1.92 | 0.0000 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 140.41 | 80.97 | 0.0000 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 2.10 | 1.51 | 0.0000 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|--------------|--------------------------------------|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.29 | 0.35 | 0.0000 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.31 | 0.33 | 0.0000 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.26 | 0.28 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | N ₂ O | - | 0.09 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.09 | 0.02 | 0.0000 | 0.0000 | 1.0000 |
| 4B | Cropland | CH ₄ | 0.09 | 0.02 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.08 | 0.01 | 0.0000 | 0.0000 | 1.0000 |
| Total | | | 801,848.54 | 464,453.44 | 0.4554 | 1.0000 | |

Table A 1.3.6 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF)

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|------------------------------------|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A1 | Energy industries: solid fuels | CO ₂ | 185,492.23 | 20,363.72 | 0.1025 | 0.2312 | 0.2312 |
| 1A1 | Energy industries: gaseous fuels | CO ₂ | 9,229.34 | 61,476.35 | 0.0661 | 0.1490 | 0.3803 |
| 1A3b | Road transportation: liquid fuels | CO ₂ | 108,570.16 | 113,364.64 | 0.0594 | 0.1340 | 0.5143 |
| 1A4 | Other sectors: gaseous fuels | CO ₂ | 70,373.19 | 73,421.79 | 0.0385 | 0.0867 | 0.6010 |
| 5A | Solid waste disposal | CH ₄ | 60,432.98 | 14,261.40 | 0.0244 | 0.0551 | 0.6561 |
| 1B1 | Coal mining and handling | CH ₄ | 21,826.68 | 483.98 | 0.0143 | 0.0323 | 0.6884 |
| 2B3 | Adipic acid production | N ₂ O | 19,934.61 | - | 0.0136 | 0.0307 | 0.7190 |
| 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ | 531.30 | 11,653.11 | 0.0134 | 0.0301 | 0.7492 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ | 38,484.20 | 12,133.91 | 0.0120 | 0.0270 | 0.7761 |
| 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ | 17,784.67 | 175.44 | 0.0119 | 0.0269 | 0.8030 |
| 1A4 | Other sectors: solid fuels | CO ₂ | 19,868.96 | 2,276.82 | 0.0109 | 0.0245 | 0.8275 |
| 1A1 | Energy industries: liquid fuels | CO ₂ | 40,883.76 | 15,767.40 | 0.0093 | 0.0210 | 0.8485 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ | 27,330.36 | 23,723.43 | 0.0093 | 0.0210 | 0.8695 |
| 3A1 | Enteric fermentation from Cattle | CH ₄ | 19,881.28 | 16,770.98 | 0.0062 | 0.0140 | 0.8835 |
| 1A1 | Energy industries: other fuels | CO ₂ | 246.13 | 5,021.63 | 0.0057 | 0.0130 | 0.8964 |
| 3D | Agricultural soils | N ₂ O | 13,610.39 | 11,466.79 | 0.0042 | 0.0095 | 0.9059 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ | 29,832.00 | 14,676.48 | 0.0031 | 0.0069 | 0.9128 |
| 1B2 | Oil and gas extraction | CH ₄ | 12,345.07 | 4,935.25 | 0.0026 | 0.0059 | 0.9187 |
| 2B2 | Nitric acid production | N ₂ O | 3,860.26 | 37.25 | 0.0026 | 0.0058 | 0.9246 |
| 1A4 | Other sectors: liquid fuels | CO ₂ | 19,607.73 | 13,284.20 | 0.0023 | 0.0051 | 0.9297 |
| 1A5 | Other: liquid fuels | CO ₂ | 5,293.44 | 1,558.20 | 0.0018 | 0.0040 | 0.9337 |
| 3B1 | Manure management from Cattle | CH ₄ | 4,732.53 | 4,226.81 | 0.0017 | 0.0039 | 0.9376 |
| 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ | 663.00 | 1,658.10 | 0.0015 | 0.0034 | 0.9410 |
| 3A2 | Enteric fermentation from Sheep | CH ₄ | 4,935.90 | 4,043.20 | 0.0014 | 0.0031 | 0.9441 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 5B | Biological treatment of solid waste | CH ₄ | 18.13 | 1,193.62 | 0.0014 | 0.0031 | 0.9473 |
| 1A3c | Railways: liquid fuels | CO ₂ | 1,455.18 | 1,939.08 | 0.0013 | 0.0029 | 0.9502 |
| 1A3d | Domestic Navigation: liquid fuels | CO ₂ | 7,611.13 | 5,443.73 | 0.0012 | 0.0027 | 0.9529 |
| 5D | Wastewater treatment and discharge | CH ₄ | 4,218.90 | 3,445.75 | 0.0012 | 0.0027 | 0.9556 |
| 1B2 | Oil and gas extraction | CO ₂ | 5,777.92 | 4,230.64 | 0.0010 | 0.0023 | 0.9580 |
| 3B2 | Manure management from Sheep | N ₂ O | 3,442.70 | 2,814.79 | 0.0010 | 0.0022 | 0.9601 |
| 1A3a | Domestic aviation: liquid fuels | CO ₂ | 1,881.31 | 1,898.39 | 0.0010 | 0.0021 | 0.9623 |
| 2C6 | Zinc production | CO ₂ | 1,358.83 | - | 0.0009 | 0.0021 | 0.9644 |
| 2C1 | Iron and steel production | CO ₂ | 5,591.54 | 2,501.24 | 0.0009 | 0.0020 | 0.9663 |
| 5B | Biological treatment of solid waste | N ₂ O | 12.97 | 728.03 | 0.0008 | 0.0019 | 0.9682 |
| 1B1 | Coal mining and handling solid fuels | CO ₂ | 1,698.56 | 268.38 | 0.0008 | 0.0019 | 0.9701 |
| 2B1 | Ammonia production | CO ₂ | 1,895.00 | 1,763.86 | 0.0008 | 0.0018 | 0.9719 |
| 1A3b | Road transportation: liquid fuels | CH ₄ | 1,246.70 | 93.16 | 0.0007 | 0.0017 | 0.9736 |
| 1A2 | Manufacturing industries and construction: other fuels | CO ₂ | 76.86 | 641.74 | 0.0007 | 0.0016 | 0.9752 |
| 1A4 | Other sectors: biomass | CH ₄ | 62.37 | 601.47 | 0.0007 | 0.0015 | 0.9767 |
| 1A4 | Other sectors: solid fuels | CH ₄ | 1,265.22 | 173.62 | 0.0007 | 0.0015 | 0.9782 |
| 5C | Incineration and open burning of waste | CO ₂ | 1,300.71 | 258.91 | 0.0006 | 0.0013 | 0.9795 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A1 | Energy industries: solid fuels | N ₂ O | 1,023.72 | 115.89 | 0.0006 | 0.0013 | 0.9807 |
| 1A3e | Other transportation: liquid fuels | CO ₂ | 224.74 | 565.53 | 0.0005 | 0.0012 | 0.9819 |
| 3G | Liming | CO ₂ | 1,015.18 | 939.13 | 0.0004 | 0.0009 | 0.9828 |
| 2F2 | Foam blowing agents | HFCs, PFCs, SF ₆ and NF ₃ | 184.49 | 450.52 | 0.0004 | 0.0009 | 0.9837 |
| 1A3b | Road transportation: liquid fuels | N ₂ O | 1,312.28 | 1,095.65 | 0.0004 | 0.0009 | 0.9846 |
| 2G3 | N ₂ O from product uses | N ₂ O | 552.57 | 645.41 | 0.0004 | 0.0009 | 0.9855 |
| 2F3 | Fire protection | HFCs, PFCs, SF ₆ and NF ₃ | 1.41 | 324.69 | 0.0004 | 0.0009 | 0.9863 |
| 3A4 | Enteric fermentation from Other livestock | CH ₄ | 292.27 | 457.93 | 0.0003 | 0.0008 | 0.9871 |
| 5D | Wastewater treatment and discharge | N ₂ O | 783.59 | 725.93 | 0.0003 | 0.0007 | 0.9878 |
| 2B8 | Petrochemical and carbon black production | CO ₂ | 4,538.91 | 2,846.51 | 0.0003 | 0.0006 | 0.9884 |
| 1A4 | Other sectors: peat | CO ₂ | 372.48 | 4.83 | 0.0002 | 0.0006 | 0.9890 |
| 2A2 | Lime production | CO ₂ | 1,462.05 | 1,052.06 | 0.0002 | 0.0005 | 0.9895 |
| 2D | Non-energy products from fuels and solvent use | CO ₂ | 552.81 | 519.22 | 0.0002 | 0.0005 | 0.9900 |
| 2C3 | Aluminium production | CO ₂ | 450.32 | 71.62 | 0.0002 | 0.0005 | 0.9906 |
| 2A1 | Cement production | CO ₂ | 7,295.26 | 4,409.79 | 0.0002 | 0.0005 | 0.9910 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|---|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 2C3 | Aluminium production | HFCs, PFCs, SF ₆ and NF ₃ | 333.43 | 15.07 | 0.0002 | 0.0005 | 0.9915 |
| 2G4 | Other product manufacture and use | N ₂ O | 41.00 | 198.47 | 0.0002 | 0.0005 | 0.9920 |
| 1A1 | Energy industries: gaseous fuels | N ₂ O | 198.91 | 289.24 | 0.0002 | 0.0005 | 0.9924 |
| 2G1 | Electrical equipment | HFCs, PFCs, SF ₆ and NF ₃ | 783.25 | 281.17 | 0.0002 | 0.0005 | 0.9929 |
| 1A1 | Energy industries: biomass | N ₂ O | 0.25 | 171.90 | 0.0002 | 0.0005 | 0.9934 |
| 2G2 | SF ₆ and PFCs from other product use | HFCs, PFCs, SF ₆ and NF ₃ | 278.70 | 318.00 | 0.0002 | 0.0004 | 0.9938 |
| 3H | Urea application to land | CO ₂ | 327.60 | 343.95 | 0.0002 | 0.0004 | 0.9942 |
| 2A3 | Glass production | CO ₂ | 405.54 | 367.98 | 0.0002 | 0.0004 | 0.9945 |
| 2B6 | Titanium dioxide production | CO ₂ | 104.63 | 181.02 | 0.0001 | 0.0003 | 0.9949 |
| 2C4 | Magnesium production | HFCs, PFCs, SF ₆ and NF ₃ | 387.17 | 112.04 | 0.0001 | 0.0003 | 0.9951 |
| 1A4 | Other sectors: solid fuels | N ₂ O | 241.68 | 30.40 | 0.0001 | 0.0003 | 0.9954 |
| 1A1 | Energy industries: biomass | CH ₄ | 0.47 | 109.01 | 0.0001 | 0.0003 | 0.9957 |
| 3F | Field burning of agricultural residues | CH ₄ | 186.57 | - | 0.0001 | 0.0003 | 0.9960 |
| 1A1 | Energy industries: other fuels | CH ₄ | 18.55 | 110.63 | 0.0001 | 0.0003 | 0.9963 |
| 1A3b | Road transportation: biomass | CO ₂ | - | 92.80 | 0.0001 | 0.0002 | 0.9965 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A2 | Manufacturing industries and construction: biomass | N ₂ O | 19.54 | 102.56 | 0.0001 | 0.0002 | 0.9968 |
| 1B1 | Coal mining and handling liquid fuels | CO ₂ | - | 91.16 | 0.0001 | 0.0002 | 0.9970 |
| 1A1 | Energy industries: other fuels | N ₂ O | 6.72 | 93.36 | 0.0001 | 0.0002 | 0.9972 |
| 1A4 | Other sectors: biomass | N ₂ O | 9.14 | 94.30 | 0.0001 | 0.0002 | 0.9975 |
| 1A1 | Energy industries: gaseous fuels | CH ₄ | 92.30 | 137.70 | 0.0001 | 0.0002 | 0.9977 |
| 1A4 | Other sectors: gaseous fuels | CH ₄ | 157.18 | 161.99 | 0.0001 | 0.0002 | 0.9979 |
| 5C | Incineration and open burning of waste | CH ₄ | 134.83 | 10.45 | 0.0001 | 0.0002 | 0.9981 |
| 1A2 | Manufacturing industries and construction: biomass | CH ₄ | 12.29 | 64.53 | 0.0001 | 0.0002 | 0.9982 |
| 1A2 | Manufacturing industries and construction: solid fuels | N ₂ O | 148.72 | 30.91 | 0.0001 | 0.0001 | 0.9984 |
| 2B10 | Other Chemical Industry | CH ₄ | 174.65 | 48.97 | 0.0001 | 0.0001 | 0.9985 |
| 2A4 | Other process uses of carbonates | CO ₂ | 640.93 | 419.65 | 0.0001 | 0.0001 | 0.9986 |
| 3J | Agriculture activities in OTs and CDs | CH ₄ | 276.73 | 206.19 | 0.0001 | 0.0001 | 0.9988 |
| 1A3c | Railways: solid fuels | CO ₂ | - | 41.15 | 0.0000 | 0.0001 | 0.9989 |
| 2F6 | Other product uses as substitutes for ODS | HFCs, PFCs, SF ₆ and NF ₃ | 36.88 | 61.71 | 0.0000 | 0.0001 | 0.9990 |
| 3F | Field burning of agricultural residues | N ₂ O | 57.66 | - | 0.0000 | 0.0001 | 0.9991 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|---|---|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A1 | Energy industries: solid fuels | CH ₄ | 49.32 | 5.69 | 0.0000 | 0.0001 | 0.9991 |
| 3A3 | Enteric fermentation from Swine | CH ₄ | 283.06 | 186.33 | 0.0000 | 0.0001 | 0.9992 |
| 1A4 | Other sectors: liquid fuels | N ₂ O | 103.32 | 82.03 | 0.0000 | 0.0001 | 0.9993 |
| 2F5 | Solvents | HFCs, PFCs, SF ₆ and NF ₃ | - | 18.67 | 0.0000 | 0.0000 | 0.9993 |
| 1A4 | Other sectors: gaseous fuels | N ₂ O | 37.47 | 38.62 | 0.0000 | 0.0000 | 0.9993 |
| 1A2 | Manufacturing industries and construction: solid fuels | CH ₄ | 47.59 | 10.86 | 0.0000 | 0.0000 | 0.9994 |
| 3J | Agriculture activities in OTs and CDs | N ₂ O | 132.00 | 93.01 | 0.0000 | 0.0000 | 0.9994 |
| 2E1 | Integrated circuit or semiconductor | HFCs, PFCs, SF ₆ and NF ₃ | 9.56 | 21.71 | 0.0000 | 0.0000 | 0.9995 |
| 1A5 | Other: liquid fuels | N ₂ O | 56.12 | 16.48 | 0.0000 | 0.0000 | 0.9995 |
| 1B2 | Oil and gas extraction | N ₂ O | 40.75 | 39.36 | 0.0000 | 0.0000 | 0.9996 |
| 1A4 | Other sectors: peat | CH ₄ | 26.35 | 0.34 | 0.0000 | 0.0000 | 0.9996 |
| 1A2 | Manufacturing industries and construction: liquid fuels | N ₂ O | 138.62 | 93.64 | 0.0000 | 0.0000 | 0.9996 |
| 5C | Incineration and open burning of waste | N ₂ O | 46.66 | 39.63 | 0.0000 | 0.0000 | 0.9997 |
| 2A4 | Other process uses of carbonates | CH ₄ | 31.10 | 5.89 | 0.0000 | 0.0000 | 0.9997 |
| 1A3d | Domestic Navigation: liquid fuels | N ₂ O | 105.00 | 72.08 | 0.0000 | 0.0000 | 0.9997 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|-----------|--|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 2C1 | Iron and steel production | CH ₄ | 36.90 | 10.68 | 0.0000 | 0.0000 | 0.9998 |
| 1A3c | Railways: liquid fuels | N ₂ O | 13.75 | 18.30 | 0.0000 | 0.0000 | 0.9998 |
| 1A2 | Manufacturing industries and construction: other fuels | N ₂ O | 0.38 | 10.05 | 0.0000 | 0.0000 | 0.9998 |
| 1A3a | Domestic aviation: liquid fuels | N ₂ O | 17.80 | 17.96 | 0.0000 | 0.0000 | 0.9998 |
| 1A1 | Energy industries: liquid fuels | CH ₄ | 43.36 | 18.08 | 0.0000 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: liquid fuels | CH ₄ | 44.43 | 32.04 | 0.0000 | 0.0000 | 0.9999 |
| 2B7 | Soda ash production | CO ₂ | 231.55 | 140.18 | 0.0000 | 0.0000 | 0.9999 |
| 1A2 | Manufacturing industries and construction: other fuels | CH ₄ | 0.16 | 6.12 | 0.0000 | 0.0000 | 0.9999 |
| 1A3e | Other transportation: liquid fuels | N ₂ O | 2.79 | 7.00 | 0.0000 | 0.0000 | 0.9999 |
| 2B8 | Petrochemical and carbon black production | CH ₄ | 29.98 | 12.16 | 0.0000 | 0.0000 | 0.9999 |
| 1A3d | Domestic Navigation: liquid fuels | CH ₄ | 3.66 | 6.28 | 0.0000 | 0.0000 | 0.9999 |
| 2C1 | Iron and steel production | N ₂ O | 17.74 | 6.19 | 0.0000 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | N ₂ O | 14.54 | 12.48 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling biomass | CH ₄ | 0.10 | 3.76 | 0.0000 | 0.0000 | 1.0000 |
| 1A2 | Manufacturing industries and construction: gaseous fuels | CH ₄ | 12.20 | 10.47 | 0.0000 | 0.0000 | 1.0000 |
| 1A3a | Domestic aviation: liquid fuels | CH ₄ | 6.37 | 1.55 | 0.0000 | 0.0000 | 1.0000 |

| IPCC Code | IPCC Category | GHG | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment | Contribution to Trend | Cumulative Total |
|--------------|---|------------------|--|---------------------------------------|------------------|-----------------------|------------------|
| 1A5 | Other: liquid fuels | CH ₄ | 3.56 | 1.03 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | CH ₄ | - | 0.94 | 0.0000 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: peat | N ₂ O | 1.47 | 0.02 | 0.0000 | 0.0000 | 1.0000 |
| 1A4 | Other sectors: liquid fuels | CH ₄ | 57.79 | 32.75 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: liquid fuels | CH ₄ | 2.46 | 1.92 | 0.0000 | 0.0000 | 1.0000 |
| 1A1 | Energy industries: liquid fuels | N ₂ O | 140.41 | 80.97 | 0.0000 | 0.0000 | 1.0000 |
| 2B8 | Petrochemical and carbon black production | N ₂ O | 2.10 | 1.51 | 0.0000 | 0.0000 | 1.0000 |
| 1A3e | Other transportation: liquid fuels | CH ₄ | 0.29 | 0.35 | 0.0000 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | N ₂ O | 0.31 | 0.33 | 0.0000 | 0.0000 | 1.0000 |
| 2B1 | Ammonia production | CH ₄ | 0.26 | 0.28 | 0.0000 | 0.0000 | 1.0000 |
| 1A3c | Railways: solid fuels | N ₂ O | - | 0.09 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | N ₂ O | 0.09 | 0.02 | 0.0000 | 0.0000 | 1.0000 |
| 1B1 | Coal mining and handling solid fuels | CH ₄ | 0.08 | 0.01 | 0.0000 | 0.0000 | 1.0000 |
| Total | | | 801,593.08 | 474,346.12 | 0.4434 | 1.0000 | |

A 1.4 QUANTITATIVE APPROACH 2 KCA FOLLOWING IPCC 2006 GUIDELINES

Following the 2006 IPCC Guidelines, the UK has also completed an Approach 2 KCA for both level and trend, which takes into account uncertainties, using the Approach 1 method for uncertainty estimates. This analysis has been performed using the data shown in **Table A 1.4.1** to **Table A 1.4.4** using the same categorisation and the same estimates of uncertainty.

The results of the level assessment (based on Approach 2) with and without LULUCF for the base year and the latest reported year are shown in **Table A 1.4.1** to **Table A 1.4.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the “Level Parameter”,

and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in level.

The results of the trend assessment (based on Approach 2) with and without LULUCF for the base year to the latest reported year to the latest reported year, are shown in **Table A 1.4.5** to **Table A 1.4.6**

The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use, Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the trend parameter, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in trend.

Any methodological improvements to the uncertainty analysis are discussed in **Annex 2**.

Table A 1.4.1 Approach 2 Level Assessment for Base year (including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|------------------------------|------------------|--|--|------------------|------------------|
| 5A | 5A Solid Waste Disposal | CH ₄ | 60432.98 | 60432.98 | 0.2397 | 0.2397 |
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 19934.61 | 19934.61 | 0.1634 | 0.4031 |
| 1A | 1A Coal | CO ₂ | 243845.39 | 243845.39 | 0.0617 | 0.4648 |
| 4B | 4B Cropland | CO ₂ | 14265.93 | 14265.93 | 0.0526 | 0.5174 |
| 1A | 1A (Stationary) Oil | CO ₂ | 95612.77 | 95612.77 | 0.0509 | 0.5683 |
| 4A | 4A Forest Land | CO ₂ | -15026.26 | 15026.26 | 0.0431 | 0.6114 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 21826.68 | 21826.68 | 0.0360 | 0.6474 |
| 5C | 5C Waste Incineration | CO ₂ | 1300.71 | 1300.71 | 0.0323 | 0.6796 |
| 4C | 4C Grassland | CO ₂ | -7111.03 | 7111.03 | 0.0291 | 0.7088 |
| 4E | 4E Settlements | CO ₂ | 7011.30 | 7011.30 | 0.0287 | 0.7375 |
| 3A | 3A Enteric Fermentation | CH ₄ | 25392.50 | 25392.50 | 0.0286 | 0.7661 |
| 1A | 1A Natural Gas | CO ₂ | 106932.90 | 106932.90 | 0.0170 | 0.7831 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 10168.33 | 10168.33 | 0.0169 | 0.8000 |
| 5D | 5D Wastewater Handling | N ₂ O | 783.59 | 783.59 | 0.0159 | 0.8159 |
| 2B | 2B Chemical industry | HFCs | 17670.77 | 17670.77 | 0.0145 | 0.8304 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 75562.72 | 75562.72 | 0.0138 | 0.8442 |
| 3D | 3D Agricultural Soils | N ₂ O | 13610.39 | 13610.39 | 0.0124 | 0.8566 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO _{2e}) | Absolute value of Base year emissions (Gg CO _{2e}) | Level Assessment | Cumulative Total |
|-----------------------|---|------------------|--|--|------------------|------------------|
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 2141.02 | 2141.02 | 0.0115 | 0.8681 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 7611.13 | 7611.13 | 0.0111 | 0.8792 |
| 2B | 2B Chemical industries | CO ₂ | 6770.09 | 6770.09 | 0.0097 | 0.8890 |
| 5D | 5D Wastewater Handling | CH ₄ | 4218.90 | 4218.90 | 0.0093 | 0.8983 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 1157.96 | 1157.96 | 0.0071 | 0.9054 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 593.57 | 593.57 | 0.0069 | 0.9123 |
| 1A3b | 1A3b DERV | CO ₂ | 33005.80 | 33005.80 | 0.0060 | 0.9183 |
| 4G | 4G Other Activities | CO ₂ | -1639.08 | 1639.08 | 0.0060 | 0.9244 |
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 989.40 | 989.40 | 0.0060 | 0.9304 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1893.15 | 1893.15 | 0.0055 | 0.9359 |
| 4 | 4 Indirect LULUCF Emissions | N ₂ O | 401.49 | 401.49 | 0.0054 | 0.9414 |
| 4D | 4D Wetland | CO ₂ | 486.95 | 486.95 | 0.0040 | 0.9454 |
| 1A3b | 1A3b DERV | N ₂ O | 322.88 | 322.88 | 0.0034 | 0.9488 |
| 1B2 | 1B2 Offshore Oil& Gas | CH ₄ | 2176.74 | 2176.74 | 0.0033 | 0.9521 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 5777.92 | 5777.92 | 0.0033 | 0.9555 |
| 3B | 3B Manure Management | CH ₄ | 4732.53 | 4732.53 | 0.0032 | 0.9587 |
| 2C | 2C Metal Industries | CO ₂ | 7400.69 | 7400.69 | 0.0032 | 0.9619 |
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 3860.26 | 3860.26 | 0.0032 | 0.9650 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1881.31 | 1881.31 | 0.0031 | 0.9681 |
| 4B | 4B Cropland | N ₂ O | 1019.85 | 1019.85 | 0.0029 | 0.9710 |
| 3B | 3B Manure Management | N ₂ O | 3442.70 | 3442.70 | 0.0027 | 0.9737 |
| 1A3 | 1A3 Other diesel | CO ₂ | 1679.92 | 1679.92 | 0.0020 | 0.9757 |
| 2A1 | 2A1 Cement Production | CO ₂ | 7295.26 | 7295.26 | 0.0019 | 0.9776 |
| 4A | 4A Forest land | N ₂ O | 231.22 | 231.22 | 0.0018 | 0.9794 |
| 3G | 3G Liming | CO ₂ | 1015.18 | 1015.18 | 0.0017 | 0.9812 |
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 552.81 | 552.81 | 0.0017 | 0.9829 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO _{2e}) | Absolute value of Base year emissions (Gg CO _{2e}) | Level Assessment | Cumulative Total |
|-----------|--|------------------|--|--|------------------|------------------|
| 3J | 3J OT & CD Agriculture | CH ₄ | 276.73 | 276.73 | 0.0016 | 0.9845 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 1416.64 | 1416.64 | 0.0014 | 0.9859 |
| 3H | 3H Urea application to agriculture | CO ₂ | 327.60 | 327.60 | 0.0013 | 0.9872 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 105.00 | 105.00 | 0.0010 | 0.9882 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 1698.56 | 1698.56 | 0.0010 | 0.9892 |
| 1A4 | 1A4 Peat | CO ₂ | 372.48 | 372.48 | 0.0010 | 0.9902 |
| 1A3b | 1A3b DERV | CH ₄ | 88.74 | 88.74 | 0.0009 | 0.9911 |
| 5C | 5C Waste Incineration | N ₂ O | 46.66 | 46.66 | 0.0009 | 0.9920 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 132.00 | 132.00 | 0.0008 | 0.9928 |
| 4E | 4E Settlements | N ₂ O | 584.68 | 584.68 | 0.0007 | 0.9935 |
| 2A2 | 2A2 Lime Production | CO ₂ | 1462.05 | 1462.05 | 0.0006 | 0.9941 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 149.16 | 149.16 | 0.0006 | 0.9946 |
| 2C | 2C Metal Industries | PFCs | 333.43 | 333.43 | 0.0005 | 0.9952 |
| 5C | 5C Waste Incineration | CH ₄ | 134.83 | 134.83 | 0.0005 | 0.9957 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 912.79 | 912.79 | 0.0005 | 0.9962 |
| 3F | 3F Field Burning | CH ₄ | 186.57 | 186.57 | 0.0004 | 0.9965 |
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 40.75 | 40.75 | 0.0004 | 0.9969 |
| 2B | 2B Chemical Industry | CH ₄ | 204.89 | 204.89 | 0.0003 | 0.9972 |
| 1A | 1A Other (waste) | CO ₂ | 247.13 | 247.13 | 0.0003 | 0.9975 |
| 2A | 2A Minerals industry | CH ₄ | 31.10 | 31.10 | 0.0003 | 0.9978 |
| 2C | 2C Metal Industries | SF ₆ | 387.17 | 387.17 | 0.0002 | 0.9980 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 640.93 | 640.93 | 0.0002 | 0.9982 |
| 1A3 | 1A3 Other diesel | N ₂ O | 16.53 | 16.53 | 0.0002 | 0.9984 |
| 2C | 2C Iron & Steel | N ₂ O | 17.74 | 17.74 | 0.0002 | 0.9985 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 81.64 | 81.64 | 0.0002 | 0.9987 |
| 2A3 | 2A3 Glass production | CO ₂ | 405.54 | 405.54 | 0.0002 | 0.9989 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.80 | 17.80 | 0.0002 | 0.9990 |
| 5B | 5B Biological treatment of solid waste | CH ₄ | 18.13 | 18.13 | 0.0002 | 0.9992 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|--|--|------------------|------------------|
| 2C | 2C Iron & Steel Production | CH ₄ | 36.90 | 36.90 | 0.0001 | 0.9993 |
| 3F | 3F Field Burning | N ₂ O | 57.66 | 57.66 | 0.0001 | 0.9994 |
| 5B | 5B Biological treatment of solid waste | N ₂ O | 12.97 | 12.97 | 0.0001 | 0.9995 |
| 4C | 4C Grassland | N ₂ O | 10.33 | 10.33 | 0.0001 | 0.9996 |
| 2B | 2B Chemical industry | PFCs | 113.90 | 113.90 | 0.0001 | 0.9997 |
| 4C | 4C Grassland | CH ₄ | 9.97 | 9.97 | 0.0000 | 0.9998 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 3.66 | 3.66 | 0.0000 | 0.9998 |
| 4D | 4D Grassland | N ₂ O | 4.13 | 4.13 | 0.0000 | 0.9999 |
| 2E | 2E Electronics Industry | HFCs | 8.73 | 8.73 | 0.0000 | 0.9999 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 6.37 | 6.37 | 0.0000 | 0.9999 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.75 | 2.75 | 0.0000 | 0.9999 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 2.10 | 2.10 | 0.0000 | 1.0000 |
| 4E | 4E Settlements | CH ₄ | 3.00 | 3.00 | 0.0000 | 1.0000 |
| 4A | 4A Forest Land | CH ₄ | 2.89 | 2.89 | 0.0000 | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.83 | 0.83 | 0.0000 | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.31 | 0.31 | 0.0000 | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.44 | 0.44 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 0.18 | 0.18 | 0.0000 | 1.0000 |
| 4B | 4B Cropland | CH ₄ | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | CO ₂ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 4F | 4F Other Land | CO ₂ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 0.00 | 0.00 | 0.0000 | 1.0000 |

Table A 1.4.2 Approach 2 Level Assessment for the latest reported year (including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------------------|--|------------------|---------------------------------------|---|------------------|------------------|
| 5A | 5A Solid Waste Disposal | CH ₄ | 14261.40 | 14261.40 | 0.1151 | 0.1151 |
| 4A | 4A Forest Land | CO ₂ | -18211.77 | 18211.77 | 0.1063 | 0.2214 |
| 4B | 4B Cropland | CO ₂ | 10971.23 | 10971.23 | 0.0824 | 0.3038 |
| 4C | 4C Grassland | CO ₂ | -8861.80 | 8861.80 | 0.0739 | 0.3777 |
| 4E | 4E Settlements | CO ₂ | 6451.34 | 6451.34 | 0.0538 | 0.4315 |
| 1A | 1A Natural Gas | CO ₂ | 158621.57 | 158621.57 | 0.0513 | 0.4828 |
| 3A | 3A Enteric Fermentation | CH ₄ | 21458.44 | 21458.44 | 0.0492 | 0.5319 |
| 1A | 1A (Stationary) Oil | CO ₂ | 45156.95 | 45156.95 | 0.0489 | 0.5808 |
| 5D | 5D Wastewater Handling | N ₂ O | 725.93 | 725.93 | 0.0300 | 0.6108 |
| 1A3b | 1A3b DERV | CO ₂ | 77521.13 | 77521.13 | 0.0289 | 0.6397 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 14166.81 | 14166.81 | 0.0276 | 0.6674 |
| 1A3b | 1A3b DERV | N ₂ O | 1000.43 | 1000.43 | 0.0217 | 0.6890 |
| 3D | 3D Agricultural Soils | N ₂ O | 11466.79 | 11466.79 | 0.0213 | 0.7104 |
| 5B | 5B Biological treatment of solid waste | CH ₄ | 1193.62 | 1193.62 | 0.0207 | 0.7311 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 843.87 | 843.87 | 0.0199 | 0.7510 |
| 1A | 1A Coal | CO ₂ | 34774.45 | 34774.45 | 0.0179 | 0.7689 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 5443.73 | 5443.73 | 0.0162 | 0.7850 |
| 5D | 5D Wastewater Handling | CH ₄ | 3445.75 | 3445.75 | 0.0155 | 0.8005 |
| 4G | 4G Other Activities | CO ₂ | -2015.60 | 2015.60 | 0.0151 | 0.8156 |
| 2B | 2B Chemical industries | CO ₂ | 4931.56 | 4931.56 | 0.0144 | 0.8301 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 1262.84 | 1262.84 | 0.0138 | 0.8439 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 35843.09 | 35843.09 | 0.0133 | 0.8572 |
| 5C | 5C Waste Incineration | CO ₂ | 258.91 | 258.91 | 0.0131 | 0.8703 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 3764.68 | 3764.68 | 0.0127 | 0.8830 |
| 1A | 1A Other (waste) | CO ₂ | 5441.36 | 5441.36 | 0.0126 | 0.8956 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------------------|---|------------------|---------------------------------------|---|------------------|------------------|
| 5B | 5B Biological treatment of solid waste | N ₂ O | 728.03 | 728.03 | 0.0115 | 0.9071 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1476.32 | 1476.32 | 0.0088 | 0.9158 |
| 4 | 4 Indirect LULUCF Emissions | N ₂ O | 243.62 | 243.62 | 0.0067 | 0.9225 |
| 1A3 | 1A3 Other diesel | CO ₂ | 2597.40 | 2597.40 | 0.0063 | 0.9289 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1898.39 | 1898.39 | 0.0063 | 0.9352 |
| 3B | 3B Manure Management | CH ₄ | 4226.81 | 4226.81 | 0.0059 | 0.9411 |
| 4D | 4D Wetland | CO ₂ | 336.78 | 336.78 | 0.0056 | 0.9467 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 4230.64 | 4230.64 | 0.0050 | 0.9516 |
| 3B | 3B Manure Management | N ₂ O | 2814.79 | 2814.79 | 0.0045 | 0.9561 |
| 1B2 | 1B2 Offshore Oil& Gas | CH ₄ | 1170.57 | 1170.57 | 0.0037 | 0.9598 |
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 519.22 | 519.22 | 0.0033 | 0.9631 |
| 3G | 3G Liming | CO ₂ | 939.13 | 939.13 | 0.0033 | 0.9664 |
| 3H | 3H Urea application to agriculture | CO ₂ | 343.95 | 343.95 | 0.0029 | 0.9692 |
| 4B | 4B Cropland | N ₂ O | 461.97 | 461.97 | 0.0027 | 0.9719 |
| 3J | 3J OT & CD Agriculture | CH ₄ | 206.19 | 206.19 | 0.0024 | 0.9744 |
| 2A1 | 2A1 Cement Production | CO ₂ | 4409.79 | 4409.79 | 0.0023 | 0.9767 |
| 4A | 4A Forest land | N ₂ O | 144.65 | 144.65 | 0.0023 | 0.9790 |
| 2C | 2C Metal Industries | CO ₂ | 2572.86 | 2572.86 | 0.0022 | 0.9812 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 483.98 | 483.98 | 0.0016 | 0.9828 |
| 5C | 5C Waste Incineration | N ₂ O | 39.63 | 39.63 | 0.0015 | 0.9844 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 351.76 | 351.76 | 0.0015 | 0.9858 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 183.51 | 183.51 | 0.0014 | 0.9873 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 72.08 | 72.08 | 0.0014 | 0.9887 |
| 4E | 4E Settlements | N ₂ O | 524.99 | 524.99 | 0.0013 | 0.9900 |
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 95.21 | 95.21 | 0.0012 | 0.9912 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 93.01 | 93.01 | 0.0011 | 0.9923 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 85.31 | 85.31 | 0.0011 | 0.9933 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|---------------------------------------|---|------------------|------------------|
| 2A2 | 2A2 Lime Production | CO ₂ | 1052.06 | 1052.06 | 0.0009 | 0.9942 |
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 39.36 | 39.36 | 0.0007 | 0.9949 |
| 4C | 4C Grassland | N ₂ O | 33.24 | 33.24 | 0.0006 | 0.9955 |
| 1A3 | 1A3 Other diesel | N ₂ O | 25.30 | 25.30 | 0.0006 | 0.9961 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 415.66 | 415.66 | 0.0004 | 0.9965 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 359.54 | 359.54 | 0.0004 | 0.9969 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.96 | 17.96 | 0.0003 | 0.9973 |
| 2A3 | 2A3 Glass production | CO ₂ | 367.98 | 367.98 | 0.0003 | 0.9976 |
| 2B | 2B Chemical industry | PFCs | 172.90 | 172.90 | 0.0003 | 0.9979 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 419.65 | 419.65 | 0.0003 | 0.9981 |
| 2B | 2B Chemical Industry | CH ₄ | 61.41 | 61.41 | 0.0002 | 0.9983 |
| 1A3b | 1A3b DERV | CH ₄ | 7.85 | 7.85 | 0.0002 | 0.9985 |
| 2E | 2E Electronics Industry | HFCs | 21.18 | 21.18 | 0.0002 | 0.9987 |
| 4C | 4C Grassland | CH ₄ | 17.78 | 17.78 | 0.0002 | 0.9988 |
| 1A3c | 1A3c Coal | CO ₂ | 41.15 | 41.15 | 0.0001 | 0.9990 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 6.28 | 6.28 | 0.0001 | 0.9991 |
| 2C | 2C Metal Industries | SF ₆ | 109.75 | 109.75 | 0.0001 | 0.9992 |
| 2C | 2C Iron & Steel | N ₂ O | 6.19 | 6.19 | 0.0001 | 0.9993 |
| 2A | 2A Minerals industry | CH ₄ | 5.89 | 5.89 | 0.0001 | 0.9994 |
| 2C | 2C Iron & Steel Production | CH ₄ | 10.68 | 10.68 | 0.0001 | 0.9995 |
| 5C | 5C Waste Incineration | CH ₄ | 10.45 | 10.45 | 0.0001 | 0.9996 |
| 4E | 4E Settlements | CH ₄ | 8.25 | 8.25 | 0.0001 | 0.9997 |
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 37.25 | 37.25 | 0.0001 | 0.9997 |
| 2C | 2C Metal Industries | PFCs | 15.07 | 15.07 | 0.0001 | 0.9998 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.27 | 2.27 | 0.0000 | 0.9998 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 3.78 | 3.78 | 0.0000 | 0.9999 |
| 1A4 | 1A4 Peat | CO ₂ | 4.83 | 4.83 | 0.0000 | 0.9999 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 1.51 | 1.51 | 0.0000 | 0.9999 |
| 4A | 4A Forest Land | CH ₄ | 2.31 | 2.31 | 0.0000 | 0.9999 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|---------------------------------------|---|------------------|------------------|
| 1A3c | 1A3c Coal | CH ₄ | 0.94 | 0.94 | 0.0000 | 1.0000 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 1.55 | 1.55 | 0.0000 | 1.0000 |
| 4D | 4D Grassland | N ₂ O | 0.30 | 0.30 | 0.0000 | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 2.29 | 2.29 | 0.0000 | 1.0000 |
| 2B | 2B Chemical industry | HFCs | 2.55 | 2.55 | 0.0000 | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.53 | 0.53 | 0.0000 | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.33 | 0.33 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 4B | 4B Cropland | CH ₄ | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 4F | 4F Other Land | CO ₂ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 3F | 3F Field Burning | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 3F | 3F Field Burning | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.00 | 0.00 | 0.0000 | 1.0000 |

Table A 1.4.3 Approach 2 Level Assessment for Base year (not including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|----------------------------|------------------|--|--|------------------|------------------|
| 5A | 5A Solid Waste Disposal | CH ₄ | 60432.98 | 60432.98 | 0.2904 | 0.2904 |
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 19934.61 | 19934.61 | 0.1980 | 0.4885 |
| 1A | 1A Coal | CO ₂ | 243845.39 | 243845.39 | 0.0748 | 0.5632 |
| 1A | 1A (Stationary) Oil | CO ₂ | 95612.77 | 95612.77 | 0.0616 | 0.6249 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 21826.68 | 21826.68 | 0.0436 | 0.6684 |
| 5C | 5C Waste Incineration | CO ₂ | 1300.71 | 1300.71 | 0.0391 | 0.7075 |
| 3A | 3A Enteric Fermentation | CH ₄ | 25392.50 | 25392.50 | 0.0346 | 0.7422 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base emissions year (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------------------|--|------------------|--|--|------------------|------------------|
| 1A | 1A Natural Gas | CO ₂ | 106932.90 | 106932.90 | 0.0206 | 0.7628 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 10168.33 | 10168.33 | 0.0204 | 0.7832 |
| 5D | 5D Wastewater Handling | N ₂ O | 783.59 | 783.59 | 0.0193 | 0.8025 |
| 2B | 2B Chemical industry | HFCs | 17670.77 | 17670.77 | 0.0176 | 0.8201 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 75562.72 | 75562.72 | 0.0167 | 0.8368 |
| 3D | 3D Agricultural Soils | N ₂ O | 13610.39 | 13610.39 | 0.0151 | 0.8518 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 2141.02 | 2141.02 | 0.0140 | 0.8658 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 7611.13 | 7611.13 | 0.0135 | 0.8793 |
| 2B | 2B Chemical industries | CO ₂ | 6770.09 | 6770.09 | 0.0118 | 0.8911 |
| 5D | 5D Wastewater Handling | CH ₄ | 4218.90 | 4218.90 | 0.0113 | 0.9023 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 1157.96 | 1157.96 | 0.0086 | 0.9110 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 593.57 | 593.57 | 0.0083 | 0.9193 |
| 1A3b | 1A3b DERV | CO ₂ | 33005.80 | 33005.80 | 0.0073 | 0.9266 |
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 989.40 | 989.40 | 0.0073 | 0.9340 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1893.15 | 1893.15 | 0.0067 | 0.9406 |
| 1A3b | 1A3b DERV | N ₂ O | 322.88 | 322.88 | 0.0042 | 0.9448 |
| 1B2 | 1B2 Offshore Oil& Gas | CH ₄ | 2176.74 | 2176.74 | 0.0040 | 0.9489 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 5777.92 | 5777.92 | 0.0040 | 0.9529 |
| 3B | 3B Manure Management | CH ₄ | 4732.53 | 4732.53 | 0.0039 | 0.9568 |
| 2C | 2C Metal Industries | CO ₂ | 7400.69 | 7400.69 | 0.0038 | 0.9607 |
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 3860.26 | 3860.26 | 0.0038 | 0.9645 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1881.31 | 1881.31 | 0.0037 | 0.9682 |
| 3B | 3B Manure Management | N ₂ O | 3442.70 | 3442.70 | 0.0033 | 0.9715 |
| 1A3 | 1A3 Other diesel | CO ₂ | 1679.92 | 1679.92 | 0.0024 | 0.9739 |
| 2A1 | 2A1 Cement Production | CO ₂ | 7295.26 | 7295.26 | 0.0023 | 0.9762 |
| 3G | 3G Liming | CO ₂ | 1015.18 | 1015.18 | 0.0021 | 0.9783 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base emissions year (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|--|--|------------------|------------------|
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 552.81 | 552.81 | 0.0021 | 0.9804 |
| 3J | 3J OT & CD Agriculture | CH ₄ | 276.73 | 276.73 | 0.0019 | 0.9824 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 1416.64 | 1416.64 | 0.0016 | 0.9840 |
| 3H | 3H Urea application to agriculture | CO ₂ | 327.60 | 327.60 | 0.0016 | 0.9856 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 105.00 | 105.00 | 0.0012 | 0.9868 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 1698.56 | 1698.56 | 0.0012 | 0.9880 |
| 1A4 | 1A4 Peat | CO ₂ | 372.48 | 372.48 | 0.0012 | 0.9892 |
| 1A3b | 1A3b DERV | CH ₄ | 88.74 | 88.74 | 0.0011 | 0.9903 |
| 5C | 5C Waste Incineration | N ₂ O | 46.66 | 46.66 | 0.0011 | 0.9914 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 132.00 | 132.00 | 0.0009 | 0.9923 |
| 2A2 | 2A2 Lime Production | CO ₂ | 1462.05 | 1462.05 | 0.0007 | 0.9931 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 149.16 | 149.16 | 0.0007 | 0.9938 |
| 2C | 2C Metal Industries | PFCs | 333.43 | 333.43 | 0.0007 | 0.9944 |
| 5C | 5C Waste Incineration | CH ₄ | 134.83 | 134.83 | 0.0006 | 0.9950 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 912.79 | 912.79 | 0.0006 | 0.9956 |
| 3F | 3F Field Burning | CH ₄ | 186.57 | 186.57 | 0.0005 | 0.9961 |
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 40.75 | 40.75 | 0.0004 | 0.9965 |
| 2B | 2B Chemical Industry | CH ₄ | 204.89 | 204.89 | 0.0004 | 0.9969 |
| 1A | 1A Other (waste) | CO ₂ | 247.13 | 247.13 | 0.0003 | 0.9972 |
| 2A | 2A Minerals industry | CH ₄ | 31.10 | 31.10 | 0.0003 | 0.9975 |
| 2C | 2C Metal Industries | SF ₆ | 387.17 | 387.17 | 0.0003 | 0.9978 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 640.93 | 640.93 | 0.0002 | 0.9980 |
| 1A3 | 1A3 Other diesel | N ₂ O | 16.53 | 16.53 | 0.0002 | 0.9983 |
| 2C | 2C Iron & Steel | N ₂ O | 17.74 | 17.74 | 0.0002 | 0.9985 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 81.64 | 81.64 | 0.0002 | 0.9987 |
| 2A3 | 2A3 Glass production | CO ₂ | 405.54 | 405.54 | 0.0002 | 0.9989 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.80 | 17.80 | 0.0002 | 0.9991 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | Absolute value of Base year emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|--|--|------------------|------------------|
| 5B | 5B Biological treatment of solid waste | CH ₄ | 18.13 | 18.13 | 0.0002 | 0.9993 |
| 2C | 2C Iron & Steel Production | CH ₄ | 36.90 | 36.90 | 0.0002 | 0.9994 |
| 3F | 3F Field Burning | N ₂ O | 57.66 | 57.66 | 0.0001 | 0.9996 |
| 5B | 5B Biological treatment of solid waste | N ₂ O | 12.97 | 12.97 | 0.0001 | 0.9997 |
| 2B | 2B Chemical industry | PFCs | 113.90 | 113.90 | 0.0001 | 0.9998 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 3.66 | 3.66 | 0.0000 | 0.9999 |
| 2E | 2E Electronics Industry | HFCs | 8.73 | 8.73 | 0.0000 | 0.9999 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 6.37 | 6.37 | 0.0000 | 0.9999 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.75 | 2.75 | 0.0000 | 1.0000 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 2.10 | 2.10 | 0.0000 | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.83 | 0.83 | 0.0000 | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.31 | 0.31 | 0.0000 | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.44 | 0.44 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 0.18 | 0.18 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | CO ₂ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 0.00 | 0.00 | 0.0000 | 1.0000 |

Table A 1.4.4 Approach 2 Level Assessment for the latest reported year (not including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|-------------------------|-----------------|---------------------------------------|---|------------------|------------------|
| 5A | 5A Solid Waste Disposal | CH ₄ | 14261.40 | 14261.40 | 0.1773 | 0.1773 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------------------|--|------------------|---------------------------------------|---|------------------|------------------|
| 1A | 1A Natural Gas | CO ₂ | 158621.57 | 158621.57 | 0.0790 | 0.2563 |
| 3A | 3A Enteric Fermentation | CH ₄ | 21458.44 | 21458.44 | 0.0757 | 0.3321 |
| 1A | 1A (Stationary) Oil | CO ₂ | 45156.95 | 45156.95 | 0.0753 | 0.4074 |
| 5D | 5D Wastewater Handling | N ₂ O | 725.93 | 725.93 | 0.0463 | 0.4537 |
| 1A3b | 1A3b DERV | CO ₂ | 77521.13 | 77521.13 | 0.0445 | 0.4982 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 14166.81 | 14166.81 | 0.0426 | 0.5408 |
| 1A3b | 1A3b DERV | N ₂ O | 1000.43 | 1000.43 | 0.0334 | 0.5742 |
| 3D | 3D Agricultural Soils | N ₂ O | 11466.79 | 11466.79 | 0.0329 | 0.6071 |
| 5B | 5B Biological treatment of solid waste | CH ₄ | 1193.62 | 1193.62 | 0.0319 | 0.6390 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 843.87 | 843.87 | 0.0307 | 0.6697 |
| 1A | 1A Coal | CO ₂ | 34774.45 | 34774.45 | 0.0276 | 0.6973 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 5443.73 | 5443.73 | 0.0249 | 0.7222 |
| 5D | 5D Wastewater Handling | CH ₄ | 3445.75 | 3445.75 | 0.0238 | 0.7460 |
| 2B | 2B Chemical industries | CO ₂ | 4931.56 | 4931.56 | 0.0222 | 0.7682 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 1262.84 | 1262.84 | 0.0213 | 0.7895 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 35843.09 | 35843.09 | 0.0205 | 0.8100 |
| 5C | 5C Waste Incineration | CO ₂ | 258.91 | 258.91 | 0.0201 | 0.8302 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 3764.68 | 3764.68 | 0.0196 | 0.8497 |
| 1A | 1A Other (waste) | CO ₂ | 5441.36 | 5441.36 | 0.0194 | 0.8692 |
| 5B | 5B Biological treatment of solid waste | N ₂ O | 728.03 | 728.03 | 0.0177 | 0.8869 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1476.32 | 1476.32 | 0.0135 | 0.9004 |
| 1A3 | 1A3 Other diesel | CO ₂ | 2597.40 | 2597.40 | 0.0097 | 0.9102 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1898.39 | 1898.39 | 0.0097 | 0.9199 |
| 3B | 3B Manure Management | CH ₄ | 4226.81 | 4226.81 | 0.0091 | 0.9290 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 4230.64 | 4230.64 | 0.0076 | 0.9366 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|---------------------------------------|---|------------------|------------------|
| 3B | 3B Manure Management | N ₂ O | 2814.79 | 2814.79 | 0.0069 | 0.9435 |
| 1B2 | 1B2 Offshore Oil& Gas | CH ₄ | 1170.57 | 1170.57 | 0.0056 | 0.9491 |
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 519.22 | 519.22 | 0.0051 | 0.9542 |
| 3G | 3G Liming | CO ₂ | 939.13 | 939.13 | 0.0050 | 0.9593 |
| 3H | 3H Urea application to agriculture | CO ₂ | 343.95 | 343.95 | 0.0044 | 0.9637 |
| 3J | 3J OT & CD Agriculture | CH ₄ | 206.19 | 206.19 | 0.0037 | 0.9674 |
| 2A1 | 2A1 Cement Production | CO ₂ | 4409.79 | 4409.79 | 0.0036 | 0.9710 |
| 2C | 2C Metal Industries | CO ₂ | 2572.86 | 2572.86 | 0.0035 | 0.9745 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 483.98 | 483.98 | 0.0025 | 0.9770 |
| 5C | 5C Waste Incineration | N ₂ O | 39.63 | 39.63 | 0.0023 | 0.9793 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 351.76 | 351.76 | 0.0023 | 0.9816 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 183.51 | 183.51 | 0.0022 | 0.9838 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 72.08 | 72.08 | 0.0021 | 0.9859 |
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 95.21 | 95.21 | 0.0018 | 0.9878 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 93.01 | 93.01 | 0.0017 | 0.9895 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 85.31 | 85.31 | 0.0016 | 0.9911 |
| 2A2 | 2A2 Lime Production | CO ₂ | 1052.06 | 1052.06 | 0.0014 | 0.9925 |
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 39.36 | 39.36 | 0.0011 | 0.9935 |
| 1A3 | 1A3 Other diesel | N ₂ O | 25.30 | 25.30 | 0.0009 | 0.9944 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 415.66 | 415.66 | 0.0007 | 0.9950 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 359.54 | 359.54 | 0.0007 | 0.9957 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.96 | 17.96 | 0.0005 | 0.9962 |
| 2A3 | 2A3 Glass production | CO ₂ | 367.98 | 367.98 | 0.0005 | 0.9967 |
| 2B | 2B Chemical industry | PFCs | 172.90 | 172.90 | 0.0004 | 0.9971 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 419.65 | 419.65 | 0.0004 | 0.9975 |
| 2B | 2B Chemical Industry | CH ₄ | 61.41 | 61.41 | 0.0003 | 0.9978 |
| 1A3b | 1A3b DERV | CH ₄ | 7.85 | 7.85 | 0.0003 | 0.9981 |
| 2E | 2E Electronics Industry | HFCs | 21.18 | 21.18 | 0.0003 | 0.9983 |

| IPCC Code | IPCC Category | Gas | 2017 emissions (Gg CO ₂ e) | Absolute value of 2017 emissions (Gg CO ₂ e) | Level Assessment | Cumulative Total |
|-----------|---|------------------|---------------------------------------|---|------------------|------------------|
| 1A3c | 1A3c Coal | CO ₂ | 41.15 | 41.15 | 0.0002 | 0.9986 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 6.28 | 6.28 | 0.0002 | 0.9988 |
| 2C | 2C Metal Industries | SF ₆ | 109.75 | 109.75 | 0.0002 | 0.9990 |
| 2C | 2C Iron & Steel | N ₂ O | 6.19 | 6.19 | 0.0002 | 0.9991 |
| 2A | 2A Minerals industry | CH ₄ | 5.89 | 5.89 | 0.0002 | 0.9993 |
| 2C | 2C Iron & Steel Production | CH ₄ | 10.68 | 10.68 | 0.0001 | 0.9994 |
| 5C | 5C Waste Incineration | CH ₄ | 10.45 | 10.45 | 0.0001 | 0.9995 |
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 37.25 | 37.25 | 0.0001 | 0.9996 |
| 2C | 2C Metal Industries | PFCs | 15.07 | 15.07 | 0.0001 | 0.9997 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.27 | 2.27 | 0.0001 | 0.9998 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 3.78 | 3.78 | 0.0000 | 0.9998 |
| 1A4 | 1A4 Peat | CO ₂ | 4.83 | 4.83 | 0.0000 | 0.9999 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 1.51 | 1.51 | 0.0000 | 0.9999 |
| 1A3c | 1A3c Coal | CH ₄ | 0.94 | 0.94 | 0.0000 | 1.0000 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 1.55 | 1.55 | 0.0000 | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 2.29 | 2.29 | 0.0000 | 1.0000 |
| 2B | 2B Chemical industry | HFCs | 2.55 | 2.55 | 0.0000 | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.53 | 0.53 | 0.0000 | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.33 | 0.33 | 0.0000 | 1.0000 |
| 1A3c | 1A3c Coal | N ₂ O | 0.09 | 0.09 | 0.0000 | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.02 | 0.02 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 3F | 3F Field Burning | CH ₄ | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 3F | 3F Field Burning | N ₂ O | 0.00 | 0.00 | 0.0000 | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.00 | 0.00 | 0.0000 | 1.0000 |

Table A 1.4.5 Approach 2 Assessment for Trend (including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------|--|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 19934.61 | 0.00 | 0.0136 | 23.0% | 0.2302 |
| 5A | 5A Solid Waste Disposal | CH ₄ | 60432.98 | 14261.40 | 0.0118 | 20.0% | 0.4303 |
| 1A | 1A Coal | CO ₂ | 243845.39 | 34774.45 | 0.0039 | 6.6% | 0.4958 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 21826.68 | 483.98 | 0.0029 | 4.9% | 0.5445 |
| 1A | 1A Natural Gas | CO ₂ | 106932.90 | 158621.57 | 0.0022 | 3.7% | 0.5819 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 1416.64 | 14166.81 | 0.0018 | 3.1% | 0.6130 |
| 5C | 5C Waste Incineration | CO ₂ | 1300.71 | 258.91 | 0.0018 | 3.0% | 0.6428 |
| 1A3b | 1A3b DERV | CO ₂ | 33005.80 | 77521.13 | 0.0015 | 2.6% | 0.6689 |
| 5B | 5B Biological treatment of solid waste | CH ₄ | 18.13 | 1193.62 | 0.0014 | 2.5% | 0.6934 |
| 4B | 4B Cropland | CO ₂ | 14265.93 | 10971.23 | 0.0014 | 2.4% | 0.7177 |
| 4E | 4E Settlements | CO ₂ | 7011.30 | 6451.34 | 0.0014 | 2.4% | 0.7415 |
| 4A | 4A Forest Land | CO ₂ | -15026.26 | -18211.77 | 0.0013 | 2.2% | 0.7634 |
| 1A3b | 1A3b DERV | N ₂ O | 322.88 | 1000.43 | 0.0012 | 2.1% | 0.7845 |
| 2B | 2B Chemical industry | HFCs | 17670.77 | 2.55 | 0.0012 | 2.0% | 0.8049 |
| 3A | 3A Enteric Fermentation | CH ₄ | 25392.50 | 21458.44 | 0.0011 | 1.8% | 0.8234 |
| 1A | 1A Other (waste) | CO ₂ | 247.13 | 5441.36 | 0.0009 | 1.5% | 0.8381 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 593.57 | 843.87 | 0.0008 | 1.4% | 0.8522 |
| 5B | 5B Biological treatment of solid waste | N ₂ O | 12.97 | 728.03 | 0.0008 | 1.4% | 0.8658 |
| 5D | 5D Wastewater Handling | N ₂ O | 783.59 | 725.93 | 0.0008 | 1.3% | 0.8792 |
| 1A | 1A (Stationary) Oil | CO ₂ | 95612.77 | 45156.95 | 0.0008 | 1.3% | 0.8925 |
| 4C | 4C Grassland | CO ₂ | -7111.03 | -8861.80 | 0.0007 | 1.2% | 0.9048 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 1157.96 | 85.31 | 0.0005 | 0.9% | 0.9136 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 10168.33 | 3764.68 | 0.0005 | 0.9% | 0.9222 |
| 3D | 3D Agricultural Soils | N ₂ O | 13610.39 | 11466.79 | 0.0005 | 0.8% | 0.9301 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------------------|---|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 989.40 | 95.21 | 0.0004 | 0.7% | 0.9372 |
| 5D | 5D Wastewater Handling | CH ₄ | 4218.90 | 3445.75 | 0.0003 | 0.5% | 0.9426 |
| 1A3 | 1A3 Other diesel | CO ₂ | 1679.92 | 2597.40 | 0.0003 | 0.5% | 0.9473 |
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 3860.26 | 37.25 | 0.0003 | 0.4% | 0.9517 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 7611.13 | 5443.73 | 0.0002 | 0.4% | 0.9554 |
| 2B | 2B Chemical industries | CO ₂ | 6770.09 | 4931.56 | 0.0002 | 0.4% | 0.9589 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 75562.72 | 35843.09 | 0.0002 | 0.4% | 0.9624 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1881.31 | 1898.39 | 0.0002 | 0.3% | 0.9656 |
| 4G | 4G Other Activities | CO ₂ | -1639.08 | -2015.60 | 0.0002 | 0.3% | 0.9685 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1893.15 | 1476.32 | 0.0002 | 0.3% | 0.9711 |
| 3B | 3B Manure Management | CH ₄ | 4732.53 | 4226.81 | 0.0001 | 0.2% | 0.9736 |
| 2C | 2C Metal Industries | CO ₂ | 7400.69 | 2572.86 | 0.0001 | 0.2% | 0.9754 |
| 3B | 3B Manure Management | N ₂ O | 3442.70 | 2814.79 | 0.0001 | 0.2% | 0.9770 |
| 3H | 3H Urea application to agriculture | CO ₂ | 327.60 | 343.95 | 0.0001 | 0.2% | 0.9785 |
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 552.81 | 519.22 | 0.0001 | 0.2% | 0.9800 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 81.64 | 351.76 | 0.0001 | 0.2% | 0.9815 |
| 3G | 3G Liming | CO ₂ | 1015.18 | 939.13 | 0.0001 | 0.1% | 0.9830 |
| 1A4 | 1A4 Peat | CO ₂ | 372.48 | 4.83 | 0.0001 | 0.1% | 0.9843 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 5777.92 | 4230.64 | 0.0001 | 0.1% | 0.9856 |
| 1A3b | 1A3b DERV | CH ₄ | 88.74 | 7.85 | 0.0001 | 0.1% | 0.9867 |
| 4D | 4D Wetland | CO ₂ | 486.95 | 336.78 | 0.0001 | 0.1% | 0.9878 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 149.16 | 183.51 | 0.0001 | 0.1% | 0.9887 |
| 4B | 4B Cropland | N ₂ O | 1019.85 | 461.97 | 0.0001 | 0.1% | 0.9896 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 1698.56 | 359.54 | 0.0001 | 0.1% | 0.9905 |
| 2C | 2C Metal Industries | PFCs | 333.43 | 15.07 | 0.0000 | 0.1% | 0.9912 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------------------|--|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 3J | 3J OT & CD Agriculture | CH ₄ | 276.73 | 206.19 | 0.0000 | 0.1% | 0.9918 |
| 4C | 4C Grassland | N ₂ O | 10.33 | 33.24 | 0.0000 | 0.1% | 0.9925 |
| 5C | 5C Waste Incineration | CH ₄ | 134.83 | 10.45 | 0.0000 | 0.1% | 0.9931 |
| 5C | 5C Waste Incineration | N ₂ O | 46.66 | 39.63 | 0.0000 | 0.1% | 0.9937 |
| 4E | 4E Settlements | N ₂ O | 584.68 | 524.99 | 0.0000 | 0.1% | 0.9942 |
| 3F | 3F Field Burning | CH ₄ | 186.57 | 0.00 | 0.0000 | 0.1% | 0.9948 |
| 1A3 | 1A3 Other diesel | N ₂ O | 16.53 | 25.30 | 0.0000 | 0.0% | 0.9952 |
| 4 | 4 Indirect LULUCF Emissions | N ₂ O | 401.49 | 243.62 | 0.0000 | 0.0% | 0.9955 |
| 1B2 | 1B2 Offshore Oil & Gas | CH ₄ | 2176.74 | 1170.57 | 0.0000 | 0.0% | 0.9959 |
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 40.75 | 39.36 | 0.0000 | 0.0% | 0.9962 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 2141.02 | 1262.84 | 0.0000 | 0.0% | 0.9965 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 105.00 | 72.08 | 0.0000 | 0.0% | 0.9968 |
| 2A | 2A Minerals industry | CH ₄ | 31.10 | 5.89 | 0.0000 | 0.0% | 0.9970 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 132.00 | 93.01 | 0.0000 | 0.0% | 0.9972 |
| 2B | 2B Chemical Industry | CH ₄ | 204.89 | 61.41 | 0.0000 | 0.0% | 0.9975 |
| 2B | 2B Chemical industry | PFCs | 113.90 | 172.90 | 0.0000 | 0.0% | 0.9977 |
| 2A2 | 2A2 Lime Production | CO ₂ | 1462.05 | 1052.06 | 0.0000 | 0.0% | 0.9979 |
| 4A | 4A Forest land | N ₂ O | 231.22 | 144.65 | 0.0000 | 0.0% | 0.9981 |
| 1A3c | 1A3c Coal | CO ₂ | 0.00 | 41.15 | 0.0000 | 0.0% | 0.9983 |
| 3F | 3F Field Burning | N ₂ O | 57.66 | 0.00 | 0.0000 | 0.0% | 0.9984 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.80 | 17.96 | 0.0000 | 0.0% | 0.9986 |
| 2C | 2C Metal Industries | SF ₆ | 387.17 | 109.75 | 0.0000 | 0.0% | 0.9988 |
| 2E | 2E Electronics Industry | HFCs | 8.73 | 21.18 | 0.0000 | 0.0% | 0.9989 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 912.79 | 415.66 | 0.0000 | 0.0% | 0.9990 |
| 2A3 | 2A3 Glass production | CO ₂ | 405.54 | 367.98 | 0.0000 | 0.0% | 0.9992 |
| 4C | 4C Grassland | CH ₄ | 9.97 | 17.78 | 0.0000 | 0.0% | 0.9993 |
| 2A1 | 2A1 Cement Production | CO ₂ | 7295.26 | 4409.79 | 0.0000 | 0.0% | 0.9994 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 3.66 | 6.28 | 0.0000 | 0.0% | 0.9995 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------|---|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 2C | 2C Iron & Steel Production | CH ₄ | 36.90 | 10.68 | 0.0000 | 0.0% | 0.9996 |
| 2C | 2C Iron & Steel | N ₂ O | 17.74 | 6.19 | 0.0000 | 0.0% | 0.9997 |
| 4E | 4E Settlements | CH ₄ | 3.00 | 8.25 | 0.0000 | 0.0% | 0.9998 |
| 4D | 4D Grassland | N ₂ O | 4.13 | 0.30 | 0.0000 | 0.0% | 0.9998 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 0.18 | 3.78 | 0.0000 | 0.0% | 0.9999 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 640.93 | 419.65 | 0.0000 | 0.0% | 0.9999 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 6.37 | 1.55 | 0.0000 | 0.0% | 0.9999 |
| 1A3c | 1A3c Coal | CH ₄ | 0.00 | 0.94 | 0.0000 | 0.0% | 1.0000 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.75 | 2.27 | 0.0000 | 0.0% | 1.0000 |
| 4A | 4A Forest Land | CH ₄ | 2.89 | 2.31 | 0.0000 | 0.0% | 1.0000 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 2.10 | 1.51 | 0.0000 | 0.0% | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 0.00 | 2.29 | 0.0000 | 0.0% | 1.0000 |
| 1A3c | 1A3c Coal | N ₂ O | 0.00 | 0.09 | 0.0000 | 0.0% | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.31 | 0.33 | 0.0000 | 0.0% | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.44 | 0.00 | 0.0000 | 0.0% | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.09 | 0.02 | 0.0000 | 0.0% | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.83 | 0.53 | 0.0000 | 0.0% | 1.0000 |
| 4B | 4B Cropland | CH ₄ | 0.09 | 0.02 | 0.0000 | 0.0% | 1.0000 |
| 4F | 4F Other Land | CO ₂ | 0.00 | 0.00 | 0.0000 | 0.0% | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 0.0% | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 0.0% | 1.0000 |

Table A 1.4.6 Approach 2 Assessment for Trend (not including LULUCF) with Key Categories Shaded in Grey

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------|--|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 2B3 | 2B3 Adipic Acid Production | N ₂ O | 19934.61 | 0.00 | 0.0136 | 25.3% | 0.2528 |
| 5A | 5A Solid Waste Disposal | CH ₄ | 60432.98 | 14261.40 | 0.0118 | 22.0% | 0.4724 |
| 1A | 1A Coal | CO ₂ | 243845.39 | 34774.45 | 0.0039 | 7.2% | 0.5444 |
| 1B1 | 1B1 Coal Mining | CH ₄ | 21826.68 | 483.98 | 0.0029 | 5.3% | 0.5979 |
| 1A | 1A Natural Gas | CO ₂ | 106932.90 | 158621.57 | 0.0022 | 4.1% | 0.6389 |
| 2F | 2F Product Uses as Substitutes for ODS | HFCs | 1416.64 | 14166.81 | 0.0018 | 3.4% | 0.6730 |
| 5C | 5C Waste Incineration | CO ₂ | 1300.71 | 258.91 | 0.0018 | 3.3% | 0.7058 |
| 1A3b | 1A3b DERV | CO ₂ | 33005.80 | 77521.13 | 0.0015 | 2.9% | 0.7344 |
| 5B | 5B Biological treatment of solid waste | CH ₄ | 18.13 | 1193.62 | 0.0014 | 2.7% | 0.7613 |
| 1A3b | 1A3b DERV | N ₂ O | 322.88 | 1000.43 | 0.0012 | 2.3% | 0.7844 |
| 2B | 2B Chemical industry | HFCs | 17670.77 | 2.55 | 0.0012 | 2.2% | 0.8068 |
| 3A | 3A Enteric Fermentation | CH ₄ | 25392.50 | 21458.44 | 0.0011 | 2.0% | 0.8271 |
| 1A | 1A Other (waste) | CO ₂ | 247.13 | 5441.36 | 0.0009 | 1.6% | 0.8432 |
| 2G | 2G Other Product Manufacture and Use | N ₂ O | 593.57 | 843.87 | 0.0008 | 1.5% | 0.8587 |
| 5B | 5B Biological treatment of solid waste | N ₂ O | 12.97 | 728.03 | 0.0008 | 1.5% | 0.8737 |
| 5D | 5D Wastewater Handling | N ₂ O | 783.59 | 725.93 | 0.0008 | 1.5% | 0.8885 |
| 1A | 1A (Stationary) Oil | CO ₂ | 95612.77 | 45156.95 | 0.0008 | 1.5% | 0.9030 |
| 1A3b | 1A3b Gasoline/ LPG | CH ₄ | 1157.96 | 85.31 | 0.0005 | 1.0% | 0.9126 |
| 1B2 | 1B2 Natural Gas Transmission | CH ₄ | 10168.33 | 3764.68 | 0.0005 | 0.9% | 0.9220 |
| 3D | 3D Agricultural Soils | N ₂ O | 13610.39 | 11466.79 | 0.0005 | 0.9% | 0.9308 |
| 1A3b | 1A3b Gasoline/ LPG | N ₂ O | 989.40 | 95.21 | 0.0004 | 0.8% | 0.9385 |
| 5D | 5D Wastewater Handling | CH ₄ | 4218.90 | 3445.75 | 0.0003 | 0.6% | 0.9444 |
| 1A3 | 1A3 Other diesel | CO ₂ | 1679.92 | 2597.40 | 0.0003 | 0.5% | 0.9496 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------------------|---|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 2B2 | 2B2 Nitric Acid Production | N ₂ O | 3860.26 | 37.25 | 0.0003 | 0.5% | 0.9544 |
| 1A3d | 1A3d Marine fuel | CO ₂ | 7611.13 | 5443.73 | 0.0002 | 0.4% | 0.9585 |
| 2B | 2B Chemical industries | CO ₂ | 6770.09 | 4931.56 | 0.0002 | 0.4% | 0.9624 |
| 1A3b | 1A3b Gasoline/ LPG | CO ₂ | 75562.72 | 35843.09 | 0.0002 | 0.4% | 0.9662 |
| 1A3a | 1A3a Aviation Fuel | CO ₂ | 1881.31 | 1898.39 | 0.0002 | 0.4% | 0.9697 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1893.15 | 1476.32 | 0.0002 | 0.3% | 0.9727 |
| 3B | 3B Manure Management | CH ₄ | 4732.53 | 4226.81 | 0.0001 | 0.3% | 0.9754 |
| 2C | 2C Metal Industries | CO ₂ | 7400.69 | 2572.86 | 0.0001 | 0.2% | 0.9774 |
| 3B | 3B Manure Management | N ₂ O | 3442.70 | 2814.79 | 0.0001 | 0.2% | 0.9791 |
| 3H | 3H Urea application to agriculture | CO ₂ | 327.60 | 343.95 | 0.0001 | 0.2% | 0.9808 |
| 2D | 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 552.81 | 519.22 | 0.0001 | 0.2% | 0.9825 |
| 1A4 | 1A4 Petroleum Coke | CO ₂ | 81.64 | 351.76 | 0.0001 | 0.2% | 0.9841 |
| 3G | 3G Liming | CO ₂ | 1015.18 | 939.13 | 0.0001 | 0.2% | 0.9857 |
| 1A4 | 1A4 Peat | CO ₂ | 372.48 | 4.83 | 0.0001 | 0.1% | 0.9872 |
| 1B2 | 1B2 Oil & Natural Gas | CO ₂ | 5777.92 | 4230.64 | 0.0001 | 0.1% | 0.9885 |
| 1A3b | 1A3b DERV | CH ₄ | 88.74 | 7.85 | 0.0001 | 0.1% | 0.9898 |
| 2G | 2G Other Product Manufacture and Use | PFCs | 149.16 | 183.51 | 0.0001 | 0.1% | 0.9908 |
| 1B1 | 1B1 Solid Fuel Transformation | CO ₂ | 1698.56 | 359.54 | 0.0001 | 0.1% | 0.9918 |
| 2C | 2C Metal Industries | PFCs | 333.43 | 15.07 | 0.0000 | 0.1% | 0.9925 |
| 3J | 3J OT & CD Agriculture | CH ₄ | 276.73 | 206.19 | 0.0000 | 0.1% | 0.9932 |
| 5C | 5C Waste Incineration | CH ₄ | 134.83 | 10.45 | 0.0000 | 0.1% | 0.9939 |
| 5C | 5C Waste Incineration | N ₂ O | 46.66 | 39.63 | 0.0000 | 0.1% | 0.9945 |
| 3F | 3F Field Burning | CH ₄ | 186.57 | 0.00 | 0.0000 | 0.1% | 0.9951 |
| 1A3 | 1A3 Other diesel | N ₂ O | 16.53 | 25.30 | 0.0000 | 0.0% | 0.9956 |
| 1B2 | 1B2 Offshore Oil& Gas | CH ₄ | 2176.74 | 1170.57 | 0.0000 | 0.0% | 0.9960 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------------------|---|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 1B2 | 1B2 Oil & Natural Gas | N ₂ O | 40.75 | 39.36 | 0.0000 | 0.0% | 0.9963 |
| 1A1 & 1A2 & 1A4 & 1A5 | 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 2141.02 | 1262.84 | 0.0000 | 0.0% | 0.9967 |
| 1A3d | 1A3d Marine fuel | N ₂ O | 105.00 | 72.08 | 0.0000 | 0.0% | 0.9969 |
| 2A | 2A Minerals industry | CH ₄ | 31.10 | 5.89 | 0.0000 | 0.0% | 0.9972 |
| 3J | 3J OT & CD Agriculture | N ₂ O | 132.00 | 93.01 | 0.0000 | 0.0% | 0.9975 |
| 2B | 2B Chemical Industry | CH ₄ | 204.89 | 61.41 | 0.0000 | 0.0% | 0.9977 |
| 2B | 2B Chemical industry | PFCs | 113.90 | 172.90 | 0.0000 | 0.0% | 0.9979 |
| 2A2 | 2A2 Lime Production | CO ₂ | 1462.05 | 1052.06 | 0.0000 | 0.0% | 0.9982 |
| 1A3c | 1A3c Coal | CO ₂ | 0.00 | 41.15 | 0.0000 | 0.0% | 0.9984 |
| 3F | 3F Field Burning | N ₂ O | 57.66 | 0.00 | 0.0000 | 0.0% | 0.9985 |
| 1A3a | 1A3a Aviation Fuel | N ₂ O | 17.80 | 17.96 | 0.0000 | 0.0% | 0.9987 |
| 2C | 2C Metal Industries | SF ₆ | 387.17 | 109.75 | 0.0000 | 0.0% | 0.9989 |
| 2E | 2E Electronics Industry | HFCs | 8.73 | 21.18 | 0.0000 | 0.0% | 0.9991 |
| 2G | 2G Other Product Manufacture and Use | SF ₆ | 912.79 | 415.66 | 0.0000 | 0.0% | 0.9992 |
| 2A3 | 2A3 Glass production | CO ₂ | 405.54 | 367.98 | 0.0000 | 0.0% | 0.9994 |
| 2A1 | 2A1 Cement Production | CO ₂ | 7295.26 | 4409.79 | 0.0000 | 0.0% | 0.9995 |
| 1A3d | 1A3d Marine fuel | CH ₄ | 3.66 | 6.28 | 0.0000 | 0.0% | 0.9996 |
| 2C | 2C Iron & Steel Production | CH ₄ | 36.90 | 10.68 | 0.0000 | 0.0% | 0.9997 |
| 2C | 2C Iron & Steel | N ₂ O | 17.74 | 6.19 | 0.0000 | 0.0% | 0.9998 |
| 1B1 | 1B1 Solid Fuel Transformation | CH ₄ | 0.18 | 3.78 | 0.0000 | 0.0% | 0.9999 |
| 2A4 | 2A4 Other process uses of carbonates | CO ₂ | 640.93 | 419.65 | 0.0000 | 0.0% | 0.9999 |
| 1A3a | 1A3a Aviation Fuel | CH ₄ | 6.37 | 1.55 | 0.0000 | 0.0% | 0.9999 |
| 1A3c | 1A3c Coal | CH ₄ | 0.00 | 0.94 | 0.0000 | 0.0% | 1.0000 |
| 1A3 | 1A3 Other diesel | CH ₄ | 2.75 | 2.27 | 0.0000 | 0.0% | 1.0000 |
| 2B8 | 2B8 Petrochemical and Carbon Black Production | N ₂ O | 2.10 | 1.51 | 0.0000 | 0.0% | 1.0000 |
| 2C | 2C Metal Industries | HFCs | 0.00 | 2.29 | 0.0000 | 0.0% | 1.0000 |

| IPCC Code | IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Trend Assessment with Uncertainty | % Contribution to Trend Uncertainty | Cumulative Total |
|-----------|---|------------------|--|---------------------------------------|-----------------------------------|-------------------------------------|------------------|
| 1A3c | 1A3c Coal | N ₂ O | 0.00 | 0.09 | 0.0000 | 0.0% | 1.0000 |
| 2B1 | 2B1 Ammonia Production | N ₂ O | 0.31 | 0.33 | 0.0000 | 0.0% | 1.0000 |
| 2F | 2F Product Uses as Substitutes for ODS | PFCs | 0.44 | 0.00 | 0.0000 | 0.0% | 1.0000 |
| 1B1 | 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.09 | 0.02 | 0.0000 | 0.0% | 1.0000 |
| 2E | 2E Electronics Industry | NF ₃ | 0.83 | 0.53 | 0.0000 | 0.0% | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | 0.00 | 0.00 | 0.0000 | 0.0% | 1.0000 |
| 2D | 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | 0.00 | 0.00 | 0.0000 | 0.0% | 1.0000 |

A 1.5 KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM

The Key Category Analysis (KCA) ranking system is an additional tool that the UK has developed to aid in the prioritisation of improvement work. The KCA ranking system works by allocating a score based on how high categories rank in the base year and most recent year level assessments and the trend assessment for the approach 1 KCA including LULUCF. For example, if CO₂ from road transport liquid fuel use is the 4th highest by the base year level assessment, 3rd highest by the most recent year level assessment and has the 5th highest trend assessment then its score would be 4+3+5=12. The categories are then ranked from lowest score to highest, with scores that are equal resolved by the most recent year level assessment.

The assessments used in this ranking exercise are only those *including* LULUCF, because if the additional *excluding* LULUCF assessments were also used, the LULUCF sectors would only be included in half of the assessments and would therefore give an unrepresentative weighting.

The results of this ranking are presented in **Table A 1.5.1**.

Table A 1.5.1 KCA Ranking

| KCA rank | IPCC Code | IPCC Category | Greenhouse Gas |
|----------|-----------|-----------------------------------|-----------------|
| 1 | 1A3b | Road transportation: liquid fuels | CO ₂ |
| 2 | 1A1 | Energy industries: solid fuels | CO ₂ |
| 3 | 1A4 | Other sectors: gaseous fuels | CO ₂ |
| 4 | 5A | Solid waste disposal | CH ₄ |
| 5 | 1A1 | Energy industries: gaseous fuels | CO ₂ |

| KCA rank | IPCC Code | IPCC Category | Greenhouse Gas |
|----------|-----------|--|---|
| 6 | 1A2 | Manufacturing industries and construction: gaseous fuels | CO ₂ |
| 7 | 1A1 | Energy industries: liquid fuels | CO ₂ |
| 8 | 1A2 | Manufacturing industries and construction: solid fuels | CO ₂ |
| 9 | 3A1 | Enteric fermentation from Cattle | CH ₄ |
| 10 | 1A2 | Manufacturing industries and construction: liquid fuels | CO ₂ |
| 11 | 4A | Forest land | CO ₂ |
| 12 | 1A4 | Other sectors: liquid fuels | CO ₂ |
| 13 | 3D | Agricultural soils | N ₂ O |
| 14 | 4B | Cropland | CO ₂ |
| 15 | 1A4 | Other sectors: solid fuels | CO ₂ |
| 16 | 1B2 | Oil and gas extraction | CH ₄ |
| 17 | 4E | Settlements | CO ₂ |
| 18 | 1B1 | Coal mining and handling | CH ₄ |
| 19 | 4C | Grassland | CO ₂ |
| 20 | 1A3d | Domestic Navigation: liquid fuels | CO ₂ |
| 21 | 2F1 | Refrigeration and air conditioning | HFCs, PFCs, SF ₆ and NF ₃ |
| 22 | 3B1 | Manure management from Cattle | CH ₄ |
| 23 | 1B2 | Oil and gas extraction | CO ₂ |
| 24 | 3A2 | Enteric fermentation from Sheep | CH ₄ |
| 25 | 1A5 | Other: liquid fuels | CO ₂ |
| 26 | 5D | Wastewater treatment and discharge | CH ₄ |
| 27 | 2C1 | Iron and steel production | CO ₂ |
| 28 | 2B9 | Fluorochemical production | HFCs, PFCs, SF ₆ and NF ₃ |
| 29 | 3B2 | Manure management from Sheep | N ₂ O |
| 30 | 1A3c | Railways: liquid fuels | CO ₂ |
| 31 | 1A1 | Energy industries: other fuels | CO ₂ |
| 32 | 1A3a | Domestic aviation: liquid fuels | CO ₂ |
| 33 | 2A1 | Cement production | CO ₂ |
| 34 | 2F4 | Aerosols | HFCs, PFCs, SF ₆ and NF ₃ |
| 35 | 2B8 | Petrochemical and carbon black production | CO ₂ |
| 36 | 4G | Harvested wood products | CO ₂ |
| 37 | 2B2 | Nitric acid production | N ₂ O |

| KCA rank | IPCC Code | IPCC Category | Greenhouse Gas |
|----------|-----------|-------------------------------------|------------------|
| 38 | 2B3 | Adipic acid production | N ₂ O |
| 39 | 5B | Biological treatment of solid waste | CH ₄ |

A 1.6 APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES

From the 2010 NIR onwards, the NIR contains a list of the Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol. The description below explains the Key Category analysis for Article 3.3 activities and any elected activities under Article 3.4.

Five categories are considered to be key: Article 3.3 Afforestation and Reforestation (CO₂), Article 3.3 Deforestation (CO₂), Article 3.4 Forest Management (CO₂), Article 3.4 Cropland Management (CO₂) and Article 3.4 Grazing Land Management (CO₂). These have been assessed according to the 2006 IPCC good practice guidance for KP (Chapter 2, Section 2.3.6). The numbers have been compared with Table A 1.4.2. The key category analysis for the latest reported year based on level of emissions (including LULUCF) is given in section 11.6.1 of the main NIR.

A 1.7 USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY

The key category analysis is used to prioritise and plan improvements. The approach the UK takes to achieve this is described in **Section 1.2.2.5. Table 1.7 to Table 1.11 in Chapter 1** show the key category summary tables.

A 1.8 TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3

Table A 1.8.1 below is Table NIR 3, containing a summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol³.

Table A 1.8.1 Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol

| KEY CATEGORIES OF EMISSIONS AND REMOVALS | GAS | CRITERIA USED FOR KEY CATEGORY IDENTIFICATION | | | COMMENTS (3) |
|---|-----------------|--|--|--|--|
| | | Associated category in UNFCCC inventory (1) is key (indicate which category) | Category contribution is greater than the smallest category considered key in the UNFCCC inventory (1), (4) (including LULUCF) | Other (2) | |
| Specify key categories according to the national level of disaggregation used (1) | | | | | |
| Afforestation and Reforestation | CO ₂ | Land converted to forest land | Yes | Associated UNFCCC category (4A) is key | The associated UNFCCC inventory category is a key category for level. |
| Deforestation | CO ₂ | Land converted to cropland, Land converted to grassland, Land converted to settlements | Yes | Associated UNFCCC categories (4B2, 4C2 and 4E2) are key. | The associated UNFCCC inventory categories are key categories for level and trend. |

³ Table NIR 3 can be found in FCCC/KP/CMP/2007/9/Add.2.

| KEY CATEGORIES OF EMISSIONS AND REMOVALS | GAS | CRITERIA USED FOR KEY CATEGORY IDENTIFICATION | | | COMMENTS (3) |
|--|-----------------|--|--|---|--|
| | | Associated category in UNFCCC inventory (1) is key (indicate which category) | Category contribution is greater than the smallest category considered key in the UNFCCC inventory (1), (4) (including LULUCF) | Other (2) | |
| Forest Management | CO ₂ | Forest land remaining forest land, Land converted to forest land | Yes | Associated UNFCCC category (4A) is key | The associated UNFCCC inventory category is a key category for level and the Forest Management category contribution is greater than the smallest UNFCCC key category. |
| Cropland Management | CO ₂ | Cropland remaining Cropland, Land converted to Cropland | Yes | Associated UNFCCC category (4B) is key. | The associated UNFCCC inventory category is a key category for level and trend and the Cropland Management category contribution is greater than the smallest UNFCCC key category. |
| Grazing Land Management | CO ₂ | Grassland remaining Grassland, Land converted to Grassland | Yes | Associated UNFCCC category (4C) is key. | The associated UNFCCC inventory category is a key category for level and trend and the Grazing Land Management category contribution is greater than the smallest UNFCCC key category. |

(1) See section 5.4 of the IPCC good practice guidance for LULUCF

(2) This should include qualitative consideration as per Section 5.4.3 of the IPCC Good Practice Guidance for LULUCF or any other criteria

(3) Describe the criteria identifying the category as key

(4) If the emissions or removals of the category exceed the emissions of the smallest category identified as key in the UNFCCC inventory (including LULUCF), Parties should indicate YES. If not, Parties should indicate NO

ANNEX 2: Assessment of Uncertainty

Uncertainty estimates are calculated using two methods: Approach 1 (error propagation) and Approach 2 (Monte Carlo simulation). These are not to be confused with Approaches 1 and 2 for Key Category Analysis, of which Approach 2 KCA uses Approach 1 uncertainties to account for uncertainty in determining Key Categories.

The uncertainty assessment estimates uncertainties according to IPCC sector in addition to presenting estimates by direct greenhouse gas. Estimated uncertainty presented for the sector breakdown used in UK Official Statistics are not reported here, since the categories are not consistent with the requirements of the UK's commitments under the UNFCCC and Kyoto Protocol.

Uncertainty parameters for new sources and sources which have been significantly revised are reviewed each year, particularly for sources which have a significant impact on overall uncertainties.

The overall method used to estimate uncertainties is described below, and the work to improve the accuracy of the uncertainty analysis continues. The key category analysis used data from the uncertainty analysis, and the results of the key category analysis are given in **Annex 1**.

A 2.1 ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1)

The IPCC 2006 Guidelines defines error propagation and Monte Carlo modelling approaches to estimating uncertainties in national greenhouse gas inventories. The results of the error propagation approach are shown in **Table A 2.1.1**. The uncertainties used in the error propagation approach are not exactly the same as those used in the Monte Carlo Simulation since the error propagation source categorisation is less detailed and has a more simplistic approach to uncertainties. The Approach 1 uncertainties assumes all parameters are normally distributed (which means it doesn't account for the skew, kurtosis or any other non-normal features of the expected distributions), and does not account for variations in uncertainty in the time series unlike the Monte Carlo approach which takes into account these factors. The parameters used for the Approach 1 uncertainties for both the base year and the most recent year are the values given for the most recent year in **Table A 2.3.1** to **Table A 2.3.4**.

A 2.1.1 Key Categories

Certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. Key source categories in this respect are identified using Approach 1 uncertainties in the Approach 2 KCA. These have been identified so that the resources available for inventory preparation may be prioritised, and the best possible estimates prepared for the most significant source categories. We have used the method described in Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (Approach 2 to identify key categories).

The results of this key category analysis can be found in **Annex 1**.

A 2.1.2 Tables of uncertainty estimates from the error propagation approach

Table A 2.1.1 Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|-------------------------------|-----------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 1A (Stationary) Oil | CO ₂ | 95,612.77 | 45,156.95 | 5.97% | 2.55% | 6.5% | 0.0000 | 1.274% | 5.632% | 0.0325% | 0.4754% | 0.0023% |
| 1A Coal | CO ₂ | 243,845.39 | 34,774.45 | 1.28% | 2.81% | 3.1% | 0.0000 | 13.238% | 4.337% | 0.3721% | 0.0783% | 0.0014% |
| 1A Natural Gas | CO ₂ | 106,932.90 | 158,621.57 | 1.01% | 1.65% | 1.9% | 0.0000 | 12.041% | 19.782% | 0.1989% | 0.2838% | 0.0012% |
| 1A Other (waste) | CO ₂ | 247.13 | 5,441.36 | 1.08% | 13.86% | 13.9% | 0.0000 | 0.661% | 0.679% | 0.0916% | 0.0104% | 0.0001% |
| 1A3 Other diesel | CO ₂ | 1,679.92 | 2,597.40 | 14.47% | 1.94% | 14.6% | 0.0000 | 0.203% | 0.324% | 0.0039% | 0.0663% | 0.0000% |
| 1A3a Aviation Fuel | CO ₂ | 1,881.31 | 1,898.39 | 19.65% | 3.24% | 19.9% | 0.0000 | 0.101% | 0.237% | 0.0033% | 0.0658% | 0.0000% |
| 1A3b DERV | CO ₂ | 33,005.80 | 77,521.13 | 1.00% | 2.00% | 2.2% | 0.0000 | 7.281% | 9.668% | 0.1456% | 0.1367% | 0.0004% |
| 1A3b Gasoline/ LPG | CO ₂ | 75,562.72 | 35,843.09 | 0.99% | 1.99% | 2.2% | 0.0000 | 0.987% | 4.470% | 0.0196% | 0.0629% | 0.0000% |
| 1A3c Coal | CO ₂ | - | 41.15 | 20.00% | 6.00% | 20.9% | 0.0000 | 0.005% | 0.005% | 0.0003% | 0.0015% | 0.0000% |
| 1A3d Marine fuel | CO ₂ | 7,611.13 | 5,443.73 | 17.71% | 1.77% | 17.8% | 0.0000 | 0.129% | 0.679% | 0.0023% | 0.1701% | 0.0003% |
| 1A4 Peat | CO ₂ | 372.48 | 4.83 | 30.00% | 10.00% | 31.6% | 0.0000 | 0.026% | 0.001% | 0.0026% | 0.0003% | 0.0000% |
| 1A4 Petroleum Coke | CO ₂ | 81.64 | 351.76 | 20.00% | 15.00% | 25.0% | 0.0000 | 0.038% | 0.044% | 0.0057% | 0.0124% | 0.0000% |
| 1B1 Solid Fuel Transformation | CO ₂ | 1,698.56 | 359.54 | 5.17% | 4.84% | 7.1% | 0.0000 | 0.078% | 0.045% | 0.0038% | 0.0033% | 0.0000% |
| 1B2 Oil & Natural Gas | CO ₂ | 5,777.92 | 4,230.64 | 4.49% | 5.40% | 7.0% | 0.0000 | 0.110% | 0.528% | 0.0060% | 0.0335% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|---|-----------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 2A1 Cement Production | CO ₂ | 7,295.26 | 4,409.79 | 1.00% | 3.00% | 3.2% | 0.0000 | 0.023% | 0.550% | 0.0007% | 0.0078% | 0.0000% |
| 2A2 Lime Production | CO ₂ | 1,462.05 | 1,052.06 | 0.00% | 5.00% | 5.0% | 0.0000 | 0.026% | 0.131% | 0.0013% | 0.0000% | 0.0000% |
| 2A3 Glass production | CO ₂ | 405.54 | 367.98 | 0.00% | 5.00% | 5.0% | 0.0000 | 0.017% | 0.046% | 0.0008% | 0.0000% | 0.0000% |
| 2A4 Other process uses of carbonates | CO ₂ | 640.93 | 419.65 | 2.00% | 3.00% | 3.6% | 0.0000 | 0.006% | 0.052% | 0.0002% | 0.0015% | 0.0000% |
| 2B Chemical industries | CO ₂ | 6,770.09 | 4,931.56 | 17.29% | 2.94% | 17.5% | 0.0000 | 0.126% | 0.615% | 0.0037% | 0.1504% | 0.0002% |
| 2C Metal Industries | CO ₂ | 7,400.69 | 2,572.86 | 1.18% | 5.09% | 5.2% | 0.0000 | 0.214% | 0.321% | 0.0109% | 0.0054% | 0.0000% |
| 2D Non Energy Products from Fuels and Solvent Use | CO ₂ | 552.81 | 519.22 | 26.78% | 27.38% | 38.3% | 0.0000 | 0.025% | 0.065% | 0.0068% | 0.0245% | 0.0000% |
| 3G Liming | CO ₂ | 1,015.18 | 939.13 | 0.00% | 20.90% | 20.9% | 0.0000 | 0.044% | 0.117% | 0.0092% | 0.0000% | 0.0000% |
| 3H Urea application to agriculture | CO ₂ | 327.60 | 343.95 | 0.00% | 50.00% | 50.0% | 0.0000 | 0.019% | 0.043% | 0.0096% | 0.0000% | 0.0000% |
| 4A Forest Land | CO ₂ | -15,026.26 | -18,211.77 | 1.00% | 35.00% | 35.0% | 0.0002 | 1.186% | 2.271% | 0.4151% | 0.0321% | 0.0017% |
| 4B Cropland | CO ₂ | 14,265.93 | 10,971.23 | 1.00% | 45.00% | 45.0% | 0.0001 | 0.338% | 1.368% | 0.1519% | 0.0193% | 0.0002% |
| 4C Grassland | CO ₂ | -7,111.03 | -8,861.80 | 1.00% | 50.00% | 50.0% | 0.0001 | 0.592% | 1.105% | 0.2958% | 0.0156% | 0.0009% |
| 4D Wetland | CO ₂ | 486.95 | 336.78 | 1.00% | 100.00% | 100.0% | 0.0000 | 0.007% | 0.042% | 0.0068% | 0.0006% | 0.0000% |
| 4E Settlements | CO ₂ | 7,011.30 | 6,451.34 | 1.00% | 50.00% | 50.0% | 0.0000 | 0.298% | 0.805% | 0.1490% | 0.0114% | 0.0002% |
| 4F Other Land | CO ₂ | - | - | 0.00% | 0.00% | 0.0% | - | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|--|-----------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 4G Other Activities | CO ₂ | -1,639.08 | -2,015.60 | 1.00% | 45.00% | 45.0% | 0.0000 | 0.133% | 0.251% | 0.0598% | 0.0036% | 0.0000% |
| 5C Waste Incineration | CO ₂ | 1,300.71 | 258.91 | 300.00% | 40.00% | 302.7% | 0.0000 | 0.062% | 0.032% | 0.0247% | 0.1370% | 0.0002% |
| 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | CH ₄ | 1,893.15 | 1,476.32 | 0.71% | 35.55% | 35.6% | 0.0000 | 0.047% | 0.184% | 0.0168% | 0.0019% | 0.0000% |
| 1A3 Other diesel | CH ₄ | 2.75 | 2.27 | 15.00% | 130.00% | 130.9% | 0.0000 | 0.000% | 0.000% | 0.0001% | 0.0001% | 0.0000% |
| 1A3a Aviation Fuel | CH ₄ | 6.37 | 1.55 | 14.18% | 55.65% | 57.4% | 0.0000 | 0.000% | 0.000% | 0.0001% | 0.0000% | 0.0000% |
| 1A3b DERV | CH ₄ | 88.74 | 7.85 | 1.00% | 130.00% | 130.0% | 0.0000 | 0.005% | 0.001% | 0.0071% | 0.0000% | 0.0000% |
| 1A3b Gasoline/ LPG | CH ₄ | 1,157.96 | 85.31 | 1.00% | 74.94% | 74.9% | 0.0000 | 0.073% | 0.011% | 0.0547% | 0.0002% | 0.0000% |
| 1A3c Coal | CH ₄ | - | 0.94 | 20.00% | 110.00% | 111.8% | 0.0000 | 0.000% | 0.000% | 0.0001% | 0.0000% | 0.0000% |
| 1A3d Marine fuel | CH ₄ | 3.66 | 6.28 | 19.14% | 124.42% | 125.9% | 0.0000 | 0.001% | 0.001% | 0.0006% | 0.0002% | 0.0000% |
| 1B1 Coal Mining | CH ₄ | 21,826.68 | 483.98 | 2.00% | 20.00% | 20.1% | 0.0000 | 1.516% | 0.060% | 0.3032% | 0.0017% | 0.0009% |
| 1B1 Solid Fuel Transformation | CH ₄ | 0.18 | 3.78 | 0.00% | 49.81% | 49.8% | 0.0000 | 0.000% | 0.000% | 0.0002% | 0.0000% | 0.0000% |
| 1B2 Natural Gas Transmission | CH ₄ | 10,168.33 | 3,764.68 | 3.00% | 20.00% | 20.2% | 0.0000 | 0.265% | 0.470% | 0.0530% | 0.0199% | 0.0000% |
| 1B2 Offshore Oil & Gas | CH ₄ | 2,176.74 | 1,170.57 | 4.54% | 18.15% | 18.7% | 0.0000 | 0.011% | 0.146% | 0.0020% | 0.0094% | 0.0000% |
| 2A Minerals industry | CH ₄ | 31.10 | 5.89 | 0.00% | 100.00% | 100.0% | 0.0000 | 0.002% | 0.001% | 0.0015% | 0.0000% | 0.0000% |
| 2B Chemical Industry | CH ₄ | 204.89 | 61.41 | 0.00% | 20.00% | 20.0% | 0.0000 | 0.007% | 0.008% | 0.0014% | 0.0000% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|---|-----------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 2C Iron & Steel Production | CH ₄ | 36.90 | 10.68 | 1.91% | 47.83% | 47.9% | 0.0000 | 0.001% | 0.001% | 0.0006% | 0.0000% | 0.0000% |
| 2D Non-energy Products from Fuels and Solvent Use | CH ₄ | - | - | 0.00% | 0.00% | 0.0% | - | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 3A Enteric Fermentation | CH ₄ | 25,392.50 | 21,458.44 | 13.73% | 0.00% | 13.7% | 0.0000 | 0.842% | 2.676% | 0.0000% | 0.5198% | 0.0027% |
| 3B Manure Management | CH ₄ | 4,732.53 | 4,226.81 | 0.00% | 8.37% | 8.4% | 0.0000 | 0.185% | 0.527% | 0.0155% | 0.0000% | 0.0000% |
| 3F Field Burning | CH ₄ | 186.57 | - | 25.61% | 0.00% | 25.6% | - | 0.013% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 3J OT & CD Agriculture | CH ₄ | 276.73 | 206.19 | 50.00% | 50.00% | 70.7% | 0.0000 | 0.006% | 0.026% | 0.0029% | 0.0182% | 0.0000% |
| 4A Forest Land | CH ₄ | 2.89 | 2.31 | 1.00% | 55.00% | 55.0% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 4B Cropland | CH ₄ | 0.09 | 0.02 | 1.00% | 55.00% | 55.0% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 4C Grassland | CH ₄ | 9.97 | 17.78 | 1.00% | 55.00% | 55.0% | 0.0000 | 0.001% | 0.002% | 0.0008% | 0.0000% | 0.0000% |
| 4E Settlements | CH ₄ | 3.00 | 8.25 | 1.00% | 55.00% | 55.0% | 0.0000 | 0.001% | 0.001% | 0.0004% | 0.0000% | 0.0000% |
| 5A Solid Waste Disposal | CH ₄ | 60,432.98 | 14,261.40 | 15.00% | 46.00% | 48.4% | 0.0002 | 2.585% | 1.779% | 1.1891% | 0.3773% | 0.0156% |
| 5B Biological treatment of solid waste | CH ₄ | 18.13 | 1,193.62 | 30.00% | 99.50% | 103.9% | 0.0000 | 0.148% | 0.149% | 0.1468% | 0.0632% | 0.0003% |
| 5C Waste Incineration | CH ₄ | 134.83 | 10.45 | 7.40% | 44.45% | 45.1% | 0.0000 | 0.008% | 0.001% | 0.0038% | 0.0001% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|---|------------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 5D Wastewater Handling | CH ₄ | 4,218.90 | 3,445.75 | 10.00% | 25.00% | 26.9% | 0.0000 | 0.125% | 0.430% | 0.0312% | 0.0608% | 0.0000% |
| 1A1 & 1A2 & 1A4 & 1A5 Other Combustion | N ₂ O | 2,141.02 | 1,262.84 | 0.66% | 65.65% | 65.7% | 0.0000 | 0.003% | 0.157% | 0.0019% | 0.0015% | 0.0000% |
| 1A3 Other diesel | N ₂ O | 16.53 | 25.30 | 15.00% | 130.00% | 130.9% | 0.0000 | 0.002% | 0.003% | 0.0025% | 0.0007% | 0.0000% |
| 1A3a Aviation Fuel | N ₂ O | 17.80 | 17.96 | 19.65% | 108.08% | 109.9% | 0.0000 | 0.001% | 0.002% | 0.0010% | 0.0006% | 0.0000% |
| 1A3b DERV | N ₂ O | 322.88 | 1,000.43 | 1.00% | 130.00% | 130.0% | 0.0000 | 0.101% | 0.125% | 0.1319% | 0.0018% | 0.0002% |
| 1A3b Gasoline/ LPG | N ₂ O | 989.40 | 95.21 | 0.99% | 74.47% | 74.5% | 0.0000 | 0.060% | 0.012% | 0.0444% | 0.0002% | 0.0000% |
| 1A3c Coal | N ₂ O | - | 0.09 | 20.00% | 110.00% | 111.8% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 1A3d Marine fuel | N ₂ O | 105.00 | 72.08 | 17.56% | 114.16% | 115.5% | 0.0000 | 0.001% | 0.009% | 0.0016% | 0.0022% | 0.0000% |
| 1B1 Fugitive Emissions from Solid Fuels | N ₂ O | 0.09 | 0.02 | 1.00% | 118.00% | 118.0% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 1B2 Oil & Natural Gas | N ₂ O | 40.75 | 39.36 | 4.77% | 105.01% | 105.1% | 0.0000 | 0.002% | 0.005% | 0.0021% | 0.0003% | 0.0000% |
| 2B1 Ammonia Production | N ₂ O | 0.31 | 0.33 | 2.00% | 50.00% | 50.0% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 2B2 Nitric Acid Production | N ₂ O | 3,860.26 | 37.25 | 0.00% | 10.00% | 10.0% | 0.0000 | 0.274% | 0.005% | 0.0274% | 0.0000% | 0.0000% |
| 2B3 Adipic Acid Production | N ₂ O | 19,934.61 | - | 2.00% | 100.00% | 100.0% | - | 1.440% | 0.000% | 1.4397% | 0.0000% | 0.0207% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|---|------------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 2B8 Petrochemical and Carbon Black Production | N ₂ O | 2.10 | 1.51 | 10.00% | 100.00% | 100.5% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 2C Iron & Steel | N ₂ O | 17.74 | 6.19 | 1.00% | 118.00% | 118.0% | 0.0000 | 0.001% | 0.001% | 0.0006% | 0.0000% | 0.0000% |
| 2D Non-energy Products from Fuels and Solvent Use | N ₂ O | - | - | 0.00% | 0.00% | 0.0% | - | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 2G Other Product Manufacture and Use | N ₂ O | 593.57 | 843.87 | 100.00% | 100.00% | 141.4% | 0.0000 | 0.062% | 0.105% | 0.0624% | 0.1488% | 0.0003% |
| 3B Manure Management | N ₂ O | 3,442.70 | 2,814.79 | 0.00% | 9.53% | 9.5% | 0.0000 | 0.102% | 0.351% | 0.0098% | 0.0000% | 0.0000% |
| 3D Agricultural Soils | N ₂ O | 13,610.39 | 11,466.79 | 0.00% | 11.16% | 11.2% | 0.0000 | 0.447% | 1.430% | 0.0499% | 0.0000% | 0.0000% |
| 3F Field Burning | N ₂ O | 57.66 | - | 25.62% | 0.00% | 25.6% | - | 0.004% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 3J OT & CD Agriculture | N ₂ O | 132.00 | 93.01 | 50.00% | 50.00% | 70.7% | 0.0000 | 0.002% | 0.012% | 0.0010% | 0.0082% | 0.0000% |
| 4 Indirect LULUCF Emissions | N ₂ O | 401.49 | 243.62 | 1.00% | 165.00% | 165.0% | 0.0000 | 0.001% | 0.030% | 0.0023% | 0.0004% | 0.0000% |
| 4A Forest land | N ₂ O | 231.22 | 144.65 | 1.00% | 95.00% | 95.0% | 0.0000 | 0.001% | 0.018% | 0.0013% | 0.0003% | 0.0000% |
| 4B Cropland | N ₂ O | 1,019.85 | 461.97 | 1.00% | 35.00% | 35.0% | 0.0000 | 0.016% | 0.058% | 0.0056% | 0.0008% | 0.0000% |
| 4C Grassland | N ₂ O | 10.33 | 33.24 | 1.00% | 115.00% | 115.0% | 0.0000 | 0.003% | 0.004% | 0.0039% | 0.0001% | 0.0000% |
| 4D Grassland | N ₂ O | 4.13 | 0.30 | 1.00% | 100.00% | 100.0% | 0.0000 | 0.000% | 0.000% | 0.0003% | 0.0000% | 0.0000% |
| 4E Settlements | N ₂ O | 584.68 | 524.99 | 1.00% | 15.00% | 15.0% | 0.0000 | 0.023% | 0.065% | 0.0035% | 0.0009% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|--|------------------|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
| 5B Biological treatment of solid waste | N ₂ O | 12.97 | 728.03 | 30.00% | 90.00% | 94.9% | 0.0000 | 0.090% | 0.091% | 0.0809% | 0.0385% | 0.0001% |
| 5C Waste Incineration | N ₂ O | 46.66 | 39.63 | 7.00% | 230.00% | 230.1% | 0.0000 | 0.002% | 0.005% | 0.0036% | 0.0005% | 0.0000% |
| 5D Wastewater Handling | N ₂ O | 783.59 | 725.93 | 10.00% | 248.00% | 248.2% | 0.0000 | 0.034% | 0.091% | 0.0841% | 0.0128% | 0.0001% |
| 2C Metal Industries | SF ₆ | 387.17 | 109.75 | 5.00% | 5.00% | 7.1% | 0.0000 | 0.014% | 0.014% | 0.0007% | 0.0010% | 0.0000% |
| 2G Other Product Manufacture and Use | SF ₆ | 912.79 | 415.66 | 0.00% | 6.17% | 6.2% | 0.0000 | 0.014% | 0.052% | 0.0009% | 0.0000% | 0.0000% |
| 2B Chemical industry | HFCs | 17,670.77 | 2.55 | 0.00% | 10.00% | 10.0% | 0.0000 | 1.276% | 0.000% | 0.1276% | 0.0000% | 0.0002% |
| 2C Metal Industries | HFCs | - | 2.29 | 5.00% | 10.00% | 11.2% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 2E Electronics Industry | HFCs | 8.73 | 21.18 | 0.00% | 47.15% | 47.1% | 0.0000 | 0.002% | 0.003% | 0.0009% | 0.0000% | 0.0000% |
| 2F Product Uses as Substitutes for ODS | HFCs | 1,416.64 | 14,166.81 | 8.23% | 8.30% | 11.7% | 0.0000 | 1.664% | 1.767% | 0.1382% | 0.2057% | 0.0006% |
| 2E Electronics Industry | NF ₃ | 0.83 | 0.53 | 0.00% | 47.15% | 47.1% | 0.0000 | 0.000% | 0.000% | 0.0000% | 0.0000% | 0.0000% |
| 2B Chemical industry | PFCs | 113.90 | 172.90 | 0.00% | 10.00% | 10.0% | 0.0000 | 0.013% | 0.022% | 0.0013% | 0.0000% | 0.0000% |
| 2C Metal Industries | PFCs | 333.43 | 15.07 | 0.00% | 20.00% | 20.0% | 0.0000 | 0.022% | 0.002% | 0.0044% | 0.0000% | 0.0000% |
| 2F Product Uses as Substitutes for ODS | PFCs | 0.44 | - | 0.00% | 25.00% | 25.0% | - | 0.0000% | 0.0000% | 0.0000% | 0.0000% | 0.0000% |
| 2G Other Product Manufacture and Use | PFCs | 149.16 | 183.51 | 0.00% | 47.15% | 47.1% | 0.0000 | 0.0121% | 0.0229% | 0.0057% | 0.0000% | 0.0000% |

| IPCC Category | Gas | Base year emissions (Gg CO ₂ e) | 2017 emissions (Gg CO ₂ e) | Activity data uncertainty (%) | Emission factor uncertainty (%) | Combined uncertainty (%) | Contribution to variance by Category in 2017 | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
|---------------|-----|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|
|---------------|-----|--|---------------------------------------|-------------------------------|---------------------------------|--------------------------|--|--------------------|--------------------|---|--|---|

| | |
|--|------|
| Percentage uncertainty in total inventory: | 3.0% |
|--|------|

| | |
|-------------------|------|
| Trend uncertainty | 2.3% |
|-------------------|------|

A 2.2 ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2)

A 2.2.1 Overview of the Method

Quantitative estimates of the uncertainties in the emissions were calculated using a Monte Carlo simulation. This corresponds to the IPCC Approach 2 method, discussed in the 2006 Guidelines (IPCC, 2006). The background to the implementation of the Monte Carlo simulation is described in detail by Eggleston *et al* (1998), with the estimates reported here revised to reflect changes in the latest inventory and improvements made in the model. This section gives a brief summary of the methodology, assumptions and results of the simulation.

The computational procedure is detailed below.

- A probability distribution function (PDF) was allocated to each unique emission factor and piece of activity data. The PDFs were mostly normal or log-normal, with more specific distributions given to a handful of sources. The parameters of the PDFs were set by analysing the available data on emission factors and activity data, and by expert judgement;
- A calculation was set up to estimate the total emissions of each gas for the years 1990 and the latest reported year;
- Using the software tool @RISK™, each PDF was sampled at least 20,000 times, such that the emission calculations performed produced a converged output distribution;
- The distribution of errors in the parameter values was calculated from the difference between 2.5 and 97.5 percentile values in the distribution, as a percentage of the distribution mean; and,
- The uncertainty in the trend between 1990 and the latest reported year, according to gas, was also estimated. This is expressed as the 95% confidence interval for the percentage reduction in emissions between the latest year and 1990.

A 2.2.2 Methodological details of the Monte Carlo model

A 2.2.2.1 Uncertainty Distributions

Nearly all of the distributions of emissions from sources in the inventory are modelled using normal or log normal distributions, with more specific distributions given to a handful of sources. The specific distributions include log-logistic, Pearson and Gamma distributions. The primary use of custom distributions is for agriculture; these are fitted distributions that reflect the results of an agriculture-specific Monte Carlo analysis done by Rothamsted Research which accounts for the various factors that influence the modelled agriculture emissions.

Emissions from landfill have been modelled using a custom distribution. Aitchison *et al.* (cited in Eggleston *et al.*, 1998) estimated the uncertainty for landfill emissions using Monte Carlo analysis and found it to be skewed. The distribution histogram was used to generate an empirical distribution of emissions. We examined the distribution and fitted a log normal distribution to Aitchison's data. The emissions are scaled according to the mean estimate of landfill emissions for each year.

There are a couple of other specific distributions for F-gases and waste water which reflect specific distributions we expect for those sources.

A 2.2.2.2 Correlations

The Monte Carlo model contains a number of correlations. If A and B are correlated, then if emissions are under or overestimated from A it would be expected to be over or underestimated by a similar amount from B.

The type and implementation of the correlations has been examined as part of a review (Abbott *et al.*, 2007). The sensitivity analysis that we have completed on the Monte Carlo model suggest that the uncertainties are not sensitive to the correlations between emission factors for fuel used, and for LULUCF sources.

A 2.2.2.2.1 Across years

In running this simulation, it was necessary to make assumptions about the degree of correlation between sources in 1990 and the latest reported year. If source emission factors are correlated this will have the effect of reducing the trend uncertainty, but will not affect uncertainties on emission totals in 1990 or the latest inventory year. The trend estimated by the Monte Carlo model is particularly sensitive to N₂O emissions from agricultural soils.

A 2.2.2.2.2 Between Sources in the same year

In many cases the same factors, or factors derived on the same basis are used for multiple sources. In these cases, we'd say that the emission factors are correlated. For example, the coal emissions factors for N₂O used for cement industry use may be the same as coal use in other industrial combustion due to lack of a more specific factor, in this case we may say the two factors are correlated. Omitting these correlations leads to an underestimate of emissions in any given year.

A 2.2.2.3 Simulation Method

Following recommendations in the 2006 IPCC Guidelines, the model uses a true Monte Carlo sampling method.

A 2.2.3 Quality Control Checks on the Monte Carlo Model Output

A number of quality control checks are completed as part of the uncertainty analysis.

A 2.2.3.1 Checks against totals of the national emissions

To ensure the emissions in the Monte Carlo model closely agree with the reported totals in the NIR, the emissions in the model were checked against the national totals both before and after the simulation was run. The central estimates from the model are expected to be similar to the reported emissions totals, but are not expected to match exactly.

A 2.2.3.2 Inter-comparison between the output of the error propagation and Monte Carlo models

A formal check to compare the output of the error propagation and Monte Carlo model is completed. The results of this comparison are discussed in **Section A 2.6**.

A 2.2.3.3 Calculation of uncertainty on the total

The uncertainty on the 1990 and the most recent year emissions was calculated using two different methods;

$$\text{i) Using } \frac{1.96s.d}{\mu}$$

$$\text{ii) Using } \frac{(97.5 \text{ percentile} - 2.5 \text{ percentile})}{2 \times \mu}$$

The first method uses the standard deviation calculated by @RISK and the mean to give an percentage uncertainty, while the second method uses the 95% confidence interval given by the percentiles quoted. When a distribution is completely normally distributed, the two methods should give the same results. However, when a distribution is skewed the two methods diverge, since the variance is dominated by outliers which aren't necessarily accounted for in the 95% confidence interval. The overall uncertainty quoted in **Table A 2.4.1** is calculated using the second method so that uncertainties in sectors that show a skewed distribution (such as agricultural soils and N₂O) are better represented.

Calculating the uncertainty using both of these methods allows us to check that the Monte Carlo analysis is behaving in the way we would expect, and that convergence of the distributions is being achieved. Comparing the results using both calculations showed that the uncertainties were almost the same for gases where the distributions used were predominantly normal, but higher for N₂O and the GWP weighted total, as expected.

A 2.3 UNCERTAINTY PARAMETERS

The following sections present the uncertainties in emissions, and the trend in emissions according to gas.

A 2.3.1 Uncertainty Parameters used

Table A 2.3.1 to **Table A 2.3.4** summarise the uncertainty parameters used for both Approach 1 and 2 uncertainties. For all of these tables the following apply:

- Uncertainties expressed as 0.5*R/E where R is the difference between 2.5 and 97.5 percentiles and E is the mean;
- Where custom distributions are used for the Approach 2 uncertainties the parameters are not used directly, but the below parameters should still be a reasonable indicator of the uncertainty in the distribution used for Approach 2;
- (r) means revised in comparison to previous NIR; and
- (a) means uncertainty for emission factors and activity cannot be separated, so one uncertainty that represents both is displayed.

Uncertainties **A2**

Table A 2.3.1 Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO₂) inventory

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A | Lubricants | 50.00% | 5.00% | 50.00% | 5.00% | It's challenging to determine the proportion of lubricant used as a fuel, hence a high activity uncertainty. |
| 1A1 | Blast Furnace Gas | 1.50% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |
| 1A1 | Coke Oven Coke | 1.00% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |
| 1A1 | Coke Oven Gas | 1.50% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |
| 1A1 | Colliery Methane | 5.00% | 5.00% | 5.00% | 5.00% | (Minor fuel in sector context) |
| 1A1 | Gas/Diesel Oil | 1.80% | 2.10% | 1.75% | 2.10% | ETS-based data, so low uncertainty. |
| 1A1 | Liquefied Petroleum Gas | 25.70% | 2.10% | 2.50% | 2.10% | The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented |
| 1A1 | Motor Gasoline | 2.50% | 2.10% | 2.50% | 2.10% | Outside of 1A3, the motor gasoline allocations are probably much more uncertain as they are reliant on the off-road model etc., so chosen 2.5%. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of motor gasoline is well understood and documented |
| 1A1 | Municipal Solid Waste | 1.00% | 15.00% | 1.00% | 15.00% | MSW quantity is known accurately. Uncertainty is in mass of fossil carbon per tonne of residual MSW. This is based on reasonable waste composition data from peer reviewed sources, adapted from landfill data. |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-----------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A1 | Naphtha | 50.00% | 5.00% | 50.00% | 5.00% | DUKES are uncertain about where naphtha is used (or not), so a high activity uncertainty has been chosen. EF uncertainty chosen as 5%. The content of naphtha is quite variable - it contains a huge range of hydrocarbons from C5 up to C70+, so the exact carbon content is variable and there are about 5 different grades of naphtha according to UKPIA. |
| 1A1 | Natural Gas | 2.80% | 2.00% | 1.00% | 2.00% | ETS-based data, so low uncertainties. |
| 1A1 | Orimulsion | 5.00% | 5.00% | 5.00% | 5.00% | (Minor fuel in sector context) |
| 1A1 | Other Bituminous Coal | 2.00% | 2.00% | 2.00% | 2.00% | ETS-based data, so low uncertainties. |
| 1A1 | Other Kerosene | 1.25% | 5.00% | 1.25% | 5.00% | ETS-based data, so low uncertainties. |
| 1A1 | Other Oil: Other | 11.90% | 5.00% | 10.00% | 5.00% | (Minor fuel in sector context) |
| 1A1 | Petroleum Coke | 7.80% | 10.00% | 5.00% | 10.00% | ETS-based data, so low uncertainties. 10% chosen for EF uncertainty as there is only a small dataset for the quality of petcoke used in the sector and the CEF could be quite variable depending on the source of the pet coke. |
| 1A1 | Refinery Gas | 50.00% | 20.00% | 25.00% | 15.00% | Comparisons between EUETS and DUKES are variable over time. Risk that in earlier years the "own use" may have been mis-reported to energy stats. High uncertainty on AD. Also a variable quality fuel, so the EF is also uncertain. |
| 1A1 | Residual Fuel Oil | 5.50% | 2.55% | 1.25% | 2.55% | ETS-based data, so low uncertainties. |
| 1A1 | Scrap Tyres | 15.00% | 10.00% | 15.00% | 10.00% | Limited reported use of this fuel; only a small amount of reporting (typically cement kilns) within EUETS and a modest number of fuel quality analyses either through the BCA/MPA (trade body) or the EUETS. Also some variability in the fossil C versus bio-C content of the tyres adds to EF uncertainty. |
| 1A2 | Blast Furnace Gas | 1.50% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A2 | Coke Oven Coke | 3.00% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |
| 1A2 | Coke Oven Gas | 3.00% | 10.00% | 1.00% | 10.00% | Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10% |
| 1A2 | Colliery Methane | 5.00% | 5.00% | 5.00% | 5.00% | (Minor fuel in sector context) |
| 1A2 | Gas/Diesel Oil | 20.00% | 2.00% | 20.00% | 2.00% | Low EF uncertainty as the composition of gas oil is well understood across the time series. The AD for stationary combustion in industrial sectors is quite uncertain, however. DUKES does not distinguish between mobile and stationary sources, and other AD data sources (e.g. EU ETS) have limited coverage of gas oil use across all of 1A2. |
| 1A2 | Liquefied Petroleum Gas | 25.70% | 2.10% | 2.50% | 2.10% | The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented |
| 1A2 | Motor Gasoline | 20.00% | 2.10% | 20.00% | 2.10% | Outside of 1A3, the motor gasoline allocations are probably much more uncertain. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of motor gasoline is well understood and documented |
| 1A2 | Municipal Solid Waste | 5.00% | 15.00% | 5.00% | 15.00% | MSW quantity is known accurately. Uncertainty is in mass of fossil carbon per tonne of residual MSW. This is based on reasonable waste composition data from peer reviewed sources, adapted from landfill data. |
| 1A2 | Natural Gas | 2.80% | 3.00% | 1.00% | 3.00% | Low EF uncertainty as gas composition is monitored and reported across much of the time series, and the fuel has narrow compositional range. AD are also well understood and low uncertainty. Gas supplier data to DUKES can be checked against periodic data matching (meter point data against industry sector information). |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-----------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A2 | Other Bituminous Coal | 5.00% | 10.00% | 5.00% | 10.00% | Limited compositional data over time (e.g. EU ETS data for coal is incomplete), so EF uncertainty reflects the range of composition for coal types in 1A2. AD uncertainty is moderate for 1A2, reflecting energy supplier reporting to BEIS. |
| 1A2 | Other Kerosene | 6.00% | 2.00% | 6.00% | 2.00% | (Minor fuel in sector context) |
| 1A2 | Other Oil: Other | 5.00% | 50.00% | 5.00% | 3.00% | (Minor fuel in sector context) |
| 1A2 | Patent Fuel | 10.00% | 3.00% | 10.00% | 3.00% | (Minor fuel in sector context) |
| 1A2 | Petroleum Coke | 25.00% | 15.00% | 20.00% | 15.00% | EF uncertainty reflects range of petcoke composition that may be used for fuel in 1A2. AD uncertainty is quite high as we have limited data from DUKES and not much AD from EU ETS on petcoke use. |
| 1A2 | Refinery Gas | 50.00% | 15.00% | 50.00% | 15.00% | (Minor fuel in sector context) |
| 1A2 | Residual Fuel Oil | 5.50% | 2.10% | 1.50% | 2.10% | Low EF uncertainty as the composition of fuel oil is well understood across the time series. The AD uncertainty is low in recent years as the fuel is not widely used other than by larger operators that report under EUETS. Moderate uncertainty in earlier years, when fewer routine annual AD sources. |
| 1A2 | Scrap Tyres | 15.00% | 10.00% | 15.00% | 10.00% | (See 1A1 comment – same applies here.) |
| 1A2 | non-fuel combustion | 50.00% | 100.00% | 50.00% | 100.00% | (Minor emission source in sector context) |
| 1A3 | Aviation Gasoline | 20.00% | 3.30% | 20.00% | 3.30% | Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split |
| 1A3 | Other Gas/Diesel Oil | 15.00% | 2.00% | 15.00% | 2.00% | (Minor fuel in sector context) |
| 1A3 | Jet Gasoline | 20.00% | 3.30% | 20.00% | 3.30% | Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split |
| 1A3 | Other Bituminous Coal | 20.00% | 6.00% | 20.00% | 6.00% | (Minor fuel in sector context) |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A3 | liquid biofuels | 5.00% (n) | 5.00% (n) | 5.00% (n) | 5.00% (n) | Activity data are not very uncertain, as it's taken from RTFO data. There is a total potential range of 10% variability in the fossil fuel carbon content of FAME (i.e. judging from the contents of the different fatty acid types used to synthesize the FAME, the highest content is around 44.8g/kg, whilst the lowest is 40.2g/kg). In reality, these are the extremes, so a lower overall uncertainty is expected. the other liquid biofuels are consumed in much smaller quantities than FAME. |
| 1A3b | Gas/Diesel Oil | 1.80% | 2.00% | 1.00% | 2.00% | Low EF uncertainty as the composition of gas oil is well understood across the time series. Low AD uncertainty as good corroboration between fuel sales data and estimates based on vehicle movement data. |
| 1A3b | Liquefied Petroleum Gas | 5.00% | 2.00% | 5.00% | 2.00% | EF uncertainty is consistent with gas oil - the makeup of LPG is well understood and documented. Not a major fuel in the sector but AD are considered moderately uncertain. |
| 1A3b | Motor Gasoline | 1.00% | 2.00% | 1.00% | 2.00% | Low EF uncertainty as the composition of petrol is well understood across the time series. Low AD uncertainty as good corroboration between fuel sales data and estimates based on vehicle movement data. |
| 1A3d | Gas/Diesel Oil | 20.00% | 2.00% | 20.00% | 2.00% | Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split |
| 1A3d | Residual Fuel Oil | 20.00% | 2.00% | 20.00% | 2.00% | Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split |
| 1A4 | Anthracite | 1.50% | 6.00% | 1.00% | 6.00% | Low AD uncertainty as tax data helps establish residential use. EF uncertainty reflects variability in anthracite composition. |
| 1A4 | Coke Oven Coke | 3.00% | 10.00% | 1.00% | 10.00% | (Minor fuel in sector context) |
| 1A4 | Gas/Diesel Oil | 30.00% | 2.00% | 30.00% | 2.00% | Low EF uncertainty as the composition of gas oil is well understood across the time series. High AD uncertainty as scarce data on use of this fuel, e.g. in mobile machinery, in 1A4. |
| 1A4 | Gas Works Gas | 5.00% | 5.00% | 5.00% | 5.00% | (Minor fuel in sector context) |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A4 | Liquefied Petroleum Gas | 25.70% | 2.10% | 2.50% | 2.10% | The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented |
| 1A4 | Motor Gasoline | 50.00% | 2.00% | 50.00% | 2.00% | Low EF uncertainty as the composition of petrol is well understood across the time series. High AD uncertainty as scarce data on use of this fuel in mobile machinery in 1A4. |
| 1A4 | Natural Gas | 2.80% | 3.00% | 2.00% | 3.00% | (As for 1A2) |
| 1A4 | Other Bituminous Coal | 3.00% | 10.00% | 3.00% | 10.00% | Chosen 3% activity uncertainty as we know that there are some limitations on the coal allocation to small-scale users. |
| 1A4 | Other Kerosene | 3.00% | 2.00% | 3.00% | 2.00% | Low AD uncertainty as tax data helps establish residential use. EF uncertainty reflects narrow range of fuel composition. |
| 1A4 | Patent Fuel | 3.30% | 3.00% | 2.00% | 3.00% | (Minor fuel in sector context) |
| 1A4 | Peat | 30.00% | 10.00% | 30.00% | 10.00% | (Minor fuel in sector context) |
| 1A4 | Petroleum Coke | 20.00% | 15.00% | 20.00% | 15.00% | Limited information on the AD of use in domestic fuels which increases uncertainty. Moderate emission factor uncertainty as there is only a small dataset for the quality of petcoke used in the sector and the CEF could be quite variable depending on the source of the pet coke. |
| 1A4 | Residual Fuel Oil | 5.50% | 2.10% | 3.00% | 2.10% | (Minor fuel in sector context) |
| 1A5 | Gas/Diesel Oil | 6.25% | 2.05% | 6.25% | 2.05% | Moderate AD uncertainty as data from very few data suppliers. EF uncertainty reflects narrow range of fuel composition. |
| 1A5 | Jet Gasoline | 10.00% | 3.00% | 10.00% | 3.00% | Activity Data comes directly from fuel users so should have high confidence. |
| 1B1 | Coke Oven Gas | 1.50% | 10.00% | 1.00% | 10.00% | (Minor fuel in sector context) |
| 1B1 | Other Bituminous Coal | 1.50% | 6.00% | 1.50% | 6.00% | EF uncertainty reflects the range of composition of coal types in SSF manufacture. AD uncertainty is quite low, reflecting the small number of operators and high level of AD reporting. |
| 1B1 | petroleum coke | 20.00% | 10.00% | 20.00% | 10.00% | (Minor fuel in sector context) |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|---------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1B2a | non-fuel combustion | 5.00% | 6.00% | 5.00% | 6.00% | (Minor source in UK context) |
| 1B2b | non-fuel combustion | 3.00% | 6.00% | 3.00% | 6.00% | (Minor source in UK context) |
| 1B2c | non-fuel combustion | 5.00% | 6.00% | 5.00% | 6.00% | (Minor source in UK context) |
| 2A1 | non-fuel combustion | 1.00% | 3.00% | 1.00% | 3.00% | EU ETS-type data collected from BCA for all sites so assume very good quality and complete. |
| 2A2 | non-fuel combustion | 10.00% | 5.00% | (a) | 5.00% | High level of reporting in EUETS for recent years and EF reflects small range of data for carbonates used in lime production. AD uncertainty higher in earlier years. |
| 2A3 | non-fuel combustion | (a) | 5.00% | (a) | 5.00% | Mostly based on ETS data. Very small sites outside EU ETS; it's not certain how well EU ETS factor will apply to these non-EU ETS sites. |
| 2A4 | non-fuel combustion | 2.00% | 3.00% | 2.00% | 3.00% | (Minor source in UK context) |
| 2B | Natural Gas | 2.80% | 1.25% | 1.75% | 1.25% | Covers both feedstock and fuel (i.e. total fuel used at the sites), so AD should be very good. |
| 2B | Coke | 1.00% | 20.00% | 1.00% | 10.00% | (Minor source in UK context) |
| 2B | petroleum coke | 1.00% | 10.00% | 1.00% | 10.00% | (Minor source in UK context) |
| 2B | OPG | (a) | 5.00% | (a) | 5.00% | Moderate uncertainty in EF reflecting good level of reporting of fuel quality in EUETS but range of variability of process off-gases that are generated and used in the chemical sector. |
| 2B | non-fuel combustion | 2.00% | 5.00% | 2.00% | 5.00% | (Minor source in UK context) |
| 2B | coke oven coke | (a) | 20.00% | (a) | 20.00% | (Minor source in UK context) |
| 2B | refinery gas | 30.00% | 5.00% | 30.00% | 5.00% | High uncertainty, as we deviate from DUKES. Low emission factor uncertainty, but not a well-characterised fuel. |
| 2C | Blast Furnace Gas | 2.00% | 10.00% | 2.00% | 10.00% | Overall uncertainty in 2C is quite low and uncertainty is more about where the carbon input (from the coking coal) ends up being emitted, and less about the overall amount of carbon emitted. |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|---------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 2C | non-fuel combustion | 2.00% | 10.00% | 2.00% | 10.00% | Overall uncertainty in 2C is quite low and uncertainty is more about where the carbon input (from the coking coal) ends up being emitted, and less about the overall amount of carbon emitted. |
| 2C | Coke | 2.00% | 10.00% | 2.00% | 10.00% | Good level of reporting from I&S operators across the time series. |
| 2C | Petroleum Coke | 10.00% | 7.50% | 10.00% | 7.50% | (Minor source in UK context) |
| 2C | coke oven coke | 2.00% | 5.00% | 2.00% | 5.00% | Activity data has low uncertainty since it's based on ETS/ISSB/DUKES. Emissions are based on regulator data, so low uncertainty. |
| 2D | Lubricants | 50.00% | 50.00% | 50.00% | 50.00% | It's challenging to determine the size of the recovered lubricant market, as this is outside the scope of energy statistics, hence a high activity uncertainty. The fraction of lubricant incidentally oxidised is also highly uncertain, so should be reflected in a high EF uncertainty. |
| 2D | Petroleum Coke | 20.00% | 30.00% | 20.00% | 30.00% | (Minor source in UK context) |
| 2D | Petroleum Waxes | 10.00% | 50.00% | 10.00% | 50.00% | (Minor source in UK context) |
| 2D | non-fuel combustion | 25.00% | 2.00% | 25.00% | 2.00% | Some uncertainty as to the proportion of HDVs requiring urea and how much is needed per vehicle. Very low EF uncertainty because carbon content of urea solution known accurately. |
| 3G | non-fuel combustion | (a) | 20.90% | (a) | 20.90% | Reflects overall uncertainty of AD and EF for carbonate application to soils. |
| 3H | non-fuel combustion | (a) | 50.00% | (a) | 50.00% | (Minor source in UK context) |
| 4A | non-fuel combustion | 1.00% | 35.00% | 1.00% | 35.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 4B | non-fuel combustion | 1.00% | 45.00% | 1.00% | 45.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 4C | non-fuel combustion | 1.00% | 50.00% | 1.00% | 50.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 4D | non-fuel combustion | 1.00% | 100.00% | 1.00% | 100.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-----------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 4E | non-fuel combustion | 1.00% | 50.00% | 1.00% | 50.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 4F | non-fuel combustion | (a) | 50.00% | (a) | 50.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 4G | non-fuel combustion | 1.00% | 45.00% | 1.00% | 45.00% | High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter. |
| 5C | Municipal Solid Waste | 300.00% | 40.00% | 1.00% | 15.00% | (Minor source in UK context) |
| 5C | Clinical waste | 300.00% | 40.00% | 5.00% | 20.00% | (Minor source in UK context) |
| 5C | Chemical waste | 300.00% | 40.00% | 10.00% | 30.00% | (Minor source in UK context) |
| 5C | non-fuel combustion | 300.00% | 40.00% | 300.00% | 40.00% | Unauthorised and widely dispersed activity, estimated from indirect data sources so high uncertainty. Significant uncertainty in the composition of material burnt. |

Table A 2.3.2 Estimated uncertainties in the activity data and emission factors used in the methane (CH₄) inventory

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-----------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A1 | | 1.50% | 50.00% | 1.00% | 50.00% | Minor source but uncertainty mainly reflects uncertainty in the EF from combustion of biomass. |
| 1A2 | | 1.50% | 50.00% | 1.00% | 50.00% | As above. |
| 1A3 | Aviation Gasoline | 20.00% | 78.50% | 20.00% | 78.50% | (Minor source in UK context) |
| 1A3 | Jet Gasoline | 20.00% | 78.50% | 20.00% | 78.50% | (Minor source in UK context) |
| 1A3 | Other Gas/Diesel Oil | 15.00% | 130.00% | 15.00% | 130.00% | (Minor source in UK context) |
| 1A3 | Other Bituminous Coal | 20.00% | 110.00% | 20.00% | 110.00% | (Minor source in UK context) |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A3b | Gas/Diesel Oil | 1.80% | 130.00% | 1.00% | 130.00% | Road transport fuel sales well documented, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in EFs for the range of vehicle (car, van, HGV) and road types. |
| 1A3b | Motor Gasoline | 1.00% | 75.00% | 1.00% | 75.00% | Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in EFs for petrol cars and road types. Lower uncertainty than diesel vehicles because consumption dominated by only one vehicle type. |
| 1A3b | Liquefied Petroleum Gas | 5.00% | 130.00% | 5.00% | 130.00% | (Minor source in UK context) |
| 1A3d | Gas/Diesel Oil | 20.00% | 130.00% | 20.00% | 130.00% | Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. Uncertainty in EF should be consistent with other 1A3 gas oil |
| 1A3d | Residual Fuel Oil | 20.00% | 130.00% | 20.00% | 130.00% | Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. |
| 1A4 | | 1.50% | 50.00% | 1.00% | 50.00% | Minor source but uncertainty mainly reflects uncertainty in the EF from combustion of biomass. |
| 1A5 | | 7.07% | 65.55% | 7.07% | 65.55% | (Minor source in UK context) |
| 1B1 | Coke Oven Gas | 1.50% | 50.00% | 1.00% | 50.00% | (Minor source in UK context) |
| 1B1 | non-fuel combustion | 2.00% | 20.00% | 2.00% | 20.00% | High EF uncertainty reflects the modelled estimates of emissions from coal mines. |
| 1B1 | wood | (a) | 50.00% | (a) | 50.00% | (Minor source in UK context) |
| 1B2a | non-fuel combustion | 5.00% | 20.00% | 5.00% | 20.00% | Good level of operator reporting. EF uncertainty reflects limited scope for verifying the factors applied in deriving operator estimates from fugitives especially. |
| 1B2b | non-fuel combustion | 3.00% | 20.00% | 3.00% | 20.00% | As above. |
| 1B2c | non-fuel combustion | 5.00% | 20.00% | 5.00% | 20.00% | As above. |
| 2A4 | | (a) | 100.00% | (a) | 100.00% | (Minor source in UK context) |
| 2B | | (a) | 20.00% | (a) | 20.00% | (Minor source in UK context) |
| 2C | Blast Furnace Gas | 2.00% | 50.00% | 2.00% | 50.00% | (Minor source in UK context) |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-----------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 2C | non-fuel combustion | 1.00% | 50.00% | 1.00% | 50.00% | (Minor source in UK context) |
| 2C | coke oven coke | 2.00% | 50.00% | 2.00% | 50.00% | Activity data has low uncertainty since it's based on ETS/ISSB/DUKES. Emissions are based on literature factors, so a high EF uncertainty. |
| 2D | | 50.00% | 50.00% | 50.00% | 50.00% | (Minor source in UK context) |
| 3A | non-fuel combustion | 13.73% | (a) | 13.73% | (a) | Based on monte carlo analysis for the agriculture model |
| 3B | non-fuel combustion | (a) (r) | 8.37% (r) | (a) (r) | 8.37% (r) | Based on monte carlo analysis for the agriculture model |
| 3F | non-fuel combustion | 25.61% | (a) | 25.61% | (a) | Based on monte carlo analysis for the agriculture model |
| 3J | non-fuel combustion | 50.00% (n) | 50.00% (n) | 50.00% (n) | 50.00% (n) | Based on monte carlo analysis for the agriculture model |
| 4A | non-fuel combustion | 1.00% | 55.00% | 1.00% | 55.00% | (Minor source in UK context) |
| 4B | non-fuel combustion | 1.00% | 55.00% | 1.00% | 55.00% | (Minor source in UK context) |
| 4C | non-fuel combustion | 1.00% | 55.00% | 1.00% | 55.00% | (Minor source in UK context) |
| 4E | non-fuel combustion | 1.00% | 55.00% | 1.00% | 55.00% | (Minor source in UK context) |
| 5A | non-fuel combustion | 15.00% | 46.00% | 15.00% | 46.00% | Moderate/high uncertainty in historical waste data, rates of decomposition and generation of methane in the modelled approach. Some extrapolation of data needed for methane utilisation, hence high uncertainty overall, across AD and EF. |
| 5B | | 30.00% | 99.50% | 30.00% | 99.50% | Scarce data for UK biological treatments. High uncertainty. |
| 5C | Municipal Solid Waste | 5.00% | 75.00% | 1.00% | 75.00% | (Minor source in UK context) |
| 5C | non-fuel combustion | 5.00% | 50.00% | 5.00% | 50.00% | (Minor source in UK context) |
| 5C | wood | 50.00% (n) | 50.00% (n) | 50.00% (n) | 50.00% (n) | (Minor source in UK context) |
| 5D | non-fuel combustion | 10.00% | 25.00% | 10.00% | 25.00% | UK industry research and model. Moderate-high uncertainty. |

Uncertainties **A2**

Table A 2.3.3 Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N₂O) inventory

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|-------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1A1 | | 1.50% | 100.00% | 1.00% | 100.00% | |
| 1A2 | | 1.50% | 100.00% | 1.00% | 100.00% | |
| 1A3 | Aviation Gasoline | 20.00% | 110.00% | 20.00% | 110.00% | |
| 1A3 | Jet Gasoline | 20.00% | 110.00% | 20.00% | 110.00% | |
| 1A3 | Other Gas/Diesel Oil | 15.00% | 130.00% | 15.00% | 130.00% | |
| 1A3 | Other Bituminous Coal | 20.00% | 110.00% | 20.00% | 110.00% | |
| 1A3b | Gas/Diesel Oil | 1.80% | 130.00% | 1.00% | 130.00% | Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in Efs for different diesel vehicle types and road types. |
| 1A3b | Motor Gasoline | 1.00% | 75.00% | 1.00% | 75.00% | Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in Efs for petrol cars and road types. Lower uncertainty than diesel vehicles because consumption dominated by only one vehicle type. |
| 1A3b | Liquefied Petroleum Gas | 5.00% | 130.00% | 5.00% | 130.00% | |
| 1A3d | Gas/Diesel Oil | 20.00% | 130.00% | 20.00% | 130.00% | Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. Uncertainty in EF should be consistent with other 1A3 gas oil |
| 1A3d | Residual Fuel Oil | 20.00% | 130.00% | 20.00% | 130.00% | Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. |
| 1A4 | | 1.50% | 100.00% | 1.00% | 100.00% | |
| 1A5 | | 7.07% | 85.15% | 7.07% | 85.15% | |
| 1B1 | | 1.50% | 118.00% | 1.00% | 118.00% | |
| 1B2a | non-fuel combustion | 5.00% | 110.00% | 5.00% | 110.00% | |
| 1B2b | non-fuel combustion | 5.00% | 110.00% | 5.00% | 110.00% | |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|---------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|---|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 1B2c | non-fuel combustion | 5.00% | 110.00% | 5.00% | 110.00% | |
| 2B1 | | 2.00% | 50.00% | 2.00% | 50.00% | Strong activity data, so low activity uncertainty. Assume a high uncertainty for the literature factor. |
| 2B2 | | 10.00% | 100.00% | (a) (r) | 10.00% (r) | Emission estimates for recent years have been based partially (1998-2008) or wholly (2009-2017) on continuous monitoring, and therefore will be subject to low uncertainty. The monitoring systems used at the 2 sites currently in operation are subject to an uncertainty of 5-10%. Uncertainty in earlier years is much higher due to more limited information |
| 2B3 | | 2.00% | 100.00% | 2.00% | 100.00% | |
| 2B8 | | 10.00% | 100.00% | 10.00% | 100.00% | |
| 2C | | 1.50% | 118.00% | 1.00% | 118.00% | |
| 2D | | 50.00% | 100.00% | 50.00% | 100.00% | |
| 2G | | 100.00% | 100.00% | 100.00% | 100.00% | |
| 3B | | (a) | 68.07% | (a) | 9.53% (r) | Based on separate monte carlo analysis for the agriculture model |
| 3D | | (a) | 53.28% | (a) | 11.16% (r) | Based on separate monte carlo analysis for the agriculture model |
| 3F | | 25.63% | (a) | 25.62% (r) | (a) | |
| 3J | | 50.00% (n) | 50.00% (n) | 50.00% (n) | 50.00% (n) | |
| 4A | non-fuel combustion | 1.00% | 95.00% (r) | 1.00% | 95.00% (r) | |
| 4B | non-fuel combustion | 1.00% | 55.00% | 1.00% | 35.00% | |
| 4C | non-fuel combustion | 1.00% | 115.00% (r) | 1.00% | 115.00% (r) | |
| 4D | non-fuel combustion | 1.00% | 100.00% | 1.00% | 100.00% | |
| 4E | non-fuel combustion | 1.00% | 20.00% | 1.00% | 15.00% | |
| 4only | non-fuel combustion | 1.00% | 165.00% | 1.00% | 165.00% | |

Uncertainties **A2**

| Category | Fuel | 1990 | | 2017 | | Justification for key sources |
|----------|------|--------------------------|---------------------------------|--------------------------|---------------------------------|-------------------------------|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| 5B | | 30.00% | 90.00% | 30.00% | 90.00% | |
| 5C | | 7.00% | 230.00% | 7.00% | 230.00% | |
| 5D | | 10.00% | 248.00% | 10.00% | 248.00% | |

Table A 2.3.4 Estimated uncertainties in the activity data and emission factors used in the F-gas inventory

| Gas | Category | 1990 | | 2017 | | Justification for key sources |
|-----------------|----------|--------------------------|---------------------------------|--------------------------|---------------------------------|--|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| HFCs | 2B9 | (a) | 10.00% | (a) | 10.00% | |
| HFCs | 2C4 | 5.00% | 10.00% | 5.00% | 10.00% | |
| HFCs | 2E1 | (a) | 47.15% | (a) | 47.15% | |
| HFCs | 2F1 | 10.00% | 10.00% | 10.00% | 10.00% | Good UK data on refrigerant supply is used to tune the model of emissions for this sector, which means that there is a high confidence in the overall estimates of a activity for this sector. Good activity data helps mitigate the uncertainty in emissions, as leakage and disposal is directly linked to refrigerant demand. |
| HFCs | 2F2 | (a) | 15.00% | (a) | 15.00% | |
| HFCs | 2F3 | (a) | 25.00% | (a) | 25.00% | |
| HFCs | 2F4a | 5.00% | 10.00% | 5.00% | 10.00% | |
| HFCs | 2F4b | (a) | 10.00% | (a) | 10.00% | |
| HFCs | 2F5 | (a) | 25.50% | (a) | 25.50% | |
| HFCs | 2F6 | (a) | 20.00% | (a) | 20.00% | |
| NF ₃ | 2E1 | (a) | 47.15% | (a) | 47.15% | |

Uncertainties **A2**

| Gas | Category | 1990 | | 2017 | | Justification for key sources |
|-----------------|----------|--------------------------|---------------------------------|--------------------------|---------------------------------|-------------------------------|
| | | Activity uncertainty (%) | Emission factor uncertainty (%) | Activity uncertainty (%) | Emission factor uncertainty (%) | |
| PFCs | 2B9 | (a) | 10.00% | (a) | 10.00% | |
| PFCs | 2C3 | (a) | 20.00% | (a) | 20.00% | |
| PFCs | 2F1 | 10.00% | 10.00% | 10.00% | 10.00% | |
| PFCs | 2F3 | (a) | 25.00% | (a) | 25.00% | |
| PFCs | 2G2e | (a) | 47.15% | (a) | 47.15% | |
| SF ₆ | 2C4 | 5.00% | 5.00% | 5.00% | 5.00% | |
| SF ₆ | 2G1 | (a) | 5.00% | (a) | 5.00% | |
| SF ₆ | 2G2a | (a) | 17.50% | (a) | 17.50% | |
| SF ₆ | 2G2b | (a) | 40.00% | (a) | 40.00% | |
| SF ₆ | 2G2e | (a) | 47.15% | (a) | 47.15% | |

A 2.3.2 General Considerations

The uncertainty parameters presented in above are based primarily on expert judgment, but where applicable will account for:

- The uncertainty range presented for data (for example the confidence interval in the 2006 IPCC guidelines for default factors)
- Monte Carlo Analysis of some of the more sophisticated models, most notably for agriculture, LULUCF and F-gases

In some cases, the individual uncertainties for the activity data and the emission factor are difficult to separate, but the uncertainty on the total emission can more easily be estimated. In these cases, the uncertainties are listed in the column marked “uncertainty in emission”.

The analysis of the uncertainties in the nitrous oxide emissions is particularly difficult because emissions sources are diverse, and few data are available to form an assessment of the uncertainties in each source. Emission factor data for the combustion sources are scarce and for some fuels are not available. The uncertainty assumed for agricultural soils (IPCC category 3D) uses a custom distribution. These parameterised functions have been defined and provided by Rothamsted Research as the best possible fit to the expected distribution of uncertainties in 1990 and the most recent year’s emissions, and are normalised in the Approach 2 methodology such that the resultant mean is consistent with the current inventory emissions in 1990 and the most recent year.

Many of the uncertainties in the emissions of HFCs, PFCs, NF₃ and SF₆ (collectively known as F-gases) are based on the recent study to update emissions and projections of F-gases (ICF, 2014). Some sources have been updated since then and the uncertainties for those sources have been revisited accordingly.

We assume that all F-gas emissions are independent between years as the technologies, gases (which have a very wide range of GWPs) used and regulations have changed drastically between the base year and the most recent year. Many HFCs in particular were not in use until the early 90s.

A 2.3.3 Uncertainty in the Trend

The uncertainty in the trend between 1990 and the most recent year is given in **Section A 2.4.2**. In running this simulation, it was necessary to make assumptions about the degree of correlation between sources and between 1990 and the most recent year. The assumptions were as follows:

- Activity data are uncorrelated;
- Emission factors of some similar fuels are correlated;
- Land Use Change and forestry emissions are correlated (e.g. 1990 4A CO₂ with 4A CO₂ for the most recent year);
- Offshore emissions are not correlated since they are based on separate studies using emission factors appropriate for the time;
- Emission factors covered by the Carbon Factors Review (Baggott et al, 2004) are not correlated;
- Process emissions from blast furnaces, coke ovens and ammonia plants are not correlated;

- Landfill emissions were partly correlated across years in the simulation. It is likely that the emission factors used in the model will be correlated, and also the historical estimates of waste arisings will be correlated since they are estimated by extrapolation from the year of the study. However, the reduction in emissions is due to flaring and utilisation systems installed since 1990 and this is unlikely to be correlated. As a simple estimate it was assumed that the degree of correlation should reflect the reduction in emissions since 1990;
- Emissions from agricultural soils and manure management are correlated in the base and inventory year;
- The emission factor used for sewage treatment was assumed to be correlated between years, though the protein consumption data used as activity data were assumed not to be correlated between years; and,
- Nitric acid production emission factors were assumed not to be correlated, since the mix of operating plants is very different in the most recent year compared with 1990 – only two of the original eight units are still operating in the latest inventory year, all of which now have differing levels of abatement fitted.

A 2.4 UNCERTAINTIES IN GWP WEIGHTED EMISSIONS

A 2.4.1 Uncertainty in the emissions

The uncertainty in the combined GWP weighted emission is given in **Table A 2.4.1**, along with uncertainties for each of the seven categorised GHGs. This is calculated as half of the 95% confidence range, i.e. the limits between which there is a 95% probability that the actual value of emissions falls. Note that the uncertainty in the GWP is not accounted for.

A 2.4.2 Uncertainty in the Trend

The uncertainty estimates for all gases are summarised in **Table A 2.4.1** under 'Range of likely % change'. This indicates the range between which there is a 95% probability that the actual trend in inventory emissions falls. Note that the uncertainty in the GWP is not accounted for.

Table A 2.4.1 Summary of Monte Carlo Uncertainty Estimates

| IPCC Source Category | Gas | 1990 Emissions | 2017 Emissions | 95% confidence interval for 1990 emissions | | Uncertainty in 1990 emissions as % of emissions in category | 95% confidence interval for 2017 emissions | | Uncertainty in 2017 emissions as % of emissions in category | % change in emissions between 1990 and 2017 | 95% confidence interval for the % change in emissions between 1990 and 2017 | |
|----------------------|-----------------------|----------------------|----------------------|--|----------------------|---|--|----------------------|---|---|---|-----------------|
| | | | | 2.5 percentile | 97.5 percentile | | 2.5 percentile | 97.5 percentile | | | 2.5 percentile | 97.5 percentile |
| | | Gg CO ₂ e | Gg CO ₂ e | Gg CO ₂ e | Gg CO ₂ e | % | Gg CO ₂ e | Gg CO ₂ e | % | % | % | % |
| TOTAL | CO ₂ (net) | 599,409 | 376,757 | 585,999 | 612,867 | 2.2% | 365,909 | 387,601 | 2.9% | -37% | -39% | -36% |
| | CH ₄ | 132,883 | 51,872 | 106,504 | 171,910 | 24.6% | 44,346 | 61,629 | 16.7% | -60% | -70% | -50% |
| | N ₂ O | 48,427 | 21,173 | 36,939 | 66,435 | 30.5% | 18,540 | 24,896 | 15.0% | -55% | -69% | -41% |
| | HFC | 14,405 | 14,194 | 12,257 | 16,535 | 14.8% | 12,890 | 15,484 | 9.1% | -1% | -17% | 19% |
| | PFC | 1,652 | 371 | 1,571 | 1,732 | 4.9% | 290 | 465 | 23.6% | -78% | -83% | -72% |
| | SF ₆ | 1,305 | 526 | 1,169 | 1,443 | 10.5% | 464 | 586 | 11.6% | -60% | -66% | -53% |
| | NF ₃ | 0.4 | 0.5 | 0.2 | 0.6 | 43.6% | 0.3 | 0.8 | 47.1% | 41% | -35% | 151% |
| | All | 798,081 | 464,893 | 764,043 | 842,085 | 4.9% | 450,970 | 479,682 | 3.1% | -42% | -45% | -39% |

Uncertainty calculated as $0.5 \cdot R/E$ where R is the difference between 2.5 and 97.5 percentiles and E is the mean calculated in the simulation.

Emissions of CO₂ are net emissions (i.e. sum of emissions and removals).

Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

A 2.5 **SECTORAL UNCERTAINTIES**

A 2.5.1 **Overview of the Method**

Sectoral uncertainties were calculated from the same base data used for the “by gas” analysis. The emissions and uncertainties per sector are presented in **Table A 2.5.1**. The estimates are presented in IPCC categories, which is consistent with the reporting format used within this submission to the UNFCCC, but we recommend that these estimates should only be considered as indicative.

Table A 2.5.1 Sectoral Uncertainty Estimates

| IPCC Source Category | 1990 Emissions | 2017 Emissions | 95% confidence interval for 2017 emissions | | Uncertainty in 2017 emissions as % of emissions in category | % change in emissions between 1990 and 2017 | 95% confidence interval for the % change in emissions between 1990 and 2017 | |
|----------------------|----------------|----------------|--|-----------------|---|---|---|-----------------|
| | | | 2.5 percentile | 97.5 percentile | | | 2.5 percentile | 97.5 percentile |
| 1A1a | 205,372 | 74,445 | 72,649 | 76,245 | 2.4% | -64% | -65% | -62% |
| 1A1b | 17,849 | 13,585 | 11,622 | 15,763 | 15.2% | -24% | -43% | 3% |
| 1A1c | 14,184 | 15,734 | 15,274 | 16,256 | 3.1% | 11% | 5% | 18% |
| 1A2a | 21,547 | 9,328 | 8,522 | 10,137 | 8.7% | -57% | -62% | -51% |
| 1A2b | 4,359 | 734 | 696 | 772 | 5.2% | -83% | -86% | -79% |
| 1A2c | 12,117 | 5,316 | 5,041 | 5,595 | 5.2% | -56% | -60% | -52% |
| 1A2d | 4,617 | 1,417 | 1,333 | 1,501 | 5.9% | -69% | -73% | -65% |
| 1A2e | 7,614 | 4,115 | 3,905 | 4,326 | 5.1% | -46% | -50% | -41% |
| 1A2f | 7,138 | 2,578 | 2,278 | 2,898 | 12.0% | -64% | -72% | -53% |
| 1A2g | 38,769 | 28,063 | 26,609 | 29,530 | 5.2% | -28% | -34% | -21% |
| 1A3a | 1,908 | 1,918 | 1,539 | 2,291 | 19.6% | 1% | -24% | 33% |
| 1A3b | 111,129 | 114,660 | 112,665 | 116,678 | 1.8% | 3% | 1% | 6% |
| 1A3c | 1,471 | 2,002 | 1,640 | 2,366 | 18.1% | 36% | 6% | 74% |
| 1A3d | 7,717 | 5,522 | 4,546 | 6,497 | 17.7% | -28% | -44% | -10% |
| 1A3e | 228 | 573 | 467 | 679 | 18.5% | 151% | 94% | 223% |
| 1A4a | 25,638 | 19,304 | 18,663 | 19,942 | 3.3% | -25% | -30% | -19% |
| 1A4b | 80,234 | 65,600 | 62,997 | 68,281 | 4.0% | -18% | -23% | -13% |

| IPCC Source Category | 1990 Emissions | 2017 Emissions | 95% confidence interval for 2017 emissions | | Uncertainty in 2017 emissions as % of emissions in category | % change in emissions between 1990 and 2017 | 95% confidence interval for the % change in emissions between 1990 and 2017 | |
|----------------------|----------------|----------------|--|-----------------|---|---|---|-----------------|
| | | | 2.5 percentile | 97.5 percentile | | | 2.5 percentile | 97.5 percentile |
| 1A4c | 6,309 | 5,300 | 3,737 | 6,858 | 29.4% | -16% | -48% | 48% |
| 1A5b | 5,355 | 1,576 | 1,455 | 1,697 | 7.7% | -71% | -74% | -67% |
| 1B1 | 23,531 | 848 | 756 | 941 | 10.9% | -96% | -97% | -96% |
| 1B2 | 18,166 | 9,197 | 6,178 | 12,242 | 33.0% | -49% | -65% | -33% |
| 2A1 | 7,295 | 4,410 | 4,269 | 4,550 | 3.2% | -40% | -40% | -39% |
| 2A2 | 1,462 | 1,052 | 1,000 | 1,105 | 5.0% | -28% | -35% | -20% |
| 2A3 | 406 | 368 | 349 | 387 | 5.1% | -9% | -10% | -9% |
| 2A4 | 672 | 426 | 410 | 442 | 3.7% | -37% | -40% | -33% |
| 2B1 | 1,895 | 1,764 | 1,727 | 1,802 | 2.1% | -7% | -11% | -3% |
| 2B2 | 3,861 | 37 | 17 | 71 | 72.6% | -99% | -100% | -97% |
| 2B3 | 19,974 | - | - | - | n/a | -100% | -100% | -100% |
| 2B6 | 105 | 181 | 163 | 199 | 10.0% | 73% | 40% | 120% |
| 2B7 | 232 | 140 | 132 | 148 | 5.6% | -39% | -45% | -33% |
| 2B8 | 4,568 | 2,859 | 1,997 | 3,739 | 30.4% | -37% | -58% | -9% |
| 2B9 | 14,418 | 175 | 158 | 193 | 9.8% | -99% | -99% | -99% |
| 2B10 | 175 | 49 | 37 | 61 | 24.4% | -72% | -80% | -60% |
| 2C | 9,395 | 2,716 | 2,582 | 2,847 | 4.9% | -71% | -73% | -69% |
| 2D | 554 | 518 | 344 | 761 | 40.3% | -6% | -52% | 111% |

| IPCC Source Category | 1990 Emissions | 2017 Emissions | 95% confidence interval for 2017 emissions | | Uncertainty in 2017 emissions as % of emissions in category | % change in emissions between 1990 and 2017 | 95% confidence interval for the % change in emissions between 1990 and 2017 | |
|----------------------|----------------|----------------|--|-----------------|---|---|---|-----------------|
| | | | 2.5 percentile | 97.5 percentile | | | 2.5 percentile | 97.5 percentile |
| 2E | 5 | 22 | 13 | 33 | 46.1% | 359% | 142% | 769% |
| 2F | 0 | 14,168 | 12,865 | 15,457 | 9.1% | 3964969% | 2628871% | 4753039% |
| 2G | 1,593 | 1,441 | 755 | 2,652 | 65.8% | -10% | -58% | 92% |
| 3A | 25,393 | 21,464 | 19,220 | 23,935 | 11.0% | -15% | -28% | -1% |
| 3B | 8,174 | 7,523 | 6,675 | 8,498 | 12.1% | -8% | -22% | 9% |
| 3D | 13,617 | 11,462 | 9,659 | 13,555 | 17.0% | -16% | -32% | 4% |
| 3F | 244 | - | - | - | n/a | -100% | -100% | -100% |
| 3G | 1,015 | 939 | 740 | 1,135 | 21.0% | -7% | -32% | 25% |
| 3H | 328 | 344 | 224 | 501 | 40.3% | 5% | -41% | 87% |
| 3J | 410 | 300 | 182 | 422 | 39.9% | -27% | -60% | 33% |
| 4 | 398 | 246 | 82 | 572 | 99.5% | -38% | -84% | 142% |
| 4A | -14,785 | -18,055 | -24,360 | -11,677 | 35.1% | 22% | 14% | 30% |
| 4B | 15,253 | 11,408 | 6,544 | 16,263 | 42.6% | -25% | -35% | -15% |
| 4C | -7,105 | -8,827 | -13,131 | -4,435 | 49.3% | 24% | 13% | 36% |
| 4D | 491 | 337 | 160 | 638 | 70.9% | -31% | -59% | -15% |
| 4E | 7,608 | 6,993 | 4,755 | 9,982 | 37.4% | -8% | -12% | -5% |
| 4F | - | - | - | - | n/a | n/a | n/a | n/a |
| 4G | -1,640 | -2,016 | -2,930 | -1,098 | 45.4% | 23% | 12% | 34% |

| IPCC Source Category | 1990 Emissions | 2017 Emissions | 95% confidence interval for 2017 emissions | | Uncertainty in 2017 emissions as % of emissions in category | % change in emissions between 1990 and 2017 | 95% confidence interval for the % change in emissions between 1990 and 2017 | |
|----------------------|----------------|----------------|--|-----------------|---|---|---|-----------------|
| | | | 2.5 percentile | 97.5 percentile | | | 2.5 percentile | 97.5 percentile |
| 5A | 60,302 | 14,227 | 8,133 | 23,049 | 52.4% | -76% | -88% | -55% |
| 5B | 31 | 1,922 | 1,188 | 3,025 | 47.8% | 6079% | 2931% | 12594% |
| 5C | 1,483 | 308 | 122 | 709 | 95.1% | -79% | -94% | -33% |
| 5D | 5,005 | 4,177 | 2,867 | 7,070 | 50.3% | -17% | -41% | 16% |
| Grand Total | 798,082 | 464,893 | 450,971 | 479,682 | 3.1% | -42% | -45% | -39% |

Note: Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

A 2.6 COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES

Comparing the results of the error propagation approach, and the Monte Carlo estimation of uncertainty by simulation, is a useful quality control check on the behaviour of the Monte Carlo model.

The reason that the error propagation approach is used as a reference is because the approach to the error propagation approach has been defined and checked by the IPCC, and is clearly set out in the IPCC 2000 Good Practice Guidance and the 2006 Guidelines. The UK has implemented the IPCC error propagation approach as set out in this guidance. The implementation of an uncertainty estimation by simulation cannot be prescriptive, and will depend on how the country constructs its model, and the correlations included within that model. Therefore, there is a greater likelihood of errors being introduced in the model used to estimate uncertainty by Monte Carlo simulation.

If all the distributions in the Monte Carlo model were normal, and the assumed correlations were identical, the estimated errors on the trend from the Monte Carlo model should approach those estimated by the error propagation approach if enough iterations are done. The error propagation approach assumes 100% correlation between EFs in the base and inventory year, and no correlation between sources, however in reality the nature and degree of correlation varies by source, and many distributions are not normal but heavily skewed, particularly those with very high uncertainty. These differences interact in various ways, but would be expected broadly to result in higher trend uncertainty, and lower uncertainty on the most recent year's total in the Monte Carlo uncertainty estimates compared to the error propagation approach. This can be seen in **Table A 2.6.1** which shows differences in the trend uncertainty between the error propagation and Monte Carlo approaches. These differences mostly arise from the fact that the error propagation approach only uses normal distributions, cannot account for different uncertainty parameters between the 1990 and the latest inventory year, cannot account for correlations between sources, and automatically assumes a correlation between the emission factor uncertainty in 1990 and the most recent year.

The central estimates of emissions generated by the Monte Carlo model in 1990, and those in the latest inventory year, are very close. We would not expect the central estimates from the two methods to be identical, but with a very large number of iterations we would expect the difference to tend to zero. It should be noted that the Approach 1 uncertainties base year is 1990 for N₂O, CH₄ and CO₂, but is 1995 for the F-gases; this differs from the Approach 2 uncertainties which uses 1990 emission for all gases for the starting year.

Table A 2.6.1 Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses

| Method of uncertainty estimation | Central estimate (Gg CO ₂ equivalent) ^b | | Uncertainty on trend, 95% CI (1990 / base year to 2017) ^a |
|----------------------------------|---|---------|--|
| | Base year | 2017 | |
| Error propagation | 801,849 | 464,453 | 2.3% |
| Monte Carlo | 801,711 | 464,893 | 2.8% |

Notes:

CI Confidence Interval

^a Calculated as half the difference between 2.5 and 97.5 percentiles, assuming a normal distribution is equal to ± 1.96 standard deviations on the central estimate.

^b Net emissions, including emissions and removals from LULUCF

ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities.

This Annex contains background information about methods used to estimate emissions in the UK GHG inventory. This information has not been incorporated in the main body of the report because of the level of detail, and because the methods used to estimate emissions cut across sectors.

This Annex provides background information on the fuels used in the UK GHG inventory, mapping between IPCC and NAEI source categories and detailed description of methods used to estimate GHG emissions, and emission factors used in those methods.

A 3.1 ENERGY

Methods for calculating emissions within the energy sector are detailed in the method statements set out in **Chapter 3**. This Annex details the emission factors used and their source, and elaborates on references commonly used within the Energy sector. The national energy balance (and how it is used) is described in **Annex 4**.

A 3.1.1 Emission factors

Emission factors used for the 2019 submission for sectors 1A and 1B can be found in the accompanying excel file: 'Energy_background_data_uk_2019.xlsx'. This can be found as one of the additional documents in on http://naei.beis.gov.uk/reports/reports?report_id=954. Note that there can be a delay between the NIR being published on the NAEI website after official submission.

A 3.1.2 Commonly used references

This section describes data sources that are used across multiple emission sources within the energy sector, and how they are used.

A 3.1.2.1 Baggott et al., 2004 – Carbon factors review

A review of the carbon factors used in the UK GHG inventory was carried out in 2004. The report detailing this study is available from:

http://naei.beis.gov.uk/reports/reports?report_id=417

This aimed to validate existing emission factors and seek new data for country specific emission factors for the UK. At the time of publication this reference provided new emission factors for:

- coal from power stations;
- fuels used in the cement industry;

- a number of petroleum based fuels;
 - natural gas; and
 - coke oven and blast furnace gas.

Since then following updates are made to the following emission factors based on new information:

1. Coal emission factors are adjusted based on the annual variations in the GCV of the fuels using methods developed as part of the 2004 analysis (Baggott et al., 2004).

$$EF_y = EF_{ref} / GCV_{ref} * GCV_y$$

Where

EF_y is the emission factor (EF) in year y

EF_{ref} is the EF in the reference year (the year for which data are available)

GCV_{ref} is the GCV in the reference year

GCV_y is the GCV in year y

2. Since the advent of EU ETS in 2005, a number of sources of emissions from coal which had previously been reliant on Baggott et al., 2004 have now been replaced with data from the ETS, where the data set was considered suitable (high proportion of source included, and high proportion of T3 plant specific data). In addition, in 2014 the use of oxidation factors from this report was reviewed, and where suitable background evidence to support the factors used were not available, the IPCC default (of 1, IPCC 2006) has been used.
3. Emission factors for petroleum based fuels (where ETS data are not available) are still largely based on Baggott et al., 2004. These were reviewed in 2014 and compared with the defaults in the 2006 IPCC Guidelines and found to be largely within the range of the 2006 Guidelines. No new data for the UK has been identified and the emission factors from Baggott et al., 2004 are still considered to be relevant country specific emission factors.
4. During 2017-18, a review of the UK's shipping inventory was conducted (Scarborough et al., 2018). This identified new carbon emission factors for marine fuels, which replace the factors identified as part of Baggott et al., 2004.
5. Emission factors for natural gas are updated annually based on analyses from the gas network operators (Personal Communications from network operators, 2015).
6. Emission factors for coke oven gas and blast furnace gas are estimated based on a carbon balance approach (as described in Chapter 3, **MS 4**).
7. The Mineral Products Association provide data for fuels used in the cement industry annually on a confidential basis, and these are validated with EU ETS data (Personal Communication, MPA, 2018).

A new review of carbon emission factors was conducted during 2017, focusing on those factors retained from the 2004 review (Brown et al., 2017). This concluded that the factors that are currently in use are slightly more conservative than more recent values identified, and that there was no new robust evidence upon which we could justify changing the current factors. This report is available here: http://naei.beis.gov.uk/reports/reports?report_id=947

A 3.1.2.2 The Pollution Inventory and other regulators' inventories

The Pollution Inventory (PI) has, since 1998, provided emission data for the six Kyoto gases and other air pollutant for installations regulated by the Environment Agency (EA) in England and Natural Resources Wales (NRW) in Wales. This is part of the UK's process for managing regulated emissions from industry processes under the IPPC permitting system. The PI does contain earlier data of carbon dioxide emissions at some sites reported from 1994 onwards. The Scottish Pollutant Release Inventory (SPRI) covers processes regulated by the Scottish Environment Protection Agency (SEPA), and contains data from 2002 and 2004 onwards. The Northern Ireland Pollution Inventory (NIPI) covers processes regulated by the Northern Ireland Environment Agency and includes data for 1999 onwards.

These data are subject to some very significant limitations:

- Emissions of each pollutant are reported for each permitted installation as a whole, so emissions data for carbon dioxide, for example, can cover emissions from fuel use as well as from an industrial process. No information is given on what the source of emissions is, so a judgement has to be made about the scope of reporting;
- Permitting arrangements have changed over time, so the reporting of data is not on a consistent basis across the time-series. In general, the tendency has been to reduce the number of permits, so that whereas in the early 1990s there might have been separate permits at an industrial installation covering the boiler plant and the chemical processes, from the late 1990s onwards the tendency would be to issue a single permit to cover both. Therefore, the problems with the scope of emissions data mentioned in the first bullet point are most severe for the second half of the GHGI time series; and,
- Since 1998, process operators need only report emissions of each pollutant if those emissions exceed a reporting threshold. For example, where emissions from an installation are less than 10,000 tonnes of CO₂, or 10 tonnes of methane, the operator does not need to report any emissions data for that substance in that year. Reporting thresholds are irrelevant for many of the sectors of interest to this study, since emissions would be many times higher than the thresholds, but the reporting thresholds do mean that it is necessary to consider whether the data available in the PI (and in the SPRI & NIPI for later years) will be complete.

Despite these limitations, these data are still a useful source of information for the UK GHG inventory. A considerable amount of effort is put into manually interpreting the individual returns and allocating these to appropriate categories for use in the inventory estimates by the Inventory Agency.

A 3.1.2.3 The Environmental and Emissions Monitoring System (EEMS) Reporting System

Emissions from upstream oil and gas production facilities, including onshore terminals, are estimated based on operator reporting via EEMS, regulated by the BEIS Offshore Inspectorate and developed in conjunction with the trade association Oil & Gas UK (formerly the UK Offshore Operators' Association, UKOOA). The EEMS data provides a detailed inventory of point source emissions estimates, based on operator returns for the years 1995-2017. However, the EEMS data for 1995 to 1997 are not complete, frequently exhibiting duplicate entries with identical submissions by operators across years. Since the 1995 – 1997 data are not considered reliable, the EEMS dataset is only used directly to inform national inventory estimates from 1998 onwards for the following sources:

- gas flaring;
- own gas combustion;
- well testing; and
- oil loading (onshore and offshore).

[Activity data are not routinely collected via EEMS for sources including: fugitive releases, direct process activities, oil storage or gas venting. The emissions from these sources are reported as annual estimates by operators and used directly within the inventory.]

These EEMS-derived activity data enable greater analysis of the oil & gas emissions and related emission factors at the installation level, providing a high degree of data transparency and improving the level of detail for performing quality checks by source, by site, by year. For those sources, this has led to an improvement in data transparency and easier query of Implied Emission Factor trends. However, the EEMS activity data are only available back to 1998, and hence the activity data back to 1990 are extrapolated using the oil and gas production time-series that were collected at that time for the purposes of energy data reporting.

A 3.1.2.4 Fynes & Sage (1994)

Fynes and Sage is a country-specific reference from the mid-1990s and it includes analysis of solid fuels typically used in the UK economy in that period, deriving mass-based emission factors that are used within the UK GHGI. In the 1990s, coal used in the UK economy was predominantly mined in the UK, whereas over the time series of the inventory there has been a decline in the share of coal from UK sources and an increase in coal imports from around the world.

For recent years, for the more significant emission sources, e.g. energy industries and manufacturing industries, the inventory agency uses EFs that are derived from EU ETS data, but for smaller emission sources in the UK that still use solid fuels (such as residential, collieries) the Fynes and Sage data are retained, as there are no EU ETS data for fuels used in these sectors. There is some uncertainty regarding how representative the EFs from Fynes and Sage may be for these smaller combustion sources, but we note that the use of coal-fired technology in sectors such as collieries and residential is predominantly in the UK coal production areas, where local supplies are still available.

A 3.1.3 Feedstocks and Non-Energy Use (NEU) of fuels

The estimation methods are described within individual sections of the NIR, but are summarised here. The general approach adopted in the UK GHG inventory is to assume that emissions from all non-energy uses of fuels are zero (i.e. the carbon is assumed to be sequestered in products such as plastics and other chemicals), except for cases where emission sources can be identified and emission estimates included in the inventory. There is one exception to this, for petroleum coke where we have no information on any non-emissive uses at all, and so we adopt the conservative approach of assuming that all petroleum coke use is emissive.

The UK Inventory Agency conducts periodic studies into the fate of fuels reported as non-energy use, in order to assess the levels of stored carbon and carbon emitted for different fuels over the time series. These detailed studies are supplemented through annual data gathering and consultation with stakeholders to maintain an accurate representation of the emitted and stored carbon in the inventory.

The assumptions and estimates for individual sources are based on a review conducted in 2013-14 (Ricardo-AEA, 2014b) which included research into UK-specific activities and data sources as well as a review of the National Inventory Reports (NIRs) of other countries.

The sections below outline the emission sources from feedstock and NEU of fuels that are included in the UK GHGI, the source data and estimation methods and a summary of the time series for each of the fuel types where there is a stored carbon component in the UK energy balance. The estimates are all presented in CRF Tables 1.Ab and 1.Ad.

Table A 3.1.1 Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics

| Fuel | IPCC | Source Category |
|--|------|---|
| <i>Liquid Fossil</i> | | |
| Naphtha, Liquid Petroleum Gases (LPG), Refinery Fuel Gas (RFG) / Other Petroleum Gases (OPG), gas oil and Ethane | 1A1a | Scrap tyre combustion in power stations (1994 to 2000 only). Fossil carbon in MSW combustion in energy from waste plant. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i> |
| | 1A1b | Other petroleum gas use in refineries (2004, 2006 to 2011, 2013 to 2017 only). <i>Re-allocated from non-energy use as EU ETS and trade association data indicates that DUKES data on OPG combustion are an under-report.</i> |
| | 1A2f | Waste solvents, waste-derived fuels containing fossil carbon, in cement kilns. Scrap tyres and waste plastics etc. combusted in cement kilns. |
| | 1A2g | Industrial combustion of waste solvents. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and solvents.</i> |
| | 2B8 | Energy recovery from process off-gases in the chemical industry. <i>Large quantities of naphtha, butane, propane, ethane, and other petroleum gases are listed in DUKES as used for non-energy applications and these fuels are known to be used extensively as chemical feedstocks. However, EU ETS and operator data indicate that process off-gases, derived from the chemical feedstocks, are a major fuel for ethylene production processes and other petrochemical sites. Emissions of carbon are reported in 2B8.</i> |
| | 5C | Fossil carbon in chemical waste incineration. Fossil carbon in MSW incineration. Fossil carbon in clinical waste incineration. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i> |

| Fuel | IPCC | Source Category |
|----------------------------------|---|--|
| Lubricants | 1A1a | Waste oil combustion in power stations. |
| | 1A2f | Waste oil combustion in cement kilns. |
| | 1A2g | Waste oil combustion in unclassified industry (including road-stone coating plant) |
| | 1A3biv | Lubricant combustion in moped engines |
| | 2D1 | Lubricant combustion in aircraft, industrial, road vehicle (except moped), marine shipping and agricultural engines. |
| | 5C | Incineration of waste oil. |
| Bitumen | n/a | <i>No known UK applications that lead to GHG emissions.</i> |
| Petroleum coke | 1A2f 1A2g 1A4b 2A4 2B6 2C1 2C3 2D4 | <p>Based on reported energy use data by specific industries within datasets such as EU ETS and also from direct dialogue with industry representatives, the Inventory Agency re-allocates a small proportion of the reported "NEU" allocation from DUKES, and reports emissions within the UK GHG inventory. This re-allocation generates emissions for the mineral processing sector (1A2f) and other industry (1A2g) and for petcoke use in the domestic sector (1A4b).</p> <p>There are also non-combustion, emissive uses of petcoke in the UK through the use of petcoke-derived anodes in the metal processing industries. Emissions from these uses of petcoke are reported in 2C1 (electrode use in electric arc furnaces) and 2C3 (anode use in aluminium manufacture). Petroleum coke is also used in the minerals (2A4) and chemicals industries (2B6) leading to further emissions. The remaining consumption of petroleum coke is also assumed to be emissive, with emissions reported under 2D4.</p> <p><i>Note that DUKES already includes allocations of petcoke use as a fuel in combustion in power stations (1A1a) and refineries (1A1b), which are included in the UK GHG inventory.</i></p> |
| Other Oil | 2D2 | Carbon released from use of petroleum waxes. Uses of petroleum waxes includes candles, with carbon emitted during use. |
| <i>Solid Fossil</i> | | |
| Coking coal (coal oils and tars) | n/a | <i>Unknown quantities of coal tar pitch are used in the manufacture of anodes for industrial processes. In the UK inventory the emissions from the use of these anodes are allocated only against petroleum coke (also used in anode production). This is a small mis-allocation of emissions between the two fuels since the carbon emitted is likely to arise from both petroleum coke and the coal tar pitch, but it is due to lack of detailed data, and does not affect the accuracy of UK inventory emissions.</i> |
| <i>Gaseous Fossil</i> | | |
| Natural Gas | 2B1 2B8 | Ammonia and methanol production leading to direct release of CO ₂ from natural gas used to provide the energy for steam reforming and from natural gas feedstock to the reformer. Carbon originating in the natural gas feedstock which is converted into methanol is assumed stored, however. |

A 3.1.3.1 Naphtha, Ethane, Gas Oil, Refinery/Other Fuel Gas (RFG/OPG) Propane and Butane (LPG)

Ethane, LPG (given separately as propane & butane in the energy statistics), gas oil, refinery / other fuel gas (RFG/OPG) and naphtha are all consumed in very significant quantities for non-energy uses, primarily as feedstock in chemical manufacturing. In the UK, several major petrochemical production facilities are supplied with Natural Gas Liquid (NGL) feedstock directly

from upstream production pipelines, and then utilise NGL fractions such as ethane, propane and butane in their manufacturing processes. In addition, several integrated refinery / petrochemical complexes in the UK use a proportion of the refinery fuel gas as a feedstock in petrochemical production.

The NEU allocations presented in DUKES reflect the reported disposals of these commodities as feedstocks to chemical and petrochemical companies. There are several sources of GHG emissions from this stock of “NEU” feedstock carbon, although a high proportion of carbon is stored into products and not emitted.

One large emission source known to occur in the UK is the use of carbon-containing process off-gases as a fuel within the chemical facilities. Whilst the exact source of the carbon cannot be traced directly to a specific feedstock commodity within the UK sectoral approach, the available information from EU ETS and from consultation with operators enables the Inventory Agency to derive estimates of the GHG emissions across the time series from this emission source.

The majority of emissions are from installations manufacturing ethylene, but a number of other chemical sites report additional emissions in the EU ETS that can be attributed to the combustion of process off-gases and residues derived from the chemical feedstock. As a result, the UK inventory emissions in 2B8 now include estimates of emissions from use of process off-gases and residues at 5 ethylene manufacturing installations and 16 other chemical manufacturing installations in the UK (some of these 16 may be using process off-gases from neighbouring ethylene plant). The derivation of a time series of emission estimates from these sources is based as far as possible on reported data by plant operators within trading scheme data and other regulatory reporting mechanisms. For the early part of the time series, data on changes in plant capacity over time is used to derive the best estimates of activity and emissions by extrapolation back from later emission estimates, whilst for later years the completeness and transparency of operator reporting is greater. Therefore, whilst the uncertainty for the emission estimates in the early part of the time series is significantly greater than for those in recent years, the Inventory Agency has made best use of the available data to derive the time series estimates of emissions from “NEU” activity. Consultation with a sector trade association has also confirmed that there are no other sector estimates of this activity, or of production data across the time series, that could be used to further improve the time series (Personal communication: Chemical Industries Association, 2014).

Other emissions included within the UK GHG inventory include emissions from the destruction of chemical products, e.g. when wastes are incinerated or used as fuels. Although emissions from incineration and combustion of wastes are estimated, we cannot relate the carbon in these wastes back to individual feedstock, so it is not possible to generate reliable UK estimates of the proportion of carbon that is ultimately emitted from each individual fuel. Incineration of wastes derived from chemical feedstocks will be reported in 1A1a (in the case of plastics etc. in municipal waste incinerated with energy recovery) and in 6C (in the case of chemical, clinical and municipal wastes incinerated without energy recovery). Waste-derived fuels, including waste solvents, waste plastics and scrap tyres are used as fuels in cement kilns and other industrial plants, and emissions reported in 1A2. Tyres contain a mixture of natural and synthetic rubbers, and so where waste tyres are used as a fuel, the emission estimates take into account that only some of the carbon emitted is derived from fossil fuels.

Some propane / butane mixtures are used as a propellant in aerosols and are emitted as VOC. The UK inventory contains estimates of these VOC emissions, combined with emissions of solvents used in aerosols.

We assume that all gas oil used for non-energy purposes is used as a feedstock material, and consultation with DECC (now BEIS) energy statisticians supports this (Personal communication: Will Spry, DECC Energy Statistics team, 2014). A possible alternative use would be in explosives, but consultation with the Health and Safety Executive, who regulate the UK explosives industry, has confirmed that no UK installations manufacture explosives using gas oil or fuel oil as a feedstock (HSE, 2013).

A 3.1.3.2 Lubricants

Lubricants are listed separately in the UK energy statistics and are used in vehicles and in machinery. The inventory includes estimates of emissions of carbon due to oxidation of lubricants during use, and also includes estimates of emissions from the combustion of waste lubricants and other oils used as fuel.

UK GHG inventory estimates of the quantities of lubricants burnt as fuels are based on data from Recycling Advisory Unit, 1999; BLF/UKPIA/CORA, 1994; Oakdene Hollins Ltd, 2001 & ERM, 2008, as well as recent research to access information regarding the UK market for waste oils and the impact of European Directives to consolidate industrial emission regulations such as the Waste Incineration Directive (Oil Recycling Association, 2010). Estimates of waste oil combustion are derived for the following source categories:

- 1A1a Power stations;
- 1A2f Cement kilns; and
- 1A2f Other (unclassified) industry.

The estimated emissions for other industry assume that waste oils are used by two sectors: road-stone coating plant and garages. Other sectors may use waste oils as a fuel or as a reductant, but research to date provides no compelling evidence that there is a significant gap in the UK inventory for waste oil use by industrial operators.

The emission trends from power station use of waste lubricants reflect the fact that the Waste Incineration Directive (WID) had a profound impact on the market for waste oil, used as a fuel. It is assumed that no waste oil was burnt in power stations for the years 2006-2008, on the basis that the classification of waste oil as a fuel would have led to users being subject to the requirements of WID. In 2009 a Quality Protocol⁴ was introduced that allowed compliant fuel produced from waste oils to be burned as non-waste and this has encouraged a resumption in the consumption of waste oil-derived fuels from 2009 onwards.

Carbon dioxide emission estimates for the oxidation of lubricants within vehicle engines and machinery, and the use of waste oils for energy are all based on a single carbon emission factor derived from analysis of the elemental composition of a series of UK-sourced samples of waste oil (Passant, 2004). The UK inventory adopts the IPCC Tier 1 methodology for lubricant use i.e. assuming that 20% of all lubricants are oxidized during use. This assumption is used for the various sub-categories of lubricant use (including road, rail, marine, off-road and air transport) given in DUKES.

A 3.1.3.3 Bitumen

In the UK, bitumen is used only for applications where the carbon is stored. By far the most important of these is the use of bitumen in road dressings. The inventory does assume that a

⁴ <http://webarchive.nationalarchives.gov.uk/20140328084622/http://www.environment-agency.gov.uk/business/topics/waste/116133.aspx>

very small proportion of the carbon in the bitumen itself is emitted as VOC during road-stone coating but does not include any estimates of direct carbon emissions from uses of bitumen. Industry consultation in 2013 (UK Petroleum Industries Association, 2013; Refined Bitumen Association, 2013) has confirmed that there are no emissive applications of bitumen in the UK. Around 85% of bitumen is used in road paving, with the remaining proportion used almost entirely in the manufacture of weather-proofing materials.

A 3.1.3.4 Coal Oils and Tars

Coal-tars and benzole are by-products of coke ovens. Consultation with the operators of coal ovens (Tata, 2013) and also the UK company that refines and processes coal tars and benzole (Koppers UK, 2013) has confirmed that all of these materials are collected, refined and processed into a range of products that are not used as fuels. The carbon within coal tars and oils are entirely used within chemical processes. In some cases, the carbon is processed into anodes used in the ferrous and non-ferrous metals industries and then used (in the UK and overseas) within emissive applications. The UK inventory already includes estimates of emissions from UK consumption of carbon anodes within these industries, using methods based on UK metal production statistics.

Based on the evidence from process operators, the Inventory Agency allocates all of the reported coal tars and oils to Non Energy Use, i.e. assuming that all carbon is stored and there are no GHG emissions from this source-activity. The Digest of UK Energy Statistics (BEIS, 2018) also report the use of tars and benzole entirely to Non Energy Use.

Coal-tar pitch is used in the manufacture of electrodes, together with petroleum coke and a proportion of the carbon ultimately emitted, but details of input materials are scarce; emissions of carbon from these sources are included in the inventory attributed to petroleum coke. This may introduce a small mis-allocation of emissions between petroleum coke and coal oils and tars, but does not affect the UK inventory emissions total.

A 3.1.3.5 Natural Gas

Natural gas is used as a chemical feedstock for the manufacture of ammonia and formerly for methanol as well, though production of the latter ceased in 2001. Emissions occur directly as a result of a) combustion of gas used to power the steam reforming process that is required for manufacture of both ammonia and methanol; b) oxidation of gas in the steam reforming, producing CO₂ which in the case of ammonia production is not needed and is instead emitted. The emissions are reported under 2B1 for ammonia and 2B8 for methanol.

Most of the emissions from feedstock use of natural gas in ammonia production are at source, i.e. waste gases containing carbon are emitted directly from the ammonia plant. Up until 2001, some was exported to a neighbouring methanol plant and here converted into methanol, and this CO₂ is treated as stored. Further CO₂ is captured and sold for use elsewhere, for example, in carbonated drinks and this CO₂ is assumed all to be emitted in the UK.

A 3.1.3.6 Other Oil (industrial spirit, white spirit, petroleum wax, miscellaneous products)

White Spirit and Special Boiling Point (SBP) spirits are used exclusively for non-energy applications, and are listed in CRF Table 1.A(d) within the category 'other oil'. They are used as solvents; SBP spirits are used for industrial applications where quick drying times are needed (e.g. adhesives and other coatings) while white spirit is used as a solvent for decorative paint, as a cleaning solvent and for other applications. Estimates of VOC emissions are included in the

UK inventory but no estimates are made of direct emissions of carbon from these products, as they are regarded as “not occurring”.

The only emissions from this group of petroleum feedstock that are included in the UK GHG inventory are the releases of carbon from petroleum waxes which are reported under 2D2. These are accounted for in the UK inventory under the fuel category “Other Oils” in CRF Table 1Ad.

A 3.1.3.7 Petroleum Coke

The evidence from industrial reporting of fuel use and from periodic surveys of fuel producers that use petroleum coke to produce domestic fuels (including smokeless fuels) indicates that the allocation of petroleum coke to combustion activities in the UK energy balance is an under-estimate across all years. Therefore, the Inventory Agency generates revised estimates for all combustion activities and effectively re-allocates some of the petroleum coke reported in DUKES as non-energy use to energy-related emission sources in the UK inventory.

Within the UK inventory, petroleum coke is included for the following energy and non-energy source categories:

- 1A1a: Power station use of petroleum coke, primarily within blends with coal at a small number of UK facilities;
- 1A1b: Refinery emissions from regeneration of catalysts;
- 1A2f: Cement industry use of petroleum coke as a fuel;
- 1A2g: Other industry use of petroleum coke as a fuel;
- 1A4b: Petroleum coke use within domestic fuels;
- 2A4: Use in brick manufacture (reported combined with other emissions e.g. from use of carbonate minerals in brickmaking);
- 2B6: Use in chemicals manufacturing;
- 2C1: Carbon emissions from electrodes used in electric arc furnaces and ladle arc furnaces and petroleum coke added to furnaces as a carbon source;
- 2C3: Carbon emissions from anode use in primary aluminium production; and
- 2D4: Petroleum coke used for non-energy applications not included elsewhere.

The UK energy balance tables in DUKES contain data on the energy use in power stations (1A1a) and refineries (1A1b), although the former are only available for 2007 onwards, and both sets of data do not always agree with the available activity data from EU ETS. The remaining energy uses in industrial combustion (1A2f, 1A2g) and the domestic sector (1A4b) are not included in DUKES. The UK Inventory Agency therefore makes independent estimates of the consumption of petroleum coke in all of these sectors.

Petroleum coke is burnt in **cement kilns** (1A2f) and has been burnt in some years at a handful of **power stations** (1A1a). A few other **large industrial sites** (1A2g) have also used the fuel. Good estimates of the consumption of petroleum coke by these large sites are available from the operators themselves, from trade associations and from EU ETS data (from 2005 onwards).

Fuel grade petroleum coke is also used as a **domestic fuel** (both smokeless and non-smokeless types, reported in 1A4b). The Inventory Agency uses data supplied by the UK fuel supply industry to estimate petroleum coke consumption for domestic fuels over the period 1990 to 2016; these estimates are broadly consistent with fuel use data published in earlier editions of DUKES for a few years in the late 1990s.

Carbon deposits build up with time on catalysts used in **refinery** processes such as catalytic cracking. These deposits need to be burnt off to regenerate the surface area of the catalyst and ensure continued effectiveness of the catalyst; emissions from this process are reported within EU ETS since 2005, with the time series estimates provided by the trade association (UKPIA, 2018) and the catalyst regeneration is treated in the inventory as use of a fuel (since heat from the process is used) and are reported under 1A1b.

Estimates of carbon released from electrodes and anodes during **metal processes** are estimated based on operator data and reported in 2C1 and 2C3. Petroleum coke content of these electrodes and anodes is estimated based on operator data and literature sources such as Best available techniques REfERENCE documents (BREF notes). EU ETS data also show that some petroleum coke is added to electric arc furnaces as a carbon source, and the emissions from this use are also reported in 2C1. EU ETS data are also used for emission estimates for brickmaking, which include a component from petroleum coke. Finally, petroleum coke is used in the manufacture of titanium dioxide, with emission estimates generated from EU ETS and other operator data.

Based on data from DUKES we believe that there is some additional non-energy use of petroleum coke for most years; we assigned this residue to 2D4 and assume that it is all eventually emitted. The total fuel assigned to sector 2 is what we report as 'excluded carbon' in the CRF, table 1A(d). The consumption estimates for industrial users of petcoke as a fuel or in industrial processes are associated with low uncertainty as they are primarily based on operator reported data within the EU ETS or other regulatory reporting mechanisms. Whilst it is conceivable that other sectors may also use petroleum coke as a fuel, there is no evidence from resources such as EU ETS and Climate Change Agreement reporting that this is the case in the UK. The remaining petroleum coke consumption given in DUKES is therefore assumed to be used in various unidentified non-energy uses, all of which are assumed to be emissive. The estimates of petroleum coke used to generate fuels for the domestic sector are associated with higher uncertainty as they are based on periodic consultation with fuel suppliers to that market, and expert judgement of stakeholders.

As well as the total UK supply figure from UK energy statistics, DUKES has data on UK production, imports and exports of petroleum coke, which together provide more information on the nature of the UK consumption of petroleum coke. These data cover three distinct types of petroleum coke – catalyst coke, produced and consumed at refineries only (so no import/export or supply of fuel to other UK sectors), and then two products made in a refinery process known as coking: fuel grade (green) coke and anode-grade coke, with the former being used as a fuel, and the latter being a calcined⁵ version of the former, used in various non-energy processes. Consultation with the DECC energy statistics team and the only UK refinery with a coking process (DECC, 2013) has confirmed that the UK produces only anode-grade coke, and exports will also be anode-grade coke, whilst imports will be fuel grade coke for use as a cost-effective fuel source or raw material for production processes under NEU.

Carbon factors for petroleum coke use are derived from industry-specific data (including EU ETS fuel analysis) in the case of cement kilns (MPA, 2018), power stations and other industrial sites (EA, 2018; SEPA, 2018). The petroleum coke factor for refinery consumption is based on trade association analysis conducted as part of the 2004 Carbon Factors Review (UKPIA, 2004)

⁵ Calcined petroleum coke is a processed petroleum coke that has a very high carbon content; the resulting fuel is somewhat similar to coke oven coke

while the factor for domestic consumption is based on compositional analysis of samples of petroleum coke sold as domestic fuels (Loader et al, 2008).

These factors do show quite a large variation from sector to sector: this is probably primarily a reflection of the different requirements of fuels for different sectors (higher quality, higher carbon for some, less so for others). The highest carbon factor is for 'petroleum coke' burnt in sector 1A1b, but this fuel is actually of a different nature from the fuel burnt as petroleum coke in sectors 1A1a, 1A2f and 1A4b. In the case of 1A1b, the fuel is a build-up of carbon on catalysts used in various refinery process units, while in the other three cases, the petroleum coke is a solid by-product of a totally different refinery process (coking) which has different characteristics.

A 3.1.3.8 Carbon Storage Fractions: Import-Export balance for Carbon-containing Materials

The analysis within the UK energy statistics or GHG inventory compilation system cannot accurately account for the variable (over time) import-export balance of carbon-containing materials in the UK economy. For example, where the Inventory Agency accounts for the carbon emissions from scrap tyres burned in cement kilns, power stations, incinerators and so on within the inventory estimates or from the incineration of plastics or synthetic fibres, there is no way of tracing the quantity that is derived from imported tyres/plastics/fibres.

The reported estimate of the fate of the reported NEU of fuels from the UK energy balance is based on an assumed "closed system", whereby we account for all emissions from carbon-containing products and fuel types that are allocated as NEU as if they are derived from the fuel statistics in the UK energy balance. In reality, the source of the carbon emitted from feedstock and NEU of fuels will partly be carbon from imported materials, with UK feedstock carbon also exported and emitted elsewhere.

A 3.1.4 Aviation (MS 7)

Table A 3.1.2 CAA aircraft types assigned to EMEP-EEA Emissions Inventory Guidebook aircraft types

| EMEP/EEA Aircraft Type | CAA Aircraft Types |
|------------------------|--|
| Airbus A300 – B4 | AIRBUS A300 (ALL FREIGHTER); AIRBUS A300-600; AIRBUS A300-600F (ALL FREIGHTER); AIRBUS A300B1/B2; AIRBUS A300B4-100/200; AIRBUS A300F4 |
| Airbus A310 | AIRBUS A310; AIRBUS A310-202; AIRBUS A310-300 |
| Airbus A318 | AIRBUS A318 |
| Airbus A319 | AIRBUS A319; AIRBUS A319 CJ (EXEC); BOMBARDIER CSERIES CS100 |
| Airbus A320 | AIRBUS A320-100/200 |
| Airbus A321 | AIRBUS A321 |
| Airbus A330-200 | AIRBUS A330-200 |
| Airbus A330-300 | AIRBUS A330-300 |
| Airbus A340-200/300 | AIRBUS A340-200; AIRBUS A340-300; AIRBUS A350-900 |
| Airbus A340-500 | AIRBUS A340-500 |
| Airbus A340-600 | AIRBUS A340-600 |

| EMEP/EEA Aircraft Type | CAA Aircraft Types |
|---------------------------|--|
| Airbus A380-800 | AIRBUS A380-800 |
| Antonov 26 | ANTONOV AN-24; ANTONOV AN26B/32; DOUGLAS DC4 SKYMASTER; NAMC YS11; VICKERS VISCOUNT 700 |
| ATR 42 - 320 | ATR42-300; BRISTOL 170 FREIGHTER; CONVAIR 240/340/440; GULF AMERICAN GULFSTREAM I; ILYUSHIN IL12/IL14 |
| ATR 42 - 45 | ATR42-500 |
| ATR 72 - 200 | ATR72; ATR72 200/500; ATR72 200/500/600; HANDLEY PAGE HERALD 200; HANDLEY PAGE HERALD 700; NORD 2501 NORTALAS |
| Avro RJ85 | AVROLINER RJ100/115; AVROLINER RJ70; AVROLINER RJ85/QT |
| BAe 1-11 | AEROSPATIALE CARAVELLE 10B/10R; AEROSPATIALE CARAVELLE 12; AEROSPATIALE CARAVELLE 6/6R; BAE(BAC)111-200; BAE(BAC)111-300/400/475; BAE(BAC)111-500; GA GULFSTREAM 3; GULF AMERICAN GULFSTREAM II; TUPOLEV TU124 |
| Bae Jetstream 31 | BAE JETSTREAM 31/32 |
| Bae Jetstream 41 | BAE JETSTREAM 41 |
| BAe146 -100/200/300 | BAE 146-100; BAE 146-200/QT; BAE 146-300 |
| Beech 1900C airline | AEROSPATIALE (NORD)262; BEECHCRAFT 1900C/D AIRLINER; BEECHCRAFT STARSHIP MODEL 2000; DOUGLAS DC3 C47 DAKOTA |
| Beech Super King Air 200B | BEECHCRAFT 200 SUPERKING AIR; BEECHCRAFT B200 SUPERKING AIR; PIAGGIO P.180 AVANTI |
| Beech Super King Air 350 | BEECHCRAFT 300 / 350 SUPER KING AIR; PIPER PA42 CHEYENNE III/IV |
| Boeing 727-100 | BOEING 727-100/100C |
| Boeing 727-200 | BOEING 727-200/200 ADVANCED; TUPOLEV TU154A/B; TUPOLEV TU154M |
| Boeing 737 100 | ANTONOV 148/158; ANTONOV AN72; ANTONOV AN72 / 74; BOEING 737-100; CONVAIR 880; GULF AMERICAN GULFSTREAM 500-550; GULF AMERICAN GULFSTREAM IV; TUPOLEV TU134 |
| Boeing 737-200 | BOEING 737-200; DASSAULT-BREGUET MERCURE; GULFSTREAM G650 |
| Boeing 737-300 | BOEING 737-300 |
| Boeing 737-400 | BOEING 737-400 |
| Boeing 737-500 | BOEING 737-500 |
| Boeing 737-600 | BOEING 737-600 |
| Boeing 737-700 | BOEING 737-700; BOEING BBJ |
| Boeing 737-800 | BOEING 737-800; BOEING 737-900; BOEING 737-900 ER |
| Boeing 747-100/300/800 | BAC/AEROSPATIALE CONCORDE; BOEING 747-100/100F; BOEING 747-300(STRETCH UP DK); BOEING 747-300M (COMBI); BOEING 747-8 (FREIGHTER); BOEING 747-8 (I); BOEING 747SP |
| Boeing 747-200 | ANTONOV AN-124; ANTONOV AN-225 MRIYA; BOEING 747-200B; BOEING 747-200B (COMBI); BOEING 747-200C/200F |
| Boeing 747-400 | BOEING 747-400; BOEING 747-400F; BOEING 747-400M (COMBI) |

| EMEP/EEA Aircraft Type | CAA Aircraft Types |
|-------------------------------|---|
| Boeing 757-200 | BOEING 757-200 |
| Boeing 757-300 | BOEING 757-300 |
| Boeing 767 200 | BOEING 767-200; BOEING 767-200ER |
| Boeing 767 300 ER | BOEING 767-300; BOEING 767-300ER/F; BOEING 767-400ER; BOEING 787-800 DREAMLINER; BOEING 787-900 DREAMLINER |
| Boeing 777-200 ER | BOEING 777-200; BOEING 777-200ER |
| Boeing 777-200 LRF | BOEING 777 FREIGHTER; BOEING 777 FREIGHTER SERIES; BOEING 777-200LR |
| Boeing 777-300 | BOEING 777-300 |
| Boeing 777-300 ER | BOEING 777-300ER |
| Canadair Regional Jet CRJ-200 | BOMBARDIER CHALLENGER 850; BOMBARDIER REGIONAL JET 100/200; DASSAULT FALCON 7X |
| Canadair Regional Jet CRJ-900 | BOMBARDIER GLOBAL 5000; BOMBARDIER GLOBAL EXPRESS; BOMBARDIER GLOBAL EXPRESS (BD700 EXEC); BOMBARDIER REGIONAL CRJ 1000/1000 ER; BOMBARDIER REGIONAL JET CRJ900; BOMBARDIER REGIONAL JET CRJ900 ER; BOMBARDIER REGIONAL JET CRJ900 ER/LR; BOMBARDIER REGIONAL JET RJ700; BOMBARDIER REGIONAL JET RJ700ER |
| Cessna 208 Caravan | Other small piston aircraft |
| Cessna Citation II | Other small jets |
| Dash 8 A | BOMBARDIER DASH 8 Q100/200; DE HAVILLAND DASH 8-100 |
| Dash 8 C | DE HAVILLAND DASH 8-300/Q300 |
| Dash 8 D | ARMSTRONG WHITWORTH ARGOSY; BOMBARDIER DASH 8 Q400; DE HAVILLAND DASH 8 Q400 |
| Dornier 328-110 | DE HAVILLAND DHC-7 DASH-7; DORNIER 328 |
| Embraer 110P2A Bandeirante | Other small turboprops |
| Embraer EMB 120 Brasilia | EMBRAER EMB 120 BRASILIA |
| Embraer ERJ145 | BOMBARDIER CHALLENGER 300; BOMBARDIER CHALLENGER 300/350; EMBRAER LEGACY 600 (BJ135); EMBRAER LEGACY 600/650 (BJ135); EMBRAER RJ135; EMBRAER RJ145; LOCKHEED JETSTAR II |
| Embraer ERJ170-ERJ175 | EMB ERJ170 (170-100); EMB ERJ175 (170-200); EMBRAER ERJ 170; EMBRAER ERJ170; EMBRAER ERJ175 |
| Embraer ERJ190 | CANADAIR CL-600-604 CHALLENGER; EMBRAER 195; EMBRAER ERJ190; EMBRAER ERJ195 |
| Falcon 2000 | BAE125-1000; CESSNA 680 CITATION SOVEREIGN; CESSNA 750 CITATION X; DASSAULT BREGUET FALCON 50; DASSAULT MYSTERE-FALCON 2000; DASSAULT MYSTERE-FALCON 900; DASSAULT MYSTERE-FALCON 900EX; DORNIER 328 JET; EMBRAER LEGACY 500 (EMB-550); GULFSTREAM G200 (IAI GALAXY); HAWKER 4000; LEARJET 60; RAYTHEON HAWKER HORIZON; YAKOVLEV YAK-40 |

| EMEP/EEA Aircraft Type | CAA Aircraft Types |
|------------------------------------|---|
| Fokker F100 | FOKKER 100; FOKKER 70 |
| Fokker F27 | BAE (HS) 748; FAIRCHILD HILLER FH 227B; FOKKER F27 100-400/600; FOKKER F27-500 |
| Fokker F28 | FOKKER F28-1000; FOKKER F28-2000; FOKKER F28-3000; FOKKER F28-4000/6000 |
| Fokker F50 | BAE ATP; FOKKER 50 |
| Let L-410 Turbolet | LET 410; MITSUBISHI MU2; SHORTS SC7 SKYLINER; SHORTS SC7 SKYVAN |
| Lockheed C-130H Hercules | AEROSPACELINES B377SUPER GUPPY; ANTONOV AN-12; CANADAIR CL-44; ILYUSHIN IL18; LOCKHEED L100 HERCULES; LOCKHEED L188 ELECTRA; SHORTS BELFAST; V953C MERCHANTMAN |
| McDonnell Douglas DC-10 | LOCKHEED L1011-1/100 TRISTAR; LOCKHEED L1011-200 TRISTAR; LOCKHEED L1011-500 TRISTAR; MCDONNELL-DOUGLAS DC10-10; MCDONNELL-DOUGLAS DC10-30; MCDONNELL-DOUGLAS DC10-40 |
| McDonnell Douglas DC8-50 | BOEING 707 ALL SERIES; BOEING 707-120/121B; BOEING 720B; MCDONNELL-DOUGLAS DC8-10/50; MCDONNELL-DOUGLAS DC8F 54/55 |
| McDonnell Douglas DC8-60/70 | ILYUSHIN IL62; MCDONNELL-DOUGLAS DC861/3 71/3; MCDONNELL-DOUGLAS DC8-62/72; MCDONNELL-DOUGLAS DC8-71/73 |
| McDonnell Douglas DC-9-10 | MCDONNELL-DOUGLAS DC9-10/15 |
| McDonnell Douglas DC-9-20/30/40/50 | MCDONNELL-DOUGLAS DC9-20; MCDONNELL-DOUGLAS DC9-30; MCDONNELL-DOUGLAS DC9-40; MCDONNELL-DOUGLAS DC9-50 |
| McDonnell Douglas MD-11 | MCDONNELL-DOUGLAS MD11 |
| McDonnell Douglas MD-82/87/88 | BOEING 717-200; MCDONNELL-DOUGLAS MD80-MD83; MCDONNELL-DOUGLAS MD87; MCDONNELL-DOUGLAS MD88; SUKHOI RRJ95 |
| McDonnell Douglas MD-83 | Mc DONNELL DOUGLAS MD90; TUPOLEV TU104; YAKOVLEV YAK-42 |
| Saab 2000 | CONVAIR 580/600/640; DOUGLAS DC6/6A/6B/6C; SAAB 2000; VICKERS VISCOUNT 800 |
| Saab 340B | SAAB FAIRCHILD 340 |
| Shorts 360-300 | SHORTS 330; SHORTS 360 |
| Swearingen Metro III | FAIRCHILD SA-227 METRO 23; FAIRCHILD SA-227 METRO III; SWEARINGEN MERLIN IIA/IIB/IIIB; SWEARINGEN MERLIN IVA; SWEARINGEN METRO II |
| Tupolev TU 204 | ILYUSHIN 76 90VD (PERM); ILYUSHIN IL76; ILYUSHIN IL86; ILYUSHIN IL96-300; TUPOLEV TU204 |

A 3.1.5 Gas leakage

An overview of the time series of estimates of gas leakage at the point of use, together with overall gas use by economic sector and appliance type is presented in **Table A 3.1.3** below.

Table A 3.1.3 Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers

| Source / Appliance type | Units | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 |
|--|-------------------------|-------------|-------------|------------|-------------|------------|-------------|-------------|-------------|
| Annual Gas Use | | | | | | | | | |
| Domestic gas fires | ktoe | 462 | 520 | 621 | 650 | 673 | 475 | 498 | 474 |
| Domestic manual ignition hobs / cookers | ktoe | 590 | 530 | 511 | 496 | 444 | 459 | 485 | 487 |
| Domestic auto-ignition hobs / cookers | ktoe | 211 | 190 | 183 | 177 | 159 | 164 | 174 | 174 |
| Domestic auto-ignition space and water heating | ktoe | 24572 | 26796 | 30491 | 31512 | 32223 | 24490 | 25617 | 24404 |
| Service sector catering (ovens and hobs) | ktoe | 596 | 762 | 772 | 774 | 705 | 570 | 590 | 572 |
| Other service sector appliances (boilers) | ktoe | 6646 | 8508 | 9818 | 9290 | 8643 | 8106 | 8412 | 8143 |
| Methane Leakage | | | | | | | | | |
| Domestic cooking and gas fires | ktCH ₄ | 1.02 | 0.94 | 0.86 | 0.85 | 0.8 | 0.78 | 0.83 | 0.84 |
| Domestic boilers and water heating | ktCH ₄ | 0.76 | 0.83 | 0.94 | 0.98 | 1 | 0.76 | 0.79 | 0.76 |
| Service sector (all sources) | ktCH ₄ | 0.83 | 1.06 | 1.09 | 1.05 | 1 | 0.91 | 0.95 | 0.93 |
| Total | ktCH₄ | 2.61 | 2.83 | 2.9 | 2.88 | 2.8 | 2.46 | 2.57 | 2.52 |

A 3.2 INDUSTRIAL PROCESSES (CRF SECTOR 2)

There is currently no additional information for this sector in this Annex.

A 3.3 AGRICULTURE (CRF SECTOR 3)

Note that the references for this section are included in **Section 17.4**.

Table A 3.3.1 Livestock Population Data by Animal Type ('000 animal places)

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 |
|--|---------|---------|---------|---------|---------|---------|---------|---------|
| Total cattle | 12,125 | 11,760 | 11,048 | 10,698 | 10,014 | 9,785 | 9,886 | 9,837 |
| - dairy cows | 2,848 | 2,603 | 2,336 | 2,003 | 1,839 | 1,906 | 1,910 | 1,901 |
| - all other cattle | 9,277 | 9,157 | 8,713 | 8,695 | 8,175 | 7,879 | 7,977 | 7,936 |
| Sheep | 45,380 | 44,174 | 43,117 | 36,138 | 31,727 | 34,034 | 34,663 | 35,577 |
| Pigs | 7,548 | 7,627 | 6,482 | 4,862 | 4,468 | 4,739 | 4,866 | 4,969 |
| Total poultry | 138,381 | 142,267 | 169,773 | 173,909 | 163,842 | 167,579 | 172,607 | 181,811 |
| - laying hens | 33,624 | 31,837 | 28,687 | 29,544 | 28,751 | 28,311 | 29,184 | 30,193 |
| - broilers | 73,944 | 77,177 | 105,689 | 111,475 | 105,309 | 107,056 | 110,639 | 117,612 |
| Total horses | 570 | 684 | 1006 | 1036 | 1024 | 978 | 963 | 954 |
| - horses kept on agricultural holdings | 202 | 273 | 287 | 346 | 312 | 283 | 268 | 258 |
| - professional horses | 62 | 62 | 70 | 91 | 91 | 87 | 87 | 87 |
| - domestic horses | 305 | 348 | 649 | 599 | 621 | 608 | 608 | 608 |
| Goat | 98 | 75 | 74 | 95 | 93 | 101 | 102 | 105 |
| Deer | 47 | 37 | 36 | 33 | 31 | 31 | 31 | 31 |

A 3.3.1 Enteric Fermentation (3A)

Table A 3.3.2 Methane Emission Factors for Livestock Emissions for 2017

| Animal type | | Enteric methane kg CH ₄ /head/year | Methane from manures kg CH ₄ /head/year |
|-------------|----------------------------|--|---|
| Cattle | Dairy cows | 122.70 | 37.24 |
| | Dairy heifers | 52.25 | 6.90 |
| | Dairy replacements >1 year | 49.25 | 6.79 |
| | Dairy calves <1 year | 42.59 | 5.42 |
| | Beef cows | 76.49 | 11.48 |
| | Beef females for slaughter | 49.74 | 6.81 |
| | Bulls for breeding | 59.67 | 9.15 |
| | Cereal fed bull | 50.49 | 10.27 |
| | Heifers for breeding | 48.89 | 6.86 |
| | Steers | 50.51 | 6.87 |
| Pigs | | 1.50 | 5.20 |

| Animal type | | Enteric methane kg CH ₄ /head/year | Methane from manures kg CH ₄ /head/year |
|-------------|-------------------|--|--|
| Sheep | Ewes | 6.48 | 0.18 |
| | Rams | 7.00 | 0.19 |
| | Lambs | 2.74 | 0.07 |
| Goats | | 5.0 | 0.13 |
| Horses | | 18.0 | 1.56 |
| Deer | | 20.0 | 0.22 |
| Poultry | Laying hens | NA | 0.028 |
| | Growing pullets | NA | 0.013 |
| | Broilers | NA | 0.013 |
| | Turkeys | NA | 0.091 |
| | Breeding flock | NA | 0.013 |
| | Ducks | NA | 0.289 |
| | Geese | NA | 0.289 |
| | All other poultry | NA | 0.013 |

A 3.3.2 Manure Management (3B)

A 3.3.2.1 Methane emissions from animal manures

Table A 3.3.3 Methane conversion factors for Manure Management Systems in the UK

^aNo differentiation is made between crusted and non-crusted slurry storage

| Manure Handling System | Methane Conversion Factor % |
|--|-----------------------------|
| Liquid ^a | 17 |
| Daily spread | 0.1 |
| Deep bedding/farm yard manure – cattle, pigs | 17 |
| Deep bedding/farm yard manure – sheep | 2.0 |
| Pasture range and paddock | 1.0 |
| Poultry manure | 1.5 |

A 3.3.2.2 Nitrous Oxide emissions from Animal Waste Management Systems**Table A 3.3.4 Nitrogen Excretion Factors, kg N animal place⁻¹ year⁻¹ for livestock in the UK (1990-2017)**

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 |
|--|------|------|------|-------|-------|-------|-------|-------|
| Dairy cows | 86.6 | 88.2 | 94.5 | 102.2 | 105.5 | 109.6 | 107.0 | 109.9 |
| Other cattle ^a | 45.3 | 45.6 | 46.4 | 45.7 | 45.5 | 45.3 | 45.1 | 44.8 |
| Sows | 23.6 | 22.5 | 21.6 | 20.1 | 18.1 | 18.1 | 18.1 | 18.1 |
| Gilts | 15.5 | 15.5 | 15.5 | 15.5 | 15.5 | 15.5 | 15.5 | 15.5 |
| Boars | 28.7 | 27.4 | 26.1 | 24.5 | 21.8 | 21.8 | 21.8 | 21.8 |
| Fatteners > 80 kg | 20.2 | 19.3 | 18.4 | 17.2 | 15.4 | 15.4 | 15.4 | 15.4 |
| Fatteners 20-80 kg | 14.6 | 13.9 | 13.2 | 12.4 | 11.1 | 11.1 | 11.1 | 11.1 |
| Weaners (<20 kg) | 4.6 | 4.4 | 4.2 | 3.9 | 3.4 | 3.4 | 3.4 | 3.4 |
| Ewes | 6.7 | 6.8 | 6.8 | 6.8 | 6.8 | 7.0 | 6.8 | 6.8 |
| Rams | 9.1 | 9.1 | 9.1 | 8.9 | 8.9 | 9.0 | 8.9 | 8.9 |
| Lambs | 3.0 | 2.9 | 3.0 | 3.2 | 3.2 | 3.4 | 3.3 | 3.3 |
| Goats | 20.6 | 20.6 | 20.6 | 20.6 | 20.6 | 20.6 | 20.6 | 20.6 |
| Horses | | | | | | | | |
| – horses kept on agricultural holdings | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |
| – professional horses | 129 | 129 | 129 | 129 | 129 | 129 | 129 | 129 |
| – domestic horses | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |
| Deer | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 |
| Laying hens | 0.87 | 0.83 | 0.80 | 0.77 | 0.70 | 0.70 | 0.70 | 0.70 |

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 |
|--------------------|------|------|------|------|------|------|------|------|
| Broilers | 0.64 | 0.59 | 0.55 | 0.49 | 0.40 | 0.40 | 0.40 | 0.40 |
| Turkeys | 1.50 | 1.59 | 1.68 | 1.76 | 1.82 | 1.82 | 1.82 | 1.82 |
| Pullets | 0.42 | 0.39 | 0.36 | 0.34 | 0.33 | 0.33 | 0.33 | 0.33 |
| Breeding flock | 1.16 | 1.13 | 1.10 | 1.07 | 1.02 | 1.02 | 1.02 | 1.02 |
| Ducks | 1.30 | 1.41 | 1.52 | 1.62 | 1.71 | 1.71 | 1.71 | 1.71 |
| Geese | 1.30 | 1.41 | 1.52 | 1.62 | 1.71 | 1.71 | 1.71 | 1.71 |
| Other poultry | 1.30 | 1.41 | 1.52 | 1.62 | 1.71 | 1.71 | 1.71 | 1.71 |

^aWeighted average for all other cattle categories

Table A 3.3.5 Distribution of Animal Waste Management Systems (%) used for Different Animal types, 2017

| Animal Type | | Liquid System | Daily Spread | Solid storage/Deep litter/Poultry litter ^b | Pasture Range and Paddock |
|-------------|------------------|---------------|--------------|---|---------------------------|
| Cattle | Dairy cows | 61.0 | 8.2 | 9.3 | 21.5 |
| | All other cattle | 18.5 | 11.7 | 21.7 | 48.2 |
| Pigs | All pigs | 35.2 | 14.7 | 39.4 | 10.7 |
| Sheep | Ewes | 0.0 | 0.0 | 8.9 | 91.1 |
| | Rams | 0.0 | 0.0 | 0.0 | 100.0 |
| | Lambs | 0.0 | 0.0 | 0.6 | 99.4 |
| Goats | | 0.0 | 0.0 | 8.2 | 91.8 |
| Deer | | 0.0 | 0.0 | 24.9 | 75.1 |
| Horses | | 0.0 | 0.0 | 30.1 | 69.9 |

| Animal Type | | Liquid System | Daily Spread | Solid storage/Deep litter/Poultry litter ^b | Pasture Range and Paddock |
|-------------|-------------|---------------|--------------|---|---------------------------|
| Poultry | All poultry | 0.0 | 35.8 | 61.4 | 2.9 |

Table A 3.3.6 Nitrous Oxide Emission Factors for Animal Waste Handling Systems

| Emission source | EF (% of total N) | Uncertainty limits (95% CI) | Data source |
|------------------------|-------------------|-----------------------------|-----------------------------|
| Cattle housing | | | |
| Slurry – solid floor | 0 | N/A | IPCC 2006 |
| Slurry – slatted floor | 0.2 | Factor of 2 | IPCC 2006 |
| FYM systems | 2.0 | Factor of 2 | UK measurement (at storage) |
| Outdoor yards | 0 | N/A | IPCC 2006 |
| Cattle manure storage | | | |
| Slurry – without crust | 0 | N/A | IPCC 2006 |
| Slurry – with crust | 0.5 | Factor of 2 | IPCC 2006 |
| Weeping wall store | 0.5 | Factor of 2 | IPCC 2006 |
| FYM heap | 0.0 | N/A | Included in housing |
| Pig housing | | | |
| Slurry – slatted floor | 0.2 | Factor of 2 | IPCC 2006 |
| FYM systems | 2.0 | Factor of 2 | UK measurement (at storage) |
| Pig manure storage | | | |
| Slurry (no crust) | 0 | N/A | IPCC 2006 |
| FYM heap | 0 | N/A | Included in housing |
| Sheep housing (FYM) | 2.0 | Factor of 2 | Based on cattle/pig |
| Sheep FYM storage | 0 | N/A | Included in housing |
| Layer housing | 0.5 | Factor of 2 | UK measurement (at storage) |
| Layer manure storage | 0 | N/A | Included in housing |
| Broiler housing | 0.5 | Factor of 2 | UK measurement (at storage) |
| Broiler litter storage | 0 | N/A | Included in housing |
| Duck housing | 2.0 | Factor of 2 | Based on cattle/pig |
| Duck manure storage | 0 | N/A | Included in housing |

A 3.3.3 Agricultural Soils (3D)**A 3.3.3.1 Inorganic Fertiliser****Table A 3.3.7 EF for direct N₂O emissions from managed soils in the UK inventory**

| Emission source | EF (% of total N) | Uncertainty | Data source |
|--|---|-------------|----------------------|
| Urea fertiliser | Non-linear function of application rate (see Section 5.5.2.1) | | Topp et al., 2016 |
| Other mineral fertilisers | Non-linear function of application rate and annual rainfall (see Section 5.5.2.1) | | Topp et al., 2016 |
| Livestock slurry | 0.7475 | SE 0.17328 | Topp et al., 2016 |
| Livestock solid manure (FYM, poultry manure) | 0.3635 | SE 0.06622 | Topp et al., 2016 |
| Sewage sludge | 1.0 | 0.3 – 3.0 | IPCC 2006 |
| Crop residues | 1.0 | 0.3 – 3.0 | IPCC 2006 |
| N mineralisation | 1.0 | 0.3 – 3.0 | IPCC 2006 |
| Histosols | 8 kg N ₂ O-N/ha | 0 - 24 | IPCC 2006 |
| Cattle urine | 0.629 | SE 0.0930 | Topp et al., 2016 |
| Cattle dung | 0.193 | SE 0.0212 | Topp et al., 2016 |
| Sheep, goat, horse and deer urine | 0.629 | SE 0.0930 | Cattle value assumed |
| Sheep, goat, horse and deer dung | 0.193 | SE 0.0212 | Cattle value assumed |
| Outdoor pig and poultry | | | IPCC 2006 |

Table A 3.3.8 Areas of UK Crops and quantities of fertiliser applied for 2017

| Crop Type | Crop area, ha | Fertiliser, ktN | Crop Type | Crop area, ha | Fertiliser, ktN |
|-----------------------------|---------------|-----------------|----------------------------|---------------|-----------------|
| Oats | 128,425 | 13.5 | Potatoes (maincrop) | 120,557 | 16.2 |
| Spring oats | 23,456 | 2.0 | Potatoes (seed or earlies) | 24,343 | 3.3 |
| Winter oats | 9,168 | 0.8 | Sugar beet | 111,269 | 10.5 |
| Spring barley | 14,031 | 1.4 | Maize | 70,196 | 4.9 |
| Spring barley (malting) | 392,939 | 43.6 | Grain maize | 7,759 | 0.5 |
| Spring barley (non-malting) | 346,924 | 33.7 | Forage maize | 119,475 | 7.4 |
| Winter barley | 54,623 | 8.5 | Rootcrops for stockfeed | 28,509 | 2.2 |

| Crop Type | Crop area, ha | Fertiliser, ktN | Crop Type | Crop area, ha | Fertiliser, ktN |
|---|---------------|-----------------|---------------------------------|---------------|-----------------|
| Winter barley (malting) | 76,614 | 10.5 | Leafy forage crops | 4,153 | 0.3 |
| Winter barley (non-malting) | 291,610 | 45.9 | Other fodder crops | 48,823 | 4.0 |
| Wheat | 8,730 | 1.5 | Vegetables (not-differentiated) | 1,717 | 0.1 |
| Wheat (milling) | 637,374 | 126.3 | Vegetables (brassicas) | 4,855 | 0.4 |
| Wheat (non-milling) | 1,145,725 | 199.3 | Vegetables (legumes) | 39,941 | 0.0 |
| Minor cereals | 51,631 | 5.7 | Vegetables (other non-legumes) | 72,558 | 5.7 |
| Oilseed rape | 5,086 | 0.9 | Other horticultural crops | 14,690 | 1.2 |
| Spring oilseed rape | 8,674 | 1.6 | Soft Fruit | 8,558 | 0.6 |
| Winter oilseed rape | 548,637 | 100.0 | Top Fruit | 24,634 | 1.8 |
| Linseed | 26,375 | 2.2 | Miscanthus | 7,366 | 0.0 |
| Field beans (harvested dry) | 191,711 | 0.2 | Willow (short rotation coppice) | 3,610 | 0.0 |
| Field peas (harvested dry) | 39,864 | 0.0 | Other field crops | 25,365 | 2.1 |
| Field beans and peas combined not Vining peas | 0.0 | 0.0 | Wine grapes | 1,992 | 0.1 |
| Fruit (mixed top & soft fruit) | 11 | 0.0 | | | |
| Permanent grass | 6,135,379 | 297.5 | Temporary grass | 1,144,430 | 112.3 |

Table A 3.3.9 Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2017

| Year | Wheat | | Spring barley | | Winter barley | | Main crop potatoes | | Oilseed rape | | Grass leys (<5yrs) | | Permanent grassland | |
|------|---------|---------|---------------|---------|---------------|---------|--------------------|---------|--------------|---------|--------------------|---------|---------------------|---------|
| | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N |
| 1990 | 2,014 | 183 | 635 | 90 | 882 | 140 | 148 | 184 | 390 | 225 | 1,606 | 166 | 5,316 | 108 |
| 1991 | 1,980 | 187 | 552 | 89 | 841 | 140 | 148 | 185 | 440 | 221 | 1,603 | 168 | 5,334 | 107 |
| 1992 | 2,067 | 185 | 515 | 89 | 784 | 141 | 151 | 175 | 421 | 196 | 1,579 | 157 | 5,286 | 94 |
| 1993 | 1,759 | 185 | 518 | 91 | 650 | 136 | 143 | 189 | 377 | 179 | 1,567 | 146 | 5,278 | 100 |
| 1994 | 1,811 | 186 | 481 | 94 | 628 | 143 | 138 | 191 | 404 | 179 | 1,456 | 170 | 5,375 | 110 |
| 1995 | 1,859 | 193 | 504 | 97 | 689 | 144 | 144 | 176 | 354 | 187 | 1,407 | 170 | 5,375 | 108 |
| 1996 | 1,977 | 185 | 518 | 93 | 749 | 140 | 149 | 171 | 356 | 190 | 1,396 | 166 | 5,347 | 104 |
| 1997 | 2,036 | 192 | 518 | 94 | 839 | 143 | 133 | 166 | 445 | 199 | 1,394 | 147 | 5,290 | 103 |
| 1998 | 2,045 | 182 | 484 | 91 | 769 | 135 | 131 | 186 | 507 | 192 | 1,302 | 156 | 5,365 | 99 |
| 1999 | 1,847 | 185 | 631 | 99 | 548 | 142 | 148 | 153 | 495 | 196 | 1,226 | 180 | 5,449 | 102 |
| 2000 | 2,086 | 188 | 539 | 106 | 589 | 146 | 138 | 157 | 395 | 189 | 1,226 | 142 | 5,363 | 90 |
| 2001 | 1,635 | 185 | 783 | 109 | 462 | 144 | 137 | 153 | 446 | 196 | 1,205 | 130 | 5,584 | 84 |
| 2002 | 1,996 | 189 | 555 | 110 | 546 | 150 | 129 | 152 | 436 | 194 | 1,243 | 135 | 5,519 | 77 |
| 2003 | 1,836 | 197 | 621 | 107 | 455 | 148 | 118 | 149 | 549 | 194 | 1,200 | 128 | 5,683 | 75 |
| 2004 | 1,990 | 190 | 587 | 101 | 420 | 144 | 121 | 164 | 498 | 171 | 1,246 | 117 | 5,620 | 71 |
| 2005 | 1,870 | 188 | 553 | 98 | 384 | 140 | 113 | 164 | 588 | 186 | 1,193 | 111 | 5,711 | 66 |
| 2006 | 1,836 | 181 | 494 | 100 | 388 | 134 | 117 | 142 | 568 | 178 | 1,137 | 106 | 5,967 | 60 |
| 2007 | 1,830 | 183 | 515 | 97 | 383 | 134 | 112 | 136 | 674 | 179 | 1,176 | 98 | 5,965 | 56 |

| Year | Wheat | | Spring barley | | Winter barley | | Main crop potatoes | | Oilseed rape | | Grass leys (<5yrs) | | Permanent grassland | |
|------|---------|---------|---------------|---------|---------------|---------|--------------------|---------|--------------|---------|--------------------|---------|---------------------|---------|
| | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N | '000 ha | kg/ha N |
| 2008 | 2,080 | 179 | 616 | 92 | 416 | 134 | 114 | 150 | 598 | 180 | 1,141 | 92 | 6,036 | 45 |
| 2009 | 1,814 | 188 | 749 | 98 | 411 | 137 | 118 | 169 | 581 | 167 | 1,262 | 88 | 6,081 | 48 |
| 2010 | 1,939 | 192 | 539 | 96 | 382 | 140 | 114 | 137 | 642 | 186 | 1,231 | 98 | 5,925 | 53 |
| 2011 | 1,969 | 194 | 611 | 99 | 359 | 140 | 120 | 157 | 705 | 186 | 1,278 | 91 | 5,877 | 51 |
| 2012 | 1,992 | 194 | 618 | 99 | 385 | 144 | 123 | 141 | 756 | 183 | 1,357 | 92 | 5,799 | 50 |
| 2013 | 1,615 | 183 | 903 | 108 | 310 | 143 | 114 | 160 | 715 | 149 | 1,390 | 94 | 5,802 | 55 |
| 2014 | 1,936 | 192 | 651 | 108 | 429 | 145 | 115 | 149 | 675 | 186 | 1,396 | 97 | 5,824 | 51 |
| 2015 | 1,832 | 193 | 659 | 105 | 442 | 148 | 105 | 162 | 652 | 190 | 1,135 | 96 | 5,886 | 49 |
| 2016 | 1,821 | 189 | 683 | 107 | 439 | 147 | 114 | 141 | 579 | 182 | 1,144 | 94 | 6,118 | 50 |
| 2017 | 1,792 | 183 | 754 | 104 | 423 | 154 | 121 | 134 | 562 | 182 | 1,144 | 98 | 6,135 | 48 |

A 3.3.3.2 Crop Residues**Table A 3.3.10 Parameter values for crop residue management.**

| Crop | Crop Harvest Index ^a | Above Ground Residue Retained after harvest | IPPC Crop Yield To Above Ground Residue Slope ^b | IPPC Crop Yield To Above Ground Residue Intercept ^b | IPCC Above To Below Ground Residue ratio |
|---|---------------------------------|---|--|--|--|
| Oats | 0.46 | 0.5 | NA | NA | 0.25 |
| Spring oats | 0.46 | 0.5 | NA | NA | 0.25 |
| Winter oats | 0.46 | 0.5 | NA | NA | 0.25 |
| Spring barley | 0.52 | 0.5 | NA | NA | 0.22 |
| Spring barley (malting) | 0.52 | 0.5 | NA | NA | 0.22 |
| Spring barley (non-malting) | 0.52 | 0.5 | NA | NA | 0.22 |
| Winter barley | 0.52 | 0.5 | NA | NA | 0.22 |
| Winter barley (malting) | 0.52 | 0.5 | NA | NA | 0.22 |
| Winter barley (non-malting) | 0.52 | 0.5 | NA | NA | 0.22 |
| Wheat | 0.50 | 0.5 | NA | NA | 0.23 |
| Wheat (milling) | 0.50 | 0.5 | NA | NA | 0.23 |
| Wheat (non-milling) | 0.50 | 0.5 | NA | NA | 0.23 |
| Minor cereals | 0.49 | 0.5 | NA | NA | 0.23 |
| Oilseed rape | 0.30 | 1 | NA | NA | 0.35 |
| Spring oilseed rape | 0.30 | 1 | NA | NA | 0.35 |
| Winter oilseed rape | 0.30 | 1 | NA | NA | 0.35 |
| Linseed and Flax | 0.38 | 0.5 | NA | NA | 0.35 |
| Linseed | 0.38 | 1 | NA | NA | 0.35 |
| Flax | 0.38 | 0.2 | NA | NA | 0.35 |
| Field beans (harvested dry) | NA | 1 | 1.13 | 0.85 | 0.19 |
| Field peas (harvested dry) | NA | 1 | 1.13 | 0.85 | 0.19 |
| Field beans and peas combined not Vining peas | NA | 1 | 1.13 | 0.85 | 0.19 |
| Potatoes | NA | 1 | 0.10 | 1.06 | 0.20 |
| Potatoes (maincrop) | NA | 1 | 0.10 | 1.06 | 0.20 |
| Potatoes (seed or earlies) | NA | 1 | 0.10 | 1.06 | 0.20 |
| Sugar beet | NA | 1 | 1.07 | 1.54 | 0.20 |
| Maize | NA | 1 | 1.03 | 0.61 | 0.22 |
| Grain maize | NA | 1 | 1.03 | 0.61 | 0.22 |
| Forage maize | NA | 0.15 | 1.03 | 0.61 | 0.22 |
| Rootcrops for stockfeed | NA | 0.15 | 1.07 | 1.06 | 0.20 |
| Leafy forage crops | NA | 0.15 | 0.30 | 0.00 | 0.35 |
| Other fodder crops | NA | 0.1 | NA | NA | 0.35 |
| Vegetables (not-differentiated) | NA | 1 | 0.30 | 0.00 | 0.35 |
| Vegetables (brassicas) | NA | 1 | 0.30 | 0.00 | 0.35 |
| Vegetables (legumes) | NA | 1 | 0.30 | 0.00 | 0.35 |
| Vegetables (other non-legumes) | NA | 1 | 0.30 | 0.00 | 0.35 |
| Other horticultural crops | NA | 1 | 0.30 | 0.00 | 0.35 |
| Soft Fruit | NA | 1 | 0.20 | 0.00 | 0.35 |

| Crop | Crop Harvest Index ^a | Above Ground Residue Retained after harvest | IPPC Crop Yield To Above Ground Residue Slope ^b | IPPC Crop Yield To Above Ground Residue Intercept ^b | IPCC Above To Below Ground Residue ratio |
|---------------------------------|---------------------------------|---|--|--|--|
| Top Fruit | NA | 1 | 0.20 | 0.00 | 0.35 |
| Miscanthus | NA | 1 | 1.00 | 0.00 | 0.35 |
| Willow (short rotation coppice) | NA | 1 | 1.00 | 0.00 | 0.35 |
| Other field crops | 0.52 | 0.5 | NA | NA | 0.22 |
| Wine grapes | NA | 1 | 0.20 | 0.00 | 0.35 |
| Fruit (mixed top & soft fruit) | NA | 1 | 0.20 | 0.00 | 0.35 |
| Field beans and peas combined | NA | 1 | 1.13 | 0.00 | 0.19 |

^aWhere 'NA' appears in the Harvest Index column, it indicates that the IPCC 2006 method was used; ^bwhere 'NA' appears in the IPCC slope or intercept column, it means that the Harvest Index approach was used

Table A 3.3.11 N concentrations in above and below ground biomass

| Crop | Below Ground N, kg N/[t DM] | Crop Residue Above Ground N, kg N/[t DM] |
|---|-----------------------------|--|
| Oats | 8.0 | 5.4 |
| Spring oats | 8.0 | 5.4 |
| Winter oats | 8.0 | 5.4 |
| Spring barley | 14.0 | 6.7 |
| Spring barley (malting) | 14.0 | 6.7 |
| Spring barley (non-malting) | 14.0 | 6.7 |
| Winter barley | 14.0 | 6.7 |
| Winter barley (malting) | 14.0 | 6.7 |
| Winter barley (non-malting) | 14.0 | 6.7 |
| Wheat | 9.0 | 6.2 |
| Wheat (milling) | 9.0 | 6.2 |
| Wheat (non-milling) | 9.0 | 6.2 |
| Minor cereals | 9.0 | 6.6 |
| Oilseed rape | 11.0 | 9.9 |
| Spring oilseed rape | 11.0 | 9.9 |
| Winter oilseed rape | 11.0 | 9.9 |
| Linseed and Flax | 11.0 | 9.9 |
| Linseed | 11.0 | 9.9 |
| Flax | 11.0 | 9.9 |
| Field beans (harvested dry) | 8.0 | 8.0 |
| Field peas (harvested dry) | 8.0 | 8.0 |
| Field beans and peas combined not Vining peas | 8.0 | 8.0 |
| Potatoes | 14.0 | 17.3 |
| Potatoes (maincrop) | 14.0 | 17.3 |
| Potatoes (seed or earlies) | 14.0 | 17.3 |
| Sugar beet | 14.0 | 24.6 |
| Maize | 7.0 | 6.0 |
| Grain maize | 7.0 | 6.0 |

| Crop | Below Ground N, kg N/[t DM] | Crop Residue Above Ground N, kg N/[t DM] |
|---------------------------------|-----------------------------|--|
| Forage maize | 7.0 | 6.0 |
| Rootcrops for stockfeed | 14.0 | 12.6 |
| Leafy forage crops | 12.0 | 26.3 |
| Other fodder crops | 14.0 | 6.7 |
| Vegetables (not-differentiated) | 12.0 | 26.1 |
| Vegetables (brassicas) | 12.0 | 38.4 |
| Vegetables (legumes) | 22.0 | 23.2 |
| Vegetables (other non-legumes) | 22.0 | 16.7 |
| Other horticultural crops | 22.0 | 26.1 |
| Soft Fruit | 11.0 | 17.7 |
| Top Fruit | 11.0 | 3.9 |
| Miscanthus | 11.0 | 0.3 |
| Willow (short rotation coppice) | 11.0 | 0.3 |
| Other field crops | 11.0 | 6.7 |
| Wine grapes | 11.0 | 3.3 |
| Fruit (mixed top & soft fruit) | 11.0 | 8.1 |
| Field beans and peas combined | 8.0 | 8.0 |

A 3.3.3.3 Mineralisation**Table A 3.3.12 Mineralised N from soils**

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| N in mineral soils that is mineralised as a result of historical land use change to Cropland (kt N/y) | 39.34 | 44.12 | 52.04 | 72.85 | 86.44 | 87.53 | 87.88 | 88.21 | 88.54 | 88.54 |
| N in mineral soils that is mineralised as a result of Cropland Management (kt N/y) | 0.0002287 | 0.0002600 | 0.0002423 | 0.0002411 | 0.0002338 | 0.0002277 | 0.0002219 | 0.0002314 | 0.0002169 | 0.0002169 |
| Direct N ₂ O emissions from mineralised N as a result of historical land use change to Cropland, kt N ₂ O/y | 0.62 | 0.69 | 0.82 | 1.14 | 1.36 | 1.38 | 1.38 | 1.39 | 1.39 | 1.39 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Direct N ₂ O emissions from mineralised N as a result of Cropland Management, kt N ₂ O/y | 0.000004 | 0.000004 | 0.000004 | 0.000004 | 0.000004 | 0.000004 | 0.000003 | 0.000004 | 0.000003 | 0.000003 |
| Indirect N ₂ O emissions from mineralised N as a result of historical land use change to Cropland and Cropland management (kt N ₂ O/y) | 0.14 | 0.16 | 0.18 | 0.26 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 |
| Total N ₂ O emissions from Mineralisation (kt N/y) | 0.76 | 0.85 | 1.00 | 1.40 | 1.66 | 1.69 | 1.69 | 1.70 | 1.70 | 1.70 |

A 3.3.3.4 Histosols


The total area used to calculate emissions from histosols is 2857 km².


A 3.4 LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4)


The following section describes in detail the methodology used in the LULUCF sector described in **Chapter 6**.


The flow chart (**Figure A 3.1**) shows the interrelationships between different data sources and the main calculation steps.


Figure A 3.1 Data flow diagrams for each land use sub-category, showing cross-linkages between sectors: (i) 4A and 4G, (ii) 4B, (iii) 4C, (iv) 4D, (v) 4E














A 3.4.1 Carbon stock changes due to afforestation and forest management (4A)

A 3.4.1.1 The Forest carbon accounting model CARBINE

Carbon uptake by the forests planted in the UK is calculated by a carbon accounting model, CARBINE, as gains and losses in pools of carbon in standing trees, litter and soil in conifer and broadleaf forests and in harvested wood products. Restocking is assumed in all forests. The method is Tier 3, as defined by IPCC (2006). Matthews *et al.* (2014) gives an overview of the CARBINE model and a comparison of its use in the 1990-2012 LULUCF inventory with the C-Flow model previously used model to forest carbon stock changes.

A key concept underlying the net exchanges of C between forests and the other C reservoirs is that it can be inferred from the changes in C stocks. So CARBINE simulates forest C stock changes represented by tree biomass growth, mortality and subsequent loss. The CARBINE model is primarily dedicated to reproducing the UK forest conditions.

The model as used for this inventory consists of three sub-models or 'compartments' which estimate carbon stocks in the forest biomass, soil, and harvested wood products. The forest biomass carbon sub-model is further compartmentalised to represent fractions due to tree stems, branches, foliage, and roots.

The main driving module of CARBINE consists of a set of computerised mathematical functions and algorithms describing the accumulation (and loss) of carbon in tree biomass of different forestry systems at the per-hectare scale. Different functions and algorithms are used to represent distinct forestry systems, defined in terms of:

- Tree species composition
- Tree growth rate (yield class)
- Management regime applied.

The tree species and growth rates represented are based on yield models originally produced by the British Forestry Commission (Matthews et al., 2016a, 2016b). The tree species covered include examples for coniferous species of spruces, pines, firs, larches, cedars, cypresses and all the major temperate and boreal broadleaf tree species. Growth rates in terms of mean annual increment (MAI) of stem volume can be represented in the range from 2 m³ ha⁻¹ yr⁻¹ up to 30 m³ ha⁻¹ yr⁻¹.

The CARBINE model uses standard estimates for wood density wood carbon content to derive stem biomass from the stem volume predictions simulated by the M1 model (Lavers and Moore, 1983; Jenkins et al., 2011; Matthews, 1993). Wood and bark density along with the carbon content differences are not taken into account. The density of bark is lower than that of wood (Aaron, 1970), but the carbon content is usually higher (Matthews, 1993), hence it is assumed that the two effects cancel out. The volume, biomass and carbon in tree foliage, branches, and coarse and fine roots are derived from the results for the stem by applying expansion factors. Species-specific biomass expansion factors are applied for these calculations.

The biomass of a component of interest is calculated by multiplying stem biomass by a corresponding expansion factor. The UK species-specific crown and root biomass expansion factors were derived from the report of Jenkins *et al.*, (2011) report. Branch biomass is calculated by subtracting foliage biomass from crown biomass. The coarse root biomass expansion factor includes an allowance for stump material. Robust information on foliage expansion factors relevant to UK conditions were not available, hence these were obtained from scientific literature. The ratio of foliage to stem changes over time, but approaches an asymptote (Matthews et al., 1991; Matthews and Duckworth, 2005). However, the asymptote in general is more suited to older, larger trees and as such is considered not to be representative of typical forests under regular management. It was decided to use a biomass expansion relationship for trees of approximately 20 cm diameter in order to better represent managed forests. It is likely that this will ultimately underestimate foliage biomass in smaller trees, and conversely over-estimate in older, larger trees. Finally, fine root biomass is calculated with a uniform expansion factor $\beta_r=0.02$ from a Liski et al., (2002) study. The expansion factors are not sensitive to stand age, management regime or growth rate. This approach was adopted for the simplicity and ease of implementation on the large scale simulations.

The mass of carbon in a forest was calculated from biomass by multiplying by the fraction of carbon in wood (0.5 assumed). As an example, the values used for these parameters for Sitka spruce (*P. Sitchensis*) are given in **Table A 3.4.1**. Sitka spruce is the most common species in UK forests (c. 30%); parameters for other tree species are given in Matthews *et al.* (2014).

Table A 3.4.1 Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce (*P. Sitchensis*), yield class 12.

| Parameter | Value |
|---|-------|
| Time of maximum mean annual increment (years) | 60 |
| Initial spacing (m) | 2 |
| First table age (years) | 20 |
| Age at first thinning (years) | 25 |
| Stemwood density (oven dried tonnes m ⁻³) | 0.33 |

| Parameter | Value |
|---|-------|
| Stemwood conversion loss | 10% |
| % Branchwood left in forest | 100% |
| % Branchwood harvested for fuel | 0% |
| % fuel from bark | 30% |
| % non-fuel products from bark | 70% |
| % small roundwood (underbark) used as fuel | 20% |
| % Pallets and fencing from small roundwood (under bark) | 20% |
| % Paper from small roundwood (under bark) | 35% |
| % Particleboard etc. from small roundwood (under bark) | 25% |
| % Fuel from sawlogs (under bark) | 30% |
| % Pallets and fencing from sawlogs (under bark) | 0% |
| % Particleboard from sawlogs (under bark) | 40% |
| % Structural timber from sawlogs (under bark) | 30% |
| Root:Stem ratio | 0.49 |
| Crown:Stem ratio | 0.32 |
| Foliage:stem ratio | 0.13 |
| Fine root:stem ratio | 0.02 |
| Foliage turnover rate (annual) | 0.2 |
| Branchwood turnover rate (annual) | 0.04 |
| Coarse Root Turnover rate (annual) | 0.02 |
| Fine Root turnover rate (annual) | 0.8 |
| Underbark/overbark ratio at 15cm DBH (varies with DBH) | 0.9 |
| Ratio of thinned stem volume that is sawlog at 15cm DBH (varies with DBH) | 0.05 |

CARBINE includes a sub-model for representing accumulation and loss of carbon in dead wood and litter. Inputs of litter are related to the standing biomass of trees and also to rates of tree mortality. Levels of tree mortality are represented implicitly in the standard Forestry Commission growth models, and explicit estimates are included in models for stands subject to no thinning, where mortality levels are high. Root and branch wood volume associated with dead trees is estimated in the same way as for living stemwood, by reference to allometric relationships. Deadwood and litter are assumed to decay according to a first order process, with rate constants that are normally set to be consistent with boreal and temperate conditions but can be adjusted for Mediterranean and tropical conditions. The other significant input of carbon to the dead wood and litter pool is due to harvesting operations (as part of either thinning or

clearfelling). The carbon in roots of harvested trees is assumed to enter the litter pool. The harvesting of stem wood is assumed to involve a conversion loss equivalent to 10% of standing stem volume, which also enters the litter pool. It is difficult to make robust assumptions about the fate of branch wood and foliage at time of harvesting. In many situations, this material will be left on-site to deteriorate and decay. Sometimes it is possible that branch wood remaining after clearfelling may be deliberately burned. There has also been an increasing interest in active harvesting of branch wood (or at least some proportion of it) to supply biomass to the Energy sector. However, currently, such practice remains very limited. For this inventory the assumption has been made that no branch wood is harvested but is left to degrade and decay on site as part of the litter pool.

The branch ATR was fixed at 4% in accordance to Canadian forest carbon accounting model CBM-CFS (Kurz et al., 2009). Deciduous species foliage turnover is assumed to be 100% (Kurz et al., 2009; Ľupek et al., 2015). Conifer species foliage ATRs were obtained by referring to relevant scientific literature. If insufficient empirical literature and data was available the species were mapped to an allometrically similar species. Coarse root annual turnover was assumed to be 2% as in the CBM-CFS (Kurz et al., 2009; Kurz and Beukema, 1996; Li et al., 2003). Fine root ATRs were mapped from the available scientific literature and the UK specific datasets provided by Vanguelova (pers. com.). The UK ATRs for fine roots were derived from Kielder forest for Sitka spruce and Alice Holt forest for oak. Lastly, root exudate ATR was set to 160% of fine root dry biomass, the upper quartile of reported exudate mass from grassland was adopted (Jones et al., 2009), because of limited understanding about forest rhizodeposition. Aboveground shed litter, foliage and branches, are accumulated in a litter layer and after partial degradation passed to the Fermenting (F) layer. Residues that are left after thinning or felling can be set to enter a litter layer. If the crop is not a forest, it is assumed that the litter and F layers are zero. The litter layer decomposition is modelled using modified ForClim-D model version (Liski et al., 2002; Perruchoud et al., 1999). Below ground litter is not included in this simulation, while the annual transfer rates are applied to foliage (C_f) and branch (C_b) litter biomass. They are expressed as a proportion relocated annually.

Branch and foliage litter transfer are set according to the model proposed by Liski et al. (2002). The transferred biomass is pooled and degraded by a fixed constant of 0.5, which is the average of constants given in the Liski et al. (2002) study

The new CARBINE Soil Carbon Accounting model (CARBINE-SCA; **Figure A 3.2**), is based on a simplified version of the ECOSSE model (Smith *et al.*, 2011), coupled with a litter decomposition model derived from the ForClim-D model (Perruchoud *et al.*, 1999; Liski *et al.*, 2002). Above-ground turnover of material such as foliage, branches and dead stemwood enters the litter pool, which is then broken down to F-material (Fermenting) as a function of temperature and rainfall, releasing CO₂. Within the soil, a number of layers exist, each with its own set of texture (Sand, Silt, Clay) characteristics. Carbon from decayed litter, dead roots, and root exudates enters each layer and is assigned to four active pools; resistant plant material (RPM), readily decomposable plant material (DPM), biological material (BIO) and humic material (HUM). A proportion of organic carbon is also assumed to be inert, and unavailable for further activity. The active pools undergo decomposition and transference, releasing CO₂. Decomposition (aerobic and anaerobic) within each pool and layer is influenced by response functions to water saturation in the soil, temperature, pH, and the presence (or not) of plant cover on the soil surface. The availability of water within each layer, and the level of saturation are largely defined from soil texture following Saxton and Rawls (2006) coupled with inputs from rainfall, (or drainage) and removal of water through evapotranspiration. In any soil layer, water above field capacity can drain to lower soil layers, complete with any dissolved organic carbon

(DOC). The rates of potential decomposition of each carbon pool and the response functions follow ECOSSE (Smith et al., 2011).

New carbon input to the soil arises from four sources:


- Recently dead root material (according to a rooting profile depth),
- Transfer from the F-material arising from the decomposition of above-ground litter,
- Secretions and exudates from the roots,
- DOC; this carbon can become available to the biological pool and enter the 'reactive material cycle'.

Turnover rates for mortality of tree components (roots, foliage etc.) are species dependent and obtained from scientific literature (see **Table A 3.4.1** for example). A full description of the model will be presented in a separate technical report.

An improved version of the soil sub-model was implemented for the 1990-2016 inventory. This included work on parameterisation of litter input from ground flora and other non-forest vegetation, assuming a decrease in the contribution of non-tree litter from that assumed in Ecosse for pasture to zero contribution at canopy closure.

A more comprehensive description of the soil sub-model will be described in a technical report.

Figure A 3.2 The CARBINE-SCA model



The harvested wood products sub-model represents wood products as long-lived and short-lived sawn timber, particleboard, paper and fuel (see also **Section 6.8**). Carbon in harvested

stemwood is allocated to these wood product categories using an assortment forecasting model that accounts for variation in product out-turn due to tree species and tree size class distribution at time of harvest (Rollinson and Gay, 1983). Wood products in primary use are assumed to decay over time with no account taken of carbon stocks in landfill or greenhouse gas emissions (due to wood products) from landfill (these are taken into account in the Waste sector).

A 3.4.1.2 Forest activity data: management

Information from the Sub-compartment database (SCDB, the Forestry Commission database of information on the growth rate and management of the Forestry Commission (FC) and Natural Resources Wales (NRW) forest estate) was used to create a distribution of species and yield class (an indication of growth rate) for the FC/NRW forest estate. For the non-FC/NRW forest estate information from the NFI survey of woodlands was analysed to estimate yield class and species by age class, and scaled to represent the whole private forest estate. Data from the Forestry Commission's new planting and wood production statistics were used to assign the areas in an age class to individual years, either as areas restocked or areas newly planted.

Management of forests is represented as one of four options: Clearfell with thinnings, clearfell without thinnings, managed but not clearfelled, and not used for timber production. For the clearfell forests restocking occurs after the rotation period. For non-clearfell productive woodlands it is assumed there is a 30 year overlap of restocking and non-restocked trees. The area of land felled each year was estimated from the wood production statistics separately for both FC/NRW and private forests. The rotation periods for forests were estimated based on information on the intended management of the FC/NRW estate. This analysis gave a target rotation period for each modelled species and yield class.

The actual rotations historically applied to the forest estate are unknown and for the private forests the area of woodland used for timber production is also unknown. In order to match production, given the age class distribution of the forest, an algorithm was implemented to adjust the assumed rotations and the percentage of private sector woodland not used for timber production. This algorithm adjusts these assumptions in order to match the modelled wood production with the timber production statistics separately for the FC/NRW and private forests. It was assumed that the forests would be felled evenly over a period +/-7 years from the target rotation period. A comprehensive description of this algorithm will be presented in a separate technical report.

Information on the management of privately owned forests that is used to inform the inventory estimates, as well as a description of how forest land AD (forest land remaining forest land; land converted to forest land) are derived is included in the National Forest Accounting Plan, which can be found at:

<https://www.gov.uk/government/publications/uk-national-forestry-accounting-plan-2021-to-2025>

A 3.4.1.3 Forestry activity data: historical and current afforestation rates

Irrespective of species assumptions, the variation in CO₂ removals from 1990 to the present is determined by the afforestation rate in earlier decades, the effect this has on the age structure in the present forest estate, and hence the average growth rate. Afforestation is assumed to occur on ground that has not been wooded for many decades, based on the assumption that if it had previously been woodland it would be in the restocking statistics rather than the new planting statistics as a result of the regulatory framework that applies to forestry in the UK.

A comparison of historical forest census data and the historical annual planting rates has been undertaken. Forest censuses were taken in 1924, 1947, 1965, 1980 and the late 1990s. The

latest census (National Forest Inventory) has only just been completed. The comparison of data sources showed that discrepancies in annual planting rates and inferred planting/establishment date (from woodland age in the forest census) are due to restocking of older (pre-1920) woodland areas and variations in the harvesting rotations. However, there is also evidence of shortened conifer rotations in some decades and transfer of woodland between broadleaved categories (e.g. between coppice and high forest). It is difficult to incorporate non-standard management in older conifer forests and broadleaved forests into the Inventory because it is not known whether these forests are on their first rotation or subsequent rotations (which would affect carbon stock changes, particularly in soils). The area of afforestation in a given year is predicted based on applying the yearly distribution from the new planting and restocking statistics to the age class inventory. Age classes prior to the availability of new planting statistics are assigned evenly to individual years. For this inventory submission the assumption was made that we can estimate the area felled for recent years based on the timber production in the year of felling. It is assumed that woodland felled is immediately restocked. As we have an estimate of the area restocked for these years, the remainder of the area for each year was assumed to be restocking or natural regeneration. For years prior to the timber production statistics (i.e. prior to 1976), an estimated ratio between restocking and afforestation was used based on the earliest data. For restocked woodland the forest area was assumed to have been restocked twice and had been managed in the same fashion and on the same rotation.

The planting data used as input to the CARBINE model come from national planting statistics from 1921 to the present (provided by the Forestry Commission) for England, Scotland and Wales and from 1900 to the present (provided by the Northern Ireland Forest Service). For England, Scotland and Wales estimates of area of woodland by species, yield class and broad age class came from analysis of the NFI (for private woodland) and the SCDB (for FC/NRW woodland).

The NFI provides woodland statistics for Great Britain, (England, Wales and Scotland), broken down by region. It comprises a digital woodland map based on comprehensive aerial photography and a field survey using 15,000 one-hectare sample squares. The digital map and field survey cover all woodland areas down to 0.5 hectares. An initial digital woodland map was published in spring 2011. The NFI woodland field survey provides direct assessments of woodland growing stock including species composition, stand structure, tree age (distribution) productivity indices, numbers of trees, and diameter and height distribution. Standing biomass (and carbon) in trees, including above and below ground biomass, can be derived from these assessments using GB-specific conversion factors and allometric equations. A complete 5-year cycle of ground survey has now been completed. NFI data do not allow the carbon stocks of deadwood or litter to be estimated. The NFI has been supplemented by an assessment of the area of small woods (woodland between 0.1 ha and 0.5 ha) to align with the minimum woodland area for UNFCCC reporting as set out in CMP.7 (Forestry Commission, 2017). The analysis of small woods area included no characterisation of the resource. Since there is currently no information on the age-distribution of the area of small woods, it was assumed to have established evenly between 1900 and 1970.

The NFI uses a lower integral open space threshold of 0.5 ha (as opposed to 1 ha), which requires a downward adjustment to areas. However, the main differences in 2010 GB woodland cover between the NFI (2982 kha) and previous estimates (2757 kha, Forestry Statistics 2010) arise from identified errors in the previous woodland survey, particularly the under-estimate of woodland areas between 0.5 and 2 hectares. Estimates of woodland loss have been assessed, which affect the total estimated woodland area in the GHGI (but are not yet reflected in the national Forestry Statistics).

We assumed that the NFI survey gives a distribution of all the non FC/NRW forest area for a base year of 2011, and the SCDB gives a distribution of all the FC/NRW forest area for a base year of 2014.

The main NFI survey includes areas of woodland >0.5 ha. An adjustment was made to the areas of woodland to account for woods between 0.1 ha and 0.5 ha. For England and Wales, the estimates are derived from a calibration of tree cover plotted in the National Tree Map (NTM) product across England and Wales⁶, using a comparison of manual photographic interpretation with the NTM product within a sample of 1 km square tiles. For Scotland, the estimates are derived from a direct evaluation of polygons in the map constructed for the Native Woodlands of Scotland Survey (NWSS)⁷, which mapped all woodland polygons in Scotland down to 0.1 hectares in size by photographic interpretation. The areas of small woods used in this inventory were based on data published in 2017 by the Forestry Commission in the report “Tree cover outside woodland in Great Britain”⁸.

An algorithm was used to obtain the area of woodland afforested each year by removing the area of felling from the age class distribution. The species were then allocated to this “residual distribution” by starting in the base year and allocating the shortest rotations first. The planting years for all restocked woodland are assigned by the algorithm to give two rotations of the same length as the assigned rotation, and are thus notional. This approach was undertaken to spin up the model in terms of soil and litter in order to reach a state consistent with land that has been forest for a long period. This algorithm will be described in detail in the same technical report as the description of allocation of the management of forests.

Conifer planting on organic soil is a subset of total conifer planting. All broadleaf planting is assumed to be on non-organic soil. As explained above, the planting rates given in **Table A 3.4.2** are derived from administrative records, information on forest age class distribution from NFI field assessments and interim assumptions about the age distribution of ‘small woods’. The planting rates given in **Table A 3.4.2** are therefore significantly different to those reported as official planting statistics supported by grant-aid. The afforestation rates for each planting type in the UK have been calculated from the data and are shown in **Table A 3.4.2**.

Table A 3.4.2 Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 based on estimates of woodland area by age from the NFI and administrative records.

| Period | Planting rate (kha annum ⁻¹) | | |
|-----------|--|--------------------------|-------------|
| | Conifers on all soil types | Conifers on organic soil | Broadleaves |
| 1501-1600 | 0.00 | 0.00 | 0.02 |
| 1601-1700 | 0.06 | 0.00 | 0.68 |
| 1701-1750 | 0.14 | 0.00 | 2.31 |

⁶ <http://www.bluesky-world.com/national-tree-map>

⁷ <http://scotland.forestry.gov.uk/supporting/strategy-policy-guidance/native-woodland-survey-of-scotland-nwss>

⁸ ⁸ <https://www.forestresearch.gov.uk/tools-and-resources/national-forest-inventory/what-our-woodlands-and-tree-cover-outside-woodlands-are-like-today-8211-nfi-inventory-reports-and-woodland-map-reports/>

| Period | Planting rate (kha annum ⁻¹) | | |
|-----------|--|--------------------------|-------------|
| | Conifers on all soil types | Conifers on organic soil | Broadleaves |
| 1751-1800 | 0.42 | 0.00 | 1.69 |
| 1801-1850 | 1.10 | 0.00 | 0.91 |
| 1851-1900 | 5.67 | 0.02 | 0.86 |
| 1901-1910 | 4.81 | 0.65 | 8.94 |
| 1911-1920 | 2.71 | 0.32 | 11.57 |
| 1921-1930 | 2.89 | 0.30 | 12.20 |
| 1931-1940 | 4.54 | 0.47 | 12.92 |
| 1941-1950 | 7.83 | 1.06 | 15.64 |
| 1951-1960 | 16.82 | 2.65 | 16.64 |
| 1961-1970 | 24.11 | 4.61 | 18.36 |
| 1971-1980 | 24.33 | 5.44 | 13.46 |
| 1981-1990 | 20.38 | 4.79 | 15.41 |
| 1991 | 12.41 | 2.93 | 11.97 |
| 1992 | 10.83 | 2.60 | 13.65 |
| 1993 | 8.58 | 2.06 | 16.74 |
| 1994 | 9.21 | 2.20 | 17.67 |
| 1995 | 8.23 | 1.98 | 15.17 |
| 1996 | 7.94 | 1.85 | 14.72 |
| 1997 | 7.39 | 1.64 | 16.14 |
| 1998 | 6.83 | 1.48 | 15.94 |
| 1999 | 6.63 | 1.39 | 16.21 |
| 2000 | 5.36 | 1.08 | 17.95 |
| 2001 | 6.01 | 1.15 | 14.41 |
| 2002 | 5.67 | 1.06 | 13.80 |
| 2003 | 5.08 | 0.93 | 12.43 |
| 2004 | 7.81 | 1.20 | 12.10 |
| 2005 | 6.46 | 0.95 | 10.91 |
| 2006 | 7.16 | 1.00 | 11.73 |
| 2007 | 6.42 | 0.87 | 9.44 |
| 2008 | 6.13 | 0.75 | 7.31 |
| 2009 | 5.99 | 0.66 | 6.55 |
| 2010 | 7.15 | 0.74 | 8.82 |
| 2011 | 9.66 | 0.93 | 11.98 |
| 2012 | 6.14 | 0.54 | 12.66 |
| 2013 | 6.12 | 0.54 | 11.66 |
| 2014 | 6.48 | 0.58 | 8.63 |
| 2015 | 2.17 | 0.23 | 4.73 |
| 2016 | 3.23 | 0.34 | 3.16 |
| 2017 | 4.85 | 0.52 | 3.65 |

The proportion of forest planting on mineral and organic soils was re-assessed in 2012, as part of the work to estimate N₂O emissions due to drainage on forest soils (Yamulki et al. 2012).

A 3.4.1.4 Allocation of CARBINE outputs to UNFCCC inventory sub-categories

The CARBINE model output was post-processed using the IPCC default 20-year transition period for Land converted to Forest to move into the Forest remaining Forest category. The area within the Land converted to Forest Land sub-category is split between cropland, pasture grassland, semi-natural grassland, settlement and other areas. Pasture grassland and semi-natural grassland are combined for Grassland reporting in the CRF. This split is based on the relative proportions of historical land use change from these categories to forest. The proportions for each country change over time because the 20-year transition period has a different start date for each reporting year.

The area and carbon stock changes in the Forest remaining Forest category are adjusted to take account of losses of forest converted to other land use categories (deforestation), as these losses, in their entirety, are not reflected in the statistics published by the Forestry Commission. Implied carbon stock changes per unit area are calculated using the unadjusted forest area and carbon stock changes. The forest area is then adjusted to reflect losses due to forest conversion and multiplied by the implied carbon stock change to obtain the adjusted carbon stock change.

The CARBINE model has not yet been implemented for forest in the Isle of Man and Guernsey (Crown Dependencies of the UK) and instead the C-Flow model is used as it was in previous submissions (UK Greenhouse Gas Inventory, 1990-2011, Annex 3.6). Transition to the use of CARBINE for the Overseas Territories and Crown Dependencies has been delayed to allow for further development and documentation of CARBINE.

A 3.4.1.5 Nitrogen fertilization of forest land

Nitrogen fertilization of forest land is assumed to occur only when absolutely necessary, i.e. new planting on 'poor' soils (mining spoil, impoverished brown field sites, or upland organic soils). In terms of the inventory, this means that N fertilisation is assumed for Settlement converted to Forest land and Grassland converted to Forest Land on organic soils. The areas of new planting with these conditions were taken from the same dataset used in the CARBINE model (see **Table A 3.4.2**) for 4.A.2. Land converted to Forest land.

Where fertilisation occurs, an application rate of 150 kg N ha⁻¹ is assumed based on Forestry Commission fertilisation guidelines (Taylor 1991). The guidelines recommend applying fertiliser on a three-year cycle until canopy closure (at approximately 10 years), but this is thought to be rather high (Skiba 2007) and unlikely to occur in reality, so two applications are adopted as a compromise. These applications occur in year 1 and year 4 after planting. The N₂O emission factor for applied nitrogen fertiliser is the default value of 1% used in the IPCC 2006 Guidelines. Emissions of N₂O from N fertilisation of forests are estimated using a Tier 1 methodology and IPCC default emission factors. The emissions have fallen since 1990 due to reduced rates of new forest planting. A GWP of 298 for N₂O is used.

A 3.4.1.6 Emissions from drainage on forest soils

Work on developing this method was undertaken by Forest Research in 2012 (Yamulki *et al.* 2012), using new GIS data on forest planting in England, Wales and Scotland. Comparable data were not available for Northern Ireland. This method was described in the 1990-2012 National Inventory Report.

The calculations use the same data on forest planting on mineral and organic soils as are used by the CARBINE model for the calculation of carbon stock changes. It is assumed that only forests planted since 1920 have been drained. The areas of forest planted on mineral soil, nutrient-rich organic soil, nutrient-poor organic soils, nutrient-rich organo-mineral soils and

nutrient-poor organo-mineral soils are estimated based on the proportion of forest cover on different soil types (Yamulki *et al.* 2012), adjusted by the amount of forest planted since 1920. The area of forest on mineral soil is adjusted further by splitting it between free-draining mineral soils and imperfectly draining (easily waterlogged) mineral soils, which require artificial drainage (based on the current guidance and policy for forest operations and management). The proportion of mineral soils requiring artificial drainage is: 34% in England, 24% in Scotland, 3% in Wales, 68% in Northern Ireland and 26% in the UK as a whole. We assumed all forest on organic and organo-mineral soils is cultivated prior to planting and therefore effectively drained.

N₂O emissions are estimated using the Tier 1 methodology and the IPCC default emission factors for drained mineral (0.06 kg N₂O-N ha⁻¹ yr⁻¹), nutrient-rich organic (0.6 kg N₂O-N ha⁻¹ yr⁻¹) and nutrient-poor organic soils (0.1 kg N₂O-N ha⁻¹ yr⁻¹) (IPCC, 2006).

A 3.4.2 Land Use Change and Soils (4B, 4C, 4E)

Changes in soil carbon due to land use change are modelled with a dynamic model of carbon stock change which is driven by matrices of change calculated from land surveys.

A 3.4.2.1 Land Use Change Matrices

For Great Britain (England, Scotland and Wales), matrices from the Monitoring Landscape Change (MLC) data from 1947 & 1980 (MLC 1986) and the Countryside Surveys (CS) of 1984, 1990, 1998 (Haines-Young *et al.* 2000) and 2007 (Smart *et al.* 2009) are used.

In Northern Ireland, matrices were calculated from the Northern Ireland Countryside Surveys of 1990, 1998 (Cooper and McCann 2002) and 2007 (Cooper, McCann and Rogers 2009). The only data available for Northern Ireland pre-1990 are land use areas from The Agricultural Census and The Forest Service (Cruickshank and Tomlinson 2000). Matrices of land use change were estimated for 1970-79 and 1980-89 using area data. The relationship between the matrix of land use transitions and initial area from recent Countryside Surveys is assumed to be the same as the relationship between the matrix and area data for each of the earlier periods – 1970-79 and 1980-89. The matrices developed in this approach were used to extrapolate areas of land use transition back to 1950 to match the start year in the rest of the UK.

The AFOLU Guidance (IPCC 2006) recommends use of six types of land for descriptive purposes: Forest, Grassland, Cropland, Settlements, Wetlands and Other Land. Areas undergoing active commercial peat extraction and areas of inland water and flooded land are reported under Wetlands in the current inventory, and the remaining land in the UK has been placed into the five other types. The more detailed habitats for the two surveys in Great Britain were combined as shown in **Table A 3.4.3** for the Monitoring Landscape Change dataset and **Table A 3.4.4** for the Countryside Survey Broad Habitats (Jackson, 2000).

Table A 3.4.3 Grouping of MLC land cover types for soil carbon change modelling

| CROPLAND | GRASSLAND | FORESTLAND | SETTLEMENTS (URBAN) | OTHER |
|---------------|---------------------|------------------|---------------------|---------------|
| Crops | Upland heath | Broadleaved wood | Built up | Bare rock |
| Market garden | Upland smooth grass | Conifer wood | Urban open | Sand/shingle |
| | Upland coarse grass | Mixed wood | Transport | Inland water |
| | Blanket bog | Orchards | Mineral workings | Coastal water |
| | Bracken | | Derelict | |
| | Lowland rough grass | | | |
| | Lowland heather | | | |

| CROPLAND | GRASSLAND | FORESTLAND | SETTLEMENTS (URBAN) | OTHER |
|----------|---------------------|------------|---------------------|-------|
| | Gorse | | | |
| | Neglected grassland | | | |
| | Marsh | | | |
| | Improved grassland | | | |
| | Rough pasture | | | |
| | Peat bog | | | |
| | Fresh Marsh | | | |
| | Salt Marsh | | | |

Table A 3.4.4 Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling

| CROPLAND | GRASSLAND | FORESTLAND | SETTLEMENTS (URBAN) | OTHER |
|-------------------------|-------------------------|-------------------------------------|------------------------------|--------------------------------|
| Arable and horticulture | Improved grassland | Broadleaved, mixed and yew woodland | Built up areas and gardens | Inland rock |
| | Neutral grassland | Coniferous woodland | Unsurveyed urban land | Supra littoral rock |
| | Calcareous grassland | | Boundary and linear features | Littoral rock |
| | Acid grassland | | | Standing open water and canals |
| | Bracken | | | Rivers and streams |
| | Dwarf shrub heath | | | Sea |
| | Fen, marsh, swamp | | | |
| | Bogs | | | |
| | Montane | | | |
| | Supra littoral sediment | | | |
| | Littoral sediment | | | |

The area data used between 1947 and the latest inventory year are shown in **Table A 3.4.5** and **Table A 3.4.6**.

The land use change data over the different periods were used to estimate annual changes by assuming that the rates of change were uniform across the period. The full set of annual land use change matrices 1990-latest inventory year are provided in in **Chapter 6** of the main NIR report, **Section 6.1.1** for the UK and **Section 6.9.2** for the Overseas Territories and Crown Dependencies.

Table A 3.4.5 Sources of land use change data used to estimate changes in soil carbon in Great Britain for different periods.

| Year or Period | Method | Change matrix data |
|----------------|---------------------|--------------------|
| 1950-1979 | Measured LUC matrix | MLC 1947->MLC1980 |
| 1980 - 1984 | Interpolated | CS1984->CS1990 |
| 1984 - 1989 | Measured LUC matrix | CS1984->CS1990 |
| 1990 - 1998 | Measured LUC matrix | CS1990->CS1998 |
| 1999-2007 | Measured LUC matrix | CS1998->CS2007 |

| Year or Period | Method | Change matrix data |
|------------------|---------------------|--------------------|
| 2008-latest year | <i>Extrapolated</i> | CS1998->CS2007 |

Table A 3.4.6 Sources of land use change data used to estimate changes in soil carbon in Northern Ireland for different periods.


| Year or Period | Method | Change matrix data |
|------------------|---------------------------------|--------------------|
| 1950 – 1969 | Extrapolation and ratio method | NICS1990->NICS1998 |
| 1970 – 1989 | Land use areas and ratio method | NICS1990->NICS1998 |
| 1990 – 1998 | Measured LUC matrix | NICS1990->NICS1998 |
| 1999-2007 | Measured LUC matrix | NICS1998->NICS2007 |
| 2008-latest year | <i>Extrapolated</i> | NICS1998->NICS2007 |

NICS = Northern Ireland Countryside Survey

The transitions between habitat types in the Countryside Surveys for the latest survey (2007) were calculated with Geographical Information System software (ArcGIS). 544 Countryside Survey squares of Great Britain were identified that coincided between the 1998 and 2007 surveys. Survey square locations are confidential. For each coincident square, the area that changed from one habitat type in 1998 to another in 2007 was calculated. There are 47 broad habitats described by the Countryside Survey. Individual surveyed squares contain a subset of these habitats and changes between habitats are called transitions. Each coincident survey square also has a 'land class' assigned to it that does not change between survey years. There are currently 45 land classes in the Land Classification of Great Britain. Land classes represent the stratification of environments across the UK. A simplified picture of the stratification is shown in **Figure A 3.3**. Northern Ireland is treated as a single uniform unit as its smaller area means that it does not display the climatic or elevation variations evident across the rest of the UK.

Transitions between broad habitats were grouped by land class. The ratio of the total area of each land class to the total area sampled within each land class is calculated so that the transitions can be up-scaled to the land class areas. Transitions can then be extracted at various scales i.e. UK or Devolved Authorities scale or 20 km by 20 km squares. These scales are required by the soil carbon and non-forest biomass models.

Figure A 3.3 Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes).



A 3.4.2.2 Soils modelling

A database of soil carbon density for the UK (Milne & Brown 1997, Cruickshank *et al.* 1998, Bradley *et al.* 2005) is used in conjunction with the land use change matrices. There are three soil surveys covering England and Wales, Scotland and Northern Ireland. The field data, soil classifications and laboratory methods for these surveys have been harmonized to reduce uncertainty in the final joint database. The depth of soil considered was also restricted to 1 m at maximum as part of this process. **Table A 3.4.7** shows total stock of soil carbon (1990) for different land types in the four devolved areas of the UK.

Table A 3.4.7 Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK

| Region Type | England | Scotland | Wales | N. Ireland | UK |
|-------------|---------|----------|-------|------------|-------|
| Forestland | 108 | 295 | 45 | 20 | 467 |
| Grassland | 995 | 2,349 | 283 | 242 | 3,870 |
| Cropland | 583 | 114 | 8 | 33 | 738 |
| Settlements | 54 | 10 | 3 | 1 | 69 |

| Region Type | England | Scotland | Wales | N. Ireland | UK |
|--------------|--------------|--------------|------------|------------|--------------|
| Other | 0 | 0 | 0 | 0 | - |
| TOTAL | 1,740 | 2,768 | 340 | 296 | 5,144 |

The dynamic model of carbon stock change requires the change in equilibrium carbon density from the initial to the final land use. The core equation describing changes in soil carbon with time for any land use transition is:

$$C_t = C_f - (C_f - C_0)e^{-kt}$$

where

C_t is carbon density at time t

C_0 is the assumed equilibrium carbon density initial land use

C_f is the assumed equilibrium carbon density after change to new land use

k is time constant of change

Differentiating this equation gives the flux f_t (emission or removal) per unit area:

$$f_t = k(C_f - C_0)e^{-kt}$$

This equation gives, for any inventory year, the land use change effects from any specific year in the past. If A_T is area in a particular land use transition in year T considered from 1950 onwards then total carbon lost or gained in an inventory year, e.g. 1990, is given by:

$$F_{1990} = \sum_{T=1950}^{t=1990} kA_T (C_f - C_0)(e^{-k(1990-T)})$$

This equation is used with k , A_T and $(C_f - C_0)$ chosen by Monte Carlo methods within ranges set by prior knowledge, e.g. literature, soil carbon database, agricultural census, LUC matrices.

In the model, the change in equilibrium carbon density from the initial to the final land use is calculated. These are calculated for each land use category as averages for Scotland, England, Wales and Northern Ireland. These averages are weighted by the area of Land Use Change occurring in four broad soil groups (organic, organo-mineral, mineral, unclassified) in order to account for the actual carbon density where change has occurred. In the UK land use change other than afforestation generally is assumed not to occur on organic soils. Changes in soil carbon stock on afforested land are modelled using the CARBINE model rather than the exponential loss model described above. Other areas of land use change on organic soils are believed to be very small and are currently not separated out from change on mineral soils.

Hence mean soil carbon density change is calculated as:

$$\bar{C}_{ijc} = \frac{\sum_{s=1}^6 (C_{sijc} L_{sijc})}{\sum_{s=1}^6 L_{sijc}}$$

This is the weighted mean, for each country, of change in equilibrium soil carbon when land use changes, where:

i = initial land use (Forestland, Grassland, Cropland, Settlements)

j = new land use (Forestland, Grassland, Cropland, Settlements)

c = country (Scotland, England, N. Ireland & Wales)

s = soil group (organic, organo-mineral, mineral, unclassified)

C_{sijc} is change in equilibrium soil carbon for a specific land use transition

The land use data (1990 to 1998) is used in the weighting. The averages calculated are presented in **Table A 3.4.8 - Table A 3.4.11**.

Table A 3.4.8 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in England

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | 0 | 25 | 32 | 83 |
| Grassland | -21 | 0 | 23 | 79 |
| Cropland | -31 | -23 | 0 | 52 |
| Settlements | -87 | -76 | -54 | 0 |

Table A 3.4.9 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Scotland

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | 0 | 47 | 158 | 246 |
| Grassland | -52 | 0 | 88 | 189 |
| Cropland | -165 | -90 | 0 | 96 |
| Settlements | -253 | -187 | -67 | 0 |

Table A 3.4.10 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Wales

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | 0 | 23 | 57 | 114 |
| Grassland | -18 | 0 | 36 | 101 |
| Cropland | -53 | -38 | 0 | 48 |
| Settlements | -110 | -95 | -73 | 0 |

Table A 3.4.11 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Northern Ireland

| From To | Forestland | Grassland | Cropland | Settlements |
|------------|------------|-----------|----------|-------------|
| Forestland | 0 | 94 | 168 | 244 |
| Grassland | -94 | 0 | 74 | 150 |

| From To | Forestland | Grassland | Cropland | Settlements |
|--------------------|------------|-----------|----------|-------------|
| Cropland | -168 | -74 | 0 | 76 |
| Settlements | -244 | -150 | -76 | 0 |

The rate of loss or gain of carbon is dependent on the type of land use transition (Table A 3.12). For transitions where carbon is lost e.g. transition from Grassland to Cropland, a 'fast' rate is applied whilst a transition that gains carbon occurs much more slowly. A literature search for information on measured rates of changes of soil carbon due to land use was carried out and ranges of possible times for completion of different transitions were selected, in combination with modelling and expert judgement (Milne and Brown, 1999; Ashman, *et al*, 2000, Salway *et al*, 2001). These are shown in Table A 3.13.

Table A 3.4.12 Rates of change of soil carbon for land use change transitions

| | | Initial | | | |
|-------|------------|-------------|-------------|-------------|-------------|
| | | Forestland | Grassland | Cropland | Settlement |
| Final | Forestland | | <i>slow</i> | <i>slow</i> | <i>slow</i> |
| | Grassland | <i>fast</i> | | <i>slow</i> | <i>slow</i> |
| | Cropland | <i>fast</i> | <i>fast</i> | | <i>slow</i> |
| | Settlement | <i>fast</i> | <i>fast</i> | <i>fast</i> | |

("Fast" & "Slow" refer to 99% of change occurring in times shown in Table A 3.4.13)

Table A 3.4.13 Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W)

| | Low (years) | High (years) |
|------------------------------|-------------|--------------|
| Carbon loss ("fast") E, S, W | 50 | 150 |
| Carbon gain ("slow") E, W | 100 | 300 |
| Carbon gain ("slow") S | 300 | 750 |

Changes in soil carbon from equilibrium to equilibrium ($C_t - C_0$) were assumed to fall within ranges based on 2005 database values for each transition (Bradley *et al*, 2005) and the uncertainty indicated by this source (up to $\pm 11\%$ of mean). The areas of land use change for each transition were assumed to fall in a range of uncertainty of $\pm 30\%$ of the mean.

A Monte Carlo approach is used to vary the rate of change, the area activity data and the values for soil carbon equilibrium (under initial and final land use) for all countries in the UK. The model of change was run 1000 times using parameters selected from within the ranges described above. The mean carbon flux for each region resulting from this imposed random variation is reported as the estimate for the Inventory. An adjustment was made to these calculations for each country to remove increases in soil carbon due to afforestation, as the CARBINE model provides a better estimate of these fluxes in the Land Converted to Forest Land category. Variations from year to year in the reported net emissions reflect the trend in land use change as described by the matrices of change.

A 3.4.2.2.1 Change in soil carbon stock due to cropland management activities

Change in soil carbon stocks due to cropland management activities is estimated using the methodology developed in Defra project SP1113 (Moxley *et al*, 2014a) which reviewed UK relevant literature on the effects of cropland management practices on soil carbon stocks and attempted to model UK specific emission factors.

Increases in inputs of fertiliser, manure and crop residues were found to increase soil carbon stocks of tillage land, but changes in the tillage regime from conventional tillage to reduced or zero tillage were found to have no significant effect in a UK context.


Using this methodology, tillage crops are divided into Medium and Low residue groups based on the data on total crop biomass. Where land receives inputs of fertiliser or manure the inputs moved up a class (e.g. cropland producing a Low residue crop which receives manure is considered to receive Medium inputs, while land producing a Medium residue crop which received manure inputs is considered to receive High inputs). If crop residues are removed from land the input level drops. A decision tree for assessing the effect of cropland management on soil carbon stocks is shown in **Figure A 3.4**.

For most cropland management activities there were insufficient UK field data to develop reliable Tier 2 stock change factors, and so Tier 1 factors have been used (for manure and residue inputs, and for differentiating perennial crops, annual crops and set-aside). However, for tillage reduction both a literature review and modelling work suggested that it did not have a significant effect on soil carbon stocks, and that the Tier 1 stock change factors over-estimated its effect under UK conditions. Therefore, a stock change factor of 1 has been used for tillage reduction.

As changes in soil carbon stocks due to changes in cropland management are smaller than changes due to land use change the IPCC default transition time of 20 years is used.

Data on the areas under the main crop types is obtained from the annual June Agricultural Censuses carried out by each UK administration (Defra; Welsh Government; Scottish Government; DAERA). Data on the areas of cropland receiving inputs of manure, fertiliser and crop residues is obtained from the annual British Survey of Fertiliser Practice (Defra).

Figure A 3.4 Decision tree for assessing the effects of cropland management activities on soil carbon stocks.



A 3.4.2.2.2 Change in soil carbon stock due to grassland management activities

Defra project SP1113 attempted to develop a methodology to allow reporting of changes in soil carbon stocks resulting from grassland management activities. There are reasonable data on the effects of management practices such as liming, reseeding and drainage on improved grassland on mineral soils. However, there are few data on the effect of many management practices if applied to semi-natural grassland or those on organo-mineral or organic soils where

there is a risk that more intensive management could increase carbon losses. As semi-natural grassland makes up a large proportion of grassland in the UK the lack of field data makes it impossible to reliably report changes in soil carbon stocks from grassland management activities. A research project commissioned by BEIS is currently underway to fill this knowledge gap; reporting on the effect of grassland management on soil carbon will be included in the inventory when new data from this project become available.

A 3.4.2.3 Future development

A new vector-based approach to tracking land use change (Levy et al. 2017) has been developed using data from the CORINE land cover map. The approach can be used to produce a set of 100 x 100 m resolution maps, where each 100 m square has an associated vector of land use over time. The maps can be aggregated into a set of distinct representative vectors with their corresponding areas. This has the potential to improve the modelling of change in soil carbon stocks resulting from land use change compared to the current statistically based land use change matrices.


Further implementation of this approach is under discussion, awaiting the outcome of related projects on the potential of earth observation to track LUC in the UK and clarity on the long-term availability of relevant datasets. As there will not be a repeat of the Countryside Survey (last undertaken in 2007), alternative data sources must be used, such as CEH land cover maps (1990, 2000, 2007), the NFI map, annual agricultural survey data or the Integrated Administration and Control System (IACS) dataset (used to administer Common Agricultural Policy payments). These all have advantages and disadvantages in terms of their coverage, spatial and temporal resolution, long-term availability and/or processing requirements.

A 3.4.3 Changes in stocks of carbon in non-forest biomass due to management and land use change (4B2, 4C2, 4E2)

A 3.4.3.1 Change in biomass carbon stock due to change in Cropland and Grassland Management.

Change in Cropland biomass carbon stocks was assessed based on agricultural census data. Areas under different crop types were taken from annual agricultural census data and assigned on one of four categories: annual crops, orchard crops, shrubby perennial crops and set aside and fallow (**Figure A 3.5**). Crop types reported in the agricultural census vary slightly for each administration. **Table A 3.4.14** shows how agricultural census crop types were grouped to assess biomass carbon stocks.

Figure A 3.5 Crop type area in the UK 1989-2017






Table A 3.4.14 Aggregation of Agricultural Census crop types for estimating biomass carbon stock changes from Cropland Management

| Devolved Administration | Annual Crops | Orchard Crops | Shrubby perennial crops | Set Aside and Fallow |
|-------------------------|--|--|---|-----------------------|
| England | Cereals, Other arable not stockfeed, Crops for stockfeeding, Vegetables for Human Consumption | Orchard Fruit | Soft fruit, Hardy nursery stock, bulbs and flowers, Area under glass or plastic covered structures. | Uncropped land |
| Scotland | Cereals, Oilseed rape, Peas for combining, Beans for combining, Linseed, Potatoes, Crops for stockfeeding, Vegetables for human consumption, Other crops | Orchard fruit | Soft fruit | Fallow , Set Aside |
| Wales | Cereals, Other arable not for stockfeeding, Crops for stockfeeding, Salad and vegetables grown in the open, Total hardy crops | Commercial orchards, Other orchards | Glasshouse | Bare fallow |
| Northern Ireland | Cereals, Other arable not for stockfeeding, Vegetables | Fruit | Ornamentals | Fallow and set aside |

The areas under each aggregated crop type were multiplied by the biomass carbon stock of each crop type using the biomass carbon stock factors in **Table A 3.4.15**. These factors were generated from a literature review. (Moxley *et al.* 2014b).

Table A 3.4.15 Biomass stock factors for UK Cropland types

| Crop Type | Total biomass Carbon Stock t C/ha | Uncertainty t C/ha (95% CI) | Root: Shoot ratio |
|----------------------------|---|--------------------------------|---------------------------------|
| Annual | 5 | 1.2 | Assume no Below Ground Biomass. |
| Orchards | 10 | 6.75 | 0.24 |
| Shrubby perennial crops | 3.7 | 2.0 | Assume no Below Ground Biomass. |
| Set Aside and Fallow | 5 | 1 | 4.0 |

Biomass carbon stock change was assumed to occur in the year in which the change in crop type was reported. Cropland biomass stock changes resulting from land use change to or from Cropland were subtracted from the changes due to change in cropland management, as they are accounted for under land use change.

Change in Grassland biomass carbon stocks was assessed based on Countryside Survey data. Grassland was separated into shrubby, non-shrubby and unvegetated Grassland based on Countryside Survey Broad Habitat types. **Table A 3.4.16** shows which Broad Habitats were allocated to which Grassland type.

Table A 3.4.16 Aggregation of Countryside Survey Broad Habitats for estimating biomass carbon stock changes from Grassland Management

| Shrubby Grassland | Non-shrubby Grassland | Unvegetated Grassland |
|---|--|--|
| Dwarf Shrub Heath Bracken Montane | Improved Pasture Improved Pasture Neutral Grassland Calcerous Grassland Acid Grassland Bogs | Littoral sediment Supra littoral sediment |

The areas under each aggregated Grassland type were multiplied by the biomass carbon stock of each crop types using the biomass carbon stock factors in **Table A 3.4.17**. These factors were generated from literature reviews (Moxley *et al.* 2014b). Only biomass carbon stock changes resulting from change between shrubby and non-shrubby Grassland were considered, as changes to and from unvegetated littoral and supra-littoral sediments were considered unlikely.

Table A 3.4.17 Biomass stock factors for UK Grassland types

| Crop Type | Total biomass Carbon Stock t C/ha | Uncertainty t C/ha (95% CI) | Root: Shoot ratio |
|-----------------------|---|--------------------------------|-------------------|
| Non-shrubby Grassland | 2.8 | 1.5 | 4.0 |

| Crop Type | Total biomass Carbon Stock t C/ha | Uncertainty t C/ha (95% CI) | Root: Shoot ratio |
|-----------------------|---|--------------------------------|-------------------|
| Shrubby Grassland. | 10 | 3.6 | 0.53 |
| Unvegetated Grassland | 0 | 0 | 0 |
| Managed hedge | 34.86 | 68.75 | 0.3 |
| Unmanaged hedge | 175.3 | 476.6 | 0.3 |

Countryside Survey data are only collected on an approximately decadal basis. The annual stock change between survey years was estimated using linear interpolation. Biomass carbon stock change was assumed to occur in the year in which the change in Grassland type occurred. Grassland biomass stock changes resulting from land use change to or from Grassland were subtracted from the changes due to change in grassland management.

Change in Grassland biomass carbon stocks due to change in hedge length are included in the estimate of change in Grassland biomass carbon stock using Countryside Survey data on hedge length and condition. Hedges were divided into managed hedges which are trimmed to prevent the growth of large trees and unmanaged hedges which do not received routine maintenance. Unmanaged hedges do not fall within the UK's definition of Forest, but may contain isolated trees and may also have some gaps in them. The biomass carbon stocks of managed and unmanaged hedges are estimated as the median of UK-relevant values in published literature, based on a literature review commissioned by BEIS (Moxley et al. 2014b) supplemented with more recent data. Full details of these values and data sources are included the Grassland Management Biomass calculation workbook.

A 3.4.3.2 Change in biomass carbon stock due to land use change.

Changes in stocks of carbon in biomass due to land use change are based on the same area matrices used for estimating changes in carbon stocks in soils (see previous section). The average biomass carbon density for Cropland, Grassland and Settlement are shown in **Table A 3.4.18**: these were derived from the distribution and biomass densities of the different crop and grassland types in each country of the UK. For Settlements the biomass stocks from Milne and Brown (1997) and land cover data from the 2007 Land Cover Map was used to assess the proportion of gardens, pasture-type grass (including sports pitches, golf courses and parks) and urban (built over) area within areas identified as Settlements.

The average change in biomass carbon density for each country is shown in **Table A 3.4.19 - Table A 3.4.22**. Changes between these equilibrium biomass carbon densities were assumed to happen in a single year.

Table A 3.4.18 Mean biomass carbon stock densities, tC/ha

| Mean C stock tC/ha | Cropland | Grassland | Settlement |
|--------------------|----------|-----------|------------|
| England | 5.02 | 3.37 | 2.77 |
| Scotland | 5.00 | 4.16 | 2.91 |
| Wales | 5.03 | 3.61 | 2.81 |
| Northern Ireland | 5.14 | 2.93 | 2.64 |

Table A 3.4.19 Weighted average change in equilibrium biomass carbon density (kg m²) for changes between different land types in England

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | | | | |
| Grassland | | 0 | -1.66 | 0.60 |
| Cropland | | 1.66 | 0 | 2.25 |
| Settlements | | -0.60 | -2.25 | 0 |

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.20 Weighted average change in equilibrium biomass carbon density (kg m²) for changes between different land types in Scotland.

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | | | | |
| Grassland | | 0 | -0.83 | 1.25 |
| Cropland | | 0.83 | 0 | 2.08 |
| Settlements | | -1.25 | -2.08 | 0 |

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.21 Weighted average change in equilibrium biomass carbon density(kg m²) for changes between different land types in Wales.

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | | | | |
| Grassland | | 0 | -1.42 | 0.80 |
| Cropland | | 1.42 | 0 | 2.22 |
| Settlements | | -0.80 | -2.22 | 0 |

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.22 Weighted average change in equilibrium biomass carbon density (kg m²) for changes between different land types in Northern Ireland.

| From To | Forestland | Grassland | Cropland | Settlements |
|-------------|------------|-----------|----------|-------------|
| Forestland | | | | |
| Grassland | | 0 | -2.21 | 0.29 |
| Cropland | | 2.21 | 0 | 2.49 |
| Settlements | | -0.29 | -2.49 | 0 |

(Transitions to and from Forestland are considered elsewhere)

Living biomass carbon stocks and Dead Organic Matter (DOM) stocks on Forest Land are modelled using CARBINE and used to calculate changes in carbon stocks due to conversions to and from Forest Land. When land is deforested to another land use, it is assumed that all living biomass and DOM is either converted to Harvested Wood Products or burnt on site in the year in which deforestation takes place. Under KP-LULUCF reporting all HWP from Article 3.3 Deforestation are assumed to be instantaneously oxidised. Increase in biomass carbon and DOM stocks on afforested land is modelled in CARBINE. Full details of CARBINE modelling of carbon stocks on Forest Land are given in **Annex A 3.4.1.1**.

A 3.4.3.3 Future development

A new vector-based approach to tracking land use change (Levy et al. 2017) as described in **Section A 3.4.2.3**.

A 3.4.4 Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G)

Deforestation is an activity that cuts across LULUCF categories, affecting net emissions and removals in all the land use categories except 4D Wetlands. The process of land use change affects carbon stock changes in biomass and soil, and the woody material left after felling either moves into the harvested wood products pool or is assumed to be burnt on-site, resulting in immediate biomass burning emissions.

Levy and Milne (2004) discuss methods for estimating deforestation since 1990 using a number of data sources. Their approach of combining Forestry Commission felling licence data for rural areas with Ordnance Survey data for non-rural areas was expanded to include new sources of information and to improve coverage of all countries in the UK. Deforestation before 1990 (which contributes to soil carbon stock change from historical land use change) is estimated from the land use change matrices described in **Section A 3.4.2**.

A 3.4.4.1 Types of deforestation activity in the UK

In Great Britain, some activities that involve tree felling require permission from the Forestry Commission, in the form of a felling licence, or a felling application within the various forestry grant schemes. There is a presumption that the felled areas will be restocked with trees, usually by replanting but sometimes by natural regeneration. However, some licences are granted without the requirement to restock – so-called unconditional felling licences. A felling licence is required unless special conditions are met⁹.

Felling for urban development, with no requirement to restock, can be allowed under planning permission but only local planning authorities hold documentation for this. Since 2006, remotely sensed data used in the NFI has included this change, but prior to this, the need for collation of data from local authorities makes estimating the national total difficult. However, in England, the Ordnance Survey (the national mapping agency) makes an annual assessment of land use change from the data it collects for map updating and provides this assessment to the Department of Communities and Local Government (DCLG)¹⁰. DCLG provides an extract of this dataset, listing annual land use change from Forest to developed land uses (1990-2008 in the latest submission). This dataset comes from a continuous rolling survey programme, both on

⁹ <https://www.gov.uk/guidance/tree-felling-licence-when-you-need-to-apply>

¹⁰ <http://www.communities.gov.uk/planningandbuilding/planningbuilding/planningstatistics/landusechange/>

the ground and from aerial photography. The changes reported each year may have actually occurred in any of the preceding 1-5 years. The survey frequency varies among areas, and can be up to 10 years for moorland/mountain areas. Consequently, for pre-2006 deforestation to Settlement a five-year moving average is applied to the data to smooth out the between-year variation appropriately, to give a suitable estimate with annual resolution.

The Countryside Survey land use change matrix (**Section A 3.4.2**) gives estimates of forest conversion to other land use categories for all countries in the UK for 1990-1998 and 1999-2007. There are known issues with Countryside Survey over-estimating the extent of Forest conversion compared with the extent estimated by the Forestry Commission. Therefore, Forest Commission data is used for Forest areas and the areas of other land uses estimated by Countryside Survey are adjusted to account for this. This is due to differences in Forest definitions, amongst other causes.

A 3.4.4.2 Compilation of activity datasets

For 1990-1999 the deforestation activity dataset is compiled from the felling licence and DCLG datasets as far as possible, using Countryside Survey (CS) data to fill gaps in the time series, to estimate deforestation in Northern Ireland (for which no direct data are available) and to estimate the conversion to different land use categories. The DCLG data are used to estimate the area of Forest Land converted to Settlement (4.E.2.1). The unconditional felling licence data are used to estimate the area of Forest Land converted to Cropland (4.B.2.1) and of Forest Land converted to Grassland (4.C.2.1). Only England has any post-1990 forest to cropland conversion: the estimated areas in Scotland, Wales and Northern Ireland are so small that they are thought to be due to survey classification error rather than genuine land use change.

The CS data are used to estimate the relative split of Forest conversion between Grassland, Cropland and Settlements (**Table A 3.4.23**), using other known data (e.g. felling licences) to correct the CS areas where datasets overlap in time. **Table A 3.4.24** shows the Corrected Forest conversion rates. A correction ratio is used to adjust the estimated deforestation areas, as the Countryside Survey is known to over-estimate deforestation as described in the section above. There are no non-CS data for Northern Ireland so the correction ratios for England or Wales are used, depending on availability. The 1990-98 correction ratios are also applied to the pre-1990 CS land use change estimates.

The annual area of forest converted to other land uses is removed from the area of 4A1 Forest Land remaining Forest Land to maintain consistency in the land area matrix.

Table A 3.4.23 Countryside Survey data for Forest conversion

| Countryside Survey land use change | | Annual rate of change, kha/yr | | | | Grassland/Cropland fractional split | | |
|---------------------------------------|-----------------------------|-------------------------------|----------|-------|-----------|--|----------|-------|
| | | England | Scotland | Wales | N Ireland | England | Scotland | Wales |
| 1990-1998 | Forest to Natural Grassland | 5.600 | 4.418 | 1.099 | 0.171 | 0.61 | 0.86 | 0.72 |
| | Forest to Pasture Grassland | 3.081 | 0.608 | 0.418 | 0.086 | 0.33 | 0.14 | 0.28 |
| | Forest to Cropland | 0.545 | 0.097 | 0.019 | 0.008 | 0.06 | 0.00 | 0.00 |
| | Forest to Settlements | 1.242 | 0.293 | 0.132 | 0.072 | | | |
| | Forest to Other Land | 0.169 | 0.231 | 0.058 | 0.025 | | | |
| 1999-2007 | Forest to Natural Grassland | 2.656 | 10.327 | 0.120 | 0.209 | 0.86 | 0.98 | 0.42 |

| Countryside Survey land use change | | Annual rate of change, kha/yr | | | | Grassland/Cropland fractional split | | |
|---------------------------------------|--------------------------------|-------------------------------|----------|-------|-----------|--|----------|-------|
| | | England | Scotland | Wales | N Ireland | England | Scotland | Wales |
| | Forest to Pasture Grassland | 0.277 | 0.186 | 0.162 | 0.102 | 0.09 | 0.02 | 0.58 |
| | Forest to Cropland | 0.141 | 0.006 | 0.001 | 0.001 | 0.05 | 0.00 | 0.00 |
| | Forests to Settlements | 0.617 | 0.098 | 0.095 | 0.142 | | | |
| | Forest to Other Land | 0.430 | 0.695 | 0.374 | 0.027 | | | |

Table A 3.4.24 Corrected Forest conversion rates

| | | Correction ratio | | | Estimated annual rate of change, kha/yr | | | |
|---------------|-----------------------------|------------------|-----------------|------------------|---|--------------------|--------------------|--------------------|
| | | England | Scotland | Wales | England | Scotland | Wales | N Ireland |
| 1990- 1998 | Grassland & Cropland | 2% ^a | | | 0.159 | 0.088 ^c | 0.026 ^c | 0.005 ^c |
| | Settlements & Other Land | 28% ^b | | | 0.390 | 0.145 ^c | 0.052 ^c | 0.027 ^c |
| 1999- 2007 | Grassland & Cropland | 20% ^a | 2% ^a | 15% ^a | 0.602 | 0.262 | 0.041 | 0.045 ^d |
| | Settlements & Other Land | 28% ^b | | | 0.296 | 0.224 ^c | 0.133 ^c | 0.048 ^c |

^a Unconditional felling licence data used for correction

^b Land Use Change Statistics used for correction

^c England correction ratio used

^d Wales correction ratio used

For 2000 onward, the area and subsequent land-use of deforestation were estimated based on a combination of data sources:

- observations on forest loss by the National Forest Inventory (internal Forestry Commission analysis) by IPCC category. This inventory includes an analysis of deforestation from 2006 to the current inventory year based on a new analysis of woodland maps (Forestry Commission, 2016);
- unconditional felling licences granted (assumed all converted to Grassland);
- analysis of the FC Sub-Compartment Database for restoration of Forest land to open habitats (assumed all converted to Grassland); and
- conversion to non-forest on private sector forest covered by long-term forest plans rather than felling licences (internal Forestry Commission report, assumed all converted to Grassland).

The revision in deforestation was only done from 2000 onwards, partly because there were no suitable data on which to base adjustments for 1990-1999, but also because a number of policy developments came into play in 2000 or shortly beforehand, which affected deforestation to restore open habitats or develop wind-farms. These include the introduction of the UK's climate change policy (2000), and the diversification in relevant forest policies in England, Scotland and Wales following the devolution of forest policy to countries in the late 1990s (Matthews *et al.* 2014). The deforestation information used in this inventory is to the same as the previous inventory, with an assumption of the deforestation rate for 2017.

Soil carbon stock changes associated with deforestation are estimated using the dynamic soil carbon model described in **Section A 3.4.2**. When deforestation occurs it is assumed that 60% of the standing biomass is removed as timber products and the remainder is burnt. In the UNFCCC inventory, KP-LULUCF reporting of deforestation assumes instantaneous oxidation of HWP. Country-specific forest biomass densities for living and dead organic matter from CARBINE are used. These densities change over time in relation to the forest age and species structure. Biomass losses are reported in the relevant carbon stock change tables, assuming a carbon fraction of 0.5 on a dry weight basis. The carbon removed as timber is reported as Harvested Wood Products (HWP) in 4G, using CARBINE to model emission from HWP (described in **Section 6.8**).

Direct and indirect greenhouse gas emissions from associated biomass burning is estimated using the Tier 1 methodology described in the IPCC 2006 guidelines (IPCC 1997 a, b, c) and the emission ratios for CH₄, CO, N₂O and NO_x from Table 3A.1.15 in the IPCC 2003 GPG for LULUCF. Only immediate losses are considered because sites are normally completely cleared, leaving no debris to decay.

A 3.4.5 Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C)

A 3.4.5.1 Activity dataset

Until the 2010 submission only wildfires on Forest land were reported due to a lack of activity data for wildfires on other land use categories. Data on Forest wildfires prior to 2010 come from the Forestry Commission and the Forest Service of Northern Ireland.

In 2010 the Fire and Rescue Service (FRS) began recording wildfires in England, Scotland and Wales on a new Incidence and Reporting Systems (IRS) which includes wildfires on all land use categories. The IRS database contains 30 attributes for each fire to which a fire appliance was called, including date, spatial location, property type description (e.g. heathland and moorland, standing crop) and an estimate of the area burnt. This dataset is available from 1st April 2009. The original dataset had >126,000 fire records but 99% of these fires were less than 1 ha in size. The IRS database is manually completed by fire service personnel and its use requires some subjective judgement. This is likely to lead to non-systematic differences in the accuracy and precision of the data. The accuracy of the locations is variable, but an assessment of a number of the larger fires suggests that the land cover type attribute is reliable. The accuracy of the FRS burnt area estimates could not be validated using aerial photography as the available imagery was not recent enough, so Landsat images were used to validate the FRS data. However, it was still difficult to find cloud-free, pre- and post-fire images for fires in 2010. In addition Landsat has been affected by image 'striping' since 2003, which affects the quality of the images and causes some data loss. There are issues with re-ignited fires or additional fires in the same area being logged in the database as separate events. Overall, the uncertainty associated with this dataset is high but should be re-assessed once a longer time series is available.

To provide data on non-Forest wildfires prior to 2010, thermal anomaly data for 2010 from the NASA-operated MODerate Resolution Imaging Spectroradiometer (MODIS) were obtained from the Fire Information Resource Management System (FIRMS) and allocated to land uses using the proportions of fire on each land use type from the Fire and Rescue Service IRS data. The correlation between MODIS data and IRS data breaks down below 25 ha, so for consistency a 25 ha threshold was set for reporting wildfires logged on the IRS.

Thermal anomalies usually represent active fires, but may also detect industrial heat sources, although these are typically masked out by the thermal anomaly processing chain. The IRS dataset records 89 fires > 25ha occurring in 2010. The FIRMS dataset records 335 fire detections for the same period, however, the FIRMS detections may contain multiple detections for a single fire event and the FIRMS detections are for a single 1 km pixel, and do not have a straightforward conversion to burnt area. Searching the IRS and FIRMS data sets for temporally and spatially coincident events, using a 2 km buffer around the IRS data, suggests that 22 fires were recorded by both the IRS and FIRMS systems. There are wide discrepancies between the two datasets, reflecting their different natures. The IRS data set records fires where a fire service response was required, so does not record controlled burning, unless the fire gets out of control. The FIRMS dataset however, responds to anomalous heat signatures, so records controlled and uncontrolled fires. However, in the UK controlled burning, which is primarily carried out for heath management, is only permitted between October and mid-April to reduce the risk of these burns running out of control (Natural England, 2014¹¹; Scottish Government, 2011¹²). As the FIRMS thermal anomaly data is only collected between March and August it will not detect most fires from controlled burning. FIRMS is only able to detect fires under cloud-free or light cloud conditions and is also only able to detect fires alight at the time of the satellite overpass. The FIRMS data are more likely to detect larger fires than smaller ones, probably due to the stronger heat signature and the longer burn time that larger fires tend to exhibit.

The IRS and FIRMS thermal anomalies give a very different perspective on the extent, timing and duration of fire events in the UK. However, the datasets did show correlation ($R^2 = 70\text{--}81\%$) for fires larger than 25 ha, which enabled an empirical relationship to be derived to extend the burnt area record back to 2001. A burnt area threshold of 25 hectares was used to extract a subset of the IRS database: this captured 75% of the IRS wildfire-burnt area in England, 86% in Scotland and 64% in Wales.

As more IRS data become available confidence should increase in the relationship between fires detected by FIRMS and fires logged in the IRS. This may allow FIRMS data to be extrapolated to fires covering less than 25 ha the inventory in future. However to extend this to small fires there would need to be reasonable confidence that the ratio of large to small fires used was valid, and also some investigation of whether the distribution of small fires across land use classes was the same as that of larger fires.

It was assumed that all fires in the IRS database were wildfires: even if they started as controlled burning, because the need for a fire appliance call-out indicates that they are no longer under control. The IRS property type descriptions were assigned to LULUCF sub-categories (**Table A 3.4.25**). There is a very small area of wildfires that occur on Settlement types, and these are included in the Grassland category as the IRS land type classification suggests that they occur on grassy areas within Settlements and there is not a separate reporting field for wildfires in Settlements in the CRF.

¹¹ <https://www.gov.uk/guidance/heather-and-grass-burning-apply-for-a-licence>

¹² <http://www.gov.scot/Resource/Doc/355582/0120117.pdf>

Table A 3.4.25 IRS database property type descriptions by LULUCF sub-category

| LULUCF sub-category | Forest | Cropland | Grassland | Settlement |
|--------------------------------------|--------------------------------------|--------------------------|----------------------------------|-----------------------------------|
| IRS property type description | Woodland/forest - conifers/softwood | Straw/stubble burning | Heathland or moorland | Domestic garden (vegetation fire) |
| | Woodland/forest - broadleaf/hardwood | Stacked/baled crop | Grassland, pasture, grazing etc. | Park |
| | | Nurseries, market garden | Scrub land | Roadside vegetation |
| | | Standing crop | Tree scrub | Railway trackside vegetation |
| | | | | Wasteland |
| | | | | Canal/riverbank vegetation |

A time series of wildfire-burnt areas for each non-forest land use type was constructed for 1990-2015 (**Figure A 3.6**). For non-forest wildfires for England, Scotland and Wales the IRS burnt areas were used for 2010 to the current inventory year and the burnt area estimated from thermal anomalies from 2000 to 2010. For 1990-2000 the average annual burnt area 2001-2011 was used.

In Northern Ireland, where no IRS data were available, it was assumed that the heathland and grassland burning rates were in the same proportions as the Scottish burning rates, using the area of heathland and grassland from the 2007 Northern Ireland Countryside Survey.

Estimates of the forest area burnt in wildfires 1990-2004 are published in different locations (FAO/ECE 2002; FAO 2005) but all originate from either the Forestry Commission (Great Britain) or the Forest Service (Northern Ireland). There is a gap in the time series 2005-2010 for Great Britain but areas of forest wildfires are reported annually for Northern Ireland. The gap was filled using the annual average areas burnt 1995-2005. These areas refer only to fire damage in state forests; no information is collected on fire damage in privately owned forests. The proportion of private-owned forest that was burnt each year was assumed to be the same as the percentage of the state forest that was burnt each year.

Figure A 3.6 Annual area of FIRMS thermal anomalies for GB for 2001 to the current inventory year (thermal anomalies were filtered to exclude those recorded over urban/industrial areas).





Figure A 3.7 shows the temporal pattern of FIRMS thermal anomalies, with peaks in hot dry years such as 2003. The FIRMS data used only includes thermal anomalies for March – August for each year, as these are the months where the IRS database recorded fires greater than 25 ha. Some FIRMS thermal anomalies were recorded outside these months due to FIRMS detecting both controlled burns and some fires less than 25 ha in size which are not included in the IRS data.

Figure A 3.7 Time series of wildfire burnt areas in the UK 1990 to the current inventory year



A 3.4.5.2 Estimation of emissions

The IPCC Tier 1 method is used for estimating emissions of CO₂ and non-CO₂ gases from wildfires (IPCC 2006). The *Calluna* heath fuel biomass consumption factor and grassland emission factors are used for heathland and moorland fires, the agricultural residues EFs for cropland and the “savannah and grassland” EFs for other grassland and settlements.

Country-specific biomass and Dead Organic Matter densities from the CARBINE model are used for estimating fuel consumption in forest fires (as discussed in the deforestation methodology section) and the ‘extra tropical forest’ EFs in the 2006 Guidelines. In line with the default value in the IPCC 2006 Guidelines for AFOLU it is assumed that 45% of the biomass is consumed in a wildfire in an unfelled temperate forest.

Emissions from all wildfires are reported under the ‘Land remaining Land’ categories (i.e. 4A1, 4B1 and 4C1) and IE reporting under 4A2, 4B2 and 4C2.

A 3.4.6 Emissions from drainage (organic soils) (4B1, 4C1)

Some Wetlands in the UK were drained many years ago for agricultural purposes and continue to emit carbon from the soil. The inventory includes emissions from areas of drained organic soils under Cropland and improved Grassland throughout the UK. The drained areas were generated from work on the UK Agricultural Greenhouse Gas Platform project (Defra project AC0114) (Anthony, ADAS pers. comm). A lack of data on drainage of semi-natural Grassland meant that the area of semi-natural Grassland on drained organic soils could not be estimated. These areas of drained organic soils have also been used for the Agricultural Sector, so there is consistency within the UK Greenhouse Gas Inventory. Work to implement the Wetlands Supplement (IPCC 2013a) guidance is developing a methodology to estimate the area of semi-natural Grassland on drained organic soils. Emissions have been estimated using Tier 1 emissions factors for drained organic soils under Cropland and Grassland taken from the AFOLU Guidelines (IPCC, 2006) applied to all depths of drained organic soil. Results from a BEIS-funded project to implement the Wetland Supplement guidance will allow more detailed estimates of emissions from drained organic soils to be developed in future inventories.

It is assumed that the area of drained organic soils has remained constant as no drainage or rewetting has occurred since 1990 as there have been no policy drivers to encourage drainage or rewetting of cropland or improved grassland. It is also assumed that land on organic soils has not been converted between land uses.

A 3.4.7 Emissions of N₂O due to disturbance associated with land use conversion and land management changes (4(III))

Methodological coverage of this activity has expanded in the IPCC 2006 Guidelines, as previously only N₂O emissions due to soil disturbance associated with land use conversion to Cropland were included. All land use conversions or land management changes that result in a loss of soil carbon, leading to N mineralization and N₂O emissions are now reported. Direct emissions from soils, and indirect emissions from nitrogen leached from soil and subsequently oxidised are included in the inventory. The UK now includes emissions resulting from the land use conversions: 4A2 Land converted to Forest Land, 4B2 Forest to Cropland and Grassland to Cropland, 4C1 Grassland remaining Grassland, 4C2 Forest to Grassland, 4E1 Settlement remaining Settlement and 4E2 Land converted to Settlement. Emissions of N₂O from 4C1 and 4E1 arise from land use change over 20 years before the inventory reporting year where loss of soil organic matter is still ongoing. Emissions of N₂O from 4B1 Cropland remaining Cropland

(resulting from land use change over 20 years before the inventory year) are calculated in the same way by the LULUCF inventory team but are included in the Agriculture sector (category 3D1).

The Tier 1 methodology described in the IPCC 2006 Guidelines is used. The activity data are the areas and soil carbon stock changes reported in the relevant categories in 4B, 4C and 4E. Some C:N ratios for UK soil/vegetation combinations are published in the Countryside Survey (with values of 11.7 to 13.4) but only for the top 15 cm of soil. However, the soil carbon stock changes reported in the inventory are from the top 1 m of soil, so these C:N ratios were not felt to be applicable. Therefore, the IPCC default C:N ratio of 15 is used for estimating mineralised N. The emission factor of 1% in the 2006 Guidelines was used to estimate N₂O emissions from mineralised N. Indirect N₂O emissions from mineralisation are also estimated from carbon stock change using Tier 1 methodology.

A 3.4.8 On-site and off-site emissions from peat extraction (4D)

On-site emissions of CO₂ and N₂O from peat extraction activities (for energy and horticultural use) and off-site emissions of CO₂ from the decomposition of horticultural peat are reported in category 4D.

A 3.4.8.1 Activity datasets

Available data sets on peat extraction vary between Northern Ireland and for Great Britain (England, Scotland and Wales). From 2002 onwards Google Earth imagery has been used to estimate the area of peat extraction and sites list in the Directory of Mines and Quarries and the BritPits online database¹³. Prior to the 2002 no Google Earth images are available, and peat extraction site areas have been estimated from other sources. **Table A 3.4.26** shows the sources of activity data used to estimate emissions from peat extraction.

Table A 3.4.26 Activity data for peat extraction sites in Northern Ireland

| Data set | Information contained | Geographic extent | Time period | Publication frequency |
|---|--|--|--------------------|---|
| Directory of Mines and Quarries (DMQ)/BritPits database | Location of active peat extraction sites | England, Scotland, Wales, Northern Ireland | 1984 - present | Online database is continuously updated |
| Google Earth | Land use images to estimate area of extraction sites identified from DMQ | UK | 2002 - date | Variable |
| Cruickshank and Tomlinson (1997) | Area with planning consent for peat extraction Local authority planning consents for peat extraction sites | England, Scotland, Wales | 1990/91 | One off compilation of data |

¹³ <https://www.bgs.ac.uk/products/minerals/britpits.html>

| Data set | Information contained | Geographic extent | Time period | Publication frequency |
|--|---|--------------------------|-------------|-----------------------------|
| Tomlinson (2010) | Estimate of peat extraction area in Northern Ireland. Volume of peat extracted (sod cutting and vacuum harvesting) | Northern Ireland | 1990 - 1991 | One off compilation of data |
| Mineral Extraction in Great Britain (Annual Minerals Raised Inquiry) | Volume of peat extracted | England, Scotland, Wales | 1947 - date | Annual |
| Cruickshank et al. 1995 | Volume of peat extracted (hand cutting) | Northern Ireland | 1990 - 1991 | One off compilation of data |

The areas of peat extractions sites listed in the BritPits database were assessed using Google Earth. Polygons were drawn around site boundaries and the area covered by the polygons was calculated in Google Earth. Change over time at individual sites was tracked to give an estimate of the extent of conversion to and from extraction sites. This method is repeated annually to incorporate changes in extraction site areas in new Google Earth images.

Any sites abandoned since 2002 where a change of land use cannot be identified are still judged to be producing on-site emissions, in line with good practice guidance. A time series was constructed using linear interpolation. The extraction area (active and abandoned where there has been no change in land use) declined between 1990 and 2015 by 40% in England, 43% on fuel peat sites in Scotland, 6% on horticultural peat sites in Scotland, 99% on fuel peat sites in Northern Ireland and 40 % on horticultural peat sites in Northern Ireland. This area was assumed to be converted to Grassland.

The 2017 version of the BritPits database lists four sites in Wales, most having ceased extraction. Only one of these is visible on Google Earth and the Mineral Extraction in Great Britain report for 2013 does not report any peat production in Wales, so it is assumed that the visible site is currently inactive but has not been converted to another land use.

Annual peat production in Great Britain (**Table A 3.4.27**) is inferred from extractor sales by volume as published in the annual Mineral Extraction in Great Britain report, formerly known as the Minerals Raised Inquiry (ONS). This gives a breakdown for horticultural and other uses of peat, which are assumed to be fuel, for English regions and for Scotland. No peat extraction is reported in Wales. Annual production is highly variable because extraction methods depend on suitable summer weather for drying peat.

Table A 3.4.27 Annual peat production, m³ for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports)

| Year | England | | Scotland | |
|------|---------------|-------|---------------|---------|
| | Horticultural | Fuel | Horticultural | Fuel |
| 1990 | 1,116,940 | 2,727 | 293,170 | 93,163 |
| 1991 | 1,202,000 | 2,000 | 241,000 | 115,000 |

| Year | England | | Scotland | |
|-------|---------------|--------|---------------|---------|
| | Horticultural | Fuel | Horticultural | Fuel |
| 1992 | 1,079,000 | 4,000 | 332,000 | 91,000 |
| 1993 | 1,069,820 | 2,180 | 306,511 | 73,489 |
| 1994 | 1,375,000 | 1,000 | 498,000 | 108,000 |
| 1995 | 1,578,000 | 2,000 | 657,000 | 44,000 |
| 1996 | 1,313,000 | 2,000 | 517,000 | 53,000 |
| 1997 | 1,227,000 | 2,000 | 332,000 | 59,000 |
| 1998 | 936,000 | 0 | 107,000 | 32,000 |
| 1999 | 1,224,000 | 0 | 392,000 | 37,000 |
| 2000 | 1,258,000 | 1,000 | 336,000 | 31,000 |
| 2001 | 1,459,000 | 1,000 | 325,000 | 30,000 |
| 2002 | 856,000 | 1,000 | 107,000 | 10,000 |
| 2003 | 1,227,000 | 1,000 | 741,000 | 38,000 |
| 2004 | 902,000 | 1,000 | 338,000 | 21,000 |
| 2005 | 927,000 | 1,000 | 556,000 | 21,000 |
| 2006 | 856,000 | 1,000 | 712,000 | 24,000 |
| 2007 | 654,000 | 0 | 221,000 | 10,000 |
| 2008 | 455,000 | 41,000 | 243,000 | 21,000 |
| 2009 | 476,000 | 0 | 390,000 | 21,000 |
| 2010 | 456,000 | 1,000 | 527,000 | 21,000 |
| 2011 | 429,000 | 0 | 369,000 | 26,000 |
| 2012 | 422,000 | 0 | 126,000 | 20,000 |
| 2013 | 661,000 | 0 | 570,000 | 24,000 |
| 2014 | 294,000 | 0 | 469,000 | 32,000 |
| 2015* | 563,000 | 0 | 417,772 | 22,528 |
| 2016* | 563,000 | 0 | 417,772 | 22,528 |
| 2017* | 563,000 | 0 | 417,772 | 22,528 |

* 2005 - 2014 average as UK Minerals Yearbook has ceased production.

A 3.4.8.2 Estimation of emissions

Default on-site emission factors for Tier 1 reporting (IPCC 2006) are used to estimate emissions. Peat extracted for horticultural use is inferred to be from oligotrophic (nutrient-poor) bogs. Peat for fuel is inferred to be from mineratrophic (nutrient-rich) fens or bogs. On-site emissions of CO₂ and N₂O from drainage are reported.

A value of 0.0641 tonnes C m⁻³ is used for Great Britain to estimate emissions from extracted horticultural peat volumes based on previous work (Thomson *et al*, 2011). This is slightly lower than the IPCC default emission factor of 0.07 tonnes C m⁻³ air-dry peat for nutrient-poor peats.

Tomlinson (2010) gives production estimates of horticultural peat production for Northern Ireland for 1990/91 and 2007/2008. These have been interpolated to produce a time series. The total emission from horticultural peat production is the sum of emissions from vacuum harvesting production, sod extraction production and mechanical extraction production.

Emissions from vacuum harvesting production =

area * annual depth of extraction * carbon fraction by volume

where

Annual depth of extraction by vacuum harvesting, m/ha = 0.1

Carbon fraction of air-dry peat by volume, tonnes C/m³ air-dry peat = 0.0641

Emissions from sod extraction production =

area * sod extraction rate * % dry matter for sods * mean % C

where

Sod extraction rate, tonnes/ha/yr = 200

Sod extraction, mean % dry matter = 35%

Mean % carbon = 49%

Emissions from mechanical extraction production =

area * extraction rate * % dry matter for mechanical extraction * mean % C

where

The mechanical extraction rate was estimated to be 206.45 tonnes/ha in 1990/91 and 243.06 tonnes/ha in 2007/08 (Tomlinson, 2010).

Mechanical extraction, mean % dry matter = 67%

Mean % carbon = 49%

A 3.4.9 Flooded Lands (4D)

Carbon stock changes on land converted to Flooded Land (reservoirs) are included in the inventory, based on the IPCC 2006 Guidance. Data on all reservoirs over 1 km² were compiled but only reservoirs established since 1990 were reported (areas of inland water under 1 km² are reported under 4D Wetlands remaining Wetlands). Activity data were compiled for England and Wales from the Public Register of Large Raised Reservoirs provided by the Environment Agency, which listed location, surface area and year built. Activity data for Scotland were compiled from the SEPA Water Body Classification database (of water bodies > 0.5 km²) and the associated Water Body data sheets. Additional information on the year of building was obtained from:

- the Gazetteer for Scotland <http://www.scottish-places.info>;
- hydro-electric power generators <http://sse.com/whatwedo/ourprojectsandassets/> <http://www.power-technology.com/projects/glendoehydropowerpla/>; and
- local authorities <http://www.argyll-bute.gov.uk>.

It was established through discussion with local experts that no new large reservoirs had been built in Northern Ireland since the 1950s.

Only five large reservoirs have been established in the UK since 1990, three in England and one each in Scotland and Wales (another five in England are sacrificial floodplains and do not fit the criteria of permanent conversion to Flooded Land). These cover a total of 1.995 kha.

The location of each reservoir was examined using the www.magic.gov.uk geographic information portal. All reservoirs were in upland locations and were assumed to be Grassland prior to their conversion to Flooded Land. (Any forest removed as part of the land conversion will have been captured under the deforestation activity methodology). A Tier 1 methodology was followed, so carbon stock changes in living biomass stock in the year of flooding were estimated, but not carbon stock change in soils. A living biomass density of 2 t dry matter/ha was used to estimate carbon stock losses. This will be updated in the next submission to be consistent with the biomass densities used in other parts of the LULUCF inventory.

A 3.4.10 Harvested Wood Products (4G)

The activity data used for calculating this activity are the annual forest planting rates. CARBINE then applies a forest management regime as given in input to the model. For a given forest stand, carbon enters the HWP pool when thinning is undertaken and when harvesting takes place. Depending on the species, first thinning occurs approximately 20 years after planting.

At thinning and harvest, the CARBINE model allocates merchantable stem volume to various wood products, while the remainder is transferred to the harvesting residue pool. The 'end-use' wood products represented are:

- Long-lived sawn timber, e.g. timber used for construction;
- Short-lived sawn timber, e.g. timber used for fencing;
- Particleboard;
- Paper; and
- Fuel.

For reporting purposes the long-lived and short-lived sawn timber are reported together in the Sawnwood category.

During wood extraction, conversion losses are assumed to be left as on-site harvest residue and enter the litter pool. The allocation of carbon to wood product categories is estimated by inputting the merchantable stem carbon (from the forest yield model) to a stand volume assortment forecasting model which estimates the volume allocated to sawn timber, roundwood and waste. This is implemented in CARBINE as a set of functions derived from the output of a more general and flexible assortment forecasting program known as ASORT (Rollinson and Gay, 1983).). The CARBINE model uses standard estimates for oven-dried wood density to derive biomass from the harvested volume (Lavers and Moore, 1983; Jenkins et al., 2011). Carbon content of all oven-dried wood is assumed to be 50% (Matthews, 1993). CARBINE assumes the wood of a tree species all has the same oven-dried wood density and carbon content, irrespective of which semi-finished wood product categories it is assigned to.

The proportions of wood produced which are allocated to different product categories are based on proportions derived from FAO data¹⁴ (prior to 1994) and forestry commission data¹⁵ (after 1993) on production of semi-finished wood products. A carbon retention curve is used to estimate product decay and return of carbon to the atmosphere. Each wood product category

¹⁴ <http://www.fao.org/faostat/en/#data/FO>

¹⁵ <https://www.forestryresearch.gov.uk/tools-and-resources/statistics/statistics-by-topic/timber-statistics/uk-wood-production-and-trade-provisional-figures/>

has its own carbon retention curve using the default half-lives in the IPCC 2013 Revised Supplementary Methods and Good Practice Guidance, taking into account the decay rate of wood products and the service life as influenced by socio-economic factors. The half-lives are: 35 years for sawn wood; 25 years for wood panels; 2 years for paper. Timber used as woodfuel is assumed to instantaneously oxidise.

In implementing the 2006 IPCC guidelines for HWP the UK has elected to report using the production approach B2, which requires disaggregation of HWP into those produced and consumed domestically and those produced and exported. In the annual Forestry Statistics publication there is data on the apparent consumption of wood products in the UK. A consistent dataset is available at the product level (i.e. sawnwood, wood panels and paper & paperboard) for 2002 onwards. The ten year average of 2002-2011 was calculated for each product type and those values were used for the years 1990-2001. This dataset was used to assign the HWP output from the CARBINE model into either consumed domestically or exported.

A 3.4.11 Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs)

The UK LULUCF inventory includes direct GHG emissions from UK Crown Dependencies (CDs) and Overseas Territories (OTs) which have joined, or are likely to join, the UK's instruments of ratification to the UNFCCC and the Kyoto Protocol. Currently, these are: Guernsey, Jersey, the Isle of Man, the Falkland Islands, the Cayman Islands, Bermuda, and Gibraltar. It should be noted that Bermuda will not ratify the 2nd Commitment Period of the Kyoto Protocol and is therefore not included in the 'GBK' submission under the Kyoto Protocol.

A web search of statistical publications was undertaken for any updates in datasets in 2017. This work builds on an MSc project to calculate LULUCF net emissions/removals for the OTs and CDs undertaken during 2007 (Ruddock 2007).

The availability of data for the different OTs and CDs is very variable, so that emission estimates can only be made for the Isle of Man, Guernsey, Jersey and the Falkland Islands. These four comprise over 95% of the area in all the OTs and CDs. Gibraltar wished to produce their own inventory and has assessed its LULUCF net emissions/removals as Not Occurring (Annex 3.6) as the country is extremely small (6 km²). A lack of suitable data for the Caribbean territories (discussed in the 1990-2006 NIR) makes it impossible to create inventories for them at the present time. Discussions have started with the Cayman Islands government with a view to reporting LULUCF emissions from this OT in the future.

A number of omissions (reported as Not Estimated) for the Overseas Territories and Crown Dependencies were identified in the 2016 UNFCCC Expert Review of the GHG Inventory. These included wildfires on forest land and grassland, flooded land, peat extraction and some carbon pools affected by deforestation. Only very limited further data has been discovered through additional investigations, and so the threshold of insignificance (Decision 24/CP.19, paragraph 37b) has been applied.

- Wildfires on forest land: no information on the occurrence of wildfires on forest land was obtained by further investigation. The average area of forest wildfires in the UK 2008-2017 was 0.632 kha/year, with an average emission of 73.6 Gg CO₂e (CO₂, CH₄ and N₂O). This equates to 0.018% of UK forests burnt every year, and the equivalent area burnt in the Crown Dependencies (the Falkland Islands have no forest cover) would be <0.001 kha/year with estimated emissions falling well below the insignificance threshold of 500 Gg CO₂e.

- **Wildfires on grassland:** no information on grassland fires in the Crown Dependencies was obtained by further investigation. There are some reports of wildfires in the Falkland Islands, caused by lightning strikes or pasture burning. The average area of grassland wildfires in the UK 2008-2017 was 10.148 kha/year, with an average emission of 13.5 Gg CO₂e (CO₂, CH₄ and N₂O). This equates to 0.076% of UK grassland burnt every year, and the equivalent area burnt in the Overseas Territories and Crown Dependencies (dominated by the large area of grassland in the Falkland Islands) would be 0.961 kha/year with estimated emissions of 1.28 Gg CO₂e falling well below the insignificance threshold of 500 Gg CO₂e. A watching brief will be kept on this potential emission source, particularly in the Falkland Islands.
- **Flooded land:** An assessment of flooded land was undertaken for the Overseas Territories and Crown Dependencies. No flooded land areas exceed the area threshold of 1 km² used for the UK, so the area of Flooded Land remaining Flooded Land has been included with the Inland Water area in the Wetlands remaining Wetlands category (0.095 kha in total).
- **Peat extraction:** Peat organic soils occur in the Falkland Islands and Isle of Man but not in the other Overseas Territories and Crown Dependencies. Information on the extent and condition of peat soils in these territories has been compiled a part of the research project on implementation of the Wetlands Supplement (see box in Chapter 6, section 6.5.8). Further work is underway to incorporate these results into the GHG Inventory. Further research on the peat soils of the Falkland Islands is also underway and will be used to improve reporting in this category in due course.
- **Deforestation:** A small area of deforestation (0.002 kha/yr) occurs in Guernsey, obtained from two habitat surveys in 1999 and 2010. The limited data only permits Tier 1 estimation of carbon stock changes in biomass and litter. The change in forest cover is a result of the changed areas losing sufficient tree cover to be reclassified as dense scrub or parkland, rather than conversion to settlement land or agriculture. Any changes in soil and dead wood are very likely to be minimal. As all emissions from deforestation in the Overseas Territories and Crown Dependencies fall well below the insignificance threshold of 500 Gg CO₂e (estimated to be 0.64 Gg CO₂ in 2017 if a similar rate of deforestation to the UK rate is assumed), the UK will continue to report these carbon stock changes as Not Occurring and will continue to report emissions from this activity that it has reported in previous years.

Information on the area of each IPCC land category, dominant management practices, land use change, soil types and climate types were compiled for each OT/CD from statistics and personal communications from their government departments (**Table A 3.4.28**). This allowed Tier 1 level inventories to be constructed for the four OT/CDs already mentioned, and a Tier 3 approach for Forest Land on the Isle of Man and Guernsey (using the C-Flow model, for information on C-Flow model please refer to 1990-2011 NIR). The assumptions and factors used for the estimation of emissions are given in **Table A 3.4.29** and **Table A 3.4.30**. The estimates have high uncertainty and may not capture all relevant activities, but given the size of the territories any missing sources are likely to be small.

Table A 3.4.28 Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies

| Territory | LULUCF category | Time period | Reference |
|------------------|--------------------|--|--|
| Isle of Man | 4A | 1970-2011 | Personal communication from Isle of Man Department of Agriculture, Fisheries and Forestry (Peter Williamson) FAO (2010) Global Forest Resources Assessment: Isle of Man |
| | 4B, 4C, 4D | 2002-2011 | Isle of Man Agricultural and Horticultural Census: completed by all farmland occupiers on an annual basis Isle of Man Digest of Economic and Social Statistics |
| | 4B, 4C | 2012 - 2013 | Isle of Man Digest of Economic and Social Statistics |
| | 4B, 4C | 2014 | The Isle of Man in Numbers |
| | 4E | 1991-1994 | Isle of Man Ecological Habitat Survey, Phase 1 Report (Sayle et al, 1995) |
| Guernsey | 4A | 1990-2010 | FAO Global Forest Resources Assessment 2010: Guernsey |
| | 4A, 4B, 4C, 4D, 4E | 1998/9, 2005, 2010 | Guernsey Habitat Survey Sustainable Guernsey 2005, 2009, Guernsey Facts and Figures 2011 |
| Jersey | 4A | 1990-2010 | FAO Global Forest Resources Assessment 2010: Jersey |
| | 4B | 1990 - 2014 | Jersey Agricultural Statistics |
| | 4A, 4B, 4C, 4D, 4E | 2006, 2008, 2012, 2015 | Jersey In Figures 2006/2008/2009/2010/2011/2012 |
| Falkland Islands | 4A | 1990-2011 | Department of Mineral Resources, personal communication FAO Global Forest Resources Assessment 2010: Falkland Islands |
| | 4B, 4C | 1991-2013 | Falkland Islands Agricultural Statistics |
| | 4E | 1990-2005 | Falkland Islands Environment and Planning Department, personal communication |
| | 4E | 1986 – 2001 with projections 2006 - 2016 | Stanley Town Plan, Environmental Planning Dept, Falkland Islands Government. |

Table A 3.4.29 Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies

| Land Use category | Sub-category | Isle of Man | Guernsey | Jersey | Falkland Islands |
|--------------------|---|-------------------|-------------------|------------------------|------------------------|
| Forest land fluxes | Living biomass, DOM, Mineral soils, Organic soils | From C-Flow model | From C-Flow model | Assumed in equilibrium | No forest on Falklands |

| Land Use category | Sub-category | Isle of Man | Guernsey | Jersey | Falkland Islands |
|--------------------------------|----------------------------|--|--|--|--|
| Crop remaining crop | Living biomass | N/A. Only for perennial crops | N/A. Only for perennial crops | N/A. Only for perennial crops | N/A. Only for perennial crops |
| | Dead organic matter | N/A | Orchards only. 10 tC/ha | N/A | N/A |
| | Mineral soils | No change in SOC | No change in SOC | No change in SOC | N/A |
| | Organic soils | N/A | N/A | N/A | Default (-5 tC/ha) |
| Land converted to Crop | Living biomass | UK shrubby grass to crop values(5 tC/ha) | UK non-shrubby grass to crop value (2.2 tC/ha) | UK non-shrubby grass to crop value (2.2 tC/ha) | UK non-shrubby grassland to crop value (2.2 tC/ha) |
| | Dead organic matter | N/A | N/A | N/A | N/A |
| | Mineral soils | Default SOC = 95 tC/ha, assume conversion from natural grassland (-1.7347 tC/ha) | Default SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha) | Default . SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha) | N/A |
| | Organic soils | N/A | N/A | N/A | Default (-5 tC/ha) |
| | N ₂ O emissions | Default (0.001817 t N ₂ O/ha) | Default (0.000995 t N ₂ O/ha) | Default (0.000995 t N ₂ O/ha) | Default (0.012571 t N ₂ O/ha) |
| Grass remaining grass | Living biomass | N/A | N/A | N/A | N/A |
| | Dead organic matter | N/A | N/A | N/A | N/A |
| | Mineral soils | No change in SOC | No change in SOC | No change in SOC | N/A |
| | Organic soils | N/A | N/A | N/A | Assume no soil C stock change |
| Land converted to grass | Living biomass | UK crop to non-shrubby grass values (-2.2 tC/ha) | UK settlement to non-shrubby grass value (0 tC/ha) | Crop to Grassland: UK crop to non-shrubby grass value (-2.2 tC/ha) Settlement to Grassland: assume increase from 0 in glasshouses (2.8 tC/ha) | Use crop to non-shrubby grassland value (-2.2 tC/ha) |

| Land Use category | Sub-category | Isle of Man | Guernsey | Jersey | Falkland Islands |
|--|---|--|--|---|-----------------------|
| | Dead organic matter | N/A | N/A | N/A | N/A |
| | Mineral soils | Default SOC = 95 tC/ha, assume conversion from cropland (1.7347 tC/ha) | Default SOC = 95 tC/ha, assume conversion from settlement, assume same soil C as for cropland (0.95 tC/ha) | Default SOC = 95 tC/ha, Cropland to Grassland: assume conversion from cropland (0.95 tC/ha) Settlement to Grassland: assume no change | N/A |
| | Organic soils | N/A | N/A | N/A | Default (-0.25 tC/ha) |
| Settlements remaining Settlements | Living biomass, DOM, Mineral soils, Organic soils | N/A | N/A | N/A | N/A |

| Land Use category | Sub-category | Isle of Man | Guernsey | Jersey | Falkland Islands |
|--|---|--|---|---|--|
| Land converted to Settlements | Living biomass | UK values, shrubby grass to settlement (-7.2 tC/ha) | UK non-shrubby grass to settlement value (0 tC/ha) | Grassland to Settlement, UK non-shrubby grass to settlement value (0 tC/ha) Cropland to Settlement: use cropland to settlement value (-2.2 tC/ha) | Use shrubby grass to settlement value (-7.2 tC/ha) |
| | Dead organic matter | N/A | N/A | N/A | N/A |
| | Mineral soils | Default SOC = 95 tC/ha, assume conversion from grassland and all soil C lost (-4.75 tC/ha) | Default SOC = 95 tC/ha, assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha) | Default . SOC = 95 tC/ha, Grassland to Settlement: assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha) Cropland to Settlement: assume 30% of land is paved over and the rest is turf grass (0.95 tC/ha) | N/A |
| | Organic soils | N/A | N/A | N/A | Default - assume cropland (-5 tC/ha) |
| | N ₂ O emissions | Default (0.004976 t N ₂ O/ha) | Default (0.00119 t N ₂ O/ha) | Default (0.00119 t N ₂ O/ha) | N/A |
| Other land remaining other land | Living biomass, DOM, Mineral soils, Organic soils | N/A | N/A | N/A | N/A |

| Land Use category | Sub-category | Isle of Man | Guernsey | Jersey | Falkland Islands |
|------------------------------|----------------------------|-------------------|---|---|------------------|
| Land converted to other land | Living biomass | N/A | Assume loss of grassland to standing water or cliff (-10 tC/ha) | Assumed loss of grassland to standing water (-2.8 t C/ha) | N/A |
| | Dead organic matter | N/A | N/A | N/A | N/A |
| | Mineral soils | N/A | Assume no change in soil stocks | N/A | N/A |
| | Organic soils | N/A | N/A | N/A | N/A |
| | N ₂ O emissions | N/A | 0 | N/A | N/A |
| Harvested wood products | | From C-Flow model | From C-Flow model | N/A | N/A |

Table A 3.4.30 IPCC Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies

| | Factor | Isle of Man/ | Guernsey/ Jersey | Falkland Islands |
|---|--|--------------|------------------|------------------|
| Biomass carbon densities, tC/ha | Cropland | 5 | 5 | 5 |
| | Grassland (shrubby) | 10 | 10 | 10 |
| | Grassland (non-shrubby) | 2.8 | 2.8 | 2.8 |
| | Settlements | 2.8 | 2.8 | 2.8 |
| | Soil C density | 95 | 95 | 87 |
| | Grass F _{lu} | 1 | 1 | 1 |
| | Grass F _{mg} | 1 | 1 | 1 |
| | Grass F _i | 1 | 1 | 1 |
| | Crop F _{lu} | 0.69 | 0.8 | 0.69 |
| | Crop F _{mg} | 1 | 1 | 1 |
| | Crop F _i | 0.92 | 1 | 0.92 |
| | C/N ratio kg N ₂ O-N/kg N | 15 | 15 | 15 |
| | N ₂ O EF | 0.01 | 0.01 | 0.01 |
| | Frac-Leach | 0.3 | 0.3 | 0.3 |
| | | 0.0075 | 0.0075 | 0.0075 |
| | Cropland Organic soils EF, tC/ha/yr | | | -5 |
| | Grassland Organic soils EF, tC/ha/yr | | | -0.25 |
| | EF2 for temperate organic crop and grassland soils, kg N ₂ O-N/ha | | | 8 |
| Forest converted to grassland (Guernsey only) | Living biomass, t DM/ha | | 120 | |


| | Factor | Isle of Man/ | Guernsey/ Jersey | Falkland Islands |
|--|---------------------------------------|--------------|------------------|------------------|
| | Dead organic matter, tC/ha | | 16 | |
| | Carbon fraction of dry matter | | 0.47 | |
| | Biomass C density, tc/ha | | 56.4 | |
| | Change in mineral soil density, tC/ha | | 0 | |

A 3.4.12 Uncertainty analysis of the LULUCF sector

The purpose of carrying out uncertainty analysis within the LULUCF inventory is to quantify where the largest sources of errors lie, and to identify areas to be targeted in future work so as to reduce the uncertainties. In the 1990-2010 inventory report a sensitivity analysis of the whole of the existing inventory methodology was undertaken, applying uncertainty quantification more widely and rigorously to all model parameters and empirical conversion factors, in order to quantify the impact of those uncertainties on the inventory. Although this analysis was carried out for the C-Flow model, which is no longer used, it is broadly applicable to the CARBINE model as both are similar forest carbon accounting models, based on the same underlying yield tables and input data. An update to this uncertainty analysis, which will include the CARBINE model, will be undertaken in 2019 and is likely to be reported in the 2020 submission.

The results of the simulations, including both input and parameter uncertainty, are that the area undergoing land use change is the single biggest uncertainty in the inventory, followed by uncertainty in the forest model parameters and the choice of model for the change in soil carbon following land use change (**Figure A 3.8**). The next five terms are all of a similar magnitude. Full details of the methodology and results are in the 1990-2010 inventory report.

The uncertainty in the land use change areas is being addressed by the development of a new vector-based approach (see **Chapter 7, Section 1.1**), combining multiple sources of land use data.

Figure A 3.8 The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions

Parameterisation of the forest model is the second largest source of uncertainty. This has been addressed with the move to CARBINE, as 19 tree species are now modelled instead of the two used in previous submissions. Results from the National Forest Inventory (NFI) and small woods dataset will also provide additional information on carbon stocks in trees (e.g. Forestry Commission 2015¹⁶). The choice of soil carbon model and its parameters are also important, because the time course of the flux following land use change may be quite different, depending on the equations used to represent this, and how carbon is distributed between fast- and slow-turnover pools. The choice of forest model is less important, largely because all the UK forest models are based on the same yield table data.

A 3.5 WASTE (CRF SECTOR 5)

A 3.5.1 Solid Waste Disposal on Land (5A)

A 3.5.1.1 Input data

Because waste sent to landfill is now evaluated using individual waste consignments by EWC code, there is no need to make assumptions regarding waste composition, other than for two waste categories. These EWC codes are 19.12.12 (residues from waste sorting) and 20.03.01 (mixed municipal waste). Wastes with these codes were allocated in accordance with the findings of a survey carried out on behalf of Defra (Resource Futures, 2012), as set out in **Table A 3.5.1**. Data on DOC, DOCf and material compositions are provided in **Table A 3.5.2**.

The model allocates waste to two types of landfill – old, closed sites which last received waste in 1979, and modern engineered landfills that came into operation from 1980. Only these latter

¹⁶ This survey is preliminary and the carbon stocks have been estimated using the same relationships and calculation parameters that underlie CARBINE; they are therefore not an independent validation of the LULUCF estimates.

sites have gas management systems. The old closed sites have no gas control. The distribution of waste between these types of site is the same as used for compiling the previous NIR.

The quantities of waste sent to landfill are shown in **Table A 3.5.3**. The amounts of methane generated, recovered, used for power generation, flared, oxidised and emitted to the atmosphere are shown in **Table A 3.5.4**.

Table A 3.5.1 Composition of waste sorting residues and mixed municipal waste

| Material | 19.12.12 (residues from waste sorting) | 20.03.01 (mixed municipal waste) |
|------------------------------|--|----------------------------------|
| Paper | 10.3% | 10.6% |
| Card | 9.1% | 7.7% |
| Plastic film | 9.4% | 8.4% |
| Dense plastics | 13.2% | 9.6% |
| Sanitary waste | 1.3% | 3.1% |
| Wood | 10.0% | 5.3% |
| Textiles and shoes | 5.9% | 5.6% |
| Glass | 1.3% | 3.0% |
| Food waste | 8.2% | 21.3% |
| Garden waste | 1.8% | 3.5% |
| Other organic | 1.3% | 2.1% |
| Metals | 3.2% | 3.7% |
| WEEE | 1.4% | 1.5% |
| Haz waste and batteries | 1.1% | 0.9% |
| Carpet, underlay & furniture | 7.0% | 5.0% |
| Other combustibles | 2.7% | 1.4% |
| Bricks, plaster and soil | 7.9% | 4.1% |
| Other non-combustible | 1.7% | 1.5% |
| Fines <10mm | 3.3% | 1.8% |
| Total | 100% | 100% |

Table A 3.5.2 DOC, DOCf and composition of waste materials

| Component | Lignin biodegradability DOCf | Non-lignin biodegradability DOCf | Moisture, %fresh matter | Lignin, % dm | Hemicellulose, %dm | Cellulose, %dm | Starch, %dm | Sugar, % dm | Fat, %dm | Proteins, %dm | Fibre, %dm | Readily soluble, %dm | Other (inert), % dm |
|---|------------------------------|----------------------------------|-------------------------|--------------|--------------------|----------------|-------------|-------------|----------|---------------|------------|----------------------|---------------------|
| Carbon contents (DOC) | | | 0% | 65.1% | 44.6% | 40.0% | 44.4% | 42.1% | 76.0% | 40.0% | 45.0% | 45.0% | 0.0% |
| Waste composition on a dry matter basis (other than moisture) | | | | | | | | | | | | | |
| Municipal solid waste | | | | | | | | | | | | | |
| Paper | 5% | 65% | 15% | 15% | 9% | 61% | | | | | | | 15.00% |
| Card | 5% | 65% | 20% | 15% | 9% | 61% | | | | | | | 15.00% |
| Nappies | 5% | 65% | 65% | | | 47% | | | | | | | 52.70% |
| Textiles and footwear | 5% | 65% | 20% | | 15% | 15% | | | | | | | 69.68% |
| Miscellaneous combustible | 5% | 65% | 20% | | 25% | 25% | | | | | | | 50.00% |
| Wood | 5% | 65% | 17% | 26% | 12% | 42% | | | | | | | 21.00% |
| Food – corrected | 15% | 70% | 70% | 6% | 4% | 27% | 13% | 7% | 14% | 15% | 14% | 0% | 0.00% |
| Garden | 10% | 65% | 55% | 20% | 16% | 20% | | | 2% | | | 26% | 17.10% |
| Soil and other organic | 5% | 65% | 30% | | 1% | 1% | | | | | | | 98.60% |
| Furniture | 5% | 65% | 12% | 1% | 10% | 11% | 0% | 0% | 0% | 0% | 0% | 0% | 77.25% |
| Mattresses | 5% | 65% | 20% | | 15% | 15% | | | | | | | 69.68% |
| Non-inert Fines | | 50% | 40% | | 25% | 25% | | | | | | | 50.00% |
| Other (100% inert) | | | | | | | | | | | | | 100.00% |

| Component | Lignin biodegradability DOCf | Non-lignin biodegradability DOCf | Moisture, %fresh matter | Lignin, % dm | Hemicellulose, %dm | Cellulose, %dm | Starch, %dm | Sugar, % dm | Fat, %dm | Proteins, %dm | Fibre, %dm | Readily soluble, %dm | Other (inert), % dm |
|--|------------------------------|----------------------------------|-------------------------|--------------|--------------------|----------------|-------------|-------------|----------|---------------|------------|----------------------|---------------------|
| Commercial & industrial waste | | | | | | | | | | | | | |
| Commercial | 5% | 65% | 37% | | 8% | 76% | | | | | | | 16.00% |
| Paper and Card | 5% | 65% | 15% | 15% | 9% | 61% | | | | | | | 15.00% |
| General industrial waste | 5% | 65% | 37% | | 8% | 76% | | | | | | | 16.00% |
| Food and Abattoir | 15% | 70% | 70% | 5% | 11% | 11% | 36% | 7% | 6% | 18% | | | 6.00% |
| Food effluent | 15% | 70% | 65% | | 55% | 7% | | | | | | | 37.40% |
| Construction and demolition | 5% | 65% | 30% | | 9% | 9% | | | | | | | 83.00% |
| Miscellaneous process waste | 5% | 65% | 20% | | 10% | 10% | | | | | | | 80.00% |
| Other waste | 5% | 65% | 20% | | 25% | 25% | | | | | | | 50.00% |
| Miscellaneous Combustible | 5% | 65% | 20% | | 25% | 25% | | | | | | | 50.00% |
| Furniture | 5% | 65% | 12% | 1% | 10% | 11% | 0% | 0% | 0% | 0% | | | 77.25% |
| Garden | 10% | 65% | 55% | 20% | 16% | 20% | | | 2% | | | 26% | 17.10% |
| Sewage sludge | 5% | 65% | 70% | | 14% | 14% | | | | | | | 72.00% |
| Textiles / Carpet and Underlay | 5% | 65% | 20% | 0% | 15% | 15% | 0% | 0% | 0% | 0% | | | 69.68% |
| Wood | 5% | 65% | 17% | 26% | 12% | 42% | | | | | | | 21.00% |
| Sanitary | 5% | 65% | 65% | 0% | 0% | 47% | 0% | 0% | 0% | 0% | | | 52.70% |
| Other | 5% | 65% | | | | | | | | | | | 100.00% |

Table A 3.5.3 Amount of waste landfilled (1945 to 2017)

| Year | England | Scotland | Wales | Northern Ireland | Total | Year | England | Scotland | Wales | Northern Ireland | Total |
|------|---------|----------|-------|------------------|-------|------|---------|----------|-------|------------------|-------|
| 1945 | 70.9 | 9.0 | 4.6 | 2.3 | 86.9 | 1982 | 77.1 | 8.5 | 4.6 | 2.5 | 92.7 |
| 1946 | 71.2 | 9.0 | 4.6 | 2.4 | 87.2 | 1983 | 77.1 | 8.5 | 4.6 | 2.6 | 92.8 |
| 1947 | 71.5 | 9.0 | 4.6 | 2.4 | 87.4 | 1984 | 77.2 | 8.5 | 4.6 | 2.6 | 92.8 |
| 1948 | 71.7 | 9.0 | 4.6 | 2.4 | 87.7 | 1985 | 77.2 | 8.4 | 4.6 | 2.6 | 92.8 |
| 1949 | 72.3 | 9.0 | 4.6 | 2.4 | 88.4 | 1986 | 77.3 | 8.4 | 4.6 | 2.6 | 92.9 |
| 1950 | 72.9 | 9.1 | 4.6 | 2.4 | 89.1 | 1987 | 77.3 | 8.3 | 4.6 | 2.6 | 92.9 |
| 1951 | 73.6 | 9.1 | 4.6 | 2.5 | 89.8 | 1988 | 77.4 | 8.3 | 4.6 | 2.6 | 92.9 |
| 1952 | 74.2 | 9.1 | 4.7 | 2.5 | 90.5 | 1989 | 77.4 | 8.3 | 4.6 | 2.6 | 92.9 |
| 1953 | 74.2 | 9.1 | 4.7 | 2.5 | 90.4 | 1990 | 77.7 | 8.3 | 4.7 | 2.6 | 93.3 |
| 1954 | 74.2 | 9.0 | 4.6 | 2.4 | 90.3 | 1991 | 77.7 | 11.3 | 4.7 | 2.6 | 96.3 |
| 1955 | 74.2 | 9.0 | 4.6 | 2.4 | 90.3 | 1992 | 77.7 | 12.2 | 4.7 | 2.6 | 97.2 |
| 1956 | 75.7 | 9.1 | 4.7 | 2.5 | 92.0 | 1993 | 77.6 | 14.0 | 4.6 | 2.6 | 98.8 |
| 1957 | 77.2 | 9.2 | 4.8 | 2.5 | 93.7 | 1994 | 77.6 | 15.9 | 4.6 | 2.6 | 100.7 |
| 1958 | 78.6 | 9.3 | 4.8 | 2.6 | 95.4 | 1995 | 81.8 | 15.0 | 4.9 | 2.8 | 104.5 |
| 1959 | 80.1 | 9.5 | 4.9 | 2.6 | 97.1 | 1996 | 80.7 | 15.0 | 4.8 | 2.8 | 103.3 |
| 1960 | 81.5 | 9.6 | 5.0 | 2.6 | 98.8 | 1997 | 81.1 | 14.0 | 4.8 | 2.8 | 102.7 |
| 1961 | 81.1 | 9.7 | 4.9 | 2.7 | 98.4 | 1998 | 75.0 | 11.9 | 4.5 | 2.6 | 93.9 |
| 1962 | 80.9 | 9.6 | 4.9 | 2.7 | 98.0 | 1999 | 69.3 | 10.9 | 4.1 | 2.4 | 86.6 |
| 1963 | 84.5 | 10.0 | 5.1 | 2.8 | 102.4 | 2000 | 67.4 | 11.2 | 4.0 | 2.3 | 84.9 |
| 1964 | 84.6 | 9.9 | 5.1 | 2.8 | 102.4 | 2001 | 71.4 | 8.9 | 4.2 | 2.4 | 86.9 |
| 1965 | 85.7 | 10.0 | 5.1 | 2.8 | 103.6 | 2002 | 66.8 | 8.2 | 3.9 | 2.3 | 81.3 |
| 1966 | 85.3 | 9.9 | 5.1 | 2.8 | 103.2 | 2003 | 65.4 | 7.9 | 3.8 | 2.2 | 79.4 |
| 1967 | 85.0 | 9.8 | 5.1 | 2.8 | 102.7 | 2004 | 64.9 | 7.8 | 3.8 | 2.2 | 78.7 |
| 1968 | 84.8 | 9.7 | 5.1 | 2.8 | 102.4 | 2005 | 60.0 | 7.1 | 3.5 | 2.0 | 72.6 |
| 1969 | 84.0 | 9.6 | 5.0 | 2.8 | 101.4 | 2006 | 61.7 | 7.1 | 4.0 | 2.0 | 74.8 |
| 1970 | 83.8 | 9.5 | 5.0 | 2.8 | 101.0 | 2007 | 60.7 | 7.4 | 3.2 | 1.9 | 73.2 |
| 1971 | 82.8 | 9.3 | 4.9 | 2.7 | 99.8 | 2008 | 53.9 | 6.1 | 2.9 | 1.6 | 64.5 |
| 1972 | 81.8 | 9.2 | 4.8 | 2.7 | 98.5 | 2009 | 44.0 | 4.7 | 2.5 | 1.1 | 52.3 |
| 1973 | 81.3 | 9.1 | 4.8 | 2.7 | 97.9 | 2010 | 43.6 | 4.6 | 2.3 | 1.0 | 51.4 |

| Year | England | Scotland | Wales | Northern Ireland | Total | Year | England | Scotland | Wales | Northern Ireland | Total |
|------|---------|----------|-------|------------------|-------|------|---------|----------|-------|------------------|-------|
| 1974 | 79.9 | 9.0 | 4.8 | 2.6 | 96.3 | 2011 | 44.7 | 4.7 | 2.2 | 1.0 | 52.5 |
| 1975 | 80.1 | 9.0 | 4.8 | 2.6 | 96.5 | 2012 | 41.8 | 4.5 | 2.2 | 1.1 | 49.6 |
| 1976 | 78.8 | 8.8 | 4.7 | 2.6 | 94.9 | 2013 | 41.1 | 4.1 | 2.2 | 1.1 | 48.4 |
| 1977 | 78.4 | 8.8 | 4.7 | 2.6 | 94.5 | 2014 | 41.3 | 4.1 | 1.5 | 1.3 | 48.2 |
| 1978 | 78.8 | 8.8 | 4.7 | 2.6 | 95.0 | 2015 | 43.9 | 4.2 | 1.3 | 1.6 | 51.0 |
| 1979 | 78.7 | 8.8 | 4.7 | 2.6 | 94.7 | 2016 | 44.7 | 3.7 | 2.0 | 1.9 | 52.3 |
| 1980 | 78.6 | 8.7 | 4.7 | 2.6 | 94.6 | 2017 | 45.4 | 3.8 | 1.8 | 1.7 | 52.8 |
| 1981 | 77.0 | 8.5 | 4.6 | 2.5 | 92.7 | | | | | | |

A 3.5.1.2 Methane emissions

The right-most column of **Table A 3.5.4** shows the current estimate of methane emitted from UK landfills, according to the approach outlined in **Chapter 7**, taking account of recovery and oxidation.

Table A 3.5.4 Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted

| Year | Waste Landfilled | Methane generated | Methane captured | | Methane used for power generation | | Methane flared | | Residual methane oxidised | | Methane emitted | |
|------|------------------|-------------------|------------------|----|-----------------------------------|----|----------------|----|---------------------------|-----|-----------------|-----|
| | Mt | Kt | kt | % | kt | % | kt | % | kt | % | kt | % |
| 1990 | 93.25 | 2,709 | 33 | 1% | 33 | 1% | 0 | 0% | 268 | 10% | 2408 | 89% |
| 1991 | 96.32 | 2,752 | 50 | 2% | 50 | 2% | 0 | 0% | 270 | 10% | 2432 | 88% |
| 1992 | 97.18 | 2,797 | 90 | 3% | 90 | 3% | 0 | 0% | 271 | 10% | 2436 | 87% |
| 1993 | 98.85 | 2,837 | 107 | 4% | 107 | 4% | 0 | 0% | 273 | 10% | 2457 | 87% |
| 1994 | 100.75 | 2,878 | 124 | 4% | 124 | 4% | 0 | 0% | 275 | 10% | 2479 | 86% |
| 1995 | 104.50 | 2,939 | 135 | 5% | 135 | 5% | 0 | 0% | 280 | 10% | 2524 | 86% |
| 1996 | 103.26 | 2,983 | 170 | 6% | 170 | 6% | 0 | 0% | 281 | 9% | 2532 | 85% |
| 1997 | 102.70 | 3,022 | 218 | 7% | 218 | 7% | 0 | 0% | 280 | 9% | 2524 | 84% |
| 1998 | 93.85 | 3,038 | 278 | 9% | 278 | 9% | 0 | 0% | 276 | 9% | 2484 | 82% |

| Year | Waste Landfilled | Methane generated | Methane captured | | Methane used for power generation | | Methane flared | | Residual methane oxidised | | Methane emitted | |
|------|------------------|-------------------|------------------|-----|-----------------------------------|-----|----------------|----|---------------------------|----|-----------------|-----|
| | Mt | Kt | kt | % | kt | % | kt | % | kt | % | kt | % |
| 1999 | 86.63 | 3,032 | 394 | 13% | 394 | 13% | 0 | 0% | 264 | 9% | 2374 | 78% |
| 2000 | 84.85 | 3,028 | 500 | 17% | 500 | 17% | 0 | 0% | 253 | 8% | 2275 | 75% |
| 2001 | 86.92 | 3,040 | 566 | 19% | 566 | 19% | 0 | 0% | 247 | 8% | 2227 | 73% |
| 2002 | 81.28 | 3,021 | 599 | 20% | 598 | 20% | 1 | 0% | 242 | 8% | 2180 | 72% |
| 2003 | 79.38 | 2,981 | 723 | 24% | 723 | 24% | 0 | 0% | 226 | 8% | 2032 | 68% |
| 2004 | 78.71 | 2,937 | 874 | 30% | 874 | 30% | 0 | 0% | 206 | 7% | 1857 | 63% |
| 2005 | 72.59 | 2,870 | 926 | 32% | 926 | 32% | 0 | 0% | 194 | 7% | 1750 | 61% |
| 2006 | 74.83 | 2,753 | 950 | 35% | 944 | 34% | 6 | 0% | 180 | 7% | 1622 | 59% |
| 2007 | 73.21 | 2,645 | 989 | 37% | 987 | 37% | 2 | 0% | 166 | 6% | 1490 | 56% |
| 2008 | 64.51 | 2,528 | 1065 | 42% | 980 | 39% | 85 | 3% | 146 | 6% | 1316 | 52% |
| 2009 | 52.33 | 2,400 | 1112 | 46% | 1015 | 42% | 97 | 4% | 127 | 5% | 1159 | 48% |
| 2010 | 51.38 | 2,278 | 1200 | 53% | 1066 | 47% | 134 | 6% | 108 | 5% | 971 | 43% |
| 2011 | 52.54 | 2,159 | 1181 | 55% | 1075 | 50% | 106 | 5% | 98 | 5% | 881 | 41% |
| 2012 | 49.55 | 2,041 | 1127 | 55% | 1042 | 51% | 85 | 4% | 91 | 4% | 822 | 40% |
| 2013 | 48.43 | 1,929 | 1152 | 60% | 1035 | 54% | 117 | 6% | 78 | 4% | 699 | 36% |
| 2014 | 48.15 | 1,820 | 1151 | 63% | 1007 | 55% | 145 | 8% | 67 | 4% | 602 | 33% |
| 2015 | 50.98 | 1,716 | 1065 | 62% | 974 | 57% | 91 | 5% | 65 | 4% | 586 | 34% |
| 2016 | 52.31 | 1,627 | 1007 | 62% | 941 | 58% | 67 | 4% | 62 | 4% | 558 | 34% |
| 2017 | 52.76 | 1544 | 918 | 59% | 857 | 56% | 61 | 4% | 63 | 4% | 563 | 36% |

Notes

- Methane generated is based on the MELMod model.
- Methane captured is the sum of methane used for power generation and methane flared.

- c. Methane used for power generation is calculated from official figures on landfill gas electricity generation (Digest of UK Energy Statistics (BEIS, 2016), in GWh/year, assuming a net calorific value for methane of 50 GJ/tonnes and a conversion efficiency between methane use and electricity export of 30% rising to 36%, which includes parasitic losses and on-site use of electricity, e.g. for gas blowers, leachate treatment and site offices.
- d. Methane flared is calculated from site-specific data provided by the Environment Agency at regulated sites for 2009 to 2013, from SEPA for 2013, from a study carried out during 2014, and from site-specific data provided voluntarily by site operators.
- e. Methane oxidised is based on the IPCC default oxidation factor of 10%, applied to methane remaining after subtraction of the amount captured.
- f. Methane emitted = (methane generated – methane captured) x (1-oxidation factor).

A 3.5.1.3 Overseas Territories and Crown Dependencies

For the overseas territories and crown dependencies, the IPCC landfill model is used. Where available, country-specific waste generation and composition data have been applied and appropriate defaults chosen e.g. taking into account climatic variation. There are no landfill emissions for Gibraltar as waste is exported. **Table A 3.5.5** below gives the parameters used.

Table A 3.5.5 Parameters used in landfill emission estimates for overseas territories and crown dependencies

| Region | Methodology | Activity data | MCF | DOC |
|-------------|---|---|--|---------------------|
| Guernsey | IPCC Landfill Model | 2005 onwards: total MSW to landfill data and percentage that is plastics, other inert. Prior to 2005: flat-lined 2005 data | IPCC default values; waste management type is unmanaged, deep (results from expert consultation, 2014) | IPCC default values |
| Jersey | N/A, all MSW is incinerated for energy from waste | N/A | N/A | N/A |
| Gibraltar | N/A, all MSW used to be incinerated, now all waste is exported to be landfilled in Spain. | N/A | N/A | N/A |
| Isle of Man | IPCC Landfill Model | 2004 onwards: all waste incinerated for energy from waste. Prior to 2004: population and IPCC default waste per capita for Western Europe | IPCC default values; waste management type is 50% unmanaged, deep and 50% managed, semi-aerobic (results from expert consultation, 2014) | IPCC default values |

| Region | Methodology | Activity data | MCF | DOC |
|------------------|---------------------|---|--|---------------------|
| Bermuda | IPCC Landfill Model | Total MSW to landfill (Environmental Statistics Compendium) | IPCC default values; no information on management system so assume unmanaged deep | IPCC default values |
| Cayman Islands | IPCC Landfill Model | 2000 onwards: Total MSW to landfill (Department of Environmental Health). Prior to 2000: flat-lined 2000 data | IPCC default values; landfill sites are lined and managed to some degree, but with limited information, "Uncategorised" considered appropriate | IPCC default values |
| Falkland Islands | IPCC Landfill Model | 1998: Halcrow Report. Other years: flat-lined after advice in personal communication from environmental officer | IPCC default values; waste management type is unmanaged, shallow (results from expert consultation, 2014) | IPCC default values |

A 3.5.2 Biological Treatment of Solid Waste (5B)**Table A 3.5.6 Activity Data: Inputs in the composting process**

| Year | Composting (Non-household) (Mg) | Composting (Household) (Mg) | MBT - Composting (Mg) |
|------|------------------------------------|--------------------------------|-----------------------|
| 1990 | 0 | 181,322 | 0 |
| 1991 | 19,283 | 181,756 | 0 |
| 1992 | 33,194 | 182,146 | 0 |
| 1993 | 48,000 | 182,638 | 0 |
| 1994 | 64,000 | 183,213 | 0 |
| 1995 | 140,000 | 183,654 | 0 |
| 1996 | 220,000 | 184,132 | 0 |
| 1997 | 315,000 | 184,643 | 0 |
| 1998 | 675,000 | 185,364 | 0 |
| 1999 | 833,044 | 185,979 | 0 |
| 2000 | 1,034,000 | 186,676 | 0 |
| 2001 | 1,663,852 | 187,441 | 67,882 |
| 2002 | 1,828,000 | 187,373 | 62,537 |
| 2003 | 1,953,000 | 187,300 | 57,192 |
| 2004 | 2,667,000 | 238,235 | 37,179 |
| 2005 | 3,424,000 | 289,058 | 88,917 |
| 2006 | 4,090,000 | 339,909 | 110,618 |
| 2007 | 4,459,000 | 390,804 | 542,678 |
| 2008 | 4,285,000 | 441,622 | 629,269 |
| 2009 | 5,265,711 | 492,393 | 438,011 |
| 2010 | 5,444,092 | 496,077 | 1,282,060 |
| 2011 | 6,053,273 | 500,008 | 1,898,570 |
| 2012 | 5,850,257 | 503,005 | 1,719,118 |
| 2013 | 5,867,640 | 505,833 | 2,138,366 |
| 2014 | 5,954,185 | 509,350 | 2,439,341 |

| Year | Composting (Non-household) (Mg) | Composting (Household) (Mg) | MBT - Composting (Mg) |
|------|------------------------------------|--------------------------------|-----------------------|
| 2015 | 6,010,218 | 513,040 | 2,740,317 |
| 2016 | 6,135,538 | 517,007 | 3,041,292 |
| 2017 | 6,286,717 | 519,898 | 3,342,268 |

Table A 3.5.7 Activity Data: Inputs in the anaerobic digestion process

| Year | Anaerobic digestion – non agricultural residue (Mg) | Anaerobic digestion - MBT (Mg) |
|------|--|-----------------------------------|
| 1990 | 0 | 0 |
| 1991 | 1,678 | 0 |
| 1992 | 1,678 | 0 |
| 1993 | 1,678 | 0 |
| 1994 | 5,435 | 0 |
| 1995 | 5,435 | 0 |
| 1996 | 5,435 | 0 |
| 1997 | 6,061 | 0 |
| 1998 | 6,061 | 0 |
| 1999 | 6,061 | 0 |
| 2000 | 6,061 | 0 |
| 2001 | 6,061 | 16,970 |
| 2002 | 56,155 | 15,634 |
| 2003 | 56,155 | 14,298 |
| 2004 | 87,513 | 37,179 |
| 2005 | 191,153 | 17,783 |
| 2006 | 203,676 | 27,655 |
| 2007 | 223,714 | 40,847 |
| 2008 | 247,633 | 66,664 |
| 2009 | 370,988 | 365,570 |
| 2010 | 721,666 | 72,262 |

| Year | Anaerobic digestion – non agricultural residue (Mg) | Anaerobic digestion - MBT (Mg) |
|------|---|--------------------------------|
| 2011 | 1,286,593 | 104,815 |
| 2012 | 1,842,418 | 795,961 |
| 2013 | 2,850,234 | 600,692 |
| 2014 | 4,286,018 | 703,455 |
| 2015 | 5,672,507 | 806,218 |
| 2016 | 7,268,381 | 908,982 |
| 2017 | 7,772,206 | 1,011,745 |

A 3.5.3 Waste Incineration (5C)

Table A 3.5.8 Activity Data: UK Waste Incineration

| Year | Municipal Waste Incineration ^a (Mt) | Clinical Waste Incineration (Mt) | Chemical Waste Incineration (Mt) | Sewage Sludge Incineration (Mt) |
|------|--|----------------------------------|----------------------------------|---------------------------------|
| 1990 | 2.093 | 0.350 | 0.290 | 0.075 |
| 1991 | 2.069 | 0.350 | 0.290 | 0.069 |
| 1992 | 1.945 | 0.330 | 0.290 | 0.072 |
| 1993 | 1.677 | 0.310 | 0.290 | 0.084 |
| 1994 | 1.148 | 0.290 | 0.289 | 0.072 |
| 1995 | 0.996 | 0.270 | 0.289 | 0.082 |
| 1996 | 1.062 | 0.250 | 0.288 | 0.088 |
| 1997 | - | 0.230 | 0.287 | 0.081 |
| 1998 | - | 0.236 | 0.287 | 0.185 |
| 1999 | - | 0.242 | 0.286 | 0.189 |
| 2000 | - | 0.248 | 0.285 | 0.194 |
| 2001 | - | 0.254 | 0.285 | 0.198 |
| 2002 | - | 0.260 | 0.284 | 0.203 |
| 2003 | - | 0.221 | 0.258 | 0.207 |
| 2004 | - | 0.182 | 0.232 | 0.212 |
| 2005 | - | 0.143 | 0.206 | 0.216 |
| 2006 | - | 0.105 | 0.180 | 0.220 |
| 2007 | - | 0.111 | 0.171 | 0.215 |

| Year | Municipal Waste Incineration ^a (Mt) | Clinical Waste Incineration (Mt) | Chemical Waste Incineration (Mt) | Sewage Sludge Incineration (Mt) |
|------|--|----------------------------------|----------------------------------|---------------------------------|
| 2008 | - | 0.115 | 0.143 | 0.192 |
| 2009 | - | 0.121 | 0.135 | 0.199 |
| 2010 | - | 0.114 | 0.148 | 0.231 |
| 2011 | - | 0.107 | 0.147 | 0.224 |
| 2012 | - | 0.107 | 0.145 | 0.209 |
| 2013 | - | 0.101 | 0.164 | 0.204 |
| 2014 | - | 0.104 | 0.172 | 0.177 |
| 2015 | - | 0.096 | 0.166 | 0.170 |
| 2016 | - | 0.095 | 0.169 | 0.148 |
| 2017 | - | 0.095 | 0.139 | 0.144 |

a Note that all MSW incinerators were either closed or converted to extract power by 1997. In the latter case they were then considered to be power generation and so emissions were reported in 1A1a.

Table A 3.5.9 Emissions Data: UK Waste Incineration

| Year | Chemical Waste Incineration | Accidental Fires | MSW Incineration ^a | Clinical Waste Incineration | Sewage Sludge Incineration | Total |
|---|-----------------------------|------------------|-------------------------------|-----------------------------|----------------------------|---------|
| Carbon Dioxide (kt CO₂) | | | | | | |
| 1990 | 381.8 | NE | 606.0 | 308.0 | NA | 1,295.7 |
| 1995 | 380.0 | NE | 357.0 | 237.6 | NA | 974.7 |
| 2000 | 304.3 | NE | NO | 218.2 | NA | 522.6 |
| 2005 | 275.9 | NE | NO | 126.2 | NA | 402.2 |
| 2010 | 168.6 | NE | NO | 100.6 | NA | 269.2 |
| 2012 | 170.6 | NE | NO | 93.8 | NA | 264.3 |
| 2013 | 171.7 | NE | NO | 89.3 | NA | 260.9 |
| 2014 | 174.0 | NE | NO | 91.7 | NA | 265.7 |
| 2015 | 164.0 | NE | NO | 84.3 | NA | 248.3 |
| 2016 | 183.3 | NE | NO | 83.4 | NA | 266.7 |
| 2017 | 174.5 | NE | NO | 83.9 | NA | 258.4 |

| Year | Chemical Waste Incineration | Accidental Fires | MSW Incineration ^a | Clinical Waste Incineration | Sewage Sludge Incineration | Total |
|--|-----------------------------|------------------|-------------------------------|-----------------------------|----------------------------|-------|
| Methane (kt CH₄) | | | | | | |
| 1990 | NE | 1.009 | 4.175 | 0.007 | 0.029 | 5.311 |
| 1995 | NE | 0.984 | 1.987 | 0.005 | 0.032 | 3.100 |
| 2000 | NE | 0.772 | NO | 0.005 | 0.076 | 0.916 |
| 2005 | NE | 0.704 | NO | 0.003 | 0.084 | 0.834 |
| 2010 | NE | 0.347 | NO | 0.002 | 0.090 | 0.462 |
| 2012 | NE | 0.331 | NO | 0.002 | 0.082 | 0.437 |
| 2013 | NE | 0.290 | NO | 0.002 | 0.080 | 0.399 |
| 2014 | NE | 0.277 | NO | 0.002 | 0.069 | 0.375 |
| 2015 | NE | 0.283 | NO | 0.002 | 0.066 | 0.379 |
| 2016 | NE | 0.287 | NO | 0.002 | 0.058 | 0.374 |
| 2017 | NE | 0.281 | NO | 0.002 | 0.056 | 0.368 |
| Nitrous oxide (kt N₂O) | | | | | | |
| 1990 | 0.029 | NE | 0.058 | 0.011 | 0.060 | 0.155 |
| 1995 | 0.029 | NE | 0.033 | 0.008 | 0.066 | 0.129 |
| 2000 | 0.029 | NE | NO | 0.007 | 0.155 | 0.191 |
| 2005 | 0.021 | NE | NO | 0.004 | 0.173 | 0.198 |
| 2010 | 0.015 | NE | NO | 0.003 | 0.185 | 0.203 |
| 2012 | 0.014 | NE | NO | 0.003 | 0.167 | 0.185 |
| 2013 | 0.016 | NE | NO | 0.003 | 0.163 | 0.183 |
| 2014 | 0.017 | NE | NO | 0.003 | 0.141 | 0.162 |
| 2015 | 0.017 | NE | NO | 0.003 | 0.136 | 0.156 |
| 2016 | 0.017 | NE | NO | 0.003 | 0.118 | 0.138 |
| 2017 | 0.014 | NE | NO | 0.003 | 0.115 | 0.132 |

a Note that all MSW incinerators were either closed or converted to extract power by 1997. In the latter case they were then considered to be power generation and so emissions were reported in 1A1a.

A 3.5.4 Wastewater Handling (5D)**A 3.5.4.1 5D1 Domestic and Commercial Waste Water Handling and Sludge Disposal****Table A 3.5.10 UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data**

| Treatment/disposal route | | unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------------|-------------------|---------|------|------|------|------|------|------|------|------|------|------|
| Total Sludge | | kt tds | 1634 | 1657 | 1682 | 1768 | 1666 | 1598 | 1596 | 1597 | 1659 | 1669 |
| Population Equivalent | | million | 68.3 | 69.2 | 70.2 | 70.5 | 69.3 | 72.9 | 72.9 | 73.0 | 73.2 | 73.4 |
| Additional Treatment | Digested | kt tds | 402 | 433 | 468 | 810 | 835 | 694 | 710 | 657 | 673 | 660 |
| | Advanced Digested | kt tds | 101 | 107 | 115 | 329 | 373 | 295 | 406 | 454 | 556 | 615 |
| | Composted | kt tds | 7 | 8 | 8 | 15 | 25 | 48 | 27 | 7 | 18 | 14 |
| Disposal route | Farmland | kt tds | 508 | 547 | 590 | 1216 | 1282 | 1287 | 1332 | 1422 | 1434 | 1479 |
| | Landfill | kt tds | 160 | 153 | 110 | 131 | 35 | 6 | 4 | 7 | 9 | 8 |
| | Incineration | kt tds | 68 | 80 | 211 | 252 | 238 | 252 | 232 | 161 | 198 | 168 |
| | Sea | kt tds | 782 | 721 | 611 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Composted | kt tds | 2 | 2 | 2 | 13 | 23 | 53 | 28 | 7 | 18 | 14 |
| | Land Reclamation | kt tds | 31 | 30 | 30 | 96 | 44 | 0 | 0 | 0 | 0 | 0 |
| | Other | kt tds | 84 | 124 | 129 | 61 | 44 | 0 | 0 | 0 | 0 | 0 |

Where tds is total dissolvable solids; this is assumed to be comparable to Biochemical Oxygen Demand (BOD)

Table A 3.5.11 UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors

| Treatment/disposal route | | unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---|-------------------------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Mechanical treatment and storage ¹ | | kt/Mt tds | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 | 2.70 |
| Additional Treatment | Digested ² | kt/Mt tds | 16.54 | 16.29 | 16.47 | 16.95 | 16.50 | 16.01 | 15.71 | 15.63 | 13.30 | 13.26 |
| | Advanced Digested | kt/Mt tds | 4.54 | 4.54 | 4.54 | 4.54 | 4.54 | 4.56 | 4.52 | 4.52 | 4.57 | 4.57 |
| | Composted | kt/Mt tds | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 | 10.00 |
| Disposal route | Farmland ³ | kt/Mt tds | 1.36 | 1.36 | 1.36 | 1.44 | 1.41 | 1.45 | 1.31 | 1.30 | 1.28 | 1.22 |
| | Landfill | kt/Mt tds | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 | 15.09 |
| | Incineration | kt/Mt tds | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| | Sea ⁴ | kt/Mt tds | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 | 60.00 |
| | Composted | kt/Mt tds | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 |
| | Land Reclamation ⁵ | kt/Mt tds | 1.36 | 1.36 | 1.36 | 1.36 | 1.36 | 1.42 | 1.29 | 1.29 | 1.27 | 1.22 |
| | Other ⁶ | kt/Mt tds | 1.19 | 1.19 | 1.19 | 1.19 | 1.19 | 1.25 | 1.14 | 1.22 | 1.20 | 1.16 |
| Total ⁷ | | kt/Mt tds | 29.82 | 28.55 | 30.99 | 13.62 | 13.61 | 12.04 | 12.14 | 11.68 | 10.94 | 10.87 |

1. All waste is mechanically treated and stored, so the emission factor is applied to total sludge.

2. Implied emission factor after methane capture.

3. Emission factor varies depending on how the waste is treated.

4. Not an IEF, this is the default IPCC factor for sea, river and lake discharge.

5. Land reclamation hasn't got associated reported emissions, so the factor is based on a weighted average of other waste to land (farmland, composting) IEFs.

6. Other hasn't got associated reported emissions, the factor is based on a weighted average of all other disposal IEFs.

7. For information, IEF when dividing total emissions by total activity.

Table A 3.5.11 UK Domestic and Commercial Waste Water Treatment (5D1) Emission Estimates (kt CH₄)

| Treatment/disposal route | | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|----------------------------------|------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Mechanical treatment and storage | | 4.41 | 4.47 | 4.54 | 4.77 | 4.50 | 4.31 | 4.31 | 4.31 | 4.48 | 4.51 |
| Additional Treatment | Digested | 6.65 | 7.06 | 7.71 | 13.74 | 13.77 | 11.11 | 11.15 | 10.27 | 8.95 | 8.75 |
| | AdvancedDigested | 0.46 | 0.49 | 0.52 | 1.49 | 1.69 | 1.34 | 1.84 | 2.06 | 2.55 | 2.81 |
| | Composted | 0.07 | 0.08 | 0.08 | 0.15 | 0.25 | 0.48 | 0.27 | 0.07 | 0.18 | 0.14 |
| Disposal route | Farmland | 0.68 | 0.73 | 0.79 | 1.72 | 1.82 | 1.90 | 1.76 | 1.85 | 1.85 | 1.82 |
| | Landfill | 2.41 | 2.30 | 1.66 | 1.97 | 0.53 | 0.09 | 0.06 | 0.10 | 0.14 | 0.12 |
| | Incineration | - | - | - | - | - | - | - | - | - | - |
| | Sea | 33.92 | 31.99 | 36.63 | - | - | - | - | - | - | - |
| | Composted | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.03 | 0.02 | 0.00 | 0.01 | 0.01 |
| | LandReclamation | 0.04 | 0.04 | 0.04 | 0.13 | 0.06 | - | - | - | - | - |
| | Other | 0.10 | 0.15 | 0.15 | 0.07 | 0.05 | - | - | - | - | - |
| Total | | 48.74 | 47.31 | 52.13 | 24.09 | 22.68 | 19.24 | 19.39 | 18.66 | 18.14 | 18.14 |

Table A 3.5.12 UK Private Waste Water Management System Emission Estimates Parameters (5D1)

| Data | Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--|------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Estimated population connected to private waste water management systems | population (thousands) | 1417.18 | 1440.63 | 1438.67 | 1465.29 | 1525.78 | 1562.08 | 1579.31 | 1584.43 | 1598.92 | 1605.40 |
| BOD value applied | g/person/day | 66.33 | 66.33 | 66.33 | 68.73 | 65.90 | 60.09 | 59.96 | 59.96 | 62.07 | 62.30 |

A 3.5.4.2 5D2 Industrial Waste Water Handling and Sludge Disposal**Table A 3.5.13 UK Industrial Waste Water Treatment Activity Data (5D2)**

| Sector | Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--|------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Organic chemical production | Mt | 1.617 | 1.617 | 1.751 | 1.752 | 1.487 | 1.539 | 1.581 | 1.668 | 1.575 | 1.616 |
| Milk-processing | million PE | 1.464 | 1.464 | 1.464 | 0.625 | 0.629 | 0.733 | 0.791 | 0.803 | 0.788 | 0.792 |
| Manufacture of fruit and vegetable products | million PE | 1.145 | 1.145 | 1.145 | 1.094 | 1.094 | 0.988 | 1.003 | 1.134 | 1.275 | 1.343 |
| Potato-processing | million PE | 0.302 | 0.302 | 0.302 | 0.289 | 0.289 | 0.261 | 0.265 | 0.299 | 0.336 | 0.355 |
| Meat industry | million PE | 0.623 | 0.623 | 0.623 | 0.618 | 0.648 | 0.634 | 0.696 | 0.723 | 0.717 | 0.760 |
| Breweries | million PE | 0.094 | 0.094 | 0.094 | 0.097 | 0.096 | 0.098 | 0.102 | 0.102 | 0.103 | 0.104 |
| Production of alcohol and alcoholic beverages | million PE | 1.931 | 1.931 | 1.931 | 1.992 | 1.967 | 2.009 | 2.093 | 2.091 | 2.105 | 2.127 |
| Manufacture of animal feed from plant products | million PE | 0.476 | 0.476 | 0.476 | 0.300 | 0.378 | 0.349 | 0.361 | 0.354 | 0.359 | 0.398 |
| Manufacture of gelatine and of glue from hides, skin and bones | million PE | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 |
| Malt-houses | million PE | 0.207 | 0.207 | 0.207 | 0.213 | 0.210 | 0.215 | 0.224 | 0.224 | 0.225 | 0.228 |
| Fish-processing industry | million PE | 0.018 | 0.018 | 0.018 | 0.006 | 0.006 | 0.006 | 0.007 | 0.007 | 0.007 | 0.007 |
| Total Food and Drink | million PE | 6.273 | 6.273 | 6.273 | 5.247 | 5.329 | 5.306 | 5.555 | 5.750 | 5.929 | 6.126 |

Where PE is population equivalent

A 3.6 UK CROWN DEPENDENCIES AND OVERSEAS TERRITORIES

A 3.6.1 Overview of Data Sources

Fuel use data for Isle of Man, Guernsey and Jersey are assumed to be included in UK national energy statistics (see **Section 1.1.2.2**), so fuel thought to be used in these territories are split out from UK total consumption unless otherwise stated in **Section A 4.2.1**.

Activity data including fuel use data for other territories are obtained from government departments for those territories, specifically:

- The Cayman Islands Government Department of Environment Sustainable Development Unit;
- The Department of Energy Bermuda; and,
- The Falkland Islands Government Policy Unit.

Activity and emissions data estimates from LULUCF sources and sinks have been researched via the FAOSTAT database, to supplement data available from the OTs. The LULUCF data for Cayman Islands from FAOSTAT (FAO, 2018) indicates zero emissions, using Tier 1 methods. The data sources and methodologies used for other sectors are described in the main methodology sections of the NIR.

A 3.6.2 Activity and Emissions Data

Table A 3.6.1 Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO₂ equivalent)

| Sector | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1. Energy | 1.48 | 1.64 | 1.75 | 1.55 | 1.47 | 1.56 | 1.48 | 1.37 | 1.43 | 1.40 |
| 2. Industrial Processes and Product Use | 0.0001 | 0.0050 | 0.0245 | 0.0515 | 0.0824 | 0.0816 | 0.0823 | 0.0812 | 0.0758 | 0.0758 |
| 3. Agriculture | 0.14 | 0.14 | 0.15 | 0.10 | 0.13 | 0.12 | 0.12 | 0.12 | 0.12 | 0.12 |
| 4. LULUCF | -0.02 | -0.04 | -0.04 | -0.03 | -0.03 | -0.02 | -0.03 | -0.01 | -0.03 | -0.03 |
| 5. Waste | 0.16 | 0.16 | 0.17 | 0.17 | 0.15 | 0.14 | 0.14 | 0.14 | 0.13 | 0.13 |
| 7. Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 1.74 | 1.91 | 2.05 | 1.84 | 1.81 | 1.88 | 1.79 | 1.69 | 1.73 | 1.69 |

Table A 3.6.2 Isle of Man, Guernsey and Jersey – Combustion activity data

| Fuel | Fuel Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-----------------------|-----------|---------|---------|---------|---------|----------|----------|----------|----------|----------|----------|
| Aviation spirit | Mt | 0.00813 | 0.00907 | 0.0123 | 0.0135 | 0.00715 | 0.00576 | 0.00569 | 0.00364 | 0.00285 | 0.00181 |
| Aviation turbine fuel | Mt | 0.0803 | 0.072 | 0.0931 | 0.107 | 0.0935 | 0.0855 | 0.0806 | 0.0855 | 0.0881 | 0.0951 |
| Burning oil | Mt | 0.225 | 0.261 | 0.35 | 0.357 | 0.363 | 0.381 | 0.373 | 0.386 | 0.408 | 0.416 |
| Coal | Mt | 0.0345 | 0.0218 | 0.0152 | 0.0103 | 0.000891 | 0.000891 | 0.000891 | 0.000972 | 0.000975 | 0.000978 |
| DERV | Mt | 0.0737 | 0.0937 | 0.133 | 0.118 | 0.125 | 0.131 | 0.132 | 0.135 | 0.139 | 0.139 |
| Fuel oil | Mt | 0.469 | 0.58 | 0.438 | 0.0777 | 0.0889 | 0.222 | 0.158 | 0.0598 | 0.0777 | 0.0454 |
| Gas oil | Mt | 0.164 | 0.174 | 0.162 | 0.181 | 0.119 | 0.0953 | 0.0873 | 0.0855 | 0.088 | 0.0879 |
| LPG | Mth | 11.7 | 13.5 | 46.4 | 25 | 22.4 | 19.1 | 17.4 | 16.9 | 16.2 | 15.3 |
| Lubricants | | | | | | | | | | | |
| MSW | Mt | 0.121 | 0.169 | 0.216 | 0.391 | 0.359 | 0.321 | 0.309 | 0.296 | 0.293 | 0.286 |
| Natural gas | Mth | 0 | 0 | 0 | 106 | 126 | 114 | 124 | 113 | 127 | 125 |
| Petrol | Mt | 0.236 | 0.228 | 0.217 | 0.219 | 0.191 | 0.175 | 0.171 | 0.169 | 0.166 | 0.164 |
| Petroleum wax | | | | | | | | | | | |
| Wood | Mt | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 | 0.00765 |

Table A 3.6.3 Isle of Man, Guernsey and Jersey – Animal numbers

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Dairy | 15,888 | 15,729 | 16,186 | 13,127 | 11,455 | 10,820 | 10,841 | 10,704 | 11,260 | 11,331 |

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------|---------|---------|---------|--------|---------|---------|---------|---------|---------|---------|
| Non-dairy | 28,663 | 28,333 | 29,176 | 16,770 | 28,615 | 26,097 | 27,498 | 26,513 | 26,338 | 24,387 |
| Sheep | 151,764 | 160,228 | 176,259 | 87,537 | 138,251 | 134,501 | 134,310 | 133,666 | 126,057 | 128,927 |
| Pigs | 4,854 | 5,411 | 4,609 | 1,148 | 4,086 | 2,605 | 2,602 | 2,861 | 2,386 | 2,342 |
| Poultry | 84,048 | 46,481 | 46,448 | 58,160 | 54,400 | 57,924 | 58,850 | 62,916 | 60,231 | 62,537 |
| Goats | 333 | 347 | 376 | 141 | 288 | 416 | 477 | 539 | 352 | 326 |
| Horses | 2,785 | 2,785 | 2,785 | 2,822 | 3,236 | 3,265 | 3,163 | 2,891 | 3,258 | 2,705 |

Table A 3.6.4 Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N₂O)

| Territory | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Isle of Man | 39,497 | 40,148 | 43,721 | 27,111 | 37,931 | 34,889 | 36,079 | 35,396 | 35,610 | 34,166 |
| Guernsey | 5,713 | 5,759 | 5,277 | 4,750 | 4,624 | 4,729 | 4,832 | 4,641 | 4,596 | 4,539 |
| Jersey | 6,361 | 6,444 | 6,605 | 5,256 | 5,394 | 5,457 | 5,451 | 5,278 | 5,255 | 5,248 |

Table A 3.6.5 Cayman Islands, Falklands Islands, and Bermuda - Emissions of Direct GHGs (Mt CO₂ equivalent)

| Sector | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---|-----------|---------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1. Energy | 1.23 | 1.25 | 1.4 | 1.59 | 1.58 | 1.55 | 1.64 | 1.69 | 1.51 | 1.61 |
| 2. Industrial Processes and Product Use | 0.0000957 | 0.00274 | 0.0146 | 0.0308 | 0.0418 | 0.0372 | 0.0369 | 0.0363 | 0.0321 | 0.0321 |
| 3. Agriculture | 0.389 | 0.367 | 0.338 | 0.312 | 0.33 | 0.323 | 0.318 | 0.313 | 0.311 | 0.319 |
| 4. LULUCF | 1.62 | 1.62 | 1.75 | 1.93 | 1.96 | 1.91 | 1.99 | 2.04 | 1.86 | 1.96 |
| 5. Waste | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7. Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Sector | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------|------|------|------|------|------|------|------|------|------|------|
| Total | 3.24 | 3.24 | 3.5 | 3.87 | 3.91 | 3.82 | 3.98 | 4.08 | 3.71 | 3.92 |

Table A 3.6.6 Cayman Islands, Falklands Islands, and Bermuda – Combustion activity data

| Fuel | Fuel Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-----------------------|-----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Aviation spirit | Mt | 0.000342 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Aviation turbine fuel | Mt | 0.0816 | 0.0373 | 0.0411 | 0.0566 | 0.071 | 0.0552 | 0.0511 | 0.0507 | 0.0506 | 0.0504 |
| Biogenic products | | | | | | | | | | | |
| Burning oil | Mt | 0.00433 | 0.00535 | 0.00616 | 0.00797 | 0.0072 | 0.00692 | 0.00715 | 0.00744 | 0.00827 | 0.00884 |
| Coal | Mt | 0.147 | 0.127 | 0.103 | 0.232 | 0.0964 | 0.13 | 0.0854 | 0.0856 | 0.0857 | 0.0859 |
| DERV | Mt | 0.145 | 0.152 | 0.266 | 0.303 | 0.315 | 0.314 | 0.315 | 0.314 | 0.314 | 0.316 |
| Fuel oil | Mt | 0.398 | 0.447 | 0.428 | 0.475 | 0.546 | 0.595 | 0.572 | 0.583 | 0.592 | 0.573 |
| Gas oil | Mt | 7.92 | 8.85 | 9.76 | 11.3 | 11.5 | 11.7 | 12.1 | 12.3 | 12.5 | 12.6 |
| LPG | Mth | 0.00042 | 0.204 | 0.204 | 0.204 | 0.192 | 0.163 | 0.166 | 0.174 | 0.167 | 0.162 |
| MSW | Mt | 0.0473 | 0.0585 | 0.0673 | 0.0882 | 0.105 | 0.0905 | 0.0966 | 0.0971 | 0.107 | 0.115 |
| Natural gas | Mth | 0.189 | 0.192 | 0.214 | 0.17 | 0.196 | 0.174 | 0.135 | 0.147 | 0.147 | 0.141 |
| Petrol | Mt | 0 | 0 | 0 | 0 | 0.000609 | 0.00126 | 0.00123 | 0.00135 | 0.00143 | 0.00149 |
| Lubricants | Mt | 0.00025 | 0.000214 | 0.000174 | 0.000413 | 0.000216 | 0.000166 | 0.000187 | 0.000251 | 0.000251 | 0.000268 |
| Petroleum waxes | Mt | 0.000342 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Table A 3.6.7 Cayman Islands, Falklands Islands, and Bermuda – Animal numbers

| Livestock Category | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Dairy Cattle | 2,161 | 1,862 | 1,911 | 1,145 | 868 | 765 | 723 | 675 | 634 | 647 |
| Non-dairy Cattle | 5,256 | 4,861 | 5,077 | 7,845 | 6,360 | 5,795 | 5,609 | 4,748 | 4,503 | 4,913 |
| Sheep | 739,999 | 717,571 | 669,905 | 580,864 | 478,625 | 486,037 | 483,135 | 482,131 | 479,752 | 490,213 |
| Goats | 405 | 867 | 1,286 | 1,704 | 2,251 | 2,080 | 1,891 | 1,812 | 2,031 | 2,337 |
| Horses | 2,217 | 2,069 | 1,703 | 1,417 | 1,269 | 1,192 | 1,202 | 1,223 | 1,252 | 1,244 |
| Swine | 1,116 | 1,174 | 1,376 | 1,384 | 1,233 | 1,248 | 1,242 | 1,058 | 1,234 | 1,301 |
| Poultry | 15,319 | 14,664 | 20,890 | 27,164 | 32,293 | 37,764 | 27,769 | 39,458 | 39,948 | 26,710 |
| Deer | 0 | 0 | 0 | 0 | 184 | 243 | 243 | 243 | 243 | 243 |

Table A 3.6.8 Cayman Islands, Falklands Islands, and Bermuda – Total emissions from Agricultural Soils (kg N₂O)

| Territory | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Bermuda | 2,409 | 2,606 | 2,803 | 2,786 | 2,801 | 2,802 | 2,802 | 2,803 | 2,803 | 2,798 |
| Cayman Islands | 2,698 | 3,212 | 3,691 | 4,173 | 4,446 | 4,566 | 4,342 | 4,359 | 3,916 | 3,862 |
| Falkland Islands | 363,328 | 350,782 | 328,265 | 288,461 | 240,569 | 245,081 | 243,913 | 242,966 | 241,529 | 239,827 |

Table A 3.6.9 Cayman Islands, Falklands Islands, and Bermuda - Amount of synthetic fertilizer applied

| Country | kg N applied |
|------------------|--------------|
| Cayman Islands | 23,080 |
| Falkland Islands | 0 |

| Country | kg N applied |
|---------|--------------|
| Bermuda | 1,480 |

Table A 3.6.10 Gibraltar – Emissions of Direct GHGs (Mt CO₂ equivalent)

| Sector | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1. Energy | 0.219 | 0.216 | 0.242 | 0.278 | 0.288 | 0.319 | 0.308 | 0.331 | 0.331 | 0.322 |
| 2. Industrial Processes and Other Product Use | 0.000 | 0.001 | 0.003 | 0.006 | 0.010 | 0.011 | 0.011 | 0.012 | 0.011 | 0.011 |
| 3. Agriculture | - | - | - | - | - | - | - | - | - | - |
| 4. LULUCF | - | - | - | - | - | - | - | - | - | - |
| 5. Waste | 0.007 | 0.009 | 0.012 | 0.001 | 0.003 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Total | 0.226 | 0.226 | 0.257 | 0.285 | 0.301 | 0.331 | 0.321 | 0.345 | 0.344 | 0.335 |

Table A 3.6.11 Gibraltar – Combustion activity data

| Fuel | Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-----------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Aviation turbine fuel | Mt | 7.90 | 6.06 | 5.27 | 8.43 | 6.55 | 8.12 | 8.64 | 8.73 | 11.34 | 11.85 |
| Charcoal | Mt | - | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Clinical waste | Mt | - | - | - | - | 1.81 | 0.30 | 0.26 | 0.27 | 0.27 | 0.27 |
| DERV | Mt | 0.82 | 0.41 | 0.92 | 2.21 | 2.45 | 2.27 | 2.74 | 3.19 | 3.76 | 3.81 |
| Fuel oil | Mt | 11.69 | 9.72 | 10.59 | 9.36 | 7.17 | - | - | - | - | - |
| Gas oil | Mt | 19.88 | 23.59 | 26.37 | 33.22 | 41.36 | 63.65 | 56.69 | 60.46 | 56.92 | 53.31 |
| LPG | Mth | 4.13 | 4.13 | 4.13 | 4.13 | 4.13 | 4.13 | 4.13 | 4.13 | 4.13 | 4.34 |
| Lubricant | kt | 0.56 | 0.54 | 0.47 | 0.45 | 0.37 | 0.29 | 0.29 | 0.27 | 0.27 | 0.27 |
| MSW | Mt | 16.40 | 19.00 | 24.09 | - | - | - | - | - | - | - |
| Petrol | Mt | 5.95 | 5.12 | 7.14 | 6.93 | 6.32 | 4.63 | 4.77 | 4.91 | 5.08 | 5.08 |

| Fuel | Unit | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-----------------------|------|------|------|------|------|------|------|------|------|-------|-------|
| Aviation turbine fuel | Mt | 7.90 | 6.06 | 5.27 | 8.43 | 6.55 | 8.12 | 8.64 | 8.73 | 11.34 | 11.85 |
| Petroleum waxes | Mt | 0.03 | 0.02 | 0.01 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |

ANNEX 4: National Energy Balance

A 4.1 UK ENERGY BALANCE

The UK energy balance is produced and published annually by the Department of Business, Energy & Industrial Strategy in the Digest of UK Energy Statistics – DUKES. This is available online from:

<https://www.gov.uk/government/collections/digest-of-uk-energy-statistics-dukes>

The aggregate energy balance for the latest year is presented below (Table 1.1 in DUKES). The following sections explain how the energy balance is used for the UK inventory for individual fuel types, and how the data are supplemented with other statistics that may lead to deviations from the DUKES statistics.

The UK energy statistics (detailed breakdown) are presented on a mass basis for liquid and solid fuels, and on a gross energy basis for gaseous fuels (including derived gases). The UK inventory is calculated using these data directly, and for the purposes of reporting in the CRF and NIR, activity data and emission factors are converted to energy units, on a net basis.

The scope of the UK energy balance, as shown below, is fuel use in the United Kingdom and its Crown Dependencies (Jersey, Guernsey and the Isle of Man), as described in the NIR section 1.1.2.2.

The fuel use estimates for Overseas Territories (OTs) are not included within DUKES, and are obtained through direct communications with the respective government contacts in each of the OTs.

Table A 4.1.1 UK Energy Balance for 2017 (thousand tonnes of oil equivalent, gross energy basis)

| | Coal | Manufactured fuel(1) | Primary oils | Petroleum products | Natural gas(2) | Bioenergy & waste(3) | Primary electricity | Electricity | Heat sold | Total |
|---------------------------|---------------|----------------------|----------------|--------------------|----------------|----------------------|---------------------|---------------|--------------|----------------|
| Supply | | | | | | | | | | |
| Production | 1,934 | - | 50,944 | - | 40,019 | 12,924 | 20,924 | - | - | 126,745 |
| Imports | 5,807 | 712 | 58,480 | 36,722 | 45,132 | 3,475 | - | 1,562 | - | 151,891 |
| Exports | -369 | -14 | -42,040 | -25,374 | -10,802 | -431 | - | -293 | - | -79,323 |
| Marine bunkers | - | - | - | -2,596 | - | - | - | - | - | -2,596 |
| Stock change(4) | +2,098 | -2 | +361 | -113 | +1,028 | - | - | - | - | +3,373 |
| Primary supply | 9,470 | 696 | 67,746 | 8,639 | 75,377 | 15,969 | 20,924 | 1,269 | - | 200,090 |
| Statistical difference(5) | -65 | +1 | -66 | -10 | +337 | - | - | -35 | - | +163 |
| Primary demand | 9,535 | 694 | 67,811 | 8,649 | 75,040 | 15,969 | 20,924 | 1,304 | - | 199,927 |
| Transfers | - | - | - | - | - | - | - | - | - | - |
| | - | +10 | -2,476 | +2,483 | +224 | -237 | -5,801 | +5,801 | - | +4 |
| Transformation | -8,134 | 379 | -65,335 | 64,539 | -27,182 | -9,587 | -15,124 | 23,071 | 1,592 | -35,779 |
| Electricity generation | -5,559 | -518 | - | -533 | -24,594 | -9,387 | -15,124 | 23,071 | - | -32,645 |
| Major power producers | -5,545 | - | - | -146 | -22,150 | -4,404 | -15,124 | 20,358 | - | -27,011 |
| Autogenerators | -14 | -518 | - | -387 | -2,445 | -4,983 | - | 2,713 | - | -5,634 |
| Heat generation | -4 | -1 | - | -52 | -2,587 | -200 | - | - | 1,592 | -1,252 |

| | Coal | Manufactured fuel(1) | Primary oils | Petroleum products | Natural gas(2) | Bioenergy & waste(3) | Primary electricity | Electricity | Heat sold | Total |
|----------------------------|----------|----------------------|--------------|--------------------|----------------|----------------------|---------------------|--------------|------------|---------------|
| Petroleum refineries | - | - | -65,795 | 65,691 | - | - | - | - | - | -104 |
| Coke manufacture | -1,435 | 1,351 | - | - | - | - | - | - | - | -84 |
| Blast furnaces | -989 | -596 | - | - | - | - | - | - | - | -1,585 |
| Patent fuel manufacture | -146 | 143 | - | -66 | - | - | - | - | - | -69 |
| Other(7) | - | - | 460 | -501 | - | - | - | - | - | -40 |
| Energy industry use | - | 458 | - | 4,315 | 4,903 | - | - | 2,041 | 322 | 12,040 |
| Electricity generation | - | - | - | - | - | - | - | 1,332 | - | 1,332 |
| Oil and gas extraction | - | - | - | 715 | 4,244 | - | - | 51 | - | 5,010 |
| Petroleum refineries | - | - | - | 3,600 | 92 | - | - | 375 | 322 | 4,390 |
| Coal extraction | - | - | - | - | 6 | - | - | 39 | - | 45 |
| Coke manufacture | - | 182 | - | - | - | - | - | 1 | - | 183 |
| Blast furnaces | - | 276 | - | - | 25 | - | - | 18 | - | 319 |
| Patent fuel manufacture | - | - | - | - | - | - | - | - | - | - |
| Pumped storage | - | - | - | - | - | - | - | 85 | - | 85 |
| Other | - | - | - | - | 536 | - | - | 139 | - | 675 |
| Losses | - | 109 | - | - | 580 | - | - | 2,283 | - | 2,972 |

| | Coal | Manufactured fuel(1) | Primary oils | Petroleum products | Natural gas(2) | Bioenergy & waste(3) | Primary electricity | Electricity | Heat sold | Total |
|----------------------------|--------------|----------------------|--------------|--------------------|----------------|----------------------|---------------------|---------------|--------------|----------------|
| Final consumption | 1,401 | 516 | - | 71,356 | 42,599 | 6,145 | - | 25,851 | 1,270 | 149,139 |
| Industry | 972 | 296 | - | 4,308 | 8,677 | 1,162 | - | 7,964 | 692 | 24,071 |
| Unclassified | - | - | - | 3,380 | 1 | 135 | - | - | - | 3,515 |
| Iron and steel | 23 | 296 | - | 5 | 331 | - | - | 230 | - | 885 |
| Non-ferrous metals | 19 | - | - | 8 | 261 | - | - | 370 | - | 659 |
| Mineral products | 439 | - | - | 174 | 1,222 | 203 | - | 524 | - | 2,562 |
| Chemicals | 43 | - | - | 115 | 1,748 | 5 | - | 1,336 | 277 | 3,524 |
| Mechanical engineering etc | 8 | - | - | 0 | 975 | 2 | - | 556 | - | 1,540 |
| Electrical engineering etc | 3 | - | - | 1 | 275 | - | - | 512 | - | 791 |
| Vehicles | 38 | - | - | 208 | 595 | - | - | 405 | - | 1,246 |
| Food, beverages etc | 54 | - | - | 110 | 1,624 | 47 | - | 945 | 3 | 2,783 |
| Textiles, leather etc | 44 | - | - | 43 | 243 | - | - | 231 | - | 561 |
| Paper, printing etc | 64 | - | - | 31 | 391 | 611 | - | 930 | - | 2,027 |
| Other industries | 232 | - | - | 38 | 609 | 159 | - | 1,801 | 412 | 3,251 |
| Construction | 4 | - | - | 195 | 401 | - | - | 125 | - | 725 |
| Transport (6) | 11 | - | - | 55,051 | - | 997 | - | 411 | - | 56,470 |
| Air | - | - | - | 12,995 | - | - | - | - | - | 12,995 |

| | Coal | Manufactured fuel(1) | Primary oils | Petroleum products | Natural gas(2) | Bioenergy & waste(3) | Primary electricity | Electricity | Heat sold | Total |
|-----------------------|------------|----------------------|--------------|--------------------|----------------|----------------------|---------------------|---------------|------------|---------------|
| Rail | 11 | - | - | 661 | - | - | - | 396 | - | 1,069 |
| Road | - | - | - | 40,468 | - | 997 | - | 15 | - | 41,480 |
| National navigation | - | - | - | 926 | - | - | - | - | - | 926 |
| Pipelines | - | - | - | - | - | - | - | - | - | - |
| Other | 419 | 172 | - | 4,507 | 33,496 | 3,986 | - | 17,476 | 578 | 60,634 |
| Domestic | 392 | 172 | - | 2,472 | 25,540 | 2,216 | - | 9,062 | 260 | 40,116 |
| Public administration | 18 | - | - | 369 | 3,111 | 72 | - | 1,696 | 97 | 5,364 |
| Commercial | 4 | - | - | 883 | 3,868 | 1,173 | - | 6,344 | 220 | 12,493 |
| Agriculture | - | - | - | 474 | 110 | 525 | - | 373 | - | 1,482 |
| Miscellaneous | 5 | - | - | 308 | 866 | - | - | - | - | 1,179 |
| Non energy use | - | 48 | - | 7,490 | 426 | - | - | - | - | 7,964 |

(1) Includes all manufactured solid fuels, benzole, tars, coke oven gas and blast furnace gas.

(2) Includes colliery methane.

(3) Includes geothermal and solar heat.

(4) Stock fall (+), stock rise (-).

(5) Primary supply minus primary demand.

(6) See paragraphs 5.42 regarding electricity use in transport and 6.66 regarding renewables use in transport.

(7) Back-flows from the petrochemical industry.

A 4.2 FUELS DATA

The fuels data are taken from DUKES - the Digest of UK Energy Statistics (BEIS, 2018), so the fuel definitions and the source categories used in the NAEI reflect those in DUKES.

IPCC Guidelines (IPCC, 2006) lists fuels that should be considered when reporting emissions. **Table A 4.2.1** lists the fuels that are used in the GHGI (based on DUKES) and indicates how they relate to the fuels listed in the IPCC Guidelines. In most cases the mapping is obvious but there are a few cases where some explanation is required.

Table A 4.2.1 Mapping of fuels used in IPCC and the NAEI

| | IPCC | NAEI |
|----------|-------------------------------------|--|
| Category | Subcategory | Subcategory |
| Liquid | Motor Gasoline | Petrol |
| | Aviation Gasoline | Aviation Spirit |
| | Jet Kerosene | Aviation Turbine Fuel ¹ (ATF) |
| | Other Kerosene | Burning Oil |
| | Gas/Diesel Oil | Gas Oil/ DERV |
| | Residual Fuel Oil | Fuel Oil |
| | Orimulsion | Orimulsion |
| | Liquefied Petroleum Gases | Liquefied Petroleum Gas (LPG) |
| | Naphtha | Naphtha |
| | Petroleum Coke | Petroleum Coke |
| | Refinery Gas | Other Petroleum Gas (OPG) |
| | Other Oil: Other Petroleum Products | Refinery Miscellaneous |
| | Lubricants | Lubricants |
| Solid | Anthracite | Anthracite |
| | Coking Coal | Coal ² |
| | Other Bituminous Coal | Coal |
| | | Slurry ³ |
| | Coke Oven Coke | Coke |
| | Patent Fuel | Solid Smokeless Fuel (SSF) |
| | Coke Oven Gas | Coke Oven Gas |
| | Blast Furnace Gas | Blast Furnace Gas |

| | IPCC | NAEI |
|-------------|-------------------------------|----------------------------------|
| Category | Subcategory | Subcategory |
| Gas | Natural Gas | Natural Gas |
| | | Sour Gas ⁴ |
| | | Colliery Methane ⁵ |
| Other Fuels | Municipal Solid Waste | Municipal Solid Waste |
| | Industrial Waste: Scrap Tyres | Scrap Tyres |
| | Waste Oils | Waste Oil |
| Peat | Peat | Peat |
| Biomass | Wood/Wood Waste | Wood |
| | Other Primary Solid Biomass | Straw |
| | | Poultry Litter, Meat & bone meal |
| | Landfill Gas | Landfill Gas |
| | Sludge Gas | Sewage Gas |
| | Charcoal | Charcoal |
| | Other liquid biofuels | Liquid Biofuels |
| | Other biogas | Biogas |

1 Includes fuel that is correctly termed jet gasoline.

2 Used in coke ovens.

3 Coal-water slurry used in some power stations

4 Unrefined natural gas used on offshore platforms and some power stations

5 IPCC Guidelines (IPCC, 2006) specifies coal seam methane is included in Natural Gas.

A 4.2.1 Reallocations of energy data and differences from UK energy statistics

The main source of energy consumption data used in the UK inventory is the Digest of UK Energy Statistics (BEIS, 2018). This annual publication gives detailed sectoral energy consumption broken down by fuel type, and covering the entire time period covered by the inventory. In many cases, these data are used directly in the inventory without modification. However, there are instances where the activity data used in the inventory are not based directly on DUKES data, but where alternative data sources provide supplementary data to inform energy use and emission estimates. In general, the UK inventory totals by fuel are kept consistent with the DUKES national totals for each fuel. There are some exceptions where the UK total may be different to that presented in DUKES due to different scopes and reporting requirements.

The reasons for any deviations from use of DUKES data in the inventory are discussed within the source category methodological descriptions in Section 3 of the main report. The main reasons for reallocations or modifications are:

- To account for differences in geographical scope (e.g. to account for energy use in OTs)
- To make best use of EU ETS data (this data is only used indirectly in producing UK energy statistics)

- To utilise other operator reported data (e.g. direct to the Inventory Agency, or via the various regulator's inventories).
- When bottom-up models are available providing fuel consumption data on a more granular level and are considered to be a higher quality estimate by the Inventory Agency.

The fuel reconciliation tables (**Table A 4.2.2 - Table A 4.2.6**) show how the deviations are applied and how the energy data for the major fuels in the UK inventory are reconciled against the energy demand data from DUKES. The tables show:

1. where fuels are re-allocated between sectors, but the overall annual fuel consumption across all UK sectors is kept consistent with the data in DUKES, and
2. where deviations are made to DUKES figures for total UK consumption of a given fuel, and in which source categories these deviations are made.

The Inventory Agency presents data below for the fuel allocations for coal, natural gas, fuel oil, gas oil (including DERV) and petroleum gases (LPG, OPG) for the latest inventory year. Together these fuels constitute the majority of the UK inventory 1A sector emissions total.

Deviations to the energy balance are made in consultation with the authors of the energy statistics.

A 4.2.1.1 Coal

Total industrial coal use within the GHG inventory is consistent with the DUKES total and in the most part, coal use at the sectoral level is consistent with the DUKES data. However, there is an apparent step change in the amount of fuel allocated to other industries within DUKES between 1999 and 2000. In addition, between 1997 and 1999, the total coal use allocated to 1A2f is less than the independent estimates for cement and lime production used within the inventory. Cement and lime production would fall into the 1A2f category for IPCC reporting. Therefore, Inventory Agency estimates have been made to construct a consistent time series for coal use. **Table A 4.2.2** below compares inventory estimates with DUKES estimates for the latest inventory year

Table A 4.2.2 Fuel reconciliation - coal use in the latest year (Mtonnes)

| DUKES Category | DUKES | GHGI | Difference | GHGI category | CRF | Comment |
|-----------------------|-------|-------|------------|-----------------------------------|------|---------|
| Major power producers | 8.702 | 8.702 | 0.000 | Power stations | 1A1a | |
| Blast furnaces | 1.301 | 1.301 | 0.000 | Blast furnaces | 2C1 | |
| Coal extraction | 0.000 | 0.000 | 0.000 | Collieries - combustion | 1A1c | |
| Autogenerators | 0.022 | | | | | |
| | | 0.013 | | Autogenerators | 1A2b | |
| | | 0.010 | | Autogeneration - exported to grid | 1A2b | |

| DUKES Category | DUKES | GHGI | Difference | GHGI category | CRF | Comment |
|-------------------------------------|--------------|--------------|--------------|--|-------------|---|
| | 0.022 | 0.022 | | Subtotal | | |
| Patent fuel manufacture etc. | 0.207 | 0.207 | 0.000 | Solid smokeless fuel production | 1B1b | |
| Coke manufacture | 1.888 | 1.826 | 0.062 | Coke Production | 1B1b | GHGI uses EU ETS data which suggests fewer emissions than in DUKES |
| Industry total | 1.344 | | | | | |
| | | 0.033 | | Iron & steel - combustion plant | 1A2a | |
| | | 0.033 | | Non-Ferrous Metal (combustion) | 1A2b | |
| | | 0.069 | | Chemicals | 1A2c | |
| | | 0.111 | | Pulp, Paper and Print | 1A2d | |
| | | 0.078 | | Food & drink, tobacco | 1A2e | |
| | | 0.580 | | Other industrial combustion | 1A2g | |
| | | 0.580 | | Cement production - combustion | 1A2f | Operator's data used in preference in the GHGI |
| | | 0.051 | | Lime production - non decarbonising | 1A2f | EU ETS used in preference in the GHGI |

| DUKES Category | DUKES | GHGI | Difference | GHGI category | CRF | Comment |
|-----------------------|---------------|---------------|--------------|--|------|--|
| | 1.344 | 1.533 | -0.188 | Subtotal | | |
| Rail | 0.015 | 0.015 | 0.000 | Rail | 1A3c | |
| Domestic - anthracite | 0.193 | 0.193 | 0.000 | Domestic combustion - anthracite | 1A4b | |
| Domestic - coal | 0.342 | 0.342 | | Domestic combustion - UK | 1A4b | |
| | | 0.000 | | Domestic combustion - crown dependencies | 1A4b | |
| | 0.342 | 0.342 | 0.000 | Subtotal | | |
| Agriculture | 0.000 | 0.000 | 0.000 | Agriculture - stationary combustion | 1A4c | |
| Commercial | 0.005 | | | | | |
| Miscellaneous | 0.007 | 0.012 | 0.000 | Miscellaneous industrial/commercial combustion | 1A4a | |
| | 0.012 | 0.012 | 0.000 | Subtotal | | |
| Public administration | 0.156 | 0.030 | 0.126 | Public sector combustion | 1A4a | Differences offset the deviations in the industrial and coke manufacture sectors |
| TOTAL | 14.183 | 14.183 | 0.000 | | | |

Notes: Rows are shaded to help illustrate reconciliation between sectors.

A 4.2.1.2 Natural Gas

Data for natural gas use is largely taken directly from DUKES and the national total is consistent between the inventory and the energy statistics, other than a small additional use of natural gas at a number of (international) gas pipeline inter-connectors and also on the Isle of Man (IoM)

which is added to the inventory, as natural gas use on IoM is not included in DUKES demand totals. Operator estimates for ammonia production (both fuel and feedstock), and ETS data for gas separation plant lead to minor reallocations of the DUKES data, these are summarised below in **Table A 4.2.3**. In addition, the NAEI model doesn't include any accounting for losses compared to DUKES tables.

Table A 4.2.3 Fuel reconciliation – natural gas use in the latest year (Mtherms)

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|-------------------------|-------|------|------------|---|------|---|
| Major power producers | 8790 | 8822 | -33 | Power stations | 1A1a | Isle of Man gas use not offset by DUKES |
| Heat generation | 0 | | 0 | | | |
| Autogenerators | 954 | | | | | |
| | | 398 | | Autogenerators | 1A2f | |
| | | 300 | | Autogeneration - exported to grid | 1A2f | |
| | | 0 | | Railways - stationary combustion | 1A4a | |
| | 954 | 699 | 255 | Subtotal | | Offset against refineries |
| Coal extraction | 2 | 2 | 0 | Collieries - combustion | 1A1c | |
| Oil and gas extraction | 1684 | | | | | |
| | | 1186 | | Upstream oil production - fuel combustion | 1A1c | |
| | | 499 | | Upstream gas production - fuel combustion | 1A1c | |
| | 1684 | 1684 | 0 | Subtotal | | |
| Petroleum refineries | 200 | 455 | -255 | Refineries - combustion | 1A1b | Offset against autogeneration |
| Blast furnaces | 10 | 0 | 10 | Blast furnaces | 1A2a | |
| Other energy industries | 213 | | | | | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|--|-------|------|------------|--|------|---|
| | | 242 | | Gas production | 1A1c | |
| | | 0 | | Nuclear fuel production | 1A1c | |
| | 213 | 242 | -30 | Subtotal | | From discussion with BEIS, interconnector sites are missing from DUKES, so we add the estimates for these across the time series, to 15_19. |
| Non-Ferrous Metal | 112 | 112 | | Non-Ferrous Metal (combustion) | 1A2b | |
| Chemicals | 939 | 939 | | Chemicals (combustion) | 1A2c | |
| Pulp, Paper and Print | 213 | 213 | | Pulp, Paper and Print (combustion) | 1A2d | |
| Food & drink, tobacco | 697 | 697 | | Food & drink, tobacco (combustion) | 1A2e | |
| | | 118 | | Ammonia production - combustion | 1A2c | Operator's data |
| All other industry (minus iron and steel) | 1852 | 1542 | | Other industrial combustion | 1A2f | |
| Other industrial combustion (colliery methane) | 0 | | | Other industrial combustion (colliery methane) | | |
| | | 47 | | Lime production - non decarbonising | 1A2f | EU ETS |
| | | 3 | | Cement production - combustion | 1A2f | Operator's data |
| All industry except iron and steel | 3814 | 3671 | 143 | Subtotal | | Offset against NEU |
| Non-energy use | 169 | 102 | | Non-energy use (stored) | | |
| | | 210 | | Ammonia production - feedstock use of gas | 2B2 | |
| | 169 | 312 | -143 | | | Offset against industry |
| Iron and steel | 147 | 147 | 0 | Iron and steel - combustion plant | 1A2a | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|------------------------------------|--------------|--------------|------------|--|------|---|
| Domestic | 10135 | 10145 | -10 | Domestic combustion | 1A4b | Isle of Man gas use not offset against DUKES |
| Public administration | 1409 | 1409 | 0 | Public sector combustion | 1A4a | |
| Commercial | 1705 | | | | | |
| Miscellaneous | 344 | | | | | |
| | | 2049 | | Miscellaneous industrial/commercial combustion | 1A4a | |
| | 2049 | 2049 | 0 | Subtotal | | |
| Agriculture | 44 | 44 | 0 | Agriculture - stationary combustion | 1A4c | |
| Autogenerators - colliery methane | 16 | | | | | |
| Coal extraction - colliery methane | 0 | | | | | |
| | | 16 | | Collieries - combustion (colliery methane) | 1A1c | |
| | 60 | 60 | 0 | Subtotal | | |
| Losses | 230 | | 230 | | | The NAEI model doesn't include any accounting for losses compared to DUKES long term trends |
| Total | 29867 | 29709 | 158 | | | |

Notes: Rows are shaded to help illustrate reconciliation between sectors.

1 Mtherm = 105.51 TJ

A 4.2.1.3 Fuel Oil

For shipping, a major research project was completed in 2017 and the results were incorporated into the 2018 submission. The estimated total fuel oil consumption derived from this research is greater than as reported for shipping in DUKES, and any deviations from the national navigation sector are considered additional and are not reconciled elsewhere in the inventory. Additional sectoral deviations are also made to account for known use of fuel oil in power stations, and the Crown Dependencies.

Table A 4.2.4 Fuel reconciliation – Fuel oils use in latest year (Mtonnes)

| DUKES Category | DUKES | GHGI | Difference | GHGI category | CRF | Comment |
|-------------------------------|--------------|--------------|---------------|--|-------------|------------------------------------|
| Major power producers | 0.101 | 0.110 | | Power Stations - UK | 1A1a | EUETS data |
| | | 0.015 | | Power Stations - crown dependencies | 1A1a | Local data sets |
| | 0.101 | 0.124 | -0.024 | Subtotal | | |
| Autogenerators | 0.015 | | 0.015 | | | |
| Industry | 0.240 | 0.011 | | Iron and steel - combustion plant | 1A2a | Reduced to offset increase in 1A1a |
| | | 0.000 | | Non-Ferrous Metal (combustion) | 1A2b | |
| | | 0.026 | | Chemicals (combustion) | 1A2c | |
| | | 0.001 | | Pulp, Paper and Print (combustion) | 1A2d | |
| | | 0.052 | | Food & drink, tobacco (combustion) | 1A2e | |
| | | 0.205 | | Other industrial combustion | 1A2g | Reduced to offset increase in 1A1a |
| | | 0.000 | | Cement production - combustion | 1A2f | Operator's data |
| | 0.240 | 0.296 | -0.055 | Subtotal | | Reduced to offset increase in 1A1a |
| Petroleum refineries | 0.208 | 0.208 | 0.000 | Refineries - combustion | 1A1b | |
| Oil and gas extraction | 0.064 | | | | | |
| Energy sector | 0.628 | 0.628 | 0.000 | Subtotal | | |
| Agriculture | 0.017 | 0.017 | 0.000 | Agriculture - stationary combustion | 1A4c | |
| Domestic | 0.000 | 0.000 | 0.000 | Domestic combustion | 1A4b | |
| Commercial | 0.062 | | | | | |
| Miscellaneous | 0.013 | 0.075 | | Miscellaneous combustion - UK | 1A4a | |

| DUKES Category | DUKES | GHGI | Difference | GHGI category | CRF | Comment |
|------------------------------|--------------|--------------|---------------|---|------|--|
| | | 0.000 | | Miscellaneous combustion - crown dependencies | 1A4a | Local data sets |
| | 0.075 | 0.075 | 0.000 | Commercial and Miscellaneous Total | | |
| Public administration | 0.025 | 0.025 | 0.000 | Public sector combustion | 1A4a | |
| National navigation | 0.070 | 0.167 | | Shipping - coastal | 1A3d | Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES. |
| | | 0.010 | | Fishing vessels | 1A4c | |
| | | 0.000 | | Shipping between UK and CDs | 1A3d | |
| | | 0.006 | | Shipping between UK and Gibraltar | 1A3d | |
| | | 0.000 | | Shipping between UK and Bermuda | 1A3d | |
| | | 0.005 | | Shipping between UK and OTs (excl. Gib and Bermuda) | 1A3d | |
| Marine bunkers | 0.777 | 0.766 | | Shipping - international IPCC definition | | |
| Shipping | 0.777 | 0.954 | -0.177 | Subtotal | | Deviation due to the implementation of the bottom-up shipping inventory, which is known to deviate from DUKES |
| TOTAL | 1.523 | 1.700 | -0.177 | | | |

Notes: Rows are shaded to help illustrate reconciliation between sectors.

A 4.2.1.4 Gas Oil

Gas oil is used in both off-road transport and machinery diesel engines, and as a fuel for stationary combustion. The varied use of this fuel and the complexity of the supply chain complicates the means of allocating consumption across the wide range of sectors that use the fuel in the inventory. DUKES provides a breakdown of gas oil consumption in different industry and other sectors, but the data resolution in DUKES does not distinguish between use of the fuel for stationary combustion and off-road machinery, a distinction which is necessary for the inventory.

The GHGI estimates consumption of gas oil and emissions for off-road machinery using a bottom-up method based on estimates of population and usage of different types of machinery. However, this has led to a situation where the total amount of gas oil consumption across sectors exceeds that which is available as given in DUKES. Therefore, consumption figures, mainly for stationary combustion in industry sectors, have had to be adjusted to obtain a total fuel balance.

The problem is extended when new sources of gas oil consumption are found. For example, the recent development of an inventory for the UK's inland waterways requires the allocation of gas oil to this sector (Walker et al, 2011). During the process of compiling the inland waterways inventory, it became clear that not all vessels with diesel engines use gas oil, but use road diesel and that this may also be the case for other off-road machinery sources, especially those that consume small amounts of fuel on an irregular basis, e.g. for private or recreational use rather than commercial use. There are also inconsistencies in terminology used to define types of fuel; it became apparent that the terms "gas oil", "red diesel" and "diesel" are used interchangeably by fuel suppliers and consumers and this confuses the situation when considering fuel allocations across different sectors.

In light of this, Task 5 of the 2011 UK GHG Inventory Improvement Programme aimed to address the allocation of gas oil and DERV in the GHGI (Murrells et al., 2011). The methodology outlined in Murrells et al. (2011) was used in the compilation of the 2011 inventory, and is summarised here. The same approach has been used in this inventory.

Several fuel suppliers and experts in the petroleum industry and at the Department for Transport were consulted to understand terminologies used, the physical differences between gas oil and DERV, and to gauge opinions on what determines where the fuels are mainly used where it is possible to use either gas oil or DERV. The study concluded that while the majority of agricultural and industrial machinery will be using low tax gas oil (red diesel), a small amount of DERV is likely to be used by private recreational boat users and by equipment with small engines used for private or small-scale commercial use on an irregular basis and the gas oil fuel supply infrastructure makes it more convenient to use DERV.

The study provided new estimates of the amount of DERV and petrol consumed by non-road transport sources with small internal combustion engines. This reduces the overestimation of gas oil consumption and relieves the pressure on how much gas oil consumption by other sources has to be adjusted to match the total amount available as given in DUKES.

The study also considered the allocation of gas oil given in DUKES to different industry and other sectors and how these can be mapped to inventory reporting categories. The detailed bottom-up method is used to estimate gas oil consumption by different off-road machinery and marine vessel types. Independent sources were used to estimate gas oil used by the rail sector while data provided by industrial sites reporting under emission trading schemes (EU ETS) were used to derive an allocation of gas oil consumption by stationary combustion sources in different industry, commercial and other sectors. Also, the UK energy statistics now include an allocation of gas oil for consumption by the oil and gas sector, but since only a partial time series was made available, the study included making estimates of gas oil for this category back to 1990.

A method of re-allocation was developed using an over-arching condition that the total sum of gas oil consumption across all sectors was consistent with the total consumption figures given in DUKES across all years. The method allowed the consumption estimates for industrial off-road machinery and stationary combustion by industry, commercial and public sector activities to

vary in order to align the total consumption estimates with DUKES on the basis that the estimates for these sources are the most uncertain.

As with fuel oil, the introduction of the results of a major research project into the shipping sector in the 2018 submission, whereby Automatic Identification System (AIS) data was used to calculate shipping movements along the coast of the UK and the Crown Dependencies, however suggested that gas oil consumption reported by DUKES is an underestimate. As a result, total gas oil use (not including DERV) deviates from DUKES as any further consumption in the national navigation sector are considered additional to DUKES and are not reconciled elsewhere in the inventory.

Table A 4.2.5 below summarised the DUKES and GHGI allocations for the latest inventory year.

Table A 4.2.5 Fuel reconciliation – Gas oil use in latest year (Mtonnes)

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|-----------------------|-------|-------|------------|-------------------------------------|------|---------|
| Petroleum refineries | 0.000 | 0.000 | 0.000 | Refineries - combustion | 1A1b | |
| Major power producers | 0.039 | 0.039 | | Power stations - UK | | |
| | | 0.002 | | Power stations - crown dependencies | | |
| | | 0.041 | -0.002 | Subtotal | | |
| Autogenerators | 0.056 | | 0.056 | | | |
| Industry | 1.740 | 0.002 | | Iron and steel - combustion plant | 1A2b | |
| | | 0.000 | | Non-Ferrous Metal | 1A2c | |
| | | 0.003 | | Chemicals | 1A2d | |
| | | 0.001 | | Pulp, Paper and Print | 1A2e | |
| | | 0.001 | | Food & drink, tobacco | | |
| | | 0.047 | | Other industrial combustion - UK | 1A2g | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|--------------------------------|--------------|--------------|---------------|--|-------------|--|
| | | 0.000 | | Other industrial combustion - crown dependencies | 1A2g | |
| | | 0.008 | | Cement production - combustion | 1A2f | Operator's data |
| | | 0.177 | | Aircraft - support vehicles | 1A3e | Inventory agency estimates |
| | | 1.526 | | Industrial off-road mobile machinery | 1A2g | Inventory agency estimates |
| | 1.740 | 1.766 | -0.026 | Subtotal | | |
| Commercial | 0.378 | | | | | |
| Miscellaneous | 0.271 | 0.029 | | Miscellaneous combustion - UK | 1A4a | Reduced to offset higher consumption elsewhere |
| | | 0.008 | | Miscellaneous combustion - crown dependencies | 1A4a | Reduced to offset higher consumption elsewhere |
| | 0.271 | 0.037 | 0.235 | Subtotal | | |
| Public administration | 0.299 | 0.019 | 0.280 | Public sector combustion | 1A4a | Reduced to offset higher consumption elsewhere |
| Agriculture | 0.328 | 1.250 | -0.922 | Agriculture - mobile machinery | 1A4c | Inventory Agency estimates which far exceed DUKES |
| Rail | 0.610 | 0.608 | 0.002 | Railways | 1A3c | Reduced to offset higher consumption elsewhere |
| Subtotal (of all above) | 3.721 | 3.721 | 0.000 | | | |
| Oil & gas extraction | 0.599 | | | Upstream Gas Production - fuel combustion | | |
| | | 0.059 | | Upstream Oil Production - fuel combustion | 1A1c | |
| | | 0.539 | | | 1A1c | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|-----------------------|--------------|--------------|---------------|--|---------|--|
| | 0.599 | 0.599 | 0.000 | Subtotal | | |
| National navigation | 0.853 | 1.217 | | Shipping - coastal | | Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES. |
| | | 0.175 | | Fishing vessels | 1A3d | |
| | | 0.006 | | Shipping between Uks and CDs | 1A3d | |
| | | 0.137 | | Shipping - naval | | |
| | | 0.046 | | Motorboats / workboats | 1A5b | |
| | | 0.002 | | Inland goods-carrying vessels | 1A3d | |
| Marine bunkers | 1.653 | 1.653 | | Shipping - international IPCC definition | 1A3d | Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES. |
| Shipping total | 2.506 | 3.236 | -0.730 | Subtotal | | Deviation due to the implementation of the bottom-up shipping inventory, which is known to deviate from DUKES |
| Road | 24.911 | | | House and garden machinery - DERV | 1A4b | |
| | | 0.011 | | Industrial off-road mobile machinery - DERV | 1A2gvii | |
| | | 0.326 | | Sailing boats with auxiliary engines | 1A3d | |
| | | 0.002 | | Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats) | 1A3d | |
| | | 0.112 | | | | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|----------------|--------|--------|------------|--|------|---------|
| | | 24.413 | | Road transport - UK | 1A3b | |
| | | 0.046 | | Road transport - crown dependencies | 1A3b | |
| | 24.911 | 24.911 | 0.000 | Subtotal | | |
| Domestic | 0.144 | | | | 1A4b | 0.000 |
| | | 0.137 | 0.137 | Domestic combustion - UK | 1A4b | |
| | | 0.007 | 0.007 | Domestic combustion - crown dependencies | 1A4b | |
| | 0.144 | 0.144 | 0.000 | Subtotal | | |
| Total | 31.880 | 32.610 | -0.730 | | | |

Notes: Rows are shaded to help illustrate reconciliation between sectors

A 4.2.1.5 Petroleum gases

For petroleum gases (LPG, OPG), the total fuel use in the inventory is greater than the national statistics in several years, to reflect information from other sources (such as EU ETS data) that indicate potential under-reports in the UK energy statistics. These modifications to the energy balance are set out in **Table A 4.2.6**. They mostly relate to refineries, use of feedstock as fuel in the petrochemicals sector, and fuel use for upstream oil and gas production.

Table A 4.2.6 Fuel reconciliation – Use of petroleum gases in the latest year (Mtherms)

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|-----------------------------------|-------|------|------------|-----------------|------|---------|
| Petroleum refineries, other gases | 943 | 1115 | | Refineries, OPG | 1A1b | |
| Autogenerators, other gases | 123 | | | | | |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|--------------------------------------|-------------|-------------|-------------|---|-------------|---|
| Sub-total | 1066 | 1115 | -49 | | | OPG total inferred from EU ETS dataset. This is larger than the reported DUKES value |
| Petroleum refineries, propane | 7 | 7 | 0 | Refineries, LPG | 1A1b | |
| Iron & steel, propane | 0 | 0 | 0 | Iron & steel combustion, LPG | 1A1b | |
| Industry, propane | 133 | | | | | |
| Industry, butane | 37 | | | | | |
| Agriculture, propane | 40 | | | | | |
| Agriculture, butane | 0 | | | | | |
| Commercial, propane | 162 | | | | | |
| Commercial, butane | 1 | | | | | |
| Public administration, propane | 7 | | | | | |
| | | 379 | | Industrial combustion, LPG - UK | 1A2g | |
| | | 0 | | Industrial combustion, LPG - crown dependencies | 1A2g | |
| Sub-total | 380 | 379 | 1 | | | |
| Industry, other gases | 0 | 523 | -523 | Chemicals, OPG | 2B8g | EU ETS higher than DUKES |

| DUKES Category | DUKES | GHGI | Difference | GHGI Category | CRF | Comment |
|-----------------------|-------------|-------------|-------------|---|------|------------------------------|
| Road, propane | 32 | 32 | 0 | Road transport - all vehicles LPG use | 1A3b | |
| Domestic, propane | 94 | 88 | | Domestic combustion, LPG - UK | 1A4b | |
| Domestic, butane | 0 | 5 | | Domestic combustion, LPG - crown dependencies | 1A4b | |
| Sub-total | 94 | 93 | 1 | | | |
| (excluded from DUKES) | | 145 | | Gas separation plant, OPG | 1A1c | EEMS. Outside scope of DUKES |
| (excluded from DUKES) | | 7 | | Gas separation plant, LPG | 1A1c | EEMS. Outside scope of DUKES |
| Total | 1579 | 2301 | -722 | | | |

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together.

1 Mtherm = 105.51 TJ

ANNEX 5: Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information.

A 5.1 ANNUAL INVENTORY SUBMISSION

No additional information.

A 5.2 SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1

No additional information.

ANNEX 6: Comparison of Emission Estimates Using Atmospheric Observations

This Annex discusses the verification of the UK estimates of the Kyoto Gases.

A 6.1 MODELLING APPROACH USED FOR COMPARISON WITH THE UK GHGI

Comparison of the UK GHGI with emission estimates made using atmospheric observations is considered to be best practice by the UNFCCC as it allows for an independent assessment of the GHG emissions from the UK using a comprehensively different approach. Significant differences in the emissions estimated using the two methods are a means of identifying areas worthy of further investigation, for example as occurred with a re-assessment of the emissions of HFC-134a from mobile air conditioning.

In order to provide a comparison with the UK Greenhouse Gas Inventory (GHGI), BEIS (Department of Business, Energy and Industrial Strategy) supported the establishment and maintenance of a high-quality remote observation station at Mace Head (MHD) on the west coast of Ireland. The station reports high-frequency concentrations of the key greenhouse gases and is under the supervision of Prof. Simon O'Doherty of the University of Bristol (O'Doherty *et al.* 2004, 2014). BEIS extended the measurement programme in 2012 with three new tall tower stations across the UK (collectively called the UK DECC (Deriving Emissions linked to Climate Change) network): Tacolneston (TAC) near Norwich; Ridge Hill (RGL) near Hereford; and Tall Tower Angus (TTA) near Dundee, Scotland which relocated to Bilsdale (BSD) in North Yorkshire in Sept 2015. Methane (CH₄), carbon dioxide (CO₂), nitrous oxide (N₂O) and sulphur hexafluoride (SF₆) are measured at all stations across the UK DECC network. The hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) are measured at MHD and TAC, and nitrogen trifluoride (NF₃) is only measured at MHD.

The Met Office, under contract to BEIS, employs the Lagrangian dispersion model NAME (Numerical Atmospheric dispersion Modelling Environment) (Ryall *et al.* 1998, Jones *et al.* 2007) driven by three-dimensional modelled meteorology to interpret the observations. NAME determines the history of the air arriving at Mace Head at the time of each observation. By estimating the underlying background trend (Northern Hemisphere mid-latitude atmospheric concentrations where the short-term impact of regional pollution has been removed from the data) and by modelling where the air has passed over on route to the observation stations on a regional scale, estimates of UK emissions are made. A methodology called Inversion Technique for Emission Modelling (InTEM) has been developed that uses a minimisation technique, Non-Negative Least Squares (NNLS) (Lawson and Hanson, 1974) to determine the emission map that most accurately reproduces the observations (Manning *et al.* 2003, 2011 and Arnold *et al.* 2017).

For each reported gas the Northern Hemisphere background trend and the UK emission estimates are presented. InTEM estimates using only Mace Head (MHD) data are presented

along with the estimates made using the full UK DECC network. In 2014 two additional tower sites, Heathfield (HFD) south of London and Bilsdale (BSD) in North Yorkshire, were established through the NERC GAUGE (Global And UK Greenhouse gas Emissions) programme and have been included in the emission estimates. HFD is currently operated by NPL, both stations are now in the UK DECC Network. For CH₄ only, a further dataset from the tall tower station at Cabauw in The Netherlands has been included courtesy of ECN, The Netherlands. When only MHD data are used a three-year inversion window is assumed (an inversion is performed for a three year period and then the period is incremented by one-year e.g. 1989 – 1991, 1990 – 1992 etc., from which a median for each year is estimated), however with the additional data from the other stations the inversion time window has been shortened to one year or smaller (2 months for CH₄, N₂O) depending on data. The geographical spread of the UK DECC (and other stations) network allows the spatial distribution of the emissions across the UK to be better constrained within InTEM. The InTEM estimates of UK emissions using the atmospheric observations are compared to the reported GHGI estimates.

A 6.2 METHANE

Figure A 6.1 Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.1 shows the Northern Hemisphere background atmospheric concentration of methane from 1990 onwards. As with all of the background plots for the different gases, it shows how the overall atmospheric concentration of the gas in question is changing in the atmosphere as a result of global emissions and atmospheric loss processes. For CH₄, the underlying background trend is positive but there is strong year-to-year variability and a strong seasonal cycle. The growth rate since 2006 has been unusually and consistently positive (average +7.2 ppb/yr).

Figure A 6.2 Verification of the UK emission inventory estimates for methane in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD+CBW, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).




The emission estimates made for the UK using the InTEM methodology are compared to the GHGI emission estimates for the period 1990 onwards. It is important to note that although the UK methane emissions are estimated to have fallen over the last 28 years, the global atmospheric concentration of methane has increased, indicating that global emissions of methane are still outperforming the global natural removal of methane from the atmosphere.

Methane has a natural (biogenic) component and it is estimated that 22% of the annual global emission is released from wetlands (Nilsson *et al.* 2001). Usually natural emissions are strongly dependent on a range of meteorological factors such as temperature and also growth and decay cycles. Such non-uniform emissions will add to the uncertainties in the modelling, although in North West Europe the natural emissions are thought to be small compared to the anthropogenic emissions (<5%, Bergamaschi *et al.* 2005). Due to the relatively strong local (within 20km) influence of biogenic emissions at some of the stations the influence of observations taken when local emissions are thought to be significant (low boundary layer heights, low wind speeds, stable atmospheres) has been reduced within InTEM by removing all observations taken during such meteorological conditions.

The GHGI trend is monotonically downwards whereas the InTEM estimates, after an initial fall, shows little change (**Figure A 6.2**). The InTEM estimates using all of the available observations (MHD+CBW+DECC+GAUGE) are consistent with the MHD+CBW estimates and agree with the reported GHGI. The InTEM 2-month estimates using the full DECC Network (**Figure A 6.3**) do not show any strong seasonal cycle in UK methane emissions.


Figure A 6.3 Verification of the UK emission inventory estimates for methane in Gg yr⁻¹ from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ($\pm 1\sigma$).



A 6.3 NITROUS OXIDE

Figure A 6.4 shows the Northern Hemisphere background atmospheric concentration of nitrous oxide (N₂O) from 1990 onwards. The background trend is monotonic and positive. In 2018, the background increased by 1.1 ppb.

Figure A 6.4 Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



The main activities in Europe resulting in the release of nitrous oxide are; agricultural practices resulting in emissions from soils (~60%), chemical industry (~20%) and combustion (~15%) (UNFCCC 1998 figures). The amount emitted from soils has significant uncertainty and has a diurnal and seasonal release cycle. It is driven by the availability of nitrogen, temperature and the soil moisture content.

Figure A 6.5 Verification of the UK emission inventory estimates for nitrous oxide in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).





Figure A 6.5 shows the InTEM and GHGI emission estimates for nitrous oxide for the period 1990 onwards for the UK. The annual InTEM estimates are similar to the GHGI estimates, with both showing declining UK totals. The InTEM estimates are higher than the GHGI post-1998 although within the 1σ uncertainty. The GHGI estimates show a sharp decline (40 Gg) between 1998 and 1999 in line with the introduction of clean technology at an adipic acid plant in Wilton, north east England. It is estimated to have cut its emissions of N₂O by 90%, from 46 thousand tonne yr⁻¹ to around 6 thousand tonne yr⁻¹. The InTEM estimates show a more gradual decline over this period (expected due to the long 3-year inversion time periods) but the overall reduction is similar. The estimates using all available observations are very similar to the MHD-only estimates. Post-2012 the InTEM estimates are notably higher than the GHGI estimates. The improved network of observations from 2013 onwards allows a strong seasonal cycle in emissions to be highlighted, Figure A 6.6 shows there is a peak in UK emissions in spring-summer and a minimum in the winter months. This is aligned with the traditional fertiliser application period.

The nature of the nitrous oxide emissions challenges the InTEM assumption of uniformity of release both in time and space. Also the point of release to the atmosphere may not be coincident with the activity generating the nitrous oxide e.g. the nitrous oxide may be transported from its source, for example by rivers, prior to its release to the atmosphere.

Figure A 6.6 Verification of the UK emission inventory estimates for nitrous oxide in Gg yr⁻¹ from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ($\pm 1\sigma$).



A 6.4 HYDROFLUOROCARBONS

A 6.4.1 HFC-134a

Figure A 6.7 Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




Figure A 6.7 shows the Northern Hemisphere background atmospheric concentration of HFC-134a from 1994 onwards. The background trend is monotonic and positive, in 2018 the background increased by 6.2 ppt.

Figure A 6.8 Verification of the UK emission inventory estimates for HFC-134a in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).




Figure A 6.8 shows the InTEM and GHGI emission estimates for the UK for HFC-134a for the period 1990 onwards. The GHGI shows a stronger increase in emission compared to the InTEM estimates. The InTEM estimates have risen at about 60% of the rate of the GHGI. Throughout the time series the trend agreement between the GHGI and InTEM is good but the InTEM estimates are consistently about 50-60% of the GHGI estimates, a difference well outside both uncertainty ranges. A similar InTEM result is obtained when the TAC observations are included (from 2013).

A 6.4.2 HFC-152a

Figure A 6.9 Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




Figure A 6.9 shows the Northern Hemisphere background atmospheric concentration of HFC-152a from 1995 onwards. The background trend shows a strong rise from the mid-1990s until 2008, then a much reduced annual increase until 2012. From 2012 onwards a small decline is observed, a result seen globally (Simmonds et al, 2016).

Figure A 6.10 Verification of the UK emission inventory estimates for HFC-152a in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).




Figure A 6.10 shows the InTEM and the GHGI emission estimates for the UK for HFC-152a for the period 1990 onwards. Between 2002-2007 and from 2011 onwards the GHGI estimates are significantly larger than those estimated through inverse modelling. The InTEM estimates are consistently on the lower side compared to the GHGI estimates with larger uncertainties. It is also interesting to note the positive trend from 2011 until 2017 in the UK GHGI, conflicting with a much flatter InTEM trend. The InTEM UK DECC network estimates are consistent with the InTEM MHD-only estimates.

A 6.4.3 HFC-125

Figure A 6.11 Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




Figure A 6.12 Verification of the UK emission inventory estimates for HFC-125 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).




Figure A 6.11 shows the Northern Hemisphere background atmospheric concentration of HFC-125 from 1998 onwards. The background trend is monotonic and exponentially increasing, in 2018 the background increased by 3.2 ppt.

InTEM emission estimates for the UK for HFC-125 for the period 1999 onwards are shown in **Figure A 6.12**. Both the InTEM and UK GHGI estimates suggest that the emissions of HFC-125 from the UK have increased significantly from the 1990s. The InTEM estimates are consistently lower than those from the GHGI (by ~50%) and the uncertainties do not overlap. The introduction of the TAC data in the InTEM estimates shows consistency with the MHD-only estimates.

A 6.4.4 HFC-143a

Figure A 6.13 shows the Northern Hemisphere background atmospheric concentration of HFC-143a from 2004 onwards. The background trend is positive, in 2018 it increased by 1.6 ppt.

Figure A 6.13 Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.14 Verification of the UK emission inventory estimates for HFC-143a in Gg yr^{-1} from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



InTEM emission estimates for the UK for HFC-143a for the period 2005 onwards are shown in **Figure A 6.14** and are compared to the GHGI estimates. UK emissions, as estimated by the GHGI, have increased year on year from the early 1990s until 2010 when a sharp decline is estimated. The InTEM estimates show a slow rise during this period but are consistently lower than the GHGI estimates. The estimates with the DECC network (i.e. including TAC data) are consistent with those from MHD-only and estimate a sharp decline in 2018, a decline that is estimated in GHGI to have started in 2016.

A 6.4.5 HFC-23

Figure A 6.15 shows the Northern hemisphere background atmospheric concentration of HFC-23 from 2008 onwards. The background trend is monotonic and positive, in 2018 the background increased by 1.2 ppt.

InTEM emission estimates for the UK for HFC-23 from 2009 are higher than the low emissions estimated by the GHGI for this period, although the InTEM uncertainties are large and mostly extend down to zero. The introduction of the TAC emissions in 2013 re-enforces this finding, albeit the emissions are very small (**Figure A 6.16**). There are clearly some intermittent emissions of HFC-23 remaining in the UK.

Figure A 6.15 Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.






Figure A 6.16 Verification of the UK emission inventory estimates for HFC-23 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$). The second plot is plotted from the year 2000 on an expanded y-axis.






A 6.4.6 HFC-32


Figure A 6.17 shows the Northern Hemisphere background atmospheric concentration of HFC-32 from 2004 onwards. The background trend is monotonic, positive and the growth rate is increasing, in 2018 the background increased by 3.4 ppt.

Figure A 6.17 Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



InTEM emission estimates for the UK for HFC-32 for 2004 onwards are shown in **Figure A 6.18**. The InTEM emission estimates are lower than the GHGI estimates. Both trends are positive but the rate of increase of the GHGI is significantly larger than the InTEM estimates. By 2017 the GHGI estimated emissions are significantly (more than 100%) larger than those estimated by InTEM. The DECC network InTEM estimates show a slight decline and then rise but are broadly consistent with the MHD-only InTEM estimates.

Figure A 6.18 Verification of the UK emission inventory estimates for HFC-32 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.4.7 HFC-43-10mee

Figure A 6.19 shows the Northern Hemisphere background atmospheric concentration of HFC-43-10mee from 2011 onwards. There is a slight positive trend in the background with an overall growth rate of ~ 0.01 ppt yr⁻¹. The UK emissions of this gas are small. The GHGI estimates are in agreement with those estimated by InTEM. The DECC network InTEM estimates are consistent with the MHD-only InTEM estimates.

Figure A 6.19 Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.20 Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.4.8 HFC-227ea

Figure A 6.21 shows the Northern Hemisphere background atmospheric concentration of HFC-227ea from 2007 onwards. There is a positive trend in the background; in 2018 it increased by 0.13 ppt. The GHGI estimates are significantly (more than 100%) higher than those obtained through inverse modelling. Both the GHGI and InTEM estimate increasing UK emissions.

Figure A 6.21 Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.22 Verification of the UK emission inventory estimates for HFC-227ea in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.4.9 HFC-365mfc

Figure A 6.23 shows the Northern Hemisphere background atmospheric concentration of HFC-365mfc from 2005 onwards. There is positive trend in the background; in 2016 it increased by 0.05 ppt. The GHGI shows a sharp decline in emissions in 2008, the MHD-only InTEM estimates, with the 3-yr inversion time periods, shows the same but understandably slower response. Post-2012 the GHGI estimates rising UK emissions, a trend reproduced by InTEM, however the InTEM estimates then show a decline starting in 2017 that is not seen in the GHGI.

Figure A 6.23 Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.24 Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.4.10 HFC-245fa

Figure A 6.25 shows the Northern Hemisphere background atmospheric concentration of HFC-245fa from 2007 onwards. There is a positive trend in the background; in 2018 it increased by 0.21 ppt. The InTEM estimates have significant uncertainty and are consistently lower than the GHGI estimates. The GHGI estimates show a significant decline in 2008 and then a steady annual increase. The DECC network InTEM estimates show a strong rise in emissions from 2014 to 2018.

Figure A 6.25 Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.26 Verification of the UK emission inventory estimates for HFC-245fa in Gg yr^{-1} from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.5 PERFLUOROCARBONS

A 6.5.1 PFC-14

Figure A 6.27 shows the Northern Hemisphere background atmospheric concentration of PFC-14 from 2004 onwards. The background trend is positive; in 2018 it increased by 0.95 ppt.

The drop in emissions in 2012 in the GHGI reflects the closure of the last significant aluminium production plant in the UK. The InTEM uncertainty ranges are large and generally extend down to zero, probably because the majority of emissions come from intermittently emitting point sources. Overall there is good agreement between the GHGI and the InTEM estimates.

Figure A 6.27 Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.28 Verification of the UK emission inventory estimates for PFC-14 in Gg yr^{-1} from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.5.2 PFC-116

Figure A 6.29 shows the Northern Hemisphere background atmospheric concentration of PFC-116 from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.1 ppt. The UK InTEM estimates are consistent with those reported in the GHGI (Figure A 6.30) given the uncertainties in both estimates.

Figure A 6.29 Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.30 Verification of the UK emission inventory estimates for PFC-116 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.5.3 PFC-218

Figure A 6.31 shows the Northern Hemisphere background atmospheric concentration of PFC-218 from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.02 ppt.

The InTEM estimates are consistent with those reported in the GHGI (**Figure A 6.32**). Interestingly the dip in UK GHGI estimates in 2008 and 2009 is shown by the InTEM estimates, although it is estimated to have occurred later in 2011. The DECC network InTEM estimates show a decline in emissions in 2016 and then a rise.

Figure A 6.31 Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.32 Verification of the UK emission inventory estimates for PFC-218 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.5.4 PFC-318

Figure A 6.33 shows the Northern Hemisphere background atmospheric concentration of PFC-318 from 2010 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.07 ppt. The UK InTEM estimates are significantly higher than the very small emissions reported in the GHGI (**Figure A 6.34**). However the estimated quantities have large uncertainties extending down to zero emissions.

Figure A 6.33 Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.





Figure A 6.34 Verification of the UK emission inventory estimates for PFC-318 in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.6 SULPHUR HEXAFLUORIDE

Figure A 6.35 shows the Northern Hemisphere background atmospheric concentration of sulphur hexafluoride (SF₆) from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.35 ppt.

The UK MHD-only InTEM estimates show a rise from 2005 until 2009 and then a decline until 2015. From 2005 until 2009 the GHGI shows a steady decline from ~0.05 Gg yr⁻¹ to ~0.02 Gg yr⁻¹, a small rise in 2010 and then a slow decline until 2015. There is a significant difference between the GHGI and InTEM estimates between 2008 and 2013. Between 2014 and 2017 the InTEM estimates with the DECC network agree with the MHD-only InTEM estimates and the GHGI.

Figure A 6.35 Monthly Northern Hemisphere trend in SF₆ estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.






Figure A 6.36 Verification of the UK emission inventory estimates for SF₆ in Gg yr⁻¹ from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ($\pm 1\sigma$). InTEM (DECC, 12-mth) estimates are shown in orange ($\pm 1\sigma$).



A 6.7 NITROGEN TRIFLUORIDE

Figure A 6.37 shows the Northern Hemisphere background atmospheric concentration of nitrogen trifluoride (NF₃) from 2013 onwards. The background trend is monotonic and positive; the growth rate in 2018 was estimated to be 0.22 ppt yr⁻¹. NF₃ is only measured at MHD, they started in May 2013. The InTEM emission estimates for the UK are 1800 kg but with an uncertainty that extends down to zero. The GHGI estimate for 2017 is 31 kg.

Figure A 6.37 Monthly Northern Hemisphere trend in NF₃ estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




A 6.8 CARBON DIOXIDE

High precision, high frequency measurements of carbon dioxide (CO₂) are made across the UK DECC network. The Northern Hemisphere background trend is positive; in 2018 it increased by 2.2 ppm. It has a strong seasonal cycle due to the influence of the biosphere.

The CO₂ observed has three principle components:

1. Northern hemisphere background (**Figure A 6.38**).
2. Anthropogenic (man-made)
3. Non-anthropogenic (natural)

Figure A 6.38 Monthly Northern Hemisphere trend in CO₂ estimated from Mace Head observations (blue line). Red line denotes annual trend. Data under grey shading are not yet ratified.



Plants both respire CO₂ and absorb it through photosynthesis. Therefore the CO₂ flux from vegetation has a strong diurnal and seasonal cycle and switches from positive to negative on a daily basis. This unknown natural (biogenic) component of the observed CO₂ is significant when compared to the anthropogenic (man-made) component and cannot be assumed negligible (minimised during the winter months). From direct CO₂ observations it is not possible to distinguish between biogenic and anthropogenic CO₂. Therefore it is difficult to use the CO₂ observations directly in an inversion to estimate anthropogenic emissions. This is because the diurnally varying biogenic CO₂ flux is at odds with a key assumption of the inversion method, namely that emissions do not strongly vary in time over the inversion time-window (monthly). Methods are under development to attempt to over-come these challenges, such as: the use of isotopic observations; through ratios with respect to anthropogenic carbon monoxide (CO); and tracking at what time of day air passes over the ground and using biogenic process models. The uncertainties associated with each of these methods are predicted to be significant.

The estimated uncertainties in the CO₂ GHGI are very small compared to inversion results. Work is on-going to seek to improve our methods of verifying inventory CO₂ emission estimates.

ANNEX 7: Analysis of EU ETS Data

A 7.1 INTRODUCTION

This annex summarises the analysis of the 2017 European Union Emissions Trading System (EU ETS) energy and emissions data that is used within the compilation of the UK GHG inventory. The EU ETS data are used to inform activity data estimates for heavy industry sectors, carbon dioxide emission factors of UK fuels within those sectors, and for comparison of fuel allocations to specific economic sectors against data presented in the *Digest of UK Energy Statistics* (DUKES), published by the Department of Business, Energy and Industrial Strategy (BEIS).

The EU ETS data are used in the UK GHGI compilation as follows:

- EU ETS raw data on energy and emission estimates are processed and checked to enable integration of the activity data, implied emission factors and installation emission estimates as far as practicable within the UK GHG inventory compilation. Emission sources reported in EU ETS are allocated to inventory fuels and source codes, outliers are identified and clarifications of data inconsistencies are sought with the regulatory agencies;
- EU ETS activity data are closely compared against the UK national energy balance (DUKES) published by BEIS, and any inconsistencies are researched, seeking to resolve these through consultation with BEIS wherever possible;
- The verified EU ETS data provides up to date high quality fuel compositional analysis of UK fuels, and these data are used to improve inventory emission estimates across the highly energy intensive sources such as power stations, refineries, cement kilns, and oil and gas sources;
- The EU ETS dataset for offshore oil and gas installations are checked to assess data consistency in emissions reporting between the EU ETS and the (more comprehensive) EEMS dataset that is used within the UK GHGI compilation;
- Overall, the Inventory Agency approach seeks to minimise data discrepancies between EU ETS and the GHGI as far as practicable, in order that the derivation of traded and non-traded emission estimates from the UK GHGI are as accurate as possible. Close consistency between the EU ETS and GHGI is an important aspect of the development of a complete and consistent evidence base for policy development and tracking progress towards UK GHG reduction targets in the non-traded sector under the EU Effort Sharing Decision.

The scope of reporting under EU ETS increased from the 2013 dataset onwards. Phase II of the EU ETS ran from 2008-2012 inclusive. Phase III reporting began in 2013, with some new emission sources and new installations reporting for the first time on their GHG emissions; in particular, the definition of combustion has now been extended to cover installations such as furnaces, driers, and other plant where heat is used directly. A handful of industrial process sources of CO₂ are also included from 2013, such as soda ash production. In the UK, the changes in reporting in Phase III are most significant for the chemicals sector, where the scope of reporting is larger than previously and now encompasses both new industrial process emission sources, and additional energy use. There is also a notable shift towards estimation methods that are based on mass balance calculations (e.g. for chemical manufacturing) within

the UK operator reporting to EU ETS. Other sectors with significant increases in reporting are food and drink manufacture, where installations such as driers, ovens etc. were included for the first time thus adding to the emissions from boilers and CHP plant that were reported in previous years, and roadstone coating, a sector which has not been present in the dataset before.

Analysis of the phase III data enabled the Inventory Agency to improve estimates of emissions from the combustion of waste residues and process off-gases within the chemical and petrochemical sectors (which are all reported under IPPU sector 2B10), as well as to generate improved estimates for the IPPU component of several specific manufacturing processes, such as for soda ash (2B7), and titanium dioxide (2B6). In addition, following a review of methodology for all IPPU sources, EU ETS data for phase II onwards has been used to improve emission estimates for glass production (2A3), brickmaking (2A4) and reductant use in electric arc furnaces (2C1).

The key findings from the analysis and use of the EU ETS data include:

- In the 2017 EU ETS dataset, a very high coverage of Tier 3 emissions data is evident for all fuel use in the power sector, and for solid, liquid and waste-derived fuels used in the cement and lime sectors. The proportion of Tier 3 data is somewhat lower for refinery fuel use, but still sufficiently high for the ETS to be considered the most reliable data available. All of the fuel quality data for these sources and fuels are therefore used within the UK GHGI, as the EU ETS fuel quality data is the most representative dataset available to inform UK carbon dioxide emission factors in the inventory;
- EU ETS emissions data from refineries are higher than estimates derived from DUKES activity data for all but two years within the time series, with a discrepancy evident in OPG emissions. Consultation with the industry trade association, UKPIA, and cross-checking with their data shows that the EU ETS data are felt to more accurately reflect estimates of CO₂, and therefore UK GHGI estimates are based on EU ETS data rather than refinery fuel use data reported in the UK energy balance;
- There are a range of other activity data discrepancies when compared to DUKES within the oil & gas, cement and lime, other industry and iron and steel sectors. Revisions to fuel allocations within the UK GHGI have been implemented for a number of sources, whilst further research is needed in some instances to clarify the issues where the reporting format of EU ETS does not map explicitly to energy balance and GHG inventory reporting requirements;
- EU ETS data for fuel use at chemical and petrochemical production facilities has helped to identify and quantify the combustion of process off-gases that are derived from Natural Gas Liquid (NGL) feedstock to petrochemical production processes, and from combustion of carbon-containing process residues. Analysis of “fuel gas” calorific values and carbon content informs the calculations to estimate emissions from NGL-derived gases and other residues.

The use of EU ETS data in the UK GHG inventory is summarised in **Table A 7.1.1**.

Table A 7.1.1 Summary of the use of EU ETS data in the UK inventory

| Category | Sub-categories | EU ETS data used | | | Comments |
|-----------------------------|---|------------------|----------|-----------|---|
| | | Factors | Activity | Emissions | |
| 1A1a | Power stations - coal, fuel oil natural gas, sour gas | ✓ | | | |
| 1A1a | Power stations – pet coke | | | ✓ | Some additional data is sourced from process operators. |
| 1A1b | Refineries – pet coke & OPG | | | ✓ | EU ETS figures only used where higher than DUKES-based emissions. |
| 1A1b | Refineries – natural gas | ✓ | | | |
| 1A1c 1B2 | Upstream oil and gas production – Gas oil, natural gas, LPG, OPG | | | ✓ | |
| 1A1c | Gas industry – natural gas | | ✓ | | |
| 1A1c 1B1b 1A2a 2C1 | Integrated steelworks | ✓ | ✓ | | Use of various EU ETS data in complex carbon balance – factors for some fuels, activity data for others |
| 1A1c | Collieries – Colliery methane | ✓ | | | |
| 1A2b | Autogenerators - coal | ✓ | | | |
| 1A2f | Lime - coal | | | ✓ | |
| 1A2f | Lime – natural gas | | ✓ | | |
| 1A2g | Industry - pet coke & waste solvents | | | ✓ | No alternative data available for this emission source. |
| 1A2g | Industry – colliery methane | ✓ | | | |
| 2A1 | Cement | | | ✓ | Data used is actually from industry trade-association, but this is based on EU ETS returns |
| 2A2 | Lime | | | ✓ | |
| 2A3 | Glass | | | ✓ | |
| 2A4 | Bricks | | | ✓ | |

| Category | Sub-categories | EU ETS data used | | | Comments |
|----------|------------------------------------|------------------|----------|-----------|----------|
| | | Factors | Activity | Emissions | |
| 2B7 | Soda ash | | | ✓ | |
| 2B8g | Ethylene & other petrochemicals | | | ✓ | |
| 2C1 | Electric arc furnaces - reductants | | | ✓ | |

A 7.2 BACKGROUND

A 7.2.1 EU ETS Data and GHG Inventories

The European Union Emissions Trading System (EU ETS) data provides annual estimates of fuel use and fuel quality data from the most energy intensive sites in the UK, and provides a source of data, or can be used to cross-check data held in the UK Greenhouse Gas Inventory (GHGI), and to inform the carbon contents of current UK fuels. The EU ETS has operated since 2005, and there are now 13 years' worth of data on fuel use and emissions across major UK industrial plant, for 2005-2017.

The data reported under the EU ETS includes quantities of fuels consumed (or other activity data for process sources of CO₂), carbon contents of fuels and other inputs, calorific values (fuels only) and emissions of carbon dioxide, all presented by installation and by emission source. Activity data are also given for many biofuels, although emissions of CO₂ from these fuels are not included in the emissions data. This is useful though, since PI/SPRI/WEI/NIPI emissions data for CO₂ often include biocarbon as well as fossil carbon, and the EU ETS data on biofuels helps to explain differences between CO₂ emissions reported in EU ETS and in those regulator inventories. EU ETS data for individual installations are treated as commercially confidential by the UK regulatory authorities and so only aggregated emissions data are reported in inventory outputs.

As part of the UK's annual reporting requirements to the MMR and UNFCCC, the UK must include a comparison of the EU ETS data against the national inventory dataset within the National Inventory Report. Furthermore, the analysis of the inventory against the EU ETS dataset is coming under increasing scrutiny due to the development of domestic GHG reduction targets that are based on non-traded¹⁷ emissions data only, and the growing need to understand the UK non-traded sector emissions for future reporting under the Effort Sharing Decision.

The EU ETS dataset helps to improve the UK GHG inventory in a number of ways:

- Identifying new sources, therefore improving completeness;

¹⁷ All GHG emissions that are regulated within the EU ETS are defined as "traded" emissions, whilst all other GHG emissions are defined as "non-traded". The EU Effort Sharing Decision will lead to the UK adopting a new target for GHG reductions by 2020 for all of the non-traded emissions (i.e. everything outside of EU ETS), and progress towards this target will be monitored through the UK GHG inventory.

- Helping assess true levels of uncertainty in fuel- and sector-specific data;
- Providing fuel quality data and oxidation factors for complex processes;
- Providing information on process-specific emissions that are not apparent from the national energy balances;
- Reducing uncertainty in the GHGI; and
- Acting as a source of quality assurance to inventory data.

In the 1990-2017 inventory cycle, the Inventory Agency has updated and extended the EU ETS analysis conducted for inventory compilation, using the 2017 EU ETS dataset, which is the fifth year of reporting under the Phase III EU ETS scope. This annex presents a comprehensive review of the thirteen years' of EU ETS data, indicating where the data have already been used in the improvement of the GHGI, as well as highlighting outstanding issues which could be investigated further, with potential for further revision and improvement of the GHGI.

The Inventory Agency has also been provided with details of the 2011-2017 EU ETS data for all offshore oil and gas installations, which are regulated by the BEIS Offshore Inspectorate. Access to these detailed data has enabled a more thorough review of the fuel/gas quality and reported emissions from combustion and flaring sources at offshore installations, and has directly improved the completeness and accuracy of the sector estimates within the UK GHGI.

The analysis of the EU ETS data for use in the UK GHGI necessitates a detailed review of the available data, in order to ensure correct interpretation and application of the available data. The study team prioritises effort to the sources and sites that are the most significant in UK GHGI terms, and/or where data reporting discrepancies have been identified from previous work. For those sectors where EU ETS data are used in the GHGI, it is important to review emission factors from all major installations to ensure that any outliers are identified and checked prior to their inclusion in inventory calculations.

Wherever possible, consistent assumptions are made when interpreting data across all years of the EU ETS. For instance ensuring that each site is allocated to the same inventory sector in each year (unless there is reason to change it – some industrial combustion plants in recent years have been converted into power stations, and so these sites do need to be allocated to different sectors in different years), and that there is consistency in the way in which site-specific names for fuels are interpreted across the entire period. The information on the EU ETS method “Tier” used for each of the data dictates whether they are used in inventory compilation. The highest tier EU ETS data are assumed to be subject to the lowest level of uncertainty, and so only tier 3 and tier 4 emission factor data are used. Occasionally there are internal inconsistencies in the EU ETS data between the data on consumption of a given fuel and emissions from the use of that fuel. These need to be resolved before the data can be used in the UK GHGI. As emissions data are verified, we cross-check the detailed emissions data against the final verified emissions for each site. As a general rule it is found that the most appropriate solution to inconsistencies is to assume that the EU ETS emissions data are correct as EU ETS reporting requirements are well regulated, and that it is the activity data that need to be amended instead.

A 7.2.2 Scope of the UK EU ETS and Implications for the GHG Inventory

There are a number of limitations to the EU ETS data that affect the data usefulness in GHG inventory compilation, including:

- The EU ETS data are only available from 2005 onwards, whilst the UK GHG inventory reports emission trends back to 1990. The additional information that EU ETS provides (e.g. year-specific emission factors for many fuels in energy intensive sectors) helps to reduce the uncertainties in inventory emission estimates for the later years, but care is needed where revisions to the time series are made back to 2005. A consistent approach to inventory compilation across the time series is a key tenet of IPCC good practice guidance, and care is needed to ensure that the use of EU ETS data does not introduce a systematic reporting step-change in the UK GHGI;
- Further to this point, it is important to note that the scope of EU ETS reporting has evolved through the years, from Phase I (2005 to 2007) into Phase II (2008 to 2012 data) and now to Phase III (2013 onwards). The comparability of EU ETS data for many sectors is poor between these three phases. For example, many cement kilns did not report to EU ETS until Phase II: several sectors including cement were reporting under Climate Change Agreements and were opted-out of EU ETS during Phase I. Therefore in several sectors, more complete coverage of EU ETS reporting is evident in Phase II and data from 2008 onwards are therefore much more useful for UK GHGI reporting. The scope of coverage of chemical industry emissions has gone through two step changes – in 2008, and again in 2013, and some sectors (such as roadstone coating) only appear for the first time in the 2013 EU ETS data. Less significantly for the GHGI, many small installations, mostly in the public sector, were removed from EU ETS at the end of phase I. It is vital that the GHGI takes full account of such changes and that UK inventory data do not include trends that merely result from the increase (or decrease) in scope of EU ETS. The changes in EU ETS scope have made the data set increasingly useful, and there are now five years' worth of Phase II data and four years of data under Phase III, hence the EU ETS dataset is now an important source of information for the UK inventory;
- In the UK during EU ETS Phases I and II, the regulators adopted a “medium” definition of the term “combustion”, and as a result there were many sectors where fuel use in specific types of combustion unit were not included in the EU ETS reporting scope until the start of Phase III (2013 onwards). Examples of this include flaring on chemical sites, and fuel use in heaters, dryers, fryers and stenters in industry sectors such as: chemicals, food and drink, textiles, paper and pulp. Hence the total fuel use and GHG emissions from these sectors have typically been under-reported within the EU ETS historically compared with the UK inventory, with many sites and sources excluded from the scope of EU ETS. However, the EU ETS data for these sectors is also incomplete both in Phase II and Phase III because small installations are not covered by EU ETS. Therefore, while the change in scope for combustion installations in phase III is a positive step, it has relatively little impact on the data used in GHGI compilation. Some Phase III data has been used to improve the estimates of emissions from combustion of process wastes / off-gases in the chemical and petrochemical sector in the recent submissions.
- Phase III also brought an increased scope for industrial process sources of CO₂, and data appeared for the first time for soda ash production, and titanium dioxide manufacture.
- When using the EU ETS data, assumptions and interpretations are required to be made regarding the fuel types used by operators; assumptions are made on a case by case basis depending on knowledge of the site or industry and expert judgement. Operators are free to describe fuels as they wish in their returns, rather than choosing from a specific list of fuels, and so assumptions occasionally need to be made where the fuel

type used is not clear from the operator's description. This issue was more significant in the earlier years of EU ETS reporting, with operators often using terms such as "Fuel 1". The assumption then made about fuel type was based both on the other data the operator provided on the fuel such as calorific value, but also by comparison with later data for the same site, since operators now tend to use more recognisable fuel names, and the use of wholly ambiguous terms is now very rare.

Note that:

- The direct use of EU ETS data (e.g. fuel use data by sector) to inform UK GHGI estimates is limited to where the EU ETS is known to cover close to 100% of sector installations. For example, the EU ETS is regarded as representative and 100% comprehensive in coverage of refineries, power stations (except in the case of some small power stations burning biomass, gas oil, or burning oil as the main fuel), integrated steelworks, cement and lime kilns, soda ash plant, titanium dioxide plant, petrochemical works and glassworks (container, flat, wool & continuous filament fibre only – small lead glass and frit producers are not included). Coverage is very close to 100% for brickworks and tileries. For many other industrial sectors (such as chemicals, non-ferrous metals, food and drink, engineering) the EU ETS is not comprehensive and therefore the data are of more limited use, mainly providing a de-minimis fuel consumption figure for these sectors;
- EU ETS Implied Emissions Factors (IEFs) can be used within the UK GHGI, but only where the evidence indicates that EU ETS data are representative of the sector as whole and provides more comprehensive and accurate data than alternative sources. The key criteria to consider in the assessment of EU ETS IEF usefulness is the percentage of annual fuel use by sector where operator estimates use Tier 3 emission factors.
- Review of the EU ETS IEFs for different fuels across different sites provides a useful insight into the level of Tier 3 reporting within different sectors, the progression of higher-Tier reporting within EU ETS through the time series and the level of variability in fuel quality for the different major fuels in the UK. As a general rule, those energy-intensive sectors with near 100% coverage in EU ETS also report a very high proportion of emission factor data at Tier 3. Those sectors with incomplete coverage tend to report most emission factor data below Tier 3. As a result, in all cases where the level of sectoral coverage is high, the quality of reported data is also sufficiently high to be used with confidence in the UK inventory.

A 7.2.3 Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration

Despite detailed research there remain some fundamental limitations in the use of EU ETS data within national inventories where the sector allocation of energy use and emissions cannot be resolved against the national energy statistics that underpin the GHG inventory compilation. One key example is that of the division between fuel use in autogeneration (or heat generation) and direct fuel use within a specific sector. For example, based on the data available from EU ETS, it is impossible to differentiate between gas use in autogeneration on a chemical installation, and gas used directly to heat chemical production processes. In this example, the allocation of EU ETS energy use and emissions between 1A2c (chemicals) and 1A2f (autogenerators) is uncertain, and therefore comparison of EU ETS and GHGI estimates is uncertain. The EU ETS data are not sufficiently detailed and transparent to enable accurate

allocation, and so in all cases fuels and emissions are allocated to the industry sector, and not to autogeneration.

It is worth noting here that the UK energy statistics are also subject to some uncertainty, however small, and that there is likely to be more uncertainty in estimates at industrial sector-level, rather than at more aggregated levels. For example, while fuel producers and suppliers will be able to quantify total fuel demand with a high level of certainty, it would be far more difficult for them to estimate fuel use by specific industrial sectors. This will be reflected in the quality of UK energy statistics which are used to estimate emissions from 1A2c etc. We consider that a high proportion of fossil fuel use by the UK chemical industry will be included in the EU ETS, on the basis that most industrial chemical processes will require sufficiently large combustion installations to exceed the threshold for EU ETS. Therefore, we consider that it is reasonable to assume that EU ETS emissions for the chemical sector should cover most of the sector and therefore be similar in magnitude to those estimated from UK energy statistics and even, given the uncertainty in fuel allocation for autogeneration, to exceed them. For other sectors such as metals, paper, and food and drink, we would assume that the level of sectoral coverage by the EU ETS would be lower, so that emission estimates based on EU ETS would probably be lower than those based on energy statistics, even taking into account the uncertainty regarding autogeneration.

A 7.3 DATA PROCESSING

BEIS provided the detailed EU ETS regulator data from the Environment Agency, Natural Resources Wales, Scottish Environment Protection Agency and Northern Ireland Environment Agency during April & May 2018, and the Inventory Agency industrial emissions experts progressed the analysis, combining the datasets to generate a UK-wide EU ETS dataset. The work built on analysis conducted in previous years, as the EU ETS has been in place since 2005, but this latest analysis, while focussing on the 2017 submission, did involve review of the data for earlier years, to ensure a consistent approach to the interpretation of energy and emissions data across the time series.

The initial step in the analysis is the allocation of all sites in the dataset to one of the economic sectors as reported within the DUKES Commodity Balance tables. Next, the reported fuels for every UK installation have to be allocated to one of the GHGI fuel names, which are also aligned with the fuel types reported within DUKES. This enables a direct comparison of EU ETS fuel totals against sector fuel allocations within DUKES and therefore used within the GHGI.

Most of the allocations have been made as part of previous years' work, and do not need to be revisited. However, several new installations included in the 2017 EU ETS data had to be allocated to DUKES' sectors, and all of the fuel data for 2017 also has had to be allocated to DUKES/GHGI fuel types. In a very small number of cases, we have revised data for earlier years, for example when it has become apparent that existing assumptions are likely to be incorrect. The allocation process does rely upon some expert judgement, with the Ricardo team using the reported EU ETS fuel names as well as the reported fuel quality data such as calorific values and carbon emission factors in order to make the fuel-type allocation for each entry in the EU ETS spreadsheet. The allocation is, occasionally, quite uncertain, particularly with the allocation of petroleum-based fuels such as the GHGI fuel categories LPG, OPG, gas oil and fuel oil, often because of the use of abbreviations or other slightly ambiguous names for fuels within the EU ETS reporting system. Cross-checking of data across the time series for each installation has been used to ensure as much consistency in fuel allocations as possible,

although in some cases, operators of installations use different fuel terminology in different years, and the possibility of the use of different fuels in different years at a site cannot always be ruled out.

The quality checking and allocation process is very resource-intensive and essentially an open-ended task for such a large dataset, and hence the Inventory Agency focuses on the highest emitters and the known “problem” sites and fuel types. Where uncertainties arise in allocations, the most important allocation decisions are copied across to the BEIS DUKES team, for their information and input, as ultimately the EU ETS analysis by the Inventory Agency is taken into account to some degree within the compilation of DUKES for the following year.

As a data verification step, the installation emissions (broken down by fuel) from the EU ETS regulator spreadsheets are then compared against the total installation emissions for 2017 on the European Union Transaction Log (EUTL) which is a central website that holds the verified EU ETS emissions totals for all EU installations in the scheme. Each year we have noted that for some sites the regulator data does not match the EUTL dataset, and therefore some “residual” emissions allocations are generated, from the difference between EUTL and regulator information. In cases where these residual emissions are large, then these are fed back to the regulator contacts, for their consideration and to request any insights into the likely fuels that the residual emissions should be allocated against. Minor residual emissions are ignored for the purposes of the analysis reported here.

A final data set is then available for fuel combustion emission sources, which includes the following data fields:

- GHGI Source Category;
- GHGI Fuel Category;
- Fuel Consumed;
- Fuel Calorific Value;
- Fuel Carbon Emission Factor; and
- Related Emissions of CO₂

The Inventory Agency then combines the data by sector and/or fuel category to provide data for comparison against GHGI emissions data, and energy statistics published in DUKES. In this way, the analysis can:

- provide improved CO₂ emission factors for highly energy-intensive industrial sectors covered by the GHGI through the use of verified data;
- provide a comparison with UK energy statistics, allowing the identification of inconsistencies between EU ETS and DUKES;
- Identify any emission sources that are not contained in the GHGI.

The analysis of the EU ETS data for all onshore facilities was completed by May 2018 and provided to the BEIS team of energy statisticians, in time for them to consider the EU ETS dataset during compilation of the UK energy balance for 2017, as published within DUKES 2018 (which was published in July 2018).

The EU ETS data for offshore oil and gas installations was provided in May 2018 and were used directly in the compilation of emission estimates for the upstream oil and gas sector, after the UK energy balance had been compiled by BEIS. Access to these EU ETS data for offshore facilities provided more fuel-specific information (GCV, carbon content) to help improve completeness and accuracy of the upstream oil and gas estimates in the UK GHGI, augmenting

the EEMS dataset which is a more comprehensive dataset (i.e. EEMS covers more emission sources than EU ETS) but does not provide the same level of fuel-specific data.

A 7.4 EU ETS DATA COVERAGE

The coverage of the EU ETS data has changed over the 13 years for which data are available. Major changes have been outlined in **Section A 7.1**, and these changes in scope have an impact on the usefulness of data for some sectors, with data generally being more complete for Phase II (2008-12) and Phase III (2013-17) of EU ETS. In addition, smaller combustion installations in the industrial, commercial and public sectors are outside the scope of EU ETS, and in fact coverage was decreased after 2007 due to the exemption of certain 'small emitters' from the UK EU ETS. For some source sectors in the GHGI, the EU ETS data therefore only includes a small proportion of the sector and the EU ETS data are not useful to directly inform the GHGI.

The following GHGI source sectors are well represented in the EU ETS data sets in the UK, with all UK installations included:

- Power stations burning coal, gas, and fuel oil as the principal fuel;
- Oil refineries;
- Coke ovens & Integrated steelworks;
- Cement kilns (from Phase II onwards); and
- Lime kilns (from Phase II onwards)
- Glassworks - container, flat, wool & continuous filament glass fibre subsectors only (from phase II onwards)
- Brickworks and other sites manufacturing heavy ceramic goods (from Phase II onwards)
- Titanium dioxide and soda ash manufacture (from Phase III onwards).

However, GHGI sectors such as industrial combustion, autogeneration, and public sector combustion are only partially represented in the EU ETS data. An indication of the actual level of coverage of the EU ETS data can be seen in . The number of sites in each sector which are included in the ETS dataset for 2005 and 2017 are given, together with the Inventory Agency's estimate of the total number of installations in that sector throughout the UK in those years.

Table A 7.4.1 Numbers of installations included in the EU ETS data

| Sector | Number of installations | | | |
|---------------------------------------|-------------------------|----------|--------|----------|
| | 2005 | | 2017 | |
| | EU ETS | UK total | EU ETS | UK total |
| Power stations (fossil fuel, > 75MWe) | 60 | 60 | 51 | 51 |
| Power stations (fossil fuel, < 75MWe) | 23 | 27 | 30 | 35 |
| Power stations (nuclear) | 12 | 12 | 9 | 9 |
| Coke ovens | 4 | 4 | 2 | 2 |
| Sinter plant | 3 | 3 | 2 | 2 |
| Blast furnaces | 3 | 3 | 2 | 2 |

| Sector | Number of installations | | | |
|---|-------------------------|-------------------|--------|-------------------|
| | 2005 | | 2017 | |
| | EU ETS | UK total | EU ETS | UK total |
| Cement kilns | 8 | 15 | 11 | 11 |
| Lime kilns | 4 | 15 | 11 | 11 |
| Refineries | 12 | 12 | 8 | 8 |
| Combustion – iron & steel industry | 11 | 200 ^a | 24 | 200 ^a |
| Combustion – other industry | 171 | 5000 ^a | 469 | 5000 ^a |
| Combustion – commercial sector | 28 | 1000 ^a | 108 | 1000 ^a |
| Combustion – public sector | 169 | 1000 ^a | 124 | 1000 ^a |
| Glassworks (flat, special, container & fibre) | 6 | 32 | 23 | 23 |
| Brickworks | 18 | 80 ^b | 52 | 52 ^b |
| Soda ash & titanium dioxide | 0 | 4 | 1 | 1 |

^a These estimates are 'order of magnitude' figures, based on expert judgement of the inventory team, to show that the number of installations in the UK is likely to be considerably higher than the number of installations reporting in the EU ETS.

^b Numbers of brickworks are not certain in 2005 but will have been significantly higher than in 2008 (when there were about 70) since many brickworks were closed or mothballed in the second half of 2007. All brickworks are believed to be covered by EU ETS in 2017.

Data are included in EU ETS for all coke ovens, refineries, sinter plant and blast furnaces. Power stations are divided into three categories in the table in order to show that, although a few stations are not included in the EU ETS data for 2017, these are all small (in most cases, very small diesel-fired plant supplying electricity to Scottish islands). In comparison, coverage is quite poor in 2005 for cement and lime kilns (due to CCA participants opting out during Phase I) and for combustion processes (due to CCA/UKETS opt-outs and the fact that numerous combustion plant are too small to be required to join the EU ETS). All cement kilns and all lime kilns are included in 2017. Coverage of glassworks and brickworks was very limited during Phase I, but since 2008 has been very good: all large glassworks have been included since 2008, and all but one brickworks were included in Phase II, with that remaining site being added for Phase III. UK totals for brickworks are subject to some uncertainty however, and may be revised in future should more data be obtained. Both soda ash plant and both plants manufacturing titanium dioxide via the chloride process have only been included in EU ETS since the start of Phase III.

For most emission sources the level of detail given in the EU ETS data matches well with the structures of the GHGI, allowing comparison of like with like. Only in the case of coke ovens and integrated steelworks is this not the case, since the EU ETS reporting format does not provide a breakdown of emissions for the sectors reported within the GHGI: i.e. estimates of emissions from coke ovens, blast furnaces and sinter plants are not provided explicitly. However, for these sectors, additional detailed analysis, including the collection of other industry data, has allowed for far greater use of EU ETS data for the inventory.

A 7.5 EU ETS DATA USE IN THE UK GHGI

The use of EU ETS data in the UK GHGI may conveniently be divided into two classes:

- Instances where activity data and, in most cases, emission totals as well are taken from EU ETS;
- Instances where emission factors only are taken from EU ETS and then used in the UK GHG Inventory with activity data from other sources such as DUKES.

A 7.5.1 Activity and Emissions Data

A 7.5.1.1 Crude Oil Refineries

The comparison of EU ETS emissions data against GHGI data based on DUKES fuel use allocations for petcoke, natural gas, fuel oil and OPG use is inconsistent to varying degrees in different years. Previous EU ETS analysis indicated that petcoke data in DUKES were too low; the BEIS energy statistics team have investigated this matter with the refinery operators and have revised data for a number of sites that had been misreporting through the DORS system used to compile DUKES. In recent years, therefore, the EU ETS and DUKES data are closely consistent for petcoke use by refineries.

Data inconsistencies between DUKES and EU ETS remain for other fuels, however. In some cases, this will be due to misallocation of fuel use data within the EU ETS analysis, where fuel names are unclear, e.g. “fuel gas” could be interpreted solely as refinery use of OPG or to also cover the use of natural gas as a back-up fuel within the refinery fuel gas system.

The fuel oil activity data in most years is around 10% higher in EU ETS than in DUKES. Natural gas is a relatively minor fuel in the sector; whilst the EU ETS allocations indicate an over-report in DUKES, there is considerable uncertainty over the allocations of gases in the EU ETS dataset, as noted above. However, DUKES data for natural gas used in autogeneration includes some fuel burnt at refineries, thus the difference between refinery fuel use as given in EU ETS, and that derived from DUKES data can be reduced by taking this into account. Consumption of naphtha reported in DUKES as “unclassified industry” is allocated to refineries as the only known consumers in the UK. However, in the case of OPG, there is typically an under-report in DUKES, although the data in DUKES is higher in two years. **Table A 7.5.1** below presents the emissions allocated to OPG for those years (2004 onwards except 2005, 2012) where UKPIA and EU ETS data indicates that DUKES data are too low. Note that the GHGI estimates also include the assumption that all of the OPG allocation to “autogenerators” within the DUKES commodity balance tables (in the column “Other gases”) is used within the refinery sector. Consultation with the BEIS DUKES team has indicated (Personal Communication, Evans, 2010) that the “Other gases” column in the Commodity Balance tables is the OPG on the refinery basis, with CHP plant on site allocated to the autogeneration line. We have therefore retained this assumption in the current analysis, including the autogenerator allocation of “other gases” within the refinery sector.

To resolve the refinery sector under-report, we have compared DUKES data against EU ETS data, and also considered the total carbon dioxide emissions for the refinery sector provided annually by UKPIA. At the installation level, the UKPIA and EU ETS data show very close consistency for recent years (typically within 1%). The close consistency of the EU ETS and UKPIA data further strengthens the case for using EU ETS data as the primary dataset to inform the UK GHG inventory, in preference to the DUKES energy statistics.

At the fuel-specific level, the greatest disparity is evident in the reporting of OPG use at refineries; the reporting disparity has therefore been resolved through a top-down emissions comparison between DUKES-derived data and the best available operator data from EU ETS (2005 onwards) and UKPIA (pre-2005), with the difference between the two then allocated to OPG use in the UK GHGI. UK inventory estimates of emissions for the sector are therefore aligned with EU ETS totals back to 2005, and with UKPIA data prior to 2005, unless the estimates derived from DUKES data are higher than those from UKPIA or EU ETS (i.e. in 2005 and 2012).

No deviations from UK energy statistics have been made prior to 2004, as the data from UKPIA and GHGI estimates based on DUKES are closely consistent with the DUKES-derived data being slightly higher; therefore a conservative approach is adopted, using DUKES-derived GHG estimates.

The time series of emissions data and the additional OPG and petroleum coke emissions data (where DUKES data are low) for the sector are shown below.

Table A 7.5.1 Refinery Emissions Data Comparison and Revision to OPG Activity

| Year | Best Operator Data ¹ | Refinery emissions total (if based on DUKES) | Additional emissions assumed from OPG | Additional emissions assumed from Pet Coke |
|------|---------------------------------|--|---------------------------------------|--|
| | kt C | kt C | kt C | kt C |
| 2000 | 4,599 | 4,717 | 0 | 0 |
| 2001 | 4,535 | 4,664 | 0 | 0 |
| 2002 | 4,767 | 5,243 | 0 | 0 |
| 2003 | 4,772 | 5,085 | 0 | 0 |
| 2004 | 4,999 | 4,925 | 74 | 0 |
| 2005 | 5,007 | 5,422 | 0 | 150 |
| 2006 | 4,910 | 4,751 | 83 | 76 |
| 2007 | 4,857 | 4,781 | 26 | 50 |
| 2008 | 4,709 | 4,470 | 118 | 121 |
| 2009 | 4,492 | 4,126 | 239 | 126 |
| 2010 | 4,632 | 4,426 | 130 | 76 |
| 2011 | 4,739 | 4,491 | 248 | 0 |
| 2012 | 4,287 | 4,300 | 0 | 0 |
| 2013 | 4,002 | 3,856 | 145 | 2 |
| 2014 | 3,678 | 3,559 | 118 | 0 |
| 2015 | 3,682 | 3,657 | 0 | 47 |
| 2016 | 3,708 | 3,586 | 65 | 56 |
| 2017 | 3,698 | 3,629 | 56 | 13 |

¹ For 2005 onwards, the EU ETS data are verified by third parties and regarded as the best available sector estimates; prior to 2005 the best available operator emissions data are from the trade association, UKPIA.

² For 2005, DUKES activity data for petroleum coke are somewhat lower than the corresponding figure in the EU ETS, so even though CO₂ emission estimates based on DUKES figures for all fuels exceed the CO₂ figure given in the EU ETS, we use the higher (EU ETS) figure for petroleum coke, with the result that for 2005, the UK inventory figure for refinery CO₂ is higher (at 5422 kt C) higher than either the operator or DUKES based totals.

There is some level of uncertainty in the allocation of fuels in EU ETS to specific “DUKES” fuels, although the OPG use in refineries seems to be reported quite consistently as “Refinery Gas”, “Refinery Off-Gas”, or “OPG/RFG”. The BEIS DUKES team have reviewed the year to year consistency of OPG use in refineries through the DORS system.

A 7.5.1.2 Oil & Gas Terminal OPG and LPG Use

The allocation of fuel use reported within EU ETS to UK energy balance fuel nomenclature is uncertain in some cases. Analysis of the EU ETS fuel use data does indicate that there are small amounts of LPG and OPG being used in the upstream oil & gas sector that are not evident within DUKES.

The DUKES team have noted previously (Personal communication, DECC, 2010) that some LPG and OPG fuels are abstracted from upstream oil and gas exploration and production sources, rather than purchased from other sources, and that no data have been collected for this source since DUKES last published data, for the year 2002.

Therefore, the data from the EU ETS from oil and gas processing terminals on LPG and OPG combustion are used directly within the UK GHG inventory for the Phase II and Phase III years of 2008 to 2017, with estimates for 2003 to 2007 derived by interpolation between the EU ETS 2008 data and the DUKES 2002 data.

A 7.5.1.3 Natural Gas Use by Downstream Gas Supply Installations

The EU ETS data includes natural gas use by large gas compressor and storage sites that operate on the UK gas transmission and distribution network, as well as the three operational LNG terminals and a small number of other downstream gas industry sites.

The gas use reported in EU ETS for these sites throughout Phase II and III has been notably higher than the allocation of gas within DUKES Commodity Balance table 4.2 (Energy Industry Use, Other). This has been evident in the traded / non-traded analysis for the gas supply sector in the UK and DA GHGI.

As this gas use arises from the downstream network, the Inventory Agency and the BEIS DUKES team consider that the DUKES data indicate a small misallocation of gas use, rather than a gap in reported gas use. For 2005 to 2017, therefore, the EU ETS data for this source are used within the UK GHG inventory, and the overall gas use data are balanced by reducing the allocation of gas use to “other industrial combustion” (IPCC source 1A2g); the EU ETS data since 2005 shows good consistency with the data from DUKES for earlier years.

The increased gas use for this sector based on EU ETS data is expected to still be a small under-report for the sector as a whole, as the EU ETS scope only includes around 35 of the larger gas compressors, LNG terminals and storage sites on the UK network, and it is likely that additional gas use on smaller sites also occurs. However, the Inventory Agency has no data to inform such estimates.

A 7.5.1.4 Other Industry OPG use

There are a number of “other industry” sites where OPG use has been allocated by the Inventory Agency from EU ETS data, where the fuel is defined as either a specific gas (e.g. ethane, propane, butane) or more generic terms such as “OPG”, “High Pressure Refinery Gas”, “Low Pressure Refinery gas”, “fuel gas” or “RFG/OPG/ROG” within the EU ETS forms.

In refinery complexes, the use of RFG for autogeneration (for the refinery and/or for co-located plant) is reported within the energy balance (allocated to “OPG”). At a number of other UK installations, commodities that are used initially as feedstocks in chemical and petrochemical production (e.g. naphtha, ethane, LPG, gas oil) are allocated to “non-energy use” in the UK energy statistics; any subsequent use of process off-gases (derived from these NEU feedstocks) as a fuel is not reflected in DUKES. Therefore the inventory agency uses other data from industry, primarily from EU ETS, to generate estimates of the use of such secondary fuels. For a small number of sites, consultation with the DUKES team, regulators and operators has clarified that the EU ETS energy and emissions data are the best available dataset for use in the UK GHGI.

In the 1990-2017 inventory cycle, EU ETS data for fuel use at petrochemical production facilities helped to identify and quantify the combustion of process off-gases that are derived from Natural Gas Liquid (NGL) feedstock to petrochemical production processes. Analysis of “fuel gas” calorific values and carbon content informs the calculations to estimate emissions from NGL-derived gases and other residues.

A 7.5.1.5 Industrial Processes

The EU ETS dataset contains data on a number of industrial processes for which alternative data sources are either unavailable or of low quality. The EU ETS data therefore constitute the most reliable set of emissions data for these processes and are used in the UK inventory. In almost all cases, the EU ETS activity data are difficult to use directly, largely because different operators provide activity data and emission factors on a different basis (e.g. some may provide input material and emission factors on a consumption basis, others will provide production data and emission factors on a production basis). Therefore, for all of the industrial process sources, the EU ETS emissions data are adopted, and activity data are generally back-calculated from the emissions using a suitable IPCC emission factor. The industrial process sources where EU ETS data have been used to generate estimates of emissions included within the UK GHGI in this submission, include:

- Emissions from the manufacture of lime. UK activity data for limestone and dolomite consumption in lime production would yield much lower emission estimates than is suggested by EU ETS returns therefore, as a conservative approach, the EU ETS data are used instead. Activity data are back-calculated using the IPCC default factor for lime production. See **Section 4.3** for further details.
- Emissions from the use of carbonates in the manufacture of glass. As with lime production, the available data on consumption of limestone and dolomite for glass production are suspect, being very inconsistent across the time series, and so EU ETS data are used in the generation of the inventory time series, as detailed in **Section 4.4**.
- Emissions from the use of clays, carbonate minerals and other additives in the manufacture of bricks and roofing tiles, as detailed in **Section 4.5**. The EU ETS data are very detailed, with separate lines for different input materials such as different types of clay, carbonate minerals used in the bricks or in scrubbers used to abate fluoride

emissions, and coke oven coke/petroleum coke used as an additive in certain bricks. UK brick production data are used as activity data.

- Estimates for emissions from the use of limestone in flue-gas desulphurisation (FGD) plant for the years 2005-2017 are taken from EU ETS data, because UK activity data (for gypsum produced from the FGD plant) are incomplete for those years. Activity data for 1990-2004 are available from non-EU ETS sources, and are back-calculated from the EU ETS CO₂ emissions for 2005 onwards assuming an emission factor of 253 kg CO₂ per tonne gypsum produced (which is based on an assumed 100% conversion of limestone and SO₂ into gypsum and CO₂).
- EU ETS Phase III saw the introduction of data for soda ash manufacturing sites and EU ETS data, and CO₂ emissions reported for earlier years in the PI, are used as the basis of UK inventory emissions data for that sector. See **Section 4.12** for more details.
- Titanium dioxide production was also included in Phase III of the EU ETS, but full data for the UK plant are not included in the data set provided, and so emission estimates are generated using an alternative, more conservative method.
- Petroleum coke is added to some electric arc furnaces as a reductant, and emissions and activity data for the period 2005-2017 are taken from EU ETS data, with emissions in earlier years being extrapolated on the basis of plant production. See **Section 4.16** for further details.

A 7.5.2 Implied Emission Factors

A 7.5.2.1 Power Stations

Table A 7.5.2 summarises EU ETS data for fuels burnt by major power stations and coal burnt by autogenerators. The percentage of emissions based on Tier 3 emission factors is given (Tier 3 factors are based on fuel analysis, and are therefore more reliable than emission factors based on default values), as well as the average emission factor for EU ETS emissions based on Tier 3 factors.

Table A 7.5.2 EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases)

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 only) |
|------|------|----------|--|
| 2005 | Coal | 99 | 615.3 |
| 2006 | | 100 | 615.0 |
| 2007 | | 100 | 614.7 |
| 2008 | | 100 | 612.4 |
| 2009 | | 100 | 607.2 |
| 2010 | | 100 | 609.0 |
| 2011 | | 100 | 608.9 |
| 2012 | | 100 | 611.7 |
| 2013 | | 100 | 612.5 |
| 2014 | | 100 | 611.8 |
| 2015 | | 100 | 607.9 |
| 2016 | | 94 | 488.5 |

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 only) |
|---------|-----------------------------------|----------------|--|
| 2017 | | 100 | 613.0 |
| 2005 | Fuel oil / Waste oil ^a | 59 | 860.3 |
| 2006 | | 66 | 873.0 |
| 2007 | | 68 | 871.1 |
| 2008 | | 91 | 869.5 |
| 2009 | | 94 | 872.7 |
| 2010 | | 95 | 873.2 |
| 2011 | | 94 | 873.9 |
| 2012 | | 95 | 873.1 |
| 2013 | | 94 | 869.2 |
| 2014 | | 93 | 868.5 |
| 2015 | | 90 | 870.3 |
| 2016 | | 86 | 872.4 |
| 2017 | | 87 | 871.5 |
| 2005 | Natural gas | 52 | 1.443 |
| 2006 | | 76 | 1.465 |
| 2007 | | 95 | 1.464 |
| 2008 | | 97 | 1.467 |
| 2009 | | 100 | 1.464 |
| 2010 | | 99 | 1.460 |
| 2011 | | 99 | 1.458 |
| 2012 | | 100 | 1.461 |
| 2013 | | 99 | 1.464 |
| 2014 | | 100 | 1.461 |
| 2015 | | 100 | 1.462 |
| 2016 | | 99 | 1.462 |
| 2017 | | 99 | 1.466 |
| 2005 | 7Coal - autogenerators | 100 | 594.3 |
| 2006 | | 100 | 596.3 |
| 2007 | | 100 | 594.5 |
| 2008 | | 100 | 581.3 |
| 2009 | | 100 | 600.6 |
| 2010 | | 100 | 599.9 |
| 2011 | | 100 | 594.9 |
| 2012 | | 100 | 598.3 |
| 2013-17 | | 0 ^b | N/A |

^a It is not possible to distinguish between fuel oil and waste oil in the EU ETS data, so all emissions have been reported under fuel oil.

^b Plant operated as a power station after 2012 and included in the figures for power stations burning coal

The EU ETS data shown are regarded as good quality data, since a high proportion of emissions are based on Tier 3 emission factors (i.e. verified emissions based on fuel analysis to ISO17025). The factors are also very consistent across the time-series, which would be expected for this sector. As shown in Section 3, the EU ETS data for power stations also cover almost all UK installations in this sector, and certainly cover all of the installations which burn coal, fuel oil and natural gas.

A few power stations burn small quantities of petroleum coke as well as coal. One supplies data to ETS for coal/petroleum coke blends i.e. there are no separate emissions data or carbon factors for the coal and the petroleum coke at that site. We therefore back-calculate the coal IEF in those blends by using an assumed default for the petcoke carbon content and more detailed activity data on the constituents of the fuel blends, obtained directly from the operator.

The EU ETS based emission factors presented above for power stations are used directly as the emission factors in the GHGI, with the exception of the 2005 figure for gas, where Tier 3 factors were only used for about half of the sector's emissions reported in EU ETS. Small quantities of sour gas were burnt at one power station in 2005-2007 and 2009 and EU ETS Tier 3 emission factors are available and therefore used. [Due to the confidentiality of the data, the emission factors are not shown]. Prior to 2005, the emission factors for these sectors are based on the methodology established by Baggott *et al*, 2004, since it has been concluded that this represents the most reliable approach.

The EU ETS factors for coal-fired autogenerators are slightly different to the factors for the power stations in that, although the EU ETS data are exclusively Tier 3, they only represent about 80-90% of total fuel used by the sector.

A 7.5.2.2 Crude Oil Refineries

Table A 7.5.3 below summarises the EU ETS data for the major fuels burnt by refineries in the UK.

The main fuels in refineries are fuel oil and OPG and emissions also occur due to the burning off of 'petroleum coke' deposits on catalysts used in processes such as catalytic cracking. In the latter case, emissions in the EU ETS are not generally based on activity data and emission factors but are instead based on direct measurement of carbon emitted. This is due to the technical difficulty in measuring the quantity of petroleum coke burnt and the carbon content. Refineries also use natural gas, although it is a relatively small source of emissions compared to other fuels.

Table A 7.5.3 Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas)

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|----------|----------|---|
| 2005 | Fuel Oil | 25 | 860.9 |
| 2006 | | 65 | 873.8 |
| 2007 | | 78 | 877.2 |
| 2008 | | 91 | 871.6 |
| 2009 | | 91 | 876.2 |

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|-------------|----------|---|
| 2010 | | 97 | 878.2 |
| 2011 | | 85 | 877.6 |
| 2012 | | 82 | 887.1 |
| 2013 | | 95 | 874.3 |
| 2014 | | 96 | 875.8 |
| 2015 | | 61 | 876.7 |
| 2016 | | 66 | 876.1 |
| 2017 | | 90 | 864.8 |
| 2005 | OPG | 56 | 1.494 |
| 2006 | | 54 | 1.468 |
| 2007 | | 65 | 1.587 |
| 2008 | | 78 | 1.482 |
| 2009 | | 78 | 1.494 |
| 2010 | | 79 | 1.509 |
| 2011 | | 68 | 1.445 |
| 2012 | | 61 | 1.463 |
| 2013 | | 77 | 1.493 |
| 2014 | | 64 | 1.508 |
| 2015 | | 62 | 1.492 |
| 2016 | | 61 | 1.470 |
| 2017 | | 66 | 1.481 |
| 2005 | Natural Gas | 0 | N/A |
| 2006 | | 43 | 1.460 |
| 2007 | | 45 | 1.462 |
| 2008 | | 98 | 1.475 |
| 2009 | | 98 | 1.480 |
| 2010 | | 97 | 1.465 |
| 2011 | | 81 | 1.447 |
| 2012 | | 63 | 1.442 |
| 2013 | | 89 | 1.459 |
| 2014 | | 87 | 1.459 |
| 2015 | | 87 | 1.465 |
| 2016 | | 81 | 1.456 |
| 2017 | | 84 | 1.462 |

There has been some variation in the proportion of Tier 3 reporting for all three fuels, which will adversely affect the quality of the emission factors, although coverage is still in excess of 50% for all fuels.

Emission factors for **fuel oil** generated from EU ETS data have been adopted in the GHGI, with the exception of data for 2005, where Tier 3 methods were used for only 25% of fuel.

Carbon factors can be derived for **OPG** based on moderate levels of Tier 3 reporting for 2005-2007 and 2011-2017 but 80% for 2008-2010, which gives us a high confidence in the representativeness of the carbon factors for 2008-10. There is some uncertainty regarding the allocation of EU ETS fuels to the OPG fuel category, and the derived emission factors do cover a wider spread of values than for many other fuels in EU ETS. However, this reflects the nature of this fuel, and the data for all years have been used in the inventory.

Carbon factors for natural gas are based on a low % of Tier 3 reporting until 2008; in 2008 to 2010 over 90% of gas use is reported at Tier 3 and over 80% in 2011 and 2013-2017. Within the UK GHGI, the EU ETS factors for 2008 to 2016 are used directly, whilst emission factors for earlier years are derived from gas network operator gas compositional analysis.

EU ETS emission data for **petroleum coke** are higher in 2005-2010, when compared against the estimates derived from DUKES activity data and the industry-recommended emission factor. This is especially noticeable for 2005, where the petroleum coke consumption given in DUKES would have to be more than 100% carbon in order to generate the carbon emissions given in the EU ETS. Consultation with BEIS energy statisticians has identified that the figures given in DUKES are subject to uncertainty and hence the EU ETS data are used directly within the UK GHGI for those years.

A 7.5.2.3 Integrated Steelworks & Coke Ovens

Table A 7.5.4 summarises EU ETS data for the major fuels burnt at integrated steelworks and coke ovens. The data exclude one independent coke oven which calculated emissions using a detailed mass balance approach which makes it more difficult to assess the data in the same way as the other installations. This site closed at the end of 2014.

**Table A 7.5.4 EU ETS data for fuels used at integrated steelworks & coke ovens
(Emission Factors in kt/Mt for solid & liquid fuels, kt/Mth for gases)**

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|-------------------------|----------|---|
| 2005 | Blast furnace gas | 0 | N/A |
| 2006 | | 100 | 6.873 |
| 2007 | | 90 | 6.920 |
| 2008 | | 92 | 6.945 |
| 2009 | | 92 | 7.029 |
| 2010 | | 100 | 6.949 |
| 2011 | | 94 | 6.990 |
| 2012 | | 96 | 6.815 |
| 2013 | | 91 | 6.766 |
| 2014 | | 91 | 6.776 |
| 2015 | | 100 | 7.653 |
| 2016 | | 100 | 7.578 |
| 2017 | | 90 | 7.219 |
| 2005 | | 0 | N/A |

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|---------------|----------|---|
| 2006 | Coke oven gas | 0 | N/A |
| 2007 | | 0 | N/A |
| 2008 | | 53 | 1.093 |
| 2009 | | 96 | 1.140 |
| 2010 | | 96 | 1.117 |
| 2011 | | 96 | 1.089 |
| 2012 | | 96 | 1.094 |
| 2013 | | 96 | 1.103 |
| 2014 | | 100 | 1.143 |
| 2015 | | 100 | 1.216 |
| 2016 | | 48 | 1.659 |
| 2017 | | 100 | 1.068 |
| 2005 | Natural gas | 0 | N/A |
| 2006 | | 3 | 1.479 |
| 2007 | | 2 | 1.478 |
| 2008 | | 0 | N/A |
| 2009 | | 58 | 1.425 |
| 2010 | | 68 | 1.441 |
| 2011 | | 64 | 1.441 |
| 2012 | | 64 | 1.443 |
| 2013 | | 27 | 1.447 |
| 2014 | | 23 | 1.445 |
| 2015 | | 0 | N/A |
| 2016 | | 12 | 1.445 |
| 2017 | | 33 | 1.446 |
| 2005 | Fuel oil | 0 | N/A |
| 2006 | | 0 | N/A |
| 2007 | | 0 | N/A |
| 2008 | | 84 | 878.3 |
| 2009 | | 89 | 884.7 |
| 2010 | | 83 | 887.6 |
| 2011 | | 88 | 888.7 |
| 2012 | | 67 | 877.3 |
| 2013 | | 33 | 846.2 |
| 2014 | | 30 | 844.7 |
| 2015 | | 32 | 845.1 |
| 2016 | | 0 | N/A |
| 2017 | | 0 | N/A |

Most of the ETS data for coke ovens and steelworks are now used in the GHGI, although not the emission factors shown above. Instead, the Inventory Agency have used the EU ETS data and other detailed, site-specific and fuel-specific data, provided by the process operators to refine the carbon balance model used to generate emission estimates for the sector. Details of the revisions to the carbon balance model can be found in the research report from the 2013-2014 inventory improvement programme (Ricardo-AEA, 2014)

A 7.5.2.4 Cement Kilns

Table A 7.5.5 summarises EU ETS data for the major fuels burnt at cement kilns.

Table A 7.5.5 EU ETS data for Fuels used at Cement Kilns (kt / Mt)

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|----------------|----------|---|
| 2005 | Coal | 100 | 678.6 |
| 2006 | | 100 | 546.2 |
| 2007 | | 100 | 664.3 |
| 2008 | | 100 | 655.8 |
| 2009 | | 97 | 658.8 |
| 2010 | | 97 | 637.8 |
| 2011 | | 100 | 645.8 |
| 2012 | | 100 | 662.4 |
| 2013 | | 100 | 694.2 |
| 2014 | | 100 | 673.9 |
| 2015 | | 100 | 675.3 |
| 2016 | | 100 | 683.6 |
| 2017 | | 100 | 683.3 |
| 2005 | Petroleum coke | 0 | 810.7 |
| 2006 | | 100 | 820.8 |
| 2007 | | 100 | 830.2 |
| 2008 | | 100 | 819.1 |
| 2009 | | 73 | 812.6 |
| 2010 | | 69 | 791.9 |
| 2011 | | 100 | 738.4 |
| 2012 | | 100 | 770.2 |
| 2013 | | 100 | 811.1 |
| 2014 | | 100 | 793.4 |
| 2015 | | 100 | 824.6 |
| 2016 | | 100 | 822.2 |
| 2017 | | 100 | 823.1 |

The EU ETS dataset also provides a detailed breakdown of cement sector process emissions from the decarbonisation of raw materials during the clinker manufacturing process. These data are useful to compare against statistics provided by the Mineral Products Association (MPA)

regarding clinker production and the non-combustion emissions associated with UK cement production. The MPA data on clinker production are commercially confidential.

The two data sets show significant differences for 2005-2007; however the EU ETS data cover only a fraction of the sector, so differences might be expected. From 2008 onwards, there is close agreement (average of 0.5% difference) between the two data sets. The coal IEF data across the time series are also fairly consistent, other than in 2006 where the ETS value is very much lower than in other years. Because of the good agreement in both activity data and emission factors for 2008 onwards, the industry-wide estimates provided by the MPA and used within the GHGI show very close comparison with the EU ETS estimates. The difference between the EU ETS and those reported to the GHGI are consistently less than 1%, as outlined below in **Table A 7.5.6**.

Table A 7.5.6 Comparison of Cement Sector Carbon Dioxide Emissions* within the UK GHGI and the EU ETS

| | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| GHGI CO ₂ emissions (kt) | 8,296 | 5,687 | 5,789 | 6,148 | 5,602 | 6,009 | 6,212 | 6,622 | 6,819 | 6,569 |
| Sum of EU ETS CO ₂ emissions (kt) | 8,259 | 5,645 | 5,722 | 6,087 | 5,556 | 5,972 | 6,205 | 6,534 | 6,800 | 6,560 |
| EU ETS / GHGI | 99.5% | 99.3% | 98.8% | 99.0% | 99.2% | 99.4% | 99.9% | 98.7% | 99.7% | 99.9% |

*The data in this table include fuel combustion emissions (reported under IPCC 1A2f) and process emissions (reported under IPCC sector 2A1) from UK cement kilns.

A 7.5.2.5 Lime Kilns

Table A 7.5.7 summarises data given in the EU ETS datasets for the major fuels burnt at lime kilns. Unlike cement kilns, which often burn a variety of fuels, many lime kilns burn just a single fuel, often natural gas. The data below exclude coke oven coke used in lime kilns at soda ash plant since these kilns were not covered by EU ETS until Phase III, and the small number of sites make the data confidential in any case.

Table A 7.5.7 EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases)

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) |
|------|-------|----------|---|
| 2005 | Coal* | - | N/A |
| 2006 | | - | N/A |
| 2007 | | 34 | 846.9 |
| 2008 | | 79 | 701.4 |
| 2009 | | 100 | 698.9 |
| 2010 | | 100 | 634.4 |
| 2011 | | 100 | 703.9 |
| 2012 | | 100 | 725.6 |
| 2013 | | 100 | 689.1 |
| 2014 | | 100 | 680.2 |
| 2015 | | 100 | 693.1 |
| 2016 | | 100 | 688.8 |
| 2017 | | 100 | 677.1 |

*Coal used in the lime industry in the UK includes a proportion of anthracitic coal, and hence some of these IEFs are notably higher than for coal used in other sectors of UK industry.

The EU ETS data for lime kilns vary across the time series, both in terms of the proportion of emissions based on Tier 3 factors, and in the emission factors themselves. EU ETS based factors are currently used for coal and petroleum coke from 2008 onwards, as the EU ETS data do include all lime kilns burning those fuels and almost all of those data are Tier 3 and hence are regarded as highly reliable.

EU ETS data for natural gas use in the lime industry does cover all installations burning this fuel, however the proportion of emissions based on Tier 3 factors is very low. Therefore the EU ETS emission factors are not used in the UK GHGI, and the emission factors for natural gas continue to be based on the methodology given in Baggott *et al*, 2004.

Table A 7.5.8 shows implied emission factors for process-related emissions from lime kilns that are used within the UK GHG inventory. The lime industry can be sub-divided into those installations where lime is the primary product, and carbon dioxide is an unwanted by-product; and those installations where both lime and carbon dioxide are utilised. The latter include kilns in the sugar industry (where carbon dioxide is used in the purification stages) and soda ash production (where carbon dioxide is combined with other chemicals to produce sodium carbonate), and in these kilns, the carbon dioxide from decarbonisation of the limestone or dolomite feedstock is assumed to be fully consumed in the process, rather than emitted to atmosphere. **Table A 7.5.8** therefore does not cover these installations. None of the emission factors in EU ETS are Tier 3, so the table shows the overall emission factors for all tiers of data.

Table A 7.5.8 EU ETS emission factor data for production of lime (kt / Mt lime produced)

| Year | Activity | EU ETS |
|------|-----------------|--------|
| 2005 | Lime production | 192.4 |
| 2006 | | 192.0 |
| 2007 | | 192.9 |
| 2008 | | 180.5 |
| 2009 | | 178.1 |
| 2010 | | 182.1 |
| 2011 | | 178.9 |
| 2012 | | 180.2 |
| 2013 | | 178.5 |
| 2014 | | 179.5 |
| 2015 | | 180.0 |
| 2016 | | 178.4 |
| 2017 | | 179.4 |

These factors compare with a theoretical emission factor based on the stoichiometry of the lime manufacturing process of 214 kt / Mt lime, assuming use of pure limestone. We note that the EU ETS factors are all lower than the theoretical emission factor and this is despite some use of dolomitic limestone in the UK industry which would be expected to further increase the emission factor above the 214 kt/Mt lime factor. The EU ETS data are subject to third party verification, and therefore the emissions data are assumed to be accurate. It is assumed that the reason for this deviation from the theoretical emission factor is due to the production activity data being inflated by either the products containing some proportion of slaked lime (i.e. hydrated product and hence containing a lower proportion of carbon than pure lime) and/or other additives to the lime product which decrease the % carbon content of the lime product.

A 7.5.2.6 Other Industrial Combustion

Table A 7.5.9 summarises EU ETS data for coal, fuel oil and natural gas used by industrial combustion installations.

At first sight, the data for coal looks like it should be reliable enough to be used in the GHGI with 92% or more of emissions based on Tier 3 factors in each year. However, it must be recalled that numerous smaller industrial consumers will not be represented in EU ETS and that the EU ETS data are not fully representative of UK fuels as a whole – see **Section A 7.4** for details. This is also true for EU ETS data for fuel oil and natural gas but here, in addition, very little of the EU ETS data are based on Tier 3 factors. Therefore, none of these data have been used directly in the compilation of the GHGI estimates.

Table A 7.5.9 EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas)

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) | GHGI Carbon Emission Factor |
|------|-------------|----------|---|-----------------------------|
| 2005 | Coal | 98 | 607.1 | 647.8 |
| 2006 | | 98 | 603.0 | 647.8 |
| 2007 | | 99 | 615.7 | 662.4 |
| 2008 | | 94 | 598.6 | 656.6 |
| 2009 | | 92 | 595.4 | 668.5 |
| 2010 | | 88 | 576.5 | 674.3 |
| 2011 | | 91 | 589.0 | 653.4 |
| 2012 | | 90 | 599.2 | 653.7 |
| 2013 | | 95 | 653.4 | 653.3 |
| 2014 | | 98 | 654.3 | 651.2 |
| 2015 | | 100 | 645.8 | 652.1 |
| 2016 | | 100 | 624.9 | 650.9 |
| 2017 | | 100 | 647.4 | 651.2 |
| 2005 | Fuel oil | 48 | 864.7 | 879.0 |
| 2006 | | 74 | 865.3 | 879.0 |
| 2007 | | 50 | 872.3 | 879.0 |
| 2008 | | 35 | 871.4 | 879.0 |
| 2009 | | 39 | 871.3 | 879.0 |
| 2010 | | 40 | 873.0 | 879.0 |
| 2011 | | 51 | 874.2 | 879.0 |
| 2012 | | 49 | 875.1 | 879.0 |
| 2013 | | 44 | 871.3 | 879.0 |
| 2014 | | 48 | 875.0 | 879.0 |
| 2015 | | 55 | 872.1 | 879.0 |
| 2016 | | 63 | 876.2 | 879.0 |
| 2017 | | 65 | 880.0 | 879.0 |
| 2005 | Natural gas | 16 | 1.593 | 1.477 |
| 2006 | | 37 | 1.470 | 1.476 |
| 2007 | | 42 | 1.466 | 1.476 |
| 2008 | | 29 | 1.496 | 1.475 |
| 2009 | | 43 | 1.499 | 1.473 |
| 2010 | | 40 | 1.503 | 1.472 |
| 2011 | | 39 | 1.466 | 1.470 |

| Year | Fuel | % Tier 3 | Average Carbon Emission Factor (Tier 3 sites only) | GHGI Carbon Emission Factor |
|------|------|----------|---|-----------------------------|
| 2012 | | 40 | 1.469 | 1.470 |
| 2013 | | 43 | 1.473 | 1.473 |
| 2014 | | 42 | 1.472 | 1.472 |
| 2015 | | 42 | 1.480 | 1.470 |
| 2016 | | 41 | 1.474 | 1.468 |

Emission factors can also be derived from EU ETS where a high percentage of Tier 3 analysis is evident, for a number of other minor fuels. Due to the very low number of sites that report data for each fuel type, these EU ETS-derived emission factors are confidential and are not tabulated here. The source/activity combinations for which EU ETS emission factor data are used within the inventory are:

- Other industrial combustion / petroleum coke
- Other industrial combustion / waste solvents
- Other industrial combustion / colliery methane

The EU ETS-derived emission factors for colliery methane for each year (2005-2017) are also applied to all other sources using these fuels.

ANNEX 8: UK Domestic Emissions Reporting Requirements

In addition to the reporting requirements of the UNFCCC, Kyoto Protocol (KP) and EU MMR, UK Greenhouse Gas Inventory statistics are published annually in a Department for Business, Energy and Industrial Strategy National Statistics release¹⁸. The geographical coverage of these estimates differs from the UNFCCC, KP and EU MMR coverage, with the totals mainly covering emissions from the UK only (i.e. excluding overseas territories and crown dependencies), although progress towards the Kyoto Protocol is still reported.

As part of the Climate Change Act 2008, the UK committed to reducing greenhouse gas emissions by at least 80 percent by 2050 (relative to the base year¹⁹), with an interim target of reducing greenhouse gas emissions by at least 34 percent by 2020, also relative to the base year. These targets are accompanied by legally binding five-year carbon budgets, which set the trajectory to reaching the targets by placing a restriction on the total amount of greenhouse gases the UK can emit over the five-year period.

Summary tables of the National Statistics release data are presented below. The data are presented in the nine categories used for the UK's National Communications to the UNFCCC and in UK Official Statistics (NC Categories). Note that the scope of emissions used for calculating Carbon Budgets differs slightly from those presented here – for example Carbon Budgets currently exclude NF₃. The 2019 UK GHG emissions statistical release included an update of the UK's performance against the second carbon budget²⁰. Note that performance against the first carbon budget was set in May 2014²¹, updated inventories do not update the first carbon budget or our performance against it.

¹⁸ <https://www.gov.uk/government/collections/final-uk-greenhouse-gas-emissions-national-statistics>

¹⁹ Under the Kyoto Protocol, the UK uses 1990 as the base year for carbon dioxide, methane and nitrous oxide emissions, and 1995 as the base year for the fluorinated gases (or F-gases: hydrofluorocarbons, perfluorocarbons and sulphur hexafluoride). To ensure consistency with our international obligations, the same base year for each greenhouse gas is used under the Climate Change Act.

²⁰ <https://www.gov.uk/government/statistics/final-uk-greenhouse-gas-emissions-national-statistics-1990-2017>

²¹ <https://www.gov.uk/government/statistics/final-statement-for-the-first-carbon-budget-period>

A 8.1 NATIONAL STATISTICS**Table A 8.1.1 Summary table of GHG emissions by NC Category, including net emissions/removals from LULUCF (Mt CO₂eq) – National Statistics coverage (UK only)**

| NC category | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|----------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Energy supply | 277.9 | 238.0 | 221.6 | 231.3 | 207.2 | 190.1 | 165.2 | 145.3 | 121.8 | 112.6 |
| Business | 114.0 | 111.8 | 115.4 | 108.9 | 94.1 | 88.6 | 86.6 | 85.1 | 81.4 | 80.1 |
| Transport | 128.1 | 129.7 | 133.3 | 136.0 | 124.5 | 120.0 | 121.3 | 123.5 | 125.9 | 125.9 |
| Public | 13.5 | 13.3 | 12.1 | 11.2 | 9.5 | 9.1 | 7.8 | 8.0 | 8.2 | 7.8 |
| Residential | 80.1 | 81.6 | 88.7 | 85.7 | 87.5 | 77.5 | 64.8 | 67.4 | 69.8 | 66.9 |
| Agriculture | 54.0 | 52.9 | 50.3 | 47.9 | 44.6 | 44.2 | 45.6 | 45.1 | 45.2 | 45.6 |
| Industrial processes | 59.9 | 50.8 | 27.1 | 20.6 | 12.6 | 12.9 | 13.0 | 12.7 | 10.6 | 10.8 |
| LULUCF ²² | 0.3 | -1.7 | -3.9 | -7.1 | -9.1 | -9.4 | -9.6 | -9.7 | -9.8 | -9.9 |
| Waste management | 66.6 | 69.1 | 62.9 | 49.0 | 29.7 | 23.1 | 20.9 | 20.6 | 20.0 | 20.3 |
| Total | 794.2 | 745.6 | 707.5 | 683.7 | 600.9 | 556.2 | 515.6 | 497.9 | 473.1 | 460.2 |

Table A 8.1.2 Summary table of GHG emissions by Gas, including net emissions/removals from LULUCF (Mt CO₂eq) – National Statistics coverage (UK only)

| Gas | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|------------------|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CO ₂ | 596.3 | 560.1 | 558.3 | 557.9 | 498.3 | 464.0 | 424.9 | 408.3 | 385.8 | 373.2 |
| CH ₄ | 132.5 | 125.9 | 108.4 | 86.9 | 63.9 | 55.5 | 53.3 | 52.7 | 51.1 | 51.5 |
| N ₂ O | 48.2 | 38.6 | 28.5 | 24.4 | 21.3 | 20.3 | 20.8 | 20.3 | 20.2 | 20.5 |
| HFCs | 14.4 | 19.1 | 9.8 | 13.0 | 16.4 | 15.7 | 15.9 | 15.9 | 15.1 | 14.1 |
| PFCs | 1.7 | 0.6 | 0.6 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 |
| SF ₆ | 1.3 | 1.3 | 1.8 | 1.1 | 0.7 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 |
| NF ₃ | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total | 794.360 | 745.6 | 707.5 | 683.7 | 600.9 | 556.2 | 515.6 | 498.0 | 473.1 | 460.2 |

²² Land use, land use change and forestry

ANNEX 9: End User Emissions

A 9.1 INTRODUCTION

This Annex explains the concept of an end user emissions (sometimes also referred to a “final user emissions”, summarises the end user calculation methodology with examples, and contains tables of greenhouse gas emissions according to the end user from 1990 to 2017.

The end user sectoral categories used are consistent with those used in the National Communications (NC) to the UNFCCC. The sectoral categories in the NC are derived from the UNFCCC reporting guidelines on national communications²³.

The purpose of the end user calculations is to allocate emissions from fuel and electricity producers to the energy users - this allows the emission estimates for a consumer of energy to include the emissions from the production of the fuel or electricity they use.

The UNFCCC does not require end user data to be included in the UK's National Inventory Report. These data have been included to provide BEIS with information for their policy support needs.

The tables in this Annex present summary data for UK greenhouse gas emissions for the years 1990-2017, inclusive. These data are updated annually to reflect revisions in the methods used to estimate emissions, and the availability of new information within the inventory. These recalculations are applied retrospectively to earlier years to ensure a consistent time series and this accounts for any differences in data published in previous reports.

Emissions presented in this chapter show emissions from the UK only, consistent with the BEIS UK statistical release.

A 9.2 DEFINITION OF END USERS

The end user²⁴ or calculations allocate emissions from fuel producers to fuel users. The end user calculation therefore allows estimates to be made of emissions for a consumer of fuel, which also include the emissions from producing the fuel the consumer has used.

The emissions included in the end user categories can be illustrated with an example of two end users - the residential sector and road transport:

- Emissions in the **residential** end user category include:
 1. All direct emissions from domestic premises, for example, from burning gas, coal or oil for space heating.

²³ See page 84 of UNFCCC Guidelines contained in FCCC/CP/1999/7 available at: <http://unfccc.int/resource/docs/cop5/07.pdf>

²⁴ An end user is a consumer of fuel for useful energy. A 'fuel producer' is someone who extracts, processes or converts fuels for the end use of end users. Clearly there can be some overlap of these categories but here the fuel uses categories of the UK BEIS publication DUKES are used, which enable a distinction to be made.


2. A portion of indirect emissions used by domestic consumers from: power stations generating electricity; emissions from refineries including refining, storage, flaring and extraction; emissions from coal mines (including emissions due to fuel use in the mining industry itself and fugitive emissions of methane from the mines); and emissions from the extraction, storage and distribution of mains gas.
- Emissions in the **road transport** end user category include:
 1. Direct emissions from motor vehicle exhausts.
 2. A portion of indirect emissions from: refineries producing motor fuels, including refining, storage, flaring and extraction of oil; the distribution and supply of motor fuels; and power stations generating the electricity used by electric vehicles.

A 9.3 OVERVIEW OF THE END USER CALCULATIONS

Fuel and electricity producers also require the use of energy which comes from other producers. Therefore, in the process of reallocating emissions to the end user, emissions are allocated from one to the other and then are reallocated to end users. This circularity results in an iterative approach being used to estimate emissions from categories of end users.

Figure A 9.1 shows a simplified view of the energy flows in the UK (the fuels used in the greenhouse gas inventory have hundreds of uses). This figure shows that while end users consuming electricity are responsible for a proportion of the emissions from power stations they are also responsible for emissions from collieries, and some of these emissions in turn come from electricity generated in power stations and from refineries.

Figure A 9.1 Simplified fuel flows for an end user calculation.



The approach for estimating end user emissions is summarised in the three steps below:

1. Emissions are calculated for each sector for each fuel.
2. Emissions from fuel and electricity producers are then distributed to those sectors that use the fuel according to the energy content²⁵ of the fuel they use (these sectors can include other fuel producers). This distribution is based on inventory fuel consumption data and DUKES electricity consumption data.
3. By this stage in the calculation, emissions from end users will have increased and those from fuel and electricity producers will have decreased. The sum of emissions from fuel producers and power stations in a particular year as a percentage of the total emissions is then calculated. If this percentage, for any year, exceeds a predetermined value (In the model used to determine emissions from end users, the value of this percentage can be adjusted. The tables presented later in this Annex were calculated for a convergence at 0.001%) the process continues at Step 2. If this percentage matches or is less than the predetermined value, the calculation is finished.

Convergence occurs as the fuel flows to the end users are much greater than fuel flows amongst the fuel producers.

While a direct solution could possibly be used it was decided to base the calculation on an iterative approach because:

- This can be implemented in the database structures already in existence for the UK greenhouse gas inventory;
- It can handle a wide range of flows and loops that occur without any of the limits that other approaches may incur; and
- The same code will cover all likely situations and will be driven by tabular data stored in the database.


A 9.4 EXAMPLE END USER CALCULATION

The following example illustrates the methodology used to calculate emissions according to end users. The units in this example are arbitrary.

The example in **Figure A 9.2** has two fuel producers, *power stations* and *collieries*, and three end users, *residential*, *industry* and *commercial*. The following assumptions have been made for simplicity:

- The only fuels used are coal and electricity;
- Coal is the only source of carbon emissions (released from burning coal in power stations to produce electricity and from burning coal in the home for space heating); and
- Commerce uses no coal and so has zero 'direct' emissions.

²⁵ If calorific data for the fuels is not available then the mass of fuel is used instead. This is the case for years prior to 1990.

Figure A 9.2 Fuel use in the example calculation

In **Figure A 9.2**, the tonnes refer to tonnes of coal burnt (black arrows), and the units refer to units of electricity consumed (blue arrows).

In this example the coal extracted by the colliery is burnt in the power station to produce electricity for the end users. Industrial and residential users also directly burn coal. Although the colliery uses electricity produced by the power station, it is not considered to be an end user. The colliery is a 'fuel producer' as it is part of the chain that extracts, processes and converts fuels for the end users.

Table A 9.4.1 summarises the outputs during this example end user calculation.

Table A 9.4.1 Example of the outputs during an end user calculation

| | | Sector | | | | | Unallocated emissions as percentage of total emission | Total emission of carbon (tonnes) |
|-----------------------------------|--------------------------------|----------|---------------|-------------|------------|------------|---|-----------------------------------|
| | | Colliery | Power Station | Residential | Industrial | Commercial | | |
| Coal use (tonnes) | Mass | 100 | 100,000 | 50,000 | 100,000 | 0 | | |
| | Energy content | 25,000 | 25,000,000 | 12,500,000 | 25,000,000 | 0 | | |
| Electricity use (arbitrary units) | Energy units | 10,000 | | 50,000 | 100,000 | 100,000 | | |
| | | | | | | | | |
| Emissions of carbon (tonnes) | Initial | 70 | 70,000 | 35,000 | 70,000 | 0 | 40.02 | 175,070 |
| | Emissions after iteration step | 1 | 2,692 | 28 | 48,476 | 96,951 | 1.55 | 175,070 |
| | | 2 | 1 | 1077 | 49,020 | 98,039 | 0.62 | 175,070 |
| | | 3 | 41 | 1 | 49,227 | 98,454 | 0.02 | 175,070 |
| | | 4 | 0 | 17 | 49,235 | 98,470 | 0.01 | 175,070 |
| | | 5 | 1 | 0 | 49,238 | 98,477 | 0 | 175,070 |
| | | 6 | 0 | 0 | 49,239 | 98,477 | 0 | 175,070 |

The initial carbon emissions are 70% of the mass of coal burnt. The emissions from the power stations are distributed to the other sectors by using the factor:

- $(\text{Electricity used by that sector}) / (\text{total electricity used minus own use by power stations})$;
- Similarly for the colliery emissions the following factor is used; and
- $(\text{Energy of coal used by that sector}) / (\text{total energy of coal consumed used minus own use by collieries})$.

At the end of iteration step one, the commerce sector has 26923 tonnes of carbon emissions allocated to it, mainly derived from power stations. Emissions allocated to the residential and industry sectors have also increased over their initial allocations. However collieries and power stations still have some emissions allocated to them (these come from each other) and so the reallocation process is repeated to reduce these allocations to zero – these two sectors are not end users. The total unallocated (in this example, equal to the total emissions from collieries and power stations) falls in each iteration until the emissions are consistently allocated across the sectors. In this example, six iterations are needed to achieve a consistent allocation across the sectors.

The sum of emissions allocated to the sectors (175,070 tonnes of carbon) remains unchanged from the initial allocation to the allocation in the sixth iteration. This check is an important quality control measure to ensure all emissions are accounted for during the end user calculations.

Figure A 9.3 Comparison of ‘direct’ and end user emissions of carbon according the sectors considered in the end user example




Figure A 9.3 compares the quantities of direct and end user carbon emitted from each sector at the end of the end user calculation. The direct emissions of carbon are from the combustion of coal in the sectors. The direct and end user emissions are from two distinct calculations and must be considered independently – in other words, the direct and end user emissions in each sector must not be summed. The sum of all the direct emissions and the sum of the end user emissions, are identical.

There are relatively large direct emissions of carbon from power stations, residential and industry sectors. The end user emissions from the power stations and the colliery are zero because these two sectors are not end users. The carbon emissions from these two sectors have been reallocated to the residential, industrial and commercial sectors. This reallocation means the end user emissions for the residential and industrial sectors are greater than their ‘direct’ emissions.

A 9.5 END USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY

The approach divides fuel user emissions into 8 categories (see column 1 of **Table A 9.5.1**). For each of these groups, source categories are distributed by the total energy consumption of a group of fuels. For example, for the coal group, the emissions of four source categories are distributed to end users according to the energy use of anthracite and coal combined.

Table A 9.5.1 Sources reallocated to end users and the fuels used

| End user group | Emission sources to be reallocated to end users | Fuels used for redistribution |
|----------------|--|-------------------------------|
| 1. Coke | Gasification processes | Coke |
| | Coke production | Blast furnace gas |
| | Iron and steel – flaring | |
| 2. Coal | Closed Coal Mines | Coal |
| | Coal storage and transport | Anthracite |
| | Collieries – combustion | |
| | Deep-mined coal | |
| | Open-cast coal | |
| 3. Natural gas | Gas leakage | Natural gas |
| | Gas production | |
| | Upstream Gas Production – flaring | |
| | Upstream Gas Production – fuel combustion | |
| | Upstream Gas Production – Gas terminal storage | |
| | Upstream Gas Production – Offshore Well Testing | |
| | Upstream Gas Production - Onshore Oil Loading | |
| | Upstream Gas Production – process emissions | |
| | Upstream Gas Production – venting | |
| | Upstream Gas production – combustion at gas separation plant | |
| | | |
| 4. Electricity | Nuclear fuel production | Electricity |
| | Power stations | |
| | Autogeneration – exported to grid | |
| | Power stations – FGD | |

| End user group | Emission sources to be reallocated to end users | Fuels used for redistribution |
|--------------------------|---|-------------------------------|
| 5. Petroleum | Upstream Oil Production – fuel combustion | Aviation spirit |
| | Upstream Oil Production –flaring | Aviation turbine fuel |
| | Upstream Oil Production –venting | Biodiesel |
| | Upstream Oil Production – Offshore Oil Loading | Bioethanol |
| | Upstream Oil Production – Offshore Well Testing | Burning oil |
| | Upstream Oil Production – Oil terminal storage | Burning oil (premium) |
| | Upstream Oil Production – Onshore Oil Loading | DERV |
| | Upstream Oil Production – process emissions | Fuel oil |
| | Petrol stations – petrol delivery | Gas oil |
| | Petrol stations – vehicle refuelling | LNG |
| | Petrol terminals – storage | LPG |
| | Petrol terminals – tanker loading | Naphtha |
| | Petroleum processes | OPG |
| | Refineries – combustion | Petrol |
| | Refineries – drainage | Petroleum coke |
| | Refineries – flares | Refinery miscellaneous |
| | Refineries – general | Vaporising oil |
| | Refineries – process | |
| | Refineries – road/rail loading | |
| | Refineries – tankage | |
| | Sea going vessel loading | |
| | Ship purging | |
| 6. Solid Smokeless Fuels | Solid Smokeless fuel production | Solid Smokeless Fuels |
| 8. Charcoal | Charcoal production | Charcoal |

Comments on the calculation methodology used to allocate emissions according to the end users are listed below:

- Emissions are allocated to end users on the basis of the proportion of the total energy produced that is used by a given sector. This approach is followed to allow for sectors such as petroleum where different products are made in a refinery;

- Some emissions are allocated to an “exports” category. This is for emissions within the UK from producing fuels, (for example from a refinery or coal mine), which are subsequently exported or sent to bunkers for use outside the UK. Therefore these emissions are part of the UK inventory even if the use of the fuel produces emissions that cannot be included in the UK inventory because it takes place outside the UK;
- No allowance is made for the emission from the production of fuels or electricity outside the UK that are subsequently imported;
- Some of the output of a refinery is not used as a fuel but used as feedstock or lubricants. This is not currently treated separately and the emissions from their production (which are small) are allocated to users of petroleum fuels. This is partly due to lack of data in the database used to calculate the inventory, and partly due to the lack of a clear, transparent way of separating emissions from the production of fuels and from the production of non-fuel petroleum products; and
- End user emissions are estimated for aviation in four categories: domestic take-off and landing, international take-off and landing, domestic cruise and international cruise. This enables both IPCC and UNECE categories to be estimated from the same end user calculation.

Our exact mapping of end user emissions to IPCC categories is shown in **Table A 9.5.2**. The NAEI source sectors and activity names are also shown, as it is necessary to subdivide some IPCC categories. This classification has been used to generate the end user tables for the greenhouse gases given in this section. As this table is for end users, no fuel producers are included in the table.

Table A 9.5.2 End user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|---|-------------------------------------|---|
| Agriculture | 1A4ci_Agriculture/Forestry/Fishing:Stationary | Agriculture - stationary combustion | Burning oil |
| | | | Coal |
| | | | Fuel oil |
| | | | Natural gas |
| | | | Straw |
| | 1A4cii_Agriculture/Forestry/Fishing:Off-road | Agriculture - mobile machinery | Gas oil |
| | | | Petrol |
| | 2D1_Lubricant_Use | Agricultural engines | Lubricants |
| | 3A1a | Enteric | Dairy - Dairy Cows |
| | 3A1b | Enteric | Other cattle - Beef females for slaughter |
| | | | Other cattle - Bulls for breeding |
| | | | Other cattle - Cereal fed bull |
| | | | Other cattle - Cows |
| | | | Other cattle - Dairy Calves Female |
| | | | Other cattle - Dairy In Calf Heifers |
| | | | Other cattle - Dairy Replacements Female |
| | | | Other cattle - Heifers for breeding |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-----------------------|----------------|---|
| | 3A2 | Enteric | Other cattle - Steers |
| | | | Sheep - Ewe |
| | | | Sheep - Lamb |
| | | | Sheep - Ram |
| | 3A3 | Enteric | Pig - Boar |
| | | | Pig - Fattening Pig < 20 kg |
| | | | Pig - Fattening Pig > 80 kg |
| | | | Pig - Fattening Pig 20 to 80 kg |
| | | | Pig - Gilt |
| | | | Pig - Sow |
| | 3A4 | Enteric | Deer |
| | | | Goats |
| | | | Horses |
| | 3B11a | Excreta | Dairy - Dairy Cows |
| | | Managed Manure | Dairy - Dairy Cows |
| | 3B11b 3B12 3B13 | Excreta | Other cattle - Beef females for slaughter |
| | | | Other cattle - Bulls for breeding |
| | | | Other cattle - Cereal fed bull |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|----------------|---|
| | | | Other cattle - Cows |
| | | | Other cattle - Dairy Calves Female |
| | | | Other cattle - Dairy In Calf Heifers |
| | | | Other cattle - Dairy Replacements Female |
| | | | Other cattle - Heifers for breeding |
| | | | Other cattle - Steers |
| | | Managed Manure | Other cattle - Beef females for slaughter |
| | | | Other cattle - Bulls for breeding |
| | | | Other cattle - Cereal fed bull |
| | | | Other cattle - Cows |
| | | | Other cattle - Dairy Calves Female |
| | | | Other cattle - Dairy In Calf Heifers |
| | | | Other cattle - Dairy Replacements Female |
| | | | Other cattle - Heifers for breeding |
| | | | Other cattle - Steers |
| | | Excreta | Sheep - Ewe |
| | | | Sheep - Lamb |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|----------------|---------------------------------|
| | | Managed Manure | Sheep - Ram |
| | | | Sheep - Ewe |
| | | | Sheep - Lamb |
| | | Excreta | Pig - Boar |
| | | | Pig - Fattening Pig < 20 kg |
| | | | Pig - Fattening Pig > 80 kg |
| | | | Pig - Fattening Pig 20 to 80 kg |
| | | | Pig - Gilt |
| | | | Pig - Sow |
| | | Managed Manure | Pig - Boar |
| | | | Pig - Fattening Pig < 20 kg |
| | | | Pig - Fattening Pig > 80 kg |
| | | | Pig - Fattening Pig 20 to 80 kg |
| | | | Pig - Gilt |
| | | | Pig - Sow |
| | 3B14 | Excreta | Poultry - Breeding flock |
| | | | Poultry - Broilers |
| | | | Poultry - Ducks |
| | | | Poultry - Geese |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------------------|
| | | | Poultry - Growing Pullets |
| | | | Poultry - Laying Hens |
| | | | Poultry - Other |
| | | | Poultry - Turkeys |
| | | Managed Manure | Poultry - Breeding flock |
| | | | Poultry - Broilers |
| | | | Poultry - Ducks |
| | | | Poultry - Geese |
| | | | Poultry - Growing Pullets |
| | | | Poultry - Laying Hens |
| | | | Poultry - Other |
| | | | Poultry - Turkeys |
| | | Wastes | Deer |
| | | | Goats |
| | | | Horses |
| | 3B21a | Dairy - Dairy Cows - Direct | Housing |
| | 3B21b | Other cattle - Beef females for slaughter - Direct | Housing |
| | | Other cattle - Bulls for breeding - Direct | Housing |
| | | Other cattle - Cereal fed bull - Direct | Housing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|-----------------------------|
| | | Other cattle - Cows - Direct | Housing |
| | | Other cattle - Dairy Calves Female - Direct | Housing |
| | | Other cattle - Dairy In Calf Heifers - Direct | Housing |
| | | Other cattle - Dairy Replacements Female - Direct | Housing |
| | | Other cattle - Heifers for breeding - Direct | Housing |
| | | Other cattle - Steers - Direct | Housing |
| | 3B22 | Sheep - Ewe - Direct | Storage |
| | | Sheep - Lamb - Direct | Storage |
| | 3B23 | Pig - Boar - Direct | Housing |
| | | Pig - Fattening Pig < 20 kg - Direct | Housing |
| | | Pig - Fattening Pig > 80 kg - Direct | Housing |
| | | Pig - Fattening Pig 20 to 80 kg - Direct | Housing |
| | | Pig - Gilt - Direct | Housing |
| | | Pig - Sow - Direct | Housing |
| | 3B24 | Deer Wastes - Direct | Excreta N managed as manure |
| | | Goats Wastes - Direct | Excreta N managed as manure |
| | | Horses Wastes - Direct | Excreta N managed as manure |
| | | Poultry - Breeding Flock - Direct | Housing |
| | | Poultry - Broilers - Direct | Housing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|--------------------------------------|
| | | Poultry - Ducks - Direct | Housing |
| | | Poultry - Geese - Direct | Housing |
| | | Poultry - Growing Pullets - Direct | Housing |
| | | Poultry - Laying Hens - Direct | Housing |
| | | Poultry - Other - Direct | Housing |
| | | Poultry - Turkeys - Direct | Housing |
| | 3B25 | Dairy - Dairy Cows - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Dairy - Dairy Cows - Indirect Leach | Storage |
| | | Deer Wastes - Indirect Leaching | N leached from manure management |
| | | Deer Wastes - Indirect Volatilisation | N volatilised from manure management |
| | | Goats Wastes - Indirect Leaching | N leached from manure management |
| | | Goats Wastes - Indirect Volatilisation | N volatilised from manure management |
| | | Horses Wastes - Indirect Leaching | N leached from manure management |
| | | Horses Wastes - Indirect Volatilisation | N volatilised from manure management |
| | | | Housing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|---------------|
| | | Other cattle - Beef females for slaughter - Indirect Deposition | Storage |
| | | | Yarding |
| | | Other cattle - Beef females for slaughter - Indirect Leach | Storage |
| | | Other cattle - Bulls for breeding - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Bulls for breeding - Indirect Leach | Storage |
| | | Other cattle - Cereal fed bull - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Cereal fed bull - Indirect Leach | Storage |
| | | Other cattle - Cows - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Cows - Indirect Leach | Storage |
| | | Other cattle - Dairy Calves Female - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Dairy Calves Female - Indirect Leach | Storage |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------|
| | | Other cattle - Dairy In Calf Heifers - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Dairy In Calf Heifers - Indirect Leach | Storage |
| | | Other cattle - Dairy Replacements Female - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Dairy Replacements Female - Indirect Leach | Storage |
| | | Other cattle - Heifers for breeding - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Heifers for breeding - Indirect Leach | Storage |
| | | Other cattle - Steers - Indirect Deposition | Housing |
| | | | Storage |
| | | | Yarding |
| | | Other cattle - Steers - Indirect Leach | Storage |
| | | Pig - Boar - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Boar - Indirect Leach | Storage |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|---------------|
| | | Pig - Fattening Pig < 20 kg - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Fattening Pig < 20 kg - Indirect Leach | Storage |
| | | Pig - Fattening Pig > 80 kg - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Fattening Pig > 80 kg - Indirect Leach | Storage |
| | | Pig - Fattening Pig 20 to 80 kg - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Fattening Pig 20 to 80 kg - Indirect Leach | Storage |
| | | Pig - Gilt - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Gilt - Indirect Leach | Storage |
| | | Pig - Sow - Indirect Deposition | Housing |
| | | | Storage |
| | | Pig - Sow - Indirect Leach | Storage |
| | | Poultry - Breeding Flock - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Breeding Flock - Indirect Leach | Storage |
| | | Poultry - Broilers - Indirect Deposition | Housing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|---------------|
| | | | Storage |
| | | Poultry - Broilers - Indirect Leach | Storage |
| | | Poultry - Ducks - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Ducks - Indirect Leach | Storage |
| | | Poultry - Geese - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Geese - Indirect Leach | Storage |
| | | Poultry - Growing Pullets - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Growing Pullets - Indirect Leach | Storage |
| | | Poultry - Laying Hens - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Laying Hens - Indirect Leach | Storage |
| | | Poultry - Other - Indirect Deposition | Housing |
| | | | Storage |
| | | Poultry - Other - Indirect Leach | Storage |
| | | Poultry - Turkeys - Indirect Deposition | Housing |
| | | | Storage |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|------------------------------------|--|
| | | Poultry - Turkeys - Indirect Leach | Storage |
| | | Sheep - Ewe - Indirect Deposition | Housing |
| | | | Storage |
| | | Sheep - Ewe - Indirect Leach | Storage |
| | | Sheep - Lamb - Indirect Deposition | Housing |
| | | | Storage |
| | | Sheep - Lamb - Indirect Leach | Storage |
| | 3D11 | Arable - Direct | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Grass - Direct | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|--|
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | 3D12a | Dairy - Dairy Cows - Direct | Spreading |
| | | FAM - direct | Manure N applied to soil |
| | | Goats FAM - Direct | Manure N applied to soil |
| | | Horses FAM - direct | Manure N applied to soil |
| | | Other cattle - Beef females for slaughter - Direct | Spreading |
| | | Other cattle - Bulls for breeding - Direct | Spreading |
| | | Other cattle - Cereal fed bull - Direct | Spreading |
| | | Other cattle - Cows - Direct | Spreading |
| | | Other cattle - Dairy Calves Female - Direct | Spreading |
| | | Other cattle - Dairy In Calf Heifers - Direct | Spreading |
| | | Other cattle - Dairy Replacements Female - Direct | Spreading |
| | | Other cattle - Heifers for breeding - Direct | Spreading |
| | | Other cattle - Steers - Direct | Spreading |
| | | Pig - Boar - Direct | Spreading |
| | | Pig - Fattening Pig < 20 kg - Direct | Spreading |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------------------------|
| | | Pig - Fattening Pig > 80 kg - Direct | Spreading |
| | | Pig - Fattening Pig 20 to 80 kg - Direct | Spreading |
| | | Pig - Gilt - Direct | Spreading |
| | | Pig - Sow - Direct | Spreading |
| | | Poultry - Breeding Flock - Direct | Spreading |
| | | Poultry - Broilers - Direct | Spreading |
| | | Poultry - Ducks - Direct | Spreading |
| | | Poultry - Geese - Direct | Spreading |
| | | Poultry - Growing Pullets - Direct | Spreading |
| | | Poultry - Laying Hens - Direct | Spreading |
| | | Poultry - Other - Direct | Spreading |
| | | Poultry - Turkeys - Direct | Spreading |
| | | Sheep - Ewe - Direct | Spreading |
| | | Sheep - Lamb - Direct | Spreading |
| | 3D12b | Sewage Sludge Application - Direct | Sewage sludge N applied to soil |
| | 3D13 | All Horses PRP - Direct | Excreta N returned at grazing |
| | | Dairy - Dairy Cows - Direct | Grazing |
| | | Deer PRP - Direct | Excreta N returned at grazing |
| | | Goats PRP - Direct | Excreta N returned at grazing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------|
| | | Other cattle - Beef females for slaughter - Direct | Grazing |
| | | Other cattle - Bulls for breeding - Direct | Grazing |
| | | Other cattle - Cereal fed bull - Direct | Grazing |
| | | Other cattle - Cows - Direct | Grazing |
| | | Other cattle - Dairy Calves Female - Direct | Grazing |
| | | Other cattle - Dairy In Calf Heifers - Direct | Grazing |
| | | Other cattle - Dairy Replacements Female - Direct | Grazing |
| | | Other cattle - Heifers for breeding - Direct | Grazing |
| | | Other cattle - Steers - Direct | Grazing |
| | | Pig - Boar - Direct | Grazing |
| | | Pig - Fattening Pig < 20 kg - Direct | Grazing |
| | | Pig - Fattening Pig > 80 kg - Direct | Grazing |
| | | Pig - Fattening Pig 20 to 80 kg - Direct | Grazing |
| | | Pig - Gilt - Direct | Grazing |
| | | Pig - Sow - Direct | Grazing |
| | | Poultry - Breeding Flock - Direct | Grazing |
| | | Poultry - Broilers - Direct | Grazing |
| | | Poultry - Ducks - Direct | Grazing |
| | | Poultry - Geese - Direct | Grazing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|------------------------------------|--|
| | | Poultry - Growing Pullets - Direct | Grazing |
| | | Poultry - Laying Hens - Direct | Grazing |
| | | Poultry - Other - Direct | Grazing |
| | | Poultry - Turkeys - Direct | Grazing |
| | | Sheep - Ewe - Direct | Grazing |
| | | Sheep - Lamb - Direct | Grazing |
| | | Sheep - Ram - Direct | Grazing |
| | 3D14 | Arable - Direct | Ammonium Nitrate Residue |
| | | | Ammonium Sulphate and Diammonium Phosphate Residue |
| | | | Calcium Ammonium Nitrate Residue |
| | | | Other Nitrogen Including Compounds Residue |
| | | | Urea Ammonium Nitrate Residue |
| | | | Urea Residue |
| | | Grass - Direct | Ammonium Nitrate Residue |
| | | | Ammonium Sulphate and Diammonium Phosphate Residue |
| | | | Calcium Ammonium Nitrate Residue |
| | | | No Nitrogen Fertiliser Applied Residue |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|--|
| | | | Other Nitrogen Including Compounds Residue |
| | | | Urea Ammonium Nitrate Residue |
| | | | Urea Residue |
| | 3D15 | Cropland management | Mineralisation |
| | 3D16 | Managed Histosols | Land area |
| | 3D21 | Arable - Indirect Deposition | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Dairy - Dairy Cows - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Deer FAM - Indirect Volatilisation | N volatilised from manure applications |
| | | Deer PRP - Indirect Volatilisation | N volatilised from grazing excreta |
| | | Goats FAM - Indirect Volatilisation | N volatilised from manure applications |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|--|
| | | Goats PRP - Indirect Volatilisation | N volatilised from grazing excreta |
| | | Grass - Indirect Deposition | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Horses FAM - Indirect Volatilisation | N volatilised from manure applications |
| | | Horses PRP - Indirect Volatilisation | N volatilised from grazing excreta |
| | | Other cattle - Beef females for slaughter - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Bulls for breeding - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Cereal fed bull - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Cows - Indirect Deposition | Grazing |
| | | | Spreading |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------|
| | | Other cattle - Dairy Calves Female - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Dairy In Calf Heifers - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Dairy Replacements Female - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Heifers for breeding - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Other cattle - Steers - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Pig - Boar - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig < 20 kg - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig > 80 kg - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig 20 to 80 kg - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Pig - Gilt - Indirect Deposition | Grazing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|---------------|
| | | | Spreading |
| | | Pig - Sow - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Breeding Flock - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Broilers - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Ducks - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Geese - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Growing Pullets - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Laying Hens - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Other - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Poultry - Turkeys - Indirect Deposition | Grazing |
| | | | Spreading |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|--|
| | | Sewage Sludge Application - Indirect Volatilisation | N volatilised from sewage sludge applications |
| | | Sheep - Ewe - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Sheep - Lamb - Indirect Deposition | Grazing |
| | | | Spreading |
| | | Sheep - Ram - Indirect Deposition | Grazing |
| | 3D22 | Arable - Indirect Leach | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Arable - Residue Indirect Leach | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|-------------------------------------|--|
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Cropland management | Mineralisation - indirect leach |
| | | Dairy - Dairy Cows - Indirect Leach | Grazing |
| | | | Spreading |
| | | Deer FAM - Indirect Leaching | N leached from manure applications |
| | | Deer PRP - Indirect Leaching | N leached from grazing excreta |
| | | Goats FAM - Indirect Leaching | N leached from manure applications |
| | | Goats PRP - Indirect Leaching | N leached from grazing excreta |
| | | Grass - Indirect Leach | Ammonium Nitrate Application |
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Grass - Residue Indirect Leach | Ammonium Nitrate Application |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|--|
| | | | Ammonium Sulphate and Diammonium Phosphate Application |
| | | | Calcium Ammonium Nitrate Application |
| | | | No Nitrogen Fertiliser Applied |
| | | | Other Nitrogen Including Compounds Application |
| | | | Urea Ammonium Nitrate Application |
| | | | Urea Application |
| | | Horses FAM - Indirect Leaching | N leached from manure applications |
| | | Horses PRP - Indirect Leaching | N leached from grazing excreta |
| | | Other cattle - Beef females for slaughter - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Bulls for breeding - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Cereal fed bull - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Cows - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Dairy Calves Female - Indirect Leach | Grazing |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|---|---------------|
| | | | Spreading |
| | | Other cattle - Dairy In Calf Heifers - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Dairy Replacements Female - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Heifers for breeding - Indirect Leach | Grazing |
| | | | Spreading |
| | | Other cattle - Steers - Indirect Leach | Grazing |
| | | | Spreading |
| | | Pig - Boar - Indirect Leach | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig < 20 kg - Indirect Leach | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig > 80 kg - Indirect Leach | Grazing |
| | | | Spreading |
| | | Pig - Fattening Pig 20 to 80 kg - Indirect Leach | Grazing |
| | | | Spreading |
| | | Pig - Gilt - Indirect Leach | Grazing |
| | | | Spreading |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------|--|---------------|
| | | Pig - Sow - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Breeding Flock - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Broilers - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Ducks - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Geese - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Growing Pullets - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Laying Hens - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Other - Indirect Leach | Grazing |
| | | | Spreading |
| | | Poultry - Turkeys - Indirect Leach | Grazing |
| | | | Spreading |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|---|---|
| | | Sewage Sludge Application - Indirect Leaching | N leached from sewage sludge applications |
| | | Sheep - Ewe - Indirect Leach | Grazing |
| | | | Spreading |
| | | Sheep - Lamb - Indirect Leach | Grazing |
| | | | Spreading |
| | | Sheep - Ram - Indirect Leach | Grazing |
| | 3F11_Field_burning | Field burning | Wheat residue |
| | 3F12_Field_burning | Field burning | Barley residue |
| | 3F14_Field_burning | Field burning | Oats residue |
| | 3F5_Field_burning | Field burning | Linseed residue |
| | 3G1_Liming - limestone | Liming | Limestone |
| | 3G2_Liming - dolomite | Liming | Dolomite |
| | 3H | Fertiliser Application | Urea Application |
| Business | non-IPCC | Agriculture - stationary combustion | Electricity |
| | 1A1ai_Public_Electricity&Heat_Production | Autogenerators | Biogas |
| | 1A2a_Iron_and_steel | Blast furnaces | Blast furnace gas |
| | | | Coke oven gas |
| | | | LPG |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------------------|-----------------------------------|-------------------|
| | | Iron and steel - combustion plant | Natural gas |
| | | | Blast furnace gas |
| | | | Coal |
| | | | Coke |
| | | | Coke oven gas |
| | | | Fuel oil |
| | | | Gas oil |
| | | | LPG |
| | | | Natural gas |
| | 1A2b_Non-Ferrous_Metals | Autogeneration - exported to grid | Coal |
| | | Autogenerators | Coal |
| | | Non-Ferrous Metal (combustion) | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |
| | 1A2c_Chemicals | Chemicals (combustion) | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|-------------------------------------|----------------|
| | 1A2d_Pulp_Paper_Print | Pulp, Paper and Print (combustion) | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |
| | 1A2e_food_processing_beverages_and_tobacco | Food & drink, tobacco (combustion) | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |
| | 1A2f_Non-metallic_minerals | Cement production - combustion | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |
| | | | Petroleum coke |
| | | | Scrap tyres |
| | | | Waste |
| | | | Waste oils |
| | | | Waste solvent |
| | | Lime production - non decarbonising | Coal |
| | | | Coke |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--------------------------------------|------------------|
| | | | Natural gas |
| | | Other industrial combustion | Scrap tyres |
| | 1A2gvii_Off-road_vehicles_and_other_machinery | Industrial off-road mobile machinery | DERV |
| | | | Gas oil |
| | | | Petrol |
| | 1A2gviii_Other_manufacturing_industries_and_construction | Autogeneration - exported to grid | Natural gas |
| | | Autogenerators | Natural gas |
| | | Other industrial combustion | Biomass |
| | | | Burning oil |
| | | | Coal |
| | | | Coke |
| | | | Coke oven gas |
| | | | Colliery methane |
| | | | Fuel oil |
| | | | Gas oil |
| | | | LPG |
| | | | Lubricants |
| | | | Natural gas |
| | | | OPG |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--|---------------------|
| | | | Petroleum coke |
| | | | SSF |
| | | | Waste solvent |
| | | | Wood |
| | 1A4ai_Commercial/Institutional | Miscellaneous industrial/commercial combustion | Coal |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Landfill gas |
| | | | MSW |
| | | | Natural gas |
| | 2B1_Chemical_Industry:Ammonia_production | Ammonia production - combustion | Natural gas |
| | 2B8a_Methanol_production | Methanol production – combustion | Natural gas |
| | 2B8g_Petrochemical_and_carbon_black_production:Other | Chemicals (combustion) | OPG |
| | 2C1b_Pig_iron | Blast furnaces | Coal |
| | 2D1_Lubricant_Use | Industrial engines | Lubricants |
| | 2D4_Other_NEU | Non Energy Use: petroleum coke | Petroleum coke |
| | 2E1_Integrated_circuit_or_semiconductor | Electronics - HFC | Non-fuel combustion |
| | | Electronics - NF3 | Non-fuel combustion |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------------------------|--------------------------|--|
| | 2F1a_Commercial_refrigeration | Commercial Refrigeration | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |
| | 2F1b_Domestic_refrigeration | Domestic Refrigeration | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |
| | 2F1c_Industrial_refrigeration | Industrial Refrigeration | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |
| | 2F1d_Transport_refrigeration | Refrigerated Transport | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|-----------------------------|--|
| | 2F1e_Mobile_air_conditioning | Mobile Air Conditioning | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |
| | 2F1f_Stationary_air_conditioning | Stationary Air Conditioning | Refrigeration and Air Conditioning - Disposal |
| | | | Refrigeration and Air Conditioning - Lifetime |
| | | | Refrigeration and Air Conditioning - Manufacture |
| | 2F2a_Closed_foam_blowing_agents | Foams | Non-fuel combustion |
| | | Foams HFCs for the 2006 GLs | Non-fuel combustion |
| | 2F2b_Open_foam_blowing_agents | One Component Foams | Non-fuel combustion |
| | 2F3_Fire_Protection | Firefighting | Non-fuel combustion |
| | 2F5_Solvents | Precision cleaning - HFC | Non-fuel combustion |
| | 2F6b_Other_Applications:Contained-Refrigerant_containers | Refrigerant containers | Non-fuel combustion |
| | 2F6b_Other_Applications:Contained-Refrigerant_Processing | F-gas handling | Non-fuel combustion |
| | 2G1_Electrical_equipment | Electrical insulation | Non-fuel combustion |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--|---------------------|
| | 2G2_Military_applications | AWACS | Non-fuel combustion |
| | 2G2_Particle_accelerators | Particle accelerators | Non-fuel combustion |
| | 2G2e_Electronics_and_shoes | Electronics - PFC | Non-fuel combustion |
| | | Electronics - SF6 | Non-fuel combustion |
| | | Sporting goods | Non-fuel combustion |
| | 2G2e_Tracer_gas | SF6 used as a tracer gas | Non-fuel combustion |
| | 2G3a_Medical_applications | N2O use as an anaesthetic | Population |
| | 5C2.2b_Non-biogenic:Other | Accidental fires - other buildings | Mass burnt |
| | non-IPCC | Chemicals (combustion) | Electricity |
| | | Food & drink, tobacco (combustion) | Electricity |
| | | Iron and steel - combustion plant | Electricity |
| | | Miscellaneous industrial/commercial combustion | Electricity |
| | | Non-Ferrous Metal (combustion) | Electricity |
| | | Other industrial combustion | Electricity |
| | | Pulp, Paper and Print (combustion) | Electricity |
| Energy Supply | 1A1ai_Public_Electricity&Heat_Production | Power stations | Burning oil |
| | | | Coal |
| | | | Fuel oil |
| | | | Gas oil |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|----------------------------------|--|-----------------|
| | | | Natural gas |
| | | | Petroleum coke |
| | 1A1b_Petroleum_Refining | Refineries - combustion | Natural gas |
| | 1A1ci_Manufacture_of_solid_fuels | Coke production | Natural gas |
| | | Solid smokeless fuel production | Coke |
| | 1A1cii_Oil_and_gas_extraction | Upstream Gas Production - fuel combustion | Gas oil |
| | | Upstream oil and gas production - combustion at gas separation plant | LPG |
| | | | OPG |
| | | Upstream Oil Production - fuel combustion | Natural gas |
| | 1A1ciii_Other_energy_industries | Collieries - combustion | Natural gas |
| | | Gas production | LPG |
| | | Nuclear fuel production | Natural gas |
| | 1B1b_Solid_Fuel_Transformation | Coke production | Coal |
| | | Solid smokeless fuel production | Coal |
| | | | Petroleum coke |
| | non-IPCC | Collieries - combustion | Electricity |
| | | Gas production | Electricity |
| | | Refineries - combustion | Electricity |
| Exports | Aviation_Bunkers | Aircraft - international cruise | Aviation spirit |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|----------------|--|-----------------------|
| | | | Aviation turbine fuel |
| | | Aircraft - international take off and landing | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft between UK and Bermuda - Cruise | Aviation turbine fuel |
| | | Aircraft between UK and Bermuda - TOL | Aviation turbine fuel |
| | | Aircraft between UK and CDs - Cruise | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft between UK and CDs - TOL | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft between UK and Gibraltar - Cruise | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft between UK and Gibraltar - TOL | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft between UK and other OTs (excl Gib. and Bermuda) - Cruise | Aviation turbine fuel |
| | | Aircraft between UK and other OTs (excl Gib. and Bermuda) - TOL | Aviation turbine fuel |
| | Marine_Bunkers | Shipping - international IPCC definition | Fuel oil |
| | | | Gas oil |
| | | Shipping between UK and Bermuda | Fuel oil |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|---|-----------------------|
| | | Shipping between UK and CDs | Fuel oil |
| | | | Gas oil |
| | | Shipping between UK and Gibraltar | Fuel oil |
| | | Shipping between UK and OTs (excl. Gib and Bermuda) | Fuel oil |
| | non-IPCC | Exports | Aviation turbine fuel |
| | | | Burning oil |
| | | | Coke |
| | | | DERV |
| | | | Electricity |
| | | | Fuel oil |
| | | | Petrol |
| | | | SSF |
| Industrial Process | 2A1_Cement_Production | Cement - decarbonising | Clinker production |
| | 2A2_Lime_Production | Lime production - decarbonising | Limestone |
| | 2A3_Glass_production | Glass - general | Dolomite |
| | | | Limestone |
| | | | Soda ash |
| | 2A4a_Other_process_uses_of_carbonates:ceramics | Brick manufacture - all types | Bricks |
| | | Brick manufacture - Fletton | Fletton bricks |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|---|---|-----------------------|
| | 2B1_Ammonia_Production | Ammonia production - feedstock use of gas | Natural gas |
| | 2B10_Chemical_Industry:Other | Chemical industry - general | Process emission |
| | 2B2_Nitric_Acid_Production | Nitric acid production | Acid production |
| | 2B3_Adipic_Acid_Production | Adipic acid production | Adipic acid produced |
| | 2B6_Titanium_dioxide_production | Chemical industry - titanium dioxide | Coke |
| | | | Petroleum coke |
| | 2B7_Soda_Ash_Production | Chemical industry - soda ash | Soda ash produced |
| | 2B8a_Methanol_production | Chemical industry - methanol | Methanol |
| | | | Natural gas |
| | 2B8b_Ethylene_Production | Chemical industry - ethylene | Ethylene |
| | 2B8c_Ethylene_Dichloride_and_Vinyl_Chloride_Monomer | Chemical Industry - ethylene dichloride | Ethylene dichloride |
| | 2B8d_Ethylene_Oxide | Chemical industry - ethylene oxide | Ethylene oxide |
| | 2B8e_Acrylonitrile | Chemical industry - acrylonitrile | Acrylonitrile |
| | 2B8f_Carbon_black_production | Chemical industry - carbon black | Carbon black capacity |
| | 2B9a1_Fluorchemical_production:By-product_emissions | Halocarbons production - by-product | Non-fuel combustion |
| | 2B9b3_Fluorchemical_production:Fugitive_emissions | Halocarbons production - fugitive | Non-fuel combustion |
| | 2C1a_Steel | Basic oxygen furnaces | Dolomite |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|---------------------------------------|---|--------------------------------------|
| | | Electric arc furnaces | Petroleum coke |
| | | | Steel production (electric arc) |
| | | Ladle arc furnaces | Steel production (electric arc) |
| | | | Steel production (oxygen converters) |
| | 2C1b_Pig_iron | Blast furnaces | Coke |
| | | | Fuel oil |
| | 2C1d_Sinter | Sinter production | Coke |
| | | | Dolomite |
| | | | Limestone |
| | 2C3_Aluminium_Production | Primary aluminium production - general | Primary aluminium production |
| | | Primary aluminium production - PFC emissions | Primary aluminium production |
| | 2C4_Magnesium_production | Magnesium cover gas | Non-fuel combustion |
| | 2C6_Zinc_Production | Non-ferrous metal processes | Coke |
| | 2G3b_N2O_from_product_uses:_Other | Other food - cream consumption | Process emission |
| | | Recreational use of N2O | Process emission |
| | 2G4_Other_product_manufacture_and_use | Chemical Industry – other process sources | Process emission |
| | non-IPCC | Blast furnaces | Electricity |
| Land Use Change | 4_Indirect_N2O_Emissions | LULUCF Indirect N2O - Atmospheric Deposition | Non-fuel combustion |
| | | LULUCF Indirect N2O - Nitrogen Leaching and Run-off | Non-fuel combustion |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--|---------------------|
| | 4A_Forest Land_Emissions_from_Drainage | Forest Land - Drainage and rewetting and other management of organic and mineral soils | Non-fuel combustion |
| | 4A1_ Forest Land remaining Forest Land | Forest Land remaining Forest Land - Biomass Burning - Wildfires | Biomass |
| | | Forest Land remaining Forest Land - Carbon stock change | Non-fuel combustion |
| | 4A2_1_Cropland converted to Forest Land | Cropland converted to Forest Land - Carbon stock change | Non-fuel combustion |
| | 4A2_2_Grassland converted to Forest Land | Grassland converted to Forest Land - Carbon stock change | Non-fuel combustion |
| | 4A2_4_Settlements converted to Forest Land | Settlements converted to Forest Land - Carbon stock change | Non-fuel combustion |
| | 4A2_Cropland converted to Forest Land | Cropland converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4A2_Grassland converted to Forest Land | Grassland converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4A2_Land converted to Forest Land_Emissions_from_Fertilisation | Direct N2O emission from N fertilisation of forest land | Non-fuel combustion |
| | 4A2_Settlements converted to Forest Land | Settlements converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4B1_Cropland Remaining Cropland | Cropland remaining Cropland - Biomass Burning - Wildfires | Biomass |
| | | Cropland remaining Cropland - Carbon stock change | Non-fuel combustion |
| | 4B1_Cropland_Remaining_Cropland | Carbon stock change | Non-fuel combustion |
| | 4B2_1_Forest Land converted to Cropland | Forest Land converted to Cropland - Biomass Burning - Controlled Burning | Biomass |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--|---------------------|
| | | Forest Land converted to Cropland - Carbon stock change | Non-fuel combustion |
| | | Forest Land converted to Cropland - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4B2_2_Grassland converted to Cropland | Grassland converted to Cropland - Carbon stock change | Non-fuel combustion |
| | | Grassland converted to Cropland - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4B2_4_Settlements converted to Cropland | Settlements converted to Cropland - Carbon stock change | Non-fuel combustion |
| | 4C_Grassland_Emissions_from_Drainage | Grassland - Drainage and rewetting and other management of organic and mineral soils | Non-fuel combustion |
| | 4C1_Grassland Remaining Grassland | Grassland remaining Grassland - Biomass Burning - Wildfires | Biomass |
| | | Grassland remaining Grassland - Carbon stock change | Non-fuel combustion |
| | | Grassland remaining Grassland - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4C2_1_Forest Land converted to Grassland | Forest Land converted to Grassland - Biomass Burning - Controlled Burning | Biomass |
| | | Forest Land converted to Grassland - Carbon stock change | Non-fuel combustion |
| | | Forest Land converted to Grassland - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4C2_2_Cropland converted to Grassland | Cropland converted to Grassland - Carbon stock change | Non-fuel combustion |
| | 4C2_3_Wetlands converted to Grassland | Wetlands converted to Grassland - Carbon stock change | Non-fuel combustion |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|--|---------------------|
| | 4C2_4_Settlements converted to Grassland | Settlements converted to Grassland - Carbon stock change | Non-fuel combustion |
| | 4D_Wetlands_Emissions_from_Drainage | Wetlands - Drainage and rewetting and other management of organic and mineral soils | Non-fuel combustion |
| | 4D1_Wetlands remaining wetlands | Peat Extraction Remaining Peat Extraction - Carbon stock change | Non-fuel combustion |
| | 4D2_Land converted to Wetlands | Grassland converted to flooded land - Carbon stock change | Non-fuel combustion |
| | | Land converted for Peat Extraction - Carbon stock change | Non-fuel combustion |
| | 4E1_Settlements remaining settlements | Settlements remaining Settlements - Carbon stock change | Non-fuel combustion |
| | | Settlements remaining Settlements - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4E2_1_Forest Land converted to Settlements | Forest Land converted to Settlements - Biomass Burning - Controlled Burning | Biomass |
| | | Forest Land converted to Settlements - Carbon stock change | Non-fuel combustion |
| | | Forest Land converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4E2_2_Cropland converted to Settlements | Cropland converted to Settlements - Carbon stock change | Non-fuel combustion |
| | | Cropland converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |
| | 4E2_3_Grassland converted to Settlements | Grassland converted to Settlements - Carbon stock change | Non-fuel combustion |
| | | Grassland converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization | Non-fuel combustion |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--------------------------------|--|---------------------|
| | 4G_Harvested Wood Products | HWP Produced and Consumed Domestically - Carbon stock change | Non-fuel combustion |
| | | HWP Produced and Exported - Carbon stock change | Non-fuel combustion |
| Public | 1A4ai_Commercial/Institutional | Public sector combustion | Burning oil |
| | | | Coal |
| | | | Coke |
| | | | Fuel oil |
| | | | Gas oil |
| | | | Natural gas |
| | | | Sewage gas |
| | non-IPCC | Public sector combustion | Electricity |
| Residential | 1A4bi_Residential_stationary | Domestic combustion | Anthracite |
| | | | Burning oil |
| | | | Charcoal |
| | | | Coal |
| | | | Coke |
| | | | Fuel oil |
| | | | Gas oil |
| | | | LPG |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|---|---|-----------------------|
| | | | Natural gas |
| | | | Peat |
| | | | Petroleum coke |
| | | | SSF |
| | | | Wood |
| | 1A4bii_Residential:Off-road | House and garden machinery | DERV |
| | | | Petrol |
| | 2D2_Non-energy_products_from_fuels_and_solvent_use:Paraffin_wax_use | Non-aerosol products - household products | Petroleum waxes |
| | 2F4a_Metered_dose_inhalers | Metered dose inhalers | Non-fuel combustion |
| | 2F4b_Aerosols:Other | Aerosols - halocarbons | Non-fuel combustion |
| | 5B1a_composting_municipal_solid_waste | Composting (at household) | Biological waste |
| | 5C2.2b_Non-biogenic:Other | Accidental fires - dwellings | Mass burnt |
| Transport | 5C2.2b_Non-biogenic:Other_Accidental fires (vehicles) | Accidental fires - vehicles | Mass burnt |
| | non-IPCC | Domestic combustion | Electricity |
| | 1A3a_Domestic_aviation | Aircraft - domestic cruise | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Aircraft - domestic take off and landing | Aviation spirit |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|-------------------------------------|---|-----------------------|
| | 1A3bi_Cars | Road transport - cars - cold start | Aviation turbine fuel |
| | | | DERV |
| | | Road transport - cars - motorway driving | Petrol |
| | | | DERV |
| | | Road transport - cars - rural driving | Petrol |
| | | | DERV |
| | | Road transport - cars - urban driving | Petrol |
| | | | DERV |
| | 1A3bii_Light_duty_trucks | Road transport - LGVs - cold start | DERV |
| | | | Petrol |
| | | Road transport - LGVs - motorway driving | DERV |
| | | | Petrol |
| | | Road transport - LGVs - rural driving | DERV |
| | | | Petrol |
| | | Road transport - LGVs - urban driving | DERV |
| | | | Petrol |
| | 1A3biii_Heavy_duty_trucks_and_buses | Road transport - buses and coaches - motorway driving | DERV |
| | | Road transport - buses and coaches - rural driving | DERV |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|----------------------------|--|---------------|
| | | Road transport - buses and coaches - urban driving | DERV |
| | | Road transport - HGV articulated - motorway driving | DERV |
| | | Road transport - HGV articulated - rural driving | DERV |
| | | Road transport - HGV articulated - urban driving | DERV |
| | | Road transport - HGV rigid - motorway driving | DERV |
| | | Road transport - HGV rigid - rural driving | DERV |
| | | Road transport - HGV rigid - urban driving | DERV |
| | 1A3biv_Motorcycles | Road transport - mopeds (<50cc 2st) - urban driving | Lubricants |
| | | | Petrol |
| | | Road transport - motorcycle (>50cc 2st) - urban driving | Petrol |
| | | Road transport - motorcycle (>50cc 4st) - motorway driving | Petrol |
| | | Road transport - motorcycle (>50cc 4st) - rural driving | Petrol |
| | | Road transport - motorcycle (>50cc 4st) - urban driving | Petrol |
| | 1A3bv_Other_road_transport | Road transport - all vehicles biofuels use | Biodiesel |
| | | | Bio-MTBE |
| | | Road transport - all vehicles LPG use | LPG |
| | 1A3c_Railways | Rail - coal | Coal |
| | | Railways - freight | Gas oil |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--------------------------------|--|-----------------------|
| | | Railways - intercity | Gas oil |
| | | Railways - regional | Gas oil |
| | 1A3d_Domestic_navigation | Inland goods-carrying vessels | Gas oil |
| | | Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats) | DERV |
| | | | Gas oil |
| | | | Petrol |
| | | Personal watercraft e.g. jet ski | Petrol |
| | | Sailing boats with auxiliary engines | DERV |
| | | Shipping - coastal | Fuel oil |
| | | | Gas oil |
| | 1A3eii_Other_Transportation | Aircraft - support vehicles | Gas oil |
| | 1A4ai_Commercial/Institutional | Railways - stationary combustion | Burning oil |
| | | | Fuel oil |
| | | | Natural gas |
| | 1A4ciii_Fishing | Fishing vessels | Fuel oil |
| | | | Gas oil |
| | 1A5b_Other:Mobile | Aircraft - military | Aviation spirit |
| | | | Aviation turbine fuel |
| | | Shipping - naval | Gas oil |

| National Communication Category | IPCC Sector | Source Name | Activity Name |
|---------------------------------|--|---|--------------------------|
| | 2D1_Lubricant_Use | Marine engines | Lubricants |
| | | Road vehicle engines | Lubricants |
| | 2D3_Non-energy_products_from_fuels_and_solvent_use:Other | Road transport - urea | Urea consumption |
| | non-IPCC | Railways - regional | Electricity |
| | | Road vehicle engines | Electricity |
| Waste Management | 5A1a_Managed_Waste_Disposal_sites_anaerobic | Landfill | Non-fuel combustion |
| | 5B1a_composting_municipal_solid_waste | Mechanical Biological Treatment - Composting | Biological waste |
| | | Total composting (non-household) | Biological waste |
| | 5B2a_Anaerobic_digestion_municipal_solid_waste | Anaerobic Digestion (other) | Biological waste |
| | | Mechanical Biological Treatment - Anaerobic Digestion | Biological waste |
| | 5C1.1b_Biogenic:Sewage_sludge | Incineration - sewage sludge | Sewage sludge combustion |
| | 5C1.2a_Non-biogenic:municipal_solid_waste | Incineration | MSW |
| | 5C1.2b_Non-biogenic:Clinical_waste | Incineration - clinical waste | Clinical waste |
| | 5C1.2b_Non-biogenic:Other_Chemical_waste | Incineration - chemical waste | Chemical waste |
| | | | |
| | 5D1_Domestic_wastewater_treatment | Sewage sludge decomposition | Non-fuel domestic |
| | | Sewage sludge decomposition in private systems | Non-fuel domestic |
| | 5D2_Industrial_wastewater_treatment | Industrial Waste Water Treatment | Non-fuel combustion |

A 9.6 DETAILED EMISSIONS ACCORDING TO END USER CATEGORIES

The end user categories in the data tables in this summary are those used in National Communications. The end user reallocation includes emissions from the UK, this is the coverage used for the UK statistical release, where the end users data are presented in more detail.

The base year for hydrofluorocarbons, perfluorocarbons, nitrogen trifluoride, and sulphur hexafluoride is 1995. For carbon dioxide, methane and nitrous oxide, the base year is 1990.

Table A 9.6.1 End user emissions from all National Communication categories, MtCO₂ equivalent

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Agriculture | 57.8 | 57.8 | 56.1 | 53.0 | 50.6 | 47.2 | 46.6 | 47.7 | 47.1 | 47.0 | 47.2 |
| Business | 248.6 | 247.8 | 218.1 | 217.1 | 212.0 | 186.6 | 175.7 | 161.0 | 148.4 | 131.7 | 125.6 |
| Energy Supply | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Exports | 9.1 | 9.1 | 13.0 | 12.8 | 16.7 | 15.9 | 13.9 | 12.2 | 11.9 | 11.9 | 12.1 |
| Industrial Process | 65.5 | 63.3 | 53.6 | 29.2 | 21.3 | 13.5 | 13.5 | 13.6 | 13.3 | 11.1 | 11.2 |
| Land Use Change | 0.3 | 0.3 | -1.7 | -3.9 | -7.1 | -9.1 | -9.4 | -9.6 | -9.7 | -9.8 | -9.9 |
| Public | 31.5 | 31.5 | 28.9 | 24.4 | 22.4 | 19.0 | 17.8 | 15.2 | 14.5 | 13.4 | 12.4 |
| Residential | 171.9 | 171.3 | 157.1 | 158.2 | 162.4 | 155.9 | 139.8 | 118.0 | 113.3 | 107.3 | 100.4 |
| Transport | 146.6 | 146.6 | 151.4 | 153.9 | 156.3 | 142.1 | 135.2 | 136.4 | 138.5 | 140.5 | 140.9 |
| Waste Management | 66.6 | 66.6 | 69.1 | 62.9 | 49.0 | 29.7 | 23.1 | 20.9 | 20.6 | 20.0 | 20.3 |
| Total greenhouse gas emissions | 798.0 | 794.4 | 745.6 | 707.5 | 683.7 | 600.9 | 556.2 | 515.6 | 498.0 | 473.1 | 460.2 |

Table A 9.6.2 End user CO₂ emissions from all National Communication categories, MtCO₂ equivalent

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-------------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Agriculture | 10.0 | 10.0 | 9.4 | 8.0 | 8.8 | 7.9 | 7.5 | 7.6 | 7.4 | 7.2 | 7.1 |
| Business | 229.8 | 229.8 | 203.3 | 202.7 | 194.9 | 166.8 | 157.2 | 142.3 | 129.8 | 114.4 | 109.1 |
| Energy Supply | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Exports | 8.4 | 8.4 | 12.1 | 12.1 | 16.0 | 15.3 | 13.2 | 11.6 | 11.4 | 11.4 | 11.5 |
| Industrial Process | 20.9 | 20.9 | 18.9 | 18.1 | 16.7 | 11.2 | 12.7 | 12.8 | 12.5 | 10.4 | 10.5 |
| Land Use Change | -2.0 | -2.0 | -3.9 | -6.0 | -8.9 | -10.7 | -10.9 | -11.1 | -11.2 | -11.3 | -11.3 |
| Public | 29.2 | 29.2 | 27.1 | 23.4 | 21.7 | 18.5 | 17.3 | 14.8 | 14.1 | 13.0 | 12.0 |
| Residential | 156.2 | 156.2 | 145.2 | 149.0 | 154.5 | 148.9 | 133.4 | 112.1 | 107.5 | 101.9 | 95.2 |
| Transport | 142.5 | 142.5 | 147.0 | 150.5 | 153.8 | 140.2 | 133.3 | 134.4 | 136.6 | 138.5 | 138.8 |
| Waste Management | 1.3 | 1.3 | 1.0 | 0.5 | 0.4 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.3 |
| Total greenhouse gas emissions | 596.3 | 596.3 | 560.1 | 558.3 | 557.9 | 498.3 | 464.0 | 424.9 | 408.3 | 385.8 | 373.2 |

Table A 9.6.3 End user CH₄ emissions from all National Communication categories, MtCO₂ equivalent

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------|-----------|------|------|------|------|------|------|------|------|------|------|
| Agriculture | 30.6 | 30.6 | 29.8 | 28.8 | 26.9 | 25.3 | 25.1 | 25.5 | 25.7 | 25.6 | 25.8 |
| Business | 15.5 | 15.5 | 11.7 | 7.4 | 4.8 | 3.7 | 2.9 | 2.9 | 2.7 | 2.3 | 2.3 |
| Energy Supply | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Exports | 0.7 | 0.7 | 0.8 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 |
| Industrial Process | 2.2 | 2.2 | 1.7 | 1.1 | 0.5 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 |
| Land Use Change | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Public | 2.1 | 2.1 | 1.7 | 0.9 | 0.6 | 0.5 | 0.4 | 0.4 | 0.4 | 0.3 | 0.3 |

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|---------------------------------------|--------------|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Residential | 14.3 | 14.3 | 10.7 | 6.7 | 4.9 | 4.5 | 4.0 | 3.6 | 3.6 | 3.2 | 3.1 |
| Transport | 2.5 | 2.5 | 2.2 | 1.5 | 1.0 | 0.8 | 0.7 | 0.7 | 0.7 | 0.6 | 0.7 |
| Waste Management | 64.5 | 64.5 | 67.3 | 61.4 | 47.6 | 28.2 | 21.5 | 19.3 | 19.0 | 18.4 | 18.6 |
| Total greenhouse gas emissions | 132.5 | 132.5 | 125.9 | 108.4 | 86.9 | 63.9 | 55.5 | 53.3 | 52.7 | 51.1 | 51.5 |

Table A 9.6.4 End user N₂O emissions from all National Communication categories, MtCO₂ equivalent

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------|-----------|------|------|------|------|------|------|------|------|------|------|
| Agriculture | 17.2 | 17.2 | 16.8 | 16.2 | 14.9 | 14.0 | 14.0 | 14.6 | 14.1 | 14.1 | 14.3 |
| Business | 1.5 | 1.5 | 1.4 | 1.3 | 1.3 | 1.2 | 1.3 | 1.2 | 1.2 | 1.1 | 1.1 |
| Energy Supply | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Exports | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Industrial Process | 23.9 | 23.9 | 14.4 | 5.4 | 3.1 | 1.5 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Land Use Change | 2.3 | 2.3 | 2.2 | 2.1 | 1.8 | 1.6 | 1.5 | 1.5 | 1.4 | 1.4 | 1.4 |
| Public | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 |
| Residential | 0.7 | 0.7 | 0.6 | 0.5 | 0.5 | 0.5 | 0.6 | 0.5 | 0.5 | 0.4 | 0.4 |
| Transport | 1.6 | 1.6 | 2.2 | 1.9 | 1.5 | 1.1 | 1.2 | 1.2 | 1.3 | 1.3 | 1.4 |
| Waste Management | 0.8 | 0.8 | 0.8 | 0.9 | 1.0 | 1.2 | 1.3 | 1.3 | 1.4 | 1.4 | 1.4 |

| End user category | Base Year | 1990 | 1995 | 2000 | 2005 | 2010 | 2013 | 2014 | 2015 | 2016 | 2017 |
|--------------------------------|-----------|------|------|------|------|------|------|------|------|------|------|
| Total greenhouse gas emissions | 48.2 | 48.2 | 38.6 | 28.5 | 24.4 | 21.3 | 20.3 | 20.8 | 20.3 | 20.2 | 20.5 |

