



National Inventory Report for 1985-2020

Hungary

Compiled by:



**HUNGARIAN
METEOROLOGICAL
SERVICE**

**Unit of National Emissions
Inventories**



2022

PREPARED BY:

Gábor KIS-KOVÁCS
Head of Unit

ENERGY

Gábor KIS-KOVÁCS

INDUSTRIAL PROCESSES AND PRODUCT USE

Krisztina LABANCZ
Erika Lilla LUDÁNYI

AGRICULTURE

Katalin LOVAS
Eszter KIS-CSATÁRI

LULUCF

Dr. Zoltán SOMOGYI
Judit SZAKÁLAS
Dr. Tamás TOBISCH

WASTE

Éva Ilona KIRÁLY
Gábor KIS-KOVÁCS

CONTENT

EXECUTIVE SUMMARY.....	6
<i>ES.1. BACKGROUND INFORMATION.....</i>	<i>6</i>
<i>ES.2. SUMMARY OF NATIONAL EMISSIONS AND REMOVAL RELATED TRENDS.....</i>	<i>6</i>
<i>ES.3. OVERVIEW OF SOURCE AND SINK CATEGORY EMISSION ESTIMATES AND TRENDS.....</i>	<i>7</i>
<i>ES.4. PRECURSORS.....</i>	<i>11</i>
1 INTRODUCTION.....	12
1.1 BACKGROUND INFORMATION AND CLIMATE CHANGE	12
1.2 INSTITUTIONAL ARRANGEMENTS.....	14
1.3 INVENTORY PREPARATION	16
1.4 DATA COLLECTION, PROCESSING AND STORAGE	18
1.5 BRIEF GENERAL DESCRIPTION OF METHODOLOGIES AND DATA SOURCES USED	21
1.6 KEY SOURCE CATEGORIES	22
1.7 QA/QC INFORMATION.....	23
1.8 UNCERTAINTY.....	27
1.9 COMPLETENESS.....	28
1.10 CONSISTENCY CHECKS.....	28
2 TRENDS IN GREENHOUSE GAS EMISSIONS.....	30
2.1 DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS FOR AGGREGATED GREENHOUSE GAS EMISSIONS	30
2.2 DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY GAS	36
2.3 DESCRIPTION AND INTERPRETATION OF EMISSION TRENDS BY CATEGORY	38
2.4 TRENDS OF INDIRECT GASES AND SO ₂	42
3 ENERGY (CRF SECTOR 1)	43
3.1 OVERVIEW OF SECTOR	44
3.2 FUEL COMBUSTION (CRF 1.A).....	48
3.3 FUGITIVE EMISSIONS FROM SOLID FUELS AND OIL AND NATURAL GAS AND OTHER EMISSIONS FROM ENERGY PRODUCTION (CRF 1.B).....	88
3.4 CO ₂ TRANSPORT AND STORAGE (CRF 1.C).....	101
3.5 REFERENCES	102
4 INDUSTRIAL PROCESSES (CRF SECTOR 2)	103
4.1 OVERVIEW OF SECTOR	103
4.2 EMISSION TRENDS	106

4.3	MINERAL PRODUCTS (CRF SECTOR 2.A).....	110
4.4	CHEMICAL INDUSTRY (CRF SECTOR 2.B).....	129
4.5	FLUOROCARBON PRODUCTION (CRF SECTOR 2.B.9)	144
4.6	METAL PRODUCTION (CRF SECTOR 2.C)	144
4.7	OTHER PRODUCTS USE (CRF SECTOR 2.D)	157
4.8	ELECTRONICS INDUSTRY (CRF SECTOR 2.E)	161
4.9	FLUORINATED SUBSTITUTES FOR OZONE DEPLETING SUBSTANCES (CRF SECTOR 2.F)	163
4.10	OTHER PRODUCTS MANUFACTURE AND USE (CRF SECTOR 2.G).....	191
4.11	OTHER PRODUCTION (CRF 2.H).....	197
4.12	REFERENCES	198
5	AGRICULTURE (CRF SECTOR 3)	201
5.1	OVERVIEW OF SECTOR	201
5.2	ENTERIC FERMENTATION (CRF SECTOR 3.A)	217
5.3	MANURE MANAGEMENT (CRF SECTOR 3.B)	238
5.4	RICE CULTIVATION (CRF SECTOR 3.C)	276
5.5	N ₂ O EMISSIONS FROM AGRICULTURAL SOILS (CRF SECTOR 3.D).....	278
5.6	PRESCRIBED BURNING OF SAVANNAS (CRF SECTOR 3.E).....	318
5.7	FIELD BURNING OF AGRICULTURAL RESIDUES (CRF SECTOR 3.F)	318
5.8	LIMING (CRF SECTOR 3.G)	321
5.9	UREA APPLICATION (CRF SECTOR 3.H)	328
5.10	OTHER CARBONATE CONTAINING FERTILIZERS (CRF SECTOR 3.I).....	332
5.11	REFERENCES	336
6	LAND-USE, LAND-USE CHANGE AND FORESTRY (CRF SECTOR 4).....	339
6.1	OVERVIEW OF SECTOR	339
6.2	LAND-USE DEFINITIONS AND CLASSIFICATION SYSTEMS	344
6.3	LAND IDENTIFICATION.....	345
6.4	GENERIC METHODOLOGICAL STEPS TO ESTIMATE EMISSIONS AND REMOVALS	361
6.5	FOREST LAND (CRF SECTOR 4.A)	369
6.6	CROPLAND (CRF SECTOR 4.B)	413
6.7	GRASSLAND (CRF SECTOR 4.C).....	421
6.8	WETLAND (CRF SECTOR 4.D).....	432
6.9	SETTLEMENTS (CRF SECTOR 4.E)	439
6.10	OTHER LAND (CRF SECTOR 4.F).....	443
6.11	UNCERTAINTY ANALYSIS OF THE LULUCF SECTOR. TIME SERIES CONSISTENCY	444

7	WASTE (CRF SECTOR 5).....	451
7.1	OVERVIEW OF SECTOR	451
7.2	SOLID WASTE DISPOSAL IN LANDFILLS (CRF SECTOR 5A)	453
7.3	BIOLOGICAL TREATMENT OF SOLID WASTE (CRF SECTOR 5B)	464
7.4	INCINERATION OF WASTE (CRF SECTOR 5C)	466
7.5	WASTEWATER TREATMENT AND DISCHARGE (CRF SECTOR 5D)	469
8	OTHER (CRF SECTOR 6)	479
9	INDIRECT CO₂ AND NITROUS OXIDE EMISSIONS	479
10	RECALCULATIONS AND PLANNED IMPROVEMENTS	480
11	KP-LULUCF	488
11.1	GENERAL INFORMATION	488
11.2	LAND-RELATED INFORMATION	499
11.3	ACTIVITY-SPECIFIC INFORMATION	504
11.4	ARTICLE 3.3	526
11.5	ARTICLE 3.4	531
11.6	OTHER INFORMATION	541
11.7	INFORMATION RELATING TO ARTICLE 6.....	541
11.8	NIR TABLES	541
11.9	REFERENCES	542
12	INFORMATION ON THE ACCOUNTING OF THE KYOTO PROTOCOL UNITS.....	545
12.1	CALCULATION OF THE COMMITMENT PERIOD RESERVE (CPR).....	546
13	INFORMATION ON CHANGES IN NATIONAL SYSTEM.....	546
14	INFORMATION ON CHANGES IN NATIONAL REGISTRY	547
15	INFORMATION ON MINIMIZATION OF ADVERSE IMPACTS IN ACCORDANCE WITH ARTICLE 3, PARAGRAPH 14	549

EXECUTIVE SUMMARY

ES.1. Background information

Pursuant to the United Nations Framework Convention on Climate Change (UNFCCC), Hungary, as a Party of the Convention, has been preparing annual inventories of greenhouse gas emissions using the IPCC methodology since 1994. The aim of a greenhouse gas (GHG) inventory is to give an as complete and accurate as possible state of the art estimation of anthropogenic emissions by sources and removal by sinks of greenhouse gases not controlled by the Montreal Protocol. In accordance with the Kyoto Protocol, the following direct greenhouse gases are taken into account: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), Sulphur hexafluoride (SF₆), and nitrogen trifluoride (NF₃). The quality of the inventory is controlled by Hungarian and international experts regularly.

The GHG inventory is compiled by the Hungarian Meteorological Service as laid down by a government decree. The participation of the National Land Centre (NLC) together with the Forest Research Institute of the University of Sopron and the National Food Chain Safety Office (NFCO) as compilers of the whole LULUCF sector is formalized by the same governmental decree. Also, other institutions and external experts are involved in the process of inventory preparation, e.g., the Hungarian Central Statistical Office, Hungarian Energy and Public Utility Regulatory Authority, and Institute of Agricultural Economics Nonprofit Kft. (AKI), just to name a few.

The main purpose of this National Inventory Report is to describe the input data and calculation methodologies on which the emissions estimates are based thus increasing the transparency of the inventory. The present report refers to the inventory time series for the years 1985-2020. The NIR provides relevant background information on institutional arrangements, QA/QC procedures and other information underlying the inventory compilation in Chapter 1. In Chapter 2 the trends for aggregated greenhouse gas emissions are discussed. The chapters following provide detailed information on each of the main source categories. Chapter 10 discusses details of recalculations and planned improvements. In the Annexes key category analysis and complementary methodological information can be found.

ES.2. Summary of National Emissions and Removal Related Trends

In 2020, total emissions of greenhouse gases in Hungary were **62.8 million tonnes** carbon dioxide equivalents (CO₂-eq) excluding the LULUCF sector. Taking into account also the mostly carbon absorbing processes in the LULUCF sector, the net emissions of Hungary were 56.0 million tonnes CO₂-eq in 2020. Being about 6 tonnes, the Hungarian per capita emissions are below the European average.

Compared to the base year (average of 1985-87), 1990, and 2005, our current emissions are lower by 43%, 34%, and 18%, respectively.

This significant reduction was partly a consequence of the regime change in Hungary (1989-90) which brought in its train radical decline in the output of the national economy. The production decreased in almost every economic sector including also the GHG relevant sectors like energy, industry and agriculture. Then, between 2005 and 2013, after a period of about 14 years of relatively stagnant

emission level (1992-2005), GHG emissions fell again quite significantly by 24 per cent. The global financial and economic crisis exerted a major impact on the output of the Hungarian economy, consequently on the level of GHG emissions as well resulting in a quite significant drop of 9% between 2008 and 2009. Then, after a smaller increase in 2010, emissions decreased further in the following four years. In contrast, the decline in economic output stopped in the first quarter of 2010, and Hungary not only reached the pre-crisis level of GDP again in 2014 but exceeded it even in 2015.

After 2013, emissions started growing again. Up to 2017, the overall increase reached 12%. After four years of increase, emissions have remained more or less at the same level between 2017 and 2019. In 2020, however, emissions fell by almost 3% to around 2016 levels, mainly due to a significant reduction in transport emissions as a consequence of COVID-19.

The most important greenhouse gas is carbon dioxide accounting for 75% of total GHG emissions. The main source of CO₂ emissions is burning of fossil fuels for energy purposes, including transport. CO₂ emissions have decreased by 45% since the middle of the 80's. Methane represents 13% in the GHG inventory. Methane is generated mainly at waste disposal sites and in animal farms, but the fugitive emissions of natural gas systems (i.e., transmission, storage, and distribution) are also important sources. CH₄ emissions are by 40% lower than in the base year. Nitrous oxide contributes 8% to the total GHG emissions. Its main sources are agricultural soils, and manure management. N₂O emissions are 55% lower compared to base year. The total emissions of fluorinated gases amount to 4%.

Table ES.1 *Trend of emissions by GHGs, excluding LULUCF (Gg CO₂-eq)*

GHG	BY	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
CO ₂	85,418	73,226	61,391	58,365	60,276	52,069	46,653	47,113	49,515	49,463	49,235	47,284
CH ₄	13,590	12,830	10,744	10,591	9,705	8,894	8,384	8,334	8,353	8,292	8,234	8,220
N ₂ O	11,135	8,377	4,750	5,405	5,608	3,714	4,515	4,781	4,781	4,848	4,850	5,013
HFCs	NO	0	36	204	754	1,250	1,821	1,895	1,964	2,055	2,159	2,189
PFCs	371	376	223	282	280	4	4	4	2	3	3	3
SF ₆	7	12	51	82	90	92	118	128	114	97	101	109
NF ₃	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
TOTAL	110,521	94,821	77,194	74,929	76,714	66,021	61,496	62,256	64,729	64,756	64,581	62,818

Base year (BY)=average of 1985-87

ES.3. Overview of Source and Sink Category Emission Estimates and Trends

By far, the biggest emitting sector was the energy sector contributing 71% to the total GHG emission in 2020. Industrial processes and product use (IPPU), and Agriculture had a similar share of 12% each. The waste sector contributed 5%. Compared to the base year, emissions significantly decreased in the energy (-45%), industrial processes and product use (-49%), and agriculture (-39%). In contrast, emissions in the waste sector have increased since 1985-87 (+5%). The land use, land-use change and forestry (LULUCF) sector shows fluctuating behavior. Looking at the most recent trends since 2005, emissions significantly dropped in the energy and industrial processes sectors by 23% and 15%, respectively. The agriculture sector seems to have recovered and could show an increase of 19% since 2005. The previous growing trend turned back in the waste sector (-19%).

The **energy sector** was responsible for 71% of total GHG emissions in 2020. Production and use of energy generate most greenhouse gases, largely CO₂. Currently, 16% of domestic primary energy supply is nuclear, 12% is renewable which means that the remaining - overwhelming - part of primary

energy demand has to be met by fossil fuels. Natural gas accounts for the largest share (47%) of incinerated, largely fossil fuels, followed by petroleum products (30%). Emissions have been positively influenced by the fact that the proportion of coal with higher specific emissions has fallen from 30% to 7% in the last 35 years, which is well below the current share of biomass in fuel consumption (14%).

The three most important sources of emissions in the energy sector are transport, energy industries, and “other sector” (mostly including residential and other buildings), each of which accounted for 20% of total national emissions in 2020. Energy use and emissions from manufacturing industries and construction contributed 8% to domestic total emissions. Fugitive (mostly methane) emissions from the domestic natural gas system represent 3% of total emissions

In recent years, the transport sector became the largest emitter, not only within the energy sector but also across all sectors, as transport accounted for 23% of total national emissions in 2019. However, as a result of preventative measures for COVID-19, emissions from the road transport-dominated sector fell sharply by 14% in 2020. Nevertheless, compared to the previous low point, 2013, transport emissions were still 25% higher. In addition, based on preliminary fuel sales data, it appears that the decline in 2020 can be considered temporary as rising emissions are expected for 2021 in this sector.

Considering energy industries, domestic electricity production increased a little by 2%. However, coal-based power production dropped by a further 9% (after a decrease of 13% last year) whereas natural gas fired power plants with more favorable specific emission levels increased their production by 4% (after an increase of 19% in 2019). Another welcome development is the sharp increase in the use of solar energy: 4% of gross electricity production now comes from solar energy. The share of nuclear power generation was at around 50% over the last few years. The relatively large share of electricity import decreased from 31% in 2014-2015 to 25% in 2020. As a result of all the above, emissions from energy industries decreased by 2% in 2020.

2020 qualified “only” as the eighth warmest year in the last 120 years, therefore the heating demand increased a little after the two warmest years. Compared to 2019, total fuel consumption in dwellings increased by 5%. While the use of natural gas increased by 8%, firewood consumption did not change significantly and the marginal coal consumption decreased further by 20%. (It is worth noting that the use of the two main energy sources often changes in opposite directions: since 2005, natural gas consumption has fallen by 25%, while firewood has increased by 33%.) Although household emissions increased by 7%, emissions from the total energy sector were 4% lower in 2020 than in 2019 due to declining transport and industrial emissions.

Table ES.2 Trend of emissions and removals by sector (including LULUCF, Gg CO₂-eq)

Sector	BY	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
Energy	80,189	69,386	59,032	56,448	57,312	49,871	44,259	45,105	46,774	46,485	46,348	44,386
Industry	15,076	11,750	8,228	8,183	9,071	6,395	6,936	6,655	7,435	7,730	7,696	7,733
Agriculture	12,030	9,994	6,006	6,141	6,133	5,655	6,752	7,068	7,071	7,119	7,113	7,297
LULUCF	-2,361	-3,099	-6,034	-849	-5,871	-4,416	-5,656	-4,496	-5,119	-4,422	-4,907	-6,821
Waste	3,225	3,690	3,929	4,157	4,198	4,100	3,549	3,428	3,449	3,421	3,425	3,402
TOTAL	108,160	91,722	71,160	74,080	70,842	61,605	55,839	57,760	59,610	60,334	59,674	55,997

Base year (BY)=average of 1985-87

The **industrial processes and product use sector** contributed 12% to total GHG emissions in 2020. The most important greenhouse gas was CO₂, contributing 66% to total sectoral GHG emissions, followed

by F-gases with 28%. In 2020, 34% of the emissions came from chemical industry, followed by 28% from product uses as ODS substitutes. Mineral industry has 17%, metal industry has 14% contribution to sectoral GHG emissions, respectively. Other product uses (containing SF₆ and N₂O) and non-energy products from fuels and solvent use have the smallest influence on the 2020 IPPU inventory with 4% and 2%, respectively. Process related industrial emissions decreased by 49% between the base year and 2020, and by 15% between 2005 and 2020.

GHG emissions from the iron and steel producing sector decreased by 9% compared to 2019 because of the decreasing production of pig iron. Production of ammonia, urea and nitric acid increased in 2020, causing 8% increase in emissions from the chemical industry sector. In the production of mineral industry, the years-long increasing trend halted in 2020, GHG emissions from this sector were 11% lower in 2020 than in 2019. Emissions from the non-energy products from fuels and solvent use increased by 16% mainly due to the increase in the lubricant consumption.

A third of industrial emissions come from the operation of equipment containing F-gases and from use of F-gas containing products. Category 2.F.1. (Refrigeration and air-conditioning) accounts for 87% of total F-gas emissions, which is still increasing. Despite the continuous regulation of high GWP gases emission has not already stopped in this sector.

Although, charging into new equipment for some gases already is forbidden, a lot of cooling systems have been operated with gases which have higher GWP and this is the reason of this trend.

In 2020, the **agriculture sector** accounted for 12% of total emissions. Emissions from agriculture include CH₄ and N₂O gases. 86 per cent of total N₂O emissions were generated in agriculture in 2020. Emissions from agriculture have decreased by 39% over the period of 1985-2020. The bulk of this reduction occurred in the years between 1985 and 1995, when agricultural production fell by more than 30 per cent, and livestock numbers underwent a drastic decline.

Between 1996 and 2008, agricultural emissions had stagnated around 6.2 Mt with fluctuations up to 4.6%. Behind this trend there were compensatory processes. While the number of livestock decreased further leading to lower emission, the use of fertilizers increased by 68% in the period 1995-2007 which caused growing nitrous oxide emissions from agricultural soils. In 2008 the significantly rising fertilizer prices led to lower fertilizer use, which resulted in some reduction in the emission levels.

Agricultural emissions decreased both in 2009 and 2010. A major reduction in emissions occurred in 2009, when 11 per cent decline in swine population also contributed to the downward trend. Agricultural emissions, after hitting the lowest point in 2010, had increased until 2018, mainly because of the increase in the inorganic fertilizer use, cattle livestock, and milk production per cow.

The GHG-emissions reflect the restructuring in the agricultural production has taken place since 2004, namely the increased ratio of crop to livestock production. Share of CH₄ emissions, which derive mainly from the animal husbandry, has decreased, while the N₂O emissions, originating primarily from the crop production has grown, since 2004.

Certain types of inorganic fertilizers as urea containing fertilizers and calcium ammonium nitrate (CAN) fertilizers contribute to the agricultural GHG-emissions not only with their nitrogen, but also their carbon content. In Hungary CAN fertilizers have become increasingly popular in the recent years, as a result N₂O and CO₂ emissions has tripled from this source since 2005.

In 2019, emissions growth temporarily slowed down, mainly due to the decreasing swine livestock and synthetic fertilizer use, but emissions increased again in 2020. The upward trend was mainly due to an increase in fertilizer use and beef cattle numbers.

The **Land Use Land-Use Change and Forestry sector** has been a net carbon sink mainly because of the huge amount of carbon uptake of forests, which in turn is due to continuous afforestation efforts and sustainable forest management. The complex dynamics of the land use and land-use changes leads to highly fluctuating estimates of sectoral removals. Over the period 1990 to 2020 our estimates indicate an average annual net removals of 4 million tonnes CO₂-eq ranging from 0.8 million tonnes in 2000 to 6.8 million tonnes CO₂ in 2020. In 2020, the net removals of forests amounted to 6.6 million tonnes CO₂ the main reason of which being the lower harvest due to the COVID pandemic. The harvested wood product pool is close to a carbon equilibrium with a small net sink in the last six years. The non-forestry land-use sectors used to be small net sinks before 2016 but they have been small sources since then.

The **waste sector** was responsible for 5% of total national GHG emissions in 2020. The largest category was solid waste disposal on land, representing 85% in 2020, followed by wastewater treatment and discharge (9%), biological treatment of solid waste (4%), and incineration of waste without energy recovery (1%). In contrast with other sectors, emissions from the waste sector are by 5% higher now than in the base year. However, the growth in emissions stopped in the last decade, and a reduction of 19% could be observed between 2005 and 2020. The degradation process in solid waste disposal sites is quite slow which means that waste that were disposed many years earlier have still an influence on current emission levels. However, the amount of disposed waste had dropped significantly since 2005 (e.g., landfilled municipal waste decreased by 50%) consequently methane emissions started to decrease as well. GHG emissions from wastewater handling have a pronounced decreasing trend due to a growing number of dwellings connected to the public sewerage network.




Figure ES. 1 Change in greenhouse gas emissions from base year (BY, 1990-2020)

ES.4. Precursors

NO_x, CO and NMVOC gases are referred to as indirect gases because they (together with SO₂) influence atmospheric warming indirectly, via secondary effects. Nitrogen oxides, carbon monoxide and (non-methane) volatile organic compounds are precursor of ozone which is itself a naturally occurring greenhouse gas. Sulphur dioxide can contribute to formation of aerosols that scatter some of the solar radiation back into space. Calculation of the emissions of these gases is required by the UNFCCC reporting guidelines. It should be noted that Hungary (as well as the other European countries) has calculated the emissions of such gases for several decades and the Geneva Convention of 1979 (CLRTAP) also laid down such obligations. Emissions are reported consistently in the above two reporting regimes. The following table shows the main trends in emissions:

Table ES. 3 Emissions of indirect gases, excluding LULUCF (Gg)

GASES	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
NO _x	245	189	187	178	146	127	119	121	119	114	107
CO	1434	964	850	695	535	469	445	445	373	363	343
NMVOC	306	210	188	172	130	126	126	123	117	117	112
SO ₂	829	613	427	43	30	24	23	28	23	17	16

The substantial reduction in sulphur dioxide emissions is attributable to the decreased use of fossil fuels in general and the decreasing share of coal with higher sulphur content. After 2000, further reductions were observed due to the introduction of SO₂ precipitators in coal-fired power stations. Reduced carbon monoxide emissions are obviously a consequence of decreased fuel uses. The decrease in NO_x emissions is relatively moderate due to the increasing significance of transport.

1 INTRODUCTION

1.1 Background information and climate change

Hungary submitted the First National Communication in 1994 when the country joined the UN Framework Convention on Climate Change (hereinafter referred to as the Convention). In conjunction with this, the greenhouse gas inventories of the preceding years were prepared. Since then, inventories have been compiled annually as required. According to the Convention, year 1990 considered as the general reference level was not adequate for Hungary as a base year because the economic output of the country was already on the descending course as a result of the ongoing transition to market economy. Instead of 1990, the average of years 1985, 1986 and 1987 (hereinafter referred to as "base year") was selected because these three years represented a certain level of stability in the fluctuating economic output. This request was accepted by the COP.

With the introduction of additional greenhouse gases, it was necessary to select the corresponding base years. (This was particularly important for HFCs because such gases had been used increasingly as replacements for ozone depleting chlorofluorocarbons since the early 1990's.) Hungary has chosen 1995 as the base year for fluoride gases. The process of inventory preparation has been improved year by year. The inventory teams did their best to meet the changing and growing requirements. Particular emphasis was placed on determining the specific emission factors for Hungary.

The regional effects of the global climate change can clearly be seen on the Hungarian observations. The annual averages of temperature in Hungary are very similar to the well-known wave of the global temperature since the beginning of the 20th century.




Figure 1.1. Linear trends in annual mean temperature (°C) over the period 1901-2020 in Hungary, based on the homogenized, interpolated dataset of the Hungarian Meteorological Service

The yearly average temperature was 11.5 °C in 2020 in Hungary. 2020 was the eighth warmest in the last 120 years, closing the hottest decade since 1901, based on controlled, homogenized and interpolated data of the Hungarian Meteorological Service. The year 2020 fits in well with the global warming trend in terms of temperature. The national average annual average temperature rises

significantly at a 90% confidence level based on a linear trend estimate from a long time series beginning in 1901. The change in the national average annual temperature over the last 120 years (1901 to 2020) has averaged +1.23°C.

According to the homogenized data, the country-wide average of yearly total precipitation was 615 mm in 2020, which is only 2 percent more than the 30-year average from the period 1981 to 2010. Over the last 120 years, between 1901 and 2020, we have seen a moderate decline, averaging 2.8%, based on the exponential trend adjusted to annual precipitation amounts.




Figure 1.2 Exponential trends in annual precipitation sum (mm) over the period 1901-2020 in Hungary, based on the homogenized, interpolated dataset of the Hungarian Meteorological Service

1.2 Institutional arrangements

The minister responsible for the environment has overall responsibility for the Hungarian Greenhouse Gas Inventory and the Hungarian National System for Climate Reporting. He is responsible for the institutional, legal and procedural arrangements for the national system and the strategic development of the national inventory. The Ministry of Environment and Water had been abolished after the elections in spring 2010, and its tasks have been taken over by the Ministry of Rural Development. The structure and duties of the ministries changed again somewhat after the elections in 2014, and the Ministry of Rural Development turned to Ministry of Agriculture which nevertheless has the same responsibilities regards environmental matters. Therefore, the designated *single national entity* is now the Ministry of Agriculture.

Contact details of the single national entity are as follows:

Ministry of Agriculture

Head office:	1055 Budapest, Kossuth Lajos tér 11.
Postal address:	1860 Budapest
Phone:	+36-1-795-2000
Fax:	+36-1-795-0200
E-mail:	info@am.gov.hu, sajto@am.gov.hu

István Nagy, Dr., Minister of Agriculture

Postal address:	1055 Budapest, Kossuth L. tér 11.
Phone:	+36-1-795-3723
Fax:	+36-1-795-0072
E-mail:	miniszter@am.gov.hu

The national system has to be operated by the minister responsible for the environment but, as prescribed by legislation, in consent and cooperation with the ministers responsible for energy policy, forest management, agricultural policy, and national budget. Within the Ministry for Innovation and Technology, i.e., the ministry responsible for energy policy, a Climate Policy Department has been established that plays a coordinating and supervisory role in the national system. The head of this department is Hungary's current UNFCCC Focal Point.

At the end of 2006, a Greenhouse Gas Inventory Division (GHG division) was established in the Hungarian Meteorological Service (HMS) for the preparation and development of the inventory. This division is responsible for most inventory related tasks, compiles the greenhouse gas inventories and other reports with the involvement of external institutions and experts on a contractual basis and supervises the maintenance of the system. In 2015, the name of the division was changed to Unit of National Emissions Inventories.

At the very end of 2009, a new government decree 345/2009 (XII.30.) on data provision relating to GHG emissions was put into force. This decree confirmed the designation of the Hungarian Meteorological Service as the compiler institute. As a new element, the participation of the Forestry Directorate of the National Food Chain Safety Office (NFCSO, Forestry Directorate) together with the National Agricultural Research and Innovation Centre (hereafter referred to as NARIC) Forest Research Institute was formalized by this decree. These two institutes were responsible for the forestry part of the LULUCF sector and for the supplementary reporting on LULUCF activities under Articles 3.3 and 3.4 of the Kyoto Protocol by making recommendations to HMS of the content of the inventory. The govt. decree had to be revised according to the changing EU regulations and reporting needs, therefore Govt. Decree 345/2009 (XII.30) was replaced by Govt. Decree 528/2013 (XII.30.).

1 January 2015, a new government decree 278/2014. (XI. 14) entered into force in Hungary designating the National Food Chain Safety Office (NFCSO) Plant Protection and Soil Conservation Directorate, together with the Hungarian Chamber of Agriculture, responsible for the development of the GHG inventory of the non-forest sectors. (This is a change from the previous system, in which the Hungarian Meteorological Service was responsible for the non-forest sectors. In order to facilitate this change, and in order to ensure a smooth transition to the application of the IPCC 2006 Guidelines, a new estimation system has been recently developed for, and together with, the NFCSO by an external expert.)

In 2019, the Hungarian National Land Centre (NLC) was established from several organizations including the Forestry Department of NFCSO. Thus, from this year onwards, together with NARIC the Forestry Department of NLC is responsible for the forestry part of the LULUCF sector.

In 2020 the Forest Research Institute became part of the University of Sopron. From this year onwards the Forest Research Institute of the University of Sopron together with the Forestry Department of NLC is responsible of the forestry part of the LULUCF sector.

The Hungarian Meteorological Service is a central office under the control of the Ministry of Agriculture. The duties of the Service are specified in a Government Decree from 2005. The financial background of operation is determined in the Finances Act. HMS has introduced the quality management system ISO 9001:2000 for the whole range of its activities in 2002 to fulfill its tasks more reliably and for the better satisfaction of its partners. The Unit of National Emissions Inventories functions as part of the Department of Climate and Ambient Air. The Unit of National Emissions Inventories of the Hungarian Meteorological Service coordinates the work with other involved ministries, government agencies, consultants, universities and companies in order to be able to draw up the yearly inventory report and other reports to the UNFCCC and the European Commission. The Unit of National Emissions Inventories can be regarded as a core expert team of four people. The division of labor and the sectoral responsibilities within the team are laid down in the QA/QC plan and other official documents of HMS. The Head of Unit coordinates the teamwork and organizes the cooperation with other institutions involved in inventory preparations. He is responsible for the compilation of CRF tables and NIR. Within the team the experts are responsible for different sectors. Besides, a QA/QC coordinator and an archive manager have been nominated.

Most parts of the inventory (energy, industrial processes and product use, and waste) are prepared by the experts of the Unit of National Emissions Inventories themselves. The agriculture sector is prepared with the involvement of the Institute of Agricultural Economics Nonprofit Kft. (AKI). The whole LULUCF sector is compiled by the institutes listed in the above-mentioned government decree. As before, and also complying with the decree mentioned above, the Forestry Department of the NLC is responsible for the GHG inventory of the forestry sector. Quality control for the forestry sector is provided by the

Forest Research Institute of the University of Sopron. Data for the estimation of non-forest related emissions is also provided by the Central Statistical Office, the Hungarian Mining Authority and National Directorate General for Disaster Management.

The following table summarizes the institutional arrangements:

<i>Function</i>	<i>Institution</i>	<i>Responsibilities</i>
	Ministry of Agriculture	
Single national entity	(in consent and cooperation with Ministry for Innovation and Technology, and Ministry of Finance)	<ul style="list-style-type: none"> • Supervision of national system • Official consideration and approval of inventory
Inventory coordination and compilation	HMS Unit of National Emissions Inventories	<ul style="list-style-type: none"> • Provision of work plan • Contracting consultants • Inventory preparation of Energy, Industry, Agriculture and Waste sectors • Compilation of the CRF and NIR • Archiving • Coordinating QA/QC activities • Reporting to UNFCCC secretariat
Inventory preparation of the LULUCF sector and LULUCF activities under the KP. (by law)	Hungarian National Land Centre (NLC) Forest Research Institute of the University of Sopron National Food Chain Safety Office (NFCSO)	<ul style="list-style-type: none"> • Data collection, choice of methods and EFs, inventory preparation • Compilation of the relevant parts of the CRF and NIR

1.3 Inventory preparation

The annual inventory cycle is carried out in accordance with the principles and procedures set out in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. As a general method of preparing the inventory, the procedures described in the IPCC Guidelines are applied and the latest CRF Reporter software is used. Usually, the sectoral experts are responsible for the choice of methods and emission factors in consultation with the head of Unit of National Emissions Inventories. According to the recommendations of the IPCC Guidelines, the calculation methods are chosen by taking into account the technologies available in Hungary whenever possible. The calculation of emissions occurs basically by using the formula: $AD \times EF$, where the activity data (AD) can be raw material or product or energy use etc. Part of the available data (e.g., production data) can directly be entered into the IPCC tables; others require previous processing and conversion. For example, energy data are not always available in the required depth and resolution. The default emission factors (EF) are being gradually replaced by country-specific emission factors characteristic of domestic technologies. Efforts are made to use the highest possible Tier method, especially in case of key categories. After preliminary quality control of the basic data, the necessary calculations are carried out with the coordination of the core team. The

sectoral data are compiled and - after repeated checks - unified by using the CRF Reporter software. QA/QC activities are described in more detail in chapter 1.7 and the full, updated QA/QC Plan (synthesizing the former QA/QC Plan, the old ISO Procedure and the old archiving manual) is included in Annex 5.

Recalculation of some data-series of the inventory can be justified by several reasons. Just to name a few, QA/QC procedures, ERT recommendations, changing for higher Tier methodologies can lead to a recalculation. As a basic rule, whenever new information emerges that improves the quality or accuracy of the emission data, the emissions are recalculated. The Hungarian Meteorological Service funds research and development projects for the improvement of the inventory whenever possible. Recalculations are always documented in the relevant chapter of the national inventory report.

The inventory cycle can be summarized with the following table based on our QA/QC plan:

Date/deadline	Item	To
From May to November	Overview of sectors to identify areas for possible improvements; Data collection, choice of methodologies, Start of calculations Repeated checks	
From September to December (and April)	Calculations from external expert	
From September to December (and April)	Calculations, checks, archiving	
08 January	Main features for National Inventory Report (CRF tables and part of NIR) for approval	National Authority
15 January	Official submission	EU
Between January and March	QC procedures including EU internal review	
08 March	National Inventory Report final version for approval	National Authority
15 March	National Inventory Report, Official submission	EU
Between March and April	QC procedures in the process of finalizing the NIR and CRF tables	
08 April	National Inventory Report for approval	National Authority
15 April	Official submission	UNFCCC
31 July	Preliminary inventory of year x-1	EU
From 15 th of April to October	Archiving, QA/QC and Development Plan	internal

A Figure presenting the inventory cycle is included in Annex 5.

To summarizing the above, the main compiler institutes are: (1) the Hungarian Meteorological Service (HMS) responsible for all sectors except for LULUCF, and (2) the National Land Centre together with

the Forest Research Institute of the University of Sopron and the National Food Chain Safety Office responsible for the LULUCF part of the inventory.

The Meteorological Service, where an inventory team is located, is authorized by law to collect the necessary data. Calculations are either carried out by the Unit of National Emissions Inventories of the HMS or by external experts on contractual basis. The inventory report is approved by two ministers: (1) minister for innovation and technology, (2) minister for agriculture (responsible for environment, agricultural policy, land administration, and forest management) before submission to the UNFCCC.

1.4 Data collection, processing and storage

Data is collected in several ways and throughout the whole yearly cycle of the inventory preparation. Sector specialists of the core team (or external experts on contractual basis) are making the data inquiry and collection in addition to the data arriving based on the reporting obligation set up by Govt. Decree 278/2014. (XI. 14) as described below in more detail). Plant specific data are collected if possible (especially in case of power stations, heating stations and industrial technologies) but statistical databases are also heavily used as source of information. The most important statistical publications are the Statistical Yearbook of Hungary, the Environmental Statistical Yearbook of Hungary and the Environmental Report of Hungary published by the Hungarian Central Statistical Office (HCSO) and the Energy Statistical Yearbook published earlier by the Energy Efficiency, Environment and Energy Information Agency. As regards energy statistics, the practice has changed in recent years. The compiler institute relies less to classic statistical publication and more to databases sent by the Hungarian Energy and Public Utility Regulatory Authority to the IEA and Eurostat. The compiler institute receives the same completed joint questionnaires that are sent to the international organizations which ensure the consistency with data reported under Regulation (EC) No 1099/2008.

Since the use of ETS data has several advantages, the inventory team was granted access (by the same Govt. decree) to the verified emissions database held earlier by the National Inspectorate for Environment and Nature (now: National Climate Protection Authority).

In addition to statistical data, contacts were established with the representatives of a number of major emitting sectors. Moreover, information from the web sites of international associations (e.g., International Iron and Steel Institute, IISI) is used as well.

For the calculation of F-gas emissions, import data from the Customs Office and Police were used together with data obtained directly from companies importing and using fluorinated gases and information from cooling industry associations, the Hungarian Monitoring and Certification Body (OMKT-HMBC), the Hungarian Electrotechnical Association (MEE) and the National Directorate General for Disaster Management, Ministry of the Interior (NDGDM).

Data reported pursuant to Article 6(1) of Regulation (EC) No 842/2006 on F-gases (for the consistency check required by the MMR) is received from Hungarian contact point responsible for the reporting under 842/2006/EC. This data provision is also included in Govt. Decree 278/2014. (XI. 14).

The Act LX of 2007 on the implementation framework of the UN Framework Convention on Climate Change and the Kyoto Protocol thereof aims to give direct data collection authorization to the Ministry of Agriculture in order to collect data for the national system for climate reporting and gives a permanent status to the system. Relevant paragraphs for data collection are the following: "The state

authorities having disposal of the data necessary to operate the National Registration System and the organizations emitting at least 100 tons of carbon dioxide equivalent per year shall provide these data for the National Registration System in accordance with the provisions of a separate legal instrument.” “The data (...) necessary to fulfill international data supply shall be provided for the National Registration System irrespective of the fact that they are qualified as individual data pursuant to the relevant provision of Act XLVI of 1993 on statistics.” This separate legal instrument, the above-mentioned government decree 278/2014. (XI. 14) on data provision relating to GHG emissions prescribes compulsory data provision for GHG inventory purposes for numerous governmental bodies and emitters. QA/QC Activities connected to data collection are regulated by the updated QA/QC Plan included in Annex 5.

The Govt. Decree 278/2014 has two annexes with quite long lists of data providers and data to be submitted. The first annex relates to information that are to be submitted by government institutions, Annex II relates to data provision obligation by firms. Information contained in these two annexes cover the basic data needs for inventory compilation. In addition to what is included in the above annexes, paragraph 5 states that:

In addition to the regular data provision as specified in the annexes, the inventory compiler institutes are entitled to request additional data in order to supplement or refine the available information, or to make the corrections required during reviews.

To facilitate exchange of information, the Hungarian Meteorological Service became "VIP customer" of the statistical office recently, which means we have a dedicated VIP contact person within the statistical office in case of additional data needs.

All the collected data, where relevant, are also used for the elaboration of the air pollutant emission inventories (NFR). Therefore, the consistency with the reporting of air pollutant emission inventories under Directive 81/2001 and the Convention on Long-range, Transboundary Air Pollution (CLRTAP) is ensured.

A copy of all data, information necessary for the compilation of the given annual inventory is stored in printed or electronic form either by the Unit of National Emissions Inventories of the HMS or by the institutions involved in inventory preparations. Significant steps were taken to create a central archive in the premises of the Hungarian Meteorological Service where all background data would be stored.

The most important paper information archived already in the Service is the following:

- Statistical Yearbooks of Hungary from the year 1961
- Environmental Statistical Yearbook of Hungary from 1996
- Energy Statistical Yearbook published by the Energy Efficiency, Environment and Energy Information Agency from 1985.
- Hungarian Statistics on Road Vehicles (in electronic format since 2000)
- National, regional and local emission survey of the Hungarian road, rail, water-borne and air transport (1995-2004) made yearly by the Institute of Transport Sciences

Lots of background data are stored by contracted expert institutions as well, which increases the security of data availability. Nevertheless, at least a copy of all important information has been transferred to the HMS. The following information is stored elsewhere:

- Data from individual industrial plants – Ministry of Agriculture;

- ETS data, registry - National Climate Protection Authority;
- Forestry statistics – Hungarian National Land Centre;
- Wastewater data – General Directorate of Water Management, Ministry of Agriculture, Ministry of Interior, HCSO.

Electronic information is stored on disks on a fileserver with a regular backup. The whole data files are backed up once a week, while the implements (those files that have been modified since the last saving) are saved two times a week. The data are stored on tape storage system. The cassettes of the data storage system are stored far from the recording system, in another room, which is air conditioned and equipped with an up-to-date fire service system. All events connected with the data saving are logged in accordance with the documents of the Quality Management System of HMS.

As HMS is a central office, strict record management, documentation and archiving rules apply in general. HMS's general record management, documentation and archiving regulation have been amended in 2011. The new regulation had been supplemented with a new chapter relating to the Unit of National Emissions Inventories. The main elements of the former proposal of the 'manual for the maintenance and management of the archiving system' as the procedures of documents and data handling had been formalized in this regulation.

A particular issue of this regulation is to ensure the integrity of the data handling in relation to the GHG inventory. The regulation has specific rules on handling confidential data as well. These rules are as follows:

Confidential data are

- accessible only for members of the Unit of National Emissions Inventories. They are not allowed to be forwarded to other institute or persons, except for the ERT
- it is not allowed to make hard copies of these documents, only one electronic copy can be made, which is stored on the server of the Unit of National Emissions Inventories;
- data stored on the server of GHG are protected by password;
- it is not allowed to take out any confidential information from the HMS, not even their copies;
- the original hard copies are not allowed to be forwarded to the Hungarian Environmental Archives; they are stored in the records of the HMS's Unit of National Emissions Inventories.

The new regulation has been endorsed by the Minister of Public Administration and Justice and has been in force since January of 2012.

The directories of the server, where the data of the Unit of National Emissions Inventories are stored have access protection, so they are available only for the staff of the Unit in charge of the different sectors of the GHG inventory. It is important to note that there are different directories for all the calculations and drafts (working folder) and for the submitted reports and incoming data which cannot be modified. Within the Unit of National Emissions Inventories of HMS, the nominated archive manager is responsible for the maintenance of the archiving system in close cooperation with the IT Department of the Service.

The most important elements of the previously planned procedural manual for management and maintenance of the archiving system (archiving manual) have been included formally into the general record management, documentation and archiving regulation of the HMS and the new QA/QC Plan of

the Unit of National Emissions Inventories of the HMS. (Instead of the introduction of a new regulation the already existing regulations have been amended and supplemented with the issues of the draft manual.). So, these two regulations define the QA/QC activities connected to data collection, processing, storage and the documentation and archiving activities of the Unit of National Emissions Inventories. Further development of the system may include the incorporation of other emission data, which are relevant to air pollution.

1.5 Brief general description of methodologies and data sources used

The IPCC Guidelines provide methodologies for estimating emissions and removals of greenhouse gases. However, the basic idea is not greenhouse gas specific, the same approach is used for other pollutants, and other emission inventories, as well (e.g. see the EMEP/EEA air pollutant emission inventory guidebook). The basic equation is as simple as this:

$$\text{Emission} = \text{AD} \times \text{EF},$$

where AD stands for activity data which represents some human activity (e.g. fuel use, industrial production, animal population, dwellings supplied with public sewerage, area of vineyard abandonment), whereas EF is the emission factor that quantifies the emission (or removal) per unit of activity. For example, in energy industry, which is the most important source category, emission factors for combusting natural gas or lignite are 56.1 t CO₂/TJ and 107.9 t CO₂/TJ, respectively; the importance of the mix of fuels used to produce energy becomes apparent at a glance.

Emission factors are usually dependent on several other factors, used technologies etc. which leads us to the concept of tiers. A tier represents a level of methodological complexity. In the Guidelines, usually three tiers are provided. Tier 1 is the basic method, where activity data are usually aggregated national statistics and the emission factors are default values representing typical process conditions. Higher tier methodologies are more demanding in terms of complexity and data requirements as they require country-specific information on the used technologies, facility level data whenever possible, or use of complex models. For key categories, i.e. categories that have a significant influence on a country's total inventory of greenhouse gases in terms of the absolute level of emissions and removals, the trend in emissions and removals, or uncertainty in emissions and removals, it is required to apply higher tier methods. Accordingly, the compilers of the Hungarian inventory aim at taking into account the technologies available in Hungary to the extent possible. For example, the emission trading system of the European Union makes possible to have access to facility level activity and verified emission data.

Although this basic equation can widely be used, in some source categories other approaches are used. For example, mass balance method is used for estimating the change in carbon content of living biomass in forests, or in case of solid waste disposal sites, a calculation method is applied which assumes that the degradable organic component in waste decays slowly throughout a few decades.

To ensure that the national inventory fulfils its main purpose, namely monitoring the country's compliance with its commitments, it has to meet certain quality standards, in other words it has to be accurate, complete, consistent, comparable and transparent (ACCCT). The first two requirements need no special explanation: an inventory is accurate, if it has no systematic bias towards under- or overestimations, whereas a complete inventory covers all relevant sources and sinks, and gases within the borders of the country. The next two criteria are closely linked to the requirements of the UNFCCC. Consistency ensures that the trends in the times-series of the inventory reflect real differences in emissions, and not caused by any methodological changes. National greenhouse gas

inventories of all countries shall be comparable; therefore, the submitted information shall be compiled in accordance with the UNFCCC reporting guidelines and the 2006 IPCC guidelines.

More detailed source specific information on used data and methodologies can be found in Chapters 3-9 in this inventory report.

1.6 Key source categories

Based on the IPCC Tier 1 methodology, 39 key categories have been identified. Two categories (2.A.4 *Other Process Uses of Carbonates – CO₂ and 4(II). Emissions and removals from drainage and rewetting and other management of organic and mineral soils – CO₂*) were dropped out in this submission compared to the last submission. On the other hand, 2G Other Product Manufacture and Use - N₂O became key.

Table.1.6.1. Key category analysis summary

KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	Level	Trend	excl	incl
1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO ₂	X	X	X	X
1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO ₂	X	X	X	X
1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO ₂	X	X	X	X
1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO ₂		X		X
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO ₂	X	X	X	X
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO ₂	X	X	X	X
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO ₂	X	X	X	X
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO ₂	X	X	X	X
1.A.3.b Road Transportation	CO ₂	X	X	X	X
1.A.3.c Railways	CO ₂		X	X	X
1.A.4 Other Sectors - Liquid Fuels	CO ₂	X	X	X	X
1.A.4 Other Sectors - Solid Fuels	CO ₂	X	X	X	X
1.A.4 Other Sectors - Solid Fuels	CH ₄		X	X	X
1.A.4 Other Sectors - Gaseous Fuels	CO ₂	X	X	X	X
1.A.4 Other Sectors - Biomass	CH ₄	X	X	X	X
1.B.1 Fugitive emissions from Solid Fuels	CH ₄		X	X	X
1.B.2.b Fugitive Emissions from Fuels - Oil and Natural Gas - Natural Gas	CH ₄	X	X	X	X
2.A.1 Cement Production	CO ₂	X		X	X
2.A.2 Lime Production	CO ₂		X	X	
2.B.1 Ammonia Production	CO ₂	X	X	X	X
2.B.2 Nitric Acid Production	N ₂ O		X	X	X
2.B.8 Petrochemical and Carbon Black Production	CO ₂	X	X	X	X

2.C.1 Iron and Steel Production	CO ₂	X	X	X	X
2.C.3 Aluminium Production	PFCs		X	X	X
2.F.1 Refrigeration and Air conditioning	F-gases	X	X	X	X
2.G Other Product Manufacture and Use	N ₂ O	X		X	X
3.A Enteric Fermentation	CH ₄	X	X	X	X
3.B Manure Management	CH ₄	X		X	X
3.B Manure Management	N ₂ O	X		X	X
3.D.1 Direct N₂O Emissions From Managed Soils	N ₂ O	X	X	X	X
3.D.2 Indirect N₂O Emissions From Managed Soils	N ₂ O	X		X	X
4.A.1 Forest Land Remaining Forest Land	CO ₂	X	X		X
4.A.2 Land Converted to Forest Land	CO ₂	X	X		X
4.B.1 Cropland Remaining Cropland	CO ₂	X	X		X
4.B.2 Land Converted to Cropland	CO ₂	X	X		X
4.E.2 Land Converted to Settlements	CO ₂		X		X
4.G Harvested Wood Products	CO ₂	X	X		X
5.A Solid Waste Disposal	CH ₄	X	X	X	X
5.D Wastewater Treatment and Discharge	CH ₄	X	X	X	X

Note: L = Level assessment; T = Trend assessment, incl = including LULUCF, excl = excluding LULUCF.

KP-LULUCF key categories could be determined on the basis of the following criteria: If the sum of emissions and removals from an activity under the Kyoto Protocol exceeds that of the smallest key category under the Convention (including LULUCF), it is good practice to identify the activity as key.

The smallest key category under the Convention was 2G Other Product Manufacture and Use - N₂O with a CO₂-eq emission of 230.69 kt. As all KP activities resulted in higher net CO₂ equivalent emissions/removals, all three categories, i.e. A.1. Afforestation and reforestation, A.2. Deforestation, and B.1. Forest management, can be considered as key.

1.7 QA/QC information

The national system has to ensure high quality of the inventory, i.e. to ensure that the inventory is transparent, consistent, comparable, complete and accurate. These principles guide the internal expert team that maintains the system. QA/QC activities are performed in two levels: based on the ISO 9001 standards and following the IPCC recommendations. The updated QA/QC Plan that entered into force in 2013 aims to integrate these two sets of requirements. The QA/QC Plan was updated again and entered into force in the beginning of 2016 in order to follow the changes of legislation and the Guidebook, and the change of the name and acronym (from “UHG” to “NELO”) of the Unit of National Emissions Inventories.

ISO activities

The Hungarian Meteorological Service introduced the quality management system ISO 9001:2000 in 2002 for the whole range of its activities which was quite unique among meteorological services. However, GHG inventory preparation was not among its activities in that time. Therefore, the scope

of our ISO accreditation had to be modified and lots of efforts have been made to bring also the national system under the umbrella of the ISO QM system. Several regulatory ISO documents were created, among others: ISO procedure on the activities of the GHG Division; QA/QC plan; registers and records for quality checks and documentation. Naturally, from that time on, HMS level QA/QC activities apply for the Unit of National Emissions Inventories as well, such as general quality objectives, application of QA/QC Manual of the HMS, QA/QC regarding contractors, etc. Further information on quality management system of the HMS is available in English at: <http://www.met.hu/en/omsz/minosegiranynitas/>

In 2012 the ISO procedure of the GHG division was reviewed, and the former QA/QC Plan with the archiving manual was integrated into it. ISO document No.: ELFO_UHG_401.01 entered into force on 4th January 2013 can be regarded as the QA/QC Plan required for inventory preparation. In addition, the records used for documentation of QA/QC and other standardized activities have also been renewed. On 21 May 2014, an update of the QA/QC Plan (No.: ELFO_UHG_401.02) entered into force in order to insert the recommendation of the review of the year before regarding the documentation of QA activities. The update of the QA/QC Plan (No.: ELFO_NELO_401.01) that entered into force in the beginning of 2016 did not contain significant changes, mainly changes of names and references to legislation and the Guidebook are reflected. The records and their functions are the following at the moment:

- NELO01: QA/QC checklist: to be filled in by sectoral experts which includes a compulsory check list, summary of results of checks, suggestions for corrective actions and planned improvements;
- NELO02: Data quality check: to be filled in case of data providers and external experts on data quality;
- NELO03: Development Plan: to be filled in every year by the end of the inventory cycle based on the outcome of all reviews and own experience;
- NELO04: Responsibility: for the specification of the sectoral responsibilities of the core team and the QA/QC coordinator
- NELO05: Data source logbook: for the standardized documentation of data sources;
- NELO06: Uncertainty and NELO07: Key category analysis; for the standardized documentation of uncertainty and key category analysis.
- NELO08: QA activities logbook: record for the documentation of QA activities.

The records and the English translation of the QA/QC Plan are presented in the Annex 5 of the NIR.

The QA/QC Plan contains detailed description of the data collection, inventory preparation and reporting processes, regulates the documentation and archiving activities in order to ensure transparency and reproducibility of the inventory the same as before, especially:

- ELFO_NELO_401.01 formalizes the data collection and inventory preparation procedure as it is described also in chapters 1.4 and 1.5 above. It is important to note that the authorization of HMS for collecting non-public data has been raised in a legally binding level by since 2009 when 528/2013. (XII.30.) Govt. Decree entered into force. In addition, Act LX of 2007 on the implementation framework of the UN Framework Convention on Climate Change and the Kyoto Protocol authorizes HMS to collect confidential data if needed as well. ELFO_NELO_401.01 prescribes that any data used by the preparation of the inventory have to be documented and archived.

- Documentation and archiving: As mentioned in chapter 1.4 above, the Hungarian Meteorological Service is a central office under the control of the Ministry of Agriculture. Strict documentation and archiving are a basic requirement by the institution. The HMS has a documentation and archiving manual valid for the whole institution, which defines that all the incoming letters and emails containing data have to be registered in the central registry system of HMS. This ensures that every document is traceable. In addition, data, data sources and calculation files and background documents for every inventory submission need to be documented and archived by the sectoral experts. The exact process of documentation and archiving (naming and location) is detailed in document ELFO_NELO_401.01.
- Data quality check. Besides self-checking, the entries of data providers and external experts are checked regularly which is an interactive process during the whole inventory cycle. Significant changes compared to previous data shall be explained. NELO02 QC record was created for standardized documentation of evaluation of data quality by the data providers which can be regarded as a continuous development. The QA/QC plan prescribes the obligation of filling in the records mentioned before, including Development Plan, where first of all the recommendations of the last years' reviews conducted by the expert review team of the UNFCCC have to be taken into consideration as much as possible every year.

Having an ISO system in place has an advantage of being subject to regular internal and external audits. During our last external audit, the activities of the Unit of National Emissions Inventories were audited as well. Our system was audited favorably in the end of March 2007; and our ISO certification has been renewed in January 2012 and a comprehensive external audit was again performed in January 2014, 2015, 2016, 2017 and 2018 as well. On the 5th April 2013 and on the 12th December 2014 and 11th October 2016 an internal audit has been performed too. In all cases the result was a few non-significant recommendations. Therefore, we can claim that the GHG inventory is subject to and our procedures are in line with ISO 9001:2008.

As part of the QA and verification activities there is an ongoing QA procedure between the NFCSO and the Forest Research Institute of the University of Sopron who are involved in the LULUCF of the inventory.

Peer-reviews on the other chapters will be conducted depending on available resources.

UNFCCC reviews

Reviews conducted by expert review teams of the UNFCCC usually have important impact on the development of quality and transparency of our emission estimates. The submitted table, MMR-IRArticle9_HU_2022JAN.xlsx summarizes the main findings in the 2021 annual inventory review report and Hungary's response to those findings. These findings will be provided in the Annexes of the March submission of the NIR. Many of the recommendations of the 2021 review processes have been implemented in the 2022 submission, while further recommendations of the reviews will be addressed when feasible in next submissions.

EU review processes

In 2012 the EU carried out a comprehensive individual technical review concentrating on the years 2005, 2008, 2009 and 2010, which can be regarded as an additional QA activity. Starting with the data reported for the year 2013, the European Commission conducts annual reviews of the national inventory data submitted by Member States in accordance with the Regulation (EU) No 525/2013 (the 'Monitoring Mechanism Regulation' or MMR).

Decision No 406/2009/EC (the 'Effort Sharing Decision' or ESD) converted the annual reporting cycle into an annual commitment cycle requiring a comprehensive review of Member States' greenhouse gas (GHG) inventories within a shorter time frame than the current UNFCCC inventory review, to enable the use of flexibilities and the application of corrective action, where necessary, at the end of each relevant year.

Regulation (EU) No 525/2013 (the 'Monitoring Mechanism Regulation' or MMR) set up at Union level a review process of the greenhouse gas inventories submitted by Member States to ensure that compliance with the ESD is assessed in a credible, consistent, transparent and timely manner.

The annual ESD reviews consist of two steps. In the first step the transparency, accuracy, consistency, completeness and comparability of Member States' inventories are checked pursuant to Articles 29 and 33 of Regulation (EU) No 749/2014 (the 'MMR Implementing Regulation').

The first step of the ESD review is performed by the European Environment Agency (EEA) European Topic Centre for Atmospheric Climate Mitigation (ETC/ACM). The main purpose of step 1 is to identify potential significant issues which are handed over to step 2 for more detailed analysis.

In the second step of the annual a Technical Expert Review Team (TERT) analyses the significant issues mainly by performing checks pursuant to Articles 32 and 33 of Regulation (EU) No 749/2014 to identify cases where inventory data are prepared in a manner that is inconsistent with the UNFCCC guidance documentation or Union rules.

The EU annual review processes contribute significantly to the quality assurance procedures. The main actions and outcomes of the annual EU review processes were as follows:

In 2015, Hungary participated in STEP 2 of the ESD trial review on a voluntary basis.

The main objectives of the trial review were to provide, through recommendations, informal feedback from a review team on emission estimates because no reviews were carried out on UN level, and to provide additional support to the improvement of the Member States GHG-inventories. During the reviews all the sectors (LULUCF excepted) had been thoroughly reviewed, and several recommendations have been formulated and the majority of the recommendations were implemented in subsequent submissions.

In May 2016, a comprehensive review was carried out by the EU for the compliance years 2013 and 2014, and for the years 2005, 2008, 2009 and 2010 pursuant to Monitoring Mechanism Regulation (EU) 525/2013 Article 27.

In 2017, 2018, and 2019 the checks performed during the annual ESD review did not identify any significant issues, therefore Hungary was not subject to a second step of the 2017-2019 annual ESD reviews.

In 2020, all EU Member States were subject of a comprehensive review. One of the main conclusions of the review was the following: *"The reviewers raised 38 issues with Hungary during the first and the second step of the 2020 comprehensive ESD review. The TERT provided recommendations for 3 of these issues. Other issues raised during the comprehensive review were clarified and are considered non-issues for the ESD review 2020."* All three recommendations have been addressed for this submission.

Quattro-lateral QA/QC cooperation

Hungary joined a quattro-lateral QA/QC cooperation with the Czech Republic, Slovakia, and Poland. In 2017 some issues from the Energy and the IPPU sector were reviewed. In 2018, the focus was put on the waste sector whereas mostly LULUCF experts had fruitful discussions in the usual summer meeting

in 2019. As an outcome of the meeting recommendations and good practices were shared to learn from the other Parties, to encourage the improvement and to enhance the quality of emission inventories. Unfortunately, due to COVID-19 situation, no further meetings could be held in the last two years.

Further QA and verification activities to be continuously performed and/or planned:

- Several consistency checks as detailed in chapter 1.10.
- Active participation in the support project organized by EU DG Climate for the „*Assistance of Member States for effective implementation of the reporting requirements under the Kyoto Protocol to the United Nations Framework Convention on Climate Change (UNFCCC)*”

Other QA/QC activities

Besides ISO requirements, other QA/QC activities are carried out, as well. For every sector of the inventory, there is a responsible person within the core team in the Meteorological Service. These sectoral responsibilities are laid down in the QC record No. NELO04. Especially in case of external experts, this responsible member of our team conducts several quality checks on the provided calculations. Moreover, this exercise can be regarded as an interactive process throughout the whole inventory cycle, since the used methodologies, early results are discussed during the process of the emission/removal calculations. This QC procedure also led to a few recalculations. The used parameters and factors, the consistency of data is checked regularly. Completeness checks are undertaken, new and previous estimates are compared every time. Data entry into the database is checked many times by a second person. If possible, activity data from different data sources are compared and thus verified. In response to our request, several data suppliers made declarations as regards quality assurance systems in place during the collection of the data and QC record NELO02 has been introduced for the documentation of evaluation of data quality by data providers. Experts involved in emission forecast consulted in many areas with inventory experts of the Hungarian Meteorological Service to reach better consistency, which in turn represented some sort of QA procedure for the inventory itself.

Nevertheless, the work continues to refine the used QA/QC procedures and implement further elements. The QA/QC Plan is under review in order to implement all changes required by the EU Monitoring Mechanism Regulation and implementation of the 2006 IPCC Guidelines.

1.8 Uncertainty

The reliability of the data for individual source categories was estimated on the basis of the 2006 IPCC Guidelines but information from the industry and expert estimates was also used. On the basis of Table 3.3 and Table 4.1 of the 2006 IPCC Guidelines we have determined the total uncertainty according to the Tier 1 method. Accordingly, the combined uncertainty as % of total national emissions (in the year 2020) is 13.3% (excluding LULUCF) and the uncertainty introduced in trend in national emissions is 3.2%.

The uncertainty values have been determined by gas as well:

% Uncertainty excluding LULUCF	
CO ₂	2.6
CH ₄	48.4
N ₂ O	145.0
F-gases	13.0

Estimation of the uncertainties including LULUCF is a planned improvement. Please find the detailed Tables presenting the whole calculation in chapter 2 of Annex of the NIR.

1.9 Completeness

GHG inventory data are provided for the base year (the average of the three years 1985–1987) and the years 1986–2020. All relevant gases, sectors and categories are included. The inventory is complete in terms of geographic coverage. The notation keys are used throughout the tables.

1.10 Consistency checks

Regulation (EU) No 525/2013 of the European Parliament and of the Council of 21 May 2013 on a mechanism for monitoring and reporting greenhouse gas emissions and for reporting other information at national and Union level relevant to climate change requires to report the results of the checks performed on the consistency of the emissions reported or of the data used to estimate emissions in preparation of the greenhouse gas inventories. The results of the consistency checks are summarized below.

Consistency with the verified emissions reported under Directive 2003/87/EC

ETS data is essential in inventory compilation, especially to derive country specific emission factors for several categories in the inventory. In addition, comparisons have been made between emissions reported under Directive 2003/87/EC and emissions reported to the UNFCCC. ETS emissions represent about one third of total emissions. Extending this comparison to source categories, there is no source category where the ETS emissions are higher than the reported values in the inventory. This is partly due to our practice of activity data reallocation where precedence is given to ETS data over the original IEA energy statistics. Good consistency can be found in case of several source categories, especially for 1.A.1.a Public electricity and heat production, 1.A.1.b Petroleum refining, 1.A.2.a Iron and steel, 2.A.1 Cement Production, 2.A.2 Lime production, 2.A.4 Other process uses of carbonates, and 2.B Chemical industry. Information on consistency is provided in a separate Annex submitted to the EU.

Consistency of the data used to estimate emissions in preparation of the greenhouse gas inventories with the data used to prepare inventories of air pollutants under Directive 2001/81/EC

As basically the same team prepares both inventories, the consistency of the used data is safeguarded. Consistently, emission estimates of carbon monoxide (CO), sulphur dioxide (SO₂), nitrogen oxides (NO_x) and volatile organic compounds, in inventories submitted by the Member State under Directive 2001/81/EC of the European Parliament and of the Council and under the UNECE Convention on Long-

range Transboundary Air Pollution are consistent with the corresponding emission estimates in greenhouse gas inventories. The difference between the total emissions of any of the above pollutants reported in both inventories is well below 5% as it is demonstrated in the table below. (The relatively larger differences in CO emissions are mainly due to emissions reported in the LULUCF sector.)

CRF	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
NOX	245	189	187	178	146	127	119	121	119	114	107
CO	1434	964	850	695	535	469	445	445	373	363	343
NMVOC	306	210	188	172	130	126	126	123	117	117	112
SO2	829	613	427	43	30	24	23	28	23	17	16
NFR	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
NOX	246	191	189	179	148	128	120	121	120	115	107
CO	1451	982	857	697	552	464	450	439	377	358	339
NMVOC	307	210	189	173	130	126	127	124	118	118	112
SO2	829	613	427	43	30	24	23	28	23	17	16
DIFF											
NOX	-0.8%	-1.0%	-0.8%	-0.9%	-1.3%	-0.5%	-0.9%	-0.6%	-0.9%	-0.7%	-0.4%
CO	-1.2%	-1.8%	-0.8%	-0.4%	-3.1%	1.0%	-1.1%	1.3%	-0.9%	1.5%	1.2%
NMVOC	-0.3%	-0.4%	-0.4%	-0.4%	-0.5%	-0.4%	-0.3%	-0.3%	-0.4%	-1.0%	0.2%
SO2	0.0%	0.0%	0.0%	-0.2%	-0.3%	-0.3%	-0.3%	-0.2%	-0.3%	-0.4%	-0.2%

Consistency of the data used to estimate emissions in preparation of the greenhouse gas inventories with the data reported pursuant to Article 19 of Regulation (EC) No 517/2014

In the case of 517/2014/EC only companies importing and exporting across the EU border are required to report. However, thanks to the Hungarian F-gas regulations, data is available also on import/export of F-gases within the EU for the preparation of the inventory. Thus, the data used for the preparation of the inventory is significantly wider than the data reported based on 517/2014/EC.

Consistency of the data used to estimate emissions in preparation of the greenhouse gas inventories with the energy data reported pursuant to Article 4 of, and Annex B to, Regulation (EC) No 1099/2008

The IEA/Eurostat joint questionnaires serve as basis of emission calculation in the energy sector and as regards non-energy use of fuels partly also in the industrial processes sector. Consistency is further enhanced by our practice that emission calculation files access directly the joint questionnaires. Where ETS data are taken into account, there might, however, be some minor differences with the energy statistics (e.g., industrial waste consumption in cement production). Nevertheless, these differences are well below 2%.

2 TRENDS IN GREENHOUSE GAS EMISSIONS

In the United Nations Framework Convention on Climate Changes, Hungary undertook to keep its CO₂ emissions in 2000 at or below the 1990 level. In the first commitment period of the Kyoto Protocol, our country committed to reduce the average greenhouse gas emission by 6% of the base year level during the five years of the first commitment period (2008 to 2012). It will be shown in the next Sections that Hungary has complied with these commitments.

2.1 Description and interpretation of emission trends for aggregated greenhouse gas emissions

The trends of the total greenhouse gas emissions may be assessed on the basis of the GWP. The table below shows the time series of net and gross emissions:

Table 2.1 Total GHG emissions (including and excluding LULUCF)

	BY	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
Total (excl LULUCF)	110,521	94,821	77,194	74,929	76,714	66,021	61,496	62,256	64,729	64,756	64,581	62,818
Total (incl LULUCF)	108,160	91,722	71,160	74,080	70,842	61,605	55,839	57,760	59,610	60,334	59,674	55,997

BY=average of 1985-87

Compared to the base year, emissions were significantly reduced in the energy (-45%), industrial processes and product use (-49%), and agriculture (-39%) sectors. In contrast, emissions in the waste sector have increased since 1985 (+5%). The land use, land-use change and forestry (LULUCF) sector shows fluctuating behavior. Looking at the most recent trends since 2005, emissions have significantly decreased in the energy and industrial processes sectors by 23% and 15%, respectively. The agriculture sector seems to have recovered and could show an increase of 19% since 2005. The previous growing trend turned back in the waste sector (-19%).

For a better understanding of the Hungarian emission trends, the time interval of the inventory should be split into three periods with different emission relevant economic processes in the background. The first period (1985-1995) would be the years of the regime change in Hungary, whereas in the second period (1995-2005) the rules of the market economy became decisive. The second period can also be characterized by the decoupling of GDP growth from the GHG emission trend which is undoubtedly an important development. By 1999, the GDP reached the pre-1990 level; however, emission levels remained significantly below the levels of the preceding years. Thus, the emissions per GDP were decreasing.




Figure 2.1 Comparison of trends in GDP and GHG emissions

In the third period, after 2005, Hungary experienced an emission reduction of 24% up to 2013, out of which 6% occurred in the first three years up to 2008: basically, due to mild winters, higher energy prices, and modernization in the chemical industry. Then in 2009, the global financial-economic crisis made its radical influence felt which can also be seen at the dropping GDP values in *Fig. 2.1*. From 2010 on a slight recovery of the economy could be observed, the emissions, in contrast, not just remained at a relatively low level but decreased again quite significantly. However, the decreasing trend stopped in 2013 and an increase of 12% could be observed altogether up to 2017. After four years of increase, emissions have remained more or less at the same level between 2017 and 2019. In 2020, however, emissions fell by almost 3% to around 2016 levels, mainly due to a significant reduction in transport emissions as a consequence of COVID-19.

Starting with the first period, the process of transition into market economy brought in its train radical and painful decline in the output of the national economy. The production decreased in almost every economic sector including also the GHG relevant sectors (energy, industry and agriculture). Consequently, GHG emissions decreased substantially in these years by around 33 million tonnes CO₂ equivalent. Between the mid 80's and the mid 90's emissions fell back in the *energy* sector by around 26%, and even more, by around 45-50% in the *industrial processes* and *agriculture* sectors.

The most significant drop in energy use occurred in the industry especially in the energy-intensive industrial sectors (manufacture of basic metals and machinery, mining etc.). The industrial output of 1992 was two third of that of 1989. Several factories were closed down, capacity utilization was reduced, consequently the production decreased more or less drastically in each industrial sector.

Some examples:

- Iron and steel production: two out of three plants were provisionally closed down;
- Aluminum: two out of three plants were closed down in 1991 (aluminium production stopped in 2006 eventually);
- Ferroalloys: ceased to exist (1991);
- Ammonia: four out of five plants were closed down (1987, 1991, 1992 and 2002);

- Nitric acid: three out of four plants were closed down (1988, 1991 and 1995).

The agricultural sector suffered a similar decline. As a result of the political and economic processes, the number of agricultural farms was reduced by more than 30%, the number of employees by more than 50%, the volume index of the gross agricultural production by more than 30%, the livestock by about 50%, and the use of fertilizers by more than 60%. As a consequence, the share of the agricultural sector in total GHG emissions decreased from 11% to 8%.




Figure 2.2 Changes in emissions due to regime change, between base year and 1995, million tonnes CO₂-eq

The small increase of emissions in the *Waste* sector is exceptional among all the sectors, and it is attributable to the slightly increasing quantities of waste generated and collected but more importantly to the applied calculation method which assumes that the degradable organic component in waste decays slowly throughout a few decades.




Figure 2.3 Changes in emissions between 1995 and 2005, million tonnes CO₂-eq

After the mid 90's, emissions seemed to have stabilized around 77 million tonnes CO₂ equivalent. However, behind the quite stable emission level opposite processes could be observed which can be illustrated by the relatively bigger changes in the energy sector. The fuel use of industry decreased further which led to about 6% share only in GHG emissions around 2005. In contrast, emissions from transport increased significantly by almost 5 million tonnes CO₂ equivalent which represented a growth of 60%.

In the third period, between 2005 and 2020, emissions fell by 13.9 million tonnes or 18%. (The decrease was even higher, 24%, if we look at the period between 2005 and 2013.) About a quarter of this decrease occurred between 2005 and 2008. The decreasing energy use by other sectors and manufacturing industries, and the diminishing process related emissions in the chemical industry were the main drivers of these changes. Most importantly, total fuel consumption in the residential sector decreased by about 16% (including a 33% drop in solid fuel and a 16% decrease in natural gas use) – mainly due to extreme mild winter in 2007 but probably the growing energy prices and the support for modernization of buildings might have played a role as well. Decreased production volumes and modernization in the chemical industry led to an emission reduction of about 45%. In contrast, emissions from energy industries and transport grew further.

Then in 2009, the Hungarian economy was hit hard by the global economic crisis that exerted a significant effect on the emission level. Emissions (excluding LULUCF) decreased by 9% (-6.2 million tonnes) between 2008 and 2009. In comparison with 2008, emissions in 2009 were lower in all major sectors. The highest relative reduction (-14%) occurred in the industrial processes and product use sector mainly due to lower production volumes especially in mineral product manufacturing (-28%). Parallel to that, also energy use decreased in manufacturing industries and construction, consequently GHG emission also fell by 28% here. Regarding absolute changes in emissions, out of the 6.2 million tonnes reduction, fuel combustion was responsible for about 4.6 million tonnes. Although the energy demand increased in the heating season due to less favorable weather conditions, the fall in the production of energy intensive sectors led to an overall decline in energy use.




Figure 2.4 Changes in emissions between 2005 and 2020, million tonnes CO₂-eq

The decline in economic output stopped in the first quarter of 2010. Mainly driven by the growth in export-oriented industrial production, the GDP grew by 1% in 2010. The change in GHG emissions was about the same. In the next three years, however, emissions decreased again altogether by 12%.

In 2011, we could see decreases in many areas but especially in the energy sector. Electricity production decreased by 4% which resulted in a similar fall in GHG emissions. Natural gas consumption of the residential sector dropped by 9%. Transport emissions fell by 5%, mineral production by a further 15%. In this overall decreasing trend, agricultural soils were the main exceptions. In agriculture, we had higher fertilizer use, and greater crop production (hence higher emissions from crop residues). In this respect, it is worth noting that the economic growth in 2011 was mainly driven by agricultural production.

In 2012, the decreasing trend in emissions continued. The decrease of 3.4 million tonnes (or -5%) can almost be explained by processes in the energy sector alone (e.g., further decrease in electricity production, a 13% drop in natural gas consumption in “other sectors”) as it will be elaborated more in chapter 2.3.

2013 was not an exception in the decreasing trend, either. Total emissions have decreased by a further 5% corresponding to 2.9 megatons in CO₂-eq. The decrease was dominated again by the energy sector. Emissions from power and heat production alone dropped no less than 2.6 Mt CO₂-eq due to significantly lower electricity production from fossil fuels.

Total emissions did not change much in 2014. In the energy sector, we could observe some counterbalancing processes. After several years of decreasing emissions, the transport sector started to show some growth. Nevertheless, the diminishing fossil fuel based electricity production, and the lower and lower energy consumption in the residential sector led to an overall decrease of emissions.

After the lowest point in 2013, emissions started growing again. Up to 2017, the overall increase reached 12%. Between 2016 and 2017, the growth rate was 4%, to which all sectors contributed to a greater or lesser extent.

Between 2017 and 2019, total emissions seemed to have stabilized. However, this stabilization of total emissions was a net result of several balancing processes: (1) increased emissions especially in transport, and (2) decreases in domestic energy use, and in the power sector.




Figure 2.5 Changes in emissions between 2017 and 2019

Then in 2020, emissions decreased by close to 3%. The transport sector due to reduced commuting, tourism and business travel contributed particularly to the decline of emissions as demonstrated by Fig. 2.6 below. Current emissions are 18% below the 2005 level.




Figure 2.6 Changes in emissions between 2019 and 2020

2.2 Description and interpretation of emission trends by gas

The following table shows the emission data for each greenhouse gas (Gg CO₂ equivalent):

Table 2.2 Trends in emissions of greenhouse gases in Hungary (excluding LULUCF Gg CO₂-eq)

CO ₂	85,418	73,226	61,391	58,365	60,276	52,069	46,653	47,113	49,515	49,463	49,235	47,284
CH ₄	13,590	12,830	10,744	10,591	9,705	8,894	8,384	8,334	8,353	8,292	8,234	8,220
N ₂ O	11,135	8,377	4,750	5,405	5,608	3,714	4,515	4,781	4,781	4,848	4,850	5,013
HFCs	NO	0	36	204	754	1,250	1,821	1,895	1,964	2,055	2,159	2,189
PFCs	371	376	223	282	280	4	4	4	2	3	3	3
SF ₆	7	12	51	82	90	92	118	128	114	97	101	109
NF ₃	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
TOTAL	110,521	94,821	77,194	74,929	76,714	66,021	61,496	62,256	64,729	64,756	64,581	62,818

Base year (BY)=average of 1985-87





Figure 2.7 Shares of emissions of greenhouse gases in the base year (BY) and in 2020

The drop of CO₂ emissions during the early 1990's was attributable to the reduction of fuel uses in conjunction with the decline of the national output. From the second half of the 1990's, emissions showed stagnating or slightly decreasing tendencies reflecting the effects of restructuring following the economic growth. The changes in the fuel-mix resulted in reduction of the specific emission levels. Between 2005 and 2013, CO₂ emissions decreased by 28 per cent, at about the same rate as during the regime change around 1990. The drop of emission accelerated after 2008 mainly driven by the global economic crisis, and the reduced fossil fuel based electricity production. Between 2013 and 2019, however, emissions increased again by 13%. Electricity production increased, transport sector showed a steadily increasing trend, and fuel consumption increased in manufacturing industry. In 2020, an exceptional year, CO₂ emissions dropped by 4%. Currently, CO₂ emissions are lower by 22% compared to 2005.

As regards CH₄ emissions, agriculture, fugitive emissions, and waste management are the trend setting sectors. Most importantly, reductions in the livestock resulted in lower emissions. Besides, emissions from waste disposal grew until 2008, but started to decrease recently. This is the reason why the resultant trend was relatively stagnating until the first half of the last decade, and why it has been slowly decreasing since then.

Due to the above factors, also N₂O emissions decreased significantly in the beginning of the period. Later it showed a slightly rising trend, followed by another drop primarily reflecting the fluctuations in agricultural output and the modernization of nitric-acid production.




Note: BY=average of 1985-87 but 1995 for F-gases

Figure 2.8 Trend of emissions by gases, including LULUCF, Gg CO₂-eq

The use of HFC gases became more intensive in the second half of the 1990's in conjunction with the restriction of the use of chlorofluorocarbons as refrigerants. The rise of emissions is obvious.

PFCs emissions are principally related to aluminium production processes. Therefore, the tendencies of PFC emissions reflect the changes in aluminium production. Following a drastic reduction in the beginning of the period, the levels showed a slow but steady increase. Then the aluminium production ceased suddenly in 2006.

SF₆ emissions primarily depend on the uses in electricity transmission, as it is mainly used in electrical equipment, first of all in switchgears for insulation and arc quenching. So, the growth of the electricity consumption results in an increasing application of SF₆, however the tendencies vary according to the manufacturing/application needs and the steep increase seems to be stopped in the recent years in SF₆ emissions too.



Note: *BY=average of 1985-87 but 1995 for F-gases

Figure 2.9 F-gases trend (1985-2020), Gg CO₂-eq

2.3 Description and interpretation of emission trends by category

The following figure shows the emissions by sources and removals by sinks for each sector. The biggest emitting sector was the energy sector contributing 71% to the total GHG emission in 2020. Industrial processes and product use, and Agriculture had a similar share of 12% each. The waste sector contributed 5%. Compared to the base year, emissions significantly decreased in the energy (-45%), industrial processes and product use (-49%), and agriculture (-39%). In contrast, emissions in the waste sector have increased since 1985-87 (+5%). The land use, land-use change and forestry (LULUCF) sector shows fluctuating behavior. Looking at the most recent trends since 2005, emissions significantly dropped in the energy and industrial processes sectors by 23% and 15%, respectively. The agriculture sector seems to have recovered and could show an increase of 19% since 2005. The previous growing trend turned back in the waste sector (-19%).




Figure 2.10 Shares of sectors in 2020

Production and use of energy generate most greenhouse gases, largely CO₂. The **energy sector** was responsible for 71% of total GHG emissions in 2020. Emissions in the energy sector decreased in the first part of the period as a result of reduced energy consumption and a more favorable fuel mix. The significant reduction in emissions between the base year and 1995 was mainly due to the economic transformation which caused sudden decrease in energy demand. (In this respect, it is perhaps worth mentioning that the decrease in fuel consumption between 2005 and 2014 was even higher!) In addition, ongoing changes in fuel-structure, i.e., gradual replacement of solid fuel by natural gas, led to further decrease of total emissions. Between 2005 and 2008 growing emissions from energy industries and transport could be observed, which were more than offset by the drastic reductions in the residential sector and manufacturing industries. And then the economic crisis came: total combustible fuel consumption decreased by 6% in 2009. Fuel consumption decreased further until 2014 by 14%, but then it increased again by 11% until 2019. In 2020, total fuel consumption decreased by 2% and it is lower by 16% than in 2005.

Currently, 16% of domestic primary energy supply is nuclear, 12% is renewable which means that the remaining - overwhelming - part of primary energy demand has to be met by fossil fuels. Natural gas accounts for the largest share (47%) of incinerated, largely fossil fuels, followed by petroleum products (30%). Emissions have been positively influenced by the fact that the proportion of coal with higher specific emissions has fallen from 30% to 7% in the last 35 years, which is well below the current share of biomass in fuel consumption (14%).

The three most important sources of emissions in the energy sector are transport, energy industries, and “other sector” (mostly including residential and other buildings), each of which accounted for 20% of total national emissions in 2020. Energy use and emissions from manufacturing industries and construction contributed 8% to domestic total emissions. Fugitive (mostly methane) emissions from the domestic natural gas system represent 3% of total emissions

In recent years, the transport sector became the largest emitter, not only within the energy sector but also across all sectors, as transport accounted for 23% of total national emissions in 2019. However, as a result of preventative measures for COVID-19, emissions from the road transport-dominated sector fell sharply by 14% in 2020. Nevertheless, compared to the previous low point, 2013, transport emissions were still 25% higher. In addition, based on preliminary fuel sales data, it appears that the decline in 2020 can be considered temporary as rising emissions are expected for 2021 in this sector.

Considering energy industries, domestic electricity production increased a little by 2%. However, coal-based power production dropped by a further 9% (after a decrease of 13% last year) whereas natural gas fired power plants with more favorable specific emission levels increased their production by 4% (after an increase of 19% in 2019). Another welcome development is the sharp increase in the use of solar energy: 4% of gross electricity production now comes from solar energy. The share of nuclear power generation was at around 50% over the last few years. The relatively large share of electricity import decreased from 31% in 2014-2015 to 25% in 2020. As a result of all the above, emissions from energy industries decreased by 2% in 2020.

2020 qualified “only” as the eighth warmest year in the last 120 years, therefore the heating demand increased a little after the two warmest years. Compared to 2019, total fuel consumption in dwellings increased by 5%. While the use of natural gas increased by 8%, firewood consumption did not change significantly and the marginal coal consumption decreased further by 20%. (It is worth noting that the use of the two main energy sources often changes in opposite directions: since 2005, natural gas consumption has fallen by 25%, while firewood has increased by 33%.) Although household emissions

increased by 7%, emissions from the total energy sector were 4% lower in 2020 than in 2019 due to declining transport and industrial emissions.

The **industrial processes and product use sector** contributed 12% to total GHG emissions in 2020. The most important greenhouse gas was CO₂, contributing 66% to total sectoral GHG emissions, followed by F-gases with 28%. In 2020, 34% of the emissions came from chemical industry, followed by 28% from product uses as ODS substitutes. Mineral industry has 17%, metal industry has 14% contribution to sectoral GHG emissions, respectively. Other product uses (containing SF₆ and N₂O) and non-energy products from fuels and solvent use have the smallest influence on the 2020 IPPU inventory with 4% and 2%, respectively. Process related industrial emissions decreased by 49% between the base year and 2020, and by 15% between 2005 and 2020.

GHG emissions from the iron and steel producing sector decreased by 9% compared to 2019 because of the decreasing production of pig iron. Production of ammonia, urea and nitric acid increased in 2020, causing 8% increase in emissions from the chemical industry sector. In the production of mineral industry, the years-long increasing trend halted in 2020, GHG emissions from this sector were 11% lower in 2020 than in 2019. Emissions from the non-energy products from fuels and solvent use increased by 16% mainly due to the increase in the lubricant consumption.

A third of industrial emissions come from the operation of equipment containing F-gases and from use of F-gas containing products. Category 2.F.1. (Refrigeration and air-conditioning) accounts for 87% of total F-gas emissions, which is still increasing. Despite the continuous regulation of high GWP gases emission has not already stopped in this sector.

Although, charging into new equipment for some gases already is forbidden, a lot of cooling systems have been operated with gases which have higher GWP and this is the reason of this trend.

In 2020, the **agriculture sector** accounted for 12% of total emissions. Emissions from agriculture include CH₄ and N₂O gases. 86 per cent of total N₂O emissions were generated in agriculture in 2020. Emissions from agriculture have decreased by 39% over the period of 1985-2020. The bulk of this reduction occurred in the years between 1985 and 1995, when agricultural production fell by more than 30 per cent, and livestock numbers underwent a drastic decline.

Between 1996 and 2008, agricultural emissions had stagnated around 6.2 Mt with fluctuations up to 4.6%. Behind this trend there were compensatory processes. While the number of livestock decreased further leading to lower emission, the use of fertilizers increased by 68% in the period 1995-2007 which caused growing nitrous oxide emissions from agricultural soils. In 2008 the significantly rising fertilizer prices led to lower fertilizer use, which resulted in some reduction in the emission levels.

Agricultural emissions decreased both in 2009 and 2010. A major reduction in emissions occurred in 2009, when 11 per cent decline in swine population also contributed to the downward trend. Agricultural emissions, after hitting the lowest point in 2010, had increased until 2018, mainly because of the increase in the inorganic fertilizer use, cattle livestock, and milk production per cow.

The GHG-emissions reflect the restructuring in the agricultural production has taken place since 2004, namely the increased ratio of crop to livestock production. Share of CH₄ emissions, which derive mainly from the animal husbandry, has decreased, while the N₂O emissions, originating primarily from the crop production has grown, since 2004.

Certain types of inorganic fertilizers as urea containing fertilizers and calcium ammonium nitrate (CAN) fertilizers contribute to the agricultural GHG-emissions not only with their nitrogen, but also their carbon content. In Hungary CAN fertilizers have become increasingly popular in the recent years, as a result N₂O and CO₂ emissions has tripled from this source since 2005.

In 2019, emissions growth temporarily slowed down, mainly due to the decreasing swine livestock and synthetic fertilizer use, but emissions increased again in 2020. The upward trend was mainly due to an increase in fertilizer use and beef cattle numbers.

The **Land Use Land-Use Change and Forestry sector** has been a net carbon sink mainly because of the huge amount of carbon uptake of forests, which in turn is due to continuous afforestation efforts and sustainable forest management. The complex dynamics of the land use and land-use changes leads to highly fluctuating estimates of sectoral removals. Over the period 1990 to 2020 our estimates indicate an average annual net removals of 4 million tonnes CO₂-eq ranging from 0.8 million tonnes in 2000 to 6.8 million tonnes CO₂ in 2020. In 2020, the net removals of forests amounted to 6.6 million tonnes CO₂ the main reason of which being the lower harvest due to the COVID pandemic. The harvested wood product pool is close to a carbon equilibrium with a small net sink in the last six years. The non-forestry land-use sectors used to be small net sinks before 2016 but they have been small sources since then.

The **waste sector** was responsible for 5% of total national GHG emissions in 2020. The largest category was solid waste disposal on land, representing 85% in 2020, followed by wastewater treatment and discharge (9%), biological treatment of solid waste (5%), and incineration of waste without energy recovery (1%). In contrast with other sectors, emissions from the waste sector are by 6% higher now than in the base year. However, the growth in emissions stopped in the last decade, and a reduction of 19% could be observed between 2005 and 2020. The degradation process in solid waste disposal sites is quite slow which means that waste that were disposed many years earlier have still an influence on current emission levels. However, the amount of disposed waste had dropped significantly since 2005 (e.g. landfilled municipal waste decreased by 50%) consequently methane emissions started to decrease as well. GHG emissions from wastewater handling have a pronounced decreasing trend due to a growing number of dwellings connected to the public sewerage network.




Figure 2.11 Sinks of LULUCF, Gg CO₂-eq

2.4 Trends of indirect gases and SO₂

[This section will be updated in the March submission]

NO_x, CO and NMVOC gases are referred to as indirect gases because they (together with SO₂) influence atmospheric warming indirectly, via secondary effects. Nitrogen oxides, carbon monoxide and (non-methane) volatile organic compounds are precursor of ozone which is itself a naturally occurring greenhouse gas. Sulphur dioxide can contribute to formation of aerosols that scatter some of the solar radiation back into space. Calculation of the emissions of these gases is required by the UNFCCC reporting guidelines. It should be noted that Hungary (as well as the other European countries) has calculated the emissions of such gases for several decades and the Geneva Convention of 1979 (CLRTAP) also laid down such obligations. Emissions are reported consistently in the above two reporting regimes.

The following table shows the main trends in emissions:

Table 2.3 Trends in emissions of indirect greenhouse gases and SO₂, including LULUCF (Gg)

GASES	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
NOX	245	189	187	178	146	127	119	121	119	114	107
CO	1434	964	850	695	535	469	445	445	373	363	343
NMVOC	306	210	188	172	130	126	126	123	117	117	112
SO2	829	613	427	43	30	24	23	28	23	17	16

The substantial reduction in sulphur dioxide emissions (-95%) is attributable to the decreased use of fossil fuels in general and the decreasing share of coal with higher sulphur content. After 2000, further reductions were observed due to the introduction of SO₂ precipitators in coal-fired power stations. Reduced carbon monoxide emissions are obviously a consequence of decreased fuel uses. The decrease in NO_x emissions is relatively moderate due to the increasing significance of transport.

3 ENERGY (CRF sector 1)

Recent key developments:

- Between 2005 and 2014, Hungary experienced an almost constant emission reduction in the energy sector. However, after a 27% reduction, the decreasing trend stopped, and total emissions increased by 12% between 2014 and 2017. Emissions remained at a similar level in 2018 and 2019, and then – mostly due to COVID-19 – decreased by 4% in 2020. Current GHG emissions are by 23 per cent lower than in 2005.
- The three most important sources of emissions in the energy sector are transport, energy industries, and “other sector” (mostly including residential and other buildings), each of which accounted for 20% of total national emissions in 2020.
- In recent years, the transport sector became the largest emitter, not only within the energy sector but also across all sectors, as transport accounted for 23% of total national emissions in 2019. However, as a result of preventative measures for COVID-19, emissions from the road transport-dominated sector fell sharply by 14% in 2020.
- The share of the energy industries in total emissions decreased to 20% in 2020 from 26% in 2005. Although domestic electricity production increased a little by 2%, coal-based power production dropped by a further 9% (after a decrease of 13% last year) whereas natural gas fired power plants with more favorable specific emission levels increased their production by 4% (after an increase of 19% in 2019). In energy and heat production, the use of conventional fossil fuels (coal, natural gas) has declined, while that of renewable and waste has increased. As a result of all the above, emissions from the energy industry decreased by 2% in 2020.
- After record warm years, the higher heating demand in 2020 was met mostly by increasing natural gas consumption leading to higher emissions in the residential sector.

Major changes compared to previous submission:

- The latest version of the Annual IEA/Eurostat Questionnaires submitted in November 2021 were used as activity data. All the changes in the energy statistics are reflected in the current inventory.
- -Emissions from all autoproducer plants have been (re-)allocated from the source category 1A2g to the relevant economic sector where they operate fully for the period 1998-2020, and to the extent possible also for earlier years.
- We have switched to the COPERT model (version 5.5.1, September 2021) for the whole time series.
- Activity data have been revised for groundwater extraction (1B2d), and the same country-specific emission factor has been applied for the whole time series.
- Another reallocation: to increase comparability, leakage and venting emissions from gas transmission and storage are reported separately on the basis of TABLE 4A.2.7 of the 2019 Refinement.

3.1 Overview of sector

Emitted gases: CO₂, CH₄, N₂O

Methods: T1, T2, T3

Emission factors: D, CS, PS

This sector covers emissions from combustion processes and fuel-related fugitive emissions from exploration, transmission, distribution and conversion of primary energy sources.

For a better understanding of the principal drivers behind fossil fuel related emission trends and variations, the main characteristics of the Hungarian Energy System will be described shortly in the following. First of all, not enough, cheap and clean domestic energy resources of good quality are available in Hungary, therefore the energy demand has to be met by import to a great extent. In 2020, primary energy production amounted to 451.7 PJ which was by 27 per cent less than in 1990 and more or less at the same level as in 2005 (-1%). Most importantly, uneconomical deep coal mines were closed down, but also crude oil and natural gas production decreased. In contrast, imports of energy with 946.3 PJ in 2020 were by 42% and by 4% larger than in 1990 and 2005, respectively. As the share of production in consumption is around 40%, our import dependency is quite significant. Domestic supply of primary energy was 1099.2 PJ in 2020 which was by 9% and 7% below the levels in 1990 and 2005, respectively. Both primary energy production and supply decreased by around 2% in 2020. Final consumption decreased also: from 842.2 PJ in 2019 to 832.4 PJ in 2020. Looking at the main sectors, transport was the most affected by the COVID pandemic as energy consumption dropped by 12% compared to the previous year. The energy consumption of industry and the tertiary sector decreased as well, while that of the population increased due to higher heating demand and the coronavirus epidemic.

(Data source: The Hungarian Energy and Public Utility Regulatory Authority (HEA): see http://mekh.hu/download/5/1e/01000/7_2_annual_national_energy_balance_2014_2020.xlsx and HCSO: see: https://www.ksh.hu/stadat_files/ene/en/ene0002.html)




Figure 3.1.1 Primary energy balance of Hungary (1990-2020)

In 2020, final domestic electricity use amounted to 41,083 GWh, which was slightly lower than in the previous year. However, electricity consumption increased by 14% since 2014. The market penetration of the nuclear electricity started in 1983 in Hungary when the first 440 MW block of the Nuclear Power Plant in Paks was put into service. In recent years, 46% (2020) to 53% (2014) of the domestic generated electricity was produced by nuclear energy whereas the share of fossil fuels decreased to 40% in 2013 and remained below that level afterwards. According to the official statistics of the Hungarian Energy and Public Utility Regulatory Authority, the share of electricity from renewable sources in gross final consumption of electricity increased from 4.4% in 2005 to 11.9% in 2020. The last few years saw significant increases in solar electricity production (from 1 GWh in 2011 to 2,459 GWh in 2020) and also wind power production increased to 655 GWh in 2020 from 10 GWh in 2005. At the same time, electricity produced from combustible fuels decreased from 21,710 GWh in 2005 to 15,358 GWh in 2020.

By far, the biggest emitting sector is the energy sector contributing 71% to the total GHG emission in 2020. *Figure 3.1.2* shows the emission trends in the sector compared to the total.




Figure 3.1.2 GHG emissions of the Energy sector compared to total (BY-2020)

The significant reduction in emissions between the base year and 1995 was mainly due to the economic transformation which caused sudden decrease in energy demand. (In this respect, it is perhaps worth mentioning that the decrease in fuel consumption after 2005 was even higher!) In addition, ongoing changes in fuel-structure, i.e., gradual replacement of solid fuel by natural gas, led to further decrease of total emissions. Some classical types of fossil fuels have disappeared or their use decreased significantly, e.g., city-gas, heavy fuel oil (by destructive technologies it has been transformed to motor fuels and partly petrol-coke is produced from it). At the same time, the market penetration of new fuel types became significant e.g., petrol-coke, bio-ethanol, LPG and compressed natural-gas (CNG) for cars and buses, biomass for firing in power plants, biogas produced by fermentation of sludge and animal carcasses etc. All these changes were taken into consideration in our emission calculations.

Between 2005 and 2014, Hungary experienced an almost constant emission reduction in the energy sector basically due to mild winters and higher energy prices. In 2009 also the global economic crisis

affected the emissions especially in the energy and manufacturing industries sectors. Then in 2010, the growth in industrial production led to somewhat increased emissions again. In the next four years, however, emissions from the energy sector decreased further and reached their lowest level in the whole time series in 2014. Altogether, emissions decreased by 27% between 2005 and 2014.

However, the decreasing trend stopped in 2015. Total emissions from the energy sector have increased by 12% or 5.1 million tonnes between 2014 and 2017 and then remained more or less at the same level in 2018 and 2019. Above all, transport related emissions started to increase again already in 2014, and they grew by 46% in the last six years (2013-19).

The residential sector produced also higher and higher emissions between 2014 and 2017, after many years of dominantly decreasing trend. Then, mostly due to the fact that 2018 and 2019 were the warmest years ever, total fuel consumption of households decreased by 13% between 2017 and 2019. While biomass use decreased by 22%, fossil natural gas consumption by only 6%.

And then in 2020, the COVID-19 pandemic caused a significant drop in transport emissions and somewhat higher energy demand in the residential sector as described above.

Carbon dioxide from fossil fuels was the largest item among greenhouse gas emissions contributing 94% to the sectoral emission. Among all sectors, the energy sector contributes the most to the total CO₂ emissions as well (89% in 2020).

As regards methane emission, its contribution is 5% and 3% to the energy sector's emissions and to the total greenhouse gas emission (without LULUCF), respectively. Primarily, this results from fugitive emissions associated with conventional oil and gas production and processing (which also includes fugitive emissions from natural gas transmission). Among methane emitters, this sector's proportion is 26% (waste and agriculture sectors dominate here, see Fig. 3.1.2).

Considering nitrous oxide emission, this sector represents 1% (without LULUCF) of the total greenhouse gas emission. Among nitrous oxide emitters, its proportion is 6% which represents the second highest contribution compared to other sectors but it is still far behind agriculture.




Figure 3.1.2 Sectoral contributions to the total emissions of the main GHG gases (2020)

The three most important sources of emissions in the energy sector are transport, energy industries, and "other sector" (mostly including residential and other buildings), each of which accounted for 20%

of total national emissions in 2020, see *Fig. 3.1.3*. In the time series there were changes in the relative contributions of the different subsectors within the energy sector, most notably the growing share of transport emissions (from 11% in the base year to 32% in 2019, dropping temporarily to 28% in 2020) and the diminishing share of manufacturing industries (from 20% in the base year to 6% in 2009 and 11% in 2020).




Figure 3.1.3 Share of the different source categories in 2020

Fugitive emissions from fuels played only a small role with 4% out of which 97% originate from oil and natural gas production, processing, transmission and distribution. Emission in subsector 1.B.1 – Fugitive emissions from solid fuels are 96% smaller than the base year caused by the huge recession of underground coal mining in Hungary. The aggregate change of sector 1B – Fugitive emissions is 59% decrease compared to the base year.




Figure 3.1.4 Changing shares of the different subsectors (BY-2020)

3.2 Fuel combustion (CRF 1.A)

The principal driver of emissions in the energy sector is fuel consumption, therefore emissions of the sector strongly depend on the amount of combusted fuel. The use of combustible fuels decreased quite considerably, by 27% between the base year and 2020. Two periods need to be emphasized in this respect. The regime change around 1990 had the first significant effect: the fuel use in 1995 was by a fifth less than in the base year. The decrease in energy use after 2005 was even more significant (-22% until 2014) where the global economic crisis must have played a role. Then, fuel use increased again by 12% between 2014 and 2017, and decreased slightly by 1% in the next two years, and by 2% in 2020.




Figure 3.2.1 Fuel consumption by main fuel types (1985-2020)

Beside the amount, also the type of the used fuels has a great influence on the emission levels. Considering fuel use in combustion processes, gases had the highest proportion (47%) in 2020, liquids and solids represented 30% and 7%, respectively. It is worth mentioning that the share of biomass in fuel combustion grew to 14%. Especially solid fuels lost their importance: their share in the fuel mix was around 30% in the base year (see Fig. 3.2.2)




Figure 3.2.2 The used fuel mix in the base year and in 2020

Calculation of greenhouse gas emissions from combustion is based on the amount of fuel used. For this purpose, the energy balance of Hungary, the fuel balance for each fuel type and the fuel consumption for each subsector compiled by the Hungarian Energy and Public Utility Regulatory Authority are used dominantly. After discussion with the energy statistics provider and following their recommendation, it was decided that, starting with the 2014 submission, the basis of the inventories would be on the IEA/Eurostat Questionnaires. (Previously, these energy statistics were available to the inventory compilers basically as hard copies of the publication series Energy Statistical Yearbook. For some years, also electronic versions (tables in Excel files) were provided. However, this publication ceased, the last yearbook contained data for 2010.)

To increase consistency of the time series, we had to make some minor amendments of the allocation of fuel consumption compared to the IEA annual questionnaires, as follows:

- Based on 2011-2017 data allocations and value-added volumes of industrial production for previous years, some gasoil consumption has been reallocated from road transport to non-road mobile machinery (1A2gvii);
- The time series of gasoil use in navigation has been improved by interpolation where the missing amounts were taken again from road transport;
- Some natural gas consumption has been reallocated between other energy industries (1A1c) and commercial/institutional (1A4a) to reflect fuel consumption in oil and gas extraction. Data on natural gas production served as basis of extrapolation here;

It has to be noted that the traditional Hungarian coal terminology as published in the Energy Statistical Yearbooks differs from that of the IPCC. The partitioning was created according to the age of coal; Table 3.2.1 shows the classification according to the Hungarian and IPCC categories. Practically this means that imported “brown coal” in the Hungarian terminology would classify as sub-bituminous coal whereas domestically produced brown coal falls under the IPCC category of lignite. Basically, most of the coal produced in Hungary can be classified as lignite. Furthermore, the Energy Statistical Yearbook dealt with anthracite, hard coal, brown coal and lignite in the fuel balance separately, while the sectoral energy consumption for coal was the aggregate of hard coal, brown coal, lignite, gas coal and coking coal. Now, as our reporting is based on the IEA statistics, the reported fuel data follow the IPCC categories consistently.

As regards carbon emissions, solid fuels caused the most problems mainly because the fuel classification had been changed. The formerly used country specific carbon emission factors were previously determined for the Hungarian categories, namely for hard coal, brown coal, and lignite. (Until the 2014 submission, the following constant values had been used based on the 2005 ETS data: 27 tC/TJ for hard coal and brown coal and 30.9 tC/TJ for lignite.) Now, new factors had to be applied for other bituminous coal, sub-bituminous coal and lignite. Most of the coal produced in Hungary can be classified as lignite in this new system irrespectively whether it stemmed from surface or underground mines, although they have different characteristics. To take into account the changing share of the higher quality lignite from underground production, a time dependent carbon emission factor (changing between 103.0 and 108.6 t CO₂/TJ) was introduced and applied for the pre-ETS years. For the lowest quality lignite from surface mines the following parameters are used: EF=112.2 t CO₂/TJ, OX=0.974. As for Hungarian brown coal EF=100.8 t CO₂/TJ and OX=0.952 is applied. It is worth noting that the share of the lower quality lignite in production increased from 20% in 1990 to 84% in 2010. For other bituminous coal and sub-bituminous coal, the IPCC default values are used.

Table 3.2.1 Comparison of Hungarian and IPCC coal terminology

Hungarian Terminology	Net Calorific Values	IPCC Category (Gross calorific value)
Hard Coal	17-33 MJ/kg	Other Bituminous Coal (>23.865 MJ/kg)
Hard Coal	17-33 MJ/kg	Sub-Bituminous Coal (17.435 MJ/kg - 23.865 MJ/kg)
Brown Coal	10-17 MJ/kg	Lignite (<17.435 MJ/kg)
Lignite (young brown coal)	3.5-10 MJ/kg	Lignite (<17.435 MJ/kg)
Gas Coal and Coking Coal		Coking Coal

(Source: Bihari, 1998; IPCC, 2006)

Fuel use and emissions of autoproducer plants (that generate electricity or heat, wholly or partly for their own use as an activity which supports their primary activity) are accounted for fully in under the relevant economic sector in the period 1998-2020 as required by the guidebook, and to the extent possible also for previous years. (In the previous submission, almost all autoproduction was allocated to the source category “other stationary combustion 1A2gviii” for all years before 2013 with a few exceptions (e.g., coke oven gas and blast furnace gas were also previously reallocated from autoproducers to iron and steel, and to manufacture of solid fuels).

The problem of the network losses in the natural-gas transmission and distribution system should be also mentioned here. These losses are partly not technical ones in the reality, but the result of accounting, e.g., due to issues as measurement accuracy, temperature or pressure conversion or theft. (For more details see description in Ch. 3.3.2.5) The point is that only about half of the losses reported in statistical publications as distribution losses was taken into consideration as real loss (i.e., that is emitted into the atmosphere as methane), while the remaining half was assumed to be fired. Thus, the natural gas consumption in the residential sector is not the same as reported in the IEA natural gas annual questionnaire because 50 per cent of the network losses on average are added to it.

Input data for the fugitive emission calculation came from the Statistical yearbook of Hungary, Energy Statistics, the Hungarian Oil and Gas Company Plc. (MOL), the Hungarian Office for Mining and from the Hungarian Energy Office.

LPG and petroleum coke were taken into account as liquid fuels which had significant influence on the IEF value of this fuel type.

3.2.1 Comparison of the sectoral approach with the reference approach

The quantity of CO₂ from energy consumption was determined both on national level (reference approach) and on sectoral level (sectoral approach). The reference approach (RA) is based on national energy balance: production, import, export, stock changes, and international bunkers. The sectoral approach (SA) allocates the emissions by source category and includes only the combusted amount of fuels. The reference approach was compared with the sectoral approach as a check of combustion-related emissions. The check was performed for all years from 1985 to 2019 and is an integral part of reporting to the UNFCCC. The analysis includes also the comparison from the base year (1985-87). The reference approach, in theory, includes all CO₂ emissions from all fossil fuel uses in a country and should be compared with a set of emissions from the sectoral approach that includes all CO₂ emissions from energy use of fossil fuels.

Emissions from feedstocks and non-energy use of fuels are taken into account in the IPPU sector (2B and 2G) in case of sectoral approach (SA), therefore the energy and carbon content of these fuels are

removed from the RA (the fraction of carbon stored is 1 for all these fuels in the 1D sector), too. Similarly, emissions from coke used for transformation in the iron and steel industry were allocated to the relevant source category of the industrial processes sector, thus removed from the reference approach.

Since the 2015 submission, more fuel has been removed from the reference approach than in previous submissions (see also Table 3.2.3):

- All coke related emissions (including blast furnace gas) in the iron and steel industry are allocated to the IPPU sector;
- More natural gas consumption in the chemical sector is allocated to the IPPU sector.

In the CRF reporting software, the RA is directly compared with the sectoral fuel combustion total. This direct comparison shows that the total fuel consumptions of the RA are mostly larger than the SA totals (*Fig. 3.2.3*) on average by 0.8%. The remaining differences – after extracting the feedstock and non-energy use of fuels – are basically statistical differences, fugitive emissions and transformation losses which are occurring during coking, briquetting or oil refining.




Figure 3.2.3 Differences between the reference and the sectoral approach as regards fuel consumption and CO₂ emissions

In 2020, the difference between the two approaches was 0.2% in energy consumption and 0.0% in CO₂ emission (*Fig. 3.2.3*). The ranges of differences are between -2.1% (2014) and 2.6% (2006) with a 0.7% mean value as regards fuel consumptions, and similarly -2.3% (2014) and 2.5% (1989) with a 0.4% mean value as regards CO₂ emissions.

3.2.2 International bunker fuels

In accordance with the reporting guidelines, emissions from international aviation were included under the category International Bunkers on the basis of the quantities of kerosene used. In the time-series of the resulting CO₂ emission, significant jumps are present at certain places, which are obviously due to the changes in kerosene consumption. Naturally, changes in kerosene consumption

reflect the travelling/transport needs. This is clearly illustrated in Figure 3.2.4 which shows the air travelling/transport performance of the past years

Consumption in international navigation was not considered, because separate data on the uses for international navigation are not included in the national statistics.

International navigation depends not only on geographical and economic but on political conditions, too. International conflicts, wars have significant impact on international navigation, which could be seen in Hungary during and after the war in Yugoslavia. The war set back the navigation on the Danube South to Hungary, and decreased also the trade in Hungary. In the last years, the sea navigation (there was only tramp navigation) has relapsed due to falling into disuse of ship-fleet. This process could be traced back to the absence of Hungarian harbor on seas and Danube-sea ships. Between 1990 and 2000 the role of transportation of goods on waterways decreased from 28.2% to 2.9% among goods transportation in other ways. (Source: webpage of Központi Közlekedési Felügyelet)




Figure 3.2.4 Air travelling and transport performance in Hungary since 2000 in selected years

3.2.3 Feedstocks and non-energy use of fuels

The 2006 IPCC Guidelines introduced significant changes regarding feedstocks and non-energy use of fuels. It is good practice now to report all the feedstock and non-energy use of fuels in the *IPPU Sector* within the source category in which the process occurs.

In addition, also chapter 1.2 of Volume 2 states: “Combustion emissions from fuels obtained directly or indirectly from the feedstock for an IPPU process will normally be allocated to the part of the source category in which the process occurs. These source categories are normally 2B and 2C.”

So, in present submission all the fuels regarded as NEU in IEA Energy Statistics are allocated into IPPU sectors and also some amount from the quantities regarded as energy use in order to follow the suggestion of IPCC2006. This is the case for natural gas use in sector 2B1 Ammonia, naphtha use in 2.B.8 Petrochemical, and coke used in 2C1 Iron and steel.

Therefore, the fuel quantities for NEU reported in CRF Table 1.A.(d) and QA/QC check Table for NEU included in Annex of the NIR are higher than the actual quantity reported in IEA Energy Statistics. Nevertheless, the differences are well-known and documented.

Table 3.2.3 Allocation of feedstocks and non-energy use of fuels

Fuel type	Allocated under IPCC sector
Other kerosene	2.B.8 -Petrochemical and Carbon Black Production
Gas/diesel oil	2.B.8 -Petrochemical and Carbon Black Production
Liquefied petroleum gases (LPG)	2.B.8 -Petrochemical and Carbon Black Production
Naphtha	2.B.8 -Petrochemical and Carbon Black Production
Bitumen	2.D Non-energy Products - Other (<i>no CO₂</i>)
Lubricants	2.D.1 - Lubricant Use
Other oil	2.D.2 - Paraffin Wax Use 2.B.8 - Petrochemical and Carbon Black Production
Coking coal	2.C.1 -Iron and Steel Production
Coke oven/gas coke	2.C.1 -Iron and Steel Production
Natural gas	2.B.1 -Ammonia Production 2.C.1 - Iron and Steel Production 2.B.8 - Petrochemical and Carbon Black Production

Carbon content of all fuels which are allocated under the Industrial Processes sector is taken as stored carbon in the 1.AD sector (and in the *reference approach*), however the calculation of emission in the IPPU sector is not based on a default carbon-stored approach, but usually plant-specific (EU ETS) data, except for Lubricant and Paraffin wax use source categories.

3.2.4 Country-specific issues: on the use of plant level EU-ETS data

It is important to note first that generally emission data are not taken directly from the ETS database and put into the CRF as they are without analysis. Instead, facility level activity data (fuel use) and carbon emission factors are used from the ETS database to calculate weighted averages of the emission factors for different fuel types. These derived country specific EFs are then applied with the fuel use from the national energy statistics. The time series of these country specific emission factors and their comparison with the default values are summarized in Table 3.2.4. Fuel uses in energy statistics and ETS are compared also to see whether the fuel use in a given category is fully covered by ETS plants or not. Fuel consumption data are compared both in natural units and in energy units to reveal any possible differences in net calorific values. Should such difference occur, emission factors need to be amended to achieve consistency in energy balance and verified emissions since national energy data serve always as activity data. It is also checked whether the oxidation factor used by the facilities is included in their EFs. Measured oxidation factors, especially in case of coal firing plants, are always taken into account.

Generally, country specific emission factors derived from the ETS data are used for lignite, blast furnace gas, mix of coal and petroleum coke used in cement industry, other oil.

Starting with the previous submission, new country-specific CO₂ emission factors were introduced also for natural gas for the entire time series. In practice, the weighted averages of EFs of all ETS installations reporting EF with Tier 3 and Tier 2b were calculated for all years separately in the period 2008-2019. The yearly averages changed between 55.2 t/TJ (2008) and 56.3 t/TJ (2015). For all years before 2008, the average of 2008-12 is used (55.6 t/TJ)

Other country-specific issues are included under the source category descriptions and methodological chapter of each category.

3.2.5 Energy Industries (CRF sector 1A1)

Emitted gases: CO₂, CH₄, N₂O

Methods: T1, T2, T3

Emission factors: D, CS, PS

Key sources:

1A1 Fuel combustion - Energy Industries - Liquid Fuels – CO₂ – L, T

1A1 Fuel combustion - Energy Industries - Solid Fuels – CO₂ – L, T

1A1 Fuel combustion - Energy Industries - Gaseous Fuels – CO₂ – T

1A1 Fuel combustion - Energy Industries - Other Fossil Fuels – CO₂ – L, T

3.2.5.1 Category description

This subsector includes facilities generating electricity, district heating stations, oil refineries and coking and briquetting plants. On an overall level, here are the largest energy consumers.




Figure 3.2.5.1 Trends of CO₂, CH₄ and N₂O emissions in the Energy Industries (1985-2020)

Public Electricity and Heat Production was responsible for about 83-85% of fuel use in energy industries. According to a publication of the Hungarian Energy and Public Utility Regulatory Authority (“Data of the Hungarian Electricity System 2020”), “The energy consumption of the electricity generation in 2020 was 342530 TJ, 1.33% less than in the previous year. In 2020, 51.74% of the used energy sources consisted of nuclear fuel. Natural gas made up 21.93%, while coal made up 11.81% of the energy sources. The renewable energy sources provide 11.74% of the total energy source consumption.”

Domestic electricity production showed an overall increasing trend up till 2008; even during the years of the regime change around 1990, whereas import suffered a more severe drop from 28% to 6-7%. In addition to the effects of the financial crisis, an interesting incident occurred in 2009 when domestic production fell back by more than 10% whereas consumption decreased only by 6%. There was a multi-week break in the natural gas supply through Ukraine, thus the electricity generation of our natural gas firing power plants had to be substituted by import electricity and by increased production of the oil-fired power plants. After 2010, until 2014, domestic electricity production decreased every year,

and it has dropped quite substantially in 2013 by 13%. In the last six years (2014-2020), however, domestic production grew again altogether by 19%. The share of import is a highly variable figure: in the previous decade, it changed between 8% (2001) and 18% (2004). After 2010, however, it grew constantly and has reached a share of 31% in 2014 and remained close to 30% afterwards but decreased to 25% in 2020.




Figure 3.2.5.2 Domestic Electricity Production and Net Import (1980-2020)

Naturally, as domestic emissions are related to domestic production, the yearly fluctuation of production is one of the decisive factors. Not less important is the way how electricity is produced, e.g., what energy source is used. In Hungary, this sector consumes the deterministic part of our solid fossil fuel production. However, some uneconomical coal-fired power plants of low efficiency were stopped, and blocks of combined-cycle-gas turbine units were installed. For example, new 150 MW combined cycle gas-turbine units were installed (Újpest, Kelenföld, Százhalombatta, Nyíregyháza Power Plants), and aged coal fired units (Inota, Bánhida) of low efficiencies were taken out of service or blocks have been converted to the combustion of biomass (Pécs, Kazincbarcika, Ajka Power Plants). The demand for fossil fuel decreased about 150 PJ in the electricity sector between 1980 and 1990 because of the penetration of the nuclear electricity into the electricity market. This means that the fossil fuel consumption of public power plants is smaller now than it was before the introduction of nuclear electricity generation, in spite of much higher domestic electricity production. As a promising new development, increasing use of renewable sources could be observed by some public power plants. These developments are demonstrated in Figure 3.2.5.3.




Figure 3.2.5.3 Share of produced electricity by fuel (1995-2020)

In 2011 there were considerable changes in several areas of the Hungarian Power System. On the generation side, AES Borsodi Energetikai Kft. (AES Borsod Heat PP Ltd), being under liquidation, ceased its electricity generation. This meant that two coal and partly biomass firing power plants were closed. However, new units were added to the system: the combined cycle power plant of E.On Erőmű Kft. (E.On Power Plant Ltd.) in Gönyű and the open cycle gas turbine power plant of BVMT Bakonyi Villamos Művek Termelő Zrt. (BVMT Bakony Power Generation Ltd.). In addition, the amendment of the operating licence of Dunamenti Erőmű Zrt. (Dunamenti Power Plant Ltd.) enabled the commercial operation of a GT3 unit.

“Since the regional supply and demand factors affect the electricity market, the utilisation of domestic power plants is strongly influenced by the fuel costs and the regional wholesale electricity prices changing country by country. The gas-fired power plants have lost significant market share also in our region due to the high and basically oil price-indexed gas prices, the drop in electricity consumption, the collapse of CO2 allowance price system and the increase of electricity generation from renewables. Consequently, the load factor of domestic power plants was low. The traders compensated the loss of domestic generation from import. Thus, the amount of import-export balance reached 18.8% of total domestic electricity consumption in 2012.”

(Source: STATISTICAL DATA OF THE HUNGARIAN POWER SYSTEM, 2012)

The above words taken from a previous edition from the already referenced Statistical Data of the Hungarian Power System 2012 seem to be valid also for recent years. There were no further large power generating units connected to the Hungarian Electricity System in the last few years. However, there was a significant change in the installed capacity of Hungarian electricity system in 2019 with an increase of almost 560 MW, which is mainly due to the new solar power plants with an installed capacity of more than 50 kW added to the system.

3.2.5.2 Methodological issues

Activity data

Energy consumption data were taken from the IEA annual questionnaires compiled by the Hungarian Energy and Public Utility Regulatory Authority. Besides, waste statistics and ETS data were taken into account.

As it can be seen in Figure 3.2.5.4, total fuel consumption (without nuclear energy) in the energy industries sector shows strong fluctuations. After a significant decrease around the political and economic regime change in 1990, we could experience some increase till 1998, then a slight decrease till 2005 and a more pronounced drop after 2008 due to the global financial crisis. After 2010, until 2014, fuel consumption has reached record low values every year. In 2015, however, the decreasing trend stopped, and we observed a small increase in energy use.

Within the inventory period, the consumption of liquid and solid fuels decreased significantly. In contrast, the consumption of natural gas increased until 2007 to a great extent then it shrunk substantially afterwards. Biomass use due to burning or co-burning in power plants became more and more important and exceeded in amount the liquid fuel use in 2005. In 2006, the greatest power plant of Hungary reduced biomass-use, because the amount of obligatory purchased electricity was less than in 2005, this is also illustrated on Figure 3.2.8. In 2007, the produced electricity increased by more than 11%, in parallel the fuel consumption (mainly natural gas) increased only by 9%, because the efficiency of natural gas combustion is better than that of the others. Biomass burning in power plants became again popular on favorable terms, which was induced by the EU carbon trading. In 2008, the produced electricity from fossil fuels and also the fossil fuel consumption of this sector decreased again, but the total generated electricity – including nuclear, waste and renewable sources – was a bit higher than in the previous year. In 2009, the electricity generation in Hungary was by 10% less than in 2008. The generation decrease of power plants of 50 MW and higher capacity was 11.6% while it was 2.8% in case of small power plants. The fuel-mix also changed in 2009: coal and natural gas consumption decreased, however liquid fuel use increased, but its contribution to total fuel consumption is very low. Use of nuclear, waste and renewable sources continued to increase. In 2010 domestic electricity production increased again by 4%.

In 2011, electricity production fell back by 4% which meant lower fuel use at power plants. Moreover, the decrease in fossil fuel use was more pronounced, whereas there was only a slight change in GHG emission irrelevant nuclear fuel use.

In 2012, gross electricity production fell back by a further 4%. Moreover, the decrease in natural gas based electricity production was the most pronounced (-12.5%), whereas the share of air pollutant neutral nuclear fuel has steadily grown in the last few years, and wind energy utilization showed a steep increase. In addition, electricity import grew significantly by 16% in 2012.

This trend continued and even intensified in 2013. Domestic electricity production has dropped by a further 13 per cent. At the same time, net import grew by 49 per cent!

The overall picture did not change in 2014, either. We experienced decreasing production levels (-3%) and increasing import (+13%). In fact, net import was never higher in the whole period (1980-2015) than in 2014, and electricity production was never lower since 1990 (see Fig. 3.2.6).

In 2015, the share of import remained at a quite high level (31%). At the same time, production increased by 3% mainly due to a 20% growth in production of natural gas fired plants.




Figure 3.2.5.4 Fuel combustion in the Energy Industries Sector (1985-2020)

Electricity production increased further in 2016 by 5%. Again, we could observe a large growth in natural gas-based power production (+27%). At the same time, the share of import remained at a quite high level (28%).

And the growth in electricity production did not stop in 2017, either (3%). Natural gas firing power plants produced 20% more electricity than in 2016. The share of import did not change significantly.

The growing trend stopped in 2018, electricity production decreased by 3% whereas import grew by 11%

In contrast to the previous year, production increased again in 2019 by 7%. At the same time, import decreased by 12% but still represents a share of 27% within domestic electricity consumption.

Gross electricity production increased further by 2% in 2020. However, production from fossil fuels increased only slightly by 0.1% whereas from renewable sources by 17.9%

Fuel consumption of oil refining showed a pronounced drop around 2000 but remained more or less at the same level afterwards. In the last three years, however, increasing fuel consumption could be observed. Currently, the share of the refinery's fuel consumption is about 14% within energy industries. Less significant is manufacture of solid fuels and other energy industries with a portion of 2-4%.

Going into more detail regarding fuel use, it can be seen that domestically produced lignite is the dominant fuel among solid fuels (Fig. 3.2.5.5). In energy industries, solid and gaseous fuels are dominant representing together around 766-77% of all fuel use. In contrast, liquid fuel use became almost negligible in electricity and heat generation. At the same time, refinery gas used in oil refinery became the most important liquid fuel type whereas the formerly dominant fuel oil almost disappeared (Fig. 3.2.5.6).




Figure 3.2.5.5 Share of different solid fuels used by energy industries (1985-2020)




Figure 3.2.5.6 Share of different liquid fuels used by energy industries (1985-2020)

Traditionally, refinery gas and heavy fuel oil were reported together in the Hungarian Energy Statistical Yearbooks. Expressed in mass units, three-four times more refinery gas is used in the refinery as fuel oil.

However, as the ETS data show, refinery gases have significantly different characteristics. Based on plant specific information from the period 2008-2020, “real” heavy fuel oil burned by the refinery has a net calorific value between 39.8 TJ/kt to 40.9 TJ/kt and a CO₂ emission factor between 79.3 t/TJ and 83.7 t/TJ. Refinery gases show in contrast more diverging values. (Even more so, if we include tail gases and purge gas, too.) We can see here calorific values between 45.6 TJ/kt and 66.1 TJ/kt with corresponding CO₂ emission factors between 32.2 t/TJ and 60.3 t/TJ. On yearly average, it can be calculated with a NCV of 48.7-52.4 kt/TJ and an EF of 49.8-55.0 t/TJ for refinery gases.

There are some differences between the classification of fuels in the plant and in the energy statistics. However, considering these fuels together, we could see a better agreement. Moreover, the calculated average CO₂ IEF based on the IEA energy statistics for the pre-ETS period (2000-2007) and the ETS period (2008-2013) is the same (i.e., 62.2 t CO₂/TJ). For the 90’s, where the share of refinery gas was

definitely lower with 40% on average. On the other hand, the use of end-gases and purge gas (as reported under the EU ETS) became more important after 2013. All the factors applied in the calculations are summarized in the following table. Please note that the factors in the table below can be regarded as implied factors for the ETS period as ETS data are directly used in the inventory.

Table 3.2.5 Country specific parameters used in the category petroleum refining

Period	Fuel	(Avg.) NCV [TJ/kt]	EF [t CO ₂ /TJ]	Comment
2008-2020	refinery gas	45.7-66.1	32.2-60.3	ETS data
2008-2020	other liquid fuel	39.8-40.9	79.3-83.7	ETS data
1985-2007	gasoil	43.0	74.1	IPCC default
1985-2007	refinery gas	49.5	57.6	IPCC default
1985-2007	fuel oil	40.2	77.4	CS / IPCC default
1985-2020	petroleum coke	32.5	106.0	EF based on ETS

Emission factors

Carbon dioxide emissions were calculated in accordance with the 2006 IPCC Guidelines. Country specific OF and EF values – taken mostly from the ETS database – were used for most solid fuels and some liquids. The used factors are summarized in Table 3.2.6.

Table 3.2.6 CO₂ emission factors used in energy industries in the 2020 inventory year

Fuel type	Emission factor (CO ₂ t/TJ)	Oxidation factor
Coking coal	94.6	1.0
Other Bituminous Coal	94.6	1.0
Lignite (imported brown coal)	<i>96.1-98.8</i>	<i>0.979-1.0</i>
Lignite (domestic brown coal)	<i>92.9-97.1</i>	<i>0.948-1.0</i>
Lignite (domestic lignite)	<i>112.57</i>	<i>0.941</i>
Coke Oven Gas	44.4	1.0
Blast Furnace Gas	<i>252.2-252.5</i>	1.0
Gas/Diesel Oil	74.1	1.0
Residual Fuel Oil	77.4	1.0
RFO in refinery	<i>81.8</i>	1.0
Refinery gases (IEF)	<i>54.5</i>	1.0
Petroleum Coke	97.5	1.0
Natural Gas (in PPs)	<i>55.8</i>	1.0
NG in coking plant	-	1.0
NG in the refinery	<i>55.7</i>	1.0
Biomass (Solid)	112.0	1.0
Biogases	56.6	1.0
Waste (IEF)	93.8	1.0

(Source: 2006 IPCC Guidelines; in bold and italics – EU ETS database of Hungary)

*For waste, only IEF is reported in summary the table, because the emission was calculated from country-specific waste amount and component data taken from Waste Information System database and the emission factors were calculated using the default or measured (from EU ETS) carbon content and fossil carbon fraction data from Table 2.4 – 2.6 in the 2006 Guidelines.

It should be noted that only those measured factors were applied where the EU ETS covers all or most of the installation of the sector. For methane and nitrous oxide, default emission factors were used generally.

As recommended by the ERT and required by the guidelines, the emissions from waste incineration for energy purposes have been allocated to the energy sector. However, emissions estimation in the energy sector is somewhat different from the methodology used in the waste incineration category. Activity data in this source category are expressed in energy consumption units (TJ) whereas in the waste sector mass and composition of waste serves as basis of calculations. For our calculations four main activity data sources were used: data from the Waste Incineration Works (FKF) of Budapest (1985-2020), the Hungarian Waste Management Information System (2004-2020), the IEA Annual Renewable Questionnaire, and the ETS data (2006-2020). The Hungarian Waste Management Information System comprises facility level data on mass and composition of waste in line with the European Waste Catalogue (EWC codes) but also on waste management methods in accordance with the Waste Framework Directive. The latter made it possible to distinguish between waste incineration on land (D10) and use of waste principally as a fuel or other means to generate energy (R1).

Incinerated waste data expressed in energy unit were either directly taken out from the IEA statistics or from the ETS database. (The latter made it possible to distinguish between fossil and biogenic waste in case of mixed waste incineration.) It should be stressed, however, that the reported TJ values are (mostly) not used for CO₂ emission estimations, especially in the case of the biggest municipal waste incineration plant in Budapest, therefore the resulting IEF values may have little significance.

As only CO₂ emissions resulting from incineration of carbon in waste of fossil origin should be included in the national CO₂ emission estimate, the fossil fraction of waste had to be determined. To do so, country-specific waste amount and composition data were needed, and the emission factors could be calculated using the default carbon content and fossil carbon fraction data from Table 2.4-2.6 in the 2006 Guidelines. In case of the two biggest incinerators, plant specific data were used. The Waste Incineration Works (FKF) of Budapest determines regularly the composition of incinerated municipal solid waste (MSW), therefore the fossil carbon fraction could easily be calculated with the help of Table 2.4 of the 2006 Guidelines. The fossil carbon fraction of MSW grew from 5% in 1990 to 17-18% around 2010 and decreased to 10-13% in 2014 and afterwards. (In 2019, it was 10.6%.) CO₂ emissions were estimated then with an oxidation factor of 1.0.

The biggest co-incinerator plant is Mátra Power Plant. Since this plant reports its verified emissions in the framework of the European emission trading, direct ETS data relating its fuel use and CO₂ emissions were taken over. (Also verified emissions of all other smaller plants reporting waste combustion under the ETS were taken into account).

All in all, waste incineration contributed around 211 Gg CO₂ to GHG emissions in this category in 2020.

CH₄ emissions from waste incineration have also been added to the inventory. Using the default emission factors (30 kg/TJ) from Table 2.2 of the 2006 Guidelines (Chapter 2: Stationary Combustion), the resulting emissions are not significant at all. The same can be stated about N₂O emissions that were estimated the same way with the default emission factor of 4 kg/TJ.

1.1.1.1 Uncertainties and time-series consistency

Practically, the accuracy and uncertainty range of the energy statistics data are determined by the accuracy of the measuring equipment (except for stock changes, which are based on expert estimates

and are not comparable with the quantity of fuels from other sources). Taking all this into account, the estimated uncertainty of the energy consumption data is $\pm 1\%$ (for biomass 5%). This is particularly likely because the quantities of fuels used by power stations were verified using the report of MVM Rt. (Hungarian Power Companies Plc.)

The estimated specific uncertainty for CO₂ is 2-5%. The uncertainty of the methane factor is significantly higher (50-150%), while that of N₂O may be of an order of magnitude.

The time series can be regarded as consistent.

3.2.5.3 Category-specific QA/QC and verification

Energy consumption data were subject of several rounds of verification before use.

National energy statistics as published in the yearbooks were compared with the statistics provided to international organizations (both prepared by the same institute). This verification pointed out some problems also previously (e.g., on coke oven/blast furnace gas use, missing refinery gas and petroleum coke consumption) which were corrected. This work has been extended, and a comprehensive consistency check between data in the IEA time series and the Hungarian Energy Statistical Yearbooks has been conducted. Based on the results of this consistency check, and after several consultations with the energy statistics provider, it was decided to build the calculations on the IEA/Eurostat questionnaires.

Verified energy use from EU ETS was compared to statistical data. It was noticed that data in metric tonnes are similar in the ETS to those in the statistics, but there are some differences in energy values due to different NCVs. Since the energy consumption in sectoral approach should be compared with those of reference approach, we kept the NCVs of the energy statistics, however the emission factors of coals were corrected for some years to achieve consistency in energy balance and verified emissions. Measured oxidation factor was also applied in the calculation for the above-mentioned reason. (As the fuel amounts in the ETS database and in the energy statistics shows good agreement, this means in practice that CO₂ emissions from solid fuel use reported under the ETS can be used directly.)

Emissions from natural gas were generally estimated using default calorific values and emission factors. For a justification of this approach, about 40 emission reports from the ETS had been analyzed. Using the same activity data as reported by these facilities, we have calculated CO₂ emissions with default parameters and compared our results with the reported CO₂ emissions from the ETS database. It turned out that the difference was minor: with default parameters, the emissions were overestimated only by 0.4%. This small difference allowed us to change our previous approach. To be more consistent with the emissions reported under the ETS regime, we have switched to country specific emission factors for 2010-2013. (For all other years, for the time being, default EFs are applied.)

A comparison between the ETS data (based on mass balance methodology) and calculations based on fuel use was also made for the coking plant. This comparison led to changes in methodology for this submission as described in recalculation part below.

As the main fuel consumption is related to public electricity and heat production, a comparison was also performed with independent dataset collected by the Hungarian Energy Office. For the main power plants the total fuel consumption's difference between the ETS and this dataset was around 1% in 2009.

As a new practice, the compiler institute receives the draft version of the energy balance (the IEA annual questionnaires) before its official submission to Eurostat. The data are checked and the comments are discussed with the energy statistics provider and many of them are taken into account in the final energy statistics.

3.2.5.4 Category-specific recalculations

Activity data have been revised on the basis of the latest IEA/Eurostat Annual Questionnaire submitted in November 2021 to Eurostat. For example, natural gas consumption in 1A1a was revised by +2PJ for 2019. No methodological change occurred.

The overall effect of the above recalculations was: +119.3 kt CO₂-eq or 0.2% of the total emissions in 2019, and 0.0 kt in the base year.

3.2.5.5 Category-specific planned improvements

None.

3.2.6 Manufacturing Industries and Construction (CRF sector 1A2)

3.2.6.1 Source category description

Emitted gases: CO₂, CH₄, N₂O

Methods: T1, T2, T3

Emission factors: D, CS, PS

Key sources:

1A2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels - CO₂ – L, T

1A2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels - CO₂ – T

1A2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels - CO₂ – L, T

This subsector covers emissions from the combustion of fuels in the industrial sector. One of the advantages of using the IEA/Eurostat questionnaires instead of the (previously used) energy statistical yearbooks is that the industrial sectors in the questionnaires and in the CRF tables can be more easily harmonized. Emissions from autoproducers have been re-allocated to the relevant end-use category to the extent possible, the remaining part that could not have been allocated elsewhere, was included in the category *1A2g.viii Other* for years before 1998. Emissions from off-road vehicles and other machinery are reported as a separate category (1.A.2.g.vii).

Emissions in the Manufacturing Industries and Construction Sector:




Figure 3.2.6.1 Trends of CO₂, CH₄ and N₂O emissions in the Manufacturing Industries and Construction Sector (1985-2020)

3.2.6.2 Methodological issues

The energy consumption data have been taken from the IEA/Eurostat questionnaires. All feedstock and non-energy use were removed from the chemicals subsector for the entire time-series, and all relating CO₂ emission originating from non-combustion processes can be found in the Industrial Processes Sector.

Part of the emissions from waste incineration for energy purposes was allocated to this source category. Activity data in energy units were taken directly from the IEA Renewable questionnaire/ETS database with preference to ETS data in case of differences. Special attention was given to the four

(currently three) big cement factories, as they incinerate large amount of waste of fossil origin (plastics, rubber etc.). Their verified ETS data (emissions and fuel use) were analyzed, from which a specific emission factor was derived: 85.5 tonne CO₂/TJ waste. This EF was used for the years 2004-2007 in case of fossil wastes. From 2008 on, ETS data (emission) of the cement factories were used directly. The ETS data made it also possible report the fossil part of mixed industrial (or municipal) waste separately. It could be seen that other industrial facilities incinerate predominantly waste of biogenic origin, mostly wood waste, therefore their CO₂ emissions did not contribute to the national total. The insignificant CH₄ and N₂O emissions were estimated for all waste (not only fossil but also biogenic) using the default emission factors of 30 kg/TJ and 4 kg/TJ, respectively.

The methodology for off-road vehicles and other machinery used in industry and construction was changed recently. Tier 2 method from the 2016 EMEP/EEA Guidebook was implemented. This method classifies the used equipment into the fuel types and layers of engine technology. The engine technology layers are stratified according to the EU emission legislation stages, and three additional layers are added to cover the emissions from engines prior to the first EU legislation stages. The used layers are as follows: <1981; 1981-1990; 1991-Stage I; Stage I; Stage II; Stage IIIA; Stage IIIB; Stage IV; Stage V. The penetration of the new technology is taken into account in the form of split (%) of total fuel consumption per engine age (irrespective of inventory year) as it can be seen for diesel-fueled non-road machinery in Table 3-3 in the Guidebook.

Activity data

Figure 3.2.6.2 illustrates the energy consumption of the sector. After 1990, following the economic changes, fuel use decreased significantly. The underlying reasons are clearly illustrated by the decreasing production data presented in the IPPU sector (Chapter 4).




Figure 3.2.6.2 Fuel combustion in the Manufacturing Industries and Construction Sector (1985-2020)

In 2009 the global economic crisis caused a drop of fuel consumption by more than 25% which led to lower emissions. In 2010, the growing industrial production increased the energy demand that did not change much either in 2011 or in 2012. In 2013, however, fuel consumption reached almost the level of the years before the economic crisis, and in 2017 it grew higher than any year after 2000. The increasing trend continued also in 2018 and 2019. In 2020, fuel consumption decreased a little by 2%.

Fig. 3.2.6.3 clearly demonstrates the dominance of natural gas (61% in 2020). Biomass use became popular especially in the last decade. (As a consequence, the relative share of methane and nitrous oxide emissions increased.) Combustion of oil products continues to lose its importance among fossil fuels. Liquid fuels represented 17% in 2020 out of which gas oil seems to be the most important.




Figure 3.2.6.3 Share of different liquid fuels used by manufacturing industries (1985-2020)

The share of solid fuels became quite low (5% in 2020). Also, the fuel mix has been changing as demonstrated by Fig. 3.2.6.4. The growing relative share of coke oven gas define the CO₂ IEF in the iron and steel category since coke oven gas has a very low (44.4 t/TJ) CO₂ emission factor. It is worth noting the relatively high IEF in food processing, beverages and tobacco which is due to the fact that dominantly (recently almost exclusively) coke was used as solid fuel by this industry. In 2016, a new co-generation power plant that uses solid fuels, including the paper mill's own residual waste from paper production, but also biomass and coal, started its operation, this is why the consumption of other bituminous coal increased as seen in the figure below.




Figure 3.2.6.4 Share of different solid fuels used by manufacturing industries (2005-2020)

Biomass cannot be considered as the most important fuel but its contribution grew slowly to 12 per cent. Within this the growing share of biogases (especially in autoproducer plants) might deserve our attention as the default emission factors are quite different for solid biomass and biogas.

Emission factors

With the introduction of county-specific CO₂ EF for natural gas, the share of emissions calculated with default CO₂ factors were significantly reduced. Special country specific emission factors are applied in the *non-metallic minerals* category (based on ETS information). The situation is somewhat complicated here as the cement factories often use mixed fuels. The applied country specific CO₂ emission factors for petroleum coke/coal mix are varying between 92.3 t/TJ and 95.0 t/TJ for the period 2008-2020.

3.2.6.3 Uncertainties and time-series consistency

Practically, the accuracy and uncertainty range of the energy statistics data are determined by the accuracy of the measuring equipment (except for stock changes, which are based on expert estimates and are not comparable with the quantity of fuels from other sources). Taking all this into account, the estimated uncertainty of the energy consumption data is 5% in consideration of the fact that uses are less easy traceable due to the high number of users.

The estimated specific uncertainty for CO₂ is 2-5%. The uncertainty of the methane factor is significantly higher (50-150%), while that of N₂O may be of an order of magnitude.

3.2.6.4 Source-specific QA/QC and verification

Energy consumption data were subject of several rounds of verification before use.

Verified energy use from EU ETS was compared to the statistical data. It was noticed that data in metric tonnes are similar in the ETS to those in the statistics, but there are some differences in energy values due to different NCVs.

Non-energy use of fuels was cross-checked with the Industrial Processes sector.

3.2.6.5 Source-specific recalculations

- Revised energy statistics was used (as submitted to the IEA/Eurostat in November 2021;
- Emissions from all autoproducer plants have been (re-)allocated from the source category 1A2g to the relevant economic sector where they operate fully for the period 1998-2020, and partly for earlier years.

All the above changes resulted in a decrease of -13.4 kt CO₂-eq (-0.0% of the national total) in 2019, and 0.0 kt in the base year.

3.2.6.6 Source-specific planned improvements

None.

3.2.7 Transport (CRF sector 1A3)

3.2.7.1 Source category description

Emitted gases: CO₂, CH₄, N₂O

Methods: T1, T2, T3

Emission factors: D, CS, M

Key sources:

1A3b Road Transportation – CO₂ - L, T;

1A3c Railways – CO₂ – T

1A3d Domestic Navigation - Liquid Fuels – CO₂ – T (only excl. LULUCF)

This sector covers all the emissions from fuels used for transportation purposes. International aviation and navigation are excluded from the national total.

Looking at the whole period of our time series, a sharp decrease of 60% in transport of goods could be observed during the regime change in the early 90's. The Hungarian transport performance expressed in freight tonne-kilometres had not reached the level of 1985 until 2005. Beside these significant changes of volume, also the structure of goods transport altered. Currently, the most important means of freight transport is road transportation with a share of 61%, followed by rail (22%), pipeline (13%) and waterway (3%). In 1990 we saw a completely different picture with railway and waterway being the dominant mode of transport representing 40% and 34%, respectively. The share of road transportation was 15% about 25 years ago. In 2020, there was a drop in transport (-10% in freight tonne-kilometres) especially due to a 13% decrease in road transportation.




Figure 3.2.7.1 Trends in goods transport (2001-2020). Source: HCSO

Passenger transport also underwent considerable changes. The stock of passenger cars had more than doubled since 1990. Within this increase, the proportion of Eastern European cars characterized by high fuel consumption and obsolete technology decreased; for example, currently more than half of the passenger cars complies with at least the Euro 4 emission standards. At the same time, the average age of the car fleet has increased again in recent years to 14.7 years in 2020. (The lowest average age

of vehicles (10.3 years) was observed in 2006, before the economic crisis.) Figure 3.2.7.2 summarizes the above-mentioned developments.




Figure 3.2.7.2 General changes in the transport sector

Electrification of the railways in Hungary eliminated mostly the solid fuel consumption. (Today there are only few lines where steam engines are used during non-scheduled vintage train trips.) Diesel oil consumption of railways decreased as well, by 80% between 1990 and 2020.

Emissions were calculated generally from the national fuel consumption data from the IEA/Eurostat annual questionnaires. However, national statistics usually does not have separate lines for the quantities of aviation gasoline used for in-country aviation and of the diesel oil used for international (river) navigation. However, aviation gasoline consumption appeared in the latest energy statistics for the years 2016-2020. Fuel consumption data (i.e., both aviation gasoline and jet kerosene) of domestic aviation are taken from the Eurocontrol database that contains data on IFR flights. We can also assume (based on personal communication with the energy statistics provider) that 0.9-1.0 kt aviation gasoline is consumed for domestic flights. It is still possible that some minor amount of aviation fuel (for VFR flights) is included elsewhere in the inventories (e.g., under road transport).

According to the information received from the energy statistics provider, natural gas use related to natural gas transport was previously included under distribution losses in the energy statistics. In the inventory, however, a complete time series of emissions from pipeline transport is included separately. Figures below illustrate the fuel consumption of the sector:




Figure 3.2.7.3 Gasoline, diesel and biomass consumption in the Transport Sector (1985-2020)




Figure 3.2.7.4 LPG, natural gas and solid fuel combustion in the Transport Sector (1985-2020)

Figure 3.2.7.5 clearly shows that in contrast to the other described sectors, transport consumption had a rising overall tendency from the mid 90's until 2008. Starting in 2009, the trend of fuel consumption has changed due to the economic crisis. Both fuel consumption and mileage of vehicles (km/year) increased until 2009 and started decreasing afterwards. The increasing fuel prices (up to 2012) could also be one of the reasons of a record low gasoline consumption in the transport sector. It is worth mentioning that the mass of domestically transported goods via road transport decreased by 44% between 2008 and 2012. However, the decreasing trend stopped, fuel consumption started to grow again and goods transport increased by 28% since 2012. Then, in 2020, there was a drop of 8% in national transport of goods, and at the same time fuel consumption decreased quite significantly by 12%.

In the second half of 2005 the Hungarian oil and gas company's refinery, MOL Danube Refinery, started to process bioethanol from vegetable raw material with high sugar content, also biodiesel have been used for blending. These bio components appear also in Fig. 3.2.7.3.

LPG has been used since 1992. It should be noted that due to the current commercial practices, in-container (household, institutional) uses are difficult to separate from traffic uses (i.e., distribution at petrol stations). This may be the reason for the sharp increase in 2003, which does not fully reflect the actual changes but is the result of a change in the approaches used for the preparation of the statistics. Accordingly, liquid fuel uses by the general public (currently including LPG only) show a significant drop – in line with the national energy statistics.




Figure 3.2.7.5 Trends of CO₂, CH₄ and N₂O emissions in the Transport Sector (1985-2020)

3.2.7.2 Methodological issues

CO₂ emission from transport was previously calculated by multiplying fuel consumption taken from Energy Statistics Yearbooks (1985-2010) by the default IPCC emission factors. In 2013, as the compiler institute received data on carbon content of gasoline and diesel oil from the refinery, the default emission factors were replaced to country specific values in road transportation (see Table 3.2.9). From the 2014 submission, activity data are basically taken from the IEA/Eurostat joint questionnaires with a few modifications.

For non-CO₂ emissions, the COPERT-5 (**C**omputer **P**rogramme to Calculate **E**mission from **R**oad **T**ransport) model, specifically version 5.5.1 was used for the whole time series. The transition to the COPERT program family was a necessary step in the area of national road transport emission calculations, since most countries use this program which ensures international comparability. By using the latest version of the model, also consistency of the time series is ensured.

The COPERT model requires quite detailed background information. To produce input data for the model for the whole time series, basically three data sources were used:

1./ The compiler institute received the COPERT input/output data from the Institute for Transport Sciences for the years 2006, 2007, 2009, 2011, and the period 2012-2020. The structure of the input data was produced in a way which fully complies with that described in the software requirement.

Generally, the input data required by the COPERT model are as follows:

- vehicle fleet [n]

- mean activity [km/year], lifetime cumulative activity [km]
- traffic situations: vehicle share [%], average speed [km/h], trip characteristic
- national annual energy consumption [tons/year]
- country-specific environmental information:
 - national monthly averages of daily minimum and maximum temperatures [°C]
 - monthly average relative humidity [%]
 - Reid vapor pressure [kPa]
 - determination of country-specific sulfur content of petrol and diesel fuels [ppm wt]
 - determination of bioethanol ETBE (Ethyl tert-butyl ether) content (biodiesel FAME (Fatty acid methyl ester) content is provided in the model because it is known from EU data)

As the input data were not obtained from the same source and were not always suitable for direct use, therefore the data were needed to be processed prior. The largest bulk of the work was processing the vehicle stock data, since this data ensures the basis for emission calculations performed by COPERT5. Therefore, it was crucial to perform an utmost precise work regarding the vehicle stock data, which was obtained from the Ministry of the Interior (BM). At the request of the KTI, vehicle data tables required to perform the task were extracted from the BM database. The vehicle stock classifications and emission categorizations were prepared using the following table:

Table 3.2.7.1 *Vehicle categories used in the COPERT model*

Category	Fuel	Engine capacity [cm ³] / Gross weight [t]
Passenger Cars	Gasoline	2-stroke (≤ 1000 cm ³)
		≤ 800 cm ³
		801 – 1400 cm ³
		1401 – 2000 cm ³
		≥ 2001 cm ³
	Petrol Hybrid	≤ 800 cm ³
		801 – 1400 cm ³
		1401 – 2000 cm ³
		≥ 2001 cm ³
	Petrol PHEV	801 – 1400 cm ³
		1401 – 2000 cm ³
		≥ 2001 cm ³
	Diesel	≤ 800 cm ³
		801 – 1400 cm ³
		1401 – 2000 cm ³
		≥ 2001 cm ³
	Diesel PHEV	≥ 2001 cm ³
	LPG Bifuel	≤ 800 cm ³
		801 – 1400 cm ³

		1401 – 2000 cm ³
		≥2001 cm ³
	CNG Bifuel	≤800 cm ³
		801 – 1400 cm ³
		1401 – 2000 cm ³
		≥2001 cm ³
Light Commercial Vehicles	Gasoline	N1-I ≤1305 t
		N1-II 1306 – 1760 t
		N1-III 1761 – 3500 t
	Diesel	N1-I ≤1305 t
		N1-II 1306 – 1760 t
		N1-III 1761 – 3500 t
Heavy Duty Trucks	Gasoline	> 3,5 t
	Diesel	Rigid ≤7,5 t
		Rigid 7,5 - 12 t
		Rigid 12 - 14 t
		Rigid 14 - 20 t
		Rigid 20 - 26 t
		Rigid 26 - 28 t
		Rigid 28 - 32 t
		Rigid >32 t
		Articulated 14 - 20 t
		Articulated 20 - 28 t
		Articulated 28 - 34 t
		Articulated 34 - 40 t
		Articulated 40 - 50 t
		Articulated 50 - 60 t
Buses	Diesel	Urban Midi ≤ 15 t
		Urban Standard 15 - 18 t
		Urban Articulated > 18 t
		Coaches Standard ≤ 18 t
		Coaches Articulated > 18 t
	Diesel Hybrid	Urban
	CNG	Urban
L-Category	Petrol	Mopeds 2-stroke <50 cm ³
		Mopeds 4-stroke <50 cm ³
		Motorcycles 2-stroke >50 cm ³

		Motorcycles 4-stroke <250 cm ³
		Motorcycles 4-stroke >750 cm ³
		Motorcycles 4-stroke 250 - 750 cm ³

In the case of traffic situations, the percentage of runtime distribution and average speed values within driving conditions (urban, rural, motorway) for each vehicle category were used based on the results of previous research carried out by the Institute for Transport Sciences.

Specifying the average speed is less important in the case of rural and highway traffic as the function takes similar values between 45-105 km/h. However, determining the average speed for urban transport is more important, because of a difference of 1 km/h in the first third of the function results in a larger difference in emissions. Naturally, the functions vary from one pollutant to another, but the influence of speed is similar in each case.

Among the trip characteristics, it is important to mention the average travel time and duration. According to available statistics, European average of 12.5 km were determined by experts. The distribution of the distances traveled varies from country to country, but typically a large proportion (80%) travel only short distances (less than 15 km). It plays a significant role in the emissions of the cold start phase. The average travel distance of 12 km average travel time of 25 minutes was used.

Detailed and accurate calculations of mean activity could not have been made in previous years. Previously, data were obtained from queries extracted from the RKF (Regular Environmental Review) database provided by the Ministry of the Environment, and subsequently corrected based on the annual fuel consumption. However, in COPERT5 it is possible to provide fuel balanced mean activity, which the program automatically counts and takes into account when calculating the emissions. From 2018, there was a development research in the Institute for Transport Sciences and the project outcomes will be used for the 2019 emission calculations. From now on, the mean activity data will be more precise and the query system calculates the mileage records of the Vehicle Inspection Database for each vehicle category.

The source of the annual fuel consumption data was the national energy statistics provided by the Hungarian Energy and Public Utility Regulatory Authority (MEKH). The data published by the MEKH will also be transmitted to EUROSTAT. Energy conversions were executed following the values given in the EMEP/EEA air pollutant emission inventory guidebook 2019.

Table 3.2.7.2 Default density and calorific values of primary fuels determined using the EMEP/EEA air pollutant emission inventory guidebook 2019

Fuel	Density [kg/m ³]	Calorific values [MJ/kg]
Gasoline	750	43.774
Diesel	840	42.695
LPG	520	46.564
CNG	175	48
Biodiesel	890	37.3
Bioethanol	794	28.8

2./ For all the years in the period 2000-2018 for which no domestic data were provided by the Institute for Transport Sciences, data purchased from Emisia SA, developer of the COPERT model, were used as inputs. As claimed by the data provider, "the vehicle fleet and activity data provided by EMISIA SA for the compilation of national emission inventories with use of the COPERT model reflect our best

knowledge of national situation in each country until 2013. These data have been updated using the road transport dataset and methodology of the TRACCS research project. More specifically, TRACCS dataset of the period 2005-2010 has been combined with the previous FLEETS research project dataset (2000-2005) and with latest official statistics available (2011-2013) to produce aligned and up to date time series for the period 2000-2013 (no projection included). The quality, completeness, and consistency of these two projects datasets, which have been extensively reviewed and cross-checked, ensure that the compiled countries data are also of good quality."

In case of larger discrepancies between the Emisia database and domestic data, preference was always given to data from domestic sources, and the time series was smoothed out. Again, whenever necessary, the mileage data were slightly modified to reflect better the domestic statistics on fuel sold.

3./ The compiler institute produced input data for the remaining years (i.e., 1985-1999). Quantification of the stock of each road vehicle type was based on Statistical yearbooks of Hungary and annual reports of Ministry of Economy and Transport about the Hungarian vehicle fleet. Also, personal communications with experts took place. Compared to recent years where about 200 vehicle categories were taken into account, the input database for the earlier part of the time series is less detailed containing 35 vehicle categories, and it probably has a higher uncertainty.

It should be noted that unleaded gasoline was sold only after 1989. Since lead is poison for catalytic converters, it was assumed that real catalyst vehicle has been used after this time.




Figure 3.2.7.6 Elimination of leaded gasoline in Hungary

(Source: Hungarian Petroleum Association (MÁSZ), Annual Reports 1996-2008)

Emissions from **in-country aviation**, which represent a very low proportion, were taken previously equal to the emission from consumption of aviation gasoline, and calculated in those years when the related data were available in the energy balance. Where aviation gasoline was not indicated in a separate line, consumption and emissions were calculated together with road traffic gasoline.

Upon receiving data from Eurocontrol, the above approach was slightly modified. Although there are no scheduled commercial domestic flights in Hungary, Eurocontrol data for the period 2005-2015 suggested that about 0.22 per cent of total jet kerosene is used for domestic flights. Using the same share back to 1985, some kerosene (i.e., 12-37 TJ) is now allocated to domestic aviation. As regards aviation gasoline, based on personal communication with the energy statistics provider, it is assumed

that altogether 0.9 kt aviation gasoline is sold in the country. This amount was amended with Eurocontrol data on international fuel consumption. For all other (pre-Eurocontrol) years (i.e., 1985-2004) the average value of 2005-2015 was used. The resulting CO₂ emission (from both aviation gasoline and jet kerosene) is 4 Gg on average (i.e., far below the significance threshold).

*(Background of the **Eurocontrol data**: At the end of 2010 the European Commission signed a framework contract with EUROCONTROL, the European organization for the safety of air navigation, regarding 'the support to the European Commission in relation to climate change policy and the implementation of the EU ETS'.*

This support project is organized in different Work Packages. One of these Work Packages pertains to the improvement of GHG and air pollutant emissions inventories sub-mitted by the 28 Member States and the European Union to the UNFCCC and to the UNECE. The main objective of the WP is to assist EU Member States improve the reporting of annual greenhouse gas (and other air pollutant) emission inventories by e.g., estimating the fuel split domestic/international using real flight data from EU-ROCONTROL.

To support the inventory process for the submission in 2017, in November 2016 MS received fuel and emissions data for the years 2005 to 2015 as calculated by EUROCONTROL using a TIER 3b methodology applying the Advanced Emissions Model (AEM).

The individual fuel burn and emission data associated with each flight are processed with AEM, Excel reports covering UNFCCC and CLRTAP are generated then made available to EEA Member states via the EUROCONTROL ftp site.

All flights having a flight plan are captured by EUROCONTROL. Military flights are excluded. Civil flights flying on Visual Flight Rules (VFR) are not known to EUROCONTROL and are therefore not part of this system.

Looking at flight types in the database, it is possible to distinguish between domestic and international flights. Three aircraft groups are currently included in the EUROCONTROL dataset: jet, turboprop and piston.)

The annual total emissions of the national **railway** were determined as an exhaust gas component based on the data received from the national energy statistics provider, the Hungarian Energy and Public Utility Regulatory Authority which calculated the quantity of the fuel used in the national railway transport. Railway transport emissions are affected by many factors. However, since the currently used method of calculation is based on the fuel consumption of the rail traction, the factors described below do not have a direct influence on the calculation.

The total length of railway lines has not changed significantly in recent years. The number of locomotives has increased by 15% since 2005. The total volume of passenger transport in terms of the number of persons transported decreased between 2001 and 2019 by about 17 percent whereas passenger km increased by 22% (mostly due to increased air transport). As far as the railways are concerned, the decrease in passenger numbers was smaller (-9%). However, expressed in passenger kilometers, the decrease was more pronounced (-23%). In 2020 alone, due to COVID-19, number passengers decreased by 28% generally, railways passengers by 31%. Considering transport of goods, rail transport has been showing some sign of growth since 2011, especially domestically, but there was also a significant drop in 2020.

Emissions from **pipeline transport** are reported separately since the 2015 submission. The calculations are based on (amended) energy statistical data and default emission factors. The IEA Annual Gas Questionnaire contains fuel consumption data only for the period 2010-2015. Comparing the IEA data to the ETS data reported by FGSZ Natural Gas Transmission Ltd (Hungary's transmission system operator), we detected an underestimation in the IEA data for 2010 which was corrected on the basis of IEA data. For the earlier years in the time series backward interpolation was carried out as follows:

- for 2005-2009 fuel consumption data were taken from the ETS database;

- for the years before 2005, the sum of natural gas production + import was used as proxy information.

Emission factors

Carbon dioxide emissions were calculated using country-specific emission factors for gasoline in road transportation, for diesel and natural gas in all relevant categories, and for kerosene in aviation. Otherwise, default factors were applied as summarized in Table 3.2.7.3 below.

Table 3.2.7.3 *Some CO₂ emission factors in the Transport Sector*

Fuel type	Emission factor (kt CO ₂ /TJ)	Source of EFs
Gasoline	69.3	2006 IPCC Guidelines
in road transport		Refinery
fossil	71.3	based on carbon content
bio	71.8	
Gas/Diesel Oil		Refinery
fossil	73.6-74.3	based on carbon content
bio	70.8	
LPG	63.1	2006 IPCC Guidelines
Residual fuel oil	77.4	2006 IPCC Guidelines
Natural Gas	55.2-56.3	ETS database
Lubricants	73.3	2006 IPCC Guidelines
Jet kerosene	72.5-73.0	Refinery

It has to be noted that the cited CO₂ emission factors in road transport are derived values based on mass of the fuels and not necessarily on their energy content. We have also slightly deviated from the NCVs reported in the IEA/Eurostat Annual Questionnaire. Originally, the net calorific value applied in the Hungarian energy statistics was usually 42 TJ/kt for both fuels. However, there were indications that the real calorific value might be different. For example, the default NCVs are 43.8 TJ/kt for gasoline and 42.7 TJ/kt for diesel in COPERT. In the 2006 IPCC Guidelines, we can find even higher values: 44.3 TJ/kt and 43 TJ/kt for gasoline and diesel, respectively. And we have also one measurement from the refinery for diesel oil: that is 43.04 MJ/kg. So, in the 2017 submission the calorific values were changed to 44 TJ/kt and 43 TJ/kt for (fossil) gasoline and diesel oil, respectively. (Meanwhile, NCVs have been revised upwards also in the energy statistics.)

The basis of the emission factor was carbon content of the fuels received from the refinery, i.e., 0.8406 t C / t gasoline and 0.86275 t C / t diesel. However, the carbon content of the fuels relates to the fuel mix E5 (i.e., 5% biofuel) so it cannot be used for the fossil part of the fuel unchanged, especially in case of gasoline. Therefore, we have changed the used emission factor for gasoline (fossil part) by taking into account the difference between the default CO₂ emission factor for gasoline (3.180 kg CO₂/kg fuel) and for the blend E5 (3.125 kg CO₂/kg fuel) (See Table 3-12 in the 2016 EMEP/EEA Guidebook). This means, we have multiplied the original EF with 3.180/3.125 for the fossil part. All the resulting EFs are included in Table 3.2.9 above.

For the years starting in 2016, the refinery provided us with carbon content and calorific value of the *fossil part* of diesel from which the following country-specific values could be derived:

Gasoil	-2015	2016	2017	2018	2019	2020
NCV (GJ/t)	43.0	42.8	42.7	42.7	42.7	42.8
CO2 EF (t/TJ)	73.6	74.0	74.3	74.3	74.2	74.1

Also, the CO₂ emissions from the *fossil part of the carbon of biodiesel* have been calculated and added to the national total. The refinery confirmed that diesel had only FAME content (HVO is expected only from 2020). Therefore, CO₂ emission was calculated with the default parameters specific for FAME, i.e., 76.5% kg C/kg FAME, and 5.4% fossil part of C of FAME. The resulting emissions were not significant: they increased from 0.3 kt in 2007 to 33.0 kt in 2020. These CO₂ emissions are reported separately as “other fossil fuels/fossil part of biodiesel”. As regards the biofuel part: in case of biodiesel (FAME), the default carbon content of 76.5% kg C/kg FAME was used with the default bio part of carbon content (1-5.4%=94.6%). In case of biogasoline, a biogenic carbon content of 55.1% was applied in the calculations.

CO₂ emissions from lubricants in 2-stroke engines are also included in the transport sector. Activity data have been taken from the COPERT database (i.e., fuel consumption of 2-stroke cars, mopeds, and motorcycles. With an assumption of a 1:40 mixing ratio, the total amount of lubricants combusted could be calculated. Using default NCV and EF, the resulting emissions remained at a moderate level as summarized in the table below.

Table 3.2.10 CO₂ emissions from lubricants in 2-stroke vehicles (with activity data)

		BY	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
Fuel consum.	kt	425.6	460.0	295.9	173.0	65.1	36.8	17.9	15.3	14.3	13.9	17.9	16.1
2-stroke cars	kt	416.1	452.0	282.2	159.9	41.8	13.7	5.1	3.3	2.6	2.3	2.3	1.9
L-Category	kt	9.5	8.1	11.0	13.1	23.3	23.1	12.8	12.0	11.7	11.6	15.6	14.3
Lubricants	kt	10.6	11.5	7.4	4.3	1.6	0.9	0.5	0.4	0.4	0.4	0.4	0.4
CO2 emissions	kt	31.1	33.6	21.6	12.6	4.7	2.7	1.3	1.1	1.0	1.0	1.3	1.2

CH₄ and N₂O emissions from road transport were calculated using the COPERT model (COPERT 5.5.1) for the whole inventory period for gasoline and diesel. For all other fuels (and categories) default IPCC emission factors were applied. For example, non-CO₂ EFs from Table 3.4.1 of the 2006 IPCC Guidelines are applied for railways.

Besides gasoil as described above, country-specific calorific value and CO₂ emission factors are applied also for kerosene as summarized in the table below.

Kerosene	-2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NCV (GJ/t)	43.4	43.5	43.4	43.4	43.3	43.3	43.3	43.3	43.5	43.3	43.3	43.3	43.3	43.4
CO2 EF (t/TJ)	72.7	72.7	72.7	72.6	73.0	72.8	72.7	72.7	72.5	72.7	72.7	72.8	72.8	72.6

3.2.7.3 Uncertainties and time-series consistency

We assume that the uncertainty of the transport-related fuel consumption data is higher than in case of stationary equipment because such data are more difficult to collect and verify. Considering the above, the estimated uncertainty of the energy consumption data is $\pm 5\%$. The estimated uncertainty of the emission factors for CO₂ is $\pm 1-5\%$. It should be noted, that in the 2006 IPCC Guidelines the uncertainty for default methane and nitrous oxide factors is much higher (200-300%).

3.2.7.4 Source-specific QA/QC and verification

IEA data were compared with the national statistics. For clarification of the differences, additional data were required from the energy statistics provider. This led to revision of the time series of fuel consumption.

We consider the technical review of the EU as a very important QA activity. In summer 2012, the EU conducted a more thorough than usual review of the inventories of all member states. After the review, Hungary was recommended to obtain the C content and net calorific values of gasoline from fuel suppliers, to develop a country-specific EF for CO₂ from gasoline that is representative for gasoline used in Hungary and to revise data accordingly.

During the 2016 review, it was recommended that CO₂ emissions from lubricants for non-energy use should be reported under 2D1 Lubricant use and emissions from energy uses (such as in 2-stroke engines) should be reported under 1A3b Road transportation. Following this recommendation, this submission contain estimates of CO₂ emissions of lubricants from 2-stroke engines taking fuel consumption data (FC_t) from the COPERT database (2-stroke passenger cars, mopeds, and motorcycles).

3.2.7.5 Source-specific recalculations

The used methodology remained basically the same. However, the following changes were implemented:

- The latest IEA Annual Questionnaires (as submitted in November 2021) was used that contained some revisions
- The latest version of the COPERT model (5.5.1, September 2021) was applied for the whole time series consistently
- Lubricants used in 2-stroke engines have been revised on the basis of new outputs from the COPERT model.

Altogether, emissions decreased insignificantly by 3.4 kt CO₂-eq (0.0% of the national total) in the base year, and increased by 2.3 kt (0.0% of the national total) in 2019.

3.2.7.6 Source-specific planned improvements

None.

3.2.8 Other Sectors (CRF sector 1A4)

3.2.8.1 Source category description

Emitted gases: CO₂, CH₄, N₂O

Methods: T1, T2

Emission factors: D, CS

Key sources:

1A4 Other Sectors - Liquid Fuels – CO₂ – L, T;

1A4 Other Sectors - Solid Fuels – CO₂ – L, T;

1A4 Other Sectors - Solid Fuels – CH₄ - T

1A4 Other Sectors - Gaseous Fuels – CO₂ – L, T;

1A4 Other Sectors - Biomass – CH₄ – L

This sector covers combustion in public institutions, by the population and in the Agriculture /Forestry/Fisheries Sector. Emissions in the Other Sectors are summarized in the figure below:




Figure 3.2.8.1 Trends of CO₂, CH₄ and N₂O emissions in the Other Sector (1985-2020)

3.2.8.2 Methodological issues

Activity data

Activity data was obtained from the IEA/Eurostat questionnaires as described in the introduction section of the chapter. Figure 3.2.8.2 illustrates the fuel consumption of the sector by fuel types.




Figure 3.2.8.2 Share of different combusted fuel types in the Other Sectors (1985-2020)

Since about two third of the fuel consumption is related to the residential category, the fuel structure is influenced principally by changes in this sector (see Fig. 3.2.8.3).




Figure 3.2.8.3 Fuel combustion in the subsectors of the Other Sector (1985-2020)

Generally, in contrast with the significant reduction of coal and oil consumption, natural gas consumption has increased significantly. The population switched from coal to natural gas combustion. Household heating oil was completely replaced by LPG (see Table 3.2.8.1).

Table 3.2.8.1 Oil and LPG consumption in the Commercial/Institutional and Residential Sectors in selected years after 1990

		1990	2000	2005	2010	2015	2016	2017	2018	2019	2020
Commercial Institutional	Oil	13,929	1372	379	0	1,064	1,015	847	978	811	812
	LPG	1,504	2209	1,081	893	658	508	415	554	508	692
Residential	Oil	35,991	1118	86	0	40	0	0	0	0	0
	LPG	13,536	12,079	7802	5640	3055	2,493	3,138	3,184	3,091	3,322

During the period 1990-2019, the length of natural gas pipe-network increased from 22,549 km to 84,906 km. The number of households supplied with natural gas increased from 1.6 million in 1990 (42%) to 3.4 million in 2010 (77%) but decreased a little to 3.3 million (73%) since 2010. Residential consumption represented 34% of total inland demand of natural gas in 2020. Piped gas is available in 91% of all settlements in Hungary, and this figure has not changed much since 2005 (but it was only 15% in 1990). 72% of households use natural gas for heating purpose as well. Although individual residential heating became more and more widespread, still 651 thousand dwellings (15% of all dwellings) are supplied with district heating and 600 thousand with hot water. Most of this heat (over 80%) is generated from natural gas use; however, the resulting emission was not accounted for here but under the Energy industries subsector.

Natural gas consumption can be influenced by several factors. One of these factors might be the weather and the resulting heating demand. Heating degree day (HDD) is a quantitative index that reflects demand for energy to heat houses and businesses. This index is derived from daily temperature observations. The inside temperature is 18°C and base temperature (the outside temperature above which a building needs no heating) is 15°C in our calculation (following the standard European methodology). Figure 3.2.8.4 illustrates the relationship between residential fuel consumption and HDD. The figure demonstrates that increased fuel use can often be explained by increased HDD values and vice versa. Heating demand decreased by 7% in 2020 (after the two warmest years) which is reflected in the higher energy consumption of the residential sector.




Figure 3.2.8.4 Comparison of residential fuel consumption and HDD between 1995 and 2020

Another factor is definitely the price. The (nominal) price of pipelined gas increased from 325 to 1360 Ft/10 m³ between 2000 and 2012. This price increase might have led to increased biomass use as a substitute fuel in the residential sector. However, the above-mentioned trends have changed in recent years. Gas prices have dropped by 26% since 2012 (but are still more than double as high as in 2005), and consumption started growing again.

So, it seems that the price elasticity of demand of natural gas and other fuels. We know that the price of natural gas was significantly higher in the period 2008-2013 than that of biomass, and in this very period natural gas consumption decreased and biomass consumption increased. After 2014, however, the trend changed due to decreased natural gas prices (the price advantage of biomass disappeared), so gas consumption started increasing again while biomass consumption decreased. This is demonstrated in Figure 3.2.8.5 below.




Figure 3.2.8.5 Price elasticity of natural gas and fuelwood (2005-2020)

The monthly natural gas consumption of an average household decreased from 125 m³ in 2003 to 70 m³ and then increased to 91 m³ in 2020. In this significant decreasing trend - beside the higher energy prices – most probably also the more energy-conscious approach of the population plays a role and is definitely greatly affected by the weather. In addition, larger decrease in biomass use indicates some fuel switch from fuelwood to natural gas in the residential sector.

Emissions from household machinery are reported separately in the category 1A4bii. Based on the latest survey of the Statistical Office, 56% of the households have garden or backyard on their own. There are 3.9 million households in Hungary; 56% of which is 2.2 million. It was assumed that for every garden 5 liters gasoline is used in a year. This would translate to 10.95 million liters or 8.2 kt gasoline. As part of the households use electronic devices, 6 kt of gasoline use was assumed for the whole time series. (The required activity data have been reallocated from the category 1A3b road transportation.) The resulting emissions are not at all significant (i.e., 19 kt CO₂). We would like to stress here that we consider the above approach as a rough estimate and as currently no other reliable information is available that would justify the introduction of any trend into the time series, constant amount is reported. In this regard it is worth mentioning that (1) all gasoline consumption in the energy statistics from the annual IEA/Eurostat questionnaires is accounted for in the inventory therefore it is merely an issue of allocation, and (2) emissions are below the threshold of significance.

In the **Agriculture** category, the trend in biogas use might deserve our attention as its share within biomass can be higher as 60% as in 2013-14, or around 40% as in 2018-2020.

In order to report separate emissions for the source category “Agriculture/Forestry/Fishing: Off-road vehicles and other machinery”, diesel oil consumption had to be split between stationary and mobile combustion. The Energy Statistical Yearbooks published around 1990 contained separate data for gasoil used in tractors and harvesters. Based on this information, a bit more than 60% could be allocated to mobile consumption in the early period of the time series. Considering the generally diminishing role of liquid fuels in stationary combustion, it is assumed that after 2001 all gasoil allocated to agriculture in the energy statistics has been used for mobile off-road machinery.

To be consistent with the air pollutant inventory submitted under the CLRTAP, the methodology for off-road vehicles and other machinery used in agriculture and forestry was changed for the 2017 submission, and the Tier 2 method from the 2019 EMEP/EEA Guidebook was implemented. This method classifies the used equipment into the fuel types and layers of engine technology. The engine technology layers are stratified according to the EU emission legislation stages, and three additional layers are added to cover the emissions from engines prior to the first EU legislation stages. The used layers are as follows: <1981; 1981-1990; 1991-Stage I; Stage I; Stage II; Stage IIIA; Stage IIIB; Stage IV; Stage V. The penetration of the new technology is taken into account in the form of split (%) of total fuel consumption per engine age (irrespective of inventory year) as it can be seen for diesel-fueled non-road machinery in Table 3-3 in the Guidebook. As domestic information on stock of agricultural machinery indicated a somewhat slower penetration of new technology (as in Denmark), original data in Table 3-3 have been modified as follows:

Table 3.2.8.2 *Used values for the split (%) of total fuel consumption per engine age (irrespective of inventory year) for diesel-fueled non-road machinery in Agriculture*

Engine age	USED	ORIGINAL in Table 3-3
0	4	8
1	4	7.6
2	4	7.2
3	4	6.79
4	6	6.39
5	6	5.99
6	6	5.59
7	6	5.18
8	6	4.78
9	6	4.38
10	6	3.98
11	4	3.57
12	3	3.17
13	3	2.77
14	3	2.37
15	3	1.97
16	3	1.9
17	3	1.83
18	3	1.76
19	3	1.69
20	3	1.62
21	2	1.55
22	1	1.48
23	1	1.41
24	1	1.34
25	1	1.28
26	1	1.21

27	1	1.14
28	1	1.07
29	2	1

As recommended by the ERT, non-CO₂ emissions were calculated separately for forestry. Separate activity data (gasoil consumption) was provided by the energy statistics provider for the first time only for 2016 therefore we allocated the same share (6%) of all gasoil consumption in this source category to forestry in previous years. As on average, this fuel consumption does not represent a significant amount (around 20 kt), the T1 method with default emission factor from the EMEP/EEA Guidebook seems appropriate here.

Emission factors

Default emission factors for CO₂ are used for liquid and for most of the solid fuels. The only exception is the residential lignite emission factor, which is the same as described under Energy Industries, because power plants that report measured carbon content of lignite, sell directly to residential consumers, too.

Since (almost) the entire quantity of liquid fuels used in residential combustion is LPG and the majority of institutional uses are also based on LPG, the IEF factor for CO₂ is very low.

For non-CO₂ emissions, default emission factors were applied (except for the category 1A4cii as described above).

3.2.8.3 Uncertainties and time-series consistency

We assume that the uncertainty of the fuel consumption data, especially biomass, in the Other Sector is higher than in case of industrial processes because such data are more difficult to collect and verify. Considering the above, the estimated uncertainty of the energy consumption data is ± 5 -20%.

The estimated specific uncertainty for CO₂ is 2-7%. The uncertainty of the methane factor is significantly higher (50-150%), while that of N₂O may be of an order of magnitude.

3.2.8.4 Source-specific QA/QC and verification

Comparing residential coal consumption data in the Hungarian Energy Statistical Yearbook and the IEA/Eurostat statistics, large discrepancies in NCV were found for the years before 1999. After discussing this issue with the energy statistics provider, the higher values from the domestic publication were kept.

3.2.8.5 Source-specific recalculations

As a general practice, the latest available energy statistics has been used for the calculations considering all modifications. This time, the IEA/Eurostat Annual Questionnaires as submitted in November 2021 were used.

Reallocation of autoproducers had an effect, especially in the 1.A.4.a Commercial/Institutional source category and to a lesser extent in 1.A.4.c Agriculture/Forestry/Fishing.

As a result, GHG emissions increased by 23.0 kt CO₂-eq (or 0.0% of total emissions) in 2019, and by 0.0 kt in the base year.

3.2.8.6 Source-specific planned improvements

None.

3.2.9 Other (CRF sector 1A5)

Following a recommendation of the EU ESD Review, we have included the first broad estimate for emissions from military aviation into this source category (1.A.5.b. Mobile). The table below contains our first estimates which is based on flight hours. Although exact flight hours of military aircraft are confidential but we learned from different press sources (referencing the Hungarian Defense Force) that the Hungarian combat fleet had altogether 14000 flight hours (including domestic and international training operations) in the last 10 years. This would mean 1400 hours per year. Based on another source, the average of flight hours was 1756 between 2011 and 2015, higher than in previous years. For our calculations, we took the highest fuel consumption factor (per hour) from Table 3.6.7 (i.e., 3283 kg fuel per hour) which resulted in a (probably somewhat conservative) kerosene use of 5.8 kt per year. First preliminary information from the energy statistics provider indicates a kerosene consumption of 4-5 kt used by the Hungarian Defense Force which more or less verifies our estimate. Emissions are then calculated with default EFs, (i.e., 71.5 t CO₂/TJ kerosene, 0.5 kg CH₄/TJ and 2 kg N₂O/TJ) resulting in 18 kt CO₂-eq. This figure is by about 40% below the threshold of significance (i.e., 30.5 kt in 2015).

For 2016, for the first time, kerosene consumption was allocated to the „Other” category also in the energy statistics so it could directly be used as activity data.

Table 3.2.9.1 The first broad estimate of emissions from military aviation. Values for 2005 were kept constant back to the base year. From 2016, energy statistics could be used directly therefore years after 2017 are not included in this Table.

		2005	2008	2009	2010	2012	2013	2014	2015	2016	2017
Flight hours	hour/ year	1400	1044	1044	1044	1756	1756	1756	1756	NE	NE
Kerosene consumption	kt	4.60	3.43	3.43	3.43	5.76	5.76	5.76	5.76	7	8
	TJ	199.48	148.75	148.75	148.75	250.20	250.20	250.20	250.20	302.96	346.6
CO₂	kt	14.26	10.64	10.64	10.64	17.89	17.89	17.89	17.89	21.66	24.78
CH₄	kt	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002
N₂O	kt	0.0004	0.0003	0.0003	0.0003	0.0005	0.0005	0.0005	0.0005	0.0006	0.0007

In accordance with the latest energy statistics, stationary natural gas consumption is also reported in the source category 1A5a in the years 2015-2020 – with the same methodology as in 1A4a.

3.3 Fugitive emissions from solid fuels and oil and natural gas and other emissions from energy production (CRF 1.B)

3.3.1 Fugitive emissions from solid fuels - (CRF sector 1.B.1)

3.3.1.1 Source category description

Emitted gas: CH₄, CO₂

Methods: T1, T2

Emission factors: CS, D

Key sources: 1B1 Solid fuels - CH₄ – T

Category *1B1a* includes fugitive CH₄ emission released during coal mining and handling. Emissions from fuels used during these activities are calculated under the combustion part of the inventory, mostly in the source category *1A1c*.

Emissions from category *1B1b* – fugitive emissions originating from solid fuel transformation are also mostly included in sector *1A1c*. The reason is that it is not possible to separate the GHG emissions from fugitive and non-fugitive sources during coking, and there is no reference in any of the Guidelines for emission estimation methodology in category *1B*. However, in this submission, CO₂ and methane emissions from *flaring of coke oven gas* have been allocated to this source category (i.e., *1B1b*).

In Hungary, only surface coal mines are present. Although underground mining was the predominant form in the 1960's and 1970's, it eventually ceased in 2018. Drastic reduction in coal production was observed between 1987 and 1988, as well as between 1989 and 1990. Underground mining decreased in both relative and absolute terms; therefore, distribution of mined coal types underwent significant changes. After 2015 only one minor underground mine was working after the closure of the mine of the last bituminous/sub-bituminous coal fired power plant.

The significant decrease of emissions is well explainable as emissions are strongly related to activity data (production of coal mined underground). So, the fall of underground coal mining described in the paragraph before and presented in Figure 3.2.1 resulted in decreasing trend of emissions.

Please note that all the coal mined in Hungary is classified now as lignite (except for very small amounts of sub-bituminous and coking coal).




Figure 3.3.1 Trend of emissions from solid fuels and underground production of coal

3.3.1.2 Methodological issues

During the 2016 in-country review it was suggested to provide more detailed description of methodologies in categories of fugitive emissions to increase transparency. A detailed table summarizes the sources of activity data and emission factors of all subsectors which can be found in A3.1 of Annexes.

Activity data

Production data were taken from the IEA Coal statistics, where both coal types and underground and surface productions are distinguished. Following the IEA Coal classification, all the coal mined in Hungary is classified as lignite, (except for very small amounts of sub-bituminous and coking coal in the years 1990-91). Mine and coal basin level data are received from the Mining and Geological Survey of Hungary (former Mining Bureau of Hungary). In the last three years, reported underground coal production differs at the two data provider. The difference was 3 kt at the most, which can be regarded as negligible, but also the production was very low, so the uncertainty is high. Since the inventory team always receives data from the Mining and Geological Survey of Hungary (former Mining Bureau of Hungary) earlier and data are more detailed, value from this dataset was used for the calculations (e.g., for the year 2017).

Table 3.3.1 *Underground and surface coal production in Hungary*

Year	SUM Production (kt)	SUM production (kt)	Surface SUM production (kt)	Underground out of which: Mecsek basin production (kt)
1985	24042	7387	16655	2639
1985-87	23338	7198	16141	2441
1986	23129	6983	16146	2325
1987	22844	7223	15621	2360
1988	20875	5634	15241	2255
1989	20030	5883	14147	2127
1990	17830	6919	10911	1819
1991	17135	6680	10455	1760
1992	15844	7815	8029	1210
1993	14832	7588	7244	950
1994	14084	7622	6462	1030
1995	14772	7834	6938	856
1996	15259	8067	7192	882
1997	15764	8828	6936	854
1998	14668	8445	6223	813
1999	14547	8425	6122	716
2000	14033	8848	5185	753
2001	13914	9174	4740	573
2002	13027	8929	4098	726
2003	13301	9871	3430	667
2004	11242	9135	2107	259
2005	9570	8321	1249	0
2006	9952	8601	1351	0
2007	9818	8421	1397	0
2008	9404	8118	1286	0
2009	8986	8078	908	0
2010	9113	8301	812	0
2011	9555	8890	665	0
2012	9290	8527	763	0
2013	9558	8941	617	0
2014	9551	8950	601	0.011*
2015	9261	9239	22	5.687*
2016	9216	9210	6	0.748*
2017	7974	7972	2	0.789*
2018	7898	7898	0	2.083*
2019	6847	6847	0	6.095*
2020	6125	6125	0	1.513*

* Surface production in Mecsek basin

Emission factors

Table 3.2.2 shows the measured methane content of coal for the mines operating since 1985 in Hungary together with the emission factors applied and defaults of the 2006 IPCC Guidelines. Data on in-situ methane content of mines in Hungary originates from research project conducted by Regional Centre for Energy Policy Research (available at: <http://www.rekk.eu/images/stories/letoltheto/uhg-ag-vol2.pdf>) included in list of References. The results are published in USGS, 2002 (please see the Reference list). The measured data is in accordance with the classification of mines regarding risk of firedamp received every year from the Mining and Geological Survey of Hungary (former Hungarian Mining Authority), which is also based on the m³ methane/ t coal value. Based on the above-mentioned references, two different emission factors are applied for underground mines - the same as in case of previous inventory submissions. One is applied for coals from Mecsek coal basin and the other for all other underground production. The former is within the range of default average emission factor from 2006 IPCC Guidelines, the latter is well below but the difference might be explained by country specific properties.

Table 3.3.2 *In-situ CH₄ content in Hungarian mines, the emission factors used and default emission factors from 2006 IPCC Guidelines*

Coal basin	Mine	In-situ CH ₄ content (m ³ /t)	
		mine-specific value	average in basin
Mecsek coal basin	Pécsbánya – Karolina	18.26	19.5
	Vasas – Észak	20.75	
Other underground coal mines	Balinka	1.29	1.00
	Lencsehegy	0.00	
	Mány I/a	0.98	
	Márkushegy	0.93	
	Bükkábrány	0.00	
Surface coal mines	Visonta	0.00	0.00
<i>defaults from 2006 IPCC Guidelines</i>	<i>Low</i>	<i>10</i>	
	<i>Average</i>	<i>18</i>	

Generally, no emissions occur in Hungary in case of surface mining based on the above-mentioned references. The reason is that the mined Hungarian lignite is relatively young in the coalification (NCV is under 10 MJ/kg). At the end of 2014 an old surface mine in the Mecsek basin was re-opened with relatively high (20.75 m³ CH₄/t coal) in-situ methane content, but the amount of mined coal was almost negligible in the first year. In the last two years amount of mined coal in this region was very low again, but underground production was also marginal, so this mine represented significant proportion in emissions, therefore the implied emission factor changed significantly. (It should be noted that all emissions of Mecsek basin were reported in the category of underground mines in the first part of the time series but under surface mines after 2014.)

Please note that the implied emission factor is changing because the activity data in CRF is the SUM coal produced underground while the emissions are mainly related to the production in Mecsek basin

where also some recovery activity occurred. Between 2005 and 2013 there was only one operating underground mine, so implied emission factor ($0.623+0.0623=0.68541$ please see above) became steady for those years.

Recovery

In 1.B.1.a Underground coal mining category, CO₂ emissions are reported from CH₄ recovery for the years 1985-1996. In this case CO₂ emissions are not direct emissions, but it is calculated from the amount of recovered CH₄ (CH₄ combusted for energy use) as follows:

$$\text{CO}_2 \text{ emissions} = (\text{Recovered CH}_4) * 44/16$$

$$(M_{\text{CO}_2} = 44 \text{ g/mol}; M_{\text{CH}_4} = 16 \text{ g/mol})$$

The yearly amount of recovered CH₄ and the stop of the recovery (due to the closure of the mines) were communicated by the Mining and Geological Survey of Hungary (former Hungarian Office for Mining, also former Mining Bureau of Hungary).

Post-mining

For post-mining activities, the same activity data and 10% of the mining emission factor is used the same as in the case of previous inventory submissions, which is in line with the suggestion for mines using pre-drainage of 2006 IPCC Guidelines chapter 4.1.3.2.

Abandoned underground mines

Activity data

It is very hard to collect detailed data on activities performed more than 50 years before that is required by the method of 2006 IPCC Guidelines, so several assumptions are applied. Chapter 4.1.5.2 of the 2006 IPCC Guidelines states that "Abandoned mines that were considered non-gassy when they were actively mined are presumed to have negligible emissions" and no emissions are to be reported from flooded mines.

So, emissions from this subsector are not significant in Hungary as coal mines are anyway "non-gassy" (please see Table 3.2.2 above) except for Mecsek basin. In addition, abandoned mines are usually flooded with water in Hungary.

In 2019, the Mecsek Mining Resources and Extraction Nonprofit Ltd. (Bányavagyon-hasznosító Nonprofit Közhasznú Kft.) was contacted to receive information on closed but not yet flooded mines in Mecsek basin to replace our previous estimates on numbers of unflooded mines.

Table 3.3.3 Revised activity data used in 1B1a – Abandoned coal mines

	1901- 1925	1926- 1950	1951- 1975	1976- 2000
1985	1	1	3	9
1986	1	1	3	9
1987	1	1	3	9
1988	1	1	3	9

	1901- 1925	1926- 1950	1951- 1975	1976- 2000
1989	1	1	3	9
1990	1	1	3	9
1991	1	1	3	9
1992	1	1	3	9
1993	1	1	3	9
1994	1		3	9
1995	1		3	8
1996	1		3	8
1997	1			8
1998	1			8
1999	1			8
2000	1			8
2001	1			7
2002	1			6
2003	1			6
2004	1			6
2005	1			5
2006	1			5
2007	1			5
2008	1			5
2009				5
2010				5
2011				4
2012				4
2013				4
2014				4
2015				4
2016				4
2017				4
2018				4
2019				4
2020				4

Emission factors

Set of Tier 1 emission factors from 2006 IPCC Guidelines Table 4.1.6 and Equation 4.1.11 is used for the calculation of emissions. Since time-series of emission factors begins with 1990 in the mentioned table of the 2006 IPCC Guidelines, Hungary had to complete with factors back to 1985. In the 2015 and 2016 submissions, constant values of 1990 were applied for previous years, but according to the hyperbolic decline curve of real emissions of abandoned coal mines it was planned to change these factors for more appropriate ones. For the 1901–1925, 1926–1950 and 1951–1975 periods emission factors were applied using the values of the original table for the next time-interval (2009–2014). For the 1976–2000 time interval this rule couldn't be applied. After discussions with experts from the European Union, Hungary asked the authors of this chapter of the guidelines to provide the missing values. The following table (Table 3.3.4) represents the completed table highlighting the new factors.

Table 3.3.4 Emission factors for abandoned coal mines

Emission factor, Million M ³ methane /mine					
Inventory year	Time interval of mine closure				
	1901–1925	1926–1950	1951–1975	1976–2000	2001–Present
1985	0.2900	0.3610	0.5420	4.0289	NA
1986	0.2880	0.3570	0.5290	2.8881	NA
1987	0.2860	0.3530	0.5180	2.2894	NA
1988	0.2840	0.3500	0.5070	1.9148	NA
1989	0.2830	0.3460	0.4960	1.6561	NA
1990	0.2810	0.3430	0.4780	1.5610	NA
1991	0.2790	0.3400	0.4690	1.3340	NA
1992	0.2770	0.3360	0.4610	1.1830	NA
1993	0.2750	0.3330	0.4530	1.0720	NA
1994	0.2730	0.3300	0.4460	0.9880	NA
1995	0.2720	0.3270	0.4390	0.9210	NA
1996	0.2700	0.3240	0.4320	0.8650	NA
1997	0.2680	0.3220	0.4250	0.8180	NA
1998	0.2670	0.3190	0.4190	0.7780	NA
1999	0.2650	0.3160	0.4130	0.7430	NA
2000	0.2640	0.3140	0.4080	0.7130	NA
2001	0.2620	0.3110	0.4020	0.6860	5.7350
2002	0.2610	0.3080	0.3970	0.6610	2.3970
2003	0.2590	0.3060	0.3920	0.6390	1.7620
2004	0.2580	0.3040	0.3870	0.6200	1.4540
2005	0.2560	0.3010	0.3820	0.6010	1.2650
2006	0.2550	0.2990	0.3780	0.5850	1.1330
2007	0.2530	0.2970	0.3730	0.5690	1.0350
2008	0.2520	0.2950	0.3690	0.5550	0.9590
2009	0.2510	0.2930	0.3650	0.5420	0.8960
2010	0.2490	0.2900	0.3610	0.5290	0.8450
2011	0.2480	0.2880	0.3570	0.5180	0.8010
2012	0.2470	0.2860	0.3530	0.5070	0.7630
2013	0.2460	0.2840	0.3500	0.4960	0.7300
2014	0.2440	0.2830	0.3460	0.4870	0.7010
2015	0.2430	0.2810	0.3430	0.4780	0.6750
2016	0.2420	0.2790	0.3400	0.4690	0.6520
2017*	0.241	0.277	0.336	0.439	-
2018*	0.239	0.275	0.333	0.432	-
2019*	0.238	0.273	0.330	0.425	-
2020*	0.237	0.272	0.327	0.419	-

*EF values for 2017-2020 were taken from the 2019 Refinement

3.3.1.3 Uncertainties and time-series consistency

Uncertainty of activity data is estimated based on chapter 4.1.3.6 of the 2006 IPCC Guidelines. Consistency with the value used as uncertainty of activity data in other subsectors in *Energy* is also taken into account where usually also IEA Energy Statistics are applied as activity data.

Unfortunately, no uncertainty is provided for measurement data used for emission factors applied in mining subsector and Tier 1 approach is used in post-mining and abandoned coal mined subsectors. So, the uncertainty of emission factor in *1B1* is estimated to be „factor of 2” based on Table 4.1.2 of 2006 IPCC Guidelines.

	AD	EF	Combined
1B1 Solid fuels (uncertainty +/-%)	5.00	200	200.06

3.3.1.4 Source-specific QA/QC and verification

General QA/QC procedures apply.

Activity data is compared with old time-series used in previous inventory submissions (data from the Mining Bureau of Hungary) and the differences are cc.1%.

Country specific emission factors are compared to defaults of 2006 IPCC Guidelines as it is presented in Table 3.2.2 above.

3.3.1.5 Source-specific recalculations

None.

3.3.1.6 Source-specific planned improvements

None.

3.3.2 Fugitive emissions from oil and natural gas activities (CRF sector 1.B.2)

3.3.2.1 Source category description

Emitted gas: CO₂, CH₄, N₂O

Methods: T1, T2, T3

Emission factors: CS, PS

Key sources:

1B2b Natural Gas - CH₄ – L;

1B2c Venting and flaring - CO₂ – T

In category *1B2* fugitive emissions arising during exploration, production, processing, transmission and distribution, and storage of oil (*1B2a*) and natural gas (*1B2b*) are reported. In addition, GHG emissions from venting and flaring activities connected to the operations mentioned before are included in a separate subcategory (*1B2c*). In subcategory *1B2d - Other* Hungary reports fugitive CH₄ emitted during extraction of thermal water and gas and fugitive CO₂ from mining of natural CO₂ occurrence.

In the past, oil production and processing was an important sector in Hungary, but production's importance is decreasing as the reserves are running out. Gas mining shows similar tendencies, although the reduction is less intensive. At the same time, natural gas consumption significantly increased compared to the 1980s but the demand is mainly covered by import. However, the increase of natural gas consumption stopped in 2005, and a slow decreasing trend could be observed until 2014. After 2015, natural gas consumption increased again.




Figure 3.3.2 Trends of emissions in CO₂-eq from 1.B.2 by subsector (Gg CO₂-eq emissions)

3.3.2.2 Methodological issues

During the 2016 in-country review it was suggested to provide more detailed description of methodologies in this category to increase transparency. Very detailed table summarizes the sources of activity data and emission factors in 1.B.2 subsectors which can be found in A3.1 of Annexes.

Activity data

Activity data is taken from IEA Energy Statistics. Data that is not included in IEA Energy Statistics has been provided by individual companies. In Hungary, the number of companies present in oil and gas sector is very limited, so full coverage might be assured. In subsector *1.B.2.c.i - Oil Flaring*, plant specific (EU ETS) data is also used, as very detailed verified data is available on flaring in oil refineries.

Emission factors

Default emission factors from 2006 IPCC Guidelines are applied. Chapter 4.2.2.3 of Vol2 contains one set of emission factors (Table 4.2.4) for “Developed Countries” and another Table (Table 4.2.5) for “Developing Countries and Countries with Economies in Transition”.

However, Hungary was regarded as a country with economy in transition in the beginning of the 1990's, the economy underwent significant changes since then. Hungary is now part of the European Union, and there is a great change regarding the application of state-of-the-art technologies and environmental investments as well.

So, in order to reflect more the real trend, emission factors from Table 4.2.5 have been applied for the years 1985-1994 and emission factors from Table 4.2.4 have been applied from the year 1995.

EU ETS data in 1.B.2.c.i – Oil Refinery Flaring

CO₂ emissions from oil refineries of Hungary are taken from EU ETS annual emission reports and oil refinery flaring data is extrapolated for the years before 2005 using the amount of “Refinery intake” as surrogate data. In this way, full coverage and consistency within the time-series has been reached.

1B2b4 Natural Gas Transmission and storage and 1B2b5 Natural Gas Distribution.

Instead of applying the T1 methodology from the 2006 IPCC Guidelines, methane emissions from natural gas transmission, storage and distribution were re-evaluated in the previous submission - based on information on technological losses determined by experts of the Hungarian Energy and Public Utility Regulatory Authority for the purpose of cost review of the domestic natural gas systems for the current price control cycle. Specific data on fugitive natural gas losses were received separately for transmission, storage, and distribution for the years 2016-2020. For the beginning of the time series (1985-1993), default emission factors from the 2019 IPCC Refinement were taken. For the intermediate years, interpolation techniques were applied.

1B2b4 Natural Gas Transmission and storage:

Data on technological losses in the transmission system received from the Authority was directly used in the inventory (26.3-29.0 Mm³/year in the period 2016-2020). Experts of the authority used basically two sets of data available from FGSZ Natural Gas Transmission Ltd, the single Hungarian transmission

system operator (TSO) of the entire domestic natural gas transmission system: on the one hand, the annual gas purchases between 2016 and 2020 to cover the deficit/losses, and secondly the own technological use (operation of compressors, etc.). Fugitive emissions are assumed to be the difference of these two. Since gas purchases showed differences and the technological loss is essentially a stable value, fugitive losses were calculated by taking the average of the purchases in this period and deducting the own use for the given year.

Please note that the new default methane emission factors in the 2019 Refinement are between 1.29 t/Mm³ ("extensive LDAR, and around 50% or more of centrifugal compressors have dry seals") and 3.36 t/million cubic meter gas consumption ("limited LDAR or less than 50% of centrifugal compressors have dry seals"). At the same time, our current numbers indicate an IEF of 1.84 t/Mm³ for this period (2016-19) which seems in better agreement with the latest science than our previous estimate. For the remaining part of the time series, the EF has been extrapolated assuming that in 1993 the higher default emission factor was valid (i.e., 3.36 t/Mm³).

Similar interpolation methodology was applied for gas storage. Actual data on technological losses (2.0 to 3.0 million cubic meter natural gas per year) indicated an average IEF of 0.17 t CH₄/Mm³ natural gas consumption. The new default T1 emission factors in the 2019 Refinement are between 0.29 and 0.67 t/Mm³, and again, it was assumed that the higher value was representative for 1993, and the emission factors were interpolated between 1993 (0.67) and 2015 (0.17).

1B2b5 Natural Gas Distribution:

In previous submissions before 2021, we applied a T1 methodology based on utility sales with a default IPCC emission factor of 1.1 t CH₄/million m³ of gas consumption. For recent submissions, our calculation is based on country-specific information on acknowledged distribution losses used for price regulation of gas distribution system operators.

Losses in distribution networks can be analyzed in the context of the 'Delta In-Out'. The Delta In-Out represents a difference observed when comparing the measurements at the intake points with the sum of downstream measurements of final customers off-take points, within a certain period. The delta in-out originates from many different factors. Based on a paper by the Council of European Energy Regulators (CEER, 2020), the full list of delta in-out components is the following:

1. Measurements frequency
2. Measurements accuracy
3. Linepack change
4. OBA changes
5. Blow-out during maintenance
6. Leakages
7. Theft

Similar to the above, experts of the Hungarian Energy and Public Utility Regulatory Authority determined the following four elements of distribution losses (see in brackets the values for 2020):

- 1.) measurement accuracy 1 (20.5 Mm³)
- 2.) measurement accuracy 2 - due to temperature or pressure conversion issues (12.2 Mm³)
- 3.) technological losses = leakages and blow-out during maintenance (56.4 Mm³)
- 4.) theft (15.8 Mm³)

From the above, technological losses were interpreted in the inventory as fugitive methane emissions, and all other elements as "real" natural gas consumption thus the latter was added as energy consumption (and the corresponding CO₂ emissions) to the source category 1A4b. (It has to be noted

that the real amount of theft might be somewhat higher; the above figure represents the acknowledged amount which is low enough to provide an incentive for companies to further reduce this loss element.) The estimated level of the main elements of the Hungarian delta in-out are summarized in the figure below:




Figure 3.3.3 Estimated main elements of the “Delta In-Out” in the Hungarian natural gas distribution system

Coming back to technological losses which are relevant for this source category. Here, the expert determined a fixed annual value during the 2016 cost review. The two main parameters in the estimation were the proportion of low-pressure pipeline sections within the distribution network (lower pressure generates less loss) and the ratio of the total length of steel pipeline sections (higher leakage on steel pipelines). The expert, of course, compared the estimates with the values recorded by the companies. The price regulation of natural gas takes place in four-year price regulation cycles pursuant to the provisions of Act XL of 2008 on Natural Gas Supply. The current cycle commenced on January 1, 2017 and expired on December 31, 2020 therefore we kept the same value for 2020, and will report a different number for 2021 in the next submission.

As the estimation of technological losses (=methane emission) serve basically the purpose of tariff determination in the natural gas system, it is changed only by the Authority if the underlying drivers change significantly. Experience shows however, that within one cycle none of the most important parameters change significantly: neither the total pipeline length, nor the segments of the pipelines operated on low pressure, nor the distribution of pipeline material. For example, total pipeline length changed only by 1% in the last four years which is most probably well below the uncertainty of the estimation of distribution losses anyway.

The general equation used for the estimation is as follows:

$$LOSS(total) = LOSS(techjust) * K(pressure) * K(material) * Q(<20) + LOSS(techjust) * K(material) * Q(20-100) + LOSS(techjust) * K(material) * Q(>100) + LOSS(techjust) * Q(notmeasured).$$

The calculation is made separately for four user categories: for example, $Q(<20)$ stands for gas consumption (m³/year) of users with less consumption than 20 m³/h. $LOSS(techjust)$ means the

justified technological loss which has currently a value of 0.6% for measured and 0.3% for unmeasured consumption. The parameter $K(\text{pressure})$ corresponds to the experience that if a distributor operates only medium or high pressure systems, the technological loss might increase by 50% (so from 0.6% to 0.9%). In contrast, losses might decrease to 0.3% with only low pressure pipelines. So,

$$K(\text{pressure}) = 1.5 - (\text{Length}[\text{lowpress}]/\text{Length}[\text{total}]).$$

Currently, the share of low pressure pipelines is 11.3%.

With the parameter $K(\text{material})$ it is taken into account that with steel pipelines the leakage might be higher by 10% whereas with plastic pipelines by 10% lower. Hence:

$$K(\text{material}) = 0.9 + 0.2 * (\text{Length}[\text{steel}]/\text{Length}[\text{total}]).$$

The share of steel pipelines is 7.5%, a rather constant value recently.

Calculating an implied emission factor would result in a much higher value as in the 2006 IPCC Guidelines (e.g., 3.63 t CH₄/Mm³ natural gas consumption in 2019 vs 1.1 t/Mm³ in the 2006 IPCC Guidelines). So, our IEF is higher than the default IPCC - even when looking at the 2019 Refinement where the new T1 default emission factors are between 0.62 and 2.92 t/Mm³. However, the 2019 Refinement suggests that the length of distribution pipeline is thought to best reflect emissions from distribution, and if pipeline data are available, they should be applied. Calculating an IEF on the basis of pipeline length, we would get 0.44 t/km which fits the interval of the new T1 EFs of the Refinement very well (0.23-1.17 t/km). (Which also means that the new T1 emission factors from the Refinement lead to quite different emission estimates depending on whether gas consumption or pipeline length is chosen as activity data.)

From the authority, we received estimated amounts of technological losses for the years 2016-2020 therefore backward extrapolation had to be carried out. In case of distribution losses, pipeline length was used as proxy data assuming that in 1993 the higher T1 emission factor from the 2019 Refinement representing a situation with "less than 50% plastic pipelines, or limited or no leak detection and repair programs" was valid.

1.B.2.d Other Fugitive emissions

Within this subsector fugitive CH₄ from groundwater extraction and fugitive CO₂ emissions from CO₂ mining is reported. No method is available in 2006 IPCC Guidelines for these activities, so country specific data has to be applied.

In the case of groundwater extraction, Geological and Geophysical Institute of Hungary provided expert estimate for the first time in 2015 based on 278/2015 Govt. Decree on data provision of Inventory preparation. This Institute is responsible for the monitoring, authorization and research of underground waters. They provided two set of data for the years 2004-2006 and noted that one method probably underestimates CH₄ emissions, while the other overestimates them.

So, the average of the two datasets has been applied and data have been extrapolated using Annual Groundwater extraction data from HCSO (replacing the previously used Eurostat data) as surrogate data.

In the case of CO₂ mining, activity data (million m³ CO₂ mined/year) is available from the Mining and Geological Survey of Hungary (former Hungarian Office for Mining and Geology) from 1987. For the years 1985 and 1986 the data from 1987 is applied as extrapolation. Due to lack of emission factor, the

EFs for fugitive emissions from natural gas production (extraction) from Table 4.2.4 (from 1995) and 4.2.5 (between 1985 and 1994 as described above) have been applied.

3.3.2.3 Uncertainties and time-series consistency

Uncertainty values from Table 4.2.4 have been aggregated using error propagation rule for the determination of emission factor uncertainty. For the uncertainty of AD, the same value is included as in other parts of the inventory for IEA Energy Statistics.

	AD	EF	Combined
1B2aOil – CH ₄	5.00	84	88.75
1B2aOil – CO ₂	5.00	44	44.70
1B2b Natural Gas – CH ₄	5.00	276	379.38
1B2b Natural Gas – CO ₂	5.00	278	266.30
1B2c Venting and flaring – CH ₄	5.00	50	39.08
1B2c Venting and flaring – CO ₂	5.00	472	465.27
1B2c Venting and flaring – N ₂ O	5.00	546	573.28
1B2d Other - CH ₄	5.00	200	200.06
1B2d Other - CO ₂	5.00	200	200.06

3.3.2.4 Source-specific QA/QC and verification

General QA/QC procedures apply. Plant specific data is verified with data in IEA Energy Statistics or with data received from the Mining and Geological Survey of Hungary (former Hungarian Mining Authority) where appropriate.

3.3.2.5 Source-specific recalculations

Emissions from flaring in the refinery was omitted by mistake in the previous inventory.

In addition, activity data have been revised for groundwater extraction, and the same country-specific emission factor: 4.5 t CH₄/million m³ groundwater, has been applied for the whole time series.

The overall effect of the recalculation was relatively small: an increase of 34.3 kt CO₂-eq (0.0% of total emissions) in the base year, and an increase of 18.5 kt CO₂-eq (0.0% of national total) in 2019.

Also, in contrast to previous submission, to increase comparability, leakage and venting emissions from gas transmission and storage are reported separately on the basis of TABLE 4A.2.7 of the 2019 Refinement.

3.3.2.6 Source-specific planned improvements

Generally, it is planned to apply the 2019 Refinement fully for fugitive emissions.

3.4 CO₂ transport and storage (CRF 1.C)

Not applicable.

3.5 References

Bihari, P., 1998: Energetics II. – university manuscript (In Hungarian: Energetika II., kézirat), *Budapesti Műszaki Egyetem*, Budapest.

CEER Paper on Regulatory Issues Related to the 'Delta In-Out' in Distribution Networks, 2020

Energy Centre, 2011: Energy Statistics Yearbook, 2010 (In Hungarian: Energiagazdálkodási Statisztikai Évkönyv, 2010), Budapest.

IEA/Eurostat: Annual Questionnaires (2019)

Intergovernmental Panel on Climate Change (IPCC), 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, *Intergovernmental Panel on Climate Change National Greenhouse Gas Inventories Programme*, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: Institute for Global Environmental Strategies, Japan.

Hungarian Central Statistical Office (HCSO), 2014: Statistical yearbook of Hungary (In Hungarian: Magyar statisztikai évkönyv, 2015), Budapest.

Hungarian Central Statistical Office (HCSO), 2014: Infrastructural supply of settlements, 2014 (In Hungarian: A települések infrastrukturális ellátottsága, 2014). Statisztikai tükör, 2015/75.

Institute for Transport Sciences Nonprofit Ltd. (2014): Determining the road, railway, airway and water transport emissions in Hungary for the year 2013 – Final Report

Hungarian Petroleum Association (MÁSZ), 1996-2014: Annual Report

Delta Informatika Zrt, 2011: Visualization system for data on national road transport at end of 2010, Budapest.

Regional Centre for Energy Policy Research (Regionális Energiagazdasági Kutatóközpont – REKK) 2004: Projection of greenhouse gas emission in Hungary until 2012 based on economical research of significant emitters (In Hungarian: Magyarország üvegházgáz kibocsátásainak előrejelzése 2012-ig a jelentős kibocsátó ágazatok közgazdasági kutatása alapján), Budapest. .
(<http://www.rekk.eu/images/stories/letoltheto/uhg-ag-vol2.pdf>)

USGS, 2002: Coalbed Gas in Hungary—A Preliminary Report By E.R. Landis, T.J.Rohrbacher, and C.E. Barker, U.S. Geological Survey and B. Fodor and G. Gombar, Hungarian Geological Survey, Minerals Management Division

MAVIR (Hungarian Independent Transmission Operator Company Ltd.), 2015: Statistical Data of the Hungarian Power System 2014, (online version).

http://www.mavir.hu/documents/10258/45985073/VER_Stat_2015_1223MAVIR.pdf/54105c7e-fc2e-439e-9779-5e468a28f5ae

Nyers, J., 2003: Observations and investigations after Zobák-mine closing down (In Hungarian: Zobák bánya bezáráshoz kapcsolódó vizsgálatok és megfigyelések), *Bányászati és Kohászati Lapok – BÁNYÁSZAT* (Hungarian Journal of Mining and Metallurgy - Mining) 136, Vol. 3, Budapest.

Available online: <http://www.ombkenet.hu/index.php/bkl-banyaszat?id=9>

4 INDUSTRIAL PROCESSES (CRF sector 2)

Major changes compared to previous submission:

2.D.1 – Lubricant use. The number of L-category vehicles was recalculated for the period 2009-2019, thus there are minor changes in the activity data and CO₂ emissions.

2.D.3 Other – Urea based catalysts. The number of Euro 6 vehicles was recalculated for 2018 and 2019, thus minor changes occur in the activity data and CO₂ emissions.

2.D.3 Other – Indirect emissions from solvent and other product uses. In March 2020, recalculation was performed for this sector. CO₂ emissions were recalculated and corrected in the 2020 and 2021 reports, while activity data (NMVOC emission from solvent and other product uses) remained unchanged. Activity data are now corrected for the whole time period.

2.D.3 Other – Indirect emissions from solvent and other product uses. Activity data for 2D3a Domestic solvent use and for 2D3d Coating and paints for 2019 was recalculated, thus minor changes occur in the activity data and CO₂ emissions of the sector.

2.F.1 – Correction of emission data from Mobile air-conditioning (2.F.1.e) sector (0.1 kt CO₂-eq).

2.F.2 – Revision of activity data (the impact of recalculation on total emissions including LULUCF is + 0.042 %).

The overall effect of the above recalculations was almost negligible: +4.8 kt CO₂-eq in 2019.

4.1 Overview of sector

Industrial Processes sector includes emissions generated by non-combustion processes related to industrial production. Emissions from the industrial processes and the agriculture are the second largest following the energy having a similar share of 12% each (see Chapter 2).

Emissions from this category comprise the following subcategories:

- Mineral Products (CRF 2.A),
- Chemical Industry (CRF 2.B),
- Metal Production (CRF 2.C),
- Non-energy Products from Fuels and Solvent use (CRF 2.D),
- Electronics Industry (2.E),
- Consumption of Halocarbons and SF₆ (CRF 2.F) and
- Other Product Manufacture and Use (CRF 2.G).


Under Mineral Products, Hungary reports the emissions from cement production (CO₂), lime production (CO₂), limestone glass (CO₂), and other mineral products including bricks and ceramics production, mineral wool production, waste gas scrubbing and soda ash use (CO₂).

Under Chemical Industry, emissions from ammonia (CO₂), nitric acid (N₂O), and petrochemical and carbon black production (CO₂, CH₄) are reported.

Under Metal Industry, emissions from pig iron (CO₂, CH₄), steel (CO₂, CH₄) ferroalloys (CO₂), aluminium (CO₂, CF₄, C₂F₆) are taken into account. Consumption of halocarbons and SF₆ means emissions from different sources, for example: refrigeration, air conditioning equipment, foam blowing, aerosols, electrical equipment. The 2.G sector contains emissions from manufacturing and use of electrical equipment and SF₆ and N₂O use in other products (SF₆ and N₂O).

Indirect GHGs are reported in an aggregated way, but the time-series are fully consistent with CLRTAP Air Pollutants Emission Inventory reporting of Hungary.

The base year is the average of 1985–1987 for CO₂, CH₄ and N₂O, and 1995 for HFCs, PFCs and SF₆.



4.1.1. Figure: Shares of subsectors within Industrial sector (Gg, CO₂-eq)

Several subsectors within Industrial Processes sector consist of emission originating from industrial facilities that are also falling under the scope of European Union Emission Trading System (EU ETS) - Directive 2003/87/EC. EU ETS data reported by the individual operators (summed together by industrial sector) is more accurate than the use of default factors, its use in inventory preparation needs special attention due to time-series consistency problems. In the Industrial Processes sector, EU ETS data is directly used in sector 2.A.1 *Cement production*, 2.A.2 *Lime production* (since 2014 submission), 2.A.3 *Glass production*, 2.A.4.d *Other mineral (Other - Waste gas scrubbing)* and partly in 2.A.4.a *Ceramics*, 2.B *Ammonia, Nitric Acid and Petrochemical production* and 2.C.1 *Iron and Steel* sectors. Consistency is ensured by the fact that before including extrapolation, the implied emission factor is always analyzed and depending on the trend either the IEF of the last year, or the average implied emission factor of the years is applied.

In the case of indirect greenhouse gases, consistency with CLRTAP/NEC reporting has been reached since 2014 submission. The calculation method of the indirect GHG and SO₂ emissions is described in detail in the Informative Inventory Report of Hungary submitted for CLRTAP reporting, available for each year at:

http://www.ceip.at/ms/ceip_home1/ceip_home/status_reporting/

Summary of tier methods in the mineral, chemical and metal industries as well as in the non-energy products uses of the IPPU sector are given in the table below.

4.1.1. Table: Tier methods, emission factors and key sources in the IPPU sector for categories A-B-C-D

Category	Tier method				Emission factor				Key sources
	CO ₂	CH ₄	N ₂ O	PFCs	CO ₂	CH ₄	N ₂ O	PFCs	
2.A.1 Cement Production	T2, T3				CS, PS				L
2.A.2 Lime Production	T2, T3				CS, PS				T
2.A.3 Glass Production	T2, T3				CS, PS				
2.A.4 Other Process Uses of Carbonates	T2, T3				CS, PS				
2.B.1 Ammonia Production	T3				CS, PS				L, T
2.B.2 Nitric Acid Prod.			T3				PS		T
2.B.8 Petrochemical and Carbon Black Prod.	T2, T3	T1			CS, PS	D			CO ₂ – L, T
2.C.1 Iron and Steel Prod.	T3	T2			PS	D			CO ₂ – L, T
2.C.2 Ferroalloy Production	T1				D				
2.C.3 Aluminium Production	T1			T2	D			D	PFCs – T
2.D Other Products Use	T1, T2				D				

QC of completeness and allocation of CO₂ from Non-Energy Uses and other fuels used in IPPU sector

Please find in A3.2 of Annexes based on Table 1.3 in Volume 3 of the 2006 IPCC Guidelines recommended for check of completeness of non-energy use (NEU) of fuels filled in for year 2017.

4.2 Emission Trends

Total emissions estimated from industrial processes were 7733 kt CO₂-eq in 2020, or 12% of the total national emissions compared to 14% in the base year. Total sectoral emissions decreased by 49% between the base year and 2020, and increased by 0.01% between 2019 and 2020.

Greenhouse gas emissions from the industrial processes sector fluctuated slightly in the beginning of the inventory period, then a considerable decline happened: emissions reached their minimum in 1992, which was mainly due to economic crisis. Later on, emissions had been fluctuating again until 2005. Since then, emissions have been showing a decreasing tendency again until 2009 and aggregated emissions decreased by 31% between 2005 and 2016. There was a slight growth in year 2010 and 2011, but GHG emissions from industrial processes sector were again lower both in 2012 than in 2013, the latter was the absolute minimal value of the whole time-series. In 2015 emissions increased again by 12% due to higher production volumes in several subsectors. In 2017 the revival of the economy can be seen also in emission data of all industrial sectors except of 2.G. However, after this increase, emissions did not change significantly in 2018, 2019 and 2020.

Figure 4.2.1 shows the trend of GHG emissions from industrial processes by subcategories from the base year to 2020. *Chemical industry* was the most important emitter in the beginning of inventory period, especially N₂O emission from nitric acid production (for details see chapter 4.4). Between 1990 and 2005 *Chemical industry*, *Mineral industry* and *Metal production* were fluctuating around the same level. After the significant fall of emission in *Chemical industry* thanks to the N₂O abatement technology introduced in *Nitric acid production* in 2007, and the hard recession of the *Mineral industry*, *Metal production* took up the leading role. The growing tendency of *Consumption of Halocarbons and SF₆* has also stopped in 2008.




Figure 4.2.1: GHG emissions from Industry sector by subsectors (Gg CO₂-eq)

The significant decrease of emissions in the period between 1989 and 1993 is strongly represented in the above figure. The reason for that is the economic transition mentioned already in previous

chapters. In the course of transition, factories were closed down, capacity utilization was reduced, consequently the production decreased more or less drastically in each industrial sector.

Some examples:

- Iron and steel production: two out of three plants were closed down;
- Aluminium: two out of three plants were closed down in 1991 and the aluminium production stopped in 2006;
- Ferroalloys: ceased to exist (1991);
- Ammonia: four out of five plants were closed down (1987, 1991, 1992 and 2002);
- Nitric acid: three out of four plants were closed down (1988, 1991 and 1995).

The privatization was slower in the industry than in other areas of the economy. Foreign investments were made rather in medium or smaller sized enterprises than in the big companies of the Hungarian industry.

One of the reasons of temporary production decrease was the modernization process of the remaining factories which was carried out that time and which by the way lead to favourable changes of specific emission factors as well. This was the situation e.g. in the cement and limestone industry. In some cases, however, also plants having more advantageous emission factors were closed, causing unfavourable changes in the national emission factor. This was the situation e.g. in the production of nitric acid before 1995.

Since the mid-1990s, emissions by industry have been showing a fluctuating behaviour reflecting the actual demands of production in the national economy.

GHG emissions from the iron and steel producing sector decreased by 9% compared to 2019 because of the decreasing production of pig iron. Production of ammonia, urea and nitric acid increased in 2020, causing 8% increase in emissions from the chemical industry sector. In the production of mineral industry, the years-long increasing trend halted in 2020, GHG emissions from this sector were 11% lower in 2020 than in 2019. Emissions from the non-energy products from fuels and solvent use increased by 16% mainly due to the increase in the lubricant consumption.

A third of industrial emissions come from the operation of equipment containing F-gases and from use of F-gas containing products. Category 2.F.1. (Refrigeration and air-conditioning) accounts for 87% of total F-gas emissions, which is still increasing. Despite the continuous regulation of high GWP gases emission has not already stopped in this sector.

Although, charging into new equipment for some gases already is forbidden, a lot of cooling systems have been operated with gases which have higher GWP and this is the reason of this trend.

4.2.1 Emission trends by gases

The most important GHG in Industrial Processes sector is carbon dioxide, contributing 66% to total GHG emissions in this sector in 2020, followed by F-gases with 29% . CH₄ and N₂O contributed 1 and 3%, respectively (*Figure 4.2.2*).




Figure 4.2.2 Shares of gases in Industry sector (Gg CO₂-eq)

The figure below (Figure 4.2.3) shows the emissions of this sector by gases. It can be seen that in 2008, N₂O emission from Industrial Processes are 99.89% below the level of the base year and dropped by 99.44% from 2007 to 2008 which is due to the introduction of a new nitric acid plant.




Figure 4.2.3 Trend by greenhouse gases in Industry sector

4.2.2 Emission trends by sources

In the base year, the chemical subsector accounted for 44% of total industrial GHG emissions, followed by metal subsector 34%, mineral subsector 19%. In 2020, 34% of the emissions came from chemical industry, followed by 28% from product uses as ODS substitutes. Mineral industry has 17%, metal industry has 14% contribution to sectoral GHG emissions, respectively. Other product uses (containing SF₆ and N₂O) and non-energy products from fuels and solvent use have the smallest influence on the

2020 IPPU inventory with 4% and 2%, respectively. (See **Figure 4.2.1** above.) Emissions by sources and by gases appear in *Table 4.2.1* for 2020.

Table 4.2.1 Emissions of Industrial processes sector in 2019 (CO₂-eq)

	CO ₂	CH ₄	N ₂ O	HFCs	PFCs	SF ₆	Total
2. Industrial processes	5120	48	264	2189	3	109	7733
A. Mineral industry	1311	NO	NO	NO	NO	NO	1311
B. Chemical industry	2580	44	33	NO	NO	NO	2657
C. Metal industry	1109	4	NO	NO	NO	NO	1113
D. Non-energy products from fuels and solvent use	120	NO	NO	NO	NO	NO	120
E. Electronic industry	NO	NO	NO	NO	NO	NO	NO
F. Product uses as ODS substitutes	NO	NO	NO	2189	3	NO	2192
G. Other product manufacture and use	NO	NO	231	NO	NO	109	340
H. Other	NO	NO	NO	NO	NO	NO	NO

4.3 Mineral Products (CRF sector 2.A)

4.3.1 Cement Production (CRF sector 2.A.1)

4.3.1.1 Source category description

Emitted gas: CO₂

Methods: T2 (1985-2004), T3 (2005-)

Emission factors: CS, PS

Key sources: 2A1 Cement Production – CO₂ – L

CO₂ is generated during cement production in the clinker production phase:

- on the one hand, during the combustion of the fuels used,
- on the other hand, during the degradation of the limestone (CaCO₃) fed into the furnace, which occurs at around 1,300°C and results in CaO (calcium oxide) and CO₂ (calcinations).

Both dry and wet technologies may be used for the preparation of the raw clinker. Wet technology is used by one of the three cement production plants in Hungary.

In this sector the emission estimation methodologies are very similar in the case of IPCC and EU ETS (Methodology of EU ETS reporting is prescribed in 601/2012/EC EU ETS Monitoring and Reporting Regulation). From 2005, in their EU ETS Annual Report, cement producing companies report the amount and CO₂ content of the used raw material and CKD supported by monthly measurement results.

Significant decrease of emissions has occurred in this sector between 2008 and 2013, but in 2014 the trend has changed (see *Figure 4.3.1*). The decrease of emissions correlates with the decrease of activity data. Activity data is reported directly by the cement producer companies via EU ETS Annual Emission Reports since the last years and verified with the data of HCSO if the latter is available. The decrease of activity data can be explained by decrease of the production, due to the continuous recession of this industrial sector. In building industry, the recession also turned up and 2014 is the first year that brought some recovery since 2010. The producing facilities were struggling to survive, which is published also on their website and reflected in volume indices (NACE Rev.2 classes CG- 2351 Manufacture of cement - Volume index of industrial gross output). Although solely 3 cement plants have been operating since 2014 (compared to 5 in 2010), cement production has been rising steadily in every year. This increasing trend stopped in 2020, GHG emissions from this sector were 9% lower in 2020 than in 2019.




Figure 4.3.1 Trend of activity data and CO₂ emission in cement production

4.3.1.2 Methodological issues

In this category, only emissions from the production processes are determined. Gases originating from fuels are included in *Energy subsector 1.A.2*.

From 2005, emissions are estimated using Tier 3 method: according to the EU ETS directive (2003/87/EC) introduced by the European Union, the factories report their CO₂ emissions. The factories calculate their CO₂ emissions based on the amount and the CO₂ content of all of the used raw materials and not recycled CKD filtered by dust collectors in a following way: CO₂ content of raw flour multiplied by the amount of raw flour minus CO₂ content of filtered dust multiplied by the amount of filtered dust. CO₂ content is analyzed by a certified laboratory. Detailed data of carbonate composition is not needed for this method.

For the period 1985-2004, emissions are estimated using Tier 2 method, which follows Eq.2.2 (Tier2 emissions based on clinker production data) of the 2006 IPCC Guidelines, fulfilling the assumptions given on page 2.10. The emission factor is a plant specific factor calculated by averaging the measured CO₂ content of raw meals for the period 2005-2010 (in case of one plant for 2005-2013) with the assumption that every CDK was recycled to the kiln.

Production data for the whole time-series were obtained directly from the factories and from the EU Emission Trading System (*Table 4.3.1*).

The UN review (ARR) of 2016 suggested to use a good practice data splicing technique given in the 2006 IPCC Guidelines (e.g. overlap technique or surrogate data), as appropriate for Hungary's national circumstances, to fill data gaps in the time series of the CO₂ IEF for the period before 2005. The suggested calculation was implemented in 2017 submission. The methodology was described in the 2017 NIR and emission changes appeared in the 2017 CRF tables. There were no recalculations in the submissions after 2017.

Table 4.3.1 Amount of raw flour used in process, clinker and cement production (kt) in Hungary and the CO₂ emission and implied emission factor in 2.A.1 sector

	Total raw flour (kt)	Total clinker (kt)	Total cement (kt)	2A1 SUM CO ₂ (kt)	2A1 CO ₂ (kt)/per clinker (kt)
1985	5044	3098	3671	1707	0.5511
1985-87	5152	3173	3889	1745	0.5498
1986	4982	3070	3845	1687	0.5497
1987	5430	3352	4151	1839	0.5487
1988	5264	3250	3871	1785	0.5492
1989	5338	3321	3857	1813	0.5459
1990	5148	3210	3933	1751	0.5453
1991	3247	2067	2563	1102	0.5329
1992	2533	1591	2246	859	0.5397
1993	3010	1907	2521	1022	0.5359
1994	3477	2211	2795	1181	0.5341
1995	3493	2214	2875	1186	0.5356
1996	3275	2034	2745	1111	0.5460
1997	3463	2185	2806	1174	0.5373
1998	3603	2262	2995	1222	0.5401
1999	3617	2271	2979	1224	0.5393
2000	3998	2532	3348	1353	0.5344
2001	4009	2522	3452	1357	0.5379
2002	4218	2687	3504	1426	0.5306
2003	4209	2696	3565	1412	0.5237
2004	3828	2495	3267	1289	0.5168
2005	3579	2353	3364	1199	0.5096
2006	3884	2533	3723	1296	0.5116
2007	3939	2577	3485	1328	0.5153
2008	3747	2468	3570	1261	0.5107
2009	2889	1883	2808	973	0.5166
2010	2181	1433	2134	735	0.5131
2011	1672	1109	1692	564	0.5081
2012	2047	1333	1478	678	0.5091
2013	1552	1018	1364	516	0.5067
2014	1537	1095	1467	566	0.5167
2015	2015	1331	C	676	0.5076
2016			C	705	
2017			C	783	
2018			C	882	

4.3.1.3 Uncertainties and time-series consistency

Time-series of emissions is consistent using country specific emission factors before 2005 – derived from measurements reported to EU ETS (detailed description can be found in the methodological part of this section).

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for determination of AD and EF in the case of cement production:

Uncertainty		AD	EF	Combined
2A1 Cement Production	CO ₂	2.5	2.5	3.54

As the use of ETS data means the use of verified data, where carbon contents should be measured in accredited laboratory (or at least a laboratory yearly validated and inter-compared with accredited laboratory as it is prescribed in 601/2012/EC Regulation on Monitoring and Reporting in EU ETS).

4.3.1.4 Source-specific QA/QC information and verification

According to the EU ETS directive (2003/87/EC) introduced by the European Union, the factories report their CO₂ emission from 2005 on. The factories calculate their CO₂ emissions on the basis of their production data, and the analysis of raw flour, and cement kiln dust (CKD), which contains CO₂ generated from all carbonates, including MgCO₃ and other. The analysis must fulfill the strict requirements of 601/2012/EU regulation which prescribes the use of ISO17025 accredited laboratories and the minimum annual frequency of analysis (modified by the Commission Regulation (EU) 743/2014 replacing Annex VII to Regulation (EU) No 601/2012 as regards minimum frequency of analyses). In addition, the annual emission reports of the factories are verified by independent EU ETS verifiers and checked by the authority responsible for EU ETS in Hungary every year.

In its 2021 review, ERT asked why Hungary's IEF for 2.A.1 is lower than many other countries requesting for complete information on the types of carbonate inputs in the NIR. As in the EU ETS only the laboratory tested CO₂ content of the raw meal is reported, it will take longer to answer this question because data need to be collected.

Cement production data from report of factories and EU ETS are always verified with the official statistical data. Usually the difference is low but there are some years where significant difference (more than 2%) appears. Hungarian Central Statistical Office (HCSO) started to investigate this problem on our proposal. According to its finding after 2011 the difference connected to one factory and this factory was asked to check the reported data. It would be important to further analyze this problem for those years which were not covered in this project however archived individual data before 2000 is not so easily accessible at HCSO.

Clinker production data was compared to Eurostat data. Unfortunately, Eurostat has incomplete time-series. 2012 is the only year when Eurostat database corresponds to our data source.

4.3.1.5 Source-specific recalculations

None.

4.3.2 Lime Production (CRF Sector 2.A.2)

4.3.2.1 Source category description


Emitted gas: CO₂

Methods: T2 (1985-2004), T3 (2005-)

Emission factors: CS, PS

Key sources: 2A3 Lime Production – CO₂ – T

This subsector includes quicklime production by limestone heating. During heat transfer, the following reaction occurs:



Here, only CO₂ is generated according to this formula. In Hungary, high-purity limestone is processed. Raw material contains high percent of CaCO₃ (96.8% in 2018) and minor amount of MgCO₃ (2.3% in 2018). CO₂ generated by combustion processes is accounted under the *Energy sector in 1.A.2*.

During the 2012 EU Technical review, a question was raised whether the autoproduction of lime of sugar producers is included. The investigation resulted that sugar producing companies have never reported technological (originating from dissociation of limestone) emissions in the EU ETS annual emission report (as they do not have this emission source in their GHG emission permit). However, the practice is right because no technological CO₂ emissions arise from Hungarian sugar producers since all of them use Ca(OH)₂ + CO₂ precipitation technology to remove impurities. This technology is described in the sector specific IPPC BAT BREF document as well (European Commission, 2006).

“2.1.4.11.3 Description of techniques, methods and equipment

Carbonation is the introduction of the milk of lime, calcium hydroxide, and carbon dioxide gas (CO₂) into a liquid to form calcium carbonate and to precipitate and remove impurities. The effect of lime and CO₂ is the precipitation of insoluble calcium salts, the flocculation of colloidal components, the chemical degradation of other molecules such as invert sugar and amides, and the absorption of non-sugars on precipitated calcium carbonate. Lime and CO₂ are normally produced in lime kilns by the thermal dissociation of limestone.)”

In addition, Hungarian BAT reference document prepared in 2005 by Hungarian Sugar Industry Research Institute for the Ministry of Environment and Water (KVVM, 2005 - available only in Hungarian) states that CO₂ emission from lime kilns in sugar production facilities are attributable solely to fuel combustion of the lime kilns since “CO₂ originating from dissociation of limestone is rebound again into CaCO₃.” (Section 4.1.2.2.2) Fuel consumption of lime kilns are reported in *Energy sector*. Precipitated CaCO₃ is used for liming of soils (in general) reported in *Agriculture sector*.

During informal review organized by EEA in November 2015, a question was raised if it was verified that all emissions from lime production was reported and all lime was produced only in lime plants which are included in the EU ETS (except sugar production as described above). Regarding this issue industrial associations in the field of mineral industry have been looked over and no other plants have

been found. In addition, lime production data in EU ETS annual emission reports and data from the Hungarian Statistical Office was also compared and strange result has been found that usually statistics are the lower. The average difference between years 2004 and 2012 is that HCSO data on lime production is 8% lower than EU ETS data. However, EU ETS data seems to be more accurate as EU ETS annual emission reports are verified every year. So, it seems that emissions are not underestimated due to incompleteness of the sector.

Please also note that emissions from lime and dolomite used in iron and steel industry are reported in sector 2.C.1 *Iron and steel*.

4.3.2.2 Methodological issues

The amount of CO₂ generated by this subsector is reported by using plant-specific (EU ETS) emission data of companies after 2005 and using a country specific IEF for extrapolation for the years before 2005.

The country specific IEF has been created taking into account that IEFs of years between 2005 and 2012 do not show a clear trend as it is presented in the following *Figure 4.3.2*, therefore the average seems to be applicable for extrapolation for the years before 2005 in order to reach consistent time series. The average of years 2005-2012 results in 0.7388 t CO₂/t lime produced which is 5.9% lower than the stoichiometric IEF of 0.785 and it is well fitting in the IEF range 0.56-0.8 applied by other countries as presented in the 2013 Synthesis and Assessment Report of UNFCCC.




Figure 4.3.2 Trend of kt CO₂/kt lime produced IEF between years 2005 and 2012

Exact carbonate contents of the raw material and the remaining carbonate content of the products determined by accredited laboratories are used for the calculations in EU ETS Annual Emission Reports (AERs). Using EU ETS data, the emissions from the minor proportion of dolomitic lime (containing MgCO₃), impurities and the eventual presence of hydraulic lime (which has the same stoichiometric ratio as lime but has a lower CaO content and the eventual recycling of lime kiln dust are also taken into account as it is required by the 2006 IPCC Guidelines. As EU ETS data for the years 2005-2012 contains the above-mentioned corrections, also the IEF used for extrapolation contains them.

In 2018, two companies produced lime in Hungary. According to their EU ETS annual reports, raw material contained high percent of CaCO_3 and minor amount of MgCO_3 . Calculation of their reported emission is based on Equation 2.7 of 2006 IPCC Guideline (page 2.21), where fraction calcination achieved for carbonate_i (F_i) equals to 1 for both carbonates, and fraction calcination achieved for lime kiln dust equals to 1. Emission factors for the carbonates are determined by monthly laboratory measurements.

The difference between the country-specific EF and the default EF is below 1 % (see Table 4.3.2).

Table 4.3.2 Comparison of country-specific and default emission factors in case of lime production

Source of EF	Value (kt CO ₂ /kt limestone)	difference to default
Default Tier 1 EF of 2006 IPCC Guidelines (Vol3 2.3.1.2 - Table 2.4) = (CO ₂ /CaO) * CaO content = 0.785* 0.95 =)	0.7458	
HU CS EF	0.7388	-0.93%

Time-series of lime production and related emissions can be seen in Figure 4.3.3.




Figure 4.3.3: Trend of production and emissions in sector 2.A.2 Lime

4.3.2.3 Uncertainties and time-series consistency

Time-series of emissions is consistent using country specific emission factors before 2005 – derived from measurements reported to EU ETS.

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF in the case of Lime production:

Uncertainty		AD	EF	Combined
2A2 Lime Production	CO ₂	2.5	2.5	3.54

4.3.2.4 Source-specific QA/QC information and verification

According to the EU ETS directive (2003/87/EC) introduced by the European Union, the factories report their CO₂ emission from 2005 on. The factories calculate their CO₂ emissions on the basis of their consumed raw material and final product, and the analysis of chemical composition of both them.

The analysis must fulfill the strict requirements of 601/2012/EU regulation which prescribes the use of ISO17025 accredited laboratories and the minimum annual frequency of analysis (modified by the Commission Regulation (EU) 743/2014 replacing Annex VII to Regulation (EU) No 601/2012 as regards minimum frequency of analyses). In addition, the annual emission reports of the factories are verified by independent EU ETS verifiers and checked by the authority responsible for EU ETS in Hungary every year.

4.3.2.5 Source-specific recalculations

None.

4.3.2.6 Source-specific planned improvements

Reported analysis of uncertainty in measurements will be included in the NIR when it will be available for all plants.

4.3.3 Glass Production (CRF sector 2.A. 3)

4.3.3.1 Source category description

Emitted gas: CO₂

Methods: T2 (1985-2004), T3(2005-)

Emission factors: CS,PS

Key sources: None

In the case of glass production, CO₂ emission is generated by adding the carbonates (mainly soda ashes) of alkali metals (Ba, Li, Na, etc.) to the melt in the course of glass melting. Glass production is also within the scope of EU Emissions Trading System. Please note that solely process emissions are reported in this sector, as combustion emissions are included in 1.A.2.

4.3.3.2 Methodological issues

Most of the glassworks are covered by EU Emission Trading System, the quantity of CO₂ emitted from carbonates reported by them is used as emissions between 2005 and 2016 submissions, and country-specific IEFs have been created for extrapolation for the years before 2005.

In inventory submissions before 2015, one single IEF was used for all glass types. As it was suggested by the ARR of 2014, new IEFs by glass types are now applied (details in Table 4.3.3). The difference

between the default and country-specific IEFs might be further investigated, however they are within the range of Tier 2 default emission factors of the 2006 IPCC Guidelines.

Table 4.3.3 Comparison of country-specific and default IEFs in the case of glass production

source/ type of emission factor	Value (t CO ₂ / t glass)
Default T1 IEF (Eq.2.13)	0.200
Default T2 IEFs (Table 2.6)	0.03 -0.25
CS IEF - Float glass	0.156
CS IEF - Container glass	0.181
CS IEF - Speciality glass	0.154
Default IEF - Fiberglass (glass wool)	0.250
OLD HU submission IEF	0.164

Quite detailed activity data is available from HCSO, so time-series by different glass-types could have been separated for the extrapolation.

During informal review organized by EEA in November 2015, a question was raised if it was verified that all emissions from glass production was reported and all glass was produced only in plants which are included in the EU ETS. Unfortunately, in glass sector it is not possible to perform comparison of activity data from Hungarian Central Statistical Office with EU ETS data, as in this case in EU ETS Annual emission reports there is no information on production, but solely on the amount of input materials (and emissions of course). So, also in this case industrial associations in the field of mineral industry has been looked over and only very small plants with technological emissions (emissions from carbonates) have been found in addition to those covered by EU ETS. However, in the National Air Emissions Database (LAIR) at least one bigger glass producer had been found that is not covered by EU ETS. Therefore, in the case of glass sector +10% had been added in order to cover the emission of plants not covered by EU ETS since the 2016 submission.

Further investigation was carried out to cover all non-ETS plants. Unfortunately, this mentioned plant with high reported production rate has not sent any data on multiple requests until now, but it confirms, that the production data in the LAIR database is not correct (the reported amount exceeds the EU ETS capacity threshold and much higher than the reported capacity of the plant). Surprisingly our data request had positive effect on LAIR database, namely the reported 2017 production data of this plant is lower with two orders of magnitude than before and this amount fits to all capacity thresholds. So, time-series of production data of this firm was corrected. It was known that Hungary has another small glasswork – producing handmade glassware. This firm has made available the amount of input material containing carbon and also emissions from them. Knowing production data of this firm for the 2005-2013 period, implied emission factor and also an average value of them were calculated for the time-series. As both of the two mentioned glassworks produce table wares, this average IEF was applied in the emission estimation in case of missing data.

In 2016 one of the plants covered by EU ETS announced capacity reduction with which it no longer falls under the EU ETS and therefore it has no more ETS reports, but this firm still reports data on production

to the LAIR database. Implied emission factor from the EU ETS emission report of the last known year of this firm was applied in the calculation of emission for the 2016 and 2017 years.

This research pointed out that the statistics did not cover some special glass products – production was accounted elsewhere -, which are part of the ETS glass sector. During the preparation of Allocation Plan of Hungary under the EU ETS, these plants made available emission data for the 1998-2004 period which are consistent with the 2005-2017 calculations. For the missing years production data of bulbs from HCSO was used as surrogate data. As the amount of glass used to produce bulb needs further investigation, activity data was not changed in the time-series.

As the non-ETS part is much lower than it was assumed before, therefore the additional 10% of the implied emission factor for the non-ETS period was replaced by a new calculation. The calculation is based on the quantity of product manufactured (glass and other special product) and the implied emission factor derived from the ETS reports for each type of glass products.

Figure 4.3.4 below shows the complete CO₂ emission from this category.




Figure 4.3.4 Trend of CO₂ emission and Glass Production (kt)

4.3.3.3 Source-specific QA/QC information and time-series consistency and uncertainty

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF in the case of glass production:

Uncertainty		AD	EF	Combined
2A3 Glass production	CO ₂	2.5	2.5	3.54

4.3.3.4 Source-specific recalculations and verification

More detailed data request was sent to HCSO in 2017 to verify the information from LAIR database. Also glass manufacturers were asked to declare their used technology and amount of used raw material. Unfortunately, HCSO has no full production time-series in mass unit for those factories which are not covered by EU ETS. Only one glass manufacturer fulfilled data request and provided detailed dataset about the amount of used raw material containing carbonate before this submission. Emission of this plant is 0.1kt CO₂/year at the most in the 2005-2016 period, which is 0.2% at the most of the

total emission of glass production in Hungary. Another small manufacturer declared last year that this small plant does not use carbonate containing raw material, they use glass pellet for manufacturing glass. In chapter 4.3.3.2 mentioned plant which was found in LAIR database was also contacted several times. Until this submission no data was provided about its technology and activity for the 2005-2016 period.

Investigation of addition of 10% to the data reported under the EU ETS for 2005 and onwards was finished this year. (This was a recommendation of the last UNFCCC review.) The non-ETS part in glass production was recalculated for the 2005-2016 period. This research pointed out that the statistics did not cover some special glass products – production was accounted elsewhere -, which are part of the ETS glass sector. As the non-ETS part is much lower than it was assumed before, therefore the additional 10% of the implied emission factor for the non-ETS period was replaced by a new calculation. The calculation is based on the quantity of product manufactured (glass and other special product) and the implied emission factor derived from the ETS reports for each type of glass products.

This recalculation resulted positive changes in the total CO₂ eq emission of Hungary only in 1999 (0.0008%), in 2003 (0.0017%) and in 2004 (0.0025%), every other year small decrease (less than 0.01%) can be observed.

4.3.3.5 Source-specific planned improvements

None.

4.3.4 Bricks and ceramics (CRF sector 2.A.4.a Other)

4.3.4.1 Source category description

Emitted gas: CO₂

Methods: T3

Emission factors: CS, PS

Key sources: None

During manufacturing of bricks, tiles and ceramic products, CO₂ emission is generated from the degradation of carbonates in raw materials (mainly from clay) on the one hand, and from burning of materials added to bricks on the other. Please note that in present submission all the fuels (also as additives) are reallocated into 1.A.2 together with all other combustion emissions from *Bricks and Ceramics production*.

4.3.4.2 Methodological issues

Tier 3 method is used to determine emission as in case of other EU ETS sectors. Plant-specific data is reported for the years 2005-2016 and a country-specific IEF is generated for the extrapolation of emissions before 2005 based on IEFs from 2005-2013 and national statistics of produced bricks and ceramics. Also in this case, the trend of IEFs is taken into consideration, in order to decide if average

or the IEF of year 2005 represents better. In the case of bricks and ceramics, the latter is applied as the trend is decreasing.




Figure 4.3.5 Trend of plant specific IEF between 2005 and 2013 in Bricks and ceramics

In previous inventory submissions – until 2015 -, +10% of the EU ETS emissions have been added supposing that not all the factories fall within the scope of EU ETS. This supposition could not be demonstrated until now, so, it was not applied in the 2015-2017 submissions.

During the 2020 review, a question arised that carbonates contained in raw materials are not described in the NIR. Brick and ceramics producing companies (in 2018, there were 22 companies) use different types of clay and refractory mass as raw material. In EU ETS, companies report the carbon and CO₂ content of their raw material supported by measurement results of certified analytical testing laboratories. CO₂ emissions from the organic carbon and carbonate content of raw materials, reported in ETS, is calculated based on the analytical results.

During the informal review organized by EEA in November 2015, a question was raised if it was verified that all emissions was reported and all bricks, tiles and ceramics are produced only in plants which are included in the EU ETS. The Hungarian Brick Association still has no information about plants not included under EU ETS, but small factories have reported activities in this field to the LAIR database. Unfortunately, product data from individual plants reported to the HCSO is confidential. Therefore, data request was sent to HCSO in 2017 and 2018 to calculate the non-ETS part of products (for years 2005-2017) comparing the list of firms reporting bricks and ceramics production to national statistics and firms reporting to the EU ETS. It is worth to mention that emission from different type of ceramics was calculated with the appropriate implied emission factors for the non-ETS part. The investigation pointed out that also that time-series is not fully consistent because statistics do not include some categories which have only confidential time-series. Also, the preparation of activity data in tonne unit from the official available statistics (e.g. bricks given in small bricks unit) is a challenge, besides the reporting requirement has changed several times and in several categories, the categorization is not permanent, as well. The results of the recalculation summarized in the following table.

4.3.4. Table: Proportion of non-ETS products and CO₂ emission in Bricks and ceramics

Year	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Proportion of non-ETS products (%)	6	6	5	12	7	4	5	6	5	4	4	3	3	3	3	3
Prop. of non-ETS CO₂ emission (%)	7	6	5	13	7	4	6	6	5	4	4	3	3	3	3	3
Emission (kt CO₂)	285.9	293.3	281.9	266.7	92.8	81.5	75.8	66.9	61.6	59.7	66.5	85.6	95.6	99.8	105.6	76.1

4.3.4.3 Uncertainties and time-series consistency

Time-series is not fully consistent, reasons are summarized in chapter 4.3.4.2 *Methodological issues*

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF of process uses of carbonates:

Uncertainty	AD	EF	Combined
2A4 Other Process Uses of Carbonates CO ₂	2.5	2.5	3.54

4.3.4.4 Source-specific QA/QC information and verification

General QA/QC procedures apply.

4.3.4.5 Source-specific recalculations

None.

4.3.4.6 Source-specific planned improvements

Time-series inconsistency appears before 2003 when the preparation of activity data in tonne units causes difficulties (i.e., brick production is given by the number of produced bricks). Alternative sources of information are hardly available, time-series inconsistency is planned to be resolved by a mathematical method in a medium term.

4.3.5 Soda Ash Use (CRF sector 2.A.4.b)

Emitted gas: CO₂

Methods: T3

Emission factors: D

Key sources: None

4.3.5.1 Methodological issues

Carbon dioxide is released when soda ash (Na₂CO₃) is heated.

During the 2013 centralized review, the ERT recommended to compare total import-export data of soda ash in Hungary and soda ash use in glass production in order to ensure that all soda ash uses are reported. (Please note that soda ash is not produced in Hungary.) Although the difference changes year by year, the sum of 2005-2016 of total import-export is higher than the sum of soda ash used in glass industry in 2005-2016. Therefore, additional reporting of CO₂ emission arising from soda ash not used in glass industry is needed in 2.A.4.b.

Activity Data

Total import/export of soda ash

Time-series of activity data is presented in *Figure 4.3.7* and *Figure 4.3.6*. As it was recommended by the ERT, total domestic soda ash consumption has been estimated “from domestic production plus net imports data available from statistics. Last year statistics were obtained from of UNComtrade (<http://comtrade.un.org>) (imports minus exports, for disodium carbonate)”.

Both HS classification code 283620 and SITC classification code 52323 for disodium carbonate results the same time-series. Hungarian Central Statistical Office publishes import-export data from year 2003 on its website. Differences between UNComtrade data and HCSO data are below 0.007% (6(t) Na₂CO₃). To have consistent and complete time-series data request was sent to HCSO for the whole period and it was recalculated in this submission. Unfortunately, still no data is available for years before 1991 on import/export of soda ash neither in UNComtrade and EUROStat databases, nor in the database of the Hungarian Central Statistical Office. Therefore, extrapolation was needed applying volume indices of total trade presented in *Table 4.3.5* as sector-specific volume indices are available only from 1999 within the databases mentioned above.

Table 4.3.5 *Volume indices of total trade of soda ash*

		1985	1986	1987	1988	1989	1990	1991	1992
Volume indices of trade (compared to previous year)	import	0.98	0.98	1.00	0.97	1.08	0.83	1.21	0.83
	export	1.02	0.97	0.94	1.00	1.04	1.05	0.99	1.15

Source: http://www.ksh.hu/docs/hun/xstadat/xstadat_hosszu/h_qkt001.html

Determination of the amount of soda ash not used in glass industry

Comparison of total domestic soda ash consumption and soda ash used in glass industry is presented in **Table 4.3.6**. The data on Na_2CO_3 used in EU ETS glass production have been extracted from the EU ETS Annual Emission Reports of the glass producing companies.

In several years, the soda ash used in glass production is higher than the total (import-export), while in other years it is lower. We assume that this changing trend might be due the volatility of the market and the stockpile of the glass producing companies.

In order to level off negative values, average values of the years 2005-2015 are taken into consideration. The average of soda ash NOT used for glass production /year is 2180 t/year, which results 0.905 Gg CO_2 /year. The average of soda ash NOT used in glass production (=2180 t) compared to the average of total import – export (=63803 t) results 3.42%.

In other words, the difference between the SUM of total import-export of soda ash and the SUM of soda ash used in glass production is 3.42% as it is presented in **Table 4.3.6**. As the amount of soda ash used for other purposes than glass production accounts for only 3-3.5% of the total amount and can be used in a variety of applications, we do not have a plan to identification of soda ash use in other industry sectors.

Table 4.3.6 SUM of total domestic soda ash consumption and soda ash used in glass industry

SUM of 2005-2015	
Total import-export (t) Na_2CO_3	701,837
(t) Na_2CO_3 in EU ETS glass	677,856
(t) Na_2CO_3 difference	23,980
(t) Na_2CO_3 difference / (t) Total	3.42%

So, for our calculations, 3.42 % of the total import-export data of the given year is considered to be the amount of soda ash not used in glass production.

This ratio seems to be applicable for extrapolation for years before 2005 as the slope of the trend is quite small: $R^2 < 0.1$ as it is presented on the figure below (*Figure 4.3.6*).




Figure 4.3.6 Trend of soda ash not used in glass production between 2005 and 2015

The following equation is applied for the entire time series and the results are presented in the below:

Soda ash NOT used in GLASS industry in year (n) = AD of 2.A.4.b=

(Total import-export of soda ash in year (n)) *3.42%

Emission factor and CO₂ emission of sector 2.A.4.b

In 2013, ERT recommended us to calculate CO₂ emissions from other soda ash use than for glass production in category 2A4b. In 2006 IPCC Guideline, Tier 1 and Tier 2 methods for calculating CO₂ emissions from other process use of carbonates are based on limestone and dolomite, thus we chose Tier 3 method for calculating emissions from Na₂CO₃ (soda ash). The emission factor is the factor for soda ash from Table 2.1 (0.41492 tonnes CO₂/tonne carbonate). As in Hungary there is no soda ash production, activity data equals to the exported minus imported amount of soda ash minus soda ash used in glass production with the assumption that 100 percent of calcination is achieved (Tier 3 method for the choice of activity data, page 2.37, 2006 IPCC Guideline).

Figure 4.3.7 summarizes the time-series of activity data and CO₂ emissions in sector 2.A.4 Soda Ash use.




Figure 4.3.7 Trend of total domestic consumption of soda ash and CO₂ emissions in sector 2.A.4.b.

4.3.5.2 Source-specific QA/QC information and verification, uncertainties and planned improvements

General QA/QC procedures apply.

The same uncertainty values have been applied as in the case of all subcategories of 2.A.4.d Other Mineral Industry.

4.3.5.3 Source-specific recalculations

None.

4.3.6 Other Process Uses of Carbonates (CRF sector 2.A.4.d)

4.3.6.1 Source category description

Emitted gas: CO₂

Methods: T2, T3

Emission factors: CS, PS, D

Key sources: None

This subsector includes processes in which calcinations (CO₂ loss) occur as a result of heating carbonates. CO₂ emissions generated by the degradation reaction are calculated while gases from fuel combustion are included in subsector 1.A.2. In this sector limestone and dolomite use for flue gas scrubbing and process emissions from mineral wool production are included.

Situation of other possible uses of limestone and dolomite in Hungary:

- carbide production is not occurring in Hungary as far as our knowledge;
- various uses during iron and steel production are included in 2.C.1;
- emissions from carbonates during production of clay-based products are included in 2.A.4.a *Bricks and ceramics*;
- emissions from carbonates during production of glass are included in 2.A.3. *Glass*, which includes also glass wool production.

We have no information of other uses of limestone and dolomite in Hungary.

4.3.6.2 Methodological issues

Flue gas desulphurization has been carried out in one power plant since 2002 and in another one since 2004. Activity data on the use of carbonates for SO₂ scrubbing is either reported by the operators directly to the HMS or to EU ETS competent authority. In EU ETS the operators are required to report also CO₂ emission from the use of carbonate for scrubbing separately in their annual emission report since 2008. So, direct, plant-specific emission data is used in 2.A.d.i. subcategory from year 2008 and emissions are calculated using stoichiometric ratios (included in Table 2.1 of Vol3. of the 2006 IPCC Guidelines: 440 kg CO₂ / ton limestone) for the years before 2008. In 2016 six plants reported emission from flue gas desulphurization to the EU ETS.

In the case of EU ETS plant specific data, emissions are also calculated by the operators using usually the stoichiometric ratio and fraction of purity of 1 (440 kg CO₂ / ton limestone). In 2013 one operator started to analyze the exact carbonate content of the limestone used in laboratory (fulfilling the requirements of 601/2013/EU Regulation). In 2014 a new entrant of EU ETS declared that its fraction of purity is lower than the theoretical one, because part of the carbonate remains in ash. So, these plants do not use a purity fraction of 1 anymore.

Process emissions from mineral wool production are small, but it is included in order to improve completeness of the inventory. Mineral wool producers report their CO₂ emissions since 2008 under the EU ETS. So, plant-specific emissions and activity data is available for these years. However, mineral

wool production has been present in Hungary since 2001 due to EuroStat Prodcum database. Therefore, extrapolation was applied for the years 2001-2008 for the estimation of emissions. Activity data was taken from EuroStat Prodcum database and from HCSO database, and IEF of process emissions of year 2013 is applied for the extrapolation due to lack of other detailed data.

Please note that in CRF waste gas scrubbing and mineral wool are reported together under 2.A.4.d, as it is not possible to add child node in this category.

Unfortunately, activity data are also different in the two subsectors. Carbonates used for waste gas scrubbing was chosen as AD in the CRF as it is much more significant than mineral wool.

Please find the detailed time-series of activity data and emission in *Table 4.3.7* below.

Table 4.3.7 Emissions from different sources and activity data within 2.A.4.d Other Carbonate Uses

	Carbonates used for waste gas scrubbing (AD in CRF) (kt)	Emission from waste gas scrubbing (Gg CO ₂)	Mineral wool production (kt)	Emission from mineral wool production (Gg CO ₂)	Sum emission 2.A.4.d (EM in CRF) (Gg CO ₂)
1985- 2000	NO	NO	NO	NO	NO
2001	NO	NO	45.0	2.1	2.1
2002	262.6	115.5	51.0	2.3	117.9
2003	315.2	138.7	57.1	2.6	141.3
2004	388.2	170.8	58.0	2.7	173.5
2005	504.8	222.1	61.0	2.8	224.9
2006	487.2	214.3	84.2	3.8	218.2
2007	493.2	217.0	99.9	4.6	221.6
2008	467.5	205.7	61.1	2.8	208.5
2009	437.8	192.6	36.7	1.7	194.3
2010	429.9	189.2	40.6	1.9	191.0
2011	478.2	210.4	51.4	2.4	212.8
2012	466.7	205.3	46.7	2.1	207.5
2013	473.7	203.9	45.4	2.1	206.0
2014	438.6	186.9	43.3	2.7	189.5
2015	440.0	191.0	48.8	2.9	194.0
2016	432.4	187.0	48.5	3.1	189.3
2017	357.7	155.3	48.8	2.8	158.1
2018	377.6	161.1	49.5	2.5	163.6
2019	337.1	139.9	49.4	2.4	142.3
2020	270.0	116.9	50.5	1.3	118.2

4.3.6.3 Uncertainties and time-series consistency

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF of process uses of carbonates:

Uncertainty		AD	EF	Combined
2A4 Other Process Uses of Carbonates	CO ₂	2.5	2.5	3.54

4.3.6.4 Source-specific QA/QC information and verification

General QA/QC procedures apply.

4.3.6.5 Source-specific recalculations

None.

4.3.6.6 Source-specific planned improvements

Further verification of activity data for mineral wool and improvement of country-specific implied emission factor is planned.

4.4 Chemical Industry (CRF sector 2.B)

The relevant processes operated in Hungary include:

- Ammonia production
- Nitric acid production
- Production of other organic chemicals: carbon black, ethylene, dichloroethylene and formaldehyde.




Figure 4.4.1 Shares of subsectors within chemical industry

Also in the case of chemical industry, the decrease of emissions after 1990 was due to the regime change (see *Figure 4.4.1*). Several factories were closed down and the production decreased drastically. Another significant drop of emissions occurred in 2007 when a state-of-the-art N₂O abatement technology has been introduced in a nitric acid plant (Nitrogénművek Zrt., 2008).

Production of the chemical industry is fluctuating since 2010, which is reflected in the volume index of this sector's gross output (corresponding period of the previous year= 100 (per cent): 2009: 83.9; 2010: 113.7; 2011: 107.7; 2012: 99.7; 2013: 105.6; 2014: 108.6; 2015: 102.4, 2016: 98.2). It is worth to mention that production of ammonia was 10% higher in 2016 meanwhile volume index of chemical industry decreased. The difference between the trend of production of chemical industry and the trend of emissions in 2.B sector might be explained by the increasing environmental performance of the chemical plants.

It is also worth to take into consideration that the 2006 IPCC Guidelines Vol. 3 Chapter 1 suggests that:

“Combustion emissions from fuels obtained directly or indirectly from the feedstock for an IPPU process will normally be allocated to the part of the source category in which the process occurs. These source categories are normally 2B and 2C.”

Therefore, all natural gas used in chemical industry for process purposes are accounted now here. Please find below the table (*Table 4.4.1*) showing the allocation of natural gas use in IEA Energy Statistics and in the HU GHG inventory.

Table 4.4.1 *The allocation of natural gas use in IEA Energy Statistics and in the HU GHG inventory*

	Natural gas use in <i>Chemical Industry</i> in IEA Energy Statistics (TJ NCV)			Natural gas allocation in <i>Chemical Industry</i> in HU GHG Inventory (TJ NCV)						
	Energy used in 2.B sector	Non-energy use	SUM	2.B.1 Ammonia	2.B.1. Hydro-gen	2.B.1 Tail gas treat-ment Nitric Acid	2.B.8 Petroche-mical	SUM 2B	Reported in 1.A sector	SUM
1985	19919	34294	54212	35317	0	1373	1722	38411	15801	54212
BY	21461	33968	55430	34371	0	1483	1720	37574	17855	55430
1986	22205	33897	56102	33820	0	1432	1718	36970	19132	56102
1987	22260	33715	55975	33976	0	1646	1719	37341	18633	55975
1988	21373	30393	51766	30583	0	1529	1706	33817	17949	51766
1989	22578	31115	53693	31182	0	1127	1701	34010	19683	53693
1990	22831	23113	45944	24342	0	993	1640	26974	18970	45944
1991	18754	14869	33623	14687	0	702	1586	16974	16649	33623
1992	17847	10099	27946	10111	0	445	1580	12136	15810	27946
1993	16622	12871	29493	12358	0	574	1675	14607	14886	29493
1994	13573	15560	29133	14683	0	1116	1866	17665	11468	29133
1995	17537	15950	33486	14536	0	981	2061	17577	15909	33486
1996	18580	17169	35749	16223	0	1669	2092	19984	15765	35749
1997	21939	12574	34513	15555	0	1678	2274	19506	15007	34513
1998	14162	12575	26736	13779	0	1476	2401	17656	9080	26736
1999	10892	11215	22107	12299	0	1357	2324	15980	6127	22107
2000	10266	13668	23935	15865	63	1567	2616	20112	3823	23935
2001	10256	13867	24123	14288	188	1599	2651	18726	5397	24123
2002	9383	8712	18095	10167	1765	1024	2536	15493	2602	18095
2003	8097	9169	17267	10114	1948	1141	2507	15709	1557	17267
2004	6368	13454	19822	13035	2028	1235	2403	18701	1121	19822
2005	8257	15055	23312	14729	2393	1392	2781	21295	2017	23312
2006	7938	14192	22130	13856	2760	1328	2944	20888	1242	22130
2007	9031	16409	25439	15115	2973	911	3304	22303	3136	25439
2008	7937	14719	22656	11486	2919	6	3634	18045	4612	22657
2009	5391	13534	18925	12730	2568	8	2555	17861	1063	18924
2010	6538	15149	21686	13699	2644	9	3128	19480	2207	21687
2011	6579	17134	23713	15788	3603	17	2868	22276	1437	23713
2012	4467	19000	23467	14022	4409	16	3147	21594	1872	23466

2013	10059	16472*	26531*	11377	4991	9	5176	21553	4978*	26531
2014	9594	20713*	30307*	15392	5552	12	4220	25176	5130*	30307
2015	9792	18596*	28388*	12682	5622	16	4046	22366	6021*	28388
2016	10308*	20406*	30713*	14342	5773	21	3712	23848	6866*	30713
2017	10879	23322	34201	16513	6474	28	3720	26735	7469*	34204
2018	10792	20395	31187	13514	6697	28	4036	24275	6909	31184
2019	10893*	20763	31656*	13690	6967	32	4122	24811	7021	31832
2020	11118	22996	34114	16706	6367	46	3549	26668	7546	34214

* Recalculated IEA values

4.4.1 Ammonia Production (CRF sector 2.B.1)

4.4.1.1 Source category description

Emitted gas: CO₂

Methods: T3

Emission factors: CS, PS

Key sources: 2B1 Ammonia Production – CO₂ – L, T

Ammonia (NH₃) production in Hungary uses natural gas. In the case of ammonia manufacture, natural gas provides both feedstock and fuel, whose carbon content is released in the form of carbon dioxide.

The same process occurs in the case of hydrogen production and the treatment of tail gas with natural gas in nitric acid plants.

In Hungary, the significant part of hydrogen synthesized is used also for ammonia production, but the hydrogen plant is operated by another company. The share of hydrogen/nitrogen-based ammonia production within all ammonia production has been about only 5%. 95% of ammonia production has been “traditional” natural gas based. In the last few years “traditional” ammonia production varies between 12% (in 2017) and 16% (in 2015). As in the new CRF Reporter Software there is no possibility to report CO₂ in sector 2.B.2, the emission from tail gas treatment with natural gas is reported here. The process (emission factor) is anyway the same as in the case of ammonia and hydrogen production.

In 2017 UNFCCC ERT disapproved of allocation of hydrogen production and tail gas treatment with natural gas of nitric acid production at 2B1. Hungary explained that only a small part (in 2016 less than 5%) of hydrogen production is NOT used for ammonia production, but the hydrogen production facilities are individual companies. It could be allocated to 2B10, but then both categories (2B1 and 2B10) will be confidential, and IEF of ammonia production will be very low. Tail gas treatment could also be moved to 2B10. However, the CO₂ emission from this source is very small (0.88 kt was in 2015) and with present methodology it can be easily compared the plant’s whole emission reported to EU ETS, also the total fuel consumption can be easily allocated from energy sector. So, for practical reasons (increasing the transparency in the national system) these categories are reported here.

The 2006 IPCC Guidelines requires also subtraction of amount of CO₂ emitted from ammonia production but used for urea production (and the reporting of urea used in agriculture and as catalyst

in vehicles). In the 2020 review, a revised estimation of CO₂ recovery for years 2005 and 2016-2018 was proposed and accepted by ESD. CO₂ recovery was recalculated based on the exported amounts of urea and the stoichiometric ratio of CO₂ to urea. The recalculation was performed for the whole time period resulted in recalculations of CO₂ emissions from 2.B.1 ammonia production. Moreover, urea balance was calculated taking into account urea imports and exports and considering urea-based CO₂ emissions reported in CRF 3H and 2D3. Export, import, production and stock change of urea are in balance in the long run. A detailed table of trade, production and consumption of urea is presented in Chapter 4.4.1.4.

4.4.1.2 Methodological issues

CO₂ emission from ammonia production is reported using Tier 3 methodology from the 2006 IPCC Guidelines. The Tier 3 method requires total fuel requirement (SUM TFR_i), which has been available from the reporting of the plants. In the 2020 review the ERT suggested to use country/plant specific carbon content factor (CCF) in this sector. Default CCF was replaced by a newly calculated country specific CCF for the years 1985-2006, and plant specific CCF values reported by the producers are used from 2007 onward. The country-specific CCF for 1985–2006 was calculated by taking the average of the CCFs for 2007–2019 because there was no significant trend in the CCF for these years. CO₂ emissions were recalculated for the whole time series. *Table 4.4.2.* presents the recalculated time-series of the different emission sources and the recovery. In 2018, the production of the urea plant fell to a quarter of that of the previous year and was almost entirely exported, thus the CO₂ recovery dropped drastically.

Table 4.4.2. CO₂ emission from the different sources in 2.B.1 sector

	Ammonia production with urea recovery	CO ₂ recovery in urea production	Hydrogen production	Tail gas treatment by nitric acid production
	Gg CO ₂	Gg CO ₂	Gg CO ₂	Gg CO ₂
1985	1719.78	244.22	NO	76.35
BY	1632.15	279.25	NO	82.50
1986	1599.51	281.26	NO	79.61
1987	1577.16	312.26	NO	91.54
1988	1361.88	338.86	NO	85.01
1989	1400.38	333.71	NO	62.65
1990	1144.99	208.67	NO	55.20
1991	719.31	97.43	NO	39.02
1992	492.36	69.93	NO	24.73
1993	620.18	67.08	NO	31.94
1994	754.99	61.54	NO	62.07
1995	746.38	61.96	NO	54.54
1996	842.41	59.78	NO	92.84
1997	803.12	61.89	NO	93.30
1998	698.13	68.17	NO	82.06
1999	610.62	73.34	NO	75.49
2000	803.46	78.81	3.52	87.16

2001	710.56	84.00	10.47	88.91
2002	472.85	92.55	98.17	56.97
2003	464.89	97.53	108.30	63.45
2004	614.28	110.60	112.76	68.67
2005	702.78	116.34	133.05	77.41
2006	659.54	111.01	153.50	73.86
2007	726.33	111.36	164.75	50.47
2008	561.75	74.12	161.59	0.31
2009	611.91	94.48	142.50	0.46
2010	657.54	101.35	146.50	0.47
2011	771.46	104.11	199.83	0.93
2012	704.07	75.82	245.24	0.91
2013	538.26	95.62	278.10	0.49
2014	748.60	108.67	309.23	0.69
2015	633.01	77.74	315.10	0.87
2016	714.44	87.34	322.72	1.16
2017	835.67	86.39	361.50	1.56
2018	729.57	26.59	374.72	1.58
2019	692.03	69.30	387.45	1.75
2020	880.69	48.33	356.99	2.56

CO₂ recovery for urea production occurs only in one plant, which has provided data on the quantity recovered in since 2013 onwards. In addition, it has provided data on the exported quantity and the share of the different sectors in which urea is used domestically. For the years before 2013 extrapolation was applied using domestic urea use as surrogate data. Time-series of ammonia production and emitted CO₂ are presented in *Figure 4.4.2*.




Figure 4.4.2 Trend of production of ammonia and CO₂ emissions

Please note that in HU CRF submissions the activity data is the sum of natural gas used for ammonia and hydrogen production and for nitric acid flue gas scrubbing (TJ NCV). HU IEF calculated based on ammonia production is almost the same value which is provided in the 2006 IPCC Guidelines for “modern plants in Europe”. The continuous decrease of implied emission factor in *Ammonia production* might be attributed to the fact, that obsolete technologies are abandoned. The existing factories have invested in several modernization and energy rationalization projects in recent years, which improved environmental performance and resulted in decrease of emissions per unit of ammonia produced.

Several environmental investments are listed on the public website of the company responsible for most of the production too, which explains the decrease of implied emission factor. The energy rationalization projects are for example:

2002 Ammonia Plant (expansion turbine) to utilize the pressure energy of the natural gas coming in pipeline to generate electricity

2003 Ammonia Plant (natural gas saturation) to reduce natural gas consumption

2005 Ammonia Plant Modernization of gas compressor

(http://www.nitrogen.hu/nat/index.php?option=com_content&view=article&id=122%3Akoernyezetvedelmi-beruhazasok&catid=9%3Akoernyezetvedelem&Itemid=19&lang=en)

From 2013, the extension of the scope of EU ETS also to ammonia production has been an incentive for further energy rationalization.

CO₂ emissions from *Hydrogen production* and *Tail Gas treatment with Natural Gas* are reported using direct plant specific data. Companies provided data on quantity of natural gas used for the whole time-series. Default CCF was replaced by a newly calculated country specific CCF for the years 1985-2006, and plant specific CCF values reported by the producers are used from 2007 onward.

The main producer of hydrogen in Hungary uses state-of-the-art technology, when CO₂ emitted by hydrogen production is recovered for the industrial production of CO gas. However, the amount of CO₂ recovered is not reported (not subtracted) in HU GHG Inventory at the moment, as the CO produced might be regarded as short term storage. So, all CO₂ is accounted for in the case of GHG inventory which causes a difference in the consistency check with EU ETS emissions (as the operator is also falling within the scope of EU ETS). The other reason of the difference in this subsector is that not all the hydrogen producers are required to report within the framework of EU ETS.

Please find the time-series of emissions from hydrogen production and tail gas treatment with natural gas in nitric acid production in the Table 4.4.2. above.

The quantities of natural gas used for the different processes have been compared with IEA Energy Statistics, as it is presented in Table 4.4.1.

4.4.1.3 Uncertainties and time-series consistency

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF in the case of *Ammonia Production*:

Uncertainty		AD	EF	Combined
2B1 Ammonia Production	CO ₂	5	5	7.07

4.4.1.4 Source-specific QA/QC information and verification

General QA/QC procedures apply. The quality and reliability of the emission data were greatly improved by using production data obtained directly from the factories. In 2013 data provided by the factory Nitrogénművek (responsible for the 90-95% of Hungarian production) was fully reviewed and the time-series has been affirmed. The decreasing IEF of CO₂ is also verified and the result is described in the methodological issues subchapter above (4.4.1.2).

Consistency with IEA Energy Statistics was checked and it is presented in *Table 4.4.1*.

As planned improvement from previous years' inventory database from ammonia production received from firms was compared with data of HCSO also obtained from firms. It became clear that national statistics include not only liquid ammonia – which was produced from natural gas or hydrogen - but also ammonia solution in water. Aqua ammonia can be produced by solution of liquid or anhydrous ammonia in water or by ammonia containing waste gas treatment with water, both processes do not release CO₂. Comparing the time-series only for anhydrous ammonia (for years 2005-2015) only one value was different. Request was sent for clarification to the company concerned and it was confirmed that value in the inventory database is correct.

In Hungary, only one plant produces urea. Produced amount and the ratio of exported and domestically sold urea are provided by the plant. According to this, the major part of domestic quantity is used as fertilizer in agriculture, other minor parts are used in AdBlue production, waste gas scrubbing and other industrial production. Annual amount of used AdBlue is calculated based on diesel fuel consumption in CRF 2D3, while urea consumption is derived from the sales statistics by products reported annually by the Agricultural Economics Research Institute in CRF 3H. Note, that the sales of a given year can be performed from the stocks, therefore the consumption and production do not belong to the same year in every case. Additional imported amounts of urea are involved in both activities. Emissions from urea-based waste gas scrubbing is well below the significance threshold, while urea-based CO₂ emissions are not calculated in other industrial sectors, therefore recovery is not needed. Export, import, production and stock change of urea are in balance in the long run. *Table 4.4.3* shows the trade, production and consumption amount of urea between 2010 and 2019. Trade values are from HCSO, the national statistical office. Values presented in 3H, 2D3 and 2A4 rows are calculated in CRF. Other industrial values are provided by the urea plant. In the 10-year-long period, the ratio of supply and demand is -4%.

Table 4.4.3. Imported, exported, produced and consumed amounts of urea

urea [t]		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
domestic usage											
3H - agriculture	t	63 051	79 007	75 725	107 834	101 612	132 875	134 426	154 403	152 346	158 106
2D3 - urea based catalysts	t	4 437	7 908	11 867	16 287	20 105	28 234	30 020	33 298	36 643	40 542
2A4 - waste gas scrubbing	t								2 313	2 118	1 446
other industrial	t	5 976	19 320	19 029	11 737	11 223	8 370	13 367	11 380	45	9 228
trade											
import	t	41 745	29 068	49 761	67 833	70 627	91 348	107 992	118 411	134 940	130 410
export	t	117 794	106 997	102 952	96 621	125 953	48 431	87 616	46 295	43 490	45 754
production	t	146 883	166 843	132 339	147 421	164 475	115 697	134 246	130 438	36 310	104 269
supply	t	70 833	88 914	79 148	118 633	109 148	158 615	154 622	202 554	127 760	188 925
demand	t	70 469	100 897	98 610	124 864	119 369	150 421	157 550	178 918	166 418	181 956
ratio of supply and demand		1%	-12%	-20%	-5%	-9%	5%	-2%	13%	-23%	4%

4.4.1.5 Source-specific recalculations

The 2006 IPCC Guidelines requires also subtraction of amount of CO₂ emitted from ammonia production but used for urea production (and the reporting of urea used in agriculture and as catalyst in vehicles). In the 2020 review, a revised estimation of CO₂ recovery for years 2005 and 2016-2018 was proposed and accepted by ESD. CO₂ recovery was recalculated based on the exported amounts of urea and the stoichiometric ratio of CO₂ to urea. The recalculation was performed for the whole time period resulted in recalculations of CO₂ emissions from 2.B.1 ammonia production.

In the 2020 review the ERT suggested to use country/plant specific carbon content factor (CCF) in this sector. Default CCF was replaced by a newly calculated country specific CCF for the years 1985-2006, and plant specific CCF values reported by the producers are used from 2007 onward. CO₂ emissions were recalculated for the whole time series.

4.4.1.6 Source-specific planned improvements

None.

4.4.2 Nitric Acid Production (CRF sector 2.B.2)

4.4.2.1 Source category description

Emitted gas: N₂O, (CO₂)

Methods: T3

Emission factors: PS

Key sources: 2B2 Nitric Acid Production - N₂O - T

Nitric acid (HNO₃) is produced by oxidizing ammonia. The process tail gas contains N₂O and NO_x. In order to control the emissions, the latter is reduced to nitrogen using natural gas and the carbon content of the natural gas is released in the form of carbon dioxide.

In 1985, 3 plants operated with 9 units. Among the old factories using obsolete technologies, one was abandoned in 1988, another in 1991, and a third in 1995. Until 2007 two production lines were operated in the country – the older one was established in 1975 and used GIAP technology which

consists of four units with four different factors. These four units represented the major part (about 80%) of the production volume. Emissions from this process were measured from 2004. The other existing technology represented only 20% and had been operational since 1984 (combined acid factory producing diluted and concentrated nitric acid). *Figure 4.4.3* below shows the operating nitric acid plants since 1985.




Figure 4.4.3 Nitric acid plants in Hungary. 1985-2017

Implementation of a new and more advanced production technology in Nitrogénművek plant was started in 2005, in the framework of a joint implementation project (one of the flexible mechanisms facilitated by the Kyoto Protocol), and it was installed in September 2007. At the same time the old production lines were closed down. Now a state-of-the-art technology is used, therefore drastic emission reduction is reported in this inventory.

At the end of year 2011 one of the former nitric acid plants has been restarted after renovation and its production has been increasing year by year. In 2017 a new nitric acid unit started to work at Nitrogénművek Zrt. increasing the capacity significantly.

4.4.2.2 Methodological issues

Measured emission data were not available for a long time. Therefore, during the first phase of the recalculation project, the default specific emission factor recommended by the IPCC (6 kg N₂O/t nitric acid) was used.

In 2004 an emission measurement system was installed at one of the factories and this has resulted in fundamental changes in the previously estimated values. N₂O meter is placed after the catalyst which measures emissions continuously. The regular monitoring report is based on daily average measurement data but the system is capable to provide data for shorter time period, e.g. hourly averages. The factory makes available its measured data to the inventory compiler.

The requirements of the set up and functioning of the continuous measurement system is prescribed in the IPPC (Integrated Pollution Prevention and Control) permit of the installation, as the plant is falling under the scope of the IPPC Directive (Directive 2008/1/EC). IPPC Directive in general is

implemented in the Hungarian law by the 314/2006. Government Decree and further requirements on the set up and functioning of continuous emission measurement systems is regulated by 6/2011. (I.14.) Ministerial Decree. The IPPC permit is issued, updated and enforced by the competent authority (Inspectorate for Environment and Nature).

In addition, the facility is also falling under the scope of E-PRTR Regulation (Regulation (EC) No 166/2006 concerning the establishment of a European Pollutant Release and Transfer Register). This means that on one hand data can be verified with data reported in E-PRTR (as all E-PRTR data is available to the public on <http://prtr.ec.europa.eu/FacilityLevels.aspx>), on the other hand the E-PRTR Regulation require also using internationally recognized measurement standards.

Therefore, on the basis of almost one year of experience with measurements, the calculated emission factors of the factories using different technologies were between 10 to 19 kg/t. For calculation of emissions of the oldest factory (established in the 1950's), which was abandoned in 1988, the highest value recommended by the 2000 Good Practice Guidance was used (19 kg N₂O/t). 14.5 kg/t was used as specific emission factor for the three other abandoned factories including the one which was abandoned in September 2007. For the combined factory, a value of 10 kg/t was used.

At the end of 2004, selective catalytic reduction was introduced in tail-gas treatment which led to emission reductions in the following years. This modernization means furthermore that the EFs before and after 2004 cannot be the same. The emission data of 2005 and 2006 are based on measurements. In the second half of 2005 a new measuring instrument was installed which might partly explain the difference between IEFs. The new factory of Nitrogénművek applies the EnviNO_x technology (please see further details below) consequently a drastic reduction of emission has been reached. N₂O emission from nitric acid production was decreased by 99% between base year and 2009.

Since 2013 nitric acid plants fall within the scope of EU ETS as well, so measurement of N₂O is also required and regulated by the EU ETS directive and 601/2013/EC Regulation that prescribes strict standards for the measurements and reporting.

Thus, the weighted average IEF ranges between 10.01 and 14.51 kg/t in the time-series of the years before 2007, depending on the production volume. In 2007. EF was 6.15 kg/t. 0.0425 kg/t in 2008. 0.108 kg/t in 2009. 0.0715 kg/t in 2010. Since the reopened plant is working IEF is slightly higher in recent years (0.087 kg/t in 2011. 0.113 in 2012. 0.25 kg/ t in 2013. 0.29 kg/t in 2014. 0.25kg/t in 2015 and 0.13kg/t in 2016. 0.19 kg/t in 2017), that can be explained by the increasing production volume of the other reopened factory with a less efficient tail-gas treatment. The sharp reduction in the last reported emission from the reopened plant was investigated because the IEF was halved for 2016. According to the information received from the plant in August 2015 during the summer repairs the DeN₂O catalyst was removed and during the assembly of the reactor 50% of the catalysts were replaced by new catalysts. With the new catalysts N₂O content of the flue gas reduced significantly.

The amount of carbon dioxide generated during the reduction reaction by the tail gas treatment is so low that it has no detectable effect on the inventory as a whole. Since 2004 process tail gas has been treated with ammonia, so CO₂ emissions are no longer an issue in this case. From 2007, further information about consumption of natural gas data was received from the factory. This was used in a new plant as a tail gas reducing agent. As in new CRF Reporter Software there is no possibility to report CO₂ in 2.B.2 sector (no possibility to add child node), this CO₂ is reported under 2.B.1 together with other non-energy uses of natural gas in *Chemical Industry*.

Production data were obtained from the factories (*Figure 4.4.4*).





Figure 4.4.4 Nitric acid production (kt) and N₂O emission in nitric acid subsector

EnviNO_x technology


The EnviNO_x process is usually located between the final tail gas heater and the tail gas turbine and contains two catalyst beds filled with iron zeolite catalysts operating at the same pressure and temperature and a device for addition NH₃ between the beds. In the first DeN₂O stage, the N₂O abatement is effected simply by the catalytic decomposition of N₂O into N₂ and O₂. Since NO_x content of the tail gas promotes the decomposition of N₂O, the required DeNO_x stage is arranged downstream of the DeN₂O stage.

In the second stage, NO_x reduction is carried out using NH₃ as a reducing agent similar to natural gas.

Reactions in the DeN₂O:



Reactions in the DeNO_x:



ATTACHMENT 1

PERFORMANCE TEST RUN SHEET

01-1418-600


Uhde		PERFORMANCE TEST RUN EnviNOx® NZRT						
DESIGNATION	UNIT	GUARANTEED	ACHIEVED					
			DAY 1	DAY 2	DAY 3	AVERAGE		
N ₂ O-REDUCTION IN TAIL GAS	%	min. 94 (initially)	99.63	99.64	99.63	99.63		
NO _x CONCENTRATION IN TAIL GAS DOWNSTREAM ENVI NOx® SYSTEM	ppm vol.	max. 25	5.7	5.6	5.7	5.7		
NH ₃ CONCENTRATION IN TAIL GAS DOWNSTREAM ENVI NOx® SYSTEM	ppm vol.	max. 5	Laboratory AI0808 0.19	Laboratory AI0808 0.47	Laboratory AI0808 0.57	0.41		
NH ₃ CONSUMPTION IN ENVI NOx® SYSTEM	mol NH ₃ / mol NO _x	max. 2.2	1.36	1.36	1.36	1.36		
NATURAL GAS HYDRO- CARBON CONSUMPTION IN ENVI NOx® SYSTEM	mol H.C. / mol N ₂ O	max. 0.2	0.077	0.078	0.077	0.077		

Figure 4.4.5 Presentation of performance of EnviNO_x technology

For a short description of the used technology can be found in a brochure prepared by ThyssenKrupp Industrial Solutions (see 4.12 References. ThyssenKrupp). Performance of EnviNO_x technology at Nitrogénművek Zrt. is presented on Figure 4.4.5 above.

4.4.2.3 Uncertainties and time-series consistency

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF in the case of nitric acid production:

Uncertainty		AD	EF	Combined
2B2 Nitric Acid Production	N ₂ O	7.5	7.5	10.61

4.4.2.4 Source-specific QA/QC information and verification

General QA/QC procedures apply. The data received directly from factories and the requirements of EU ETS since 2013 greatly improved the quality of data.

The significantly decreasing IEF after 2007 was verified and the results are also described in the methodological issues subchapter above. Also the significantly decreased IEF for 2016 was investigated and it is described in chapter 4.4.2.2.

4.4.2.5 Source-specific recalculations

None.

4.4.2.6 Source-specific planned improvements

None.

4.4.3 Petrochemical and Carbon Black Production (CRF sector 2.B.8)

4.4.4 Source category description

Emitted gas: CO₂, CH₄

Methods: T1, T2, T3

Emission factors: CS, D, PS

Key sources: 2B8 Petrochemical and carbon black production - CO₂ - L, T

During petrochemical production processes, mainly oil products are used as feedstock or other non-energy purposes. Most of the carbons contained in these raw materials are stored in the products too, however during the conversion processes some carbon is emitted in the form of CO₂ or CH₄.

Usually, it is very hard to distinguish the energy and non-energy uses of fuels during the complex processes in petrochemical production. Therefore, the suggestion of the 2006 IPCC Guidelines is to be followed:

„Combustion emissions from fuels obtained directly or indirectly from the feedstock for an IPPU process will normally be allocated to the part of the source category in which the process occurs.”

„If surplus methane or hydrogen from the steam cracking of naphtha is combusted within the petrochemical site for another process then the emissions are reported as emissions in IPPU 2B8. On the other hand, if the gases are passed to a nearby refinery for fuel use then the associated emissions would be reported under 1A1b. Petroleum Refining.”

In the case of Hungary, no gases are passed for fuel use out of the petrochemical companies, but all are used inside. Therefore, all emissions are reported here in 2.B.8 including the natural gas reported by the petrochemical and carbon black production companies. In addition, all oil products considered as non-energy use (NEU) in IEA Energy Statistics are considered here in sector 2.B.8, except for lubricants and paraffin waxes (reported in 2.D.1-2).

In Hungary production of ethylene dichloroethylene (DCE for the purpose of production of further petrochemical products like TDI/MDI) and carbon black are present. Ethylene is made from naphtha. Very few and well-known companies are operating in this sector in Hungary.

From 2018, CO₂ emissions have been reported in the ETS system from formaldehyde production. As the company have been producing formaldehyde from 1998, CO₂ emissions are reported calculated on a new subsector sheet 2.B.8.g.i formaldehyde production from 1998 onward.

Production of bulk organic chemicals fall within the scope of EU ETS since 2013, so the availability of detailed plant-specific data is even more improved. However. EU ETS data has been available already since 2008 for most of the sources of these companies.

4.4.4.1 Methodological issues

CO₂ emissions are reported using plant-specific data from the year 2008 using the very detailed and good quality data from reporting of companies within the EU ETS framework. The good quality of data reported under EU ETS is ensured by the strict monitoring and reporting requirements of Regulation 601/2012/EC including the obligation of the control by independent, accredited verifiers. EU ETS Annual emission reports are available for HMS for GHG Inventory preparation purposes.

2.B.8.b Ethylene production

In Hungary, only one company produces ethylene. According to its EU ETS Annual Report, during the production the company uses three different gaseous fuels. Reported CO₂ emission is calculated under the standard methodology (Commission Regulation No 601/2012, Article 24) supported by monthly laboratory measurements.

2.B.8.c ETM/VCM production

In Hungary, only one company produces VCM and TDI. For VCM natural gas, for TDI natural gas, OTD and tar are used in the production process. Reported CO₂ emission is calculated under the standard methodology (Commission Regulation No 601/2012, Article 24) supported by monthly laboratory measurements.

2.B.8.f Black carbon production

In Hungary, only one company produces black carbon using quench oil, natural gas and negligible amount of other materials. Reported CO₂ emission is calculated under the mass balance methodology (Commission Regulation No 601/2012, Article 25) supported by monthly laboratory measurements.

2.B.8.g.i Formaldehyde production

From 2018, CO₂ emissions have been reported in the ETS system from formaldehyde production. As the company have been producing formaldehyde from 1998, CO₂ emissions are reported calculated on a new subsector sheet 2.B.8.g.i formaldehyde production from 1998 onward. Reported CO₂ emission is calculated under the mass balance methodology (Commission Regulation No 601/2012, Article 25) supported by monthly laboratory measurements.

For the years before 2008 extrapolation is applied using the appropriate petrochemical production data from HCSO as surrogate data.

Please note that extrapolation of emissions is made based on quantities of petrochemical products (ethylene + DCE + carbon black), and the activity data in CRF in 2.B.8 is also kilotons of products.

Time-series of non-energy use of oil products (as raw material in petrochemical production) in IEA EnStat have been taken into account, as all emissions from these sources are excluded from *Energy* sector and reported here. So, in order to be consistent with both EU ETS reports of petrochemical companies and IEA Energy Statistics (EnStat), emissions are reported in 2.B.8 based on EU ETS reporting, and it is supposed that all non-energy use of oil from IEA EnStat and some amount of natural gas (reported by the companies) were used for the production. The related energy sources were subtracted from *Energy* sector.

In this way, IEF compared to petrochemical products is stable (see red line on *Figure 4.4.6* below), while IEF compared to NEU Oil products in IEA EnStat is changing as it is possible to see at the same figure (blue line). But this problem seems to originate from the original time-series of the NEU Oils in the IEA EnStat, as the trend of petrochemical products (t) per NEU Oils (TJ) is changing, too (see green line on the figure).

Unfortunately, it seems that the allocation of NEU Oils within the IEA EnStat is not consistent across the years, so this question is to be clarified with the authority responsible for the preparation of the energy statistics.




Figure 4.4.6 Trend of implied emission factors in 2.B.8 compared to petrochemical production and the amount of non-energy use of fuels

As EU ETS annual emission reports contain only all CO₂ emission sources. CH₄ is reported based on default Tier 1 emission factors from the 2006 IPCC Guidelines for the whole time-series.

4.4.4.2 Uncertainties and time-series consistency

Uncertainties are estimated based on the minimum requirements of EU ETS Monitoring and Reporting Regulation (601/2012/EU) for the determination of AD and EF in the case of nitric acid production:

Uncertainty		AD	EF	Combined
2B8 Petrochemical and carbon black production	CO ₂	7.5	7.5	10.61
2B8 Petrochemical and carbon black production	CH ₄	3	10	10.44

1.1.1.2 Source-specific QA/QC information and verification

General QA/QC procedures apply. Time-series in IEA EnStat production data from HCSO and the EU ETS emission reporting data have been thoroughly compared as it is described above.

1.1.1.3 Source-specific recalculations

CH₄ emission for 2016 was replaced due to calculation error.

CO₂ emission for 2017 was recalculated, because flaring reported by the ethylene producing company under the EU ETS was not included for the year 2017.

4.4.4.3 Source-specific planned improvements

Further investigation of consistency of the trend of non-energy use of oils in IEA EnStat is needed.

4.5 Fluorocarbon Production (CRF sector 2.B.9)

Fluorocarbons are not produced in Hungary.

4.6 Metal Production (CRF sector 2.C)

4.6.1 Iron and Steel Production (CRF sector 2.C.1)

4.6.1.1 Source category description

Emitted gas: CO₂, CH₄

Methods: T2, T3

Emission factors: CS, PS, D

Key sources: 2C1 Iron and Steel Production - CO₂ – L, T

In this subsector, gases emitted by the iron/steel industry (sinter, iron and steel production) are calculated. During sintering (agglomeration), a mixture of iron ore, coke or carbon and limestone are agglomerated by heat transfer to obtain a material suitable for feeding into the furnace. During iron production, coke and carbonate-containing slag-forming additives are added to the agglomerated ore, and the mixture is reduced at a high temperature. This reaction releases CO and CO₂. Therefore, CO₂ is produced from two sources during the process: 1) from fuel, which also serves as a reducing agent, and 2) from carbonate-containing slag-forming agent (limestone or dolomite). The gases arising in the blast furnace during the production of the pig iron are recovered as blast furnace gas (BFG) and used partly for energy purposes.

During steel production, the carbon content of iron is reduced from 4-5% to cca. 1%. (1% in the 2006 IPCC Guidelines). Also, this is released in form of CO₂. Basic oxygen furnace (BOF also known as LD converter) technology for production of steel uses the hot, molten pig iron with scrap iron, additives and quicklime. Electric arc furnace (EAF) technology uses mainly scrap iron with additives and the heat is provided by electric arc formed between graphite electrodes. The consumption of graphite electrodes results CO₂ emission as well.

In Hungary, all the activities connected to iron and steel industry are present: production of coke, sintering, production of pig iron and production of steel using basic oxygen furnace (BOF) and electric arc furnace (EAF) technology, too. Except for the EAF steel production, all the activities mentioned before are located in one single plant, which is however operated by different operators, so it cannot be regarded as an integrated iron and steel plant.

Processes within iron and steel production are very complex, using several fuels either for energy or for non-energy purposes. It is very hard to distinguish energy and non-energy use also in this case, so the recommendation of the 2006 IPCC Guidelines Vol. 3 Bo1.1 is followed that states:

“During these activities, emissions may occur from both the fuel combustion and industrial process stages. However, it is often impractical or impossible to report separately the two types of emissions. Accordingly, the following rule has been formulated to simplify reporting:

Combustion emissions from fuels obtained directly or indirectly from the feedstock for an IPPU process will normally be allocated to the part of the source category in which the process occurs. These source categories are normally 2B and 2C. However, if the derived fuels are transferred for combustion in another source category, the emissions should be reported in the appropriate part of Energy Sector source categories (normally 1A1 or 1A2).

Two examples may help illustrate the definition.

1. If blast furnace gas is combusted entirely within the Iron and Steel industry (whether for heating blast air, site power needs or for metal finishing operations) the associated emissions are reported in the IPPU source subcategory 2C1. If part of the gas is delivered to a nearby brick works for heat production or a main electricity producer, then the emissions are reported in source subcategories (1A2f or 1A1a).”

The example mentioned above is presented in Hungary, one part of the blast furnace gas is used in own processes, while the other part is delivered to an electricity producer. Amount of CO₂ emission from blast furnace gas which is not taken into account in 2.C.1 emission is reported in 2.C.1 as **“Recovery”** – only for transparency purposes - and it is delivered to emissions of 1.A sector.

Please find the trend of emissions and the trend of production in *Iron and Steel* sector in *Figure 4.6.1* below.




Figure 4.6.1 Trend of production and emissions in Iron and steel sector

As it is possible to observe, drastic reduction of the production occurred around 1990 and between 2012 and 2013, too, but in 2014 it seems that recession has stopped. In 2015 iron and steel production reached almost the level before the 2008 economic crisis. Metal industry (particularly in iron and steel production) realized 26% emission reduction in 2016. Amount of both pig iron and steel products

decreased, which is the outcome of unfavorable process in export markets. Steel product to building industry could also profit from quickening of housing (more than two-and-one-half times housing permission was granted in 2016 compared to the previous year) with higher production, meanwhile the total amount of steel was lower in 2016 than 2015. Emission increased by 38% in metal industry subsector in 2017, where the favourable EU export market situation and competitiveness of Dunafer Zrt. in this market resulted from the efficiency improvement measures taken by the company between 2014 and 2016. However, the rising slowed down in 2018.

4.6.1.2 Methodological issues

Earlier only the emissions from carbon content reduction of the input materials during steel production and emission from consumption of graphite electrodes (2.C.1.1 subsector) were reported within this sector and all the other emissions were included elsewhere. The actual allocation of emissions is summarized in *Table 4.6.1*. All cell comments of IE cells in CRF tables have also been updated accordingly.

Table 4.6.1 Allocation of emissions connected to Iron and steel production

IPCC sector code	Activity	Emission source	From the 2015 submission - Emission reported in
1.A.1.a	Combustion of blast furnace gas recovered from Pig Iron production	combustion	1.A.1.a
1.A.1.c	Production of coke	combustion	1.A.1.c (including coke oven gas)
1.A.2.a	Combustion needed for iron and steel production	combustion	1.A.2.a (including coke oven gas)
2.C.1.d-e	Sinter	Coke consumption during sintering	2.C.1.d-e Sinter-Pellet
		Limestone and dolomite use	
2.C.1.b	Pig Iron	Combustion	2.C.1.b Pig Iron
		Consumption of Natural gas for non-energy purposes	
		Limestone and dolomite use	
		Consumption of coke in the blast furnace (after deduction of the amount of recovered blast furnace gas delivered outside for energy production purposes)	
2.C.1.a	Steel	Reduction of carbon content (from 4% to 1 %)	2.C.1.a Steel
		Emission from graphite electrode during EAF steel production	

Emission factors

In the case of CO₂ emissions, default emission factors from the 2006 IPCC Guidelines or plant-specific emission and activity data were used which is available from both direct reporting to the inventory compiler (HMS) and from EU ETS reports. In the case of CH₄, default emission factors from the 2006 IPCC Guidelines are applied.

Activity data

Iron and steel production data were obtained from the reports of the International Iron and Steel Institute, World Steel Association (WORLDSTEEL) and the similar European agency (EUROFER).

Data on consumption of coke, natural gas, coke oven gas in the blast furnace is extracted from IEA Energy Statistics of Hungary as well as the amount of blast furnace gas (BFG) recovered and used. Amounts of limestone and dolomite, other additives including graphite electrode in EAF steel production are available either from direct reporting of the companies to HMS or from EU ETS reports since 2005. Detailed plant-specific data is available from the EU ETS annual emission reports of the companies.

However, the precise allocation among the subsectors (e.g. the amount of coke used in blast furnace or by sintering) is sometimes not available in IEA Energy statistics. In these cases, extrapolation based on the shares of subsectors in plant specific data is applied, but the sum of the subsectors is always the same as the IEA time-series. Please see the example of coke in *Table 4.6.2*.

4.6.1.3 Steel (CRF sector 2.C.1.a)

Carbon dioxide is released from carbon content of pig iron and graphite electrode of the electric arc furnace (EAF) during steel production are reported in *2.C.1.1 Steel* subsector.

During basic oxygen steel production, the carbon content of the pig iron is converted from 4% to 1%. This means that 3% of carbon content present in pig iron is emitted as CO₂ and 1% of carbon content is stored in the steel.

Carbon content of the pig iron might originate from the coke consumed in the blast furnace, iron ore and additives. In order to avoid double counting these emissions should be subtracted from subsector *2.C.1.b Pig Iron*.

The default carbon content of pig iron is: 4% (2006 IPCC Guidelines). In the case of carbon content of steel these guidelines specify it as 1%.

In the case of EAF steel production, data is available in the EU ETS emission reports on carbon content reduction during the process from 2008, therefore plant-specific data is used for the reporting of emissions from EAF steel production and extrapolation of the average of years 2008-2012 is applied using EAF steel production as surrogate data. The factor calculated as average of years 2008-2012 is 0.055 compared to the default 0.05 t CO₂/ t EAF steel.

In the case of EAF steel production the input material is usually scrap iron and other unknown material. This feedstock does contain carbon but it is not originating directly from the use of coke of the blast

furnace of the given year. So, EAF steel production is included in „Steel produced (kt)” data of the formula above, but the carbon content reduction of EAF steel production is not subtracted from 2.C.1.b.

Quicklime used in BOF furnaces is not produced on-site, as it is declared by the operator.

During the investigation of EU ETS data it was discovered, that for 2008, CO₂ emissions for EAF steel production was reported years later by one of the EAF companies under the EU ETS. As estimated 2008 emission data were used in the calculated extrapolation for 1985-2006, emission was recalculated for years 1985-2006 and 2008.

4.6.1.4 Pig Iron (CRF sector 2.C.1.b.)

Emission from use of coke, natural gas, coke oven gas (COG) and own use of blast furnace gas is reported in 2.C.1.b.

In addition, use of limestone, dolomite and other ores and additives is also included here, so the whole process of pig iron production is aimed to be reported here.

The blast furnace gas (BFG) recovered and delivered for energy production purposes is reported in 1.A sector, while the CO₂ emissions during the carbon content reduction in steel production process (originating from the coke) is reported in 2.C.1.a. One part of the blast furnace gas is used in own processes, while the other part is delivered to an electricity producer. Amount of CO₂ emission from blast furnace gas which is not taken into account in 2.C.1 emission is reported in 2.C.1 as **“Recovery”** – only for transparency purposes – and it is delivered to emissions of 1.A sector.

Time-series of coke, BFG, COG and natural gas are available in IEA EnStat, but the precise allocation is not included (or not consistent). But plant specific data on the exact allocation is available from the year 2004, therefore extrapolations are applied where needed (please see the example in above on allocation of coke consumption from IEA EnStat and within subsectors of 2.C.1). In addition, plant-specific net calorific values had been prioritized, where available.

CO₂ emissions from coke are reported using plant-specific data from 2008. For the years before 2008 extrapolations are applied using default emission factors. In the case of CH₄ emissions, emissions from natural gas, COG, limestone and dolomite use are calculated using default factors from the 2006 IPCC Guidelines. Emission from other ores and additives is reported using plant-specific data from 2007 (first year where available) and extrapolation is applied using the average of the implied emission factor of the last two years and pig iron production as surrogate data.

During the investigation of EU ETS data it was discovered that part of the blast furnace gas produced during the BOF pig iron production was not transferred offsite but was flared, therefore this quantity cannot be included in the recovery. CO₂ emissions and recovery was recalculated from 2007 onwards.

Table 4.6.2 Time-series of coke consumption (kt) in IEA EnStat and in 2.C.1

IEA EnStat (kt)			2.C.1 (kt)			
	Coke consumption in blast furnaces (Transfor-mation)	Coke consumption in <i>Iron and steel</i>	SUM IEA	2.C.1.b <i>Pig Iron</i> coke used in BF	2.C.1.d <i>Sinter</i> coke used in sinter	SUM 2.C.1
1985	1 471	209	1 680	1 565	115	1 680
B.Y.	1 447	144	1 591	1 483	109	1 591
1986	1 525	112	1 637	1 525	112	1 637
1987	1 346	111	1 457	1 358	99	1 457
1988	1 292	141	1 433	1 335	98	1 433
1989	1 129	102	1 231	1 147	84	1 231
1990	1 040	16	1 056	984	72	1 056
1991	737	115	852	794	58	852
1992	656	158	814	758	56	814
1993	778	43	821	765	56	821
1994	891	50	941	877	64	941
1995	870	43	913	851	62	913
1996	815	63	878	818	60	878
1997	562	83	645	601	44	645
1998	597	100	697	649	48	697
1999	590	157	747	696	51	747
2000	639	108	747	696	51	747
2001	566	126	692	645	47	692
2002	606	112	718	669	49	718
2003	549	161	710	662	48	710
2004	570	170	740	689	51	740
2005	596	123	719	671	48	719
2006	601	111	712	665	47	712
2007	620	135	755	709	46	755
2008	599	123	722	668	54	722
2009	593	38	631	599	41	640
2010	686	52	738	689	49	738
2011	687	53	740	685	55	740
2012	661	49	710	661	49	710
2013	356	27	383	347	36	383
2014	462	30	492	443	49	492
2015	635	50	685	639	46	685
2016	452	36	488	447	41	488
2017	655	51	706	662	44	706
2018	662	52	714	666	48	714
2019	592	46	638	589	49	638

2020	535	42	577	533	44	577
------	-----	----	-----	-----	----	-----

Please find the activity data used and the resulting emissions in 2.C.1.b Pig Iron subsector in Table 4.6.3 below.

Table 4.6.3 Trend of activity data and emissions in 2.C.1 b Pig Iron subsector

	Pig Iron produced	Coke consumption in BF	BFG recovered and used outside for energy	COG used in BF	NatGas Consump- tion in BF	Lime- stone used in BF	Dolomite used in BF	SUM CO ₂ Emission in 2.C.1.b	SUM CH ₄ emission in 2.C.1.b
	kt	TJ	TJ	TJ	TJ	kt	kt	kt	kt
1985	2 095	42 475	2 189	283	3 948	57	2	3898.98	0.43
B.Y.	2 085	40 234	2 364	436	3 930	57	2	3616.83	0.41
1986	2 054	41 388	2 690	411	3 871	56	2	3648.10	0.42
1987	2 107	36 837	2 214	616	3 971	58	2	3303.39	0.37
1988	2 093	36 231	1 868	679	3 945	57	2	3327.66	0.37
1989	1 954	31 123	1 443	694	3 683	53	2	2897.18	0.32
1990	1 697	26 699	1 446	629	3 198	46	2	2426.70	0.27
1991	1 314	21 541	1 090	656	2 476	36	2	2005.57	0.22
1992	1 176	20 580	1 503	618	2 216	32	1	1815.57	0.21
1993	1 407	20 783	1 615	576	2 652	38	2	1812.03	0.21
1994	1 595	23 820	1 936	794	3 006	44	2	2062.99	0.24
1995	1 515	23 111	1 674	876	2 855	41	2	2061.35	0.23
1996	1 496	22 226	1 553	746	2 820	41	2	1985.46	0.23
1997	1 140	17 470	1 972	746	2 149	31	1	1375.76	0.18
1998	1 259	19 204	3 859	716	2 373	34	1	1077.06	0.20
1999	1 310	20 649	3 939	614	2 469	36	2	1203.20	0.21
2000	1 340	20 719	4 141	673	2 526	37	2	1160.52	0.21
2001	1 226	19 194	3 685	273	2 311	33	1	1071.70	0.19
2002	1 335	19 946	3 800	118	2 516	36	2	1137.44	0.20
2003	1 333	19 535	3 711	311	2 512	36	2	1125.97	0.20
2004	1 351	20 237	4 236	358	2 952	14	0	1080.08	0.21
2005	1 338	19 589	4 063	369	2 453	25	0	1034.86	0.20
2006	1 336	19 619	4 065	906	2 186	20	0	1041.58	0.20
2007	1 394	20 881	4460*	1 031	1 869	24	0	1008.28*	0.21
2008	1 289	19 685	4310*	774	1 477	28	0	954.49*	0.20
2009	1 050	17 743	3947*	300	605	36	0	812.49*	0.18
2010	1 325	20 618	5238	927	1 757	66	0	859.94*	0.21
2011	1 315	20 354	4692*	831	1 692	61	0	886.96*	0.21
2012	1 228	19 501	4561*	744	490	48	4	762.63*	0.20
2013	628	10 121	2489*	380	530	14	5	444.95*	0.10

2014	801	13067	3494*	599	816	38	0.8	532.22*	0.13
2015	1247	18941	4844*	1130	1584	30	0	788.34*	0.19
2016	863	13424	3250*	688	699	22	0.02	558.36*	0.14
2017	1313	19556	4928*	1018	1590	27	0	819.67*	0.20
2018	1355	19654	4885	1207	1916	29	0	895.71	0.20
2019	1151	17266	4227	892	1373	21	0	785.30	0.17
2020	930	15628	3995	809	1117	52	0	701.71	0.16

* Recalculated in the 2020 submission

4.6.1.5 Sinter and pellet (CRF sector 2.C.1.d-e.)

Amount of sinter or pellet produced is not available. However, the amount of coke and natural gas, limestone and dolomite and other ores and additives used during sintering is available from direct reporting of the company from the year 2004. For the years before 2004 activity data was extrapolated knowing material consumption and pig iron production rate after 2004, emission extrapolation is applied using the implied emission factor of the last year or the average of the years available depending on the trend of the IEF.

CO₂ emissions from coke, natural gas, limestone, dolomite and "Other ores and additives" use are reported.

In addition, CH₄ is estimated using default EF for coke (10 kg/ TJ coke), due to lack of data on the amount of sinter or pellet produced. CH₄ is reported from coke combustion in sinter plant and from 2016 submission also the CH₄ emission from natural gas use in sinter plant calculated with default EF has been included due to the recommendation received during the informal review organised by the EU in November 2015. However, this recalculation causes less than 0.01 Gg increase in emissions.

In recent years coke and natural gas consumption are quite stable, meanwhile limestone and dolomite use has strong fluctuation which can be seen also in the emission data.

Please find the activity data used and the resulting emissions in 2.C.1.d-e Sinter subsector in the table below (Table 4.6.4).

4.6.1.6 Uncertainties and time-series consistency

Uncertainty values are estimated based on maximum uncertainties determined in EU ETS 601/2012/EC Regulation for Iron and Steel production. Uncertainties for CH₄ are estimated based on 2006 IPCC Guidelines.

Uncertainty		AD	EF	Combined
2C1 Iron and Steel Production	CH ₄	10	10	14.14
2C1 Iron and Steel Production	CO ₂	7.5	5	9.01

4.6.1.7 Source-specific QA/QC information and verification

Please note that in Hungary a quite wide range of emission sources are allocated in present 2.C.1 sector, which might cause differences compared to other countries, although we believe that it is in

line with reporting of the 2006 IPCC Guidelines as it is described in the introduction of this chapter above.

For example, during the trial review performed by EU in November 2015 it was noted that the IEF is high compared to other countries both in 2.C.1.b and 2.C.1.d-e subsectors. The high IEF can be explained by the fact that several types of emissions, including emissions from BOF and limestone and dolomite use are also included in this category.

In addition, it was also noted that in 1993, 1998 and 2013, the change in CO₂ emissions (total of 1.A.2.a and 2.C.1. compared to the previous year) deviated distinctly from the change in pig iron production (compared to the previous year). The explanation is that in HU Inventory BFG delivered outside from iron and steel factory and used for energy purposes is reported in 1.A.1.a sector. If BFG used for energy purposes (and reported in 1.A.1.a) is also taken into consideration, trends are much more parallel (see Figure 4.6.2 below). In this case the deviations from the trend of pig iron production in the years mentioned also disappear, except for the year 1993. In this year, there might be some problem with pig iron production data, as the trend of emissions and the trend of fuel used in IEA EnStat are in agreement (see blue columns and green line in Figure 4.6.2 below). The amount of BFG used within iron and steel production and delivered outside for energy purposes is based on IEA EnStat allocation.




Figure 4.6.2 Comparison of trends of pig iron production, fuel use and the emissions allocated in different sectors in HU GHG inventory

4.6.1.8 Source-specific recalculations

In 2.C.1.a – Steel production, from 2009 a double count was performed regarding the CO₂ emissions from EOF production. Corrected CO₂ emissions are reported from 2009 onward.

in 2.C.1.d – Sinter and pellet, from 2016 a double count was performed regarding the CO₂ emissions from produced sinter. Corrected CO₂ emissions are reported from 2016 onward.

4.6.1.9 Source-specific planned improvements

None.

Table 4.6.4 *Trend of activity data and emissions in 2.C.1 d. Sinter subsector*

	Coke consumption in sinter plant	NatGas consumption in sinter plant	Limestone used in sinter plant	Dolomite used in sinter plant	SUM emission 2.C.1d-e
	TJ	TJ	kt	kt	kt CO ₂ -eq
1985	3012	207	175	164	552.92
B.Y.	2853	206	174	163	532.59
1986	2935	203	172	161	539.88
1987	2612	208	176	165	504.99
1988	2569	207	175	164	498.49
1989	2207	193	163	153	441.91
1990	1893	168	142	133	380.91
1991	1528	130	110	103	302.49
1992	1460	116	98	92	282.03
1993	1472	139	118	110	303.84
1994	1687	158	133	125	346.70
1995	1637	150	127	119	333.53
1996	1574	148	125	117	324.17
1997	1157	113	95	89	241.74
1998	1250	124	105	99	263.61
1999	1339	129	110	103	279.07
2000	1339	132	112	105	281.70
2001	1241	121	103	96	259.62
2002	1287	132	112	104	274.89
2003	1273	132	111	104	272.96
2004	1333	113	78	137	280.52
2005	1264	157	91	124	273.85
2006	1228	125	104	108	265.50
2007	1147	127	118	103	263.00
2008	1408	137	137	78	293.31
2009	1086	108	91	49	214.35
2010	1338	120	137	65	281.98
2011	1462	128	139	100	310.75
2012	1296	131	110	100	279.04
2013	943	138	35	75	191.04
2014	1340	140	110	76	262.70

2015	1317	131	93	115	272.52
2016	1191	130	36	88	202.43*
2017	1273	124	63	131	257.50*
2018	1350	132	68	132	265.56*
2019	1350	138	76	123	261.88
2020	1239	131	119	108	258.71

* Recalculated in the 2021 submission

4.6.2 Ferroalloy Production (CRF sector 2.C.2)

4.6.2.1 Source category description

Emitted gas: CO₂

Methods: T1

Emission factors: D

Key sources: None

Upon smelting alloying additive and iron, together with slag-forming additives, a reduction reaction occurs which results in release of CO₂.

Ferroalloy production was present in Hungary only between 1985 and 1990.

4.6.2.2 Methodological issues

Fuels were included in sector 1.A.2.A and only technological CO₂ emissions were calculated here. The production data were obtained from the HCSO.

Default Tier 1 emission factors from the 2006 IPCC Guidelines have been applied together with the new EF for CH₄ emission.

4.6.2.3 Uncertainties and time-series consistency

Uncertainties are estimated based on 2006 IPCC Guidelines.

Uncertainty		AD	EF	Combined
2C2 Ferroalloys Production	CH ₄	5	37.5	37.83
2C2 Ferroalloys Production	CO ₂	5	37.5	37.83

4.6.2.4 Source-specific QA/QC information and verification

General QA/QC procedure apply.

4.6.2.5 Source-specific recalculations

None.

4.6.2.6 Source-specific planned improvements

None.

4.6.3 Aluminium Production (CRF sector 2.C.3)

4.6.3.1 Source category description

Emitted gases: CO₂, PFCs (CF₄, C₂F₆)

Methods: T1, T2

Emission factors: D

Key sources: 2C3 Aluminium Production – PFCs – T

During alumina electrolysis, CO₂ is released from carbon anode. At the same time, fluorinated hydrocarbons are produced from cryolite as a result of anode effect when aluminium oxide concentration is low in the electrolyte of the reduction cell. From the beginning of 2006 this technology is no longer in use in Hungary.

4.6.3.2 Methodological issues

PFC emissions were calculated using Tier 1 methodology for CO₂ and Tier 2 methodology for PFCs recommended by the 2006 IPCC Guidelines.

Production data, including data on the sites already abandoned, were obtained directly from the factories. After the major political changes, two electrolysis plants were abandoned. The resulting changes in the volume of aluminium production (Söderberg process) are shown in the table below (*Table 4.6.5*).

Very detailed, equipment-level data is also available from the factories on production: anode effect minutes per cell-day that makes possible the use of Tier 2 method for PFCs. Default slope coefficients from Table 4.16 of the 2006 IPCC Guidelines are applied.

The trend of emissions is also included in *Table 4.6.5* below. CO₂-eq emissions of PFC are calculated using new GWP values from IPCC 4th AR as it is required.

Table 4.6.5 Amount of aluminium produced (kt) and trend of CO₂ and PFC emissions

	Production of aluminium (kt)	CO ₂ Emission (kt CO ₂)	CF ₄ emission (kt CO ₂ -eq)	C ₂ F ₆ emission (kt CO ₂ -eq)
1985	73.86	125.57	333.35	34.09
B.Y.	73.75	125.37	336.50	34.58

1986	73.87	125.59	337.46	34.67
1987	73.51	124.96	338.67	34.99
1988	74.64	126.89	329.96	33.67
1989	75.19	127.82	357.52	36.90
1990	75.13	127.72	340.18	35.54
1991	62.88	106.89	293.23	30.37
1992	26.82	45.59	165.55	14.49
1993	27.88	47.39	178.93	15.66
1994	29.65	50.40	195.12	17.07
1995	31.91	54.25	204.80	17.92
1996	33.47	56.89	195.68	17.12
1997	33.67	57.25	195.06	17.07
1998	33.71	57.31	209.94	18.37
1999	33.64	57.19	214.99	18.81
2000	33.85	57.55	258.68	22.63
2001	34.59	58.80	243.64	21.32
2002	35.29	60.00	247.63	21.67
2003	35.04	59.56	231.16	20.23
2004	34.35	58.39	245.58	21.49
2005	31.78	54.03	255.15	22.32
2006-	NO	NO	NO	NO

4.6.3.3 Uncertainties and time-series consistency

Uncertainties are estimated based on 2006 IPCC Guidelines.

Uncertainty		AD	EF	Combined
2C3 Aluminium Production	CO ₂	2	10	10.20
2C3 Aluminium Production	PFC	2	99	99.02

4.6.3.4 Source-specific QA/QC information and verification

The factory operated an accredited quality assurance system. We have seen very well kept production records. The necessary data were given to us from these records. The company could provide data from almost 20 years of production without any difficulty.

4.6.3.5 Source-specific recalculations

None.

4.6.3.6 Source-specific planned improvements

None.

4.6.4 Zinc Production (CRF sector 2.C.6)

Notation keys for activity data and CO₂ emission were changed from “IE” to “NO” in the whole time-series due to recommendation of ERT (2016 in-country review). The last zinc mine was closed in 1985 and was flooded in 1986, since then only zinc processing is occurring in Hungary with fuel related emissions (it is included in 1A sector).

4.7 Other Products Use (CRF sector 2.D)

4.7.1 Source category description

Emitted gas: CO₂

Methods: T1, T2

Emission factors: D

Key sources: None

In this sector, CO₂ emitted during the use of lubricants (2.D.1) and paraffin waxes (2.D.2) are included. In these products carbon is mostly stored, however some carbon is oxidized and emitted in the form of CO₂ during their use. In addition, CO₂ emitted during urea-based catalyst in vehicles is reported in 2.D.3 subsector as it was suggested by EU experts; however, these emissions are very low.

Since 2016 submission indirect CO₂ emissions from the oxidation of NMVOC has also been included, but solely from those sectors that had been reported before the 2015 submission, too. This is in line with the recommendation of EU WG1 of February 2016 that states:

“According to paragraph 29 of the UNFCCC reporting guidelines for GHG inventories (Annex I to decision 24/CP.19) “Annex I Parties may report indirect CO₂ from the atmospheric oxidation of CH₄, CO and NMVOCs. For Parties that decide to report indirect CO₂ the national totals shall be presented with and without indirect CO₂”.

Para 37(b) the UNFCCC reporting guidelines states: “Once emissions from a specific category have been reported in a previous submission, emissions from this specific category shall be reported in subsequent GHG inventory submissions.”

Reporting of indirect CO₂ emissions is not mandatory (“may” in paragraph 29), however in combination with paragraph 37(b) those countries that included indirect CO₂ emissions in the past in their GHG inventories shall continue to report indirect CO₂ emissions in their inventory.

In the case of Hungary, indirect CO₂ from the oxidation of NMVOC from old NFR sector 3 (NFR09 codes) had been reported before 2015 submission. This corresponds to NFR sector 2.D.3.a Domestic solvent use, 2.D.3.d Coating applications, 2.D.3.e Degreasing, 2.D.3.f Dry cleaning, 2.D.3.g Chemical Products Use, 2.D.3.h Printing and 2.D.3.i Other subsectors at the moment. Indirect GHGs reported in an aggregated way under 2.D sector are taken from the last submission of CLRTAP Air Pollutants Emission Inventory of Hungary. In this way consistency is ensured with the other reporting obligation. In 2019 submission a new source was added - edible and non-edible oil extraction. Extraction of oil from oil seeds is performed either mechanically or through the use of solvents, or both. NMVOC emissions (diffuse and point sources) from solvent used were obtained from the National Air Emissions Database (LAIR) for the 2006-2017 period and implied emission factor was used for the 1985-2005 period knowing the amount of oil seeds processed. This indirect CO₂ emission was added to this category.

4.7.2 Methodological issues

CO₂ emission from lubricants and paraffin wax use are reported using Tier 1 method and default emission factors from the 2006 IPCC Guidelines (see *Table 4.7.1*).

Activity data is taken from IEA Energy Statistics.

In the case of urea based catalyst in vehicles (2.D.3), emissions are reported using Tier 1 method and Eq.3.2.2 from Vol. 2 of 2006 IPCC Guidelines. Activity data is taken from COPERT model.

Table 4.7.1 Default emission factors applied in 2.D.1 and 2.D.2 sectors

	Lubricant use Paraffin wax use
CC Lubricant = carbon content (default) kg C / GJ NCV (=t/TJ)	20
ODU - oxidised during use factor (default)	0.2

Indirect CO₂ emissions from the oxidation of NMVOC from subsectors mentioned above have been included using default 0.6 t C/ t NMVOC value from 2006 IPCC Guidelines (Volume 1. Chapter 7. p. 7.6). Please note that the same values have been used in case of submissions before 2015 of Hungary. The trend of direct and indirect CO₂ emissions of the whole 2D sector is presented in *Figure 4.7.1* below.

Please note that the trend of emissions from lubricant and paraffin wax use is consistent with the trend of lubricant and paraffin wax use in IEA EnStat.




Figure 4.7.1 Trend of CO₂ emissions in sector 2.D

4.7.3 Source-specific QA/QC information and verification

General QA/QC procedures apply.

4.7.4 Source-specific recalculations

2.D.3 – NMVOC emission from solvent and other product uses subcategory was recalculated due to revised estimates in NFR: including new source - edible and non-edible oil extraction. Extraction of oil from oil seeds is performed either mechanically or through the use of solvents, or both. NMVOC emissions (diffuse and point sources) from solvent used were obtained from the National Air Emissions Database (LAIR) for the 2006-2017 period and implied emission factor was used for the 1985-2005 period knowing the amount of oil seeds processed. This indirect CO₂ emission was added to this category.

The recalculation has a very small effect on total CO₂ emission of Hungary, also changes within the subcategory „2D3 Other – Indirect CO₂ emissions from solvent and other product uses” are lower than 1.5% for the whole time-series.

In the 2020 submission the 2.D.3 – NMVOC emission from solvent and other product uses subcategory was recalculated again. NMVOC emissions have been changed in every year because of corrected activity data in sub-sectors 2.D.3.d, 2.D.3.g and 2.D.3.h and changing from Tier1 to Tier2 methodology in sector 2.D.3.f (Dry cleaning). These recalculations are described in the 2020 IIR report of Hungary.

From the 2020 submission, NMVOC emissions from dry cleaning are reported using Tier 2 approach based on technology-dependent emission factors and the quantity of material cleaned. Activity data are available from the year 2004, for earlier years an extrapolation was made based on an estimated factor of emission per capita using population data.

In the 2021 submission, in sub-sectors 2.D.3.a, d-i – Indirect emissions from solvent use, calculation of CO₂ from default carbon content applied for NMVOC emissions was corrected for the whole time series.

In the 2022 submission, the following recalculations were made:

2.D.1 – Lubricant use. The number of L-category vehicles was recalculated for the period 2009-2019, thus there are minor changes in the activity data and CO₂ emissions.

2.D.3 Other – Urea based catalysts. The number of Euro 6 vehicles was recalculated for 2018 and 2019, thus minor changes occur in the activity data and CO₂ emissions.

2.D.3 Other – Indirect emissions from solvent and other product uses. In March 2020, recalculation was performed for this sector. CO₂ emissions were recalculated and corrected in the 2020 and 2021 reports, while activity data (NMVOC emission from solvent and other product uses) remained unchanged. Activity data are now corrected for the whole time period.

2.D.3 Other – Indirect emissions from solvent and other product uses. Activity data for 2D3a Domestic solvent use and for 2D3d Coating and paints for 2019 was recalculated, thus minor changes occur in the activity data and CO₂ emissions of the sector.

With the update to a new version of the COPERT 5 model (5.3.0), including also some revisions on stock and mileage:

- lubricant use in 2-stroke engines (reported in the energy sector) has been revised which affected the remaining part of all lubricants accounted here in the IPPU sector;
- amount of urea-based additive consumed for use in catalytic converters calculated from energy consumption in heavy duty vehicles has also been slightly revised.

The resulting change in emissions was less than 1 kt, far below the threshold of significance.

4.7.5 Source-specific planned improvements

NMVOC emissions from the Solvent and other product uses sector are further investigated.

4.8 Electronics industry (CRF sector 2.E)

Emitted gas: SF₆

Methods: T3

Emission factors: PS

Key sources: None

Several electronics manufacturing processes utilize fluorinated compounds for plasma etching, cleaning chambers and temperature control. The specific electronic industry sectors are semiconductor, thin-film-transistor flat-panel display and photovoltaic (PV) manufacturing. In Hungary emission from this sector is appears only between 2001 and 2005.

During the search for potential emission sources of fluorinated gases from electronics industry, no NF₃ use has been identified in Hungary before 2019, but a small quantity of SF₆ has been used between 2001 and 2005 by a semiconductor manufacturer company. So, SF₆ is reported in 2.E sector solely between years 2001 and 2005 based on the data provision of a semiconductor manufacturer company. They also declared that the SF₆ has been acquired domestically, so the amount was allocated from the time-series of annual sales of SF₆ for other use in order to avoid double-counting. This fact is in line with Table 6.7 of Volume 3 of the 2006 IPCC Guidelines, where some amount of “Si design capacities” from Hungary is listed for the years 2003-2005.

Table 4.8.1.: Emission from category 2.E

	Emission (kt CO ₂ .eq)
2001	1.19
2002	1.19
2003	1.19
2004	1.19
2005	1.19

In response to a recommendation from review conducted in 2019 Hungary should give an explanation how emissions were determined only between 2001 and 2005 and for only one company.

In Hungary, manufacturing of raw materials for the electronics industry is not significant, mostly only assembly activities take place in this sector. In order to calculate the emission of this sector for earlier years, potential sources have been identified by searching of potential emission sources. ERT recommended that Hungary provide an explanation how these potential sources were determined.

For the earlier years, Hungary got in touch with companies in 2014 which are potential emission sources of greenhouse gases in this category. Only one company said that SF₆ was used between 2001 and 2005, after 2005 the company had not used and had not bought this type of gas (this emission from 2.E category between 2001 and 2005 is below the threshold of significance. In response to the review 2020, this company confirmed that this gas was last used 15 years ago as production of that kind of product was removed from the product range and introduced new substances for cleaning panels.

Other companies said that greenhouse gases are not used during manufacturing. According to the Hungarian Photovoltaic and Solar Collector Association there are no manufacturing of photovoltaic panels in Hungary, but there was a company where technology of slicing of solar cells was used by way of an experiment but the company did not use any type of F-gases.

During the review 2020 ERT would like to know how new researches into new potential emission sources for the electronic industry are progressing. During the Web Conference of review 2020 Hungary informed the ERT that one plant has used from 2001 to 2005 SF₆ and Hungary said that Member State Experts had a list of companies which manufacture/assemble semiconductors, PV cells or other electrical equipment. These 30 plants cover the period between 1996 and 2018 according to the Hungarian Central Statistical Office (HCSO). According to this list of HCSO in 2018 and 2017 there were only 8 company in Hungary which is manufactured several kinds of electrical equipment. During the review member state experts could get in touch these plants. All of these companies confirmed, that they do not use PFCs and SF₆ during manufacturing because they only assemble accessories.

However, after the review the search for potential sources continued. As a result of this, it turned out that since 2019, solar cell production has been going on in a company by test operation and this manufacturer uses NF₃. This quantity was not reported in national statistics until now. Member state got in touch this company and clarified that this company uses NF₃ in production to periodically clean the surface of our deposition equipment. NF₃ is used through a completely closed and continuously monitored system for leaks. After use the remaining and reacted gas is neutralized by a gas burner, and the combustion product of this is treated with alkaline and electrostatic air purifiers. During these processes remaining and reacted NF₃ gas is completely removed.

As there is only one manufacturer in the country, this data is included as confidential data in CRF and also NIR.

The company provided information about the amount of NF₃ used and the amount of solar cells produced. According to the manufacturer, during the manufacturing the technology is completely safety. Due to this fact and because there is no emission factor per square meter produced in the case of PV production, Hungarian inventory does not include the emission of this activity.

Nevertheless, since 2020 the factory will no longer be operating only as planned in the test plant, Hungary will pay increased losses for the most accurate calculation, in which we will be happy to receive any help later.

Nevertheless, as this plant will no longer only operate as test operation after 2020, Hungary will pay more attention to this potential source in order to the most accurate calculation.

4.9 Fluorinated substitutes for ozone depleting substances (CRF sector 2.F)

4.9.1 Overview of sector (2.F)

Emitted gases: HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a, HFC-152a, HFC-227ea, HFC-236fa, HFC-365mfc, PFC116 (C₂F₆), PFC218 (C₃F₈), PFC-5-1-14 (C₆F₁₄)

Methods: T1, T2

Emission factors: CS, D

Key sources: Refrigeration and Air-Conditioning (2.F.1)

HFCs or their blends are mainly used in household, commercial, industrial, transport refrigeration and air conditioning equipment; fire suppression and explosion protection equipment; in aerosol products; by solvent cleaning; as foam blowing agents and other applications.

Category 2.F includes Refrigeration and Air Conditioning (2.F.1.), Foam blowing agents (2.F.2.), Fire protection (2.F.3.), Aerosols/Metered Dose Inhalers (2.F.4.) and Solvents (2.F.5.). These categories use HFCs (fluorinated hydrocarbons) and PFCs (perfluorocarbons) which are alternatives to ozone depleting substances (CFCs, HCFCs, etc.) being phased out under the Montreal Protocol. HFCs are chemicals contain hydrogen, carbon and fluorine, while PFCs contain only carbon and fluorine. HFCs and PFCs might be used alone or mixed in blends. PFCs were started to be used as an ingredient of cooling blends in 1997. In 1998 and 1999, some quantities were also used for adhesive tape production. Please note that PFCs are also emitted during aluminium production to be reported in sector 2.C.3 that used to be the main source of PFCs in the beginning of the time-series, but stopped in 2005.

In Hungary HFCs, PFs, SF₆ and NF₃ are not produced, these gases are all imported for the whole timeseries. HFCs and PFCs, SF₆ and NF₃ are included under the UNFCCC as they have high global warming potentials (GWPs). New GWPs from the IPCC 4th Assessment Report are applied as it is required by 24/CP.19 UNFCCC Guidelines (Table 4.9.1). The applicable GWPs are determined based on the effects of greenhouse gases over a 100-year time horizon as provided by the IPCC in its Fourth Assessment Report.

Table 4.9.1 The list of relevant F-gases in Hungary with GWP values to be used are defined in Annex III of Decision 24/CP.19

Greenhouse gas	Global warming potentials (GWP)	Greenhouse gas	Global warming potentials (GWP)
HFC-23	14800	HFC-245ca	693
HFC-32	675	HFC-245fa	1030
HFC-125	3500	HFC-365mfc	794
HFC-134a	1430	PFC-14	7390
HFC-143a	4470	PFC-116	12200
HFC-152a	124	PFC-218	8830
HFC-227ea	3220	SF ₆	22800
HFC-236fa	9810	NF ₃	17200

Trend

Use of HFCs started in 1990, first in mobile air-conditioners and household refrigerators. Furthermore, F-gases in medical aerosols has been used since 1992 (and using of it is still significant today). Then the use of HFCs and PFCs as a refrigerant in household refrigerators has strongly declined (for example the only Hungarian producer of household refrigerators uses exclusively R600 (isobutane) for years), and commercial, industrial refrigeration and air-conditioning became more and more important.

Coverage of all gases and all 2.F.1 subsectors are ensured by the fact that both a Hungarian and an EU-level Regulation are in force that require quite detailed data provision. The scope (list of gases to be reported) of Govt. Decree 14/2015. on fluorinated gases is the same as the scope of 517/2014/EC Regulation of the EU on F-gases, which is the same as the UNFCCC (before February 2015, the Govt. Decree 310/2008. and the Regulation No 517/2014/EC were in force in Hungary).




Figure 4.9.1 Trend of emission from 2.F.1 between 1990 and 2020 (kt CO₂-eq)

Emissions of F-gases are continuously increasing. In average a four-fifths (it is between 60-100% for the whole time series) of category 2.F come from the operation of refrigeration and air-conditioning equipment containing F-gases. In 2020, this category accounts for 91% of total 2.F emissions. The second largest subsector is 2.F.2. (Foam blowing agent), which is accounts for 7 percent of total 2.F emissions. Despite the continuous regulation of high GWP gases emission has not already stopped in this sector. Due to charging into new equipment a lot of cooling systems have been operated with gases which have higher GWP and this is the reason of this trend.

Emissions are to be reported by gas in the inventory (Figure 4.9.1.), so the blends/preparations containing different F-gases need to be proportionated.

Trend of emissions of fluorinated greenhouse gases is presented in **Table 4.9.2**.

Table 4.9.2 Emission from 2.F by subcategories between 1990 and 2020

	2.F.1.	2.F.2.	2.F.3.	2.F.4.	2.F.5.	SUM
kt CO ₂ -equivalent						
1990	0.002	0.000	0.000	0.000	0.000	0.00
1991	15.136	0.000	0.000	0.000	0.000	15.14
1992	15.478	0.000	0.000	10.038	0.000	25.52
1993	15.747	0.000	0.000	10.530	0.000	26.28
1994	19.657	0.000	0.000	11.078	0.000	30.73
1995	24.373	0.000	0.000	11.679	0.000	36.05
1996	29.538	0.000	0.000	12.532	0.000	42.07
1997	52.686	0.000	0.000	13.445	0.000	66.13
1998	76.899	0.000	0.000	14.520	33.327	124.75
1999	150.852	0.000	0.000	13.874	50.745	215.47
2000	188.598	0.000	0.354	16.081	0.000	205.03
2001	221.255	0.000	0.563	17.835	4.314	243.97
2002	306.418	0.000	0.952	19.295	4.314	330.98
2003	383.520	50.768	1.758	21.041	0.000	457.09
2004	456.951	110.370	3.530	22.727	0.000	593.58
2005	523.414	183.823	5.986	43.364	0.000	756.59
2006	627.426	205.803	6.699	58.925	0.000	898.85
2007	774.988	161.055	6.733	53.918	0.000	996.69
2008	904.672	154.169	6.602	52.060	0.000	1117.50
2009	960.257	158.270	7.359	46.466	0.000	1172.35
2010	1069.703	128.964	8.319	46.282	0.000	1253.27
2011	1197.639	136.360	8.623	48.735	0.000	1391.36
2012	1264.830	137.975	8.557	51.387	0.000	1462.75
2013	1346.489	145.538	8.375	56.231	0.000	1556.63
2014	1611.041	156.917	8.132	57.585	0.000	1833.67
2015	1644.252	127.334	7.954	45.223	0.000	1824.76
2016	1723.428	135.215	7.638	33.458	0.000	1899.74
2017	1783.907	140.156	7.397	34.035	0.000	1965.49
2018	1865.284	150.122	7.105	34.545	0.000	2057.06
2019	1965.641	153.716	7.197	35.017	0.000	2161.57
2020	2000.731	149.398	6.90932	34.9698	0.000	2192.01

Methodological issues

Following table summarizes the methodologies used for each subcategory in 2.F. Detailed information of these methods are explained in the following chapters.

Table 4.9.3. : Overview of methods, emitted gases and emission factors used in category 2.F.

	CRF Code	Emitted gases	Method	EF
Refrigeration and Air-Conditioning	2.F.1	HFCs, PFCs	Tier2	CS, D
Foam blowing agents	2.F.2	HFCs	Tier2	CS
Fire protection	2.F.3	HFCs	Tier1	D
Aerosols	2.F.4	HFCs	Tier2	CS, D
Solvents	2.F.5	HFCs, PFCs	Tier2	CS

4.9.2 Refrigeration and Air-Conditioning (2.F.1)

4.9.2.1 Source Category Description (2.F.1)

Emission from 2.F.1. is continuously increasing, its trend is depicted on **Figure 4.9.3**. The major share 42% in the range of actual emissions for year 2020 belongs to the subcategory 2.F.1.a. Subcategory 2.F.1.c, 2.F.1.e, 2.F.1.f. 2.F.1.d and 2.F.1.b accounts for 22%, 18%, 13%, 5 % and 0.0% of total 2.F.1 emissions, respectively.




Figure 4.9.2 Emission from 2.F.1 by subcategories

Table 4.9.4 Emission from 2.F.1 by subcategories

YEARS	2.F.1.a	2.F.1.b	2.F.1.c	2.F.1.d	2.F.1.e	2.F.1.f	SUM
kt CO ₂ -equivalent							
1990	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1991	0.00	0.00	0.00	0.00	15.13	0.00	15.14
1992	0.00	0.03	0.00	0.00	15.45	0.00	15.48
1993	0.00	0.04	0.00	0.00	15.70	0.00	15.75
1994	0.00	0.06	0.00	0.00	19.59	0.00	19.66
1995	0.00	0.08	0.00	0.00	24.29	0.00	24.37
1996	0.57	0.10	0.47	0.04	28.07	0.29	29.54
1997	10.60	0.11	8.58	0.92	29.07	3.41	52.69
1998	24.76	0.12	10.68	2.80	34.57	3.97	76.90
1999	60.13	0.13	30.77	6.17	43.03	10.62	150.85
2000	72.10	0.13	34.71	7.08	61.75	12.82	188.60
2001	90.49	0.14	39.46	8.88	67.78	14.51	221.26
2002	136.33	0.15	53.46	13.53	84.33	18.62	306.42
2003	166.72	0.91	71.80	16.14	102.34	25.61	383.52

YEARS	2.F.1.a	2.F.1.b	2.F.1.c	2.F.1.d	2.F.1.e	2.F.1.f	SUM
	kt CO ₂ -equivalent						
2004	211.08	0.91	92.61	20.02	99.48	32.85	456.95
2005	231.65	6.34	105.69	21.64	119.92	38.18	523.41
2006	284.50	5.83	126.46	26.63	138.84	45.16	627.43
2007	352.94	5.85	150.50	33.37	179.64	52.68	774.99
2008	414.15	6.23	181.94	38.78	199.65	63.92	904.67
2009	442.14	4.47	192.35	40.77	208.03	72.50	960.26
2010	487.59	2.67	220.49	44.06	222.96	91.94	1069.70
2011	557.34	2.63	248.70	50.30	225.85	112.82	1197.64
2012	587.84	2.86	259.54	51.62	235.51	127.47	1264.83
2013	624.53	2.29	283.11	53.64	248.26	134.66	1346.49
2014	771.66	1.71	334.49	65.09	264.61	173.48	1611.04
2015	742.68	1.24	361.95	59.58	288.65	190.15	1644.25
2016	789.18	1.27	354.59	64.02	329.19	185.18	1723.43
2017	798.28	0.49	389.11	64.75	352.67	178.62	1783.91
2018	843.28	0.15	370.38	79.11	365.63	206.73	1865.28
2019	799.16	0.02	460.27	88.08	381.35	236.76	1965.64
2020	839.83	0.01	440.79	93.49	358.90	267.70	2000.73

Share of emissions within category 2.F.1 is shown in the following Figure. Use of HFCs started in the beginning of 1990s, first in mobile air-conditioners and household refrigerators. Then the use of HFCs as a refrigerant in household refrigerators has strongly declined (for example the only Hungarian producer of household refrigerators uses exclusively R600 (isobutane) for years), and commercial, industrial refrigeration and air-conditioning became more and more important.




Figure 4.9.3 Share of emissions within category 2.F.1 (%)

4.9.2.2 Methodological issues (2.F.1)

Table 4.9.5: Emission factors used for calculations

Subcategory	QG	Emitted gases	Lifetime [years]	Emission factor	
				Lifetime (x) [%]	Initial charge remaining (p) [%]
Commercial Refrigeration	2.F.1.a	HFCs, PFCs	15	35.0	50.0
Domestic Refrigeration	2.F.1.b	HFC	15	0.3	95.6
Industrial Refrigeration	2.F.1.c	HFCs, PFCs	20	25.0	70.0
Transport Refrigeration	2.F.1.d	HFCs, PFCs	15	50.0	20.0
Mobile Air-Conditioning	2.F.1.e	HFCs	15	15.0	16.7
Stationary Air-Conditioning	2.F.1.f	HFCs, PFCs	12	15.0	80.0

Emission factors, summarized in previous table have been determined based on expert estimates and 2006 IPCC Guidelines. *Emission factor x* is the average annual leakage and average annual emissions during servicing (%). *Emission factor p* is the residual charge of HFC in equipment being disposed (expressed in percentage of full charge, %).

The leakage rate of 1 percent of production was determined with the help of experts a few years ago. After the 2021 review, Hungary tried to collect more data. According to one of Hungary's largest car manufacturers, the leakage rate is zero. So, Hungary is asking other companies to provide the best estimate for this parameter (for the next submission).

Although, IPCC default range is 6–9 years (lifetime) in 2006 IPCC Guidelines (vol. 3, p.7.52) for subcategory 2.F.1.d, the experience in Hungary shows that it is significantly longer.

History of improvements

Before submission 2019, Hungary used the mass-balance approach for the estimation of emissions from the whole 2.F.1 category. So, Equation 7.9 from Vol.3 of the 2006 IPCC Guidelines was applied, where data on annual sales of new refrigerant; total charge of new equipment. original total charge of retiring equipment and amount of intentional destruction is needed.

During the 2017 submission, the ERT recommended that instead of Tier1, Hungary use the Tier 2 method for 2.F.1. category. Instead of dividing sector 2.F.1. to sub-categories based on a study, Hungary should collect more data by sub-categories. In order to follow the recommendations and to obtain the most reliable data for the estimates of HFCs and PFCs emissions from this category including emissions from F-gases imported and exported in bulk and also in equipments, the following changes were implemented for submission 2019:

For years the mass-balance approach and the 'top-down' approach were applied for all sub-categories in 2.F.1. For the following applications a 'bottom-up' approach had been applied relying on statistics and expert estimations:

- Domestic Refrigeration (2.F.1.b)
- Mobile Air -Conditioning (2.F.1.e)

The amount of HFC-134a used for fill and refill products in subcategories Mobile Air-Conditioning and Domestic refrigerator was subtracted from the amount of imported and exported chemicals in bulk (for HFC-134a), as these amounts are already accounted for.

For the other four sub-categories, Hungary used chemical sales which is based on the amount of exported and imported F-gases. In these sub-categories Hungary used the same approach as before completed with an estimation of exported and imported F-gases in equipments, so disposal emissions from prefilled equipment appear in the inventory. So, Hungary was used the combination of Tier 1 and Tier 2 approach for the estimation of emissions until submission 2021.

Up to submission 2020, recovery of refrigerants was not considered in the calculations. Previous submission, according to the ERT recommendation recovery efficiency was developed for subcategories of 2F1. Therefore, the amount of emission from disposal was recalculated based on the amount of fluid in operating systems. For these sub-categories we used the same allocation of refrigerants as in previous submissions except for HFC-134a. For the latter, the proportion of the amounts in operating systems as contained in the F-gas database was taken into account.

The calculation of recovery efficiency was based on data from the F-gas database, which has been operating since 2010. Generally, according to the Hungarian experts, recovery efficiency has been increasing over the past years. As before 2010 Hungary has no further information about the efficiency of disposal, it was supposed that before 2010 recovery was negligible. To consider the imported products and equipment containing F-gases in the calculations the following method was applied previous submissions: the Hungarian 'F-gas database' includes the amount of imported F-gases in prefilled equipment since 2015. The most relevant blend contained in imported equipment is R410A (50% HFC-32 and 50% HFC-125) based on the Hungarian 'F-gas database'. Imported F-gases contained in equipment is calculated according to the database. The rate of the amount of fluid in operation systems and the amount of imported R410A in equipment are taken into consideration in the calculations. The calculated percentages for the amounts of F-gases contained in net imported products are 22% and 7% for HFC-32 and 125, respectively.

Activity data

Annual sales data is calculated as the difference of import and export of bulk chemicals. Documented, consistent time-series of import-export exists since 1992, thanks to the fact that the former Ministry for Environment, Nature and Water collected this data together with annual sales data of ozone depleting substances directly from the wholesaler companies. HMS has always been in a strong cooperation with the Ministry, so this data was used for the calculation of the inventory, together with the additional information collected directly by HMS when it was necessary.

By entry into force of Govt. Decree 310/2008 (XII.20.) the task of data collection was transferred to the Hungarian Monitoring Body for Certification (HMBC) as it is described in 2.F General chapter above. HMS receives the data needed from the HMBC database for the preparation of the inventory still through the ministry responsible for environment (Ministry for Agriculture). Consistency of the time series is ensured by the fact that it was checked that the wholesale companies reporting to the Ministry of that time are the same companies that report to the HMBC database too (except for the natural changes of the market, like cessations and entries of course).

The Hungarian Monitoring and Certification Body was also appointed for certification of persons required by 842/2006/EC ("EU F-gas Regulation"). HMBC and the database were maintained by the Association of Cooling and Air Conditioning Businesses. Govt. Decree 310/2008. (XII.20.) was replaced by Govt. Decree 14/2015 (II.20) that moves the responsibilities of HMBC to the newly established National Climate Protection Authority. With the adoption of Government Decree No 14/2015. (II.10.) on the conditions of activities with fluorinated greenhouse gases and ozone depleting substance the Authority was established on February 18, 2015 and its tasks were defined. HMBC database was renamed as 'F-gas database'.

Examining data exist in F-gas database, Hungary has realized that data by sub-application are accurate only from 2017. Amount of HFCs and PFCs banked in existing systems per sub-application serves as the basis for the calculations.

New method

In order to develop a Tier 2 methodology, further data collection and detailed data analysis were required. In addition to the above, it was recommended to Hungary uses the emission factor (Tier2a) approach in this sector. As, availability of data has not expanded enough to Hungary use top-down approach in every subcategory (in 2.F.1), data on chemical sales also plays an important role in the calculations. In subcategories, where top-down approach was improved in earlier years, there was some changes in the calculation method in order to meet the emission factor approach. These changes and the emission factors will be specified in the relevant chapters.

In general, the equation 7.10-7.14 of Chapter 7.5.2.1 on Choice of method from 2006 IPCC Guidelines was used:

$$E_{total, t} = E_{charge, t} + E_{lifetime, t} + E_{end-of-life, t}$$

where:

- $E_{charge, t}$: the emission during system assembly (as in Hungary manufacturing of refrigerators and air-conditioning systems is not appears, only assembly losses are taken into account),
- $E_{lifetime, t}$: the amount of F-gas emitted during system operation,
- $E_{end-of-life, t}$: the amount of HFC and PFC emission at system disposal in year t.

$E_{containers}$ have been left out of the equation, due to emission related to management of refrigerant containers is typically less than 0.05% in Hungary (according to a Hungarian expert of the biggest F-gas distributor). Hungary would like to improve this value by several experts later.

Previous equation consists of the following terms:

1. Refrigerant charge emissions of new equipment:

$$E_{charge,t} = M_t \cdot \frac{k}{100}$$

where:

- M_t : amount of HFC charged into new equipment in year t (per sub-application), kg
- k : emission factor of assembly losses of the HFC charged into new equipment (per sub-application), %

2. Emission during lifetime (operation and servicing):

$$E_{lifetime,t} = B_t \cdot \frac{x}{100}$$

where:

- B_t : amount of HFC banked in existing systems in year t (per sub-application), kg
- x : annual emission rate of HFC of each sub-application bank during operation accounting for average annual leakage and average annual emissions during servicing , %

3. Emissions at end-of-life:

$$E_{end-of-life,t} = M_{t-d} \cdot \frac{p}{100} \cdot \left(1 - \frac{\eta_{rec}}{100}\right)$$

where:

- M_{t-d} : amount of HFC emitted at system disposal in year t (per sub-application), kg
- p : residual charge of HFC in equipment being disposed of expressed in percentage of full charge , %
- η_{rec} : recovery efficiency at disposal, %

To make a consistent timeseries the following assumptions and methods was used:

- Hungary uses the real quantity of banked existing systems per sub-application between 2017 and 2019 for subcategories 2.F.1.a/c/d/f
- It is important to highlight, according to the legislation, all applications that are subject to leak testing are included is also required to register in the F-gas database. Refrigeration and air-conditioning equipment containing 3 kg or more of F-gases are subject to leak testing. Due to this fact smaller mobile air-conditioning systems are not obligatory to reported. Some undertaking report air-conditioning equipment in database, but this value is negligible.
- For that 3 years, rate of gases existing systems per sub-application can be quantifiable. **Table 4.9.6** includes these values.

- As less variability can be detected in the data, Hungary was used the average of that 3 years in further calculations.
- Refrigerant quantities in equipment are available as aggregated data for the whole 2.F.1 sector from 2010. These data are still unreliable at the beginning of the time series, as significant changes took place in the institutions responsible for data collection, so this data was used only from 2012. According to this, Hungary provided a timeseries by subcategory based on the percentages summarized in **Table 4.9.6**, to estimate the amount of gas in equipment.
- For years before 2012, data based on the balance of import and export. Since 2010, database contains data on how much gas was used for filling and refilling into new appliances. Hungary calculated ratios from these values between 2010 and 2015. From these 6 years an average was obtained. A share of 36% is charged to new equipment (installation) and 64% for servicing.
- These data are in proportion to the total amount of gas aggregated, not for each gas.
- 1% of the amount calculated as total charge of new equipment was taken into account as prompt emission.

Table 4.9.6 Rates of gases existing systems per sub-application

	2.F.1.a	2.F.1.c	2.F.1.d	2.F.1.	SUM
HFC-23	0%	84%	0%	16%	100%
HFC-32	9%	38%	1%	53%	100%
HFC-125	41%	34%	3%	21%	100%
HFC-134a	14%	66%	1%	19%	100%
HFC-143a	77%	16%	6%	1%	100%
HFC-152a	4%	30%	4%	62%	100%
PFC-116	1%	92%	0%	8%	100%
PFC-218	44%	15%	38%	3%	100%

The following table contains the values of recovery efficiency:

Table 4.9.7 Recovery efficiencies between 2010 and 2020 by gas (%)

Recovery efficiency	2010	2011	2012	2013	2014	2015	2016	2017-2020
HFC-23	5.0	10.1	15.1	20.1	25.1	30.2	35.2	40.2
HFC-32	8.3	16.6	24.9	33.1	41.4	49.7	58.0	66.3
HFC-125	7.0	14.1	21.1	28.2	35.2	42.2	49.3	56.3
HFC-134a	4.5	9.0	13.5	18.0	22.5	27.0	31.4	35.9
HFC-143a	6.2	12.4	18.6	24.8	31.0	37.2	43.5	49.7
HFC-218	0.7	1.3	2.0	2.7	3.3	4.0	4.7	5.3

Determination of original total charge of retiring equipment is the total charge of new equipment lifetime years ago multiplied by the emission factor initial charge remaining (p), according to the equation 3 in this chapter. An average lifetime of 12 years has been applied based on Table 7.9 and chapter 7.5.2.2 on Choice of Emission Factors from Vol. 3 of 2006 IPCC Guidelines.

These efficiencies were calculated using data from the F-gas database, which was created in 2010. According to national experts, recovery efficiencies have been increasing over the past few years. As Hungary has no further information about the efficiency of the disposal of refrigerants prior to 2010, recovery was assumed to be negligible prior to this time. For 2017–2019, the average of these three years was used to estimate the recovery efficiencies and, assuming negligible recovery prior to 2010, calculated a linear trend between 2010 and 2017.

4.9.2.3 Domestic refrigeration (2.F.1.b)

In Hungary, there is only one relevant household refrigerator manufacturer. Although the ban of the placing on the market of domestic refrigeration appliances containing F-gases with a GWP >150 entered into force on 1 January 2015 according to the Regulation 517/2014 (Annex III) in this factory instead of HFC-134a, R600 was used since 2008 for filling refrigerators.

The following method was applied to calculate emission from domestic refrigerators:

- The number of manufactured refrigerators and the amount of refrigerants filled in new products are available directly from the producer.
- Equations in Chapter 4.9.2. was used, emission factors are summarized in *Table 4.9.5*.
- Number of prefilled (imported) products is available from HCSO statistics.
- For calculating the charge of manufactured and prefilled domestic refrigerators, a value of 20 g is used.
- In this sub-category the lifetime of domestic refrigerators is 15 years and a value of 0.3% annual leakage rate was determined (according to the table 7.9).
- The refrigerant stock is determined according to the number of sold refrigerators in Hungary. The amount of F-gases used for servicing (refill) is calculated from the amount of filled fluid multiplied by the annual leakage rate.
- Emissions from manufacturing are not applicable (NA) because the factory used an isolated system for filling new refrigerators.
-

For sub-category 2.F.1.b, disposal loss factor of 100% is used until submission 2019 to avoid underestimation of emission from this category. For last submission Hungary was developed the value of this factor. In Hungary there are programmes in place in order to reform labor-saving devices since 2014 and the successfulness of this exchange program was summarized in studies. The programs are declared by Ministry for Innovation and Technology. The main target of them to households use more energy-saver devices, like refrigerators and washing machines in order to people can exchange their appliances in a more economy way. The number of changes are summarized in several studies, which are the base of used percentages by Hungary to calculate emissions from domestic refrigerators.

Due to the programme has started in 2014, data are reliable for years between 2015 and 2018. An average amount (piece of 30000) of exchanged refrigerators was multiplied by the average charge of refrigerant (20 g). The rate of this value and the amount of fluid in operating systems was used. According to the information recovery before 2005 was negligible, between 2005 and 2010 values were interpolated. So according to the ERT recommendation recovery was taken into account.

4.9.2.4 Mobile Air-conditioning (2.F.1.e)

As for Hungary, a database about the number of cars equipped with air-conditioning systems is not available, it's necessary to use more activity data for the calculations. The number of manufactured cars, the stock of road vehicles, the number of road vehicles registered for the first time in Hungary and the rate of cars equipped and not-equipped with air-conditioning systems was used in the calculations. The rate for the whole time series is based on the European vehicle categories. The following table (**Table 4.9.8**) includes the used percentages of vehicles with air-conditioning systems in manufactured cars and for stock of road vehicles.

Table 4.9.8 Percentages of cars equipped with air-conditioning systems.

Year	new cars equipped with air-conditioning systems	registered cars equipped with air-conditioning systems	Year	new cars equipped with air-conditioning systems	registered cars equipped with air-conditioning systems
1991	5%	5%	2006	95%	37%
1992	5%	5%	2007	95%	45%
1993	30%	5%	2008	95%	50%
1994	30%	5%	2009	100%	53%
1995	30%	7%	2010	100%	56%
1996	60%	8%	2011	100%	57%
1997	60%	8%	2012	100%	59%
1998	60%	10%	2013	100%	61%
1999	60%	12%	2014	100%	62%
2000	85%	16%	2015	100%	64%
2001	85%	17%	2016	100%	72%
2002	85%	20%	2017	100%	74%
2003	85%	23%	2018	100%	76%
2004	85%	28%	2019	100%	78%
2005	85%	33%	2020	100%	80%

The following method was applied for calculate emission from mobile air-conditioning for cars:

- The number of manufactured cars is available directly from the manufacturers. The emission from manufactured cars is calculated by multiplying percentages of new cars with air-conditioning systems and the refrigerant charge of cars.
- The average refrigerant charge is between 700 g and 550 g (in the earlier years more refrigerant was required for mobile air-conditioners in cars).
- As the use of HFC-1234yf is more and more significant, the inventory includes the emission of HFC-1234yf from submission 2017. *Figure 4.9.4.* shows the rate of HFCs in automobiles according to an expert estimation.
- A value of 1% for leakage rate of manufacturing was used in the calculations.

- The refrigerant stock are determined according to the number of registered cars on the road, which are available from national statistics. Cars are divided according to their classification to European emission standards.
- The annual quantities of refrigerants in operating systems are calculated by multiplying the number of cars and the percentages of vehicles with air-conditioners and the average charge.
- Amount of F-gases used for servicing (refill) is calculated from the number of passenger cars multiplied by a leakage rate of 15% (following the recommendation of national experts which is in accordance with the 2006 IPCC Guidelines (from table 7.9 on page 7.52 of the 2006 IPCC Guidelines).
- Finally, the average lifetime is 14 years in the calculations. So, disposal emission was determined using the annual total charge of vehicles 14 years before.
- In the category mobile air-conditioning systems (2.F.1.e), emissions also from buses and railways were taken into account in this submission (until now the inventory included solely emissions from automobiles in this sub-category). In addition, activity data were revised because of clarifying the number of air-conditioned cars.
- Equations in Chapter 4.9.2. was used, emission factors are summarized in *Table 4.9.5*.

During the review in 2021, the ERT recommended that Hungary explain the main reason of the reduction of emissions for 2.F.1.e. Refrigerant charge in cars, was updated from 0.7 kg for the whole time series to 0.55 kg for 2004–2019. According to one of the biggest manufacturers, the typical charge 15 years ago was 0.7 kg, but as of 2020 was 0.4 kg, resulting in an average refrigerant charge of 0.55 kg for the intermediate period of 2004– 2019.




Figure 4.9.4 Rate of quantity of HFC charged for automobiles between 2000 and 2020 (%)

Calculation method for automobiles:

- the activity data is available from national statistics (number of buses: number of manufactured buses, stock of road buses/coaches. number of road vehicles registered for the first time)
- moreover, some information (number of air-conditioned coaches and charge of the mobile air-conditioners, the refrigerant type in these vehicles) are available directly from the biggest transport companies, like Volánbusz Transport Company and BKK (Centre for Budapest Transport)
- emissions from buses are calculated from 1995, because before this year Hungary has not relevant data, moreover according to existing data it is negligible because of small rate of air-conditioned vehicles (1-2%)
- the refrigerant used in automobiles is HFC-134a
- the average charge is 8kg for long-distance buses, 10kg for local buses according to the companies
- a value of 15% for leakage rate was used for a year
- a value of 16.7% for initial charge remaining (p) was used
- the average lifetime is 14 years in the calculations, so disposal emission was determined using the annual total charge of vehicles 14 years before
- amount of F-gases used for servicing (refill) is calculated from the quantity of refrigerant charge in air-conditioners, number of air-conditioned buses multiplied by a leakage rate of 15% (according to the 2006 IPCC Guidelines (from table 7.9 on page 7.52 of the 2006 IPCC Guidelines)

Calculation method for trains:

- data on stock of trains and the used refrigerant type and the refrigerant quantity are available from the company which has the main activity in railway passenger transportation
- for every kind of trains the type and the quantity of charged refrigerant data is available directly from the aforesaid company (12 kg, 41 kg and 27 kg)
- in mobile air-conditioners of trains HFC-134a, HFC-422D are charged (90% is HFC-134a)
- emission from disposal is not occurring
- leakage rate is 15 % for this sub-category

Up to now, emission of HFC-1234yf was not considered in the calculations. In this submission, an expert estimation was applied to refine the emission from automobiles.

Calculation method for trams:

- data on stock of trams and the used refrigerant type and the refrigerant quantity are available from the company which has the main activity in urban public transport;

- trams (so-called Combino) were put into service in 2006 but these were basically non air-conditioned, air conditioning systems were installed into the vehicle only in 2008;
- trams (CAF) were put into service in 2015-16, and from 2019;
- mobile air-conditioners of trams in Hungary use R-407C and R-410A, both which contain HFC-32;
- emission from disposal is not occurring;
- leakage rate is 15 % for this sub-category.

The ERT noted that information about trams is not included in the NIR and section 4.9.2.4 contains only information on the methodology and emission estimates from air-conditioning systems in road vehicle and trains. The ERT also noted that HFC-32 emissions from stock for subcategory 2.F.1.e mobile air conditioning were greater for 2017 (0.11 t HFC-32) than for 2016 and 2018 (0.01 t HFC-32 in both cases). Moreover, Hungary clarified that there were copy errors in the compilation file for 2008–2019. These errors were corrected for the 2022 submission.




Figure 4.9.5 Emission of different gases from 2.F.1.e Mobile Air-Conditioning (kt CO₂-eq)

4.9.2.5 Uncertainties and time-series consistency

Uncertainties are estimated taking into account the uncertainty of a legally binding data provision of the companies.

Uncertainty	AD	EF	Combined
2.F.1 Refrigeration and Air Conditioning Equipment - HFC+PFC	10	10	14.14

In response to the recommendation of 2019 review, methodology for automobiles and trains are provided in this submission of the NIR (Ch. 4.9.1.2.3).

4.9.3 Foam Blowing (CRF sector 2.F.2)

Emitted gases: HFC 134a, HFC-152a, HFC-227ea, HFC-365mfc

Methods: T2

Emission factors: CS

Key sources: none

4.9.3.1 Source category description (2.F.2.)

Hydrofluorocarbons have been used in foam blowing industry, mainly as replacements for CFCs and HCFCs. There are 2 types of foam blowing, which are open-cell and close-cell products. The main characteristic of the open-cell foam is emission occur during the manufacturing process and shortly after it.

On the other hand, HFC emission from close-cell foams occurs throughout the products lifetime (during use). These products are polyurethane foams and extruded polystyrene foams (XPS) and these are used mainly for insulating applications.

4.9.3.2 Methodological issues (2.F.2.)

Country specific method is applied using activity data derived from PRODCOM statistics and emission factors from the 2006 IPCC Guidelines. This method has been developed due to the requirement of ERT during 2012 review and has been checked during the EU MS Support Project.

The new method of 2006 IPCC Guidelines is basically different because it encourages the inclusion of emissions from decommissioning and recovery. Please note that no decommissioning losses and destroyed quantity are reported currently but at the moment no product have yet reached the estimated (default) end of lifetime and no specific information is available on any destructed quantity or recycling technology.

Emissions of different gases from 2.F.2 Foam blowing sector are summarized in **Figure 4.9.6**.




Figure 4.9.6 Emission of different gases from 2.F.2 Foam blowing (kt CO₂-eq)

At the moment, we have no information on the use of HFC-245fa as foam blowing agent in Hungary. Until 2013 there was no reporting on any import or export of this gas into Hungary and the emission estimation method in 2.F.2 subsector does not include this gas either.

4.9.3.3 Activity data (2.F.2)

As it is stated in both Guidelines and NIRs of other parties: „it is extremely difficult to collect activity data...” Indeed, no direct data or statistics are available on the HFCs imported in products, neither on the amount of HFCs present in products. So, there was only the possibility to start from the viewpoint of the foam products, as it was discovered that in fact PRODCOM statistics (Statistics on the production of manufactured goods published on the website of EUROStat and Hungarian Central Statistical Office) contain both import-export and production data of two foam types.

These are: 22214120 - Cellular plates, sheet, film, foil and strip of polymers of styrene (containing XPS) and 22214150 - Cellular plates, sheets, film, foil and strip of polyurethanes (PUR).

In order to get the amount of HFC blown into foam products, the percentage of blowing agent within foam products, proportion of HFCs within foam blowing agents and proportion of type of HFC is also needed, as it follows:

Chemical used in Foam Manufacture (HFC filled in new products) = domestic production of year t of foam product (t) * blowing agent/ foam type (%) * HFC blowing agent / all blowing agents (%) * HFC-type / all HFC (%)

(**Chemical used in Foam Manufacture** data is to be multiplied by “first year loss” EF in order to calculate the **emissions from manufacturing**.)

Chemical emitted during the lifetime of closed cell foams (HFC charged into the product) = production + import - export of the foam type (t) * blowing agent/ foam type (%) * HFC blowing agent / all blowing agents (%) * HFC-type / all HFC (%)

(**Chemical emitted during the lifetime of closed cell foams** data is to be accumulated as many years as the lifetime of the foam product and to be multiplied by “annual loss” EF in order to calculate the **emissions from stocks**.)

In this way, **Chemical used in foam manufacture** and **Chemical emitted during the lifetime of closed cell foams** required by eq. 7.7 of the 2006 IPCC Guidelines has been expressed, so default EF-s from Table 7.5 and 7.6 could be used. In this method both the amounts imported in bulk (**Chemical used in foam manufacture**) and in products (within **Chemical emitted during the lifetime**) are accounted. The double usage of statistical data of foam production does not result double count in emissions, since in the first case it is used to determine the emissions from manufacturing occurred in Hungary even if the product is exported and in the second case it is needed to determine the amount of foam products remaining in the country responsible for the emissions from stocks.

The following **Table 4.9.9** summarizes the values and their references used in the calculation.

Table 4.9.9 Summary of factors used by the calculation in 2.F.2

	XPS	PUR	Reference
domestic production of year t of foam product (t)	Prodcom Statistics		http://epp.eurostat.ec.europa.eu/portal/page/portal/prodcom/data/databases
production+import-export of the foam type (t)			
blowing agent/ foam type (%)	6%	8%	Revised IPCC1996 page2.59 (6-15%) and IPCC/TEAP study (please see References)
HFC blowing agent / all blowing agents (%)	40% and decreasing until 20%	20% and decreasing until 10%	DG Climate F-gases Reg.Review Study and IPCC/TEAP study (please see References) and suggestion received during EU MS Support Project

Within PRODCOM 22214120 polystyrene foams category, only XPS (extruded polystyrene) type foam might be blown with HFCs. The proportion of XPS foam within polystyrene foams in the Hungarian market is estimated to be 10% by the Hungarian Association of EPS Insulating Foam Producers and another expert architect.

PRODCOM data of PUR production of 2005 to 2008 and XPS production data of 2004 and 2005 were averaged in order to avoid negative production+import-export values in the years 2006 to 2008 and to reflect better the trend. (Production data of the mentioned years are summed and divided by number of years).

HFC are used as blowing agent in foams mainly after 2003 as substitutes of ODS after the ban of CFCs and HCFCs under Montreal Protocol. Nowadays also HFCs are substituted by materials with less GWP (CO₂, Hydrocarbons, HFO, etc.) The background study of F-gases Regulation Review of DG Climate states the ratio of HFCs among the blowing agents is 40% for XPS and 20% for PUR between 2003 and 2011.

HFC use in foam blowing started in 2003 based on data reported by the intermediate material producer company (BASF).

The DG Climate study estimated that the final year of significant HFC use in foam blowing would be in 2011. However, our assumption for Hungary is 2015 (instead of 2011) based on suggestion of the IPCC/TEAP study (IPCC/TEAP. 2005), which seemed more realistic.

During the EU MS Support Project, the expert noted that the elimination of HFC blowing agent by 2015 is still not realistic and suggested to apply 20% for XPS foams and 10% for PUR foams after 2011 as well. HFC emissions of year have been recalculated based on this suggestion in sector 2.F.2.Foam.

Table 4.9.10 Proportions of HFC foam blowing agents applied by calculation in 2.F.2

		2007	2008	2009	2010	2011-2020
% of HFC blowing agent usage/All blowing agent usage in the case of	XPS products	35.6	31.1	26.7	22.2	20.0
	PUR products	17.8	15.6	13.3	11.1	10.0

The proportion of the different types of HFCs is based on the historical data reported by the intermediate material producer company (BASF). The average result is 10% HFC-227a and 90% HFC-134a. HFC-365mfc is reported the same as HFC-227ea because the company reported the use of a blend containing precise proportion of HFC -227ea and HFC-365mfc. So, HFC-365mfc is reported by multiplying the amount for HFC-227ea by this appropriate proportion. This method is the same as HFC-365mfc had been reported as cross cutting info in previous inventory submissions.

The proportion of soft foams and hard foams within PUR foams is also based on the historical data reported by the intermediate material producer company (BASF). The average result is: 10% soft foam and 90% hard foam. All soft foam is accounted as open cell foam (using eq. 7.8 of the 2006 IPCC Guidelines) and all hard foam is accounted as closed cell foam (using eq.7.7 of the 2006 IPCC Guidelines).

Please note that in addition to the abovementioned method, also a directly reported experimental usage of HFC-152a solely in the years 2006 and 2007 is included within the soft foam subcategory.

4.9.3.4 Emission factors (2.F.2)

Default emission factors from Table 7.6 of the 2006 IPCC Guidelines are used for XPS and general default emission factors from 7.5 of the 2006 IPCC Guidelines are used for PUR as the proportion of the different types of PUR foams is not known. Used values are summarized in **Table 4.9.11**.

Table 4.9.11 Default emission factors used from 2006 IPCC Guidelines

	XPS	PUR
lifetime	50 years	20 years
first year loss	40%	10%
annual loss	3%	4.5%

In the case of soft foams (all accounted as open-cell foam) equation 7.8 of 2006 IPCC Guidelines is used, so ALL the filled amount is emitted during manufacture.

Please note that the IEF (in CRF) is changing through the years due to the fact that EF of the Guidelines is determined by foam type. while IEF in the CRF is determined by HFC type.

4.9.3.5 Source specific recalculations, QA/QC activities. uncertainties and planned improvements (2.F.2.)

It is a planned improvement and also a further recommendation of MS Support Project to get in touch directly with producers and to verify whether HFC blowing agents have been used in Hungary at all, the country specific proportion of foam types and HFC types used.

Uncertainty	AD	EF	Combined
2.F.2 Foam Blowing – HFC	50	21	54.23

4.9.4 Fire Extinguishers (CRF sector 2.F.3)

Emitted gases: HFC 125, HFC-227ea, HFC-134a

Methods: T1

Emission factors: D

Key sources: none

4.9.4.1 Source Category Description (2.F.3.)

Until the beginning of 1990s, halon systems were the second wide-spread fire extinguishers after dry chemical powder extinguishing systems. Because halons do not contain hydrogen, these are inflammable. According to the Montreal Protocol these substances were phased out. Nowadays fire protection equipment is filled with HFCs as partial replacement for halons. In Hungary, mainly HFC-125 and HFC-227ea gases are the most widespread in fire extinguishing systems.




Figure 4.9.7 Emission of different gases from 2.F.3 Fire Extinguishers from year 2000 (kt CO₂-eq)

4.9.4.2 Methodological issues (2.F.3)

Activity data for this subcategory is collected by the Fire Protection Department of the National Directorate General for Disaster Management. Ministry of the Interior as part of the yearly national statistical data collection program. This data is available from 2000 and ensures the full coverage of the country on installed fire protection equipment. In addition, it includes the amount of import in products; however, it is not possible to separate from import in bulk.

The consumptions of the years are accumulated as a “bank”, and emissions of a given year is calculated as a certain (default) percent of this “bank”.

After submission 2020, Hungary has contacted with an expert who highlighted that lifetime of these devices in Hungary is usually 20 years (in agreement with IPCC Guidelines - the Guideline recommends to use years between 15 and 20 for lifetime years.). Moreover, Hungarian legislations includes this fact

for fire extinguishers. Therefore, for submission 2021 March this factor had been changed (from 15 to 20 years), so the first year when the disposal emission appears will be in 2020. In addition, as there are no accurate statistics on the recovery rate in this sector, this factor is still considered to be zero, based on the IPCC Guidelines (Chapter 7.6.2.2) and Hungary try to make further efforts to estimate recovery efficiency.

The following default emission factors have been applied:

Annual operational emission % per installed base	4%
Lifetime years	20

(source: 2006 IPCC Guidelines chapter 7.6.2.2. Tier 1 methodology)

4.9.4.3 Recalculations, QA/QC activities, uncertainties and planned improvement

2.F.3 – Recalculation was implemented in this submission as there the lifetime years was determined as 15, while in Hungary - according to the legislations - the lifetime of fire extinguishers including F-gases is 20 years. However, an additional error has been recognized after submission 2020, which caused changes for later years as well. The effect of recalculations is summarized in the following table.

Table 4.9.12 Changes in the emissions from 2.F.3. due to recalculations for period 2016-2018

YEAR	Submission 2020 [Gg CO ₂ -eq]	Submission 2021 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]
2016	7.32	7.64	0.32
2017	7.40	7.40	-0.01
2018	6.28	7.11	0.82

Uncertainties are estimated based on the 2006 IPCC Guidelines chapter 7.6.4 regarding activity data and chapter 7.6.2.2 regarding the emission factor. where it is stated: “factor range of 2 to 6 percent (that is 4 % +/-2%)”

Uncertainty	AD	EF	Combined
2.F.3 Fire extinguishers - HFC	20	2	15.13

4.9.5 Aerosols and Metered Dose Inhalers (CRF sector 2.F.4)

Emitted gas: HFC-152a, HFC-134a

Methods: T2

Emission factors: CS, D

Key sources: none

This category includes metered-dose inhalers (MDI) which are used in medical products and general-purpose aerosols. Most aerosol packages contain mainly hydrocarbons (HC) as propellants, but in a small fraction also HFCs are used, especially HFC-134a in industrial, household and medical applications. The graph below (**Figure 4.9.8**) shows the emission of F-gases from this sub-category. The main substance is HFC-134a. Emission of HFC-152a from general-purpose aerosols is almost negligible.




Figure 4.9.8 Trend of emission in sector 2.F.4 Aerosols and MDI

4.9.5.1 Metered-dose inhalers (2.F.4.a)

Emitted gas: HFC-134a

Methods: T2

Emission factors: CS

Key sources: none

4.9.5.1.1 Source Category Description (2.F.4.a)

Metered-dose inhalers are used in medical appliances, primarily for the treatment of asthma and COPD (Chronic Obstructive Pulmonary Disease). As for medical aerosols, F-gases are today only used in MDIs, not in nebulizers. These medical products with HFC propellant first reached the Hungary market in 1992.

4.9.5.1.2 Methodological issues (2.F.4.a)

In the MDI sector the Tier2a method, i.e. the user-based approach was applied. Activity data of metered-dose inhalers subcategory was the consumption of MDIs. So. data on production was not used here. The emissions are calculated from the number of MDIs (HFC quantities) sold per year in Hungary. Emissions from aerosols should be calculated using the same method for each year in the time series. For the early years in the time series. the splicing technique. the overlap method was applied to form a complete time series (find detailed description in the following chapter).

4.9.5.1.3 Activity data (2.F.4.a)

Data of annual sales from National Institute of Pharmacy and Nutrition was used between 2006 and 2015. The number of total consumption and the filled quantity in every aerosol was available. The typical charge contained in product is between 5 and 18 g. Before 2006 and after 2015 the overlap technique was applied to complete the time series. To estimate emissions for these years, the relationship between the consumption and the number of asthma and COPD (patients) cases was used applying the method of the 2006 IPCC Guidelines chapter 5.3.3.1.

Equation 5.1. from the 2006 IPCC Guidelines was applied.

$$y_t = x_t \cdot \frac{\sum_{i=m}^m y_i}{\sum_{i=m}^m x_i}$$

where: y_t – the estimated AD (the amount of the HFC quantities) in year t

x_t – the number of patients in year t

y_i – the amount of the HFC quantities in year i

x_i – the number of patients in year i.

As the difference between the average of the overlap for these 10 years (2006-2015) (19.6%) and the overlap of the first year (2006) (16.3%) is significant and the latter is more representative. we have chosen to use the overlap for the first year. For years between 2016 and 2018 the overlap of year 2015 was used (21.2%). The graph (**Figure 4.9.9**) below shows the relationship between the quantity of the gas and the number of patients.




Figure 4.9.9 Activity data of the 2.F.4.a subcategory

In the CRF table the activity data „In operation system” (the annual average stocks) and „Emission from stocks” is the same amount, because the time period between sales and use is short (product lifetime is one year). Hungary uses notation key ‘NO’ in CRF for ‘filled into new manufactured products’, as Hungary uses the user-based approach.

4.9.5.1.4 Emission factors (2.F.4.a)

In the calculations, instead of the manufactured inhalers, the purchased aerosols were used. MDIs are prescription medicine and in Hungary it is rigorously regulated. In the calculations, it was assumed that the emission level corresponded to 100% of usage (i.e. purchased aerosols), because the time period between sales and use is short. Moreover, inhaled gases are emitted into the atmosphere without undergoing any changes (according to the IPCC specifications (2006 IPCC Guidelines, Vol. 3, p. 7.28)). In consequence, a country-specific emission factor was used (EF=1).

4.9.5.1.5 Recalculations, QA/QC activities, uncertainties and planned improvements (2.F.4.a)

There was no recalculation in this year.

4.9.5.2 Other aerosols (2.F.4.b)

Emitted gas: HFC-134a, HFC-152a

Methods: T2

Emission factors: D

Key sources: none

4.9.5.2.1 Source Category Description (2.F.4.b)

Other aerosols include personal care products, household products, special cleaning sprays etc. In these cans HFC-134a, HFC-152 and HFC-227 are the most prevalent gases.

4.9.5.2.2 Methodological issues (2.F.4.b)

For 2F4b (other aerosols) the Tier 2a method was applied from IPCC2006 Guidelines, which means that 'half of the chemical charge escapes within the first year and the remaining charge escapes during the second year' (chapter 7.3.2.2. of the IPCC2006). A 50% emissions in use of aerosols is assumed.

4.9.5.2.3 Emission factors (2.F.4.b)

In line with Chapter 7.3.2.1 of the IPCC2006 the equation 7.6 was used in the course of emission calculations:

$$Emission_t = S_t \cdot EF + S_{t-1} \cdot (1 - EF),$$

where S_t is the sum of the purchases of the current year and the previous year, and EF is the emission factor ($EF = 0.5$). So, the sum of half of the purchases of the current year and half of the purchases of the previous year was considered. The equation above was applied to each chemical individually.

4.9.5.2.4 Activity data (2.F.4.b)

In subsector Aerosols, annual sales data is directly reported by the producers. So, the activity data was the consumption of purchased aerosols and the HFC quantities of aerosol cans.

In Hungary manufacturing of aerosols is not occurring. While the amount of "filled into new manufactured products" was not estimated (NE), the amount of "in operating system" was filled in as the HFC quantity of sold aerosol cans in the current year and the half the sales of the previous year.

4.9.5.2.5 Recalculations, QA/QC activities, uncertainties and planned improvements (2.F.4.b)

There was no recalculation in this year.

Uncertainties are estimated based on chapter 7.3.3 of 2006 IPCC Guidelines and taking into account that activity data is provided on one hand by individual companies, on the other hand national statistics are used.

For HFC-134a and HFC-152a in 2.F.4.b subcategory 'NE' was reported in CRF for 'filled into new manufactured products' as activity data. During the review 2019, ERT's question was referred to why this notation key was used by Hungary. In response to the recommendation Hungary has changed notation key 'NE' to 'NO' because in Hungary manufacturing of aerosols is not occurring.

Uncertainty	AD	EF	Combined
2.F.4 Aerosol + MDI(HFCs)	10	50	50.99

4.9.6 Recalculations, QA/QC activities, uncertainties and planned improvements (2.F.)

There was a correction of emission data from Mobile air-conditioning (2.F.1.e) sector (the impact of recalculation on total emissions is 0.1 kt CO₂-eq).

Subcategory 2.F.2 was recalculated, because of the revision of activity data (the impact of recalculation on total emissions including LULUCF is + 0.042 %).

For submission 2021 March, 3 subcategories of 2.F. was recalculated. The most significant change was caused by recalculation of subcategory Refrigeration and Air-Conditioning (2.F.1) due to improvements in the methodology.

A project to create a national greenhouse gas database is launched under the leadership of Ministry for Innovation and Technology. This will be a data entry and decision support IT system based on the emissions inventory. Recently, the methodological development of the F-Gas methodology has been completed, during which the methodology of the calculations for categories 2.F and 2.G has been verified. Most of the results found here have already been incorporated into this year's inventory.

4.10 Other products manufacture and use (CRF sector 2.G)

4.10.1 SF₆ use in Electronics industry (CRF sector 2.G.1)

Emitted gases: SF₆

Methods: T1

Emission factors: D

Key sources: none

4.10.1.1 Methodological issues (2.G.1)

SF₆ is mainly used as an insulation gas in electrical equipment, such as switchboards, switchgears. It was further used in the past as intermediate gas in double-glass heat insulation windows and production of optical bodies, etc. and in electronics industry for several years. In Hungary SF₆ is not used as a cover gas in coloured metal foundries.

The application of the 2006 IPCC Guidelines causes a major change within this sector although still the basic Tier 1 method is applied. Old Tier 1 calculation method accounted only for potential emissions, while the new Tier 1 method estimated actual emissions. As it was expected, emissions are lower using the new method (see **Figure 4.10.1**).




Figure 4.10.1 Trend of SF₆ emissions in 2.G.1 Electrical equipment sector (t)

4.10.1.2 Activity data (2.G.1)

Annual sales data is the basis for the calculation in the new method, as well. Data was collected from both manufacturers and the Hungarian Electrotechnical Association. The latter was appointed to data

collection by 310/2008. Govt. Decree for data collection on import-export of SF₆. This is the same time-series of annual sales data that had been used by the old calculation of potential emissions in previous submissions. However, during the review of time-series it came out that the largest equipment manufacturer did not report any export before 1998. Therefore, export data has been extrapolated using the import/export proportion of the last reported year (1998).

310/2008. Govt. Decree has now been replaced by 14/2015 (II.14) Govt. Decree that moves the responsibilities and the handling of the database to the newly established National Climate Protection Authority.

4.10.1.3 Emission factors (2.G.1)

Default emission factor from Tables 8.2, 8.3 and 8.4 of the 2006 IPCC Guidelines are applied. As there is no information in the country what percent of electrical equipment are sealed pressure electrical equipment (MV switchgear) or closed pressure electrical equipment (HV switchgear) or gas insulated transformers. always the higher EFs are taken into consideration as conservative estimation.

Table 4.10.1 Emission Factors used in 2.G.1 sector

Manufacturing Emission Factor	0.085
Use Emission Factor	0.026
Fraction of SF ₆ remaining at retirement	0.930
Lifetime (years)	35

4.10.1.4 Recalculations, QA/QC activities, uncertainties and planned improvements

General QA/QC procedures apply. Emission factors (and resulting time-series) have been verified with those included in "*Update on global SF₆ emissions trends from electrical equipment – Edition 1.1 Ecofys Emission Scenario Initiative on Sulphur Hexafluoride for Electric Industry (ESI-SF₆)*". Latter EFs are lower than the EFs in the 2006 IPCC Guidelines but it is planned to potentially include in the calculation method this more up-to-date information after further verification. Uncertainties are estimated based on Table 8.5 of 2006 IPCC Guidelines, taking the highest value as conservative estimation. Activity data's uncertainties are estimated taking into account that activity data is provided by individual companies based on a legally binding data provision requirement.

Uncertainty	AD	EF	Combined
2.G Other Product Manufacture and Use - SF ₆	3	40	40.11

4.10.2 Other applications (CRF sector 2.G.2.)

Emitted gas: SF₆

Methods: T1, T2

Emission factors: D

Key sources: none

4.10.2.1 Source category description (2.G.2)

SF₆ used mainly as an insulating medium in the radar systems of AWACS, in equipment in university and research particle accelerator, industrial and medical accelerators, adiabatic applications and in sound-proof windows. SF₆ used for sound-proof window production (only in the past due to the ban introduced by 842/2006/EC), scientific research and other non-defined purposes is included in sector 2.G.2.

4.10.2.2 Methodological issues

Equation 8.23 from the 2006 IPCC Guidelines is applied, so emissions are considered as 'prompt' emissions that is recommended in the case of any other applications. This means that emissions are distributed within two years, *"because both sales and emissions are assumed to be continuous over the year; that is, chemical sold in the middle of year t-1 is not fully emitted until the middle of year t."*(Chapter 8.3.2.2 of the 2006 IPCC Guidelines). Only emission from sound-proof windows is estimated separately.

Recommendation by ERT already had been drafted at review 2017. This was about reporting of emission from use of double-glazed sound-proof windows. The application of SF₆ in windows began in 1975, but in Hungary it may appeared later. The average lifetime is 25 years because the annual leakage rate of 1 percent is assumed. Emission was calculated according to the 2006 Guidelines (equations 8.20 to 8.22). Annual leakage rate of 1 percent is assumed and a rate of 33 percent for the filling operations.

As Hungary has no data about the whole amount of targeted windows, emission was established from produced data. Data about manufactured double-glazed windows are available from 1996, moreover Hungary has information about domestic trade from one of the biggest SF₆ wholesaler. For years 2007 and 2008 data are also available from the wholesaler by companies. From these data it could be concluded how much SF₆ companies were used for windows.

According to the years 2007 and 2008 the used SF₆ gas for windows is a percent of 7.5 of the whole amount of sold SF₆ gas. So, before 2007 this rate was considered in the calculations. This amount of SF₆ gas was filled to the sound-proof windows in Hungary.

The stock was established by summarized the annual filled quantity of the gas.

SF₆ is reported in 2.E sector between years 2001-2005 based on the data provision of a semiconductor manufacturer company. They also declared that the SF₆ has been acquired domestically, so the amount was allocated from the time-series of annual sales of "SF₆ for other use" in order to avoid double-counting. The SF₆ wholesaler company reports the list of their customer too. So, the intended use of SF₆ might be determined based on the sector of the activity of the customers. The activity data (and

consequently the emissions calculated with present methodology. too) show strong interannual variations throughout the whole time series.

4.10.2.3 Recalculations, QA/QC activities, uncertainties and planned improvement

General QA/QC procedures apply.

Uncertainties are estimated together with sector 2.G.2.

Emission from sound-proof windows was taken into account separately in response to the ERT's recommendation previous submission. In the 2021 submission, recalculation was implemented because during last year's submission sound-proof windows were double-counted in category 2.G.2. That is, So, the estimated amount of SF₆ charged into windows was not subtracted from the total amount of SF₆ used in 2.G.2 during the previous recalculation.

4.10.3 Use of N₂O (CRF sector 2.G.3)

Emitted gas: N₂O

Methods: T3

Emission factors: CS, PS

Key sources: none

4.10.3.1 Source category description

This sub-sector includes emissions of N₂O from different product uses and the manufacturing (and other) losses from the production of these products. One of the two main important purposes is bulk N₂O use as an anaesthetic gas. Another is the use by household whipped cream cartridges. In Hungary, making whipped cream in siphons using N₂O cartridges was highly popular at the beginning of the time-series.

N₂O from these products is emitted directly into the atmosphere, so all the filling of these products used should be considered as emission.

The largest manufacturer of the region of bulk N₂O is operating in Hungary. The manufacturers of the whipped cream cartridges acquire the bulk N₂O also from this manufacturer.

4.10.3.2 Methodological issues

Emissions are reported using plant-specific data. Production and domestic sales data for both bulk N₂O and N₂O in whipped cream cartridges are available from the manufacturers for the whole time-series (presented in Figure 4.10.3). N₂O used for the preparation of whipped cream cartridges (2.G.3.b.i) is subtracted from bulk domestic N₂O use (2.B.b.ii), as the manufacturer declared that they acquire the gas from the manufacturer of bulk N₂O.

Manufacturing losses for the whole time-series is also available in the case of whipped cream cartridges. While in the case of bulk N₂O production data on losses is available only from 2008. In this year, extrapolation was performed in order to include emissions from losses for the years before 2008

too, for the improvement of the consistency of the time-series. Extrapolation was performed using data on losses from 2008 (as trend of the losses is decreasing later in time) and N₂O production as surrogate data. Please find the trend of production and emissions on the following figure (**Figure 4.10.2**).




Figure 4.10.2 Trend of N₂O production and N₂O emissions from Product Uses

In 2014, an expert from the manufacturer of whipped cream chargers (cartridges) containing N₂O provided an estimate for the share of imported products on the Hungarian market, as well. In the 2014 submission, this amount has been included in the time-series. So, imports are estimated only within the subsector whipped cream, however in the case of other subsector (2.G.3.b.ii bulk use mainly for anaesthetics) imports are even less significant, as there is no notable bulk import as the Hungarian producer is the most important producer of the whole region. In addition, the wholesalers having a valid wholesaling authorisation for products containing nitrous-oxide issued by the National Institute of Pharmacy (the agency responsible for licensing and control of drugs) have been identified. Most of them have already declared that they acquire N₂O domestically (from the Hungarian producer).

The strong interannual variations are due to the interannual variations of the reported data, which is related to the production of the company and the volatility of the market.

The significant decrease on inland consumption (and consequently the emissions from inland consumption) was also reported by the companies which was investigated in this submission. The producer of N₂O has reported separately the consumption data of the firm producing N₂O cartridges and other inland consumption for the last four years. The previous assumption - “inland consumption” is equal to the total inland consumption – was not correct, because most of the N₂O leave Hungary in cartridges produced by a Hungarian company. Upon the new data request the producer company has provided the separated consumption data back to 2002.

In case of „Propellant for pressure and aerosol products” emission during charging of cartridges intended for export was not covered before. As the inland consumption is less year after year meanwhile export rises dramatically as it turned out from the corrected inland consumption data. it was necessary to include charging loss of it. Unfortunately, data about loss of N₂O charging of export cartridges is available since 2009. Until 2002 this emission could be calculated from the new consumption data of the company (amount of N₂O in cartridges for inland consumption is known for the whole time-series) combined with implied emission factor for 2009. It was assumed that N₂O cartridges were made only for inland consumption for the period 1985-1990. Before 2002 until 1990 linear interpolation was used for calculating other N₂O consumption (bulk use mainly for anaesthetics), and the rest of the inland consumption was assumed to be the consumption of the company producing cartridges. Inclusion of this new source resulted significant changes of emission in this subcategory. Data sources are summarized in the following table.

Table 4.10.2 Summary table of data sources in 2.G.3 sector

Data source for different periods	1985-1990	1991-2001	2002-2007	2008-2013	2014-
1 - N ₂ O production	Directly from the producer				
2 – Inland consumption of bulk N ₂ O	Directly from the producer				Sum of “3” and “4”
3 - Inland consumption for producing cartridges	Equal to “5”	Difference of “2” and “4”	Directly from the producer	Directly from the company	
4 - Other inland consumption	Difference of “2” and “5”	Linear interpolation between 1990 and 2002		Directly from the producer	
5 – N ₂ O amount used for charging cartridges for inland consumption	Directly from the company				
6 - Emission from charging cartridges for inland consumption	Directly from the company				
7 - Emission from charging cartridges for export	Assumed to be null	Difference of “3” and “5” (using average EF from period 2009-2014)			Directly from the producer

4.10.3.3 Uncertainties and time-series consistency

Production data is quite reliable because they are obtained directly from manufacturers. based on a legally binding data provision requirement, therefore the uncertainties are estimated as follows:

Uncertainty	AD	EF	Combined
2.G Other Product Manufacture and Use - N ₂ O	3	3	4.24

4.10.3.4 Recalculations, QA/QC activities, uncertainties and planned improvement

General QA/QC procedures apply.

Further investigation of data regarding imported products (especially whipped cream cans) might be performed. However, this source is reported only by few countries at the moment and the amount are expected to be insignificant compared to emissions reported under 2.G.3.b.ii. (production and domestic use of bulk N₂O) by Hungary.

There was no recalculation in this submission.

4.11 Other production (CRF 2.H)

4.11.1 Source category description

Emitted gas: NMVOC

In accordance with the NRF report of Hungary, indirect NMVOC emissions from category 2.H are reported in CRF tables. From category 2.H only NMVOC emissions are reported. NMVOC is emitted from 2.H.1 (Pulp and paper) and 2.H.2 (Food and beverages industry).

The calculation methodology for NMVOC emissions is described in detail in the current IIR report.

In review 2020 TERT notes, that Hungary reports certain emission data and recovery in subcategory 2.H but without creating the relevant subcategory 2.H.3 Other. This leads to problems in the assessment tool.

For these indirect emissions, Hungary has created subcategories 2.H.1. 2.H.2 and 2.H.3. and these cells has filled with appropriate values for submission 2021 March.

4.12 References

Intergovernmental Panel on Climate Change (IPCC). 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. *Intergovernmental Panel on Climate Change National Greenhouse Gas Inventories Programme*. Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: Institute for Global Environmental Strategies. Japan.

European Asphalt Pavement Association (EAPA). 2011: Asphalt in Figure 2010. http://www.eapa.org/usr_img/Asphalt%20in%20figures%20Version%2022-12-2011.pdf

ThyssenKrupp Industrial Solutions: An environmental star: EnviNOx. Brochure. Available at: http://www.thyssenkrupp-industrial-solutions.com/fileadmin/documents/brochures/tkis_EnviNOx_brochure.pdf

Nitrogénművek Zrt.. 2008: JI project on N₂O emission reduction at the Nitric Acid plant of Nitrogénművek Zrt. Annual Report. (N₂O kibocsátás-csökkentési együttes végrehajtási project a Nitrogénművek Zrt. Salétromsavüzemében. Éves Jelentés. – only in Hungarian) http://klima.kvvm.hu/documents/116/Nitrog_nm_vekZrt_ves_jelent_s2008publikus.pdf

European Commission. 2006. Integrated Pollution Prevention and Control (IPPC). Best Available Techniques in the Food, Drink and Milk Industries. Available at: http://eippcb.jrc.ec.europa.eu/reference/BREF/fdm_bref_0806.pdf

Hungarian Ministry for Environment and Water (Környezetvédelmi és Vízügyi Minisztérium). 2005: Guideline for BAT BREF in case of sugar production. (Útmutató az elérhető legjobb technika meghatározásához a cukorgyártás terén). Budapest. Available at: http://ippc.kormany.hu/download/4/e9/70000/cukor_utmutato.pdf

EMEP/EEA 2013 – Air Pollutant Emission Inventory Guidebook

Regulation (EC) No 166/2006 concerning the establishment of a European Pollutant Release and Transfer Register) <http://prtr.ec.europa.eu/>

Preparatory study for the F-gas Review. Schwarz et al. (2011) Öko-Recherche GmbH) available at: http://ec.europa.eu/clima/policies/f-gas/docs/2011_study_en.pdf

UNEP – ECA Network meeting 2011. Budapest documents: <http://www.unep.org/ozonaction/ecanetwork/Activities2011/ThematicmeetinginBudapest/tabid/56112/Default.aspx>

Personal communication to Mr. Róbert Tóth (*Ministry for Environment, Nature and Water; HMS*). Mr. Attila Zoltán. Mrs. Erika Barna (*HMBC*)

2.F.2. Sector:

DG Climate F-gases Regulation Review Study – SKM Enviros: Further Assessment of Policy Options for the Management and Destruction of Banks of ODS and F-Gases in the EU FINAL REPORT Revised Version 2 March 2012

http://ec.europa.eu/clima/policies/ozone/research/docs/ods_f-gas_destruction_report_2012_en.pdf

IPCC/TEAP. 2005 - Bert Metz. Lambert Kuijpers. Susan Solomon. Stephen O. Andersen. Ogunlade Davidson. José Pons. David de Jager. Tahl Kestin. Martin Manning. and Leo Meyer (Eds) Safeguarding

the Ozone Layer and the Global Climate System: Issues Related to Hydrofluorocarbons and Perfluorocarbons. Cambridge University Press. UK. pp 478. ISBN: 9780521682060. Chapter 7 FOAM
<http://www.ipcc.ch/pdf/special-reports/sroc/sroc07.pdf>

Emissions. Activity Data. and Emission Factors of Fluorinated Greenhouse Gases (F-Gases) in Germany 1995-2002 Adaptation to the Requirements of International Reporting and Implementation of Data into the Centralised System of Emissions (ZSE) by Dr. Winfried Schwarz Öko-Recherche Büro für Umweltforschung und –beratung GmbH. Frankfurt/Main
<http://www.umweltdaten.de/publikationen/fpdf-l/2903.pdf>

Finnish 2002 Inventory of HFC, PFC and SF6 Emissions Teemu Oinonen. 2004 The Finnish Environment 686. Environmental Protection <http://www.ymparisto.fi/default.asp?contentid=73110&lan=en>

NL Agency - NL Energy and Climate - Secretary Team NBTM - Monitoring Protocols 12-022 HFCs from Other Use HFC <https://www.agentschapnl.nl/sites/default/files/12-022%20HFCs%20from%20Other%20Use.pdf>

PlasticsEurope - GUA - The potential of plastic insulation to realise energy savings and de-coupling in Europe - Final report - January 2006 <http://www.plasticseurope.org/document/gua---the-potential-of-plastic-insulation-to-realise-energy-savings-and-de-coupling-in-europe---january-2006.aspx>

Helmut Winterling, Nathalie Sonntag: Polystyrene Rigid Foam (EPS, XPS) (Kunststoffe 10/2011)
http://www.plasticsportal.net/wa/plasticsEU~en_GB/function/conversions:/publish/common/upload/technical_journals/plastics_trendreports/Rigid_polystyrene_foam.pdf

COM(2011) 581 final REPORT FROM THE COMMISSION On the application, effects and adequacy of the Regulation on certain fluorinated greenhouse gases (Regulation (EC) No 842/2006)
http://ec.europa.eu/clima/policies/f-gas/docs/report_en.pdf

Meeting Report of the Joint IPCC/TEAP Expert Meeting on Options for the Limitation of Emissions of HFCs and PFCs Petten. The Netherlands 26-28 May 1999
<http://www.unep.ch/ozone/Joint-IPCC-TEAP-HFC-PFC.shtml>

ISOPA- Socio- Economic Information on the European Polyurethanes Industry
<http://www.isopa.org/isopa/uploads/Documents/documents/Soc-Econ%20Final%206-10-2003.pdf>

AFEAS (Alternative Fluorocarbons Environmental Acceptability Study). 1998: Production, Sales and Atmospheric Release of Fluorocarbons Through 1997. AFEAS. Washington, DC.

UNEP 2011. HFCs: A Critical Link in Protecting Climate and the Ozone Layer. United Nations Environment Programme (UNEP). 36pp
Available on-line;http://www.unep.org/dewa/Portals/67/pdf/HFC_report.pdf

Personal communication to Bodnár György (Hungarian Association of EPS Insulating Foam Producers) and Borzák Balarám Béla (building insulation architecture expert) and Buzási Vera (Hungarian Plastic Industry Association)

2.F.4. method references

General info on medical aerosols: IPCC/TEAP Special Report: Safeguarding the Ozone Layer and the Climate System. 2005. Chapter 8: Medical Aerosols.

Prevalence of asthma:

Masoli. M.. Fabian. D.. Holt. S. and Beasley. R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. Allergy. 59. 469–478 (2004)

German data (for benchmark): The time series has been published until 2008 so far: <http://www.umweltdaten.de/publikationen/fpdf-l/3999.pdf> (page 37)

Approach: Schwarz. W.; Gschrey. B.; Leisewitz. A.; Herold. A.; Gores. S.; Papst. I.; Usinger. J.; Oppelt. D.; Croiset. I.; Pedersen. P.H.; Colbourne. D.; Kauffeld. M.; Kaar. K.; Lindborg. A.: Preparatory study for a review of Regulation (EC) No. 842/2006 on certain fluorinated greenhouse gases. September 2011. The approach is described in the Annex volume (p.142): http://ec.europa.eu/clima/policies/f-gas/docs/2011_study_annex_en.pdf

5 AGRICULTURE (CRF sector 3)

5.1 Overview of sector

Agriculture production contributed to the greenhouse gas emission through the following processes:

- 3.A Enteric Fermentation by domestic livestock (CH₄).
- 3.B Manure Management (CH₄ and N₂O).
- 3.C Rice Cultivation (CH₄).
- 3.D Agricultural Soils (N₂O).
- 3.F Field Burning of Agricultural Residues (CH₄ and N₂O).
- 3.G Liming (CO₂).
- 3.H Urea application (CO₂).
- 3.I Other carbon containing fertilizers (CO₂).

Category 3.E Prescribed Burning of Savannas is not relevant to Hungary therefore notation key 'NO' is used relating to all associated emissions in the CRF Tables. In spite of this the NIR contains a chapter on 3.E, following a recommendation from the UNFCCC annual review conducted in 2013.

The main greenhouse gas emissions from Agriculture are CH₄ and N₂O. Although CO₂ emissions from carbonate containing materials are also reported in the Agriculture sector, these emissions are less significant compared with non-CO₂ emissions. Other CO₂ emissions associated with agricultural production as energy consumption of agricultural activities (heat production, agricultural vehicles, and machinery) are reported in the Energy sector (1.AA.4C Energy, Agriculture/Forestry/Fishing), while CO₂ emissions from agricultural soils are included in the LULUCF sector.

To give an overview of Hungarian agriculture the main characteristics are as follows:

Due to national conditions agriculture played a definitive role in the Hungarian economy in the past and even today. The share of agriculture in the GDP was 4.1 per cent in 2020 (HCSO, 2021). The agricultural land area was 53 per cent of the total (HCSO, 2021b). According to the data of the General Agricultural Census, 2020 (HCSO, 2021c),

According to the 2020 General Agricultural Census, the number of agricultural holdings in 2020 was 241,000 a reduction of 31% since 2010, with those using agricultural land less than those keeping livestock. In the crop structure, the share of cereals has decreased and that of industrial, fodder crops and vegetables has increased. The largest share of agricultural land was used by those with 5 hectares or more but less than 300 hectares. The reduction in subsistence farming has led to a concentration of livestock production. Between 2010 and 2020, the cattle population increased by 32%. The number of pigs has fallen by 7.4% compared to 2010, and the sow herd has shrunk by a quarter. There are still many farms with at most a few poultry, but production is more concentrated than in 2010.

The main characteristics for current trends are as follows:

In Hungary, agricultural production practically stopped growing in the late 1980's. This was followed by a dramatic drop in the 1990s, as a result of the economic and political transition taking place in the country. The gross value of agricultural production decreased, by 20 to 40 per cent from the level of the 1980s. The drop was smaller for crop production (10-30%) than for animal husbandry. The output of the latter was only two thirds or less of the level of 1990 (Laczka and Soós, 2003). The volume index of gross agricultural production in 1993 reached a minimum of 69.1 per cent of 1990 level. The crop production has fluctuated considerably since 1993. It fell in 2002-2003 and 2007 due to drought. In contrast, the agricultural production was relatively high due to the significantly high crop production in 2004 and 2008. Animal husbandry remained at a low level between 1993 and 2004 and has been decreasing steadily since the year of the European Union accession (2004) (Laczka, 2007). In recent years swine population decreased furtherly, while cattle population increased as a result of the state incentives to promote the recovery of livestock sector. In 2020 the gross production of agriculture decreased by 2.4%. The largest falls in production were in fruit crops, fodder crops and in animal livestock. (HCSO, 2021)

5.1.1 Emission trends

In 2020, the agriculture sector contributed 12% to Hungary's total GHG emissions (excluding LULUCF). The trend in emissions (Figure 5.1.1) shows a decrease of 39.3% over the period BY-2020 as a result of a drop in activity data (Figure 5.1.2). The bulk of this decline occurred between 1985 and 1995, when agricultural production fell by more than 30 per cent, and livestock numbers underwent a drastic decrease. Between 1996 and 2008, agricultural emissions had stagnated around 6.2 Mt CO₂-eq with fluctuations of up to 5%. Agricultural emissions decreased both in 2009 and 2010, hitting the lowest point in 2010. There was a slight increase in emissions in 2011 reflecting the higher fertilizer use and crop production. Since 2012 agricultural emissions has increased mainly, due to the increasing cattle livestock, milk production and fertilizer use. The increasing N-input from crop residues also contributed to the upward trend in emissions.




Figure 5.1.1 Trends in emissions from Agriculture BY-2020

Note: emissions from 3.C, 3.F, 3.G, 3.H and 3.I are small, but not zeros.




Figure 5.1.2 Main drivers of Agricultural emissions BY-2020

5.1.2 Emission trends by gas

From the BY to 2020, CH₄, N₂O and CO₂ emissions from agriculture decreased by 50, 29 and 41 per cent, respectively.

The decrease in CH₄ emissions is even more significant than N₂O, because CH₄ emissions are driven by the type and numbers of livestock. In Hungary's case, the amounts are largely determined by cattle and swine population. In 2020 cattle and swine accounted for 76 and 13% of combined total of emissions of CH₄ from enteric fermentation and manure management, respectively. After the sudden drop of livestock population at the beginning of the '90s it remained at that low level. Thus, CH₄ emissions had dropped by 46% from the base year level of 5,538 Gg CO₂-eq to 3,000 Gg CO₂-eq in 1994, when reached a plateau. In 2004, which is the year of the European Union accession for Hungary, animal livestock started to decrease moderately again, leading to the lowest level of CH₄ emissions at 2,370 Gg CO₂-eq in 2010 represent a reduction of 57% on the level of the BY. Since 2012 cattle populations increased resulting in a moderate increase in the emissions to 2,755 Gg CO₂-eq in 2020.

Emissions of N₂O show similar trends to those of CH₄ because the change of the regime resulted in a significant reduction in emissions. Agricultural N₂O emissions were 6,085 Gg CO₂-eq in the BY and decreased by 54% to 1993 to reach the lowest level in emissions at 2,811 Gg CO₂-equivalent. But unlike the livestock sector, there was a slight recovery in the crop production and nitrogen fertilizer use at the beginning of the second half of the 90s, resulting in a moderate increase in the emissions in the period between 1993 and 2004. Subsequently, in spite of the slightly increasing trends in nitrogen fertilizer use, N₂O emissions fluctuated, rather than increased, because the effect of the decreasing animal livestock overbalanced the increasing emissions from synthetic fertilizers. As a result, emissions amounted to 4,303 Gg CO₂-eq in 2020, representing a reduction of 29% cent on the BY level. N fertilizer use produces the bulk of agricultural N₂O emissions (28 per cent of the total emissions of the Agriculture in 2020).

Agricultural CO₂ emissions have dropped by 41% over the inventory period, which is the effect of the fall in urea use and liming. However, Agricultural CO₂ emissions are of low importance in the overall emissions, accounting for 0.5 per cent of the national total CO₂ emissions (excluding LULUCF).

The trends in emissions by gas are presented in Table 5.1.1 . Trends by gas and sub-categories are shown in Figure 5.1.3, Figure 5.1.4 and Figure 5.1.5.

Table 5.1.1 Emissions of CH₄, N₂O and CO₂ from Agriculture BY-2020

Year	GHG emissions (Gg)		
	CH ₄	N ₂ O	CO ₂
BY	222	20	407
1990	197	16	385
1991	185	12	274
1992	158	10	237
1993	136	9	195
1994	120	10	159
1995	118	10	117
1996	117	10	105
1997	114	10	95
1998	114	11	96
1999	117	11	95
2000	115	11	104
2001	111	12	108
2002	111	12	117
2003	111	11	127
2004	106	12	151
2005	104	11	142
2006	100	12	144
2007	101	11	149
2008	98	12	91
2009	96	11	98
2010	95	11	106
2011	94	11	129
2012	98	11	142
2013	99	12	174
2014	102	13	169
2015	105	13	203
2016	107	14	209
2017	107	14	223
2018	109	14	214
2019	109	14	217
2020	110	14	239
Share of Hungarian total in BY	41%	55%	0.5%
Share of Hungarian total in 2020	34%	86%	0.5%
Trend BY-2020	-50%	-29%	-41%




Figure 5.1.3 CH₄ emissions from Agriculture BY-2020

Note: emissions from 3.C and 3.F are small, but not zeros




Figure 5.1.4 N₂O emissions from Agriculture BY-2020

Note: emissions from 3.F are small, but not zeros




Figure 5.1.5 CO₂ emissions from Agriculture BY-2020

5.1.3 Emission trends by sub-category

Agricultural GHG emissions amounted to 12,030 Gg CO₂-eq in the BY and 7,297 Gg CO₂-eq in 2020, which means a reduction of 39%. Total emissions from the Agriculture sector in 2010 at 5,655 Gg CO₂-eq was the lowest level in the whole time series.

Table 5.1.2 shows the trends in GHG emissions by source categories as well as their contribution to the overall national emissions (excluding LULUCF). The most important category is 3.D Agricultural Soils at 6.2%, followed by 3.A Enteric Fermentation at 3.3% and 3.B Manure Management accounting for 1.7% of national total GHG emissions in 2020. CRF category 3.C Rice Cultivation accounts for less than one-tenth of a per cent of the national total. As it reveals from the Figure 5.1.3-Figure 5.1.5 emissions from all categories are decreasing except 3.I. The reason for the declining trend in the emissions from Enteric Fermentation and Manure Management is the decrease in livestock numbers, especially cattle and swine. The total emissions from the Livestock, which is equal to the combined total of emissions from 3.A Enteric fermentation and 3.B Manure management (Indirect emissions included) expressed in CO₂ equivalents was 6,337 Gg CO₂-eq in the BY. This decreased by 45 per cent to reach 3,496 Gg CO₂-eq in 1994 and subsequently decreased by 5 per cent to 3,169 Gg CO₂-eq in 2020. Livestock accounted for 43% of GHG emissions in agriculture in 2020. The biggest contributor to Livestock emissions is cattle, with 31% share of agricultural total emissions.

Over the period BY-1993 emissions from 3.D Agricultural soils had dipped sharply from 5,144 Gg CO₂-eq in the base year to 2,223 Gg CO₂-eq represents a decrease of 53% from the BY to the 1993 level, when the state subsidies on fertilizers were halted and the amount of animal manure decreased due to the decreasing animal livestock. Emissions totaling 3,868 Gg CO₂-eq in 2020 represent a reduction of 25% from the base year level. The slight increase in emissions from agricultural soils over the period 1993-2020 is a result of compensatory processes, the slight and steady increase in emissions from nitrogen fertilizers were partly overbalanced by the decreasing emissions from organic manure. N₂O emissions from 3.D in the BY and 2020 accounted for 43% and 53%, respectively of the total agricultural emissions, reflecting the restructuring of the Hungarian agriculture.

The trends of emissions from liming, urea application and carbon-containing fertilizers slightly differ from of the other sectors. Emissions from all of the CO₂ relevant sectors decreased at the beginning of the time series, but emissions from urea application and carbon-containing fertilizers started to increase at the end of the '90s due to the increasing fertilizer use, while emissions from liming after a slight increase to mid-2000s decreased again continuously. The reason for the second drop in the emissions from urea use was the economic crisis in 2008, when the price of the urea increased thus the urea application fell sharply. Emissions from urea and carbon-containing fertilizers increased again over the period 2009-2020.

Table 5.1.2 GHG emissions BY-2020 from agriculture by subcategories

Year	GHG emissions (Gg CO ₂ -eq)								
	3	3.A	3.B	3.C	3.D	3.F	3.G	3.H	3.I
BY	12,030	4,151	2,186	81	5,144	60.73	130	229	48
1990	9,994	3,668	2,027	81	3,833	1.11	191	171	23
1991	8,469	3,480	1,866	61	2,788	0.59	188	80	7
1992	7,249	3,015	1,583	34	2,380	0.43	163	57	16
1993	6,395	2,548	1,395	34	2,223	0.37	128	55	12
1994	6,287	2,248	1,248	34	2,598	0.43	100	50	8
1995	6,006	2,202	1,240	27	2,421	0.35	58	51	8
1996	6,027	2,168	1,270	21	2,463	0.20	47	49	9
1997	6,002	2,116	1,233	15	2,542	0.20	38	51	7
1998	6,195	2,088	1,270	15	2,726	0.21	34	56	6
1999	6,264	2,111	1,319	15	2,724	0.21	24	60	10
2000	6,141	2,084	1,299	22	2,631	0.29	24	65	15
2001	6,335	2,002	1,267	16	2,942	0.21	23	69	16
2002	6,364	1,978	1,304	14	2,950	0.27	23	76	18
2003	6,190	1,957	1,316	17	2,771	0.31	30	80	17
2004	6,459	1,902	1,230	19	3,157	0.27	34	91	26
2005	6,133	1,884	1,173	18	2,915	0.26	33	87	22
2006	6,093	1,813	1,145	16	2,975	0.22	28	91	24
2007	6,063	1,819	1,152	18	2,925	0.28	24	95	31
2008	6,096	1,791	1,101	17	3,096	0.28	17	46	27
2009	5,732	1,764	1,036	18	2,815	0.33	16	43	39
2010	5,655	1,756	1,028	14	2,752	0.06	14	46	45
2011	5,864	1,748	1,012	18	2,956	0.25	18	58	54
2012	5,901	1,819	1,030	20	2,890	0.33	18	56	69
2013	6,294	1,870	1,020	18	3,212	0.30	18	79	77
2014	6,538	1,928	1,057	16	3,367	0.27	19	75	75
2015	6,752	1,978	1,088	19	3,464	0.26	18	97	87
2016	7,068	2,022	1,084	20	3,733	0.32	13	99	97
2017	7,071	2,030	1,059	19	3,740	0.34	11	113	99
2018	7,119	2,065	1,069	20	3,751	0.37	10	112	92
2019	7,113	2,072	1,071	18	3,735	0.31	8	116	93
2020	7,297	2,095	1,074	20	3,868	0.33	5	136	97
Share of Hungarian total in BY	10.9%	3.8%	2.0%	0.1%	4.7%	0.1%	0.1%	0.2%	0.04%
Share of Hungarian total in 2020	11.6%	3.3%	1.7%	0.0%	6.2%	0.0%	0.01%	0.2%	0.2%
Trend BY-2020	-39%	-50%	-51%	-75%	-25%	-99%	-96%	-40%	103%

5.1.4 Key Categories

Key category analysis is presented in Chapter 1.6. Table 1.2 contains the key categories of the agriculture sector.

5.1.5 Methodological issues

Methodologies of the 2006 IPCC Guidelines have been implemented throughout the agricultural inventory. However, in some cases, where there are gaps in the 2006 IPCC Guidelines, methodologies provided in the 2019 Refinement were applied.

IPCC Tier 2 methods were used for the following categories:

- 3.A Enteric Fermentation in Cattle.
- CH₄ emissions from 3.B Manure Management associated with all livestock categories, except Rabbits.
- N₂O emissions from 3.B for Cattle, Swine, Poultry (Laying hens and Broilers) and Indirect emissions.
- N₂O emissions from 3.D.1.5 Mineralization/immobilization associated with loss/gain of soil organic matter.
- Indirect N₂O emissions from 3B and 3D are reported in line with the national NH₃ and NO_x inventories, which also meets the requirements of a Tier 2 methodology.

For the other categories IPCC Tier 1 methods were applied. Country-specific emission factors were used whenever sufficient information was available, otherwise the IPCC default factors were used. See the individual categories for further details.

5.1.6 Uncertainties and time series consistency

The following chapter gives an overview of uncertainty estimates for CH₄, N₂O and CO₂ emissions from Agriculture.

Uncertainty estimates were performed using the Tier 1 approach based on the error propagation. Error propagation was calculated independently for the lower (2.5 percentile) and for the upper (97.5 percentile) range to treat the asymmetric confidence ranges. Uncertainties were combined in accordance with 2006 IPCC Guidelines Equation 3.1 and 3.2. The results of the Tier 1 approach are shown in Table 5.1.4.

The uncertainty of the activity data was calculated on the basis of the available data of the Hungarian Central Statistical Office (hereafter HCSO), the 2019 EMEP/EEA Guidebook and expert judgement; the uncertainty of the emission factors was calculated following the 2006 IPCC Guidelines. The uncertainty of the livestock population data for 2020 is presented according to the uncertainty assessment of the HCSO, in Table 5.1.3. The overall weighted mean of the uncertainties in the livestock population is ± 2.3 per cent. The uncertainty in the Swine population is the lowest at 0.6 per cent, while the uncertainty in Mules and Asses populations is the highest at approximately 48.5 per cent.

In the Hungarian agricultural GHG inventory, the uncertainties of N₂O emissions from agricultural soils are the highest. These high values derive from the uncertainties of the emission factors. The uncertainty and the distribution of these emission factors (EF₁, EF₄ and EF₅) strongly influence the uncertainty and the distribution of the agricultural emissions as well as the overall uncertainty of the Hungarian GHG inventory. For these emission factors default confidence limit ranges and lognormal distributions have been applied according to the 2006 IPCC Guidelines.

Table 5.1.3 Uncertainty of animal population data for 2020 (HCSO)

Livestock categories	2019 Dec	2020 Jun	2020 Dec	Annual mean	Uncertainty of the annual mean u(AD _i)	Weighted annual mean
95% Confidence Interval (+/- 1,000 head)					%	1,000 head
Dairy Cattle	16.00	7.35	12.00	6.21	2.60	238
Non-Dairy Cattle	32.90	16.85	24.68	13.29	1.93	689
Buffalo	0.30	0.50	0.38	0.28	3.58	8
Sheep	115.30	90.68	103.77	59.66	5.98	998
Goats	9.60	12.65	11.04	7.31	13.02	56
Horses	6.40	7.75	8.00	4.64	8.01	58
Mules and Asses	2.80	3.63	3.50	2.13	48.53	4
Swine	41.80	20.40	31.35	16.57	0.59	2831
Poultry	1,440.21	1,437.27	1,512.22	888.26	2.35	37835
Rabbit	21.90	32.55	23.00	18.11	1.49	1218
Overall (weighted mean)					2.31	

Table 5.1.4 Uncertainties of activity data, emission factors and emissions for key and particularly significant* categories by Tier 1 approach

3 Agriculture	GHG	Uncertainty of activity data	Uncertainty of Emission Factor	Combined uncertainty of emissions
		%		
3.A Enteric Fermentation	CH ₄	±0	±13	±13
3.A.1 Enteric Fermentation/ Cattle	CH ₄	±0	±14	±14
3.A.2 Enteric Fermentation/ Sheep	CH ₄	±7	±40	±41
3.B Manure Management	CH ₄	±0	±14	±14
3.B.1 Manure Management/ Cattle	CH ₄	±0	±14	±14
3.B.3 Manure Management/ Swine	CH ₄	±1	±30	±30
3.B Manure Management	N ₂ O	±0	-36/+131	-36/+130
3.B.13 Manure Management/ Other	N ₂ O	±21	-50/+100	-54/+102
3.B Manure Management/ Indirect	N ₂ O	±0	-84/+399	-84/+399
3D Agricultural Soil Emissions	N ₂ O	±0	-66/+187	-66/+187
3.D.a.1 Direct Soil Emissions/ Synthetic Fertilizer	N ₂ O	±5	-70/+200	-70/+200
3.D.a.4 Direct Soil Emissions/ Crop residues	N ₂ O	±25	-70/+200	-74/+202
3.D.3 Indirect Emissions	N ₂ O	±0	-71/+272	-71/+272

*Note: In accordance with the 2006 IPCC Guidelines particularly significant categories are those which contribute together more than 60% to the key category.

5.1.7 Quality Assurance and Quality Control

The agricultural greenhouse gas inventory is compiled by the HMS in strong cooperation with the Institute of Agricultural Economics Nonprofit Kft. (AKI). The used activity data is mainly derived from the official database of the HCSO, in cases where HCSO's data are not available the EUROSTAT's, NFCSO's or the Research Institute for Agricultural Economics' data are applied.

Data and documentation are archived by the Unit of National Emissions Inventories of the Hungarian Meteorological Service. The annual sector specific QA/QC procedures are as follows:

- Check of activity data for transcription and rounding errors, comparison with original data sources.
- Re-check of activity data, comparison with the latest submission of the activity data (following the revision of the data by data supplier).
- Check of reasons for data gaps.
- Verification of activity data with other data sources if it is possible.
- Consistency check of time series of the activity data and the estimated emissions (reasons for jumps).
- Consistency check, following the methodological changes of the data collection.
- Check of the time series consistency of the applied livestock characterization.
- Cross-check of data sources of the activity data if it is possible (e.g., total annual milk yield per cow, and total dairy cow population).
- Cross-check of the applied activity data between the different sub-categories.
- Check of emission factors, comparison with the IPCC default ones and comparison with the values applied by other countries (especially EU member states) according to the EU's NIR and S&A report of the UNFCCC.
- Check of the methodologies used for the development of county-specific emission factors, comparison with the IPCC methodologies or other methodologies if it is available.
- Check of the correct use of the units in the calculation sheets.
- Check for transcription errors between the calculation sheets and the CRF tables.
- Consistency check of sub-categories with totals.
- Check of recalculation differences.
- Listing of QA/QC findings and the actions taken in the spreadsheets.
- Recording of sources of activity data and equations in the spreadsheets.

Details of other source-specific quality checks can be found in the respective sub-chapters.

Since 2011 the Unit of National Emissions Inventories of the HMS has also been participated in the preparation of the Air Pollution Emission Inventory under the Convention on Long-range Transboundary Air Pollution of the United Nations Economic Commission for Europe (UNECE/LRTRAP). (As a party to the UNECE/LRTRAP Convention Hungary is required to report annually data on emissions of air-pollutants covered in the Convention.) This provides an additional opportunity to cross-check the activity data and emissions with the GHG-inventory to ensure the consistency between the two inventories.

Hungary as a member state of the EU has additional reporting obligations (e.g., Nitrate Directive and Nutrient Balance) arising from different Community policies. In some cases, the same data and

coefficients are required for the background calculation of these reports. As an additional QA procedure, these data and methodologies are compared in the course of regular expert meetings.

Checks and reviews of national emission inventories reported by EU Member States under the Monitoring Mechanism Regulation (MMR) is also considered as a quality assurance activity.

In-depth reviews required by the 406/2009/EC EU Effort Sharing Decision are performed in every two years by external experts contracted by the EU, which covers the full inventory. First review was performed in 2012. The last comprehensive audit was in 2020, during which there were no recommendations for the agriculture sector.

All the recommendations from the 2021 UNFCCC review were implemented for the 2022 submission.

5.1.8 Recalculations

The main reason for changes in the emissions from 3. Agriculture sector are the manure management data updates. The 2020 Agricultural Census and the recently available annual statistics on agricultural wastes used in biogas plants have made it possible to reflect the current practices and the effects of anaerobic digestion. As a result of this inventory improvements animal manure used in “digesters” has been reported at the first time in this submission.

Data from the nitrate database, which aims to monitor the implementation of the European Nitrate Directive (91/676/EEC), also provides data on manure management, grazing and since 2015 on manure management related NH_3 abatement technologies. Data from the nitrate database was also used to revise the data on manure management for the GHG- and the air pollutant emission inventory. Revision of NH_3 emissions affected the direct and indirect N_2O emissions from manure management as well as agricultural soils.

Reallocation of manure to “digesters” for the period 2004-2019 and the change in the proportion of covered liquid slurry for the whole time series caused the most significant changes in CH_4 and N_2O emissions from 3. Agriculture.

The overall effect of recalculation in the 3. Agriculture sector resulted in an increase in the emissions in the range of 0.4 to 12.4 kt CO_2 eq (0.0%-0.1%) in the period BY to 2002 and a decrease in the range of 2.7 to 35.2 kt CO_2 eq (0.0%-0.5%) between 2003 and 2019. The largest decrease, 35.2 kt CO_2 -eq occurred in 2015 and the largest increase, 12.4 kt CO_2 -eq in the base year (BY).

Reasons for recalculations by CRF sectors are as follows:

3.A Enteric Fermentation

3.A.1 Enteric Fermentation -Dairy Cattle, 2004-2019

The HCSO has changed the methodology of data collection on milk fat and milk protein since 2019 and 2020, respectively. This change led to time series inconsistency, as the HCSO has not revised the data, retrospectively in line with the new methodology. To ensure the time series consistency the milk fat and protein content data were revised based on the Institute of Agricultural Economics' Market Price Information System (MPIS) data. The revised nutritional values of raw milk used to derive the gross energy intake (GE) for dairy cattle, resulted in an increase of 0.15% and 1.1 kt CO_2 -eq on average in the CH_4 emissions from 3.A.1 Enteric Fermentation - Dairy Cattle for the

2004-2019 trend. The change ranged between -5.2 kt CO₂-eq in 2004 and 7.7 kt CO₂-eq in 2016.

3.A.1 Enteric Fermentation - Non-Dairy Cattle, heifers 1-2 yr, 2016-2019

A data linking error was corrected, because in the previous submissions the livestock number of heifers, 1-2 yrs for the year 2015 was used for the period 2016-2019 as well. Due to this recalculation emissions increased by 0.4%-2.2% for the period 2016-2019.

The overall effect of recalculations on 3.A Enteric Fermentation was a slight decrease in the emissions between 2004 and 2009 (except 2005) and increase in 2005 and over the period 2010-2019. The increase is the most significant (23.8 kt CO₂-eq) in 2019.

Revisions made in Sector 3.A, through gross energy intake for dairy cows and animal numbers for other cattle, also affected emissions from manure management.

3.B.1 CH₄ Emissions from Manure Management

3.B.1.1 and 3.B.1.3 CH₄ Emissions from Manure Management - Cattle and Swine, across the whole time series

Proportions of pig and cattle liquid/slurry covered by “natural crust” was revised for the whole time series. The former expert judgement was replaced by annual survey data, which are available from the year 2015. Between 2001 and 2015, interpolation was used for cover types other than natural crust; and for natural crust, 2015 data were used for the period before 2015, as well. This recalculation resulted in an increase in the MCF of liquid/slurry of cattle and swine. The increase is larger for cattle and minor for pigs.

3.B.1.1, 3.B.1.3 and 3.B.1.4 CH₄ / 3.B.2.1, 3.B.2.3 and 3.B.2.4 N₂O Emissions from Manure Management - Cattle, Swine and Poultry for the period 2004-2019

CH₄ and N₂O emissions from manure management of cattle, swine, and poultry for the period 2004-2019 was revised considering proportions of manure used in anaerobic digesters. In the previous submissions the anaerobic digested manure was allocated in line with the housing, irrespective of the manure use. In this submission, the anaerobic digested manure was reallocated to ‘digesters’, as data on the agricultural wastes used in anaerobic digesters had become available.

3.B.1.2 CH₄ Emissions from Manure Management - Sheep, 2014-2019

There are minor recalculations to CH₄ emissions from manure management of sheep due to the revision on the proportion of manure excreted during grazing considering the length of grazing period. This revision resulted in an insignificant (less than 0.5 kt CO₂-eq) increase in the CH₄ emissions from 3.B.1.2.

The overall changes in the CH₄ emissions from 3.B.1. Manure management ranged from a decrease of 25.5 kt CO₂-eq (4.1%) in 2013 to an increase of 15.7 kt CO₂-eq (1.3%) in the BY. Revision of “natural crust” resulted in an increase in the BY emissions, 3.0, 10.9 and 1.7 kt CO₂-eq for dairy cattle, non-dairy cattle, and swine, respectively. In contrast, reallocation of anaerobic digested manure to “digesters” resulted in a decrease in the emissions for the period 2004 to 2019, partly offset by the increase in emissions due to revised natural crust. The

highest net decrease in the emissions due to anaerobic digestion were 9.4 kt CO₂-eq (in 2013), 5.3 kt CO₂-eq (in 2019) and 17.1 kt CO₂-eq (in 2014), for dairy-cattle, non-dairy cattle, and swine, respectively.

3.B.2 N₂O emissions from Manure Management

3.B.2 Direct N₂O emissions from manure management, whole time series

Revision of “natural crust” for the whole time series and the anaerobic digested manure for the period 2004-2019 resulted in an overall decrease of 1.3% and 4.1 kt CO₂-eq on average on the BY and 1990-2019 trend. Decrease in the emissions ranged from 1.1 kt CO₂-eq in 2000 to 18.3 kt CO₂-eq in 2019.

3.B.2.5 Indirect N₂O emissions from manure management, whole time series

Atmospheric Deposition, whole time series

Revision of NH₃ emissions from 3.B Manure Management for the reporting to the UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP) (UNECE, 1999) and the National Emissions Ceiling Directive (EP, CEU, 2016) resulted in changes in the indirect N₂O emissions from manure management due to atmospheric deposition. The revision of NH₃ emissions aimed to take account of emission abatement measures, as a result of new data collections recently launched.

The following abatement measures were considered in the revision of NH₃ emissions by stages of manure and animals:

Animal housing

- Non-leaking drinking systems for broilers
- Aviary system for layers
- Partially slatted floor for piglets after weaning and growers-finishers

Manure storage

- “Low technology” floating covers (e.g., chopped straw, peat, bark, etc.)
- “Tight lid”, roof or tent structure
- Plastic sheeting (floating cover)
- Natural crust

Revision of the NH₃ emissions from manure management lead to an 1.1% (1.5 kt CO₂-eq) decrease in indirect N₂O emissions due to atmospheric deposition on average over the time series.

Nitrogen Leaching and Run-off, 2004-2019

Revision of data on manure management system distribution, mainly reallocation of manure to digesters, resulted in additional changes in the indirect N₂O emissions due to leaching and run-off. Changes ranged from a decrease of 0.08 kt CO₂ eq in 2019 (6.4%) to an increase of 0.02 kt CO₂ eq in 2016 (1.3%).

The overall indirect N₂O emissions from manure management decreased by 1.1% and 1.5 kt CO₂ eq on average in the BY and 1990-2019 trend.

The overall impact of revisions for the 3.B Manure Management sector CH₄ and N₂O emissions is an increase in emissions in the range of 1.2-13.3 kt CO₂-eq (0.1%-0.7%) between BY and 2005, and a decrease of between 0.3 and 41.8 kt CO₂-eq (0.0%-3.8%) for the years 2006-2019.

3.D Agricultural Soils, whole time series

3.D.1 Direct N₂O Emissions from Managed Soils, whole time series

3.D.1.2.a Animal Manure Applied to Soils

Due to the interlinking between the 3.B and 3.D CRF sectors through the N-balance, revisions to the 3.B resulted in revised emissions from 3.D.1.2a Animal manure applied to soils and 3.D.2 Indirect N₂O emissions from agricultural soils for the whole time series. Correction to the livestock number of “heifers 1-2 yrs” for the years 2016-2019, as mentioned in the paragraph related to the CRF Sector 3.A, resulted in some increase in the F_{AM}. The net effect of these recalculations is an insignificant reduction in the direct emissions from 3.D.1.2.a Direct N₂O emissions from agricultural soils - animal manure applied to soils over the period BY-2008 and an increase ranging from 0.02 kt CO₂ eq to 8.4 kt CO₂ eq over the period 2010-2019.

3.D.1.2.c Other Organic Fertilizers Applied to Soils

Data on composted sewage sludge and municipal waste have been revised to include only use on agricultural land. In the previous submissions, the total amount of composted sewage sludge and waste was used, which was not in line with the Hungarian regulation, because in Hungary, the quality of compost that can be applied is regulated by law, and application is subject to a permit, so not all compost can be used on agricultural land without restriction.

At the same time, as a result of our improvements in the utilization of agricultural waste in anaerobic digestion, digestate from anaerobic digestion, which was not included in previous submissions, is now included.

As a result of the revision of activity data, emissions from 3Da2c decreased by -0.5 to 4.4 kt CO₂ eq between BY and 2009, while they increased by 0.1 to 9.2 kt CO₂ eq between 2010 and 2019.

3.D.1.3 Urine and Dung Deposited by Grazing Animals

The update of the AWMS data also resulted in a minor change in the N excreted on grazing (F_{PRP}) for the period 2014-2019. Revision of F_{PRP} resulted in changes in the emissions ranging from -6.1 kt CO₂ eq (4.8%) in 2016 to 4.8 kt CO₂ eq (3.6%) in 2019.

3.D.1.4 Crop Residues

The AWMS data update for the period 2014-2019, also resulted in a change in N input from crop residues (F_{CR}) due to a revision of the straw used for bedding. Recalculation to the F_{CR} resulted in changes in the emissions ranging from -0.03 kt CO₂ eq in 2016 to 2.3 kt CO₂ eq in 2019.

The net effect of recalculations for 3.D.1 is a change in emissions ranging from -5.6 kt CO₂ eq in 2003 to 13.3 kt CO₂ eq in 2018.

Due to revisions in N inputs from the above-mentioned sources-categories 3.D Direct N₂O emissions from Agricultural Soils N₂O emissions decreased by 0.5-5.6 kt CO₂ eq between BY and 2009 and increased by 0.3-13.3 kt CO₂ eq over the period 2010-2019.

3.D.2 Indirect N₂O Emissions from Managed Soils, whole time series

The above-mentioned recalculations in 3.D.1 Direct N₂O Emissions from Managed Soils, which are not detailed again, have also led to changes in indirect N₂O emissions from agricultural soils due to atmospheric deposition and nitrogen leaching and run-off.

The recalculation of NH₃ emissions from 3.D also contributed to the changes in indirect N₂O emissions from agricultural soils due to atmospheric deposition. The main driver behind the decrease in the emissions in the period 2001-2019 is accounting for NH₃ abatement technologies in solid manure application. NH₃ emission abatement technologies for slurry application have already been accounted for in the previous submissions of the air pollutant as well as the GHG-emission inventory. While the manure NH₃ abatement technologies for solid manure application are reported for the first time in this year's submissions. In Hungary, measures on manure incorporation entered into force in 2001, so these technologies were taken into account in the estimation of NH₃ emissions from this date. The reduced NH₃ emissions resulted in a reduction of indirect N₂O emissions due to atmospheric deposition ranging from 0.09 kt CO₂ eq (0.6%) in 2002 to 15.2 kt CO₂ eq (8.7%) in 2019. In contrast, over the BY and 1990-2000 period, due to the decreased N-loss from the animal manure the emission increased, slightly. This increase is less than 2 kt CO₂ eq.

Changes in emissions from nitrogen leaching and run-off because of the aforementioned revisions are negligible (less than 0.3 kt CO₂ eq).

The net effect of these recalculations are changes in 3.D.2 Indirect N₂O emissions from agricultural soils ranging from -14.9 kt CO₂ eq in 2019 to 1.9 kt CO₂ eq in the BY.

The overall effect of recalculations for 3.D Agricultural Soils was a decrease ranging up to 8.7 kt CO₂ eq (0.3%) in 2012, except 2013, when emissions increased negligibly.

5.1.9 Planned improvements

Participation in the EU review mechanisms, which is part of the QA/QC processes for compiling EU inventory, provides an opportunity for examination of individual IPCC sectors and particular issues relating to methodologies, country-specific emission factors and coefficients. Issues of planned improvements will be assigned largely in accordance with the outcome of the EU and UNFCCC review processes.

N-excretion rates for the main animal categories as Cattle, Swine and Poultry also planned to be updated depending on the availability and processing of the data from the animal feeding monitoring program. In parallel, we also plan to revise the volatile solid excretion rates for pigs and poultry, in line with the data from the feed monitoring program.

5.2 Enteric fermentation (CRF sector 3.A)

Enteric fermentation in animals is considered as significant source of CH₄. The most important process of generation is anaerobic cellulose degradation in the rumen of ruminants. Some CH₄ is generated in the colon of horses and rabbits, and in the caecum of poultry. In Hungary, the leading CH₄ emitters are cattle and sheep, with the most important category being non-dairy cattle. In addition to the number of animals, the level of production and feeding practices are the factors which primarily influencing the amount of CH₄ from enteric fermentation.

In 2020 76% of the total CH₄ emissions from agriculture derived from this source category.

5.2.1 Source Category Description

Emitted gas: CH₄

Methods: T1, T2

Emission factors: D, CS

Key source: Yes

Particularly significant sub-categories: Cattle

Figure 5.2.1 presents the estimates of CH₄ emissions for 3.A *Enteric Fermentation* by livestock categories. Emissions amounted to 166 Gg in the base year and have reduced by 51 per cent to 84 Gg in 2020 due to the decrease in cattle livestock. The bulk of this decrease occurred between 1985 and 1994, during which Hungary experienced a period of unprecedented drop in the agricultural production resulting in a dramatic decrease in animal populations. Despite the continuous decrease in the livestock populations, emissions stagnated in the years between 1995 and 2000, because the improving cow productivity overbalanced the effect of declining cattle population (Figure 5.2.2). In the period 2000 to 2010 emissions slightly decreased again reaching their lowest level in 2010. This decrease reflects the further decline in Cattle livestock. Since 2011, emissions started to increase, following the slightly rising cattle population and milk production. Emissions from 3.A mostly depend on cattle population and milk production. Enteric fermentation in Cattle produced 82% of emissions from 3.A in 2020.




Figure 5.2.1 Trend in emissions from 3.A Enteric Fermentation by livestock categories BY-2020




Figure 5.2.2 Dairy Cattle population and daily milk production per cow BY-2020

5.2.2 Methodological issues

Emissions from enteric fermentation were calculated using the Tier 1 method of 2006 IPCC Guidelines, except for the Dairy Cattle and the Non-Dairy Cattle categories, where country-specific emission factors were used in accordance with the Tier 2 method of 2006 IPCC Guidelines.

5.2.2.1 Activity Data - Livestock Population

The HCSO provides national livestock survey data to the emission estimate.

Following a recommendation of the centralized review conducted in 2014 the HCSO has provide livestock data rounded to the nearest hundred instead of nearest thousand. The HCSO has been producing two censuses of animal numbers per year since 2009. One survey is conducted in June and the other in December. The annual average population for a year t was calculated by using the chronological mean of surveys data, as follows:

$$\text{NoA}_t = (0.5 * \text{NoA}_{\text{Dec},t-1} + \text{NoA}_{\text{June},t} + 0.5 * \text{NoA}_{\text{Dec},t}) / 2 \quad (\text{Equation 5.1.})$$

Where:

NoA_t = chronological mean of the annual population of a livestock category in a year t [1'000 head]

$\text{NoA}_{\text{Dec},t-1}$ = population of a livestock category in December of the year $t-1$ [1'000 head]

$\text{NoA}_{\text{June},t}$ = population of a livestock category in June of the year t [1'000 head]

$\text{NoA}_{\text{Dec},t}$ = population of a livestock category in December of the year t [1'000 head]

The method delineated above was suggested by the HCSO's expert (Tóth, 2004) to smooth out the seasonal changes in the livestock population.

Until the end of 2008 the HCSO collected data on animal livestock population three times a year, namely April, August and December. For the calculation of the annual average population for the years before 2009 the chronological mean was used similarly, based on the three surveys data.

The annual average livestock populations calculated as the *chronological means of the total animal populations* are reported in the CRF tables. Trends in livestock populations are provided in Table 5.2.1.

In response to a recommendation from the UNFCCC Review, 2019 data provided in Table 5.2.1 was cross-checked with the CRF Tables.

In the case of Non-dairy Cattle and Poultry enhanced characterization for livestock populations were used according to the requirements of the IPCC methodology. The annual average populations for these livestock were determined by sub-categories. Detailed livestock data by sub-categories for Cattle and Poultry are shown in Table 5.2.2 and Table 5.2.3.

Table 5.2.1 Livestock populations and trends BY-2020 (1'000 head)

Year	Dairy cattle	Non-dairy cattle	Sheep	Swine	Buffalo	Goats	Horses	Mules and Asses	Poultry	Other (Rabbit)
BY	589.5	1233.6	2498.3	8963.0	0.1	19.3	98.7	5.0	81738.8	2536.5
1990	563.6	1053.0	1958.3	8708.5	0.1	35.1	79.8	4.5	70325.6	2587.2
1991	526.6	1017.8	2008.7	7809.1	0.1	39.3	84.0	4.3	58827.4	2629.5
1992	479.5	833.9	1867.3	6237.4	0.1	50.0	78.8	4.3	52168.4	2389.5
1993	436.3	648.5	1457.7	5805.4	0.1	60.6	74.6	4.3	43429.1	2149.5
1994	408.5	554.4	1089.0	5006.9	0.1	71.3	85.1	4.3	44477.4	1909.4
1995	394.5	548.8	997.7	5023.0	0.2	76.1	74.6	4.3	44874.5	1669.4
1996	389.4	545.9	930.0	5493.5	0.3	80.9	73.5	4.3	38537.7	1148.9
1997	387.8	520.8	900.9	5012.7	0.4	85.7	75.6	4.3	40416.6	1071.3
1998	381.3	493.8	954.5	5246.7	0.5	90.5	76.7	4.3	42707.6	1051.8
1999	385.0	488.5	980.7	5609.0	0.6	95.3	77.7	4.3	40260.3	1040.4
2000	362.8	479.2	1192.2	5146.2	0.7	96.6	77.8	3.6	48562.1	942.5
2001	353.0	443.3	1162.8	4823.3	0.8	107.2	67.5	3.5	51074.0	1087.2
2002	344.5	433.7	1138.2	5050.0	0.9	96.7	63.2	3.4	51333.7	1179.7
2003	330.0	433.2	1226.5	5077.5	1.0	94.5	62.5	3.3	52486.2	1088.8
2004	309.3	424.3	1380.2	4385.0	1.1	84.5	64.5	3.2	50492.0	1181.7
2005	299.8	419.7	1446.7	4021.7	1.2	77.8	67.0	3.0	46404.7	1002.7
2006	275.2	428.2	1358.2	3943.7	1.3	81.2	64.8	2.3	44653.3	1084.3
2007	267.5	442.3	1300.7	4039.0	1.4	71.5	59.0	2.1	43159.7	1055.0
2008	263.8	436.2	1269.7	3664.7	1.4	72.8	58.3	2.0	45032.7	903.5
2009	257.5	444.3	1260.8	3248.0	1.5	65.0	59.8	1.9	44789.3	871.3
2010	244.5	454.0	1203.0	3208.0	2.5	79.3	65.5	3.1	46587.0	916.3
2011	250.6	440.2	1159.1	3131.3	3.7	83.8	73.0	3.5	46283.8	949.1
2012	256.0	474.8	1179.3	2981.5	3.4	86.0	76.2	3.5	43063.7	1367.1
2013	248.5	518.7	1204.9	2943.9	3.7	85.2	66.1	2.7	41674.3	1560.1
2014	252.0	538.5	1222.6	3064.9	3.7	76.7	63.0	2.1	42683.1	1643.2
2015	252.1	562.8	1193.9	3127.0	3.7	79.5	61.3	2.5	44459.1	1610.4
2016	247.0	592.2	1189.3	3020.8	5.4	84.0	56.5	3.2	44907.6	1300.4
2017	244.8	617.7	1160.3	2847.6	6.1	85.0	53.7	4.1	42711.1	1149.9
2018	242.7	635.8	1145.6	2865.1	6.6	75.3	51.9	4.6	43136.9	1204.6
2019	243.9	660.0	1100.3	2796.4	6.9	69.0	51.6	5.0	42875.1	1195.7
2020	238.4	688.7	997.9	2831.1	7.8	56.1	58.0	4.4	37835.1	1218.4
Trend BY-2020	-60%	-44%	-60%	-68%	7650%	190%	-41%	-13%	-54%	-52%
Trend 2005-2020	-20%	64%	-31%	-30%	546%	-28%	-13%	48%	-18%	22%

Table 5.2.2 Livestock population and trends for non-dairy cattle (1'000 head)

Year	<1 year		1-2 year		>2 year			
	Bovines for slaughter and other calves (male)	Bovines for slaughter and other calves (female)	Bovines (male)	Heifers for slaughter and other heifers	First calf heifers	Mature Non-Dairy (male)	Heifers for slaughter	Beef Cow
BY	256.9	264.3	226.0	277.7	72.0	20.4	19.6	96.8
1990	212.6	241.2	169.6	256.9	65.6	17.1	15.7	74.4
1991	204.7	237.8	162.2	251.9	61.8	16.4	14.9	67.9
1992	164.1	206.5	110.7	219.5	55.1	13.1	11.0	54.0
1993	128.7	162.9	86.2	170.9	44.7	9.7	7.0	38.5
1994	109.1	143.9	68.3	151.2	41.2	8.0	5.0	27.8
1995	107.4	143.4	65.9	149.1	42.7	7.9	4.9	27.5
1996	105.5	139.3	70.1	144.3	43.8	7.8	4.8	30.3
1997	99.5	133.0	63.5	138.8	47.3	7.4	4.3	27.0
1998	98.7	131.8	41.5	137.5	49.5	6.9	3.7	24.3
1999	97.4	130.1	47.8	135.7	44.3	6.8	3.6	23.0
2000	96.0	132.6	36.2	136.7	41.9	5.8	2.7	27.4
2001	88.0	125.9	29.4	131.4	37.1	4.8	2.7	24.0
2002	85.0	124.7	27.0	130.0	37.2	4.7	2.2	22.9
2003	87.8	121.4	26.6	124.4	36.0	4.5	2.3	30.1
2004	81.5	113.7	25.3	122.4	34.2	6.0	2.7	38.5
2005	84.7	109.2	22.6	119.1	32.8	5.8	2.0	43.4
2006	84.6	106.5	30.3	116.9	30.6	5.5	2.5	51.3
2007	86.6	106.2	37.0	116.4	33.0	6.2	2.2	54.7
2008	78.9	109.5	32.1	114.7	32.0	6.0	2.3	60.6
2009	81.5	108.2	31.7	120.2	32.5	6.5	2.0	61.7
2010	75.7	108.2	35.0	120.7	35.5	7.2	3.2	68.5
2011	74.5	105.6	26.4	115.7	35.6	7.0	2.6	72.8
2012	86.9	113.5	31.8	117.5	35.6	7.0	4.2	78.3
2013	89.1	119.6	41.5	130.1	35.3	8.2	4.3	90.5
2014	90.2	122.9	44.2	131.2	37.0	8.4	2.7	101.9
2015	90.3	129.9	43.2	135.7	38.3	9.2	3.7	112.5
2016	98.6	133.3	38.3	138.1	39.1	10.1	4.7	129.9
2017	104.0	138.9	37.8	137.4	37.8	10.5	5.2	146.2
2018	107.8	140.7	43.0	140.4	34.0	11.3	3.3	155.3
2019	107.4	146.7	44.9	146.4	33.3	11.1	4.3	166.0
2020	115.9	149.0	51.4	142.3	35.7	13.6	5.7	175.2
Trend BY-2020	-55%	-44%	-77%	-49%	-50%	-33%	-71%	81%
Trend 2005-2020	37%	36%	127%	19%	9%	134%	187%	303%

Table 5.2.3 Livestock population and trends for Poultry

Year	Animal Population 1,000 head					
	Laying hens	Broilers	Turkey	Ducks	Geese	Guinea Fowls
BY	24,484.7	50,939.4	1,420.2	2,717.6	1,814.1	362.8
1990	22,735.0	40,178.1	1,772.6	2,463.6	2,926.5	249.8
1991	23,460.1	29,487.6	1,252.7	2,216.7	2,167.5	242.6
1992	20,187.3	27,392.8	916.7	1,969.9	1,459.2	242.6
1993	19,314.4	19,289.5	1,080.1	2,008.4	1,494.1	242.6
1994	17,092.6	21,666.5	1,288.8	2,339.1	1,854.9	235.5
1995	15,732.5	23,349.4	1,599.1	2,144.6	1,833.9	215.0
1996	16,368.0	16,430.5	1,979.1	1,955.3	1,616.4	188.3
1997	15,491.1	18,816.0	2,156.9	2,139.8	1,634.8	178.0
1998	15,824.0	20,158.3	2,156.9	2,725.7	1,623.8	219.0
1999	15,255.0	17,749.4	2,084.3	3,222.1	1,689.9	259.6
2000	13,744.3	24,223.7	4,029.8	3,249.5	3,080.3	234.4
2001	15,396.5	25,290.0	3,449.3	3,790.2	2,915.5	232.5
2002	16,051.5	23,327.7	3,789.8	4,490.0	3,474.3	200.3
2003	16,384.8	23,645.2	3,495.8	4,770.7	3,986.3	203.3
2004	15,398.8	23,187.2	4,637.3	3,898.0	3,177.3	193.3
2005	14,232.3	22,058.3	4,036.5	3,704.0	2,183.2	190.3
2006	14,424.7	20,268.5	4,270.3	3,117.3	2,387.3	185.2
2007	13,063.8	20,359.0	4,430.8	2,780.5	2,374.5	151.0
2008	13,376.3	21,865.8	4,071.2	3,070.0	2,487.8	161.5
2009	12,732.3	22,364.5	3,422.3	3,736.3	2,384.8	149.3
2010	12,544.5	23,163.5	3,365.0	5,155.0	2,211.3	147.8
2011	11,453.4	23,878.3	3,152.8	5,208.1	2,455.5	135.9
2012	11,088.8	22,003.7	3,023.6	4,489.2	2,311.0	147.4
2013	11,839.9	19,959.2	2,432.8	4,533.1	2,774.7	134.6
2014	11,291.9	21,505.5	2,692.7	4,781.3	2,280.7	131.1
2015	11,722.5	22,963.7	2,928.3	4,687.6	2,027.5	129.6
2016	11,246.6	23,307.9	3,022.3	4,854.5	2,354.1	122.3
2017	10,748.7	22,990.4	2,888.4	3,952.7	2,016.1	114.8
2018	10,891.7	22,118.3	2,834.1	4,898.8	2,290.3	103.8
2019	10,732.0	22,176.7	2,825.0	4,768.1	2,270.9	102.6
2020	9,312.4	21,176.9	3,013.8	2,829.6	1,404.9	97.6
Trend BY-2020	-62%	-58%	112%	4%	-23%	-73%
Trend 2005-2020	-35%	-4%	-25%	-24%	-36%	-49%

5.2.2.2 Emission Factors

5.2.2.2.1 Cattle

CH₄ emissions from enteric fermentation in Dairy Cattle and Non-dairy Cattle categories were calculated using the Tier 2 method (2006 IPCC Guidelines, Equation 10.21):

$$EF = (GE * (Y_m/100) * 365) / 55.65 \quad (\text{Equation 5.2})$$

Where:

EF	CH ₄ emission factor [kg CH ₄ head ⁻¹ yr ⁻¹]
GE	gross energy intake [MJ head ⁻¹ day ⁻¹]
Y _m	methane conversion rate [MJ MJ ⁻¹]
365	days of year [day yr ⁻¹]
55.65	energy content of methane [MJ kg ⁻¹ CH ₄]

5.2.2.2.1.1 Gross Energy Intake in Dairy Cattle

Tier 2 emission estimate requires feed intakes expressed in terms of gross energy (MJ/head/day), which is the amount of energy an animal needs for maintenance, activity, lactation, and pregnancy. In the calculation of net energy requirements Holstein-Friesian and Hungarian Simmental cattle were distinguished, which are the most widespread dairy cattle breeds in Hungary.

To calculate the daily net energy requirements of cows the Hungarian Nutrition Codex (2004) was generally applied, which contains standards of animal feeding for Hungary. In Hungary, the American energy requirement system was adapted in 1986 with some minor changes, thus the Hungarian and the IPCC equations for the calculation of net energy requirements are basically very similar. The main difference between the Hungarian and the IPCC methodology is that the Hungarian system does not differentiate the net energy for maintenance and activity, thus both energy requirements are taken into account in the net energy for maintenance. To ensure the closest conformity with the IPCC methodology and to avoid underestimate of emissions it was decided to take also into account the net energy for activity based on the IPCC methodology, using the Eq. 10.4 of 2006 IPCC Guidelines.

Calculation of net energy for lactation according to the Hungarian standards also differs from the IPCC methodology. For this reason, it was determined based on both equations. Use of Hungarian standards indicated higher values than the IPCC methodology. Thus, the net energy for lactation was calculated using the Hungarian standards for the inventory purposes, because it was assumed that it is more reliable for the Hungarian species.

The net energy requirement for pregnancy was also determined based on the standards of the Hungarian Nutrition Codex (2004) as well as the IPCC methodology, and it revealed that there is no difference between the outcomes of the two methodologies. As a consequence, Eq. 10.4 of 2006 IPCC Guidelines was applied for the sake of simplicity.

The Equation 10.16 of 2006 IPCC Guidelines was applied to transform net energy requirements into gross energy intake. The value of digestible energy (DE%) was calculated as weighted average of digestibility of components in the diet to the use of the aforementioned equation. Composition of the diet were taken from the dataset of the Farm Accountancy Data Network (FADN). This dataset provides annual data on the composition of the diet per 1000 kg milk basis, from which the feed intake can be calculated using the annual milk yields. This statistical data was combined with expert judgement on the composition of the diet and the seasonal changes in the feeding practices. Digestibility values for

the different fodder crops in the diet were taken from the 'feed database' provided in the Hungarian Nutrition Codex (2004). This database contains results of laboratory measurements for feeds used for animal nutrition in Hungary.

Parameters and equations used to estimate the gross energy intake for dairy cattle and their sources are listed in Table 5.2.4.

Calculation of net energy requirements requires further statistical data and parameters, which are summarized in Table 5.2.4. Net energy for maintenance depends on the average body mass of dairy cattle, which was determined for each year of the time series based on the change of livestock composition and characteristics of species. In the 70-ies the Hungarian cattle herd consisted mainly of double used cows (Hungarian Simmental) and partly dairy cattle having smaller body mass (Jersey, Ayrshire). Since 1970 this cattle herd has been changed, continuously, crossing the above-mentioned species with Holstein Friesian cattle. In 1985 the Hungarian cattle herd consisted mainly of Holstein-Friesian and Holstein-Friesian Cross-bred, but the Hungarian Simmental also had an importance. Proportions of Jersey and Ayrshire can be considered as negligible. Since 1985 proportion of Hungarian Simmental species has been dropped and as a result the annual milk yield increased from 4671 kg to 6429 kg in the period BY-2005, together with this change the average body weight in the herd also increased. The annual average body mass was calculated from the typical body mass of the two main species and their proportions in the certain year based on HCSO statistics. The typical body mass of Holstein-Friesian and Hungarian Simmental is assumed to be 650 and 550 kg in the calculation. The resulted body weights by years are shown in Table 5.2.7.

Table 5.2.4 Parameters and equations used to estimate the GE for Dairy Cattle

Activity data, parameters and coefficients	Unit	Source	Values/ Notes
Weight	kg	Kovács, 2013	Calculated annually, based on the ratio and the body mass of typical Hungarian breeds.
C_{pregnancy}		Table 10.7 of 2006 IPCC GLs	0.1
Digestible energy intake (DE)	%	Kovács, 2013	Calculated annually, based on feeding statistics from FADN and laboratory measurements (Hungarian Nutrition Codex, 2004).
C_a		Table 10.5 of 2006 IPCC GLs	0 for stall, 0.17 for pasture
Proportion for grazing		HCSO, agricultural surveys, NFCSO's Nitrate database	See also Chapter 6.3.
$NE_m = 2.96 + FM * 4.25 + W * 0.06$ where, FM = farming method (1 = stalled; 2 = farming on good pasture; 3 = farming on average pasture) W = live weight of Cow, kg	MJ/day	Hungarian Nutrition Codex, 2004	Country-specific methodology according to the Hungarian net energy requirements standards. Calculated separately for Holstein-Friesian and Hungarian Simmental
NE_a	MJ/day	Eq. 10.4 of 2006 IPCC GLs	calculated
$NE_i = NE_{l,milk} * \text{kg of milk per day}$ $NE_{l,milk} = 1.45 + 38.45 * \text{Milk fat} + 3.02 * \text{Milk protein}$ where, Milk fat = Fat content of milk, % Milk protein = Protein content of milk, %	MJ/day	Hungarian Nutrition Codex, 2004	Country-specific methodology according to the Hungarian net energy requirements standards.
Ne_p	MJ/day	Eq. 10.13 of 2006 IPCC GLs	calculated
REM		Eq.10.14 2006 IPCC GLs	calculated
GE	MJ/day	Eq. 10.16 of 2006 IPCC GLs	calculated

Net energy for lactation depends on the amount of daily milk production and fat content of milk. The daily average milk yield was calculated based on the HCSO's annual milk yield statistics. In 2020 the daily average milk production was 23.1 kg of milk per cow (*Table 5.2.7*). Data on fat content of milk was taken from the Eurostat statistics for the period 1998-2011, while for the period 1985-1997 the average of the values calculated for the period 1998-2003 were assumed due to lack of statistical data. Since 2004 this data has been taken from the Market Price Information System of the Institute of Agricultural Economics Nonprofit Kft. (hereafter AKI). (The legal background of the price transmission is laid down in the Commission Implementing Regulation (EU) 2019/1746 and in the national regulation Degree of Ministry of Rural Development 127/2013. According to the national regulation notification is mandatory for dairy processors exceeding 10,000 tons of raw milk delivery in the previous year.)

Background data to estimate emissions from cattle are shown in *Table 5.2.5*.

Table 5.2.5 Background data to estimate emissions from dairy cattle

Year	Body Mass	Milk Yield (MY)	Digestibility (DE)	Neutral Detergent Fiber (NDF)	Forage in the diet
	kg	kg head ⁻¹ yr ⁻¹	%	%	%
BY	628	4,671	68.5	41.8%	78%
1990	633	5,031	69.3	40.7%	72%
1991	636	4,711	69.3	41.0%	72%
1992	639	4,780	69.4	41.3%	71%
1993	641	4,757	69.4	41.4%	71%
1994	641	4,716	69.4	41.5%	71%
1995	641	4,991	69.9	40.2%	68%
1996	640	5,064	69.9	39.8%	68%
1997	640	5,112	70.0	39.6%	68%
1998	641	5,513	70.3	39.0%	66%
1999	639	5,454	70.2	38.6%	66%
2000	641	5,886	70.5	38.4%	64%
2001	641	6,051	70.6	37.9%	64%
2002	641	6,155	70.6	37.6%	64%
2003	642	6,154	70.6	37.6%	64%
2004	642	6,131	70.4	37.4%	64%
2005	642	6,429	70.5	37.1%	63%
2006	642	6,706	70.6	36.8%	62%
2007	643	6,874	70.7	36.7%	63%
2008	643	6,972	70.5	36.6%	65%
2009	642	6,815	70.4	36.6%	65%
2010	642	6,877	70.3	36.3%	66%
2011	640	6,835	70.0	35.8%	68%
2012	639	7,104	70.1	35.4%	67%
2013	641	7,135	70.2	35.6%	66%
2014	641	7,443	70.4	35.4%	64%
2015	642	7,702	70.0	35.4%	69%
2016	643	7,768	69.9	35.5%	71%
2017	643	8,038	70.1	35.1%	70%
2018	643	8,031	69.6	36.9%	75%
2019	643	8,048	70.1	35.2%	73%
2020	643	8,447	70.3	36.5%	73%

5.2.2.2.1.2 Methane Conversion Rate for Dairy Cattle

Following a recommendation from the annual review conducted in 2013 country-specific value of Y_m for Dairy Cattle were developed based on the data on composition of diet used for the estimation of GE. Laboratory measurements on Y_m , similarly to most of the other country, is unavailable in Hungary. Therefore, until the 2019 submission country-specific values were calculated based on conclusions of the related publication of Soliva (2006). As a result of a question raised during the UNFCCC Review, 2019, for this submission the Y_m values were revised according to the latest scientific publications.

According to the most recently published scientific publications Y_m values depends on the feed type, quality and the animal characteristics as breed and genetics. In Hungary the most typical dairy cattle breed is Holstein-Friesian (87.5%) while the remaining Jersey, and Ayrshire, which breeds were mainly examined in the studies of Niu et al. (2018) and Hellwing et al. (2016). In the study of Niu et. al 68% of the data examined related to Holstein cows. In the publication of Hellwing et al. (2016) Holstein and Jersey cows were studied.

In line with these publications and the 2019 Refinement to the 2006 IPCC Guidelines (IPCC, 2019) annual value of the Y_m were determined based on the country-specific values of milk yield (MY), digestibility (DE) and Neutral Detergent Fiber (NDF) in percentage of the dry matter intake (DMI).

Hungarian cattle are considered as low producing cows ($MY < 5000 \text{ kg yr}^{-1}$) at the beginning of the time series and medium producing cows ($5000 \text{ kg yr}^{-1} < MY < 8500 \text{ kg yr}^{-1}$) since 1996. Proportions of concentrate in the diet have changed in a range between 21% and 36%; while digestibility has changed within a narrow range, between 68.3% and 70.7% over the time series, due to the good-quality forage crops. (Hungarian cattle's diet contains mainly maize silage as forage.)

According to Niu et. al (2018) information on neutral detergent fiber (NDF) improve the prediction of the enteric methane (CH_4) production from cattle. Therefore, to get more accurate estimate of Y_m the values of NDF were determined based on the diet for the whole time series. Percentage of NDF in DMI has decreased from 42% to 35% across the time series.

The annual values of Y_m were determined as it is shown in Table 5.2.6 .

Table 5.2.6 Assumptions made to estimate Methane Conversion Rates (Y_m) for Dairy Cattle

Milk Yield (MY) [kg head ⁻¹ yr ⁻¹]	Digestibility (DE) [%]	Neutral Detergent Fiber (NDF) [%]	Enteric Conversion Factor (Y_m) [%]	Methane
<5000	68%<DE<70%	40%<NDF	6.5	
5000<MY<8500	68%<DE<70%	37%<NDF	6.3	
5000<MY<8500	DE<70%	35%<NDF≤37%	6.2	
5000<MY<8500	DE≤70%	NDF≤35%	6.1	

Values of the derived parameters used to calculate the emissions from the enteric fermentation and manure management of dairy cows, including the methane conversion factor for 3.A.1, are shown in Table 5.2.7.

*Table 5.2.7 Gross energy Intakes, Methane Conversion, and Nitrogen Excretion Rates for Dairy Cattle
BY-2020*

Year	Gross Energy Intake	Enteric methane conversion factor Y_m	Nitrogen excretion N_{ex}
	MJ/head/day	%	kgN/head/yr
BY	254	6.50	76.4
1990	255	6.30	83.0
1991	246	6.50	81.1
1992	246	6.50	81.8
1993	244	6.50	81.7
1994	243	6.50	81.6
1995	247	6.50	88.3
1996	249	6.30	89.2
1997	250	6.30	90.1
1998	257	6.20	94.2
1999	257	6.20	94.0
2000	264	6.20	97.3
2001	267	6.20	98.7
2002	270	6.20	99.9
2003	271	6.20	100.4
2004	269	6.20	103.2
2005	275	6.20	106.2
2006	280	6.10	109.1
2007	285	6.10	110.8
2008	288	6.10	112.0
2009	286	6.10	109.9
2010	289	6.10	110.1
2011	290	6.10	109.2
2012	297	6.10	111.7
2013	298	6.10	112.1
2014	302	6.10	115.3
2015	308	6.10	118.9
2016	313	6.20	119.9
2017	316	6.10	123.3
2018	319	6.20	118.6
2019	315	6.10	124.5
2020	322	6.10	131.5

5.2.2.2.2 Gross Energy Intake for Non-Dairy Cattle

Gross energy intakes for non-dairy cattle were derived from the study of Kovács, 2013, where the typical Hungarian diets for each sub-category of non-dairy cattle was determined. Besides, the seasonal changes in the diets were also taken into account for each sub-category. In the calculation the available data, the Hungarian dietary standards and expert judgments were combined to get the most reliable results. Similarly, to the dairy cattle values of net energy requirements and net energy for lactation for other cattle were calculated according to the Hungarian standards. Table 5.2.8 summarizes the parameters and equations used to estimate the gross energy intake for non-dairy cattle.

Table 5.2.8 Parameters and equations to estimate gross energy intakes for Non-dairy cattle

Activity data, parameters and coefficients	Unit	Sources	Values/ Notes
Weight	kg	Kovács, 2013	Calculated based on the livestock composition.
Weight Loss			NO
WG (daily weight gain)	kg	Kovács, 2013	1 for male<1 year, 0.73 for female<1 year, 0.65 for heifers, 0.9 for bovines 1-2 years, 0 for mature
C, Coefficient for Eq. 10.6 of GI (IPCC, 2006)		2006 IPCC GLs	0.8 for females, 1.2 for bulls, 0 for mature
C _{pregnancy}		Table 10.7 of 2006 IPCC GLs	0.1
Digestible energy intake (DE%)	%	Kovács, 2013	Calculated based on fed diets and laboratory measurements
C _a		Table 10.5 of 2006 IPCC GLs	0 for stall, 0.17 for pasture
proportion for grazing		HCSO, agricultural surveys, NFCSO's Nitrate database	
NE _m	MJ/day	Hungarian Nutrition Codex, 2004	Country-specific methodology according to the Hungarian standards of net energy requirements
NE _a	MJ/day	Eq. 10.4 of 2006 IPCC GLs	calculated
NE _i	MJ/day	Hungarian Nutrition Codex, 2004	Country-specific methodology according to the Hungarian standards of net energy requirements
Ne _g	MJ/day	Eq. 10.6 of 2006 IPCC GLs	calculated
Ne _p	MJ/day	Eq. 10.13 of 2006 IPCC GLs	calculated
REM		Eq. 10.14 of 2006 IPCC GLs	calculated
REG		Eq. 10.15 of 2006 IPCC GLs	calculated
GE	MJ/day	Eq. 10.16 of 2006 IPCC GLs	calculated

Net energy for maintenance depends on the live weight, which was determined based on the study of Kovács, 2013. The typical body mass for each sub-category as well as the resulted gross energy intake and the emission factors for the BY and the year 2020 are shown in Table 5.2.9 and Table 5.2.10.

Table 5.2.9 Gross energy intakes and emission factors by non-dairy cattle subcategories for the base year (BY)

BY	<1 year		1-2 year		>2 year			
	Bovines for slaughter and other calves (male)	Bovines for slaughter and other calves (female)	Bovines (male)	Heifers for slaughter and other heifers	First calf heifers	Mature Non-Dairy (male)	Heifers for slaughter	Beef Cow

Live weight	kg	195	170	415	370	515	575	530	600
Digestible Energy	%	69	71	62	62	69	66	67	69
N-excretion	kg N / head/ year	42	41	40	31	61	56	53	70
Gross Energy Intake	MJ/ head/ day	94	92	156	160	200	192	185	157
Concentrate ratio	%	31%	33%	10%	12%	18%	17%	17%	16%
Y_m	%	5.51	5.47	7.03	7.00	6.87	6.90	6.89	6.90
Emission Factor for 3.A	kg CH ₄ / head/ year	24	23	72	73	90	87	84	71

Table 5.2.10 Gross energy intakes and emission factors by non-dairy cattle subcategories for the year 2020

		<1 year		1-2 year			>2 year		
2020		Bovines for slaughter and other calves (male)	Bovines for slaughter and other calves (female)	Bovines (male)	Heifers for slaughter and other heifers	First calf heifers	Mature Non-Dairy (male)	Heifers for slaughter	Beef Cow
Live weight	kg	195	170	415	370	515	575	530	600
Digestible Energy	%	69	70	61	62	68	65	65	67
N-excretion	kg N/ head/ year	44	42	46	41	66	60	57	75
Gross Energy Intake	MJ/ head/ day	94	94	161	163	192	199	191	162
Concentrate ratio	%	30%	33%	10%	12%	18%	17%	17%	17%
Y_m	%	5.53	5.48	7.03	7.01	6.87	6.90	6.89	6.90
Emission Factor for 3.A	kg CH ₄ / head/ year	24	24	74	75	87	90	86	73

Methane conversion rate for non-dairy cattle

Methane conversion rate for non-dairy cattle was calculated by linear interpolation in dependency of the proportion of concentrate in the dry matter intakes. In case of 'Bovines < 1 year' for the period of

consuming only milk methane conversion rate zero was assumed in accordance with the 2006 IPCC Guidelines. The time of consuming only milk for juveniles was assumed to be 60 days, which is in line with the Hungarian standards.

5.2.2.2.3 Other livestock categories

Detailed information required to develop the Tier 2 emission factor is not available for other important livestock category in Hungary, such as sheep. Therefore, the Tier 1 methodology for enteric fermentation for all livestock categories other than cattle is applied. The emission factors used are the IPCC default ones provided for developed countries in the Table 10.10 of the 2006 IPCC Guidelines. In the case of Rabbit and Poultry the IPCC methodologies do not provide emission factors. Emissions from enteric fermentation in rabbits are relatively small, accounting for 0.2 percent of the total emissions from enteric fermentations in all livestock, so development of a country-specific emission factor does not seem to be reasonable. The emission factor provided by the Italian NIR, 2008 is used, because Italy is the nearest neighbor of Hungary, who reports emissions from rabbits. It is assumed that the Hungarian housing and feeding practices do not differ from the Italian ones. Emission factor for poultry was taken from the literature, due to lack of IPCC default values. Sources of emission factors per livestock species are summarized in *Table 5.2.11*.

Table 5.2.11 Emission factors used for the calculation of the methane emissions from enteric fermentation

Animal category	CH ₄ -emission factor [kg head ⁻¹ yr ⁻¹]	Comments
Buffalo	55	IPCC default value for developed countries
Sheep	8	IPCC default value for developed countries
Goats	5	IPCC default value for developed countries
Horses	18	IPCC default value for developed countries
Asses & Mules	10	IPCC default value for developed countries
Swine	1.5	IPCC default value for developed countries
Poultry	0.015	expert judgement, according to Minonzio et al. (1998)
Rabbits	0.08	expert judgement, according to the NIR of Italy, 2008

5.2.3 Uncertainties and time series consistency

Uncertainty of activity data (animal population) was estimated based on the confidence intervals for each animal category and livestock survey provided by the HCSO. The uncertainty of the mean annual averages was estimated according to the error propagation rules. (See Table 5.1.3) For the uncertainty of the country specific EFs $\pm 20\%$ were assumed, while for the default EFs $\pm 40\%$ were applied in accordance with the 2006 IPCC Guidelines. The combined uncertainty in emissions from the 3.A sector is ± 13 per cent. CH₄ EF for dairy cattle was estimated based on data for milk production and GE, for which uncertainties could be estimated to be less than $\pm 3\%$ and 10%, respectively. Data on milk

production is readily available while the GE is checked against cattle feeding requirements arising from the biology of ruminants (e.g., ratio of crude protein, dry matter intake and proportion of silage in the diet).

5.2.4 Source specific QA/QC information

Consistency of Animal Populations

Since the centralized review conducted in 2014 the HCSO has provide animal populations rounded to the nearest hundred instead of nearest thousand (according to the recommendation of the ARR, 2014 para 47). Until 2014 the HCSO provided animal populations rounded to the nearest thousand; data for each livestock subcategory as well as the total livestock population were rounded by the HCSO.

It was not feasible to provide new animal populations rounded to the nearest hundreds for the full time series. The HCSO provided revised animal numbers backward to 2011.

In the case of animal categories for which enhanced livestock characterization is used rounding can cause slight differences between the rounded totals and the sum of the rounded values of subcategories.

As the IPCC methodology requires annual average animal populations, this fact can cause further seeming discrepancies in the case of enhanced livestock characterization, because of the error propagation.

To avoid inconsistencies in NIR tables arising from rounding and the use of annual average animal populations the following correction is used:

- The HCSO has provided animal numbers rounded to the nearest hundred since the inventory year 2011. However, using rounded values to the nearest hundred instead of nearest thousand the error propagation cannot be completely avoided; only reduced.
- An adjustment of animal numbers of Non-dairy cattle and Swine subcategories was applied to the chronological means of totals to eliminate the differences between the chronological means of totals and the sum of the chronological means by subcategories for those years when a slight difference occurred. It is worth noting, that the aforementioned discrepancy also exists in the case of the inventory year 2011, because the chronological means contain the livestock population in December of the previous year (i.e., 2010).

In the next figures we present an example of processing and adjustment of HCSO's livestock populations to get the required activity data to the emission estimate. The method used to derive

Dairy cattle and Non-dairy cattle average annual populations for the year 2011 are outlined in the following steps below:

- STEP1 HCSO provides annual population survey data by subcategories (*Figure 5.2.3*).

- STEP2 Chronological means are calculated for each subcategory, as well as the total (*Figure 5.2.4*). See also Section 5.2.2.1.
- STEP3 Adjustment to the HCSO's total Cattle livestock is applied (*Figure 5.2.5*).

<i>STEP 1 Data provided by the HCSO (1'000 head)</i>			
	12/1/2010	6/1/2011	12/1/2011
Bovines less than one year old			
Calves for slaughter, male	40.0	46.7	47.8
Calves for slaughter, female	12.0	10.9	12.0
Other calves, male	26.0	30.6	29.5
Other calves, female	92.0	93.0	98.5
Bovines aged between one and two			
Male	30.0	25.6	24.5
Female for slaughter (heifers)	18.0	7.2	4.9
Other heifers	105.0	106.1	108.1
Bovines of two years and over			
Male	8.0	6.7	6.4
Female for slaughter (heifers)	3.0	2.6	2.2
Other heifers	37.0	35.2	34.8
Cows, dairy	193.0	206.0	196.9
Cows, beef	70.0	71.9	77.2
Cows, dual purpose	47.0	49.3	54.7
Cattle, total	682.0	691.7	697.4
<i>Cattle, SUM OF SUBCATEGORIES</i>	<i>681.0</i>	<i>691.8</i>	<i>697.5</i>
<i>Difference</i>	<i>-1.00</i>	<i>0.10</i>	<i>0.10</i>
Dairy Cattle	240.0	255.3	251.6
Other Cattle	442.0	436.4	445.8
Cattle, total	682.0	691.7	697.4
<i>Cattle, SUM OF SUBCATEGORIES</i>	<i>681.0</i>	<i>691.8</i>	<i>697.5</i>
<i>Difference</i>	<i>-1.00</i>	<i>0.10</i>	<i>0.10</i>

Figure 5.2.3 Cattle populations survey data provided by the HCSO for the inventory year 2011

	2011			
Bovines less than one year old				
Calves for slaughter, male	45.3			
Calves for slaughter, female	11.5			
Other calves, male	29.2			
Other calves, female	94.1			
Bovines aged between one and two				
Male	26.4			
Female for slaughter (heifers)	9.3			
Other heifers	106.3			
Bovines of two years and over				
Male	7.0			
Female for slaughter (heifers)	2.6			
Other heifers	35.6			
Cows, dairy	200.48			
Cows, beef	72.8			
Cows, dual purpose	50.08			
Cattle, total	690.7			
SUM OF CHRONOLOGICAL MEANS	690.5			
Difference for Cattle	-0.17			
Dairy Cattle	250.6	reported in the CRF Table		
Other Cattle	440.2	reported in the CRF Table		
Cattle, total	690.7	consistent with the HCSO's total		
Other Cattle SUM OF CHRONOLOGICAL MEANS	440.0			
Difference for Non-dairy Cattle	-0.18			

Figure 5.2.4 Chronological means for Cattle, 2011

STEP 3 Adjustment and conversion to activity data	2011		
	Original	Adjusted	Difference
<1 year			
Calves, male	74.5	74.5	0.03
Calves, female	105.6	105.6	0.04
1-2 year			
Bovines (male)	26.4	26.4	0.01
Heifers for slaughter and other heifers	115.7	115.7	0.05
>2 year			
Mature Non-Dairy (male)	7.0	7.0	0.00
Mature Non-Dairy (female)	2.6	2.6	0.00
First calf heifers	35.6	35.6	0.01
Beef Cow	72.8	72.8	0.03
Non-dairy Cattle	440.2	440.2	0.00
Non-dairy Cattle, SUM OF SUBCATEGORIES	440.0	440.2	0.17
Dairy Cattle	250.6	250.6	0.00
Cattle, total	690.7	690.7	0.00

Figure 5.2.5 Adjustment of average annual populations for Non-dairy cattle, 2011

The country specific value of the gross energy intake for Dairy Cattle was verified using values reported by the EU member states. Verification revealed the Hungarian value of 315 MJ head⁻¹ d⁻¹ for the year

2019 was consistent with the reported values by other EU member states. The average gross energy intake for the EU-28 member states was slightly lower, 312 MJ head⁻¹ d⁻¹ according to the EU's NIR 2021 submission. The milk yield in Hungary was slightly higher than in the EU-28. The milk production for the year 2019 was 20.2 kg for the EU-28, while 22.0 kg for Hungary. The feed digestibility was 71.3% in the EU-28, while 70.1% in Hungary.

5.2.5 Source-specific recalculations

In sector 3.A, recalculations were carried out to ensure consistency of the time series in terms of nutritional values of raw milk and parameters calculated from them, and to correct a previous data entry error.

The changes by source are as follows:

3.A.1 Enteric Fermentation - Dairy Cattle, 2004-2019

The HCSO has changed the methodology of data collection on milk fat and milk protein since 2019 and 2020, respectively. This change led to time series inconsistency, as the HCSO has not revised the data, retrospectively in line with the new methodology. To ensure the time series consistency the milk fat and protein content data were revised based on the Institute of Agricultural Economics' Market Price Information System (MPIS) data. The revised nutritional values of raw milk used to derive the gross energy intake (GE) for dairy cattle, resulted in an increase of 0.15% and 1.1 kt CO₂-eq on average in the CH₄ emissions from 3.A.1 Enteric Fermentation - Dairy Cattle for the 2004-2019 trend. The change ranged between -5.2 kt CO₂ eq in 2004 and 7.7 kt CO₂ eq in 2016.

Impact of recalculations in 3.A.1 Enteric Fermentation - Dairy Cattle is shown in *Table 5.2.12*.

Table 5.2.12 Impact of recalculations in 3.A.1 Enteric Fermentation - Dairy Cattle

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
2004	852	846	-5.2	-0.6%
2005	836	838	1.8	0.2%
2006	774	770	-3.8	-0.5%
2007	764	763	-1.6	-0.2%
2008	762	760	-1.7	-0.2%
2009	739	737	-2.1	-0.3%
2010	704	708	4.0	0.6%
2011	723	726	3.2	0.4%
2012	755	759	4.0	0.5%
2013	734	739	5.4	0.7%
2014	756	761	5.1	0.7%
2015	773	777	4.1	0.5%
2016	777	785	7.7	1.0%
2017	769	774	4.8	0.6%
2018	782	787	5.2	0.7%
2019	765	769	3.7	0.5%

3.A.1 Enteric Fermentation - Non-Dairy Cattle, heifers 1-2 yr, 2016-2019

A data linking error was corrected, because in the previous submissions the livestock number of heifers, 1-2 years for the year 2015 was used for the period 2016-2019 as well. Due to this recalculation emissions increased by 0.4%-2.2% for the period 2016-2019.

Impact of recalculations in 3.A.1 Enteric Fermentation - Non-Dairy Cattle is shown in *Table 5.3.13*

Table 5.2.13 Impact of recalculations in 3.A.1 Enteric Fermentation - Non-Dairy Cattle

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
2016	818	822	4.4	0.5%
2017	852	855	3.1	0.4%
2018	871	880	8.9	1.0%
2019	898	918	20.1	2.2%

The overall effect of recalculations on 3.A Enteric Fermentation was a slight decrease in the emissions between 2004 and 2009 (except 2005) and increase in 2005 and over the period 2010-2019. The increase is the most significant (23.8 kt CO₂-eq) in 2019 (*Table 5.2.14*).

Table 5.2.14 Impact of recalculations in 3.A Enteric Fermentation

Year	Submission 2021 [Gg CO ₂ - eq]	Submission 2022 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
2004	1,907	1,902	-5.2	-0.3%
2005	1,882	1,884	1.8	0.1%
2006	1,817	1,813	-3.8	-0.2%
2007	1,821	1,819	-1.6	-0.1%
2008	1,792	1,791	-1.7	-0.1%
2009	1,767	1,764	-2.1	-0.1%
2010	1,752	1,756	4.0	0.2%
2011	1,745	1,748	3.2	0.2%
2012	1,815	1,819	4.0	0.2%
2013	1,864	1,870	5.4	0.3%
2014	1,923	1,928	5.1	0.3%
2015	1,974	1,978	4.1	0.2%
2016	2,010	2,022	12.1	0.6%
2017	2,022	2,030	7.9	0.4%
2018	2,051	2,065	14.0	0.7%
2019	2,048	2,072	23.8	1.2%

5.2.6 Planned improvements

See Section 5.1.9

5.3 Manure management (CRF sector 3.B)

Emitted gases: CH₄, N₂O

Methods: T1, T2

Emission factors: D, CS

Key source: Yes

Particularly significant sub-categories, CH₄: Swine and Cattle

Particularly significant sub-categories, N₂O: Other AWMS and Indirect emissions

Animal manure is an important source of CH₄ and N₂O. The amount of CH₄ and N₂O emitted from the manure to the atmosphere depends on the conditions of manure management and use as well as on the composition of released excrements. CRF category 3.B comprises direct and indirect emissions during storage and treatment of manure before it is applied to land.

5.3.1 Source Category Description

In 2020 23% of agricultural CH₄ and 10% of agricultural N₂O emissions arose from the 3.B Manure management. The bulk of emissions were generated in cattle and swine husbandry (in 2020 they accounted for 536 and 295 Gg CO₂-eq, which equates to 50% and 27% of total GHG emissions from 3.B, respectively), due to the considerable share of deep bedding and liquid manure. The main sources of CH₄ emissions from 3.B are Swine and Cattle manure (Figure 5.3.1), and most of N₂O emissions are generated in the solid and 'other' systems containing 'Cattle and Swine deep bedding' and 'Poultry manure with or without litter'. Indirect emissions contributed 12% to the N₂O emissions from this sector. The uncovered manure tanks and the cattle housing are the main sources of the significant amount of N₂O emissions from volatilization of N in form of NH₃ and NO_x.




Figure 5.3.1 Emissions from 3.B Manure management by sources BY-2020

Emissions from 3.B Manure management have decreased by 50% since the BY (Table 5.3.1). Considering CH₄ and N₂O emissions separately, they have decreased by 49% and 53% over the inventory period, respectively. The significant decrease in the emissions reflects the decreasing swine and cattle livestock between 1985 and 1994. In the period 1995-2003 emissions fluctuated to some extent on yearly basis, following the annual changes in swine population. Emissions have decreased again in the period 2004 to 2010 reflecting again the falling swine livestock numbers over that period. Since 2011 emissions from 3.B slightly increased due to the increasing non-dairy cattle livestock. CH₄ and N₂O emissions from 3.B are shown in Table 5.3.1 and

Table 5.3.2.

Indirect N₂O emissions due to leaching decreased continuously over the time series. The decrease in the emission levels reflects the drop in livestock population and the effect of measures to reduce nitrate leaching during manure storage.

Table 5.3.1 Trend in CH₄ emissions from 3.B Manure Management by livestock categories

Year	CH ₄ emissions from 3.B				
	Dairy-Cattle	Non-Dairy Cattle	Sheep	Swine	Other livestock
BY	14.85	10.79	0.74	20.47	3.51
1990	14.00	9.16	0.58	20.06	3.22
1991	12.60	8.79	0.60	18.08	3.04
1992	11.40	7.11	0.55	14.52	2.65
1993	10.28	5.52	0.43	13.58	2.45
1994	9.57	4.70	0.32	11.78	2.36
1995	9.26	4.66	0.30	11.88	2.26
1996	9.19	4.69	0.28	13.59	2.18
1997	9.17	4.50	0.27	12.96	2.17
1998	9.19	4.17	0.28	14.14	2.24
1999	9.30	4.14	0.29	15.74	2.17
2000	8.89	4.00	0.35	15.01	2.38
2001	8.75	3.67	0.34	14.77	2.47
2002	8.64	3.59	0.34	16.18	2.56
2003	8.28	3.59	0.36	16.99	2.58
2004	7.82	3.57	0.41	14.89	2.47
2005	7.79	3.51	0.43	13.88	2.17
2006	7.31	3.61	0.40	13.77	2.08
2007	7.27	3.75	0.38	14.25	1.90
2008	7.20	3.72	0.37	12.95	1.82
2009	6.92	3.85	0.37	11.48	1.68
2010	6.59	4.03	0.35	11.38	1.67
2011	6.69	3.93	0.34	11.11	1.60
2012	7.04	4.34	0.35	10.73	1.50
2013	6.58	4.81	0.36	10.46	1.38
2014	6.78	5.20	0.37	10.78	1.40
2015	7.05	5.64	0.36	10.89	1.43
2016	7.06	6.12	0.36	10.38	1.41
2017	7.04	6.26	0.35	9.79	1.34
2018	7.24	6.38	0.34	10.01	1.36
2019	7.19	6.57	0.32	9.84	1.35

2020	7.27	6.80	0.29	9.98	1.22
Share in 3B CH4 in BY	29.5%	21.4%	1.5%	40.6%	7.0%
Share in 3B CH4 in 2020	28.4%	26.6%	1.1%	39.1%	4.8%
Trend BY-2020	-51%	-37%	-61%	-51%	-65%

Table 5.3.2 Trend in N₂O emissions from 3.B Manure Management by sources

Year	N ₂ O emissions from 3.B						
	Direct					Indirect	
	Dairy-Cattle	Non-Dairy Cattle	Sheep	Swine	Other livestock	Atmospheric deposition	Nitrogen leaching and run-off
BY	0.48	0.51	0.14	0.57	0.99	0.99	0.04
1990	0.49	0.44	0.11	0.53	0.90	0.90	0.04
1991	0.45	0.43	0.11	0.48	0.81	0.81	0.04
1992	0.41	0.35	0.10	0.39	0.69	0.69	0.03
1993	0.38	0.27	0.08	0.36	0.60	0.60	0.03
1994	0.35	0.23	0.06	0.31	0.55	0.55	0.02
1995	0.37	0.23	0.06	0.31	0.56	0.56	0.02
1996	0.37	0.24	0.05	0.34	0.55	0.55	0.02
1997	0.37	0.23	0.05	0.31	0.53	0.53	0.02
1998	0.38	0.22	0.05	0.33	0.55	0.55	0.02
1999	0.38	0.21	0.06	0.35	0.56	0.56	0.02
2000	0.37	0.21	0.07	0.33	0.58	0.58	0.02
2001	0.37	0.19	0.07	0.30	0.57	0.57	0.02
2002	0.36	0.19	0.07	0.32	0.57	0.57	0.02
2003	0.34	0.19	0.08	0.32	0.58	0.58	0.02
2004	0.33	0.19	0.09	0.28	0.55	0.55	0.02
2005	0.33	0.19	0.09	0.26	0.52	0.52	0.02
2006	0.31	0.19	0.09	0.25	0.50	0.50	0.01
2007	0.31	0.20	0.09	0.26	0.50	0.50	0.01
2008	0.30	0.20	0.09	0.23	0.49	0.49	0.01
2009	0.29	0.20	0.09	0.21	0.46	0.46	0.01
2010	0.27	0.21	0.09	0.20	0.47	0.47	0.01
2011	0.27	0.21	0.09	0.20	0.46	0.46	0.01
2012	0.28	0.23	0.09	0.19	0.45	0.45	0.01
2013	0.27	0.25	0.09	0.18	0.44	0.44	0.01
2014	0.28	0.26	0.10	0.18	0.45	0.45	0.01
2015	0.28	0.28	0.09	0.18	0.46	0.46	0.01
2016	0.28	0.30	0.09	0.18	0.46	0.46	0.00*
2017	0.28	0.31	0.09	0.16	0.45	0.45	0.00*
2018	0.26	0.32	0.08	0.16	0.45	0.45	0.00*
2019	0.28	0.33	0.08	0.15	0.45	0.45	0.00*
2020	0.28	0.34	0.07	0.15	0.44	0.44	0.00*

Share in 3B N ₂ O in BY	15.3%	16.5%	4.5%	18.5%	31.9%	31.9%	1.3%
Share in 3B N ₂ O in 2020	19.2%	23.0%	4.8%	10.5%	30.2%	30.2%	0.3%
Trend BY- 2020	-41%	-35%	-50%	-73%	-56%	-56%	-91%

*We note that the '0.00' values are due to the rounding of small numbers rather than inferring those emissions are zeros.

5.3.2 Methodological issues

5.3.2.1 Calculation method

CH₄ emissions from manure management were estimated using Tier 2 methodology, except Rabbit, which contribution is less than 1% to the source category. Direct N₂O emissions were calculated using Tier 2 method for the important livestock categories in Hungary, such as Dairy cattle, Non-dairy cattle, Swine and Poultry. For these livestock categories country-specific nitrogen excretion rates, but IPCC default values of emission factors were applied. For the other livestock categories Tier 1 method was adopted. Indirect N₂O emissions were estimated based on the national air pollutant emissions inventory (i.e., reported NH₃ and NO_x emissions), which meet the requirement of the IPCC Tier 2 method. A detailed description of the methods applied for the calculation of NH₃ and NO_x emissions is given in the report 'Hungary's Informative Report, 2022' – Submission under the UNECE/CLRTAP.

5.3.2.2 Activity Data

5.3.2.2.1 Animal Waste Management System Distribution

Activity data on allocation of manure to animal waste management systems is based on processing and synthesizing of statistics from the HCSO's General Agricultural Censuses conducted in 2000, 2010 and 2020, Farm Structure Surveys, conducted in 2003, 2005, 2007, 2013, 2016, annual data for the period 2004-2020 from the Nitrate Database, statistics on agricultural wastes treated in biogas plants provided by the Hungarian Energy and Public Utility Regulatory Authority. Expert judgments were drawn on to the further stratification of primarily data e.g., to stratify deep litter by mixing and frequency of removals.

In Hungary the first comprehensive study on animal waste management system distribution for emission inventory purposes was carried out by Ráky in 2003 based on the HCSO's General Agricultural Census 2000. This study focused on product producer farms and provides data by farm-size structure. The results of the HCSO's General Agricultural Census 2010 and 2020 provided comprehensive information on the manure management system distribution again. The census provide data on housing practices for cattle, swine and laying hens, and in addition on grazing for all animal species. The surveyed housing systems are as follows:

Cattle

- Tied and loose housing solid systems
- Tied and louse housing liquid systems
- Other solid and liquid systems

Swine

- Partly slatted floors (liquid systems)

- Fully slatted floors (liquid systems)

- Solid systems

- Deep litter

- Other

Poultry

- Deep litter

- Cage with manure belt

- Cage with pit

- Battery cage with stilt house

- Other battery cage

- Aviary (without litter)

- Other

Farm Structure Survey data was applied to get representative activity data from the different datasets published by farm size structure, and it was applied as surrogate data to the interpolation of the 2000-2010 and 2010-2020 time series. Farm structure survey conducted in 2013 and 2016 contained a more detailed data collection on grazing than former surveys. This data on proportion of grazing animals as well as grazing period was also taken into account in the inventory preparation.

Agricultural census is taken every 10 years, thus for the recent years statistics from the Nitrate Database provides the most reliable data on animal waste management system distribution. Annual statistics from the Nitrate Database are supplied by the National Food Chain Safety Office (NFCO) to the inventory. Data collection for the Nitrate Database is based on the Decree of the Ministry of Agriculture and Rural Development No. 59/2008 (IV. 29). The Annex 6 of the Decree contains a questionnaire, which is the basis of the data collection. Data supply obligation is prescribed for farmers, whose animal production exceeds the household requirements. The first version of this Decree (Government Decree No. 49/2001 (IV. 3)) entered into force in 2001. The collected data have been stored in a database since 2003. This database contains data on cattle and swine by sub-categories, poultry (laying hens, cocks and broilers, ducks, geese, turkey), sheep and goats, horse. Six different management systems were distinguished: liquid, solid, deep litter, grazing, farmyard/paddock and other. Amendments of this decree in 2008 resulted in a minor change in the structure of the data collection. Until 2007 only the livestock numbers for six housing systems were collected, while since 2008 the amount of the manure has also been surveyed. In 2009 a more detailed livestock characterization was introduced for cattle and swine. At the same time sheep and goats were separated into two different categories. The former paper questionnaires were replaced by on-line forms in 2014. This measure contributed to the improvement of compliance with data provision obligations. In 2013, Hungary revised the area of the so-called 'Nitrate Vulnerable Zones' (hereafter NVZs). Thus, the areas designated as NVZs increased to approximately 68-69% of the country from the former 47%, further increasing the number of farms under the data provision obligations. In 2016 the

data provision obligations of farmers were amended. The new regulations were developed in line with the data needs of emission inventories. The former six categories of management systems were improved by more detailed categories.

The number of the received questionnaire has been increasing since 2003, although the representativeness of this sample varies between different years and livestock categories. The dataset is most representative for cattle and poultry, about 80-90 per cent of these livestock are covered. This data collection covers about 70-75% of the pig livestock. It is least representative for goats and horse with 10-15 per cent coverage.

The applied data sources sometimes contain information on housing practices rather than manure management storage systems in many cases, therefore additional qualitative information was needed to define the relationship between the housing and manure management systems. Two studies (Mészáros, 2005 and Pazsiczky et. al, 2006) were applied to get additional information. Data on sheep and goats were updated by a survey for sheep and goats in 2011 (Borka et al., 2010).

The 2006 IPCC Guidelines require data on liquid manure to be disaggregated by manure storage coverage. The Nitrate Database has provided data on covered manure tanks, including formation of natural crust from the year 2015 onwards. This database was used to determine the proportion of covered and uncovered manure stores for the period 2015-2020. In Hungary, regulations on the construction of manure storage facilities have been in place since 2001, and covered slurry storage facilities have been built since then. Therefore, before 2001, only natural crust cover was considered, other types of cover were considered as 0. Between 2001 and 2015, linear interpolation was used. Table 5.3.3 shows the proportions of the covered cattle and swine slurry tanks in Hungary.

Table 5.3.3 Proportions of covered cattle and swine slurry tanks

	1990	1995	2000	2005	2010	2015	2020
Cattle	52%	52%	52%	54%	57%	59%	54%
Swine	53%	53%	53%	55%	56%	58%	64%

Expert judgments of Mészáros, 2015 were drawn on to the further stratification of deep litter by mixing and frequency of removals.

Anaerobic digestion

In Hungary, the first biogas plant utilizing animal manure was established in 2004, so the inventory takes into account the amount of manure treated in the biogas plants from 2004 onwards. For the period before that, we report notation key 'NO' in the CRF tables. Detailed data on agricultural wastes treated in biogas plants have been collected by the Hungarian Energy and Public Utility Regulatory Authority based on Regulation No 11/2017. (VIII. 25.) since 2017. However, according to the Hungarian Statistical Act (Act CLV of 2016 on Official Statistics), these detailed feedstock statistics are confidential, i.e., some feedstocks are used by less than three plants. Based on the amount of manure used in biogas plants, we determine the proportion of slurry and farmyard manure (FYM) treated in biogas plants for the calculation of emissions from 3.B Manure management and based on this data collection we also calculate the emissions from the application of biogas digestate for the 3Da2c sector. In order not to violate the statistical law, but to comply with the transparency requirement of the GHG inventory, in

Table 5.3.4 we report the proportion of slurry and solid manure used in the biogas plant, by animal species.

Table 5.3.4 Proportions of slurry/solid manure used for biogas production

	2000	2005	2010	2015	2020
Swine, slurry	0%	<i>0.3%</i>	<i>2.9%</i>	6.2%	5.9%
Swine, solid	0%	<i>0.0%</i>	<i>0.2%</i>	0.4%	0.5%
Cattle, slurry	0%	<i>4.2%</i>	<i>30.0%</i>	39.9%	25.9%
Cattle, solid	0%	<i>0.1%</i>	<i>0.9%</i>	1.9%	1.7%
Broiler, solid	0%	<i>0.0%</i>	<i>0.0%</i>	0.0%	0.0%
Turkey, solid	0%	<i>0.0%</i>	<i>0.4%</i>	1.0%	1.0%
Laying hen, solid	0%	<i>0.0%</i>	<i>0.1%</i>	0.3%	0.5%

Values in italics are estimates based on “other biogas” production and feedstock consumption statistics.

The amount of “other biogas” production and the biogas production potential from the feedstock were verified, ensuring consistency of data between the energy, waste, and agriculture sectors in terms of emissions from anaerobic digestion. Based on this verification, we found that on average about 25% of other biogas produced in Hungary comes from animal manure. We used this verification for gap filling for the period 2004-2015 and estimated the amount of manure used in biogas plants based on the amount of “other biogas” produced as follows. Based on available data, it was found that the manure N used for biogas production averaged 4.16 tones N per TJ. The manure N used for each year based on biogas production was thus estimated and distributed among the different animal species based on the 2018 data.

Trends in data on animal waste management system distribution

The most significant change occurred in the poultry manure management. From 2000 to 2020, the proportion of the liquid manure had dropped from 26 per cent to almost zero for laying hens. Previously, the semi-solid manure was diluted by water and handled as liquid manure, but recently the semi-solid manure is rather dried than diluted and handled as solid manure. Thus, the liquid manure technology has been replaced by the drying technology as a result of environmental restrictions (Pazsiczky et. al, 2006). The other notable change in the poultry manure management is the decrease of the proportion of grazing for geese. As a result of the Avian influenza scare, the animals are kept in stalls rather than pastures.

For pigs, and in particular dairy cattle, there is an increase in slurry farming practices, which is also linked to an increase in the concentration of production. A growing proportion of the herd is concentrated on large farms where slurry farming is the dominant practice. At the same time, since 2004 part of the slurry has been used in biogas plants.

The share of manure used in biogas plants peaked in 2018 and has declined slightly in recent years, on one hand due to a decrease in other biogas production (see also 5B2) and the other hand due to a decrease in the share of animal manure in other biogas production compared to other agricultural and food waste.

In case of non-dairy cattle, the AWMS data show an increase in grazing. This can be explained by the increase in the proportion of beef cattle in the non-dairy cattle population, as in Hungary the most common type of beef cattle farming is deep-litter housing and grazing.

Activity data for the base year and 2020 are presented in Table 5.3.5 and Table 5.3.6, respectively. In case of cattle and swine extrapolation and surrogate data were used to complete the time series. For the other livestock categories data for the year 2000 were used for the period 1985-1999 due to lack of information.

Table 5.3.5 Animal waste management distributions for the base year per livestock categories

BY	Liquid	Solid	Pasture	Anaerobic digesters	Other	Deep litter	Yard	Poultry manure with bedding	Poultry manure without bedding
Dairy Cattle	3.6%	40.9%	8.0%	NO	47.5%	44.0%	3.4%	-	-
Non-Dairy Cattle	2.5%	39.8%	16.1%	NO	41.6%	38.3%	3.3%	-	-
Swine	39.5%	59.0%	0.0%	NO	1.5%	0.0%	1.5%	-	-
Poultry	8.2%	22.0%	0.1%	NO	69.7%	-	-	58.0%	11.7%
Sheep	0.8%	44.9%	54.2%	NO	0.0%	-	-	-	-
Goats	0.8%	55.7%	43.4%	NO	0.0%	-	-	-	-
Horses	0.0%	60.0%	40.0%	NO	0.0%	-	-	-	-

Table 5.3.6 Animal waste management distributions for the year 2020 by livestock categories

2020	Liquid	Solid	Pasture	Anaerobic digesters	Other	Deep litter	Yard	Poultry manure with bedding	Poultry manure without bedding
Dairy Cattle	17.2%	29.3%	4.9%	7.1%	41.6%	33.9%	7.7%	-	-
Non-Dairy Cattle	1.8%	16.1%	25.2%	1.7%	55.2%	45.8%	9.4%	-	-
Swine	64.1%	22.3%	0.0%	4.1%	9.5%	7.4%	2.1%	-	-
Poultry	0.0%	20.3%	0.0%	0.4%	79.2%	-	-	65.1%	14.1%
Sheep	0.0%	56.4%	43.6%	NO	0.0%	-	-	-	-
Goats	0.0%	48.6%	51.4%	NO	0.0%	-	-	-	-
Horses	0.0%	60.3%	39.7%	NO	0.0%	-	-	-	-

5.3.2.2.2 Livestock Number

Livestock population data provided by the HCSO are used for the estimation. For more details on the calculation of the annual average population and the activity data see section 5.2.2.1. The enhanced livestock characterization was used for the key categories according to the IPCC methodology. The livestock population data for swine by sub-categories are shown in Table 5.3.7.

Table 5.3.7 Swine population and trends from the BY to 2020

Year	Animal Population 1,000 head						
	Piglets under 20 kg	Young pigs, 20-50 kg	Pigs for fattening over 50 kg	Breeding boars	Breeding sows	Guilts not yet mated	Sows mated for the first time
BY	2,015.2	1,718.2	4,341.0	25.2	690.9	76.1	96.4
1990	1,953.1	2,626.3	3,239.6	27.2	657.5	116.3	88.5
1991	1,612.3	2,349.7	3,090.6	25.1	563.0	104.0	64.4
1992	1,310.1	1,844.5	2,436.3	20.4	486.7	81.7	57.8
1993	1,222.7	1,744.1	2,245.4	18.0	446.0	77.2	52.0
1994	1,050.0	1,499.5	1,958.4	15.4	372.5	66.4	44.8
1995	1,107.2	1,458.5	1,921.6	15.3	404.7	64.6	51.1
1996	1,256.8	1,523.9	2,146.8	15.7	429.8	67.5	53.0
1997	1,187.5	1,302.1	2,039.4	14.3	356.3	56.7	56.3
1998	1,247.4	1,407.0	2,073.5	14.0	364.0	65.0	75.8
1999	1,281.3	1,503.1	2,299.8	14.8	396.9	56.3	56.8
2000	1,208.0	1,302.8	2,143.5	14.2	359.8	56.7	61.2
2001	1,260.5	1,108.0	1,984.5	12.5	342.2	54.7	61.0
2002	1,361.2	1,136.7	2,043.0	12.8	368.0	60.3	68.0
2003	1,281.8	1,157.8	2,150.9	12.0	362.3	56.0	56.7
2004	1,064.3	1,015.0	1,885.3	9.8	309.2	50.0	51.3
2005	998.7	916.5	1,701.9	10.0	291.5	50.7	52.5
2006	976.0	933.1	1,635.4	8.7	282.2	54.8	53.5
2007	1,015.3	934.2	1,700.3	7.8	279.0	52.0	50.3
2008	877.5	848.3	1,595.3	6.8	249.7	46.3	40.7
2009	757.4	795.4	1,374.1	6.0	226.5	45.0	43.5
2010	763.4	751.7	1,374.1	6.5	225.2	42.2	44.7
2011	751.5	748.6	1,326.9	5.7	217.6	43.4	37.8
2012	707.1	726.7	1,256.8	5.0	205.8	42.0	38.1
2013	724.0	683.6	1,250.4	4.8	194.2	44.1	42.8
2014	761.3	725.0	1,288.5	4.9	198.5	43.5	43.2
2015	784.2	741.4	1,308.3	4.7	201.1	44.8	42.5
2016	710.9	665.9	1,370.3	4.2	184.9	43.7	40.9
2017	683.0	636.8	1,271.5	3.4	175.0	41.0	36.9
2018	695.9	633.9	1,275.1	3.0	176.6	44.1	36.4
2019	690.2	626.4	1,228.9	2.7	167.9	45.3	35.0
2020	730.6	574.1	1,277.3	2.7	164.2	46.1	36.1
Trend BY-2020	-63%	-78%	-61%	-90%	-75%	-60%	-59%
Trend 2005-2020	-27%	-37%	-25%	-73%	-44%	-9%	-31%

5.3.2.2.3 Annual Average Nitrogen Excretion Rates (N_{ex})

For the values of annual average nitrogen excretion rates country specific (Tier 2) coefficients derived based on the Equation 10.31 of the 2006 IPCC Guidelines were used for Dairy Cattle, Non-dairy Cattle, Swine, Laying hens and Broilers.

In the calculation for breeding sows three different stages as gestation, lactation, and the period 'between weaning and mating' were distinguished. The daily nitrogen intake/retention was determined for each period and annual values were calculated as the weighted mean using the length of the periods as weighting factors. According to the Hungarian practices the length of gestation and lactation are 114 and 21 days, respectively. While the period between two successive farrowing decreased gradually across the time series. Annual values are provided in Table 5.3.11.

For broilers four-phase feeding was assumed for the period 2005-2018 and three phases from 2004 backwards. Time series consistency was ensured based on the time series overlap approach of the 2006 IPCC Guidelines (Volume 1, Chapter 5). Therefore, for the years 2005-2007 the three- and four-phase feeding systems were assumed, parallelly.

There was no need to distinguish between different stages in case of laying hens.

Nitrogen intake

To the above equation Nitrogen intakes were determined from the crude protein content of the dietary components for all subcategories of these animals. The crude protein intakes were multiplied by 0.16, which is the fraction of nitrogen in protein, to convert the protein content into nitrogen. In the case of cattle and swine subcategories (breeding sows and breeding boars excepted) crude protein content in the diet was calculated from the feed ingredients. Data on crude protein contents of each component were taken from the so-called 'feed database' containing the laboratory measurements of all kinds of feed used for animal nutrition in Hungary. The feed database is available in the Hungarian Nutrition Codex, 2004. In the case of cattle, nitrogen intakes were determined in conjunction with the examination of gross energy intake (see also section 5.2.2.2).

In respect of breeding swine, laying hens and broilers data on crude protein content of the feed (CP%) in proportion of dry matter intake (DMI) was provided from the animal feed monitoring system operated by the AKI. Therefore, the nitrogen intake for a certain animal subcategory and stage was calculated using the Equation 10.32A of the 2019 Refinement:

$$N_{intake(T,i)} = DMI_{T,i} \cdot \frac{CP\%_{T,i}}{6.25}$$

Equation 5.3

Where:

$N_{intake(T,i)}$ = daily N consumed per animal of category (T) and stage (i), kg N animal⁻¹ day⁻¹

$DMI_{T,i}$ = dry matter intake per animal in a certain stage (kg DMI animal day⁻¹)

CP%_{T,i} = per cent crude protein in dry matter

6.25 = conversion from kg of dietary protein to kg of dietary N, kg feed protein (kg N)⁻¹

In Hungary a feed monitoring system started to operate in 2016, with a retrospective data collection for the year 2005. For the year 1990 standards of the DMI and CP% intakes taken from the Hungarian Nutrition Codex, 1990 were applied and interpolation was used to complete the time series. According to the expert opinions and the depth interviews of the AKI, the Hungarian Nutrition Codex provided

the most appropriate values of DMI and CP% for swine. While for broiler and laying hens the breeder's guides seemed to be the most reliable sources before 2005. However, as the research of the AKI revealed, in the years between 2005-2007, the crude protein content of the laying hen and broiler diet could be slightly lower than as it is suggested in the breeder's guides. Therefore, time series overlapping was applied to avoid the time series inconsistency arising from the use of data from two different sources.

Table 5.3.8 shows the trends in the crude protein content in the diet of breeding sows and breeding boars. These trends in the CP% for sows are the result of two opposite effects. The rising productivity resulted in an increase in the protein demand and the nitrogen intake. This rising trend was maintained to 2010 after which the amino acid supplements lead to decreasing trends in the crude protein content in the breeding sow diet. Trends in the CP% for breeding boar shows a slight increase over the inventory period.

The trend in the N-intake of broiler is also driven by the abovementioned two contrary effects. The growing living weight result in an increase in the protein demand and the nitrogen intake. In 2007 the N-intake reached a peak. After that N-intake decreased gradually, due to the amino acid supplements. Finally, the N-intake started to increase slightly again in the last two years. The CP% of the diet which was the source of the N-intake estimate are provided in *Table 5.3.9*

The overall slightly decreasing trend in the N-intake of laying hens reflects the improvement of the feeding practices and the importance of the amino acid supplements *Table 5.3.10*.

*Table 5.3.8 Crude protein content in the diet of breeding sow and breeding boar in proportion of DMI
BY, 1990-2018*

Year	gestating sow	lactating sow	weighted average for breeding sow	breeding boar
	CP(%)			
BY	13.8	16.1	14.1	15.0
1990	13.8	16.1	14.1	15.0
1991	13.8	16.2	14.1	15.1
1992	13.9	16.2	14.2	15.1
1993	13.9	16.3	14.2	15.2
1994	13.9	16.4	14.2	15.2
1995	14.0	16.5	14.3	15.3
1996	14.0	16.5	14.3	15.3
1997	14.0	16.6	14.3	15.4
1998	14.1	16.7	14.4	15.4
1999	14.1	16.8	14.4	15.5
2000	14.1	16.8	14.5	15.5
2001	14.2	16.9	14.5	15.6
2002	14.2	17.0	14.5	15.6
2003	14.2	17.1	14.6	15.7
2004	14.3	17.1	14.6	15.7
2005	14.3	17.2	14.7	15.8
2006	14.3	17.2	14.7	15.8

Year	gestating sow	lactating sow	weighted average for breeding sow	breeding boar
	CP(%)			
2007	14.3	17.2	14.7	15.9
2008	14.3	17.2	14.7	15.9
2009	14.3	17.2	14.7	16.0
2010	14.3	17.2	14.7	16.0
2011	14.3	17.0	14.6	16.1
2012	14.0	17.1	14.4	16.1
2013	13.9	17.2	14.3	16.2
2014	13.7	16.9	14.1	16.2
2015	13.6	16.7	14.0	16.3
2016	13.4	16.7	13.8	16.3
2017	13.3	16.9	13.8	16.4
2018	13.4	16.9	13.9	16.4

Table 5.3.9 Crude protein content in the diet of broilers in proportion of DMI BY, 1990-2018

Year	CP% (breeder's recommendation)			Year	CP% (AERI's data collection)			
	starter	grower I.	finisher		starter	grower I.	grower II.	finisher
BY	23.0	20.0	18.5	2005	22.1	20.3	20.2	19.4
1990	23.0	20.0	18.5	2006	22.0	20.3	20.0	19.2
1991	23.0	20.0	18.5	2007	22.0	20.3	19.9	19.1
1992	23.0	20.0	18.5	2008	22.0	20.3	19.7	19.0
1993	23.0	20.0	18.5	2009	21.9	20.3	19.6	18.9
1994	23.0	20.0	18.5	2010	21.9	20.3	19.5	18.7
1995	22.9	20.0	18.6	2011	21.9	20.2	19.2	18.6
1996	22.9	20.1	18.6	2012	21.6	20.1	19.0	18.4
1997	22.8	20.1	18.7	2013	21.4	19.9	19.1	18.3
1998	22.8	20.2	18.8	2014	21.3	19.8	19.0	18.5
1999	22.7	20.2	18.8	2015	21.1	19.9	18.9	18.3
2000	22.7	20.3	18.9	2016	21.2	19.8	19.0	18.2
2001	22.6	20.3	19.0	2017	21.4	20.0	19.3	18.3
2002	22.6	20.4	19.1	2018	21.5	20.1	19.4	18.5
2003	22.5	20.4	19.1					
2004	22.5	20.5	19.2					
2005	22.4	20.5	19.3					
2006	22.4	20.6	19.3					
2007	22.3	20.6	19.4					

Table 5.3.10 Crude protein content in the diet of laying hens in proportion of DMI BY, 1990-2018

Year	CP% (breeder's recommendation)	Year	CP% (AERI's data collection)
BY	17.4	2005	17.2
1990	17.4	2006	17.2
1991	17.5	2007	17.2
1992	17.6	2008	17.1
1993	17.6	2009	17.1
1994	17.7	2010	17.1
1995	17.8	2011	17.4
1996	17.6	2012	17.5
1997	17.5	2013	16.8
1998	17.4	2014	16.7
1999	17.4	2015	16.7
2000	17.4	2016	16.7
2001	17.4	2017	16.6
2002	17.4	2018	16.4
2003	17.4		
2004	17.4		
2005	17.4		
2006	17.3		
2007	17.2		

N retention

N retained by gestating sows and lactating sows were calculated using the Equation 10.33A and 10.33B of the 2019 Refinement:

$$N_{\text{retention},i} = N_{\text{gain},i} + N_{\text{piglets},i}$$

Equation 5.4

Where:

$N_{\text{retention},i}$ = amount of N retained by the sow in the stage i (head · day⁻¹)

$N_{\text{gain},i}$ = amount of N retained in the sow in the stage i (head · day⁻¹)

$N_{\text{piglets},i}$ = amount of N in piglets in the stage i (heads · day⁻¹)

i = stage ($i=1$ gestation, $i=2$ lactation, $i=3$ period 'between weaning and mating')

$$N_{\text{piglets},i} = 0.0256 \cdot \text{LITSIZE}_i \cdot \text{WG}_{\text{piglets},i}$$

Equation 5.5

Where:

LITSIZE_i = litter size, in the stage i , heads;

$WG_{piglets,i}$ = weigh gain of piglets, in the stage i , head-day-1;

0.0256 = N-content of weight gain (kg/kg) Lfl, 2013

For sows in the period between two successive farrowing and breeding boars the nitrogen retention was calculated based on the daily weight gain.

Background data as litter size, weaning weight and days between two successive farrowing are provided in Table 5.3.11.

Data was compiled by the HMS, based on the annual yearbooks of 'Results of Pig Breeding' 1985-2018, published by the NFCSO. Piglets weight at birth was assumed to be 1.3 kg.

Table 5.3.11 Background data to the calculation of nitrogen retention rate of breeding sows

Year	piglet weight at weaning	number of piglets at birth	number of piglets at weaning	period between two successive farrowing
	kg	heads	heads	days
BY	5.9	9.8	8.9	178.4
1990	6.3	10.1	8.8	178.1
1991	6.2	10.1	8.9	178.8
1992	6.2	10.2	8.8	181.2
1993	6.2	9.8	8.5	181.2
1994	6.2	9.8	8.4	182.2
1995	6.4	10.0	8.4	181.3
1996	6.5	10.1	8.5	181.0
1997	6.5	10.1	8.5	182.9
1998	6.2	10.2	9.1	181.6
1999	6.2	10.3	9.3	178.6
2000	6.3	10.4	9.3	180.2
2001	6.4	10.3	9.2	173.4
2002	6.5	10.2	9.1	173.3
2003	6.3	10.2	9.2	172.1
2004	6.4	10.2	9.2	170.7
2005	6.6	10.2	9.2	170.6
2006	6.7	10.3	9.2	170.4
2007	6.8	10.3	9.4	169.1
2008	7.0	10.4	9.4	168.9
2009	7.3	10.4	9.5	167.5
2010	7.6	10.5	9.5	166.6
2011	7.7	10.6	9.6	166.0
2012	7.7	10.7	9.8	162.6
2013	7.7	10.9	9.9	162.2
2014	7.7	11.1	10.0	162.0
2015	7.8	11.2	10.1	161.9
2016	7.8	11.3	10.2	161.8
2017	7.8	11.5	10.3	161.4
2018	7.9	11.4	10.2	161.8

Source: NFCSO

N retention for laying hens was calculated from the production data using the Equation 10.33D of the IPCC Refinement:

$$N_{\text{retention}} = \left[N_{\text{LW}} \cdot \text{DWG} + \left(\frac{N_{\text{egg}} \cdot \text{EGG}}{1000} \right) \right]$$

Equation 5.6

Where:

$N_{\text{retention}}$ = daily nitrogen retention of laying hens, kg N·head⁻¹·day⁻¹;

N_{LW} = average content of nitrogen in live weight, kg N·kg head⁻¹. Default value of 0.028 provided in the 2019 Refinement was applied;

DWG = average daily weight gain, kg·head⁻¹·day⁻¹;

N_{EGG} = average content of nitrogen in eggs, kg N·kg egg⁻¹. Default value of 0.0185 provided in the 2019 Refinement was used.

EGG = egg mass production, g egg·head⁻¹·day⁻¹.

Data on egg production was obtained from the HCSO (Table 5.3.12). Average daily weight gain (DWG) was calculated from the daily weight gain of the typical laying hen breeds, as Tetra, Lohman and Hy-Line. Data on the distribution of typical breeds in Hungary were provided by the Hungarian Poultry Board. The average egg weight was calculated similarly, based on the egg weight of the typical laying hen breeds.

Table 5.3.12 Background data to the calculation of nitrogen retention rate of laying hens

Year	Egg production [egg·head ⁻¹ ·year ⁻¹]
BY	206
1990	206
1991	189
1992	206
1993	218
1994	227
1995	220
1996	200
1997	212
1998	207
1999	202
2000	208
2001	213
2002	212
2003	210
2004	212
2005	208
2006	205
2007	218
2008	215
2009	215
2010	218
2011	214
2012	217
2013	208
2014	214
2015	218

Year	Egg production
	[egg· head ⁻¹ ·year ⁻¹]
2016	225
2017	227
2018	233

Source: HCSO, 2019

Nitrogen retention for broilers was calculated using the Equation 10.33E of the 2019 Refinement:

$$N_{\text{retention}} = \frac{(BW_{\text{Final}} - BW_{\text{Initial}}) \cdot N_{\text{gain}}}{\text{production_period}}$$

Equation 5.7

Where:

N retention= amount of N retained in animal (head) day⁻¹

BW_{Final} = Live weight of the animal at the end of the stage (kg)

BW_{Initial} = Live weight of the animal at the beginning of the stage (kg)

N_{gain} = the fraction of N (kg) retained per kg BW gain (kg kg⁻¹)

Production period = length of time from chick to slaughter (fattening duration)

N_{gain} was assumed to be 0.0304 kg kg⁻¹ based on Haenel et al., 2018 and Haenel és Dämmgen, 2009. This value relates to the 40 days fattening duration, but in order to ensure the time series consistency this value was applied for the whole inventory period. Data on BW_{Final} was obtained from the slaughterhouse statistics of the AERI. This statistic provides data on living weight before slaughtering. The value of BW_{initial} was estimated to be 0.042 kg based on expert judgement. Fattening duration was estimated to be 49, 42 and 40 days for the years 1994, 2007 and 2018 based on the Breeders Management Manuals of Arbor Acres and Aviagen, respectively; and interpolation was used to complete the timeseries.

Table 5.3.13 Background data to the calculation of the nitrogen retention rate in broilers

Year	BW _{final}	Production period (fattening duration)
	kg	day
BY	1.9	49.0
1990	1.9	49.0
1991	1.9	49.0
1992	1.9	49.0
1993	1.9	49.0
1994	1.9	49.0
1995	1.9	48.5
1996	1.9	47.9
1997	1.9	47.4
1998	1.9	46.8
1999	2.0	46.3
2000	2.0	45.8
2001	2.0	45.2
2002	2.0	44.7
2003	2.1	44.2
2004	2.0	43.6
2005	2.0	43.1
2006	2.1	42.5

Year	BW _{final}	Production period (fattening duration)
		kg day
2007	2.1	42.0
2008	2.1	41.8
2009	2.2	41.6
2010	2.2	41.3
2011	2.3	41.1
2012	2.3	40.9
2013	2.3	40.7
2014	2.4	40.4
2015	2.4	40.2
2016	2.4	40.0
2017	2.5	40.0
2018	2.5	40.0

Values of fraction of annual N-intakes that is retained by animals ($N_{\text{retention}}$) and their sources are summarized in Table 5.3.14. The resulted values of N-excretion for Dairy Cattle and Non-dairy Cattle are provided in Table 5.2.7, Table 5.2.9 and Table 5.2.10, respectively, while values of N excretion for Swine are presented in Table 5.3.15.

Table 5.3.14 $N_{\text{retention}}$ rates and their sources

Animal species	$N_{\text{retention}}$	Source
Dairy Cattle	0.20	2006 IPCC GLs
Non-Dairy Cattle	0.07	2006 IPCC GLs
Swine		
Piglets under 20 kg	0.48	Fébel and Gundel, 2007
Young pigs, 20-50 kg	0.34	Fébel and Gundel, 2007
Pigs for fattening over 50 kg	0.34	Fébel and Gundel, 2007
Breeding sows, weighted mean (2020)	0.34	country-specific (calculated based on the Hungarian production data, annually)
Gestating Sows	0.30	country-specific
Lactating Sows	0.42	country-specific
Sows between weaning and mating	0.26	country-specific
Breeding boars (2020)	0.08	country-specific (calculated based on the Hungarian production data)
Guilts not yet mated	0.34	Fébel and Gundel, 2007
Sows mated for the first time	0.34	Fébel and Gundel, 2007
Laying hens	0.27	country-specific (calculated based on the Hungarian production data, annually)
Broilers	0.55	country-specific (calculated based on the Hungarian production data, annually)

Table 5.3.15 Annual average Nitrogen excretion rates (N_{ex}) for Swine

Sub-categories	N_{ex}
	[kg head ⁻¹ year ⁻¹]
Piglets under 20 kg	3.0
Young pigs, 20-50 kg	8.6
Pigs for fattening over 50 kg	12.3
Breeding sows (weighted average, BY)	15.9
<i>Gestating Sows</i>	13.3
<i>Lactating Sows</i>	33.7
<i>Sows between weaning and mating</i>	13.4
Breeding sows (weighted average, 2020)	15.8
<i>Gestating Sows</i>	12.6
<i>Lactating Sows</i>	38.1
<i>Sows between weaning and mating</i>	12.9
Breeding boars (BY)	24.4
Breeding boars (2020)	22.1
Guilts not yet mated	9.9
Sows mated for the first time	13.8
Swine, weighted average (BY)	9.9
Swine, weighted average (2020)	9.3

For other livestock categories the default values of nitrogen excretion provided in Table 10.19 of the 2006 IPCC Guidelines were used except Buffalo for which the EMEP/EEA Guidebook (EEA, 2019) were applied (Table 3.9). It should be noted that in the case of nitrogen excretion rate of Buffalo the 2006 IPCC Guidelines refer to the EEA, 2002, thus the use of the 2019 EMEP/EEA Guidebook, which is the most up-to-date emission inventory guidebook of the EEA, seems to be reasonable. Nitrogen excretion rates for 'Other animals' and the related body weights are shown in Table 5.3.16 and Table 5.3.17.

Table 5.3.16 Annual average Nitrogen excretion rates (N_{ex}) for 'Other livestock'

Animal Category	N_{ex}	Source
	[kg head ⁻¹ year ⁻¹]	
Buffalo	82*	2019 EMEP/EEA GB / 2006 IPCC GLs
Sheep	16	2006 IPCC GLs, Eastern Europe
Goats	18	2006 IPCC GLs, Eastern Europe
Horses	41	2006 IPCC GLs, Eastern Europe
Asses & Mules	14	2006 IPCC GLs, Eastern Europe
Poultry (2020)	0.73	Weighted average for 2020
<i>Laying hens (2020)</i>	0.76	<i>Country-specific, calculated periodically</i>

Animal Category	N _{ex} [kg head ⁻¹ year ⁻¹]	Source
Broilers (2020)	0.56	Country-specific, calculated periodically
Turkey	1.84	2006 IPCC GLs, Eastern Europe
Ducks	0.82	2006 IPCC GLs, Eastern Europe
Geese	0.55**	2019 EMEP/EEA GB
Guinea Fowls	0.36	as default for Broilers
Rabbit	8.1	2006 IPCC GLs

*2006 IPCC GLs refer to the 2002 EMEP/EEA GB. Therefore, the 2019 EMEP/EEA GB as the more updated version of the GB was applied.

**There is no value provided in the 2006 IPCC GLs

Table 5.3.17 Typical animal mass (TAM) for other livestock

Livestock	Weight [kg]	Source/Note
Buffalo	380	Table 10A-6 of 2006 IPCC GLs
Sheep	48.5	Table 10A-9 of 2006 IPCC GLs
Goats	38.5	Table 10A-9 of 2006 IPCC GLs
Horses	377	Table 10A-9, Developed, 2006 IPCC GLs
Asses and Mules	130	Table 10A-9, Developed, 2006 IPCC GLs
Poultry	2.2	Weighted average for 2020
Laying hens (2020)	2.0*	Country-specific
Broiler (2020)	1.5*	Country-specific
Turkey	6.8	Table 10A-9 of 2006 IPCC GLs
Ducks	2.7	Table 10A-9 of 2006 IPCC GLs
Geese	3.5	2019 EMEP/EEA Guidebook
Guinea fowls	0.9	Default for Broiler due to lack of information
Rabbit	1.6	Table 10A-9 of 2006 IPCC GLs

*Please note that Tier 2 is applied, therefore TAM is not used in the calculation. These values are reported for information.

5.3.2.3 Estimation of CH₄ emissions

5.3.2.3.1 Emission factors for CH₄

As Manure Management is a key source the Tier 2 method was applied to calculate the CH₄ emission factors, except Rabbit, for which a default value of 0.08 is given in the Table 10A-9 of 2006 IPCC Guidelines was used. According to the Equation 10.23 of 2006 IPCC Guidelines, development of country-specific emission factors involves determining a weighted average methane conversion factor (MCF) using the estimates of the manure managed in each AWMS and the volatile solid excretion (VS), which means the organic material in livestock manure. The CH₄ emission factor also depends on the maximum methane producing capacity (B₀) for the livestock categories. The values of these components in the above-mentioned equation were calculated as it is delineated in the following sub-sections.

5.3.2.3.1.1 Volatile solid excretion per day (VS)

Country-specific values of VS for Cattle, Laying hens and Broilers were calculated according to the Equation 10.24 of 2006 IPCC Guidelines. Values needed for this calculation are the gross energy intake (GE), and its fractional digestibility, DE. The estimation of these values is detailed in Chapter 6.2.2.2. Metabolizable and digestible energy of feed for Laying hens and Broilers were calculated similarly, based on feeding practices. Forage composition parameters were taken from the Hungarian Nutrition Codex, 2004.

For the ash content of the manure the IPCC default value (8%) was applied due to lack of country-specific values. Similarly, the urinary energy which is also required to the Equation 10.24 was calculated as 0.04·GE according to the 2006 IPCC Guidelines.

Table 5.3.18 contain the values of volatile solid excretion rate and the emission factors for non-dairy cattle sub-categories for the BY and 2020, respectively. Following the recommendations of the 2020 and 2021 UNFCCC reviews (A.10, 2020 and A.1 2021), Table 5.3.18 has been restructured and the unit error have been corrected. Tables 5.3.17 and 5.13.18 of the 2020 and 2021 submissions have been merged here, and the values previously included in that table are presented in a single table in this submission.)

Table 5.3.18 Volatile solid excretion rate and CH₄-Emission Factor for Non-Dairy Cattle in the BY and 2020

Non-dairy cattle sub-categories		BY		2020	
		VS excretion	CH ₄ -Emission Factor	VS excretion	CH ₄ -Emission Factor
		kg DM·head ⁻¹ ·day ⁻¹	kg·head ⁻¹ ·yr ⁻¹	kg DM·head ⁻¹ ·day ⁻¹	kg·head ⁻¹ ·yr ⁻¹
<1 year	Bovines for slaughter and other calves (male)	1.6	5	1.6	7
	Bovines for slaughter and other calves (female)	1.5	4	1.6	7
1-2 year	Bovines (male)	3.2	13	3.4	16
	Heifers for slaughter and other heifers	3.3	11	3.4	11
>2 year	First calf heifers	3.4	14	3.5	13
	Mature Non-Dairy (male)	3.6	15	3.9	15
	Heifers for slaughter	3.5	14	3.7	14
	Beef Cow	2.8	10	3.0	11

For the other livestock categories, as Swine, Buffalo, Sheep, Horses, Asses and Mules, Ducks and Geese the IPCC default values provided for Eastern Europe or Developed countries in the Table 10A-6-10A-9 in the 2006 IPCC Guidelines was used. IPCC default values for geese and guinea fowls are not available;

hence values for ducks and broilers were used, respectively. In the case of swine, the default VS for market swine provided in the Table 10A-7 of the 2006 IPCC Guidelines was applied. According to the 2006 IPCC Guidelines the body mass of breeding swine and market swine are 180 and 50 kg, respectively. In Hungary, the average body mass of swine is about 64 kg, thus in the absence of country-specific value of VS the use of the market swine VS is reasonable.

5.3.2.3.1.2 *Maximum CH₄ producing capacity (B₀) values*

Due to lack of country-specific data default values listed in Tables 10A-4-10A-9 of the 2006 IPCC Guidelines were applied.

5.3.2.3.1.3 *Methane conversion factors (MCF)*

Default MCFs for different manure management systems by average annual temperatures provided in Table 10.17 of 2006 IPCC Guidelines were used. The annual mean temperature in most parts of Hungary is between 10 and 11 °C. Thus, MCFs values provided for cool climate zone were applied for Pasture/Range/Paddock, Solid and both Poultry manure.

The choice of MCFs for liquid manure and deep litter required the disaggregation of livestock categories by annual average temperatures. However, Hungary does not have either large animal populations or multiple climate regions. To the further stratification the annual mean temperature and animal livestock data by counties were used. The detailed climate data (i.e., annual mean temperatures for 19 counties of Hungary) were taken from the HMS climate database, while the detailed livestock data from the Farm Structure Survey, conducted in 2013. The resulted proportion of animal population by average annual temperature and livestock categories are provided in Table 5.3.19.

Table 5.3.19 Distribution of main livestock categories by average annual temperatures

Average annual temperature	Proportion of animal population				
	Dairy Cattle	Other Cattle	Swine	Laying Hens	Broiler
11	67%	67%	81%	69%	64%
≤10	33%	33%	19%	31%	36%

Beyond the average annual temperature, the IPCC methodology differentiates between liquid manure with 'natural crust cover' and without, as well as 'deep litter < 1 month' and 'deep litter > 1 month'. According to the expert opinion of the NARIC Institute of Agricultural Engineering (Mészáros, 2015), duration of deep litter is generally longer than one month in Hungary.

To determine the MCF for pig and cattle slurry, the proportion of covered and uncovered manure stores was determined (see also Section 5.3.2.2.1) and the IPCC default values for these were weighted according to the proportion of covered and uncovered manure tanks.

Anaerobic digestion

The technological specificities of the use of animal manure in biogas plants were investigated by AKI in 2021. The study was financed by the Ministry of Agricultural and a report an “The Hungarian biogas plants technological survey” was submitted to the Ministry. The AKI’s project found that in Hungary, most manure is treated in agricultural biogas plants and not in centralized ones. Consequently, emission losses during transport and storage of manure are not typical. Thus, slurry used in biogas plants is not stored on livestock farms, either on the site of the biogas facilities. In the biogas plant, manure is fed continuously, so the manure is transported via pipeline to the biogas plant, which is mostly located on the farm. Most of the solid manure treated in biogas plants comes from calves and is stored for up to two weeks before being used in biogas plants.

The MCFs for manure treated in anaerobic digesters were determined based on the 2019 Refinement tier 2 methodology using equations provided in the Annex 10A.4 of the 2019 Refinement. The 2019 Refinement provides a much more detailed and scientifically more improved methodology for calculating the MCF for anaerobic digestion than the 2006 IPCC Guidelines. I.e., it separates the MCF from the digester during pre-storage and storage of digestate, allowing open and covered storage to be taken into account. Therefore, we believe, that the 2019 Refinement methodology reflects better the Hungarian circumstances than the 2006 IPCC Guidelines.

The following assumptions were used in the calculations, and the parameters required for the calculations were determined as follows:

For the leakage rate of digester (L_{dig}) the default value of 0.01 was assumed, presuming high quality biogas digesters. This assumption is also supported by the verification calculation carried out, as the estimated biogas yields are in good agreement with the amount of biogas produced, confirming that there should not be any significant loss from the digester. The value of 0.046 is used for relative amount of residual gas (μ_{rg}), because in Hungary the mesophilic digestion is dominant. In accordance with the report of the AKI the share of gastight storage of the digestate (x_{gts}) was 21% in Hungary, in 2020. No data was available on the leakage rate of the storage ($L_{sto,gt}$), therefore $L_{sto,gt} = L_{dig}$ was assumed. Methane conversion factor for the non-gastight storage of digestate, (MCF_{ngts}) a value of 11% was assumed, similarly to the country-specific value of open cattle slurry tank with natural crust, because the viscosity of the digestate is similar to the cattle liquid manure and the co-fermented crop residues allow the formation of natural crust. Values of methane conversion factor for the non-gastight storage of digestate (MCF_{ps}) were calculated based on the values of the MCFs for manure not treated in biogas plants, considering the significantly lower length of manure storage.

IPCC Guidelines provide no methane conversion factor for Yard therefore the MCF of Solid was applied for yard manure. Methane conversion factors used in the inventory are provided in Table 5.3.20.

Table 5.3.20 Methane conversion factors for manure management systems

Manure Management System	MCF [%]
Pasture range and paddock	1
Solid storage and dry lot	2
Liquid system (2020)	
Cattle	14.2
Swine	13.7
Poultry	18.4

Other AWMS	
Anaerobic digestion (2020)	
Cattle	2.2
Swine	1.5
Poultry	1.4
Cattle deep bedding	18.3
Swine deep bedding	18.6
Yard	2
Poultry manure with litter	1.5
Poultry manure without litter	1.5

1.1.1.4 Estimation of direct N₂O emissions from Manure Management

Default emission factors from 2006 IPCC Guidelines were used. In the case of cattle and swine liquid manure covered and uncovered manure tanks were distinguished, similarly to the selection of MCF values (see section above for further details). Mixing of cattle and swine deep bedding does not occur in practice in Hungary (expert judgement, Fenyvesi, 2015). Therefore, for cattle and swine deep bedding 'no mixing' was assumed.

In the IPCC Guidelines emission factor is unavailable for Yard, therefore the emission factor for solid manure was applied. Emission factors used in the inventory to estimate N₂O emissions from manure management are listed in Table 5.3.21. In response to a recommendation from the UNFCCC review 2019, Table 5.3.15 was supplemented with the sources of the emission factor indicating where weighted averages of the IPCC default values were applied.

Table 5.3.21 Emission factors used for the estimation of N₂O emissions

Manure management system	N ₂ O-N emission factors [kg N ₂ O-N kg ⁻¹ N _{ex}]	Source
Solid storage and dry lot	0.005	Table 10.21 of the 2006 IPCC Guidelines
Liquid system		
Cattle (2020)	0.003	weighted average depending on the ratio of covered to uncovered slurry tanks
Swine (implied, 2020)	0.003	weighted average
Sows (2020)	0.003	weighted average depending on the ratio of covered to uncovered slurry tanks
Fattening pigs (2020)	0.003	weighted average depending on the ratio of covered to uncovered slurry tanks

Poultry	0.000	Table 10.21 of the 2006 IPCC Guidelines (without natural crust cover)
Other AWMS		
Cattle deep bedding	0.010	Table 10.21 of the 2006 IPCC Guidelines (No mixing)
Swine deep bedding	0.010	Table 10.21 of the 2006 IPCC Guidelines (No mixing)
Yard	0.005	as solid due to lack of default value provided in the 2006 IPCC Guidelines
Poultry manure with litter	0.001	Table 10.21 of the 2006 IPCC Guidelines
Poultry manure without litter	0.001	Table 10.21 of the 2006 IPCC Guidelines

5.3.2.4 Estimation of indirect N₂O emissions from Manure Management

5.3.2.4.1 Indirect N₂O emissions through volatilization losses from manure management

Following the 2006 IPCC Guidelines, indirect N₂O emissions due to volatilization of N from manure management were calculated using the Tier 2 methodology.

The country-specific value of fraction of N that is volatilized as NH₃ and NO_x (Frac_{GASMS}) was calculated based on the NH₃ and NO_x emissions from 3.B Manure management reported to the UNECE under the LRTAP Convention. Hungary applies the 2019 EMEP/EEA Emission Inventory Guidebook to calculate emissions of air pollutants from agriculture. Tier 2 and Tier 3 methodology is applied for Cattle, Sheep, Swine, Laying hens and Broiler and Tier 1 for other livestock categories. NO_x emissions are calculated based on Tier 1 method. Derivation of the amount of manure nitrogen that is lost due to the volatilization of NH₃ and NO_x based on the Hungarian air pollutant emission inventory under the UNECE/LRTAP Convention is demonstrated in Table 5.3.22 for the year 2020. Time series of volatilization losses were calculated similarly for all years of the inventory period (Table 5.3.23).

Table 5.3.22 Volatilized N as NH₃ and NO_x from manure management systems for 2020

CRF code	Long name	NO _x (as NO ₂)	NH ₃	Total N volatilized [kg N]
3B1a	Manure management - Dairy cattle	0.14	7.23	5,992,297
3B1b	Manure management - Non-Dairy Cattle	0.15	7.27	6,031,656
3B2	Manure management - Sheep	0.01	1.62	1,341,533
3B3	Manure management - Swine	0.04	7.61	6,276,888
3B4a	Manure management - Buffalo	0.00	0.03	27,640
3B4d	Manure management - Goats	0.00	0.02	18,685

CRF code	Long name	NO _x (as NO ₂)	NH ₃	Total N volatilized [kg N]
3B4e	Manure management - Horses	0.00	0.41	334,794
3B4f	Manure management - Mules and asses	0.00	0.01	7,601
3B4gi	Manure management - Laying hens	0.12	1.44	1,226,068
3B4gii	Manure management - Broilers	0.57	4.04	3,503,998
3B4giii	Manure management - Turkeys	0.08	1.69	1,414,659
3B4giv	Manure management - Other poultry	0.07	1.71	1,428,026
3B4h	Manure management - Other animals (Rabbit)	0.00	0.57	471,954
3B	Manure management	1.18	33.65	28,075,800

*We note that the '0.00' values are due to the rounding of small numbers rather than inferring those emissions are zeros.

Table 5.3.23 NH₃-N and NO_x-N volatilization losses of manure management systems 1990-2020

Year	Total N volatilized	Frac _{GasMS}
	kg N	%
BY	63,133,402	22%
1990	57,141,728	22%
1991	51,699,038	21%
1992	44,054,276	21%
1993	38,278,031	21%
1994	35,134,016	21%
1995	35,513,648	22%
1996	34,816,168	22%
1997	33,927,278	22%
1998	35,103,031	22%
1999	35,561,008	22%
2000	37,080,190	23%
2001	36,035,342	23%
2002	36,429,539	23%
2003	36,744,763	23%
2004	34,986,737	22%
2005	33,021,887	22%
2006	31,934,750	22%
2007	31,655,178	22%
2008	30,913,255	22%
2009	29,427,854	22%
2010	29,627,032	22%
2011	28,975,307	22%
2012	28,392,793	21%
2013	27,757,496	21%
2014	28,666,512	21%
2015	29,364,556	21%

Year	Total N volatilized	Frac _{GasMS}
	kg N	%
2016	29,379,947	21%
2017	28,531,714	21%
2018	28,644,287	21%
2019	28,864,057	22%
2020	28,075,800	21%

To ensure transparency of reporting the volatilization losses, agricultural NH₃ emissions and NO_x emissions from 3.B Manure management are reported in CRF Table 6.

To estimate the indirect N₂O emissions from volatilization default value of 0.01 kg N₂O-N (kg NH₃-N + kg NO_x-N volatilized)⁻¹ for EF₄ given in Table 11.3 of 2006 IPCC Guidelines was used.

5.3.2.4.2 Indirect N₂O emissions through N-leaching and run-off from manure storage

In Hungary there are strict environmental obligations concerning the manure storage arising from the Nitrates Directive (91/676/EEC). The current regulations require to prevent/decrease N-leaching. Therefore, N-leaching from housing and manure storage systems not falling within the scope of nitrate regulations, and nitrogen leaching before the compliance deadlines of the regulations are reported here.

Information concerning Hungarian animal housing and manure storage systems was derived from the Nitrate Database, data from the national Farm Structure Survey (FSS), Codes of Good Agricultural Practice (GAP) and nitrate regulations.

National regulations concerning N-leaching from manure storage systems are as follows:

- Government Decree 27/2006. (II. 7.) on the protection of waters against pollution caused by nitrates from agricultural sources (Nitrate Decree), amending Government Decree 49/2001(IV.3);
- Decree 59/2008 (IV. 29.) FVM on the detailed rules of the action program required for the protection of waters against pollution caused by nitrates from agricultural sources and on the procedures for data provision and registration (hereafter GAP Decree; database established from data provided under the regulation is called Nitrate Database);
- Decree 43/2007. (VI.1.) FVM on Designation of nitrate vulnerable zones by MePAR blocks (MePAR is the abbreviation of the Hungarian Land Parcel Information System).

The watertight construction of animal housings and manure storage systems is generally required under the GAP Decree. The use of impermeable barriers on the underlying strata, to prevent nitrate leaching is needed to ensure the compliance with the Nitrate Decree.

Areas and farms falling within the scope of Nitrate Decree are as follows:

- Farms operating on designated nitrate vulnerable zones and producing more than household needs. (Household needs is here defined as five livestock unit (LSU) or three LSU in the case of poultry; LSU is here defined as 500 kg live weight);

- Farms under the Environmental Permitting Regulations (EPR);
- Large livestock farms (defined in Decree 41/1997/. (V. 28.) FM);
- Areas of manure storage systems and manure processing.




Figure 5.3.2. Maps of areas under the regulation of Nitrate Decree

(Legend: nitrate vulnerable zones designated in 2007 are marked with green color, while areas under the regulations of Nitrate Decree since 2013 are marked with red color)

The compliance deadline for nitrate regulations was 31 December 2014 for liquid/ slurry, and 22 December 2015 for solid manure storage systems in accordance with the amended Nitrate Decree.

(The Government Decree 49/2001(IV.3) contained earlier deadlines, between 2006 and 2014, which were extended in the amended regulation.)

Arising from the nitrate regulations amount of nitrogen leached was determined using the following assumptions:

- Measures for preventing N-leaching from manure management systems should have been started in 2001.
- Measures to improve manure management systems in order to prevent/ decrease N-leaching should have been finished according to the extended deadlines in the amended Nitrate Decree (i.e., at the end of 2014 or 2015).
- After the compliance deadline N-leaching could only occur on small farms (<5 LSU; or <3 LSU for poultry).

To summarize, N-leaching could occur from solid manure and deep litter of animals housed on small farms; and from solid manure, deep litter of animals housed on large farms before the compliance deadlines of nitrate regulations.

In 2000, one year before the introduction of the new legislation, a comprehensive survey was conducted to measure the proportion of farms where actions were needed in order to ensure the compliance with the nitrate regulations. The results of the survey were published in the study of Ráky (2003). For the year 2000 livestock on farms where N leaching could occur was estimated based on this study in the emission estimate.

For each year between 2000 and 2016 the livestock housed on farms with less than five or three LSU was determined based on FSS data. Besides, the fraction of non-compliance was estimated using linear interpolation between 2000 and the deadline of full compliance. (Smooth implementation of the nitrate regulations was assumed.) For the period 1985-1999 data for the year 2000 was applied due to lack of reliable statistics.

Annual amount of manure nitrogen that leached from manure management systems was determined according to the Eq 10.28 of the 2006 IPCC Guidelines, modified for the Hungarian legal circumstances as follows:

$$N_{leaching-MMS} = \sum_S \left[\sum_T \left[\left(N_{(T)} \circ Nex_{(T)} \circ \left(\frac{Frac_{small} + Frac_{NC}}{100} \right)_{(T)} \circ MS_{(T,S)} \right) \circ \left(\frac{Frac_{leachMS}}{100} \right)_{(T,S)} \right] \right]$$

(Equation 5.8.)

Where:

$N_{leaching-MMS}$ = amount of manure nitrogen that leached from manure management systems, kg N yr⁻¹

$N_{(T)}$ = number of head of livestock species/category T in the country

$Nex_{(T)}$ = annual average N excretion per head of species/category T, kg N animal⁻¹ yr⁻¹

$MS_{(TS)}$ = fraction of total annual nitrogen excretion for each livestock species/category T that is managed in manure management system S in the country, dimensionless

$Frac_{leachMS}$ = percent of managed manure nitrogen losses for livestock category T due to runoff and leaching during solid and liquid storage of manure.

$Frac_{small}$ = percent of livestock on small farms (<5 LU; or <3 LU for poultry);

$Frac_{NC}$ = percent of livestock on large farms (>5 LU; or >3 LU for poultry), not meeting the requirements of nitrate regulations.

In the case of Buffalo, Goat, Horse, Mule and Asses, Guinea Fowls and Rabbit, which livestock categories have small share to the total emissions, the whole livestock was assumed to be farmed on small farms ($Frac_{small}=100$).

Time series of percent of livestock on small farms ($Frac_{small}$) or large farms, which do not meet the requirements of nitrate regulation ($Frac_{NC}$), by livestock categories, are shown in Table 5.3.24.

Table 5.3.24 Fractions of livestock on small farms ($Frac_{Small}$) or large farms, which do not meet the requirements of nitrate regulation ($Frac_{NC}$) for 2000-2020

Year	Swine	Dairy Cattle	Non-Dairy Cattle	Sheep	Laying hens	Geese	Broiler	Turkey	Ducks
	Solid/Deep litter	Solid/Deep litter	Solid/Deep litter	Solid	Solid	Solid	Solid	Solid	Solid
2000	49	100	100	53	72	21	29	29	97
2001	47	95	94	51	71	20	28	27	92
2002	44	89	89	48	70	19	26	26	87
2003	42	83	82	46	69	18	24	24	81
2004	39	78	76	44	67	17	22	22	76
2005	36	72	70	43	65	16	20	21	71
2006	33	65	64	39	66	15	18	19	66
2007	30	59	58	35	68	15	17	17	61
2008	29	53	52	34	61	14	15	16	56
2009	27	47	47	32	55	13	13	14	51
2010	26	42	42	30	47	12	11	12	46
2011	24	35	36	27	46	10	10	11	40
2012	22	29	29	25	45	8	8	10	35
2013	20	23	23	22	43	7	6	8	29
2014	18	16	16	20	38	6	4	6	22
2015	16	9	10	17	32	4	2	4	15
2016	14	3	3	15	26	3	0	2	9
2017	12	2	2	15	23	3	0	2	9
2018	11	2	2	15	19	3	0	1	10
2019	9	2	2	15	16	3	0	1	10
2020	8	2	2	15	13	3	0	1	11

The 2006 IPCC Guidelines provide a range of 1-20% for the default value of $Frac_{LeachMS}$, but there is no animal and manure management specific default values available in the IPCC methodology. However, the difference of $Frac_{LossMS}$ and $Frac_{GasMS}$ given in Tables 10.23 and 10.22 seems to be as a proxy for $Frac_{LeachMS}$, but according to the footnote of the Table 10.23, this difference contains the leaching losses as well as the N_2 emissions, therefore cannot be used as default values in the estimation.

in the submissions before 2021, the default ($EF_{leachateN}=12.0$ as a proportion of TAN entering storage) provided for solid manure in the Table A1.12 of the 2016 EMEP/EEA Guidebook was applied for $Frac_{LeachMS}$, due to lack of country-specific measurements on the leaching losses from manure management systems and considering the low share of this emission. In line with the 2006 IPCC Guidelines, as well as the 2016 EMEP/EEA Guidebook N leaching from liquid/slurry was not assumed as a source of leachate. In the most up to date EMEP methodology, in the 2019 EMEP/EEA Guidebook N-leaching is no longer taken into account in the N-flow, thus the related EF was omitted from the Guidebook. Therefore, to calculate the indirect N_2O emissions from Manure Management due to leaching/runoff in this submission the methodology and the default value of $Frac_{LeachMS}$ provided in the 2019 Refinement was applied (2% for solid and 3.5% for deep litter).

In accordance with the Table 11.3 of 2006 IPCC Guidelines the default value of 0.0075 kg N₂O-N (kg N leaching and run off)⁻¹ for EF₅ was applied to estimate the indirect emissions from leaching/ run off.

5.3.3 Uncertainties and time series consistency

5.3.3.1 CH₄ emissions

Uncertainty of activity data (animal population) was estimated based on the confidence intervals for each animal species and livestock survey provided by the HCSO. The uncertainty of the mean annual averages was estimated according to the error propagation rules.

Uncertainty of EFs for CH₄ emissions from manure management was assumed to be ±20% for Cattle and ±30% for all livestock categories except rabbit, for which ±50% was applied. The 2006 IPCC Guidelines provide ±30% for T1 and ±20% for T2 methods, thus the estimated uncertainties are in line with the IPCC values. The Tier1 uncertainty analysis gives an overall uncertainty of ±14 % for the CH₄ emission from manure management.

5.3.3.2 Direct N₂O emissions

Uncertainties of ±25% are assumed relating to the N excretion of dairy cattle, non-dairy cattle, and swine, for which country-specific values are used. Uncertainties of ±30% are assumed for Poultry because country-specific N excretion rates are applied for Broiler and Laying hens, and ±50% for the other livestock categories in accordance with the 2006 IPCC Guidelines. The uncertainty of the manure management system usage (MS_{T,S}) data was assumed to be ±25% in accordance with the default value provided by 2006 IPCC Guidelines. The uncertainty of the EFs are -50%/+100% according to the 2006 IPCC Guidelines, therefore the lower combined uncertainty of the direct N₂O emissions from Manure management is 36% and the upper one is 67%.

5.3.3.3 Indirect N₂O emissions

Currently, Hungary does not have uncertainty assessment on the reported air pollutant emissions. However, uncertainties in emission factor (EF₄) are likely to dominate these emissions, thus uncertainties in the volatilized nitrogen are comparatively less important in terms of emissions. Consequently, due to lack of country-specific uncertainties the default uncertainty ranges for the Frac_{GasMS}, and default uncertainty of the emission factor (EF₄), taken from the 2006 IPCC Guidelines were applied. The lower combined uncertainty of the indirect N₂O emissions from Manure management is 84% and the upper one is 399%.

The overall combined uncertainty in the N₂O emissions from 3.B is -36%/+130%.

5.3.4 Source specific QA/QC

Following the recommendation of the 2021 UNFCCC review (A.7, 2021), the following information has been added to this chapter:

the CH₄ IEF for swine for 1985-2000 is lower than the lowest value in the IPCC default range due to the higher proportion of solid manure in Hungary than accounted for in the IPCC default value for Eastern

Europe. For 1985–2000, the proportion of solid manure ranged from 59.1 to 45.7 per cent, while tables 10A-7–10A-8 of the 2006 IPCC Guidelines assume a proportion of 42 per cent for Eastern Europe. Additionally, the IPCC default MMS usage assumes 3 per cent anaerobic lagoon with methane correction factors of 66 and 68 per cent, but anaerobic lagoons do not exist in Hungary. The extremely high methane correction factors for anaerobic lagoons significantly increase the IPCC default EF.

Nitrogen excretion rates for Cattle and Swine were verified using different calculation methodologies and compared with values used by other countries (EU Member States). For Dairy Cattle two different methodologies were used to verify the annual value of N excretion. Firstly, the N excretion was estimated based on the body mass and milk yields according to the methodology provided in Fébel and Gundel, 2007. This methodology indicates a value of 124 kg N/head/year for the year 2019 and 149 kg N/head/year for the year 2020. This methodology also justifies a significant increase in the N-excretion factor in the inventory between 2019 and 2020, due to a significant increase in milk production.

In case of Non-Dairy Cattle and Swine the methodology provided in Fébel and Gundel, 2007 indicates lower values. However, the difference for Swine is insignificant. Nitrogen excretion rates were compared with the values reported by other EU Member States. This verification revealed that the Hungarian values are in the range of values reported by EU Member States. The N_{ex} for non-dairy cattle was on average 52 kg N/head/year for the EU MS for the year 2019, which is very close to the Hungarian figure. The EU average of N-excretion rates for pigs was 10.8 kg N/head/year for the year 2019. The Hungarian value is slightly lower, but the Hungarian value is well within the range of EU values. The slightly lower value in Hungary is probably due to the lower fattening weights and the use of amino acid supplements.

Nitrogen excretion rate for Laying hens was compared with the default values provided in the 2006 IPCC Guidelines, 2019 Refinement and the 2019 EMEP/EEA Guidebook (EEA, 2019). Default IPCC values are given in kg N (1000 kg animal mass)⁻¹ day⁻¹, hence the 'typical animal mass' (TAM) is required to determine the annual N excretion. The table below summarizes the default values and the resulted annual N excretion.

Table 5.3.25 Comparison of N excretion rates for laying hens between the IPCC Guidelines and the EMEP/EEA Guidebook

	2006 IPCC Guidelines		2019 Refinement		EMEP/EEA Guidebook (EEA, 2019)
	Eastern Europe	Western Europe	Eastern Europe	Western Europe	
Daily N excretion rate (kgN (1000 kg animal mass)⁻¹ day⁻¹)	0.82	0.96	0.81	0.87	0.96*
TAM (kg)	1.8	1.8	1.9	1.9	2.2
Annual N excretion (kg N head⁻¹ yr⁻¹)	0.54	0.63	0.56	0.6	0.77

*Calculated from the values provided in Table 3.9 and Table A1.5 of the 2019 EMEP/EEA Guidebook.

The value of 0.76 kg N head⁻¹ yr⁻¹ used in the Hungarian inventory for the year 2018 is higher than the IPCC default values. The higher, 2.0 kg weight of laying hens in the Hungarian inventory justifies the higher annual N excretion rates. The 2019 EMEP/EEA Guidebooks also uses higher default weight than the IPCC Guidelines, resulting in similarly higher annual N excretion rate. The Table above reveals that the value of 0.76 kg N head⁻¹ yr⁻¹ used in the Hungarian inventory is in the range of the default values provided in the different emission inventory guidelines.

Nitrogen excretion rates for broiler was also compared with the default values provided in the different emission inventory guidebooks. This comparison reveals that the most up-to-date guidebooks suggests higher values than the 2006 IPCC Guidelines due to the increase in the weight and the recently published defaults ranged from 0.4 to 0.5 kg N head⁻¹·yr⁻¹. However, the default daily N excretion rates combined with the country-specific TAM result in values ranging from 0.60 to 0.62 kg N head⁻¹·yr⁻¹, thus the value of 0.56 N head⁻¹·yr⁻¹ for the year 2018 in the Hungarian inventory can be considered as consistent with the default values provided in the recently published emission inventory guidebooks.

Table 5.3.26 Comparison of N excretion rates for broilers between the IPCC Guidelines and the EMEP/EEA Guidebook

	2006 IPCC Guidelines		2019 Refinement		EMEP/EEA Guidebook (EEA, 2019)*
	Eastern Europe	Western Europe	Eastern Europe	Western Europe	
Daily N excretion rate (kgN (1000 kg animal mass)⁻¹ day⁻¹)	1.1	1.1	1.12	1.14	1.1
TAM (kg)	0.9	0.9	1.1	1.2	1
Annual N excretion (kg N head⁻¹ yr⁻¹)	0.36	0.36	0.45	0.50	0.40

*Values are taken from the Manure Management N-flow tool.xlsx provided to the 2019 EMEP/EEA Guidebook (EEA, 2019)

N₂O emissions are calculated and reported consistently with the NH₃ and NO_x inventory under the UNECE/LRTAP convention. To calculate the NH₃ and NO_x emissions the 2019 EMEP/EEA Guidebook was applied.

5.3.5 Source-specific recalculations

The main reasons for changes in the emissions from 3. Agriculture sector for this submission are the manure management data updates. The 2020 Agricultural Census and the recently available annual statistics on agricultural wastes used in biogas plants have made it possible to reflect the current practices and the effects of anaerobic digestion. As a result of this inventory improvements animal manure used in “digesters” has been reported at the first time in this submission. By reporting manure treated in biogas plants, Hungary is responding to several UNFCCC review recommendations over the past three years.

In addition, the revisions made in Sector 3.A, through gross energy intake for dairy cows and animal numbers for other cattle, also affected emissions from manure management, and these changes will not be repeated in the following chapters.

The recalculations within category 3.B by gas and subcategory are as follows:

3.B.1 CH₄ Emissions from Manure Management

3.B.1.1 and 3.B.1.3 CH₄ Emissions from Manure Management - Cattle and Swine, across the whole time series

Proportions of pig and cattle liquid/slurry covered by “natural crust” was revised for the whole time series. The former expert judgement was replaced by annual survey data, which are available from the year 2015. Between 2001 and 2015, interpolation was used for cover types other than natural crust, and for natural crust, 2015 data were used for the period before 2015, as well. This recalculation resulted in an increase in the MCF of liquid/slurry of cattle and swine. The increase is larger for cattle and minor for pigs (*Table 5.3.27*).

3.B.1.1, 3.B.1.3 and 3.B.1.4 CH₄ / 3.B.2.1, 3.B.2.3 and 3.B.2.4 N₂O Emissions from Manure Management - Cattle, Swine and Poultry for the period 2004-2019

CH₄ and N₂O emissions from manure management of cattle, swine, and poultry for the period 2004-2019 was revised considering proportions of manure used in anaerobic digesters. In the previous submissions the anaerobic digested manure was allocated in line with the housing, irrespective of the manure use. In this submission, the anaerobic digested manure was reallocated to ‘digesters’, as data on the agricultural wastes used in anaerobic digesters had become available.

3.B.1.2 CH₄ Emissions from Manure Management - Sheep, 2014-2019

There are minor recalculations to CH₄ emissions from manure management of sheep due to the revision on the proportion of manure excreted during grazing considering the length of grazing period. This revision resulted in an insignificant (less than 0.5 kt CO₂-eq) increase in the CH₄ emissions from 3.B.1.2.

The overall changes in the CH₄ emissions from 3.B.1. Manure Management (*Table 5.3.28*) ranged from a decrease of 25.5 kt CO₂-eq (4.1%) in 2013 to an increase of 15.7 kt CO₂-eq (1.3%) in the BY. Revision of “natural crust” resulted in an increase in the BY emissions, 3.0, 10.9 and 1.7 kt CO₂-eq for dairy cattle, non-dairy cattle, and swine, respectively. In contrast, reallocation of anaerobic digested manure to “digesters” resulted in a decrease in the emissions for the period 2004 to 2019, partly offset by the increase in emissions due to revised natural crust. The highest net decrease in the emissions due to anaerobic digestion were 9.4 kt CO₂-eq (in 2013), 5.3 kt CO₂-eq (in 2019) and 17.1 kt CO₂-eq (in 2014), for dairy-cattle, non-dairy cattle, and swine, respectively.

Table 5.3.27 Changes in the emissions between the 2021 and 2022 submissions, by sources due to recalculations to CH₄ emissions from 3.B Manure Management (kt CO₂-eq)

Year	Dairy Cattle	Non-Dairy Cattle	Swine	Poultry	Sheep
BY	3.0	10.9	1.7	0.0	0.0
1990	2.9	10.1	1.7	0.0	0.0
1991	2.6	9.8	1.5	0.0	0.0
1992	2.3	8.5	1.2	0.0	0.0
1993	2.1	6.6	1.1	0.0	0.0
1994	1.9	5.8	1.0	0.0	0.0
1995	1.8	5.7	1.0	0.0	0.0
1996	1.8	5.5	1.1	0.0	0.0
1997	1.8	5.3	1.0	0.0	0.0
1998	1.8	5.2	1.1	0.0	0.0
1999	1.8	5.2	1.2	0.0	0.0
2000	1.7	5.2	1.1	0.0	0.0
2001	1.9	5.0	1.1	0.0	0.0
2002	2.1	4.9	0.7	0.0	0.0
2003	2.2	4.8	0.1	0.0	0.0
2004	0.3	4.5	-1.1	0.0	0.0
2005	2.0	4.4	-1.5	0.0	0.0
2006	0.5	4.3	-2.1	0.0	0.0
2007	0.3	4.0	-3.4	0.0	0.0
2008	-1.2	3.5	-5.2	0.0	0.0
2009	-2.9	3.3	-6.9	0.0	0.0
2010	-2.5	3.0	-8.3	0.0	0.0
2011	-5.8	2.0	-11.5	0.0	0.0
2012	-3.7	2.5	-10.3	0.0	0.0
2013	-9.4	0.5	-16.6	0.0	0.0
2014	-9.0	0.9	-17.1	0.0	0.1
2015	-8.5	1.4	-16.9	0.1	0.2
2016	-7.6	1.5	-16.8	0.1	0.4
2017	-8.2	-1.4	-15.7	0.0	0.2
2018	-5.5	-3.3	-11.8	-0.1	0.1
2019	-3.0	-5.3	-9.6	-0.2	0.0

Table 5.3.28 The overall impact of recalculations on CH₄ emissions from 3.B Manure Management

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ -eq]	Percentage change
BY	1,243	1,259	15.7	1.3%
1990	1,161	1,175	14.6	1.3%
1991	1,064	1,078	13.9	1.3%
1992	894	906	12.0	1.3%
1993	797	807	9.8	1.2%
1994	709	718	8.6	1.2%
1995	700	709	8.5	1.2%
1996	740	748	8.4	1.1%
1997	718	727	8.1	1.1%

Year	Submission 2021 [Gg CO ₂ - eq]	Submission 2022 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
1998	743	751	8.1	1.1%
1999	783	791	8.2	1.0%
2000	758	766	8.1	1.1%
2001	742	750	8.1	1.1%
2002	775	783	7.7	1.0%
2003	788	795	7.1	0.9%
2004	726	729	3.6	0.5%
2005	690	695	4.9	0.7%
2006	677	679	2.6	0.4%
2007	688	689	0.9	0.1%
2008	655	652	-2.9	-0.4%
2009	614	607	-6.6	-1.1%
2010	608	601	-7.9	-1.3%
2011	607	592	-15.3	-2.5%
2012	610	599	-11.5	-1.9%
2013	615	590	-25.5	-4.1%
2014	638	613	-25.0	-3.9%
2015	658	634	-23.7	-3.6%
2016	656	633	-22.3	-3.4%
2017	644	619	-25.0	-3.9%
2018	654	633	-20.7	-3.2%
2019	650	632	-18.2	-2.8%

3.B.2 N₂O emissions from Manure Management

3.B.2 Direct N₂O emissions from manure management, whole time series

Revision of “natural crust” for the whole time series and the anaerobic digested manure for the period 2004-2019 resulted in an overall decrease of 1.3% and 4.1 kt CO₂-eq on average on the BY and 1990-2019 trend. Decrease in the emissions ranged from 1.1 kt CO₂-eq in 2000 to 18.3 kt CO₂-eq in 2019 (*Table 5.3.29*).

Table 5.3.29 Impact of recalculations on 3.B.2 Direct N₂O Emissions from Manure Management

Year	Submission 2021 [Gg CO ₂ - eq]	Submission 2022 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
BY	621	619	-2.0	-0.3%
1990	575	573	-1.9	-0.3%
1991	537	535	-1.8	-0.3%
1992	462	461	-1.5	-0.3%
1993	402	401	-1.3	-0.3%
1994	360	358	-1.1	-0.3%
1995	358	357	-1.1	-0.3%
1996	353	351	-1.1	-0.3%
1997	342	341	-1.1	-0.3%
1998	349	348	-1.1	-0.3%
1999	355	354	-1.1	-0.3%
2000	354	353	-1.1	-0.3%
2001	343	342	-1.2	-0.3%

Year	Submission 2021 [Gg CO ₂ - eq]	Submission 2022 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
2002	346	345	-1.3	-0.4%
2003	345	344	-1.3	-0.4%
2004	334	332	-1.7	-0.5%
2005	321	319	-1.8	-0.6%
2006	314	312	-1.9	-0.6%
2007	314	311	-2.4	-0.8%
2008	304	301	-3.3	-1.1%
2009	292	288	-4.1	-1.4%
2010	290	285	-4.7	-1.6%
2011	288	282	-6.3	-2.2%
2012	301	295	-5.7	-1.9%
2013	307	298	-9.2	-3.0%
2014	316	308	-8.3	-2.6%
2015	322	315	-7.4	-2.3%
2016	318	311	-6.6	-2.1%
2017	316	304	-11.6	-3.7%
2018	316	301	-14.6	-4.6%
2019	321	303	-18.3	-5.7%

3.B.2.5 Indirect N₂O Emissions from Manure Management, *whole time series*

Atmospheric Deposition, whole time series

Revision of NH₃ emissions from 3.B Manure Management for the reporting to the UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP) (UNECE, 1999) and the National Emissions Ceiling Directive (EP, CEU, 2016) resulted in changes in the indirect N₂O emissions from manure management due to atmospheric deposition. The revision of NH₃ emissions aimed to take account of emission abatement measures, as a result of new data collections recently launched.

The following abatement measures were considered in the revision of NH₃ emissions by stages of manure and animals:

Animal housing

- Non-leaking drinking systems for broilers
- Aviary system for layers
- Partially slatted floor for piglets after weaning and growers-finishers

Manure storage

- "Low technology" floating covers (e.g., chopped straw, peat, bark, etc.)
- "Tight lid", roof or tent structure
- Plastic sheeting (floating cover)
- Natural crust

Revision of the NH₃ emissions from manure management lead to an 1.1% (1.5 kt CO₂-eq) decrease in indirect N₂O emissions due to atmospheric deposition on average over the time series.

Nitrogen Leaching and Run-off, 2004-2019

Revision of data on manure management system distribution, mainly reallocation of manure to digesters, resulted in additional changes in the indirect N₂O emissions due to leaching and run-off. Changes ranged from a decrease of 0.08 kt CO₂ eq in 2019 (6.4%) to an increase of 0.02 kt CO₂ eq in 2016 (1.3%).

The overall indirect N₂O emissions from manure management decreased by 1.1% and 1.5 kt CO₂ eq on average in the BY and 1990-2019 trend. Changes in the emissions are shown in *Table 5.3.30*.

Table 5.3.30 The overall impact of recalculations on N₂O emissions from 3.B.2.5 Indirect N₂O emissions from Manure Management

Year	Submission 2020 [Gg CO ₂ -eq]	Submission 2021 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
BY	308	308	-0.4	-0.1%
1990	279	279	-0.3	-0.1%
1991	253	253	-0.1	0.0%
1992	216	216	0.2	0.1%
1993	187	187	0.1	0.1%
1994	172	172	0.3	0.2%
1995	173	174	0.4	0.2%
1996	170	170	0.3	0.2%
1997	165	166	0.4	0.2%
1998	171	171	0.3	0.2%
1999	173	174	0.2	0.1%
2000	180	180	0.3	0.2%
2001	175	175	0.2	0.1%
2002	177	177	-0.1	-0.1%
2003	178	178	-0.5	-0.3%
2004	170	169	-0.7	-0.4%
2005	160	159	-0.8	-0.5%
2006	155	154	-1.0	-0.7%
2007	154	152	-1.4	-0.9%
2008	150	148	-1.8	-1.2%
2009	143	141	-2.0	-1.4%
2010	144	142	-2.3	-1.6%
2011	141	138	-2.9	-2.1%
2012	138	136	-2.7	-1.9%
2013	136	132	-3.6	-2.6%
2014	140	136	-3.9	-2.8%
2015	144	139	-4.3	-3.0%
2016	144	139	-4.9	-3.4%
2017	140	135	-5.2	-3.7%
2018	141	135	-5.2	-3.7%
2019	142	136	-5.3	-3.7%

The overall impact of revisions for the 3.B Manure Management sector CH₄ and N₂O emissions is an increase in emissions in the range of 1.2-13.3 kt CO₂-eq (0.1%-0.7%) between BY and 2005, and a decrease of between 0.3 and 41.8 kt CO₂-eq (0.0%-3.8%) for the years 2006-2019 (*Table 5.3.31*).

Table 5.3.31 The overall impact of recalculations on CH₄ and N₂O emissions from 3.B Manure Management

Year	Submission 2021 [Gg CO ₂ - eq]	Submission 2022 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
BY	2,172	2,186	13.3	0.6%
1990	2,015	2,027	12.4	0.6%
1991	1,854	1,866	12.0	0.6%
1992	1,572	1,583	10.6	0.7%
1993	1,386	1,395	8.6	0.6%
1994	1,241	1,248	7.8	0.6%
1995	1,232	1,240	7.8	0.6%
1996	1,262	1,270	7.6	0.6%
1997	1,226	1,233	7.4	0.6%
1998	1,262	1,270	7.3	0.6%
1999	1,312	1,319	7.3	0.6%
2000	1,292	1,299	7.3	0.6%
2001	1,260	1,267	7.1	0.6%
2002	1,298	1,304	6.4	0.5%
2003	1,311	1,316	5.3	0.4%
2004	1,229	1,230	1.2	0.1%
2005	1,171	1,173	2.3	0.2%
2006	1,146	1,145	-0.3	0.0%
2007	1,155	1,152	-2.9	-0.2%
2008	1,109	1,101	-7.9	-0.7%
2009	1,049	1,036	-12.7	-1.2%
2010	1,042	1,028	-14.8	-1.4%
2011	1,036	1,012	-24.5	-2.4%
2012	1,049	1,030	-19.9	-1.9%
2013	1,058	1,020	-38.2	-3.6%
2014	1,094	1,057	-37.2	-3.4%
2015	1,124	1,088	-35.3	-3.1%
2016	1,117	1,084	-33.8	-3.0%
2017	1,100	1,059	-41.8	-3.8%
2018	1,110	1,069	-40.5	-3.6%
2019	1,113	1,071	-41.8	-3.8%

5.3.6 Planned improvements

Depending on the data availability further revision of feeding data for swine (other sub-categories than breeding animals), cattle, poultry and revision of inventory parameters according to the feeding monitoring program of AKI is planned.

5.4 Rice cultivation (CRF sector 3.C)

5.4.1 Source Category Description

Emitted gas: CH₄

Methods: T1

Emission factors: D

Key source: none

Hungary is situated on the north edge of the rice production area. According to this the climatic conditions are unfavorable. The production area of rice involves the poorer quality soils. Since the production volume is very low in Hungary, the contribution of rice cultivation to the greenhouse gas emissions is minimal, only 0.8% of the entire CH₄ emissions from agriculture sector.

5.4.2 Methodological issues

In Hungary the rice is cultivated on poorer quality soil, without organic amendments, the fields are intermittently flooded. Aeration is applied as a pest control during the cultivation. (Apáti, 2003).

Methane emissions from rice cultivation were calculated according to the Equation 5.1 of 2006 IPCC Guidelines. As CH₄ emissions from rice cultivation are not a key category in Hungary, the Tier 1 methodology with default emission factors was applied. The adjusted daily emission factor to the above equation was calculated based on Equation 5.2 of 2006 IPCC Guidelines. The required values of baseline emission factor (EF_c), water regime (SF_w), water regime in the pre-season (SF_p) to this equation were taken from Tables 5.11-5.13 of 2006 IPCC Guidelines. The adjusted CH₄ emission scaling factor for organic amendment (SF_o) was calculated using the Eq. 5.3 of 2006 IPCC Guidelines. The value of conversion factor (CFOA) was taken from Table 5.14 of 2006 IPCC Guidelines. Due to lack of detailed information 'straw incorporated shortly (<30 days) before cultivation' was assumed as a conservative approach.

In response to a recommendation from the in-country review conducted in 2016 this section was supplemented with the *Table 5.4.1* to present values of parameters used for calculating the EFs for irrigated rice cultivation.

Table 5.4.1 Parameters used for calculating the Emission Factors for CRF category 3.C

Parameters	Value	Unit	Source	Notes
EF_{i,j,k}, daily emission factor	1.86	kg CH ₄ ha ⁻¹ d ⁻¹	Eq. 5.2 of 2006 Gl.	
t_{i,j,k}, cultivation period, day	145	day	Calculated	Sowing: third ten days of April, or first ten days of May. Harvesting: between 10th of September and 10th of October
EF_c, baseline emission factor	1.3	kg CH ₄ ha ⁻¹ d ⁻¹	Table 5.11 of 2006 Gl.	
SF_w, water regime	0.78		Table 5.12 of 2006 Gl.	
SF_p, water regime, pre-season	1.22		Table 5.13 of 2006 Gl.	
SF_o, organic amendment	1.51		Eq. 5.3 and Table 5.14 for CFOA of 2006 Gl.	
SF_{s,r}, soil type, rice cultivar	1			There is no default value provided in the 2006 IPCC Gl. However, it should take into account if it is available according to the IPCC methodology.

As activity data, the total size of the production area was taken from the HCSO's statistics.

5.4.3 Uncertainties and time series consistency

For the uncertainty of the activity data, $\pm 5\%$ has been estimated by expert judgement. Uncertainties of scaling factors and the baseline emission factor were taken from the 2006 IPCC Guidelines. (SF_w $\pm 26\%$; SF_o $-4\%/+5\%$; SF_p $-14\%/+15\%$, EF_c $-63\%/+69\%$) Combination of uncertainties listed above resulted in $69\%/+75\%$ combined uncertainty for the adjusted daily emission factor. Therefore, the overall lower and upper uncertainty of 69% and 76% can be calculated for the emissions from rice cultivation.

5.4.4 QA/QC Information

See 6.1.5.

5.4.5 Source-specific recalculations

There were no recalculations in this category.

5.4.6 Planned improvements

There are no further improvements planned.

5.5 N₂O emissions from Agricultural soils (CRF sector 3.D)

5.5.1 Source Category Description

Emitted gas: N₂O

Methods: T1

Emission factors: D

Key source: Yes

Particularly significant sub-categories: Inorganic N fertilizers, Crop residue

In 2020 agricultural soils emitted 90% of the total N₂O emissions from the agriculture sector, and 77% of the national total N₂O emissions are generated in agricultural soils (Table 5.5.1). Emissions from agricultural soils contributed 6.2 percent (3,868 Gg CO₂-eq) to the national total GHG emissions in 2020).

The overall trend in emissions is decreasing. However, trends in emissions from crop production related sectors as 3.D.a.4 and 3.D.a.5 are different from the other subsectors. Emissions from 3.D.a.4 fluctuated significantly depending on crop production, which is determined by the weather conditions. Trends for 3.D.a.5 also seem to be fluctuating. However, emission from this source is low and uncertain.

Table 5.5.1 Trends in emissions from 3. D Agricultural Soils by subcategories

Year	N ₂ O emissions (Gg N ₂ O)									
	3.D.a	3.D.a.1	3.D.a.2	4.D.a.3	4.D.a.4	3.D.a.5	3.D.a.6	3.D.b	3.D.b.1	3.D.b.2
BY	15.75	9.25	3.24	0.76	2.50	0.010	NO	1.51	1.06	0.46
1990	11.65	5.63	2.97	0.65	2.39	0.012	NO	1.21	0.85	0.37
1991	8.50	2.20	2.77	0.63	2.88	0.012	NO	0.86	0.61	0.25
1992	7.24	2.33	2.38	0.56	1.97	0.012	NO	0.74	0.52	0.22
1993	6.78	2.53	2.09	0.45	1.70	0.012	NO	0.68	0.47	0.21
1994	8.01	3.49	1.88	0.38	2.25	0.012	NO	0.71	0.47	0.24
1995	7.43	3.00	1.86	0.36	2.19	0.012	NO	0.70	0.48	0.22
1996	7.57	3.19	1.81	0.35	2.20	0.012	NO	0.70	0.48	0.22
1997	7.84	3.24	1.75	0.35	2.50	0.012	NO	0.69	0.48	0.21
1998	8.41	3.90	1.78	0.35	2.37	0.008	NO	0.74	0.50	0.23
1999	8.40	4.12	1.81	0.35	2.12	0.008	NO	0.74	0.52	0.22
2000	8.06	4.05	1.82	0.38	1.80	0.008	NO	0.77	0.52	0.24
2001	9.08	4.32	1.79	0.37	2.60	0.008	NO	0.79	0.53	0.26
2002	9.08	4.76	1.82	0.36	2.14	0.008	NO	0.82	0.55	0.27
2003	8.49	4.54	1.83	0.37	1.75	0.008	NO	0.81	0.55	0.26
2004	9.76	4.60	1.77	0.38	3.00	0.008	NO	0.83	0.54	0.29
2005	9.03	4.09	1.69	0.38	2.87	0.000	NO	0.75	0.51	0.25
2006	9.21	4.54	1.67	0.37	2.63	NO	NO	0.77	0.52	0.25
2007	9.03	5.03	1.66	0.36	1.98	NO	NO	0.79	0.52	0.27
2008	9.67	4.62	1.62	0.35	3.07	NO	NO	0.72	0.45	0.27
2009	8.78	4.32	1.56	0.35	2.54	NO	NO	0.67	0.42	0.25

Year	N ₂ O emissions (Gg N ₂ O)									
	3.D.a	3.D.a.1	3.D.a.2	4.D.a.3	4.D.a.4	3.D.a.5	3.D.a.6	3.D.b	3.D.b.1	3.D.b.2
2010	8.57	4.42	1.57	0.35	2.23	NO	NO	0.67	0.42	0.25
2011	9.23	4.74	1.57	0.35	2.56	0.006	NO	0.69	0.43	0.26
2012	9.01	4.92	1.60	0.36	2.11	0.015	NO	0.69	0.42	0.27
2013	10.02	5.39	1.65	0.37	2.61	0.015	NO	0.75	0.46	0.29
2014	10.52	5.36	1.70	0.38	3.06	0.014	NO	0.78	0.46	0.31
2015	10.80	5.95	1.73	0.39	2.72	0.014	NO	0.82	0.50	0.32
2016	11.68	6.35	1.70	0.40	3.20	0.014	NO	0.85	0.52	0.34
2017	11.67	6.67	1.67	0.43	2.89	0.014	NO	0.88	0.54	0.34
2018	11.71	6.67	1.65	0.44	2.94	0.014	NO	0.87	0.54	0.34
2019	11.66	6.54	1.65	0.46	2.99	0.014	NO	0.87	0.54	0.34
2020	12.07	6.97	1.65	0.48	2.97	0.014	NO	0.91	0.56	0.35
Share of Hungarian total N ₂ O emissions in BY	42%	25%	9%	2%	7%	0.03%	NO	4%	3%	1%
Share of Hungarian total N ₂ O, in 2020	72%	41%	10%	3%	18%	0.1%		5%	3%	2%
Trend BY-2020	-23%	-25%	-49%	-37%	19%	37%		-40%	-47%	-24%

The total emissions from 3.D Agricultural soils have reduced by 25 per cent of the BY levels until 2020. A significant drop had occurred in the period 1985-1993 due to the significant decrease in synthetic fertilizer use and livestock population which resulted in less N-input. After reaching the lowest point of the emission levels in 1993 there was a slight increase until 1998 due to a small rise in synthetic fertilizer use. After that emission levels remained quasi stable in the period 1998-2013 as a result of compensatory processes between the different sources of N input. As the Figure 5.5.1 reveals emissions are primarily driven by the amount in synthetic fertilizer used. At the beginning of the time series the second most important source was the organic manure. In contrast, in the recent years N in crop residues has exceeded the amount of organic N, reflecting the restructuring in the Hungarian agricultural. Namely, the animal husbandry has declined in importance in the agricultural production, while crop production has become more meaningful. For more details on trends see also Section 5.1.1.




Figure 5.5.1 Trends in nitrogen input to soils

5.5.2 Methodological issues

5.5.2.1 Direct soil (CRF sector 3.D.a)

Direct soil emissions are the main source of N_2O in the Hungarian inventory. In 2020, 72% of the national total N_2O emissions originated from this sector, which includes N inputs from synthetic N-fertilizer (3.D.a.1), organic manures as animal manure use and sewage sludge application (3.D.a.2), emissions from urine and dung N deposited on pasture (3.D.a.3) and crop residues (3.D.a.4). Emissions from N mineralization associated with loss of SOM (3.D.a.5) are rather low, but also was taken into account to meet the principle of completeness. In response to the recommendation of the ESD review, 2016, N_2O emissions from 3.D.2.c Other organic fertilizers applied to soils (compost) have been reported since the 2018 submission. Organic soils are protected, thus not cultivated in Hungary. Therefore, emissions from cultivation of histosols are not reported.

Emissions from these sources were calculated using Tier 1 methodology based on the Equation 11.1 of 2006 IPCC Guidelines. The N_2O -N was converted to N_2O by the factor (44/28) in line with the IPCC methodology. Terms of the Equation 11.1 were determined as it is detailed in the following sub-sections, while amounts of various N inputs to soils are provided in Table 5.5.7.

5.5.2.1.1 N input from synthetic fertilizer use (F_{SN})

Annual amount of synthetic N fertilizer applied to soils was estimated from the total amount of synthetic fertilizer sold to end users, annually. Collection of this data is executed according to the National Statistical Data Collection Program (OSAP) by the AKI. Although, this is sale statistics instead of consumption data, but so comprehensive survey on fertilizer consumption there is not available in Hungary. Moreover, this sale statistics contain the sold fertilizers by product line, which enables us to determine the amount of Nitrogen applied to soils by fertilizer types, thus the detailed and more

accurate calculation of volatilization and indirect emissions. Data on synthetic fertilizer applied (F_{SN}) for the period 1985-2020 are provided in Table 5.5.7.

5.5.2.1.2 Applied organic fertilizers (F_{ON})

The amount of organic N inputs applied to soils other than by grazing animals was calculated using Equation 11.3 of 2006 IPCC Guidelines. In Hungary, this includes animal manure, sewage sludge and composted sewage sludge, composted municipal solid waste and digestate from anaerobic digesters.

Animal manure applied to soils (F_{AM})

Annual amount of animal manure N applied to soils (F_{AM}) was calculated using the equation 11.4 of the 2006 IPCC Guidelines. In Hungary manure is not used as feed, fuel or for construction, therefore fractions ($Frac_{FEED}$, $Frac_{FUEL}$, $Frac_{CNST}$) in the equation were assumed to be zero.

Consequently, annual amount of animal manure N applied to soils (F_{AM}) is corresponded to the managed manure nitrogen available for application to managed soils (N_{MMS_Avb}), which was calculated based on Eq. 10.34 of 2006 IPCC Guidelines. The first term of this equation is the managed manure N taking into account the losses in the manure management systems. Data and information on the calculation of managed manure N, volatilization, leaching and N_2O -N losses from manure management systems are provided in Chapter 5.3.

In the equation 10.34 reference is also made to $Frac_{LOSSMS}$, for which default values are provided in Table 10.23. According to the footnote b of this table N_2 emissions are also taken into account in the N losses from the manure management systems. Thus, the losses in equation 10.34 covers the N_2 emissions for most manure management systems.

Hungary uses Tier 2 and country-specific values to calculate the volatilization and leaching losses from the manure management systems, therefore the default values provided in Table 10.22 and 10.23 are not used in the inventory. Nevertheless, accepting the concept that the default total N losses from manure management systems covers the N_2 emissions, the N_2 emissions were subtracted to get the N content of animal manure applied to soils.

As the IPCC methodology does not provide emission factors to calculate N_2 emissions from the storage of manure the default EFs given in the Table 3.10 of the 2019 EMEP/EEA Guidebook was applied. ($EF_{solid}=0.3$ kg N_2 / kg TAN and 0.003 kg N_2 / kg TAN.) N_2 emissions are calculated in the N-flow tool used to calculate NH_3 emissions and published with the 2019 EMEP/EEA Guidebook (EEA, 2019).

The time series of N_2 emissions are shown in Table 5.5.2.

Table 5.5.2 N_2 emissions calculated to estimate F_{AM} , for the BY and the period 1990-2020

Year	N_2 emissions [kg] from manure management systems
BY	29,114,449
1990	26,199,192
1991	23,932,415
1992	20,594,325
1993	17,642,028
1994	16,212,676
1995	16,365,051
1996	15,852,307
1997	15,561,855
1998	15,952,601
1999	15,810,298
2000	16,524,508
2001	16,219,023
2002	16,100,270
2003	16,053,955
2004	15,644,376
2005	14,936,710
2006	14,452,473
2007	14,186,008
2008	14,096,113
2009	13,571,887
2010	13,601,806
2011	13,325,715
2012	13,089,437
2013	12,962,070
2014	13,230,922
2015	13,505,837
2016	13,429,119
2017	13,143,738
2018	12,823,578
2019	12,864,535
2020	12,391,018

The second term of the Equation 10.34 of the 2006 IPCC Guidelines is the N input from bedding materials. Straw N amounts depend on livestock population and the housing systems. The 2006 IPCC Guidelines provide default values on the nitrogen contained in organic bedding materials for Cattle and Swine, which were used in our calculation. For other livestock categories default values taken from the Table 3.7 of the 2019 EMEP/EEA Guidebook were applied. Data used in the calculations to estimate N from bedding materials with their sources is provided in Table 5.5.3.

Table 5.5.3 Nitrogen in bedding materials by animal category and manure management systems

Animal category	N content of bedding materials by manure management systems [kg N head ⁻¹ yr ⁻¹]		Source
	Solid	Deep Litter	
Dairy Cattle	7	13	p.10.66 of 2006 IPCC GLs.
Non-Dairy Cattle	4	8	p.10.66 of 2006 IPCC GLs.
Buffalo	6	-	2019 EMEP/EEA GB
Sheep	0.08	-	2019 EMEP/EEA GB
Goats	0.08	-	2019 EMEP/EEA GB
Horses	2	-	2019 EMEP/EEA GB
Mules	2	-	2019 EMEP/EEA GB
Swine	0.9	1.8	p.10.66 of 2006 IPCC GLs
Poultry	0.022*	-	Expert judgments

*Poultry manure with bedding

The time series of the resulted N inputs from bedding are shown in Table 5.5.5.

Sewage N (F_{SEW})

Data on annual amount of total sewage N that is applied to agricultural soils is available in the Urban Wastewater Information System (UWIS) since 2011. For the period 1994-2010 data were taken from the EUROSTAT statistics. The EUROSTAT provides data on sewage sludge disposal for agricultural use, but this statistic contains the sewage sludge disposal for recultivation as well as agricultural purposes. 40% of the reported disposed sewage sludge based on expert judgment was assumed to be applied on agricultural lands and the remaining 60% for recultivation. Activity data was extrapolated for the period 1988-1994. For the years 1988 backwards application of sewage was assumed to be 'not occurring', because of the low proportion of wastewater treatment in Hungary. The N-content of sewage sludge was assumed to be 4.2% in the calculation. Following a recommendation from the centralized review 2019, country-specific value on the N-content of the sewage sludge based on the measured data provided by the NFCSO were used in the estimate. Data on applied organic fertilizers (F_{ON}) was determined in coordinated with the Waste sector.

Other organic waste including compost N (F_{COMP})

In this category, composted sewage sludge, composted municipal solid waste (MSW) and digestate from anaerobic digesters are reported.

Composted sewage sludge

For the calculation of emissions from the application of sewage sludge compost, the amount of composted sewage sludge reported as activity data in the NFR sector 5B1 is used. The Wastewater Sludge Processing and Use Strategy 2014-2023 (General Directorate of Water Management, 2013) shows that 38% of composted sewage sludge is used in agriculture. The N content of the sewage sludge compost was assumed to be 2% based on the Table 4.1 of the Vol. 5. Ch. 4 of the 2006 IPCC Guidelines.

Composted MSW

Activity data was taken from the NFR sector 5B1. The IPCC default parameters on moisture content and N in dry matter given in Table 4.1 of the Vol. 5. Ch. 4 of the 2006 IPCC GIs was used. According to the NHKV (National Coordination of Waste Management and Asset Management Plc.) reports for the last years, loss during composting approximately 25% and 50% of the sewage sludge compost generated is used for agricultural purposes (NHKV, 2020).

Digestate (other than animal manure and crop residues)

As a result of inventory improvements for this year, anaerobic digestion has been included in the agricultural emission inventories, and in parallel, emissions from the application of digestate are also taken into account under CRF sector 3Da2c. The N content of the biogas compost applied is calculated on the basis of the N content of feedstock. As biogas feedstock statistics are only available from 2017 onwards, the N consumption per TJ energy production was determined for the previous years using the feedstock consumption in the period 2017-2020. N per TJ are estimated to 4.16 kg N per TJ based on average of N in feedstock and energy production in 2017-2020.

The resulted activity data for the period 1985-2020 are provided in *Table 5.5.4*

Table 5.5.4 Nitrogen applied to soils as organic waste including compost

Year	Composted sewage sludge N	Composted MSW N	Digestate (other than animal manure and crop residues) N	Total N applied
kg N				
BY	152,000	-	-	152,000
1990	152,000	-	-	152,000
1991	152,000	-	-	152,000
1992	152,000	-	-	152,000
1993	152,000	-	-	152,000
1994	152,000	-	-	152,000
1995	212,800	-	-	212,800
1996	220,400	54,000	-	274,400
1997	197,600	57,000	-	254,600
1998	174,800	54,000	-	228,800
1999	243,200	54,000	-	297,200
2000	228,000	51,000	-	279,000
2001	205,200	51,000	-	256,200
2002	281,200	141,000	-	422,200
2003	425,600	141,000	-	566,600
2004	182,058	117,000	212,069	511,127
2005	399,932	123,000	202,159	725,092
2006	326,414	174,000	255,672	756,086
2007	388,672	192,000	495,489	1,076,161
2008	469,573	255,000	971,158	1,695,731
2009	683,795	270,000	1,454,755	2,408,550
2010	624,935	444,000	1,779,796	2,848,731
2011	618,408	549,000	2,647,893	3,815,301
2012	685,342	548,619	2,190,061	3,424,022
2013	708,761	561,991	4,047,153	5,317,905

Year	Composted sewage sludge N	Composted MSW N	Digestate (other than animal manure and crop residues) N	Total N applied
			kg N	
2014	739,125	708,262	3,995,623	5,443,009
2015	754,845	691,786	3,872,741	5,319,372
2016	773,726	881,907	3,894,543	5,550,175
2017	779,684	927,369	4,615,595	6,322,649
2018	749,154	931,353	4,275,537	5,956,043
2019	695,815	1,058,415	4,019,431	5,773,662
2020	704,836	1,149,123	4,299,867	6,153,826

5.5.2.1.3 Urine and dung from grazing animals (F_{PRP})

The term F_{PRP} is estimated using Equation 11.5. For the required values of the equation see Chapter 5.3. Annual amount of urine and dung nitrogen deposited by grazing animals on pasture, range and paddock (F_{PRP}) for the period 1985-2020 are provided in Table 5.5.7.

5.5.2.1.4 Crop residue N including forage/ pasture renewal (F_{CR})

Nitrogen input from crop residues was estimated in accordance with the Tier 1 methodology, Equation 11.7A of the 2006 IPCC Guidelines. Activity data on crop yields and annual area of harvested crops were taken from the HCSO. To estimate the N added to soils from crop residues and forage/pasture renewal mainly default parameters from the Table 11.2 of the 2006 IPCC Guidelines were used. Since yield statistics are reported as field-dry weight a correction factor was applied to estimate dry matter yields in accordance with Equation 11.7 of 2006 IPCC Guidelines. In the case of wheat parameters provided for grains were used, because the default values given for wheat in the 2006 Guidelines are inappropriate for Hungarian wheat species. For rapeseed and sunflower seed, for which default values are unavailable in the 2006 IPCC Guidelines, country-specific values of ratio of above-ground residues, dry matter to harvested yield crop and N content of above-ground residues for crop were used, while N-contents of below-ground biomass for these crops were calculated using default values provided for 'beans and pulses' in the 2006 IPCC Guidelines. Dry matter contents of forage crops as lucerne-hay, red clover-hay, silo maize and grass hay were sourced from the Hungarian Nutrition Codex, 2004. Input factors used to estimate the N added to soils from crop residues are provided in Table 5.5.6.

The 2006 IPCC method accounts for the effect of residue burning or other removal of residues. Annual areas of burning for cereals ($Area_{burnt(T)}$) were estimated based on expert judgement. It was taken into account for the years before 1990, because burning of crop residues has been banned since 1986 in Hungary. A decreasing proportion of illegal field burning for cereals was assumed for the period between 1986 and 1990. Since this submission emissions from rice field burning due to plant protection reasons have already been reported, for the estimation of area of burnt rice fields see chapter 5.7.

Equation 11.7 requires fractions of total area of crops that is renewed annually. For annual crops $Frac_{Renew}=1$ was assumed, while for Lucerne hay (Alfalfa) and Red clover hay 25%, as the area of these forage crops are renewed on average every four years. In addition, $Frac_{Renew}=0.2$ was assumed for the forage/pasture renewal, assuming five-year renewal frequency based on expert judgement (Monori, 2015).

In the fraction of above-ground residues of crops removed annually ($Frac_{Remove}$), straw used as bedding materials was taken into account. Proportion of straw used as bedding materials were subtracted here, to avoid double counting, as this N is taken into account in the term of F_{AM} . This fraction was calculated consistently with the 3.B Manure management and the 3.D.a.2 Annual amount of animal manure N applied to soils (F_{AM}). In response to a recommendation from the 2017 UNFCCC review, this section has been supplemented with the following information on the derivation of the value $Frac_{Remove}$.

According to the 2006 IPCC Guidelines survey of experts in country is required to obtain data on $Frac_{Remove}$. If data for $Frac_{Remove}$ are not available, assume no removal. While, in accordance with the p. 10.64 of the IPCC Guidelines where organic forms of bedding material (straw, sawdust, chippings, etc.) are used, the additional nitrogen from the bedding material should also be considered as part of the managed manure N applied to soils. However, the 2006 IPCC Guidelines referring to the EEA, 2002 states, that the volatilization losses from bedding is zero. It should be noted that the most up-to-date version of the EMEP/EEA Emission Inventory Guidebook does consider the bedding material as a source of ammonia. Consequently, the use of Tier 2 for the volatilization losses entails the detailed characterization of the flow of nitrogen through the manure management. In housing systems with bedding the bedding material is generally straw and the bedding material should be considered as a part of the nitrogen budget.

As Hungary uses a N-flow approach to calculate the emissions from 3.B and 3.D, which is in line with the IPCC Guidelines, the N_2O emissions from straw used for bedding is reported in CRF 3.D.a.2 Animal manure applied to soils, and this amount of N was taken into account in the value of $Frac_{Remove}$. The value of $Frac_{Remove}$ was calculated for all year from the N content of straw used for bedding divided by the sum of the N content of the above-ground biomass of grain crops of which straw is used for bedding (wheat, barley, rye and oats). For other crops the value of $Frac_{Remove}$ was zero.

Table 5.5.5 Nitrogen in bedding materials and $Frac_{Remove}$ for the BY and 1990-2020

Year	N input from bedding materials [kg]	N content of above-ground biomass of grain crops used as bedding material [kg]	$Frac_{Remove}$ (for Wheat, Barley, Rye and Oat)
BY	18,570,123	45,816,497	41%
1990	16,288,098	54,154,798	30%
1991	14,942,494	53,999,172	28%
1992	12,581,264	38,806,951	32%
1993	10,876,230	32,929,235	33%
1994	9,783,694	48,318,199	20%

Year	N input from bedding materials [kg]	N content of above-ground biomass of grain crops used as bedding material [kg]	Frac _{Remove} (for Wheat, Barley, Rye and Oat)
1995	9,689,709	44,110,126	22%
1996	9,784,246	37,435,227	26%
1997	9,433,348	48,700,129	19%
1998	9,366,668	45,877,198	20%
1999	9,541,831	28,788,874	33%
2000	9,400,809	34,985,052	27%
2001	8,875,423	47,729,049	19%
2002	8,872,012	38,049,930	23%
2003	8,740,600	30,688,130	28%
2004	8,192,855	53,247,162	15%
2005	7,747,942	45,797,614	17%
2006	7,497,936	40,484,154	19%
2007	7,513,447	38,025,633	20%
2008	7,282,139	50,820,104	14%
2009	7,040,603	40,763,617	17%
2010	7,083,717	35,366,925	20%
2011	7,031,319	37,476,309	19%
2012	7,084,451	37,608,560	19%
2013	7,150,026	44,180,338	16%
2014	7,341,000	46,829,467	16%
2015	7,485,597	47,625,106	16%
2016	7,563,525	50,157,836	15%
2017	7,386,764	46,362,040	16%
2018	7,355,804	44,676,710	16%
2019	7,288,569	46,831,375	16%
2020	7,119,336	45,684,704	16%

There is not comprehensive survey on the amount of crop residues burned as fuel in Hungary. Thus, no removal for burning of fuel was assumed. Amount of N in crop residues, including N-fixing crops, and from forage/ pasture renewal, returned to soils are shown in Table 5.5.7.

Table 5.5.6 Parameters used to estimate emissions from crop residues

Crops	Dry matter fraction of harvested product (DRY)	Slope	Intercept	N content of above-ground residues (N _{AG})	Ratio of below-ground residues to above-ground biomass (R _{BG-BIO})	N content of below-ground residues (N _{BG})
Wheat ¹	0.880	1.09	0.88	0.0060	0.22	0.009
Maize (corn)	0.870	1.03	0.61	0.0060	0.22	0.007
Rice	0.890	0.95	2.46	0.0070	0.16	0.009
Barley	0.890	0.98	0.59	0.0070	0.22	0.014
Rye	0.880	1.09	0.88	0.0050	0.22	0.011
Oats	0.890	0.91	0.89	0.0070	0.25	0.008
Bean	0.900	0.36	0.68	0.0100	0.19	0.010

Crops	Dry matter fraction of harvested product (DRY)	Slope	Intercept	N content of above-ground residues (N_{AG})	Ratio of below-ground residues to above-ground biomass (R_{BG-BIO})	N content of below-ground residues (N_{BG})
Peas	0.910	1.13	0.85	0.0080	0.19	0.008
Soya-bean	0.910	0.93	1.35	0.0080	0.19	0.008
Green peas	0.910	1.13	0.85	0.0080	0.19	0.008
Potatoes	0.220	0.10	1.06	0.0190	0.20	0.014
Sugar beat	0.220	0.10	1.06	0.0190	0.20	0.014
Sunflower ²	0.800	NA	NA	0.0057	0.19	0.008
Rape ²	0.700	NA	NA	0.0033	0.19	0.008
Lucerne-hay ³	0.864	0.29	0.00	0.027	0.40	0.019
Red Clover-hay ³	0.855	0.29	0.00	0.027	0.40	0.019
Maize (silo) ³	0.317	1.03	0.61	0.006	0.22	0.007
Meadows ³	0.874	0.18	0.00	0.015	0.54	0.012

¹2006 IPCC default for 'grains' was applied, as data for wheat are inappropriate for Hungarian species.

²Dry matter content and R_{AG} are country-specific based on Zsembeli et. al, 2011. $R_{AGsunflower}=3.0$, $R_{AGrape}=2.0$.

³Values of DRY are country-specific, sourced from Hungarian Nutrition Codex, 2004.

5.5.2.1.5 N mineralization associated with loss of SOM (F_{SOM})

F_{SOM} refers to the amount of N mineralized from loss in soil organic C in mineral soils through land-use change or management practices. To estimate the N mineralized as consequence of this loss of soil carbon the Equation 11.8 of 2006 IPCC Guidelines was applied. The activity data was the carbon loss from management changes under 4.B.1 cropland remaining cropland/ mineral soils.

CRF category 4.B.1 covers conversions of set-aside croplands and non-set-aside-croplands to each other in Hungary. Among these conversions non-set-aside croplands conversions to non-set-aside croplands and set-aside croplands conversions to non-set-aside croplands leads to carbon losses. These carbon losses calculated in the LULUCF sector based on the detailed land-use matrices were used as activity data to calculate the N-losses due to mineralization. (See also Section 6.6.2). According to the 2006 IPCC Guidelines, usage of disaggregated land-use categories of 4.B.1 cropland remaining cropland, in the calculation meets the requirement of Tier 2 methodology.

The default C:N ratio of the soil organic matter of 10 was used. The resulted annual values for F_{SOM} are provided in Table 5.5.7.

Table 5.5.7 Amount of N inputs to soils BY-2020

Year	Synthetic Fertilizers (F _{SN})	Organic N applied as fertilizer (F _{ON})			Urine and dung inputs to grazed soils (F _{PRP})	Crop Residues (F _{CR})	N mineralization associated with loss of SOM (F _{SOM})
		Animal manure (F _{AM})	Sewage (F _{SW})	Compost (F _{COMP})			
1000t N							
BY	589	206	NO	NO	36	159	0.45
1990	358	189	0.2	0.2	30	152	0.77
1991	140	176	0.2	0.2	30	183	0.77
1992	148	151	0.3	0.2	27	125	0.77
1993	161	132	0.3	0.2	22	108	0.77
1994	222	119	0.4	0.2	18	143	0.76
1995	191	118	0.6	0.2	17	139	0.76
1996	203	115	0.5	0.3	16	140	0.76
1997	206	110	0.4	0.3	16	159	0.76
1998	248	113	0.5	0.2	16	151	0.49
1999	262	114	0.4	0.3	17	135	0.49
2000	258	115	0.5	0.3	18	114	0.48
2001	275	113	0.4	0.3	18	165	0.48
2002	303	115	0.5	0.4	17	136	0.48
2003	289	115	0.5	0.6	17	111	0.48
2004	293	112	0.6	0.5	18	191	0.48
2005	260	106	0.9	0.7	19	182	0.03
2006	289	104	0.9	0.8	18	168	NO
2007	320	104	0.8	1.1	17	126	NO
2008	294	101	1.0	1.7	17	196	NO
2009	275	96	1.1	2.4	16	162	NO
2010	281	96	1.0	2.8	16	142	NO
2011	302	95	0.9	3.8	16	163	0.36
2012	313	98	0.8	3.4	16	134	0.93
2013	343	99	0.6	5.3	16	166	0.92
2014	341	102	0.6	5.4	17	195	0.92
2015	378	104	0.6	5.3	17	173	0.92
2016	404	102	0.7	5.6	17	204	0.92
2017	424	99	0.7	6.3	18	184	0.92
2018	424	99	0.7	6.0	19	187	0.92
2019	416	99	0.7	5.8	19	190	0.92
2020	443	98	0.7	6.2	19	189	0.91


5.5.2.1.6 Area of drained/managed organic soils (F_{OS})

Cultivation of Histosols is not occurring in Hungary, therefore notation key 'NO' is reported for the N₂O emission in CRF Table 3.D. Following the recommendation in the 2013 annual review, and the 2016 in-country review, the NIR has been supplemented with the following justification.

In the Hungarian soil classification system Peat soils and Ameliorated peat soils could be identified as WRB Histosols. It should be noted that one of the features of the Hungarian genetic soil classification system is that the name of certain types of soils hints at the condition of the soil formation and not

necessarily at their current characteristics. Thus, the translation of this name into English can cause some misunderstanding. I. e. words like 'swampy', or 'peat' in the name of the Hungarian genetic soil types does not definitely mean that the soil falls in the group of Histosols.

Total areas of Peat and Ameliorated peat soils are 41,612 and 90,685 ha based on AGROTOPO, the Hungarian agro-topographical map at scale of 1: 100,000. The delineated areas of Peat and Ameliorated Peat soils are shown on **Figure 5.5.2**.



Source: Research Institute for Soil Science and Agricultural Chemistry (Hungary)

Note: 'lecsapolt' = Ameliorated peat soils, 'síkláp'=Peat soils


Figure 5.5.2 Peat soils and Ameliorated peat soils in Hungary

Peat soils form and can be restored under wetland conditions, which are 'ex lege' protected in Hungary in accordance with the Article 23 Paragraph (2) of the Act No LIII of 1996. (This law is in force currently, but Wetlands was protected decades ago.) Hungary also signed the Ramsar convention in 1971, thus the protection of wetlands is also encouraged based on this convention. As a consequence, areas of wetland soils are mainly national parks and landscape protection areas. **In summary, cultivation of organic soils/ Histosols is prohibited by law in Hungary.**

Before the 1950's attempts were made on the utilization of peat lands, by draining. The results of this activity are the Ameliorated peat soils. After draining, the organic carbon content of these soils declined resulting from the oxidization of organic matters during more than 60 years of continuous cultivation. Consequently, as it is proved by measurements, these cultivated Ameliorated peat soils

have an average humus content of 6%, which do not meet the definition of 'Histosols' or 'organic soils' used in the IPCC Guidelines.

The FAO database on GHG emissions indicates 229.2 kha cultivated organic soil for Hungary based on the Harmonized World Soil Map and the Global Land Cover 2000 dataset, but this area exceeds the total area of Peat soils and Ameliorated peat soils in Hungary (i.e., 132.3 kha altogether) delineated based on higher resolution national soil map regardless of land cover and land use. This significant overestimation indicates the high uncertainty in the estimated area from the harmonized international databases. Areas of cultivated organic soils based on FAO GHG emission database are shown on Figure 5.5.3



Source: FAO

Figure 5.5.3 Cultivated Histosols in Hungary based on FAO GHG emission database

The data of the Hungarian Soil Protection and Monitoring System (hereafter referred to as TIM) prove this fact, namely there are no cultivated organic soils in Hungary. Arising from the definition of Histosols in the 2006 IPCC Guidelines, the organic carbon contents of agricultural soils were analyzed to distinguish between Histosols (organic soils) and mineral soils.

The organic matter content of agricultural soils was derived from the measurements of the TIM. A summary about the TIM is also available in English, in Várallyay, 2005. The measured humus content data for **1014 sample points** was received from the Plant, Soil and Agri-environmental Directorate of

the National Food Chain Safety Office. The humus content data based on laboratory measurements are available for the year 1998 and 2000. According to the data of the TIM, the humus content of the soil does not reach 20%, the lower limit of humus content in organic soils, in any of the TIM points. The Table 5.5.8 summarizes the measured humus content data for Ameliorated peat soils and Peat soils.

Table 5.5.8 Measured humus content for Ameliorated peat soils and Peat soils based on TIM

Code	EOV_X	EOV_Y	Humus content	Humus content	Average humus content	Hungarian genetic soil types	IPCC soil type
			1998	2000	1998, 2000		
			%	%	%		
I0908	489320	258154	5.69	4.97	5.33	Ameliorated peat soils	Wetland
I1614	531672	149870	6.01	6.7	6.36	Ameliorated peat soils	Wetland
I1707	590004	203609	5.36	6.04	5.70	Ameliorated peat soils	Wetland
I3008	520145	273846	5.69	4.61	5.15	Ameliorated peat soils	Wetland
I3819	512187	219795	6.77	5.3	6.04	Ameliorated peat soils	Wetland
I4019	526785	166020	5.65	5.9	5.78	Peat soils	Wetland
I4715	901096	276942	2.83	2.44	2.64	Ameliorated peat soils	Wetland
S4920	504239	148173	5.25	5.85	5.55	Peat soils	Wetland
S5008	502087	262733	5.24	5.6	5.42	Ameliorated peat soils	Wetland
S6303	675522	115685	5.11		5.11	Peat soils	Wetland
S6515	901822	285368	12.05	11	11.53	Ameliorated peat soils	Wetland

Source: Soil and Agri-environmental Directorate of the National Food Chain Safety Office (NFC SO)

In 2009, the European Commission extended the periodic Land Use/Land Cover Area Frame Survey (LUCAS) to sample and analyze the main properties of topsoil in 23 Member States of the European Union (EU). The LUCAS Topsoil Data (EC JRC, 2013) providing measured organic carbon data for 20,000 sample points in Europe for the year 2009 were also analyzed and confirmed that organic soils are not cultivated in Hungary.




Figure 5.5.4 Organic carbon contents of soil samples in Hungary based on LUCAS Topsoil Survey, 2009

In the course of the LUCAS Topsoil Survey 497 samples were collected in Hungary, out of them 314 samples from annual croplands, 6 from permanent crops, 60 from woodlands, 4 from shrublands and 104 from grasslands. For more information on the survey and representativity issues see also Tóth G. et al, 2013. As the data on organic carbon content of soil samples revealed, only two samples have 12.06 and 12.15 per cent organic carbon content which might be classified into the group of Histosols, but none of them was from cultivated cropland. (The point IDs of these samples are 49422714 and 48782686, respectively.) The resulted organic carbon contents expressed in terms of g/kg are shown on Figure 5.5.4. Photos of sample plots where the samples having about 12% organic carbon content were collected are shown in Figure 5.5.5. According to the LUCAS survey, the land cover of these plots is grassland without tree/ shrub cover (Code: E20). Therefore, cannot be disturbance leading to loss of organic matter stored in the soil.



Figure 5.5.5 Photos of sample plots where soil samples with 12% organic carbon content were collected in the course of LUCAS Topsoil Survey, 2009

As a consequence of the facts due to the domestic legislation on one hand and based on data from soil surveys on the other, it can be confirmed that organic soils are not cultivated in Hungary. Soils in

Hungary, which are classified as Histosols in international soil databases (e.g., FAO HWSD) are either on protected wetlands (Peat soils) or, if on managed croplands, they have lost most of their carbon content (Ameliorated peat soils) and cannot be classified as Histosols.

5.5.2.2 Indirect Emissions (CRF 3.D.b)

In addition to the direct emissions of N_2O from managed soils, emissions of N_2O also take place through two indirect pathways. The first of these pathways is the volatilization of N as NH_3 and NO_x , and the subsequent deposition of these gases and their products onto soils and water surfaces. The sources of N volatilization are not confined to agricultural fertilizers and manures, but also include fossil fuel combustion, biomass burning, and processes in the chemical industry. In the Hungarian inventory reporting of volatilization of N and the deposition of NH_3 and NO_x is confined to agricultural sources.

The second pathway of indirect emissions is the leaching and runoff from land of N from agricultural inputs.

5.5.2.2.1 Indirect N_2O emissions through atmospheric deposition of N volatilized

In response to a recommendation from the in-country review conducted in 2016 this section was supplemented with information on the EMEP/EEA methodology used to derive $Frac_{GASF}$ and $Frac_{GASM}$ including the parameters and equation used.

The Hungarian national system takes advantage of parallel inventory preparation and reporting of air pollutants under the LRTAP Convention ensuring efficiency and consistency in the compilation of emission inventories, because a wide range of substances using common datasets and inputs. Hungary applies the most up to date EMEP/EEA Emission Inventory Guidebook to calculate the agricultural NH_3 and NO_x emissions. A detailed description of the method applied for NH_3 and NO_x is given in the report 'Hungary's Informative Report 2019 submitted under the UNECE Convention on Long-range Transboundary Air Pollution' (CLRTAP) (UNECE, 1999) and the National Emissions Ceilings Directive (EP and CEU, 2016).

The N_2O emissions from atmospheric deposition of N volatilized from managed soil was calculated based on Tier 1 methodology, following the Equation 11.9 of 2006 IPCC Guidelines. The activity data are the same as those under 3.D.a.

The method requires values for the fractions of N that are lost through volatilization ($Frac_{GASF}$, $Frac_{GASM}$) and the emission factor (EF_4). The volatilization rates for Hungary are country-specific based on the reported NH_3 and NO_x emissions.

Country-specific volatilization fraction of synthetic fertilizers ($Frac_{GASF}$) includes the NH_3 -N and NO_x -N losses from fertilizers calculated by fertilizer types.

Country-specific volatilization fraction of applied organic N fertilizer materials and urine and dung N deposited by grazing animals ($Frac_{GASM}$) includes:

- NH_3 -N and NO_x -N losses from livestock manure application on agricultural soils.
- NH_3 -N and NO_x -N losses from dung and urine deposited by grazing livestock.
- NH_3 -N and NO_x -N losses from sewage sludge applied to soils.
- NH_3 -N and NO_x -N losses from compost applied to soils.

To calculate the NH_3 -N losses the reported NH_3 emissions were multiplied by 14/17. In the air pollutant inventory NO_x is reported as NO_2 ; therefore, the NO_2 emissions were multiplied by 14/46 to get the NO_x -N losses.

NH_3 -N and NO_x -N volatilization losses from mineral fertilizer application

The parallel and consistent emission inventory compilation enables the use of country-specific data, which is more accurate than the use of the IPCC default value of $Frac_{GASF}$.

NH_3 and NO_x emissions from Sector 3 Agriculture are estimated according to the 2019 EMEP/EEA Guidebook. For the calculation of NH_3 -N losses from synthetic fertilizers the tier 2 methodology was applied. This method uses specific NH_3 emission factors for different types of synthetic fertilizers depending on the soil acidity and climate. To summarize, NH_3 emissions can be calculated by means of the following equation:

$$E_{fert_{NH_3}} = \sum_{i=1} \sum_{j=1} m_{fert_{i,j}} \cdot EF_{i,j} \quad (\text{Equation 5.9})$$

Where:

$E_{fert_{NH_3}}$ = NH_3 emission from fertilization ($kg\ a^{-1}\ NH_3$)

m_{fert_i} = mass of fertilizer type i consumed nationally ($kg\ a^{-1}\ N$)

$EF_i = EF$ for fertilizer type i in region j ($kg\ NH_3\ (kg\ N)^{-1}$)

Definitions of climate zones of the 2019 EMEP/EEA Guidebook are the same as those of 2006 IPCC Guidelines. According to the Guidelines, cool climate zone has an annual mean temperature below 15°C. The annual mean temperature in most parts of Hungary is between 10 and 11 °C, hence the emission factors given for cool climate zone were applied for the whole country.

Proportion of soil with normal pH and high pH was determined based on the most up-to-date high resolution (250m) soils map (Tóth, G. et al., 2015). Emission factors provided by soil pH in the EMEP/EEA Guidebook were weighted by the resulted proportions and weighted national average emission factors, given in Table 5.5.9, were calculated for each fertilizer types.

Table 5.5.9 Country specific emission factors for ammonia emission from fertilizers

Fertilizers	IEF _i [kg NH ₃ kg ⁻¹ N]
Ammonium nitrate	0.025
Anhydrous ammonia	0.028
Ammonium phosphate, NP mixtures	0.074
Ammonium sulphate	0.134
Calcium ammonium nitrate	0.013
Other straight N compounds	0.015
Nitrogen solutions	0.096
Urea	0.160
NK mixtures	0.025
NPK mixtures	0.074
Implied EF (2020)	0.048

Mass of fertilizer *i* consumed nationally was derived from the sales statistics by product line.

Detailed data on fertilizer consumption by fertilizer types is not published in the Hungary's IIR, 2020 either this report because of data confidentiality. However, main driver in the trend of NH₃ emissions from inorganic fertilizers is the urea consumption and the time series of urea use is published in this report.

For the calculation of NO_x emissions, the tier 1 methodology of the 2019 EMEP/EEA Guidebook was applied, by means of the following equation:

$$E_{\text{pollutant}} = AR_{N_applied} \cdot EF_{\text{pollutant}}$$

(Equation 5.10)

Where:

$E_{\text{pollutant}}$ = amount of pollutant emitted (kg a⁻¹)

$AR_{N_applied}$ = amount of N applied in fertilizer or organic waste (kg a⁻¹)

$EF_{\text{pollutant}}$ EF of pollutant (kg kg⁻¹)

Emissions were calculated as a fixed percentage of total fertilizer nitrogen applied to soil. For all types of mineral fertilizer, the default emission factor of 0.04 kg NO₂ per kg applied fertilizer-N was used (EEA, 2019).

In 2020 the value of $Fra_{C_{GASF}}$ was 0.05, which is lower than the IPCC default value, because of the low proportion of Urea in the total fertilizer use.

NH₃-N and NO_x-N volatilization losses from organic N fertilizers and N deposited by grazing animals

Similarly, to $Frac_{GASF}$, $Frac_{GASM}$ is also an annual implied value of N-losses referring to NH₃-N as well as NO_x-N losses from organic manure that is volatilized as NH₃ and NO_x.

NH₃-N volatilization losses from livestock manure application

For the calculation of NH₃ emissions from manure application, we partly used the EMEP/Gb Tier 1 and Tier 2 emission factors, on the other hand, the Tier 2 emission factors have been adjusted according to the specific NH₃ reduction technologies and their penetration, taking into account the measures for solid/liquid manure application. Information on the derivation of correction factors to account for NH₃ reduction technologies for the unabated emission factor is described in the 2021 submission of Hungary's Informative Inventory Report, 2021 (HMS, 2021). Emission factors and the level of the applied methodologies to calculate NH₃ emissions from manure application are shown in Table 5.5.10.

Table 5.5.10 Emission factors for NH₃ emissions from animal manure application

Livestock	Manure type	EF spreading [kg NH ₃ -N (kg TAN) ⁻¹]	Applied methodology
Cattle	slurry	0.32	Tier 3
	solid	0.37	Tier 3
Fattening pigs	slurry	0.2	Tier 3
	solid	0.24	Tier 3
Sows	slurry	0.14	Tier 3
	solid	0.24	Tier 3
Sheep	solid	0.9	Tier 2
Horses, Mules and Asses	solid	0.9	Tier 1
Laying hens	solid/ slurry	0.24	Tier 3
Broilers	solid	0.20	Tier 3
Turkey	solid	0.54	Tier 1
Ducks	solid	0.54	Tier 1
Geese	solid	0.45	Tier 1

NO_x-N emissions from animal manure spreading

NO_x emissions were calculated using the default emission factors for 3.D.a.2 Animal manure applied to soils. The default emission factors were calculated on the basis that all manure is stored before surface application without rapid incorporation.

NH₃-N and NO_x-N volatilization losses from sewage sludge application

As with the application of animal manure, the emissions from the application of sewage sludge were calculated taking into account NH₃ mitigation technologies. Therefore, the 'base' emission factor (0.13 kg NH₃ per kg N applied) provided in the EMEP/EEA Guidebook (EEA, 2019) was adjusted according to

the annual penetration of the applied NH₃ abatement technologies. For more information see HMS, 2021.

NO_x emissions were estimated using the default emission factor of 0.04 kg NO₂ per sewage sludge Nitrogen (EMEP/EEA, 2019).

NH₃-N and NO_x-N volatilization losses from compost application

For the calculation of NH₃ emissions the default emission factor provided for other organic waste (0.08 kg NH₃ per kg N applied) was applied (EEA, 2019).

NO_x emissions were estimated using the default emission factor of 0.04 kg NO₂ per N applied (EMEP/EEA, 2019).

Derivation of N losses from mineral fertilizer and applied organic N fertilizer materials including grazing from the Hungarian air pollutant emission inventory is demonstrated in Table 5.5.11 for the year 2020. The time series of the volatilization losses were calculated similarly for all years of the inventory period.

Annual NH₃-N and NO_x-N volatilization losses from synthetic fertilizers and organic N fertilizers (including grazing) for the BY and the period from 1990 to 2020 are provided in Table 5.5.12 together with the resulted values of Frac_{CGASF} and Frac_{CGASM}.

Table 5.5.11 Derivation of NH₃-N and NO_x-N volatilization losses from synthetic and organic N fertilizers (including grazing) for the year 2020

NFR code	Long name	NO _x (as NO ₂)	NH ₃	Total N volatilized [kg N]
3Da1	Inorganic N-fertilizers (includes also urea application)	17.73	21.35	22,976,166
3Da2a	Animal manure applied to soils	3.92	11.35	10,540,196
3Da2b	Sewage sludge applied to soils	0.03	0.04	44,211
3Da2c	Other organic fertilizers applied to soils (including compost)	0.25	0.50	486,037
3Da3	Urine and dung deposited by grazing animals	0.77	1.98	1,865,196
3D	Agricultural soils	22.70	35.22	35,911,806

Table 5.5.12 $\text{NH}_3\text{-N}$ and $\text{NO}_x\text{-N}$ volatilization losses from synthetic and organic N fertilizers (including grazing) BY and 1990 to 2020

Year	N losses from mineral fertilizer	N losses from applied organic N fertilizer materials and grazing	Frac _{GASF}	Frac _{GASM}
	kg N	kg N		
BY	32,967,447	34,261,580	0.06	0.14
1990	22,932,366	31,048,941	0.06	0.14
1991	10,113,426	28,963,640	0.07	0.14
1992	8,586,793	24,729,681	0.06	0.14
1993	8,377,202	21,742,060	0.05	0.14
1994	10,222,155	19,619,855	0.05	0.14
1995	10,617,988	19,712,592	0.06	0.15
1996	10,531,507	19,874,288	0.05	0.15
1997	11,036,074	19,288,854	0.05	0.15
1998	12,072,679	19,919,170	0.05	0.15
1999	12,906,199	20,308,640	0.05	0.15
2000	12,623,718	20,759,579	0.05	0.15
2001	13,455,514	20,248,860	0.05	0.15
2002	14,825,530	20,284,291	0.05	0.15
2003	14,526,641	20,208,140	0.05	0.15
2004	15,391,037	19,209,320	0.05	0.15
2005	14,113,368	18,099,447	0.05	0.14
2006	15,687,551	17,363,281	0.05	0.14
2007	16,298,925	16,909,373	0.05	0.14
2008	12,219,732	16,288,201	0.04	0.14
2009	11,281,073	15,322,083	0.04	0.13
2010	11,406,120	15,162,709	0.04	0.13
2011	12,582,043	14,737,100	0.04	0.13
2012	12,693,250	14,328,646	0.04	0.12
2013	15,168,532	14,065,223	0.04	0.12
2014	15,275,607	14,211,483	0.04	0.11
2015	17,828,675	14,239,785	0.05	0.11
2016	18,635,667	14,170,509	0.05	0.11
2017	20,768,060	13,676,666	0.05	0.11
2018	20,753,782	13,311,669	0.05	0.11
2019	21,087,678	13,128,431	0.05	0.11
2020	22,976,166	12,935,640	0.05	0.10

5.5.2.2.2 Leaching and runoff

The N_2O emissions from the N lost through leaching and runoff was calculated using the Tier 1 methodology and Equation 11.10 of 2006 IPCC Guidelines. The activity data are the same as those under 3.D.a.

In accordance with the 2006 IPCC Guidelines for humid regions and in regions where irrigation is used, the default value of $F_{\text{LEACH-(H)}}$, 0.3 was applied. For dryland regions, where precipitation is lower than evapotranspiration throughout most of the year, F_{LEACH} was assumed to be zero. According to the IPCC methodology the determination of proportion of irrigated areas and humid regions are required. Thus, the Equation 11.10 of the 2006 IPCC Guidelines can be considered as the same as Equation 5.11.

$$N_2O_{(L)}-N=(F_{\text{SN}}+F_{\text{ON}}+F_{\text{PRP}}+F_{\text{CR}}+F_{\text{SOM}})\cdot(F_{\text{Cirr}}+F_{\text{Cwet}})\cdot F_{\text{LEACH-H}}\cdot EF_5$$

(Equation 5.11)

Where:

N_2O-N =annual amount of N_2O-N produced from leaching and run-off of N additions to managed soils in regions where leaching/runoff occurs, kg N_2O-N

F_{SN} =annual amount of synthetic fertilizer N applied to soils in regions where leaching/runoff occurs, kg $N\text{ yr}^{-1}$

F_{ON} =annual amount of managed animal manure, compost, sewage sludge and other organic N additions applied to soils in regions where leaching/runoff occurs, kg $N\text{ yr}^{-1}$

F_{PRP} = annual amount of urine and dung N deposited by grazing animals in regions where leaching/runoff occurs, kg $N\text{ yr}^{-1}$

F_{CR} = amount of N in crop residues (above- and below-ground), including N-fixing crops, and from forage, pasture renewal, returned to soils annually in regions where leaching/runoff occurs, kg $N\text{ yr}^{-1}$

F_{SOM} = annual amount of N mineralized in mineral soils associated with loss of soil C from soil organic matter as a result of changes to land use or management in regions where leaching/runoff occurs, kg $N\text{ yr}^{-1}$

F_{Cirr} = fraction of irrigated agricultural areas

F_{Cwet} = fraction of humid agricultural areas

$F_{\text{LEACH-(H)}}$ =fraction of all N added to/mineralized in managed soils in regions where leaching/runoff occurs

EF_5 =emission factor for N_2O emissions from N leaching and runoff, kg N_2O-N (kgN leached and runoff) $^{-1}$

Derivation of fraction of irrigated areas (F_{Cirr})

Proportions of irrigated areas were derived annually from HCSO's statistics on irrigated, utilized, and total agricultural areas. The 2006 IPCC Guidelines require to distinguish the drip irrigated areas within the irrigated areas, because according to the methodology, N-leaching is assumed to be unlikely on drip irrigated areas. The HCSO publishes the total irrigated areas annually, which include the drip irrigated areas as well for the period 1989-2014. Unfortunately, the drip irrigated areas are not reported separately in this statistic. For the period before 1989 data on the total irrigated areas and within this the drip irrigated areas are available, separately. Therefore, since the 2016 submission it has been decided to account the total irrigated areas, i.e., areas of drip irrigation are not subtracted from the total irrigated areas, due to lack of reliable and consistent statistics on drip irrigation for the whole time series. As the fraction of drip irrigation was 1.5% of the total irrigated areas on average for the period 1985-1989, inclusion of drip irrigation in the emission estimation probably causes a negligible overestimation of emissions.

Because it is assumed that N is applied only on utilized agricultural areas, annual fractions of irrigated areas were calculated as a fraction of utilized agricultural areas. (Calculation of irrigated areas as fraction of the total agricultural areas could result in an underestimation of the amount of N leached and run-off.) Utilized areas were taken from the HCSO's censuses for the year 2000, 2010 and 2013. Annual areas for years between censuses data have been based on linear interpolation between data points. For the period before 1990 the total agricultural area was assumed to be utilized, as agricultural areas started to be abandoned as a result of the change of the regime in the 90's.

In 2019 108.3 thousand hectares (2.3% of the utilized agricultural areas) were irrigated. Although, the National Water Authority indicates a total area with water right permit 190.0 ha (about 4% of total agricultural areas) for 2014, according to the HCSO's statistics 57% of the irrigable areas were irrigated in 2019. In 2020, the HCSO did not carry out a survey on irrigation, so the proportion of irrigated areas for 2020 was also based on the 2019 data.

In Hungary the fraction of irrigated areas is significantly lower than the average of the EU Member States because of the limited and outdated irrigation system. In 2010 5.8% of the Member States utilized agricultural areas were irrigated. In contrast this proportion was 2.4% for Hungary in that year according to the EUROSTAT statistics. Because of this low proportion of irrigation, the improvement of water management efficiency and irrigation systems are among the priorities of the Hungarian Rural Development Program for the period 2014-2020. Additionally, the large inter-annual fluctuations in the harvested crop productions also reflect the high dependence on weather conditions (e.g., droughts) partially due to the low proportion of irrigated areas.

The resulted fractions of irrigated areas are shown in Table 5.5.13.

Table 5.5.13 Derivation of activity data on irrigated agricultural areas, from the BY to 2019

Year	Total irrigated areas	Total agricultural areas	Utilized agricultural areas (UAA)	Irrigated areas as % of UAA
	ha	1000 ha	1000 ha	
BY	147,871	6186	6186	2.4%
1990	216,937	6132	6132	3.5%
1991	148,669	6116	5989	2.5%
1992	177,808	6091	5839	3.0%
1993	180,088	6080	5702	3.2%
1994	160,384	6064	5562	2.9%
1995	146,541	6048	5422	2.7%
1996	126,344	6028	5278	2.4%
1997	81,908	6008	5137	1.6%
1998	93,431	5990	4997	1.9%
1999	44,822	5972	4858	0.9%
2000	125,866	5745	4555	2.8%
2001	104,172	5729	4598	2.3%
2002	117,035	5698	4629	2.5%
2003	148,642	5667	4660	3.2%
2004	120,596	5632	4686	2.6%
2005	75,161	5604	4718	1.6%
2006	78,193	5570	4744	1.6%
2007	121,064	5536	4769	2.5%

Year	Total irrigated areas ha	Total agricultural areas 1000 ha	Utilized agricultural areas (UAA) 1000 ha	Irrigated areas as % of UAA
2008	80,149	5503	4794	2.0%
2009	107,106	5471	4820	2.2%
2010	114,550	5261	4686	2.4%
2011	101,046	5256	4681	2.2%
2012	124,944	5257	4682	2.7%
2013	118,934	5259	4657	2.6%
2014	130,400	5266	4663	2.8%
2015	124,300	5266	4663	2.7%
2016	108,233	5286	4680	2.3%
2017	108,595	5305	4697	2.3%
2018	111,401	5298	4691	2.4%
2019	108,300	5271	4667	2.3%

Derivation of fraction of humid regions ($Frac_{wet}$)

To estimate the fraction of humid regions is also required to calculate the emissions from N-leaching. Proportion of humid regions was determined based on the analysis of the 30-year climate means (1981-2010) of the monthly precipitation and evaporation data from the HMS climate database.

According to the definition of the $Frac_{LEACH-(H)}$ in the 2006 IPCC Guidelines, the determination of 'rainy seasons' are required based on the data on precipitation and Pan Evaporation (E_{PAN}). The 2006 IPCC Guidelines define the 'rainy seasons' as periods when $rainfall > 0.5 \cdot Pan \text{ Evaporation}$, which criteria is equal to that $P/E_{PAN} > 50\%$, where P is the monthly precipitation.

Table 5.5.14 Data for the derivation of 'rainy seasons' to calculate emissions from 3.D.2.2

Month	Potential Evaporation (PE)	Pan (E_{PAN})	Evaporation	Precipitation (P)	P/E_{pan}	P/PE
January	22.9	-		33.2	-	145%
February	29.4	-		32.8	-	112%
March	54	-		35.3	-	65%
April	83.7	93.8		44.4	47%	53%
May	115.2	128		63.7	50%	55%
Jun	129.1	141.6		73.6	52%	57%
July	157.8	172.3		65	38%	41%
August	148.1	148		63.3	43%	43%
September	94.2	93.1		54.7	59%	58%
October	61.2	62.7		42.6	68%	70%
November	33	-		49.8	-	151%
December	22.3	-		47	-	211%

Source: HMS

Pan Evaporation data is registered from April to October in Hungary. To avoid underestimation of emission it has been decided to examine the Potential Evaporation instead of Pan Evaporation for the remainder months, from November to March. Analysis of data in Table 5.5.14 reveals Jun and the period from September to March can be considered as 'rainy seasons' in Hungary, according to the definition of the 2006 IPCC Guidelines.

According to the 2006 IPCC Guidelines N-leaching could occur where:

$$\sum(\text{rain in rainy season}) - \sum(\text{PE in same period}) > \text{soil water holding capacity} \quad (\text{Equation 5.12})$$

Where:

PE = potential evaporation

Because the soil water holding capacity is generally greater than zero, the following equation can be derived from Equation 5.12:

$$\sum(P \text{ in rainy season}) - \sum(\text{PE in same period}) > 0 \quad (\text{Equation 5.13})$$

Where:

P=precipitation

Evaporation is the process whereby liquid water is converted to water vapor and removed from the evaporating surface. Water evaporates from different surfaces, such as water, soils, and wet vegetation. On agricultural areas the soil and plants are the evaporating surfaces. Thus, evaporation on agricultural areas depends on the weather conditions, soil properties, management practices and crop type. Consequently, PE could be highly different within a country and cannot be expressed with a representative value. To analyze the climatic conditions of leaching and run-off the 30 year means of monthly precipitation and reference evapotranspiration (ET_0) from station data as high-resolution gridded data over Hungary were determined by the HMS. It is important to note that the 2006 IPCC Guidelines on p.11.23 also use the potential evapotranspiration instead of potential evaporation to distinguish between dryland and humid regions.

The FAO Penman-Monteith method was used to determine ET_0 for each month. The definition and concept of ET_0 is as follows:

'The evapotranspiration rate from a reference surface, not short of water is called the reference crop evapotranspiration or reference evapotranspiration and is denoted as ET_0 . The reference surface is a hypothetical grass reference crop with specific characteristics. The concept of the reference evapotranspiration was introduced to study the evaporative demand of the atmosphere independently of crop type, crop development and management practices. As water is abundantly available at the reference evapotranspiration surface, soil factors do not affect evapotranspiration (ET). Relating ET to a specific surface provides a reference to which ET from other surfaces can be

related. It obviates the need to define a separate ET level for each crop and stage of growth. ET_o values measured or calculated at different locations or in different seasons are comparable as they refer to the ET from the same reference surface. The only factors affecting ET_o are climatic parameters. Consequently, ET_o is a climatic parameter and can be computed from weather data. ET_o expresses the evaporating power of the atmosphere at a specific location and time of the year and does not consider the crop characteristics and soil factors. The method has been selected because it closely approximates grass ET_o at the location evaluated, is physically based, and explicitly incorporates both physiological and aerodynamic parameters.' (FAO, ET_o calculator.)

Consequently, in the Equation 5.13 the PE was replaced with the ET_o , and the data of P/ET_o for June and $\sum P/\sum ET_o$ for the period September to March were generated with a spatial resolution of 30 arc-seconds (≈ 1 km) to the analysis. The resulted maps are shown on Figure 5.5.6 and Figure 5.5.7. Subsequently, areas where in the 'rainy seasons', namely June, and the September-March period in Hungary, $\sum P/\sum ET_o > 1$ were determined from GIS analysis of the gridded climate data and the resulted areas were superimposed on the CORINE 2012 land cover database.




Figure 5.5.6 The ratio of average precipitation and average reference evapotranspiration (P/ET_o) for June




Figure 5.5.7 The ratio of total average precipitation and total average reference evapotranspiration ($\Sigma P / \Sigma ET_0$) for the period September to March

From the CORINE 2012 land cover database croplands, grasslands and agricultural mosaics (200<CLC codes<300) were considered to be agricultural lands.

As a result of the GIS analysis of climate and land cover maps (Figure 5.5.8 and Figure 5.5.9) fraction of agricultural lands where N-leaching could occur because of the potential existence of precipitation surplus is 1,516 ha in June, which equates to 0.02% of the agricultural lands in the CORINE database, and 659,439 ha (10.65%) in the September-March period (Table 5.5.15). As humid regions analyzed for the September-March period include the affected areas in June, the total fraction of humid regions is 10.65%. In other words, the resulted areas for Jun and September-March period were not added, because the humid regions in Jun are also humid in the September to March period.

Table 5.5.15 Resulted areas from GIS analysis of climate and CLC, 2012 land cover databases

	Area	
	ha	As % of the total area of the country
Total area of humid regions in June	22,460	0.24%
Total area of humid regions in September-March period	544,345	5.86%
Agricultural lands from CORINE, 2012	6,190,940	67%
	Area	
	ha	As % of the total area of agricultural lands
Humid agricultural lands in June	1,516	0.02%
Humid agricultural lands in the September-March period	659,436	10.65%
Total area of humid agricultural lands in 'rainy seasons' for Hungary	659,436	10.65%




Figure 5.5.8 Humide ($P/ET_o > 1$) agricultural areas by CORINE land cover codes in June




Figure 5.5.9 Humide ($\Sigma P / \Sigma ET_o > 1$) agricultural areas by CORINE land cover codes in the September-March period

5.5.3 Uncertainties and time-series consistency

Implementation of the methodologies provided in the 2006 IPCC Guidelines resulted in significant reduction in the uncertainties in 3.D N₂O Emissions from Agricultural Soils. Uncertainties in this category are driven by uncertainties related to the emission factors. For the default emission factor for direct emissions (EF₁), a range from -70% up to +200% is assigned by the 2006 IPCC Guidelines. This uncertainty range is significantly narrower than the former one provided in the GPG (IPCC, 2000), leading to significant reduction in the overall uncertainty in the N₂O emissions.

For the uncertainties in the activity data as F_{SN}, F_{ON}, F_{PRP}, F_{CRP}, F_{SOM} ±5%, ±36%, ±30%, ±25%, ±91% were calculated, respectively. The resulted combined uncertainty in the activity data for 3.D.a is ±8.6%, which is negligible comparing with the uncertainty in the emission factor. The estimated combined uncertainties in the emissions from 3.D.a were -71%/+200%.

To estimate uncertainties in indirect emissions the same values of uncertainties were applied for activity data as in the calculation of direct emissions. These uncertainties were combined with the uncertainties in the Fra_{C_{GAS}M} (±50%) and Fra_{C_{GAS}F} (±75%). Uncertainty in Fra_{C_{GAS}F} was estimated based on the EMEP/EEA Guidebook (EEA, 2016). For the EF₄ the default uncertainty range provided in the 2006 IPCC Guidelines was applied. The resulting uncertainty for the indirect emission from agricultural soils ranges from -71% to +200%.

5.5.4 QA/QC Information and verification

Direct N₂O emissions

The main driver of the GHG-emissions from the agriculture sector is the N-fertilizer use in Hungary. Therefore, the verification of the amount of N-fertilizer applied is very essential in the QA/QC process in the Agriculture sector.

The amount of the N-fertilizer applied has been compared with the international statistics, namely FAO and IFA (International Fertilizer Industry Association). There is not any difference between the reported N-fertilizer used in the FAO statistics and the GHG-inventory. However, the IFA reports higher N-fertilizer use for the years before 2007. The reasons for it have already been investigated by the experts of the HCSO and the Research Institute of Agricultural Economics and IFA's.

The fertilizer consumption data used in the GHG inventory derives from the HCSO's official statistics. HCSO gets these data from the data collection of the Research Institute of Agricultural Economics. The Research Institute of Agricultural Economics collects data on the sold amount of the different types of fertilizers. The IFA used an expert judgement for the estimation of fertilizer consumption data made by the Yara's (a Norwegian chemical company) experts, recently. The IFA's methodology for expert judgement is based on the sowing area of the main crops, such as cereals, maize and sunflower and so on. The estimation considered the area and the fertilizer need requirements of these crops. Unfortunately, the fertilizer consumption in Hungary is generally lower than the suggested amount, due to the high price of the fertilizer. So, this methodology of IFA resulted in an overestimation.

The HCSO's, the Research Institute of Agricultural Economics' and the YARA's experts consulted on this issue in 2012, and consequently the IFA revised the applied methodology. So, as an outcome of this consultation the IFA's data for the years 2007 onwards are not higher than the official statistic of Hungary.

Indirect N₂O emissions due to volatilization

NH₃-N and NO_x-N losses are calculated in compliance to the obligations under UNECE/CLRTAP. To estimate the NH₃ and NO_x emissions from 3.D methodologies of 2019 EMEP/EEA Guidebook were applied.

Indirect N₂O emissions due to leaching and run off

In response to a recommendation of the in-country review, conducted in 2016, this section was supplemented with verification information.

Verification of the applied data

Dry and humid/ irrigated regions were distinguished using the IPCC definitions, based on climate data and the HCSO's statistics on irrigated areas. Climate data was taken from the HMS's climate database. The HMS fulfill international and national quality standards concerning data collection and processing.

Statistics on irrigated areas was verified using data collection of different institutes and data on areas with irrigation license and drought affected areas.

In the frame of the National Statistical Data Collection Program (OSAP) the HCSO collects data on irrigated areas, annually. The HCSO provides another statistic on irrigated areas based on data collection of the Hungarian General Directorate of Water Management (OVF). Generally, the HCSO (OSAP) data indicates slightly higher irrigated areas (see Figure 5.5.10). However, in the last two years the OVF's data were higher. The HCSO's, OSAP data was used to estimate the proportion of irrigated areas until 2013, and the OVF's data for the years 2014-2019 to avoid underestimation of emissions.




Figure 5.5.10 Irrigated areas and areas with irrigation license, 2003-2019

In Hungary the water use for agricultural purposes must be permitted; statistics on areas with irrigation license is published by the HCSO. Data on irrigated areas was compared to the areas with irrigation license to check the reliability of the applied data. Areas of irrigated lands highly depends on the weather. On average 48% of areas with irrigation license was irrigated in the period between 2003 and 2019. This proportion was the highest in 2015, when 63% of areas with irrigation license was irrigated due to the dry weather. Areas with irrigation license did not exceed the 4.8% of the utilized agricultural areas (Table 5.5.16). In 2020, the HCSO did not conduct a survey on irrigation, so no data are available for 2020. Therefore, indirect N₂O emissions from leaching and run off cannot be significantly underestimated due to underestimation of irrigated areas.

Table 5.5.16 Irrigated areas from different sources and areas with irrigation license in proportion to the utilized agricultural areas (2003-2015)

	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Irrigated areas, HCSO (OSAP)	1.6%	2.4%	2.2%	2.7%	2.6%	2.1%	1.7%	2.1%	2.2%	2.1%	2.2%
Irrigated areas, OVF	1.4%	1.2%	1.6%	2.3%	2.1%	2.8%	2.7%	2.2%	2.3%	2.4%	2.3%
Areas with irrigation license	4.7%	3.7%	3.9%	4.1%	3.6%	4.8%	4.2%	4.1%	4.2%	4.3%	4.1%

Verification with other countries

According to the European field experiments 2-8% or 1-16% of the applied N was lost by leaching (Füleky, 2014). Therefore, the IPCC default value of 30% for $\text{Frac}_{\text{LEACH-H}}$ is probably too high even for countries with higher annual precipitation than Hungary. Thus, accounting lower proportion of N addition than the default 30% in the light of climate data as well as the amount of N applied seems to be reasonable. In spite of this $\text{Frac}_{\text{LEACH-H}}$ was assumed to be 30% in this inventory submission due to lack of country-specific value. Although, taking into account the proportion of humid and irrigated areas the resultant fraction of all N added to/mineralized in managed soils for 2019 and 2020 is 4%, which is in line with the results of the European field experiments.

For the general procedure of the QC see 6.1.5.

5.5.5 Source-specific recalculations

3.D.1 Direct N₂O Emissions from Managed Soils, whole time series

3.D.1.2.a Animal Manure Applied to Soils

Due to the interlinking between the 3.B and 3.D CRF categories through the N-balance, revisions to the 3.B resulted in revised emissions from 3.D.1.2a Animal Manure Applied to Soils and 3.D.2 Indirect N₂O Emissions from Managed Soils for the whole time series. Correction to the livestock number of “heifers 1-2 yrs” for the years 2016-2019, as mentioned in the paragraph related to the CRF Sector 3.A, resulted in some increase in the F_{AM} . The net effect of these recalculations is an insignificant reduction in the emissions from 3.D.1.2.a Animal Manure Applied to Soils over the period BY-2008 and an increase ranging from 0.02 kt CO₂ eq to 8.4 kt CO₂ eq over the period 2010-2019 (Table 5.5.17).

Table 5.5.17 Impact of recalculation on 3.D.1.2.a Animal Manure Applied to Soils

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
BY	966	964	-1.6	-0.2%
1990	886	884	-1.6	-0.2%
1991	825	823	-1.9	-0.2%
1992	708	706	-2.0	-0.3%
1993	621	620	-1.8	-0.3%
1994	559	558	-1.8	-0.3%
1995	553	550	-2.1	-0.4%
1996	539	537	-2.0	-0.4%
1997	519	517	-2.1	-0.4%
1998	530	528	-2.1	-0.4%
1999	538	536	-2.0	-0.4%
2000	541	539	-2.1	-0.4%
2001	531	530	-1.9	-0.4%
2002	540	539	-1.6	-0.3%
2003	541	540	-1.2	-0.2%
2004	524	523	-1.0	-0.2%
2005	498	497	-0.9	-0.2%
2006	489	489	-0.7	-0.1%
2007	487	486	-0.4	-0.1%
2008	471	471	-0.2	0.0%
2009	450	450	0.0	0.0%
2010	450	450	0.2	0.1%
2011	445	446	0.5	0.1%
2012	458	458	0.3	0.1%
2013	463	463	0.6	0.1%
2014	475	478	3.1	0.7%
2015	482	488	5.6	1.2%
2016	470	478	8.4	1.8%
2017	459	465	6.2	1.3%
2018	458	462	3.7	0.8%
2019	461	462	0.7	0.2%

3.D.1.2.c Other Organic Fertilizers Applied to Soils

Data on composted sewage sludge and municipal solid waste (MSW) have been revised to include only use on agricultural land. In the previous submissions, the total amount of composted sewage sludge and MSW was used, which was not in line with the Hungarian regulation, because in Hungary, the quality of compost that can be applied is regulated by law, and application is subject to a permit, so not all compost can be used on agricultural land without restriction.

At the same time, as a result of our improvements in the utilization of agricultural waste in anaerobic digesters, digestate from anaerobic digestion, which was not included in previous submissions, is now included.

Table 5.5.18 summarises the changes in activity data for 3.D.a.2.c by source.

Table 5.5.18 Revision of N-inputs from other organic wastes (kg N)

Year	N-input from							
	Composted sewage sludge		Composted MSW		Digestate		Compost, total	
	2021 submission	2022 submission	2021 submission	2022 submission	2021 submission	2022 submission	2021 submission	2022 submission
BY	400,000	152,000	0	0	0	0	400,000	152,000
1990	400,000	152,000	0	0	0	0	400,000	152,000
1991	400,000	152,000	0	0	0	0	400,000	152,000
1992	400,000	152,000	0	0	0	0	400,000	152,000
1993	400,000	152,000	0	0	0	0	400,000	152,000
1994	400,000	152,000	0	0	0	0	400,000	152,000
1995	560,000	212,800	0	0	0	0	560,000	212,800
1996	580,000	220,400	144,000	54,000	0	0	724,000	274,400
1997	520,000	197,600	152,000	57,000	0	0	672,000	254,600
1998	460,000	174,800	144,000	54,000	0	0	604,000	228,800
1999	640,000	243,200	144,000	54,000	0	0	784,000	297,200
2000	600,000	228,000	136,000	51,000	0	0	736,000	279,000
2001	540,000	205,200	136,000	51,000	0	0	676,000	256,200
2002	740,000	281,200	376,000	141,000	0	0	1,116,000	422,200
2003	1,120,000	425,600	376,000	141,000	0	0	1,496,000	566,600
2004	479,099	182,058	312,000	117,000	0	212,069	791,099	511,127
2005	1,052,454	399,932	328,000	123,000	0	202,159	1,380,454	725,092
2006	858,983	326,414	464,000	174,000	0	255,672	1,322,983	756,086
2007	1,022,822	388,672	512,000	192,000	0	495,489	1,534,822	1,076,161
2008	1,235,719	469,573	680,000	255,000	0	971,158	1,915,719	1,695,731
2009	1,799,460	683,795	720,000	270,000	0	1,454,755	2,519,460	2,408,550
2010	1,644,566	624,935	1,184,000	444,000	0	1,779,796	2,828,566	2,848,731
2011	1,627,390	618,408	1,464,000	549,000	0	2,647,893	3,091,390	3,815,301
2012	1,803,530	685,342	1,462,984	548,619	0	2,190,061	3,266,514	3,424,022
2013	1,865,160	708,761	1,498,643	561,991	0	4,047,153	3,363,803	5,317,905
2014	1,945,066	739,125	1,888,698	708,262	0	3,995,623	3,833,764	5,443,009
2015	1,986,434	754,845	1,844,762	691,786	0	3,872,741	3,831,196	5,319,372
2016	2,036,120	773,726	2,351,752	881,907	0	3,894,543	4,387,872	5,550,175
2017	2,051,801	779,684	2,472,984	927,369	0	4,615,595	4,524,785	6,322,649
2018	1,971,457	749,154	2,483,607	931,353	0	4,275,537	4,455,064	5,956,043
2019	1,831,093	695,815	2,822,440	1,058,415	0	4,019,431	4,653,533	5,773,662

As a result of the revision of activity data, emissions from 3Da2c decreased by -0.5 to 4.4 kt CO₂ eq between BY and 2009, while they increased by 0.1 to 9.2 kt CO₂ eq between 2010 and 2019 (Table 5.5.19).

Table 5.5.19 Impact of recalculation on 3.D.1.2.c Other Organic Fertilizers Applied to Soils

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
BY	1.87	0.7	-1.16	-62%
1990	1.87	0.71	-1.16	-62%
1991	1.87	0.71	-1.16	-62%
1992	1.87	0.71	-1.16	-62%
1993	1.87	0.71	-1.16	-62%
1994	1.87	0.71	-1.16	-62%
1995	2.62	1.00	-1.63	-62%
1996	3.39	1.28	-2.11	-62%
1997	3.15	1.19	-1.95	-62%
1998	2.83	1.07	-1.76	-62%
1999	3.67	1.39	-2.28	-62%
2000	3.45	1.31	-2.14	-62%
2001	3.17	1.20	-1.97	-62%
2002	5.23	1.98	-3.25	-62%
2003	7.01	2.65	-4.35	-62%
2004	3.70	2.39	-1.31	-35%
2005	6.46	3.40	-3.07	-47%
2006	6.20	3.54	-2.65	-43%
2007	7.19	5.04	-2.15	-30%
2008	8.97	7.94	-1.03	-11%
2009	11.80	11.28	-0.52	-4%
2010	13.25	13.34	0.09	1%
2011	14.48	17.87	3.39	23%
2012	15.30	16.03	0.74	5%
2013	15.75	24.90	9.2	58%
2014	17.95	25.5	7.54	42%
2015	17.94	24.9	6.97	39%
2016	20.55	26.0	5.44	26%
2017	21.19	29.6	8.42	40%
2018	20.86	27.9	7.03	34%
2019	21.79	27.0	5.25	24%

3.D.1.3 Urine and Dung Deposited by Grazing Animals

The update of the AWMS data also resulted in a minor change in the N excreted on grazing (F_{PRP}) for the period 2014-2019. Revision of F_{PRP} resulted in changes in the emissions ranging from -6.1 kt CO₂ eq (4.8%) in 2016 to 4.8 kt CO₂ eq (3.6%) in 2019.

3.D.1.4 Crop Residues

The AWMS data update for the period 2014-2019, also resulted in a change in N input from crop residues (F_{CR}) due to a revision of the straw used for bedding. Recalculation to the F_{CR} resulted in changes in the emissions ranging from -0.03 kt CO₂ eq in 2016 to 2.3 kt CO₂ eq in 2019.

Due to revisions in N inputs from the above-mentioned sources-categories 3.D.1 Direct N₂O Emissions from Managed Soils N₂O emissions decreased by 0.5-5.6 kt CO₂ eq between the BY and 2009 and increased by 0.3-13.3 kt CO₂ eq over the period 2010-2019 (Table 5.5.20).

Table 5.5.20 Overall impact of recalculation on 3.D.1 Direct N₂O emissions from Agricultural soils

Year	Submission 2020 [Gg CO ₂ -eq]	Submission 2021 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
BY	4,696	4,693	-2.8	-0.1%
1990	3,474	3,471	-2.7	-0.1%
1991	2,535	2,532	-3.0	-0.1%
1992	2,162	2,159	-3.2	-0.1%
1993	2,023	2,020	-2.9	-0.1%
1994	2,389	2,386	-3.0	-0.1%
1995	2,217	2,213	-3.7	-0.2%
1996	2,260	2,256	-4.1	-0.2%
1997	2,341	2,337	-4.1	-0.2%
1998	2,511	2,507	-3.8	-0.2%
1999	2,509	2,504	-4.3	-0.2%
2000	2,407	2,403	-4.2	-0.2%
2001	2,710	2,706	-3.9	-0.1%
2002	2,711	2,707	-4.9	-0.2%
2003	2,536	2,531	-5.6	-0.2%
2004	2,912	2,909	-2.3	-0.1%
2005	2,695	2,691	-4.0	-0.1%
2006	2,748	2,745	-3.3	-0.1%
2007	2,693	2,690	-2.6	-0.1%
2008	2,883	2,882	-1.2	0.0%
2009	2,616	2,615	-0.5	0.0%
2010	2,553	2,553	0.3	0.0%
2011	2,746	2,750	3.9	0.1%
2012	2,683	2,684	1.0	0.0%
2013	2,978	2,987	9.7	0.3%
2014	3,126	3,135	8.5	0.3%
2015	3,210	3,219	8.5	0.3%
2016	3,472	3,479	7.7	0.2%
2017	3,466	3,479	12.5	0.4%
2018	3,478	3,491	13.3	0.4%
2019	3,461	3,475	13.1	0.4%

3.D.2 Indirect N₂O Emissions from Managed Soils, whole time series

The above-mentioned recalculations in 3.D.1 Direct N₂O Emissions from Managed Soils, which are not detailed again, have also led to changes in 3.D.2 Indirect N₂O Emissions from Managed Soils due to atmospheric deposition and nitrogen leaching and run-off.

The recalculation of NH₃ emissions from 3.D also contributed to the changes in indirect N₂O emissions from managed soils due to atmospheric deposition. The main driver behind the decrease in the emissions in the period 2001-2019 is accounting for NH₃ abatement technologies in solid manure application. NH₃ emission abatement technologies for slurry application have already been accounted for in the previous submissions of the air pollutant as well as the GHG-emission inventory. While the manure NH₃ abatement technologies for solid manure application are reported for the first time in this year's submissions. In Hungary, measures on manure incorporation entered into force in 2001, so these technologies were taken into account in the estimation of NH₃ emissions from this date. The reduced NH₃ emissions resulted in a reduction of indirect N₂O emissions due to atmospheric deposition ranging from 0.09 kt CO₂ eq (0.6%) in 2002 to 15.2 kt CO₂ eq (8.7%) in 2019. In contrast, over the BY and 1990-2000 period, due to the decreased N-loss from the animal manure the emission increased, slightly. This increase is less than 2 kt CO₂ eq.

Changes in emissions from nitrogen leaching and run-off because of the aforementioned revisions are negligible (less than 0.3 kt CO₂ eq).

The net effect of these recalculations are changes in 3.D.2 Indirect N₂O Emissions from Managed Soils ranging from -14.9 kt CO₂ eq in 2019 to 1.9 kt CO₂ eq in the BY (*Table 5.5.21*).

Table 5.5.21 Overall impact of recalculation on 3.D.2 Indirect N₂O Emissions from Managed Soils

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
BY	449	451	1.9	0.4%
1990	360	362	1.7	0.5%
1991	255	256	1.6	0.6%
1992	220	221	1.3	0.6%
1993	202	203	1.2	0.6%
1994	210	212	1.1	0.5%
1995	207	208	1.1	0.5%
1996	207	208	1.1	0.5%
1997	205	206	1.1	0.5%
1998	218	220	1.1	0.5%
1999	219	220	1.2	0.5%
2000	227	228	1.1	0.5%
2001	235	236	0.4	0.2%
2002	245	244	-1.1	-0.4%
2003	243	241	-2.4	-1.0%
2004	251	248	-3.0	-1.2%
2005	228	224	-4.2	-1.9%
2006	235	230	-5.1	-2.2%
2007	241	235	-6.0	-2.5%
2008	221	214	-6.8	-3.1%

Year	Submission 2021 [Gg CO ₂ -eq]	Submission 2022 [Gg CO ₂ -eq]	Difference [Gg CO ₂ - eq]	Percentage change
2009	207	200	-7.4	-3.6%
2010	207	199	-8.2	-4.0%
2011	215	206	-8.5	-4.0%
2012	216	206	-9.7	-4.5%
2013	234	225	-9.5	-4.0%
2014	243	232	-10.9	-4.5%
2015	258	245	-12.4	-4.8%
2016	267	254	-13.4	-5.0%
2017	275	262	-13.6	-4.9%
2018	274	261	-13.7	-5.0%
2019	275	260	-14.9	-5.4%

The overall effect of recalculations for 3.D Agricultural Soils was a decrease ranging up to 8.7 kt CO₂ eq (0.3%) in 2012, except 2013, when emissions increased negligibly.

Table 5.5.22 Overall impact of recalculation on 3.D N₂O Emissions from Agricultural Soils

Year	Submission 2020 [Gg CO ₂ - eq]	Submission 2021 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
BY	5,145	5,144	-0.9	0.0%
1990	3,834	3,833	-1.1	0.0%
1991	2,790	2,788	-1.4	-0.1%
1992	2,382	2,380	-1.8	-0.1%
1993	2,224	2,223	-1.7	-0.1%
1994	2,600	2,598	-1.9	-0.1%
1995	2,423	2,421	-2.7	-0.1%
1996	2,466	2,463	-3.0	-0.1%
1997	2,545	2,542	-3.0	-0.1%
1998	2,729	2,726	-2.7	-0.1%
1999	2,728	2,724	-3.1	-0.1%
2000	2,634	2,631	-3.1	-0.1%
2001	2,945	2,942	-3.5	-0.1%
2002	2,956	2,950	-6.0	-0.2%
2003	2,779	2,771	-8.0	-0.3%
2004	3,162	3,157	-5.3	-0.2%
2005	2,923	2,915	-8.2	-0.3%
2006	2,983	2,975	-8.5	-0.3%
2007	2,933	2,925	-8.6	-0.3%
2008	3,104	3,096	-8.0	-0.3%
2009	2,823	2,815	-8.0	-0.3%
2010	2,760	2,752	-7.9	-0.3%
2011	2,961	2,956	-4.6	-0.2%
2012	2,898	2,890	-8.7	-0.3%
2013	3,212	3,212	0.3	0.0%
2014	3,369	3,367	-2.4	-0.1%
2015	3,468	3,464	-3.9	-0.1%
2016	3,739	3,733	-5.7	-0.2%

Year	Submission 2020 [Gg CO ₂ - eq]	Submission 2021 [Gg CO ₂ - eq]	Difference [Gg CO ₂ -eq]	Percentage change
2017	3,742	3,740	-1.1	0.0%
2018	3,752	3,751	-0.5	0.0%
2019	3,737	3,735	-1.8	0.0%

5.5.6 Planned improvements

See Section 5.1.9

5.6 Prescribed Burning of Savannas (CRF Sector 3.E)

Category 3.E Prescribed Burning of Savannas is not relevant to Hungary therefore notation keys 'NO' is used relating to all associated emissions in CRF Tables.

5.7 Field burning of agricultural residues (CRF Sector 3.F)

5.7.1 Source Category Description

Emitted gases: CH₄, N₂O

Key source: none

In Hungary, the first legislation in order to control field burning of agricultural residues entered into force in 1986. According to the regulation No. 21/1986. (VI. 2.) of the Council of Ministers a burning permit was required from the local authority for crop residue burning. This legislation had been in force until 2001, when the Government Decree No. 21/2001. (II. 14.) was issued. The new decree banned the field burning of agricultural crop residues, unless otherwise provided by law. Plant health emergency was the special exception, when burning of crop residues had been allowed. This Government Decree was amended at the end of 2010. The Government Decree No. 306/2010. (XII.23.) is currently in force, which explicitly ban the field burning of crop residues. According to this, field burning of standing crops and crop residues are prohibited unless otherwise provided by law. The only exception is if there is a plant disease on the agricultural field that can only be eliminated by field burning. In this case the plant protection authority – the county government office – in principle may issue a burn permit. In practice such permits are issued rarely. According to the information and data provided by the plant protection authority burn permits have been issued for only rice lands due to infection of *Pyricularia oryzae* or *Helminthosporium oryzae*.

In line with the legal legislation, it was assumed that field burning of crop residues has been not allowed in Hungary since 1986. According to the estimation of the regional inspectors of the Central (Budapest) Soil and Plant Protection Service, less than 1% of the area sown by crops (i.e., not the entire arable area) is affected by illegal burning (Sári 2003, verbal communication), therefore it was taken into account only between 1985 and 1989, and it was considered as negligible in the period after 1990.

However, field burning due to plant protection reason is practically negligible, to ensure the consistency with the reporting to the UNECE on CRTAP and NECD, emissions from this this source have been reported for the first time in the 2020 submission.

5.7.2 Methodological issues

Until the middle of the 1980s, field burning was quite widespread. In the lack of reliable and quantitative information, it was assumed that the rate of field burning in crop cultivation areas had been gradually decreasing between 1985 and 1989 and was essentially eliminated in 1990.

Accordingly, for the mentioned period between 1985 and 1990 the following values for crops were used as the proportion of biomass burnt on field: $\text{Frac}_{\text{BURN}} = 0.11, 0.09, 0.07, 0.04$ and 0.02 (it meant for all plants produced: $\text{Frac}_{\text{BURN}} = 0.05, 0.04, 0.03, 0.02$ and 0.01). To the emission estimation Equation 2.27 of 2006 IPCC Guidelines was applied. As regards other parameters required for the calculation (dry matter, product/by-product ratio, C to N ratio), the default values given in Table 2.5 and Table 2.6 of 2006 IPCC Guidelines were used.

To estimate the emissions from rice field burning the plant protection authority provided data on the areas for which burn permits were issued for the period 2010-2016. Due to unavailability of data for other years the time series from the BY to 2009 was gap-filled by calculating an average proportion of the rice cropping area affected by plant diseases from the available data. The burnt areas for the years 2017-2020 were estimated similarly.

IPCC default dry matter fraction and residue to crop ration was applied. Fraction burned on fields was estimated as 100% of the biomass available. Default combustion factor (C_f) from the Table 2.6 of the 2006 IPCC GIs was used.

In 2020, 989 ha was burnt In Hungary, which equates to 0.02% of the areas covered by crops in 2020.

Table 5.7.1 Activity data to calculate emissions from 3F field burning due to plant protection reasons

Year	Rice field burnt (ha)
BY	3,985
1990	3,974
1991	2,980
1992	1,656
1993	1,656
1994	1,656
1995	1,325
1996	1,031
1997	726
1998	937
1999	748
2000	1,067
2001	775
2002	696
2003	848
2004	932
2005	882
2006	799
2007	867
2008	839
2009	898
2010	257
2011	882
2012	1,037

Year	Rice field burnt (ha)
2013	1,089
2014	993
2015	932
2016	1,034
2017	916
2018	971
2019	877
2020	989

5.7.3 Uncertainties and time-series consistency

Uncertainty in activity data was assumed as 40%. Uncertainty in the EF as 50%. Uncertainty in the emissions is estimated to be $\pm 51.2\%$.

5.7.4 QA/QC Information

See 6.1.5.

5.7.5 Source-specific recalculations

There was no recalculation.

5.7.6 Planned improvements

There are no further improvements planned.

5.8 Liming (CRF Sector 3.G)

Emitted gases: CO₂

Methods: T1

Emission factors: D

Key sources: none

5.8.1 Source Category Description

Liming is a small source of CO₂ emissions in Hungary, which occur mainly from Limestone CaCO₃. The use of Dolomite CaMg(CO₃)₂ is rather low in Hungary, especially in the last decade. In Hungary the recommended methods for liming are as follows:

- Meliorative liming: chemical improvement of acidic soils, applying 5-15 tones carbonate lime per ha.
- Maintenance liming: low dose liming, improving the efficiency of meliorative liming and preventing the recurrence of soil acidity, with a dose of 1-2 tones carbonate lime per ha.
- Lime fertilization: improvement of the Ca supply of the regularly cultivated soil layer, preventing the development of soil acidity. Recommended doses: 1-2 tons per hectare.

In Hungary before the regime change meliorative liming was subsidized by the state, while maintenance liming and lime fertilization were not promoted. As a result, the maintenance liming and lime fertilization were rather rare activity in the large agricultural co-operatives and state farms. In the 90's, after the liquidation of the large-scale agricultural companies, the regular state subsidization for soil melioration was practically removed, and the meliorative liming decreased significantly. Currently, liming with doses higher than 2 tons per hectare must be licensed with the Soil Conservation Authority (County Governmental Offices dept. of Plant Protection and Soil Conservation under the supervision of the National Food Chain Safety Office). The licenses are valid for five years (i.e., the licensed amount of carbonate lime is allowed to apply to soils within a five-year period). Licenses contain information on the amount and the type of the applied chemical amendment.

The most frequent substances used for the improvement of acidic soils in Hungary are as follows:

- Hard and soft limestone powder.
- Beet potash: the by-product of sugar production (especially at the beginning of the inventory period; currently only one sugar factory is operating in Hungary, thus the amount of beet potash used for soil melioration is low.).
- Bog or lake lime.
- Industrial waste products (e.g., lime sludge).

CRF sector 3.G is a minor source of emissions in Hungary, accounting for 0.1% and 0.01% of the national total emissions in the BY and 2020, respectively. Emissions from 3.G decreased by 96 per cent between the BY and 2020. The bulk of this decrease occurred in the early 90's, reflecting the dramatic drop in the agricultural production and the effect of suspension of state support for soil reclamation. After the period of change in the regime (i.e., the 90s), agricultural production started to recover and the ownership feeling strengthened, promoting the soil conservation activities. In this period the

carbonate lime usage slightly increased, leading to moderately growing emissions until peaking in 2004, when a slight decrease started again. In 2007, the economic downturn resulted in a further reduction in the applied lime and the resulted emissions, and emissions continued to decline until 2019. As Figure 5.8.1 reveals, this remarkably decreasing trend in the recent years can be explained by the significantly increasing trend in the use of 'Other carbonate containing fertilizers' (CRF category 3.H). Trends in emissions from 3.G Liming are shown in Figure 5.8.2.




Figure 5.8.1 Trends in activity data for 3.G and 3.I




Figure 5.8.2 Trends in CO₂ emissions from 3.G Liming

As a result of a recommendation arising from the in-country review conducted in 2017, this chapter was supplemented with detailed information on the activity data used and time-series consistency issues.

5.8.2 Methodological issues

5.8.2.1 Calculation method

Emissions from additions of carbonate limes were estimated using the Equation 11.12 of the 2006 IPCC Guidelines which is the Tier 1 method. Because of the relatively negligible share of liming in the national total GHG emissions the use of simple methodology is reasonable.

5.8.2.2 Activity Data

National usage statistics for carbonate lime is not available for the full timeseries in Hungary, which would be used optimally in accordance with the 2006 IPCC Guidelines, as activity data to determine the CO₂ emissions from additions of carbonate lime. Therefore, method was developed to acquire activity data from statistics providing direct inference on lime application to resolve data gap for the beginning of the inventory period.

Meliorative liming (high dose liming)

For the period 1985-2006 annual statistics on soil melioration by soil type (acidic soils, salt-affected soils, and sandy soils) was used to derive activity data to the emission estimate. For the period 1985-1999 areas of the improved soils were published annually in the statistical pocketbooks of the Hungarian Central Statistical Office. While for the years between 2000 and 2006 detailed statistics was available at the website of the AERI, including the areas as well as the amounts and the agent content of the applied chemical amendments by soil types. Nevertheless, the consistency and the representativity of the time series are ensured, as both institutions used the same data sources (regular national agricultural surveys that cover agricultural enterprises as well as private farms). Therefore, for the beginning of the inventory period experts of the Karcag Research Institute of the University of Debrecen, Centre for Agricultural and Applied Economic Sciences made expert judgment using the available statistics to provide activity data to the emission estimate. The main assumptions of the expert judgment were as follows:

- 67% of meliorated acidic soils were improved using limestone containing amendments.
- 27% of meliorated acidic soils were reclaimed by dolomite.
- 50% of meliorated salt-affected soils were reclaimed with limestone containing amendments.
- Liming was not assumed for sandy soils, as high organic matter containing amendments are added to these soils to increase their fertility, not carbonate containing materials.
- The average doses of lime and dolomite was applied (8 tones CaCO₃ and 7 tones CaMg(CO₃)₂ per hectares).

For the years between 2000 and 2006 the published usage data were taken into account, while the direct relationship between the reclaimed areas and the lime usage provided the basis for the expert

judgment for the previous period (i.e., the overlap technique was used to ensure the time-series consistency). See also Section 5.8.4

Statistics used to derive annual amounts of limestone and dolomite content of liming matters for the period 1985-1999 are provided in Table 5.8.1. Data in bold show the original statistical data to distinguish from estimated amounts shown in italics in the Table.

Table 5.8.1 Statistics and calculated amount of limestone and dolomite use for the period 1985-2006

Year	Area (ha)				Amount of agent (t)		Amount (t)		
	Acidic	Salt-affected	Sandy	Total	Limestone	Dolomite	Limestone	Limestone	Dolomite
					on Acidic soils		on Salt-affected soils	Total	
1985	30,709	3,068	2,718	36,495	164,600	58,040	12,272	176,872	58,040
1986	34,476	3,538	1,507	39,521	184,791	65,160	14,152	198,943	65,160
1987	61,084	4,633	1,224	66,941	327,410	115,449	18,532	345,942	115,449
1988	57,867	4,753	2,095	64,715	310,167	109,369	19,012	329,179	109,369
1989	57,696	1,046	1,197	59,939	309,251	109,045	4,184	313,435	109,045
1990	73,013	2,073	3,025	78,111	391,350	137,995	8,292	399,642	137,995
1991	31,352	202	305	31,859	168,047	59,255	808	168,855	59,255
1992	25,342	569	49	25,960	135,833	47,896	2,276	138,109	47,896
1993	7,555	50	488	8,093	40,495	14,279	200	40,695	14,279
1994	13,580	1,754	0	15,334	72,789	25,666	7,016	79,805	25,666
1995	10,346	100	15	10,461	55,455	19,554	400	55,855	19,554
1996	13,873	0	0	13,873	74,359	26,220	0	74,359	26,220
1997	11,861	82	85	12,028	63,575	22,417	328	63,903	22,417
1998	1,206	5	23	1,234	6,464	2,279	20	6,484	2,279
1999	20	0	0	20	107	38	0	107	38
2000	9,894	266	1,751	11,911	56,283	14,805	231	56,514	14,805
2001	11,173	90	504	11,767	53,613	24,091	445	54,058	24,091
2002	10,097	20	383	10,500	50,715	25,645	145	50,860	25,645
2003	11,309	1,142	3,371	15,822	52,608	14,354	1,713	54,321	14,354
2004	6,443	1,318	2,171	9,932	19,864	14,737	1,954	21,818	14,737
2005	7,310	1,079	1,920	10,310	48,514	3,607	651	49,165	3,607
2006	7,279	1,077	764	9,120	15,660	2,892	1,702	17,362	2,892

No data on land reclamation are available after 2006; hence other sources had to be used to estimate the limestone and dolomite contents of the applied liming matters. In Hungary liming with higher doses of lime than 2 tons per hectare must be licensed by the Soil Conservation Authority (County Governmental offices dept. of Plant Protection and Soil conservation, National Food Chain Safety Office). Since 2007 the NFCSO has recorded the content of the issued license in an electronic database and provided data annually, on the quantity and type of the used liming matters and the affected areas for emission inventory purposes. (For the first year of the data collection the dataset was incomplete, therefore the County offices of the NFCSO collected supplementary data on the agent content of carbonate lime from the hard copies of the issued licenses. Thus, for 2007 only the agent contents of the applied liming matters are available.) Table 5.8.2 shows the quantities of liming material permitted

per year based on the permits issued for liming (please note that the lime permits are for five years and the quantities permitted must be used by the farmer within five years).

Table 5.8.2 Carbonate lime usage for the period 2000-2020

Year	Liming maters							Amount (t)	
	Beet potash	Dolomite	Lime sludge	Hard limestone powder	Calcareous moorland	Bog lime	Soft limestone powder	Total CaCO ₃ content	Total CaMg(CO ₃) ₂
2000	22,505	17,723	NO	48,038	NO	NO	NO	56,514	17,723
2001	36,767	30,170	NO	34,313	NO	10,721	NO	53,732	30,170
2002	55,276	32,000	NO	18,457	NO	4,812	NO	50,715	32,000
2003	62,850	18,884	NO	29,237	NO	833	NO	54,070	18,884
2004	28,552	18,998	NO	9,430	NO	162	NO	21,818	18,998
2005	34,364	5,312	285	42,905	NO	NO	NO	48,969	5,312
2006	31,303	4,228	45	2,898	NO	NO	NO	17,318	4,228
2007	NO	133	NO	22,892	NO	NO	NO	20,603	133
2008	NO	77	NO	12,870	2,239	NO	NO	12,898	77
2009	7,323	1,439	NO	16,319	2,639	NO	713	20,527	1,439
2010	1,157	491	NO	28,231	2,014	NO	450	28,362	491
2011	3,088	807	NO	45,955	2,007	NO	463	46,173	807
2012	1,452	278	NO	20,843	NO	NO	NO	20,527	278
2013	4,197	570	NO	10,321	3,693	NO	NO	13,011	570
2014	1,738	836	NO	31,391	NO	NO	NO	30,690	836
2015	NO	851	NO	21,309	NO	4	NO	20,246	851
2016	60	1,112	NO	11,176	NO	NO	NO	10,648	1,112
2017	6	192	NO	4,297	NO	NO	NO	5,032	192
2018	1,545	60	NO	2,003	NO	NO	NO	2,675	60
2019	NO	NO	NO	1,780	NO	NO	NO	21,210	NO
2020	NO	NO	NO	NO	NO	NO	NO	NO	NO

As mentioned, for the period 2000-2006 very detailed statistics are available on land reclamation which is consistent with the data used for the period 1985-1999 as well as the statistics for the period 2007-2016. Data for the period 2000-2006 are provided in the structure of the statistics for the period 1985-1999 as well as 2000-2017 to demonstrate the time-series consistency. Carbonate containing chemical amendments could also contain various amounts of inert materials, or other non-carbonated ingredients, which was also taking into account in the estimate. The assumed agent contents of liming maters appearing in the statistics are summarized in Table 5.8.3. The CaCO₃ content of liming maters were determined consistently between data from the literature and statistics for the period 2000-2006.

Table 5.8.3 CaCO₃ content of the applied liming maters

Liming materials	CaCO ₃ content
Beet potash	45%
Bog lime	50%
Calcareous moorland	25%
Hard limestone powder	90%
Lime sludge	6%
Soft limestone powder	80%

Lime fertilization/ maintenance liming (low dose liming)

For the period 1985-1999 average doses of limestone and dolomite usage were assumed therefore additional liming as lime fertilization were not estimated. Besides, as mentioned, lime fertilization and maintenance liming were not typical in this period arising from the agricultural subsidization system. For the period 2000-2006 statistics on lime fertilization were available, while for the period 2007 to 2016 expert judgment was applied. The agent content of limestone applied for lime fertilization was assumed to be 53% of liming maters applied for soil reclamation.

Resulted activity data

Since meliorative and maintenance liming are periodic rather than annual activities, besides the liming licenses relates to five-year periods, activity data was calculated as five-year moving average of the statistical and the estimated usage data delineated above to smooth out inter-annual inequities.

5.8.2.3 Emission factors

IPCC default values of 0.12 for limestone, 0.13 for dolomite and 0.20 for urea were used.

5.8.3 Uncertainties and time-series consistency

Uncertainties in the activity data used to estimate emissions from 3.G, were estimated to be $\pm 10\%$. Uncertainties in the emission factors were assumed to be ranging from -20% to 0%. Thus, the estimated combined uncertainties in the emissions for 3.G is $\pm 22\%$.

5.8.4 QA/QC Information

Time series consistency

In response to a recommendation from the in-country review conducted in 2016, this section has been supplemented with further information on addressing the time-series consistency issues arising from the use of different sources of activity data through the time series.

IPCC 2006 Guidelines provide different splicing techniques to ensure time-series consistency. In the case of 3.G the applied methodology to form a complete time series is the overlap technique provided in the Chapter 5.3 of the 2006 IPCC Guidelines.

The areas of carbonate lime usage are available *consistently* for the full period of 1985-2006 from the same data collection. For the period 2000-2006 the areas as well as the amount of the carbonate lime usage are available, therefore the period of 2000-2006 is suitable for overlapping. To resolve data gap relationship observed between the areas and the applied amount (i.e., doses) during the period of overlap was applied. (In accordance with the 2006 IPCC Guidelines, the average of annual doses was used instead of the dose calculated from the total amount and the total areas over the period 2000-2006.)

Data on carbonate lime usage for soil *melioration* is available since 2000. However, data was sourced from different statistics. For the period 2000-2006 melioration statistics, while for the period since 2007 data acquired from liming licenses was applied. As the subject of liming licenses is liming with higher doses than 2 tons per hectare, which is the same as the definition of meliorative liming in the former statistics, consequently the data on meliorative liming are consistent for the full time series.

Data on lime fertilization for the period 2007-2016 are estimated using expert judgement based on the statistics of the previous period.

The following further source specific QA/QC procedures have been carried out:

- The missing data were elaborated by scientific experts from the Karcag Research Institute of University of Debrecen.
- Activity data is checked for plausibility.

5.8.5 Source-specific recalculations

There are no recalculations for this source category in this submission.

5.8.6 Planned improvements

Considering that agricultural CO₂ emissions are of minor importance in Hungary improvements are not planned.

5.9 Urea Application (CRF Sector 3.H)

Emitted gases: CO₂

Methods: T1

Emission factors: D

Key sources: none

5.9.1 Source Category Description

Adding urea to soils during fertilization leads to a loss of CO₂ that was fixed in the industrial production process. Reporting of CO₂ emissions from urea application is a new element of the 2006 IPCC Guidelines. This source category has been introduced because the CO₂ removal from atmosphere during urea manufacturing is estimated under the Industrial Processes and Product Use sector (IPPU).

CRF sectors 3.H is a minor sources of CO₂ emissions in Hungary, accounting for 0.3% of the national total CO₂ emissions in 2020. The overall trend in emissions from 3.H shows an overall decrease of 40% (Figure 5.9.1). CO₂ emissions from 3.H increased at the beginning of the inventory period until they peaked in 1988 and dropped by 82 % to 1996, reflecting the effects of the change in the regime and the suspension of agricultural subsidies. Between 1997 and 2007 emissions moderately increased, reaching a peak in 2007 due to the slightly rising fertilizer use.

The decline in the emission level and the drop in the urea use reflect the impact of the economic downturn in 2008, when the urea prices increased significantly on the world market as well as in Hungary. Besides, Péti Nitrogénművek Ltd., the only fertilizer producer in Hungary, came to a halt of the production due to the uncertain market conditions on 18th of October 2008. The production started again on 26th of February in 2009. (55% of the urea used in Hungary is produced by this company and 45% is import.) The loss of production also contributed to that the urea prices remained at a high level in Hungary in 2009 leading to further decline in urea consumption.

Fertilizer prices increased more than 60% because of the economic recession in 2008. In 2009 the fertilizer prices started to slightly decrease, but it remained at a high level in the first half of 2009, especially in spring, when the demand for fertilizers is the highest. Thus, the fertilizer consumption was significantly lower than in other years, additionally farmers favored the other fertilizers with lower N-content and lower price than the urea which resulted in further decline in the urea use in 2009. Emissions have since increased.




Figure 5.9.1 Trends in the CO₂ emissions from 3.H Urea use

5.9.2 Methodological issues

5.9.2.1 Calculation method

Emissions from CO₂ emissions from urea fertilization were estimated using the Equation 11.23 of the 2006 IPCC Guidelines, which is the basic Tier 1 method. Because of the relatively negligible share of CO₂ in total agricultural GHG emissions the use of simple methodologies is reasonable.

5.9.2.2 Activity Data

Annual consumption of urea was derived from the sales statistics by products reported annually by the AKI. AKI's statistics contain the amount of Urea and other ammonium solutions (UAN) and urea ammonium sulphate (UAS) fertilizers. To calculate CO₂ emissions from urea application the annual activity data was derived as the sum of the amount of urea and the urea contents of UAN and UAS fertilizers (Table 5.9.1).

Table 5.9.1 Urea applied to soils, BY-2020

Year	3.H	
	Solid Urea	Urea in UAN and UAS
	tones	
BY	279,275	33,045
1990	223,511	9,875
1991	104,262	4,705
1992	73,913	4,301
1993	69,907	5,123
1994	61,536	7,291
1995	59,840	9,460
1996	58,906	7,958
1997	60,953	8,267
1998	64,530	11,709
1999	66,621	15,407
2000	64,955	23,191
2001	69,235	24,719
2002	76,284	27,236
2003	87,506	21,576
2004	105,415	18,289
2005	101,268	17,973
2006	102,229	21,930
2007	103,190	25,887
2008	30,114	32,394
2009	27,382	31,692
2010	35,211	27,840
2011	47,913	31,094
2012	41,874	33,850
2013	65,337	42,496
2014	52,618	48,993
2015	75,425	57,450
2016	74,761	59,665
2017	87,758	66,645
2018	74,769	78,460
2019	76,175	81,931
2020	92,969	92,989

5.9.2.3 Emission factors

IPCC default value of 0.20 was used.

5.9.3 Uncertainties and time-series consistency

Uncertainties in the activity data used to estimate emissions from 3.H were estimated to be $\pm 5\%$ for urea. Uncertainties in the emission factors were assumed to be ranging from -20% to 0% . Thus, the estimated combined uncertainties in the emissions for 3.H were $\pm 21\%$.

5.9.4 QA/QC Information

HCSO publishes the nitrogen content of synthetic fertilizers sold, which report is also based on the RIAE's data collection. Consequently, data on the total annual amount of synthetic N fertilizer applied to soils (F_{SN}) under the category 3.D are consistent with data used to estimate emissions for CRF 3.H and 3.I.

Emissions from 3.H were cross-checked with the IPPU sector (CRF 2.B.1), to calculate emissions from urea manufacturing consistently.

5.9.5 Source-specific recalculations

There was no recalculation.

5.9.6 Planned improvements

Considering that agricultural CO₂ emissions are of minor importance in Hungary improvements are not planned.

5.10 Other carbonate containing fertilizers (CRF Sector 3.I)

Emitted gases: CO₂

Methods: T1

Emission factors: D

Key sources: none

5.10.1 Source Category Description

As some types of fertilizers contain liming matters to reduce the soil acidity and improve plant growth, CO₂ emissions from carbonate containing fertilizers has also been reported under the 3.I Other sector to ensure the completeness of the agricultural inventory.

Under the category *3.I Other Carbon containing fertilizers* CO₂ emissions from calcium ammonium nitrate (CAN) are reported. According to the sale statistics CAN fertilizers sold in Hungary are predominantly the so-called 'Pétisó', which is a mixture of ammonium nitrate and very fine dolomite powder (NH₄NO₃ + CaMg (CO₃)₂). This fertilizer is the main product of the 'Nitrogénművek Zrt.' (Information about this fertilizer is available on the website of the producer company:

<https://genezispartner.hu/termek-es-szolgaltatasok/mutragyak/nitrogen-mutragyak/genezis-petiso/>

CRF sectors 3.I is a minor sources of CO₂ emissions in Hungary, accounting for 0.06% and 0.2% of the national total CO₂ emissions in the BY and 2020, respectively. However, this is the only one source-category in the Agriculture sector, from which emissions increased significantly, compared to the BY. CO₂ emissions from 3.I increased by 90 per cent between the BY (48 Gg) and 2020 (97 Gg).

Emissions from carbonate containing fertilizers decreased sharply until 1991, reflecting the effect of suspension of state support of fertilizers. After the period of change in the regime emissions fluctuated annually, depending on the fertilizer's prices. Emissions reached the lowest level at -87% (6 Gg) in 1998. Emissions from other carbonate containing fertilizers grew steadily from 1999. The increase over the period 2008-2017 was largely, due to the increase in the production volume of Pétisó, thanks to investments at the producer company. Figure 5.10.1 shows the trend in CO₂ emissions from 3.I Other carbonate containing fertilizers for the BY and the period 1990 to 2020.




Figure 5.10.1 Trends in CO₂ emissions from 3.1 Other carbonate containing fertilizers

5.10.2 Methodological issues

5.10.2.1 Calculation method

Emissions from carbonate containing fertilizers were estimated using the Equation 11.12 of the 2006 IPCC Guidelines, because carbonate containing fertilizers contains dolomite in Hungary.

5.10.2.2 Activity Data

Annual consumption of fertilizers by fertilizer types such as carbon-containing fertilizers were derived from sales statistics by products reported annually by the Agricultural Economics Research Institute (AERI). AERI's statistics contain the amount of soled CAN fertilizers.

The activity data of the emission estimate was the average dolomite content of the soled CAN fertilizers, which was estimated as 25% of the amount of CAN fertilizers based on the chemical formula of 'Pétisó'.

Activity data used to estimate CO₂ emissions from categories 3.1 are summarized in Table 5.10.1.

Table 5.10.1 Activity data for 3.I

Year	3.I Carbon containing- fertilizers
	CAN fertilizers (tones)
BY	403,704
1990	190,768
1991	55,681
1992	137,037
1993	100,290
1994	71,020
1995	63,094
1996	76,704
1997	54,993
1998	52,513
1999	85,051
2000	126,920
2001	135,283
2002	149,058
2003	143,067
2004	221,375
2005	181,847
2006	202,130
2007	259,592
2008	230,307
2009	323,794
2010	379,529
2011	452,342
2012	580,401
2013	648,440
2014	632,993
2015	734,119
2016	816,018
2017	828,278
2018	769,126
2019	777,275
2020	818,144

5.10.2.3 Emission factors

IPCC default value of 0.13 for dolomite was used, as carbon content of calcium ammonium nitrate (CAN) fertilizers, which is reported here, is dolomite (see also the section above).

5.10.3 Uncertainties and time-series consistency

Uncertainties in the activity data used to estimate emissions from 3.I were estimated to be $\pm 5\%$ for carbon-containing fertilizers. Uncertainties in the emission factors were assumed to be ranging from -20% to 0%. Thus, the estimated combined uncertainties in the emissions for 3.I are $\pm 21\%$, respectively.

5.10.4 QA/QC Information

HCSO publishes the nitrogen content of synthetic fertilizers sold, which report is also based on the RIAE's data collection. Consequently, data on the total annual amount of synthetic N fertilizer applied to soils (F_{SN}) under the category 3.D are consistent with data used to estimate emissions for CRF 3.H and 3.I.

5.10.5 Source-specific recalculations

There was no recalculation.

5.10.6 Planned improvements

Considering that agricultural CO₂ emissions are of minor importance in Hungary improvements are not planned.

5.11 References

Agency for Environmental Protection and Technical Services (APAT, Italy) (2008): Italian Greenhouse Gas inventory 1990-2006. National Inventory Report, 2008, Rome.

Apáti, F. (2003): A magyar rizságazat technológiai és ökonómiai elemzése [Technological and economical analysis of the Hungarian Rice sector]. Debreceni Egyetem Agrár és Műszaki Tudományok Centruma [Centre for Agricultural Science University of Debrecen], Acta-Agraria 2003-10, <http://www.date.hu/acta-agraria/2003-10/apati.pdf>

(last version: 20-January-2005) (in Hungarian with English summary)

Borka, G., Németh, Tímea, Kukovics, S. 2010. Analysis of GHG emissions from sheep sector compared with the features of cattle sector. In: Kukovics, S. – Jávorski, A. (edited) (2010): Possibilities for development in sheep sector. JUHINNOV PLATFORM, K-OVI-CAP Bt. – University of Debrecen Centre for Agricultural Sciences, Érd-Debrecen, 553 p. (ISBN 978-963-08-0624-4), pp. 315-354, pp. 532-546. (In Hungarian)

European Commission (2013): Land Use/Land Cover Area Frame Survey (LUCAS) 2009 TOPSOIL data

EEA (European Environment Agency) (2016): EMEP/EEA Air Pollutant Emission Inventory Guidebook, 2016. Technical Report No 21/2016. Copenhagen.

EEA [European Environmental Agency], (2013): CORINE Land Cover database, 2012.

EUROSTAT, Agriculture statistics and Agri-Environmental Indicators, 2015

Fébel, H.Ms. and Gundel, J.: A takarmányozás és a környezetvédelem kapcsolata. [Connection between nutrition and environmental protection]. Állattenyésztés és Takarmányozás. 2007. 56:427-456. (In Hungarian, with English summary)

Füleky, Gy (2014): Nitrate movement in the soils of Hungarian field trials. Hungarian Agricultural Research. Vol. 23, No. 4. pp. 4-13. http://www.agrarlapok.hu/sites/default/files/har_2014-4_teljes.pdf

[Haenel et al. \(2018\): Gaseous and particulate emissions from German agriculture 1990 – 2016 \(RMD\), Thünen Report 57, p. 274](#)

HAENEL H-D, DÄMMGEN U (2009a) Consistent time series of data to model volatile solids and nitrogen excretions of poultry. 3a. Broilers. Fattening procedures and animal properties. Landbauforsch, 59(1), pp. 61-86

HCSO [Hungarian Central Statistical Office] (2021): <https://www.ksh.hu/docs/hun/xftp/idoszaki/mezo/2020/index.html#azagriumabrutthozzadottrtk41tadta2020ban>
HCSO [Hungarian Central Statistical Office] (2021b): https://www.ksh.hu/stadat_files/mez/hu/mez0008.html

HMS (2021): Hungary's Informative Inventory Report 2021. Submission under the UNECE Convention on Long-range Transboundary Air Pollution and Directive (EU) 2016/2284 on the reduction of national emissions of certain atmospheric pollutants. Reports, DRAFT, Budapest

IPCC (Intergovernmental Panel on Climate Change) (2006): 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

<http://www.ipcc-nggip.iges.or.jp/public/2006gl/>

Kovács, A (2013): Kutatási jelentés ÜHG kalkulációkhoz Állattenyésztési ágazatokra 1985-2011 időszakra [Project report on GHG emission estimates on livestock for the period 1985-2011] On behalf of Hungarian Meteorological Service. (in Hungarian).

Laczka, É. and Soós, L. (2003): Some Characteristics of the Hungarian Agriculture in the 1990s. Hungarian Statistical Review, Special number 8. 2003. pp. 3-19.
http://www.ksh.hu/statszemle_archive/2003/2003_K8/2003_K8_003.pdf

Laczka, É. (2007): A Magyar mezőgazdaság az EU-csatlakozás körüli években, 2000-2005. [Hungarian Agriculture in the years of the European Union Accession, 2000-2005.] Hungarian Statistical Review, Vol. 85. No. 1 pp. 5-20. (In Hungarian, with English summary.)
http://www.ksh.hu/statszemle_archive/2007/2007_01/2007_01_001.pdf

Magyar Takarmánykódex Bizottság [Hungarian Nutrition Codex Commission] (2004): Magyar Takarmánykódex II. [Hungarian Nutrition Codex II.] 535 p. (in Hungarian)

Mészáros, Gy. (2005): Tanácsadási füzetek a nemzeti vidékfejlesztési terv intézkedéseihez. I-II. A környezetterhelés csökkentési lehetőségei. [Guidebooks to the implementation of the National Rural Development Plan I.-II. Potential to reduce the environmental pressure] (in Hungarian) Institute of Agricultural Engineering, Gödöllő

Mészáros, Gy. (2015): Expert judgement

Monori, I. (2015): Expert judgement, personal communication.

Pazsiczki, I. et al (2006): Trágyatárolással összefüggő technikai ismeretek a Nitrát Direktívában (ND) megfogalmazott kritériumok alapján és elvárások figyelembevételével. [Technological knowledge referring to manure management in light of requirements of Nitrates Directive] Institute of Agricultural Engineering, Gödöllő. On behalf of Ministry of Agriculture and Rural Development (in Hungarian)

Ráki, Z. (2003): Az állattartás épületkapacitása, kapacitáskihasználása és a nagyobb telepek műszaki állapota [Building capacity, capacity utilization of animal management and the technical status of larger farms]. Budapest. (unpublished, in Hungarian)

Sári, D. (2003) Expert judgement, verbal communication.

Soliva, R. (2006): Report to the attention of IPCC about the data set and calculation method used to estimate methane formation from enteric fermentation of agricultural livestock population and manure management in Swiss agriculture. Institute of Animal Science

Animal Nutrition ETH Zurich Switzerland.

Systemexpert Tanácsadó Kft. [Systemexpert Consulting LTD] (2007): A 2007-es magyarországi üvegházgáz-leltár IPCC és UNFCCC követelmények szerinti felülvizsgálata és a felhasznált módszertan áttekintése. [Review of the 2007 greenhouse gas inventory of Hungary according to the requirements of the IPCC and UNFCCC and of the applied methodology] Budapest. 27 p. (unpublished, in Hungarian)

Tóth G., Hengl. T. Hermann T., Kocsis M., Tóth B., Makó A., Berényi Üveges J. 2015. Magyarország mezőgazdasági területeinek talajtulajdonság-térképei (Soil property maps of the agricultural land of Hungary)

EUR 27539; doi 10.2788/673710

<http://publications.jrc.ec.europa.eu/repository/bitstream/JRC98136/lb-na-27539-hu-n.pdf>

http://airterkep.nebih.gov.hu/gis_portal/talajvedelem/kiadv.htm

Tóth P., HCSO (2004). Expert judgement, verbal communication.

Velthof, G.L. (2007). Development and application of the integrated nitrogen model

MITERRA-EUROPE

Zsembeli et. al. (2011): Determination of country-specific values of the crop residue rate, dry matter content and Nitrogen content of crop residue of oilseed rape and sunflower. (Project report) On behalf of Hungarian Meteorological Service

6 Land-Use, Land-Use Change and Forestry (CRF sector 4)

6.1 Overview of sector

The greenhouse gas inventory of the Land-use, Land-use Change and Forestry (LULUCF) sector covers both CO₂ emissions and removals due to gains and losses in the relevant carbon pools of the predefined six IPCC land-use categories and non-CO₂ emissions from biomass burning and disturbance of soils associated with land-use conversions.

Table 6.1.1. below reports all comments from the most recent Annual Review Report (ARR; unedited draft at the time of the submission of this NIR) and how they have been addressed.

Table 6.1.1. Comments in draft ARR of the annual review in 2021 and how they have been addressed in this NIR.

ID of comment in draft ARR	How the comment has been addressed
L.1	The activity data was corrected where necessary; see the text in NIR sections 6.7.2.3 and 6.8.2
L.2	We provide an extended version of NIR Table 6.5.2; see also the text in NIR section 6.5.1
L.3	The development of the uncertainty analyses is in progress for the forestry sector, Hungary can provide updated results later. Concerning the non-forest categories of the LULUCF sector, we report the uncertainty analysis for the first time in section 6.11.
L.4	We have found the source of these minor errors and they have been corrected in this submission.
L.5	As agreed with the ERT, we continue to apply a 20-year transition period, and there is no need to recalculate the time series. Rather, the text is modified to include the statement that Hungary reports zero carbon stock changes for the years before 1985. We also note here that this also means that the emission estimates for the years 1985-2004 might include bias, therefore, we continue to use the shading of the bars for these years on the respective charts. See sections 6.1.1 and 6.3.2.
L.6	See updated NIR section 6.5.1 and 6.5.2.
L.7	(see above)
L.8	The notation key was changed from “NO” to “NE” for the DOM (DW and LI) and mineral soils pools for forest land remaining forest land in CRF table 4.A
L.9	The figures in tables 6.5.3 and 6.5.11 and CRF table 4.A for category 4.A.2 are now consistent and address the problem that occurred in the underlying database for inventory year 2017 (i.e. which resulted in some figures for 2017 in NIR table 6.5.11 showing a slight increase from the figures in the previous year).
L.10	Hungary corrected the necessary AD and updated the emission and removal estimates and the NIR (Table 6.7.3, section 6.7.2).
L.11	Following the recommendation by the ARR, Hungary now reports on carbon stock changes in lands under peat extraction (i.e., lands for its for which the standard land-use categories based on the 2006 IPCC Guidelines (e.g. peat extraction and flooded land remaining flooded land) are not applicable, for instance the mineral soils carbon stock changes under wetlands remaining wetlands with grass vegetation) under “other wetlands” with a notification in the documentation box, together with a clear explanation in the section 6.8.2 of the NIR, where in the CRF tables the emissions from those lands are reported.
L.12	The source of the erroneous high emission factor value was identified, the correct value from Hahn, 1984) was implemented, and the entire time series of emissions was recalculated. See NIR section 6.8.2.1.
L.13	
L.14	
L.15	In section 6.5.1.1, we provide information on our forest-related databases as well as on the question of change our data source to the Hungarian National Forest Inventory.
L.16	We have corrected the necessary values and table 6.5.5 (P. 379 NIR 2021) and CRF Table

ID of comment in draft ARR	How the comment has been addressed
	4A include the same values.
L.17	Hungary continued its efforts to increase the accuracy of the carbon stock change estimates for mineral soils. The description of the estimation process for soil carbon stock changes in NIR section 6.4.1 is updated so that, to increase transparency, the SOC used to estimate these changes for the different land-use transition categories are now included in Table 6.4.2 of the NIR. The revision of the SOC values and the associated SOC-change values due to land-use change is under way and is planned to be implemented in the next submission. See also NIR section 6.6.2.3.
L.18	See comments for L.12 and L.13 above, and revised section 6.8.2.1 of the NIR.
L.19	Hungary improved text of section 6.4.3 of the NIR to increase transparency.

6.1.1 Emission trends

We firstly note that the reported area of the non-forest land-use conversion categories, which is a key input for the estimation of emissions and removals and which should include areas under conversion for the (default) period of 20 years, excludes areas that were converted before 1985. This is because no estimates are available before 1985 with regard to either the nature or the direction of trends of conversion areas before that year. Therefore, data for these years could only be generated using assumptions applied in simple mathematical models such as extrapolating the trend of later conversions backwards or keeping conversion rates in 1985 constant beforehand. However, there is no guarantee that any of these assumptions would yield realistic time series. The conversion rates generated by these models thus yield emission and removal estimates with very high uncertainties. Therefore, we decided not to generate either area or emission and removal data for the years before 1985. An important consequence of this lack of data before 1985 is that we assume that carbon stock changes due to land-use changes are zero before 1985, and that the reported trend of both the areas and the calculated emissions and removals in the conversion categories before 2005 may involve artefacts. To avoid false conclusions arising from these artefacts, we only analyse trends beginning 2005. However, for reasons of transparency, and to also comply with the request of the ARR of 2017, the data before 2005 are also shown in all relevant graphs in the LULUCF and KP-LULUCF related sections of the NIR, but they are under a *shaded box* (like the one in Figure 6.1.1). We also report on the entire time series (i.e., for all years beginning 1985) of the area as well as emissions and removals of all LULUCF and KP-LULUCF categories in the CRF tables.

The estimates show that the LULUCF sector in Hungary was a net sink in the last decade. In 2022, total removals in the sector corresponded to over 14.5 per cent of total GHG emissions in Hungary (excluding LULUCF), with a rather high variability. This share has substantially increased since the base period because of the large drop of emissions in the non-land-use sectors. In general, Forest Land (FL) is the largest contributor to this net carbon sink, Wetlands (WL) and Settlements (SE) are net sources, whereas Grassland (GL) and Cropland (CL) are net sources in some years and net sinks in others (Figure 6.1.1). Depending on a category, the bulk of the emissions may be in either the “remaining” sub-category (e.g., Forest Land remaining Forest Land, or FL-FL) or in a land converted to another category (e.g., Land converted to Grassland, or L-GL).




Figure 6.1.1 Trends in estimated emissions/removals from the LULUCF sector by land use and land use change subcategories since 1985. (CL-CL: Cropland remaining Cropland; L-CL: Land converted to Cropland; GL-GL: Grassland remaining Grassland; L-GL: Land converted to Grassland; WL-WL: Wetland remaining Wetland; L-WL: Land converted to Wetland; L-SE: Land converted to Settlements; L-OL: Land converted to Other Land, FL-FL: Forest land remaining Forest land; L-FL: Land converted to Forest land. In other sub-categories, no emissions and removals are estimated, see text. The data under the grey box is shaded out for reasons explained above.

Most removals are generated by biomass gains in the FL-FL and the Land converted to Forest Land (L-FL) categories. The net sink in the FL-FL category is mainly due to the fact that the forest area has been increasing, and that the total woody increment of the growing stock in forest land has been larger than the annual harvest for the last three decades (see Figures 6.5.1, 6.5.4 and 11.1).

Although the reported levels of emissions and removals from the non-forest land-use categories are smaller than in the FL, soils in the Cropland and Grassland categories have added to the net sinks in the last decade or so (Figure 6.1.2) which reflects trends in land-use changes (see also section 6.3 and 6.6.4).




Figure 6.1.2. Emissions and removals in the most important land-use change categories due to soil carbon stock changes. (SA: set-aside.) The data under the grey box is shaded out for reasons explained above.

6.1.2 Key categories

Key category analysis is presented in Chapter 1.6, whereas Table 1.2 contains the key categories of the LULUCF sector.

6.1.3 Completeness

In this submission, Hungary reports on carbon stock changes as well as greenhouse gas emissions and removals from Forest Land (CRF 4.A), Cropland (CRF 4.B), Grassland (CRF 4.C), Wetland (CRF 4.D) and Settlements (CRF 4.E). N₂O emissions from N in mineral soils that is mineralized/immobilized in association with loss of soil C are reported in CRF Table 3.D for CL-CL. CRF Table 4(III) reports N₂O emissions for all other land-use and land-use change categories. (Hungary does not report N₂O emissions from immobilization associated with gain of organic matter resulting from change of land-use or management of mineral soils because we apply a combination of Tier1/Tier 2 to estimate carbon stock changes in soils.) N₂O emissions from fertilization in Wetlands (CRF 4(I)) do not occur in Hungary; N₂O emissions from fertilization in other land-use categories, where relevant, are reported under the Agriculture sector (CRF 3). In addition, CO₂ emissions from liming are reported in CRF table 3G, whereas CO, CH₄, N₂O and NO_x emissions from biomass burning are reported in CRF table 4(V). Indirect N₂O emissions due to N mineralisation associated with loss of soil organic matter resulting from change of land-use or management of mineral soils are also reported. (Of all the possible sources of such emissions, only those ones due to leaching/runoff occur.)

Apart from a few cases, emissions from OL (CRF 4F) are not reported because, consistent with the national definition of this category, it contains unmanaged land for which it is only required to report area data, and because rather small areas are sometimes converted to OL, if any. CH₄ emissions from drainage of soils and Wetlands are not reported, either, because this is an optional reporting category, therefore, the notation key NA and NO were used in CRF Table 4 (II).

6.1.4 Recalculations

In this submission, we have implemented a number of recalculations. There are two main reasons for the recalculations. One is related to that, in preparation for the last year of the second commitment period of the Kyoto Protocol, we wanted to make sure that our estimates are as accurate as possible. For this reason, we continued our work, which was started in 2020, to revise all forest-related data and developed a new calculation system that automates all calculations. This means that the *entire* time series data are recalculated for each inventory year by MS Excel formulas instead of assessing figures of the current inventory year only and adding them to the former time series as was the case earlier (except for recalculations when the complete time series were recalculated). Copying the earlier time series data had led to some minor errors because in some cases formulas had not been applied consistently throughout the time series or because of copying errors. Furthermore, the new system automatically generates CRF tables by MS Excel Visual Basic macros and in this way, eliminates ‘copy-paste’ errors.

We also note that, in some cases, e.g., concerning the land-use change matrix, while correct numbers were developed and used for the GHG inventory in previous reporting years, several incorrect values were uploaded to the CRF Reporter system due to technical reasons, and we have made efforts to correct these errors this year. Finally, we now report for all categories and sub-categories (in CRF tables 4.A.1, 4.A.2, 4.B.1 etc.) areas under forest management in two sub-categories, i.e., forest subcompartments and “other” (i.e., other subcompartments), and areas in these CRF tables are now consistent with those in CRF table 4.1.

With respect to the scale of the recalculations, see the category level differences in the respective sections below.

6.1.5 Methodology

The description of the methodological details in the subsequent sections follows the structure of the national inventory reports as outlined in the Appendix to Annex I (Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual greenhouse gas inventories) of Decision 24/CP.19 (Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention).

In estimating emissions and removals, we have been using the 2006 IPCC Guidelines (mainly for reporting under the UNFCCC) and the IPCC 2013 KP Supplement (under the second Commitment Period of the Kyoto Protocol, IPCC 2014a and IPCC 2014b) as a methodological basis since 2016.

In general, we apply Tier 2 methodology with country specific data where any such data is available. We also apply “best estimates”, i.e., we have made use of all data and information that exist within the country in relation to the GHG inventory. In all other cases, we refer to the source of the data applied (i.e., the 2006 IPCC GL for default data).

Due to the complexity of the LULUCF sector, the Hungarian national circumstances and data availability, the methodology of the estimation sometimes differs for the various land-use and land-use change categories, the various pools and emissions. Therefore, this report is completed with methodology matrices at the beginning of the section of each major land-use and land-use change sector which include information on the Tier applied, type of data (e.g., default or country-specific) for emission factors, where and how “included elsewhere”, if any, is treated, other major methodological information, or if a pool (e.g., organic soils) or non-CO₂ emissions are estimated or not. Subsequent methodological sections only provide more detailed methodological description for those pools and non-CO₂ emissions for which estimation has been done.

The estimated emissions and removals are generally only reported in the CRF tables, but they are often also shown in graphs for the major land-use and land-use change categories. In contrast, the time series of major activity data are often reported in this NIR.

6.2 Land-use definitions and classification systems

The land use categories applied in the Hungarian inventory are consistent with the requirements of IPCC (2006), Volume 4, Chapters 2 and 3. Consistent also with the definitions of national land-use categories, the following definitions are used for the various land-use categories:

Forest land is defined as land spanning more than 0.5 hectares with trees higher than five meters and a canopy cover of more than 30 percent, or trees able to reach these thresholds, *in situ*. It does not include land that is predominantly under agricultural or urban land-use but, in addition to areas covered by trees, it includes areas under regeneration, roads and other areas that have no tree cover but are under forest management (see section 6.5.1 for details).

Cropland contains arable lands, kitchen gardens, orchards and vineyards, as well as set-aside croplands. *Arable lands* are any land area under regular cultivation irrespective of the rate or method of soil cultivation and whether the area is under crop production or not due to any reason, such as temporary inland waters or fallow. Areas under tree nurseries (including ornamental and orchard tree nurseries, vineyard nurseries, forest tree nurseries excluding those for the own requirements of forestry companies grown in the forest), permanent crops (e.g., alfalfa and strawberries), herbs and aromatic crops are included. *Kitchen gardens* are areas around residential houses where, in addition to meeting the owners' demand, owners may produce some surplus of low amount which is usually traded. *Orchards* are land under fruit trees and bushes that may include several fruit species (e.g.: apples, pears, cherries, etc.). Included are non-productive orchards and orchards of systematic layout in kitchen gardens if the area is 200 m² or above in case of berries and 400 m² or above in case of fruit trees. *Vineyards* are areas where grapes are planted in equal row width and planting space and include non-productive areas and vineyards in kitchen gardens (e.g., trellises) if grapes are planted in equal row width and planting space, and the size of the area is at least 200 m². *Set-aside cropland* is land that is temporarily unmanaged but not converted to any other land-use.

Grassland is land under grass (artificial planting included) that includes meadows where the production is utilized by cutting, irrespective of whether it is used for grazing sometimes, and pasture that is utilized for grazing irrespective of whether it is cut sometimes or not. Grassland includes areas with trees which are utilized for grazing and set-aside grasslands (i.e., unmanaged grasslands) which are not in use for agricultural purposes.

Wetland includes the wetlands and water bodies as defined by the CORINE land-cover databases and contains inland marshes (low-lying land usually flooded in winter, and more-or-less saturated by water all year round), peat bogs (peat land consisting mainly decomposed moss and vegetable matter), water courses (natural or artificial water-courses including those serving as water drainage) and water bodies (natural or artificial lakes, ponds etc.).

Settlements are areas matching the 'Artificial surfaces' category of the CORINE land-cover database, which comprises the urban and other residential areas, industrial, commercial and transport units, as well as mines, dump and construction sites and artificial non-agricultural vegetated areas.

Other Land includes areas matching the 'Open spaces with little or no vegetation' category of the CORINE land-cover database, which comprises all area that is not included in any of the above categories.

6.3 Land identification

This chapter describes data sources, the national adaptation of the IPCC land-use categories and the resulting land-use change matrices that are used to estimate emissions and removals from the LULUCF sector.

Note that the reported total area of all land-use categories is equal to the total official land area of Hungary as published by the annual HCSO's land-use statistics (i.e., 9,303,266 ha) in each reporting year. (There are very little changes in the annually reported total land area in land-use statistics, which are due to movements of the borders of Hungary based on bilateral agreements with neighbouring countries, and improvements of mapping techniques. To avoid inconsistency, the longer-term average of the annually published total areas is reported for each inventory year in this GHG inventory.)

6.3.1 Methodology of land identification

The development of the annual land-use and land-use change data in Hungary involves elements of both Approach 1 and 2. The system of identification of IPCC land-use categories, which is based on Hungarian statistical categories as well as the main data sources (together with a reference with respect to the Approach it allows for), is reported in Table 6.3.1.

Table 6.3.1. *The system of identification of IPCC land-use categories in Hungary based on national statistical categories and data sources to meet respective data requirements. Acronyms used: HCSO: Hungarian Central Statistical Office; NFD: National Forestry Database; CLC: Corine Land Cover; HLC85: satellite-based land-use change database of FÖMI (Institute of Geodesy, Cartography and Remote Sensing; see text for details). CLC-change surveys were done for 1990-2000, 2000-2006, 2006-2012 and 2012-2018.*

IPCC land-use categories	Category used in the respective database	Primary data sources (and associated Approach)
Forest Land	Land under Forest Management	NFD (maintained by the Forestry Department of the National Land Centre, NFL, Approach 2)
Cropland	Arable land	HCSO's land-use statistics, and General Agricultural Censuses of 1991, 2000 and 2010, Vineyard and Orchard Censuses of 2001 and 2012 (Approach 1)
	Kitchen gardens	
	Orchards	
	Vineyards	
	Set-aside Cropland	HCSO's land-use statistics, General Agricultural Censuses of 1991, 2000 and 2010, CLC-change (Approach 1)
Grassland	Grassland (meadows and pastures)	HCSO's land-use statistics, General Agricultural Censuses of 1991, 2000 and 2010, CLC-change (Approach 1)
	Set-aside Grassland (Unmanaged Grassland)	HCSO's land-use statistics, General Agricultural Censuses of 1991 and 2000, CLC-change (Approach 1)

Wetlands	Wetlands and water bodies	CLC2012, HLC-change1985-1990, CLC-change (Approach 1/2)
Settlements	Artificial surfaces	CLC2012, HLC-change1985-1990, CLC-change (Approach 1/2)
Other Land	all areas not included above	HLC85, CLC90, CLC-change (Approach 1/2)

Areas undergoing land-use change are identified using both national statistics and information on *changes* (between two consecutive surveys) from the Corine Land Cover (CLC) database. For *Forest Land*, the main source of national statistics is the National Forestry Database that includes information on conversions both from and to forests. Forestry statistics are in general available since 1985 and are detailed in Sections 6.5.1-6.5.2. However, for forest land converted to other land-uses, the rate of 1989 is the first data available, and it is used for all years 1985-1989. For mapping the main CLC land-cover categories to the IPCC categories, see Tables 6.3.2-6.3.5.

Table 6.3.2 Classification of the **CLC 1990** land-cover categories into IPCC land-use categories

CLC land-cover categories (Simplified nomenclatures)	IPCC category
311, 312, 313, 324 (310)	Forest land
211, 212, 213, 221, 222 (210, 220)	Cropland
231, 321 (230)	Grassland
111, 112, 121, 122, 123, 124, 131, 132, 133, 141, 142 (100)	Settlements
411, 412, 511, 512 (400, 500)	Wetlands
331, 332, 333 (330)	Other land

For other land-use change categories, it was necessary to use the statistics of the Hungarian Central Statistical Office (HCSO) and other datasets while ensuring consistency between them. The annual statistics on land-use by the HCSO is published (in Hungarian) at its website (http://www.ksh.hu/docs/hun/xstadat/xstadat_eves/i_omf001a.html). The HCSO's land-use statistics record the whole official area of the country divided into the following nine land-use categories: Arable land, Kitchen gardens, Orchards, Vineyards, Grassland, Forest, Reed, Fishpond and Uncultivated land area. Lands not in use for agricultural purposes in the year of the statistic (including set-aside areas (SA), unmanaged grassland (UGL), Settlements and some parts of Wetlands) are reported aggregately as Uncultivated land area. The data acquisition is based on questionnaires, and some land-use data are available since 1853. There have been changes in the methodology since the beginning of the data collection (Keckés, 1997), but the data set was adjusted considering these methodological changes to achieve consistency over time. The adjustment, which was implemented in consultation with the HCSO's expert, included the following steps and assumptions:

- Between 1985 and 1990 the system of landowners and data collection can be considered to be

in steady state, therefore, the annual data was accepted without adjustment.

- Significant changes occurred in land ownership in the period 1990-2000 (i.e., after political changes in the country), making the HCSO statistics less accurate. Therefore, except for orchards and vineyards, the annual dataset for all categories was replaced with values that were interpolations between the statistics of two General Agricultural Censuses of 1991 and 2000. For the vineyards and orchard category, the results of the more detailed and reliable census on vineyards and orchards were accepted instead of the results of the general agricultural census.
- For the period 2000-2010, the annual Cropland and Grassland areas were interpolated between the areas reported for the years of General Agricultural Censuses conducted in 2000 and 2010. Vineyard and Orchard areas were interpolated between the years for which the most detailed survey data are available (2001 and 2012). For the period after 2012, an extrapolation is applied until new data is available.

Concerning the CLC data, the CLC-change 1990-2000 and CLC-change 2000-2006 databases (FÖMI, 2004; FÖMI, 2009a), as well as the CLC-change 2006-2012 (FÖMI, 2014) and CLC-change 2012-2018 databases were supplemented with the database (HCL85 and HLC-changes 1985-1990) on land-use changes of FÖMI (FÖMI, 2009b) that was developed for 1985-1990 using satellite images according to the requirements of the LULUCF GHG inventory, in order to get higher accuracy. Unlike the HCSO reports, the CLC data sets often include statistics on land cover change. It was assumed that, for any period between two CLC assessments (1990, 2000, 2006, 2012 and 2018) and where that was necessary, the difference between the area of the various land-cover categories corresponds to the change in the respective IPCC land-use change category.

For *non-set-aside Cropland (non-SA CL)*, *non-set-aside Grassland (non-SA GL)*, *Settlements (SE)*, *Wetlands (WL)* and *Other Land (OL)*, the above databases directly include the statistics necessary for the land-use change matrix. Separating set-aside lands is necessary for the estimation of carbon stock changes in soils. For this separation, the differences between Category 330 of the CLC databases and the Uncultivated land area category of the HCSO statistics (which include SE and WL together with set-asides) were taken as the *total area* of set-aside agricultural areas. This area (available since 1984) was then split into *set-aside croplands (SA CL)* and *unmanaged grasslands* (i.e., set-aside grassland, SA GL, for both total areas and annual changes) using *expert judgment*.

For land-use changes that were estimated using the CLC database, it was necessary to map the CLC codes to the respective IPCC categories. The CLC code 411 represents inland marshes, which contains 'Low-lying land usually flooded in winter and more-or-less saturated by water all year round' in accordance with the CLC's nomenclature (<http://www.eea.europa.eu/publications/COR0-landcover>). Therefore, conversions listed in Table 6.3.3 below can be the results of the change in total annual precipitation. The analysis of the total annual precipitation supports this assumption, because the total annual precipitation before the acquisition date of the satellite images, on which the CLC2000 data sets are based, highly exceeds the precipitation of the other years.

Table 6.3.3 Areas classified as 'Grassland converted to Wetlands'

Period	CLC code	Explanation
1990-2000	231-411	Pastures converted to inland marshes
	231-512	Pastures converted to water bodies
	321-411	Natural grasslands converted to inland marshes
	321-512	Natural grasslands converted to water bodies
2000-2006		
	231-512	Pastures converted to water bodies

	321-411	Natural grasslands converted to inland marshes
	321-512	Natural grasslands converted to water bodies
2006-2012, 2012-2018		
	231-512	Pastures converted to water bodies

The Settlements converted to Wetland category mainly contains the area of sandpits and gravel pits. The area of these conversions is small, and the emissions from these land-use change conversions are probably zero (Table 6.3.4).

Table 6.3.4. Areas classified as 'Settlements converted to Wetlands'

Period	CLC code	Explanation
1990-2000		
	131-512	Mineral extraction sites converted to water bodies
	133-511	Construction sites converted to water courses
2000-2006		
	131-512	Mineral extraction sites converted to water bodies
	133-512	Construction sites converted to water bodies
2006-2012, 2012-2018		
	131-512	Mineral extraction sites converted to water bodies
	132-512	Dump sites converted to water bodies
	133-512	Construction sites converted to water bodies

Conversions in Table 6.3.5 below also include conversions of water courses and water bodies which are not covered by soil and biomass, therefore, they could not be source of anthropogenic CO₂ emissions.

Table 6.3.5. Areas classified as 'Wetland converted to Settlements'

Period	CLC code	Explanation
1990-2000		
	411-142	Inland marshes converted to sport and leisure facilities
	511-142	Water courses converted to sport and leisure facilities
	412-133	Peat bogs converted to construction sites
2000-2006	511-133	Water courses converted to construction sites
	411-122	Inland marshes converted to road and rail networks and associated land
	411-133	Inland marshes converted to construction sites
	411-142	Inland marshes converted to sport and leisure facilities
	512-122	Water bodies converted to road and rail networks and associated land
	512-131	Water bodies converted to mineral extraction sites
	512-133	Water bodies converted to construction sites

2006-2012, 2012-2018		
	411-122	Inland marshes converted to road and rail networks and associated land
	411-131	Inland marshes converted to Mineral extraction sites
	511-133	Water courses converted to construction sites

The classification of two CLC2018 categories, i.e., 2.4.2 Complex cultivation patterns and 2.4.3 Land principally occupied by agriculture, with significant areas of natural vegetation, represented a challenge as they involve rather small areas that could be classified, e.g., either as cropland or grassland. To estimate the share of these land-uses within the above two categories, data from the Lechner Knowledge Center (formerly FÖMI), obtained from high resolution studies, was used. For example, the shares of converting areas under code 2.4.3 to SE, CL, GL, FL, WL and OL were 0.299, 0.587, 0.009, 0.104, 0.000 and 0.000, respectively.

6.3.2 Land-use statistics and land-use change matrices

For the development of the area of the other land-use and land-use change categories, it was necessary to consider that the area of a category for a year may somewhat be different in the various statistical sources. Inconsistencies mainly occur because of differences in definition (including differences in the definition of land-use vs land cover) and data collection methodology. For example, the HCSO's and CLC forest land data only refer to areas that are covered by trees, whereas the forestry statistics also include areas that are managed in the forestry sector but are not covered by trees (see various sections on forests in Chapters 6.5.1-6.5.2 and 11 for more details).

Also, it was necessary to consider the possible uncertainty of the various data sources that is not only affected e.g. by the methodology used but also the size of the land pieces that are converted annually from one land-use category to another. This size shows large differences, e.g., the total area of the FL-L category is the smallest, whereas that of CL-L is about ten times larger. Even this latter area is only 0.169 percent of the total area. Also considering that the size of the converted units is small, it can generally be stated that the use of remote sensing currently involves rather large uncertainties. Therefore, we mostly rely on data from administrative statistics where possible, which also involves uncertainties but probably less.

To develop the most accurate overall area estimates for the entire AFOLU sector based on all the above, the statistical sources on the various land-use and land-use change categories were treated hierarchically during the compilation of the annual land-use change matrices. This means that, if data from different sources were available, that from a data source deemed more accurate was used. For example, since the forestry statistics are regarded as the most accurate and forests are the most important for the overall accuracy in the LULUCF sector, the forestry statistics were used for all areas related to forests. The hierarchy of the available sources established is as follows:

- Forestry statistics
- CLC data
- HCSO land-use statistics.

In estimating the annual area of the non-forestry land-use *change* categories, first, periodic land-use changes (for the periods 1986-1992, 1992-2000, 2000-2006, 2006-2012 and 2012-2018) were estimated using the CLC database. Annual land-use change values were then calculated from periodic ones using interpolation (until 2018) and extrapolation (after 2018), using also expert judgment, so that, for each land-use category, the net sum of the annual changes in the time period is equal to the difference between the *area* at the end and at the beginning of the given period in the CLC database. (For the forestry data

where annual land-use change data is available, this data was used without any further adjustment.)

The arising net *changes* were in the second step compared with the net *changes* that could be calculated from the HCSO's land-use statistics calculated for the similar periods. The differences between the net change in the HCSO's land-use statistics and the land-cover change datasets were taken to happen due to conversions involving set-aside grassland and cropland areas.

The total area in each land-use change category was calculated applying the default assumption that transitions take place in a period of 20 years, and the areas in the conversion categories are not converted again to any other land-use category during this 20-year transition period. All land in each conversion category is moved to the respective 'remaining' category in the 21th year after the conversion.

Note that, as mentioned in Section 6.1.1 above, the reported area of the land-use conversion categories for 1985-2004, which should include areas under conversion for the (default) period of 20 years, assumes that the area of land that was converted before 1985 is zero as we have no accurate information on conversion areas before 1985. This is equivalent to reporting zero carbon stock changes for the years before 1985, and the trend of both the areas and the calculated emissions and removals before 2005 most probably involve artefacts. Thus, although we report on the entire time series of both areas and emissions and removals in the CRF tables, only data beginning 2005 should typically be analysed on the graphs with LUC information in this report (and this is the reason that data for the years 1985-2004 are shaded out on the respective graphs).

The resulting time series of land-use data are shown in Figure 6.3.1, whereas areas in the 'land remaining' and 'land converted to' subcategories are reported in Figures 6.3.2 and 6.3.3.




Figure 6.3.1. The evolution of the area of the land-use categories. Note that CL and GL categories are split into non-SA and SA subcategories in order that carbon stock changes can include to and from non-SA – SA conversions, and N_2O emissions due to such conversions can also be estimated.




Figure 6.3.2. The evolution of the area of the 'land remaining' categories. Note that the CL and GL categories are split into non-SA and SA sub-categories as on Figure 6.3.1.




Figure 6.3.3. The evolution of the area of the 'land converted to' categories. The data under the grey box is shaded out for reasons explained in section 6.1.1.

The above area estimates of the various land-use and land-use change categories include *all* areas in the land-use change categories for the default transition period of 20 years. For reasons of transparency, Figure 6.3.4 demonstrates *annual* conversion areas, and Table 6.3.6 below reports matrices of the *annual* land-use changes over the period 1985 to 2020. Also, for reasons of transparency, the data in the table

includes forests that were identified as new forests in previous inventories (“found forests”, FF). For further information on FF, see Section 6.5.2.




Figure 6.3.4. Annual area of the ‘land converted to’ categories. The data under the grey box is shaded out for reasons explained in section 6.1.1.

Finally, we note also that land-use and land-use change categories are further subdivided by climate, soil, management and input for the estimation of soil carbon stock changes, see relevant sections, e.g., Section 6.4.1 for more details.

Table 6.3.6. Annual land-use change matrix for the period of 1985-2020.

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
1984	1 741 288	5 289 600	202 647	1 264 900	25 000	251 775	525 612	2 444
FL	1 740 962	95	0	21	0	0	210	0
non-SA CL	2 778	5 280 646	0	5 338	0	0	838	0
SA-CL	8 388	7 640	186 619	0	0	0	0	0
non-SA GL	1 864	4 910	0	1 240 924	16 811	0	391	0
SA-GL	1 515	0	0	0	23 187	298	0	0
WL	16	0	0	0	0	251 745	14	0
SE	118	9	0	117	0	23	525 344	0
OL	0	0	0	0	0	0	0	2 444
1985	1 755 640	5 293 300	186 619	1 246 400	39 997	252 067	526 798	2 444
FL	1 755 314	95	0	21	0	0	210	0
non-SA CL	2 778	5 284 346	0	5 338	0	0	838	0
SA-CL	5 223	540	180 856	0	0	0	0	0
non-SA GL	1 864	4 910	0	1 228 224	11 011	0	391	0
SA-GL	558	0	0	0	39 142	298	0	0
WL	11	0	0	0	0	252 041	14	0
SE	84	9	0	117	0	23	526 563	0
OL	0	0	0	0	0	0	0	2 444
1986	1 765 833	5 289 900	180 856	1 233 700	50 152	252 363	528 018	2 444
FL	1 765 507	95	0	21	0	0	210	0
non-SA CL	2 778	5 281 530	4 753	0	0	0	838	0
SA-CL	5 730	0	175 126	0	0	0	0	0
non-SA GL	1 864	7 366	0	1 222 162	1 918	0	391	0
SA-GL	711	0	0	0	49 143	298	0	0
WL	12	0	0	0	0	252 337	14	0
SE	90	9	0	117	0	23	527 777	0
OL	0	0	0	0	0	0	0	2 444
1987	1 776 691	5 289 000	179 879	1 222 300	51 061	252 658	529 232	2 444
FL	1 776 365	95	0	21	0	0	210	0
non-SA CL	2 778	5 279 930	5 453	0	0	0	838	0
SA-CL	5 774	0	174 105	0	0	0	0	0
non-SA GL	1 864	7 366	0	1 209 762	2 918	0	391	0
SA-GL	724	0	0	0	50 039	298	0	0
WL	12	0	0	0	0	252 632	14	0
SE	90	9	0	117	0	23	528 991	0
OL	0	0	0	0	0	0	0	2 444
1988	1 787 607	5 287 400	179 558	1 209 900	52 957	252 954	530 446	2 444
FL	1 787 281	95	0	21	0	0	210	0
non-SA CL	2 778	5 279 130	4 653	0	0	0	838	0
SA-CL	7 989	0	171 570	0	0	0	0	0
non-SA GL	1 864	7 366	0	1 197 162	3 118	0	391	0
SA-GL	1 395	0	0	0	51 264	298	0	0
WL	15	0	0	0	0	252 924	14	0
SE	114	9	0	117	0	23	530 182	0
OL	0	0	0	0	0	0	0	2 444
1989	1 801 435	5 286 600	176 223	1 197 300	54 382	253 246	531 636	2 444

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
1989	1 801 435	5 286 600	176 223	1 197 300	54 382	253 246	531 636	2 444
FL	1 800 822	180	0	40	0	0	393	0
non-SA CL	2 778	5 280 045	2 938	0	0	0	838	0
SA-CL	7 172	0	169 051	0	0	0	0	0
non-SA GL	1 864	7 366	0	1 185 442	2 237	0	391	0
SA-GL	1 147	0	0	0	52 937	298	0	0
WL	14	0	0	0	0	253 218	14	0
SE	105	9	0	117	0	23	531 381	0
OL	0	0	0	0	0	0	0	2 444
1990	1 813 902	5 287 600	171 989	1 185 600	55 174	253 539	533 017	2 444
FL	1 812 085	454	0	98	0	0	1 266	0
non-SA CL	2 778	5 237 950	30 021	16 013	0	0	838	0
SA-CL	7 354	0	164 636	0	0	0	0	0
GL	1 864	0	0	1 155 932	27 414	0	391	0
SA-GL	1 202	0	0	0	53 672	300	0	0
WL	14	0	0	0	0	253 511	14	0
SE	107	9	0	117	0	23	532 760	0
OL	0	0	0	0	0	0	0	2 444
1991	1 825 404	5 238 413	194 656	1 172 160	81 086	253 834	535 269	2 444
FL	1 823 956	512	0	108	0	0	827	0
non-SA CL	2 778	5 188 704	30 079	16 013	0	0	838	0
SA-CL	8 163	0	186 493	0	0	0	0	0
non-SA GL	1 864	0	0	1 142 482	27 424	0	391	0
SA-GL	1 447	0	0	0	79 339	299	0	0
WL	15	0	0	0	0	253 804	14	0
SE	116	9	0	117	0	23	535 004	0
OL	0	0	0	0	0	0	0	2 444
1992	1 838 339	5 189 225	216 572	1 158 720	106 763	254 127	537 075	2 444
FL	1 838 011	13	0	83	0	0	233	0
non-SA CL	3 349	5 131 728	46 503	6 707	0	0	938	0
SA-CL	3 356	0	213 216	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 138 312	10 550	0	297	0
SA-GL	70	0	0	0	106 095	597	0	1
WL	18	0	0	0	0	254 101	8	0
SE	244	28	0	178	0	16	536 609	0
OL	0	0	0	0	0	0	0	2 444
1993	1 846 338	5 140 038	259 719	1 145 280	116 645	254 714	538 086	2 445
FL	1 846 120	28	0	27	0	0	163	0
non-SA CL	3 349	5 082 525	46 519	6 707	0	0	938	0
SA-CL	1 498	0	258 221	0	0	0	0	0
non-SA GL	984	8 269	0	1 124 928	10 802	0	297	0
SA-GL	0	0	0	0	116 048	597	0	1
WL	13	0	0	0	0	254 693	8	0
SE	176	28	0	178	0	16	537 688	0
OL	0	0	0	0	0	0	0	2 445
1994	1 852 141	5 090 851	304 739	1 131 840	126 850	255 305	539 094	2 446

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
1994	1 852 141	5 090 851	304 739	1 131 840	126 850	255 305	539 094	2 446
FL	1 851 783	53	0	61	0	0	244	0
non-SA CL	3 349	5 033 313	46 543	6 707	0	0	938	0
SA-CL	4 410	0	300 329	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 111 454	10 528	0	297	0
SA-GL	284	0	0	0	125 968	597	0	1
WL	21	0	0	0	0	255 276	8	0
SE	282	28	0	178	0	16	538 590	0
OL	0	0	0	0	0	0	0	2 446
1995	1 861 421	5 041 664	346 872	1 118 400	136 496	255 889	540 077	2 447
FL	1 860 804	140	0	141	0	0	335	0
non-SA CL	3 349	4 984 039	46 631	6 707	0	0	938	0
SA-CL	5 460	0	341 412	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 097 934	10 608	0	297	0
SA-GL	497	0	0	0	135 400	597	0	1
WL	24	0	0	0	0	255 857	8	0
SE	320	28	0	178	0	16	539 536	0
OL	0	0	0	0	0	0	0	2 447
1996	1 871 746	4 992 476	388 043	1 104 960	146 009	256 471	541 114	2 447
FL	1 871 224	192	0	90	0	0	240	0
non-SA CL	3 349	4 934 800	46 682	6 707	0	0	938	0
SA-CL	6 590	0	381 452	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 084 545	10 558	0	297	0
SA-GL	727	0	0	0	144 684	597	0	1
WL	27	0	0	0	0	256 436	8	0
SE	361	28	0	178	0	16	540 531	0
OL	0	0	0	0	0	0	0	2 447
1997	1 883 569	4 943 289	428 135	1 091 520	155 242	257 049	542 014	2 448
FL	1 883 167	89	0	42	0	0	271	0
non-SA CL	3 349	4 885 716	46 579	6 707	0	0	938	0
SA-CL	5 342	0	422 793	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 071 153	10 509	0	297	0
SA-GL	473	0	0	0	154 171	597	0	1
WL	23	0	0	0	0	257 017	8	0
SE	316	28	0	178	0	16	541 477	0
OL	0	0	0	0	0	0	0	2 448
1998	1 893 962	4 894 102	469 372	1 078 080	164 680	257 630	542 991	2 449
FL	1 892 515	98	0	332	0	0	1 017	0
non-SA CL	3 349	4 836 519	46 588	6 707	0	0	938	0
SA-CL	8 725	0	460 647	0	0	0	0	0
non-SA GL	1 291	8 269	0	1 057 423	10 799	0	297	0
SA-GL	1 160	0	0	0	162 920	599	0	1
WL	32	0	0	0	0	257 589	8	0
SE	439	28	0	178	0	16	542 331	0
OL	0	0	0	0	0	0	0	2 449
1999	1 907 512	4 844 915	507 235	1 064 640	173 719	258 204	544 591	2 450

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
1999	1 907 512	4 844 915	507 235	1 064 640	173 719	258 204	544 591	2 450
FL	1 906 326	112	0	93	0	0	982	0
non-SA CL	3 349	4 787 318	46 602	6 707	0	0	938	0
SA-CL	8 602	0	498 633	0	0	0	0	0
non-SA GL	1 598	8 269	0	1 044 222	10 254	0	297	0
SA-GL	828	0	0	0	172 292	598	0	1
WL	32	0	0	0	0	258 164	8	0
SE	434	28	0	178	0	16	543 935	0
OL	0	0	0	0	0	0	0	2 450
2000	1 921 170	4 795 727	545 235	1 051 200	182 546	258 778	546 160	2 451
FL	1 919 873	153	0	251	0	0	893	0
non-SA CL	5 638	4 768 881	17 396	1 847	0	0	1 965	0
SA-CL	8 641	0	536 594	0	0	0	0	0
non-SA GL	2 597	2 985	0	1 020 123	24 957	0	538	0
SA-GL	0	0	0	0	182 058	487	0	0
WL	11	0	0	0	0	258 732	35	0
SE	185	1	0	119	0	30	545 825	0
OL	0	0	0	0	0	0	0	2 451
2001	1 936 944	4 772 020	553 990	1 022 340	207 016	259 249	549 256	2 451
FL	1 935 088	317	0	260	0	0	1 280	0
non-SA CL	5 638	4 747 392	15 177	1 847	0	0	1 965	0
SA-CL	11 167	0	542 823	0	0	0	0	0
non-SA GL	3 057	2 985	0	991 254	24 506	0	538	0
SA-GL	0	0	0	0	206 528	488	0	0
WL	13	0	0	0	0	259 201	35	0
SE	218	1	0	119	0	30	548 888	0
OL	0	0	0	0	0	0	0	2 451
2002	1 955 180	4 750 696	558 000	993 480	231 034	259 719	552 706	2 451
FL	1 953 928	54	0	93	0	0	1 105	0
non-SA CL	5 638	4 726 331	14 915	1 847	0	0	1 965	0
SA-CL	5 775	0	552 225	0	0	0	0	0
GL	2 076	2 985	0	962 561	25 321	0	538	0
SA-GL	0	0	0	0	230 547	487	0	0
WL	8	0	0	0	0	259 675	35	0
SE	148	1	0	119	0	30	552 408	0
OL	0	0	0	0	0	0	0	2 451
2003	1 967 573	4 729 371	567 139	964 620	255 868	260 192	556 052	2 451
FL	1 966 187	109	0	175	0	0	1 103	0
non-SA CL	5 638	4 704 952	14 969	1 847	0	0	1 965	0
SA-CL	2 311	0	564 829	0	0	0	0	0
non-SA GL	1 446	2 985	0	933 619	26 033	0	538	0
SA-GL	0	0	0	0	255 384	484	0	0
WL	6	0	0	0	0	260 151	35	0
SE	103	1	0	119	0	30	555 799	0
OL	0	0	0	0	0	0	0	2 451
2004	1 975 690	4 708 047	579 798	935 760	281 416	260 665	559 440	2 451

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
2004	1 975 690	4 708 047	579 798	935 760	281 416	260 665	559 440	2 451
FL	1 974 831	149	0	56	0	0	654	0
non-SA CL	5 638	4 683 588	15 009	1 847	0	0	1 965	0
SA-CL	1 944	0	577 854	0	0	0	0	0
non-SA GL	1 379	2 985	0	904 878	25 980	0	538	0
SA-GL	0	0	0	0	280 925	491	0	0
WL	6	0	0	0	0	260 624	35	0
SE	98	1	0	119	0	30	559 191	0
OL	0	0	0	0	0	0	0	2 451
2005	1 983 896	4 686 722	592 863	906 900	306 905	261 145	562 384	2 451
FL	1 982 569	116	0	54	0	0	1 157	0
non-SA CL	5 638	4 662 296	14 976	1 847	0	0	1 965	0
SA-CL	8 011	0	584 852	0	0	0	0	0
non-SA GL	2 483	2 985	0	876 020	24 875	0	538	0
SA-GL	0	0	0	0	306 418	487	0	0
WL	10	0	0	0	0	261 099	35	0
SE	177	1	0	119	0	30	562 057	0
OL	0	0	0	0	0	0	0	2 451
2006	1 998 887	4 665 398	599 828	878 040	331 293	261 616	565 753	2 451
FL	1 997 534	91	0	202	0	0	1 061	0
non-SA CL	0	4 659 448	0	4 703	0	0	1 246	0
SA-CL	20 696	0	579 132	0	0	0	0	0
non-SA GL	0	5 183	0	845 052	27 521	0	285	0
SA-GL	964	0	0	0	330 212	117	0	0
WL	0	0	0	0	0	261 611	5	0
SE	0	47	0	188	0	53	565 465	0
OL	0	0	0	0	0	0	0	2 451
2007	2 019 194	4 664 769	579 132	850 145	357 733	261 781	568 062	2 451
FL	2 018 042	380	0	138	0	0	635	0
non-SA CL	0	4 654 838	3 981	4 703	0	0	1 246	0
SA-CL	11 820	0	567 313	0	0	0	0	0
non-SA GL	0	0	0	816 945	32 915	0	285	0
SA-GL	689	0	0	0	356 927	117	0	0
WL	0	0	0	0	0	261 776	5	0
SE	279	47	0	188	0	53	567 495	0
OL	0	0	0	0	0	0	0	2 451
2008	2 030 830	4 655 264	571 294	821 973	389 842	261 946	569 666	2 451
FL	2 029 340	184	0	336	0	0	970	0
non-SA CL	0	4 638 599	10 716	4 703	0	0	1 246	0
SA-CL	9 038	0	562 256	0	0	0	0	0
non-SA GL	0	4 147	0	788 662	28 879	0	285	0
SA-GL	776	0	0	0	388 949	117	0	0
WL	0	0	0	0	0	261 941	5	0
SE	194	47	0	188	0	53	569 184	0
OL	0	0	0	0	0	0	0	2 451
2009	2 039 347	4 642 977	572 972	793 889	417 829	262 111	571 690	2 451

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
2009	2 039 347	4 642 977	572 972	793 889	417 829	262 111	571 690	2 451
FL	2 036 995	670	0	526	0	0	1 155	0
non-SA CL	0	4 626 477	10 551	4 703	0	0	1 246	0
SA-CL	8 150	0	564 822	0	0	0	0	0
non-SA GL	0	2 609	0	760 095	30 901	0	285	0
SA-GL	483	0	0	0	417 229	117	0	0
WL	0	0	0	0	0	262 106	5	0
SE	766	47	0	188	0	53	570 636	0
OL	0	0	0	0	0	0	0	2 451
2010	2 046 394	4 629 803	575 372	765 512	448 130	262 276	573 327	2 451
FL	2 044 792	388	0	140	0	0	1 075	0
non-SA CL	0	4 623 854	0	4 703	0	0	1 246	0
SA-CL	5 035	6 192	564 145	0	0	0	0	0
non-SA GL	0	2 609	0	757 506	5 113	0	285	0
SA-GL	751	0	0	0	447 262	117	0	0
WL	0	0	0	0	0	262 271	5	0
SE	83	47	0	188	0	53	572 955	0
OL	0	0	0	0	0	0	0	2 450
2011	2 050 662	4 633 090	564 145	762 537	452 375	262 441	575 566	2 450
FL	2 048 948	248	0	852	0	0	614	0
non-SA CL	0	4 627 141	0	4 703	0	0	1 246	0
SA-CL	6 296	9 665	548 184	0	0	0	0	0
non-SA GL	0	2 609	0	757 551	2 093	0	285	0
SA-GL	280	0	0	0	451 978	117	0	0
WL	0	0	0	0	0	262 436	5	0
SE	107	47	0	188	0	53	575 171	0
OL	0	0	0	0	0	0	0	2 450
2012	2 055 632	4 639 709	548 184	763 294	454 071	262 606	577 321	2 450
FL	2 054 386	270	0	274	0	0	702	0
non-SA CL	0	4 635 722	2 024	980	0	0	983	0
SA-CL	4 471	0	543 713	0	0	0	0	0
non-SA GL	0	9 336	0	747 935	5 694	0	329	0
SA-GL	270	0	0	14 948	438 853	0	0	0
WL	0	0	0	0	0	262 589	17	0
SE	326	62	0	222	0	182	576 528	0
OL	0	0	0	0	0	0	0	2 450
2013	2 059 453	4 645 389	545 737	764 359	444 547	262 771	578 559	2 450
FL	2 056 663	711	0	448	0	0	1 631	0
non-SA CL	0	4 643 500	0	980	0	0	910	0
SA-CL	3 820	0	541 917	0	0	0	0	0
non-SA GL	0	8 093	0	752 095	3 914	0	256	0
SA-GL	272	0	0	13 632	430 643	0	0	0
WL	83	0	0	0	0	262 670	17	0
SE	593	62	0	222	0	157	577 524	0
OL	0	0	0	0	0	0	0	2 450
2014	2 061 432	4 652 366	541 917	767 378	434 557	262 828	580 338	2 450

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
2014	2 061 432	4 652 366	541 917	767 378	434 557	262 828	580 338	2 450
FL	2 059 732	521	0	413	0	0	766	0
non-SA CL	0	4 650 397	122	980	0	12	854	0
SA-CL	885	0	541 032	0	0	0	0	0
non-SA GL	0	1 759	0	759 251	6 156	12	200	0
SA-GL	71	0	0	7 255	427 232	0	0	0
WL	0	0	0	0	0	262 811	17	0
SE	131	62	0	222	0	157	579 766	0
OL	0	0	0	0	0	0	0	2 450
2015	2 060 819	4 652 739	541 154	768 121	433 387	262 993	581 604	2 450
FL	2 057 990	1 341	0	576	0	0	911	0
non-SA CL	0	4 631 140	19 883	980	0	12	723	0
SA-CL	729	0	540 425	0	0	0	0	0
non-SA GL	0	1 628	0	766 411	0	12	69	0
SA-GL	9	0	0	21 692	411 686	0	0	0
WL	0	0	0	0	0	262 976	17	0
SE	0	62	0	222	0	157	581 162	0
OL	0	0	0	0	0	0	0	2 450
2016	2 058 728	4 634 171	560 309	789 882	411 686	263 158	582 883	2 450
FL	2 054 853	2 367	0	878	0	0	630	0
non-SA CL	0	4 614 630	17 669	980	0	29	864	0
SA-CL	2 307	0	558 001	0	0	0	0	0
non-SA GL	0	1 786	0	787 858	0	29	210	0
SA-GL	75	0	0	20 958	390 653	0	0	0
WL	39	0	0	0	0	263 102	17	0
SE	0	62	0	222	0	157	582 442	0
OL	0	0	0	0	0	0	0	2 450
2017	2 057 273	4 618 844	575 670	810 896	390 653	263 317	584 162	2 450
FL	2 053 896	785	0	1 817	0	0	776	0
non-SA CL	0	4 609 093	7 809	980	0	0	962	0
SA-CL	1 303	0	574 366	0	0	0	0	0
non-SA GL	0	5 143	0	800 614	5 139	0	0	0
SA-GL	38	0	0	2 980	387 635	0	0	0
WL	0	0	0	0	0	263 317	0	0
SE	0	62	0	222	0	157	583 721	0
OL	0	0	0	0	0	0	0	2 450
2018	2 055 237	4 615 082	582 176	806 614	392 774	263 474	585 459	2 450
FL	2 052 606	934	0	687	0	0	1 010	0
non-SA CL	0	4 613 140	0	980	0	0	962	0
SA-CL	1 540	0	580 636	0	0	0	0	0
non-SA GL	0	1 855	0	804 451	0	0	308	0
SA-GL	79	0	0	0	392 696	0	0	0
WL	14	0	0	0	0	263 309	17	135
SE	43	62	0	222	0	189	583 443	1 499
OL	0	0	0	0	0	0	0	2 450
2019	2 054 281	4 615 991	580 636	806 340	392 696	263 497	585 740	4 084

Table 6.3.6. (ctd.)

	FL	non-SA CL	SA-CL	non-SA GL	SA-GL	WL	SE	OL
2019	2 054 281	4 615 991	580 636	806 340	392 696	263 497	585 740	4 084
FL	2 052 778	465	0	492	0	0	546	0
non-SA CL	0	4 614 049	0	980	0	0	962	0
SA-CL	3 832	0	576 804	0	0	0	0	0
non-SA GL	0	1 855	0	804 177	0	0	308	0
SA-GL	305	0	0	1	392 390	0	0	0
WL	0	0	0	0	0	263 346	17	135
SE	89	62	0	222	0	157	583 710	1 499
OL	0	0	0	0	0	0	0	4 084
2020	2 057 004	4 616 431	576 804	805 872	392 390	263 503	585 544	5 718

6.4 Generic methodological steps to estimate emissions and removals

In this section, general methodological description is provided for those methodological elements that are used for more than one land-use and/or land-use change sub-categories. Activity data and emission/removal factors are usually sub-category specific, and their description can be found in the respective sections below.

Concerning pools, the 2006 IPCC Guidelines (its Table 1.1. in Chapter 1 of Volume 4) define carbon pools in a generic manner. In Hungary, pools are defined in a bit different, and more specific way to match them with available data in order that the estimation is as accurate and precise as possible and practicable. These definitions, which are the same in sections both 7 and 11 of the NIR, i.e., those for reporting under the UNFCCC and under the Kyoto Protocol, respectively, are (see also section 6.5.3):

Above-ground biomass (AB): all biomass of living trees, including bark, branches, twigs and leaves that can be found above the height of potential cutting of the stem at its bottom by a chainsaw. This height is usually a few cm above ground; only 1-2 cm for small trees, 5-10 cm for trees at thinning age, and can be 10-30 cm for trees of the age of the final harvest. Note that, in the Hungarian forests, the understory and shrub layers usually have very little biomass.

Below-ground biomass (BB): all living parts of the living trees below that above-mentioned potential cutting height. These parts thus include stumps (up to the heights defined above for AB), coarse roots (i.e., roots thicker than 2 cm) and fine roots.

Litter (LI): all dead plant mass, whether above-ground or below-ground, that is smaller than around 1 cm in diameter (in case of branches) and 2 cm (in case of dead roots). Note that the above diameter thresholds were chosen to match definitions in the quantitative assessment of the carbon content of the litter for the Hungarian forests (Heil et al., 2012).

Deadwood (DW): all dead plant mass, above-ground and below-ground, that is not litter (i.e., above the 1 cm threshold for standing and lying dead trees, including stumps, and 2 cm for dead roots).

Soil (SO): includes the organic carbon in the topsoil down to a depth of 30 cm that excludes deadwood and litter. Inorganic carbon, as well as organic carbon in the below-ground deadwood and litter pools are excluded, but organic carbon in the topsoil layer is included. (Carbon stocks below 30 cm usually do exist in the Hungarian soils, however, consistent with IPCC (2006), they are assumed to be in carbon equilibrium.)

6.4.1 Soil carbon stock changes

Soil carbon stocks may change due to conversion of land to other land-use, conversion of land within a land-use sub-category to another sub-category (non-set-aside to set-aside or vice versa), or change of the management within a specific land-use sub-category over time. These two types of changes are together referred below to occur in soil carbon stock change sub-categories. For the entire land-use sector, the sum of all soil carbon stock changes is estimated using the below formula:

$$\Delta C = \sum_i \Delta C_i$$

where

ΔC = total carbon stock changes in mineral soils due to land conversion or changes of soil management, tC; and

i = a soil carbon stock change category (by land-use change, climate type, soil type, and/or management and input type as appropriate).

Except for those situations that are mentioned explicitly (including the Tier 1 assumption for forests, i.e., assuming no carbon stock change), the estimation of ΔC_i follows the Tier 1/2 approach in which ΔC_i was estimated using the first formula in Equation 2.25 of the 2006 IPCC GL:

$$\Delta C_i = (SOC_0 - SOC_{0-T})_i / D$$

where

ΔC_i = annual area-specific soil organic carbon stock change in a soil carbon stock change sub-category, $tCha^{-1}yr^{-1}$;

SOC_0 = SOC soil organic carbon stock in sub-category i in the inventory year, tC;

SOC_{0-T} = SOC soil organic carbon stock in sub-category i T years prior to the inventory year, tC;

T = number of years over a single inventory time period, yr, T = 1 yr; and

D = default time period for transition between equilibrium SOC values, yr (the default value of 20 years is applied).

For estimating SOC (for both the inventory year and T year before) for a soil carbon stock change sub-category i for any year, the second formula in Equation 2.25 (i.e., Formulation B (in Box 2.1, p. 2.34) of Section 2, Volume 4) of the 2006 IPCC GL was used together with Tier 1/2 data:

$$SOC = A * SOC_{REF} * F_{LU} * F_{MG} * F_I$$

where

A = land area of sub-category i in the inventory year, ha

SOC_{REF} = area-specific reference soil organic carbon, $tCha^{-1}$

F_{LU} , F_{MG} and F_I are specific land-use, management and input stock change factors for which default values are used. (Note that F_{LU} changes from year to year due to land-use change, whereas F_{MG} and F_I change due to changes in management.)

The A land area values are respective values in the land-use change matrix in the inventory year and include all area in the year in a 'remaining' category, or all areas for conversion category i that have been in the category for a maximum period of default length of 20 years (see section 6.2 for details).

According to the above formulas, the SOC values were all estimated from the SOC_{REF} values for which country-specific values were developed in the course of a research project (Zsembeli et. al, 2013). The estimation was based on the Hungarian Soil Protection and Monitoring System (hereafter referred to as TIM). Based on physiographical and soil-ecological units, 877 representative observation points were selected in this system on agricultural lands (Figure 6.4.1). The representative sampling sites had been selected by regional soil experts on the basis of all available soil information (profile descriptions, results of laboratory analysis, long-term field observations, maps, etc.) and on their local experiences. The soil carbon stocks were determined from humus content (H, %) values (Füleky-Filep, 1999), which were measured for the uppermost 30 cm of the soil (in several layers), using a standard conversion value of 0.58 tC/ha / H%:

$$SOC = H\% * 0.58.$$




Figure 6.4.1. Sample plots of the Soil Protection and Monitoring System (TIM) by climate zones.

The soil types of the investigated TIM points and the area of the different soil types were determined using a Hungarian digital soil map (called AGROTOPO map). This map used the Hungarian national soil classification system which classifies soils by genetic types. The 79 national soil types identified from the TIM and the soil map cannot directly be allocated to the IPCC soil types, therefore, they were converted into the soil types of the FAO soil classification system in a dedicated study (Michéli, 1999). Then, the FAO (WRB) soil types were converted to the IPCC soil types using the IPCC soil carbon tools. Altogether, 14 different WRB soil types were identified that corresponded to 3 IPCC soil types, i.e., high activity clay soil, sandy soil and aquic soils.

The sample plots were also classified into the IPCC climate zones by the Hungarian Meteorological Service, using the methodology of the GPG (IPCC, 2003), based on the climate map of the Hungarian Meteorological Service. Note that, as the mean annual temperature in Hungary is about 10 degrees Celsius and that there is no big topographical variation within the country, the average difference between the sites in the cold and the warm IPCC climate zones is only a few tenth of a degree. The warm climate zone in Hungary is mainly situated in a lowland called the Great Hungarian Plain (GHP). The lowlands along the rivers were inundated almost every spring before the river flows were regulated. (The Danube and Tisza, i.e., the main rivers of Hungary that cross the GHP, have been mostly regulated since the 19th century in Hungary.) The regular flooding had resulted in the formation of wetlands and high organic content of the soil, but most of these wetlands were drained one or two centuries ago. The other typical types of vegetation were forest and forest steppe which, centuries ago, also contributed to the evolution of the organic carbon content of the soil. Therefore, the soil organic carbon content varies less due to climate, and more due to soil type, land-use and the history of soil formation.

This needs to be duly considered because, in the methodology applied as described above, SOC changes are calculated as differences in SOC that are estimated as averages of soils under different land-use or management (which could also be interpreted as equilibrium SOC values at different time points, i.e., before and after change in land-use or management). Considering these averages, the Guidelines (Volume 4, Chapter 2.3.3.1, page 2.38 of IPCC, 2006) require that sites included in a soil carbon stock change sub-category should have similar histories and management as well as similar ecological properties (including topographic position and soil physical properties) for both time points.

Unfortunately, the Zsembeli et al. (2013) study only partially complied with these requirements.

To improve accuracy for FL, CL and GL, which are the most important land-use categories with respect to land-use changes, the average SOC values were re-calculated in 2015 based on the following considerations. First, SOC values were separately computed for the major 'from-to' conversion sub-categories for which activity data (i.e., area) is available using data from areas for which the conversion is *possible* (see Table 6.4.2 below). This is necessary because, in Hungary, conversion may not be possible in all sub-categories and soils types. For example, there are some croplands on aquatic soils in Hungary but there are no forests on such soils, therefore, it would be inappropriate to calculate the average SOC for the entire cropland and for the entire forest area and then estimate carbon stock changes for cropland-forest land conversions and vice versa using average SOC values for cropland and forest land. Instead, the average SOC for cropland *for the case of forest-cropland conversions* was calculated for only about 84.6% of all cropland, i.e., only for non-aquic sites that could be used to establish forests and on which forests actually occur, that is, where forest-to-cropland and cropland-to-forest conversions are possible. For similar reasons, for forest-grassland conversions, the SOC content of forest soils were re-calculated considering forest and grassland areas between which conversion is possible.

To check where conversions are possible, we analysed the distribution of the area of the various soil carbon stock change sub-categories by climate and soil type (Table 6.4.1). The data shows that CL and GL have similar distributions, while the distribution of FL somewhat differs from that of the others, and that SOC_{REF} (i.e., SOC for forests) is significantly different for the various soil and climate types. The data suggests that forests occupy the poorer sites within the possible CL-FL and GL-FL conversion paths, and that SOC differences due to conversions should not be calculated along an *average* CL – *average* FL carbon stock change trajectory.

Table 6.4.1. *The relative distribution of area and estimated SOC_{REF} by soil and climate type for those areas that could be converted to and from other land-uses.*

soil type	climate type	Land use type			SOC _{REF} (tCha ⁻¹)
		FL	CL	GL	
HAC	CD	0,54	0,38	0,39	48
	WD	0,36	0,56	0,56	58
sandy	CD	0,10	0,03	0,02	15
	WD	0,01	0,03	0,03	21

Based on the above considerations, we developed corrected conversion-specific SOC values for land-use categories (Table 6.4.2 (a)). An important element of this development is to only consider non-aquatic situations for conversions because the vast majority of conversions occur on soil with typically no groundwater (at least not in the soil layers where roots and the carbon can be found whose change has to be accounted for). Although it is not currently possible to remove all possible bias with this methodology due to lack of data, and further analysis is necessary, we consider the corrected SOC estimates “as far as practicable”. Using these SOC values, it was possible to develop what might be called a specific soil carbon stock change conversion matrix (Table 6.4.2 (b)) which, for FL, CL and GL, includes conversion-specific differences between corrected average SOC values.

For the SOC of land under Settlement, 80% of the SOC of the pre-conversion category was used, based on the Tier 1 approach by the 2006 IPCC GL. For Wetland (which is only involved in very small conversions in terms of area), no methodology is provided in the 2006 GL. For the sake of completeness only, the SOC of Wetland is set to equal to that of Grassland. The changes of SOC to and from Other land is set to zero. Using these considerations, SOC changes involving SE and WL could also developed and are included in the below matrix.

Table 6.4.2. (a) Area-specific SOC of sub-categories considering the possible combination of „from” and „to” land-use (non-SA: non-set-aside; SA: set-aside), and (b) the matrix of the calculated average annual area-specific SOC change values (over $D=20$ years, $tC\ ha^{-1}yr^{-1}$). (Values for OL are not reported as conversions to and from OL are NO. Values for WL are also reported only for the sake of completeness.)

(a)

LU sub-category	SOC, tC/ha
FL	48,1
FL (to-from CL)	51,7
FL (to-from GL)	52,1
FL (other)	48,1
non-SA CL	49,8
non-SA CL non-aquic	40,8
non-SA GL	65,2
non-SA-GL non-aquic	52,2
OL	49,1
SA-CL	61,4
SA-CL non-aquic	50,3
SA-GL	65,2
SA-GL non-aquic	52,2
SE - FL	38,5
SE - non-SA CL	39,9
SE - SA CL	49,1
SE - non-SA GL	52,1
SE - SA GL	52,1
SE - WL	52,1
SE - OL	39,3
WL	65,2

(b)

FROM	TO						
	FL	non-SA CL	SA CL	non-SA GL	SA GL	SE	WL
FL		-11,1	-1,5	2,6	2,6	-9,6	20,3
non-SA CL	11,1		11,8	18,7	18,7	-9,9	18,7
SA CL	1,5	-11,8		7,0	7,0	-12,3	7,0
non-SA GL	-2,6	-18,7	-7,0		0,0	-13,7	0,0
SA GL	-2,6	-18,7	-7,0	0,0		-13,7	0,0
SE	9,6	9,9	12,3	13,7	13,7		13,7
WL	-20,3	-18,7	-7,0	0,0	0,0	-13,7	

Once the conversion-specific average SOC change values are fixed, the calculations of the carbon stock changes in the various conversion sub-categories depend on, and are thus sensitive, to the estimated area of these sub-categories. Uncertainties in this regard are, however, reduced for longer periods (for which the area of the conversions can be more accurately estimated) and for trends.

6.4.2 N₂O emissions from mineral soils

According to the IPCC 2006 Guidelines, N mineralizes in mineral soils when there is loss of soil organic C stocks through land-use change or management practices, and this loss also leads to N₂O emissions. For each land-use and land-use change sub-category and for each year when carbon is lost from mineral soils, these emissions were estimated, for each climate, soil, management and input type as appropriate, using the following Equations of the 2006 IPCC GL:

Equation on page 11.10:

$$N_2O = N_2O-N * 44/28$$

where

N_2O = N₂O emissions, kg N₂O yr⁻¹

N_2O-N = annual direct N₂O-N emissions produced from managed soils, kg N₂O-N yr⁻¹;

Equation 11.1:

$$\text{Direct } N_2O-N = F_{SOM} * EF_1$$

where

F_{SOM} = annual amount of N in mineral soils that is mineralized, in association with loss of soil C from soil organic matter as a result of changes to land-use or management, kg N yr⁻¹ (note that this value is also reported in the CRF table as AD);

EF_1 = emission factor for N₂O emissions from N inputs, kg N₂O-N (kg N input)⁻¹ (the value 0.01 was taken from Table 11.1 of the 2006 IPCC GL); and

Equation 11.8:

$$F_{SOM} = \Delta C_{\text{Mineral}} / R * 1000$$

where

$\Delta C_{\text{Mineral}}$ = average annual loss of soil carbon for each land-use type (LU), tonnes C; and

R = C:N ratio of the soil organic matter. Due to lack of more specific data for the area, the default value of 15 is used for situations involving land-use change from Forest Land or Grassland to Cropland, and the default value of 10 is used for situations involving management changes on CL-CL (page 11.16 of the 2006 IPCC GL).

The method of estimating indirect N₂O emissions from N mineralisation associated with loss of soil organic matter resulting from change of land-use or management on mineral soils due to leaching/runoff is also based on the methodology as suggested by the 2006 IPCC GL, and uses Equation 11.10 (only the part of the equation for leaching/runoff; for each csmi sub-category):

$$\text{Indirect } N_2O_{(L)}-N = F_{SOM} * \text{Frac}_{\text{LEACH-(H)}} * EF_5$$

where

F_{SOM} = as above;

$\text{Frac}_{\text{LEACH-(H)}}$ = fraction of all N added to/mineralised in managed soils in regions where leaching/runoff occurs that is lost through leaching and runoff, kg N (kg of N additions)⁻¹ (the value 0,3 and 0 were

taken from Table 11.3 of the 2006 IPCC GL for cold & dry and warm & dry, respectively);
 EF_5 = emission factor for N_2O emissions from N leaching and runoff, kg N_2O-N (kg N leached and runoff)⁻¹ (the value of 0.0075 for EF_5 was taken from Table 11.1 of the 2006 IPCC GL).

6.4.3 Non- CO_2 emissions from wildfires

Except for slash burning in forests, and in accordance with Government Decrees No. 21/2001(II.14) and No. 306/2010. (XII.23.), the on-site burning of living biomass is prohibited in Hungary. Therefore, the controlled burning of biomass is reported as “not occurring” for Hungary for land-use categories other than Forest Land, and only emissions from wildfires in Forest Land, Cropland and Grassland are reported. Except for forests (where emissions from wildfires are separately reported for FL-FL and L-FL), all such emissions are reported in the “remaining land” categories that include any emissions in the conversion categories.

In estimating these emissions, the Tier 1 method and Equation 2.27 of the 2006 IPCC GL were used as follows:

$$L_{\text{fire}} = A * M_B * C_f * G_{\text{ef}} * 10^{-3}$$

where:

L_{fire} = amount of greenhouse gas emissions from fire, tonnes of each GHG

A = area burnt, ha

M_B = mass of fuel available for combustion, tonnes ha^{-1}

C_f = combustion factor, dimensionless

G_{ef} = greenhouse-gas specific emission factor g (kg.d.m.)⁻¹.

Data on the areas affected by wildfires (A) is derived from the statistics of the National Directorate General for Disaster Management. Data on the areas affected by wildfires has been collected since 1998, but in the system of data collection a methodological change has been introduced in 2007, therefore, more details and complete data are only available since then. To avoid inconsistency arising from the methodological changes, data for the period 1998-2007 had to be adjusted. For the period 1998-2007 the average of the areas affected by wildfires over the period 2007-2009 were applied to the trends. For the period 1985-1997 the average of the emissions since 1998 are reported, due to lack of data. Note that the average for the period from 1998-2021 is not applied for AD. This is because one way to estimate gaps in a time series is to extrapolate activity data and then estimate emissions. However, if the emission factors don't change in time, this method would yield the same result as the one applied by Hungary, which is also in line with the 2006 IPCC GL (Section 5.3 of Volume 1). Hungary prefers the technique of extrapolating emissions rather than AD because, by reporting NE for the AD, it can be more transparent and indicate the fact that there is actually a gap in the activity data time series that causes a gap in the emissions time series that could only be filled using extrapolation.

The amount of M_B and C_f is sector-specific, see the relevant sections for details.

6.4.4 Conversion-related carbon stock changes of the biomass pools

Conversion-related biomass carbon stock changes occur due to land conversions (e.g., converting land to Forest land) but, in some cases, also on land remaining land (e.g., with converting perennial crops to annual ones and when wetland areas are opened for peat extraction). The estimation of carbon stock changes in these cases is done using Equation 2.16 of the 2006 IPCC GL:

$$\Delta C = A_{\text{conv}} * (B_{\text{After}} - B_{\text{Before}}) * CF$$

where

ΔC = carbon stock change, tonnes C yr⁻¹

A_{conv} = the area undergoing conversion, ha yr⁻¹

B_{After} = biomass after the conversion, t biomass d.m. ha⁻¹

B_{Before} = biomass before the conversion, t biomass d.m. ha⁻¹

CF = conversion factor, tonnes C tonnes biomass⁻¹.

Note that “biomass” here means the sum of above-ground and below-ground biomass with the exception of annual croplands where, consistent with the 2006 IPCC GL, only the above-ground biomass is considered for the “from annual cropland” conversions. The method of the estimation of the above variables is described in the relevant sections.

6.4.5 Conversion-related carbon stock changes of the dead organic matter pools

The carbon stock of both litter and deadwood may change due to conversion of land to other land-use. For these pools, carbon stock changes are estimated as the sum of gains and losses using a modified form of Equation 2.23 of the 2006 IPCC GL:

$$\Delta C_{\text{DOM}} = A_{\text{new}} * (C_{\text{new}} - C_{\text{old}}) * T_{\text{old-to-new}}$$

where

ΔC_{DOM} = annual carbon stock changes in litter or deadwood, tCha⁻¹yr⁻¹;

A_{new} = area undergoing conversion from old to new land-use category, ha;

C_{new} = area-specific equilibrium carbon stocks in the new land-use category, tC ha⁻¹;

C_{old} = area-specific equilibrium carbon stocks in the old land-use category, tC ha⁻¹;

$T_{\text{old-to-new}}$ = time period of the transition from old to new land-use category, year. In case $C_{\text{new}} > C_{\text{old}}$, the default value of 20 years is applied, whereas a one-time loss is assumed in the year of the conversion in case $C_{\text{new}} < C_{\text{old}}$.

6.5 Forest Land (CRF sector 4.A)

This section describes forests and forestry in Hungary, as well as methodologies of estimating GHG emissions and removals in the forestry sector. The description covers all information related to reporting under the UNFCCC, but this information is used together with supplementary information in Section 11 where some information is reported in more details. Thus, the consideration of both sections may be necessary to understand methods and data in this section.

6.5.1 Category description

Forest land is managed in Hungary by a well-developed and relatively stringent planning and inspecting system. A general description of this system together with a general description of the Hungarian forests can be found at http://www.nfk.gov.hu/download.php?id_file=40461. Additional information on the Forest Monitoring and Observation System can be found at https://nfk.gov.hu/download.php?id_file=40336.

Forests in Hungary are predominantly managed in units of relatively homogenous tree cover, i.e., stands (or forest sub-compartments), with a mean area of about four ha. The geographical location of all known stands, which are sometimes called sub-compartments, can be identified (in Hungarian) at <http://erdoterkep.nebih.gov.hu/>. Further data and information, mainly in Hungarian, can also be found at www.nfk.gov.hu/erdeszeti_foosztaly. Finally, additional information concerning data, methods and demonstrating specific procedures can be found at http://www.nfk.gov.hu/Supplementary_Information_news_547.

Forest management has a long history in the country, and most forests are more-or-less intensively managed. The area of forests that could be considered as “unmanaged” under the UNFCCC is negligible. There are some forests where no forestry operations have taken place for about two decades to a century. These are called forest reserves, however, their strictly protected so called “core zones” only occupy a few hundred ha in the entire country and even these forests are managed in one way or another as we also consider forest monitoring, inspecting, forest protection, forest tourism and game management as forest management activities, and these may take place even in forest reserves. Therefore, all reported forests of Hungary are considered as managed under the UNFCCC.

Forest land is subdivided into sub-categories under the UNFCCC and the KP according to these provisions. The definitions that are generally applied to identify the areas of these sub-categories, and to estimate emissions and removals in these sub-categories, are the following (see also section 6.2 above):

“**Forest land**”, as also reported in section 6.2 above, is defined in Hungary as land spanning at least 0.5 hectares with forest trees (actually or potentially) higher than five meters at maturity and a canopy cover at maturity of (actually or potentially) more than 30 percent. It does not include land that is predominantly under agricultural or urban land-use. Both the *Forest Land remaining Forest Land, FL-FL* and the *Land converted to forest land (L-FL)* sub-categories include areas covered by trees, as well as areas under regeneration, roads and other areas that are under forest management but are not covered by trees (called hereinafter ‘other’ sub-compartment including areas serving game feeding, forest railways, glades, lanes, timber yards etc.; see Table 6.5.1 below).

“**Afforestation**” (since 1990 under the KP, or AR, which includes “**reforestation**”) is an activity that leads to the conversion of non-forest land to forest land. From a domestic administrative point of view, the afforestation process can take place in a period of 3-15 years, i.e., until the young stand is deemed established itself successfully, depending on tree species and site. Notwithstanding this, the default transition period of 20 years is used in the GHG inventory to include areas in the L-FL category, consistent with the default IPCC methodology. Note that, usually, the area of newly established forests

included in the L-FL category under the UNFCCC, which contains all administered forestations, is different from that of the *AR since 1990* category under the KP in some years, the difference being that AR only includes areas where the requirement of “direct human induced activity” is fully met in our databases (see also Chapter 11). Also, unlike in L-FL (where both forest sub-compartments and other sub-compartments are included, beginning 1985), areas under AR only include forest sub-compartments and only afforestations since 1990.

“**Deforestation**” is an activity (under the KP) that leads to the conversion of forest land to non-forest land. In Hungary, such conversions take place within one year. Partly because of this reason, we account for all emissions due to deforestation in each conversion sub-category in the year of the deforestation itself. All deforested land in each year is registered both in the *forest land converted to other land-uses (FL-L)* category under the UNFCCC and all “deforestation since 1990” (D) category. Note that both area covered by trees and all other land that is moved from Forest land to another LU category are reported under both the UNFCCC and the KP. See Chapter 11 for more details.

Using the above definitions, forest land covers a bit more than one fifth of the terrestrial area of the country. The *total area of land under forest management*, which is considered in the land-use change matrix as forest land area, includes both forest sub-compartments that are at least potentially covered by trees and un-stocked areas (other sub-compartments) that indirectly serve forest management purposes. The area of forest land using this definition was 2,057 thousand ha by the end of 2020. To be consistent with the land-use change matrix, forest land in the CRF tables also equals to this forest land area. However, due to historical reasons and because the area actually covered by trees is smaller than this, we also report other area statistics, too. The *total area of all forest sub-compartments*, which is the potentially stocked area, amounted to 1,942 thousand ha in 2020. (As the biomass carbon stock changes take place in the forest sub-compartments, the correct implied emission factors and m³/ha data should reflect the area of forest sub-compartments.) The *area actually covered by trees*, which is the actually stocked area and which appears in several official Hungarian statistics, amounted to 1,873 thousand ha in 2020. This area is calculated from that of the forest sub-compartments by adjusting for gaps and overlaps in the canopy closure, which are measured during surveys as “canopy closure” (%). Finally, the area under Article 3.4 Forest Management (FM) under the KP amounted to 1,764 kha (forest sub-compartments only) and 1,879 kha (forest+other sub-compartments) in 2020 (**Table 6.5.1**).

Table 6.5.1. *The area of forest land, forest compartments and land covered by trees, as well as that under the FM (ha) over time.*

Reporting year	Total forest area				
	under UNFCCC (forest and other subcompartments)	under UNFCCC (forest subcompartments)	under UNFCCC (area covered by trees)	under forest management (FM) under KP since 1990 (forest+other subcompartments)	under forest management (FM) under KP since 1990 (forest subcompartments)
	ha	ha	ha	ha	ha
1985	1 755 640	1 643 276	1 493 135		
1986	1 765 833	1 650 576	1 505 764		
1987	1 776 691	1 658 660	1 513 582		
1988	1 787 607	1 667 352	1 526 395		
1989	1 801 435	1 674 815	1 530 587		
1990	1 813 902	1 681 467	1 551 375	1 809 420	1 676 985
1991	1 825 404	1 694 546	1 563 585	1 813 234	1 682 377
1992	1 838 339	1 708 804	1 570 750	1 820 590	1 691 054
1993	1 846 338	1 713 763	1 589 760	1 824 425	1 691 850
1994	1 852 141	1 719 146	1 599 669	1 826 627	1 693 631
1995	1 861 421	1 727 223	1 608 811	1 831 479	1 697 280
1996	1 871 746	1 737 818	1 616 716	1 836 250	1 702 323
1997	1 883 569	1 748 358	1 627 588	1 841 379	1 706 167
1998	1 893 962	1 758 645	1 642 288	1 842 572	1 707 255
1999	1 907 512	1 773 247	1 656 399	1 844 558	1 710 293
2000	1 921 170	1 787 372	1 657 827	1 847 774	1 713 976
2001	1 936 944	1 803 922	1 689 401	1 850 521	1 717 499
2002	1 955 180	1 823 377	1 686 740	1 855 065	1 723 261
2003	1 967 573	1 836 429	1 723 805	1 856 427	1 725 282
2004	1 975 690	1 844 988	1 749 246	1 856 373	1 725 672
2005	1 983 896	1 853 642	1 769 988	1 858 397	1 728 143
2006	1 998 887	1 869 452	1 805 802	1 861 680	1 732 245
2007	2 019 194	1 890 866	1 825 953	1 867 793	1 739 465
2008	2 030 830	1 903 360	1 840 171	1 872 208	1 744 739
2009	2 039 347	1 912 917	1 853 170	1 877 205	1 750 776
2010	2 046 394	1 922 108	1 862 002	1 877 986	1 753 700
2011	2 050 662	1 927 702	1 861 033	1 880 612	1 757 653
2012	2 055 632	1 933 604	1 861 691	1 884 422	1 762 393
2013	2 059 453	1 938 139	1 863 679	1 887 546	1 766 231
2014	2 061 432	1 941 016	1 867 062	1 888 103	1 767 687
2015	2 060 819	1 940 720	1 869 325	1 887 572	1 767 473
2016	2 058 728	1 939 342	1 869 189	1 885 745	1 766 359
2017	2 057 273	1 940 052	1 869 213	1 883 771	1 766 550
2018	2 055 237	1 939 175	1 867 479	1 880 678	1 764 617
2019	2 054 281	1 938 544	1 867 558	1 878 365	1 762 628
2020	2 057 004	1 941 579	1 872 778	1 879 069	1 763 644

The total area of forests has changed considerably since 1930, from about 11% to 21% today, because of systematic afforestations of well over 800 thousand ha and very little deforestation. The reason for these area dynamics is mainly that the country is much less forested than other countries (or e.g., the European Union). Also, the Hungarian Forest Law is really rather rigorous, and it is also rather strictly implemented and inspected with respect to deforestations. Forest owners who make a deforestation are obliged to cover the costs of a new afforestation of the same area to offset that deforestation, and these costs are always used to make the afforestation elsewhere. The area of forest sub-compartments deforested (i.e., areas with tree cover before the conversion), which is the main source of emissions due to biomass loss, was under 1000 ha/year until 2015, which is only about 0.07% of the forest area and about 2.3% of the average rate of afforestation. After that year, a three-four times larger area, including area not covered by trees for which emissions from soils are also estimated, has been moved to the other land-use categories annually (**Table 6.5.2**).

Table 6.5.2. The area of, and emissions from, conversion of forest land to other land-use categories. The annual area has been slightly fluctuating, e.g., because of varying rate of highway building. (Emissions from biomass and soils are also reported here for information only. Emissions from other sources are also estimated and reported, see tables below.) AC_t – cumulative area at the end of the given inventory year; AC_{t-1} – cumulative area at the end of the previous inventory year; AA_t – annual area in the given inventory year; AA_{t-20} – annual area 20 years before the given year.

Inventory year	Conversions from FL to other land use					
	Annual area (ha): AA_t		Cumulative area (ha), calculated as: $AC_t = AC_{t-1} + AA_t - AA_{t-20}$		CO ₂ emissions (Gg CO ₂)	
	forest subcompartments	forest and other subcompartments	forest subcompartments	forest and other subcompartments	from biomass	from soils
1985	326	326	326	326	41	0.6
1986	326	326	652	652	41	1.1
1987	326	326	978	978	41	1.7
1988	326	326	1 304	1 304	41	2.2
1989	326	326	1 631	1 631	41	2.8
1990	613	613	2 243	2 243	77	3.8
1991	240	1 817	2 483	4 060	30	7.0
1992	126	1 447	2 609	5 508	16	9.4
1993	329	329	2 937	5 836	41	9.7
1994	218	218	3 156	6 054	27	10.1
1995	358	358	3 514	6 412	45	10.6
1996	346	617	3 859	7 029	43	11.2
1997	522	522	4 381	7 551	65	11.9
1998	402	402	4 783	7 953	50	12.5
1999	395	1 447	5 179	9 400	49	13.9
2000	719	1 187	5 898	10 586	90	15.8
2001	521	1 297	6 419	11 883	65	17.3
2002	638	1 856	7 056	13 740	79	19.9
2003	593	1 252	7 650	14 992	74	21.9
2004	944	1 387	8 593	16 379	117	23.8
2005	411	859	8 678	16 911	51	24.7
2006	509	1 327	8 861	17 912	63	26.5
2007	245	1 353	8 780	18 939	30	27.7
2008	294	1 152	8 748	19 765	27	28.9
2009	450	1 490	8 872	20 929	58	29.8
2010	208	2 351	8 467	22 667	28	31.2
2011	276	1 604	8 503	22 454	46	30.6
2012	782	1 713	9 160	22 720	132	27.8
2013	532	1 246	9 363	23 637	62	28.8
2014	1 891	2 791	11 037	26 210	85	30.4
2015	1 383	1 699	12 061	27 551	117	31.6
2016	2 116	2 829	13 832	29 763	151	34.3
2017	1 711	3 875	15 021	33 117	168	38.0
2018	2 218	3 378	16 836	36 092	217	36.2
2019	2 307	2 631	18 748	37 277	228	37.1
2020	1 191	1 503	19 220	37 593	134	38.1

Some of the above net increase of the forest area is also due to the fact that surveys that are done for forest management planning purposes have identified new forests each year for most of the last three decades, and the area of these forests (called “**found forests**”) was added to the area under forest management. We include them in the Article 3.4 FM category under the KP (see also Chapter11).

The history of these forests is usually unknown. The mean age of the found forests has been above 20 years (see below), but the age of the individual stands at the time of finding them varies (which makes it impossible to accurately estimate the point of time of the conversions). We decided not to estimate the carbon stock changes (typically sinks) of these predominantly young forests *before finding them*, rather, to treat them as an error (i.e., an underestimation) in our estimates for these historical years for the following reasons: (1) to be conservative; (2) because the total area of these forests is not small but not large, either; (3) to avoid estimations with rather high uncertainties (e.g., due to the unknown pre-conversion land-use, no biomass or soil carbon stock before conversion can be estimated); and (4) it is not practicable to deal with rather small and uncertain amounts that had occurred in the past and the assumption about which does not really add to our understanding of the relevant trends of emissions and removals. (Note that, after finding forests, their carbon stock changes are estimated and reported.)

The trends of all forests have been rather steady for the Hungarian forestry since 1985, affecting both area and standing volume. **Figure 6.5.1** demonstrates these statistics for the FL-FL and L-FL categories.




Figure 6.5.1. The area and standing volume of stocked forest on land remaining forest land (FL-FL) and land converted to forest land (L-FL). Note that the values of L-FL are rather small but not zero (see text below).

The data demonstrate that Hungarian forests have had a positive carbon stock change balance for the last almost four decades, mainly due to the above-mentioned large-scale afforestations, but also due to the distribution of the forest area by species as well as over age and site fertility classes, because much less wood has been removed from the forests than the woody increment, and because of the success to manage forests in a sustainable way in other respects, too.

6.5.1.1 Databases used for estimating activity data

Considering methodological issues in general, activity data (i.e. area and growing stock volume statistics) for estimating GHG emissions and removals were taken mainly from the *National Forestry Database* (NFD, see a detailed description about forestry-related databases at http://www.nfk.gov.hu/download.php?id_file=40461). NFD stores data on forest sub-compartments by species, age class and origin (seed or sprout) as well as on other sub-compartments. NFD data have two

main sources: forest planning (modelled growing stock volume and annual increment, see Chapter 6.5.2) and inspection (harvested volume, data related to forest land conversions).

Besides NFD, Hungary has another database on forests containing data of the National Forest Inventory (NFI). Sampling methodology of the NFI was developed from that of the Growth Monitoring System (described also in the document linked above) which means that the same grid of 2.8 x 2.8 km is applied in the NFI, as well. Each point of the grid serves as a southwest ‘corner point’ of a cluster (called ‘tract’). Each tract consists of four corner points of a square being placed 200 m from each other. Sampling is carried out in concentric circular plots (of 3 m, 7 m and 12,6 m in radius) around the corners points. Corresponding the size of the plot, diameter of all trees reaching 7, 12 and 20 cm diameter at breast height is measured. Height of individual trees is assessed from diameter-height curves created for each plot by species. The amount of deadwood is also assessed on the plots (see Chapter 6.5.5.2.2). NFI sampling is periodic, each tract is visited every 10 years. A detailed description of the NFI is available at: <https://nfi.nfk.gov.hu/>.

Hungary chose to apply the NFI only for assessing the deadwood pool whereas living biomass and various area data necessary to estimate carbon stock changes of other pools are calculated from data of the NFD. The reasons for that are the following:

- consistent time series from 1985 is available only from the NFD;
- afforestation and deforestation is registered (in a spatially explicit way) only in the NFD;
- data on forest fire is available only in the NFD (in a spatially explicit way);
- the data updating system of the NFD provides annual data;
- NFD stores sub-compartment polygons ‘wall-to-wall’, thus not only changes of area but also those of the growing stock can be tracked in a spatially explicit way.

However, there are some disadvantages of NFD:

- uncertainty of growing stock volume statistics cannot be calculated directly from field data due to two reasons:
 - o forest planners apply ‘preferential sampling’ in most cases which means that sampling plots and trees are not chosen randomly or systematically;
 - o sampling data are not stored in the NFD thus cannot be used for uncertainty assessment;
- all growing stock and increment statistics are modelled by yield tables and yield tables assume (refer to) specific harvesting regimes – if thinning intensity or frequency is different from that the statistics may be biased.

Thus, uncertainty of growing stock volumes of NFD must be analysed indirectly (see https://nfk.gov.hu/download.php?id_file=40460).

Growing stock volume statistics of the NFI are not based on yield tables and sampling data are stored in the database and can be used for uncertainty assessment. Land conversions may be detected but only with very high uncertainty. The NFI was started only in 2010 whereas the Growth Monitoring System was begun in 1993 (the first 5-year-long sampling cycle ended in 1998).

Recently, Hungary has carried out a project on the question of applying the NFI instead of the NFD data (which would lead to higher removal values in case of the biomass pool). This project led to a stakeholder decision (made by Ministry of Agriculture) that due to the large differences between growing stock statistics of the NFI and those of the NFD further investigations are necessary in order to clarify the reason of these differences.

Most emission/removal factors, e.g., wood density, are available by species or species group as country specific data (arising from appropriate research projects). Some data are taken from literature, while only IPCC default values were available for other factors (see below). Expert judgment is rarely applied, and they are mentioned each time when such an expert judgment is used.

The following sections describe land identification and how carbon stock changes as well as non-CO₂ emissions are estimated.

6.5.2 Land identification

The basis of land identification is the field-based, GIS-supported, annual forest planning process whose main aims are:

- to sample the main stand attributes (such as height, diameter, basal area, age, canopy closure) by species and
- to map sub-compartment polygons and in this way
- to create forest management plans.

As a main role, each sub-compartment is surveyed every 10 years. From sampling data growing stock volume and annual increment are modelled by yield tables. The obtained statistics together with sub-compartment polygons (as `sdo_geometries`) are stored in the NFD that is an Oracle database with spatial extension. Between two sampling occasions stand statistics are updated by forest sub-compartment which means that annual increment is added and annual harvested volume is subtracted from the growing stock year-by-year.

Besides storing the polygons of forest and other sub-compartments polygons of afforested and deforested compartments are saved in a separate database table that also contains the date of land conversion as well as the pre- or post-land-use category. However, such data on land-conversions related to forest land have been available only since 2008. The reason for that is that – just like in most other countries – the forest mapping system in Hungary had been designed and run in the last several decades to be predominantly be able to capture the (entire) *area* that is deemed to be forest according to laws and regulations in effect at any given point in time, and not to capture *changes* of this area. Therefore, any *changes* were only registered before 2008 as a result of different *mechanisms* that were *required by domestic law*, such as subsidizing afforestations, or inspecting the implementation of the Forest Act in effect, i.e., closely monitoring deforestations. The forest mapping system was thus not explicitly designed to capture forest area changes, although it especially aimed at identifying and minimizing deforestations, and in fact effectively resulted in a limited extent of forest area reductions that could not be captured. However, as mentioned above, to meet international reporting requirements, conversions involving forests have been registered at the stand level since 2008.

By the `sdo_geom` package of Oracle it is possible to track temporal and spatial changes of all polygons. In this way, changes in forest land can be tracked in a spatially-explicit way which can be regarded as ‘wall-to-wall’ since total forest land is covered by spatially-explicit polygons.

As mentioned before, our system keeps finding new forests in some years. All FF, irrespective of the sub-category, are subject to land-use change, but since they are found older than 20 years, all FF areas are included in the FL-FL category when they are found.

The vast majority of the increase of the forest area in the FF sub-category over the past decades is due to the following processes and causes:

- natural expansion of the forest area, i.e., natural establishment of stands, sometimes resulting in an increase of the size of an area due to surpassing the thresholds of “forest” as detailed above,
- re-classification of land (i.e., areas of former “croplands”, “grasslands” or “settlements” etc. that were found during a survey to be covered by trees, possibly due to unregistered earlier afforestation, or where the above thresholds of “forests” had been surpassed since the previous survey), and
- geodesic re-measurements of the area of previously existing stands at subsequent surveys.

The identified changes of the total forest area in any inventory year are thus only partly physical and actual increases of the “forest” area but are partly due to the continuous development of the ability of the forest mapping and the land-use inventory in general to identify forests with increasing accuracy.

Most stands in the FF category have been identifiable individually since 2008 (see Table 6.5.3 below and also Figure 6.5.2). A complete assessment of FF with respect to the area and carbon stock changes is presented in Chapter 11.2.2.

The origin of found forests is unknown, however it can be assessed based from stand characteristics. In order to estimate the origin we applied the following procedure:

- 1.) We investigated whether there were any planned forests in the 20-m-wide ‘buffer zone’ of the given found forest. If there were not any we assumed that the given found forest had been planted. Otherwise (i.e., for all other found forests):
- 2.) We investigated whether there were any planned forests of the same age and stand type (see Table 6.5.2.1) like the given found forest in the 20-m-wide buffer zone. If there were any such stands and the area of the found forest was smaller than 0.5 ha then we assumed that the found forest was a result of a geodesic remeasurement. Otherwise:
- 3.) All stands of hybrid poplars and willows were regarded as earlier human-induced conversions. Otherwise:
- 4.) Irrespective their age all black locust stands were regarded as spontaneous forestation if there were any planned black locust stands in the nearby (in the 20-m-wide buffer zone). Note that very young black locust stands may expand by root sprouts. Otherwise:
- 5.) If there were any planned forests of the same type in the buffer zone which were older by at least a given ‘threshold’ number of years than the given found forest we assumed spontaneous expansion of the stand. Otherwise:
- 6.) If there were any planned forests of the same type in the buffer zone which were younger than the given found forest we assumed spontaneous expansion of the stand, too, because we assumed that the neighbouring planned forest (from which the given found forest had spread) had been harvested earlier. Otherwise:
- 7.) We assumed that the given found forest was originated from human-induced afforestation activities. As it can be concluded from the previous ‘steps’ of the procedure no planned stands of the same type occurred nearby these found forests (again, in the 20-m-wide buffer zone) or the planned stand (of the same type) was older than the found forest but the difference between stand ages ‘is not enough’ for supposing spontaneous expansion.


Table 6.5.2.1. *Stand types and threshold ages (i.e., minimum age when spontaneous expansion may occur either from seeds or from sprouts) used for assessing the origin of found forests. For black locust, hybrid poplar and willow stands no such age was applied, see the assessment procedure described in the text.*

Stand type	Threshold age (year)
Sessile oak-hornbeam	40
Pedunculate oak-hornbeam	40
Sessile oak	40
Pedunculate oak	40
Pubescent oak	40
Turkey oak	40
Beech	40
Black locust	
Other hard broadleaves	30
Hornbeam	30
Maple	30
Ash	30
Hybrid poplar and willow	
Other soft broadleaves	30
Indigenous poplar	10
Willow	10
Alder	30
Lime	30
Birch	20
Scots pine	20
Black pine	20
Norway spruce	20
Other coniferous	25

The result of the assessment procedure is summarized in Table 6.5.2.2.

Table 6.5.2.2. The proportion of found forests by type of origin.

Inventory year	Spontaneous expansion	Human-induced conversion	Geodesic remeasurment
	proportion (%)		
2008	34	65	1
2009	46	51	3
2010	44	63	1
2011	35	52	2
2012	46	48	2
2013	50	52	3
2014	45	48	3
2015	42	56	2
2016	55	43	3
2017	44	55	1
2018	0	0	0
2019	0	0	0
2020	49	48	3

*Figure 6.5.2. The distribution of found forests 2008-2020, with an enlarged portion of a specific region directly showing found forest areas for several years/periods.*

One important issue with FF is the need to meet a specific requirement of the 2006 IPCC Guidelines (section 4.2.1.1) when using the stock difference method (Equation 2.5 of Chapter 2, Volume 4). According to this requirement, when estimating biomass carbon stock changes (and we apply this stock change method, see below), “subsequent inventories must also allow identical area coverage in order to get reliable results”. The area of the sub-category FF has been relatively large in some inventory years, i.e., on average about half of annual afforestations (see Figure 11.1), therefore, it must be, and is indeed kept, for each inventory year separate from the area of forests that were known at the beginning of the inventory year. (Note that, from a statistical and database management point of view, only those areas can be regarded as “forest” in any inventory year that the forest inventory system “knows” that they exist.) Under the UNFCCC these FF become parts of the forest land in the next inventory year because the definition of FL-FL starts over in each calendar year.

Table 6.5.3. The algorithm of allocating area to the various land-use and land-use change categories, together with the estimated area (ha) by inventory year for the last several years. In the formulas, t_1 means the beginning of the inventory year (i.e., the end of the preceding year), whereas t_2 means the end of the inventory year. The light-yellow color in some cells of the table (with column title “from DB”) shows that the data in those cells are taken from the database (i.e., they are the result of other calculations), whereas data in white cells are calculated in this table. FL = Forest Land; FL-FL = Forest land remaining forest land; L-FL = Land converted to forest land; FF = found forest; D = deforestation. Δ is used to denote changes of the value of a land-use sub-category between two points of time, or changes estimated by using another methodology if the data is taken from the database. (The table is for demonstration only and may include rounding errors; for precise numbers, and for data by geographical locations, see the respective CRF tables.)

AREA, ha													
FL			new D (forest +other subc.)	new FF (forest subc.)	L-FL (cumulative, forest subc.)						FL-FL (forest+other subc.)		
FL = FL-FL + L-FL + new FF													
t1	t2	Δ	Δ	Δ	t1	new	moved to FL-FL	D on L-FL	Δ	t2	t1	t2, w/o FF	t2, w/ FF
from DB; t2 of prev. year	from DB	t2-t1	from DB	from DB	from DB; t2 of prev. year	from DB	from DB	from DB	new - moved	t1 + Δ	FL - L-FL	FL - L-FL - new FF	FL - L-FL
2 019 194	2 030 830	11 636	1 152	5 567	167 556	7 221	8 484	0	-1 264	166 293	1 851 638	1 858 970	1 864 537
2 030 830	2 039 347	8 517	1 490	6 487	166 293	3 520	7 285	0	-3 765	162 527	1 864 537	1 870 332	1 876 819
2 039 347	2 046 394	7 048	2 351	3 132	162 527	6 267	6 494	0	-227	162 300	1 876 819	1 880 962	1 884 094
2 046 394	2 050 662	4 267	1 604	4 229	162 300	1 641	6 334	0	-4 693	157 607	1 884 094	1 888 825	1 893 054
2 050 662	2 055 632	4 971	1 713	5 522	157 607	1 162	6 739	0	-5 577	152 031	1 893 054	1 898 079	1 903 602
2 055 632	2 059 453	3 821	1 246	4 370	152 031	697	3 045	0	-2 348	149 683	1 903 602	1 905 400	1 909 771
2 059 453	2 061 432	1 978	2 791	3 348	149 683	1 422	2 713	0	-1 292	148 391	1 909 771	1 909 693	1 913 041
2 061 432	2 060 819	-613	1 699	841	148 391	245	3 946	300	-4 001	144 390	1 913 041	1 915 587	1 916 428
2 060 819	2 058 728	-2 091	2 829	496	144 390	242	6 237	387	-6 382	138 009	1 916 428	1 920 223	1 920 720
2 058 728	2 057 273	-1 455	3 875	1 664	138 009	756	7 841	165	-7 249	130 759	1 920 720	1 924 850	1 926 514
2 057 273	2 055 237	-2 037	3 378	0	130 759	1 341	7 681	113	-6 453	124 307	1 926 514	1 930 930	1 930 930
2 055 237	2 054 281	-955	2 631	0	124 307	1 676	8 178	94	-6 596	117 711	1 930 930	1 936 571	1 936 571
2 054 281	2 057 004	2 723	1 503	1 795	117 711	2 431	9 169	158	-6 896	110 815	1 936 571	1 944 394	1 946 190

6.5.3 Methodology to estimate biomass carbon stock changes

For the estimation of carbon stock changes of the **biomass carbon pools**, we apply the **definitions** in Section 6.4. Additionally, we define “**wood volume**”, or simply “**volume**” as the total above-ground volume of trees taller than two meters. Note that, in Hungary, the metric “merchantable volume” is not used, and the total above-ground volume is estimated from measured mean breast-height diameter or basal area, mean tree height and age using country-specific yield tables.

With respect to **carbon stock changes in the biomass pools**, only those of trees are estimated using different approaches for the various categories under the UNFCCC and under the KP as the carbon stock of the understory vegetation is negligible. For FL-FL, the basis for all approaches is that, first, we

calculate carbon *stocks* for *all* forests for year *t* and year *t-1*. From this, the *difference* of carbon stocks is calculated (i.e., we use the stock difference method, Equation 2.5 of Chapter 2, Volume 4 of the 2006 IPCC Guidelines). However, this must be corrected to obtain carbon stock change by excluding the stock of the *newly found* forests. For the formulas applied, see below; they are also highlighted in Table 7.3.6 (and in Table 11.4 for FM under the KP). Carbon stock *changes* are separately calculated for the other forest-related categories, i.e., L-FL using a category-specific method (see section 7.3.2), for FL-L under the UNFCCC (the latter being equal to D under the KP, using the stock difference method, see sections 6.5.2 and 6.5.3) as well as for AR under the KP (also using the stock difference method, see sections 11.3.1.1 and 11.3.1.2).

In Hungary, the stock difference method is used because, due to the nature of the Hungarian forestry statistics, estimates of total above-ground volume of all forests in the country are available annually. The NFD contains aggregate annual statistics on total growing stocks by species and age classes. These statistics are produced by a bottom-up approach, i.e., growing stocks of stands are aggregated by species and age classes. The stock difference method is also used because, due to the relatively high uncertainties related to annual increment, harvest and mortality statistics, this method is deemed more accurate than the gain-loss method. Systematic errors, i.e., most types of bias, are considerably reduced when consecutive growing stock values are deducted to obtain stock changes. (Note, however, that since growing stocks and their changes incorporate the effects of all processes mentioned above, no particular inferences on emissions and removals can be made separately for any of these processes.)

To estimate carbon stock changes of *all forests*, the first part of Equation 2.8 of the 2006 IPCC Guidelines is used:

$$\Delta C_B = (C_{t2} - C_{t1}) / (t_2 - t_1)$$

where

ΔC_B = carbon stock changes of biomass (tonnes C)

C_t = carbon stock at time *t* (tonnes C)

t_1 and t_2 = (final day of) two consecutive years.

To estimate biomass carbon stocks, the second part of Equation 2.8 of the 2006 IPCC Guidelines has been *adapted* to the Hungarian conditions (by excluding BEF term from the equation, see below) in the following form:

$$C_t = [V_t * D] * (1 + R) * CF$$

where:

V_t = growing stock (i.e., above-ground volume of all trees) at time *t* (m³)

D = basic wood density, tonnes m⁻³

R = root-to-shoot ratio, dimensionless

CF = carbon fraction of biomass, tonnes C tonnes biomass⁻¹.

Growing stock in the NFD is calculated from measured mean basal area, height and age of sample trees using yield tables. Volume data of the yield tables are assessed by the volume functions by Kiraly (1978) which are in turn based on the volume tables by Sopp et al. (1974). These functions were derived from field measurement of many felled trees, and directly provide total aboveground volume information which includes the volume of stem, all branches, twigs and bark, i.e., all above-ground parts of the trees. Thus, no (biomass or volume) expansion factors are included in the calculations (i.e., their value is taken to be equal to 1).

Growing stock is estimated during the continuous survey of the forest planning from various stand

measures (such as height, diameter, basal area, and density) depending on species, age, site and stand quality (see Chapters 6.5.1.1. and 6.5.2). More accurate methods are usually used for stands of higher volume stocks. Given that, as mentioned above, growth and yield functions are used in years between surveys to update volume stocks of each stand, volume and carbon stocks are available for each stand and for each inventory year. (Note that, according to Somogyi (2008b), the growth of trees accelerated in Hungary recently. Yield tables cannot be regularly updated, however, this growth increase is equivalent to a small underestimation of the updated volume stock changes and net CO₂ removals, thus, the approach is conservative.)

For inventory years prior to 2008, we only identified the total area of FF, and conducted a sampling of management plans to establish their specific growing stock (m³/ha). From these values, total growing stock of FF could be estimated for each inventory year by using the total FF area. The mean growing stock of all FF that were identified before 2007 is 129.6 m³/ha. For years 2008 and later, we are able to directly estimate the volume stock of all FF land, and thus we are able to report that the average growing stock is 126.9, 122.0, 123.4, 123.2, 127.7, 116.7, 134.3, 150.2, 141.1, 159.9 and 81.2 m³ha⁻¹ for the years of 2008-2017 and 2020, respectively. The mean age of FF is 25.5, 22.2, 24.7, 22.9, 22.8, 22.0, 23.7, 27.8, 25.5 and 26.9 years for the years of 2008-2017, respectively. In 2018 and 2019, no new forests were found, whereas the mean age of FF found in 2020 is 15.9.

Concerning wood density, a new set of basic wood densities was introduced in 2010. This dataset (Table 6.5.4), which is used across all reporting years, includes basic wood densities based on a thorough revision of data reported in literature combined with re-measurements of wood densities for some species in a dedicated project (Somogyi, 2008a). Note that, to be consistent with the approach to use total aboveground volumes, the basic wood densities applied were measured from samples taken from all above-ground parts of the sample trees, including branches and bark.

Table 6.5.4. Basic wood density values for the main tree species and species groups in Hungary (Somogyi, 2008a).

Species or species group	Basic wood density (t/m ³)
Quercus robur	0.57
Quercus pertaea	0.61
Other quercus	0.55
Quercus cerris	0.64
Fagus silvatica	0.59
Carpinus betulus	0.58
Robinia pseudoacacia	0.59
Acer sp.	0.52
Ulmus sp.	0.58
Fraxinus sp.	0.56
Other hard broadleaves	0.5
Hybrid poplars	0.34
Indigenous poplars	0.36
Salix sp.	0.36
Alnus sp.	0.43
Tilia sp.	0.48
Other soft broadleaves	0.48
Pinus silvestris	0.42
Pinus nigra	0.47
Picea abies	0.39
Larix decidua	0.49
Other conifers	0.37

With respect to below-ground biomass, a general value for the root-to-shoot ratio (R) is applied. Due to lack of proper country-specific data, and in order to be consistent with previous estimates, IPCC default values (Table 4.4 of Chapter 4 of Volume 4 of the 2006 IPCC Guidelines) were considered by expert judgment). Considering that the majority of the forests in Hungary is young, and that the average volume stocks of FL-FL (calculated on the basis of volume and the area of forest sub-compartments) is 177 m³ ha⁻¹ (in 1990) and 211 m³ ha⁻¹ (in 2020), which demonstrates that stands predominantly grow on medium-quality sites, a conservative value of R of 0.25 is used for all species. The IPCC default values have relatively high uncertainty, but we believe that the probable value for the Hungarian forests is significantly higher than 0.25, which is thus a conservative value as long as forests are net sinks.

Concerning the carbon fraction of dry wood, the IPCC default values, i.e., 0.48 and 0.51 tonnes C tonnes biomass⁻¹ (Table 4.3 of Chapter 4 of Volume 4 of the 2006 IPCC Guidelines) were used for broadleaves and coniferous species, respectively. (These values, just like wood density and R, were consistently applied for the entire time series.)

Note that losses of carbon in biomass on land converted to forests are also reported in section 6.5.5 below using the methodology described in section 6.4.4 above.

6.5.4 Forest Land remaining Forest Land (CRF sector 4.A.1)

6.5.4.1 Category description

The main inventory estimates for the FL-FL category can be found in Table 6.5.5, whereas Table 6.5.6 summarizes methodological information for this category. (Note that, in order to be consistent with the CRF tables, only the area of forest and other sub-compartments is reported in this table.)

Table 6.5.5. *The area of forest and other sub-compartments as well as emissions (+) and removals (-) in the FL-FL sub-category by gas and inventory year.*

Inventory year	Emissions (+) and removals (-) in the FL-FL sub-category by gas and inventory year					
	Area	CO ₂	CH ₄	CO	N ₂ O	NO _x
	(ha)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1985	1 748 164	-312	0.678	10.653	0.027	0.351
1986	1 750 861	-3 211	0.692	10.896	0.028	0.358
1987	1 753 827	-3 491	0.662	10.401	0.027	0.343
1988	1 756 259	-3 722	0.646	10.177	0.026	0.335
1989	1 762 801	-2 423	0.645	10.153	0.026	0.334
1990	1 768 774	-2 872	0.597	9.416	0.024	0.310
1991	1 773 942	-2 205	0.572	8.961	0.023	0.296
1992	1 780 140	-2 754	0.515	8.005	0.021	0.265
1993	1 785 094	-5 221	0.433	6.575	0.017	0.220
1994	1 788 184	-5 612	0.435	6.541	0.017	0.220
1995	1 793 517	-5 357	0.458	6.923	0.018	0.232
1996	1 797 602	-1 394	0.500	7.570	0.020	0.254
1997	1 801 572	-1 689	0.494	7.361	0.019	0.249
1998	1 804 219	-2 634	0.475	7.022	0.018	0.238
1999	1 809 551	-332	0.316	7.035	0.017	0.202
2000	1 813 966	675	0.497	7.237	0.028	0.317
2001	1 817 339	-875	0.429	6.863	0.024	0.274
2002	1 821 574	-42	0.422	6.704	0.023	0.270
2003	1 822 625	-2 308	0.398	6.560	0.022	0.254
2004	1 823 592	-1 199	0.287	6.421	0.016	0.183
2005	1 832 045	-3 513	0.662	6.383	0.037	0.422
2006	1 841 327	-1 212	0.279	6.192	0.015	0.178
2007	1 851 638	-1 399	0.577	5.719	0.032	0.368
2008	1 864 537	-2 896	0.262	5.788	0.014	0.167
2009	1 876 819	-1 922	0.248	5.316	0.014	0.158
2010	1 884 094	-1 892	0.276	6.075	0.015	0.178
2011	1 893 054	-1 750	0.633	6.994	0.035	0.420
2012	1 903 602	-2 529	0.536	6.745	0.030	0.342
2013	1 909 771	-1 939	0.382	6.903	0.021	0.249
2014	1 913 041	-2 929	0.463	6.650	0.026	0.301
2015	1 916 428	-3 758	0.680	6.436	0.038	0.442
2016	1 920 720	-2 941	0.334	6.368	0.018	0.215
2017	1 926 514	-3 343	0.611	6.522	0.034	0.404
2018	1 930 930	-2 815	0.347	6.765	0.019	0.224
2019	1 936 571	-3 510	0.521	6.493	0.029	0.346
2020	1 946 190	-5 332	0.367	5.633	0.020	0.249

Table 6.5.6. Methodological summary for the Forest Land remaining Forest Land category.
(CS=country specific; D: default; EJ: expert judgment; IE: included elsewhere; AD: activity data;
EF: emission/removal factor)

Category	Type of information	Carbon stock changes					Table(S) I, II, V
		AGB	BGB	DW	LI	SOIL	
FL-FL	E/R	CS	D/EJ	AD: CS; EF:CS	Not estimated (demonstrated that not a source)	Mineral: Not estimated (demonstrated that not a source)	Fertilization: IE
						Organic: AD: CS	Drainage and re-wetting: NO
						EF: D	Biomass burning (slash burning + wildfires): AD: CS; EF: D
	Uncertainty	Results of the Tier 2 (Monte Carlo) analysis under FM are applicable	N/A				Results of the Tier 2 (Monte Carlo, where applicable) analysis under FM are applicable

6.5.4.2 CO₂ emissions and removals

The methodology to estimate emissions and removals in the forestry sector is based on that of the 2006 IPCC Guidelines. Whenever it was possible, country specific data was used (Tier 2), and IPCC default values (Tier 1) and expert judgment were only used in a few cases. Emissions and removals from the biomass and soil carbon pools are quantified and reported, however, due to lack of data, only assumptions are applied with respect to the other carbon pools to comply with requirements for completeness.

6.5.4.2.1 Biomass

Carbon stock changes of the biomass pool in the FL-FL category are calculated from those of the entire Forest Land, D and L-FL sub-categories. The methodology applied for the various sub-categories is as described in section 6.4 above. The input data from these categories, together with the formulas that are used to derive the FL-FL estimates are included in Table 6.5.7 below. The calculation method used ensures that the stocks of the new FF are excluded from calculating net removals. However, the net removals of all FF are included in the net removals of the FL-FL category. This is consistent with the method of KP-LULUCF reporting where net removals of all FF are included in those of FM (but the annual stock is excluded, see Chapter 11.3.1.1.2 and Table 11.9). Removals for the areas that are found in the inventory year are not estimated based on a dedicated survey in the newly found forests, rather, using an area specific mean net removal value (just like an “implied emission factor”) of the entire FL-FL category (calculated as the ratio of the total net removals and total area of forest sub-compartments), multiplied by the total area of found forests.

Table 6.5.7 Algorithms and data sources of calculating carbon stock changes for FL-FL under the UNFCCC, together with sample data for the last few years. For the calculation of emissions and removals from other categories in the table, see also the respective sections. The light yellow color in some cells of the table (with column title “from DB”) shows that the data in those cells are taken from the database (i.e., they are the result of other calculations), whereas data in white cells are calculated in this table from the respective cells. NE means net emissions, and IEF means “implied emission factor”. Symbol Δ is used to denote changes estimated as either differences between the value of a land-use category at two time points or using another methodology. All other notations are as in Table 6.5.3. (The table is for demonstration only and may include rounding errors; for precise numbers, see the respective CRF tables.)

Inventory year	Δ C of BIOMASS under the UNFCCC, GgCO ₂													
	FL		new FF (identified in the inventory year)		FL-L, new		FL		L-FL			L-FL net	FL-L	
	gross Δ	IEF	stock	IEF	Δ	IEF	net Δ = NR	IEF	gains	IEF	losses	Δ	NR	IEF
	from DB	NR/area (Gg CO ₂ /ha)	from DB	stock /area (Gg CO ₂ /ha)	from DB	Δ /area (Gg CO ₂ /ha)	gross Δ FL - new FF stock - D	NR/area (Gg CO ₂ /ha)	from DB	gains /area (Gg CO ₂ /ha)	from DB, only for information in this table	gains+ losses, only for information in this table	net Δ FL - L-FL gains (includes NR of all FF)	NR/area (Gg CO ₂ /ha)
2008	-5 048	-0.002652	876	0.1573	27	0.0922	-4 199	-0.002206	-1 260	-0.155056	123	-1 137	-2 939	-0.001576
2009	-4 139	-0.002163	980	0.1510	58	0.1289	-3 217	-0.001682	-1 202	-0.142745	63	-1 140	-2 015	-0.001074
2010	-3 603	-0.001875	478	0.1527	28	0.1336	-3 153	-0.001640	-1 216	-0.137042	100	-1 116	-1 937	-0.001028
2011	-3 566	-0.001850	644	0.1524	46	0.1659	-2 967	-0.001539	-1 144	-0.126015	31	-1 113	-1 823	-0.000963
2012	-4 578	-0.002368	872	0.1579	132	0.1682	-3 838	-0.001985	-1 098	-0.117352	24	-1 074	-2 740	-0.001439
2013	-3 692	-0.001905	630	0.1441	62	0.1156	-3 124	-0.001612	-1 086	-0.107155	16	-1 071	-2 038	-0.001067
2014	-4 632	-0.002386	555	0.1659	85	0.0447	-4 161	-0.002144	-1 097	-0.102811	23	-1 074	-3 064	-0.001602
2015	-5 059	-0.002607	156	0.1855	117	0.0844	-5 020	-0.002587	-1 075	-0.085556	4	-1 071	-3 945	-0.002059
2016	-4 159	-0.002144	87	0.1747	151	0.0716	-4 224	-0.002178	-1 041	-0.074622	4	-1 036	-3 183	-0.001657
2017	-4 776	-0.002462	329	0.1975	168	0.0981	-4 615	-0.002379	-1 003	-0.062471	12	-991	-3 612	-0.001875
2018	-3 919	-0.002021	0	NA	217	0.0977	-4 135	-0.002133	-973	-0.054772	23	-951	-3 162	-0.001638
2019	-4 589	-0.002367	0	NA	228	0.0987	-4 817	-0.002485	-942	-0.047133	28	-914	-3 875	-0.002001
2020	-6 289	-0.003239	180	0.1003	134	0.1125	-6 243	-0.003215	-911	-0.040845	42	-869	-5 332	-0.002740

The resulting carbon stock changes in FL-FL, in combination with those in L-FL (see below) demonstrate that the biomass of the forests in Hungary has been a sink for the last three decades. This is consistent with the fact that the total current annual increment (CAI) for the country has been estimated to be much higher than the annual harvests for all historical years (Figure 6.5.3).

The net volume stock changes, and thus the net carbon stock changes display some variability which is a partly consequence of the relatively stable CAI estimates and the rather variable harvest estimates, thus, the reported variability is considered to partly represent true variability. However, the applied survey method (i.e., continuous forest planning with a mean recycle length of about ten years, and the use of yield tables for updating between two consecutive surveys etc.) both reduces and removes and also adds some variability. Other components of the true variability, which are related to the annual variation of true increment of the stands due to weather may also be reflected in the data, but they cannot be separately captured by our estimation system. However, the reported inter-annual variability of the FL-FL carbon stock change estimates is not an artefact, and reflects relatively small changes in relatively large carbon stocks (these carbon stock changes being, in the last couple of years, less than one percent of the total biomass carbon stocks), mainly due to volume increment and harvest.




Figure 6.5.3 Annual harvest and current annual increment (CAI) in Hungary for the entire time series. Data source: National Forestry Database.

6.5.4.2.2 Dead organic matter

Of all dead organic matter (DOM), we can report on the carbon stock changes of the deadwood (DW) pool based on regression models created from deadwood sampling data of the NFI. The regression models and calculation method of deadwood stock-change are described in Chapter 6.5.5.2 for land converted to forest land. The same models were applied for forest land remaining forest land as follows: first the models were applied for the total forest land (i.e. not only for stands of 1-20 year-old) by stand type and deadwood stock change was calculated using the same equation as for land converted to forest land and then the amount of deadwood stock-change of land converted to forest land was subtracted from that.

The estimates show that the total amount of the above-ground and below-ground deadwood has been slowly but steadily increasing since 1985 (Figure 6.5.4).




Figure 6.5.4. The amount of the sum of the above-ground and below-ground deadwood stock in the Hungarian forests (except for afforested land). Data source: National Forest Inventory.

This increase of the amount of DW stock, and also that of LI, in the Hungarian forests is mainly due to two reasons. One is the high level of sustainability of the management of existing forests, which means that, for many years, less or much less wood has been harvested than what is grown (Figure 6.5.3). The difference between increment and harvests is large enough to claim not only sustained yield as detailed above, but also to assume that a lot of the uncut trees die due to the well-known process of self-thinning in stands where density has become high, or small-scale disturbances due to wind-break and snow-break which result in the increase of the amount of DW and LI. Also, for the last several decades, close-to-nature forest management has been promoted in Hungary, and clear-cuts have been restricted, especially after the adoption of the most recent Forest Act of 2009. This Act requests that semi-natural forests be managed in an increasingly natural way, which includes leaving more DW and LI in the forest after harvests than before, that gaps in the canopy be created and maintained, and that species mixture be enhanced. As a result of the implementation of these requirements, we can safely assume the accumulation of both DW and LI in the Hungarian forests for the last several decades.

Another reason of the increase of the amount of both DW and LI in all forests is that about one-third of all forests are afforestations since 1930 (i.e., before 1985), and most of these forests are still in their intensive growing phase, which means that carbon stocks of DOM pools have not saturated yet.

Finally, no major disturbances or other processes have occurred that could have resulted in substantial emissions from the DOM. Therefore, although no quantitative estimates can be made on the increase of the amount of DOM, the Tier 1 assumption can safely be made for LI, at least on average in the long run, that this pool is not a source.

The application of this assumption, as in most other countries, is due to the prohibitive costs and the very high variability which would require that measurements are taken on a very high number of plots, however, limits due to the practicability principle would not necessarily result in the reduction of the uncertainty to an acceptable level.

Our neighbouring country, Austria e.g., demonstrated in its NIR of 2020¹ that the uncertainty of the LI and soil pools of forest land is very high (for LI, $\pm 162\%$ at the country level and $\pm 144\%$ at the foothills that are most similar to Hungary), and that Monte Carlo simulations showed that these two pools have by far the highest contributions (3/4 to 4/5) to the total uncertainties of the emissions/removals of the total forest land subcategory.

There are very few examples for a LI monitoring that could be used for the actual estimation of carbon stock changes in the LI pool. One example is the one of the Netherlands where five different datasets were combined for the purposes of the GHG inventory², yet, it was concluded that: “The results of the Monte Carlo analysis consistently showed a carbon sink in litter; however the magnitude was very uncertain (Figure 4.5). Therefore, the more conservative estimate was used to set the accumulation of carbon in litter in Forest Land remaining Forest Land to zero.... Consequently, under the KP accounting the litter carbon pool under Forest Management is considered to be not a source.”

Some countries apply modelling (e.g., the YASSO model) to estimate soil and LI carbon stock changes. However, to correctly apply such a model, proper parametrization or validation is required, and while we are also thinking about applying such a model, we are not yet ready to do it especially considering the still high associated uncertainty of applying such a model.

The above-mentioned monitoring in the Netherlands demonstrates that, in correlation with the increase of the biomass carbon stocks, the carbon stocks of both the DW and the LI pools are also increasing. Given the similarities between the forests of the two countries (not necessarily concerning forest types but other relevant characteristics such as age distribution which determine the trends over large forested areas), the above correlation is also believed to apply to Hungary, and since our biomass carbon stocks have been increasing for several decades now, it follows that the litter carbon stocks have also been increasing.

As mentioned above, there is evidence from our NFI that the DW C stock is increasing in forest land remaining forest land. LI is a pool that contains dead mass of plants from two main sources: one is leaves and the other is dead woody parts such as twigs and cones that have currently not been accounted for in the deadwood pool. This means that when there is a carbon flow into the DW pool, then there must also be a carbon flow into the LI pool, i.e., the inflows are correlated, and an increasing deadwood C stock may lead to an increasing litter C stock. In the case of Hungary, LI is not removed from the forests, which means that even if emissions from decay are different from the DW and LI pools, the net carbon

¹ page 411 in: AUSTRIA'S NATIONAL INVENTORY REPORT 2020. Submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. URL: <https://unfccc.int/sites/default/files/resource/aut-2020-nir-15apr20.zip>

² pages 38-40 in: Greenhouse gas reporting for the LULUCF sector in the Netherlands. Methodological background, update 2019. E.J.M.M. Arets, J.W.H. van der Kolk, G.M. Hengeveld, J.P. Lesschen, H. Kramer, P.J. Kuikman & M.J. Schelhaas. Statutory Research Tasks Unit for Nature & the Environment, Wageningen, February 2019. URL: https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&cad=rja&uact=8&ved=2ahUKewih77Gb7NnsAhXK_qQKHWkcBN8QFjAAegQIAhAC&url=https%3A%2F%2Fedepot.wur.nl%2F472433&usg=AOvVaw0FMmuFuzUNAMeVTRMNMZkk

stock changes of these two pools should also correlate to a degree high enough to state that if the DW pool is a sink (and it is), then, as a minimum, the LI pool is not a source, either.

To conclude, we repeat that the main reason for the increase of the DW C stock in Hungary is the ageing of the forests (in general, there is less litter fall in young forests than in older ones) and the low intensity of harvest relative to the current annual increment. When harvests are less than current annual increment, self-thinning kicks in sooner or later, producing more and more of both DW and LI. On the other hand, there are no processes in the Hungarian forests that would lead to the loss of LI (or DW). The Hungarian forests have been sustainably managed (at least with respect to area and biomass) for decades, and transition to the forest land remaining forest land of formerly afforested areas, as well as that of FF (both also to land under FM), ensure that the increase of both the area and the carbon stock in the FL-FL (and FM) category is larger than losses (note that losses due to deforestation are accounted for in the FL-L / D category). Therefore, by also considering the practicability principle, and applying the combination of literature review and sound scientific reasoning, we have demonstrated that the LI pool is not a source.

6.5.4.2.3 Soils

As noted above, the Tier 1 assumptions applied for mineral soils of FL-FL, i.e., that these mineral soils are in carbon equilibrium. However, because the amount of dead organic matter increases, the DOM input into the soils increases which ensures that, even if slowly, the carbon stocks of the soils increase. Although there are some events in some forests that lead to emissions (e.g., natural disturbances, harvests etc.), there is a net carbon sink in large areas. A recent evaluation of permanent soil sampling points in forests showed that carbon stocks of forests where final harvest and subsequent soil preparation has not been made, the mean annual rate of carbon stocks amounts to 0.357 tC/ha*yr with a half-width CI of 0.013 tC/ha*yr. (Note that these numbers are under review at the time of the publication of this NIR.) Based on the combination of these quantitative estimates and reasoning (see details in Chapter 11) we conclude that, overall, mineral soils under the FL-FL category can be considered to be not a source.

With respect to organic soils, we conducted a dedicated project (Illés et al. 2013) to identify the forest area on organic soils. In this project, we measured the depth and carbon content of various layers in a sample of about 130 stands where it was suspected that we identify organic soils. The results of the project show that, in fact, the total area of forests on organic soils in the country amounts to 6.46 kha (the distribution of stands of organic soil is demonstrated on Figure 6.5.5). There is no rewetting of forest areas in Hungary. The emissions from the current organic soils, which were typically drained several decades ago, are calculated by multiplying this area by the default emission factor given in Table 2.1 of the Wetlands Supplement for Drained Forest land, temperate zone, i.e., 2.6 tCO₂ha⁻¹.




Figure 6.5.5. The distribution of forest stands on organic soil in Hungary (green areas; based on Illés et al., 2013).

6.5.4.2.4 Harvested Wood Products

For the estimation of the carbon stock changes of the harvested wood products (HWP) pool, the following data sources were used:

- international databases:
 - UNECE/FAO TIMBER database, 1964-2010, as of January 2012;
 - Joint Forest Sector Questionnaire (JFSQ) by ITTO, 2011-14;
- domestic forestry databases:
 - Halász, A., 1960 ("Erdőgazdaságunk, faiparunk és faellátásunk helyzete és fejlődése 1920-1958-ig"; Közgazdasági és jogi könyvkiadó, Budapest, 333 p., in Hungarian);
 - Halász, A., 1966 "Faellátásunk helyzete és fejlődése"; Mezőgazdasági Könyvkiadó Vállalat Budapest, 322 p., in Hungarian);
 - Halász, A., 1994 ("A magyar erdészet 70 éve számokban 1920-1990; FM Erdőrendezési Szolgálat, Budapest, 204 p., in Hungarian);
 - OSAP (Országos Statisztikai Adatfelvételi Program, or National Statistical Data Collection Program, by the National Land Centre); and
 - production and export/import data of the Hungarian Central Statistical Office.

Data are available since 1900 for the production categories as well as by domestic removals, import and export, however, due to concerns of accuracy, only data since 1964 have been used. Some data as examples are shown in Table 6.5.8.

The methodology used was first published by Király and Kottek (2014). Exports and imports were treated according to Equations 2.8.1 (for industrial roundwood) and 2.8.2 (for wood pulp) of the IPCC 2013 KP Supplement. The amounts of volume that are accounted for as input to the HWP pool exclude firewood as its carbon stock is accounted for using the instantaneous oxidation method.

Annual volumes of wood products were converted to amounts of carbon using the default conversion factors from Table 2.8.1 of the IPCC 2013 KP Supplement. To estimate net carbon stock changes of the HWP pool, the Tier 2 first order decay calculation method was used, i.e., Equation 12.1 from the 2006 IPCC Guidelines, together with default half life time values as required by Equation 2.8.5 of the IPCC 2013 KP Supplement, i.e., two years for paper, 25 years for wood panels and 35 years for sawn wood. Instantaneous oxidation assumed for wood in solid waste disposal sites.

Table 6.5.8. Volume or mass of wood in selected inventory years by product and production categories used in the calculation of the carbon stock changes of the HWP pool.

Wood product category	Type of quantity	Unit	Calendar year													
			1990	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Industrial roundwood	Removals	1000 m ³	3 518	2 822	2 365	2 746	3 018	2 987	3 169	3 119	3 065	2 950	2 862	3 038	2 892	2 457
	Import	1000 m ³	958	261	195	262	250	208	207	224	284	315	225	353	275	260
	Export	1000 m ³	1 159	725	691	875	881	858	975	871	680	683	634	547	624	728
Wood pulp	Production	1000 m.t.	46	0	0	0	4	6	2	24	19	22	21	21	28	51
	Import	1000 m.t.	152	107	91	88	109	94	106	131	142	130	141	149	140	147
	Export	1000 m.t.	3	0	0	0	4	6	2	24	10	0	0	0	2	4
Coniferous sawnwood	Production	1000 m ³	331	89	88	13	122	90	33	36	73	61	65	61	62	52
Non-Coniferous sawnwood	Production	1000 m ³	767	118	87	77	100	153	76	84	95	112	109	111	104	83
Veneer sheets	Production	1000 m ³	14	34	27	28	95	46	37	63	14	38	41	46	47	35
Plywood	Production	1000 m ³	14	19	17	5	38	43	26	61	47	34	46	58	67	49
Particle board (including OSB)	Production	1000 m ³	317	606	257	487	243	226	133	349	418	412	662	698	669	693
Hardboard	Production	1000 m ³	0	120	112	152	167	160	177	166	168	124	177	209	201	212
MDF (medium density fibreboard)	Production	1000 m ³	0	0	0	0	0	4	0	1	1	0	0	0	0	0
Fibreboard, compressed	Production	1000 m ³	50	0	0	0	0	0	0	0	0	0	0	0	0	0
Other board	Production	1000 m ³	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Paper and paperboard	Production	1000 m.t.	443	424	435	640	696	641	675	765	793	823	852	846	873	864

Note that, in Hungary, the estimated carbon stock changes in the HWP pool are relatively small due to the fact that the annual amounts of carbon entering this pool (i.e., through non-firewood wood products from harvests) and exiting it (i.e., through products ending their life cycle) are typically about the same.

See also section 11.5.2.5 for other details.

6.5.4.3 Non-CO₂ emissions

Estimated non-CO₂ emissions include those from burning of slash on-site and those from wildfires. Non-CO₂ emissions from these sources are not significant and are only reported for the sake of completeness and that of time series consistency with previous years. CO₂ emissions from these sources are accounted for in the biomass pool, because we apply the stock-change method. The carbon in the non-CO₂ emissions, i.e., the carbon of CO and CH₄, has been accounted for in the carbon stock change of the biomass pool, these gases are nevertheless reported because of their high global warming potential, because the double counting of the carbon is negligible and also to comply with the 2006 IPCC GL.

6.5.4.3.1 Non-CO₂ emissions from burning of slash

The estimation of these emissions is done according to section 6.4.3 with the following modification:

$$M_b = V_b * D$$

where

V_b = volume burnt, m^3 (only includes biomass, reported in Table 6.5.9);

D = wood density, $kg\ biomass\ m^{-3}$ (values used here are the same as those used to estimate carbon stock changes in biomass, see Table 6.5.4 above); and

$$V_b = V_H * C_f$$

where

V_H = total harvest, m^3 of wood removed from forest (taken from harvest statistics), and

C_f = combustion factor, dimensionless, for which we use average country-specific values by species (*Rumpf, 2013*). These values are based on expert solicitation and are in line with in legislature on burning in forests.

Finally, when estimating non- CO_2 emissions from the above data, default IPCC G_{ef} values were used in Equation 2.27 (see Section 6.4.3).

Table 6.5.9. *The amount of harvested volume, slash burnt and forest fires based on all available data.*

Reporting year	The amount of controlled burning and forest fires based on all available data					
	Harvested volume (m3)	Slash burned on site (t)	Number of wildfires in forest	Area of forest subcompartments burnt in forest and agricultural fires EFFIS (ha)	Area of forest subcompartments burnt in forest fires (ha)	Wood volume burnt in forest fires (m3)
1985	8 345 562	99 560	NE	NE	NE	NE
1986	8 500 991	101 835	NE	NE	NE	NE
1987	8 193 145	97 202	NE	NE	NE	NE
1988	7 960 397	95 109	NE	NE	NE	NE
1989	8 031 779	94 892	NE	NE	NE	NE
1990	7 415 162	88 002	NE	NE	NE	NE
1991	7 255 202	83 750	NE	NE	NE	NE
1992	6 589 097	74 816	NE	NE	NE	NE
1993	5 723 745	61 452	NE	NE	NE	NE
1994	5 717 468	61 131	NE	NE	NE	NE
1995	6 049 151	64 698	NE	NE	NE	NE
1996	6 603 733	70 750	NE	NE	NE	NE
1997	6 713 101	68 793	NE	NE	NE	NE
1998	6 578 931	65 630	NE	NE	NE	NE
1999	6 900 612	65 749	229	756	756	3 000
2000	7 287 456	67 632	811	1 595	1 595	80 000
2001	7 010 979	64 142	419	na	1 223	57 000
2002	7 013 167	62 653	382	1 227	1 226	57 000
2003	7 085 514	61 312	375	845	1 054	49 000
2004	7 094 753	60 013	104	247	354	2 000
2005	7 167 426	59 651	150	3 531	3 530	170 000
2006	7 005 190	57 872	97	625	625	3 000
2007	6 609 099	53 444	139	4 636	2 057	160 660
2008	7 024 025	54 091	54	2 404	402	2 730
2009	6 773 537	49 679	87	6 463	845	7 000
2010	7 424 046	56 780	7	878	239	5 324
2011	8 080 206	65 363	569	8 055	1 189	149 651
2012	7 731 605	63 038	712	14 115	4 303	120 918
2013	7 874 792	64 518	259	1 955	407	36 457
2014	7 517 408	62 152	367	4 454	756	79 768
2015	7 354 188	60 153	421	4 730	1 593	192 394
2016	7 338 350	59 516	161	974	218	24 049
2017	7 576 110	60 950	575	1 454	1 222	150 353
2018	7 766 751	63 224	530	906	275	22 832
2019	7 315 179	60 687	2 088	6 541	844	115 887
2020	6 580 366	52 646	1 239	2 895	629	61 530

6.5.4.3.2 Non-CO₂ emissions from wildfires

Wildfires are very erratic in nature and are not a really significant phenomenon in Hungary. Beginning 1999, the Fire Department started to provide data on the number and area of forest wildfires, however, until 2006, these numbers are not deemed accurate, and the emissions based on these are only rough estimates.

In 2006, Hungary joined the European Forest Fire Information System (EFFIS, <http://effis.jrc.ec.europa.eu/>), and a new database was established. Thus, beginning 2007, the Fire Department locates the fires, surveys the affected area and, subsequently, the Forest Authority identifies the affected forest sub-compartments. This identification is done on site, during a survey after the fire. The Forest Authority also develops data, based on area estimates by the National Directorate for Disaster Management of the Ministry of the Interior, on the growing stock that was burnt in fires. This way, the activity data is double-checked, and the emissions can be accurately calculated based on the estimated amount of standing volume burnt.

The amount of wood volume burnt in wildfires between 1999-2006 are calculated by the ratio of fire-affected area for this period and the burned growing stock per unit area of wildfires of 2007-2008.

With the exclusion of some areas affected by forest fires that are subsequently considered and reported as Deforestation (D), the vast majority of burnt areas remain under forest management by law, and the Forest Authority prescribes the reforestation/regeneration of these areas and inspects their success.

The estimation of the amount of emissions is done according to section 6.4.3 with the modification that is also applied for slash burned, see above. The amount of V_b , i.e., the amount of volume burnt in the areas affected is reported in Table 6.5.9 above (i.e., $C_f = 1$). Finally, when estimating non-CO₂ emissions from the above data, default IPCC G_{ef} values were used in Equation 2.27 (see Section 6.4.3).

6.5.5 Land converted to Forest land (CRF sector 4.A.2)

6.5.5.1 Category description

In Hungary, mainly former croplands are afforested. Converting grasslands to forests occurs less frequently (in about 15% of all conversions), whereas converting other areas to forests is marginal, therefore, we predominantly report carbon stock changes from converting croplands and grasslands to forest land.

Land converted to forest land includes areas that, apart from soils, do not contain much carbon in either of the carbon pools before they are afforested. These areas are subject to the effect of intensive photosynthesis after the afforestation which, possibly after some pool-specific lag, usually results in the increase of carbon in all carbon pools. An exception to this is perennial cropland where perennial biomass may be present before the afforestation. Concerning soils, cropland usually stores less carbon than forest land. Thus, converting land to forest land generally increases the amount of carbon in all pools, although at different rates, due to tree growth after the afforestation.

Figure 6.5.6 reports estimated emissions and removals for the various pools, whereas Table 6.5.10 summarizes methodological information. See also Section 11 for other details.




Figure 6.5.6. The sources of emissions and removals in the various pools in the L-FL category. The data under the grey box is shaded out for reasons explained in section 6.1.1.

Table 6.5.10. Methodological summary for Land converted to Forest Land. (CS=country specific; D: default; EJ: expert judgment; IE: included elsewhere; AD: activity data; EF: emission/removal factor)

Category	Type of information	Carbon stock changes					Table(5) I, II, V
		AGB	BGB	DW	LI	SOIL	
L-FL	E/R	Post-conversion: CS	D/EJ	AD: CS; EF:CS	AD: CS; EF:CS	Mineral: AD: CS	Fertilization: IE
						EF: CS	Drainage and re-wetting: NO
						Organic: not occurring	Biomass burning: NO
	Uncertainty	Tier 2 (Monte Carlo)		NE			

6.5.5.2 CO₂ emissions and removals

6.5.5.2.1 Biomass

2 CO₂ emissions and removals from the biomass pool in L-FL are estimated from carbon stock changes due to gains in the trees appearing after the afforestations and losses of biomass before the conversion, if any. Whereas a country-specific method is applied for the estimation of gains, Equation 16 of the 2006 IPCC GL are used to estimate losses (see section 6.4.4). For both procedures, the estimation of the conversion area is necessary.

Area of conversion

The area is taken from the “initial planting of afforestations” statistics of the NFCSO Forestry Directorate by target stand-type (*Quercus* sp., *Quercus cerris* and other hard broadleaved, *Fagus*,

Robinia, Hybrid Poplar and Salix sp., Indigenous poplars and other soft broadleaved, and Conifers). For the period 2008-2020 the data is taken from the AR database. Data for 1990-2007 was taken from a historical dataset of the Forest Authority that is primarily used to have a subsidy-supporting roll. The area of L-FL cannot be identified on sub-compartment-level in this period. Therefore, for the entire period, modeling was used to develop the growing stock, increment and removal data based on total annual conversion area and age-mean volume function (see below).

Table 6.5.11 below demonstrates the evolution of total area of the category over time (similar tables are used for the estimation by the above species groups). The table shows the area *entering* the category as new afforestation in the second column (under year 1). The area in a year is then rolled over to 19 additional inventory years (in the subsequent columns and rows). The total area in the category in a year (in the last column) is thus the sum of all areas in all age classes. Each area is moved to the FL-FL category after the period of the default 20 years except for some L-FL areas that have been deforested and that are moved to the D category.

Table 6.5.11. The total area of land that is successfully converted to forestland (for all species combined) by year of conversion (blue cells), and total land in the category. Incoming areas are reported in the first year of conversion, and areas reported in the column “area of successfully converted land ... in age class ... 20” are transferred to the FL-FL category the next year. Note that, in rare cases, afforested areas are deforested and the values in the below matrix show the areas after the deforestation.

Inventory year	area of successfully converted land (i.e. area actually covered by trees, ha) in age class (yr)																				Total in inventory year (ha)
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1985	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7 476
1986	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	14 972
1987	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	22 864
1988	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	31 348
1989	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	38 633
1990	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	0	45 127
1991	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	0	51 461
1992	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	0	58 200
1993	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	0	61 244
1994	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	0	63 957
1995	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	0	67 904
1996	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	0	74 144
1997	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	0	81 997
1998	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	0	89 742
1999	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	0	97 961
2000	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	0	107 203
2001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	0	119 605
2002	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	0	133 606
2003	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	0	144 948
2004	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	7 476	152 098
2005	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	7 496	151 851
2006	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	7 892	157 561
2007	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	8 484	167 556
2008	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	7 285	166 293
2009	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	6 494	162 527
2010	6 267	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	6 334	162 300
2011	1 641	6 267	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	6 739	157 607
2012	1 162	1 641	6 267	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	3 045	152 031
2013	697	1 162	1 641	6 267	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	2 713	149 683
2014	1 422	697	1 162	1 641	6 267	3 520	7 221	17 888	13 206	7 228	7 150	11 342	14 001	12 402	9 242	8 219	7 745	7 854	6 240	3 946	148 391
2015	245	1 422	688	1 140	1 639	6 252	3 519	7 195	17 865	13 150	7 214	7 134	11 318	13 976	12 378	9 215	8 212	7 741	7 849	6 237	144 390
2016	242	244	1 419	673	1 126	1 635	6 251	3 503	7 185	17 819	13 124	7 200	7 115	11 272	13 932	12 360	9 190	8 194	7 685	7 841	138 009
2017	756	240	229	1 406	673	1 126	1 628	6 230	3 489	7 177	17 813	13 113	7 194	7 103	11 264	13 932	12 355	9 175	8 179	7 681	130 759
2018	1 341	749	240	228	1 405	669	1 119	1 622	6 219	3 487	7 158	17 813	13 105	7 183	7 090	11 261	13 925	12 342	9 172	8 178	124 307
2019	1 676	1 341	747	238	228	1 403	669	1 119	1 619	6 219	3 486	7 132	17 807	13 094	7 175	7 076	11 260	13 923	12 329	9 169	117 711
2020	2 428	1 665	1 321	747	237	222	1 403	669	1 095	1 619	6 200	3 479	7 132	17 798	13 085	7 170	7 060	11 251	13 914	12 320	110 815

A conversion of non-forest land to forest land, i.e., an afforestation activity, is deemed to have begun

when soil preparation has been started. Typically, the first (“initial”) planting of the propagation material on the area happens in a short time after soil preparation is done. Beating up may be carried out depending on the success rate of the initial planting.

Post-conversion biomass

Carbon stock changes in the biomass pool of the newly established trees are estimated using simplified regression-type models of growing stock over age on unit areas of afforestation. The models were developed using a sample of young stands of varying age (which is known based on the year of the afforestation) for which volume was known. This volume was available either from direct assessment or from yield tables (in this last case, height was measured). The models are available separately for the above 7 target stand-types (Figure 6.5.7 below is an example of data and the regression obtained for *Quercus* sp.). We used 3rd degree polynomial regressions for species of long rotation age (such as beech and oak), and linear regressions for species of short rotation age (such as Black locust). The regression curves were developed by also using stands somewhat older than 20 years to increase the robustness of the regression curve, but they are only used for stands between ages 1 and 20 years in the L-FL category. The curves were forced to start from the pole. All coefficients of determination are above 0.9, and the regression coefficients are significant for each species. (Note that these curves represent rather constant growing conditions, and it is only worth checking the validity of these curves, or revise them, rather rarely.)




Figure 6.5.7. An example of the regression curve of stand volume over age using data of *Quercus* sp.

Volume stock *change* from a specific year of age to the next one is equal to the *difference* between the volume stocks of the respective consecutive ages as estimated from the regression curves. To get total carbon stock change, these differences for the various age classes and species were multiplied by the area of the same classes (whereas conversion factors by species were also used to develop amounts of carbon from amounts of volume). The estimated changes reflect the effects of tree growth, artificial thinnings and self-thinnings. The calculated changes are smoothed ones, i.e., they do not represent any

inter-annual variation due to e.g., variation of growing conditions. More importantly, however, the above procedure ensures that the volume *stocks* of the respective classes are not directly applied in the calculations, and no transition of volumes are directly included (either for L-FL or FL-FL) in the calculation of carbon stock changes when stands are moved from the L-FL category to the FL-FL category.

In estimating *carbon* stock changes in these forests, the conversion of volume to carbon happens the same way as described above when discussing the second part of Equation 2.8 of the 2006 IPCC Guidelines.

Pre-conversion biomass

Emissions arising from removing perennial biomass carbon during conversions to forest are estimated using data from a study that estimated the amount of carbon lost by removing all above-ground biomass due to conversion *for a unit area*. This loss was measured to be 4.7 tC/ha in case of orchards and 9.39 tC/ha in case of vineyards. The average age of the converted areas is 15 and 15.9 years, respectively, which are half of the length of the rotation period of these perennials (Juhos and Tőkei, 2013). The respective amounts of carbon lost from annual biomass, which are IPCC defaults, are 4.7 and 2.94 tC ha⁻¹ (see the respective sections on Cropland and Grassland for more details). The amount of *total loss* of carbon is estimated by multiplying the above values with the the area actually converted. (In practice, carbon stock changes from biomass of perennials on CL-FL are estimated by first estimating all carbon stock changes from cropland with perennials to all other categories, the methodology of which is reported in section 6.6.2.1.1, and then multiplying it with the proportion of the area of perennial CL-FL to the total area of perennial CL converted to all other land-use category. For forest land, this proportion varies between 12 and 19%.)

Note that, according to the default method (Equation 2.12), the pre-conversion below-ground biomass is not considered a loss, and indeed it remains in the ground and adds to the carbon stocks of the deadwood, litter and soil pools.

6.5.5.2.2 Dead organic matter

Carbon stock changes in the DW pool are estimated the following way:

$$\Delta C_{DW} = \sum_{i,j} A_{i,j} * \Delta v_{i,j} * (1+R) * D * CF$$

where

$A_{i,j}$ = area of species group i ($i = 1 \dots 7$) in age class j ($j = 1 \dots 20$), ha;

$\Delta v_{i,j}$ = annual deadwood volume stock increase of species group i in age class j , tC * ha⁻¹ * yr⁻¹,

R = root-to-shoot ratio (same as for biomass, see above);

D = wood density as for woody biomass (same as for biomass, see above);

CF = carbon fraction as for woody biomass (same as for biomass, see above).

The area-specific $\Delta v_{i,j}$ values are based on data collected from 7629 sampling points of the NFI. The measurements were made in all forests (mostly in the FL-FL category) between 2010 and 2017. (The amount of data from plots in afforested areas was not enough to develop reliable regression fitting, see below.) Data were collected for standing dead trees with diameter breast height equal to or larger than 7cm, and for lying dead trees with a minimum length of 1m and a diameter of at least 10 cm. (This means that only above-ground values were measured with definitions that are a bit different from those for living trees, but the differences in the definitions render the carbon stock change estimates conservative.) Area-specific measurements were aggregated in seven tree stand types by age, and so

seven lines were fitted to the data over age (Figure 6.5.8). The data show that the increase of DW carbon stock lasts until about 30 years (in case of fast-growing species like poplars) to well over 100 years (in case of slow-growing species like oaks); beyond available data, to be conservative, zero increase was assumed. However, as areas are in the L-FL category only for 20 years, we use data for only the 1-20 years of age part of the regression lines for each species. Note that most regression lines do not start at zero stocks because the analyses were done for regenerated stands in the FL-FL category where some stocks remain in the stands after a regeneration. For L-FL, however, we assumed that the initial stocks are zero and the increase (change over time) is the same as in the FL-FL category (i.e., it is a bit underestimated).




Figure 6.5.8. Data and fitted regression line for the seven tree species groups over age. Beyond the oldest age categories with data, horizontal lines, i.e., $\Delta v = 0$, were used in the calculations.

Concerning afforestations, it is a general experience that the DOM pools do not contain carbon on cropland and grassland before the conversion. This is corroborated by the fact that, for sanitary and other reasons, biomass (dead or alive) of all crop is removed during harvests, and there is usually not enough time on abandoned croplands for the woody biomass to develop substantial dead organic matter before the land is converted. Somogyi et al. (2013) also measured zero dead organic matter carbon stocks on pre-conversion land in their study.

Concerning the assumption that the rate of DW increase is the same on L-FL and FL-FL, our argumentation is based on the theory that if the initial stock is not zero, then the net change is the result of input to the pool and the loss from the decay of the initial stock. This means that the rate of increase on an afforested land, where the initial stock is zero, should be larger than on an FL-FL land. It follows

that, while also limiting the accounting for the increase to the first 20 years of the afforestation, our approach is conservative.

For LI, the estimation is done by applying the methodology described in section 6.4.5 above. We assume that the carbon stock in the litter pool before the conversion is zero. For the equilibrium carbon stock in the forest land after the conversion, we apply the same value, 8.78 t/ha (Heil et al., 2012), that is used for the estimation of emissions due to deforestation events (i.e., conversions of forest land to other land-uses), see section 6.5.6.1.2. We assume that the equilibrium litter carbon stock level is reached in 20 years.

6.5.5.2.3 Soil

The estimation of carbon stock changes in soils is done according to section 6.4.1.

The results of the estimation are corroborated by recent estimates according to which converting land from cropland to forest does not entail any net emissions from soil, rather, net removals (see Somogyi, 2005, Somogyi-Horváth, 2006a, Somogyi-Horváth, 2006b, and Somogyi et al., 2013), and that converting grassland to forest may lead to some emissions (see Horvath, 2006). However, because most of the huge amounts of marginal lands that are afforested are former croplands, and also because of biodiversity concerns, the overwhelming majority of conversions occur on abandoned croplands (see above), overall, no major emissions from soils are expected due to conversion of land to forest land (Figure 6.5.9).




Figure 6.5.9. Emissions and removals from converting land to forest land (non-SA: non-set-aside; SA: set-aside). The data under the grey box is shaded out for reasons explained in section 6.1.1.

Concerning organic soils, there are no afforestations on such soils, therefore, no emissions occur from this source.

6.5.5.3 Non-CO₂ emissions

6.5.5.3.1 Emissions from wildfires

In Hungary, very few forest fires occur in the Land converted to Forest Land category. However, following the former recommendation of the inventory review, non-CO₂-emissions are separately reported for L-FL and FL-FL (see above).

6.5.5.3.2 Direct and indirect N₂O emissions from mineral soils

The estimation of both direct emissions from mineral soils associated with loss of carbon resulting from change of land-use or management and indirect N₂O emissions from leaching/runoff is done according to section 6.4.2, using the appropriate carbon loss data for the sub-category. Note that, consistent with what is reported for soils above, emissions only occur when grasslands are converted to forests.

6.5.6 Forest Land converted to other land-uses (CRF sector 4.B.2.1, 4.C.2.1, 4.E.2.1)

The area of forest land converted to other land-uses has been small in Hungary for the last decades, and the conversions only include conversions to cropland, grassland and settlements. Conversions from forest land to any other land-use type are generally prohibited by the Forest Act and can take place only after the Forest Authorities grant the specific permission. All areas of conversions are surveyed, and emissions are estimated using the land conversion database of the Forest Authorities (see at http://www.nfk.gov.hu/Official_statistics_news_546, in Hungarian). However, these statistics are only available since 1985, and the average of the period 1985-1989 is used for the previous years for which estimates are also needed to run up the calculation of the estimation of emissions from soils.

The area of FL-L is the same as for the D category and has been identified in a detailed procedure that is reported in section 11.1.3.2. For further information on deforestation in Hungary, see Section 11.3.1.1.

6.5.6.1 CO₂ emissions and removals

Table 6.5.12 reports CO₂ emissions and removals estimated for the biomass (above-ground plus below-ground), deadwood, litter and soil pools, whereas Table 6.5.13 reports methodological information for this sub-category.

Table 6.5.12 Total emissions from biomass (above-ground plus below-ground), deadwood, litter and soils from the FL-L category. The light-yellow color in some cells of the table (with column title “from DB”) shows that the data in those cells are taken from the database (i.e., they are the result of other calculations), whereas data in white cells are calculated in this table. All notations are as for Tables 6.5.3 and 6.5.6. (The table is for demonstration only and may include rounding errors; for precise numbers, and for data by geographical locations, see the respective CRF tables.)

Inventory year	Emissions and Removals from FL-L, GgCO ₂				
	biomass	mineral soils	organic soils	litter	dead-wood
	from DB	from DB	from DB	from DB	from DB
2008	27	29	NO	9	3
2009	58	30	NO	14	5
2010	28	31	NO	7	2
2011	46	31	NO	9	3
2012	132	28	NO	25	9
2013	62	29	NO	17	6
2014	85	32	NO	61	22
2015	117	33	NO	45	17
2016	151	36	NO	68	26
2017	168	40	NO	55	21
2018	217	38	NO	71	27
2019	228	39	NO	74	29
2020	134	38	NO	38	15

Table 6.5.13. Methodological summary. (CS=country specific; D=default; NO: Not occurring; NE: not estimated)

Category	Type of information	Carbon stock changes					Table(5) I, II, V
		AGB	BGB	DW	LI	SOIL	
FL-L	E/R	Post-conversion: 0	0	CS	CS	D	Drainage: NO
		Pre-conversion: CS	CS				Biomass burning: NO
	Uncertainty	NE					

6.5.6.1.1 Biomass

For biomass in the forest sub-compartments, the methodology described in section 6.4.4 is applied. For all conversions of FL to CL, GL, WL, SE and OL, it is assumed that the forest biomass carbon stock after the conversion is zero, so all carbon in the biomass due to deforestation, estimated as described in section 6.5.3, is completely emitted as CO₂. (Note that, depending on the post-conversion land-use, there may be non-forest biomass after the conversion, which is estimated following the methodology described in the respective section on the post-conversion land-use.)

For the biomass pool in the other sub-compartments (see section 11.1.3.2), the emission factors B_{Before} (in the equation in section 6.4.4. Conversion-related carbon stock changes of the biomass pools) are the following: for the utilization of forest types, the annual B_{Before} emission factor is the same that is used for the “forest sub-compartments”; for the other types we assumed that the pre-conversion biomass is equal to the default biomass of the land-use category which the particular areas were classified to before the conversion. The post-conversion value corresponds to the biomass on the “converted-to” land-use category.

6.5.6.1.2 Dead organic matter

Emissions from deadwood and litter are estimated by multiplying the area of annual deforestations by the average stock value of these pools. In these calculations, just like with biomass, we assume that all deadwood and litter are completely removed from the area, i.e., all carbon in these pools are emitted in the year of the deforestation.

The total deforested area applied in these calculations includes, in addition to the area of the forest sub-compartments, the area of the fraction of the other sub-compartments that is re-classified as FL (see section 11.1.3.2). However, due to the nature of the non-forest land-use categories, no litter or deadwood is assumed to be present in the other sub-compartments either before or after the conversion, therefore, emissions from the litter and deadwood pools in these areas is assumed to be zero and the activity data in the below calculations is the sum of the area of forest sub-compartments and those parts of the other sub-compartments that were classified as FL.

The area-specific value of the amount of deadwood comes from the National Forest Inventory as described in section 6.5.5.2.2. Since the amount of deadwood in the sampled years do not show a decreasing trend either overall (see Figure 6.5.4) or by species (see Figure 6.5.8), we believe that the average area-specific data calculated from these samples are suitable for estimating the emissions from deadwood for the entire time series.

In estimating total carbon stock changes from the estimated volume, we used the methodology of stock change as detailed above and applied the assumption that the average wood density of the deadwood is the same as for the woody biomass.

Considering litter, the dedicated case study by Heil, Kovács and Szabó (2012) provided an estimate of the mean litter content (excluding coarse litter between about 1 cm and 10 cm) of the Hungarian forests. In this study it was found that the average amount of carbon in litter is 8.78 tC/ha.

We note that, because of the small scale of deforestations each year, and because litter and especially DW are relatively small carbon pools, this simple but anyway Tier 2 approach can be regarded as an accurate methodology as far as practicable. We also note that the above value is considerably smaller than the IPCC default values reported in Table 2.2 of Chapter 2 of the 2006 IPCC Guidelines for mature forests (i.e., 27-28 tC/ha). This is partly because of the differences in the definition of litter in this NIR and the 2006 IPCC Guidelines (i.e., most fine woody debris is excluded from our definition of the litter),

and partly because the IPCC default values are for mature forests whereas most of our forests are not mature, and possibly partly due to other methodological differences such as the carbon content of litter.

It is additionally assumed that neither deadwood nor litter are produced any more after Forest land to non-forest land conversions, thus, no removals are accounted for in these pools after the conversions.

6.5.6.1.3 Soil

The estimation of carbon stock changes in mineral soils was done according to section 6.4.1. No forest land on organic soils is converted to other land uses.

For each piece of land converted (including both forest and other sub-compartments), the same amounts of annual carbon stock changes are accounted for 20 consecutive years for all conversion types. Carbon stock changes for all conversion types are added up to get the aggregated emissions for the entire FL-L category.

For the forest sub-compartments (see section 11.1.3.2), the post-conversion land use type is known, therefore, values in Table 6.4.2 above can directly be used. Concerning the other sub-compartments, in addition to the assumption for the re-classification of the area to areas by land use types, we had to make assumptions concerning the conversion type because it is not known for these other sub-compartments what pre-conversion land-use type is converted to what post-conversion land-use type. Because of these uncertainties, the assumption applied is that the difference between the amount of area-specific carbon before and after the conversion for a specific pre-conversion land-use type is the largest value of all the possible conversion types. For example, for an area within the other sub-compartments classified as Wetland (WL) before the conversion, we applied the emission factor for the WL – non-SA CL conversion in Table 6.4.2 because it involves the largest area-specific C loss.

The FL-L areas identified and the resulting total CO₂ emissions from all pools are included in Table 6.5.14.

Table 6.5.14. *The area, as well as CO₂ emissions from soils on land converted from forest to other land-uses.*

Inventory year	FL converted to CL			FL converted to SE			FL converted to GL			All conversions from FL to other land use		
	Area (ha)		CO ₂ emissions (GgCO ₂ eq)	Area (ha)		CO ₂ emissions (GgCO ₂ eq)	Area (ha)		CO ₂ emissions (GgCO ₂ eq)	Area (ha)		CO ₂ emissions (GgCO ₂ eq)
	all	forest subcomp-artments		all	forest subcomp-artments		all	forest subcomp-artments		all	forest subcomp-artments	
1985	94.8	94.8	0.211	210.5	210.5	0.407	20.9	20.9	-0.062	326.1	326.1	0.556
1986	94.8	94.8	0.423	210.5	210.5	0.813	20.9	20.9	-0.124	326.1	326.1	1.111
1987	94.8	94.8	0.634	210.5	210.5	1.220	20.9	20.9	-0.188	326.1	326.1	1.666
1988	94.8	94.8	0.847	210.5	210.5	1.626	20.9	20.9	-0.251	326.1	326.1	2.222
1989	94.8	94.8	1.059	210.5	210.5	2.033	20.9	20.9	-0.312	326.1	326.1	2.780
1990	180.0	180.0	1.463	392.6	392.6	2.791	40.3	40.3	-0.428	612.9	612.9	3.826
1991	453.6	59.9	2.480	1265.6	167.0	5.236	97.8	12.9	-0.715	1817.0	239.8	7.000
1992	511.8	44.4	3.628	827.3	71.8	6.834	107.9	9.4	-1.020	1447.1	125.6	9.442
1993	12.7	12.7	3.660	233.1	233.1	7.284	82.7	82.7	-1.218	328.6	328.6	9.727
1994	28.4	28.4	3.727	162.5	162.5	7.598	27.3	27.3	-1.236	218.2	218.2	10.089
1995	53.2	53.2	3.849	244.1	244.1	8.070	60.5	60.5	-1.362	357.8	357.8	10.557
1996	140.4	78.7	4.168	335.4	188.1	8.718	140.9	79.0	-1.701	616.7	345.9	11.185
1997	192.1	192.1	4.604	239.6	239.6	9.180	90.3	90.3	-1.890	522.0	522.0	11.894
1998	88.9	88.9	4.801	271.4	271.4	9.705	41.7	41.7	-1.969	402.0	402.0	12.537
1999	98.1	26.8	5.020	1016.9	277.9	11.669	331.9	90.7	-2.769	1446.9	395.4	13.920
2000	111.8	67.8	5.260	981.6	594.9	13.565	93.1	56.4	-2.987	1186.6	719.1	15.837
2001	152.8	61.4	5.596	893.0	358.6	15.290	251.2	100.9	-3.570	1297.0	520.9	17.316
2002	317.0	108.9	6.300	1279.7	439.5	17.762	259.6	89.2	-4.191	1856.4	637.5	19.870
2003	54.3	25.7	6.415	1104.7	523.4	19.895	93.1	44.1	-4.406	1252.1	593.3	21.904
2004	109.0	74.2	6.651	1102.7	750.5	22.025	175.0	119.1	-4.828	1386.7	943.8	23.848
2005	148.7	71.2	6.763	654.4	313.2	22.883	55.7	26.7	-4.900	858.8	411.1	24.746
2006	115.8	44.4	6.803	1156.6	443.4	24.710	54.2	20.8	-4.968	1326.7	508.6	26.545
2007	90.7	16.4	6.786	1061.1	192.5	26.353	201.7	36.6	-5.390	1353.5	245.5	27.749
2008	379.6	96.8	7.412	634.7	161.9	27.173	137.5	35.1	-5.703	1151.9	293.8	28.882
2009	183.8	55.5	7.601	970.3	293.0	28.640	335.9	101.5	-6.424	1490.0	450.0	29.817
2010	670.4	59.4	8.679	1154.8	102.3	30.112	526.1	46.6	-7.590	2351.3	208.3	31.201
2011	388.5	66.8	8.544	1074.6	184.9	29.744	140.5	24.2	-7.695	1603.5	275.9	30.593
2012	247.6	113.1	7.968	614.1	280.4	29.332	851.6	388.9	-9.478	1713.2	782.4	27.822
2013	270.2	115.4	8.550	702.4	299.9	30.238	273.5	116.8	-9.947	1246.1	532.1	28.841
2014	711.2	482.0	10.081	1631.3	1105.6	33.074	448.4	303.9	-10.966	2790.8	1891.4	32.189
2015	520.6	423.7	11.137	766.0	623.4	34.082	412.5	335.7	-11.836	1699.1	1382.8	33.384
2016	1340.8	1003.1	13.833	911.4	681.8	35.194	576.4	431.2	-12.847	2828.6	2116.2	36.180
2017	2367.3	1045.1	18.714	630.3	278.3	35.948	877.8	387.5	-14.707	3875.4	1710.9	39.956
2018	785.0	515.4	20.296	775.6	509.2	36.923	1817.0	1193.0	-18.971	3377.6	2217.6	38.248
2019	934.0	818.8	22.195	1010.1	885.6	36.910	687.0	602.3	-19.808	2631.1	2306.8	39.297
2020	465.2	368.5	22.999	546.4	432.8	36.068	491.9	389.7	-20.778	1503.4	1191.0	38.289

6.5.6.2 Non-CO₂ emissions

6.5.6.2.1 Emissions from fires

The estimation of non-CO₂ emissions from fires is based on the fact that deforestations in Hungary are done by clear-cutting the areas and removing most biomass from there.

As deforestations rarely occur in the country, the probability that wildfires affect these areas is negligible. In the last years, no wildfires occurred on land that later (in the same year) were converted to other land-use. Therefore, emissions from wildfires are reported as not occurring in this category.

On the other hand, controlled burning (burning of slash) occurs in this category. The methodology to estimate emissions from this source is the same as described in section 6.4.3. Activity data is available for both the area and the volume of forest land converted to other land-use.

6.5.6.2.2 Direct and indirect N₂O emissions from mineral soils

The estimation of both direct emissions from mineral soils associated with loss of carbon resulting from change of land-use or management and indirect N₂O emissions from mineral soils due to leaching/runoff is done according to section 6.4.2.

6.5.7 Category-specific uncertainties and time-series consistency

For this category, we are planning to update in future the uncertainty analysis conducted in 2012. The main objective of that uncertainty analysis, complying with the 2006 IPCC Guidelines, was to identify possible major sources of errors, and to indicate, based on the prioritization of the uncertainty of the estimates, where efforts on development should concentrate in future inventories. The uncertainty analysis focused on the uncertainty of carbon stock change estimates of the biomass of forests for the categories under the Kyoto Protocol, therefore, the detailed results can be found in Chapter 11 (Section 11.3.1.5). As the methods of quantitative estimation are similar to respective categories under the UNFCCC, and because KP and UNFCCC categories significantly overlap, we regard the results reported there relevant for the uncertainties of emissions and removals under the UNFCCC, and only some additional notes are made here.

In general, information on uncertainties includes, among others, information on completeness, accuracy, and non-quantifiable elements. Concerning *completeness*, some minor emissions and removals could not be estimated because of the reasons provided in the respective sections above (and/or were approximated by assumptions), however, it is highly probable that their exclusion only results in conservative estimation, i.e., overestimation of net emissions, and underestimation of net removals.

The reported estimated emissions and removals are generally considered accurate and precise as far as practicable and are based on the best available data and methods. Where uncertainty seems to be high, and for *non-quantifiable factors*, the principle of conservativeness is always applied. Conservative estimates are used for volume stock change, the root-to-shoot ratio, and carbon stock changes in the soil, litter and deadwood pools. Where no country-specific values are available, IPCC default values are used. Whenever more accurate methods could be identified, these were applied (see section 6.1.4 on recalculations above).

It is probable that total forest area is somewhat underestimated, which is shown by the fact that the forest

inventory has identified new forest areas (“found forests”) most years for the last two decades or so. As long as forests in Hungary are a sink, this underestimation of the forest area can only lead to the underestimation of net removals, especially because found forests are predominantly young and have a positive carbon balance. Nevertheless, the detection and monitoring of forest area has been continuously improving and will continue to improve.

It is also probable that, due to conservativeness built into the methods of the national forest inventory to comply with traditional requirements for sustained yield, both volume stocks and volume stock changes are underestimated. This assessment is also supported by preliminary statistical results of a sample-based inventory (i.e., the National Forest Inventory) which indicate higher volume stocks and higher annual volume increment than the continuous forest inventory that is currently used for the purposes of the GHG inventory. Finally, wood harvests also seem to be underestimated a bit due to illegal cuttings which, according to some expert judgments, may account for up to 250,000 m³ annually. This amount is additional to, but small relative to, the annual official figure of annual harvests of around 7-8 million m³. Although this means that actual wood harvests are somewhat underestimated, so is volume stock increment but to a larger extent, thus, net volume stock changes net removals are most probably underestimated.

We have continuously been improving not only our stand statistics, but also our country-specific emission factors. As reported in the previous NIRs, the accuracy was improved earlier, among others, by introducing new, more realistic, country-specific basic wood density values, slash fraction estimates, soil C/N values and the biomass of orchards and vineyards that have been removed during conversions of cropland to forest land and grassland.

Uncertainty cannot always be quantified partly because the error distributions are unknown due to lack of measured data, and partly because calculation errors or assumptions cannot be quantified. However, calculation errors during the development of the GHG inventory are highly unlikely due to the double-checking of the data processing as described in the next section.

For carbon stock changes in biomass, the system of calculations allows for the use of a simple sensitivity analysis. This is especially true if only the major sources of CO₂ emissions and removals are considered, which represent the bulk of all emissions and removals. The reason for this is that the equation used for the calculation is simple: only volume stock changes, wood density, root-to-shoot ratio, and carbon fraction factors are involved.

With respect to net *annual* CO₂ emissions (i.e., net removals), actual values may deviate from estimated values as the stock volume inventory for the whole country is not able to capture all inter-annual variability of timber growth and harvests, which can be high due to the variability of meteorological conditions. Note that, as mentioned above, the inter-annual variability of the estimated net removals in the Forest Land sector is due to a number of reasons, including the continuously, although slowly, changing structure of the forests by species, site fertility and age, and the annual variability of weather, harvests and mortality. All these effects have rather different delayed effects on the inventory estimates, and these effects may be rather small relative to the total volume stocks. These effects can reinforce or naturalize each other when combined.

It can be concluded that in general, with regard to carbon stock change estimation in the forestry sector, errors are rather limited, and it is expected that current estimates rather well reflect actual emissions and removals associated with forest land.

With regard to non-CO₂ emissions, the estimation is accurate and precise as far as practicable for the years for which we have data on wildfires and controlled burning, as well as for forests on organic soils. Data collection considerably improved in the last few years.

Finally, both methods and data are applied consistently throughout the entire reporting period. This

results in a consistent time series of both the area and the GHG information.

This year, we conducted an uncertainty analysis for the entire LULUCF section focussing on the non-forest categories, with the forestry categories only serving as references to which the importance of the other categories are compared. See section 6.11 for more details.

6.5.8 Category-specific QA/QC and verification

The calculations to obtain emission and removal estimates are generally based on the activity data taken from the National Forest Database, and the databases of the Forest Authorities on afforestations and deforestations. These databases are the most accurate ones in the country on the forests. The first complete and country-wide inventory was accomplished in 1976. Forests have been continuously monitored since that year, and the responsible authorities have been applying computer-based information technology for data management since the early '80-s. The database is updated annually, and the data is checked by many people at subsequent procedures from field assessment to data processing. The constant development of field methods and informatics, the improvement of checks, and the increasing requirements to ensure the quality of work resulted in the increasing accuracy of the Database in recent years.

The GHG inventory has currently been completed by the Forestry Department of the National Land Authority, i.e., the institute that runs the National Forest Database and other mentioned databases.

As a quality assurance, double-checking of the data processing of the calculations involved in the GHG inventory and the correct application of IPCC assumptions and methodologies have been performed at the national level by the Hungarian Forest Research Institute of Sopron University for years. The separation of the two roles (i.e., the preparation and the QA of the GHG inventory) improved the data quality. Final checks and integration of the data into the GHG inventory was performed by the Hungarian Meteorological Service, i.e., the institute responsible for the entire national GHG inventory.

Data verification was and is continuously conducted concerning activity data (see the comparison of volume stock changes with trends of wood volume increment and harvest in Section 6.5.4.2.1, and also previous NIRs of Hungary). All information used for the development of the GHG information is archived by the inventory agency. Thus, the correctness of the estimation methodology is in principle *verifiable*.

For other activities, see section 6.6.5 where activities are listed that have also been conducted in the forestry sector.

6.5.9 Category-specific recalculations

As noted in section 6.1.4, we have done a number of recalculations in the forestry sector to improve the accuracy of our estimates and to correct upload errors. These recalculations for FL-FL and L-FL as well as for the HWP pool, their reason and scale are detailed in Tables 6.5.15, 6.5.16. and 6.5.17., respectively. Note that the scale of difference expressed in percent may be misleading due to the reference which in the case of (gross or net) removals can be close to zero and does not necessarily represent a useful reference for the calculation of percentages.

Table 6.5.15. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for the FL-FL category for some recent inventory years (Note that zero values mean that there is no recalculation for that year).

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.A.1 Carbon stock change, Total area	ktC	1865	1877	1884	1893	1904	1910	1913	1916	1921	1927	1931	1937	revision of area data and revision of reporting practice
2022		ktC	1768	1781	1791	1803	1816	1823	1826	1834	1842	1850	1858	1870	
	difference, %	%	-5.2	-5.1	-4.9	-4.8	-4.6	-4.6	-4.6	-4.3	-4.1	-3.9	-3.8	-3.4	
2021	4.A.1 Carbon stock change, Area of mineral soil	ktC	1858	1870	1878	1887	1897	1903	1907	1910	1914	1920	1925	1930	revision of area data and revision of reporting practice
2022		ktC	1761	1774	1785	1796	1809	1816	1819	1827	1836	1844	1851	1864	
	difference, %	%	-5.2	-5.1	-4.9	-4.8	-4.6	-4.6	-4.6	-4.3	-4.1	-4.0	-3.8	-3.4	
2021	4.A.1 Carbon stock change, Area of organic soil	ktC	0	0	0	0	0	0	0	0	0	0	0	0	NA
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	0	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
2021	4.A.1 Carbon stock change, Gains	kt C	906	664	588	575	849	631	938	1099	881	1028	866	1060	revision of area data and the ratio of success initial planting
2022		kt C	790	524	516	477	690	529	799	1025	802	912	768	957	
	difference, %	%	-12.8	-21.1	-12.2	-17.0	-18.8	-16.2	-14.9	-6.8	-8.9	-11.3	-11.4	-9.7	
2021	4.A.1 Carbon stock change, Losses	kt C	0	0	0	0	0	0	0	0	0	0	0	0	revision of area data and the ratio of success initial planting
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	0	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
2021	4.A.1 Carbon stock change, Net change	kt C	906	664	588	575	849	631	938	1099	881	1028	866	1060	see above
2022		kt C	790	524	516	477	690	529	799	1025	802	912	768	957	
	difference, %	%	-12.8	-21.1	-12.2	-17.0	-18.8	-16.2	-14.9	-6.8	-8.9	-11.3	-11.4	-9.7	
2021	4.A.1 Carbon stock change, Net carbon stock change in dead wood	kt C	59	57	55	54	61	56	58	73	52	60	42	162	revision of area data
2022		kt C	59	57	55	54	61	55	57	73	52	60	42	49	
	difference, %	%	-0.5	-0.5	-0.5	-0.5	-0.4	-0.4	-0.4	-0.3	-0.3	-0.3	-0.3	-69.6	

Table 6.5.16. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for the L-FL category. (Note that zero means, there is no recalculation for that year).

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.A.2 Carbon stock change, Total area	ktC	166	163	162	158	152	150	148	144	138	131	124	118	revision of area data and revision of reporting practice
2022		ktC	263	259	255	248	240	237	235	227	217	207	197	184	
	difference, %	%	58.1	59.3	57.2	57.2	57.9	58.2	58.7	57.2	57.1	58.2	58.9	56.5	
2021	4.A.2 Carbon stock change, Gains	kt C	335	320	324	305	292	289	292	287	278	269	261	253	revision of ratio of successful initial planting
2022		kt C	344	328	332	312	299	296	299	293	284	274	265	257	
	difference, %	%	2.6	2.5	2.4	2.4	2.4	2.4	2.4	2.2	2.0	1.6	1.6	1.4	
2021	4.A.2 Carbon stock change, Losses	kt C	-34	-17	-27	-8	-6	-4	-6	0	-1	-4	0	-8	revision of area data
2022		kt C	-34	-17	-27	-8	-6	-4	-6	0	-1	-3	0	-8	
	difference, %	%	0.0	0.1	0.1	-0.3	-0.1	-0.1	-4.2	0.0	44.4	-9.7	0.0	0.0	
2021	4.A.2 Carbon stock change, Net change	kt C	302	303	297	296	286	285	286	286	277	266	255	246	see above
2022		kt C	310	311	304	304	293	292	293	292	283	270	259	249	
	difference, %	%	2.8	2.7	2.6	2.5	2.5	2.4	2.3	2.2	1.9	1.8	1.6	1.5	
2021	4.A.2 Carbon stock change, Net carbon stock change in dead wood	kt C	11	11	11	11	11	10	10	10	9	9	9	8	revision of area data
2022		kt C	11	11	11	11	11	11	11	10	9	9	9	8	
	difference, %	%	2.8	2.6	2.6	2.4	2.4	2.4	2.4	2.1	1.8	1.7	1.5	1.4	
2021	4.A.2 Carbon stock change, Net carbon stock change in litter	kt C	73	71	71	69	67	0	0	0	61	57	55	52	revision of area data
2022		kt C	73	71	71	69	67	0	0	0	61	57	55	52	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	
2021	4.A.2 Carbon stock change, Mineral soils	kt C	46	49	52	52	53	53	53	51	49	48	23	21	revision of area data and correction in formulas
2022		kt C	32	31	32	31	31	30	29	28	26	24	22	20	
	difference, %	%	-31.3	-35.2	-38.6	-40.5	-42.4	-44.0	-44.7	-45.8	-47.5	-49.8	-50.9	-2.9	

Table 6.5.17. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for the HWP pool.

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.G.B Total HWP from Domestic, Gains	ktC	0	0	0	0	0	0	0	0	0	0	0	315964	revision of production data
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	332178	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.1	
2021	4.G.B Total HWP from Domestic, Losses	ktC	0	0	0	0	0	0	0	0	0	0	0	-238974	revision of production data
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	-240766	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	
2021	4.G.B Total HWP from Domestic, Annual change in stock	ktC	0	0	0	0	0	0	0	0	0	0	0	77	revision of production data
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	91	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	18.7	
2021	4.G.B Total HWP from Domestic, HWP in use	kt C	0	0	0	0	0	0	0	0	0	0	0	-282	revision of production data
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	-335	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	18.7	
2021	-3, Gains	ktC	0	0	0	0	0	0	0	0	0	0	0	35567	revision of production data
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	46700	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	31.3	
2021	-3, Losses	ktC	0	0	0	0	0	0	0	0	0	0	0	-27209	revision of production data
2022		ktC	0	0	0	0	0	0	0	0	0	0	0	-28934	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.3	
2021	-3, Annual change in stock		0	0	0	0	0	0	0	0	0	0	0	8	revision of production data
2022			0	0	0	0	0	0	0	0	0	0	0	18	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	112.6	
2021	-3, HWP in use		0	0	0	0	0	0	0	0	0	0	0	-31	revision of production data
2022			0	0	0	0	0	0	0	0	0	0	0	-65	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	112.6	

6.5.10 Category-specific planned improvements

We will continue the revision of our databases that are the sources of activity data to ensure that estimates are as accurate as possible. We are also planning to revise the entire estimation system for any change required by reporting under the Paris Agreement. Finally, we are planning to implement the results of the recently run revision of soil carbon stock changes in our next submission.

6.6 Cropland (CRF sector 4.B)

6.6.1 Description of category

Although the area of croplands decreased in the last three decades (roughly 800,000 hectares were abandoned or converted to another land-use category, after which only parts of this loss was regained), croplands with their 56% proportion of the total area of the country still represent the main land-use category in Hungary (see Figure 6.3.1 above). All lands with annual crops, orchards and vineyards (i.e., perennial woody crops) and kitchen gardens are classified as cropland. Set-aside croplands are also reported in this category (Figure 6.6.1.) The distribution of emissions and removals by sub-categories is reported in Figure 6.6.2.




Figure 6.6.1. The distribution of Cropland area since 1985.




Figure 6.6.2. Emissions and removals in Cropland 1985-2020. The data under the grey box is shaded out for reasons explained in section 6.1.1.

6.6.2 Cropland remaining Cropland

Figure 6.6.3 reports emissions and removals by pool and gas, whereas Table 6.6.1 reports methodological information for this sub-category. Figure 6.6.4 is reported to demonstrate that most removals in the Cropland category (including the CL-CL category) arise due to changes in land-use either between sub-categories or within a sub-category, i.e., from non-set-aside to set-aside land and back.




Figure 6.6.3. Emissions and removals in CL-CL 1985-2020. The data under the grey box is shaded out for reasons explained in section 6.1.1.




Figure 6.6.4. The trend of annual area of the most important land-use change categories involving croplands.

Table 6.6.1. Methodological summary for CL-CL (CS=country specific; D: default; AD: activity data; EF: emission/removal factor; NO=not occurring).

METHODOLOGY MATRICES TO BE INCLUDED IN NIR									
CL-CL	Type of information	Carbon stock changes						Table (4)III, (4)IV	Table (4)V
		BIOMASS		DOM		SOIL			
		annual	perennial	DW	LI	mineral	organic		
	E/R	Tier 1: 0	AD: CS; EF: CS and D	Tier 1: 0		AD: CS; EF: CS/D	NO	D	Wildfires: AD: CS; EF: D
									Biomass burning NO
Uncertainty	Tier 2		NE		Tier 2		Tier 2		

6.6.2.1 Biomass

Consistent with the 2006 IPCC GL, carbon stock changes of biomass are only estimated for perennial woody crops of orchards and vineyards (the biomass of annual crops is assumed to be in equilibrium). This chapter thus reports on emissions and removals from biomass in areas that remained orchards or vineyards in the inventory year. (Carbon stock changes from loss of perennial biomass due to land-use change are accounted for in the respective conversion sub-categories.)

Carbon stock change of biomass (ΔC_{Biom}) was estimated applying Equation 2.15 of the 2006 IPCC GL:

$$\Delta C_{\text{Biom}} = \Delta C_{\text{G}} + \Delta C_{\text{Conversion}} - \Delta C_{\text{loss}}$$

where

ΔC_{Biom} = annual change in carbon stocks of biomass, tonnes C yr⁻¹

ΔC_G = annual increase in carbon stocks due to biomass growth, tonnes C yr⁻¹

ΔC_L = annual decrease in carbon stocks due to biomass loss, tonnes C yr⁻¹

$\Delta C_{\text{Conversion}}$ = annual carbon loss due to converting perennials to other land-use, tonnes C yr⁻¹.

The estimation is done separately for orchards and vineyards and for growth and losses, and for losses due to conversions where perennials are converted to annual croplands or other land-use.

6.6.2.1.1 Growth and loss

Carbon stock changes due to growth and loss were estimated using Equation 2.7 of the 2006 IPCC GL:

$$\Delta C_{\text{Biom}} = \Delta C_G - \Delta C_L$$

Orchards and vineyards are assumed to be grown in rotations of 30 and 31.8 years, respectively. For the entire period, ΔC_G was estimated using Equation 2.9 of the 2006 IPCC GL:

$$\Delta C_G = A_{\text{perennials}} * G_{\text{TOTAL}} * CF$$

where

$A_{\text{perennials}}$ = area of orchards and vineyards at the beginning of the inventory year, ha (data is taken from the statistics of the HCSO),

G_{TOTAL} = county-specific net biomass accumulation rate (0.313 and 0.626 t biomass ha⁻¹ yr⁻¹ for orchards and vineyards, respectively), and

CF = carbon fraction (the default value of 0.5 tC t biomass⁻¹ is used).

G_{TOTAL} was estimated in the detailed study of Juhos and Tőkei (2013). Consistent with section 6.5.5.2.1, as it was not possible to measure below-ground biomass, G_{TOTAL} only includes above-ground biomass (which is the application of the default assumption, according to which there is no change in below-ground biomass of perennial trees in agricultural systems).

The annual decrease in carbon stocks from biomass loss due to regenerating perennials was estimated using Equation 2.16 of the 2006 IPCC GL:

$$\Delta C_L = A_{\text{regenerated_perennials}} * B_{\text{Before}} * CF$$

where

$A_{\text{regenerated_perennials}}$ = the area of regenerated orchard or vineyard in the inventory year =
= $(A_{\text{perennials}} - A_{\text{conv}}) / \text{RPL}$, ha

A_{conv} = area if orchards and vineyards that are converted to other land-use categories in the inventory year (see below), ha

RPL = length of rotation period, 30 years (orchards) and 31.8 years (vineyards), and

B_{Before} = biomass of the regenerated orchard or vineyard at the end of the rotation period, t biomass, which is equal to G_{TOTAL} (tC ha⁻¹ yr⁻¹) * RPL (years). (Since all biomass is considered lost during the regeneration, the “fraction of biomass lost in disturbance” term, or fd , in the original equation of the 2006 IPCC GL is taken to be equal to 1.)

Note that the above methodology implies that removals due to slight increases of the area of perennials (this has only happened in a few years and on small areas) are all accounted for in the CL-CL category (i.e., the assumption is applied that all increases of the area of perennials are due to conversions from annual cropland).

6.6.2.1.2 Conversions

Total emissions from biomass from converting orchards and vineyards are estimated applying the methodology described in section 6.4.4 (i.e., Equation 2.16 of the 2006 IPCC GL) with

$B_{\text{After}} = 0$, and
 B_{Before} and CF as above.

A_{conv} was estimated using an estimated proportion of converted perennials, P_P , that remained in the Cropland category:

$$A_{\text{conv}} = A_{\text{perennials}} * P_P.$$

P_P for Cropland was estimated from the land statistics database (see section 6.3.2 for details).

The areas removed are calculated from actual reduction, if any, of the total areas and thus the estimation of these areas is fully consistent with that of areas where growth occurs.

6.6.2.2 Dead organic matter

For DOM, the Tier 1 method is applied. This method assumes that the dead wood and litter stocks are not present in Cropland or are at equilibrium as in agroforestry systems and orchards. Thus, the carbon stock changes are not reported for these pools.

6.6.2.3 Mineral soils

The method and emission factors used are those described in section 6.4.1. For CL-CL, what may cause changes of the mineral soil carbon stocks are the following: changes in management, changes in input, and converting non-set aside to set-aside and back. Using country-specific SOC_{REF} and default F_{LU} , F_{MG} and F_{I} values, the effect of all these changes and conversions is included in the calculations.

We highlight here that CL-CL is a category in Hungary that is split into non-set-aside and set-aside CL. There is difference in the assumed carbon stocks of these sub-categories (see Table 6.4.2 of the NIR) and when land is converted from one sub-category to the other, the soil SOC changes with a dynamics of over a 20-year period according to the IPCC default method. There were quite large areas converted to the set-aside sub-category even before 2011 (but in some years, later, too), and the overall effect of all these factors is reflected in the IEF values for the entire CL-CL category. In other words, even if the total area of the CL-CL category remains, there may be net emissions or removals from the soil if the distribution of the area of the SA and non-SA sub-categories changes over time.

Concerning the management of soils, the most recent data on the share of no-till and other reduced impact methodologies is available from 2016, which are incorporated in the GHG calculations.

6.6.2.4 Non-CO₂ emissions

The amount of non-CO₂ emissions is estimated according to section 6.4.2 (for N₂O emissions from soils) and 6.4.3 (for emissions from wildfires).

For the mass of available fuel (M_B) in the wildfire calculation, no proper country-specific values have been derived yet, therefore, a default value of 10 t d.m. ha⁻¹ was assumed. This is a value taken from Table 2.4 of the 2006 IPCC GL for maize, which is quite a representative crop in Hungary, and can be considered as a conservative value as wheat and other crops of less biomass are also abundant. For the combustion factor C_f , and for the GHG-specific emission factors, default data (for maize residues in Table 2.6, and those in Table 2.5 of the 2006 IPCC GL, respectively) are used.

6.6.3 Land converted to Cropland

Figure 6.6.5 reports emissions and removals, whereas Table 6.6.2 reports methodological information for this sub-category. Note that non-CO₂ emissions from wildfires, if any, are reported as IE and accounted for in the CL-CL category. Emissions from burning is NO.




Figure 6.6.5. Emissions and removals in L-CL since 1985. The data under the grey box is shaded out for reasons explained in section 6.1.1.

Table 6.6.2. Methodological summary for L-CL. (CS=country specific; D: default; IE: included elsewhere; AD: activity data; EF: emission/removal factor)

L-CL	Type of information	"FROM" category	BIOMASS	DOM		SOIL		Table (4)III, (4)IV	Table (4)V	
				DW	LI	mineral	organic			
	E/R	FL	AD: CS; EF: CS	AD: CS; EF: CS		AD: CS; EF: CS/D		NO	D	slash burning: I (FL-FL)
		GL	AD: CS; EF: D	Tier 1: 0		AD: CS; EF: CS/D		NO		IE (CL-CL)
		WL	NO	NO		NO		NO		IE (CL-CL)
		SE	Tier 1: 0	Tier 1: 0		AD: CS; EF: CS/D		NO		IE (CL-CL)
		OL	NO	NO		NO		NO		IE (CL-CL)
Uncertainty	Tier 2		NE		Tier 2		Tier 2			

6.6.3.1 Forest Land converted to Cropland

For the methodology to estimate carbon stock changes in the biomass, DOM and soil pools, see Sections 6.5.6.1.1, 6.5.6.1.2 and 6.5.6.1.3, respectively. Note that the total emissions from these pools were split between FL-CL and other conversions by the area of these conversions. The share of FL-CL to all FL-L is, on average, 22% but increased to 61% in 2017.

6.6.3.2 Grassland converted to Cropland

6.6.3.2.1 Biomass

Equations 2.15 of the 2006 IPCC GL were applied as follows:

$$\Delta C_B = \Delta C_G + \Delta C_{\text{CONVERSION}} - \Delta C_L$$

where:

ΔC_B = biomass carbon stock change due to land-use conversion, tC year⁻¹,

ΔC_G = annual increase in carbon stocks in biomass due to growth on the 'converted to' land, tonnes C yr⁻¹,

ΔC_L = annual decrease in biomass carbon stocks due to losses, tonnes C yr⁻¹,

$\Delta C_{\text{CONVERSION}}$ = initial change in carbon stocks in biomass on the 'converted to' land, tonnes C yr⁻¹, estimated using Equation 2.16 of the 2006 IPCC GL as described in section 6.4.4.

For A_{Conv} , data from the annual land-use change matrix was used. B_{before} was estimated from the proportion of Grassland area of cold dry and warm dry climate types ($P_{\text{CD}} = 0.41$, $P_{\text{WD}} = 0.59$) and respective specific default Grassland biomass (in order that all biomass is accounted for, the following total above- and below-ground biomass values were taken from Table 6.4 of the 2006 IPCC GL: $B_{\text{CD}} = 6.5$ t biomass ha⁻¹ and $B_{\text{WD}} = 6.1$ t biomass ha⁻¹, respectively). The resulting weighted biomass is: $B_{\text{before}} = P_{\text{CD}} * B_{\text{CD}} + P_{\text{WD}} * B_{\text{WD}} = 6.26$ t biomass ha⁻¹ (see section 6.7.2.1 for more details). In accordance with the Tier1 assumption, B_{after} in the equation is 0, and the carbon fraction is the default value of 0.47 tC t biomass⁻¹ (page 6.29 of the 2006 IPCC GL). For ΔC_G , the value of 4.7 tC ha⁻¹ was used, whereas ΔC_L was assumed to be equal to 0.

6.6.3.2.2 Mineral soils

The method and emission factors used are those described in section 6.4.1.

6.6.3.3 Wetlands converted to Cropland

This land-use change is not occurring in Hungary.

6.6.3.4 Settlements converted to Cropland

For the rather small conversion areas in this category, only carbon stock changes in mineral soils are estimated. The method and emission factors used are those described in section 6.4.1.

6.6.3.5 Other Land converted to Cropland

This land-use change is not occurring in Hungary.

6.6.4 Uncertainties and time-series consistency

See section 6.11.

6.6.5 Category-specific QA/QC and verification

Emissions/removals were estimated by the National Food Chain Safety Office, whereas the QA/QC was done by an external expert. This division of tasks made it possible to separate the work related to emission estimation and the QA/QC procedures.

The LULUCF QC measures are based on the QC procedures as described by Chapter 5 of the 2006 IPCC Guidelines.

The main checks carried out are related to the following issues:

Activity data:

- Methodological issues of the collection of the land-use / land-cover data.
- The differences between the different land-use datasets.
- Consistency of the activity data. In the case of inconsistency (methodological change in the data collection) the dataset is adjusted in consultation with the data provider.
- Data inputs for transcription errors.
- The units of activity data in the calculation sheets throughout the emission calculation.
- Consistency of the total area of Hungary in the land-use change matrices and the CRF tables.
- The comparison of activity data with data from other sources, if possible.

Methodology:

- The applied methodologies and emission factors against the 2006 IPCC GL.
- The correctness of the equations and factors in the calculation sheets.
- The consistency of the applied methodology throughout the entire time series.

Emissions and removals:

- Reported emissions for transcription errors between the calculation sheets and the CRF tables.
- Recalculation differences and reasons for recalculations.

6.6.6 Category-specific recalculations

This year, only a few recalculations of small effect were made for the CL category as detailed in Table 6.6.3.

Table 6.6.3. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for (a) the CL-CL and (b) the L-CL category for some recent inventory years.

(a)

Sub mission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	CL-CL		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.B.1 Carbon stock change, Total area	ktC	0	0	0	0	5083	5085	5090	5095	5102	5106	5113	5118	revision of the LUC matrix
2022		ktC	0	0	0	0	5083	5084	5087	5093	5099	5103	5108	5113	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	-0,1	-0,1	-0,1	-0,1	-0,1	-0,1	
2021	4.B.1 Carbon stock change, Area of mineral soil	ktC	0	0	0	0	5083	5085	5090	5095	5102	5106	5113	5118	revision of the LUC matrix
2022		ktC	0	0	0	0	5083	5084	5087	5093	5099	5103	5108	5113	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	-0,1	-0,1	-0,1	-0,1	-0,1	-0,1	
2021	4.B.1 Carbon stock change, Losses	kt C	0	0	0	0	0	-40	0	0	0	0	-54	-61	revision of the LUC matrix as well as calculation error
2022		kt C	0	0	0	0	0	-40	0	0	0	0	-53	-62	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	-2,2	1,7	
2021	4.B.1 Carbon stock change, Mineral soils	kt C	0	0	0	305	281	255	227	200	184	167	145	132	revision of the LUC matrix as well as calculation error
2022		kt C	0	0	0	297	274	249	222	195	179	162	140	112	
	difference, %	%	0,0	0,0	0,0	-2,5	-2,4	-2,4	-2,5	-2,8	-2,8	-3,0	-3,5	-14,9	

(b)

Sub mission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.B.2 Carbon stock change, Total area	ktC	0.0	0.0	0.0	0.0	0.0	0.0	106.9	100.9	95.5	91.2	88.8	83.3	revision of the LUC matrix
2022		ktC	0.0	0.0	0.0	0.0	0.0	0.0	107.2	101.2	95.8	91.5	89.1	83.6	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.4	
2021	4.B.2 Carbon stock change, Area of mineral soil	ktC	0.0	0.0	0.0	0.0	0.0	0.0	106.9	100.9	95.5	91.2	88.8	83.3	revision of the LUC matrix
2022		ktC	0.0	0.0	0.0	0.0	0.0	0.0	107.2	101.2	95.8	91.5	89.1	83.6	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.4	
2021	4.B.2 Carbon stock change, Net change	kt C	11.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	revision of the LUC matrix as well as calculation errors
2022		kt C	-1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
		difference, %	%	-109.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
2021	4.B.2 Carbon stock change, Net carbon stock change in dead organic matter	kt C	-1.3	-0.8	-3.3	-1.0	-1.7	-1.4	-1.9	-4.9	-11.5	-13.2	-9.9	-9.4	revision of the LUC matrix as well as calculation error
2022		kt C	-1.4	-0.9	-3.5	-1.1	-1.8	-1.5	-6.0	-5.1	-12.2	-14.0	-6.3	-10.0	
		difference, %	%	5.2	5.4	5.5	5.7	5.8	5.9	214.6	6.1	6.2	6.3	6.2	
2021	4.B.2 Carbon stock change, Mineral soils	kt C	-82.4	-80.3	-76.3	-78.2	-80.0	-80.9	-80.9	-76.1	-71.6	-67.4	-65.3	-60.7	revision of the LUC matrix
2022		kt C	-82.4	-80.3	-76.3	-78.4	-80.4	-81.8	-81.8	-77.1	-72.7	-68.8	-66.7	-62.2	
		difference, %	%	0.0	0.0	0.0	0.2	0.4	0.7	1.1	1.3	1.6	2.0	2.1	

6.6.7 Category-specific planned improvements

Planned improvements include the implementation of specific elements of the result of the Tier 2 uncertainty estimation, including the development based on new soil carbon stock change factors.

6.7 Grassland (CRF sector 4.C)

6.7.1 Description of category

In 1985, the livestock of grazing animals included 2 million cattle, 1 million geese and 3 million sheep. The decade beginning 1980 both saw the highest number of grazing livestock in the country and was the period of the most intensive management of the Hungarian grasslands with respect to fertilizer doses and irrigation. The number of grazing animals and the intensity of grassland management started to decrease after about the mid-1980's and reached its bottom in the middle of the 1990's. All this also affected the area of grassland which considerably decreased after 1985, but started to increase again beginning 2011, and amounted to roughly 9 percent of the official area of Hungary in 2020 (Figure 6.7.1). Of the emissions and removals due to changes to and from grasslands, those that are accounted for in the Grassland category are reported in Figure 6.7.2.




Figure 6.7.1. The area of the Grassland category since 1985.




Figure 6.7.2. Emissions and removals in the Grassland category since 1985. The data under the grey box is shaded out for reasons explained in section 6.1.1.

6.7.2 Grassland remaining Grassland

Figure 6.7.3 reports emissions and removals, whereas Table 6.7.1 reports methodological information for this sub-category.




Figure 6.7.3. Emissions and removals in the GL-GL category since 1985. The data under the grey box is shaded out for reasons explained in section 6.1.1.

Table 6.7.1. Methodological summary for GL-GL (CS=country specific; D: default; AD: activity data; EF: emission/removal factor).

GL-GL	Type of information	Carbon stock changes						Table (4)III, (4)IV	Table (4)V
		BIOMASS		DOM		SOIL			
		annual	perennial	DW	LI	mineral	organic		
	E/R	Tier 1: 0	Tier 1: 0	Tier 1: 0	Tier 1: 0	AD: CS; EF: CS/D	NO	D	Wildfires: AD: CS; EF: D Biomass burning: NO
Uncertainty	Tier 2		NE		Tier 2		Tier 2		

6.7.2.1 Biomass

Grasslands are meadows and pastures some of which are grazed or harvested annually, while others are unmanaged, and where tree cover is non-existent or very low. Due also to its relatively small area and dynamics, the biomass of GL-GL is not a key category. Therefore, we adapt the Tier 1 method of the 2006 IPCC GL which assumes no change in the biomass carbon stocks. In line with this, 'NO' is reported for the biomass of this category. Note that, due to the increase of set-aside grassland, a rather slow increase in woody biomass might have been occurring, thus, applying the Tier 1 method might slightly underestimate carbon removals.

Note that a domestic study (Tasi et al. 2016) reports highly variable above-ground biomass, but no estimates for the below-ground biomass. The change of grassland management, such as from non-set-aside grassland to set-aside grassland and vice versa, does not lead to a change in grassland productivity in biomass. This information together with expert judgment suggests that the total biomass is around the default IPCC value. Therefore, we apply the IPCC default value for the biomass of grasslands.

6.7.2.2 Dear organic matter

As the dead organic matter pool and its carbon stock changes are relatively small, the Tier 1 method is applied which assumes that the dead wood and litter stocks are at equilibrium, and the carbon stock changes for these pools are thus not estimated.

6.7.2.3 Soils

Some direct local results have already been published concerning CO₂-emission from grasslands (Nagy et al. 2007, Zsembeli et al. 2006). However, in lack of country-wide monitoring results the Tier 2 IPCC method is applied which, together with emission factors, is described in section 6.4.1. Data demonstrate that high activity clay mineral soils are dominant, similar to the case of croplands. (These include salt affected soils that are very characteristic to Hungary and that are partly utilized as grasslands, mainly depending on the extent of salinization.)

Concerning management, sufficient statistics are lacking for the period 1985-2002. As an approximation, the management, hence the quality of grasslands is determined for this period based on the number of grazing animals and the level of management costs for each soil type and climate region, taking into consideration the spatial distribution of the number of livestock by species. The spatial distribution of quality, utilization and load of grasslands were estimated and overlaid on the genetic soil maps and climatic zone maps mentioned in section 6.4.1. Based on this, the following two broad categories are used to characterize the management of the Hungarian grasslands: nominally managed (non-degraded) grasslands (with no input), which includes pastures, rangelands and other unmanaged grasslands, and improved grasslands with medium input. The area of the latter can be calculated from HCSO data based on the area of grasslands treated with chemical fertilizers (Table 6.7.2) and that of irrigated grasslands which are available for some inventory years since 2003. The proportion of irrigated grasslands is less than 0.1 per cent, therefore, the area of grasslands treated with chemical fertilizers is considered to represent improved grasslands in Hungary, and the rest is taken as nominally managed.

Table 6.7.2. The area of grasslands treated with chemical fertilizer as well as the amount of active ingredient applied to grasslands for recent years (until 2019, i.e., the last year with datasource: HCSO: https://www.ksh.hu/stadat_files/mez/hu/mez0042.html).

Year	Grasslands treated with chemical fertilizer [ha]	Amount of active ingredient used (kg/ha)		
		N	P	K
2003	22361			
2004	21290			
2005	no data			
2006	no data			
2007	10114			
2008	16412			
2009	8962			
2010	8774			
2011	11441			
2012	11494	50.3	1.2	1.4
2013	12387	51.9	2.7	3.3

2014	11675	68.3	2	2
2015	13233	50.5	2	3.9
2016	NA	57.8	4.2	5.5
2017	NA	52.6	5.9	5.1
2018	NA	49.4	5.6	6.2
2019	NA	49.1	6.8	6.1

The management of grasslands was reduced due to the introduction of Agro-environmental Management Program in 2002-2003 and was limited to slightly intensive planted grasslands. This program resulted in the natural succession of pastures that is characterized by the propagation of weeds and soil degradation. The management of grasslands is limited to their grazing and cutting.

All the above information formed a sufficiently good basis for the expert judgment that was necessary to develop the required proportions (Table 6.7.3).

Table 6.7.3. *The distribution of grasslands in Hungary by management and input.*

Calendar year	Proportion of total grassland area (%)	
	nominally managed (non-degraded): no input	improved: medium input
1985	0,641	0,359
1986	0,639	0,361
1987	0,627	0,373
1988	0,626	0,374
1989	0,631	0,369
1990	0,654	0,346
1991	0,657	0,343
1992	0,681	0,319
1993	0,741	0,259
1994	0,817	0,183
1995	0,859	0,141
1996	0,886	0,114
1997	0,920	0,080
1998	0,939	0,061
1999	0,952	0,048
2000	0,957	0,043
2001	0,970	0,030
2002	0,973	0,027
2003	0,977	0,023
2004	0,977	0,023
2005	0,981	0,019
2006	0,984	0,016
2007	0,988	0,012
2008	0,980	0,020
2009	0,989	0,011
2010	0,988	0,012
2011	0,988	0,012
2012	0,988	0,012
2013	0,987	0,013
2014	0,985	0,015
2015	0,983	0,017
2016	0,990	0,013
2017	0,990	0,010
2018	0,989	0,011
2019	0,990	0,010
2020	0,985	0,011

Note that carbon stock changes for mineral soils (for both grasslands and elsewhere) depend on both the area (activity data) and the appropriate emission factor that in turn depends on the management practice. Since the distribution shows a change over time, and the factor F_{mg} is different for the non-degraded and the improved grasslands, it follows that carbon stock changes are not zero (but rather small anyways). The proportions in Table 6.7.3 are expert judgments for the last several years, and we will correct the historical values when better data will be available at the next census.

For the above climate, soil, management and input categories, the applied reference soil organic carbon stocks were the same as those reported in Table 6.4.2 above. The land-use factor (F_{LU}) is 1.0 for all grasslands, whereas the selected management factors (F_{MG}) are reported in Table 6.7.4 and the level of input (F_I) was assumed to be 1.0 for both the nominally managed grasslands and the improved grassland (IPCC 2006).

Note that the above methodology may result in zero carbon stock change estimate in some years. To demonstrate that it is the result of a calculation, it is reported as zero and not as “NO”.

Table 6.7.4. Management factors (F_{MG}) applied for Grassland

Management regime	F_{MG}
Nominally managed (non-degraded)	1.00
Improved	1.14

6.7.2.4 Non-CO₂ emissions

The amount of non-CO₂ emissions is estimated according to section 6.4.2 (for N₂O emissions from soils) and 6.4.3 (for emissions from wildfires).

For the mass of available fuel (M_B) in the wildfire calculation, no proper country-specific values have been derived yet, therefore, the value of 6.26 t d.m. ha⁻¹ was assumed, which is the same that we use in the GL-CL conversions (see section 6.6.3.2.1 above). For the combustion factor C_f , the conservative value of 1 is used (Table 2.6 of the 2006 IPCC GL fails to report appropriate values), whereas for the Gef emission factors default values in Table 2.5 of the 2006 IPCC GL are used.

6.7.3 Land converted to Grassland

Figure 6.7.2 reports emissions and removals, whereas Table 6.7.5 reports methodological information for this sub-category.




Figure 6.7.4. Emissions and removals in the L-GL category since 1985. The data under the grey box is shaded out for reasons explained in section 6.1.1.

Table 6.7.5. Methodological summary for L-GL. (CS=country specific; D: default; IE: included elsewhere; AD: activity data; EF: emission/removal factor)

L-GL	Type of information	"FROM" category	BIOMASS	DOM		SOIL		Table (4)III, (4)IV	Table (4)V
				DW	LI	mineral	organic		
	E/R	FL	AD: CS; EF: CS	AD: CS; EF: CS	AD: CS; EF: CS/D	NO	D	Wildfires: IE (GL-GL)	
		CL	AD: CS; EF: D	Tier 1: 0	AD: CS; EF: CS/D	NO			
		WL	NO	NO	NO	NO			
		SE	Tier 1: 0	Tier 1: 0	AD: CS; EF: CS/D	NO			
		OL	NO	NO	NO	NO			
Uncertainty	Tier 2		NE	Tier 2		Tier 2			

6.7.3.1 Forest Land converted to Grassland

For the methodology to estimate carbon stock changes in the biomass, DOM and soil pools, see Sections 6.5.6.1.1 and 6.4.4, 6.5.6.1.2 and 6.5.6.1.3, respectively. The total emissions from these pools estimated using this methodology was split between FL-GL and other conversions by the area of these conversions. The share of FL-GL to all FL-L varies between about 4 and 50% and was 33% in 2020.

6.7.3.2 Cropland converted to Grassland

6.7.3.2.1 Biomass

Carbon stock changes in biomass in this category are the sum of those from converting Cropland with annual crops to Grassland and those from converting Cropland with perennials to Grassland.

For annual crops, the methodology of estimating carbon stock changes is the same as reported in sections 6.4.4 and 6.6.3.2.1, and symbols applied there are used here, too.

For A_{Conv} , data from the annual land-use change matrix was used.

The value of B_{before} for annual croplands, the default 10 t biomass/ha, was taken from text to Table 5.9 of the 2006 IPCC GL, whereas B_{after} was estimated from the proportion of Cropland area of cold dry and warm dry climate types ($P_{CD} = 0.41$, $P_{WD} = 0.59$) and respective specific default Grassland biomass (total above- and below-ground biomass, Table 6.4 of the 2006 IPCC GL: $B_{CD} = 6.5$ t biomass ha^{-1} and $B_{WD} = 6.1$ t biomass ha^{-1} , respectively): $B_{after} = P_{CD} * B_{CD} + P_{WD} * B_{WD}$ (see section 6.7.2.1 for more details). In accordance with the Tier 1 assumption, B_{after} in the equation is 0, and the carbon fraction is the default value of 0.47 tC t biomass $^{-1}$.

For ΔC_G , the same biomass value of 6.26 tC ha^{-1} , together with the default carbon fraction of 0.47 tC t biomass $^{-1}$ was used as for the pre-conversion biomass of B_{before} in the CL-GL category, whereas ΔC_L was assumed to be equal to 0.

When converting perennial Cropland to Grassland, the methodology of estimating carbon stock changes is consistent with the one reported in section 6.6.2.1.1. P_P for Grassland was also estimated from the CORINE land cover change database (see Table 6.7.6 and section 6.2 for details).

Table 6.7.6. *The area of cropland-grassland conversions since 1985.*

Year	Area (ha)			
	Annual cropland converted to grassland	Vineyard converted to Grassland	Orchard converted to Grassland	Total CL-GL
1985	4898	352	88	5338
1986	4551	360	427	5338
1987	0	0	0	0
1988	0	0	0	0
1989	0	0	0	0
1990	0	0	0	0
1991	3814	0	332	4145
1992	3814	0	332	4145
1993	3294	0	851	4145
1994	3294	0	851	4145
1995	3294	0	851	4145
1996	3294	0	851	4145
1997	3294	0	851	4145
1998	3294	0	851	4145
1999	3294	0	851	4145
2000	3516	0	630	4145
2001	1567	0	281	1847
2002	1310	64	171	1544
2003	1310	64	171	1544
2004	1310	64	171	1544
2005	1310	64	171	1544
2006	1310	64	171	1544
2007	1310	64	171	1544
2008	1310	64	171	1544
2009	1310	64	171	1544
2010	1310	64	171	1544
2011	1627	191	101	1919
2012	465	0	24	489
2013	315	16	0	331
2014	800	0	41	841
2015	931	20	29	980
2016	932	0	48	980
2017	932	0	48	980
2018	932	0	48	980
2019	932	0	48	980
2020	304	14	2	320

6.7.3.2.2 Mineral soils

The method and emission factors used are those described in section 6.4.1.

6.7.3.3 Wetlands converted to Grassland

This land-use change is not occurring in Hungary.

6.7.3.4 Settlements converted to Grassland

The land cover change databases indicate rather small areas of Settlement converted to Grassland (ranging between 23 and 191 ha/yr). These areas are predominantly biological re-cultivation of abandoned surface mines. In general, the biological re-cultivation results in an increase in the carbon stocks. For the sake of consistency and completeness, carbon stock changes in mineral soils are estimated using the method and emission factors that are reported in section 6.4.1.

6.7.3.5 Other Land converted to Grassland

This land-use change is not occurring in Hungary.

6.7.4 Uncertainties and time-series consistency

See section 6.11.

6.7.5 Category-specific QA/QC and verification

See section 6.6.5.

6.7.6 Category-specific recalculations

This year, only a few recalculations of small effect were made for the GL category as detailed in Table 6.7.7.

Table 6.7.7. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for (a) the GL-GL and (b) the L-GL category for some recent inventory years. (The key NO --> v means that while NO was reported for 2019, a value was reported in 2020.)

(a)

Sub mission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.C.1 Carbon stock change, Total area	kg ha	0	0	0	0	1117	1113	1111	1116	1121	1125	1127	1132	revision of the LUC matrix
2022		kg ha	0	0	0	0	1117	1114	1113	1117	1123	1128	1129	1134	
		difference, %	%	0,0	0,0	0,0	0,0	0,0	0,1	0,2	0,2	0,2	0,2	0,2	
2021	4.C.1 Carbon stock change, Area of mineral soil	kg ha	0	0	0	0	1117	1113	1111	1116	1121	1125	1127	1132	revision of the LUC matrix
2022		kg ha	0	0	0	0	1117	1114	1113	1117	1123	1128	1129	1134	
		difference, %	%	0,0	0,0	0,0	0,0	0,0	0,1	0,2	0,2	0,2	0,2	0,2	
2021	4.C.1 Carbon stock change, Mineral soils	kg C	0	0	0	0	0	1	1	1	-4	0	0	0	revision of the LUC matrix as well as calculation errors
2022		kg C	0	0	0	0	0	1	1	1	-2	-2	0	-1	
		difference, %	%	0,0	0,0	0,0	0,0	0,0	0,1	0,2	0,2	-41,6	NO --> v	-345,3	

(b)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	L-GI		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.C.2 Carbon stock change, Total area	ktC	0.0	0.0	0.0	0.0	0.0	0.0	89.2	83.9	78.6	73.7	69.8	64.5	revision of the LUC matrix
2022		ktC	0.0	0.0	0.0	0.0	0.0	0.0	89.4	84.1	78.8	73.9	70.0	64.7	
	<i>difference, %</i>	%	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.3	0.3	0.3	
2021	4.C.2 Carbon stock change, Area of mineral soil	ktC	0.0	0.0	0.0	0.0	0.0	0.0	89.2	83.9	78.6	73.7	69.8	64.5	revision of the LUC matrix
2022		ktC	0.0	0.0	0.0	0.0	0.0	0.0	89.4	84.1	78.8	73.9	70.0	64.7	
	<i>difference, %</i>	%	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.3	0.3	0.3	
2021	4.C.2 Carbon stock change, Net change	kt C	0.0	0.0	0.0	0.0	0.0	-3.1	0.0	0.0	0.0	0.0	-47.3	-21.7	revision of the LUC matrix as well as calculation errors
2022		kt C	0.0	0.0	0.0	0.0	0.0	-3.1	0.0	0.0	0.0	0.0	-47.1	-21.9	
	<i>difference, %</i>	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.3	0.5	
2021	4.C.2 Carbon stock change, Net carbon stock change in dead organic matter	kt C	-0.5	-1.5	-2.6	-0.4	-5.7	-1.4	-1.2	-3.8	-4.9	-4.9	-13.7	-6.9	revision of the LUC matrix as well as calculation errors
2022		kt C	-0.5	-1.6	-2.7	-0.4	-6.1	-1.5	-3.8	-4.1	-5.2	-5.2	-14.6	-7.3	
	<i>difference, %</i>	%	5.2	5.4	5.5	5.7	5.8	5.9	214.6	6.1	6.2	6.3	6.2	6.2	
2021	4.C.2 Carbon stock change, Mineral soils	kt C	88.3	91.7	95.7	87.0	78.7	74.5	70.2	66.1	61.7	57.8	54.5	50.4	revision of the LUC matrix as well as calculation errors
2022		kt C	88.3	91.7	95.7	87.0	78.9	74.8	70.8	66.7	62.5	58.6	55.4	51.3	
	<i>difference, %</i>	%	0.0	0.0	0.0	0.0	0.2	0.4	0.8	1.0	1.4	1.4	1.6	1.8	

6.7.7 Category-specific planned improvements

Planned improvements for this category include the implementation of the results of the uncertainty estimation.

6.8 Wetland (CRF sector 4.D)

6.8.1 Description of category

Wetlands account for only about 2.8 per cent of the total area of Hungary (Figure 6.3.1) and include inland marshes, peat bogs, and natural and artificial water courses and water bodies. Wetlands are *ex lege* protected in Hungary that is among the signatories of the Ramsar Convention. The preservation and the sustainable use of Wetlands became standard practice decades ago. In 2017, altogether 29 wetland areas (with a total area of 256,948 ha) were included the Ramsar List of Wetlands of International Importance and managed accordingly (<http://www.ramsar.org/sites/default/files/documents/library/sitelist.pdf>).

The methodology of identifying Wetlands (see section 6.3) does not allow for the separation of managed and unmanaged Wetlands (the latter having a rather small share), but the area of Wetlands could be split into remaining and 'converted to' sub-categories. As wetlands are mainly precipitation dominated, their extent to a certain degree depends on the seasonal and annual variability in precipitation, and due to the nature of land-use and land-use change statistics, this variability could not be entirely reflected in the annual land-use change matrices.

Anthropogenic emissions from wetlands are not significant in Hungary because both the managed and total Wetland area have been small and quite constant for decades, therefore, the Tier 1 method is applied for the estimation of the emissions. To ensure completeness, emissions from both land conversions to water bodies (as conversions to 'flooded land') and peat extraction are reported. The Hungarian Mining Authority (HMA) provides data on the establishment of new peat extraction sites and on the amount of peat extracted annually. The effect of peat bogs conversion to peat extraction sites seems to be insignificant, because peat mining is a very rare activity due to the strict natural protection law. No new extraction sites have been established since 2006.

6.8.2 Wetland remaining Wetland

Table 6.8.1 reports methodological information for this sub-category.

Table 6.8.1. Methodology summary for WL-WL (CS=country specific; D: default; AD: activity data; EF: emission/removal factor; NO: not occurring).

	Type of information	Carbon stock changes		CO2 emissions, on-site	N2O emissions, on-site	CO2 emissions, off-site	Table (4)I, II, III, IV	Table (4)V
		BIOMASS	DOM	SOIL				
WL-WL	E/R	AD: CS; EF: D	Tier 1: 0	AD: CS; EF: D	AD: CS; EF: CS/D	AD: CS; EF: D	Direct N2O emissions from N inputs to managed soils: NO; Emissions from peat extraction: D; Direct and indirect N2O emissions from N mineralization: D	Wildfires: NO Biomass burning: NO
	Uncertainty	NE		NE			NE	

In WL-WL, the main sources of emissions are related to peat extraction. Biomass carbon stock changes unrelated to peat extraction, which are negligible, are not estimated. For the sake of completeness, soil carbon stock changes in “other wetlands” (i.e., wetlands with predominantly grassy vegetation) are estimated using the methodology described in section 6.4.1. Note that this methodology may result in a zero carbon stock change estimate in some inventory years. To demonstrate that it is the result of a calculation, it is reported as zero and not as “NO”.

6.8.2.1 Carbon stock changes as well as CO₂ emissions (on-site and off-site)

According to Equations 7.3. and 7.4 of the 2006 IPCC GL, one source of CO₂ emissions (for all production phases) from peatlands is emissions from peatland extraction (both on-site and off-site) and from biomass clearing:

$$\text{CO}_2\text{-C}_{\text{WW peat on-site}} = [(A_{\text{peatRich}} * \text{EF}_{\text{CO}_2 \text{ peatRich}}) + (A_{\text{peatPoor}} * \text{EF}_{\text{CO}_2 \text{ peatPoor}})] / 1000 + \Delta\text{C}_{\text{WW peat B}}$$

where

$\text{CO}_2\text{-C}_{\text{WW peat on-site}}$ = on-site CO₂-C emissions from peat deposits, kt C yr⁻¹

A_{peatRich} = area of nutrient-rich peat soils managed for peat extraction, ha

A_{peatPoor} = area of nutrient-poor peat soils managed for peat extraction, ha

$\text{EF}_{\text{CO}_2 \text{ peatRich}}$ = CO₂ emission factors for nutrient-rich peat soils managed for peat extraction or abandoned after peat extraction, tonnes C ha⁻¹ yr⁻¹

$\text{EF}_{\text{CO}_2 \text{ peatPoor}}$ = CO₂ emission factors for nutrient-poor peat soils managed for peat extraction or abandoned after peat extraction, tonnes C ha⁻¹ yr⁻¹

$\Delta\text{C}_{\text{WW peat B}}$ = CO₂-C emissions from change in carbon stocks in biomass due to vegetation clearing, kt C yr⁻¹.

Data for A_{peatRich} and A_{peatPoor} was obtained from the HMA for the period 1995-2020 (Table 6.8.2). For the years 1985 to 1994 data on area conversions are not available, therefore, proxy data, i.e., data of 1995 was used for the estimation. For the emission factors, IPCC default values, i.e., 1.1 and 0.2 tonnes C ha⁻¹ yr⁻¹, from Table 7.4 of Volume 4 of the 2006 IPCC GL, were used for nutrient-rich and nutrient-poor peats, respectively, irrespective of the current status (in operation vs abandonment) of the various peat extraction sites.

Table 6.8.2. Area of land converted annually to peat extraction (ha)

Year	Area of land converted annually to peat extraction (ha)	
	Mire	Peat
1995	NO	169.32
1996	NO	68.37
1997	12.16	73.87
1998	NO	802.15
1999	205.15	211.97
2000	88.67	28.13
2001	NO	NO
2002	NO	105.45

2003	NO	NO
2004	4.12	NO
2005	NO	34.45
2006	NO	18.53
2007	NO	NO
2008	NO	NO
2009	NO	NO
2010	NO	NO
2011	NO	NO
2012	NO	NO
2013	NO	NO
2014	NO	NO
2015	NO	NO
2016	NO	NO
2017	NO	NO
2018	NO	NO
2019	17.9	NO
2020	11.5	NO

We note here that the expert review team (ERT) noted during the review in 2017 that the various situations that occur after extraction has ceased, such as abandonment, restoration or land conversion, result in different levels of emissions according to the 2006 IPCC Guidelines and the Wetlands Supplement. The ERT also noted the small impact of these lands on the national level of emissions but encouraged Hungary to undertake further research on site-specific information on the practices taking place at peat extraction sites after extraction ceases, if resource allocation allows the country to do so. Indeed, while this may be a task for further years, we currently have neither appropriate data nor data collection capacity to improve our estimates, and consider the impact mentioned above highly insignificant.

Concerning the methodology to differentiate between nutrient rich and nutrient poor sub-categories, the basis for the classification is the type of organic material in the soil. The database we have allows us to apply two classes here: “mire” that we regard as being poor in nutrients, and “peat” that we regard as being rich in nutrients.

Carbon stock change in biomass due to vegetation clearing was estimated using Equation 2.16 (see Section 6.4.4). In Hungary, the typical biomass of peat bogs, i.e., B_{before} in this equation, is grass as demonstrated in different studies (e.g., Hubayné, 2005 and Dömsödy, 2006). Therefore, B_{before} was estimated from the proportion and average specific biomass for cold dry and warm dry climate types: $B_{\text{before}} = P_{\text{CD}} * B_{\text{CD}} + P_{\text{WD}} * B_{\text{WD}}$ ($P_{\text{CD}} = 0.41$, $P_{\text{WD}} = 0.59$, $B_{\text{CD}} = 6.5 \text{ t biomass ha}^{-1}$ and $B_{\text{WD}} = 6.1 \text{ t biomass ha}^{-1}$, respectively; see more details in Section 6.7.3.3). B_{after} in the equation is 0, and the carbon fraction is the IPCC default value of $0.47 \text{ tC t biomass}^{-1}$.

Off-site emissions from managed peatlands were estimated using Equation 7.5 (by modifying it, i.e., deleting the division by 1000, to correct for appropriate dimensions):

$$\text{CO}_2\text{-C}_{\text{WW peat off-site}} = W_{\text{tdry_peat}} * C_{\text{fractionwt_peat}}$$

where

$\text{CO}_2\text{-C}_{\text{WW peat off-site}}$ = off-site $\text{CO}_2\text{-C}$ emissions from peat removed for horticultural use, tC yr^{-1}

$W_{\text{tdry_peat}}$ = air-dry weight of extracted peat, tonnes yr^{-1}

$C_{\text{fractionwt_peat}}$ = carbon fraction of air-dry peat by weight, tonnes C (tonnes of air-dry peat) $^{-1}$.

$W_{t_{dry_peat}}$ was estimated from the annual statistics (based on measured values) of the amount of peat extracted (provided by the HMA, Table 6.8.3) and the density of the peat, also provided by the HMA by extraction site. The density value (i.e., 0.2 tonnes biomass m^{-3}) is used to convert volume extracted to biomass extracted, and is taken from Hahn (1984). For $C_{fraction_{wt_peat}}$, the area-weighted value of 0.42 tonnes C (tonnes of air-dry peat) $^{-1}$ is used, which was calculated based on the area of nutrient rich and nutrient poor sites and respective data from Table 7.5 of the 2006 IPCC GL. Note that while the conversion factors are constant, the amount of peat extracted varies from year to year due to the high variation of demand for peat.

Table 6.8.3. Annual amount of peat extracted.

Year	Amount of peat extracted	
	Estimated by HMA, 1000 m^3	Converted to mass, tonnes
1985	464	92 800
1986	797	159 400
1987	860	172 000
1988	795	159 000
1989	704	140 800
1990	637	127 400
1991	395	79 000
1992	263	52 600
1993	464	92 800
1994	275	55 000
1995	321	64 200
1996	202	40 400
1997	346	69 200
1998	240	48 000
1999	313	62 600
2000	330	66 000
2001	355	71 000
2002	341	68 200
2003	247	49 400
2004	273	54 600
2005	294	58 800
2006	297	59 400
2007	226	45 200
2008	188	37 600
2009	323	64 600
2010	169,5	33 900
2011	268,7	53 740
2012	185,7	37 140
2013	285,2	57 040
2014	166	33 200
2015	286	57 200
2016	217	43 400
2017	273,6	54 720
2018	137,6	27 520
2019	130,6	26 120
2020	212,2	42 440

6.8.2.2 N₂O emissions

The 2006 IPCC GL provides a Tier 1 methodology to estimate N₂O emissions due to peat extraction. These emissions were only estimated for nutrient rich sites using Equation 7.7:

$$N_2O_{WW \text{ peatExtraction}} = (A_{\text{peatRich}} * EF_{N_2O \text{ peatRich}}) * 44/28 * 10^{-6}$$

where

$N_2O_{WW \text{ peatExtraction}}$ = N₂O emissions due to peat extraction, Gg N₂O yr⁻¹

A_{peatRich} = area of nutrient rich peat extraction sites, ha (see above)

$EF_{N_2O-N \text{ peatRich}}$ = emission factor for drained nutrient-rich wetlands, kg N₂O–N ha⁻¹yr⁻¹ for which the IPCC default value of 1.8 kg N₂O–N ha⁻¹yr⁻¹ from Table 7.6 was used (the multiplier 10⁻⁶ is necessary in the equation to obtain the result in units of Gg N₂O yr⁻¹).

6.8.3 Land converted to Wetland

Table 6.8.4 reports methodological information for this sub-category.

Table 6.8.4. Methodology summary for Land converted to Wetland (CS=country specific; D: default; AD: activity data; EF: emission/removal factor; NO: not occurring).

L-WL	Subcategory	"FROM" category	BIOMASS	DOM		SOIL		Table (4)I, III, IV	Table (4)V
				DW	LI	mineral	organic		
	E/R, land converted to peatland	not estimated (insignificant amounts; if any, included in E/R in land converted to flooded land)							
	E/R, land converted to flooded land	FL	NO	NO	NO	NO	NO	Direct and indirect N2O emissions from N inputs to managed soils: NO; Direct and indirect N2O emissions from N mineralization/ immobilization: NO	Wildfires: NO; Biomass burning NO
		CL	NO	NO	NO	NO	NO		
		GL	AD: CS; EF: D	Tier 1: 0		AD: CS; EF: CS/D	NO		
		SE	Tier 1: 0	Tier 1: 0		AD: CS; EF: CS/D	NO		
		OL	NO	NO		NO	NO		
	Uncertainty	Tier 2		NE		Tier 2		Tier 2	

6.8.3.1 Grassland converted to Wetland

The general methodology of identifying the area of Land converted to Wetland is described in section 6.3.1.

6.8.3.1.1 Biomass

Equation 7.10 of the 2006 IPCC GL was applied as follows:

$$\Delta C_{LW \text{ floodLB}} = A_{\text{Conversion}} * (B_{\text{after}} - B_{\text{before}}) * CF$$

where:

$\Delta C_{LW flood LB}$ = biomass carbon stock change due to land-use conversion to Wetland, tC year⁻¹

$A_{conversion}$ = annual area of land converted to Wetland, ha year⁻¹

B_{after} = carbon stocks of biomass after the conversion to Wetland, tonnes C ha⁻¹

B_{before} = carbon stocks in biomass before the conversion to Wetland, tonnes C ha⁻¹

CF = carbon fraction, tC (t biomass)⁻¹.

To estimate the amount of biomass cleared in the year of conversion, the annual areas of Cropland or Grassland converted to Wetland need to be used because these are the types of conversions that have so far occurred in the time series since 1985.

B_{after} is zero, and B_{before} was estimated the same way as described for wetland prepared for peat extraction above. For more details see "Grassland converted to Cropland" in Chapter 6.6.3.2.

6.8.3.2 Settlements converted to Wetland

There area of Settlements converted to Wetland is very small. The CLC codes which were classified into this category are reported in Table 6.3.2. This land-use change category mainly contains the area of sandpits, gravel pits, and extraction and construction area which are not covered by soil and do not contain biomass. Therefore, emissions from these land-use change conversions are most probably, and taken to be, zero.

6.8.4 Uncertainties and time-series consistency

See section 6.11.

6.8.5 Category-specific QA/QC and verification

This year, with the exception of emissions from peat extraction, only a few recalculations of small effect were made for the WL category as detailed in Table 6.8.5.

Table 6.8.5. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for (a) WL-WL, (b) peat extraction, separately, and (c) L-WL for some recent inventory years. (The key NO --> v means that while NO was reported for 2019, a value was reported in 2020.)

(a)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	WL-WL		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.D.1 Carbon stock change, Total area	ktC	0	0	0	0	0	0	255	255	256	257	257	258	revision of the LUC matrix
2022		ktC	0	0	0	0	0	0	255	255	256	256	257	257	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.3	
2021	4.D.1 Carbon stock change, Net change	kt C	0	0	0	0	0	0	0	0	0	0	0	0	revision of the LUC matrix
2022		kt C	0	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	0	
	difference, %	%	NO --> v	NUM	NUM	NUM	NUM	NUM	NUM	NUM	NUM	NUM	NUM	0.0	
2021	4.D.1 Carbon stock change, Mineral soils	kt C	0	0	0	0	0	0	0	0	0	-1	0	0	revision of the LUC matrix as well as calculation errors
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	0	
	difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-41.7	NO --> v	-344.8	NO --> v	

(b)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	Pest extraction		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.D.1 Pest Extraction, Area	km ²	0	0	0	0	0	0	0	0	0	0	0	0	NA
2022		km ²	0	0	0	0	0	0	0	0	0	0	0	0	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	
2021	4.D.1 Pest Extraction, CO ₂	kt	238	405	215	338	235	358	211	223	270	340	177	182	revision of emission factor
2022		kt	64	106	59	89	64	94	57	94	69	87	45	42	
	difference, %	%	-73,1	-73,8	-72,8	-73,6	-73,0	-73,7	-72,8	-57,7	-74,3	-74,5	-74,9	-76,9	

(c)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	L-WL		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.D.2 Carbon stock change, Total area	km ²	0	0	0	0	0	0	0	0	0	0	0	6	revision of the LUC matrix
2022		km ²	0	0	0	0	0	0	0	0	0	0	0	6	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,4	
2021	4.D.2 Carbon stock change, Losses	kt C	0	0	0	0	0	0	0	0	0	0	0	0	revision of the LUC matrix
2022		kt C	0	0	0	0	0	NO	NO	NO	NO	NO	0	NO	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	NUM	NUM	NUM	NUM	NUM	0,0	NUM	
2021	4.D.2 Carbon stock change, Mineral soils	kt C	0	0	0	0	0	0	0	0	1	1	1	1	revision of the LUC matrix as well as calculation errors
2022		kt C	0	0	0	0	0	0	0	0	1	1	1	1	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,3	-2,0	0,3	-0,2	

6.8.6 Category-specific recalculations

This year, no recalculations were made for the WL category.

6.8.7 Category-specific planned improvements

Currently, there are no planned improvements for this sector.

6.9 Settlements (CRF sector 4.E)

6.9.1 Description of category

Settlements account for 6.3 per cent of the area of Hungary, and they are the sources of only a tiny fraction of all domestic emissions.

6.9.2 Settlements remaining Settlements

As this category is not a key category, the Tier 1 assumption of no change in carbon stocks in all pools is applied.

6.9.3 Land converted to Settlements

Table 6.9.1 reports methodological information for this sub-category.

Table 6.9.1. Methodology summary for Land converted to Settlements (CS=country specific; D: default; AD: activity data; EF: emission/removal factor; NO: not occurring).

L-SE	Type of information	"FROM" category	BIOMASS	DOM		SOIL		Table (4)I, III, IV	Table (4)V		
	E/R	FL	AD: CS; EF: CS	AD: CS; EF: CS		AD: CS; EF: CS/D				Direct and indirect N2O emissions from N inputs to managed soils: NO; Direct and indirect N2O emissions from N mineralization: D	Wildfires: NO; Biomass burning NO
				DW	LI	mineral	organic				
CL				AD: CS; EF: D	Tier 1: 0		AD: CS; EF: CS/D				
GL				AD: CS; EF: D	Tier 1: 0		AD: CS; EF: CS/D				
WL				Tier 1: 0	Tier 1: 0		AD: CS; EF: CS/D				
OL	NO	NO		NO		NO					
Uncertainty	Tier 2			NE		Tier 2		Tier 2			

6.9.3.1 Forest land converted to Settlements

The share of emissions from FL-SE to all FL-L varies between about 29 and 91%, and it was 55% in 2020. For the methodology to estimate carbon stock changes in the biomass, DOM and soil pools, see Sections 6.5.6.1.1, 6.5.6.1.2 and 6.5.6.1.3, respectively.

6.9.3.2 Cropland converted to Settlements

6.9.3.2.1 Biomass

Carbon stock changes in biomass in this category are the sum of those from converting Cropland with annual crops to Settlement and those from converting Cropland with perennials to Settlement.

For annual crops, the methodology of estimating carbon stock changes is the same as reported in Sections 6.4.4 and 6.6.3.2.1, and symbols applied there are used here, too.

For A_{Conv} , data from the annual land-use change matrix was used.

For B_{before} for annual croplands, the default 10 t biomass/ha, was taken from text to Table 5.9 of the 2006 IPCC GL. In accordance with the Tier 1 assumption, B_{after} in the equation is 0, and the carbon fraction is the default value of 0.47 tC t biomass⁻¹. ΔC_L was assumed to be equal to 0.

When converting Cropland with perennials to Settlements, the methodology of estimating carbon stock changes is the same as reported in section 6.6.2.1.1. P_P for Settlements was also estimated from the CORINE land cover change database (see Table 6.9.2 and section 6.2 for details).

Table 6.9.2. *The distribution of the area of cropland annually converted to Settlements.*

Year	Annual area of conversions (ha)			
	Cropland converted to Settlements	Vineyard converted to Settlements	Orchard converted to Settlements	Total CL-SE
1985	755	1	5	761
1986	725	5	3	733
1987	944	5	5	953
1988	764	5	3	772
1989	926	6	2	934
1990	1000	8	0	1009
1991	985	10	0	995
1992	987	10	0	997
1993	871	17	0	888
1994	826	16	0	843
1995	937	18	0	955
1996	1069	21	0	1090
1997	1068	21	0	1089
1998	771	15	0	786
1999	827	16	0	843
2000	824	43	0	867
2001	1859	101	0	1959
2002	2133	74	28	2234
2003	2375	87	33	2495
2004	2157	78	29	2264
2005	1915	77	29	2021
2006	1590	57	21	1668
2007	1157	39	15	1210
2008	1395	52	20	1467
2009	1202	44	17	1263
2010	1236	48	18	1302
2011	1046	18	34	1098
2012	1358	41	0	1398
2013	1040	0	0	1040
2014	992	37	0	1030
2015	854	20	14	889
2016	651	26	0	678
2017	867	35	0	902
2018	902	0	0	902
2019	751	30	0	781
2020	947	2	13	962

6.9.3.2.2 Mineral soils

The method and emission factors used are those described in section 6.4.1.

6.9.3.3 Grassland converted to Settlements

6.9.3.3.1 Biomass

The methodology of estimating carbon stock changes applied for Grassland converted to Settlement is the same as reported in section 6.6.3.2.1. B_{before} was estimated from the proportion of Grassland area of cold dry and warm dry climate types ($P_{\text{CD}} = 0.41$, $P_{\text{wD}} = 0.59$) and respective specific default Grassland biomass (total above- and below-ground biomass, Table 6.4 of the 2006 IPCC GL: $B_{\text{CD}} = 6.5 \text{ t biomass ha}^{-1}$ and $B_{\text{wD}} = 6.1 \text{ t biomass ha}^{-1}$, respectively). The weighted value of $B_{\text{before}} = P_{\text{CD}} * B_{\text{CD}} + P_{\text{wD}} * B_{\text{wD}}$, whereas, in accordance with the Tier 1 assumption, B_{after} is 0.

6.9.3.3.2 Mineral soils

The method and emission factors used are those described in section 6.4.1.

6.9.3.4 Wetland converted to Settlements

6.9.3.4.1 Biomass

Wetland converted to Settlements typically includes inland marshes (rarely peat bogs) the biomass of which is typically grass (Dömsödi, 2006) and water bodies with no biomass. Therefore, the emissions from biomass were estimated using the methodology that is applied for the Grassland converted to Settlement, see Section 6.9.3.3.1 which, for water bodies, may somewhat overestimate the amount of biomass lost and is thus conservative.

6.9.3.4.2 Organic soils

For these conversions, Equation 2.26 was used to estimate the annual carbon loss:

$$L_{\text{organic from water-bodies}} = A_{\text{entire category}} * P_{\text{water-bodies}} * EF_{\text{water-bodies}}$$

and

$$L_{\text{organic from marshes-bogs}} = A_{\text{entire category}} * P_{\text{marshes-bogs}} * EF_{\text{marshes-bogs}}$$

where

L_{organic} = annual carbon loss from organic soils of water bodies and marshes-bogs, respectively, from converting Wetland to Settlements, tCyr^{-1}

$A_{\text{entire category}}$ = area of the entire category of Wetland converted to Settlements, ha

$P_{\text{water-bodies}}$ = proportion of the area of water bodies relative to $A_{\text{entire category}}$, %

$P_{\text{marshes-bogs}}$ = proportion of the area of marshes-bogs, relative to $A_{\text{entire category}}$, %

$EF_{\text{water-bodies}}$ = emission factor for water bodies, $\text{tCha}^{-1}\text{yr}^{-1}$

EF_{water-bodies} = emission factor for marshes and bogs, tCha⁻¹yr⁻¹.

The P values were identified according to section 6.3.1, whereas for the emission factors the default IPCC (2006) values of 0.25 tCha⁻¹yr⁻¹ (cold temperate), 2.5 tCha⁻¹yr⁻¹ (warm temperate, Table 6.3) and 10.0 tCha⁻¹yr⁻¹ (Table 5.6) were used, respectively.

6.9.4 Uncertainties and time-series consistency

See section 6.11.

6.9.5 Category-specific QA/QC and verification

See section 6.6.5.

6.9.6 Category-specific recalculations

This year, only a few recalculations of small effect were made in the SE category as detailed in Table 6.9.3.

Table 6.9.3. The reason and scale of the difference between the value of the estimates in this submission and the previous submission for (a) the SE-SE and (b) the L-SE category for some recent inventory years.

(a)

Sub mission year	Category, quantity	Unit	Inventory year												Reason for recalculation
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	4.E.1 Carbon stock change, Total area	kha	0	0	0	0	528	528	529	530	531	532	534	534	revision of the LUC matrix
2022		kha	0	0	0	0	528	528	529	530	531	532	533	533	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	-0.1	-0.1	-0.1	-0.1	-0.1	-0.1	
2021	4.E.1 Carbon stock change, Area of mineral soil	kha	0	0	0	0	528	528	529	530	531	532	534	534	revision of the LUC matrix
2022		kha	0	0	0	0	528	528	529	530	531	532	533	533	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	-0.1	-0.1	-0.1	-0.1	-0.1	-0.1	

(b)

Sub mission year	Category, quantity	Unit	Inventory year											Reason for recalculation	
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018		2019
2021	4.E.2 Carbon stock change, Total area	kha	0.0	0.0	0.0	0.0	0.0	0.0	50.8	51.1	51.3	51.5	50.8	50.8	revision of the LUC matrix
2022		kha	0.0	0.0	0.0	0.0	0.0	0.0	51.5	51.9	52.0	52.3	52.5	52.5	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	1.5	1.5	1.5	1.5	3.4	
2021	4.E.2 Carbon stock change, Area of mineral soil	kha	0.0	0.0	0.0	0.0	0.0	0.0	50.8	51.1	51.3	51.5	50.8	50.8	revision of the LUC matrix
2022		kha	0.0	0.0	0.0	0.0	0.0	0.0	51.5	51.9	52.0	52.3	52.5	52.5	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	1.5	1.5	1.5	1.5	3.4	
2021	4.E.2 Carbon stock change, Losses	kt C	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-8.9	0.0	revision of the LUC matrix as well as calculation errors
2022		kt C	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-13.4	0.0	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	51.0	
2021	4.E.2 Carbon stock change, Net change	kt C	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-8.9	0.0	revision of the LUC matrix as well as calculation errors
2022		kt C	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-13.4	0.0	
		difference, %	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	51.0	
2021	4.E.2 Carbon stock change, Mineral soils	kt C	-29.3	-29.8	0.0	-30.1	-30.1	-30.4	-30.7	-30.8	-30.7	-30.7	-30.1	-30.1	revision of the LUC matrix as well as calculation errors
2022		kt C	-29.3	-29.8	0.0	-30.2	-30.1	-30.4	-31.1	-31.2	-31.1	-31.1	-31.0	-31.0	
		difference, %	%	0.0	0.0	0.0	0.1	0.2	0.2	1.3	1.3	1.3	1.4	2.9	

6.9.7 Category-specific planned improvements

No improvements are planned in this category.

6.10 Other Land (CRF sector 4.F)

The Other Land category includes the sparsely vegetated areas, which account for only 0.03 percent of the total area of the country (see Figure 6.3.1). The area in the Other Land category is unmanaged (excludes unmanaged grasslands) with very little conversions from and to Other Land.

6.10.1 Uncertainty and time-series consistency

See section 6.11.

6.10.2 Sector specific QA/QC and verification

See section 6.6.5.

6.10.3 Sector specific recalculation

This year, no recalculations were made in the OL category.

6.10.4 Sector specific planned improvements

One of the main plans is to conduct an uncertainty analysis sometimes in the future.

6.11 Uncertainty analysis of the LULUCF sector. Time series consistency

An uncertainty analysis of the emission and removal estimates for the Cropland category using a Tier 1 approach was reported in the NIR of 2014. This year, we replace that with the first uncertainty analysis so far for the entire non-forestry land-use and land-use change sectors. The analysis was done using a standard Tier 2 Monte Carlo (MC) approach according to Chapter 3 of Volume 1 of the 2006 IPCC GL.

Important to note is that the analysis was done for the inventory years 2005 and 2020. While the selection of the latter is straightforward, that of the former is not, and is due to the fact that, as mentioned above, no AD is available before 1985, therefore, neither the time series is consistent before 2005, nor bias due to artifacts cannot be ruled out. (Time series consistency after 2005 is ensured, see section 6.3.) Because of the years that could thus be selected, the analysis of the trends is based on a rather short period, i.e., 15 years.

Concerning the error distributions of the various EFs (the values of which is always assumed to be constant), some uncertainty information was taken from the 2006 IPCC GL when no respective country-specific data was available. When this uncertainty information was used, which is usually provided by the IPCC GL in the form of plus and minus percent uncertainty (U+% and U-%, respectively), we used triangular probability density functions, or PDFs, with parameters

$$a = EF - EF \cdot U\% / 100$$

and

$$c = EF + EF \cdot U\% / 100$$

for lower and upper limits of the variables, respectively, and the EF value itself for the preferred value. We also applied expert judgment in some cases due to lack of appropriate data. Otherwise, which was the majority of the cases, we collected country-specific data. Concerning carbon stock change in mineral soils, which are the most important sources of emissions and removals, we could develop estimates based on measurements in the TIM database referred to above (see section 6.4.1). These estimates are under publication now. The relevant data of all PDFs is reported in **Table 6.11.1**.

Table 6.11.1. Data related to emission factor uncertainty assumed in the uncertainty estimation.

Variable name	Unit	Mean value	SD	a	c	Source of uncertainty information
SOCref, cold dry, HAC	tC/ha	48		43,2	52,8	exp. judg.
SOCref, warm dry, HAC	tC/ha	58,0		52,2	63,8	exp. judg.
SOCref, cold dry, sandy	tC/ha	15,0		13,5	16,5	exp. judg.
SOCref, warm dry, sandy	tC/ha	21,0		18,9	23,1	exp. judg.
SOCref, cold dry, aquic	tC/ha	116,0		104,4	127,6	exp. judg.
SOCref, warm dry, aquic	tC/ha	132,0		118,8	145,2	exp. judg.
SOC reduction factor (relative to respective categories) for SE	dimensionless	0,80		0,640	0,960	exp. judg.
ΔSOC, FL to non-SA CL	tC/ha	-11,1	0,38			Somogyi, 2021; exp. judg.
ΔSOC, FL to SA CL	tC/ha	-1,5	0,38			Somogyi, 2021; exp. judg.
ΔSOC, FL to non-SA GL	tC/ha	2,6	2,00			Somogyi, 2021; exp. judg.
ΔSOC, FL to SA GL	tC/ha	2,6	2,00			Somogyi, 2021; exp. judg.
ΔSOC, FL to SE	tC/ha	-9,6	0,50			Somogyi, 2021; exp. judg.
ΔSOC, FL to WL	tC/ha	20,3	2,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA CL to SA CL	tC/ha	11,8	0,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA CL to non-SA GL	tC/ha	18,7	0,52			Somogyi, 2021; exp. judg.
ΔSOC, non-SA CL to SA GL	tC/ha	18,7	2,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA CL to SE	tC/ha	-9,9	0,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA CL to WL	tC/ha	18,7	2,50			Somogyi, 2021; exp. judg.
ΔSOC, SA CL to non-SA GL	tC/ha	7,0	2,50			Somogyi, 2021; exp. judg.
ΔSOC, SA CL to SA GL	tC/ha	7,0	2,50			Somogyi, 2021; exp. judg.
ΔSOC, SA CL to SE	tC/ha	-12,3	0,50			Somogyi, 2021; exp. judg.
ΔSOC, SA CL to WL	tC/ha	7,0	2,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA GL to SA GL	tC/ha	0,0	0,50			Somogyi, 2021; exp. judg.
ΔSOC, non-SA GL to SE	tC/ha	-13,7	2,00			exp. judg.
ΔSOC, non-SA GL to WL	tC/ha	0,0	0,52			Somogyi, 2021; exp. judg.
ΔSOC, SA GL to SE	tC/ha	-13,7	2,00			exp. judg.
ΔSOC, SA GL to WL	tC/ha	0,00	0,50			Somogyi, 2021; exp. judg.
ΔSOC, SE to WL	tC/ha	13,67	2,00			Somogyi, 2021; exp. judg.
FLU, non-SA CL, temperate, dry, CL-CL	dimensionless	0,8	0,072			IPCC GL Vol. 4. Ch. 4. Table 5.5
FLU, SA CL, temperate, dry, CL-CL	dimensionless	0,93	0,102			IPCC GL Vol. 4. Ch. 4. Table 5.5
FMG, full till, temperate, dry, CL-CL	dimensionless	1		0,8	1,2	exp. judg.
FMG, reduced till, temperate, dry, CL-CL	dimensionless	1,02	0,061			IPCC GL Vol. 4. Ch. 4. Table 5.5
FMG, no till, temperate, dry, CL-CL	dimensionless	1,1	0,055			IPCC GL Vol. 4. Ch. 4. Table 5.5
FI, low, temperate, dry, CL-CL	dimensionless	0,95	0,124			IPCC GL Vol. 4. Ch. 4. Table 5.5
FI, medium, temperate, dry, CL-CL	dimensionless	1		0,8	1,2	IPCC GL Vol. 4. Ch. 4. Table 5.5
FI, high without manure, temperate, dry, CL-CL	dimensionless	1,04	0,135			IPCC GL Vol. 4. Ch. 4. Table 5.5
FLU, GL-GL	dimensionless	1		0,5	1,5	Table 6.2 of Chapter 6
FMG, nominally managed, GL-GL	dimensionless	1		0,8	1,2	exp. judg.
FMG, improved, GL-GL	dimensionless	1,14		1,0	1,3	Table 6.2 of Chapter 6
FI, improved, GL-GL	dimensionless	1		0,9	1,1	as per Table 6.2 of Chapter 6
R = C/N ratio	tN / tC	15		10,5	19,5	page 11.16 of IPCC 2006 GL
R = C/N ratio, for CL-CL only	tN / tC	10		7,0	13,0	CL-CL only, p. 11.16 of 2006IPCCGL
EF1 of Equation 11.1	tN2O-N / tN	0,01		0,0	0,0	Table 11.1 of IPCC 2006 GL
FracLEACH(-H)	t N / t N additions	0,3		0,1	0,8	Table 11.3 of IPCC 2006 GL
FracLEACH(-H), for Warm & Dry	t N / t N additions	0,3		0,1	0,8	Table 11.3 of IPCC 2006 GL
EF5 of Equation 11.10	tN2O-N / tN leaching/runoff	0,0075		0,0	0,0	Table 11.1 of IPCC 2006 GL

Table 6.11.1. Data related to emission factor uncertainty assumed in the uncertainty estimation (ctd.).

Variable name	Unit	Mean value	SD	a	c	Source of uncertainty information
length of rotation period, orchards	years	30		21,0	39,0	exp. judg.
length of rotation period, vineyards	years	31,8		22,3	41,3	exp. judg.
carbon fraction, orchards	tC/t biomass	0,5		0,5	0,5	as per Table 4.3 of the GL
carbon fraction, vineyards	tC/t biomass	0,5		0,5	0,5	as per Table 4.3 of the GL
average biomass carbon (measured, i.e., average carbon before conversion, AB + BB) and the age for which it refers to, orchards	tC/ha	2,35		1,6	3,1	exp. judg.
average biomass carbon (measured, i.e., average carbon before conversion, AB + BB) and the age for which it refers to, vineyards	tC/ha	4,43		3,1	5,8	exp. judg.
the age for which the average biomass carbon refers to, orchards	years	15		10,5	19,5	exp. judg.
the age for which the average biomass carbon refers to, vineyards	years	15		10,5	19,5	exp. judg.
combustion factor	dimensionless	0,8		0,0	1,6	exp. judg.
emission factor, CH ₄	g/kg dm burnt	2,7		1,8	3,6	exp. judg. based on Table 2.5
emission factor, N ₂ O	g/kg dm burnt	0,07		0,0	0,1	exp. judg. based on Table 2.5
GWP, CH ₄	tCO ₂ eq / t CH ₄	25	4,2	16,8	33,3	exp. judg. based on several literature sources
GWP, N ₂ O	tCO ₂ eq / t N ₂ O	298	50,2	199,7	396,3	exp. judg. based on several literature sources
Biomass removed when preparing sites, cold dry, L-CL	t/ha	6,5	2,4			Table 6.4 (total AG+BG: BG also due to tillage)
Biomass removed when preparing sites, warm dry, L-CL	t/ha	6,1	2,288			Table 6.4 (total AG+BG: BG also due to tillage)
C conversion factor, L-CL	tC/t biomass	0,47		0,5	0,5	as per Table 4.3 of the GL
Cropland (annual): Biomass carbon present after conversion, L-CL	t biomass/ha	10	3,750			as per Table 6.4 of the GL
emission factor, CH ₄	g/kg dm burnt	2,3		1,4	3,2	Table 2.5
emission factor, N ₂ O	g/kg dm burnt	0,21		0,1	0,3	Table 2.5
emission factor, poor sites, WL-WL	tC/ha*yr	0,2		0,0	0,4	exp. judg. based on Table 7.4
emission factor, rich sites, WL-WL	tC/ha*yr	1,1		0,0	2,2	exp. judg. based on Table 7.4
carbon fraction, poor sites, WL-WL	tC/t biomass	0,45		0,4	0,5	exp. judgment based on other Cf U'ss
carbon fraction, rich sites, WL-WL	tC/t biomass	0,4		0,4	0,4	exp. judgment based on other Cf U'ss
emission factor, rich sites, WL-WL	kgN ₂ O-Nha-1yr-1	1,8		0,6	3,0	exp. judg. based on Table 7.6
emission factor from water bodies, cold dry, WL-WL	tC/ha*yr	0,25		0,0	0,5	Table 6.3
emission factor from water bodies, warm dry, WL-WL	tC/ha*yr	2,5		0,3	4,8	Table 6.3
Emission factor: from inland marshes and peat bogs, WL-WL	tC/ha*yr	10		1,0	19,0	Table 5.6
basic density of peat, WL-WL	t dry peat/m ³	0,2	0,100			exp. judg.
Proportions of FL, CL and GL by climate and soil types	dimensionless	0,536	0,055	0,4	0,6	exp. judg.
Proportions of FL, CL and GL by climate and soil types	dimensionless	0,357	0,036	0,3	0,4	exp. judg.
Proportions of FL, CL and GL by climate and soil types	dimensionless	0,098	0,010	0,1	0,1	exp. judg.
Proportions of FL, CL and GL by climate and soil types	dimensionless	0,009	0,001	0,0	0,0	exp. judg.

Concerning AD (i.e., area data), expert judgment had to be used because the current complicated land identification system that produces them, especially the land-use change data, does not really allow one to mathematically or numerically derive uncertainty-related information. The basis for the expert judgment was area data for the forest-related conversions and their uncertainties that were estimated in a dedicated project as reported in section 11.3.1.5. This area data is collected in the forestry surveys, and since it is deemed rather accurate, an uncertainty of 5% is used in this analysis. In contrast, the accuracy of the non-forest related conversions is assumed to be lower primarily due to two main factors. One is the uncertainty of identifying land-use changes by the CLC system, assumed to be 20%, and the uncertainty due to the fact that this system provides land-use change information for six-year periods (section 6.3), and the annual data during such periods is assumed to have an *uncertainty* of plus and minus 30 percent for all non-forest related LUC categories.

Note that we assumed different percentage *errors* for both the different land-use change classes and each year. However, for each year, we calculated the area of the land remaining land categories from the area in the previous year and the area of all land-use changes affecting these categories in that year, the sum of which thus always adds up to the total area of the country. We also note that the land-use change matrix of the totals of the land-use change sub-categories is also recalculated for each inventory year using the perturbed annual areas of the land-use change categories for the default transition periods of 20 years. (Since some annual land-use change areas in a sub-category can be larger than the median value, and some can be smaller than that, the area of the 20-year land-use change sub-categories changes much less over time in relative terms than that of the annual changes.)

Finally, the uncertainty of some other AD as well as some key emission sources from the forestry sector were also used in the uncertainty analysis as reported in **Table 6.11.2**. The data for the forestry sector (with uncertainties reduced) was used to analyze and demonstrate the relative importance of the non-forestry sectors within the entire LULUCF sector when drawing conclusions.

Table 6.11.2. Uncertainty information for some AD and emission estimates from some key sources from the forestry sector.

Variable name	Unit	Mean value in 2020	U%	a	c	Source of uncertainty information
Share of area by tillage (in proportion to TOTAL CL-CL area): full till	dimensionless	0,902		0,895	0,905	exp. judg.
reduced till	dimensionless	0,089		0,085	0,092	exp. judg.
area of pCL-FL orchards	ha/yr	143	10	129	157	exp. judg.
area of pCL-FL vineyards	ha/yr	11	10	10	13	exp. judg.
FL-FL Biomass net CO2 emissions	tCO2/yr	-5 331 902	30	-3 732 331	-6 931 473	exp. judg.
L-FL tree biomass CO2 sink	tCO2/yr	-910 683	15	-774 080	-1 047 285	exp. judg.
ΔCFL-CL - BIOMASS	tC/yr	-6 566	20	-5 253	-7 879	exp. judg.
ΔCFL-CL - DEADWOOD	tC/yr	-1 267	51	-621	-1 913	exp. judg.
ΔCFL-CL - LITTER	tC/yr	-3 242	51	-1 589	-4 896	exp. judg.
ΔCFL-GL - BIOMASS	tC/yr	-13 547	20	-10 838	-16 257	exp. judg.
ΔCFL-GL - DEADWOOD	tC/yr	-1 340	50	-670	-2 010	exp. judg.
ΔCFL-GL - LITTER	tC/yr	-3 429	50	-1 714	-5 143	exp. judg.
ΔCFL-SE - BIOMASS	tC/yr	-16 430	20	-13 144	-19 717	exp. judg.
ΔCFL-SE - DEADWOOD	tC/yr	-1 488	20	-1 191	-1 786	exp. judg.
ΔCFL-SE - LITTER	tC/yr	-3 808	20	-3 047	-4 570	exp. judg.
L-FL DEADWOOD CO2	tCO2/yr	-28 729	50	-14 364	-43 093	exp. judg.
L-FL LITTER CO2	tCO2/yr	-178 353	50	-89 177	-267 530	exp. judg.

Based on all measured and assumed data, and assuming normal error distributions for variables for which this PDF type is known and triangular distributions for variables with uncertainty information (including the land-use change areas) and other variables, we run the MC analysis using the same software in which the inventory calculations are done, thus ensuring that all calculations in the MC analysis are consistent with the inventory calculations. The MC simulations were repeated 10,000 times. For the uncertainty estimates, the 2.5 and 97.5 percentiles of the simulated sub-category or category-level distributions were used as the lower and upper limits of the confidence intervals, respectively. The contribution of the subcategory and pool/gas levels to the total variance and the uncertainty of trends were also calculated, again according to the 2006 IPCC GL.

The results of the analysis are reported in **Table 6.11.3**.

Table 6.11.3. Summary results of the LULUCF-level uncertainty analysis. (N/A means that the emissions in the given sub-category were not occurring. Important sub-categories are highlighted. For Forest land, only the biomass pool is considered.

Category	Sub-category	Pool / gas	sub-category	2005				2020				Trend			
				E/R	Uncertainty, %		Contribution to variance, %	E/R	Uncertainty, %		Contribution to variance, %	Mean value, %	Uncertainty, %		Contribution to variance, %
					(-)	(+)			(-)	(+)			(-)	(+)	
FL-FL	NA	Biomass	NA	-3 510 401	-23	23	9,1	-5 326 892	-23	23	61,0	51	-84	122	1,1
L-FL	NA	Biomass	NA	-1 457 051	-36	42	5	-1 150 200	-35	44	10	-22	-73	90	0
CL-CL	All	Biomass	perennials	30 787	-32	47	<0.1	5 664	-72	70	<0.1	-81	-16	13	<0.1
CL-CL	NA	Soil, CO2	mineral soil	-186 371	-1 236	1 128	73	118 412	-555	508	17	-80	-286	309	16
CL-CL	NA	CH4	wildfires	1 038	-86	168	<0.1	214	-86	170	<0.1	-79	-6	9	<0.1
CL-CL	NA	N2O	wildfires	321	-86	172	<0.1	66	-86	171	<0.1	-79	-6	9	<0.1
CL-CL	NA	soil, non-CO2	NA	154 878	-74	128	0,5	71 800	-74	123	0,3	-54	-10	13	<0.1
L-CL	FL-CL	Biomass	NA	8 841	-15	15	<0.1	24 110	-16	15	<0.1	173	-33	40	1
L-CL	GL-CL	Biomass	NA	-18 879	-230	217	<0.1	-11 994	-222	216	<0.1	-38	-49	67	0,2
L-CL	All	Soil	mineral soil	360 823	-175	187	6,3	215 904	-172	186	6	-40	-66	63	0,2
CL-CL	NA	soil, non-CO2	NA	35 658	-99	350	0,2	21 386	-97	345	0,2	-39	-24	752	23
GL-GL	NA	Soil	mineral	7 242	-140	182	<0.1	121	-140	182	<0.1	-98	0	0	<0.1
GL-GL	NA	CH4	wildfires	5 973	-55	76	<0.1	794	-56	76	<0.1	-87	-4	5	<0.1
GL-GL	NA	N2O	wildfires	6 422	-58	83	<0.1	854	-57	85	<0.1	-87	-4	5	<0.1
GL-GL	NA	soil, non-CO2	NA	796	-93	297	<0.1	13	-93	297	<0.1	-98	0	0	<0.1
L-GL	CL-GL	Biomass	NA	17 452	-148	153	<0.1	6 614	-206	212	<0.1	-62	-191	198	4
L-GL	WL, SE, OL - GL	Biomass	NA	-1 271	-63	53	<0.1	-2 378	-65	53	<0.1	87	-61	90	2
L-GL	All	Soil	mineral	-304 923	-201	188	5	-172 590	-197	183	5	-44	-63	65	0,2
L-GL	NA	Non-CO2 emissions	NA	6 792	-100	732	<0.1	2 320	-100	1 119	<0.1	-42	-46	1 024	50
WL-WL	peatland remaining	biomass	NA	373	-53	55	<0.1	124	-53	55	<0.1	-67	-6	7	<0.1
WL-WL	NA	Soil	NA	6 231	-75	76	<0.1	3 201	-62	61	<0.1	-49	-24	82	0,4
WL-WL	NA	on-site, N2O	N2O from soils	1 252	-52	54	<0.1	484	-52	53	<0.1	-61	-8	9	<0.1
WL-WL	NA	emissions, off-site	NA	89 310	-98	107	0,1	64 761	-98	106	0,2	-28	-63	84	0,1
L-WL	GL-WL, CL-WL	biomass	NA	5 245	-54	65	<0.1	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
L-SE	FL-SE	biomass	NA	74 474	-34	39	<0.1	77 104	-20	22	<0.1	4	-603	928	0,3
L-SE	CL-SE, GL-SE, WL-SE	Biomass	NA	41 320	-61	70	<0.1	20 135	-61	69	<0.1	-51	-23	30	<0.1
L-SE	All	Soil	mineral	105 138	-60	62	<0.1	112 760	-59	62	0,2	8	-163	189	<0.1
L-SE	All	Soil	organic	1 938	-64	66	<0.1	2 009	-65	68	<0.1	3	-286	335	<0.1
L-SE	NA	Soil	N2O from soils	11 076	-78	172	<0.1	11 795	-77	174	<0.1	7	-163	169	<0.1
Total			non-forest only	506 846	-455	423	86	684 852	-96	90	29	-37	-1 836	1 674	99
TOTAL			LULUCF	-4 471 796	-50	50	100	-5 775 371	-24	23	100	29	<-100	>100	100

Concerning the uncertainty ranges in the above table, one often finds that they are large, e.g., >100% in absolute values. This is only partly because of the high uncertainties, however, also (and often mainly) partly because of the nature of the percentage calculations. If a central net emission estimate, C , is close to zero, then the U^- and U^+ absolute uncertainty values, when converted to percentages by dividing them with C and multiplying by 100, become large or very large. This is an inherent difficulty of the uncertainty analysis for the LULUCF sector where net emissions (i.e., the net of emissions and removals, often not insignificant in themselves) and certain emissions can be rather small.

According to the results, and also considering the above, both the ratio of the net emissions and the uncertainty values considerably depend on the inventory year. In both inventory years selected, FL-FL and L-FL are important categories, however, the largest contribution in 2005 comes from the (then large) carbon stock changes from mineral soils from CL-CL, with carbon stock changes from mineral soils from L-CL being the third. In 2020, since the sink in CL-CL dropped considerably, its contribution dropped, too, whereas that of L-CL has remained about the same. In contrast, as the sink from FL increased, its relative importance increased, too.

Considering trends, it is N_2O emissions from soils that seem to play the most important role, however, one has to consider here that such emissions only occur in some years (i.e., these emissions are zero in others), and that the current calculation procedure has difficulties in considering years with zero emissions, thus, most probably, these emissions are much less important than what the raw numbers suggest. When analysing trends, one must also take into consideration the relatively high inter-annual variability. In other words, changing either the start year or the end year of the analysis might have a rather large impact on the result of the analysis.

As a conclusion of the uncertainty analysis, overall, it is emissions and removals from forests that need to be focused on in the development of the LULUCF GHG inventory. Additionally, the estimation of the emissions from cropland (both CL-CL and L-CL) and grasslands (L-GL) deserves due attention, both concerning AD and EF.

7 Waste (CRF sector 5)

Recent key developments:

- In contrast with other sectors, emissions from the waste sector are by 5% higher now than in the base year.
- However, the growth in emissions had stopped in the last decade, and a reduction of 19% could be observed between 2005 and 2020.
- Amount of disposed municipal waste decreased by 50% between 2005 and 2020.
- Emissions from wastewater handling have a pronounced decreasing trend due to a growing number of dwellings connected to the public sewerage network.

Major changes from previous submission:

- Data of biogas produced have been revised. Biogas from manure management has been subtracted from biogas accounted for under 5.B.2.b as it is accounted for under the Agriculture sector.
- Protein consumption data and the amount of wastewater undergoing tertiary treatment has been updated for year 2019.

7.1 Overview of sector

This section discusses the emissions from solid waste disposal (CH_4), biological treatment of solid waste including composting and anaerobic digestion at biogas facilities (CH_4 , N_2O), waste incineration (CO_2 , CH_4 , and N_2O), and domestic and industrial wastewater treatment (CH_4 and N_2O). One peculiarity of the sector is that most part of the carbon-dioxide emissions is generated from biological (biogenic) sources and this CO_2 emissions are either reported as carbon stock change in the LULUCF sector or do not need to be accounted for (e.g. annual crops).

The waste sector with 3,401.6 Gg CO_2 equivalent represented 5% of total national GHG emissions in 2020. In the base year, total GHG emissions from the waste sector amounted to 3,225.20 Gg CO_2 equivalent which accounted for 3% of total national GHG emissions. The largest category was solid waste disposal on land, representing 85% in 2020, followed by wastewater treatment and discharge (9%), biological treatment of solid waste (4%), and incineration of waste without energy recovery (1%). In contrast with other sectors, emissions from the waste sector are slightly higher now than in the base year. However, the growth in emissions stopped in the last decade, and a reduction of 18% could be observed between 2005 and 2020. The degradation process in solid waste disposal sites is quite slow which means that waste that were disposed many years earlier have still an influence on current emission levels. However, the amount of disposed waste had dropped significantly since 2005 (e.g. landfilled municipal waste decreased by 50%) consequently methane emissions started to decrease as well. GHG emissions from wastewater handling have a pronounced decreasing trend due to a growing

number of dwellings connected to the public sewerage network. All these developments are summarized in Figure 7.1.1.




Figure 7.1.1 Trend of emissions of the different categories in waste sector

A major but decreasing part of **municipal solid wastes (MSW)** is treated by managed disposal and a smaller part by reuse, incineration or other means. The average specific municipal household waste generation rate decreased from 1.3 to 1.0 kg/capita/day in the last few years. The total amount of MSW was 3,546 Gg in 2020. Out of this, 1,171 Gg (33%) was recovered by recycling and composting, 601 Gg (17%) was incinerated for energy purposes, and 1,770 Gg (50%) went to landfills and 4 Gg (0.1%) waste was treated in other ways. (In previous years, before 2010, 30-228 Gg waste was treated in other ways which meant mostly mechanical biological treatment (MBT) that produced refuse-derived fuel that could be used in power plants and cement factories.)

Figure 7.1.2 summarizes recent changes in generation and treatment of municipal waste for the period 2004-2020. The following beneficial trends could be observed:

- The increase of waste generation stopped around 2006, and started to decrease quite significantly afterwards (-25% between 2006 and 2020);
- Share of landfilling decreased from 84% to 50% between 2004 and 2020. However, in comparison with the Western-European situation, the share of waste disposal is still relatively high;
- Importance of both recycling (including export) and composting increased significantly; currently they represent 28% and 9%, respectively.

Please note that the above general information relates only to municipal solid waste whereas in the calculations also other waste types (e.g. industrial, construction) are included.




Figure 7.1.2 Main trends in municipal waste handling

7.2 Solid waste disposal in landfills (CRF sector 5A)

Emitted gas: CH₄

Key source category: Level, Trend 1

7.2.1 Source category description

In case of managed disposal, the waste is disposed in landfills where it is compacted and covered. Under these circumstances, anaerobic degradation occurs, during which methane and carbon dioxide is emitted. In advanced disposal sites, the generated methane is recovered by incineration or flaring. Degradation requires several decades and occurs at varying rates. Since waste disposal is continuous, gas generation can also be considered continuous on a country scale.

The CO₂ generated in landfills is of biogenic origin and is thus excluded from the inventory. Under the conditions prevailing in landfills, CO₂ generated from wastes containing carbon of fossil origin is insignificant and direct incineration does not occur in landfills. Illegally disposed wastes are not considered here, partly as they are disposed in batches, in thin layers without compaction, in a fashion well-penetrable for oxygen. Therefore, degradation is aerobic and only carbon dioxide is produced. In accordance with the IPCC Guidelines, no CO₂ emission has to be included in this category.

7.2.2 Methodological issues

Emissions were calculated using the default first order decay methodology. For the calculations, the IPCC Waste Model from the 2006 IPCC Guidelines was used with the “waste by composition” option. The FOD method produces a time-dependent emission profile which reflects the true pattern of the degradation process.

Activity data

The FOD method requires a quite long time series. The default first year in the IPCC Waste Model is 1950. As the eldest data which could be found in statistical publications were for 1975, extrapolation had to be made. For this purpose, a similar pattern as in Figure 7.2.1 had been used. This figure was taken from a university textbook sponsored by the Ministry of Education and Culture.




Figure 7.2.1 The loosening trend of municipal solid waste in Budapest. The solid line denotes the amount of waste while the dotted line shows the decrease of volume-density. Source: (<http://www.hik.hu/tankonyvtar/site/books/b108/>).

Before 2001, the amount of removed solid waste was reported in volume units (m³), therefore these data had to be converted to mass unit using the gravimetric density (t/m³) as an important physical characteristic of the waste. Between 1975 and 2000, the value of this parameter decreased from 0.3 t/m³ to 0.2 t/m³ based on the data of the Statistical Office. Both international and national studies suggested that the mass of municipal solid waste increased hardly while waste volumes increased drastically all over the world, which is reflected in decrease of the gravimetric density. These changes are attributable to the increasing amounts of paper and plastic in the packaging sector. In other words, this is the so-called loosening trend in MSW which can be seen clearly in Fig. 7.2.1. To summarize the above, the following densities can be used for conversion from volume to waste units:

Table 7.2.1 Waste densities suitable for conversion

	1975-1985	From 1990	2000
Density (t/m ³)	0.3	0.22	0.2

(As the statistical office publishes data in mass units from 1990, practically there was no need for the compiler institute to do this conversion for the recent years.)

For the period 1950-1975, the following assumptions were made. The first data found in statistical publication was from 1975, i.e. 6,241 thousand m³. This value was converted using a density of 0.3 t/m³ which resulted in a mass value of 1,872 kilotonnes. The IPCC Guidelines suggest using surrogates, e.g. population in cases where domestic data do not cover the last 50 years. In this submission amount of disposed waste between 1950 and 1975 is assumed to be proportional to urban population. Urban

population increased by more than 50 per cent between 1950 and 1975 based on information from the statistical office. GDP might have been an alternative but then the increase would have been steeper. (GDP grew by 128 per cent, whereas urban population changed by 31 per cent between 1960 and 1975. A little experiment carried out a few years ago showed, however, that the model is not that sensitive for early years. Even when halving the landfilled amount in 1950, the resulting change in emissions was minus 2% to 3% between 2000 and 2011.)

The next published data for landfilled waste we found was 9,952 thousand m³ for 1980. Using the same conversion, this amount is equivalent to 2,986 kilotonnes. For the years between 1975 and 1980, simple interpolation was carried out. The next data was from 1985, i.e. 13,791 thousand m³. Using the same density, it equaled to 4,137 kt from which the incinerated amount (244 kt) was subtracted. Again, an interpolation was made between 1980 and 1985. Then, from 1986, the now yearly published statistical data were converted from volume to mass with a diminishing waste density (from 0.29 t/m³ in 1986 to 0.24 t/m³ in 1989).

From 1990, yearly data in mass units published by the central statistical office was used.

From 2005, data from the Waste Management Information System maintained by the Ministry of Environment and Water were analyzed and used for calculations. This database contains very detailed information on waste management practices in Hungary. The Waste Management Information System (EHIR) can be accessed via internet as well (<http://web.okir.hu/en/ehir>). Data availability has been improved significantly, at least for recent years.

For activity data collection, the main data sources were the following:

- From 1975: Statistical Yearbooks
- 1990-2004: Statistical Yearbooks, Environmental Statistical Yearbooks, Eurostat;
- 2005- Waste Management Information System, Statistical Yearbooks, Eurostat

Beside municipal waste, also industrial waste disposal is taken into account in emission estimations. In the waste information system, disposed waste is categorized by waste types in line with European legislation. The waste types in the database are defined both on the basis of the six-digits European Waste Catalogue (EWC) codes and on the basis of the statistical waste nomenclature European Waste Classification (EWC-Stat), which is a substance-oriented nomenclature and which has been specially created for EU waste statistics (See Eurostat Manual on Waste statistics).

For this submission, we used mainly the latter categorization with only a few exceptions. Especially the following waste categories were included in the calculations:

- Health care and biological wastes (W05)
- Paper and cardboard wastes (W072)
- Wood wastes (W075)
- Textile wastes (W076)
- Animal and vegetal wastes (W091 Animal and mixed food waste +W092 Vegetal wastes +W093 Animal faeces, urine and manure);
- Mixed ordinary wastes (W101 Household and similar wastes + W102 Mixed and undifferentiated materials + W103 Sorting residues)
- Mineral waste from construction & demolition (W121).

As exceptions from the above, EWC codes are applied for the following categories:

- EWC 200307 (bulky waste): it is assumed that half of it is wood waste;

- EWC 200201 (biodegradable waste from garden and park wastes): this made possible to include garden / park waste as a separate category into the IPCC Waste model.

Sludges: Here we take also into account data reported by wastewater treatment plants (and collected by the General Directorate of Water Management). In contrast to the “normal” waste management data, sludges are reported not in fresh amount but in tonnes of dry matter.

The categories above are considered as degradable, all other categories are considered as inert.

Currently, about half of the disposed waste can be considered as degradable (i.e. falls into the above waste categories) within which mixed ordinary (mostly municipal) waste is the dominant category representing 53% of all disposed waste in 2020. The other half was inert waste where combustion wastes dominated with a share of 36% in 2020. The amount of all disposed waste decreased quite significantly after 2005 as demonstrated with the following figure.




Figure 7.2.2 Decreasing trend in waste disposal based on data from the EHIR database (2005-2020)

As reliable information on waste treatment in the EHIR database is only available from 2005, data on disposed waste needed to be extrapolated back to 1950. The following proxy data were used for this extrapolation.

- For paper, wood, textile, animal and vegetal (and other industrial wastes):
 - 1990-2004: primer energy use with the assumption that energy efficiency measures might go hand in hand with improved material usage in industry (i.e., who is wasting energy might be wasting materials too).
 - 1960-1990: volume index of GDP was used as proxy. For the preceding 10 years (1950-1959) the amount was kept constant.
- For clinical waste:
 - 1960-2004: Number of active hospital beds. Similarly, as above, the amount was kept constant for the preceding 10 years.
- For construction and demolition waste:

- 1989-2004: volume indices of value added in the construction sector were used as proxy;
- 1960-1989: similar to industrial waste, the volume indices of GDP were used as proxy. For the preceding 10 years (1950-1959) the amount was kept constant.

Other parameters used in the calculations:

As regards **waste composition** of mixed (municipal) waste, up to the inventory year 2014, statistics only existed for the waste collected in Budapest and in good quality only from 1990. Having no other choice, these yearly data were used for the entire country. Again, as the FOD method requires data starting in 1950, further assumptions had to be made. For 1950, the regional default values representative for Eastern Europe were taken from Table 2.3 of the 2006 IPCC Guidelines (i.e. food 30.1%, paper 21.8%, wood 7.5%, textiles 4.7% etc.), and interpolation was carried out between these and the measured values for 1980.

In the Hungarian statistics, the following waste composition categories have been used for a longer period of time: paper, plastic, textile, glass, metal, degradable organic, hazardous waste, other non-organic. Recently, hygienic waste (e.g. nappies) has been added to the categories. These categories are mainly in line with the requirements of the models. We have added wood to MSW composition categories with a share of 2.5% for the period 1990-2004, and data from the waste statistics were used directly from 2005. Emissions from garden/park waste were calculated separately.

As detailed information on waste composition became available from about 70 waste disposal sites from the 2014 inventory year onwards, it was decided to use the weighted average of the reported waste composition data as country-wide average in this submission. Comparing the estimated DOC values calculated from the different waste composition data, we can see a quite good agreement for 2014. However, in recent years, DOC seems to be higher in the capital than in other parts of the country mainly due to higher share of disposed degradable organic waste (food and green waste) in Budapest. The result of this comparison is shown in the figure below.





Figure 7.2.3 Comparison of DOC estimated from waste composition in Budapest vs country-wide average

In the IPCC waste model, the following waste categories were used:

- Food = degradable organic waste in mixed ordinary waste + separately collected animal and vegetal wastes (i.e., W091, W092, W093);
- Garden/park waste = biodegradable waste from garden and park wastes (EWC 200201);
- Paper = paper, cardboard and composite part of mixed ordinary waste + separately collected paper and cardboard wastes (W072);
- Wood = wood in mixed ordinary waste + separately collected wood wastes (W075) + half of bulky waste (EWC 200307);
- Textiles = textiles in mixed ordinary waste + separately collected textile wastes (W076);
- Nappies = share of hygienic waste in mixed ordinary waste + separately collected health care and biological wastes (W05);
- Sludge = mainly sludges from domestic wastewater plants.
- Construction and demolition waste = mineral waste from construction & demolition (W121).

Default parameters of the IPCC waste model typical of **dry temperate climate** were used. In accordance with Table 3.4 in the 2006 IPCC Guideline, a climate zone can be defined as dry when the mean annual precipitation (MAP) is lower than the potential evapotranspiration (PET). Now, the climatological average of MAP/PET is between 0.48 and 1.11 with a mean of 0.6. Values higher than 1 occur only on 0.04% of our territory (see the map below with greenish and blue colors representing $P/PET < 1$ and $P/PET > 1$, respectively.)



The methane generation rate constants (k) were between 0.04 and 0.06 depending on waste type with an average value of 0.05. The default 50% fraction of methane in developed gas was kept, and so was the 6 month of delay time. Basically, the default values given in the IPCC 2006 Guidelines were chosen for DOC, too. (In previous submissions we used one category (“degradable organic waste”) for food and non-food (e.g. garden waste) fraction of the municipal solid waste, therefore a value (0.16)

between the default values representative for food (0.15) and garden (0.2) was chosen for degradable organic carbon (DOC) content. As garden waste is reported now separately, there is no need any more for changing the default DOCs.)

Table 7.2.2 Used DOC content and methane generation rate constant of different waste categories

	DOC IPCC 2006	DOC Used values	Methane generation rate constant (k)
Paper	0.4	0.4	0.04
Textiles	0.24	0.24	0.04
Food	0.15	0.15	0.06
Garden	0.2	0.2	0.05
Wood	0.43	0.43	0.02
Sewage sludge	0.05	0.5*	0.06
Hygienic waste	0.24	0.24	0.05
Construction and demolition	0.04	0.04	0.05
DOCF	0.5	0.5	

*Calculated on dry matter basis

In earlier submissions, constant methane correction factor of 1.0 valid for well managed landfills was used for the entire time series. This approach could be regarded as overly conservative as it did not take into account the modernization process in solid waste disposal practices and available information on landfill sites. Let us quote a study "Landfills in Hungary" under a research framework "Organising for EU Enlargement: A Challenge for the Member States and the Candidate States"

State-of-the-art: the Hungarian landfills

In Hungary the typical form of managing waste is disposal: 85 per cent of collected waste is disposed of into landfills. According to a PHARE project designed to assess all landfills in the country, there were 2700 operational landfills, out of these only 728 were registered landfills serving all the municipalities in Hungary in 2002 (European Commission, 2001). Only 6 are so far in line with the *acquis* and a further 67 seem to be aligned to a large extent. In particular, a great number of low capacity local landfills do not conform to the *acquis* and there are a large number of illegal ones. Measures are being implemented to close down all the illegal or non-EU compliant landfills. The objective laid down in the National Waste Management Plan is to establish regional collection and management systems with a maximum of 100-120 landfills. At the end of 2004, the regional collection and management systems planned in the framework of ISPA programme cover one-third of the country. A further objective is that the abovementioned cover-rate should reach 100 per cent by 2009.]

In 2002, a comprehensive survey of landfill sites was carried out with the support of PHARE. During this project, stock was taken of no less than 2,667 landfill sites of which 1,300 were already closed. Out of the operating 1,367 sites, only 42 met current environmental requirements. It was suggested, though, that further 216 sites could operate temporarily till 2009, and the rest should be closed. One of the outcomes of the project was a database of landfills with several attributes such as depth, volume, insulation, cover etc. The database contained information among others on controlling, lining, compacting, leachate drainage, biogas collection. Summarizing the data based on total volume of disposed waste (and not on number of landfills), 15% of the disposal could be classified as managed (controlled), 16% as unmanaged shallow, and the remaining 69% as unmanaged deep.

Based on the above information, it didn't seem to be appropriate anymore to allocate all waste disposals to the managed category for the entire time series. Instead, all disposed waste is allocated now to the uncategorized category between 1950 and 1974. For the next period, between 1975 and 2000, the outcome of the above mentioned PHARE project is used, i.e. 85% of the disposed waste is considered as unmanaged (mostly deep), and the remaining 15% as managed. From 2001 on, all disposals are regarded as managed reflecting also the fact that a domestic act on waste management came into force in 2000 (Act No. XLIII of 2000 on waste management).

Naturally, changes did not occur from one day to another. Still, the development was quite rapid. Hungary started a modernization program relating disposal sites: the number of SWDSs decreased from about 2700 to 701 in 2000, then to 340 in 2005, to 213 in 2008, and to 69 in 2011. Currently (2012), 72 disposal sites are in operation.

Parallel to the closure of obsolete sites, the general level of management of the remaining disposal sites must have been improved.

Some domestic statistics indicate that only about 4% of municipal waste was still disposed uncontrolled in the early 2000's, therefore we decided to use $MCF=1$ for all years after 2000 which might be a little conservative estimate for the transition years.

As a consequence of this new approach, the formerly used parameters, especially the constant MCF value of 1.0 had to be replaced as follows:

- 1950-1974: $MCF=0.6$ for uncategorized SWDS, $OX=0$.
- 1975-1985: $MCF=0.77$ representing 15% managed, 16% unmanaged shallow and 69% unmanaged deep disposal. $OX=0$. (This means $MCF=0.72$ in the unmanaged category, and $MCF=1.0$ in the managed category.)
- 1986-2000: $MCF=0.77-0.81$ keeping the same share of managed/unmanaged sites but gradually decreasing shallow disposal. (This means $MCF=0.72-0.78$ in the unmanaged, and $MCF=1.0$ in the managed category.) $OX=0$
- 2001-2003 $MCF=1.0$ $OX=0$
- 2004-2009 $MCF=1.0$, $OX=0.05-0.1$.
- 2009- $MCF=1.0$, $OX=0.1$.

As for the oxidation factor, previously the default zero value was applied for the entire time series. However, based on the IPCC Guidelines, the use of the oxidation value of 0.1 is justified for covered, well managed SWDS to estimate both diffusion through the cap and escape by cracks/fissures.

The Hungarian Waste Information System that serves as our main source of information for activity data from 2004 contains two categories for disposals:

D1 Deposit into or onto land, e.g. landfill

D5 Specially engineered landfill, e.g. placement into lined discrete cells which are capped and isolated from one another and the environment

Landfills categorized as D5 can be regarded as well-managed modern landfills that comply with the EU Landfill Directive and where daily and temporary soil covers are applied therefore an oxidation value of 0.1 is justified. The following table shows, how the share of disposal into well-managed landfills increased in the last years.

Table 7.2.3 *The ratio of managed vs. well-managed landfills*

	D1	D5
2004	50%	50%
2005	48%	52%
2006	34%	66%
2007	36%	64%
2008	44%	56%
2009	29%	71%
2010	17%	83%
2011	35%	65%
2012	12%	88%
2013	2%	98%
2014	1%	99%
2015	1%	99%
2016	1%	99%
2017	0%	100%
2018	0%	100%
2019	0%	100%

Also managed landfills (D1) are covered in Hungary but not necessarily immediately after waste was deposited. In some instances the application of a soil cover oxidation might be delayed owing to modernization efforts or post-closure management. In 2017, for example, all managed (D1 and D5) landfills in Hungary were covered. For the period before 2003 Hungary does not have information on covering of managed landfills and therefore 0 per cent oxidation is assumed.

A 2009-report on the implementation of the EU Landfill Directive indicates that in the period 2007-2009 all managed landfills already met the requirements from the EU Landfill Directive and are therefore covered. Based on the conclusion of the above mentioned report from 2009 onwards OX=0.1 is assumed.

The amount of recovered CH₄ was calculated on the basis of energy production data obtained from the Energy Centre Hungary. These data in energy unit (TJ) were converted to mass unit as the amount of recovered methane by using the net calorific value from Table 1.2 in the 2006 IPCC Guidelines (Volume 2, Chapter 1), which is 50.4 TJ/Gg. Data collection has also been started on flaring. Disposal sites were contacted and asked for amount of flared landfill gas and methane content when available. In case site-specific methane content was not available, it was assumed that the share of methane in biogas was 50%. The collected data are summarized in the table below.

Table 7.2.4 *Data on flaring and biogas utilization (2001-2020)*

	2001	2002	2003	2004	2005	2006	2007	2008	2009
Flaring Mm3	3.060	3.115	2.868	2.893	3.230	7.753	7.353	9.500	4.357
CH₄ kt	1.0970	1.1167	1.0282	1.0371	1.1580	2.7794	2.6362	3.4058	1.5620
Biogas TJ	-	-	-	-	2	46	85	86	119
CH₄ kt					0.04	0.91	1.69	1.71	2.36
	2010	2011	2012	2013	2014	2015	2016	2017	2018
Flaring Mm3	4.050	4.271	6.283	7.929	7.438	2.107	0.692	0.958	0.750
CH₄ kt	1.4519	1.5310	2.2525	2.8994	2.7642	0.8566	0.2045	0.225	0.269
Biogas TJ	199	462	190	471	576	674	771	631	530
CH₄ kt	3.95	9.17	3.77	9.35	11.43	13.37	15.30	12.52	10.52

	2019	2020
Flaring Mm3	0.877	0.910
CH4 kt	0.314	0.326
Biogas TJ	438	410
CH4 kt	8.7	8.1

Recovered methane has been subtracted from the calculated emissions.

Please note that the earliest available data on the amount of CH₄ flared are from 2001 therefore NE was reported previously for earlier years. It is most possible that flaring activity did not occur before 2001 as landfill gas production also started only in 2005. Therefore, we have changed the notation key to “NO”.

The following figure summarizes the used activity data and the results of our calculations.




Figure 7.2.4 Summary of activity data and the resulting emissions

7.2.3 Uncertainties and time-series consistency

Uncertainty can be estimated using Table 3.5 of the 2006 Guidelines. Accordingly, the following values were obtained:

Quantity of disposed municipal solid wastes	>±10%
Degradable organic carbon	±20%
Fraction of Degradable Organic Carbon Decomposed	±20%
CH ₄ correction factor (=1)	-10 %, +0 %
CH ₄ content of landfill gases (0.5)	±5%
CH ₄ recovery	one order of magnitude
Half-life	±25%

The time series can be regarded as consistent.

(In the past, complete and obligatory data reporting on the collection of municipal solid waste did not exist in Hungary and the published data were estimations partly based on representative surveys. During the initial part of the calculation period, the authority procedures for waste recording were not uniform. In this system, which was based on self-reporting (self-registering), data were processed at varying detail and quality levels due to the lack of legal and technical regulations related to individual waste types. In addition, an overall central registry of industrial waste was missing and the rules related to such wastes were not laid down in any legal instruments).

7.2.4 QA/QC information

The compiler institute has now direct access to the Waste Management Information System maintained by the ministry responsible for environment. Data from different sources are compared. Our most detailed data source is the Waste Management Information System (EHIR) maintained now by the Ministry of Agriculture. This contains among others data on amount, type, consistency, management practices, mode of treatment. Converting these data to an Excel file, we get about 16,000 rows for one year. After analyzing these data, comparisons are made with the aggregated data published by the Hungarian Statistical Office, and also by EUROSTAT. Should we detect any problem, both the statistical office and the ministry can be contacted. The calculations with the IPCC Waste Spreadsheet Model have been saved and archived for future reviews.

7.2.5 Recalculation

There was no recalculation for this submission.

7.2.6 Planned improvements

It is planned to analyze the latest most detailed waste composition analysis available.

7.3 Biological treatment of solid waste (CRF sector 5B)

Emitted gases: CH₄, N₂O

Key source: none

As composting is showing a growing tendency recently, GHG emissions were calculated and reported for this category also in the submissions in the first commitment period using the IPCC 2006 methodology.

7.3.1 Methodological issues

The Tier 1 method from 2006 Guidelines was used with default emission factors.

Table 7.3.1 Activity data and emissions from biological treatment of solid waste

	Composting			CH ₄ (kt)	N ₂ O (kt)	Biogas facilities	
	MSW (kt) wet weight	dry weight	Sludge (kt)			Biogas (TJ)	CH ₄ (kt)
1985			20	0.20	0.01		
1986			20	0.20	0.01		
1987			20	0.20	0.01		
1988			20	0.20	0.01		
1989			20	0.20	0.01		
1990			20	0.20	0.01		
1991			20	0.20	0.01		
1992			20	0.20	0.01		
1993			20	0.20	0.01		
1994			20	0.20	0.01		
1995			28	0.28	0.02		
1996	18	7	29	0.36	0.02		
1997	19	8	26	0.34	0.02		
1998	18	7	23	0.30	0.02		
1999	18	7	32	0.39	0.02		
2000	17	7	30	0.37	0.02	6	0.00
2001	17	7	27	0.34	0.02	4	0.00
2002	47	19	37	0.56	0.03	4	0.00
2003	47	19	56	0.75	0.04	62	0.05
2004	39	16	24	0.40	0.02	107	0.08
2005	41	16	53	0.69	0.04	102	0.08
2006	58	23	43	0.66	0.04	129	0.10
2007	64	26	51	0.77	0.05	250	0.20
2008	85	34	62	0.96	0.06	490	0.38
2009	90	36	90	1.26	0.08	734	0.57
2010	148	59	82	1.41	0.08	898	0.70
2011	183	73	81	1.55	0.09	1336	1.05
2012	183	73	90	1.63	0.10	1105	0.87
2013	187	75	93	1.68	0.10	2042	1.60
2014	236	94	97	1.92	0.12	2018	1.58
2015	231	92	99	1.92	0.11	1954	1.53
2016	294	118	102	2.19	0.13	1965	1.54
2017	309	124	103	2.26	0.14	2297	1.80
2018	310	124	99	2.23	0.13	2127	1.67
2019	353	141	92	2.33	0.14	2142	1.68
2020	383	153	93	2.46	0.15	2108	1.65

The amount of composted municipal waste was received from the Hungarian Central Statistical Office. In 2020, 383.041 Gg waste was composted which represented 11% of all generated MSW.

As regards the amount of composted sludge, the time series was constructed using the following data sources:

- Data published by the statistical office;
- Composting related information from the Waste Management Information System (the same database that is used for SWDS);
- Data from the Wastewater Information System or for recent years from the Urban Wastewater Information System (the same databases that are used for emission calculations for wastewater treatment);
- For the period 1985-1993 we used a constant value corresponding to the amount reported for 1994.

In 2020 92.74 kt (dm) sludge was composted.

As generally the calculations were carried out on **dry weight basis**, and the corresponding emission factors from Table 4.1 in the 2006 IPCC Guidelines were applied (i.e., 10 g CH₄/kg dry waste and 0.6 g N₂O/kg dry waste), some of the original data had to be converted to dry weight. Sludge data in the Waste Management Information System are categorized on the basis of their dry matter content which allowed this conversion. For composted municipal waste, 60% moisture content was assumed (see Remarks in Table 4.1 in the IPCC Guidelines).

Our starting point for estimating methane emission from anaerobic digestion at biogas facilities was produced biogas from the energy statistics. The energy values (TJ) were then converted to mass of methane (kt) using the default calorific value of biogas, i.e., 50.4 TJ/Gg. Emissions of CH₄ due to unintentional leakages at biogas facilities were then assumed to be 5% as suggested by the 2006 IPCC Guidelines.

Although not used in the calculation, previously we also reported activity data (annual waste amount treated in kt dm). For the period 2015-2018, a very detailed database on various feedstock used for anaerobic digestion was analyzed. This database contained information on more than 40 types of feedstock, including fresh weight and dry matter content which could then be used directly. (Unfortunately, we found some data problems for 2015-17 therefore for this submission only data from 2018 was used.) For the remaining part of the time series, data on produced biogas (TJ) taken from the IEA/Eurostat Annual Questionnaire was used with a conversion factor of 7.3 TJ/kt dm (which was derived from the data for 2018).

Please note that there is no information on flaring activity for 5.B.1 Composting therefore NE was reported previously for flaring. As we believe, flaring is not occurring in composting plants, we have replaced the notation key with NO.

7.3.2 Uncertainties and time-series consistency

No category specific information is available.

7.3.3 Source-specific QA/QC and verification

The used data from Eurostat was compared with data from the Hungarian Central Statistical Office.

7.3.4 Recalculations

Data of biogas produced have been revised. Biogas from manure management has been subtracted from biogas accounted for under 5.B.2.b as it is accounted for under the *Agriculture* sector. In agreement with the agriculture expert, and on the basis of feedstock data from the period 2017-2020, it is assumed that 25% of all biogas is produced from manure. (The average share of manure was 26.4% in the analyzed period, so 25% seemed to be a safe estimate.)

7.3.5 Planned improvements

None.

7.4 Incineration of waste (CRF sector 5C)

Emitted gases: CO₂, CH₄, N₂O

Key source: none

7.4.1 Source category description

This subsector covers only emissions from thermal waste treatment without energy recovery (D10). Emissions from waste incineration for energy purposes (R1) are allocated to the energy sector.

During waste incineration, mainly CO₂ is emitted out of which only the fossil part contributes to the total emissions. (Biogenic CO₂ emissions were calculated as well but these were included only as memo items). Methane emissions are insignificant and N₂O generation is also minimal.

In 1986 a decree on the protection of air quality came into force, under which waste incineration (of any kind) required authorization. In 2001, decree 21/2001 (II.14) came into force explicitly prohibiting the open burning of waste, including in household furnaces. The same prohibition was included in the current Government Decree on air protection (306/2010 (XII. 23.)). Based on a Government Decree on air protection (306/2010 (XII. 23.)), open burning of waste (or incineration of waste in an installation that does not comply with the legislation setting the conditions for incineration of waste), with the exception of household waste paper and incineration of untreated non-hazardous wood waste is prohibited.




Figure 7.4.1 Activity data in waste incineration

7.4.2 Methodological issues

For estimating CO₂ emissions, the Tier 2 method was applied as country-specific data on waste amount, composition and management practices was used. The very detailed Hungarian Waste Management Information System made it possible to disaggregate the activity data (amount of incinerated waste) into different waste types according to the European Waste Catalogue (EWC codes) and European Waste Classification (EWC-Stat). It might be an interesting fact that 82 to 97 per cent of all incinerated waste in this source category was hazardous waste of which most part was liquid. Therefore, it was decided that all incineration would be reported in the “Hazardous Waste” category in the CRF. However, different emission factors were applied in the emission calculations for the following waste types:

- clinical waste: defined as EWC-Stat code W05 (Health care and biological wastes)
- fossil liquid waste: defined as liquid wastes with EWC-Stat codes W011 (spent solvents) and W013 (used oils);
- hazardous (non-liquid, non-clinical) waste: defined as all other hazardous waste;
- other waste: all other incinerated waste

Having these country-specific waste amount and composition data, the carbon content of the incinerated waste and the fossil fraction thereof could be determined by using default values from Table 2.5 and Table 2.6 in the 2006 Guidelines (Volume 5. Ch. 2). For liquid waste, the default carbon content of 80% was applied from Table 5.2 of the 2006 IPCC Guidelines. For hazardous waste, the average of 5-50 per cent was used, i.e. 0.275. The same value was applied for all other wastes.

The CH₄ emission factor of 300 kg/kt was derived from the default emission factor of 30 kg/TJ for industrial wastes (Table 2.3 in Chapter 2) with an assumed calorific value of 10 TJ/kt.

The following table summarizes the used emission factors for the different waste types.

Table 7.4.1 *The used GHG emission factors*

Waste type	Carbon EF	Ref.	CH4	Ref.	N2O	Ref.
Fossil liquid	0.8	Table 5.2	0.56	Page 5.20	9.8	Table 5.5 (waste oil)
Clinical	0.25	Table 2.6	300	Chapter 2	100	Table 5.6 (industrial waste)
Hazardous (non-liquid, non-clinical)	0.275	Table 2.6	300	Chapter 2	100	Table 5.6 (industrial waste)
Other waste	0.275		300	Chapter 2	100	Table 5.6 (industrial waste)

The CH₄ emission factor of 300 kg/kt was derived from the default emission factor of 30 kg/TJ for industrial wastes (Table 2.3 in Chapter 2) with an assumed calorific value of 10 TJ/kt.

7.4.3 Uncertainties and time-series consistency

Consistency of the time series needs to be investigated, as constant values are used for the years before 2004.

7.4.4 QA/QC information

Data taken from the Hungarian Waste Management Information System for the calculations are compared with the relevant data published by the domestic statistical office and by Eurostat. As most part of waste incineration occurs with energy recovery, even if the resulting emissions are accounted for in the energy sector it is worth mentioning here that the IEA and ETS data were cross-checked, and also the biggest incinerator plant is contacted once in a while for verification purposes.

7.4.5 Recalculations

No recalculation has been made for this submission.

7.4.6 Source-specific planned improvements

None.

7.5 Wastewater treatment and discharge (CRF sector 5D)

Emitted gas: CH₄, N₂O

Key source: CH₄: Level

7.5.1 Source category description

This sector covers emissions generated during municipal and industrial wastewater treatment. When the wastewater is treated anaerobically, methane is produced. Wastewater handling can also be a source of nitrous oxide, therefore N₂O emissions from human sewage are also part of the inventory.

7.5.2 Methodological issues

While estimating the methane emissions of wastewater handling, the key parameter is the fraction of wastewater treated anaerobically. Methane emissions from wastewater treatment were calculated using partly basic statistical data, partly very detailed facility level information on wastewater discharge together with the specific emission factors recommended by the 2006 IPCC Guidelines. For recent years, wastewater data (COD values for the industrial sector, proportion of different treatment methods) based on measurements conducted by the authorities and emitters were obtained from the regional inspectorates for environment, nature and water. Besides, the inventory compilers consulted with experts, visited a few wastewater plants and checked the calculations of the neighboring countries as well.

General background information

Based on data of the Wastewater Information System that contains about 1500 emission reports per year, from an analysis of the period 2005-2013 the following conclusions could be drawn:

- 140 to 240 million cubic meter wastewater from industrial facilities was discharged into rivers and seas. On average, 60% of this amount had either no treatment or only mechanical treatment beforehand (trend decreasing);
- The average COD content of the above, only partially treated wastewater was as low as 0.05 kg/m³.
- On average, about 30 million m³ industrial wastewater was collected via the public sewerage system, and treated in centralized plants, consequently domestic and industrial wastewater treatment could not be separated entirely. Around 80% of this amount went into the public sewerage system after at least biological treatment.
- The average COD content of the above, mostly treated industrial wastewater was definitely higher with 0.6 to 0.8 kg/m³ in 2008-2012.
- Domestic and commercial wastewater treatment plants, (that also treat industrial wastewater), discharge yearly 440 to 580 million m³ into open water.
- The share of the collected wastewater treated at least biologically or at more advanced treatment plants increased from 35% in 1990 to 65% in 2005, and to 96-98% in 2010-2017;
- In line with the above development, the average BOD₅ content of the discharged wastewater decreased from 0.15 kg/m³ in 2005 to 0.02 kg/m³ in 2013.

Activity data

For domestic wastewater, the activity data - the quantity of total organic waste (TOW) - was calculated by multiplying the population of the country by the IPCC default value of Biochemical Oxygen Demand that is $BOD_5 = 60 \text{ g/person/day}$ (Table 6.4 in Volume 5 Chapter 6 of the 2006 IPCC Guidelines). This default BOD value was confirmed by Hungarian experts of the Ministry of Environment and Water as well and was used uniformly for the entire times series and for the whole country.

Total organics produced by industrial facilities is partly taken into account with the default correction factor $I (=1.25)$ corresponding to additional industrial BOD discharged into sewers thus accounted for in the domestic category. From 2015 on, total organic load as reported by wastewater treatment plants to the statistical office are also taken into consideration. The average difference between the calculated and the reported total BOD values was 3% with a growing tendency.

As described above, the degree of utilization of modern, centralized WWT plants increased, and tertiary treatment became the dominant technology.




Figure 7.5.1 The evolution of the different treatment systems (1990–)

Source of data: Hungarian Central Statistical Office:

http://www.ksh.hu/docs/eng/xstadat/xstadat_annual/i_uw005.html

The activity data in the industrial wastewater category were the total output of wastewater [1000m³/year], the emitted total organic wastewater [kg COD/year] and in some cases the organic load (kg BOD/day) which were collected by the regional inspectorates and further processed by the (former) Research Institute for Environmental and Water Management (VITUKI). However, limited data were available on the industrial wastewater generation in individual sectors, especially for the initial years of the calculation period.

Activity data for the industrial wastewater category were partly taken from the Wastewater Information System database. For earlier years, before 2008, activity data were extrapolated using proxy data, i.e. volume of water supplied to other than household consumers published by the

statistical office. (see http://www.ksh.hu/docs/eng/xstadat/xstadat_annual/i_uw004.html) For the period 1985-1990, constant values are used.

For a few years now, we have received a very detailed database containing all domestic wastewater treatment plants (around 800) collected by General Directorate of Water Management, an independently operating institute and a central government body under the direction and supervision of the Minister of Interior.

This database contains among others information on the amount of treated wastewater by treatment type, but also other inventory relevant parameters such as BOD, COD, and Nitrogen content of both the influent and the effluent, and information on sludge treatment.

Few years ago, we had also the possibility to look also into detailed reports of wastewater facilities of different industrial plants. Special emphasis was given to industries with high COD output, e.g. food and beverage, paper and pulp, chemical industry etc. By analyzing organic load data before treatment, we were able to introduce the following country-specific data on industrial wastewater as summarized in Table 7.5.1 below:

Table 7.5.1 *The used data for industrial wastewater*

	BOD
	[kg/m3]
Pulp and paper	2
Starch	1.14
Sugar	3.4
Pharmaceutical	1.5
Beer	1.5
Meat	1
Dairy products	1.5
Vegetable oils	0.85
Wine	5.27
Fruits	2.9
Chemical industry	0.25
Coke production*	5
Oil refinery*	1

**refers to COD*

As for industrial wastewater, generally COD values are used, the above BOD values were converted using a conversion factor of 2.4. Please note, as we expect more data from individual facilities, the above data might be subject of changes.

Emission factors

For the calculation of the emission factor (EF), default maximum CH₄ producing capacities of 0.25 kg CH₄/kg COD and 0.6 kg CH₄/kg BOD were used for industrial and domestic wastewater, respectively.

The choice of a proper methane conversion factor (MCF) was somewhat more difficult. To calculate the weighted average of MCF, additional information was collected on the share of population with no connection to the public sewerage system. Using these additional activity data, the following assumptions were made:

- In accordance with the 2006 IPCC Guidelines, for people using septic systems or any other domestic means (no connection to public sewerage network), it can be assumed that half of

the BOD settles, therefore $MCF=0.5$ was chosen. (Table 6.3 in the 2006 Guidelines). In the base year, the portion of population (or dwellings) connected to public sewerage system was 38%, now it's around 81%. It must be noted, however, that the percentage of dwellings connected to public sewerage network is still below the Central-European average.

- As a refinement of the above, for those dwellings where neither public nor domestic sewerage exists and probably latrines are used, $MCF=0.1$ was used in accordance with the above referenced table from the 2006 Guidelines.
- Annual data on the number of dwellings connected to public sewerage systems are provided by the Hungarian Central Statistical Office for 1990–2020. Data on the proportion of the population connected thereto are available for 2011 onward. Prior to 2011 the share of population connected or not connected to a sewerage system was estimated based on the number of all dwellings using a correction factor of 5%.
- Usually, collected wastewater undergoes aerobic treatment in treatment plants. Default MCF for centralized, aerobic treatment plant is zero. (Table 6.3 in the 2006 Guidelines) Still, an MCF value of 0.05 was applied to secondary treatment up to 2004, thus to allow some emissions in case of incidental overload, and more importantly, to reflect modernization in the sector. (In a previous submission, the default MCF value of 0.1 was applied for direct discharge into rivers and lakes. This was abolished following a recommendation of a European review team, since it can be assumed that flowing rivers (such as the Danube) are not oxygen-deficient.)
- For industrial wastewater, MCF of 0.05 (i.e. the middle of the range for aerobic treatment from Table 6.8 in the Guidelines) is used from 2000, and the highest value of the range ($MCF=0.1$) for the beginning of the time series (until 1995), and interpolated values between 1995 and 2000. We applied the general assumptions that industrial wastewater is treated aerobically and that, in the case of anaerobic sludge treatment, CH_4 generated is recovered as sewage sludge gas with an assumed leakage of 5 per cent.

Methane recovery

Based on the energy statistics, sewage sludge gas utilization started in 2001 in Hungary. As unintentional leakage might occur during anaerobic digestion of sewage sludge, some CH_4 emission are added to this category. The same methodology was used as for the category 5B Biological treatment of solid waste, i.e., 5% leakage was assumed. (Please note that the total amount of CH_4 for energy recovery is reported under category 5.D.1 domestic wastewater as this value is calculated on the basis of energy statistics and the data available include sewage sludge gas from both municipal and industrial wastewater plants. For this reason notation key IE is reported for 5.D.2 Industrial wastewater.)




Figure 7.5.2 Sewage gas production (TJ)

As regards flaring, we were not aware of any official database therefore individual wastewater treatment plants were contacted. About 10 WWTPs reported flaring activity. The aggregate amount of flared methane expressed both in Nm³ and kt is included in Figure 7.5.3. Methane density was assumed as 0.717 kg/m³. As it can be seen, the amount of flared methane is far below 1 kt. (Please note that flared methane is not part of the emission estimates.)

Please note that flaring did not occur on domestic wastewater treatment plants in years 1990-2000 therefore notation key NO is reported. As there is no information on flaring activity for subcategory 5.D.1 for years 2001-2003 the notation key NE is reported.

Please note that there is no information on flaring activity for subcategory 5.D.2 Industrial wastewater therefore NE is reported from 2001 on (previously no sewage gas production occurred).




Figure 7.5.3 Flared methane in WWTPs (2004-2020)

The above considerations, used parameters and the resulting emissions are summarized in the Figure below. It seems obvious that the decrease in emission is mostly due to the growing share of households connected to the public sewerage system that increased from 43.7% in 1990 to 86.9% in 2020.




Figure 7.5.4 Domestic wastewater treatment (1985-2020)

As regards emissions from sludge treatment, our approach is as follows. It is assumed that whenever anaerobic digestion of sludge takes place, the generated methane (reported as sludge gas in the energy statistics) is recovered and used for energy purposes. The amount of methane recovered for energy is reported in CRF Table 5D but this amount was not subtracted from the total emissions as no additional methane emission from sludge digestion was taken into account in the above calculations.

It has to be emphasized that emissions from sludge treatment (besides leakage at biogas facilities) are taken into account in other emission categories:

- Landfilled sewage sludge is accounted for in the 5A Solid waste disposal in landfills category;
- Composted sewage sludge is taken into account in the 5B Biological treatment of solid waste category;

N₂O emissions from wastewater has been significantly revised for the previous submission. Considering direct emissions, the calculation method for nitrous oxide emissions from advanced centralized wastewater treatment plants remained the same, i.e., the method described in Box 6.1 in the 2006 IPCC Guidelines was used. We have, however, changed our approach for indirect N₂O emissions on the basis of newly collected data. Based on measurements of the total nitrogen content of incoming wastewater, we have re-evaluated the total N content in wastewater. It turned out that there is no need to apply the parameters for non-consumed protein and for industrial a commercial co-discharged protein.

For example, modern (third category) domestic wastewater plants measured 30.9 kt nitrogen in the influent in 2015 based on information collected by General Directorate of Water Management and further processed by the compiler institute. We know that 85% of all wastewater went in such advanced WWTPs. We also know that 83% of the population is connected. Based on protein consumption (100.1 g/cap/day), the estimated N in the effluent would be 58 kt (without applying the

parameters $F_{non-con}$ and $F_{ind-com}$). Using this figure with the degree of utilization of modern centralized WWT plants (85%) and the connection rate of the population (83%), we would get $55 \times 85\% \times 83\% = 40.2$ kt that is higher by more than 20% than the measured value, therefore it seems that there is no need for these two additional F parameters. (We got similar results for the period 2011-2018 where measured data were available.)

Consequently, we have recalculated the basic activity data (i.e., available nitrogen in the effluent) by using the same protein consumption data but not taking into account the default parameters for non-consumed protein and for industrial a commercial co-discharged protein. And we went one step further. As we have measured data for the nitrogen content in both the influent and the effluent (see Table 7.5.2), their difference was considered as removed nitrogen in the treatment process that won't be discharged to aquatic environments anymore, therefore it was subtracted from the calculated value of N in effluent.

Table 7.5.2 Measured N content in wastewater in advanced WWT plants (tertiary treatment)

	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total N in influent (kt)	18.36	23.59	24.68	28.58	30.86	33.82	34.13	39.01	36.11	36.67
Total N in effluent (kt)	4.28	4.75	4.63	4.85	5.34	5.45	5.62	6.57	6.10	6.53

For a comparison, the table below summarizes the overall performance of all domestic wastewater treatment plants in Hungary on the basis of data collected by the statistical office. As it can be seen, BOD removal efficiency is 97%, N -removal efficiency is around 82%. (These data are not yet used in the calculations.)

Table 7.5.2a Summary table of the overall performance of domestic WWTPs in Hungary

	2015	2016	2017	2018	2019	2020
BOD_in_kt	213.7	233.0	244.6	245.2	247.9	247.3
COD_in_kt	404.9	419.3	441.2	444.9	447.3	452.0
COD/BOD	1.89	1.80	1.80	1.81	1.80	1.83
BOD_effl_kt	6.3	6.8	7.2	7.6	8.0	7.8
BOD_removal	0.97	0.97	0.97	0.97	0.97	0.97
N_{in_kt}	39.2	43.2	42.9	43.6	44.4	42.1
$N_{effluent_kt}$	7.4	7.4	7.4	7.9	8.0	7.8
Removal (%)	81%	83%	83%	82%	82%	81%

Our results are summarized in Table 7.5.3 and Figure 7.5.5. Total N was estimated from the statistics "per capita amount of available proteins" with the parameter "fraction of nitrogen in protein, default = 0.16 kg N /kg protein". The used emission factor for N_2O effluent was also a default one from Table 6.11 of the 2006 IPCC Guidelines (i.e., 0.005 kg N_2O - N /kg- N). In the reported N_2O emissions in the CRF Table 5.D, however, also emissions from advanced treatment plants are included (see " N_2O plants" in Table 7.5.3) which was calculated applying Equation 6.9 from the Guidelines with country-specific and

yearly changing values for P (human population) and T_{plant} (degree of utilization of modern, centralized WWT plants), and defaults for F_{ind-com} and EF_{plant}. This is why the IEF is somewhat higher than 0.005.

Table 7.5.3 Protein consumption and all the resulting N₂O emissions

Year	Protein [g/day]	Total N [kt]	N_removed [kt]	N_effluent [kt]	N2O plants [kt]	N2O effluent [kt]	N2O Total [kt]
1985	106.1	65.7	0.0	65.7	0.00	0.52	0.52
BY	107.3	66.1	0.0	66.1	0.00	0.52	0.52
1986	106.4	65.6	0.0	65.6	0.00	0.52	0.52
1987	109.4	67.1	0.0	67.1	0.00	0.53	0.53
1988	107.8	65.9	0.0	65.9	0.00	0.52	0.52
1989	108.8	66.2	0.0	66.2	0.00	0.52	0.52
1990	104.7	63.4	0.5	62.9	0.00	0.49	0.49
1991	100.5	60.9	0.5	60.4	0.00	0.47	0.48
1992	101.1	61.2	0.4	60.8	0.00	0.48	0.48
1993	94.8	57.4	0.5	56.9	0.00	0.45	0.45
1994	91.3	55.2	0.3	54.8	0.00	0.43	0.43
1995	87.0	52.5	0.3	52.2	0.00	0.41	0.41
1996	84.4	50.9	0.4	50.5	0.00	0.40	0.40
1997	87.7	52.8	0.3	52.4	0.00	0.41	0.41
1998	87.2	52.4	1.0	51.4	0.00	0.40	0.40
1999	89.5	53.6	1.1	52.4	0.00	0.41	0.41
2000	96.6	57.7	2.2	55.5	0.00	0.44	0.44
2001	93.9	55.9	2.4	53.5	0.00	0.42	0.42
2002	93.5	55.6	3.7	51.8	0.00	0.41	0.41
2003	103.0	61.0	7.7	53.3	0.01	0.42	0.43
2004	101.0	59.7	8.1	51.6	0.01	0.41	0.41
2005	105.4	62.2	9.3	52.9	0.01	0.42	0.42
2006	104.6	61.6	6.7	54.9	0.01	0.43	0.44
2007	101.3	59.5	8.8	50.7	0.01	0.40	0.41
2008	100.6	59.0	9.5	49.5	0.01	0.39	0.40
2009	99.5	58.3	9.9	48.4	0.01	0.38	0.39
2010	95.8	56.0	12.7	43.3	0.01	0.34	0.35
2011	93.6	54.6	14.1	40.5	0.02	0.32	0.33
2012	93.4	54.2	18.8	35.3	0.02	0.28	0.30
2013	92.1	53.3	20.1	33.2	0.02	0.26	0.28
2014	95.7	55.2	23.7	31.5	0.02	0.25	0.27
2015	100.1	57.6	25.5	32.1	0.03	0.25	0.28
2016	103.2	59.2	28.4	30.9	0.03	0.24	0.27
2017	106.5	60.9	28.5	32.4	0.03	0.25	0.28
2018	109.9	62.8	32.4	30.3	0.03	0.24	0.27
2019	111.1	63.4	30.0	33.4	0.03	0.26	0.29
2020	111.1	63.4	30.1	33.2	0.03	0.26	0.29




Figure 7.5.5 Increasing share of tertiary treatment leads to decreasing N2O emissions

7.5.3 Uncertainties and time-series consistency

Based on the above considerations, the uncertainty of the calculation of the emissions from household wastewater is relatively high. In the industrial sector, data became more reliable in the recent years as a result of the new reporting requirements. However, they do not cover all the emitters, although the most important wastewater emitting sectors are included.

Uncertainty of the emissions from household wastewater treatment:

Per human populations	-5 % to +5 %
BOD/capita	-30 % to +30 %,
Maximum methane production capacity B0	-30 % to +30 %

Uncertainty of the emissions from industrial wastewater treatment:

Quantity of industrial wastewater:	-25 % to +25 %
Wastewater /unit of production COD/ unit of wastewater:	-50 % to +100 %
Maximum CH4 production capacity Bo :	-30 % to + 30 %

Uncertainty of N2O emissions

Emission factor	order of 2
Per capita protein consumption	±10%
Used factors	±20%

Source: according to the recommendations of the Revised Guidelines and 2006 Guidelines, on the basis of expert estimates

The time series of emissions are most probably consistent.

7.5.4 QA/QC information

The data collected by the environmental authorities were checked by an independent institution (VITUKI) that further processed the data. Data from the database of facility level wastewater information that are used ultimately by the inventory compiler institute undergoes basic checks, e.g. duplications are removed, outliers are analyzed and corrected whenever necessary.

7.5.5 Recalculation

Protein consumption data and the amount of wastewater undergoing tertiary treatment has been updated for year 2019. The resulting increase in emission (1.6 kt CO₂-eq) is negligible.

7.5.6 Planned improvements

More analyses of the industrial wastewater treatment facilities are planned to confirm or modify data in Table 7.5.2 above. More importantly, should the Refinement of the 2006 IPCC Guidelines be accepted, we plan to implement the revised methodology in a future submission.

8 OTHER (CRF SECTOR 6)

CRF Table 6 comprises NO_x emissions from 3B Manure management and total NH₃ emissions from 3. Agriculture to ensure the consistency with the reporting to the UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP) and the EU Directive on the reduction of national emissions of certain atmospheric pollutants.


9 INDIRECT CO₂ AND NITROUS OXIDE EMISSIONS

Not applicable in this submission.

10 RECALCULATIONS AND PLANNED IMPROVEMENTS

Since the 2015 submission, the methodologies provided in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories have been used in a consistent manner. The switch to the new methodological framework led to recalculations in every sector for the whole time series. The most fundamental changes were implemented for the 2015 submission. This time, rather the usual refinements have been carried out. Recalculations of some data-series of the inventory occur and can be justified by several reasons. Just to name a few, QA/QC procedures, ERT recommendations, changing for higher Tier methodologies can lead to a recalculation. As a basic rule, whenever new information emerges that improves the quality or accuracy of the emission data, the emissions are recalculated.

The overall effect of all recalculations described below was +9.96 kt CO₂-eq or 0.0% and +120,16 kt CO₂-eq or 0.2% of the total emissions in the base year and 2019, respectively. The figure below shows the total emissions from the base year until the last year assessed in the last two submissions (without LULUCF). The almost negligible effect of the recalculations can hardly be seen on this figure.



ENERGY

General: The latest versions of the Annual IEA/Eurostat Questionnaires submitted in November 2021 were used as activity data. All the changes in the energy statistics are reflected in the current inventory.

1.A.3.b: We have switched to the latest version of the COPERT model (version 5.5.1, September 2021) for the whole time series. This affected slightly the estimated fuel use in the different vehicle categories. In addition, the model provided revised N₂O emission estimates.

1.A.3.a: Activity data (aviation gasoline and kerosene used for national aviation) have been revised based on latest Eurocontrol data (2005-2018). This revision affected slightly also backward extrapolated data.

1.B.2.c Emissions from flaring in the refinery was omitted by mistake in the previous inventory.

1.B.2.d: Activity data have been revised for groundwater extraction, and the same country-specific emission factor has been applied for the whole time series.

Reallocations:

1.A.2.g: Emissions from all autoproducer plants have been (re-)allocated from the source category 1A2g to the relevant economic sector where they operate fully for the period 1998-2020, and to the extent possible also for earlier years.

1.B.2: to increase comparability, leakage and venting emissions from gas transmission and storage are reported separately on the basis of TABLE 4A.2.7 of the 2019 Refinement.

The overall effect of the above recalculations are as follows:

- emissions from combustion activities decreased by 3.40 kt CO₂-eq (-0.0% of total emissions) in the base year, and increased by 131.21 kt CO₂-eq (0.2% of total emissions) in 2019.
- fugitive emissions increased by 34.34 kt CO₂-eq in the base year, and by 18.53 kt CO₂-eq (0.0% of total emissions) in 2019.

INDUSTRIAL PROCESSES AND PRODUCT USE

2.D.1 – Lubricant use. The number of L-category vehicles was recalculated for the period 2009-2019, thus there are minor changes in the activity data and CO₂ emissions.

2.D.3 Other – Urea based catalysts. The number of Euro 6 vehicles was recalculated for 2018 and 2019, thus minor changes occur in the activity data and CO₂ emissions.

2.D.3 Other – Indirect emissions from solvent and other product uses. In March 2020, recalculation was performed for this sector. CO₂ emissions were recalculated and corrected in the 2020 and 2021 reports, while activity data (NMVOC emission from solvent and other product uses) remained unchanged. Activity data are now corrected for the whole time period.

2.D.3 Other – Indirect emissions from solvent and other product uses. Activity data for 2D3a Domestic solvent use and for 2D3d Coating and paints for 2019 was recalculated, thus minor changes occur in the activity data and CO₂ emissions of the sector.

2.F.1 – Correction of emission data from Mobile air-conditioning (2.F.1.e) sector (0.1 kt CO₂-eq).

2.F.2 – Revision of activity data (the impact of recalculation on total emissions including LULUCF is + 0.042 %).

The overall effect of the above recalculations was almost negligible: +4.8 kt CO₂-eq in 2019.

AGRICULTURE

The main reason for changes in the emissions from 3. Agriculture sector are the manure management data updates. The 2020 Agricultural Census and the recently available annual statistics on agricultural wastes used in biogas plants have made it possible to reflect the current practices and the effects of anaerobic digestion. As a result of this inventory improvements animal manure used in “digesters” has been reported at the first time in this submission.


Data from the nitrate database, which aims to monitor the implementation of the European Nitrate Directive (91/676/EEC), also provides data on manure management, grazing and since 2015 on manure management related NH_3 abatement technologies. Data from the nitrate database was also used to revise the data on manure management for the GHG- and the air pollutant emission inventory. Revision of NH_3 emissions affected the direct and indirect N_2O emissions from manure management as well as agricultural soils.

Reallocation of manure to “digesters” for the period 2004-2019 and the change in the proportion of covered liquid slurry for the whole time series caused the most significant changes in CH_4 and N_2O emissions from 3. Agriculture.

The overall effect of recalculation in the 3. Agriculture sector resulted in an increase in the emissions in the range of 0.4 to 12.4 kt CO_2 eq (0.0%-0.1%) in the period BY to 2002 and a decrease in the range of 2.7 to 35.2 kt CO_2 eq (0.0%-0.5%) between 2003 and 2019. The largest decrease, 35.2 kt CO_2 -eq occurred in 2015 and the largest increase, 12.4 kt CO_2 -eq in the base year (BY).

Impact of recalculations is shown in Figure 10.1.

Figure 10.1 Impact of recalculations in 3. Agriculture between annual submissions 1990-2019



Reasons for recalculations by CRF sectors are as follows:

3.A Enteric Fermentation

3.A.1 Enteric Fermentation -Dairy Cattle, 2004-2019

The HCSO has changed the methodology of data collection on milk fat and milk protein since 2019 and 2020, respectively. This change led to time series inconsistency, as the HCSO has not revised the data, retrospectively in line with the new methodology. To ensure the time series consistency the milk fat and protein content data were revised based on the Institute of Agricultural Economics' Market Price Information System (MPIS) data. The revised nutritional values of raw milk used to derive the gross energy intake (GE) for dairy cattle, resulted in an increase of 0.15% and 1.1 kt CO₂-eq on average in the CH₄ emissions from 3.A.1 Enteric Fermentation - Dairy Cattle for the 2004-2019 trend. The change ranged between -5.2 kt CO₂-eq in 2004 and 7.7 kt CO₂-eq in 2016.

3.A.1 Enteric Fermentation - Non-Dairy Cattle, heifers 1-2 yr, 2016-2019

A data linking error was corrected, because in the previous submissions the livestock number of heifers, 1-2 yrs for the year 2015 was used for the period 2016-2019 as well. Due to this recalculation emissions increased by 0.4%-2.2% for the period 2016-2019.

The overall effect of recalculations on 3.A Enteric Fermentation was a slight decrease in the emissions between 2004 and 2009 (except 2005) and increase in 2005 and over the period 2010-2019. The increase is the most significant (23.8 kt CO₂-eq) in 2019.

Revisions made in Sector 3.A, through gross energy intake for dairy cows and animal numbers for other cattle, also affected emissions from manure management.

3.B.1 CH₄ Emissions from Manure Management

3.B.1.1 and 3.B.1.3 CH₄ Emissions from Manure Management - Cattle and Swine, across the whole time series

Proportions of pig and cattle liquid/slurry covered by "natural crust" was revised for the whole time series. The former expert judgement was replaced by annual survey data, which are available from the year 2015. Between 2001 and 2015, interpolation was used for cover types other than natural crust, and for natural crust, 2015 data were used for the period before 2015, as well. This recalculation resulted in an increase in the MCF of liquid/slurry of cattle and swine. The increase is larger for cattle and minor for pigs.

3.B.1.1, 3.B.1.3 and 3.B.1.4 CH₄ / 3.B.2.1, 3.B.2.3 and 3.B.2.4 N₂O Emissions from Manure Management - Cattle, Swine and Poultry for the period 2004-2019

CH₄ and N₂O emissions from manure management of cattle, swine, and poultry for the period 2004-2019 was revised considering proportions of manure used in anaerobic digesters. In the previous submissions the anaerobic digested manure was allocated in line with the housing, irrespective of the manure use. In this submission, the anaerobic digested manure was reallocated to 'digesters', as data on the agricultural wastes used in anaerobic digesters had become available.

3.B.1.2 CH₄ Emissions from Manure Management - Sheep, 2014-2019

There are minor recalculations to CH₄ emissions from manure management of sheep due to the revision on the proportion of manure excreted during grazing considering the length of grazing period. This revision resulted in an insignificant (less than 0.5 kt CO₂-eq) increase in the CH₄ emissions from 3.B.1.2.

The overall changes in the CH₄ emissions from 3.B.1. Manure management ranged from a decrease of 25.5 kt CO₂-eq (4.1%) in 2013 to an increase of 15.7 kt CO₂-eq (1.3%) in the BY. Revision of “natural crust” resulted in an increase in the BY emissions, 3.0, 10.9 and 1.7 kt CO₂-eq for dairy cattle, non-dairy cattle, and swine, respectively. In contrast, reallocation of anaerobic digested manure to “digesters” resulted in a decrease in the emissions for the period 2004 to 2019, partly offset by the increase in emissions due to revised natural crust. The highest net decrease in the emissions due to anaerobic digestion were 9.4 kt CO₂-eq (in 2013), 5.3 kt CO₂-eq (in 2019) and 17.1 kt CO₂-eq (in 2014), for dairy-cattle, non-dairy cattle, and swine, respectively.

3.B.2 N₂O emissions from Manure Management

3.B.2 Direct N₂O emissions from manure management, whole time series

Revision of “natural crust” for the whole time series and the anaerobic digested manure for the period 2004-2019 resulted in an overall decrease of 1.3% and 4.1 kt CO₂-eq on average on the BY and 1990-2019 trend. Decrease in the emissions ranged from 1.1 kt CO₂-eq in 2000 to 18.3 kt CO₂-eq in 2019.

3.B.2.5 Indirect N₂O emissions from *manure management, whole time series*

Atmospheric Deposition, whole time series

Revision of NH₃ emissions from 3.B Manure Management for the reporting to the UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP) (UNECE, 1999) and the National Emissions Ceiling Directive (EP, CEU, 2016) resulted in changes in the indirect N₂O emissions from manure management due to atmospheric deposition. The revision of NH₃ emissions aimed to take account of emission abatement measures, as a result of new data collections recently launched.

The following abatement measures were considered in the revision of NH₃ emissions by stages of manure and animals:

Animal housing

- Non-leaking drinking systems for broilers
- Aviary system for layers
- Partially slatted floor for piglets after weaning and growers-finishers

Manure storage

- “Low technology” floating covers (e.g., chopped straw, peat, bark, etc.)
- “Tight lid”, roof or tent structure
- Plastic sheeting (floating cover)
- Natural crust

Revision of the NH_3 emissions from manure management lead to an 1.1% (1.5 kt CO_2 -eq) decrease in indirect N_2O emissions due to atmospheric deposition on average over the time series.

Nitrogen Leaching and Run-off, 2004-2019

Revision of data on manure management system distribution, mainly reallocation of manure to digesters, resulted in additional changes in the indirect N_2O emissions due to leaching and run-off. Changes ranged from a decrease of 0.08 kt CO_2 eq in 2019 (6.4%) to an increase of 0.02 kt CO_2 eq in 2016 (1.3%).

The overall indirect N_2O emissions from manure management decreased by 1.1% and 1.5 kt CO_2 eq on average in the BY and 1990-2019 trend.

The overall impact of revisions for the 3.B Manure Management sector CH_4 and N_2O emissions is an increase in emissions in the range of 1.2-13.3 kt CO_2 -eq (0.1%-0.7%) between BY and 2005, and a decrease of between 0.3 and 41.8 kt CO_2 -eq (0.0%-3.8%) for the years 2006-2019.

3.D Agricultural Soils, whole time series

3.D.1 Direct N_2O Emissions from Managed Soils, whole time series

3.D.1.2.a Animal Manure Applied to Soils

Due to the interlinking between the 3.B and 3.D CRF sectors through the N-balance, revisions to the 3.B resulted in revised emissions from 3.D.1.2a Animal manure applied to soils and 3.D.2 Indirect N_2O emissions from agricultural soils for the whole time series. Correction to the livestock number of "heifers 1-2 yrs" for the years 2016-2019, as mentioned in the paragraph related to the CRF Sector 3.A, resulted in some increase in the F_{AM} . The net effect of these recalculations is an insignificant reduction in the direct emissions from 3.D.1.2.a Direct N_2O emissions from agricultural soils - animal manure applied to soils over the period BY-2008 and an increase ranging from 0.02 kt CO_2 eq to 8.4 kt CO_2 eq over the period 2010-2019.

3.D.1.2.c Other Organic Fertilizers Applied to Soils

Data on composted sewage sludge and municipal waste have been revised to include only use on agricultural land. In the previous submissions, the total amount of composted sewage sludge and waste was used, which was not in line with the Hungarian regulation, because in Hungary, the quality of compost that can be applied is regulated by law, and application is subject to a permit, so not all compost can be used on agricultural land without restriction.

At the same time, as a result of our improvements in the utilization of agricultural waste in anaerobic digestion, digestate from anaerobic digestion, which was not included in previous submissions, is now included.

As a result of the revision of activity data, emissions from 3Da2c decreased by -0.5 to 4.4 kt CO_2 eq between BY and 2009, while they increased by 0.1 to 9.2 kt CO_2 eq between 2010 and 2019.

3.D.1.3 Urine and Dung Deposited by Grazing Animals

The update of the AWMS data also resulted in a minor change in the N excreted on grazing (F_{PRP}) for the period 2014-2019. Revision of F_{PRP} resulted in changes in the emissions ranging from -6.1 kt CO₂ eq (4.8%) in 2016 to 4.8 kt CO₂ eq (3.6%) in 2019.

3.D.1.4 Crop Residues

The AWMS data update for the period 2014-2019, also resulted in a change in N input from crop residues (F_{CR}) due to a revision of the straw used for bedding. Recalculation to the F_{CR} resulted in changes in the emissions ranging from -0.03 kt CO₂ eq in 2016 to 2.3 kt CO₂ eq in 2019.

The net effect of recalculations for 3.D.1 is a change in emissions ranging from -5.6 kt CO₂ eq in 2003 to 13.3 kt CO₂ eq in 2018.

Due to revisions in N inputs from the above-mentioned sources-categories 3.D Direct N₂O emissions from Agricultural Soils N₂O emissions decreased by 0.5-5.6 kt CO₂ eq between BY and 2009 and increased by 0.3-13.3 kt CO₂ eq over the period 2010-2019.

3.D.2 Indirect N₂O Emissions from Managed Soils, whole time series

The above-mentioned recalculations in 3.D.1 Direct N₂O Emissions from Managed Soils, which are not detailed again, have also led to changes in indirect N₂O emissions from agricultural soils due to atmospheric deposition and nitrogen leaching and run-off.

The recalculation of NH₃ emissions from 3.D also contributed to the changes in indirect N₂O emissions from agricultural soils due to atmospheric deposition. The main driver behind the decrease in the emissions in the period 2001-2019 is accounting for NH₃ abatement technologies in solid manure application. NH₃ emission abatement technologies for slurry application have already been accounted for in the previous submissions of the air pollutant as well as the GHG-emission inventory. While the manure NH₃ abatement technologies for solid manure application are reported for the first time in this year's submissions. In Hungary, measures on manure incorporation entered into force in 2001, so these technologies were taken into account in the estimation of NH₃ emissions from this date. The reduced NH₃ emissions resulted in a reduction of indirect N₂O emissions due to atmospheric deposition ranging from 0.09 kt CO₂ eq (0.6%) in 2002 to 15.2 kt CO₂ eq (8.7%) in 2019. In contrast, over the BY and 1990-2000 period, due to the decreased N-loss from the animal manure the emission increased, slightly. This increase is less than 2 kt CO₂ eq.

Changes in emissions from nitrogen leaching and run-off because of the aforementioned revisions are negligible (less than 0.3 kt CO₂ eq).

The net effect of these recalculations are changes in 3.D.2 Indirect N₂O emissions from agricultural soils ranging from -14.9 kt CO₂ eq in 2019 to 1.9 kt CO₂ eq in the BY.

The overall effect of recalculations for 3.D Agricultural Soils was a decrease ranging up to 8.7 kt CO₂ eq (0.3%) in 2012, except 2013, when emissions increased negligibly.

WASTE

5.B.2.b: AD (biogas produced) for 2019 have been revised. Biogas from manure management has been subtracted from biogas accounted for under 5.B.2.b as it is accounted for under the Agriculture sector.

5.D.1: Protein consumption data for 2019 have been revised. The amount of wastewater undergoing tertiary treatment has been revised for 2019.

5.F: The reported data have been corrected as the amount was erroneously given in kt C instead of kt CO₂.

The overall effect of all recalculations was below the threshold of significance.

11 KP-LULUCF

11.1 General information

According to relevant provisions, Parties to the Kyoto Protocol (KP) must submit information on land-use, land-use change and forestry (LULUCF) that is supplementary to what is contained in the report under the UNFCCC (i.e., Section 6). These provisions set principles to govern the treatment of LULUCF activities; require a consistent definition for terms such as “forest”, as well as definitions for activities under Article 3.3 and agreed activities under Article 3.4; and describe how modalities, rules and guidelines are implemented relating to the accounting of activities under Articles 3.3 and 3.4. Hungary follows both these provisions and the good practice guidance concerning the methodology for estimating GHG emissions and removals as provided in IPCC (2013).

Hungary started to report LULUCF-related information in 2006 in its Initial Report under Article 7, paragraph 4, of the Kyoto Protocol (http://unfccc.int/files/national_reports/application/pdf/hungaryareport_v4fin_c3.pdf) where, among others, Hungary reported the election of an activity under Art. 3.4, i.e., Forest Management (FM) for the first commitment period, and broadly defined both FM and “forest”. Hungary also published its initial report for the second commitment period (https://unfccc.int/files/national_reports/annex_i_ghg_inventories/national_inventories_submissions/application/zip/hun-2016-ir-15jun16.zip).

This part of the NIR provides supplementary information for the actual inventory year under the KP based on the above legal documents. *Information on forests not contained in this chapter, and/or reference from this Chapter, can be found in Chapter 6 of the NIR.*

Hungary only elected FM under Art. 3.4 for the first commitment period (it is obligatory to continue to report on FM in the second commitment period), and no other activity has been elected for the second commitment period. Therefore, this part of the NIR mainly covers issues related to the forestry sector. Information on other land-use related activities (e.g., cropland management) is limited to relevant information about land-use conversions.

The below table reports all comments from the most recent Annual Review Report (ARR; unedited draft at the time of the submission of this NIR) and how they have been addressed.

Table 21.1 Comments in ARR of the annual review in 2021 and how they have been addressed.

ID of comment in draft ARR	How the comment has been addressed
KL.1	Resolved.
KL.2	Resolved.
KL.3	We corrected and extended the table 6.5.1. We also provided detailed information about “found forest”, please see updated NIR section 6.5.2.
KL.4	Resolved.
KL.5	We extended the demonstration for soil and DOM pool in the NIR section 6.5.5.2.2. and 6.5.4.2.2., but as we write in the section 11.3.1.2, Hungary does not have and won’t have in the near future reliable monitoring for litter. Therefore, we must assume that like the deadwood and soil pools the litter pool is not a source.
KL.6	See updated NIR section 6.4.1 and 6.5.4.2.3.
KL.7	See above (by KL.5).
KL.8	See updated NIR section 11.5.2.3.
KL.9	We changed the notation key from “NA” to “NE” for the mineral soil pool in the CRF table 4(KP-I).B.1.

KL.10	We provided an extended version of Table 11.6.
KL.11	See updated NIR section 6.1.4.
KL.12	See updated NIR section 6.5.2.
KL.13	We provided the corrected Table 6.5.17.
KL.14	We changed the notation key from “NE” to “NO” in the CRF table 4(KP-II)3 under FM for carbon stock change resulting from N ₂ O emission.

11.1.1 Definition of forest and any other criteria

As reported in our both Initial Reports (i.e., 2006 and 2016), Hungary has elected elements and single minimum values for „forest” according to Table 11.1. We note that these elements have not been changed since, and are the same as for, the first commitment period.

Table 11.3 Definition of “forest” with prescribed characteristics and the justification of the chosen values.

Characteristics	Chosen value	Justification
Single minimum land area	0.5 ha	identical with value reported to FAO earlier
Single minimum width of forest area	10 m	defined by the methodology of current forest planning
A single minimum tree crown cover value	30%	identical with value reported to FAO earlier
A single minimum tree height value	5 meters	identical with value reported to FAO earlier

Concerning the **minimum size** of land area, it is the minimum size, by law, of forest stands. The mean size of stands in the country is around four ha. There are also patches of areas covered by trees of forest species in the country that are smaller than 0.5 ha, however, these patches are not surveyed currently.

Concerning **minimum width**, our forests are most often much wider than that, i.e., the chosen value occurs quite rarely. The width of 10 m typically allows for only 3-4 rows of trees.

Concerning **minimum crown cover**, the vast majority of the forests are on sites that allow for closed canopy closure already in young stands, and this closure is usually kept well above 50% until final harvest and regeneration. There are some stands in the country on sites where forests would not necessarily occur under natural conditions (and thus have low crown closure), however, the proper and intensive management of even these stands ensures that they would usually have more than 50% crown closure. None of these stands would be cultivated if the management of these stands were not profitable, which requires a relatively high crown closure.

The above also holds true for **minimum tree height**. It only happens on very few extreme sites that trees cannot reach a mean height of five meters at maturity.

In addition to managerial aspects, the above elected definitions match those applied in the forest inventory and monitoring: the definition was elected also to attain the highest possible accuracy in reporting. Moreover, the selected values are consistent with those reported to FAO and used in other international statistics.

The above elements of the definition of “forest” under the KP are exactly the same as those under the

UNFCCC. Note, however, that additional information is needed to define “forest” under the KP, e.g., *when* a certain piece of land becomes “forest” due to an afforestation or reforestation activity, and which areas are accounted for under FM. These additional pieces of information are detailed in the following sections.

With respect to origin, forests in Hungary are:

- planted, when the propagation material (seeds, cuttings or seedlings) is artificially put in the soil;
- semi-natural, when the propagation material comes from the mature, harvested trees of managed forests during an assisted natural regeneration, or when parts of these trees (roots and stumps) serve as sources of the regenerating shoots of a new generation of trees after harvest; and
- natural, when the entire regeneration process, which includes the production of propagation material of any sort, happens due to natural processes.

Based on the above, Hungary defines “**planted forests**” under the provisions of Decision 2.CMP/7 as forest plantations with artificially regenerated, short rotation (<40 years) species and long rotation species of coppice origin. All other forests, whether managed or not (such as forest reserves, see section 11.1.3.3 below), are considered “**natural forests**”.

11.1.2 Elected activities under Article 3, paragraph 4, of the Kyoto Protocol

As stated in both of our Initial Reports, as well as above, Hungary only elected FM under Article 3, paragraph 4 for the first commitment period, and no other activities under Art. 3.4 have been elected for the second commitment period.

11.1.3 Description of how the definitions of each activity under Article 3.3 and each elected activity under Article 3.4 have been implemented and applied consistently over time

Under the UNFCCC, emissions and removals from forests must be reported for “managed forests”. As reported in Chapter 6, all forests can be regarded as “managed”. Beginning 2016³, all forests in each new inventory year that are identified in the reporting year and are classified as “found forests” (FF) are included under FM.

The total area within the forestry sector reported under the KP is split among the following categories for each inventory year:

- AR: land under afforestation or reforestation since 1990
- D: land that has been deforested since 1990
- FM: all other forest land that was known to exist 31 December 1989 less D plus FF.

³ This decision is consistent with the suggestion of the ARR of 2013.

In the remaining parts of this section, first we define each activity, then provide details of the methodology of the estimation of emissions and removals. Note that for the development of carbon stock changes of most pools, most methodological information is discussed in Sections 6.4 and 6.5. Both the definitions and methods of estimation are consistently applied throughout the period 1990-2020.

The evolution of the area of AR, D and FF is demonstrated in Figure 11.1, whereas that of FM is demonstrated in Figure 11.3 below.




Figure 11.1. The evolution of the cumulative area of AR, FF and D since 1990.

11.1.3.1 Definition and identification of “AR since 1990”

In general, AR is an activity with the objective to establish “forest” as defined above on land that was not covered by such a “forest” before. The category “AR since 1990” includes all forest that has been established through direct human induced activity since 1990. AR land has so far only been deforested in Hungary in a few cases. As required by relevant provisions, AR can only include forest that can be demonstrated to have originated due to direct human induced activity. As a combined effect of these processes, in most years, the area of this category increases less than the L-FL category which includes increases of forest area due to both human-induced and natural causes.

In Hungary, afforestations are done in three steps. The first step is to do site preparation and, after that, to plant the propagation material in the area (initial planting). The second step lasts for a period of one to several years, depending on species and site fertility, when the newly established stand is tended and when beating-up is done if deemed necessary. Finally, the third and last step occurs when the afforestation is deemed “mature” by authorities. At this point, the stand is inspected, and, if it is found to have established itself and is expected to be able to survive, grow and develop to a fully mature forest, it is regarded as a “certified forest” (however, under the UNFCCC, it is only moved to the FL-FL category 20 years after the planting has taken place, see section 6). The whole process from site preparation to certification can last from one to 10-15 years, depending on species, site, weather and other factors, see **Table 11.7** below. In any event, an area becomes part of AR when the first step is completed.

Note that we began to identify areas of “AR since 1990” by considering the database of the above certificates. It was found during the analysis that some of the certified areas have not yet entered, or could not be identified in, the NFD (i.e., in the database of stands), which contains growing stock information, and which is used for the estimation of emissions and removals. This may mean e.g., that the afforestation proved to be unsuccessful. Therefore, we only included the smaller of the two sets in the “AR since 1990” category, i.e., the one for which we have data in the NFD, and for which a proof exists that the afforestation was indeed successful.

In relation to the KP, which sets a specific cut-off point (1 January 1990) in requesting countries to account for afforestations/reforestations, it is important to precisely define afforestations considering this cut-off point. In Hungary, in order to be conservative, “afforestations since 1990” are those, and only those, areas where both site preparation, as well as the planting of the propagation material started to happen after 1 January 1990. In a similar fashion, new AR areas by an inventory year are those, and only those, areas where both site preparation as well as the planting of the propagation material started to happen after 1 January of the inventory year. In general, site preparation and planting do occur in the same season anyway, shortly one after the other.

It is also important to define the cut-off point after which an afforestation counts as an area “subject to 3.4 FM”. Indeed, due to provisions of the Hungarian Forest Act, all afforestations become subject to FM right away as they enter the AR category. Thus, all forests under AR since 1990 are subject to 3.4 FM.

We also note here that the category “AR since 1990” includes the areas of stands that were actually afforested (i.e., the area of forest sub-compartments), but excludes adjacent roads or other areas that are not covered by trees, see section 11.2.2 below.

Finally, we note again that the statistically captured forest area keeps increasing at a rate that is higher than the area of land under AR. This is because not all increases are due to direct human induced activities, and that we find forests (i.e., FF) each year and some of these forests are not the result of direct human-induced activities.

11.1.3.2 Definition and identification of “D since 1990”

D areas are those that have been clear-cut and removed from areas under forest management in order that the area can be used for non-forestry purposes (i.e., for road building and other land-use).

An area enters the D category right away, i.e., in the year, of the clear-cut which is made in order that the area can be used for non-forestry purposes. In Hungary, deforestations have not been done frequently since 1990 nor were they done before that.

The *location* of D areas has only been registered since 2008, i.e., the beginning of the first commitment period under the KP, as it was of no importance for the forest inventory earlier, and the exact location of most deforestations prior to 1 January 2008 are thus not known. Because all deforestations have to be identified under the KP, we set up a system to identify at least the total area of deforestations from all available information even before 2008. This system allows for estimating and accounting for all emissions from deforestations. (Note that all possible emissions from biomass are accounted for on D land in the year of the deforestation, therefore, no such emissions can be expected on any of the land that was deforested before 2008, and no removals are accounted for, either, on this land in order to be conservative. Therefore, and because the annual rate of deforestations is higher after 2008 than before that year, we believe that the ignorance of the location of land deforested before 2008 presents no risk at all of underestimation of emissions from the biomass pools during the commitment period. In contrast, emissions from the other pools require information on the area of deforestations before 2008, but here

too, knowing the location is not required, only the post-conversion land-use category needs to be known, see later.)

The total area of deforestations was established based on statistical data collection back to 1990 using the certificates of the deforestations as all conversions from forest area must be approved by forestry authorities. For this authorization, the follow-up of the utilization of the deforested areas, but also to estimate emissions, ID-s, the type by utilization (within the “forest land” land-use category until the deforestation) and polygons of the deforested sub-compartments (whether forest or other sub-compartments), together with the date of deforestation are recorded in a specific database.

However, it was suspected that these certificates are fully available only since 2003. Therefore, a sample-based study was conducted that indeed showed that the total area of the deforestations before 2003 that could be retrieved from the National Forestry Database, which contains data of forest stands only, was higher than the one that could be developed from the hard copy files of the certificates. This means that in fact some certificates, thus, some deforestation areas could not be identified by only using these certificates. Therefore, the area established by the certificates before 2003 was multiplied by a factor of 1.18, which was established in the above study and was deemed representative for the whole country, to estimate the area of all deforestations before 2003.

Beginning 2020, as a follow-up of the review in that year, we started to separately report on areas of the so called “forest compartments” (i.e., areas with forest stands) and “other compartments” that usually do not, but sometimes can, contain trees. **Table 11.2** below reports the proportion of the “other” sub-compartments by utilization type (under forest management) between 2008 and 2020 for the D category, and for FM as a comparison. To report further details, we report Figure 11.2 below which is a map demonstrating examples for types of sub-compartments in four inserts with enlarged parts of the map. In these parts, we have colored the polygons that were deforested in 2020 by sub-compartment and type by utilization for the other sub-compartments to clearly distinguish the forest sub-compartments from the other sub-compartments. For the type of utilization of the other sub-compartments, see the legend.

Table 11.2. Share of area of other sub-compartments by utilization type between 2008-2020 for (a) FM and (b) D.

(a)

Utilization type on deforested "other sub-compartments"	Share of area on FM												
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Christmas tree plantation	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Area for twig production for decoration	1%	1%	1%	1%	1%	1%	1%	1%	0%	0%	0%	0%	0%
Forest research area	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Plant nursery	2%	1%	1%	1%	1%	1%	1%	1%	1%	0%	0%	0%	0%
Lane	19%	19%	18%	18%	18%	18%	18%	18%	18%	18%	19%	19%	19%
Glade	28%	28%	28%	28%	27%	27%	26%	26%	26%	26%	26%	26%	25%
Area serving game feeding	10%	10%	10%	10%	9%	9%	9%	9%	9%	9%	9%	9%	8%
Area covered by scrubs	9%	10%	11%	12%	12%	12%	13%	13%	13%	13%	14%	14%	14%
Not managed forest	0%	3%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Park	1%	1%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Infertile area	16%	15%	17%	17%	17%	17%	17%	17%	17%	18%	18%	19%	19%
Artificial water body	1%	1%	0%	0%	0%	0%	0%	0%	1%	1%	1%	1%	1%
Forestry office building	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%	1%
Private forest road	7%	7%	7%	7%	7%	7%	7%	8%	8%	8%	8%	8%	8%
Forest railway	0%	0%	0%	1%	1%	1%	1%	2%	2%	2%	1%	0%	0%
Timber yard	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Mine	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Lake or watercourse	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%	3%
Other facilities under forest management	3%	1%	1%	1%	1%	1%	1%	1%	1%	2%	1%	1%	1%
Forest ski resort	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Unknown	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%

(b)

Utilization type on deforested "other sub-compartments"	Share of area on D												
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Christmas tree plantation	0%	0%	0%	0%	0%	0%	1%	1%	0%	8%	0%	0%	0%
Area for twig production for decoration	0%	20%	0%	0%	0%	0%	0%	4%	0%	18%	0%	0%	0%
Forest research area	0%	0%	0%	5%	4%	1%	0%	2%	0%	0%	1%	0%	1%
Plant nursery	2%	15%	0%	1%	1%	2%	2%	0%	7%	41%	0%	0%	0%
Lane	7%	4%	4%	2%	4%	5%	8%	3%	2%	1%	0%	1%	2%
Glade	65%	10%	12%	19%	37%	23%	27%	38%	8%	4%	11%	22%	25%
Area serving game feeding	7%	15%	18%	29%	8%	20%	20%	26%	7%	2%	24%	6%	16%
Area covered by scrubs	7%	6%	3%	2%	12%	24%	20%	14%	12%	2%	3%	3%	29%
Not managed forest	0%	2%	1%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Park	0%	3%	32%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Infertile area	5%	5%	15%	35%	4%	9%	8%	4%	6%	1%	17%	1%	12%
Artificial water body	0%	0%	1%	0%	0%	2%	0%	0%	0%	0%	0%	0%	1%
Forestry office building	0%	1%	1%	0%	1%	0%	0%	2%	0%	0%	3%	0%	1%
Private forest road	0%	1%	0%	2%	0%	0%	1%	3%	1%	5%	0%	1%	4%
Forest railway	0%	0%	0%	0%	3%	5%	6%	0%	0%	16%	38%	58%	0%
Timber yard	0%	0%	0%	0%	0%	0%	0%	0%	1%	0%	0%	0%	0%
Mine	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Lake or watercourse	0%	0%	0%	1%	2%	1%	0%	0%	38%	0%	0%	1%	2%
Other facilities under forest management	0%	10%	8%	1%	3%	5%	3%	0%	15%	0%	2%	7%	8%
Forest ski resort	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
Unknown	7%	9%	5%	2%	21%	2%	4%	1%	2%	0%	0%	0%	0%




Figure 11.2. The spatial distribution of deforested land since 2008, with an enlarged portion directly showing deforested areas in a specific region for specific types of utilization of areas undergoing deforestation in 2020.

Based on the above, we have been able to report the complete time series of the estimated total area of D since 1990 (**Table 11.3**). Concerning activity data for the calculation of the various emissions and removals from deforestation, it must be considered that, in addition to emissions from the biomass and DOM pools which are assumed to occur in the year of deforestation, emissions from soils also take place. These emissions are estimated using the default transition period of 20 years. Therefore, three

types of areas within the D category have to be used. The first is the total area since 1990 which is now reported in the CRF tables using two sub-categories, i.e., the forest sub-compartments and all other sub-compartments (a total of 41,13 kha in 2018). The second is the annual area of all sub-compartments which was 3378 ha in 2018. The third area statistics is the area, which also includes forest and other sub-compartments, that is considered to be in transition. **Table 11.3** below reports statistics for all three area types by year. (Note that the third statistics is not necessarily equal to the area in transition in the FL-L category because for this category, unlike with D, the accumulation of annual areas in transition starts in 1985 rather than 1990.)

Table 11.3. Complete time series of the estimated area of the D since 1990 category for the first and second commitment periods under the Kyoto Protocol.

Inventory year	Deforestation since 1990				
	Cumulative area (ha)			Area in transition (forest + other subcompartments, ha)	
	forest sub-compartments	other sub-compartments	total	forest sub-compartment	other sub-compartment
2008	8 422	11 017	19 439	8 422	11 017
2009	8 872	12 057	20 929	8 872	12057.1
2010	9 080	14 200	23 280	8 467	14200
2011	9 356	15 528	24 884	8 503	13 951
2012	10 138	16 459	26 597	9 160	13 560
2013	10 670	17 173	27 843	9 363	14 274
2014	12 562	18 072	30 634	11 037	15 173
2015	13 945	18 389	32 333	12 061	15 490
2016	16 061	19 101	35 162	13 832	15 931
2017	17 772	21 266	39 037	15 021	18 096
2018	19 989	22 426	42 415	16 836	19 256
2019	22 296	22 750	45 046	18 748	18 529
2020	23 487	23 062	46 549	19 220	18 374

To estimate emissions and removals, an important step was to categorize all types of the other sub-compartments into land use categories by utilization type as if they belonged to one of the general land-use types (**Table 11.4**). This categorization makes it possible to estimate the amount of carbon in each pool before and after the conversion by the respective land-use types for the other sub-compartments.

Table 11.4. Reclassification of the utilization type on deforested other sub-compartments.

Utilization type on deforested "other sub-compartments"	LU category
Christmas tree plantation	FL
Area for twig production for decoration	CL
Forest research area	FL
Plant nursery	CL
Lane	GL
Glade	GL
Area serving game feeding	CL
Area covered by scrubs	GL
Not managed forest	FL
Park	FL
Infertile area	OL
Artificial water body	WL
Forestry office building	SE
Private forest road	SE
Forest railway	SE
Timber yard	SE
Mine	SE
Lake or watercourse	WL
Other facilities under forest management	SE
Forest ski resort	GL
Unknown	FL

Finally, the demonstration that regenerated areas under FM are not accounted for as D can be found in section 11.4.2 below.

11.1.3.3 Definition and identification of "FM since 1990"

The definition of "forest management" in Hungary is well described in the Forest Act. The relevant forest act that was mainly in effect for the period of 1990-2008 was passed by Parliament in 1996 (Act LIV of 1996 on Forests and the Protection of Forests, see at http://www.nfk.gov.hu/download.php?id_file=40588). Article 7 of this Act stated that "For the purposes of this Act, forest management shall be qualified as the entire range of activities aimed at maintaining, guarding and protecting forests, ensuring their public function, increasing forest assets, and exercising the forest usufructs in accordance with the provisions of Article 2." The relevant section of Article 2, in turn, reads: "Forests should be used and exploited in such a manner and at such a rate, which allows the prospects of management to endure also for future generations (hereinafter referred to as: sustainable forestry), so that the forests preserve their biological diversity, naturalness, fertility, ability to regenerate, viability, furthermore, that they satisfy the protective and economic needs in harmony with the requirements of society, and fill their role of serving the purposes of nature conservation and environmental protection, health and welfare, tourism, research and education." The most recent forest act was passed in 2009 (Act XXXVII of 2009 on Forests, Protection of Forests and Forest Management), which further reinforced provisions to protect forests and avoid deforestations and initiated a transition to close-to-nature forestry at an increased rate. (The text of the Act, currently in Hungarian can be found at http://net.jogtar.hu/jr/gen/hjegy_doc.cgi?docid=A0900037.TV.)

"Forest management" in general includes all kinds of activities in the forest from protecting forests to their multi-purpose utilization of a wide variety of social and ecological functions and services of the forests. All these activities often require that all forests are managed rather intensively, although the

intensity is quite different in the various stands depending on site, species, and the local objective of managing the stand. Managing forests involves preparing forest management plans, afforesting, intensive thinning, harvesting and regenerating as well as forest protection, maintenance of roads and road building, inspecting of forestry operations and others. The intensity of management is characterized by the length of the operational cycle of returning to each forest sub-compartment (of about four ha in average as mentioned above), which varies from about a few weeks (in afforested or regenerated areas where tending is necessary) to a year (in young poplar stands for tending) to five years (between pre-commercial thinnings in young stands of fast growing species) to maximum 15(-20) years (between thinnings in older stands of slow growing species). Forest management planning covers all forests, and forest management plans are made for 10(-12) years. That all forests (in the sense of the above “forest” definition) are managed in one way or another in Hungary is partly an economic and practical necessity because of the high rate of wood utilization, and because the density of the population, which requires all kinds of products and services from the forests, is quite high according to official statistics (105 capita km⁻², HCSO 2018).

We also note that there are practically no remnants of virgin forests, old growth forests or other primary forests in the country. There are some 70 so-called forest reserves in Hungary, whose total area amounts to some 12 kha. Forest operations in these reserves are limited to a so-called protection zone (altogether about 8 kha), which thus makes up most of the area of these reserves, and which surrounds the so-called core zone (altogether about 4 kha) where no traditional operation is conducted whatsoever. However, there is usually some activity even within these core areas such as protection by fencing, wildlife management, forest protection, research and education, and tourism. All protected forests are also included in the so called “Natura 2000” protection network of the European Union that involves various protection measures.

Activities that are carried out in all Hungarian forests also include preparing forest management plans, surveying/monitoring and inspecting stands regularly.

Because one or several of the above activities are carried out in each known stand each year, all forests in Hungary are regarded as “*managed since 1990*”.

The above also means that Hungary applies a *broad definition* of “Forest Management” under Art. 3.4 of the KP.

Land under the “FM since 1990” activity is identified by establishing FM in 31 December 1989 (which equaled the total FL at that point), and then subtracting D areas and adding FF areas in subsequent years, if any. It thus excludes D areas, but includes all land that, with the exception of AR, increased forest area (see also Chapter 6.2). In most cases, FF are young and are thus in their intensive growing phase. Note that as very little information was available on the origin of these forests, it was deemed to be impossible to demonstrate “direct human induced activity” in their establishment (see Chapter 6.5.2.1), therefore, we excluded these forests from FM in the first commitment period. However, as it seems improbable that these forests are unmanaged, and to comply with the requirements of ARR 2013, we now include all FF in our FM estimates.

11.1.3.4 Separating AR from FM

As stated above, as soon as site preparation and planting or seeding of propagation material is done, all AR lands become “forest” from the viewpoint of the definition of “forest” under the KP. From a domestic administrative point of view, when an AR land becomes a “forest” under the Hungarian regulations, it right away becomes an area subject to FM. Thus, since the category “AR since 1990” includes all areas that have been afforested since 1990, these areas could also be regarded as 3.4 FM. All of these areas are, however, excluded from FM areas to avoid double counting.

Full consistency with reporting under the UNFCCC is achieved by first establishing the area of AR and then developing FM as all forests (“FL” in the report under the UNFCCC) in 1 January 1990 minus all “AR since 1990” minus all “D since 1990” plus FF (see below). In this way, AR since 1990 that would otherwise classify as FM is automatically excluded from FM.

11.1.3.5 Separating D from FM

This issue is covered under section 11.4.2 below.

11.1.4 Description of precedence conditions and/or hierarchy among Article 3.4 activities, and how they have been consistently applied in determining how land was classified

As Hungary only elected FM under Article 3.4, no precedence or hierarchy issues arise.

11.2 Land-related information

The information below is supplementary to that reported in Section 6.2.

11.2.1 Spatial assessment unit used for determining the area of the units of land under Article 3.3

The spatial assessment unit in Hungary that is applied for the purposes of reporting under the KP is 1 ha. As a part of forest planning this is ensured by forest mapping that includes information of stands as small as 0.5 ha, i.e., areas that are smaller than 1.0 ha. Individual stands that are larger than 0.5 ha are also mapped at a spatial assessment unit of minimum 0.5 ha.

11.2.2 Methodology used to develop the land transition matrix

The land transition matrix is developed the following way:

- Areas under annual AR activities are identified on a per stand basis each year, and the area of these stands are summed up.
- Areas under D activity are identified since 1 Jan 2008 on a per stand basis each year, and the area of these stands is summed up.
- Both before and since 2008, all additional changes in the forest area were also identified that were not due to AR or D activities (i.e., FF).
- The total (known) forest area at the end of each year (since 1990) is identified on the basis of the NFD that includes appropriate records of each known stand in the country.
- By identifying the total forest area as well as all additions to, and reductions from, the forest area of the previous year, the constant elements (i.e., FM) can be identified. Land under FM was first identified at 31 December 1989. FM area has subsequently been reduced by the area of the deforested stands and increased by the area of FF.

As noted above, this procedure ensures the consistency of land identification under all KP activities, as well as FL under the UNFCCC. We identified all changes in the land-use statistics and classified them so that, eventually, all land can be accounted for in the respective categories since 1990. (See also section 6.2 above.)

To demonstrate that the land-use and land-use change information as reported under the UNFCCC is consistent with information under the various activities under the KP, below is a summary of the method of establishing the area of FM with the relevant data at the country level.

Note that, as discussed in Chapter 6.2 above, total area of “Forest land” as reported in the CRF table under the UNFCCC is more than the total area of all *stands* (the difference being forest roads and other areas (sub-compartments) not covered by trees). The reason for reporting total forest land area under the UNFCCC is that it is only possible to account for all land area of the country in the CRF tables under the UNFCCC if this area (i.e., the area of Forest Land) is to be consistently reported together with the area of all other land-uses so that the total of all these areas add up to the total land area of the country. Note that, for KP reporting purposes, we can only use and report, for AR, the total area of *stands*, or *sub-compartments*, which is included in the above “forest land” but excludes areas outside of the stands such as roads. (The area of stands also includes areas *within* the stands that are occasionally, and mostly only temporarily, not covered by trees, however, these are reported under the KP for all categories.)

We note that, in order to be conservative, emissions from all carbon pools are estimated for the total area of the FL-L category, i.e., the total of the area of stands plus the area not covered by trees.

The time series data of the total area all forests, along with that of the land that is strictly covered by trees ("calculated area covered by trees") is reported in Table 6.5.1 of the NIR. **Table 11.5** below summarizes changes of area under AR and D, whereas Figure 11.3 below is a draft graphical representation of all changes of the area of all mandatory and elected activities under the KP (using the area of forest sub-compartments). These changes represent actual changes (for AR, D and FM) due to the activities under Articles 3.3 and 3.4 of the KP, but also include those processes mentioned above that have resulted in the creation of the FF category.

Table 11.5 The size of annual land conversions for (a) D, (b) AR and (c) FF for the years of the first and second commitment periods.

(a) D

Inventory year	FL converted to Cropland	FL converted to Grassland	FL converted to Settlement	All conversions from FL to other land use
	Area (forest and other subcompartments, ha)			
2008	380	138	635	1 152
2009	184	336	970	1 490
2010	670	526	1 155	2 351
2011	388	140	1 075	1 604
2012	248	852	614	1 713
2013	270	274	702	1 246
2014	383	241	878	1 501
2015	521	413	766	1 699
2016	1 341	576	911	2 829
2017	2 367	878	630	3 875
2018	785	1 817	776	3 378
2019	934	687	1 010	2 631
2020	465	492	546	1 503

(b) AR

Inventory year	Cropland converted FL	Grassland converted to FL	Settlement converted to FL	All conversions to FL from other land use
	Area (forest subcompartments, ha)			
2008	6 674	389	157	7 221
2009	3 179	273	68	3 520
2010	5 434	322	511	6 267
2011	1 408	210	23	1 641
2012	1 010	134	17	1 162
2013	539	119	39	697
2014	82	36	13	130
2015	178	40	26	245
2016	229	13	0	242
2017	616	129	0	746
2018	1 098	243	0	1 341
2019	1 401	223	40	1 663
2020	2 068	315	48	2 431

(c) FF

Inventory year	Found Forests
	Area (forest+other subcompartments, ha)
2008	5 567
2009	6 487
2010	3 132
2011	4 230
2012	5 522
2013	4 370
2014	3 348
2015	841
2016	496
2017	1 664
2018	0
2019	0
2020	1 795




Figure 11.3. Graphical demonstration of changes of the area of the various activities under Articles 3.3 and 3.4 of the KP since 1990 (numbers after the activities are in kha). The area denoted by the dashed lines shows the development of the area over time identified by the NFD in each inventory year except for the AR area. For any given inventory year (i.e., at any vertical intersection of the graph) before 2020, the distance between the dashed lines shows the area of FM including that part of the FF that was identified up to that inventory year. Data under various activities are total areas of sub-compartments (they may be slightly different from respective numbers as reported elsewhere due to rounding-off, and so they do not represent official statistics). See text for other details.

Based on the definitions and the graph as outlined above, the areas of the sub-compartments under Article 3.3 and 3.4 activities are derived as shown by the formulas and data in **Table 11.6** (only rounded numbers are used for the entire area of the various activities; for precise numbers, and for data by geographical locations, see the KP CRF table).

Table 11.6. The evolution of the areas of forest sub-compartments (a) and other sub-compartments (b) under the relevant land-use categories under the KP, together with data for total forests and found forests for the first commitment period, as well as the algorithm (i.e., formulas, in the heading) of developing the data. The table shows all changes according to the formulas in which t1 means the beginning of the inventory year (i.e., the end of the preceding year), whereas t2 means the end of the year. The light-yellow color in some cells of the table (with column title “from DB”) shows that the data in those cells are taken from the database (i.e., they are the result of other compilations), whereas data in white cells are calculated in this table. All other notations are as in Tables 6.5.3. (The table is for demonstration only and may include rounding-off errors; for precise numbers, and for data by geographical locations, see the respective CRF tables.)

(a)

Inventory year	AREA (forest sub-compartments), ha													
	All Forest Land (FL)			D since 1990			FF since 1990			AR since 1990			FM since 1990	
	FL = FM + AR(cum.) + FF(cum.)													
	t1	t2	Δ	t1	Δ	t2	t1	Δ	t2	t1	new AR	t2	t1	t2
	from DB; t2 of prev. year	from DB	t2-t1	from DB; t2 of prev. year	from DB	t1 + Δ	from DB; t2 of prev. year	from DB	t1 + Δ	from DB; t2 of prev. year	from DB	t1 + Δ	FL89t2 - D + FF; t2 of prev. year	t1 - ΔD + ΔFF
2008	1 890 866	1 903 360	12 494	8 128	294	8 422	72 779	5 567	78 346	151 401	7 221	158 621	1 739 465	1 744 739
2009	1 903 360	1 912 917	9 557	8 422	450	8 872	78 346	6 487	84 833	158 621	3 520	162 141	1 744 739	1 750 776
2010	1 912 917	1 922 108	9 191	8 872	208	9 080	84 833	3 132	87 965	162 141	6 267	168 408	1 750 776	1 753 700
2011	1 922 108	1 927 702	5 594	9 080	276	9 356	87 965	4 229	92 194	168 408	1 641	170 049	1 753 700	1 757 653
2012	1 927 702	1 933 604	5 902	9 356	782	10 138	92 194	5 522	97 717	170 049	1 162	171 211	1 757 653	1 762 393
2013	1 933 604	1 938 139	4 535	10 138	532	10 670	97 717	4 370	102 087	171 211	697	171 908	1 762 393	1 766 231
2014	1 938 139	1 941 016	2 878	10 670	1 891	12 562	102 087	3 348	105 434	171 908	1 422	173 329	1 766 231	1 767 687
2015	1 941 016	1 940 720	-296	12 562	1 383	13 945	105 434	841	106 276	173 329	245	173 247	1 767 687	1 767 473
2016	1 940 720	1 939 342	-1 378	13 945	2 116	16 061	106 276	496	106 772	173 247	242	172 983	1 767 473	1 766 359
2017	1 939 342	1 940 052	710	16 061	1 711	17 772	106 772	1 664	108 436	172 983	756	173 502	1 766 359	1 766 550
2018	1 940 052	1 939 175	-877	17 772	2 218	19 989	108 436	0	108 436	173 502	1 341	174 558	1 766 550	1 764 617
2019	1 939 175	1 938 544	-631	19 989	2 307	22 296	108 436	0	108 436	174 558	1 676	175 916	1 764 617	1 762 628
2020	1 938 544	1 941 579	3 035	22 296	1 191	23 487	108 436	1 795	110 232	175 916	2 431	177 935	1 762 628	1 763 644

(b)

Inventory year	AREA (other sub-compartments), ha													
	All Forest Land (FL)			D since 1990			FF since 1990			AR since 1990			FM since 1990	
	L = FM + AR(cum.) + FF(cum.)													
	t1	t2	Δ	t1	Δ	t2	t1	Δ	t2	t1	new AR	t2	t1	t2
	from DB; t2 of prev. year	from DB	t2-t1	from DB; t2 of prev. year	from DB	t1 + Δ	from DB; t2 of prev. year	from DB	t1 + Δ	from DB; t2 of prev. year	from DB	t1 + Δ	FL89t2 - D + FF; t2 of prev. year	t1 - ΔD + ΔFF
2008	128 328	127 470	-858	10 159	858	11 017	21 613	0	21 613	0	0	0	128 328	127 470
2009	127 470	126 430	-1 040	11 017	1 040	12 057	21 613	0	21 613	0	0	0	127 470	126 430
2010	126 430	124 286	-2 143	12 057	2 143	14 200	21 613	0	21 613	0	0	0	126 430	124 286
2011	124 286	122 959	-1 327	14 200	1 328	15 528	21 613	1	21 614	0	0	0	124 286	122 959
2012	122 959	122 029	-931	15 528	931	16 459	21 614	0	21 614	0	0	0	122 959	122 029
2013	122 029	121 315	-714	16 459	714	17 173	21 614	0	21 614	0	0	0	122 029	121 315
2014	121 315	120 415	-899	17 173	899	18 072	21 614	0	21 614	0	0	0	121 315	120 415
2015	120 415	120 099	-316	18 072	316	18 389	21 614	0	21 614	0	0	0	120 415	120 099
2016	120 099	119 386	-712	18 389	712	19 101	21 614	0	21 614	0	0	0	120 099	119 386
2017	119 386	117 222	-2 165	19 101	2 165	21 266	21 614	0	21 614	0	0	0	119 386	117 222
2018	117 222	116 062	-1 160	21 266	1 160	22 426	21 614	0	21 614	0	0	0	117 222	116 062
2019	116 062	115 737	-324	22 426	324	22 750	21 614	0	21 614	0	0	0	116 062	115 737
2020	115 737	115 425	-312	22 750	312	23 062	21 614	0	21 614	0	0	0	115 737	115 425

The above calculation demonstrates that (1) all land is accounted for; (2) double counting is avoided; (3) all areas that are not in forest sub-compartments but are included in the “forestry area” (i.e., 2,054,281 - 1,938,544 = 115,737 ha in 2020, see also Table 6.5.1) are included in, and accounted for, under “Other” of the KP CRF table (Table NIR 2. LAND TRANSITION MATRIX).

11.2.3 Maps and/or database to identify the geographical locations, and the system of identification codes for the geographical locations

Hungary applies **Reporting Method 1** of IPCC (2013). This means that, in reporting area as well as emissions and removals, we identify regions for which we developed total areas under the various KP activities.

Two geographical locations are separated under the requirement of Annex II of 2/CMP.8 that the geographical location of the boundaries that encompass the lands subject to activities under Article 3.3 and FM under Article 3.4 must be reported. These locations are called North-Hungary and South-Hungary (see Figure 11.4 below). These geographical locations are separated along the borders of municipalities (which in turn follow partly other administrative, partly natural borders), and were found appropriate for the purposes of this reporting. (The boundaries are the same as in the first commitment period.) The identification codes used in the CRF tables are the following: North-Hungary, 1; South-Hungary, 2. „North” consist of the North Hungarian Mountains, the agglomeration of Budapest, the Trans-danubian Mountains (north to Lake Balaton) and the Little Hungarian Plain. The Great Hungarian Plains and the Trans-danubian Hills (South to Lake Balaton) belongs to „South”.




Figure 11.4. Map of Hungary with forests (green patches) and the border of the two geographical locations (red line).

For each year, all area (i.e., each stand) is allocated to one of the above geographical locations, thus, aggregate data (e.g., volume stocks, volume stock changes etc.) for these locations can be developed for each year. The identification system of sub-compartments is made up of three elements which are registered for every sub-compartment. These elements are: the municipality (village, or town) to which the sub-compartment is administered; the compartment (a larger piece of forest, e.g., a hillside or a valley) within the municipality; and the sub-compartment (which is part of a compartment). The sub-

compartment is the basic unit of forest management, its mean size being approximately four ha. The number of municipalities was 3166 in 1990 and 3195 in 2017, so the borders of the municipalities are considerably stable over time. (The borders of municipalities are declared and mapped by the Lechner Knowledge Center, Hungary, <https://lechnerkozpont.hu/en>). Since every sub-compartment exactly belongs to one and only one municipality, and municipalities are unambiguously mapped, and data for the geographical locations can be developed from the above stand level data by appropriately summing them up (see below).

11.3 Activity-specific information

11.3.1 Methods for the estimation of carbon stock changes and GHG emissions and removals

11.3.1.1 Description of the methodologies and the underlying assumptions used

11.3.1.1.1 Definition of pools as applied in Hungary

For all carbon pools, we apply the same definitions under the Kyoto as those under the UNFCCC. See Sections 6.4 and 6.5.3 for details.

11.3.1.1.2 Methodological issues

As AR and FM are different from L-FL and FL-FL, emissions and removals to be reported on land under AR and D are different from those in the respective categories under the UNFCCC. Therefore, these emissions and removals must be estimated using specific procedures. However, it is mainly the land to be accounted for that is different, and the methodology of the estimation is in general the same as that described in section 6.4 and 6.5. This methodology is accurate and precise as far as practicable.

In the case of Hungary, the methodology is pool and GHG-dependent. The coverage of emissions and removals estimation and its main methodological elements are detailed in Tables 11.7 and 11.8, respectively.

Table 11.7. Emissions and removals by source on land under (a) AR, (b) D and (c) FM.**(a)**

Inventory year	Emissions and Removals from FM since 1990, ktCO ₂					
	biomass	mineral soils	organic soils	litter	dead-wood	burning + wildfires (CH ₄ +N ₂ O)
2008	-3 053	demonstrated that not a source	62	demonstrated that not a source	demonstrated that not a source	11
2009	-2 082		62			10
2010	-1 933		62			11
2011	-1 780		62			26
2012	-2 738		62			22
2013	-1 878		62			15
2014	-3 042		62			18
2015	-3 812		62			27
2016	-3 011		62			13
2017	-3 313		62			24
2018	-2 891		62			13
2019	-3 690		62			20
2020	-5 055		62			14

(b)

Inventory year	Emissions and Removals from AR since 1990, ktCO ₂ eq						
	post-conversion biomass	pre-conversion biomass	mineral soils	organic soils	litter	dead-wood	burning + wildfires (CH ₄ +N ₂ O)
2008	-1 146	123	demonstrated that not a source	NO	demonstrated that not a source	-42	0.240
2009	-1 135	63		NO		-43	0.219
2010	-1 220	100		NO		-45	0.381
2011	-1 187	31		NO		-45	1.353
2012	-1 100	24		NO		-46	0.341
2013	-1 246	16		NO		-46	0.627
2014	-1 120	23		NO		-45	1.422
2015	-1 193	4		NO		-46	1.556
2016	-1 192	4		NO		-46	1.168
2017	-1 287	12		NO		-46	2.106
2018	-1 232	23		NO		-47	1.557
2019	-1 109	28		NO		-47	2.337
2020	-1 172	42		NO		-48	2.074

(c)

Inventory year	Emissions and Removals from D since 1990, ktCO ₂ eq					
	biomass	mineral soils	organic soils	litter	dead-wood	burning + wildfires (CH ₄ +N ₂ O)
2008	27	31	NO	9	3	0.042
2009	58	34	NO	14	5	0.097
2010	28	39	NO	7	2	0.043
2011	46	37	NO	9	3	0.095
2012	132	36	NO	25	9	0.229
2013	62	84	NO	17	6	0.132
2014	85	89	NO	61	22	0.178
2015	117	93	NO	45	17	0.170
2016	151	99	NO	68	26	0.264
2017	168	111	NO	55	21	0.294
2018	217	117	NO	71	27	0.362
2019	228	120	NO	74	29	0.296
2020	134	121	NO	38	15	0.208

Table 11.8. Methodological summary for (a) FM, (b) AR, (c) D. (CS=country specific; D: default; EJ: expert judgment; IE: included elsewhere; AD: activity data; EF: emission/removal factor; NO: not occurring)

(a)

Category	Type of information	Carbon stock changes					Non-CO ₂ emissions
		AGB	BGB	DW	LI	SOIL	
FM	E/R	CS	D/EJ	Not estimated (demonstrated that not a source)	Not estimated (demonstrated that not a source)	Mineral: Not estimated (demonstrated that not a source);	N ₂ O (N fertilization): IE
						Organic: AD: CS;	N ₂ O (drainage and re-wetting): IE
						EF: D	C (liming): not occurring
							Burning: D, CS
	Uncertainty	Tier 2 (Monte Carlo)		N/A			Tier 2 (Monte Carlo, where applicable)

(b)

Category	Type of information	Carbon stock changes					Non-CO ₂ emissions
		AGB	BGB	DW	LI	SOIL	
AR	E/R	Post-conversion: CS	D/EJ	CS	CS	Mineral: Not estimated (demonstrated that not a source);	N ₂ O (fertilization): IE Drainage and re-wetting: NO
		Pre-conversion: CS	D			Organic: not occurring	C (liming): not occurring
							Burning: D, CS
	Uncertainty	Tier 2 (Monte Carlo)		N/A			Tier 2 (Monte Carlo) where applicable

Table 11.8 (ctd.). Methodological summary for (a) FM, (b) AR, (c) D. (CS=country specific; D: default; EJ: expert judgment; IE: included elsewhere; AD: activity data; EF: emission/removal factor)

(c)

Category	Type of information	Carbon stock changes					Non-CO ₂ emissions
		AGB	BGB	DW	LI	SOIL	
D	E/R	Post-conversion: 0	0	CS	CS	D	N ₂ O (disturbance):
		Pre-conversion: CS	CS				mineral soils: D;
							Organic soils: NO
	Uncertainty						C (liming): D
							Burning: D, CS
		Tier 2 (Monte Carlo)					Tier 2 (Monte Carlo) where applicable

Biomass

Carbon stock changes of trees are estimated using the stock change method (in a fashion similar to categories under the UNFCCC), which automatically ensures that all processes, i.e., all changes due to gains, i.e., growth, and all changes due to losses, i.e., harvest, natural disturbances like fires etc., are taken into account. The estimation of emissions and removals on lands under the AR and D activities are directly estimated from the carbon stocks of consecutive calendar years.

The forests included in the AR category are identified and mapped at the sub-compartment (stand) level. Growing stocks and stock changes in the afforested areas are estimated by using field measurements and applying yield tables by appropriate species and site classes. These yield tables (which are true yield tables and different from the volume stock functions applied for the L-FL category) are planned to be updated once information is available that the growing conditions may have deteriorated. (We note here that, according to Somogyi, 2008, the growth of trees has so far accelerated in Hungary. Not adjusting the yield tables for this acceleration is thus equivalent to underestimating the removals, thus, it is conservative.)

The parameters of the equation used for the estimation are as detailed in section 6.5.3. In lack of country-specific measurements, the same root-to-shoot value of 0.25 is assumed for stands of land under AR (i.e., for young forests) as for all other forests. This can be regarded as rather conservative because young trees usually have higher root-to-shoot ratios than mature trees. As forests in the AR category are net sinks, this assumption leads to an underestimation of removals on AR land.

It must also be noted here that the NFD is designed to provide information on the actual *situation* (i.e., stocks) of stands *in each year* (see Chapters 6.5.1.1 and 6.5.2). However, the borders of the stands change annually mainly due to forest planning (e.g., in order to better comply with site patterns or management purposes). Although it is possible to track changes in position of border lines of sub-compartment polygons, stock changes cannot be calculated on sub-compartment level since sub-compartments may be divided into smaller parts or they may be merged year-by-year because of forest planning. Due to the fact that growing stock volume data are stored on sub-compartment level such changes in sub-compartment polygons cannot be tracked by stock volume data (i.e., stock volume data *within* a sub-compartment is not spatially explicit). Thus, it is not possible to keep track stock *changes* at the stand level, rather, only at higher administrative units (including the geographical locations). This means that the carbon stock *changes* cannot typically be estimated bottom-up from the stand level, rather, they are calculated from carbon stocks of consecutive years at aggregate levels (i.e., species and species groups), and thus estimated bottom-up from the stand level for categories of AR and D. The same applies to all forests, for which of course the estimation of carbon stocks is split for the two geographical locations.

Consistent with section 6.3, the emissions and removals for lands under FM are indirectly estimated from those of all forest land (FL-FL + L-FL) as well as AR and D. (Emissions and removals from FL-

FL and L-FL in an inventory year exclude carbon stocks of FF found in that inventory year.) This procedure is applied, among others, in order that the estimates under the UNFCCC and under the KP are consistent, that carbon stock changes are neither underestimated nor overestimated, and that double counting is avoided. With this approach, total net removals (NR, i.e., net gains) for FM are calculated using NR of FL under the UNFCCC (FL-FL plus L-FL, which includes NR of FF, but excludes NE (net emissions) of deforested land) and NR on land under AR (Table 11.9).

$$\begin{aligned} \text{Total NR of forests under FM in 2020} = & \\ & + \text{Total NR of FL-FL in 2020} \\ & + \text{Total NR of L-FL in 2020} \\ & - \text{NR of AR in 2020} \end{aligned}$$

Table 11.9 The development of emissions and removals in FM land, together with the algorithm (i.e., formulas) of the calculations, since 2008. The light-yellow color in the table shows that the data is taken from the database (i.e., it is the result of other calculations), whereas data in white cells are calculated in this table. All other notations are as in Tables 6.5.3 and 7.3.7. (The table is for demonstration only and may include rounding; for precise numbers, and for data by geographical locations, see the respective CRF tables.)

Inventory year	ΔC of biomass UNDER THE KP, converted to emissions in ktCO₂					
	FL (=gross ΔFL - new FF stock - D)		AR since 1990		FM since 1990	
	NR	IEF	Δ	IEF	Δ	IEF
	from DB	NR/area (Gg CO ₂ /ha)	from DB	NR/area (Gg CO ₂ /ha)	FL - AR - FF C stock found in year	NR/area (Gg CO ₂ /ha)
2008	-4 199	-0.00221	-1 146	-0.00723	-3 053	-0.00175
2009	-3 217	-0.00168	-1 135	-0.00700	-2 082	-0.00119
2010	-3 153	-0.00164	-1 220	-0.00724	-1 933	-0.00110
2011	-2 967	-0.00154	-1 187	-0.00698	-1 780	-0.00101
2012	-3 838	-0.00198	-1 100	-0.00642	-2 738	-0.00155
2013	-3 124	-0.00161	-1 246	-0.00725	-1 878	-0.00106
2014	-4 161	-0.00214	-1 120	-0.00646	-3 042	-0.00172
2015	-5 020	-0.00259	-1 193	-0.00688	-3 812	-0.00216
2016	-4 224	-0.00218	-1 192	-0.00689	-3 011	-0.00170
2017	-4 615	-0.00238	-1 287	-0.00742	-3 313	-0.00188
2018	-4 135	-0.00213	-1 232	-0.00706	-2 891	-0.00164
2019	-4 817	-0.00248	-1 109	-0.00631	-3 690	-0.00209
2020	-6 243	-0.00322	-1 172	-0.00659	-5 055	-0.00287

For AR, we have also developed a methodology to account for emissions from pre-conversion biomass losses due to afforestation. This is necessary as some of the afforestations take place in former orchards and vineyards. (The majority of the AR area is nevertheless done in former cropland with annual crops and grasslands with no woody vegetation.) To estimate these emissions, the country-specific loss rates of 9.4 and 18.8 tdm/ha are used for orchards and vineyards, respectively, consistent with data used under the UNFCCC (see section 6.6.2.1.1 on accounting for gains in carbon stocks of perennial crops on croplands). This loss, which is a mean value for all types of orchards and vineyards converted to forest, is assumed to be accumulated in 30 and 31.8 years, respectively, so the mean annual accumulation rates are 0.3 and 0.59 tdm/ha*yr.

For AR, we also estimated the loss of carbon from the pre-conversion biomass of other vegetation (predominantly the remains of annual crops, and herb vegetation on abandoned croplands and grasslands). The estimation was done the same way as described in the sub-section Pre-conversion biomass of Section 6.5.5.2.1.

Dead organic matter

For the deadwood and litter pools, too, the same approach was taken for the categories under the KP as for categories under the UNFCCC. See methodological details in section 6.5.6.2.2 for D. For FM and AR, the option is applied that it is demonstrated that the dead organic pools in these categories are not a source, see section 11.3.1.2 below.

Soil

For soils, the approach described in detail in section 6.5.4.2.3 was taken for D, however, only emissions were accounted for, whereas removals were not. For FM and AR, the option is applied that it is demonstrated that the dead organic pools in these categories are not a source, see section 11.3.1.2 below.

Non-CO₂ emissions

Non-CO₂ emissions are estimated based on harvests statistics, and experience that almost all natural forest fires occur on FM land, and only very few on AR land. The methodology is the same as described in the various sections of Chapter 6. The resulting data are reported in Table 11.7.

We note here, too, that, as a follow-up of the review of our report in 2017, we added the estimates of indirect N₂O emissions from leaching/runoff on both AR and FM land, using the same methodology that is described in section 6.4.2.

11.3.1.2 Justification when omitting any carbon pool or GHG emissions/removals from activities under Article 3.3 and elected activities under Article 3.4

For FM and AR, Hungary does not explicitly quantify emissions and removals for three forest carbon pools, i.e., soil, deadwood and litter, but demonstrates that these pools are not a source. To demonstrate that soils are not a source, a conservative approach is taken based on the IPCC 2006 GL methodology using country-specific and other data. The demonstration for DW and LI is based on expert judgment which is a practicable method in our situation (see below).

Demonstration for FM and AR that the soil carbon pool is not a source

This demonstration, which is separately done for AR and FM land, is necessary because, until this point, there has not been any forest soil carbon monitoring program in Hungary. The below demonstration involves all available country-specific data and information, i.e., Tier 2 elements. This data notwithstanding, we continue to apply the conservativeness approach used before, i.e., we always apply the information from various options, when there is any, that leads to higher emission estimates and lower removal estimates. Overall, the data suggests that the demonstration can be done with a high certainty.

(Note that, under the UNFCCC, it was not possible to estimate soil carbon stock changes for FL-FL, but it was possible for the L-FL category using a Tier 1 method. However, this estimation was regarded as not accurate enough to develop an acceptable carbon stock change estimate under the KP. Nevertheless, the estimates for the L-FL category will be cited below to further support the demonstration.)

A major research project was run 2009-2011 with the aim to develop more country-specific data for the demonstration, and all information from that project is used to support the demonstration.

The results of the project were published in a peer-reviewed research journal (Somogyi et al., 2013), therefore, only a summary of the most important arguments is presented.

The demonstration is based on an approach that the forest area is stratified into strata of rather different emissions or removals so that both the area and specific emission or removal factors of the strata, which mainly depend on the types of forestry operations conducted in the strata, can be identified. Stratification is used to most efficiently use information and data that is available in the country, including forestry statistics that are developed each year. The strata that are defined in this demonstration are based on relevant KP activities and available country-specific data.

The strata applied are the following:

for AR:

- areas where afforestations and reforestations occurred since 1990 on cropland, and
- areas where afforestations and reforestations occurred since 1990 on grassland,

for FM since 1990:

- land where final cutting and artificial regeneration following professional standards occur,
- land where final cutting and natural regeneration following professional standards occur,
- land where no final cutting occurs, only thinnings and other operations that cause no disturbance to the soil.

The area of each above stratum is known each year from the national forestry database. The area of AR on cropland is calculated using the relative amount of land that was cropland and that was grassland prior to the afforestation. The data shows a high share of cropland as a predominant land-use before afforestations with a mean value of 86.4% in the period 2008-2020. The mean share of grassland converted to forest land in the period 2008-2020 is 9.8%.

For FM land, there are specific statistics available for the above first two FM strata, from which the area of the third stratum is deducted from the total FM area.

Concerning the area-specific emission and removal factors, field measurements (in the above-mentioned project and an earlier one), modelling, literature review, expert judgment and reasoning are applied.

For the AR land since 1990 that was converted from cropland and grassland, equations from local case studies (Horváth, 2006, Somogyi, 2005 and Somogyi et al. 2013) were earlier used for demonstration. These implied an area-specific carbon stock increase of $0.555 \text{ tCyr}^{-1}\text{ha}^{-1}$ in 20 years for CL-FL and an area-specific emission rate of $0.13 \text{ tCyr}^{-1}\text{ha}^{-1}$ for GL-FL for the default period of 20 years that is used for mineral soils for the estimation of carbon stock changes in land-use conversions. When country-specific carbon stock change values are used (see section 6.4.1 and Table 6.4.2), the area-specific carbon stock increase is the same as for the non-SA CL conversion and for the GL-FL conversion, but only 0.075 for the non-SA CL-FL conversion. Since these latter conversions also occur, to be consistent with the IPCC default method, and to simplify the demonstration, we use the IPCC method (section 6.4.1) with this latter country-specific data for the demonstration. This method makes the resulting estimate for AR, which is still a net removal value, even more conservative than before (see also Table 11.11 below).

Concerning land under FM since 1990 where final cutting and artificial regeneration following professional standards occur, artificial regeneration means that a stand is replaced by a new one by applying operations that closely resemble those of conversions. These operations may include disturbances associated with final cutting and skidding of timber, soil preparation, erosion (on steep slopes), and planting or seeding. The amount of loss may depend on tree species, site and the technologies applied.

Until 2011, we used an area-specific emission value of 6 tCha⁻¹ for the specific carbon loss for this stratum. It was assumed that all emissions due to disturbing soils take place in the year of the start of the regeneration, i.e., the above specific value is applied to the total area of the harvested forests in the inventory year.

To check the plausibility of the above emission value, we compared it to the respective IPCC default factor. According to the 2006 IPCC GL, if a forest land is converted to a full-till cropland without additional input of organic carbon (when forests are regenerated, no additional organic carbon input is applied), it loses some 18% of the original (i.e., reference) carbon stock, for which we assume the mean value (Table 11.10) that results from classification of the area by climate type and soil type, and from applying IPCC default soil carbon stock values (IPCC, 2006, see section 6.4.1 for details). Based on the above default loss rate and reference carbon stock, the overall loss in a conversion is equal to $48.09 \times 0.18 = 8.6 \text{ tCha}^{-1}$, thus, not much higher than the above value of 6 tCha⁻¹, but smaller than the Tier 2 estimate of forest land – non-set-aside cropland conversions for 20 years, which (according to Table 6.4.2 (b)) is 11.1 tCyr⁻¹ha⁻¹.

Table 11.10. *The distribution and carbon stock of forest soils in Hungary by climate and soil types (for details, see section 6.4.1).*

Soil characteristics	WD HAC	CD HAC	WD sandy	CD sandy	Total
Distribution of area (%)	35.7%	53.6%	0.9%	9.8%	100.0%
SOC _{ref} (tC/ha)	58	48	21	15	48.09

Beginning the reporting year of 2011, we started to use a different specific carbon loss for this stratum which we consider more appropriate for the Hungarian conditions, and which is based on the recent project by Somogyi et al. (2011, 2013). In this project, several case studies were conducted to estimate the potential area-specific emissions. To model these emissions, the carbon stocks of paired stands before and after regeneration (1-15 years of age) were compared, and differences were regarded as carbon stock changes. In stands of slow growing species, sessile oak was used again, whereas intensively growing poplars were used to represent fast growing species. According to results, there are indeed areas where carbon stocks decrease after afforestation, with a rather high variability.

There are, however, several other factors to consider, too. One is that carbon stocks also increase due to the transfer of carbon from the dead roots of trees of the mature stand, which were harvested before the regeneration, to the soil pool. In a mature stand, it is not uncommon to have 320 m³ of above-ground wood volume (this value was only chosen for the sake of demonstration). If basic wood density is 0.5 tm⁻³ (a good approximation of national average), then the above-ground biomass is 160 tha⁻¹, which translates to a carbon stock of 80 tCha⁻¹. After applying a root-to-shoot ratio of 0.25 (that we consistently apply for the belowground biomass pool, see above), we get a carbon stock of 20 tCha⁻¹ in the roots of the mature trees. Because the root-to-shoot ratio is a conservative one, this estimate is again a rather conservative estimate, but it must also be considered that some of this carbon can be found in the coarse roots and stump. Most of the carbon that is transferred from the roots to the soil is found in the topsoil layers, and the transfer takes place a few years after felling the trees, i.e., after the death of the roots. The full decomposition of most of this dead-wood-turned-soil-carbon may take decades.

Note that the emissions from dead roots due to decomposition are fully accounted for in the biomass pool as both gains due to increments and losses due to harvests and mortality are taken into account when estimating carbon stock changes of the biomass pool by using the stock change method.

That carbon stocks of the soil do not decline much, rather, increase after regenerations, show that emissions from soils due to direct human induced disturbances from soil preparation are more than offset

by the transfer of carbon from the dead roots to the soil. However, this transfer could not be measured separately from emissions which thus remain rather uncertain.

It must also be highlighted that most forest soils in Hungary, just like those elsewhere, are deeper than the 30 cm for which the estimation / demonstration of carbon stock changes must be done according to the IPCC methodology. The set depth of 30 cm is rather artificial and has nothing to do with soil processes. Also, soil preparation may mix up various soil layers, which may also result in an increase of soil carbon in plains where soil layers of relatively large organic content are covered by layers of lower soil organic carbon content (discussed to some extent by Somogyi et al., 2013). Therefore, estimates and the demonstration for the 30 cm layer may not have to do anything with actual processes.

Considering all the above, combined with the specific soil conditions of Hungary, we reduced the mean area-specific emissions that is applicable for conversions for this demonstration. To stay conservative, however, we keep this rate at a still highly conservative value of **5 tCha⁻¹** until further evidence.

For land under FM since 1990 where harvesting and natural regeneration is made following professional standards, natural regeneration means that the area is regenerated exclusively through the propagation material that is locally produced by the trees of the mature stand. With a few exceptions, when seeds or seedlings from elsewhere are planted under the mature stand, and which sometimes involves some, but not intensive soil preparation, this type of regeneration usually makes it unnecessary to do any soil preparation, thus, only some small amounts of carbon may be lost due to inevitable damages caused by removing timber from the area. However, this loss, if any, is assumed to be quickly offset by the growth of the dense new generation of trees, if not offset right away by inputs from deadwood (mostly dead branches of harvested trees) and dead roots (of the same harvested trees) originating from the harvest of the mature stand.

Because of the above, this stratum is assumed to have no overall emissions, i.e., a specific carbon loss of **0 tCha⁻¹**.

Here we again present one specific result from the above-mentioned research project (Somogyi, 2013, Figure 11.5). We measured carbon stock changes of soils in several stands of Black locust, which is the most widespread tree species in Hungary. It seems that in stands where artificial regeneration took place, carbon stocks declined, however, regenerating the stands from roots, which is considered one form of natural regeneration in Hungary, resulted in both increase (in two case studies) and decrease (in one case study), or at least a much higher C stock than in the comparable stand after artificial regeneration (in the fourth case study). Thus, the above assumption is supported by some evidence.




Figure 11.5. Soil carbon stock before (“reference”) and after regeneration (artificial regeneration: “artificial reg.”, and regeneration from roots: “r. from roots”) of chronosequences of Black locust stands in four case studies. (Somogyi et al., 2013).

Finally, the stratum of **all other land under FM since 1990** includes stands that are between regeneration and the beginning of the subsequent regeneration and final cutting and that may only be affected by normal silvicultural operations such as thinnings. This stratum is by far the biggest one by area, and it includes all forests that cannot be classified into any of the previous categories. In these forests, the predominant process is the slow but steady growth of trees together with the associated slow but steady sequestration of carbon in the soil. These stands may occasionally and locally be disturbed by abiotic or biotic natural agents, or by thinnings, and some carbon may thus additionally be lost due to natural decomposition of dead biomass. However, these disturbances generally only affect trees but not the soil, the roots of the harvested trees slowly decompose and some of their parts become part of the soil, and the overall balance of all these processes is a net gain.

Therefore, these areas will be assumed to have a rather small but positive net carbon stock change per unit area. The value assumed until 2020, based on a literature review (see below and Somogyi et al., 2013), was a net removal of 0.05 tCha^{-1} . In 2020, however, we have modified the removal factor to represent the “central” estimate of this factor for the Hungarian conditions (instead of representing a conservative value). This figure, which is 0.357 tC/ha*yr with a half-width CI of 0.013 tC/ha*yr , comes from a not-yet-published analysis of soil C stock estimates from 186 plots from our TIM soil monitoring system (see section 6.4.1) between 1992 and 2016 that allow for the calculation of differences of C stocks in two consecutive surveys. Based on the revised figure, which is a preliminary one and of course has uncertainty, but which is nevertheless a kind of “central” estimate, the sink side of the balance is much higher than the conservative one that we reported in our NIRs earlier.

The summary of the data for AR and FM, and all of the above strata for 2020 are found in **Table 11.11**. The data in the table should only be regarded as values whose only role is to establish the sign of the net results for the demonstration. In other words, the estimated values, including the total carbon stock change value, are not regarded as accurate, and are not intended to be the basis for accounting, rather, they are only intended to serve the demonstration of the correctness of the assumption that soils are not a source.

Table 11.11. The area, emission and removal data for the various AR and FM strata and for their total in 2020. For the calculations for years preceding this inventory year, see our previous NIRs. See text for other details.

Forest Land Stratum under the KP		Estimated area	Emission (+) and removal (-) factor (IEF in <i>italics</i>)	Total emissions (+) or removals (-)
		(kha)	(tC ha ⁻¹)	(ktC)
Land under AR since 1990 (that is still in the transition phase from pre-conversion land use to forest land use)	that was converted from cropland	$104.6 \cdot 0.831 = 86.9$	estimated using the same stock change method that is applied to mineral soils for all conversion categories, i.e. according to section 6.4.1 of the NIR	-16.2
	that was converted from grassland	$104.6 \cdot 0.169 = 17.7$		1.9
	Total	104.6	-0.14	-14.4
Land under FM since 1990	where final cutting and artificial regeneration is made following professional standards	14.1	5	70.4
	where harvesting and natural regeneration is made following professional standards	3.4	0	0
	that are between regeneration and the beginning of the subsequent regeneration and final cutting, and that may be affected by normal silvicultural operations such as thinnings	$1761.8 - 14.1 - 3.4 = 1744.3$	-0.181	-315.7
	Total	1761.8	-0.139	-245.3

The result of the calculations for the current inventory year is a sink for both AR and FM for both 2020 and previous years (**Table 11.12**), so it can safely be stated that, overall, the mineral soils of forests of the FM land have not been a source for all years of the entire first and second commitment periods. For reasons of transparency, **Table 11.12** below reports not only the time series for AR and FM estimated using the above methodology but also that for L-FL using the Tier 1 methodology and the estimated areas of non-SA CL-FL and SA CL-FL conversions, the sum of which is larger than the area of AR.

Table 11.12. *The time series of net removals of soils for AR and FM using the methodology in the demonstration that AR and FM soils are not a source, and for L-FL using the Tier 1 methodology under the UNFCCC.*

Inventory year	Total emissions (+) or removals (-) (ktC)		
	AR	FM	L-FL
2008	-20	-235	-32
2009	-21	-235	-31
2010	-21	-227	-32
2011	-21	-218	-31
2012	-20	-224	-31
2013	-20	-226	-30
2014	-20	-234	-29
2015	-19	-236	-28
2016	-18	-236	-26
2017	-18	-233	-24
2018	-17	-228	-22
2019	-15	-233	-20
2020	-14	-245	-18

To further support the confidence in the above derivation for FM, i.e., why the above reasoning leads to highly conservative estimates, and therefore, highly certain conclusions that soils are not a source in lands under all KP activities, we note the following additional arguments:

- Concerning the value applied for artificially regenerated FM land, the assumed value of 5 tCha⁻¹ for the emissions in this land is the absolute maximum that one could assume based on the idea of completely converting forest to any another land-use. However, even if regenerating (including tilling once) may mean high disturbance, no till certainly occurs continuously after the regeneration is done, which means that repeated emissions of ploughing do not occur in forests, thus, total carbon stock losses must be much smaller in forest land remaining forest land than converting a forest land to cropland. Also, there are many types of artificial regeneration applied, including ones that do not involve any types of high-disturbance operations like ploughing. Currently, however, no statistics exist with respect to the share of the various regeneration types. According to experience, the operations leading to high emissions have been continuously replaced by less intensive ones (even due to economic reasons). The selected area-specific emission estimate of 5 tCha⁻¹ is with high probability a rather high overestimation, and it is applied for the sake of the demonstration only.
- It is documented in many scientific publications that forests accumulate C in their soil. We selected a rather comprehensive study published recently by Berg et al. (2007) that states that “The amount of carbon sequestered in humus increases in forests and it appears that the average rate for Sweden is of the magnitude 100 to 200 kg C ha⁻¹yr⁻¹.” (Note that this accumulation occurred in the humus layer of podsol soils, the depth of which never reached 12 cm.) Hungary is situated in a warmer region and has definitely higher tree growth rates, which involve higher ecosystem turnovers. Therefore, the assumed sequestration rate of 181 kg C ha⁻¹ yr⁻¹ is highly plausible.
- We highlight the fact that carbon stock change estimates are rather uncertain for both Hungary and any other country. This means that the uncertainty range of the above estimates, which cannot be quantified at the moment, is in our view so wide that makes no accounting reasonable. This is one major reason we opted for the demonstration. Our demonstration is, however, heavily biased on the emission side, thus increasing the confidence in the final conclusion that soils are not a source.
- Finally, we note that, although we do not use either our Tier 1 or Tier 2 estimates for accounting, the method of the above demonstration is capable of serving one important aim, which is the final goal of preparing greenhouse gas inventories, i.e., to identify sources of emissions due to direct human induced activity in order that the impacts of these activities can be reduced. By having

identified such processes broadly in our demonstration, we are now able to develop policies to reduce the emissions mentioned above.

In summary, by applying a method whereby all steps included conservative or even highly conservative estimates, we can conclude that the sum of all emissions and removals is negative emissions for both AR and FM, i.e., we demonstrated that the Hungarian forests are not a source. By applying conservative values, and demonstrating how and why they are conservative ones, leads to a high level of confidence in the conclusion.

Finally, once again, all the above is only meant to demonstrate that the Hungarian forests are not a source. The final result of the reasoning is not meant to be interpreted as an accurate scientific estimate of the rate of removals, or values used for accounting emissions or removals under the KP.

Demonstration that the deadwood and litter carbon pools are not a source

Concerning DW, the reported estimates for both AR and FL-FL directly demonstrate that this pool is not a source for AR and FM, respectively. For FM this means that we assume that FL-FL is a good representation of FM. In a similar fashion, we assume that L-FL, for which we report estimated net sink for the litter pool, is representative for AR.

For the litter pool on FM land, we currently do not have a monitoring that could provide accurate estimates for the amount of carbon stock *change*. Therefore, we apply the result of the demonstration in section 6.5.5.2.2, which is based on some measurements but mainly on sound scientific knowledge and reasoning, assuming again also that the results for FL-FL hold true for FM. Based on the above we conclude that neither the deadwood nor the litter pool are a source in both AR and FM land.

11.3.1.3 Information on whether or not indirect and natural GHG emissions and removals have been factored out

According to the report of the IPCC Expert Meeting on Revisiting the Use of Managed Land as a Proxy for Estimating National Anthropogenic Emissions and Removals (5-7 May 2009, Sao Paulo, Brazil), there are currently no scientifically sound methods to separate out indirect and natural GHG emissions and removal (IPCC, 2010). On the other hand, this is not necessarily needed if appropriate proxies are used. The above-mentioned meeting, among others, stated that, although not perfect, the currently applied proxy, i.e., the so called “managed land” proxy is one that approximates the effects of direct human induced activities.

We also note that, especially for FM, this separation is taken care of by the various steps of the *accounting*, thus, no additional separation is necessary, and we have indeed not have done any.

11.3.1.4 Changes in data and methods since the previous submission (recalculations)

Recalculations were made for all categories under the KP this year (Table 11.13). The reasons for this are the same or similar as those for the UNFCCC reporting. For other details, see Chapter 6.1.4. Partly due to these recalculations, a technical correction was made for the FM category (see later).

Table 11.13. The scale of the difference between the value of the estimates in this submission and the previous submission for the various categories under the KP: (a) AR, (b) D, (c) FM, (d)-(f) HWP.

(a)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	Afforestation and reforestation		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Area subject to the activity	kha	0	0	0	0	0	0	172	0	0	0	0	0	recalculation due to revision of area data
2022		kha	0	0	0	0	0	0	173	0	0	0	0	0	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,7	0,0	0,0	0,0	0,0	0,0	
2021	Carbon stock change in above-ground biomass, Gains	kt C	0	0	0	0	0	0	0	257	255	277	266	238	recalculation due to correction of formulas
2022		kt C	0	0	0	0	0	0	0	260	260	281	269	242	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,3	1,8	1,2	1,0	1,6	
2021	Carbon stock change in above-ground biomass, Losses	kt C	-33	-16	-27	-8	-6	-4	-6	0	-1	-4	0	-8	recalculation due to revision of area data
2022		kt C	-33	-16	-27	-8	-6	-4	-6	0	-1	-3	0	-8	
	difference, %	%	0,0	0,1	0,1	-0,3	-0,2	-0,1	2,4	0,0	47,9	-9,7	0,0	0,0	
2021	Carbon stock change in above-ground biomass, Net change	kha	217	231	239	251	234	268	239	256	255	274	260	231	see above
2022		kha	217	231	239	251	234	268	239	259	259	277	263	234	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	-0,1	1,3	1,6	1,4	1,0	1,7	
2021	Carbon stock change in below-ground biomass, Gains	kha	0	0	0	0	0	0	0	64	64	69	67	60	recalculation due to correction of formulas
2022		kha	0	0	0	0	0	0	0	65	65	70	67	61	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,3	1,8	1,2	1,0	1,6	
2021	Carbon stock change in below-ground biomass, Losses	kha	0	0	0	0	-1	0	0	0	0	0	0	0	recalculation due to revision of area data
2022		kha	0	0	0	0	-1	0	0	0	0	0	0	0	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	38,9	0,0	0,0	1,4	0,0	0,8	
2021	Carbon stock change in below-ground biomass, Net change	kha	0	0	0	0	0	0	61	64	64	69	66	59	see above
2022		kha	0	0	0	0	0	0	61	65	65	70	67	60	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	-0,2	1,3	1,8	1,2	1,0	1,6	

(b)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	Deforestation		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Area subject to the activity	kha	0	0	0	25	27	28	29	31	34	38	41	44	recalculation due to revision of area data
2022		kha	0	0	0	25	27	28	31	32	35	39	42	45	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	4,4	4,2	3,8	3,4	3,1	2,9	
2021	Carbon stock change in above-ground biomass, Losses	kt C	-6	-14	-7	-11	-29	-14	-19	-26	-33	-39	-49	0	recalculation due to revision of area data and correction of area data
2022		kt C	-8	-19	-37	-14	-40	-15	-21	-26	-33	-45	-50	0	
	difference, %	%	42,9	39,0	438,8	36,8	39,4	8,0	13,7	1,3	1,2	15,7	1,0	0,0	
2021	Carbon stock change in above-ground biomass, Net change	kt C	-6	-14	-7	-10	-29	-14	-19	-26	-33	-39	-49	0	recalculation due to revision of area data and correction of area data
2022		kt C	-8	-19	-37	-14	-40	-15	-21	-26	-33	-45	-50	0	
	difference, %	%	45,1	39,0	438,8	36,9	39,4	8,0	13,7	1,3	1,3	15,7	1,0	0,0	
2021	Carbon stock change in above-ground biomass, Losses	kha	-1	-3	-2	-2	-7	-3	-5	-6	-8	-9	-12	0	recalculation due to revision of area data and correction of area data
2022		kha	-2	-5	-9	-3	-10	-4	-5	-6	-8	-11	-12	0	
	difference, %	%	43,0	42,5	498,4	38,8	39,7	8,2	13,9	1,4	1,2	16,6	1,0	0,0	
2021	Carbon stock change in above-ground biomass, Net change	kha	-1	-3	-2	-2	-7	-3	-5	-6	-8	-9	-12	0	recalculation due to revision of area data and correction of area data
2022		kha	-2	-5	-9	-3	-10	-4	-5	-6	-8	-11	-12	0	
	difference, %	%	43,0	42,5	498,4	38,8	39,7	8,2	13,9	1,4	1,2	16,6	1,0	0,0	
2021	Carbon stock change in above-ground biomass, Net carbon stock	kha	-4	-6	-16	-4	-11	-5	-6	-12	-19	-18	-20	0	recalculation due to revision of area data and correction of area data
2022		kha	-3	-5	-9	-3	-9	-5	-17	-12	-19	-17	-20	0	
	difference, %	%	-14,6	-19,2	-44,3	-20,7	-18,8	-4,7	174,6	-0,8	-0,6	-8,7	-0,5	0,0	
2021	Carbon stock change in above-ground biomass, Net carbon stock		-1	-2	-5	-1	-3	-1	-2	-4	-6	-5	-6	-6	recalculation due to revision of area data and correction of area data
2022			-1	-2	-3	-1	-3	-2	-6	-5	-7	-6	-8	-8	
	difference, %	%	5,3	0,4	-30,4	-0,5	2,3	20,7	247,5	26,0	26,3	16,0	25,8	26,2	

(c)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	Forest management		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Area subject to the activity	kha	1877	1882	1883	1890	1895	1900	1903	1903	1902	1901	1898	1896	recalculation due to revision of area data
2022		kha	1872	1877	1878	1881	1884	1888	1888	1888	1886	1884	1881	1878	
	difference, %	%	-0,3	-0,2	-0,3	-0,5	-0,6	-0,7	-0,8	-0,8	-0,8	-0,9	-0,9	-0,9	
2021	Carbon stock change in above-ground biomass, Gains	kt C	654	429	409	369	540	383	673	787	597	660	542	713	recalculation due to revision of area data and correction of formulas
2022		kt C	666	454	422	388	597	410	664	832	657	723	631	805	
	difference, %	%	1,8	5,9	3,0	5,4	10,6	7,0	-1,5	5,6	10,1	9,5	16,4	12,9	
2021	Carbon stock change in above-ground biomass, Net change	kt C	654	429	409	369	540	383	673	787	597	660	542	713	recalculation due to revision of area data and correction of formulas
2022		kt C	666	454	422	388	597	410	664	832	657	723	631	805	
	difference, %	%	1,8	5,9	3,0	5,4	10,6	7,0	-1,5	5,6	10,1	9,5	16,4	12,9	
2021	Carbon stock change in above-ground biomass, Gains	kha	164	107	102	92	135	96	168	197	149	165	135	178	recalculation due to revision of area data and correction of formulas
2022		kha	167	114	105	97	149	102	166	208	164	181	158	201	
	difference, %	%	1,8	5,9	3,0	5,4	10,6	7,0	-1,5	5,6	10,1	9,5	16,4	12,9	
2021	Carbon stock change in above-ground biomass, Net change	kha	164	107	102	92	135	96	168	197	149	165	135	178	recalculation due to revision of area data and correction of formulas
2022		kha	167	114	105	97	149	102	166	208	164	181	158	201	
	difference, %	%	1,8	5,9	3,0	5,4	10,6	7,0	-1,5	5,6	10,1	9,5	16,4	12,9	

(d)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	HWP - sawnwood1		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Initial stock	kt C	0	6286	6163	6042	5923	5833	5747	5672	5600	5530	5460	5390	recalculation due to correction in formulas
2022		kt C	0	6325	6222	6149	6084	5991	5902	5824	5748	5676	5603	5531	
	difference, %	%	0,0	0,6	1,0	1,8	2,7	2,7	2,7	2,7	2,7	2,6	2,6	2,6	
2021	Gains	kt C	0	0	0	0	0	0	0	0	0	0	0	37	recalculation due to correction in formulas
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	38	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,8	
2021	Losses	kt C	0	-126	-123	-121	-118	-116	-115	-113	-112	-110	-109	-107	recalculation due to correction in formulas
2022		kt C	0	-126	-124	-122	-121	-120	-118	-116	-115	-113	-112	-110	
	difference, %	%	0,0	0,3	0,8	1,4	2,2	2,7	2,7	2,7	2,7	2,6	2,6	2,6	
2021	Net change	kt C	0	-86	-102	-72	-62	-90	-86	-75	-72	-70	-70	-70	recalculation due to correction in formulas
2022		kt C	0	-87	-103	-73	-65	-94	-89	-78	-75	-73	-73	-72	
	difference, %	%	0,0	0,4	1,0	2,3	4,3	3,5	3,6	4,0	4,1	4,2	4,1	3,0	

(e)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	HWP - paper and paperboard1		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Initial stock	kt C	0	0	0	0	0	0	0	0	0	0	0	82	recalculation due to correction of production data
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	91	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	11,4	
2021	Gains	kt C	0	0	0	0	0	0	0	0	0	0	0	35	recalculation due to correction of production data
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	46	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	31,3	
2021	Losses	kt C	0	0	0	0	0	0	0	0	0	0	0	-27	recalculation due to correction of production data
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	-29	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	6,3	
2021	Net change	kt C	0	0	0	0	0	0	0	0	0	0	0	8	recalculation due to correction of production data
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	18	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	112,7	

(f)

Submission year	Category, quantity	Unit	Inventory year												Reason for recalculation
	HWP - wood panels1		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
2021	Initial stock	kt C	0	2754	2679	2605	2534	2565	2657	2745	2817	2971	3134	3285	recalculation due to correction in formulas
2022		kt C	0	2858	2944	3000	3041	3057	3136	3211	3271	3412	3563	3707	
	difference, %	%	0,0	3,8	9,9	15,1	20,0	19,2	18,0	17,0	16,1	14,8	13,7	12,8	
2021	Gains	kt C	0	0	0	0	0	0	0	0	0	0	0	241	recalculation due to correction in formulas
2022		kt C	0	0	0	0	0	0	0	0	0	0	0	245	
	difference, %	%	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,8	
2021	Losses	kt C	0	-77	-75	-73	-71	-71	-72	-75	-77	-80	-85	-89	recalculation due to correction in formulas
2022		kt C	0	-79	-80	-82	-84	-85	-86	-88	-90	-93	-97	-101	
	difference, %	%	0,0	1,9	6,8	12,5	17,5	19,6	18,6	17,5	16,5	15,4	14,2	13,2	
2021	Net change	kt C	0	28	91	66	53	30	92	89	72	154	163	152	recalculation due to correction in formulas
2022		kt C	0	26	86	56	41	16	79	76	59	141	151	144	
	difference, %	%	0,0	-5,2	-5,6	-13,9	-23,6	-45,7	-14,6	-14,7	-17,7	-8,1	-7,4	-4,9	

11.3.1.5 Uncertainty estimation

Uncertainties are associated with each step of the estimation of emissions and removals. Some of the uncertainties are already assessed above, and uncertainties are also covered to some extent in Chapter 6.5.7. Uncertainties are further assessed in a detailed procedure below. This section describes methods and results of uncertainty estimation both for categories under the Kyoto Protocol and those under the UNFCCC as it seems more practicable to describe similar systems once and highlight differences.

One of the objectives of the uncertainty analysis is to demonstrate that emissions are not underestimated. It is therefore underlined here, too, that, whenever the inherent uncertainties of our estimation procedure justify that, we always take a conservative approach to avoid the underestimation of emissions and to minimize those sources of uncertainties that we are aware of.

Another, by far not unimportant, aspect of dealing with uncertainties is to identify and quantify them in order that the inventory can be developed so that the more important and/or less certain estimates can be improved first. However, we note that 2022 is the last year when an inventory under the KP must be submitted, therefore, it only makes sense to consider this aspect with respect to future inventories under the UNFCCC and the Paris Agreement. Nevertheless, we continue to report on uncertainties in this section of the NIR, too, to also indicate the efforts we have made earlier and this year to ensure that our inventory under the KP is as accurate as possible, with as low uncertainty as practicable.

One principle in the above identification and quantification is that we should first identify and quantify various amounts, and then prioritize uncertainties so that the total uncertainty of the entire inventory and its main categories could effectively be reduced by practicable policies and measures.

Concerning identification, we believe that the most important sources of uncertainties in the estimation of GHG emissions and removals due to the various KP activities include the following (the ones that are regarded less important, based on the magnitude of their size, are in brackets):

- identification of land under the various 3.3 and 3.4 activities over time,
- growing stock and its changes,
- basic wood density,
- root-to-shoot ratio,

- (carbon fraction of wood),
- carbon loss from soils, and carbon stock changes in the deadwood and litter pools due to forestry operations,
- (forest fires and other disturbances within their normal, i.e., usual, range),
- forest fires and other disturbances outside their normal range (such events, however, have not occurred in our forests in the last decades).

We note here that the uncertainty of some forest characteristics, e.g., the size of the area of land under the various activities, is rather unimportant *in the process of estimating emissions and removals* in our system because they do not directly enter the algorithm of the GHG estimation. However, when estimating the stand-level values during surveys, the area is used to upscale sampling plot information (or unit area information in case of using yield tables). Whether a land is identified or not, i.e., whether carbon stock changes on that land must be estimated or not, is also important, see the first bullet point above. In this respect, we believe that our data collection system can be regarded as conservative and may in this sense result in an underestimation of removals and overestimation of emissions as demonstrated in section 11.2.2 above.

With respect to the estimation related to the biomass on FM land, data from the forest monitoring system is used, the primary objective of which has been to obtain accurate information on the status and development of all forests in the country, and to assist forest management by developing forest management plans at the sub-compartment and forest enterprise level. The forest inventory was designed to collect data at the stand level, but to provide accurate estimates at various aggregate levels. To achieve highest efficiency while also considering practicability, different levels of accuracy are applied in the survey of individual sub-compartments depending on the age of the trees and the estimated amount and value (quality) of their growing stock.

Due to needs for accurate emission and removal estimates from D, the data collection system has been developed since 2008 so that an accurate and detailed field survey is applied to areas to be deforested, thus, a fairly high accuracy has been achieved with respect to the biomass lost in deforestations.

Concerning the estimation of carbon stock changes on AR lands, volume is estimated using yield tables, as well as ground surveys. Where the volume of the stand makes it practical to take field measurements, sampling and actual measurements are applied according to the forest monitoring protocol. The same way, where the growth of the stands is still slow and, due to the height of the trees and the thickness of the stand, the model estimates of yield tables are used as it is simply impractical to take field measurements. Because of all the above, the emission and removal estimates for biomass on AR lands can be regarded as accurate and precise as far as practicable, but with somewhat higher uncertainty than for FM or D. Also, as mentioned before, a root-to-shoot ratio is assumed for the AR stands that can be considered low for young stands, thus, below-ground biomass values are most probably underestimated. As long as AR land is a net sink, this yields a conservative estimation. Concerning the dead organic matter pools, a conservative approach was also used.

For FM, we conducted (in 2012) a thorough uncertainty analysis based on the above list of the most important sources of uncertainties. It focuses on source and sink categories in the various activities under the KP. We assume that the uncertainty estimates developed apply to the respective categories under the UNFCCC. (Concerning the estimation of the uncertainty of L-FL under the UNFCCC, a different method should have had to be applied as the methodology of estimating removals because the L-FL category is different from that for AR. However, we focused on the estimation of the uncertainty of the estimates at the activity level under the KP, and assumed that, some methodological differences notwithstanding, similar uncertainties will apply to both AR and L-FL.)

The analysis involves calculations of the emissions and removals at the same levels that are used for the GHG inventory. To obtain information on the error distributions, we applied some calculations at the stand level (see below), too. The quantifiable uncertainties were calculated using a (Tier 2) Monte Carlo (MC) analysis. The methods of the uncertainty calculations are demonstrated first for forest area, then

for carbon stock changes by pool and emissions by sources.

Forest area

As greenhouse gas information in general is related to the area of the various categories, it is important to estimate the error of area identification. This was done by assuming that the location of the borders of the stands as polygons have a maximum error of 6 m. A dedicated study showed that, in calculating the error of the size of the area, it did not matter if we used actual polygons or rectangles of the same size (see Figure 11.6 below). Thus, we could simulate the errors of the area for all stand assuming a normal distribution and using the mean size of the areas as the mean of the distribution, and the maximum and minimum areas as their range.




Figure 11.6. Possible largest, mean and smallest area of a stand if perimeters are assumed to be off (thin lines) from those in the database (i.e., those in the middle, thick lines) by a maximum 6 m in both directions (left). To simplify calculations, we used rectangles to actually estimate the error of the area (right).

Biomass pools

For all activities, we have calculated the uncertainties of the GHG inventory in the MC analysis using aggregated volume stock changes and error estimates at the *species level* (i.e., for 22 subdivisions). The estimation of the errors at this level using stand level volume stock information is described in a dedicated document that can be found at

http://www.nfk.gov.hu/download.php?id_file=40459. In essence, the National Forestry Database (NFD) contains a volume stock per unit area (m³/ha) data for each species of each of the circa 500 thousand stands. For each species, total volume stock is obtained by multiplying this species data with the area of the stands. The uncertainty of the total volume stock of a species thus depends on the uncertainty of the volume stock per unit area and that of the area data. The uncertainty of the volume stock per unit area data arises from sampling errors (when field surveys are made once in 10-12 years) and errors using yield tables (when volume stocks are updated for each year between consecutive surveys). The size of the two errors combined was assessed in a study using detailed field measurements in 642 stands. The study concluded that for stands of each slow growing species, the uncertainty of the volume stock per unit area for *individual stands* older than 40 years was typically between ±30-40%. For these stands, the uncertainty in the MC analysis was assumed to be ±40%. For stands younger than 40 years, the assumed uncertainty was ±80%. For one fast growing species, i.e., Black locust, similar values were assumed for ages above 20 years and below, respectively. The resulting overall uncertainty at the species level can be found in Table 11.10.

For basic wood density, we used a $\pm 10\%$ uncertainty based on Somogyi (2008), assuming a triangular distribution whereas default data in Table 3A.1.8 of Annex 3A.1 of the GPG for LULUCF 2003 and Table 4.3 of the 2006 IPCC GL and triangular distributions were used for other factors such as root-to-shoot ratio and carbon fraction.

Deadwood and litter

Emissions from these pools are estimated for deforested areas. The mean amount of deadwood (11.51 m³/ha) and its uncertainty ($\pm 8\%$) was estimated using a statistical sampling as part of the National Forest Inventory (<http://portal.nebih.gov.hu/en/erdoleltar/>). Other parameters to estimate carbon content are the same as for biomass above.

The amount of carbon stored in the litter pool (8.78 t C/ha) and its uncertainty ($-94/+308\%$, a rather asymmetrical interval) was derived by the literature review and expert judgment by Heil et al. (2012).

Soil

As reported above, emissions from soils are only estimated for deforested land. For the area of this land, the same uncertainty was assumed as above.

The uncertainties of the F_{LU} , F_I and F_{MG} factors were taken from Table 3.3.4 of the GPG for LULUCF (IPCC, 2003). For the uncertainty of the SOC_{ref} , country-specific values were used (Zsembeli et al. 2011, see Table 11.5). For all these factors, triangular distributions were assumed.

The estimation of N₂O emissions due to disturbances required the application of C:N ratio as well as the EF1 emission factor. The uncertainty of the former was set to be between $-48.6/+172.5\%$ (i.e., a rather asymmetrical range) based on the expert judgment by Heil et al. (2012). The uncertainty of the EF1 factor ($80/+380\%$, again, very asymmetric) was taken from pages 3.47-3.48. of the GPG for LULUCF (IPCC, 2003). Because of these asymmetrical values, it was not possible to apply the triangular distributions, and, as the emissions are small, we applied Approach 1 error propagation methods to estimate the resulting uncertainties.

Burning slash and wildfires

The estimation of the uncertainty of emissions from burning slash and wildfires is based on the formulas as reported in previous NIRs. (This means that these formulas are not exactly the same as in section 6.4.2 because the uncertainty estimation was done when the methodological basis was the GPG for LULUCF (IPCC, 2003). However, this probably does not much affect the developed uncertainty estimates). Preliminary estimates of uncertainties of the various factors were provided by Rumpf (2013). As the fraction of harvested volume burnt could not be modeled using a triangular distribution, the error propagation method was used to estimate its effect. Based on expert judgment, the uncertainty of the fraction of the amount burnt in wildfires is estimated to be $\pm 20\%$ (Debreceni, 2011). The uncertainty of factors that are the same way necessary to calculate biomass carbon as with other categories is as above. Finally, the uncertainty of the emission ratios is from Table 3A1.15 of the Annex of the GPG for LULUCF (IPCC, 2003; CH₄: $\pm 25\%$, CO: $\pm 33.3\%$, N₂O: $\pm 28.6\%$, NO_x: $\pm 22.31\%$), whereas that of the N/C is assumed to be $\pm 100\%$ based on the default value of Table A1-1 of Annex 1 of the IPCC Revised 1996 Guidelines. The uncertainty of the fraction oxidized on site was set to $\pm 10\%$.

Concerning error distributions, normal distribution was assumed for the volume data, whereas triangular distributions were assumed for the emission factors.

All input data that were applied for the Monte Carlo analysis are summarized in **Table 11.14**. The uncertainty of the combination of various pools was calculated using the error propagation method reported by IPCC (2006).

Table 11.14. *Input data for the uncertainty analysis.*

pool	gas	variable	KP category	UNFCCC category	assumed type of the pdf of errors	uncertainty value	source
area	CO ₂	area of forest subcompartment	AR, D, FM	L-FL, FL-L, FL-FL	normal	+/- 6 m in border lines	expert judgement (Mezei 2011)
biomass	CO ₂	m ³ /ha values of tree species on forest subcompartment level	AR, D, FM	L-FL, FL-L, FL-FL	normal	+/- 40 % (stands older than 40 years old), +/- 80 % (younger stands)	analyses of forest planning sampling data
	CO ₂	wood density	AR, D, FM	L-FL, FL-L, FL-FL	triangular	+/- 10 %	Somogyi (2008)
	CO ₂	carbon fraction	AR, D, FM	L-FL, FL-L, FL-FL	triangular	+/- 4.17 % (deciduous species), +/- 7.84 % (conifers)	GL for LULUCF 2006 Table 4.3
	CO ₂	root-to-shoot ratio	AR, D, FM	L-FL, FL-L, FL-FL	triangular	-50 / +100 %	GPG for LULUCF 2003 Annex 3A.1 Table 3A.1.8
	CO ₂	carbon content of orchards and vineyards	AR (losses)	L-FL (losses)	triangular	+/- 40 %	expert judgement (Juhos and Tökei 2012)
deadwood	CO ₂	area of forest subcompartment	D	-	normal	+/- 6 m in border lines	expert judgement (Mezei 2011)
	CO ₂	m ³ /ha value on country level	D	-	normal	+/- 8 %	data of National Forest Monitoring and Observation System
	CO ₂	carbon fraction	D	-	triangular	+/- 10 %	GPG 2003 Appendix 3A.1 Table 3a1.4
	CO ₂	wood density	D	-	triangular	+/- 10 %	Somogyi (2008)
litter	CO ₂	t C/ha	D	-	N/A	-94/+308 %	expert judgement (Heil et al. 2012)
slash burning	CH ₄ , CO, N ₂ O, NO _x	m ³ /ha values of tree species on forest subcompartment level	AR, D, FM	FL-FL	normal	+/- 40 % (stands older than 40 years old), +/- 80 % (younger stands)	analyses of forest planning sampling data
	CH ₄ , CO, N ₂ O, NO _x	burned slash fraction on forest subcompartment level	AR, D, FM	FL-FL	N/A	-100 % / +98-269 % (depending on the tree species; in the case of beech +2608 %, however, it means very little absolute volume value because only 1000 m ³ beech wood is burnt on site on country level)	expert judgement (Rumpf 2012)
wildfires	CH ₄ , CO, N ₂ O, NO _x	burned fraction of the total standing volume on subcompartment level	AR, FM	FL-FL	triangular	+/- 20 %	expert judgement (Debreceni 2011)
slash burning, wildfires	CH ₄ , CO, N ₂ O, NO _x	fraction oxidized on site	AR, D (slash burning only), FM	FL-FL	triangular	+/- 10 %	expert judgement (Kottek and Tobisch 2012)
	CH ₄ , CO, N ₂ O, NO _x	carbon fraction	AR, D (slash burning only), FM	FL-FL	triangular	+/- 4.17 % (deciduous species), +/- 7.84 % (conifers)	GL for LULUCF 2006 Table 4.3
	CH ₄ , CO, N ₂ O, NO _x	wood density	AR, D (slash burning only), FM	FL-FL	triangular	+/- 10 %	Somogyi (2008)
	CH ₄ , CO, N ₂ O, NO _x	emission ratio	AR, D (slash burning only), FM	FL-FL	triangular	CH ₄ : +/- 25 %, CO: +/- 33.3 %, N ₂ O: +/- 28.6 %, NO _x : +/- 22.31 %	GPG 2003 Annex 3A1 Table 3A.1.15
	N ₂ O, NO _x	N/C ratio	AR, D (slash burning only), FM	FL-FL	triangular	+/- 100 %	GPG_1996annex1ri, Table A1-1
soil	CO ₂ , N ₂ O	FLU, FI, FMG on country level	D	FL-L	triangular	FMG: +/- 9 %; FLU: +/- 10 %; FI: +/- 7 %	GPG Table 3.3.4 and area of climate and soil types (Zsembeli et al. 2011)
	CO ₂ , N ₂ O	SOCref on country level	D	FL-L	triangular	WD-HAC: +/- 85 %; CD-HAC: +/- 103 %; WD-SANDY: +/- 113 %; CD-SANDY: +/- 87 %	study of Zsembeli et al. (2011)
	N ₂ O	C/N ratio in forest soils	D	FL-L	N/A	-48.6 / +172.5 %	expert judgement (Heil et al. 2012)
	N ₂ O	EF1	D	FL-L	N/A	-80 / +380 %	GPG p. 3.47-3.48

Results

We report all results in **Table 11.15** below. According to the results, the combined uncertainty of the net removal estimates of categories under the KP amount to between about $\pm 15\%$ (for D) and $\pm 30\%$ (for FM), and the uncertainty of the activity data (volume stock change, volume and area) is the source of roughly the half of all uncertainties except for FM where it has a larger share. For AR we estimated uncertainties somewhere in between the above estimates.

As the absolute value of total emissions from D are smaller than that of the removals from AR and FM by a factor of two, the uncertainty of emissions from D is considered satisfactory. The confidence interval of the emissions from D is rather asymmetrical mainly due to the asymmetrical confidence interval of the uncertainty of the carbon stock change estimate from litter. The overall uncertainty of the emissions from D is also mainly affected by the litter uncertainty, but the biomass and soil uncertainties are also considerable. Although the factors used to estimate emissions from litter and soil can be considered country-specific, they are mainly based on expert judgment (Heil et al. 2012) but also partly representative sampling (Zsembeli et al. 2011).

For both AR and FM, the combined uncertainty practically comes from that of the biomass stock change due to the fact that other emissions are very small. Concerning the uncertainty of the biomass stock change estimates, they are affected by the uncertainty of the area, volume stock change, wood density, root-to-shoot ratio and carbon fraction estimates. Of all these, the uncertainty of the area is very small (0.03 % at the country level), and that of the wood density, root-to-shoot ratio and carbon fraction cannot really be affected by any policy, nor it is practicable to obtain more accurate estimates.

The uncertainty of the volume stock change at the stand level is due to sampling errors, measurement errors, and errors resulting from the use of yield tables which represent average forest conditions. The resulting uncertainty of the volume stock changes at the level of various species or species group varies between 15-290%. The results suggest that efforts should be taken to reduce the uncertainty of data at the stand level. The distribution of the uncertainty could also be studied in relation to the age as well as other characteristics of the stands (e.g., the mixing rates, heterogeneity of the stand structure etc.)

Table 11.15. Aggregate results of the Monte Carlo analysis for AR (a), D (b) and FM (c).**(a) AR**

Sink/source	Gas	E/R	E/R	Activity data		Emission factor		Combined		Contribution to overall uncertainty
		Gg	GgCO ₂ eq.	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	%
biomass (stock-change)	CO ₂	-1256.35	-1256.353	-12.1	11.5	-17.65	11.83	-21.4	16.5	>99
slash burning	CH ₄	0.015256	0.320375	-3.2	3.1	-66.52	97.75	-66.6	97.8	<1
slash burning	CO	0.133504	0	-3.2	3.1	-67.32	98.35	-67.4	98.4	<1
slash burning	N ₂ O	0.000105	0.032513	-3.2	3.1	-83.34	110.66	-83.4	110.7	<1
slash burning	NO _x	0.003792	0	-3.2	3.1	-83.84	110.46	-83.9	110.5	<1
wildfires	CH ₄	0.03601	0.756219	-18.4	19.3	-18.09	24.00	-25.8	30.8	<1
wildfires	CO	0.315125	0	-18.4	19.3	-25.81	30.39	-31.7	36	<1
wildfires	N ₂ O	0.000248	0.076745	-18.4	19.3	-76.11	87.80	-78.3	89.9	<1
wildfires	NO _x	0.008951	0	-18.4	19.3	-75.90	90.46	-78.1	92.5	<1
TOTAL			-1255.167							100

(b) D

Sink/source	Gas	E/R	E/R	Activity data		Emission factor		Combined		Contribution to overall uncertainty
		Gg	GgCO ₂ eq.	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	%
Deadwood	CO ₂	2.397653	2.397653	-8.7	9.2	-9.52	12.10	-12.9	15.2	0.045
litter	CO ₂	8.879759	8.879759	-2.3	2.3	-93.67	307.49	-93.7	307.5	71.385
biomass (stock-change)	CO ₂	45.75307	45.75307	-9.9	10.5	-11.14	15.71	-14.9	18.9	22.245
slash burning	CH ₄	0.005822	0.12227	-9.2	10	-51.18	93.27	-52	93.8	0.002
slash burning	CO	0.050951	0	-9.2	10	-51.79	93.57	-52.6	94.1	0.000
slash burning	N ₂ O	4E-05	0.012409	-9.2	10	-63.84	101.71	-64.5	102.2	0.000
slash burning	NO _x	0.001447	0	-9.2	10	-64.04	101.61	-64.7	102.1	0.000
soil	CO ₂	12.99638	12.99638	-2.3	2.3	-29.91	19.46	-30	19.6	6.192
soil	N ₂ O	0.000833	0.258115	-4.9	4.8	-57.79	552.18	-58	552.2	0.131
TOTAL			70.420							100

(c) FM

Sink/source	Gas	E/R	E/R	Activity data		Emission factor		Combined		Contribution to overall uncertainty
		Gg	GgCO ₂ eq.	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	CI_lower, %	CI_upper, %	%
biomass (stock-change)	CO ₂	-1560.13	-1560.134	-28.3	27.1	-13.83	11.91	-31.5	29.6	>99
slash burning	CH ₄	1.118271	23.4837	-0.7	0.7	-39.09	39.09	-39.1	221.3	<1
slash burning	CO	9.785924	0	-0.7	0.7	-39.49	39.49	-39.5	221.4	<1
slash burning	N ₂ O	0.007688	2.383262	-0.7	0.7	-48.79	48.79	-48.8	223.4	<1
slash burning	NO _x	0.277964	0	-0.7	0.7	-49.20	49.20	-49.2	223.3	<1
wildfires	CH ₄	0.486317	10.21265	-6.7	5.8	-20.64	20.91	-21.7	22.3	<1
wildfires	CO	4.255726	0	-6.7	5.8	-27.29	27.49	-28.1	28.1	<1
wildfires	N ₂ O	0.003343	1.036439	-6.7	5.8	-78.01	78.08	-78.3	87.7	<1
wildfires	NO _x	0.120882	0	-6.7	5.8	-78.31	78.39	-78.6	86.2	<1
TOTAL			-1523.018							100

11.3.1.6 Information on other methodological issues

It is important to highlight that we always use the best methods and data that is currently available. This often, but not always, represents Tier 2 or 3. In order not to underestimate emissions and overestimate removals, a highly conservative approach is applied in all steps of the inventory whenever the application of higher Tiers is not possible. This approach is characterized by always selecting data and methods that overestimate emissions and underestimate removals.

Generally, the area, harvest and forest fire statistics are based on annual nationwide assessments, whereas the emission factors and models applied do not consider the inter-annual variability of the physical processes. Therefore, the estimated emissions and removals partly, but not completely, reflect the inter-annual variability of the true processes. (The annual stock data mainly reflect actual harvests, but partly only modelled increment data.) It also needs to be underlined that the net removal values for either FM or AR represent rather small changes (i.e., net removals) relative to rather large stocks (i.e., the total carbon stocks of the biomass of all forests in the respective categories). It is due to the nature of such relatively small net values that they have a rather high inter-annual variability and are not a result of some artefacts.

We consistently use the same methods for estimating carbon stock change and non-CO₂ greenhouse gas emissions for the whole 1990-2020 period, and data reported under the KP is consistent with those under the UNFCCC.

With respect to the methodological Tiers applied in this report, at least the same or higher Tiers are applied for the categories under the KP as in our report under the UNFCCC. In general, higher tier, or at least methods of higher accuracy, are applied with respect to the identification and estimation of areas in the various land-use and land-use change categories under the KP. In general, Tier 2/3 is applied for AR, D and FM land: the land area identification is country-specific, and so is the estimation of volume, as well as that of the species-specific biomass factors to convert volume to above-ground biomass. For the expansion of above-ground to total biomass, a conservative Tier 1 factor is applied which may result in a bias in the estimation, but this bias is conservative as it is towards lower net removals.

Concerning QA/QC, estimations and QC have been done by the Forestry Department of the National Land Center, whereas the QA activities have been done by the Hungarian Forest Research Institute, in a similar fashion to the system applied for the preparation of the GHG inventory under the UNFCCC.

Almost all forestry data that have been used for the development of the GHG emission and removal estimates are collected, processed, aggregated and archived by the Forestry Department of the National Land Center. Experts of the Department participated in a training earlier on the requirements and methods of developing the GHG inventory for the forestry sector. This system ensures that all background data are collected and processed at the required quality, and the number of possible sources of errors and uncertainties are reduced. On the other hand, the expert of the Hungarian Forest Research Institute, which has been involved in the QA activities, used to develop the GHG inventory for the country, and is a lead author of various IPCC methodological Guidelines and a former reviewer for the UNFCCC thus, he is knowledgeable about the needs, method and challenges of the development of the inventory. Some data and experience of the Hungarian Forest Research Institute of the Sopron University, as well as the Institute of Site Fertility of the Sopron University, were also incorporated in the GHG inventory.

11.3.1.7 The year of the onset of an activity, if after 2008

The CRF tables under KP, as well as data and calculations as demonstrated above, clearly and transparently report both the areas and the associated emissions and removals under Article 3.3 that have entered the accounting system. For Art. 3.4 FM, activities on all land are assumed to be started before the beginning of the first commitment period. As a consequence, the Hungarian accounting system fully complies with paragraph 23 in Annex to Decision 2/CMP.7.

11.4 Article 3.3

11.4.1 Information that demonstrates that activities under Article 3.3 began on or after 1 January 1990 and before 31 December 2013 and are direct human-induced

For D and AR, field certificates of conversions exist by stand for the majority of the stands. These are archived and documented. Such certificates are only prepared for conversions that are inspected and proved to have taken place, i.e., where human activity has indeed occurred. These certificates are in general documented since 1 January 1990. Also, forest management plans are prepared for all stands in the AR category (see under section 11.5.1).

11.4.2 Information on how harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation

In Hungary, all forests must be regenerated after clearing mature stands by law (as defined by all Forest Acts since 1879, the latest one passed in 2017). Regeneration usually means that a cut-and-regeneration sequence of operations is applied, which involves that most of the area that is cut in a year is void of mature trees for many years. Moreover, regeneration may start one or two years after the final cut is made. When the regeneration is established, it may take years, even a decade, for the seedlings to reach a height of one-two meters, and a full crown closure. In general, less time is needed to reach a crown closure of 30% (i.e., the minimum requirement to meet the definition of forests), but more time may be needed in parts of the regenerations where the first attempt is not successful (where seedlings cannot establish themselves due to, e.g., bad weather conditions, weed competition, browsing by game and others). In general, the rate of closure and whether an area is cleared (deforested) or is under regeneration can only be monitored in the field.

There are country specific professional standards (as defined in the Implementation Rules of the Forest Act, 2009, practically unchanged for years) that set the time limits when regenerations (and afforestations) are deemed as successful. According to these Rules, regeneration must be started not later than two years after clear-cutting. In case of shelterwood and selection cutting, the total cover of the upper canopy layer and that of seedlings and saplings must not be lower than 60 % during the regeneration. If necessary, seedlings must be planted until 15th April of the year after harvesting. "Success" of regeneration means that it is believed that, except for rare extreme events, trees will continue to normally develop after the regeneration has been deemed successful so it can already be regarded a forest. This stage is defined by the following criteria:

- species composition is within the limits as requested by the forest management plan
- an even distribution of trees over the entire area
- healthy tree individuals overall

- the number of trees with main shoots is more than a species-specific minimum value, usually between four and eight thousand trees per hectare
- no invasive tree species is widespread in the stand
- minimum height of the main species (e.g., *Quercus petraea* and *Quercus pubescens*) reaches 1 m or 1.5 m (all other species).

This stage is to be reached by time limits that are also defined by the above Rules. The time limits depend on species and site conditions and can vary quite substantially (see Table 11.16 below). All areas that had to be regenerated have always been regenerated within these limits so far. In case the regeneration of an area is unsuccessful, it becomes part of the D category.

Table 11.16. *Time limits of completing regenerations and afforestations (years after the area becomes subject to regeneration, e.g., after clear-cutting).*

Species and origin	Time limit (years) for regeneration type: shelterwood cutting or selection cutting
<i>Quercus pubescens</i> <i>Quercus virgiliana</i>	12
<i>Quercus petraea</i> <i>Quercus robur</i> <i>Quercus cerris</i> <i>Quercus frainetto</i> <i>Fagus silvatica</i>	10
Other species	8
	Time limit (years) for other types of regeneration
<i>Quercus pubescens</i> , seed origin <i>Quercus virgiliana</i> , seed origin	14
<i>Quercus petraea</i> , seed origin <i>Quercus robur</i> , seed origin <i>Quercus frainetto</i> , seed origin <i>Fagus silvatica</i> , seed origin	12
Coniferous sp. Other hard broadleaves, seed origin	10
Other species, seed origin	8
Any species of shoot origin	5

All AR and D areas, as well as those under regeneration are identified by categorizing the above-mentioned forest compartments. These compartments have been surveyed since 1 Jan 2008 for all information that is relevant for assigning them to the respective Kyoto Forest categories (AR or D and, in case of regenerations, FM), as well as their location within each geographical area. It is also possible to identify each compartment in both the underlying database of this report (which is part of the documentation) and on the forest management maps since 2008.

Harvests on afforested areas have so far mainly been final cuttings in stands that have reached their rotation age. By law, in case an area is regenerated that was afforested or reforested earlier but after 1989, the same rules apply as for all other forests. These rules require that harvested forests must be regenerated. All areas under regeneration are continuously surveyed by the Forest Authorities, and tough penalties are applied to those forest management units that violate relevant provisions.

11.4.3 Information on the size and geographical location of forest areas that have lost forest cover but which are not yet classified as deforested

In Hungary, the Forest Authorities disclose a report each year on the current status of forests and forestry. This report includes the area of stands under regeneration. As Table 11.17 below demonstrates, this area varies around 120 kha on average but increased somewhat later (and started to decline after 2016). The same reports also state the area of final harvests each year which varied around 20-25 kha in the last three decades. From these numbers one can conclude that the average time a stand is regarded as “under regeneration” is about five-to-six years. For areas “under regeneration”, the same thresholds and criteria are in effect as for an afforested area (see section 11.4.2 and Table 11.16 above, and section 6.5.5 of the NIR). Thus, the above mean length of period of five-six years is regarded as a normal value for regenerations. (Note here, too, that individual stands can be classified “under regeneration” for a much shorter or longer time depending on species, site fertility, weather and other local conditions that determine the success of the regeneration.)

Table 11.17. *The total area of stands under regeneration as reported by annual reports on forests and forestry.*

Reporting year	Area of stands under regeneration (ha)
1985	120 043
1986	126 120
1987	128 265
1988	130 333
1989	132 956
1990	132 816
1991	136 330
1992	135 582
1993	133 522
1994	127 611
1995	120 067
1996	116 716
1997	115 768
1998	112 926
1999	110 286
2000	112 814
2001	113 825
2002	115 740
2003	117 197
2004	117 855
2005	118 989
2006	119 854
2007	120 419
2008	123 717
2009	125 344
2010	127 783
2011	131 453
2012	141 205
2013	149 997
2014	155 822
2015	165 357
2016	166 030
2017	165 052
2018	160 352
2019	161 342
2020	158 848

11.4.4 Information related to the natural disturbances provision under article 3.3

As reported in our Initial Report in 2016, Hungary does not intend to apply the provisions to exclude emissions from natural disturbances for the accounting for afforestation and reforestation under Article 3.3 (and forest management under Article 3.4) of the KP during the second commitment period.

11.4.5 Information on Harvested Wood Products under article 3.3

As requested by para 26 of Annex to 2/CMP.7, carbon stock changes in the HWP pool are reported and accounted for in the Hungarian inventory. The methodology of estimation is described in Section 11.5.2.5. In applying the methodology, it was assumed that, due to lack of data, we are unable to separate harvest from AR and FM and all harvesting is allocated to land under forest management and that all forests in Hungary are managed. Therefore, following the guidance on page 2.118 of the IPCC 2013 KP Supplement, “in case it is not possible to differentiate between the harvest from AR and FM, it is conservative and in line with good practice to assume that all HWP entering the accounting framework originate from FM”, we report carbon stock changes together for the two categories. In contrast, harvest from D is separated and excluded, and emissions from this harvest is treated as results of instantaneous oxidation.

11.5 Article 3.4

11.5.1 Information that demonstrates that activities under Article 3.4 have occurred since 1 January 1990 and are human-induced

As mentioned above, all forests are rather intensively managed in Hungary. The basis for the management is forest management plans that are prepared for all forests of the country, i.e., all stands of both the AR and the FM category. These plans, which are parts of the underlying documentation, contain information, among others, on the status of the stand during the survey, long-term objectives, plans for short-term operations (for as long as a maximum 10-year period) and information on the last harvesting operations. These plans thus demonstrate that activities under Article 3.4 have occurred since 1 January 1990 and are human-induced.

11.5.2 Information relating to Forest Management

11.5.2.1 Conversion of natural forest to planted forest

Hungary does not apply the Carbon Equivalent Forest Conversion provision (paragraphs 37 – 39) of Annex to Decision 2/CMP.7 contained in document FCCC/KP/CMP/2011/10/Add.1, p.19. Nevertheless, below we report the required information:

- plantations in Hungary are stands of exotic species (e.g., *Robinia pseudoacacia*, a species of North-American origin) or hybrids (e.g., *Populus x euramericana* clones), or that are intensively managed monocultures of indigenous species (e.g., *Salix* sp.) where the management is done for biomass production with a rotation age is much shorter than in stands that are managed for industrial wood production, protection or social services;
- natural forests are the rest of the forests;
- in general, based on the relevant provisions of the Forest Act of 2009, naturalness must not be decreased due to forest management.

11.5.2.2 Forest Management Reference Level

As reported in our Submission of information on forest management reference levels (URL: https://unfccc.int/files/meetings/ad_hoc_working_groups/kp/application/pdf/awgkp_hungary_2011.pdf), the forest management reference level (FMRL) for the second commitment period for Hungary, a member state of the EU, was developed in cooperation with the Joint Research Centre (JRC) of the European Commission in 2011. First, annual net emissions for FM were estimated for 2000-2008 and projected until 2020, assuming a 'business as usual' scenario, for the total of the above-ground and below-ground biomass carbon pools using two models of EU modelling groups, i.e., the G4M (Global Forestry Model) (from the International Institute for Applied Systems Analysis, or IIASA) and the EFISCEN (European Forest Information Scenario Model of the European Forest Institute) models. Then, the emissions and removals estimated by the models in this run for the period 2000 to 2008 were calibrated/adjusted using an offset, defined as the difference between the average of the historical forest management net emissions for 2000–2008, included in the National GHG Inventory of 2011, and the average of the mean values from the two models for the same period. (The offset was applied to the model results in order to make the projection and the historical forest management values more consistent.)

Note that, at the request of the team that reviewed the original FMRL submission, the models were re-run during the technical assessment of the FMRL submission of Hungary, producing a somewhat different output. Then, the above calibration was also repeated, yielding the FMRL value that was officially approved by the Report of the technical assessment of the forest management reference level submission of Hungary submitted in 2011 (FCCC/TAR/2011/HUN), and is used in Appendix to Decision 2/CMP.7.

The only forest pools included in the construction of the FMRL were the above- and below-ground biomass pools. Later, emissions from organic soils, and non-CO₂ emissions from wildfires were added, but the dead organic matter (litter and dead wood) and the mineral soil organic carbon pools were not included as they are demonstrated that they are not a source. The contribution of HWP to the FMRL of Hungary was estimated using the approach proposed in document FCCC/KP/AWG/2010/18/Add.1, chapter II, annex I (see section 11.5.2.5 below).

Due to a number of methodological changes since the above estimation and review, a technical correction of the FMRL has become necessary. This is described in the next section.

11.5.2.3 Technical Correction of the FMRL

Pursuant to Paragraphs 14 and 15 of Annex to Decision 2/CMP.7 (Land-use, land-use change and forestry) contained in document FCCC/KP/CMP/2011/10/Add.1, p.15, a technical correction (TC) was necessary for the above FMRL. TC in this reporting year is due to: (1) change of the method used for GHG reporting as compared to the methods used for the development of the original FMRL and its assessment; (2) change of the area under FM over time due to the inclusion of found forests (FF) into the FM category; (3) changes in the estimates of various components of the total FM emissions. Below we discuss the first two of these issues in the above order, then the combination of these with the third item.

Change of the method used for GHG reporting as compared to the methods used for the FMRL assessment

This step of the TC includes two elements.

(a) The method used for GHG reporting for biomass for FM has been different than that for the development of the FMRL, since Hungary applies the IPCC stock-difference method for FM while the FMRL was assessed by applying two process models (i.e., G4M and EFISCEN). Because of this, Hungary applied an ex-post calibration/adjustment already at the time of the FMRL submission using an offset, defined as the difference between the average of the historical FM net emissions for 2000–2008 (estimated using the stock change method) and the average of the mean values from the model for the same period. Hungary has continued to use the stock change method since this ex-post calibration. Even if there were no additional change of method or data for the historical time series for FM had not been changed since the first ex-post calibration, this calibration would have to be kept. However, since historical data before 2009 was recalculated (due to the inclusion of emissions from FF 2000-2008 into the FM category, but also due to revision of historical emissions of the FM category itself), the TC of the above offset is necessary.

(b) A further TC for the second commitment period (CP) is necessary to take into account the carbon stock changes of FF since 2008. With respect to these emissions from FF, they were included in the model estimate for 2000-2008 but were excluded from the FM estimate reported for 2000-2008, which necessitates a TC (in the form of a correction of the ex-post calibration, see below). Also, emissions from FF that occurred after 2008 were not included in the original model projection for the CP. Therefore, the FMRL has to be technically corrected.

Change of the area under FM over time. The difference in the reported FM area and the area used in the FMRL assessment for 2020

There is a difference in the area assumed in the above model that was used to develop the FMRL, and the reported area for both the period 2000-2008 and the CP. This means that the FMRL should be technically corrected to take these differences into consideration.

However, it must first be highlighted that the difference is partly due to the fact that two different area types are used in the various reports. While the statistics “forest area covered by trees” was reported for, and used in, the model (for reasons of modelling which could only consider areas covered by trees), the statistics “area of forest sub-compartments” are reported and used for FM in our submissions (see Table 6.5.1 above). To remove the difference arising from using two different area types, we converted the above “forest area covered by trees” as used by the model to “area of forest sub-compartments” using data for Forest Land for which we have data for both statistics for 2000-2020. The latter is, on average, by about 4,27 percent larger than the former one, with small annual variation. By adjusting the area used in the model by this amount one gets almost about the same area of forest sub-compartment for the period 2000-2008, with identical values for 2005 (rows #6 and #7 in **Table 11.18** below). The calculation demonstrates that all forest area, including that of FF during 2000-2008 (about 34 kha), was included in the model estimates. The adjustment also explains away $1702-1631=71$ of the $1764-1631=131$ kha difference in 2020 (using rows #6 and #3, and rows #7 and #3 of **Table 11.18**, respectively).

The remaining difference of area of about 60 kha in 2020 result from two factors plus an error term. One factor is deforestation for which (concerning areas covered by trees) the value assumed by the model (about 20 kha since 2008, i.e., the difference between values for 2020 and 2008 in row #3) turned out to be higher than the actually estimated number (concerning forest sub-compartments, about 13 kha since 2008, in that last column of row #8 in **Table 11.18**).

The other factor affecting the above remaining difference between areas is due to the fact that, as a follow-up of the recommendation by the ARR2013, we integrated the FF sub-category into the FM since 1990 category beginning 2014. As forests had also been found after 2008, the area of FM was continually increasing (unless the annual D was larger than FF, see **Table 11.5**, or when no forests were found in a year). The estimated amount of FF between 2005, i.e., about the middle year of the 2000-2008 period, and 2020 is 44 kha (row #11 in **Table 11.18**). This amount plus the adjusted modelled value of 1710 in 2020 (row #10 in **Table 11.18**) make 1754 kha. This is by 10 kha smaller than the estimated FM area of 1764 kha in 2020 (row #7 in **Table 11.18**). We regard this remaining difference as a small error that can partly be attributed to the fact that changes are non-linear over time with respect to both D and FF which creates variation around modelled values in the short term, i.e., for any single inventory year. The area of forest sub-compartments estimated for the model using the above procedure may also include error that could have contributed to the remaining difference.

Table 11.18. Derivation of revised FM area for the TC. All areas are kha.

Row #	Category	Source or equation using row#	2000	2005	2008	2010	2015	2020	
1	FM area used by G4M: area covered by trees	Hungary FMRL Submission, 2011	1 649	1 646	1 644	1 642	1 627	1 610	
2	FM area used by EFISCEN: area covered by trees		1 662	1 659	1 657	1 657	1 655	1 651	
3	Average of models: area covered by trees	= average of #1 & #2	1 656	1 653	1 651	1 650	1 641	1 631	
4	Estimated FL area: area covered by trees	Table 6.5.1 of the NIR	1 658	1 770	1 840	1 862	1 869	1 873	
5	Estimated FL area: forest subcompartments	Table 6.5.1 of the NIR	1 787	1 854	1 903	1 922	1 941	1 942	
6	Revised FM area used by model model: forest subcompartments	= #3 * mean value of (#5 / #4)	1 728	1 725	1 723	1 722	1 713	1 702	
7	Estimated FM area: forest subcompartments	Table 6.5.1 of the NIR	1 714	1 728	1 745	1 754	1 767	1 764	change since 2008
8	estimated cumulative D (=FL-L) from FM since 2000: forest subcompartments	Table 6.5.2 of the NIR	0,72	3,8	4,9	5,5	10,1	17,9	13,0
9	assumed (in revised model) cumulative D from FM since 2000: forest subcompartments	= #6 value in a year - #6 value in 2000		3,3	5,4	6,5	15,8	27,2	21,8
10	Adjusted model of FM area since 2000: forest subcompartments	= #6 in 2000 - #8	1 728	1 724	1 723	1 723	1 718	1 710	
11	estimated FF since 2000: forest subcompartments	NFDB; Table 11.2 (d) of the NIR	5,6	23	35	45	63	67	32
12	calculated FF since 2000 = difference between estimated and adjusted FM area: forest subcompartments	= #10 - #7	-14	4	22	31	49	53	32

The technical correction

All details of the TC for 2020 can be found in **Table 11.19** below. Its first section reports data that was already reported in our FMRL submission and its Revision. Then, the table reports emissions and removals from FM as estimated in this reporting year. Finally, the last part of the table lists all possible factors that might trigger TC according to Table 2.7.1 of the 2013 IPCC KP Supplement, reports the TC, and calculates the technically corrected FMRL.

For TC components that are not relevant in Hungary in an inventory year, NA is reported.

For the reduction of the area of FM due to deforestation, zero TC is made. In theory, it would be possible to estimate the effect of this smaller D value on the FMRL in terms of emissions. However, it is a rather difficult task in practice, it is not practicable to do in our case, and as the 2013 IPCC KP Supplement noted (page 2.94), “Some Parties did not consider the impact of future deforestation rate on the evolution of the FM area, assuming this has a conservative impact on the FMRL value.” To be conservative, i.e., to avoid reducing the FMRL value by using the larger D rate, we decided not to attempt to conduct this estimation.

For the other terms, the TC is either equal to estimated emissions in the second part of **Table 11.19**, or it is calculated according to the formula reported in the the respective row in the table.

The revision of the ex-post calibration (for biomass) due to the inclusion of FF in FM before 2009 and due to recalculations is reported *for the period 2000-2008*. This revision of the ex-post calibration could be directly done using the revised historical FM values as they included the FF emissions and recalculations.

In contrast, the correction of the carbon stock changes due to the inclusion of FF found *since 2008* was possible by developing annual emissions for the respective forest sub-compartments using volume data queried from the National Forestry Database. From this data, the annual volume stock change of all FF could be calculated according to the generic stock change formula by the 2006 IPCC GL, and then projection could be made for the 2013-2020 period using simple linear extrapolation. The annual carbon stock change was in turn calculated from the annual volume stock change by multiplying it by the conversion factors applied for forests for similar calculations, i.e., as required in the equations in section 6.5.3, these conversion factors include wood density, root-to-shoot ratio and carbon fraction.

Finally, the TC of the FRML due to addition of a new pool (i.e., the organic soil) and due to recalculations of several other emission estimates (including the revision of the HWP contribution earlier due to the revised methodology as reported in the IPCC 2013 KP Supplement, see section 11.5.2.5 below; see also **Table 11.10**) is included in **Table 11.19** as explained there.

The total TC for 2020 is the sum of all the above corrections.

The result of the calculations is that the total TC for 2020 is -334 ktCO₂e, and the technically corrected FMRL is -1 334 ktCO₂eq.

Table 11.19. Elements of the Technical Correction and the technically corrected FMRL. All numbers are ktCO₂eq.

Row #	FMRL or its components				average 2000- 2008	2000	2005	2010	2015	2020	average 2013-2020	TC based on average 2000- 2008 or 2013- 2020 values
1	FMRL as approved earlier	Step 1: models' results (only include CO2 from biomass)	EFISCEN		-1 394	-1 413	-1 406	-1 300	-365	522	-103	
2			G4M		-2 225	-2 055	-2 382	-2 020	-1 981	-1 611	-1 845	
3			Average of models		-1 809	-1 734	-1 894	-1 660	-1 173	-545	-974	
4		Step 2: ex-post processing	Offset	biomass	53						53	
5				non-biomass pools and GHG sources (excludes organic soils)	28						28	
6				Total offset	82						82	
7			Adjusted model results		-1 728	-1 652	-1 812	-1 578	-1 091	-463	-892	
8		Step 3: adding HWP using first-order decay function	Offset due to difference in accounting between instantaneous oxidation and first order decay function from HWP		-175						-108	
9			Adjusted model results		-1 903						-1 000	
10	FM estimate (e) or projection (p) in reporting year 2022	biomass (e); includes FF 2000-2008 but excludes FF since 2008			-1 885	240	-3 956	-1 933	-3 812	-5 055	-3 336	
11		biomass (p) of FF since 2008			NA	NA	NA	-119	-271	-422	-316	
12		CO2 emissions from organic soils (e)			62	62	62	62	62	62	62	
13		non-CO2 emissions (e)			17	20	27	12	15	15	15	
14		HWP (e)			-20	16	-69	62	-76	57	-45	
15	Technical correction (TC; relative to FMRL as approved earlier, row #9) in reporting year 2022 due to:	Changes in the method used for GHG reporting of FM or Forest Land remaining Forest Land (FL - FL) after the adoption of FMRL: see #18 and #19 below										IE
16		Changes in any of the following methodological elements used to establish the FMRL (as reported in the FMRL submission) after the adoption of FMRL:										NA
17		reduction of area of the FM land in 2008 due to D: loss of sink included in the original FMRL (conservative estimate)										0
18		change of Offset in Step 2: ex-post processing because of revised historical FM time series (due to incorporating the FF category into the FF category as a follow-up of AR2013, and due to recalculations) = (#10 - #3) - #4										-129
19		inclusion of emissions from biomass due to addition of new FF area to the FM after 2008: emissions not included in the original FMRL = #11										-316
20		recalculation of historical harvesting rates										NA
21		climate data assumed by models for projecting										NA
22		new pool (i.e., organic soils) not included in original FMRL = #12										62
23		recalculation of historical non-CO2 emissions = #13 - #5										-13
24		revision of HWP estimates = #14 - #8										63
25		Other possible methodological inconsistency										NA
26		all factors combined:										-334
27	Technically corrected FMRL in reporting year 2022				= FMRL approved earlier + total TC in reporting year 2022 =							-1 334

Explanation of the difference between the accounted quantity and the projected FMRL during the commitment period

As described in the report on FMRL (*Report of the technical assessment of the forest management reference level submission of Hungary submitted in 2011*, hereinafter called FMRL report) the original FMRL was calculated from the mean value of results of two models referring to the period 2013-2020: G4M and EFISCEN (which used data from other models: GLOBIOM and PRIMES) that was modified by an offset in order to increase accuracy.

Hungary provided historical data on various forest (FM)-related attributes, such as area, increment, harvest rate, carbon inflow in the HWP pool for the period 2000-2008. Furthermore, as also communicated to the ERTs earlier, there were no specific policies before 2010 which could have affected energy wood production. However, it was foreseen that the country *'moves towards a wider market economy, production is expected to accelerate. This will affect the demand for wood as raw material and wood for bioenergy'* - as written in paragraph 23 on page 8 of the above FMRL report. The projected harvesting rate increased by 10.1 % between 2013 and 2020 (**Table 11.20**).

Among the input variables for the calculation of the FRML, the applied models were most sensitive to the harvesting rate which means that *'an increase or a decrease of only 10 per cent in the assumed harvest value can result in significantly different results and direction (from sink to source and vice versa) [of FM removals], which indicates the importance of a sound projection of the future harvesting rates'* (par. 19 on page 7).

The actual harvests turned out to be lower than expected. Annual differences between the projected and implemented harvested volume statistics are shown in **Table 11.20**. Between 2013 and 2020, the total difference was 7.1 million m³ which means that the real harvested volume was 11 % lower than projected. According to Table 7 of the FMRL report a 10 % decrease in the assumed (projected) harvested volume means a considerable 70 % increase in FM removals.

The reason for the lower harvesting rate is that the market economy did not grow as it was expected because power stations used less wood than it had been assumed earlier. Moreover, there were two years (2019 and 2020) which were greatly affected by the COVID-19 pandemic.

Table 11.20. Projected and historical harvested volume.

Inventory year	Projected	Implemented	Difference
	harvested volume		
	thousand m3		
2013	7 904	7 875	29
2014	8 018	7 517	501
2015	8 132	7 354	778
2016	8 246	7 338	908
2017	8 360	7 576	784
2018	8 474	7 767	707
2019	8 588	7 315	1 273
2020	8 702	6 580	2 121
TOTAL	66 423	59 323	7 100

The net removals (from biomass) from FM by 2020 also increased relative to what was predicted by the models in 2011 by the larger net sink of FF. Additional differences between the projected (in 2011) and estimated (in 2020) net removals from FM might stem from a number of causes. These include the

effects of climate change and the fertilization effect (neither of which have been estimated), relatively warm years in the decade 2011-2020 (still having a positive effect on the growth of trees), the low level of natural disturbances (although elected, there was not necessary to apply the natural disturbance provision) and that the level of informal harvests (i.e., illegal cutting) could be reduced and kept at a low level, partly due to the implementation during the 2nd CP of Regulation (EU) No 995/2010 of the European Parliament and of the Council of 20 October 2010 Laying down the obligations of operators who place timber and timber products on the market. The remaining differences are due to inherent and unavoidable uncertainties in both model predictions and estimation of the annual historical emissions and removals.

11.5.2.4 Information related to the natural disturbances provision under Article 3.4

As reported in the Initial Report, Hungary did not elect the option to apply the natural disturbances provision.

11.5.2.5 Information on Harvested Wood Products under Article 3.4

From a methodological point of view, historical emissions and removals from HWP under FM are treated similarly than those under the UNFCCC, see Section 6.5.4.2.4.

The estimation was done with annual historical production data, specific half-lives for product types, application of the first-order decay function using equation 12.1 from the 2006 IPCC Guidelines, with default half-lives of two years for paper, 25 years for wood panels and 35 years for sawn wood and instantaneous oxidation assumed for wood in solid waste disposal sites. Historical data dated back to 1964. It was assumed that, except for wood harvested in deforestations, all harvested wood is allocated to forest management and that all forests in Hungary are managed. The estimates include exports.

As a result of the above procedure, the net emission estimates from the HWP pool in the FM category under the KP are only different from those under the UNFCCC in that while the latter includes harvested wood products produced from all harvests from all forests, the former excludes harvested wood products from the Deforestation category.

Concerning the contribution of the HWP pool to the FMRL, data was developed for all years until 2020 using a projection with the below steps (following the example provided in Box 2.8.2 of the IPCC 2013 KP Supplement):

1. For harvests, the same projection of an increasing trend was used as those in the development of the projection of net removals of the forests under FM (Figure 11.7).
2. Annual changes (in percent) of the projected total annual wood harvest rates were calculated between subsequent years.
3. The averages of the historical inflow rates of the most recent five years before the projected years (i.e., 2005-2009) were calculated for the sawnwood, wood-based panel and paper and paper board categories.
4. These averages were increased using the annual changes under item 2 above to get projected inflow values for each HWP pool (Figure 11.7).
5. The projected inflow values were used in Equation 2.8.5 of the IPCC 2013 KP Supplement to estimate carbon stocks, as well as gains and losses.




Figure 11.7. Historical rates of inflow that were used to develop the projected ones, and the historical and projected (in 2011) trend of rates of inflow and total wood harvests.

For the technical correction of the FMRL (see section 11.5.2.3 above), the average of the projected carbon stock changes for years 2013-2020 (projected as described above) was used.

For the sake of transparency, some additional information is provided below to demonstrate how the provisions in paragraph 16 of the Decision 2/CMP.7 are observed:

- "Emissions that occur during the second commitment period from harvested wood products removed from forests prior to the start of the second commitment period shall also be accounted for."

These emissions are only relevant for the non-firewood wood products and are estimated using the first order decay approach that accounts for wood removed from forests prior to the start of the second commitment period.

- "In the case the forest management reference level is based on a projection, a Party may choose not to account for the emissions from harvested wood products originating from forests prior to the start of the second commitment period, ..."

Hungary's FMRL is based on a projection, and Hungary has chosen not to account for the emissions from HWP originating from forests prior to the start of the second CP (i.e., Hungary has chosen one option of the "may" clause above). Mathematically, for any particular year in the commitment period, estimates of the emissions from the HWP pool from harvests before the start of the second CP are included in both the FMRL (in form of a projection) and the annual total emissions from the FM category (in the annual estimates) and, under the assumptions of the construction of the FMRL, the difference between sums of the two estimates taken for the entire CP should result in zero credits/debits. This approach is one way to not account for the the emissions from HWP originating from forests prior to the start of the second CP.

- "... and shall ensure consistency in the treatment of the harvested wood products pool in the second commitment period in accordance with paragraph 14 above."

Consistency is ensured by the application of the above-described estimation and accounting methodologies throughout the entire CP.

- "Emissions from harvested wood products already accounted for during the first commitment period on the basis of instantaneous oxidation shall be excluded."

This requirement is met by only including during the second CP emissions from the non-firewood harvested wood product sub-categories (i.e., sawnwood, wood-based panels, as well as paper and paperboard) that have been produced from harvests after the start of the second CP.

- "The treatment of harvested wood products in the construction of a projected forest management reference level shall be on the basis of provisions outlined in paragraph 29 below and shall not be on the basis of instantaneous oxidation."

This requirement is fully met by applying the first order decay functions, and other methodological elements as described in the IPCC 2013 KP Supplement.

11.5.3 Information relating to Cropland Management, Grazing Land Management, Wetland drainage and Rewetting, and Revegetation, if elected, for the base year

Hungary did not elect either Cropland Management, nor Grazing Land Management, nor Wetland drainage and Rewetting, nor Revegetation, therefore, this is a non-issue.

11.5.4 Information relating to Forest Management

11.5.4.1 That the definition of forest for this category conforms with the definition in item 11.1 above

FM land only includes managed forest areas that are included in the FL category, for which the definition of "forest" is applied as required by the Forest Act, as it is demonstrated above in section 11.1.

11.5.4.2 That forest management is a system of practices for stewardship and use of forest land aimed at fulfill relevant ecological (including biological diversity), economic and social functions of the forest in a sustainable manner (paragraph 1(f) of the annex to decision 16/CMP.1 (land-use, land-use change and forestry))

All the principles defined in paragraph 1(f) of the annex to decision 16/CMP.1 (land-use, land-use change and forestry) are among the principles of forestry of Hungary as set by law. The text of the most recent Forest Act (in Hungarian) can be found at http://net.jogtar.hu/jr/gen/hjegy_doc.cgi?docid=A0900037.TV.

11.5.4.3 Emissions and removals from Forest Management

The methodology is described in section 11.3.1.1, General methodological notes, whereas the estimated emissions and removals are reported in the KP CRF tables.

11.6 Other information

Not applicable.

11.7 Information relating to Article 6

Not applicable.

11.8 NIR tables

TABLE NIR 1. SUMMARY TABLE

Activity coverage and other information relating to activities under Article 3, paragraph 3, forest management under Article 3.4, and elected activities under Article 3.4

Activity	CHANGE IN CARBON POOL REPORTED ⁽¹⁾							GREENHOUSE GAS SOURCES REPORTED ⁽²⁾									
	Above-ground biomass	Below-ground biomass	Litter	Dead wood	Soil		HWP ⁽⁴⁾	Fertilization ⁽⁵⁾	Drained, rewetted and other soils ⁽⁶⁾		Nitrogen mineralization in mineral soils ⁽⁶⁾	Indirect N ₂ O emissions from managed soil ⁽⁵⁾	Biomass burning ⁽⁶⁾				
					Mineral	Organic ⁽³⁾			N ₂ O	CH ₄ ⁽⁷⁾	N ₂ O		N ₂ O	N ₂ O	CO ₂ ⁽⁸⁾	CH ₄	N ₂ O
Article 3.3 activities																	
Afforestation and reforestation	R	R	NR	NR	NR	NO	IE	IE	NO	NO	NO	NO	IE	R			
Deforestation	R	R	R	R	R	NO	IO	IE	NO	NO	R	R	IE	R			
Article 3.4 activities																	
Forest management	R	R	NR	NR	NR	R	R	IE	NO	NO	NO	NO	IE	R			
Cropland management	NA	NA	NA	NA	NA	NA			NA		NA		NA	NA	NA	NA	NA
Grazing land management	NA	NA	NA	NA	NA	NA			NA		NA		NA	NA	NA	NA	NA
Revegetation	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Wetland drainage and rewetting	NA	NA	NA	NA	NA	NA		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Table NIR 2. LAND TRANSITION MATRIX

Areas and changes in areas between the previous and the current inventory year^{(1),(2)}

	ARTICLE 3.3 ACTIVITIES		ARTICLE 3.4 ACTIVITIES					Other ⁽⁴⁾	Total area at the end of the previous inventory year ⁽²⁾
	Afforestation and reforestation	Deforestation	Forest management ⁽⁵⁾	Cropland management (if elected)	Grazing land management (if elected)	Revegetation (if elected)	Wetland drainage and rewetting (if elected)		
	(kha)								
Article 3.3 activities									
Afforestation and reforestation	175,50	0,41							175,92
Deforestation		45,05							45,05
Article 3.4 activities									
Forest management		1,09	1877,27						1878,37
Cropland management ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Grazing land management ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Revegetation ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Wetland drainage and rewetting ⁽³⁾ (if elected)	NA		NA	NA	NA	NA	NA		NA
Other ⁽⁴⁾	2,43	NO	1,80	NA	NA	NA	NA	7199,71	7203,94
Total area at the end of the current inventory year	177,94	46,55	1879,07	NA	NA	NA	NA	7199,71	9303,27

TABLE NIR 3. SUMMARY OVERVIEW FOR KEY CATEGORIES FOR LAND USE, LAND-USE CHANGE AND FORESTRY ACTIVITIES UNDER THE KYOTO PROTOCOL

KEY CATEGORIES OF EMISSIONS AND REMOVALS	Gas	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			Comments ⁽⁴⁾
		Associated category in UNFCCC inventory ⁽¹⁾ is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory ⁽²⁾ (including LULUCF)	Other ⁽³⁾	
Specify key categories according to the national level of disaggregation used ⁽¹⁾					
Afforestation and Reforestation	CO ₂	Land converted to forest land	Yes	NA	as key in the UNFCCC inventory.
Deforestation	CO ₂	to grassland/Land converted to settlements	Yes	NA	as key in the UNFCCC inventory.
Forest Management	CO ₂	Forest land remaining forest land	Yes	NA	as key in the UNFCCC inventory.

11.9 References

Central Agricultural Office, Forestry Directorate web page:

<http://portal.nebih.gov.hu/erdeszeti-igazgatosag>

Corine Land Cover Programme: European Environment Agency:
<http://www.eea.europa.eu/publications/COR0-landcover>.

Dömsödi, J. 2006. Földhasználat (Land-use), in Hungarian Environmental Report of Hungary (HCSO, 2006)

Hahn, Gy. 1984. Magyarország tőzeg- és lápföldvagyona. Földtani Kutatás XXVII.3:85-94.

Hungarian Central Statistical Office website:

http://www.ksh.hu/docs/hun/xstadat/xstadat_eves/i_omf001a.html

HCSO Statistical Yearbooks for Agriculture. URL: <https://www.ksh.hu/mezogazdasag>.

Genetic soil types of Hungary (Agrotopo 2000). URL: <https://www.mta-taki.hu/hu/osztalyok/kornyezetinformatikai-osztaly/agrotopo>;

http://airterkep.nebih.gov.hu/gis_portal/talajvedelem/pdf/LB-NA-27539-HU-N_online.pdf

Heil, B., Kovács, G., Szabó, O., 2012. Szakértői becslés a Kiotói Jegyzőkönyv által előírt éves gázleltár-jelentéshez. NYME research report (in Hungarian), pp. 99.

Institute of Geodesy, Cartography and Remote Sensing FÖMI (2004): Final Report CLC2000-Hungary.

Institute of Geodesy, Cartography and Remote Sensing FÖMI (2009a): Final Report CLC2006-Hungary.

Institute of Geodesy, Cartography and Remote Sensing FÖMI (2009b): Üvegházhatású gázok kibocsátási leltárhoz kapcsolódó részfeladatok elvégzése. [Sub-project for making GHG-inventory] (in Hungarian).

Institute of Geodesy, Cartography and Remote Sensing FÖMI (2014): GIO Land Monitoring – Hungary, Final report.

Hubayné Horváth, N., (2005): Felhagyott tőzegkitermelő-helyek természeti értékei és optimális hasznosítása [Natural value of cutaway peatlands and their optimal after-use]. PhD thesis. Corvinus University of Budapest. (in Hungarian with English summary).

Illés, G. et al. 2013. Jelentés a magyarországi erdőterületeken lévő szerves talajok elterjedésének, szénkészletének és a szénkészlet változását célzó becslések első eredményeiről (Report on the results of estimating the distribution, carbon stock and carbon stock changes of organic soils under forests in Hungary). The project was supported by the National Food Chain Safety Office, Forestry Directorate. Erdészeti Tudományos Intézet, Budapest. In Hungarian.

IPCC, 2003. Good Practice Guidance for Land-use, Land-Use Change and Forestry. Penman J., Gytarsky M., Hiraishi T., Krug, T., Kruger D., Pipatti R., Buendia L., Miwa K., Ngara T., Tanabe K., Wagner F. (Eds). Intergovernmental Panel on Climate Change (IPCC), IPCC/IGES, Hayama, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan. URL: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>

IPCC, 2009. Revisiting the Use of Managed Land as a Proxy for Estimating National

Anthropogenic Emissions and Removals, Meeting Report, 5 -7 May, 2009, INPE, São José dos Campos, Brazil, Pub. IGES, Japan 2009. URL: http://www.ipcc-nggip.iges.or.jp/public/mtdocs/pdfiles/0905_MLP_Report.pdf

IPCC, 2014a. 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol, Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds) Published: IPCC, Switzerland. URL: <http://www.ipcc-nggip.iges.or.jp/public/kpsg/index.htm>

IPCC, 2014b. 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds). Published: IPCC, Switzerland. URL: <http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html>

Juhos K. és Tőkei, L. (2012). A hazai szőlőkben és gyümölcsösökben tárolt szén mennyisége. [Carbon stock of vineyard and orchards in Hungary]. Report based on a project supported by the National Food Chain Safety Office, Forestry Directorate. Corvinus University of Budapest Budapesti Corvinus Egyetem Kertészettudományi Kar Talajtan és Vízgazdálkodás Tanszék (in Hungarian).

Kecskés Csaba (1997): Módszertani változások a földterületi statisztikában [Methodological changes in land-use statistics]. Statisztikai szemle. Vol 75. No. 12. 1047-1061 (in Hungarian, with English summary).

Király É., Kotték P. (2014): A hazai faipari termékekben tárolt szén mennyiségének és készletváltozásának becslése A 2013 IPCC Supplementary Guidance módszertana alapján. Erdészettudományi Közlemények 2014/I. pp. 95-110 (in Hungarian with English summary).

Király L. (1978): Új eljárások a hosszú lejáratú erdőgazdasági üzemtervek készítésében. Kandidátusi értekezés, Budapest.

Michéli, E. (1999): A FAO-talajvilágterképe és osztályozási rendszere [Soil map and classification system of the FAO]. In: Talajtan (szerk. Stefanovits P., Filep Gy., Füleky Gy.), Mezőgazda Kiadó., Budapest., ISBN 963 286 045 49239 13 5 (in Hungarian).

Nagy, Z., Pintér, K., Czóbel, Sz., Balogh, L., Horváth, L., Fóti, Sz., Barcza, Z., Weidinger, T, Csintalan, Zs, Dinh, N.Q., Grosz, B. and Tuba, Z., (2007). The carbon budget of semi-arid grassland in a wet and a dry year in Hungary. Agriculture, Ecosystems and Environment. 121 (2007) 21–29.

Rumpf, J.(2013): A fakitermelések vágásterületén maradó összes apadék – és az elégetésre kerülő rész – aránya, fafajonként. Report based on a project supported by the National Food Chain Safety Office, Forestry Directorate (in Hungarian).

Füleky Gy., Filep Gy. (1999): A Talaj szerves anyagi [The humus content of soil]. In: Talajtan (eds. Stefanovits P., Filep Gy., Füleky Gy.), Mezőgazda Kiadó. Budapest. ISBN 963 286 045 4 (in Hungarian).

Somogyi Z. (2006): How to demonstrate if a pool is not a source? – case study: forest soils in Hungary. URL: http://www.scientia.hu/cv/2006/Z_Somogyi_27_Nov_2006.pdf.

Somogyi, Z. et multiple authors (2011): A hazai erdőtalajok szénkészlet változásának becslése (Estimating soil carbon stock changes in domestic forests). Research report for the Ministry of Agriculture (in Hungarian), Budapest, pp. 194.

Somogyi, Z., Bidló, A., Csiha, I., Illés, G., 2013. Carbon balance of forest soils of an entire country: a Hungarian country-specific model based on local case studies. European Journal of Forest Research

132:825-840.

Somogyi, Z., Horváth, B. (2006): Az 1930 óta telepített erdők szénlekötéséről. Erdészeti Lapok CLI. 9. 257-259.

Somogyi, Z., Horvath, B., (2006): Detecting C-stock changes in soils of afforested areas in Hungary. Presentation at the workshop Development of Models and Forest Soil Surveys for Monitoring of Soil Carbon. April 5-8, 2006 at Koli, Finland. URL: www.metla.fi/tapahtumat/2006/soil2006

Somogyi Z. (2008): A hazai erdők üvegház hatású gáz leltára. Erdészeti kutatások 2007-2008. Vol. 92. 154. URL: <http://www.erti.hu/ek.php?id=1&fn=vol92>

Somogyi Z., J. Merganic, J. Merganicova, Illés G. (2005): CarniInvent WP 8 Final Report – D8.5 Guidelines and improved standards for monitoring and verification of carbon removals in afforestation/reforestation joint implementation projects. Result of the monitoring case study in the test site in Hungary. URL: www.joanneum.at/carboinvent/D_8_5.pdf

Sopp L. (1974): Fatömegszámítási táblázatok. Mezőgazdasági Kiadó, Budapest.

Tasi J., Halász A., Bajnok M.: Különböző típusú gyepek terméspotenciálja és takarmányozási minősége. Előadás. Gödöllő, 2016. október 6. MTA Állatnemesítési, Tenyésztési, Takarmányozási és Gyepgazdálkodási Bizottságának tudományos ülése.

Zsembeli, J., Tuba G. and Kovács Gy. (2006): Development and extension of CO₂-emission measurements for different soil surfaces. Cereal Research Communications. Vol. 34. No. 1. 359-362.

Zsembeli, J. Czimbalmos, R. Kovács, Gy. Somlyai, L. (2011): Calculation of Organic Carbon Stocks of the Soil Types of Hungary on the Base of Their Humus Content. University of Debrecen, Centre for Agricultural and Applied Economic Sciences Research Institutes and Study Farm, Karcag Research Institute. pp. 3

Zsembeli J, Czimbalmos R., Kovács Gy. et al. (2013): Revised calculation of organic carbon stocks of the soil types of Hungary on the base of their humus content (Research report.)

12 INFORMATION ON THE ACCOUNTING OF THE KYOTO PROTOCOL UNITS

Annual Submission Item	Reference / Information
15/CMP.1 annex I.E paragraph 11: Standard electronic format (SEF)	The SEF Report is submitted as a separate file created by the UNFCCC SEF Report Tool 3.8.3 The filename is: [RREG1_HU_2021_2_1.zip].
15/CMP.1 annex I.E paragraph 12: List of discrepant transactions	No discrepant transactions occurred in 2021. R2 report has been generated with UNFCCC SEF Report Tool 3.8.3. The filename is: [RREG2_HU_2021_1.xlsx].
15/CMP.1 annex I.E paragraph 13 & 14: List of CDM notifications	No CDM notifications occurred in 2021. R3 report has been generated with UNFCCC SEF Report Tool 3.8.3. The filename is: [RREG3_HU_2021_1.xlsx].
15/CMP.1 annex I.E paragraph 15: List of non-replacements	No non-replacements occurred in 2021. R4 report has been generated with UNFCCC SEF Report Tool 3.8.3. The filename is: [RREG4_HU_2021_1.xlsx].
15/CMP.1 annex I.E paragraph 16: List of invalid units	No invalid units exist at 31 December 2021. R5 report has been generated with UNFCCC SEF Report Tool 3.8.3. The filename is: [RREG5_HU_2021_1.xlsx].
15/CMP.1 annex I.E paragraph 17 Actions and changes to address discrepancies	No discrepancies have occurred in the reporting period.
15/CMP.1 annex I.E Publicly accessible information	Publicly available information accessible on the website of the Hungarian National Registry are the following: - Account information detailed in 13/CMP.1 par. 45 are available at: http://ec.europa.eu/clima/policies/ets/registry/docs/hu_accinfo_en.xls - Article 6 project information detailed in 13/CMP.1 par. 46 are available at: http://ji.unfccc.int/JI_Parties/DB/BBOE0EE02Y77126OGTQ91OS4GBWMZN/viewDFP - Holding and transaction information detailed in 13/CMP.1 par. 47 are available at: https://ec.europa.eu/assets/clima/ets/registry/sef/hu_cp2_sef_2020.xlsx

	- List of legal entities authorized by party detailed in 13/CMP.1 par. 48 is available at: http://ec.europa.eu/clima/policies/ets/registry/docs/hu_legal_en.xls
15/CMP.1 annex I.E paragraph 18 CPR Calculation	The commitment period reserve is calculated in accordance with 11/CMP.1 (and paragraph 18 of decision 1/CMP.8), based on the inventory of 2020 (NIR submission 2022).

12.1 Calculation of the commitment period reserve (CPR)

The commitment period reserve is calculated in accordance with decision 11/CMP.1 (and paragraph 18 of decision 1/CMP.8):

"Each Party included in Annex I shall maintain, in its national registry, a commitment period reserve which should not drop below 90 per cent of the Party's assigned amount calculated pursuant to Article 3, paragraphs 7 and 8, of the Kyoto Protocol, or 100 per cent of eight times its most recently reviewed inventory, whichever is lowest."

At the time of the preparation of this document the "most recently reviewed inventory" is the inventory of 2019 (National Inventory Submission 2021). However, the inventory of 2020 (National Inventory Submissions 2022) is already available and by the time this document will be assessed, the inventory of 2020 might already be the "most recently reviewed inventory", so CPR is calculated based on 2020's data. (Please note that the above choice of the most recently reviewed inventory has no effect on the CPR.)

Calculations:

(a) On the basis of assigned amount:

90% of the assigned amount of Hungary

$434,486,280 \times 0.9 = 391,037,652$ tonnes CO₂-eq

(b) On the basis of the inventory of 2020 (NIR 2022)

eight times the inventory of 2020:

$62,818,386 \times 8 = 502,547,085$ tonnes CO₂-eq

Based on the above calculations, the commitment period reserve amounts to **391,037,652 tonnes CO₂-eq**.

13 INFORMATION ON CHANGES IN NATIONAL SYSTEM

There have been no *fundamental* changes in the Hungarian national system. Nevertheless, the Forest Research Institute, that has been contributing to inventory preparation for many years, became part of the University of Sopron in 2020. So, from this year onwards, the Forest Research Institute of the University of Sopron together with the Forestry Department of NLC is responsible of the forestry part of the LULUCF sector. Also, the Institute of Agricultural Economics Nonprofit Kft. (AKI) plays a greater role in the preparation of the agriculture part of the inventory.

14 INFORMATION ON CHANGES IN NATIONAL REGISTRY

The following changes to the national registry of HU have occurred in 2021. Note that the 2021 SIAR confirms that previous recommendations have been implemented and included in the annual report.

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(a) Change of name or contact	None
15/CMP.1 annex II.E paragraph 32.(b) Change regarding cooperation arrangement	There was a change in the cooperation arrangement during the reported period as the United Kingdom of Great Britain and Northern Ireland no longer operate their registry in a consolidated manner within the Consolidated System of EU registries, CS EUR.
15/CMP.1 annex II.E paragraph 32.(c) Change to database structure or the capacity of national registry	There has been 6 new EUCR releases (versions 12.4, 13.0.2, 13.2.1, 13.3.3, 13.5.1 and 13.5.2) after version 11.5 (the production version at the time of the last Chapter 14 submission). No changes were applied to the database, whose model is provided in Annex A. No change was required to the application backup plan or to the disaster recovery plan. No change to the capacity of the national registry occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(d) Change regarding conformance to technical standards	The changes that have been introduced with versions 12.4, 13.0.2, 13.2.1, 13.3.3, 13.5.1 and 13.5.2 compared with version 11.5 of the national registry are presented in Annex B. It is to be noted that each release of the registry is subject to both regression testing and tests related to new functionality. These tests also include thorough testing against the DES and are carried out prior to the relevant major release of the version to Production (see Annex B). No other change in the registry's conformance to the technical standards occurred for the reported period.
15/CMP.1 annex II.E paragraph 32.(e) Change to discrepancies procedures	No change of discrepancies procedures occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(f) Change regarding security	No changes regarding security were introduced.
15/CMP.1 annex II.E paragraph 32.(g) Change to list of publicly available information	No change to the list of publicly available information occurred during the reported period.

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(h) Change of Internet address	No change to the registry internet address during the reported period.
15/CMP.1 annex II.E paragraph 32.(i) Change regarding data integrity measures	No change of data integrity measures occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(j) Change regarding test results	No change during the reported period.

Contact information of the registry administrator

The primary contact is:

Name: dr. Anna Fetter Mrs. Ilyés

Position: head of department

Organization: National Climate Protection Authority (NCPA)

Address: 1011 Budapest, Iskola utca 13., Hungary

Tel: +36-1-795-9423

E-mail: anna.fetter.ilyesne@itm.gov.hu

Further contacts are:

Name: Ágnes Gulyás-Béky

Position: head of unit - registry administrator

Organization: National Climate Protection Authority (NCPA)

Address: 1011 Budapest, Iskola utca 13., Hungary

Tel: +36-1-795-8019

E-mail: agnes.gulyas-beky@itm.gov.hu

15 INFORMATION ON MINIMIZATION OF ADVERSE IMPACTS IN ACCORDANCE WITH ARTICLE 3, PARAGRAPH 14

Information on how Hungary as a Party included in Annex I of the Convention is striving, under Article 3, paragraph 14, of the Kyoto Protocol, to implement its commitments mentioned in Article 3, paragraph 1, of the Kyoto Protocol in such a way as to minimize adverse social, environmental and economic impacts on developing country Parties, particularly those identified in Article 4, paragraphs 8 and 9, of the Convention.

Firstly, being an EU Member State, the Hungarian climate policy is largely determined by EU legislation. Therefore, the information provided by the European Union on the subject matter in its respective reports is relevant in case of Hungary. In accordance with Article 3, paragraph 1 of the Kyoto Protocol Hungary is committed to limit its anthropogenic carbon dioxide equivalent emissions of greenhouse gases listed in Annex A of the Protocol to such level that they are in line with Hungary's reduction targets while aiming at further emission reduction. Hungary is guided by the principle that ambitious national reduction targets shall be supported by a climate policy ensuring that adverse impacts on developing countries, such as carbon leakage are avoided. Hungary fully supports the endeavors, measures and implements regulations of the European Union targeting the avoidance of such impacts and fostering sustainable development, while in the same time also a specific policy framework has been put into practice. The policy framework is laid down in Hungary's second National Climate Change Strategy (NCCS-II) for the period 2018-2030, with an outlook to 2050. The NCCS-II – as a review of the first climate change strategy – was adopted by the 23/2018 Parliamentary resolution on 30 October 2018. Similarly to other multisectoral, horizontal strategies, NCCS-II is a strategy paper to facilitate sectoral planning. It sets out an individual set of goals and specific action lines, but does not “overwrite” any of the sectoral development efforts. According to the topics of mitigation–adaptation–awareness raising, NCCS-II includes the Hungarian Decarbonisation Roadmap laying down the (HDR) goals, priorities and action lines to reduce GHG emission. NCCS-II also covers the assessment of the expected effects of climate change in Hungary and its natural, social and economic consequences. It also includes National Adaptation Strategy (NAS) which is based on the climate vulnerability assessment of ecosystems and industries. The duties of Hungarian decarbonisation and climate change adaptation are supplemented by the Climate Awareness Raising Programme (“Partnership for Climate” Awareness-Raising Plan). The NCCS II. defines long-, mid- and short-term goals and action lines in the field of mitigation, adaptation and awareness raising. In January 2020, a “climate package” was adopted by the Hungarian Government including the first Climate Change Action Plan (CCAP) in order to implement the NCCS-II, and the draft long term climate strategy called National Clean Development Strategy which sets the basic principles for the country's 2050 climate neutrality goal. The planning of the 1st CCAP was carried out in parallel with the preparation of the National Energy and Climate Plan (NECP), and the renewal of the National Energy Strategy. In February 2020 the government has adopted another action plan regarding climate change: the Climate and Nature Protection Action Plan, which consists eight points. The final version of the long-term National Clean Development Strategy was adopted and submitted to the UNFCCC in September 2021, and the second Climate Change Action Plan is currently under governmental approval.

A 22-party Hungarian consortium led by the Hungarian Ministry for Innovation and Technology successfully applied for funding under the 2019 Call for strategic integrated projects within the framework of the EU LIFE programme. The 9-year long „LIFE-IP North-HU-Trans” project, with a total

budget of EUR 14.8 million, aims to ensure the successful implementation of the National Energy and Climate Plan, with special emphasis on the sustainable and just transition of Hungary's single largest coal region located in North Hungary. The central element of the project is the climate-friendly transformation of the Mátra Power Plant (hereinafter: MPP) with parallel efforts towards job preservation and economic diversification in the region. MPP is the second largest power plant in the Hungarian electricity production system. It accounts for nearly 50% of the total energy sector GHG emissions, and about 10% of the total national GHG emissions in Hungary. Therefore, the above strategic priorities of the NECP cannot be achieved without the gradual phase-out of the lignite-fired units of the MPP. On the other hand, MPP is also a major employer in the region providing a livelihood for 2,100 employees and almost 1,000 local companies. A complex, multidisciplinary-multistakeholder solution is being realized in order to ensure the sustainable and just transition.

Through its retraining and corporate mobilization programmes, the project will help address the situation of the miners, workers and enterprises concerned. It also contributes to the design, testing and evaluation of innovative prototypes in power plant decarbonisation, recultivation of post-mining landscapes, energy efficiency and energy community measures for the affected households and the promotion of regional green transport solutions. Furthermore, the project provides technical support for the mobilisation of other national and EU funds for ensuring the sustainable and just transition in the region.

The Hungarian Ministry for Innovation and Technology and the Swedish Environmental Protection Agency implemented a project between November 2019 and July 2021 financed by the EU Technical Support Instrument with the aim of strengthening the implementation of Hungary's National Climate Change Strategy. One of the main goals of the project was to support Hungary in establishing a coherent monitoring framework for climate and energy policies, actions, strategies at national-level. This framework is intended to enable feedback on the progress towards the climate-neutrality. The setting up of a coherent climate and energy policy monitoring framework will be one of the tasks of 2022-2023 after the Government approves the second Climate Change Action Plan.