



REPORT

M-1271 | 2019

Greenhouse Gas Emissions 1990-2017, National Inventory Report



COLOPHON

Executive institution

The Norwegian Environment Agency

Project manager for the contractor**Contact person in the Norwegian Environment Agency**

Kathrine Loe Bjønness

M-no

1271

Year

2019

Pages

534

Publisher

The Norwegian Environment Agency

Author(s)

Norwegian Environment Agency, Statistics Norway, Norwegian Institute of Bioeconomy Research

Title - Norwegian and English

Greenhouse Gas Emissions 1990-2017, National Inventory Report

Summary - sammendrag

Norges utslippsrapportering av klimagasser for perioden 1990-2017 til FN

4 emneord

Rapportering, klimagasser, utslipp, opptak

4 subject words

NIR, greenhouse gases, emissions, removals

Front page photo

Photo: Roger Hardy, Samfoto

Preface

The United Nations Framework Convention on Climate Change (UNFCCC) was adopted in 1992 and entered into force in 1994. According to Articles 4 and 12 of the Convention, Parties are required to develop and submit to the UNFCCC national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol on an annual basis.

To comply with the above requirement, Norway has prepared the present 2019 National Inventory Report (NIR). The NIR and the associated Common Reporting Format (CRF) tables have been prepared in accordance with the revised UNFCCC Reporting Guidelines on Annual Inventories as adopted by the COP by its Decision 24/CP.19. The methodologies used in the calculation of emissions are consistent with the *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The structure of this report is consistent with the UNFCCC guidelines for inventory reporting.

This National Inventory Report also includes supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol. This supplementary information comprises chapter 11 with emissions and removals from Land Use, Land-Use Change and Forestry under the Kyoto Protocol. Chapter 12 includes information on Kyoto units, chapter 13 includes information on changes in national systems, chapter 14 includes information on changes in national registries and chapter 15 includes information on minimization of adverse impacts.

The Norwegian Environment Agency, a directorate under the Norwegian Ministry of Climate and Environment, is responsible for the reporting. Statistics Norway has been the principle contributor while the Norwegian Institute of Bioeconomy Research is responsible for chapters 6 and 11 and all information regarding Land Use, Land Use Change and Forestry.

Oslo, April 12th, 2019.

Siri Sorteberg

Acting Director, Department of Climate

Norwegian Environment Agency

Table of contents

E.S. Executive Summary.....	1
Part I: Annual Inventory Submission	7
1 Introduction	8
1.1 Background information on GHG inventories and climate change	8
1.2 A description of the national inventory arrangements.....	10
1.2.1 Institutional, legal and procedural arrangements	10
1.2.2 Overview of inventory planning, preparation and management	10
1.2.3 Quality assurance, quality control and verification	11
1.2.4 Changes in the national inventory arrangements since previous submission	17
1.3 Inventory preparation, data collection, processing and storage	18
1.4 Brief general description of methodologies (including tiers used) and data sources used	19
1.4.1 Introduction.....	19
1.4.2 The main emission model	19
1.4.3 The LULUCF model	20
1.4.4 Data sources	21
1.5 Brief description of key categories	22
1.6 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals .	25
1.6.1 Approach 1 uncertainty analysis	25
1.6.2 Approach 2 uncertainty analysis	25
1.7 General assessment of completeness	29
2 Trends in Greenhouse Gas Emissions	32
2.1 Description and interpretation of emission trends for aggregated GHG emissions.....	32
2.2 Description and interpretation of emission trends by sector	35
2.2.1 Energy	36
2.2.2 Industrial processes and product use	41
2.2.3 Agriculture.....	44
2.2.4 Land Use, Land-Use Change and Forestry (LULUCF) and KP-LULUCF	46
2.2.5 Waste.....	48
2.3 Description and interpretation of emission trends by gas	51

National Inventory Report 2019 - Norway

2.3.1	Carbon dioxide (CO ₂)	53
2.3.2	Methane (CH ₄)	55
2.3.3	Nitrous oxide (N ₂ O)	58
2.3.4	Perfluorocarbons (PFCs)	61
2.3.5	Sulphur hexafluoride (SF ₆).....	62
2.3.6	Hydrofluorocarbons (HFCs)	64
2.4	Emission trends for indirect greenhouse gases and SO ₂	66
3	Energy (CRF sector 1)	67
3.1	Overview of sector	67
3.2	Energy Combustion	73
3.2.1	Overview	73
3.2.2	Energy industries, 1A1 (Key category for CO ₂ and CH ₄)	92
3.2.3	Manufacturing industries and construction, 1A2 (Key category for CO ₂ and CH ₄).....	98
3.2.4	Transport – Civil Aviation, 1A3a (Key category for CO ₂)	102
3.2.5	Transport – Road Transportation, 1A3b (Key category for CO ₂)	104
3.2.6	Transport – Railways, 1A3c	118
3.2.7	Transport – Navigation, 1A3d (Key category for CO ₂ and CH ₄)	119
3.2.8	Transport – Other transportation, 1A3e	122
3.2.9	Motorized equipment (incl. in 1A2, 1A4 and 1A5)	123
3.2.10	Other Sectors, 1A4 (Key category for CO ₂ and CH ₄)	125
3.2.11	A new boating survey was published in 2018. Plans are being made to implement the results in the model for pleasure crafts. Other emissions from energy combustion, 1A5	128
3.3	Fugitive Emissions from Coal Mining and Handling, 1B1a (Key category for CH ₄)	132
3.3.1	Description	132
3.3.2	Methodological issues	133
3.3.3	Activity data.....	134
3.3.4	Emission factors	134
3.3.5	Uncertainties and time-series consistency.....	136
3.3.6	Category-specific QA/QC and verification	137
3.3.7	Category-specific recalculations	137
3.3.8	Category-specific planned improvements.....	137
3.4	Fugitive Emissions from Oil and Natural Gas – 1B.....	138

National Inventory Report 2019 - Norway

3.4.1	Overview	138
3.4.2	Fugitive Emissions from Oil, 1.B.2.a (Key category for CO ₂)	145
3.4.3	Fugitive Emissions from Natural Gas, 1.B.2.b (Key category for CH ₄)	149
3.4.4	Fugitive Emissions from Venting and Flaring, 1.B.2.c (Key category for CO ₂ and CH ₄)	151
3.5	CO ₂ capture and storage at oil and gas production fields, 1C (Key Category for CO ₂)	158
3.5.1	Overview	158
3.5.2	CO ₂ capture and storage at the oil and gas production field Sleipner Vest	158
3.5.3	CO ₂ capture and storage at Hammerfest LNG/the gas-condensate production field Snøhvit	164
3.6	Cross-cutting issues	175
3.6.1	Sectoral versus reference approach	175
3.6.2	Quality controls within reference and sectoral approach	177
3.6.3	Feedstocks and non-energy use of fuels	179
3.7	Memo items	181
3.7.1	International bunkers	181
3.7.2	CO ₂ emissions from biomass	183
4	Industrial processes and product use (CRF sector 2)	184
4.1	Overview of sector	184
4.2	Mineral industry – 2A	186
4.2.1	Cement Production, 2A1 (Key category for CO ₂)	187
4.2.2	Lime Production, 2A2 (Key category for CO ₂)	189
4.2.3	Glass production, 2A3	192
4.2.4	Ceramics, 2A4a	192
4.2.5	Other uses of soda ash, 2A4b	193
4.2.6	Non-metallurgical magnesium production, 2A4c	195
4.2.7	Other process use of carbonates, 2A4d	196
4.3	Chemical industry – 2B	199
4.3.1	Ammonia Production, 2B1 (Key category for CO ₂)	200
4.3.2	Production of Nitric Acid, 2B2 (Key category for N ₂ O)	202
4.3.3	Silicon carbide, 2B5a (Key category for CO ₂)	204
4.3.4	Calcium carbide, 2B5b	209
4.3.5	Titanium dioxide production, 2B6 (Key category for CO ₂)	210
4.3.6	Methanol, 2B8a	211

National Inventory Report 2019 - Norway

4.3.7	Ethylene, 2B8b	212
4.3.8	Ethylene dichloride and vinyl chloride monomer, 2B8c	213
4.3.9	Other, production of fertilizers, 2B10	215
4.4	Metal industry – 2C	217
4.4.1	Steel, 2C1a.....	218
4.4.2	Production of Ferroalloys, 2C2 (Key category for CO ₂)	219
4.4.3	Aluminium production, 2C3 (Key Category for CO ₂ and PFC)	223
4.4.4	Magnesium production, 2C4 (Key category for SF ₆).....	230
4.4.5	Zinc production, 2C6.....	231
4.4.6	Anode production, 2C7i	232
4.4.7	Nickel production, 2C7ii	233
4.5	Non-energy products from fuels and solvent use – 2D	235
4.5.1	Lubricant use, 2D1 (Key category for CO ₂)	235
4.5.2	Paraffin wax use, 2D2	239
4.5.3	Solvent use, 2D3a	241
4.5.4	Road paving with asphalt, 2D3b	245
4.5.5	Other, 2D3d (use of urea as a catalyst)	246
4.6	Electronics industry – 2E	248
4.6.1	Integrated circuit or semiconductor, 2E1.....	248
4.7	Product uses as substitutes for ODS – 2F (key category for HFCs)	250
4.7.1	Refrigeration and air conditioning, 2F1	251
4.7.2	Other applications, 2F6.....	255
4.8	Other product manufacture and use – 2G	257
4.8.1	Electric equipment, 2G1.....	257
4.8.2	SF ₆ and PFC from other product use, 2G2	259
4.8.3	Use of N ₂ O in anaesthesia, 2G3a.....	260
4.8.4	Propellant for pressure and aerosol products, 2G3b.1.	261
4.8.5	Other use of N ₂ O, 2G3b.2.....	262
4.9	Other – 2H	263
4.9.1	Pulp and paper, 2H1	263
4.9.2	Food and beverages industry, 2H2.....	264
5	Agriculture (CRF sector 3)	266

National Inventory Report 2019 - Norway

5.1	Overview	266
5.2	Livestock population characterisation	268
5.2.1	Data sources	268
5.2.2	Method for estimating number of cattle	270
5.2.3	Method for estimating number of sheep	272
5.2.4	Deviations from FAO statistics	272
5.3	Nitrogen in animal manure as basis for emission estimates	274
5.4	Emissions from enteric fermentation in domestic livestock - 3A (Key category for CH ₄)	277
5.4.1	Category description	277
5.4.2	Uncertainties and time-series consistency	281
5.4.3	Category specific QA/QC and verification	281
5.4.4	Category-specific recalculations	282
5.4.5	Category-specific planned improvements	282
5.5	Emissions from manure management - 3B (Key categories for CH ₄ and N ₂ O)	283
5.5.1	Category description	283
5.5.2	Uncertainties and time-series consistency	296
5.5.3	Category specific QA/QC and verification	297
5.5.4	Category-specific recalculations	298
5.5.5	Category-specific planned improvements	298
5.6	Direct and indirect N ₂ O emissions from agricultural soils - 3D (Key categories for N ₂ O)	299
5.6.1	Category description	299
5.6.2	Uncertainties and time-series consistency	311
5.6.3	Category-specific QA/QC and verification	312
5.6.4	Category-specific recalculations	313
5.6.5	Category-specific planned improvements	313
5.7	Emissions from field burning of agricultural residues – 3F	313
5.7.1	Category description	313
5.7.2	Uncertainties and time-series consistency	314
5.7.3	Category-specific QA/QC and verification	314
5.7.4	Category-specific recalculations	314
5.7.5	Category-specific planned improvements	314
5.8	Emissions from liming – 3G (Key category for CO ₂)	315

National Inventory Report 2019 - Norway

5.8.1	Category description	315
5.8.2	Uncertainties and time-series consistency.....	315
5.8.3	Category-specific recalculations	315
5.8.4	Category-specific planned improvements.....	315
5.9	Emissions from urea application – 3H	316
5.9.1	Category description	316
5.9.2	Uncertainties and time-series consistency.....	316
5.9.3	Category-specific recalculations	316
5.9.4	Category-specific planned improvements.....	316
6	Land-use, land-use change and forestry (CRF sector 4).....	317
6.1	Sector Overview.....	317
6.1.1	Emissions and removals	317
6.1.2	Activity data.....	321
6.1.3	Uncertainties	323
6.1.4	Key categories.....	326
6.1.5	Completeness	328
6.1.6	Quality assurance and quality control (QA/QC) for LULUCF.....	328
6.2	Land-use definitions and classification system	330
6.2.1	Land-use definitions	330
6.2.2	Consistency in areas and reporting categories	332
6.2.3	Sink/source categories.....	333
6.3	Land area representation and the National Forest Inventory	335
6.3.1	Current NFI design.....	335
6.3.2	Classification of mineral and organic soil areas	337
6.3.3	Changes in the NFI design	338
6.3.4	Inter- and extrapolation for area and living biomass estimates	340
6.3.5	Uncertainties in areas and living biomass	343
6.3.6	QA/QC for the NFI data.....	344
6.4	Forest land – 4A	346
6.4.1	Forest land remaining forest land, 4A1.....	346
6.4.2	Land converted to forest land, 4A2	356
6.4.3	Completeness	359

National Inventory Report 2019 - Norway

6.5	Cropland – 4B	360
6.5.1	Cropland remaining cropland, 4B1.....	360
6.5.2	Land converted to cropland, 4B2	364
6.5.3	Completeness	367
6.6	Grassland – 4C	368
6.6.1	Grassland remaining grassland, 4C1.....	368
6.6.2	Land converted to grassland, 4C2	374
6.6.3	Completeness	376
6.7	Wetlands – 4D	377
6.7.1	Wetlands remaining wetlands, 4D1	377
6.7.2	Land converted to wetlands, 4D2	379
6.7.3	Completeness	380
6.8	Settlements – 4E	381
6.8.1	Settlements remaining settlements, 4E1	381
6.8.2	Land converted to settlements, 4E2.....	382
6.8.3	Completeness	383
6.9	Other land – 4F	384
6.9.1	Other land remaining other land, 4F1	384
6.9.2	Land converted to other land, 4F2	384
6.9.3	Completeness	385
6.10	Harvested wood products – 4G	386
6.10.1	Methodological issues	386
6.10.2	Uncertainties and time-series consistency.....	388
6.10.3	QA/QC and verification	388
6.10.4	Recalculations	388
6.10.5	Planned improvements.....	388
6.10.6	Completeness	388
6.11	Direct N ₂ O emissions from managed soils – 4(l)	389
6.11.1	Inorganic fertilizer on forest land.....	389
6.11.2	Organic fertilizer on forest land	391
6.11.3	Organic fertilizer on settlements	391
6.11.4	Uncertainties	392

National Inventory Report 2019 - Norway

6.11.5	QA/QC and verification	392
6.11.6	Recalculations	392
6.11.7	Planned improvements	392
6.11.8	Completeness	392
6.12	Emissions and removals from drainage, rewetting and other management of soils – 4(II)	393
6.12.1	N ₂ O emissions from drainage of organic soils (key category)	393
6.12.2	CH ₄ emissions from drainage of organic soils (key category)	394
6.12.3	Uncertainties	394
6.12.4	QA/QC and verification	395
6.12.5	Recalculations	395
6.12.6	Planned improvements	395
6.12.7	Completeness	395
6.13	Direct N ₂ O emissions from N mineralization and immobilization – 4(III)	396
6.13.1	Methodological issues	396
6.13.2	Recalculations	396
6.13.3	Planned improvements	396
6.13.4	Completeness	396
6.14	Indirect N ₂ O emissions from managed soils – 4(IV)	397
6.14.1	Atmospheric deposition	397
6.14.2	Nitrogen leaching and run-off	397
6.14.3	Uncertainties	398
6.14.4	QA/QC and verification	398
6.14.5	Recalculations	398
6.14.6	Planned improvements	398
6.14.7	Completeness	398
6.15	Biomass Burning – 4(V)	399
6.15.1	Fires on forest land	399
6.15.2	Completeness	402
6.16	Recalculations for LULUCF	403
6.16.1	Forest Land	405
6.16.2	Cropland	405
6.16.3	Grassland	406

National Inventory Report 2019 - Norway

6.16.4	Wetlands	406
6.16.5	Settlements	407
6.16.6	Other Land	407
6.16.7	Harvested wood products	407
6.16.8	Direct N ₂ O emission from managed soils 4(I)	407
6.16.9	Emissions and removal from drainage of organic soils 4(II)	408
6.16.10	Direct N ₂ O emissions from N mineralization and immobilization 4(III)	408
6.16.11	Indirect N ₂ O emissions from managed soils 4(IV)	408
6.16.12	Biomass burning 4(V).....	409
7	Waste (CRF sector 5)	410
7.1	Overview	410
7.2	Managed Waste Disposal on Land – 5A1	411
7.2.1	Anaerobic managed waste disposal sites, 5A1a (Key category for CH ₄)	411
7.3	Unmanaged Waste Disposal Sites – 5A2	419
7.4	Biological treatment of Solid Waste – 5B (Key category for CH ₄)	420
7.4.1	Composting and Anaerobic digestion of organic waste – 5B1 and 5B2	420
7.5	Waste incineration – 5C	425
7.5.1	Description	425
7.5.2	Methodological issues	425
7.5.3	Activity data.....	426
7.5.4	Emission factors	427
7.5.5	Uncertainties and time-series consistency.....	428
7.5.6	Source specific QA/QC and verification	428
7.5.7	Recalculations	428
7.5.8	Planned improvements.....	428
7.6	Wastewater treatment and discharge – 5D (Key category for CH ₄)	429
7.6.1	Overview	429
7.6.2	Methodological issues	430
7.6.3	Activity data.....	433
7.6.4	Emission factors	435
7.6.5	Uncertainties and time-series consistency.....	436
7.6.6	Source specific QA/QC and verification	436

National Inventory Report 2019 - Norway

7.6.7	Recalculations	436
7.6.8	Planned improvements	436
7.7	Other emissions sources from the waste sector – 5E	437
7.7.1	Description	437
7.7.2	Recalculations	437
7.7.3	Planned improvements	437
8	Other (CRF sector 6) (if applicable)	438
9	Indirect CO₂ and nitrous oxide emissions	439
9.1	Description of sources of indirect emissions in GHG inventory	439
9.2	Methodological issues	442
9.3	Uncertainties and time-series consistency	442
9.4	Category-specific QA/QC and verification	442
9.5	Category-specific recalculations	442
9.6	Category-specific planned improvements	442
10	Recalculations and improvements	443
10.1	Explanations and justifications for recalculations, including in response to the review process	443
10.2	Specific description of the recalculations	444
10.2.1	Energy	444
10.2.2	Industrial processes and product use	445
10.2.3	Agriculture	446
10.2.4	Land use, Land–Use Change and Forestry	449
10.2.5	Waste	455
10.2.6	KP-LULUCF	456
10.3	Implications for emissions levels and trends, including time-series consistency	458
10.4	Implemented and planned improvements, including in response to the review process	461
10.4.1	Work plan for further improvement of the Reference and Sectoral Approach	467
10.4.2	Work plan for further improvement of the Institutional Arrangements	470
Part II: Supplementary information required under article 7, paragraph 1		473
11	KP-LULUCF	474
11.1	General information	474
11.1.1	Relation between UNFCCC land classes and KP activities	475
11.1.2	Definitions of elected activities under Article 3.4	478

National Inventory Report 2019 - Norway

11.1.3	Description of how the definitions of each activity under Article 3.3 and 3.4 have been applied consistently over time	479
11.1.4	Hierarchy among Article 3.4 activities and how they have been consistently applied in determining how land was classified.....	479
11.2	Land-related information	480
11.2.1	Spatial assessment units used for determining the area of the units of land under Article 3.3 480	
11.2.2	Methodology used to develop the land transition matrix	480
11.2.3	Maps and/or database to identify the geographical locations and the system of identification codes for the geographical locations.....	480
11.3	Activity specific information	483
11.3.1	Methods for carbon stock change and GHG emission and removal estimates	483
11.3.2	Uncertainty estimates	483
11.3.3	Changes in data and methods since the previous submission (recalculations)	484
11.3.4	Omissions of carbon pool or GHG emissions/removals from activities under Article 3.3 and elected activities under Article 3.4.....	486
11.3.5	Provisions for natural disturbances.....	486
11.3.6	Emissions and removals from the harvested wood product pool.....	486
11.3.7	Information on whether emissions and removals have been factored out	488
11.4	Article 3.3	489
11.4.1	Activities under Article 3.3 began on or after 1 January 1990 and before 31 December of the last year of the commitment period and are directly human-induced.....	489
11.4.2	How harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation	489
11.5	Article 3.4	490
11.5.1	Activities under Article 3.4 occurred since 1 January 1990 and are human-induced	490
11.5.2	Information relating to Cropland Management, Grazing Land Management, Revegetation, and Wetland Drainage and Rewetting, if elected, for the base year	490
11.5.3	Emissions and removals from Forest Management, Cropland Management, and Grazing land Management under Article 3.4 are not accounted for under activities under Article 3.3	491
11.5.4	Conversion of natural forests to planted forests	491
11.5.5	Methodological consistency between the reference level and forest management reporting and technical corrections	491
11.5.6	Information about emissions or removals resulting from the harvest and conversion of forest plantations to non-forest land.....	493
11.6	Other information	494

National Inventory Report 2019 - Norway

11.6.1	Key category analysis for Article 3.3 activities and any elected activities under Article 3.4....	494
11.7	Information relating to Article 6	494
12	Information on accounting of Kyoto units.....	495
12.1	Background information	495
12.2	Summary of information reported in the SEF tables	495
12.3	Discrepancies and notifications	497
12.4	Publicly accessible information.....	497
12.5	Calculation of the commitment period reserve (CPR)	499
13	Information on changes in the National System.....	500
13.1	Changes in the National Greenhouse Gas Inventory System	500
14	Information on changes in national registry	501
15	Information on minimization of adverse impacts in accordance with Art. 3.14	504
16	References	508

ANNEX (I – XI) (in separate electronic document)

Annex I:	Key categories
Annex II:	Uncertainty analysis
Annex III:	Energy balance sheets
Annex IV:	CO₂ capture and storage
Annex V:	National system in Norway
Annex VI:	CRF summary 2 reports
Annex VII:	SEF and Registry Changes
Annex VIII:	QA/QC of point sources
Annex IX:	Agriculture, method description
Annex X:	Overview of notation keys NE and IE
Annex XI:	Reference and sectoral approach

Authors

This NIR has been prepared by the core institutions in the national greenhouse gas inventory system in Norway, namely the Norwegian Environment Agency, Statistics Norway and the Norwegian Institute of Bioeconomy Research. The respective authors are listed below.

Norwegian Environment Agency

Kathrine Loe Bjønnes (editor), Ketil Flugsrud, Ella Giske Havnevik, Eilev Gjerald, Hege Haugland, Mari A. Hjorthol, Britta Marie Hoem, Nina Holmengen, Julien Jabot, Tor Egil Tønnesen Kjenn, Kristin Madsen Klokkeide, Hans H. Kolshus, Anne-Grethe Kolstad, Vilde Fluge Lillesund, Mona Marstrander Rødland, Carina Heimdal Waag.

Statistics Norway

Trude Bothner Melby, Mona Irene Andersen Engedal, Eirik Knutsen, Peder Gjølstad Røhnebæk, Bjørn Thorsdalen, Trond Sandmo, Lene Skyrudsmoen og Berit Storbråten.

Norwegian Institute of Bioeconomy Research

Aaron Smith, Gry Alfredsen, Johannes Breidenbach, Lise Dalsgaard, Christophe Moni, Gunnhild Søgaard.

National Inventory Report 2019

E.S. Executive Summary

E.S.1. Background information on greenhouse gas (GHG) inventories and climate change

The 1992 United Nations Framework Convention on Climate Change (UNFCCC) requires that the Parties to the Convention develop, update and submit to the UNFCCC annual inventories of greenhouse gas emissions by sources and removals by sinks. This report documents the Norwegian National Inventory Report (NIR) 2019 for the period 1990-2017.

The report and the associated Common Reporting Format (CRF) tables have been prepared in accordance with the revised UNFCCC Reporting Guidelines on Annual Inventories as adopted by the COP by its Decision 24/CP.19. The methodologies used in the calculation of emissions are consistent with the *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. As recommended by the IPCC Guidelines, country specific methods have been used where appropriate.

Emissions of the following greenhouse gases are covered in this report: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), perfluorocarbons (PFCs), hydrofluorocarbons (HFCs) and sulphur hexafluoride (SF₆). Norway does not have any emissions of nitrogen trifluoride (NF₃) to report. In addition, the inventory includes calculations of emissions of the precursors NO_x, NMVOC, and CO, as well as for SO₂. Indirect CO₂ emissions originating from the fossil part of CH₄ and NMVOC are calculated and reported.

E.S.2 Summary of national emission and removal-related trends

In 2017, the total emissions of greenhouse gases in Norway amounted to 52.7 million tonnes CO₂ equivalents, without emissions and removals from Land-Use, Land-Use Change and Forestry (LULUCF). From 1990 to 2017, the total emissions increased by 2.9 %. Norway has experienced economic growth since 1990, with only minor setbacks in the early 1990s. The economic growth partly explains the general growth in CO₂ emissions since 1990. In addition, the offshore petroleum sector has expanded significantly during the past 20 years.

The total GHG emissions, without LULUCF, decreased by 1.7 % between 2016 and 2017. In 2017, CO₂ contributed to 82.9 % of the total emission figures, while methane and nitrous oxide contributed to 9.5 and 4.5%, respectively. PFCs, HFCs and SF₆ together accounted for 3.0 % of the total GHG emissions.

In 2017, the total net removal from the LULUCF sector was 25.0 million tonnes CO₂ equivalents. The land-use category forest land was the main contributor to the total amount of sequestration with 29.1 million tonnes of CO₂. The net greenhouse gas emissions, including all sources and sinks, were 27.7 million tonnes CO₂ equivalents in 2017, a decrease of 32.8 % from the net figure in 1990.

E.S.3 Overview of source and sink category emission estimates and trends

Figure E.S. 1 shows the overall trend in the total emissions by gas for the period 1990-2017. The proportion of CO₂ emissions of the national total greenhouse gas emissions has increased from about 69.0 % in 1990 to almost 82.9 % in 2017. The increased proportion of CO₂ relative to other gases is due to growth in the CO₂ emissions during this period, as well as a reduction in emissions of N₂O, PFCs and SF₆ gases because of implemented environmental measures and/or technological improvements and closures of industrial plants.




Figure E.S. 1 Total emissions of greenhouse gases by sources and removals from LULUCF in Norway, 1990-2017 (Mtonnes CO₂ equivalents).

Source: Statistics Norway/Norwegian Environment Agency/Norwegian Institute of Bioeconomy Research

National Inventory Report 2019 - Norway

Table E.S. 1 Emissions of greenhouse gases in Norway during the period 1990-2017. Units: CO₂ in Mtonnes (Mt), CH₄ and N₂O in ktonnes (kt) and other gases in ktonnes CO₂ eq. (kt CO₂ eq.).

Gas	CO ₂	CH ₄	N ₂ O	PFC	SF ₆	HFC
Year	Mt	kt		kt CO ₂ eq		
1990	35.3	232.0	13.7	3894.8	2098.5	0.04
1995	38.7	235.3	12.4	2314.0	579.8	92.0
2000	42.5	227.9	12.8	1518.5	891.4	383.3
2005	44.0	219.2	13.7	955.3	296.1	614.3
2008	45.4	213.1	10.5	896.0	59.8	806.1
2009	43.9	214.5	8.6	438.3	55.7	856.1
2010	46.2	215.2	8.3	238.4	68.6	1064.5
2011	45.5	208.9	8.3	262.6	54.3	1105.8
2012	45.0	207.3	8.4	200.5	53.5	1140.8
2013	44.9	208.4	8.3	181.0	56.3	1155.2
2014	44.9	212.0	8.3	178.9	50.1	1235.6
2015	45.3	207.6	8.4	146.4	69.8	1232.9
2016	44.5	203.7	8.2	186.2	63.6	1363.6
2017	43.7	200.9	8.0	131.0	58.8	1402.8

Source: Statistics Norway/Norwegian Environment Agency

Table E.S. 2 Emissions in million tonnes CO₂ equivalents in 1990, 2016, 2017 and changes (%) between 1990-2017 and 2016-2017 (without LULUCF).

Year	CO ₂	CH ₄	N ₂ O	PFCs	SF ₆	HFCs	Total
1990	35.3	5.8	4.1	3.9	2.1	0.00004	51.2
2016	44.5	5.1	2.4	0.2	0.1	1.4	53.6
2017	43.7	5.0	2.4	0.1	0.1	1.4	52.7
Changes 1990-2017	23.7 %	-13.4 %	-41.5 %	-96.6 %	-97.2 %	3187970.5 %	2.9 %
Changes 2016-2017	-1.7 %	-1.4 %	-1.8 %	-29.7 %	-7.6 %	2.9 %	-1.7 %

Source: Statistics Norway/Norwegian Environment Agency

About 52 % of the methane emissions in 2017 originated from agriculture, and 22 % originated from landfills. The total methane emissions decreased by 1.4 % from 2016 to 2017.

In 2017, agriculture and nitric acid production contributed to 74 % and 13 % of the total N₂O emissions, respectively. Due to technical improvements in production of nitric acid, and despite the increased production, the total emissions of N₂O have decreased by 42 % since 1990.

The PFC emissions decreased by 29.7 % from 2016 to 2017, and the emissions have, in total, been reduced by 96.6 % since 1990. PFC emissions originate primarily from the production of aluminium, where technical measures have been undertaken to reduce them. CO₂ emissions from aluminum production have increased since 1990 due to increased production levels.

SF₆ emissions have been reduced by 97.2 % from 1990 to 2017, mainly because of technological improvements and the closure of a magnesium production plant and a magnesium recycling foundry.

HFC emissions increased by 2.9 % in 2017 compared to 2016. Emissions in 1990 were insignificant and then increased significantly from mid-1990s until 2002. The increase in HFCs emissions has been moderated by the introduction of a tax on HFCs in 2003.

The net removal from the LULUCF sector was 25.0 million tonnes CO₂-equivalents in 2017. Since 1990, there has been an increase in carbon stored in living biomass, dead organic matter and in soils in Norway, increasing net sequestration of CO₂ by 151 % since 1990. The increase in carbon stored is a result of an active forest management policy over the last 60 to 70 years. The annual harvest rate have been much lower than the annual increments, thus causing an accumulation of wood and other tree components.

Figure E.S. 2 shows the various IPCC sectors' share of the total greenhouse gas emissions in Norway in 2016.




Figure E.S. 2 Emissions by IPCC sector in 2017, excluding LULUCF.

Source: Statistics Norway/Norwegian Environment Agency

The most important sector in Norway, with regards to the emissions of greenhouse gases (GHG), is the energy sector, accounting for 73 % of the total Norwegian emissions. The energy sector includes the energy industries (including oil and gas extraction), the transport sector, energy use in manufacturing and constructing, fugitive emissions from fuels and energy combustion in other sectors. Road traffic and offshore gas turbines (electricity generation and pumping of natural gas) are the largest single contributors, while coastal navigation and energy commodities used for the production of raw materials are other major sources.

Figure E.S. 3 shows the percentage change in emissions of greenhouse gases from 1990 to 2017 for the various IPCC sectors, compared to emissions in 1990. The development for each of the sectors

since 1990 with regards to greenhouse gas emissions, and the most important sources, are described briefly in the following.




Figure E.S. 3 Changes in GHG emissions, relative to 1990, by IPCC sector 1990-2017. Index 1990 = 1.

Source: Statistics Norway/Norwegian Environment Agency

From 1990 to 2017, the increase in the emissions from the energy sector amounted to 29 %, mainly due to higher activity in the offshore and transport sectors. The energy sector's emissions decreased by 2.2 % from 2016 to 2017. Between 1990 and 2017, there have been temporary emission reductions in e.g. 1991, 1995, 2000, 2002 and 2005 and again in 2008 and 2009, when the energy sector emissions decreased due to lower economic activity.

Emissions from Transport showed an overall increase of 24.2 % from 1990 to 2017, with a decrease of 8.9 % from 2016 to 2017. The share of transport in the total GHG emissions has increased from 19.6 % in 1990 to 23.7 % in 2017. Road transportation accounts for 70.3 % of emissions from the transport sub-sector, while emissions from navigation and civil aviation accounts for 20.4 and 8.9 %, respectively. Due to the fact that most railways are electrified in Norway, emissions of GHG from this source are insignificant

Industrial processes and other product use sector contributed to almost 16.4 % of the total national emissions of greenhouse gases in 2017. Production of metals and chemicals are the main sources of process-related industrial emissions of both CO₂ and other greenhouse gases such as N₂O (fertilizer production) and PFCs (aluminium production). Between 1990 and 2017, emissions from industrial processes experienced an overall decrease by 40.5 %. This is mainly due to reduced PFC emissions from the production of aluminium and SF₆ from the production of magnesium.

The agricultural sector contributed in 2017 to 8.5 % to the total emissions of greenhouse gases, corresponding to 4.5 million tonnes CO₂ equivalents. Emissions from agriculture increased by 0.2 % between 2016 and 2017 and decreased by 4.8 % between 1990 and 2017. The dominant sources of

GHGs are agricultural soils (N_2O) and enteric fermentation (CH_4) from domestic animals. These sources contributed to about 36.3 and 52.1 % to the sector's emissions, respectively.

The waste sector contributed to 2.3 % of total Norwegian greenhouse gas emissions in 2017. GHG emissions from the waste sector were relatively stable during the 1990s. From 1998, the emissions declined, and in 2017, they were 46.5 % lower than in 1990. Total waste volumes have increased significantly over the period, but this has been offset by increased recycling and incineration of waste as well as increased flaring of methane from landfills. Several measures introduced in the 1990s have resulted in smaller amounts of waste disposed at disposal sites. With a few exceptions, it was then prohibited to dispose easy degradable organic waste at landfills in Norway. In 1999, a tax was introduced on waste delivered to final disposal sites. From July 1 2009, it was banned to deposit biodegradable waste to landfills. This will result in further reduction of methane emissions.

E.S.4 Other information (precursors and SO_2)

Nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOC) and carbon monoxide (CO) are not greenhouse gases, but they have an indirect effect on the climate through their influence on greenhouse gases, in particular ozone. Sulphur dioxide (SO_2) also has an indirect impact on climate, as it increases the level of aerosols with a subsequent cooling effect. Therefore, emissions of these gases are to some extent included in the inventory.

The overall NO_x emissions have decreased by approximately 19.9 % from 1990 to 2017, primarily because of stricter emission regulations directed towards road traffic, which counteracted increased emissions from oil and gas production and from navigation. From 2016 to 2017, the total NO_x emissions decreased by 4.3 %.

NMVOC emissions experienced an increase in the period from 1990 to 2001, mainly because of the rise in oil production and the loading and storage of oil. However, the emissions decreased by 63.3 % from 2001 to 2017, and were, in 2017, 52.0 % lower than in 1990. From 2016 to 2017, NMVOC emissions decreased by 2.8 %.

Over the period 1990-2017, emissions of CO decreased by 50.2 %. This is primarily explained by the implementation of new emissions standards for motor vehicles.

Emissions of SO_2 were reduced by 69.9 % from 1990 to 2017. This can mainly be explained by a reduction in sulphur content of all oil products and lower process emissions from ferroalloys and aluminium productions, as well as refineries.

Part I: Annual Inventory Submission

1 Introduction

1.1 Background information on GHG inventories and climate change

The 1992 United Nation Framework Convention on Climate Change (UNFCCC) was ratified by Norway on 9 July 1993 and entered into force on 21 March 1994. One of the commitments of the Convention is that Parties are required to report their national inventories of anthropogenic emissions by sources and removals by sinks of the greenhouse gases CO₂, CH₄, N₂O as well as fluorinated greenhouse gases (HFCs, PFCs, NF₃ and SF₆), using methodologies agreed upon by the Conference of the Parties to the Convention (COP).

In compliance with its reporting requirements, Norway has submitted to the UNFCCC national emission inventory reports on an annual basis since 1993. The National Inventory Report 2019 together with the associated Common Reporting Format (CRF) tables are Norway's contribution to the 2019 round of reporting and it covers emissions and removals for the period 1990-2017.

The 2019 NIR contains supplementary information required under Article 7, paragraph 1 of the Kyoto Protocol:

- Information on anthropogenic greenhouse gas emissions by sources and removals by sinks from land use, land-use change and forestry (LULUCF) activities under Article 3, paragraph 3, and elected activities under Article 3, paragraph 4, of the Kyoto Protocol.
- Information on Kyoto units (emission reduction units, certified emission reductions, temporary certified emission reductions, long-term certified emission reductions, assigned amount units and removal units).
- Changes in national systems in accordance with Article 5, paragraph 1.
- Changes in national registries.
- Minimization of adverse impacts in accordance with Article 3, paragraph 14.

The national inventory report is prepared in accordance with the revised UNFCCC Reporting Guidelines on Annual Inventories as adopted by the COP by its Decision 24/CP.19. The methodologies used in the calculation of emissions and removals are consistent with the *2006 IPCC Guidelines for National Greenhouse Gas Inventories*.

As recommended by the IPCC Guidelines, country specific methods have been used where appropriate and where they provide more accurate emission data.

The greenhouse gases or groups of gases included in the national inventory are the following:

- Carbon dioxide (CO₂);
- Methane (CH₄);
- Nitrous oxide (N₂O);
- Hydrofluorocarbons (HFCs);
- Perfluorocarbons (PFCs);
- Sulphur hexafluoride (SF₆)

Norway has examined whether there are activities that would result in emissions of trinitrogenfluoride (NF₃) and our assessment is that there are no emissions of NF₃ in Norway.

Aggregated emissions and removals of greenhouse gases expressed in CO₂ equivalents are also reported. We have used Global Warming Potentials (GWP) calculated on a 100-year time horizon, as provided by the IPCC in the Fourth Assessment Report.

Indirect CO₂ emissions originating from the fossil part of CH₄ and NMVOC are calculated according to the reporting guidelines to the UNFCCC, and are included in the inventory. This includes emissions from fuel combustion and non-combustion sources, such as fugitive emissions from loading of crude oil, oil refineries, distribution of oil products, and from solvents and other product use.

The report also contains calculations of emissions of the precursors and indirect greenhouse gases NO_x, NMVOC, CO and SO₂, which should be included according to the reporting guidelines. However, we have in this submission not included detailed descriptions of the calculation methodologies for these gases. This information is available in the report *Informative Inventory Report (IIR) 2019. Norway* (Norwegian Environment Agency 2018).

Since the introduction of annual technical reviews of the national inventories by independent experts in 2000, Norway has undergone many desk/centralized/in-country reviews. The recommendations from these reviews have resulted in many improvements to the inventory. For the latest implemented improvements and planned improvements, see chapter 10.

1.2 A description of the national inventory arrangements

1.2.1 Institutional, legal and procedural arrangements

The Norwegian CO₂ emission inventory has been produced for more than three decades, and was gradually expanded with other emission components. It started as a collaboration between Statistics Norway and the Norwegian Environment Agency, and the reporting to the UNFCCC has evolved based on this greenhouse gas emission inventory. The Norwegian Environment Agency, Statistics Norway and the Norwegian Institute of Bioeconomy Research (NIBIO) are the institutions in the national greenhouse gas inventory system in Norway. Statistics Norway is responsible for the calculation of emissions from the Energy, IPPU, Agriculture and Waste source categories. The Norwegian Institute of Bioeconomy Research is responsible for the calculations of emission and removals from Land Use, Land-Use Change and Forestry (LULUCF).

The Norwegian Environment Agency was appointed as the national entity through the budget proposition to the Norwegian parliament (Stortinget) for 2006. These institutional arrangements have been continued for the second commitment period of the Kyoto Protocol, as described in the budget proposition to the Norwegian parliament in 2015 (Prop. 1S (2014-2015)).

To ensure that the institutions comply with their responsibilities, Statistics Norway and NIBIO have signed agreements with Norwegian Environment Agency as the national entity. Through these agreements, the institutions are committed to implementing the QA/QC and archiving procedures, providing documentation, making information available for review, and delivering data and information in a timely manner to meet the deadline for reporting to the UNFCCC.

1.2.2 Overview of inventory planning, preparation and management

The Norwegian Environment Agency, Statistics Norway, and the Norwegian Institute of Bioeconomy Research are the institutions of the national greenhouse gas inventory system, and work together to fulfill the requirements for the national system.

The allocation of responsibilities for producing estimates of emissions and removals, QA/QC and archiving is presented in more detail in section 1.2.3, section 1.3 and Annex V. An overview of institutional responsibilities and cooperation is shown in Figure 1.1.




Figure 1.1 Overview of institutional responsibilities and cooperation

1.2.3 Quality assurance, quality control and verification

1.2.3.1 Quality assurance and quality control (QA/QC)

Several quality assurance and quality control procedures for the preparation of the national emission inventory have been established in Norway during the past years. Statistics Norway made its first emission inventory for some gases in 1983 for the calculation year 1973. The emission estimation methodologies and the QA/QC procedures have been developed continuously since then.

Norway has implemented a formal quality assurance/quality control plan. The detailed description of this is found in Annex V. All three institutions annually prepare a QA/QC report, according to the plan. These reports document to what extent the QA/QC procedures have been followed. These reports are available to the expert review teams (ERT).

Based on these reports, the three institutions collaborate on which actions to take to further improve the QA/QC of the inventory.

This chapter describes general QA/QC procedures. For source specific QA/QC, see each source sector for detailed descriptions. The QA/QC work has several dimensions, of which accuracy and timeliness are the most essential. As these two aspects may be in conflict, the QA/QC improvements in recent years have focused on how to implement an effective QA/QC procedure and how to obtain a more efficient dataflow in the inventory system. Transparency is also an important issue that steadily receives more attention.

The established QA/QC procedures include the following:

- The Norwegian Environment Agency is the national entity designated to be responsible for the reporting of the national inventory of greenhouse gases to the UNFCCC. This includes coordination of the QA/QC procedures;
- Statistics Norway and the Norwegian Institute of Bioeconomy Research are responsible for the quality control system with regard to technical activities of the emission inventory preparation in their respective institutions;

- General inventory level QC procedures, as listed in table 6.1 in chapter 6, volume 1, of the 2006 IPCC Guidelines (IPCC 2000), are performed every year;
- Source category-specific QC procedures are performed for key categories and some non-key categories with regard to emission factors, activity data and uncertainty estimates.

1.2.3.2 QA Procedures

According to the IPCC Good practice guidance, good practice for QA procedures requires an objective review to assess the quality of the inventory and to identify areas where improvements should be made. Furthermore, it is good practice to use QA reviewers that have not been involved in preparing the inventory. In Norway, the Norwegian Environment Agency is responsible for reviewing the inventory with regard to quality and areas for improvement.

Norway has performed several studies comparing inventories from different countries (Kvingedal et al. 2000). Annex V gives more information concerning the quality assurance of emission data in the Norwegian emission inventory.

1.2.3.3 General QC procedures

The Norwegian emission inventory is produced in several steps. Statistics with preliminary emission estimates are published by Statistics Norway 4-5 months after the end of the inventory year. These data are based on preliminary statistics and indicators and data that have been subjected to a less thorough quality control. The more final emission statistics, which forms the basis for the emission inventory reported to the UNFCCC (for all source categories except LULUCF) is produced about one year after the inventory year. At this stage, final statistics are available for almost all emission sources. Recalculations of the inventory are performed annually to ensure that methodological changes and refinements are implemented for the whole time series. In itself, this stepwise procedure is a part of the QA/QC procedure since all differences in data are recorded and verified.

General quality control procedures are performed for each of the steps above, but with different levels of detail and thoroughness as mentioned. The national emission model was revised in 2002 in order to facilitate the QC of the input data rather than the emission data only. Input data include emissions reported from large plants, activity data, emission factors and other estimation parameters.

In the following, the procedures listed in table 6.1 in chapter 6 of the 2006 IPCC Guidelines (IPCC 2000) are described, as well as how these checks are performed for the Norwegian greenhouse gas emission inventory.

Check that assumptions and criteria for the selection of activity data and emissions factors are documented

Thorough checks of emission factors and activity data and their documentation are performed for existing emission sources. When new sources appear (for example a new industrial plant) or existing sources for the first time are recognised as a source, the Norwegian Environment Agency delivers all relevant information to Statistics Norway. This information is then thoroughly checked by the inventory team at Statistics Norway. All changes in methodologies or data are documented and kept up to date.

Check for transcription errors in data input and references

Activity data are often statistical data. Official statistical data undergo a systematic revision process, which may be manual or computerised. The revision significantly reduces the number of errors in the statistics used as input to the inventory. Furthermore, all input data (reported emissions, emission factors and activity data) for the latest inventory year are routinely compared to those of the previous inventory year, using automated procedures. Large changes are automatically flagged for further, manual QC. In addition, implied emission factors are calculated for emissions from stationary combustion at point sources. The IEFs are subjected to the same comparison between the years t and $t-1$. The most thorough checks are made for the gases and categories with the largest contribution to total emissions.

Check that emissions are calculated correctly

When possible, estimates based on different methodologies are compared. An important example is the metal production sector, where CO₂ estimates reported by the plants are compared with estimates based on the Good Practice methodology corrected for national circumstances. In this case, both production based and reducing agent based calculations are performed to verify the reported value. The Norwegian Environment Agency and Statistics Norway control and verify emission data reported to the Norwegian Environment Agency by industrial enterprises, registered in the database Forurensning. First, the Norwegian Environment Agency checks the data received from these plants, and if errors are discovered, they may then ask the plants responsible to submit new data. Subsequently, Statistics Norway makes, where possible, occasional comparable emission calculations based on activity data sampled in official statistics, and deviations are explained through contact with the plants. Regarding more detailed information about the QC of data reported by industrial plants, see Annex V and VIII.

Check that parameter and emission units are correctly recorded and that appropriate conversion factors are used

All parameter values are compared with values used in previous years and with any preliminary figures available. Whenever large deviations are detected, the value of the parameter in question is first checked for typing errors or unit errors. Changes in emissions from large plants are compared with changes in activity level. If necessary, the primary data suppliers (e.g. the Norwegian Institute of Bioeconomy Research, The Norwegian Petroleum Directorate, Norwegian Public Roads Administration, various plants etc.) are contacted for explanations and possible corrections.

Check the integrity of database files

Control checks of whether appropriate data processing steps and data relationships are correctly represented are made for each step of the process. Furthermore, it is verified that data fields are properly labelled, have correct design specifications and that adequate documentation of database and model structure and operation are archived.

Check for consistency in data between source categories

Activity data and other parameters that are common to several source categories should be evaluated for consistency. An example is recovery of landfill gas. A fraction of this gas is flared, and emissions are reported in the Waste source category. Another fraction is recovered for energy

purposes, and this gas is an input to the energy balance with emissions reported in the Energy source category. Consistency checks ensure that the amount landfill gas subtracted from source category 5A (Managed waste disposal on land), equals the amount added to source category 1A (Energy combustion) and source category 5C (Waste incineration) (the amount of gas flared).

Consistency is also checked for activity data that is used in both the Agriculture and LULUCF sectors. This is the case for the area of organic soils on croplands and grasslands, which is used to estimate CO₂ emissions in the LULUCF sector (source categories 4.B and 4.C) and N₂O emissions in the agriculture sector (source category 3D16). Within agriculture (source categories 3A, 3B and 3D), the same activity data on animal numbers and characteristics is used as far as possible.

Check that the movement for inventory data among processing steps is correct

Statistics Norway has established automated procedures to check that inventory data fed into the model does not deviate too much from the estimates for earlier years, and that the calculations within the model are correctly made. Checks are also made that emissions data are correctly transcribed between different intermediate products. The model is constructed so that it gives error messages if factors are lacking, which makes it quite robust to miscalculations.

Check that uncertainties in emissions and removals are estimated correctly

An approach 2 uncertainty analysis for greenhouse gases is undertaken annually, see further information in section 1.6.2 and Annex II.

Undertake review of internal documentation

For some sources, expert judgements dating some years back are used with regard to activity data/emission factors. In most of the cases these judgements have not been reviewed since then, and may not be properly documented, which may be a weakness of the inventory. The procedures have improved the last few years, and the requirements for internal documentation to support estimates are now quite strict; all expert judgements and assumptions made by the Statistics Norway staff should be documented. This should increase reproducibility of emissions and uncertainty estimates.

Check of changes due to recalculations

Emission time series are recalculated every year to ensure time series consistency. The recalculated emission data for a year are compared with the corresponding estimates from the year before. For example, CO₂ data calculated for 1990 in 2017 are compared with the 1990 CO₂ data calculated in 2016. The intention is to explain all major differences as far as possible. Changes may be due to revisions in energy data, new plants, correction of former errors and new emission methodologies.

Undertake completeness checks

Estimates are reported for all source categories and for all years to the best of our knowledge with the exception of a few known data gaps, which are listed in section 1.7. There may, of course, exist sources of greenhouse gases which are not covered. However, emissions from potentially additional sources are likely to be very small or negligible. During the implementation of the 2006 IPCC Guidelines, a systematic evaluation of all potential new sources was performed.

Compare estimates to previous estimates

Internal checks of time series for all emission sources are performed every year when an emission calculation for a new year is implemented. It is examined whether any detected inconsistencies are due to data and/or methodology changes. For example, in 2017 Statistics Norway/the Norwegian Environment Agency calculated emission data for 2016 for the first time. These data were compared with the 2015 estimates for detection of any considerable deviations. There may be large deviations that are correct, caused for instance by the shutdown of large industrial plants or the launch of new ones.

1.2.3.4 Source category-specific QC procedures

Statistics Norway and the Norwegian Environment Agency have carried out several studies on specific emission sources, e.g. emissions from road, sea, and air transport, emissions from landfills as well as emissions of HFCs and SF₆. These projects are repeated in regular intervals when new information is available. During the studies, emission factors have been assessed and amended in order to represent the best estimates for national circumstances, and a rationale for the choice of emission factor is provided. The emission factors are often compared with factors from literature. Furthermore, activity data have been closely examined and quality controlled, as have the uncertainty estimates.

The QC procedures with regard to emission data, activity data and uncertainty estimates for the different emission sources are described in the QA/QC-chapters of the relevant source-categories. The source category-specific analyses have primarily been performed for key categories on a case-by-case basis, which is described as being good practice. The QC procedures are described Annex V: "National Greenhouse Gas Inventory System in Norway" and Annex VIII: "QA/QC performed for GHG emissions from industrial point sources included in the national GHG inventory".

1.2.3.5 Verification studies

In general, the final inventory data provided by Statistics Norway and the Norwegian Institute of Bioeconomy Research are checked and verified by Norwegian Environment Agency. Some verification studies, which have been performed previously, are briefly described in the following.

Emission estimates for a source are often compared with estimates performed with a different methodology. In particular, Norway has conducted a study on verification of the Norwegian emission inventory (Kvingedal et al. 2000). The main goals of that work were to investigate the possibility of using statistical data as indicators for comparing emission estimates between countries on a general basis, and to test the method on the Norwegian national emission estimates. In the report, Norwegian emission data were compared with national data for Canada, Sweden and New Zealand. It was concluded that no large errors in the Norwegian emission inventory were detected. The process of verification did, however, reveal several smaller reporting errors; emissions that had been reported in other categories than they should have been. These errors were corrected. We do realize that this method of verification only considers consistency and completeness compared with what other countries report. It is not a verification of the scientific value of the inventory data themselves.

In 2002, a project funded by the Nordic Council of Ministers compared emissions of greenhouse gases from the agricultural sector in the national emission inventories with the emissions derived from the IPCC default methodology and the IPCC default factors.

In 2006, as part of the improvements for the Initial report, the Norwegian Environment Agency performed a major QA/QC exercise on the time series from 1990 to 2004 of greenhouse gas (GHG) emissions from the largest industrial plants in Norway. A first time series of emission data as well as activity data was established for each plant based on existing data sources. It was then possible to identify lack of emission data and activity data for any year or time series and possible errors in the reported data. Possible errors were typically identified if there were discrepancies between reported activity data (consumption of raw materials, production volumes etc.) and emissions, or if there were large variations in the existing time series of emissions. The emission data were supplemented and/or corrected if possible by supply of new data from the company, supplementary data from Norwegian Environment Agency paper archives, verification of reported emission data by new calculations based on reported activity data and calculation of missing emissions (if sufficient activity data were present). A final time series of greenhouse gas emissions from 1990 to 2004 were established and the main documentation from this work is contained in Excel spread sheets and in a documentation report (SFT 2006). This approach is described in Annex VIII.

From 2005 and especially from 2008, Norway's use of plant specific data has been strengthened by the availability of data from the EU ETS. The Norwegian Environment Agency conducted the verification of the annual reports up until the inventory year 2012. Since then, verification has been performed by an accredited third party. As a data source, the EU ETS provides better quality data, and these data are checked against the emissions reported under the regular permits and the reports submitted as part of the voluntary agreement. More details are found in Annex VIII.

In 2009, a new model for calculating the emissions of NMVOC from the use of solvents and other product uses was developed. The emission factors were evaluated and revised through a cooperation project between the Nordic countries. The results from the new model were compared against the similar results in Sweden and the United Kingdom; see Holmengen and Kittilsen (2009) for more details.

In 2011, the Norwegian University of Life Sciences (NMBU) published a comparison of the methodologies used for calculating CH₄ emissions from manure management in Sweden, Finland, Denmark and Norway (Morken & Hoem 2011).

In a project in 2012 at the Norwegian University of Life Sciences (NMBU) that updated the Norwegian nitrogen excretion factors and the values for manure excreted for different animal species, comparisons were made with the corresponding factors used in Sweden, Denmark and Finland and with IPCC default factors as a verification of the Norwegian factors (Karlengen et al. 2012). Comparisons were also made of the emission factors used for calculating enteric methane. In 2015, the equations for calculating emissions from enteric fermentation were evaluated and updated.

In 2015, IEFs for many of the IPPU source categories have been compared with what other Annex I countries have reported using a tool developed by the UNFCCC.¹

A Technical committee on agricultural emission was established in Norway by the end of 2017. The aim is to enhance the knowledge about possible ways to improve the emission inventory in order to

¹ http://unfccc.int/ghg_data/ghg_data_unfccc/items/4146.php

better reflect mitigation measures, and to compare the methodology used with methods in other similar countries. Members of the committee are from relevant ministries and other governmental institutions. Members of the secretariat are from expert agencies and institutes (Statistics Norway, NIBIO, Norwegian Environment Agency, and the Norwegian Agriculture Agency). The final report will be delivered in July 2019.

1.2.3.6 Quality assurance and quality control (QA/QC) and verification for the LULUCF sector

The Norwegian Institute of Bioeconomy Research (NIBIO) implements the QA/QC plan described for the National Greenhouse Gas Inventory System in Annex V. A LULUCF-specific plan for QA/QC was developed internally at NIBIO. The LULUCF-specific plan has two objectives: 1) to ensure that emission estimates and data contributing to the inventory are of high quality, and 2) to facilitate an assessment of the inventory in terms of quality and completeness. These objectives are in accordance with chapter 6 of the 2006 IPPC guidelines for quality assurance and quality control.

The QA/QC plan for the LULUCF sector is based on the general Tier 1 QC procedures and includes two check lists (one for the source-category compiler and one for the LULUCF inventory compiler), an annual timeframe of the outlined QC activities, and a target for when to elicit QA reviews. In general, QA is initiated if a new method or model is implemented.

Internal structures at NIBIO have changed slightly every year with regard to the LULUCF reporting. Existing QC procedures are evaluated and improved upon each year in order to ensure that the methods and calculations used are subjected to an internal QC prior to reporting. The CRF tables go through internal QC by more than one person before the database is submitted to the national focal point. Furthermore, after the overall compilation of estimates from all sectors, there is an exchange of CRF tables from the focal point to NIBIO, and an additional QC is performed. Improving the QA/QC procedures is an ongoing process that will be further improved in future submissions.

1.2.3.7 Confidentiality issues

In general, the data contained in the Norwegian emission inventory are available to the public, both emission estimates, activity data and emission factors. Data that are confidential according to the Statistics Act are replaced by non-confidential data collected by the Norwegian Environment Agency for most sources. Confidentiality is still an issue for some of the data collected by Statistics Norway when there are few entities reporting for a source category. In order to comply with confidentiality issues, emission estimates for these sources are aggregated. This is especially prominent in source category 2F, where emissions from 2F2-5 are aggregated in category 2F6 due to confidentiality.

1.2.4 Changes in the national inventory arrangements since previous submission

Statistics Norway, one of the three parts in the Norwegian National System, has undergone a reorganization of staff and work areas between its two offices/locations; Oslo and Kongsvinger. The experts compiling the emission inventory for all sectors except LULUCF, was up to 2018 located in Oslo. This group of experts has through 2018 been replaced by a new staff located in Kongsvinger. The long term goal of this relocation is to improve data quality by increasing the contact and collaboration between the departments producing the input (activity) data and the inventory compilers.

1.3 Inventory preparation, data collection, processing and storage

The institutions in the national inventory system; the Norwegian Environment Agency, Statistics Norway, and the Norwegian Institute of Bioeconomy Research, have agreed on a “milestone” production plan. This production plan reflects national publishing obligations etc. The plan is described in Annex V (Norway’s National System) and is supplemented by internal production plans in each of the three institutions.

The three institutions of the national system have defined areas of responsibility for data collection, this is further described in Annex V.

Statistics Norway is responsible for the collection and development of activity data, and compiling of the data used in the models that produce emission estimates for the source categories Energy, IPPU, Agriculture and Waste. Statistics Norway also operates these models. The Norwegian Environment Agency is responsible for the emission factors, for providing data from specific industries and sources and for considering the quality, and assuring necessary updating, of emissions models like e.g. the road traffic model and calculation of methane emissions from landfills. Emission data are used for a range of national applications and for international reporting. The Norwegian Institute of Bioeconomy Research is responsible for the estimated emissions from the LULUCF sectors, collects almost all data and calculates the emissions.

The collected data are subjected to the Quality Assessment and Quality Control (QA/QC) routines described in section 1.2.3.3 and Annex V, as well as source specific routines as described under each source chapter. They are all (except data regarding LULUCF) subsequently processed by Statistics Norway into a format appropriate to enter the emission models. The models are designed in a manner that accommodates both the estimation methodologies reflecting Norwegian conditions and those recommended internationally.

All three institutions are responsible for archiving the data they collect and the estimates they calculate with associated methodology documentation and internal documentation on QA/QC. Due to the differences in the character of data collected, Norway has chosen to keep archiving systems in the three institutions, which means that not all information is archived at a single location. These archiving systems are, however, consistent, and operate under the same rules. Although the data are archived separately, all can be accessed efficiently during a review. In addition, the Norwegian Environment Agency has established a library with the most important methodology reports.

1.4 Brief general description of methodologies (including tiers used) and data sources used

1.4.1 Introduction

Norway has an integrated inventory system for producing inventories of the greenhouse gases included in the Kyoto Protocol and the air pollutants SO₂, NO_x, non-methane volatile organic compounds (NMVOC), ammonia, CO, particulate matter, heavy metals and persistent organic pollutants reported under the LRTAP Convention. The data flow and QA/QC procedures are to a large extent common to all pollutants.

The emission estimation methodologies are being improved continuously. Statistics Norway and the Norwegian Environment Agency have carried out several studies on specific emission sources. Often, such projects are connected to an evaluation of emission reduction measures. An important consequence of Statistics Norway's work is increased environmental relevance of the statistical system. As far as possible, data collection relevant to the emission inventories is integrated into other surveys and statistics.

1.4.2 The main emission model

The model was developed by Statistics Norway (Daasvatn et al. 1992; 1994). It was redesigned in 2003 in order to improve reporting to the UNFCCC and LRTAP, and to improve QA/QC procedures.

Several emission sources – e.g. road traffic, agriculture, air traffic and solvents – are covered by more detailed side models. Aggregated results from these side models are used as input to the general model.

The general emission model is based on equation (1.1).

$$(1.1) \quad \text{Emissions (E)} = \text{Activity level (A)} \cdot \text{Emission Factor (EF)}$$

For emissions from *combustion*, the activity data is use of energy products. In the Norwegian energy accounts, the use of energy products is allocated to industries (economic sectors). In order to calculate emissions to air, energy use must also be allocated to technical sources (e.g. equipment). This makes it possible to match activity data with relevant emission factors.

The energy use data are combined with a corresponding matrix of emission factors. In principle, there should be one emission factor for each combination of fuel, industry, source, and pollutant. However, in a matrix with a cell for each combination, most of the cells would be empty (no consumption), while on the other hand, the same emission factor would apply to many cells.

Emissions of some pollutants from major manufacturing plants (point sources) are available from measurements or other plant-specific calculations (collected by the Norwegian Environment Agency). When such measured data are available, they are usually considered to give better representation of the actual emission, and the estimated values are replaced by the measured ones:

$$(1.2) \quad \text{Emissions } (E) = [(A - A_{PS}) \cdot EF] + E_{PS}$$

where A_{PS} and E_{PS} are the activity and the measured emissions at the point sources, respectively. Emissions from activities for which no point source estimate is available ($A - A_{PS}$) are still estimated with the regular emission factor.

Non-combustion emissions are generally calculated in the same way, by combining appropriate activity data with emission factors. Some emissions are measured directly and reported to the Norwegian Environment Agency from the plants, and some may be obtained from current reports and investigations. The emissions are fitted into the general model using the parameters industry, technical source, and pollutant. The fuel parameter is not relevant here. The source sector categories are based on EMEP/NFR and UNFCCC/CRF categories, with further subdivisions where more detailed methods are available.

The model uses approximately 220 *industries* (economic sectors). The classification is common with the basis data in the energy balance/accounts, and is almost identical to that used in the national accounts, which is aggregated from the European NACE classification (Statistics Norway 2008). The large number of sectors is an advantage in dealing with important emissions from manufacturing industries. The disadvantage is an unnecessary disaggregation of sectors with very small emissions. To make the standard sectors more appropriate for calculation of emissions, a few changes have been made, e.g. "Private households" is defined as a sector.

1.4.3 The LULUCF model

The Norwegian Institute of Bioeconomy Research is in charge of estimating emissions and removals from Land use, Land-Use Change and Forestry (LULUCF) where most of the categories have area statistics as activity data. A software based calculation system that primarily uses the data analysis software R, was developed for the implementation of the IPCC good practice guidance for the LULUCF sector. The system uses input data from different sources and creates final output datasets. These final datasets include all the data needed for the tables in the common reporting format (CRF) for both the Climate Convention and the Kyoto Protocol.

The National Forest Inventory (NFI) database contains data on areas for all land uses and land-use conversions as well as carbon stocks in living biomass. The NFI is used to estimate total areas of forest land, cropland, grassland, wetlands, settlements, other land, and land-use transitions between these categories. The data from the NFI are complemented with other data (e.g. timber harvest, horticulture, crop types, fertilizer use, drainage of forest soil, and forest fires) collected by Statistics Norway, Norwegian Agricultural Authority, Food Safety Authority, The Norwegian Directorate for Nature Management, and The Directorate for Civil Protection and Emergency Planning.

The sampling design of the NFI is based on a systematic grid of geo-referenced sample plots covering the entire country. The NFI utilizes a 5-year cycle based on a re-sampling method of the permanent plots (interpenetrating panel design). Up until 2010 the estimates were based on detailed information from sample plots in lowlands outside Finnmark county. Since 2010 the NFI has been expanded to include mountainous areas and Finnmark county in order to monitor the land use, land-use changes, and forestry activities in the whole country. All areas were included for the first time in

the estimates for the LULUCF sector in the 2012 submission. Area estimates have been bridged in a consistent manner.

The estimates of carbon stocks and their changes in living biomass are based on single tree measurements of trees larger than 50 mm at 1.3 m height (DBH) on sample plots within forest and other wooded land. Biomass is calculated using single tree allometric biomass models developed in Sweden for Norway spruce and Scots pine (Marklund 1988; Petersson & Ståhl 2006) and Norwegian models for birch (Smith 2016; Smith 2014). These models provide biomass estimates for various tree biomass components: stem, bark, living branches, dead branches, foliage, stumps, and roots. These components are used to calculate above- and belowground biomass.

The dynamic soil model Yasso07 is used to calculate changes in carbon stock in dead organic matter and in soil for forest land remaining forest land (Tuomi et al. 2009; 2011). Estimates are made for individual NFI plots for the entire time-series. The Yasso07 model provides an aggregated estimate of carbon stock change for the total of litter, dead wood, and soil organic matter. All data used as input to the models is provided by the NFI. Auxiliary data used for estimation of C emissions from cropland, grassland, wetlands, and settlements were provided by Statistics Norway, Norwegian Meteorological Institute, as well as other data sources at the Norwegian Institute of Bioeconomy Research.

1.4.4 Data sources

The data sources used in the Norwegian inventory are outlined in the following:

Activity levels: These normally originate from official statistical sources available internally in Statistics Norway and other material available from external sources. When such information is not available, research reports are used or extrapolations are made from expert judgments.

Emission factors: These originate from reports on Norwegian conditions and are either estimated from measurements or elaborated in special investigations. However, international default data are used in cases where national emission factors are highly uncertain or lacking (e.g. N₂O from agriculture, CH₄ and N₂O from stationary combustion) or when the source is insignificant in relation to other sources.

Aggregated results from the side models: The operation of the side models in the inventory requires various sets of additional parameters pertinent to the emission source at hand. These data sets are as far as possible defined in official registers, public statistics and surveys, but some are based on assumptions.

Emission figures for point sources: For large industrial plants these are figures reported to the Norwegian Environment Agency by the plants' responsible (based on measurements or calculations at the plants).

1.5 Brief description of key categories

According to the IPCC definition, key categories are those that add up to 90 % of the total uncertainty in level and/or trend. In the Norwegian greenhouse gas emission inventory key categories are primarily identified by means of a Approach 2 method. A description of the methodology as well as background tables and the results from the analyses is presented in Annex 1. In this chapter a summary of the analysis and the results are described.

According to the IPCC Good Practice Guidance (IPCC 2000) it is good practice to give the results at the Approach 2 level if available. The advantage of using an Approach 2 methodology is that uncertainties are taken into account and the ranking shows where uncertainties can be reduced. However, in the 2006 IPCC guidelines it is suggested that good practice reporting should include key categories from both Approach 1 and Approach 2.

The Approach 2 and Approach 1 analyses were performed at the level of IPCC source categories and each greenhouse gas from each source category was considered separately with respect to total GWP weighted emissions, except land-use, land-use change and forestry (LULUCF).

The results from the key category analyses are summarized in Table 1.1 (excluding LULUCF). In addition we have also included a category identified on qualitative criteria. Altogether there are 45 key categories. Key categories in LULUCF were identified in separate analyses and are summarized in Table 1.2.

The complete analyses are included in Annex 1 together with background data and the complete analysis including LULUCF.

The Approach 1 analysis included in the NIR uses a different aggregation level for some source categories than in the Approach 1 analysis generated in the CRF reporter.

Table 1.1 Summary of identified emission key categories. Excluding LULUCF.

IPCC Category Code	IPCC Category	Greenhouse gas	Identification criteria ¹				Method
1A1,1A2,1A4	Stationary combustion, Solid Fuels	CO ₂	L1	T1			Tier 2
1A1,1A2,1A4	Stationary combustion, Biomass	CH ₄	L1		L2		Tier 2
1A1,1A2,1A4	Stationary combustion, Gaseous Fuels	CO ₂	L1	T1	L2	T2	Tier 2
1A1,1A2,1A4	Stationary combustion, Gaseous Fuels	CH ₄			L2	T2	Tier 2
1A1,1A2,1A4	Stationary combustion, Liquid Fuels	CO ₂	L1	T1	L2	T2	Tier 2
1A1,1A2,1A4	Stationary combustion, Other Fuels	CO ₂	L1	T1	L2	T2	Tier 2
1A3a	Civil Aviation	CO ₂	L1	T1	L2	T2	Tier 3
1A3b	Road Transportation	CO ₂	L1	T1	L2	T2	Tier 1a
1A3d	Navigation	CO ₂	L1	T1	L2	T2	Tier 2
1A3d	Navigation	CH ₄		T1	L2	T2	Tier 2
1A4	Other Sectors, mobile combustion	CO ₂	L1	T1	L2		Tier 2
1A5b	Mobile	CO ₂	L1				Tier 2
1B1a	Coal Mining	CH ₄		T1	L2	T2	Tier 1
1B2a	Oil (incl. oil refineries, gasoline distribution)	CO ₂	L1	T1	L2	T2	Tier 2
1B2a	Oil (incl. oil refineries, gasoline distribution)	CH ₄	L1		L2		Tier 2
1B2b	Natural Gas	CH ₄			T2		Tier 2
1B2c	Venting and Flaring	CO ₂	L1	T1	L2	T2	Tier 2
1B2c	Venting and Flaring	CH ₄	L1	T1	L2	T2	Tier 2

National Inventory Report 2019 - Norway

IPCC Category Code	IPCC Category	Greenhouse gas	Identification criteria ¹	Method
1C	CO ₂ Transport and storage	CO ₂	Q	CS (Tier 2)
2A1	Cement Production	CO ₂	L1 T1	Tier 3
2A2	Lime Production	CO ₂	L1 T1	Tier 3
2B1	Ammonia Production	CO ₂	L1 T1	Tier 2
2B2	Nitric Acid Production	N ₂ O	L1 T1 L2 T2	Tier 3
2B5	Carbide Production	CO ₂	L1 T1 T2	Tier 2
2B6	Titanium dioxide production	CO ₂	L1	Tier 2
2C2	Ferroalloys production	CO ₂	L1 L2	Tier 2/3
2C3	Aluminium production	CO ₂	L1 T1 L2 T2	Tier 2/3
2C3	Aluminium production	PFCs	L1 T1 L2 T2	Tier 2
2C4	Magnesium production	SF ₆	L1	Tier 2
2D1	Lubricant use	CO ₂	T1	Tier 2
2F	Product uses as substitutes for ODS	HFCs	L1 T1 L2 T2	Tier 2
3A	Enteric Fermentation	CH ₄	L1 T1 L2 T2	Tier 1/2
3B1	Manure management - Cattle	CH ₄	L1	Tier 2
3B	Manure Management	N ₂ O	L2 T2	Tier 2
3Da1	Direct emissions from managed soils -Inorganic N fertilizers	N ₂ O	L1 L2 T2	Tier 1
3Da2	Direct emissions from managed soils - Organic N fertilizer	N ₂ O	L1 L2 T2	Tier 1
3Da3	Direct emissions from managed soils - Urine and dung deposited by grazing animals	N ₂ O	L1 L2	Tier 1
3Da4	Direct emissions from managed soils - Crop residues	N ₂ O	L2 T2	Tier 1
3Da6	Direct emissions from managed soils - Cultivation of organic soils	N ₂ O	L1 L2	Tier 1
3Db1	Indirect emissions from managed soils - Atmospheric deposition	N ₂ O	L2	Tier 1
3Db2	Indirect emissions from managed soils - Nitrogen leaching and run-off	N ₂ O	L2	Tier 1
3G	Liming	CO ₂	L1 T1	Tier 1
5A1a	Managed Waste Disposal sites. Anaerobic.	CH ₄	L1 T1 L2 T2	Tier 2
5B	Biological treatment of Solid Waste	CH ₄	T2	Tier 1
5D	Wastewater treatment and discharge	CH ₄	L2 T2	Tier 1

¹ "L" refers to level and "T" to trend analyses. Numbers refer to approaches. "Q" refers to qualitative criteria.

Both the Approach 1 level analysis and trend analysis in Table 1.1 identified one new source each. The level analysis identified CH₄ from Stationary combustion of biomass (1A1,1A2,1A4) while the trend analysis identified CO₂ from Cement production (2A1). In addition the level analysis removed CO₂ in Petrochemical and carbon black production (2B8) as a source as well as N₂O in Manure management (3B). The Approach 1 trend analysis also removed CO₂ in Mobile (1A5B) and CO₂ in Petrochemical and carbon black production (2B8) as sources.

The Approach 2 trend analysis (excluding LULUCF) for 2017 identified one new source and the removal of two sources. The new source is N₂O from Manure management (3B) and the sources removed are CO₂ in Other sectors, mobile combustion (1A4) and N₂O in Biological treatment of solid waste (5B).

In total the analysis does not include any additions of new categories, but it has resulted in the removal of two. The categories removed are Biological treatment of solid waste for N₂O (5B), and Petrochemical and carbon black production for CO₂ (2B8).

From the LULUCF analyses, 27 key categories were identified by both the Approach 1 and Approach 2 level analyses (Table 1.2). There were no new key categories added this year, but the three sources that were no longer categorized as key categories are living biomass on grassland converted to forest land, mineral soils on cropland converted to settlement, and CH₄ emissions related to drained organic soils from cropland.

Table 1.2 Summary of identified LULUCF key categories.

IPCC Category Code	IPCC Category	Greenhouse gas	Identification criteria ¹	Method
4(II)Forest	Forest land drained organic soils - Drained organic soil	CH ₄	L2	Tier 1
4(II)Forest	Forest rem forest- drained organic soils (SSB) - Drained organic soil	N ₂ O	L1 T1 L2 T2	Tier 1
4(III)	Direct N ₂ O from N mineralization/immobilization - Mineralization/immobilization	N ₂ O	L2 T2	Tier 1
4.A.1	Forest remaining forest - Litter + dead wood + Mineral soil	CO ₂	L1 T1 L2 T2	Tier 3
4.A.1	Forest remaining forest - Living biomass	CO ₂	L1 T1 L2 T2	Tier 3
4.A.1	Forest remaining forest, drained organic soils - Organic soil	CO ₂	L1 T1 L2 T2	Tier 1
4.A.2.1	Cropland to Forest - DOM	CO ₂	L2 T2	Tier 2
4.A.2.2	Grassland to Forest - DOM	CO ₂	T1 L2 T2	Tier 2
4.A.2.2	Grassland to Forest - Mineral soil	CO ₂	T2	Tier 2
4.A.2.4	Settlements to Forest - DOM	CO ₂	L1 T1 L2 T2	Tier 2
4.B.1	Cropland remaining cropland - Organic soil	CO ₂	L1 T1 L2 T2	Tier 1
4.B.2.1	Forest to Cropland - DOM	CO ₂	L1 T1 L2 T2	Tier 2
4.B.2.1	Forest to Cropland - Living biomass	CO ₂	L2 T2	Tier 3
4.B.2.1	Forest to Cropland - Mineral soil	CO ₂	T2	Tier 2
4.B.2.1	Forest to Cropland - Organic soil	CO ₂	L2 T2	Tier 1
4.B.2.3	Wetland to Cropland - Organic soil	CO ₂	L2	Tier 1
4.C.1	Grassland remaining grassland – Living biomass	CO ₂	L2	Tier 2
4.C.2.1	Forest to Grassland - DOM	CO ₂	L1 T1 L2 T2	Tier 2
4.C.2.1	Forest to Grassland - Living biomass	CO ₂	L2 T2	Tier 3
4.C.2.1	Forest to Grassland - Mineral soil	CO ₂	L1 T1 L2 T2	Tier 2
4.D.1	Wetland Peat extraction - on+off-site - Organic soil	CO ₂	L2	Tier 1 & Tier 2
4.E.1	Settlements remaining settlements - Organic soil	CO ₂	L1 T1 L2 T2	Tier 1
4.E.2.1	Forest to Settlement - DOM	CO ₂	L1 T1 L2 T2	Tier 2
4.E.2.1	Forest to Settlement - Living biomass	CO ₂	L1 T1 L2 T2	Tier 3
4.E.2.1	Forest to Settlement - Mineral soil	CO ₂	T1 L2 T2	Tier 2
4.E.2.1	Forest to Settlement - Organic soil	CO ₂	L1 L2	Tier 1
4.G	Harvested wood Products - HWP	CO ₂	L1 T1 L2 T2	Tier 2

¹ "L" refers to level and "T" to trend analyses. Numbers refer to approaches.

1.6 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals

1.6.1 Approach 1 uncertainty analysis

The uncertainties in the emission levels for 2017 have been investigated by an approach 1 analysis. The results are given in Table 1.3 and Table 1.4. Note that the figures may differ slightly from data elsewhere in the NIR because the uncertainty analysis was performed before the data were finalized.

Table 1.3 Approach 1 uncertainties in emission levels. Each gas and total GWP weighted emissions. Excluding the LULUCF sector. 2017.

2017	μ (mean)	Uncertainty 2σ (% of mean)
Total	52.7 mill. tonnes	3
CO ₂	43.8 mill. tonnes	3
CH ₄	5.0 mill. tonnes	14
N ₂ O	2.4 mill. tonnes	41
HFC	1.4 mill. tonnes	49
PFC	131 ktonnes	20
SF ₆	59 ktonnes	42

Table 1.4 Approach 1 uncertainties in emission levels. Each gas and total GWP weighted emissions. Including the LULUCF sector. 2017.

2017	μ (mean)	Uncertainty 2σ (% of mean)
Total	27.7 mill. tonnes	15
CO ₂	18.3 mill. tonnes	22
CH ₄	5.2 mill. tonnes	13
N ₂ O	2.7 mill. tonnes	37
HFC	1.4 mill. tonnes	50
PFC	131 ktonnes	21
SF ₆	59 ktonnes	42

1.6.2 Approach 2 uncertainty analysis

The uncertainty in the Norwegian greenhouse gas emission inventory has been investigated by an approach 2 analysis and the results are given in Table 1.5 to Table 1.8. The approach 2 analysis is also further described in Annex II.

The uncertainty in the Norwegian emission inventory was initially investigated systematically in three reports (SFT/Statistics Norway 1999, Statistics Norway 2000, Statistics Norway 2001c). The first two reports focused on the uncertainty in the greenhouse gas emissions, based on approach 2 analyses,

and the last report investigated the uncertainty in the emission estimates of long-range air pollutants. The analysis of greenhouse gases was repeated in Statistics Norway (2010) and more thoroughly in (Flugsrud & Hoem 2011). The report *Uncertainties in the Norwegian Greenhouse Gas Emission Inventory* (Rypdal & Zhang 2000) includes more detailed documentation of the analysis method used in all analyses. Both approach 1 and 2 uncertainty analyses are now performed annually.

The national greenhouse gas (GHG) emission inventory is compiled from estimates based on emission factors and activity data and direct measurements by plants. All these data and parameters will contribute to the overall inventory uncertainty. The uncertainties and probability distributions of the inventory input parameters have been assessed based on available data and expert judgements. Finally, the level and trend uncertainties of the national GHG emission inventory have been estimated using Monte Carlo simulation. The methods used in the analysis correspond to an IPCC Approach 2 method, as described in the IPCC guidelines (IPCC 2006). Analyses have been made both excluding and including the sector LULUCF (Land Use, Land-Use Change and Forestry).

Table 6.2 from the IPCC good practice guidance is included in Annex II as Table AII-4. Column G in Table 6.2 is estimated as uncertainty for source category divided by total GHG emissions.

1.6.2.1 Uncertainty in emission levels

The estimated uncertainties of the levels of total emissions and in each gas are shown in Table 1.5 and Table 1.6.

Table 1.5 Uncertainties in emission levels. Each gas and total GWP weighted emissions. Excluding the LULUCF sector.

1990	μ (mean)	Fraction of total emissions	Uncertainty 2σ (% of mean)
Total	51.2 mill. tonnes	1	4
CO ₂	35.3 mill. tonnes	0.69	3
CH ₄	5.8 mill. tonnes	0.11	16
N ₂ O	4.1 mill. tonnes	0.08	26
HFC	44 tonnes	0.00	50
PFC	3.9 mill. tonnes	0.08	20
SF ₆	2.1 mill. tonnes	0.04	1
2017	μ (mean)	Fraction of total emissions	Uncertainty 2σ (% of mean)
Total	52.7 mill. tonnes	1	3
CO ₂	43.8 mill. tonnes	0.83	3
CH ₄	5.0 mill. tonnes	0.10	14
N ₂ O	2.4 mill. tonnes	0.05	41
HFC	1.4 mill. tonnes	0.03	49

National Inventory Report 2019 - Norway

PFC	131 ktonnes	0.00	20
SF ₆	59 ktonnes	0.00	42

Table 1.6 Uncertainties in emission levels. Each gas and total GWP weighted emissions. Including the LULUCF sector.

1990	μ (mean)	Fraction of total emissions	Uncertainty 2σ (% of mean)
Total	41.2 mill. tonnes	1	6
CO ₂	25.0 mill. tonnes	0.61	8
CH ₄	5,9 mill. tonnes	0.14	16
N ₂ O	4.3 mill. tonnes	0.11	25
HFC	44 tonnes	0.00	50
PFC	3.9 mill. tonnes	0.09	20
SF ₆	2.1 mill. tonnes	0.05	1
2017	μ (mean)	Fraction of total emissions	Uncertainty 2σ (% of mean)
Total	27.7 mill. tonnes	1	15
CO ₂	18.3 mill. tonnes	0.66	22
CH ₄	5.2 mill. tonnes	0.19	13
N ₂ O	2.7 mill. tonnes	0.10	37
HFC	1.4 mill. tonnes	0.05	50
PFC	131 ktonnes	0.00	21
SF ₆	59 ktonnes	0.00	42

The total national emissions of GHG (LULUCF sector excluded) in 1990 are estimated with an uncertainty of 4 % of the mean. The main emission component CO₂ is known with an uncertainty of 3 % of the mean. The total uncertainty level was 3 % of the mean in 2017. There have been major changes in uncertainty level for the different emission components between the two years. The highest uncertainty change between 1990 and 2017 is in the uncertainty estimates for the SF₆ emissions, which has increased from 1 to 42 % of the mean. However, the SF₆ emissions are strongly reduced because magnesium production was closed down. The figures for the emission of SF₆ from magnesium production was quite well known, but now a larger part of the SF₆ emissions comes from sources with higher uncertainty. For N₂O there is also a considerable increase in the uncertainty between the years. One reason for the change can be found in that N₂O from the production of synthetic fertilizer with a quite low uncertainty contributes to a smaller part of the total N₂O emissions in 2017 than in 1990. For the other gases there are only smaller changes in the uncertainty from 1990 to 2017.

By including the LULUCF sector the results from the analysis show a total uncertainty of 6 % of the mean in 1990 and 15 % in 2017. This is due to the fact that the uncertainty in the LULUCF sector in general is higher than in most other sectors.

1.6.2.2 Uncertainty in emission trend

The estimated uncertainties of the trends of total emissions and each gas are shown in Table 1.7 and Table 1.8.

Table 1.7 Uncertainty of emission trends. 1990-2017. Excluding the LULUCF sector.

	% change $((\mu_{2017}-\mu_{1990})*100/\mu_{1990})$	Uncertainty $(2*\sigma*100/\mu_{1990})$
Total	3	3
CO ₂	24	3
CH ₄	-13	11
N ₂ O	-42	5
HFC ¹
PFC	-97	20
SF ₆	-97	1

Table 1.8 Uncertainty of emission trends. 1990-2017. Including the LULUCF sector.

	% change $((\mu_{2017}-\mu_{1990})*100/\mu_{1990})$	Uncertainty $(2*\sigma*100/\mu_{1990})$
Total	-33	7
CO ₂	-27	10
CH ₄	-13	11
N ₂ O	-37	5
HFC ¹
PFC	-97	20
SF ₆	-97	1

¹ The base year emissions of HFCs are so close to zero that figures for % change and uncertainty are meaningless.

The result shows that the increase in the total GHG emissions from 1990 to 2017 is 3 %, with an uncertainty in the trend of ± 3 percentage points, when the LULUCF sector is not included. This means that the 2017 emissions are likely between 0 and 6 % above the 1990 emissions (a 95 percent confidence interval).

With the sector LULUCF included in the calculations there has been a decrease in the total emissions figures on -33 %, with a trend uncertainty of ± 7 percentage points.

1.7 General assessment of completeness

An assessment of the completeness of the emission inventory should, according to the IPCC Good Practice Guidance, address the issues of spatial, temporal and sectoral coverage along with all underlying source categories and activities. Confidentiality is an additional element of relevance, which has been addressed in Section 1.2.3.6.

The inventory includes emissions on the archipelago Svalbard as well as on mainland Norway. In particular, emissions from coal mining on Svalbard is included.

The revised UNFCCC Reporting Guidelines on Annual Inventories as adopted by the COP by its Decision 24/CP.19 specifies that a Party may consider that a disproportionate amount of effort would be required to collect data for a gas from a specific category that would be insignificant in terms of the overall level and trend in national emissions and in such cases use the notation key NE. The Party should in the NIR provide justifications for exclusion in terms of the likely level of emissions. An emission should only be considered insignificant if the likely level of emissions is below 0.05 % of the national total GHG emissions (specified in a footnote to total GHG emissions without LULUCF for the latest reported inventory year) and does not exceed 500 kt CO₂-equivalents. The total national aggregate of estimated emissions for all gases and categories considered insignificant shall remain below 0.1 % of the national total GHG emissions.

Norway has used the emissions for 2017 as reported in this NIR as the basis for national total GHG emissions. The national total GHG emissions without LULUCF in 2017 is reported to be 52 712 543 tonnes CO₂-equivalents. The threshold for an individual emission to be considered insignificant is therefore 26 356 tonnes CO₂-equivalents while the total threshold to be considered insignificant is 52 713 tonnes CO₂-equivalents.

The emissions (excluding LULUCF) that Norway has considered as insignificant and their likely level of emissions are presented in Table 1.9. The individual emissions excluded are all below the individual threshold and the total emissions excluded are also below the total threshold. *Table 1.10* provide information on emissions reported as NE for the LULUCF-sector.

Table 1.9. Emissions considered insignificant and reported as NE (excluding LULUCF).

CRF code	Description of emission source	Gases	Likely level of emissions (tonnes CO ₂ -equivalents)
1C1a	CO ₂ transport and storage; Pipelines	CO ₂	Less than 200 tonnes. See chapter 3.5.
3A4, 3B4	Other animals: Enteric fermentation and manure management	CH ₄ , N ₂ O	See chapter 6.2. Includes ostrich, llama, etc. Emissions from ostrich were reported in previous submissions, and were less than 500 t CO ₂ -eq when population was highest. Other animals have smaller populations.
3D	Agricultural soils	CH ₄	No methodology, see note to CRF Table3s2.
5C2	Open burning of waste	CO ₂ , CH ₄ , N ₂ O	Order of 1200 t CO ₂ -eq. by estimate from 1999.
5D2	Wastewater treatment: Industrial wastewater	N ₂ O	Order of 1000 t CO ₂ -eq. by estimate in 2017. Emissions are estimated to be stable over the period 1990-2017.
	Total		Estimated emissions less than 3000 t CO ₂ -eq.

Source: Statistics Norway and Norwegian Environment Agency

For the LULUCF sector the notation key NE was used to report the following sources, either because they were non-mandatory or considered negligible.

Table 1.10. Emissions reported as NE for LULUCF.

CRF code	Description of emission source	Gases	Explanation
4D1	Wetlands – flooded land remaining flooded land	CO ₂	It is not mandatory because there is no default method for estimating carbon stock changes for this source.
4D2	Wetlands – land converted to peat extraction	CO ₂	Emissions from organic soils are not estimated because they are considered a negligible source.
4(II)	Emissions from drainage and rewetting	CO ₂ , CH ₄ and N ₂ O	It is not mandatory to estimate emissions from rewetting organic or mineral soils.
4(V)	Biomass burning – controlled burning- forest land remaining forest land land converted to forest land	CH ₄ and N ₂ O	We assume emission from controlled forest fires are negligible because very few fire drills are performed and a consistent time-series is not available.
4(V)	Biomass burning- wildfires – grassland	CH ₄ and N ₂ O	We assume emission from wildfires on grasslands are negligible because these very rarely occur.

Source: Norwegian Institute for Bioeconomy Research

2 Trends in Greenhouse Gas Emissions

2.1 Description and interpretation of emission trends for aggregated GHG emissions

In 2017, total greenhouse gas (GHG) emissions in Norway were 52.7 million tonnes of carbon dioxide equivalents, which is a decrease of 0.9 million tonnes compared to 2016². Between 1990 and 2017, the total GHG emissions increased by approximately 1.5 million tonnes, equivalent to an increase of 3 %. Emissions reached their peak at 57.0 million tonnes in 2007.

The net GHG emissions, including all sources and sinks, were 27.7 million tonnes of CO₂ equivalents in 2017. The total emissions distribution among the main IPCC sectors from 1990 to 2017 is illustrated in Figure 2.1.




Figure 2.1. Total emissions of greenhouse gases by sources and removals from LULUCF in Norway, 1990-2017 (Mtonnes CO₂ equivalents). Source: Statistics Norway/Norwegian Environment Agency/Norwegian Institute of Bioeconomy Research

Table 2.1 presents the total emissions including indirect CO₂ emissions³ and the distribution among the main CRF sectors from 1990 to 2017. Total indirect CO₂ emissions and net removal from LULUCF are also presented in this table.

² In this NIR, if not specified otherwise, total emission figures include indirect CO₂ emissions but not emissions and removals from land use, land-use change and forestry (LULUCF).

³ Non-CO₂, carbon-containing gases (methane (CH₄), CO or NMVOC) will eventually be oxidised to CO₂ in the atmosphere. The CO₂ emissions formed are termed "indirect CO₂ emissions".

Table 2.1. Total emissions of greenhouse gases by sources and removals in Norway 1990-2017. Million tonnes of CO₂ equivalents.

Year	Energy	Industrial processes and product use	Agriculture	LULUCF	Waste	Total with indirect CO ₂ and without LULUCF	Total with indirect CO ₂ and with LULUCF	Indirect CO ₂ emissions
1990	29.8	14.5	4.7	-10.0	2.2	51.2	41.2	0.6
1995	33.0	11.6	4.6	-13.8	2.1	51.3	37.5	0.9
2000	36.4	12.1	4.5	-24.4	1.8	54.8	30.4	1.0
2005	38.7	10.6	4.5	-25.1	1.6	55.4	30.2	0.5
2008	40.0	9.7	4.3	-27.7	1.5	55.6	27.9	0.4
2009	39.9	7.4	4.3	-30.3	1.5	53.2	22.8	0.3
2010	41.5	8.2	4.2	-26.5	1.5	55.5	29.0	0.3
2011	40.7	8.2	4.2	-28.0	1.5	54.6	26.7	0.3
2012	40.2	8.2	4.2	-24.5	1.5	54.1	29.6	0.3
2013	40.0	8.3	4.3	-25.2	1.4	54.0	28.8	0.3
2014	40.0	8.4	4.4	-24.1	1.4	54.1	30.1	0.4
2015	40.2	8.5	4.4	-23.2	1.3	54.5	31.2	0.4
2016	39.3	8.6	4.5	-23.8	1.3	53.6	29.8	0.3
2017	38.4	8.6	4.5	-25.0	1.2	52.7	27.7	0.3

Source: Statistics Norway/Norwegian Environment Agency/Norwegian Institute of Bioeconomy Research

Figure 2.2 illustrates the yearly evolution of GHG emissions from the IPCC sectors (not including LULUCF) in percentage change, relative to 1990.




Figure 2.2. Changes in emissions of greenhouse gases, relative to 1990, illustrated by UNFCCC source categories, 1990-2017. Index 1990 = 1. Source: Statistics Norway/Norwegian Environment Agency

Norway has experienced economic growth since 1990, generating a general growth in emissions. In addition, the offshore petroleum sector has expanded significantly for the past 20 years. Together this has resulted in higher CO₂ emissions from energy use, both in energy industries and transport. Looking at the overall trend from 1990 to 2017, emissions increased by 3 %.

The downward trend in GHG emissions from the industry sector can be explained, in the early 1990's, by the implementation of policies and measures in the metal industry, resulting in less emission intensive production methods and later in the 2000's by close-downs and production reductions mainly in the metal industry as well.

Emissions from agriculture have decreased by 4.8 % since 1990 due to reductions of activity in the agriculture sectors.

The downward trend in GHG emissions from the waste sector is due to reductions of waste amounts disposed at disposal sites.

In the next two chapters, emission trends are explained both by sectors and by gas for the period 1990-2017.

2.2 Description and interpretation of emission trends by sector

Figure 2.3 illustrates the 2017 distribution of Norwegian GHG emissions by IPCC classification of sources. The energy sector is by far the most important source of emissions, contributing to 72.9 % of the national GHG emissions.




Figure 2.3. Distribution of GHG emissions in Norway in 2017 by sector, excluding LULUCF.
Source: Statistics Norway/Norwegian Environment Agency

Figure 2.4 displays GHG emissions trends by sectors between 1990 and 2017. The Energy sector is divided into its five main sub-sectors: fuel combustion in energy industries, fuel combustion in manufacturing industries and construction, fuel combustion in transport, fuel combustion in other sectors⁴, and fugitive emissions from fuels.

While emissions have decreased for most of the sectors, emissions from energy industries and transport have significantly increased since 1990.

⁴ Includes CRF key categories 1A4 (stationary combustion in agriculture, forestry, fishing, commercial and institutional sectors and households, motorized equipment and snow scooters in agriculture and forestry, and ships and boats in fishing) and 1A5 (fuel used in stationary and mobile military activities).




Figure 2.4. Development of emissions of all GHG (Mtonnes CO₂ eq.) from the different sectors, excluding LULUCF, 1990-2017. Source: Statistics Norway/Norwegian Environment Agency

2.2.1 Energy

Figure 2.5 displays the distribution of GHG emissions in 2017 by the main sub-sectors within the energy sector.




Figure 2.5. Greenhouse gas emissions in 2017 from the energy sector distributed among the different source categories. Source: Statistics Norway/Norwegian Environment Agency

The major sources of emissions within the energy sector are energy industries and transport, contributed to 40.6 % and 32.5 % of emissions from the energy sector in 2017, respectively. The remaining emissions are nearly equally shared between the sub-sectors: energy use in manufacturing industries and construction (CRF 1A2), other fuel combustion (CRF 1A4 and 1A5) and fugitive emissions from fuels (CRF 1B).

The Norwegian electricity production is dominated by hydroelectric power. Thus, emissions from energy industries origin almost completely from fuel combustion in oil and gas extraction and related activities. Electricity is normally used in manufacturing processes and for heating purposes.

The total emissions of GHG from the energy sector over the period 1990-2017 are listed in Table 2.2.

Table 2.2. Total emissions of greenhouse gases (Mtonnes CO₂-eq.) from the energy sector in Norway, 1990-2017. CO₂ transport and storage is presented in ktonnes CO₂-eq

Year	Energy Industries	Energy in Manufacturing Industries and Construction	Transport	Other fuel combustion	Fugitive Emissions from Fuels	CO ₂ transport and storage	Total
1990	7.3	4.3	10.0	4.7	3.5	0	29.8
1995	9.1	4.7	10.9	4.6	3.6	0	33.0
2000	11.0	4.6	12.1	3.9	4.9	0.01	36.4
2005	13.5	4.3	13.3	3.9	3.7	0.01	38.7
2008	13.8	4.5	13.9	3.6	4.1	0.1	40.0
2009	14.8	4.1	13.7	3.9	3.4	0.1	39.9
2010	15.0	4.5	14.3	4.0	3.6	0.1	41.5
2011	14.6	4.3	14.6	3.5	3.6	0.1	40.7
2012	14.3	4.1	14.8	3.4	3.5	0.1	40.2
2013	14.4	4.2	14.7	3.2	3.6	0.03	40.0
2014	15.1	3.9	14.7	2.8	3.3	0.04	40.0
2015	15.6	4.0	14.3	2.8	3.5	0.04	40.2
2016	15.1	3.9	13.7	3.2	3.4	0.01	39.3
2017	15.6	4.0	12.5	3.1	3.2	0.01	38.4

Source: Statistics Norway/Norwegian Environment Agency

Emission changes from 1990 to 2017, relative to 1990, presented for various sub-sectors within the energy sector, are illustrated in Figure 2.6.




Figure 2.6. Changes in emissions of greenhouse gases, relative to 1990, for the various sub-sectors within the energy sector, 1990-2017. Index 1990 = 1. Source: Statistics Norway/Norwegian Environment Agency

The GHG emissions from the energy sector increased by 29.0 % from 1990 to 2017, primarily due to increased activity in oil and gas extraction and transport, specifically road transportation.

Since 2010, the energy sector's emissions decreases except for 2015. From 2016 to 2017, emissions decreased by 2.2 %.

Emissions from fuel combustion in **Energy Industries** were 114.0 % higher in 2017 than in 1990. They increased by 3.0 % from 2016 to 2017. The main emission source in the Energy Industries sub-sector, oil and gas extraction, has played an important role in the national economy in recent decades. On the offshore oil and gas installations, electricity and pumping power is principally produced by gas turbines, and to a lesser extent, diesel engines.

In 2017, emissions from energy use in oil and gas extraction contributed to 24.4 % of the national GHG emissions. In 1990, the corresponding contribution was 11.6 %. The growth can be explained by the increase of oil and gas production and the increase of energy demand in extraction, due to aging of oil fields and transition from oil to gas.

Electricity production is largely dominated by hydroelectric generation. Between 1990 and 2017, important exceptions are gas fired electricity power plants, waste incineration power plants and a small coal combustion plant (6 MW) on the island of Spitsbergen.

Industrial emissions related to fuel combustion⁵ originate to a large extent from the production of raw materials and semi-manufactured goods, e.g. alloys, petrochemicals, paper and minerals.

⁵ Includes mainly emissions from use of oil or gas for heating purposes. Does not include consumption of coal as feedstock and reduction medium, which is included in the industrial process category.

Emissions from **Manufacturing Industries and Construction** have remained relatively stable since 1990, with a small decrease of 0.2 Mtonnes CO₂eq from 1990 to 2017. Between 2016 and 2017, emissions have increased by 3.0 %.

Emissions from **Transport** showed an overall increase of 24.2 % from 1990 to 2017, with a decrease of 8.9 % from 2016 to 2017. The highest emissions from transport since 1990 was 14.8 million tonnes in 2012. The share of transport in the total GHG emissions has increased from 19.6 % in 1990 to 23.7 % in 2017. In 2017, road transportation accounts for 70.3 % of emissions from the transport sub-sector, while emissions from navigation and civil aviation accounts for 20.4 and 8.9 %, respectively. Due to the fact that most railways are electrified in Norway, emissions of GHG from this source are insignificant.

GHG emissions from road transportation increased by 22.1 % from 1990 to 2017 and contributed to 16.6 % of the national GHG emissions in 2017. This trend is mainly due to the increase of activity in goods transport, as a response to higher economic activity. From 2016 to 2017, emissions decreased by 9.5 %. In addition to a reduced activity, the decreased emissions observed since 2007 could for the first years after be explained by the switch from petrol to diesel driven personal cars, due to the implementation of a CO₂ differentiated tax in 2007. However, in the later years a blending requirement of biofuels have increased consumption of bio diesel and bio ethanol and hence reduced CO₂ emissions. In addition, the sales of electric vehicles have gradually increased since 2011, and added up to 21% of personal cars and 2% of light duty vehicles in 2017, due to economic incentives.

Navigation contributed to the national total GHG emissions by 4.8 % in 2017. Emissions from navigation increased mainly due to an increase of activity related to the oil and gas extraction sector. Since the year 2012, the emissions have been reduced by 33.0 %. During the period 1990-2017, emissions have increased by 23.7 %.

Civil aviation contributed to 2.1 % of the national GHG emissions in 2017. Emissions from civil aviation have increased by 58.4 % since 1990, but the substitution of older planes by new and more energy efficient planes has played an important role to limit the emission growth.

GHG emission trends from the main transport modes are illustrated in Figure 2.7 and Table 2.3.

National Inventory Report 2019 - Norway




Figure 2.7. Emissions in million tonnes CO₂ equivalents from the most important modes of transport, 1990-2017. Source: Statistics Norway/Norwegian Environment Agency

Table 2.3. Total emissions of greenhouse gases from the transport sector in Norway, 1990-2017. Million tonnes CO₂ equivalents.

Year	Civil Aviation	Road transportation	Railways	Navigation	Total Transport
1990	0.70	7.18	0.10	2.05	10.04
1995	0.89	7.27	0.11	2.67	10.95
2000	1.09	8.14	0.07	2.80	12.10
2005	0.97	9.29	0.06	2.94	13.26
2008	1.11	9.67	0.06	3.03	13.87
2009	1.10	9.52	0.05	3.05	13.72
2010	1.07	9.76	0.07	3.41	14.31
2011	1.14	9.71	0.07	3.69	14.61
2012	1.16	9.75	0.07	3.79	14.77
2013	1.14	9.76	0.06	3.72	14.68
2014	1.26	9.91	0.05	3.50	14.71
2015	1.18	9.94	0.05	3.11	14.27
2016	1.10	9.69	0.05	2.85	13.70
2017	1.11	8.77	0.05	2.54	12.47

Source: Statistics Norway/Norwegian Environment Agency

The sub-sector “**Other fuel combustion**” (Table 2.2) includes, in particular, fuel combustion in agriculture, forestry and fisheries, residential sector and commercial/institutional sectors (CRF categories 1A4). The total emissions from CRF sector 1A4 were 2.9 million tonnes of CO₂ equivalents

in 2017. Emissions decreased by 35.6 % from 1990 to 2017, and decreased by 3.0 % from 2016 to 2017.

In 2017, GHG emissions from residential sources (stationary and mobile) accounted for 31.7 % of emissions from the “other fuel combustion” sub-sector. Emissions from the residential subsector have been reduced by 48.3 % since 1990, mainly due to the electrification of heating infrastructures. However, temperature variations and changes in electricity prices have at times reversed this trend.

Emissions from stationary combustion in the residential subsector are climate-dependent. Indeed, mild winter can lead to relatively lower consumption of fuels and thus reduced emissions. Whereas dry and cold winter can lead to relatively higher emissions. Since 1990, emissions from stationary combustion in the residential sector have decreased by 69.7 %, mainly due to decreased energy consumption.

Emissions from commercial/institutional sources (mobile and stationary) have decreased by 12.8 % since 1990. Emissions from commercial/institutional stationary sources decreased by 53.9 % from 1990 to 2017, whereas emissions from mobile sources increased by 85.7 % between 1990 and 2017.

The sub-sector “**Fugitive emissions from fuels**” in Table 2.2 refers to emissions from oil and gas activities such as flaring of natural gas, leakages and venting of methane. Indirect CO₂ emissions from NMVOC emitted during the loading and unloading of oil tankers are also accounted for in this sub-sector. Fugitive emissions from fuels contributed to 6.1 % of the national GHG emissions in 2017 and to 8.4 % of the GHG emissions within the energy sector. Fugitive emissions from fuels have decreased by 6.8 % since 1990 with a decrease of 3.8 % between 2016 and 2017.

The reduced emissions from flaring since 1990 are partly explained by the introduction of tax on gas flared off shore from 1991 and implemented technical measures. The amount of gas flared may fluctuate from year to year due to variation of startups, maintenance and interruption in operation.

2.2.2 Industrial processes and product use

The industrial processes and other product use (IPPU) sector accounted for 16.4 % of the national GHG emissions in 2017. The emissions from this sector decreased by 40.5 % from 1990 to 2017. Emissions remained stable between 2016 and 2017.

Metal Industry is the main source of emissions within the IPPU sector in the period 1990-2017. It contributed indeed to 56.1 % of the GHG emissions from the IPPU sector in 2017. The other main contributing sources in 2017 were Product uses as substitutes for ODS, Mineral Industry and Chemical Industry. They contributed to 16.3, 11.9 and 11.2 % of the GHG emissions from the IPPU sector, respectively.

Figure 2.8 shows the variations in the contribution to GHG emissions from 1990 to 2017 of the different IPPU sub-sectors. Table 2.4 provides figures for the total GHG emissions from the IPPU sector for the same period.




Figure 2.8. Total greenhouse gas emissions (Mtonnes CO₂-eq.) in the IPPU sub-sector in Norway, 1990-2017⁶. Source: Statistics Norway/Norwegian Environment Agency

During the first half of the 20th century, a large-scale industrialization took place in Norway. Many industrial communities appeared around the large hydroelectric resources particularly in the western parts of the country. Typical products were raw materials and semi-manufactured goods such as aluminium and ferroalloys. The main energy source has always been hydroelectricity. However, fossil fuels have been used as reducing agents or raw materials. Greenhouse gases are then emitted as process related gases.

9.2 % of national GHG emissions came from **Metal industry** in 2017, whose emissions increased by 0.7 % from 2016 to 2017.

The largest contributors to the GHG emissions from Metal Production in 2017 are productions of ferroalloys and aluminium. Emissions from those productions constituted more than 97 % of emissions from Metal industry in 2017. The large decrease in emissions in 2009 reflects low production levels of ferroalloys, due to lower economic activity and economic recession.

In 1990, PFCs emissions from aluminium production contributed to 7.6 % of the national GHG emissions, while in 2017, it has been reduced to 0.2 %. Emissions of PFCs have decreased by 96.6 % since 1990 and by 29.7 % between 2016 and 2017.

Since 2010, production of ferroalloys has been the most important source of GHG emissions within the metal production category. The GHG emissions from ferroalloys production amounted to 2.6 million tonnes of CO₂ equivalents in 2017 and accounted for 5.0 % of the national total GHG emissions. Emissions from production of ferroalloys increased by 3.3 % from 1990 to 2017, with a 1.1

⁶ Under Other production, Norway reports the two source categories: pulp and paper and food and drink.

% increase from 2016. The large increase in emissions from 2009 to 2010 (50.2 %) is due to a low production level in 2009. The production level in 2009 is also lower than 2008 and reflects the lower economic activity due to the economic recession.

Table 2.4. Total greenhouse gas emissions (Mtonnes CO₂-eq.) from the IPPU sub-sectors in Norway, 1990-2017.

Year	Mineral industry	Chemical Industry	Metal industry	Non-energy products from fuels and solvent use	Electronics Industry	Product uses as ODS substitutes	Other product manufacture and use	Other	Total
1990	0.7	3.3	10.1	0.3	0.0	0.00004	0.1	0.03	14.5
1995	1.0	2.8	7.3	0.2	0.001	0.1	0.1	0.05	11.6
2000	1.0	2.9	7.4	0.2	0.001	0.4	0.2	0.1	12.1
2005	0.9	2.8	5.9	0.2	0.001	0.6	0.1	0.1	10.6
2008	1.0	2.0	5.5	0.2	0.001	0.8	0.1	0.1	9.7
2009	1.0	1.3	3.8	0.2	0.001	0.9	0.1	0.1	7.4
2010	1.0	1.4	4.3	0.2	0.001	1.1	0.1	0.1	8.2
2011	1.0	1.3	4.4	0.2	0.001	1.1	0.1	0.1	8.2
2012	1.0	1.3	4.4	0.2	0.001	1.1	0.1	0.1	8.2
2013	1.1	1.2	4.5	0.2	0.001	1.2	0.1	0.1	8.3
2014	1.1	1.1	4.7	0.2	0.001	1.2	0.1	0.1	8.4
2015	1.0	1.2	4.6	0.2	0.001	1.2	0.1	0.1	8.5
2016	1.0	1.1	4.8	0.2	0.001	1.4	0.1	0.1	8.6
2017	1.0	1.0	4.8	0.2	0.001	1.4	0.1	0.1	8.6

Source: Statistics Norway/Norwegian Environment Agency

In 1990, SF₆ from magnesium foundries accounted for 4.0 % of the national total GHG emissions. Emissions decreased until the closure of all plants in 2007. Reductions in SF₆ emissions over the period are, in the early 90s, mainly due to improvements in the production processes, in 2002, due to the closing down of production of cast magnesium and in 2006, due to the closing down of secondary magnesium production.

Emissions from **Mineral Industry** were 1.0 million tonnes in 2017, which accounted for 1.9 % of the national GHG emissions. Emissions increased by 40.9 % from 1990 to 2017, mainly due to the increase of clinker and lime production in more recent years. Emissions from Mineral industry increased by 5.5 % from 2016 to 2017.

In 2017, the CO₂ process emissions from cement production were 1.5 % of the national GHG emissions. They have increased by 20.7 % since 1990, due to increased production of clinker, with an increase of 11.9 % from 2016 to 2017.

The Chemical Industry includes primarily N₂O from nitric acid production and CO₂ from production of ammonia and carbides. The GHG emissions from this sub-sector amounted to 1.0 million tonnes of CO₂ equivalents in 2017, which represented 1.8 % of the national GHG emissions. Emissions have decreased by 70.2 % since 1990, mainly due to the reduction of emissions from the production of nitric acid, ammonia and carbides. Emissions have decreased by 9.8 % since 2016 mainly due to the decrease of ammonia production.

2.2.3 Agriculture

In 2017, 8.5 % of the national GHG emissions originated from agriculture, corresponding to 4.5 million tonnes of CO₂ equivalents. Emissions from agriculture have decreased by 4.8 % since 1990 and increased by 0.2 % since 2016.

The largest sources of GHGs within the agriculture sector are “enteric fermentation” (CH₄) and “agricultural soils” (N₂O). In 2017, these sub-sectors represented 52.1 % and 36.3 % of the agriculture sector, respectively, while “manure management” represented 9.6 %.

The main driver behind the emission trend in agriculture is the development in the number of animals for the significant animal groups. The main reasons for the decreasing trend in GHG emissions are the reduction of nitrogen content in the synthetic fertilizers used, use of more concentrate and more effective milk production which led to reduction of the number of dairy cows.

Enteric fermentation contributed to 2.3 million tonnes of CO₂ equivalents in 2017, corresponding to 4.4 % of the national GHG emissions. This sub-sector constituted almost 90 % of the overall CH₄ emissions from agriculture for the period 1990-2017.

The emissions of N₂O from **agricultural soils** amounted to 1.6 million tonnes of CO₂ equivalents in 2017. This accounted for 67.7 % of the national N₂O emissions in 2017 and 3.2 % of the national GHG emissions.

In 2017, emissions CH₄ and emissions of N₂O from **manure management** amounted to 0.3 million and 0.1 million tonnes of CO₂ equivalents, respectively. This accounted for 0.8 % of the Norwegian GHG emissions.

Table 2.5. Greenhouse gas emissions (Mtonnes CO₂-eq.) from the agricultural sub-sectors in Norway, 1990-2017. Urea application is in ktonnes CO₂-eq.

Year	Enteric Fermentation	Manure Management	Agricultural Soils	Field burning of agricultural residues	Liming	Urea application	Total
1990	2.4	0.3	1.7	0.04	0.2	0.001	4.8
1995	2.4	0.3	1.6	0.02	0.2	0.001	4.7
2000	2.3	0.3	1.6	0.01	0.1	0.0001	4.6
2005	2.3	0.4	1.6	0.01	0.1	0.0001	4.5
2008	2.3	0.4	1.6	0.01	0.1	0.001	4.5
2009	2.3	0.4	1.6	0.004	0.1	0.001	4.5
2010	2.3	0.4	1.5	0.004	0.1	0.0003	4.4
2011	2.2	0.4	1.6	0.003	0.1	0.0003	4.4
2012	2.2	0.4	1.6	0.003	0.1	0.0002	4.3
2013	2.2	0.4	1.6	0.003	0.1	0.0002	4.3
2014	2.2	0.4	1.6	0.004	0.1	0.0002	4.3
2015	2.3	0.4	1.6	0.004	0.1	0.0002	4.4
2016	2.3	0.4	1.6	0.004	0.1	0.0002	4.5
2017	2.3	0.4	1.6	0.004	0.1	0.0001	4.5

Source: Statistics Norway/Norwegian Environment Agency




Figure 2.9. Total greenhouse gas emissions (Mtonnes CO₂-eq.) in the agriculture sub-sectors in Norway, 1990-2017. Source: Statistics Norway/Norwegian Environment Agency

2.2.4 Land Use, Land-Use Change and Forestry (LULUCF) and KP-LULUCF

The LULUCF sector differs from the other sectors in that it can function as both a source of atmospheric emissions and a sink of emissions through the removal of atmospheric CO₂. The balance of the two is net emissions or removals in the LULUCF sector.

In 2017, the net removal in the LULUCF sector was 24.9 million tonnes CO₂ equivalents, which correspond to almost half of the national GHG emissions that year. The average annual net sequestration from the LULUCF sector has been 22.3 million tonnes CO₂ equivalents per year for the period 1990-2017.

The calculated changes in carbon depend upon several factors such as growing conditions, harvest levels, management practices and land use changes.

The area distribution of the land-use categories for Norway in 1990 and 2017 is illustrated in Figure 2.10. The figure shows that the net changes in land-area distribution in Norway from 1990 to 2017 have been relatively small; only the area of settlements has increased slightly, while the other land-use categories have decreased. Details on gross changes between the respective categories may be found in the land transition matrix, Table 6.2 in the LULUCF chapter.




Figure 2.10 Area (%) distribution between the IPCC land-use categories, 1990 and 2017.

Source: The Norwegian Institute of Bioeconomy Research

Figure 2.11 illustrates net emissions and removals of CO₂-eq by land use-category. As can be seen, all land-use categories other than forest land and harvested wood products showed net emissions in 2017. In total, the emissions were calculated to about 4.1 million tonnes of CO₂ equivalents, of which the main emissions came from the land-use categories cropland and settlements. Emissions from settlements increased by more than two times from 1990 to 2017, and are, in 2017, responsible for the second largest emissions from the LULUCF sector, with 2.1 million tonnes of CO₂ equivalents.

Forest land was the major contributor to the net sequestration of CO₂ in the sector. In 2017, the total net removals from forest land were 29.1 million tonnes of CO₂. Within this category, land converted to forest land contributed with 0.51 million tonnes of CO₂.

The figure clearly shows that the net removals from forest land has increased from 1990 to 2017. During this time period, the total net sequestration of CO₂ from forest land increased by 152 %. The explanation for this growth is an increase in standing volume and gross increment, while the amount of CO₂ emissions due to harvesting and natural losses has been quite stable. The increase in living carbon stock is due to an active forest management policy over the last 60–70 years. The combination of the policy to re-build the country after the Second World War II and the demand for timber led to a great effort to invest in forest tree planting in new areas. These areas are now at their most productive age and contribute to the increase in living biomass and hence the carbon stock.




Figure 2.11 Net CO₂ emissions and removals (kt CO₂-equivalents per year) from the LULUCF sector by land-use category (forest land, cropland, grassland, wetlands, settlements, other land, and harvested wood products) from 1990 to 2017, including emissions of N₂O and CH₄.

Source: Norwegian Institute of Bioeconomy Research

In chapter 11, supplementary information on Norway's commitment to report on and account for emissions and removals from Land Use, Land-Use Change and Forestry under the Kyoto Protocol (KP-LULUCF) is provided. All emissions and removals are estimated according to the 2013 Kyoto Protocol supplement (IPCC 2014a).

Reporting on activities under Article 3.3 (Afforestation/reforestation and Deforestation) and forest management (Article 3.4) is mandatory for all Parties under the Kyoto Protocol. In addition, any activity elected in the first commitment period (2008-2012) is mandatory in the second commitment

period (2013-2020). For the second commitment period, Norway has also elected the voluntary activities Cropland Management and Grazing Land Management in the accounting under Article 3.4.

Areas where afforestation and reforestation and deforestation activities have occurred in Norway are small compared to the area of forest management. As illustrated in Table 2.6, estimated C sequestration for the activity forest management is substantial, whereas net emissions occur from deforestation, cropland and grazing land management. In addition, C sequestration from afforestation/reforestation is estimated.

Table 2.6. CO₂, N₂O and CH₄ emissions (kt CO₂ eq yr⁻¹) and CO₂ removals of all pools for Article 3.3 and 3.4 under the Kyoto Protocol for the base year (1990) and for each of the first five years of the second commitment period.

Year	Net emissions (kt CO ₂ -eq yr ⁻¹)				
	Afforestation/ reforestation	Deforestation	Forest management	Cropland management	Grazing land management
1990			-12482.45	1786.98	-73.19
2013	-569.07	2279.38	-29118.20	1769.69	16.15
2014	-561.63	2158.26	-27923.20	1776.96	11.64
2015	-532.84	2157.39	-27035.01	1776.20	10.04
2016	-505.87	2230.14	-27719.40	1773.53	10.20
2017	-489.06	2218.46	-28914.54	1763.53	12.04

Source: Norwegian Institute of Bioeconomy Research

The accounting of emissions and removals from LULUCF towards Norway's commitment under the Kyoto protocol will be in accordance with Decision 2/CMP.7. The final quantity of emissions and removals for each year of the commitment period to be accounted towards Norway's commitment will be determined at the end of the commitment period, i.e. in 2022, when emissions and removals for the year 2020 have been reported. Until the year of accounting, emissions and removals from the Kyoto Protocol activities may be recalculated due to changes in activity data and/or methodology.

Preliminary accounting quantities from land use, land-use change and forestry for the first five years of the second commitment period under the Kyoto Protocol indicate that Norway will have a net removals slightly above 0.25 million tonnes of CO₂- equivalents in total for these five years. The preliminary accounting quantities from the activities, calculated according to Decision 2/CMP.7, comprise emissions (million tonnes of CO₂-equivalents) of 11 from deforestation and 0.4 from grazing land management; and removals of 2.6 from afforestation and reforestation, 9 from forest management and 0.07 from cropland management.

2.2.5 Waste

The waste sector, with emissions of 1.2 million tonnes of CO₂ equivalents in 2017, accounted for 2.3 % of the national GHG emissions.

This sector includes emissions from landfills (CH₄), wastewater handling (CH₄ and N₂O), biological treatment of solid waste and small-scale waste incineration (CO₂ and CH₄). Waste incineration with utilization of energy is included in the Energy sector.

Solid waste disposal on land (landfills) is the main sub-sector within the waste sector. It accounted for 81.5 % of the sector's total emissions in 2017. Whereas wastewater handling accounted for 12.2 % and biological treatment of solid waste for 6.2 %. Small-scale waste incineration accounted for 0.1 %.

GHG emissions from the waste sector have generally decreased since 1990. In 2017, emissions were 46.5 % lower than in 1990 and 4.6 % lower than in 2016. The total amount of waste generated increased by more than 60.4 % from 1995 to 2016, but due to the increase in material recycling and a ban against disposing biodegradable waste to landfills, methane emissions have decreased leading to a decrease in total emissions of greenhouse gases from the waste sector.

The distribution of the waste emissions by sub-sector is presented in Figure 2.12 and Table 2.7.




Figure 2.12. Total emissions of greenhouse gases (Mtonnes CO₂-eq.) in Norway from the waste sub-sectors, 1990-2017. Source: Statistics Norway/Norwegian Environment Agency

Table 2.7 shows the decrease of methane emissions (landfills) since 1990. The reduction is due to a smaller amount of waste disposed at disposal sites. With a few exceptions, it was then prohibited to dispose easy degradable organic waste at landfills in Norway. In 1999, a tax was introduced on waste delivered to final disposal sites. Since July 2009, it is banned to deposit biodegradable waste to landfills. This results in further reduction of methane emissions.

Table 2.7. Emissions (Mtonnes CO₂-eq.) from the waste sub-sector in Norway, 1990-2017. Incineration and open burning of waste is presented in ktonnes CO₂-eq.

Year	Solid waste disposal	Biological treatment of solid waste	Incineration and open burning of waste	Wastewater treatment and discharge	Total
1990	2.1	0.0	0.0002	0.2	2.2
1995	1.9	0.0	0.001	0.2	2.1
2000	1.6	0.0	0.001	0.1	1.8
2005	1.4	0.1	0.001	0.1	1.6
2008	1.3	0.1	0.001	0.1	1.5
2009	1.3	0.1	0.001	0.1	1.5
2010	1.3	0.1	0.001	0.1	1.5
2011	1.3	0.1	0.001	0.1	1.5
2012	1.2	0.1	0.006	0.1	1.5
2013	1.2	0.1	0.001	0.1	1.4
2014	1.2	0.1	0.001	0.1	1.4
2015	1.1	0.1	0.001	0.1	1.3
2016	1.0	0.1	0.001	0.1	1.3
2017	1.0	0.1	0.001	0.1	1.2

Source: Statistics Norway/Norwegian Environment Agency

2.3 Description and interpretation of emission trends by gas

As shown in Figure 2.13, CO₂ is by far the largest contributor to the total GHG emissions, followed by CH₄, N₂O, and then the fluorinated gases PFCs, SF₆ and HFCs. In 2017, the relative contributions to the national total from the different gases were: CO₂ 82.9 %, CH₄ 9.5 %, N₂O 4.5 % and fluorocarbons (PFCs, SF₆ and HFCs) 3.0 %. The relative share of the gases has been quite stable since 2010, the relative share of CO₂ has increased by approximately 1 % each year during the period 2005-2010, from 79.4 % in 2005 up to 83.3 % in 2010.




Figure 2.13. Distribution of emissions of greenhouse gases in Norway by gas, 2017.

Source: Statistics Norway/Norwegian Environment Agency

Table 2.8 presents emission figures for all greenhouse gases, expressed in absolute emission figures and total CO₂ equivalents.

Table 2.8. Emissions of greenhouse gases in Norway, 1990-2017. Units: CO₂ in Mtonnes (Mt), CH₄ and N₂O in ktonnes (kt) and other gases in ktonnes CO₂ eq. (kt CO₂ eq.).

Gas	CO ₂	CH ₄	N ₂ O	PFC	SF ₆	HFC
Year	Mt	kt		kt CO ₂ eq		
1990	35.3	232.0	13.7	3894.8	2098.5	0.04
1995	38.7	235.3	12.4	2314.0	579.8	92.0
2000	42.5	227.9	12.8	1518.5	891.4	383.3
2005	44.0	219.2	13.7	955.3	296.1	614.3
2008	45.4	213.1	10.5	896.0	59.8	806.1
2009	43.9	214.5	8.6	438.3	55.7	856.1
2010	46.2	215.2	8.3	238.4	68.6	1064.5
2011	45.5	208.9	8.3	262.6	54.3	1105.8
2012	45.0	207.3	8.4	200.5	53.5	1140.8
2013	44.9	208.4	8.3	181.0	56.3	1155.2
2014	44.9	212.0	8.3	178.9	50.1	1235.6
2015	45.3	207.6	8.4	146.4	69.8	1232.9
2016	44.5	203.7	8.2	186.2	63.6	1363.6
2017	43.7	200.9	8.0	131.0	58.8	1402.8

Source: Statistics Norway/Norwegian Environment Agency

Table 2.9 presents the emissions in million tonnes per greenhouse gas and the changes in % for each greenhouse gas for the period 1990–2017, and for 2016-2017.

Table 2.9. Emissions in Mtonnes CO₂ equivalents and changes in % for each greenhouse gas.

Year	CO ₂	CH ₄	N ₂ O	PFCs	SF ₆	HFCs	Total
1990	35.3	5.8	4.1	3.9	2.1	0.00004	51.2
2016	44.5	5.1	2.4	0.2	0.1	1.4	53.6
2017	43.7	5.0	2.4	0.1	0.1	1.4	52.7
Changes 1990-2017	23.7 %	-13.4 %	-41.5 %	-96.6 %	-97.2 %	3187970.5 %	2.9 %
Changes 2016-2017	-1.7 %	-1.4 %	-1.8 %	-29.7 %	-7.6 %	2.9 %	-1.7 %

Source: Statistics Norway/Norwegian Environment Agency

As presented in Table 2.8 and Table 2.9, CO₂ emissions increased significantly from 1990 to 2017 with 8.4 million tonnes. Emissions of CH₄ and N₂O decreased by 0.8 and 1.7 million tonnes CO₂ equivalents, respectively. During the same period, PFCs and SF₆ emissions significantly decreased with 3.8 and 2.0 million tonnes CO₂ equivalents, respectively, while HFCs has increased from almost 0 to 1.4 million tonnes CO₂ equivalents.

The fluorocarbons constituted a larger fraction of the GHG emission total in the early 1990s than in 2017, while CO₂ represented a smaller share in 1990 than in 2017.

The Figure 2.14 illustrates the changes in % for the different greenhouse gases for the period 1990 to 2017.




Figure 2.14. Changes in emissions of greenhouse gases, relative to 1990, by gas, 1990-2017. Index 1990 = 1. Source: Statistics Norway/Norwegian Environment Agency

Figure 2.14 shows that the overall increasing total emission trend of CO₂ has been weakened by decreased emissions of fluorinated gases due to SF₆ and PFCs emissions reduction.

2.3.1 Carbon dioxide (CO₂)

The Norwegian CO₂ emissions originate primarily from energy industries, transport and industrial processes. Since generation of electricity is almost exclusively hydroelectric, emissions from stationary combustion are dominated by industrial sources and internal energy use.

The distribution of CO₂ emissions among various categories is shown in Figure 2.15.

National Inventory Report 2019 - Norway




Figure 2.15. Distribution of CO₂ emissions in Norway by various source categories in 2017.

Source: Statistics Norway/Norwegian Environment Agency

Table 2.10 lists CO₂ emissions from each source category for the period 1990-2017. The variations in the contribution of CO₂ emissions 1990 to 2017 are displayed in Figure 2.16.

Table 2.10. CO₂ emissions (million tonnes) from different source categories, 1990-2017.

Year	Energy Industries	Manufacturing Industries and Construction	Transport	Other Fuel Combustion	Fugitive emissions from fuel and CO ₂ transport and storage	Industrial processes and product use	Other sources	Total
1990	7.2	4.2	9.9	4.5	2.9	6.4	0.2	35.3
1995	9.0	4.7	10.8	4.4	2.8	6.9	0.2	38.7
2000	10.9	4.5	11.9	3.6	3.9	7.5	0.1	42.5
2005	13.4	4.3	13.1	3.6	2.7	6.7	0.1	44.0
2008	13.7	4.5	13.7	3.3	3.2	6.9	0.1	45.4
2009	14.6	4.1	13.5	3.6	2.6	5.4	0.1	43.9
2010	14.9	4.4	14.1	3.8	2.8	6.3	0.1	46.2
2011	14.5	4.3	14.4	3.2	2.7	6.3	0.1	45.5
2012	14.2	4.1	14.5	3.1	2.7	6.3	0.1	45.0
2013	14.2	4.2	14.4	2.9	2.7	6.4	0.1	44.9
2014	15.0	3.9	14.4	2.6	2.3	6.5	0.1	44.9
2015	15.4	4.0	14.0	2.6	2.6	6.6	0.1	45.3
2016	15.0	3.9	13.4	3.0	2.5	6.7	0.1	44.5
2017	15.4	4.0	12.2	2.9	2.4	6.7	0.1	43.7

Source: Statistics Norway/Norwegian Environment Agency

Since 1990, the total emissions of CO₂ have increased by 23.7 %, or by 8.4 million tonnes. The increases of natural gas use in gas turbines in the oil and gas extraction industry have been the most important contributor to the overall CO₂ increase.

In 2017, the total Norwegian emissions of CO₂ were 43.7 million tonnes. It has decreased by 1.7 % or 0.8 million tonnes since 2016.




Figure 2.16. CO₂ emissions (Mtonnes) in Norway, 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

CO₂ emissions from energy industries have increased by 113.8 % since 1990 as a result of large increases in production volume of oil and gas and the export of natural gas in pipelines. In 2017, emissions from energy industries increased by 0.5 million tonnes or 3.0 % compared to 2016.

CO₂ emissions from transport have increased by 23.8 % since 1990 and contributed to 28.0 % of the total CO₂ emissions in 2017. CO₂ emissions from this sector is dominated by road transportation which accounted for 70.8 % of the CO₂ emissions from transport in 2017. CO₂ emissions from road transportation increased by 22.9 % between 1990 and 2017 although emissions from personal cars powered by gasoline decreased by 59.0 % during this period. CO₂ emissions from road transportation decreased by 9.6 % between 2016 and 2017.

CO₂ emissions from industrial processes have increased by 4.9 % since 1990, and contributed to 15.4 % of total CO₂ emissions in 2017. Metal production accounted for 70.0 % of the CO₂ emissions from industrial processes in 2017.

2.3.2 Methane (CH₄)

In 2017, 52.0 % of methane emissions originated from agriculture, 26.0 % from the energy sector and 21.6 % originated from the waste sector. Methane emissions are dominated by releases from enteric fermentation in the agriculture sector, by fugitive emissions from oil and gas extraction in the energy sector and by landfills in the waste sector.

Figure 2.17 illustrates the distribution of Norwegian CH₄ emissions in 2017.




Figure 2.17. Distribution of Norwegian CH₄ emissions by major sources in 2017.

Source: Statistics Norway/Norwegian Environment Agency

The methane figures from 1990 to 2017, distributed among the different categories are displayed in Table 2.11.

Table 2.11. Emissions of CH₄ (ktonnes) in Norway, 1990-2017.

Years	Energy	Enteric fermentation	Other sources in Agriculture	Waste	Other Sources	Total
1990	37	97	11	87	0.5	232
1995	47	95	11	82	0.5	235
2000	54	94	10	70	0.8	228
2005	56	93	10	59	0.8	219
2008	56	90	10	57	0.8	213
2009	55	91	10	57	0.7	214
2010	57	91	10	56	0.8	215
2011	54	88	10	55	0.8	209
2012	54	88	10	54	0.8	207
2013	55	89	11	53	0.8	208
2014	60	90	11	51	0.8	212
2015	57	91	11	48	0.8	208
2016	54	92	11	46	0.7	204
2017	52	93	11	43	0.7	201

Source: Statistics Norway/Norwegian Environment Agency

The total methane emissions decreased by 1.4 % from 2016 to 2017. Since 1990, CH₄ emissions have decreased by 13.4 %. Table 2.11 and Figure 2.18 show that this decrease is primarily due to the decrease of emissions from waste treatment, which more than compensated the growth in the energy sector, specially in the oil and gas industry.

The waste volumes have grown during the period 1990-2017, but this effect has been more than offset by the increase of recycling, incineration of waste and burning of methane from landfills.




Figure 2.18. CH₄ emissions (ktonnes) for major Norwegian sources, 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

2.3.3 Nitrous oxide (N₂O)

Figure 2.19 shows that, in 2017, 73.9 % of the Norwegian N₂O emissions are of agricultural origin, agricultural soils being the most prominent contributor within the agriculture sector. Industrial processes is the second contributor, with 12.7 %. Nitric acid production is the main source of N₂O emissions within industrial processes accounted for more than 69 % of the sector.

The energy sector accounted for 8.6 % and the waste sector for 4.8 %. Emissions are dominated by road transport in the energy sector and by waste water treatment and discharge in the waste sector.




Figure 2.19. Distribution of Norwegian N₂O emissions by major sources in 2017.

Source: Statistics Norway/Norwegian Environment Agency

Reductions of emissions during the period 1990-2017 are mainly due to decreased emissions from nitric acid production. Changes in the production processes of nitric acid led to the decrease of N₂O emissions, first in the beginning of the 1990s, and then since 2006. Technological improvements in the production process have significantly brought the emissions down during the last ten years.

During the period 1990–2017 the total N₂O emissions decreased by 41.5 %. From 2016 to 2017, emissions decreased by 1.8 %. Details are presented in Table 2.12 and Figure 2.20.

National Inventory Report 2019 - Norway

Table 2.12. Emissions of N₂O (ktonnes) in Norway by major sources, 1990-2017.

Years	Agriculture	Energy	Industrial processes and product use	Waste	Total
1990	0.6	7.0	6.0	0.20	13.7
1995	0.7	5.7	5.9	0.24	12.4
2000	0.7	6.0	5.9	0.27	12.8
2005	0.6	6.8	5.9	0.32	13.7
2008	0.6	3.6	5.9	0.36	10.5
2009	0.6	2.0	5.7	0.35	8.6
2010	0.7	1.8	5.5	0.35	8.3
2011	0.7	1.6	5.7	0.34	8.3
2012	0.7	1.6	5.7	0.37	8.4
2013	0.7	1.4	5.8	0.39	8.3
2014	0.7	1.3	5.9	0.36	8.3
2015	0.7	1.4	6.0	0.36	8.4
2016	0.7	1.1	6.0	0.38	8.2
2017	0.7	1.0	5.9	0.38	8.0

Source: Statistics Norway/Norwegian Environment Agency




Figure 2.20. N₂O emissions for major Norwegian sources, 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

2.3.4 Perfluorocarbons (PFCs)

Aluminium production is the only source of PFC emissions in 2017. In 2017, perfluorocarbons tetrafluoromethane (CF₄) and hexafluoroethane (C₂F₆) emissions from Norwegian aluminium plants were reported at 14.9 and 1.7 tonnes respectively, corresponding to a total of 0.13 million tonnes of CO₂ equivalents. Total PFCs total emissions have decreased by 96.6 % since 1990 following a steady downward trend as illustrated in Figure 2.21. Since 1990, emissions of CF₄ have decreased by 96.8 %, while the emission of C₂F₆ have decreased by 95.3 %.

Improvement of technology and process control in aluminium production led to a significant emissions decrease. In 1990, PFCs emissions were 4.48 tonne CO₂ equivalents per tonne aluminium produced. It was reduced to 0.70 tonne CO₂ equivalents per tonne aluminium produced in 2007 and to 0.10 tonne CO₂ equivalents per tonne aluminium produced in 2017. Total PFCs emissions decreased by 0.06 million tonnes between 2016 and 2017.




Figure 2.21. Emissions (million tonnes CO₂-eq) of PFCs in Norway, 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

Table 2.13. Emissions of PFCs in Norway in 1990-2017 in tonnes. PFC218 is in kg and total is in million tonnes of CO₂-eq.

Year	PFC14 (CF ₄) (t)	PFC116 (C ₂ F ₆) (t)	PFC218 (C ₃ F ₈) (kg)	Total CO ₂ eq. (Mt)
1990	467.4	36.2	0.0	3.89
1995	283.3	18.1	0.0	2.31
2000	186.4	11.6	0.0	1.52
2005	116.7	7.6	0.0	0.96
2008	104.7	10.1	0.0	0.90
2009	49.8	5.8	0.0	0.44
2010	27.4	3.0	7.4	0.24
2011	29.9	3.4	7.3	0.26
2012	22.9	2.6	6.4	0.20
2013	20.7	2.3	2.1	0.18
2014	20.3	2.4	0.2	0.18
2015	16.7	1.9	0.0	0.15
2016	21.2	2.5	0.0	0.19
2017	14.9	1.7	0.0	0.13

Source: Statistics Norway/Norwegian Environment Agency

2.3.5 Sulphur hexafluoride (SF₆)

Until 2006, the largest source of SF₆ emissions in Norway was magnesium production. The consumption of SF₆ was reduced through the 1990s due to improvements in technology and process management, and to reductions in production levels. In 2017, the SF₆ emissions were 97.2 % lower than in 1990. Until 2002, SF₆ emission reductions were mainly due to the improved technology and process control within the metal industries. In 2002, production of cast magnesium closed down and production of secondary magnesium closed down in 2006.


The main other use of SF₆ is in gas insulated switchgears (GIS) and other high-voltage applications. Since the signing of a voluntary agreement in 2002, emissions from these sources have decreased.

The increase in emissions in 2015 was due to decommissioning of sound insulated windows that was produced in Norway in the 1980ies and 90ies.

Table 2.14. SF_6 emissions (tonnes), in Norway 1990-2017.

Year	GIS	Magnesium and Aluminium Industry	Other	Total
1990	2.2	89.7	0.1	92.0
1995	3.6	21.3	0.5	25.4
2000	4.5	32.4	2.3	39.1
2005	2.3	10.0	0.6	13.0
2008	2.1	0.0	0.6	2.6
2009	1.9	0.0	0.6	2.4
2010	2.5	0.0	0.6	3.0
2011	2.0	0.0	0.4	2.4
2012	1.9	0.0	0.4	2.3
2013	2.0	0.0	0.4	2.5
2014	1.8	0.0	0.4	2.2
2015	1.8	0.0	1.3	3.1
2016	1.5	0.0	1.3	2.8
2017	1.2	0.0	1.4	2.6

Source: Statistics Norway/Norwegian Environment Agency


Figure 2.22. Emissions of SF_6 (tonnes) in Norway 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

2.3.6 Hydrofluorcarbons (HFCs)

The total actual emissions from HFCs used as substitutes for ozone depleting substances amounted to 1.4 million tonnes of CO₂ equivalents in 2017. It is an increase of 2.9 % compared to 2016. The emissions in 1990 were insignificant. Indeed, emissions have been multiplied by more than 15 since 1995.

The application category refrigeration and air conditioning contributes by far to the largest part of the HFCs emissions. The other categories foam/foam blowing and fire extinguishing contributes to small amounts to the overall emissions.

Figure 2.23 displays the development of HFCs emissions since 1990 and Table 2.15 presents HFCs emission values for different HFCs from 1990 to 2017. The trend is due to the strong demand for substitution of ozone depleting substances. The increase in HFCs emissions has been moderated by the introduction of a tax on HFCs in 2003.

Table 2.15. Emissions of HFCs (tonnes), unspecified mix of HFCs and total (Mtonnes CO₂-eq.) in Norway, 1990-2017.

Year	HFC23	HFC32	HFC125	HFC134a	HFC143a	HFC152a	HFC227ea	HFC134	HFC143	Unspecified mix of HFCs Mtonnes CO ₂ eq.	Total in Mtonnes CO ₂ eq.
1990	0.0	0.0	0.0	0.0	0.0	0.4	NO	NO	NO	0.0	0.0
1995	0.0	0.4	5.2	36.8	4.1	1.3	NO	NO	NO	0.003	0.1
2000	0.1	2.0	34.5	77.0	28.7	1.8	NO	NO	NO	0.02	0.4
2005	0.2	6.1	56.1	123.1	44.8	4.4	NO	0.8	1.1	0.03	0.6
2008	0.1	12.5	68.1	202.6	52.0	0.9	NO	2.7	1.4	0.03	0.8
2009	0.1	15.9	73.1	229.3	50.4	0.9	NO	2.2	1.3	0.03	0.9
2010	0.1	19.8	93.5	262.7	69.3	4.0	NO	2.0	1.1	0.03	1.1
2011	0.2	22.6	98.2	278.4	65.0	1.7	NO	1.8	1.0	0.1	1.1
2012	0.5	25.5	97.5	303.5	60.6	2.3	0.1	1.7	0.9	0.1	1.1
2013	0.4	31.1	95.8	327.7	57.4	1.5	0.2	1.5	0.8	0.1	1.2
2014	0.3	34.6	100.4	335.3	69.4	1.1	0.2	1.4	0.8	0.1	1.2
2015	0.3	39.5	110.6	324.6	66.9	0.6	0.2	1.3	0.7	0.05	1.2
2016	0.3	42.4	119.0	357.5	78.0	3.0	0.2	1.2	0.6	0.1	1.4
2017	0.3	44.7	120.7	361.0	79.4	0.1	0.1	1.1	0.5	0.1	1.4

Source: Statistics Norway/Norwegian Environment Agency

National Inventory Report 2019 - Norway




Figure 2.23. Emissions of HFCs (Mtonnes CO₂-eq.) in Norway, 1990-2017.

Source: Statistics Norway/Norwegian Environment Agency

2.4 Emission trends for indirect greenhouse gases and SO₂

Nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOC) and carbon monoxide (CO) are not greenhouse gases but have an indirect effect on the climate through their influence on greenhouse gases and in particular ozone. Sulphur dioxide (SO₂) also has an indirect impact on climate, as it increases the level of aerosols with a subsequent cooling effect. Therefore, emission trends of these gases are to some extent included in the inventory.

The overall **NO_x emissions** decreased with approximately 19.9 % from 1990 to 2017. This can primarily be explained by stricter emission regulations with regard to road traffic, which has led to a 35.0 % reduction of emissions from transport since 1990. These reductions counteracted increased emissions from e.g. oil and gas production. From 2016 to 2017, the total NO_x emissions decreased by 4.3 %, mainly due to reduction in the transport and stationary combustion.

NMVOC emissions experienced an increase in the period from 1990 to 2001, mainly because of the rise in oil production. However, NMVOC emissions decreased by 63.3 % from 2001 to 2017, and were, in 2017, 52.0 % lower than in 1990. This decrease has been achieved through the implementation of measures to increase the recycling of oil vapour offshore at loading and storage terminals. From 2016 to 2017, the emissions of NMVOC decreased by 2.8 %.

CO emissions have decreased by 50.2 % over the period 1990-2017. This is explained primarily by the implementation of new emission standards for motor vehicles.

SO₂ emissions were reduced by 69.9 % from 1990 to 2017. This can mainly be explained by a reduction in sulphur content of all oil products and lower process emissions from ferroalloys and aluminium productions as well as refineries.




Figure 2.24. Emissions (ktonnes) of NO_x, NMVOC, SO₂, and CO in Norway, 1990-2017.

Source: Statistics Norway/ Norwegian Environment Agency

3 Energy (CRF sector 1)

3.1 Overview of sector

The Energy sector, including fugitive emissions, accounted for 73 % of the Norwegian greenhouse gas emissions in 2017. In 1990, the Energy sector's share of the total greenhouse gas emissions was 58 %.

Road traffic and offshore gas turbines (electricity generation and pumping of natural gas in pipelines) are the sector's largest single contributors to the sector's emissions and the latter is the sector that has increased the most since 1990. Other important sources in the Energy sector are coastal navigation, energy use in the production of raw materials, as well as oil and gas operations, which also give rise to significant amounts of fugitive emissions.

GHG emissions in the Energy sector have increased by 29 % from 1990 to 2017 (Figure 3.2), primarily due to increased activity in the sectors of oil and gas extraction and transport, specifically road transport. Between 1990 and 2017, there have been temporary emission reductions in the sector. Among these temporary reductions, emissions decreased by 3 % from 2007 to 2009 and by 4 % from 2010 to 2014. The former decrease is partly due to the fact that a new gas terminal started up in 2007 but had start-up problems during the first years and was only fully operational in 2009. The financial crisis also contributed to lower emissions in 2009.

The growth in emissions from 2009 to 2010 was mainly due to increased emissions from gas fired power plant and district heating. The latter due to the increase of fuel oils used during one of the coldest winters since the 1950s. The emission reduction from 2010 to 2014 was mainly due to reversed trends in the same sector. In 2016 and 2017, increased use of biofuels in road transport also contributed to the reduction in total emissions.

Figure 3.1 and Figure 3.2 show the trend and the relative changes to 1990, in GHG emissions for the different Energy sectors. The main emitting sectors are the energy industries sector (combustion in oil and gas production, refineries, electricity production and district heating) and the transport sector (civil aviation, road transportation, railways, navigation). Both sectors have increased since 1990, especially the energy industries sector, which has more than doubled since 1990.

The manufacturing industries and construction sector, the other fuel combustion sector⁷ and the fugitive emissions from fuel sector experienced small fluctuations between 1990 and 2017. In 2017, emissions from the manufacturing industries sector and from the fugitive sector are almost as they were in 1990. On the other hand, the other fuel combustion sector underwent a decrease of 35 % between 1990 and 2017.

⁷ Other fuel combustion sector includes both the sectors Other Combustion (CRF 1A4) and Other (CRF 1A5)




Figure 3.1. Greenhouse gas emissions from energy sectors and fugitive emissions. 1990-2017. Million tonne CO₂ equivalents. Source: Statistics Norway/Norwegian Environment Agency




Figure 3.2. Relative changes to 1990 in GHG emissions for the energy sector including fugitive emissions. 1990-2017. Index 1990=1. Source: Statistics Norway/Norwegian Environment Agency

Transport

In 2017, the transport sector's total GHG emissions was 12.5 million tonnes CO₂ equivalents of which road transportation contributed to 70 %, navigation to 20 %, civil aviation to 9 % and railways to less than 1 %.

Figure 3.3 and Figure 3.4 show the trend and the relative changes of transport emissions from 1990 to 2017. They show that emissions from road transportation, navigation and aviation have increased during the period, while emissions from railways have decreased. Since 1990, emissions from civil aviation have increased by 58, navigation by 24 % and road transportation by 22 %. Emissions from road transport have decreased the last couple of years, mainly due to use of biofuels. Emissions from railways have decreased by 53 % compared with 1990, mainly due to railways electrification.




Figure 3.3. Greenhouse gas emissions from the most important transport sectors. 1990-2017. Million tonnes CO₂ equivalents. Source: Statistics Norway/ Norwegian Environment Agency




Figure 3.4. Relative changes to 1990 in GHG emissions for the most important transport sectors. Civil aviation, road transportation, navigation and other transportation. 1990-2017. Index 1990=1.

Source: Statistics Norway/Norwegian Environment Agency

Key source categories

Section 1.5 describes the overall results of the approach 2 key category analysis performed for the years 1990 and 2017. Table 3.1 gives the key categories in the energy sector in terms of total level and/or trend uncertainty for 1990 and/or 2017 in CRF order.

Table 3.1 Key categories in the Energy sector in 2017.

CRF code	Source Category	Gas	Key category according to approach	Method
1A1,1A2,1A4	Stationary combustion, Solid Fuels	CO ₂	1	Tier 2
1A1,1A2,1A4	Stationary combustion, Liquid Fuels	CO ₂	2	Tier 2
1A1,1A2,1A4	Stationary combustion, Gaseous Fuels	CO ₂	2	Tier 2
1A1,1A2,1A4	Stationary combustion, Gaseous Fuels	CH ₄	2	Tier 2
1A1,1A2,1A4	Stationary combustion, Other Fuels	CO ₂	2	Tier 2
1A1,1A2,1A4	Stationary combustion, Biomass	CH ₄	2	Tier 2
1A3a	Civil Aviation	CO ₂	2	Tier 3
1A3b	Road Transportation	CO ₂	2	Tier 1a
1A3d	Navigation	CO ₂	2	Tier 2
1A3d	Navigation	CH ₄	2	Tier 2
1A4	Other sectors - Mobile Fuel Combustion	CO ₂	2	Tier 2
1A5b	Other – Mobile	CO ₂	1	Tier 2
1B1a	Coal Mining and Handling	CH ₄	2	CS, Tier 2
1B2a	Fugitive emissions from oil	CO ₂	2	Tier 2
1B2a	Fugitive emissions from oil	CH ₄	2	Tier 2
1B2b	Fugitive emissions from natural gas	CH ₄	2	CS, Tier 2
1B2c	Venting and Flaring	CO ₂	2	Tier 2
1B2c	Venting and Flaring	CH ₄	2	Tier 2
1C	CO ₂ transport and storage	CO ₂	Qualitative	CS, Tier 2

Source: Statistics Norway/Norwegian Environment Agency

In addition to source categories defined as key categories according to the approach 2 key category analysis, two source categories are defined as key according to approach 1 key category analysis. They are CO₂ Stationary combustion, solid fuels (1A) and CO₂ from Military, mobile (1A5b).

An important issue, which is also elaborated in this sector, concerns the capture and storage of CO₂ emissions at the offshore oil and gas field Sleipner Vest and Hammerfest LNG (Snøhvit gas-condensate field). These unique operations are discussed in detail in section 3.5.

Emission allocation

Generally, energy combustion for energy purposes is reported in 1.A Fuel Combustion Activities, while flaring and other fugitive emissions are reported in 1.B Fugitive Emissions from Fuels. Emissions from waste incineration at district heating plants are accounted for under the energy sector, as the energy is utilized. Methane from landfills and other biogas used for energy purposes are also accounted for in this sector. Emissions from flaring in the energy sectors are reported in 1.B.2c Flaring and described in section 3.4, as this energy combustion is not for energy purposes. Emissions

from burn off of coke at catalysts in refineries are reported in 1.B.2.a iv for the same reason as for flaring. Coal and coke used as reducing agents and gas used for production of ammonia (non-energy part) are accounted for in sector 2 IPPU (chapter 4). Flaring in manufacturing industries is also reported in 2 IPPU. Flaring of landfill gas and other biogas is reported in sector 5 Waste (chapter 7). The same applies to emissions from accidental fires, etc. Emissions from burning of crop residues and agricultural waste are accounted for in sector 3 Agriculture (chapter 5).

A more detailed description of the delimitation of energy combustion is given in section 3.2.1.1.

Mode of presentation

The elaboration of the energy sector in the following starts with a general description of emissions from the energy combustion sources (section 3.2). Then followed by a description of fugitive emissions (sections 3.3 and 3.4) and a discussion on the capture and storage of CO₂ emissions at the oil and gas field Sleipner Vest and Hammerfest LNG (Snøhvit gas-condensate field) (section 3.5).

Cross-cutting issues are elaborated in section 3.6 and comprise the following elements:

- Comparison between the sectoral and reference approach
- Feedstock and non-energy use of fuels
- Indirect CO₂ emissions from CH₄ and NMVOC

Finally, the memo items of international bunker fuels and CO₂ emissions from biomass are addressed in section 3.7.

In the case of energy combustion, emissions from the individual combustion sources are discussed after a comprehensive presentation of the energy combustion sector as a whole (section 3.2). The purpose for such an arrangement is to avoid repetition of methodological issues which are common among underlying source categories, and to enable easier cross-reference.

3.2 Energy Combustion

3.2.1 Overview

This section describes the general methodology for calculation of GHG emissions from the combustion of fossil fuels and biomass. All known combustion activities within energy utilisation in various industries and private households are included.

The GHG emissions from fuel combustion (1A) accounted for 67 % of national total emissions in 2017. The emissions increased by 34 % between 1990 and 2017. The increase is primarily due to activity growth in oil and gas extraction, which comprises the major part of energy industries sector, and in transport, mainly road transport.

Emissions from source category 1A decreased by 2 % from 2016 to 2017. In 2017, emissions from the sector energy industries (CRF 1A1) increased and the manufacturing and construction sector (CRF 1A2) increased by 3 %. The transport sector (CRF 1A3) decreased by 9 %, primarily due to increased use of biofuels. Emissions in the other combustion sector (CRF 1A4 and CRF 1A5) decreased by 2 %.

The fuel combustion sector is dominated by emissions of CO₂, which in 2017 contributed 98 % to the totals of this sector (CRF 1A).

This sector hosts ten source categories defined as keys according to approach 2 key category analysis and two as key category from the approach 1 analysis, which, along with the non-key categories, are presented in detail in the following sections.

Table 3.3 presents the shares of estimated and reported emissions used in the inventory for the different sectors and for the different greenhouse gases in 2017. It shows that a large share of GHG emissions from Energy industries and Manufacturing Industries and Construction included in the Norwegian GHG Inventory are taken from annual reports sent by each plant to the Norwegian Environment Agency.⁸ Such annual reports are:

- reports as required by their regular permit
- reports as required by the permit under the EU emission trading system (EU ETS)
- reports as required by a voluntary agreement

Annex VIII QA/QC of point sources includes references to documents that in detail describe requirements for measuring and reporting, specifically for the EU ETS and the voluntary agreement.

3.2.1.1 Methodological issues

Emissions from fuel combustion are estimated at the sectoral level in accordance with the IPCC sectoral approach Tier 1/Tier 2/Tier 3. Total fuel consumption is, in many cases, more reliable than the breakdown to sectoral consumption.

The general methodology for estimating emissions from fuel combustion is multiplication of fuel consumption by source and sector by an appropriate emission factor, as shows in equation (3.1). Exceptions are road traffic and aviation, where more detailed estimation models are used; involving

⁸ Former Norwegian Pollution Control Authority and Climate and Pollution Agency

additional activity data (see sections 3.2.5 and 3.2.4, respectively). The total amounts of fuel consumption is taken from the Norwegian energy balance (see Annex III). The mean theoretical energy content of fuels and their density are listed in Table 3.2.

The general method for calculating emissions from energy consumption is:

$$(3.1) \quad Emissions(E) = Activity\ level(A) \times Emission\ Factor\ (EF)$$

Emissions of pollutants from major manufacturing plants (point sources) are available from measurements or other plant-specific calculations. When such measured data are available it is possible to replace the estimated values by the measured ones:

$$(3.2) \quad Emissions(E) = [(A - A_{PS}) \times EF] + E_{PS}$$

Where A_{PS} and E_{PS} are the activity and the measured emissions at the point sources, respectively. Emissions from activity for which no point source estimate is available ($A - A_{PS}$) are still estimated with the default emission factor. See section 1.4.2 for more information about the main emission model.

Table 3.2 Average energy content (NCV) and density of fuels*.

Energy product	Theoretical energy content	Unit	Density Tonne/m ³
Coal	28.1	GJ/tonne	:
Coke	28.5	GJ/tonne	:
Petrol coke	35	GJ/tonne	:
Crude oil	42.3	GJ/tonne	0.85
Motor gasoline	43.9	GJ/tonne	0.74
Aviation gasoline	43.9	GJ/tonne	0.74
Kerosene (heating)	43.1	GJ/tonne	0.81
Jet kerosene	43.1	GJ/tonne	0.81
Auto diesel	43.1	GJ/tonne	0.84
Marine gas oil/diesel	43.1	GJ/tonne	0.84
Light fuel oils	43.1	GJ/tonne	0.84
Heavy distillate	43.1	GJ/tonne	0.88
Heavy fuel oil	40.6	GJ/tonne	0.98
Bitumen	40.2	GJ/tonne	:
Lubricants	40.2	GJ/tonne	:
Natural gas (dry gas) (land)	35.5	GJ/1000 Sm ³	0.74 ¹
Natural gas (rich gas) (off shore)	40.3	GJ/1000 Sm ³	0.85 ¹
LPG	46.1	GJ/tonne	0.53
Refinery gas	48.6	GJ/tonne	:
Blast furnace gas	6.1-10.0	GJ/1000 Sm ³	:
Fuel gas ³	50	GJ/tonne	:
Landfill gas ^{2,4}	50.4	GJ/tonne	:
Biogas ^{2,4}	50.4	GJ/tonne	:
Fuel wood ²	16.8	GJ/tonne	0.5
Ethanol ²	26.8	GJ/tonne	0.79
Biodiesel ²	36.8	GJ/tonne	0.88
Wood waste ²	16.8	GJ/tonne	:
Black liquor ²	7.2 - 9.2	GJ/tonne	:
Wood pellets ²	17.3	GJ/tonne	:
Wood briquettes ²	15.5	GJ/tonne	:
Charcoal	29.5	GJ/tonne	:
Municipal waste	11.5	GJ/tonne	:
Special waste	40.6	GJ/tonne	0.98

* The theoretical energy content of a particular energy commodity may vary; Figures indicate mean values.

¹ kg/Sm³. Sm³ = standard cubic meter (at 15 °C and 1 atmospheric pressure).

² Non-fossil emissions, not included in the inventory CO₂ totals

³ In this inventory, fuel gas is a hydrogen-rich excess gas from petrochemical industry

⁴ Landfill gas and other types of biogas are reported as methane content in the energy balance

Source: Energy statistics, Statistics Norway and Norwegian Environment Agency

For offshore activities and some major manufacturing plants (in particular refineries, gas terminals, cement industry, production of plastics, ammonia production, and methanol production), emissions of one or more compounds reported by the plants to the Norwegian Environment Agency are used, as described in equation (3.2) (see Table 3.3). In these cases, the energy consumption of the plants in question is subtracted from the total energy use before the general method is used to calculate the remaining emissions of the compound in question, in order to prevent double counting.

Emissions are reported to the Norwegian Environment Agency under a number of different reporting obligations. Most of the CO₂ emissions (except metal production, etc.) are reported as part of the Emissions Trading System (ETS).

In the general equation (3.2), E_{PS} represents the reported emission data, while A_{PS} represents the energy consumption at the plants. Note that for most plants, reported emissions are used only for some of the substances. For the remaining substances in the inventory, the general method with standard emission factors is used.

Reported figures are used for a relatively small number of plants, but as they contribute to a large share of the total energy use, a major part of the total emissions are based on such reported figures. Table 3.3 gives an overview of the shares of estimated and reported emissions used in the inventory for the different sectors for the greenhouse gases CO₂, CH₄ and N₂O in 2017.

In 2017, 92 % of the CO₂ emissions from Energy Industries (oil and gas extraction and production, refineries, gas terminals, gas fired power plants and district heating plants) were based on reported emissions and 40 % of the CO₂ emissions from Manufacturing Industries and Construction.

Table 3.3. Share of total CO₂, CH₄ and N₂O emissions in the energy sector based on estimated and reported emission estimates for 2017.

	CO ₂		CH ₄		N ₂ O	
	Estimated	Reported	Estimated	Reported	Estimated	Reported
A. Fuel Combustion Activities (Sectoral Approach)	54 %	46 %	77 %	23 %	96 %	4 %
1. Energy Industries	8 %	92 %	16 %	84 %	80 %	20 %
a. Public Electricity and Heat Production	63 %	37 %	71 %	29 %	72 %	28 %
b. Petroleum Refining	0 %	100 %	66 %	34 %	100 %	0 %
c. Manufacture of Solid Fuels and Other Energy Industries	1 %	99 %	0 %	100 %	100 %	0 %
2. Manufacturing Industries and Construction	60 %	40 %	100 %	0 %	95 %	5 %
a. Iron and Steel	15 %	85 %	100 %	0 %	100 %	0 %
b. Non-Ferrous Metals	99 %	1 %	100 %	0 %	100 %	0 %
c. Chemicals	18 %	82 %	97 %	3 %	59 %	41 %
d. Pulp, Paper and Print	100 %	0 %	100 %	0 %	100 %	0 %
e. Food Processing, Beverages and Tobacco	100 %	0 %	100 %	0 %	100 %	0 %
f. Non-metallic minerals	32 %	68 %	100 %	0 %	100 %	0 %
g. Other (Oil drilling, construction, other manufacturing)	100 %	0 %	100 %	0 %	100 %	0 %
3. Transport	100 %		100 %		100 %	
a. Civil Aviation	100 %		100 %		100 %	
b. Road Transportation	100 %		100 %		100 %	
c. Railways	100 %		100 %		100 %	
d. Navigation	100 %		100 %		100 %	
e. Other Transportation (pipeline transport)	(IE)		(IE)		(IE)	
4. Other Sectors	100 %		100 %		100 %	
a. Commercial/Institutional	100 %		100 %		100 %	
b. Residential	100 %		100 %		100 %	
c. Agriculture/Forestry/Fisheries	100 %		100 %		100 %	
5. Other (Military)	100 %		100 %		100 %	

Source: Statistics Norway, Norwegian Environment Agency

Delimitation toward industrial processes etc.

The energy combustion sector borders to several other source categories. This section presents a more detailed description of the demarcation with other sectors used in the inventory, compared to section 3.1.

Energy consumption reported as activity data in the emission inventories is generally delimited in the same way as emissions. In cases where different substances are handled differently, the delimitation of energy consumption follows the delimitation of CO₂ emissions.

Flaring is not reported as energy use under energy combustion (CRF 1A). Instead, flaring is reported under the following source categories:

- Flaring in refineries and in exploration/extraction is reported under fugitive emissions (CRF 1B).

- Flaring in manufacturing industries is reported under industrial processes (CRF 2), particularly under chemical industry (CRF 2B). (In the energy balance, flaring in manufacturing is reported as "losses".)
- Flaring of landfill gas is reported under waste incineration (CRF 5C).

Emissions from *reducing agents* are reported under industrial processes (CRF 2). This contrasts with the delimitation in the energy balance, where use as reducing agents is reported as energy consumption.

In some special cases, CO₂ emissions from combustion are reported under other source categories, while emissions of other substances are reported in energy combustion (CRF 1A):

- CO-rich excess gas from metallurgical plants burnt on-site is reported under industrial processes (CRF 2), according to IPCC guidelines (IPCC 2006). (Gas which is sold to other plants is reported under energy combustion (CRF 1A)).
- Coal used as fuel in some metallurgical plants which also use coal as a reducing agent is reported under industrial processes (CRF 2).
- CO₂ from coke that is burnt off from catalytic crackers in refineries is reported under fugitive emissions (CRF 1B). This also applies to CO₂ from coke calcining kilns. This combustion is currently reported as energy use of CO₂-rich gas ("other gas") in the energy balance.

In these cases, energy consumption reported in the inventories follows the delimitation of the CO₂ emissions. This gives meaningful implied emission factors for CO₂, while IEFs for other substances may be skewed.

At a small number of plants, CO₂ emissions are reported in the ETS system from *derived fuels* which are not included as energy use in the energy balance. The carbon in the fuels is likely reported as feedstock in the energy balance. These cases are handled in two different ways. Both methods should give correct total CO₂ emissions, but the correspondence to reported energy data is different. In both cases, no emission of other substances from these fuels is currently estimated.

- For methanol production, CO₂ emissions from several fuels not included in the energy balance are reported as process emissions under Methanol production (CRF 2B8).
- In other cases, emissions from derived fuels are included in the total combustion CO₂ which is entered into the inventory for the plants. Thus, emissions are larger than the corresponding energy use reported in the inventory. As far as it is currently known, this method is only used when emissions from derived fuels are small relative to total fuel use in the source category, mainly in energy in manufacturing of chemicals (CRF 1A2c). The method leads to higher implied emission factors relative to standard range.

Emissions from *paraffin wax* are reported under Other Industrial processes (CRF 2G).

Combustion of *solid waste* and *hazardous waste* is reported under the energy section (district heating (CRF 1A1a) and in several manufacturing industries (CRF 1A2). No significant combustion of solid or hazardous waste occurs without energy recovery.

Combustion of *landfill gas* with energy recovery is reported under the energy section (mainly in Commercial/Institutional (CRF 1A4a)). Flaring is reported under waste incineration (CRF 5C), as mentioned above.

Emissions reported by plants: Energy data

Energy data for plants with reported emissions (A_{PS} in equation (3.2)) should be consistent both with the energy balance that is used for activity total A and with the reported emission data. Consistency with emission data means that the energy data should correspond to the same activity as the reported emissions.

In most cases, figures on plant energy use in the inventory are based on data reported from the plants to Statistics Norway. This ensures consistency with the energy balance.

In the emission trading system (ETS), emissions are, in most cases, reported together with data on the corresponding energy use. Usually, the energy data reported in the ETS are the same as those reported by the plants to Statistics Norway. However, for some plants, some of the energy data differ between reports to Statistics Norway and to the ETS. This leads to problems of consistency.

- In a few cases, the inventory uses plant energy data from the ETS instead of data from the energy balance of Statistics Norway. In these cases, the difference is significant, and the ETS data are deemed to be the most reliable. The emission inventory will be inconsistent with the energy balance. Currently, this applies to CO-rich excess gas in iron and steel production for 2008 and later.

In other cases, with mainly small emissions, the inconsistency between energy data from Statistics Norway (A_{PS}) and reported emissions data (E_{PS}) may lead to deviations in implied emission factors. However, the deviations are usually small, and generally, this should not be regarded as an important issue.

Emissions reported by plants: Allocation to combustion/processes

CH₄ emissions from an oil refinery are reported as a plant total, which includes both combustion and process emissions. These emissions have to be allocated to the two emission categories. Emissions from combustion are calculated from energy use with standard factors and the remaining part of reported emissions is reported in the inventory as process emissions.

Emissions reported by plants: Allocation to fuels

The following discussion is relevant for cases where emissions are reported with a fuels split. This applies to greenhouse gases reported to the UNFCCC, and to emission statistics in Statistics Norway's Statbank. In other reporting, emissions are aggregated over fuels.

For some plants and substances, emissions are reported by fuel, but in most cases reported combustion emissions are entered as a plant total. Emissions are then allocated to fuels based on standard EFs using equation (3.3):

$$(3.3) \quad E_{PS,f} = E_{PS} \times \frac{A_{PS,f} \times EF_f}{\sum_f A_{PS,f} \times EF_f}$$

where the subscript f denotes the fuel type.

This means that any deviations in data will be distributed across all fuels at the plant. Typical situations include:

- Plants with atypical fuels which differ from standard emission factors
- Plants with errors or other inconsistencies in energy data

In such cases, implied emission factors may deviate from the standard range also for other fuels than the one really affected.

Plants/substances which are entered by fuel currently include among others:

- CO₂ emissions from natural gas in almost all activities
- CO₂ emissions from cement production, 2008 and later
- CO₂ emissions from iron and steel production, 2008 and later
- CO₂ and several other substances from oil and gas production, offshore and onshore

Except for the cases listed above, fuel specific CO₂ emissions from the emission trading system reports (ETS) are not entered into the inventory, only the total plant emission is used.

3.2.1.2 Activity data

The energy balance defines the total energy consumption for which emissions are accounted. However, as explained above, a large part of the total emissions are based on reports from plants that use much energy, i.e. offshore activities and energy-intensive industries on shore. Energy consumption from these plants is included in the energy balance. These consumptions are then subtracted from the energy balance before calculating the remaining emissions. Emissions are estimated using the standard method of multiplying energy use by emission factors described in equation (3.2).

The energy consumption data used in the emission calculations are, with few exceptions, taken from the annual energy balance compiled by Statistics Norway. The energy balance surveys the flow of the different energy carriers within Norwegian territory. These accounts include energy carriers used as raw materials and reducing agents, which are subtracted from the energy balance and are not included in the data used to estimate emissions from combustion. Figure 3.5 shows a flowchart of data and work flow in the energy balance system, with data input, production system and output. There is also an overview of agreements and data collection.

National Inventory Report 2019 - Norway




Figure 3.5. Data and work flow in the energy balance system.

Source: Statistics Norway

As some emissions vary with the combustion technology, a distribution between different sources is required. The total use of the different oil products is based on the Norwegian sales statistics for petroleum products. For other energy carriers, the total use of each energy carrier is determined by summing up reported/estimated consumption within the different sectors.

A short summary of the determination of amounts used by the main groups of energy carriers and of the distribution between emission sources is given below. The following paragraphs give also an explanation of the differences between energy accounts and the energy balance sheets, including the differences involved in Norway's submissions to international organizations. Energy balance sheets for all years in the reporting period are presented in Annex III of this report.

The independent collection of different energy carriers conducted by Statistics Norway, as described below, enables a thorough verification of the emission data reported by the entities to the Norwegian Environment Agency and Norwegian Petroleum Directorate that are included in the inventory.

Natural gas

Most of the combustion of natural gas is related to extraction of oil and gas on the Norwegian continental shelf. The amounts of gas combusted, distributed between gas turbines and flaring, are reported annually to Statistics Norway by the Norwegian Petroleum Directorate (NPD). These figures include natural gas combusted in gas turbines on the various oil and gas fields as well as on Norway's

four gas terminals onshore. However, as explained above, emission figures of CO₂ from the largest gas consumers, e.g. off shore activities, gas terminals, and petrochemical industry, are figures reported by the plants. The data are considered to be of high quality, due to the Norwegian system of CO₂ taxation on fuel combustion.

The remaining combustion of natural gas is given by Statistics Norway's annual survey on energy use in manufacturing industries and by sales figures from distributors. Some manufacturing industries use natural gas in direct-fired furnaces; the rest is burnt in boilers and, in some cases, flared.

LPG and other gases

Consumption of LPG in manufacturing industries is reported by the plants to Statistics Norway in the annual survey on energy use⁹. Figures on use of LPG in households and construction are based on sales figures, collected annually from the oil companies. Use in agriculture are prior to 2005 taken from agriculture statistics (SN2005). From 2005 and onwards, total consumption is given by the annual sales statistics for petroleum products and distributed to agriculture industry using the share of direct sales in 2009-2012. Until further work is done, the same distribution formula is applied to all these years.

Use of refinery gas is reported to Statistics Norway from the refineries. The distribution between direct-fired furnaces, flaring and boilers for the years prior to 2009, is based on information collected from the refineries in the early 1990's. From 2009, the energy consumption are reported according to the energy use at each plant. Emissions from energy combustion for energy purposes are reported under Petroleum refining (CRF 1A1b), emissions from flaring under fugitive emissions from Flaring (CRF 1B2c2) and emissions from cracker are reported under Refining/Storage (CRF 1B2a4). Section 3.4 (CRF 1B2a4) describes the methodology for estimating emissions from cracker. The distribution of emissions from combustion at refineries among the different categories is based on the same proportion for the whole time series. Comparisons made and previously reported to ERTs, shows consistency with what has been reported by the plants.

At some industrial plants, excess gas from chemical and metallurgical industrial processes is burnt, partly in direct-fired furnaces and partly in boilers. These amounts of gases are reported to Statistics Norway. A petrochemical plant generates *fuel gas* derived from ethane and LPG. Most of the gas is burnt on-site, but some fuel gas is also sold to several other plants. All use of fuel gas is reported as energy consumption in the inventory.

Several metallurgical plants generate *CO-rich excess gas* that is either burnt on-site or sold to adjacent plants. Two ferroalloy plants sell parts of their CO-rich gas to other plants (an ammonia producer, a district heating plant, iron and steel producers and mineral industries), where it is used for energy purposes. Thus, these amounts are reported as energy consumption in the inventory.

One sewage treatment plant utilizes biogas extracted at the plant, and reports quantities combusted (in turbines) and calculated CO₂ emissions. Other emissions are estimated by Statistics Norway, using the same emission factors as for combustion of natural gas in turbines.

⁹ <https://www.ssb.no/en/energi-og-industri/statistikker/indenergi>

Oil products

The total use of the different oil products is based on Statistics Norway's annual sales statistics for petroleum products¹⁰. The statistics are based on annual reports from the oil companies and import data from the external trade statistics at Statistics Norway. This is also the data source for consumption in industries that do not collect their own data. For the time series from 1990 to 2009, monthly sales data are used in i EA/EB. These data are also reported to Statistics Norway by the oil companies, but they do not contain as much information as the annual reports. In the monthly sales data, industrial distribution is specified by the oil companies, and there is no information on individual buyers, such as organisation number, name or address. The data are considered very reliable since all major oil companies selling oil products report to these statistics and have an interest in the quality of the data. The statistics are corrected for direct import by other importers or companies.

The use of sales statistics provides a total for the use of oil products. The use in the different sectors must sum up to this total. This is not the case for the other energy carriers. The method used for oil products defines use as identical to sales. Nevertheless, in practice, there will be annual changes in consumer stocks, which are not accounted for. In the statistics on sales of petroleum there is a breakdown of sales by industry. Direct sales to end users are linked to industries or households using the organisation number or other identifiers in the data from the oil companies, while sales to distributors of solid, liquid and gaseous fuels remain attributed to the distributors. However, in energy balance all consumption must be broken down, also that which is sold via distributors. Thus, the breakdown by industry is therefore different in energy balance and in the statistics on sales of petroleum products. The method for this breakdown is described in the report "Energy Accounts and Energy balance – Documentation of statistics production since statistics year 1990" (SSB 2018). Stationary combustion takes place in boilers and, in some manufacturing industries, in direct-fired furnaces. Small ovens can also be used, mainly in private households.

Mobile combustion is distributed among different sources, described in more detail under the transport sector (sections 3.2.4 to 3.2.9).

In addition to oil products included in the sales statistics, figures on use of waste oil are given in Statistics Norway's industry statistics. Statistics Norway also collects additional information directly from a few companies using of waste oil as fuel.

Coal, coke and petrol coke

Use of coal, coke and petrol coke in manufacturing industries is annually reported from the plants to Statistics Norway. The statistics cover all main consumers and are considered of high quality. Combustion of coal and cokes takes place partly in direct-fired furnaces, partly in boilers. The minor quantities burnt in small ovens in private households are estimated based on sales figures. In addition, an insignificant figure of coal use in the agricultural sector has formerly been collected from the farmers. Since 2002, coal has not been used in Norwegian agriculture.

Biofuels

Use of wood waste and black liquor in manufacturing industries is taken from Statistics Norway's annual survey on energy use in these sectors. For the years before 2005 and for 2012, the use of

¹⁰ <https://www.ssb.no/en/energi-og-industri/statistikker/petroleumsalg/aar>

wood in households is based on the annual survey on consumer expenditure which gives the amount of wood burnt. The statistics cover purchase in physical units and estimates for self-harvest of wood. The survey figures refer to quantities *acquired*, which do not necessarily correspond to *use*. The survey gathers monthly data that cover the preceding twelve months; the figure used in the emission calculations (taken from the energy balance), is the average of the survey figures from the year in question and the following year. For the period 2005-2011, the figures are based on responses to questions relating to wood-burning in Statistics Norway's Travel and Holiday Survey. The figures from the survey refer to quantities of wood used. The survey gathers quarterly data that cover the preceding twelve months. The figure used in the emission calculations is the average of 5 quarterly surveys. Since 2013 the figure used in the emission calculations is the average of 3 quarterly surveys. Figures on some minor use in agriculture and in construction have been derived from earlier surveys for these sectors. Combustion of wood product takes place in boilers and in small ovens in private households. Consumption figures for wood pellets and wood briquettes are estimated based on annual information from producers and distributors. Data on use of peat for energy purposes are not available, but according to the Energy Farm, the center for Bioenergy in Norway, such use is very limited (Hohle 2005).

The amount of biofuels (biodiesel and bioethanol) for road transportation are reported separately in the CRF tables. Figure 3.10 shows the consumption of biofuels in the transport sector. The amount of fuels sold is collected from the fuel marketing companies.

Waste

District heating plants and incineration plants annually report combusted amounts of waste (boilers) to Statistics Norway and the Norwegian Environment Agency. Amounts used in manufacturing industries are also reported to Statistics Norway.

According to the Norwegian Pollution Act, each incineration plant has to report emission data for SO₂, NO_x, CO, NH₃, particles, heavy metals and dioxins, and the amount of waste incinerated to the county governor. The county governor then reports this information to the Norwegian Environment Agency. If emissions are not reported, the general method used to estimate emissions from waste incineration is to multiply the amount of waste used by an appropriate emission factor. Normally a plant specific emission factor is built for the component in question. This factor is based on the ratio between previous emission figures and quantities of waste burnt. This factor is then multiplied with the amount of waste incinerated that specific year.

Energy balance sheets vs energy accounts

There are two different ways of presenting energy balances: *Energy balance* sheets (EBS) and *energy accounts*. The energy figures used in the emission calculations are mainly based on the energy balance sheets. The energy balance sheets for the reporting period are presented in Annex III.

The *energy accounts* follow the energy consumption in Norwegian economic activity in the same way as the National accounts. All energy used by Norwegian enterprises and households is to be included. Energy used by Norwegian transport trades and tourists abroad is also included, while the energy used by foreign transport industries and tourists in Norway is excluded.

The *energy balance sheet* follows the flow of energy within Norway. This means that the figures only

include energy sold in Norway, regardless of the users' nationality. This includes different figures between the energy sources balance sheet and the energy account, especially for international shipping and aviation.

The energy balance sheet has a separate item for energy sources consumed for transportation purposes. The energy accounts place the consumption of all energy under the relevant consumer sector, regardless of whether the consumption refers to transportation, heating or processing.

In response to previous review comments, the energy balance has been further disaggregated on energy products.

The consumption of natural gas in the sector is divided among three flows in the energy balance:

- 8.3 – Thermal power plants: Auto producer generation (only segregated for 2007 onwards)
- 10 – Losses: Flaring
- 13 – Net consumption in manufacturing: Remaining natural gas.

Figures from the energy sources balance sheet are reported to international organizations such as the OECD and the UN. The energy balance sheet should therefore usually be comparable with international energy statistics.

Important differences between figures presented in the energy balance sheet (EBS) and figures used in the emission calculations (EC) are:

- *Fishing*: EC use only fuel sold in Norway, whereas EBS also includes an estimate for fuel purchased abroad
- *Air transport*: EC use only Norwegian domestic air traffic (excluding military), while EBS includes all fuel sold in Norway for air transport, including military and fuel used for international air transport
- *Coal/coke for non-energy purposes*: This consumption is included in net domestic consumption in EBS, whereas EC include only energy used for combustion in the calculation of emissions from energy.

3.2.1.3 Emission factors

The standard emission factors used in the absence of more specific ones are addressed as *general*.

CO₂

Emission factors for CO₂ are independent of technology and are based on the average carbon content of fuels used in Norway. The general emission factors for CO₂ used in the emission inventory are listed in Table 3.4, followed by a more detailed description of the factors used for offshore operations and gas terminals.

The factor of 2.34 kg/Sm³ is the default factor used for rich gas combusted in turbines at offshore installations. However, the latest years and specifically after ETS was introduced, field specific EFs have been used in the estimation of CO₂ emissions from combustion of rich gas. More information is given below under *Offshore operations*.

Table 3.4 General emission factors for CO₂.

Energy product	Emission factors	
	Tonne CO ₂ /tonne fuel	Tonne CO ₂ /TJ fuel
Coal	2.52	89.68
Coke	3.19	111.93
Petrol coke	3.59	102.57
Crude oil	3.2	75.65
Motor gasoline	3.13	71.3
Aviation gasoline	3.13	71.3
Kerosene (heating)	3.15	73.09
Jet kerosene	3.15	73.09
Auto diesel	3.17	73.55
Marine gas oil/diesel	3.17	73.55
Light fuel oils	3.17	73.55
Heavy distillate	3.17	73.55
Heavy fuel oil	3.2	78.82
Natural gas (dry gas) (kg/Sm ³) (land)	1.99	56.08
Natural gas (rich gas) (kg/Sm ³) (off shore)	2.34	58.09
LPG	3	65.08
Refinery gas	2.8	57.61
Blast furnace gas	:	198
Fuel gas ³	2.5	50
Landfill gas ^{2,4}	2.75	54.78
Biogas ^{2,4}	2.75	54.78
Fuel wood ²	1.8	107.14
Ethanol ²	1.91	70.84
Biodiesel ²	2.85	76.86
Wood waste ²	1.8	100-110.77
Black liquor ²	1.8	195.65-250
Charcoal	3.299	111.83
Municipal waste	0.55	52.36
Special waste	3.2	78.82

¹ The emission factor for natural gas used in the emission inventory varies as indicated in Tables 3.5 and 3.6.

² Non-fossil emissions, not included in the inventory CO₂ totals.

³ In this inventory, fuel gas is a hydrogen-rich excess gas from petrochemical industry

⁴ Landfill gas and other types of biogas are reported as methane content in the energy balance

Source: Statistics Norway, Norwegian Petroleum Industry Association, SFT (1990), SFT (1996), Climate and Pollution Agency (2011b)

Offshore operations

For all years up to 2002, emissions of CO₂ from gas combustion off shore are calculated by Statistics Norway on the basis of activity data reported by the oil companies to the Norwegian Petroleum Directorate and the Norwegian Environment Agency and the emission factors shown in Table 3.5. For

the years 2003 and onwards,, the data used in the inventory are emissions reported directly by the field operators. The latter are obliged to report these and other emissions annually to the Norwegian Petroleum Directorate and the Norwegian Environment Agency.

The CO₂ emission factor used for all years leading up to 1998 and for all fields except one is one average (standard) factor based upon a survey carried out in the early 1990s (OLF 1993). From 1999 and onwards, the employed emission factors reflect increasingly field specific conditions, as individual emission factors have been reported directly from fields. The measurement frequency varies among the installations. An increasing number uses continuous gas chromatography analysis. Table 3.5 displays the time series of such emission factors, expressed as averages, and based on data reported in EPIM Environment Hub. It is the database in which field operators report emissions data.

Since 2008, off shore gas combustion has been included in the European emission trading system (ETS).

Table 3.5. Average emission factors of CO₂ from the combustion of natural gas in turbines at offshore gas and oil fields.

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017
Gas turbines offshore t CO ₂ /TJ	58.1	56.8	61.5	60.8	58.6	58.6	58.6	58.7	58.2	57.9	58.2	57.9

Source: Norwegian Environment Agency/Norwegian Petroleum Directorate/Environmental Web/EPIM Environment Hub (EEH)

Gas terminals

There are four gas terminals in Norway. The eldest started up before 1990, and then one started up in 1996 and two in 2007.

The CO₂ emission factors for combustion of natural gas on gas terminals are based on continuous or daily plant-specific measurements.

Since 2005, the terminals have been included in the emission trading system (ETS). The average CO₂ emission factors for fuel gas at one gas terminal are shown in Table 3.6. The natural gas used at the terminal originates from three different gas fields and the emission factors in the table reflect the average carbon content in the respective gases. The gas terminal also uses gas from the CO₂ Removal and increased ethane recovery unit (CRAIER) as fuel in a boiler for production of steam. The boiler is connected to a gas treatment unit. The CRAIER unit makes it possible for the gas terminal to receive gas with high content of CO₂ and reduce the CO₂ content in the sales gas to a level that is low enough for the gas market. The CO₂ content in the CRAIER gas burnt in the boiler has varied between 1.6-1.7 tonne CO₂ per tonne gas corresponding to approximately 100 tonnes CO₂ per TJ.

Table 3.6. Average emission factor for CO₂ from the combustion of fuel gas at one gas terminal.

	1990	1995	2000	2005	2010	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Average content of CO ₂ in natural gas t CO ₂ /TJ	56.95	61.80	57.58	56.32	56.32	56.11	55.90	56.11	55.90	55.68	55.47	55.47	55.43	55.22	55.22	55.40

Source: Norwegian Environment Agency

CH₄ and N₂O

For CH₄ and N₂O, information on emission factors is generally very limited, because, unlike the CO₂ emission factors, they depend on the source of the emissions and the sector where the emissions take place. CH₄ and N₂O emission factors for stationary combustion are default factors from IPCC (2006). Net calorific values from the energy balance have been used in order to combine the factors to primary energy data in physical units. Methane emission factor from fuel wood is taken from SINTEF (1995). Due to lack of data, some emission factors are used for sector/source combinations different from those they have been estimated for.

The general CH₄ and N₂O emission factors used in the emission inventory for this source are listed in Table 3.7 and Table 3.9, respectively. Table 3.8 and Table 3.10 display the cases where emission factors other than the general ones have been used in the calculations.

Table 3.7. General emission factors for CH₄, stationary combustion. Unit: kg CH₄ / TJ.

	Direct-fired furnaces	Gas turbines	Boilers	Small stoves	Flares
Coal	1		300	300	
Coke	10		300	300	
Petrol coke	3		10		
Kerosene (heating)			10	10	
Marine gas oil/diesel	10		10		
Light fuel oils			10	10	
Heavy distillate	10		10	10	
Heavy fuel oil	10		10		
Natural gas (dry gas) (land)	5	25.63	5		6.76
Natural gas (rich gas) (off shore)	4.4	22.58	4		5.96
LPG			5	5	
Refinery gas	1		1		5.76
Blast furnace gas	0.67		0.67		
Fuel gas	1		1		1.08
Landfill gas	5		5		7.34
Fuel wood				300	
Wood pellets			11	300	
Wood briquettes			11		
Wood waste			11		
Black liquor			3		
Charcoal	200			203	
Municipal waste			30		
Special waste	30		30		

Numbers in bold have exceptions for some sectors, see Table 3.8.

Source: IPCC (2006), SFT (1996), SINTEF (1995) and (OLF 1994)

National Inventory Report 2019 - Norway

Table 3.8. Exceptions from the general factors for CH₄, stationary combustion. Unit: kg CH₄/TJ.

Emission factor	Fuel	Source	Sectors
3	Kerosene (heating), marine diesel; light fuel oil, heavy distillate	Direct fired furnaces	Energy industry and manufacturing of product
3	heavy fuel oil	Direct fired furnaces, boilers	Energy industry and manufacturing of product
1	LPG	Boilers	Energy industry and manufacturing of product
1.14	Natural gas	Direct fired furnaces, boilers	Extraction of oil and gas
1	Natural gas	Direct fired furnaces, boilers	Energy industry and manufacturing of product
0	Blast furnace gas	Boilers	Refinery
1	Landfill gas, Bio gas	Gas turbines, boilers	Energy industry and manufacturing of product
30	Wood waste	Boilers	Energy industry and manufacturing of product
300	Wood briquettes	Boilers	Private households

Sources: IPCC (2006), SFT (1996), SINTEF (1995) and (OLF 1994)

Table 3.9. General emission factors for N₂O, stationary combustion. Unit: kg N₂O/TJ.

	Direct-fired furnaces	Gas turbines	Boilers	Small stoves	Flares
Coal	1.50		1.50	1.50	
Coke	1.50		1.50	1.50	
Petrol coke	0.60		0.60		
Kerosene (heating)			0.60	0.60	
Marine gas oil/diesel	0.60	0.60	0.60		
Light fuel oils			0.60	0.60	
Heavy distillate	0.60		0.60	0.60	
Heavy fuel oil	0.60		0.60		
Natural gas (dry gas) (land)	0.10	0.10	0.10		0.56
Natural gas (rich gas) (off shore)	0.09	0.09	0.09		0.50
LPG			0.10	0.10	
Refinery gas	0.10		0.10		0.49
Blast furnace gas	0.07		0.07		
Fuel gas	0.10		0.10		0.48
Landfill gas	0.10	0.10	0.10		0.03
Fuel wood				4	
Wood pellets			4	4	
Wood briquettes			4		
Wood waste			4		
Black liquor			2		
Charcoal	4			1	
Municipal waste			4		
Special waste	4		4		

Numbers in bold have exceptions for some sectors, see Table 3.10.

Source: IPCC (2006), SFT (1996), SINTEF (1995) and OLF (1994)

Table 3.10. Exceptions from the general factors for N₂O, stationary combustion. Unit: kg N₂O/TJ.

Emission factor	Fuel	Source	Sectors
0.11	Natural gas	Direct-fired furnaces, gas turbines, boilers	Extraction of oil and gas

Source: Statistics Norway

3.2.1.4 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are presented and discussed in Annex II, as well as under the individual underlying source categories described in the following.

In general, the total energy use is less uncertain than the energy use in each sector. For some sectors, (e.g. the energy and manufacturing industries) the energy use is well known. However, in the case of households and service sectors energy use is more uncertain. The energy use in the most uncertain sectors has been adjusted in the official energy statistics, so that the sum of the energy use in all sectors equals the total sales.

The current method is based on uncertainty estimates for the individual source categories. The main categories are:

- *Use of oil products*: Total amounts are given by the petroleum sales statistics. The uncertainty for total sales are considered to be low due to reliable and complete sales statistics, CO₂-tax and other taxes. The project undertaken for the RA&SA also underlines that this statistics is reliable. However, the allocation of the total consumption to individual sources is more uncertain.
- *Reported emissions from other fuels*, primarily natural gas: Uncertainty data for emissions and energy use are provided in ETS reports. A comparison undertaken as part of the RA&SA project shows that there is good correspondence between the energy consumption by plants covered by the EU ETS and the voluntary agreement and Statistics Norway's own statistics. This also indicates that the energy use in manufacturing industry in the inventory is reliable.

These groups comprise today of about 95 % of CO₂ from energy and 88 % in 1990.

The analyses have not uncovered any major completeness problems in the consumption data. Thus, we have chosen to use the within-source uncertainties in the uncertainty analysis, and to discuss the RA/SA problems in a separate section.

Time series consistency is obtained by the continuous effort to recalculate the entire time series whenever a new source is included in the inventory or new information or methodologies are obtained. However, data availability both for activity data and reported emissions have generally improved over time and new data are included in the emission estimates when deemed of better quality. This causes a degree of time series inconsistency, but the entire time series are considered when new data are included, and efforts made to take the new information into account for all years.

When it comes to activity data, the statistics that form the basis for the energy consumption are not always complete from 1990 onwards. For instance, the waste statistics that form the basis for the waste incineration started in 1995. For the years prior to this, activity data have been backwards extrapolated to ensure consistency in emission estimates.

Emissions reported from the plants are in most cases of good quality, but it may be unfeasible to obtain the estimates for the entire time series. In cases where the reported emissions are deemed to add to accuracy or level of detail in the emission inventory, and the reported figures are unavailable for parts of the time series, reported figures are used although this introduces a certain level of inconsistency. However, emissions for the rest of the time series is calculated based on fuel consumption and standard emission factors, and checks have been made to ensure that the two methodologies gives comparable emission estimates. Times series consistency is thus considered to be met.

3.2.1.5 Category-specific QA/QC and verification

Emission sources in the energy sector are subjected to the QA/QC procedures described in Section 1.2.3 and in Annex VIII QAQC of point sources. Several documentation reports have been published describing the methodologies used for road traffic (Holmengen & Fedoryshyn 2015) and navigation (Tornsjø 2001) and (Flugsrud *et al.* 2010). The methodology for aviation is described in an internal document from Statistics Norway (Thovsen 2018).

The energy statistics that form the basis for the energy balance and energy accounts are subject to individual QA/QC procedures which are not directly linked to the emission inventory system. For the survey on energy use in manufacturing industries, data are edited in a top-down manner, where large units are edited first. The responses from the plants are subject to a set of automated controls that flag outliers and other possible errors (Statistics Norway 2012). The statistics on sales of petroleum products are checked by comparing total sales for each company with additional information from the company. In addition, the companies check that the complete statistics correspond with their own figures. The companies receive tables containing their sales figures, total sales and market shares (Statistics Norway 2015).

Plant specific emission data included in the greenhouse gas inventory are as explained above based on three different reports. Firstly, the annual report that each plant with a permit from the Norwegian Environment Agency has a legal obligation to submit. This report covers all activity at the plant. Emissions data from the largest plants are included in the national greenhouse gas inventory. Secondly, from 2005, we have also received an annual report from entities included in the ETS. In connection with establishing the ETS the plants estimates were quality checked for the time series and specific emphasis on the years 1998-2001. During this process a consistent time series were established for the period from 1990. Thirdly, the Norwegian Environment Agency also receives emission data through a voluntary agreement first established in 1997 between the authority and the industry. From 2005, the agreement covers sectors that are not yet included in the ETS. Data received by the Norwegian Environment Agency through the different reporting channels described above are controlled thoroughly by the Norwegian Environment Agency and Statistics Norway. Especially the emission data plants included in the ETS and in the voluntary agreement are verified extensively. See Annex VIII QAQC of point sources.

3.2.1.6 Category-specific recalculations

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under energy combustion for the whole time-series 1990-2017. See Chapter 10 Recalculations for more details.

3.2.1.7 Category-specific planned improvements

The Norwegian Emission Inventory was subjected to an in-country review in 2018. The team of experts expressed special concerns about the size of the differences in energy use and emissions as estimated by the reference and the sectoral approach. These concerns were addressed in a Saturday Paper. Chapter 10 Recalculations provide information and plans on how the Norwegian inventory team will be working with and respond to the issues raised in the Saturday Paper.

3.2.2 Energy industries, 1A1 (Key category for CO₂ and CH₄)

3.2.2.1 Description

Energy industries include emissions from electricity and heat generation and distribution, extraction and production of oil and natural gas, coal production, gas terminals and oil refineries. Norway produces electricity mainly from hydropower and therefore, emissions from electricity production are small compared to most other countries. Due to the large production of oil and gas, emissions from combustion in energy production are high.

It is important to specify that only emissions from energy combustion for energy purposes are included in section 3.2 Energy Combustion and therefore in the source category Energy industries (CRF 1A1). Emissions from combustion not for energy purposed e.g. flaring are included in section 3.3, 3.4 and 7.5.

Emissions from drilling at moveable offshore installations are included in section 3.2. Emissions from these installations, while not in operation (during transport, etc.), are included with 1A3d Navigation.

In 2017, GHG emissions from the energy industries accounted for 40 % of the energy sector total emissions and 30 % of the total emissions in Norway. Emissions increased by 114 % during the period 1990-2017, primarily due to the increased activity in the oil and gas extraction sector. In 2009, however, the increase was due to approximately one million ton higher CO₂ emissions from gas fired electricity power plants, while the 2.3 % reduction between 2011 and 2012 is the result of decreased emissions from the same sector.

According to the approach 2 key category analysis for 1990 and 2017, this sector is, in conjunction with sectors 1A2 and 1A4, a key category with respect to:

- Emissions of CO₂ from the combustion of liquid fuels, gaseous fuels and other fuels in level in 1990 and 2017, and trend
- Emissions of CH₄ from the combustion of biomass in level in 1990 and 2017
- Emissions of CH₄ from the combustion of gaseous fuels in level in 2017 and in trend

In addition to source categories defined as key categories according to the approach 2 key category analysis, this sector is, in conjunction with sectors 1A2 and 1A4, is defined as key according to approach 1 key category analysis with respect to emissions of CO₂ from combustion of solid fuels.

3.2.2.2 Methodological issues

A description of the general method used for estimating emissions from fuel combustion is given in section 3.2.1.1 and (Statistics Norway 2013). However, most of the reported emissions in this source category are from the annual report from the entities to the Norwegian Environment Agency and the

Norwegian Petroleum Directorate. The guidelines for estimating and reporting emissions are lengthy and in Norwegian, so instead of attaching these to the NIR URLs are provided in section 3.2.1.1 and in Annex VII.

In the case of waste incineration, further specifications on the methodology are given below.

Oil refineries

The emissions from oil refineries are based on annual report from each refinery to the Norwegian environment agency. The reports up to 2004 are taken from the mandatory reporting obligation that is a part of the plants permits given by the authorities and from 2005, emission data are taken from the emission trading system. The distribution of emissions between flaring and energy utilisation of refinery gas in the whole period from 1990 is based on plant and year specific figures. Emissions from energy utilization are reported in petroleum refining (CRF 1A1b) and from flaring in fugitive emissions from flaring (CRF 1B2c).

One of the refineries has a catalytic cracker. Emissions from coke burn off on the catalyst at the cracker are, since they are not for energy purposes, reported in Fugitive Emissions from Oil (CRF 1B2a).

Waste incineration – CO₂ and CH₄

Net CO₂ emissions from wood/ biomass burning are not considered in the Norwegian inventory, because the amount of CO₂ released during burning is the same as that absorbed by the plant during growth. Carbon emitted in compounds other than CO₂, e.g. as CO, CH₄ and NMVOC is also included in the CO₂ emission estimates. This double counting of carbon is in accordance with the IPCC guidelines (IPCC 2006).

Waste incineration – N₂O

Emissions of N₂O are derived from the emissions of NO_x, which are reported from each plant to the Norwegian Environment Agency. More specifically, an estimated amount of 2.5 % of this NO_x is subtracted and reported to UNFCCC as N₂O (SFT 1996). Accordingly, the net NO_x emissions constitute 97.5 % of the emissions reported by the plants. For some years, emissions of NO_x have not been reported for a number of plants. In these cases, specific emission factors for the plants have been made, based upon earlier emissions and amounts of waste incinerated. These new factors have been used to estimate the missing figures.

Public electricity and heat production (CRF 1A1a) – Varying IEFs

The emission sources included in Public electricity and heat production – liquid fuels are consumption of refinery gas at gas fired power plants, consumption of fuel oils, LPG, etc. at district heating plants and consumption of fuel oils in the production of electricity sector.

Emissions from consumption of *refinery gas* included in the inventory are taken from the ETS reports and adjusted for the backflow of fuel gas to refinery. The removed amount of CO₂ is included in Petroleum refining (CRF 1A1b). The adjustment for backflow is due to the fact that the amount and composition of the gas are measured before a separation facility that removes excess hydrogen together with some hydrocarbons.

Emissions from district heating plants and the electricity sector are based on data from the energy balance and default emission factors. Consumption of other liquid fuels is entered as totals in the table below and in the excel spreadsheet due to confidentiality.

The energy liquid carriers used in this sector are refinery gas and other liquid fuels mainly fuel oils and LPG. The change in IEFs from 2010 to 2011 was due to changes in fuel mix between years. The NCV for refinery gas is about 11 % higher than that for other liquid fuels, and the emission factor is 20 % lower. This change in energy mix explains the reduction in the IEF for liquid fuels used in this source category from 2010 to 2011.

3.2.2.3 Activity data

Electricity and heat generation and distribution

The energy producers annually report their use of different energy carriers to Statistics Norway. There is only some minor use of oil products at plants producing electricity from hydropower. Combustion of coal at Norway's only dual purpose power plant at Svalbard/Spitsbergen is of a somewhat larger size. The amount of waste combusted at district heating plants is reported annually both to Statistics Norway and the Norwegian Environment Agency, see Table 3.11. Data are considered to be of high quality.

Table 3.11. Amount of waste combusted at waste incineration plants. 1990-2017. Unit: 1000 tonnes.

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017
Amount of waste incinerated	385	447	586	732	1036	1246	1426	1526	1600	1636	1615	1657

Source: Statistics Norway, Norwegian Environment Agency

Extraction of oil and natural gas

Production of oil and natural gas is the dominating sector for emissions from combustion in the energy industries in Norway. The Norwegian Petroleum Directorate reports annually the amounts of gas combusted in turbines and diesel burnt in turbines and direct-fired furnaces on the oil and gas fields. The data are considered of high quality due to the CO₂ tax on fuel combustion. The activity data are used for 1990-2002. From 2003 onwards, reported emission figures from the field operators are reported into the EPIM Environment Hub (EEH), previously Environmental Web.

The guidelines for estimating and reporting emissions are lengthy and in Norwegian, so instead of attaching these to the NIR URLs are provided in references. Annex VIII describes QA/QC performed for plant specific emission data use in the inventory.

EPIM Environment Hub (EEH) (offshore activities) is described in guidance documents (Norsk olje&gass 2012).

Coal production

Norway's coal production takes place on Svalbard. The only coal producing company reports its coal consumption and some minor use of oil products annually. In addition to emissions related to Norway's own coal production, emissions from Russian activities are also included in the Norwegian emission inventory. As Russian activity data are scarce, emissions from an estimated quantity of coal

combusted in Russian power plants are calculated. Since 1999, there has been only one such plant; in earlier years there were two of those.

Gas terminals

Norway has four gas terminals, where natural gas from the Norwegian continental shelf is landed, treated and distributed. Annual figures on natural gas combusted in turbines and flared are reported to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. Emissions included in inventory for this category are from the gas terminals annual report to the Norwegian Environment Agency.

Oil refineries

The oil refineries annually report their use of different energy carriers to Statistics Norway. Refinery gas is the most important, but there is also some use of LPG and oil products. Emissions included in inventory for this category are from the refineries annual report to the Norwegian Environment Agency. Emissions from the catalytic cracker at one refinery are reported in Refining/Storage (CRF 1B2a4).

3.2.2.4 Emission factors

The emission factors used for energy industries are presented in section 3.2.1.3. For some industries and components, more information about the derivation of the emission factors is given below.

Gas in electricity generation

The CO₂ implied emission factor for use of natural gas in electricity generation varies significantly over the period, from the regular factor of 56.1 t/TJ to over 60 t/TJ. The highest value is in a year with very low emissions (7.2 kt CO₂). In the years with high emissions (>100 kt CO₂), the highest IEF is 58.4 t/TJ.

The variation is primarily a result of the economics of gas power production. Thus, the relative contributions of plants with different plant-specific factors (as based on reports to the Emissions Trading System) also vary significantly. This accounts for the changes in the time series.

Coal in electricity and heat production

The CO₂ factor for solid fuels in electricity generation is low, at 89.7 t/TJ. The emissions in this category is from use of coal at Svalbard. The coal mined at Svalbard has a low carbon emission factor.

The CO₂ factor for solid fuels in heat generation is high and variable, ranging from 164 to 202 t/TJ. The emissions in this category are from blast furnace gas which is sold from a ferroalloy plant to heat distributors. The emissions are based on reports from the plants, from 2008 onwards as part of the Emissions Trading System.

Waste incineration

The emission factors for CO₂, CH₄ and N₂O from combustion of waste (fossil part only) are displayed in Table 3.4, Table 3.7 and Table 3.9, respectively. Emission factors for CH₄ have been calculated by SFT (1996).

The CO₂ emission factor for the fossil part of waste combusted in waste incineration plants in Norway was revised in 2014 (Fedoryshyn 2015). The new factor is based on the assumption that 2.708 tonnes CO₂ per tonne plastic are combusted (based upon the same composition of polymers combusted as in Danish calculations (Denmark NIR 2010 (Nielsen et al. 2010)) and that 20 % of the combusted waste was fossil in 2009 (Norwegian Climate and Pollution Agency 2011). The new factor is a time series that is based on the mean annual change in the fossil share of combusted waste. This change is calculated using the data from Waste accounts Statistics (Statistics Norway) in the period of 1995-2011. For years when data from Waste accounts are not available, the CO₂ emission factor is held constant: in 1994 and before, the 1995 factor is used, while 2011 factor is used in the years after 2011. The energy content of waste used in the new calculation is 11.5 GJ per tonne waste and is based on a report from Avfall Norge (Marthinsen et al. 2010).

Extraction of oil and natural gas

The CO₂ emission factor for gas combustion offshore that has been used for all years leading up to 1998 and for all fields except one is an average factor based upon a survey carried out in the early 1990's (OLF 1993; OLF 1994). From 1999 onwards, the emission factors employed reflect increasingly field specific conditions (see also section 3.2.1.3).

The carbon content of gas burnt varies considerably between the various oil and gas fields. These changes are reflected in the reported emissions. Up to the early 1990s, most of the gas was used in the Ekofisk area, which has a below average carbon content. From around 2000, fields with higher carbon content came into production. Since the last few years, there has been a shift towards fields with somewhat lower carbon content, again.

Oil refineries

The CO₂ emission factor for combustion of refinery gas is based on daily or weekly plant-specific measurements. The refinery gas consists of hydrogen and various hydrocarbons. The composition is variable, leading to changing emissions factors measured as tonne CO₂/tonne fuel or tonne CO₂/TJ. High hydrogen content leads to low emission factors as measured in tonne CO₂/TJ. As an example, a gas with 40 % hydrogen and 60 % hydrocarbons with an average carbon number of 2 gives an emission factor of 50 tonne CO₂/TJ. In the Norwegian inventory, the emission factor varies in the range 45-60 tonne CO₂/TJ.

3.2.2.5 Uncertainties and time series consistency

The uncertainty analysis performed for the energy industries (Annex II) has shown that the uncertainty in the activity data is ± 3 % of the mean for oil, ± 4 % for gas and ± 5 % of the mean for coal/coke and waste.

In the case of the emission factors for CO₂, the uncertainty is ± 3 % of the mean for oil, ± 7 % for coal/coke and gas and ± 30 % of the mean for waste.

Emission factors for CH₄ and N₂O are very uncertain. Distributions are strongly skewed with uncertainties which lie below and above the mean by a factor of 2 and 3, respectively.

The EU ETS emission estimates are available for all years since 2005. The information included in the ETS cannot reasonably be obtained for the time series 1990-2004. Thus, the use of this relatively new

data source introduces a degree of inconsistency in the time-series. However, the energy consumption reported under the ETS system is consistent with the energy consumption reported to Statistics Norway for individual plants. In addition, the CO₂ emission estimates are consistent with the emissions reported to EPIM Environment Hub for offshore activities and through the regular permits for land-based industries. These are the data sources used for emissions, for the years prior to the introduction of the EU ETS scheme. It has thus been assumed that time-series consistency is not significantly affected and that the emission trend is reliable.

3.2.2.6 Category-specific QA/QC and verification

The energy industries are subjected to the general QA/QC procedures described in section 1.2.3 and in Annex VIII QAQC point sources. The category-specific QA/QC described in section 3.2.1.5 is also valid for Energy Industries.

Some category-specific QA/QC activities were conducted in the following industries:

Extraction of oil and natural gas

From 2003 onwards, field specific emission figures reported from the companies are used directly in the emission model. These figures are compared with emissions calculated on the basis of field specific activity data and emission factors.

Oil refineries

The CO₂ emissions reported from the refineries are compared with the emissions estimated by Statistics Norway on the basis of activity data and emission factors for the different energy carriers used.

Results from the above studies have so far shown that emission estimates are consistent with the reported figures.

3.2.2.7 Category-specific recalculations

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under energy combustion for the whole time-series 1990-2017.

1A1ai Public electricity and heat production

- Previously reported activity data where replaced by directly reported emissions. The change resulted in decreased emissions of Co2 and CH4 in the period 2008 to 2016.

3.2.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.2.3 Manufacturing industries and construction, 1A2 (Key category for CO₂ and CH₄)

3.2.3.1 Description

A description of the general method used for estimating emissions from fuel combustion is given in section 3.2.1.1 and in Statistics Norway (2013). Emissions from the sector of manufacturing industries and construction include industrial emissions originating to a large extent from the production of raw materials and semi-manufactured goods (e.g. iron and steel, non-ferrous metals, chemicals (e.g. ammonia, methanol, plastics), fertilizers, pulp and paper, mineral industries, food processing industries, building and construction industry). These emissions are related to fuel combustion only, i.e. emissions from use of oil or gas for heating purposes. Consumption of coal as feedstock and reduction medium is not included in this sector, but is accounted for under the industrial processes sector (CRF 2).

Emissions from this sector contributed to 10 % of the national GHG total in 2017. Emissions from the sector decreased by 5 % from 1990 to 2017.

According to the Approach 2 key category analysis for 1990 and 2017, this sector is, in conjunction with sectors 1A1 and 1A4, a key category with respect to:

- Emissions of CO₂ from the combustion of liquid fuels, gaseous fuels and other fuels in level in 1990 and 2017, and trend
- Emissions of CH₄ from the combustion of biomass in level in 1990 and 2017
- Emissions of CH₄ from the combustion of gaseous fuels in level in 2017 and in trend

In addition to source categories defined as key categories according to the Approach 2 key category analysis, this sector is, in conjunction with sectors 1A1 and 1A4, is defined as key according to Approach 1 key category analysis with respect to emissions of CO₂ from combustion of solid fuels.

3.2.3.2 Methodological issues

A description of the general method used for estimating emissions from fuel combustion is given in section 3.2.1.1. For many plants the emission figures are based on reported figures from the plants to the Norwegian Environment Agency. Indeed, in 2016, these plants accounted for 43 % of the CO₂ emissions from the sector (Table 3.3). The general calculation method, amount of fuel combusted multiplied with a fuel specific emissions factor, is valid for both estimates performed by Statistics Norway and emissions reported by the plants to the Norwegian Environment Agency in this sector.

The reports are from the mandatory reporting obligation that is a part of the plants permits given by the authorities and from 2005, the emission data are from the emission trading system. The ETS was first a voluntary system, 2005-2007, and then as a part of EU ETS, since 2008. From 1997, there have been different voluntary agreements between national authority and the industry. The agreement from 1997 covered the aluminum producers and included, since 2005, industry not included in the ETS. Industry has, in the different voluntary agreements, committed themselves to reduce their greenhouse gas emissions as a group. As part of the agreements, industry has every year reported detailed AD and emissions to the Norwegian Environment Agency. The voluntary agreement has

involved industry i.e. ferroalloy, aluminum, ammonia. From 2013 most of these industries are also part of the ETS.

Figures on energy use are based on data reported from the plants to Statistics Norway. Some of the energy figures used to calculate reported emissions may deviate from the figures in the energy balance. This may, in some cases, cause inaccuracies in IEFs, but generally, this should not be regarded as an important issue.

The guidelines for estimating and reporting emissions are lengthy and in Norwegian, so instead of attaching these to the NIR, URLs are provided in the reference section. Annex VIII describes QA/QC performed for plant specific emission data use in the inventory.

EU ETS

The guidelines for the EU ETS emission reports (Miljødirektoratet 2015) are consistent with the European Union's guidance documents (European Commission). A description of annual normal permit and reporting to the Norwegian Environment Agency is available at the Miljødirektoratet webpage (Miljødirektoratet 2016).

Ammonia production

Emissions from production of ammonia is reported in this section, as far as emissions from combustion from energy utilization is concerned, while emissions from production of hydrogen from wet gas is reported under process emissions (CRF 2B1), see Section 4.3.1. Emissions included in the inventory are from the plant's annual report to the Norwegian Environment Agency.

The emissions from fuel combustion included in this section are liquid petroleum gas of different composition and CO rich blast furnace gas from a producer of ferroalloy. The activity data and emission factors for the different fuels combusted are shown in section 3.2.3.4.

Motorized equipment

Motorized equipment used in manufacturing and construction have been included in this category (CRF 1A2g). Methodologies, activity data and emissions factors are detailed in section 3.2.9.

3.2.3.3 Activity data

Statistics Norway carries out annual surveys on energy use in manufacturing industries, which supply most of the data material for the calculation of combustion emissions in these sectors. The energy use survey covers 90 % of the energy use in this sector. For the remaining companies, figures are estimated based on data from the sample together with data on economic turnover, taking into account use of different energy carriers in the same industries and size groups. A change in methodology from 1998 has had minor consequences for the time series, since the energy use is mainly concentrated in a few major plants within the industry, from which data have been collected both in the current and in the earlier method. The data on energy use in manufacturing industries are considered to be of high quality.

Information on use of waste oil and other hazardous waste is also collected through the energy use statistics.

For the construction industry, the figures on use of the different energy carriers are partly taken from the annual sales statistics for petroleum products and are partly projected from earlier surveys; energy data are considered rather uncertain.

In some sectors, auto diesel is mainly used in machinery and off-road vehicles, particularly in mining and construction. This amount of fuel is based on reported consumption of duty-free auto diesel in the manufacturing industries and on reported sales of duty-free auto diesel to construction. The methods for calculating emissions are discussed in section 3.2.9.

3.2.3.4 Emission factors

Emission factors used in this source category are presented in detail section 3.2.1.3. This section provides information on sectors with variable or deviating implied emission factors in the CRF tables.

Chemical industry (1A2c) – liquid fuels

The IEF for liquid fuels in chemical industry is outside the range of regular liquid fuels due to the use of hydrogen-rich fuel gas.

The liquid energy carriers used in this sector are fuel gas and other liquid fuels, like fuel oils, LPG and oxy gas. Emission sources included in the use of liquid fuels in Chemical industry, are consumption of fuel gas in different chemical productions, e.g. production of ethylene, propylene, polypropylene, polyethylene, and consumption of liquid fuels like fuel oils, LPG and oxy gas. Emissions from consumption of *fuel gas* included in the inventory are taken from ETS reports. Emissions reported by the ETS entities are considered being accurate and lead to a lower IEF since 2008.

Emissions of other liquid fuels included in the inventory are mainly based on data from the energy balance and default emission factors. One exception is emissions from oxy gas from one ETS report.

The ETS reports from *one plant* until 2010 did not report fuel specific emissions. Instead, emissions are reported based on mass balance calculations. For these years, emissions were allocated to fuels based on fuel consumption data reported to Statistics Norway.

The low IEF is due to a high share of hydrogen rich fuel gas (e.g. 68 % in 2011), but activity data are confidential.

Ammonia

The LPGs used as fuels in the ammonia production is mainly a mix of propane/butane with the emission factor of 3.01 tonne CO₂ per tonne gas and ethane with an emission factor of 2.93 tonne CO₂ per tonne gas. For a few years, a small amount of a light fuel gas (composition of 60 % H₂ and 40 % CH₄) from a producer of plastic is used with an emissions factor of 2.4 t CO₂ per tonne gas.

Chemical industry (1A2c) – solid fuels

The CO₂ IEF for solid fuels in chemical industry is outside the range of regular solid fuels due to the use of CO-rich blast furnace gas as a fuel. This gas is sold from a ferroalloy producer and is mainly used as fuel in ammonia production and is reported under solid fuels. The gas has an average plant-specific emission factor of 198 t CO₂/TJ, but inconsistencies in the energy statistics lead to implied emission factors in the range of 190-264 tonne CO₂/TJ. The default emission factor for blast furnace gas in the 2006 guidelines is 70.8 tonnes C/TJ, or 260 tonnes CO₂/TJ (IPCC 2006).

Pulp and paper (1A2d) – biomass

The CO₂ IEF for biomass in the pulp and paper industry varies significantly due to changes in the relative amounts of different fuels. The emissions are primarily from black liquor with plant-specific emission factors in the range of 200-250 t CO₂/TJ and from wood waste with an emission factor of 111 t CO₂/TJ. In 2013, a large plant using black liquor closed down. This led to a large shift to wood waste in the fuel composition, with a corresponding drop in the IEF.

Non-metallic minerals (1A2f) – biomass

The CH₄ IEF for biomass in the minerals industry varies significantly due to changes in the relative amounts of different fuels. The emissions are primarily from charcoal with an emission factor of 200 kg CH₄/TJ and from wood waste with an emission factor of 30 kg CH₄/TJ. Most of the fuel consumption is wood waste, but in some years the use of charcoal leads to strong increases in the average IEF, in particular for 2003. Emissions of CO₂ and N₂O are similar for the fuels, and the IEFs for these gases vary little among years.

3.2.3.5 Uncertainties and time series consistency

Uncertainties in the activity data and the emission factors in the manufacturing industries and construction are as presented in section 3.2.2.5. A more detailed description is presented in Annex II.

The EU ETS emission estimates are available for all years from 2005. For the time period 1990-2004 there are no data from ETS. Thus, the use of this relatively new data source introduces a degree of inconsistency in the time-series. However, the energy consumption reported under the ETS system is consistent with the energy consumption reported to Statistics Norway for individual plants. In addition, the CO₂ emission estimates are consistent with the emissions reported through the regular permits for land-based industries. These are the data sources used for emissions for the years prior to the introduction of the EU ETS scheme. It is thus assumed that time-series consistency is not significantly affected and that the emission trend is reliable.

No other time series inconsistencies are known for this sector.

3.2.3.6 Category-specific QA/QC and verification

QC of plant specific data performed by the inventory compilers in the Norwegian Environment Agency before handing over the data to Statistics Norway to be included in the inventory is quite extensive. The QC is described in section 1.2.3 of the NIR and also in Annex VIII QAQC of point sources, section 5 Current QA/QC procedures and data sources. This is an annual QC.

3.2.3.7 Category-specific recalculations

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under energy combustion for the whole time-series 1990-2017.

3.2.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.2.4 Transport – Civil Aviation, 1A3a (Key category for CO₂)

3.2.4.1 Description

In 2017, emissions from this source category amounted to 9 % of the total emissions from transport and 2 % of the GHG national total. From 1990 to 2017, these emissions increased by 58 % due to activity growth. Emission fluctuations over time have been dictated by the activity growth rates. During the period 1990-2017, the average annual growth in emissions was almost 2 %. The growth amounted to 6 % between 1990 and 1999, and -0.4 % between 1999 and 2016. This indicates that the growth in emissions from domestic aviation was substantial higher in the 90ies than it has been since.

According to the approach 2 key category analysis, Civil aviation is a key category with respect to CO₂ emissions in level both in 1990 and in 2017, and in trend. Emissions of CH₄ and N₂O from this source category are insignificant.

3.2.4.2 Methodological issues

In 2018, the method for calculating emissions from aviation was revised. The method is still in development and will be reviewed until the next submission. The calculation methodology applied is described in an internal document at Statistics Norway, Thovsen (2018). According to the IPCC Good Practice Guidance the methodology used is Tier 3a based on the detailed methodology in the EMEP/EEA (2013). The new method is based on Eurocontrols "Advanced Emission Model"- AEM, combined with data from all aircraft movements to and from Norwegian airports. The new calculation method is based on a "bottom up" calculation of jet kerosene consumption and emissions from aviation based on traffic data, emission factors and energy use factors for aircraft types (kg / km). These calculations make a distribution basis for the majority (> 95%) of total sales of jet kerosene within the categories of use (domestic / foreign), nationality (Norwegian / foreign companies) and flight phase (LTO / Cruise).

The remaining jet kerosene and aviation gasoline are distributed based on assumptions about place of use and nationality in invoice information in sales data from the oil companies. The invoice information also contains information that forms the basis for the economic distribution of all consumption. There is also a distribution of consumption on the type of aircraft (helicopter, jet engine, small aircraft), which is needed to calculate emissions.

The calculation method described is only valid from 2010 and onwards due to missing traffic data for previous years. The time series 1990 to 2009 is adjusted by adding some industries that have previously been missing in the activity data, where there is sufficient information to rewrite consumption within these industries. This will have little effect on the overall distribution between domestic and foreign aviation. No further adjustments have been made to the domestic / foreign distribution. All movements below 1000 metres are included in the "Landing Take Off" (LTO) cycle. Movements over 1000 metres are included in the cruise phase. All emissions from international aviation are excluded from national totals, and are reported separate (see section 3.7.1.3).

3.2.4.3 Activity data

The types of fuel used in aircrafts are both jet kerosene and aviation gasoline. The latter is used mostly in small aircrafts. The total sales of jet kerosene and aviation gasoline are retrieved from the sales statistics of petroleum products, and is believed to cover the actual sales of fuel at Norwegian

airports. Helicopter data is collected from several Norwegian airlines as the data source with aircraft movements has incomplete helicopter data.

Domestic consumption prior to 1995 is estimated by extrapolation on the basis of domestic kilometres flown and is more uncertain.

3.2.4.4 Emission factors

The emission factors used in the emission inventory for civil aviation are presented in Table 3.12 and Table 3.13.

The Norwegian Petroleum Industry Association provides CO₂ emission factors for the combustion of jet fuel and gasoline (Finstad et al. 2002). The CO₂ emission factor used for aviation gasoline is 71.3 tonne CO₂ per TJ and has been applied to all small aircraft. All other aircraft use jet fuel (kerosene) with an emission factor of 73.1 tonne CO₂ per TJ (table 3.4).

For N₂O, a default emission factor is used for all aircraft (IPCC) and is valid for both LTO and the cruise phase.

For NMVOC and CH₄ only aggregated emission factors (kg/tonne fuel used) are used in the Norwegian inventory. The emission factors are calculated based on total emission divided by total fuel consumption from a bottom-up analysis based on EEA data. The VOC emission factors are aircraft specific as given in EEA (2013). For CH₄, the LTO emission factors are calculated annually in the new aviation model. Emission factors prior to 2010 are constant, equal the emission factor in 2010. Studies indicate that only insignificant amounts of methane is emitted during the cruise phase, therefore no methane is calculated for the cruise phase.

Table 3.12. General emission factors for aviation

Source	CH ₄ kg/TJ		N ₂ O kg/TJ
	Aviation gasoline	Jet kerosene	Aviation gasoline/ Jet Kerosene
Charter/scheduled flights			
LTO (0-1000 m)		4.13	2.3
Cruise (Above 1000)			2.3
Helicopters			
LTO (0-1000 m)	43,1	74.2	2.3
Cruise (Above 1000)	0	0	2.3
Small aircraft			
LTO (0-1000 m)	10.7	10.9	2.3
Cruise (Above 1000)	0	0	2.3

Bold numbers are different for different years, see Table 3.13.

Source: EMEP/EEA (2016) and Thovsen (2018)

Table 3.13. Time series of variable CH₄ emission factors from the combustion of jet kerosene and aviation gasoline in aviation

Source	Fuel	CH ₄ Emission Factor (kg/TJ)			
		1990-2010	2015	2016	2017
LTO (0-1000 m)					
Charter/scheduled flights	Jet kerosene	4.34	4.39	4.31	4.12
Small aircraft	Aviation gasoline	9.8	11.9	11.4	10.7

Source: EMEP/EEA (2016) and Thovsen (2018)

3.2.4.5 Uncertainties and time series consistency

Activity data

The uncertainty in the activity data for civil aviation is estimated to be $\pm 20\%$ of the mean primarily due to the difficulty in separating domestic emissions from emissions from fuel used in international transport (Rypdal & Zhang 2000). As described above, data before 1995 are more uncertain than for later years. This may also, to a certain degree, affect the time series consistency.

Emission factors

The uncertainty in the CO₂ emission factors is $\pm 3\%$. The uncertainty in the CH₄ and N₂O emission factors lies below and above the mean by a factor of 2 and 3, respectively.

3.2.4.6 Category-specific QA/QC and verification

There is no category-specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

3.2.4.7 Category-specific recalculations

Due to incorporation of the new aviation model, emissions from aviation has been recalculated for the whole time series. New emission factors from EMEP/EEA, combined with updated flight data and total consumption of aviation fuel from the sales statistics of petroleum products has resulted in decreased emissions from CO₂, CH₄, N₂O and NMVOC in the latter part of the time series. The new factors have been applied from 2010 - 2017, in combination with flight data for the respective reference years. Emission factors for the years prior to 2010 have been set to factors equal to the calculated factor in 2010.

3.2.4.8 Category-specific planned improvements

The new aviation model is still under development and will be thoroughly quality controlled this year to ensure good data quality and documentation.

3.2.5 Transport – Road Transportation, 1A3b (Key category for CO₂)

Road traffic accounted for 70 % of the total GHG emissions from transport and for 17 % of the national GHG total in 2017.

During the period 1990-2017 an increase in emissions of 22 % took place in road transportation.

According to the approach 2 key category analysis for 1990 and 2017, this sector is a key category with respect to emissions of CO₂ in level in 1990 and 2017, and trend.

Passenger cars (PC): Since 1990, emissions from PCs have decreased by 3.6 %, while vehicle kilometers for PC have increased by 47 % and the number of PCs has grown by 69 %. During the period the vehicles have become more fuel efficient and there has been a switch from petrol to diesel driven personnel cars. The switch has specifically been higher since 2007, due to the CO₂ differentiated tax on new personnel cars implemented that year. However, recently the shift in sales has been back to petrol and to electric vehicles. In addition, the consumption of biodiesel and bioethanol has increased since 2006 with a particularly large jump in 2016 and 2017, see Figure 3.10, and hence contributes to the CO₂ emission decrease.

Emissions from **light commercial vehicles** (LCV) and **heavy duty vehicles** (HDV) increased by 120 and 63 %, respectively, during the period 1990-2016.

PC's contribution to total CO₂ emissions from road traffic decreased from 65.9 % in 1990 to 53.2 % in 2016. Light commercial vehicles (LCV) and heavy duty vehicles (HDV) increased their contribution to total emissions for road traffic from 8.8 to 15.1 %, and 23.0 to 29.4 %, respectively, from 1990 to 2016.

The increase in LCV's share of the total emissions from road traffic illustrates the increase of goods transport since 1990 as a consequence of increased trade and consumption of goods due to economic growth.

HDVs consist of trucks and buses but it is specifically trucks that are responsible for the increase of emissions from 1990. This increase is due to economic growth which led to increased activity in the building and construction sector but also to the fact that the trucks have larger motors and are heavier in general.




Figure 3.5. Emissions of CO₂. PC petrol and diesel, LCV and HDV.

Source: Statistics Norway/Norwegian Environment Agency




Figure 3.6. Vehicle kilometer. PC petrol and diesel, LCV and HDV.

Source: Statistics Norway/Norwegian Environment Agency




Figure 3.7. Relative change to 1990 in total CO2 emissions from PC, LCV and HDV. Index 1990=1
Source: Statistics Norway/Norwegian Environment Agency




Figure 3.8. Relative change to 1990 in total vehicle km. PC, LCV, HDV. Index 1990=1
Source: Statistics Norway/Norwegian Environment Agency




Figure 3.9. Relative change to 1990 in number of passenger cars, CO₂ emissions and vehicle kilometers. Index 1990=1. Source: Statistics Norway/Norwegian Environment Agency

3.2.5.1 Methodological issues

Total emissions of CO₂ have been estimated directly from total consumption of each fuel. The consumption of gasoline, including bioethanol, for road traffic has been estimated as total sales minus consumption for other uses, i.e. a top-down approach. Other uses for gasoline are e.g. small boats, snow mobiles and motorized equipment. For auto diesel, the total consumption in road traffic is all auto diesel, including bio-diesel, charged with auto diesel tax. Other uses of auto diesel, excluding bio-diesel are e.g. motorized equipment in agriculture and construction. Consumption on CNG is based on a survey reported by suppliers of CNG. Consumption of LPG is estimated based on figures from the sales statistics on petroleum products and figures from “Drivkraft Norge”, a Norwegian association for the fuel and energy sector in Norway.

Estimates of emissions of CH₄ and N₂O are estimated by the HBEFA model (INFRAS 2017). The model uses a mileage approach:

$$\text{Emissions} = \text{mileage} * \text{emission per km}$$

The model results are used directly, without any adjustment for discrepancies between estimated consumption in the model and registered fuel sale.

The HBEFA model provides emission factors and possibilities for calculating emissions for segments and sub-segments for six vehicle classes: passenger cars, light commercial vehicles, heavy commercial vehicles, urban buses, coaches and motorcycles (including mopeds). The segments are based on engine volume for passenger cars and motorcycles, total weight for heavy commercial vehicles, urban buses and coaches, and gross weight for light commercial vehicles. The segments are

further disaggregated into sub segments based on fuel type and technology type (e.g. Euro-1 – Euro-6). The segments used for Norway in the HBEFA model are presented in Table 3.14.

The model combines the number of vehicles within each segment with driving lengths for the same segments to produce annual national mileage per sub segment. For heavy goods vehicles, the vehicle number is corrected for vehicles driving with trailers, and the driving is split into three load classes (empty, half loaded and fully loaded).

The annual national mileage is split between shares driven in different traffic situations. The traffic situations are a combination of area (urban/rural), road type (e.g. trunk road and access road), speed limit and level of service (free flow, heavy, saturated, and stop and go). The traffic situations are further disaggregated by gradients, where the amount of driving on roads with slopes ranging from - 6 % to 6 % is specified for each traffic situation.

Hot emission factors are provided on the disaggregated level of sub segments and traffic situations with different gradients, and emissions are estimated after these steps of disaggregation.

The HBEFA model provides emission factors for cold emissions and evaporative emissions (soak, running losses and diurnal), in addition to hot emission factors. In order to calculate cold and evaporative emissions, information on diurnal variation in curves of traffic, trip length distributions, parking time distributions and driving behaviour distributions must be provided, in addition to variation in mean air temperature and humidity.

National Inventory Report 2019 - Norway

Table 3.14. Segments used for Norway in the HBEFA

Vehicle class	Segment	Fuel type	Segment split based on
Passenger car	PC petrol <1,4L	Petrol	Engine volume
	PC petrol 1,4-<2L	Petrol	Engine volume
	PC petrol >=2L	Petrol	Engine volume
	PC Hybrid petrol/el medium	Petrol	-
	PC diesel <1,4L	Diesel	Engine volume
	PC diesel 1,4-<2L	Diesel	Engine volume
	PC diesel >=2L	Diesel	Engine volume
	PC Hybrid diesel/el medium	Diesel	-
	PC LPG	LPG	-
	PC PHEV Petrol	Petrol/electricity	-
	PC PHEV Diesel	Diesel/electricity	-
Light commercial vehicles	LCV petrol M+N1-I	Petrol	Tare weight
	LCV petrol N1-II	Petrol	Tare weight
	LCV petrol N1-III	Petrol	Tare weight
	LCV diesel M+N1-I	Diesel	Tare weight
	LCV diesel N1-II	Diesel	Tare weight
	LCV diesel N1-III	Diesel	Tare weight
Heavy goods vehicles	RT petrol	Petrol	-
	RigidTruck <7,5t	Diesel	Gross weight
	RigidTruck 7,5-12t	Diesel	Gross weight
	RigidTruck >12-14t	Diesel	Gross weight
	RigidTruck >14-20t	Diesel	Gross weight
	RigidTruck >20-26t	Diesel	Gross weight
	RigidTruck >26-28t	Diesel	Gross weight
	RigidTruck >28-32t	Diesel	Gross weight
	RigidTruck >32t	Diesel	Gross weight
	Tractor for AT <=7,5t	Diesel	Gross weight
	Tractor for AT>7,5-14t	Diesel	Gross weight
	Tractor for AT>14-20t	Diesel	Gross weight
	Tractor for AT>20-28t	Diesel	Gross weight
	Tractor for AT >34-40t	Diesel	Gross weight
	Tractor for AT >40-50t	Diesel	Gross weight
	Tractor for AT >50-60t	Diesel	Gross weight
Coach	Coach Std <=18t	Diesel	Gross weight
	Coach 3-Axes >18t	Diesel	Gross weight
Urban bus	Ubus Midi <=15t	Diesel	Gross weight
	Ubus Std >15-18t	Diesel	Gross weight
	Ubus Artic >18t	Diesel	Gross weight
	Ubus Std >15-18t CNG	CNG	Gross weight
	Ubus Artic >18t CNG	CNG	Gross weight
Motorcycles and mopeds	Moped <=50cc (v<50kmh)	Petrol	Engine volume
	MC 2S <=150cc	Petrol	Engine volume
	MC 2S >150cc	Petrol	Engine volume
	MC 4S <=150cc	Petrol	Engine volume
	MC 4S 151-250cc	Petrol	Engine volume
	MC 4S 251-750cc	Petrol	Engine volume
	MC 4S >750cc	Petrol	Engine volume

3.2.5.2 Activity data

All activity data are, as far as possible, updated for every year of the inventory. Data are taken primarily from official registers, public statistics and surveys. However, some of the data are based on assumptions. Many of the data sources are less comprehensive for the earliest years in the inventory. The sources of activity data are listed below:

- *Total fuel consumption*: the total amounts of fuels consumed are corrected for off-road use (in boats, snow scooters, motorized equipment, etc.). These corrections are estimated either from assumptions about the number of units, annual operation time, and specific fuel consumption, or from assumptions about and investigations of the fraction of consumption used off-road in each sector. Statistics Norway's sales statistics for petroleum products supplies the data for total fuel consumption (Statistics Norway, Annually). See *Figure 3.10*, which shows the fuel consumption split between fossil petrol and diesel and biofuels (biodiesel and bioethanol). Consumption of biofuels is included in the inventory from 2004. In 2017, the share of bioethanol and biodiesel in fuel sales was respectively 7 and 19 %.
- *Number of vehicles*: the number of vehicles in the various categories and age groups is taken from the statistics on registered vehicles, which receives data from the official register of the Norwegian Directorate of Public Roads. The model input is number of vehicles per vehicle class for each inventory year, and the share of vehicles for any given combination of segment and fuel type. This data are combined with information on the introduction of technology classes to provide number of vehicles within each sub segment. The information on introduction of technology classes are for recent years, based on information from the official register of the Norwegian Directorate of Public Roads and on legislation for the years in which the information in the register is insufficient.
 - The HBEFA model distinguishes between two types of buses: urban buses mainly used for urban driving, and coaches, mainly used for rural and motorway driving. Due to lack of specific information to make this split in the national vehicle register, the distinction between urban buses and coaches are based on a methodology used in Sweden (Swedish Environmental Protection Agency 2011), where the split is made based on the ratio p/w . Here, p is equal to the maximum allowed number of passengers (number of seats plus number of allowed standing passengers), and w is equal to the gross vehicle weight. These data are available in the national vehicle register. Buses with a p/w -value above 3.7 are classified as urban buses, whereas buses with a p/w -value below 3.75 are classified as coaches.
- *Average annual mileage*: Mileages for passenger cars, light commercial vehicles, heavy goods vehicles, coaches and urban buses are, from 2005 onwards, based on odometer readings taken during annual or biannual roadworthiness tests. The readings are collected by the Directorate of Public Roads and further processed by Statistics Norway (Statistics Norway 2010b). For earlier years, most figures are determined from surveys by Statistics Norway or the Institute of Transport Economics. In some instances, assumptions are needed.
 - The statistics on number of vehicles depict the vehicle fleet per December 31st of the inventory year, while the statistics on mileages represents annual driving for the entire year, including

vehicles that have been scrapped or in other ways been in the vehicle fleet for only parts of the inventory year. To adjust for this discrepancy for the years 2005-2016, mean annual driving lengths for each vehicle category have been adjusted upwards in such a way that the totals correspond to the total annual traffic activity from the statistics on annual driving lengths.

- The average annual mileages vary as a function of age, with older vehicles generally driving shorter annual distances than newer vehicles. The correction of driving as a function of vehicle age is based on odometer readings taken during the roadworthiness test. The functions are calculated as the mean of the years 2005-2016, and the same correction curve is used for all years.
 - Motorcycles and mopeds are not subject to roadworthiness tests in Norway. Average annual mileage are taken from a report on transport volumes in Norway (Vågane & Rideng 2010). Due to lack of data, corrections of annual mileage as a function of age for motor cycles and mopeds are taken from a Swedish survey (Björketun & Nilsson 2007) under the assumption that annual mileage as a function of age are comparable in Norway and Sweden.
- *Load data* are taken from the Road goods transport survey (Statistics Norway 2010b).
 - *Transformation patterns* are calculated using information from Statistics Norway' Road goods transport survey on use of trailers and trailer size (Statistics Norway 2010b).
 - *Traffic situations*: The Directorate of Public Roads has data on the annual number of vehicle-kilometres driven on national and county roads. Data are allocated by speed limits, road type, area type (urban/ rural), and vehicle size (small/ large). Traffic on municipal roads is estimated by Statistics Norway based on road lengths, detailed population data, traffic on adjoining roads, etc. The HBEFA model has emission factors for different situations of traffic flow (free flow, heavy traffic, saturated traffic, and stop and go). Assumptions have been made as to this distribution for the different combinations of area type, road type and speed limits for Norway. Effects of road gradients are included, based primarily on Swiss data supplied to the HBEFA.
 - *Ambient conditions* (air temperature and humidity) are included in the model to calculate cold and evaporative emissions. An average of five larger Norwegian cities has been used for spring, summer, autumn and winter separately. Data are based on measurements from the Norwegian meteorological institute.
 - *Trip length and parking time distributions* are calculated from the Norwegian Travel survey (Vibe 1993). The distributions are given on hourly basis.




Figure 3.10. Consumption of gasoline, auto diesel and biofuel for road transportation. 1990-2017. PJ
Source: Statistics Norway

3.2.5.3 Emission factors

Emission factors (except CO₂) are taken from the Handbook of Emission Factors (HBEFA; (INFRAS 2017)). Factors are given as emission per vehicle kilometres for detailed combinations of sub segments and traffic situations.

CO₂

Emission factors for CO₂ are given by fuel type in Table 3.4. The factor for fossil motor gasoline is 71.3 tonne CO₂ per TJ, while the factor for auto diesel is 73.55 tonne CO₂ per TJ. The CO₂ factors used for ethanol is 70.84 tonne CO₂ per TJ and for biodiesel 76.86 tonne CO₂ per TJ.

Table 3.15 shows average CO₂ emissions per year and vehicle category, as calculated by the use of HBEFA.

National Inventory Report 2019 - Norway

Table 3.15. Average CO₂ emission from different vehicle classes, including cold start emissions and evaporation. 1990-2017. Unit: g/km.

Motor gasoline					Auto diesel		
	Passenger cars	Light commercial vehicles	Heavy duty vehicles	Motorcycles	Passenger cars	Light commercial vehicles	Heavy duty vehicles
1990	209	184	482	71	189	215	834
1995	201	187	482	77	175	216	794
2000	187	190	482	84	158	215	810
2005	180	184	483	82	154	203	850
2006	178	182	483	82	154	200	870
2007	177	182	483	82	152	199	891
2008	176	182	482	82	149	198	902
2009	175	181	482	83	148	198	906
2010	174	181	481	83	146	199	911
2011	173	181	481	83	144	199	923
2012	171	181	481	83	142	198	934
2013	169	181	481	84	141	197	944
2014	167	181	481	84	140	196	954
2015	164	181	481	85	140	195	964
2016	162	183	481	85	139	195	977
2017	158	184	481	86	138	194	990

Source: The Norwegian road emission model that is operated by Statistics Norway.

CH₄ and N₂O

Table 3.16. General CH₄ and N₂O emission factors from use of natural gas and LPG for passenger cars and heavy duty vehicles.

Source	Fuel	CH ₄ kg/TJ	N ₂ O kg/TJ
Passenger cars	Natural gas	7.36	0.72
	LPG	22.4	0.97
Heavy duty vehicles	Natural gas	689	9.0

Source: HBEFA (INFRAS 2017), COPERT 5, IPCC (2006)

National Inventory Report 2019 - Norway

Table 3.17. Average N₂O emission factors from road traffic including cold start emissions and evaporation. 1990-2017. Unit: g/km.

	Motor gasoline				Auto diesel		
	Passenger cars	Other light duty vehicles	Heavy duty vehicles	Motorcycles	Passenger cars	Other light duty vehicles	Heavy duty vehicles
1990	0,0072	0.0068	0.0071	0.0013	0.0000	0.0000	0.0087
1995	0,0092	0,0083	0,0071	0,0014	0,0002	0,0004	0,0086
2000	0,0101	0,0113	0,0071	0,0016	0,0024	0,0030	0,0089
2005	0,0054	0,0104	0,0071	0,0015	0,0039	0,0040	0,0076
2006	0,0051	0,0100	0,0071	0,0015	0,0040	0,0041	0,0075
2007	0,0048	0,0097	0,0071	0,0016	0,0042	0,0042	0,0081
2008	0,0045	0,0092	0,0071	0,0016	0,0043	0,0043	0,0094
2009	0,0043	0,0086	0,0071	0,0016	0,0043	0,0043	0,0114
2010	0,0040	0,0079	0,0071	0,0016	0,0043	0,0043	0,0157
2011	0,0036	0,0075	0,0071	0,0016	0,0044	0,0044	0,0211
2012	0,0032	0,0069	0,0071	0,0016	0,0044	0,0044	0,0249
2013	0,0028	0,0063	0,0071	0,0016	0,0044	0,0044	0,0272
2014	0,0023	0,0057	0,0071	0,0016	0,0044	0,0044	0,0318
2015	0,0020	0,0052	0,0071	0,0016	0,0044	0,0044	0,0351
2016	0,0017	0,0047	0,0071	0,0016	0,0044	0,0044	0,0380
2017	0,0014	0,0041	0,0071	0,0016	0,0044	0,0044	0,0402

Source: The Norwegian road emission model that is operated by Statistics Norway

Table 3.18. Average CH₄ emission factors from road traffic including cold start emissions and evaporation. 1990-2017. Unit: g/km.

	Motor gasoline				Auto diesel		
	Passenger cars	Other light duty vehicles	Heavy duty vehicles	Motorcycles	Passenger cars	Other light duty vehicles	Heavy duty vehicles
1990	0.1149	0.1143	0.0935	0.2128	0,0065	0.0065	0.0217
1995	0,0919	0,0989	0,0935	0,1851	0,0050	0,0063	0,0202
2000	0,0588	0,0714	0,0934	0,2041	0,0035	0,0045	0,0134
2005	0,0354	0,0465	0,0934	0,2894	0,0018	0,0026	0,0099
2006	0,0320	0,0430	0,0934	0,2811	0,0016	0,0022	0,0094
2007	0,0295	0,0402	0,0934	0,2713	0,0013	0,0019	0,0086
2008	0,0275	0,0378	0,0934	0,2631	0,0011	0,0016	0,0077
2009	0,0259	0,0359	0,0935	0,2557	0,0010	0,0014	0,0068
2010	0,0244	0,0345	0,0935	0,2507	0,0009	0,0012	0,0057
2011	0,0231	0,0337	0,0935	0,2464	0,0008	0,0011	0,0049
2012	0,0216	0,0328	0,0935	0,2429	0,0007	0,0010	0,0043
2013	0,0202	0,0323	0,0935	0,2388	0,0007	0,0009	0,0040
2014	0,0188	0,0320	0,0935	0,2344	0,0007	0,0008	0,0033
2015	0,0177	0,0318	0,0935	0,2280	0,0006	0,0007	0,0029
2016	0,0165	0,0317	0,0935	0,2219	0,0006	0,0006	0,0027
2017	0,0151	0,0305	0,0935	0,2142	0,0006	0,0006	0,0023

Source: The Norwegian road emission model that is operated by Statistics Norway

NO₂ from gasoline fuelled PC: The N₂O EF in the HBEFA is from the COPERT IV model. In addition to the "normal" reduction of the EF according to the Euro-classes, the N₂O EF is influenced by the sulphur content. Indeed, a lower sulphur content of gasoline leads to a reduced deactivation of the catalyst and reduced N₂O formation. This finding is backed up by several international peer-reviewed papers.

The sulphur content in petrol was 0.3 % in 2004 and 0.05 % in 2005. This sharp drop in sulphur content explains the decrease in N₂O EF between 2004 and 2005. Similar development in the N₂O EF can also be seen in countries, which also use the HBEFA model, e.g. Switzerland and Sweden.

CH₄ and N₂O from biofuels/biomass in road transport

In the inventory, the same emission factors for CH₄ and N₂O are used for biofuels as for corresponding fossil fuels. Thus, the average IEF for biomass in road transport is a function of the fractions of ethanol and biodiesel in the biofuel mix. Initially, the small biofuel amounts were almost exclusively biodiesel, but in recent years ethanol has had a growing share of the mix.

3.2.5.4 Uncertainties and time series consistency

The uncertainty in the activity data and the CO₂ emissions from road transportation is found to be ± 5 % and ± 3 % of the mean, respectively. In the case of CH₄ and N₂O, the uncertainty in the emission factors lies on ± 45 and ± 65 , respectively (Gustafsson 2005). A detailed description of the uncertainty analysis is given in Annex II.

The total consumption of petrol and auto diesel, and hence the CO₂ emissions from these fuels, are well known. The uncertainty for petrol and auto diesel is related to allocation to non-road use.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category. The data quality is generally better for the latter part of the time series.

3.2.5.5 Category-specific QA/QC and verification

The comparison of bottom-up estimates of fuel consumption from HBEFA with total sales (category-specific QA/QC) reveals a discrepancy of 5-15 %. This is deemed to be a reasonable difference. This discrepancy is handled differently for different emission components. The total consumption of each type of fuel is the most important parameter in relation to the reporting requirements of the UNFCCC, as this forms the basis for the calculation of CO₂ from road traffic. One kilogram of gasoline or auto diesel yields a fixed amount of CO₂ irrespective of vehicle type.

The methodology used for calculating N₂O and CH₄ emissions from road transport has been discussed in previous reviews. Emissions are calculated based on vehicle kilometres driven and not by fuel consumption. Calculations of CH₄, N₂O and many other components reported to CLRTAP (e.g. NO_x and particulates), depends on more detailed information about vehicle types and driving patterns, and thus, a more detailed model (for example HBEFA) should be applied. The relationship between emissions and fuel consumption must be considered differently for the emission components that depends directly on the composition and quantity of fuel (CO₂, SO₂ and heavy metals) and those who, to a larger extent, depend on the type of vehicle and driving mode (e.g. NO_x, CH₄, N₂O, NH₃, CO, particles).

Fuel consumption is not an input to HBEFA, where emissions are calculated based on mileage and number of vehicles in each sub-segment of vehicle classes, as well as other data sets, such as cold start and age distribution of mileage. Fuel consumption is however calculated in the model similarly to emission calculations. Biofuels are not handled as separate fuels in HBEFA. The estimated fuel consumption for the country as a whole can be compared with fuel sales from statistics on deliveries of petroleum products and the energy balance. After the revision in the energy balance, the comparison shows that the fuel consumption calculated in HBEFA is more equal to the fuel consumption in the energy balance, especially for petrol. The difference in consumption of auto diesel is fluctuating in the time series, but is less than before the revision. From 1999 the consumption in HBEFA is lower than in the energy balance.

It is not known why there is a discrepancy between the consumption of energy balance and bottom-up calculations in HBEFA, but there are several possible explanations as to why fuel sold does not match the fuel consumption calculated from road transport emission model:

1. *Fuel purchased by foreign vehicles:* Foreign vehicles is not included in the vehicle register statistics, even though they drive on Norwegian roads. Similarly, no fuel bought by Norwegian vehicles abroad is sampled. It is likely that there is no systematic "fuel tourism" across the Norwegian border, as there are no significant price differences between fuel prices in Norway and Sweden. The current calculations are based on the assumption that driving in Norway by foreign vehicles equals the driving of Norwegian vehicles abroad.
2. *Driving patterns:* There may be elements in the driving patterns that cause fuel consumption per kilometre per vehicle to be higher than what the model calculates. One possible reason here is that the fuel consumptions stated in the vehicle type approvals are used as part of the input to the model, and there is an ongoing discussion about whether these systematically underestimates consumption. These data are however available only for the latter part of the series, and cannot explain the discrepancies in the 1990s.
3. *Non-road use:* The allocation of fuels to non-road use is associated with some uncertainty.

Whether the emission calculations should be corrected for differences in fuel consumption depends on the pollutants in questions. For those components that are directly dependent on the amount of fuel (CO₂, SO₂, heavy metals), it will always be appropriate to use the fuel consumption from the energy balance as a basis for calculation. For the other emission components, the decision on whether to correct for total fuel consumption or not will depend on what is causing the discrepancy between fuel consumption calculated in the model and fuel consumption in the energy balance. If the reason is that the total mileage is underestimated in the model, and that the energy balance represents a "truer" picture of the consumption of fuels, emissions should be corrected. If the discrepancy, however, is due to an underestimation of the fuel consumption per kilometre, the emission estimates should not be corrected unless one finds a clear correlation between changes in consumption per kilometre and emissions per kilometre for the relevant emission components. As long as the reason for the discrepancy stay unknown, an assessment of data quality in the various input data is crucial to determining whether emissions should be reconciled against fuel sales or not.

In the previous road transport emission model (SFT 1993), (SFT 1999d), the emissions of all substances were corrected to account for the discrepancy between the energy balance and the model calculations, because the energy balance was considered the most secure data source. When

HBEFA was introduced as the computational model, a new data source was also introduced, namely the mileage statistics at Statistics Norway. These statistics are based on data from periodical technical inspections, and goes back to 2005. This important new data source is considered to be of good quality, and it has changed the assessment of whether the emissions shall be corrected for the consumption of energy balance or not. There is no reason to believe that the total driving lengths are underestimated, and we consider it likely, that the reason for the discrepancy lies in the estimates of fuel consumption per kilometer. We have not found any reason to believe that the reasons for the discrepancies in fuel consumption are directly correlated with driving behaviour. It has therefore been assessed that HBEFA estimates of pollutants that are not directly related to fuel consumption should not be reconciled with fuel consumption.

There are currently no comprehensive statistics on foreign vehicles driving in Norway. One possible explanation for the discrepancy between the calculated fuel consumption in HBEFA and sold quantity of fuel is that foreign driving in Norway exceeds Norwegian of vehicles driving abroad. There has been an issue that the proportion of heavy vehicles with foreign vehicles increases. However, we see no clear increasing trend in the difference between the model results and sales. Better data related to foreign driving in Norway and the Norwegian driving vehicles abroad would strengthen or refute the current assumption that these two balance each other out.

3.2.5.6 Category-specific recalculations

1A3bi-iv Road transport

- Revised activity data. In the revised Energy balance some use of gasoline and diesel are moved from road transport to off-road. Due to the revision, CO₂ emissions are lower in the years 1990-2016 in road transport. There were small changes in CH₄ and N₂O. Activity data for LPG is revised resulting in higher use of LPG in road transport.

3.2.5.7 Category-specific planned improvements

The emission estimation methodology for this source category is currently undergoing improvement, see table 10.9.

3.2.6 Transport – Railways, 1A3c

3.2.6.1 Description

Railway traffic in Norway uses mainly electricity. Auto diesel is used at a small number of lines, for shunting etc. There is also a minor consumption of coal in museum railways. In 2017, GHG emissions from this source category accounted for 0.4 % of the total emissions from transport. Emissions from railways decreased by 53 % from 1990 to 2017.

3.2.6.2 Methodological issues

The general estimation methodology for calculating combustion emissions from consumption figures and emission factors is used in this source category.

3.2.6.3 Activity data

Consumption figures for auto diesel used in rail transport is based on sales statistics for petroleum products. Consumption of coal is estimated based on information from different museum railways;

the same figure is used for all years from 1990.

3.2.6.4 Emission factors

The emission factors used in this source category are displayed in Table 3.4 for CO₂ and Table 3.20 for CH₄ and N₂O.

General emission factors for coal are used in the calculations.

3.2.6.5 Uncertainties and time series consistency

The consumption data are considered to be of high quality. Their uncertainty is estimated to be ± 5 % of the mean. The uncertainty in the emission factors for CO₂ is ± 3 % of the mean, whereas for CH₄ and N₂O, the uncertainty is below and above the mean by a factor of 2 and 3, respectively.

A general assessment of time series consistency has not revealed any inconsistencies in the emission estimates for this category, but there is, as described in section 3.2.6.6 differences before and after 1998 in results from QA/QC checks.

3.2.6.6 Category-specific QA/QC and verification

Foreign railways in Norway are expanding, and the sale statistics for petroleum products include sales to foreigners. Therefore, the sale statistics is considered a better source for fuel consumption than the consumption data from the Norwegian State Railways.

3.2.6.7 Category-specific recalculations

- Revised activity data. In the revised Energy balance the source for activity data has changed for this source category. The use of diesel has increased most of the time series, causing higher emissions.

3.2.6.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.2.7 Transport – Navigation, 1A3d (Key category for CO₂ and CH₄)

3.2.7.1 Description

According to UNFCCC, Norwegian national sea traffic is defined as ships moving between two Norwegian ports. In this connection, installations at the Norwegian part of the continental shelf are defined as ports. Emissions from fishing are described in section 3.2.10.

Greenhouse gas emissions from navigation constituted 5 % of the national GHG total in 2017 and 20 % of emissions from transport. Emissions from shipping have increased by 24 % from 1990 to 2017. The increased emissions in the 90ies can, to a large extent, be explained by the growing activity in the oil and gas sector in general but especially by the fast growing production of crude oil and hence the increasing demand for ships transporting the oil from the oil fields to land. Due to the decreasing production of crude oil since 2001, the demand for transport of crude oil has been reduced. Nevertheless, this reduction has been counteracted by growth in demand in other segments of transport.

Navigation is a key category with respect to CO₂ emissions in level both in 1990 and in 2017 and, for CH₄, in level in 2017 and in trend.

3.2.7.2 Methodological issues

Emissions from navigation are estimated according to the Tier 2 IPCC methodology. Emissions from moveable installations used in oil and gas exploration and extraction are split between energy industries (CRF 1A1) described in section 3.2.2 and navigation: Emissions from drilling are reported under energy industries while emissions from transport and other activities are reported under navigation. Emissions from international marine bunkers are excluded from the national totals and are reported separately (see section 3.7.1.2), in accordance with the IPCC guidelines (IPCC 2006).

Annual emissions are estimated from sales of fuel in domestic shipping, using average emission factors in the calculations.

For 1993 and 1998, (Tornsjø 2001), 2004 and 2007, emissions have also been estimated based on a bottom-up. Fuel consumption data were collected for all categories of ships (based on the full population of Norwegian ships in domestic transport); freight vessels (bulk and tank by size), oil loading vessels, supply/standby ships, tug boats, passenger vessels, fishing vessels, military ships and other ships. Emissions were estimated from ship and size specific emission factors and fuel use. From this information, average emission factors were estimated for application in the annual update based on fuel sales. This approach is unfortunately too resource demanding to conduct annually.

3.2.7.3 Activity data

The annual sales statistics for petroleum products give figures on the use of marine gas oil, heavy distillates and heavy fuel oil in domestic navigation. Information on fuel used in the ship categories in the bottom-up analysis is mainly given by data from the Business Sector's NO_x-fund for 2007 and by earlier Statistics Norway analyses for 1993 and 1998 (Tornsjø 2001), and 2004. Data on fuel consumed by public road ferries are available from the Directorate of Public Roads.

Fuel sales to the oil and gas extraction sector includes stationary and mobile consumption at offshore facilities as well as consumption at supply ships and other supporting vessels. These sales are split between navigation and energy industries. Information on use for drilling, stationary combustion etc., has been taken from the oil companies' reports to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. These reports are found in *EPIM Environment Hub*, a database operated by the Norwegian Oil Industry Association (OLF), Norwegian Petroleum Directorate and the Norwegian Environment Agency. Consumption for these activities is reported under Energy industries (CRF 1A1c-ii). Only the remaining part of sales, assumed to be for drilling rigs during transit, supply ships, etc., is included with Navigation.

For marine gas oil, the amount used for navigation is equal to total sales figures except bunkers, after the deduction of estimated stationary use, mainly in oil and gas extraction, but also some minor use in manufacturing industries and construction.

Use of natural gas in navigation, which was introduced in 2003 and has increased considerably from 2007, is based on sales figures reported to Statistics Norway from the distributors.

3.2.7.4 Emission factors

CO₂

For CO₂ the following standard emission factors based on carbon content are used:

- Marine gas oil/diesel and special distillate: 73.55 tonne per TJ
- Heavy fuel oil: 78.82 tonne per TJ

CH₄ and N₂O

For liquid fuels, the general/standard emission factors for CH₄ and N₂O used in the emission inventory are taken from IPCC/OECD: 0.23 kg CH₄/tonne fuel and 0.08 kg N₂O/tonne fuel.

In the case of oil drilling, the employed factors are as follows:

- CH₄: 0.8 kg/tonne marine gas oil/diesel; 1.9 kg/tonne heavy fuel oil
- N₂O: 0.02 kg/tonne marine gas oil/diesel.

Some natural gas is combusted in ferry transportation and offshore supply; the CH₄ emission factors used are based on the emission factors in

Table 3.19. From the year 2000, when the first vessel that used LNG as fuel started operating, a mean factor for all skips weighted after consumption data for the different ship categories (ferries and supply ships) are calculated. Ferry consumption data used in the calculations are given by the Directorate of Public Roads (Norddal 2010).

Table 3.19. Methane emission factors for vessels using LNG as fuel gas

Vessel category	Methane emission factor (kg CH ₄ /tonne LNG)	Methane emission factor (kg CH ₄ / TJ)
Ferry (currently lean burn engines only)	44	917
Offshore supply (Currently dual fuel engines only)	80	1668

Source: MARINTEK (2010), and estimations from Statistics Norway

The IPCC factor for N₂O from liquid fuels is also used for LNG.

3.2.7.5 Uncertainties and time series consistency

An important source of uncertainty is assumed to be estimation of fuel used by fishing vessels. There is also an uncertainty connected to the fuel use for other domestic sea traffic due to uncertainty in the sale statistics for petroleum products. Important sources of uncertainty are also delimitation of national sea traffic and the emission factors.

The uncertainty in the activity data for navigation is assessed to be ±20 %. With regard to emission factors the uncertainty for ships and fishing vessels is ±3 % of the mean for CO₂. For CH₄ and N₂O the corresponding uncertainties lie in the ranges -50 to +100 and -66 to +200 (see also Annex II).

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

3.2.7.6 Category-specific QA/QC and verification

As mentioned, emission estimates for ships have been made bottom-up for 1993 and 1998 (Tornsjø 2001) and for 2004 and 2007. These results have been compared with top down data (from sales) on

fuel consumption used in the annual estimates. The outcome showed that data from sales were only 1 % higher than data from reported consumption in 2007. For 2004, the data sales were 27 % higher than the consumption data in the bottom-up analysis. This can be explained by the fact that the bottom-up method does not cover all ships, but it may also be that the domestic/international distinction is not specified precisely enough in the sales statistics. Another element, which has not been taken into account, is possible changes in stock. For the years 1993 and 1998, a deviation of -12 and -15 % respectively has been found. In the calculations, sales figures are used, as they are assumed to be more complete and are annually available.

3.2.7.7 Category-specific recalculations

Important recalculations were made in 1A3d National navigation and 1A4ciii National Fishing. Energy consumption for National Navigation has increased for all years, partly because of reallocation from International Navigation and partly because of a reallocation of energy previously reported as consumption under National Fishing. This reallocation is a consequence of new information on the use of marine gas oils from tax data. The former estimation method is believed to have overestimated the emissions from National Fishing with a correspondingly underestimation of National Navigation. The recalculations affect all components, especially CO₂.

3.2.7.8 Category-specific planned improvements

The Norwegian Coastal Administration started in 2011 a project with the aim to use the Automatic Identification System (AIS) to estimate the supply of pollutants from ships to sea. The Norwegian Environment Agency was co-financing the project. In 2015 the delimitation of activities was improved in order to correspond better to definitions in the inventory. A project in collaboration with Statistics Norway and the Norwegian Coastal Administration is currently looking into how these analyses can be used to improve or verify the inventory for navigation.

3.2.8 Transport – Other transportation, 1A3e

3.2.8.1 Description

In old submissions, this source category included emissions from motorized equipment. Since the previous submission, emissions have been reported under the accurate sectors according to the guidelines (IPCC 2006) i.e., CRF 1A2, 1A4 and 1A5.

3.2.8.2 Pipelines

Figures on natural gas used in turbines for pipeline transport at two separate facilities are reported annually from the Norwegian Petroleum Directorate to Statistics Norway. However, energy generation for pipeline transport also takes place at the production facilities. Specific data on consumption for transport are not available. Thus, the consumption at the two pipeline facilities does not give a correct picture of the activity in this sector. For the integrated facilities, the total CO₂ emissions from each facility are reported under the ETS system and are of high quality. The emissions might be split into production and transport using surrogate data, but the accuracy for the two fractions would be much lower than for the total. As a consequence, all emissions from pipelines have been reported under 1A1 Energy Industries.

3.2.9 Motorized equipment (incl. in 1A2, 1A4 and 1A5)

3.2.9.1 Description

The category *motorized equipment* comprises all mobile combustion sources except road, sea, air, and railway transport. Equipment used in agricultural and construction sector is the most important categories. Other categories include mines and quarries, forestry, snow scooters, small boats and miscellaneous household equipment.

Emissions from motorized equipment are estimated using a common methodology but are reported under several source categories:

- Manufacturing and construction: IPPC 1A2g-vii
- Commercial and institutional: IPPC 1A4a-ii
- Households: IPPC 1A4b-ii
- Agriculture/Forestry/Fishing: IPCC 1A4c-ii
- Military: IPCC 1A5b

Primarily consumption of gasoline and auto diesel is considered. A small amount of fuel oil used for equipment in construction is also accounted for.

3.2.9.2 Methodological issues

Emissions are estimated through the general methodology described in section 3.2.1.1, involving consumption figures and appropriate emission factors.

3.2.9.3 Activity data

Gasoline and auto diesel are handled differently. They are both based on data from the energy balance. Auto diesel used in off-road vehicles are tax-free from 1994, and tax-free auto diesel in the years 1990-1993 are extrapolated based on the split between diesel with or without tax in 1995-1998.

Small boats (leisure): The consumption of gasoline and tax-free auto diesel is estimated based on a model using data on size of the fleet, type of fuel, 2- and 4-stroke engine, size of engine. The data is collected from a survey (Båtlivsundersøkelsen) in 2010, and the time series are extrapolated.

Other motorized equipments on tax-free auto diesel: is given as the difference between total sales of tax-free diesel and estimated use in railway transportation and small boats (leisure).

Snow scooters: A portion of 1 % of the gasoline consumption (including bioethanol) in households (mobile combustion) is assigned to snow scooters in the years 1990-2017.

Other motorized equipments on gasoline (e.g. chainsaws and lawn mowers): 2 % of the gasoline consumption (including bioethanol) in households (mobile combustion) is assigned to other motorized equipments in the years 1990-2017. 97 % of the consumption in forestry is assigned to other motorized equipments.

3.2.9.4 Emission factors

The emission factors used are presented in Table 3.20 and Table 3.21.

Emission factors for tractors have been used for tax-free auto diesel consumption in agriculture and

forestry, while emission factors for construction machinery have been used for tax-free auto diesel consumption in all other industries and households.

The emission factors used in the emission model are calculated from the basic factors in Winther and Nielsen (2006), weighted by the age and engine rating distribution of the tractor and construction machinery populations, as well as assumptions on motor load and operating hours and the introduction scheme for emission regulations by the EU (Stage I, II, III and IV).

Emission factors for snow scooters are adapted from the factors for mopeds and motorcycles in the road traffic emission model.

Table 3.20. General emission factors for other mobile sources.

Source	Fuel	CH ₄ kg/TJ	N ₂ O kg/TJ
Railway	Auto diesel	4.18	27.84
	Coal	9.96	1.42
Small boats 2 stroke	Motor gasoline	116.17	0.46
Small boats 4 stroke	Motor gasoline	38.72	1.82
	Auto diesel	4.18	0.70
Motorized equipment 2 stroke	Motor gasoline	136.67	0.46
Motorized equipment 4 stroke	Motor gasoline	50.11	1.59
	Auto diesel	3.94	3.23
	Light fuel oils	3.94	30.16

Snow scooters have the same emission factors as those for Mopeds, see Table 3.17 and Table 3.18

Bold figures have exceptions for some sectors, see Table 3.21.

Sources: Bang (1993), (SFT 1999d) and Statistics Norway (2014b)

Table 3.21. Exceptions from the general factors for greenhouse gases and precursors for other mobile sources.

Component	Emission factor (kg/TJ)	Fuel	Source	Sectors
CH ₄	141.23	Motor gasoline	Motorized equipment 2 stroke	Agriculture
CH ₄	84.28	Motor gasoline	Motorized equipment 4 stroke	Agriculture
CH ₄	178.65	Motor gasoline	Motorized equipment 2 stroke	Forestry and logging
CH ₄	187.94	Motor gasoline	Motorized equipment 2 stroke	Private households
CH ₄	127.61	Motor gasoline	Motorized equipment 4 stroke	Private households
CH ₄	4.18	Auto diesel	Motorized equipment 4 stroke	Private households
N ₂ O	3.06	Auto diesel	Motorized equipment 4 stroke	Agriculture and forestry
N ₂ O	1.86	Motor gasoline	Motorized equipment 4 stroke	Agriculture and forestry, Fishing, Energy sectors, Mining/Manufacturing

Sources: Bang (1993), (SFT 1999d) and Statistics Norway (2002)

3.2.9.5 Uncertainties and time series consistency

The estimates of consumption are considered quite uncertain. However, the total consumption of gasoline and auto diesel is well known (see also Annex II).

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

3.2.9.6 Category-specific QA/QC and verification

There is no category-specific QA/QC procedure for this sector. For a description of the general QA/QC procedure (see Section 1.2.3).

3.2.9.7 Category-specific recalculations

Recalculations are described in chapter 10.

3.2.9.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.2.10 Other Sectors, 1A4 (Key category for CO₂ and CH₄)

3.2.10.1 Description

The source category *Other Sectors* includes stationary combustion in agriculture, forestry, fishing, commercial and institutional industries and households, motorized equipment in agriculture and forestry, snow scooters, fishing vessels and pleasure crafts.

In 2017, the emissions from this sector amounted to 2.9 million tonnes CO₂-equivalents and constitute of 6 % of national total GHG. The sectors emissions decreased by 35 % from 1990 to 2017. Throughout the period 1990-2017, emissions have fluctuated although with a decreasing trend. The low decreasing trend is mainly due to reduced consumption of fuel oil in the commercial, institutional and households sectors.

According to the Approach 2 key category analysis for 1990 and 2017, this sector is, in conjunction with sectors 1A1 and 1A2, a key category with respect to:

- Emissions of CO₂ from the combustion of liquid fuels, gaseous fuels and other fuels in level in 1990 and 2017, and trend
- Emissions of CH₄ from the combustion of biomass in level in 1990 and 2017.
- Emissions of CH₄ from the combustion of gaseous fuels in level in 2017 and in trend

This sector is also an Approach 2 key category with respect to CO₂ emissions in mobile fuel combustion in level in 1990 and 2017.

In addition to source categories defined as key categories according to the Approach 2 key category analysis, this sector is, in conjunction with sectors 1A1 and 1A4, defined as key according to Approach 1 key category analysis with respect to emissions of CO₂ from combustion of solid fuels.

3.2.10.2 Activity data

Motorized equipment

Activity data are as described in section 3.2.9.

Households

Use of fuelwood in households for the years from 2005 to 2011 and after 2013 is based on responses to questions relating to wood-burning in Statistics Norway's Travel and Holiday Survey. The figures in the survey refer to quantities of wood *used*. The survey quarterly gathers data that cover the

preceding twelve months. For the period 2005 to 2011 the figure used in the emission calculations is the average of 5 quarterly surveys. Since 2013 the figure used in the emission calculations is the average of 3 quarterly surveys. For the years before 2005 and for 2012, figures are based on the amount of wood burnt from the annual survey on consumer expenditure. The statistics cover purchase in physical units and estimates for self-harvest. The survey figures refer to quantities *acquired*, which not necessarily correspond to *use*. The survey gathers monthly data that cover the preceding twelve months; the figure used in the emission calculations (taken from the energy accounts) is the average of the survey figures from the year in question and the following year. Combustion takes place in small ovens in private households.

Figures on use of coal and coal coke are derived from information from the main importer. Formerly, Norway's only coal producing company had figures on coal sold for residential heating in Norway. From about 2000, this sale has been replaced by imports from abroad. The volumes of imported charcoal for grocery trade, garden centres, retail sale of furniture and other shops that sell items for the home are placed under household consumption. For the period 2002-2005 the LPG consumption in households is estimated by distributors. From 2005 and onwards, total consumption is given by the annual sales statistics for petroleum products and distributed using the shares of direct sales in 2009-2012, as for agriculture. The consumption is split in use for transport (in passenger cars) and stationary use by estimates from distributors. The estimated use in transport is currently constant for a larger part of the time series, as new estimates have not been attained for a number of years. Consumption of light heating oil and heating kerosene in households is calculated using consumption figures collected as part of Statistics Norway's survey of consumer expenditure (FBU) (SN, 2013). This survey was conducted for the years 1993, 1994, 1995, 2001, 2004, 2006, 2009 and 2012, and households were asked to report the volumes of oil and kerosene they had procured during the past 12 months. The intervening years are calculated using changes in sales of light heating oil and heating oil from the statistics on sales of petroleum products, adjusted for consumption in manufacturing and mining (SN, 2018d; SN 2018e). This is because consumption in manufacturing and mining is less dependent on temperatures than household consumption. Use of natural gas has been based on sales figures reported to Statistics Norway from the distributors. Statistics Norway has a separate model for calculating fuel used in pleasure crafts by households. Here, the consumption of motor gasoline and off-road diesel for boats is calculated for 2-stroke and 4-stroke engines. The number of boats and fuel consumption figures are calculated using a model developed by Civitas. The model is based on a boating survey. The survey estimates the stock of boats for 2010. For 2011 and onwards, the stock is projected using annual sales figures from Norboat and an estimate of the number of boats scrapped per year from 2016. The model was revised by the Norwegian Environment Agency in the beginning of 2018.

Agriculture

The figures for consumption of off-road diesel in agriculture come from Statistics Norway's Sample Survey of Agriculture and Forestry (LU). This is a form-based sample survey where agricultural holdings report how much diesel they have used in their business activity. Questions regarding energy are only included in LU every 3–4 years. Activity data for intervening years is calculated by using the percentage change in quantities in the aggregate accounts for diesel of the Budget Committee for Agriculture as calculated by the Budget Committee for Agriculture (Nibio).

Figures on LPG consumption prior to 2005 are taken from agriculture statistics. From 2005 and onwards, total consumption is given by the annual sales statistics for petroleum products and distributed to agriculture industry using the share of direct sales in 2009-2012. Until further work is done, the same distribution formula is applied to all these years. A figure on the minor use of coal was previously collected annually from the only consumer. Since 2002, however, there has been no use of coal in the Norwegian agricultural activities. Use of natural gas in agriculture, which has increased considerably since it first was registered in 2003, is based on sales figures reported to Statistics Norway from the distributors. The survey was first carried out in 2004, but data on inland consumption of natural gas had been collected since 1994. Prior to 1994 the consumption was insignificant.

Consumption of petroleum products that is not mentioned above (off-road diesel, heating kerosene, light heating oil, heavy gas oil and LPG) is covered by the annual statistics on sales of petroleum products. Distributor sales are broken down by industry according to distribution formulas.

Fishing

Consumption of petroleum products (off-road diesel, heating kerosene, light heating oil, marine gas oil, heavy gas oil and LPG) is covered by the annual statistics on sales of petroleum products. Distributor sales are broken down by industry according to distribution formulas. Monetary figures on refunds of and exemptions from the basic fee on mineral oil are used for distributing consumption of marine gas oil that is not sold directly to industries. Only industries where substantial amounts of marine gas oil are consumed are included: Fishing, extraction of crude petroleum and natural gas, domestic coastal transport and international sea transport. It is assumed that the distribution of refunds and exemptions from the fee is representative for the distribution of consumption of marine gas oil, even though all mineral oils are covered by the fee. The figures are not consistent and do not cover all the relevant industries until 2014. Thus, the figures for 2014 are used for the years 2010-2014.

Commercial and institutional sectors

Consumption of petroleum products is retrieved from the statistics on sales of petroleum products. For stationary petroleum products like light heating oil and heating kerosene, more sales are made directly to users than via distributors. The distribution of the direct sales will then be used as distribution formulas, with the assumption that the direct sales have the same industrial classification as the distributor sales.

From 2005 and onwards, total consumption of LPG is given by the annual sales statistics for petroleum products and distributed using the shares of direct sales in 2009-2012, as for agriculture. It is assumed that LPG consumption in the transport industries, as well as sale and maintenance of vehicles, support activities for transportation and rental and leasing activities is used for transport. For the years prior to 2005 the source of LPG consumption is statistics on the construction industry.

Consumption of natural gas is collected in a separate survey. When necessary, assumptions are made in order to break down consumption in accordance with the detailed industrial classification. Calculated emissions from combustion of biogas at a sewage treatment plant are included for all years since 1993.

3.2.10.3 Emission factor

The emission factors used in this source category are presented in sections 3.2.1.3 and 3.2.9.4.

3.2.10.4 Uncertainties

Uncertainty in *fishing* is described together with navigation in section 3.2.7.

Uncertainty is reduced for sectors where use was previously given as a residual, e.g. use of heating kerosene and heavy distillates in households, and total use of fuel oil in commercial and institutional industries. Generally, the uncertainty, applies mainly to the distribution of use between industries – the total use is defined as equal to registered sales, regardless of changes in stock.

The uncertainty in the activity data for this source category is $\pm 20\%$ of the mean for solid and liquid fuels, and $\pm 30\%$ of the mean for biomass and waste (see Annex II).

3.2.10.5 Category-specific QA/QC and verification

There is no category-specific QA/QC procedure for this sector. For a description of the general QA/QC procedure (see section 1.2.3).

3.2.10.6 Category-specific recalculations

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under energy combustion for the whole time-series 1990-2017. See Chapter 10 Recalculations for more details.

3.2.10.7 Category-specific planned improvements

3.2.11 A new boating survey was published in 2018. Plans are being made to implement the results in the model for pleasure crafts. Other emissions from energy combustion, 1A5

This source includes emissions from fuel use in military stationary and mobile activities, and the use of lubricants in mobile combustion.

3.2.11.1 Description

Military

Emissions of CO₂ from the other mobile sub-sector (1A5b) appear to be a key category according to approach 1 key source analysis.

Emissions from non-fuel use of liquid fuels

In the energy balance, small amounts of gasoline, autodiesel and residual fuel oil is reported as used for non-fuel purposes. A fraction of this consumption is estimated to result in emissions to air, while the remaining fractions remain in products or enter waste streams.

Lubricants in mobile combustion

Two-stroke petrol engines are lubricated by adding oil to the petrol. The oil is thus combusted, and converts to CO₂. As lubricant, oil in two-stroke petrol is not included in the Norwegian energy statistics, a separate estimation must be performed in order to obtain completeness.

3.2.11.2 Activity data and Emission factors

Military

Figures on fuel oil are annually collected directly from the military administration, while for other energy carriers figures from the sales statistics for petroleum products are used. Emission factors used for stationary activities are presented in section 3.2.1.3 and emission factors used for mobile activities are presented in the corresponding transport sectors (see sections 3.2.4 to 3.2.9). The stationary and mobile emissions from the Norwegian military activities for the years 1990-2017 are presented in Table 3.22.

Table 3.22. Stationary and mobile emissions from military activities. 1990-2017.

CO₂ in 1000 tonnes, CH₄ and N₂O in tonnes

	1A5a Military – stationary			1A5b Military – mobile		
	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
1990	IE	IE	IE	266.09	.	8.45
1995	48.06	6.75	0.43	406.55	8.08	12.59
2000	33.90	4.61	0.28	139.26	6.72	4.17
2005	26.56	3.61	0.22	221.25	0.6	6.94
2010	32.72	4.44	0.27	204.62	3.31	6.28
2011	31.09	4.02	0.22	167.06	5.09	5.13
2012	25.32	3.23	0.18	190.45	4.26	5.86
2013	22.74	2.89	0.16	91.54	5.03	2.58
2014	11.07	1.51	0.09	85.69	6.03	2.29
2015	7.72	1.05	0.06	144.73	5.03	4.20
2016	5.79	0.79	0.05	198.05	4.60	5.91
2017	7.95	1.08	0.06	207.65	4.87	6.20

Source: Statistics Norway

Emissions from non-fuel use of liquid fuels

Activity data: Total non-fuel domestic consumption of fuel oils was obtained from the Norwegian energy balance. The quality of the data is mixed throughout the time series. Annual surveys of feedstock use were performed for 1993-2001. For 2002 a different type of survey was used. Results from this survey are used for 2002-2009. For 2010 new estimates were made based on expenditure data, and these results have been used for subsequent years. The changes in method have led to breaks in the time series.

In this inventory, no attempt has been made to splice the different methods in the energy balance.

In addition to gas/diesel oil, gasoline and residual fuel oil has been recorded earlier in the time series.

Times series for the non-fuel use of these fuels are given in section 3.6.3 on feedstocks etc.

Fraction emitted: Gas/diesel oil and gasoline: 0.5, and residual oil: 0.11.

Emission factors: General emission factors for liquid fuels were obtained from Table 3.4, Table 3.8 and Table 3.9. The selected factors are 3 kg CH₄/TJ and 0,6 kg N₂O/TJ.

Lubricants in mobile combustion

The amount of combusted lubricant oil is proportionate to the consumed two-stroke petrol. The blend ratio is assumed to be falling linearly from 3 % in 1990 to 2 % in 2012, based on Internet search (retailers and discussion fora 2014, pages in Norwegian only). Parts of the two-stroke petrol are blended abroad (petrol retailers pers. comm., 2014), and the estimated CO₂ emission from this lubricant oil is hence included in the emission estimates for petrol. The share being blended abroad is not known, and is assumed to be 50 %.

The amount of oil giving emissions not already accounted for is estimated by multiplying the two-stroke petrol consumption by the oil blend ratio and the share of petrol being blended in Norway:

$$(3.4) \quad E = A \times R \times D$$

where:

E = emission

A = consumed two-stroke petrol

R = blend ratio (oil:petrol)

D = share of two-stroke petrol being blended domestically

CH₄ and N₂O

The conversion from tonnes of consumed lubricant to tonnes of emitted CO₂, is performed based on IPCC default factors for energy content (NCV) and carbon content per unit of energy.

Table 3.23. Conversion factors used to estimate CO₂ emissions.

Factor	Value	Unit
Net calorific value (NCV)	0.0402	TJ/tonne
Carbon content (CC)	20	Tonne C/TJ

Source : IPCC (2006)

N₂O and CH₄ emissions have been estimated as fixed fractions of the CO₂ emission, based on IPCC default factors.

Table 3.24. Conversion factors used to estimate CH₄ and N₂O emissions.

Factor	Value	Unit
CH ₄	0.00286	Tonne CO ₂ eq/tonne CO ₂ emitted
N ₂ O	0.00254	Tonne CO ₂ eq/tonne CO ₂ emitted

Source : IPCC (2006)

3.2.11.3 Uncertainties**Military**

There have been large variations in annual sales of military aviation kerosene as stock changes have not been taken into account. The actual annual use of kerosene and hence emissions is therefore uncertain.

Lubricants in mobile combustion

The uncertainty in the emissions estimate from lubricant use in two-stroke petrol engines is assumed to be moderate. The total consumption of gasoline is well known, while the amount going to two-stroke petrol engines is estimated. The uncertainty in the activity data is assumed to be 20 %, based on the uncertainty in the road traffic estimation (see section 3.2.5). The uncertainty of the carbon content is an IPCC default value, and the NCV uncertainty is assumed to be equally large. Based on these uncertainties, the overall uncertainty of the emissions from lubricating oil used in two-stroke petrol engines is estimated to be 30 %.

3.2.11.4 Category-specific recalculations

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under energy combustion for the whole time-series 1990-2017. See Chapter 10 Recalculations for more details.

3.2.11.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.3 Fugitive Emissions from Coal Mining and Handling, 1B1a (Key category for CH₄)

3.3.1 Description

Coal has been shipped from Svalbard since 1907. There were in 2016 two coal mines at Spitsbergen (the largest island in the Svalbard archipelago) operated by a Norwegian company and from 2017 it is only one Norwegian mine left. The second mine was opened in 2001. As the Norwegian GHG inventory, according to official definitions, shall include emissions from all activities at Svalbard, also emissions from Russian coal production have been estimated and included in the Norwegian greenhouse gas inventory. Until 1998, there was production in two Russian coal mines, Barentsburg and Pyramiden, but since then, production takes place only in the Barentsburg mine. The Norwegian mines and Pyramiden are defined as surface mines, whereas Barentsburg is an underground mine.

Abandoned underground mines is included in the inventory. The emissions is reduced from about 10,600 tonne in 1990 to 5,400 tonne CO₂ in 2017 that is a decrease of 49 %.

In 2005 there was a fire in one of the Norwegian coal mines and this caused that the production was almost halved from 2004 to 2005 as Figure 3.11 illustrates it. The emissions from this fire are included in the inventory. The CO₂ emissions from the fire are estimated to approximately 3,000 tonne.

Russian production has since 2001 been considerably smaller than the production in the Norwegian mines. In 2008 a fire started in the Russian mine at Barentsburg. Shortly after the fire started, the mine was filled with water and hence there were no significant emissions from the fire. This is the reason why emissions from the fire are not estimated. The production in 2008 and 2009 was therefore very small. In autumn 2010, ordinary production was restarted. Russian activity data are more uncertain than the Norwegian, which causes a correspondingly higher uncertainty in the emission figures.

At Svalbard there were a smouldering fire in the mine Pyramiden, the Russian mine that was closed down in 1998. At an inspection in 2005, no emissions were registered, which indicates that the fire had burnt out. Due to lack of data, emissions for earlier years from this fire have not been estimated. However, Norwegian authorities assume that these emissions were limited.

Emissions from NMVOC and particles from handling of coal are included.

Figure 3.11 shows that the production of coal at Svalbard has decreased 71 % from 1990 to 2017 due to close down of one of the Norwegian mines. There was a peak in the production in 2007 when the production was nearly five times higher than in 1990. The production increased 80 % from 2000 to 2001 due to the start up of a new Norwegian mine. The production of coal was 27 % lower in 2016 than in 2015. It is the production from the Norwegian surface mines that was reduced by 30 %. This was due to that the mines were preliminary stopped this year.

The emissions from mining were in 2016 estimated to 38,000 tonnes CO₂ equivalents. The emissions decreased by 13 % from 2015 and 2016. Total production of coal in 2016 was 1.0 million tonne.

CH₄ from coal mining is defined as key category according to Approach 2 in both level and trend and only in trend according to Approach 1.




Figure 3.11. Coal productions in Norway excluded abandoned underground mines. 1990-2017. Relative change in production and GHG emissions. 1990=1.

Source: Statistics Norway/Norwegian Environment Agency

3.3.2 Methodological issues

CO₂

Indirect CO₂ emissions from methane and NMVOC oxidized in the atmosphere are calculated by multiplying the calculated CH₄ and NMVOC emissions with, respectively, the factors 2.75 tonne CO₂ per tonne CH₄ and 2.2 tonne CO₂ per tonne NMVOC. (see Chapter 9 for more information about indirect CO₂).

CH₄

Emissions of methane from coal mining on Svalbard are calculated by multiplying the amount of coal extracted (raw coal production) with country specific emission factors (Tier 2). The calculations are performed by Statistics Norway.

NMVOC

NMVOC emissions from handling of coal are estimated by multiplying the amount of coal extracted (raw coal production) with Tier 2 emission factors from EMEP/EEA Guidebook (2013).

Abandoned underground mines

Methane emissions from abandoned underground mines have been calculated with a Tier 1 methodology from the 2006 IPCC Guidelines, using the following formula:

$$CH_4 \text{ emissions} = \text{Number of abandoned coal mines remaining unflooded} \\ * \text{Fraction of gassy coal mines} * \text{Emission factor} * \text{Conversion factor}$$

The conversion factor is the density of CH₄ and converts volume of CH₄ to mass of CH₄. The conversion factor (density) has a value of $0.67 \cdot 10^{-6} \text{ Gg m}^{-3}$.

3.3.3 Activity data

Figures on Norwegian production (raw coal production) are reported by the plant to Statistics Norway. Russian figures are reported to the Norwegian authorities on Svalbard; these figures are, however, regarded as highly uncertain, consisting of a mixture of figures on production and shipments.

Abandoned underground mines

Information on the history of mining at Svalbard was obtained from the Directorate of Mining with the Commissioner of Mines at Svalbard in 2014. The information from the directorate included assessment of degree of flooding. Where no information about flooding is available, the mines are included in the number of abandoned mines remaining unflooded, in order to avoid underestimation. Table 3.25 gives an overview of the number of abandoned mines remaining unflooded for different time periods of abandonment, as well as the used fractions of gassy mines for each time period.

Table 3.25 Number of mines abandoned from 1901-present.

Time of abandonment	Number of abandoned mines remaining unflooded	Fraction of gassy mines
1901-1925	6	0.5
1926-1950	3	0.3
1951-1975	7	0.4
1976-2000	6	0.3
2001-present	0	0.0

Source: Directorate of Mining (2014)

It is assumed that all historic coal mining activities in Norway has taken place at Svalbard.

3.3.4 Emission factors

CH₄

For Norwegian coal production a country specific emission factor of CH₄ from extraction of coal was determined in 2000 in two separate studies performed by (IMC Technical Services Limited 2000) and Bergfald & Co AS (2000).

The emissions of methane from coal mining were in the study measured in two steps. First, coal was sampled and the methane content in coal was analyzed (IMC Technical Services Limited 2000). The sampling process started after a long period (a week) of continuous production. Small samples of coal were removed directly from the coalface as soon as possible after a cut was taken. This was to minimize degassing losses in the samples if the face or heading had been standing for a long time.

The samples yielded an estimate of seam gas content of 0.535-1.325 m³ methane per tonne coal derived from an average content of 0.79 m³ per tonne. This factor includes the total possible methane emissions from coal mining, loading and transport on shore and on sea. The factor also includes the possible emission from handling and crushing of coal at the coal power plant.

Secondly, the methane content in ventilation air from the underground coal mines at Spitsbergen was measured (Bergfald & Co AS 2000). From the Norwegian mines the methane content in the ventilation air was measured to 0.1-0.4 m³ methane per tonne coal.

Considering the measurements it was therefore decided to use 0.54 kg methane per tonne coal as emission factor when calculating methane emissions from coal mining in Norway.

According to IPCC's Good Practice Guidance, the Norwegian mines at Spitsbergen have characteristics that should define the mines as underground mines, whereas the emission factor we use is more characteristic for surface mines. The low content of methane is explained with the mine's location 300-400 meters *above* sea level. Furthermore, the rock at Spitsbergen is porous and therefore methane has been aired through many years.

For the Russian mine in Barentsburg, the emission factor for CH₄ has been estimated in the same manner as the Norwegian factor, based on measurements by Bergfald & Co AS (2000). This is an underground mine, which causes considerably higher emissions than from the Norwegian mines; we use the factor 7.16 kg methane per tonne coal for this mine. Pyramiden, the Russian mine that was closed down in 1998 is, however, situated more like the Norwegian mines; accordingly we use the same emission factor for this as for the Norwegian mines.

NMVOC

Emission factors for NMVOC are taken from EMEP/EEA Guidebook (2013). The Tier 2 factors used are 3 kg NMVOC per tonne coal for surface mines and 0.2 kg NMVOC per tonne coal for underground mines.

Abandoned underground mines

The fraction of gassy mines is determined by the Norwegian Environment Agency based on information about geological characteristics of the different geographic areas of Svalbard, obtained from Bergfald & Co AS (2000) and Directorate Mining with the Commissioner of Mines at Svalbard.

Default emission factors from the tier 1 methodology of the 2006 IPCC Guidelines are used (Table 3.26).

Table 3.26 Emission factors used for calculating emissions from abandoned underground mines. Million m^3 CH_4 /mine.

Inventory year	Time period of abandonment				
	1901-1925	1926-1950	1951-1975	1976-2000	2001-present
1990	0.281	0.343	0.478	1.561	NA
1991	0.279	0.34	0.469	1.334	NA
1992	0.277	0.336	0.461	1.183	NA
1993	0.275	0.333	0.453	1.072	NA
1994	0.273	0.33	0.446	0.988	NA
1995	0.272	0.327	0.439	0.921	NA
1996	0.27	0.324	0.432	0.865	NA
1997	0.268	0.322	0.425	0.818	NA
1998	0.267	0.319	0.419	0.778	NA
1999	0.265	0.316	0.413	0.743	NA
2000	0.264	0.314	0.408	0.713	NA
2001	0.262	0.311	0.402	0.686	5.735
2002	0.261	0.308	0.397	0.661	2.397
2003	0.259	0.306	0.392	0.639	1.762
2004	0.258	0.304	0.387	0.62	1.454
2005	0.256	0.301	0.382	0.601	1.265
2006	0.255	0.299	0.378	0.585	1.133
2007	0.253	0.297	0.373	0.569	1.035
2008	0.252	0.295	0.369	0.555	0.959
2009	0.251	0.293	0.365	0.542	0.896
2010	0.249	0.29	0.361	0.529	0.845
2011	0.248	0.288	0.357	0.518	0.801
2012	0.247	0.286	0.353	0.507	0.763
2013	0.246	0.284	0.35	0.496	0.73
2014	0.244	0.283	0.346	0.487	0.701
2015	0.243	0.281	0.343	0.478	0.675
2016	0.242	0.279	0.34	0.469	0.652
2017	0.241	0.277	0.336	0.461	0.631

Source: IPCC (2006)

3.3.5 Uncertainties and time-series consistency

The uncertainty in the activity data concerning Norwegian coal production is regarded as being low. The uncertainty in Russian data is regarded being considerably higher.

Today, country specific factors based on measurements are used in the calculations. We assume that the uncertainty in the EF is much lower than that reported in Rypdal and Zhang (2000), when an IPCC default emission factor was used. In Rypdal and Zhang (2000) the uncertainty in the EF was estimated by expert judgments to as much as -50 to +100 %.

The EF we use for the Norwegian mines is an average of the measurement of methane in coal sampled in the study (IMC Technical Services Limited 2000). This average EF is two to eight times higher than the methane content measured in ventilation air by Bergfald & Co AS (2000). This should

indicate that the chosen emission factor is rather conservative.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

For abandoned underground mines the same data source is used for the entire time series, and no time series inconsistencies are identified for the calculation of CH₄ emissions from.

3.3.6 Category-specific QA/QC and verification

Independent methods to estimate the EFs used in the calculations are described above in this chapter.

Statistics Norway and the Norwegian Environment Agency carry out internal checks of the emission time-series and corrections are made when errors are detected; see Section 1.2.3 for general QA/QC procedures.

For abandoned underground mines no category-specific QA/QC routines are in place for the emission estimates.

3.3.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

3.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.4 Fugitive Emissions from Oil and Natural Gas – 1B

3.4.1 Overview

Production of oil and gas on the Norwegian continental shelf started on 15 June 1971 when the Ekofisk field came in production, and in the following years a number of major discoveries were made. The Ekofisk field is still in production and is expected to produce oil maybe for additional 40 years. This illustrates the huge amount of oil and gas in that field area. There has been almost a quantum jump in the development of the production technology in the off shore sector since the production activity started. An illustration of this is that the expected recovery factor at Ekofisk was 17 % when the production started and today they expect the recovery to be approximately 50 %. In 2014 there were 78 fields in production on the Norwegian continental shelf included 4 fields that came into production in 2014. Additional 4 fields are being developed and started production in 2015, two fields in 2016 and 5 in 2017. One field on the Norwegian continental shelf closed down in 2014, five in 2016, one in 2017 and three in 2018. By turn of the year 2018/2019 there were 83 fields in production and nine 13 fields were under approved for production.

The overall trend is that the production of oil, gas and NGL and condensate is decreasing since top was reached in 2004. Figure 3.12 below shows the net sale production of oil, gas and NGL and condensate in the period 1974-2017. The total production of oil and gas was 91 % higher in 2017 than in 1990 and 2.4 % higher than in 2016. The production of oil decreased by 1.9 % in 2017 and gas production increased 6.3 %. Maximum production of oil and gas that was reached in 2004 was approximately 264 mill Sm³ oil equivalents. This was an increase since 1990 of 111 %. In 2017 the total production was 10 % lower than the all-time high production in 2004. The maximum production of oil was reached in 2000 and in 2017 the production was 43.4 % lower than in 2000. Production data also shows that the production of gas in 2010 was then for the first time higher than the production of oil and in 2017 the sale gas production was about 34.7 % higher than the sale production of oil. For more information about the Norwegian petroleum sector (OED 2017).




Figure 3.12. Net sale production of oil, gas and NGL and condensate. 1974-2016. Million Sm³ oe.

Source: Statistics Norway

The sector Fugitive Emissions from Oil and Natural Gas (2B) includes fugitive emissions mainly connected to oil and gas exploration and production and gas terminals and refineries. Emissions from combustion for energy purposes in the source categories mentioned above are reported in source category 1.A.c. This is emissions from combustion of natural gas and diesel in turbines, motors and boilers. See below for description of methodology for reported fugitive emissions.

As response to the 2009 annual review report sale production of oil, NGL and condensate are reported in the CRF in source category 1.B.2.a.2 Production oil and sale production of gas in 1.B.2.b.2 Production/processing gas.

3.4.1.1 Fugitive emissions 1990-2017

Fugitive emissions from oil, natural gas and venting and flaring contribute 6.0 % to the total GHG emissions in Norway in 2017 and with 8.2 % of the total GHG emissions in the energy sector. This includes emissions from burn off of coke on the catalysts at one refinery. Without the latter source category fugitives emissions from what we define as oil and gas exploration and production contribute 3.7 % to the total GHG emissions in Norway in 2017 and with 5.0 % of the total GHG emissions in the energy sector.

Figure 3.13 below shows the trend in fugitive emissions from oil and gas production, venting and flaring including burn off of coke at catalytic cracker while Figure 3.14 shows relative change in emissions for the same emission sources. The total sector emissions decreased by 4.0 % from 1990 to 2017 and the emissions decreased by 3.6 % from 2016 to 2017. The emissions from flaring off shore in 2017 decreased specifically due to decreased flaring at an oil field that started in 2016 (down 0.1 million tonn CO₂ or about 30 %). In addition the flaring was reduced with 10-30 000 t CO₂ at several oil and gas fields in 2017.

The fugitive emissions *excluded* emissions from burn off of coke at catalytic cracker at refinery, which are connected to oil and gas exploration and production, decreased by 15.7 % between 1990-2017 while the production of oil and gas increased by 91 %. The different development in emissions and production is mainly explained by measures taken to reduce NMVOC emissions from storage and loading of crude oil offshore and onshore and that flaring of gas is for most years lower than in 1990. More information about flaring off shore is given below.

From Figure 3.13 you can also see that the total emissions from the source category increased substantially from 2006 to 2007-08 and that the emissions today are at 2005 level. The peak emissions in 2007-08 were due to that the LNG plant that started up in 2007 had some start-up problems that gave high emissions. From 2009 the plant came into more regular production.

CO₂ emissions from the burn off of coke at catalytic cracker, that is reported in sector 2.B.2.a.iv Refining/Storage, increased by about 24.4 % in 2017.

Figure 3.13 shows the emissions from source categories in absolute values and Figure 3.14 shows the relative change in emissions compared to 1990. The total emissions for the two source categories with highest emissions, flaring and fugitives from oil including burn off of coke at catalytic cracker (Figure 3.13) contribute over the years to between 60 and 80 % of the sector total. However, emissions from transport that is indirect CO₂ emissions of NMCO₂ and CH₄ from storage and loading of crude oil offshore and onshore is reduced substantially due to measures implemented. The

reduction was compensated with increased emissions from catalytic cracker. Emissions from venting have increased in orders of magnitude from 1990, especially from 2002, and the emissions were about 0.6 million CO₂ equivalents in 2017. See 3.4.4.2 for more information about venting off shore.




Figure 3.13. Fugitive emissions from oil and gas production included catalytic cracker at refinery. Million tonne CO₂ equivalents.

Source: Statistics Norway/Norwegian Environment Agency




Figure 3.14. Relative change in fugitive emissions in CO₂ equivalents from oil and gas included catalytic cracker. 1990=1.

Source: Statistics Norway/Norwegian Environment Agency

3.4.1.2 Emissions from flaring in oil and gas exploration and production

In 2017, CO₂ emissions from flaring off shore contributed with 1.3 % to the total GHG emissions in Norway. The CO₂ emissions from flaring were more than 50 % lower in 2017 than it was in 1990. While the oil and gas production were about 91 % higher, see Figure 3.15. The reduced CO₂ emission from flaring is partly explained by the introduction of tax on gas flared off shore from 1991. The amount of gas flared may fluctuate from year to year due to variation of start-ups, maintenance and interruption in operation. In principle it is allowed to flare from safety reasons only. To minimize emissions from venting and flaring technical measures have been implemented. The venting rate is low due to strict security regulations. The giant leap in emissions from flaring in 1999-2001 was due to that several oil/gas fields came into production in that period. The even higher increase in emissions from flaring in 2007-08 was due to start-up problems at a new LNG plant.




Figure 3.15. Relative change in CO₂ emissions from flaring off shore and total production of oil and gas. 1990-2017. 1990=1.

Source: Statistics Norway/Norwegian Environment Agency

3.4.1.3 Number of exploration and development wellsbores

Figure 3.16 shows the number of exploration wellbores on the Norwegian continental shelf started up in the period 1990-2017. The activity for exploration has been high most of the years with 1994, 1999, 2002-2004 and especially 2005 as years with low activity. In average 37 exploration wells have been started each year from 1990. The timeserie for all exploration wellbores is reported in CRF Reporter in 1B2a1 Exploration and therefore IE is reported in 1B2b1 Exploration.

The total numbers of development wellbores (production, observation and injection) are shown in Figure 3.17.




Figure 3.16. Exploration wellbores. Number of wildcats and appraisal wells started. 1990-2017.
Source: Norwegian petroleum directorate




Figure 3.17. Production wellbores. Number of production, observation and injection wells. 1990-2017.
Source: Norwegian petroleum directorate

3.4.1.4 Overall description of methodology for fugitive emissions from fuels

Table 3.27 gives an overview over methodology (tier), EF and AD for each source category within the sector used in the calculations of the fugitive emissions of CO₂, CH₄, N₂O and NMVOC. The table shows if the EF and/or AD used in the calculation are CS or PS. The notation R/E in the table indicates that emission estimates is based on reporting from the entities (R) or calculated (E) by Statistics Norway; see e.g. Section 3.4.4.2 about flaring. Basically emissions estimates up to about 2002 are carried out by Statistics Norway.

Fugitive and vented emissions from oil and gas activities are included in 1.B.2.c Venting. Flaring are reported in 1B.2.c ii and includes emissions from flaring at oil and gas fields off shore, gas terminals and refineries.

Fugitive emissions (gas leaks) from the following source categories are included in 1.B.2.c Venting and therefore the notation key is IE in CRF:

- exploration and production of oil
- exploration, production/processing and transmission of gas.

ERT's have raised question to why we are not separating fugitive and vented emissions in the inventory. From our knowledge, fugitive emissions from off shore activities represent about 10 % of total fugitive and vented emission. This assumption is based on a survey performed in 2016 (Norwegian Environment Agency 2016). The inventory for fugitive and vented emissions are mostly based on reports from the field operators, see Table 3.28, and we have today not enough information to separate the emissions between the two sources. From our judgment, the accuracy of the emissions will not improve if the emissions were distributed between the source categories 1B2a ii and 1B2b ii. The reporting is from our understanding also in accordance with the reporting guidelines.

Table 3.28 shows the shares of total CO₂, CH₄ and N₂O emissions in the sector that is based on reported and estimated estimates in 2016. From the table you can see that about 90 % of the CO₂ and CH₄ emissions in the sector, included coal mining, are based on reports from the plants, mainly off shore installations. N₂O is based on estimates performed by Statistics Norway.

Sector 1.B.2.a Oil:

- CO₂: 86 % of the emissions in the source category are based on *reports*. The emissions are from catalytic cracker at one oil refinery and indirect CO₂ emissions from loading and storage of crude oil. The emissions from the latter source category are estimated based on reported emission of NMVOC and CH₄.
- CH₄: 100 % is based on *reports* from refineries and oil and gas installations.

1.B.2.b Natural gas:

- CO₂: 100 % is *estimated* and is indirect CO₂ based on mostly reported CH₄ emissions from gas terminals
- CH₄: 72 % of the emissions is based on *reported* emissions from gas terminals.

1.B.2.c Venting and flaring:

- CO₂: 92 % of the emissions are based on *reports* mostly from the oil and gas installations.
- CH₄: 99 % of the emissions are based on *reported* emissions from the oil and gas installations.

Table 3.27. Fugitive emissions from oil and natural gas. Emission sources, compounds, methods, emission factors and activity data included in the Norwegian GHG Inventory.

B Fugitive emissions from fuels	CO ₂	CH ₄	N ₂ O	NMVOC	Method	Emission factor	Activity data
1.B.2.a Oil							
i. Exploration	IE	IE	NO	IE	Tier II	CS	PS
ii. Production	IE	IE	NA	IE	Tier II	CS	PS
iii. Transport	E	R/E	NA	R/E	Tier II	CS	PS
iv. Refining/Storage	R/E	R	NA	R	Tier I/II	CS	PS
v. Distribution of oil products	E	NA	NA	R/E	Tier I	C/CS	CS/PS
vi. Other	NO	NO	NA	NO			
1.B.2.b Natural gas							
i. Exploration	IE	IE	NA	IE	Tier II	CS	PS
ii. Production	IE	IE	NA	IE	Tier II	CS	PS
iii. Processing	IE	IE	NA	IE	Tier II	CS	PS
iv. Transmission	IE	IE	NA	IE	Tier II	CS	PS
v. Distribution	IE	E	NA	IE	Tier II	OTH	CS/PS
vi. Other	E	R	NA	R	Tier II	CS	PS
1.B.2.c Venting							
i. Oil	IE	IE	NA	IE	Tier II	CS/PS	PS
ii. Gas	IE	IE	NA	IE	Tier II	CS/PS	PS
iii. Combined	R/E	R/E	NA	R/E	Tier II	CS/PS	PS
Flaring							
i. Oil (well testing)	R/E	E	E	R/E	Tier II	CS	PS
ii. Gas							
- Gas and oil fields	R/E	R/E	E	R/E	Tier II	CS	PS
- Gas terminals	R	R	E	R/E	Tier I	CS	CS
- Refineries	R	R	R/E	E	Tier I	CS	CS
iii. Combined	IE	IE	IE	IE	Tier I	CS	CS

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated by Statistics Norway (Activity data * emission factor). IE = Included elsewhere, NO = Not occurring, CS = Country specific, PS = Plant specific, Tier = the qualitative level of the methodology used, C=Corinair, OTH=Other.

Table 3.28. Fugitive emissions from oil and natural gas. Share of total CO₂, CH₄ and N₂O emissions in the sector based on estimated and reported emission estimates for 2017.

	CO ₂		CH ₄		N ₂ O	
	Estimated	Reported	Estimated	Reported	Estimated	Reported
1B Fugitive emissions from fuels	10 %	90 %	12 %	88 %	100 %	0 %
1.B.1.a Coal Mining	100 %	0 %	100 %	0 %	100 %	0 %
1.B.2.a Oil	10 %	90 %	0 %	100 %	100 %	0 %
1.B.2.b Natural gas	100 %	0 %	36 %	64 %	100 %	0 %
1.B.2.c Venting and flaring	7 %	93 %	0 %	100 %	100 %	0 %

3.4.2 Fugitive Emissions from Oil, 1.B.2.a (Key category for CO₂)

3.4.2.1 Description

1.B2a covers emissions from loading and storage of crude oil, refining of oil and distribution of gasoline.

Included in the inventory is emission from loading and storage of crude oil produced at the Norwegian continental shelf. This means also those oil fields that is on both the Norwegian and UK continental shelf and is loaded on the Norwegian side of the shelf is included as a whole in the Norwegian inventory and opposite.

Loading, unloading and storage of crude oil on the oil fields offshore and at oil terminals on shore causes direct emissions of CH₄ and indirect emissions of CO₂ from oxidized NMVOC and CH₄. Non-combustion emissions from Norway's two oil refineries (a third was closed down in 2000) include CO₂, CH₄ and NMVOC. It is important to have in mind that included in source category 1.B.2.a.iv is CO₂ from burn off of coke on the catalyst at the catalytic cracker at one refinery, see Section 3.2.2.2. Gasoline distribution causes emissions of NMVOC, which lead to indirect CO₂ emissions.

Loading and storage of crude oil, distribution of gasoline, direct CO₂ emissions from burn off of coke on catalytic cracker at a refinery are according to Approach 1 and 2 *key category* in level and trend for CO₂ and only in level for CH₄.

3.4.2.2 Methodological issues

Loading and storage of crude oil off shore and on shore

The general method for calculating emissions of CH₄ and NMVOC from loading and storage of crude oil are:

field specific amount of crude oil loaded and stored multiplied with field specific emission factors.

For the years 1990-2002 the emissions of CH₄ and NMVOC is calculated by Statistics Norway. The calculation is based on the field specific amounts of crude oil loaded and stored multiplied with field specific emission factors. Field specific activity data and emission factors (the latter only to the Norwegian Environment Agency) used in the calculation were annually reported by the field operators to Statistics Norway and the Norwegian Environment Agency. Since year 2000 an increasing share of the shuttle tankers have had installed vapor recovery units (VRU), and emissions from loading of crude oil on shuttle tankers with and without VRU are calculated separately for each field. In addition emission figures were annually reported to the Norwegian Environment Agency and used in the QC of the emission figures calculated by Statistics Norway.

From 2003, emission of CH₄ and NMVOC from loading and storage of crude oil on shuttle tankers included in the GHG Inventory are based on reported emission figures from the oil companies. Emissions, activity and emissions factors with and without VRU are reported from each field operator into the database EPIM Environment Hub (EEH), previously Environmental Web. The database is operated by the Norwegian Petroleum Directorate, the Norwegian Environment Agency and ¹The Norwegian Oil Industry Association. The method for calculating the emissions is the same as for 1990-2002.

An agreement was established 25 June 2002 between the Norwegian Pollution Control Authority (now Norwegian Environment Agency) and VOC Industrisamarbeid (a union of oil companies operating on the Norwegian continental shelf) aiming to reduce NMVOC emissions from loading and storage of crude oil off shore. So in addition, *also from 2003*, the emission of CH₄ and NMVOC from loading and storage of crude oil on shuttle tankers is reported annually to the Norwegian Environment Agency by the "VOC Industrisamarbeid" in the report "VOC Industrisamarbeid. NMVOC reduksjon bøyelasting norsk sokkel" (VOC Cooperation. Reduction of NMVOC from buoy loading on the Norwegian continental shelf). The report include e.g. details of ships buoy loading and which oil fields the oil has been loaded /stored at, amount of oil loaded, EFs with and without VRU. The method for calculating the emissions is the same as for 1990-2002.

Norway considers that the method for calculating the CH₄ and NMVOC emissions from loading and storage of crude oil is consistent for the period 1990-2014.

Only emissions from loading and storage of the Norwegian part of oil production are included in the inventory.

For the two Norwegian oil terminals on shore, the emissions from loading of crude oil are reported annually from the terminals to the Norwegian Environment Agency. At one of the terminals VRU for recovering NMVOC was installed in 1996. The calculation of the emissions of CH₄ and NMVOC at both terminals is based upon the amount of crude oil loaded and oil specific emission factor dependent of the origin of the crude oil loaded.

The reported indirect CO₂ emissions from the oxidation of CH₄ and NMVOC in the atmosphere for this source category is calculated by Statistics Norway, see Chapter 9.

Refining/Storage – 1.B.2.A.iv

The direct emissions of CO₂, CH₄ and NMVOC included in the inventory are reported by the refineries to the Norwegian Environment Agency. The direct CO₂ emissions reported in this sector originate from the burn off of coke on the catalyst and from the coke calcining kilns at one refinery. The emissions from the catalytic cracker are included in the Norwegian ETS and the emissions reported in source category 1.B.2.a. iv is from the ETS and is therefore regarded being of high quality. The CO₂ emissions from catalytic cracker and calcining kilns are calculated from the formula:

$$\text{tonne CO}_2 \text{ per year} = ((\text{Nm}^3 \text{ RG per year} * \text{volume\% CO}_2) / 100 * (\text{molar weight of CO}_2 / 22.4)) / 1000$$

- the amount of stack gas (RG) is measured continuously
- the density of the stack gas is 1.31 kg/Nm³
- volume percentage of CO₂ is based on continuously measurements. However, if the refinery can document that the volume percentage of CO₂ is not fluctuating more than 2 % from last year report it is not mandatory to have continuous measurements.

Statistics Norway calculates the indirect CO₂ from oxidized CH₄ and NMVOC.

Gasoline distribution – 1.B.2.a.v

NMVOC emissions from gasoline distribution are calculated from the amount of gasoline sold and emission factors for loading of tankers at gasoline depot, loading of tanks at gasoline stations and loading of cars.

3.4.2.3 Activity data

Loading and storage of crude oil off shore and on shore

The amount of oil buoy loaded and oil loaded from storage tankers is reported by the field operators in an annual report to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. The amount of oil loaded on shuttle tankers with or without VRU is separated in the report.

Before 2003, Statistics Norway gathered data on amounts of crude oil loaded at shuttle tankers and stored at storage vessels from the Norwegian Petroleum Directorate. The data from each field are reported monthly by the field operators to the Norwegian Petroleum Directorate on both a mass and a volume basis. The allocation of the amount of crude oil loaded at shuttle tankers and stored at storage vessels with or without VRU is from the annually report the field operators are committed to deliver to the Norwegian Environment Agency and the Norwegian Petroleum Directorate.

The amount of oil loaded at on shore oil terminals is also reported to the Norwegian Environment Agency and the Norwegian Petroleum Directorate.

The amount of crude oil buoy loaded and loaded from storage tankers off shore and crude oil loaded and unloaded at on shore oil terminals is reported for all years in source category 1.B.2.a.iii, as recommended by ERT in previous review reports.

Refining – 1.B.2.a.iv

The crude oil refined included in the CRF is crude oil converted in refineries from the Energy balance.

Gasoline distribution – 1.B.2.a.v

Gasoline sold is annually collected in Statistics Norway's sale statistics for petroleum products.

3.4.2.4 Emission factors

Loading and storage of crude oil off shore and on shore

From 1990 to 2002 emission factors used in the calculation of CH₄ and NMVOC emissions from loading and storage of crude oil offshore and on shore are field/plant specific and were reported to the Norwegian Environment Agency and the Norwegian Petroleum Directorate in an annual report. The Norwegian Environment Agency forwarded the emission factors to Statistics Norway that calculated the emissions.

The evaporation rate varies from field to field and over time, and the emission factors are dependent on the composition of the crude oil as indicated by density and Reid vapour pressure (RVP). The VOC evaporation emission factors are obtained from measurements, which include emissions from loading and washing of shuttle tankers. For some fields the emission factors are not measured, only estimated. The CH₄ content of the VOC evaporated is also measured so that total emissions of VOC are split between CH₄ and NMVOC.

The emission factors that the field operator use in their calculations is reported to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. They report emissions factor with and without VRU and the split between CH₄ and NMVOC. The emission factors are reported by the field operators into the database EPIM Environment Hub (EEH), previously Environmental web.

Loading on shore: The emission factors are considerably lower at one of Norway's two oil terminals than at the other, because the oil is transported by ship and therefore the lightest fractions have already evaporated. At the other terminal the oil is delivered by pipeline. The latter terminal has installed VRU, which may reduce NMVOC emissions from loading of ships at the terminal by about 90 %. NMVOC emissions at this terminal are estimated to be more than 50 % lower than they would have been without VRU. However, the VRU technology is not designed to reduce methane and ethane emissions.

Refining/Storage – 1.B.2.A.iv

The CO₂ emissions from the burn off of coke from the catalytic cracker are calculated as described above under Methodological issues. The CO₂ IEF in CRF is calculated from the emissions from catalytic cracker at one refinery and the amount of crude oil refined at three refineries up to 2002 and thereafter two refineries. This may indicate a low IEF compared to other party's IEF, and, if so, it explains the low IEF.

The emission factor used in the calculation of methane emissions from the largest refinery is based upon measurements using DIAL (Differential absorption LIDAR). A new measurement program was initiated in 2009. An annual EF is deduced from the measured methane emissions and the crude oil throughput. The average EF for the period 2009-2013 is used for the years before the current program was initiated, *i.e.* 1990-2008.

Gasoline distribution – 1.B.2.a.v

Emission factor for NMVOC from filling gasoline to cars used in the calculations are from (EEA 2001) and is 1.48 kg NMVOC/tonne gasoline.

3.4.2.5 Uncertainties and time-series consistency

The uncertainty in the emission factors of methane from *oil loading* (Statistics Norway 2000) and NMVOC (Statistics Norway 2001c) is estimated to be ± 40 % and in the activity data ± 3 %.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

3.4.2.6 Source-specific QA/QC and verification

Statistics Norway gathers data for the amount of crude oil loaded off and on shore from the Norwegian Petroleum Directorate. These data are reported monthly by the field operators to the Norwegian Petroleum Directorate. The activity data are quality controlled by comparing them with the figures reported in the field operator's annual report to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. We have not found any discrepancy of significance between the data from the two data sources.

Statistics Norway's calculated emissions for 1990-02 are compared with the emission data that the field operators report to the Norwegian Environment Agency and the Norwegian Petroleum Directorate. We have not found any discrepancy of significance between the two emission calculations.

From 2003 the Norwegian Environment Agency annual compare data annually reported into the EW by the oil field operators with data from the report "VOC Cooperation. Reduction of NMVOC from

buoy loading on the Norwegian continental shelf". If discrepancies are found between the two sets of data they are investigated and corrections are made if appropriate. If errors are found, the Norwegian Environment Agency contacts the plant to discuss the reported data and changes are made if necessary.

3.4.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

3.4.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.4.3 Fugitive Emissions from Natural Gas, 1.B.2.b (Key category for CH₄)

3.4.3.1 Description

Sector 1.B.2.b covers fugitive emissions of CH₄ and NMVOC and indirect emissions of CO₂ from the two gas terminals and emissions from *distribution* of natural gas. For 1.B.2.b.i Exploration and ii Production/Processing, see section 3.4.1.

The Norwegian gas system has two main parts: The extraction and export sector, including processing terminals and transmission pipelines handling large gas volumes, and a much smaller domestic network. Emissions from transmission, distribution and storage within the main extraction/export system is reported in 1.B.2.b v Other leakage. Emissions from the domestic system is reported in 1.B.2.b iv Distribution.

The rationale for this allocation is that emissions from transmission and storage in the extraction and export sector cannot be split from emissions from extraction and processing emissions at integrated facilities. The emissions from the domestic system might be split. However, the data in 1.B.2.b.iv *Transmission and storage* would then be misleading, as they would cover only a small fraction of Norwegian emissions for this activity. Thus, emissions from 1.B.2.b.iv are reported as "included elsewhere".

CH₄ from natural gas is according to Approach 2 *key category* with respect to trend.

3.4.3.2 Methodological issues

Gas terminals

Fugitive emissions of CH₄ and NMVOC from gas terminals are annually reported from the terminals to the Norwegian Environment Agency.

The emissions are calculated based on the number of sealed and leaky equipment units that is recorded through the measuring and maintenance program for reducing the leakage. The number of sealed and leaky equipment units is collected two times a year and the average number of the counting is used in the calculation. It is assumed in the calculation that a leakage has lasted the whole year if not the opposite is documented.

Gas distribution

Norway has chosen to calculate data for gas transmission and distribution based on the default emission factor from the IPCC 2006 Guidelines. This was decided as conclusion to the discussion with the expert review team during the review of NIR 2016

Only emissions of CH₄ are reported in 1.B.2.b.v. CO₂ emissions are reported as "included elsewhere". According to the energy statistics, the total consumption data refer to amounts fed into the domestic transmission and distribution systems. The same activity data are used for calculating emissions from combustion of the natural gas. Any carbon leakage before combustion would thus be included as CO₂ in the combustion emissions in 1A. This applies to both direct emissions of CO₂ and indirect CO₂ emissions from CH₄ leakage. Direct emissions of CO₂ are likely to be very small: Using the default values from IPCC (2006) they would be 25 tonnes or less throughout the time series. There was no activity in this sector until 1994.

3.4.3.3 Activity data

Activity data are sampled through the terminals measuring and maintenance program which aim is to reduce leakage.

Gas distribution

Data on use of natural gas from the energy statistics are used. From the total domestic consumption including energy sectors the following consumption is excluded:

- Consumption in the gas extraction and processing industry, offshore and onshore, whose emissions from transport is included in 1.B.2.b.vi *Other leakage*
- Consumption for methanol production, whose emissions from transport is included in 2.B.8.a *Methanol*. The plant has its own gas pipeline from an offshore gas field, and emissions from transmission is included with other process emissions at the plant.

The remaining consumption of natural gas is distributed to final consumption by pipeline or LNG systems. An increasing fraction of the consumption is LNG.

The same activity data are used for transmission and distribution. The factors from IPCC (2006) shown below actually refer to amount of *marketable gas* (transmission/storage) and *utility sales* (distribution).

3.4.3.4 Emission factors

Gas distribution

Emission factors from IPCC (2006) are used for the emission estimates, as shown in Table 3.29. The factors refer to pipeline distribution. As no tier 1 methodology was available for LNG distribution, and data for a tier 2 or 3 approach could not be obtained within the available time frame, the factors are used for all Norwegian activity as defined above.

Table 3.29. Emission factors for gas distribution.

Category	Subcategory	Emission source	Value	Selected value	Uncertainty
Gas Transmission and Storage	Transmission	Fugitives	6.6 E-05 to 4.8 E-04	2.73 E-04	±100%
		Venting	4.4 E-05 to 3.2 E-04	1.82 E-04	±75%
	Storage	All	2.5 E-05	2.5 E-05	-20 - 500%
Gas Distribution	All	All	1.1 E-03	1.1 E-03	-20 - 500%

Source: IPCC (2006), vol 2 Energy, table 4.2.4.

3.4.3.5 Uncertainties and time-series consistency

The uncertainty in the emission factors for fugitive methane from natural gas is estimated to be - 50/+100 % and in the activity data ± 3 %.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

3.4.3.6 Source-specific QA/QC and verification

Reported emissions are compared with previous years' emissions.

3.4.3.7 Category-specific recalculations

1B2B5 Distribution

- Updated activity data resulted in increased emissions of CH₄ for the period 1994 to 2016.

3.4.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.4.4 Fugitive Emissions from Venting and Flaring, 1.B.2.c (Key category for CO₂ and CH₄)

3.4.4.1 Description

Included in sector 1.B.2.c *Flaring* are emissions from flaring of gas off shore from extraction and production, at gas terminals and at refineries and the emissions is reported in sector 1.B.2.c.ii. Emission of CO₂, CH₄ and N₂O from flaring of oil when well testing is reported in sector 1.B.2.c.i.

Sector 1.B.2.c *Venting* includes emissions of CO₂, CH₄ and NMVOC from exploration and production drilling of gas and oil. The major source is cold vent and leakage of CH₄ and NMVOC from production drilling.

The sector 1.B.2.c *Venting* includes emissions of CH₄ and NMVOC and hence indirect CO₂ emissions from cold venting and diffuse emissions from extraction and exploration of oil and gas. Since most oil and gas production occur at combined production fields of oil and gas it is not appropriate to split the

emissions between oil and gas production. To divide the emissions from venting between gas and oil production will improve the accuracy of the inventory.

Venting and other emissions connected to CCS is reported in 1C. See Section 3.5 and Annex IV CO₂ capture and storage at the oil and gas production field Sleipner Vest and Hammerfest LNG (Snøhvit gas-condensate field) for description of this source.

Most of the emissions in sector *1.B.2.c Flaring* come from flaring of natural gas offshore (during both well testing, extraction, production and pipeline transport) and at gas terminals and flaring of refinery gas at the refineries. There is some flaring of oil in connection with well testing – amounts flared and emissions are reported to the Norwegian Petroleum Directorate and the Norwegian Environment Agency.

CO₂ and CH₄ from venting and flaring is according to Approach 1 and 2 *key category* with respect to both level and trend.

3.4.4.2 Methodological issues

Venting

Emissions of CH₄ and NMVOC from cold venting and diffuse emissions for each field are reported annually to the Norwegian Environment Agency from the field operator. The emissions are calculated by multiplying the amount of gas produced with an emission factor. The indirect CO₂ emissions are calculated by Statistics Norway.

Research have been conducted by Norwegian oil and gas industry consultancies to develop new country-specific methods and EFs for oil and gas operators regulated by NPD for the estimation of emissions from fugitives and cold venting sources. These new methods and EFs will be used by oil and gas operators to estimate and report emission data for 2017 and onward. This means that the time series must be assessed. Before we do this we (the authorities and the industry) need to gain some experience with the new method. Our assessment today is that the inventory will reflect the new method hopefully from 2020.

Flaring

Flaring of gas off shore - CO₂

The general method for calculating CO₂ emissions from flaring off shore is the amount of gas flared at each field multiplied by field specific emissions factors.

Gas specific data about the gas flared is not available for all flares and years. Therefore the method used for calculating emissions for this source category is not exactly the same for all years.

Estimations of CO₂ 1990-2007.

For the period 1990-2007 the emissions is estimated from the amount of gas flared per field and emission factor based on EU ETS data for 2013. See information below in sub-chapter Emission factors about the emission factors that are used.

Estimations of CO₂ after 2007.

The EU ETS data are reported annually to the Norwegian Environment Agency. From 2008, emissions of CO₂ from flaring used in the inventory is estimated in this way

- Reported EU ETS emissions from flares based on CMR data are used unchanged

- Fields where some flares are with and some are without CMR data: then an average EF for the field based on the CMR data for 2014 is calculated and used for the flares using default EF. For the first years with EU ETS this method is often used for the fields as a whole and thereafter up to 2014 in a decreasing scope
- Gas fields with flaring but without any CMR data in 2014. Then the average emissions factor for 2014 of 2.694 CO₂ per Sm³ based on all CMR data is used.
- For the years after 2014, the same procedure is used for each new year, but the time series is not recalculated annually.

We consider that the method is consistent for all years.

Estimations of CH₄ and N₂O from flaring of gas off shore

Estimated emissions of CH₄ from flaring of gas off shore is calculated by Statistics Norway for 1990-2002 and is thereafter based on reported emission data from the field operators to the Norwegian Petroleum Directorate and the Norwegian Environment Agency. N₂O emissions from flaring is estimated by Statistics Norway for all years.

Well testing

Emissions of CH₄ and N₂O from flaring of *oil in well testing* is estimated for all years by Statistics Norway based on the amount of oil well tested reported annually by the field operators to the Norwegian Petroleum Directorate and the Norwegian Environment Agency. The same emission factors are used for the whole period. CO₂ emissions from well testing is based on the plants annual report.

Gas terminals

Emissions of CO₂ from flaring at the four *gas terminals* that is included in the inventory are reported from the plant.

Refineries

The *refineries* reports annually CO₂ emissions from flaring to the Norwegian Environment Agency. The emissions are calculated by multiplying the amount of gas flared with plant specific emission factors. See additional information section 3.2.1.2.

3.4.4.3 Activity data

Venting

Amount of gas produced or handled at the platforms are reported from the Norwegian Petroleum Directorate to Statistics Norway and used in the QC of the reported emissions.

Flaring

Amounts of gas flared at offshore oil and gas installations are reported on a monthly basis by the operators to the Norwegian Petroleum Directorate.

Amounts of gas flared at the four gas terminals are reported to the Norwegian Petroleum Directorate and the Norwegian Environment Agency.

Amounts of refinery gas flared are found by distributing the total amounts of refinery gas between different combustion technologies by using an old distribution key, based on data collected from the

refineries in the early 1990s. This distribution is confirmed in 2003.

3.4.4.4 Emission factors

Venting

The emission factors used in the calculation of vented emissions is the default emission factors listed in Table 3.30 or field specific factors. Some of the EFs in the table are more accurate (more decimals) than those given in this table in previous submissions. The reference for the default factors is Aker Engineering (1992).

Table 3.30. Default emission factors for cold vents and leakage at oil and gas fields off shore. Emissions are given per Sm³ of processed gas.

	NMVOC	CH ₄	
	Emission factor	Emission factor	Calculation method
Emission source	[g/Sm ³]	[g/ Sm ³]	
Glycol regeneration	0.065	0.265	Tonne per start up
Gas dissolved in liquid from K.O. Drum	0.004	0.0025	
Gas from produced water system	0.03	0.03	
Seal oil systems	0.015	0.010	
Leaks through dry compressor gaskets	0.0014	0.0012	
Start gas for turbines ¹	0.4	0.36	
Depressurization of equipment	0.005	0.016	
Instrument flushing and sampling	0.00021	0.00005	
Purge and blanket gas ¹	0.032	0.023	
Extinguished flare	0.014	0.015	
Leaks in process	0.007	0.022	
Depressurization of annulus	0.000005	0.000005	Tonne per well
Drilling	0.550	0.250	

¹ The gas source is standard fuel gas.

Source: Aker Engineering (1992)

Flaring

Flaring off shore – CO₂

It is mandatory for oil and gas field operators included in the EU ETS to use field or flare specific emissions factor in the calculation of CO₂. If not flare specific factor is used the default emissions factor is 3.73 kg CO₂ per Sm³. The default emission factor is often considerable higher than measured emission factors. This has motivated the field operators to establish flare and field specific emissions factors. So in 2013, there are flare specific factors for a majority of the flares.

The field specific factors are estimated in a model developed by the Christian Michelsen Research (CMR) institute. The estimations are based on measurements with ultrasound of mass and volume on each flare.

There is several flares on a field but flare specific emissions factor are not estimated for all flares. For each field it is estimated a field specific emissions factor based on the flares with measurement data. For 2013, it is also calculated an average emissions factor of 2.637 kg CO₂ per Sm³ for all flares at all fields with measurements data.

Emissions factors 1990-2007

An annual emission factor is estimated from the field specific CMR measurements from 2013 weighted with the amount of flared gas for each field. The amount of gas for 1990-99 are from the Norwegian Petroleum Directorate and from Environmental Web/EPIM Environment Hub (EEH) for 2000-2015.

Emissions factors after 2007

For the years after 2007 there is information in the EU ETS about each single flare. At most fields there are a mixture of flares with CMR emission factors and default factors.

The emission factors used for calculation of emissions after 2007 is explained in sub-chapter “*Estimations of CO₂ after 2007*” above.

Table 3.31 presents the average EF for flaring off shore for the period 1990-2016.

Gas terminals

In Table 3.31, the CO₂ emission factors for flaring at one gas terminals are shown. The CO₂ emissions from flaring at that gas terminal were in 2016 a little bit more than 50,000 tonne.

Well testing

Emission factors used in the calculations for well testing are shown in Table 3.30. During the review of the 2008 inventory submission the expert review team raised question to that CH₄ and N₂O from well testing off shore were not included in the inventory. Norway then estimated the emissions of CH₄ and N₂O and presented the result for the expert review team. The emission estimates was for the first time included in the inventory in the 2010 submission.

Table 3.31. Emission factors for flaring of natural gas at off shore oil fields and one gas terminal on shore. 1990-2017

	Average emission factor for flaring at one gas terminal	Average emission factor for flaring off shore
	<i>tonne CO₂/tonne natural gas</i>	<i>kg CO₂/Sm³ natural gas</i>
1990	2.7	2.70
1991	2.7	2.66
1992	2.7	2.73
1993	2.7	2.80
1994	2.7	2.79
1995	2.7	2.69
1996	2.7	2.66
1997	2.7	2.69
1998	2.7	2.74
1999	2.7	2.75
2000	2.7	2.73
2001	2.7	2.65
2002	2.7	2.68
2003	2.7	2.63
2004	2.7	2.63
2005	2.7	2.62
2006	2.69	2.63
2007	2.67	2.66
2008	2.67	2.64
2009	2.67	2.85
2010	2.65	2.89
2011	2.76	2.93
2012	2.75	2.80
2013	2.62	2.71
2014	2.59	2.77
2015	2.53	3.02
2016	2.59	3.14
2017	2.54	3.14

Source: Norwegian Environment Agency/Norwegian Petroleum Directorate/Statistics Norway

Table 3.32. Emission factors for flaring in connection with well testing

Compounds (unit)	unit/tonne flared oil	Source	unit/kSm ³ flared natural gas	Source
CO ₂ (tonnes)	3.20	SFT (1990)	2.34	SFT (1990)
CH ₄ (tonnes)	0.0004 ¹	Same factors as for fuel oil used for boilers in manufacturing	0.00024	(IPCC 1997a)
N ₂ O (tonnes)	0.00003 ¹	OLF (2009)	0.00002	OLF (2009)
NM VOC (tonnes)	0.0033	OLF (2009)	0.00006	OLF (2009)
CO (tonnes)	0.018	OLF (2009)	0.0015	OLF (2009)

¹The Norwegian Oil Industry Association

3.4.4.5 Uncertainties and time-series consistency

The uncertainty in the amount of gas flared is in Rypdal and Zhang (2000) regarded as being low, ± 1.4 %, due to that there is a tax on gas flared and there is requirement by law that the gas volume flared is measured (Norwegian Petroleum Directorate 2001). The uncertainty in the CO₂ emission factor for flaring is ± 10 (Statistics Norway 2000).

The uncertainty in the amount of gas flared is in regarded as being low, ± 1.4 %, based on data reported in the emission trading scheme (Climate and Pollution Agency 2011a) and assumptions in Rypdal and Zhang (2000). The uncertainty in the CO₂ emission factor for flaring is ± 4.5 (Climate and Pollution Agency 2011a) and Rypdal and Zhang (2000).

The uncertainty in CH₄ and NMVOC emissions from venting and, hence, in the indirect emissions of CO₂, is much higher than for flaring.

All uncertainty estimates for this source are given in Annex II.

3.4.4.6 Source-specific QA/QC and verification

Statistics Norway gathers activity data used in the calculation from the Norwegian Petroleum Directorate. The figures are quality controlled by comparing them with the figures reported in the field operators annually report to the Norwegian Environment Agency and the Norwegian Petroleum Directorate and time series are checked.

Statistics Norway and the Norwegian Environment Agency perform internal checks of the reported data for venting from the field operators. Some errors in the time-series are usually found and the field operators are contacted and changes are made. The same procedure is followed to check the amount of gas reported as flared. The quality of the activity data is considered to be high due to that there is a tax on gas flared off shore. The Norwegian Petroleum Directorate has a thorough control of the amount of gas reported as flared. The oil and gas sector is included in the EU ETS from 2008.

3.4.4.7 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.5 CO₂ capture and storage at oil and gas production fields, 1C (Key Category for CO₂)

3.5.1 Overview

This chapter describes emissions related to the capture, transport and storage of CO₂ from natural gas produced at the gas-condensate field Sleipner Vest, including gas from the Gudrun field and the gas field Snøhvit.

Emissions occur primarily from venting of captured CO₂ when the injection facilities are not operating. Smaller emissions occur from a number of minor sources such as leakage from compressors. No emissions are reported from pipeline transport or from the CO₂ reservoirs.

The emissions are reported under 1C *CO₂ Transport and storage*. The emissions were until NIR 2016 reported in 1B2c together with indirect CO₂ emissions from CH₄ and NMVOC from venting and other fugitive emissions. The reporting in CRF Table 1.C also includes data on total CO₂ capture and injected amounts.

3.5.2 CO₂ capture and storage at the oil and gas production field Sleipner Vest

3.5.2.1 Description

The natural gas in the Sleipner Vest offshore gas-condensate field in the North Sea contains about 9 % CO₂. The CO₂ content has to be reduced to about 2.5 % to meet sales gas specifications. The CO₂ removed amounts to about 0,7 million tonnes per year.

When Sleipner Vest was planned around 1990 the considerations were influenced by the discussions about strategies to reduce greenhouse gas emissions and a possible national tax on CO₂-emissions (introduced in 1991 and extended in 1996). It was therefore decided that the removed CO₂ should be injected for permanent storage into a geological reservoir. The selection of an appropriate reservoir is essential for the success of geological storage of CO₂. In the search for a suitable reservoir the operators were looking for a saline aquifer with reasonable high porosity and a cap rock above to prevent leakage. Furthermore, the CO₂ should be stored under high pressure – preferably more than 800 meters below the surface. Under these conditions CO₂ is buoyant and less likely to move upwards than CO₂ in gaseous form.

The Utsira Formation aquifer, which is located above the producing reservoirs at a depth of 800 – 1000 meters below sea level, was chosen for CO₂ storage because of its shallow depth, its large extension (which guarantees sufficient volume), and its excellent porosity and permeability (which is well suited for high injectivity). The formation is overlain by a thick, widespread sequence of Hordaland Group shales, which should act as an effective barrier to vertical CO₂ leakage, see Figure 3.18.




Figure 3.18. CO₂ capture from Sleipner Vest well stream and storage at Sleipner.

Source: Statoil

The reservoir was characterised by reservoir information such as seismic surveys and information from core drillings.

In the Sleipner case it was very important to locate the injection well and the storage site such that the injected CO₂ could not migrate back to the Sleipner A platform (SLA) and the production wells. This will both prevent corrosion problems in the production wells and minimise the risk of CO₂ leakage through production wells. The injection point is located 2.5 km east of the Sleipner A platform. Migration evaluations have been based on the Top Utsira map (see Figure AVI-2 in Annex IV) with the CO₂ expected to migrate vertically to the sealing shales and horizontally along the saddle point of the structure. This will take the CO₂ away from other wells drilled from the Sleipner platform. A more detailed description of the reservoirs suitability for long term CO₂ storage is given in Annex IV.

The field and the injection program has been in operation since 1996. Statoil monitors the injected CO₂ with respect to leakages by 4 D seismic surveys.

Investigations carried out so far show that the injected CO₂ is kept in place without leaking out. In case unexpected CO₂ movements take place beyond the capture rock in the future it can be registered by the monitoring techniques. Table 3.33 gives the amount of CO₂ injected since the project started in 1996.

Table 3.33. CO₂ from the Sleipner field injected in the Utsira formation.

Year	CO ₂ (ktonnes)	Year	CO ₂ (ktonnes)	Year	CO ₂ (ktonnes)
1996	70	2004	750	2012	842
1997	665	2005	858	2013	702
1998	842	2006	820	2014	658
1999	971	2007	921	2015	707
2000	933	2008	814	2016	632
2001	1 009	2009	860	2017	679
2002	955	2010	743		
2003	914	2011	929		

Source: Statoil/The Norwegian Environment Agency

When the injection is stopped due to maintenance or any unplanned reasons, the captured CO₂ is vented to the atmosphere. The amount of CO₂ vented to the atmosphere is included in the greenhouse gas inventory reported under 1C1a. The emissions from venting are presented in Table 3.34.

Table 3.34. Emissions of CO₂ vented from the Sleipner Vest CO₂ –injection plant due to inaccessibility of the injection facility.

Year	CO ₂ (ktonnes)	Year	CO ₂ (ktonnes)	Year	CO ₂ (ktonnes)
1996	81.0	2004	21.4	2012	5.9
1997	29.0	2005	6.2	2013	5.0
1998	4.2	2006	2.5	2014	5.4
1999	9.1	2007	6.4	2015	0.8
2000	8.3	2008	13.6	2016	4.6
2001	3.1	2009	4.6	2017	0.8
2002	87.6	2010	0.9		
2003	23.9	2011	2.4		

Source: The Norwegian Environment Agency

The status by 31.12.2017 is that 17.3 million tonnes CO₂ have been injected and stored in the Utsira Formation and 0.25 million tonnes CO₂ have been vented. Figure 3.19 shows the yearly injected and vented volumes for the entire injection period on Sleipner.




Figure 3.19. Injected and vented CO₂ at Sleipner Vest.

Source: Norwegian Environment Agency

Diffuse emissions from the CO₂-capture plant (amineabsorber) and the CO₂-compressor are estimated to about 1,0 kt CO₂/year and these figures are included in CRF Table1.C.

The compressorized CO₂ is transported by pipeline to the well head (injection well). The transport distance is 350 m and is controlled by pressure monitoring.

3.5.2.2 Methodological issues

The reported data covers emissions to the atmosphere e.g. when the injection system is out of operation. These emissions are determined by continuous metering of the gas stream by VCONE-meter. The reported amounts of CO₂ injected in the Utsira formation are based on continuous metering of the gas stream by orifice meter. The composition of the CO₂-stream is stable, about 98% CO₂ and the remaining 2% mainly methane and heavier hydrocarbons.

The diffuse emissions are estimated on the basis of equipment specific leakage factors. CO₂-dectectors are monitoring almost all potential leakages sources (e.g. flanges).

The Sleipner CO₂-injection project is considered as the first industrial-scale, environmentally driven CO₂-injection project in the world. In order to document what happens with the CO₂ a European research project initially called SACS ("The saline aquifer carbon dioxide storage project") was organized around it. The SACS project ended in 2002 and was succeeded by the ongoing EU-co-funded CO2STORE and ECO2. The projects have run parallel to the development of Sleipner Vest and have special focus on monitoring and simulation. Research institutes and energy companies from several countries participate in the projects. The core of the projects has been to arrive at a reasoned view of whether carbon dioxide remains in the Utsira sand and whether developments in this formation can be monitored. The migration of carbon dioxide through the aquifer is recorded by seismic surveys. Base line 3D seismic data were acquired in 1994, prior to injection, and the first repeat survey was acquired in 1999, when some 2.28 mill tonnes of CO₂ had been injected into the reservoir. This was followed by 4-D seismic surveys in 2001, 2002, 2004, 2006, 2008, 2010, 2013 and

2016. The monitoring methodology and the results of the monitoring are described in Annex IV written by Statoil.




Figure 3.20 Results of seismic monitoring Sleipner Vest, 1998-2016. Accumulated amplitudes on the differences between 1994 and respectively 2001, 2004, 2006, 2008, 2010, 2013 and 2016. Source: Statoil

The stored CO₂ has been monitored using time lapse seismic to confirm its behaviour and evaluate

- whether any of it has leaked into the overburden seal, the ocean or the atmosphere, or
- whether any of it has migrated towards the Sleipner installations, potentially leading to corrosion problems for well casing

The results show that neither of these eventualities has occurred. is no sign of CO₂ above the top of Utsira Formation.

Results from the projects are published in several reports and articles such as:

- EU (2002)
- Arts et al. (2005)
- Chadwick et al. (2004)
- Chadwick et al. (2005)

A more detailed list of publications and presentations is given in Annex IV. The project has confirmed that sound waves reflect differently from carbon dioxide and salt water. Comparing seismic data collected before and after injection started has allowed researchers to show how CO₂ deep inside the

Utsira formation migrates (see Figure AVI-5 in Annex IV). It is held under the layer of shale cap rock, 80 metres thick, which covers the whole formation. This extends for several hundred kilometres in length and about 150 kilometres in width.

The time-lapse seismic data clearly image the CO₂ within the reservoir, both as high amplitude reflections and as a pronounced velocity pushdown (see Figure 3.20 and Figure AIV4 in Annex IV).

The data also resolve a vertical CO₂ chimney, which is regarded the primary feeder of CO₂ in the upper part of the bubble.

Flow simulation models, which match the 4D seismic data reasonably well, have been used to predict the CO₂ behaviour, see Figure 3.21.




Figure 3.21. Flow simulation of CO₂ Sleipner Vest.

Source: Statoil

The results from the simulations indicate that the cap rock shales provide a capillary seal for the CO₂ phase.

There is no seismic indication of faults within the upper part of the reservoir, and no indications of leakage into the capture rock.

The time-lapse seismic images clearly show the development of the CO₂ plume, and have been used to calculate the amount of CO₂ in the reservoir. The volume calculated from the observed reflectivity and velocity pushdown is consistent with the injected volume.

Other monitoring methods Statoil is running are monitoring the injected CO₂, gravimetric monitoring, pressure measurements and well monitoring. For more details see Annex IV.

3.5.2.3 Uncertainties

The reported data covers emissions to the atmosphere e.g. when the injection system is out of operation. The accuracy in these measurements made by VCONE-meter is $\pm 5\%$. The orifice meter used to meter the amount of CO₂ injected in the Utsira formation have $\pm 3\%$ accuracy. So far there has not been detected any leakage from the storage.

3.5.2.4 Category-specific QA/QC and verification

The results are promising and the injected gas remains in place. Storage of CO₂ is regulated by the Pollution Control Act and the specific regulations of geological storage of CO₂ (entered into force January 1, 2016). Pursuant to the Pollution Control Act and the specific regulations, the operator shall hold a permit. According to the permit conditions Statoil shall monitor the CO₂-storage. Statoil reports annually the amount of CO₂ injected and emitted to The Norwegian Environment Agency. The injected CO₂ is so far proven to be removed from the atmosphere and hence, it is not reported as emissions in the emission inventory. When the injection of CO₂ is stopped for maintenance purposes, the operator pays a CO₂-tax for the emissions. From 2013 these emissions are included in the EU-ETS. In the national emissions inventory the amount of CO₂ vented is reported under 1C2a - Injection.

3.5.2.5 Category-specific recalculations

There have been no recalculations performed for this source category this year.

3.5.2.6 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.5.3 CO₂ capture and storage at Hammerfest LNG/the gas-condensate production field Snøhvit

3.5.3.1 Description

The natural gas in the Snøhvit gas-condensate subsea field in the Barents Sea contains about 5-7.5 % CO₂. Prior to the LNG production the CO₂ has to be removed to avoid it freezing out in the downstream liquefaction process. The facilities for separation and injection of CO₂ are placed onshore at the Hammerfest LNG process plant at Melkøya.

An amine absorption unit performs the separation. The recovered CO₂ is condensed and recompressed before transported by a subsea pipeline and re-injected into Tubåen and Stø reservoir. A schematic of the CO₂ re-injection system is shown in Figure 3.22 About 0.73 Mtonnes CO₂ are removed from the feed gas every year at full production. During the expected lifetime of the field, about 23 million tonnes CO₂ from the feed gas will be removed and re-injected.




Figure 3.22. Snøhvit Field overview.

Source: Statoil

Reservoir

Several geological structures in the Snøhvit area were evaluated for disposal of CO₂. The four structures identified as possible candidates for CO₂ storage were Marcello, 7122/2-1 structure, 7122/7-1 Goliath and the water bearing Tubåen Formation on the Snøhvit and Albatross fields. Marcello and the 7122/2-1 structure were considered as immature for CO₂ storage for the Snøhvit CO₂ storage project because the reservoir data was not sufficiently detailed and there are no current plans for exploration drilling (ref: Plan for Development and Operation). The Tubåen formation was chosen as the primary storage location.

Hammerfest LNG (former Snøhvit LNG Statoil) was granted a permit pursuant to the Pollution Control Act to inject 730 000 tonnes of CO₂ per year into the geological formation, Tubåen in 2004. The permit was issued by the Norwegian Environment Agency. The production started in 2008.

In March 2011, the injection point was moved from Tubåen to the Stø reservoir, due to lower injectivity in Tubåen than expected.

The Snøhvit Fields are not very complex structurally. Two well-defined fault directions, E-W and N-S, define most of the major structures. Minor internal faulting is present within the major structures.

Tubåen formation is a saline aquifer lying around 100-200 metres below the gas cap at Snøhvit. Tubåen formation is water filled and has a thickness between 45 and 75 metres. Core samples show that the formation consists of relatively pure quartz sand. The porosity and permeability are 10-16% and 200-800 md, respectively. The formation is bounded by large faults on all sides. Formation depth is 2600 m below sea level.

Stø water zone formation, which is the bottom of the current producing gas reservoir, was perforated for injection. The water zone has a thickness of 42 metres. Core samples show that the formation consists of relatively sand. The porosity and permeability are 15% and 400md, respectively (Table 3.35) Formation depth is 2450 m below sea level.

The geophysical, geological and petrophysical evaluations are based on 19 exploration wells and 10 development wells within the area. The data available from these wells are generally of good quality, including logs, core data and pressure data.

The reservoir was characterised by reservoir information such as seismic surveys and information from core drilling.

Table 3.35. Key parameters for injection well F-2 H and Tubåen reservoir at the Snøhvit field. Stø reservoir pressure is being depleted by field production.

Key Parameters	Tubåen	Stø
Initial reservoir pressure	288 bar	255 bar
Initial temperature	98 C	98 C
Porosity	10-16%	15%
Permeability	200-800 md	400 md
Reservoir depth	2600 m	2450 m
Water depth at F-template	330m	330m
Length pipeline from Melkøya	152km	152km

Location of the CO₂ injection well F-2 H.

The CO₂ injection well is located at the F-segment at the western part of the Snøhvit reservoir (Figure 3.23). The injection pipeline is 152 km long (Figure 3.22). A new injection well, located in the G-segment, has been established in 2016.




Figure 3.23. Location of the CO₂ well at the Snøhvit field.

Source: Statoil

At the beginning, to keep the CO₂ as deep as possible, it was decided to perforate the mid and lower part of Tubåen as shown in Figure 3.24. Since injection was changed to Stø, additional perforations were done in the bottom of Stø as shown in Figure 3.24.




Figure 3.24. Cross-section of F-segment where CO₂ is injected, Snøhvit field formation.

Source: Statoil

CO₂ injected and vented

The status by 31.12.2017 is that 1 087 ktonnes CO₂ have been injected into the Tubåen Formation and 3 936 ktonnes have been injected into the Stø Formation, and 561 ktonnes CO₂ have been vented (Table 3.36). CO₂ venting occurs when the CO₂ reinjection system has to be shut down. The maximum vent rate is almost equal to the CO₂ removal flow rate. A separate vent stack for the CO₂ is provided at the plant.

Table 3.36. Injected and vented CO₂ Hammerfest LNG/Snøhvit field.

Year	CO ₂ injected (ktonnes)	CO ₂ vented (ktonnes)	Year	CO ₂ injected (ktonnes)	CO ₂ vented (ktonnes)
2007	0	71	2013	469	27
2008	197	93	2014	587	37
2009	308	50	2015	679	39
2010	460	93	2016	750	4
2011	403	87	2017	680	4
2012	490	55			

The following Figure 3.25 shows the yearly injected at in the Tubåen/Stø formation at the Snøhvit field and vented volumes for the injection period at Hammerfest LNG. These figures are reported to the Norwegian Environment Agency on a yearly basis.




Figure 3.25. Injected and vented CO₂ at the Snøhvit field and Hammerfest LNG.

Source: Statoil

3.5.3.2 Methodological issues

CO₂ injection well specification

The completion design basis for the CO₂ injector at Tubåen/Stø depth is a perforated 7" liner. A downhole pressure and temperature gauge is installed.

CO₂ re-injection system




Figure 3.26. Schematic of the CO₂ injection system in the Snøhvit area.

Source: Statoil

CO₂ is most likely re-injected as a single phase (liquid condition in the pipeline from the export pump to the well head, transformed to supercritical condition in the reservoir where the temperature is higher).

CO₂ venting to atmosphere

The reported data covers CO₂ emissions to the atmosphere, e.g. when the injection system is out of operation. These emissions are measured by a venturi flow meter.

Flow metering of the well stream to the CO₂ injector is measured by an orifice meter.

Gas composition of injected or vented gas from the CO₂ injector is controlled by analyses. This is primarily done as a quality assurance of the CO₂ removal system (system 22). Analyses have shown that composition is 99.549 weight % CO₂, 0.0066 weight % H₂S, 0.331% CH₄ and 0.088 weight % NMVOC. It has been agreed that in the reports to the environmental authorities, ventilated gas shall be reported as 100% weight CO₂.

Diffuse sources and pipeline transport

Diffuse emissions from the CO₂-capture plant are estimated to about 1,0 kt CO₂/year and these figures are included in CRF Table1.C.

For pipeline transport of CO₂, emissions are reported as Not Estimated. The IPCC tier 1 method gives a medium emission factor of 0.0014 Gg per year and per km of transmission pipeline. The pipeline is 152 km long, corresponding to an emission estimate of 0.2 kt CO₂. However, based on our best knowledge we anticipate that this would overestimate emissions from the pipeline.

Firstly, the main sources of emissions are likely to be equipment at the ends of the pipeline. In the Snøhvit case, emissions from equipment at the input end is included with reported emissions from the Hammerfest plant.

Secondly, the pipeline and injection well are continuously monitored by pressure monitoring (downhole well, choke, export pump). The pipeline and injection well are also subject to-acoustic deep water survey and visual inspection by Remote Operated Vehicle in order to detect any sign of corrosion or irregularities which may cause leakages. 2D and 3D seismic surveys are carried out on a regular basis. Based on 3D seismic data 4D seismic is used to monitor CO₂ movement in vertical and horizontal direction, detect leakages or unexpected migration of CO₂ in the geological formation.

Reservoir monitoring by seismic

4D seismic monitoring was carried out in 2011, 2012 and 2014 in order to monitor the CO₂ plume migration inside the Stø formation and its movement towards the gas zone. Strong focus has also been on optimizing the reservoir simulation model in order to match the 4D observation. Reservoir simulation model is the main tool for predicting CO₂ flow in the future.

The strong 4D signal is mainly related to the fluid replacement effect, CO₂ replacing water. Some of the 4D signal close to the injector is also most likely related to thermal fracturing because of cold CO₂ injection. The CO₂ follows the Stø2 layer and does not seem to migrate up into Stø 3 due to the much lower permeability in Stø3 compared to Stø 2.




Figure 3.27 The upper figures show the differences from 2009 to 2012. The lower figures show 4D amplitude maps on CO₂ plume for 2009-2011 (left) and 2009-2012 (right).
Source: Statoil




Figure 3.28. Seismic 4D amplitude map from 2011, showing a clear anomaly around the CO₂ injector Pressure/temperature gauge, reservoir modeling and prediction of reservoir performance in Tubåen.

The pressure development in the injection well is monitored on a daily basis by using data from the pressure and temperature (PT) gauge installed in the well. Due to problems during drilling there is diameter restriction in the well and the PT gauge had to be installed about 600 m above the reservoir. Actual bottom hole pressure is estimated based on gauge measurements and CO₂ PVT (pressure, volume, temperature). An Eclipse 300 Compositional simulation model is used for prediction pressure development in the well. In this model CO₂ is injected into the water filled Stø reservoir. Using this model, it has proven to be easy to match the CO₂ plume size/shape geometry in this model with time-lapses seismic data. A weakness of the model is that it does not include temperature and other advanced simulation physical effects. Temperature effects are likely in the near well area as CO₂ at 21 °C is injected into a reservoir of initially 91 °C.

Since mid 2011 CO₂ in liquid phase has been injected to Stø water saturated formation. The well has shown that its ability to receive injected CO₂ is stable. This is confirmed by weekly monitoring.

As can be seen from Figure 3.29, the reservoir pressure (green line) has depleted since May 2011 until December 2014. This is due to production of the gas zone above the water zone, from the gas zone.




Figure 3.29. History pressures and volume injection into Stø formation.
Source: Statoil

Gravimetric monitoring

A baseline gravity and seafloor subsidence monitoring survey was carried out over the Snøhvit and Albatross fields in June 2007. The closest benchmark is 419 m from the CO₂ injection well. A total of 76 sea floor benchmarks were deployed at the start of the survey, and relative gravity and depth was measured. A new gravity monitoring was carried out in spring 2011. Comparison of 2011 and 2007 gravity measurements confirmed the prognoses.

3.5.3.3 Category-specific QA/QC and verification

Operators for CO₂-storage projects have to apply for a permit pursuant to the Pollution Control Act. In accordance with the permit provisions, Statoil has implemented system for monitoring the CO₂-storage. So far there is no sign of emissions to the water column or the atmosphere from the injected CO₂. Hence the CO₂ injected is not reported as emissions in the emission inventory. Statoil pays a CO₂-tax for the emissions when the injection facility is out of operation due to maintenance etc.

From 010113 these emissions are also regulated under the emission trade scheme (EU-ETS). The emissions of CO₂ and the amount of CO₂ injected are reported to the Norwegian Environment Authority. In the emissions inventory the amount of CO₂ vented at Hammerfest LNG (Snøhvit CO₂ storage project) – is reported under 1C2a - Injection 1B2c.

Statoil performs internal QA/QC for the ongoing CO₂ studies.

3.5.3.4 Category-specific recalculations

There have been no recalculations performed for this source category this year.

3.5.3.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

3.5.3.6 Activities and future plans

Stø formation was perforated in April 2011 and is currently injecting in this zone. Injection was monitored every week by a fall-off test performed during stable conditions. During 2014 monitoring was done on a monthly basis by the fall-off test. Injection of CO₂ has been stable and there are no well integrity issues related to operation of the well.




Figure 3.30. CO₂ injector current completion.

Source: Statoil

The challenge of production CO₂ from Snøhvit field has led to a great effort to find solutions that makes the CO₂ injection as robust as possible. The authorities have been kept informed about the situation and the activities and measures planned. A monitoring program covering the period 2011-2020 has been submitted to the environmental authorities.

A new injector well has been established and was put into operation 4Q 2016. This well has been the primary injection well in 2017.

Based on the experience using 4D seismic monitoring in 7120/F-2H it is very likely that 4D seismic monitoring will work well for the new CO₂ injector that is located in the G-segment.

3.5.3.7 CO₂ projects outside Statoil ASA using Snøhvit data

The EU project CO2ReMoVe plans to perform a complete performance and risk assessment for the Snøhvit project by complementing the work done under the CASTOR umbrella. Particular attention will be paid to potential vertical CO₂ migration to the upper gas field and lateral migration, potential flow through deteriorated wells and through undetected faults. The geochemical interaction between CO₂, fluids and rock and coupling with geomechanical effects will be investigated.

Data from Snøhvit is released to the FME SUCCESS Centre (Centre for Environmental Friendly Energy Research; Subsurface CO₂ Storage- Critical Elements and Superior Strategy). Based on this information, specific research tasks may be defined.

3.6 Cross-cutting issues

3.6.1 Sectoral versus reference approach

In the review of the Norwegian greenhouse gas inventory submitted in 2018 the ERT expert team expressed concerns about the size of the differences in energy use and emissions as estimated by the reference and the sectoral approach. In response, it is an ongoing project aimed at reducing the gap between the RA and the SA to an acceptable level and providing a better explanation of the remaining differences. The results from the project are expected to be ready for the 2021 inventory. The project is described in more detail in chapter 10, section 10.4.1 and 10.4.2. The result of the estimation with the two methods is shown in Table 3.37.

There are large differences between the output from RA and SA, both for the energy consumption data and the CO₂ emissions. The difference between the fuel consumption in the RA and SA ranges from about -23 % to + 40 %. The differences for CO₂ emissions ranges between -16 and +45 %. The highest discrepancy for CO₂ is in 1999-2001 and in 2004-2006. For 2017, the difference for CO₂ is 4.1 %. The large discrepancies are primarily due to statistical differences in the energy balance.

A detailed analysis of the relationship between the RA and SA and the energy balance is given in annex XI. The main conclusion is that the difference between the energy consumption in RA and SA is primarily due to statistical differences in the energy balance. In addition, a number of other smaller differences were identified. The remaining difference between RA and SA after adjusting for these items is within +/- 4 %. The reference approach may be an important tool for verification of the sectoral approach used in the inventory. The analyses undertaken in the present and the previous NIRs have shown that the difference between RA and SA is mainly due to the statistical difference in the energy balance, and that important parts of the consumption block in the EB are unlikely to have major completeness issues. If the statistical differences are due to problems in the supply block of the balance, then resolving these problems will only affect the RA, but not the SA and the reported emissions.

A large statistical difference relative to national consumption is not unreasonable in Norway, given the large production and export share. With reference to the IPCC Guidelines 2006, Volume 2 Energy, chapter 6.8 "It should be noted that for countries that produce and export large amounts of fuel, the uncertainty on the residual supply may be significant and could affect the Reference Approach".

Table 3.37 Comparison of fuel consumption and CO₂ emission data between the Reference Approach¹ (RA) and the Sectoral Approach (SA). 1990-2017.

Year	Fuel consumption			CO ₂ emissions		
	RA, apparent consumption (PJ)	SA (PJ)	Difference RA-SA (%)	RA (Gg)	SA (Gg)	Difference RA-SA (%)
1990	294	379	-22.5	21 599	25 809	-16.3
1991	382	377	1.5	26 893	25 531	5.3
1992	343	386	-11.1	23 749	26 160	-9.2
1993	356	399	-10.6	24 652	26 947	-8.5
1994	367	418	-12.2	25 933	28 310	-8.4
1995	389	425	-8.4	26 887	28 842	-6.8
1996	386	461	-16.2	27 200	31 512	-13.7
1997	410	466	-12.0	28 695	31 579	-9.1
1998	518	466	11.0	36 075	31 593	14.2
1999	563	465	21.0	39 982	31 859	25.5
2000	633	453	39.7	44 841	30 934	45.0
2001	584	479	21.9	39 950	33 038	20.9
2002	491	486	1.0	34 370	33 280	3.3
2003	535	507	5.5	37 316	34 688	7.6
2004	620	507	22.3	44 146	34 624	27.5
2005	607	501	21.1	42 875	34 429	24.5
2006	651	527	23.7	45 863	35 480	29.3
2007	524	531	-1.3	35 583	35 624	-0.1
2008	596	533	11.8	40 646	35 208	15.4
2009	556	542	2.5	37 928	35 819	5.9
2010	631	564	11.9	43 230	37 137	16.4
2011	498	548	-9.3	34 373	36 362	-5.5
2012	546	544	0.4	37 177	35 879	3.6
2013	584	544	7.2	40 137	35 731	12.3
2014	592	551	7.5	40 600	35 915	13.0
2015	601	553	8.6	41 026	36 002	14.0
2016	499	541	-7.7	33 261	35 213	-5.5
2017	523	533	-2.0	35 915	34 509	4.1

¹ Apparent energy consumption (excluding non-energy use, reductants and feedstocks).

Source: Statistics Norway/Norwegian Environment Agency

3.6.2 Quality controls within reference and sectoral approach

For several years there has been a problem regarding statistical difference between the supply and use of petroleum products in the Norwegian Energy Balance. This should not be unexpected from a country exporting almost 90 % of its petroleum products. Just minor discrepancies between production and export on the supply side of the balance may result in significant imbalances with the use side figures. There has, however, been a tendency for a positive bias in the statistical difference for a long time, which has caused uncertainty whether the domestic use of petroleum products might have been underestimated. Therefore a project was launched in 2012 in order to address the bias and make corrections if possible.

In Norway, most of the produced petroleum products are *primary*¹¹, while most of the domestic use relates to *secondary* petroleum products. Hence, separate energy balances for *primary and secondary* petroleum products were elaborated in the project. To further increase the transparency, more detailed product categories and one transfer item were elaborated as well. New data on primary petroleum products were identified and collected, in order to establish alternative export figures and new revision controls. No alternative data on secondary petroleum products was found, and hence these products were not prioritized.

The new export data is consistent with the production figures, and most discrepancies can be discussed with the data owner without breaking the confidentiality rules. Hence, the new data provides a solid basis for quality control. A similar crosschecking of the original export data from the external trade statistics (ETS) is tedious, or for some products almost impossible. Moreover, corrections must be made in the energy balance to obtain consistency between the export and production data.

All new data is readily available from the Norwegian Petroleum Directorate, one terminal and one pretreatment facility, respectively, and suitable for routine revision control. Compiling them for use in the energy balance is relatively simple. One dataset is not distributed by destination country, and in international reporting of country specific figures this data should instead be used to adjust the current ETS export data.

Based on new and original data, two alternative detailed energy balances for primary petroleum products were established, for revision purposes. Most causes of statistical difference for primary petroleum products were found due to the new data and revision methods, and the statistical differences for these products were significantly reduced.

The major part of the recommendations from the analysis are now implemented in the energy balance.

- New annual statistics on sales of petroleum products, based on detailed information on every delivery sent by the companies (SA).
- New collection of supplementary data, including detailed shipment data from the Norwegian Petroleum Directorate, export from one terminal and one crude petroleum pretreatment plant (RA).

¹¹ *Primary* means unrefined (incl. pretreatment like fractionation and stabilization). *Secondary* means refined into finished products or semi-manufactured products for use as raw material in manufacturing

- New revision methods for production, export and stock change micro data on all primary petroleum products (excl. dry gas), applying both former and new data. The supplementary data are as well used to estimate the export of crude oil according to a new method. The crude oil figures are cross checked at most detailed level against Statistics Norway's external trade statistics (RA).
- Cross check of dry gas micro data against Statistics Norway's external trade statistics (RA).
- Stock data from one previously missing terminal (RA).
- The improvements are implemented in the energy balance for 2015, and partly for 2014.
- LPG from one refinery was found to be propylene, which is not a typical energy product. The propylene is used as raw material in chemical industry. The use as raw material is not part of the energy balance, and hence a statistical difference occurs. This will be solved in the new energy balance (RA).

The new revision method has revealed underreporting of shipments of Norwegian NGL/LPG products from one UK sales point and a small underreporting of shipments of crude oil from another UK sales point, rendering the statistical difference for these product categories still on the positive side on average.

All improvements on statistical differences concern primary petroleum products, while no methods for reducing the statistical difference of refined petroleum products were found due to lack of supplementary data.

Table 3.38 Statistical difference for primary petroleum products in the energy balance. PJ. 2015.

		NGL/LPG	Natural gas	Crude oil
1	Primary energy production	484.2	4551.9	3318.8
1.1.1	Prod. of prim. energy carriers	409.5	4536.2	3318.8
1.1.2	Flaring	-	15.7	-
1.2	Prod. of sec. energy carriers	74.7	-	-
2	Imports	37.8	-	76.0
3	Exports	402.9	4354.0	2845.9
4	International bunkers	-	1.6	-
5	Changes in stocks (+ = net decrease, - = net increase)	-1.1	-3.9	41.1
6	Total energy supply (1+2-3-4+5)	119.0	192.4	590.1
7	Transformation	34.8	13.9	562.0
8	Energy industries own use	0.6	194.5	-
9	Losses in transport and distribution	-	0.2	-
11	Non energy use	36.7	21.2	-
13	Statistical differences (6+7-8-9-10)	34.6	-55.9	28.0
12	Final energy consumption, excl. non-energy use	12.3	18.6	-

New platform for energy balance and energy accounts: Statistics Norway has had a project running over several years on improving the energy balance and the energy accounts. The main phase was published in 2017 with results for 2010-2016 (Statistics Norway (Annually-a), Hendriks et al. (2017)). The project entails both new technical solutions and methodological changes. The has streamlined national and international reporting of energy statistics, and increased consistency in the

different reportings. The relationships with other statistics has been improved, in particular with respect to emissions to air from energy use. It has also improved transparency between energy and emission statistics.

3.6.3 Feedstocks and non-energy use of fuels

Emissions from the use of feedstock are according to the IPCC guidelines generally accounted for in the industrial processes sector in the Norwegian inventory. By-products from processes like blast furnace gas and fuel gas from ethylene cracking that are sold and combusted are accounted for and reported under the energy sector.

Table 1Ad Feedstocks and non-energy use of fuels in the CRF is filled in with fuels that are used as feedstock or any other non-energy use according to the energy balance. The data in the energy balance are adjusted with respect to reducing agents that are considered as fuel use in the energy balance, but accounted as IPPU emissions in the inventory.

The table also includes information of the amount of carbon excluded. The excluded amount corresponds to the fuel quantity except for emissions that are included as energy combustion in 1A. These emissions comprise the following categories, which are further described in section 3.2.11:

- A fraction of non-fuel use of gasoline, gas/diesel oil and residual fuel oil is assumed to be emitted to air. The emissions are reported under 1A5a *Non-fuel use*.
- Emissions from use of lubricants in 2-stroke engines are reported under 1A5b. Emissions from other lubricants is on the other hand reported in IPPU under 2D2 *Lubricants* and are as such accounted as carbon excluded.

The following table gives balances relevant fuels. The table shows that the fuel amounts reported in as fuel combustion in 1A and the carbon excluded from the Reference approach add up to the total fuel quantity for feedstock and other non-fuel use.

National Inventory Report 2019 - Norway

Table 3.39 Balances for fuels for which emissions from feedstock and non-fuel use are partly reported in the Energy sector. 1990-2017

		1990	1995	2000	2005	2010	2015	2016	2017
1AD Feedstock etc., reported Fuel quantity									
Gasoline	TJ	-	11	15	-	-	-	-	-
Gas Diesel Oil	TJ	-	-	-	-	-	-	-	-
Residual fuel oil	TJ	-	570	611	-	-	-	-	-
Lubricants	TJ	4 422	3 457	3 578	3 734	1 648	1 769	2 090	2 010
Reported in Energy Sectoral Approach (1A), Activity data									
1A5a Non-fuel use	TJ	-	124	138	348	0.1	0.1	0.1	0.1
of which Gasoline	TJ	-	6	8	-	-	-	-	-
of which Gas Diesel Oil	TJ	-	55	63	247	0.1	0.1	0.1	0.1
of which Residual fuel oil	TJ	-	63	67	101	-	-	-	-
1A5b Lubricants	TJ	84	78	69	58	40	39	39	39
Remainder, excluded from Reference Approach									
Gasoline	TJ	-	6	8	-	-	-	-	-
Gas Diesel Oil	TJ	-	55	63	247	0.1	0.1	0.1	0.1
Residual fuel oil	TJ	-	507	544	-	-	-	-	-
Lubricants	TJ	4 338	3 379	3 509	2 676	1 608	1 730	2 051	1 971
Remainder, excluded from Reference Approach as reported in 1AD									
Gasoline	kt C	-	0.11	0.15	-	-	-	-	-
Gas Diesel Oil	kt C	-	1.11	1.26	4.95	0.002	0.002		
Residual fuel oil	kt C	-	10.91	11.7	17.57	-	-	-	-
Lubricants	kt C	45.56	30.60	22.57	17.56	15.57	9.58	11.54	11.34

3.7 Memo items

3.7.1 International bunkers

3.7.1.1 Description

Emissions from international marine and aviation bunker fuels are excluded from the national totals, as required by the IPCC Guidelines (IPCC 2006). The estimated emission figures are reported separately and are presented in Table 3.40.

In 2017 CO₂ emissions from ships and aircraft in international traffic bunkered in Norway amounted to a total of 2.2 million tonnes, which corresponds to 4.2 % of the total Norwegian CO₂ emissions.

During the period 1990-2017, emissions of CO₂ from marine bunkers decreased by 80 %. The emissions have varied greatly in this period and reached a peak in 1997. Thereafter there has been a descending trend in emissions.

The CO₂ emissions from international air traffic bunkered in Norway was in 2017 1.7 million tonne. The emissions in 2017 was almost three times as high as in 1990. In 2017 the emissions were 5 % higher than in 2016.

Table 3.40 Emissions from ships and aircraft in international traffic bunkered in Norway, 1990-2017. 1000 tonnes. CO₂ in Mtonnes.

	Aviation							Marine						
	CO ₂	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂	CO ₂	CH ₄	N ₂ O	NO _x	CO	NM VOC	SO ₂
1990	0.6	0.0	0.0	2.5	1.1	0.2	0.1	2.3	0.2	0.1	41.1	2.2	1.7	13.8
1991	0.6	0.0	0.0	2.3	1.2	0.3	0.1	2.0	0.1	0.1	35.9	2.0	1.5	11.6
1992	0.6	0.0	0.0	2.4	1.4	0.3	0.1	2.2	0.2	0.1	38.6	2.1	1.6	12.4
1993	0.6	0.0	0.0	2.4	1.5	0.4	0.1	2.1	0.2	0.1	38.1	2.1	1.6	11.7
1994	0.6	0.0	0.0	2.3	1.6	0.5	0.1	2.1	0.1	0.1	37.0	2.0	1.6	9.7
1995	0.6	0.0	0.0	2.1	1.5	0.5	0.1	2.3	0.2	0.1	40.8	2.2	1.7	12.4
1996	0.7	0.0	0.0	2.5	1.7	0.5	0.1	2.6	0.2	0.1	46.4	2.5	1.9	14.7
1997	0.7	0.0	0.0	2.8	1.7	0.4	0.1	3.2	0.2	0.1	57.2	3.1	2.4	19.2
1998	0.8	0.0	0.0	2.9	1.6	0.4	0.1	3.1	0.2	0.1	56.1	2.8	2.3	15.6
1999	0.9	0.0	0.0	3.4	1.7	0.3	0.1	2.9	0.2	0.1	51.9	2.6	2.2	13.2
2000	0.9	0.0	0.0	3.2	1.4	0.1	0.1	2.8	0.2	0.1	51.4	2.6	2.1	11.3
2001	0.8	0.0	0.0	3.0	1.3	0.1	0.1	2.8	0.2	0.1	50.5	2.5	2.1	13.4
2002	0.7	0.0	0.0	2.7	1.1	0.1	0.1	2.3	0.2	0.1	41.2	2.1	1.7	7.7
2003	0.7	0.0	0.0	2.8	1.1	0.1	0.1	2.2	0.2	0.1	38.5	2.0	1.6	9.7
2004	0.8	0.0	0.0	3.1	1.2	0.1	0.1	2.1	0.2	0.1	37.9	1.9	1.6	9.5
2005	0.9	0.0	0.0	3.6	1.3	0.1	0.1	2.6	0.2	0.1	45.6	2.4	2.0	10.2
2006	1.1	0.0	0.0	4.4	1.6	0.1	0.1	2.4	0.2	0.1	41.5	2.2	1.8	5.7
2007	1.1	0.0	0.0	4.6	1.6	0.1	0.1	2.4	0.2	0.1	41.8	2.2	1.8	7.0
2008	1.1	0.0	0.0	4.6	1.5	0.1	0.1	2.2	0.2	0.1	37.2	2.0	1.7	7.2
2009	1.1	0.0	0.0	4.5	1.4	0.1	0.1	2.1	0.2	0.1	35.0	1.9	1.6	6.1
2010	1.3	0.0	0.0	5.5	1.7	0.1	0.1	1.6	0.1	0.0	25.6	1.5	1.2	5.5
2011	1.4	0.0	0.0	6.0	1.8	0.2	0.1	1.6	0.1	0.0	25.0	1.5	1.2	4.4
2012	1.5	0.0	0.0	6.6	1.9	0.2	0.1	1.6	0.1	0.0	23.7	1.4	1.2	4.0
2013	1.7	0.0	0.1	7.3	2.1	0.2	0.1	1.4	0.1	0.0	20.2	1.2	1.0	3.5
2014	1.8	0.0	0.1	7.7	2.2	0.2	0.1	0.8	0.1	0.0	12.1	0.8	0.6	2.0
2015	1.7	0.0	0.1	7.2	2.0	0.2	0.1	0.7	0.1	0.0	9.9	0.6	0.5	0.9
2016	1.6	0.0	0.1	6.9	1.9	0.2	0.1	0.5	0.0	0.0	7.1	0.4	0.4	0.6
2017	1.7	0.0	0.1	7.2	2.0	0.2	0.1	0.5	0.0	0.0	7.5	0.5	0.4	0.6

Source: Statistics Norway/Norwegian Environment Agency.

Differences between the IEA (International Energy Agency) data and the data reported to UNFCCC in sectoral data for marine shipping and aviation are due to the fact that different definitions of domestic use are employed. In the Norwegian inventory, domestic consumption is based on a census in accordance with the IPCC good practice guidance. On the other hand, the IEA makes its own assessment with respect to the split between the domestic and the international market.

3.7.1.2 Shipping

Methodological issues

Emissions are calculated by multiplying activity data with emission factors. The sales statistics for petroleum products, which is based on reports from the oil companies to Statistics Norway, has figures on sales for bunkers of marine gas oil, heavy distillates and heavy fuel oil. The same emission factors as in the Norwegian national calculations are used.

Activity data

Sales figures for international sea transport from Statistics Norway's sales statistics for petroleum products are used for marine gas oil, heavy distillates and heavy fuel oil.

Emission factors

Emission factors used for shipping are described under *Navigation* in Section 3.2.7.

3.7.1.3 Aviation

Methodological issues

The new calculation method for aviation is based on a "bottom up" calculation of jet kerosene consumption and emissions from aviation based on traffic data, emission factors and energy use factors for aircraft types (kg / km). Since traffic data also includes foreign flights, it is possible to link calculations of foreign and domestic aviation directly to activity data. Figures included in international aviation (bunker) is emissions from jet kerosene consumption from flights departing from a Norwegian airport and arriving in a different country.

Activity data

Statistics Norway annually collects data on sales of jet kerosene and aviation gasoline in the sales statistics of petroleum products in addition to use of fuel from the air traffic companies, including specifications on domestic use and purchases of fuel in Norway and abroad. Activity data on flight movements are collected annually from Avinor.

Emission factors

Emission factors used for *Aviation* are described under *Aviation* in Section 3.2.4.

3.7.1.4 Precursors

Emissions of NO_x from international sea traffic in 2016 were about 9.8 ktonnes, a decrease of 12 % from 2015.

NO_x emissions from international aviation amounted to 6.4 ktonnes in 2016, a decrease of 2 % from 2015.

Apart from NO_x from marine bunkers, emissions of precursors from international aviation and sea transport are small compared to the total national emissions of these gases.

3.7.2 CO₂ emissions from biomass

Emissions are estimated from figures in the energy accounts on use of wood, wood waste and black liquor. According to the guidelines, these CO₂ emissions are not included in the national total in the Norwegian emission inventory but are reported as memo items in the CRF.

Emission factors for biomass are shown in Table 3.4. Details are given in the sector chapters where necessary.

4 Industrial processes and product use (CRF sector 2)

4.1 Overview of sector

The chapter provides descriptions of the methodologies used to calculate emissions of greenhouse gases from industrial processes and product use (IPPU). Only non-combustion emissions are included in this chapter. Emissions from fuel combustion in Industry are reported in Chapter 3 (Energy).

Norway has a long experience of using GHG emissions from industrial point sources in the national GHG inventory. The Norwegian Environment Agency has been given the authority to manage and enforce the Pollution Control Act, the Product Control Act and the Greenhouse Gas Emission Trading Act. The Norwegian Environment Agency grants permits, establishes requirements and sets emission limits, and carries out inspections to ensure compliance. This is one of the core responsibilities of the agency.

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. Chapter 2 of Annex VIII describes a major QA/QC exercise in 2006 on consistent time series from 1990 to 2004 from the largest industrial plants in Norway. Several data sources were used to fill data gaps and establish a consistent time series. If no reference is made to data filling in the various source categories, any data filling was done as part of the the QA/QC exercise in 2006. Chapter 3 of Annex VIII describes the current QA/QC procedures and the data sources used.

A large share of the GHG emissions from industrial processes included in the Norwegian GHG inventory are from annual reports sent by each plant to the Norwegian Environment Agency. Such annual reports are reports as required by their regular permit, reports as required by the permit under the EU emission trading system (EU ETS) and reports as required by the voluntary agreement up to the year 2012 when the agreement terminated. The rest of the emissions included in the inventory are calculated by Statistics Norway. The calculations are based on emission factors and activity data. The emission factors are collected from different sources, while the activity data used in calculations carried out by Statistics Norway is from official statistics is normally collected by Statistics Norway.

Indirect emissions of CO₂ from oxidized CH₄ and NMVOC for some source categories are included in the IPPU sector. The indirect emissions of CO₂ are calculated by Statistics Norway and are based on the emissions of CH₄ and NMVOC. See chapter 9 for more details. The IPPU sector contributed to about 119 000 tonnes of indirect CO₂ in 1990 and about 106 000 tonnes of indirect CO₂ in 2017. The majority of these emissions are reported in 2D (Non-energy products from fuels and solvent use).

Table 4.1 gives an overview of the Norwegian IPPU sector. The GHG emissions from IPPU in 2017 were about 8.6 million tonnes CO₂-equivalents, or 16.4 % of the total GHG emissions in Norway. The corresponding percentage in 1990 was 28.3 %. The emissions from this source category have decreased by 40.5 % from 1990 to 2017 and increased by 0.1 % from 2016 to 2017. The decrease from 1990 to 2017 is mainly due to reduced PFC emissions from production of aluminium and SF₆ from production of magnesium. The reduction in the SF₆ emissions is due to the closing down of production of cast magnesium in 2002, improvements in the GIS-sector and an almost end in the use

of SF₆ as tracer gas. In June 2006, the magnesium recycling foundry also closed down. In addition, N₂O emissions from nitric acid production have decreased substantially since 1990.

Table 4.1 Emission trends for IPPU categories (ktonnes CO₂-equivalents).

Category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990- 2017 (%)	Trend 2016- 2017 (%)
2A	727.7	5.0 %	971.6	1 025.3	11.9 %	40.9 %	5.5 %
2B	3 250.6	22.4 %	1 075.1	969.6	11.2 %	-70.2 %	-9.8 %
2C	10 113.3	69.8 %	4 805.0	4 838.7	56.1 %	-52.2 %	0.7 %
2D	287.5	2.0 %	207.6	201.7	2.3 %	-29.8 %	-2.8 %
2E	0.0	0.0 %	1.1	1.1	0.0 %	NA	0.0 %
2F	0.0	0.0 %	1 363.6	1 402.8	16.3 %	3 195 305.3 %	2.9 %
2G	87.5	0.6 %	84.1	79.0	0.9 %	-9.7 %	-6.1 %
2H	31.3	0.2 %	115.6	113.6	1.3 %	263.3 %	-1.7 %
Total	14 497.9		8 623.9	8 631.9		-40.5 %	0.1 %

Source: Statistics Norway and the Norwegian Environment Agency

Metal industry (2C) is the largest category within IPPU and the emissions are mainly from production of ferroalloys and aluminium. The other main contributing sectors are Product uses as substitutes for ODS (2F), Chemical Industry (2B), and Mineral Industry (2A).

Table 4.2 shows the source categories in IPPU that have been identified as key categories from either approach 1 or 2 in the 1 key category analysis.

Table 4.2 Key categories in the sector Industrial processes and product use.

CRF code	Source category	Gas	Key category according to approach	Method
2A1	Cement production	CO ₂	1	Tier 3
2A2	Lime production	CO ₂	1	Tier 3
2B1	Ammonia production	CO ₂	1	Tier 2
2B2	Nitric Acid Production	N ₂ O	2	Tier 3
2B5	Carbide production	CO ₂	2	Tier 2
2B6	Titanium dioxide production	CO ₂	1	Tier 2
2C2	Ferroalloys production	CO ₂	2	Tier 2/3
2C3	Aluminium production	CO ₂	2	Tier 2/3
2C3	Aluminium production	PFCs	2	Tier 2
2C4	Magnesium production	SF ₆	1	Tier 2
2D1	Lubricant use	CO ₂	1	Tier 2
2F	Product uses as substitutes for ODS	HFCs	2	Tier 2

Sources: Statistics Norway and the Norwegian Environment Agency

4.2 Mineral industry – 2A

The sector category Mineral industry includes CO₂ emissions in the source categories cement production, lime production, glass production, ceramics, other uses of soda ash, non metallurgical magnesia production and other process use of carbonates. Table 4.3 shows that components included in the inventory, the tier method used and whether the source categories are key categories or not.

Table 4.3 Mineral industry. Component included in the inventory, tier of method and key category.

Source category	CO ₂	Tier	Key category
2A1. Cement production	R	Tier 3	Yes
2A2. Lime production	R	Tier 3	Yes
2A3. Glass production	R	Tier 3	No
2A4a. Ceramics	R	Tier 3	No
2A4b. Other uses of soda ash	E	Tier 1	No
2A4c. Non metallurgical magnesium production	R	Tier 3	No
2A4d. Other process use of carbonates	R	Tier 2	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occuring. IE = Included Elsewhere.

Table 4.4 shows the trends for 2A as a whole and for the various source categories. The CO₂ emissions from this sector category were a little more than 1 million tonnes in 2017, this accounts for 11.9 % of the total emission from the IPPU-sector. The emissions from this sector have increased with 40.9 % from 1990-2017, mainly due to increased production of clinker and lime in more recent years. The emissions from this sector category increased by 5.5 % from 2016 to 2017.

Table 4.4 Emission trends for 2A Mineral industry (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2A1. Cement production	634.3	4.4 %	684.5	765.6	8.9 %	20.7 %	11.9 %
2A2. Lime production	49.8	0.3 %	218.6	214.9	2.5 %	331.0 %	-1.7 %
2A3. Glass production	5.6	0.0 %	6.6	6.6	0.1 %	18.6 %	-0.3 %
2A4a. Ceramics	3.7	0.0 %	0.0	0.0	0.0 %	NA	NA
2A4b. Other uses of soda ash	9.0	0.1 %	3.0	3.4	0.0 %	-62.7 %	11.6 %
2A4c. Non metallurgical magnesium production	0.0	0.0 %	40.8	14.7	0.2 %	NA	-64.0 %
2A4d. Other process use of carbonates	25.3	0.2 %	18.0	20.1	0.2 %	-20.3 %	11.8 %
Total 2A.	727.7	5.0 %	971.6	1 025.3	11.9 %	40.9 %	5.5 %

Source: Statistics Norway and Norwegian Environment Agency

Table 4.5 and Table 4.6 show that most of the limestone and dolomite uses within the IPPU sector are within 2A. In addition to uses in the IPPU sector, there are reported emissions in 3G from the use of limestone and dolomite.

Table 4.5 Balance in ktonnes for the use of limestone for which IPPU emissions are reported.

Limestone use	2014	2015	2016	2017
2A1 - Cement production	1 653	1 526	1 556	1 740
2A2 - Lime production	498	504	490	481
2A3 – Glass production	0	0	0	0
2A4a – Ceramics	1	0	0	0
2A4c – Non-metallurgical magnesium production	0	0	0	0
2A4d – Other process uses of carbonates	0	0	0	0
2C2 - Production of ferroalloys	71	61	51	32
2H1 – Pulp and paper	19	20	22	19
Total limestone	2 242	2 112	2 119	2 272

Sources: Statistics Norway and the Norwegian Environment Agency

Table 4.6 Balance in ktonnes for the use of dolomite for which IPPU emissions are reported.

Dolomite use	2014	2015	2016	2017
2A2 - Lime production	22	19	27	27
2A4c – Non-metallurgical magnesium production	164	147	90	32
2A4d – Other process uses of carbonates	16	15	11	16
2A3 - Glass production	8	11	11	11
2C2 - Production of ferroalloys	3	4	6	14
Total dolomite	214	196	145	100

Sources: Statistics Norway and the Norwegian Environment Agency

A QA/QC exercise was undertaken for the 2016 NIR to elaborate a mass balance of the limestone and dolomite used in the country. There was no information that indicated that there were other emissive uses of limestone and dolomite that were not reported, and the QA/QC exercise confirmed this. The uses with no emissions are according to the Geological Survey of Norway slurry and crushed rock for filling.

A potential use of limestone is in flue gas desulphurization (FGD), but this is not used in Norway. In Norway, the industry primarily uses the sea water scrubbing technology. This combined with closures of some industrial plants, increasingly strict requirements on the sulphur content in various oil products, the introduction of a SO₂ tax and requirements for industry to reduce its emissions have decreased the SO₂ emissions.

4.2.1 Cement Production, 2A1 (Key category for CO₂)

4.2.1.1 Category description

Two plants in Norway produce cement and they are covered by the EU ETS. Production of cement gives rise to both non-combustion and combustion emissions of CO₂. The emissions from combustion is reported in Chapter 3 Energy. The non-combustion emissions originate from the raw material calcium carbonate (CaCO₃). The resulting calcium oxide is heated to form clinker and then crushed to form cement.



CO₂ from cement production is defined as a key category according to the approach 1 analysis.

4.2.1.2 Methodological issues

The emissions of CO₂ from clinker production included in the GHG inventory are reported by the two producers in their annual report under their regular permit and under the EU ETS to the Norwegian Environment Agency. Before entering the EU ETS, the plants used a tier 2 methodology while they now use a tier 3 methodology. The plants report data on the types and quantities of carbonates consumed to produce clinker, as well as their emission factors and oxidation factors. The reported emissions include Cement Kiln Dust (CKD).

4.2.1.3 Activity data

The amount of clinker, CKD and other carbonates that the plants use in their calculation are reported by the plants to the Norwegian Environment Agency. The annual total clinker production is reported in the CRF Table 2(I).A-Hs1 and Table 4.7 shows the clinker production for some selected years in the time series in total and for each of the two plants.

Table 4.7 Norwegian clinker production (ktonnes).

Year	Total production	Production in plant 1	Production in plant 2
1990	1 244.1	1 055.8	188.3
1995	1 682.9	1 181.0	501.9
2000	1 649.6	1 201.2	448.4
2005	1 454.3	979.3	475.0
2008	1 534.0	1 049.2	484.8
2009	1 528.3	1 038.8	489.5
2010	1 433.8	1 008.5	425.3
2011	1 415.4	1 000.3	415.1
2012	1 399.1	1 002.7	396.3
2013	1 399.8	1 047.3	352.5
2014	1 374.9	1 006.0	368.8
2015	1 284.1	953.7	330.4
2016	1 306.3	953.6	352.7
2017	1 461.5	1 009.2	452.3

Source: Norwegian Environment Agency

4.2.1.4 Emission factors

CO₂

The emission factors used are plant specific. The factors are dependent on the chemical composition of the clinker i.e. the content of Ca and Mg. The fraction of CaO from non-carbonate sources like ashes is subtracted. The emission factors are calculated particularly for the two Norwegian plants. Prior to entering the EU ETS, the emission factors did not vary much and tended to be around 0.530 tonne CO₂ per tonne clinker for one plant (Tokheim 2006) and 0.541 tonne CO₂ per tonne clinker as recommended by SINTEF (1998e) for the other plant. The IPCC default emission factor is 0.52 tonne CO₂/tonne clinker. After entering the EU ETS, the plants face stricter requirements concerning how their EF are determined and the EFs may vary more from one year to another. The same emission factors are used for CKD as for clinker production.

Table 4.8. Emission factors (tonne CO₂/tonne clinker) and conversion factors for clinker production.

Year	Emission factors	Conversion factors
2015	0.519691 - 0.5235334	0.96214 - 0.970692
2016	0.520148 - 0.5221745	0.9655105 - 0.975824
2017	0.50826982 - 0.523415	0.9626982 - 0.970254

Source: Norwegian Environment Agency

Until 2009, both plants have used a conversion factor of 1. This means that all Ca and Mg have been assumed to be carbonates. From 2010, the largest plant has reported conversion factors that are less than 1 (0.948 or higher). The smaller plant has continued to use a conversion factor of 1 until 2015. Table 4.8 shows the emission factors and conversion for the two plants for some years.

4.2.1.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

The two plants have reported their emissions to the Norwegian Environment Agency for many years. Cement production was included in the EU ETS in 2005. After entering the EU ETS, the plants face stricter requirements concerning how AD and EF are determined and the EFs will vary more from one year to another. The reduction in IEF from 2009 to 2010 is a consequence of lower EFs in 2010 for both plants. The EF for the plant producing about 70% of the total production decreased the most, pushing the IEF for total production down. This explains the inter-annual variations in the IEF in the end of the time series.

4.2.1.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the Norwegian Environment Agency's inventory team.

Statistics Norway occasionally calculates alternative emission figures for CO₂ and compares them with the emission figures reported by the plants to the Norwegian Environment Agency to check if they are reasonable. The calculations are based on the clinker production (reported annually from the plants to the Statistic Norway. The calculated emission figures have agreed quite well with emissions figures reported by the plants.

4.2.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.2.2 Lime Production, 2A2 (Key category for CO₂)

4.2.2.1 Category description

Three plants that produce lime in Norway reported CO₂ emissions from processes to the Norwegian Environment Agency and all three plants are covered by the EU ETS. The large increase in CO₂

emissions from lime production from 1990 is due to increased production at existing plants and the establishment of a new plant in 2007 with large production. CO₂ from lime production is defined as a key category according to the approach 1 analysis.

4.2.2.2 Methodological issues

All three plants calculate the emissions of CO₂ based on the input of limestone and dolomite and plant specific emission factors for CO₂ from limestone and dolomite respectively. The emissions are reported to the Norwegian Environment Agency. This is in accordance with the reporting requirements of the EU ETS and is in line with the tier 3 method of the IPCC 2006 GL. The activity data is corrected for lime kiln dust (LKD).

4.2.2.3 Activity data

The activity data used for the reported emissions is the input of limestone and dolomite and this is reported annually to the Norwegian Environment Agency. Nearly all production in Norway consists of quicklime, but there is also some dolomitic lime. Table 4.9 shows the lime production and consumption in 2A for some of the years in the time series.

Table 4.9 Lime consumption (ktonnes) in 2A2.

Year	Total consumption	Consumption plant 1	Consumption plant 2	Consumption plant 3
1990	116.3	65.0	51.3	0.0
1995	162.7	88.2	74.5	0.0
2000	158.9	80.2	78.7	0.0
2005	197.6	100.3	97.3	0.0
2008	338.1	102.7	82.4	152.9
2009	324.3	85.3	92.7	146.3
2010	576.5	107.5	105.6	363.3
2011	524.1	106.0	115.1	303.0
2012	531.6	94.0	113.1	324.6
2013	518.1	94.3	111.3	312.6
2014	521.5	99.6	90.4	331.4
2015	524.8	79.6	106.5	338.7
2016	519.1	93.7	114.5	310.9
2017	510.8	102.5	115.3	293.0

Source: Norwegian Environment Agency

The amounts of dolomite used in lime production are shown in Table 4.6. Even though the emissions are calculated based on limestone and dolomite consumption, Norway reports final lime production values as AD in CRF Table 2(I).A-Hs1 in order to assist with comparability across Parties.

4.2.2.4 Emission factors

The plants use an emission factor of 0.474 tonnes CO₂ per tonne dolomite used. The range of emission factors for limestone are plant specific and are shown in Table 4.10 for some years. The plants used a conversion factor of 1 up to and including 2007 (one plant) or 2008 (two plants). This means that all Ca and Mg have been assumed to be carbonates. Since then, the plants have reported a range of conversion factors that are less than 1 and these are shown in Table 4.10 for some years.

Table 4.10 Range for emission factors (tonne CO₂/tonne limestone) and conversion factors for lime production.

Year	Emission factors	Conversion factor
2015	0.4373 - 0.43760732	0.9550 - 0.972
2016	0.43337606 - 0.4378	0.9532 - 0.97257987
2017	0.4328894 - 0.4341	0.9639 - 0.97032381

Source: Norwegian Environment Agency

4.2.2.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

Figure 4.1 shows the IEFs for lime production for both consumption and production as AD. Using final lime production values as AD results in IEFs closer to the default IPCC EF, but also to less stable IEFs as it varies more than if consumption is used as AD.




Figure 4.1 IEF (tonne CO₂ per tonne limestone) with consumption or production as AD.

Source: Norwegian Environment Agency

4.2.2.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by Norwegian Environment Agency's inventory team.

4.2.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.2.3 Glass production, 2A3

4.2.3.1 Category description

Three plants producing glass or glass fibre are included in the emission inventory, based on emission reports to the Norwegian Environment Agency. All three plants are covered by the EU ETS.

4.2.3.2 Methodological issues

Two plants producing glass wool and one plant producing glass fibre report emission figures on CO₂ to the Norwegian Environment Agency. The two glass wool producing plants report emissions from the use of soda ash, limestone and dolomite, while the glass fibre producer reports emissions from the use of limestone and dolomite.

4.2.3.3 Activity data

The aggregated use of soda ash, limestone and dolomite is reported as activity data in CRF Table 2(I).A-Hs while details for the use of soda ash and dolomite are shown in Table 4.11 and Table 4.6.

4.2.3.4 Emission factors

The emission factors used are 0.41492 tonnes CO₂/tonne soda ash (2006 IPCC GL), 0.477 tonnes CO₂/tonne dolomite (EU ETS) and 0.44 tonnes CO₂/tonne limestone (EU ETS).

4.2.3.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.2.3.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by Norwegian Environment Agency's inventory team.

4.2.3.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.2.4 Ceramics, 2A4a

4.2.4.1 Category description

One plant that produced bricks until 2014 is included in the emission inventory, based on emission reported to the Norwegian Environment Agency. The plant was covered by the EU ETS.

4.2.4.2 Methodological issues

The plant reported emission figures of CO₂ to the Norwegian Environment Agency. The emissions are calculated by multiplying the amount of limestone and clay used in its production with emission factors.

4.2.4.3 Activity data

The amount of limestone and clay used in the production of bricks was reported each year from the plant to the Norwegian Environment Agency. The amounts of limestone used are reported in CRF Table 2(l).A-Hs1.

4.2.4.4 Emission factors

The EF of 0.44 tonnes CO₂ per tonne limestone used by the brick producing plant is the standard EF used in the EU ETS for limestone. The plant used an emission factor of 0.088 tonnes CO₂ per tonne clay used.

4.2.4.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

The emissions reported under 2A4a include emissions from the use of clay, but the AD in the CRF is limestone only. The use of clay has decreased since 1996 and this explains the overall decrease in IEF for 2A4a. Calculations show that the IEF for CO₂ from limestone and dolomite use only is more stable than when the emissions from the of clay also are included.

4.2.4.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions were covered by the EU ETS and their emissions were verified annually. In addition, the emissions were checked both by the case handler and by the Norwegian Environment Agency's inventory team.

4.2.4.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.4.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.2.5 Other uses of soda ash, 2A4b

4.2.5.1 Category description

There are no data on soda ash in Norway in production statistics (PRODCOM) from Statistics Norway, so all soda ash is imported. Soda ash is used and reported in 2A3 (glassworks), 2C3 (aluminium production) and 2C7a(ii) (nickel production). The import of soda ash is higher than the sum of the amounts consumed in these industries. This use is assumed to be emissive and the corresponding CO₂-emissions are estimated and reported here under 2A4b.

4.2.5.2 Methodological issues

The emission figures for CO₂ are estimated by multiplying the amount of soda ash assumed to be emissive with an emission factor.

4.2.5.3 Activity data

The activity data is total import of soda ash minus consumption in glass wool, nickel and aluminium production. The amounts of soda ash are reported in CRF Table 2(I).A-Hs1 and are shown in Table 4.11.

Table 4.11 Balance for soda ash use for Norway (ktonnes).

Year	Import	2A4b (other uses of soda ash)	2A3 (Glassworks)	2C3 (Aluminium production)	2C7ii (Nickel production)
1990	45.1	21.7	4.2	0.9	18.3
1995	55.0	24.5	4.2	0.9	25.3
2000	49.1	17.0	5.3	0.9	25.8
2005	63.8	21.3	5.4	0.9	36.1
2007	53.9	16.7	3.5	0.9	32.7
2008	59.6	22.9	3.5	0.9	32.3
2009	41.4	1.8	3.5	0.9	35.1
2010	34.9	-	3.5	0.9	33.6
2011	48.7	10.7	3.6	0.9	33.4
2012	42.1	-	3.7	0.9	38.1
2013	51.8	11.1	4.0	0.9	35.8
2014	47.5	5.1	3.4	1.1	37.8
2015	44.1	2.6	4.0	1.2	36.2
2016	47.8	7.2	3.7	0.9	36.0
2017	47.5	8.1	3.7	0.7	34.9

Source: Statistics Norway and Norwegian Environment Agency

4.2.5.4 Emission factors

The emission factor for soda ash use is 0.41492 tonnes CO₂/tonne soda ash from the IPCC 2006 Guidelines (IPCC 2006).

4.2.5.5 Uncertainties and time-series consistency

As we do not have sufficient information to determine where the rest of the imported soda ash has been consumed, there is some uncertainty as to whether all soda ash consumption in fact is emissive. There is also some uncertainty associated with the foreign trade statistics, as well as with the assumption that the CO₂ is emitted the same year as the soda ash are imported. According to the IPCC Guidelines 2006, there is negligible uncertainty associated with the emission factor, given that the correct emission factor is applied (IPCC 2006).

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.2.5.6 Category-specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. However, when the calculation first was included in the inventory, a comparison was made between figures on net import of soda ash in

foreign trade statistics and in the Norwegian Product Register. Import figures from the Product Register for the period 2000-2011 never constituted more than 41 % of the amounts imported according to foreign trade statistics. Thus, it was assumed that the net import in the foreign trade statistics is a good proxy for the total quantity of soda ash used in Norway.

4.2.5.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.5.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category. In the future, we might examine what these other uses of soda ash actually are in order to confirm whether they are emissive or not.

4.2.6 Non-metallurgical magnesium production, 2A4c

4.2.6.1 Category description

One plant whose main activity is producing magnesium oxide from limestone and dolomite is included in the emission inventory. The plant was established in 2005 and is covered by the EU ETS.

4.2.6.2 Methodological issues

The plant reports emission figures of CO₂ to the Norwegian Environment Agency. The emissions are calculated by multiplying the amount of limestone and dolomite used in its production with emission factors.

4.2.6.3 Activity data

The amount of limestone and dolomite used in the production is reported each year from the plant to the Norwegian Environment Agency. The aggregate amounts of limestone and dolomite used are reported in CRF Table 2(l).A-Hs1 and Table 4.12 shows the usage for some selected years in the time series.

Table 4.12 Usage (kt) of limestone and dolomite in the non-metallurgical magnesium production.

Year	limestone use	dolomite use
2005	0.0	1.4
2008	1.5	23.9
2009	0.8	9.7
2010	1.4	0.0
2011	0.0	0.0
2012	0.0	14.2
2013	0.0	124.5
2014	0.0	163.7
2015	0.0	147.3
2016	0.0	90.2
2017	0.0	32.46

Source: Norwegian Environment Agency

4.2.6.4 Emission factors

The plant has used the emission factor (EF) equal to the standard EF used in the EU ETS for limestone before it entered the EU ETS and uses plant specific EFs after it has entered the EU ETS. The plant does not use limestone every year, and the EFs for 2006, 2008, 2009 and 2010 are in the range of 0.41-0.4504. The EF for the dolomite used is equal to the standard EF used in the EU ETS (0.45) for 2005, 2006 and 2007 before it entered the EU ETS. From 2008 the plant has used plant specific EFs in the range of 0.46-0.495096639.

With the exception of 2012 and 2015-2017, the plant has used a conversion factor of 1. This means that for most years, all Ca and Mg have been assumed to be carbonates.

4.2.6.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.2.6.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the Norwegian Environment Agency's inventory team.

4.2.6.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.6.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.2.7 Other process use of carbonates, 2A4d

4.2.7.1 Category description

The emissions from five plants are reported here under 2A4d. The CO₂ emissions from two plants producing leca are included in the emission inventory, based on emission reports to the Norwegian Environment Agency. One of the plants stopped its production in 2004 and the existing plant is covered by the EU ETS. The third plant has neutralized sulphuric acid waste primarily with limestone and fly ash, but uses now only fly ash. The use of fly ash reduces the CO₂ emissions compared with when limestone is used. The CO₂ emissions from two plants producing rock wool are also included in the emission inventory, based on emission reports to the Norwegian Environment Agency.

4.2.7.2 Methodological issues

The two plants producing leca report their use of dolomite and the corresponding CO₂ emissions to the Norwegian Environment Agency. The leca producer that still is in production also reports some minor emissions from the use of clay and these are included in the inventory. For the plant neutralizing sulphuric acid waste, the emissions are calculated by multiplying the amounts of limestone and ash used to neutralize sulphuric acid waste with emission factors. The two plants

producing rock wool report their use of dolomite and limestone and the corresponding CO₂ emissions to the Norwegian Environment Agency.

4.2.7.3 Activity data

The activity data are primarily the use of dolomite and limestone. The aggregate amounts of limestone and dolomite used by the plants included in 2A4d are reported in the CRF Table 2(I).A-Hs1 and are shown in Table 4.13 for some selected years in the time series.

Table 4.13 Use (kt) of limestone and dolomite in 2A4d (other process use of carbonates).

Year	limestone use	dolomite use
1990	49.572	5.338
1995	42.963	7.759
2000	73.126	7.376
2005	59.012	13.098
2008	60.050	9.567
2009	3.818	13.235
2010	0.000	8.677
2011	33.205	14.413
2012	5.915	18.906
2013	33.932	17.719
2014	0.000	16.336
2015	0.000	15.263
2016	0.000	11.452
2017	0.000	15.556

Source: Norwegian Environment Agency

In addition, clay is used by the leca producer that still is in production and ash is used by the plant neutralizing sulphuric acid waste.

4.2.7.4 Emission factors

An emission factor (EF) of 0.48 t CO₂/t dolomite was used by the leca producer that closed down in 2004. At that time, there was no standard EF for dolomite. We assume that the EF of 0.48 is derived from the standard factors for CaCO₃ and MgCO₃ and an assumption of the ratio of these in the dolomite. For the leca producer that still is in production, the EF (for dolomite) for 1990-2011 is 0.477, and has since then used standard EF from the EU ETS and plant-specific EFs in the range of 0.466-0.48. The EFs for the use of clay from 2013 and onwards ranges between 0.01596492-0.01974294. The plant that neutralizes sulphuric acid waste uses an emission factor of 0.44 t CO₂/t limestone. For fly ash, IEFs for the years 2010-2016 have been calculated by the Norwegian Environment Agency to be in the range of 40 to 140.2 kg CO₂/t fly ash based on activity data and emissions. The IEF of 68.5 kg CO₂/t fly ash for 2010 has been used for the years 1997-2009. The two plants producing rock wool use emission factors of 0.44 t CO₂/t limestone and 0.481 t CO₂/t dolomite.

4.2.7.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category. The IEF is relatively stable 1990-2008, variations may be due to emissions from the use of clay and ash that are not included as AD in the CRF. The IEF since 2008 varies

more and this is primarily due to a shift from using limestone to ash at the plant that neutralizes sulphuric acid waste.

4.2.7.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The existing plant producing leca is covered by the EU ETS and the emissions are verified annually. The emissions are checked both by the case handler and by the Norwegian Environment Agency's inventory team. The reported emissions from the plant that neutralizes sulphuric acid waste occurs under its regular permit and are checked both by the case handler and by the agency's inventory team.

4.2.7.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.2.7.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3 Chemical industry – 2B

Several activities are included under Chemical Industry. Nearly all emissions figures from this industry included in the inventory are reported figures from the plants to the Norwegian Environment Agency. Table 4.14 shows the GHGs that are emitted from each source category, tier of methodology and if the source category is key category or not.

Table 4.14 Chemical industry. Components included in the inventory, tier of method and key category

Source category	CO ₂	CH ₄	N ₂ O	NM VOC	Tier	Key category
2B1. Ammonia production	R	NA	NA	NA	Tier 2	Yes
2B2. Nitric acid production	NA	NA	R	NA	Tier 3	Yes
2B5a. Silicon carbide production	R+E	R/E	NA	NA	Tier 2	Yes
2B5b. Calcium carbide production	R	NA	NA	R	Tier 1	No
2B6. Titanium dioxide production	R	NA	NA	NA	Tier 2	Yes
2B8a. Methanol production *	R	R+E	R	R+E	Tier 2	No**
2B8b. Ethylene production *	R+E	R	R	R	Tier 2	No**
2B8c. Ethylene dichloride and vinyl	R+E	R	NA	R	Tier 2	No**
2B10. Other (production of	NA	NA	R+E	NA	Tier 2	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

*Minor N₂O emissions from 2B8a and 2B8b are reported under 2B10 Petrochemical N₂O in the CRF, but are included in Table 4.15 below. ** In the key category analysis, 2B8a, 2B8b and 2B8c have been aggregated to 2B8, which has not been identified as a key category.

Table 4.15 shows the trends for 2B as a whole and for the various source categories. The GHG emissions from this sector category were a little less than 1 million tonnes in 2017, this is 11.2 % of the total emission from the IPPU-sector. The emissions from this sector decreased by 70.2 % from 1990, mainly due to lower emissions from the production of nitric acid, ammonia and carbide. The emissions decreased by 9.8 % from 2016 to 2017.

Table 4.15 Emission trends for 2B Chemical industry (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2B1. Ammonia production	500.1	3.4 %	384.6	260.5	3.0 %	-47.9 %	-32.3 %
2B2. Nitric acid production	1	13.7 %	241.3	210.5	2.4 %	-89.4 %	-12.8 %
2B5a. Silicon carbide production	230.2	1.6 %	51.6	58.8	0.7 %	-74.5 %	14.0 %
2B5b. Calcium carbide	178.1	1.2 %	0.0	0.0	0.0 %	NA %	NA
2B6. Titanium dioxide production	201.1	1.4 %	147.6	261.1	3.0 %	29.8 %	76.9 %
2B8a. Methanol production	0.0	0.0 %	127.1	66.7	0.8 %	NA	-47.5 %
2B8b. Ethylene production	71.1	0.5 %	41.8	28.6	0.3 %	-59.8 %	-31.6 %
2B8c. Ethylene dichloride and vinyl chloride production	18.7	0.1 %	13.5	15.3	0.2 %	-18.1 %	13.5 %
2B10. Other (production of fertilizers)	58.0	0.4 %	67.7	68.1	0.8 %	17.5 %	0.7 %
2B. Total	3	22.4 %	1 075.1	969.6	11.2 %	-70.2 %	-9.8 %

Source: Statistics Norway and Norwegian Environment Agency

4.3.1 Ammonia Production, 2B1 (Key category for CO₂)

4.3.1.1 Category description

In Norway ammonia is produced by catalytic steam reforming of wet fuel gas (containing ethane, propane and some buthane). This is one of the steps in the production of fertilizers. Hydrogen is needed to produce ammonia, and wet fuel gas is the basis for the production of hydrogen. A substantial amount of CO₂ is recovered from the production process.

CO₂ from ammonia production is defined as a key category according to the approach 1 analysis.

Figure 4.2 shows the time series for the gross CO₂ emissions, amount of recovered CO₂ and the net CO₂ emissions. The variations from 1998 to 1999 and 1999 to 2000 are likely to be a result of the plant upgrading production capacity and energy efficiency in 1999-2000. The increase in emisisions from 2014 to 2015 is due to an expansion in production capacity in which imported ammonia is replaced with domestic ammonia production. The emisisions decreased however in 2016 and 2017 since domestic ammonia production decreased.




Figure 4.2 CO₂ emissions from production of ammonia.

Source: Norwegian Environment Agency

4.3.1.2 Methodological issues

The CO₂ emission figures in the Norwegian emission inventory model are based on annual reports from the plant. The plant calculates the emissions by multiplying the amount of each gas used with gas specific emission factor.

The plant has reported consistent figures back to 1990. A part of the CO₂, which is generated during the production process, is captured and sold to other objectives et cetera soft drinks, and therefore deducted from the emission figures for this source. The amount of CO₂ recovered by the plant is determined by using the amount of CO₂ from the compressor unit minus the amount of CO₂ emitted. In both cases, these are measured by orifice plate with a dp-cell. This then goes to tank and then

transported by trucks or boats. In accordance with the footnote 5 in CRF table 2(l)-A-H, the amount recovered that is not exported is included in 2H2 Food and Drink.

4.3.1.3 Activity data

The total amount of gas consumed is annually reported by the plant to the Norwegian Environment Agency. The use of the different gases varies from one year to another. As a part of the official Industrial statistics, gas consumed is also reported to Statistics Norway that uses these figures for the QA/QC calculations by an alternative method.

4.3.1.4 Emission factors

The plant emission factors used in the calculations of emissions are calculated based on the composition of the gases consumed. The plant states that the composition is based on daily analysis and that the composition of each gas (their emission factor) is stable.

4.3.1.5 Uncertainties and time-series consistency

The amount of gas is measured by using turbine meters and the meters are controlled by the Norwegian Metrology Service. The uncertainty in the measurement of propane and butanes is calculated to ± 0.2 and ethane ± 0.13 %. The mix of propane/butanes is as average 60 % propane and 40 % butanes.

There are some large inter-annual variations in the IEF. The variations from 1998 to 1999 and 1999 to 2000 are likely to be a result of the plant upgrading production capacity and energy efficiency in 1999-2000. Figure 4.2 shows that there was a large drop in emissions and recovery in 1999. We do not have explanations for the variations from 1996 to 1997 and 1997 to 1998. The IEF of 1.8 in 1997 indicates that the emissions may have been overestimated or the production may have been underestimated. It is challenging to investigate this further as more data is not available and since the data quality at that time is poorer than now. Since the plant has reported under the voluntary agreement for 2008-2012 and under the EU ETS from 2013, the data quality has improved as

Figure 4.3 shows a relatively stable IEF for the end of the time series.




Figure 4.3 IEF for process emissions of CO₂ from ammonia production (tonne CO₂/tonne ammonia).

Source: Norwegian Environment Agency

4.3.1.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant has reported under the voluntary agreement and the emissions are now covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

The figures reported from the plant are occasionally compared to calculations done by Statistics Norway based on total amount of gas consumed and an emission factor on 3 tonne CO₂/tonne LPG. The calculated emissions figures have agreed quite well with emissions figures reported by the enterprise.

4.3.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category. We have investigated the issue of the IEFs to the extent possible and the IEF of 1.8 in 1997 indicates that the emissions may have been overestimated or the production may have been underestimated.

4.3.2 Production of Nitric Acid, 2B2 (Key category for N₂O)

4.3.2.1 Category description

There are two plants in Norway producing nitric acid and these plants are covered by the EU ETS. Nitric acid is used as a raw material in the manufacture of nitrogenous-based fertilizer. The production of nitric acid (HNO₃) generates nitrous oxide (N₂O) and NO_x as by-products of high temperature catalytic oxidation of ammonia (NH₃). N₂O from nitric acid production is defined as a key category according to the approach 2 analysis.

Table 4.16 compares the Norwegian plant-specific production technologies compared with the technologies described in table 3.3 in the IPCC 2006 Guidelines (IPCC 2006).

Table 4.16 Production process and default factors for nitric acid production.

Production process	N ₂ O Emission Factor (relating to 100 percent pure acid)
A. Plants with NSCR ¹² (all processes)	2 kg N ₂ O/tonne nitric acid ±10%
B. Plants with process-integrated or tailgas N ₂ O destruction	2.5 kg N ₂ O/tonne nitric acid ±10%
C. Atmospheric pressure plants (low pressure)	5 kg N ₂ O/tonne nitric acid ±10%
D. Medium pressure combustion plants	7 kg N ₂ O/tonne nitric acid ±20%
E. High pressure plants	9 kg N ₂ O/tonne nitric acid ±40

Source: IPCC (2006)

The two plants have together five production lines. Four of the production lines are a mix of technology C and D in Table 4.16 and the last one is technology B. One production line was rebuilt in 1991 and in 2006 two lines were equipped with the technology – N₂O decomposition by extension of

¹² A Non-Selective Catalytic Reduction (NSCR)

the reactor chamber. Since then, all production lines have to a certain extent been equipped with this technology.

Figure 4.4 shows that the production specific N₂O emissions were reduced substantially in the early 1990s and again from 2006. The reduced emissions in the early 1990s were due to rebuilding of one production line in 1991 and that a larger part of the production came from that line. The reduced emissions from 2006 are due to the installation of the earlier mentioned technology. There was a large increase in production of about 43 percent from 2009 to 2010 that came after a decrease in production of about 26 percent from 2008 to 2009. The low production level in 2009 reflects the lower economic activity due to the economic recession.

4.3.2.2 Methodological issues

N₂O

The two plants report the emissions of N₂O to the agency. The N₂O emissions have been continuously measured since 1991 at one production line and from 2000 at another. The emissions at the three other production lines were based on monthly and weekly measurements, but are from 2008 based on continuous measurements.

4.3.2.3 Activity data

The plants report the amounts of N₂O in the gas, based on continuous measurements. The plants also report the production of HNO₃ to the agency. The production and emission levels (in total and by plant) are shown for some years in *Table 4.17*.

Table 4.17. Production (ktonnes HNO₃) and N₂O emissions (ktonnes)

Year	Total production	Production plant 1	Production plant 2	Total emissions	Emissions plant 1	Emissions plant 2
2015	1729.7	1373.5	356.2	0.844	0.656	0.189
2016	1669.2	1310.8	358.4	0.810	0.631	0.179
2017	1733.7	1334.7	398.9	0.706	0.524	0.183

Source: Statistics Norway and Norwegian Environment Agency

4.3.2.4 Emission factors

Not relevant.

4.3.2.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II. The uncertainty in the measurements was in 2000 estimated by the plant to ± 7 . However, in the 2006 report to the Norwegian Environment Agency one plant reports that the uncertainty in measurement of N₂O is calculated to $\pm 1-3$ %.

The inter-annual changes of IEFs are likely to be explained by variations in the level of production between the lines with different IEFs. Figure 4.4 shows that the IEF for nitric acid production has been substantially decreased from 1990 to 2017. The low production level in 2009 reflects the lower economic activity due to the economic recession.




Figure 4.4 Relative change in total emissions, total production and IEF for nitric acid production. 1990=100
Source: Norwegian Environment Agency

4.3.2.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.



4.3.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.3 Silicon carbide, 2B5a (Key category for CO₂)

4.3.3.1 Category description

Silicon carbide (SiC) has been produced at three plants until 2006 when one plant was closed down. The plants were included into the EU ETS from 2013. SiC is produced by reduction of quartz (SiO₂) with petrol coke as a reducing agent.



In the production of silicon carbide, CO₂ and CO is released as a by-product from the reaction between quartz and carbon. Methane (CH₄) may be emitted from petrol coke during parts of the process and sulphur origin from the petrol coke.

The large decrease in emissions since 1990 is due to reduced production and that one plant was closed down in 2006. The fluctuation in emissions over the years is due to variation in production of crude silicon carbide. There was a large production decrease from 2002 to 2003 and an increase from 2009 to 2010 and this is due to a low production level in 2009. The production level in 2009 is also lower than 2008 and reflects the lower economic activity due to the economic recession.

CO₂ from carbide production is defined as a key category according to the approach 2 analysis.

4.3.3.2 Methodological issues

The emissions are calculated based on crude silicon carbide production as activity data and emission factors. This is regarded as being a Tier 2 method in IPCC (2006).

CO₂

Emission figures are reported annually by the three plants to the agency.

CO₂ from process is calculated based on the following equation:

$$(4.3) \quad CO_2 = \sum Activity\ data * Emission\ factor$$

The three production sites have used amount of produced crude silicon carbide as activity data in the calculation of CO₂ emissions.

NMVOC

Emission figures are reported to the Norwegian Environment Agency by the plants. The emissions are calculated by multiplying annual production of silicon carbide by an emission factor.

Indirect emission of CO₂ is calculated by Statistics Norway based on the emission of CH₄.

CH₄

The emission of CH₄ from production of silicon carbide is calculated based on the following equation:

$$(4.4) \quad CH_4 = \sum Activity\ data_i * Emission\ factor_i$$

The three production sites have used amount of produced crude silicon carbide as activity data and a plant specific emission factor.

4.3.3.3 Activity data

The activity data used by the plants for the calculation of CO₂, CH₄ and NMVOC are the amount of produced crude silicon carbide. The production is shown in Table 4.18 for some years. For the calculations of indirect CO₂, the AD is the amount of CH₄.

Table 4.18. Norwegian crude silicon carbide production (ktonnes).

Year	Production
1990	83.4
1995	83.6
2000	66.7
2005	35.0
2008	28.9
2009	18.9
2010	27.2
2011	25.3
2012	15.8
2013	16.9
2014	17.9
2015	17.6
2016	18.7
2017	21.2

Source: Statistics Norway and Norwegian Environment Agency

4.3.3.4 Emission factors

CO₂

All three sites use the country-specific emission factor that is the basis for the IPCC (2006) default factor of 2.62 ton CO₂/tonne crude silicon carbides, see Table 4.19.

CH₄

For calculation of methane emissions the country-specific emission factor 4.2 kg CH₄/tonne crude SiC is used, see Table 4.19. Documentation of the choice and uncertainties of the emission factor is given under Uncertainties.

Table 4.19 Emission factor for CO₂ and CH₄ used for silicon carbide production.

Component	Emission factor	Source
CO ₂	2.62 tonnes CO ₂ /tonnes crude SiC	IPCC 2006
CH ₄	4.2 kg CH ₄ /tonnes crude SiC	CS

NMVOC

From 2007 and onwards the emission factor is based on measurements made once a year. The emission factors for one of the plants is stable at around 10.8 t NMVOC/kt SiC while the emission factor at the other plant is less stable and increasing. The concerned plant has responded that the variations are within the expected variations. For previous years, the emission factor for one of the plants is more or less constant whereas the emission factor for the second plant varies.

4.3.3.5 Uncertainties and time-series consistency

CO₂

Activity data

The three production sites use the amount of produced crude silicon carbide as activity data. The uncertainty of the activity data is related to the uncertainty of the weighing equipment and is calculated to be $\pm 3\%$.

Emission factor

The emission factor of 2.62 tonne CO₂/tonnes SiC has an estimated uncertainty range of -16% to $+7\%$. This can be explained due to variations in raw materials as well as process variations, and is based on previous development of country specific emissions factors (SINTEF 1998d).

The carbon content in coke is varying, normally from 85 to 92 % carbon. The coke is also varying in the content of volatile components, e.g hydrocarbons. There are also variations in the process itself. The Acheson process is a batch process, and the reactions include many part reactions that differ from batch to batch, because of variations in the mix of quartz and coke, the reactivity of the coke etc. The process variations described above is the reason why the factor presented in tonne CO₂/tonne coke used is not constant. For one plant, the factor is in the range 1.07-1.27. For the other plant, one also has to consider the closed plant, because the input and output from them are somewhat mixed together. The factor for them is in the range 0.99-1.24. This implies that the output of SiC will have some variation from batch to batch.

Prior to 2006, the emissions were based on a mass-balance method (input of reducing agents). The justification of changing method is that the IEF tonne CO₂ /tonne coke varies over the years due to variation in carbon content in coke and that this variation is larger or in the same order of variation than the production of crude silicon carbide. In addition, there is a relatively large difference in the carbon consumption data in the early 1990s due to the use of purchase data as a proxy for carbon consumption. The silicon carbide production data in the early 1990s especially is considered being more accurate than the coke consumption.

Emissions

The total uncertainty of the resulting emissions of CO₂, based on uncertainties in activity data and emission factor, is calculated to be in the range of -20% to $+10\%$.

CH₄

Activity data

The three production sites use the amount of produced crude silicon carbide as activity data. The uncertainty of the activity data given as this production figure is calculated to be $\pm 3\%$.

Emission factor

The emission factor of 4.2 kg CH₄/tonne SiC is used, and the uncertainty level is estimated to be $\pm 30\%$.

The calculation of emission factor and the uncertainty level is explained below. The production of SiC is a batch process with duration of about 43 hours. The CH₄-concentration (ppm) is monitored continuously the first 6.5 hours. After this, only control monitoring is carried out. The results show that the concentration of CH₄ is peaking in the first hour of the process, giving a CH₄ concentration 10

– 15 times higher than in the last 36 hours of the process. A typical level of the concentration of CH₄ is given in Figure 4.5. If the CH₄-concentration is averaged over the total batch time of 43 hours, this will give an emissions factor of 4.2 kg CH₄/tonne SiC, i.e. 3.5 kg CH₄/tonne petrol coke.




Figure 4.5. Concentration of CH₄ for one batch of SiC.

To establish the uncertainty level, the following assessments were done:

- The uncertainty in monitoring of concentration is normally $\pm 5\%$ (expert judgment).
- The uncertainty of monitoring of the amount of gas is within $\pm 15\%$ (type of monitoring equipment).
- The uncertainty of the production of SiC for each batch is stable, and is assessed to be within a level of $\pm 5\%$.
- The uncertainties of raw materials and process variation add $\pm 5\%$.

If these uncertainties are added, the estimate result of total uncertainties for the resulting emissions of CH₄ is $\pm 30\%$.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.3.3.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants have reported under the voluntary agreement and the emissions are now covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.3.7 Category-specific recalculations

Correction of error. The activity data, CO₂ emissions and CH₄ emissions for the years 2010-2016 have been recalculated for one plant. The concerned plant has for these years mixed the reporting of GHG emissions to the Norwegian Environment Agency with internal reporting. For some years total crude production was used instead of total pure production and for some other years an emission factor

other than 2.62 was used. This has now been corrected. The recalculations vary from a reduction in emissions of almost 2 300 tonnes CO₂ in 2013 to an increase in emissions of about 2 700 tonnes CO₂ in 2015. The changes in CH₄ emissions are minor.

4.3.3.8 Category-specific planned improvements


There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.4 Calcium carbide, 2B5b

4.3.4.1 Category description


One plant in Norway was producing calcium carbide until 2003 and the emissions from this source were about 178 000 tonnes CO₂ in 1990. The production of calcium carbide generates CO₂ emissions when limestone is heated and when petrol coke is used as a reducing agent.

The reaction



which takes place when limestone (calcium carbonate) is heated.

The reactions



where petrol coke is used as a reducing agent to reduce the CaO to calcium carbide.

4.3.4.2 Methodological issues

The CO₂ figures in the inventory are based on emission figures reported from the plant to the agency. The emission estimates are based on the amount of calcium carbide produced each year and an emission factor estimated by SINTEF (1998d). Some of the carbon from petrol coke will be sequestered in the product, but not permanently. Thus, this carbon is included in the emission estimate.

4.3.4.3 Activity data

The amount of calcium carbide produced is reported by the plant to the agency.

4.3.4.4 Emission factors

The emission factor used by the plants in the calculation of CO₂ has been estimated to be 1.69 tonne/tonne CaC₂ by SINTEF (1998d). An additional 0.02 t CO₂ /t CaC₂ from fuel is reported in the Energy chapter.

4.3.4.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.3.4.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII.

4.3.4.7 Category specific recalculations

There have been no recalculations performed for this source category this year.

4.3.4.8 Category-specific planned improvements

Since the plant is closed down there is no further planned activity to review historical data.

4.3.5 Titanium dioxide production, 2B6 (Key category for CO₂)

4.3.5.1 Category description

One plant producing titanium dioxide slag is included in the Norwegian Inventory and it was included in the EU ETS in 2013. The plant also produced pig iron as a by-product. The titanium dioxide slag and pig iron are produced from the mineral ilmenite and coal is used as a reducing agent. Various components included CO₂ are emitted during the production process.

CO₂ from titanium dioxide production is defined as a key category according to the approach 1 analysis.

4.3.5.2 Methodological issues

The method that is used for all years can be defined as a calculation based on carbon balance. This method accounts for all the carbon in the materials entering the process and subtracts the CO₂ captured in the products.

4.3.5.3 Activity data

The carbon inputs are dominated by coal, but there is also some pet coke, electrodes, carbides and some masses. Table 4.20 shows the carbon inputs for 2017.

Table 4.20 Carbon inputs (tonnes) for titanium dioxide production in 2017.

Activity data	Amount (tonne)
Melting mass	134
Carbides	418
Coal	89 043
Clay	172
Slag	179 831
Iron	73 589
Antracites	3 743
Ore	292 400
Electrode mass (tonne dry	1 873
Pet coke	2 739

Source: Norwegian Environment Agency

4.3.5.4 Emission factors

The mass balance for each year requires emission factors. The emission factors for the most important carbon inputs for 2017 are shown in Table 4.21.

Table 4.21 Emission factors for some of the carbon inputs in 2017.

Activity data	EF (tonne CO ₂ /tonne)
Coal	2.70
Iron	0.15
Antracites	3.19
Electrode mass (tonne dry weight)	3.50
Pet coke	3.19

Source: Norwegian Environment Agency

4.3.5.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.3.5.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant has reported under the voluntary agreement and the emissions are now covered by the EU ETS and the emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.5.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.5.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.6 Methanol, 2B8a

4.3.6.1 Category description

One plant established in 1997 produces methanol and it is covered by the EU ETS. Natural gas and oxygen are used in the production of methanol. The conversion from the raw materials to methanol is done in various steps and on different locations at the plant. CH₄, N₂O and NMVOC are emitted during the production process. The emissions reported in this category includes flaring and combustion of fuels derived from the natural gas feedstock.

The CO₂ emissions from other energy combustion are included under 1.A.2.C. Indirect emissions of CO₂ are calculated by Statistics Norway based on the emission of CH₄ and NMVOC, see chapter 9 for details about EFs.

4.3.6.2 Methodological issues

The plant reports emission figures of CO₂, CH₄, N₂O and NMVOC to the Norwegian Environment Agency. The reported emissions from flaring and combustion of derived fuels are based on the amounts of gas multiplied by emission factors while the diffuse CH₄ and NMVOC emissions are estimated through the use of the measuring method DIAL (Differential Absorption LIDAR) about every third year since 2002. The plant was divided into various process areas and measurements were taken for at least two days for all process areas. The DIAL method results in an emission factor

per operating hour and this forms the basis for the plant's reported diffuse NMVOC and CH₄ emissions from the production of methanol. This method has been used from 2008 and onwards. This method therefore results in a fixed emission level for some years. The time series for the years 1997-2007 are based on the results from 2008 together with the production levels of methanol for these years.

Minor N₂O emissions are reported under 2B10 Petrochemical N₂O in the CRF. The NMVOC emissions included in the inventory are based on the reported emissions from the plant as these appear to be consistent.

4.3.6.3 Activity data

The annual emissions from flaring and combustion of derived fuels are based on the reported combusted amounts. The activity data used to calculate the indirect CO₂ emissions are the diffuse emissions of CH₄ and NMVOC.

4.3.6.4 Emission factors

CO₂

The plant concerned is part of the EU ETS and the EFs for flaring and combustion of derived fuels are reported annually since 2008.

4.3.6.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

As the reported emissions have varied greatly (e.g. emissions from flaring were much higher in 2000 than in 1999 and 2001), IEFs based on production figures will also fluctuate.

4.3.6.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant is covered by the EU ETS and the emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.6.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.6.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.7 Ethylene, 2B8b

4.3.7.1 Category description

Two plants report emissions under this source category and they are both covered by the EU ETS. One of the plants produces ethylene while the other produces polyethylene and polypropylene.

The majority of the emissions reported here are from flaring. In addition, CH₄ and NMVOC emissions are reported from leakages in the process. Indirect emissions of CO₂ from CH₄ and NMVOC are also

calculated and reported. Minor N₂O emissions are reported under 2B10 Petrochemical N₂O in the CRF.

4.3.7.2 Methodological issues

CO₂, CH₄ and NMVOC

Direct emissions are annually reported to the agency. CO₂ from flaring is based on gas specific emissions factors and activity data. CH₄ and NMVOC emissions reported are based on measurements.

Indirect emissions of CO₂ calculated by Statistics Norway are based on the emission of CH₄ and NMVOC.

4.3.7.3 Activity data

For CO₂ from flaring, the annual emissions from flaring are based on the combustion of natural gas in the flare. The activity data used to calculate the indirect CO₂ emissions are the diffuse emissions of CH₄ and NMVOC.

4.3.7.4 Emission factors

CO₂

The plants report the emission factors used as part of their reporting under the EU ETS.

4.3.7.5 Uncertainties and time-series consistency

Uncertainty estimates are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.3.7.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.7.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.7.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.8 Ethylene dichloride and vinyl chloride monomer, 2B8c

4.3.8.1 Category description

A plant producing vinyl chloride reports CO₂ process emissions that stem from recycling hazardous waste to hydrochloric acid. CH₄ and NMVOC emissions are reported from leakages in the process.

4.3.8.2 Methodological issues

CO₂, CH₄ and NMVOC

The plant has annually reported process emissions to the agency. From 2013 and onwards, the reported CO₂ emissions are based on measurements of the amounts of gas multiplied by emission factors. For the years prior to 2013, the reported CO₂ emissions are based on the amount of hazardous waste recycled to hydrochloric acid multiplied by emission factors. The CH₄ and NMVOC emissions are reported annually to the Norwegian Environment Agency and are based on measurements.

Indirect emissions of CO₂ calculated by Statistics Norway are based on the emissions of CH₄ and NMVOC.

4.3.8.3 Activity data

The plant has reported the amounts of gas relevant for estimating CO₂ emissions since 2013 and prior to 2013 the amounts of recycled hazardous waste. The amounts of CH₄ are reported in the CRF.

4.3.8.4 Emission factors

The emission factors used to estimate CO₂ emissions for the years 2013-2017 are in the range of 0.040503-0.0469792 tonnes CO₂/tonne gas respectively. The emission factors used to estimate CO₂ emissions for the years 2010, 2011 and 2012 are 1.1, 1.1 and 1.08 tonnes CO₂/tonne recycled hazardous waste respectively. The emission factors are all plant specific.

The reported diffuse emissions of CH₄ are based on a measurement derived emission factor of 5 kg CH₄ per operating hour. See chapter 9 for details concerning the EFs used for indirect CO₂ emissions from CH₄ and NMVOC.

4.3.8.5 Uncertainties and time-series consistency

Although two different methods have been used to estimate the CO₂ emissions from recycling hazardous waste, we have no indications that the time series is not consistent. Uncertainty estimates for greenhouse gases are given in Annex II.

4.3.8.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant is covered by the EU ETS and the emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.3.8.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.8.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.3.9 Other, production of fertilizers, 2B10


4.3.9.1 Category description

A plant producing fertilizers has since 2011 reported N₂O emissions from its production to the agency. Urea nitrate is added to the process to reduce the formation of NO_x emissions and this process forms N₂O emissions.

4.3.9.2 Methodological issues

According to the plant, the formation of NO_x is reduced through the use of urea nitrate and cyanic acid. The process forms N₂O, see formulas below.

Reduction of NO_x through Urea nitrate and cyanic acid:



N₂O formation:



The emissions of N₂O are based on measurements of gas volumes and samples are taken for analysis by gas chromatograph. The plant has reported N₂O emissions since 2011 and the Norwegian Environment Agency has estimated the emissions for the years 1990-2010 based on production levels and assumptions about the IEF. There are many factors that influence the emissions and these have varied over time. Such factors are production levels, composition of phosphates, use of urea etc. The plant's reporting of emissions for the years 2011-2013 results in an average IEF of 0.27 kg N₂O per tonne produced fertilizer. This IEF was used to estimate the emissions for the years 2007-2010 as the factors influencing the emissions were similar to 2011-2013. In the years 2002-2006, the plant used more of one type of phosphate than in the period 2007-2013 and the IEF is therefore assumed to be 25% lower. The use of the phosphate type was even larger in the years 1990-2001 and the IEF is therefore assumed to be 50% lower than for the years 2007-2013.

4.3.9.3 Activity data

See description in chapter 4.3.9.2. Although there are several factors that influence the emissions, the production of fertilizers is included as activity data in CTF table 2(I).A-Hs1.

4.3.9.4 Emission factors

See description in chapter 4.3.9.2.

4.3.9.5 Uncertainties and time-series consistency

The estimates for the years 1990-2010 are very uncertain since there are many factors that could influence the emissions. Chapter 4.3.9.2 describes how the emissions for 1990-2010 were estimated and explains the differences in IEF over time. Uncertainty estimates for greenhouse gases are given in Annex II.

4.3.9.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The emissions in this category are not covered by the EU ETS, but the emissions have been reported since 2011 and are considered and tracked by the agency's inventory team.

4.3.9.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.3.9.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4 Metal industry – 2C

The Metal industry in Norway includes plants producing iron and steel, ferroalloys, aluminum, magnesium, zinc, anodes and nickel, see Table 4.22. Nearly all emissions figures from the production of metals included in the inventory are figures reported annually from the plants to the agency.

Table 4.22 Metal industry. Components included in the inventory, tier of method and key category.

Source category	CO ₂	CH ₄	PFCs	SF ₆	Tier	Key category
2C1a. Iron and steel production	R	NA	NA	NA	Tier 3	No
2C2. Ferroalloys production *	R	R	NA	NA	Tier 2/3	Yes
2C3. Aluminium production	R	NA	R	R	Tier 2/3	Yes
2C4. Magnesium production	E	NA	NA	R	Tier 2	Yes
2C6. Zinc production	R + E	NA	NA	NA	Tier 2	No
2C7i. Anode production	R	NA	NA	NA	Tier 2	No
2C7ii. Nickel production	R	NA	NA	NA	Tier 2	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

* Small N₂O emissions from 2C2 are reported under 2C7i Ferroalloys N₂O in the CRF, but are included in Table 4.23 below.

Table 4.23 shows the trends for the sector Metal Production (2C) as a whole and for the various source categories. The GHG emissions from this sector category were about 4.8 million tonnes in 2017, this is 56.1 % of the total emission from the IPPU-sector. The largest contributors to the GHG emissions from Metal industry in 2017 are Ferroalloy production and Aluminum production. The emissions from this sector decreased by 52.2 % from 1990. The reduction since 1990 is due to decreased PFC and SF₆ emissions that again were due to improvement in technology aluminum production, the close down of a magnesium plant in 2006 and generally lower production volumes. The emissions increased by 0.7 % from 2016 to 2017. There was a large increase in emissions from 2009 to 2010, this is mainly due to a low production level for ferroalloys in 2009. The production level in 2009 is also lower than 2008 and reflects the lower economic activity due to the economic recession.

Table 4.23 Emission trends for 2C Metal industry (kt CO₂ equivalents).

Source category	1990	1990, % IPPU	2016	2017	2017, % IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2C1a. Iron and steel production	12.4	0.1 %	26.8	27.4	0.3 %	121.7 %	2.3 %
2C2. Ferroalloys production	2 560.0	17.7 %	2 616.9	2 644.5	30.6 %	3.3 %	1.1 %
2C3. Aluminium production	5 313.8	36.7 %	2 061.5	2 073.0	24.0 %	-61.0 %	0.6 %
2C4. Magnesium production	2 172.8	15.0 %	0.0	0.0	0.0 %	NA	NA
2C6. Zinc production	3.0	0.0 %	2.2	2.7	0.0 %	-7.0 %	27.0 %
2C7i. Anode production	43.8	0.3 %	82.8	76.6	0.9 %	74.7 %	-7.5 %
2C7ii. Nickel production	7.6	0.1 %	14.9	14.5	0.2 %	90.8 %	-2.8 %
2C. Total	10 113.3	69.8 %	4 805.0	4 838.7	56.1 %	-52.2 %	0.7 %

Source: Statistics Norway and Norwegian Environment Agency

4.4.1 Steel, 2C1a

4.4.1.1 Category description

Norway includes one plant producing steel that is covered by the EU ETS and the activity data in the CRF is steel produced.

4.4.1.2 Methodological issues

The total emissions from steel production cover emissions from industrial processes and from combustion, but only the process emissions are reported in this sub-category. Emission figures of CO₂ annually reported to the Norwegian Environment Agency are used in the Norwegian GHG Inventory. This reporting includes both the reporting under the EU ETS and reporting as required under its regular emission permit. The emission figures are based on mass balance calculations.

4.4.1.3 Activity data

The process CO₂ emissions stem from an Electric Arc Furnace (EAF) where scrap iron is melted with other carbon materials. The emissions from the scrap iron are calculated based on the use of each types of scrap iron and the appurtenant content of carbon in each type of scrap iron. The types of scrap iron are according to the UK steel protocol and the carbon content in the types of scrap used varies from 0.15 % up to 4 %. The other input materials to the EAF are coal, lime and the metals ferromanganese, ferrosilicon and silicomanganese and electrodes. The outputs are steel, dust and slag. The net emissions from the mass balance are the process emissions.

4.4.1.4 Emission factors

Since a mass balance is used, it is the carbon contents of the carbon materials that go into the mass balance that are used. For the scrap iron, all ten types of scrap iron have their own carbon content.

4.4.1.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category. The process emissions prior to 2005 have to a large extent been estimated based on the process emissions per ton steel produced in 1998 and 2005, this explains the increasing variation in the CO₂ IEF for steel after 2005 since the emissions from 2005 and onwards are based on annual reported data from the EU ETS.

4.4.1.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant is covered by the EU ETS and the emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.4.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.


4.4.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4.2 Production of Ferroalloys, 2C2 (Key category for CO₂)

4.4.2.1 Category description

There are 12 plants producing ferroalloys in Norway and the plants were included in the EU ETS in 2013. One plant closed down in 2001, two plants were closed down during 2003 and two in 2006. The plant that was out of production in 2006 started up again in 2007. Ferrosilicon, silicon metal, ferromanganese and silicon manganese are now produced in Norway. Ferrochromium was produced until the summer in 2001. Ferro silicon with 65 to 96 % Si and silicon metal with 98-99 % Si is produced. The raw material for silicon is quartz (SiO₂). SiO₂ is reduced to Si and CO using reducing agents like coal, coke and charcoal.



The waste gas CO and some SiO burns to form CO₂ and SiO₂ (silica dust).

In ferroalloy production, raw ore, carbon materials and slag forming materials are mixed and heated to high temperatures for reduction and smelting. The carbon materials used are coal, coke and some bio carbon (charcoal and wood). Electric submerged arc furnaces with graphite electrodes or consumable Söderberg electrodes are used. The heat is produced by the electric arcs and by the resistance in the charge materials. The furnaces used in Norway are open, semi-covered or covered.

The CO is a result of the production process. In open or semi- closed furnaces the CO reacts with air and forms CO₂ before it is emitted. This is due to high temperature and access to air in the process. In a closed furnace the CO does not reach to CO₂ as there are no access to air (oxygen) in the process. The waste gas is then led from furnace and used as an energy source or flared and is reported under the relevant Energy sectors. The technical specification of the furnaces is irrelevant since emissions are calculated using a mass balance or calculated by multiplying the amount of reducing agents in dry weight with country specific EFs.

Several components are emitted from production of ferroalloys. Emission of CO₂ is a result of the oxidation of the reducing agent used in the production of ferroalloys. In the production of FeSi and silicon metal NMVOC and CH₄ emissions originates from the use of coal and coke in the production processes. From the production of ferro manganes (FeMn), silicon manganes (SiMn) and ferrochromium (FeCr) there is only CO₂ emissions.

Measurements performed at Norwegian plants producing ferroalloys indicate that in addition to emissions of CO₂ and CH₄ also N₂O is emitted. Due to the CRF, the N₂O emissions are reported in 2C7i.

The large increase in emissions from 2009 to 2010 is due to a low production level for ferroalloys in 2009. The production level in 2009 is also lower than 2008 and reflects the lower economic activity due to the economic recession.

CO₂ emissions from production of ferroalloys is defined as a key category according to the approach 2 analysis.

4.4.2.2 Methodological issues

CO₂

The methods used in the calculation of CO₂ emissions from production of ferroalloy is in accordance with the method recommended by the IPCC (2006). Emissions are reported by each plant in an annual report to the agency.

The plants have used one of the two methods below for calculating CO₂-emissions:

1. Mass balance; the emissions for CO₂ is calculated by adding the total input of C in raw materials before subtracting the total amount of C in products, wastes and sold gases (Tier 3).
2. Calculate emission by multiplying the amount of reducing agents in dry weight with country specific emission factors for coal, coke, petrol coke, electrodes, anthracite, limestone and dolomite (Tier 2).

Each plant has for consistency just used one method for the entire time series.

Indirect emissions of CO₂ are calculated based on the emission of CH₄ and are reported in this sub-category.

CH₄ and N₂O

The emissions of CH₄ and N₂O are calculated by multiplying the amount of ferroalloy produced with an emission factor. Emissions are reported by each plant in an annual report to the agency.

Plants producing ferro manganese, silicon manganese and ferrochromium do not emit emissions of CH₄ and N₂O.

NMVOC

The emissions are estimated by Statistics Norway from the consumption of reducing agents and an emission factor.

4.4.2.3 Activity data

CO₂

Calculation of emissions is based on the consumption of gross reducing agents and raw materials (carbonate ore, limestone and dolomite). Note that the use of limestone and dolomite and the corresponding emissions are included here under 2C2.

Table 4.24 shows the amount of reducing agents used as activity data in the CRF for some selected years. The reducing agents include the use of bio carbon and the use increased from about 2001.

Table 4.24 Tonnes of reducing agents in the ferroalloys production for some selected years.

Activity data	1990	2000	2010	2015	2016	2017
Coal (dry weight)	395 255	544 946	360 291	490 798	496 292	501 551
Coke (dry weight)	379 028	450 096	328 013	349 898	366 508	374 025
Electrodes	34 748	48 137	48 813	51 547	50 052	49 804
Petrol coke	8 423	12 935	7 793	15 220	18 210	8 537
Pulverised coke	-	0	9 708	14 012	13 305	20 132
Bio carbon	16 565	17 451	104 013	144 639	164 535	166 190
Total	834 019	1 073 565	858 631	1 066 114	1 108 903	1 120 239
Bio as % av total	2 %	2 %	12 %	14 %	15 %	15 %

Source: Norwegian Environment Agency

CH₄ and N₂O

The gross production of different ferroalloys is used in the calculation.

NMVOC

The gross amount of reducing agents that are used for the calculation of NMVOC emissions are annually reported to Statistics Norway from each plant.

4.4.2.4 Emission factors

CO₂

The carbon content of each raw materials used in the Tier 3 calculation is from carbon certificates from the suppliers. The carbon in each product, CO gas sold et cetera is calculated from the mass of product and carbon content. In the Tier 2 calculation the emission factors are as listed in Table 4.25.

Table 4.25 Emission factors from production of ferroalloys. Tonnes CO₂/tonne reducing agent or electrode

	Coal	Coke	Electrodes	Petrol coke	Carbonate ore	Dolomite Limestone
Ferro silicon	3.08	3.36	3.36	--	--	--
Silicon metal	3.12	3.36	3.54	--	--	--
Ferro chromium	--	3.22	3.51	--	--	--
Silicon	--	3.24	3.51	3.59	0.16- 0.35	0.43-0.47
Ferro	--	3.24	3.51	3.59	0.16- 0.35	0.43-0.47

Source: SINTEF (1998b), SINTEF (1998c), SINTEF (1998a)

CH₄ and N₂O

Measurements performed at Norwegian plants producing ferro alloys indicate emissions of N₂O in addition to CH₄. The emissions of CH₄ and N₂O are influenced by the following parameters:

- The silicon level of the alloy (65, 75, 90 or 98 % Si) and the silicon yield
- The method used for charging the furnace (batch or continuously)
- The amount of air used to burn the gases at the top controlling the temperature in off gases.

Measurement campaigns at silicon alloy furnaces have been performed since 1995, and these measurements are the base for the values in the BREF document for silicon alloys. The results of the measurements, that the emissions factors in the Norwegian CH₄ and N₂O are based upon, are presented in SINTEF (2004). A summary of the report is given in the publication "Reduction of

emissions from ferroalloy furnaces" (Grådahl et al. 2007). The main focus for the studies has been NO_x emissions. However, the emissions of CH₄ and N₂O have also been measured.

Full scale measurements have been performed at different industrial FeSi/Si furnaces. The average CH₄ and N₂O concentrations in the ferroalloy process are with some exceptions a few ppm. For N₂O and CH₄ the exception is during spontaneous avalanches in the charge (i.e. collapse of large quantities of colder materials falling into the crater or create cavities) occur from time to time, see Figure 7 in Grådahl et al. (2007). In the avalanches the N₂O emissions go from around zero to more than 35 ppm. The avalanches are always short in duration. There are also increased N₂O emissions during blowing phenomenon.

The EF used in the inventory represents the longer-term average N₂O and CH₄ concentration measurements outside the peaks in concentrations. The peaks in concentration occur due to avalanches (sudden fall of large amount of colder charge into the furnace) that occur from time to time is not fully reflected in the EFs. The EFs used we regard as conservative particular for the early 1990s when the avalanches were more frequent than the latest years.

All companies apply sector specific emission factors in the emission calculation, see Table 4.26. The factors are developed by the Norwegian Ferroalloy Producers Research Organisation (FFF) and standardized in meeting with The Federation of Norwegian Process Industries (PIL) (today named Federation of Norwegian Industries) in February 2007.

NMVOC

Statistics Norway uses an emission factor of 1.7 kg NMVOC/tonne coal or coke in the calculations (Limberakis et al. 1987).

Table 4.26 Emission factors for CH₄ and N₂O from production of ferroalloys.

	Si-met			FeSi-75%			FeSi-65%		
Alloy, charging routines and temperature	Batch-charging	Sprinkle-charging ¹	Sprinkle-charging and >750°C ²	Batch-charging	Sprinkle-charging ¹	Sprinkle-charging and >750°C ²	Batch-charging	Sprinkle-charging ¹	Sprinkle-charging and >750°C ²
kg CH ₄ per tonne metal	0.1187 (M)	0.0881 (M)	0.1000 (E)	0.0890 (E)	0.0661 (E)	0.0750 (E)	0.0772 (E)	0.0573 (E)	0.0650 (E)
kg N ₂ O per tonne metal	0.0433 (E)	0.0214 (E)	0.0252 (E)	0.0297 (E)	0.0136 (E)	0.0161 (E)	0.0117 (E)	0.0078 (E)	0.0097 (E)

¹ Sprinkle-charging is charging intermittently every minute.

² Temperature in off-gas channel measured where the thermocouple cannot 'see' the combustion in the furnace hood.

M=measurements and E= estimates based on measurements

4.4.2.5 Uncertainties and time-series consistency

The uncertainty in activity data and emission factors have been calculated to ±5 % and ±7 % respectively, see Annex II.

The IEF (tonne CO₂/tonne reducing agent) for the ferroalloys production has a downward trend from around the year 2001. This is due to the increased use of bio carbons. Fluctuations in the IEF can also be due to variations in use of the various reducing agents, amounts of sold CO and production of ferro alloy products.

4.4.2.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants have reported under the voluntary agreement and the emissions are now covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

Statistics Norway makes in addition occasional quality controls (QC) of the emission data on the basis of the consumption of reducing agents they collect in an annual survey and average emission factors.

4.4.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4.3 Aluminium production, 2C3 (Key Category for CO₂ and PFC)

4.4.3.1 Category description

One open mill in Norway has handled secondary aluminium production, but it closed down in 2001. Minor emissions of SF₆ in the period 1992-2000 are therefore included in the inventory.

There are seven plants in Norway producing primary aluminium and they were included into the EU ETS in 2013. Both prebaked anode and the Soederberg production methods are used. In the Soederberg technology, the anodes are baked in the electrolysis oven, while in the prebaked technology the anodes are baked in a separate plant. In general, the emissions are larger from the Soederberg technology than from the prebaked technology.

Production of aluminium leads to emission of CO₂ and perfluorocarbons (PFCs). The emission of CO₂ is due to the electrolysis process during the production of aluminium.

There has been a substantial reduction in the total PFC emissions from the seven Norwegian aluminium plants in the period from 1990 to 2017. This is a result of the sustained work and the strong focus on reduction of the anode effect frequency in all these pot lines and that there has been a shift from Soederberg to prebaked technology. The focus on reducing anode effect frequency started to produce results from 1992 for both technologies. For prebaked technology the PFC emissions in kg CO₂ equivalents per tonne aluminium were reduced from 2.99 in 1990 to 2.30 in 1991 and 1.12 in 1992 and respective values for Soederberg were 6.45, 6.09 and 5.78. In 2017 the specific PFC emissions for prebaked and Soederberg were 0.10 and 0.12 kg CO₂-equivalent, see Figure 4.6 and Table 4.27.




Figure 4.6 PFC in kg CO₂ equivalent per tonne aluminium.

Source: Norwegian Environment Agency

Table 4.27 Shares of the technologies used in aluminum production and the PFC IEFs.¹³

Year	Share of production from Soederberg technology	Share of production from pre-baked technology	PFC IEF Soderberg	PFC IEF pre-baked
1990	43%	57 %	6.45	2.99
1995	39 %	61 %	5.81	0.78
2000	39 %	61 %	3.26	0.35
2005	20 %	80 %	2.32	0.28
2008	15 %	85 %	1.33	0.53
2009	8 %	92 %	0.21	0.41
2010	8 %	92 %	0.31	0.21
2011	8 %	92 %	0.33	0.23
2012	7%	93%	0.29	0.15
2013	8%	92%	0.26	0.15
2014	8%	92%	0.10	0.16
2015	8%	92%	0.12	0.12
2016	8%	92%	0.11	0.15
2017	7%	93%	0.12	0.10

Source: Norwegian Environment Agency

In 1990, 57 % of the aluminium production in Norway was produced with prebaked technology and the share of aluminium production from prebaked was increased to 93 % in 2017. Two new plants with prebaked technology were established in 2002 and plants using Soederberg technology were closed down in the period 2002-2009. The shares of the two technologies and their PCF IEFs are shown in Table 4.27. The PFCs emissions from production of aluminium have decreased by 96.6 % from 1990 to 2017.

¹³ PFC in kg CO₂ equivalents per tonne aluminium

The PFC emissions per tonne aluminium produced in Norway was 4.48 kg CO₂ equivalents in 1990 and 0.11 kg CO₂ equivalents in 2017. This is a reduction of 97.7 % from 1990 to 2017.

An increase in production capacity is also included in the modernisation, leading to higher total emissions of CO₂. PFCs and CO₂ emissions from aluminium production are both identified as key categories according to the approach 2 analysis.

4.4.3.2 Methodological issues

CO₂

The inventory uses the emission figures reported to the Norwegian Environment Agency calculated by each plant.

For the years including 2012, the aluminium industry calculated the CO₂ emissions separate for each technology on the basis of consumption of reducing agents. This includes carbon electrodes, electrode mass and petroleum coke. The emissions factors are primarily calculated from the carbon content of the reducing agents.

The following methods were used up to 2012:

CO₂ from Prebake Cells

$$(4.9) \quad Q = A \cdot C \cdot 3.67$$

Where

Q is the total yearly emissions of CO₂

A is the yearly net consumption of anodes

C is % carbon in the anodes

3.67 is the mol-factor CO₂/C

CO₂ from Soederberg Cells

$$(4.10) \quad Q = S \cdot 3.67 \cdot (K \cdot C1 + P \cdot C2)$$

Where

Q is the total yearly emissions of CO₂

S is the yearly consumption of Soederberg paste

K is the share of coke in the Soederberg paste

P is the share of pitch in the Soederberg paste

K+P=1

C1 is the fraction of carbon in the coke. Fraction is % Carbon/100

C2 is the fraction of carbon in the pitch. Fraction is % Carbon/100

From 2013 and onwards, the CO₂ emissions from Soederberg cells and from Prebake cells are calculated using the mass balance methodology that considers all carbon inputs, stocks, products and other exports from the mixing, forming, baking and recycling of electrodes as well as from electrode consumption in electrolysis. We have no indications that this has resulted in an inconsistent time series.

PFCs


Perfluorinated hydrocarbons (PFCs), e.g. tetrafluoromethane (CF₄) and hexafluoroethane (C₂F₆), are produced during anode effects (AE) in the Prebake and Soederberg cells, when the voltage of the cells increases from the normal 4-5V to 25-40V. During normal operating condition, PFCs are not

produced. The fluorine in the PFCs produced during anode effects originates from cryolite. Molten cryolite is necessary as a solvent for alumina in the production process.

Emissions of PFCs from a pot line (or from smelters) are dependent on the number of anode effects and their intensity and duration. Anode effect characteristics will be different from plant to plant and also depend on the technology used (Prebake or Soederberg).

During electrolysis two per fluorocarbon gases (PFCs), tetrafluormethane (CF₄) and heksafluorethane (C₂F₆), may be produced in the following reaction:

Reaction 1



Reaction 2



The national data are based on calculated plant specific figures from each of the Norwegian plants. A Tier 2 method is used in the calculations, which are based on a technology specific relationship between anode effect performance and PFCs emissions. The PFCs emissions are then calculated by the so-called slope method, where a constant slope coefficient (see Table 4.28), is multiplied by the product of anode effect frequency and anode effect duration (in other words, by the number of anode effect minutes per cell day), and this product is finally multiplied by the annual aluminum production figure (tonnes of Al/year). The formula for calculating the PFCs is:

kg CF₄ per year = S_{CF4} • AEM • MP and

kg C₂F₆ per year = kg CF₄ per year • F_{C2F6/CF4}

Where:

S_{CF4} = "Slope coefficient" for CF₄, (kg PFC/t_{Al}/anode effect minutes/cell day

AEM = anode effect minutes per cell day

MP = aluminium production, tonnes Al per year

F_{C2F6/CF4} = weight fraction of C₂F₆/CF₄

Table 4.28 Technology specific slope and overvoltage coefficients for the calculation of PFCs emissions from aluminium production.

Technology ^a	"Slope coefficient" ^{b, c} (kg PFC/t _{Al})/ (anode effect/cell day)		Weight fraction C ₂ F ₆ /CF ₄	
	S _{CF₄}	Uncertainty (±%)	F _{C₂F₆/CF₄}	Uncertainty (±%)
CWPB	0.143	6	0.121	11
SWPB	0.272	15	0.252	23
VSS	0.092	17	0.053	15
HSS	0.099	44	0.085	48

a. Centre Worked Prebake (CWPB), Side Worked Prebake (SWPB), Vertical Stud Søderberg (VSS), Horizontal Stud Søderberg (HSS).

b. Source: Measurements reported to IAI, US EPA sponsored measurements and multiple site measurements.

c. Embedded in each slope coefficient is an assumed emission collection efficiency as follows: CWPB 98%, SWPB 90%, VSS 85%, HSS 90%. These collection efficiencies have been assumed based on measured PFC collection fractions, measured fluoride collection efficiencies and expert opinion.

"Slope coefficient": The connection between the anode parameters and emissions of PFC.

Measurements of PFCs at several aluminium plants have established a connection between anode parameters and emissions of CF₄ and C₂F₆. The mechanisms for producing emissions of PFC are the same as for producing CF₄ and C₂F₆. The two PFC gases are therefore considered together when PFC emissions are calculated. The C₂F₆ emissions are calculated as a fraction of the CF₄ emissions.

The Tier 2 coefficients for Centre Worked Prebaked cells (CWPB) are average values from about 70 international measurement campaigns made during the last decade, while there are fewer data (less than 20) for Vertical Stud Soederberg cells (VSS). The main reason for the choice of the Tier 2 method is that the uncertainties in the facility specific slope coefficients is lower than the facility specific based slope coefficients in Tier 3. This means that there is nothing to gain in accuracy of the data by doing measurements with higher uncertainties.

"Slope coefficient" is the number of kg CF₄ per tonne aluminium produced divided by the number of anode effects per cell day. The parameter cell day is the average number of cells producing on a yearly basis multiplied with the number of days in a year that the cells have been producing.

Sulphur hexafluoride (SF₆)

SF₆ used as cover gas in the aluminium industry is assumed to be inert, and SF₆ emissions are therefore assumed to be equal to consumption. At one plant SF₆ was used as cover gas in the production of a specific quality of aluminium from 1992 to 1996. The aluminium plant no longer produces this quality, which means that SF₆ emissions have stopped.

4.4.3.3 Activity data

The consumption of reducing agents and electrodes and the production of aluminium is reported annually to the agency and the latter is reported as activity data in the CRF tables. The mass balance summarising the consumption of reducing agents in 2017 is shown in Table 4.29.

Table 4.29. Mass balance for aluminium production in 2017.

Reducing agent	Amount (tonne)	CO2 emissions (tonne)
Petroleum coke	278 235	989.0
Prebaked anodes	630 304	752.3
Pitch	71 129	241.8
Coke	32 555	115.8
Soda	712	0.3
Anode remnant	137 171	-151.9
Waste	3 776	-5.1
Tar	63	-0.1
Total	1 153 947	1942.0

Source: Norwegian Environment Agency

PFCs

The basis for the calculations of PFCs is the amount of primary aluminium produced in the pot lines and sent to the cast house. Thus, any remelted metal is not included here.

4.4.3.4 Emission factors

There are many factors for the carbon contents of the reducing agent that are used for the mass balances reported by the plants. The contents in tonne C/tonne reducing agent for the most important reducing agents in 2017 are 0.97-0.9786 for petroleum coke, 0.960704-0.975 for prebaked anodes and 0.927734-0.9281657 for pitch.

The PFC emissions are calculated using the Tier 2 recommended values by IAI (2005) for CF₄ (the slope coefficients of 0.143 kg CF₄/tonne Al/anode effect minutes per cell day for CWPB and 0.092 for VSS). The amount of C₂F₆ is calculated from the Tier 2 values for CF₄, where the weight fraction of C₂F₆ to CF₄ is set equal to 0.121 for CWPB and 0.053 for VSS. This is consistent with the 2006 IPCC GL. All values are technology specific data, recommended by IAI. Our facility specific measured data that we have used until today are all in agreement with these data, within the uncertainty range of the measurement method employed.

4.4.3.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

PFCs

The uncertainties in the so-called Tier 2 slope coefficients from IAI is lower (6% and 17% for CWPB and VSS cells, respectively), compared to the measured facility specific based slope coefficients, where the uncertainties are around 20%, even when the most modern measuring equipment is used (the continuous extractive-type Fourier Transform Infrared (FTIR) spectroscopic system). Control measurements in two Hydro Aluminium plants (Karmøy and Sunndal) done by Jerry Marks in November 2004, showed that the measured values for CWPB and VSS cells were well within the uncertainty range of the Tier 2 slope coefficients.

Chapter 4.4.3.1 explains the downward trend of the IEF for PCF emissions, but there are also some inter-annual changes that can be explained. The reduced IEF for Soederberg from 2002 to 2003 is due to the fact that one plant using this technology closed down and had no production in 2003. This plant produced 18% of the aluminium produced with this technology in 2002 and had an IEF in 2002

that was the highest among all the plants producing with Soederberg technology in that year. The reduced IEF for Soederberg from 2008 to 2009 is due to the fact that another plant using this technology closed down in 2009. This plant produced 56% of the aluminium produced with this technology in 2008 and the production in 2009 was minor. The plant's IEF in 2008 was the highest among all the plants producing with Soederberg technology in that year. The variations in the IEF for prebaked between 2014 and 2017 are due variations in the anode effect frequency for one plant.

CO₂

The implied emission factor for CO₂ is relatively stable over the time series. The largest inter-annual changes in the IEF are from 2009 to 2010 and from 2010 to 2011 and can be explained by production problems at one plant in 2010. The concerned plant produced about 18% of the total aluminium in 2010 and uses the prebaked technology. Its CO₂ IEF in 2010 was unusually high since the consumption of anodes per tonne aluminium produced were 22 % higher in 2010 than in comparable years.

With the inclusion of the aluminium and anode production in the EU ETS system from 2013, a new methodology was introduced for the calculation of CO₂ emissions from anode production in integrated aluminium and anode plants. For one plant, it was no longer possible to split CO₂ process emission between aluminium and anode production and all the emissions from this plant are reported in 2C3.

4.4.3.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants have reported under the voluntary agreement and the emissions are now covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

As a quality control, it is checked that the reports are complete. Each figure is compared with similar reports from previous years and also analysed taking technical changes and utilisation of production capacity during the year into account. If errors are found the Norwegian Environment Agency contacts the plant to discuss the reported data and changes are made if necessary.

The Norwegian Environment Agency has annual meetings with the aluminium industry where all plants are represented. This forum is used for discussion of uncertainties and improvement possibilities. The agency's auditing department are regularly auditing the aluminium plants. As part of the audits, their system for monitoring, calculation and reporting of emissions are controlled.

The emission figures reported by the plants are also occasionally controlled by Statistics Norway. Statistics Norway make their own estimates based on the consumption of reducing agents and production data collected in an annual survey and average emission factors.

4.4.3.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4.4 Magnesium production, 2C4 (Key category for SF₆)

4.4.4.1 Category description

There was previously one plant in Norway producing magnesium. The plant closed down the production of primary magnesium in 2002 and the production of cast magnesium was closed down in 2006. From the mid-1970s, both the magnesium chloride brine process and the chlorination process were used for magnesium production. Since 1991, only the chlorination process was in use.

Production of magnesium leads to process related CO₂ and CO emissions. During the calcinations of Dolomite (MgCa(CO₃)₂) to magnesium oxide, CO₂ is emitted. During the next step, magnesium oxide is chlorinated to magnesium chloride and coke is added to bind the oxygen as CO and CO₂. SO₂ is emitted due to the sulphur in the reducing agent used.

In the foundry, producing cast magnesium, SF₆ is used as a cover gas to prevent oxidation of magnesium. The Norwegian producers of cast magnesium has assessed whether SF₆ used as a cover gas reacts with other components in the furnace. The results indicate that it is relatively inert, and it is therefore assumed that all SF₆ used as cover gas is emitted to the air.

The emissions decreased due to improvements in technology and in process management. The primary magnesium production stopped in 2002 and only secondary production is retained and this production has no CO₂ emissions from processes. During 2006 also the production of remelting Mg stopped and since then there were no emissions from this source.

SF₆ emissions from magnesium foundries is defined as a key category according to the approach 1 analysis.

4.4.4.2 Methodological issues

CO₂

The Norwegian emission inventory uses production data as activity data. The CO₂ emissions are therefore calculated by using annual production volume and the emission factor recommended by SINTEF (SINTEF 1998e). This is considered to be in line with the tier 2 method in the IPCC 2006 Guidelines (IPCC 2006).

SF₆

The consumption of the cover gas SF₆ is used as the emission estimates in accordance with the tier 2 method in the IPCC 2006 Guidelines (IPCC 2006). The plant reported the emissions each year to the agency.

4.4.4.3 Activity data

In the GHG emission inventory we have used production volumes as activity data in the calculation of CO₂. The plant reported the consumption of SF₆ to the agency.

4.4.4.4 Emission factor

An emission factor of 4.07 tonnes CO₂/tonnes produced magnesium is used to calculate the annual emissions of CO₂ (SINTEF 1998e).

4.4.4.5 Uncertainties and time-series consistency

The uncertainty estimates are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.4.4.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII.

4.4.4.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.4.8 Category-specific planned improvements

Since the plant is closed down there is no further planned activity to review historical data.

4.4.5 Zinc production, 2C6

4.4.5.1 Category description

One plant in Norway produces zinc and the plant's main products are zinc, zinc-alloys and aluminiumfluoride. The plant has reported process emission of CO₂ from the use of ore materials for the year 2012 and onwards.

4.4.5.2 Methodological issues

CO₂

The tier 1 method and a default EF from the 2006 IPCC GL is not applicable since the plant uses an electrolytic process and the 2006 IPCC GL states that this does not result in non-energy CO₂ emissions. However, the plant reports some process emissions of CO₂ from the use of ore materials using a mass balance approach for both roasting and sintering. This is 2 method that is more accurate than the tier 1 method because it takes into account the materials and the variety of furnace types used rather than assuming industry-wide practices.

Emission figures have been reported by the plant to the Norwegian Environment Agency for the year 2012 and onwards and the agency has estimated the emissions for the years 1990-2011 by correlating the annual production levels of zinc with the ratio between process and combustion emissions in 2012. For the years 1990-1993 with no production data available, the emissions have been set equal to the emissions in 1994.

4.4.5.3 Activity data

The plant has reported the amounts of ore materials used for the year 2012 and onwards and it ranges from about 144 000 to 298 000 tonnes.

4.4.5.4 Emission factors

The plant has reported emission factors for the ore materials used for the year 2012 and onwards. The emission factors (t C/t ore material) ranges from 0.00148 to 0.0062.

4.4.5.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.4.5.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant reports annually through its permit and the agency's inventory team tracks emissions and AD for the plant.

4.4.5.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.5.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4.6 Anode production, 2C7i

4.4.6.1 Category description

Four plants in Norway produce anodes and they were included into the EU ETS in 2013. Three plants produce prebaked anodes and one plant produces coal electrodes. These are alternatives to the use of coal and coke as reducing agents in the production process for aluminium and ferroalloys. The anodes and coal electrodes are produced from coal and coke. The production of anodes and coal electrodes leads to emissions of CO₂.

4.4.6.2 Methodological issues

The emissions of CO₂ from the production of anodes are calculated by each plant and the method is based on the Aluminium Sector Greenhouse Gas Protocol by the International Aluminium Institute (IAI 2005).

The emissions are calculated from the consumption of anthracite and petrol coke. In addition, pitch is included in production. The calculations of CO₂ from processes are uptime in hours multiplied with an emission factor for each feedstock. When calcinations of anthracite the EF are 167 kg CO₂ per uptime hour and for petrol coke the EF is 238 kg CO₂. In addition, there are emissions from energy use that is reported in the Energy sector.

From 2012, there was a methodological challenge for integrated anode and aluminum production plants since reported EU ETS data do not provide information to split emissions on the two processes. Equation 4.21 from the 2006 IPCC Guidelines are not used for calculating these emissions in the EU ETS system, where emissions are calculated based on a carbon mass balance approach without information on ash and sulphur content. Therefore, emissions from two integrated plants that previously were reported under 2C7ai were included under 2C3.

4.4.6.3 Activity data

See methodological issues.

4.4.6.4 Emission factors

See methodological issues.

4.4.6.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.4.6.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants have reported under the voluntary agreement and the emissions are now covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.4.6.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.6.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.4.7 Nickel production, 2C7ii

4.4.7.1 Category description

One plant in Norway produces nickel. During the production of nickel, CO₂ is emitted from the use of soda ash.

4.4.7.2 Methodological issues

CO₂ emission figures are annually reported from the plant to the agency.

4.4.7.3 Activity data

The activity data is the annual amounts of soda ash used in the production process, see Table 4.11.

4.4.7.4 Emission factors

An emission factor of 0.41492 tonnes CO₂/tonne soda ash is used for the calculations.

4.4.7.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.4.7.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plant reports as required by its regular permit and has also reported under the voluntary agreement. The agency's inventory team tracks emissions and AD for the plant.

4.4.7.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.4.7.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.5 Non-energy products from fuels and solvent use – 2D

Norway reports the source categories lubricants use, paraffin wax, solvent use, road paving with asphalt and asphalt roofing under the category 2D, see Table 4.30.

Table 4.30 Non-energy products from fuels and solvent use. Components included in the inventory, tier of method and key category.

Source category	CO ₂	Tier	Key category
2D1. Lubricants use	E	Tier 2	Yes
2D2. Paraffin wax use	E	Tier 1	No
2D3a. Solvent use	E	Tier 2	No
2D3b. Road paving with asphalt	E	Tier 1	No
2D3d. Other (use of urea)	E	Tier 1	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

Table 4.31 shows the emission trends for 2D as a whole and for the various source categories. The GHG emissions from this sector category were about 0.2 million tonnes in 2017, this is 2.3 % of the total emission from the IPPU-sector. The emissions from this sector decreased by 29.8 % from 1990 and the emissions decreased by 2.8 % from 2016 to 2017.

Table 4.31 Emission trends for 2D Non-energy products from fuels and solvent use (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2D1. Lubricants use	167.1	1.2 %	42.3	41.6	0.5 %	-75.1 %	-1.7 %
2D2. Paraffin wax use	6.2	0.0 %	46.0	39.8	0.5 %	538.7 %	-13.5 %
2D3a. Solvent use	114.1	0.8 %	103.6	103.2	1.2 %	-9.6 %	-0.4 %
2D3b. Road paving with asphalt	0.0	0.0 %	0.0	0.0	0.0 %	58.0 %	8.4 %
2D3d. Other (use of urea)	0.0	0.0 %	15.6	17.1	0.2 %	NA	9.3 %
2D. Total	287.5	2.0 %	207.6	201.7	2.3 %	-29.8 %	-2.8 %

Source: Statistics Norway and Norwegian Environment Agency

4.5.1 Lubricant use, 2D1 (Key category for CO₂)

4.5.1.1 Category description

Lubricants are mostly used in transportation and industrial applications, and are partly consumed during their use. It is difficult to determine which fraction of the consumed lubricant is actually combusted, and which fraction is firstly resulting in NMVOC and CO emissions and then oxidised to CO₂. Hence, the total amount of lubricants lost during their use is assumed to be fully oxidized and these emissions are directly reported as CO₂ emissions.

Emissions from waste oil handling are reported in the Energy Sector (energy recovery) and in the Waste sector (incineration).

CO₂ emissions from lubricants use is defined as a key category according to the approach 1 analysis.

4.5.1.2 Methodological issues

The CO₂ emissions from lubricant use are estimated by multiplying sold amounts of lubricants (m³) by density, country specific oxidation factors, default NCV value (TJ/tonne), default C content (tonne/TJ) and the mass ratio of CO₂/C:

$$(4.11) \quad E_p = A_p * d * NCV * ODU_p * CC * 44/12$$

where:

E_p = CO₂ emission from product group p

A_p = Sold amount of lubricant from product group p (activity data)

d = Density

NCV = Net calorific value for lubricants

ODU_p = Fraction being oxidized during use from product group p

CC = Carbon content

The method is applied to subgroups of lubricants, as does the tier 2 method in the 2006 guidelines. However, even though the lubricant product groups in the Norwegian inventory are more detailed than in the tier 2 method, no distinction is made between lubricant oil and lubricant wax in the activity data. Thus, tier 1 factors are applied for NCV and CC .

It is assumed that all lubricant consumption and oxidation occurs within the sales year.

4.5.1.3 Activity data

The sold amount of lubricant by product group is given in Statistics Norway's statistics on sales of petroleum products, see Table 4.32. This statistics is based on reporting from the oil companies, and divides the lubricant into five product groups (numbered 204 – 208, see Table 4.32 and Table 4.33).

Table 4.32 Sold amounts of lubricants, except to foreign navigation (1.000 m³), 1990 – 2017.

Year	201	202	203	204	205	206	207	208
1990	99 637	0	0	0	0	0	0	0
1995	40 583	0	0	23 270	0	22 726	0	0
2000	0	29 369	12 734	9 160	13 724	18 594	0	0
2005	0	0	0	13 215	10 751	5 919	33 671	4 233
2009	0	0	0	10 809	12 573	6 642	34 104	4 967
2010	0	0	0	10 412	12 189	4 147	35 434	5 514
2011	0	0	0	9 432	12 897	7 763	35 661	6 230
2012	0	0	0	9 405	11 665	4 188	31 168	5 813
2013	0	0	0	10 161	12 515	5 195	37 047	5 944
2014	0	0	0	5 655	13 627	4 898	28 504	5 973
2015	0	0	0	4 419	11 612	2 317	25 416	5 188
2016	0	0	0	5 230	13 352	3 300	29 586	5 922
2017	0	0	0	5 254	12 640	3 251	28 937	5 961

Source: Statistics Norway

Historically, all lubricant was allocated to product group 201. From 1995 product group 204 and 206 were separated out, and from 1998 the remainder of 201 was split into the product groups 202, 203 and 295. Product groups 207 and 208, which were established in 2003, are reallocations of group 202 and 203.

Table 4.33 Lubricant product groups in the sales of petroleum statistics.

Product group	Product group (text)
201	Lubricants
202	Auto motor and gear oil
203	Navigation and aviation motor and gear oil
204	Industrial lubricants
205	Hydraulic oils
206	Process and transformer oil
207	Motor oil
208	Gear oil

Source: Statistics Norway

The sales statistics does not distinguish between lubricant wax and lubricant oil, and hence the default average (tier 1) carbon content (CC) factor was used.

4.5.1.4 Emission factors

ODU factors

The factors for oxidation during use (ODU) for all product groups 204 to 208 are shown in

Table 4.34. The factors were found by contacting a broad selection of users and purchasers of lubricant oils, as well as branch organisations and interest groups. We have here assumed that loss during use corresponds to oxidation during use, as described above. As the former product groups 201 – 203 are not covered in the report (Weholt et al. 2010), ODUs for these product groups were estimated. The ODU for product group 202 and 203 is simply the average of the ODUs for product number 207 and 208. For product group 201 the ODU in 1990 to 1994 was estimated as the weighted average of ODU for product group 202 to 206, based on sold amounts in 1998. In 1995 to 1997 it was estimated from product group 202, 203 and 205 in 1998.

Table 4.34 Oxidation during use (ODU) factors.

Product group	ODU factor	Source (L = literature, E = estimated)
201 (1990 to 1994)	0.67	E
201 (1995 to 1997)	0.17	E
202	0.175	E
203	0.175	E
204	0.75	L
205	0.15	L
206	0.90	L
207	0.25	L
208	0.10	L

Source: Weholt et al. (2010)

The statistics on sold lubricant include oil combusted in two-stroke petrol engines, and hence considerations must be made in order to avoid double counting. However, the report (Weholt et al. 2010), which is quite detailed when describing the elaboration of ODU factors, does not mention consumption in two-stroke petrol engines. We therefore assume that consumption in two-stroke petrol engines are omitted in the ODU factors, and thus no correction for double counting is necessary.

Other factors

The figures on sold lubricants are given in m³, and must be converted to tonnes. The density varies between different lubricant types, and based on sources available on the Internet it is estimated to 0.85 m³/tonne as an average for all lubricant types, see Exxonmobile (2009) and Neste_Oil (2014). The conversion from tonnes of consumed lubricant to tonnes of emitted CO₂ is performed based on IPCC default factors for energy content (NCV) and carbon content per unit of energy see Table 4.35. This conversion method implicitly adjusts for the content of non-hydrocarbons.

Table 4.35 Other factors.

Factor	Value	Unit	Source
Density (<i>d</i>)	0.85	m ³ /tonne	Producers
Net calorific value (NCV)	0.0402	TJ/tonne	IPCC 2006 GL
Carbon content (CC)	20	Tonne C/TJ	IPCC 2006 GL

4.5.1.5 Uncertainties and time-series consistency

The uncertainty in the estimated emissions from lubricant use (except in two-stroke petrol engines) is assumed to be rather low. The uncertainty in the activity data is assumed to be 5 %, see Table 4.36, in line with the IPCC guidelines for countries with well developed energy statistics. Also the uncertainty of the carbon content is an IPCC default value, and the NCV uncertainty is assumed to be equally large. The uncertainty estimate for the density is based on an expert judgement of the available data on the Internet.

The uncertainty of the country specific ODU estimate is set much lower than for the IPCC default value. This is partly due to the thorough evaluation in the report (Weholt et al. 2010), and partly due to estimations based on the ODUs from this report combined with sales and waste collection statistics, which states that 85 to 90 % of all waste lubricant oil is collected by Statistics Norway (Statistics Norway & SOE Norway 2014). This rather high collection percentage seems reasonable, due to a refund scheme for waste oil combined with strict control of the collected amounts. Higher ODUs would increase this percentage, and vice versa.

Table 4.36 Uncertainty estimates (%).

Parameter	Uncertainty
Activity data (<i>A</i>)	5
Oxidation during use (<i>ODU</i>)	5
Density (<i>d</i>)	3
Net calorific value (NCV)	3
Carbon content (CC)	3

Based on these uncertainties, the overall uncertainty of the emissions from lubricating oil (except from use in two-stroke petrol engines) is estimated at 20 %.

The split of lubricants between different product groups in the activity data have varied throughout the time series, and the level of detail is lower at the beginning of the time series. This might potentially introduce some time series inconsistencies. However, this variation is taken into account for the used ODU factors, and no significant time series inconsistencies are thus expected.

4.5.1.6 Category-specific QA/QC and verification

Emissions from lubricant use are calculated in Excel sheets before being included in the main model. Activity data for the calculations of emissions from lubricants are subject to checks for consistency compared to previous years. Major discrepancies are examined. Periodically, sales statistics are compared to waste statistics as a quality control of level. In addition, the emission estimates are subject to the general QA/QC procedures (see chapter 1.2.3) when included in the main model.

4.5.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.5.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.5.2 Paraffin wax use, 2D2

4.5.2.1 Category description

Paraffin waxes are produced from crude oil and used in a number of different applications, including candles, tapers and the like. Combustion of such products results in emissions of fossil CO₂. Emissions from the incineration of products containing paraffin wax, such as wax coated boxes, are covered by emissions estimates from waste incineration.

4.5.2.2 Methodological issues

Emissions of CO₂ from the burning of candles, tapers and the like are calculated using a modified version of equation 5.4 for Waxes – Tier 1 Method of the 2006 IPCC Guidelines:

$$(4.12) \text{ Emissions} = PC * PF * CC_{Wax} * 44/12$$

Where:

- Emissions = CO₂ emissions from waxes, tonne CO₂
- PC = total candle consumption, TJ
- PF = fraction of candles made of paraffin waxes
- CC_{Wax} = carbon content of paraffin wax (default), tonne C/TJ (Lower Heating Value basis)
- 44/12 = mass ratio of CO₂/C

Consumption figures on paraffin waxes are multiplied by the default net calorific values (NCV). Net consumption in calorific value is then converted to carbon amount, using the value for carbon content (Lower Heating Value basis) and finally to CO₂ emissions, using the mass ratio of CO₂/C.

4.5.2.3 Activity data

Statistics Norway collects data on import, export and sold produce of “Candles, tapers and the like (including night lights fitted with a float)”. Using these data, net consumption of paraffin waxes and other candle waxes (including stearin) can be calculated.

4.5.2.4 Emission factors

Parameter values used in the emissions calculations are given in Table 4.37.

Table 4.37 Parameters employed when calculating emissions.

Parameters	Factor	Unit	References
Net calorific value (NCV)	40.20	TJ/Gg	2006 IPCC
Carbon content (CC_{wax} , Lower Heating Value basis)	20.00	tonnes C/TJ = kg C/GJ	2006 IPCC
Mass ratio of CO_2/C	3.67	-	
Fraction of paraffin wax (PF)	0.66	-	

The assumption of 0.66 as the fraction of all candles being made of paraffin waxes is based on estimates obtained from one major candle and wax importer (estimating ca. 0.5) and one Norwegian candle manufacturer (estimating ca 0.8). The importer estimated the fraction to be ca. 5 % higher in 1990. However, since this possible change is considerably smaller than the difference between the two fraction estimates, we have chosen to set this factor constant for the whole time series. The fraction of paraffin waxes has probably varied during this period, as it, according to the importer, strongly depends on the price relation between paraffin wax and other, non-fossil waxes. However, at present we do not have any basis for incorporating such factor changes.

Furthermore, we assume that practically all of the candle wax is burned during use, so that emissions due to incineration of candle waste are negligible.

4.5.2.5 Uncertainties and time-series consistency

According to the 2006 IPCC Guidelines, the default emission factors are highly uncertain. However, the default factor with the highest uncertainty is made redundant in our calculations, due to the level of detail of our activity data.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.5.2.6 Category-specific QA/QC and verification

There is no specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

4.5.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.5.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.5.3 Solvent use, 2D3a

4.5.3.1 Category description

The use of solvents leads to emissions of non-methane volatile organic compounds (NMVOC) which is regarded as an indirect greenhouse gas. The NMVOC emissions will over a period of time in the atmosphere oxidise to CO₂, which is included in the total greenhouse gas emissions reported to UNFCCC. As explained in chapter 9, the indirect CO₂ emissions from oxidized CH₄ and NMVOC are calculated from the content of fossil carbon in the compounds.

Solvents and other product use are non-key categories.

4.5.3.2 Methodological issues

The general model used is a simplified version of the detailed methodology described in chapter 6 of the EMEP/CORINAIR Guidebook 2007 (EEA 2007). It represents a mass balance *per substance*, where emissions are calculated by multiplying relevant activity data with an emission factor. For better coverage, point sources reported from industries to the Norwegian Environment Agency and calculated emissions from a side model for cosmetics are added to the estimates. A detailed description of method and activity data is available in Holmengen and Kittilsen (2009).

It is assumed that all products are used the same year as they are registered, and substances are not assumed to accumulate in long-lived products. In other words, it is assumed that all emissions generated by the use of a given product during its lifetime take place in the same year as the product is declared to our data source, the Norwegian Product Register. In sum, this leads to emission estimates that do not fully reflect the actual emissions taking place in a given year. Emissions that in real life are spread out over several years all appear in the emission estimate for the year of registration. However, this systematic overestimation for a given year probably more or less compensates for emissions due to previously accumulated amounts not being included in the estimate figures.

No official definition of solvents exists, and a list of substances to be included in the inventory on NMVOC emissions was thus created. The substance list used in the Swedish NMVOC inventory (Skårman et al. 2006) was used as a basis. This substance list is based on the definition stated in the UNECE Guidelines¹⁴. The list is supplemented by NMVOC reported in the UK's National Atmospheric Emissions Inventory (AEA 2007). The resulting list was comprised by 678 substances. Of these, 355 were found in the Norwegian Product Register for one or more years in the period 2005-2007.

Cosmetics

Cosmetics are not subject to the duty of declaration. The side model is based on a study in 2004, when the Climate and Pollution Agency (now called Norwegian Environment Agency) calculated the consumption of pharmaceuticals and cosmetics (SFT 2005a). The consumption was calculated for product groups such as shaving products, hair dye, body lotions and antiperspirants. The consumption in tonnes each year is calculated by using the relationship between consumption in Norwegian kroner and in tonnes in 2004. Figures on VOC content and emission factors for each product group were taken for the most part from a study in the Netherlands (IVAM 2005), with some supplements from the

¹⁴ "Volatile compound (VOC) shall mean any organic compound having at 293.15 degrees K a vapor pressure of 0.01 kPa or more, or having a corresponding volatility under the particular conditions of use."

previous Norwegian solvent balance (the previous NMVOC emission model).

NMVOC and CO₂

The use of solvents leads to emissions of non-methane volatile organic compounds (NMVOC) which is regarded as an indirect greenhouse gas. The NMVOC emissions will over a period of time in the atmosphere oxidise to CO₂, which is included in the total greenhouse gas emissions reported to UNFCCC.

4.5.3.3 Activity data

The data source is the Norwegian Product Register. Any person placing dangerous chemicals on the Norwegian market for professional or private use has a duty of declaration to the Product Register, and import, export and manufacturing is reported annually. The only exception is when the amount of a given product placed on the market by a given importer/producer is less than 100 kg per year.

The information pertained in the data from the Product Register makes it possible to analyse the activity data on a substance level, distributed over product types (given in UCN codes; Product Register 2007), industrial sectors (following standard industrial classification (NACE; Statistics Norway (2014c)), including private households (no NACE), or a combination of both. As a consequence, the identification of specific substances, products or industrial sectors that have a major influence on the emissions is greatly facilitated.

Cosmetics

The side model for cosmetics is updated each year with data on from the Norwegian Association of Cosmetics, Toiletries and Fragrance Suppliers (KLF).

Point sources

Data from nine point sources provided by the Norwegian Environment Agency is added to the emissions estimates. The point sources are reported from the industrial sector "Manufacture of chemicals and chemical products" (NACE 24). In order to avoid double counting, NMVOC used as raw materials in this sector are excluded from the emission estimates from the Product Register data.

4.5.3.4 Emission factors

Emission factors are specific for combinations of product type and industrial sector. Emission factors are gathered from the Swedish model for estimating NMVOC emissions from solvent and other product use (Skårman et al. 2006). The emission factors take into account different application techniques, abating measures and alternative pathways of release (e.g. waste or water). These country-specific emission factors apply to 12 different industries or activities that correspond to subdivisions of the four major emission source categories for solvents used in international reporting of air pollution (EEA 2007).

It is assumed that the factors developed for Sweden are representative for Norwegian conditions, as we at present have no reasons to believe that product types, patterns of use or abatement measures differ significantly between the two countries. Some adjustments in the Swedish emission factors were made when the model was first developed by Holmengen and Kittilsen (2009) and several improvements of single emissions factors have been made in the following years.

In accordance with the Swedish model, emission factors were set to zero for a few products that are assumed to be completely converted through combustion processes, such as EP-additives soldering

agents and welding auxiliaries. Quantities that have not been registered to industrial sector or product type are given emission factor 0.95 (maximum). Emission factors may change over time, and such changes may be included in this model. However, all emission factors are at the moment constant for all years.

4.5.3.5 Uncertainties and time-series consistency

Uncertainty in emission factors

The emission factors are more detailed in the new NMVOC model than in the previous model, as this model can take into account that emissions are different in different sectors and products, even when the substance is the same. However, for this to be correct, a thorough evaluation of each area of use is desirable, but not possible within a limited time frame. Thus, the emission factor is set with general evaluations, which leads to uncertainty.

The emission factors are gathered from several different sources, with different level of accuracy. The uncertainties in emission factors depend on how detailed assessment has been undertaken when the emission factor was established. Some emission factors are assumed to be unbiased, while others are set close to the expected maximum of the range of probable emission factors. This, together with the fact that the parameter range is limited, gives us a non-symmetrical confidence interval around some of the emission factors. For each emission factor we thus have two uncertainties; one negative (n) and one positive (p). These are aggregated separately, and the aggregated uncertainty is thus not necessarily symmetrical.

Uncertainty in activity data

For the activity data, the simplified declarations and the negative figures due to exports lead to known overestimations, for which the uncertainty to a large extent is known. A more elaborate problem in calculations of uncertainty is estimating the level of omissions in declaration for products where the duty of declaration does apply. In addition, while declarations with large, incorrect consumption figures are routinely identified during the QA/QC procedure, faulty declarations with small consumption figures will only occasionally be discovered. There is however no reason to believe that the Product Register data are more uncertain than the data source used in the previous model (statistics on production and external trade), as similar QA/QC routines are used for these statistics.

The errors in activity data are not directly quantifiable. Any under-coverage in the Product Register is not taken into account. The activity data from the Swedish Product register has an uncertainty of about 15 % (Skårman et al. 2006). The Norwegian Product Register is assumed to be comparable to the Swedish, and thus the uncertainty in the activity data is assumed to be 15 %. For some products, simplified declarations give an indication of maximum and minimum possible amounts. In these cases, the maximum amount is used, and the positive uncertainty is set to 15 % as for other activity data, while the negative uncertainty is assumed to be the interval between maximum and minimum amount. All activity data are set to zero if negative.

A detailed description of the uncertainty analysis is available in Holmengen and Kittilsen (2009). The variance of total emission was estimated from the variance estimates obtained for emission factors and activity data, using standard formulas for the variance of a sum and the variance of a product of independent random variables. The aggregated uncertainties in level and trend are given in Table 4.38 and Table 4.39.

Table 4.38 Uncertainty estimates for level in NMVOC emissions, 2005-2007. Tonnes and %.

Uncertainty in level	Negative (n)	Negative (n) (% of total emissions)	Positive (p)	Positive (p) (% of total emissions)
2005	2 288	4.58	1 437	2.88
2006	1 651	3.70	1 103	2.47
2007	1 299	2.79	1 168	2.51

Table 4.39 Uncertainty estimates for trend in NMVOC emissions, 2005-2007. Tonnes.

Uncertainty in trend	Negative (n)	Positive (p)	95% confidence interval for change
2005-2006	2 135	1 067	(-7 366, -4 164)
2006-2007	1 420	947	(407, 2 774)
2005-2007	1 882	1 076	(-5 286, -2 328)

Time series consistency

The activity data from the Norwegian Product Register is only available from 2005 onwards. For the years from 1990 to 2000, data from the previous solvent balance has been used. The two time series have been spliced by interpolation. This introduces a degree of time series inconsistency. However, the results from the previous solvent balance were evaluated and updated with new knowledge from the current model in Holmengen and Kittilsen (2009). Thus, overall time series consistency is deemed to be satisfactory.

4.5.3.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V. Large between-year discrepancies in the time series of substance quantities are routinely identified and investigated, in order to correct errors in consumption figures. Large within-year discrepancies between minimum and maximum quantities in simplified declarations are routinely identified and investigated, in order to prevent overestimation for substances where consumption figures are given in intervals. Large within-year discrepancies between totals for industrial sectors (NACE) and totals for products (UCN) are routinely identified and investigated, in order to detect erroneous or incomplete industrial sectoral and product type distribution.

4.5.3.7 Category-specific recalculations

Correction of activity data. The activity data for 2016 in the previous submission was based on figures from 2015. In this submission the activity data for 2016 has been updated with new information from the Norwegian Product Register. The correction resulted in a reduced indirect CO₂ emissions from NMVOC by 3.2 %.

4.5.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.5.4 Road paving with asphalt, 2D3b

4.5.4.1 Category description

Indirect CO₂ emissions from NMVOC emissions from road paving with asphalt are included in the inventory.

4.5.4.2 Methodological issues

The emissions from road paving are calculated in accordance with a Tier 1 approach (EEA 2013).

$$E_{\text{pollutant}} = AR_{\text{production}} * EF_{\text{pollutant}}$$

where

E pollutant = the emission of the specified pollutant

AR production = the activity rate for the road paving with asphalt

EF pollutant = the emission factor for this pollutant

4.5.4.3 Activity data

The activity data used is the annual weight of asphalt used for road paving in Norway, collected by the Contractors Association - Building and Construction annually (EBA 2014).

4.5.4.4 Emission factors

The share of bitumen in the asphalt is set to be 0,05 for all years, based on information from a road technology Institute, a centre for research and development, quality control and documentation of asphalt (<http://www.asfaltteknisk.no/>). The emissions of NMVOC are calculated using an emission factor of 16 g NMVOC / tonne asphalt (EEA 2013).

4.5.4.5 Uncertainties and time series consistency

The activity data and emission factor used are uncertain. The annual emissions are however low. Activity data on asphalt used are available from 1995 onwards. For the years 1990-1994, the emission figure for 1995 is used. This introduces some degree of time series inconsistency in methodology. The annual variability in emissions throughout the entire time series is however insignificant, and this inconsistency is thus deemed acceptable.

4.5.4.6 Category-specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

4.5.4.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.


4.5.4.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.5.5 Other, 2D3d (use of urea as a catalyst)

4.5.5.1 Category description

Urea is used as a catalyst to reduce NO_x emissions, in Norway primarily from road transport and shipping. When urea is injected upstream of a hydrolysis catalyst in the exhaust line, the following reaction takes place:



The ammonia formed by this reaction is the primary agent that reacts with nitrogen oxides to reduce them to nitrogen.

There were no emissions from the use of urea as a catalyst in 1990, and the use of urea and thus emissions have increased significantly the last few years.

4.5.5.2 Methodological issues

Emissions are calculated based on equation 3.2.2 of Volume 2 of the 2006 IPCC Guidelines:

$$\text{Emissions} = \text{Activity} * 12/60 * \text{Purity} * 44/12$$

where

Emissions = CO₂ emissions from urea-based additive in catalytic converters (Gg CO₂)

Activity = amount of urea-based additive consumed for use in catalytic converters

Purity = the mass fraction (= fraction of urea in the urea-based additive)

The fraction 12/60 converts the emission figure from urea (CO(NH₂)₂) to carbon (C), while 44/12 converts C to CO₂.

Emissions are calculated as the sum of emissions from each purity.

4.5.5.3 Activity data

No official statistics cover sale, production, or use of urea as a catalyst in Norway. There is no national production of urea used as a catalyst, as the urea produced in Norway is used for fertilizers only. There are many importers of urea used as a catalyst, and the urea is often imported in smaller containers, and not in bulk. Information from the largest importer of urea shows that urea is imported to Norway in at least three different purities: 32.5 % for use in road transport, 40 % for use in shipping, and 100 % for dilution before use. The statistics on external trade does not have a clear split on urea used for fertilizers and urea used as catalyst, nor does it split on different purities.

Based on these considerations, import data from the largest producer together with estimates of marked shares have been used to calculate the total consumption of urea used as a catalyst each year. The first year of activity is considered to be 2008, as very few vehicles had the technology prior to this year.

4.5.5.4 Emission factors

There are no emission factors used for this calculation. All carbon in the urea used is converted to CO₂.

4.5.5.5 Uncertainties and time series consistency

There are no emission factors as such in these calculations, and the purity of the different solutions are deemed to be reliable. However, the calculations are based on activity data where expert judgement is an important parameter, and there is a certain degree of uncertainty.

The same source of activity data and the same parameters have been used for all years, and the time series consistency is thus deemed to be satisfactory.

4.5.5.6 Category-specific QA/QC and verification

In the development of the emission estimates, activity data used (import data from the largest importer) were compared with import data from the statistics on external trade.

4.5.5.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.5.5.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.6 Electronics industry – 2E

Norway reports the source category integrated circuit or semiconductor under the category 2E, see Table 4.40 and Table 4.41.

Table 4.40 Electronics industry. Components emitted and included in the Norwegian inventory.

Source category	SF ₆	HFCs	PFCs	NF ₃	Tier	Key category
2E1. Integrated circuit or semiconductor	E	NO	NO	NO	1	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

Table 4.41 Emission trends for 2E Electronics industry (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2E1. Integrated circuit or semiconductor	0.0	0.0	1.1	1.1	0.0 %	NA	0.0

Source: Statistics Norway and Norwegian Environment Agency

4.6.1 Integrated circuit or semiconductor, 2E1

4.6.1.1 Category description

There are SF₆ emissions from the use in the manufacturing of semiconductors. There were no emissions from the production of integrated circuit or semiconductors in 1990, but the emissions in 2017 were 1 140 tonnes of CO₂-equivalents, see Table 4.41.

4.6.1.2 Methodological issues

The method is described in a report from SFT (1999c) and there have been emissions of SF₆ from this source since 1995. Data on sales to semiconductor manufacturers were collected for 1998, and total sales amounted to 90 kg. The report projected that sales would increase to 100 kg, but would then remain in that range in the next decade. No new data have been collected, and the projection from the 1999 report has been prolonged.

4.6.1.3 Activity data

The report from 1999 assumed that 50% of the gas reacts in the etching process and the remaining 50% are emitted. Hence 45 kg are reported as emissions until 1998 and 50 kg from 1999 onwards.

4.6.1.4 Emission factors

The leakage rate for the production of semiconductors is shown in Table 4.42.

Table 4.42 Yearly rate of leakage of SF₆ from the production of semiconductors.

Emission source	Leakage rate (% of input of SF ₆)
Production of semiconductors	50

Source: SFT (1999c)

4.6.1.5 Uncertainties and time series consistency

An uncertainty estimate is given in Annex II.

A general assessment of the time series consistency has not revealed any time series inconsistencies in the emission estimates for this source category.

4.6.1.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V. Since the emissions have been assumed to be constant since 1999, there is no specific QA/QC procedure for this source category.

4.6.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.6.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.7 Product uses as substitutes for ODS – 2F (key category for HFCs)

Norway reports the source category HFCs and PFCs from refrigeration and air conditioning and other applications under the category 2F. See Table 4.43 and Table 4.44 for details.

Table 4.43 Product uses as substitutes for ODS. Components included in the inventory, tier of method and key category.

Source category	HFCs	PFCs	SF ₆	NF ₃	Tier	Key category
2F1-2F6. Refrigeration and air conditioning, foam blowing agents, fire protection, aerosols, solvents, other applications.	E	E	NO	NO	*	Yes**

*Mainly estimated using Tier 2a (emissions calculated at a disaggregated level, emission factor approach). Exceptions are mobile air conditioning that is estimated using Tier 2b (b=mass balance approach) and fire protection, aerosols and solvents that are estimated using Tier 1a (emissions calculated at an aggregated level, emission factor approach).

**In the key category analysis, 2F1 and 2F6 have been aggregated.

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

Table 4.44 Emission trends for 2F Product uses as substitutes for ODS (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2F1-2F6. Refrigeration and air conditioning, foam blowing agents, fire protection, aerosols, solvents, other applications.	0.0	0.0	1 363.6	1 402.8	16.3 %	3 195 305.3 %	2.9 %

Source: Statistics Norway and Norwegian Environment Agency

Table 4.44 shows the emission trends for 2F as a whole. The GHG emissions from this sector category were a little less than 1.4 million tonnes in 2017, this is 16.3 % of the total emission from the IPPU-sector. The emissions were 44 tonnes CO₂-equivalents in 1990 and have increased substantially over the years. The emissions increased by 2.9 % from 2016 to 2017. The majority of the emissions are reported in 2F1 and these include minor emissions of PFC-218 in the years 2010-2014.

HFCs and PFCs are mainly used as substitutes for ozone depleting substances (CFCs and HCFCs) that are being phased out according to the Montreal Protocol. They are used in varied applications, including refrigeration and air conditioning equipment, as well as in foam blowing, fire extinguishers, aerosol propellants and analysing purposes. There is no production of HFCs and PFCs in Norway. However, PFCs are emitted as a by-product during the production of aluminium. Due to, i.e., high taxation, the use of PFCs in product-applications has been very low. PFC-218 has been used as a commercial cooling agent.

The amounts of imported and exported gases are found in registers from the Norwegian Directorate of Customs and Excise. All import of F-gases is covered in these registers, as Norway lays a tax on the import of F-gases (Ministry of Finance 2014). In January 2003 a tax on import and production of HFC and PFC was introduced. In July 2004 this tax was supplemented with a refund for the destruction of

used gas. From 1st of January 2014, the tax increased by about 100 NOK to NOK 330 (approximately EUR 40) per tonne CO₂ equivalents of gas imported as of 2014. In May 2010, EU regulation (EC) No 842/2006 on certain fluorinated greenhouse gases was included in Norwegian legislation.

Also practically all export of F-gases is covered, as commodities with F-gases have their own commodity code (HS-code). The registered export of F-gases from Norway is very low, and any underestimation of the export of F-gases would thus be very slight and eventually lead to over-estimation (and not under-estimation) of the emissions.

The imported and exported gases are allocated to sectors based on commodity codes and information identifying each company. In some cases (sector 2F1) the type of gas is used as additional information. Uncertainties in the distribution by sector do not affect the total amount of F-gases to be emitted over time, as the emissions over time are determined by the total amount of F-gases to be distributed. Thus, under-estimation in one sector would eventually lead to an equivalent over-estimation in another sector at some point of time.

The HFC emissions from 2F1 and 2F6 is defined as a key category according to the approach 2 analysis.

4.7.1 Refrigeration and air conditioning, 2F1

4.7.1.1 Category description

HFCs and PFCs are mainly used as substitutes for ozone depleting substances (CFCs and HCFCs) that are being phased out according to the Montreal Protocol. Emissions from refrigeration and air conditioning equipment are reported under this source category.

4.7.1.2 Methodological issues

Actual emissions of HFCs and PFCs are calculated using the Tier 2 methodology. This methodology takes into account the time lag in emissions from long lived sources, such as refrigerators and air-conditioning equipment. The chemicals slowly leak out from seams and ruptures during the lifetime of the equipment. The leakage rate, or emission factor, varies considerably depending on type of equipment and its maintenance.

An emission factor approach is used to estimate emissions from all categories except mobile air conditioning. For mobile air conditioning a hybrid of mass-balance and emission factor approach is used. There is no production of new cars in Norway; hence it is assumed that HFCs imported in bulk for use in mobile air conditioning is used to refill the systems after leakage. This is the mass-balance approach, and it leads to very high product life factors (up to 100 %) in the beginning of the time series because no bank of chemicals was yet accumulated. A restriction is however set in the model: Where the imported bulk is lower than 10 % of the bank, the emission factor approach is applied. This model assumption means that the product life factor will never be lower than 10 (using an emission factor of 10 %) and is occurring towards the end of the time series.

In the CRF, there are reported amounts of HFC-143 in 2005 and 2006 and of HFC-134 in 2004 and 2008 filled into new manufactured products in commercial refrigeration. According to an expert on refrigeration and HFCs, these two gases are not used regularly in Norway but is imported to be used in equipment testing. For other years, NO is therefore considered to be the appropriate notation key for these two gases for amounts filled into new manufactured products. This is confirmed by our

data, and any new import of these two gases will be reflected in the registers from the Norwegian Directorate of Customs and Excise as both gases are covered by the tax on HFCs.

The reported stock emissions for perfluoropropane (C₃F₈) consumed in commercial refrigeration in CRF table 2(II).B-H have a declining trend. The use of PFCs is being phased out and replaced by other gases and only small amounts of PFCs have been imported in bulk in the last few years. About the same amount of PFCs that are reported as imported are reported as collected for destruction, so there has not been a build-up of stock that would generate emissions (i.e. the entire amount of gas remained in stock from the previous year is assumed to be collected for destruction). Since 2014, the AD for annual average stock and stock emissions for C₃F₈ are therefore reported as “NO”.

4.7.1.3 Activity data

There is no production of HFC or PFC in Norway. Hence all emissions of these chemicals originate from chemicals imported in *bulk* or in products. The methodology requires that annual imported amounts of each chemical are obtained by source category. Various data sources are used:

Amounts of chemicals imported in bulk were up to 2009 obtained from the Norwegian Climate and Pollution Agency (now Norwegian Environment Agency). After 2009, bulk data are collected from the Norwegian Directorate of Customs and Excise. Time series for imported and exported amounts of chemicals in *products* are based on collected data for some years and data prior to and between these years are estimated. For the years 1995-1997 data were collected through a survey performed in 1999 (SFT 1999b). Data on imports were collected from customs statistics for the years 2005-2006 and 2010-2012. After 2011 the import data has been collected annually.

Amounts of chemicals destructed after collection from retired equipment are annually reported to Statistics Norway from the company in charge of the collection. A more thorough description of the activity data is available in Bjønnes (2013). A provisional distribution of chemicals by application category was used for 2012, based on the 2011 distribution. The totals per gas, however, were collected from the Norwegian Directorate of Customs and Excise.

4.7.1.4 Emission factors

Leakage rates and product lifetimes used in the calculations are shown in Table 4.45.

Table 4.45 Emission factors¹ for HFCs and PFCs from 2F1 Refrigeration and Air conditioning.

Source category	Lifetime (years)	Production/initial emission (% of initial charge)	Lifetime emission (% of initial charge/year)
2.F.1.a. Commercial Refrigeration			
Stand-alone Commercial	10	NO	3.5
Medium and Large Commercial	15	2	10
2.F.1.b Domestic Refrigeration	15	NO	0.5
2.F.1.c Industrial Refrigeration	15	2	10
2.F.1.c. Transport Refrigeration	9	1	20
2.F.1.e Mobile Air-Conditioning	12	NO	10
2.F.1.f Stationary Air-Conditioning	15	1	4

¹IPCC (2006), IPCC (1997b)

It is important to note that subapplication 2.F.1.a, Commercial refrigeration, is calculated at a more detailed level. Two groups of equipment that differs substantially in their life cycle and emission patterns, and hence emission factors, are taken into account:

- Stand-alone commercial applications includes equipment like vending machines and moveable refrigerators and freezers typically used for keeping beverages and ice cream cold in supermarkets, office buildings, schools etc. There is currently no production of this kind of equipment in Norway. All emissions take place during the operating phase (emissions from stocks/lifetime emissions) or at decommissioning. The IPCC 2006 Guidelines recommends an operation emission factor between 1 and 15 % for this application category, and between 0.1 and 0.5 % for domestic refrigeration. Because the units imported to Norway are small, sealed units and thus similar to the refrigerators and freezers for domestic use, an emission factor in the lower end of IPCCs recommendation is believed to best reflect the actual emissions.
- Medium and large commercial refrigeration equipment is normally built and filled with fluorinated substances on site. They will thus have emissions both in the production phase and from operation/use the subsequent years. The IPCC 2006 Guidelines recommends an operation emission factor between 10 and 35 % for this application category. The lower emission factor is used in the Norwegian calculations. The reasoning behind this is that the tax on imports of fluorinated substances is assumed to result in a high level of maintenance of the equipment and low leakage rates.

This means that the implied emission factor named “Product life factor” as calculated in the CRF, will vary for this group as the share of stock for the two groups of equipment are not constant over time. In order to provide better transparency, Table 4.46 provides information on the relative share of stock for the two categories, aggregated for all substances in CO₂ equivalents. For most fluorinated gases, the majority of stock is comprised by medium and large equipment, hence the product life factor is close to 10. Important exceptions are HFC-32 and HFC-134a.

Table 4.46 Relative share of emissions from imported and domestically filled commercial refrigeration applications.

Year	Share imported	Share domestically filled
1990	0.0	100.0
1995	0.5	99.5
2000	0.6	99.4
2005	0.8	99.2
2008	1.7	98.3
2009	2.3	97.7
2010	2.9	97.1
2011	3.8	96.2
2012	5.0	95.0
2013	6.6	93.4
2014	8.9	91.1
2015	9.1	90.9
2016	9.6	90.4
2017	11.2	88.8

Source: Statistics Norway

4.7.1.5 Uncertainties and time series consistency

The uncertainties of the different components of the national greenhouse gas inventory have been evaluated in detail in 2006 by Statistics Norway (See annex II). Both the leakage rate (emission factor) and the stored amount of chemicals (activity data) are considered quite uncertain. The total uncertainties for the emission estimates by the consumption of halocarbons are estimated to be ± 50 % for both HFC and PFC.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.7.1.6 Category-specific QA/QC and verification

In addition to the general QA/QC procedures described in Annex V, the activity data, emissions figures and the model used to estimate emissions are checked by several manual and automatic controls performed during and after the production of the figures.

Firstly, the activity data on imports and exports of chemicals undergo automatic checks before they are used as input for the emission calculations. Double counting and missing values are flagged and checked at a detailed level, i.e. for each observation of amount of chemical by importer/exporter and goods.

Aggregated figures for amount of chemical imported/exported for each year are then checked manually as they are entered into excel sheets containing time series for each CRF equipment/emission source. Potential errors like missing values or major differences in figures between years are checked and corrected if necessary. Aggregated figures on imports per gas are also compared with tables from the Norwegian Directorate of Customs and Excise, in order to check that our computerized coding has not altered the total amounts of import per gas. These controls would probably benefit from being automatic in the future.

Both activity data and emissions per chemical and type of equipment/emission source undergo controls in the excel sheets where the emissions calculations take place: Potential emissions are compared with actual emissions and IEFs are calculated and checked. Currently, these controls are only applied to the most important emission sources. The controls would probably also benefit from being automatic in the future.

The estimated emissions are finally subjected to several controls in order to identify errors in activity data and/or the calculation model. This includes the flagging of:

- Emissions from combinations of industrial sector and type of chemical that have not occurred previous years
- Emissions from combinations of industrial sector and type of chemical that occurred the previous year, are not occurring this year
- Large or small emissions compared with the previous years:
 - By type of chemical and sector
 - By type of chemical and CRF/NFR emission source
 - Recalculations

4.7.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.7.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.7.2 Other applications, 2F6

4.7.2.1 Category description

Due to confidentiality restrictions, Norwegian emissions from categories 2.F.2 (foam blowing), 2.F.3 (fire extinguishers), 2.F.4 (aerosols/metered dose inhalers (MDI)) and 2.F.5 (solvents) are reported in the CRF tables using the notation key "IE" and aggregated under 2.F.6 (Other applications using ODS substitutes) and not disaggregated by substance. Note however, that the calculations are made for each subsector.

In response to a recommendation from the technical expert review team, more transparent information on the uses and the levels of emissions per capita compared to other Parties was included in the NIR. This information is shown below.

More than 95 % of the Norwegian emissions reported in 2F6 since 1995, in terms of CO₂-equivalents¹⁵, were from:

- i. Foam blowing agents (2.F.2), i.e. emissions of HFC-134a and HFC-152a from the use of hard foam/ closed cells-products.
 - i. For HFC-134a the per capita emissions were in the range of 0-1.9 kg CO₂-eq before 1998 and 2.0-3.9 kg CO₂-eq in the period 1998 to 2012. Per capita emissions in comparable countries were in the range of 0-11.71 kg CO₂-eq in 2012.
 - ii. For HFC-152a the per capita emissions were in the range of 0-1.74 kg CO₂-eq in the period 1990-2012. Per capita emissions in comparable countries were in the range of 0-1.74 kg CO₂-eq in 2012.
- ii. Aerosol (2.F.4), i.e. emissions from the use of HFC-134a in metered dose inhalers (2.F.4.a). The per capita emissions have grown from 0-1.9 kg CO₂-eq per capita before 2011, to 2.0-3.9 kg CO₂-eq per capita in 2011 and 4.0 to 5.9 kg CO₂-eq per capita in 2012. Per capita emissions in comparable countries were in the range of 0.24-12.91 kg CO₂-eq in 2012.
- iii. Fire extinguishers (2.F.3), both in use and in the waste phase, of the gases HFC-125, HFC-134a and HFC-227ea). The emissions have increased from 0-1.9 kg CO₂-eq per capita before 2011, to 2.0-3.9 kg CO₂-eq per capita in 2011 and 2012. Comparable countries had emissions in the range of 0.59-6.78 kg CO₂-eq per capita in 2012.

As can be seen from the list above, the Norwegian per capita emission for each of these three sectors in 2012 was well within the range of the selected comparable countries (Austria, Denmark, Finland, Ireland, Sweden, United Kingdom and United States). For the other categories included in the aggregated 2.F.6 amount, the emitted amounts were zero or close to zero. This explains the difference from the other comparable countries in the overall 2F2 to 2F6 amount. The increase in the

¹⁵ Note that the reported emissions in sector 2F6 are given in CO₂-eq.

reported Norwegian aggregated 2F6 emission since 2009 is due to 2F4 (metered dose inhalers, HFC-134a, from stocks).

4.7.2.2 Methodological issues

See description for source category 2F1.

4.7.2.3 Activity data

See description for source category 2F1.

4.7.2.4 Emission factors

Leakage rates and product lifetimes used in the calculations are shown in Table 4.47.

Table 4.47 Emission factors¹ for HFCs from products and lifetime of products.

Source category	Lifetime (years)	Production/initial emission (% of initial charge)	Lifetime emission (% of initial charge/year)
2.F.2 Foam			
2.F.2a Closed cells	20	5	4.5
2.F.2b Open cells	NO	NO	NO
2.F.3 Fire protection	15	2	5
2.F.4 Aerosols			
2.F.4.a Metered Dose Inhalers	2	NO	50
2.F.4.b Other aerosols	2	NO	50
2.F.5 Solvents	2	NO	50

¹IPCC (2006), IPCC (1997b)

4.7.2.5 Uncertainties and time series consistency

See description for source category 2F1.

4.7.2.6 Category-specific QA/QC and verification

See description for source category 2F1.

4.7.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.7.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.8 Other product manufacture and use – 2G

Norway reports the source categories electric equipment, SF₆ and PFCs from other product use, medical applications, propellant for pressure and aerosol cans and other use of N₂O under the category 2G, see Table 4.48.

Table 4.48 Other product manufacture and use. Components included in the inventory, tier of method and key category.

Source category	HFCs	PFCs	SF ₆	NF ₃	N ₂ O	Tier	Key category
2G1. Electric equipment	NO	NO	E	NO	NA	Tier 1	No
2G2. SF ₆ and PFCs from other	NA	NO	E	NO	NA	Tier 1	No
2G3a. Use of N ₂ O in anaesthesia	NA	NA	NA	NA	E	Tier 1	No
2G3b.1. Propellant for pressure	NA	NA	NA	NA	E	Tier 1	No
2G3b.2. Other use of N ₂ O	NA	NA	NA	NA	E	Tier 1	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

Table 4.49 shows the emission trends for 2G as a whole and for the various sub-categories. The GHG emissions from this sector category were about 80 kt in 2017, this is 0.9 % of the total emission from the IPPU-sector. The emissions from this sector decreased by 9.7 % from 1990 and the emissions decreased by 6.1 % from 2016 to 2017.

Table 4.49 Emission trends for 2G Other product manufacture and use (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2G1. Electric equipment	51.1	0.4 %	34.8	26.9	0.3 %	-47.5 %	-22.9 %
2G2. SF ₆ and PFCs from other product use	2.2	0.0 %	27.7	30.8	0.4 %	1 281.0 %	11.5 %
2G3a. Use of N ₂ O in anaesthesia	34.2	0.2 %	19.7	19.3	0.2 %	-43.4 %	-1.5 %
2G3b.1. Propellant for pressure and aerosol cans	0.0	0.0 %	1.6	1.6	0.0 %	NA	0.0 %
2G3b.2. Other use of N ₂ O	0.0	0.0 %	0.4	0.4	0.0 %	NA	0.0 %
2G. Total	87.5	0.6 %	84.1	79.0	0.9 %	-9.7 %	-6.1 %

Source: Statistics Norway and Norwegian Environment Agency

As part of the transformation to new reporting guidelines, Norway has examined whether there are activities that would result in emissions of trinitrogenfluoride (NF₃). Our assessment is that there are no emissions of NF₃ in Norway.

4.8.1 Electric equipment, 2G1

4.8.1.1 Category description

SF₆ is used as an insulation medium in high tension electrical equipment including gas insulated switchgear (GIS) and circuit breakers. There is no production of SF₆ in Norway. In March 2002 a voluntary agreement was signed between the Ministry of Environment and the most important users

and producers of GIS. According to this agreement emission from this sector should be reduced by 13 % in 2005 and 30 % in 2010 with 2000 as base year. For the following up of this agreement, the users (electricity plants and –distributors) and producer (one factory) report annually to the government. This voluntary agreement terminated successfully in 2010, but a continuation is being discussed. Although the voluntary agreement has terminated, the users still report annually to the government.

4.8.1.2 Methodological issues

The general methodology for estimating SF₆ emissions was revised in a SFT report (SFT 1999c), while the sector specific methodology for GIS has been revised in the 2010 reporting based on new information from the agreement.

Emissions from production of GIS (one factory) were included for the first time in 2003. The company has, as part of the voluntary agreement with the Ministry of the Environment, made detailed emission estimates back to 1985. These emissions constitute a significant part of national emissions of SF₆. In recent years emissions rates have been considerably reduced due to new investments and better routines. The company now performs detailed emission calculations based on accounting of the SF₆ use throughout the whole production chain.

Emissions from a small number of GIS users that are not part of the agreement are calculated with emission factors from Table 4.50. SF₆ emissions from manufacturing are included in emissions from stocks due to confidentiality.

4.8.1.3 Activity data

Data is collected from companies that use SF₆ in various processes. The calculations take into account imports, exports, recycling, accumulation in bank, technical lifetimes of products, and different rates of leakage from processes, products and production processes. From 2003 onwards emission estimates reported directly from users and producers, according to the voluntary agreement, are important input.

4.8.1.4 Emission factors

Leakage rates and product lifetimes used in the calculations are shown in Table 4.50.

Table 4.50 Product lifetimes and leakage rates from products containing SF₆.

Product emission source	Yearly rate of leakage	Product lifetime (years)
Sealed medium voltage switchgear	0.1	30
Electrical transformers for measurements	1	30

Source: SFT (1999c)

4.8.1.5 Uncertainties and time series consistency

An uncertainty estimate is given in Annex II. The uncertainty of 60 % is an expert judgement (Rypdal & Zhang 2000).

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.8.1.6 Category-specific QA/QC and verification

The current methodology was established in the SFT report (SFT 1999c), with emissions from GIS calculated from stock data estimates and leakage factors. It was revised in 2004 when data from the voluntary agreement on GIS became available, with emissions estimated from reported data on refilling (Hansen 2007).

4.8.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.8.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.8.2 SF₆ and PFC from other product use, 2G2

4.8.2.1 Category description

This source category includes SF₆ emissions from other product use.

4.8.2.2 Methodological issues

The method for other sources is described in a SFT report (SFT 1999c). For tracer gas, medical use, and other minor uses, the activity data are annual consumption as estimated in the SFT report. However, for tracer gas some major research projects expired in 2001 and 2006, respectively, and the consumption has been reduced. For sound-insulating windows and footwear, the emissions are calculated from estimated stock of SF₆ in the products, and from production of windows. Footwear with SF₆ was imported, and the use ended in 2001. There was no production of sound-insulating windows from 2008.

4.8.2.3 Activity data

Data is collected from direct consultations with importers and exporters of bulk chemicals and products containing SF₆. The activity data are annual additions of SF₆ to the product stock, as estimated by SFT (1999c). The calculations take into account imports, exports, recycling, accumulation in bank, technical lifetimes of products, and different rates of leakage from processes, products and production processes.

4.8.2.4 Emission factors

Leakage rates and product lifetimes used in the calculations are shown in Table 4.51 and Table 4.52.

Table 4.51 Yearly rate of leakage of SF₆ from different processes.

Emission source	Leakage rate (% of input of SF ₆)
Secondary magnesium foundries	100
Tracer gas in the offshore sector	0
Tracer gas in scientific experiments	100
Medical use (retinal surgery)	100
Production of sound-insulating windows	2 ¹
Other minor sources	100

¹ 1 % after 2002

Source: SFT (1999c)

Table 4.52 Product lifetimes and leakage rates from products containing SF₆.

Product emission source	Yearly rate of leakage	Product lifetime (years)
Sound-insulating windows	1	30
Footwear (trainers)	25	9
Other minor sources

Source: SFT (1999c)

4.8.2.5 Uncertainties and time series consistency

An uncertainty estimate is given in Annex II. The uncertainty of 60% is an expert judgement (Rypdal & Zhang 2000).

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.8.2.6 Category-specific QA/QC and verification

The current methodology was established in a SFT report (SFT 1999c), with emissions from GIS calculated from stock data estimates and leakage factors. It was revised in 2004 when data from the voluntary agreement on GIS became available, with emissions estimated from reported data on refilling (Hansen 2007).

4.8.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.8.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.8.3 Use of N₂O in anaesthesia, 2G3a

4.8.3.1 Category description

N₂O is used in anaesthesia procedures in hospitals, by dentists and by veterinarians.

4.8.3.2 Methodological issues

N₂O is used in anaesthesia procedures and will lead to emissions of N₂O. For the year 1998 and annually from 2000, the emissions are given by data on sales of N₂O for medical uses from the three major producers and importers in Norway. The data include N₂O used as anaesthesia in hospitals, by dentist and by veterinarians. For the year 1999, sales figures have been interpolated between 1990 and 2000. For the years prior to 1998, annual consumption is estimated on basis of sales figures for 1998 and the number of births and number of bednights in hospitals for each year to estimate consumption. For the years 1990-1998, no N₂O is assumed used by dentists and veterinarians as the amounts they used in 2000 were very small.

4.8.3.3 Activity data

For this source actual sale of N₂O is used for the year 1998, and annually from 2000. For the calculations of use prior to 1998, annual number of births and bednights in hospitals are taken from the Statistical yearbook of Norway.

4.8.3.4 Emission factors

The figures are based on sales of N₂O.

4.8.3.5 Uncertainties and time-series consistency

The figures are uncertain. There may be small importers not included in Statistics Norway's telephone survey with 2000 and the investigation done by the Norwegian Environment Agency in 2014, but the emissions are small, so it is believed that the uncertainty is at an acceptable level.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.8.3.6 Category-specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

4.8.3.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.8.3.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.8.4 Propellant for pressure and aerosol products, 2G3b.1.

4.8.4.1 Category description

N₂O is used as a propellant in spray boxes and this use will lead to emissions of N₂O. It is also used in research work, for instance in the food industry and at universities. There is no production of N₂O for these purposes in Norway.

4.8.4.2 Methodological issues

Information on sale volumes has been reported by the plants to Statistics Norway. It is assumed that all propellant is released to air.

4.8.4.3 Activity data

Information has been gathered from the plants indicating that there is no production or sale of N₂O for use as a propellant in Norway. The N₂O is already in the spray cans when imported. There was no import of these spray cans prior to 1993. For the years 1994-2002 the number of cans imported in 1994 have been used as activity data, while the number of cans imported in 2003 has been used as activity data for all years since.

4.8.4.4 Emission factors

Not relevant.

4.8.4.5 Uncertainty and time-series consistency

The figures for one year are used for all years. It is believed that all figures from all major importers are included in the inventory.

A general assessment of time series consistency has not revealed any time series inconsistencies in the

emission estimates for this category.

4.8.4.6 Category-specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

4.8.4.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.8.4.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.8.5 Other use of N₂O, 2G3b.2

4.8.5.1 Category description

Small amounts of N₂O are used for research work and for drag-racing.

There were no emissions of N₂O from use in research and for drag racing in 1990. The use has been estimated to 407 tonnes CO₂ equivalents from the year 1993 and onwards.

4.8.5.2 Methodological issues

Data on imported amounts in 2002 has been used for all years and it is assumed that all propellant is released to air.

4.8.5.3 Activity data

Data on imported amounts in 2002 has been used for all years.

4.8.5.4 Emission factors

Not relevant.

4.8.5.5 Uncertainty and time-series consistency

The figures for one year are used for all years. A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.8.5.6 Category-specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Annex V for the description of the general QA/QC procedure.

4.8.5.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.8.5.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.9 Other – 2H

Under Other production, Norway reports the two source categories pulp and paper and food and beverages industry, see Table 4.53.

Table 4.53 Other production. Components included in the inventory, tier of method and key category.

Source category	CO ₂	NM VOC	Tier	Key category
2H1. Pulp and paper	R	NA	Tier 2	No
2H2. Food and beverages industry	R	E	Tier 2	No

R = emission figures in the national emission inventory are based on figures reported by the plants. E = emission figures are estimated. NA = Not Applicable. NO = Not Occurring. IE = Included Elsewhere.

Table 4.54 shows the emission trends for 2H as a whole and for the various sub-categories. The GHG emissions from this sector category were 114 kt in 2017, this is 1.3 % of the total emission from the IPPU-sector. The emissions from this sector increased by 263.3 % from 1990 and the emissions decreased by 1.7 % from 2016 to 2017.

Table 4.54 Emission trends for 2H Other (kt CO₂ equivalents).

Source category	1990	1990, % of IPPU	2016	2017	2017, % of IPPU	Trend 1990-2017 (%)	Trend 2016-2017 (%)
2H1. Pulp and paper	10.5	0.1 %	10.2	8.0	0.1 %	-23.8 %	-22.0 %
2H2. Food and beverages industry	20.8	0.1 %	105.3	105.6	1.2 %	408.1 %	0.2 %
2H. Total	31.3	0.2 %	115.6	113.6	1.3 %	263.3 %	-1.7 %

Source: Statistics Norway and Norwegian Environment Agency

4.9.1 Pulp and paper, 2H1

4.9.1.1 Category description

There are CO₂ emissions from non-combustion from two plants in this sector and they are covered by the EU ETS. The emissions originate from the use of limestone. Emissions from combustion are included in Chapter 3.

4.9.1.2 Methodological issues

The CO₂ emissions are calculated by multiplying the amount of limestone by an emission factor.

4.9.1.3 Activity data

Activity data is reported by the plants to the agency. The amount of limestone is calculated from purchased amount, adjusted for the amount of limestone in storage in the beginning and end of the year. The aggregate amounts of limestone used by the plants included in 2H1 are reported in the CRF Table 2(l).A-Hs2 and are shown in Table 4.5 for some selected years in the time series.

4.9.1.4 Emission factors

The emission factor used in the calculation is 0.44 tonne CO₂ per tonne limestone.

4.9.1.5 Uncertainties and time-series consistency

Uncertainty estimates are given in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.9.1.6 Category-specific QA/QC and verification

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII. The plants are covered by the EU ETS and their emissions are verified annually. In addition, the emissions are checked both by the case handler and by the agency's inventory team.

4.9.1.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.9.1.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

4.9.2 Food and beverages industry, 2H2

4.9.2.1 Category description

This source category includes CO₂ from carbonic acid mainly used in breweries, domestic use of captured CO₂, imported CO₂ and CO₂ from production of bio protein.

Some CO₂ from the production of ammonia (2B1) is captured and in Norway mainly used as carbonic acid in carbonated beverages. The emissions reported here in 2H2 include CO₂ bound in products and imported CO₂. The emissions are reported in this source category, although the largest part of the emissions takes place after the bottles is opened and not in the breweries. Exported CO₂ from this source is not included in the Norwegian emission inventory.

One plant produced bio protein in the years 2001-2005. Natural gas was used to feed the bacteria cultures that produced the bio protein and this was used as animal fodder.

4.9.2.2 Methodological issues

For carbonic acid, the CO₂ figures are based on the sales and export statistics from the ammonia producing plant and import statistics from Statistics Norway's External trade in goods statistics.

For the production of bio protein, the plant reported emissions of about 2 000 – 11 000 tonnes CO₂ and these are included in the national inventory.

4.9.2.3 Activity data

For carbonic acid, the CO₂ figures are based on the sales and export statistics from the ammonia producing plant and import statistics from Statistics Norway's External trade in goods statistics, see Table 4.55.

Table 4.55 Sold CO₂ (minus exports) and imported CO₂ (tonnes).

Year	Sold CO ₂ (minus exports)	Imported CO ₂	Domestic use of CO ₂ (2H2)
1990	20 000	787	20 787
1995	34 000	2 374	36 374
2000	50 000	2 597	52 597
2005	52 974	18 433	71 407
2007	50 676	28 512	79 188
2008	63 636	13 974	77 610
2009	61 414	13 664	75 078
2010	76 000	8 675	84 675
2011	76 557	14 750	91 307
2012	81 399	13 560	94 959
2013	78 000	13 249	91 249
2014	83 680	7 712	91 392
2015	88 602	11 144	99 746
2016	80 436	24 912	105 348
2017	45 807	59 805	105 612

Sources: Statistics Norway and the Norwegian Environment Agency

For the production of bio protein, the activity data is the amount of natural gas used in the process.

4.9.2.4 Emission factors

The activity data for carbonic acid is CO₂, so emission factors are not relevant.

4.9.2.5 Uncertainties and time-series consistency

See the uncertainty in the activity data for the ammonia plant (2B1) in Annex II.

A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

4.9.2.6 Category-specific QA/QC and verification

NMVOC and CO₂

The general QA/QC methodology is given in Annex V and the specific QA/QC carried out for Industrial processes is described in Annex VIII.

4.9.2.7 Category-specific recalculations

There have been no recalculations performed for this source category this year.

4.9.2.8 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5 Agriculture (CRF sector 3)

5.1 Overview

About 8.5 % of the total Norwegian emissions of greenhouse gases (GHG) originated from agriculture in 2017. This corresponds to 4.47 million tonnes CO₂-eq. Emissions from agriculture were in 2017 about 5 % lower than in 1990, and about 0.2 % higher than in 2016.

The sector's clearly biggest sources of GHG's were enteric fermentation (CH₄) from domestic animals contributing with 52 % of the sector's emissions, and N₂O from agricultural soils contributing with 36 %. Manure management contributed with about 10 %. CO₂ emissions in the agriculture sector, mainly from liming and a minor part from urea application, contributed with 2 %. There are also some minor emissions of the greenhouse gases N₂O and CH₄ arising from the burning of crop residues on the fields.

Table 5.1 Emissions from Agriculture categories in 1990, 2016 and 2017 (ktonnes CO₂-equivalents).

Category	1990	1990, % of Agriculture	2016	2017	2017, % of Agriculture	Trend 1990-2017 (%)	Trend 2016-2017 (%)
3A Enteric fermentation	2 415	51 %	2 305	2 329	52 %	-4 %	1 %
3B Manure management	339	7 %	426	430	10 %	27 %	1 %
3D Agricultural soils	1 672	36 %	1 642	1 622	36 %	-3 %	-1 %
3F Field burning of agricultural residues	36	1 %	4	4	0 %	-89 %	-1 %
3G Liming	231	5 %	82	84	2 %	-64 %	2 %
3H Urea application	0.6	0 %	0.2	0.1	0 %	-83 %	-52 %
Total	4 694	100 %	4 459	4 469	100 %	-5 %	0.2 %

Source: Statistics Norway and the Norwegian Environment Agency

Agriculture contributes particularly to CH₄, N₂O and NH₃ emissions. Domestic animals are the major source of CH₄ emissions from agriculture. Both enteric fermentation and manure management contribute to emissions of CH₄. Manure management also generates emissions of N₂O.

Microbiological processes in soil lead to emissions of N₂O. Both direct and indirect N₂O from soil processes are described in the IPCC methodology and are included in the Norwegian inventory. Direct N₂O emissions arising from the use of fertilizer (manure, synthetic fertilizer, sewage sludge and other organic fertilizers applied to soils), emissions from pastures, crop residues and cultivation of organic soils are included. Indirect N₂O emissions from atmospheric deposition and nitrogen leaching and run-off from both the manure management systems and from agricultural soils are also included.

Manure storage and the use of fertiliser (manure, synthetic fertilizer, sewage sludge and other organic fertilizers applied to soils) also generate emissions of ammonia (NH₃) and NO_x that gives

indirect N₂O from atmospheric deposition. NH₃ volatilized from grazing animals are also included in the estimations of indirect N₂O.

As indicated in chapter 1.5, the Approach 2 key category analysis performed in 2019 for the years 1990 and 2017 has revealed key categories in terms of total level and/or trend uncertainty in the agriculture sector as shown in Table 5.2. The key categories according to Approach 1 key category analysis are also provided in Table 5.2.

Table 5.2 Key categories in the sector Agriculture.

IPCC	Source category	Gas	Key category according to approach	Method
3A	Enteric fermentation	CH ₄	2	Tier 1/2
3B1	Manure management - Cattle	CH ₄	1	Tier 2
3B	Manure management	N ₂ O	2	Tier 2
3Da1	Direct emissions from managed soils - Inorganic N fertilizers	N ₂ O	2	Tier 1
3Da2	Direct emissions from managed soils - Organic N fertilizers	N ₂ O	2	Tier 1
3Da3	Direct emissions from managed soils – Urine and dung deposited by grazing animals	N ₂ O	2	Tier 1
3Da4	Direct emissions from managed soils - Crop residues	N ₂ O	2	Tier 1
3Da6	Direct emissions from managed soils - Cultivation of organic soils	N ₂ O	2	Tier 1
3Db1	Indirect emissions from managed soils – Atmospheric deposition	N ₂ O	2	Tier 1
3Db2	Indirect emissions from managed soils – Nitrogen leaching and run-off	N ₂ O	2	Tier 1
3G	Liming	CO ₂	1	Tier 1

5.2 Livestock population characterisation

5.2.1 Data sources

The animal population data used in the estimations on a disaggregated level are provided in Annex IX, Table AIX-1. The same data for number of animals of the various animal groups is used in all the different calculations of emissions.

The main sources of the livestock statistics are the register of production subsidies (sheep for breeding, goats, breeding pigs, poultry for egg production and beef cows), statistics of approved carcasses (animals for slaughter) and the Cow Recording System at TINE BA¹⁶ (TINE BA Annually) (heifers for breeding and dairy cows). These sources cover 80-100 % of the animal populations. The estimated shortage of coverage is compensated in the total number of animals used in the emission estimates. The coverage in the data sources is shown in Table 5.3.

¹⁶ TINE BA is the sales and marketing organisation for Norway's dairy cooperative and covers most of the milk production and the meat production induced by milk production.

Table 5.3 Estimated coverage of animal populations in the the data sources used. 2017

	Statistics Norway, production subsidies	Statistics Norway, statistics of approved carcasses ⁴	TINE	Other
Dairy cows		100 ²	96.4 ¹	
Replacement heifers			96.4 ¹	
Young cattle for slaughter		100	100 ³	
Beef cows	99.8	100 ²		
Sheep	99.6	100		
Goats	100			
Laying hens	99.9			
Chics for breeding	100			
Chicken for slaughter		100		
Other poultry for breeding	100			
Other poultry for slaughter		100		
Sows	97.8			
Young pigs for breeding	100			
Pigs for slaughter		100		
Horses	Unkown ⁵			Unkown ⁵
Fur-bearing animals	100			
Deer	100			
Reindeer				100 ⁶

Source: Estimations by Statistics Norway and the Cow Recording System (dairy cows and heifers).

¹ Share of livestock herds.

² Data source only for slaughter weight

³ Data source only for slaughter age

⁴ Figure refers to share of slaughtered animals, excluding home slaughter. Animals dead from other causes also excluded

⁵ Total number of horses used in the inventory is based on data from productions subsidies (roughly 50 % of total number) and an additional estimation of number of horses outside agriculture by NIBIO.

⁶ Norwegian Agriculture Agency

The statistics of approved carcasses covers close to 100 % of all slaughtered animals. Home slaughter is not included, but the extent of home slaughter is very low due to legal restrictions. Even animals consumed by producers are in most cases registered at the slaughterhouses.

The number of dairy cows and heifers for breeding derive from the Cow Recording Systems (TINE BA Annually). Between 98 and 99 % of all dairy cows are registered here, and in addition, the number used in the inventory is adjusted for this missing part. The adjustment is based on the percentage of herds controlled by the cow recording system.

The registers are updated annually. In addition to the animals included in these registers, an estimate of the number of other horses is obtained from the Norwegian Institute of Bioeconomy Research (NIBIO)¹⁷. The number of reindeer is obtained from the Norwegian Reindeer Husbandry Administration.

For the categories of animals living shorter than a full year or two, generally animals for slaughter, lifetime is taken into account to get a yearly average for the number of animals.

5.2.2 Method for estimating number of cattle

For dairy cows, additional information from the Cow Recording System concerning annual milk production and proportion of concentrate in the diet is used (TINE BA Annually). The Cow Recording System also supplies annual information about slaughter age for heifers and bulls and data for estimating live weight of dairy cows and heifers for breeding, and also the age of young cows at their first calving. (Moen, *pers. comm.*¹⁸).

For heifers and bulls for slaughter, animal numbers are based on data from statistics of approved carcasses which provide data on numbers slaughtered and slaughter weights. Combined with slaughter age from the Cow Recording System (TINE BA Annually), this gives a precise estimation of animal life time for each animal slaughtered. There are several reasons for this way of counting the animals:

1. The coefficients for N and VS excretion (3B) and formation of methane (3A) is based on the development of the animals through their lifetime (weight gain, age, feed consumption). The information on these parameters is most precise when they are connected to the animals in this way.
2. When using the number of animals slaughtered, there is an almost perfect coherence between the definition of the categories counted and the representation of the coefficients.
3. Counting the number of animals slaughtered gives a very precise value of the number of animals.
4. Counting the number of animals as number of live animals at specific dates (which is the only other alternative) takes away the possibility to group the animals in the five categories of young cattle that is being used in the definitions now because this statistic only gives the number of young cattle as a total.

One principal draw-back of this method for estimating animal population is that emissions in all stages of these animals' lives will be accounted for in the year of slaughter, even though the

¹⁷ Former named the Agricultural Economics Research Institute (NILF).

¹⁸ Moen, O. (*annually*): Personal information, email from Oddvar Moen, Tine Rådgivning annually.

emissions in the early stages of the lives of these animals to a large extent took place in the previous year. In a stable population of animals, this error is automatically adjusted for. Since animal populations are relatively stable, this error is considered much smaller compared to errors related to estimating animal year based on animal populations in the register of production subsidies which was previously used. The data sources used also ensure a better coherence between animal numbers, life time and weight. Estimated animal years for cattle are provided in Table 5.4.

The number of milk cows calving for the first time (=heifers for replacement) and their average age at time of calving is reported by the Cow Recording System (TINE BA Annually) on request from Statistics Norway. These data date back to 2004. For the years 1990-2003, average fraction (number of heifers)/(number of milk cows) for the years 2004-2011 is used to estimate number of heifers based on number of milk cows. Number of heifers for replacement in beef production is collected from annual reports from Animalia (Norwegian Meat and Poultry Research Center (www.animalia.no)). Figures exist from 2007. For previous years, the number is estimated with the same method as for heifers for milk production.

Table 5.4 Estimated animal years for cattle

	Heifer for replacement	Heifers for slaughter	Bulls for slaughter	Beef cows ¹	Dairy cows
1990	311 279	47 020	289 945	8 193	325 896
1995	299 284	47 103	284 237	20 334	310 346
2000	280 121	63 512	285 349	42 324	284 880
2005	255 862	57 619	263 170	54 841	255 663
2008	240 399	54 831	238 111	60 401	238 550
2009	247 902	53 397	235 689	63 803	235 480
2010	239 839	53 410	230 872	67 110	232 294
2011	239 007	48 778	223 536	68 539	224 721
2012	235 891	42 863	217 050	71 834	229 767
2013	239 386	47 294	220 401	70 969	225 163
2014	244 601	67 624	208 979	73 894	221 032
2015	238 485	64 814	206 328	77 408	217 576
2016	241 173	64 361	217 885	84 372	215 015
2017	245 716	43 501	250 630	88 332	217 318

¹ Counted animals

Source: Cow Recording System (TINE BA Annually)(dairy cows), slaughter statistics and estimations by Statistics Norway

5.2.3 Method for estimating number of sheep

In the estimations of emissions from manure management, the sheep population is divided between sheep > one year, and sheep < one year. Data from both the register of production subsidies and slaughter statistics is used in estimating the number of animals.

Sheep more than one year old is estimated as the number of sheep registered 1. of January deducted for the number of sheep slaughtered Jan.-May the same year. The sheep slaughtered later in the year are counted as living the whole year.

Sheep less than one year old is estimated as number of sheep under one year registered 1. of January + number of lambs slaughtered June-December *143/365. Lambs slaughtered before June are assumed to be registered as sheep under one year 1. of Jan. Practically all lambs slaughtered after June are born in the spring. An expert judgment suggests an average lifetime of 143 days for slaughtered lambs born in the spring (UMB, *pers. comm*¹⁹).

In the estimations of enteric methane, sheep is split in four categories: sheep > 1 year, sheep < 1 year for breeding, lambs slaughtered June-Dec. and lambs slaughtered Jan.-May. Sheep over one year is estimated as explained above. Sheep under one year for breeding is estimated as the number of sheep under one year registered 1. of January deducted for the number of lambs slaughtered Jan.-May. For the numbers of slaughtered lambs, slaughter statistics are used. The numbers of slaughtered lambs were previously estimated. A more detailed description of the method can be found in Annex IX.

5.2.4 Deviations from FAO statistics

There are some differences between the number of animals used in these calculations and the FAO statistics. The general reason that animal statistics used in the emission inventory differ from the statistics delivered to FAO is that the statistics are used for different purposes. Animal statistics used in the inventory has to be categorized so that the categories fit the recommended methodology and the various emission factors used in the emission estimations. The figures reported to the FAO are provided by the Norwegian Institute of Bioeconomy Research (NIBIO)²⁰. NIBIO makes an overall estimation for the agricultural sector, which is the basis for the annual negotiations for the economic support to the sector. This estimate includes a grouping of all agricultural activities, comprising area, number of animals and production data. Differences include:

- Different emphasis on the dates for counting, 31.07 and 31.12
- NIBIO does not register pigs under 8 weeks, whilst Statistics Norway does. For the number of animals for slaughter, Statistics Norway uses the statistics of approved carcasses, which together with data on slaughter age gives a far better figure on estimated animal years (average population through the year) compared to figures for registered animals at specific dates which is used in the FAO statistics.
- For the number of dairy cows and heifers for replacement, Statistics Norway uses statistics from the Cow Recording System (TINE BA Annually), which is presumed to give a more

¹⁹ UMB (2001): Expert judgement by Department of Animal Science, Ås: Norwegian University of Life Sciences.

²⁰ Former named the Agricultural Economics Research Institute (NILF).

accurate figure on number of animal years of dairy cows than the figures from Statistics Norway.

Emissions from other animal groups than included in the estimations (ostrich, donkey, lama and alpaca) are expected to be very small and decreasing. Emissions from ostrich have earlier been included in the estimations, but the number of ostrich has had a decreasing trend and are now very limited (39 in 2013). At the most the number of ostrich was 2113 in 1999. The total emissions from ostrich were less than 500 tonnes of CO₂ equivalents when the animal population was at its highest.

5.3 Nitrogen in animal manure as basis for emission estimates

Access to nitrogen is vital for all plant growth; hence nitrogen is added to the soil from i.a. animal manure. This causes emissions to air at various points of compounds containing nitrogen. Of the nitrogen compounds emitted to air from animal manure, N_2O , NO_x and NH_3 are estimated.

According to the IPPC and LRTAP guidelines, process emissions of nitrogen compounds from use of animal manure are calculated from the following sources:

1. Manure management systems (N_2O , NH_3 and NO_x)
2. Application of manure on soil (N_2O , NH_3 and NO_x)
3. Droppings from animals on pastures (N_2O , NH_3 and NO_x)
4. Leakage of nitrogen through manure management systems and soils (N_2O)
5. Deposition of nitrogen from emissions of NH_3 and NO_x (N_2O)


Though the nitrogen flow is continuously depending on its surroundings (soil characteristics, temperature, moisture etc.) and the preceding supplies and losses of N, the emission estimates of each of the sources above are generally done independently of emissions from the other sources mentioned. Figure 5.1 gives an overview of the manure nitrogen flows in the Norwegian greenhouse gas inventory.

The following decides the amounts of N that are used as the basis for the respective emission calculations:

- The amount of N in manure systems is calculated as total N in manure adjusted for the N that is dropped on pastures.
- N_2O emitted during spreading is calculated from the amounts of N in manure spread to land. This means that N lost through leaching in manure storage and as N_2O , NH_3 and NO_x in manure housing and storage is deducted. However, N lost as N_2O , NH_3 and NO_x during spreading, as well as indirect emissions of N_2O due to atmospheric deposition, are not deducted.
- NH_3 emitted during and after spreading of manure is based on the amounts of TAN in manure spread to land minus N lost through leaching in manure storage and as N_2O , NH_3 and NO_x in manure housing and storage. NO_x emitted during and after spreading of manure also has the same basis. For NH_3 emissions, N lost as N_2O and NO_x during spreading, as well as indirect emissions of N_2O due to atmospheric deposition is not deducted. Similarly, for NO_x emissions, N lost as N_2O and NH_3 during spreading, as well as indirect emissions of N_2O due to atmospheric deposition is not deducted.
- Emissions of N_2O , NH_3 and NO_x from pasture are calculated independently of each other, and are based on the amounts of N (or TAN for NH_3 emissions) estimated in manure dropped during grazing.
- N_2O lost through leaching due to spreading is based on total N in manure spread to land minus N lost through leaching in manure storage and as N_2O , NH_3 and NO_x in manure housing and storage. N_2O lost through leaching due to grazing is based on total N excreted on pastures. N_2O lost through leaching during storage of manure is based on the amounts of

N estimated for the particular management systems that are susceptible to leaching. N lost through emissions of NH_3 from housing is not deducted.

- The nitrogen in NH_3 and NO_x volatilised during housing, storage, pasture and spreading of manure is the basis for the calculation of N_2O emissions from atmospheric deposition. How the amounts of N are estimated in the various emission estimates, is described in more details in the respective chapters below.



¹ For estimation of NH_3 and NO_x emissions from manure storage systems, emissions of NH_3 from housing are deducted from N excreted in housing. N_2O emissions (direct and indirect) are estimated directly from N excreted in housing.

² Emissions of N_2O , NH_3 and NO_x that have occurred prior to spreading of manure on managed soils (during housing and storage) are deducted before emissions of N_2O , NH_3 and NO_x from application to soils are estimated.

Figure 5.1 Overview of the manure nitrogen flows in the Norwegian greenhouse gas inventory.

5.4 Emissions from enteric fermentation in domestic livestock - 3A (Key category for CH₄)

5.4.1 Category description

An important end product from the ruminal fermentation is methane (CH₄). The amount of CH₄ produced from enteric fermentation is dependent on several factors, like animal species, production level, quantity and quality of feed ingested and environmental conditions. According to IPCC the method for estimating CH₄ emissions from enteric fermentation requires three basic items:

- The livestock population must be divided into animal subgroups, which describe animal type and production level.
- Estimate the emission factors for each subgroup in terms of kilograms of CH₄ per animal per year.
- Multiply the subgroup emission factors by the subgroup populations to estimate subgroup emissions, and sum across the subgroups to estimate total emission.

Enteric fermentation is a key category both for level and trend assessment.

Enteric fermentation contributed with 2 329 ktonnes CO₂ equivalents in 2017, which is 52 percent of the GHG emissions from this sector. Emissions decreased by 4 % in the period 1990-2017 and increased by 1 % in 2016-2017.

5.4.1.1 Methodological issues

A Tier 2 methodology is used for calculating CH₄ from enteric fermentation for the main emission sources cattle and sheep.

For dairy cattle gross energy intake, GE and the methane conversion rate, Y_m are based on milk yield and the percentage of concentrate in the diet. This Tier 2 methodology used is described in more detail in Annex IX, section 2.

For beef cattle the Tier 2 method is also presented further in Annex IX. The method is based on data from one year, and there have been some changes in the composition of the breed of the beef population in Norway since 1990. But we lack data for a good variable that we could use to get a trend for GE and Y_m for beef cow. This is expected to be a minor source of error, since the population of beef cow was of less significance earlier. Since the change in the composition of the population has been an increase of heavier breed, does this mean that there is an overestimation of the emissions for the earlier years.

The methodology for calculating CH₄ from enteric fermentation for the other animal categories is in accordance with the Tier 1 method from the IPCC guidelines (IPCC 2006). The numbers of animals of each category and average emission factors of tonnes CH₄ per animal and year for each category of animals are used to calculate the emissions.

5.4.1.2 Activity data

Emissions are estimated from the animal population. How the animal population is estimated is described in Section 5.2 and Annex IX.

The Tier 2 method of calculation which is implemented for cattle and sheep requires subdividing the cattle and sheep populations by animal type, physiological status (dry, lactating or pregnant) live weight and age. Table 5.5 describes the animal categories used for cattle and sheep in the calculations.

Table 5.6 and Table 5.7 gives important input parameters in the estimations of enteric methane from cattle.

Table 5.5 Categories of cattle and sheep used in the Norwegian calculations of methane emission from enteric fermentation.

Categories of cattle and sheep
Dairy cows
Beef cows
Replacement heifers
Finisher heifers, < one year at time of slaughter
Finisher heifers, > one year at time of slaughter
Finisher bulls, < one year at time of slaughter
Finisher bulls, > one year at time of slaughter
Breeding sheep, > one year
Breeding sheep, < one year
Slaughter lamb, < one year. Jan- May
Slaughter lamb, < one year. Jun- Dec

Table 5.6 Important parameter inputs in the calculations of methane emissions from cattle over 1 year.

	Annual milk production, dairy cows. kg/animal/year	Proportion of feed concentrate in the rations of mature dairy cows. %	Carcass weight at time of slaughter, heifer > 1 year. kg	Age at time of slaughter, heifers > 1 year. Months	Carcass weight at time of slaughter, bulls > 1 year. Kg	Age at time of slaughter, bulls > 1 year. Months
1990	6 320	39.1	185	21.6	255	19.7
1995	6 326	36.8	200	22.2	276	19.7
2000	6 156	36.4	202	22.3	269	18.8
2005	6 723	37.7	216	22.8	296	19.0
2008	7 144	39.8	213	22.5	298	18.2
2009	7 276	40.1	219	22.8	301	18.0
2010	7 373	41.0	221	22.8	302	18.0
2011	7 309	41.9	210	22.5	297	17.7
2012	7 475	42.9	205	22.7	294	17.7
2013	7 691	43.4	209	22.8	298	17.5
2014	7 711	43.4	244	22.8	302	17.3
2015	7 958	43.6	256	23.2.8	310	17.4
2016	8 062	43.6	260	23.2	317	18.2
2017	7902	43.6	251	23.0	313	18.4

Source: Cow Recording System (TINE BA Annually) (dairy cows) and estimations by Statistics Norway

For dairy cows GE is based on milking yield due to annual milk production and proportion of concentrate in the diet.

Table 5.7 Important parameter inputs in the calculations of methane emissions from cattle under 1 year.

	Heifers < 1 year. Carcass weight	Heifers < 1 year. Average age, months	Bulls < 1 year. Carcass weight	Bulls < 1 year. Average age, months
1990	56.30	6.46	75.81	6.43
1995	69.65	7.00	93.79	6.94
2000	65.00	6.05	82.05	5.88
2005	92.87	7.86	115.60	7.46
2008	92.49	7.89	116.49	7.53
2009	93.28	8.02	118.42	7.56
2010	93.23	8.09	116.05	7.50
2011	94.71	8.15	117.61	7.50
2012	95.88	7.92	119.66	7.56
2013	101.58	8.15	122.50	7.59
2014	106.02	8.18	124.47	7.52
2015	108.90	8.125	125.19	7.52
2016	106.89	7.86	126.29	7.52
2017	103.52	8.23	125.80	7.63

Source: Cow Recording System (TINE BA Annually) and estimations by Statistics Norway

Table 5.8 Important parameter inputs in the calculations of methane emissions from sheep. 2017

	Carcass weight. kg	Age at slaughter. Months	Conversion factor for methane. %
Breeding sheep > 1 year	30.3	NA	6.5
Breeding sheep < 1 year	29	NA	4.5
Lamb for slaughter, Jan.-May	18.62	11	4.5
Lamb for slaughter, June-Dec.	18.44	4.7	4.5

Source: Carcass weight from slaughter statistics, Statistics Norway, and age at slaughter and conversion factors from Volden and Nes, 2006, see Annex IX section 2.

5.4.1.3 Emission factors

For cattle and sheep, the following basic equation is used to calculate the CH₄ emission factor for the subgroups (Tier 2):

$$EF = (GE \cdot Y_m \cdot 365 \text{ days/yr}) / (55.65 \text{ MJ/kg CH}_4)$$

Where:

EF = emission factor, kg CH₄/head/yr

GE = gross energy intake, MJ/head/day

Y_m = CH₄ conversion rate, which is the fraction of gross energy in feed converted to CH₄.

This equation assumes an emission factor for an entire year (365 days). In some circumstances the animal category may be alive for a shorter period or a period longer than one year and in this case the emission factor will be estimated for the specific period (e.g., lambs living for only 143 days and for beef cattle which are slaughtered after around 540 days, varying from year to year). Further description of the determination of the variables GE and Y_m for the different animal categories and the values used in the calculations are given in Annex IX.

The emissions from hens and pullets, domestic reindeer, deer and fur-bearing animals are also included in the Norwegian calculations. The Norwegian University of Life sciences has investigated and documented the national emission factors for poultry. Only hens and pullets have emissions of significance (Svihus 2015). For reindeer the emission factor 14.0 kg/animal/year is used and for deer 20.0 kg/animal/year. Both factors are expert judgments from the University of Life Sciences (Karlengen et al. 2012) and have been estimated based on the methodology described for cervidae in IPCC (2006). Danish emission factors are used for goat since they are considered to reflect Norwegian feed intake and circumstances (Karlengen et al. 2012). Emission factor for fur-bearing animals has been developed by scaling emission factor for pigs, which are assumed most similar with regard to digestive system and feeding. The scaling is done by comparing average weights for fur-bearing animals and pigs and the factor is set to 0.1 kg/animal/year.

For the other animal categories the Tier 1 default emission factors for each kind of animal (IPCC 2006) is used. The factors used are shown in Table 5.9.

Table 5.9 Emission factors for CH₄ from enteric fermentation and different animal types estimated with the Tier 1 method

Animal	Emission factor (Tonnes/animal/year)	Source
Horses	0.018	(IPCC 2006)
Goats	0.013	(Karlengen et al. 2012)
Pigs	0.0015	(IPCC 2006)
Hens	0.00002	(Svihus 2015)
Pullets	0.0000036	(Svihus 2015)
Reindeer	0.014	(Karlengen et al. 2012)
Deer	0.02	(Karlengen et al. 2012)
Fur-bearing animals	0.0001	Estimate by Statistics Norway

Source: IPCC (2006), Karlengen et al. (2012), Svihus (2015).

5.4.2 Uncertainties and time-series consistency

Activity data

The data is considered to be known within ± 5 %. There is also uncertainty connected to the fact that some categories of animals are only alive part of the year and the estimation of how long this part is.

Emission factors

Although the emissions depend on several factors and therefore vary between different individuals of one category of animal, average emission factors for each category are used in the tier 1 methodology for all animal categories except cattle and sheep, where a tier 2 methodology is used.

The standard deviation of the emission factors is considered to be ± 40 %, which is the estimate from the IPCC guidelines (IPCC 2006). An uncertainty estimate of ± 25 % is used for the emission factors for cattle and sheep in the Tier 2 methodology (Storlien & Harstad 2015).

5.4.3 Category specific QA/QC and verification

In 2001, a project was initiated to improve the estimate of the exact number of animal populations. This was completed in 2002. In 2012, a further revision of the numbers of bulls and heifers was implemented. The revised data on animal populations form the basis for the emission calculations for all years. In 2005-2006, Statistics Norway and the Climate and Pollution Agency carried out a project in cooperation with the Norwegian University of Life Sciences, which resulted in an update of the emission estimations for cattle and sheep using a tier 2 method. In 2015, the equations of this model were updated for cattle based on Norwegian data from the Cow recording system/NorFor (Storlien & Harstad 2015).

The Norwegian University of Life sciences has further investigated and documented the national emission factor of 20 g CH₄ per head used for laying hens in a project in 2015 (Svihus 2015). New emission factors for poultry in Norway were estimated in Svihus (2015). Only hens and pullets have measurable emissions. In the previous submissions, only hens and turkeys were considered. The new emissions factors have increased total emissions from poultry between 1 and 10 tonnes of CH₄ per

year. Total emissions were 98 tonnes in 2015, which shows this an insignificant source of CH₄ emissions.

In 2015, a project at the Norwegian University of Life sciences NMBU investigated the basic equations used to calculate the emission factors for enteric methane for cattle in the tier 2 methodology. The results of this project were implemented in the 2016 submission.

5.4.4 Category-specific recalculations

There have been no recalculations performed for this source category this year.

5.4.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5.5 Emissions from manure management - 3B (Key categories for CH₄ and N₂O)

5.5.1 Category description

The relevant greenhouse gases emitted from this source category are CH₄ (IPCC 3Ba) and N₂O (IPCC 3Bb). Emissions from cattle are most important in Norway for both components.

N₂O emissions from manure management is key category according to the Approach 2 key category analysis and CH₄ emissions from cattle manure management is key category according to Approach 1 key category analysis.

CH₄ emissions due to manure management amounted to 282 ktonnes CO₂ equivalents in 2017 whilst N₂O emissions amounted to 148 ktonnes CO₂ equivalents.

Manure management emitted 430 ktonnes of CO₂ equivalents in 2017, which are approximately 10 % of the GHG emissions from agriculture.

Emissions of GHGs from manure management increased by 27 % in the period 1990-2017, and by 1 % from 2016 to 2017.

Organic material in manure is transformed to CH₄ in an anaerobic environment by microbiological processes. Emissions from cattle (manure) are most important in Norway. The emissions from manure depend on several factors; type of animal, feeding, manure management system and weather conditions (temperature and humidity).

During storage and handling of manure (i.e. before the manure is added to soils), some nitrogen is converted to N₂O. The fraction converted to N₂O depends on the system and duration of manure management. Liquid system is the most widespread storage system, and consequently the most important source. Indirect emissions of N₂O (atmospheric deposition and leaching) from manure storage are also estimated.

Emissions of NH₃ from manure depend on several factors, e.g. type of animal, nitrogen content in fodder, manure management, climate, time of spreading of manure, cultivation practices and characteristics of the soil. In the IPCC default method, a NH₃ volatilisation fraction of 20 % is used for the total N excretion by animals in the country. However, in the Norwegian emission inventory, yearly updated NH₃ volatilisation estimations are used, because this is expected to give more correct values for Norway. The estimated national volatilization fractions from spreading of manure ($\text{frac}_{\text{gasm}}$) have differed between 13-14 % since 1990.

Like for NH₃, emissions from NO_x from animal manure lead to indirect emissions of N₂O from deposition. The amounts of N from NO_x emissions from manure management systems, spreading of manure and droppings on pastures are however small.

Section 5.3 gives more information about nitrogen in animal manure as basis for emission estimates and an overview of the manure nitrogen flows in the Norwegian greenhouse gas inventory.

5.5.1.1 Methodological issues

CH₄

For sheep, goat, horse, deer, reindeer, mink and fox, IPCC Tier 1 methods are used for the estimations of emission of CH₄ from manure management (IPCC 2006). The emission factors used are based on country specific expert judgements (Karlengen et al. 2012) where such exists (horse, mink and fox, deer and reindeer), while for sheep and goat the IPCC default emission factors are used.

For cattle, swine and poultry, emissions of methane from manure are estimated using the following equations, in accordance with the IPCC Tier 2 method (IPCC 2006).

$$CH_4 \text{ Emissions} = EF * Population$$

$$EF_i = VS_i * 365 \text{ days/year} * B_{0i} * 0.67 \text{ kg/m}^3 * \sum_{(jk)} MCF_{jk} * MS_{ijk}$$

EF_i = annual emission factor for defined livestock population *i*, in kg

VS_i = daily VS excreted for an animal within defined population *i*, in kg

B_{0i} = maximum CH₄ producing capacity for manure produced by an animal within defined population *i*, m³/kg of VS

MCF_{jk} = CH₄ conversion factors for each manure management system *j* by climate region *k*

MS_{ijk} = fraction of animal species/category *i*'s manure handled using manure system *j* in climate region *k*

The factors VS, B₀ and MCF are average factors meant to represent the whole country. The populations of animals are consistent with the animal data used elsewhere in the inventory (see chapter 5.2 and Annex IX for further details). For young cattle, this implies that the VS production is estimated for the whole average life time/time until first calving and not per animal year. The amount of volatile solids (VS) for other cattle is estimated directly as kg/animal/year. The VS factors are based on the same data sources used in the estimations of nitrogen excretion factors, that are used in estimations of N₂O from manure (Karlengen et al. 2012). For beef cows, national factors for VS and nitrogen excretion factors were estimated by the Norwegian University of Life Sciences (NMBU) in 2018. National feeding data for beef cow under Norwegian circumstances from Wetlesen et. al. (2018), the same as used for the estimation of enteric methane, has been used as a basis for the estimations. The method is described more in detail in Aspeholen Åby et al (2019), (published in NIR 2019, Annex IX, section 4). For swine and poultry, country specific estimates of NMBU for the percentage of the manure in dry matter that are volatile solids are used. Background data used for the estimations of VS are given in Table 5.10 and in Annex IX.

The factor B₀ represents the maximum potential production of methane under optimum conditions. For dairy cows, the B₀ factors are based on Norwegian research and for pigs the factor is based on literature studies (Morken et al. 2013), for other cattle and poultry the default IPCC factors are used.

National Inventory Report 2019 - Norway

Table 5.10 Norwegian factors for amount of manure (in dry matter - d.m.), VS and Bo used to estimate CH₄ from manure management with the IPCC Tier 2 method. 2017.

	Manure (kg dry matter per animal)	VS %	VS, kg per animal	VS, total, tonnes	Bo
Non-Dairy Cattle			963	381 128	0.18
Beef cows			1461	129 053	0.18
Replacement heifer			966	110 030	0.18
Finisher heifer			766	18 515	0.18
Finisher bulls			728	123 529	0.18
Dairy cows			1 525	331 401	0.23
Poultry					
Hens	13.15	0.9	11.84	51 664	0.39
Chicks bred for laying hens, animal places	3.10	0.9	2.79	3 059	0.36
Chicks for, slaughter animal places	4.08	0.9	3.67	35 882	0.36
Ducks for breeding	30.00	0.9	27.00	83	0.36
Ducks for slaughter, animal places	8.12	0.9	7.31	405	0.36
Turkey and goose for breeding	30.00	0.9	27.00	473	0.36
Turkey and goose for slaughter, animal places	17.23	0.9	15.51	6 434	0.36
Swine					
Piglets 4-10 weeks	18.41	0.9	16.57	4 426	0.30
Young pigs for breeding	113.00	0.9	101.70	4 170	0.30
Pigs for slaughter, animal places/årsdyr	131.34	0.9	118.21	56 921	0.30
Sows	307.90	0.9	277.11	13 050	0.30
Boar	307.90	0.9	277.11	221	0.30

Sources:

Manure, dry matter poultry and swine: Karlengen et al. (2012)

VS%, poultry: expert estimate Birger Svihus NMBU, email 03.01.2013

VS%, swine: expert estimate Nils Petter Kjos, NMBU, email 03.01.2013.

VS per animal, cattle: Estimates based on Karlengen et al. (2012)

Bo: Morken et al. (2013) for dairy cattle and swine and IPCC (2006) for other animal groups.

For MCF, standard IPCC factors from 2006 IPCC Guidelines (IPCC 2006) are used for the different manure management systems.

Table 5.11 Norwegian factors for MCF used to estimate CH₄ from manure management with the IPCC Tier 2 method

	MCF
Pit storage below animal confinement >1month ¹	0.17
Pit storage below animal confinement <1month ¹	0.03
Liquid / slurry without cover	0.17
Liquid / slurry with cover	0.1
Solid storage	0.02
Cattle and swine deep bedding	0.17
Dry lot	0.01
Poultry manure	0.015
Pasture range and paddock, cattle	0.02
Pasture range and paddock, horses, goats and sheep	0.01

¹The share of the manure stored over and under one month before spreading is based on expert judgement by J. Morken, Norwegian University of Life Sciences, 06.08.14.

Sources: IPCC (2006)

N₂O

In Norway, all animal excreta that are not deposited during grazing are managed as manure. N₂O emissions from manure are estimated in a revised nitrogen model which closely follows the stepwise approach proposed in the EMEP/EEA 2016 guidelines (EEA 2016). As the model is based on the nitrogen mass balance approach specified by EMEP/EEA, it allows estimates to be made of all the main nitrogen species, namely NH₃, NO and N₂ in addition to N₂O. The estimations are made in accordance with the IPCC tier 2 method (IPCC 2006), using Norwegian values for N in excreta from different animals according to Table 5.12. The rationale for the Norwegian values for N in excreta is provided in Karlengen (2012). For beef cow, the nitrogen excretion factors were estimated by the Norwegian University of Life Sciences (NMBU) in 2018 based on national feeding data for beef cow (Wetlesen et al 2018). The method is described more in detail in Aspeholen Åby et al. (2019) (published in NIR 2019, Annex IX, section 3.2.2). The N-excretion factors for cattle, poultry and pigs have been scientifically investigated, while the remaining categories have been given by expert judgements (Karlengen et al. 2012). Based on typical Norwegian feedstock ratios, the excretion of nitrogen (N) were calculated by subtracting N in growth and products from assimilated N. Comparisons have also been made with emission factors used in other Nordic countries and IPCC default factors.

The factors for cattle are based on equations using animal weight, production (milking cows), life time (young cattle) and protein content in the fodder as activity data.

The Nordic feed evaluation system (NorFor) was used to develop the nitrogen factors for dairy cows and young cattle. Excretions of N in the manure were calculated as the difference between their intake, and the sum of what is excreted in milk, fetus and deposited in the animal itself. The procedure used for calculating the excretion of faeces and N consisted of two steps:

1. Simulations in "NorFor" were conducted to gain values for the faeces/manure characteristics covering a wide variation of feed characteristics (N content) and production intensities (milk yield/meat production)
2. The results from the simulations were used to develop regression equations between faeces/manure characteristics and parameters related to the diet (N content) and animal characteristics (milk yield, weight, age etc).

Calculations of N-factors based on these equations have been made back to 1990 for cattle. For beef cattle we only have national feeding data from one year. There have been some changes in the composition of the breed of beef cow population in Norway since 1990. But we lack data for a good variable that we could use to get a trend for beef cow. It is expected that this is a minor source of error, since the population of beef cow was of less significance earlier. Since the change in the composition of the population has been an increase of heavier breed, does this mean that there is an overestimation of the emissions for the earlier years. For poultry and pigs, N-factors have been estimated for 2011 in Karlengen et al. (2012). The factors used until this update were estimated in 1988 (Sundstøl & Mroz 1988), and are regarded as still valid for 1990. A linear interpolation has been used for the years between 1990 and 2011. For the remaining animal categories, N in excreta is considered constant throughout the time series. More background data for the calculations is provided in Annex IX, Section 3.1. The factors are shown in Table 5.12. The factors for total N excreted are used in the estimations of N₂O emissions, whilst ammonium N is used in the estimations of NH₃ and NO_x emissions. N₂O emissions from MMS are estimated based on total N excreted, while all other N emissions from MMS are estimated based on N available prior a specific MMS stage.

Norwegian values are also used for the fraction of total excretion per species for each management system (MS) and for pasture. The fractions are updated every year and are provided in Table 5.14 and Table 5.15. This is described in section 5.5.1.2.

Table 5.12 N in excreta from different animal categories¹. 2017. kg/animal/year unless otherwise informed in footnote.

	Total N	Ammonium N
Dairy cattle	128.6	73.3
Beef cows	93.0	52.6
Replacement heifers ²	86.8	47.8
Heifers for slaughter ²	68.1	41.4
Bull for slaughter ²	72.2	43.9
Sows	24.4	15.3
Boars	24.4	15.3
Piglets	1.4	0.9
Fattening pigs ³	3.2	2.1
Young pigs for breeding	9.7	6.5
Laying hens	0.7	0.3
Chickens reared for laying ³	0.05	0.02
Broilers ³	0.03	0.01
Turkeys for slaughter ³	0.5	0.2
Ducks and geese for slaughter ³	0.1	0.03
Turkeys, ducks and geese reared for laying	2.0	0.8
Horses	50.0	25.0
Dairy goats	16.9	10.1
Other goats	8.5	5.1
Sheep over 1 year old	11.6	6.4
Sheep under 1 year old	7.7	4.3
Mink	4.3	1.7
Foxes	9.0	3.6
Deer	12.0	5.4
Reindeer	6.0	2.7

¹ Includes pasture.

² Factors for excreted nitrogen apply for the whole life time of animals, and nitrogen is calculated when animals are slaughtered/replaced.

³³ Per animal. For these categories, life time is less than a year. This means that the number of animals bred in a year is higher than the number of stalls (pens).

Source: Karlengen et al. (2012), Aspeholen Åby et al (2019), and estimations by Statistics Norway.

NH₃

Ammonia volatilised from manure storage represents, together with NO_x, the activity data in the estimations of indirect N₂O emissions from atmospheric deposition, source 3Bb5 (atmospheric deposition from manure storage). A model based on the stepwise approach proposed by the EEA (2016) is used for calculating the emissions of ammonia from manure management. The principle of the model is illustrated in Figure 5.2.




Figure 5.2 The principle of the revised nitrogen model

The revised nitrogen model closely follows the stepwise approach proposed by the EEA (2016) guidelines, with all the 15 steps proposed in the former being followed in the Norwegian model. Emissions of NH_3 are determined from buildings and yards and from manure storage systems. Total NH_3 emissions from manure management are estimated by multiplying the amount of manure nitrogen (ammonium N) by the different emission factors for the housing and storage systems, taking into account the effect of any abatement measures and improved practices. The amount of ammonium nitrogen in the manure is estimated by the number of animals and ammonium nitrogen excretion factors for each type of animal (see Table 5.12).

NO_x

Nitrous oxide volatilised from manure storage represents, together with NH_3 , the activity data in estimations of indirect N_2O emissions from atmospheric deposition, source 3Bb5 (atmospheric deposition from manure storage).

NO_x emissions from manure management systems are estimated according to the methodology in EEA (2016). In Norway, all animal excreta that are not deposited during grazing are managed as manure. Norwegian values for N in excreta from different animals according to Table 5.12 are used. Norwegian values are also used for the fraction of total excretion per species for each management system (MS) and for pasture, see Table 5.14 and Table 5.15. The fractions are updated every year.

Indirect N₂O from manure management, 3Bb5

Deposition of nitrogen from manure management is assumed to correspond to the amount of nitrogen in the NH₃ and NO_x that volatilises from manure storage systems. The N₂O emissions are calculated by multiplying the amount of N from deposition with the IPCC default emission factor of 0.01 kg N₂O-N/(kg NH₃-N+NO_x-N volatilized) (IPCC 2006).

Storage systems that are not watertight may cause leaching of manure nitrogen. It is assumed that leaching occurs from the storage systems solid storage (not including dry manure in manure cellars), cattle and swine deep bedding, dry lot and poultry manure. The fractions that are assumed leached are based on expert judgement, see Table 5.13.

Table 5.13 FraC_{leach} for storage systems that are assumed to have leaching.

	FraC _{leach} , %
Solid storage ¹	25
Cattle and swine deep bedding	15
Dry lot	25
Poultry manure	25

¹ Solid manure stored in manure cellars are not assumed to have leaching.

Source: Expert judgement by Statistics Norway and the Norwegian Agricultural Agency²¹.

The IPCC default emission factor of 0.0075 kg N₂O-N/kg N lost to leaching/runoff is used (IPCC 2006).

5.5.1.2 Activity data

CH₄, N₂O, NO_x and NH₃

Emissions are estimated from the animal population. How the animal population is estimated is described in Section 5.2 and Annex IX.

Surveys for assessing use of manure management systems (MMS) have been carried out in 2000, 2003 and 2013.

In addition to collecting data on the MMS, the surveys aim to determine the fraction of manure from each animal category that is deposited in pastures as opposed to in animal housing, which is summarized in Table 5.14.

Norway has developed a single model for estimating N₂O, NH₃ and NO_x emissions, which replaces two separate calculation models which were used for previous years. The previous models treated the basic data from the 2013 Manure Survey differently for estimating NH₃ emissions compared to

²¹ Email from Jon Magnar Haugen, the Norwegian Agricultural Agency, 25.11.2015.

N₂O and NO_x emissions due to different manure storage categories being used. For the development of a single model, manure quantities used in the previous N₂O model have had to be reassigned between different types of manure storage systems, in addition to re-categorization of manure types between slurry, FYM and solid manure. The fraction of total excretion per animal category for each management system is presented in Table 5.15.

Table 5.14 Percent of total excretion per species processed by a MMS (i.e. deposited in housing) and deposited on pasture. 2017

	% manure to pasture	% manure to MMS
Dairy cattle	16 %	84 %
Beef cows	31 %	69 %
Youngcattle	31 %	69 %
Swine	0 %	100 %
Laying hens	0 %	100 %
Broilers	0 %	100 %
Turkeys	0 %	100 %
Other poultry	0 %	100 %
Horses	26 %	74 %
Goats	37 %	63 %
Sheep	62 %	38 %
Fur animals	0 %	100 %
Deer	100 %	0 %
Reindeer	100 %	0 %

Source: Data for storage systems from Statistics Norway (Gundersen & Heldal 2015), data for pasture times from (TINE BA Annually) (Dairy cattle, goat), Statistics Norway's Sample Survey 2001 (Statistics Norway 2002)

Table 5.15 Fraction of total excretion per animal category for each management system used in the estimations of NH₃ and NO_x. 2017.

	In-house slurry pit [pit storage below animal confinements]	Tank without cover [Liquid/slurry]	Tank with cover [Liquid/slurry]	Heaps [solid storage]	In-house deep litter [Cattle and swine deep bedding]	Dry lot
Dairy cattle	0.76	0.00	0.24	0.00	0.00	0.00
Beef cows	0.53	0.00	0.17	0.14	0.12	0.04
Young cattle	0.74	0.00	0.19	0.03	0.03	0.01
Swine	0.64	0.20	0.11	0.03	0.02	0.00
Laying hens	0.38	0.00	0.00	0.62	0.00	0.00
Broilers	0.38	0.00	0.00	0.62	0.00	0.00
Turkeys	0.38	0.00	0.00	0.62	0.00	0.00
Other poultry	0.38	0.00	0.00	0.62	0.00	0.00
Horses	0.51	0.00	0.00	0.39	0.05	0.04
Goats	0.51	0.00	0.00	0.39	0.05	0.04
Sheep	0.75	0.01	0.00	0.07	0.14	0.03
Fur animals	0.32	0.00	0.00	0.68	0.00	0.00
Deer	NA	NA	NA	NA	NA	NA
Reindeer	NA	NA	NA	NA	NA	NA

Source: Data for storage systems from Statistics Norway (Gundersen & Heldal 2015)

Data on storage systems for years other than those covered by the manure survey is not available. Separate estimations of the effects on emissions of the assumed changes in storage systems since 1990 show that these assumed changes do not significantly impact the emissions. For the intermediate years 2004-2012 between the surveys of 2003 and 2013, the distribution of management system has been estimated using a linear interpolation of changes between 2003 and 2013, for each system. The surveys on management systems do not include pasture. Data for pasture times for dairy cattle and dairy goat has however been annually updated in the Cow Recording System (TINE BA Annually) until 2013, while for the other animals, data from Sample survey of agriculture and forestry for 2001 at Statistics Norway (2002) is used.

In the CH₄ estimations, the share of the manure stored more respective less than one month in pit storage below animal confinement before spreading is based on expert judgement²². It is assumed that 1/6 of the manure is stored less than 1 month, the rest more than 1 month.

In the manure surveys of 2000 and 2013, the manure of each management system is distributed by all combinations of the following regions and productions:

Regions:

- South-Eastern Norway
- Hedmark and Oppland
- Rogaland
- Western Norway

²² Personal communication John Morken, NMBU, 6.8.2014

- Trøndelag
- Northern Norway

Production²³:

- Cattle
- Swine
- Sheep
- Goats and horses
- Poultry

5.5.1.3 Emission factors

CH₄

The calculated average emission factors for different animal types are shown in Table 5.16 and Table 5.17. Except for sheep and goats, they are country specific factors which may deviate from the IPCC default values.

Table 5.16 CH₄ emission factors for manure management used in the IPCC tier 1 method. kg/animal/year.

	Emission factor ¹	Source
Sheep > 1 year	0.19	IPCC (2006)
Sheep < 1 year	0.19	IPCC (2006)
Dairy goats	0.13	IPCC (2006)
Other goats	0.13	IPCC (2006)
Horses	2.95	(Karlengen et al. 2012)
Mink, males	0.27	(Karlengen et al. 2012)
Mink, females	0.54	(Karlengen et al. 2012)
Fox, males	0.43	(Karlengen et al. 2012)
Fox, females	0.87	(Karlengen et al. 2012)
Reindeer	0.36	(Karlengen et al. 2012)
Deer	0.9	(Karlengen et al. 2012)

¹ Includes pasture.

²³ The grouping of animals is different in the two surveys. Cattle is one category in the 2000 survey and three categories in the 2013 survey. Goats are grouped with sheep in the 2000 survey, but with horses in the 2013 survey. Horses are grouped with other animals in the 2000 survey. Fur bearing animals are not included in the 2013 survey

Table 5.17 Average CH₄ emission factors for manure management used in the IPCC tier 2 method.
kg/animal/year. 2017

	Emission factor ¹
Dairy cows	22.61
Bulls ²	6.93
Heifers ² for slaughter	7.29
Heifers for breeding ²	9.20
Beef cows (mature non-dairy cattle)	13.11
Sows	6.34
Piglets	0.38
Young pigs for breeding	2.33
Pigs for slaughter ³	2.71
Hens	0.046
Chicks bred for laying hens	0.01
Chicks for slaughter ³	0.013
Ducks for breeding	0.098
Ducks for slaughter ³	0.026
Turkey and goose for breeding	0.098
Turkey and goose for slaughter ³	0.056

¹ Includes pasture.

² Factors apply for the whole life time of animals.

³ Per animal place. This means that the factor includes all animals bred in on place (pen) during the year

Source: Karlengen et al. (2012), IPCC (2006), Morken et al. (2013), Aspehølen Åby et al (2019) and estimations by Statistics Norway.

N₂O

The IPCC default values for N₂O emission factors from manure management are used (Table 5.18). These are consistent with the 2006 IPCC Guidelines (IPCC 2006).

Table 5.18 Emission factors for direct N₂O emissions from manure management per manure management system

Manure management system	Emission factor, kg N ₂ O-N/kg N excreted
Pit storage below animal confinement	0.002
Liquid / slurry without cover	0
Liquid / slurry with cover	0.005
Solid storage	0.005
Dry lot	0.02
Cattle and swine deep bedding	0.01
Dry lot	0.02
Poultry manure	0.001
Pasture range and paddock (cattle, pigs and poultry)	0.02
Pasture range and paddock (other animals)	0.01

Source: IPCC (2006)

NH₃

Emission factors vary with production and storage system; in the model there is no variation between regions for the manure management systems. The factors used are shown in Table 5.19. All emission factors in Table 5.19 are sourced from EEA (2016), since measurements of NH₃ losses in animal housing and manure storage have so far not been carried out in Norway.

Table 5.19 NH₃ emissions factors for various storage systems and productions. % losses of N of ammonium N.

	Housing		Storage	
	Slurry	Solid manure	Slurry	Solid
Dairy cattle	20 %	19 %	20 %	27%
Suckling cows	20 %	19 %	20 %	27%
Young cattle	20 %	19 %	20 %	27%
Swine	28 %	27 %	14 %	45%
Laying hens	41 %	41 %	14 %	14 %
Broilers	28 %	28 %	17 %	17 %
Turkeys	35 %	35 %	24 %	24 %
Other poultry	57 %	57 %	24 %	24 %
Horses	22 %	22 %	35 %	35 %
Goats	22 %	22 %	28 %	28 %
Sheep	22 %	22 %	28 %	28 %
Fur animals	27 %	27 %	9 %	9 %
Deer	20 %	20 %	20 %	27%
Reindeer	20 %	20 %	20 %	27%

Source: EMEP/EEA air pollutant emission inventory Guidebook 2016: 3.B Manure management (EEA 2016).

As recommended in EEA (2016), the effect of the abatement measures and improved practices are described using a reduction factor, i.e. a proportional reduction in the emission estimate for the unabated situation. These reduction factors are summarized in Table 5.20.

Table 5.20 NH₃ abatement measures for manure management systems and corresponding emission reduction, % applied to NH₃ emissions factors presented in Table 5.19

Abatement measure	Housing	Storage			
	In-house slurry pit, below slatted floor	Tank with cover, "tight" lid	Tank with cover, plastic sheeting (floating cover)	Natural crust	"Low technology" floating cover
NH ₃ emission reduction, %	50	80	60	40	40

Source: Rösemann et al. (2017), UNECE (2014).

To estimate losses, these unabated or amended emission factors are in turn multiplied with the amount of manure (based on number of animals and N-factors per animal) entering each of the different housing or storage category, as determined by the Statistics Norway survey of different

storage systems in 2000 (Gundersen & Rognstad 2001), the Sample survey of agriculture and forestry 2003 (Statistics Norway 2004) and data from the report *Use of inorganic and organic fertilisers in agriculture 2013* (Gundersen & Heldal 2015). The changes in storage systems from 2003 to 2013 have been linearly interpolated in the intermediate years. From 1990-2002 and from 2013, the number of animals is the only activity data that differs from year to year.

NO_x

The emissions factors used for NO_x emissions in manure management systems are shown in Table 5.20.

Table 5.20 NO_x emission factors for manure management per manure management system.

	EF, proportion of TAN
Slurry storage	0.0001
Solid storage	0.01

Source:EEA (2016)

N excretions is estimated as total ammoniacal nitrogen (TAN), which is the same N excretion factor that is used in the estimations of NH₃ from manure management systems.

5.5.2 Uncertainties and time-series consistency

Uncertainties estimates are provided in Annex II.

Activity data

CH₄

The data for the number of animals is considered to be known within ± 5 %. Other activity data are the different kinds of manure treatment (which will determine the emission factor), which have been assessed by expert judgments. This will contribute to the uncertainty.

N₂O and NH₃

The data for the number of animals is considered to be known within ± 5 %.

For the emissions from manure management, Norwegian data for N in excreta is used (Table 5.12). The nitrogen excretion factors are uncertain, but the range is considered to be within ± 15 % (Rypdal 1999). The uncertainty has not been estimated for the revised nitrogen excretion factors from Karlengen et al (2012), and in the key category analysis is the uncertainty estimate for the country specific nitrogen excretion factors from 1999 still used as the best available estimate. This can be considered as a conservative estimate of the uncertainty since it is expected that the new nitrogen excretion factors have a lower uncertainty. The uncertainty is connected to differences in excretion between farms in different parts of the country, the fact that the survey farms may not have been representative, general measurement uncertainty and the fact that fodder and fodder practices have changed since the factors were determined.

There is also an uncertainty connected to the division between different storage systems for manure, which is considered to be within ± 10 %, and the division between storage and pasture, which is considered to be within ± 15 %.

Emission factors**CH₄**

The emission factors are considered to have the uncertainty range ± 20 % for cattle, poultry and pigs (Tier 2) and ± 30 % for other animals (Tier 1) (IPCC 2006).

N₂O

For the emission of N₂O from different storage systems, IPCC default emission factors are used. They have an uncertainty range of a factor of 2 (IPCC 2006).

NH₃

All emission factors for NH₃ which have been used for both housing and storage are sourced from EEA (2016). As stated in EEA (2016), uncertainties with regard to NH₃ EFs vary considerably. EEA (2016) concludes that the overall uncertainty for the United Kingdom NH₃ emissions inventory, as calculated using a Tier 3 approach, was ± 21 % (Webb and Misselbrook, 2004), while that for the Netherlands, also calculated using a Tier 3 approach, was ± 17 % (van Gijlswijk et al., 2004).

NO_x

Uncertainty has not been assessed. According to EEA (2016), uncertainty is high, ranging between -50 % to + 100 %.

5.5.3 Category specific QA/QC and verification

In 2011, the Norwegian University of Life Sciences (NMBU) published a comparison of the methodologies used for calculating CH₄ emissions from manure management in Sweden, Finland, Denmark and Norway (Morken & Hoem 2011).

In a project in 2012 at the Norwegian University of Life Sciences (NMBU) that updated the Norwegian nitrogen excretion factors and the values for manure excreted for the different animal species, comparisons were made with the corresponding factors used in Sweden, Denmark and Finland and with IPCC default factors as a verification of the Norwegian factors (Karlengen et al. 2012).

A project with the aim to revise the Norwegian CH₄ conversion factors (MCF) for the manure storage systems in use was conducted at the Norwegian University of Life Sciences (NMBU) in 2013. The maximum CH₄ producing capacity (B₀) was also revised for cattle manure.

The methodology for estimating methane from manure management was revised in the 2014 submission. The emissions of methane from manure for cattle, pigs and poultry were estimated with tier 2 method in accordance with IPCC GPG (IPCC 2000). The population of animals was brought into consistency with the animal data used elsewhere in the inventory.

In 2014, a new manure survey for 2013 was carried out by Statistics Norway (Gundersen & Heldal 2015). The results are implemented in the estimations of CH₄ and N₂O emissions from manure. Statistics Norway's detailed manure survey gave more extended activity data which is better related to emission source categories, for manure management and spreading.

As part of the revision of the nitrogen model undertaken in 2018, a review was undertaken of the Norwegian emissions factors compared to those used in other Nordic and Northern European countries. The review demonstrated that the Norwegian EFs compare well with those used in other Nordic and Northern European countries. In the majority of cases the EFs used in the Norwegian model lie within the range of EFs used by the example countries, and in most cases are at the upper

end of the range which demonstrates the conservative approach taken in the absence of country specific EFs.

5.5.4 Category-specific recalculations

N₂O emissions from manure management are based on estimations from the new model described section 5.5.1.1. This model is made for calculating the emissions of ammonia and other nitrogen species from manure management. The main changes is the inclusion of added N in animal bedding and the consequent immobilization of TAN in that bedding. The model also introduces a wider number of options for storage of manure which introduces updated emission factors. These factors are in general higher, but give a more accurate estimate of emissions for storage.

For CH₄ from manure management the different options for manure storage were also integrated in the model for estimating CH₄ emissions. The new VS factor for beef cattle also lead to higher CH₄ emissions for these animal group. See Chapter 10 for more details.

5.5.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5.6 Direct and indirect N₂O emissions from agricultural soils - 3D (Key categories for N₂O)

5.6.1 Category description

The emissions of N₂O from agricultural soils in Norway in 2017 amounted to 1.6 Mtonnes calculated in CO₂-equivalents. They accounted for about 36 % of the total Norwegian GHG emissions from agriculture in 2017.

The emissions had minor fluctuations in the period 1990-2017. During the period 1990-2017, emissions decreased by 3 %. From 2016 to 2017, the emissions decreased by 1 %.

Table 5.21 Emission trends for 3D Direct and indirect N₂O emissions from agricultural soils (kt CO₂ equivalents).

Source category	1990	1990, % of Agriculture	2016	2017	2017, % of Agriculture	Trend 1990-2017 (%)	Trend 2016-2017 (%)
3Da1. Inorganic N fertilizers	516	11 %	479	460	10 %	-11 %	-4 %
3Da2. Organic N fertilizers	236	5 %	276	277	6 %	17 %	1 %
3Da3. Urine and dung deposited by grazing animals	195	4 %	185	188	4 %	-4 %	2 %
3Da4. Crop residues	105	2 %	74	72	2 %	-31 %	-3 %
3Da6. Cultivation of organic soils	363	8 %	376	375	8 %	3 %	0 %
3Db1. Indirect N ₂ O Emissions from managed soils - Atmospheric deposition	96	2 %	96	96	2 %	-0,3 %	0 %
3Db2. Indirect N ₂ O Emissions from managed soils - Nitrogen leaching and run-off	161	3 %	156	153	3 %	-5 %	-2 %
3D. Total	1672	36 %	1642	1622	36 %	-3 %	-1 %

Source: Statistics Norway and Norwegian Environment Agency

Different sources of N₂O from agricultural soils are distinguished in the IPCC methodology, namely:

- Direct emissions from agricultural soils (from use of synthetic fertilisers, animal excreta nitrogen, sewage sludge and other organic fertilisers applied to soils, droppings from grazing animals, crop residues and cultivation of soils with a high organic content);
- N₂O emissions indirectly induced by agricultural activities (N losses by volatilization, leaching and run-off).

The use of synthetic fertilisers, animal excreta nitrogen and sewage sludge used as fertiliser, and other organic fertilisers applied to soils also results in emissions of NH₃ and NO_x that gives indirect N₂O emissions from volatilization. NH₃ volatilized from grazing animals are also included in the estimations of indirect N₂O.

Section 5.3 gives more information about nitrogen in animal manure as basis for emission estimates and an overview of the manure nitrogen flows in the Norwegian greenhouse gas inventory.

Emissions of N₂O from agricultural soils are key categories because of uncertainty, both in level and trend.

5.6.1.1 Inorganic N fertilisers, 3Da1

Methodological issues

N₂O

IPCC Tier 1 methodologies and default emission factors (IPCC 2006) are used for estimating direct N₂O emissions from managed soils.

The direct emissions of N₂O from use of synthetic fertilisers are calculated from data on total annual amount of fertiliser sold in Norway and its nitrogen content, subtracting the amount of synthetic fertiliser applied in forest. The resulting amount that is applied on agricultural fields is multiplied with the IPCC default emission factor (IPCC 2006).

NH₃

The calculations of NH₃ emissions from the use of synthetic fertiliser are based on the amounts of nitrogen supplied and emission factors for the percentage of nitrogen emitted as NH₃ during spreading.

NO_x

The sum of all nitrogen applied to soil has been multiplied with the default tier 1 emission factor to estimate the nitric oxide (NO) emission from crop production. Thereafter the amount of NO is translated to amount of NO₂.

Activity data

N₂O

The Norwegian Food Safety Authority calculates a total value for annual consumption of synthetic fertilisers in Norway based on sale figures (Norwegian Food Safety Authority Annually). This data is corrected for the amount fertiliser used in forests which is provided yearly by the Norwegian Institute of Bioeconomy Research (NIBIO).

The calculation of emissions from use of nitrogen fertiliser is based on sales figures for each year. A strong price increase for nitrogen fertiliser caused a stock building in 2008 and corresponding lower sales in 2009. In addition, new fertilisation standards may have brought about a reduction of the use of fertilisers. To correct for this, a transfer of fertiliser use has been made from 2008 to 2009.

NH₃

For the calculation of the emission of NH₃, the amount of total N in the fertilizer is used (the same as for estimating N₂O). For the calculation of the emission of NH₃, we need a specification of the use of different types of synthetic fertiliser since the NH₃ emission factor vary between the different types. This is given by the Norwegian Food Safety Authority for the years from 2000. Due to lack of data for the years before 2000, we have to assume that the percentual distribution between the usage of different fertiliser types is the same as in 1994 for these years.

NO_x

For the calculation of the emission of NO_x, the amount of total N in the fertilizer is used (the same as for estimating N₂O).

Emission factors

N₂O

The IPCC default emission factor of 0.01 kg N₂O-N/kg N applied (IPCC 2006) has been used.

NH₃

The percentage of nitrogen emitted as NH₃ during spreading ($\text{frac}_{\text{gasf}}$) are used as emission factor. More information about the calculation of $\text{frac}_{\text{gasf}}$ and the NH₃ emission factors (g NH₃/kg N applied) for the different types of fertilisers is provided in Annex IX, section 4.

NO_x

The tier 1 default emission factor of 0.04 kg NO₂/ kg fertilizer-N applied (EEA 2016) has been used.

5.6.1.2 Animal manure applied to soils, 3Da2

Methodological issues

N₂O

IPCC Tier 1 methodologies and default emission factors (IPCC 2006) are used for estimating direct N₂O emissions from animal manure applied to managed soils.

In Norway, all animal excreta that are not deposited during grazing are used as manure and applied to soils. Further, it is assumed that animals do not emit N₂O themselves. NH₃ emissions in housing and storage, and N₂O emissions in housing and storage and manure application are all estimated individually and the emission estimates are based on the same nitrogen pool.

The emission of N₂O from manure used as fertiliser is calculated by multiplying the total amount of N in manure used as fertiliser with the IPCC default emission factor (IPCC 2006). This amount is equivalent to total N excreted from the animals deducted for the amount dropped during grazing.

NH₃

NH₃ emissions from spreading of manure depend on several factors, e.g. climate and time of spreading of manure, type of cultivation and cultivation practices and characteristics of the soil.

Emissions of ammonia are calculated for spreading of manure on cultivated fields and meadow. The total amount of manure nitrogen that is spread is estimated by the number of animals and nitrogen excretion factors for each type of animal, and is thereafter distributed on different spreading methods based on national data. The nitrogen basis for the estimated amounts of nitrogen that volatilises as NH₃ during spreading takes into account the amount of nitrogen in the NH₃, NO_x, and N₂ that volatilises during housing and storage, as well as the N lost as N₂O and leaching during storage. Total emissions from spreading are estimated by emission factors for each different spreading method used multiplied by the amount of manure nitrogen spread with the respective method.

NO_x

The sum of all nitrogen applied to soil has been multiplied with the default tier 1 emission factor to estimate the nitric oxide emission from crop production. Thereafter the amount of NO is translated to amount of NO₂.

Activity data**N₂O, NH₃ and NO_x**

The amount of N in manure systems is calculated as total N in manure adjusted for the N that is dropped on pastures.

Emissions of N₂O, NH₃ and NO_x during spreading are calculated from the amounts of N excreted minus losses due to housing and storage emissions.

There are several sources of activity data on spreading of manure. The main sources are manure surveys performed in 2000 and in 2013 by Statistics Norway (Gundersen & Rognstad 2001) and (Gundersen & Heldal 2015), various sample surveys of agriculture and forestry 1990-2007 and the annual animal population.

Data from the manure survey only exists for 2000 and 2013, while the data from the sample surveys has been updated for several, but not all, years. The manner of spreading the manure affects the NH₃ emissions estimates, while the N₂O emission estimations are insensitive to methods of spreading.

How the amount of nitrogen in animal manure is estimated is further described in section 5.3 and 5.5.1.1. How the animal population is estimated is described in Section 5.2 and Annex IX.

Nitrogen factor are estimated by Karlengen et al (2012). In the estimations of NH₃ losses, the factors of N excretion correspond to the estimated nitrogen excreted in the urine.

Table 5.22 Sources for activity data for emissions from animal manure applied to soils.

	Sources
Number of animals	Statistics Norway (applications for productions subsidies, no. and weight of approved carcasses), The Cow Recording System (TINE BA Annually)
Nitrogen factors for manure, Annex IX, section 3.	Karlengen et al (2012), various sources, compiled by Statistics Norway
Distribution between manure storage systems	Sample Survey of agriculture and forestry 2003 (Statistics Norway 2004), manure survey in 2000 and 2013 (Gundersen & Rognstad 2001) and (Gundersen & Heldal 2015)
Pasture times for different animal categories	TINE BA (Annually) (Dairy cows, goat), Statistics Norway's Sample Survey 2001 (Statistics Norway 2002) (non-dairy cattle, sheep), expert judgements.

Table 5.23 Parameters included in the estimation of NH_3 emissions from manure.

	Sources
Area where manure is spread, split on cultivated field and meadow	Statistics Norway (Sample Surveys of Agriculture, various years), (Gundersen & Rognstad 2001) (Gundersen & Heldal 2015)
Area and amount where manure is spread, split on spring and autumn	Gundersen and Rognstad (2001) and Gundersen and Heldal (2015)
Addition of water to manure	(Gundersen & Rognstad 2001), and Gundersen and Heldal (2015), expert judgements, Statistics Norway's Sample Survey 2006 (Statistics Norway 2007)
Spreading techniques	(Gundersen & Rognstad 2001), Gundersen and Heldal (2015), expert judgements
Usage and time of harrowing and ploughing	(Gundersen & Rognstad 2001), Gundersen and Heldal (2015), expert judgements, Statistics Norway's Sample Surveys of Agriculture

Emission factors N_2O

The IPCC default emission factor of 0.01 kg N_2O -N/kg N applied (IPCC 2006) has been used.

 NH_3

Emission factors for spreading of stored manure vary with spreading method (Gundersen & Rognstad 2001; Gundersen & Heldal 2015), water contents (Statistics Norway 2007), type and time of treatment of soil (Gundersen & Rognstad 2001; Gundersen & Heldal 2015), time of year of spreading (Gundersen & Rognstad 2001; Gundersen & Heldal 2015; Statistics Norway 2007), cultivation and region.

The basic factors used are shown in Table 5.24.

Table 5.24 Emissions factors for $\text{NH}_3\text{-N}$ for various methods of spreading of manure. % of ammonium N.

Meadow			Spring	Summer	Autumn
kg NH ₃ -N/kg TAN					
Spreading method	Added water				
Broadcast spreading	< 100%		0.4	0.7	0.7
	> 100%		0.24	0.35	0.35
Trailing hose	< 100%		0.3	0.5	0.4
	> 100%		0.18	0.25	0.2
Injection			0.15	0.30	0.05
Dry manure			0.7	0.9	0.7
Arable land		Incorporation time	Spring	Summer	Autumn
kg NH3-N/kg TAN					
Spreading method	Added water	Hours			
Broadcast spreading	< 100%	0-1	0.08	0.08	0.12
		1-4	0.20	0.20	0.30
		4-12	0.33	0.33	0.45
		12+	0.50	0.50	0.45
	> 100%	0-1	0.04	0.04	0.06
		1-4	0.10	0.10	0.15
		4-12	0.17	0.17	0.28
		12+	0.25	0.25	0.28
Trailing hose	< 100%	0-1	0.03	0.03	0.05
		1-4	0.12	0.12	0.17
		4-12	0.23	0.23	0.35
		12+	0.50	0.50	0.45
	> 100%	0-1	0.02	0.02	0.02
		1-4	0.06	0.06	0.09
		4-12	0.12	0.12	0.22
		12+	0.25	0.25	0.28
Dry manure			0.70	0.70	0.70

Source: Karlsson S. and Rodhe L. (2002), Morken and Nesheim (2004), Rösemann et al. (2017).

The factors in Table 5.24 are combined with data from the Sample survey of agriculture and forestry 2006 (Statistics Norway 2007) and a time series on mixture of water in manure. Emission factors for NH_3 emissions from spreading of manure are connected to activity data that is updated for the whole time series when new information is available, i.e. number of animals (amount of manure), time of spreading and type of cultivation of the areas where the manure is spread.

NO_x

The tier 1 default emission factor of 0.04 kg NO_2 /kg fertilizer-N applied ((EEA 2016)) has been used.

5.6.1.3 Sewage sludge applied to soils, 3Da2

Methodological issues

N_2O

IPCC Tier 1 methodologies and default emission factors (IPCC 2006) are used for estimating direct N_2O emissions from managed soils.

Data for the N_2O emission arising from sewage sludge applied on fields has been calculated by multiplying the amount of nitrate in the sewage sludge applied with the IPCC default emission factor.

NH_3

To calculate NH_3 emissions from sewage sludge used as fertiliser, the total N in sewage sludge is multiplied by $\text{frac}_{\text{gas}} \cdot \text{Frac}_{\text{gas}}$. Frac_{gas} is volatilised N in animal manure applied and dung and urine deposited by grazing animals as fraction of total N in the manure applied and dung and urine deposited (kg $\text{NH}_3\text{-N} + \text{NO}_x\text{-N}$)/(kg N applied or deposited).

NO_x

The sum of all nitrogen applied to soil has been multiplied with the default tier 1 emission factor to estimate the nitric oxide emission from crop production. Thereafter the amount of NO is translated to amount of NO_2 .

Activity data

N_2O , NH_3 and NO_x

Statistics Norway (waste water statistics) annually gives values for the amount of sewage sludge, and the fraction of the sewage sludge that are applied on fields. The N-content in the sludge is given in Statistics Norway (2001), and the same value of 2.82 % is used for all years.

Emission factors

N_2O

The IPCC default emission factor of 0.01 kg $\text{N}_2\text{O-N}$ /kg N applied (IPCC 2006) has been used.

NH_3

The estimated national volatilization fractions (frac_{gas}) have differed between 14-15 % since 1990.

NO_x

The tier 1 default emission factor of 0.04 kg NO_2 /kg fertilizer-N applied (EEA 2016) has been used.

5.6.1.4 Other organic fertilizers applied to soils, 3Da2

Methodological issues

N₂O

Emissions of N₂O from other organic fertilisers applied to soils are estimated by multiplying estimated amounts of N in organic fertilisers with the IPCC default emission factor for N applied to agricultural soils.

NH₃

Emissions of NH₃ from other organic fertilisers applied to soils are estimated by multiplying estimated amounts of N in organic fertilisers with the $\text{frac}_{\text{gas}}^{\text{N}}$ -factor. $\text{frac}_{\text{gas}}^{\text{N}}$ is volatilised N in animal manure applied and dung and urine deposited by grazing animals as fraction of total N in the manure applied and dung and urine deposited (kg NH₃-N + NO_x-N)/(kg N applied or deposited)

NO_x

The amount of nitrogen in other organic fertilisers applied to soil has been multiplied with the default tier 1 emission factor to estimate the nitric oxide emission from crop production. Thereafter the amount of NO is translated to amount of NO₂.

Activity data

N₂O, NH₃ and NO_x

The annual amount of nitrogen in other organic fertilisers applied in agriculture during the period 1990-2013 was assessed in 2014 (Aquateam COWI AS 2014). Other organic fertilizer consists of compost and organic fertilizer from composting and biogas plants based on food waste, in addition of meat and bone meal (PAP) and sludge from land based aquaculture facility in agriculture. This was a practically non-existent source of nitrogen before 2000. Since then, it has varied over the years. One important reason for the inter-annual variations is changes in regulations for the usage of meat and bone meal as fertilizer on agriculture land, which has resulted in significant inter-annual variations.

This is a small emission source, emissions in 2015 were estimated to less than 0.01 ktonnes N₂O, or approx. 0.005 % of total GHG emissions.

Emission factors

N₂O

The IPCC default emission factor of 0.01 kg N₂O-N/kg N applied (IPCC 2006) has been used.

NH₃

The estimated national volatilization fractions have differed between 14-15 % since 1990.

NO_x

The tier 1 default emission factor of 0.04 kg NO₂/kg fertilizer-N applied (EEA 2016) has been used.

5.6.1.5 Urine and dung deposited by grazing animals, 3Da3

Methodological issues

N₂O

The fraction of the total amount of animal manure produced that is droppings on pastures is given by national data for the distribution of manure to different storage systems and data for pasture times

(Table 5.14). The amount of N deposited during grazing is multiplied with the IPCC default emission factor (IPCC 2006).

NH₃

Animal population data, data for pasture times, and factors for the nitrogen amount in excreta for different animal categories give the nitrogen amounts for the animal categories on pastures. Specific emission factors by animal category are used.

Activity data

N₂O and NH₃

Data for time on pasture and share of animals on pasture are collected from the Sample Survey in Statistics Norway 2001 (Statistics Norway 2002) and from TINE BA (TINE BA Annually). The data from TINE BA comprises pasture data for goats and milking cows. All other pasture data is from the Sample survey 2001 (Statistics Norway 2002).

How the amount of nitrogen in animal manure and the fraction of nitrogen amount that goes to pasture is estimated is further described in section 5.3 and 5.5.1. How the animal population is estimated is described in Section 5.2 and Annex IX.

Emission factors

N₂O

The emissions of N₂O from animals on pastures are calculated using the IPCC default emission factors of 0.02 kg N₂O-N/kg N for cattle, poultry and pigs, and of 0.01 kg N₂O-N/kg N for other animal groups (IPCC 2006).

NH₃

The emission factors used for the calculation of the NH₃ emissions from grazing animals are shown in Table 5.25.

These are the same as the emission factors recommended in EEA (2016).

Table 5.25 Ammonia emission factors from droppings from grazing animals on pasture. % of TAN.

	NH ₃ loss, % of TAN
Dairy cattle	10 %
Suckling cows	10 %
Young cattle	6 %
Swine	25 %
Horses	35 %
Goats	9 %
Sheep	9 %
Fur animals	9 %
Deer	10 %
Reindeer	10 %

Source: EEA (2002). For deer and reindeer EF for dairy cattle used

5.6.1.6 N₂O from crop residues, 3Da4

Methodological issues

N₂O emissions associated with crop residue decomposition are estimated using the IPCC tier 1 approach (IPCC 2006) but with some national factors. Some country specific factors are given for fraction of dry matter, fraction of total area that is renewed annually, ratio of above-ground and below ground residues to harvested yield, N content of above-ground and below-ground residues and fraction of above ground residues removed from the field. The national factors are documented in Grønlund et al. (2014). In the development of national factors, residues from perennial grass and grass-clover mixtures were prioritized, in addition to the cereal species; wheat, barley and oats, which combined constitute about 85 percent of the total agricultural crop residues. For other productions, the IPCC default factors (IPCC 2006) are assumed to be sufficiently representative.

The factors were calculated from the sale statistics for clover seeds, area statistics of meadows of different age classes, area statistics of renewed meadow, and research results on clover and N content in meadow, and yield and N content of straw in Norway.

Based on area statistics on renewed meadows the $Frac_{Renew}$ has been estimated to 0.1.

About 75 percent of the meadows have been renewed with a mixture of grass and clover seeds, but only about 55 percent of 1 and 2 year old meadow areas can be considered as grass-clover mixtures with more than 5 percent clover. The mean clover share in the grass-clover mixtures has been estimated to about 20 percent. The clover share is lower in older meadow, but the content in the first years is more representative for the total crop residues produced during the lifetime of the meadow.

Above-ground crop residues contain both leaves and stubbles, while below ground residues are assumed to contain only roots. The N contents of above-ground and below-ground crop residues (N_{AG} and N_{BG}) have been estimated to 0.015 and 0.011 respectively for meadow without clover and 0.019 and 0.016 respectively for meadow with 20 percent clover share. A possible higher clover share in the beginning of the 1990s has not had a significant influence on N fractions of grass-clover mix in meadows.

Straw harvested for purposes as feed, beddings and energy ($FRAC_{Remove}$) has been estimated to 0.13 of the total straw production.

For wheat, barley and oats the ratio of above-ground residues (straw) to harvested grain yield (R_{AG}) has been estimated to 0.95, 0.76 and 0.92 respectively, and the N fraction in the straw (N_{AG}) has been estimated to 0.0042, 0.005 and 0.033 respectively (Grønlund et al. 2014). The fraction of crop residue burned on field was updated in 2012 by the Norwegian Agricultural Authorities²⁴. This reduced the fraction for 2011 from 7.5 to 4 %.

$$F_{CR} = \sum_T \left[Crop_{(T)} * Frac_{DM(T)} * (1 - Frac_{BURN(T)}) * Frac_{RENEW(T)} * [R_{AG(T)} * N_{AG(T)} * (1 - Frac_{REMOVE(T)}) + R_{BG(T)} * N_{BG(T)}] \right]$$

F_{CR} = N in crop residue returned to soils (tonnes)

²⁴ Johan Kollerud, Norwegian Agricultural Agency, unpublished material 2012.

$Crop_T$ = Annual crop production of crop (tonnes)

$Frac_{DM}$ = Dry matter content

$Frac_{BURN}$ = Fraction of crop residue burned on field

$Frac_{RENEW}(T)$ = fraction of total area under crop T that is renewed annually

$RAG(T)$ = ratio of above-ground residues dry matter (AGDM(T)) to harvested yield for crop T (kg d.m.)⁻¹,

$NAG(T)$ = N content of above-ground residues for crop T, kg N (kg d.m.)⁻¹

$Frac_{REMOVE}$ = Fraction of crop residue removed for purposes as feed beddings and energy

$RBG(T)$ = ratio of below-ground residues to harvested yield for crop T, kg d.m. (kg d.m.)⁻¹

$NBG(T)$ = N content of below-ground residues for crop T, kg N (kg d.m.)⁻¹

Table 5.26 Factors used for the calculation of the nitrogen content in crop residues returned to soils.

	Share of meadows	$Frac_{DM}$	$Frac_{RENEW}$	R_{AG}	N_{AG}	$Frac_{REMOVE}$	R_{BG}	N_{BG}
Perennial grasses	0.45	0.9	0.1	0.3	0.015	0	1.04	0.011
Grass-clover mixtures	0.55	0.9	0.1	0.3	0.019	0	1.04	0.013
Wheat		0.85	1	0.95	0.0042	0.13	0.47	0.009
Rye		0.85	1	1.1	0.005	0.13	0.46	0.011
Rye wheat		0.85	1	1.09	0.006	0.13		0.009
Barley		0.85	1	0.76	0.005	0.13	0.39	0.014
Oats		0.85	1	0.92	0.0033	0.13	0.48	0.008
Rapeseed		0.85	1	1.1	0.006	0.15	0.46	0.009
Potatoes		0.22	1	0.1	0.019	0	0.22	0.014
Roots for feed		0.22	1	0.1	0.019	0		0.014
Green fodder (non-N fix)		0.9	1	0.3	0.015	0	0.70	0.012
Vegetables		0.22	1	0.1	0.019	0	0.22	0.014
Peas		0.91	1	1.1	0.008	0	0.40	0.008
Beans		0.91	1	1.1	0.008	0	0.40	0.008

Source: Grønlund et al. (2014)

Activity data

As activity data for the estimations of emissions of N₂O from crop residues are annual crop yield statistics from Statistics Norway used.

Emission factors

The IPCC default emission factor of 0.01 kg N₂O-N/kg N applied (IPCC 2006) has been used.

5.6.1.7 N₂O from mineralization/immobilization associated with loss/gain of soil organic matter, 3Da5

Methodological issues

Cropland remaining cropland result in positive SOC stock changes in the mineral soil pool; thus no N₂O emissions are reported from this sub-category. For more information, see NIR Chapter 6.13 Direct N₂O from N mineralization and immobilization – 4(III).

5.6.1.8 N₂O from cultivation of organic soils, 3Da6

Methodological issues

Large N₂O emissions occur as a result of cultivation of organic soils (histosols) due to enhanced mineralization of old, N-rich organic matter (IPCC 2006). The emissions are calculated using the IPCC default emission factors, and an estimation of the area of cultivated organic soil in Norway.

Activity data

The area estimate of cultivated organic soils is given from the Norwegian Institute of Bioeconomy Research (NIBIO) and are consistent with the area used in the LULUCF sector and includes all areas with organic soils of cropland remaining cropland, grassland remaining grassland, land converted to cropland and land converted to grassland. More information about the methodology used for estimation of this area is given in the LULUCF Chapter 6.3.2.

Emission factors

Emissions occurring as a result of cultivation of organic soils are calculated using the IPCC default emission factor of 13 kg N₂O-N/ha per year for cropland and 1.6 kg N₂O-N/ha per year for grassland (IPCC 2014).

5.6.1.9 Indirect N₂O emissions from atmospheric deposition, 3Db1

Methodological issues

Atmospheric deposition of nitrogen compounds fertilises soils and surface waters, and enhances biogenic N₂O formation. Deposition of nitrogen are assumed to correspond to the amounts of NH₃ and NO_x that volatilises during the spreading of synthetic fertilisers, manure, sewage sludge and other organic fertilisers, and NH₃ volatilisation from pastures. The N₂O emissions are calculated by multiplying the amount of N from deposition with the IPCC default emission factor (IPCC 2006).

Activity data

Information about the estimation of NH₃ and NO_x from spreading of synthetic and organic fertilisers and the estimation of NH₃ volatilisation from pastures are given in NIR section 5.6.1.1- 5.6.1.5.

Emission factors

N₂O

The IPCC default emission factor of 0.01 kg N₂O-N/(kg NH₃-N+NO_x-N volatilized) (IPCC 2006) is used to calculate indirect emissions of N₂O from volatilized NH₃ and NO_x.

5.6.1.10 Indirect N₂O emissions from leaching and run-off, 3Db2

Methodological issues

A considerable amount of fertiliser nitrogen is lost from agricultural soils through leaching and run-off. Fertiliser nitrogen in ground water and surface waters enhances biogenic production of N₂O as the nitrogen undergoes nitrification and denitrification. The fraction of the fertiliser and manure nitrogen lost to leaching and surface runoff may vary depending on several factors. A default value of 30 % is proposed (IPCC 2006), but in the Norwegian inventory a national factor of 22 % is used as that is believed to give better results under Norwegian conditions (Bechmann et al. 2012). This estimation was based on data from the Agricultural Environmental monitoring program (JOVA). The overall $\text{Frac}_{\text{leach}}$ estimated in this study was 22 % of the N applied. This value is a median of $\text{Frac}_{\text{leach}}$ for every year during the monitoring period and for each of eight catchments with different production systems. The JOVA-program includes catchment and field study sites representing typical situations in Norwegian agriculture with regard to production system, management, intensity, soil, landscape, region and climate. Data from plot-scale study sites confirmed the level of N leaching from the agricultural areas within the JOVA catchments. The amount of nitrogen lost to leaching is multiplied with the IPCC default emission factor to calculate the emission of N₂O (IPCC 2006).

Nitrogen sources included are inorganic fertilisers, manure, sewage sludge and other organic fertilisers spread on fields, crop residues, and droppings from grazing animals.

Activity data

Information about the estimation of the nitrogen amounts from spreading of synthetic and organic fertilisers and from pastures and crop residues are given in NIR section 5.6.1.6.

Emission factors

N₂O

The IPCC default emission factor of 0.0075 kg N₂O-N/kg N lost to leaching/runoff is used (IPCC 2006).

5.6.2 Uncertainties and time-series consistency

5.6.2.1 Activity data

There are several types of activity data entering the calculation scheme:

Sales of nitrogen fertiliser: The data is based on sales figures during one year (The Norwegian Food Safety Authority). The uncertainty in the sales figures is within ± 5 % (Rypdal & Zhang 2000). In addition, there is a possible additional error due to the fact that sale does not necessarily equal consumption in a particular year due to storage.

Amount of nitrogen in manure: The figures are generated for each animal type, by multiplying the number of animals with a nitrogen excretion factor. The nitrogen excretion factors are uncertain. However, due to monitoring of nitrogen leakage in parts of Norway, the certainty has been improved over time. The range is considered to be within ± 15 % (Rypdal & Zhang 2000). The uncertainty is connected to differences in excreted N between farms in different parts of the country, that the surveyed farms may have not been representative, general measurement uncertainty and the fact that fodder and feeding practices have changed since the factors were determined.

The uncertainty connected to the estimate of the amount of manure is higher than for the amount of synthetic fertiliser used.

Fate of manure: There is significant uncertainty connected to the allocation of manure between what is used as fertiliser and droppings on pastures.

Atmospheric deposition of agricultural NH₃ emissions: The data is based on national figures for NH₃ emission from agriculture. These are within ± 30 % (Rypdal & Zhang 2000)

Leakage of nitrogen: The upper limit for the leakage is the applied nitrogen. The uncertainty is roughly about ± 70 % (Rypdal & Zhang 2000).

5.6.2.2 Emission factors

N₂O

Uncertainty estimates used for the N₂O emission factors are given in Annex II.

NH₃

The uncertainty in the estimate of NH₃ emissions from use of fertiliser is assessed to be about ± 20 % (Rypdal & Zhang 2001). This uncertainty could be lower if better data on fertiliser composition were obtained. The uncertainty is higher for animal manure, ± 30 % (Rypdal & Zhang 2001). This is due to uncertainties in several parameters including fraction of manure left on pastures, amount of manure, conditions of storage, conditions of spreading and climate conditions (Rypdal & Zhang 2001). Other factors that could lead to uncertainty are variation in storage periods, variation in house types and climate, and variation in manure properties.

5.6.3 Category-specific QA/QC and verification

In 2006, the methodology used for estimating N₂O from crop residues has been changed to the method Tier 1b (IPCC 2000). The new method is more detailed and is supposed to better reflect the real emissions than the earlier used national method. In 2014, the methodology was further enhanced with emphasis on nitrogen in residues in grass and in grain production (Grønlund et al. 2014).

There was a strong price increase for nitrogen fertiliser, which caused a stock building in 2008 and corresponding lower purchases in 2009. The calculation of N₂O emissions from use of nitrogen fertiliser is based on sales figures for each year. To correct for this, a transfer of fertiliser from 2008 to 2009 was made in the calculations.

New factors for nitrogen excretion from animals and a revision of animal statistics has been made in 2012, to better reflect the actual nitrogen excretion from each animal category and to have a more correct linkage between the nitrogen excretion factors used and the different animal categories.

In the project in 2012, when the Norwegian University of Life Sciences (NMBU) updated the Norwegian nitrogen excretion factors for the different animal species, comparisons were made with the corresponding factors used in Sweden, Denmark and Finland and with IPCC default factors as a verification of the Norwegian factors (Karlengen et al. 2012).

A new Frac_{leach} factor was estimated in a study by Bioforsk (Norwegian Institute for Agricultural and Environmental Research) in 2012 (Bechmann et al. 2012). The updated factor is based on data from the Agricultural Environmental monitoring program (JOVA).

A project with the aim to revise the Norwegian CH₄ conversion factors (MCF) for the manure storage systems in use was conducted at the Norwegian University of Life Sciences (NMBU) in 2013. The maximum CH₄ producing capacity (B₀) was also revised for cattle manure.

An update of the manure distribution between different manure management systems has been made for the N₂O emissions estimates based on the results of a survey conducted by Statistics Norway in 2013-2014 (Gundersen & Heldal 2015). Data from the manure survey of 2013 was implemented in the estimations of N₂O and CH₄ emissions from manure in the 2015 submission, and in the 2016 submission for NH₃.

In the 2015 submission, the area of cultivated organic soils has been revised back to 1990 based on an assessment by the Norwegian Institute of Bioeconomy Research. The new area estimates better reflect the land use changes measured in the national forest inventory. In connection with the implementation of the 2006 IPCC guidelines in the 2015 submission, the emission factor was reassessed and the Nordic factor of 13 kg N₂O-N/ha cropland and 9.5 kg N₂O-N/ha grassland was implemented.

5.6.4 Category-specific recalculations

In the 2019 submission the emission factor for grassland was revised from 9.5 N₂O-N/ha to 1.6 N₂O-N/ha due to a change in categorization. The Norwegian Institute of Bioeconomy Researchs changed the categorization of grassland from temperate deep-drained nutrient rich to temperate shallow drained. To be consistent with this the emission factor was changed and the time series were recalculated.

The new model for calculating NH₃ emissions and other N-species from animal manure is basis for the estimations of N₂O from animal manure applied to soils, urine and dung deposited by grazing animals and the animal manure part of indirect N₂O emissions from managed soils. See chapter 10 for more details.

5.6.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5.7 Emissions from field burning of agricultural residues – 3F

5.7.1 Category description

Burning of agricultural residues gives emissions of standard non-fossil combustion products. The source contributes with 0.1 % of the agricultural greenhouse gas emissions, and the trend has been decreasing with 90 % since 1990.

5.7.1.1 Methodological issues

CH₄, N₂O

Emissions from the burning of crop residues are being calculated in accordance with a Tier 1 approach (EEA 2013):

$$E_{\text{Pollutant}} = AR_{\text{residue_burnt}} * EF_{\text{Pollutant}}$$

$E_{\text{Pollutant}}$ = emission (E) of pollutant

$AR_{\text{residue_burnt}}$ = activity rate (AR), mass of residue burnt (dry matter)

$EF_{\text{Pollutant}}$ = emission factor (EF) for pollutant

5.7.1.2 Activity data

The calculation of the annual amount of crop residue burned on the fields is based on crop production data for cereals and rapeseed from Statistics Norway, and estimates of the fraction burned ($Frac_{\text{BURN}}$) made by the Norwegian Crop Research Institute and Statistics Norway (chapter 5.6.1.6). For cereals, a water content of 15 % is used (Statistics Norway). The activity data is consistent with the data used in the estimations of N_2O from crop residues.

5.7.1.3 Emission factors

Table 5.27 Emission factors for agricultural residue burning.

Components	Emission factors	Unit	Source
CH ₄	2.7	kg/ tonnes crop residue (d.m.) burned	(IPCC 2006)
N ₂ O	0.07	kg/ tonnes crop residue (d.m.) burned	(IPCC 2006)

5.7.2 Uncertainties and time-series consistency

Uncertainty estimates are given in Annex II.

5.7.3 Category-specific QA/QC and verification

In 2002, the emissions of CH₄ and N₂O, from agricultural residual burning were included in the Norwegian inventory. The time series were included, but it should be noted that the figures for the earlier years have a higher uncertainty than the more recent years. The amount of crop residues burned in Norway has been investigated by questionnaires in 2004 and 2012.

5.7.4 Category-specific recalculations

There have been no recalculations performed for this source category this year.

5.7.5 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5.8 Emissions from liming – 3G (Key category for CO₂)

5.8.1 Category description

Liming of agricultural soils and lakes gives emissions of CO₂. The source contributed with about 2 % of the agricultural greenhouse gas emissions in year 2017, and the emissions has decreased with 64 % since 1990.

CO₂ emissions from liming is key category according to Approach 1 key category analysis.

It is common to lime Norwegian soils because of the low buffer capacity of most soils. The use of limestone is more popular than dolomite. Also, for several years many lakes in the southern parts of Norway have been limed to reduce the damages from acidification. Estimated emissions from liming on agricultural lands have reduced since 1990, whereas liming of lakes has been relatively constant.

5.8.1.1 Methodological issues

A Tier 1 method was used with specific emission factors for limestone and dolomite.

5.8.1.2 Activity data

Statistics on consumption of liming applied to agricultural soils are derived from the Norwegian Food Safety Authority. The statistics are based on reports from commercial suppliers of lime. The amount of lime applied to lakes was collected from the Norwegian Environment Agency. It was not possible to separate the amount of lime originating as limestone or dolomite for lakes for the whole time-series.

5.8.1.3 Emission factors

The default emission factor values provided by IPCC are 0.12 Mg CO₂-C Mg⁻¹ for limestone and 0.13 Mg CO₂-C Mg⁻¹ for dolomite. For limestone this is equal to emissions of 0.44 Mg CO₂ per Mg CaCO₃ applied. The emission factors are based on the stoichiometry of the lime types.

For emissions estimates for liming on lakes, the emissions factor for limestone is used (0.12 Mg CO₂-C Mg⁻¹), as only the total amount of lime was available.

5.8.2 Uncertainties and time-series consistency

The amount of limestone and dolomite used is expected to be known with an uncertainty on ±5 percent and the emission factor with an uncertainty of ±10%.

5.8.3 Category-specific recalculations

There have been no recalculations performed for this source category this year.

5.8.4 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

5.9 Emissions from urea application – 3H

5.9.1 Category description

Urea application on agriculture soils is a minor source of CO₂ emissions in the inventory and contributes with about 0.002 % of the agriculture greenhouse gas emissions in 2017.

5.9.1.1 Methodological issues

Application of urea results in an emission of CO₂. Norway uses a Tier 1 methodology.

Annual CO₂ emissions from urea fertilisation are estimated according to equation 11.13 (IPCC 2006):

$$\text{CO}_2\text{-C Emission} = M \cdot \text{EF},$$

where:

CO₂-C Emission = annual C emissions from urea application, tonnes C yr⁻¹

M = annual amount of urea fertilisation, tonnes urea yr⁻¹

EF = emission factor, tonne of C (tonne of urea)⁻¹

5.9.1.2 Activity data

Amount of urea used is received from Norwegian Food Safety Authority annually; total sale of synthetic fertiliser, and is the same figure for the amount of urea used in the estimations of NH₃ from use of synthetic fertilisers. The amount used is very small, and consequently this is a very small source of CO₂ emissions.

5.9.1.3 Emission factors

The default emission factor of 0.20 is used (IPCC 2006).

5.9.2 Uncertainties and time-series consistency

Activity data

The uncertainty that applies to use of mineral fertilisers on ±5 percent are used.

Emission factor

Using the Tier 1 method, it is assumed all C in the urea is lost as CO₂ from the atmosphere. This is a conservative approach (IPCC 2006). No uncertainty estimate is found, and Norway uses an uncertainty of ±10%.

5.9.3 Category-specific recalculations

There have been no recalculations performed for this source category this year.

5.9.4 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6 Land-use, land-use change and forestry (CRF sector 4)

This chapter provides estimates of emissions and removals from Land Use, Land-Use Change and Forestry (LULUCF), documentation of the implementation of guidelines given in *2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006)* (hereinafter referred to as IPCC 2006 Guidelines), the 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol (IPCC 2014a), and selected parts of the *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands* (IPCC 2014b) (hereinafter referred to as IPCC 2013 Wetlands supplement).

All analyses in this chapter, except the key category analysis (Table 6.6), have been conducted by the Norwegian Institute of Bioeconomy Research (NIBIO).

6.1 Sector Overview

6.1.1 Emissions and removals

The LULUCF sector is unique compared to the other reported sectors in that it can function as both a source of atmospheric emissions through disturbance and decomposition of organic matter and a sink of emissions through the removal of atmospheric CO₂ by plants, fungi, and bacteria. The balance of the two is net emissions or removals in the LULUCF sector.

The LULUCF sector had a net removal of 24 991 kt CO₂-equivalents in 2017. These removals are substantial, and equal to approximately half of the total emissions from all other sectors than LULUCF in the Norwegian GHG accounting. The average annual net sequestration from the LULUCF sector was about 22 286 kt CO₂-equivalents per year for the period 1990-2017.

Forest land is responsible for the vast majority of the CO₂ removals in the sector in 2017. In 2017 the net removals were 29 077 kt CO₂-equivalents (Figure 6.1). Wetlands also serve as a net sink in some years, due to biomass sequestration in trees on such areas. However, in 2017, there were net emissions from wetlands of 19 kt CO₂-equivalents. Cropland was the land use category with the largest emissions in the beginning of the inventory period, with 1 845 kt CO₂-equivalents in 1990, and the emissions have increased to 2 046 kt CO₂-equivalents in 2017. The main source of emissions in the grassland category is organic soils, and the emissions are estimated to be 44 kt CO₂-equivalents in 1990 increasing to 213 kt CO₂-equivalents in 2017. Emissions from settlements have become about 2.5 times greater during the inventory period, with an increase from 769 kt CO₂-equivalents in 1990 to 2 034 kt CO₂-equivalents in 2017. Cropland followed closely by Settlements are the land use categories that are responsible for the largest emissions of the six land use categories in the LULUCF sector in 2017. Emissions from other land were 0.95 kt CO₂-equivalents in 2017. The decay from the harvested wood products (HWP) pool was larger than the input to the pool from 2009-2016. In 2017, the removals from HWP were -237 kt CO₂, a reduction by 322 kt CO₂ since last year.




Figure 6.1 Net CO₂ emissions and removals (kt CO₂-equivalents per year) from the LULUCF sector by land-use category (forest land, cropland, grassland, wetlands, settlements, other land, and harvested wood products) from 1990 to 2017, including emissions of N₂O and CH₄. Source: Norwegian Institute of Bioeconomy Research.

Forest land was the major contributor to the net sequestration of CO₂ in the sector. In 2017, the total net removals from forest land were 29 424 kt CO₂ (Figure 6.2). Emissions from forest land occurred primarily from organic soils (677 kt CO₂ from organic soils on forest land remaining forest land and land converted to forest land). Mineral soil constituted a removal of 72 kt CO₂. Living biomass was the primary contributor to sequestration with 75 % (21 635 kt CO₂) of the total removals for forest land remaining forest land. The dead wood and litter pools contributed 5 % (1 304 kt CO₂) and 22 % (6 458 kt CO₂) to the total C sequestration on forest land remaining forest land, respectively. Land converted to forest land contributed with removals of 514 kt CO₂, primarily due to sequestration in living biomass and the litter pool.




Figure 6.2 Emissions and removals of CO₂ on forest land from organic and mineral soil, dead wood, litter, and living biomass, 1990–2017. Source: Norwegian Institute of Bioeconomy Research.

Since 1989, the carbon stocks in living biomass in the LULUCF sector have increased significantly by around 40 % (Table 6.1). This increase is mainly due to the increase in the growing stock on forest land (Figure 6.3).

Table 6.1 Carbon stocks in 1989 and 2015 and differences in C stocks compared to 1989 as a total for all land-use categories, including associated uncertainties. The estimates are based on the sample plots in the lowlands outside Finnmark (>16 000 plots). SE = standard error.

Year	C stock (kt)	C stock difference to 1989 (kt)	2 SE (%) of C stock difference to 1989
1989	316 856	-	-
2015*	449 038	132 182	6

*The estimates are based on the last five years sampled in the National Forest Inventory (2013-2017). The estimate is therefore valid for the mid-year 2015.

Annual variation in CO₂ removals on Forest land

Forest land covers around one third of the mainland area of Norway, and is the largest land-use category that is considered managed. Other land is the largest land-use category irrespective of management status. The carbon stock in living biomass on forest land has increased throughout the inventory period (Figure 6.3).




Figure 6.3 Development of the carbon stock in living biomass on forest land remaining forest land from 1990–2017. Source: Norwegian Institute of Bioeconomy Research.

The steady increase in carbon stock in living biomass is caused by several factors. One main factor is an active forest management policy over the last 60–70 years which resulted in increasing increment of timber volume (and biomass). The combination of the policy to rebuild the country after World War II and the demand for timber, led to a large national effort to invest in forest tree planting in new areas, mainly on the west coast of the country, and replanting after harvest on existing forest land. In the period 1955–1992, more than 60 million trees were planted annually with a peak of more than 100 million planted annually in the 1960s. These trees are now at their most productive age and contribute to the increase in living biomass and hence the carbon stock. At the same time, levels of annual fellings are much lower than the annual increment, causing an accumulation of tree biomass (Figure 6.4). The number of planted trees has been decreasing since 1992, and since 2003, only about 20 million trees have been planted annually. The last few years, the number of planted trees has slightly increased, and in 2015 31 million trees were planted (Landbruksdirektoratet, 2017). The lower number of planted trees compared to the post-war decades, together with a changed age structure of the forest, may result in a relative decrease in biomass accumulation, and hence a decrease in the future carbon sequestration.




Figure 6.4 Forest felling, annual increment and volume, 1919–2017. The two last years are extrapolated for volume (without bark) and annual increment. Source: Norwegian Institute of Bioeconomy Research and Statistics Norway.

6.1.2 Activity data

The main data source used for the LULUCF sector is the National Forest Inventory (NFI). Data from the NFI is used to estimate the total areas of forest land, cropland, wetlands, settlements and other land, as well as the land-use transitions between these categories. Land area accounting for the inventory has been done according to an Approach 3, as described in chapter 3 of the IPCC Guidelines (IPCC 2006). The NFI data are also used to calculate changes of carbon stocks in living biomass and as input values for modeling changes in the carbon stock in dead organic matter (DOM) and mineral soil for forest land remaining forest land. DOM is the combined pool of litter and dead wood.

The NFI utilizes a 5-year cycle based on re-sampling of permanent plots. The sample plots are distributed across the country in order to reduce the periodic variation between years, and each year 1/5 of the plots are inventoried. The same plots are inventoried again after five years, and all plots are assessed during a 5 year period. The current system with permanent plots was put in place between 1986 and 1993, and made fully operational for the cycle covering the years 1994 through 1998. Because the re-sampling method was not fully implemented before 1994, the method used to calculate annual emissions and removals is not the same throughout the time-period, and the methods have been bridged. See section 6.3.1 for a detailed description of the method.

The annual changes in the C stock depend upon several factors, such as harvest levels and variable growing conditions due to temperature and precipitation. All these factors influence the reported annual changes of CO₂ removals from the atmosphere, but among these the harvest level is the most important.

The annual fluctuation seen in CO₂ sequestered from dead organic matter and soil are influenced by annual variation in the C input data to the Yasso07 model. Carbon input to the Yasso07 model is from

standing biomass, dead organic matter from natural mortality, and harvest residues including stumps and roots from harvested trees. All these factors are influenced by the same natural and man-made factors as stated for living biomass and all cause annual changes.

The NFI data are complemented with auxiliary data for several other sink/source categories, e.g. horticulture, arable crop types, grassland management, synthetic N fertilization, drainage of forest soil, and forest fires. These data are acquired from Statistics Norway, the Norwegian Agricultural Authority, the Norwegian Food Safety Authority, the Norwegian Environment Agency, and The Directorate for Civil Protection and Emergency Planning. Detailed descriptions of these data are provided under their relevant emission categories.

6.1.2.1 Land-use changes 1990–2017

Net land-use changes in Norway from 1990 to 2017 have been very small. Only the area of settlements has increased slightly, while the other land-use categories have decreased or remained constant (Figure 6.5).




Figure 6.5 Area distribution of the IPCC land-use categories for 1990 and 2017.

Source: Norwegian Institute of Bioeconomy Research

The small land-use changes are also illustrated by the land-use conversion matrix for the whole inventory period from 1990 to 2017 (Table 6.2). A key finding from these data is that the changes in land-use from 1990 to 2017 are quite small; with approximately 0.8 % of the total land in a "land-conversion" category and the rest in a "remaining" category. The largest changes were in forest land and settlements. There have been land-use conversions from all categories to forest land and to settlements. Whereas the net area of settlements decreased, the net area of forest land decreased. The classification of land-use change is almost directly transferable to the activities reported under the Kyoto Protocol, which is illustrated by the land-use matrix in Table 6.2. More details about the activities reported under the Kyoto Protocol, as well as the definition of human-induced land-use change, are given in chapter 11. Under the convention reporting we apply a 20-year transition period, which means that areas reside in conversion classes for 20 years before they are transferred to the remaining class.

Table 6.2 Land-use change matrix for the IPCC land-use categories from 1990 to 2017. The 20 year transition period is ignored in this table, in order to give a full picture of all conversions that happened in the period. This results in differences to the specific area estimates in the CRF. The column Total here are nonetheless the same as the sum of lands to and remaining in a category reported in the CRF.*

Year	Land-use (kha)							
	Land-use	2017						Total*
	Land-use	Forest land	Cropland	Grassland	Wetlands	Settlements	Other land	Total*
1990	Forest land	12038	22	31	4	107	0	12200
	Cropland	14	904	0	0	20	0	938
	Grassland	17	3	200	0	9	1	231
	Wetlands	13	4	1	3723	2	0	3742
	Settlements	23	2	1	0	559	0	585
	Other land	34	0	0	1	3	14645	14682
	Total*	12138	934	233	3728	699	14646	32378

**Differences of totals and column or row sums are due to rounding.*

6.1.3 Uncertainties

Uncertainties of area estimates are based on standard sampling methodology. The areas of the largest land-use categories, other land remaining other land and forest land remaining forest land, can be estimated with a precision (2 standard errors) of < 2 % (Table 6.3). Land-use changes are generally small in Norway. The largest change category is forest land converted to settlements. The uncertainty estimate for this area estimate is approximately 18 %. Due to the small number of NFI sample plots in several of the other land-use conversion categories, the relative size of the uncertainty estimates can be quite large. However, the absolute size of the uncertainty in those classes is nonetheless small.

The uncertainties of carbon stock change (CSC) estimates in living biomass in forest land, grasslands, wetlands, and other lands, were estimated as described in section 6.3. Estimated uncertainties are based on the sampling error. As for area estimates, the relative uncertainty estimates for CSC were quite large for small land-use categories, whereas their absolute size was comparably small (Table 6.3). For living biomass on cropland converted to settlements and settlements converted to croplands, the uncertainty was based on Tier 1 defaults. Uncertainty estimates for CSC estimates for the dead organic matter (DOM) pool were based on expert judgement by considering the uncertainty in the living biomass estimates.

Table 6.3 Uncertainties of living biomass and dead organic matter (DOM) pools shown as total aggregated uncertainty (U_{total}) based on the uncertainties of the carbon stock change (CSC) per hectare and the area estimates. Area uncertainty is for combined mineral and organic soils. See Table 6.4 for area uncertainty of mineral soils which is used to obtain the total uncertainty of DOM. 2 SE means two times the standard error.

Code	Land-use class	Area ^a	CSC	U_{total}	CSC	U_{total}
		2 SE %	Living biomass (2 SE % or default uncertainty)		DOM (2 SE % or default uncertainty)	
4A1	Forest land remaining forest land ^b	2	12	13	14	15
4A2	Cropland to forest land	53	71	87	100	119
4A2	Grassland to forest land	52	100	113	200	201
4A2	Other land to forest land	67	88	111	135	151
4A2	Settlements to forest land	39	54	67	100	107
4A2	Wetlands to forest land	57	121	134	100	142
4B1	Cropland remaining cropland ^c	0	75	75	NA	NA
4B2	Forest land to cropland	42	109	117	128	135
4B2	Grassland to cropland	125	NO	NO	NA	NA
4B2	Settlements to cropland	200	75	201	NA	NA
4B2	Wetlands to cropland	97	NA	NA	NA	NA
4C1	Grassland remaining grassland	13	100	101	NA	NA
4C2	Forest land to grassland	35	194	194	107	113
4C2	Wetlands to grassland	200	150	201	NA	NA
4D1	Wetlands remaining wetlands	5	44	44	NA	NA
4D2	Forest land to wetlands	101	169	197	182	201
4D2	Other land to wetlands	200	NA	NA	NA	NA
4E2	Cropland to settlements	49	75	90	NA	NA
4E2	Forest land to settlements	18	48	51	100	102
4E2	Grassland to settlements	67	75	101	NA	NA
4E2	Other land to settlements	136	NA	NA	NA	NA
4E2	Wetlands to settlements	106	150	184	NA	NA
4F2	Grassland to other land	201	NA	NA	NA	NA

^a The area uncertainty is the same for living biomass and DOM. ^b Includes a safety margin for model errors of 1 percentage-points. DOM is combined for litter, dead wood, and mineral soil and therefore the same as in Table 6.4. ^c Area uncertainty of 0 % is based on SSB data and pertains to orchards. Total area uncertainty for cropland remaining cropland is 7% based on NFI estimates.

Uncertainties for mineral soil CSC factors on land-use conversion categories were assumed to be 50 % for conversions between forest land, cropland, and grassland. We assumed a lower uncertainty for these conversions than for the others because the SOC stocks were based on national measurements or national data of soil types applied to IPCC default values. For conversions to and from land-use classes, where SOC stock measurements were not available, we assumed the uncertainty to be 100 % (Table 6.4). Uncertainties in the C loss from drained organic soils were calculated using the error ranges supplied in the IPCC 2013 Wetlands supplement (IPCC 2014b) for all drained organic soils on cropland, grassland, forest land, and land under peat extraction. The uncertainty of the emission factors was then combined with the uncertainty of the area estimates determined by the sampling error. For two smaller classes (managed wetlands – peat extraction and orchards on croplands), the uncertainty of the area estimates is based on expert judgement as their

areas are not estimated by the NFI. The uncertainty in the soil type classification method, i.e. the inaccuracy of the soil maps, was ignored.

Table 6.4 Uncertainties of the mineral soil and drained organic soil pools shown as total aggregated uncertainty (U_{total}) based on the uncertainties of the carbon stock change (CSC) and the area estimates. 2 SE means two times the standard error.

Code	Land-use class	CSC	Area	U_{total}	CSC	Area	U_{total}
		Mineral soil (2 SE % or default uncertainty)			Drained organic soil (2 SE % or default uncertainty)		
4A1	Forest land remaining forest land ^a	14	2	15	40	50	64
4A2	Cropland to forest land	50	64	81	40	94	102
4A2	Grassland to forest land	50	52	72	NA	NA	NA
4A2	Other land to forest land	100	67	121	NA	NA	NA
4A2	Settlements to forest land	100	39	107	40	200	201
4A2	Wetlands to forest land	90	102	136	40	69	80
4B1	Cropland remaining cropland	50	7	51	19	25	31
4B2	Forest land to cropland	50	44	67	19	141	142
4B2	Grassland to cropland	50	125	135	NA	NA	NA
4B2	Settlements to cropland	100	200	201	NA	NA	NA
4B2	Wetlands to cropland	NA	NA	NA	19	97	99
4C1	Grassland remaining grassland	91	13	92	50	125	135
4C2	Forest land to grassland	50	35	61	NA	NA	NA
4C2	Wetlands to grassland	NA	NA	NA	50	200	201
4D1	Wetlands remaining wetlands ^b	NA	NA	NA	NA	NA	110
4D2	Forest land to wetlands	90	144	170	40	141	147
4D2	Other land to wetlands	100	200	201	NA	NA	NA
4E2	Cropland to settlements	100	49	111	NA	NA	NA
4E2	Forest land to settlements	100	19	102	19	66	69
4E2	Grassland to settlements	100	70	122	100	200	201
4E2	Other land to settlements	100	136	169	NA	NA	NA
4E2	Wetlands to settlements	90	151	176	19	141	142
4F2	Grassland to other land	100	201	201	NA	NA	NA

^a Uncertainty for mineral soil on forest remaining forest is combined for litter, dead wood, and mineral soil.

^b The sub-category peat extraction includes on-site and off-site emissions; specific uncertainties for areas and CSC are therefore not given.

Default uncertainty estimates were also used for N₂O and CH₄ emissions from drained organic soils, for direct and indirect N₂O emissions, and for biomass burning.

Table 6.5 Uncertainties of N₂O and CH₄ emissions for direct and indirect N₂O emissions and for drained organic soils shown as total uncertainty (U_{total}) based on the uncertainties of the emission factor (EF) and the activity data (AD). 2 SE means two times the standard error.

Code	Source	Land-use class	Gas	U_{total}	EF	AD
			Default uncertainty			
4(I)	Direct N ₂ O from inorganic N inputs	Forest land	N ₂ O	201	200	20
4(I)	Direct N ₂ O from organic N inputs	Settlements	N ₂ O	201	200	20
4(II)	Drained organic soils	Forest land	N ₂ O	64	41	50
4(II)	Drained organic soils	Wetlands - Peat extraction	N ₂ O	151	113	100
4(II)	Drained organic soils	Cropland	CH ₄	74	70	23
4(II)	Drained organic soils	Forest land	CH ₄	180	173	50
4(II)	Drained organic soils	Grassland	CH ₄	117	48	107
4(II)	Drained organic soils	Wetlands - Peat extraction	CH ₄	128	NA	NA
4(III)	Direct N ₂ O	N mineralization/ immobilization	N ₂ O	224	200	100
4(IV)	Indirect N ₂ O from managed soils	Atmospheric deposition	N ₂ O	447	400	200
4(IV)	Indirect N ₂ O from managed soils	Leaching and runoff	N ₂ O	287	233	167
4(V)	Biomass burning	Wildfires in forest	N ₂ O	70	70	28
4(V)	Biomass burning	Wildfires in forest	CH ₄	70	70	28

In the cases where the uncertainty of the activity data estimate was not derived from the NFI and the uncertainty of the CSC was based on expert judgment, the total uncertainty was derived by combining the two uncertainties. The specific methods and assumptions are described further for each of the sinks/sources under the sections of the individual land-use categories.

6.1.4 Key categories

A sink or source can be a key category either with respect to the level (size of the emission) or the trend (change in the size between 1990 and 2017). The key category analysis for the Norwegian inventory is performed by Statistics Norway. All of the reported sinks and sources were included in the analysis for the LULUCF sector. The CSC estimates for living biomass, dead organic matter (DOM), mineral soils and organic soils, were considered for each specific land-use conversion (e.g. forest land converted to cropland). The key category analysis was performed using an Approach 1 and Approach 2 approach for the whole greenhouse gas inventory.

From the analyses, 27 key categories were identified by a combination of Approach 1 and Approach 2 level analyses (Table 6.6). Forest land remaining forest land (FF) is the most important category in the LULUCF sector. Living biomass in FF is identified as the largest key category; followed by litter + dead wood + mineral soil; forest land converted to settlements DOM; and cropland remaining cropland organic soil. Living biomass was also a key category for forest land converted to settlements, grassland or cropland, and for grassland remaining grassland. Carbon stock change estimates for dead organic matter (DOM) on all lands converted from forest land, except for other land and wetlands, were also identified as key categories. CO₂ emissions from drained organic soils were a key category for the remaining categories for cropland, forest land, settlements, and peat

extraction on-site and off-site emissions combined; N₂O and CH₄ emissions from drained organic soils on forest land were also key categories. For the mineral soil pools on land in conversions, forest conversion to grassland, settlements, and cropland were key categories, as were grassland converted to forest land. Forest land converted to settlements is an important land use change category (largest area change), and all four sources were determined as key categories. N₂O emission from mineralization and immobilization due to soil management is also a key category due to the inclusion of all land-use conversions. There were no new sources that became key categories this year. The sources no longer classified as key categories this year are living biomass on grassland converted to forest, mineral soil on cropland converted to settlement, and CH₄ emissions related to drained organic soils from cropland.

Table 6.6 Approach 2 key category analysis results for the LULUCF sector showing level assessments for 1990 and 2017, and the trend assessment for 1990–2017. Key categories are indicated by bold values and the larger the value the more important is the key category.

Code	Sink/source category	Gas	Level assess 1990	Level assess 2017	Trend assess 1990–2017
4.A.1	Forest remaining forest - Living biomass	CO ₂	9.67	14.53	15.82
4.A.1	Forest remaining forest - Litter + dead wood + Mineral soil	CO ₂	3.00	6.13	7.44
4.E.2.1	Forest to Settlement - DOM	CO ₂	0.33	4.52	6.83
4.B.1	Cropland remaining cropland - Organic soil	CO ₂	4.03	2.61	1.20
4.A.1	Forest remaining forest - drained organic soils	CO ₂	2.95	2.10	1.19
4.C.2.1	Forest to Grassland - DOM	CO ₂	0.02	1.97	3.07
4.A.2.2	Grassland to Forest - DOM	CO ₂	0.09	1.43	2.17
4.B.2.1	Forest to Cropland - DOM	CO ₂	0.04	1.30	2.01
4.E.2.1	Forest to Settlement - Living biomass	CO ₂	1.90	1.21	0.54
4.A.2.4	Settlement to Forest - DOM	CO ₂	0.05	1.13	1.73
4.C.2.1	Forest to Grassland - Living biomass	CO ₂	0.46	1.11	1.41
4(II)Forest	Forest remaining forest- drained organic soils	N ₂ O	1.21	0.92	0.58
4.E.2.1	Forest to Settlement - Mineral soil	CO ₂	0.06	0.78	1.18
4.E.1	Settlements remaining settlements - Organic soil	CO ₂	1.00	0.77	0.49
4.B.2.1	Forest to Cropland - Living biomass	CO ₂	0.57	0.77	0.80
4.C.2.1	Forest to Grassland - Mineral soil	CO ₂	0.01	0.66	1.03
4G	Harvested Wood Products - HWP	CO ₂	3.81	0.61	1.76
4.E.2.1	Forest to Settlement - Organic soil	CO ₂	.	0.61	.
4(II)Forest	Forest land - drained organic soils	CH ₄	0.70	0.52	0.31
4.B.2.3	Wetland to Cropland - Organic soil	CO ₂	.	0.50	.
4.C.1	Grassland remaining grassland - Living biomass	CO ₂	0.78	0.47	0.17
4.A.2.1	Cropland to Forest - DOM	CO ₂	0.04	0.41	0.61
4.B.2.1	Forest to Cropland - Organic soil	CO ₂	0.05	0.38	0.56
4(III)	Direct N ₂ O from N mineralization/immobilization	N ₂ O	0.02	0.38	0.58
4.D.1	Wetland Peat extraction - on+off-site - Organic soil	CO ₂	0.60	0.35	0.11
4.A.2.2	Grassland to Forest - Mineral soil	CO ₂	0.02	0.34	0.52
4.B.2.1	Forest to Cropland - Mineral soil	CO ₂	0.01	0.25	0.39

6.1.5 Completeness

The following sources were not reported because they are not mandatory: carbon stock change in living biomass, DOM, and net carbon stock change in soils on flooded land remaining flooded land (CRF table 4.D.1.2) and CO₂, N₂O, and CH₄ emissions from rewetting of organic soils on forest land, cropland, grassland, and wetlands (CRF table 4(II)). Furthermore, the following sources were not reported because emissions are considered negligible: N₂O and CH₄ from controlled burning on forest land and from wild fires on grasslands (CRF table 4(V)), and the area of organic soil and net carbon stock change in organic soils on land converted to peat extraction (CRF table 4.D.2.1). For sources that were not reported, the notation key NE is used in the CRF.

6.1.6 Quality assurance and quality control (QA/QC) for LULUCF

NIBIO implements the QA/QC plan described for the National Inventory System in Annex V. In addition, a LULUCF-specific plan for QA/QC was developed internally at NIBIO. The LULUCF-specific plan has two objectives: 1) to ensure that emission estimates and data contributing to the inventory are of high quality, and 2) to facilitate an assessment of the inventory in terms of quality and completeness. These objectives are in accordance with chapter 6 of the 2006 IPCC guidelines for quality assurance and quality control.

The QA/QC plan for the LULUCF sector is based on the general Tier 1 QC procedures and includes two check lists (one for the source-category compiler and one for the LULUCF inventory compiler), an annual timeframe of the outlined QC activities, and a target for when to elicit QA reviews.

Specifically, the QC is performed on the following 12 points:

1. Documentation of assumptions and selection criteria
2. Transcription errors
3. Emission calculations
4. Labeling of parameter units, conversion factors, and unit transfers
5. Database integrity
6. Consistency within sectors and source categories
7. Transfer of estimated emissions between inventory staff
8. Uncertainty estimation and calculations
9. Review of internal documentation
10. Time-series consistency
11. Completeness
12. Comparison to previous estimates

Several QA projects have been undertaken for the LULUCF reporting. In general, QA is initiated if a new method or model is implemented. Below are some examples of previously elicited QA activities.

Two external quality-assurance actions were undertaken in 2012. First, elicitation by the Norwegian Institute for Forest and Landscape (now NIBIO) of a qualified researcher to evaluate and improve the methodologies applied for emission estimates from cropland and grassland. This work resulted in

substantial method revisions for most source categories due to the lack of methods evaluation since their development documented by (Rypdal et al. 2005). Moreover, a detailed documentation and justification of the new methods are provided in the report *Emissions and methodologies for cropland and grassland used in the Norwegian national greenhouse gas inventory* (Borgen & Hylen 2013). The second external QA was a smaller task performed on the final emission estimates for mineral soil on grassland remaining grassland, which was elicited from an expert at Colorado State University. This task provided a review of the emission calculations (the application of the new Tier 1 method) and of the method and activity data documentation. The methods were developed in accordance with the 2006 IPCC guidelines and implemented in the National GHG inventory in 2013.

Work was done to make a QA of the Yasso07-model estimates for mineral soil on forest land in 2014 – 2015. In this project, modelled and measured soil C stocks were compared on two field sites over time. Results from these sites and the overall estimation methodology for the relevant pools on forest land were discussed at two seminars with three contracted external experts from Finland, Denmark, and Norway (Dalsgaard, L. et al. 2017). In addition, a comparison of Yasso07 (current methodology) and field estimates of soil C stocks was carried out (Dalsgaard, L et al. 2016).

With the implementation of the IPCC 2006 guidelines, an external QA was elicited on the HWP calculations. The QA was performed by an expert from the Swedish University of Agricultural Sciences before the NIR 2015 submission.

Internal structures at NIBIO for the work on the LULUCF reporting have changed slightly every year. One important aim of the changes is to improve the QC procedures and to ensure that methods and calculations are put through an internal QC before reporting. The CRF tables went through internal QC by more than one person before the database was submitted to the national focal point. Furthermore, after the overall compilation of estimates from all sectors, there was an exchange of CRF tables from the focal point to NIBIO, and an additional QC was performed. Improving the QA/QC procedures is an ongoing process that will be further improved in future submissions.

6.2 Land-use definitions and classification system

6.2.1 Land-use definitions

National Forest Inventory (NFI) data are used to estimate total area of forest land, cropland, grassland, wetlands, settlements, and other land, and the land-use transitions between these. The rationale of using the NFI as activity data for all land-use categories is that it covers the whole country by sample plots. In addition, the NFI is the the most reliable data set available that can be used to determine transitions between different land-use categories during the whole reporting period in a consistent manner. The land-use categories are defined in accordance with the IPCC guidelines (IPCC 2003; IPCC 2006). They are described below, using the national terminology. In order for a unit to be classified as separate from an adjacent land-use category, it must have a minimum area and width of 0.1 ha and 4 m, which is consistent among all land-use categories in Norway. The NFI land cover and land-use categories and their transcription into IPCC land-use categories are illustrated in Table 6.7. All forest land, cropland, grassland (and settlement), are managed lands. A very small part of wetlands is managed for peat extraction. The vast majority of wetlands (and all of other lands) are unmanaged. Because managed and unmanaged lands are reported for the specific categories, total unmanaged lands are reported as IE.

Table 6.7 NFI land cover and land-use categories and their correspondence to the UNFCCC land-use categories.

Land use Land cover	Forestry (no other use or restrictions)	City, urban area Settlements of different kinds	Cabin area	Recreation area	Military training field	Protected Area, Nature Reserve	Roads/Railroad Airport	Power line	Other
Productive forest land (1)	Forest land	Settlements	Forest land	Forest land	Forest land	Forest land	Settlements	Settlements	Settlements
Non-productive forest land (2)	Forest land	Settlements	Forest land	Forest land	Forest land	Forest land		Settlements	Settlements
Other wooded land, Crown cover 5-10% (3)	Other land		Other land	Other land	Other land	Other land		Other land	
Wooded mire, Crown cover 5-10%	Wetlands					Wetlands			
Coastal calluna heath	Other land	Settlements				Other land	Settlements		
Bare rocks, shallow soil	Other land		Other land	Other land	Other land	Other land			Other land
Mire without tree cover	Wetlands					Wetlands			
Lakes and rivers (not sea)					Wetlands	Wetlands			Wetlands
Grazing land, not regularly cultivated									Grassland
Arable land, regularly cultivated						Cropland			Cropland
Other areas, gravel pits, mines, gardens, halting places, skiing slopes, forest roads etc.	Settlements	Settlements		Settlements	Settlements		Settlements	Settlements	Settlements

1) Productive forest land is defined as forest with crown cover that exceeds 10 % and that hosts a potential yield of stem-wood including bark of $> 1 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$.

2) Non-productive forest land is defined as forest with crown cover that exceeds 10 % and that hosts a potential yield of stem-wood including bark of $< 1 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$.

3) Other wooded land is defined as land with sparse tree cover with crown cover between 5 and 10 % and hosts trees that have the potential to reach a height of 5 m, or with a combined cover of shrubs, bushes, and trees above 10 %. It is classified as other wooded land when found on mineral soil (organic layer $< 40 \text{ cm}$ deep) and as wooded mire if found on organic soil (organic layer $> 40 \text{ cm}$ deep).

Forest land (4A) is defined in the National Forest Inventory (NFI). The values used in the NFI are in accordance with the range of parameters in the definition from the Global Forest Resources Assessment (FRA) 2005. Forest land is land with tree crown cover > 10 %. The trees have to be able to reach a minimum height of 5 m at maturity in situ. Minimum area and width for forest land considered in the Norwegian inventory is 0.1 ha and 4 m. Forest roads are considered as settlements. The minimum area and width is consistent among all land-use categories in Norway. Young natural stands and all plantations established for forestry purposes, as well as forest land, which is temporarily unstocked as a result of e.g. harvest or natural disturbance, are included under forest land. All forest in Norway is managed either for wood harvesting, protection and protective purposes, recreation, and/or to a greater or lesser extent, hunting and berry picking. On more marginal and less productive forest land the various management practices may be less intense, but still be present. Hence, all forest in Norway is considered managed.

Cropland (4B) is defined as lands that are annually cropped and regularly cultivated and plowed. Both annual and perennial crops are grown. It also encompasses grass leys that are in rotations with annual crops, which may include temporarily grazed fields that are regularly cultivated. This category includes arable land that was previously annually cropped and regularly plowed, but has since been abandoned. These areas remain in the cropland category until they have a regrowth of trees that make them unsuitable for plowing. All cropland is considered managed.

Grassland (4C) is defined as areas utilized for grazing on an annual basis. More than 50 % of the area should be covered with grass and it may be partly covered with trees, bushes, stumps, rocks etc. The grass may be mechanically harvested but the soil cannot be plowed. Land with tree cover may be classified as grassland if grazing is considered more important than forestry, even if it meets the forest definition. According to the agricultural statistics that are used for determining grassland management practices, grasslands include two categories; grazing lands and surface-cultivated grass. All grassland is considered managed according to these categories.

Wetlands (4D) are defined as lakes, rivers, mires, and other areas regularly covered or saturated by water for at least part of the year. Mires may be stocked by trees but with a tree coverage that do not meet the forest definition. Most wetlands are assumed to be unmanaged. Wetlands used for peat extraction and flooded lands caused by human constructed dams are considered managed.

Settlements (4E) include all types of built-up land: houses, gardens, villages, towns, cities, parks, golf courses, sport recreation areas, power lines within forests, areas close to cabins (< 5m), industrial areas, gravel pits and mines. All settlements are considered managed.

Other land (4F) is defined in the NFI as waste land, such as bare rocks, ice, and shallow soils that may have particularly unfavorable climatic conditions. In accordance with the IPCC definition, other land can also include unmanaged land areas that do not fall into any of the other five land-use categories, for example, heath lands, other wooded land (i.e. land with sparse tree cover on mineral soil), and open areas.

Table 6.8 Management status of different land-use categories. An area is only classified as belonging to one land-use category. The predominant national land cover and land use determines the assigned category.

Land-use category	Management status
Forest land	Managed
Cropland	Managed
Grassland	Managed
Wetlands	Unmanaged and managed (small area)
Settlements	Managed
Other land	Unmanaged

6.2.2 Consistency in areas and reporting categories

6.2.2.1 Area consistency

Up to the 2010 submission, the area of the different land-use categories were based on sample plots below the coniferous limit. In order to determine the land use at higher altitudes and in Finnmark county, the NFI included the first complete set of sample plots for these areas in the period 2005–2010. This allows for assessment of the extent of forest area, other wooded land, and other land uses in these areas. The plots are incorporated in the ordinary management plan for the NFI. On plots without previous measurements, land use and biomass development was estimated back to 1990 (back-casting) using data from the NFI (Anton-Fernandez & Astrup 2012), maps and aerial photographs for settlements, grassland, and cropland. This was done to improve the area estimates of 1990 for all new plots included in the system.

The definitions of land cover and land-use categories have been consistent for most categories since the permanent plots were established in the period 1986-1993. There have, however, been some changes in definitions throughout this period that have affected the land-use change matrix. The most important change relates to the forest definition. In 2005, the NFI forest definition was adjusted to the (IPCC 2003) definition for forest land, replacing a similar but not identical definition. The change of the forest definition did not result in an inconsistency, however, because the new forest definition was also applied to NFI data acquired before 2005. Also the category grassland had not been defined in the land-use classification in the first cycle of the NFI with permanent sample plots (6th NFI, 1986 - 1993). The land-use classes assessed in the 7th NFI (1994-1998) have been utilized for the corresponding plots in the 6th NFI. The Norwegian Mapping Authority provided the value for the total land area of Norway.

6.2.2.2 Land-use changes prior to 1990

The NFI was not designed to assess land use change categories prior to 1990, and the forest inventory at that time did not cover the whole country. To be able to make a rough indication of the overall trend in forest area, the areas of productive forest land according to national classification is presented in Table 6.9. The data are taken from the Census of Agriculture and Forestry 1967, 1979 and 1989. Because no data from permanent sample plots exists before 1986 and relatively small changes have been detected on forest land, we have chosen not to take into account land-use changes that may have occurred prior to 1990. This implies that CSC in living biomass on land converted to forest land may be underestimated, but the potential changes in living biomass are included in forest land remaining forest land.

Table 6.9 Area estimates of productive Forest land (kha) in the years 1967, 1979, and 1989.

Region	1967	1979	1989
Eastern and Southern Norway	3 903	4 085	4 289
Western Norway	689	770	895
Mid-Norway (Trøndelag)	974	976	997
Northern Norway	916	829	1 439
Total	6 482	6 660	7 620

Source: Statistics Norway 1969, 1983, 1992

6.2.3 Sink/source categories

Changes in C stocks are reported for the five main pools under the UNFCCC: living biomass (gains and losses), litter, dead wood, mineral soils, and organic soils. For all land-use classes except for forest land, litter and dead wood are summarized and reported as a part of the dead organic matter pool. The pools are defined as follows:

Living biomass: For all land-use categories except cropland, living biomass is defined as the biomass of living trees with a breast height diameter > 50 mm. Table 6.10 describes in more detail on which land-use categories living biomass is measured in the NFI. The tree biomass is the sum of the biomass estimates of the tree fractions stem wood, stem bark, living branches, dead branches, needles or leaves as well as stump and roots down to a root diameter of 2 mm (see section 6.4.1). On cropland, carbon stock changes in living biomass are calculated on areas with fruit trees.

Table 6.10 Measurements of tree parameters in the NFI given Norwegian land cover and land-use classes. Green cells indicate measurement of trees (a = measurements since 2007, and b = measurements since 2010). Grey cells indicate that trees are not measured on that land-use class. Not all land use and land cover combinations exist (see Table 6.7).

Land cover	Land use								
	Forestry (no other use or restrictions)	City urban area Settlement s of different kinds	Cabin area	Recreation area	Military training field	Protected Area, Nature Reserve	Roads/Railr oad Airport	Power line	Other
Productive forest land (1)								b	
Non-productive forest land (2)								b	
Other wooded land, Crown cover 5-10% (3)								b	
Wooded mire, Crown cover 5- 10%								b	
Coastal calluna heath								b	
Bare rocks, shallow soil								b	
Mire without tree cover									
Lakes and rivers (not sea)									
Grazing land, not regularly cultivated	a	a	a	a	a	a	a	a	a
Arable land, regularly cultivated									
Other areas, gravel pits, mines, gardens, halting places, skiing slopes, forest roads etc.									

Litter: For forest land remaining forest land, the changes in the dead organic matter pool are the changes resulting from the input and decomposition of all dead organic material (woody and non-woody, aboveground and belowground; C input) regardless of size and stage of decomposition. Only the most recalcitrant material (humus) originating from root decomposition is allocated to the soil

pool. The changes in the litter and the dead wood pools, respectively, are allocated according to the origin of the model C input (aboveground or belowground elements), the chemical quality and the size of the C input elements – see details in chapter 6.4.1.1. For land converted to or from forest land, the litter pool entails dead organic material in various stages of decay found above the mineral forest soil and developed primarily from leaves/needles, twigs, and woody material (L, F, and H horizons in the Canadian soil classification). Due to the field sampling and laboratory methodology, this includes living fine roots and excludes particles > 2 mm after sample preparation.

Dead wood: For forest land remaining forest land the estimates for CSC in the dead wood pool are modeled (see above for *litter*). For land converted to or from forest land, the dead wood pool entails dead organic material (standing and lying dead wood in various stages of decay) aboveground (dimension > 10 cm) and belowground (dimension > 5 mm). Estimates were found through expert judgement and dimensional limits are approximate.

Mineral and organic soils: Because soil mapping was historically covered by three different entities, the separation of organic and mineral soils differs somewhat between forest land, cropland, and grassland. On forest land, organic soils mapping was performed by the NFI and was defined as having an organic layer deeper than 40 cm with no requirement on minimum C content. On cropland and grassland, organic soils were partially mapped by a national soil survey and were defined as soil with a topsoil organic layer of at least 40 cm and with a carbon content greater than or equal to 10%. So far only 59 % of all croplands and 6% of all grasslands are classified. The remaining non-covered area for cropland (41%) and grassland (94%) classification were derived from the national land resource map AR5²⁵ based on old economic maps where organic soils were defined by a topsoil organic layer deeper than 20 cm with no requirement on minimum carbon content.

As strict correspondence with the official organic soil definition of the world reference base (WRB) would be impossible (Table 6.11), the general term “organic soil” used consistently throughout this report refers to the three definitions mentioned above.

Table 6.11 Organic soil definition for the three entities that participated to the carbon mapping in Norway compared with WRB definition.

Entities	Land use covered	Topsoil organic horizon	
		Thickness	C content %
NFI	Forest	≥ 40cm	-
National soil survey	59 % cropland 6 % grassland	≥ 40 cm	≥10 %
AR5	41 % cropland 84 % grassland	≥ 20 cm	-
WRB (Histosol)	All	≥ 10cm	Rare saturation: >20 % in top 20 cm*
			Frequent saturation, clay = 0 %: ≥12 % in top 20 cm*
			Frequent saturation, clay > 60 %: ≥18 % in top 20 cm *

*The top 20 cm may partially include mineral horizon if the organic layer thickness < 20 cm.

²⁵http://kilden.skogoglandskap.no/?topic=arealinformasjon&layers=ar5_bonitet&X=7260874.41&Y=125323.37&zoom=0&lang=nb&bgLayer=graatone_cache&layers_opacity=0.75

6.3 Land area representation and the National Forest Inventory

The area representation applied in the LULUCF reporting is based on the Norwegian National Forest Inventory (NFI; see section 6.3.1 below). Land accounting is based on an Approach 3 according to IPCC 2006 guidelines. Under the convention reporting we apply a 20-year transition period. Hence, land stays in a conversion class for 20 years (transition period) before it is transferred to a remaining class.

6.3.1 Current NFI design

The NFI can be characterized as a single-phase, permanent, systematic, and stratified survey. An interpenetrating panel design is used, where 1/5th of the sample plots that are evenly distributed across the country (the so-called “panel”) using a Latin square design are measured each year. The Norwegian Institute of Bioeconomy Research is responsible for operating the NFI. Inventory work was started in 1919 with regular inventory cycles. The 11th inventory cycle started in 2015 and will be completed in 2019.

The NFI divides Norway into four strata: lowlands (below the coniferous limit) except Finnmark county, mountain areas (above the coniferous limit) except Finnmark, lowlands in Finnmark, and mountain areas in Finnmark. The lowland strata contain the most productive forests, while the forests in the other strata consist mainly of low productive birch forests. The arctic island groups Svalbard and Jan Mayen are not covered by trees or bushes, and are therefore not considered in the NFI.

NFI sample plots are placed on the intersections of grid lines to ensure a systematic distribution of the plots (Figure 6.6). The distance between neighboring plots is different in the strata. A 3x3 km (Easting x Northing) grid is used in the lowlands including Finnmark county, a 3x9 km grid is used in the mountains not located in Finnmark and a 9x9 km grid is used in the mountainous area of Finnmark county (Figure 6.6).




Figure 6.6 The sample plots are covering all land-use categories. In the example map to the left, plots are placed in the systematic 3x3 km grid. On the right-hand side, we see the distribution of land use-categories in the south eastern part of Norway below the coniferous tree line (only 3x3 grid).




Figure 6.7 Spatial distribution (approximate locations) of the NFI sample plots in the four strata. The sample plots in Finnmark county located on the 3x3 km grid are covering lowlands, while the sample plots on the 9x9 km grid cover mountainous areas.

As can be seen from the estimate of all land-use categories for the year 2010, more than 94 % of the living biomass stock is allocated in the lowland forests outside Finnmark (Table 6.). The mountain forest outside Finnmark, the mountain forest in Finnmark county, and the lowlands in Finnmark account for 3.7 %, 1.6 %, and 0.4 % of the carbon in living biomass, respectively.

Table 6.12 Area and estimates of carbon stocks in living biomass in 2010 (the reference year is based on observations from 2008-2012) by stratum and associated uncertainties (SE = standard error).

Stratum	Area (kha)	C stock (kt)	2 SE (%) C stock	Percent (%) of total C stock
Lowlands outside Finnmark	14 989	423 533	2.9	94.3
Mountain forest outside Finnmark	12 528	16 738	12.3	3.7
Lowlands in Finnmark	135	1 773	21.4	0.4
Mountain forest in Finnmark	4 727	7 164	24.9	1.6
All	32 378	449 208	2.9	100.0

A plot that has measured trees in the current inventory is always revisited in the next inventory. Plots that were not visited in the field in the most recent inventory are monitored using aerial images, which are acquired approximately every five years for the entire country. From the aerial images, the plot is assessed for land-use changes and the occurrence of trees. If it is not possible to determine the land-use category with certainty or if there is an indication that the sample plot may be tree covered, the sample plot is visited in the field. Exceptions are cropland and settlements, which are not visited in the field in order to measure tree parameters.

Among other attributes, the positions, diameter at breast height (DBH, diameter measured at 1.3 m above ground) and tree species of all trees with DBH >50 mm are recorded on circular sample plots with a radius of 8.92 m (250 m²). On plots with 10 trees or less, all tree heights are measured using hypsometers. On plots with more than 10 trees, heights are measured on a relascope-selected subsample with a target sample size of 10 trees per plot (NFLI 2008). The heights of the unmeasured trees are estimated using tariffs (models) calibrated at the plot-level with data from measured trees (Breidenbach et al. 2013).

The area of a stratum A_h was estimated by multiplying the proportion of points on the 3x3 km grid that belong to the stratum h with Norway's land area. The representation factor, also known as the design weight or the inverse of the sampling probability, determines how much area of Norway one sample plot represents. The representation factor of a sample plot is given by A_h/n_h , where n_h is the number of sample plots on the grid that is specific to the stratum.

If a sample plot covers two land use classes, the sample plot, and consequently also the representation factor, is divided between the plot parts according to the proportion of the land-use classes covering the plots. A land-use class must cover at least 20 % of the sample plot in order to be considered. Land-use class cover is recorded in 10 % steps on divided sample plots.

6.3.2 Classification of mineral and organic soil areas

In order to identify the soil type (mineral or organic) for all land-use classes, additional sources to the NFI data are necessary for all sample plots without tree cover. Since the 2015 NIR submission, we have used a baseline 1990 map, classifying all areas as organic or mineral soil for all land uses, and overlaid it with the NFI plots. This enabled geo-referencing of the areas of organic soils for each land-use class and tracking of land-use changes on mineral or organic soils.

Two maps were first combined to obtain spatial soil type information for cropland, grassland, and settlements. The Norwegian agricultural soil classification database contains detailed soil profiles of 59 % of croplands and 6% of grasslands. Information of soil type on the rest of the land area was derived from the national land resource map AR5²⁶. The soil type information on forest land, wetlands, and other land were derived from NFI registrations.

Figure 6.8 displays all organic soils and includes non-drained organic soils on forest land and wetlands. On cropland, grassland, and settlements all organic soils were considered drained.




Figure 6.8 Map of NFI plots on organic soil per land-use class for the 1990 baseline.

6.3.3 Changes in the NFI design

The NFI consisted of temporary sample plots before 1986. Between 1986 and 1993, all lowland sample plots outside Finnmark were permanently marked. All sample plots located within one to three neighboring counties (“fylke”) were measured within one year. Annual estimates representative for the whole country were therefore complicated in those years. The current system with interpenetrating panels was made fully operational in the cycle covering the years 1994 through 1998.

Today, all sample plots are circular with a size of 250 m². However, in the inventory period between 1986 and 1993, concentric sample plots were used in some counties. All trees with a DBH \geq 50 mm

²⁶http://kilden.skogoglandskap.no/?topic=arealinformasjon&layers=ar5_bonitet&X=7260874.41&Y=125323.37&zoom=0&lang=nb&bgLayer=graatone_cache&layers_opacity=0.75

were measured on a circular sample plot with an area of 100 m². Trees with a DBH larger than 19.9 cm were measured on a larger sample plot. The area of the large sample plots was 200 m² in Østfold, Akershus, Oslo, and Nord-Trøndelag counties, whereas the area was 250 m² in all other counties. The sample weights of the trees on these plots were modified to bridge the methodological change in the subsequent inventory period where all sample plots had a size of 250 m².

The sample plots in the mountain stratum outside Finnmark were established between 2005 and 2009. The first re-measurements of these plots were consequently started in 2010. The sample plots in the two Finnmark strata were established between 2005 and 2011 with first re-measurements starting in 2012. This made special methods for estimating changes on the plot-level necessary, as described in section 6.2.2.1. Almost 95 % of the carbon stock in living biomass is, however, found in the lowland stratum outside Finnmark. The land use classes observed on the plots in Finnmark and the mountain plots established between 2005-2009 (2005-2011 for Finnmark) were back-casted to 1990. Plots were assigned to one of the two Finnmark strata in the inventory cycle by assessing their location on the 3x3 km grid. In a number of western and northern Norwegian municipalities outside Finnmark, a height threshold separating lowland and mountain areas was set by local forest authorities. In the other parts of Norway, the stratum decision was made using auxiliary information and in the field. The strata classification can therefore be described as a two-phase procedure.

Prior to 2005, the tree heights of three trees per species were measured on each sample plot. Since 2005, 10 trees per plot are measured as described above.

Until 1994 no differentiation between grasslands and croplands was made; both were considered agricultural land. Since 1994, this difference is made and the areas of croplands and grasslands were back-casted.

In the first years of establishing the mountain plots, there were no aerial photos available. Maps (N50) were used to determine which plots could be forested. Those were visited in the field, while the rest of the plots were given a land use class derived from an overlay of the maps (plot center). A plot with a center in a land use class was assumed to be that land use class on its full area (circle of 250 m²). If the plot center was in water, the sample plot was not visited, even if it was close to forest. Between 2011 and 2015, all plots were checked against aerial images. Land use changes occurred to and from all land use classes. Most commonly, however, parts of the plots in water were then considered other lands or wetland (mire), but in some cases forest or other land use classes. The changed land use class and possible measurements of trees were back-casted to 1990 in those cases.

Until 2005, trees were not measured on land use classes other than forest. Sample plots were only split if one of the parts was forest. In other cases the plots were categorized as fully covering the land use class of the center coordinate. Since 2005, trees are also measured on other land use classes than forest. First "other wooded land" (part of the land use category other lands) was included, further land use classes in 2007 ("grassland" and the rest of "other land" and "cabin areas") and since 2010 also trees below "power lines" (part of the settlements category) are measured. This resulted in splitting a number of sample plots between other land use classes than forest, which resulted in area changes of those land use classes. The changed land use class and possible measurements of trees were back-casted to 1990 in those cases.

In 2005, the forest definition was changed to be in accordance with the FAO definition, which includes the consideration of crown cover. Until 2005, the forest definition was based on the

(potential) productivity of the land, which had to be larger or equal to $0.1 \text{ m}^3 \text{ ha}^{-1} \text{ year}^{-1}$ to be considered as forest. (In addition, a threshold of $1 \text{ m}^3 \text{ ha}^{-1} \text{ year}^{-1}$ is used to separate unproductive and productive forest. This is nonetheless not relevant for LULUCF reporting.) Since 2005, the crown cover on a 0.1 ha area around the plot center has to be more than 10 % for a land to be forest. After harvests, areas may be temporarily unstocked but will remain forest in the NFI classification system unless the land use changes later. For young productive forest, also the stem number is considered. All plots with forest according to the new definition that were visited for the first time in the NFI period 2005-2009 (2005-2011 for Finnmark) were back-casted. All plots (not only new plots) in forest according to the new definition were considered forest also in 1990. Exceptions were plots with a human induced afforestation. Exceptions were also made in productive forest if the plot had been assessed as “non forest” in a previous cycle and the age of the trees clearly allowed determining a year of change from some land use class to forest.

6.3.4 Inter- and extrapolation for area and living biomass estimates

The NFI consists of five panels each of which consists of approximately $1/5^{\text{th}}$ of all 22 008 sample plots. Panel #1 was installed²⁷ in 1994 and the other panels in the following years, such that panel #5 was installed in 1998 (Figure 6.9). After the panels were installed, all plots were re-measured every 5 years. However, all sample plots were visited for the first time and permanently marked between 1986 and 1993. In this period before the panels were installed, the measurement intervals for the sample plots within a panel varied. For example, for panel #1 in 1994, the sample plots were previously measured between one and eight years before.

All estimates are based on linear interpolation of areas and carbon stocks between panel-wise estimates. The first estimate for each panel is for 1989, based on sample plots measured between 1986 and 1993 in the respective panel. Towards the end of the reporting period, estimates were extrapolated based on the last two estimates per panel. This way, the rate of land-use changes is projected based on observations of the last 10 years (Figure 6.9). The extrapolation results in recalculations of the estimates of the last four years in the forthcoming reports as new data become available, and interpolation can be used instead of extrapolation.

An example to illustrate the method: while no extrapolation was necessary for panel #4 in the 2014 reporting (2012 as the final year), four years of extrapolation were necessary for panel #5 (Figure 6.9). Measurements on panel #5 for the year 2013 became available in the 2015 reporting (2013 as the final year), which resulted in a recalculation of the years 2009-2012 for panel #5.

The annual estimate reported is the sum of one estimate in the panel that was measured in the reporting year and the interpolated or extrapolated estimates of the other panels in the reporting year.

²⁷ Installation in this context means that all sample plots within the panel were visited in one year. All sample plots (in the lowlands outside Finnmark) were visited and marked for the first time between 1986 and 1993.




Figure 6.9 Estimated forest land remaining forest land area covering mineral soils within the five NFI panels illustrated with data from 1989 until 2012. Diamonds indicate the measurement year of the sample plots in the respective panel (1-5). The estimated area was interpolated between two measurement years and extrapolated in the years after the last measurement year in each panel. Areas of lands converted to forest land that will change their category to forest land remaining forest land after 2009 are not considered in the graphic.

More formally, the area of a land-use class (A_{LUC}) in a given measurement year (diamonds in Figure 6.9) is the sum over all $i=1, \dots, n_p$ sample plots within a panel

$$A_{LUC} = \sum_i p_{LUC,i} \cdot rf_i$$

where $p_{luc,i}$ is the proportion (0,...,1) of a land-use class covering a sample plot and rf is the representation factor (the area of Norway which the sample plot represents).

Linear interpolation of stocks means constant changes (gains and losses) between two measurements. Biomass losses (fellings) are mainly due to harvests and are observed over five years in each panel. In order to reflect the annual variability in harvests, the constantly interpolated or extrapolated biomass losses have been adjusted according to harvest statistics provided by Statistics Norway (Figure 6.10). This results in annual variability of the net carbon changes. The adjustment according to the harvest statistics was carried out for the land-use categories land converted to forest land and forest land remaining forest land.

The change of biomass stocks (gains or losses) within a land-use class in a given measurement year (diamonds in Figure 6.10) is the sum of changes over all sample plots within a panel

$$c_{LUC} = \sum_i p_{LUC,i} r f_i c_{LUC,i}$$

where $c_{LUC,i}$ is the mean annual change of the biomass stock per hectare on a sample plot per land-use class. The change $c_{LUC,i}$ can either be a gain (positive change) or a loss (negative change) of biomass. Biomass gains or losses were multiplied with the default factor of 0.5 in order to obtain estimates of carbon gains or losses.




Figure 6.10 Biomass losses in forest land remaining forest land observed on the five panels illustrated with data from 1989 until 2012. Left-hand side: Diamonds indicate the measurement year of the sample plots in the respective panel. The estimated biomass loss was interpolated between two measurement years and extrapolated in the years after the last measurement year in each panel. Interpolation and extrapolation are based on a constant function. Right-hand side: The constant interpolation or extrapolation is adjusted according to harvest statistics (thick black line).

The harvest statistics are based on Statistics Norway's table 03895²⁸ describing the timber use by assortment and municipality. This table was supplemented by the estimated use of domestic fire wood obtained from Statistics Norway (personal communication). The harvest statistics for the last reporting year are preliminary.

²⁸ Statistics Norway. (2016). 03895: Commercial removals of industrial roundwood, by assortment [Avvirking for salg, etter sortiment] (m³) (K).

6.3.5 Uncertainties in areas and living biomass

Standard errors of area and biomass change estimates used in the key category analysis were estimated based on a 5-year moving average estimate for the mid-year of the last NFI cycle. For example, in NIR 2018, the last NFI cycle includes measurements from the years 2012-2016, which means that the mid year was 2014. Model-related uncertainties resulting from interpolation and extrapolation are ignored. Also model-related uncertainties resulting from the use of biomass models to estimate single tree biomass from diameter and height measurements were ignored since they can be assumed to be small for CSC compared to the sampling error (Breidenbach et al. 2013). Furthermore, the uncertainty resulting from using an estimated instead of measured tree height for some trees on the sample plots was ignored. Also this source of variation can be assumed to be negligible compared to that of sampling. For the most important gain category, living biomass in forest land remaining forest land, 5 % points were added to the standard error (2 SE) by expert judgment to consider the ignored uncertainties.

For the estimation of sampling errors, the estimates of land-use class areas are stratified estimates of land-use class proportions multiplied with Norway's land area (including lakes). Random sampling is assumed in all estimates. The variances can therefore be assumed to be conservative estimates.

The estimated proportion of a land-use class within a stratum is given by

$$p_h = 1/n_h \sum_i y_{hi}$$

where $h = (1, \dots, 4)$ is the stratum identifier, n is the number of sample plots, y is the proportion of the NFI sample plot that belongs to a land-use class, which is 1 if the sample plot falls completely into the class, and $i = 1, \dots, n_h$.

The estimated variance of the proportion is given by

$$\text{var}(p_h) = \frac{s_h^2}{n_h}$$

with $s_h^2 = \frac{1}{n_h-1} \sum_i (y_{hi} - \bar{y}_{hi})^2$. The area estimate of a land-use class (A_{LUC}) over all strata is then given by the stratified estimator

$$A_{LUC} = A \frac{1}{N} \sum_h N_h p_h$$

where A is Norway's land area, N is the land area divided by the NFI plot size, N_h is the stratum area divided by the plot size, and p_h is the proportion of the respective land-use class. The estimated variance of the area estimate is given by

$$\text{var}(A_{LUC}) = A \sum_h \left(\frac{N_h}{N} \right)^2 \text{var}(p_h).$$

Similar to the area estimates, estimates of sampling errors of carbon gains or losses are based on the full set of NFI sample plots. The estimate of the total biomass gain or loss within a stratum is given by the ratio estimator

$$T_h = \frac{N_h}{n_h} \sum_{i=1}^{n_h} y_{hi}$$

where n_h is the number of sample plots within a stratum and y is the average annual gain or loss that occurred during the last five years on a sample plot. An estimate of the variance is given by

$$var(T_h) = N_h^2 \frac{s_h^2}{n_h}.$$

The total biomass gain or loss estimate (T) over all strata and its variance ($var(T)$) is the sum over T_h and $var(T_h)$, respectively.

Post-stratification did not improve the precision of biomass gain and loss estimates. We tested climatic zones, counties, and forest districts as possible post-strata.

The estimation of biomass or carbon stocks is not required in the CRF. In this report, stocks were calculated in analogy to the biomass change estimates.

The uncertainties of carbon estimates are given by

$$U(C) = \sqrt{U(T)^2 + U(CF)^2}$$

where $U(T)$ is the uncertainty of the total biomass gain or loss estimate in percent of the estimate

$U(T) = \frac{2\sqrt{var(T)}}{T} 100$ and $U(CF)=2\%$ is the relative uncertainty in the carbon fraction.

6.3.6 QA/QC for the NFI data

Fieldwork is conducted by NFI field staff. Qualification requirements are forestry or natural management education at the college level or higher. Before a new employee can work independently, a training period of at least three weeks is conducted. All field staff undergo a week long course prior to each field season. There are currently about 25 employees who perform fieldwork in the period from May to October. It has been a stable situation with few changes in the field personnel, and on average the field workers have more than 10 years experience.

All data collection is done on handheld computers with software developed particularly for the purpose. The field computer program has a number of features built in for quality assurance:

- The program ensures that everything that must be recorded is recorded.
- A series of tests on the logical values of measurements.
- Categorical variables are recorded with the help of menus.

For plots that have been previously registered, the field computer contains data from the previous record. Depending on the character of the variable, quality checks are handled in three different ways:

- The old value is displayed and can be confirmed or amended.
- The old value is hidden, but a warning is given if the new value is not logical compared to the

old value.

- The old value is displayed as information before a new registration is done.

Data is sent by e-mail to the data reception center at the main office once a week. The data reception center keeps track of which sample plots have been registered and which plots remain, thereby ensuring that no plots are omitted. The data is then read into a database and further quality checks are made. Incorrect data or questions are returned to the field worker for clarification.

Each field worker is usually visited by a supervisor for one day in the field. Control registrations are carried out by an experienced field worker who makes a second registration for approximately 5 % of all sample plots. The control data is then analyzed to document the quality of field recordings, partly to clarify misunderstandings, and to correct for any systematic errors. Results of control entries are published in a separate report.

During the winter months, there is a systematic review of the data with additional error testing and inspection of all codes and logic. This happens before the data is read into the final table structure.

The database is a relational database that is designed to ensure data quality. Primary keys and foreign keys prevent double accounting and ensure coherence in the data.

6.4 Forest land – 4A

6.4.1 Forest land remaining forest land, 4A1

Forest land remaining forest land covers slightly more than 12 million hectares (ha). Forest ownership in Norway is dominated by private ownership, with many small properties. There were 127 544 forest holdings with more than 2.5 ha of productive forest land in Norway in 2016 (SSB 2015). Due to the ownership structure and specific terrain conditions, Norwegian forestry is diversified and characterized by small-scale activity. The average size of clear-cuttings was estimated to be 1.9 ha in 2003 (Statistics Norway 2004). Approximately 90 % of the harvesting is fully mechanized.

Forest land is the most important land-use category with respect to biomass sequestration in Norway. According to the Approach 2 key category analysis (Section 6.1.4), forest land is a key category for sequestration in living biomass, dead wood, litter, and mineral soils, and emissions from organic soils, because of the uncertainty in both the level and trend.

6.4.1.1 Methodological issues

Living biomass (key category)

The stock change method is used. The method implemented corresponds to Tier 3, which uses a combination of NFI data and models to estimate changes in biomass.

The reported CSC refer to the biomass of all living trees observed on an NFI sample plot with a stem diameter larger than 50 mm at breast height (DBH). Thus, shrubs and non-woody vegetation are not included in the estimates. Since tree coordinates are measured on NFI plots, each tree can be attributed to a land use category. Single tree allometric regression models developed by Smith et al. (2016; 2014), Marklund (1988), and Petersson and Ståhl (2006) are applied to DBH and height measurements from the NFI for estimating the tree biomass. For consistency with estimates reported under the Kyoto Protocol, the tree biomass is defined as the sum of aboveground and belowground biomass. The aboveground biomass of a tree is the sum of the estimates of the fractions of stem, stump, bark, living branches, and dead branches. The belowground biomass is the estimate of the fraction of stump and roots minus the estimate of the fraction of stump. Table 6. lists the models used to estimate the biomass of the different tree fractions. The biomass models are defined for Norway spruce (*Picea abies*), Scots pine (*Pinus sylvestris*), and birch (*Betula pendula* and *Betula pubescens*). These species constitute approximately 92 % of the standing forest volume (Larsson & Hysten 2007). Other broad-leaved species constitute most of the remaining eight percent. The birch biomass models are applied to all broad-leaved species. The living biomass is estimated consistently based on the same biomass models from the base year 1990 onwards.

Table 6.13 Biomass models for estimating living biomass. In Marklund's (1988) models, the notation "G (model number)" indicates Norway spruce and "T (model number)" Scots pine.

Component	Reference and specific model
Dead branches	Marklund (1988), G20, T22. Smith et al. (2014) birch/deciduous.
Living branches	Marklund (1988), G12, T14, Smith et al. (2014) birch/deciduous. Includes needles.
Foliage	Smith et al. (2014) birch/deciduous.
Bark	Marklund (1988), G8, T10. Smith et al. (2014) birch/deciduous.
Stem	Marklund (1988), G5, T6. Smith et al. (2014) birch/deciduous.
Stump	Marklund (1988), G26, T28.
Stump and roots (>2 mm)	Petersson and Ståhl (2006), B i (for Norway spruce, and Scots pine). Smith et al. (2016) birch/deciduous.

Dead organic matter (key category)

The model used to estimate C stock changes in soils provides a change estimate for total soil organic carbon (SOC), which includes the dead wood, litter, and soil pools. This methodology is used for the forest area on mineral soil only. The estimate of total SOC entails all stages of decomposition and all C input elements regardless of size and origin (input aboveground or belowground). The total SOC change estimate was allocated to the dead wood, litter, and soil pools, respectively. This was done by allocating specific chemical model pools to the reporting pools and by using the information about the dimension of the C input elements as well as its origin as either aboveground or belowground C input (Figure 6.11). Only the changes in the H pool (humus; Figure 6.12) (1.9 %) originating from the belowground C input elements of all sizes were allocated to the changes in the UNFCCC soil sink/source category. The remaining change in the total soil organic C stock was attributed to dead wood (16.5 %) and to litter (81.6 %). The same allocation percentages were used for all years since 1990. See below for a description of the Yasso07-model used for the simulations on mineral soils.

Origin	Aboveground					Belowground				
Chemical component	A	W	E	N	H	A	W	E	N	H
Non woody	LITTER									
Fine woody										
Coarse woody										
	DEAD WOOD									
										SOIL

Figure 6.11 Conceptual definitions of soil pools based on the chemical composition of Yasso07 output for total soil C stock change. AWENH is defined as: Acid soluble, Water soluble, Ethanol soluble, Non-soluble, and Humus.

Mineral soils (key category)

Choice of method

A Tier 3 method was applied. The emissions and removals of total soil organic C (dead wood, litter, and soil pools) from forest land on mineral soil are estimated using the decomposition model Yasso07 (Tuomi et al. 2008; Tuomi et al. 2009; Tuomi et al. 2011a; Tuomi et al. 2011b). The Norwegian application is described in Dalsgaard et al. (2016). Yasso07 represents processes for mineral soils down to a depth of 1 m and operates using five chemical soil C pools (Figure 6.12). Decomposition (CO₂ release) and fluxes among the chemical C pools are regulated by climatic input data and parameters governing decomposition, transformation, and fractionation of C inputs. The

model is applied to the time series for each individual NFI plot. It is run on an annual time step, but only estimates for the NFI registration years are used. The term “entry” below refers to any combination of an NFI plot and registration year.

For each NFI plot in the category forest land remaining forest land, C changes per hectare since the last measurement of trees on the plot were calculated using Yasso07, as described below. The calculated change was then up-scaled to country-wide estimates using the same method as for living biomass, which is described in section 6.3.4.




Figure 6.12 Flow diagram for Yasso07. Fluxes significantly different from 0 are indicated by the arrows (Liski et al. 2009).

For each entry (ca. 11 400 NFI plots) annual living tree C input to the model is estimated from tree registrations. On plots where the time series was not complete, back-casting was applied (see section 6.2.2.1). Tree biomass models were used to estimate biomass components (Table 6.13) and annual turnover rates for roots and branches were applied to estimate the annual C input (Table 6.4 and Table 6.15).

Tree C input generated annually from natural mortality and residues from diffuse harvest (i.e. harvest not including commercial thinning or final harvest) was estimated on all entries as a percentage of the standing biomass. Data from the 8th NFI (2000-2004) and the 9th NFI (2005-2009) were used to establish look-up tables for this purpose (Anton-Fernandez & Astrup 2012). Registrations of mortality and harvest on NFI plots started in 1994. The look-up tables are grouped by tree species (broadleaved or conifer), site-index (up to six classes), and age (up to nine classes). Harvest residues from commercial thinning and final harvest were estimated from plot specific registrations (since 1994) of harvested volume. This C input was relevant on a total of 1 753 entries.

The look-up tables mentioned above also contain factors (percentages) describing the biomass development between two inventories. These were used to establish a time series of living biomass and harvest residues (commercial thinning and final harvest) back to 1960 (back-cast). Field registrations of the 6th inventory (1986-1993) on prior land use and forest management activities were used to establish eight rules covering all relevant NFI plots. For young stands where harvest must have taken place during the back-cast period, harvested biomass and biomass of the old stand back in time was estimated using old NFI inventories, where standing volume was generally lower

than found in current inventories. Estimation of C input from the back-cast time-series (including from mortality and diffuse harvest) followed the same procedures as for the NFI time-series but using average distribution (from NFI) to biomass components as individual tree information is not available in the back-casted part of the time series. The 1960-1990 time-series is used to reduce the effect of the equilibrium assumption on the reported values of soil C change in the inventory period (see below).

Table 6.14 Biomass models used in Yasso07 simulations. When models from Marklund (1988) are used, the notation "G" is used for Norway spruce and "T" for Scots pine.

Component	Reference and specific model
Dead branches	Marklund (1988), G20, T22, Smith et al. (2014) birch/deciduous
Living branches	Marklund (1988), G12 and G16, T14 and T18, Smith et al. (2014) birch/deciduous
Foliage	Marklund (1988), G16, T18, Smith et al. (2014) birch/deciduous
Bark	Marklund (1988), G8, T10, Smith et al. (2014) birch/deciduous
Stem	Marklund (1988), G5, T6, Smith et al. (2014) birch/deciduous
Stump	Marklund (1988), G26, T28 (for Scots pine), Smith et al. (2016) birch/deciduous ^a
Roots (> 5 cm)	Marklund (1988), G28, T31, Smith et al. (2016) birch/deciduous ^a
Roots (2 mm–5 cm)	Petersson and Ståhl (2006), Bi (for Norway spruce, Scots pine)
Roots (< 2 mm)	Marklund 1988, G28, G26, T31, T28, Smith et al. (2016) birch/deciduous ^a
	0.3 × foliage biomass; (Kjønaas et al. Manuscript)

^a No distinct diameter limit is inferred between the two classes of deciduous coarse roots.

Table 6.15 Annual turnover rates applied for tree C input estimation. Compiled in Peltoniemi et al. (2004) and de Wit et al. (2006).

Component	Norway spruce	Scots pine	Broadleaved	Reference
Foliage	0.143	0.33	1	Tierney and Fahey (2002)
Live and dead branches, roots > 2 mm	0.0125	0.027	0.025	Muukkonen and Lethonen (2004) DeAngelis et al. (1981) Lethonen et al. (2004)
Roots < 2 mm	0.6	0.6	0.6	Matamala et al. (2003)

The C input generated from the ground vegetation is estimated using models based on plot tree species and age (Muukkonen & Mäkipää 2006; Muukkonen et al. 2006). Distinction is made among Norway spruce, Scots pine, and deciduous (birch spp.), with an age span of 0-200 years (Norway spruce and Scots pine) or 0-100 years (deciduous). Output of aboveground biomass is generated for four layers of ground vegetation: i) moss, ii) lichens, iii) herbs and grasses, and iv) shrubs. For shrubs and herbs and grasses, it is assumed that belowground biomass is twice the aboveground biomass (e.g. Poeplau (2016)). A compilation of studies documenting the above-to belowground-ratio for biomass and the annual turnover rates for ground vegetation litter Table 6.16 can be found in Peltoniemi et al. (2004).

Table 6.16 Annual turnover rates for litter from ground vegetation.

Component	Moss	Lichens	Herbs and grasses	Dwarf shrubs
Aboveground	0.33	0.1	1	0.25
Belowground	-	-	0.33	0.33

The chemical composition of tree C input was based on data used in the development of the Yasso07 model. For ground vegetation litter, the values in Peltoniemi et al. (2004) were used (Table 6.17).

Table 6.17 The fraction of C input made up of acid soluble (A), water soluble (W), ethanol soluble (E), and insoluble (N). See also Figure 6.12. If more than one value was available these were averaged by species and by chemical fraction and normalized to a sum of 1 across all four fractions.

Component ^a	A	W	E	N
<u>Stem</u>				
Norway spruce	0.63, 0.7	0.03, 0.005	0, 0.005	0.33, 0.28
Scots pine	0.66, 0.68	0.03, 0.015	0, 0.015	0.29, 0.28
Deciduous	0.65, 0.78	0.03, 0	0	0.32, 0.22
<u>Roots (<2mm)</u>				
Norway spruce	0.5508	0.1331	0.0665	0.2496
Scots pine	0.5791	0.1286	0.0643	0.228
Deciduous	as foliage	as foliage	as foliage	as foliage
<u>Foliage</u>				
Norway spruce	0.4826	0.1317	0.0658	0.3199
Scots pine	0.5180	0.1773	0.0887	0.2160
Deciduous	0.4079, 0.46	0.198, 0.1929	0.099, 0.0964	0.2951, 0.2507
<u>Living and dead branches</u>				
Norway spruce	as stem	as stem	as stem	as stem
Scots pine ^b	0.3997-0.5307	0.0105-0.0295	0.0382-0.1309	0.411-0.4608
Deciduous	as stem	as stem	as stem	as stem
Roots > 2 mm	as branches	as branches	as branches	as branches
Stumps	as stem	as stem	as stem	as stem
Bark	as foliage	as foliage	as foliage	as foliage
<u>Ground vegetation^c</u>				
Moss	0.74	0.0867	0.0433	0.13
Lichens	0.836	0.0747	0.0373	0.052
Herbs and grasses	0.27	0.4667	0.2333	0.03
Shrubs	0.56	0.2067	0.1033	0.13

^a The majority of values are from the Yasso07 user manual (Liski et al. 2009). ^b 25 observations were available. The range is given. ^c From Peltoniemi et al. (2004): W is 2/3 of "extractable"; E is 1/3 of "extractable".

C input was either non-woody (foliage, fine roots, all ground vegetation input), fine-woody (living and dead branches, coarse roots and bark), or coarse-woody (stems and stumps). The dimensions entering Yasso07 for each of the three size-groups are 0, 2, and 10 cm, respectively. Mean C input for all entries are found in Table 6.18.

Table 6.18 Mean values for C input and predicted soil C (AWENH denotes the chemical pools in Yasso07 to which C is distributed).

C input and model estimated soil C stocks (mean values)	Non-woody	Fine-woody	Coarse wood mortality	Coarse wood harvest	Total
C input (kg C m ⁻² yr ⁻¹)*	0.170	0.055	0.008	0.005	0.238
Equilibrium stock (kg C m ⁻²)	3.8	1.0	0.4	0.1	5.3
Equilibrium stock (kg C m ⁻²), A pool	0.37	0.10	0.07	0.01	0.54
Equilibrium stock (kg C m ⁻²), W pool	0.05	0.01	0.01	<0.01	0.07
Equilibrium stock (kg C m ⁻²), E pool	0.05	<0.01	<0.01	<0.01	0.06
Equilibrium stock (kg C m ⁻²), N pool	1.70	0.48	0.28	0.04	2.50
Equilibrium stock (kg C m ⁻²), H pool	1.67	0.41	0.07	0.01	2.16
Predicted stock* (kg C m ⁻²)	3.8	1.1	0.4	0.1	5.5

*Across all entries in the time-series, excluding back-cast entries.

For each NFI plot, start values for the five chemical C pools (Figure 6.12) were found by a pre-simulation or spin-up. This was done in two steps: 1) running the model in 5000 annual time steps to

equilibrium in all chemical pools²⁹ and 2) running the model for a C input time series 1960-1990 specifically constructed for this purpose (see above). C input for the equilibrium spin-up was the mean C input estimated for the historical time series (1990-2016), grouped by tree species and site-index. For the back-cast period as well as for the inventory period, total SOC was estimated for each entry, i.e. each time where a registration was available. Plot specific total SOC was found as follows: individual plot estimated annual C input for each entry in the time-series was used as input. Stock from the previous entry was used as the start value. A loop was applied to drive the model in as many years as is found between the entries (mostly five years but this deviates in some cases in the early inventory years). For the first entry, a loop of five years was applied following the spin-up stock. C input as well as the simulated soil organic C stocks are kept in units of kg C m⁻². The Graphical User Interface parameter set for Yasso07 was applied (Tuomi et al. 2011b). To arrive at the reported timeseries, interpolation was done in the same way as for the living biomass estimates.

For spin-up as well as for the time-series, the applied weather data for Norway (Engen-Skaugen et al. 2008) was specifically produced for the NFI grid. Weather data for the equilibrium spin-up was the plot-specific climatic normal for the time period 1961-1990. For the time series simulations, plot specific weather data using the mean for 1991-2008³⁰ was applied.

The estimate of total SOC changes between entries in the time-series have been distributed to the dead wood, litter, and soil sink/source categories described above under the section on dead organic matter (see also Figure 6.11.).

Activity data

A variety of input data were used for mineral soils. This includes area representation for plots (as described for the NFI), basic NFI registrations (as described for the living biomass) as well as site-index and stand age, complementary models and parameters including biomass models, turnover rates, chemical C input composition, and C input dimensions. Climate data were available from the Norwegian Meteorological Institute. The usage and values of input data are described under *Choice of method* above.

The input data from the NFI database used for the Yasso07 simulations did not account for the fact that certain plots of land were converted to forest land before the current NFI design was completed (see section 6.3) and are to be included as forest land remaining forest land in 2010. These small areas were found by interpolation (see section 6.4.1.1) and were assigned an emission/sequestration rate equal to the mean in the relevant years for the area covered by the methodology described above.

Assumptions/justification

The NFI definition of mineral soil is based on the depth of the organic layer (< 0.4 m). We assume that the decomposition processes on these areas are represented by the model structure and the parameters of the Yasso07 model found from data on mineral soils throughout the world. A more detailed delineation between mineral and organic soils (based on soil taxonomical classification) is currently not possible.

²⁹ Increasing the spin-up time to 15000 annual time steps resulted in C stock changes of 0.03% lower and C stocks of 0.09% higher than with the chosen standard of 5000 steps (mean values across all plots).

³⁰ For technical reasons climate data is currently not available for 2009-2016.

The allocation to the dead wood, litter, and soil pools assumes that there was no transport of humus (H) from the aboveground pools to the mineral soil since 1990. Thus, changes in soil organic C originating from aboveground litter in all stages of decomposition are assumed to be found in the organic layer above the mineral soil. While this is not strictly to be expected in reality, all soil organic C is accounted for and assumptions related to the distribution to the dead wood, litter, and soil pools do not affect the total emissions/removals. The assumptions result in a very small part of the total change to be allocated to the soil pool. According to field studies, changes in the mineral soil are very slow and are often not significantly different from zero (Emmett et al. 2007; Peltoniemi et al. 2004).

Drained organic soils (key category)

CO₂ emissions from drained organic soils on forest land is a key category. Due to the lack of national emissions factors, a Tier 1 method was chosen.

Activity data

To estimate the area of drained organic soils on forest land, statistics on subsidies for draining forest soils to enhance productivity was used. There was an increase in the area drained annually in the 1950's, with a peak of approximately 13 kha yr⁻¹ in the early 1960's. Since then, drainage to promote forest production became much less practiced and the establishment of drainage ditches on mires with the aim of forest production was prohibited by law in 2007. Up until 2007, a total accumulated area of 241 kha was reported as drained.

Areas of drained soils for forest production were provided by Statistic Norway and are based on registration of subsidies provided for the implementation of drainage or ditches in connection with planting activities. The areas may be categorized as either forest or peatlands. The drained areas for both categories were summarized and accumulated for the years 1950 to 1989. The total accumulated area from 1990 and onwards is used for the reporting under forest land remaining forest land. However, from 1990 and onwards, only forest areas were included in the statistics. Peatlands drained after 1990 are included in land converted to forest land, but the total area in the conversion category is derived from the NFI.

The activity data is further stratified into vegetation zones as suggested in the IPCC 2013 Wetlands supplement. All Norwegian forest land is considered boreal. The vegetation registration in the NFI database was studied in order to determine the distribution of drained organic soils to nutrient rich and nutrient poor. A ditch registration was performed on all NFI plots when the permanent sample plots were established (between 1986 and 1993). The plots were classified as ombrotrophic if one of the three conditions were met: 1) peat soils isolated from natural rivers, streams or springs with spruce and birch forest; 2) hummocks dominated with *Calluna vulgaris* and sphagnum mosses on the bottom, or 3) if hummocks were missing, the vegetation was dominated by *Trichophorum cespitosum*, *Eriophorum vaginatum*, and *Carex pauciflora*. The remaining plots with a ditch registration were classified as minerotrophic peatlands. According to the IPCC 2013 Wetlands supplement, minerotrophic peat soils can be classified as nutrient rich and ombrotrophic as nutrient poor. The results showed that 69 % of all drained plots are nutrient rich and 31 % are nutrient poor. This distribution was applied for estimation of CO₂, N₂O, and CH₄ on forest land remaining forest land.

During the recording of ditches between 1986 and 1993, only ditches less than 25 years of age were recorded. Thus, the NFI contains ditch registration only for ditches established after approximately

1965. A large proportion of the area was drained before this, and that is why the area of drained organic forest soils is not based on the NFI registration but instead on subsidy statistics.

Emission factors

There are no national data on the CO₂ losses due to drainage of organic soils in forest land. We hence used the default emission factors from the IPCC 2013 Wetlands supplement as these represent the most up to date information. The mean national EF derived using the nutrient class distribution described above is 0.72 t C-CO₂ yr⁻¹ ha⁻¹.

Undrained organic soils

Organic soils on forest land not subject to drainage were assumed to be in equilibrium. No methods are available for the estimation of the C emissions or removals on these areas. Based on NFI registrations since 1990, final harvest or thinning was registered on about 8 % of the forest area (NFI definition of forest, i.e. including areas in conversion in UNFCCC terminology) on organic soil not subject to drainage and on 22 % of the forest area on mineral soils. Thus, the forestry activity on areas with undrained organic forest soils is relatively low. A study was carried out to survey existing empirical evidence on C emissions/removals from undrained organic forest soils. A total number of 30+ publications reporting on open and tree-covered bogs and fens in countries including Finland, Sweden, Canada, and Russia were included in the survey. The overall conclusion was that these areas have been long term C sinks (for millennia; based on peat column studies) and contemporary rates (short term studies 1-10 years) indicate that they on average and in most years act as sinks, but that they in some (dry) years may act as a source. Where comparisons have been made between open and tree-covered areas, there were no indications that open areas had higher accumulation rates than tree-covered areas. Comprehensive studies include Tolonen and Turunen (1996), Turunen et al. (2002), Roulet et al. (2007), and Nilsson et al. (2008).

6.4.1.2 Uncertainties and time-series consistency

Living biomass

The estimation of uncertainties for C stock changes in living biomass on forest land is described in section 6.3.5 and estimated uncertainties are presented in Table 6.3.

The calculations of carbon stock changes in living biomass are conducted according to the stock change method and are based on data obtained from the NFI. More details are described in section 6.3.4.

Dead organic matter and mineral soils

The uncertainties for dead organic matter and soil organic matter used in the key category analyses are based on Monte Carlo simulations of the national level of total soil organic C change (i.e. soil + litter + dead wood). One thousand simulation loops were run using the same calculation procedures as described above for forest land remaining forest land – mineral soils, but with variability introduced to a number of parameters (Table 6.19).

Table 6.19 Characteristics of the parameters used in the Monte Carlo simulations.

Parameter	Distribution	Mean	Standard deviation (% of mean)	References
Coarse woody litter dimension (cm)	Normal	10	20 %	Expert judgment
Branch and coarse root turnover (yr ⁻¹)	Normal	0.0125; 0.027; 0.025 ^a	20 %; 25 % ^b	Peltoniemi et al. (2006); expert judgment
Fine root turnover (yr ⁻¹)	Lognormal	0.6 ^c		(Brunner et al. 2013) (Hansson K et al. 2013); expert judgment
Foliage turnover (yr ⁻¹)	normal; uniform ^b	0.143; 0.33; 0.9-1.0 ^a	15 %	Peltoniemi et al. (2006); expert judgment
Ground vegetation turnover (yr ⁻¹)	Normal	0.33; 0.1; 1.0; 0.25 (aboveground) ^d 0.33; 0.33 (belowground) ^e	40 %	Peltoniemi et al. (2006)
Biomass ratio for ground vegetation, below-to-above	Normal	2	20 %	Peltoniemi et al. (2006)

^a Spruce, pine and deciduous respectively. ^b Conifers and deciduous respectively. ^c In lognormal: mean -0.51 and standard deviation 0.3. ^d Moss, lichens, herbs/grasses and shrubs respectively. ^e Herbs, grasses and shrubs.

Uncertainty around the Yasso07 model parameters was described in a number of parameter sets (Tuomi et al. 2011b) where covariance among model parameters are taken into consideration. For the C input parameters, a number of parameters were selected because they were assumed to have particularly large uncertainties. The C input parameters were assumed to be independent of each other, but in cases where differences among species or specific components could not be documented, parameter values were drawn from the same distribution. Most of the parameters were assumed to be normally distributed and negative values were avoided by truncated distributions (negative values replaced by 0). The simulations were run with the Yasso07 model, with spin-up loops coded in Fortran, and the litter estimation run with the R software (R Core Team 2015). The result was an uncertainty estimate of the Yasso07 simulated C stock changes reported in 2014 of 15.5 %, which applied to both the DOM and mineral soil pools. The uncertainty is not likely to diverge with the values reported in the current year's NIR. The simulations are illustrated in Figure 6.13.

Uncertainties in the biomass models (Table 6.) and the diffuse harvest and mortality frequencies underlying the C input estimates to Yasso07 are currently ignored; mainly for technical reasons. However, we believe that most of the uncertainty associated with the current methodology is captured.




Figure 6.13 Results of the 1000 Monte Carlo simulation runs (blue lines) and 95 % confidence intervals (red lines and circles).

Drained organic soils

Default uncertainties of the emissions factors from the IPCC 2013 Wetlands Supplements were applied, and uncertainties of the areas were estimated by sample error. See Table 6.4.

6.4.1.3 QA/QC and verification

The Tier 1 QC procedures were followed for all source categories. Since the method to estimate C stock changes in living biomass was not generally changed, external QA was not necessary. The area estimates were carried out by two independent experts using two different statistical software systems based on the same database. Similarly, the carbon change estimates were compared on a sample basis.

The NFI database has QA/QC procedures as explained in section 6.1.6. For estimation of C changes in mineral soils on forest land, all input was kept strictly to one unit (kg C m^{-2}). An area based unit makes it easier to compare estimates with those from other studies and regions. Specific attention was given to unit conversions particularly when data were moved from one platform to another. The input data was screened for inconsistencies, i.e. occurrence of null-data/missing data, length of input objects etc. Plot specific C input, scaled in the expected manner with total plot standing biomass and plot specific soil organic C changes, had the expected dynamics (i.e. on average C change on the plot level was negative or low in young stands vs. medium age stands). Average litter input and total SOC output time series were compared with the time series used in the previous year's NIR. The estimated C stocks were low compared to field measurements (de Wit & Kvindesland 1999). Studies with an earlier version of Yasso (de Wit et al. 2006), showed that the model estimated about 40 % of the measured forest soil C stock in southeast Norway. This was suggested to be due, in part, to an overestimation of decomposition rates for recalcitrant organic matter. Comparison of the current model methodology and measured soil C stocks confirmed that Yasso07 underestimates measured

stocks and that it may be related to specific soil types and moisture conditions (Dalsgaard, L et al. 2016). The area-based estimates of C change from the current application of Yasso07 were in the range observed in Liski et al. (2005) and Häkkinen et al. (2011). Conclusions from a validation project on soil C changes are found in Dalsgaard et al. (2017). The data from NFI dead wood registrations were supplemented by assumptions generally based on statistics and published quantitative factors (see footnote under 6.4.2.1) and were used to make an alternative calculation (validation) for a reference stock for C in dead wood in forests.

The programming methodology (programming software “R”) was characterized by: 1) a step-by-step development of functions, 2) checking the reproducibility of new functions (new code), and 3) close cooperation among programmers/developers. Code development and code control was done by different people.

In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.4.1.4 Recalculations

Living biomass

The time-series was recalculated due to updates in the NFI database and the extrapolation method.

Dead organic matter and mineral soils

The time-series was recalculated due to updates in the NFI database and the extrapolation method.

Drained organic soils

In the 2019 submission, a correction of the proportion of nutrient rich and nutrient poor organic soils lead to a slight decrease of the carbon stock change.

6.4.1.5 Planned improvements

Dead organic matter and soils

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.4.2 Land converted to forest land, 4A2

Land converted to forest land occurs from all land-uses, but with the largest areas from other land, settlements, and grassland. In the case of settlements, there are many types of settlements that have been converted to forest land in Norway. These can roughly be divided into four groups: power lines, roads, extraction (i.e. gravel, sand, and mining), and other. Estimates of C stock changes are provided for living biomass, dead organic matter (DOM), mineral soils, and organic soils for all relevant conversions. Conversion from other land to forest is almost exclusively due to natural succession.

6.4.2.1 Methodological issues

Living biomass

When a stand of trees reaches the predetermined minimum width, size, and crown cover in the forest definition, the stand is measured by the NFI. Estimates of the carbon stock change in this category are carried out as for the category forest land remaining forest land (see section 6.4.1.1).

Dead organic matter (key category)

Grassland, settlement, and cropland to forest are identified as key category for Level assessment 1990, Level assessment 2017 and/or Trend assessment 1990-2017 (see Table 6.6).

Choice of method

A Tier 2 method is used for estimating C changes in dead organic matter (DOM) for land converted to forest land. The method is based on a C stock change rate multiplied by the area under each land-use conversion.

Carbon stock change factors

Carbon stock change factors were estimated specifically for cropland, grassland, wetlands, settlements, and other land converted to forest land. The C change rates were calculated as the sum of the rates for the dead wood and litter pools and based on a C stock estimate that was assumed to be reached within 20 years, according to the default value stock change dependency. A reference stock for forest litter (61 t C ha^{-1}) was estimated as the average C density (t C ha^{-1}) in the L (litter), F (fermentation), and H (humus) layer of 893 forest soil profiles classified as mineral soil types or dry organic soils (Folisols) (de Wit & Kvindesland 1999; Esser & Nyborg 1992; Strand et al. 2016). Profiles were classified according to the Canadian soil classification system and the soil types were Podzols (443), Brunisols (158), Gleysols (76), Regosols (95), Hemic Folisols (35), and Nonsoils (20). Due to the field registration methodology, an LFH layer was not distinguished for Folisols, rather the whole C profile was assigned to the litter pool. Bulk density was found from Norwegian forest soils (Strand et al. 2016). An average reference stock for C in dead wood in forest (5 t C ha^{-1} , Stokland pers. comm) was based on expert judgment³¹.

For all land-use conversions, except from other land, we assumed that the full litter stock of 61 t C ha^{-1} would develop over 20 years, resulting in a change rate of $3.05 \text{ t C ha}^{-1} \text{ yr}^{-1}$ and 10 % of the reference dead wood stock resulting in a change rate of $0.025 \text{ t C ha}^{-1} \text{ yr}^{-1}$. The major part of the conversions from other land to forest land is on wooded land of low productivity. For this conversion, the annual stock change rate was limited to a 5 % relative build up compared to the stock on the previous land use, which resulted in a change rate for litter of $0.15 \text{ t C ha}^{-1} \text{ yr}^{-1}$ and for dead wood of $0.013 \text{ t C ha}^{-1} \text{ yr}^{-1}$ (Table 6.20).

³¹ Based on a series of assumptions: a mean dead wood volume of $8.3 \text{ m}^3 \text{ ha}^{-1}$ (NFI registration), a weighted volume to biomass factor of 0.44 distributed to decay classes 1-5 from NFI registrations, dry biomass densities from Næsset 1999 (for individual decay classes), 50% C, expansion factors to estimate stump and belowground deadwood from NFI data. Further, a constant annual harvest since ca. 1900 of 10 mill m^3 stem wood was assumed (based on Statistics Norway, see Figure 6.4) from which belowground deadwood from harvested trees was estimated. It was also assumed that decomposition rates were identical for all dimensions, climatic regions, and belowground decomposition equaling aboveground decomposition. Dimensions of < 10 cm and dead wood older than 101 years were ignored. The result was $4.5\text{-}5.5 \text{ t C ha}^{-1}$ depending on the decomposition rate (Næsset 1999, Melin et al. 2009). To complement these calculations, Yasso07 simulations showed an overall mean of 4 t C ha^{-1} in forest originating from coarse woody litter.

Table 6.20 Annual stock change rates ($t\ C\ ha^{-1}\ yr^{-1}$) for land converted to forest land.

	Soil	DOM	Litter	Dead Wood	Total
(t C ha ⁻¹ yr ⁻¹)					
Cropland	-1.30	3.08	3.05	0.03	1.78
Grassland	-2.05	3.08	3.05	0.03	1.03
Wetlands	-1.50	3.08	3.05	0.03	1.58
Settlements	0.57	3.08	3.05	0.03	3.65
Other land	0.14	0.17	0.15	0.013	0.31

Activity data

The total areas of land converted to forest land were estimated by NFI data, which also corresponded to mineral soil areas.

Mineral soils (key category)

Grassland converted to forest land is identified as a key category for, Level assessment 2017 and Trend assessment 1990-2017 (see Table 6.6).

Choice of method, C stock change factors, and activity data

We used a Tier 2 method based on soil organic carbon (SOC) stock change rates multiplied by the area pertaining to each land-use change. The SOC stock change rates were derived by subtracting the mean national soil C stock for the previous land use from the stock of the current land use and dividing this difference by 20 years according to the IPCC methodology. The mean SOC stocks for forest land and cropland were based on measurements. For wetlands, the IPCC default SOC reference stock value was used. For grasslands, the IPCC default stock was modified using national data (see section on Land converted to grassland). No data were available for other land and it was assumed that conversion from other land resulted in a small positive stock change as described above for DOM.

The national forest mean SOC stock estimate was $57\ t\ C\ ha^{-1}$ based on the same forest soil database ($n=893$) as described above for the DOM pool. Upscaling to a depth of 30 cm was made on the basis of field registrations and bulk density was estimated from the function of Baritz et al. (2010). Only mineral soil horizons were included. For non-soils where no differentiation between LFH and mineral horizons were made, all C in the profile was assumed to belong to the IPCC soil pool. The mean SOC stock estimate for cropland was $83\ t\ C\ ha^{-1}$, for grassland $98\ t\ C\ ha^{-1}$, and for wetlands $87\ t\ C\ ha^{-1}$. The resulting SOC change rates are shown in Table 6.20. Due to the lack of data, we assumed that the mean SOC stock for settlements was equal to 80 % of the relevant land use and thus a 20 % SOC loss over 20 years. The areas of land converted to forest land on mineral soils were obtained from the NFI.

Drained organic soils

For conversions to forest land on organic soils, we used a Tier 1 methodology applying the default emission factor for boreal and nutrient rich vegetation zone provided in the IPCC 2013 Wetlands supplement of $0.93\ t\ C\ ha^{-1}$. We assumed that organic soils previously used for grassland, cropland, wetlands, and settlements are drained. The activity data (areas) for organic soils converted to forest land was derived from the NFI.

6.4.2.2 Uncertainties and time-series consistency

Generally, the uncertainties related to emission estimates for all sinks/sources were rather large, partly due to the uncertainty of the area estimate. Uncertainties are shown in Table 6.3 for living biomass and DOM and in Table 6.4 for mineral and drained organic soils.

The time-series were consistently estimated.

6.4.2.3 QA/QC and verification

The internal QA/QC plan was completed as relevant for all source categories under land converted to forest. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.4.2.4 Recalculations

The recalculations described in section 6.4.1.4 apply. Areas and living biomass were updated with the availability of new data from the NFI as part of the extrapolation method. Recalculations of the emissions from organic soils were also due to an update in the area statistics from Statistics Norway.

In 2019, the rectification of the proportion of nutrient rich vs. nutrient poor organic soil from 79-21 % to 69-31% induced a decrease of the CO₂ emissions under forest.

6.4.2.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.4.3 Completeness

The reporting of emissions and removals from forest land is complete.

6.5 Cropland – 4B

Agricultural cropland in Norway includes annual crops, temporary grass leys, and horticulture. Most of the area for agriculture is used for annual crops, primarily consisting of grass leys (55 %) used as forage or green manure, cereals (37 %) and a smaller area with root crops (2 %) where potatoes and rutabagas are the most important crops. Consequently, carbon is not stored for long time intervals in aboveground biomass. An exception is horticultural crops, where fruit trees can store large amounts of C. However, the area of perennial woody crops is a small fraction of the cropland area (approximately 0.2 %).

Substantial amounts of C reside in the soil, which is affected by agricultural management practices such as tillage, crop residues input and organic manure application (Paustian et al. 2000). Dead organic matter is not an important source category for cropland in Norway, since agroforestry systems are uncommon. This is with the exception of forest land converted to cropland, where emissions are reported. Over the time-series the total area of cropland has decreased on a national scale, despite that conversion to cropland also occurs, primarily from forest land but also from wetlands on organic soils. CO₂ emissions from living biomass, DOM, mineral and organic soils on croplands are reported in CRF Table 4.B as described below, and CH₄ emissions from organic soils on croplands are reported in CRF Tabel 4(II) as described in section 6.12.2.

6.5.1 Cropland remaining cropland, 4B1

The following emission sources were reported under cropland remaining cropland: C stock changes in living biomass of perennial horticultural crops (fruit trees), C emission from mineral soils due to agricultural management (crop rotations, C inputs, and tillage) and C emission caused by cultivation of organic soils. By far, the vast majority of emissions are caused by the cultivation of organic soils, which is a key category because of the uncertainty in the level and trend (see section 6.1.4). Net C gains are reported for living biomass and mineral soils.

6.5.1.1 Methodological issues

Annual changes in C stocks on cropland remaining cropland can be estimated as the sum of changes in living biomass and soils by $\Delta C_{CC} = \Delta C_{LB} + \Delta C_{SO}$. Norway applies the Tier 1 steady state assumptions for dead organic matter because agroforestry is generally not practiced. Thus, the agricultural systems have small amounts of dead organic matter. Living biomass is reported for fruit trees and emissions from soils are reported for mineral soils and organic soils.

Living biomass

Changes in C in living biomass are only considered for perennial woody crops, i.e. fruit trees. Perennial berry bushes are not considered due to the small area of approximately 300 ha (Borgen & Hylen 2013). Orchards may be felled but are considered to remain cropland. It is likely that orchards are converted to annual crops, leys or vegetables, or are replanted with fruit trees. Annual changes in the area of fruit trees fluctuate, leading to both net emissions and removals during the inventory period. However, C stock changes are relatively small.

Choice of method, emission factors, and activity data

Due to a lack of national data on biomass and carbon content in Norwegian fruit trees, we apply the Tier 1 gain-loss method. In the default method the change in C stock in living biomass (ΔC_{LB}) is equal to the C gain (ΔC_G) minus the C loss (C_L) by $\Delta C_{LB} = \Delta C_G - \Delta C_L$.

Statistics Norway collects data every year on the areas of fruit trees (apples, plums, cherries, sweet cherries and pears). The data are collected as a questionnaire survey with the objective of providing information about yields and production area. We use these data collected for the whole time-series 1990-2017. The area of fruit trees has generally decreased since 1990.

The IPCC default value for biomass accumulation in the temperate climate is $2.1 \text{ t C ha}^{-1} \text{ yr}^{-1}$, and the corresponding value for C loss when plantations are terminated is $63 \text{ t C ha}^{-1} \text{ yr}^{-1}$. The default age for fruit trees to reach maturity and cease accumulating C is 30 years.

Assumptions/justification

Given the default method, we assume that: 1) all orchard trees are less than 30 years old, and that growth accumulates at the default growth rate; and 2) all felled orchards are plantations with mature trees around 30 years of age. These assumptions may not be representative for Norway, as Norwegian fruit trees may mature in 20-25 years. However, the activity data does not provide information on the age of the plantations when felled.

Dead organic matter

The Tier 1 method was used assuming no carbon stock change in the dead organic matter pool on cropland remaining cropland and the notation key NO is reported in the CRF tables.

Mineral soils

The majority (roughly 94 %) of agricultural production occurs on mineral soils. Management practices have changed relatively little since 1990. Carbon inputs from animal manure have slightly increased in some parts of the country resulting in C uptake, i.e. positive carbon stock changes.

Choice of method

The Tier 2 method estimates annual changes in soil organic C (SOC) according to Equation 2.25 (IPCC, 2006a), where the annual change in SOC is given by $\Delta \text{SOC} = (\text{SOC}_0 - \text{SOC}_{0-T})/D$, where D is the time dependency of the stock change factors. SOC_0 is the stock the last year of the inventory period and SOC_{0-T} is the C stock at the beginning of the inventory period. The default value for D was adjusted to 30 years, given the slower decomposition rates under the cool temperate climate in Norway (Borgen et al. 2012). The SOC stock is calculated as the product of the soil C reference stock (SOC_{REF}), the stock change factor for a given management and climate regime (F), and the associated area (A) given by $\text{SOC} = \text{SOC}_{\text{REF}} \times F \times A$. We used the reference stock and stock change factors estimated by the Introductory Carbon Balance Model (ICBM) in a study where CO_2 emissions were estimated for Norwegian cropland for 1999-2009 (Borgen et al. 2012). The ICBM is an ecosystem model from Sweden developed by Andrén et al. (2004). Soil C reference stocks were estimated for 31 different climatic zones (agrozones) assuming that continuous grass ley cropping was the reference condition. Stock change factors were calculated for eight rotations with and without manure application for each of the 31 agrozones, resulting in a total of 496 stock change factors. The rotations were 1:2 ley-grain, 1:1 ley-grain, 2:1 ley-grain, continuous grain (with and without straw removal), continuous ley, 1:2 roots-grain, and 1:2 roots-ley, where 1:2 means 1 year of root crops and 2 years of ley and so on.

Further details of the model application and the stratification are given in Borgen et al. (2012). We calculated annual SOC changes per agrozone and summarized the CSC for the whole country.

Stock change factors and soil C reference stocks

The stock change factors represent the annual response of SOC to a change in management from a reference condition and can be calculated as $F = \text{SOC}/\text{SOC}_{\text{REF}}$. The soil C reference stocks were estimated by solving the ICBM model for steady state conditions using C input equal to continuous ley cropland for each Norwegian agrozone. Average national stock change factors for the 16 crop rotations and average soil C reference stocks per agrozone are listed in Borgen et al. (2012).

Activity data

Area per crop type and manure statistics are collected annually from the Norwegian Agricultural Authority (NAA) to determine the area under each of the 16 crop rotations. Norway was divided into 31 agrozones based on a combination of counties (*fylke*) and climate-based production zones (defined by NAA for subsidy application) as described in Borgen et al. (2012), and the statistics are given for each agrozone. Within each agrozone, the relation between the major crops of small grains (cereal and oilseeds), root crops (potato and rutabaga), and grass ley were used to allocate the areas under each of the eight crop rotations. The activity data of the quantity of manure applied to fields with annual crops per county were received from Statistics Norway. These data correspond to the data used for estimating non-CO₂ emissions related to animal manure for the Agricultural sector. As it is not possible to differentiate between manure applied to perennial grass leys on cropland from the grass leys on grassland, we only include the manure applied to annual crops. Further information on assumptions regarding the manure distribution is given in Borgen et al. (2012). The areas of cropland remaining cropland on mineral soils were estimated by the NFI for the whole time series.

Assumptions

The IPCC Tier 1 and 2 methods assume that the SOC change resulting from a change in management is linear between two steady states. Soil C changes are likely to be more dynamic, and it has been argued that the lower tier methods overestimate net C sequestration, particularly where the soil was not in a steady state at the beginning of the inventory (Sanderman & Baldock 2010). However, at the present time, this method provides an acceptable approximation.

Organic soils (key category)

Organic soils make the largest contribution of CO₂ emissions within the source categories for cropland. It is a key category with a relatively large uncertainty. The Norwegian definition of organic soils for cropland is described in detail in section 1.2.3 Table 6.11.

Choice of method and emission factor

A Tier 1 method is used for estimation of CO₂ emissions from organic soils on cropland. The IPCC Tier 1 method necessitates the use of the default emission factor (EF) to be multiplied by the area (A) of organic cultivated soil according to Equation 2.26 $C_{\text{LOSS}} = A \times \text{EF}$ (IPCC 2006). Since the 2015 submission, Norway has used the default EFs from the IPCC 2013 Wetlands supplement (IPCC 2014b) for boreal/temperate cropland of 7.9 t CO₂-C ha⁻¹ yr⁻¹. For Norway we considered the default value from IPCC to provide a more robust and suitable estimate than the EF based on expert judgment which was used until NIR 2014.

Activity data

The area of agricultural organic soil was estimated as described in section 6.3.2.

6.5.1.2 Uncertainties and time-series consistency

Estimation of uncertainty is related to the tier level of the methodology used for each sink/source category and land-use category. For cropland remaining cropland, Tier 1 and 2 methods were applied. The IPCC guidelines include uncertainty estimates for default emission/removal factors.

For cropland remaining cropland, the total uncertainty is equal to the propagation of the uncertainty related to the living biomass (U_{CC_LB}), mineral soils (U_{CC_MS}), and organic soils (U_{CC_OS}) and is given by

$$U_{CC} = \sqrt{U_{CC_LB}^2 + U_{CC_MS}^2 + U_{CC_OS}^2}.$$

For each source category, the uncertainty is a combination of the uncertainties related to the emission factors U_{EF} and the activity data U_A , which is the uncertainty used in the KCA, and can be calculated by

$$U = \sqrt{U_A^2 + U_{EF}^2}.$$

The uncertainty of the activity data may include errors in census returns as well as differences in the definition between agencies, sampling design, and interpretation of samples. The activity data used under cropland, i.e. areas per crop types and manure production were collected through the subsidy application scheme administrated by NAA and compiled by SSB. The data is based on a total national census. The NAA performs quality control on 5 % of farms to determine if areas are provided correctly. These sample checks show very few errors. The area reported is based on a factor value multiplied by the last year's area, thus errors in previous years may accumulate. However, according to expert judgment given by SSB, the uncertainty of the activity data is estimated to be approximately 0 %.

Living biomass

Sources of uncertainty for the Tier 1 method for living biomass includes the degree of accuracy in the C accumulation and loss rates and the land-use activity data. The IPCC default uncertainty error ranges for aboveground woody biomass accumulation in the temperate climate is ± 75 % based on expert judgment. Uncertainty of the activity data was estimated by SSB as approximately 0 %. The areas of orchards are used directly from the NAA/SSB data and are not related to the NFI database. The uncertainty of the C biomass accumulation per unit area is therefore equal to the total uncertainty of the C changes in living biomass on cropland remaining cropland.

Mineral and organic soils

Uncertainty related to emission estimates from soils on cropland can currently only be precisely quantified for the area estimates, which is based on the NFI data. For the area of mineral soils on cropland remaining cropland, the uncertainty estimate was 7 % (Table 6.4). For the mineral soil estimates, the areas per crop type that are used to determine the areas under individual crop rotations were collected and compiled by the Norwegian Agriculture Authority (NAA) and Statistics Norway (SSB). Since the data are based on a census, it was assumed not to increase the area uncertainty. The uncertainties related to the stock change factors estimated by ICBM were estimated at ± 50 % based on expert judgment, making the total uncertainty 51 %. For organic soils the area

uncertainty was 25 % and default uncertainties were used from the guidelines. Total uncertainty for CO₂ emissions from drained organic soils on cropland was 32 % (Table 6.4).

6.5.1.3 QA/QC and verification

The standard Tier 1 QC procedures described in section 6.1.6 were performed for both living biomass and soil estimates. No external QA was performed on the Tier 1 method for estimating C changes in living biomass stocks in orchard trees. Before the 2013 submission, when the Tier 2 for mineral soils on cropland remaining cropland was implemented, quality assurance was done through the standardized peer-review process.

In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.5.1.4 Recalculations

Recalculations for CSC in living biomass were made for the year 2016 only due to an update in the activity data from Statistics Norway. The area times-series of both mineral and organic soils were also slightly recalculated due to the annual NFI updates and extrapolation of area data. The identification of mistakes during the 2019 QA lead to recalculations for the CSC on mineral soil. In brief, the carbon stocks of silty loam soils were added twice by mistake when summing the carbon stocks of all soil categories. A conceptual mistake of little impact lead to an erroneous distribution of areas between crop rotations categories.

6.5.1.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.5.2 Land converted to cropland, 4B2

Emissions and removals on land converted to cropland are reported from the C stock changes in living biomass, dead organic matter, mineral soils, and organic soils. Carbon stock changes in dead organic matter on other land-use conversions than those to and from forest land can be considered insignificant and are reported with the notation key NO in the CRF-reporter.

Land conversion to cropland primarily occurs from forest land and less so from grassland, wetlands, and settlements. There were no conversions from other land to cropland during the inventory period, and NO is reported in the CRF Table 4.B. Conversion of land to cropland usually results in a net loss of carbon from living biomass and soils to the atmosphere (IPCC 2003). However, the reference soil C stock on settlements and forests is relatively small compared to cropland, and thus net C sequestration is reported, while losses are estimated in the DOM pools for forest land converted to cropland.

6.5.2.1 Methodological issues

Living biomass

For forest land and wetlands converted to cropland, we used the Tier 3 method described for forest land to estimate C stock changes in living biomass. For settlements converted to cropland a Tier 1 method that assumes the immediate gain of the default C stock on settlements of 4.7 t C ha^{-1} changed area (IPCC 2006, p. 8.18) was applied. The category grassland converted to cropland is very small. Nonetheless, as of 2007, changes in the woody biomass are monitored also on grassland. No change in the living biomass was observed for those sample plots converted from grassland to cropland. In addition, the default emission factors for grassland and cropland are so similar that NO is reported for the whole time series. Formally, this is a Tier 2 method.

Dead organic matter (key category)

Carbon stock changes in the dead organic matter (DOM) pool on forest land converted to cropland is a key category for Level assessment 2016 and/or Trend assessment 1990-2016 (see Table 6.6).

Choice of method, C stock change factors, and activity data

A Tier 2 method was used to estimate C stock changes in DOM from forest land converted to cropland. No changes have been made in the method since the 2015 submission. The method is based on a C stock change rate multiplied by the area of forest land converted to cropland as described under land converted to forest land – dead organic matter. The mean C change rate was $-3.3 \text{ t C ha}^{-1} \text{ yr}^{-1}$ based on the assumption that all litter and dead wood in an average Norwegian forest would be lost over 20 years. Areas of land converted to cropland were estimated using the NFI data.

For grassland, wetlands, and settlements converted to cropland, we used the Tier 1 method that assumes no carbon stock change in the DOM pool. Emissions are reported as NO.

Mineral soils (key category)

The sink/source category mineral soil on forest land converted to cropland was identified as a key category for Level assessment 1990 and Trend assessment 1990-2016 (see Table 6.6).

Choice of method and C stock change factors

We used a Tier 2 method for estimating C stock changes in mineral soil on land converted to cropland. The same method was used for all land-use conversions and described under forest land in section 6.4.2.1. It is based on annual stock change rates multiplied by the area. The stock change rates were derived from the difference between the mean stock of the previous land use and the cropland stock divided by 20 years according to IPCC default methodology. For settlements we assumed the stock was equal to 80 % of the cropland stock, i.e. a 20 % relative increase in SOC over 20 years. For forest land converted to cropland, the stock change rate for the mineral soil was positive, indicating an uptake of SOC. However, the loss rates in the DOM pool were larger and the net result for the two pools combined was a net C loss (Table 6.21).

Table 6.21 Annual stock change rates ($t\ C\ ha^{-1}\ yr^{-1}$) for land converted to cropland.

	Soil	DOM	Litter	Dead Wood	Total
(t C ha ⁻¹ yr ⁻¹)					
Forest land	1.3	-3.30	-3.05	-0.25	-2.00
Grassland	-0.75	0			-0.75
Settlements	0.83	0			0.83

The mean national SOC stock for cropland was estimated based on 1 418 soil profiles made throughout the country from 1980 to 2012. The data are a compilation of several different sampling projects where soil profiles were examined using an auger, and soil type and thickness were recorded at different horizons. The organic carbon concentration was measured by dry combustion analysis. To estimate the national mean C stock, the C density was calculated per soil horizon and summarized down to 30 cm depth based on the bulk density function for Norwegian cropland from Riley (1996) and assumed zero weight percentage of gravel. The mean national C stock for Norwegian cropland was 83 t C ha⁻¹. The forest stock equal to 57 t C ha⁻¹ was also based on measurements (see section 6.4.2.1), whereas the grassland (98 t C ha⁻¹) and the wetlands (87 t C ha⁻¹) stocks were derived from IPCC default reference values (see their respective section for details).

Activity data

Areas of land converted to cropland on mineral soils were estimated using the NFI data and the 1990 baseline map of soil types.

Organic soils (key category)

Forest land and wetlands converted to cropland on organic soils were determined to be key categories for Level assessment 2017 (see Table 6.6).

Choice of method and emission factor

We used a Tier 1 method to estimate emissions from organic soils on land converted to croplands. The default emission factor of 7.9 t C ha⁻¹ yr⁻¹ was applied, assuming similar emissions as for cropland remaining cropland, and regardless of the previous land use.

Activity data

All areas were derived as described in section 6.3.2.

6.5.2.2 Uncertainties and times-series consistency

Uncertainties were estimated as described in section 6.1.3 and are shown in Table 6.3 for living biomass and DOM, and in Table 6.4 for mineral and organic soils.

6.5.2.3 QA/QC and verification

The Tier 1 QC procedures were performed during the estimation of C stock changes for land converted to cropland. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.5.2.4 Recalculations

Carbon stock changes of all pools were recalculated because of the revised area data. For conversions from forest to cropland, the recalculations described in section 6.4.1.4 apply.

6.5.2.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.5.3 Completeness

The reporting of emissions from cropland is complete.

6.6 Grassland – 4C

Grasslands cover a very small (approximately 0.7 %) part of Norway. According to the IPCC guidelines, grasslands are defined as grass areas that have insufficient woody biomass to be classified as forest land and that are not considered cropland (IPCC 2006). However, if grazing as a land use is considered more important than forestry, the NFI (National Forest Inventory) classifies a plot as grassland even if the forest definition is met. Grasslands also include range lands and pastures where some mechanical surface harvesting for fodder may take place. The Norwegian interpretation of the IPCC land-use category grassland, which is based on available data, is that grasslands are generally mechanically harvested or grazed, but cannot be plowed. They may be cultivated more or less intensively by the use of fertilization, mechanical harvesting, and utilization of improved species.

In the national agricultural statistics collected through the subsidy application scheme, two types of grassland areas can be identified. These are surface-cultivated grass pastures (*overflatedyrka eng*) and unimproved grazing land (*innmarksbeite*). Surface-cultivated pastures tend to have shallow topsoil layers, often with surface rocks. They can be mechanically harvested but cannot be plowed. Unimproved grazing lands cannot be mechanically harvested (or plowed) and are considered semi-natural landscapes. Furthermore, unimproved grazing land is defined as areas covered by a minimum of 50 % grasses or grazable herbs and enclosed by a fence or a natural barrier. An additional requirement for both grassland types is that the area must be grazed or harvested at least once a year to be eligible for subsidy support.

6.6.1 Grassland remaining grassland, 4C1

For grassland remaining grassland, C stock changes are reported for living biomass and mineral and organic soils. Grassland remaining grassland is a relatively small key category with respect to living biomass according to the level assessment of 1990 and 2017.

6.6.1.1 Methodological issues

Emissions due to changes in dead organic matter are assumed negligible for this category, because little dead wood and litter are generated in grassland systems. Assuming that C stock change in DOM is in a steady state condition is in accordance with the IPCC 2006 guidelines, and the notation key NO is used in the CRF tables.

Living biomass

A Tier 2 method was used. Living biomass on grassland has been measured in the NFI since 2007. The average C stock change (gains and losses) per hectare and year was calculated based on the sample plots measured since 2007. The C gain and loss averages were then multiplied with the area of grassland remaining grassland to obtain the total gain and loss estimates, respectively.

Mineral soils

Choice of method

The default Tier 1 approach was used for estimating CO₂ emissions from grassland remaining grassland on mineral soils. The default IPCC methodology estimates soil C changes based on default stock change factors specific to management and climate regimes and soil C reference stocks specific to climate and soil type. The annual changes in SOC can be calculated as the difference between the

SOC stock in the last year (SOC_0) and in the beginning (SOC_{0-T}) of the inventory period (T) divided by the time dependency of the stock change factors (D) and is given by

$$\Delta SOC = (SOC_0 - SOC_{0-T})/D \quad \text{Equation 2.25 (IPCC 2006).}$$

The time dependency of the stock change factors is by default 20 years. If T is larger than D, then T replaces D and T is equal to the length of the inventory period. This is relevant for the emissions estimated from 2011 and onwards. SOC stocks for any year of the inventory can be calculated as the product of the soil C reference stock (SOC_{REF}), the stock change factors (F), and the area under a given management practice (A) according to

$$SOC = SOC_{REF} \times F \times A \quad \text{Equation 2.25 (IPCC 2006).}$$

The C reference stock is the soil C stock under the reference condition, which in the default method is native uncultivated soil. The reference stock is specific to climate zone and soil. Exposed bedrock should be assigned a reference stock of zero, however, this is not specifically accounted for.

Activity data

Areas of the two grassland management types were collected by Statistics Norway. These data were collected from farmers' applications for subsidies. Areas of unimproved and improved grassland are given per farm unit. The total area of grassland remaining grassland on mineral soils came from the NFI database. The percentages under each management type were taken from data by Statistics Norway (SSB) and applied to the area of mineral soil. Due to different methodological approaches, the area estimated by NFI is larger than the area reported by SSB (Table 6.). The difference is larger in the beginning of the inventory period than later, which is partly because the area of unimproved grassland in the SSB data only accounted for fertilized pasture from 1990 to 1997, whereas all unimproved pastures were included in the later years. In general, the area of extensively-managed grassland (unimproved) has increased, while intensively managed (improved) grazing lands have decreased.

Table 6.22 Areas (ha) of unimproved, improved, and total grasslands in Norway from 1990 to 2017.

	Area (ha) from Statistics Norway (SSB)			Area (ha) from the NFI database
Year	Unimproved grassland	Improved grassland	Total grassland	Grassland remaining grassland (mineral soil)
1990	81 357	27 180	108 537	225 792
1991	85 453	26 973	112 426	225 107
1992	89 735	27 153	116 888	224 423
1993	94 215	25 975	120 190	223 738
1994	98 422	26 050	124 471	222 874
1995	100 719	26 447	127 166	221 979
1996	103 008	26 672	129 681	220 997
1997	107 900	25 478	133 378	219 699
1998	111 474	29 179	140 653	218 113
1999	121 606	29 517	151 123	216 888
2000	129 133	28 997	158 129	215 752
2001	132 293	28 244	160 536	214 833
2002	135 408	28 067	163 474	214 274
2003	137 061	27 382	164 443	213 932
2004	139 083	26 951	166 033	213 048
2005	142 407	26 770	169 177	211 895
2006	145 588	26 110	171 698	210 381
2007	149 207	25 375	174 582	208 831
2008	150 810	24 327	175 137	207 010
2009	152 352	22 455	174 806	205 707
2010	155 136	20 704	175 839	204 601
2011	156 452	20 119	176 571	203 676
2012	156 407	20 128	176 535	202 895
2013	156 436	19 953	176 389	202 250
2014	155 782	19 851	175 633	201 640
2015	156 106	19 767	175 873	201 292
2016	155 986	19 517	175 503	201 034
2017	160 973	19 242	180 215	225 792

The grassland areas per management type were stratified into eight regions (Figure 6.14). The area data from SSB are available on a municipality level facilitating the stratification. Soil maps were collected to stratify the areas according to soil type and to assign specific C reference stocks based on the distribution of soil type within each region.




Figure 6.14 Eight regions of Norway used to stratify grassland activity data for the Tier 1 application.

Stock change factors and soil C reference stocks

The default stock change factors developed by Ogle et al. (2004) were used; see Table 6.2 (IPCC 2006). The land-use factor for grassland is one ($F_{LU} = 1$). There are four management factors (F_{MG}): unimproved/nominal (non-degraded), moderately degraded, severely degraded, and improved grasslands. There are two input factors (F_I): nominal and high input level. For the two types of grassland management identified (unimproved and improved) we assigned the following management factors: $F_{MG} = 1$ for nominally managed (non-degraded) grassland for permanent unimproved grass (i.e. *innmarksbeite*), and $F_{MG} = 1.14$ for improved grassland with surface cultivated grassland (i.e. *overflatedryka eng*). The latter factor is assigned to grassland that is sustainably managed with moderate grazing pressure and that receives one improvement of fertilization, species improvement, or irrigation. The input factor is not modified due to a lack of detailed activity data for manure management on the two grass types. Under Norwegian conditions, it is a reasonable assumption that most grassland receives only one improvement in the form of fertilizers annually, as grazing areas are seldom reseeded (except in cases of severe frost damage) and also irrigation is generally not practiced.

To assign the soil C reference stock, an analysis was made of the national soil classification (World Reference Base, WRB, soil taxonomy) database developed by the Norwegian Institute of Bioeconomy Research. The percentage of the total grassland area that has been sampled until now varies between the eight strata defined. The results of the analysis were that high-activity clay (HAC) soils predominate in all climate zones, but spodic soils make up almost one third of the area in region 2 (Figure 6.15).




Figure 6.15 Distribution of soil types on grassland areas for the eight strata. The IPCC soil types are high-activity clay soils (HAC): leptosols, fluvisol, phaeosem, albeluvisol, luvisol, umbrisol, cambisol, regosol; wetland soils: gleysols; sandy soils: arenosols; and spodic soils: podzol.

The soil C reference stock (SOC_{REF}) for the cold temperate moist climate zone in 0-30 cm depth are 95 t C ha⁻¹, 71 t C ha⁻¹, 115 t C ha⁻¹, and 87 t C ha⁻¹ for HAC, sandy, spodic, and wetland soils, respectively; see Table 2.3 (IPCC 2006). Soil C stock changes were first calculated per stratum and soil type. The final stock changes were given by multiplying the C stocks per stratum and soil type with the fractions for each soil type.

Organic soils

Organic soils on grassland remaining grassland are responsible for minor CO₂ emissions similar to those from mineral soils.

Choice of method

We used the Tier 1 method described for organic soils in cropland remaining cropland (section 6.5.1).

Activity data

The area of organic soil on grassland remaining grassland was derived in the procedure described in section 6.3.2.

Emission factor

The default EF for shallow-drained, nutrient-rich grassland of 3.6 t C ha⁻¹ yr⁻¹ was applied (IPCC 2014b). The emission factor was changed after studying the orthophotos of the NFI plots of grassland on organic soils. The majority of the plots were either with too many trees to allow proper cultivation and thus overgrown drains, or on fairly thin soil layers (visible bedrock in some places) such as surface cultivated grasslands. We found the shallow drainage factor to be more appropriate.

6.6.1.2 Uncertainties and time-series consistency

The uncertainties were estimated for all sink/source categories under grassland remaining grassland and included in the key category analysis.

For living biomass, the uncertainty estimate of the C stock change and the area were based on the sample variance and estimated as described in section 6.1.3 and is shown in Table 6.3.

For the mineral soil pool, a Tier 1 uncertainty assessment was made considering the uncertainty related to the C stock estimate (the stock change factors) using default values and the activity data using the sample variance. Firstly, we estimated the uncertainty of the SOC stock estimate (U_C) by propagating the uncertainty of the stock change factors and SOC reference stock. The errors of the stock change factors are provided in Table 6.2 (IPCC 2006). For the improved grassland management stock change factor, the uncertainty is $\pm 11\%$. The stock change factor for nominally managed grassland has no associated uncertainty as it is the reference condition. The default C reference stock has an uncertainty of $\pm 90\%$, according to Table 2.3 (IPCC 2006). Secondly, the uncertainty of the activity data was combined with that of the C stock change per hectare. The uncertainty in the activity data (U_A) covers both uncertainty in the estimates of the grassland management type (SSB data) and uncertainty in the areas of grassland remaining grassland determined in the NFI. The first source of uncertainty, which is related to the estimation of the grassland management system, was estimated to be close to zero by SSB. According to the sample validations routinely performed by the collection agency (NAA), farmers are unlikely to make errors (or false reporting) and very few of these errors exist. The second source of uncertainty in the activity data, i.e. of the area estimate of grassland remaining grassland, was determined by the sample error and equal to 14% (Table 6.4). Although the area included organic soils, we assume that the uncertainty for the mineral soil area is similar. Uncertainties of the area estimates are quantified as described in section 6.1.3. The total uncertainty for the mineral soil estimate was propagated using equation 5.2.1 of the Good Practice Guidance (IPCC 2003) and equal to 91% .

The uncertainty for organic soils is based on default values for the emission factor and on the sample error for the area estimate. Uncertainty estimates for both mineral and organic soils are shown in Table 6.4.

6.6.1.3 QA/QC and verification

The Tier 1 QC procedures were performed both for living biomass, mineral soil, and organic soil emission estimates. The Tier 1 method used for mineral soils was elicited for external QA before the 2013 submission. All necessary documentation was supplied to an international expert for an evaluation of the method application and description. The expert emphasized the need to keep the area of grassland remaining grassland constant at the beginning and end of each inventory period when recalculating the entire time-series. Furthermore, quality checks were implemented to ensure that the total land area per stratum remains constant over the time-series. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.6.1.4 Recalculations

The recalculations described in section 6.4.1.4 apply. The whole time-series was recalculated for all sources due to the updates in the NFI data.

6.6.1.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.6.2 Land converted to grassland, 4C2

Emissions from land converted to grassland were primarily caused by net C losses in the DOM pool on forest land converted to grassland. There were land-use conversions from forest land, wetlands and settlements to grassland. For forest land converted to grassland, C emissions were estimated from changes in living biomass, DOM, and soils (mineral and organic). All the area of wetlands converted to grassland was on organic soils. Emissions were therefore estimated for stock changes in living biomass and organic soils for this land use conversion. All the area of settlements converted to grassland was on mineral soils.

Forest land converted to grassland is identified as a key category with respect to living biomass, DOM, and mineral soils, in the Level assessment 2016 and Trend assessment 1990-2016 (see Table 6.6).

6.6.2.1 Methodological issues

Living biomass (key category)

The choice of method, activity data, and assumptions related to the estimation of C stock changes in living biomass on land converted to grassland are identical to those described under forest land. For settlement to grassland conversions, an initial increase to the Tier 1 default C stock of 4.25 t C ha^{-1} (i.e. $8.5 \text{ t dry matter ha}^{-1}$) (IPCC 2006, Table 6.4, p.6.27) is assumed in the first year of conversion.

Dead organic matter (key category)

Carbon stock changes in DOM were reported with a Tier 2 method for forest land converted to grassland. For wetlands converted to grassland and settlements converted to grassland we apply the Tier 1 method that assume no net change in the C pool of dead organic matter, thus the notation key NO is used in the CRF-tables.

Method choice, C stock change factors, and activity data

A Tier 2 or Tier 1 method was used (see above). The areas of land converted to grassland were estimated using the NFI data. The C stock change rate estimate of the DOM pool on forest land converted to grassland was $-3.30 \text{ t C ha}^{-1}$ based on change rates of $-3.05 \text{ t C ha}^{-1}$ and $-0.25 \text{ t C ha}^{-1}$ for the litter and dead wood pools, respectively. The change rates were estimated assuming that a C stock of 61 t C ha^{-1} reduces to zero in 20 years (default value). The estimation of the litter and dead wood stocks are described under forest land.

Mineral soils (key category)

A Tier 2 method is used to estimate C stock changes on land converted to grassland (as well as all other land-use conversion).

Choice of method, C stock change factors, and activity data

The Tier 2 method is based on the multiplication of a C stock change rate with the pertaining area. Carbon stock change rates were estimated as the difference between the soil C stocks per land-use class before and after land-use change, divided by 20 years. The C change rate for forest land

converted to grassland was $2.05 \text{ t C ha}^{-1} \text{ yr}^{-1}$. The estimate of the SOC stock for forest land was 57 t C ha^{-1} based on the measurements as described in section 6.4.2.1. The stock for grassland was based on IPCC default reference stocks per soil type and national area distribution of the soil types. Both stocks were estimated for 30 cm soil depth.

The mean national SOC stock estimate for grassland was 98 t C ha^{-1} and was derived by multiplying the IPCC default stock change factors with the SOC reference stock for average Norwegian grassland. This estimate is based on the national ratio of improved and unimproved grassland management practices and the national distribution of IPCC defined soil types for the grassland area. More specifically, a mean stock change factor was calculated as $F = 0.82 \times 1 + 0.18 \times 1.14 = 1.03$, based on the long-term mean distribution of unimproved and improved grassland (82 % and 18 %, respectively) and the default stock change factors of 1 and 1.14 for unimproved and improved grasslands. A mean SOC reference stock was estimated assuming the following distribution: 85 % high-activity clay soil, 2 % sandy soils, 9 % spodic soil, and 4 % wetland soils (i.e. gleysols), resulting in an estimate of $\text{SOC}_{\text{REF}} = (0.85 \times 95 + 0.02 \times 71 + 0.09 \times 115 + 0.04 \times 87) \text{ t SOC ha}^{-1} = 96 \text{ t SOC ha}^{-1}$ (see section 6.6.1) for details. The mean national C stock for grassland was $1.03 \times 96 \text{ t C ha}^{-1} = 98 \text{ t C ha}^{-1}$.

The C change rate for settlements converted to grassland was $0.98 \text{ t C ha}^{-1} \text{ yr}^{-1}$. It is based on the assumption that settlements reference mineral soil C stocks are 80% of those of grassland (same assumption as for settlements converted to forest).

The areas of land converted to grassland were estimated using the NFI data. To get the area of mineral soil on forest land converted to grassland, the area of organic soils was subtracted from the total area.

Organic soils

Emissions from organic soils on land converted to grassland were estimated using the Tier 1 method. Only wetlands on organic soils have been converted to grassland and these areas were assumed drained to enable grassland production.

Method choice, emissions factors, and activity data

The Tier 1 method was used applying a default emissions factor of $3.6 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for shallow-drained grasslands in the temperate zone from the IPCC 2013 Wetlands supplement. The NFI database was used to estimate the areas of wetlands converted to grassland on organic soils.

6.6.2.2 Uncertainties and time-series consistency

The total uncertainties for living biomass, DOM, mineral, and organic soils are shown in Table 6.3 and Table 6.4. All methods were applied consistently for the entire time-series.

6.6.2.3 QA/QC and verification

The standard Tier 1 QC procedures were performed during the estimation of C stock changes for land converted to grassland. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.6.2.4 Recalculations

The recalculations described in section 6.4.1.4 apply. All emissions of land converted to grassland were recalculated in the current year's submission due to the updates in the NFI data for living biomass and areas.

6.6.2.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.6.3 Completeness

The reporting for grassland is complete.

6.7 Wetlands – 4D

Wetlands in Norway cover almost 12 % of the total land area. Most of the wetlands in Norway are unmanaged mires, bogs and fens, as well as lakes and rivers. Carbon stock changes in living biomass are reported for wooded mires. Managed wetlands include peat extraction areas and reservoirs (dams). For peat extraction sites, both on-site and off-site emissions are reported. On lands converted to wetlands, emissions and removals are reported for living biomass, DOM, and soils. There is no default method for estimating carbon stock changes for flooded land remaining flooded land, we have therefore no estimates for this source.

6.7.1 Wetlands remaining wetlands, 4D1

The NFI contains data on C stock changes in living biomass (trees) on wooded mires, for which the associated emissions and removals have been reported. Carbon stock changes in other sources (DOM and soils) in unmanaged wetlands have not been estimated. Emissions caused by soil C changes during peat extraction have been accounted for according to the 2006 IPCC guidelines (IPCC 2006) and IPCC 2013 Wetlands supplement (IPCC 2014b). On-site and off-site CO₂ emissions from peat extraction (reported as organic soils in CRF table 4.D) is a key category with respect to the 1990 level assessment (Table 6.6).

6.7.1.1 Methodological issues

Living biomass – wooded mires

Wooded wetlands are classified as forest if the definition of forest land is met. If this is not the case, such areas are considered under wetlands remaining wetlands as the subgroup wooded mire. Wooded mires are not considered managed lands, and hence we only report CSC in the living biomass. CSC in DOM, mineral and organic soil are reported as NO.

To estimate CSC in living biomass, we applied the Tier 3 method, which was used for all reported biomass estimates, except for cropland remaining cropland, and land converted to settlements. The method is described in detail in section 6.4.1. The areas of wetlands remaining wetlands and C stocks on wooded mires, are based on the NFI.

Peat extraction (key category)

For wetlands subject to peat extraction we use a Tier 1 approach for on-site emissions and a Tier 2 approach for off-site emissions. Under the default method, the activity data do not distinguish between peatlands under peat extraction, and those being converted for peat extraction (IPCC 2006; IPCC 2014b). The area of land converted to peat extraction is therefore reported as NE. The emissions from removals of trees during clearing are included under living biomass on wooded mires, and reported as IE. Other changes in C stocks in living biomass on managed peat lands are assumed to be zero (IPCC 2006).

The area utilized for peat extraction is estimated to be 2.00 kha for the whole time series 1990 - 2017. On-site emissions caused by peat extraction are thus constant over the inventory period. Soil C stock changes are estimated to be -5.6 kt C yr⁻¹, which is equal to emissions of 20.5 kt CO₂ yr⁻¹. On-site emissions of N₂O and CH₄ are estimated to 0.0009 kt N₂O yr⁻¹ and 0.0658 kt CH₄ yr⁻¹, respectively. Off-site emissions vary with years, and were 40.3 kt CO₂ in 2017, and an average of 40.3 kt CO₂ yr⁻¹ over the time-series (1990-2017). Total emissions from peat extraction, including on-site emissions,

represents 62.7 kt CO₂-equivalents in 2017, and an average of 62.7 kt CO₂-equivalents yr⁻¹ over the inventory period.

Choice of method, activity data, and emission factor

For wetlands subject to peat extraction, on-site emissions are estimated with default emission factors from the IPCC 2013 Wetlands Supplement (boreal / temperate zone), and is hence considered a Tier 1 method. Off-site CO₂ emissions are estimated using a national emission factor of 0.05 t C / m³ based on expert judgment, and it is therefore a Tier 2 method. We assume a peat dry matter density of 0.1 t m⁻³, and C content of 50 %.

Table 6.23 Emission factors used for estimation of on- and off-site emissions from peat extraction and their estimated uncertainties.

Gas	Emission factor (EF)	EF uncertainty (% 2 SE)	Activity data uncertainty %	Total uncertainty %
On-site				
CO ₂	2.8 t CO ₂ -C ha ⁻¹ yr ⁻¹	50	100	110
CH ₄ LAND	6.1 kg CH ₄ ha ⁻¹ yr ⁻¹	80	100	128
CH ₄ DITCH	542 kg CH ₄ ha ⁻¹ yr ⁻¹	81	100	129
Frac _{ditch}	0.05			
N ₂ O	0.30 kg N ₂ O-N ha ⁻¹ yr ⁻¹	113	100	151
Off-site				
CO ₂	0.05 t C m ⁻³ air-dry peat	50	50	71

The reported peat extraction area is based on a 2015 study. Norwegian peat producers were surveyed to provide information on current and previous peat extraction production areas. In addition, a supplemental digital orthophoto sampling of identified ditched marsh areas where peat extraction was likely to have occurred was sampled. Off-site emissions from extracted peat volume were based partly on data from the Norwegian Food Safety Authority (1990 – 2007) and partly on information from a peat producer survey conducted in 2015 (covering the years 2008 – 2015). The two data series are not complete for all years, and extrapolation has been done to ensure a consistent time-series. The peat extraction activity data is described in detail in Sjøgaard and Økseter (2017).

6.7.1.2 Uncertainties and time-series consistency

The estimation of the uncertainty of the area and the C stock of wooded mire is described in section 6.3.5.

For the key category analyses (KCA), on- and off-site CO₂ emission uncertainties were combined by the weighted sum of the variances (square of Total uncertainty % in Table 6.) assuming no (0) correlation between on- and off-site emissions. The assumption of no correlation is based on the fact that on-site emissions will occur for many years even if no peat extraction would be conducted and off-site emissions would be 0. The weights (w) were given by squared proportions of the total emissions ($w = (40/53)^2 = 0.57$ for off-site emissions and $w = (13/53)^2 = 0.06$ for on-site emissions). Also the uncertainties for CH₄ on land between and in ditches were combined for the KCA. It was the weighted average of the variance of the emission factors for land between and ditch and their covariance assuming a direct correlation of 1. The correlation of 1 was chosen because of the increase in one area due to an increase in the other, and vice versa. The weight was given by the squared ditch proportion. In sum the uncertainty is assumed to be 110 % for CO₂, 128 % for CH₄, and

151 % for N₂O emissions. Uncertainties for CO₂ emissions estimated from drained organic soils on wetlands used for peat extraction are shown in Table 6.4, and for CH₄ and N₂O in Table 6.5.

6.7.1.3 QA/QC and verification

The QA/QC performed on the NFI area estimates was made for the wooded mire areas. The general QC procedures were performed on all sources under wetlands remaining wetlands. In addition, extensive QA by a national expert was performed for the off-site CO₂ emission factor.

6.7.1.4 Recalculations

The recalculations described in section 6.4.1.4 apply. The estimates of C stock changes in living biomass on wooded mires were recalculated due to the extrapolation method for the area estimate and C stock change in living biomass estimates. For 4.D.1.1 Peat Extraction Remaining Peat Extraction carbon stock change in organic soils, no peat volume data was available for the year 2017. Therefore, the average peat volume from 1990-2016 was used.

6.7.1.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.7.2 Land converted to wetlands, 4D2

Conversion of land to wetlands is most likely a slow process, unless in the form of flooding of land, which enables rapid inundation. Flooding can be human-induced (e.g. created by dams for hydropower production), or non human-induced (e.g. beaver dams). Only a few small-scale hydropower dams have been created in streams in the last 20-30 years and the total area is less than 4 kha. We consider emissions from this conversion category as negligible and report it using the notation key NO. We report C stock changes in living biomass, DOM, and soils for forest land converted to other wetlands. The area of land converted to peat extraction and soil related emissions are reported as NE because it is considered negligible.

6.7.2.1 Methodological issues

Emissions from land converted to wetlands were estimated for living biomass, DOM, mineral and organic soils.

Living biomass

Carbon stock changes in the living biomass pool were estimated using the Tier 3 approach (section 6.4.1.1), where gains and losses are recorded in the NFI. No changes occurred on non-forested land-use categories converted to wetlands.

Dead organic matter

A Tier 2 method was used to estimate C stock changes in DOM on forest land converted to wetlands. The stock change rate was estimated at -3.30 t C ha⁻¹ yr⁻¹, based on the assumption that all litter and dead wood in an average Norwegian forest are decomposed over 20 years after conversion. The derivation of the C stock estimates for dead wood and litter are described in section 6.4.2.1.

Table 6.24 Annual stock change rates ($\text{t C ha}^{-1} \text{ yr}^{-1}$) for forest land converted to wetlands. Other land converted to wetlands was assumed to have all C pools in a steady state condition.

	Mineral Soil	DOM	Litter	Dead Wood	Total
	($\text{t C ha}^{-1} \text{ yr}^{-1}$)				
Forest land converted to wetlands	1.5	-3.30	-3.05	-0.25	-1.80

Soils

Changes in SOC in mineral soil were estimated using a Tier 2 method. The C stock change rate for forest land converted to wetlands was estimated based on a measured national mean SOC stock of the mineral soil layer at a depth of 30 cm for forest as $57 \text{ t C ha}^{-1} \text{ yr}^{-1}$ and the IPCC default soil C reference stock for wetland soils in a temperate climate of $87 \text{ t C ha}^{-1} \text{ yr}^{-1}$; Table 2.3 (IPCC 2006). For organic soils we used the default emission factor for nutrient poor, boreal climate on forest land, which is $0.25 \text{ t C ha}^{-1} \text{ yr}^{-1}$. The conversion of other land to wetlands is not likely to result in any change in SOC, and the notation key NO is reported in the CRF.

6.7.2.2 QA/QC and verification

In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.7.2.3 Recalculations

The recalculations described in section 6.4.1.4 apply. Recalculations due to changes in NFI related data on areas and biomass were performed.

6.7.2.4 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.7.3 Completeness

The reporting for emissions and removals occurring on wetlands is complete.

6.8 Settlements – 4E

Settlements is a diverse land-use class consisting of for example residential areas, roads, recreation areas, powerlines within forests, gravel pits, mines, and industrial areas. The land-use class is especially important for the sub-group of land converted to settlements because of the increase in the size of the area of this land-use class since 1990.

6.8.1 Settlements remaining settlements, 4E1

On settlements remaining settlements we report carbon stock changes in living biomass, DOM, mineral and organic soils using Tier 1 methods. Organic soils on settlements remaining settlements is a key category with respect to the 1990 and 2017 level assessments.

6.8.1.1 Methodological issues

Living biomass

To estimate CSC in the living biomass pool a Tier 1 method is used, assuming no stock change and NO is reported. This is because trees are traditionally not measured on settlements in the NFI, due to the relatively small amounts of living biomass on settlements (Løken 2012).

In a specific study, trees were measured in land-use classes where trees are usually not measured in the NFI, including those within settlements (Løken 2012). A panel of NFI plots visited in 2009 containing almost 900 plots within settlements was used in the study. Settlements cover slightly more than 2 % of the Norwegian land area, but have a relatively low biomass density and contain only approximately 0.4 % of the total biomass stock (Løken 2012).

DOM and mineral soils

Carbon stock changes in DOM and mineral soil pools are also estimated using a Tier 1 method. This implies an assumption that no CSC occurs, and hence the notation key NO is used.

Organic soils (key category)

According to the IPCC 2006 guidelines, emissions from settlements on drained organic soils can be assumed to be similar to those on croplands (IPCC 2006). Emissions from organic soils in settlements are thus reported with Tier 1 using the default emission factor for croplands, which is $7.9 \text{ t C ha}^{-1} \text{ yr}^{-1}$.

6.8.1.2 Uncertainties and time-series consistency

Uncertainties are shown in Table 6.3 and estimated as described in section 6.1.3.

6.8.1.3 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.8.1.4 Recalculations

Emissions from organic soils were recalculated due to the updates in the NFI area.

6.8.1.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.8.2 Land converted to settlements, 4E2

The conversion of land to settlements is a significant source of emissions, primarily due to forest land conversion, which causes large losses in all C pools.

Forest land converted to settlements is identified as a key category with respect to living biomass, DOM, mineral soil, and for organic soil for Level assessment 1990, Level assessment 2017, and/or Trend assessment 1990-2017 (see Table 6.6).

6.8.2.1 Methodological issues

Living biomass (key category)

For lands converted to settlements, except for croplands, tree measurements are usually available before the conversion, if the area was tree covered. While trees are not measured on settlements, the NFI records which of the trees are remaining on the converted sample plot the first time the sample plot is visited after the conversion. Diameter and height measurements are, however, not carried out. Based on the information of which trees were removed, the carbon stock change on the converted sample plots is calculated using the last biomass measurement before conversion assuming no increment. The carbon stock of the last measurement minus the carbon stock of the removed trees is then used as the carbon stock of the plot assuming no changes in the future. For forest land, wetlands, and other land converted to settlements, this constitutes a Tier 3 method. The recording of which trees are remaining on a converted sample plot started in 2005. In the time-series before 2005, we assume that all trees were removed in the year when the land-use change was observed. An example of a situation where land is converted to settlements with remaining trees, is a forested sample plot of which the biggest part is converted to a house, while some of the trees are still alive inside what is now a garden.

For grassland, tree measurements are available since 2007 and a Tier 2 method is applied. Sample plots converted from grassland to settlements since 2007 did not have living tree biomass. Therefore, the Tier 1 default factor of 4.25 t ha^{-1} (i.e. $8.5 \text{ t dry matter ha}^{-1}$) (IPCC 2006, Table 6.4, p.6.27) was assumed as a loss in the first year of conversion. For cropland converted to settlements, no tree measurements are available in the NFI and the carbon stock changes are reported according to a Tier 1 method, assuming immediate loss of the default C stock of 4.7 t ha^{-1} on cropland (IPCC 2006, p. 8.18).

Dead organic matter (key category)

We used a Tier 2 method to estimate C stock changes in DOM on forest land converted to settlements for mineral soil areas. The method is based on C stock change rates multiplied by the area. The change rate for DOM was $-3.30 \text{ t C ha}^{-1} \text{ yr}^{-1}$ based on the change rates for litter and dead wood (Table 6.25). We assumed that the C stocks in litter and dead wood of an average Norwegian forest were completely lost over a 20 year period; see section 6.4.2.1 for the estimation of the stocks.

Table 6.25 Annual stock change rates ($\text{t C ha}^{-1} \text{ yr}^{-1}$) for Land converted to settlements.

Land converted to settlements	Soil	DOM	Litter	Dead Wood	Total
	$(\text{t C ha}^{-1} \text{ yr}^{-1})$				
Forest land	-0.57	-3.30	-3.05	-0.25	-3.87
Cropland	-0.83	0			-0.83
Grassland	-0.98	0			-0.98
Wetlands	-0.87	0			-0.87

Mineral soil (key category)

Changes in SOC were estimated using a Tier 2 method. The method is based on C stock change rates multiplied by the area as described under forest land. The C stock change rates were based on mean soil C stocks per land-use class. The assumption was made that upon conversion to settlements a 20 % C loss relative to the previous land use occurs over 20 years (IPCC 2006). The mean soil C stock for forest land and cropland were based on measurements as described in the respective chapters and on the IPCC default value for grassland and wetlands. The mean national SOC stock estimates were 57 t C ha^{-1} for forest land, 83 t C ha^{-1} for cropland, 98 t C ha^{-1} for grassland, and 87 t C ha^{-1} for wetlands. We assumed no SOC change when other land was converted to settlements.

Organic soil (key category)

CO_2 emission from drained organic soils on forest land converted to settlements was identified as a key category. Emissions were calculated using the Tier 1 method. According to IPCC (2006), we assume the emission factor for land converted to settlements corresponds to the cropland emission factor of 7.9 t C ha^{-1} .

6.8.2.2 Uncertainties and time-series consistency

Uncertainties are shown in Table 6.3 for living biomass and DOM and in Table 6.4 for organic and mineral soils. The time-series was consistently calculated.

6.8.2.3 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.8.2.4 Recalculations

The recalculations described in section 6.4.1.4 apply for changes from forests and tree-stocked land use categories. The time-series was recalculated due to updates in the NFI database and the extrapolation method.

6.8.2.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.8.3 Completeness

The reporting for emissions and removals occurring on settlements is complete.

6.9 Other land – 4F

The land use category other land covers approximately 45 % of the total land area in Norway, and reflects the large amount of mountainous and rocky terrain. Land-use changes to other land only occurred from grassland and settlements throughout the inventory period. It is only mandatory to report CSC for land converted to other land.

6.9.1 Other land remaining other land, 4F1

Reporting of emissions from other land remaining other land is not mandatory. Given the size of the area of other land, we analyzed the NFI data to determine the area usage and location of the land-use class. The vast majority of the other land is located above the alpine forest limit and only 21 % is located below (Table 6.26). Area of lands which have soil cover and are located below the alpine tree limit could potentially become forest land. Approximately 7 % of other land fulfills these criteria.

Table 6.26 Distribution of other land related to the alpine location and vegetation.

Location & area usage	Percentage (%)
<i>Area above the alpine forest limit</i>	
Other wooded land	4
Bare land	75
<i>Area below the alpine forest limit</i>	
Other wooded land	6
Coastal calluna heath land	1
Bare land	14
Total area of other land	100

6.9.2 Land converted to other land, 4F2

Only a small area of grassland and settlements on mineral soils was converted to other land during the inventory period. Carbon stock changes are reported for all sources. There were no areas of forest land, croplands, and wetlands converted to other land and these have been reported as NO.

6.9.2.1 Methodological issues

The area estimates are based on the NFI data. The Tier 3 method described under forest land was used for estimating C stock changes in living biomass. No CSC changes in the living biomass are recorded and NO is reported for the whole times series. For DOM, we use the Tier 1 method that assumes no CSC and report NO.

To estimate SOC changes in mineral soils on grassland converted to other land we used a Tier 2 method with a soil C stock change rate of $-0.25 \text{ t C ha}^{-1} \text{ yr}^{-1}$, equal to a 5 % loss relative to the SOC stock of grassland over the 20 year period. The change rate was multiplied with the area estimate determined by the NFI.

6.9.2.2 Uncertainties and time-series consistency

Uncertainties are estimated as described in section 6.1.3 and are shown in Table 6.3 for living biomass and DOM and in Table 6.4 for mineral and organic soils.

6.9.2.3 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure.

6.9.2.4 Recalculations

The time-series for C stock changes for living biomass, mineral, and organic soils were recalculated due to updates in the NFI database.

6.9.2.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.9.3 Completeness

The reporting for emissions and removals occurring on other land is complete.

6.10 Harvested wood products – 4G

Harvested wood products (HWP) prolong the period carbon is bound in timber after it is removed from the forest. Included in the HWP accounting is the carbon pool inflow in sawnwood, wood-based panels, and paper and paperboard. That is, HWP do not include all wood material that leaves the harvest site, only those parts of the harvest used for the three above mentioned default HWP categories. In the base year 1990, the total HWP pool was 1000 kt CO₂ according to the current calculation approach.

Net annual removals from HWP in use in 2017 were -145 kt CO₂ for HWP produced and consumed domestically and -91 kt CO₂ for HWP produced and exported. In 2017, the total removals from the HWP pool were -236 kt CO₂.

In the period 2009-2016, there has been a clear trend toward less storage (see Figure 6.16) in the HWP pool. A similar trend was found for other European countries, including the peak in the period around 2008-2010. In 2017 HWP again provided removals: sawnwood -415 kt CO₂ (-347 in 2016), wood based panels 35 kt CO₂ (25 in 2016), paper and paperboards 144 kt CO₂ (407 in 2016). Hence, the effect is due to less emissions from paper and paperboards and increased removals from sawnwood.




Figure 6.16 CO₂ emissions and removals in HWP (kt CO₂). Source: Norwegian Institute of Bioeconomy Research.

6.10.1 Methodological issues

Choice of method

Emissions and removals reported for HWP are estimated using a Tier 2 method. For consistency reasons the calculations are based on chapter 2.8 in the 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol (IPCC 2014a). The 2013 IPCC KP supplement approach fulfills the requirements in footnote 12 in Table 4.Gs.1. The Tier 2 default options are applied, including the three default HWP categories sawnwood, wood-based panels, and paper and paperboard and their associated half-lives and conversion factors (IPCC 2014a).

To improve the transparency and in order to be in accordance with Table 4.Gs.1, the Norwegian estimates differentiate between domestically produced and consumed HWP and wood products that are produced and exported.

All harvested wood in Norway originates from existing forest lands. The activity data used starts in 1961 and is based on FAO statistics. Calculations have been performed using data from 1961 to 2016. We calculated the historic pool from 1950-1960 according to IPCC (2014a). Only emissions from 1990 and onwards are reported.

The estimation of carbon stocks (C) and annual carbon stock changes (ΔC) for each HWP category was estimated using Eq. 2.8.5 (IPCC 2014a)

$$C(i+1) = e^{-k} \times C(i) + \left[\frac{(1 - e^{-k})}{k} \right] \times Inflow(i)$$

$$\Delta C(i) = C(i+1) - C(i)$$

Where, i = year; $C(i)$ = the carbon stock (kt C) in the particular HWP category at the beginning of year i , with $i = (1961, \dots, 2015)$, but only emission estimates for $i = (1990, \dots, 2015)$ are reported; k is the decay constant for the first-order decay for HWP category (i.e. sawnwood, wood-based panels or paper and paperboards) given in units year^{-1} as $k = \ln(2)/HL$, where HL is the half-life of the HWP pool, which is constant and given in the unit years. $Inflow(i)$ = the inflow to the particular HWP category during year i ; $\Delta C(i)$ = carbon stock change of the HWP category during year i , kt C year^{-1} .

The approximation of the carbon stocks in HWP pools at initial time $C(i = t_0 = 1961)$ was calculated according to Eq. 2.8.6

$$C_{(t_0)} = \frac{Inflow_{average}}{k}$$

where

$$Inflow_{average} = \frac{(\sum_{i=t_0}^{t_4} Inflow(i))}{5}$$

The C stock changes for each of the three HWP categories (sawnwood, wood-based panels, paper and paperboard) were estimated and summed to provide the total for Norway.

Activity data

All the activity data are from the FAO forestry statistics (<http://www.fao.org/faostat/en/#data/FO>). The initial unit is m^3 , except for the pulp and paper where the unit is metric ton. Conversion to carbon was performed using the default conversion factors given by the IPCC (IPCC 2014a). Exported and domestically consumed HWP is calculated and reported separately. The inflow data of domestically produced and consumed are based on consumption (Production – Export). Imported HWP is not included in the calculations (Production approach).

Assumptions

It is assumed that the Tier 2 method reflects the carbon flow in the HWP pool. The assumption of first-order decay (i.e. exponential decay) implies that loss from the stock of products is estimated as a constant fraction of the amount of stock (IPCC 2006).

It is assumed that the default half-lives are representative values for Norway.

6.10.2 Uncertainties and time-series consistency

The reported uncertainty estimates for half-lives are ± 50 % according to IPCC 2006. According to the 2013 IPCC KP Supplement (2014) an overall estimate of the HWP activity data (i.e. sawnwood, wood-based panels, and paper and paperboard) from e.g. FAO, result in an estimated uncertainty of the reported values between -25 % to +5 %.

6.10.3 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure.

6.10.4 Recalculations

Each year, when the new activity data are added from the FAO database, the activity data for the previous five years are checked and updated, if needed. The following activity data, which is used as data for inflow for 2014-2016, were updated:

- Wood-based panels 2016: export and domestically consumed. These updates in the activity data did not result in recalculation of the total emissions from HWP for the year 2016.

6.10.5 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.10.6 Completeness

The reporting for emissions and removals from harvested wood products is complete.

6.11 Direct N₂O emissions from managed soils – 4(I)

Direct N₂O emissions from managed soils are estimated from N inputs of inorganic and organic origin. N inputs from inorganic N fertilizer applied to forest land are reported. Inorganic fertilizer is not applied to managed wetlands and is hence reported as NO. Any inorganic fertilizer applied in the land use category settlements is included in the agriculture sector and reported as IE in the LULUCF CRF tables. Emissions from the use of organic fertilizers on settlements are reported. Livestock are generally not grazing managed wetlands (peat extraction areas and flooded lands), and is hence reported as NO. N inputs from organic and inorganic N fertilizer on cropland and grassland are reported as IE and included in the agriculture sector.

6.11.1 Inorganic fertilizer on forest land

N₂O is produced in soils as a by-product of nitrification and denitrification. Fertilizer input is particularly important for this process. However, fertilization of forest land is limited in Norway. The area fertilized was 24 km² in 1990 and the area as well as the net amount of N applied have decreased during the inventory period. In 2016 and 2017 however, there was a large increase in area to approximately 91 km² and a corresponding increase in net amount of N applied (Table 6.27). Reported emissions are presented in Table 6.27.

Table 6.27 Estimated emissions from fertilization of forest land 1990-2017.

Year	Fertilizer input (t N)		Net amount N applied (t N)	N ₂ O emissions (t N ₂ O)
	Mineral soil	Organic soil		
1990	177	59	236	2.4
1991	326	67	392	3.9
1992	253	102	356	3.6
1993	181	67	248	2.5
1994	169	67	236	2.4
1995	160	60	220	2.2
1996	199	37	235	2.4
1997	232	19	252	2.5
1998	243	23	265	2.7
1999	218	44	261	2.6
2000	135	22	156	1.6
2001	154	19	173	1.7
2002	178	8	187	1.9
2003	85	1	87	0.9
2004	76	2	78	0.8
2005	53	31	84	0.8
2006	34	4	38	0.4
2007	81	1	82	0.8
2008	106	1	107	1.1
2009	113	1	114	1.1
2010	73	0.2	73	0.7
2011	85	0	85	0.8
2012	112	0.1	112	1.1
2013	170	0.2	170	1.7
2014	59	0	59	0.6
2015	91	0	91	0.9
2016	1236	0	1236	12.4
2017	1350	0	1350	13.5

6.11.1.1 Methodological issues

Choice of method

The estimate is based on a Tier 1 method with a default emission factor. Emissions are calculated according to

$$N_2O \text{ direct-}N_{\text{fertilizer}} = F_{SN} \times EF \times 44/28,$$

where F_{SN} is the annual amount of synthetic fertilizer nitrogen applied (kt N) to forest soil and EF is the emission factor for N₂O emissions from N inputs, kg N₂O-N/kg N input.

Activity data

Statistics Norway supplied unpublished data on the application of synthetic fertilizer. The statistics include the area applied with fertilizer, the amounts and types of fertilizer applied for the period 1995-2015, but only the area and amount of fertilizer applied is available for 2016-2017 due to a

change in the data acquisition procedures. For the period 1990–1994, only data for the total fertilized area is available. Data from the period 1995–2004 were used to estimate the amount of N-fertilizer applied for the period 1990–1994.

The amount of fertilizer applied is given as total weight. The nitrogen content depends on the type of fertilizer. Yara supplied sales numbers for forest fertilization. From 1993 to 1994 and onwards, calcium ammonium nitrate based fertilizer has dominated the market for fertilization of forest on mineral soils (Pers. comm. Ole Stampe, Yara Norge AS, 2013). The N-content of calcium ammonium nitrate is 27 % (weight percent). According to Statistics Norway, this fertilizer is applied to approximately 97 % of the fertilized forest land in Norway; it is currently uncertain what fertilizer is applied to the remaining small area percent.

Emission factor

The default emission factor is 1 % of applied N (Table 11.1 IPPC 2006). The emission factor is highly uncertain, with uncertainty range from 0.003 to 0.03 (IPCC 2006).

6.11.2 Organic fertilizer on forest land

In Norway livestock grazes the outer fields during the summer months. The outer fields encompass land classified as other land and forest land. Emissions from organic N fertilizer applied by animal manure when livestock graze in the forest land remaining forest land and land converted to forest land is reported with the notation key IE and emissions are reported in the agriculture sector starting in the 2019 submission. These emissions were previously reported in both the LULUCF and agricultural sector in Norway (see 6.11.6).

6.11.3 Organic fertilizer on settlements

Direct N₂O emissions from application of organic N fertilizer in settlements have been reported in the LULUCF sector since NIR 2015. Previously, emissions from the application of sewage sludge on urban lawns, road-side grass-strips, and parks were reported in the waste sector. Emissions have increased slightly from 0.009 kt N₂O-N yr⁻¹ in 1990 to 0.019 kt N₂O-N yr⁻¹ in 2017 (equivalent to 2.68 to 5.66 kt CO₂ yr⁻¹).

6.11.3.1 Methodological issues

Choice of method

A Tier 1 method was used applying the default emission factor (IPCC 2006). To derive N inputs from organic fertilizer, the total dry matter amount of all types of sewage sludge applied was multiplied by an N content of 2.82 % (Statistics Norway 2001).

Activity data

Data of total amount (dry matter) of sewage sludge are derived from Statistics Norway (SSB) and cover the following distribution types: parks and green areas, soil fertilizer production, cover on landfills, other use and unknown use. The data is collected every year by SSB, and a consistent time series from 1990 was available.

6.11.4 Uncertainties

The uncertainty related to the default emissions factor for N₂O from N additions from mineral and organic fertilizer is provided by the IPCC as the range of 0.003 - 0.03 equal to ± 200 %. In addition, we assume that the activity data have ± 20 % uncertainty associated with the estimation of inorganic N applied to forest land and organic N applied to settlements. The activity data and the method used to estimate the organic N input to forest land are more uncertain and an error of ± 50 % was assumed. The total uncertainties (of the emission factor and the activity data and method) were used in the KCA for each of the three sources.

6.11.5 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure.

6.11.6 Recalculations

Recalculations for this source were due to the removal from the LULUCF sector of organic N fertilizer inputs on forest land, due to double counting. From the 2019 submission it is only included in the agriculture sector. Emissions from forest land for the year 2016 were 0.0843 kt N₂O (NIR 2018) and 0.0124 kt N₂O (NIR 2019).

6.11.7 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.11.8 Completeness

The reporting for Direct N₂O emissions from managed soils is complete.

6.12 Emissions and removals from drainage, rewetting and other management of soils – 4(II)

Rewetting of organic and mineral soils is not practiced in large scale in Norway, and drainage and rewetting (WDR) is not an elected KP activity. Thus, in CRF Table 4(II) we report only emissions from drained organic soils (including peat extraction) as these are mandatory to report. CO₂ emissions from drained organic soils are reported as IE in CRF table 4(II) because they are included in CRF tables 4.A-4.D as C stock changes in the organic soils pool. In CRF table 4(II) we report CH₄ and N₂O emissions from forest land and from wetlands used for peat extraction, and CH₄ emissions from cropland and grassland. NE is reported for rewetting of mineral and organic soils on forest land, cropland, grassland, and wetlands (except for rewetted mineral soils which is reported as NO). According to the IPCC guidelines, N₂O emissions from drained organic agricultural soils (cropland and grassland) are reported in the agriculture sector. Both CH₄ and N₂O emissions are key categories for forest land, in the 1990 and 2017 level assessments for CH₄ and in both level and trend assessment for N₂O emissions. CH₄ emissions from cropland is also a key category, in the 2017 level assessment.

Please note that CRF tables *Table4* and *Summary2* are inconsistent due to some emissions of CH₄ and N₂O that cannot be reported in the CRF by detailed area types, according to footnote 4 in Table 4(II) and Table 4(IV). These emissions are entered into the tables only at more aggregated levels. The level of reporting is due to properties of the CRF system, follows decision 24/CP.19, and is not caused by lack of data in the Norwegian emission inventory. The UNFCCC Secretariat did confirm the inconsistency in the sums of the subtotals in 2015.

6.12.1 N₂O emissions from drainage of organic soils (key category)

6.12.1.1 Methodological issues

For the estimation of N₂O emission from drained organic soils on all land uses we use a Tier 1 method based on the 2006 IPCC guidelines (IPCC 2006). The area is multiplied with an emission factor. To make use of the most recent scientific knowledge we apply the emission factors from the IPCC 2013 Wetlands supplement (IPCC 2014b).

Activity data

The area of drained forest soil was provided by Statistic Norway and stratified into boreal nutrient rich and boreal nutrient poor vegetation zones, as described in section 6.4.1.1. For the reporting under 4(II), all forest land, including land converted to forest land, is included in the estimate.

The area of land under peat extraction was estimated as described under section 6.7.1.1.

Emissions factors

The default emission factors from the IPCC 2013 Wetlands supplement were used. All Norwegian forest land is considered boreal and we used the same distribution of nutrient rich and nutrient poor, as described under forest land; organic soils (69 % nutrient rich and 31 % nutrient poor) which gives an average national EF of 2.28 kg N₂O-N yr⁻¹. For the area in the conversion classes, we used the nutrient rich EF (3.2 kg N₂O-N yr⁻¹). N₂O emissions from wetlands used for peat extraction were estimated with the emission factor of 0.3 kg N₂O-N yr⁻¹ (IPCC 2014b); see Table 6.23.

6.12.2 CH₄ emissions from drainage of organic soils (key category)

6.12.2.1 Methodological issues

To estimate CH₄ emissions, we used the Tier 1 method applying the EFs of the IPCC 2013 Wetlands supplement (IPCC 2014b). The method accounts for methane fluxes both in the drainage ditches and on the land using the following equation

$$\text{CH}_4 = A \times ((1 - \text{Frac}_{\text{ditch}}) \times \text{EF}_{\text{CH}_4_{\text{land}}} + \text{Frac}_{\text{ditch}} \times \text{EF}_{\text{CH}_4_{\text{ditch}}})$$

where, A is the area of drained organic soil; $\text{Frac}_{\text{ditch}}$ is the fraction of the area occupied with ditches; and $\text{EF}_{\text{CH}_4_{\text{land}}}$ $\text{EF}_{\text{CH}_4_{\text{ditch}}}$ are the emissions factor for the land and the ditch, respectively.

There is no information available in Norway to provide an accurate estimate for the fraction of the area occupied with ditches ($\text{Frac}_{\text{ditch}}$), we therefore used the default values of 2.5 % for forest land, and 5 % for cropland, grassland, and peat extraction (IPCC 2014b).

Activity data

Activity data of the area of drained forest soil was provided by Statistic Norway and stratified into boreal nutrient rich and boreal nutrient poor vegetation zones, as described in section 6.4.1.1. For the reporting under 4(II), all forest land, including land converted to forest land, was reported. The area of land under peat extraction was estimated as described under section 6.7.1.1. For cropland and grasslands, the estimation of the areas of drained organic soils were as described in section 6.5.1.1 and section 6.6.1.1, and the areas of land converted to cropland and grassland, respectively, are also included in these estimates.

Emission factor

For forest, the default EFs for CH₄ from land ($\text{EF}_{\text{CH}_4_{\text{land}}}$) from the IPCC 2013 Wetlands supplement, given the same distribution of nutrient rich and nutrient poor forest land as for the N₂O and CO₂ estimation, resulted in a mean national EF of 3.55 kg CH₄ yr⁻¹. For cropland the EF is 0 and for grassland we used the factor for shallow-drained, nutrient rich grassland of 39 kg CH₄ yr⁻¹. For peat extraction on wetlands the emission factor is 6.1 kg CH₄ yr⁻¹ for the boreal zone (Table 6.23).

The emission factors for CH₄ from the ditches or drains ($\text{EF}_{\text{CH}_4_{\text{ditch}}}$) were 217 kg CH₄ yr⁻¹ for forest land, 1165 kg CH₄ yr⁻¹ for cropland and 572 kg CH₄ yr⁻¹ for grassland, and 542 kg CH₄ yr⁻¹ for peat extraction land.

6.12.3 Uncertainties

The uncertainties associated with the emission factors of the IPCC 2013 Wetlands supplement are summarized in Table 6.5.

To derive the total uncertainty of the emission estimate we aggregated the uncertainty for the emission factor and the area estimate, respectively. For land converted to forest land and the cropland or grassland categories, the area uncertainties were calculated as the sample error in the NFI. We assumed a 50 % uncertainty for the area of drained forest soils from Statistics Norway and 100 % for the area with peat extraction.

6.12.4 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure. In 2019, a complete quality assurance procedure on the Tier 1 for organic soils was performed.

6.12.5 Recalculations

All times series were recalculated due to the updates in the NFI area data except for peat extraction.

In 2019 the simultaneous correction of the emission factor for shallow-drained, nutrient rich grassland from 36 to 39 kg CH₄ yr⁻¹ as well as the rectification of the proportion of non-ditch area from 0,095 to 0,95 lead to the doubling of the methane emissions in grassland.

The rectification of the proportion of nutrient rich vs nutrient poor organic soil from 79-21 % to 69-31% induced a slight increase and decrease of the CH₄ and N₂O emissions respectively under forest.

Additionally, the CH₄ emission factor for cropland converted to forestland was corrected from “forest temperate” to “forest boreal nutrient rich” with little impact on the emissions.

6.12.6 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.12.7 Completeness

The reporting for emissions and removals from drainage and rewetting of organic and mineral soils is complete.

6.13 Direct N₂O emissions from N mineralization and immobilization – 4(III)

In the 2006 IPCC Guidelines direct N₂O emissions are estimated from N mineralization-immobilization turnover associated with loss of soil organic matter resulting from change of land use or management of mineral soils on all types of land use. Previously, only land-use changes to cropland were considered to result in N mineralization-immobilization. We estimate N₂O losses from all land uses that have negative C stock changes in the mineral soil pool, and the areas reported in CRF table 4(III) correspond to the areas with negative C stocks changes and not the total area for all land-use conversions. N₂O emissions from N mineralization-immobilization is a small key category in the 2017 level assessment and the 1990-2017 trend assessment.

6.13.1 Methodological issues

6.13.1.1 Choice of method

To estimate N₂O emissions from N mineralization we first calculate the net annual amount of N mineralized in mineral soils resulting from SOC loss (F_{SOM}) from

$$F_{\text{SOM}} = \Delta C \times 1 / \text{CN} \quad \text{Eq. 11.8; (IPCC 2006)}$$

where ΔC is the average annual C loss from mineralization of soil for each land-use type (in kt C yr⁻¹) and CN is the C/N ratio of cropland soils. To estimate the N₂O emissions from N mineralization we multiply F_{SOM} with the default emission factor ($\text{EF} = 0.01 \text{ kg N}_2\text{O-N (kgN}^{-1}\text{)}$). We consider the method a Tier 1 because we used the default C/N ratio ($\text{CN} = 15$), although most SOC losses were derived using a Tier 2 method.

Certain land-uses (e.g. forest land remaining forest land and cropland remaining cropland) and land-use changes (e.g. settlements converted to cropland or forest land) result in positive SOC stock changes in the mineral soil pool, thus no N₂O emissions are reported from these sub-categories.

6.13.1.2 Activity data

Activity data used for this source is the annual average C losses, which are those reported in the CSC tables 4.A-4.F for each land-use class. The CSC change is estimated as described under the mineral soil pool for each land-use class.

6.13.2 Recalculations

The area changes caused by the NFI updates caused only minor recalculations for this source.

6.13.3 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.13.4 Completeness

Reporting from the source 4(III) Direct N₂O emissions from N mineralization and immobilization is complete.

6.14 Indirect N₂O emissions from managed soils – 4(IV)

Indirect N₂O emissions occur through two pathways: 1) the volatilization of N as NH₃ and NO_x and subsequent deposition of N compounds (atmospheric deposition), and 2) the leaching and runoff of N from land that has been subjected to excess N application from organic or inorganic fertilizers, as well as N mineralized due to soil C loss. CRF table 4(IV) has the two sub-categories 1) atmospheric deposition and 2) nitrogen leaching and runoff. The 2006 IPCC methodology for estimation of indirect emissions includes N inputs from several sources (Eq. 11.9 and 11.10), however, the sources are split between the reporting in the LULUCF and the agriculture sector. The indirect emissions reported in the LULUCF sector under atmospheric deposition are derived from the N inputs coming from synthetic N fertilizer on forest land (F_{SN}) and organic N fertilizer on forest land and settlements (F_{ON}). For the sub-category N leaching and runoff, N inputs arrive as synthetic and organic N fertilizers as for atmospheric deposition, but also from N mineralization immobilization in mineral soils associated with loss of soil C (F_{SOM}). Indirect emissions caused by N inputs from crop residues, urine and dung application from livestock, and N fertilizers on agricultural lands (cropland and grassland) are reported in the agriculture sector.

Please note that CRF tables *Table 4* and *Summary 2* are inconsistent due to some emissions of CH₄ and N₂O that cannot be reported in the CRF by detailed area types, according to footnote 4 in Table 4(II) and Table 4(IV). These emissions are entered into the tables only at more aggregated levels. The level of reporting is due to properties of the CRF system and follows decision 24/CP.19, and is not caused by lack of data in the Norwegian emission inventory. The UNFCCC Secretariat has confirmed the inconsistency in the sums of the subtotals.

6.14.1 Atmospheric deposition

Indirect emissions reported under atmospheric deposition are estimated from synthetic N fertilizer input on forest land (F_{SN}) and organic fertilizer N inputs on settlements (F_{ON}). Emissions are rather small; 0.0008 kt N₂O in 2017 (0.24 kt CO₂-equivalents).

6.14.1.1 Methodological issues

Method choice

We used the Tier 1 method of the 2006 IPCC guidelines dictating that a fraction (Frac_{GASM} or Frac_{GASF}) of the organic and inorganic N inputs (F_{ON} and F_{SN}), respectively, is considered volatilized and multiplied by the emission factor for atmospheric deposition (EF) according to

$$\text{N}_2\text{O-N} = (\text{Frac}_{\text{GASF}} \times F_{\text{SN}} + F_{\text{ON}} \times \text{Frac}_{\text{GASM}}) \times \text{EF}_{\text{vol}} \quad \text{Eq.11.9; (IPCC 2006)}.$$

All parameters are default values: Frac_{GASM} = 0.2, Frac_{GASF} = 0.1, and EF = 0.01 kg N₂O-N (kg N)⁻¹.

Activity data

The N inputs from synthetic and organic N fertilizer were derived as described in section 6.11.

6.14.2 Nitrogen leaching and run-off

Indirect emissions from leaching and runoff were estimated from the N inputs of synthetic fertilizers on forest land and organic fertilizer on settlements and from N mineralized due to soil organic matter decomposition.

6.14.2.1 Methodological issues

Method choice

The Tier 1 method was applied where the fraction of all N added to the soils (F_{LEACH}) is multiplied with the default emission factor $EF_{\text{leach}} = 0.0075 \text{ kg N}_2\text{O-N (kg N leaching/runoff)}^{-1}$ by

$$\text{N}_2\text{O}_{(\text{L})-\text{N}} = (F_{\text{SN}} + F_{\text{ON}} + F_{\text{SOM}}) \times F_{\text{LEACH}} \times EF_{\text{leach}}$$

where F_{SN} is the N input from synthetic fertilizer, F_{ON} is the N input from organic fertilizer, and F_{SOM} is the input from N mineralized decomposition of mineral soils. We applied the default value for F_{LEACH} of 0.3.

Activity data

The activity data were derived as described in section 6.11 for organic and inorganic fertilizer and in section 6.13 for N mineralized during soil C loss.

6.14.3 Uncertainties

The uncertainty associated with the default emission factor for N_2O emissions from volatilization and deposition is $\pm 400 \%$ (IPCC 2006) and has a major influence on the emissions from atmospheric deposition. The EF for leaching has $\pm 233 \%$ uncertainty (IPCC 2006). In addition, the default values for the fraction of N that is volatilized from synthetic and organic fertilizer and the fraction that is lost by leaching, have high uncertainties. According to the 2006 IPCC guidelines, the uncertainties are $\pm 200 \%$, $\pm 150 \%$, and $\pm 167 \%$ for F_{GASF} , F_{GASM} , and F_{LEACH} , respectively. Furthermore, the estimated N inputs (F_{SN} , F_{ON} , and F_{SOM}) also have uncertainties either due to the activity data or methods as mentioned in the previous sections. Aggregating the individual uncertainties, we derive a total uncertainty of $\pm 447 \%$ for emissions due to atmospheric deposition and $\pm 300 \%$ from leaching and run-off (Table 6.5).

6.14.4 QA/QC and verification

The QA/QC plan was performed according to the Tier 1 procedure.

6.14.5 Recalculations

Recalculations for nitrogen leaching and run-off resulted in reduced N_2O emissions in 2016 from 0.0516 (NIR 2018) to 0.0343 (NIR 2019) due to organic N fertilizers in forest land now being reported in the agriculture sector and no longer used in the estimation of Nitrogen leaching and run-off.

6.14.6 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.14.7 Completeness

Reporting from source 4(IV) Indirect N_2O emissions from managed soils is complete.

6.15 Biomass Burning – 4(V)

Emissions of CO₂, CH₄, and N₂O due to biomass burning are reported for all land-use classes. For cropland and grassland, burning should be reported for woody biomass which is not common on these land-use classes in Norway. Agroforestry is not normally practiced and woody biomass is found mostly in fruit tree orchards and these are generally not burned. Burning of woody biomass in wetlands, settlements, and on other land does not occur either. We therefore report NO for all gasses in all land-use classes, except for forest land. Controlled fires on forest land is reported as NE as very few fire drills are performed and we expect the emissions to be negligible. Wildfires on grasslands are also reported as NE due to the lack of data but also because emissions are considered negligible as wildfires rarely occur on Norwegian grasslands.

6.15.1 Fires on forest land

Prescribed burning of forest takes place in Norway only connected to firefighting rehearsals, comprising a very small area (approximately 15 ha yr⁻¹). Thus, these emissions are reported as NE and assumed negligible. The area subject to wildfires varies considerably from year to year due to natural factors (e.g. variations in precipitation). According to the IPCC 2006 guidelines, emissions of CO₂ from biomass burning in forest land remaining forest land need to be accounted for, however, CO₂ emissions caused by biomass burning are included in the estimate of C stock change in living biomass derived from the stock-change method. Hence, estimates of CO₂ emissions from wildfires are reported as IE.

6.15.1.1 Methodological issues

Emissions of N₂O and CH₄ from forest wildfires are relatively small, and the emissions of CO₂ contribute to the major part of the total emissions (Table 6.28).

Choice of method

There are no national data on emission factors for non-CO₂ gases from forest fires. N₂O and CH₄ emissions from forest wildfires are estimated based on a Tier 1 method from the 2006 IPCC guidelines. The following equation 2.27 is used:

$$L_{\text{fire}} = A * M_B * C_f * G_{\text{ef}} * 10^{-3}$$

Where: L_{fire} = amount of greenhouse gas (CO₂, CH₄, and N₂O) from fire (tonnes), A = area burnt (ha), M_B = mass of fuel available for combustion (tonnes ha⁻¹), C_f = combustion factor (dimensionless), and G_{ef} = emission factor (g kg⁻¹ dry matter burnt). Activity data (area burnt) is based on country level estimates. Values used for M_B and C_f are derived by taking into account estimates of the mass and the amount consumed of unproductive forest, productive forest, dead wood, and humus. The quantification of national estimates for biomass burned and carbon released is based on expert judgment.

Table 6.28 Estimates of CO₂, CH₄, and N₂O emissions (kt) from forest fire from 1990 to 2017, and sum of emissions in kt CO₂-equivalents.

Year	CO ₂ (kt)	CH ₄ (kt)	N ₂ O (kt)	SUM CO ₂ -eqv. (kt)
1990	9.74983	0.73015	0.48146	10.96144
1991	16.39590	1.22786	0.80966	18.43342
1992	15.04645	1.12681	0.74302	16.91627
1993	2.78617	0.20865	0.13759	3.13240
1994	3.16571	0.23708	0.15633	3.55911
1995	1.25425	0.09393	0.06194	1.41011
1996	8.77266	0.65697	0.43321	9.86284
1997	9.41687	0.70521	0.46502	10.58710
1998	3.45390	0.25866	0.17056	3.88311
1999	0.71337	0.05342	0.03523	0.80202
2000	1.49074	0.11164	0.07362	1.67599
2001	0.60668	0.04543	0.02996	0.68207
2002	2.89344	0.21669	0.14288	3.25301
2003	6.08545	0.45573	0.30051	6.84170
2004	1.22235	0.09154	0.06036	1.37425
2005	3.57510	0.26773	0.17654	4.01938
2006	32.43060	2.42868	0.60148	34.46076
2007	2.53236	0.18964	0.12505	2.84706
2008	51.50211	3.85691	2.54327	57.90229
2009	8.89877	0.66641	0.43944	10.00462
2010	14.61807	1.09473	0.72187	16.43467
2011	1.94002	0.14529	0.09580	2.18111
2012	0.76764	0.05749	0.03791	0.86304
2013	0.53202	0.03984	0.02627	0.59814
2014	5.94011	0.44485	0.29333	6.67829
2015	0.86648	0.06489	0.04279	0.97416
2016	14.26019	1.06792	0.70419	16.03231
2017	5.05824	0.37880	0.24979	5.68683

Activity data

Data of burned areas due to wild forest fires are available from the Norwegian Directorate for Civil Protection (DSB) for 1993–2017. Data are available for the number of fires and the area of productive and unproductive forests that burned. There were only data available for the number of fires between 1990 and 1992, and these data were used to estimate the area burned based on the ratio for subsequent years. This method may be very inaccurate because the size of fires is very variable. The number of fires was higher in 1990-1992 than later and it has assumed that the area burned was proportionally higher (Rypdal et al. 2005).

Standing volume for unproductive and productive forest were based on average numbers and accounted for 23 and 109 m³ ha⁻¹, respectively (Granhus et al. 2012). In biomass this is equal to 12 and 55 t ha⁻¹, respectively. The IPCC (2003) estimate that 50 % of the carbon is released during fires is appropriate, because this is assumed to be the C content of woody biomass.

In addition to the lack of data on the tree biomass, there are no exact data on the amount of biomass burned per area. Normally, only the needles/leaves, parts of the humus, and smaller branches would

burn. The mass of trees burned constitute 25 % of the biomass, which is consistent with IPCC (2003). It is also likely that about 1 m³ dead wood per ha will be affected by the fire due to its dryness. It is difficult to assess how much of the humus is burned, and this is much dependent on forest type. There is about 7 500 kg humus per ha and we assume that 10 % of this is burned. This percentage, however, is very dependent on the vegetation type. The CO₂ estimates provided in (Table 6.28) are for comparison only and to enable estimation of N₂O and CH₄ emissions, and thus, not used in the reported CO₂ emissions.

Table 6.29 Information on forest fires in Norway for 1990–2017 and estimated CO₂ emissions.

Year	Number of fires	Unproductive forest (ha)	Productive forest (ha)	Area burned (ha)
1990	578	679.6	256.4*	936.0*
1991	972	1 142.8	431.2*	1 574.0*
1992	892	1 048.8	395.7*	1 444.5*
1993	253	135.5	88.3	223.8
1994	471	123.6	108.1	231.7
1995	181	77.6	35.5	113.1
1996	246	169.7	343.8	513.5
1997	533	605.8	260.6	866.4
1998	99	164.7	110.3	275.0
1999	148	73.4	12.7	86.1
2000	99	142.6	29.3	171.9
2001	117	84.3	5.2	89.5
2002	213	124.7	95.8	220.5
2003	198	905.6	36.8	942.4
2004	119	84.6	32.3	116.9
2005	122	252.4	93.2	345.6
2006	205	3 222.2	606.7	3 828.9
2007	65	22.2	106.1	128.3
2008	171	1 210.2	1 963.6	3 173.8
2009	109	1 257.9	70.8	1 328.7
2010	62	165.9	602.8	768.7
2011	49	47.8	73.4	121.2
2012	24	35.1	24.9	60.0
2013	40	30.8	15.6	46.4
2014	133	681.7	87.7	769.4
2015	40	141.6	2.0	143.6
2016	341	1 685.9	197.9	1 883.8
2017	301	405.8	119.4	525.2

Source: Norwegian Directorate for Civil Protection (DSB) *Area estimated in Rypdal et al. (2005).

Emission factors

The IPCC (2006) emission ratios of 4.70 g/kg and 0.26 g/kg dry matter burnt are used for methane and nitrous oxide, respectively.

6.15.1.2 Uncertainties

The total uncertainty for wildfires was estimated at 75 % based on the default uncertainty for the non-CO₂ emission factor of 70 % IPCC (2003) and expert judgment on the activity data.

6.15.1.3 Recalculations

There were no recalculations for this source for the 2019 submission.

6.15.1.4 Planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

6.15.2 Completeness

Reporting of emissions from source 4(V) Biomass burning is complete.

6.16 Recalculations for LULUCF

All emissions related to the Norwegian National Forest Inventory (NFI) estimated areas are recalculated every year as well as for the 2019 submission. This is due to the interpolation and extrapolation method used to estimate the areas. Since NIR 2014, areas have been estimated using linear interpolation between the 1/5th of the NFI sample plots that are surveyed every year. One fifth of the area estimates are therefore based on measured sample plots, and four fifths of the estimate is based on interpolation or extrapolation in the final years. Extrapolation affects the four final reporting years and requires a recalculation of these years in the subsequent submissions. There will therefore always be recalculation in the last few years of the time series. The area-related emissions are affected by this and comprise all carbon stock changes estimated for living biomass, litter, dead wood, DOM, mineral and organic soils (CRF Tables 4.A-4.F). In addition, the following non-CO₂ emission sources are also indirectly affected by the area estimates: 4(II) Emissions and removals from drainage and rewetting of organic soils, and source 4(III) Direct N₂O emissions from N mineralization/immobilization. In addition to the interpolation and extrapolation method affecting area estimates, NFI sample plots are surveyed either in the field (if forested) or using aerial images. Corrections of previously attributed land-use categories that are not due to actual land-use changes therefore occur if new information becomes available. This can result in revised area estimates among all categories for the full time series. Another factor affecting recalculations is the calculation for changes in soil C, litter, and deadwood in afforestation and reforestation and deforestation for the inventory year 2015 for which a correction was made. For living biomass, a correction of the time series transition ("new remaining" starts in 2010, not 2011 as before) and the attribution of gains on lands-converted-to-not-Forest (gains are now correctly reported under the previous remaining class) affected the estimates on all land use categories with reported living biomass. Also an updated Statistics Norway harvests statistic affected the estimates of living biomass losses on all land use categories. The implementation of a Tier 1 method for living biomass changes affected the categories GS,SG and CS,SC. (Default factors for C and G are similar (4.7 t vs. 4.25 t). Therefore, no change was assumed for the GC category).

For the NIR 2019 the whole time series was recalculated for all C emission sources and sinks due to revised activity data (areas) and a few methodological changes. It was necessary to utilize preliminary data to produce emission and removal estimates for some select sources due to an earlier reporting schedule for the 2019 submission. Preliminary data was obtained for: timber harvest statistics, area of orchard trees, areas of grassland, proportions of crop types, organic N fertilizer inputs from animals, and organic N fertilizer inputs on settlements. No updated data were available at the writing for these data. The largest recalculations occurred for the last four years (2013-2016). Total changes in emissions (including non-CO₂ emissions) caused by recalculations for the LULUCF sector are shown in Figure 6.17.

Quantitatively, the largest changes in estimated C stock changes occurred for forest land and grasslands. The estimated CO₂ uptake on forest land for the year 2016 was 616 kt smaller in the 2019 submission compared to the 2018 submission. The estimated CO₂ emissions in grasslands for 2016 were 4.4 kt smaller (Table 6.). The removals between the estimates of total GHG emissions from the LULUCF sector were 577 kt CO₂-eq more in the 2019 NIR compared to the 2018 NIR submission for the recalculated year 2016 (Table 6.30).




Figure 6.17 Recalculations illustrated for total emissions (kt CO₂-equivalents per year) estimated for the LULUCF sector in the 2019 submission compared to the 2018 submission.

Table 6.30 Recalculated GHG emissions (kt CO₂-equivalents yr⁻¹) for 2016 per land-use category in the LULUCF sector.

Land-use category	Emissions for 2016 (kt CO ₂ -equivalents yr ⁻¹)		
	NIR 2019	NIR 2018	Absolute difference
4.A Forest land	-28 210.6	-28 827.0	616.4
4.B Cropland	2 058.9	2 038.5	20.4
4.C Grassland	191.9	196.3	-4.4
4.D Wetlands	18.4	21.6	-3.2
4.E Settlements	2 066.2	2 112.8	-46.6
4.F Other land	0.95	0.95	0
4.G HWP	85.00	85.00	0
4(IV)Total sum	10.5	16.1	-5.6
Total sum	-23 778.8	-24 355.8	577.0

The reasons for the recalculations within each land-use category and sink/source category and the effects for the last recalculated year of the inventory (2016) are described below.

6.16.1 Forest Land

6.16.1.1 Forest land remaining forest land

- Total CSC for forest land remaining forest land in 2016 was 190 kt C larger in the NIR 2018 submission compared with the NIR 2019 submission.
- Net change in living biomass decreased by 235 kt C from a C uptake of 5 823 kt C in NIR 2018 to 5 588 kt C in NIR 2019. Carbon uptake in dead wood, litter, and mineral soils increased by 6 kt C (from 362 to 367 kt C), 28 kt C (from 1 792 to 1 820 kt C), and 1 kt C (41 to 42 kt C), respectively. Recalculations were caused by the updates in the area estimates in addition to the general methodological changes described above.
- Emission from organic soil decreased consistently by 16 kt C between NIR 2018 and NIR 2019 for all reported years. For the year 2016, the value decreased by 16.4 kt C from 189.85 kt C to 173.45 kt C. This change resulted from the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively.

6.16.1.2 Land converted to forest land

- Compared to last year's reporting, the total C uptake for the year 2016 on land converted to forest land was 2 kt C larger in the NIR 2019 submission (from 143 to 145 kt C). This was mainly due to the larger C uptake in living biomass, which increased by 8 kt C from 46 kt C (NIR 2018) to 53 kt C (NIR 2019), which occurred primarily on settlements converted to forest land (7 kt C) and other land converted to forest land (1 kt C). Recalculations in the litter pool comprised the largest share (-7 kt C), while the deadwood (-0.06 kt C), mineral soil (1.4 kt C), and organic soil (-0.1 kt C) pools comprised a much smaller proportion. All recalculations were caused by the updated NFI areas.

6.16.2 Cropland

6.16.2.1 Cropland remaining cropland

- Total C stock loss for the year 2016 on cropland remaining cropland increased by 12 kt C from 416 kt C in the 2018 submission to 428 kt C in the 2019 submission. This was primarily due to changes in mineral soil, C stock changes which strongly decreased by 11.1 kt C between NIR 2018 and NIR 2019 from 22.1 kt C to 11.0 kt C, respectively. This was mostly due to the correction of an error. Prior correction, carbon stock change for silty loam soils were counted twice when summing the changes for all soil types. A second error correction, dealing with the attribution of area of grain-ley rotations had little impact on the final C stock change. Briefly, for each agrozone there were three types of grain-ley rotations fixed by the grain percentage of the total grain-Ley area. Agrozones with: >65% grain were attributed rotation with 1 third Ley and 2 third Grain (1:2LG rotation). Similarly 35-65% grain gave 1:1 LG rotation, and <35% grain gave 2:1 LG rotation (See Borgen et al., 2012). For more consistency, the choice of the grain percentage thresholds, 35% and 65 % in the previous version, were changed to 33.3% and 66.7%. Not doing so results in the attribution for some rotation of small negative surface areas when the grain percentage fall between 33.3% and 35% or between 65% and 66.7%.
- Recalculations in organic soils were caused by the area updates by the NFI extrapolation method. Emissions increased by 0.6 kt C from 442.3 kt C (NIR 2018) to 442.9 kt C (NIR 2019).

6.16.2.2 Land converted to cropland

- For land converted to cropland the recalculations of the 2016 estimate resulted in smaller C losses of 5.7 kt C from 114.6 kt C (NIR 2018) to 108.9 kt C (NIR 2019). All recalculations resulted from the modification of the area inputs.
- For organic soils, the biggest change was attributed to wetland converted to cropland where a decrease of 1 kha resulted in a 10.1 kt C reduction of the emission from 37.7 kt C (NIR 2018) to 26.7 kt C (NIR 2019). For forest land to cropland and grassland to cropland a small reduction of area (< 0.2 kha) generated a small reduction in the emission (<2 kt C), which for 2016 equaled 0.6 kt C from 15.6 to 15.0 kt C for forest land to cropland and 0 kt C for grassland to cropland.
- Net C stock losses on living biomass increased by 5 kt C and DOM decreased by 0.8 kt C, due to the extrapolation method used on the NFI living biomass and area data.

6.16.3 Grassland

6.16.3.1 Grassland remaining grassland

- Total C stock changes on grassland remaining grassland were recalculated and resulted in decreased emissions in 2016 by 0.11 kt C, from 9.6 kt C to 9.7 kt C. This was mostly caused by increased removals in living biomass of 0.17 kt C. Recalculations for mineral soil pool did not generate carbon stock change differences higher than 0.05 kt C between NIR 2018 and NIR 2019. There were no changes in organic soils.

6.16.3.2 Land converted to grassland

- Total C stock loss estimates for 2016 for land converted to grasslands was 1.7 kt C smaller in the 2019 submission, primarily due to updates in the NFI affecting forest land converted to grassland.
- Most C stock gains occurred on forest land converted to grassland with 1.1 kt C from 61.2 kt C (NIR 2018) to 60.0 kt C (NIR 2019).
- C stock losses in living biomass and DOM on forest land converted to grassland were decreased by 0.2 kt C and 2.5 kt C, respectively, from 26 kt C to 25.9 kt C for living biomass and from 92.6 kt C to 90 kt C for DOM. The changes were caused by the extrapolation method of the NFI data used for biomass and area estimates.
- Emissions from mineral soils were also recalculated due to the NFI area updates, and resulted in a minor decrease in the C uptake of 1.6 kt C from 58.6 kt C (NIR 2018) to 57.0 kt C (NIR 2019).

6.16.4 Wetlands

6.16.4.1 Wetlands remaining wetlands

- Carbon stock uptake in living biomass on wooded mires was recalculated due to the NFI extrapolation method. The recalculation resulted in a slight decrease of 1.0 kt C, from 20 kt C (NIR 2018) to 19 kt C (NIR 2019).

6.16.4.2 Land converted to wetlands

- A minor recalculation for forest land converted to wetlands in the living biomass pool resulted in a reduction of 1.9 kt C.

6.16.5 Settlements

6.16.5.1 Settlements remaining settlements – organic soils

- Emission from organic soil increased consistently by ~0.7 kt C between NIR 2018 and NIR 2019 for most reported years. For 2016 the emission increased by 5.7 kt C from 54.1 kt C to 59.8 kt C between NIR 2018 and NIR 2019. This change was directly linked to an increase of surface estimates by 0.8 kha between NIR 2018 and NIR 2019.

6.16.5.2 Land converted to settlements

- Total C stock losses from the category for 2016 were reduced by 18 kt C in NIR 2018 (515 kt C) compared to NIR 2019 (497 kt C). This was a result of the combined effect of less living biomass, DOM, and mineral soils losses, and increased losses from organic soils.
- Living biomass losses on forest land converted to settlements were reduced by 7 kt C from 143 kt C (NIR 2018) to 136 kt C (NIR 2019). DOM losses were reduced 13.4 kt C from 248 kt C (NIR 2018) to 235 kt C (NIR 2019).
- Mineral soils losses were reduced 3 kt C on land converted to settlements from 63 kt C (NIR 2018) to 60 kt C (NIR 2019).
- Emissions from organic soils increased by 2 kt C from 58 kt C (NIR 2018) to 60 kt C (NIR 2019), due to a smaller organic soils area from the NFI. This occurred mostly on forest land converted to settlements. In 2016, a surface decrease of 0.8 kha led to a decrease of emission of 6.2 kt C from 55.5 to 49.3 kt C between NIR 2018 and NIR 2019. For wetland converted to settlement, in 2016, a slight surface increase of 0.2 kha led to a 1.7 kt C increase of the emission from 2.14 kt C to 3.84 kt C between NIR 2018 and NIR 2019.

6.16.6 Other Land

6.16.6.1 Land converted to other land

- There were no recalculations for this source for neither land converted to other land nor grassland converted to other land.

6.16.7 Harvested wood products

- The FAO activity data for exported wood-based panels 2016 had been updated since last year. Hence, the export and domestically consumed (= production – export) activity data changed from 106 297 m³ to 106 300 m³ and 329 703 m³ to 329 700 m³ respectively. This did not change the total emissions from 2016.

6.16.8 Direct N₂O emission from managed soils 4(l)

- There were recalculations for Direct N₂O emissions from managed soils for forest land and settlements for the year 2016. N inputs were reduced from 7 141 678 kg N/yr (NIR 2018) to 2 569 239 kg N/yr (NIR 2019). N₂O emissions were reduced by 21.5 kt CO₂ eq (0.072 kt N₂O)

from 32.8 kt CO₂ eq (0.11 kt N₂O NIR 2018) to 8.9 kt CO₂ eq (0.03 kt N₂O NIR 2019). This was due to double counting in both the LULUCF and agricultural sector of organic N fertilizers applied to forest lands by grazing animals. The source is now being reported only in the agriculture sector.

6.16.9 Emissions and removal from drainage of organic soils 4(II)

- In forest land, CH₄ emission increased by 0.12 kt (3 kt CO₂ eq) from 2.11 kt (52.75 kt CO₂ eq) to 2.23 kt (55.75 kt CO₂ eq) while N₂O decreased by 0.11 kt (32.78 kt CO₂ eq) from 1.03 kt (306.94 kt CO₂ eq) to 0.92 kt (274.16 kt CO₂ eq) for year the 2016 between NIR 2018 and NIR 2019. This change was mostly due to the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively. For methane only, the emission factor for cropland converted to forestland, erroneously set to temperate zone (7.8625 kg of CH₄ ha⁻¹ yr⁻¹), was corrected to match that of the boreal nutrient rich zone (7.375 kg of CH₄ ha⁻¹ yr⁻¹). However the induced change was only visible with four decimals numbers. Finally, the modification of the area of drained organic soil also induced some slight recalculation not visible in the year 2016.
- In cropland, CH₄ emission decreased by 0.07 kt (1.75 kt CO₂ eq) from 3.65 kt (91.25 kt CO₂ eq) to 3.58 kt (89.5 kt CO₂ eq) for year 2016 between NIR 2018 and NIR 2019. This change was directly related to a 1.3 kha decrease of area.
- In grassland, CH₄ emission increased by more than twofold for all reported dates between NIR 2018 and NIR 2019 (e.g. from 0.07 to 0.15 kt CH₄ in 2016). The correction of an emission factor was responsible for these changes. This emission factor was calculated from the emission factors of ditch vs non-ditch area weighted by the respective proportion of these areas. Between NIR 2018 and NIR 2019, the emission factor for drained organic soil outside of the ditch was reevaluated from 36 kg to 39 kg of CH₄ ha⁻¹ yr⁻¹ while, the proportion of non-ditch area that was mistakenly set to 9.5% instead of 95% was corrected. As a consequence the resulting emission factor changed from 30.06 kg CH₄ ha⁻¹ yr⁻¹ to 60.55 kg CH₄ ha⁻¹ yr⁻¹.
- There were no recalculations for Wetlands peat extraction lands for 2016.

6.16.10 Direct N₂O emissions from N mineralization and immobilization 4(III)

- The area changes caused by NFI updates caused only minor recalculations for this source in the year 2016 and throughout the timeseries.

6.16.11 Indirect N₂O emissions from managed soils 4(IV)

- Recalculations for atmospheric deposition resulted in reduced N₂O emissions from 0.0022 kt N₂O (NIR 2018) to 0.0008 (NIR 2019). Recalculations for nitrogen leaching and run-off also resulted in reduced N₂O emissions from 0.0516 kt N₂O (NIR 2018) to 0.0343 kt N₂O (NIR 2019) for the year 2016. The reductions for these two sources was due to organic N fertilizer inputs from grazing animals being reported in the agriculture sector starting in the 2019 submission.

6.16.12 Biomass burning 4(V)

- There were no recalculations for this source in the year 2016.

7 Waste (CRF sector 5)

7.1 Overview

This sector includes emissions from landfills (CRF 5A), Biological treatment of solid waste (CRF 5B), Incineration and open burning of waste (CRF 5C), and Wastewater treatment and discharge (CRF 5D). Waste incineration from plants with energy utilization is accounted for under Energy industries (CRF 1A1). Waste incineration included in CRF 5C are emissions of greenhouse gases other than CO₂ from methane flared at landfills, and emissions from combustion of hospital waste in hospital incinerators.

The emissions of greenhouse gases from the waste sector decreased by 46.5 % (1.0 million tonnes CO₂ equivalents) from 1990 to 2017. The reductions were mainly due to decreased CH₄ emissions from landfills. Indeed, they decreased by 52.6 % from 1990 to 2017, corresponding to 1.1 million tonnes CO₂ equivalents. Emissions from Industrial wastewater decreased by 0.05 million tonnes CO₂ equivalents during the same period. Emissions from domestic wastewater handling and biological treatment of waste increased their emissions by 0.02 and 0.07 million tonnes CO₂ equivalents, respectively.

Solid waste disposal on land (i.e. in landfills) is the main emission category within the waste sector, accounting for, in 2017, 81.5 % of the sector's total emissions. Wastewater handling in domestic and industrial sectors account for 9.0 and 3.0 % of the sectors emission. Biological treatment of waste accounts for 6.2 % of emissions from the waste sector. Emission from waste incineration, without energy recovery, are only minor emissions.

The waste sector accounted for 2.3 % of the total GHG emissions in Norway in 2017. Table 7.1 presents the key categories included in the Waste sector.

Table 7.1 Key categories in level or trend in the Waste sector

IPCC	Source category	Gas	Key category according to approach	Method
5A1	Managed Waste Disposal on Land	CH ₄	2	Tier 2
5D	Wastewater treatment and discharge	CH ₄	2	Tier 2
5B	Biological treatment of Solid Waste	CH ₄	2	Tier 2

Source: Statistics Norway/Norwegian Environment Agency

7.2 Managed Waste Disposal on Land – 5A1

7.2.1 Anaerobic managed waste disposal sites, 5A1a (Key category for CH₄)

7.2.1.1 Description

CH₄ and non-fossil CO₂ are emitted during biological decomposition of waste. This transformation of organic matter takes place in several steps. During the first weeks or months, decomposition is aerobic, and the main decomposition product is CO₂. When there is no more oxygen left, the decomposition becomes anaerobic, and methane emissions start to increase. After a year or so, CH₄ emissions reach a peak, after that the emissions will decrease over some decades (SFT 1999a), (NCASI 2004).

The emissions of methane from landfills have decreased since 1990 and specifically after 1998 due to reduction of the amount of degradable waste disposed at disposal sites. This emissions reduction is the result of several policies and measures introduced in the waste sector, particularly in the 1990s. With few exceptions, notably the mixed waste from households in municipalities with a source separation of food waste, it was then prohibited to dispose easy degradable organic waste, sewage sludge included, at landfills in Norway.

From 1999 to 2014, a tax was introduced on waste delivered to final disposal sites. In 2014, this tax was 294 NOK per tonne waste. From July 1st 2009, it was prohibited to deposit biodegradable waste to landfills. This results in further reduction of methane emissions.

In addition to the above described policies and measures, landfills receiving biodegradable waste (waste containing degradable organic carbon (DOC)) are required to collect and treat landfill gas. In 2016, 71 landfills, which had installed a landfill gas extraction system, reported extraction of gas. 7 960 tonnes of methane were recovered representing a 7 % decrease compared to 2015. Methane extraction increased until 1998, and then underwent fluctuations between 1999 and 2008. The fluctuations were due to instability in the pipeline systems. Due to needs for maintenance of the pipeline system, methane extraction was reduced.

Since 2008, extraction has had a decreasing trend. This can be explained by the increased amounts of waste recycled. The total amount of waste generated has increased by almost 60 % from 1995 to 2016, but due to the increase in material recycling and energy utilization, the amount disposed at landfills has dropped substantially since 2008. As a consequence of the prohibition against depositing of biodegradable waste of July 1st 2009, there has been a strong decrease in waste depositing.

Since building the necessary treatment capacity would take time, temporary exemptions were granted in certain cases during a transitional period. Many permits for disposal of biodegradable waste had been given for one year extra, some extended out 2010, and a few within 2011. The transitional period ended on December 31st 2012.

In 2016, methane emissions from managed waste disposal sites were 1.0 million tonnes CO₂-equivalents.

Emissions of CH₄ from solid waste disposal are key category in level assessment for both 1990 and 2016 and in trend assessment due to uncertainty in AD and EF. It should be noted that the IEF for CH₄ varies due to variation of the amount of extracted CH₄ from the landfills.

There are no known semi anaerobic disposal sites in Norway, according to expert judgment (Skullerud, Pers. Comm)³², only managed anaerobe disposal sites.

Figure 7.1 shows the relative change (Index 1995=100%) in methane emissions from landfills, extraction of methane, solid waste disposed at landfills and total amount of waste generated in Norway.




Figure 7.1 Relative change in methane emissions from solid waste disposal, annual MSW at the SWDS, methane extracted from landfills and total amount of waste generated in Norway. 1990-2017 Index 1995 = 1.

Source: Statistics Norway/Norwegian Environment Agency

7.2.1.2 Methodological issues

The model proposed in the IPCC 2006 Guidelines has been used.

This model starts with the calculation of the amount of decomposing $DDOC_m$ (mass of decomposable organic carbon = the part of DOC (degradable organic carbon) that will decompose (degrade) under anaerobic conditions) contained in the material being landfilled.

As this is a first order reaction, the amount of product formed will always be proportional to the amount of reactant. This means that it is of no concern to the process when the $DDOC_m$ came into the landfill. As far as we know the amount of $DDOC_m$ in the landfill at the start of each years, all years can be considered individually in the calculations. This simplifies the model.

With the start of the reaction set to be January 1st the year following the landfilling, the “motor” of the new calculating model has been made out of these two very simple equations:

$$(7.1) \quad DDOC_{mdiss} = (DDOC_{ma(ly)} + DDOC_{md}) \times (1 - e^{-k})$$

³² Håkon Skullerud 2014: Personal communication by telephone. Statistics Norway

$$(7.2) \quad DDOC_{ma} = (DDOC_{ma(ly)} + DDOC_{md}) \times e^{-k}$$

Equation (7.1) calculates DDOC_{mass} decomposing (DDOC_{mdiss}) as the sum of the not decomposed DDOC mass accumulated from the previous years (DDOC_{ma(ly)}), and the DDOC mass landfilled during the previous year (DDOC_{md}).

Equation (7.2) calculates the DDOC mass accumulated as not decomposed (DDOC_{ma}), which is then used as (DDOC_{ma(ly)}) for the following year's calculations using equation (7.1).

The mass of decomposable organic carbon landfilled (DDOC_{md}) is estimated using equation (7.3).

$$(7.3) \quad DDOC_{md} = W \times MCF \times DOC \times DOC_f$$

The amount of decomposed DDOC_m being estimated, CH₄ produced, and CH₄ emitted are estimated using the equations stated below ((7.4)-(7.9)).

If the reaction of decomposition is set to start during the year of landfilling and not on January 1st of the following year, equations (7.1) and (7.2) need to be adjusted.

Equation (7.1) must be replaced by equations (7.4) and (7.5).

$$(7.4) \quad DDOC_{mdi} = DDOC_{md} \times \left(1 - e^{-k \times \left(\frac{13-M}{12}\right)}\right)$$

$$(7.5) \quad DDOC_{mdiss} = DDOC_{ma(ly)} \times (1 - e^{-k}) + DDOC_{mdi}$$

Equation (7.2) must be replaced by equations (7.6) and (7.7).

$$(7.6) \quad DDOC_{ml} = DDOC_{md} \times e^{-k \times \left(\frac{13-M}{12}\right)}$$

$$(7.7) \quad DDOC_{ma} = DDOC_{ma(ly)} \times e^{-k} + DDOC_{ml}$$

The amount of methane produced from decomposition of DDOC is estimated using equation (7.8).

$$(7.8) \quad CH_{4\,prod} = DDOC_{mdiss} \times F \times \frac{16}{12}$$

Methane emissions are estimated from the amount of methane produced and the amount of methane recovered. Equation (7.9) details the calculations.

$$(7.9) \quad CH_{4\,emitted\,in\,year\,T} = \left(\sum CH_{4\,prod}(T) - R(T)\right) \times (1 - OX)$$

Where, in equations (7.1)-(7.9):

W : amount landfilled

MCF : Methane Correction Factor

M : Month number for reaction start. (January 1, year after landfilling, M=13)

DOC : Degradable Organic Carbon

DOC_f : Fraction of DOC decomposing, anaerobic conditions

DDOC_m : Mass of Decomposable Organic Carbon, anaerobic conditions

$DDOC_{md}$: DDOC mass landfilled
$DDOC_{ml}$: DDOC mass left not decomposed from $DDOC_m$ landfilled, year of landfilling
$DDOC_{ma}$: DDOC mass left not decomposed at end of year
$DDOC_{ma(ly)}$: DDOC mass accumulated from last year
$DDOC_{mdi}$: DDOC mass decomposed from $DDOC_m$ landfilled, year of landfilling
$DDOC_{mdiss}$: DDOC mass decomposed in calculation year
CH_4_{prod}	: CH_4 produced
F	: Fraction of CH_4 by volume in generated landfill gas
16/12	: Conversion factor from C to CH_4
R(T)	: Recovered CH_4 in year of calculation
OX	: Oxidation factor (fraction).

7.2.1.3 Activity data

The methane is formed by decomposition of biological waste in landfills. The decomposition time varies from material to material. Easily degradable waste (food, etc.) has shortest decomposition time, while wood waste has the longest decomposition time. Other materials do not emit methane at all, either because they are inorganic (metal, glass, etc.) or because they break down extremely slowly (plastic). It is therefore of vital importance for the calculations that the waste quantities used as input to the model are correct, both total quantity and the distribution by material.

Data over the amount of different waste materials is taken from Statistics Norway's waste accounts. The waste accounts consist of data from several sources, such as special surveys, register data and statistics, indirect data sources such as production statistics, foreign trade statistics and different factors combined with activity data. Data from all these sources are combined in the waste accounts, which give an overview of waste quantities in Norway, divided into type of product, material, industry and method of treatment. Waste incineration in the waste accounts includes export, and is thus not comparable with the emission inventory as a substantial amount is exported to Sweden for incineration.

From 2012 onwards, data for the categories food waste, plastics, wood and paper are taken directly from the waste accounts. The amount of sludge deposited are taken from statistics on discharges and treatment of municipal waste water. In addition, there is a category "other" in the waste accounts, of which content is not known. Due to the prohibition to deposit biodegradable waste to landfills it is assumed that no methane is formed from these materials.

Historic data up until 2011 have been recalculated from the former waste category basis, to a waste material basis. The amount of each material type deposited is estimated based on surveys and sorting analyses. The model is based on types of waste materials for instance food waste (incl. garden waste), paper, wood and textiles. All sources of waste, MSW, industrial, commercial, construction and demolition waste are accounted for in these annual surveys.

Municipal landfills

Historical data for years before 1973 on municipal solid waste deposited are based upon:

1. New statistics on municipal waste, divided into household waste and industrial waste (1974 to 1997),
2. Estimates based on population,
3. Assumption that less people were connected to public waste management during the forties and fifties.

Since 1974, the amount of municipal waste is based upon questionnaires and linear interpolations. Surveys were held in 1974, 1980 and 1985. The amount of waste going to landfills is allocated to material based on sorting analyses. For the period 1995-2011 the amounts of waste is taken from the waste accounts, with three adaptations:

- Wood content in sludge deposited at industrial sites is added to the amount of deposited wood from the waste accounts,
- Textiles are supposed to consist of 50 % plastic (SFT 2005b). The plastic fraction of deposited textiles is therefore subtracted from the amount of deposited textiles and added to deposited plastic,
- The material category "Other materials" is assumed to contain degradable organic matter with an average half-life. This degradable share is added to the amount of paper. The amount is estimated by $0.2 * \text{landfilled 'other materials' from manufacturing} + 0.5 * \text{'other combustible' in landfilled mixed waste from all sectors}$.

Contaminated soils are assumed not to develop methane in landfills. The same applies to waste used as cover material, due to excess oxygen availability.

No bio-degradable hazardous waste is landfilled in Norway.

No organic waste is imported for landfilling, as it is prohibited.

Due to lack of data, linear interpolation of the amount of waste deposited has been applied to the period 1985-1995.

Table 7.2 Amounts deposited in municipal SWDS, 1945-2017. 1 000 tonnes.

Year	Food	Paper	Wood	Textile	Sewage sludge	Plastics
1945	75	148	120	3	7	11
1950	116	228	171	4	10	17
1955	131	256	207	5	11	19
1960	171	335	258	6	14	25
1965	258	422	270	8	18	50
1970	279	463	307	9	20	54
1975	305	513	318	10	22	59
1980	343	584	300	11	23	66
1985	357	635	280	11	24	68
1990	342	461	280	22	21	144
1995	327	286	279	33	17	219
2000	253	249	194	29	13	189
2005	218	195	169	26	4	164
2006	223	217	165	26	6	171
2007	223	227	166	28	2	186
2008	205	216	160	27	2	180
2009	138	143	106	18	3	126
2010	71	69	54	9	2	65
2011	29	33	23	3	2	28
2012	0	1	8	0	1	3
2013	0	1	0	0	2	2
2014	1	1	0	0	5	2
2015	3	2	2	0	3	3
2016	1	0	5	0	1	3
2017*	1	0	5	0	8	3

*Figures for the last inventory year are set equal to the previous year because the waste accounts are not updated in time for the emission inventory calculations. Source: Statistics Norway/Norwegian Environment Agency

Industrial disposal sites

Historical data for industrial waste for years before 1970 are estimated by extrapolation using the same trend as for municipal waste. After 1970, literature studies and information from the industrial waste study from the years 1993, 1996, 1999 and 2003 have been used. Linear interpolation is used for the years where data are missing.

Data from each landfill site with methane recovery units are reported by the landfills via an electronic web portal and the Norwegian Environment Agency assembles these data in their own

database. Further, these data are imported into the national model for calculating methane from landfills.

Table 7.3 Waste amounts deposited in industrial SWDS, 1945- onwards . 1 000 tonnes.

Year	Food	Paper	Wood	Textile	Plastics
1945	476	339	74 160	756	180
1950	735	524	101 108	1 167	277
1955	825	588	128 056	1 310	311
1960	1 077	767	155 004	1 710	406
1965	1 776	888	181 952	1 869	888
1970	2 000	1 000	208 900	2 000	1 000
1975	2 000	1 000	208 900	2 000	1 000
1980	2 000	1 000	173 872	2 000	1 000
1985	2 000	1 000	138 844	2 000	1 000
1990	2 000	1 000	103 817	2 000	1 000
1995	1936	899	88 800	1 957	759
2000	0	200	677	0	0
2001	0	38	0	0	0
2005- onwards	0	0	0	0	0

Source: Statistics Norway/Norwegian Environment Agency

7.2.1.4 Emission factors

All parameters used in the Norwegian model are IPCC defaults values for Northern Europe. Table 7.4 shows some of the variables used in the calculations of methane emissions from solid waste disposals both municipal and industrial.

Table 7.4 Variables used in the calculations of methane from landfills.

Variables	Type of waste				
	Food waste	Paper	Wood	Textiles	Sewage sludge
$t_{1/2}$ (half life time) (years)	3.7	11.6	23.1	11.6	3.7
DOC (Mg/Mg)	0.150	0.400	0.430	0.24	0.05
DOC _f (Part of DOC decomposing)	0.5	0.5	0.5	0.5	0.5
Ox. Methane oxidized in top layer	0.1	0.1	0.1	0.1	0.1
F. Part of methane in generated landfill gas	0.5	0.5	0.5	0.5	0.5

Source: IPCC (2006)

In Norway, all SWDS are considered well managed and covering including oxidising material shall apply according to the regulation on SWDS.

7.2.1.5 Uncertainties and time-series consistency

The amount of different waste materials is considered to be known within ± 20 %. The emission factors used are considered to have the uncertainty range ± 30 %. More information about the uncertainty estimates for this source is given in Annex II.

The importance of the uncertainties in calculations of methane from landfills will decrease with decreased source contribution and improved IPCC default parameter values, but most likely it will still remain among the main uncertainties in the Norwegian GHG inventory.

The methodology Statistics Norway/the Norwegian Environment Agency use to calculate methane emissions from landfills is consistent for the whole time series. The quality of the activity data used in the model has been improved in the last years. This is also the case regarding the data for recovered methane.

In 2014, a major revision of the methodologies of the waste accounts took place. The time series for waste amounts has not been recalculated to take this new information into account. There are several reasons for this, among others that many sources for the statistics do not have numbers for earlier years. Since 2012, publication divides wastes into new categories, different from the previous categories. The category “mixed waste” is no longer separated into its different material types. See Statistics Norway’s documentation of the waste accounts for more details about the revisions (Statistics Norway Annually-b). This change in the waste accounts introduces a certain degree of inconsistency in the time series of the activity data used for the calculation of methane emissions from municipal landfills. However, due to the measures described in 7.2.1.1, the amount of biological waste deposited at SWDS is currently very low, and the effect of the alterations in the waste accounts are thus considered to be negligible.

7.2.1.6 Source specific QA/QC and verification

Internal checks of time series for all emission sources are conducted every year along with the new inventory production.

Internal checks are carried out for time series of waste data, methane recovered at landfill sites and calculated methane emissions from the model. Corrections are made if necessary.

The Norwegian Environment Agency contacts landfill sites to discuss outliers and major changes in the trend of methane recovery. Corrections are made if necessary.

7.2.1.7 Recalculations

Activity data has been updated. Updated figures of sludge deposited at MSWDS for 2014 has replaced earlier used figures from 2013. This revision does not influence on the emissions estimated for 2014.

7.2.1.8 Planned improvements

There are no improvements planned for this sector.

7.3 Unmanaged Waste Disposal Sites – 5A2

In Norway, landfilling of solid waste has been regulated and controlled for some decades, and unmanaged landfills date from before 1970. Furthermore, the methane emissions for all years have been calculated from the total amounts of landfilled materials. Therefore, unmanaged waste disposal sites are not occurring and hence Norway does not separately report emissions from unauthorized/unmanaged SWDSs.

7.4 Biological treatment of Solid Waste – 5B (Key category for CH₄)

7.4.1 Composting and Anaerobic digestion of organic waste – 5B1 and 5B2

7.4.1.1 Description

This section covers the biological treatment of solid waste.

Composting is an aerobic process. A large fraction of the degradable organic carbon (DOC) in the waste material is oxidized into carbon dioxide (CO₂). CH₄ is also formed during the process, in anaerobic sections of the compost, but is largely oxidized in the aerobic sections of the compost. Composting can also produce emissions of N₂O.

Anaerobic digestion of organic waste expedites the natural decomposition of organic material without oxygen, i.e. biogas production.

In the Norwegian inventory, emissions from compost production and biogas production without energy recovery are included in this category. CH₄, N₂O and CO₂ are emitted during this process. CO₂ emissions from compost production are biogenic.

According to the approach 2 key category analysis emissions of CH₄ from Biological treatment of solid waste (CRF 5B) is a key categorie in trend assessment (1990-2016).

7.4.1.2 Methodological issues

Emissions from composting of municipal waste have been calculated according to the Tier 1 default methodological guidance available in the guidelines (IPCC 2006).

CH₄ emissions from biological treatment

$$(7.10) \text{ CH}_4 \text{ Emissions} = \sum_i (M_i * EF_i) * 10^{-3}$$

Where:

CH₄ Emissions : total CH₄ emissions in inventory year, Gg

M_i : mass of organic waste treated by biological treatment type *i*, Gg

EF : emission factor for treatment *i*, g CH₄/kg waste

i : composting or anaerobic digestion

In Norway, composting of solid biological waste includes composting of:

- organic waste from households and other sources,
- garden and park waste (GPW),
- sludge,
- home composting of garden and vegetable food waste.

CH₄ emissions from anaerobic digestion at biogas facilities are estimated based on the amount of waste treated at biogas facilities multiplied by the IPCC default emission factor. Norway is currently improving the data quality for both the amount of waste treated in biogas facilities, and the amount of energy produced. When the data is available, Norway will consider to use them in the calculation of the emissions.

Composting is performed with simple technology in Norway; this implies that temperature, moisture and aeration are not consistently controlled or regulated. During composting, a large fraction of the degradable organic carbon (DOC) in the waste material is converted into CO₂. Anaerobic sections are inevitable and cause emissions of CH₄. In the same manner, aerobic biological digestion of N leads to emission of N₂O (IPCC 2006).

N₂O emissions from composting

$$(7.11) \text{ } N_2O \text{ Emissions} = \sum_i (M_i * EF_i) * 10^{-3}$$

Where:

N₂O Emissions : total N₂O emissions in inventory year, Gg

M_i : mass of organic waste treated by biological treatment type i, Gg

EF : emission factor for treatment i, g N₂O/kg waste

Emissions from compost production are considered to be complete; calculations include composting at all nationally registered sites and best available estimated data for home composting.

7.4.1.3 Activity data

All Norwegian waste treatment plants are obligated to statutory registration and reporting of all waste entering and leaving the plants. All waste streams are weighed, categorized according to a waste type and type of treatment. Data is available for all years since 1995.

Activity data for the years since 1995 are collected from Statistics Norway's, waste statistics. Data for 1991 is also available from the waste statistics. For the year 1990, activity data for 1991 have been used, while AD for 1995 is used for 1992 to 1994.

Table 7.5 Amount of waste biologically treated at composting and biogas facilities, 1990-2017. Tonnes of waste on wet basis.

Year	Composting	Anaerobic digestion
1990	21 000	0
1995	57 000	0
2000	234 000	0
2005	319 232	4 768
2006	317 076	29 924
2007	408 706	31 294
2008	393 000	62 000
2009	354 877	83 123
2010	359 384	86 616
2011	296 000	105 000
2012	407 000	57 000
2013	441 000	77 000
2014	328 000	80 000
2015	309 000	289 000
2016	357 000	226 000
2017*	357 000	226 000

*Figures for the last inventory year have been set equal to the previous year because the waste accounts have not been updated in time for the emission inventory calculations.

Source: Statistics Norway

Home composting

The last waste category involved in composting is home composting of garden waste and vegetable waste. The activity data for this category is available for the years 2009 to 2012 from Statistics Norway. The amount of organic waste from households composted for the period 1990-2008 is estimated assuming that 3 % of all households composts their garden and vegetable food waste (Lystad 2005). The average value of the period 2009-2012, 2.6 %, has been used for the following period.

Table 7.6 Number of households with home composting and amount of organic waste composted, 1990-2017. Tonnes.

Year	Number of households with home composting	Amount of organic waste composted
1990	53 114	8 200
1995	55 980	10 234
2000	58 846	12 607
2005	61 107	15 764
2010	57 307	14 310
2011	57 479	13 703
2012	54 786	12 852
2013	58 848	14 135
2014	59 569	14 200
2015	60 356	14 390
2016	61 193	14 326
2017	61 927	14 448

Source: Statistics Norway

7.4.1.4 Emission factors

Emissions from composting, and anaerobic digestion in biogas facilities, will depend on the composition of waste composted, the amount and type of supporting material used (such as wood chips and peat), the temperature, the moisture content and the aeration during the process.

Table 7.7 gives the default factors for CH₄ and N₂O emissions from biological treatment for the Tier 1 method used in the Norwegian inventory (IPCC 2006). The CO₂ produced and emitted during composting is short-cycled C and is therefore regarded as CO₂ neutral (Boldrin et al. 2009).

Table 7.7 Composting emission factors. kg/tonnes.

	Composting	Anaerobic digestion at biogas facilities	Home composting
CH ₄	4	0.8*	4
N ₂ O	0.3	NO	0.3

*Amount of waste on a wet basis

Source: IPCC (2006)

7.4.1.5 Uncertainties and time-series consistency

The amount of waste biological treated at composting and biogas facilities is considered to be known within ± 20 %. The amount of waste composted at home is considered to be known within ± 100 %. The emission factors used are considered to have the uncertainty range ± 100 %. More information about the uncertainty estimates for this source is given in Appendix D.

The methodology Statistics Norway/the Norwegian Environment Agency use to calculate emissions from biological treatment of solid waste is consistent for the whole time series.

7.4.1.6 Source specific QA/QC and verification

Internal checks of time series for all emission sources are conducted every year along with the new inventory production.

Internal checks are carried out for time series of waste data and calculated methane emissions. Corrections are made if necessary.

7.4.1.7 Recalculations

Composting facilities

Activity data for the previous year, 2016, have been updated because the Norwegian waste accounts not has released updated figures in time for the emission inventory calculations. The correction has resulted in an increase of CH₄, N₂O and NH₃ emissions of 22 % compared to the previous calculation.

Biogas facilities

The activity data was updated for 2016 and led to a reduction of CH₄ emissions by 22 %.

7.4.1.8 Planned improvements

Norway is currently improving the data quality for both the amount of waste treated in biogas facilities, and the amount of energy produced. When the data become available, Norway will consider using them in the calculation of the emissions.

7.5 Waste incineration – 5C

7.5.1 Description

Emissions from waste incineration in district heating plants are reported under energy industries (CRF 1A1a), **as the energy** is utilized, and therefore described in Chapter 3. In 2017, there were 18 waste incineration plants where household waste was incinerated. In addition, some incineration plants burn waste other than household waste, mainly wooden waste, paper, pasteboard and cardboard. These emissions are reported and described under energy (CRF 1A1a). Waste, other than household waste, is also used as energy source in some manufacturing industries. These emissions are reported and described in the relevant subsectors under energy in manufacturing industries (CRF 1A2). Flaring off-shore and in refineries are included under fugitive emissions (CRF 1B2c). Flaring in chemical industry is included under chemical process emissions (CRF 2B8a). In this chapter, the focus will be on waste reported in IPCC sector 5C. This includes emissions from flaring of landfill gas at waste treatment plants, flaring of biogas in industry, incineration of municipal waste without energy recovery and hospital waste.

In Norway, the open burning of private yard waste is under different restrictions according to the respective municipality. These restrictions include what can be burned, but also the quantity, how, when and where it can be burned. In some municipalities, a complete ban is imposed. There is no registration of private waste burning and the activity data on this subject are difficult to estimate. Citizens are generally encouraged to compost their yard waste or to dispose of it through one of the many waste disposal/recycling sites. Emissions from open burning of waste are considered to be insignificant and have therefore not been included in the inventory.

7.5.2 Methodological issues

Landfill gas

Emissions from flaring of landfill gas by landfills are estimated. However, CO₂ emissions from flaring of landfills are not included in the inventory, as these are considered as being of biogenic origin. Emissions have been estimated by multiplying the amount of gas flared with the emission factors shown in Table 7.10.

Municipal waste incineration – CO₂ and CH₄

Net CO₂ emissions from wood/ biomass burning are not considered in the Norwegian inventory, because the amount of CO₂ released during burning is the same as that absorbed by the plant during growth. Carbon emitted in compounds other than CO₂, e.g. as CO, CH₄ and NMVOC is also included in the CO₂ emission estimates. This double counting of carbon is in accordance with the IPCC guidelines (IPCC 2006).

Municipal waste incineration – N₂O

Emissions of N₂O are derived from the emissions of NO_x, which are reported from each plant to the Norwegian Environment Agency. More specifically, an estimated amount of 2.5 % of this NO_x is subtracted and reported to UNFCCC as N₂O (SFT 1996). Accordingly, the net NO_x emissions constitute 97.5 % of the emissions reported by the plants. For some years, emissions of NO_x have not been reported for a number of plants. In these cases, specific emission factors for the plants have been

made, based upon earlier emissions and amounts of waste incinerated. These new factors have been used to estimate the missing figures.

Hospital waste

Emissions from combustion of hospital waste have been calculated based on an emission factor and the amount hospital waste incinerated. Since 2006, hospital wastes have been incinerated in municipal waste incineration plants and emissions are then reported under energy industries (CRF 1A1a).

7.5.3 Activity data

Landfill gas

The total amount of landfill gas extracted each year is reported by landfill owners to the Norwegian Environment Agency and to the Statistics Norway. The data are based on measurements of both amount of gas and CH₄ content. Most landfill owners are required to measure continuously, and as a minimum to report: Hours of operation, amount of gas extracted, volume percentage of CH₄, and amount of CH₄ used for flaring, heat, and electricity. The landfill operator reports the percentage of methane, along with the total amount of landfill gas (volume) to the Norwegian Environment Agency. The amount of recovered methane is then calculated.

Statistics Norway subtracts the amount utilized for district heating and thermal power, which is given by the energy statistics in Statistics Norway. Information on the amount flared is given by the Norwegian Environment Agency and by surveys in Statistics Norway.

Emissions from the amount of landfill gas flared is included under 5C1a, municipal solid waste. Emissions from landfill gas used for district heating and used in other sectors are reported in the relevant subsectors under Energy (CRF 1A1 and 1A4).

Table 7.8 Amount of landfill gas flared and used for energy purposes. Tonnes.

Year	5C. Flared	1A1a Public electricity and heat production	1A4a, Other sectors, commercial/institutional
1990	879	0	67
1995	6 098	208	2 533
2000	15 471	3 654	3 350
2005	10 331	187	12 777
2006	11 695	177	9 103
2007	10 420	1 767	10 558
2008	12 452	3 061	9 103
2009	12 818	4 752	7 845
2010	10 079	1 393	8 636
2011	8 581	1 600	7 745
2012	7 143	1 154	7 312

National Inventory Report 2019 - Norway

2013	6 097	971	5 789
2014	4 826	1 742	4 334
2015	4 027	902	857
2016	3 511	2 036	903
2017*	4 223	1 075	3 292

* Figures for the last inventory year have been set equal to the previous year for 5C1a and 1A4a. These figures will be updated in next year submission.

Source: Statistics Norway/ Norwegian Environment Agency

Natural gas

The amount of natural gas flared by the production of methanol is, as recommended by the ERT, reported under 2B8.

Hospital waste

The amount of hospital waste was reported to Statistics Norway for the years 1998 and 1999. For the period 1990-1997, the average for 1998 and 1999 has been used. After 1999, as there has been no collection of hospital waste data, and due to the lack of better information, the waste amount of 1999 has been used to calculate the emissions for the subsequent years.

Hospital incinerators have gradually been closed down, mainly due to new emission limit values. Since 2006, no hospital incinerator has been running and hospital waste has been incinerated in incinerators for municipal waste. Therefore, emissions are included under energy (CRF 1A1a).

Table 7.9 Estimated amount of hospital waste incinerated in hospital incinerators. 1 000 tonnes.

Year	Hospital waste incinerated
1990	0.63
1995	0.48
2000	0.24
2001	0.24
2002	0.14
2003	0.14
2004	0.14
2005	0.14
2006 onwards	0

Source: Statistics Norway

7.5.4 Emission factors

Table 7.10 presents the emission factors used for calculating emissions from flaring, cremation and hospital waste.

Table 7.10 Emission factors for flare, cremation and hospital waste incineration.

Component	Flare Landfill gas ¹ and biogas	Hospital waste
	kg/tonnes	Tonnes/tonnes
CO ₂	0	0.3
CH ₄	0.37	0.00023
N ₂ O	0.0015	0.000035

Source: ¹SFT (1996)

7.5.5 Uncertainties and time-series consistency

Activity data

Uncertainty estimates for greenhouse gases are presented and discussed in Annex II.

No new data on the amount of hospital waste has been reported since 1999. The amount of hospital waste incinerated the subsequent years may vary from the data reported in 1999 currently used in the inventory for the period 2000-2005. Uncertainty has been estimated to $\pm 30\%$.

Emission factors

Uncertainty estimates for greenhouse gases are presented and discussed in Annex II.

If the composition of the hospital waste is different from the one, which the emission factors are based on, the calculated emissions will be incorrect. Combustion engineering and processes also influence the emissions. See Annex II.

7.5.6 Source specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Section 1.2 for a description of the general QA/QC procedures of the Norwegian emission inventory.

7.5.7 Recalculations

5C11a

The activity data for the amount of landfill gas flared is updated. The change in data are due to new information from the energy statistics in Statistics Norway. This information has been used in connection with the amount reported from the landfills to the Norwegian Environment Agency and has resulted in higher activity data than in the previous years.

5C12a Municipal waste incineration

There has been a reallocation in connection to combustion of waste without energy utilisation. One plant previously included in 1A1a has now been reallocated to 5C1a. The relocation results in increased emissions in the period 1995-2016.

7.5.8 Planned improvements

7.6 Wastewater treatment and discharge – 5D (Key category for CH₄)

7.6.1 Overview

In 2016, wastewater handling accounted for 11.5 % of the emissions in the waste sector. CH₄ and N₂O emissions from Wastewater handling have been reduced by 18.4 % between 1990 and 2016, emissions been relatively stable since 1998.

Wastewater can be a source of methane (CH₄) when treated or disposed anaerobically. It can also be a source of nitrous oxide (N₂O). Carbon dioxide (CO₂) from wastewater is not considered in the IPCC Guidelines because of its biogenic origin and should not be included in national total emissions.

Sludge is produced in all wastewater handling. It consists of solids that are removed from the wastewater. This sludge must be treated further before it can be safely disposed of. In Norway, some of the wastewater sludge is treated aerobically, emissions are then included in composting (CRF 5B). Some facilities treat sludge anaerobically, producing biogas. During this process, CH₄ is produced. Emissions from the use of the produced CH₄ are included in the energy and industry sectors. Emissions of CH₄ from such facilities, due to unintentional leakages during process disturbances or other unexpected events, are included in this source category – 5D.

N₂O emissions from sewage sludge applied on fields are not treated in this chapter and are included in the LULUCF sector according to the guidelines (IPCC 2006).

According to the Approach 2 key category analysis, CH₄ emissions from wastewater handling are key category in level assessment in 1990 and trend assessment (1990-2016).

The Norwegian wastewater treatment system is characterized by a few big and advanced aerobic wastewater treatment plants (WWTP) and many smaller WWTP. In 2016, 65 % of Norway's population was connected to high-grade treatment plants with biological and/or chemical treatment. Furthermore, 18 % of the population was connected to mechanical or other types of treatment, 15 % of the population was connected to small wastewater facilities (less than 50 pe) and the remaining 2 % had direct discharges.

There is almost 2 700 wastewater facilities with a capacity of more than 50 population equivalents (pe) in Norway which treated wastewater from 85 % of the population in 2016.

The source category 5D includes estimation of CH₄ and N₂O emission from wastewater handling; i.e. wastewater collection and treatment. CH₄ is produced during anaerobic conditions and treatment processes, while N₂O may be emitted as a bi-product from nitrification and denitrification processes under anaerobic as well as aerobic conditions.

It is not possible to fully distinguish between emissions from industrial and domestic wastewater, as Norwegian industries, to a great extent, are coupled to the municipal sewer system. Wastewater streams from households and industries are therefore mixed in the sewer system prior to further treatment at centralised WWTP.

Industrial wastewater may be treated on-site or released into domestic sewer systems. If it is released into the domestic sewer system, the emissions are included in the domestic wastewater emissions (CRF 5D1). Norway estimates CH₄ emissions from on-site industrial wastewater treatment not connected to domestic sewer systems. Only industrial wastewater with significant carbon loading

that is treated under intended or unintended anaerobic conditions will produce CH₄. Industries which have been considered are:

- Pulp and paper industry,
- Chemical industry,
- Food processing industries.

As a response to previous reviews, Norway has initiated collection of activity data from Norwegian industry to enhance completeness of emissions from wastewater handling. Norway has conducted investigations on industries with separate wastewater facilities in the chemical industry, and has concluded that no company in this industry has anaerobic treatment of wastewater. In the food processing industry, all identified plants have aerobic treatment except from one. In this plant, the methane generated is flared.

Two companies in the pulp and paper industry have been identified as running anaerobic wastewater treatment facilities. The methane emissions generated from this treatment are either flared or used for energy purposes. Emissions from energy recovery have been included in energy combustion for Manufacturing Industries and construction (sector 1A2d) pulp, paper and print, for the years 1991-2016 as recovery began only in 1991.

Emissions from flaring have been included in the waste incineration sector (CRF 5C).

7.6.2 Methodological issues

7.6.2.1 Domestic wastewater

CH₄

[CH₄ from domestic waste water treatment plants](#)

Methane emissions from domestic wastewater have been calculated according to the IPCC default methodology (2006):

$$(7.12) \text{ CH}_4 \text{ Emissions} = \left[\sum_j (U_i \times T_{i,j} \times EF_j) \right] (TOW - S) - R$$

Where:

U _i :	fraction of the population I income group i
T _{i,j}	Degree of utilisation of treatment/discharge j, for each income group i
EF _j	Emission factor for treatment/discharge j, kg CH ₄ / kg BOD
TOW:	Maximum methane-producing capacity (kg CH ₄ /kg BOD)
S	Organic component removed as sludge (kg BOD / yr)
R	Amount of CH ₄ recovered in the inventory yr.

In Norway, all domestic wastewater treatment plants are aerobic and are considered well managed. Therefore, only direct discharge has been considered in the calculation of CH₄ emissions. Also, both amount of organic component removed as sludge and amount recovered CH₄ have not been considered in the calculation.

Equation (7.12) can then be simplified by equation (7.13):

$$(7.13) \text{CH}_4 \text{Emissions} = EF \times TOW$$

With:

$$(7.14) EF = B_0 \times MCF \quad \text{and} \quad (7.15) TOW = N \times BOD$$

Where:

- B_0 Maximum CH_4 producing capacity, kg CH_4 /kg BOD
- MCF: Weighted average methane correction factor by population
- N: Population total in Norway, 1000 persons
- BOD: Organic load in biochemical oxygen demand, kg BOD/1000 persons/year

Unintentional leakage of CH_4 from biogas facilities

According to IPCC (2006), emissions of CH_4 from biogas facilities may occur unintentionally due to leakages during process disturbances or other unexpected events. Emissions have been estimated as a fraction of the produced biogas using equation (7.16).

$$(7.16) \text{CH}_4 = \text{CH}_{4 \text{ generated}} \times 0.005$$

N₂O

Emissions of nitrous oxide from domestic and commercial wastewater treatment have been estimated for both the part of the population connected to large wastewater treatment plants (WWTP) (>50 pe) and the part of the population not connected to large WWTP. The former includes the part of industries connected to the large WWTP while the latter includes N_2O emissions from human sewage, which are not treated in sewage treatment.

N₂O emissions from large WWTP

N_2O emissions from the part of the population and industries connected to large treatment plants (>50 pe) have been estimated based on nitrogen content in wastewater effluent and N_2O emissions occurring as a by-product in biological nitrogen-removal plants. This method is assumed to be more precise than the IPCC Tier 1 method based on annual per capita protein intake.

N_2O emissions from domestic wastewater nitrogen effluent have been calculated by multiplying the total amount of nitrogen supplied to the residents by the IPCC default emission factor of 0.005 kg N_2O -N/kg sewage-N produced. Emissions in N needs to be converted into N_2O using the conversion ratio of N into N_2O : 44/28. Emissions have been estimated using equation (7.17).

$$(7.17) \text{N}_2\text{O} = N_{\text{supplied to pipelines}} \times 0.005 \times \frac{44}{28}$$

N_2O emissions in **biological nitrogen removal-plants** have been estimated using equation (7.18), assuming that 2 % of the nitrogen removed from the plants will form N_2O .

$$(7.18) \text{N}_2\text{O} = N_{\text{removed}} \times 0.02 \times \frac{44}{28}$$

The amount of N removed is multiplied by 2 % and emissions are then converted using the conversion ratio of N into N_2O : 44/28.

Data on the amount of nitrogen removed in biological steps of the actual wastewater treatment plants is gathered from Statistics Norway's wastewater statistics.

[N₂O emissions from other domestic wastewater handling](#)

For the part of the population that is not connected to large treatment plants, the N₂O emissions have been estimated as recommended by the IPCC review team, using a Tier 1 method. This method is based on annual per capita protein intake assuming consumed and non consumed protein.

Emissions are calculated using the equation (7.19):

$$(7.19) \quad N_2O_{(s)} = Protein \times Fncp \times Frac_{NPR} \times NR_{PEOPLE} \times EF_6 \times \frac{44}{28}$$

Where:

N ₂ O _(s) :	N ₂ O emissions from human sewage (kg N ₂ O –N/ yr)
Protein:	Annual per capita protein intake (kg/person/yr)
NR _{PEOPLE} :	Number of people not connected to treatment plants
EF ₆ :	Emissions factor (default 0.005 (0.002-0.12) kg N ₂ O –N/kg sewage- N produced)
Frac _{NPR} :	Fraction of nitrogen in protein (default = 0.16 kg N/kg protein).
Fncp	Factor for non-consumed protein added to wastewater

7.6.2.2 Industrial wastewater

Organic material in industrial wastewater is often expressed in terms of COD (chemical oxygen demand). CH₄ emissions from on-site industrial wastewater treatment are estimated based on the amount COD released into recipient. Emissions of methane from industrial wastewater are calculated according to the IPCC default methodology described in equation (7.20):

$$(7.20) \quad CH_4 = COD \times B_0 \times MCF$$

Where:

<i>COD</i> :	chemical oxygen demand (industrial degradable organic component in wastewater)
<i>B₀</i> :	Maximum methane-producing capacity (kg CH ₄ /kg COD)
<i>MCF</i> :	Methane correction factor

Emissions from the following industries are included in the Norwegian inventory:

- Pulp and paper industry
- Chemical industry
- Food processing industries

CH₄ and N₂O emissions from industries connected to large treatment plants (>50 pe) are included in domestic waste water handling sector (CRF 5D1).

7.6.3 Activity data

7.6.3.1 Domestic wastewater

CH₄ emissions from domestic WWTP

Norwegian population data are extracted from Statistics Norway's population statistics. A country-specific value of 21.9 kg BOD/person/year has been used for the degradable organic component value in the waste (D), for all years (Berge & Mellem 2013).

Unintentional leakage of CH₄ from biogas facilities

Productions of biogas from biogas facilities (CH₄_{generated}) are reported to the Norwegian Environment Agency.

N₂O emissions from large and small WWTP

Data for the amount of nitrogen released into recipients are extracted from Statistics Norway's waste water statistics.

Data for the number of people in Norway connected to waste water treatment plants are extracted from the waste water statistics at Statistics Norway (2014a).

In 1990, 75 % of the Norwegian population was connected to WWTP. The population connected to large WWTP (>50 pe) is available for the whole period 1990-2016, while the population connected to small WWTP (<50 pe) is only available after 2002. Knowing the total Norwegian population connected for 1990, population connected to small WWTP has been estimated by interpolation for the period 1990-2002.

N₂O emissions from other domestic wastewater handling

Protein is annual per capita protein intake (kg/person/year). The Directorate for Health and Social Affairs has estimated the amount of daily per capita protein intake for Norway for 1997 (Johansson L. Solvoll 1999). In 1997, the daily per capita protein intake for Norway amounted to 86 gram, corresponding to 31.39 kilos per person per year.

No similar survey has been performed since then. Nevertheless, for the years 1990, 1995, 1999, 2000 and the period 2003-2016, the Norwegian Directorate for Health has estimated the potential protein intake for the population (Directorate for Health and Social Affairs 2013). Potential protein intake has been estimated using equation (7.21):

$$(7.21) \text{ Potential protein intake} = \text{production} + \text{import} - \text{export}$$

This estimation assumes that all the products are eaten and does not take into consideration the food ending up as waste. To avoid overestimations, potential protein intake are not used directly as protein intake in the inventory. Indeed, the trend of potential protein intake has been extrapolated to the protein intake estimated for 1997 (1.39 kilos per person) so as to build the time series. These estimations rely on recommendations from the Directorate for Health and Social Affairs (Johansson, pers. Comm.³³). Table 7.11 presents the potential protein intake in both g/person/day and kg/person/year and the estimated annual protein intake per capita.

³³ Johansson, L. (2005): Personal information by telephone, Directorate for Health and Social Affairs

Table 7.11 Potential protein intake, and estimated protein intake, in g/person/day, kg/person/year, for the years 1990-2017.

Year	Potential protein intake g/person/day	kg/person/year	Index 1997 =100	Estimated protein intake kg/person/year
1990	94	34.3	100.5	31.6
1995	93	33.9	99.5	31.2
2000	95	34.7	101.6	31.9
2005	100	36.5	107.0	33.6
2006	98	35.8	104.9	32.9
2007	105	38.3	112.3	35.3
2008	104	38.0	111.2	35.0
2009	102	37.2	109.1	34.2
2010	100	36.5	107.0	33.6
2011	100	36.5	107.0	33.6
2012	100	36.5	107.0	33.6
2013	101	36.9	108.0	33.9
2014	99	36.1	105.9	33.2
2015	99	36.1	105.9	33.2
2016	100	36.5	107.0	33.6
2017*	100	36.5	107.0	33.6

Numbers in bold come from the Norwegian Directorate for Health and Social Affairs

*Figures for the last inventory year are set equal to the previous year because data from the Norwegian Directorate for Health and Social Affairs are not updated in time for the emission inventory calculations.

Non-consumed protein has been taken into account in the estimation of protein added to wastewater. It has been set to 10 % of consumed protein. Therefore, Fncp, the factor for non-consumed protein added to wastewater has been set equal to 1.1.

7.6.3.2 Industrial wastewater

CH₄ emissions from industrial WWTP

The amount COD released into recipient is reported by industries to the Norwegian Environment Agency.

Table 7.12 Reported COD released into recipient by industries, in tonn/year, for the years 1990-2017.

Year	COD released into recipient from industrial WWTP (tonn)
1990	143 748
1995	125 066
2000	91 276
2005	87 472
2006	85 975
2007	83 965
2008	79 723
2009	77 377
2010	69 656
2011	84 997
2012	72 079
2013	64 511
2014	60 948
2015	56 442
2016	55 611
2017	53 102

Source: Statistics Norway

N₂O emissions from industrial WWTP

N₂O emissions from industries connected to large treatment plants (>50 pe) are included in domestic wastewater handling sector (CRF 5D1) while N₂O emissions from industries with their own WWTP are not estimated and therefore, not included in this inventory.

7.6.4 Emission factors**CH₄ emissions from domestic and industrial WWTP**

The default emission factor for B₀ of 0.6 kg CH₄/kg BOD is used (IPCC 2006). The methane correction factor (MCF) is, according to good practice, given by the fraction of BOD that will ultimately degrade anaerobically. MCF assumed to be equal to 1 for the population connected to tanks with anaerobic conditions. Information on the part of the population connected to tanks with anaerobic conditions are taken from Statistics Norway (wastewater statistics). It corresponds to the population connected to the WWTP categorized as "Sealed tank" and a fraction of the population connected to the category "Separate toilet system". These are the treatment methods assumed to be anaerobic and hence to emit CH₄. In 2000 it is assumed that 1,1% of the population is connected to that type of treatment and in 1990, it is assumed that 2 % of the population was connected to anaerobic treatment systems. For the period 1990-2000, it is assumed that the share gradually decreased. The estimated trend is consistent with the estimated factors for the period 2000-2017. Table 7.13 gives an overview of the part of the population connected to tanks with anaerobic conditions used in the Norwegian emission inventory.

Table 7.13 Part of the population connected to tanks with anaerobic conditions for the period 1990-2017.

	1990	1995	2000	2005	2010	2011
MCF	2.0%	1.5%	1.1%	0.9%	0.9%	0.8%
	2012	2013	2014	2015	2016	2017
MCF	0.8%	0.9%	0.8%	0.8%	0.8%	0.8%

Source: Statistics Norway

Unintentional leakage of CH₄ from biogas facilities

Unintentional leakages are generally between 0 and 10 % of the amount of CH₄ generated. In the absence of further information, 5 % is used as a default value for estimating the CH₄ emissions from unintentional leakage from Norwegian biogas facilities.

N₂O emissions from large WWTP

N₂O emissions in **biological nitrogen removal-plants** have been estimated assuming that 2 % of the nitrogen removed from the plants will form N₂O. This country-specific emission factor is given in SFT (1990). This assumption is based on measurements in plants and comparisons to factors used in Sweden. This emission factor is used for all plants except for one.

It has been hypothesized that one plant had a much higher performance, i.e. a lower percentage of processed N emitted as N₂O. In 2011, N₂O emissions were measured at various spots within the treatment plant, as well as the concentrations of N₂O in the liquid phase throughout, including the exit water. The results verified that the performance of this process, with respect to N₂O emission, is much better than the emission factor used for the other treatment plants. On the average, the emission of N₂O -N to air from the entire plant (through the chimney) amounted to 0.2 % of the processed N. If the N₂O lost as dissolved N₂O in the exit water is included, the percentage increases

to 0.3 (Bakken et al. 2012). For this WWTP, it has then been assumed that 0.3 % of the nitrogen removed from plants will form N₂O. This emission factor has been used for that plant for all years since 1996, the year when the nitrification and denitrification reactors were fully operational. For the period 1990-1996, the default emission factor of 2 % has been used.

N₂O emissions from other domestic waste water handling

For the part of the population that is not connected to large WWTP, IPCC default emission factors have been used:

- EF₆: 0.005 kg N₂O/kg sewage-N produced has been used.
- Frac_{NPR}, the fraction of nitrogen in protein, has been set equal to 0.16 kg N/kg protein.

N₂O emissions from on-site industrial WWTP

N₂O emissions from industries with their own WWTP are not included in this inventory. A quantitative assessment has been conducted with emissions from other European countries and has shown that emissions are far under the estimating threshold. Therefore N₂O emissions from on-site industrial WWTP have not been included in the inventory.

7.6.5 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are presented and discussed in Annex II. A general assessment of time series consistency has not revealed any time series inconsistencies in the emission estimates for this category.

7.6.6 Source specific QA/QC and verification

There is no source specific QA/QC procedure for this sector. See Section 1.2 for the description of the general QA/QC procedure.

7.6.7 Recalculations

There has been two recalculations in 5D2, industrial wastewaterhandling:

- The activity data was updated in the period of 2000-2016. The new information resulted in a reduction of emissions of CH₄. The decline is marginal in the beginning of the periode and largest in 2010-2013 with a reduction of emissions around 10 %.
- The activity data used for estimateing NMVOC from industrial wastewater was also updated for 2016, leading to an increase in the emissions for NMVOC of 31 %.

7.6.8 Planned improvements

Emissions from the amount of CH₄ that is flared or recovered for energy use has been reported to the energy sector and should be subtracted from total emissions. Following up 2016 review, Norway has begun to investigate possible double counting between emissions from waste water treatment and emissions from flaring included in sector 1A2 (manufacturing industries and construction). The results of this investigation will be presented in the next submission.

7.7 Other emissions sources from the waste sector – 5E

7.7.1 Description

No GHG emissions are currently included in this category.

7.7.2 Recalculations

No recalculation has been made in this category.

7.7.3 Planned improvements

There are currently no planned improvement for this source category.

8 Other (CRF sector 6) (if applicable)

9 Indirect CO₂ and nitrous oxide emissions

9.1 Description of sources of indirect emissions in GHG inventory

According to the reporting guidelines to the Climate Convention, all emissions of carbon from fossil compounds are to be included in the national emission inventory. When methane or NMVOC are oxidised in the atmosphere, indirect CO₂ emissions are formed. The emissions of CH₄, CO and NMVOC from some sources will partly be of fossil origin and should therefore be included. Fossil carbon in fuels combusted are included in the emission inventory due to the fact that the CO₂ emissions factors take into account the fossil carbon in the fuel and that complete combustion is assumed. These indirect CO₂ emissions are included in the Norwegian emission inventory. However, indirect CO₂ emissions from non-combustion sources originating from the fossil part of CH₄, CO and NMVOC are taken into account separately, calculated on the basis of average carbon content.

Indirect emissions of N₂O from NO_x and NH₃ from energy, industrial processes and waste are included as memo items in the inventory. For agriculture, only indirect emissions from burning of crop residues are included as memo items. Indirect emissions from manure management, fertilizers, and etcetera are included in the inventory proper.

In NIR 2015 Norway wrote the following “Indirect CO₂ emissions from CO is not included in the inventory this year. We assume that indirect CO₂ emissions should have been included for silicium carbide, magnesium production and well testing off shore. We estimate the emission to vary between 15-90 000 t CO₂, the lower part of the interval the latest year.” We looked into this issue and concluded in NIR 2016 not to estimate indirect CO₂ emissions from the three sources and the arguments are as followed

- Well testing
The CO₂ emission factor for well testing of oil in 2013/14 is 3.17 and 3.20 kg CO₂/kg oil and for natural gas 2.34, 2.5 and 3.73 kg CO₂ per Sm³. This indicate that all carbon in the oil and gas is included in the emission factors and hence additional estimation of indirect CO₂ emissions of CO would imply double counting.
- Production of magnesium, silicon and calcium carbide
Estimated CO₂ emissions from the three source categories included in the inventory includes all carbon that is put into the processes and carbon in products and dust is deducted. Therefore, indirect CO₂ emissions from CO are not estimated to avoid double counting.

Fossil carbon in the emissions of CH₄ and NMVOC from several non-combustion sources are included in the Norwegian emission inventory. See Table 9.1.

Table 9.1 Source categories in the inventory where indirect CO₂ emissions is calculated for CH₄ and NMVOC.

1.B.1.a: Coal Mining and Handling
1.B.2.a.3: Oil and Natural Gas and Other Emissions from Energy Production; Oil; Transport
1.B.2.a.4: Oil and Natural Gas and Other Emissions from Energy Production; Oil; Refining/Storage
1.B.2.a.5: Oil and Natural Gas and Other Emissions from Energy Production; Oil; Distribution of Oil Products
1.B.2.b.2: Oil and Natural Gas and Other Emissions from Energy Production; Natural Gas; Production
1.B.2.c: Oil and Natural Gas and Other Emissions from Energy Production; Venting and Flaring
2.B.5: Carbide Production
2.B.8.a: Petrochemical and Carbon Black Production; Methanol
2.B.8.b: Petrochemical and Carbon Black Production; Ethylene
2.B.8.c: Petrochemical and Carbon Black Production; Ethylene Dichloride and Vinyl Chloride Monomer
2.C.2: Ferroalloys Production
2.D.3: Solvent use

Indirect CO₂ emissions have been included in the Norwegian emission inventory for many years. Indirect CO₂ emissions are included in the emission estimates for each source category at the most disaggregated level, and are thus included in the sums named "Total CO₂ equivalent emissions without land use, land-use change and forestry" and "Total CO₂ equivalent emissions with land use, land-use change and forestry" in the summary tables in the CRF Reporter. Thus, in order to achieve correct totals including indirect CO₂, table 6 of the CRF Reporter does not include indirect CO₂ emissions, as this would have led to double counting in the summary table totals "including indirect CO₂". The indirect CO₂ emissions are given in Table 9.2 for transparency.

National Inventory Report 2019 - Norway

Table 9.2 Indirect CO₂ emissions from CH₄ and NMVOC, 1990-2017. Kilotonnes.

	Energy	IPPU	Agriculture	LULUCF	Waste	Total
1990	475.66	119.19	NA	NA	NE	594.85
1991	513.17	104.84	NA	NA	NE	618.01
1992	607.16	108.31	NA	NA	NE	715.47
1993	672.82	108.84	NA	NA	NE	781.67
1994	712.76	116.07	NA	NA	NE	828.83
1995	767.22	113.31	NA	NA	NE	880.53
1996	774.68	119.60	NA	NA	NE	894.28
1997	790.42	115.79	NA	NA	NE	906.21
1998	781.17	116.42	NA	NA	NE	897.59
1999	815.89	113.65	NA	NA	NE	929.53
2000	883.57	108.69	NA	NA	NE	992.26
2001	930.26	110.80	NA	NA	NE	1041.06
2002	788.97	112.54	NA	NA	NE	901.50
2003	670.41	113.41	NA	NA	NE	783.82
2004	580.59	116.02	NA	NA	NE	696.61
2005	442.06	104.93	NA	NA	NE	546.99
2006	364.58	96.97	NA	NA	NE	461.55
2007	364.59	97.25	NA	NA	NE	461.84
2008	276.08	93.66	NA	NA	NE	369.74
2009	253.49	79.21	NA	NA	NE	332.69
2010	233.22	94.53	NA	NA	NE	327.75
2011	210.12	96.42	NA	NA	NE	306.54
2012	204.86	100.03	NA	NA	NE	304.88
2013	219.17	103.47	NA	NA	NE	322.64
2014	269.15	103.66	NA	NA	NE	372.81
2015	252.07	109.98	NA	NA	NE	362.05
2016	236.30	106.29	NA	NA	NE	342.59
2017	220.59	106.42	NA	NA	NE	327.01

9.2 Methodological issues

The indirect CO₂ emissions from oxidised CH₄, CO and NMVOC are calculated from the content of fossil carbon in the compounds. For CH₄ and CO the factors for indirect emissions are simply calculated on basis of mass of molecules. For NMVOC the average carbon fraction is also taken into account. The default value for carbon fraction, 0.6, is used. This leads to the emission factors 2.75 kg CO₂/kg CH₄, 1.57 kg CO₂/kg CO and 2.2 kg CO₂/kg NMVOC. The NMVOC factor is for all other source categories than NMVOC from loading and storage of crude oil off shore. There we use the emission factor 3.0 kg CO₂ per kg NMVOC from CMR Instrumentation (2011) (Måle- og beregningsprogram for bestemmelse av utslipp av NMVOC i forbindelse med bøyelasting 2011, only in Norwegian) (Measuring program for emissions of NMVOC from loading of crude oil).

9.3 Uncertainties and time-series consistency

Uncertainty estimates for greenhouse gases are given in Annex II.

9.4 Category-specific QA/QC and verification

The general QA/QC methodology is given in chapter 1.2.3.

9.5 Category-specific recalculations

The emission factor for indirect CO₂ from NMVOC from distribution of oil products has been altered from the standard factor 2.2 to 3.13 kg CO₂ per kg NMVOC, which is the same as the combustion factor.

9.6 Category-specific planned improvements

There are no planned activities this year that will improve the data quality or the documentation for this source category.

10 Recalculations and improvements

10.1 Explanations and justifications for recalculations, including in response to the review process

The Norwegian greenhouse gas emission inventory has in 2018 been recalculated for the entire time series 1990-2016 for all components and sources, to account for new knowledge on activity data and emission factors, and to correct for discovered errors in the calculations. There is a continuous process for improving and correcting the inventory and the documentation of the methodologies employed, based on questions and comments received in connection with the annual reviews performed by the expert review teams (ERTs) under the UNFCCC. The figures in this inventory are, as far as possible, consistent through the whole time series.

The driving force for making improvements in the emission inventory is to meet the reporting requirements in the revised UNFCCC Reporting Guidelines. In addition, it is important for decision makers and others to have accurate emission estimates as basis for making decisions of what measures to introduce to reduce emissions.

10.2 Specific description of the recalculations

10.2.1 Energy

The Norwegian Energy Balance has been rebuilt in a new data system and with considerable changes in methodology and data input. This has led to changes in most categories under *energy combustion* for the whole time-series 1990-2016. The aggregated results of the recalculations are illustrated in Figure 10.1 below. The new data system is thoroughly described in the report "Energy Accounts and Energy balance – Documentation of statistics production since statistics year 1990" (SSB 2018).




Figure 10.1 Recalculations illustrated for total emissions (kt CO₂-equivalents per year) estimated for the Energy sector in the 2019 submission compared with the 2018 submission

1A1a Public Electricity and Heat Production

- Previously reported activity data where replaced by directly reported emissions. The change resulted in decreased emissions of CO₂ and CH₄ in the period 2008 to 2016.

1A3b Road transport

- Revised activity data. In the revised Energy balance some use of gasoline and diesel are moved from road transport to off-road (in boats, snow scooters, motorized equipment, etc.). Due to the revision, CO₂ emissions are lower in the years 1990-2016 in road transport. There were small changes in CH₄ and N₂O. Activity data for LPG is revised resulting in higher use of LPG in road transport.

1A3dii National navigation and 1A4ciii Fishing

- Revised activity data. Important recalculations were made in 1A3d National navigation and 1A4ciii National Fishing. Energy consumption for National Navigation has increased for all years, partly because of reallocation from International Navigation and partly because of a reallocation of energy previously reported as consumption under National Fishing. This reallocation is a consequence of new information on the use of marine gas oils from tax data. The former estimation method is believed to have overestimated the emissions from National Fishing with a correspondingly underestimation of National Navigation. The recalculations affect all components, especially CO₂.

1A2 and 1A4 Motorized equipment

- Revised activity data. In the revised Energy balance the use of gasoline and diesel has increased in off-road, causing higher emissions in these categories, except for in 1A4c-ii (agriculture/forestry/fishing).

In addition to recalculations due to the new Energy Balance, the following sectors were updated:

1A3a Domestic aviation

- Accuracy: New emission factors from EMEP/EEA, combined with updated flight data and total consumption of aviation fuel from the sales statistics of petroleum products has resulted in decreased emissions from CO₂, CH₄, N₂O and NMVOC in 2016. The new factors have been applied from 2010 - 2017, in combination with flight data for the respective reference years. Emission factors for the years prior to 2010 have been set to factors equal to the calculated factor in 2010.

1B2B5 Distribution

- Updated activity data resulted in increased emissions of CH₄ for the period 1994 to 2016.

10.2.2 Industrial processes and product use

2B5a Silicon carbide.

Correction of error. The activity data, CO₂ emissions and CH₄ emissions for the years 2010-2016 have been recalculated for one plant. The concerned plant has for these years mixed the reporting of GHG emissions to the Norwegian Environment Agency with internal reporting. For some years total crude production was used instead of total pure production and for some other years an emission factor other than 2.62 was used. This has now been corrected. The recalculations vary from a reduction in emissions of almost 2 300 tonnes CO₂ in 2013 to an increase in emissions of about 2 700 tonnes CO₂ in 2015. The changes in CH₄ emissions are minor.

2D31 Non energy products from fuels and solvent use

Correction of activity data. The activity data for 2016 in the previous submission was based on figures from 2015. In this submission the activity data for 2016 has been updated with new information from the Norwegian Product Register. The correction resulted in a reduced indirect CO₂ emissions from NMVOC by 3.2 %.

10.2.3 Agriculture

A new model for calculating the nitrogen emissions from animal manure has been implemented for estimating the emission figures for 1990-2017. The model was updated to meet the most recent requirements of the emission inventory guidelines of UNECE for ammonia and other nitrogen species (EMEP/EEA 2016) and the requirements in 2006 IPCC Guidelines (IPCC 2006). As the updated model is based on the nitrogen mass balance approach specified by EMEP/EEA, it allows estimates to be made of all the main nitrogen species, namely NH₃, N₂O, NO and N₂, in a single model. In addition, the project also aimed to update the ammonia emission factors used, in order to reflect current national and international knowledge and best practice.

These results affects all N₂O emissions that originate from animal manure, both direct and indirect emissions.

- Revision of emission factors: In the figures for 1990-2017 revised VS and Nex factors for mature non dairy cattle (beef cow) were used see Annex. For 2016 figures the Nex factor changed from 65 to 93 kg per year, and the VS factor changed from 970 to 1461 kg per year.

Table 10.1 Differences between NH₃ emissions from animal manure using old and new model

New model				Old model			% change		
Year	Manure management	Animal manure applied to soils	Urine and dung deposited by grazing animals	Manure management	Animal manure applied to soils	Urine and dung deposited by grazing animals			
	3 B	3 D a 2 a	3 D a 3	3 B	3 D a 2 a	3 D a 3	3 B	3 D a 2 a	3 D a 3
1990	10 700	13 390	1 325	5 857	15 202	1 002	82.7	-11.9	32.2
1995	9 970	11 647	1 261	5 683	12 988	954	75.4	-10.3	32.3
2000	10 279	11 678	1 367	5 832	12 851	1 029	76.3	-9.1	32.9
2005	11 110	12 613	1 429	5 495	14 602	1 021	102.2	-13.6	40.0
2011	11 377	12 789	1 331	5 760	15 152	917	97.5	-15.6	45.1
2012	11 484	12 885	1 309	5 806	15 240	912	97.8	-15.5	43.5
2013	11 686	13 055	1 303	6 149	15 412	927	90.0	-15.3	40.6
2014	11 702	13 030	1 306	6 184	15 419	930	89.2	-15.5	40.5
2015	11 732	13 065	1 326	6 119	15 498	943	91.7	-15.7	40.6
2016	11 879	13 230	1 359	6 164	15 701	963	92.7	-15.7	41.1

3B Manure management

- Revision of emission factor. The new model has an increased number of options for storage of manure which results in updated and more accurate emission factors for NH₃. The updated emission factors are higher than those used previously for both housing and storage and results in higher emissions of NH₃. The differences between total NH₃ emissions from animal manure in previous- and new model is showed in table 10.1. A higher NH₃ emission reduces the nitrogen basis and this leads to lower N₂O emissions from manure management systems as showed in Table 10.2.
- N₂O: The main reason for lower N₂O is the reassigning of manure quantities between different types of storage systems, in addition to re-categorization of manure types between slurry, farmyard manure and solid manure. Updated emission factors were specified for the newly created storage systems. In addition, emission factors for swine have been reviewed to incorporate conclusions about crust formation.

Table 10.2 Differences between direct N₂O emissions from animal manure using old and new model

New model				Old model			% change		
	Manure management	Animal manure applied to soils	Urine and dung deposited by grazing animals	Manure management	Animal manure applied to soils	Urine and dung deposited by grazing animals			
Year	3 B	3 D a 2 a	3 D a 3	3 B	3 D a 2 a	3 D a 3	3 B	3 D a 2 a	3 D a 3
1990	141	782	653	428	952	624	-66.9	-17.8	4.7
1995	164	762	631	410	921	594	-60.0	-17.3	6.2
2000	186	766	669	411	920	627	-54.7	-16.7	6.6
2005	223	814	676	450	974	638	-50.4	-16.4	5.9
2011	262	842	600	455	995	582	-42.4	-15.4	3.3
2012	272	855	589	462	1 013	561	-41.0	-15.6	4.9
2013	284	874	589	457	1 034	563	-37.9	-15.5	4.5
2014	287	875	593	460	1 045	546	-37.6	-16.3	8.5
2015	294	876	602	463	1 047	554	-36.5	-16.3	8.8
2016	302	886	619	469	1 058	565	-35.6	-16.3	9.5

3Da2 Animal manure applied to soils N₂O

- The new model includes added N in animal bedding and the consequent immobilization of ammoniacal nitrogen (TAN) in that bedding. In spite of this addition to the N basis in this stage, less nitrogen is available for spreading due to increased losses during housing and storage stages. This leads to lower N₂O emissions from animal manure applied to soils.

3Da3 Direct emissions from managed soils- urine and dung deposited by grazing animals. N₂O

- The ammonia emissions from grazing animals is higher due to higher Nex factor for beef cow, and a higher N₂O emission from this stage is estimated. More information about the revised Nex factor for beef cattle is given in section 3.2.2 in NIR Annex IX. The revision of the Nex factor for beef cattle results in recalculations for all manure related emissions of NH₃ and N₂O, for both 3B and 3D emission sources. For grazing animals the percentage of the animal year spent on pasture is used to calculate total N excreted on pasture per year. This grazing time factor for sheep younger than one year is revised due to expert opinion³⁴, which leads to a higher proportion of N emission on pasture for sheep.

3Db1 Indirect N₂O emissions from managed soils - atmospheric deposition

- Figures were revised due to new model and 10.4 % higher N₂O figures were estimated for 2016.

³⁴ Nortura (2018): Email contact Finn Avdem, November 2018.

3Db2 N₂O Leaching and run-off

- Figures were revised due to new model and 3.6 % lower N₂O figures were estimated for 2016.

3B Manure management CH₄

- Revision of emission factor: The estimation of methane emissions from manure in mature non dairy cattle (beef cow) is revised due to new VS factor for beef, see section 3.2.2 in NIR Annex IX. This was developed to be consistent with the estimations of enteric methane which was revised the previous year. The factor has changed from a variable factor 969 (2016) depending on yield, animal weight and protein content in feed, to a constant factor 1461. This has increased the estimated emission per animal and year from 7.7 to 13.1 kg CH₄.
- Updated activity data: Updated data on the distribution of manure on different storage types was implemented in the model to be consistent with the new nitrogen model. This results in a higher emission for dairy cattle, young cattle and swine. The CH₄ emission from manure management for 2016 had a total increase by 6.2 per comparing old and new estimations.

3Da6 Cultivation of organic soils

- Revision of emission factor. The estimation of nitrous oxide from cultivation of organic soils is revised due to a change in emission factor for N₂O from cultivation of organic soils. A change in emission category for Grassland from temperate deep-drained nutrient rich to temperate shallow drained was done to be consistent with LULUCF reporting, and to better represent Norwegian soil conditions. This factor was changes from for Grassland from 9.5 to 1.6 kg N₂O-N / ha. This has decreased the estimated emissions with 4.4 % in 2016.

10.2.4 Land use, Land–Use Change and Forestry

All emissions related to the Norwegian National Forest Inventory (NFI) estimated areas are recalculated every year as well as for the 2019 submission. This is due to the interpolation and extrapolation method used to estimate the areas. Since NIR 2014, areas have been estimated using linear interpolation between the 1/5th of the NFI sample plots that are surveyed every year. One fifth of the area estimates are therefore based on measured sample plots, and four fifths of the estimate is based on interpolation or extrapolation in the final years. Extrapolation affects the four final reporting years and requires a recalculation of these years in the subsequent submissions. There will therefore always be recalculation in the last few years of the time series. The area-related emissions are affected by this and comprise all carbon stock changes estimated for living biomass, litter, dead wood, DOM, mineral and organic soils (CRF Tables 4.A-4.F). In addition, the following non-CO₂ emission sources are also indirectly affected by the area estimates: 4(II) Emissions and removals from drainage and rewetting of organic soils, and source 4(III) Direct N₂O emissions from N mineralization/immobilization. In addition to the interpolation and extrapolation method affecting area estimates, NFI sample plots are surveyed either in the field (if forested) or using aerial images. Corrections of previously attributed land-use categories that are not due to actual land-use changes therefore occur if new information becomes available. This can result in revised area estimates

among all categories for the full time series. Another factor affecting recalculations is the calculation for changes in soil C, litter, and deadwood in afforestation and reforestation and deforestation for the inventory year 2016 for which a correction was made. For living biomass, a correction of the time series transition (“new remaining” starts in 2010, not 2011 as before) and the attribution of gains on lands-converted-to-not-Forest (gains are now correctly reported under the previous remaining class) affected the estimates on all land use categories with reported living biomass. Also an updated Statistics Norway harvests statistic affected the estimates of living biomass losses on all land use categories. The implementation of a Tier 1 method for living biomass changes affected the categories GS,SG and CS,SC. (Default factors for C and G are similar (4.7 t vs. 4.25 t). Therefore, no change was assumed for the GC category).

For the NIR 2019 the whole time series was recalculated for all C emission sources and sinks due to revised activity data (areas, time series revision) and a few methodological changes. It was necessary to utilize preliminary data to produce emission and removal estimates for some select sources due to an earlier reporting schedule for the 2019 submission. Preliminary data was obtained for: timber harvest statistics, area of orchard trees, areas of grassland, proportions of crop types, organic N fertilizer inputs from animals, and organic N fertilizer inputs on settlements. No updated data were available at this writing for these data. The largest recalculations occurred for the last four years (2013-2016). Total changes in emissions (including non-CO₂ emissions) caused by recalculations for the LULUCF sector are shown in Figure 10.2.




Figure 10.2 Recalculations illustrated for total emissions (kt CO₂-equivalents per year) estimated for the LULUCF sector in the 2019 submission compared to the 2018 submission.

Quantitatively, the largest changes in estimated C stock changes occurred for forest land and grasslands. The estimated CO₂ uptake on forest land for the year 2016 was 616 kt smaller in the 2019 submission compared to the 2018 submission. The estimated CO₂ emissions in grasslands for 2016 were 4.4 kt smaller (Table 10.3). The removals between the estimates of total GHG emissions from the LULUCF sector were 577 kt CO₂-eq more in the 2019 NIR compared to the 2018 NIR submission for the recalculated year 2016 (Table 10.3).

Table 10.3 Recalculated GHG emissions (kt CO₂-eq yr⁻¹) for 2016 per land-use category in the LULUCF sector.

Land-use category	Emissions for 2016 (kt CO ₂ -eq yr ⁻¹)		
	NIR 2019	NIR 2018	Absolute difference
4.A Forest land	-28 210.6	-28 827.0	616.4
4.B Cropland	2 058.9	2 038.5	20.4
4.C Grassland	191.9	196.3	-4.4
4.D Wetlands	18.4	21.6	-3.2
4.E Settlements	2 066.2	2 112.8	-46.6
4.F Other land	0.95	0.95	0
4.G HWP	85.00	85.00	0
4(IV)Total sum	10.5	16.1	-5.6
Total sum	-23 778.8	-24 355.8	577.0

The reasons for the recalculations within each land-use category and sink/source category and the effects for the last recalculated year of the inventory (2016) are described below.

4A1 Forest land remaining forest land

- Total CSC for forest land remaining forest land in 2016 was 190 kt C larger in the NIR 2018 submission compared with the NIR 2019 submission.
- Net change in living biomass decreased by 235 kt C from a C uptake of 5 823 kt C in NIR 2018 to 5 588 kt C in NIR 2019. Carbon uptake in dead wood, litter, and mineral soils increased by 6 kt C (from 362 to 367 kt C), 28 kt C (from 1 792 to 1 820 kt C), and 1 kt C (41 to 42 kt C), respectively. Recalculations were caused by the updates in the area estimates in addition to the general methodological changes described above.
- Emission from organic soil decreased consistently by 16 kt C between NIR 2018 and NIR 2019 for all reported years. For the year 2016, the value dropped by 16.4 kt C from 189.85 kt C to 173.45 kt C. This change resulted from the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively.

4A2 Land converted to forest land

- Compared to last year's reporting, the total C uptake for the year 2016 on land converted to forest land was 2 kt C larger in the NIR 2019 submission (from 143 to 145 kt C). This was mainly due to the larger C uptake in living biomass, which increased by 8 kt C from 46 kt C (NIR 2018) to 53 kt C (NIR 2019), which occurred primarily on settlements converted to forestland (7 kt C) and other land converted to forest land (1 kt C). Recalculations in the litter pool comprised the largest share (-7 kt C), while the deadwood (-0.06 kt C), mineral soil (1.4 kt C), and organic soil (-0.1 kt C) pools comprised a much smaller proportion. All recalculations were caused by the updated NFI areas.

4B1 Cropland remaining cropland

- Total C stock loss for the year 2016 on cropland remaining cropland increased by 12 kt C from 416 kt C in the 2018 submission to 428 kt C in the 2019 submission. This was primarily due to changes in mineral soil, C stock changes which strongly decreased by 11.1 kt C between NIR 2018 and NIR 2019 from 22.1 kt C to 11.0 kt C, respectively. This was mostly due to the correction of an error. Prior correction, carbon stock change for silty loam soils were counted twice when summing the changes for all soil types. A second error correction, dealing with the attribution of area of grain-ley rotations had little impact on the final C stock change. Briefly, for each agrozone there were three types of grain-ley rotations fixed by the grain percentage of the total grain-ley area. Agrozones with: >65% grain were attributed rotation with 1 third Ley and 2 third Grain (1:2LG rotation). Similarly 35-65% grain gave 1:1 LG rotation, and <35% grain gave 2:1 LG rotation (See Borgen et al., 2012). For more consistency, the choice of the grain percentage thresholds, 35% and 65 % in the previous version, were changed to 33.3% and 66.7%. Not doing so results in the attribution for some rotation of small negative surface areas when the grain percentage fall between 33.3% and 35% or between 65% and 66.7%.
- Recalculations in organic soils were caused by the area updates by the NFI extrapolation method. Emissions increased by 0.6 kt C from 442.3 kt C (NIR 2018) to 442.9 kt C (NIR 2019).

4B2 Land converted to cropland

- For land converted to cropland the recalculations of the 2016 estimate resulted in smaller C losses of 5.7 kt C from 114.6 kt C (NIR 2018) to 108.9 kt C (NIR 2019). All recalculations resulted from the modification of the area inputs.
- For organic soils, the biggest change was attributed to wetland converted to cropland where a decrease of 1 kha resulted in a 10.1 kt C reduction of the emission from 37.7 kt C (NIR 2018) to 26.7 kt C (NIR 2019). For forest land to cropland and grassland to cropland a small reduction of area (< 0.2 kha) generated a small reduction in the emission (<2 kt C), which for 2016 equaled 0.6 kt C from 15.6 to 15.0 kt C for forest land to cropland and 0 kt C for grassland to cropland.
- Net C stock losses on living biomass increased by 5 kt C and DOM decreased by 0.8 kt C, due to the extrapolation method used on the NFI living biomass and area data.

4C1 Grassland remaining grassland

- Total C stock changes on grassland remaining grassland were recalculated and resulted in decreased emissions in 2016 by 0.11 kt C, from 9.6 kt C to 9.7 kt C. This was mostly caused by increased removals in living biomass of 0.17 kt C. Recalculations for mineral soil pool did not generate carbon stock change differences higher than 0.05 kt C between NIR 2018 and NIR 2019. There were no changes in organic soils.

4C2 Land converted to grassland

- Total C stock loss estimates for 2016 for land converted to grasslands was 1.7 kt C smaller in the 2019 submission, primarily due to updates in the NFI affecting forest land converted to grassland.

- Most C stock gains occurred on forest land converted to grassland with 1.1 kt C from 61.2 kt C (NIR 2018) to 60.0 kt C (NIR 2019).
- C stock losses in living biomass and DOM on forest land converted to grassland were decreased by 0.2 kt C and 2.5 kt C, respectively, from 26 kt C to 25.9 kt C for living biomass and from 92.6 kt C to 90 kt C for DOM. The changes were caused by the extrapolation method of the NFI data used for biomass and area estimates.
- Emissions from mineral soils were also recalculated due to the NFI area updates, and resulted in a minor decrease in the C uptake of 1.6 kt C from 58.6 kt C (NIR 2018) to 57.0 kt C (NIR 2019).

4D1 Wetlands remaining wetlands

- Carbon stock uptake in living biomass on wooded mires was recalculated due to the NFI extrapolation method. The recalculation resulted in a slight decrease of 1.0 kt C, from 20 kt C (NIR 2018) to 19 kt C (NIR 2019).

4D2 Land converted to wetlands

- A minor recalculation for forest land converted to wetlands in the living biomass pool resulted in a reduction of 1.9 kt C.

4E1 Settlements remaining settlements – organic soils

- Emission from organic soil increased consistently by ~0.7 kt C between NIR 2018 and NIR 2019 for most reported years. For 2016 the emission increased by 5.7 kt C from 54.1 kt C to 59.8 kt C between NIR 2018 and NIR 2019. This change was directly linked to an increase of surface estimates by 0.8 kha between NIR 2018 and NIR 2019.

4E2 Land converted to settlements

- Total C stock losses from the category for 2016 were reduced by 18 kt C in NIR 2018 (515 kt C) compared to NIR 2019 (497 kt C). This was a result of the combined effect of less living biomass, DOM, and mineral soils losses, and increased losses from organic soils.
- Living biomass losses on forest land converted to settlements were reduced by 7 kt C from 143 kt C (NIR 2018) to 136 kt C (NIR 2019). DOM losses were reduced 13.4 kt C from 248 kt C (NIR 2018) to 235 kt C (NIR 2019).
- Mineral soils losses were reduced 3 kt C on land converted to settlements from 63 kt C (NIR 2018) to 60 kt C (NIR 2019).
- Emissions from organic soils increased by 2 kt C from 58 kt C (NIR 2018) to 60 kt C (NIR 2019), due to a smaller organic soils area from the NFI. This occurred mostly on forest land converted to settlements. In 2016, a surface decrease of 0.8 kha led to a decrease of emission of 6.2 kt C from 55.5 to 49.3 kt C between NIR 2018 and NIR 2019. For wetland converted to settlement, in 2016, a slight surface increase of 0.2 kha led to a 1.7 kt C increase of the emission from 2.14 kt C to 3.84 kt C between NIR 2018 and NIR 2019.

4F2 Land converted to other land

- There were no recalculations for this source for neither land converted to other land nor grassland converted to other land.

4G Harvested wood products

- The FAO activity data for exported wood-based panels 2016 had been updated since last year. Hence, the export and domestically consumed (= production – export) activity data changed from 106 297 m³ to 106 300 m³ and 329 703 m³ to 329 700 m³ respectively. This did not change the total emissions from 2016.

4(I) Direct N₂O emission from managed soils

- There were recalculations for Direct N₂O emissions from managed soils for forest land and settlements for the year 2016. N inputs were reduced from 7 141 678 kg N/yr (NIR 2018) to 2 569 239 kg N/yr (NIR 2019). N₂O emissions were reduced by 21.5 kt CO₂ eq (0.072 kt N₂O) from 32.8 kt CO₂ eq (0.11 kt N₂O NIR 2018) to 8.9 kt CO₂ eq (0.03 kt N₂O NIR 2019). This was due to double counting in both the LULUCF and agricultural sector of organic N fertilizers applied to forest lands by grazing animals. The source is now being reported only in the agriculture sector.

4(II) Emissions and removal from drainage of organic soils

- In forest land, CH₄ emission increased by 0.12 kt (3 kt CO₂ eq) from 2.11 kt (52.75 kt CO₂ eq) to 2.23 kt (55.75 kt CO₂ eq) while N₂O decreased by 0.11 kt (32.78 kt CO₂ eq) from 1.03 kt (306.94 kt CO₂ eq) to 0.92 kt (274.16 kt CO₂ eq) for year the 2016 between NIR 2018 and NIR 2019. This change was mostly due to the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively. For methane only, the emission factor for cropland converted to forestland, erroneously set to temperate zone (7.8625 kg of CH₄ ha⁻¹ yr⁻¹), was corrected to match that of the boreal nutrient rich zone (7.375 kg of CH₄ ha⁻¹ yr⁻¹). However the induced change was only visible with four decimals numbers. Finally, the modification of the area of drained organic soil also induced some slight recalculation not visible in the year 2016.
- In cropland, CH₄ emission decreased by 0.07 kt (1.75 kt CO₂ eq) from 3.65 kt (91.25 kt CO₂ eq) to 3.58 kt (89.5 kt CO₂ eq) for year 2016 between NIR 2018 and NIR 2019. This change was directly related to a 1.3 kha decrease of area.
- In grassland, CH₄ emission increased by more than twofold for all reported dates between NIR 2018 and NIR 2019 (e.g. from 0.07 to 0.15 kt CH₄ in 2016). The correction of an emission factor was responsible for these changes. This emission factor was calculated from the emission factors of ditch vs non-ditch area weighted by the respective proportion of these areas. Between NIR 2018 and NIR 2019, the emission factor for drained organic soil outside of the ditch was reevaluated from 36 kg to 39 kg of CH₄ ha⁻¹ yr⁻¹ while, the proportion of non-ditch area that was mistakenly set to 9.5% instead of 95% was corrected. As a consequence the resulting emission factor changed from 30.06 kg CH₄ ha⁻¹ yr⁻¹ to 60.55 kg CH₄ ha⁻¹ yr⁻¹.
- There were no recalculations for Wetlands peat extraction lands for 2016.

4(III) Direct N₂O emissions from N mineralization and immobilization

- The area changes caused by NFI updates caused only minor recalculations for this source in the year 2016 and throughout the timeseries.

4(IV) Indirect N₂O emissions from managed soils

- Recalculations for atmospheric deposition resulted in reduced N₂O emissions from 0.0022 kt N₂O (NIR 2018) to 0.0008 (NIR 2019). Recalculations for nitrogen leaching and run-off also resulted in reduced N₂O emissions from 0.0516 kt N₂O (NIR 2018) to 0.0343 kt N₂O (NIR 2019) for the year 2016. The reductions for these two sources was due to organic N fertilizer inputs from grazing animals being reported in the agriculture sector starting in the 2019 submission.

Biomass burning 4(V)

- There were no recalculations for this source in the year 2016.

10.2.5 Waste

5B1A composting facilities, Municipal solid waste

- Revised activity data. Activity data for the previous year, 2016, have been updated because the Norwegian waste accounts has not released updated figures in time for the emission inventory calculations. The correction has resulted in an increase of CH₄, N₂O and NH₃ emissions of 22 % compared to the previous calculation.

5B2A biogas facilities

- Revised activity data. Activity data was updated for 2016 and led to a reduction of CH₄ emissions by 22 %.

5C11a Municipal biogenic waste incineration

- The activitydata for the amount of landfill gas flared is updated. The change in data are due to new information from the energy statistics in Statistics Norway. This information has been used in connection with the amount reported from the landfills to the Norwegian Environment Agency and has resulted in higher activitydata then in the previous years.

5C12a Municipal waste incineration

- Reallocation. Combustion of waste without energy utilisation at one plant was previously included in 1A1a, but has now been reallocated to 5C1a. The relocation results in increased emissions in the periode 1995-2016

5D2 Industrial Wastewater

- Revised activity data. Activity data was updated in the period of 2000-2016. The new information resulted in a reduction of emissions of CH₄. The decline is marginal in the beginning of the periode and largest in 2010-2013 with a reduction of emissions around 10 %.

- The activity data used for estimating NMVOC from industrial wastewater was also updated for 2016, leading to an increase in the emissions for NMVOC of 31 %.

10.2.6 KP-LULUCF

Recalculation for the year 2016 for KP-LULUCF for each activity is shown in Table 10.4.

Table 10.4 Recalculated GHG emissions (kt CO₂-eq yr⁻¹) for 2016 per activity of KP-LULUCF.

KP activity	Emissions for 2016 (kt CO ₂ -eq yr ⁻¹)		
	NIR 2019	NIR 2018	Absolute difference
Afforestation	-505.87	-519.18	13.31
Deforestation	2 230.14	2 325.99	-95.85
Forest management	-27 719.40	-28 396.13	676.73
Cropland management	1 773.53	1 769.89	3.64
Grazing land management	10.20	0.37	9.83
Total	-24 211.40	-24 819.06	607.66

Total recalculations for the KP-LULUCF submissions for the year 2016³⁵ resulted in reduced C uptake of 608 kt CO₂-eq from -24 819 kt CO₂-eq (NIR 2018) to -24 211 kt CO₂-eq (NIR 2019) including non-CO₂ emissions. The majority of the change was due to the reduction in CO₂ uptake for FM (reduced by 677 kt CO₂-eq). The CO₂ uptake for AR decreased by 13 kt CO₂-eq, while emissions reduced by 96 kt CO₂-eq for D including non-CO emission. Emissions increased by 4 kt CO₂-eq for CM and by 10 kt CO₂-eq for GM, including non-CO₂ emissions. Recalculation for CH₄ emissions were minor and increased total KP-LULUCF CH₄ emissions by 3 kt CO₂-eq, from 158 kt CO₂-eq to 161 kt CO₂-eq. N₂O emissions were also recalculated and decreased by 42 kt CO₂-eq from 343 kt CO₂-eq to 304 kt CO₂-eq. Recalculations of CO₂ emissions were more important and these are described below for each KP activity.

4(KP-I) A.1 Afforestation and reforestation

- Total CO₂ sequestration for AR decreased by 13 kt CO₂ from 542 kt CO₂ to 529 kt CO₂, which was primarily due to the increased C uptake of 2 kt CO₂ in the aboveground living biomass (from 165 kt CO₂ to 167 kt CO₂). Belowground biomass sequestration decreased 4 kt CO₂ (from 49 kt CO₂ to 45 kt CO₂). C uptake in the litter pool decreased by 8 kt CO₂ from 433 kt CO₂ to 425 kt CO₂. C losses from the mineral soil pool were increased by 3 kt CO₂ from 86 kt CO₂ to 89 kt CO₂. All recalculations were due to the changes in the NFI area and living biomass data. NFI harvest data are used to distribute the known total harvests to the reported activities such as afforestation.

4(KP-I) A.2 Deforestation

- Total CO₂ emissions for deforestation were reduced by 95 kt CO₂ from 2 298 kt CO₂ to 2 203 kt CO₂. This was caused by the combination of reduced emissions from litter (17 kt CO₂ from 1 307 kt CO₂ to 1 290 kt CO₂), dead wood (1.4 kt CO₂ from 107 kt CO₂ to 106 kt CO₂), and below ground biomass (14.5 kt CO₂ from 148.7 kt CO₂ to 134.2 kt CO₂). In addition a slight

³⁵ Note that recalculations for KP-LULUCF, as described in the following, refer to absolute values of the activities, and not the value that is accounted for under KP2.

increase in the sequestration in mineral soils (0.7 kt CO₂; from 137.5 kt CO₂ to 138 kt CO₂), a smaller emission from aboveground biomass (58 kt CO₂ from 581 kt CO₂ to 523 kt CO₂) and a decrease in emissions from organic soils (4 kt CO₂ from 292 kt CO₂ to 288 kt CO₂).

Recalculations for all C pools were due to the updates in the NFI data.

4(KP-I) B.1 Forest management

- Carbon removals in the FM activity reduced by 714 kt CO₂. The majority of the reduction occurred in the living biomass; removals aboveground biomass was reduced by 631 kt CO₂ from 17 184 kt CO₂ to 16 553 kt CO₂ and belowground by 261 kt CO₂ from 4 108 kt CO₂ to 3 847 kt CO₂. The removals in the litter, dead wood, and mineral soil pools all increased due to a larger C uptake of 95 kt CO₂ (from 6 566 kt CO₂ to 6 661 kt CO₂), 19 kt CO₂ (from 1 326 kt CO₂ to 1 345 kt CO₂), and 2 kt CO₂ (from 151 to 153 kt CO₂), respectively. Emissions from organic soils decreased 60 kt CO₂ from 696 kt CO₂ to 636 kt CO₂. This change resulted from the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively.

4(KP) B.2 Cropland management

- Total CO₂ emissions for the CM activity increased 5 kt CO₂ (from 1 681 kt CO₂ to 1 686 kt CO₂). Removals in aboveground biomass increased by 0.6 kt CO₂ from 2.5 kt CO₂ to 3.1 kt CO₂ as did belowground biomass which increased 0.3 kt CO₂ from 1.0 kt CO₂ to 1.3 kt CO₂. The removals in the mineral soil pool decreased 40.7 kt CO₂ from 75.7 kt CO₂ to 35.0 kt CO₂. The emissions in the organic soil pool decreased 34 kt CO₂ from 1 760 kt CO₂ to 1 726 kt CO₂. Recalculation differences in mineral soils were due to corrections described under “4B1 Cropland remaining cropland” above and organic soils estimates were affected by the NFI extrapolation method.

4(KP) B.3 Grazing land management

- Recalculations for the GM activity resulted in increased emissions by 8 kt CO₂ from -6 kt CO₂ to 2 kt CO₂. This was primarily the result of the combined effect of decreased C uptake in above- and belowground living biomass (4.5 kt CO₂ for aboveground and 2 kt CO₂ for belowground) and slightly increased emissions from organic soils 0.0001 kt CO₂. Recalculation for living biomass was due to the updates in the NFI database. The change in the NFI database affected the area of organic soils under GM.

10.3 Implications for emissions levels and trends, including time-series consistency

Table 10.5 shows the effects of recalculations on the emission figures for CO₂, CH₄ and N₂O 1990-2016. Table 10.6 shows the effect of recalculations on the emission figures for HFCs, PFCs and SF₆ 1990-2016. Table 10.7 shows the effects of recalculations for the trends in emissions 1990-2016.

Table 10.5. Recalculations in 2019 submission to the UNFCCC compared to the 2018 submission. CO₂, CH₄ and N₂O. 1000 tonnes CO₂ equivalents.

	CO ₂			CH ₄			N ₂ O		
	Previous subm.	Latest subm.	Diff (%)	Previous subm.	Latest subm.	Diff (%)	Previous subm.	Latest subm.	Diff (%)
1990	35 704.4	35 323.0	-1.1	5 788.4	5 801.2	0.2	4 210.8	4 092.8	-2.8
1991	34 069.3	33 817.5	-0.7	5 754.9	5 770.7	0.3	4 053.0	3 937.6	-2.8
1992	34 831.9	34 737.0	-0.3	5 834.3	5 850.8	0.3	3 519.2	3 403.8	-3.3
1993	36 531.6	36 223.8	-0.8	5 930.6	5 940.2	0.2	3 706.8	3 606.2	-2.7
1994	38 446.0	38 087.3	-0.9	5 990.0	6 001.9	0.2	3 777.4	3 676.5	-2.7
1995	38 477.2	38 703.8	0.6	5 865.8	5 882.6	0.3	3 807.5	3 706.0	-2.7
1996	41 593.8	41 823.6	0.6	5 942.6	5 958.5	0.3	3 826.0	3 724.3	-2.7
1997	41 698.9	41 920.5	0.5	5 937.5	5 957.8	0.3	3 812.2	3 716.5	-2.5
1998	41 956.7	42 210.8	0.6	5 735.7	5 757.9	0.4	3 898.8	3 807.7	-2.3
1999	42 709.5	42 962.4	0.6	5 605.9	5 630.1	0.4	4 121.2	4 023.2	-2.4
2000	42 215.9	42 515.3	0.7	5 672.6	5 698.0	0.4	3 916.6	3 825.6	-2.3
2001	43 559.8	43 866.8	0.7	5 709.6	5 735.3	0.5	3 847.1	3 756.3	-2.4
2002	42 667.0	42 984.0	0.7	5 557.6	5 581.3	0.4	4 089.2	3 999.4	-2.2
2003	43 921.8	44 323.5	0.9	5 663.7	5 692.5	0.5	3 947.2	3 853.1	-2.4
2004	44 337.0	44 645.6	0.7	5 651.7	5 682.0	0.5	4 099.8	3 999.8	-2.4
2005	43 560.7	43 951.2	0.9	5 451.0	5 480.1	0.5	4 168.9	4 069.8	-2.4
2006	43 923.3	44 496.3	1.3	5 323.4	5 356.2	0.6	3 836.8	3 737.3	-2.6
2007	45 851.9	46 239.0	0.8	5 441.5	5 471.4	0.5	3 666.1	3 566.0	-2.7
2008	44 903.0	45 382.1	1.1	5 296.9	5 327.1	0.6	3 224.6	3 124.9	-3.1
2009	43 205.4	43 890.6	1.6	5 334.6	5 362.2	0.5	2 674.1	2 575.6	-3.7
2010	45 823.3	46 229.2	0.9	5 353.1	5 380.1	0.5	2 588.5	2 486.0	-4.0
2011	44 982.6	45 511.9	1.2	5 196.6	5 222.6	0.5	2 579.9	2 480.9	-3.8
2012	44 560.8	45 022.2	1.0	5 158.4	5 182.5	0.5	2 588.5	2 490.8	-3.8
2013	44 302.7	44 946.6	1.5	5 183.9	5 211.1	0.5	2 557.1	2 465.1	-3.6
2014	43 952.7	44 890.5	2.1	5 269.6	5 300.6	0.6	2 559.6	2 471.6	-3.4
2015	44 663.7	45 303.5	1.5	5 163.0	5 191.1	0.5	2 595.4	2 506.3	-3.4
2016	44 031.6	44 462.5	0.9	5 078.8	5 093.1	0.3	2 518.6	2 438.8	-3.2

National Inventory Report 2019 - Norway

Table 10.6. Recalculations in 2019 to the UNFCCC submission compared to the 2018 submission. HFCs, PFCs and SF₆. 1000 tonnes CO₂ equivalents.

	HFCs			PFCs			SF ₆		
	Previous subm.	Latest subm.	Diff (%)	Previous subm.	Latest subm.	Diff (%)	Previous subm.	Latest subm.	Diff (%)
1990	0.0	0.0	-	3 894.8	3 894.8	-	2 098.5	2 098.5	-
1991	9.9	9.9	-	3 456.7	3 456.7	-	1 983.5	1 983.5	-
1992	20.0	20.0	-	2 637.2	2 637.2	-	672.6	672.6	-
1993	31.6	31.6	-	2 648.3	2 648.3	-	703.8	703.8	-
1994	49.9	49.9	-	2 342.5	2 342.5	-	837.6	837.6	-
1995	92.0	92.0	-	2 314.0	2 314.0	-	579.8	579.8	-
1996	129.5	129.5	-	2 107.6	2 107.6	-	547.7	547.7	-
1997	191.5	191.5	-	1 882.7	1 882.7	-	553.2	553.2	-
1998	244.1	244.1	-	1 712.0	1 712.0	-	693.3	693.3	-
1999	316.0	316.0	-	1 600.0	1 600.0	-	833.7	833.7	-
2000	383.3	383.3	-	1 518.5	1 518.5	-	891.4	891.4	-
2001	473.3	473.3	-	1 531.3	1 531.3	-	754.8	754.8	-
2002	578.2	578.2	-	1 658.8	1 658.8	-	227.3	227.3	-
2003	557.6	557.6	-	1 051.1	1 051.1	-	216.5	216.5	-
2004	597.1	597.1	-	1 016.7	1 016.7	-	262.2	262.2	-
2005	614.3	614.3	-	955.3	955.3	-	296.1	296.1	-
2006	678.0	678.0	-	859.1	859.1	-	200.4	200.4	-
2007	715.3	715.3	-	951.2	951.2	-	70.5	70.5	-
2008	806.1	806.1	-	896.0	896.0	-	59.8	59.8	-
2009	856.1	856.1	-	438.3	438.3	-	55.7	55.7	-
2010	1 064.5	1 064.5	-	238.4	238.4	-	68.6	68.6	-
2011	1 105.8	1 105.8	-	262.6	262.6	-	54.3	54.3	-
2012	1 140.8	1 140.8	-	200.5	200.5	-	53.5	53.5	-
2013	1 155.2	1 155.2	-	181.0	181.0	-	56.3	56.3	-
2014	1 228.4	1 228.4	-	178.9	178.9	-	50.1	50.1	-
2015	1 225.7	1 225.7	-	146.4	146.4	-	69.8	69.8	-
2016	1 363.6	1 363.6	-	186.2	186.2	-	63.6	63.6	-

National Inventory Report 2019 - Norway

Table 10.7. Trends in emissions 1990-2016. 2019 submission compared to 2018 submission. GHG. % change 1990-2016.

	Total GHG	CO₂	CH₄	N₂O	HFCs	PFCs	SF₆
This submission	4.7	25.9	-12.2	-40.4	3106161.8	-95.2	-97.0
Previous submission	3.0	23.3	-12.3	-40.2	3106145.1	-95.2	-97.0

10.4 Implemented and planned improvements, including in response to the review process

The Norwegian Environment Agency co-ordinates the development and improvements of the inventory's different sectors. The recommendations from the review process are recorded in a spread sheet together with the needs recognized by the Norwegian inventory experts to form a yearly inventory improvement plan. Needs identified by use of the data for purposes other than reporting is also included. The overall aim of inventory improvement is to improve the accuracy and reduce uncertainties associated with the national inventory estimates. Each issue is assigned to a sector/theme and the overview tracks where the issue has originated from and the organization/person responsible for following up the recommendations. The overview is discussed among the agencies and each issue is given a priority and a deadline. Each organization in the inventory preparation therefore has responsibility for the development of the inventory. The issues are prioritized on the basis of the recommendations from the ERT and available human and financial resources.

The national greenhouse gas inventory has undergone substantial improvements over the recent years, and the inventory is considered to be largely complete and transparent. There was an in-country review of the inventory in 2018, but the latest annual inventory review report is ARR2016³⁶. Many of the recommendations in the ARR2016 have already been followed up and are reflected in the 2018 NIR. Implemented improvements since the 2018 NIR and how these are related to the review process are described in Table 10.8.

Table 10.8 Implemented improvements in response to the review process.

Sector/issue	ERT recommendation/ self-initiated	Source	Implementation	Internal Identification code
General				
Energy				
Energy, road transportation (1A3b)	Derive updated AD representative of annual consumption of LPG in road transportation in order to confirm that there is not an underestimation of emissions; alternatively demonstrate that the current approach of keeping AD flat does not lead to an underestimation of emissions in 2014.	ARR2016, E.26	Activity data were revised as part of the new energy balance.	EK.38
IPPU				

³⁶ <http://unfccc.int/resource/docs/2017/arr/nor.pdf>

Sector/issue	ERT recommendation/ self-initiated	Source	Implementation	Internal Identification code
IPPU- general	Update the existing documentation on time series consistency approaches applied for the IPPU sector, by category, to improve transparency on how Norway has ensured time series consistency	PMF 2018, finding I.20	Chapter 2 of Annex VIII describes a major QA/QC exercise in 2006 on consistent time series from 1990 to 2004 from the largest industrial plants in Norway. Chapter 3 of Annex VIII describes the current QA/QC procedures and the data sources used.	IK.74
IPPU- general	Improve consistency in the presentation of information on specific methods, AD, and EFs where emissions are aggregated from plant-specific reporting, in particular for key categories	PMF 2018, finding I.21	More information on methods, AD and EFs are included for the key categories. See for instance sections 4.2.1.3, 4.2.1.4, 4.2.2.3, 4.2.2.4, 4.3.2.3, 4.3.3.3, 4.3.5.4, 4.4.3.3, 4.4.3.4,	IK.75
2A3- Glass production	Report the AD for limestone consumption for glass production in the NIR table 4.5 (total balance of limestone)	PMF 2018, finding I.22	Table 4.5 in the NIR includes a row for 2A3 and it is clear that there was no consumption of limestone in 2014-2017	IK.76
2B1- Ammonia	Report the specific methods used to estimate CO2 recovered in the NIR	PMF 2018, finding I.23	Section 4.3.1.2 describes the method used to estimate CO2 recovered.	IK.77
2B5 - Carbide	Correct the AD for the reported time series in the 2019 submission	PMF 2018, finding I.24	QA/QC checks of the IEF identified irregularities in the IEF. A follow up with the facility indicated that reported production for some years was based on pure production rather than crude production. The AD has been corrected in the CRF and the time series for the IEF is improved.	IK.78
Agriculture				
Agriculture, 3A Enteric fermentation - dairy cow	The description of how the calculation of enteric CH4 emissions from dairy cows is made needs improvement to allow replication of the calculations.	PMF2018, A.6	The method description in NIR Annex IX, section 2.2.1 has been revised according to the ERT comments, in order to enhance the transparency.	JK.61

Sector/issue	ERT recommendation/ self-initiated	Source	Implementation	Internal Identification code
Agriculture, 3A Enteric fermentation - heifers and bulls for slaughter	In page 273, section 5.2.2 of the 2018 NIR an explanation related to the method for accounting for heifers and bulls for slaughter implies that the emissions will be calculated once over the entire lifetime of the animal. During the in-country review, Norway provided explanations for this practice that needs to be incorporated in the NIR.	PMF2018, A.9	NIR section 5.2.2 provides an explanation for the method used for estimating number of heifers and bulls for slaughter.	JK.64
Agriculture, General-Livestock characterisation	The 2018 NIR refers to a completeness level for livestock population without describing how this condition is dealt with. During the in-country review Norway provided clarification related to the matter. These explanations should be incorporated into the NIR to confirm that emissions cover 100% of the estimated population of livestock.	PMF2018, A.11	In NIR section 5.2.1, the explanation of the completeness level for the livestock population is revised and enhanced.	JK.65
Agriculture, General-Livestock characterisation-sheep	The sheep population requires a better description to make it more transparent. The ERT [recommends][encourages] that the methods to estimate the population of sheep that was presented during the in-country review is incorporated in the NIR or its annexes.	PMF2018, A.12	In NIR Annex IX, section 1.2, a detailed description of the livestock characterisation for sheep is given.	JK.66
Agriculture, 3B3 Swine	The ERT [recommends][encourages] that a better description of the "other swine" (does it include sow and boars?) population is needed in the CRF and not clear in the NIR where the consistency of the numbers needs to be	PMF2018, A.13	The description of the different swine categories has been updated, and NIR Annex IX Table AIX-1 gives consistent data with the CRF.	JK.67

Sector/issue	ERT recommendation/ self-initiated	Source	Implementation	Internal Identification code
	checked and adjusted to align with the CRF.			
LULUCF³⁷				
Waste				
Waste, Wastewater treatment and discharge, (5D)	The ERT [recommends][encourages] Norway to clarify in the next submission that these values represent the proportions of population using the WWTP and not the MCF values	PMF 2018, W15	Norway has updated the methodology description in the NIR .	AK.47
KP-LULUCF³⁸				

Improving the inventory and the reporting is a continuous process and there are still some issues previously identified that need to be addressed. Table 10.9 gives an overview of the planned improvements.

³⁷ Self-initiated improvements, not related to ARR findings, are described in the recalculations for LULUCF, chapter 10.2.4.

³⁸ Self-initiated improvements, not related to ARR findings, are described in the recalculations for KP-LULUCF, chapter 10.2.6.

Table 10.9 Plan for improvements for the Norwegian GHG inventory.

Sector/ issue	ERT recommendation/ self-initiated	Source	Plan for improvement	Internal Identifi- cation code
General				
General, Uncertainty analysis	The ERT [recommends][encourages] that Norway updates and improves its uncertainty analysis through a comprehensive revision and update of the uncertainty parameters applied in the base year and for the methods now used in the Norway inventory in the latest year, and to report on progress in the next submission.	PMF2018, G.11	The reporting in the NIR 2020 will contain updated uncertainties for activity data and emission factors for the most recent year (2018). Furthermore, the consequences of using the same set of uncertainties for base year (1990) and the most recent year will be evaluated. A description of how we want to handle this issue will be included in the NIR.	GK.48
Energy				
Energy, reference approach (1A(b))	The ERT recommends that Norway report on the time frame and progress of the revised energy balance system in the 2017 submission, highlighting the resulting reduction in statistical differences for solid fuels	ARR2014, §26 ARR2016, E.4/E.17	Statistics Norway revised the energy balance in 2017-2018 and the results were incorporated into the emission inventory in 2018. The results are reflected in the current NIR and CRF reporting. However, further work is required in order to resolve the issues, particularly with respect to statistical differences and RA/SA discrepancies. This project is described in section 10.4.1	EK.63

National Inventory Report 2019 - Norway

Sector/ issue	ERT recommendation/ self-initiated	Source	Plan for improvement	Internal Identifi- cation code
Energy, reference approach (1A(b))	The ERT recommends that the Party transparently describe the technical solution that aims to improve the link between the energy balance and IEA reporting, including providing any preliminary results in the 2017 submission, and then improve the alignment of the energy balance and IEA reporting for the 2018 submission	ARR2014, §26 ARR2016, E.5/E.18	Statistics Norway revised the energy balance in 2017-2018 and the results were incorporated into the emission inventory in 2018. The results are reflected in the current NIR and CRF reporting. However, further work is required in order to resolve the issues, particularly with respect to statistical differences and RA/SA discrepancies. This project is described in section 10.4.1	EK.31a
Energy, feedstocks, reductants and non-energy use of fuels (1A(d) etc)	E.7: Improve QC procedures to ensure consistency of the information reported on feedstocks, reductants and NEU in different CRF tables E.8: Review and revise the reporting in CRF table 1.A(d) and improve QC procedures to ensure consistency of the reporting E.20: The ERT recommends that Norway report on the time frame and progress of the revised energy balance system including any improvements in the consistency of the information on feedstocks, reductants and non-energy use of fuels reported in the CRF tables	ARR2014, §29 and previous ARRs ARR2016, E.7/E.20, E.8	Statistics Norway revised the energy balance in 2017-2018 and the results were incorporated into the emission inventory in 2018. The results are reflected in the current NIR and CRF reporting However, further work is required in order to resolve the issues, particularly with respect to statistical differences and RA/SA discrepancies. This project is described in section 10.4.1	EK.8
IPPU				
Agriculture				
LULUCF				
Waste				
Waste, Wastewater treatment and discharge, Industrial wastewater (5D2)	Investigate possible overestimation of emissions. In addition, the ERT recommends Norway to apply the IPCC 2006 Guidelines to estimate CH ₄ emissions from industrial wastewater (EQUATION 6.4) considering that the amount of CH ₄	ARR 2016, W9	Norway has included this recommendation in its improvement plan	AK.39

Sector/ issue	ERT recommendation/ self-initiated	Source	Plan for improvement	Internal Identifi- cation code
	that is flared or recovered for energy use should be subtracted from total emissions. The ERT also noted that emissions from CH ₄ recovery for energy generation should be reported in the Energy Sector taking into account the avoidance of double counting emissions from flaring and energy used			
Waste, Wastewater treatment and discharge, (5D)	The ERT noted that in CRF table 5.D the notation key "NE" is used for total organic product (kt DC/year) for domestic, industrial and other wastewater. Nevertheless in the NIR it is stated that biochemical oxygen demand and COD data are available and used to estimate emissions. The ERT recommends that Norway present total organic product data in the NIR and in CRF table 5.D	ARR 2016, W8	Organic product data will be included in the next submission	AK.44
KP-LULUCF				

10.4.1 Work plan for further improvement of the Reference and Sectoral Approach

The Norwegian Emission Inventory was subjected to an in-country review in 2018. The team of experts expressed concerns about the size of the differences in energy use and emissions as estimated by the reference and the sectoral approach. This section provide information and plans on how the Norwegian inventory team will be working with this issue.

Firt of all, the goal of the plans made in response to the concerns is:

"For the reporting in 2021, the differences in energy use and emissions between the Reference Approach (RA) and the Sectoral Approach (SA) are reduced to an acceptable level. This level of acceptance is well documented and reasoned for in the NIR of 2021."

The work plan to address the difference between the Reference approach and the Sectoral approach has been adopted. The following overall roles and responsibilities have been agreed for this work:

- **The steering group.** A steering group consisting of the Directors of the departments in both NEA and SN has been established for the follow up of this plan. The group has the overall responsibility for meeting the requirements of the plan, and for sufficient resource allocation

and financing. The head of sections in both NEA and SN will be responsible for planning of meetings and the reporting to the steering group.

- **Statistics Norway (SN)** holds the responsibility for calculating emissions for both the reference and sectoral approach in the Norwegian emission inventory. SN will be responsible for the practical implementation of the work plan.
- **Norwegian Environment Agency (NEA)** is the appointed national entity responsible for reporting the greenhouse gas emissions for Norway, and for coordination of the three institutions in the Norwegian National System. Inventory experts at NEA will be contributing as discussion partners where needed, especially on the issue of defining the accepted level of difference between the reference and sectoral approach. NEA also holds the responsibility of arranging the biannual National System Meetings, where progress towards reducing the difference between RA and SA will be discussed and reported as described below.

The work is organized in three main groups with the responsibility of work programs as described in Table 10.10. Reporting of work progress, updating of plans and evaluation of resources will be addressed four times a year according to Table 10.11. A summary of the progress of the work and updated plans will be provided annually in the Norwegian National Inventory Report.

Table 10.10 Organisation of the project in three groups with related tasks

Task group name	Who	Short description of tasks/milestones
The RASA-framework and reporting	The Emission Inventory team at SN	<ul style="list-style-type: none"> • The appointed RASA-responsible(s) in the emission inventory team has a clear understanding of the reporting requirements as described in 2006 IPCC Guidelines • A new system/routine for reporting RASA in CRF and NIR is established • Updated tables and analysis in the CRF and NIR reported • All differences in RASA that are not due to statistical differences are removed or explained
The Energy Balance and statistical differences	The Energy Balance team at SN	<ul style="list-style-type: none"> • A clear documentation (illustrations) of the data and work flows of the Energy Balance is provided in NIR • Statistical differences are reduced to an accepted level, as defined by project group working on this issue ("Level of accepted differences") <ul style="list-style-type: none"> ○ Evaluate inconsistencies between production and export data on refined petroleum products. Gather updated data, or perform correction ○ Evaluate the coverage of the sales statistics and the import data in

		<p>the foreign trade statistics. Gather updated data, or perform correction</p> <ul style="list-style-type: none"> ○ Evaluate the need for quality control of other energy products, like gas and coal ○ Perform a new consolidation of the energy balance, if the problems with statistical differences are still not solved (due November 2020). Main focus on 2010 to 2018 <ul style="list-style-type: none"> • The Energy Balance data and System have gone through a thorough quality check • Regular contact with important data providers is established to ensure good quality of data
Level of accepted differences	Both teams at SN and the inventory team at NEA	<ul style="list-style-type: none"> • The level of accepted differences in energy use and emissions is evaluated and documented in NIR

Table 10.11 Reporting of plans and progress

Reporting "format"	When	Participants	Suggested agenda	Documents
National system meeting	April	Head of sections in NEA and SN, relevant members of inventory teams/leaders of the three project groups	Status of work and presentation of plans for this year	
Written report by e-mail	June	From leaders of the three project groups to Head of sections in NEA and SN	Status of work, evaluation of progress and resources, updated plans	
National system meeting	September	Head of sections in NEA and SN, relevant members of inventory teams/leaders of the three project groups	Status of work, evaluation of progress and resources, updated plans	
Steering group meeting	December	Directors of the departments in NEA and SN, Head of sections in NEA and SN, relevant members of inventory teams/leaders of the three project groups	Summary of work, evaluation of progress and resources, presentation of plans for next year	Draft text to chapter 10.4 for NIR (summary of progress and plans)

The following list provides status for the activities mentioned in the response to Saturday paper:

- Publish an updated energy balance. Brief consolidating performed. *Progress: Revised energy balance was published in November 2018.*
- A clear documentation (illustrations) of the data and work flows of the Energy Balance is provided in NIR. *Progress: Flowchart of the data and work flow in the Energy Balance is included in the Energy chapter, 3.2.1.2.*
- Assuring that all data from the old energy balance system that were meant to be transferred to the new system, are correctly transferred. Thorough consolidating. *Progress: This work has started and continuous spring 2019. A few errors have been found, which will be corrected when publishing the energy balance in June 2019. The RA-SA project in 2013 – 2015 led to several findings, mostly new data sources. These data sources are now incorporated in the new energy balance. The work ensuring that all the findings are included in the energy balance is soon completed. No deficiencies have been found so far.*
- Identify and propose criteria. *Progress: Not started.*
- Evaluate inconsistencies between the production and export data on refined petroleum products. Gather updated data or perform correction. *Progress: In November 2018, SN had a meeting with the Norwegian Tax administration and after the meeting, SN received a file with export data. A comparison with export data from foreign trade statistics revealed a difference of 20 % in export of motor gasoline. This deviation was discussed in a meeting between the foreign trade statistics and the energy balance team in March 2019. A further examination of this deviation will be undertaken in 2019. Similar comparisons will be performed for the other refined petroleum products. Import and export of fuels used in navigation was also discussed, revealing that fuels used by the ship itself is not customs declared. If the ships fill fuel abroad, this energy consumption will not be included in the energy balance.*
- Evaluate the coverage of the sales statistics and the import data in the foreign trade statistics. Gather updated data or perform correction. *Progress: The coverage of the sales statistics has been evaluated and the population has been controlled. The evaluation revealed a lack of sales data form biodiesel in 2017. This will be included when publishing the energy balance in June 2019.*

10.4.2 Work plan for further improvement of the Institutional Arrangements

During the in-country review of the Norwegian Emission Inventory in 2018, the team of experts expressed special concerns about the change of staff (as described in chapter 13) and potential consequences on the quality of the emission inventory. The following text and tables describe how SN and NEA have been working, and will be working during the next cycle, to ensure that the quality of the emission inventory is not compromised.

Since the review in September 2018 both the Energy balance team and the Emission inventory team has kept up with the knowledge and transfer plans established in SN in relation to the reorganizing of staff. During the fall and winter, there has been regular contact between the inventory team in SN and NEA, to discuss any unresolved issues in the inventory. A revised energy balance was published in November 2018 in a cooperation between the new Energy balance team and an experienced expert. Final emission figures from the period 1990 - 2017 were published in December 2018,

basically with the new inventory team. Work related to the CRF-tables and documentation in the NIR has been performed by the new inventory team, with guidance from experienced inventory experts.

In addition, a plan for capacity building and increased contact between the more experienced staff in NEA and the new staff in SN has been set up in order to counteract the risks of loss of institutional memory. This plan is summarized in Table 10.12 below.

Table 10.12 Plan for capacity building in 2018-2019

	Activity	Status by April 2019
November 2018	<ul style="list-style-type: none"> • Participation of relevant SN-staff in a seminar about emission inventory for the Oil and Gas Industry, arranged by NEA for developing countries. 	Completed
January 2019	<ul style="list-style-type: none"> • Workshop on RASA. Arranged by SN to make further plans for the project that will run in 2019 and 2020 • Workshop on Key Source Analysis and Uncertainty Analysis arranged by NEA. Capacity building 	Completed
March 2019	<ul style="list-style-type: none"> • Workshop on SF₆ arranged by NEA. Participants from both SN and the most important data provider ("Brukergruppa for SF₆"). Ensuring correct understanding of the data needs and emission model 	Completed
April 2019	<ul style="list-style-type: none"> • Nordic Inventory Meeting in Helsinki: Four persons from the new inventory team will attend the meeting in Helsinki to learn, discuss and get to know other inventory experts • Workshop on tools for collaboration. Capacity building and improved use/routines: <ul style="list-style-type: none"> ○ Sharepoint ○ Jira ○ Skype 	
May 2019	<ul style="list-style-type: none"> • HBFA meeting in Zurich: The inventory compiler responsible for road traffic will attend the meeting, travelling with representative from NEA • CEIP LRTAP meeting in Thessaloniki 13-15. May: Three persons from the inventory team will attend the meeting, traveling with representative from NEA 	
June 2019	<ul style="list-style-type: none"> • Workshop on indirect emissions arranged by NEA 	
September 2019	<ul style="list-style-type: none"> • Two-days seminar for all members of the National System: 	

National Inventory Report 2019 - Norway

	<ul style="list-style-type: none"> ○ Roles and responsibilities of the member institutions. The agreements and obligations ○ Other relevant issues: To be decided ○ Teambuilding. Sosializing and getting to know each other ● GHG management institute courses: Several of the new inventory compilers are planning to attend courses under the direction of GHG management institute during summer/fall. 	
November 2019	<ul style="list-style-type: none"> ● Data flow of reported emissions from NEA in SN 	

Part II: Supplementary information required under article 7, paragraph 1

11 KP-LULUCF

11.1 General information

Norway provides supplementary information under Article 7 of the Kyoto Protocol (KP) for the Land Use, Land-Use Change and Forestry sector. The information provided in this chapter is in accordance with relevant CMP decisions such as Decision 16/CMP.1, 2/CMP.7, 2/CMP.8, 6/CMP.9 and the 2006 IPCC Guidelines, the 2013 IPCC KP Supplement (IPCC, 2014a), and the IPCC 2013 Wetlands Supplement (IPCC, 2014b).

In the first commitment period of the Kyoto Protocol (2008-2012) Norway reported on emissions and removals from the obligatory activities Deforestation (D) and Afforestation/reforestation (AR) under Article 3.3, in accordance with Paragraph 6 of the Annex to Decision 16/CMP.1. In addition, Norway decided to elect the voluntary activity Forest Management (FM) under Article 3.4 for inclusion in its accounting.

For the second commitment period (2013-2020) Norway reports, in accordance with paragraph 7 of decision 2/CMP.7, Annex I, emissions and removals from Article 3.3 activities and from Forest Management under Article 3.4. In addition, Norway has elected the voluntary activities Cropland Management (CM) and Grazing Land Management (GM) in its accounting under Article 3.4 and reports emissions and removals from all sources and sinks under these KP Article 3.4 activities. In the second commitment period Norway has chosen commitment-period accounting.

This chapter covers information on emissions and removals from activities under Article 3.3 and 3.4 of the Kyoto Protocol for the years 2013, 2014, 2015, 2016, and 2017, and is in accordance with Annex II of decision 2/CMP.8. Reported emissions and removals from areas under the KP activities includes the following sources and sinks: carbon stock changes in aboveground biomass, belowground biomass, litter, dead wood, mineral soils and organic soils, direct N₂O emissions from N fertilization (for AR, D, and FM), emissions and removals from drained and rewetted organic soils, N₂O mineralization in mineral soils, indirect N₂O emissions from managed soils, and N₂O and CH₄ emissions from biomass burning.

Areas where afforestation and reforestation and deforestation activities have occurred in Norway are small compared to the area of forest management. Estimated C sequestration for the activity FM is substantial, and there is also a C uptake as a result of AR. The activities deforestation and cropland management are sources of net emissions for all reported years. Grazing land management had net removals in the base year 1990, but has small net emissions in the reported years 2013-2017. Table 11.1 shows the emissions and removals for each KP activity for the base year 1990 (where relevant), and for each year of the second commitment period. During the second commitment period (from 2013 to 2017) emissions from CM were relatively stable, lower than that of the base year (1990) and remained within the range of strong variation observed during the first commitment period. This last point advocates for an absence of trend for the whole reporting period for CM emissions. Compared to the base year which had removals, emissions from GM were observed for all years of the second commitment period. This increase in emission is caused mostly by emissions from the mineral soil pool. The majority of the emissions from D occur in the litter pool.

Table 11.1 CO₂, N₂O, and CH₄ emissions (kt CO₂ equivalents yr⁻¹) and CO₂ removals of all pools for Article 3.3 and 3.4 under the Kyoto Protocol for the base year (1990) and for each year of the second commitment period.

Year	Net emissions/removals (kt CO ₂ -equivalents yr ⁻¹)				
	Afforestation/ reforestation	Deforestation	Forest management	Cropland management	Grazing land management
1990			-12482.45	1786.98	-73.19
2013	-569.07	2279.38	-29118.20	1769.69	16.15
2014	-561.63	2158.26	-27923.20	1776.96	11.64
2015	-532.84	2157.39	-27035.01	1776.20	10.04
2016	-505.87	2230.14	-27719.40	1773.53	10.20
2017	-489.06	2218.46	-28914.54	1763.53	12.04

Source: Norwegian Institute of Bioeconomy Research

11.1.1 Relation between UNFCCC land classes and KP activities

The land classification under the convention can be directly translated into activities under the KP with two exceptions. First, land-use changes reported under the convention includes human-induced and non-human induced land-use change, whereas only human-induced land-use changes are reported under KP. Second, the 20-year transition period for land-use changes is not applied under KP, which means that land cannot leave a land-use change category. However, we do apply appropriate methods to estimate the emissions or removals from land that has been in a conversion category in the reporting to the UNFCCC for more than 20 years.

The correspondence between the national land cover and land-use categories (Table 6.7) and the KP activities is illustrated by a translation matrix (Table 11.2). Briefly, land classified as the activity D is the sum of forest land converted to cropland, grassland, wetlands, settlements, and other land (direct human-induced land-use change). Analogously, land classified for the activity AR is the sum of cropland, grassland, wetlands, settlements, and other land converted to forest land, but only where the conversions are directly human-induced (Table 11.2). Once land is classified as D, it stays in D even if subsequent afforestation takes place. Land classified as the activity FM is forest land that has remained forest land since 1990 and land conversions to or from forest that are not caused by human activity. Cropland management entails the activities on land that has remained cropland since 1990 and non-forest related land conversion to or from cropland since 1990. Land classified as grazing land management is land that has remained grassland since 1990 and land-use conversion to or from grassland, with the exception of those related to forest land or cropland.

Table 11.2 Land-use change matrix with classification of the KP activities and the corresponding land-use classes. The following notations are used for classification of land-use changes. AR: Article 3.3 Afforestation/Reforestation, D: Article 3.3 Deforestation, FM: Article 3.4 Forest management, CM: Article 3.4 Cropland management, GM: Article 3.4 Grazing land management, and O: Other activities. In the case of non-human induced land-use transition, the activity in brackets () is assigned.

		Reporting year					
Base year	Land-use	Forest land	Cropland	Grassland	Wetlands	Settlements	Other land
1990	Forest land	FM	D	D	D (FM)	D	FM
	Cropland	AR	CM	CM	CM	CM	CM
	Grassland	AR	CM	GM	GM	GM	GM
	Wetlands	AR (FM)	CM	GM	O	O	O
	Settlements	AR	CM	GM	O	O	O
	Other land	AR (FM)	CM	GM	O	O	O

Specifically, the annual change in the area of D is not exactly equal to the annual change in the area of FM (Table 11.3), because only human-induced land-use changes are reported under the KP. Also, areas of AR and D do not exactly equal the areas of lands converted to forest land (LF) and forest land converted to lands (FL), respectively, under the Convention reporting. The difference between the sum of AR and FM and the sum of LF and forest land remaining forest land under the Convention is equal to the non-human induced changes from other land to forest land.

Furthermore, since 2011, an additional reason for the lack of correspondence between AR and LF, and between D and FL, is the application of the 20-year transition period in the Convention reporting, where areas are classified in transition (as land in conversion) for 20 years before they enter a remaining land-use category. This means that the area of land converted to forest land in 1990, 1991, and 1992 under the Convention will enter the forest land remaining forest land category in 2011, 2012, and 2013, respectively. However, for KP-LULUCF reporting, the areas reported for the activities AR and D remain AR and D for the whole reporting period and are thus not reported as a FM activity after 20 years. A full time-series of the areas considered for the activities AR, D, FM, CM, and GM from 1989 to 2017 is presented in Table 11.3.

National Inventory Report 2019 - Norway

Table 11.3 Time-series of area estimates (kha) for the activities Afforestation/Reforestation (AR), Deforestation (D), Forest management (FM), Cropland management (CM), and Grazing land management (GM) for 1989-2017.

	Area (kha)				
Year	Afforestation/ Reforestation (AR)	Deforestation (D)	Forest management (FM)	Cropland management (CM)	Grazing land management (GM)
1989	0	0	12200.34	938.11	230.86
1990	1.94	4.16	12196.44	937.83	230.23
1991	3.88	8.33	12192.53	937.55	229.59
1992	5.82	12.49	12188.63	937.27	228.96
1993	7.77	16.66	12184.72	936.98	228.32
1994	9.71	20.82	12180.81	936.70	227.69
1995	11.76	25.85	12176.04	936.61	226.69
1996	13.79	31.07	12171.09	936.39	225.85
1997	15.96	37.1	12165.4	936.17	225.05
1998	18.3	43.04	12159.91	936.08	223.93
1999	20.57	49.21	12154.38	936.07	222.70
2000	23.02	55.25	12148.79	935.69	221.84
2001	25.49	61.43	12143.31	935.65	220.88
2002	28.01	67.11	12138.25	935.18	220.14
2003	29.87	72.8	12132.98	934.72	219.94
2004	31.89	78.43	12127.61	934.12	219.78
2005	34.32	84.75	12121.54	933.75	219.26
2006	36.84	91.31	12115.16	932.86	218.82
2007	39.33	98.42	12108.96	932.32	218.32
2008	42.68	106.06	12102.45	931.96	217.64
2009	45.98	114.06	12095.87	931.87	216.75
2010	48.32	121.23	12090.65	932.36	216.57
2011	50.29	127.67	12088.6	933.58	216.22
2012	52.32	133.35	12087.39	934.41	215.77
2013	53.69	138.49	12086.89	934.94	215.42
2014	55.08	143.45	12086.43	935.35	215.17
2015	56.95	148.82	12085.28	935.81	214.86
2016	59.01	154.57	12082.55	935.96	215.00
2017	61.04	160.69	12079.4	936.65	215.36

11.1.2 Definitions of elected activities under Article 3.4

Forest management is defined according to forest lands described in chapter 6.2.1. The values used in the National Forest Inventory are in accordance with the range of parameters in the definition from the Global Forest Resources Assessment (FRA) 2005 and IPCC 2003 (Table 11.4). Forest land is land with tree-crown cover of more than 10 %. The trees have to be able to reach a minimum height of 5 m at maturity in situ. Minimum area and width for forest land considered in the Norwegian inventory is 0.1 ha and 4 m, respectively, which is a discrepancy from the definition in FRA 2005 (0.5 ha and 20 m). Furthermore, forest roads are considered as settlements. Young natural stands and all plantations established for forestry purposes, as well as forests that are temporarily unstocked, e.g. as a result of harvest or natural disturbances, are included under forest management.

Table 11.4 Parameters for the definition of forest land in IPCC 2003, the Global Forest Resources Assessment (FRA) 2005, and in the National Forest Inventory (NFI).

Parameters	IPCC 2003	FRA 2005	Values used (NFI)
Minimum land area	0.05 – 1 ha	0.5 ha	0.1 ha
Minimum crown cover	10 – 30%	>10%	>10%
Minimum height	2 – 5 m	5 m	5 m
Minimum width		20 m	4 m

Cropland management is defined as the activities that occur on cropland and cropland is defined as described in chapter 6.2.1. Croplands are areas that are annually cropped and regularly cultivated and plowed. Both annual and perennial crops are grown. It also encompasses grass leys that are in rotations with annual crops, which may include temporarily grazed fields that are regularly cultivated. This category also includes arable land that has been annually cropped and regularly plowed, but has since then been abandoned. These areas remain in the cropland management category until they have a regrowth of trees that make them unsuitable for plowing. In addition, to the areas classified as cropland remaining cropland, cropland management also includes all non-forest conversion to or from cropland.

Grazing land management is defined as the activities that occur on grassland, which is described in chapter 6.2.1. Grasslands are areas utilized for grazing on an annual basis. More than 50 % of the area should be covered with grass and it can be partly covered with trees, bushes, stumps, rocks etc. The grass may be mechanically harvested but the soil cannot be plowed. Land with tree cover may be classified as grassland if grazing is considered more important than forestry even if the forest definition is met. In addition to the areas classified as grassland remaining grassland, grazing land management also includes all non-forest and non-cropland conversion to or from grassland.

11.1.3 Description of how the definitions of each activity under Article 3.3 and 3.4 have been applied consistently over time

The Norwegian National Forest Inventory (NFI) provides data on land use, land-use change and forestry for the greenhouse gas reporting related to Article 3.3 and Article 3.4. A detailed description of the NFI can be found in chapter 6, section 6.3.

Estimates of areas subject to Afforestation/Reforestation (AR), Deforestation (D), Forest Management (FM), Cropland Management (CM) and Grazing land Management (GM) are based on the NFI, which has been carried out continuously since 1986. Land use obtained between 1986 and 1993 serves as the baseline for the area and living biomass estimates on 31 December 1989. Because no data from permanent sample plots exists before 1986 and relatively small changes have been detected for forest land as well as the other land-use classes, we have chosen not to take into account changes that may have occurred prior to 1990.

All forests in Norway are considered managed and this includes recreational areas, protected areas, and nature reserves. All forests in Norway are used either for wood harvesting, protecting and protective purposes, recreation, and/or to a greater or smaller extent, for hunting and picking berries, and are therefore subject to the FM activity.

11.1.4 Hierarchy among Article 3.4 activities and how they have been consistently applied in determining how land was classified

As Norway has included FM, CM, and GM under Article 3.4 of the Kyoto Protocol in the accounting for the second commitment period, it is necessary to determine the hierarchy among Article 3.4 activities. Forest management takes precedence over both cropland and grazing land management. Norway has further decided that cropland management takes precedence over grazing land management, because it covers a larger area and it is more important in terms of emissions per area. Thus, the hierarchy is as follows: forest management > cropland management > grazing land management. In practice, this means that grassland converted to cropland will change activity from grazing land to cropland management, but cropland converted to grassland will remain as cropland management activity. Article 3.3 activities (AR and D) always take precedence over Article 3.4 activities.

11.2 Land-related information

11.2.1 Spatial assessment units used for determining the area of the units of land under Article 3.3

The activity data used for determining the area of the units of land under Article 3.3 are the 250 m² large NFI sample plots (see detailed description in chapter 6.3). A land conversion will be recorded as soon as 20 % or more of the plot area is converted to another land use class. Sample plots are split between two land use classes if one of the land use classes covers at least 20 % of the plot area. Since 1986, all plots are classified according to a national land cover and land-use classification system, which is consistently translated to KP activities.

The NFI database provides activity data for the entire country. However, there is no time-series of field observations in Finnmark County and the mountain forest stratum before 2005. For plots in Finnmark County and the mountain forest stratum, information from maps, registers, and old and new aerial photographs were used to determine the land use of each plot in the base year 1990. The models used to back-cast the living biomass on these sample plots were based on the methods described in the LULUCF chapter (chapter 6). All land-use changes, except for one, were observed in the lowland forest stratum outside Finnmark.

11.2.2 Methodology used to develop the land transition matrix

The land-use transition matrix (Table 11.2) is based upon changes in the land-use category of the sample plots surveyed in a given year. Changes in land use are recorded for the year the land use is observed. A full NFI cycle, i.e. plots observed over a 5-year period, are used for estimating areas of land-use categories. Extrapolation is used in the last 4 years of the reporting period (see section 6.3.4).

11.2.3 Maps and/or database to identify the geographical locations and the system of identification codes for the geographical locations

All the NFI plots are geo-referenced and each plot has a unique identification code. According to the IPCC good practice guidance, the coordinates of these plots are classified information. However, a list of sample plots can be provided to the expert review team upon request. The approximate spatial distribution of the areas subject to the activities under Article 3.3 and to the activity FM under Article 3.4 is given in Figure 11.1. Figure 11.2 displays the approximate location of the activities FM, CM, and GM under Article 3.4.




Figure 11.1 Spatial distribution (approximate location of sample plots) of Article 3.3 activities afforestation and deforestation, and the Article 3.4 activity forest management from 1990 to 2017. Symbol sizes for plots with afforestation and deforestation activities are increased to improve the visibility of these activities.




Figure 11.2 Spatial distribution (approximate location of sample plots) of elected Article 3.4 activities for 2017 in Norway. Symbol sizes for plots with cropland or grazing land management activities are increased to improve the visibility of these activities.

11.3 Activity specific information

11.3.1 Methods for carbon stock change and GHG emission and removal estimates

Methods and activity data used to calculate the emissions reported under KP-LULUCF are in general identical to those applied in the reporting under the Convention (chapter 6). They are in accordance with the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006) and we refer to chapter 6 for detailed descriptions. In this chapter we provide information about methods specific for reporting under KP. All relevant methods are in accordance with the 2013 IPCC KP supplement (IPCC, 2014a) and the IPCC 2013 Wetlands Supplement (IPCC, 2014b).

11.3.1.1 Differences in the methodologies used for the KP and the Convention reporting

For AR and D, the methods used to estimate carbon stock changes were identical to those used for the corresponding land-use change. However, there was one difference in the carbon stock change rate for dead wood used for other land converted to forest land. The rate $0.025 \text{ t C ha}^{-1} \text{ yr}^{-1}$ was used in KP to better reflect a human-induced conversion, while $0.013 \text{ t C ha}^{-1} \text{ yr}^{-1}$ was used in the Convention reporting. Carbon stock changes (CSC) in living biomass must be divided between above- and belowground for all KP activities. For cropland management, the Tier 1 method for living biomass does not provide this division. We assumed that 30 % of the loss or gains occur belowground and 70 % aboveground. No other methodological differences exist for CSC estimation in any pools between the Convention and the KP reporting. Further, for ARD areas after a period of 20 years (with transition from one assumed equilibrium to another) the CSC in soil, deadwood, and litter are assigned emissions reflecting their current land use, while their area is still included in the appropriate ARD categories.

To estimate direct and indirect N_2O emissions under FM and AR, respectively, we used a multiplication factor based on the percentage of the area under AR or FM of the total forested land (AR + FM area). The multiplication factor is calculated annually. The same approach was applied for biomass burning.

Methods used to estimate N_2O from N mineralization-immobilization due to soil C loss and emissions and removal from drained and rewetted organic soils were also identical to those used in the reporting under the Convention.

11.3.2 Uncertainty estimates

Sampling errors for proportions (areas) and totals (carbon stock change) are estimated according to standard sampling methodology based on the 5 most recent years of NFI data (see section 6.3.4). The sample plots in the NFI are systematically distributed. Since we have assumed random sampling, the variances are conservative estimates. Uncertainties in terms of standard errors related to the estimates of area are shown in Table 11.5. Uncertainties in terms of standard errors related to the estimates of net C stock changes are shown in Table 11.6.

Table 11.5 Uncertainty of annual area estimates.

Activity	Area 2SE (%)
Afforestation/Reforestation	23
Deforestation	14
Forest management	2
Cropland management	7
Grazing land management	13

Uncertainties in C stock changes are dependent on area uncertainties and the variability in the C stock changes per unit area. Uncertainties for the C stock change estimates in living biomass are based on standard sampling methodology for the estimates of totals, except for CM where default uncertainties are given and for GM where it is based on expert judgement. Uncertainties for the C stock change estimates per hectare in the dead wood, litter, and soil pools were based on expert judgment, except for FM. Uncertainties in area estimates and per hectare estimates were combined to arrive at the final estimates presented in Table 11.6. For FM, the estimates for dead wood, litter, and the soil pool were estimated using Yasso07-model and a Monte-Carlo method was applied to determine the associated uncertainty (section 6.4.1.2). Assumptions behind the expert judgments used for AR and D are described in chapter 6, see section 6.4.2.1.

Table 11.6 Uncertainties of annual total C stock changes.

Activity	AG and BG living biomass 2SE (%)	Dead wood + litter 2SE (%)	Mineral soils 2SE (%)	Organic soils 2SE (%)
Afforestation/Reforestation	53	100 – 200	50 – 100	50
Deforestation	49	100 - 182	50 – 100	19
Forest management	13	15	15	50
Cropland management	75*	NO	50	19
Grazing land management	101**	NO	91	50

* Uncertainties for living biomass in cropland management (fruit orchards) are based on the default method. ** Expert judgement.

11.3.3 Changes in data and methods since the previous submission (recalculations)

Recalculation for the year 2016 for KP-LULUCF for each activity is shown in Table 11.7.

Table 11.7 Recalculated GHG emissions (kt CO₂-equivalents yr⁻¹) for 2016 per activity of KP-LULUCF.

KP activity	Emissions for 2016 (kt CO ₂ -equivalents yr ⁻¹)		
	NIR 2019	NIR 2018	Absolute difference
Afforestation	-505.87	-519.18	13.31
Deforestation	2 230.14	2 325.99	-95.85
Forest management	-27 719.40	-28 396.13	676.73
Cropland management	1 773.53	1 769.89	3.64
Grazing land management	10.20	0.37	9.83
Total	-24 211.40	-24 819.06	607.66

Total recalculations for the KP-LULUCF submissions for the year 2016³⁹ resulted in reduced C uptake of 608 kt CO₂-eq from -24 819 kt CO₂-eq (NIR 2018) to -24 211 kt CO₂-eq (NIR 2019) including non-CO₂ emissions. The majority of the change was due to the reduction in CO₂ uptake for FM (reduced by 677 kt CO₂-eq). The CO₂ uptake for AR decreased by 13 kt CO₂-eq, while emissions reduced by 96 kt CO₂-eq for D including non-CO emission. Emissions increased by 4 kt CO₂-eq for CM and by 10 kt CO₂-eq for GM, including non-CO₂ emissions. Recalculation for CH₄ emissions were minor and increased total KP-LULUCF CH₄ emissions by 3 kt CO₂-eq, from 158 kt CO₂-eq to 161 kt CO₂-eq. N₂O emissions were also recalculated and decreased by 42 kt CO₂-eq from 343 kt CO₂-eq to 304 kt CO₂-eq. Recalculations of CO₂ emissions were more important and these are described below for each KP activity.

11.3.3.1 Afforestation and reforestation – KP. A.1

- Total CO₂ sequestration for AR decreased by 13 kt CO₂ from 542 kt CO₂ to 529 kt CO₂, which was primarily due to the increased C uptake of 2 kt CO₂ in the aboveground living biomass (from 165 kt CO₂ to 167 kt CO₂). Belowground biomass sequestration decreased 4 kt CO₂ (from 49 kt CO₂ to 45 kt CO₂). C uptake in the litter pool decreased by 8 kt CO₂ from 433 kt CO₂ to 425 kt CO₂. C losses from the mineral soil pool were increased by 3 kt CO₂ from 86 kt CO₂ to 89 kt CO₂. All recalculations were due to the changes in the NFI area and living biomass data. NFI harvest data are used to distribute the known total harvests to the reported activities such as afforestation.

11.3.3.2 Deforestation – KP. A.2

- Total CO₂ emissions for deforestation were reduced by 95 kt CO₂ from 2 298 kt CO₂ to 2 203 kt CO₂. This was caused by the combination of reduced emissions from litter (17 kt CO₂ from 1 307 kt CO₂ to 1 290 kt CO₂), dead wood (1.4 kt CO₂ from 107 kt CO₂ to 106 kt CO₂), and below ground biomass (14.5 kt CO₂ from 148.7 kt CO₂ to 134.2 kt CO₂). In addition a slight increase in the sequestration in mineral soils (0.7 kt CO₂; from 137.5 kt CO₂ to 138 kt CO₂), a smaller emission from aboveground biomass (58 kt CO₂ from 581 kt CO₂ to 523 kt CO₂) and a decrease in emissions from organic soils (4 kt CO₂ from 292 kt CO₂ to 288 kt CO₂). Recalculations for all C pools were due to the updates in the NFI data.

11.3.3.3 Forest management – KP.B.1

- Carbon removals in the FM activity reduced by 714 kt CO₂. The majority of the reduction occurred in the living biomass; removals aboveground biomass was reduced by 631 kt CO₂ from 17 184 kt CO₂ to 16 553 kt CO₂ and belowground by 261 kt CO₂ from 4 108 kt CO₂ to 3 847 kt CO₂. The removals in the litter, dead wood, and mineral soil pools all increased due to a larger C uptake of 95 kt CO₂ (from 6 566 kt CO₂ to 6 661 kt CO₂), 19 kt CO₂ (from 1 326 kt CO₂ to 1 345 kt CO₂), and 2 kt CO₂ (from 151 to 153 kt CO₂), respectively. Emissions from organic soils decreased 60 kt CO₂ from 696 kt CO₂ to 636 kt CO₂. This change resulted from the correction of the proportion of nutrient poor vs nutrient rich organic soil, which were changed from 21% vs 79% to 31% vs 69%, respectively.

³⁹ Note that recalculations for KP-LULUCF, as described in the following, refer to absolute values of the activities, and not the value that is accounted for under KP2.

11.3.3.4 Cropland management – KP.B.2

- Total CO₂ emissions for the CM activity increased 5 kt CO₂ (from 1 681 kt CO₂ to 1 686 kt CO₂). Removals in aboveground biomass increased by 0.6 kt CO₂ from 2.5 kt CO₂ to 3.1 kt CO₂ as did belowground biomass which increased 0.3 kt CO₂ from 1.0 kt CO₂ to 1.3 kt CO₂. The removals in the mineral soil pool decreased 40.7 kt CO₂ from 75.7 kt CO₂ to 35.0 kt CO₂. The emissions in the organic soil pool decreased 34 kt CO₂ from 1 760 kt CO₂ to 1 726 kt CO₂. Recalculation differences in mineral soils were due to corrections described under “4B1 Cropland remaining cropland” above and organic soils estimates were affected by the NFI extrapolation method.

11.3.3.5 Grazing land management – KP. B.3

- Recalculations for the GM activity resulted in increased emissions by 8 kt CO₂ from -6 kt CO₂ to 2 kt CO₂. This was primarily the result of the combined effect of decreased C uptake in above- and belowground living biomass (4.5 kt CO₂ for aboveground and 2 kt CO₂ for belowground) and slightly increased emissions from organic soils 0.0001 kt CO₂. Recalculation for living biomass was due to the updates in the NFI database. The change in the NFI database affected the area of organic soils under GM.

11.3.4 Omissions of carbon pool or GHG emissions/removals from activities under Article 3.3 and elected activities under Article 3.4

No omissions were made of any C pools or GHG emissions.

11.3.5 Provisions for natural disturbances

Norway does not apply the provisions for natural disturbances to its accounting in the second commitment period.

11.3.6 Emissions and removals from the harvested wood product pool

The reporting of emissions and removals from the HWP pool under the KP is done in accordance with Decision 2/CMP.7, Annex § 16 and 27-32, and Decision 2/CMP.8 Annex II, § 2(g)(i-vii). Emissions from HWP in solid waste disposal sites are reported in the waste sector. As the FMRL is not based on a projection (but the 1990 base year), it is not relevant to provide further information in this regard. There is no double accounting from the HWP pool in the second commitment period because emissions/removals were not accounted under the first commitment period according to the Marrakesh Accords (Decision 11/CP.7), thus there is no need to exclude these emissions/removals from the accounting under the second commitment period. For reporting under deforestation, the Tier 1 method is applied and carbon stock changes in the HWP pool are reported as zero (NO).

Norway uses the Tier 2 method to estimate carbon stock change in the harvested wood products pool. The calculations follow (IPCC, 2014a) including: the three default HWP categories sawnwood,

wood-based panels, and paper and paperboard along with their associated half-lives and conversion factors.

All the activity data are obtained from FAO forestry statistics⁴⁰. The initial unit is m³, except for the pulp and paper, where the unit is metric ton. Exported and domestically consumed HWP is calculated and reported separately. The inflow data of domestically produced and consumed HWP are based on consumption (Production – Export), since including export could result in double counting. Imported HWP is not included in the calculations (Production approach).

The following are specifics from IPCC (2014a) and applicable only to the reporting of HWP under KP and do not apply for the Convention reporting:

The annual fraction of feedstock for HWP production originating from domestic harvest is estimated applying Eq. 2.8.1 (IPCC, 2014a)

$$f_{IRW}(i) = \frac{IRW_p(i) - IRW_{EX}(i)}{IRW_p(i) + IRW_{IM}(i) - IRW_{EX}(i)}$$

where $f_{IRW}(i)$ = fraction of industrial roundwood for the domestic production of HWP originating from domestic forests in year i ; $IRW_p(i)$ = domestic production of industrial roundwood in year i ; $IRW_{IM}(i)$ = import of industrial roundwood in year i ; $IRW_{EX}(i)$ = export of industrial roundwood in year i .

The annual fraction of feedstock for paper and paperboard production originating from domestically produced wood pulp is estimated applying Eq. 2.8.2 (IPCC, 2014a)

$$f_{PULP}(i) = \frac{PULP_p(i) - PULP_{EX}(i)}{PULP_p(i) + PULP_{IM}(i) - PULP_{EX}(i)}$$

where $f_{PULP}(i)$ = fraction of domestically produced pulp for the domestic production of paper and paperboard in year i ; $PULP_p(i)$ = production of wood pulp in year i ; $PULP_{IM}(i)$ = import of wood pulp in year i ; $PULP_{EX}(i)$ = export of wood pulp in year i .

The annual fraction of feedstock for HWP originating from forest activities under Article 3.3 and 3.4 (FM or AR or D) in year i is calculated from the total harvest (kt C) applying Eq. 2.8.3 (IPCC, 2014a)

$$f_j(i) = \frac{harvest(i)}{harvest_{Total}(i)}$$

where $f_j(i)$ = fraction of harvest originating from the particular activity j in year i , j = activity FM or AR or D in year i (aboveground C losses in living biomass as reported in the CRF tables 4(KP-I)A.1, 4(KP-I)A.2, and 4(KP-I)B.1).

The annual HWP resulting from domestic harvests related to activities under Article 3.3 and 3.4 was estimated as the product of the production of the commodity, the annual fraction of the feedstock, and the fraction of the domestic feedstock for each of the HWP categories applying IPCC 2014 Eq. 2.8.4.

⁴⁰ FAO data available from: <http://www.fao.org/faostat/en/#data/FO>

The carbon stock change of the HWP pool was estimated for each of the KP activities AR and FM by

$$HWP_j(i) = [HWP_p(i) \times f_{DP}(i) \times f_j(i)]$$

where $HWP_j(i)$ = the reported estimates in the CRF tables = HWP resulting from domestic harvest associated with activity j in year i , in $\text{m}^3 \text{yr}^{-1}$ or Mt yr^{-1} , $HWP_p(i)$ = production of the particular HWP commodities (i.e. sawnwood, wood-based panels, and paper and paperboard) in year i , in $\text{m}^3 \text{yr}^{-1}$ or Mt yr^{-1} , $f_{DP}(i)$ is the fraction of domestic feedstock for the production of the particular HWP category originating from domestic forest in year i , and $f_{DP}(i) = f_{IRW}(i)$ for HWP categories 'sawnwood' and 'wood-based panels', $f_{DP}(i) = (f_{IRW}(i) \times f_{PULP}(i))$ for HWP category 'paper and paperboard' with: $f_{IRW}(i) = 0$ if $f_{IRW}(i) < 0$ and $f_{PULP}(i) = 0$ if $f_{PULP}(i) < 0$, where: $f_j(i)$ = fraction of domestic feedstock for the production of the particular HWP category originating from domestic forests in the activity j = FM or AR.

For land subjected to deforestation, gains and losses in the HWP pool is reported as NO to resemble instantaneous oxidation.

Harvests (h) in a reporting year were reported as

$$h = l \cdot f$$

where l are the reported losses of the aboveground living biomass in the year of interest and the activity considered, and $f = 0.564$ is the stem fraction. The stem fraction is the average proportion of stem biomass of the total biomass and is calculated from all registered trees on NFI plots in the season prior to harvest independent of tree species.

11.3.7 Information on whether emissions and removals have been factored out

Emissions and removals have not been factored out.

11.4 Article 3.3

11.4.1 Activities under Article 3.3 began on or after 1 January 1990 and before 31 December of the last year of the commitment period and are directly human-induced

The NFI covers the period of consideration. The permanent plots were established between 1986 and 1993. Since then the plots have been monitored continuously beginning with the first re-inventory in 1994 (see chapter 6.3). By repeatedly assessing the land cover and land use on each plot, the NFI records land-use changes to and from forest land.

In order to be included as AR and D activities under Article 3.3, land-use changes must be directly human-induced. For AR and D, land-use changes are considered directly human-induced in the following two cases: (1) all conversions to forest land from land-use categories which are considered managed (cropland, grassland, and settlements); and (2) conversions from wetlands or other land (non-managed lands) to forest land, when actual evidence of management is present. Such evidence consists of planting and ditching, which can both be documented via the current status of the forest in combination with aerial photos. Land-use changes from wetlands or other land to forest land is considered to be the natural expansion of the forest if no direct evidence of management is present. Land-use changes between forest land, wetlands, or other land can either be reported as FM in cases of non-human induced changes, or reported as AR or D for human-induced changes (see Table 11.2).

11.4.2 How harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation

Young natural stands and all plantations established for forestry purposes, as well as forests that are temporarily unstocked as a result of e.g. harvest or natural disturbances, are included under forest management and not treated as deforestation. The NFI teams assess land cover and land use according to national criteria (see Table 6.10) that are defined in the field protocol (NFLI, 2008). They are also trained to distinguish between forest management operations and land-use change. As a general rule, land will be considered temporarily unstocked if the stumps and ground vegetation are still present, and there is no construction work done on the area. The area is considered deforested if the ground vegetation is removed (e.g. if the area is leveled), and/or other construction work is done on the area.

11.5 Article 3.4

11.5.1 Activities under Article 3.4 occurred since 1 January 1990 and are human-induced

The NFI covers the period of consideration for all activities elected (FM, CM, and GM). The permanent plots were installed from 1986 until 1993. From 1994 and onwards the plots have been monitored continuously. As described above, certain criteria apply.

11.5.2 Information relating to Cropland Management, Grazing Land Management, Revegetation, and Wetland Drainage and Rewetting, if elected, for the base year

To identify the areas included in the cropland management (CM) and grazing land management (GM) activities in the base year (1990), we define the management practices identically to those on cropland and grassland as defined under the Convention. The management practices on the cropland land-use class are the same as those that take place on land included under the CM activity. This is the same for the grassland land-use class and the GM activity. The only difference is that CM or GM can include land that was cropland or grassland in 1990 and has since been converted to a non-forest category (e.g. settlements). Under the KP reporting, land can only leave an activity if it enters another activity on a higher hierarchical level. Therefore, the following land use and land-use change classes⁴¹ are considered under CM and GM:

$$CM = \underline{CC} + \underline{GC} + \underline{WC} + \underline{SC} + \underline{CS} + CG + CW + CO,$$

$$GM = \underline{GG} + \underline{WG} + \underline{SG} + OG + \underline{GO} + \underline{GS} + GW.$$

Conversion categories that have occurred in Norway are in bold and underlined. Due to the 20 year conversion rule applied under the Convention, areas of some land-use change classes were not identical to those reported under the Convention. Under the Convention, areas in the categories land converted to cropland and land converted to grassland will be transferred to CC or GG after 20 years. Under the KP, these areas will therefore automatically stay in CM or GM even after 20 years. However, areas of cropland or grassland converted to other land-uses would also be transferred to the remaining category of that land-use under the 20 year rule. We therefore did not apply the 20 year rule for the CS, GS, and GO land-use change classes that are included in CM or GM. This is illustrated in the CRF tables (4(KP-I)B.2 and 4(KP-I)B.3) in the sub-division under Norway for CM and GM.

⁴¹ The land-use change class abbreviations are: C = Cropland, G = Grassland, W = Wetlands, S = Settlements, and O = Other land. The first letter in the abbreviation is the starting class and the second letter is the class in which the land remains or is converted to. For example, the abbreviation CC means "Cropland remaining Cropland" and GC means "Grassland converted to Cropland".

11.5.3 Emissions and removals from Forest Management, Cropland Management, and Grazing land Management under Article 3.4 are not accounted for under activities under Article 3.3

Neither the NFI used to track land areas, nor the methodologies applied to estimate emissions and removals from activities under Article 3.4 allow any double counting.

11.5.4 Conversion of natural forests to planted forests

This is not applicable for Norway.

11.5.5 Methodological consistency between the reference level and forest management reporting and technical corrections

Norway has chosen 1990 as the base year for the forest management reference level (FMRL). The FMRL presented in the appendix to decision 2/CMP.7 has been recalculated. Hence, a technical correction is required.

The corrected FMRL and the technical correction were obtained in the following way:

- First, all FM-related net C stock changes in 1990 (kt CO₂ equivalents.) were added to obtain the corrected FMRL.
- Second, the technical correction was obtained by subtracting the original FMRL from the corrected FMRL. The technical correction is the same for all years. See Table 11.8 for more details and the current technical correction.

The biggest differences compared to the original FMRL are due to:

- Increased net carbon stock gains in living biomass due to a changed interpolation procedure.
- Reduced net uptake in the dead organic matter and mineral soil pools since Yasso07 is now used on a NFI plot scale.
- The inclusion of HWP.
- Increased emissions from drained organic soils since the default Tier 1 emission factors have increased from the 2003 good practice guidance (IPCC, 2003) to the IPCC 2013 Wetlands Supplement (IPCC, 2014b).
- The use of new country-specific biomass functions for birch by Smith et al. (2014) and Smith et al. (2016).

Further details on the methodological changes have been described in the relevant sections of chapter 6 LULUCF.

Table 11.8 Components of the original and corrected Forest Management Reference Level (FMRL).

Source/sink	Original FMRL (kt CO ₂ -eq./year)	Corrected FMRL (kt CO ₂ -eq./year)
Living biomass ^a	-6 420	-9 686.20
Dead organic matter ^b	-2 040	-2 579.17
Mineral soils ^c	-3 060	-49.24
Biomass burning (Wildfires – N ₂ O and CH ₄) ^d	2	1.29
Fertilization ^e	1	0.97
Drainage of (organic) soils under Forest management ^f	150	899.13
HWP ^g	NE	-1 069.16
N ₂ O emissions due to land-use conversions and management change in mineral soils ^h	NE	NO
Sum	-11 370 ⁱ	-12482.45 ^j
Forest Management Reference Level (FMRL)	-11 400 ^k	
Technical correction		-1082.45
Corrected FMRL		-12482.45

^a All Norwegian forests including mountain forest and Finnmark were considered in the original FMRL. Sum of "Above" (cell S11) and "Below-ground biomass Net change" (cell V11) in the CRF table "4(KP-I)B.1 1990" converted to CO₂ equivalents. (C stock change x 44/12).

^b Below the coniferous limit in the original FMRL. All Norwegian forests including mountain forest and Finnmark in the corrected FMRL. Sum of "Litter" (cell W11) and "dead wood" (cell X11) in the CRF table "4(KP-I)B.1 1990" converted to CO₂ equivalents.

^c Below the coniferous limit and denoted "Soil organic matter" in the original FMRL. The value for the corrected FMRL is obtained from cell Y11 of the CRF table "4(KP-I)B.1 1990" converted to CO₂ equivalents.

^d Sum of CH₄ (cell I27) and N₂O (cell J27) converted to CO₂ equivalents. for "Forest management" in the CRF table "4(KP-II)4 1990". GWP were 25 for CH₄ and 298 for N₂O (see http://www.ipcc.ch/publications_and_data/ar4/wg1/en/errataserrata-errata.html#table214).

^e Direct and indirect N₂O emissions from N fertilization (cell D13) in the CRF table "4(KP-II)1 1990" converted to CO₂ equivalents for "Forest management".

^f Only included CO₂ and N₂O in the original FMRL. Also contains CH₄ in the corrected FMRL. "Organic soils" in the table "4(KP-I)B.1 1990" (cell Z11) and "Drained organic soils" in the CRF table "4(KP-II)2 1990" (cells E17 and F17) converted to CO₂ equivalents.

^g Cell M32 in the CRF table "4(KP-I)C 1990".

^h This source is now included but was 0 for 1990 in the current reporting (cell E17 in the CRF table "4(KP-II)3 1990").

ⁱ Sum refers to the FMRL as given in Table 3 in the submission to the UNFCCC from March 2011, available: http://unfccc.int/files/meetings/ad_hoc_working_groups/kp/application/pdf/awgkp_norway_2011.pdf. The actual value is -11367 but the values were reported in Mt and rounded to the second decimal, ie. 11.4 Mt CO₂-equivalents./year cf. footnote ^k.

^j Since the 2016 submission, this value is taken from cell E11 of the current CRF table "4(KP) 1990" directly, rather than adding the values in this table. Therefore, the last digit of the value is not exactly equal to the sum of the sources/sinks above. A possible difference of the last digit is due to rounding.

^k FMRL as inscribed in the appendix to the annex to Decision 2/CMP.7, ie. -11.400 (kt CO₂-equivalents./year).

11.5.6 Information about emissions or removals resulting from the harvest and conversion of forest plantations to non-forest land

This is not applicable for Norway.

11.6 Other information

11.6.1 Key category analysis for Article 3.3 activities and any elected activities under Article 3.4.

According to the IPCC guidelines, the key-category analysis for KP can be based on the assessment made for the Convention inventory reporting (see chapter 1.5 for details). Additionally, the key categories are reported in CRF table NIR 3. Both Approach 1 and Approach 2 assessments are made for the whole inventory including the LULUCF sector (Table 1.1; Table 1.2). The key-category analysis is made specific to sink/source categories per individual land-use conversion (e.g. forest land converted to cropland instead of land converted to cropland). The analysis can, therefore, not be directly translated into the KP activities, but by combining the information in Table 6.6 and the relation between Convention land-use categories and KP activities shown in Table 11.2, we can derive the key categories. Any sink/source under the AR, D, CM or GM activities was considered as a key category if at least one of the land-use transitions within the activity was identified as a key category in the analysis.

11.7 Information relating to Article 6

There are no Article 6 activities concerning the LULUCF sector in Norway.

12 Information on accounting of Kyoto units

12.1 Background information

Norway's Standard Electronic Format (SEF) reports for 2018 (for the first and the second commitment period) are reported as annex VII to this document and will be made available at the UNFCCC website. The name of the files are SEF_NO_2018_CP1.xlsx and SEF_NO_2018_CP2.xlsx.

12.2 Summary of information reported in the SEF tables

The tables below show the amount of different units; AAUs, ERUs, CERs, tCERs, ICERs and RMUs, from CP1 and CP2 within the registry, and on which account type these units are present at the end of 2018.

Table 12.1 AAUs from CP1

AAUs	# CP1 UNITS
Party holding account	5 984 774
Entity holding account	85 225
Article 3.3/3.4 net source cancellation accounts	1 824 462
Other cancellation accounts	8 433 893
Retirement account	253 134 092
TOTAL amount	269 462 446

Table 12.2 ERUs from CP1 and CP2

ERUs	# CP1 UNITS	# CP2 UNITS
Party holding account	NO	738 305
Entity holding account	NO	84 848
Other cancellation accounts	1 098 212	
Retirement account	2 605 670	NO
TOTAL amount	3 703 882	823 153
TOTAL CP1 + CP2 ERUs	4 527 035	

Table 12.3 CERs from CP1 and CP2

CERs	# CP1 UNITS	# CP2 UNITS
Party holding account	341	21 433 553
Entity holding account	654	634 399
Other cancellation accounts	19 701 689	
Voluntary cancellation accounts		176 634
Retirement account	9 260 279	NO
TOTAL amount	28 962 963	22 244 586
TOTAL CP1 + CP2 CERs	51 207 549	

Table 12.4 tCERs from CP1 and CP2

tCERs	# CP1 UNITS	# CP2 UNITS
Party holding account	NO	NO
Entity holding account	NO	NO
Other cancellation accounts	35 424	
Voluntary cancellation accounts		NO
Retirement account	NO	NO
TOTAL amount	35 424	NO
TOTAL CP1 + CP2 tCERs	35 424	

Table 12.5 RMUs from CP1 and CP2

RMUs	# CP1 UNITS	# CP2 UNITS
Party holding account	NO	NO
Entity holding account	NO	NO
Article 3.3/3.4 net source cancellation accounts	9 947 523	NO
Other cancellation accounts	7 333 333	
Voluntary cancellation accounts		NO
Retirement account	1 824 462	NO
TOTAL amount	19 105 318	NO
TOTAL CP1 + CP2 RMUs	19 105 318	

The registry did not contain any ICERs from CP1 or CP2.

The total amount of the units in the Norwegian registry at the end of 2018 corresponded to 344 337 772 tonnes CO₂ eq.

The following account types did not contain any units:

- ICER Replacement Account for Expiry (CP1 and CP2)
- ICER Replacement Account Non-Submission Report (CP1 and CP2)
- ICER Replacement Account Reversal in Storage (CP1 and CP2)
- Mandatory Cancellation Account CP2
- Net Source Cancellation Account CP2
- Non-compliance Cancellation Account (CP1 and CP2)
- Retirement Account CP2
- tCER Replacement Account for Expiry (CP1 and CP2)

The following account types did not exist in the registry:

- Article 3.1 ter and quarter ambition increase cancellation account
- Article 3.7 ter cancellation account
- Cancellation account for remaining units after carry-over
- ICER cancellation account for expiry
- ICER cancellation account for reversal of storage
- ICER cancellation account for non-submission of certification report
- Previous Period Surplus Reserve account (PPSR)
- tCER cancellation account for expiry

12.3 Discrepancies and notifications

Table 12.6 Discrepancies and notifications

Annual Submission Item	Reporting information
15/CMP.1 annex I.E paragraph 12: List of discrepant transactions	No discrepant transaction occurred in 2018.
15/CMP.1 annex I.E paragraph 13 & 14: List of CDM notifications	No CDM notifications occurred in 2018.
15/CMP.1 annex I.E paragraph 15: List of non-replacements	No non-replacements occurred in 2018.
15/CMP.1 annex I.E paragraph 16: List of invalid units	No invalid units exist as of 31 December 2018.
15/CMP.1 annex I.E paragraph 17 Actions and changes to address discrepancies	No discrepant transactions occurred in 2018.

We have not submitted the R2- R5 reports since none of these events have occurred in the registry, and these reports would thus be empty.

12.4 Publicly accessible information

Information relating to the Norwegian registry, which is deemed to be public information, can be accessed via the [Kyoto Protocol Public Reports](http://www.miljodirektoratet.no/no/Tema/klima/CO2_kvoter/Klimakvoteregisteret/public_reports/) page in the national registry. The SEF reports may also be downloaded from the registry web site,

http://www.miljodirektoratet.no/no/Tema/klima/CO2_kvoter/Klimakvoteregisteret/public_reports/.

In accordance with the requirements of the Annex to Decision 13/CMP.1, all required information for a Party with an active Kyoto registry is provided with the exceptions as outlined below:

Account Information (Paragraph 45) and Account holders authorised to hold Kyoto units in their account (Paragraph 48)

In line with the data protection requirements of Regulation (EC) No 45/2001 and Directive 95/46/EC and in accordance with Article 110 and Annex XIV of Commission Regulation (EU) No 389/2013, the information on account representatives, account holdings, account numbers, legal entity contact information, all transactions made and carbon unit identifiers, held in the EUTL, the Union Registry and any other KP registry (required by paragraph 45 and paragraph 48) is considered confidential. This information is therefore not publicly available.

JI projects in Norway (Paragraph 46)

No information on Article 6 (Joint Implementation) projects is publicly available, as conversion to an ERU under an Article 6 project did not occur in Norway in 2018.

Holding and transaction information of units (Paragraph 47)

General remarks

Holding and transaction information is provided on a holding type level due to more detailed information on transactions being considered confidential according to Article 110 of Commission Regulation (EU) no 389/2013, ref. paragraph 47(a), 47(d), 47(f) and 47(l).

Article 110 of Commission Regulation (EU) no 389/2013 provides that “Information, including the holdings of all accounts, all transactions made, the unique unit identification code of the allowances and the unique numeric value of the unit serial number of the Kyoto units held or affected by a transaction, held in the EUTL, the Union Registry and other KP registry shall be considered confidential except as otherwise required by Union law, or by provisions of national law that pursue a legitimate objective compatible with this Regulation and are proportionate”.

Paragraph 47(b)

No AAUs were issued on the basis of the assigned amount pursuant to Article 3, paragraphs 7 and 8 in 2018.

Paragraph 47(c)

Norway does not host JI projects. Therefore no ERUs have been issued on the basis of Article 6 projects.

Paragraph 47(e)

No RMUs were issued in 2018.

Paragraph 47(g)

No ERUs, CERs, AAUs or RMUs were cancelled on the basis of activities under Article 3.3 and 3.4 in 2018.

Paragraph 47(h)

No ERUs, CERs, AAUs and RMUs were cancelled following determination by the Compliance Committee that the Party does not comply with its commitment under Article 3, paragraph 1 in 2018.

Paragraph 47(i)

No ERUs, CERs, AAUs and RMUs were cancelled in 2018.

Paragraph 47(j)

No ERUs, CERs, AAUs nor RMUs were retired in 2018.

Paragraph 47(k)

No ERUs, CERs or AAUs were carried over from previous commitment periods in 2018.

Paragraph 47(a) bis

Norway did not have any previous period surplus reserve account (PPSR) at the beginning of 2018, and therefore had no AAUs in such an account.

Paragraph 47(h) bis

No AAUs were cancelled under Article 3, paragraphs 1 ter and 1 quarter in 2018.

Paragraph 47(h) ter

No AAUs were cancelled under Article 3, paragraph 7 ter in 2018.

12.5 Calculation of the commitment period reserve (CPR)

The reporting of the calculation of the commitment period reserve, pursuant to decision 11/CMP.1, 15/CMP.1, 1/CMP.8 and 3/CMP.11 is as follows:

The commitment period reserve is the lower of the two values given by 90 percent of the assigned amount and eight times 100 percent of the total emissions in the most recently reviewed inventory.

The assigned amount for Norway for the Kyoto Protocol's second commitment period has been set to 348 914 303 t CO₂ equivalents through the review of our report to facilitate the calculation of its assigned amount (FCCC/IRR /2016/NOR).

90 % of the assigned amount:

The review report (FCCC/IRR/2016/NOR) determined that with the appropriate rounding rules, 90 % of the assigned amount equals:

= 314 022 874 ton CO₂ equivalents

100 % of eight times its most recently reviewed inventory:

The common approach/guidance is that for this approach of calculating the CPR, the most recent inventory submission should be used because, at the end of the review, this becomes the most recently reviewed inventory. Hence, the emissions without LULUCF in 2017 (52 712 543 ton CO₂ equivalents) are used. With the appropriate rounding rules, 100 % of eight times the most recently reviewed inventory equals:

= 421 700 344 ton CO₂ equivalents

For Norway, the lowest number is equivalent to 90 % of Norway's assigned amount. Norway's commitment period reserve is therefore:

= 314 022 874 ton CO₂ equivalents

13 Information on changes in the National System

13.1 Changes in the National Greenhouse Gas Inventory System

Statistics Norway, one of the three parts in the Norwegian National System, has undergone a reorganization of staff and work areas between its two offices/locations; Oslo and Kongsvinger. The experts compiling the emission inventory for all sectors except LULUCF, was up to 2018 located in Oslo. This group of experts has through 2018 been replaced by a new staff located in Kongsvinger. The long term goal of this relocation is to improve data quality by increasing the contact and collaboration between the departments producing the input (activity) data and the inventory compilers.

14 Information on changes in national registry

Table 14.1 shows the changes to the national registry of Norway that have occurred in 2018.

Table 14.1 Changes to the national registry of Norway in 2018.

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(a) Change of name or contact	None
15/CMP.1 Annex II.E paragraph 32.(b) Change regarding cooperation arrangement	No change of cooperation arrangement occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(c) Change to database structure or the capacity of national registry	<p>The version of the EUCR released after 8.0.8 (the production version at the time of the last Chapter 14 submission) introduced minor changes in the structure of the database.</p> <p>These changes were limited and only affected EU ETS functionality. No change was required to the database and application backup plan or to the disaster recovery plan. The database model is provided in Annex A.</p> <p>No change to the capacity of the national registry occurred during the reported period.</p>
15/CMP.1 annex II.E paragraph 32.(d) Change regarding conformance to technical standards	<p>Changes introduced since version 8.0.8 of the national registry are listed in Annex B.</p> <p>Each release of the registry is subject to both regression testing and tests related to new functionality. These tests also include thorough testing against the DES and were successfully carried out prior to the relevant major release of the version to Production (see Annex B).</p> <p>No other change in the registry's conformance to the technical standards occurred for the reported period.</p>

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(e) Change to discrepancies procedures	No change of discrepancies procedures occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(f) Change regarding security	No changes regarding security occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(g) Change to list of publicly available information	No change to the list of publicly available information occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(h) Change of Internet address	The registry internet address changed during the reported period. The new URL is https://unionregistry.ec.europa.eu/euregistry/XX/index.xhtml
15/CMP.1 annex II.E paragraph 32.(i) Change regarding data integrity measures	No change of data integrity measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(j) Change regarding test results	Changes introduced since version 8.0.8 of the national registry are listed in Annex B. Both regression testing and tests on the new functionality were successfully carried out prior to release of the version to Production. The site acceptance test was carried out by quality assurance consultants on behalf of and assisted by the European Commission.
The previous Annual Review recommendations	See below

The latest annual review report for Norway is FCCC/ARR/2016/NOR. In this review report, the recommendation concerning the national registry made in the previous review report was considered as resolved and no additional finding concerning the national registry was made during the review of the 2016 technical review. The latest Standard Independent Assessment Reports for Norway are SIAR/2018/1/1 for part 1 (completeness) and SIAR/2017/2/1 for part 2 (substance).

There was no recommendations concerning the Norwegian national registry in the SIAR reports.
There are no outstanding recommendations for the registry, Table 14.2 is therefore empty.

Table 14.2 Follow-up to recommendations for the registry.

Reference	Recommendation description	Response

15 Information on minimization of adverse impacts in accordance with Art. 3.14

Norway strives to follow a comprehensive approach to climate change mitigation since policy development started around 1990, addressing all sources as well as sinks, in order to minimise adverse effects of climate policies and measures on the economy. In developing environmental, as well as economic and energy policy, Norway endeavours to include the polluter pays principle and to have a market-based approach where prices reflect costs including externalities.

As regards emissions of greenhouse gases, costs of externalities are reflected by levies and by participation in the European Emissions Trading Scheme (EU ETS). These instruments place a price on emissions of greenhouse gases. Norway believes that the best way to reduce emissions on a global scale, in line with the two degree target and striving for 1.5 degree limit, would ideally be to establish a global price on emissions. Pursuing a global price on emissions would be the most efficient way to ensure cost-effectiveness of mitigation actions between different countries and regions, and secure equal treatment of all emitters and all countries. This will help minimise adverse impacts of mitigation efforts. For more information about levies on energy commodities and the design of the EU ETS, see Chapter 4. of the seventh National Communication.

Norway is involved in several initiatives that contribute to technology transfer and capacity building to developing countries in shifting the energy mix away from high emission sources to more renewable energy systems and low-emission sources. These initiatives are reported here as relevant activities under Article 3.14 of the Kyoto Protocol.

National strategy for green competitiveness

The government presented a national strategy for green competitiveness in October 2017. The aim of the strategy is to provide more predictable framework conditions for a green transition in Norway, while maintaining economic growth and creating new jobs. In October the government also appointed an expert commission to analyze Norway's exposure to climate risk.

Cooperation on carbon capture and storage

Both the International Energy Agency (IEA) and the Intergovernmental Panel on Climate Change have pointed out that Carbon capture and storage (CCS) will be an important mitigation tool. In order for CCS to become a viable mitigation tool, countries and companies need to invest in technology development and demonstration of CO₂ capture and storage projects.

CCS is one of five priority areas for enhanced national climate action. Norway strives to disseminate information and lessons learned from projects in operation in the petroleum sector, new large scale projects under planning and from research, development and demonstration projects. The information and lessons learned are shared both through international fora, and through bilateral cooperation with developing and developed countries.

Norway has a long experience with CCS. Since 1996, CO₂ from natural gas production on the Norwegian shelf has been captured and reinjected into sub-seabed formations. The CCS projects from natural gas on the Sleipner, Gudrun and Snøhvit petroleum fields are the only CCS projects currently in operation in Europe and the only projects in the offshore industry.

The Technology Centre Mongstad (TCM) is the world's largest facility of its kind for testing and improving CO₂ capture technologies. TCM has been operating since 2012, providing an arena for targeted development, testing and qualification of CO₂ capture technologies on an industrial scale

Norway also provides funding for CCS projects abroad in cooperation with other countries and through existing programmes and institutions.

In Norway, funding for CCS research is provided through the CLIMIT programme. The CLIMIT programme is a national programme for research, development and demonstration of technologies for capture, transport and storage of CO₂ from fossil-based power production and industry. The programme supports projects in all stages of the development chain, from long-term basic research to build expertise to demonstration projects for CCS technologies. Projects under the CLIMIT programme have yielded important results for the development of CCS in Norway and internationally.

In addition, a Centre for Environment-friendly Energy Research for CCS, NCCS, has been established. The centre is co-financed by the Research Council of Norway, industry and research partners.

The Norwegian Government has an ambition to realize a cost effective solution for full-chain CCS if this makes sense in a global technology development and transfer perspective.

In order for CCS to play an effective role in climate change mitigation, international cooperation on developing and commercialising new technology is essential. Norway collaborates with other countries through a number of regional and international forums. Examples of such forums are North Sea Basin Task Force, Clean Energy Ministerial, Mission Innovation and The Carbon Sequestration Leadership Forum. Norway furthermore provides funding for CCS projects abroad in cooperation with other countries and through existing programmes and institutions. For example, Norway is currently supporting CCS projects in South Africa and Mexico.

Cooperation with developing countries related to fossil fuels – “Oil for Development”

The Norwegian Oil for Development (OfD) programme, which was launched in 2005, aims at assisting developing countries, at their request, in their efforts to manage petroleum resources in a way that generates economic growth and promotes the welfare of the whole population in an environmentally sound way. A description of the OfD program can be found at: <https://www.norad.no/en/front/thematic-areas/oil-for-development/>. The programme is currently engaged in 14 countries, mainly in Africa.

The operative goal of the program is "economically, environmentally and socially responsible management of petroleum resources which safeguards the needs of future generations."

Petroleum plays an important role in an increasing number of developing countries. Oil and gas hold the promise of becoming vital resources for economic and social development. Unfortunately, in many cases it proves difficult to translate petroleum resources into welfare for the people. Decades

of experience in the oil and gas sector has given Norway valuable expertise on how to manage petroleum resources in a sustainable way. The Norwegian expertise can be useful for developing countries with proven petroleum resources, or countries that are in the exploration phase.

OfD takes a holistic approach meaning that management of petroleum resources, revenues, environment and safety are addressed in a coherent manner. Norwegian public institutions enter into long-term agreements with public institutions in partner countries. Assistance is directed towards three main outcomes: 1) policy makers set goals, define and assign responsibilities, 2) the authorities regulating the petroleum sector carry out their assigned responsibilities and 3) policy makers and regulatory authorities are held accountable for their management of the petroleum sector.

OfD assistance is tailor-made to the particular needs of each partner country. It may cover the designing and implementing legal frameworks, mapping of resources, environmental impact assessments, handling of licenses, establishing preparedness to handle accidents and oil spills, health, safety and environmental legislation, petroleum fiscal regimes and petroleum sovereign wealth fund issues as well as initiatives related to transparency, anti-corruption and climate change.

A Steering Committee formulates strategic direction, guidelines and priorities for the OfD. The Steering Committee consists of the Ministry of Foreign Affairs (Chair), the Ministry of Petroleum and Energy, the Ministry of Finance and the Ministry of Climate and Environment, The Ministry of Transport and Communications. The OfD secretariat resides in the Norwegian Agency for Development Cooperation (Norad). The OfD secretariat is responsible for coordination and implementation of the program. Norwegian embassies play a key role in the program, as they are responsible for the overall bilateral relations, have competence on the the local situation and have extensive development cooperation responsibilities. Key implementing institutions are the Norwegian Petroleum Directorate, the Norwegian Environment Agency, the Petroleum Safety Authority, the Norwegian Coastal Administration, the Norwegian Tax Administration and Statistics Norway. A range of research institutions, international organizations and consultancies are also involved. Furthermore, several national and international NGOs are contributing to the OfD initiative. These organizations are in particular involved in building civil society capacity on issues related to governance and petroleum activities in OfD partner countries. Moreover, Norway gives priority to the Extractive Industries Transparency Initiative (EITI). OfD also cooperates with Statistics Norway and coordinates its activities with the Office of the Auditor General of Norway.

There have been no major changes to these policies and activities in 2018.

Cooperation with developing countries related to renewable energy – “Clean energy for Development”

Energy has been at the core of Norway’s development assistance policy for several years. There has been a steady increase in funds allocated to clean energy activities over a period of years, both within bilateral and multilateral development assistance. For various reasons priorities of the ODA changed significantly in 2015. Overall spending to clean energy for development was reduced and amounted to about NOK 500 million in 2017. The budget for 2019 is increasing again to 1 142 million NOK, thus fulfilling the commitment to double the budget compared to 2017.

Seven countries receive most of the funding (Liberia, Mozambique, Myanmar, Nepal, Tanzania, and Uganda), but Norway is also engaged on a smaller scale in some other countries.

The overall objective of Norway's contribution to renewable energy is to contribute to access SDG 7 and the Paris agreement.

Norway's interventions in renewable energy is also seen as a contribution to reduce further development of coal power.

The main focus on the investments will be directed towards interventions that contribute to an enabling environment for commercial and private investment in the energy sector. Important activities are policy dialogue, sector reforms, legislation, institutional cooperation, planning and regional cooperation. The public power infrastructure, such as the distribution and transmission system is also important for private investments to take place and as such also an important area for Norway's development cooperation. Increased access is supported by grant funding for extension of the electricity grid as well as off-grid solutions.

Further, Norway provide support for feasibility studies, training, infrastructure in order to reduce risk as incentives for private investors in power production. Norway is currently looking into a further development of economic instruments directed towards the private sector, i.e. guarantees.

The Norwegian Investment Fund for Developing Countries (Norfund) is providing risk financing as equity and loans to clean energy projects together with private investors. Over time, Norfund is investing half of its capital in clean energy. The current investments in clean energy from Norfund is approx. NOK 10 billion. The Government has increased fund allocations to Norfund significantly over the past years.

Consequence assessments

Norway has issued Instructions for Official Studies and Reports (Utredningsinstruksen), laid down by Royal Decree. These Instructions deal with consequence assessments, submissions and review procedures in connection with official studies, regulations, propositions and reports to the Storting. The Instructions are intended for use by ministries and their subordinate agencies. The Instructions form part of the Government's internal provisions and deviation may only be allowed pursuant to a special resolution. The provisions make it mandatory to study and clarify financial, administrative and other significant consequences in advance.

In addition, Norway has a legal framework that deals specifically with environmental impact assessments. The purpose is to promote sustainable development for the benefit of the individual, society and future generations. Transparency, predictability and participation for all interest groups and authorities involved are key aims, and it is intended that long-term solutions and awareness of effects on society and the environment will be promoted.

16 References

- AEA (2007): *UK's national atmospheric inventory* Available at: <http://naei.defra.gov.uk/>
- Aker Engineering (1992): Direct hydrocarbon gas emissions from production and riser platforms: Aker Engineering
- Andrén, O., Kätterer, T. & Karlsson, T. (2004): ICBM regional model for estimations of dynamics of agricultural soil carbon pools. *Nutrient Cycling in Agroecosystems*, 70 (2): 231-239
- Anton-Fernandez, C. & Astrup, R. (2012): Empirical harvest models and their use in regional business-as-usual scenarios of timber supply and carbon stock development. *Scandinavian Journal of Forest Research*, 27 (4): 379-392
- Aquateam COWI AS (2014): Forbruk av annen organisk gjødsel for å beregne utslipp av N₂O fra jordbruket. *Commissioned report by Norwegian Environment Agency (Miljødirektoratet)*, Report nr: 14041. Project nr: O-14110. 29 pp
- Arts, R., Chadwick, A. & Eiken, O. (2005): Recent time-lapse seismic data show no indication of leakage at the Sleipner CO₂-injection site In *Proceedings of the 7th International Conference on Greenhouse Gas Technologies, Vancouver, Canada, 2004*, pp. 653-660. Oxford, UK: Elsevier
- Aspehølen Åby, B., Wetlesen, M. & Aass, L. (2017). *Enteric methane emissions from beef cows and sheep in Norway. Method description*. Greenhouse Gas Emissions 1990-2016, Annexes to NIR 2017. (Norwegian Environment Agency). 1Department of Animal and Aquacultural Sciences, Norwegian University of Life Sciences. Unpublished manuscript.
- Bakken, L. R., Marsteng, A. K. & Nadeem, S. (2012): Emissions of N₂O, CH₄, and CO₂ from VEAS waste water treatment plant: Department of Plant and Environmental Sciences, Norwegian University of Life Sciences
- Bang, J. R. (1993): Utslipp fra dieseldrevne anleggsmaskiner, arbeidsredskaper, traktorer og lokomotiver (Emissions from diesel-powered construction machinery, tools, tractors and locomotives). Oslo: National Institute of Technology (TI-Teknologisk institutt)
- Bang, J. R. (1996): *Utslipp av NMVOC fra fritidsbåter og bensindrevne motorredskaper (Emissions of NMVOC from leisure craft and gasoline-powered equipment)*. Oslo: National Institute of Technology (TI-Teknologisk institutt)
- Baritz, R., Seufert, G., Montanarella, L. & Van Ranst, E. (2010): Carbon concentration and stocks in forest soils of Europe. *Forest Ecology and Management*, 260 (3): 262-277
- Bechmann, M., Greipsland, I., Riley, H. & Eggestad, H. (2012): Nitrogen losses from agricultural areas - a fraction of applied fertilizer and manure. 31 pp
- Berge, G. & Mellem, K. B. (2013): Kommunale avløp. Ressursinnsats, utslipp, rensing og slamdisponering 2012. Gebyrer 2013. Oslo: Statistics Norway
- Bergfald & Co AS (2000): Metangassutslipp fra Norsk og Russisk gruvedrift på Svalbard (Emissions of methane from coal mining at Svalbard). Oslo: Bergfald & Co AS
- Bjørnnes, K. L. (2013): Emissions of HFCs and PFCs from product use in Norway. Documentation of methodologies: Statistics Norway, Documents 24/2013. <http://www.ssb.no/en/natur-og-miljo/artikler-og-publikasjoner/attachment/126338?ts=13f85bebb88>
- Björketun, U. & Nilsson, G. (2007): Skaderisker för motorcyklister (The risk of injury to motorcyclists), VTI report 566. Linköping: The Road Traffic Inspectorate (Vägtrafikinspektionen)
- Boldrin, A., Andersen, J. K. & Christensen, T. H. (2009): Environmental assessment of garden waste management in Århus Kommune (Miljøvurdering af haveaffald i Århus kommune) Technical University of Denmark - Miljø 3R (<https://www.aarhus.dk/~media/Subsites/AVA/Om-AVA/Bibliotek/Rapporter-og-undersogelser/Aarhus-haveaffald-engelsk-rapport.pdf>)

- Borgen, S. K., Grønlund, A., Andrén, O., Kätterer, T., Tveito, O. E., Bakken, L. R. & Paustian, K. (2012): CO₂ emissions from cropland in Norway estimated by IPCC default and Tier 2 methods. *Greenhouse Gas Measurement and Management*, 2 (1): 1-17
- Borgen, S. K. & Hylen, G. (2013): Emissions and methodologies for cropland and grassland used in the Norwegian national greenhouse gas inventory, Report 11/2013: Norwegian Forest and Landscape Institute (Norsk institutt for skog og landskap). 45 pp
- Breidenbach, J., Antón-Fernández, C., Petersson, H., McRoberts, R. E. & Astrup, R. (2013): Quantifying the model-related variability of biomass stock and change estimates in the Norwegian National Forest Inventory. *Forest Science*, 60 (1): 25-33
- Brunner, I., Bakker, M. R., Björk, R. G., Hirano, Y., Lukac, M., Aranda, X., Børja, I., Eldhuset, T. D., Helmissaari, H. S., Jourdan, C., et al. (2013): Fine-root turnover rates of European forests revisited: an analysis of data from sequential coring and ingrowth cores. *Plant Soil*, 362 (1-2): 357-372
- Chadwick, R. A., Arts, R., Eiken, O., Kirby, G. A., Lindeberg, E. & Zweigel, P. (2004): 4D Seismic Imaging of an Injected CO₂ Plume at the Sleipner Field, Central North Sea. In Davies, R. J. (ed.) vol. 29 *3D seismic technology: application to the exploration of sedimentary basins*, pp. 311-320. London, UK: Geological Society of London
- Chadwick, R. A., Arts, R. & Eiken, O. (2005): 4D seismic quantification of a growing CO₂ plume at Sleipner, North Sea. In Doré, A. G. & Vining, B. A. (eds) *Proceedings of the 6th Petroleum Geology Conference on "Petroleum Geology: North-West Europe and Global Perspectives"*, pp. 1385-1399 London, UK: Geological Society of London
- CMR (2011): Måle- og beregningsprogram for bestemmelse av utslipp av NMVOC i forbindelse med bøyelasting. Only in Norwegian
- Daasvatn, L., Flugsrud, K., Høie, H., Rypdal, K. & Sandmo, T. (1992): Modell for beregning av nasjonale utslipp til luft. Dokumentasjon (Model for calculation of national air emissions. Documentation), Interne notater 92/17: Statistics Norway (SSB-Statistisk sentralbyrå)
- Daasvatn, L., Flugsrud, K., Hunnes, O. K. & Rypdal, K. (1994): Beregning av regionaliserte utslipp til luft. Beskrivelse av modell og metoder for estimering (Calculation of emissions to air on a regional basis. Description of a model and estimation methods), Notater 94/16: Statistics Norway (SSB-Statistisk sentralbyrå)
- Dalsgaard, L., Astrup, R., Antón-Fernández, C., Borgen, S. K., Breidenbach, J., Lange, H., Lehtonen, A. & Liski, J. (2016): Modeling Soil Carbon Dynamics in Northern Forests: Effects of Spatial and Temporal Aggregation of Climatic Input Data. *PLOS ONE*
- Dalsgaard, L., Lange, H., Strand, L., Callesen, I., Borgen, S. K., Liski, J. & Astrup, R. (2016): Underestimation of boreal forest soil carbon stocks related to soil classification and drainage. *Canadian Journal of Forest Research*, 46 (12): 1413-1425
- Dalsgaard, L., Kjønnaas, O. J. & Lange, H. (2017): Forest soil carbon changes from measurements and models. Site-specific comparisons and implications for UNFCCC reporting. Ås: Norwegian Forest and Landscape Institute
- Dalsgaard, L., Kjønnaas, O. J. & Lange, H. (2017): Forest soil carbon changes from measurements and models. Site-specific comparisons and implications for UNFCCC reporting. Ås: Norwegian Institute for Bioeconomy Research. 112 pp
- de Wit, H. A. & Kvindesland, S. (1999): Carbon stocks in Norwegian forest soils and effects of forest management on carbon storage. Rapport fra skogforskningen. Supplement 14: Norwegian Forest Research Institute (Skogforsk). 52 pp
- de Wit, H. A., Palosuo, T., Hylen, G. & Liski, J. (2006): A carbon budget of forest biomass and soils in southeast Norway calculated using a widely applicable method. *Forest Ecology and Management*, 225 (1-3): 15-26
- DeAngelis, D. L., Gardner, R. H. & Shugart Jr, H. H. (1981): Productivity of forest ecosystems studied during IBP: the woodlands data set. In Reichle, D. E. (ed.) *Dynamic properties of Forest Ecosystems*, pp. 567-672. Cambridge: Cambridge University Press

- Directorate for Health and Social Affairs (2013): Utviklingen i norsk kosthold 2012, Matforsyningsstatistikk og Forbruksundersøkelser. 12/2103. IS-2116, <http://helsedirektoratet.no/publikasjoner/utviklingen-i-norsk-kosthold-langversjon-2013/Publikasjoner/IS-2116%20 langversjon.pdf>
- Dämmgen, U., Lüttich, M., Döhler, H., Eurich-Menden, B. & Osterburg, B. (2002): GAS-EM – a procedure to calculate gaseous emissions from agriculture. *Landbauforsch Völkenrode*, 52: 19-42
- EBA (2014): Expert judgement by Contractors Association - Building and Construction (EBA), Oslo, Norway.
- EEA (1996): EMEP/CORINAIR Atmospheric Emission Inventory Guidebook - 1st edition: EEA-European Environment Agency
- EEA (2001): EMEP/CORINAIR Atmospheric Emission Inventory Guidebook - 3rd edition, Technical report No 30: EEA-European Environment Agency
- EEA (2007): EMEP/CORINAIR Atmospheric Emission Inventory Guidebook - 2007, Technical report No 16/2007: EEA-European Environment Agency
- EEA (2013): EMEP-EEA air pollutant emission inventory guidebook, <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (aviation factors in section 1.A.3.a Aviation annex.zip).
- EEA (2016): EMEP-EEA air pollutant emission inventory guidebook, <http://www.eea.europa.eu/publications/emep-eea-guidebook-2016>: EEA
- Emmett, B. A., Reynolds, B., Chamberlain, P. M., Rowe, E., Spurgeon, D., Brittain, S. A., Frogbrook, Z., Hughes, S., Lawlor, A. J., Poskitt, J., et al. (2007): Countryside Survey: Soils Report from 2007, CS Technical Report No. 9/07
- Engen-Skaugen, T., Haugen, J. E. & Hanssen-Bauer, I. (2008): Dynamically downscaled climate scenarios available at the Norwegian Meteorological Institute—per December 2008, Met.no report no. 24/2008: Norwegian Meteorological Institute (MET-Meteorologisk institutt)
- Esser, J. M. & Nyborg, Å. (1992): Jordsmonn i barskog - en oversikt for Norge., NIJOS rapport 3/92. Ås, Norway: Norwegian Institute of Land Inventory (Norsk institutt for jord- og skogkartlegging)
- EU (2002): Final technical report of the SACS2 project, issued 30. Aug. 2002, EU project NNE-1999-00521
- European Commission *Monitoring, reporting and verification of EU ETS emissions* European Union. Available at: http://ec.europa.eu/clima/policies/ets/monitoring/index_en.htm
- Exxonmobile (2009). Available at: http://www.oil.ie/files/documents/Product_Catalogue_Esso-Mobil_2009_No_Restriction.pdf
- Fedoryshyn, N. (2015): Beregning av CO₂-faktor for utslipp fra fossil del av avfall brent i forbrenningsanlegg. Dokumentasjon av metode. (Calculation of CO₂ factor for emissions from fossil fraction of incinerated waste. Documentation of method). Statistics Norway
- Finstad, A., Flugsrud, K. & Rypdal, K. (2002): Utslipp til luft fra norsk luftfart 1989-2000 (Emissions to Air from Norwegian Air Traffic 1989-2000), Reports 2002/08: Statistics Norway (SSB-Statistisk sentralbyrå). 32 pp
- Flugsrud, K. & Hoem, B. (2011): Uncertainties in the Norwegian greenhouse gas emission inventory, Reports 35/2011: Statistics Norway (SSB-Statistisk sentralbyrå). 51 pp
- Granhus, A., Hysten, G. & Nilsen, J.-E. Ø. (2012): Skogen i Norge. Statistikk over skogforhold og skogressurser i Norge registrert i perioden 2005-2009. . *Ressursoversikt fra Skog og landskap* 03/12. Ås: Norwegian Forest and Landscape Institute
- Grønland, A., Sturite, I., Riley, H., Fystro, G. & Lunnan, T. (2014): Nitrogen i restavlinger. 18 pp
- Grådahl, S., Johansen, S. T., Ravary, B., Andresen, B. & Tveit, H. (2007): Reduction of emissions from ferroalloy furnaces. In *Innovations in the Ferro Alloy Industry*, pp. 479-488. India: Macmillan
- Gundersen, G. I. & Rognstad, O. (2001): Lagring og bruk av husdyrgjødsel (Storage and use of manure), Reports 2001_39: Statistics Norway (SSB-Statistisk sentralbyrå)

- Gundersen, G. I. & Heldal, J. (2015): Bruk av gjødselressurser i jordbruket 2013. Metodebeskrivelse og resultater fra en utvalgsbasert undersøkelse (Use of inorganic and organic fertilisers in agriculture 2013). 84 pp
- Gustafsson, T. (2005): Improved structures for uncertainty analysis in the Swedish Greenhouse Gas Emission Inventory. *Commissioned report by the Swedish Environmental Protection Agency*, SMED Report No 69 2005: SMED-Swedish Methodology for Environmental Data
- Hansen, K. L. (2007): Emission from consumption of HFCs, PFCs and SF6 in Norway, Documents 2007/8: Statistics Norway (SSB-Statistisk sentralbyrå). 48 pp
- Hansson K, Helmisaari H-S, Sah SP & H, L. (2013): Fine root production and turnover of tree and understory vegetation in Scots pine, silver birch and Norway spruce stands in SW Sweden. *Forest Ecology and Management*, 309 (58-65)
- Hendriks, S. H., Fedoryshyn, N., Kittilsen, M. O. & Kolshus, K. (2017): Energiregnskap og -balanse. Dokumentasjon av statistikkproduksjonen fra statistikkår 2010 og fremover. (Energy accounts and balance. Documentation of statistics production for the years 2010 and forwards), Documents 2017/26: Statistics Norway (SSB - Statistisk sentralbyrå). 49 pp
- Hohle, E. E. (ed.) (2005): *Bioenergi. Miljø, teknikk og marked (Bio energy. Environment, technology and market)*, vol. 2. Brandbu: Energy Farm (Energigården-Senter for Bioenergi)
- Holmengen, N. & Kittilsen, M. O. (2009): Estimating emissions of NMVOC from solvent and other product use. Revised model, Reports 2009/14: Statistics Norway (SSB-Statistisk sentralbyrå). 77 pp
- Holmengen, N. & Fedoryshyn, N. (2015): Utslipp fra veitrafikken i Norge. Dokumentasjon av beregningsmetoder, data og resultater (Emissions from road traffic in Norway. Documentation of estimation methodologies, data and results): Statistics Norway
- Hutchings, N. J., Sommer, S. G., Andersen, J. M. & Asman, W. A. H. (2001): A detailed ammonia emission inventory for Denmark. *Atmospheric Environment*, 35 (11): 1959-1968
- Häkkinen, M., Heikkinen, J. & Mäkipää, R. (2011): Soil carbon stock increases in the organic layer of boreal middle-aged stands. *Biogeosciences*, 8: 1279-1289
- IAI (2005): The aluminium sector greenhouse gas protocol: International Aluminium Institute, IAI
- INFRAS (2017): Handbook emission factors for road transport (HBEFA), Version 3.3: INFRAS
- IPCC (1997a): Greenhouse Gas Inventory. Reference Manual (Volume 3). *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*
- IPCC (1997b): Greenhouse Gas Inventory. Workbook (Volume 2). *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*
- IPCC (2000): Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories: Prepared by the National Greenhouse Gas Inventories Programme, J Penman, D Kruger, I Galbally, T Hiraishi, B Nyenzi, S Emmanul, L Buendia, R Hoppaus, T Martinsen, J Meijer, K Miwa and K Tanabe. Published for the IPCC by the Institute for Global Environmental Strategies, Japan
- IPCC (2003): Good Practice Guidance for Land Use, Land-Use Change and Forestry: [Penman, J., M.Gytarsky, T. Hiraishi, T. Krug, D. Kruger, R. Pipatti, L. Buendia, K. Miwa, T. Ngara, K. Tanabe and F. Wagner (eds.)]. IPCC National Greenhouse Gas Inventories Programme. Institute for Global Environmental Strategies, Hayama, Kanagawa, Japan
- IPCC (2006): 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan
- IPCC (2014a): 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol. In Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. & Troxler, T. G. (eds). Switzerland: IPCC
- IPCC (2014b): 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. In Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. & Troxler, T. G. (eds). Switzerland: IPCC

- IVAM (2005): VOC emissions from cosmetics and cleaning agents. Amsterdam, the Netherlands: IVAM research and consultancy on sustainability
- Johansson L. Solvoll, K. (1999): Landsomfattende kostholdsundersøkelse blant menn og kvinner i alderen 16-79 år, Norkost 1997, Rapport nr. 2/1999: Statens råd for ernæring og fysisk aktivitet
- Karlengen, I. J., Svihus, B., Kjos, N. P. & Harstad, O. M. (2012): Husdyrgjødsel; oppdatering av mengder gjødsel og utskillelse av nitrogen, fosfor og kalium. Sluttrapport. (Manure; an update of amounts of manure and excretion of nitrogen, phosphorus and potassium. Final report). Ås: Departement of Animal and Aquacultural Sciences, Norwegian University of Life Sciences (Institutt for husdyr- og akvakulturvitenskap, NMBU-Norges miljø- og biovitenskapelige universitet)
- Kjønaas, O. J., Skre, O., Strand, L. T., Børja, I., Clarke, N., de Wit, H. A., Eldhuset, T. D. & Lange, H. (Manuscript): Understorey vegetation matters: Above- and belowground carbon and nitrogen pools and accumulation rates in a Norwegian Norway spruce forest chronosequence
- Kvingedal, E., Tornsjø, B., Holtskog, S., G., H. & Rypdal, K. (2000): Verification of the Norwegian Emission Inventory. Comparing emission intensity values with similar countries, TA-1736/00: Norwegian Pollution Control Authority (SFT-Statens forurensingstilsyn), Statistics Norway (SSB-Statistisk sentralbyrå)
- Larsson, J. Y. & Hylen, G. (2007): Skogen i Norge. Statistikk over skogforhold og skogressurser i Norge registrert i perioden 2000-2004 (Statistics of forest conditions and forest resources in Norway), Viten fra Skog og landskap 1/07: Norwegian Forest and Landscape Institute (Norsk institutt for skog og landskap). 91 pp
- Lehtonen, A., Sievänen, R., Mäkelä, A., Mäkipää, R., Korhonen, K. T. & Hokkanen, T. (2004): Potential litterfall of Scots pine branches in southern Finland. *Ecological Modelling*, 180 (2-3): 305-315
- Limberakis, E., Vay, J. & Gronberg, S. (1987): Ferroalloy industry particulate emissions: Source category report, EPA/600/S7-86/039. Washington, D.C: U.S. Environmental Protection Agency, EPA
- Liski, J., Palosuo, T., Peltoniemi, M. & Sievänen, R. (2005): Carbon and decomposition model Yasso for forest soils. *Ecological Modelling*, 189 (1-2): 168-182
- Liski, J., Tuomi, M. & Rasinmäki, J. (2009): Yasso07 user-interface manual: Finnish Environment Institute (SYKE-Suomen ympäristökeskus/ Finlands miljöcentral). 14 pp
- Lystad, H. (2005): Hjemmekompostering i kommunal regi – NRF medlemsundersøkelse. *NRF - Samarbeidsforum for avfallshåndtering*
- Løken (2012): Den totale biomasse av trær i Norge. Ås: Norwegian Forest and Landscape Institute
- MARINTEK (2010): Emissions factors for CH₄, NO_x, particulates and black carbon for domestic shipping in Norway, revision 1. *Commissioned report by Norwegian Climate and Pollution Agency (Klif-Klima- og forurensningsdirektoratet)*, Report no. 22232.00.02. TA-2746: Authors: Nielsen, J.B. and D. Stenersen, MARINTEK-Norwegian Marine Technology Research Institute. 35 pp
- Marklund, L. G. (1988): Biomassafunktioner för tall, gran och björk i Sverige: biomass functions for pine, spruce and birch in Sweden, Rapport 45: Swedish University of Agricultural Sciences (SLU-Sveriges lantbruksuniversitet, Institutionen för skogstaxering). 73 pp
- Marthinsen, J., Sandberg, K. & Johansen, M. (2010): Fornybar andel i avfall til norske forbrenningsanlegg - oppdatering: Avfall Norge. 6/42 pp
- Matamala, R., Gonzalez-Meler, M. A., Jastrow, J. D., Norby, R. J. & Schlesinger, W. H. (2003): Impacts of Fine Root Turnover on Forest NPP and Soil C Sequestration Potential. *Science*, 302 (5649): 1385-1387
- Miljødirektoratet (2015): *Rapportering og verifikasjon av utslipp* Miljødirektoratet. Available at: http://www.miljodirektoratet.no/no/Tema/klima/CO2_kvoter/Klimakvoter-for-industrien/Rapportering-og-verifikasjon-av-utslipp/

- Miljødirektoratet (2016): *Årlig rapportering av egenkontrollen for landbasert industri* Available at: <http://www.miljodirektoratet.no/no/Tjenester-og-verktoy/Skjema/landbasert/>
- Morken, J. (2003): Evaluering av ammoniakkutslippsmodellen (Evaluation of the ammonia emission model). Internal Note. Department of Agricultural Engineering. Norwegian University of Life Sciences (NMBU-Norges miljø- og biovitenskapelige universitet)
- Morken, J. & Nesheim, L. (2004): Utnytting og tap av næringsstoff i husdyrgjødsel - framtidige utfordringar. *Grønn kunnskap*, 8 (3): 51-64
- Morken, J. & Hoem, B. (2011): Models for calculating methane emission from manure management in Norway, IMT-Rapport 43/2011: Norwegian University of Life Sciences (NMBU-Norges miljø- og biovitenskapelige universitet). 18 pp
- Morken, J., Ayoub, S. & Sapci, Z. (2013): Revision of the Norwegian model for estimating methane emission from manure management, IMT-Rapport 54/2013: Norwegian University of Life Sciences (NMBU-Norges miljø- og biovitenskapelige universitet). 10 pp
- Muukkonen, P. & Lehtonen, A. (2004): Needle and branch biomass turnover rates of Norway spruce (*Picea abies*). *Canadian Journal of Forest Research*, 34: 2517-2527
- Muukkonen, P. & Mäkipää, R. (2006): Empirical biomass models of understory vegetation in boreal forests according to stand and site attributes. *Boreal Environment Research*, 11: 355-369
- Muukkonen, P., Mäkipää, R., Laiho, R., Minkinen, K., Vasander, H. & Finer, L. (2006): Relationship between biomass and percentage cover in understory vegetation of boreal coniferous forests. *Silva Fennica*, 40 (2): 231-245
- NCASI (2004): Critical review of forest products decomposition in municipal solid waste landfills. *Technical Bulletin No. 872*. Research Triangle Park, N.C.: National Council for Air and Stream Improvement, Inc. 71 pp
- Neste_Oil (2014): <http://www.nesteoil.com/default.asp?path=1,41,535,547,14077,5604> (Products and services/Lubricants and car chemicals/Basic concepts of lubricants).
- NFLI (2008): Landsskogtakseringens feltinstruks 2008, Håndbok fra Skog og landskap 05/2008: Norwegian Forest and Landscape Institute (Norsk institutt for skog og landskap)
- Nielsen, O.-K., Lyck, E., Mikkelsen, M. H., Hoffmann, L., Gyldenkerne, S., Winther, M., Nielsen, M., Fauser, P., Thomsen, M., Plejdrup, M. S., et al. (2010): Denmark's National Inventory Report 2010. Emission Inventories 1990-2008 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol, NERI Technical Report No 784. Aarhus University, Denmark: National Environmental Research Institute of Denmark (DMU-Danmarks Miljøundersøgelser). 1178 pp
- Nilsson, M., Sagerfors, J., Buffam, I., Laudon, H., Eriksson, T., Grelle, A., Klemetsson, L., Weslien, P. E. R. & Lindroth, A. (2008): Contemporary carbon accumulation in a boreal oligotrophic minerogenic mire – a significant sink after accounting for all C-fluxes. *Global Change Biology*, 14 (10): 2317-2332
- Norddal, T. (2010): Miljørapport for innenriks ferjetrafikk 2009: Norwegian Maritime Authority (Sjøfartsdirektoratet), Norwegian Public Roads Administration (Statens vegvesen), Federation of Norwegian Coastal Shipping (Rederienes Landsforening)
- Norsk olje&gass (2012): *Veiledning til den årlige utslippsrapporteringen*: Norsk olje&gass. Available at: <http://www.norskoljeoggass.no/no/Publikasjoner/Miljorapporter/Veiledning-utslippsrapportering-2012/>
<https://www.norskoljeoggass.no/contentassets/cd872e74e25a4aadac1a6e820e7f5f95/044-retningslinje-for-utslippsrapportering.pdf>
- Norwegian Climate and Pollution Agency (2011): CO₂-faktor for fossil del av avfallet som blir brent i avfallsforbrenningsanlegg i Norge (CO₂ emission factor for the fossil part of waste combusted in waste incineration plants in Norway), Note, Eilev Gjerald, 20.01.2011: Norwegian Climate and Pollution Agency (Klif-Klima- og forurensningsdirektoratet)

- Norwegian Environment Agency (2016): Vellykket samarbeid om å redusere utslipp til luft fra norsk sokkel. Available at: <http://www.miljodirektoratet.no/no/Nyheter/Nyheter/2016/Juni-2016/Vellykket-samarbeid-om-a-redusere-utslipp-til-luft-fra-norsk-sokkel/>:
- Norwegian Environment Agency (2018): Informative Inventory Report (IIR) 2018. Norway. Air Pollutant Emissions 1990-2016 Report M-967/2018
- Norwegian Food Safety Authority (Annually): Mineralgjødselstatistikk: Norwegian Food Safety Authority (Mattilsynet)
- OED (2017): *Norwegian Petroleum - Everything you need to know about Norwegian Petroleum activities*. Available at: <http://www.norskpetroleum.no/en/>
- Ogle, S. M., Conant, R. T. & Paustian, K. (2004): Deriving Grassland Management Factors for a Carbon Accounting Method Developed by the Intergovernmental Panel on Climate Change. *Environmental Management*, 33 (4): 474-484
- OLF (1993): OLF Environmental Programme. Summary report Phase II. Stavanger: Norwegian Oil Industry Association (OLF-Oljeindustriens landsforening). 134 pp
- OLF (1994): Anbefalte retningslinjer for utslippsberegning. Identifisering, kvantifisering og rapportering av forbruks- og utslippsdata fra aktiviteter i norsk oljevirkksomhet (Recommended guidelines for emission calculations. Identification, quantification and reporting of data on consumption and emissions from activities in the Norwegian oil and gas sector). Stavanger: Norwegian Oil Industry Association (OLF-Oljeindustriens landsforening)
- OLF (2009): Veiledning til den Årlige Utslippsrapporteringen: Norwegian Oil Industry Association (OLF-Oljeindustriens landsforening). 80 pp
- Olivier, J. G. J. (1991): Inventory of aircraft Emissions: A review of Recent Literature, RIVM Report 736301008. Bilthoven, Netherlands: National Institute of Public Health and Environmental Protection (RIVM-Rijksinstituut voor Volksgezondheid en Milieu). 131 pp
- Paustian, K., Six, J., Elliott, E. T. & Hunt, H. W. (2000): Management options for reducing CO₂ emissions from agricultural soils. *Biogeochemistry*, 48 (1): 147-163
- Peltoniemi, M., Mäkipää, R., Liski, J. & Tamminen, P. (2004): Changes in soil carbon with stand age – an evaluation of a modelling method with empirical data. *Global Change Biology*, 10 (12): 2078-2091
- Peltoniemi, M., Palosuo, T., Monni, S. & Mäkipää, R. (2006): Factors affecting the uncertainty of sinks and stocks of carbon in Finnish forests soils and vegetation. *Forest Ecology and Management*, 232 (1–3): 75-85
- Petersen, S. O. & Olesen, J. E. e. (2002): Greenhouse gas inventories for agriculture in the nordic countries. In *Proceedings from an international workshop, Helsingør, Denmark 24-25 Jan. 2002*, p. 157: DIAS report, Plant Production, no. 81
- Petersson, H. & Ståhl, G. (2006): Functions for below-ground biomass of *Pinus sylvestris*, *Picea abies*, *Betula pendula* and, *Betula pubescens* in Sweden. *Scandinavian Journal of Forest Research*, 21 (S7): 84-93
- R Core Team (2015): *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing
- Refsgard Andersen, H. (1990): Ammekoens energibehov og foderoptagelseskapaet. (In danish). Landhusholdningsselskabets forl. Copenhagen
- Riley, H. (1996): *Estimation of physical properties of cultivated soils in southeast Norway from readily available soil information*. Norwegian Journal of Agricultural Science/ Supplement, vol. no. 25 1996. Ås: Norwegian Agricultural Advisory Centre. 51 s. : ill. pp
- Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T. R., Humphreys, E. R. & Bubier, J. (2007): Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland. *Global Change Biology*, 13 (2): 397-411
- Rypdal, K. (1999): Evaluation of uncertainty in the Norwegian emission inventory, SFT-report 99:01. TA-1609/99: Norwegian Pollution Control Authority (SFT-Statens forurensingstilsyn), Statistics Norway (SSB-Statistisk sentralbyrå). 58 pp

- Rypdal, K. & Zhang, L.-C. (2000): Uncertainties in the Norwegian greenhouse Gas Emission Inventory, Reports 2000/13: Statistics Norway (SSB-Statistisk sentralbyrå). 44 pp
- Rypdal, K. & Zhang, L.-C. (2001): Uncertainties in Emissions of Long-Range Air Pollutants, Reports 2001/37: Statistics Norway (SSB-Statistisk sentralbyrå). 49 pp
- Rypdal, K., Bloch, V. V. H., Flugsrud, K., Gobakken, T., Hoem, B., Tomter, S. & Aalde, H. (2005): Emissions and removals of greenhouse gases from land use, land-use change and forestry in Norway. *Commissioned report by Norwegian Pollution Control Authority (SFT-Statens Forurensningstilsyn) and Ministry of Agriculture and Food (Landbruks- og matdepartementet)*, NIOS rapport 11/05: Center for International Climate and Environmental Research (CICERO-Senter for klimaforskning, Norwegian Institute of Land Inventory (NIOS-Norsk institutt for jord- og skogkartlegging), Statistics Norway (SSB-Statistisk sentralbyrå). 105 pp
- Sanderman, J. & Baldock, J. A. (2010): Accounting for soil carbon sequestration in national inventories: a soil scientist's perspective. *Environmental Research Letters*, 5 (3): 1-6
- SFT (1990): Klimagassregnskap for Norge. Beskrivelse av utslippsmengder, drivhusstyrke og utslippsfaktorer. Bidrag til den interdepartementale klimautredningen (Greenhouse gas inventory for Norway. Emission figures, global warming potentials and emission factors. Contribution to the interministerial climate report). Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (1993): Emissions from road traffic in Norway – Methods for estimation, input data and emission estimates. *Authors: Bang, J., Figenbaum, E., Flugsrud, K., Larssen, S., Rypdal, K., Torp, C.*, SFT report 93:12. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (1996): Utslipp ved håndtering av kommunalt avfall (Emissions from municipal waste management). *Authors: Sandgren J., Heie, A. and Sverud, T.*, SFT-report 96:16. TA-1366/96. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (1999a): Beregningsmodell for utslipp av metangass fra norske deponier (Model for estimation of CH₄ from landfills in Norway). *Commissioned report by Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn). Authors: Jensen, J. E. F., Williksen, T. and Bartnes, J., Norconsult AS*, SFT-report 99:16. TA-1693/99. Oslo. 70 pp
- SFT (1999b): Calculations of emissions of HFCs and PFCs in Norway. Tier 2 method. *Authors: Haukås, H. T. and Holtskog, S. (Statistics Norway), Pettersen, M. V. (Norwegian Pollution Control Authority)*, SFT-report 99:03. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (1999c): Materialstrømsanalyse av SF₆. Beregning av potensielt og faktisk utslipp over tid (Material flow analysis of SF₆. Calculation of potential and actual emissions over time), SFT-report 99:14. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (1999d): Utslipp fra veitrafikk i Norge. Dokumentasjon av beregningsmetode, data og resultater (Emissions from road traffic in Norway - Method for estimation, input data and emission estimates). *Authors: Bang, J., Flugsrud, K., Haakonsen, G., Holtskog, S., Larssen, S., Maldum, K.O., Rypdal, K. and A. Skedsmo*, SFT-report 99:04 (Updated SFT report 93:12). TA-1622/99. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (2005a): Kartlegging av omsetning av enkelte miljøskadelige stoffer i legemidler og kosmetikk (Survey of turnover of some environmentally hazardous substances in pharmaceuticals and cosmetics), TA-2128/05. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (2005b): Methane emissions from solid waste disposal sites, TA-2079/05. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn)
- SFT (2006): Documentation of methodology and results: QA/QC performed for Greenhouse Gas Emissions for Industrial plants included in the National Inventory. Oslo: Norwegian Pollution Control Authority (SFT-Statens forurensningstilsyn). 42 pp

- SINTEF (1995): Round robin test of a wood stove-emissions. *Author: Karlsvik, E.*, SINTEF Report STF12 A95012. Trondheim: SINTEF, Thermal Energy and Fluid Machinery. 62 pp
- SINTEF (1998a): Omregningsfaktorer for CO₂-utslipp fra metallurgisk industri og sementproduksjon. Delprosjekt 1. CO₂-utslipp fra forskjellige typer reduksjonsmaterialer (Conversion factors for CO₂ emissions from metal manufacturing and cement production. Part 1. CO₂ emissions from various types of reducing agents). *Author: Raaness, O.*, Revised edition, SINTEF Report STF24 A98550. Trondheim: SINTEF
- SINTEF (1998b): Omregningsfaktorer for CO₂-utslipp fra metallurgisk industri og sementproduksjon. Delprosjekt 2. Produksjon av ferrosilisium og silisiummetall i Norge (Conversion factors for CO₂ emissions from metal manufacturing and cement production. Part 2. Production of ferro silicon and silicon metal in Norway). *Author: Monsen, B.*, Revised edition, SINTEF Report STF24 A98537. Trondheim: SINTEF
- SINTEF (1998c): Omregningsfaktorer for CO₂-utslipp fra metallurgisk industri og sementproduksjon. Delprosjekt 3. Produksjon av ferromangan, silikomangan og ferrokrom i Norge (Conversion factors for CO₂ emissions from metal manufacturing and cement production. Part 3. Production of ferro manganese, silicon manganese and ferro chromium in Norway). *Authors: Monsen, B. and Olsen, S.E.*, SINTEF Report STF24 A98548. Trondheim: SINTEF
- SINTEF (1998d): Omregningsfaktorer for CO₂-utslipp fra metallurgisk industri og sementproduksjon. Delprosjekt 4. Utslipp av CO₂ ved produksjon av silisiumkarbid og kalsiumkarbid (Conversion factors for CO₂ emissions from metal manufacturing and cement production. Part 4. CO₂ emissions from production of silicon carbide and calcium carbide). *Authors: Raaness, O. and Olsen, S.*, SINTEF Report STF24 A98549. Trondheim: SINTEF
- SINTEF (1998e): Omregningsfaktorer for CO₂-utslipp fra metallurgisk industri og sementproduksjon. Delprosjekt 5. Produksjon av magnesium (Conversion factors for CO₂ emissions from metal manufacturing and cement production. Part 5. Magnesium production). *Authors: Olsen, K., Støre, T. and Tunold, R.*, SINTEF Report STF24 A98574. Trondheim: SINTEF
- SINTEF (2004): Utslipp av klimagasser fra Si-prosessen hos Elkem Thamshavn og Holla Metall med fokus på NO_x reduksjon (Emissions of GHG from Si-process at Elkem Thamshavn and Holla Metall with focus on reduction of emissions of NO_x), SINTEF Report STF24 F03555: SINTEF
- Skårman, T., Danielsson, H., Henningsson, E. & Östmann, M. (2006): Revised Method for Estimating Emissions of NMVOC from Solvent and Other Product Use in Sweden. *Commissioned report by the Swedish Environmental Protection Agency*, SMED Report No 75 2006: SMED Report No 75 2006
- Smith, A., Granhus, A., Astrup, R. (2016): Functions for estimating belowground and whole tree biomass of birch in Norway. *Scandinavian Journal of Forest Research*, 31 (6): 568-582
- Smith, A., Granhus, A., Astrup, R., Bollandsås, O.M., Petersson, H., (2014): Functions for estimating aboveground biomass of birch in Norway. *Scandinavian Journal of Forest Research*, 29 (6): 565-578.
- SSB (2015): *Structure in silviculture, 2013. Struktur i skogbruket, 2013 (In Norwegian)*. Published september 30th 2014: Statistics Norway (SSB). Available at: <http://ssb.no/jord-skog-jakt-og-fiskeri/statistikker/stskog/aar> (accessed: 10. March)
- Statistics Norway (2001): Natural resources and the environment 2001, Statistical Analyses 47: Statistics Norway (SSB-Statistisk sentralbyrå). 293 pp
- Statistics Norway (2002): Sample survey of agriculture and forestry 2001: Statistics Norway (SSB-Statistisk sentralbyrå)
- Statistics Norway (2004): Sample survey of agriculture and forestry 2003: Statistics Norway (SSB-Statistisk sentralbyrå)
- Statistics Norway (2007): Sample survey of agriculture and forestry 2006: Statistics Norway (SSB-Statistisk sentralbyrå)

- Statistics Norway (2008): Standard Industrial Classification, NOS report D 383: Statistics Norway. http://www.ssb.no/a/publikasjoner/pdf/nos_d383/nos_d383.pdf (English version: <http://www.ssb.no/en/klasse/#!/klassifikasjoner/6>)
- Statistics Norway (2010a): The Norwegian Emission Inventory 2010. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. In Sandmo, T. (ed.), Documents 2010/21: Statistics Norway (SSB-Statistisk sentralbyrå). 236 pp
- Statistics Norway (2010b): *Road goods transport*: Statistics Norway (SSB-Statistisk sentralbyrå). Available at: http://www.ssb.no/english/subjects/10/12/20/lbunasj_en/ (accessed: 2010)
- Statistics Norway (2012): *Energy use in the manufacturing sector, 2012*. Statistics Norway (SSB-Statistisk sentralbyrå). Available at: <https://www.ssb.no/en/energi-og-industri/statistikker/indenergi/aar/2013-06-27?fane=om#content>
- Statistics Norway (2013): The Norwegian Emission Inventory 2013. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. In Sandmo, T. (ed.), Documents 2013/30: Statistics Norway (SSB-Statistisk sentralbyrå). 262 pp
- Statistics Norway (2014a): *Discharges and treatment of municipal waste water*: Statistics Norway. Available at: <https://www.ssb.no/statistikkbanken/selecttable/hovedtabellHjem.asp?KortNavnWeb=avlut&CMSSubjectArea=natur-og-miljo&PLanguage=1&checked=true>
- Statistics Norway (2014b): The Norwegian Emission Inventory 2014. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. In Sandmo, T. (ed.), Documents 2014/35: Statistics Norway (SSB-Statistisk sentralbyrå). 278 pp
- Statistics Norway. (2014c). Standard Industrial Classification (SIC2007). In *Statistics Norway* (SSB-Statistisk sentralbyrå). Available at: <http://stabas.ssb.no/itemsFrames.asp?ID=8118001&Language=en&VersionLevel=ClassVersion>.
- Statistics Norway & SOE Norway (2014): Estimations made in Statistics Norway's hazardous waste statistics, <http://www.ssb.no/en/natur-og-miljo/statistikker/spesavf>, and published on the Internet site of State of the Environment Norway, <http://www.miljostatus.no/Tema/Avfall/Farlig-avfall/Spillolje/#D> (click 'Les mer om tallene' below the figure).
- Statistics Norway (2015): *Sales of petroleum products, 2015, final figures*: Statistics Norway (SSB-Statistisk sentralbyrå)
- Statistics Norway (2016a): The Norwegian Emission Inventory 2016. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. In Sandmo, T. (ed.): Statistics Norway
- Statistics Norway (2016b): Statistical differences for primary energy products. Norwegian Energy Balance 2011-2014. <http://www.ssb.no/energi-og-industri/artikler-og-publikasjoner/statistical-differences-for-primary-energy-products>. In Skullerud, H. F. (ed.): Statistics Norway (SSB-Statistisk sentralbyrå)
- Statistics Norway (Annually-a): *Production and consumption of energy, energy balance*: Statistics Norway (SSB-Statistisk sentralbyrå). Available at: <https://www.ssb.no/en/energi-og-industri/statistikker/energibalanse/aar-enderlige>
- Statistics Norway (Annually-b): *Waste accounts*. Statistics Norway (SSB-Statistisk sentralbyrå). Available at: <http://www.ssb.no/en/natur-og-miljo/statistikker/avfregno/aar/2015-06-16?fane=om#content>
- Storlien, T. M. & Harstad, O. M. (2015): Enteric methane emissions from the cattle population in Norway. Method description. (Included in NIR 2016, Annex IX): Norwegian University of Life Sciences. Department of Animal and Aquacultural Sciences

- Strand, L. T., Callesen, I., Dalsgaard, L. & de Wit, H. A. (2016): Carbon and nitrogen stocks in Norwegian forest soils — the importance of soil formation, climate, and vegetation type for organic matter accumulation. *Canadian Journal of Forest Research*, 46 (12): 1459-1473
- Sundstøl, F. & Mroz, Z. (1988): Utskillelse av nitrogen og fosfor i gjødsel og urin fra husdyr i Norge (Nitrogen and phosphorus in manure and urine from domestic animals in Norway), Report no. 4 from the project "Agricultural policy and environmental management". Ås: Agricultural University of Norway (Norges landbrukshøgskole)
- Svihus, B. (2015): Production of methane from enteric fermentation in poultry in Norway: Norwegian University of Life sciences. 2 pp
- Swedish Environmental Protection Agency (2011): National inventory report 2011 Sweden - Annexes. Stockholm, Sweden: Swedish Environmental Protection Agency (Naturvårdsverket)
- Søgaard, G. & Økseter, R. (2017): Klimagassutslipp fra torvproduksjon i Norge. Metode, datagrunnlag og utslippfaktorer benyttet i klimagassregnskapet under FNs klimakonvensjon (UNFCCC). Ås, Norway: NIBIO
- TI (1991): Tiltak 11. Reduksjon av VOC-utslipp fra totaktsmotorer. (Reduction of VOC-emissions from two stroke engines). *Commissioned report by Norwegian Pollution Control Authority (SFT-Statens Forurensningstilsyn)*. Oslo: National Institute of Technology (TI-Teknologisk Institutt)
- Tierney, G. L. & Fahey, T. J. (2002): Fine root turnover in a northern hardwood forest: a direct comparison of the radiocarbon and minirhizotron methods. *Canadian Journal of Forest Research*, 32 (9): 1692-1697
- TINE BA (Annually): Nøkkeltall fra Kukontrollen: TINE
- Thovsen, K. (2018): Documentation of calculation method for emissions and energy use in aviation (in Norwegian). Unpublished document Statistics Norway
- Tokheim, L. A. (2006): Usikkerhet i utslippsfaktorer ifm. rapportering av CO₂-utslipp, Report 9A-R06-002: Norcem Brevik
- Tolonen, K. & Turunen, J. (1996): Accumulation rates of carbon in mires in Finland and implications for climate change. *The Holocene*, 6 (2): 171-178
- Tornsjø, B. (2001): Utslipp til luft fra innenriks sjøfart, fiske og annen sjøtrafikk mellom norske havner. (Emissions to Air from Fishing Fleet and Sea Traffic between Norwegian Harbours), Reports 2001/6: Statistics Norway (SSB-Statistisk sentralbyrå). 36 pp
- Tuomi, M., Vanhala, P., Karhu, K., Fritze, H. & Liski, J. (2008): Heterotrophic soil respiration—Comparison of different models describing its temperature dependence. *Ecological Modelling*, 211 (1–2): 182-190
- Tuomi, M., Thum, T., Järvinen, H., Fronzek, S., Berg, B., Harmon, M., Trofymow, J. A., Sevanto, S. & Liski, J. (2009): Leaf litter decomposition—Estimates of global variability based on Yasso07 model. *Ecological Modelling*, 220 (23): 3362-3371
- Tuomi, M., Laiho, R., Repo, A. & Liski, J. (2011a): Wood decomposition model for boreal forests. *Ecological Modelling*, 222 (3): 709-718
- Tuomi, M., Rasinmäki, J., Repo, A., Vanhala, P. & Liski, J. (2011b): Soil carbon model Yasso07 graphical user interface. *Environmental Modelling & Software*, 26 (11): 1358-1362
- Turunen, J., Tomppo, E., Tolonen, K. & Reinikainen, A. (2002): Estimating carbon accumulation rates of undrained mires in Finland – application to boreal and subarctic regions. *The Holocene*, 12 (1): 69-80
- Vibe, N. (1993): Norske reisevaner. Dokumentasjonsrapport for den landsomfattende reisevaneundersøkelsen 1991-92, TØI rapport 183/1993. Oslo: Institute of Transport Economics (TØI-Transportøkonomisk institutt). 67 pp
- Vågane, L. & Rideng, A. (2010): Transportytelser i Norge 1946-2009, TØI Report 1090/2010. Oslo: Institute of Transport Economics (TØI-Transportøkonomisk institutt)
- Weholt, Ø., Aakre, S. & Dehlin, F. (2010): Utvikling av en modell for å beregne årlig generert mengde spillolje (developing a model for estimating annual generated amounts of waste oil). TA -

2739/2010 (in Norwegian), Norwegian Climate and Pollution Agency (current Norwegian Environment Agency).

Winther, M. & Nielsen, O.-K. (2006): Fuel use and emissions from non-road machinery in Denmark from 1985-2004 - and projections from 2005-2030, Environmental project no. 1092 2006: National environmental research institute, Danish ministry of the environment

Norwegian Environment Agency

Telephone: +47 73 58 05 00 | **Fax:** +47 73 58 05 01

E-mail: post@miljodir.no

Web: www.miljodirektoratet.no

Postal address: Postboks 5672 Torgarden, N-7485 Trondheim

Visiting address Trondheim: Brattørkaia 15, 7010 Trondheim

Visiting address Oslo: Grensesvingen 7, 0661 Oslo

The Norwegian Environment Agency is working for a clean and diverse environment. Our primary tasks are to reduce greenhouse gas emissions, manage Norwegian nature, and prevent pollution.

We are a government agency under the Ministry of Climate and Environment and have 700 employees at our two offices in Trondheim and Oslo and at the Norwegian Nature Inspectorate's more than sixty local offices.

We implement and give advice on the development of climate and environmental policy. We are professionally independent. This means that we act independently in the individual cases that we decide and when we communicate knowledge and information or give advice.

Our principal functions include collating and communicating environmental information, exercising regulatory authority, supervising and guiding regional and local government level, giving professional and technical advice, and participating in international environmental activities.